

Computer algebra independent integration tests

4-Trig-functions/4.5-Secant/4.5.0-a-sec-^m-b-trg-ⁿ

Nasser M. Abbasi

July 17, 2021

Compiled on July 17, 2021 at 3:56pm

Contents

1	Introduction	13
1.1	Listing of CAS systems tested	13
1.2	Results	14
1.3	Performance	18
1.4	list of integrals that has no closed form antiderivative	19
1.5	list of integrals solved by CAS but has no known antiderivative	20
1.6	list of integrals solved by CAS but failed verification	20
1.7	Timing	21
1.8	Verification	21
1.9	Important notes about some of the results	21
1.9.1	Important note about Maxima results	21
1.9.2	Important note about FriCAS and Giac/XCAS results	22
1.9.3	Important note about finding leaf size of antiderivative	22
1.9.4	Important note about Mupad results	23
1.10	Design of the test system	24
2	detailed summary tables of results	25
2.1	List of integrals sorted by grade for each CAS	25
2.1.1	Rubi	25
2.1.2	Mathematica	25
2.1.3	Maple	26
2.1.4	Maxima	26

2.1.5	FriCAS	27
2.1.6	Sympy	27
2.1.7	Giac	28
2.1.8	Mupad	28
2.2	Detailed conclusion table per each integral for all CAS systems	30
2.3	Detailed conclusion table specific for Rubi results	90

3	Listing of integrals	103
3.1	$\int \sec(a + bx) dx$	103
3.2	$\int \sec^2(a + bx) dx$	106
3.3	$\int \sec^3(a + bx) dx$	109
3.4	$\int \sec^4(a + bx) dx$	112
3.5	$\int \sec^5(a + bx) dx$	115
3.6	$\int \sec^6(a + bx) dx$	118
3.7	$\int \sec^7(a + bx) dx$	121
3.8	$\int \sec^8(a + bx) dx$	125
3.9	$\int \sec^{\frac{7}{2}}(a + bx) dx$	128
3.10	$\int \sec^{\frac{5}{2}}(a + bx) dx$	132
3.11	$\int \sec^{\frac{3}{2}}(a + bx) dx$	135
3.12	$\int \sqrt{\sec(a + bx)} dx$	138
3.13	$\int \frac{1}{\sqrt{\sec(a+bx)}} dx$	141
3.14	$\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$	144
3.15	$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$	148
3.16	$\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx$	152
3.17	$\int (c \sec(a + bx))^{7/2} dx$	156
3.18	$\int (c \sec(a + bx))^{5/2} dx$	160
3.19	$\int (c \sec(a + bx))^{3/2} dx$	163
3.20	$\int \sqrt{c \sec(a + bx)} dx$	167
3.21	$\int \frac{1}{\sqrt{c \sec(a+bx)}} dx$	170
3.22	$\int \frac{1}{(c \sec(a+bx))^{3/2}} dx$	173
3.23	$\int \frac{1}{(c \sec(a+bx))^{5/2}} dx$	177
3.24	$\int \frac{1}{(c \sec(a+bx))^{7/2}} dx$	181
3.25	$\int \sec^{\frac{4}{3}}(a + bx) dx$	185
3.26	$\int \sec^{\frac{2}{3}}(a + bx) dx$	188

3.27	$\int \sqrt[3]{\sec(a + bx)} dx$	191
3.28	$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx$	194
3.29	$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx$	197
3.30	$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx$	200
3.31	$\int (c \sec(a + bx))^{4/3} dx$	203
3.32	$\int (c \sec(a + bx))^{2/3} dx$	206
3.33	$\int \sqrt[3]{c \sec(a + bx)} dx$	209
3.34	$\int \frac{1}{\sqrt[3]{c \sec(a+bx)}} dx$	212
3.35	$\int \frac{1}{(c \sec(a+bx))^{2/3}} dx$	215
3.36	$\int \frac{1}{(c \sec(a+bx))^{4/3}} dx$	218
3.37	$\int \sec^n(a + bx) dx$	221
3.38	$\int (c \sec(a + bx))^n dx$	224
3.39	$\int \sec^2(x)^{7/2} dx$	227
3.40	$\int \sec^2(x)^{5/2} dx$	230
3.41	$\int \sec^2(x)^{3/2} dx$	233
3.42	$\int \sqrt{\sec^2(x)} dx$	236
3.43	$\int \frac{1}{\sqrt{\sec^2(x)}} dx$	239
3.44	$\int \frac{1}{\sec^2(x)^{3/2}} dx$	242
3.45	$\int \frac{1}{\sec^2(x)^{5/2}} dx$	245
3.46	$\int \frac{1}{\sec^2(x)^{7/2}} dx$	248
3.47	$\int (a \sec^2(x))^{7/2} dx$	252
3.48	$\int (a \sec^2(x))^{5/2} dx$	257
3.49	$\int (a \sec^2(x))^{3/2} dx$	261
3.50	$\int \sqrt{a \sec^2(x)} dx$	265
3.51	$\int \frac{1}{\sqrt{a \sec^2(x)}} dx$	268
3.52	$\int \frac{1}{(a \sec^2(x))^{3/2}} dx$	271
3.53	$\int \frac{1}{(a \sec^2(x))^{5/2}} dx$	275
3.54	$\int \frac{1}{(a \sec^2(x))^{7/2}} dx$	279
3.55	$\int (a \sec^3(x))^{5/2} dx$	283
3.56	$\int (a \sec^3(x))^{3/2} dx$	287

3.57	$\int \sqrt{a \sec^3(x)} dx$	291
3.58	$\int \frac{1}{\sqrt{a \sec^3(x)}} dx$	295
3.59	$\int \frac{1}{(a \sec^3(x))^{3/2}} dx$	299
3.60	$\int \frac{1}{(a \sec^3(x))^{5/2}} dx$	303
3.61	$\int (a \sec^4(x))^{7/2} dx$	307
3.62	$\int (a \sec^4(x))^{5/2} dx$	311
3.63	$\int (a \sec^4(x))^{3/2} dx$	314
3.64	$\int \sqrt{a \sec^4(x)} dx$	317
3.65	$\int \frac{1}{\sqrt{a \sec^4(x)}} dx$	320
3.66	$\int \frac{1}{(a \sec^4(x))^{3/2}} dx$	323
3.67	$\int \frac{1}{(a \sec^4(x))^{5/2}} dx$	327
3.68	$\int ((b \sec(c + dx))^p)^n dx$	331
3.69	$\int (a(b \sec(c + dx))^p)^n dx$	334
3.70	$\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx$	337
3.71	$\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx$	341
3.72	$\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx$	345
3.73	$\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx$	349
3.74	$\int \sqrt{b \sec(c + dx)} dx$	353
3.75	$\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx$	356
3.76	$\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx$	359
3.77	$\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx$	363
3.78	$\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx$	367
3.79	$\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx$	371
3.80	$\int \sec^3(c + dx) (b \sec(c + dx))^{3/2} dx$	375
3.81	$\int \sec^2(c + dx) (b \sec(c + dx))^{3/2} dx$	379
3.82	$\int \sec(c + dx) (b \sec(c + dx))^{3/2} dx$	383
3.83	$\int (b \sec(c + dx))^{3/2} dx$	387
3.84	$\int \cos(c + dx) (b \sec(c + dx))^{3/2} dx$	391
3.85	$\int \cos^2(c + dx) (b \sec(c + dx))^{3/2} dx$	394
3.86	$\int \cos^3(c + dx) (b \sec(c + dx))^{3/2} dx$	397
3.87	$\int \cos^4(c + dx) (b \sec(c + dx))^{3/2} dx$	400
3.88	$\int \cos^5(c + dx) (b \sec(c + dx))^{3/2} dx$	404
3.89	$\int \cos^6(c + dx) (b \sec(c + dx))^{3/2} dx$	408

3.90	$\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx$	412
3.91	$\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx$	416
3.92	$\int (b \sec(c + dx))^{5/2} dx$	420
3.93	$\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx$	423
3.94	$\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx$	427
3.95	$\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx$	430
3.96	$\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx$	433
3.97	$\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx$	437
3.98	$\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx$	441
3.99	$\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx$	445
3.100	$\int (b \sec(c + dx))^{7/2} dx$	449
3.101	$\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	453
3.102	$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	457
3.103	$\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	461
3.104	$\int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	465
3.105	$\int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	469
3.106	$\int \frac{1}{\sqrt{b \sec(c+dx)}} dx$	472
3.107	$\int \frac{\cos(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	475
3.108	$\int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	479
3.109	$\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	483
3.110	$\int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	487
3.111	$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	491
3.112	$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	495
3.113	$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	499
3.114	$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	503
3.115	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	507
3.116	$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	510
3.117	$\int \frac{1}{(b \sec(c+dx))^{3/2}} dx$	514
3.118	$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	518
3.119	$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	522

3.120	$\int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	526
3.121	$\int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	530
3.122	$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	534
3.123	$\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	538
3.124	$\int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	542
3.125	$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	546
3.126	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	549
3.127	$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	553
3.128	$\int \frac{1}{(b \sec(c+dx))^{5/2}} dx$	557
3.129	$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	561
3.130	$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$	565
3.131	$\int \frac{1}{(b \sec(c+dx))^{7/2}} dx$	569
3.132	$\int \sec^{\frac{9}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	573
3.133	$\int \sec^{\frac{7}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	578
3.134	$\int \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	582
3.135	$\int \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx$	586
3.136	$\int \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} dx$	589
3.137	$\int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx$	592
3.138	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx$	595
3.139	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$	598
3.140	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx$	602
3.141	$\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx$	606
3.142	$\int \sec^{\frac{7}{2}}(c+dx) (b \sec(c+dx))^{3/2} dx$	610
3.143	$\int \sec^{\frac{5}{2}}(c+dx) (b \sec(c+dx))^{3/2} dx$	615
3.144	$\int \sec^{\frac{3}{2}}(c+dx) (b \sec(c+dx))^{3/2} dx$	619
3.145	$\int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{3/2} dx$	623
3.146	$\int \frac{(b \sec(c+dx))^{3/2}}{\sqrt{\sec(c+dx)}} dx$	626

3.147	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx$	629
3.148	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^5(c+dx)} dx$	632
3.149	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^7(c+dx)} dx$	635
3.150	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^9(c+dx)} dx$	639
3.151	$\int \frac{(b \sec(c+dx))^{3/2}}{\sec^{11}(c+dx)} dx$	643
3.152	$\int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2} dx$	647
3.153	$\int \sec^3(c+dx)(b \sec(c+dx))^{5/2} dx$	651
3.154	$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^{5/2} dx$	655
3.155	$\int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx$	659
3.156	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^3(c+dx)} dx$	662
3.157	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^5(c+dx)} dx$	665
3.158	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^7(c+dx)} dx$	668
3.159	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^9(c+dx)} dx$	671
3.160	$\int \frac{(b \sec(c+dx))^{5/2}}{\sec^{11}(c+dx)} dx$	675
3.161	$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	679
3.162	$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	683
3.163	$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx$	686
3.164	$\int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx$	689
3.165	$\int \frac{1}{\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}} dx$	692
3.166	$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)}} dx$	695
3.167	$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}} dx$	699
3.168	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$	703

3.169	$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$	707
3.170	$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$	710
3.171	$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$	713
3.172	$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{\frac{3}{2}}} dx$	716
3.173	$\int \frac{1}{\sqrt{\sec(c+dx)} (b \sec(c+dx))^{\frac{3}{2}}} dx$	719
3.174	$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{\frac{3}{2}}} dx$	723
3.175	$\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{\frac{3}{2}}} dx$	727
3.176	$\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$	731
3.177	$\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$	735
3.178	$\int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$	738
3.179	$\int \frac{\sec^{\frac{5}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$	741
3.180	$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$	744
3.181	$\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{\frac{5}{2}}} dx$	747
3.182	$\int \frac{1}{\sqrt{\sec(c+dx)} (b \sec(c+dx))^{\frac{5}{2}}} dx$	751
3.183	$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{\frac{5}{2}}} dx$	755
3.184	$\int \sec^2(c+dx) \sqrt[3]{b \sec(c+dx)} dx$	759
3.185	$\int \sec(c+dx) \sqrt[3]{b \sec(c+dx)} dx$	762
3.186	$\int \sqrt[3]{b \sec(c+dx)} dx$	765
3.187	$\int \cos(c+dx) \sqrt[3]{b \sec(c+dx)} dx$	768
3.188	$\int \cos^2(c+dx) \sqrt[3]{b \sec(c+dx)} dx$	771
3.189	$\int \sec^2(c+dx)(b \sec(c+dx))^{\frac{4}{3}} dx$	774
3.190	$\int \sec(c+dx)(b \sec(c+dx))^{\frac{4}{3}} dx$	777
3.191	$\int (b \sec(c+dx))^{\frac{4}{3}} dx$	780
3.192	$\int \cos(c+dx)(b \sec(c+dx))^{\frac{4}{3}} dx$	783
3.193	$\int \cos^2(c+dx)(b \sec(c+dx))^{\frac{4}{3}} dx$	786
3.194	$\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	789
3.195	$\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	792

3.196	$\int \frac{1}{\sqrt[3]{b \sec(c+dx)}} dx$	795
3.197	$\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	798
3.198	$\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	801
3.199	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	804
3.200	$\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	807
3.201	$\int \frac{1}{(b \sec(c+dx))^{4/3}} dx$	810
3.202	$\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	813
3.203	$\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	816
3.204	$\int \sec^m(c+dx)(b \sec(c+dx))^{4/3} dx$	819
3.205	$\int \sec^m(c+dx)(b \sec(c+dx))^{2/3} dx$	822
3.206	$\int \sec^m(c+dx)\sqrt[3]{b \sec(c+dx)} dx$	825
3.207	$\int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$	828
3.208	$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx$	831
3.209	$\int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx$	835
3.210	$\int \sec^m(c+dx)(b \sec(c+dx))^n dx$	839
3.211	$\int \sec^2(c+dx)(b \sec(c+dx))^n dx$	842
3.212	$\int \sec(c+dx)(b \sec(c+dx))^n dx$	845
3.213	$\int (b \sec(c+dx))^n dx$	848
3.214	$\int \cos(c+dx)(b \sec(c+dx))^n dx$	851
3.215	$\int \cos^2(c+dx)(b \sec(c+dx))^n dx$	854
3.216	$\int \cos^3(c+dx)(b \sec(c+dx))^n dx$	857
3.217	$\int \sec^{5/2}(c+dx)(b \sec(c+dx))^n dx$	860
3.218	$\int \sec^{3/2}(c+dx)(b \sec(c+dx))^n dx$	863
3.219	$\int \sqrt{\sec(c+dx)}(b \sec(c+dx))^n dx$	866
3.220	$\int \frac{(b \sec(c+dx))^n}{\sqrt{\sec(c+dx)}} dx$	869
3.221	$\int \frac{(b \sec(c+dx))^n}{\sqrt[3]{\sec(c+dx)}} dx$	872
3.222	$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^n \sqrt[5]{\sec(c+dx)}} dx$	876
3.223	$\int (d \sec(a+bx))^{7/2} \sin(a+bx) dx$	880
3.224	$\int (d \sec(a+bx))^{5/2} \sin(a+bx) dx$	883
3.225	$\int (d \sec(a+bx))^{3/2} \sin(a+bx) dx$	886
3.226	$\int \sqrt{d \sec(a+bx)} \sin(a+bx) dx$	889

3.227	$\int \frac{\sin(a+bx)}{\sqrt{d \sec(a+bx)}} dx$	892
3.228	$\int (d \sec(a+bx))^{5/2} \sin^3(a+bx) dx$	895
3.229	$\int (d \sec(a+bx))^{9/2} \sin^3(a+bx) dx$	899
3.230	$\int (d \csc(a+bx))^{9/2} \sqrt{c \sec(a+bx)} dx$	902
3.231	$\int (d \csc(a+bx))^{7/2} \sqrt{c \sec(a+bx)} dx$	906
3.232	$\int (d \csc(a+bx))^{5/2} \sqrt{c \sec(a+bx)} dx$	909
3.233	$\int (d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)} dx$	913
3.234	$\int \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} dx$	916
3.235	$\int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx$	919
3.236	$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{3/2}} dx$	924
3.237	$\int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx$	928
3.238	$\int (d \csc(a+bx))^{9/2} (c \sec(a+bx))^{3/2} dx$	933
3.239	$\int (d \csc(a+bx))^{7/2} (c \sec(a+bx))^{3/2} dx$	936
3.240	$\int (d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2} dx$	940
3.241	$\int (d \csc(a+bx))^{3/2} (c \sec(a+bx))^{3/2} dx$	943
3.242	$\int \sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2} dx$	947
3.243	$\int \frac{(c \sec(a+bx))^{3/2}}{\sqrt{d \csc(a+bx)}} dx$	950
3.244	$\int \frac{(c \sec(a+bx))^{3/2}}{(d \csc(a+bx))^{3/2}} dx$	954
3.245	$\int \frac{(c \sec(a+bx))^{3/2}}{(d \csc(a+bx))^{5/2}} dx$	960
3.246	$\int (d \csc(a+bx))^{9/2} (c \sec(a+bx))^{5/2} dx$	964
3.247	$\int (d \csc(a+bx))^{7/2} (c \sec(a+bx))^{5/2} dx$	968
3.248	$\int (d \csc(a+bx))^{5/2} (c \sec(a+bx))^{5/2} dx$	972
3.249	$\int (d \csc(a+bx))^{3/2} (c \sec(a+bx))^{5/2} dx$	976
3.250	$\int \sqrt{d \csc(a+bx)} (c \sec(a+bx))^{5/2} dx$	979
3.251	$\int \frac{(c \sec(a+bx))^{5/2}}{\sqrt{d \csc(a+bx)}} dx$	983
3.252	$\int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{3/2}} dx$	986
3.253	$\int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{5/2}} dx$	990
3.254	$\int \frac{(d \csc(a+bx))^{9/2}}{\sqrt{c \sec(a+bx)}} dx$	996
3.255	$\int \frac{(d \csc(a+bx))^{7/2}}{\sqrt{c \sec(a+bx)}} dx$	999
3.256	$\int \frac{(d \csc(a+bx))^{5/2}}{\sqrt{c \sec(a+bx)}} dx$	1003
3.257	$\int \frac{(d \csc(a+bx))^{3/2}}{\sqrt{c \sec(a+bx)}} dx$	1006

3.258	$\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx$	1010
3.259	$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx$	1015
3.260	$\int \frac{1}{(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} dx$	1019
3.261	$\int \frac{1}{(d \csc(a+bx))^{5/2} \sqrt{c \sec(a+bx)}} dx$	1025
3.262	$\int \frac{(d \csc(a+bx))^{11/2}}{(c \sec(a+bx))^{3/2}} dx$	1029
3.263	$\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{3/2}} dx$	1033
3.264	$\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{3/2}} dx$	1037
3.265	$\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{3/2}} dx$	1040
3.266	$\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{3/2}} dx$	1044
3.267	$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{3/2}} dx$	1050
3.268	$\int \frac{1}{\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} dx$	1054
3.269	$\int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{3/2}} dx$	1060
3.270	$\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$	1064
3.271	$\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx$	1070
3.272	$\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{5/2}} dx$	1073
3.273	$\int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{5/2}} dx$	1077
3.274	$\int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{5/2}} dx$	1083
3.275	$\int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx$	1087
3.276	$\int \frac{1}{\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{5/2}} dx$	1092
3.277	$\int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{5/2}} dx$	1096
3.278	$\int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{5/2}} dx$	1102
3.279	$\int \frac{1}{(d \csc(a+bx))^{7/2} (c \sec(a+bx))^{5/2}} dx$	1106
3.280	$\int \csc^n(e+fx) \sec^m(e+fx) dx$	1112
3.281	$\int \csc^n(e+fx) (a \sec(e+fx))^m dx$	1115
3.282	$\int (b \csc(e+fx))^n \sec^m(e+fx) dx$	1118
3.283	$\int (b \csc(e+fx))^n (a \sec(e+fx))^m dx$	1121
3.284	$\int (b \csc(e+fx))^n \sec^5(e+fx) dx$	1124
3.285	$\int (b \csc(e+fx))^n \sec^3(e+fx) dx$	1127
3.286	$\int (b \csc(e+fx))^n \sec(e+fx) dx$	1130

3.287	$\int \cos(e + fx)(b \csc(e + fx))^n dx$1133
3.288	$\int \cos^3(e + fx)(b \csc(e + fx))^n dx$1136
3.289	$\int \cos^5(e + fx)(b \csc(e + fx))^n dx$1139
3.290	$\int (b \csc(e + fx))^n \sec^6(e + fx) dx$1142
3.291	$\int (b \csc(e + fx))^n \sec^4(e + fx) dx$1145
3.292	$\int (b \csc(e + fx))^n \sec^2(e + fx) dx$1148
3.293	$\int (b \csc(e + fx))^n dx$1151
3.294	$\int \cos^2(e + fx)(b \csc(e + fx))^n dx$1154
3.295	$\int \cos^4(e + fx)(b \csc(e + fx))^n dx$1157
3.296	$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx$1160
3.297	$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$1163
3.298	$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$1166
3.299	$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx$1170

4	Listing of Grading functions		1175
4.0.1	Mathematica and Rubi grading function1175
4.0.2	Maple grading function1177
4.0.3	Sympy grading function1182
4.0.4	SageMath grading function1185

Chapter 1

Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [299]. This is test number [115].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.3 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12.1 on windows 10.
3. Maple 2021.1 (64 bit) on windows 10.
4. Maxima 5.44 on Linux. (via sagemath 9.3)
5. Fricas 1.3.7 on Linux (via sagemath 9.3)
6. Giac/Xcas 1.7 on Linux. (via sagemath 9.3)
7. Sympy 1.8 under Python 3.8.8 using Anaconda distribution on Ubuntu.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 under windows 10 (64 bit)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	solved	Failed
Rubi	% 100.00 (299)	% 0.00 (0)
Mathematica	% 100.00 (299)	% 0.00 (0)
Maple	% 75.25 (225)	% 24.75 (74)
Maxima	% 31.10 (93)	% 68.90 (206)
Fricas	% 35.45 (106)	% 64.55 (193)
Sympy	% 7.02 (21)	% 92.98 (278)
Giac	% 11.71 (35)	% 88.29 (264)
Mupad	% 26.09 (78)	% 73.91 (221)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	86.62	1.34	12.04	0.00
Maple	35.12	10.37	29.77	24.75
Maxima	23.08	8.03	0.00	68.90
Fricas	33.11	2.34	0.00	64.55
Sympy	7.02	0.00	0.00	92.98
Giac	8.36	3.34	0.00	88.29
Mupad	0.00	26.09	0.00	73.91

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	74	100.00 %	0.00 %	0.00 %
Maxima	206	100.00 %	0.00 %	0.00 %
Fricas	193	92.75 %	7.25 %	0.00 %
Sympy	278	59.35 %	40.65 %	0.00 %
Giac	264	99.24 %	0.00 %	0.76 %
Mupad	221	100.00 %	0.00 %	0.00 %

Table 1.4: Time and leaf size performance for each CAS



1.3 Performance

The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.06	79.00	1.00	70.00	1.00
Mathematica	0.21	63.33	0.95	56.00	0.87
Maple	0.95	192.73	2.32	122.00	1.58
Maxima	0.80	168.53	2.44	42.00	0.96
Fricas	0.80	76.20	1.57	51.00	1.07
Sympy	39.54	42.38	1.02	36.00	1.05
Giac	0.84	54.74	1.93	44.00	1.08
Mupad	0.78	61.22	1.19	46.00	1.05

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used columns from the above table.



1.4 list of integrals that has no closed form antiderivative

{

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {280, 281, 282, 283, 298}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (SageMath uses Maxima), then any integral where Maxima needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore Maxima result below is lower than what could result if Maxima was run directly and each question Maxima asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the Timofeev test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be indentified by looking at the output of the integration in each section for Maxima. The exception message will indicate of the error is due to the interactive question being asked or not.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima `abs_integrate` was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/X-CAS results

There are Few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail With error `Exception raised: NotImplementedError`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and XCAS syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the builtin system function `LeafSize` is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy and Giac antiderivatives is determined using the following function, thanks to user `slelievre` at <https://>

ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which is called directly from Python, the following code is used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative, Maple was used to determine the leaf size of Mupad output by post processing.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
 2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
 3. integer. Leaf size of result.
 4. integer. Leaf size of the optimal antiderivative.
 5. number. CPU time used to solve this integral. 0 if failed.
 6. string. The integral in Latex format
 7. string. The input used in CAS own syntax.
 8. string. The result (antiderivative) produced by CAS in Latex format
 9. string. The optimal antiderivative in Latex format.
 10. integer. 0 or 1. Indicates if problem has known antiderivative or not
 11. String. The result (antiderivative) in CAS own syntax.
 12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
- The following field present only in Rubi and Mathematica Tables*
13. integer. 1 if result was verified or 0 if not verified.
- The following fields present only in Rubi Tables*
14. integer. Number of rules used.
 15. integer. Integrand leaf size.
 16. real number. Ratio of field 14 over field 15
 17. integer. 1 if result was verified or 0 if not verified.
 18. String of form "{n,n,...}" which is list of the rules used by Rubi

High level overview of the CAS independent integration test build system

Chapter 2

detailed summary tables of results

2.1 List of integrals sorted by grade for each CAS

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85,

86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 231, 233, 235, 237, 238, 240, 242, 247, 249, 251, 253, 254, 256, 262, 264, 266, 268, 270, 271, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 296, 297, 299 }

B grade: { 41, 42, 294, 295 }

C grade: { 230, 232, 234, 236, 239, 241, 243, 244, 245, 246, 248, 250, 252, 255, 257, 258, 259, 260, 261, 263, 265, 267, 269, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 298 }

F grade: { }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 11, 39, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 66, 67, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 229, 231, 233, 236, 238, 240, 242, 247, 249, 250, 251, 252, 254, 256, 262, 264, 267, 269, 271 }

B grade: { 9, 10, 12, 13, 14, 15, 16, 40, 41, 42, 228, 230, 232, 234, 239, 241, 243, 245, 246, 248, 255, 257, 259, 261, 263, 265, 272, 274, 276, 278, 287 }

C grade: { 17, 18, 19, 20, 21, 22, 23, 24, 55, 56, 57, 58, 59, 60, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 235, 237, 244, 253, 258, 260, 266, 268, 270, 273, 275, 277, 279 }

F grade: { 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 68, 69, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 280, 281, 282, 283, 284, 285, 286, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 39, 40, 41, 42, 43, 44, 45, 46, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 66, 67, 135, 137, 138, 139, 140, 141, 145, 147, 148, 149, 150, 151, 155, 157, 158, 159, 160, 164, 165, 166, 167, 171, 172, 173, 174, 175, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 228, 229, 287, 288, 289 }

B grade: { 47, 48, 49, 132, 133, 134, 136, 142, 143, 144, 146, 152, 153, 154, 156, 161, 162, 163, 168, 169, 170, 176, 177, 178 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 55, 56, 57, 58, 59, 60, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

2.1.5 FriCAS

A grade: { 2, 4, 5, 6, 7, 8, 39, 40, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 61, 62, 63, 64, 65, 66, 67, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 228, 229, 231, 233, 238, 240, 242, 247, 249, 254, 256, 262, 287, 288, 289 }

B grade: { 1, 3, 41, 42, 251, 264, 271 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 55, 56, 57, 58, 59, 60, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 230, 232, 234, 235, 236, 237, 239, 241, 243, 244, 245, 246, 248, 250, 252, 253, 255, 257, 258, 259, 260, 261, 263, 265, 266, 267, 268, 269, 270, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

2.1.6 Sympy

A grade: { 1, 43, 44, 45, 46, 51, 52, 53, 54, 137, 138, 139, 147, 164, 165, 166, 171, 172, 173, 174, 181 }

B grade: { }

C grade: { }

F grade: { 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 47, 48, 49, 50, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92,

93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 140, 141, 142, 143, 144, 145, 146, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 167, 168, 169, 170, 175, 176, 177, 178, 179, 180, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

2.1.7 Giac

A grade: { 2, 3, 4, 5, 6, 7, 8, 39, 43, 44, 45, 46, 47, 48, 49, 50, 51, 53, 54, 61, 62, 63, 64, 65, 226 }

B grade: { 1, 40, 41, 42, 52, 223, 224, 225, 227, 229 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 55, 56, 57, 58, 59, 60, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 228, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 12, 20, 43, 51, 61, 62, 63, 64, 74, 133, 135, 137, 138, 139, 140, 141, 143, 145, 147, 148, 149, 150, 151, 152, 153, 155, 157, 158, 159, 160, 162, 164, 165, 166, 167, 169, 171, 172, 173, 174, 175, 177, 179, 180, 181, 182, 183, 223, 224, 225, 226, 227, 228, 229, 231, 233, 238, 240, 242, 247, 249, 251, 254, 256, 262, 264, 271, 287, 288, 289 }

C grade: { }

F grade: { 9, 10, 11, 13, 14, 15, 16, 17, 18, 19, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 52, 53, 54, 55, 56, 57, 58, 59, 60, 65, 66, 67, 68, 69, 70, 71, 72, 73, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98,

99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 134, 136, 142, 144, 146, 154, 156, 161, 163, 168, 170, 176, 178, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 230, 232, 234, 235, 236, 237, 239, 241, 243, 244, 245, 246, 248, 250, 252, 253, 255, 257, 258, 259, 260, 261, 263, 265, 266, 267, 268, 269, 270, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	19	18	28	36	44	11
normalized size	1	1.00	1.00	1.73	1.64	2.55	3.27	4.00	1.00
time (sec)	N/A	0.004	0.002	0.028	0.405	0.752	1.925	1.911	0.397
Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	10	18	0	10	10
normalized size	1	1.00	1.00	1.10	1.00	1.80	0.00	1.00	1.00
time (sec)	N/A	0.009	0.004	0.540	0.720	0.905	0.000	0.965	0.098
Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	34	38	46	61	0	48	36
normalized size	1	1.00	1.00	1.12	1.35	1.79	0.00	1.41	1.06
time (sec)	N/A	0.014	0.010	0.560	0.475	0.807	0.000	2.935	0.111

Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	23	24	22	31	0	22	21
normalized size	1	1.00	0.88	0.92	0.85	1.19	0.00	0.85	0.81
time (sec)	N/A	0.011	0.047	0.518	0.329	0.766	0.000	1.386	0.077

Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	42	57	71	74	0	63	58
normalized size	1	1.00	0.76	1.04	1.29	1.35	0.00	1.15	1.05
time (sec)	N/A	0.025	0.076	0.538	0.351	0.870	0.000	0.561	0.124

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	35	34	34	41	0	34	31
normalized size	1	1.00	0.85	0.83	0.83	1.00	0.00	0.83	0.76
time (sec)	N/A	0.015	0.110	0.523	0.422	0.822	0.000	0.461	0.089

Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	52	76	91	84	0	73	79
normalized size	1	1.00	0.68	1.00	1.20	1.11	0.00	0.96	1.04
time (sec)	N/A	0.039	0.135	0.532	0.430	0.900	0.000	0.359	0.158

Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	43	44	44	51	0	44	39
normalized size	1	1.00	0.81	0.83	0.83	0.96	0.00	0.83	0.74
time (sec)	N/A	0.017	0.186	0.516	0.341	0.895	0.000	0.785	0.082

Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	59	358	0	0	0	0	-1
normalized size	1	1.00	0.69	4.21	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.172	5.094	0.000	0.777	0.000	0.000	0.000

Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	46	213	0	0	0	0	-1
normalized size	1	1.00	0.74	3.44	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.059	3.073	0.000	0.800	0.000	0.000	0.000

Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	45	101	0	0	0	0	-1
normalized size	1	1.00	0.78	1.74	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.037	3.227	0.000	0.595	0.000	0.000	0.000

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	36	133	0	0	0	0	33
normalized size	1	1.00	1.00	3.69	0.00	0.00	0.00	0.00	0.92
time (sec)	N/A	0.017	0.020	2.433	0.000	0.775	0.000	0.000	0.126

Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	36	133	0	0	0	0	-1
normalized size	1	1.00	1.00	3.69	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.017	0.026	2.403	0.000	0.595	0.000	0.000	0.000

Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	49	179	0	0	0	0	-1
normalized size	1	1.00	0.79	2.89	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.044	2.851	0.000	0.566	0.000	0.000	0.000

Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	55	202	0	0	0	0	-1
normalized size	1	1.00	0.89	3.26	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.062	2.987	0.000	0.634	0.000	0.000	0.000

Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	61	199	0	0	0	0	-1
normalized size	1	1.00	0.72	2.34	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.101	3.018	0.000	0.693	0.000	0.000	0.000

Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	62	354	0	0	0	0	-1
normalized size	1	1.00	0.63	3.61	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.170	1.007	0.000	0.823	0.000	0.000	0.000

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	51	128	0	0	0	0	-1
normalized size	1	1.00	0.73	1.83	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.070	0.836	0.000	0.685	0.000	0.000	0.000

Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	48	322	0	0	0	0	-1
normalized size	1	1.00	0.73	4.88	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.040	0.041	0.982	0.000	0.943	0.000	0.000	0.000

Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	98	0	0	0	0	35
normalized size	1	1.00	1.00	2.58	0.00	0.00	0.00	0.00	0.92
time (sec)	N/A	0.019	0.021	0.915	0.000	0.680	0.000	0.000	0.196

Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	306	0	0	0	0	-1
normalized size	1	1.00	1.00	8.05	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.027	0.030	1.059	0.000	0.783	0.000	0.000	0.000

Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	59	131	0	0	0	0	-1
normalized size	1	1.00	0.82	1.82	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.060	0.817	0.000	0.629	0.000	0.000	0.000

Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	60	321	0	0	0	0	-1
normalized size	1	1.00	0.83	4.46	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.074	0.973	0.000	0.886	0.000	0.000	0.000

Problem 24	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	66	153	0	0	0	0	-1
normalized size	1	1.00	0.66	1.53	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.102	0.939	0.000	0.761	0.000	0.000	0.000

Problem 25	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	55	0	0	0	0	0	-1
normalized size	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.051	0.634	0.000	0.894	0.000	0.000	0.000

Problem 26	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	55	0	0	0	0	0	-1
normalized size	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.037	0.611	0.000	0.772	0.000	0.000	0.000

Problem 27	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.033	0.039	0.675	0.000	0.719	0.000	0.000	0.000

Problem 28	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.084	0.553	0.000	0.682	0.000	0.000	0.000

Problem 29	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	55	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.084	0.599	0.000	0.730	0.000	0.000	0.000

Problem 30	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	55	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.068	0.538	0.000	0.756	0.000	0.000	0.000

Problem 31	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	57	0	0	0	0	0	-1
normalized size	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.034	0.046	0.536	0.000	0.541	0.000	0.000	0.000

Problem 32	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	57	0	0	0	0	0	-1
normalized size	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.043	0.594	0.000	0.648	0.000	0.000	0.000

Problem 33	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	-1
normalized size	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.035	0.042	0.745	0.000	0.526	0.000	0.000	0.000

Problem 34	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	-1
normalized size	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.055	0.532	0.000	0.817	0.000	0.000	0.000

Problem 35	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	57	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.055	0.538	0.000	0.898	0.000	0.000	0.000

Problem 36	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	57	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.072	0.447	0.000	0.735	0.000	0.000	0.000

Problem 37	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	61	0	0	0	0	0	-1
normalized size	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.053	1.458	0.000	0.868	0.000	0.000	0.000

Problem 38	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	61	0	0	0	0	0	-1
normalized size	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.031	0.046	1.317	0.000	0.818	0.000	0.000	0.000

Problem 39	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	74	72	42	49	0	59	-1
normalized size	1	1.00	1.48	1.44	0.84	0.98	0.00	1.18	-0.02
time (sec)	N/A	0.017	0.304	0.437	0.895	0.442	0.000	0.994	0.000

Problem 40	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	68	64	30	43	0	53	-1
normalized size	1	1.00	1.89	1.78	0.83	1.19	0.00	1.47	-0.03
time (sec)	N/A	0.012	0.158	0.353	0.589	0.713	0.000	0.430	0.000

Problem 41	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	A	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	52	55	18	34	0	44	-1
normalized size	1	1.00	2.36	2.50	0.82	1.55	0.00	2.00	-0.05
time (sec)	N/A	0.009	0.063	0.292	0.763	0.805	0.000	0.345	0.000

Problem 42	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	A	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	3	3	44	21	3	17	0	35	-1
normalized size	1	1.00	14.67	7.00	1.00	5.67	0.00	11.67	-0.33
time (sec)	N/A	0.006	0.010	0.367	0.458	0.486	0.000	0.731	0.000

Problem 43	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	14	11	4	10	6	12
normalized size	1	1.00	1.00	1.27	1.00	0.36	0.91	0.55	1.09
time (sec)	N/A	0.007	0.007	0.397	0.522	0.756	0.403	0.315	0.159

Problem 44	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	23	21	25	10	27	16	-1
normalized size	1	1.00	0.79	0.72	0.86	0.34	0.93	0.55	-0.03
time (sec)	N/A	0.011	0.015	0.341	0.482	0.634	1.046	0.497	0.000

Problem 45	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	31	29	37	18	44	25	-1
normalized size	1	1.00	0.72	0.67	0.86	0.42	1.02	0.58	-0.02
time (sec)	N/A	0.015	0.026	0.316	0.418	0.631	10.677	0.426	0.000

Problem 46	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	37	35	49	24	60	34	-1
normalized size	1	1.00	0.65	0.61	0.86	0.42	1.05	0.60	-0.02
time (sec)	N/A	0.019	0.040	0.362	0.337	0.536	147.491	0.434	0.000

Problem 47	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	78	74	2175	65	0	79	-1
normalized size	1	1.00	0.93	0.88	25.89	0.77	0.00	0.94	-0.01
time (sec)	N/A	0.041	0.140	0.448	3.536	0.477	0.000	0.331	0.000

Problem 48	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	72	66	1111	56	0	67	-1
normalized size	1	1.00	1.11	1.02	17.09	0.86	0.00	1.03	-0.02
time (sec)	N/A	0.031	0.132	0.360	1.000	0.826	0.000	0.410	0.000

Problem 49	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	55	57	324	39	0	42	-1
normalized size	1	1.00	1.20	1.24	7.04	0.85	0.00	0.91	-0.02
time (sec)	N/A	0.022	0.054	0.297	1.028	0.668	0.000	0.598	0.000

Problem 50	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	46	23	38	55	0	31	-1
normalized size	1	1.00	1.84	0.92	1.52	2.20	0.00	1.24	-0.04
time (sec)	N/A	0.015	0.009	0.385	0.888	0.637	0.000	0.334	0.000

Problem 51	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	16	6	16	15	11	15
normalized size	1	1.00	1.00	1.23	0.46	1.23	1.15	0.85	1.15
time (sec)	N/A	0.029	0.006	0.396	0.915	0.618	0.509	0.403	0.211

Problem 52	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	27	23	14	24	37	58	-1
normalized size	1	1.00	0.75	0.64	0.39	0.67	1.03	1.61	-0.03
time (sec)	N/A	0.018	0.020	0.299	1.002	0.562	1.179	0.605	0.000

Problem 53	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	36	31	22	32	60	84	-1
normalized size	1	1.00	0.65	0.56	0.40	0.58	1.09	1.53	-0.02
time (sec)	N/A	0.027	0.030	0.311	0.679	0.809	10.569	0.540	0.000

Problem 54	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	42	37	28	38	80	110	-1
normalized size	1	1.00	0.57	0.50	0.38	0.51	1.08	1.49	-0.01
time (sec)	N/A	0.035	0.030	0.339	0.563	0.732	150.570	0.700	0.000

Problem 55	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	59	223	0	0	0	0	-1
normalized size	1	1.00	0.50	1.91	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.101	0.758	0.000	0.829	0.000	0.000	0.000

Problem 56	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	43	87	0	0	0	0	-1
normalized size	1	1.00	0.66	1.34	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.035	0.039	0.549	0.000	0.702	0.000	0.000	0.000

Problem 57	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	32	191	0	0	0	0	-1
normalized size	1	1.00	0.76	4.55	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.019	0.706	0.000	0.550	0.000	0.000	0.000

Problem 58	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	31	76	0	0	0	0	-1
normalized size	1	1.00	0.70	1.73	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.043	0.577	0.000	0.567	0.000	0.000	0.000

Problem 59	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	43	198	0	0	0	0	-1
normalized size	1	1.00	0.59	2.71	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.102	0.546	0.000	0.516	0.000	0.000	0.000

Problem 60	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	59	114	0	0	0	0	-1
normalized size	1	1.00	0.50	0.97	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.103	0.523	0.000	0.580	0.000	0.000	0.000

Problem 61	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	54	53	61	76	0	67	589
normalized size	1	1.00	0.33	0.33	0.37	0.47	0.00	0.41	3.61
time (sec)	N/A	0.039	0.179	0.557	1.136	0.680	0.000	0.410	4.684

Problem 62	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	42	41	43	58	0	49	119
normalized size	1	1.00	0.36	0.35	0.37	0.50	0.00	0.42	1.02
time (sec)	N/A	0.030	0.100	0.360	0.753	0.629	0.000	0.899	2.355

Problem 63	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	30	29	25	34	0	22	36
normalized size	1	1.00	0.49	0.48	0.41	0.56	0.00	0.36	0.59
time (sec)	N/A	0.022	0.063	0.295	0.440	0.603	0.000	0.743	0.562

Problem 64	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	6	13	0	6	6
normalized size	1	1.00	1.00	0.93	0.40	0.87	0.00	0.40	0.40
time (sec)	N/A	0.016	0.006	0.395	0.700	0.582	0.000	0.699	0.111

Problem 65	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	23	22	25	27	0	39	-1
normalized size	1	1.00	0.64	0.61	0.69	0.75	0.00	1.08	-0.03
time (sec)	N/A	0.015	0.025	0.413	0.652	0.682	0.000	0.446	0.000

Problem 66	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	38	41	58	43	0	0	-1
normalized size	1	1.00	0.44	0.48	0.67	0.50	0.00	0.00	-0.01
time (sec)	N/A	0.032	0.043	0.378	0.694	0.673	0.000	0.000	0.000

Problem 67	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	55	57	88	55	0	0	-1
normalized size	1	1.00	0.42	0.43	0.67	0.42	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.091	0.597	0.694	0.675	0.000	0.000	0.000

Problem 68	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	69	0	0	0	0	0	-1
normalized size	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.053	0.100	1.832	0.000	0.863	0.000	0.000	0.000

Problem 69	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	71	0	0	0	0	0	-1
normalized size	1	1.00	0.86	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.050	0.078	1.656	0.000	0.888	0.000	0.000	0.000

Problem 70	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	69	152	0	0	0	0	-1
normalized size	1	1.00	0.71	1.57	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.217	1.201	0.000	0.668	0.000	0.000	0.000

Problem 71	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	69	356	0	0	0	0	-1
normalized size	1	1.00	0.73	3.75	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.061	0.175	0.915	0.000	0.897	0.000	0.000	0.000

Problem 72	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	51	130	0	0	0	0	-1
normalized size	1	1.00	0.74	1.88	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.042	0.087	0.786	0.000	0.621	0.000	0.000	0.000

Problem 73	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	47	314	0	0	0	0	-1
normalized size	1	1.00	0.75	4.98	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.041	0.052	0.915	0.000	0.990	0.000	0.000	0.000

Problem 74	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	98	0	0	0	0	35
normalized size	1	1.00	1.00	2.58	0.00	0.00	0.00	0.00	0.92
time (sec)	N/A	0.020	0.022	0.812	0.000	0.702	0.000	0.000	0.232

Problem 75	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	39	303	0	0	0	0	-1
normalized size	1	1.00	1.00	7.77	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.030	0.037	0.874	0.000	0.624	0.000	0.000	0.000

Problem 76	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	51	123	0	0	0	0	-1
normalized size	1	1.00	0.76	1.84	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.057	0.894	0.000	0.751	0.000	0.000	0.000

Problem 77	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	57	315	0	0	0	0	-1
normalized size	1	1.00	0.81	4.50	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.065	0.961	0.000	0.767	0.000	0.000	0.000

Problem 78	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	63	145	0	0	0	0	-1
normalized size	1	1.00	0.66	1.53	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.093	1.000	0.000	1.145	0.000	0.000	0.000

Problem 79	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	71	325	0	0	0	0	-1
normalized size	1	1.00	0.72	3.32	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.251	0.940	0.000	0.800	0.000	0.000	0.000

Problem 80	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	64	152	0	0	0	0	-1
normalized size	1	1.00	0.67	1.60	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.187	0.768	0.000	0.641	0.000	0.000	0.000

Problem 81	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	64	356	0	0	0	0	-1
normalized size	1	1.00	0.65	3.63	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.177	0.762	0.000	0.665	0.000	0.000	0.000

Problem 82	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	49	122	0	0	0	0	-1
normalized size	1	1.00	0.73	1.82	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.078	0.663	0.000	0.716	0.000	0.000	0.000

Problem 83	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	48	320	0	0	0	0	-1
normalized size	1	1.00	0.73	4.85	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.034	0.034	0.844	0.000	0.796	0.000	0.000	0.000

Problem 84	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	39	98	0	0	0	0	-1
normalized size	1	1.00	1.00	2.51	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.030	0.020	0.746	0.000	0.582	0.000	0.000	0.000

Problem 85	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	309	0	0	0	0	-1
normalized size	1	1.00	1.00	7.54	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	0.022	0.770	0.000	0.784	0.000	0.000	0.000

Problem 86	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	52	129	0	0	0	0	-1
normalized size	1	1.00	0.74	1.84	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.049	0.824	0.000	0.654	0.000	0.000	0.000

Problem 87	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	58	319	0	0	0	0	-1
normalized size	1	1.00	0.81	4.43	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.049	0.843	0.000	0.700	0.000	0.000	0.000

Problem 88	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	64	151	0	0	0	0	-1
normalized size	1	1.00	0.65	1.54	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.077	0.944	0.000	0.650	0.000	0.000	0.000

Problem 89	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	72	331	0	0	0	0	-1
normalized size	1	1.00	0.72	3.31	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.075	0.155	0.946	0.000	0.601	0.000	0.000	0.000

Problem 90	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	61	152	0	0	0	0	-1
normalized size	1	1.00	0.62	1.55	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.193	0.805	0.000	0.732	0.000	0.000	0.000

Problem 91	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	61	348	0	0	0	0	-1
normalized size	1	1.00	0.63	3.59	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.181	0.823	0.000	0.611	0.000	0.000	0.000

Problem 92	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	51	128	0	0	0	0	-1
normalized size	1	1.00	0.73	1.83	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.032	0.018	0.786	0.000	0.706	0.000	0.000	0.000

Problem 93	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	50	324	0	0	0	0	-1
normalized size	1	1.00	0.74	4.76	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.024	0.957	0.000	0.579	0.000	0.000	0.000

Problem 94	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	98	0	0	0	0	-1
normalized size	1	1.00	1.00	2.39	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	0.017	0.813	0.000	0.922	0.000	0.000	0.000

Problem 95	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	38	311	0	0	0	0	-1
normalized size	1	1.00	0.93	7.59	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	0.043	0.820	0.000	0.878	0.000	0.000	0.000

Problem 96	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	54	131	0	0	0	0	-1
normalized size	1	1.00	0.75	1.82	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.054	0.824	0.000	0.497	0.000	0.000	0.000

Problem 97	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	60	321	0	0	0	0	-1
normalized size	1	1.00	0.83	4.46	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.053	0.945	0.000	0.645	0.000	0.000	0.000

Problem 98	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	66	153	0	0	0	0	-1
normalized size	1	1.00	0.66	1.53	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.078	0.974	0.000	1.390	0.000	0.000	0.000

Problem 99	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	74	333	0	0	0	0	-1
normalized size	1	1.00	0.74	3.33	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.074	0.182	1.033	0.000	0.768	0.000	0.000	0.000

Problem 100	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	62	354	0	0	0	0	-1
normalized size	1	1.00	0.63	3.61	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	0.089	0.915	0.000	0.797	0.000	0.000	0.000

Problem 101	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	69	152	0	0	0	0	-1
normalized size	1	1.00	0.69	1.52	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.087	0.992	0.000	0.544	0.000	0.000	0.000

Problem 102	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	61	356	0	0	0	0	-1
normalized size	1	1.00	0.63	3.67	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.238	1.050	0.000	0.689	0.000	0.000	0.000

Problem 103	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	51	130	0	0	0	0	-1
normalized size	1	1.00	0.71	1.81	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.077	0.892	0.000	1.575	0.000	0.000	0.000

Problem 104	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	48	319	0	0	0	0	-1
normalized size	1	1.00	0.74	4.91	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.040	0.081	0.963	0.000	0.688	0.000	0.000	0.000

Problem 105	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	98	0	0	0	0	-1
normalized size	1	1.00	1.00	2.39	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.021	0.017	0.846	0.000	0.611	0.000	0.000	0.000

Problem 106	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	306	0	0	0	0	-1
normalized size	1	1.00	1.00	8.05	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.019	0.016	0.905	0.000	0.637	0.000	0.000	0.000

Problem 107	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	60	126	0	0	0	0	-1
normalized size	1	1.00	0.87	1.83	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.075	1.020	0.000	0.649	0.000	0.000	0.000

Problem 108	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	60	318	0	0	0	0	-1
normalized size	1	1.00	0.90	4.75	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.088	1.168	0.000	0.551	0.000	0.000	0.000

Problem 109	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	66	148	0	0	0	0	-1
normalized size	1	1.00	0.68	1.53	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.088	0.978	0.000	0.770	0.000	0.000	0.000

Problem 110	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	70	328	0	0	0	0	-1
normalized size	1	1.00	0.74	3.45	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.196	0.986	0.000	0.675	0.000	0.000	0.000

Problem 111	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	69	152	0	0	0	0	-1
normalized size	1	1.00	0.69	1.52	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.173	0.893	0.000	0.645	0.000	0.000	0.000

Problem 112	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	64	356	0	0	0	0	-1
normalized size	1	1.00	0.64	3.56	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.056	0.964	0.000	0.635	0.000	0.000	0.000

Problem 113	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	56	125	0	0	0	0	-1
normalized size	1	1.00	0.78	1.74	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.094	0.862	0.000	0.592	0.000	0.000	0.000

Problem 114	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	51	324	0	0	0	0	-1
normalized size	1	1.00	0.75	4.76	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.058	1.019	0.000	0.569	0.000	0.000	0.000

Problem 115	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	98	0	0	0	0	-1
normalized size	1	1.00	1.00	2.39	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.023	0.738	0.000	0.665	0.000	0.000	0.000

Problem 116	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	311	0	0	0	0	-1
normalized size	1	1.00	1.00	7.59	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.022	0.033	0.778	0.000	0.544	0.000	0.000	0.000

Problem 117	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	59	131	0	0	0	0	-1
normalized size	1	1.00	0.82	1.82	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.049	0.789	0.000	0.844	0.000	0.000	0.000

Problem 118	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	60	323	0	0	0	0	-1
normalized size	1	1.00	0.87	4.68	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.031	0.953	0.000	0.604	0.000	0.000	0.000

Problem 119	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	66	153	0	0	0	0	-1
normalized size	1	1.00	0.67	1.56	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.099	0.897	0.000	0.822	0.000	0.000	0.000

Problem 120	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	73	333	0	0	0	0	-1
normalized size	1	1.00	0.75	3.43	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.082	0.978	0.000	0.643	0.000	0.000	0.000

Problem 121	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	64	152	0	0	0	0	-1
normalized size	1	1.00	0.64	1.52	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.145	0.899	0.000	0.662	0.000	0.000	0.000

Problem 122	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	64	351	0	0	0	0	-1
normalized size	1	1.00	0.64	3.51	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.117	1.029	0.000	0.570	0.000	0.000	0.000

Problem 123	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	51	130	0	0	0	0	-1
normalized size	1	1.00	0.71	1.81	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.069	0.879	0.000	0.730	0.000	0.000	0.000

Problem 124	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	51	324	0	0	0	0	-1
normalized size	1	1.00	0.75	4.76	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.054	1.059	0.000	0.630	0.000	0.000	0.000

Problem 125	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	98	0	0	0	0	-1
normalized size	1	1.00	1.00	2.39	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.022	0.016	0.765	0.000	0.545	0.000	0.000	0.000

Problem 126	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	38	311	0	0	0	0	-1
normalized size	1	1.00	0.93	7.59	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.038	0.748	0.000	0.502	0.000	0.000	0.000

Problem 127	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	62	131	0	0	0	0	-1
normalized size	1	1.00	0.86	1.82	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.033	0.742	0.000	0.631	0.000	0.000	0.000

Problem 128	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	60	321	0	0	0	0	-1
normalized size	1	1.00	0.83	4.46	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.019	0.832	0.000	0.673	0.000	0.000	0.000

Problem 129	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	66	153	0	0	0	0	-1
normalized size	1	1.00	0.68	1.58	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.063	0.041	0.836	0.000	0.777	0.000	0.000	0.000

Problem 130	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	73	333	0	0	0	0	-1
normalized size	1	1.00	0.74	3.40	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.126	0.938	0.000	0.683	0.000	0.000	0.000

Problem 131	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	66	153	0	0	0	0	-1
normalized size	1	1.00	0.66	1.53	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.021	0.898	0.000	0.884	0.000	0.000	0.000

Problem 132	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	64	131	1656	229	0	0	-1
normalized size	1	1.00	0.60	1.22	15.48	2.14	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.150	1.199	1.058	0.668	0.000	0.000	0.000

Problem 133	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	52	294	43	0	0	126
normalized size	1	1.00	0.64	0.74	4.20	0.61	0.00	0.00	1.80
time (sec)	N/A	0.017	0.093	1.069	0.830	0.628	0.000	0.000	2.548

Problem 134	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	50	114	661	199	0	0	-1
normalized size	1	1.00	0.69	1.58	9.18	2.76	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.058	1.068	0.967	0.770	0.000	0.000	0.000

Problem 135	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	39	54	30	0	0	46
normalized size	1	1.00	1.00	1.22	1.69	0.94	0.00	0.00	1.44
time (sec)	N/A	0.012	0.019	1.076	1.053	0.815	0.000	0.000	0.288

Problem 136	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	52	65	111	0	0	-1
normalized size	1	1.00	1.00	1.58	1.97	3.36	0.00	0.00	-0.03
time (sec)	N/A	0.007	0.013	0.977	1.025	0.693	0.000	0.000	0.000

Problem 137	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	32	26	98	5	0	24
normalized size	1	1.00	1.00	1.33	1.08	4.08	0.21	0.00	1.00
time (sec)	N/A	0.002	0.016	1.034	0.501	0.601	0.584	0.000	0.204

Problem 138	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	41	13	30	36	0	32
normalized size	1	1.00	1.00	1.28	0.41	0.94	1.12	0.00	1.00
time (sec)	N/A	0.007	0.038	1.157	0.629	0.667	13.387	0.000	0.225

Problem 139	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	45	54	25	158	82	0	41
normalized size	1	1.00	0.71	0.86	0.40	2.51	1.30	0.00	0.65
time (sec)	N/A	0.014	0.071	1.297	0.795	0.784	110.948	0.000	0.439

Problem 140	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	52	42	48	0	0	45
normalized size	1	1.00	0.64	0.74	0.60	0.69	0.00	0.00	0.64
time (sec)	N/A	0.016	0.116	1.387	1.047	0.876	0.000	0.000	0.522

Problem 141	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	55	74	49	202	0	0	52
normalized size	1	1.00	0.56	0.76	0.50	2.06	0.00	0.00	0.53
time (sec)	N/A	0.025	0.120	1.361	0.782	0.684	0.000	0.000	0.706

Problem 142	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	64	131	1742	236	0	0	-1
normalized size	1	1.00	0.58	1.19	15.84	2.15	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.117	0.940	1.184	0.824	0.000	0.000	0.000

Problem 143	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	45	52	299	44	0	0	127
normalized size	1	1.00	0.62	0.72	4.15	0.61	0.00	0.00	1.76
time (sec)	N/A	0.017	0.093	0.877	1.002	0.819	0.000	0.000	1.361

Problem 144	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	50	114	691	202	0	0	-1
normalized size	1	1.00	0.68	1.54	9.34	2.73	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.065	0.904	0.920	0.853	0.000	0.000	0.000

Problem 145	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	32	39	54	31	0	0	47
normalized size	1	1.00	0.97	1.18	1.64	0.94	0.00	0.00	1.42
time (sec)	N/A	0.011	0.021	1.011	0.729	0.659	0.000	0.000	0.217

Problem 146	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	33	52	68	112	0	0	-1
normalized size	1	1.00	0.97	1.53	2.00	3.29	0.00	0.00	-0.03
time (sec)	N/A	0.007	0.019	1.277	0.781	1.097	0.000	0.000	0.000

Problem 147	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	24	32	26	99	5	0	25
normalized size	1	1.00	0.96	1.28	1.04	3.96	0.20	0.00	1.00
time (sec)	N/A	0.003	0.026	0.984	0.774	0.794	18.080	0.000	0.196

Problem 148	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	32	41	13	31	0	0	33
normalized size	1	1.00	0.97	1.24	0.39	0.94	0.00	0.00	1.00
time (sec)	N/A	0.007	0.047	1.105	0.687	0.651	0.000	0.000	0.245

Problem 149	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	45	54	28	161	0	0	42
normalized size	1	1.00	0.69	0.83	0.43	2.48	0.00	0.00	0.65
time (sec)	N/A	0.015	0.082	1.191	0.669	0.847	0.000	0.000	0.431

Problem 150	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	45	52	45	51	0	0	46
normalized size	1	1.00	0.62	0.72	0.62	0.71	0.00	0.00	0.64
time (sec)	N/A	0.017	0.144	1.099	0.737	0.437	0.000	0.000	0.435

Problem 151	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	55	74	53	207	0	0	53
normalized size	1	1.00	0.54	0.73	0.52	2.05	0.00	0.00	0.52
time (sec)	N/A	0.028	0.164	1.187	1.020	0.732	0.000	0.000	0.590

Problem 152	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	57	62	705	63	0	0	205
normalized size	1	1.00	0.49	0.53	6.08	0.54	0.00	0.00	1.77
time (sec)	N/A	0.024	0.225	0.942	1.177	0.552	0.000	0.000	4.700

Problem 153	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	52	311	48	0	0	129
normalized size	1	1.00	0.59	0.68	4.09	0.63	0.00	0.00	1.70
time (sec)	N/A	0.017	0.089	0.947	0.667	0.533	0.000	0.000	1.367

Problem 154	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	50	114	747	208	0	0	-1
normalized size	1	1.00	0.64	1.46	9.58	2.67	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.080	0.957	0.938	0.599	0.000	0.000	0.000

Problem 155	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	39	54	33	0	0	66
normalized size	1	1.00	0.91	1.11	1.54	0.94	0.00	0.00	1.89
time (sec)	N/A	0.012	0.031	1.090	1.196	0.721	0.000	0.000	0.725

Problem 156	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	33	52	72	114	0	0	-1
normalized size	1	1.00	0.92	1.44	2.00	3.17	0.00	0.00	-0.03
time (sec)	N/A	0.008	0.034	0.961	1.043	0.615	0.000	0.000	0.000

Problem 157	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	24	32	26	101	0	0	27
normalized size	1	1.00	0.89	1.19	0.96	3.74	0.00	0.00	1.00
time (sec)	N/A	0.003	0.020	1.015	0.763	0.845	0.000	0.000	0.116

Problem 158	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	41	13	33	0	0	35
normalized size	1	1.00	0.91	1.17	0.37	0.94	0.00	0.00	1.00
time (sec)	N/A	0.007	0.069	1.126	0.991	0.751	0.000	0.000	0.328

Problem 159	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	54	32	167	0	0	44
normalized size	1	1.00	0.65	0.78	0.46	2.42	0.00	0.00	0.64
time (sec)	N/A	0.015	0.112	0.980	0.691	0.826	0.000	0.000	0.368

Problem 160	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	52	49	55	0	0	48
normalized size	1	1.00	0.59	0.68	0.64	0.72	0.00	0.00	0.63
time (sec)	N/A	0.017	0.167	0.969	0.876	0.737	0.000	0.000	0.499

Problem 161	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	50	114	661	205	0	0	-1
normalized size	1	1.00	0.69	1.58	9.18	2.85	0.00	0.00	-0.01
time (sec)	N/A	0.019	0.061	0.867	1.075	0.715	0.000	0.000	0.000

Problem 162	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	39	59	33	0	0	51
normalized size	1	1.00	1.00	1.22	1.84	1.03	0.00	0.00	1.59
time (sec)	N/A	0.011	0.021	0.817	0.962	0.440	0.000	0.000	0.269

Problem 163	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	52	65	114	0	0	-1
normalized size	1	1.00	1.00	1.58	1.97	3.45	0.00	0.00	-0.03
time (sec)	N/A	0.007	0.014	0.832	1.068	0.637	0.000	0.000	0.000

Problem 164	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	24	32	26	101	5	0	27
normalized size	1	1.00	1.00	1.33	1.08	4.21	0.21	0.00	1.12
time (sec)	N/A	0.002	0.012	0.746	0.525	0.831	9.464	0.000	0.296

Problem 165	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	41	13	33	36	0	35
normalized size	1	1.00	1.00	1.28	0.41	1.03	1.12	0.00	1.09
time (sec)	N/A	0.007	0.031	0.937	0.940	0.790	19.620	0.000	0.313

Problem 166	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	45	54	25	165	82	0	44
normalized size	1	1.00	0.71	0.86	0.40	2.62	1.30	0.00	0.70
time (sec)	N/A	0.014	0.064	0.941	0.946	0.613	30.226	0.000	0.428

Problem 167	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	45	52	42	51	0	0	48
normalized size	1	1.00	0.64	0.74	0.60	0.73	0.00	0.00	0.69
time (sec)	N/A	0.016	0.089	1.049	0.666	0.674	0.000	0.000	0.516

Problem 168	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	50	114	670	205	0	0	-1
normalized size	1	1.00	0.64	1.46	8.59	2.63	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.062	0.821	0.771	0.950	0.000	0.000	0.000

Problem 169	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	39	67	33	0	0	51
normalized size	1	1.00	0.91	1.11	1.91	0.94	0.00	0.00	1.46
time (sec)	N/A	0.012	0.037	0.817	0.847	0.677	0.000	0.000	0.266

Problem 170	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	33	52	65	114	0	0	-1
normalized size	1	1.00	0.92	1.44	1.81	3.17	0.00	0.00	-0.03
time (sec)	N/A	0.007	0.027	0.749	0.867	0.706	0.000	0.000	0.000

Problem 171	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	24	32	26	101	5	0	27
normalized size	1	1.00	0.89	1.19	0.96	3.74	0.19	0.00	1.00
time (sec)	N/A	0.003	0.026	0.759	0.713	0.997	26.545	0.000	0.252

Problem 172	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	41	13	33	36	0	39
normalized size	1	1.00	0.91	1.17	0.37	0.94	1.03	0.00	1.11
time (sec)	N/A	0.008	0.043	0.861	1.036	0.729	15.078	0.000	0.422

Problem 173	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	54	25	165	82	0	44
normalized size	1	1.00	0.65	0.78	0.36	2.39	1.19	0.00	0.64
time (sec)	N/A	0.014	0.069	0.982	0.884	0.635	29.988	0.000	0.369

Problem 174	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	45	52	42	51	65	0	48
normalized size	1	1.00	0.59	0.68	0.55	0.67	0.86	0.00	0.63
time (sec)	N/A	0.017	0.098	1.024	0.723	0.709	121.592	0.000	0.327

Problem 175	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	55	74	49	208	0	0	55
normalized size	1	1.00	0.51	0.69	0.46	1.94	0.00	0.00	0.51
time (sec)	N/A	0.027	0.121	1.033	0.972	0.724	0.000	0.000	0.615

Problem 176	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	53	114	688	205	0	0	-1
normalized size	1	1.00	0.68	1.46	8.82	2.63	0.00	0.00	-0.01
time (sec)	N/A	0.020	0.048	0.816	0.837	0.936	0.000	0.000	0.000

Problem 177	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	32	39	67	33	0	0	51
normalized size	1	1.00	0.91	1.11	1.91	0.94	0.00	0.00	1.46
time (sec)	N/A	0.012	0.033	0.832	0.721	0.546	0.000	0.000	0.263

Problem 178	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	33	52	65	114	0	0	-1
normalized size	1	1.00	0.92	1.44	1.81	3.17	0.00	0.00	-0.03
time (sec)	N/A	0.008	0.022	0.783	0.904	0.972	0.000	0.000	0.000

Problem 179	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	24	32	26	101	0	0	27
normalized size	1	1.00	0.89	1.19	0.96	3.74	0.00	0.00	1.00
time (sec)	N/A	0.003	0.020	0.809	0.823	0.948	0.000	0.000	0.323

Problem 180	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	35	41	13	33	0	0	39
normalized size	1	1.00	1.00	1.17	0.37	0.94	0.00	0.00	1.11
time (sec)	N/A	0.007	0.026	0.842	0.904	0.764	0.000	0.000	0.392

Problem 181	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	48	54	25	165	82	0	64
normalized size	1	1.00	0.70	0.78	0.36	2.39	1.19	0.00	0.93
time (sec)	N/A	0.015	0.052	0.910	0.965	0.934	110.564	0.000	0.628

Problem 182	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	48	52	42	51	0	0	48
normalized size	1	1.00	0.63	0.68	0.55	0.67	0.00	0.00	0.63
time (sec)	N/A	0.017	0.067	1.131	1.037	0.706	0.000	0.000	0.422

Problem 183	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	58	74	49	208	0	0	55
normalized size	1	1.00	0.54	0.69	0.46	1.94	0.00	0.00	0.51
time (sec)	N/A	0.028	0.058	1.120	1.139	0.678	0.000	0.000	0.459

Problem 184	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.033	0.072	0.519	0.000	0.656	0.000	0.000	0.000

Problem 185	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	60	0	0	0	0	0	-1
normalized size	1	1.00	1.13	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.048	0.433	0.000	0.695	0.000	0.000	0.000

Problem 186	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	-1
normalized size	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.042	0.542	0.000	0.595	0.000	0.000	0.000

Problem 187	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.065	0.585	0.000	0.542	0.000	0.000	0.000

Problem 188	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	59	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.043	0.152	1.059	0.000	0.829	0.000	0.000	0.000

Problem 189	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.077	0.435	0.000	0.829	0.000	0.000	0.000

Problem 190	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	60	0	0	0	0	0	-1
normalized size	1	1.00	1.09	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.052	0.432	0.000	0.619	0.000	0.000	0.000

Problem 191	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	57	0	0	0	0	0	-1
normalized size	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.006	0.441	0.000	0.639	0.000	0.000	0.000

Problem 192	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	56	0	0	0	0	0	-1
normalized size	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	0.006	0.637	0.000	0.686	0.000	0.000	0.000

Problem 193	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.046	0.012	1.147	0.000	0.645	0.000	0.000	0.000

Problem 194	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.057	0.373	0.000	0.721	0.000	0.000	0.000

Problem 195	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	60	0	0	0	0	0	-1
normalized size	1	1.00	1.13	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.046	0.444	0.000	0.572	0.000	0.000	0.000

Problem 196	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	55	0	0	0	0	0	-1
normalized size	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.025	0.053	0.449	0.000	0.616	0.000	0.000	0.000

Problem 197	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	0.084	0.451	0.000	0.695	0.000	0.000	0.000

Problem 198	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.045	0.068	0.934	0.000	0.680	0.000	0.000	0.000

Problem 199	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	60	0	0	0	0	0	-1
normalized size	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.008	0.439	0.000	0.869	0.000	0.000	0.000

Problem 200	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	58	0	0	0	0	0	-1
normalized size	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.014	0.427	0.000	1.017	0.000	0.000	0.000

Problem 201	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	57	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.004	0.380	0.000	0.725	0.000	0.000	0.000

Problem 202	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	58	0	0	0	0	0	-1
normalized size	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	0.018	0.446	0.000	0.776	0.000	0.000	0.000

Problem 203	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	60	0	0	0	0	0	-1
normalized size	1	1.00	1.03	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.047	0.087	1.027	0.000	0.865	0.000	0.000	0.000

Problem 204	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	83	0	0	0	0	0	-1
normalized size	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.095	0.829	0.000	0.834	0.000	0.000	0.000

Problem 205	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.142	0.734	0.000	0.536	0.000	0.000	0.000

Problem 206	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.139	0.727	0.000	0.737	0.000	0.000	0.000

Problem 207	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.135	0.675	0.000	0.749	0.000	0.000	0.000

Problem 208	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	83	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.136	0.582	0.000	0.735	0.000	0.000	0.000

Problem 209	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	83	0	0	0	0	0	-1
normalized size	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.179	0.635	0.000	0.732	0.000	0.000	0.000

Problem 210	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	76	0	0	0	0	0	-1
normalized size	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.068	2.425	0.000	0.704	0.000	0.000	0.000

Problem 211	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	71	0	0	0	0	0	-1
normalized size	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.044	0.049	1.469	0.000	0.579	0.000	0.000	0.000

Problem 212	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	65	0	0	0	0	0	-1
normalized size	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	0.038	1.692	0.000	0.748	0.000	0.000	0.000

Problem 213	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	61	0	0	0	0	0	-1
normalized size	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.045	1.069	0.000	0.756	0.000	0.000	0.000

Problem 214	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	68	0	0	0	0	0	-1
normalized size	1	1.00	0.91	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.049	0.097	2.382	0.000	0.865	0.000	0.000	0.000

Problem 215	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	71	0	0	0	0	0	-1
normalized size	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.077	4.023	0.000	0.904	0.000	0.000	0.000

Problem 216	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	73	0	0	0	0	0	-1
normalized size	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.107	6.418	0.000	0.926	0.000	0.000	0.000

Problem 217	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.099	0.885	0.000	0.952	0.000	0.000	0.000

Problem 218	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.091	1.007	0.000	0.660	0.000	0.000	0.000

Problem 219	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.099	0.887	0.000	0.888	0.000	0.000	0.000

Problem 220	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.113	0.880	0.000	1.028	0.000	0.000	0.000

Problem 221	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.137	0.969	0.000	0.754	0.000	0.000	0.000

Problem 222	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	81	0	0	0	0	0	-1
normalized size	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.173	0.987	0.000	1.000	0.000	0.000	0.000

Problem 223	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	23	28	0	154	77
normalized size	1	1.00	1.00	0.85	1.15	1.40	0.00	7.70	3.85
time (sec)	N/A	0.035	0.058	0.137	0.722	0.876	0.000	1.645	1.591

Problem 224	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	23	28	0	111	39
normalized size	1	1.00	1.00	0.85	1.15	1.40	0.00	5.55	1.95
time (sec)	N/A	0.036	0.042	0.122	0.390	0.762	0.000	1.592	0.252

Problem 225	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	23	18	0	62	18
normalized size	1	1.00	1.00	0.94	1.28	1.00	0.00	3.44	1.00
time (sec)	N/A	0.036	0.033	0.122	0.614	0.965	0.000	1.492	0.096

Problem 226	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	23	23	0	22	23
normalized size	1	1.00	1.00	0.94	1.28	1.28	0.00	1.22	1.28
time (sec)	N/A	0.033	0.037	0.187	0.470	0.645	0.000	0.317	0.235

Problem 227	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	23	28	0	35	28
normalized size	1	1.00	1.00	0.85	1.15	1.40	0.00	1.75	1.40
time (sec)	N/A	0.033	0.052	0.150	0.377	0.733	0.000	1.758	0.243

Problem 228	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	32	357	36	42	0	0	50
normalized size	1	1.00	0.78	8.71	0.88	1.02	0.00	0.00	1.22
time (sec)	N/A	0.049	0.223	0.909	0.640	0.796	0.000	0.000	0.588

Problem 229	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	36	38	44	0	257	95
normalized size	1	1.00	0.98	0.84	0.88	1.02	0.00	5.98	2.21
time (sec)	N/A	0.049	0.108	0.676	0.344	0.910	0.000	1.831	4.337

Problem 230	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	122	534	0	0	0	0	-1
normalized size	1	1.00	0.95	4.17	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.204	1.506	1.278	0.000	0.919	0.000	0.000	0.000

Problem 231	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	56	54	0	67	0	0	85
normalized size	1	1.00	0.81	0.78	0.00	0.97	0.00	0.00	1.23
time (sec)	N/A	0.097	0.120	1.194	0.000	0.889	0.000	0.000	1.338

Problem 232	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	109	281	0	0	0	0	-1
normalized size	1	1.00	1.17	3.02	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.143	0.886	1.229	0.000	0.835	0.000	0.000	0.000

Problem 233	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	42	0	36	0	0	36
normalized size	1	1.00	1.00	1.35	0.00	1.16	0.00	0.00	1.16
time (sec)	N/A	0.049	0.066	1.222	0.000	0.783	0.000	0.000	0.366

Problem 234	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	68	155	0	0	0	0	-1
normalized size	1	1.00	1.28	2.92	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.093	0.675	1.415	0.000	0.804	0.000	0.000	0.000

Problem 235	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	270	270	171	275	0	0	0	0	-1
normalized size	1	1.00	0.63	1.02	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.140	1.368	1.004	0.000	0.000	0.000	0.000	0.000

Problem 236	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	80	186	0	0	0	0	-1
normalized size	1	1.00	0.86	2.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.147	0.692	1.022	0.000	1.421	0.000	0.000	0.000

Problem 237	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	322	322	222	514	0	0	0	0	-1
normalized size	1	1.00	0.69	1.60	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.212	1.822	1.104	0.000	0.000	0.000	0.000	0.000

Problem 238	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	104	57	64	0	85	0	0	110
normalized size	1	1.00	0.55	0.62	0.00	0.82	0.00	0.00	1.06
time (sec)	N/A	0.162	0.294	1.079	0.000	1.508	0.000	0.000	2.175

Problem 239	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	166	166	114	980	0	0	0	0	-1
normalized size	1	1.00	0.69	5.90	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.268	1.141	1.159	0.000	0.885	0.000	0.000	0.000

Problem 240	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	54	0	58	0	0	61
normalized size	1	1.00	0.65	0.78	0.00	0.84	0.00	0.00	0.88
time (sec)	N/A	0.103	0.142	1.036	0.000	0.961	0.000	0.000	0.759

Problem 241	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	125	99	498	0	0	0	0	-1
normalized size	1	1.00	0.79	3.98	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.205	0.519	1.146	0.000	1.019	0.000	0.000	0.000

Problem 242	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	42	0	36	0	0	36
normalized size	1	1.00	1.00	1.35	0.00	1.16	0.00	0.00	1.16
time (sec)	N/A	0.048	0.065	1.148	0.000	0.574	0.000	0.000	0.304

Problem 243	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	66	497	0	0	0	0	-1
normalized size	1	1.00	0.74	5.58	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.143	0.376	1.091	0.000	0.602	0.000	0.000	0.000

Problem 244	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	327	327	64	646	0	0	0	0	-1
normalized size	1	1.00	0.20	1.98	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.219	0.248	1.062	0.000	0.000	0.000	0.000	0.000

Problem 245	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	69	512	0	0	0	0	-1
normalized size	1	1.00	0.73	5.45	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.153	0.415	1.013	0.000	0.815	0.000	0.000	0.000

Problem 246	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	166	166	92	555	0	0	0	0	-1
normalized size	1	1.00	0.55	3.34	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.265	1.455	1.194	0.000	0.636	0.000	0.000	0.000

Problem 247	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	57	64	0	89	0	0	112
normalized size	1	1.00	0.54	0.60	0.00	0.84	0.00	0.00	1.06
time (sec)	N/A	0.162	0.193	1.124	0.000	0.966	0.000	0.000	2.319

Problem 248	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	87	300	0	0	0	0	-1
normalized size	1	1.00	0.66	2.29	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.204	0.693	1.164	0.000	0.670	0.000	0.000	0.000

Problem 249	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	54	0	58	0	0	64
normalized size	1	1.00	0.65	0.78	0.00	0.84	0.00	0.00	0.93
time (sec)	N/A	0.102	0.214	1.042	0.000	0.842	0.000	0.000	0.835

Problem 250	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	68	188	0	0	0	0	-1
normalized size	1	1.00	0.73	2.02	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.141	0.602	1.251	0.000	0.511	0.000	0.000	0.000

Problem 251	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	42	0	58	0	0	66
normalized size	1	1.00	1.00	1.27	0.00	1.76	0.00	0.00	2.00
time (sec)	N/A	0.049	0.115	0.986	0.000	0.784	0.000	0.000	0.859

Problem 252	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	70	190	0	0	0	0	-1
normalized size	1	1.00	0.71	1.94	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.150	0.532	1.072	0.000	0.692	0.000	0.000	0.000

Problem 253	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	329	329	280	536	0	0	0	0	-1
normalized size	1	1.00	0.85	1.63	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.217	1.800	1.192	0.000	0.000	0.000	0.000	0.000

Problem 254	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	45	54	0	79	0	0	99
normalized size	1	1.00	0.65	0.78	0.00	1.14	0.00	0.00	1.43
time (sec)	N/A	0.100	0.266	1.207	0.000	0.594	0.000	0.000	1.829

Problem 255	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	104	976	0	0	0	0	-1
normalized size	1	1.00	0.81	7.62	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.193	1.085	1.228	0.000	0.656	0.000	0.000	0.000

Problem 256	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	33	42	0	51	0	0	49
normalized size	1	1.00	1.00	1.27	0.00	1.55	0.00	0.00	1.48
time (sec)	N/A	0.048	0.115	1.155	0.000	0.766	0.000	0.000	0.799

Problem 257	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	80	494	0	0	0	0	-1
normalized size	1	1.00	0.90	5.55	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.139	0.483	1.199	0.000	0.676	0.000	0.000	0.000

Problem 258	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	270	270	55	316	0	0	0	0	-1
normalized size	1	1.00	0.20	1.17	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.131	0.129	1.242	0.000	0.000	0.000	0.000	0.000

Problem 259	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	66	509	0	0	0	0	-1
normalized size	1	1.00	1.25	9.60	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.088	0.223	1.013	0.000	0.864	0.000	0.000	0.000

Problem 260	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	322	322	66	658	0	0	0	0	-1
normalized size	1	1.00	0.20	2.04	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.204	0.265	0.909	0.000	0.000	0.000	0.000	0.000

Problem 261	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	84	523	0	0	0	0	-1
normalized size	1	1.00	0.88	5.51	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.141	0.363	1.008	0.000	1.346	0.000	0.000	0.000

Problem 262	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	57	54	0	88	0	0	125
normalized size	1	1.00	0.52	0.49	0.00	0.80	0.00	0.00	1.14
time (sec)	N/A	0.154	0.288	1.097	0.000	2.421	0.000	0.000	3.695

Problem 263	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	119	542	0	0	0	0	-1
normalized size	1	1.00	0.88	4.01	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.200	1.459	1.226	0.000	1.160	0.000	0.000	0.000

Problem 264	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	45	42	0	59	0	0	70
normalized size	1	1.00	1.36	1.27	0.00	1.79	0.00	0.00	2.12
time (sec)	N/A	0.053	0.136	1.097	0.000	0.858	0.000	0.000	1.294

Problem 265	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	105	286	0	0	0	0	-1
normalized size	1	1.00	1.07	2.92	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.148	0.827	1.193	0.000	0.723	0.000	0.000	0.000

Problem 266	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	327	327	199	959	0	0	0	0	-1
normalized size	1	1.00	0.61	2.93	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.215	0.672	1.139	0.000	0.000	0.000	0.000	0.000

Problem 267	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	84	193	0	0	0	0	-1
normalized size	1	1.00	0.91	2.10	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.143	0.658	1.225	0.000	1.215	0.000	0.000	0.000

Problem 268	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	322	322	223	520	0	0	0	0	-1
normalized size	1	1.00	0.69	1.61	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.213	2.019	0.988	0.000	0.000	0.000	0.000	0.000

Problem 269	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	89	220	0	0	0	0	-1
normalized size	1	1.00	0.66	1.63	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.207	0.573	0.882	0.000	0.978	0.000	0.000	0.000

Problem 270	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	371	371	246	548	0	0	0	0	-1
normalized size	1	1.00	0.66	1.48	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.297	2.283	0.848	0.000	0.000	0.000	0.000	0.000

Problem 271	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	45	42	0	67	0	0	93
normalized size	1	1.00	1.36	1.27	0.00	2.03	0.00	0.00	2.82
time (sec)	N/A	0.054	0.162	1.112	0.000	4.016	0.000	0.000	1.762

Problem 272	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	101	977	0	0	0	0	-1
normalized size	1	1.00	0.75	7.24	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.202	1.772	1.206	0.000	1.032	0.000	0.000	0.000

Problem 273	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	329	329	55	1239	0	0	0	0	-1
normalized size	1	1.00	0.17	3.77	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.214	0.204	1.226	0.000	0.000	0.000	0.000	0.000

Problem 274	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	80	507	0	0	0	0	-1
normalized size	1	1.00	0.85	5.39	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.147	0.695	1.171	0.000	0.000	0.000	0.000	0.000

Problem 275	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	322	322	70	658	0	0	0	0	-1
normalized size	1	1.00	0.22	2.04	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.203	0.226	1.197	0.000	0.000	0.000	0.000	0.000

Problem 276	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	79	522	0	0	0	0	-1
normalized size	1	1.00	0.83	5.49	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.142	0.361	0.984	0.000	2.397	0.000	0.000	0.000

Problem 277	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	371	371	81	686	0	0	0	0	-1
normalized size	1	1.00	0.22	1.85	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.291	0.373	0.864	0.000	0.000	0.000	0.000	0.000

Problem 278	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	90	536	0	0	0	0	-1
normalized size	1	1.00	0.67	3.97	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.206	0.652	0.870	0.000	1.011	0.000	0.000	0.000

Problem 279	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	406	406	95	712	0	0	0	0	-1
normalized size	1	1.00	0.23	1.75	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.390	0.403	0.894	0.000	0.000	0.000	0.000	0.000

Problem 280	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	278	0	0	0	0	0	-1
normalized size	1	1.00	3.43	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.074	2.276	2.007	0.000	5.816	0.000	0.000	0.000

Problem 281	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	280	0	0	0	0	0	-1
normalized size	1	1.00	3.26	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.086	0.590	1.840	0.000	1.697	0.000	0.000	0.000

Problem 282	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	281	0	0	0	0	0	-1
normalized size	1	1.00	3.35	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.087	0.323	1.885	0.000	3.475	0.000	0.000	0.000

Problem 283	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	283	0	0	0	0	0	-1
normalized size	1	1.00	3.18	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.094	0.186	1.867	0.000	1.387	0.000	0.000	0.000

Problem 284	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	-1
normalized size	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.051	0.040	1.244	0.000	1.182	0.000	0.000	0.000

Problem 285	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	51	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.051	0.038	1.199	0.000	1.631	0.000	0.000	0.000

Problem 286	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	-1
normalized size	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.034	1.164	0.000	1.665	0.000	0.000	0.000

Problem 287	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	23	66	29	29	0	0	28
normalized size	1	1.00	0.96	2.75	1.21	1.21	0.00	0.00	1.17
time (sec)	N/A	0.034	0.023	0.297	0.711	0.860	0.000	0.000	0.325

Problem 288	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	45	0	58	49	0	0	66
normalized size	1	1.00	0.87	0.00	1.12	0.94	0.00	0.00	1.27
time (sec)	N/A	0.052	0.137	3.865	0.710	0.878	0.000	0.000	0.632

Problem 289	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	81	0	86	73	0	0	134
normalized size	1	1.00	1.04	0.00	1.10	0.94	0.00	0.00	1.72
time (sec)	N/A	0.064	0.562	4.433	0.687	1.084	0.000	0.000	1.411

Problem 290	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	77	0	0	0	0	0	-1
normalized size	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.611	1.283	0.000	1.064	0.000	0.000	0.000

Problem 291	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	77	0	0	0	0	0	-1
normalized size	1	1.00	1.07	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.079	0.529	1.221	0.000	1.111	0.000	0.000	0.000

Problem 292	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	75	0	0	0	0	0	-1
normalized size	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.078	0.468	1.124	0.000	0.865	0.000	0.000	0.000

Problem 293	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	65	0	0	0	0	0	-1
normalized size	1	1.00	0.90	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.107	1.510	0.000	1.168	0.000	0.000	0.000

Problem 294	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	165	0	0	0	0	0	-1
normalized size	1	1.00	2.29	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.478	2.680	0.000	2.136	0.000	0.000	0.000

Problem 295	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	246	0	0	0	0	0	-1
normalized size	1	1.00	3.42	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.076	0.622	3.063	0.000	1.318	0.000	0.000	0.000

Problem 296	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	92	0	0	0	0	0	-1
normalized size	1	1.00	1.14	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.115	1.888	0.886	0.000	1.616	0.000	0.000	0.000

Problem 297	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	90	0	0	0	0	0	-1
normalized size	1	1.00	1.11	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.097	1.718	0.922	0.000	0.875	0.000	0.000	0.000

Problem 298	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	326	0	0	0	0	0	-1
normalized size	1	1.00	4.02	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.096	3.183	0.793	0.000	1.052	0.000	0.000	0.000

Problem 299	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	115	0	0	0	0	0	-1
normalized size	1	1.00	1.42	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.107	1.020	0.730	0.000	1.204	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [270] had the largest ratio of [.4400]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	6	0.167
2	A	2	2	1.00	8	0.250
3	A	2	2	1.00	8	0.250
4	A	2	1	1.00	8	0.125
5	A	3	2	1.00	8	0.250
6	A	2	1	1.00	8	0.125
7	A	4	2	1.00	8	0.250
8	A	2	1	1.00	8	0.125
9	A	4	3	1.00	10	0.300
10	A	3	3	1.00	10	0.300

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
11	A	3	3	1.00	10	0.300
12	A	2	2	1.00	10	0.200
13	A	2	2	1.00	10	0.200
14	A	3	3	1.00	10	0.300
15	A	3	3	1.00	10	0.300
16	A	4	3	1.00	10	0.300
17	A	4	3	1.00	12	0.250
18	A	3	3	1.00	12	0.250
19	A	3	3	1.00	12	0.250
20	A	2	2	1.00	12	0.167
21	A	2	2	1.00	12	0.167
22	A	3	3	1.00	12	0.250
23	A	3	3	1.00	12	0.250
24	A	4	3	1.00	12	0.250
25	A	2	2	1.00	10	0.200
26	A	2	2	1.00	10	0.200
27	A	2	2	1.00	10	0.200
28	A	2	2	1.00	10	0.200
29	A	2	2	1.00	10	0.200
30	A	2	2	1.00	10	0.200
31	A	2	2	1.00	12	0.167
32	A	2	2	1.00	12	0.167
33	A	2	2	1.00	12	0.167
34	A	2	2	1.00	12	0.167
35	A	2	2	1.00	12	0.167
36	A	2	2	1.00	12	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
37	A	2	2	1.00	8	0.250
38	A	2	2	1.00	10	0.200
39	A	5	3	1.00	8	0.375
40	A	4	3	1.00	8	0.375
41	A	3	3	1.00	8	0.375
42	A	2	2	1.00	8	0.250
43	A	2	2	1.00	8	0.250
44	A	3	3	1.00	8	0.375
45	A	4	3	1.00	8	0.375
46	A	5	3	1.00	8	0.375
47	A	6	4	1.00	10	0.400
48	A	5	4	1.00	10	0.400
49	A	4	4	1.00	10	0.400
50	A	3	3	1.00	10	0.300
51	A	2	2	1.00	10	0.200
52	A	3	3	1.00	10	0.300
53	A	4	3	1.00	10	0.300
54	A	5	3	1.00	10	0.300
55	A	7	4	1.00	10	0.400
56	A	5	4	1.00	10	0.400
57	A	4	4	1.00	10	0.400
58	A	4	4	1.00	10	0.400
59	A	5	4	1.00	10	0.400
60	A	7	4	1.00	10	0.400
61	A	3	2	1.00	10	0.200
62	A	3	2	1.00	10	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
63	A	3	2	1.00	10	0.200
64	A	3	3	1.00	10	0.300
65	A	3	3	1.00	10	0.300
66	A	5	3	1.00	10	0.300
67	A	7	3	1.00	10	0.300
68	A	3	3	1.00	12	0.250
69	A	3	3	1.00	14	0.214
70	A	5	4	1.00	21	0.190
71	A	5	4	1.00	21	0.190
72	A	4	4	1.00	21	0.190
73	A	4	4	1.00	19	0.210
74	A	2	2	1.00	12	0.167
75	A	3	3	1.00	19	0.158
76	A	4	4	1.00	21	0.190
77	A	4	4	1.00	21	0.190
78	A	5	4	1.00	21	0.190
79	A	5	4	1.00	21	0.190
80	A	5	4	1.00	21	0.190
81	A	5	4	1.00	21	0.190
82	A	4	4	1.00	19	0.210
83	A	3	3	1.00	12	0.250
84	A	3	3	1.00	19	0.158
85	A	3	3	1.00	21	0.143
86	A	4	4	1.00	21	0.190
87	A	4	4	1.00	21	0.190
88	A	5	4	1.00	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
89	A	5	4	1.00	21	0.190
90	A	5	4	1.00	21	0.190
91	A	5	4	1.00	19	0.210
92	A	3	3	1.00	12	0.250
93	A	4	4	1.00	19	0.210
94	A	3	3	1.00	21	0.143
95	A	3	3	1.00	21	0.143
96	A	4	4	1.00	21	0.190
97	A	4	4	1.00	21	0.190
98	A	5	4	1.00	21	0.190
99	A	5	4	1.00	21	0.190
100	A	4	3	1.00	12	0.250
101	A	5	4	1.00	21	0.190
102	A	5	4	1.00	21	0.190
103	A	4	4	1.00	21	0.190
104	A	4	4	1.00	21	0.190
105	A	3	3	1.00	19	0.158
106	A	2	2	1.00	12	0.167
107	A	4	4	1.00	19	0.210
108	A	4	4	1.00	21	0.190
109	A	5	4	1.00	21	0.190
110	A	5	4	1.00	21	0.190
111	A	5	4	1.00	21	0.190
112	A	5	4	1.00	21	0.190
113	A	4	4	1.00	21	0.190
114	A	4	4	1.00	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
115	A	3	3	1.00	21	0.143
116	A	3	3	1.00	19	0.158
117	A	3	3	1.00	12	0.250
118	A	4	4	1.00	19	0.210
119	A	5	4	1.00	21	0.190
120	A	5	4	1.00	21	0.190
121	A	5	4	1.00	21	0.190
122	A	5	4	1.00	21	0.190
123	A	4	4	1.00	21	0.190
124	A	4	4	1.00	21	0.190
125	A	3	3	1.00	21	0.143
126	A	3	3	1.00	21	0.143
127	A	4	4	1.00	19	0.210
128	A	3	3	1.00	12	0.250
129	A	5	4	1.00	19	0.210
130	A	5	4	1.00	21	0.190
131	A	4	3	1.00	12	0.250
132	A	4	3	1.00	23	0.130
133	A	3	2	1.00	23	0.087
134	A	3	3	1.00	23	0.130
135	A	3	3	1.00	23	0.130
136	A	2	2	1.00	23	0.087
137	A	2	2	1.00	23	0.087
138	A	2	2	1.00	23	0.087
139	A	3	3	1.00	23	0.130
140	A	3	2	1.00	23	0.087

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
141	A	4	3	1.00	23	0.130
142	A	4	3	1.00	23	0.130
143	A	3	2	1.00	23	0.087
144	A	3	3	1.00	23	0.130
145	A	3	3	1.00	23	0.130
146	A	2	2	1.00	23	0.087
147	A	2	2	1.00	23	0.087
148	A	2	2	1.00	23	0.087
149	A	3	3	1.00	23	0.130
150	A	3	2	1.00	23	0.087
151	A	4	3	1.00	23	0.130
152	A	3	2	1.00	23	0.087
153	A	3	2	1.00	23	0.087
154	A	3	3	1.00	23	0.130
155	A	3	3	1.00	23	0.130
156	A	2	2	1.00	23	0.087
157	A	2	2	1.00	23	0.087
158	A	2	2	1.00	23	0.087
159	A	3	3	1.00	23	0.130
160	A	3	2	1.00	23	0.087
161	A	3	3	1.00	23	0.130
162	A	3	3	1.00	23	0.130
163	A	2	2	1.00	23	0.087
164	A	2	2	1.00	23	0.087
165	A	2	2	1.00	23	0.087
166	A	3	3	1.00	23	0.130

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
167	A	3	2	1.00	23	0.087
168	A	3	3	1.00	23	0.130
169	A	3	3	1.00	23	0.130
170	A	2	2	1.00	23	0.087
171	A	2	2	1.00	23	0.087
172	A	2	2	1.00	23	0.087
173	A	3	3	1.00	23	0.130
174	A	3	2	1.00	23	0.087
175	A	4	3	1.00	23	0.130
176	A	3	3	1.00	23	0.130
177	A	3	3	1.00	23	0.130
178	A	2	2	1.00	23	0.087
179	A	2	2	1.00	23	0.087
180	A	2	2	1.00	23	0.087
181	A	3	3	1.00	23	0.130
182	A	3	2	1.00	23	0.087
183	A	4	3	1.00	23	0.130
184	A	3	3	1.00	21	0.143
185	A	3	3	1.00	19	0.158
186	A	2	2	1.00	12	0.167
187	A	3	3	1.00	19	0.158
188	A	3	3	1.00	21	0.143
189	A	3	3	1.00	21	0.143
190	A	3	3	1.00	19	0.158
191	A	2	2	1.00	12	0.167
192	A	3	3	1.00	19	0.158

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
193	A	3	3	1.00	21	0.143
194	A	3	3	1.00	21	0.143
195	A	3	3	1.00	19	0.158
196	A	2	2	1.00	12	0.167
197	A	3	3	1.00	19	0.158
198	A	3	3	1.00	21	0.143
199	A	3	3	1.00	21	0.143
200	A	3	3	1.00	19	0.158
201	A	2	2	1.00	12	0.167
202	A	3	3	1.00	19	0.158
203	A	3	3	1.00	21	0.143
204	A	3	3	1.00	21	0.143
205	A	3	3	1.00	21	0.143
206	A	3	3	1.00	21	0.143
207	A	3	3	1.00	21	0.143
208	A	3	3	1.00	21	0.143
209	A	3	3	1.00	21	0.143
210	A	3	3	1.00	19	0.158
211	A	3	3	1.00	19	0.158
212	A	3	3	1.00	17	0.176
213	A	2	2	1.00	10	0.200
214	A	3	3	1.00	17	0.176
215	A	3	3	1.00	19	0.158
216	A	3	3	1.00	19	0.158
217	A	3	3	1.00	21	0.143
218	A	3	3	1.00	21	0.143

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
219	A	3	3	1.00	21	0.143
220	A	3	3	1.00	21	0.143
221	A	3	3	1.00	21	0.143
222	A	3	3	1.00	21	0.143
223	A	2	2	1.00	19	0.105
224	A	2	2	1.00	19	0.105
225	A	2	2	1.00	19	0.105
226	A	2	2	1.00	19	0.105
227	A	2	2	1.00	19	0.105
228	A	3	2	1.00	21	0.095
229	A	3	2	1.00	21	0.095
230	A	5	4	1.00	25	0.160
231	A	2	2	1.00	25	0.080
232	A	4	4	1.00	25	0.160
233	A	1	1	1.00	25	0.040
234	A	3	3	1.00	25	0.120
235	A	12	9	1.00	25	0.360
236	A	4	4	1.00	25	0.160
237	A	13	10	1.00	25	0.400
238	A	3	2	1.00	25	0.080
239	A	6	5	1.00	25	0.200
240	A	2	2	1.00	25	0.080
241	A	5	5	1.00	25	0.200
242	A	1	1	1.00	25	0.040
243	A	4	4	1.00	25	0.160
244	A	13	10	1.00	25	0.400

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
245	A	4	4	1.00	25	0.160
246	A	6	5	1.00	25	0.200
247	A	3	3	1.00	25	0.120
248	A	5	5	1.00	25	0.200
249	A	2	2	1.00	25	0.080
250	A	4	4	1.00	25	0.160
251	A	1	1	1.00	25	0.040
252	A	4	4	1.00	25	0.160
253	A	13	10	1.00	25	0.400
254	A	2	2	1.00	25	0.080
255	A	5	4	1.00	25	0.160
256	A	1	1	1.00	25	0.040
257	A	4	4	1.00	25	0.160
258	A	12	9	1.00	25	0.360
259	A	3	3	1.00	25	0.120
260	A	13	10	1.00	25	0.400
261	A	4	4	1.00	25	0.160
262	A	3	3	1.00	25	0.120
263	A	5	5	1.00	25	0.200
264	A	1	1	1.00	25	0.040
265	A	4	4	1.00	25	0.160
266	A	13	10	1.00	25	0.400
267	A	4	4	1.00	25	0.160
268	A	13	10	1.00	25	0.400
269	A	5	5	1.00	25	0.200
270	A	14	11	1.00	25	0.440

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
271	A	1	1	1.00	25	0.040
272	A	5	5	1.00	25	0.200
273	A	13	10	1.00	25	0.400
274	A	4	4	1.00	25	0.160
275	A	13	10	1.00	25	0.400
276	A	4	4	1.00	25	0.160
277	A	14	11	1.00	25	0.440
278	A	5	5	1.00	25	0.200
279	A	15	11	1.00	25	0.440
280	A	2	2	1.00	17	0.118
281	A	2	2	1.00	19	0.105
282	A	2	2	1.00	19	0.105
283	A	2	2	1.00	21	0.095
284	A	2	2	1.00	19	0.105
285	A	2	2	1.00	19	0.105
286	A	2	2	1.00	17	0.118
287	A	2	2	1.00	17	0.118
288	A	3	2	1.00	19	0.105
289	A	3	2	1.00	19	0.105
290	A	2	2	1.00	19	0.105
291	A	2	2	1.00	19	0.105
292	A	2	2	1.00	19	0.105
293	A	2	2	1.00	10	0.200
294	A	2	2	1.00	19	0.105
295	A	2	2	1.00	19	0.105
296	A	2	2	1.00	23	0.087

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
297	A	2	2	1.00	23	0.087
298	A	2	2	1.00	23	0.087
299	A	2	2	1.00	23	0.087

Chapter 3

Listing of integrals

3.1 $\int \sec(a + bx) dx$

Optimal. Leaf size=11

$$\frac{\tanh^{-1}(\sin(a + bx))}{b}$$

[Out] arctanh(sin(b*x+a))/b

Rubi [A] time = 0.00, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3770}

$$\frac{\tanh^{-1}(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x], x]

[Out] ArcTanh[Sin[a + b*x]]/b

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]

Rubi steps

$$\int \sec(a + bx) dx = \frac{\tanh^{-1}(\sin(a + bx))}{b}$$

Mathematica [A] time = 0.00, size = 11, normalized size = 1.00

$$\frac{\tanh^{-1}(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x], x]

[Out] ArcTanh[Sin[a + b*x]]/b

fricas [B] time = 0.75, size = 28, normalized size = 2.55

$$\frac{\log(\sin(bx + a) + 1) - \log(-\sin(bx + a) + 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a), x, algorithm="fricas")

[Out] 1/2*(log(sin(b*x + a) + 1) - log(-sin(b*x + a) + 1))/b

giac [B] time = 1.91, size = 44, normalized size = 4.00

$$\frac{\log\left(\left|\frac{1}{\sin(bx+a)} + \sin(bx + a) + 2\right|\right) - \log\left(\left|\frac{1}{\sin(bx+a)} + \sin(bx + a) - 2\right|\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a), x, algorithm="giac")

[Out] 1/4*(log(abs(1/sin(b*x + a) + sin(b*x + a) + 2)) - log(abs(1/sin(b*x + a) + sin(b*x + a) - 2)))/b

maple [A] time = 0.03, size = 19, normalized size = 1.73

$$\frac{\ln(\sec(bx + a) + \tan(bx + a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a),x)`

[Out] `1/b*ln(sec(b*x+a)+tan(b*x+a))`

maxima [A] time = 0.41, size = 18, normalized size = 1.64

$$\frac{\log(\sec(bx + a) + \tan(bx + a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a),x, algorithm="maxima")`

[Out] `log(sec(b*x + a) + tan(b*x + a))/b`

mupad [B] time = 0.40, size = 11, normalized size = 1.00

$$\frac{\operatorname{atanh}(\sin(a + bx))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(a + b*x),x)`

[Out] `atanh(sin(a + b*x))/b`

sympy [A] time = 1.93, size = 36, normalized size = 3.27

$$\left\{ \begin{array}{ll} \frac{\log(\tan(a+bx)+\sec(a+bx))}{b} & \text{for } b \neq 0 \\ \frac{x(\tan(a)\sec(a)+\sec^2(a))}{\tan(a)+\sec(a)} & \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a),x)`

[Out] `Piecewise((log(tan(a + b*x) + sec(a + b*x))/b, Ne(b, 0)), (x*(tan(a)*sec(a) + sec(a)**2)/(tan(a) + sec(a)), True))`

3.2 $\int \sec^2(a + bx) dx$

Optimal. Leaf size=10

$$\frac{\tan(a + bx)}{b}$$

[Out] $\tan(b*x+a)/b$

Rubi [A] time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3767, 8}

$$\frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^2, x]$

[Out] $\text{Tan}[a + b*x]/b$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \sec^2(a + bx) dx &= -\frac{\text{Subst}(\int 1 dx, x, -\tan(a + bx))}{b} \\ &= \frac{\tan(a + bx)}{b} \end{aligned}$$

Mathematica [A] time = 0.00, size = 10, normalized size = 1.00

$$\frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2,x]

[Out] Tan[a + b*x]/b

fricas [A] time = 0.91, size = 18, normalized size = 1.80

$$\frac{\sin(bx + a)}{b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2,x, algorithm="fricas")

[Out] sin(b*x + a)/(b*cos(b*x + a))

giac [A] time = 0.96, size = 10, normalized size = 1.00

$$\frac{\tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2,x, algorithm="giac")

[Out] tan(b*x + a)/b

maple [A] time = 0.54, size = 11, normalized size = 1.10

$$\frac{\tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2,x)

[Out] tan(b*x+a)/b

maxima [A] time = 0.72, size = 10, normalized size = 1.00

$$\frac{\tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2,x, algorithm="maxima")

[Out] tan(b*x + a)/b

mupad [B] time = 0.10, size = 10, normalized size = 1.00

$$\frac{\tan(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/cos(a + b*x)^2,x)
```

```
[Out] tan(a + b*x)/b
```

```
sympy [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int \sec^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)**2,x)
```

```
[Out] Integral(sec(a + b*x)**2, x)
```

3.3 $\int \sec^3(a + bx) dx$

Optimal. Leaf size=34

$$\frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\tan(a + bx) \sec(a + bx)}{2b}$$

[Out] 1/2*arctanh(sin(b*x+a))/b+1/2*sec(b*x+a)*tan(b*x+a)/b

Rubi [A] time = 0.01, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3770}

$$\frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\tan(a + bx) \sec(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^3, x]

[Out] ArcTanh[Sin[a + b*x]]/(2*b) + (Sec[a + b*x]*Tan[a + b*x])/(2*b)

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x] * (b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sec^3(a + bx) dx &= \frac{\sec(a + bx) \tan(a + bx)}{2b} + \frac{1}{2} \int \sec(a + bx) dx \\ &= \frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\sec(a + bx) \tan(a + bx)}{2b} \end{aligned}$$

Mathematica [A] time = 0.01, size = 34, normalized size = 1.00

$$\frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\tan(a + bx) \sec(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^3,x]

[Out] ArcTanh[Sin[a + b*x]]/(2*b) + (Sec[a + b*x]*Tan[a + b*x])/(2*b)

fricas [B] time = 0.81, size = 61, normalized size = 1.79

$$\frac{\cos (b x+a)^2 \log (\sin (b x+a)+1)-\cos (b x+a)^2 \log (-\sin (b x+a)+1)+2 \sin (b x+a)}{4 b \cos (b x+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3,x, algorithm="fricas")

[Out] 1/4*(cos(b*x + a)^2*log(sin(b*x + a) + 1) - cos(b*x + a)^2*log(-sin(b*x + a) + 1) + 2*sin(b*x + a))/(b*cos(b*x + a)^2)

giac [A] time = 2.94, size = 48, normalized size = 1.41

$$\frac{\frac{2 \sin (b x+a)}{\sin (b x+a)^2-1}-\log (|\sin (b x+a)+1|)+\log (|\sin (b x+a)-1|)}{4 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3,x, algorithm="giac")

[Out] -1/4*(2*sin(b*x + a)/(sin(b*x + a)^2 - 1) - log(abs(sin(b*x + a) + 1)) + log(abs(sin(b*x + a) - 1)))/b

maple [A] time = 0.56, size = 38, normalized size = 1.12

$$\frac{\sec (b x+a) \tan (b x+a)}{2 b}+\frac{\ln (\sec (b x+a)+\tan (b x+a))}{2 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^3,x)

[Out] 1/2*sec(b*x+a)*tan(b*x+a)/b+1/2/b*ln(sec(b*x+a)+tan(b*x+a))

maxima [A] time = 0.47, size = 46, normalized size = 1.35

$$\frac{\frac{2 \sin (b x+a)}{\sin (b x+a)^2-1}-\log (\sin (b x+a)+1)+\log (\sin (b x+a)-1)}{4 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3,x, algorithm="maxima")

[Out] $-1/4*(2*\sin(b*x + a)/(\sin(b*x + a)^2 - 1) - \log(\sin(b*x + a) + 1) + \log(\sin(b*x + a) - 1))/b$

mupad [B] time = 0.11, size = 36, normalized size = 1.06

$$\frac{\operatorname{atanh}(\sin(a + bx))}{2b} - \frac{\sin(a + bx)}{2b(\sin(a + bx)^2 - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^3,x)

[Out] $\operatorname{atanh}(\sin(a + b*x))/(2*b) - \sin(a + b*x)/(2*b*(\sin(a + b*x)^2 - 1))$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3,x)

[Out] Integral(sec(a + b*x)**3, x)

3.4 $\int \sec^4(a + bx) dx$

Optimal. Leaf size=26

$$\frac{\tan^3(a + bx)}{3b} + \frac{\tan(a + bx)}{b}$$

[Out] $\tan(b*x+a)/b+1/3*\tan(b*x+a)^3/b$

Rubi [A] time = 0.01, antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3767}

$$\frac{\tan^3(a + bx)}{3b} + \frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^4, x]$

[Out] $\text{Tan}[a + b*x]/b + \text{Tan}[a + b*x]^3/(3*b)$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] := -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}\{c, d\}, x \ \&\& \ \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \sec^4(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 + x^2) dx, x, -\tan(a + bx)\right)}{b} \\ &= \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A] time = 0.05, size = 23, normalized size = 0.88

$$\frac{\frac{1}{3} \tan^3(a + bx) + \tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Sec}[a + b*x]^4, x]$

[Out] $(\tan[a + b*x] + \tan[a + b*x]^3/3)/b$

fricas [A] time = 0.77, size = 31, normalized size = 1.19

$$\frac{(2 \cos(bx + a)^2 + 1) \sin(bx + a)}{3b \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4,x, algorithm="fricas")`

[Out] $1/3*(2*\cos(b*x + a)^2 + 1)*\sin(b*x + a)/(b*\cos(b*x + a)^3)$

giac [A] time = 1.39, size = 22, normalized size = 0.85

$$\frac{\tan(bx + a)^3 + 3 \tan(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4,x, algorithm="giac")`

[Out] $1/3*(\tan(b*x + a)^3 + 3*\tan(b*x + a))/b$

maple [A] time = 0.52, size = 24, normalized size = 0.92

$$\frac{\left(-\frac{2}{3} - \frac{(\sec^2(bx+a))}{3}\right) \tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^4,x)`

[Out] $-1/b*(-2/3-1/3*\sec(b*x+a)^2)*\tan(b*x+a)$

maxima [A] time = 0.33, size = 22, normalized size = 0.85

$$\frac{\tan(bx + a)^3 + 3 \tan(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4,x, algorithm="maxima")`

[Out] $1/3*(\tan(b*x + a)^3 + 3*\tan(b*x + a))/b$

mupad [B] time = 0.08, size = 21, normalized size = 0.81

$$\frac{\tan(a + bx) (\tan(a + bx)^2 + 3)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(a + b*x)^4,x)`

[Out] `(tan(a + b*x)*(tan(a + b*x)^2 + 3))/(3*b)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**4,x)`

[Out] `Integral(sec(a + b*x)**4, x)`

3.5 $\int \sec^5(a + bx) dx$

Optimal. Leaf size=55

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{8b} + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} + \frac{3 \tan(a + bx) \sec(a + bx)}{8b}$$

[Out] $3/8*\operatorname{arctanh}(\sin(b*x+a))/b+3/8*\sec(b*x+a)*\tan(b*x+a)/b+1/4*\sec(b*x+a)^3*\tan(b*x+a)/b$

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3770}

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{8b} + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} + \frac{3 \tan(a + bx) \sec(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^5, x]

[Out] $(3*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]])/(8*b) + (3*\operatorname{Sec}[a + b*x]*\operatorname{Tan}[a + b*x])/(8*b) + (\operatorname{Sec}[a + b*x]^3*\operatorname{Tan}[a + b*x])/(4*b)$

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sec^5(a + bx) dx &= \frac{\sec^3(a + bx) \tan(a + bx)}{4b} + \frac{3}{4} \int \sec^3(a + bx) dx \\ &= \frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec^3(a + bx) \tan(a + bx)}{4b} + \frac{3}{8} \int \sec(a + bx) dx \\ &= \frac{3 \tanh^{-1}(\sin(a + bx))}{8b} + \frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec^3(a + bx) \tan(a + bx)}{4b} \end{aligned}$$

Mathematica [A] time = 0.08, size = 42, normalized size = 0.76

$$\frac{3 \tanh^{-1}(\sin(a + bx)) + \tan(a + bx) \sec(a + bx) (2 \sec^2(a + bx) + 3)}{8b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^5,x]

[Out] (3*ArcTanh[Sin[a + b*x]] + Sec[a + b*x]*(3 + 2*Sec[a + b*x]^2)*Tan[a + b*x])/ (8*b)

fricas [A] time = 0.87, size = 74, normalized size = 1.35

$$\frac{3 \cos(bx + a)^4 \log(\sin(bx + a) + 1) - 3 \cos(bx + a)^4 \log(-\sin(bx + a) + 1) + 2(3 \cos(bx + a)^2 + 2) \sin(bx + a)}{16b \cos(bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5,x, algorithm="fricas")

[Out] 1/16*(3*cos(b*x + a)^4*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^4*log(-sin(b*x + a) + 1) + 2*(3*cos(b*x + a)^2 + 2)*sin(b*x + a))/(b*cos(b*x + a)^4)

giac [A] time = 0.56, size = 63, normalized size = 1.15

$$\frac{2(3 \sin(bx+a)^3 - 5 \sin(bx+a))}{(\sin(bx+a)^2 - 1)^2} - 3 \log(|\sin(bx + a) + 1|) + 3 \log(|\sin(bx + a) - 1|)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5,x, algorithm="giac")

[Out] -1/16*(2*(3*sin(b*x + a)^3 - 5*sin(b*x + a))/(sin(b*x + a)^2 - 1)^2 - 3*log(abs(sin(b*x + a) + 1)) + 3*log(abs(sin(b*x + a) - 1)))/b

maple [A] time = 0.54, size = 57, normalized size = 1.04

$$\frac{(\sec^3(bx + a)) \tan(bx + a)}{4b} + \frac{3 \sec(bx + a) \tan(bx + a)}{8b} + \frac{3 \ln(\sec(bx + a) + \tan(bx + a))}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^5,x)

[Out] $1/4*\sec(b*x+a)^3*\tan(b*x+a)/b+3/8*\sec(b*x+a)*\tan(b*x+a)/b+3/8/b*\ln(\sec(b*x+a)+\tan(b*x+a))$

maxima [A] time = 0.35, size = 71, normalized size = 1.29

$$\frac{2(3 \sin(bx+a)^3 - 5 \sin(bx+a))}{\sin(bx+a)^4 - 2 \sin(bx+a)^2 + 1} - 3 \log(\sin(bx+a) + 1) + 3 \log(\sin(bx+a) - 1)$$

$$16b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^5,x, algorithm="maxima")`

[Out] $-1/16*(2*(3*\sin(b*x + a)^3 - 5*\sin(b*x + a))/(\sin(b*x + a)^4 - 2*\sin(b*x + a)^2 + 1) - 3*\log(\sin(b*x + a) + 1) + 3*\log(\sin(b*x + a) - 1))/b$

mupad [B] time = 0.12, size = 58, normalized size = 1.05

$$\frac{3 \operatorname{atanh}(\sin(a + bx))}{8b} + \frac{\frac{5 \sin(a+bx)}{8} - \frac{3 \sin(a+bx)^3}{8}}{b(\sin(a + bx)^4 - 2 \sin(a + bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(a + b*x)^5,x)`

[Out] $(3*\operatorname{atanh}(\sin(a + b*x)))/(8*b) + ((5*\sin(a + b*x))/8 - (3*\sin(a + b*x)^3)/8)/(b*(\sin(a + b*x)^4 - 2*\sin(a + b*x)^2 + 1))$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^5(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**5,x)`

[Out] `Integral(sec(a + b*x)**5, x)`

3.6 $\int \sec^6(a + bx) dx$

Optimal. Leaf size=41

$$\frac{\tan^5(a + bx)}{5b} + \frac{2 \tan^3(a + bx)}{3b} + \frac{\tan(a + bx)}{b}$$

[Out] $\tan(b*x+a)/b+2/3*\tan(b*x+a)^3/b+1/5*\tan(b*x+a)^5/b$

Rubi [A] time = 0.01, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3767}

$$\frac{\tan^5(a + bx)}{5b} + \frac{2 \tan^3(a + bx)}{3b} + \frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Sec[a + b*x]^6,x]`

[Out] $\text{Tan}[a + b*x]/b + (2*\text{Tan}[a + b*x]^3)/(3*b) + \text{Tan}[a + b*x]^5/(5*b)$

Rule 3767

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rubi steps

$$\begin{aligned} \int \sec^6(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 + 2x^2 + x^4) dx, x, -\tan(a + bx)\right)}{b} \\ &= \frac{\tan(a + bx)}{b} + \frac{2 \tan^3(a + bx)}{3b} + \frac{\tan^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A] time = 0.11, size = 35, normalized size = 0.85

$$\frac{\frac{1}{5} \tan^5(a + bx) + \frac{2}{3} \tan^3(a + bx) + \tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] `Integrate[Sec[a + b*x]^6,x]`

[Out] $(\tan[a + b*x] + (2*\tan[a + b*x]^3)/3 + \tan[a + b*x]^5/5)/b$

fricas [A] time = 0.82, size = 41, normalized size = 1.00

$$\frac{(8 \cos(bx + a)^4 + 4 \cos(bx + a)^2 + 3) \sin(bx + a)}{15 b \cos(bx + a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^6,x, algorithm="fricas")`

[Out] $1/15*(8*\cos(b*x + a)^4 + 4*\cos(b*x + a)^2 + 3)*\sin(b*x + a)/(b*\cos(b*x + a)^5)$

giac [A] time = 0.46, size = 34, normalized size = 0.83

$$\frac{3 \tan(bx + a)^5 + 10 \tan(bx + a)^3 + 15 \tan(bx + a)}{15 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^6,x, algorithm="giac")`

[Out] $1/15*(3*\tan(b*x + a)^5 + 10*\tan(b*x + a)^3 + 15*\tan(b*x + a))/b$

maple [A] time = 0.52, size = 34, normalized size = 0.83

$$\frac{\left(-\frac{8}{15} - \frac{(\sec^4(bx+a))}{5} - \frac{4(\sec^2(bx+a))}{15}\right) \tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^6,x)`

[Out] $-1/b*(-8/15-1/5*\sec(b*x+a)^4-4/15*\sec(b*x+a)^2)*\tan(b*x+a)$

maxima [A] time = 0.42, size = 34, normalized size = 0.83

$$\frac{3 \tan(bx + a)^5 + 10 \tan(bx + a)^3 + 15 \tan(bx + a)}{15 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^6,x, algorithm="maxima")`

[Out] $1/15*(3*\tan(b*x + a)^5 + 10*\tan(b*x + a)^3 + 15*\tan(b*x + a))/b$

mupad [B] time = 0.09, size = 31, normalized size = 0.76

$$\frac{\frac{\tan(a+bx)^5}{5} + \frac{2\tan(a+bx)^3}{3} + \tan(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/cos(a + b*x)^6,x)`

[Out] `(tan(a + b*x) + (2*tan(a + b*x)^3)/3 + tan(a + b*x)^5/5)/b`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^6(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**6,x)`

[Out] `Integral(sec(a + b*x)**6, x)`

3.7 $\int \sec^7(a + bx) dx$

Optimal. Leaf size=76

$$\frac{5 \tanh^{-1}(\sin(a + bx))}{16b} + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} + \frac{5 \tan(a + bx) \sec^3(a + bx)}{24b} + \frac{5 \tan(a + bx) \sec(a + bx)}{16b}$$

[Out] 5/16*arctanh(sin(b*x+a))/b+5/16*sec(b*x+a)*tan(b*x+a)/b+5/24*sec(b*x+a)^3*tan(b*x+a)/b+1/6*sec(b*x+a)^5*tan(b*x+a)/b

Rubi [A] time = 0.04, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3770}

$$\frac{5 \tanh^{-1}(\sin(a + bx))}{16b} + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} + \frac{5 \tan(a + bx) \sec^3(a + bx)}{24b} + \frac{5 \tan(a + bx) \sec(a + bx)}{16b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^7, x]

[Out] (5*ArcTanh[Sin[a + b*x]])/(16*b) + (5*Sec[a + b*x]*Tan[a + b*x])/(16*b) + (5*Sec[a + b*x]^3*Tan[a + b*x])/(24*b) + (Sec[a + b*x]^5*Tan[a + b*x])/(6*b)

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sec^7(a + bx) dx &= \frac{\sec^5(a + bx) \tan(a + bx)}{6b} + \frac{5}{6} \int \sec^5(a + bx) dx \\
&= \frac{5 \sec^3(a + bx) \tan(a + bx)}{24b} + \frac{\sec^5(a + bx) \tan(a + bx)}{6b} + \frac{5}{8} \int \sec^3(a + bx) dx \\
&= \frac{5 \sec(a + bx) \tan(a + bx)}{16b} + \frac{5 \sec^3(a + bx) \tan(a + bx)}{24b} + \frac{\sec^5(a + bx) \tan(a + bx)}{6b} + \frac{5}{16} \int \sec dx \\
&= \frac{5 \tanh^{-1}(\sin(a + bx))}{16b} + \frac{5 \sec(a + bx) \tan(a + bx)}{16b} + \frac{5 \sec^3(a + bx) \tan(a + bx)}{24b} + \frac{\sec^5(a + bx) \tan(a + bx)}{6b}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 52, normalized size = 0.68

$$\frac{15 \tanh^{-1}(\sin(a + bx)) + \tan(a + bx) \sec(a + bx) (8 \sec^4(a + bx) + 10 \sec^2(a + bx) + 15)}{48b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^7, x]

[Out] (15*ArcTanh[Sin[a + b*x]] + Sec[a + b*x]*(15 + 10*Sec[a + b*x]^2 + 8*Sec[a + b*x]^4)*Tan[a + b*x])/(48*b)

fricas [A] time = 0.90, size = 84, normalized size = 1.11

$$\frac{15 \cos(bx + a)^6 \log(\sin(bx + a) + 1) - 15 \cos(bx + a)^6 \log(-\sin(bx + a) + 1) + 2(15 \cos(bx + a)^4 + 10 \cos(bx + a)^2 + 8) \sin(bx + a)}{96 b \cos(bx + a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7, x, algorithm="fricas")

[Out] 1/96*(15*cos(b*x + a)^6*log(sin(b*x + a) + 1) - 15*cos(b*x + a)^6*log(-sin(b*x + a) + 1) + 2*(15*cos(b*x + a)^4 + 10*cos(b*x + a)^2 + 8)*sin(b*x + a))/(b*cos(b*x + a)^6)

giac [A] time = 0.36, size = 73, normalized size = 0.96

$$\frac{\frac{2(15 \sin(bx+a)^5 - 40 \sin(bx+a)^3 + 33 \sin(bx+a))}{(\sin(bx+a)^2 - 1)^3} - 15 \log(|\sin(bx + a) + 1|) + 15 \log(|\sin(bx + a) - 1|)}{96 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7,x, algorithm="giac")

[Out] $-1/96*(2*(15*\sin(b*x + a)^5 - 40*\sin(b*x + a)^3 + 33*\sin(b*x + a))/(\sin(b*x + a)^2 - 1)^3 - 15*\log(\text{abs}(\sin(b*x + a) + 1)) + 15*\log(\text{abs}(\sin(b*x + a) - 1)))/b$

maple [A] time = 0.53, size = 76, normalized size = 1.00

$$\frac{(\sec^5(bx + a)) \tan(bx + a)}{6b} + \frac{5(\sec^3(bx + a)) \tan(bx + a)}{24b} + \frac{5 \sec(bx + a) \tan(bx + a)}{16b} + \frac{5 \ln(\sec(bx + a) + \tan(bx + a))}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^7,x)

[Out] $1/6*\sec(b*x+a)^5*\tan(b*x+a)/b+5/24*\sec(b*x+a)^3*\tan(b*x+a)/b+5/16*\sec(b*x+a)*\tan(b*x+a)/b+5/16/b*\ln(\sec(b*x+a)+\tan(b*x+a))$

maxima [A] time = 0.43, size = 91, normalized size = 1.20

$$\frac{2(15 \sin(bx+a)^5 - 40 \sin(bx+a)^3 + 33 \sin(bx+a))}{\sin(bx+a)^6 - 3 \sin(bx+a)^4 + 3 \sin(bx+a)^2 - 1} - 15 \log(\sin(bx + a) + 1) + 15 \log(\sin(bx + a) - 1)$$

96 b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7,x, algorithm="maxima")

[Out] $-1/96*(2*(15*\sin(b*x + a)^5 - 40*\sin(b*x + a)^3 + 33*\sin(b*x + a))/(\sin(b*x + a)^4 - 3*\sin(b*x + a)^2 + 3*\sin(b*x + a)^2 - 1) - 15*\log(\sin(b*x + a) + 1) + 15*\log(\sin(b*x + a) - 1))/b$

mupad [B] time = 0.16, size = 79, normalized size = 1.04

$$\frac{5 \operatorname{atanh}(\sin(a + bx))}{16b} - \frac{\frac{5 \sin(a+bx)^5}{16} - \frac{5 \sin(a+bx)^3}{6} + \frac{11 \sin(a+bx)}{16}}{b(\sin(a + bx)^6 - 3 \sin(a + bx)^4 + 3 \sin(a + bx)^2 - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^7,x)

[Out] $(5*\operatorname{atanh}(\sin(a + b*x)))/(16*b) - ((11*\sin(a + b*x))/16 - (5*\sin(a + b*x)^3)/6 + (5*\sin(a + b*x)^5)/16)/(b*(3*\sin(a + b*x)^2 - 3*\sin(a + b*x)^4 + \sin(a + b*x)^6 - 1))$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^7(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)**7,x)
```

```
[Out] Integral(sec(a + b*x)**7, x)
```

3.8 $\int \sec^8(a + bx) dx$

Optimal. Leaf size=53

$$\frac{\tan^7(a + bx)}{7b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{\tan^3(a + bx)}{b} + \frac{\tan(a + bx)}{b}$$

[Out] $\tan(b*x+a)/b+\tan(b*x+a)^3/b+3/5*\tan(b*x+a)^5/b+1/7*\tan(b*x+a)^7/b$

Rubi [A] time = 0.02, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {3767}

$$\frac{\tan^7(a + bx)}{7b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{\tan^3(a + bx)}{b} + \frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^8, x]$

[Out] $\text{Tan}[a + b*x]/b + \text{Tan}[a + b*x]^3/b + (3*\text{Tan}[a + b*x]^5)/(5*b) + \text{Tan}[a + b*x]^7/(7*b)$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}\{c, d\}, x] \&\& \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \sec^8(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 + 3x^2 + 3x^4 + x^6) dx, x, -\tan(a + bx)\right)}{b} \\ &= \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{\tan^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A] time = 0.19, size = 43, normalized size = 0.81

$$\frac{\frac{1}{7} \tan^7(a + bx) + \frac{3}{5} \tan^5(a + bx) + \tan^3(a + bx) + \tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^8,x]

[Out] (Tan[a + b*x] + Tan[a + b*x]^3 + (3*Tan[a + b*x]^5)/5 + Tan[a + b*x]^7/7)/b

fricas [A] time = 0.89, size = 51, normalized size = 0.96

$$\frac{(16 \cos(bx + a)^6 + 8 \cos(bx + a)^4 + 6 \cos(bx + a)^2 + 5) \sin(bx + a)}{35 b \cos(bx + a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^8,x, algorithm="fricas")

[Out] 1/35*(16*cos(b*x + a)^6 + 8*cos(b*x + a)^4 + 6*cos(b*x + a)^2 + 5)*sin(b*x + a)/(b*cos(b*x + a)^7)

giac [A] time = 0.78, size = 44, normalized size = 0.83

$$\frac{5 \tan(bx + a)^7 + 21 \tan(bx + a)^5 + 35 \tan(bx + a)^3 + 35 \tan(bx + a)}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^8,x, algorithm="giac")

[Out] 1/35*(5*tan(b*x + a)^7 + 21*tan(b*x + a)^5 + 35*tan(b*x + a)^3 + 35*tan(b*x + a))/b

maple [A] time = 0.52, size = 44, normalized size = 0.83

$$\frac{\left(-\frac{16}{35} - \frac{(\sec^6(bx+a))}{7} - \frac{6(\sec^4(bx+a))}{35} - \frac{8(\sec^2(bx+a))}{35} \right) \tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^8,x)

[Out] -1/b*(-16/35-1/7*sec(b*x+a)^6-6/35*sec(b*x+a)^4-8/35*sec(b*x+a)^2)*tan(b*x+a)

maxima [A] time = 0.34, size = 44, normalized size = 0.83

$$\frac{5 \tan(bx + a)^7 + 21 \tan(bx + a)^5 + 35 \tan(bx + a)^3 + 35 \tan(bx + a)}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^8,x, algorithm="maxima")

[Out] 1/35*(5*tan(b*x + a)^7 + 21*tan(b*x + a)^5 + 35*tan(b*x + a)^3 + 35*tan(b*x + a))/b

mupad [B] time = 0.08, size = 39, normalized size = 0.74

$$\frac{\frac{\tan(a+bx)^7}{7} + \frac{3\tan(a+bx)^5}{5} + \tan(a+bx)^3 + \tan(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/cos(a + b*x)^8,x)

[Out] (tan(a + b*x) + tan(a + b*x)^3 + (3*tan(a + b*x)^5)/5 + tan(a + b*x)^7/7)/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^8(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**8,x)

[Out] Integral(sec(a + b*x)**8, x)

3.9 $\int \sec^2(a + bx) dx$

Optimal. Leaf size=85

$$\frac{2 \sin(a + bx) \sec^{\frac{5}{2}}(a + bx)}{5b} + \frac{6 \sin(a + bx) \sqrt{\sec(a + bx)}}{5b} - \frac{6 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b}$$

[Out] 2/5*sec(b*x+a)^(5/2)*sin(b*x+a)/b+6/5*sin(b*x+a)*sec(b*x+a)^(1/2)/b-6/5*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticE(sin(1/2*b*x+1/2*a),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b

Rubi [A] time = 0.04, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3768, 3771, 2639}

$$\frac{2 \sin(a + bx) \sec^{\frac{5}{2}}(a + bx)}{5b} + \frac{6 \sin(a + bx) \sqrt{\sec(a + bx)}}{5b} - \frac{6 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(7/2), x]

[Out] (-6*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/(5*b) + (6*Sqrt[Sec[a + b*x]]*Sin[a + b*x])/(5*b) + (2*Sec[a + b*x]^(5/2)*Sin[a + b*x])/(5*b)

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] *(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^(n)*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{7}{2}}(a+bx) dx &= \frac{2 \sec^{\frac{5}{2}}(a+bx) \sin(a+bx)}{5b} + \frac{3}{5} \int \sec^{\frac{3}{2}}(a+bx) dx \\
&= \frac{6\sqrt{\sec(a+bx)} \sin(a+bx)}{5b} + \frac{2 \sec^{\frac{5}{2}}(a+bx) \sin(a+bx)}{5b} - \frac{3}{5} \int \frac{1}{\sqrt{\sec(a+bx)}} dx \\
&= \frac{6\sqrt{\sec(a+bx)} \sin(a+bx)}{5b} + \frac{2 \sec^{\frac{5}{2}}(a+bx) \sin(a+bx)}{5b} - \frac{1}{5} \left(3\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \right. \\
&= -\frac{6\sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{\sec(a+bx)}}{5b} + \frac{6\sqrt{\sec(a+bx)} \sin(a+bx)}{5b} + \frac{2 \sec^{\frac{5}{2}}(a+bx) \sin(a+bx)}{5b}
\end{aligned}$$

Mathematica [A] time = 0.17, size = 59, normalized size = 0.69

$$\frac{\sec^{\frac{5}{2}}(a+bx) \left(7 \sin(a+bx) + 3 \sin(3(a+bx)) - 12 \cos^{\frac{5}{2}}(a+bx) E\left(\frac{1}{2}(a+bx) \middle| 2\right) \right)}{10b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(7/2), x]

[Out] (Sec[a + b*x]^(5/2)*(-12*Cos[a + b*x]^(5/2)*EllipticE[(a + b*x)/2, 2] + 7*Sin[a + b*x] + 3*Sin[3*(a + b*x)])/(10*b)

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\sec(bx+a)^{\frac{7}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(7/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(7/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx+a)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(7/2),x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(7/2), x)

maple [B] time = 5.09, size = 358, normalized size = 4.21

$$2\sqrt{-\left(-2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} \left(12 \operatorname{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^(7/2),x)

[Out] $\frac{2}{5} * (-(-2 * \cos(1/2 * b * x + 1/2 * a)^2 + 1) * \sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} / (8 * \sin(1/2 * b * x + 1/2 * a)^6 - 12 * \sin(1/2 * b * x + 1/2 * a)^4 + 6 * \sin(1/2 * b * x + 1/2 * a)^2 - 1) / \sin(1/2 * b * x + 1/2 * a)^3 * (12 * \operatorname{EllipticE}(\cos(1/2 * b * x + 1/2 * a), 2^{(1/2)}) * (\sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} * \sin(1/2 * b * x + 1/2 * a)^4 - 24 * \cos(1/2 * b * x + 1/2 * a) * \sin(1/2 * b * x + 1/2 * a)^6 - 12 * \operatorname{EllipticE}(\cos(1/2 * b * x + 1/2 * a), 2^{(1/2)}) * (\sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} * \sin(1/2 * b * x + 1/2 * a)^2 + 24 * \sin(1/2 * b * x + 1/2 * a)^4 * \cos(1/2 * b * x + 1/2 * a) + 3 * (\sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} * \operatorname{EllipticE}(\cos(1/2 * b * x + 1/2 * a), 2^{(1/2)}) - 8 * \sin(1/2 * b * x + 1/2 * a)^2 * \cos(1/2 * b * x + 1/2 * a) * (-2 * \sin(1/2 * b * x + 1/2 * a)^4 + \sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} / (2 * \cos(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} / b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(7/2),x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{1}{\cos(a + b * x)}\right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(a + b*x))^(7/2),x)

[Out] int((1/cos(a + b*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**(7/2),x)

[Out] Timed out

3.10 $\int \sec^{\frac{5}{2}}(a + bx) dx$

Optimal. Leaf size=62

$$\frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} + \frac{2\sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{3b}$$

[Out] $2/3*\sec(b*x+a)^{(3/2)}*\sin(b*x+a)/b+2/3*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}*\sec(b*x+a)^{(1/2)}/b$

Rubi [A] time = 0.03, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3768, 3771, 2641}

$$\frac{2 \sin(a + bx) \sec^{\frac{3}{2}}(a + bx)}{3b} + \frac{2\sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(5/2), x]

[Out] $(2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2]*\text{Sqrt}[\text{Sec}[a + b*x]])/(3*b) + (2*\text{Sec}[a + b*x]^{(3/2)}*\text{Sin}[a + b*x])/(3*b)$

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{5}{2}}(a+bx) dx &= \frac{2 \sec^{\frac{3}{2}}(a+bx) \sin(a+bx)}{3b} + \frac{1}{3} \int \sqrt{\sec(a+bx)} dx \\
&= \frac{2 \sec^{\frac{3}{2}}(a+bx) \sin(a+bx)}{3b} + \frac{1}{3} \left(\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx \\
&= \frac{2 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{\sec(a+bx)}}{3b} + \frac{2 \sec^{\frac{3}{2}}(a+bx) \sin(a+bx)}{3b}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 46, normalized size = 0.74

$$\frac{2 \sec^{\frac{3}{2}}(a+bx) \left(\sin(a+bx) + \cos^{\frac{3}{2}}(a+bx) F\left(\frac{1}{2}(a+bx) \middle| 2\right) \right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(5/2), x]

[Out] (2*Sec[a + b*x]^(3/2)*(Cos[a + b*x]^(3/2)*EllipticF[(a + b*x)/2, 2] + Sin[a + b*x]))/(3*b)

fricas [F] time = 0.80, size = 0, normalized size = 0.00

$$\text{integral}\left(\sec(bx+a)^{\frac{5}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(5/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx+a)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(5/2), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(5/2), x)

maple [B] time = 3.07, size = 213, normalized size = 3.44

$$\frac{2 \left(-2 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \operatorname{EllipticF} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \right)}{3 \sqrt{-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + \sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^(5/2), x)`

[Out] $-2/3 * (-2 * (\sin(1/2 * b * x + 1/2 * a))^2)^{(1/2)} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} * \operatorname{EllipticF}(\cos(1/2 * b * x + 1/2 * a), 2^{(1/2)}) * \sin(1/2 * b * x + 1/2 * a)^2 + (\sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} * \operatorname{EllipticF}(\cos(1/2 * b * x + 1/2 * a), 2^{(1/2)}) - 2 * \sin(1/2 * b * x + 1/2 * a)^2 * \cos(1/2 * b * x + 1/2 * a) * ((2 * \cos(1/2 * b * x + 1/2 * a)^2 - 1) * \sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} / (-2 * \sin(1/2 * b * x + 1/2 * a)^4 + \sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} / (2 * \cos(1/2 * b * x + 1/2 * a)^2 - 1)^{(3/2)} / \sin(1/2 * b * x + 1/2 * a) / b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^(5/2), x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{1}{\cos(a + bx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(a + b*x))^(5/2), x)`

[Out] `int((1/cos(a + b*x))^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^{\frac{5}{2}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**(5/2), x)`

[Out] `Integral(sec(a + b*x)**(5/2), x)`

3.11 $\int \sec^{\frac{3}{2}}(a + bx) dx$

Optimal. Leaf size=58

$$\frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \frac{2 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

[Out] $2 \sin(bx+a) \sec(bx+a)^{(1/2)}/b - 2 (\cos(1/2 bx + 1/2 a)^{(1/2)})^2 / \cos(1/2 bx + 1/2 a) \text{EllipticE}(\sin(1/2 bx + 1/2 a), 2^{(1/2)}) \cos(bx+a)^{(1/2)} \sec(bx+a)^{(1/2)}/b$

Rubi [A] time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3768, 3771, 2639}

$$\frac{2 \sin(a + bx) \sqrt{\sec(a + bx)}}{b} - \frac{2 \sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(3/2), x]

[Out] $(-2 \sqrt{\cos[a + b*x]} \text{EllipticE}[(a + b*x)/2, 2] \sqrt{\sec[a + b*x]})/b + (2 \sqrt{\sec[a + b*x]} \sin[a + b*x])/b$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P
i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]
]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), I
nt[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] &&
IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n * Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{3}{2}}(a + bx) dx &= \frac{2\sqrt{\sec(a + bx)} \sin(a + bx)}{b} - \int \frac{1}{\sqrt{\sec(a + bx)}} dx \\
&= \frac{2\sqrt{\sec(a + bx)} \sin(a + bx)}{b} - \left(\sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} \right) \int \sqrt{\cos(a + bx)} dx \\
&= -\frac{2\sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{\sec(a + bx)}}{b} + \frac{2\sqrt{\sec(a + bx)} \sin(a + bx)}{b}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 45, normalized size = 0.78

$$\frac{2\sqrt{\sec(a + bx)} \left(\sin(a + bx) - \sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right) \right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(3/2), x]

[Out] (2*Sqrt[Sec[a + b*x]]*(-(Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]) + Sin[a + b*x]))/b

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\sec(bx + a)^{\frac{3}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(3/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(3/2), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(3/2), x)

maple [A] time = 3.23, size = 101, normalized size = 1.74

$$\frac{2 \left(\sqrt{\frac{1 - \cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \operatorname{EllipticE} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) - 2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \cos \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{\sin \left(\frac{bx}{2} + \frac{a}{2} \right) \sqrt{2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^(3/2), x)`

[Out] `-2*((sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^(3/2), x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{1}{\cos(a + bx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(a + b*x))^(3/2), x)`

[Out] `int((1/cos(a + b*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^{\frac{3}{2}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**(3/2), x)`

[Out] `Integral(sec(a + b*x)**(3/2), x)`

3.12 $\int \sqrt{\sec(a + bx)} dx$

Optimal. Leaf size=36

$$\frac{2\sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}*\sec(b*x+a)^{(1/2)}/b$

Rubi [A] time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3771, 2641}

$$\frac{2\sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[a + b*x]], x]

[Out] $(2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2]*\text{Sqrt}[\text{Sec}[a + b*x]])/b$

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{\sec(a + bx)} dx &= \left(\sqrt{\cos(a + bx)} \sqrt{\sec(a + bx)}\right) \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\ &= \frac{2\sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{\sec(a + bx)}}{b} \end{aligned}$$

Mathematica [A] time = 0.02, size = 36, normalized size = 1.00

$$\frac{2\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)}F\left(\frac{1}{2}(a+bx)\middle|2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[a + b*x]], x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/b

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{\sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\sec(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(sec(b*x + a)), x)

maple [B] time = 2.43, size = 133, normalized size = 3.69

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{-2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1}\text{EllipticF}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^(1/2), x)

[Out] -2*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(sec(b*x + a)), x)

mupad [B] time = 0.13, size = 33, normalized size = 0.92

$$\frac{2 \sqrt{\cos(a + bx)} \sqrt{\frac{1}{\cos(a+bx)}} F\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(a + b*x))^(1/2),x)

[Out] (2*cos(a + b*x)^(1/2)*(1/cos(a + b*x))^(1/2)*ellipticF(a/2 + (b*x)/2, 2))/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\sec(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**(1/2),x)

[Out] Integral(sqrt(sec(a + b*x)), x)

$$3.13 \quad \int \frac{1}{\sqrt{\sec(a+bx)}} dx$$

Optimal. Leaf size=36

$$\frac{2\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b}$$

[Out] 2*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticE(sin(1/2*b*x+1/2*a),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b

Rubi [A] time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3771, 2639}

$$\frac{2\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[Sec[a + b*x]], x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/b

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\sec(a+bx)}} dx &= \left(\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)}\right) \int \sqrt{\cos(a+bx)} dx \\ &= \frac{2\sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{\sec(a+bx)}}{b} \end{aligned}$$

Mathematica [A] time = 0.03, size = 36, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[Sec[a + b*x]], x]

[Out] (2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[Sec[a + b*x]])

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\sqrt{\sec(bx+a)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(1/2), x, algorithm="fricas")

[Out] integral(1/sqrt(sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\sec(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(1/sqrt(sec(b*x + a)), x)

maple [B] time = 2.40, size = 133, normalized size = 3.69

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{-2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1}\text{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right)}{\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(1/2), x)

[Out] 2*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),

$2^{(1/2)} / (-2 * \sin(1/2 * b * x + 1/2 * a)^4 + \sin(1/2 * b * x + 1/2 * a)^2)^{(1/2)} / \sin(1/2 * b * x + 1/2 * a) / (2 * \cos(1/2 * b * x + 1/2 * a)^2 - 1)^{(1/2)} / b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{\frac{1}{\cos(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(1/2),x)

[Out] int(1/(1/cos(a + b*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)**(1/2),x)

[Out] Integral(1/sqrt(sec(a + b*x)), x)

$$3.14 \quad \int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx$$

Optimal. Leaf size=62

$$\frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} + \frac{2\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)}F\left(\frac{1}{2}(a+bx)\middle|2\right)}{3b}$$

[Out] 2/3*sin(b*x+a)/b/sec(b*x+a)^(1/2)+2/3*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticF(sin(1/2*b*x+1/2*a),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b

Rubi [A] time = 0.03, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3769, 3771, 2641}

$$\frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} + \frac{2\sqrt{\cos(a+bx)}\sqrt{\sec(a+bx)}F\left(\frac{1}{2}(a+bx)\middle|2\right)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(-3/2), x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/(3*b) + (2*Sin[a + b*x])/(3*b*Sqrt[Sec[a + b*x]])

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx &= \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} + \frac{1}{3} \int \sqrt{\sec(a+bx)} dx \\
&= \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}} + \frac{1}{3} \left(\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx \\
&= \frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{\sec(a+bx)}}{3b} + \frac{2 \sin(a+bx)}{3b\sqrt{\sec(a+bx)}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 49, normalized size = 0.79

$$\frac{\sqrt{\sec(a+bx)} \left(\sin(2(a+bx)) + 2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(-3/2), x]

[Out] (Sqrt[Sec[a + b*x]]*(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Sin[2*(a + b*x)]))/(3*b)

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{1}{\sec(bx+a)^{\frac{3}{2}}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(3/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(-3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(3/2), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(-3/2), x)

maple [B] time = 2.85, size = 179, normalized size = 2.89

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\left(4\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)}{3\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)\sqrt{2\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(3/2), x)

[Out] -2/3*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(4*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(3/2), x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(-3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(3/2), x)

[Out] int(1/(1/cos(a + b*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec^{\frac{3}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(b*x+a)**(3/2),x)
```

```
[Out] Integral(sec(a + b*x)**(-3/2), x)
```

$$3.15 \quad \int \frac{1}{5 \sec^2(a+bx)} dx$$

Optimal. Leaf size=62

$$\frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} + \frac{6\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{5b}$$

[Out] 2/5*sin(b*x+a)/b/sec(b*x+a)^(3/2)+6/5*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticE(sin(1/2*b*x+1/2*a),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b

Rubi [A] time = 0.03, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3769, 3771, 2639}

$$\frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} + \frac{6\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(-5/2), x]

[Out] (6*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]*Sqrt[Sec[a + b*x]])/(5*b) + (2*Sin[a + b*x])/(5*b*Sec[a + b*x]^(3/2))

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P
i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Simp[(Cos[c + d*x]*(
b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c +
d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n
]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx &= \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} + \frac{3}{5} \int \frac{1}{\sqrt{\sec(a+bx)}} dx \\
&= \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)} + \frac{1}{5} \left(3\sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} \right) \int \sqrt{\cos(a+bx)} dx \\
&= \frac{6\sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{\sec(a+bx)}}{5b} + \frac{2 \sin(a+bx)}{5b \sec^{\frac{3}{2}}(a+bx)}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 55, normalized size = 0.89

$$\frac{\sqrt{\sec(a+bx)} \left(\sin(a+bx) + \sin(3(a+bx)) + 12\sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right) \right)}{10b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(-5/2), x]

[Out] (Sqrt[Sec[a + b*x]]*(12*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + Sin[a + b*x] + Sin[3*(a + b*x)]))/(10*b)

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{1}{\sec(bx+a)^{\frac{5}{2}}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(5/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(-5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(5/2),x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(-5/2), x)

maple [B] time = 2.99, size = 202, normalized size = 3.26

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\left(-8\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 8\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 5\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\sin\left(\frac{bx}{2} + \frac{a}{2}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(5/2),x)

[Out] -2/5*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(-8*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^6+8*sin(1/2*b*x+1/2*a)^4*cos(1/2*b*x+1/2*a)-3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))-2*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(5/2),x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(5/2),x)

[Out] int(1/(1/cos(a + b*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec^{\frac{5}{2}}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(b*x+a)**(5/2),x)
```

```
[Out] Integral(sec(a + b*x)**(-5/2), x)
```

$$3.16 \quad \int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx$$

Optimal. Leaf size=85

$$\frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{10 \sin(a+bx)}{21b \sqrt{\sec(a+bx)}} + \frac{10 \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{21b}$$

[Out] 2/7*sin(b*x+a)/b/sec(b*x+a)^(5/2)+10/21*sin(b*x+a)/b/sec(b*x+a)^(1/2)+10/21*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticF(sin(1/2*b*x+1/2*a),2^(1/2))*cos(b*x+a)^(1/2)*sec(b*x+a)^(1/2)/b

Rubi [A] time = 0.04, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {3769, 3771, 2641}

$$\frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{10 \sin(a+bx)}{21b \sqrt{\sec(a+bx)}} + \frac{10 \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{21b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(-7/2), x]

[Out] (10*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*sqrt[Sec[a + b*x]])/(21*b) + (2*Sin[a + b*x])/(7*b*Sec[a + b*x]^(5/2)) + (10*Sin[a + b*x])/(21*b*sqrt[Sec[a + b*x]])

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d^n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^{\frac{7}{2}}(a+bx)} dx &= \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{5}{7} \int \frac{1}{\sec^{\frac{3}{2}}(a+bx)} dx \\
&= \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{10 \sin(a+bx)}{21b \sqrt{\sec(a+bx)}} + \frac{5}{21} \int \sqrt{\sec(a+bx)} dx \\
&= \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{10 \sin(a+bx)}{21b \sqrt{\sec(a+bx)}} + \frac{1}{21} (5 \sqrt{\cos(a+bx)} \sqrt{\sec(a+bx)}) \int \frac{1}{\sqrt{\cos(a+bx)}} dx \\
&= \frac{10 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{\sec(a+bx)}}{21b} + \frac{2 \sin(a+bx)}{7b \sec^{\frac{5}{2}}(a+bx)} + \frac{10 \sin(a+bx)}{21b \sqrt{\sec(a+bx)}}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 61, normalized size = 0.72

$$\frac{\sqrt{\sec(a+bx)} \left(26 \sin(2(a+bx)) + 3 \sin(4(a+bx)) + 40 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \right)}{84b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(-7/2), x]

[Out] (Sqrt[Sec[a + b*x]]*(40*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + 26*Sin[2*(a + b*x)] + 3*Sin[4*(a + b*x)]))/(84*b)

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{1}{\sec(bx+a)^{\frac{7}{2}}}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(7/2), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(-7/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(7/2),x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(-7/2), x)

maple [B] time = 3.02, size = 199, normalized size = 2.34

$$\frac{2\sqrt{\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\left(48\left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 120\left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 128\left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 21\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}\right)}{21\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(7/2),x)

[Out] -2/21*((2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(48*cos(1/2*b*x+1/2*a)^9-120*cos(1/2*b*x+1/2*a)^7+128*cos(1/2*b*x+1/2*a)^5-72*cos(1/2*b*x+1/2*a)^3+5*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+16*cos(1/2*b*x+1/2*a))/(-2*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(2*cos(1/2*b*x+1/2*a)^2-1)^(1/2)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(7/2),x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(-7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(7/2),x)

[Out] int(1/(1/cos(a + b*x))^(7/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec^{\frac{7}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)**(7/2), x)

[Out] Integral(sec(a + b*x)**(-7/2), x)

3.17 $\int (c \sec(a + bx))^{7/2} dx$

Optimal. Leaf size=98

$$-\frac{6c^4 E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{5b\sqrt{\cos(a+bx)}\sqrt{c\sec(a+bx)}} + \frac{6c^3 \sin(a+bx)\sqrt{c\sec(a+bx)}}{5b} + \frac{2c \sin(a+bx)(c\sec(a+bx))^{5/2}}{5b}$$

[Out] $2/5*c*(c*\sec(b*x+a))^{(5/2)}*\sin(b*x+a)/b-6/5*c^4*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})/b/\cos(b*x+a)^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+6/5*c^3*\sin(b*x+a)*(c*\sec(b*x+a))^{(1/2)}/b$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3771, 2639}

$$\frac{6c^3 \sin(a+bx)\sqrt{c\sec(a+bx)}}{5b} - \frac{6c^4 E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{5b\sqrt{\cos(a+bx)}\sqrt{c\sec(a+bx)}} + \frac{2c \sin(a+bx)(c\sec(a+bx))^{5/2}}{5b}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(7/2), x]

[Out] $(-6*c^4*\text{EllipticE}[(a+b*x)/2, 2])/(5*b*\text{Sqrt}[\text{Cos}[a+b*x]]*\text{Sqrt}[c*\text{Sec}[a+b*x]]) + (6*c^3*\text{Sqrt}[c*\text{Sec}[a+b*x]]*\text{Sin}[a+b*x])/(5*b) + (2*c*(c*\text{Sec}[a+b*x])^{(5/2)}*\text{Sin}[a+b*x])/(5*b)$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x] * (b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n * Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int (c \sec(a + bx))^{7/2} dx &= \frac{2c(c \sec(a + bx))^{5/2} \sin(a + bx)}{5b} + \frac{1}{5} (3c^2) \int (c \sec(a + bx))^{3/2} dx \\
&= \frac{6c^3 \sqrt{c \sec(a + bx)} \sin(a + bx)}{5b} + \frac{2c(c \sec(a + bx))^{5/2} \sin(a + bx)}{5b} - \frac{1}{5} (3c^4) \int \frac{1}{\sqrt{c \sec(a + bx)}} dx \\
&= \frac{6c^3 \sqrt{c \sec(a + bx)} \sin(a + bx)}{5b} + \frac{2c(c \sec(a + bx))^{5/2} \sin(a + bx)}{5b} - \frac{(3c^4) \int \sqrt{\cos(a + bx)}}{5\sqrt{\cos(a + bx)} \sqrt{c}} dx \\
&= -\frac{6c^4 E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5b\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{6c^3 \sqrt{c \sec(a + bx)} \sin(a + bx)}{5b} + \frac{2c(c \sec(a + bx))^{5/2} \sin(a + bx)}{5b}
\end{aligned}$$

Mathematica [A] time = 0.17, size = 62, normalized size = 0.63

$$\frac{c(c \sec(a + bx))^{5/2} \left(7 \sin(a + bx) + 3 \sin(3(a + bx)) - 12 \cos^2(a + bx) E\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{10b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(7/2), x]

[Out] (c*(c*Sec[a + b*x])^(5/2)*(-12*Cos[a + b*x]^(5/2)*EllipticE[(a + b*x)/2, 2] + 7*Sin[a + b*x] + 3*Sin[3*(a + b*x)]))/(10*b)

fricas [F] time = 0.82, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \sec(bx + a)} c^3 \sec(bx + a)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))*c^3*sec(b*x + a)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(7/2), x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(7/2), x)

maple [C] time = 1.01, size = 354, normalized size = 3.61

$$2(-1 + \cos(bx + a))^2 \left(3i \sin(bx + a) \operatorname{EllipticF}\left(\frac{i(-1 + \cos(bx + a))}{\sin(bx + a)}, i\right) (\cos^3(bx + a)) \sqrt{\frac{1}{\cos(bx + a) + 1}} \sqrt{\frac{\cos(bx + a)}{\cos(bx + a) + 1}} - 3i \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(7/2),x)

[Out]
$$-2/5/b*(-1+\cos(b*x+a))^2*(3*I*\sin(b*x+a)*\operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)*\cos(b*x+a)^3*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}-3*I*\sin(b*x+a)*\cos(b*x+a)^3*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}*\operatorname{EllipticE}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)+3*I*\sin(b*x+a)*\operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)*\cos(b*x+a)^2*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}-3*I*\sin(b*x+a)*\cos(b*x+a)^2*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}*\operatorname{EllipticE}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)+3*\cos(b*x+a)^3-2*\cos(b*x+a)^2-1)*\cos(b*x+a)*(\cos(b*x+a)+1)^2*(c/\cos(b*x+a))^{7/2}/\sin(b*x+a)^5$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(7/2),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(7/2),x)

[Out] int((c/cos(a + b*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sec(b*x+a))**(7/2),x)
```

```
[Out] Timed out
```

3.18 $\int (c \sec(a + bx))^{5/2} dx$

Optimal. Leaf size=70

$$\frac{2c^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{3b} + \frac{2c \sin(a + bx) (c \sec(a + bx))^{3/2}}{3b}$$

[Out] $2/3*c*(c*\sec(b*x+a))^{3/2}*\sin(b*x+a)/b+2/3*c^2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a),2^{(1/2)})*\cos(b*x+a)^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}/b$

Rubi [A] time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3771, 2641}

$$\frac{2c^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{3b} + \frac{2c \sin(a + bx) (c \sec(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{5/2}, x]$

[Out] $(2*c^2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(3*b) + (2*c*(c*\text{Sec}[a + b*x])^{3/2}*\text{Sin}[a + b*x])/(3*b)$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3768

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Csc}[c + d*x])^{(n-1)})/(d*(n-1)), x] + \text{Dist}[(b^2*(n-2))/(n-1), \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3771

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n-1)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (c \sec(a + bx))^{5/2} dx &= \frac{2c(c \sec(a + bx))^{3/2} \sin(a + bx)}{3b} + \frac{1}{3}c^2 \int \sqrt{c \sec(a + bx)} dx \\
&= \frac{2c(c \sec(a + bx))^{3/2} \sin(a + bx)}{3b} + \frac{1}{3} \left(c^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)} \right) \int \frac{1}{\sqrt{\cos(a + bx)}} \\
&= \frac{2c^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{3b} + \frac{2c(c \sec(a + bx))^{3/2} \sin(a + bx)}{3b}
\end{aligned}$$

Mathematica [A] time = 0.07, size = 51, normalized size = 0.73

$$\frac{2c^2 \sqrt{c \sec(a + bx)} \left(\tan(a + bx) + \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(5/2),x]

[Out] (2*c^2*Sqrt[c*Sec[a + b*x]]*(Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Tan[a + b*x]))/(3*b)

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \sec(bx + a)} c^2 \sec(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))*c^2*sec(b*x + a)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(5/2), x)

maple [C] time = 0.84, size = 128, normalized size = 1.83

$$\frac{2(-1 + \cos(bx + a)) \left(i \cos(bx + a) \sin(bx + a) \sqrt{\frac{1}{\cos(bx+a)+1}} \sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}} \text{EllipticF}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)}, i\right) - \cos(bx + a) \right)}{3b \sin(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^(5/2),x)`

[Out] $-2/3/b*(-1+\cos(b*x+a))*(I*\cos(b*x+a)*\sin(b*x+a)*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}*\text{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)-\cos(b*x+a)+1)*\cos(b*x+a)*(\cos(b*x+a)+1)^2*(c/\cos(b*x+a))^{5/2}/\sin(b*x+a)^3$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(5/2),x)`

[Out] `int((c/cos(a + b*x))^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(a + bx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))**(5/2),x)`

[Out] `Integral((c*sec(a + b*x))**(5/2), x)`

3.19 $\int (c \sec(a + bx))^{3/2} dx$

Optimal. Leaf size=66

$$\frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - \frac{2c^2 E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}}$$

[Out] $-2*c^2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})/b/\cos(b*x+a)^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+2*c*\sin(b*x+a)*(c*\sec(b*x+a))^{(1/2)}/b$

Rubi [A] time = 0.04, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3771, 2639}

$$\frac{2c \sin(a + bx) \sqrt{c \sec(a + bx)}}{b} - \frac{2c^2 E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(-2*c^2*\text{EllipticE}[(a + b*x)/2, 2])/(b*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (2*c*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sin}[a + b*x])/b$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3768

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Csc}[c + d*x])^{(n - 1)})/(d*(n - 1)), x] + \text{Dist}[(b^2*(n - 2))/(n - 1), \text{Int}[(b*\text{Csc}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 3771

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (c \sec(a + bx))^{3/2} dx &= \frac{2c\sqrt{c \sec(a + bx)} \sin(a + bx)}{b} - c^2 \int \frac{1}{\sqrt{c \sec(a + bx)}} dx \\
&= \frac{2c\sqrt{c \sec(a + bx)} \sin(a + bx)}{b} - \frac{c^2 \int \sqrt{\cos(a + bx)} dx}{\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} \\
&= -\frac{2c^2 E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{b\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{2c\sqrt{c \sec(a + bx)} \sin(a + bx)}{b}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 48, normalized size = 0.73

$$\frac{2c\sqrt{c \sec(a + bx)} \left(\sin(a + bx) - \sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(3/2),x]

[Out] (2*c*Sqrt[c*Sec[a + b*x]]*(-(Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2]) + Sin[a + b*x]))/b

fricas [F] time = 0.94, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \sec(bx + a)} c \sec(bx + a), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))*c*sec(b*x + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(3/2), x)

maple [C] time = 0.98, size = 322, normalized size = 4.88

$$2(\cos(bx+a)+1)^2(-1+\cos(bx+a))^2 \left(i \cos(bx+a) \sin(bx+a) \operatorname{EllipticE}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)}, i\right) \sqrt{\frac{1}{\cos(bx+a)+1}} \sqrt{\dots} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^(3/2),x)`

[Out] $2/b * (\cos(b*x+a)+1)^2 * (-1+\cos(b*x+a))^2 * (I*\cos(b*x+a)*\sin(b*x+a)*\operatorname{EllipticE}(I * (-1+\cos(b*x+a))/\sin(b*x+a), I) * (1/(\cos(b*x+a)+1))^{1/2} * (\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2} - I*\cos(b*x+a)*\sin(b*x+a) * (1/(\cos(b*x+a)+1))^{1/2} * (\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2} * \operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I) + I*\sin(b*x+a) * \operatorname{EllipticE}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I) * (1/(\cos(b*x+a)+1))^{1/2} * (\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2} - I*\sin(b*x+a) * \operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I) * (1/(\cos(b*x+a)+1))^{1/2} * (\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2} - \cos(b*x+a)+1) * \cos(b*x+a) * (c/\cos(b*x+a))^{3/2} / \sin(b*x+a)^5$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx+a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x+a))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{c}{\cos(a+bx)} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a+b*x))^(3/2),x)`

[Out] `int((c/cos(a+b*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(a+bx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sec(b*x+a))**(3/2),x)
```

```
[Out] Integral((c*sec(a + b*x))**(3/2), x)
```

3.20 $\int \sqrt{c \sec(a + bx)} dx$

Optimal. Leaf size=38

$$\frac{2\sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{b}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}/b$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3771, 2641}

$$\frac{2\sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sec[a + b*x]], x]

[Out] $(2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]])/b$

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{c \sec(a + bx)} dx &= \left(\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}\right) \int \frac{1}{\sqrt{\cos(a + bx)}} dx \\ &= \frac{2\sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{b} \end{aligned}$$

Mathematica [A] time = 0.02, size = 38, normalized size = 1.00

$$\frac{2\sqrt{\cos(a+bx)}F\left(\frac{1}{2}(a+bx)\middle|2\right)\sqrt{c\sec(a+bx)}}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sec[a + b*x]], x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[c*Sec[a + b*x]])/b

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c\sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c\sec(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(c*sec(b*x + a)), x)

maple [C] time = 0.92, size = 98, normalized size = 2.58

$$\frac{2i\sqrt{\frac{c}{\cos(bx+a)}}(-1+\cos(bx+a))\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\text{EllipticF}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)}, i\right)(\cos(bx+a)+1)^2}{b\sin(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(1/2), x)

[Out] -2*I/b*(c/cos(b*x+a))^(1/2)*(-1+cos(b*x+a))*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticF(I*(-1+cos(b*x+a))/sin(b*x+a), I)*(cos(b*x+a)+1)^2/sin(b*x+a)^2

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*sec(b*x + a)), x)

mupad [B] time = 0.20, size = 35, normalized size = 0.92

$$\frac{2 \sqrt{\cos(a + bx)} \sqrt{\frac{c}{\cos(a+bx)}} F\left(\frac{a}{2} + \frac{bx}{2} \middle| 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(1/2),x)

[Out] (2*cos(a + b*x)^(1/2)*(c/cos(a + b*x))^(1/2)*ellipticF(a/2 + (b*x)/2, 2))/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sec(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(1/2),x)

[Out] Integral(sqrt(c*sec(a + b*x)), x)

$$3.21 \quad \int \frac{1}{\sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=38

$$\frac{2E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}}$$

[Out] $2*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})/b/\cos(b*x+a)^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3771, 2639}

$$\frac{2E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[c*Sec[a + b*x]], x]

[Out] $(2*\text{EllipticE}[(a + b*x)/2, 2])/(b*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{c \sec(a+bx)}} dx &= \frac{\int \sqrt{\cos(a+bx)} dx}{\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}} \\ &= \frac{2E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{b\sqrt{\cos(a+bx)}\sqrt{c \sec(a+bx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 38, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(a+bx)\middle|2\right)}{b\sqrt{\cos(a+bx)}\sqrt{c\sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[c*Sec[a + b*x]], x]

[Out] (2*EllipticE[(a + b*x)/2, 2])/(b*Sqrt[Cos[a + b*x]]*Sqrt[c*Sec[a + b*x]])

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{c\sec(bx+a)}}{c\sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))/(c*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c\sec(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(1/2), x, algorithm="giac")

[Out] integrate(1/sqrt(c*sec(b*x + a)), x)

maple [C] time = 1.06, size = 306, normalized size = 8.05

$$2\left(i\cos(bx+a)\sin(bx+a)\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\text{EllipticF}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)}, i\right) - i\cos(bx+a)\sin(bx+a)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sec(b*x+a))^(1/2), x)

[Out] 2/b*(I*cos(b*x+a)*sin(b*x+a)*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)*EllipticF(I*(-1+cos(b*x+a))/sin(b*x+a), I)-I*cos(b*x+a)*sin(b*x+a)*EllipticE(I*(-1+cos(b*x+a))/sin(b*x+a), I)*(1/(cos(b*x+a)+1))^(1/2)*(cos

$(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}+I*\sin(b*x+a)*\text{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)*(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}-I*\sin(b*x+a)*\text{EllipticE}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)*(1/(\cos(b*x+a)+1))^{(1/2)}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{(1/2)}-\cos(b*x+a)^2+\cos(b*x+a))*(c/\cos(b*x+a))^{(1/2)}/\sin(b*x+a)/c$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(c*sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c/cos(a + b*x))^(1/2),x)

[Out] int(1/(c/cos(a + b*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))**(1/2),x)

[Out] Integral(1/sqrt(c*sec(a + b*x)), x)

$$3.22 \quad \int \frac{1}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=72

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{c \sec(a+bx)}}{3bc^2} + \frac{2 \sin(a+bx)}{3bc\sqrt{c \sec(a+bx)}}$$

[Out] 2/3*sin(b*x+a)/b/c/(c*sec(b*x+a))^(1/2)+2/3*(cos(1/2*b*x+1/2*a)^2)^(1/2)/cos(1/2*b*x+1/2*a)*EllipticF(sin(1/2*b*x+1/2*a),2^(1/2))*cos(b*x+a)^(1/2)*(c*sec(b*x+a))^(1/2)/b/c^2

Rubi [A] time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3769, 3771, 2641}

$$\frac{2\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{c \sec(a+bx)}}{3bc^2} + \frac{2 \sin(a+bx)}{3bc\sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(-3/2), x]

[Out] (2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2]*Sqrt[c*Sec[a + b*x]])/(3*b*c^2) + (2*Sin[a + b*x])/(3*b*c*Sqrt[c*Sec[a + b*x]])

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d^n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(c \sec(a + bx))^{3/2}} dx &= \frac{2 \sin(a + bx)}{3bc\sqrt{c \sec(a + bx)}} + \frac{\int \sqrt{c \sec(a + bx)} dx}{3c^2} \\
&= \frac{2 \sin(a + bx)}{3bc\sqrt{c \sec(a + bx)}} + \frac{(\sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}) \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{3c^2} \\
&= \frac{2\sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{3bc^2} + \frac{2 \sin(a + bx)}{3bc\sqrt{c \sec(a + bx)}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 59, normalized size = 0.82

$$\frac{\sec^2(a + bx) \left(\sin(2(a + bx)) + 2\sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{3b(c \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(-3/2),x]

[Out] (Sec[a + b*x]^2*(2*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + Sin[2*(a + b*x)]))/(3*b*(c*Sec[a + b*x])^(3/2))

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{c \sec(bx + a)}}{c^2 \sec(bx + a)^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))/(c^2*sec(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^{-3/2}, x)

maple [C] time = 0.82, size = 131, normalized size = 1.82

$$\frac{2(\cos(bx+a)+1)^2(-1+\cos(bx+a))\left(i\sin(bx+a)\operatorname{EllipticF}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)},i\right)\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\right)}{3b\left(\frac{c}{\cos(bx+a)}\right)^{\frac{3}{2}}\cos(bx+a)^2\sin(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sec(b*x+a))^{3/2},x)

[Out]
$$-2/3/b*(\cos(b*x+a)+1)^2*(-1+\cos(b*x+a))*(I*\sin(b*x+a)*\operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a),I)*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}-\cos(b*x+a)^2+\cos(b*x+a))/(c/\cos(b*x+a))^{3/2}/\cos(b*x+a)^2/\sin(b*x+a)^3$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx+a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^{3/2},x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^{-3/2}, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c/cos(a + b*x))^{3/2},x)

[Out] int(1/(c/cos(a + b*x))^{3/2}, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(a+bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(c*sec(b*x+a))**(3/2),x)
```

```
[Out] Integral((c*sec(a + b*x))**(-3/2), x)
```


$$3.23 \quad \int \frac{1}{(c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5bc^2\sqrt{\cos(a+bx)}\sqrt{c\sec(a+bx)}} + \frac{2\sin(a+bx)}{5bc(c\sec(a+bx))^{3/2}}$$

[Out] $2/5*\sin(b*x+a)/b/c/(c*\sec(b*x+a))^{(3/2)}+6/5*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticE}(\sin(1/2*b*x+1/2*a),2^{(1/2)})/b/c^2/\cos(b*x+a)^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3769, 3771, 2639}

$$\frac{6E\left(\frac{1}{2}(a+bx)\middle|2\right)}{5bc^2\sqrt{\cos(a+bx)}\sqrt{c\sec(a+bx)}} + \frac{2\sin(a+bx)}{5bc(c\sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{(-5/2)}, x]$

[Out] $(6*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*c^2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(5*b*c*(c*\text{Sec}[a + b*x])^{(3/2)})$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3769

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 3771

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{n*}\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int \frac{1}{(c \sec(a + bx))^{5/2}} dx &= \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}} + \frac{3 \int \frac{1}{\sqrt{c \sec(a+bx)}} dx}{5c^2} \\
&= \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}} + \frac{3 \int \sqrt{\cos(a + bx)} dx}{5c^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} \\
&= \frac{6E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{5bc^2 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{2 \sin(a + bx)}{5bc(c \sec(a + bx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.07, size = 60, normalized size = 0.83

$$\frac{\sqrt{c \sec(a + bx)} \left(\sin(a + bx) + \sin(3(a + bx)) + 12\sqrt{\cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{10bc^3}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(-5/2), x]

[Out] (Sqrt[c*Sec[a + b*x]]*(12*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + Sin[a + b*x] + Sin[3*(a + b*x)]))/(10*b*c^3)

fricas [F] time = 0.89, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{c \sec(bx + a)}}{c^3 \sec(bx + a)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))/(c^3*sec(b*x + a)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(-5/2), x)

maple [C] time = 0.97, size = 321, normalized size = 4.46

$$2 \left(-3i \cos(bx + a) \sin(bx + a) \sqrt{\frac{1}{\cos(bx+a)+1}} \sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)}, i\right) + 3i \cos(bx + a) \sin(bx + a) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sec(b*x+a))^(5/2), x)

[Out]
$$-2/5/b * (-3*I*\cos(b*x+a)*\sin(b*x+a)*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I)+3*I*\cos(b*x+a)*\sin(b*x+a)*\operatorname{EllipticE}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I)*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}-3*I*\sin(b*x+a)*\operatorname{EllipticF}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I)*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}+3*I*\sin(b*x+a)*\operatorname{EllipticE}(I*(-1+\cos(b*x+a))/\sin(b*x+a), I)*(1/(\cos(b*x+a)+1))^{1/2}*(\cos(b*x+a)/(\cos(b*x+a)+1))^{1/2}+\cos(b*x+a)^4+2*\cos(b*x+a)^2-3*\cos(b*x+a))/(c/\cos(b*x+a))^{5/2}/\cos(b*x+a)^3/\sin(b*x+a)$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(5/2), x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c/cos(a + b*x))^(5/2), x)

[Out] int(1/(c/cos(a + b*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(a + bx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))**(5/2),x)

[Out] Integral((c*sec(a + b*x))**(-5/2), x)

$$3.24 \quad \int \frac{1}{(c \sec(a+bx))^{7/2}} dx$$

Optimal. Leaf size=100

$$\frac{10\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{c \sec(a+bx)}}{21bc^4} + \frac{10 \sin(a+bx)}{21bc^3 \sqrt{c \sec(a+bx)}} + \frac{2 \sin(a+bx)}{7bc(c \sec(a+bx))^{5/2}}$$

[Out] $2/7*\sin(b*x+a)/b/c/(c*\sec(b*x+a))^{(5/2)}+10/21*\sin(b*x+a)/b/c^3/(c*\sec(b*x+a))^{(1/2)}+10/21*(\cos(1/2*b*x+1/2*a)^2)^{(1/2)}/\cos(1/2*b*x+1/2*a)*\text{EllipticF}(\sin(1/2*b*x+1/2*a), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}/b/c^4$

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3769, 3771, 2641}

$$\frac{10 \sin(a+bx)}{21bc^3 \sqrt{c \sec(a+bx)}} + \frac{10\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right) \sqrt{c \sec(a+bx)}}{21bc^4} + \frac{2 \sin(a+bx)}{7bc(c \sec(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(-7/2), x]

[Out] $(10*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(21*b*c^4) + (2*\text{Sin}[a + b*x])/(7*b*c*(c*\text{Sec}[a + b*x])^{(5/2)}) + (10*\text{Sin}[a + b*x])/(21*b*c^3*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(c \sec(a + bx))^{7/2}} dx &= \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} + \frac{5 \int \frac{1}{(c \sec(a + bx))^{3/2}} dx}{7c^2} \\
&= \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} + \frac{10 \sin(a + bx)}{21bc^3 \sqrt{c \sec(a + bx)}} + \frac{5 \int \sqrt{c \sec(a + bx)} dx}{21c^4} \\
&= \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} + \frac{10 \sin(a + bx)}{21bc^3 \sqrt{c \sec(a + bx)}} + \frac{(5 \sqrt{\cos(a + bx)} \sqrt{c \sec(a + bx)}) \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{21c^4} \\
&= \frac{10 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{c \sec(a + bx)}}{21bc^4} + \frac{2 \sin(a + bx)}{7bc(c \sec(a + bx))^{5/2}} + \frac{10 \sin(a + bx)}{21bc^3 \sqrt{c \sec(a + bx)}}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 66, normalized size = 0.66

$$\frac{\sqrt{c \sec(a + bx)} \left(26 \sin(2(a + bx)) + 3 \sin(4(a + bx)) + 40 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \middle| 2\right) \right)}{84bc^4}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(-7/2), x]

[Out] (Sqrt[c*Sec[a + b*x]]*(40*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + 26*Sin[2*(a + b*x)] + 3*Sin[4*(a + b*x)]))/(84*b*c^4)

fricas [F] time = 0.76, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{c \sec(bx + a)}}{c^4 \sec(bx + a)^4}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(c*sec(b*x + a))/(c^4*sec(b*x + a)^4), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(-7/2), x)

maple [C] time = 0.94, size = 153, normalized size = 1.53

$$\frac{2(\cos(bx+a)+1)^2(-1+\cos(bx+a))\left(-5i\sin(bx+a)\operatorname{EllipticF}\left(\frac{i(-1+\cos(bx+a))}{\sin(bx+a)},i\right)\sqrt{\frac{1}{\cos(bx+a)+1}}\sqrt{\frac{\cos(bx+a)}{\cos(bx+a)+1}}\right)}{21b\left(\frac{c}{\cos(bx+a)}\right)^{\frac{7}{2}}\cos(bx+a)^4\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sec(b*x+a))^(7/2),x)

[Out] 2/21/b*(cos(b*x+a)+1)^2*(-1+cos(b*x+a))*(-5*I*sin(b*x+a)*EllipticF(I*(-1+cos(b*x+a))/sin(b*x+a),I)*(1/(cos(b*x+a)+1))^(1/2)*(cos(b*x+a)/(cos(b*x+a)+1))^(1/2)+3*cos(b*x+a)^4-3*cos(b*x+a)^3+5*cos(b*x+a)^2-5*cos(b*x+a))/(c/cos(b*x+a))^(7/2)/cos(b*x+a)^4/sin(b*x+a)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(7/2),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(-7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c/cos(a + b*x))^(7/2),x)

[Out] int(1/(c/cos(a + b*x))^(7/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(a + bx))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(c*sec(b*x+a))**(7/2),x)
```

```
[Out] Integral((c*sec(a + b*x))**(-7/2), x)
```


3.25 $\int \sec^{\frac{4}{3}}(a + bx) dx$

Optimal. Leaf size=51

$$\frac{3 \sin(a + bx) \sqrt[3]{\sec(a + bx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*sec(b*x+a)^(1/3)*sin(b*x+a)/b/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{3 \sin(a + bx) \sqrt[3]{\sec(a + bx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(4/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*Sec[a + b*x]^(1/3)*Sin[a + b*x])/(b*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \sec^{\frac{4}{3}}(a + bx) dx = \sqrt[3]{\cos(a + bx)} \sqrt[3]{\sec(a + bx)} \int \frac{1}{\cos^{\frac{4}{3}}(a + bx)} dx$$

$$= \frac{{}_3F_2\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a + bx)\right) \sqrt[3]{\sec(a + bx)} \sin(a + bx)}{b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 1.08

$$\frac{3\sqrt{-\tan^2(a + bx)} \csc(a + bx) \sqrt[3]{\sec(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sec^2(a + bx)\right)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(4/3), x]

[Out] (3*Csc[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[a + b*x]^2]*Sec[a + b*x]^(1/3)*Sqrt[-Tan[a + b*x]^2])/(4*b)

fricas [F] time = 0.89, size = 0, normalized size = 0.00

$$\text{integral}\left(\sec(bx + a)^{\frac{4}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(4/3), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(4/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(4/3), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(4/3), x)

maple [F] time = 0.63, size = 0, normalized size = 0.00

$$\int \sec^{\frac{4}{3}}(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^(4/3), x)`

[Out] `int(sec(b*x+a)^(4/3), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^(4/3), x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^(4/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{1}{\cos(a + bx)} \right)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(a + b*x))^(4/3), x)`

[Out] `int((1/cos(a + b*x))^(4/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^{\frac{4}{3}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**(4/3), x)`

[Out] `Integral(sec(a + b*x)**(4/3), x)`

3.26 $\int \sec^{\frac{2}{3}}(a + bx) dx$

Optimal. Leaf size=51

$$\frac{3 \sin(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)} \sqrt[3]{\sec(a + bx)}}$$

[Out] -3*hypergeom([1/6, 1/2], [7/6], cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(1/3)/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{3 \sin(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)} \sqrt[3]{\sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(2/3), x]

[Out] (-3*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[a + b*x]^2]*Sin[a + b*x])/(b*Sec[a + b*x]^(1/3)*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \sec^{\frac{2}{3}}(a + bx) dx = \cos^{\frac{2}{3}}(a + bx) \sec^{\frac{2}{3}}(a + bx) \int \frac{1}{\cos^{\frac{2}{3}}(a + bx)} dx$$

$$= -\frac{{}_3F_2\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}; \cos^2(a + bx)\right) \sin(a + bx)}{b\sqrt[3]{\sec(a + bx)} \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 1.08

$$\frac{3\sqrt{-\tan^2(a + bx)} \csc(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sec^2(a + bx)\right)}{2b\sqrt[3]{\sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(2/3), x]

[Out] (3*Csc[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(2*b*Sec[a + b*x]^(1/3))

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral}\left(\sec(bx + a)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(2/3), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(2/3), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(2/3), x)

maple [F] time = 0.61, size = 0, normalized size = 0.00

$$\int \sec^{\frac{2}{3}}(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^(2/3),x)`

[Out] `int(sec(b*x+a)^(2/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^(2/3),x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^(2/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{1}{\cos(a + bx)} \right)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(a + b*x))^(2/3),x)`

[Out] `int((1/cos(a + b*x))^(2/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^{\frac{2}{3}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**(2/3),x)`

[Out] `Integral(sec(a + b*x)**(2/3), x)`

3.27 $\int \sqrt[3]{\sec(a + bx)} dx$

Optimal. Leaf size=53

$$\frac{3 \sin(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a + bx)\right)}{2b\sqrt{\sin^2(a + bx)} \sec^{\frac{2}{3}}(a + bx)}$$

[Out] $-3/2*\text{hypergeom}([1/3, 1/2], [4/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/\sec(b*x+a)^{(2/3)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{3 \sin(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a + bx)\right)}{2b\sqrt{\sin^2(a + bx)} \sec^{\frac{2}{3}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(1/3), x]

[Out] $(-3*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(2*b*\text{Sec}[a + b*x]^{(2/3)}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \sqrt[3]{\sec(a+bx)} dx = \sqrt[3]{\cos(a+bx)} \sqrt[3]{\sec(a+bx)} \int \frac{1}{\sqrt[3]{\cos(a+bx)}} dx$$

$$= -\frac{{}_3F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right) \sin(a+bx)}{2b \sec^{\frac{2}{3}}(a+bx) \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.04, size = 53, normalized size = 1.00

$$\frac{3\sqrt{-\tan^2(a+bx)} \csc(a+bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \sec^2(a+bx)\right)}{b \sec^{\frac{2}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(1/3), x]

[Out] (3*Csc[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(b*Sec[a + b*x]^(2/3))

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\text{integral}\left(\sec(bx+a)^{\frac{1}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(1/3), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx+a)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^(1/3), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(1/3), x)

maple [F] time = 0.68, size = 0, normalized size = 0.00

$$\int \sec^{\frac{1}{3}}(bx+a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^(1/3),x)`

[Out] `int(sec(b*x+a)^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^(1/3),x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{1}{\cos(a + bx)} \right)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(a + b*x))^(1/3),x)`

[Out] `int((1/cos(a + b*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{\sec(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**(1/3),x)`

[Out] `Integral(sec(a + b*x)**(1/3), x)`

$$3.28 \quad \int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx$$

Optimal. Leaf size=53

$$\frac{3 \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a+bx)\right)}{4b\sqrt{\sin^2(a+bx)} \sec^{\frac{4}{3}}(a+bx)}$$

[Out] $-3/4*\text{hypergeom}([1/2, 2/3], [5/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/\sec(b*x+a)^{(4/3)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{3 \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a+bx)\right)}{4b\sqrt{\sin^2(a+bx)} \sec^{\frac{4}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(-1/3), x]

[Out] $(-3*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/ (4*b*\text{Sec}[a + b*x]^{(4/3)}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \frac{1}{\sqrt[3]{\sec(a+bx)}} dx = \cos^{\frac{2}{3}}(a+bx) \sec^{\frac{2}{3}}(a+bx) \int \sqrt[3]{\cos(a+bx)} dx$$

$$= -\frac{{}_3F_2\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a+bx)\right) \sin(a+bx)}{4b \sec^{\frac{4}{3}}(a+bx) \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.08, size = 53, normalized size = 1.00

$$\frac{3\sqrt{-\tan^2(a+bx)} \csc(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sec^2(a+bx)\right)}{b \sec^{\frac{4}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(-1/3), x]

[Out] (-3*Csc[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(b*Sec[a + b*x]^(4/3))

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\sec(bx+a)^{\frac{1}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(1/3), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(-1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(1/3), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(-1/3), x)

maple [F] time = 0.55, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(1/3), x)

[Out] int(1/sec(b*x+a)^(1/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(1/3), x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(-1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(1/3), x)

[Out] int(1/(1/cos(a + b*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{\sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)**(1/3), x)

[Out] Integral(sec(a + b*x)**(-1/3), x)

$$3.29 \quad \int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx$$

Optimal. Leaf size=53

$$-\frac{3 \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a+bx)\right)}{5b\sqrt{\sin^2(a+bx)} \sec^{\frac{5}{3}}(a+bx)}$$

[Out] $-3/5*\text{hypergeom}([1/2, 5/6], [11/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/\sec(b*x+a)^{(5/3)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$-\frac{3 \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a+bx)\right)}{5b\sqrt{\sin^2(a+bx)} \sec^{\frac{5}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(-2/3), x]

[Out] $(-3*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/((5*b*\text{Sec}[a + b*x])^{(5/3)}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2)]/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/((Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \frac{1}{\sec^{\frac{2}{3}}(a+bx)} dx = \sqrt[3]{\cos(a+bx)} \sqrt[3]{\sec(a+bx)} \int \cos^{\frac{2}{3}}(a+bx) dx$$

$$= -\frac{{}_3F_2\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a+bx)\right) \sin(a+bx)}{5b \sec^{\frac{5}{3}}(a+bx) \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.08, size = 55, normalized size = 1.04

$$-\frac{3\sqrt{-\tan^2(a+bx)} \csc(a+bx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sec^2(a+bx)\right)}{2b \sec^{\frac{5}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(-2/3), x]

[Out] (-3*Csc[a + b*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(2*b*Sec[a + b*x]^(5/3))

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\sec(bx+a)^{\frac{2}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(2/3), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(-2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(2/3), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(-2/3), x)

maple [F] time = 0.60, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(2/3),x)

[Out] int(1/sec(b*x+a)^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(-2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(2/3),x)

[Out] int(1/(1/cos(a + b*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec^{\frac{2}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)**(2/3),x)

[Out] Integral(sec(a + b*x)**(-2/3), x)

$$3.30 \quad \int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx$$

Optimal. Leaf size=53

$$\frac{3 \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a+bx)\right)}{7b\sqrt{\sin^2(a+bx)} \sec^{\frac{7}{3}}(a+bx)}$$

[Out] -3/7*hypergeom([1/2, 7/6], [13/6], cos(b*x+a)^2)*sin(b*x+a)/b/sec(b*x+a)^(7/3)/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.02, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{3 \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a+bx)\right)}{7b\sqrt{\sin^2(a+bx)} \sec^{\frac{7}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^(-4/3), x]

[Out] (-3*Hypergeometric2F1[1/2, 7/6, 13/6, Cos[a + b*x]^2]*Sin[a + b*x])/(7*b*Sec[a + b*x]^(7/3)*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \frac{1}{\sec^{\frac{4}{3}}(a+bx)} dx = \cos^{\frac{2}{3}}(a+bx) \sec^{\frac{2}{3}}(a+bx) \int \cos^{\frac{4}{3}}(a+bx) dx$$

$$= -\frac{{}_3F_2\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a+bx)\right) \sin(a+bx)}{7b \sec^{\frac{7}{3}}(a+bx) \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.07, size = 55, normalized size = 1.04

$$-\frac{3\sqrt{-\tan^2(a+bx)} \csc(a+bx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \sec^2(a+bx)\right)}{4b \sec^{\frac{7}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^(-4/3), x]

[Out] (-3*Csc[a + b*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(4*b*Sec[a + b*x]^(7/3))

fricas [F] time = 0.76, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{1}{\sec(bx+a)^{\frac{4}{3}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(4/3), x, algorithm="fricas")

[Out] integral(sec(b*x + a)^(-4/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx+a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(4/3), x, algorithm="giac")

[Out] integrate(sec(b*x + a)^(-4/3), x)

maple [F] time = 0.54, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(b*x+a)^(4/3), x)

[Out] int(1/sec(b*x+a)^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec(bx + a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)^(4/3), x, algorithm="maxima")

[Out] integrate(sec(b*x + a)^(-4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(a+bx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(a + b*x))^(4/3), x)

[Out] int(1/(1/cos(a + b*x))^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sec^{\frac{4}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(b*x+a)**(4/3), x)

[Out] Integral(sec(a + b*x)**(-4/3), x)

3.31 $\int (c \sec(a + bx))^{4/3} dx$

Optimal. Leaf size=54

$$\frac{3c \sin(a + bx) \sqrt[3]{c \sec(a + bx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)}}$$

[Out] 3*c*hypergeom([-1/6, 1/2], [5/6], cos(b*x+a)^2)*(c*sec(b*x+a))^(1/3)*sin(b*x+a)/b/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$\frac{3c \sin(a + bx) \sqrt[3]{c \sec(a + bx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a + bx)\right)}{b \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(4/3), x]

[Out] (3*c*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[a + b*x]^2]*(c*Sec[a + b*x])^(1/3)*Sin[a + b*x])/(b*Sqrt[Sin[a + b*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int (c \sec(a + bx))^{4/3} dx = \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \frac{1}{\left(\frac{\cos(a+bx)}{c}\right)^{4/3}} dx$$

$$= \frac{3c {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(a + bx)\right) \sqrt[3]{c \sec(a + bx)} \sin(a + bx)}{b\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.05, size = 57, normalized size = 1.06

$$\frac{3\sqrt{-\tan^2(a + bx)} \cot(a + bx)(c \sec(a + bx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sec^2(a + bx)\right)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(4/3),x]

[Out] (3*Cot[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[a + b*x]^2]*(c*Sec[a + b*x])^(4/3)*Sqrt[-Tan[a + b*x]^2])/(4*b)

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left((c \sec(bx + a))^{1/3} c \sec(bx + a), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(4/3),x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^(1/3)*c*sec(b*x + a), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{4/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(4/3),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(4/3), x)

maple [F] time = 0.54, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{4/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^(4/3),x)`

[Out] `int((c*sec(b*x+a))^(4/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec (bx + a))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^(4/3),x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^(4/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{c}{\cos (a + bx)} \right)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(4/3),x)`

[Out] `int((c/cos(a + b*x))^(4/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec (a + bx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))**(4/3),x)`

[Out] `Integral((c*sec(a + b*x))**(4/3), x)`

3.32 $\int (c \sec(a + bx))^{2/3} dx$

Optimal. Leaf size=54

$$-\frac{3c \sin(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a + bx)\right)}{b\sqrt{\sin^2(a + bx)} \sqrt[3]{c \sec(a + bx)}}$$

[Out] $-3*c*\text{hypergeom}([1/6, 1/2], [7/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(c*\sec(b*x+a))^{1/3}/(\sin(b*x+a)^2)^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$-\frac{3c \sin(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a + bx)\right)}{b\sqrt{\sin^2(a + bx)} \sqrt[3]{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(2/3), x]

[Out] $(-3*c*\text{Hypergeometric2F1}[1/6, 1/2, 7/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(b*(c*\text{Sec}[a + b*x])^{1/3}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int (c \sec(a + bx))^{2/3} dx = \left(\frac{\cos(a + bx)}{c} \right)^{2/3} (c \sec(a + bx))^{2/3} \int \frac{1}{\left(\frac{\cos(a+bx)}{c} \right)^{2/3}} dx$$

$$= -\frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(a + bx)\right) (c \sec(a + bx))^{2/3} \sin(a + bx)}{b \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.04, size = 57, normalized size = 1.06

$$\frac{3 \sqrt{-\tan^2(a + bx)} \cot(a + bx) (c \sec(a + bx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sec^2(a + bx)\right)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(2/3), x]

[Out] (3*Cot[a + b*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[a + b*x]^2]*(c*Sec[a + b*x])^(2/3)*Sqrt[-Tan[a + b*x]^2])/(2*b)

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(c \sec(bx + a)\right)^{\frac{2}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(2/3), x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^(2/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(2/3), x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(2/3), x)

maple [F] time = 0.59, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^(2/3),x)`

[Out] `int((c*sec(b*x+a))^(2/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^(2/3),x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^(2/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(2/3),x)`

[Out] `int((c/cos(a + b*x))^(2/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(a + bx))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))**(2/3),x)`

[Out] `Integral((c*sec(a + b*x))**(2/3), x)`

3.33 $\int \sqrt[3]{c \sec(a + bx)} dx$

Optimal. Leaf size=56

$$-\frac{3c \sin(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a + bx)\right)}{2b\sqrt{\sin^2(a + bx)}(c \sec(a + bx))^{2/3}}$$

[Out] $-3/2*c*\text{hypergeom}([1/3, 1/2], [4/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/(c*\sec(b*x+a))^{(2/3)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$-\frac{3c \sin(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a + bx)\right)}{2b\sqrt{\sin^2(a + bx)}(c \sec(a + bx))^{2/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{(1/3)}, x]$

[Out] $(-3*c*\text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(2*b*(c*\text{Sec}[a + b*x])^{(2/3)}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)(x_)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n - 1)}*((\text{Sin}[c + d*x]/b)^{(n - 1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[n]$

Rubi steps

$$\int \sqrt[3]{c \sec(a+bx)} dx = \sqrt[3]{\frac{\cos(a+bx)}{c}} \sqrt[3]{c \sec(a+bx)} \int \frac{1}{\sqrt[3]{\frac{\cos(a+bx)}{c}}} dx$$

$$= -\frac{3 \cos(a+bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(a+bx)\right) \sqrt[3]{c \sec(a+bx)} \sin(a+bx)}{2b\sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.98

$$\frac{3\sqrt{-\tan^2(a+bx)} \cot(a+bx) \sqrt[3]{c \sec(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \sec^2(a+bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(1/3),x]

[Out] (3*Cot[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[a + b*x]^2]*(c*Sec[a + b*x])^(1/3)*Sqrt[-Tan[a + b*x]^2])/b

fricas [F] time = 0.53, size = 0, normalized size = 0.00

$$\text{integral}\left((c \sec(bx + a))^{\frac{1}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/3),x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^(1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/3),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(1/3), x)

maple [F] time = 0.74, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^(1/3),x)`

[Out] `int((c*sec(b*x+a))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^(1/3),x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(1/3),x)`

[Out] `int((c/cos(a + b*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{c \sec(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))**(1/3),x)`

[Out] `Integral((c*sec(a + b*x))**(1/3), x)`

$$3.34 \quad \int \frac{1}{\sqrt[3]{c \sec(a+bx)}} dx$$

Optimal. Leaf size=56

$$-\frac{3c \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a+bx)\right)}{4b\sqrt{\sin^2(a+bx)}(c \sec(a+bx))^{4/3}}$$

[Out] $-3/4*c*\text{hypergeom}([1/2, 2/3], [5/3], \cos(b*x+a)^2)*\sin(b*x+a)/b/(c*\sec(b*x+a))^{4/3}/(\sin(b*x+a)^2)^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$-\frac{3c \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a+bx)\right)}{4b\sqrt{\sin^2(a+bx)}(c \sec(a+bx))^{4/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{-1/3}, x]$

[Out] $(-3*c*\text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(4*b*(c*\text{Sec}[a + b*x])^{4/3}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x])*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $! \text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $! \text{IntegerQ}[n]$

Rubi steps

$$\int \frac{1}{\sqrt[3]{c \sec(a+bx)}} dx = \left(\frac{\cos(a+bx)}{c} \right)^{2/3} (c \sec(a+bx))^{2/3} \int \sqrt[3]{\frac{\cos(a+bx)}{c}} dx$$

$$= \frac{3 \cos^2(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(a+bx)\right) (c \sec(a+bx))^{2/3} \sin(a+bx)}{4bc \sqrt{\sin^2(a+bx)}}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 0.98

$$\frac{3\sqrt{-\tan^2(a+bx)} \cot(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sec^2(a+bx)\right)}{b\sqrt[3]{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(-1/3), x]

[Out] (-3*Cot[a + b*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(b*(c*Sec[a + b*x])^(1/3))

fricas [F] time = 0.82, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(c \sec(bx+a))^{\frac{2}{3}}}{c \sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(1/3), x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^(2/3)/(c*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx+a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(1/3), x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(1/3), x)

maple [F] time = 0.53, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(c*sec(b*x+a))^(1/3), x)`

[Out] `int(1/(c*sec(b*x+a))^(1/3), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(c*sec(b*x+a))^(1/3), x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^(-1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(c/cos(a + b*x))^(1/3), x)`

[Out] `int(1/(c/cos(a + b*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{c \sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(c*sec(b*x+a))**(1/3), x)`

[Out] `Integral((c*sec(a + b*x))**(-1/3), x)`

$$3.35 \quad \int \frac{1}{(c \sec(a+bx))^{2/3}} dx$$

Optimal. Leaf size=56

$$\frac{3c \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a+bx)\right)}{5b \sqrt{\sin^2(a+bx)} (c \sec(a+bx))^{5/3}}$$

[Out] $-3/5*c*\text{hypergeom}([1/2, 5/6], [11/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(c*\sec(b*x+a))^{(5/3)}/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$\frac{3c \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a+bx)\right)}{5b \sqrt{\sin^2(a+bx)} (c \sec(a+bx))^{5/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{(-2/3)}, x]$

[Out] $(-3*c*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/((5*b*(c*\text{Sec}[a + b*x])^{(5/3)}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)(x_)]*(b_*))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/((\text{Sin}[c + d*x]/b)^n, x)], x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \frac{1}{(c \sec(a + bx))^{2/3}} dx = \sqrt[3]{\frac{\cos(a + bx)}{c}} \sqrt[3]{c \sec(a + bx)} \int \left(\frac{\cos(a + bx)}{c}\right)^{2/3} dx$$

$$= -\frac{3 \cos^2(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(a + bx)\right) \sqrt[3]{c \sec(a + bx)} \sin(a + bx)}{5bc \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.05, size = 57, normalized size = 1.02

$$-\frac{3\sqrt{-\tan^2(a + bx)} \cot(a + bx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sec^2(a + bx)\right)}{2b(c \sec(a + bx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(-2/3), x]

[Out] (-3*Cot[a + b*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(2*b*(c*Sec[a + b*x])^(2/3))

fricas [F] time = 0.90, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(c \sec(bx + a))^{1/3}}{c \sec(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(2/3), x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^(1/3)/(c*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(2/3), x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(1/3)/(c*sec(b*x + a)), x)

maple [F] time = 0.54, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sec(b*x+a))^(2/3),x)

[Out] int(1/(c*sec(b*x+a))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(-2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c/cos(a + b*x))^(2/3),x)

[Out] int(1/(c/cos(a + b*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(a + bx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))**(2/3),x)

[Out] Integral((c*sec(a + b*x))**(-2/3), x)

$$3.36 \quad \int \frac{1}{(c \sec(a+bx))^{4/3}} dx$$

Optimal. Leaf size=56

$$\frac{3c \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a+bx)\right)}{7b\sqrt{\sin^2(a+bx)}(c \sec(a+bx))^{7/3}}$$

[Out] $-3/7*c*\text{hypergeom}([1/2, 7/6], [13/6], \cos(b*x+a)^2)*\sin(b*x+a)/b/(c*\sec(b*x+a))^{7/3}/(\sin(b*x+a)^2)^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$\frac{3c \sin(a+bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a+bx)\right)}{7b\sqrt{\sin^2(a+bx)}(c \sec(a+bx))^{7/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^{(-4/3)}, x]$

[Out] $(-3*c*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[a + b*x]^2]*\text{Sin}[a + b*x])/(7*b*(c*\text{Sec}[a + b*x])^{7/3}*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $!\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $!\text{IntegerQ}[n]$

Rubi steps

$$\int \frac{1}{(c \sec(a + bx))^{4/3}} dx = \left(\frac{\cos(a + bx)}{c} \right)^{2/3} (c \sec(a + bx))^{2/3} \int \left(\frac{\cos(a + bx)}{c} \right)^{4/3} dx$$

$$= -\frac{3 \cos^3(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(a + bx)\right) (c \sec(a + bx))^{2/3} \sin(a + bx)}{7bc^2 \sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.07, size = 57, normalized size = 1.02

$$\frac{3\sqrt{-\tan^2(a + bx)} \cot(a + bx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \sec^2(a + bx)\right)}{4b(c \sec(a + bx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(-4/3), x]

[Out] (-3*Cot[a + b*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[a + b*x]^2]*Sqrt[-Tan[a + b*x]^2])/(4*b*(c*Sec[a + b*x])^(4/3))

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(c \sec(bx + a))^{2/3}}{c^2 \sec(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(4/3), x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^(2/3)/(c^2*sec(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(4/3), x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(4/3), x)

maple [F] time = 0.45, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sec(b*x+a))^(4/3), x)

[Out] int(1/(c*sec(b*x+a))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(bx + a))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))^(4/3), x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(-4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c/cos(a + b*x))^(4/3), x)

[Out] int(1/(c/cos(a + b*x))^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(a + bx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sec(b*x+a))**(4/3), x)

[Out] Integral((c*sec(a + b*x))**(-4/3), x)

3.37 $\int \sec^n(a + bx) dx$

Optimal. Leaf size=70

$$\frac{\sin(a + bx) \sec^{n-1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

[Out] -hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*sec(b*x+a)^(-1+n)*sin(b*x+a)/b/(1-n)/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3772, 2643}

$$\frac{\sin(a + bx) \sec^{n-1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^n, x]

[Out] -((Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*Sec[a + b*x]^(-1 + n)*Sin[a + b*x])/(b*(1 - n)*Sqrt[Sin[a + b*x]^2]))

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \sec^n(a + bx) dx = \cos^n(a + bx) \sec^n(a + bx) \int \cos^{-n}(a + bx) dx$$

$$= -\frac{{}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) \sec^{-1+n}(a + bx) \sin(a + bx)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.05, size = 61, normalized size = 0.87

$$\frac{\sqrt{-\tan^2(a + bx)} \csc(a + bx) \sec^{n-1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{n+2}{2}; \sec^2(a + bx)\right)}{bn}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^n, x]

[Out] (Csc[a + b*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[a + b*x]^2]*Sec[a + b*x]^(-1 + n)*Sqrt[-Tan[a + b*x]^2])/(b*n)

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}(\sec(bx + a)^n, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^n,x, algorithm="fricas")

[Out] integral(sec(b*x + a)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^n,x, algorithm="giac")

[Out] integrate(sec(b*x + a)^n, x)

maple [F] time = 1.46, size = 0, normalized size = 0.00

$$\int \sec^n(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^n,x)`

[Out] `int(sec(b*x+a)^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec(bx + a)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^n,x, algorithm="maxima")`

[Out] `integrate(sec(b*x + a)^n, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{1}{\cos(a + bx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(a + b*x))^n,x)`

[Out] `int((1/cos(a + b*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sec^n(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**n,x)`

[Out] `Integral(sec(a + b*x)**n, x)`

3.38 $\int (c \sec(a + bx))^n dx$

Optimal. Leaf size=73

$$\frac{c \sin(a + bx)(c \sec(a + bx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

[Out] -c*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*(c*sec(b*x+a))^(n-1)*sin(b*x+a)/b/(1-n)/(sin(b*x+a)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{c \sin(a + bx)(c \sec(a + bx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^n,x]

[Out] -((c*Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*(c*Sec[a + b*x])^(n - 1)*Sin[a + b*x])/(b*(1 - n)*Sqrt[Sin[a + b*x]^2]))

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int (c \sec(a + bx))^n dx = \left(\frac{\cos(a + bx)}{c}\right)^n (c \sec(a + bx))^n \int \left(\frac{\cos(a + bx)}{c}\right)^{-n} dx$$

$$= -\frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) (c \sec(a + bx))^n \sin(a + bx)}{b(1-n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A] time = 0.05, size = 61, normalized size = 0.84

$$\frac{\sqrt{-\tan^2(a + bx)} \cot(a + bx) (c \sec(a + bx))^n {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{n+2}{2}; \sec^2(a + bx)\right)}{bn}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^n,x]

[Out] (Cot[a + b*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[a + b*x]^2]*(c*Sec[a + b*x])^n*sqrt[-Tan[a + b*x]^2])/(b*n)

fricas [F] time = 0.82, size = 0, normalized size = 0.00

$$\text{integral}((c \sec(bx + a))^n, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^n,x, algorithm="fricas")

[Out] integral((c*sec(b*x + a))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^n,x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^n, x)

maple [F] time = 1.32, size = 0, normalized size = 0.00

$$\int (c \sec(bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^n,x)`

[Out] `int((c*sec(b*x+a))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec (bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^n,x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^n, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos (a + bx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^n,x)`

[Out] `int((c/cos(a + b*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec (a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a)**n,x)`

[Out] `Integral((c*sec(a + b*x)**n, x)`

3.39 $\int \sec^2(x)^{7/2} dx$

Optimal. Leaf size=50

$$\frac{1}{6} \tan(x) \sec^2(x)^{5/2} + \frac{5}{24} \tan(x) \sec^2(x)^{3/2} + \frac{5}{16} \tan(x) \sqrt{\sec^2(x)} + \frac{5}{16} \sinh^{-1}(\tan(x))$$

[Out] 5/16*arcsinh(tan(x))+5/24*(sec(x)^2)^(3/2)*tan(x)+1/6*(sec(x)^2)^(5/2)*tan(x)+5/16*(sec(x)^2)^(1/2)*tan(x)

Rubi [A] time = 0.02, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4122, 195, 215}

$$\frac{1}{6} \tan(x) \sec^2(x)^{5/2} + \frac{5}{24} \tan(x) \sec^2(x)^{3/2} + \frac{5}{16} \tan(x) \sqrt{\sec^2(x)} + \frac{5}{16} \sinh^{-1}(\tan(x))$$

Antiderivative was successfully verified.

[In] Int[(Sec[x]^2)^(7/2), x]

[Out] (5*ArcSinh[Tan[x]])/16 + (5*Sqrt[Sec[x]^2]*Tan[x])/16 + (5*(Sec[x]^2)^(3/2)*Tan[x])/24 + ((Sec[x]^2)^(5/2)*Tan[x])/6

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 215

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSinh[(Rt[b, 2]*x)/Sqrt[a]]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \sec^2(x)^{7/2} dx &= \text{Subst} \left(\int (1+x^2)^{5/2} dx, x, \tan(x) \right) \\
&= \frac{1}{6} \sec^2(x)^{5/2} \tan(x) + \frac{5}{6} \text{Subst} \left(\int (1+x^2)^{3/2} dx, x, \tan(x) \right) \\
&= \frac{5}{24} \sec^2(x)^{3/2} \tan(x) + \frac{1}{6} \sec^2(x)^{5/2} \tan(x) + \frac{5}{8} \text{Subst} \left(\int \sqrt{1+x^2} dx, x, \tan(x) \right) \\
&= \frac{5}{16} \sqrt{\sec^2(x)} \tan(x) + \frac{5}{24} \sec^2(x)^{3/2} \tan(x) + \frac{1}{6} \sec^2(x)^{5/2} \tan(x) + \frac{5}{16} \text{Subst} \left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \tan(x) \right) \\
&= \frac{5}{16} \sinh^{-1}(\tan(x)) + \frac{5}{16} \sqrt{\sec^2(x)} \tan(x) + \frac{5}{24} \sec^2(x)^{3/2} \tan(x) + \frac{1}{6} \sec^2(x)^{5/2} \tan(x)
\end{aligned}$$

Mathematica [A] time = 0.30, size = 74, normalized size = 1.48

$$\frac{1}{96} \cos(x) \sqrt{\sec^2(x)} \left(\frac{1}{8} (198 \sin(x) + 85 \sin(3x) + 15 \sin(5x)) \sec^6(x) - 30 \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + 30 \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(Sec[x]^2)^(7/2), x]

[Out] (Cos[x]*Sqrt[Sec[x]^2]*(-30*Log[Cos[x/2] - Sin[x/2]] + 30*Log[Cos[x/2] + Sin[x/2]] + (Sec[x]^6*(198*Sin[x] + 85*Sin[3*x] + 15*Sin[5*x]))/8))/96

fricas [A] time = 0.44, size = 49, normalized size = 0.98

$$\frac{15 \cos(x)^6 \log(\sin(x) + 1) - 15 \cos(x)^6 \log(-\sin(x) + 1) + 2(15 \cos(x)^4 + 10 \cos(x)^2 + 8) \sin(x)}{96 \cos(x)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(7/2), x, algorithm="fricas")

[Out] -1/96*(15*cos(x)^6*log(sin(x) + 1) - 15*cos(x)^6*log(-sin(x) + 1) + 2*(15*cos(x)^4 + 10*cos(x)^2 + 8)*sin(x))/cos(x)^6

giac [A] time = 0.99, size = 59, normalized size = 1.18

$$\frac{5 \log(\sin(x) + 1)}{32 \operatorname{sgn}(\cos(x))} - \frac{5 \log(-\sin(x) + 1)}{32 \operatorname{sgn}(\cos(x))} - \frac{15 \sin(x)^5 - 40 \sin(x)^3 + 33 \sin(x)}{48 (\sin(x)^2 - 1)^3 \operatorname{sgn}(\cos(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(7/2),x, algorithm="giac")

[Out] $5/32 \cdot \log(\sin(x) + 1)/\operatorname{sgn}(\cos(x)) - 5/32 \cdot \log(-\sin(x) + 1)/\operatorname{sgn}(\cos(x)) - 1/48 \cdot (15 \cdot \sin(x)^5 - 40 \cdot \sin(x)^3 + 33 \cdot \sin(x))/((\sin(x)^2 - 1)^3 \cdot \operatorname{sgn}(\cos(x)))$

maple [A] time = 0.44, size = 72, normalized size = 1.44

$$\frac{15 \left(\cos^6(x) \right) \ln \left(-\frac{-1 + \cos(x) + \sin(x)}{\sin(x)} \right) - 15 \left(\cos^6(x) \right) \ln \left(-\frac{-\sin(x) - 1 + \cos(x)}{\sin(x)} \right) - 15 \left(\cos^4(x) \right) \sin(x) - 10 \left(\cos^2(x) \right) \sin(x)}{6 \left(\cos(2x) + 1 \right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((sec(x)^2)^(7/2),x)

[Out] $-1/48 \cdot (15 \cdot \cos(x)^6 \cdot \ln(-(-1 + \cos(x) + \sin(x))/\sin(x)) - 15 \cdot \cos(x)^6 \cdot \ln(-(-\sin(x) - 1 + \cos(x))/\sin(x)) - 15 \cdot \cos(x)^4 \cdot \sin(x) - 10 \cdot \cos(x)^2 \cdot \sin(x) - 8 \cdot \sin(x)) \cdot \cos(x) \cdot (1/\cos(x)^2)^{7/2}$

maxima [A] time = 0.89, size = 42, normalized size = 0.84

$$\frac{1}{6} \left(\tan(x)^2 + 1 \right)^{\frac{5}{2}} \tan(x) + \frac{5}{24} \left(\tan(x)^2 + 1 \right)^{\frac{3}{2}} \tan(x) + \frac{5}{16} \sqrt{\tan(x)^2 + 1} \tan(x) + \frac{5}{16} \operatorname{arsinh}(\tan(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(7/2),x, algorithm="maxima")

[Out] $1/6 \cdot (\tan(x)^2 + 1)^{5/2} \cdot \tan(x) + 5/24 \cdot (\tan(x)^2 + 1)^{3/2} \cdot \tan(x) + 5/16 \cdot \operatorname{sqrt}(\tan(x)^2 + 1) \cdot \tan(x) + 5/16 \cdot \operatorname{arcsinh}(\tan(x))$

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{1}{\cos(x)^2} \right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(x)^2)^(7/2),x)

[Out] int((1/cos(x)^2)^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)**2)**(7/2),x)

[Out] Timed out

3.40 $\int \sec^2(x)^{5/2} dx$

Optimal. Leaf size=36

$$\frac{1}{4} \tan(x) \sec^2(x)^{3/2} + \frac{3}{8} \tan(x) \sqrt{\sec^2(x)} + \frac{3}{8} \sinh^{-1}(\tan(x))$$

[Out] 3/8*arcsinh(tan(x))+1/4*(sec(x)^2)^(3/2)*tan(x)+3/8*(sec(x)^2)^(1/2)*tan(x)

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4122, 195, 215}

$$\frac{1}{4} \tan(x) \sec^2(x)^{3/2} + \frac{3}{8} \tan(x) \sqrt{\sec^2(x)} + \frac{3}{8} \sinh^{-1}(\tan(x))$$

Antiderivative was successfully verified.

[In] Int[(Sec[x]^2)^(5/2), x]

[Out] (3*ArcSinh[Tan[x]])/8 + (3*Sqrt[Sec[x]^2]*Tan[x])/8 + ((Sec[x]^2)^(3/2)*Tan[x])/4

Rule 195

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])
```

Rule 215

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[(Rt[b, 2]*x)/Sqrt[a]]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 4122

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)^2])^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\int \sec^2(x)^{5/2} dx &= \text{Subst} \left(\int (1+x^2)^{3/2} dx, x, \tan(x) \right) \\
&= \frac{1}{4} \sec^2(x)^{3/2} \tan(x) + \frac{3}{4} \text{Subst} \left(\int \sqrt{1+x^2} dx, x, \tan(x) \right) \\
&= \frac{3}{8} \sqrt{\sec^2(x)} \tan(x) + \frac{1}{4} \sec^2(x)^{3/2} \tan(x) + \frac{3}{8} \text{Subst} \left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \tan(x) \right) \\
&= \frac{3}{8} \sinh^{-1}(\tan(x)) + \frac{3}{8} \sqrt{\sec^2(x)} \tan(x) + \frac{1}{4} \sec^2(x)^{3/2} \tan(x)
\end{aligned}$$

Mathematica [A] time = 0.16, size = 68, normalized size = 1.89

$$\frac{1}{16} \cos(x) \sqrt{\sec^2(x)} \left(\frac{1}{2} (11 \sin(x) + 3 \sin(3x)) \sec^4(x) - 6 \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + 6 \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(Sec[x]^2)^(5/2), x]

[Out] (Cos[x]*Sqrt[Sec[x]^2]*(-6*Log[Cos[x/2] - Sin[x/2]] + 6*Log[Cos[x/2] + Sin[x/2]] + (Sec[x]^4*(11*Sin[x] + 3*Sin[3*x]))/2))/16

fricas [A] time = 0.71, size = 43, normalized size = 1.19

$$\frac{3 \cos(x)^4 \log(\sin(x) + 1) - 3 \cos(x)^4 \log(-\sin(x) + 1) + 2(3 \cos(x)^2 + 2) \sin(x)}{16 \cos(x)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(5/2), x, algorithm="fricas")

[Out] -1/16*(3*cos(x)^4*log(sin(x) + 1) - 3*cos(x)^4*log(-sin(x) + 1) + 2*(3*cos(x)^2 + 2)*sin(x))/cos(x)^4

giac [B] time = 0.43, size = 53, normalized size = 1.47

$$\frac{3 \log(\sin(x) + 1)}{16 \operatorname{sgn}(\cos(x))} - \frac{3 \log(-\sin(x) + 1)}{16 \operatorname{sgn}(\cos(x))} - \frac{3 \sin(x)^3 - 5 \sin(x)}{8 (\sin(x)^2 - 1)^2 \operatorname{sgn}(\cos(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(5/2), x, algorithm="giac")

[Out] $\frac{3}{16} \log(\sin(x) + 1) / \operatorname{sgn}(\cos(x)) - \frac{3}{16} \log(-\sin(x) + 1) / \operatorname{sgn}(\cos(x)) - \frac{1}{8} \frac{(3 \sin(x)^3 - 5 \sin(x))}{(\sin(x)^2 - 1)^2 \operatorname{sgn}(\cos(x))}$

maple [B] time = 0.35, size = 64, normalized size = 1.78

$$\frac{\left(3 \cos^4(x) \ln\left(-\frac{-1 + \cos(x) + \sin(x)}{\sin(x)}\right) - 3 \cos^4(x) \ln\left(-\frac{-\sin(x) - 1 + \cos(x)}{\sin(x)}\right) - 3 \left(\cos^2(x) \sin(x) - 2 \sin(x)\right) \cos(x) \sqrt{2}\right)}{2 (\cos(2x) + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((sec(x)^2)^(5/2), x)`

[Out] $-1/8 * (3 * \cos(x)^4 * \ln(-(-1 + \cos(x) + \sin(x)) / \sin(x)) - 3 * \cos(x)^4 * \ln(-(-\sin(x) - 1 + \cos(x)) / \sin(x)) - 3 * \cos(x)^2 * \sin(x) - 2 * \sin(x)) * \cos(x) * (1 / \cos(x)^2)^(5/2)$

maxima [A] time = 0.59, size = 30, normalized size = 0.83

$$\frac{1}{4} (\tan(x)^2 + 1)^{\frac{3}{2}} \tan(x) + \frac{3}{8} \sqrt{\tan(x)^2 + 1} \tan(x) + \frac{3}{8} \operatorname{arsinh}(\tan(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((sec(x)^2)^(5/2), x, algorithm="maxima")`

[Out] $1/4 * (\tan(x)^2 + 1)^{(3/2)} * \tan(x) + 3/8 * \operatorname{sqrt}(\tan(x)^2 + 1) * \tan(x) + 3/8 * \operatorname{arcsinh}(\tan(x))$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \left(\frac{1}{\cos(x)^2}\right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(x)^2)^(5/2), x)`

[Out] `int((1/cos(x)^2)^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (\sec^2(x))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((sec(x)**2)**(5/2), x)`

[Out] `Integral((sec(x)**2)**(5/2), x)`

3.41 $\int \sec^2(x)^{3/2} dx$

Optimal. Leaf size=22

$$\frac{1}{2} \tan(x) \sqrt{\sec^2(x)} + \frac{1}{2} \sinh^{-1}(\tan(x))$$

[Out] 1/2*arcsinh(tan(x))+1/2*(sec(x)^2)^(1/2)*tan(x)

Rubi [A] time = 0.01, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4122, 195, 215}

$$\frac{1}{2} \tan(x) \sqrt{\sec^2(x)} + \frac{1}{2} \sinh^{-1}(\tan(x))$$

Antiderivative was successfully verified.

[In] Int[(Sec[x]^2)^(3/2), x]

[Out] ArcSinh[Tan[x]]/2 + (Sqrt[Sec[x]^2]*Tan[x])/2

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 215

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSinh[(Rt[b, 2]*x)/Sqrt[a]]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)^2])^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \sec^2(x)^{3/2} dx &= \text{Subst} \left(\int \sqrt{1+x^2} dx, x, \tan(x) \right) \\ &= \frac{1}{2} \sqrt{\sec^2(x)} \tan(x) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \tan(x) \right) \\ &= \frac{1}{2} \sinh^{-1}(\tan(x)) + \frac{1}{2} \sqrt{\sec^2(x)} \tan(x) \end{aligned}$$

Mathematica [B] time = 0.06, size = 52, normalized size = 2.36

$$\frac{1}{2} \cos(x) \sqrt{\sec^2(x)} \left(\tan(x) \sec(x) - \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(Sec[x]^2)^(3/2), x]

[Out] (Cos[x]*Sqrt[Sec[x]^2]*(-Log[Cos[x/2] - Sin[x/2]] + Log[Cos[x/2] + Sin[x/2]] + Sec[x]*Tan[x]))/2

fricas [B] time = 0.80, size = 34, normalized size = 1.55

$$\frac{-\cos(x)^2 \log(\sin(x) + 1) - \cos(x)^2 \log(-\sin(x) + 1) + 2 \sin(x)}{4 \cos(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(3/2), x, algorithm="fricas")

[Out] -1/4*(cos(x)^2*log(sin(x) + 1) - cos(x)^2*log(-sin(x) + 1) + 2*sin(x))/cos(x)^2

giac [B] time = 0.34, size = 44, normalized size = 2.00

$$\frac{\log(\sin(x) + 1)}{4 \operatorname{sgn}(\cos(x))} - \frac{\log(-\sin(x) + 1)}{4 \operatorname{sgn}(\cos(x))} - \frac{\sin(x)}{2(\sin(x)^2 - 1) \operatorname{sgn}(\cos(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(3/2), x, algorithm="giac")

[Out] 1/4*log(sin(x) + 1)/sgn(cos(x)) - 1/4*log(-sin(x) + 1)/sgn(cos(x)) - 1/2*sin(x)/((sin(x)^2 - 1)*sgn(cos(x)))

maple [B] time = 0.29, size = 55, normalized size = 2.50

$$\frac{\left(\cos^2(x)\right) \ln\left(\frac{-1+\cos(x)+\sin(x)}{\sin(x)}\right) - \left(\cos^2(x)\right) \ln\left(\frac{-\sin(x)-1+\cos(x)}{\sin(x)}\right) - \sin(x) \cos(x) \sqrt{2} \sqrt{\frac{1}{\cos(2x)+1}}}{\cos(2x) + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((sec(x)^2)^(3/2), x)

[Out] -1/2*(cos(x)^2*ln(-(-1+cos(x)+sin(x))/sin(x))-cos(x)^2*ln(-(-sin(x)-1+cos(x))/sin(x))-sin(x))*cos(x)*(1/cos(x)^2)^(3/2)

maxima [A] time = 0.76, size = 18, normalized size = 0.82

$$\frac{1}{2} \sqrt{\tan(x)^2 + 1} \tan(x) + \frac{1}{2} \operatorname{arsinh}(\tan(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(3/2), x, algorithm="maxima")

[Out] 1/2*sqrt(tan(x)^2 + 1)*tan(x) + 1/2*arcsinh(tan(x))

mupad [F] time = 0.00, size = -1, normalized size = -0.05

$$\int \left(\frac{1}{\cos(x)^2}\right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(x)^2)^(3/2), x)

[Out] int((1/cos(x)^2)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (\sec^2(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)**2)**(3/2), x)

[Out] Integral((sec(x)**2)**(3/2), x)

3.42 $\int \sqrt{\sec^2(x)} dx$

Optimal. Leaf size=3

$$\sinh^{-1}(\tan(x))$$

[Out] arcsinh(tan(x))

Rubi [A] time = 0.01, antiderivative size = 3, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {4122, 215}

$$\sinh^{-1}(\tan(x))$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[x]^2], x]

[Out] ArcSinh[Tan[x]]

Rule 215

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[(Rt[b, 2]*x)/Sqrt[a]]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 4122

Int[((b_)*sec[(e_) + (f_)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \sqrt{\sec^2(x)} dx &= \text{Subst} \left(\int \frac{1}{\sqrt{1+x^2}} dx, x, \tan(x) \right) \\ &= \sinh^{-1}(\tan(x)) \end{aligned}$$

Mathematica [B] time = 0.01, size = 44, normalized size = 14.67

$$\cos(x)\sqrt{\sec^2(x)} \left(\log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) - \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[x]^2], x]

[Out] Cos[x]*(-Log[Cos[x/2] - Sin[x/2]] + Log[Cos[x/2] + Sin[x/2]])*Sqrt[Sec[x]^2]

fricas [B] time = 0.49, size = 17, normalized size = 5.67

$$-\frac{1}{2} \log(\sin(x) + 1) + \frac{1}{2} \log(-\sin(x) + 1)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(1/2),x, algorithm="fricas")

[Out] -1/2*log(sin(x) + 1) + 1/2*log(-sin(x) + 1)

giac [B] time = 0.73, size = 35, normalized size = 11.67

$$\frac{\log\left(\left|\frac{1}{\sin(x)} + \sin(x) + 2\right|\right)}{4 \operatorname{sgn}(\cos(x))} - \frac{\log\left(\left|\frac{1}{\sin(x)} + \sin(x) - 2\right|\right)}{4 \operatorname{sgn}(\cos(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/4*log(abs(1/sin(x) + sin(x) + 2))/sgn(cos(x)) - 1/4*log(abs(1/sin(x) + sin(x) - 2))/sgn(cos(x))

maple [B] time = 0.37, size = 21, normalized size = 7.00

$$-2 \cos(x) \sqrt{2} \sqrt{\frac{1}{\cos(2x) + 1}} \operatorname{arctanh}\left(\frac{-1 + \cos(x)}{\sin(x)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((sec(x)^2)^(1/2),x)

[Out] -2*cos(x)*(1/cos(x)^2)^(1/2)*arctanh((-1+cos(x))/sin(x))

maxima [A] time = 0.46, size = 3, normalized size = 1.00

$$\operatorname{arsinh}(\tan(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((sec(x)^2)^(1/2),x, algorithm="maxima")

[Out] arcsinh(tan(x))

mupad [F] time = 0.00, size = -1, normalized size = -0.33

$$\int \sqrt{\frac{1}{\cos(x)^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(x)^2)^(1/2), x)`

[Out] `int((1/cos(x)^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\sec^2(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((sec(x)**2)**(1/2), x)`

[Out] `Integral(sqrt(sec(x)**2), x)`

$$3.43 \quad \int \frac{1}{\sqrt{\sec^2(x)}} dx$$

Optimal. Leaf size=11

$$\frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

[Out] $\tan(x)/(\sec(x)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {4122, 191}

$$\frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

Antiderivative was successfully verified.

[In] `Int[1/Sqrt[Sec[x]^2], x]`

[Out] `Tan[x]/Sqrt[Sec[x]^2]`

Rule 191

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]`

Rule 4122

`Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]`

Rubi steps

$$\int \frac{1}{\sqrt{\sec^2(x)}} dx = \text{Subst} \left(\int \frac{1}{(1+x^2)^{3/2}} dx, x, \tan(x) \right) \\ = \frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

Mathematica [A] time = 0.01, size = 11, normalized size = 1.00

$$\frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[Sec[x]^2],x]

[Out] Tan[x]/Sqrt[Sec[x]^2]

fricas [A] time = 0.76, size = 4, normalized size = 0.36

$$-\sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(1/2),x, algorithm="fricas")

[Out] -sin(x)

giac [A] time = 0.31, size = 6, normalized size = 0.55

$$\operatorname{sgn}(\cos(x)) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(1/2),x, algorithm="giac")

[Out] $\operatorname{sgn}(\cos(x)) \sin(x)$

maple [A] time = 0.40, size = 14, normalized size = 1.27

$$\frac{\sin(x)\sqrt{2}}{2\sqrt{\frac{1}{\cos(2x)+1}} \cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sec(x)^2)^(1/2),x)

[Out] $\sin(x)/(1/\cos(x)^2)^{(1/2)}/\cos(x)$

maxima [A] time = 0.52, size = 11, normalized size = 1.00

$$\frac{\tan(x)}{\sqrt{\tan(x)^2 + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(1/2),x, algorithm="maxima")

[Out] $\tan(x)/\sqrt{\tan(x)^2 + 1}$

mupad [B] time = 0.16, size = 12, normalized size = 1.09

$$\frac{\sqrt{2} \sin(2x)}{2\sqrt{2} \cos(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1/cos(x)^2)^(1/2),x)`

[Out] `(2^(1/2)*sin(2*x))/(2*(2*cos(x)^2)^(1/2))`

sympy [A] time = 0.40, size = 10, normalized size = 0.91

$$\frac{\tan(x)}{\sqrt{\sec^2(x)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(sec(x)**2)**(1/2),x)`

[Out] `tan(x)/sqrt(sec(x)**2)`

$$3.44 \quad \int \frac{1}{\sec^2(x)^{3/2}} dx$$

Optimal. Leaf size=29

$$\frac{2 \tan(x)}{3\sqrt{\sec^2(x)}} + \frac{\tan(x)}{3 \sec^2(x)^{3/2}}$$

[Out] 1/3*tan(x)/(sec(x)^2)^(3/2)+2/3*tan(x)/(sec(x)^2)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4122, 192, 191}

$$\frac{2 \tan(x)}{3\sqrt{\sec^2(x)}} + \frac{\tan(x)}{3 \sec^2(x)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(Sec[x]^2)^(-3/2), x]

[Out] Tan[x]/(3*(Sec[x]^2)^(3/2)) + (2*Tan[x])/(3*Sqrt[Sec[x]^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^2(x)^{3/2}} dx &= \text{Subst} \left(\int \frac{1}{(1+x^2)^{5/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{3 \sec^2(x)^{3/2}} + \frac{2}{3} \text{Subst} \left(\int \frac{1}{(1+x^2)^{3/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{3 \sec^2(x)^{3/2}} + \frac{2 \tan(x)}{3 \sqrt{\sec^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 23, normalized size = 0.79

$$\frac{(9 \sin(x) + \sin(3x)) \sec(x)}{12 \sqrt{\sec^2(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[(Sec[x]^2)^(-3/2), x]

[Out] (Sec[x]*(9*Sin[x] + Sin[3*x]))/(12*Sqrt[Sec[x]^2])

fricas [A] time = 0.63, size = 10, normalized size = 0.34

$$-\frac{1}{3} (\cos(x)^2 + 2) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(3/2), x, algorithm="fricas")

[Out] -1/3*(cos(x)^2 + 2)*sin(x)

giac [A] time = 0.50, size = 16, normalized size = 0.55

$$-\frac{1}{3} \text{sgn}(\cos(x)) \sin(x)^3 + \text{sgn}(\cos(x)) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(3/2), x, algorithm="giac")

[Out] -1/3*sgn(cos(x))*sin(x)^3 + sgn(cos(x))*sin(x)

maple [A] time = 0.34, size = 21, normalized size = 0.72

$$\frac{\sin(x) (\cos^2(x) + 2) (\cos(2x) + 1) \sqrt{2}}{12 \cos(x)^3 \sqrt{\frac{1}{\cos(2x)+1}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sec(x)^2)^(3/2), x)`

[Out] `1/3*sin(x)*(cos(x)^2+2)/cos(x)^3/(1/cos(x)^2)^(3/2)`

maxima [A] time = 0.48, size = 25, normalized size = 0.86

$$\frac{2 \tan(x)}{3 \sqrt{\tan(x)^2 + 1}} + \frac{\tan(x)}{3 (\tan(x)^2 + 1)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(sec(x)^2)^(3/2), x, algorithm="maxima")`

[Out] `2/3*tan(x)/sqrt(tan(x)^2 + 1) + 1/3*tan(x)/(tan(x)^2 + 1)^(3/2)`

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\left(\frac{1}{\cos(x)^2}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1/cos(x)^2)^(3/2), x)`

[Out] `int(1/(1/cos(x)^2)^(3/2), x)`

sympy [A] time = 1.05, size = 27, normalized size = 0.93

$$\frac{2 \tan^3(x)}{3 (\sec^2(x))^{\frac{3}{2}}} + \frac{\tan(x)}{(\sec^2(x))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(sec(x)**2)**(3/2), x)`

[Out] `2*tan(x)**3/(3*(sec(x)**2)**(3/2)) + tan(x)/(sec(x)**2)**(3/2)`

$$3.45 \quad \int \frac{1}{\sec^2(x)^{5/2}} dx$$

Optimal. Leaf size=43

$$\frac{8 \tan(x)}{15\sqrt{\sec^2(x)}} + \frac{4 \tan(x)}{15 \sec^2(x)^{3/2}} + \frac{\tan(x)}{5 \sec^2(x)^{5/2}}$$

[Out] $1/5*\tan(x)/(\sec(x)^2)^{(5/2)}+4/15*\tan(x)/(\sec(x)^2)^{(3/2)}+8/15*\tan(x)/(\sec(x)^2)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4122, 192, 191}

$$\frac{8 \tan(x)}{15\sqrt{\sec^2(x)}} + \frac{4 \tan(x)}{15 \sec^2(x)^{3/2}} + \frac{\tan(x)}{5 \sec^2(x)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(Sec[x]^2)^(-5/2), x]

[Out] Tan[x]/(5*(Sec[x]^2)^(5/2)) + (4*Tan[x])/(15*(Sec[x]^2)^(3/2)) + (8*Tan[x])/(15*Sqrt[Sec[x]^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^2(x)^{5/2}} dx &= \text{Subst} \left(\int \frac{1}{(1+x^2)^{7/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{5 \sec^2(x)^{5/2}} + \frac{4}{5} \text{Subst} \left(\int \frac{1}{(1+x^2)^{5/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{5 \sec^2(x)^{5/2}} + \frac{4 \tan(x)}{15 \sec^2(x)^{3/2}} + \frac{8}{15} \text{Subst} \left(\int \frac{1}{(1+x^2)^{3/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{5 \sec^2(x)^{5/2}} + \frac{4 \tan(x)}{15 \sec^2(x)^{3/2}} + \frac{8 \tan(x)}{15 \sqrt{\sec^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 31, normalized size = 0.72

$$\frac{(150 \sin(x) + 25 \sin(3x) + 3 \sin(5x)) \sec(x)}{240 \sqrt{\sec^2(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[(Sec[x]^2)^(-5/2), x]

[Out] (Sec[x]*(150*Sin[x] + 25*Sin[3*x] + 3*Sin[5*x]))/(240*Sqrt[Sec[x]^2])

fricas [A] time = 0.63, size = 18, normalized size = 0.42

$$-\frac{1}{15} (3 \cos(x)^4 + 4 \cos(x)^2 + 8) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(5/2), x, algorithm="fricas")

[Out] -1/15*(3*cos(x)^4 + 4*cos(x)^2 + 8)*sin(x)

giac [A] time = 0.43, size = 25, normalized size = 0.58

$$\frac{1}{5} \operatorname{sgn}(\cos(x)) \sin(x)^5 - \frac{2}{3} \operatorname{sgn}(\cos(x)) \sin(x)^3 + \operatorname{sgn}(\cos(x)) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(5/2), x, algorithm="giac")

[Out] $1/5*\text{sgn}(\cos(x))*\sin(x)^5 - 2/3*\text{sgn}(\cos(x))*\sin(x)^3 + \text{sgn}(\cos(x))*\sin(x)$

maple [A] time = 0.32, size = 29, normalized size = 0.67

$$\frac{\sin(x) \left(3 \left(\cos^4(x) \right) + 4 \left(\cos^2(x) \right) + 8 \right) (\cos(2x) + 1)^2 \sqrt{2}}{120 \cos(x)^5 \sqrt{\frac{1}{\cos(2x)+1}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sec(x)^2)^(5/2), x)`

[Out] $1/15*\sin(x)*(3*\cos(x)^4+4*\cos(x)^2+8)/\cos(x)^5/(1/\cos(x)^2)^(5/2)$

maxima [A] time = 0.42, size = 37, normalized size = 0.86

$$\frac{8 \tan(x)}{15 \sqrt{\tan(x)^2 + 1}} + \frac{4 \tan(x)}{15 (\tan(x)^2 + 1)^{\frac{3}{2}}} + \frac{\tan(x)}{5 (\tan(x)^2 + 1)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(sec(x)^2)^(5/2), x, algorithm="maxima")`

[Out] $8/15*\tan(x)/\text{sqrt}(\tan(x)^2 + 1) + 4/15*\tan(x)/(\tan(x)^2 + 1)^(3/2) + 1/5*\tan(x)/(\tan(x)^2 + 1)^(5/2)$

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(x)^2} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1/cos(x)^2)^(5/2), x)`

[Out] `int(1/(1/cos(x)^2)^(5/2), x)`

sympy [A] time = 10.68, size = 44, normalized size = 1.02

$$\frac{8 \tan^5(x)}{15 (\sec^2(x))^{\frac{5}{2}}} + \frac{4 \tan^3(x)}{3 (\sec^2(x))^{\frac{5}{2}}} + \frac{\tan(x)}{(\sec^2(x))^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(sec(x)**2)**(5/2), x)`

[Out] $8*\tan(x)**5/(15*(\sec(x)**2)**(5/2)) + 4*\tan(x)**3/(3*(\sec(x)**2)**(5/2)) + \tan(x)/(\sec(x)**2)**(5/2)$

$$3.46 \quad \int \frac{1}{\sec^2(x)^{7/2}} dx$$

Optimal. Leaf size=57

$$\frac{16 \tan(x)}{35\sqrt{\sec^2(x)}} + \frac{8 \tan(x)}{35 \sec^2(x)^{3/2}} + \frac{6 \tan(x)}{35 \sec^2(x)^{5/2}} + \frac{\tan(x)}{7 \sec^2(x)^{7/2}}$$

[Out] $1/7*\tan(x)/(\sec(x)^2)^{(7/2)}+6/35*\tan(x)/(\sec(x)^2)^{(5/2)}+8/35*\tan(x)/(\sec(x)^2)^{(3/2)}+16/35*\tan(x)/(\sec(x)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4122, 192, 191}

$$\frac{16 \tan(x)}{35\sqrt{\sec^2(x)}} + \frac{8 \tan(x)}{35 \sec^2(x)^{3/2}} + \frac{6 \tan(x)}{35 \sec^2(x)^{5/2}} + \frac{\tan(x)}{7 \sec^2(x)^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[(Sec[x]^2)^(-7/2), x]

[Out] Tan[x]/(7*(Sec[x]^2)^(7/2)) + (6*Tan[x])/(35*(Sec[x]^2)^(5/2)) + (8*Tan[x])/(35*(Sec[x]^2)^(3/2)) + (16*Tan[x])/(35*Sqrt[Sec[x]^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^2(x)^{7/2}} dx &= \text{Subst} \left(\int \frac{1}{(1+x^2)^{9/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{7 \sec^2(x)^{7/2}} + \frac{6}{7} \text{Subst} \left(\int \frac{1}{(1+x^2)^{7/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{7 \sec^2(x)^{7/2}} + \frac{6 \tan(x)}{35 \sec^2(x)^{5/2}} + \frac{24}{35} \text{Subst} \left(\int \frac{1}{(1+x^2)^{5/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{7 \sec^2(x)^{7/2}} + \frac{6 \tan(x)}{35 \sec^2(x)^{5/2}} + \frac{8 \tan(x)}{35 \sec^2(x)^{3/2}} + \frac{16}{35} \text{Subst} \left(\int \frac{1}{(1+x^2)^{3/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{7 \sec^2(x)^{7/2}} + \frac{6 \tan(x)}{35 \sec^2(x)^{5/2}} + \frac{8 \tan(x)}{35 \sec^2(x)^{3/2}} + \frac{16 \tan(x)}{35 \sqrt{\sec^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 37, normalized size = 0.65

$$\frac{(1225 \sin(x) + 245 \sin(3x) + 49 \sin(5x) + 5 \sin(7x)) \sec(x)}{2240 \sqrt{\sec^2(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[(Sec[x]^2)^(-7/2), x]

[Out] (Sec[x]*(1225*Sin[x] + 245*Sin[3*x] + 49*Sin[5*x] + 5*Sin[7*x]))/(2240*Sqrt[Sec[x]^2])

fricas [A] time = 0.54, size = 24, normalized size = 0.42

$$-\frac{1}{35} (5 \cos(x)^6 + 6 \cos(x)^4 + 8 \cos(x)^2 + 16) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(7/2), x, algorithm="fricas")

[Out] -1/35*(5*cos(x)^6 + 6*cos(x)^4 + 8*cos(x)^2 + 16)*sin(x)

giac [A] time = 0.43, size = 34, normalized size = 0.60

$$-\frac{1}{7} \operatorname{sgn}(\cos(x)) \sin(x)^7 + \frac{3}{5} \operatorname{sgn}(\cos(x)) \sin(x)^5 - \operatorname{sgn}(\cos(x)) \sin(x)^3 + \operatorname{sgn}(\cos(x)) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(7/2),x, algorithm="giac")

[Out] $-1/7*\text{sgn}(\cos(x))*\sin(x)^7 + 3/5*\text{sgn}(\cos(x))*\sin(x)^5 - \text{sgn}(\cos(x))*\sin(x)^3 + \text{sgn}(\cos(x))*\sin(x)$

maple [A] time = 0.36, size = 35, normalized size = 0.61

$$\frac{\sin(x) \left(5 \left(\cos^6(x) \right) + 6 \left(\cos^4(x) \right) + 8 \left(\cos^2(x) \right) + 16 \right) (\cos(2x) + 1)^3 \sqrt{2}}{560 \cos(x)^7 \sqrt{\frac{1}{\cos(2x)+1}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sec(x)^2)^(7/2),x)

[Out] $1/35*\sin(x)*(5*\cos(x)^6+6*\cos(x)^4+8*\cos(x)^2+16)/\cos(x)^7/(1/\cos(x)^2)^(7/2)$

maxima [A] time = 0.34, size = 49, normalized size = 0.86

$$\frac{16 \tan(x)}{35 \sqrt{\tan(x)^2 + 1}} + \frac{8 \tan(x)}{35 (\tan(x)^2 + 1)^{\frac{3}{2}}} + \frac{6 \tan(x)}{35 (\tan(x)^2 + 1)^{\frac{5}{2}}} + \frac{\tan(x)}{7 (\tan(x)^2 + 1)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(sec(x)^2)^(7/2),x, algorithm="maxima")

[Out] $16/35*\tan(x)/\text{sqrt}(\tan(x)^2 + 1) + 8/35*\tan(x)/(\tan(x)^2 + 1)^(3/2) + 6/35*\tan(x)/(\tan(x)^2 + 1)^(5/2) + 1/7*\tan(x)/(\tan(x)^2 + 1)^(7/2)$

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{1}{\cos(x)^2}\right)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1/cos(x)^2)^(7/2),x)

[Out] int(1/(1/cos(x)^2)^(7/2), x)

sympy [A] time = 147.49, size = 60, normalized size = 1.05

$$\frac{16 \tan^7(x)}{35 (\sec^2(x))^{\frac{7}{2}}} + \frac{8 \tan^5(x)}{5 (\sec^2(x))^{\frac{7}{2}}} + \frac{2 \tan^3(x)}{(\sec^2(x))^{\frac{7}{2}}} + \frac{\tan(x)}{(\sec^2(x))^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(sec(x)**2)**(7/2),x)
```

```
[Out] 16*tan(x)**7/(35*(sec(x)**2)**(7/2)) + 8*tan(x)**5/(5*(sec(x)**2)**(7/2)) +  
2*tan(x)**3/(sec(x)**2)**(7/2) + tan(x)/(sec(x)**2)**(7/2)
```

3.47 $\int (a \sec^2(x))^{7/2} dx$

Optimal. Leaf size=84

$$\frac{5}{16}a^{7/2} \tanh^{-1}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{5}{16}a^3 \tan(x) \sqrt{a \sec^2(x)} + \frac{5}{24}a^2 \tan(x) (a \sec^2(x))^{3/2} + \frac{1}{6}a \tan(x) (a \sec^2(x))^{5/2}$$

[Out] $5/16*a^{(7/2)}*\operatorname{arctanh}(a^{(1/2)}*\tan(x)/(a*\sec(x)^2)^{(1/2)})+5/24*a^2*(a*\sec(x)^2)^{(3/2)}*\tan(x)+1/6*a*(a*\sec(x)^2)^{(5/2)}*\tan(x)+5/16*a^3*(a*\sec(x)^2)^{(1/2)}*\tan(x)$

Rubi [A] time = 0.04, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4122, 195, 217, 206}

$$\frac{5}{16}a^3 \tan(x) \sqrt{a \sec^2(x)} + \frac{5}{24}a^2 \tan(x) (a \sec^2(x))^{3/2} + \frac{5}{16}a^{7/2} \tanh^{-1}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{1}{6}a \tan(x) (a \sec^2(x))^{5/2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(a*\operatorname{Sec}[x]^2)^{(7/2)}, x]$

[Out] $(5*a^{(7/2)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tan}[x])/\operatorname{Sqrt}[a*\operatorname{Sec}[x]^2]])/16 + (5*a^3*\operatorname{Sqrt}[a*\operatorname{Sec}[x]^2]*\operatorname{Tan}[x])/16 + (5*a^2*(a*\operatorname{Sec}[x]^2)^{(3/2)}*\operatorname{Tan}[x])/24 + (a*(a*\operatorname{Sec}[x]^2)^{(5/2)}*\operatorname{Tan}[x])/6$

Rule 195

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^{(n_+)})^{(p_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(x_+*(a_+ + b_+*x_+^{n_+})^{p_+})/(n_+*p_+ + 1), x] + \operatorname{Dist}[(a_+*n_+*p_+)/(n_+*p_+ + 1), \operatorname{Int}[(a_+ + b_+*x_+^{n_+})^{(p_+ - 1)}, x], x] /;$ Free Q[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1*\operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2]*x)/\operatorname{Rt}[a, 2]])/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]), x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_+ + (b_+)*(x_+)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b_+*x^2), x], x, x/\operatorname{Sqrt}[a_+ + b_+*x^2]] /;$ FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
 \int (a \sec^2(x))^{7/2} dx &= a \operatorname{Subst} \left(\int (a + ax^2)^{5/2} dx, x, \tan(x) \right) \\
 &= \frac{1}{6} a (a \sec^2(x))^{5/2} \tan(x) + \frac{1}{6} (5a^2) \operatorname{Subst} \left(\int (a + ax^2)^{3/2} dx, x, \tan(x) \right) \\
 &= \frac{5}{24} a^2 (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{6} a (a \sec^2(x))^{5/2} \tan(x) + \frac{1}{8} (5a^3) \operatorname{Subst} \left(\int \sqrt{a + ax^2} dx, x, \tan(x) \right) \\
 &= \frac{5}{16} a^3 \sqrt{a \sec^2(x)} \tan(x) + \frac{5}{24} a^2 (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{6} a (a \sec^2(x))^{5/2} \tan(x) + \frac{1}{16} (5a^4) \operatorname{Subst} \left(\int \sqrt{a + ax^2} dx, x, \tan(x) \right) \\
 &= \frac{5}{16} a^3 \sqrt{a \sec^2(x)} \tan(x) + \frac{5}{24} a^2 (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{6} a (a \sec^2(x))^{5/2} \tan(x) + \frac{1}{16} (5a^4) \operatorname{Subst} \left(\int \sqrt{a + ax^2} dx, x, \tan(x) \right) \\
 &= \frac{5}{16} a^{7/2} \tanh^{-1} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right) + \frac{5}{16} a^3 \sqrt{a \sec^2(x)} \tan(x) + \frac{5}{24} a^2 (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{6} a (a \sec^2(x))^{5/2} \tan(x) + \frac{1}{16} (5a^4) \operatorname{Subst} \left(\int \sqrt{a + ax^2} dx, x, \tan(x) \right)
 \end{aligned}$$

Mathematica [A] time = 0.14, size = 78, normalized size = 0.93

$$\frac{1}{96} \cos^7(x) (a \sec^2(x))^{7/2} \left(\frac{1}{8} (198 \sin(x) + 85 \sin(3x) + 15 \sin(5x)) \sec^6(x) - 30 \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + 30 \log \left(\cos \left(\frac{x}{2} \right) + \sin \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^2)^(7/2), x]

[Out] (Cos[x]^7*(a*Sec[x]^2)^(7/2)*(-30*Log[Cos[x/2] - Sin[x/2]] + 30*Log[Cos[x/2] + Sin[x/2]] + (Sec[x]^6*(198*Sin[x] + 85*Sin[3*x] + 15*Sin[5*x]))/8))/96

fricas [A] time = 0.48, size = 65, normalized size = 0.77

$$\frac{\left(15 a^3 \cos(x)^6 \log \left(-\frac{\sin(x)-1}{\sin(x)+1} \right) - 2 \left(15 a^3 \cos(x)^4 + 10 a^3 \cos(x)^2 + 8 a^3 \right) \sin(x) \right) \sqrt{\frac{a}{\cos(x)^2}}}{96 \cos(x)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(7/2), x, algorithm="fricas")

[Out] $-1/96*(15*a^3*\cos(x)^6*\log(-(\sin(x) - 1)/(\sin(x) + 1)) - 2*(15*a^3*\cos(x)^4 + 10*a^3*\cos(x)^2 + 8*a^3)*\sin(x))*\sqrt{a/\cos(x)^2}/\cos(x)^5$

giac [A] time = 0.33, size = 79, normalized size = 0.94

$$\frac{1}{96} \left(15 a^3 \log(\sin(x) + 1) \operatorname{sgn}(\cos(x)) - 15 a^3 \log(-\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \frac{2(15 a^3 \operatorname{sgn}(\cos(x)) \sin(x)^5 - 40 a^3 \operatorname{sgn}(\cos(x)) \sin(x)^3 + 33 a^3 \operatorname{sgn}(\cos(x)) \sin(x))}{(\sin(x)^2 - 1)^3} \sqrt{a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sec(x)^2)^(7/2),x, algorithm="giac")`

[Out] $1/96*(15*a^3*\log(\sin(x) + 1)*\operatorname{sgn}(\cos(x)) - 15*a^3*\log(-\sin(x) + 1)*\operatorname{sgn}(\cos(x)) - 2*(15*a^3*\operatorname{sgn}(\cos(x))*\sin(x)^5 - 40*a^3*\operatorname{sgn}(\cos(x))*\sin(x)^3 + 33*a^3*\operatorname{sgn}(\cos(x))*\sin(x))/(\sin(x)^2 - 1)^3)*\sqrt{a}$

maple [A] time = 0.45, size = 74, normalized size = 0.88

$$\frac{\left(15 \left(\cos^6(x) \right) \ln \left(-\frac{-1+\cos(x)+\sin(x)}{\sin(x)} \right) - 15 \left(\cos^6(x) \right) \ln \left(-\frac{-\sin(x)-1+\cos(x)}{\sin(x)} \right) - 15 \left(\cos^4(x) \right) \sin(x) - 10 \left(\cos^2(x) \right) \sin(x) \right)}{48}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sec(x)^2)^(7/2),x)`

[Out] $-1/48*(15*\cos(x)^6*\ln(-(-1+\cos(x)+\sin(x))/\sin(x))-15*\cos(x)^6*\ln(-(-\sin(x)-1+\cos(x))/\sin(x))-15*\cos(x)^4*\sin(x)-10*\cos(x)^2*\sin(x)-8*\sin(x))*\cos(x)*(a/\cos(x)^2)^(7/2)$

maxima [B] time = 3.54, size = 2175, normalized size = 25.89

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sec(x)^2)^(7/2),x, algorithm="maxima")`

[Out] $1/96*(2040*a^3*\cos(3*x)*\sin(2*x) + 360*a^3*\cos(x)*\sin(2*x) - 360*a^3*\cos(2*x)*\sin(x) - 60*a^3*\sin(x) + 4*(15*a^3*\sin(11*x) + 85*a^3*\sin(9*x) + 198*a^3*\sin(7*x) - 198*a^3*\sin(5*x) - 85*a^3*\sin(3*x) - 15*a^3*\sin(x))*\cos(12*x) - 60*(6*a^3*\sin(10*x) + 15*a^3*\sin(8*x) + 20*a^3*\sin(6*x) + 15*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\cos(11*x) + 24*(85*a^3*\sin(9*x) + 198*a^3*\sin(7*x) - 198*a^3*\sin(5*x) - 85*a^3*\sin(3*x) - 15*a^3*\sin(x))*\cos(10*x) - 340*(15*a^3*\sin(8*x) + 20*a^3*\sin(6*x) + 15*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\cos(9*x) + 60*(198*a^3*\sin(7*x) - 198*a^3*\sin(5*x) - 85*a^3*\sin(3*x) - 15*a^3*\sin(x))*\cos(8*x)$

$$\begin{aligned}
& 8*x) - 792*(20*a^3*\sin(6*x) + 15*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\cos(7*x) - \\
& 80*(198*a^3*\sin(5*x) + 85*a^3*\sin(3*x) + 15*a^3*\sin(x))*\cos(6*x) + 2376*(5* \\
& a^3*\sin(4*x) + 2*a^3*\sin(2*x))*\cos(5*x) - 300*(17*a^3*\sin(3*x) + 3*a^3*\sin(\\
& x))*\cos(4*x) + 15*(a^3*\cos(12*x)^2 + 36*a^3*\cos(10*x)^2 + 225*a^3*\cos(8*x)^ \\
& 2 + 400*a^3*\cos(6*x)^2 + 225*a^3*\cos(4*x)^2 + 36*a^3*\cos(2*x)^2 + a^3*\sin(1 \\
& 2*x)^2 + 36*a^3*\sin(10*x)^2 + 225*a^3*\sin(8*x)^2 + 400*a^3*\sin(6*x)^2 + 225 \\
& *a^3*\sin(4*x)^2 + 180*a^3*\sin(4*x)*\sin(2*x) + 36*a^3*\sin(2*x)^2 + 12*a^3*\co \\
& s(2*x) + a^3 + 2*(6*a^3*\cos(10*x) + 15*a^3*\cos(8*x) + 20*a^3*\cos(6*x) + 15* \\
& a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\cos(12*x) + 12*(15*a^3*\cos(8*x) + 20*a \\
& ^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\cos(10*x) + 30*(20*a^ \\
& 3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\cos(8*x) + 40*(15*a^3* \\
& \cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\cos(6*x) + 30*(6*a^3*\cos(2*x) + a^3)*\cos(4 \\
& *x) + 2*(6*a^3*\sin(10*x) + 15*a^3*\sin(8*x) + 20*a^3*\sin(6*x) + 15*a^3*\sin(4 \\
& *x) + 6*a^3*\sin(2*x))*\sin(12*x) + 12*(15*a^3*\sin(8*x) + 20*a^3*\sin(6*x) + 1 \\
& 5*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\sin(10*x) + 30*(20*a^3*\sin(6*x) + 15*a^3*\s \\
& in(4*x) + 6*a^3*\sin(2*x))*\sin(8*x) + 120*(5*a^3*\sin(4*x) + 2*a^3*\sin(2*x))* \\
& \sin(6*x))*\log(\cos(x)^2 + \sin(x)^2 + 2*\sin(x) + 1) - 15*(a^3*\cos(12*x)^2 + 3 \\
& 6*a^3*\cos(10*x)^2 + 225*a^3*\cos(8*x)^2 + 400*a^3*\cos(6*x)^2 + 225*a^3*\cos(4 \\
& *x)^2 + 36*a^3*\cos(2*x)^2 + a^3*\sin(12*x)^2 + 36*a^3*\sin(10*x)^2 + 225*a^3* \\
& \sin(8*x)^2 + 400*a^3*\sin(6*x)^2 + 225*a^3*\sin(4*x)^2 + 180*a^3*\sin(4*x)*\sin \\
& (2*x) + 36*a^3*\sin(2*x)^2 + 12*a^3*\cos(2*x) + a^3 + 2*(6*a^3*\cos(10*x) + 15 \\
& *a^3*\cos(8*x) + 20*a^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\c \\
& os(12*x) + 12*(15*a^3*\cos(8*x) + 20*a^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3* \\
& \cos(2*x) + a^3)*\cos(10*x) + 30*(20*a^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\c \\
& os(2*x) + a^3)*\cos(8*x) + 40*(15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\cos(6 \\
& *x) + 30*(6*a^3*\cos(2*x) + a^3)*\cos(4*x) + 2*(6*a^3*\sin(10*x) + 15*a^3*\sin(\\
& 8*x) + 20*a^3*\sin(6*x) + 15*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\sin(12*x) + 12*(\\
& 15*a^3*\sin(8*x) + 20*a^3*\sin(6*x) + 15*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\sin(1 \\
& 0*x) + 30*(20*a^3*\sin(6*x) + 15*a^3*\sin(4*x) + 6*a^3*\sin(2*x))*\sin(8*x) + 1 \\
& 20*(5*a^3*\sin(4*x) + 2*a^3*\sin(2*x))*\sin(6*x))*\log(\cos(x)^2 + \sin(x)^2 - 2* \\
& \sin(x) + 1) - 4*(15*a^3*\cos(11*x) + 85*a^3*\cos(9*x) + 198*a^3*\cos(7*x) - 19 \\
& 8*a^3*\cos(5*x) - 85*a^3*\cos(3*x) - 15*a^3*\cos(x))*\sin(12*x) + 60*(6*a^3*\cos \\
& (10*x) + 15*a^3*\cos(8*x) + 20*a^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\cos(2* \\
& x) + a^3)*\sin(11*x) - 24*(85*a^3*\cos(9*x) + 198*a^3*\cos(7*x) - 198*a^3*\cos(\\
& 5*x) - 85*a^3*\cos(3*x) - 15*a^3*\cos(x))*\sin(10*x) + 340*(15*a^3*\cos(8*x) + \\
& 20*a^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\sin(9*x) - 60*(19 \\
& 8*a^3*\cos(7*x) - 198*a^3*\cos(5*x) - 85*a^3*\cos(3*x) - 15*a^3*\cos(x))*\sin(8* \\
& x) + 792*(20*a^3*\cos(6*x) + 15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\sin(7*x \\
&) + 80*(198*a^3*\cos(5*x) + 85*a^3*\cos(3*x) + 15*a^3*\cos(x))*\sin(6*x) - 792* \\
& (15*a^3*\cos(4*x) + 6*a^3*\cos(2*x) + a^3)*\sin(5*x) + 300*(17*a^3*\cos(3*x) + \\
& 3*a^3*\cos(x))*\sin(4*x) - 340*(6*a^3*\cos(2*x) + a^3)*\sin(3*x))*\sqrt{a}/(2*(6 \\
& *\cos(10*x) + 15*\cos(8*x) + 20*\cos(6*x) + 15*\cos(4*x) + 6*\cos(2*x) + 1)*\cos(\\
& 12*x) + \cos(12*x)^2 + 12*(15*\cos(8*x) + 20*\cos(6*x) + 15*\cos(4*x) + 6*\cos(2 \\
& *x) + 1)*\cos(10*x) + 36*\cos(10*x)^2 + 30*(20*\cos(6*x) + 15*\cos(4*x) + 6*\cos \\
& (2*x) + 1)*\cos(8*x) + 225*\cos(8*x)^2 + 40*(15*\cos(4*x) + 6*\cos(2*x) + 1)*\co
\end{aligned}$$

$s(6*x) + 400*\cos(6*x)^2 + 30*(6*\cos(2*x) + 1)*\cos(4*x) + 225*\cos(4*x)^2 + 36*\cos(2*x)^2 + 2*(6*\sin(10*x) + 15*\sin(8*x) + 20*\sin(6*x) + 15*\sin(4*x) + 6*\sin(2*x))*\sin(12*x) + \sin(12*x)^2 + 12*(15*\sin(8*x) + 20*\sin(6*x) + 15*\sin(4*x) + 6*\sin(2*x))*\sin(10*x) + 36*\sin(10*x)^2 + 30*(20*\sin(6*x) + 15*\sin(4*x) + 6*\sin(2*x))*\sin(8*x) + 225*\sin(8*x)^2 + 120*(5*\sin(4*x) + 2*\sin(2*x))*\sin(6*x) + 400*\sin(6*x)^2 + 225*\sin(4*x)^2 + 180*\sin(4*x)*\sin(2*x) + 36*\sin(2*x)^2 + 12*\cos(2*x) + 1)$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{a}{\cos(x)^2} \right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^2)^(7/2), x)

[Out] int((a/cos(x)^2)^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**2)**(7/2), x)

[Out] Timed out

3.48 $\int (a \sec^2(x))^{5/2} dx$

Optimal. Leaf size=65

$$\frac{3}{8}a^{5/2} \tanh^{-1}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{3}{8}a^2 \tan(x) \sqrt{a \sec^2(x)} + \frac{1}{4}a \tan(x) (a \sec^2(x))^{3/2}$$

[Out] $3/8*a^{(5/2)}*\operatorname{arctanh}(a^{(1/2)}*\tan(x)/(a*\sec(x)^2)^{(1/2)})+1/4*a*(a*\sec(x)^2)^{(3/2)}*\tan(x)+3/8*a^2*(a*\sec(x)^2)^{(1/2)}*\tan(x)$

Rubi [A] time = 0.03, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4122, 195, 217, 206}

$$\frac{3}{8}a^2 \tan(x) \sqrt{a \sec^2(x)} + \frac{3}{8}a^{5/2} \tanh^{-1}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{1}{4}a \tan(x) (a \sec^2(x))^{3/2}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^2)^(5/2), x]

[Out] $(3*a^{(5/2)}*ArcTanh[(Sqrt[a]*Tan[x])/Sqrt[a*Sec[x]^2]])/8 + (3*a^2*Sqrt[a*Sec[x]^2]*Tan[x])/8 + (a*(a*Sec[x]^2)^{(3/2)}*Tan[x])/4$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 4122

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFac
tors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1),
x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\int (a \sec^2(x))^{5/2} dx &= a \operatorname{Subst} \left(\int (a + ax^2)^{3/2} dx, x, \tan(x) \right) \\
&= \frac{1}{4} a (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{4} (3a^2) \operatorname{Subst} \left(\int \sqrt{a + ax^2} dx, x, \tan(x) \right) \\
&= \frac{3}{8} a^2 \sqrt{a \sec^2(x)} \tan(x) + \frac{1}{4} a (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{8} (3a^3) \operatorname{Subst} \left(\int \frac{1}{\sqrt{a + ax^2}} dx, x, \tan(x) \right) \\
&= \frac{3}{8} a^2 \sqrt{a \sec^2(x)} \tan(x) + \frac{1}{4} a (a \sec^2(x))^{3/2} \tan(x) + \frac{1}{8} (3a^3) \operatorname{Subst} \left(\int \frac{1}{1 - ax^2} dx, x, \frac{\tan(x)}{\sqrt{a \sec^2(x)}} \right) \\
&= \frac{3}{8} a^{5/2} \tanh^{-1} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right) + \frac{3}{8} a^2 \sqrt{a \sec^2(x)} \tan(x) + \frac{1}{4} a (a \sec^2(x))^{3/2} \tan(x)
\end{aligned}$$

Mathematica [A] time = 0.13, size = 72, normalized size = 1.11

$$\frac{1}{16} \cos^5(x) (a \sec^2(x))^{5/2} \left(\frac{1}{2} (11 \sin(x) + 3 \sin(3x)) \sec^4(x) - 6 \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + 6 \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^2)^(5/2), x]

[Out] (Cos[x]^5*(a*Sec[x]^2)^(5/2)*(-6*Log[Cos[x/2] - Sin[x/2]] + 6*Log[Cos[x/2] + Sin[x/2]] + (Sec[x]^4*(11*Sin[x] + 3*Sin[3*x]))/2))/16

fricas [A] time = 0.83, size = 56, normalized size = 0.86

$$\frac{\left(3 a^2 \cos(x)^4 \log \left(-\frac{\sin(x)-1}{\sin(x)+1} \right) - 2 \left(3 a^2 \cos(x)^2 + 2 a^2 \right) \sin(x) \right) \sqrt{\frac{a}{\cos(x)^2}}}{16 \cos(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(5/2), x, algorithm="fricas")

[Out] -1/16*(3*a^2*cos(x)^4*log(-(sin(x) - 1)/(sin(x) + 1)) - 2*(3*a^2*cos(x)^2 + 2*a^2)*sin(x))*sqrt(a/cos(x)^2)/cos(x)^3

giac [A] time = 0.41, size = 67, normalized size = 1.03

$$\frac{1}{16} \left(3a^2 \log(\sin(x) + 1) \operatorname{sgn}(\cos(x)) - 3a^2 \log(-\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \frac{2(3a^2 \operatorname{sgn}(\cos(x)) \sin(x)^3 - 5a^2 \operatorname{sgn}(\cos(x)) \sin(x))}{(\sin(x)^2 - 1)^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(5/2),x, algorithm="giac")

[Out] 1/16*(3*a^2*log(sin(x) + 1)*sgn(cos(x)) - 3*a^2*log(-sin(x) + 1)*sgn(cos(x)) - 2*(3*a^2*sgn(cos(x))*sin(x)^3 - 5*a^2*sgn(cos(x))*sin(x))/(sin(x)^2 - 1)^2)*sqrt(a)

maple [A] time = 0.36, size = 66, normalized size = 1.02

$$\frac{\left(3 \left(\cos^4(x) \right) \ln \left(-\frac{-1+\cos(x)+\sin(x)}{\sin(x)} \right) - 3 \left(\cos^4(x) \right) \ln \left(-\frac{-\sin(x)-1+\cos(x)}{\sin(x)} \right) - 3 \left(\cos^2(x) \right) \sin(x) - 2 \sin(x) \right) \cos(x) \left(\frac{1}{\cos(x)} \right)}{8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^2)^(5/2),x)

[Out] -1/8*(3*cos(x)^4*ln(-(-1+cos(x)+sin(x))/sin(x))-3*cos(x)^4*ln(-(-sin(x)-1+cos(x))/sin(x))-3*cos(x)^2*sin(x)-2*sin(x))*cos(x)*(a/cos(x)^2)^(5/2)

maxima [B] time = 1.00, size = 1111, normalized size = 17.09

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(5/2),x, algorithm="maxima")

[Out] 1/16*(176*a^2*cos(3*x)*sin(2*x) + 48*a^2*cos(x)*sin(2*x) - 48*a^2*cos(2*x)*sin(x) - 12*a^2*sin(x) + 4*(3*a^2*sin(7*x) + 11*a^2*sin(5*x) - 11*a^2*sin(3*x) - 3*a^2*sin(x))*cos(8*x) - 24*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))*cos(7*x) + 16*(11*a^2*sin(5*x) - 11*a^2*sin(3*x) - 3*a^2*sin(x))*cos(6*x) - 88*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*cos(5*x) - 24*(11*a^2*sin(3*x) + 3*a^2*sin(x))*cos(4*x) + 3*(a^2*cos(8*x)^2 + 16*a^2*cos(6*x)^2 + 36*a^2*cos(4*x)^2 + 16*a^2*cos(2*x)^2 + a^2*sin(8*x)^2 + 16*a^2*sin(6*x)^2 + 36*a^2*sin(4*x)^2 + 48*a^2*sin(4*x)*sin(2*x) + 16*a^2*sin(2*x)^2 + 8*a^2*cos(2*x) + a^2 + 2*(4*a^2*cos(6*x) + 6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(8*x) + 8*(6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(6*x) + 12*(4*a^2*cos(2*x) + a^2)*cos(4*x) + 4*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))

```

*sin(8*x) + 16*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(6*x))*log(cos(x)^2 + s
in(x)^2 + 2*sin(x) + 1) - 3*(a^2*cos(8*x)^2 + 16*a^2*cos(6*x)^2 + 36*a^2*co
s(4*x)^2 + 16*a^2*cos(2*x)^2 + a^2*sin(8*x)^2 + 16*a^2*sin(6*x)^2 + 36*a^2*
sin(4*x)^2 + 48*a^2*sin(4*x)*sin(2*x) + 16*a^2*sin(2*x)^2 + 8*a^2*cos(2*x)
+ a^2 + 2*(4*a^2*cos(6*x) + 6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(8*x)
+ 8*(6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*cos(6*x) + 12*(4*a^2*cos(2*x)
+ a^2)*cos(4*x) + 4*(2*a^2*sin(6*x) + 3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(
8*x) + 16*(3*a^2*sin(4*x) + 2*a^2*sin(2*x))*sin(6*x))*log(cos(x)^2 + sin(x)
^2 - 2*sin(x) + 1) - 4*(3*a^2*cos(7*x) + 11*a^2*cos(5*x) - 11*a^2*cos(3*x)
- 3*a^2*cos(x))*sin(8*x) + 12*(4*a^2*cos(6*x) + 6*a^2*cos(4*x) + 4*a^2*cos(
2*x) + a^2)*sin(7*x) - 16*(11*a^2*cos(5*x) - 11*a^2*cos(3*x) - 3*a^2*cos(x)
)*sin(6*x) + 44*(6*a^2*cos(4*x) + 4*a^2*cos(2*x) + a^2)*sin(5*x) + 24*(11*a
^2*cos(3*x) + 3*a^2*cos(x))*sin(4*x) - 44*(4*a^2*cos(2*x) + a^2)*sin(3*x))*
sqrt(a)/(2*(4*cos(6*x) + 6*cos(4*x) + 4*cos(2*x) + 1)*cos(8*x) + cos(8*x)^2
+ 8*(6*cos(4*x) + 4*cos(2*x) + 1)*cos(6*x) + 16*cos(6*x)^2 + 12*(4*cos(2*x)
) + 1)*cos(4*x) + 36*cos(4*x)^2 + 16*cos(2*x)^2 + 4*(2*sin(6*x) + 3*sin(4*x)
) + 2*sin(2*x))*sin(8*x) + sin(8*x)^2 + 16*(3*sin(4*x) + 2*sin(2*x))*sin(6*
x) + 16*sin(6*x)^2 + 36*sin(4*x)^2 + 48*sin(4*x)*sin(2*x) + 16*sin(2*x)^2 +
8*cos(2*x) + 1)

```

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{a}{\cos(x)^2} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^2)^(5/2), x)

[Out] int((a/cos(x)^2)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec^2(x))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**2)**(5/2), x)

[Out] Integral((a*sec(x)**2)**(5/2), x)

3.49 $\int (a \sec^2(x))^{3/2} dx$

Optimal. Leaf size=46

$$\frac{1}{2}a^{3/2} \tanh^{-1}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{1}{2}a \tan(x) \sqrt{a \sec^2(x)}$$

[Out] $1/2*a^{(3/2)}*\operatorname{arctanh}(a^{(1/2)}*\tan(x)/(a*\sec(x)^2)^{(1/2)})+1/2*a*(a*\sec(x)^2)^{(1/2)}*\tan(x)$

Rubi [A] time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4122, 195, 217, 206}

$$\frac{1}{2}a^{3/2} \tanh^{-1}\left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}}\right) + \frac{1}{2}a \tan(x) \sqrt{a \sec^2(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^2)^(3/2), x]

[Out] $(a^{(3/2)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[a]*\operatorname{Tan}[x])/\operatorname{Sqrt}[a*\operatorname{Sec}[x]^2]])/2 + (a*\operatorname{Sqrt}[a*\operatorname{Sec}[x]^2]*\operatorname{Tan}[x])/2$

Rule 195

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(x*(a + b*x^n)^p)/(n*p + 1), x] + Dist[(a*n*p)/(n*p + 1), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 4122

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x]] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \int (a \sec^2(x))^{3/2} dx &= a \operatorname{Subst} \left(\int \sqrt{a + ax^2} dx, x, \tan(x) \right) \\ &= \frac{1}{2} a \sqrt{a \sec^2(x)} \tan(x) + \frac{1}{2} a^2 \operatorname{Subst} \left(\int \frac{1}{\sqrt{a + ax^2}} dx, x, \tan(x) \right) \\ &= \frac{1}{2} a \sqrt{a \sec^2(x)} \tan(x) + \frac{1}{2} a^2 \operatorname{Subst} \left(\int \frac{1}{1 - ax^2} dx, x, \frac{\tan(x)}{\sqrt{a \sec^2(x)}} \right) \\ &= \frac{1}{2} a^{3/2} \tanh^{-1} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right) + \frac{1}{2} a \sqrt{a \sec^2(x)} \tan(x) \end{aligned}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 1.20

$$\frac{1}{2} a \cos(x) \sqrt{a \sec^2(x)} \left(\tan(x) \sec(x) - \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) + \log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[(a*Sec[x]^2)^(3/2), x]
```

```
[Out] (a*Cos[x]*Sqrt[a*Sec[x]^2]*(-Log[Cos[x/2] - Sin[x/2]] + Log[Cos[x/2] + Sin[x/2]] + Sec[x]*Tan[x]))/2
```

fricas [A] time = 0.67, size = 39, normalized size = 0.85

$$\frac{\left(a \cos(x)^2 \log \left(-\frac{\sin(x)-1}{\sin(x)+1} \right) - 2 a \sin(x) \right) \sqrt{\frac{a}{\cos(x)^2}}}{4 \cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sec(x)^2)^(3/2), x, algorithm="fricas")
```

```
[Out] -1/4*(a*cos(x)^2*log(-(sin(x) - 1)/(sin(x) + 1)) - 2*a*sin(x))*sqrt(a/cos(x)^2)/cos(x)
```

giac [A] time = 0.60, size = 42, normalized size = 0.91

$$\frac{1}{4} \left(\log(\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \log(-\sin(x) + 1) \operatorname{sgn}(\cos(x)) - \frac{2 \operatorname{sgn}(\cos(x)) \sin(x)}{\sin(x)^2 - 1} \right) a^{3/2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(3/2),x, algorithm="giac")

[Out] $\frac{1}{4} * (\log(\sin(x) + 1) * \text{sgn}(\cos(x)) - \log(-\sin(x) + 1) * \text{sgn}(\cos(x)) - 2 * \text{sgn}(\cos(x)) * \sin(x) / (\sin(x)^2 - 1)) * a^{3/2}$

maple [A] time = 0.30, size = 57, normalized size = 1.24

$$\frac{\left((\cos^2(x)) \ln\left(-\frac{-1+\cos(x)+\sin(x)}{\sin(x)}\right) - (\cos^2(x)) \ln\left(-\frac{-\sin(x)-1+\cos(x)}{\sin(x)}\right) - \sin(x) \right) \cos(x) \left(\frac{a}{\cos(x)^2}\right)^{\frac{3}{2}}}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^2)^(3/2),x)

[Out] $-1/2 * (\cos(x)^2 * \ln(-(-1+\cos(x)+\sin(x))/\sin(x)) - \cos(x)^2 * \ln(-(-\sin(x)-1+\cos(x))/\sin(x)) - \sin(x)) * \cos(x) * (a/\cos(x)^2)^{3/2}$

maxima [B] time = 1.03, size = 324, normalized size = 7.04

$$\frac{(8a \cos(3x) \sin(2x) - 8a \cos(x) \sin(2x) + 8a \cos(2x) \sin(x) - 4(a \sin(3x) - a \sin(x)) \cos(4x) - (a \cos(4x) - a \cos(2x)) \sin(4x)) \sqrt{a}}{2 \cos^2(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(3/2),x, algorithm="maxima")

[Out] $-1/4 * (8 * a * \cos(3 * x) * \sin(2 * x) - 8 * a * \cos(x) * \sin(2 * x) + 8 * a * \cos(2 * x) * \sin(x) - 4 * (a * \sin(3 * x) - a * \sin(x)) * \cos(4 * x) - (a * \cos(4 * x)^2 + 4 * a * \cos(2 * x)^2 + a * \sin(4 * x)^2 + 4 * a * \sin(4 * x) * \sin(2 * x) + 4 * a * \sin(2 * x)^2 + 2 * (2 * a * \cos(2 * x) + a) * \cos(4 * x) + 4 * a * \cos(2 * x) + a) * \log(\cos(x)^2 + \sin(x)^2 + 2 * \sin(x) + 1) + (a * \cos(4 * x)^2 + 4 * a * \cos(2 * x)^2 + a * \sin(4 * x)^2 + 4 * a * \sin(4 * x) * \sin(2 * x) + 4 * a * \sin(2 * x)^2 + 2 * (2 * a * \cos(2 * x) + a) * \cos(4 * x) + 4 * a * \cos(2 * x) + a) * \log(\cos(x)^2 + \sin(x)^2 - 2 * \sin(x) + 1) + 4 * (a * \cos(3 * x) - a * \cos(x)) * \sin(4 * x) - 4 * (2 * a * \cos(2 * x) + a) * \sin(3 * x) + 4 * a * \sin(x)) * \sqrt{a} / (2 * (2 * \cos(2 * x) + 1) * \cos(4 * x) + \cos(4 * x)^2 + 4 * \cos(2 * x)^2 + \sin(4 * x)^2 + 4 * \sin(4 * x) * \sin(2 * x) + 4 * \sin(2 * x)^2 + 4 * \cos(2 * x) + 1)$

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{a}{\cos(x)^2} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a/cos(x)^2)^(3/2),x)
```

```
[Out] int((a/cos(x)^2)^(3/2), x)
```

```
sympy [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int (a \sec^2(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sec(x)**2)**(3/2),x)
```

```
[Out] Integral((a*sec(x)**2)**(3/2), x)
```


3.50 $\int \sqrt{a \sec^2(x)} dx$

Optimal. Leaf size=25

$$\sqrt{a} \tanh^{-1} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right)$$

[Out] arctanh(a^(1/2)*tan(x)/(a*sec(x)^2)^(1/2))*a^(1/2)

Rubi [A] time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4122, 217, 206}

$$\sqrt{a} \tanh^{-1} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right)$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Sec[x]^2], x]

[Out] Sqrt[a]*ArcTanh[(Sqrt[a]*Tan[x])/Sqrt[a*Sec[x]^2]]

Rule 206

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^2)^(p_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \sqrt{a \sec^2(x)} dx &= a \operatorname{Subst} \left(\int \frac{1}{\sqrt{a + ax^2}} dx, x, \tan(x) \right) \\ &= a \operatorname{Subst} \left(\int \frac{1}{1 - ax^2} dx, x, \frac{\tan(x)}{\sqrt{a \sec^2(x)}} \right) \\ &= \sqrt{a} \tanh^{-1} \left(\frac{\sqrt{a} \tan(x)}{\sqrt{a \sec^2(x)}} \right) \end{aligned}$$

Mathematica [A] time = 0.01, size = 46, normalized size = 1.84

$$\cos(x) \sqrt{a \sec^2(x)} \left(\log \left(\sin \left(\frac{x}{2} \right) + \cos \left(\frac{x}{2} \right) \right) - \log \left(\cos \left(\frac{x}{2} \right) - \sin \left(\frac{x}{2} \right) \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Sec[x]^2],x]

[Out] Cos[x]*(-Log[Cos[x/2] - Sin[x/2]] + Log[Cos[x/2] + Sin[x/2]])*Sqrt[a*Sec[x]^2]

fricas [A] time = 0.64, size = 55, normalized size = 2.20

$$\left[-\frac{1}{2} \sqrt{\frac{a}{\cos(x)^2}} \cos(x) \log \left(-\frac{\sin(x) - 1}{\sin(x) + 1} \right), -\sqrt{-a} \arctan \left(\frac{\sqrt{-a} \sqrt{\frac{a}{\cos(x)^2}} \cos(x) \sin(x)}{a} \right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(1/2),x, algorithm="fricas")

[Out] [-1/2*sqrt(a/cos(x)^2)*cos(x)*log(-(sin(x) - 1)/(sin(x) + 1)), -sqrt(-a)*arctan(sqrt(-a)*sqrt(a/cos(x)^2)*cos(x)*sin(x)/a)]

giac [A] time = 0.33, size = 31, normalized size = 1.24

$$\frac{1}{4} \sqrt{a} \left(\log \left(\left| \frac{1}{\sin(x)} + \sin(x) + 2 \right| \right) - \log \left(\left| \frac{1}{\sin(x)} + \sin(x) - 2 \right| \right) \right) \operatorname{sgn}(\cos(x))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^2)^(1/2),x, algorithm="giac")

[Out] 1/4*sqrt(a)*(log(abs(1/sin(x) + sin(x) + 2)) - log(abs(1/sin(x) + sin(x) - 2)))*sgn(cos(x))

maple [A] time = 0.38, size = 23, normalized size = 0.92

$$-2 \cos(x) \sqrt{\frac{a}{\cos(x)^2}} \operatorname{arctanh}\left(\frac{-1 + \cos(x)}{\sin(x)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sec(x)^2)^(1/2),x)`

[Out] `-2*cos(x)*(a/cos(x)^2)^(1/2)*arctanh((-1+cos(x))/sin(x))`

maxima [A] time = 0.89, size = 38, normalized size = 1.52

$$\frac{1}{2} \sqrt{a} \left(\log(\cos(x)^2 + \sin(x)^2 + 2 \sin(x) + 1) - \log(\cos(x)^2 + \sin(x)^2 - 2 \sin(x) + 1) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sec(x)^2)^(1/2),x, algorithm="maxima")`

[Out] `1/2*sqrt(a)*(log(cos(x)^2 + sin(x)^2 + 2*sin(x) + 1) - log(cos(x)^2 + sin(x)^2 - 2*sin(x) + 1))`

mupad [F] time = 0.00, size = -1, normalized size = -0.04

$$\int \sqrt{\frac{a}{\cos(x)^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a/cos(x)^2)^(1/2),x)`

[Out] `int((a/cos(x)^2)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \sec^2(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sec(x)**2)**(1/2),x)`

[Out] `Integral(sqrt(a*sec(x)**2), x)`

$$3.51 \quad \int \frac{1}{\sqrt{a \sec^2(x)}} dx$$

Optimal. Leaf size=13

$$\frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

[Out] $\tan(x)/(a*\sec(x)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {4122, 191}

$$\frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/\text{Sqrt}[a*\text{Sec}[x]^2], x]$

[Out] $\text{Tan}[x]/\text{Sqrt}[a*\text{Sec}[x]^2]$

Rule 191

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] := \text{Simp}[(x*(a + b*x^n)^{(p + 1)})/a, x] /;$ $\text{FreeQ}\{a, b, n, p\}, x\} \ \&\& \ \text{EqQ}[1/n + p + 1, 0]$

Rule 4122

$\text{Int}[(b_)*\sec[(e_ + (f_)*(x_))]^2]^{(p_)}, x_Symbol] := \text{With}\{\{ff = \text{FreeFactors}[\text{Tan}[e + f*x], x]\}, \text{Dist}[(b*ff)/f, \text{Subst}[\text{Int}[(b + b*ff^2*x^2)^{(p - 1)}, x], x, \text{Tan}[e + f*x]/ff], x]\} /;$ $\text{FreeQ}\{b, e, f, p\}, x\} \ \&\& \ !\text{IntegerQ}[p]$

Rubi steps

$$\int \frac{1}{\sqrt{a \sec^2(x)}} dx = a \text{Subst} \left(\int \frac{1}{(a + ax^2)^{3/2}} dx, x, \tan(x) \right) \\ = \frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

Mathematica [A] time = 0.01, size = 13, normalized size = 1.00

$$\frac{\tan(x)}{\sqrt{a \sec^2(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a*Sec[x]^2],x]

[Out] Tan[x]/Sqrt[a*Sec[x]^2]

fricas [A] time = 0.62, size = 16, normalized size = 1.23

$$\frac{\sqrt{\frac{a}{\cos(x)^2}} \cos(x) \sin(x)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(1/2),x, algorithm="fricas")

[Out] sqrt(a/cos(x)^2)*cos(x)*sin(x)/a

giac [A] time = 0.40, size = 11, normalized size = 0.85

$$\frac{\sin(x)}{\sqrt{a} \operatorname{sgn}(\cos(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(1/2),x, algorithm="giac")

[Out] sin(x)/(sqrt(a)*sgn(cos(x)))

maple [A] time = 0.40, size = 16, normalized size = 1.23

$$\frac{\sin(x)}{\sqrt{\frac{a}{\cos(x)^2}} \cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^2)^(1/2),x)

[Out] sin(x)/(a/cos(x)^2)^(1/2)/cos(x)

maxima [A] time = 0.92, size = 6, normalized size = 0.46

$$\frac{\sin(x)}{\sqrt{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(1/2),x, algorithm="maxima")

[Out] $\sin(x)/\sqrt{a}$

mupad [B] time = 0.21, size = 15, normalized size = 1.15

$$\frac{\sqrt{2} \sin(2x)}{2\sqrt{a} \sqrt{2\cos(x)^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(1/(a/\cos(x)^2)^{(1/2)}, x)$

[Out] $(2^{(1/2)}*\sin(2*x))/(2*a^{(1/2)}*(2*\cos(x)^2)^{(1/2)})$

sympy [A] time = 0.51, size = 15, normalized size = 1.15

$$\frac{\tan(x)}{\sqrt{a} \sqrt{\sec^2(x)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/(a*\sec(x)**2)**(1/2), x)$

[Out] $\tan(x)/(\sqrt{a}*\sqrt{\sec(x)**2})$

$$3.52 \quad \int \frac{1}{(a \sec^2(x))^{3/2}} dx$$

Optimal. Leaf size=36

$$\frac{2 \tan(x)}{3a\sqrt{a \sec^2(x)}} + \frac{\tan(x)}{3(a \sec^2(x))^{3/2}}$$

[Out] $1/3*\tan(x)/(a*\sec(x)^2)^{(3/2)}+2/3*\tan(x)/a/(a*\sec(x)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4122, 192, 191}

$$\frac{2 \tan(x)}{3a\sqrt{a \sec^2(x)}} + \frac{\tan(x)}{3(a \sec^2(x))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^2)^(-3/2),x]

[Out] Tan[x]/(3*(a*Sec[x]^2)^(3/2)) + (2*Tan[x])/(3*a*Sqrt[a*Sec[x]^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)]^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \sec^2(x))^{3/2}} dx &= a \operatorname{Subst} \left(\int \frac{1}{(a + ax^2)^{5/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{3 (a \sec^2(x))^{3/2}} + \frac{2}{3} \operatorname{Subst} \left(\int \frac{1}{(a + ax^2)^{3/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{3 (a \sec^2(x))^{3/2}} + \frac{2 \tan(x)}{3a \sqrt{a \sec^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 27, normalized size = 0.75

$$\frac{(9 \sin(x) + \sin(3x)) \sec^3(x)}{12 (a \sec^2(x))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^2)^(-3/2), x]

[Out] (Sec[x]^3*(9*Sin[x] + Sin[3*x]))/(12*(a*Sec[x]^2)^(3/2))

fricas [A] time = 0.56, size = 24, normalized size = 0.67

$$\frac{(\cos(x)^3 + 2 \cos(x)) \sqrt{\frac{a}{\cos(x)^2}} \sin(x)}{3 a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(3/2), x, algorithm="fricas")

[Out] 1/3*(cos(x)^3 + 2*cos(x))*sqrt(a/cos(x)^2)*sin(x)/a^2

giac [B] time = 0.60, size = 58, normalized size = 1.61

$$\frac{2 \left(3 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^2 \operatorname{sgn} \left(-\tan\left(\frac{1}{2}x\right)^2 + 1 \right) - 4 \operatorname{sgn} \left(-\tan\left(\frac{1}{2}x\right)^2 + 1 \right) \right)}{3 a^2 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(3/2),x, algorithm="giac")

[Out] $\frac{2}{3} * (3 * (1/\tan(1/2*x) + \tan(1/2*x))^{2} * \operatorname{sgn}(-\tan(1/2*x)^2 + 1) - 4 * \operatorname{sgn}(-\tan(1/2*x)^2 + 1)) / (a^{3/2} * (1/\tan(1/2*x) + \tan(1/2*x))^{3})$

maple [A] time = 0.30, size = 23, normalized size = 0.64

$$\frac{\sin(x) (\cos^2(x) + 2)}{3 \cos(x)^3 \left(\frac{a}{\cos(x)^2}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^2)^(3/2),x)

[Out] $\frac{1}{3} * \sin(x) * (\cos(x)^2 + 2) / \cos(x)^3 / (a / \cos(x)^2)^{3/2}$

maxima [A] time = 1.00, size = 14, normalized size = 0.39

$$\frac{\sin(3x) + 9 \sin(x)}{12 a^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(3/2),x, algorithm="maxima")

[Out] $\frac{1}{12} * (\sin(3*x) + 9 * \sin(x)) / a^{3/2}$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\left(\frac{a}{\cos(x)^2}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a/cos(x)^2)^(3/2),x)

[Out] int(1/(a/cos(x)^2)^(3/2), x)

sympy [A] time = 1.18, size = 37, normalized size = 1.03

$$\frac{2 \tan^3(x)}{3 a^{\frac{3}{2}} (\sec^2(x))^{\frac{3}{2}}} + \frac{\tan(x)}{a^{\frac{3}{2}} (\sec^2(x))^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*sec(x)**2)**(3/2),x)
```

```
[Out] 2*tan(x)**3/(3*a**(3/2)*(sec(x)**2)**(3/2)) + tan(x)/(a**(3/2)*(sec(x)**2)*  
*(3/2))
```

$$3.53 \quad \int \frac{1}{(a \sec^2(x))^{5/2}} dx$$

Optimal. Leaf size=55

$$\frac{8 \tan(x)}{15a^2 \sqrt{a \sec^2(x)}} + \frac{4 \tan(x)}{15a (a \sec^2(x))^{3/2}} + \frac{\tan(x)}{5 (a \sec^2(x))^{5/2}}$$

[Out] 1/5*tan(x)/(a*sec(x)^2)^(5/2)+4/15*tan(x)/a/(a*sec(x)^2)^(3/2)+8/15*tan(x)/a^2/(a*sec(x)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4122, 192, 191}

$$\frac{8 \tan(x)}{15a^2 \sqrt{a \sec^2(x)}} + \frac{4 \tan(x)}{15a (a \sec^2(x))^{3/2}} + \frac{\tan(x)}{5 (a \sec^2(x))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^2)^(-5/2), x]

[Out] Tan[x]/(5*(a*Sec[x]^2)^(5/2)) + (4*Tan[x])/(15*a*(a*Sec[x]^2)^(3/2)) + (8*Tan[x])/(15*a^2*Sqrt[a*Sec[x]^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \sec^2(x))^{5/2}} dx &= a \operatorname{Subst} \left(\int \frac{1}{(a + ax^2)^{7/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{5 (a \sec^2(x))^{5/2}} + \frac{4}{5} \operatorname{Subst} \left(\int \frac{1}{(a + ax^2)^{5/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{5 (a \sec^2(x))^{5/2}} + \frac{4 \tan(x)}{15a (a \sec^2(x))^{3/2}} + \frac{8 \operatorname{Subst} \left(\int \frac{1}{(a+ax^2)^{3/2}} dx, x, \tan(x) \right)}{15a} \\
&= \frac{\tan(x)}{5 (a \sec^2(x))^{5/2}} + \frac{4 \tan(x)}{15a (a \sec^2(x))^{3/2}} + \frac{8 \tan(x)}{15a^2 \sqrt{a \sec^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 36, normalized size = 0.65

$$\frac{(150 \sin(x) + 25 \sin(3x) + 3 \sin(5x)) \cos(x) \sqrt{a \sec^2(x)}}{240a^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^2)^(-5/2), x]

[Out] (Cos[x]*Sqrt[a*Sec[x]^2]*(150*Sin[x] + 25*Sin[3*x] + 3*Sin[5*x]))/(240*a^3)

fricas [A] time = 0.81, size = 32, normalized size = 0.58

$$\frac{(3 \cos(x)^5 + 4 \cos(x)^3 + 8 \cos(x)) \sqrt{\frac{a}{\cos(x)^2}} \sin(x)}{15 a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(5/2), x, algorithm="fricas")

[Out] 1/15*(3*cos(x)^5 + 4*cos(x)^3 + 8*cos(x))*sqrt(a/cos(x)^2)*sin(x)/a^3

giac [A] time = 0.54, size = 84, normalized size = 1.53

$$\frac{2 \left(15 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^4 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) - 40 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^2 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) + 48 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) \right)}{15 a^2 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(5/2),x, algorithm="giac")

[Out] $\frac{2}{15} * (15 * (1/\tan(1/2*x) + \tan(1/2*x))^4 * \text{sgn}(-\tan(1/2*x)^2 + 1) - 40 * (1/\tan(1/2*x) + \tan(1/2*x))^2 * \text{sgn}(-\tan(1/2*x)^2 + 1) + 48 * \text{sgn}(-\tan(1/2*x)^2 + 1)) / (a^{5/2} * (1/\tan(1/2*x) + \tan(1/2*x))^5)$

maple [A] time = 0.31, size = 31, normalized size = 0.56

$$\frac{\sin(x) \left(3 \left(\cos^4(x) \right) + 4 \left(\cos^2(x) \right) + 8 \right)}{15 \cos(x)^5 \left(\frac{a}{\cos(x)^2} \right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^2)^(5/2),x)

[Out] $\frac{1}{15} * \sin(x) * (3 * \cos(x)^4 + 4 * \cos(x)^2 + 8) / \cos(x)^5 / (a / \cos(x)^2)^{5/2}$

maxima [A] time = 0.68, size = 22, normalized size = 0.40

$$\frac{3 \sin(5x) + 25 \sin(3x) + 150 \sin(x)}{240 a^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(5/2),x, algorithm="maxima")

[Out] $\frac{1}{240} * (3 * \sin(5*x) + 25 * \sin(3*x) + 150 * \sin(x)) / a^{5/2}$

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{a}{\cos(x)^2} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a/cos(x)^2)^(5/2),x)

[Out] int(1/(a/cos(x)^2)^(5/2), x)

sympy [A] time = 10.57, size = 60, normalized size = 1.09

$$\frac{8 \tan^5(x)}{15 a^{\frac{5}{2}} \left(\sec^2(x) \right)^{\frac{5}{2}}} + \frac{4 \tan^3(x)}{3 a^{\frac{5}{2}} \left(\sec^2(x) \right)^{\frac{5}{2}}} + \frac{\tan(x)}{a^{\frac{5}{2}} \left(\sec^2(x) \right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*sec(x)**2)**(5/2),x)
```

```
[Out] 8*tan(x)**5/(15*a**(5/2)*(sec(x)**2)**(5/2)) + 4*tan(x)**3/(3*a**(5/2)*(sec(x)**2)**(5/2)) + tan(x)/(a**(5/2)*(sec(x)**2)**(5/2))
```

$$3.54 \quad \int \frac{1}{(a \sec^2(x))^{7/2}} dx$$

Optimal. Leaf size=74

$$\frac{16 \tan(x)}{35a^3 \sqrt{a \sec^2(x)}} + \frac{8 \tan(x)}{35a^2 (a \sec^2(x))^{3/2}} + \frac{6 \tan(x)}{35a (a \sec^2(x))^{5/2}} + \frac{\tan(x)}{7 (a \sec^2(x))^{7/2}}$$

[Out] $1/7*\tan(x)/(a*\sec(x)^2)^{(7/2)}+6/35*\tan(x)/a/(a*\sec(x)^2)^{(5/2)}+8/35*\tan(x)/a^2/(a*\sec(x)^2)^{(3/2)}+16/35*\tan(x)/a^3/(a*\sec(x)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4122, 192, 191}

$$\frac{16 \tan(x)}{35a^3 \sqrt{a \sec^2(x)}} + \frac{8 \tan(x)}{35a^2 (a \sec^2(x))^{3/2}} + \frac{6 \tan(x)}{35a (a \sec^2(x))^{5/2}} + \frac{\tan(x)}{7 (a \sec^2(x))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^2)^(-7/2), x]

[Out] Tan[x]/(7*(a*Sec[x]^2)^(7/2)) + (6*Tan[x])/(35*a*(a*Sec[x]^2)^(5/2)) + (8*Tan[x])/(35*a^2*(a*Sec[x]^2)^(3/2)) + (16*Tan[x])/(35*a^3*Sqrt[a*Sec[x]^2])

Rule 191

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(x*(a + b*x^n)^(p + 1))/a, x] /; FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 192

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> -Simp[(x*(a + b*x^n)^(p + 1))/(a*n*(p + 1)), x] + Dist[(n*(p + 1) + 1)/(a*n*(p + 1)), Int[(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, n, p}, x] && ILtQ[Simplify[1/n + p + 1], 0] && NeQ[p, -1]

Rule 4122

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^2)^(p_), x_Symbol] :> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[(b*ff)/f, Subst[Int[(b + b*ff^2*x^2)^(p - 1), x], x, Tan[e + f*x]/ff], x] /; FreeQ[{b, e, f, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \sec^2(x))^{7/2}} dx &= a \operatorname{Subst} \left(\int \frac{1}{(a + ax^2)^{9/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{7(a \sec^2(x))^{7/2}} + \frac{6}{7} \operatorname{Subst} \left(\int \frac{1}{(a + ax^2)^{7/2}} dx, x, \tan(x) \right) \\
&= \frac{\tan(x)}{7(a \sec^2(x))^{7/2}} + \frac{6 \tan(x)}{35a(a \sec^2(x))^{5/2}} + \frac{24 \operatorname{Subst} \left(\int \frac{1}{(a+ax^2)^{5/2}} dx, x, \tan(x) \right)}{35a} \\
&= \frac{\tan(x)}{7(a \sec^2(x))^{7/2}} + \frac{6 \tan(x)}{35a(a \sec^2(x))^{5/2}} + \frac{8 \tan(x)}{35a^2(a \sec^2(x))^{3/2}} + \frac{16 \operatorname{Subst} \left(\int \frac{1}{(a+ax^2)^{3/2}} dx, x, \tan(x) \right)}{35a^2} \\
&= \frac{\tan(x)}{7(a \sec^2(x))^{7/2}} + \frac{6 \tan(x)}{35a(a \sec^2(x))^{5/2}} + \frac{8 \tan(x)}{35a^2(a \sec^2(x))^{3/2}} + \frac{16 \tan(x)}{35a^3 \sqrt{a \sec^2(x)}}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 42, normalized size = 0.57

$$\frac{(1225 \sin(x) + 245 \sin(3x) + 49 \sin(5x) + 5 \sin(7x)) \cos(x) \sqrt{a \sec^2(x)}}{2240a^4}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^2)^(-7/2), x]

[Out] (Cos[x]*Sqrt[a*Sec[x]^2]*(1225*Sin[x] + 245*Sin[3*x] + 49*Sin[5*x] + 5*Sin[7*x]))/(2240*a^4)

fricas [A] time = 0.73, size = 38, normalized size = 0.51

$$\frac{(5 \cos(x)^7 + 6 \cos(x)^5 + 8 \cos(x)^3 + 16 \cos(x)) \sqrt{\frac{a}{\cos(x)^2}} \sin(x)}{35a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(7/2), x, algorithm="fricas")

[Out] 1/35*(5*cos(x)^7 + 6*cos(x)^5 + 8*cos(x)^3 + 16*cos(x))*sqrt(a/cos(x)^2)*sin(x)/a^4

giac [A] time = 0.70, size = 110, normalized size = 1.49

$$\frac{2 \left(35 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^6 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) - 140 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^4 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) + 336 \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^2 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) - 320 \operatorname{sgn}\left(-\tan\left(\frac{1}{2}x\right)^2 + 1\right) \right)}{35 a^{\frac{7}{2}} \left(\frac{1}{\tan\left(\frac{1}{2}x\right)} + \tan\left(\frac{1}{2}x\right) \right)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(7/2),x, algorithm="giac")

[Out] 2/35*(35*(1/tan(1/2*x) + tan(1/2*x))^6*sgn(-tan(1/2*x)^2 + 1) - 140*(1/tan(1/2*x) + tan(1/2*x))^4*sgn(-tan(1/2*x)^2 + 1) + 336*(1/tan(1/2*x) + tan(1/2*x))^2*sgn(-tan(1/2*x)^2 + 1) - 320*sgn(-tan(1/2*x)^2 + 1))/(a^(7/2)*(1/tan(1/2*x) + tan(1/2*x))^7)

maple [A] time = 0.34, size = 37, normalized size = 0.50

$$\frac{\sin(x) \left(5 \cos^6(x) + 6 \cos^4(x) + 8 \cos^2(x) + 16 \right)}{35 \cos(x)^7 \left(\frac{a}{\cos(x)^2} \right)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^2)^(7/2),x)

[Out] 1/35*sin(x)*(5*cos(x)^6+6*cos(x)^4+8*cos(x)^2+16)/cos(x)^7/(a/cos(x)^2)^(7/2)

maxima [A] time = 0.56, size = 28, normalized size = 0.38

$$\frac{5 \sin(7x) + 49 \sin(5x) + 245 \sin(3x) + 1225 \sin(x)}{2240 a^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^2)^(7/2),x, algorithm="maxima")

[Out] 1/2240*(5*sin(7*x) + 49*sin(5*x) + 245*sin(3*x) + 1225*sin(x))/a^(7/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{a}{\cos(x)^2} \right)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a/cos(x)^2)^(7/2), x)`

[Out] `int(1/(a/cos(x)^2)^(7/2), x)`

sympy [A] time = 150.57, size = 80, normalized size = 1.08

$$\frac{16 \tan^7(x)}{35 a^{\frac{7}{2}} (\sec^2(x))^{\frac{7}{2}}} + \frac{8 \tan^5(x)}{5 a^{\frac{7}{2}} (\sec^2(x))^{\frac{7}{2}}} + \frac{2 \tan^3(x)}{a^{\frac{7}{2}} (\sec^2(x))^{\frac{7}{2}}} + \frac{\tan(x)}{a^{\frac{7}{2}} (\sec^2(x))^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)**2)**(7/2), x)`

[Out] `16*tan(x)**7/(35*a**(7/2)*(sec(x)**2)**(7/2)) + 8*tan(x)**5/(5*a**(7/2)*(sec(x)**2)**(7/2)) + 2*tan(x)**3/(a**(7/2)*(sec(x)**2)**(7/2)) + tan(x)/(a**(7/2)*(sec(x)**2)**(7/2))`

3.55 $\int (a \sec^3(x))^{5/2} dx$

Optimal. Leaf size=117

$$\frac{154}{585}a^2 \tan(x)\sqrt{a \sec^3(x)} + \frac{2}{13}a^2 \tan(x) \sec^4(x)\sqrt{a \sec^3(x)} + \frac{22}{117}a^2 \tan(x) \sec^2(x)\sqrt{a \sec^3(x)} - \frac{154}{195}a^2 \cos^{\frac{3}{2}}(x)E\left(\frac{x}{2}\right)$$

[Out] $-154/195*a^2*\cos(x)^{(3/2)}*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\text{EllipticE}(\sin(1/2*x), 2^{(1/2)})*(a*\sec(x)^3)^{(1/2)}+154/195*a^2*\cos(x)*\sin(x)*(a*\sec(x)^3)^{(1/2)}+154/585*a^2*(a*\sec(x)^3)^{(1/2)}*\tan(x)+22/117*a^2*\sec(x)^2*(a*\sec(x)^3)^{(1/2)}*\tan(x)+2/13*a^2*\sec(x)^4*(a*\sec(x)^3)^{(1/2)}*\tan(x)$

Rubi [A] time = 0.05, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4123, 3768, 3771, 2639}

$$\frac{2}{13}a^2 \tan(x) \sec^4(x)\sqrt{a \sec^3(x)} + \frac{22}{117}a^2 \tan(x) \sec^2(x)\sqrt{a \sec^3(x)} + \frac{154}{585}a^2 \tan(x)\sqrt{a \sec^3(x)} - \frac{154}{195}a^2 \cos^{\frac{3}{2}}(x)E\left(\frac{x}{2}\right)$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^3)^(5/2), x]

[Out] $(-154*a^2*\cos[x]^{(3/2)}*\text{EllipticE}[x/2, 2]*\text{Sqrt}[a*\text{Sec}[x]^3])/195 + (154*a^2*\cos[x]*\text{Sqrt}[a*\text{Sec}[x]^3]*\sin[x])/195 + (154*a^2*\text{Sqrt}[a*\text{Sec}[x]^3]*\tan[x])/585 + (22*a^2*\text{Sec}[x]^2*\text{Sqrt}[a*\text{Sec}[x]^3]*\tan[x])/117 + (2*a^2*\text{Sec}[x]^4*\text{Sqrt}[a*\text{Sec}[x]^3]*\tan[x])/13$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> -Simp[(b*Csc[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rule 4123

```
Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] :> Dist[(b
^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPar
t[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] &
& !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
\int (a \sec^3(x))^{5/2} dx &= \frac{(a^2 \sqrt{a \sec^3(x)}) \int \sec^{\frac{15}{2}}(x) dx}{\sec^{\frac{3}{2}}(x)} \\
&= \frac{2}{13} a^2 \sec^4(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{(11a^2 \sqrt{a \sec^3(x)}) \int \sec^{\frac{11}{2}}(x) dx}{13 \sec^{\frac{3}{2}}(x)} \\
&= \frac{22}{117} a^2 \sec^2(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{2}{13} a^2 \sec^4(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{(77a^2 \sqrt{a \sec^3(x)}) \int \sec^{\frac{7}{2}}(x) dx}{117 \sec^{\frac{3}{2}}(x)} \\
&= \frac{154}{585} a^2 \sqrt{a \sec^3(x)} \tan(x) + \frac{22}{117} a^2 \sec^2(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{2}{13} a^2 \sec^4(x) \sqrt{a \sec^3(x)} \tan(x) \\
&= \frac{154}{195} a^2 \cos(x) \sqrt{a \sec^3(x)} \sin(x) + \frac{154}{585} a^2 \sqrt{a \sec^3(x)} \tan(x) + \frac{22}{117} a^2 \sec^2(x) \sqrt{a \sec^3(x)} \tan(x) \\
&= \frac{154}{195} a^2 \cos(x) \sqrt{a \sec^3(x)} \sin(x) + \frac{154}{585} a^2 \sqrt{a \sec^3(x)} \tan(x) + \frac{22}{117} a^2 \sec^2(x) \sqrt{a \sec^3(x)} \tan(x) \\
&= -\frac{154}{195} a^2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)} + \frac{154}{195} a^2 \cos(x) \sqrt{a \sec^3(x)} \sin(x) + \frac{154}{585} a^2 \sqrt{a \sec^3(x)} \tan(x)
\end{aligned}$$

Mathematica [A] time = 0.10, size = 59, normalized size = 0.50

$$-\frac{2}{585} a \sec(x) (a \sec^3(x))^{3/2} \left(-45 \tan(x) + 231 \cos^{\frac{11}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) - 231 \sin(x) \cos^5(x) - 77 \sin(x) \cos^3(x) - 55 \sin(x) \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[(a*Sec[x]^3)^(5/2), x]
```

```
[Out] (-2*a*Sec[x]*(a*Sec[x]^3)^(3/2)*(231*Cos[x]^(11/2)*EllipticE[x/2, 2] - 55*Cos[x]*Sin[x] - 77*Cos[x]^3*Ssin[x] - 231*Cos[x]^5*Ssin[x] - 45*Tan[x]))/585
```

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{a \sec(x)^3} a^2 \sec(x)^6, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(a*sec(x)^3)*a^2*sec(x)^6, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a \sec(x)^3\right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(5/2),x, algorithm="giac")

[Out] integrate((a*sec(x)^3)^(5/2), x)

maple [C] time = 0.76, size = 223, normalized size = 1.91

$$2(\cos(x)+1)^2(-1+\cos(x))^2\left(231i(\cos^7(x))\sin(x)\sqrt{\frac{1}{\cos(x)+1}}\sqrt{\frac{\cos(x)}{\cos(x)+1}}\text{EllipticE}\left(\frac{i(-1+\cos(x))}{\sin(x)},i\right)-231i(\cos^7(x))\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^3)^(5/2),x)

[Out] 2/585*(cos(x)+1)^2*(-1+cos(x))^2*(231*I*cos(x)^7*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(-1+cos(x))/sin(x),I)-231*I*cos(x)^7*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x),I)+231*I*cos(x)^6*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticE(I*(-1+cos(x))/sin(x),I)-231*I*cos(x)^6*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x),I)-231*cos(x)^7+154*cos(x)^6+22*cos(x)^4+10*cos(x)^2+45)*cos(x)*(a/cos(x)^3)^(5/2)/sin(x)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a \sec(x)^3\right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(5/2),x, algorithm="maxima")

[Out] integrate((a*sec(x)^3)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{a}{\cos(x)^3} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^3)^(5/2),x)

[Out] int((a/cos(x)^3)^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec^3(x))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**3)**(5/2),x)

[Out] Integral((a*sec(x)**3)**(5/2), x)

3.56 $\int (a \sec^3(x))^{3/2} dx$

Optimal. Leaf size=65

$$\frac{10}{21}a \sin(x)\sqrt{a \sec^3(x)} + \frac{2}{7}a \tan(x) \sec(x)\sqrt{a \sec^3(x)} + \frac{10}{21}a \cos^{\frac{3}{2}}(x)F\left(\frac{x}{2}\middle|2\right)\sqrt{a \sec^3(x)}$$

[Out] 10/21*a*cos(x)^(3/2)*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x),2^(1/2))*(a*sec(x)^3)^(1/2)+10/21*a*sin(x)*(a*sec(x)^3)^(1/2)+2/7*a*sec(x)*(a*sec(x)^3)^(1/2)*tan(x)

Rubi [A] time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4123, 3768, 3771, 2641}

$$\frac{10}{21}a \sin(x)\sqrt{a \sec^3(x)} + \frac{2}{7}a \tan(x) \sec(x)\sqrt{a \sec^3(x)} + \frac{10}{21}a \cos^{\frac{3}{2}}(x)F\left(\frac{x}{2}\middle|2\right)\sqrt{a \sec^3(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^3)^(3/2),x]

[Out] (10*a*cos(x)^(3/2)*EllipticF[x/2, 2]*Sqrt[a*Sec[x]^3])/21 + (10*a*Sqrt[a*Sec[x]^3]*Sin[x])/21 + (2*a*Sec[x]*Sqrt[a*Sec[x]^3]*Tan[x])/7

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := -Simp[(b*cos[c + d*x])*(b*csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*csc[c + d*x])^n*sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rule 4123

```
Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b
^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPar
t[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] &
& !IntegerQ[p]
```

Rubi steps

$$\begin{aligned}
 \int (a \sec^3(x))^{3/2} dx &= \frac{(a\sqrt{a \sec^3(x)}) \int \sec^2(x) dx}{\sec^{\frac{3}{2}}(x)} \\
 &= \frac{2}{7} a \sec(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{(5a\sqrt{a \sec^3(x)}) \int \sec^{\frac{5}{2}}(x) dx}{7 \sec^{\frac{3}{2}}(x)} \\
 &= \frac{10}{21} a \sqrt{a \sec^3(x)} \sin(x) + \frac{2}{7} a \sec(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{(5a\sqrt{a \sec^3(x)}) \int \sqrt{\sec(x)} dx}{21 \sec^{\frac{3}{2}}(x)} \\
 &= \frac{10}{21} a \sqrt{a \sec^3(x)} \sin(x) + \frac{2}{7} a \sec(x) \sqrt{a \sec^3(x)} \tan(x) + \frac{1}{21} \left(5a \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)} \right) \int \frac{1}{\sqrt{\cos(x)}} dx \\
 &= \frac{10}{21} a \cos^{\frac{3}{2}}(x) F\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)} + \frac{10}{21} a \sqrt{a \sec^3(x)} \sin(x) + \frac{2}{7} a \sec(x) \sqrt{a \sec^3(x)} \tan(x)
 \end{aligned}$$

Mathematica [A] time = 0.04, size = 43, normalized size = 0.66

$$\frac{2}{21} a \sec(x) \sqrt{a \sec^3(x)} \left(3 \tan(x) + 5 \cos^{\frac{5}{2}}(x) F\left(\frac{x}{2} \middle| 2\right) + 5 \sin(x) \cos(x) \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[(a*Sec[x]^3)^(3/2), x]
```

```
[Out] (2*a*Sec[x]*Sqrt[a*Sec[x]^3]*(5*Cos[x]^(5/2)*EllipticF[x/2, 2] + 5*Cos[x]*Sin[x] + 3*Tan[x]))/21
```

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{a \sec(x)^3} a \sec(x)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sec(x)^3)^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(a*sec(x)^3)*a*sec(x)^3, x)
```


giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a \sec(x)^3 \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(3/2),x, algorithm="giac")

[Out] integrate((a*sec(x)^3)^(3/2), x)

maple [C] time = 0.55, size = 87, normalized size = 1.34

$$\frac{2(\cos(x)+1)^2(-1+\cos(x)) \left(5i(\cos^3(x)) \sin(x) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) - 5(\cos^3(x)) + \right)}{21 \sin(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^3)^(3/2),x)

[Out] -2/21*(cos(x)+1)^2*(-1+cos(x))*(5*I*cos(x)^3*sin(x)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)*EllipticF(I*(-1+cos(x))/sin(x),I)-5*cos(x)^3+5*cos(x)^2-3*cos(x)+3)*cos(x)*(a/cos(x)^3)^(3/2)/sin(x)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a \sec(x)^3 \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(3/2),x, algorithm="maxima")

[Out] integrate((a*sec(x)^3)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{a}{\cos(x)^3} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^3)^(3/2),x)

[Out] int((a/cos(x)^3)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec^3(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**3)**(3/2),x)

[Out] Integral((a*sec(x)**3)**(3/2), x)

3.57 $\int \sqrt{a \sec^3(x)} dx$

Optimal. Leaf size=42

$$2 \sin(x) \cos(x) \sqrt{a \sec^3(x)} - 2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)}$$

[Out] $-2 \cos(x)^{(3/2)} * (\cos(1/2*x)^2)^{(1/2)} / \cos(1/2*x) * \text{EllipticE}(\sin(1/2*x), 2^{(1/2)}) * (a * \sec(x)^3)^{(1/2)} + 2 * \cos(x) * \sin(x) * (a * \sec(x)^3)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4123, 3768, 3771, 2639}

$$2 \sin(x) \cos(x) \sqrt{a \sec^3(x)} - 2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Sec[x]^3], x]

[Out] $-2 * \text{Cos}[x]^{(3/2)} * \text{EllipticE}[x/2, 2] * \text{Sqrt}[a * \text{Sec}[x]^3] + 2 * \text{Cos}[x] * \text{Sqrt}[a * \text{Sec}[x]^3] * \text{Sin}[x]$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x] * (b*Csc[c + d*x])^(n - 1)) / (d*(n - 1)), x] + Dist[(b^2*(n - 2)) / (n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n * Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b^IntPart[p] * (b*(c*Sec[e + f*x])^n)^FracPart[p]) / (c*Sec[e + f*x])^(n*FracPar

t[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] & & !IntegerQ[p]

Rubi steps

$$\begin{aligned}
 \int \sqrt{a \sec^3(x)} dx &= \frac{\sqrt{a \sec^3(x)} \int \sec^{\frac{3}{2}}(x) dx}{\sec^{\frac{3}{2}}(x)} \\
 &= 2 \cos(x) \sqrt{a \sec^3(x)} \sin(x) - \frac{\sqrt{a \sec^3(x)} \int \frac{1}{\sqrt{\sec(x)}} dx}{\sec^{\frac{3}{2}}(x)} \\
 &= 2 \cos(x) \sqrt{a \sec^3(x)} \sin(x) - \left(\cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)} \right) \int \sqrt{\cos(x)} dx \\
 &= -2 \cos^{\frac{3}{2}}(x) E\left(\frac{x}{2} \middle| 2\right) \sqrt{a \sec^3(x)} + 2 \cos(x) \sqrt{a \sec^3(x)} \sin(x)
 \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 0.76

$$2 \cos(x) \sqrt{a \sec^3(x)} \left(\sin(x) - \sqrt{\cos(x)} E\left(\frac{x}{2} \middle| 2\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Sec[x]^3], x]

[Out] 2*Cos[x]*Sqrt[a*Sec[x]^3]*(-(Sqrt[Cos[x]]*EllipticE[x/2, 2]) + Sin[x])

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{a \sec(x)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(a*sec(x)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \sec(x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(a*sec(x)^3), x)

maple [C] time = 0.71, size = 191, normalized size = 4.55

$$2(\cos(x)+1)^2(-1+\cos(x))^2 \left(i \cos(x) \sin(x) \operatorname{EllipticE}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} - i \cos(x) \sin(x) \operatorname{EllipticE}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^3)^(1/2),x)

[Out] 2*(cos(x)+1)^2*(-1+cos(x))^2*(I*cos(x)*sin(x)*EllipticE(I*(-1+cos(x))/sin(x), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)-I*cos(x)*sin(x)*EllipticF(I*(-1+cos(x))/sin(x), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)+I*sin(x)*EllipticE(I*(-1+cos(x))/sin(x), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)-I*sin(x)*EllipticF(I*(-1+cos(x))/sin(x), I)*(1/(cos(x)+1))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)-cos(x)+1)*cos(x)*(a/cos(x)^3)^(1/2)/sin(x)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \sec(x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^3)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*sec(x)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \sqrt{\frac{a}{\cos(x)^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^3)^(1/2),x)

[Out] int((a/cos(x)^3)^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \sec^3(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sec(x)**3)**(1/2),x)
```

```
[Out] Integral(sqrt(a*sec(x)**3), x)
```

$$3.58 \quad \int \frac{1}{\sqrt{a \sec^3(x)}} dx$$

Optimal. Leaf size=44

$$\frac{2 \tan(x)}{3\sqrt{a \sec^3(x)}} + \frac{2F\left(\frac{x}{2} \middle| 2\right)}{3 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}}$$

[Out] $2/3 * (\cos(1/2*x)^2)^{(1/2)} / \cos(1/2*x) * \text{EllipticF}(\sin(1/2*x), 2^{(1/2)}) / \cos(x)^{(3/2)} / (a * \sec(x)^3)^{(1/2)} + 2/3 * \tan(x) / (a * \sec(x)^3)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4123, 3769, 3771, 2641}

$$\frac{2 \tan(x)}{3\sqrt{a \sec^3(x)}} + \frac{2F\left(\frac{x}{2} \middle| 2\right)}{3 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Sec[x]^3],x]

[Out] $(2 * \text{EllipticF}[x/2, 2]) / (3 * \text{Cos}[x]^{(3/2)} * \text{Sqrt}[a * \text{Sec}[x]^3]) + (2 * \text{Tan}[x]) / (3 * \text{Sqrt}[a * \text{Sec}[x]^3])$

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n * Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rule 4123

```
Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b
^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPar
t[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] &
& !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a \sec^3(x)}} dx &= \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx}{\sqrt{a \sec^3(x)}} \\ &= \frac{2 \tan(x)}{3\sqrt{a \sec^3(x)}} + \frac{\sec^{\frac{3}{2}}(x) \int \sqrt{\sec(x)} dx}{3\sqrt{a \sec^3(x)}} \\ &= \frac{2 \tan(x)}{3\sqrt{a \sec^3(x)}} + \frac{\int \frac{1}{\sqrt{\cos(x)}} dx}{3 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} \\ &= \frac{2F\left(\frac{x}{2} \middle| 2\right)}{3 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} + \frac{2 \tan(x)}{3\sqrt{a \sec^3(x)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 31, normalized size = 0.70

$$\frac{2 \left(\tan(x) + \frac{F\left(\frac{x}{2} \middle| 2\right)}{\cos^{\frac{3}{2}}(x)} \right)}{3\sqrt{a \sec^3(x)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/Sqrt[a*Sec[x]^3], x]
```

```
[Out] (2*(EllipticF[x/2, 2]/Cos[x]^(3/2) + Tan[x]))/(3*Sqrt[a*Sec[x]^3])
```

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{a \sec(x)^3}}{a \sec(x)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*sec(x)^3)^(1/2), x, algorithm="fricas")
```


[Out] integral(sqrt(a*sec(x)^3)/(a*sec(x)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \sec(x)^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(a*sec(x)^3), x)

maple [C] time = 0.58, size = 76, normalized size = 1.73

$$\frac{2(-1 + \cos(x)) \left(-i \sin(x) \operatorname{EllipticF} \left(\frac{i(-1 + \cos(x))}{\sin(x)}, i \right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} + \cos^2(x) - \cos(x) \right) (\cos(x) + 1)^2}{3 \cos(x)^2 \sin(x)^3 \sqrt{\frac{a}{\cos(x)^3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^3)^(1/2),x)

[Out] 2/3*(-1+cos(x))*(-I*sin(x)*EllipticF(I*(-1+cos(x))/sin(x),I)*(1/(cos(x)+1)))^(1/2)*(cos(x)/(cos(x)+1))^(1/2)+cos(x)^2-cos(x))*(cos(x)+1)^2/cos(x)^2/sin(x)^3/(a/cos(x)^3)^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \sec(x)^3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(a*sec(x)^3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{\frac{a}{\cos(x)^3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a/cos(x)^3)^(1/2),x)

```
[Out] int(1/(a/cos(x)^3)^(1/2), x)
```

```
sympy [F] time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{1}{\sqrt{a \sec^3(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(a*sec(x)**3)**(1/2), x)
```

```
[Out] Integral(1/sqrt(a*sec(x)**3), x)
```

$$3.59 \quad \int \frac{1}{(a \sec^3(x))^{3/2}} dx$$

Optimal. Leaf size=73

$$\frac{14 \sin(x)}{45a\sqrt{a \sec^3(x)}} + \frac{14E\left(\frac{x}{2} \middle| 2\right)}{15a \cos^{\frac{3}{2}}(x)\sqrt{a \sec^3(x)}} + \frac{2 \sin(x) \cos^2(x)}{9a\sqrt{a \sec^3(x)}}$$

[Out] $14/15*(\cos(1/2*x)^2)^{(1/2)}/\cos(1/2*x)*\text{EllipticE}(\sin(1/2*x), 2^{(1/2)})/a/\cos(x)^{(3/2)}/(a*\sec(x)^3)^{(1/2)}+14/45*\sin(x)/a/(a*\sec(x)^3)^{(1/2)}+2/9*\cos(x)^2*\sin(x)/a/(a*\sec(x)^3)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4123, 3769, 3771, 2639}

$$\frac{14 \sin(x)}{45a\sqrt{a \sec^3(x)}} + \frac{2 \sin(x) \cos^2(x)}{9a\sqrt{a \sec^3(x)}} + \frac{14E\left(\frac{x}{2} \middle| 2\right)}{15a \cos^{\frac{3}{2}}(x)\sqrt{a \sec^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^3)^(-3/2), x]

[Out] $(14*\text{EllipticE}[x/2, 2])/(15*a*\text{Cos}[x]^{(3/2)}*\text{Sqrt}[a*\text{Sec}[x]^3]) + (14*\text{Sin}[x])/(45*a*\text{Sqrt}[a*\text{Sec}[x]^3]) + (2*\text{Cos}[x]^2*\text{Sin}[x])/(9*a*\text{Sqrt}[a*\text{Sec}[x]^3])$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.))^(p_), x_Symbol] := Dist[(b
^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPar
t[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] &
& !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \frac{1}{(a \sec^3(x))^{3/2}} dx &= \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\sec^{\frac{2}{3}}(x)} dx}{a\sqrt{a} \sec^3(x)} \\ &= \frac{2 \cos^2(x) \sin(x)}{9a\sqrt{a} \sec^3(x)} + \frac{\left(7 \sec^{\frac{3}{2}}(x)\right) \int \frac{1}{\sec^{\frac{2}{3}}(x)} dx}{9a\sqrt{a} \sec^3(x)} \\ &= \frac{14 \sin(x)}{45a\sqrt{a} \sec^3(x)} + \frac{2 \cos^2(x) \sin(x)}{9a\sqrt{a} \sec^3(x)} + \frac{\left(7 \sec^{\frac{3}{2}}(x)\right) \int \frac{1}{\sqrt{\sec(x)}} dx}{15a\sqrt{a} \sec^3(x)} \\ &= \frac{14 \sin(x)}{45a\sqrt{a} \sec^3(x)} + \frac{2 \cos^2(x) \sin(x)}{9a\sqrt{a} \sec^3(x)} + \frac{7 \int \sqrt{\cos(x)} dx}{15a \cos^{\frac{3}{2}}(x) \sqrt{a} \sec^3(x)} \\ &= \frac{14E\left(\frac{x}{2} \middle| 2\right)}{15a \cos^{\frac{3}{2}}(x) \sqrt{a} \sec^3(x)} + \frac{14 \sin(x)}{45a\sqrt{a} \sec^3(x)} + \frac{2 \cos^2(x) \sin(x)}{9a\sqrt{a} \sec^3(x)} \end{aligned}$$

Mathematica [A] time = 0.10, size = 43, normalized size = 0.59

$$\frac{33 \sin(x) + 5 \sin(3x) + \frac{84E\left(\frac{x}{2} \middle| 2\right)}{\cos^{\frac{3}{2}}(x)}}{90a\sqrt{a} \sec^3(x)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^3)^(-3/2), x]

[Out] ((84*EllipticE[x/2, 2])/Cos[x]^(3/2) + 33*Sin[x] + 5*Sin[3*x])/(90*a*Sqrt[a*Sec[x]^3])

fricas [F] time = 0.52, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{a} \sec(x)^3}{a^2 \sec(x)^6}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(a*sec(x)^3)/(a^2*sec(x)^6), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \sec(x)^3)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(3/2),x, algorithm="giac")

[Out] integrate((a*sec(x)^3)^(-3/2), x)

maple [C] time = 0.55, size = 198, normalized size = 2.71

$$\frac{2 \left(5 \left(\cos^6(x) \right) - 21i \cos(x) \sin(x) \operatorname{EllipticF} \left(\frac{i(-1+\cos(x))}{\sin(x)}, i \right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} + 21i \cos(x) \sin(x) \operatorname{EllipticE} \left(\frac{i(-1+\cos(x))}{\sin(x)}, i \right) \sqrt{\frac{1}{\cos(x)+1}} \sqrt{\frac{\cos(x)}{\cos(x)+1}} \right)}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^3)^(3/2),x)

[Out]
$$-2/45*(5*\cos(x)^6-21*I*\cos(x)*\sin(x)*\operatorname{EllipticF}(I*(-1+\cos(x))/\sin(x),I))*(1/(\cos(x)+1))^{1/2}*(\cos(x)/(\cos(x)+1))^{1/2}+21*I*\cos(x)*\sin(x)*\operatorname{EllipticE}(I*(-1+\cos(x))/\sin(x),I)*(1/(\cos(x)+1))^{1/2}*(\cos(x)/(\cos(x)+1))^{1/2}-21*I*\sin(x)*\operatorname{EllipticF}(I*(-1+\cos(x))/\sin(x),I)*(1/(\cos(x)+1))^{1/2}*(\cos(x)/(\cos(x)+1))^{1/2}+21*I*\sin(x)*\operatorname{EllipticE}(I*(-1+\cos(x))/\sin(x),I)*(1/(\cos(x)+1))^{1/2}*(\cos(x)/(\cos(x)+1))^{1/2}+2*\cos(x)^4+14*\cos(x)^2-21*\cos(x))/\cos(x)^5/\sin(x)/(a/\cos(x)^3)^{3/2}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \sec(x)^3)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(3/2),x, algorithm="maxima")

[Out] integrate((a*sec(x)^3)^(-3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{a}{\cos(x)^3}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a/cos(x)^3)^(3/2), x)

[Out] int(1/(a/cos(x)^3)^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(a \sec^3(x)\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)**3)**(3/2), x)

[Out] Integral((a*sec(x)**3)**(-3/2), x)

$$3.60 \quad \int \frac{1}{(a \sec^3(x))^{5/2}} dx$$

Optimal. Leaf size=117

$$\frac{26 \tan(x)}{77a^2 \sqrt{a \sec^3(x)}} + \frac{26F\left(\frac{x}{2} \middle| 2\right)}{77a^2 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} + \frac{2 \sin(x) \cos^5(x)}{15a^2 \sqrt{a \sec^3(x)}} + \frac{26 \sin(x) \cos^3(x)}{165a^2 \sqrt{a \sec^3(x)}} + \frac{78 \sin(x) \cos(x)}{385a^2 \sqrt{a \sec^3(x)}}$$

[Out] 26/77*(cos(1/2*x)^2)^(1/2)/cos(1/2*x)*EllipticF(sin(1/2*x),2^(1/2))/a^2/cos(x)^(3/2)/(a*sec(x)^3)^(1/2)+78/385*cos(x)*sin(x)/a^2/(a*sec(x)^3)^(1/2)+26/165*cos(x)^3*sin(x)/a^2/(a*sec(x)^3)^(1/2)+2/15*cos(x)^5*sin(x)/a^2/(a*sec(x)^3)^(1/2)+26/77*tan(x)/a^2/(a*sec(x)^3)^(1/2)

Rubi [A] time = 0.06, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 4, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {4123, 3769, 3771, 2641}

$$\frac{26 \tan(x)}{77a^2 \sqrt{a \sec^3(x)}} + \frac{2 \sin(x) \cos^5(x)}{15a^2 \sqrt{a \sec^3(x)}} + \frac{26 \sin(x) \cos^3(x)}{165a^2 \sqrt{a \sec^3(x)}} + \frac{26F\left(\frac{x}{2} \middle| 2\right)}{77a^2 \cos^{\frac{3}{2}}(x) \sqrt{a \sec^3(x)}} + \frac{78 \sin(x) \cos(x)}{385a^2 \sqrt{a \sec^3(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^3)^(-5/2),x]

[Out] (26*EllipticF[x/2, 2])/(77*a^2*Cos[x]^(3/2)*Sqrt[a*Sec[x]^3]) + (78*Cos[x]*Sin[x])/(385*a^2*Sqrt[a*Sec[x]^3]) + (26*Cos[x]^3*Sin[x])/(165*a^2*Sqrt[a*Sec[x]^3]) + (2*Cos[x]^5*Sin[x])/(15*a^2*Sqrt[a*Sec[x]^3]) + (26*Tan[x])/(77*a^2*Sqrt[a*Sec[x]^3])

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_.)])^(n_))^(p_), x_Symbol] :> Dist[(b^IntPart[p]*(c*Sec[e + f*x])^n)^FracPart[p]]/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] & !IntegerQ[p]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a \sec^3(x))^{5/2}} dx &= \frac{\sec^{\frac{3}{2}}(x) \int \frac{1}{\sec^{\frac{15}{2}}(x)} dx}{a^2 \sqrt{a} \sec^3(x)} \\
 &= \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a} \sec^3(x)} + \frac{\left(13 \sec^{\frac{3}{2}}(x)\right) \int \frac{1}{\sec^{\frac{11}{2}}(x)} dx}{15a^2 \sqrt{a} \sec^3(x)} \\
 &= \frac{26 \cos^3(x) \sin(x)}{165a^2 \sqrt{a} \sec^3(x)} + \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a} \sec^3(x)} + \frac{\left(39 \sec^{\frac{3}{2}}(x)\right) \int \frac{1}{\sec^{\frac{7}{2}}(x)} dx}{55a^2 \sqrt{a} \sec^3(x)} \\
 &= \frac{78 \cos(x) \sin(x)}{385a^2 \sqrt{a} \sec^3(x)} + \frac{26 \cos^3(x) \sin(x)}{165a^2 \sqrt{a} \sec^3(x)} + \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a} \sec^3(x)} + \frac{\left(39 \sec^{\frac{3}{2}}(x)\right) \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx}{77a^2 \sqrt{a} \sec^3(x)} \\
 &= \frac{78 \cos(x) \sin(x)}{385a^2 \sqrt{a} \sec^3(x)} + \frac{26 \cos^3(x) \sin(x)}{165a^2 \sqrt{a} \sec^3(x)} + \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a} \sec^3(x)} + \frac{26 \tan(x)}{77a^2 \sqrt{a} \sec^3(x)} + \frac{\left(13 \sec^{\frac{3}{2}}(x)\right) \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx}{77a^2 \sqrt{a} \sec^3(x)} \\
 &= \frac{78 \cos(x) \sin(x)}{385a^2 \sqrt{a} \sec^3(x)} + \frac{26 \cos^3(x) \sin(x)}{165a^2 \sqrt{a} \sec^3(x)} + \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a} \sec^3(x)} + \frac{26 \tan(x)}{77a^2 \sqrt{a} \sec^3(x)} + \frac{13 \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx}{77a^2 \cos^{\frac{3}{2}}(x)} \\
 &= \frac{26F\left(\frac{x}{2} \middle| 2\right)}{77a^2 \cos^{\frac{3}{2}}(x) \sqrt{a} \sec^3(x)} + \frac{78 \cos(x) \sin(x)}{385a^2 \sqrt{a} \sec^3(x)} + \frac{26 \cos^3(x) \sin(x)}{165a^2 \sqrt{a} \sec^3(x)} + \frac{2 \cos^5(x) \sin(x)}{15a^2 \sqrt{a} \sec^3(x)} + \frac{26 \tan(x)}{77a^2 \sqrt{a} \sec^3(x)} + \frac{13 \int \frac{1}{\sec^{\frac{3}{2}}(x)} dx}{77a^2 \cos^{\frac{3}{2}}(x)}
 \end{aligned}$$

Mathematica [A] time = 0.10, size = 59, normalized size = 0.50

$$\frac{\cos(x)\sqrt{a\sec^3(x)}\left(19122\sin(2x) + 4406\sin(4x) + 826\sin(6x) + 77\sin(8x) + 24960\sqrt{\cos(x)}F\left(\frac{x}{2}\middle|2\right)\right)}{73920a^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^3)^(-5/2), x]

[Out] (Cos[x]*Sqrt[a*Sec[x]^3]*(24960*Sqrt[Cos[x]]*EllipticF[x/2, 2] + 19122*Sin[2*x] + 4406*Sin[4*x] + 826*Sin[6*x] + 77*Sin[8*x]))/(73920*a^3)

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{a\sec(x)^3}}{a^3\sec(x)^9}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(a*sec(x)^3)/(a^3*sec(x)^9), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a\sec(x)^3)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^3)^(5/2), x, algorithm="giac")

[Out] integrate((a*sec(x)^3)^(-5/2), x)

maple [C] time = 0.52, size = 114, normalized size = 0.97

$$\frac{2(-1 + \cos(x))\left(77(\cos^8(x)) - 77(\cos^7(x)) + 91(\cos^6(x)) - 91(\cos^5(x)) - 195i\sin(x)\text{EllipticF}\left(\frac{i(-1+\cos(x))}{\sin(x)}, i\right)\right)}{1155\cos(x)^8\sin(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^3)^(5/2), x)

[Out] $2/1155*(-1+\cos(x))*(77*\cos(x)^8-77*\cos(x)^7+91*\cos(x)^6-91*\cos(x)^5-195*I*\sin(x)*\text{EllipticF}(I*(-1+\cos(x))/\sin(x),I)*(1/(\cos(x)+1))^{1/2}*(\cos(x)/(\cos(x)+1))^{1/2}+117*\cos(x)^4-117*\cos(x)^3+195*\cos(x)^2-195*\cos(x))*(\cos(x)+1)^2/\cos(x)^8/\sin(x)^3/(a/\cos(x)^3)^{5/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \sec(x)^3)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)^3)^(5/2),x, algorithm="maxima")`

[Out] `integrate((a*sec(x)^3)^(-5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{a}{\cos(x)^3}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a/cos(x)^3)^(5/2),x)`

[Out] `int(1/(a/cos(x)^3)^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \sec^3(x))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)**3)**(5/2),x)`

[Out] `Integral((a*sec(x)**3)**(-5/2), x)`

3.61 $\int (a \sec^4(x))^{7/2} dx$

Optimal. Leaf size=163

$$a^3 \sin(x) \cos(x) \sqrt{a \sec^4(x)} + \frac{1}{13} a^3 \sin^2(x) \tan^{11}(x) \sqrt{a \sec^4(x)} + \frac{6}{11} a^3 \sin^2(x) \tan^9(x) \sqrt{a \sec^4(x)} + \frac{5}{3} a^3 \sin^2(x) \tan^7(x) \sqrt{a \sec^4(x)}$$

```
[Out] a^3*cos(x)*sin(x)*(a*sec(x)^4)^(1/2)+2*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)+3*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^3+20/7*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^5+5/3*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^7+6/11*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^9+1/13*a^3*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^11
```

Rubi [A] time = 0.04, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {4123, 3767}

$$a^3 \sin(x) \cos(x) \sqrt{a \sec^4(x)} + \frac{1}{13} a^3 \sin^2(x) \tan^{11}(x) \sqrt{a \sec^4(x)} + \frac{6}{11} a^3 \sin^2(x) \tan^9(x) \sqrt{a \sec^4(x)} + \frac{5}{3} a^3 \sin^2(x) \tan^7(x) \sqrt{a \sec^4(x)}$$

Antiderivative was successfully verified.

```
[In] Int[(a*Sec[x]^4)^(7/2), x]
```

```
[Out] a^3*Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x] + 2*a^3*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x] + 3*a^3*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^3 + (20*a^3*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^5)/7 + (5*a^3*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^7)/3 + (6*a^3*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^9)/11 + (a^3*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^11)/13
```

Rule 3767

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rule 4123

```
Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \int (a \sec^4(x))^{7/2} dx &= \left(a^3 \cos^2(x) \sqrt{a \sec^4(x)} \right) \int \sec^{14}(x) dx \\ &= - \left(\left(a^3 \cos^2(x) \sqrt{a \sec^4(x)} \right) \text{Subst} \left(\int (1 + 6x^2 + 15x^4 + 20x^6 + 15x^8 + 6x^{10} + x^{12}) dx, x, - \right. \right. \\ &= a^3 \cos(x) \sqrt{a \sec^4(x)} \sin(x) + 2a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan(x) + 3a^3 \sqrt{a \sec^4(x)} \sin^2(x) \tan^3(x) \end{aligned}$$

Mathematica [A] time = 0.18, size = 54, normalized size = 0.33

$$\frac{\sin(x) \cos(x) (2380 \cos(2x) + 1093 \cos(4x) + 378 \cos(6x) + 92 \cos(8x) + 14 \cos(10x) + \cos(12x) + 2048) (a \sec^4(x))^{7/2}}{6006}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^4)^(7/2), x]

[Out] (Cos[x]*(2048 + 2380*Cos[2*x] + 1093*Cos[4*x] + 378*Cos[6*x] + 92*Cos[8*x] + 14*Cos[10*x] + Cos[12*x])*(a*Sec[x]^4)^(7/2)*Sin[x])/6006

fricas [A] time = 0.68, size = 76, normalized size = 0.47

$$\frac{(1024 a^3 \cos(x)^{12} + 512 a^3 \cos(x)^{10} + 384 a^3 \cos(x)^8 + 320 a^3 \cos(x)^6 + 280 a^3 \cos(x)^4 + 252 a^3 \cos(x)^2 + 231 a^3) \sin(x)}{3003 \cos(x)^{11}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(7/2), x, algorithm="fricas")

[Out] 1/3003*(1024*a^3*cos(x)^12 + 512*a^3*cos(x)^10 + 384*a^3*cos(x)^8 + 320*a^3*cos(x)^6 + 280*a^3*cos(x)^4 + 252*a^3*cos(x)^2 + 231*a^3)*sqrt(a/cos(x)^4)*sin(x)/cos(x)^11

giac [A] time = 0.41, size = 67, normalized size = 0.41

$$\frac{1}{3003} (231 a^3 \tan(x)^{13} + 1638 a^3 \tan(x)^{11} + 5005 a^3 \tan(x)^9 + 8580 a^3 \tan(x)^7 + 9009 a^3 \tan(x)^5 + 6006 a^3 \tan(x)^3 + 3003 a^3 \tan(x)) \sqrt{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(7/2), x, algorithm="giac")

[Out] 1/3003*(231*a^3*tan(x)^13 + 1638*a^3*tan(x)^11 + 5005*a^3*tan(x)^9 + 8580*a^3*tan(x)^7 + 9009*a^3*tan(x)^5 + 6006*a^3*tan(x)^3 + 3003*a^3*tan(x))*sqrt(a)

maple [A] time = 0.56, size = 53, normalized size = 0.33

$$\frac{(1024(\cos^{12}(x)) + 512(\cos^{10}(x)) + 384(\cos^8(x)) + 320(\cos^6(x)) + 280(\cos^4(x)) + 252(\cos^2(x)) + 231)\left(\frac{a}{\cos(x)}\right)^{7/2}}{3003}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^4)^(7/2),x)

[Out] 1/3003*(1024*cos(x)^12+512*cos(x)^10+384*cos(x)^8+320*cos(x)^6+280*cos(x)^4+252*cos(x)^2+231)*(a/cos(x)^4)^(7/2)*sin(x)*cos(x)

maxima [A] time = 1.14, size = 61, normalized size = 0.37

$$\frac{1}{13} a^{7/2} \tan(x)^{13} + \frac{6}{11} a^{7/2} \tan(x)^{11} + \frac{5}{3} a^{7/2} \tan(x)^9 + \frac{20}{7} a^{7/2} \tan(x)^7 + 3 a^{7/2} \tan(x)^5 + 2 a^{7/2} \tan(x)^3 + a^{7/2} \tan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(7/2),x, algorithm="maxima")

[Out] 1/13*a^(7/2)*tan(x)^13 + 6/11*a^(7/2)*tan(x)^11 + 5/3*a^(7/2)*tan(x)^9 + 20/7*a^(7/2)*tan(x)^7 + 3*a^(7/2)*tan(x)^5 + 2*a^(7/2)*tan(x)^3 + a^(7/2)*tan(x)

mupad [B] time = 4.68, size = 589, normalized size = 3.61

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^4)^(7/2),x)

[Out] (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*2048i)/(7*(exp(x*2i) + 1)^7*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*1536i)/((exp(x*2i) + 1)^8*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*10240i)/(3*(exp(x*2i) + 1)^9*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*4096i)/((exp(x*2i) + 1)^10*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*30720i)/(11*(exp(x*2i) + 1)^11*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) - (a^3*(a/(exp(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i) + exp(x*8i) + 1)*1024i

```
)/((exp(x*2i) + 1)^12*(exp(x*2i) + 2*exp(x*4i) + exp(x*6i))) + (a^3*(a/(exp
(-x*1i)/2 + exp(x*1i)/2)^4)^(1/2)*(4*exp(x*2i) + 6*exp(x*4i) + 4*exp(x*6i)
+ exp(x*8i) + 1)*2048i)/(13*(exp(x*2i) + 1)^13*(exp(x*2i) + 2*exp(x*4i) + e
xp(x*6i)))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sec(x)**4)**(7/2),x)
```

```
[Out] Timed out
```

3.62 $\int (a \sec^4(x))^{5/2} dx$

Optimal. Leaf size=117

$$a^2 \sin(x) \cos(x) \sqrt{a \sec^4(x)} + \frac{1}{9} a^2 \sin^2(x) \tan^7(x) \sqrt{a \sec^4(x)} + \frac{4}{7} a^2 \sin^2(x) \tan^5(x) \sqrt{a \sec^4(x)} + \frac{6}{5} a^2 \sin^2(x) \tan^3(x) \sqrt{a \sec^4(x)}$$

[Out] a^2*cos(x)*sin(x)*(a*sec(x)^4)^(1/2)+4/3*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)+6/5*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^3+4/7*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^5+1/9*a^2*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^7

Rubi [A] time = 0.03, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {4123, 3767}

$$a^2 \sin(x) \cos(x) \sqrt{a \sec^4(x)} + \frac{1}{9} a^2 \sin^2(x) \tan^7(x) \sqrt{a \sec^4(x)} + \frac{4}{7} a^2 \sin^2(x) \tan^5(x) \sqrt{a \sec^4(x)} + \frac{6}{5} a^2 \sin^2(x) \tan^3(x) \sqrt{a \sec^4(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^4)^(5/2), x]

[Out] a^2*Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x] + (4*a^2*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x])/3 + (6*a^2*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^3)/5 + (4*a^2*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^5)/7 + (a^2*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^7)/9

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int (a \sec^4(x))^{5/2} dx &= \left(a^2 \cos^2(x) \sqrt{a \sec^4(x)} \right) \int \sec^{10}(x) dx \\
&= - \left(\left(a^2 \cos^2(x) \sqrt{a \sec^4(x)} \right) \text{Subst} \left(\int (1 + 4x^2 + 6x^4 + 4x^6 + x^8) dx, x, -\tan(x) \right) \right) \\
&= a^2 \cos(x) \sqrt{a \sec^4(x)} \sin(x) + \frac{4}{3} a^2 \sqrt{a \sec^4(x)} \sin^2(x) \tan(x) + \frac{6}{5} a^2 \sqrt{a \sec^4(x)} \sin^2(x) \tan^3(x)
\end{aligned}$$

Mathematica [A] time = 0.10, size = 42, normalized size = 0.36

$$\frac{1}{315} \sin(x) \cos(x) (130 \cos(2x) + 46 \cos(4x) + 10 \cos(6x) + \cos(8x) + 128) (a \sec^4(x))^{5/2}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^4)^(5/2),x]

[Out] (Cos[x]*(128 + 130*Cos[2*x] + 46*Cos[4*x] + 10*Cos[6*x] + Cos[8*x])*(a*Sec[x]^4)^(5/2)*Sin[x])/315

fricas [A] time = 0.63, size = 58, normalized size = 0.50

$$\frac{(128 a^2 \cos(x)^8 + 64 a^2 \cos(x)^6 + 48 a^2 \cos(x)^4 + 40 a^2 \cos(x)^2 + 35 a^2) \sqrt{\frac{a}{\cos(x)^4}} \sin(x)}{315 \cos(x)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(5/2),x, algorithm="fricas")

[Out] 1/315*(128*a^2*cos(x)^8 + 64*a^2*cos(x)^6 + 48*a^2*cos(x)^4 + 40*a^2*cos(x)^2 + 35*a^2)*sqrt(a/cos(x)^4)*sin(x)/cos(x)^7

giac [A] time = 0.90, size = 49, normalized size = 0.42

$$\frac{1}{315} (35 a^2 \tan(x)^9 + 180 a^2 \tan(x)^7 + 378 a^2 \tan(x)^5 + 420 a^2 \tan(x)^3 + 315 a^2 \tan(x)) \sqrt{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(5/2),x, algorithm="giac")

[Out] 1/315*(35*a^2*tan(x)^9 + 180*a^2*tan(x)^7 + 378*a^2*tan(x)^5 + 420*a^2*tan(x)^3 + 315*a^2*tan(x))*sqrt(a)

maple [A] time = 0.36, size = 41, normalized size = 0.35

$$\frac{(128(\cos^8(x)) + 64(\cos^6(x)) + 48(\cos^4(x)) + 40(\cos^2(x)) + 35)\cos(x)\sin(x)\left(\frac{a}{\cos(x)^4}\right)^{\frac{5}{2}}}{315}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^4)^(5/2), x)

[Out] 1/315*(128*cos(x)^8+64*cos(x)^6+48*cos(x)^4+40*cos(x)^2+35)*cos(x)*sin(x)*(a/cos(x)^4)^(5/2)

maxima [A] time = 0.75, size = 43, normalized size = 0.37

$$\frac{1}{9}a^{\frac{5}{2}}\tan(x)^9 + \frac{4}{7}a^{\frac{5}{2}}\tan(x)^7 + \frac{6}{5}a^{\frac{5}{2}}\tan(x)^5 + \frac{4}{3}a^{\frac{5}{2}}\tan(x)^3 + a^{\frac{5}{2}}\tan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(5/2), x, algorithm="maxima")

[Out] 1/9*a^(5/2)*tan(x)^9 + 4/7*a^(5/2)*tan(x)^7 + 6/5*a^(5/2)*tan(x)^5 + 4/3*a^(5/2)*tan(x)^3 + a^(5/2)*tan(x)

mupad [B] time = 2.35, size = 119, normalized size = 1.02

$$\frac{128a^{5/2}\left(e^{x46i}1i + e^{x48i}9i + e^{x50i}36i + e^{x52i}84i + e^{x54i}126i\right)}{315\left(\frac{e^{-x2i}}{2} + \frac{e^{x2i}}{2} + 1\right)\left(e^{x48i} + 7e^{x50i} + 21e^{x52i} + 35e^{x54i} + 35e^{x56i} + 21e^{x58i} + 7e^{x60i} + e^{x62i}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^4)^(5/2), x)

[Out] (128*a^(5/2)*(exp(x*46i)*1i + exp(x*48i)*9i + exp(x*50i)*36i + exp(x*52i)*84i + exp(x*54i)*126i))/(315*(exp(-x*2i)/2 + exp(x*2i)/2 + 1)*(exp(x*48i) + 7*exp(x*50i) + 21*exp(x*52i) + 35*exp(x*54i) + 35*exp(x*56i) + 21*exp(x*58i) + 7*exp(x*60i) + exp(x*62i)))

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a \sec^4(x)\right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**4)**(5/2), x)

[Out] Integral((a*sec(x)**4)**(5/2), x)

3.63 $\int (a \sec^4(x))^{3/2} dx$

Optimal. Leaf size=61

$$a \sin(x) \cos(x) \sqrt{a \sec^4(x)} + \frac{1}{5} a \sin^2(x) \tan^3(x) \sqrt{a \sec^4(x)} + \frac{2}{3} a \sin^2(x) \tan(x) \sqrt{a \sec^4(x)}$$

[Out] a*cos(x)*sin(x)*(a*sec(x)^4)^(1/2)+2/3*a*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)+1/5*a*sin(x)^2*(a*sec(x)^4)^(1/2)*tan(x)^3

Rubi [A] time = 0.02, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {4123, 3767}

$$a \sin(x) \cos(x) \sqrt{a \sec^4(x)} + \frac{1}{5} a \sin^2(x) \tan^3(x) \sqrt{a \sec^4(x)} + \frac{2}{3} a \sin^2(x) \tan(x) \sqrt{a \sec^4(x)}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^4)^(3/2), x]

[Out] a*Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x] + (2*a*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x])/3 + (a*Sqrt[a*Sec[x]^4]*Sin[x]^2*Tan[x]^3)/5

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b^IntPart[p]*(c*Sec[e + f*x])^n)^FracPart[p]]/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] & !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int (a \sec^4(x))^{3/2} dx &= \left(a \cos^2(x) \sqrt{a \sec^4(x)} \right) \int \sec^6(x) dx \\ &= - \left(\left(a \cos^2(x) \sqrt{a \sec^4(x)} \right) \text{Subst} \left(\int (1 + 2x^2 + x^4) dx, x, -\tan(x) \right) \right) \\ &= a \cos(x) \sqrt{a \sec^4(x)} \sin(x) + \frac{2}{3} a \sqrt{a \sec^4(x)} \sin^2(x) \tan(x) + \frac{1}{5} a \sqrt{a \sec^4(x)} \sin^2(x) \tan^3(x) \end{aligned}$$

Mathematica [A] time = 0.06, size = 30, normalized size = 0.49

$$\frac{1}{15} \sin(x) \cos(x) (6 \cos(2x) + \cos(4x) + 8) (a \sec^4(x))^{3/2}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^4)^(3/2), x]

[Out] (Cos[x]*(8 + 6*Cos[2*x] + Cos[4*x])*(a*Sec[x]^4)^(3/2)*Sin[x])/15

fricas [A] time = 0.60, size = 34, normalized size = 0.56

$$\frac{(8a \cos(x)^4 + 4a \cos(x)^2 + 3a) \sqrt{\frac{a}{\cos(x)^4}} \sin(x)}{15 \cos(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(3/2), x, algorithm="fricas")

[Out] 1/15*(8*a*cos(x)^4 + 4*a*cos(x)^2 + 3*a)*sqrt(a/cos(x)^4)*sin(x)/cos(x)^3

giac [A] time = 0.74, size = 22, normalized size = 0.36

$$\frac{1}{15} (3 \tan(x)^5 + 10 \tan(x)^3 + 15 \tan(x)) a^{3/2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(3/2), x, algorithm="giac")

[Out] 1/15*(3*tan(x)^5 + 10*tan(x)^3 + 15*tan(x))*a^(3/2)

maple [A] time = 0.30, size = 29, normalized size = 0.48

$$\frac{(8(\cos^4(x)) + 4(\cos^2(x)) + 3) \cos(x) \sin(x) \left(\frac{a}{\cos(x)^4}\right)^{3/2}}{15}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^4)^(3/2), x)

[Out] 1/15*(8*cos(x)^4+4*cos(x)^2+3)*cos(x)*sin(x)*(a/cos(x)^4)^(3/2)

maxima [A] time = 0.44, size = 25, normalized size = 0.41

$$\frac{1}{5} a^{3/2} \tan(x)^5 + \frac{2}{3} a^{3/2} \tan(x)^3 + a^{3/2} \tan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(3/2),x, algorithm="maxima")

[Out] $1/5*a^{3/2}*tan(x)^5 + 2/3*a^{3/2}*tan(x)^3 + a^{3/2}*tan(x)$

mupad [B] time = 0.56, size = 36, normalized size = 0.59

$$\frac{4a^{3/2}\sin(x)}{5\cos(x)^3} + \frac{a^{3/2}\sin(x)}{5\cos(x)^5} - \frac{8a^{3/2}\sin(x)^3}{15\cos(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^4)^(3/2),x)

[Out] $(4*a^{3/2}*sin(x))/(5*cos(x)^3) + (a^{3/2}*sin(x))/(5*cos(x)^5) - (8*a^{3/2}*sin(x)^3)/(15*cos(x)^3)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec^4(x))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**4)**(3/2),x)

[Out] Integral((a*sec(x)**4)**(3/2), x)

3.64 $\int \sqrt{a \sec^4(x)} dx$

Optimal. Leaf size=15

$$\sin(x) \cos(x) \sqrt{a \sec^4(x)}$$

[Out] `cos(x)*sin(x)*(a*sec(x)^4)^(1/2)`

Rubi [A] time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4123, 3767, 8}

$$\sin(x) \cos(x) \sqrt{a \sec^4(x)}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[a*Sec[x]^4],x]`

[Out] `Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x]`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 3767

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rule 4123

`Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b ^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]`

Rubi steps

$$\begin{aligned} \int \sqrt{a \sec^4(x)} dx &= \left(\cos^2(x) \sqrt{a \sec^4(x)} \right) \int \sec^2(x) dx \\ &= - \left(\left(\cos^2(x) \sqrt{a \sec^4(x)} \right) \text{Subst} \left(\int 1 dx, x, -\tan(x) \right) \right) \\ &= \cos(x) \sqrt{a \sec^4(x)} \sin(x) \end{aligned}$$

Mathematica [A] time = 0.01, size = 15, normalized size = 1.00

$$\sin(x) \cos(x) \sqrt{a \sec^4(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Sec[x]^4], x]

[Out] Cos[x]*Sqrt[a*Sec[x]^4]*Sin[x]

fricas [A] time = 0.58, size = 13, normalized size = 0.87

$$\sqrt{\frac{a}{\cos(x)^4}} \cos(x) \sin(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(1/2), x, algorithm="fricas")

[Out] sqrt(a/cos(x)^4)*cos(x)*sin(x)

giac [A] time = 0.70, size = 6, normalized size = 0.40

$$\sqrt{a} \tan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(1/2), x, algorithm="giac")

[Out] sqrt(a)*tan(x)

maple [A] time = 0.40, size = 14, normalized size = 0.93

$$\cos(x) \sin(x) \sqrt{\frac{a}{\cos(x)^4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sec(x)^4)^(1/2), x)

[Out] cos(x)*sin(x)*(a/cos(x)^4)^(1/2)

maxima [A] time = 0.70, size = 6, normalized size = 0.40

$$\sqrt{a} \tan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)^4)^(1/2),x, algorithm="maxima")

[Out] sqrt(a)*tan(x)

mupad [B] time = 0.11, size = 6, normalized size = 0.40

$$\sqrt{a} \tan(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(x)^4)^(1/2),x)

[Out] a^(1/2)*tan(x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \sec^4(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sec(x)**4)**(1/2),x)

[Out] Integral(sqrt(a*sec(x)**4), x)

$$3.65 \quad \int \frac{1}{\sqrt{a \sec^4(x)}} dx$$

Optimal. Leaf size=36

$$\frac{x \sec^2(x)}{2\sqrt{a \sec^4(x)}} + \frac{\tan(x)}{2\sqrt{a \sec^4(x)}}$$

[Out] 1/2*x*sec(x)^2/(a*sec(x)^4)^(1/2)+1/2*tan(x)/(a*sec(x)^4)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4123, 2635, 8}

$$\frac{x \sec^2(x)}{2\sqrt{a \sec^4(x)}} + \frac{\tan(x)}{2\sqrt{a \sec^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[a*Sec[x]^4],x]

[Out] (x*Sec[x]^2)/(2*Sqrt[a*Sec[x]^4]) + Tan[x]/(2*Sqrt[a*Sec[x]^4])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x] * (b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b^IntPart[p]*(b*(c*Sec[e + f*x])^n)^FracPart[p])/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{a \sec^4(x)}} dx &= \frac{\sec^2(x) \int \cos^2(x) dx}{\sqrt{a \sec^4(x)}} \\ &= \frac{\tan(x)}{2\sqrt{a \sec^4(x)}} + \frac{\sec^2(x) \int 1 dx}{2\sqrt{a \sec^4(x)}} \\ &= \frac{x \sec^2(x)}{2\sqrt{a \sec^4(x)}} + \frac{\tan(x)}{2\sqrt{a \sec^4(x)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 23, normalized size = 0.64

$$\frac{\tan(x) + x \sec^2(x)}{2\sqrt{a \sec^4(x)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[a*Sec[x]^4],x]

[Out] (x*Sec[x]^2 + Tan[x])/(2*Sqrt[a*Sec[x]^4])

fricas [A] time = 0.68, size = 27, normalized size = 0.75

$$\frac{(\cos(x)^3 \sin(x) + x \cos(x)^2) \sqrt{\frac{a}{\cos(x)^4}}}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^4)^(1/2),x, algorithm="fricas")

[Out] 1/2*(cos(x)^3*sin(x) + x*cos(x)^2)*sqrt(a/cos(x)^4)/a

giac [A] time = 0.45, size = 39, normalized size = 1.08

$$-\frac{1}{2} \sqrt{a} \left(\frac{\pi \left\lfloor \frac{x}{\pi} + \frac{1}{2} \right\rfloor - x}{a} - \frac{\tan(x)}{(\tan(x)^2 + 1)a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^4)^(1/2),x, algorithm="giac")

[Out] -1/2*sqrt(a)*((pi*floor(x/pi + 1/2) - x)/a - tan(x)/((tan(x)^2 + 1)*a))

maple [A] time = 0.41, size = 22, normalized size = 0.61

$$\frac{\cos(x) \sin(x) + x}{2 \cos(x)^2 \sqrt{\frac{a}{\cos(x)^4}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a*sec(x)^4)^(1/2), x)`

[Out] `1/2*(cos(x)*sin(x)+x)/cos(x)^2/(a/cos(x)^4)^(1/2)`

maxima [A] time = 0.65, size = 25, normalized size = 0.69

$$\frac{x}{2\sqrt{a}} + \frac{\tan(x)}{2(\sqrt{a}\tan(x)^2 + \sqrt{a})}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)^4)^(1/2), x, algorithm="maxima")`

[Out] `1/2*x/sqrt(a) + 1/2*tan(x)/(sqrt(a)*tan(x)^2 + sqrt(a))`

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{\frac{a}{\cos(x)^4}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a/cos(x)^4)^(1/2), x)`

[Out] `int(1/(a/cos(x)^4)^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{a \sec^4(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)**4)**(1/2), x)`

[Out] `Integral(1/sqrt(a*sec(x)**4), x)`

$$3.66 \quad \int \frac{1}{(a \sec^4(x))^{3/2}} dx$$

Optimal. Leaf size=86

$$\frac{5x \sec^2(x)}{16a\sqrt{a \sec^4(x)}} + \frac{5 \tan(x)}{16a\sqrt{a \sec^4(x)}} + \frac{\sin(x) \cos^3(x)}{6a\sqrt{a \sec^4(x)}} + \frac{5 \sin(x) \cos(x)}{24a\sqrt{a \sec^4(x)}}$$

[Out] $5/16*x*\sec(x)^2/a/(a*\sec(x)^4)^{(1/2)}+5/24*\cos(x)*\sin(x)/a/(a*\sec(x)^4)^{(1/2)}$
 $+1/6*\cos(x)^3*\sin(x)/a/(a*\sec(x)^4)^{(1/2)}+5/16*\tan(x)/a/(a*\sec(x)^4)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 86, normalized size of antiderivative = 1.00,
 number of steps used = 5, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} =$
 0.300, Rules used = {4123, 2635, 8}

$$\frac{5x \sec^2(x)}{16a\sqrt{a \sec^4(x)}} + \frac{5 \tan(x)}{16a\sqrt{a \sec^4(x)}} + \frac{\sin(x) \cos^3(x)}{6a\sqrt{a \sec^4(x)}} + \frac{5 \sin(x) \cos(x)}{24a\sqrt{a \sec^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^4)^(-3/2), x]

[Out] $(5*x*Sec[x]^2)/(16*a*Sqrt[a*Sec[x]^4]) + (5*Cos[x]*Sin[x])/(24*a*Sqrt[a*Sec[x]^4])$
 $+ (Cos[x]^3*Sin[x])/(6*a*Sqrt[a*Sec[x]^4]) + (5*Tan[x])/(16*a*Sqrt[a*Sec[x]^4])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*
 (b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /;
 FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b
 ^IntPart[p])*(b*(c*Sec[e + f*x])^n)^FracPart[p]/(c*Sec[e + f*x])^(n*FracPart[p]),
 Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] && !IntegerQ[p]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(a \sec^4(x))^{3/2}} dx &= \frac{\sec^2(x) \int \cos^6(x) dx}{a\sqrt{a} \sec^4(x)} \\
&= \frac{\cos^3(x) \sin(x)}{6a\sqrt{a} \sec^4(x)} + \frac{(5 \sec^2(x)) \int \cos^4(x) dx}{6a\sqrt{a} \sec^4(x)} \\
&= \frac{5 \cos(x) \sin(x)}{24a\sqrt{a} \sec^4(x)} + \frac{\cos^3(x) \sin(x)}{6a\sqrt{a} \sec^4(x)} + \frac{(5 \sec^2(x)) \int \cos^2(x) dx}{8a\sqrt{a} \sec^4(x)} \\
&= \frac{5 \cos(x) \sin(x)}{24a\sqrt{a} \sec^4(x)} + \frac{\cos^3(x) \sin(x)}{6a\sqrt{a} \sec^4(x)} + \frac{5 \tan(x)}{16a\sqrt{a} \sec^4(x)} + \frac{(5 \sec^2(x)) \int 1 dx}{16a\sqrt{a} \sec^4(x)} \\
&= \frac{5x \sec^2(x)}{16a\sqrt{a} \sec^4(x)} + \frac{5 \cos(x) \sin(x)}{24a\sqrt{a} \sec^4(x)} + \frac{\cos^3(x) \sin(x)}{6a\sqrt{a} \sec^4(x)} + \frac{5 \tan(x)}{16a\sqrt{a} \sec^4(x)}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 38, normalized size = 0.44

$$\frac{(60x + 45 \sin(2x) + 9 \sin(4x) + \sin(6x)) \sec^6(x)}{192 (a \sec^4(x))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^4)^(-3/2), x]

[Out] (Sec[x]^6*(60*x + 45*Sin[2*x] + 9*Sin[4*x] + Sin[6*x]))/(192*(a*Sec[x]^4)^(-3/2))

fricas [A] time = 0.67, size = 43, normalized size = 0.50

$$\frac{(15x \cos(x)^2 + (8 \cos(x)^7 + 10 \cos(x)^5 + 15 \cos(x)^3) \sin(x)) \sqrt{\frac{a}{\cos(x)^4}}}{48 a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^4)^(3/2), x, algorithm="fricas")

[Out] 1/48*(15*x*cos(x)^2 + (8*cos(x)^7 + 10*cos(x)^5 + 15*cos(x)^3)*sin(x))*sqrt(a/cos(x)^4)/a^2

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)^4)^(3/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,x);OUTPUT:sym2poly/r2sym(const gen & e,const index_m &
i,const vecteur & l) Error: Bad Argument Value

maple [A] time = 0.38, size = 41, normalized size = 0.48

$$\frac{8 \sin(x) \left(\cos^5(x)\right) + 10 \left(\cos^3(x)\right) \sin(x) + 15 \cos(x) \sin(x) + 15x}{48 \cos(x)^6 \left(\frac{a}{\cos(x)^4}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a*sec(x)^4)^(3/2),x)`

[Out] $1/48*(8*\sin(x)*\cos(x)^5+10*\cos(x)^3*\sin(x)+15*\cos(x)*\sin(x)+15*x)/\cos(x)^6/$
 $(a/\cos(x)^4)^(3/2)$

maxima [A] time = 0.69, size = 58, normalized size = 0.67

$$\frac{15 \tan(x)^5 + 40 \tan(x)^3 + 33 \tan(x)}{48 \left(a^{\frac{3}{2}} \tan(x)^6 + 3 a^{\frac{3}{2}} \tan(x)^4 + 3 a^{\frac{3}{2}} \tan(x)^2 + a^{\frac{3}{2}}\right)} + \frac{5x}{16 a^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)^4)^(3/2),x, algorithm="maxima")`

[Out] $1/48*(15*\tan(x)^5 + 40*\tan(x)^3 + 33*\tan(x))/(a^{(3/2)}*\tan(x)^6 + 3*a^{(3/2)}*$
 $\tan(x)^4 + 3*a^{(3/2)}*\tan(x)^2 + a^{(3/2)}) + 5/16*x/a^{(3/2)}$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{a}{\cos(x)^4}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a/cos(x)^4)^(3/2),x)`

[Out] `int(1/(a/cos(x)^4)^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \sec^4(x))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)**4)**(3/2), x)

[Out] Integral((a*sec(x)**4)**(-3/2), x)

$$3.67 \quad \int \frac{1}{(a \sec^4(x))^{5/2}} dx$$

Optimal. Leaf size=132

$$\frac{63x \sec^2(x)}{256a^2 \sqrt{a \sec^4(x)}} + \frac{63 \tan(x)}{256a^2 \sqrt{a \sec^4(x)}} + \frac{\sin(x) \cos^7(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{9 \sin(x) \cos^5(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{21 \sin(x) \cos^3(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{21 \sin(x) \cos(x)}{128a^2 \sqrt{a \sec^4(x)}}$$

[Out] 63/256*x*sec(x)^2/a^2/(a*sec(x)^4)^(1/2)+21/128*cos(x)*sin(x)/a^2/(a*sec(x)^4)^(1/2)+21/160*cos(x)^3*sin(x)/a^2/(a*sec(x)^4)^(1/2)+9/80*cos(x)^5*sin(x)/a^2/(a*sec(x)^4)^(1/2)+1/10*cos(x)^7*sin(x)/a^2/(a*sec(x)^4)^(1/2)+63/256*tan(x)/a^2/(a*sec(x)^4)^(1/2)

Rubi [A] time = 0.05, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {4123, 2635, 8}

$$\frac{63x \sec^2(x)}{256a^2 \sqrt{a \sec^4(x)}} + \frac{63 \tan(x)}{256a^2 \sqrt{a \sec^4(x)}} + \frac{\sin(x) \cos^7(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{9 \sin(x) \cos^5(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{21 \sin(x) \cos^3(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{21 \sin(x) \cos(x)}{128a^2 \sqrt{a \sec^4(x)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sec[x]^4)^(-5/2), x]

[Out] (63*x*Sec[x]^2)/(256*a^2*Sqrt[a*Sec[x]^4]) + (21*Cos[x]*Sin[x])/(128*a^2*Sqrt[a*Sec[x]^4]) + (21*Cos[x]^3*Sin[x])/(160*a^2*Sqrt[a*Sec[x]^4]) + (9*Cos[x]^5*Sin[x])/(80*a^2*Sqrt[a*Sec[x]^4]) + (Cos[x]^7*Sin[x])/(10*a^2*Sqrt[a*Sec[x]^4]) + (63*Tan[x])/(256*a^2*Sqrt[a*Sec[x]^4])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)]/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 4123

Int[((b_.)*((c_.)*sec[(e_.) + (f_.)*(x_)])^(n_))^(p_), x_Symbol] := Dist[(b^IntPart[p])*(b*(c*Sec[e + f*x])^n)^FracPart[p]]/(c*Sec[e + f*x])^(n*FracPart[p]), Int[(c*Sec[e + f*x])^(n*p), x], x] /; FreeQ[{b, c, e, f, n, p}, x] &

& !IntegerQ [p]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(a \sec^4(x))^{5/2}} dx &= \frac{\sec^2(x) \int \cos^{10}(x) dx}{a^2 \sqrt{a \sec^4(x)}} \\
 &= \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{(9 \sec^2(x)) \int \cos^8(x) dx}{10a^2 \sqrt{a \sec^4(x)}} \\
 &= \frac{9 \cos^5(x) \sin(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{(63 \sec^2(x)) \int \cos^6(x) dx}{80a^2 \sqrt{a \sec^4(x)}} \\
 &= \frac{21 \cos^3(x) \sin(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{9 \cos^5(x) \sin(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{(21 \sec^2(x)) \int \cos^4(x) dx}{32a^2 \sqrt{a \sec^4(x)}} \\
 &= \frac{21 \cos(x) \sin(x)}{128a^2 \sqrt{a \sec^4(x)}} + \frac{21 \cos^3(x) \sin(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{9 \cos^5(x) \sin(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{(63 \sec^2(x)) \int \cos^2(x) dx}{128a^2 \sqrt{a \sec^4(x)}} \\
 &= \frac{21 \cos(x) \sin(x)}{128a^2 \sqrt{a \sec^4(x)}} + \frac{21 \cos^3(x) \sin(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{9 \cos^5(x) \sin(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}} + \frac{63 \tan(x)}{256a^2 \sqrt{a \sec^4(x)}} \\
 &= \frac{63x \sec^2(x)}{256a^2 \sqrt{a \sec^4(x)}} + \frac{21 \cos(x) \sin(x)}{128a^2 \sqrt{a \sec^4(x)}} + \frac{21 \cos^3(x) \sin(x)}{160a^2 \sqrt{a \sec^4(x)}} + \frac{9 \cos^5(x) \sin(x)}{80a^2 \sqrt{a \sec^4(x)}} + \frac{\cos^7(x) \sin(x)}{10a^2 \sqrt{a \sec^4(x)}}
 \end{aligned}$$

Mathematica [A] time = 0.09, size = 55, normalized size = 0.42

$$\frac{(2520x + 2100 \sin(2x) + 600 \sin(4x) + 150 \sin(6x) + 25 \sin(8x) + 2 \sin(10x)) \cos^2(x) \sqrt{a \sec^4(x)}}{10240a^3}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sec[x]^4)^(-5/2), x]

[Out] (Cos[x]^2*Sqrt[a*Sec[x]^4]*(2520*x + 2100*Sin[2*x] + 600*Sin[4*x] + 150*Sin[6*x] + 25*Sin[8*x] + 2*Sin[10*x]))/(10240*a^3)

fricas [A] time = 0.67, size = 55, normalized size = 0.42

$$\frac{(315x \cos(x)^2 + (128 \cos(x)^{11} + 144 \cos(x)^9 + 168 \cos(x)^7 + 210 \cos(x)^5 + 315 \cos(x)^3) \sin(x)) \sqrt{\frac{a}{\cos(x)^4}}}{1280a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^4)^(5/2),x, algorithm="fricas")

[Out] 1/1280*(315*x*cos(x)^2 + (128*cos(x)^11 + 144*cos(x)^9 + 168*cos(x)^7 + 210*cos(x)^5 + 315*cos(x)^3)*sin(x))*sqrt(a/cos(x)^4)/a^3

giac [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^4)^(5/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,x);OUTPUT:sym2poly/r2sym(const gen & e,const index_m & i,const vecteur & l) Error: Bad Argument Value

maple [A] time = 0.60, size = 57, normalized size = 0.43

$$\frac{128 \sin(x) (\cos^9(x)) + 144 \sin(x) (\cos^7(x)) + 168 \sin(x) (\cos^5(x)) + 210 (\cos^3(x)) \sin(x) + 315 \cos(x) \sin(x) + 1280 \cos(x)^{10} \left(\frac{a}{\cos(x)^4}\right)^{\frac{5}{2}}}{1280 \cos(x)^{10} \left(\frac{a}{\cos(x)^4}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*sec(x)^4)^(5/2),x)

[Out] 1/1280*(128*sin(x)*cos(x)^9+144*sin(x)*cos(x)^7+168*sin(x)*cos(x)^5+210*cos(x)^3*sin(x)+315*cos(x)*sin(x)+315*x)/cos(x)^10/(a/cos(x)^4)^(5/2)

maxima [A] time = 0.69, size = 88, normalized size = 0.67

$$\frac{315 \tan(x)^9 + 1470 \tan(x)^7 + 2688 \tan(x)^5 + 2370 \tan(x)^3 + 965 \tan(x)}{1280 \left(a^{\frac{5}{2}} \tan(x)^{10} + 5 a^{\frac{5}{2}} \tan(x)^8 + 10 a^{\frac{5}{2}} \tan(x)^6 + 10 a^{\frac{5}{2}} \tan(x)^4 + 5 a^{\frac{5}{2}} \tan(x)^2 + a^{\frac{5}{2}}\right)} + \frac{63 x}{256 a^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(a*sec(x)^4)^(5/2),x, algorithm="maxima")

[Out] 1/1280*(315*tan(x)^9 + 1470*tan(x)^7 + 2688*tan(x)^5 + 2370*tan(x)^3 + 965*tan(x))/(a^(5/2)*tan(x)^10 + 5*a^(5/2)*tan(x)^8 + 10*a^(5/2)*tan(x)^6 + 10*a^(5/2)*tan(x)^4 + 5*a^(5/2)*tan(x)^2 + a^(5/2)) + 63/256*x/a^(5/2)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{a}{\cos(x)^4}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(a/cos(x)^4)^(5/2), x)`

[Out] `int(1/(a/cos(x)^4)^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(a \sec^4(x))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(a*sec(x)**4)**(5/2), x)`

[Out] `Integral((a*sec(x)**4)**(-5/2), x)`

3.68 $\int ((b \sec(c + dx))^p)^n dx$

Optimal. Leaf size=81

$$\frac{\sin(c + dx) \cos(c + dx) ((b \sec(c + dx))^p)^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 - np); \frac{1}{2}(3 - np); \cos^2(c + dx)\right)}{d(1 - np)\sqrt{\sin^2(c + dx)}}$$

[Out] $-\cos(d*x+c)*\text{hypergeom}([1/2, -1/2*n*p+1/2], [-1/2*n*p+3/2], \cos(d*x+c)^2)*((b*\sec(d*x+c))^p)^n*\sin(d*x+c)/d/(-n*p+1)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {4123, 3772, 2643}

$$\frac{\sin(c + dx) \cos(c + dx) ((b \sec(c + dx))^p)^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 - np); \frac{1}{2}(3 - np); \cos^2(c + dx)\right)}{d(1 - np)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c + d*x])^p]^n, x]$

[Out] $-\left(\left(\text{Cos}[c + d*x]*\text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1 - n*p)}{2}, \frac{(3 - n*p)}{2}, \text{Cos}[c + d*x]^2\right]*((b*\text{Sec}[c + d*x])^p)^n*\text{Sin}[c + d*x]\right)\right)/\left(d*(1 - n*p)*\text{Sqrt}[\text{Sin}[c + d*x]^2]\right)$

Rule 2643

$\text{Int}[(b*.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(n + 1)}{2}, \frac{(n + 3)}{2}, \text{Sin}[c + d*x]^2\right])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n - 1)}*((\text{Sin}[c + d*x]/b)^{(n - 1)}*\text{Int}[1/((\text{Sin}[c + d*x]/b)^n, x)], x] /;$ $\text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[n]$

Rule 4123

$\text{Int}[(b*.)*((c_.)*\sec[(e_.) + (f_.)*(x_.)]^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{IntPart}[p]*(b*(c*\text{Sec}[e + f*x])^n)^{\text{FracPart}[p]})/(c*\text{Sec}[e + f*x])^{(n*\text{FracPart}[p])}], \text{Int}[(c*\text{Sec}[e + f*x])^{(n*p)}, x], x] /;$ $\text{FreeQ}\{b, c, e, f, n, p\}, x] \&$

& !IntegerQ [p]

Rubi steps

$$\begin{aligned} \int ((b \sec(c + dx))^p)^n dx &= ((b \sec(c + dx))^{-np} ((b \sec(c + dx))^p)^n) \int (b \sec(c + dx))^{np} dx \\ &= \left(\left(\frac{\cos(c + dx)}{b} \right)^{np} ((b \sec(c + dx))^p)^n \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{-np} dx \\ &= - \frac{\cos(c + dx) {}_2F_1 \left(\frac{1}{2}, \frac{1}{2}(1 - np); \frac{1}{2}(3 - np); \cos^2(c + dx) \right) ((b \sec(c + dx))^p)^n \sin(c + dx)}{d(1 - np)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 69, normalized size = 0.85

$$\frac{\sqrt{-\tan^2(c + dx)} \cot(c + dx) ((b \sec(c + dx))^p)^n {}_2F_1 \left(\frac{1}{2}, \frac{np}{2}; \frac{1}{2}(np + 2); \sec^2(c + dx) \right)}{dnp}$$

Antiderivative was successfully verified.

[In] Integrate[((b*Sec[c + d*x])^p)^n,x]

[Out] (Cot[c + d*x]*Hypergeometric2F1[1/2, (n*p)/2, (2 + n*p)/2, Sec[c + d*x]^2]*((b*Sec[c + d*x])^p)^n*Sqrt[-Tan[c + d*x]^2])/(d*n*p)

fricas [F] time = 0.86, size = 0, normalized size = 0.00

$$\text{integral} \left(((b \sec(dx + c))^p)^n, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((b*sec(d*x+c))^p)^n,x, algorithm="fricas")

[Out] integral(((b*sec(d*x + c))^p)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int ((b \sec(dx + c))^p)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((b*sec(d*x+c))^p)^n,x, algorithm="giac")

[Out] integrate(((b*sec(d*x + c))^p)^n, x)

maple [F] time = 1.83, size = 0, normalized size = 0.00

$$\int \left((b \sec(dx + c))^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((b*sec(d*x+c))^p)^n,x)

[Out] int(((b*sec(d*x+c))^p)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left((b \sec(dx + c))^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((b*sec(d*x+c))^p)^n,x, algorithm="maxima")

[Out] integrate(((b*sec(d*x + c))^p)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\left(\frac{b}{\cos(c + dx)} \right)^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((b/cos(c + d*x))^p)^n,x)

[Out] int(((b/cos(c + d*x))^p)^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left((b \sec(c + dx))^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((b*sec(d*x+c))**p)**n,x)

[Out] Integral(((b*sec(c + d*x))**p)**n, x)

3.69 $\int (a(b \sec(c + dx))^p)^n dx$

Optimal. Leaf size=83

$$\frac{\sin(c + dx) \cos(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 - np); \frac{1}{2}(3 - np); \cos^2(c + dx)\right) (a(b \sec(c + dx))^p)^n}{d(1 - np)\sqrt{\sin^2(c + dx)}}$$

[Out] $-\cos(d*x+c)*\text{hypergeom}([1/2, -1/2*n*p+1/2], [-1/2*n*p+3/2], \cos(d*x+c)^2)*(a*(b*\sec(d*x+c))^p)^n*\sin(d*x+c)/d/(-n*p+1)/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {4123, 3772, 2643}

$$\frac{\sin(c + dx) \cos(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 - np); \frac{1}{2}(3 - np); \cos^2(c + dx)\right) (a(b \sec(c + dx))^p)^n}{d(1 - np)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a*(b*\text{Sec}[c + d*x])^p)^n, x]$

[Out] $-\left(\left(\text{Cos}[c + d*x]*\text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1 - n*p)}{2}, \frac{(3 - n*p)}{2}, \text{Cos}[c + d*x]^2\right]*(a*(b*\text{Sec}[c + d*x])^p)^n*\text{Sin}[c + d*x]\right)/\left(d*(1 - n*p)*\text{Sqrt}[\text{Sin}[c + d*x]^2]\right)\right)$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(n + 1)}{2}, \frac{(n + 3)}{2}, \text{Sin}[c + d*x]^2\right])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x \&\& \text{!IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n - 1)}*((\text{Sin}[c + d*x]/b)^{(n - 1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}\{b, c, d, n\}, x \&\& \text{!IntegerQ}[n]$

Rule 4123

$\text{Int}[(b_*)*((c_*)*\sec[(e_*) + (f_*)*(x_*)]^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{IntPart}[p]*(b*(c*\text{Sec}[e + f*x])^n)^{\text{FracPart}[p]})/(c*\text{Sec}[e + f*x])^{(n*\text{FracPart}[p])}], \text{Int}[(c*\text{Sec}[e + f*x])^{(n*p)}, x], x] /; \text{FreeQ}\{b, c, e, f, n, p\}, x \&$

& !IntegerQ[p]

Rubi steps

$$\begin{aligned} \int (a(b \sec(c + dx))^p)^n dx &= ((b \sec(c + dx))^{-np} (a(b \sec(c + dx))^p)^n) \int (b \sec(c + dx))^{np} dx \\ &= \left(\left(\frac{\cos(c + dx)}{b} \right)^{np} (a(b \sec(c + dx))^p)^n \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{-np} dx \\ &= - \frac{\cos(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 - np); \frac{1}{2}(3 - np); \cos^2(c + dx)\right) (a(b \sec(c + dx))^p)^n \sin(c + dx)}{d(1 - np)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 71, normalized size = 0.86

$$\frac{\sqrt{-\tan^2(c + dx)} \cot(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{np}{2}; \frac{1}{2}(np + 2); \sec^2(c + dx)\right) (a(b \sec(c + dx))^p)^n}{dnp}$$

Antiderivative was successfully verified.

[In] Integrate[(a*(b*Sec[c + d*x])^p)^n,x]

[Out] (Cot[c + d*x]*Hypergeometric2F1[1/2, (n*p)/2, (2 + n*p)/2, Sec[c + d*x]^2]*(a*(b*Sec[c + d*x])^p)^n*Sqrt[-Tan[c + d*x]^2])/(d*n*p)

fricas [F] time = 0.89, size = 0, normalized size = 0.00

$$\text{integral}\left(\left((b \sec(dx + c))^p a\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*sec(d*x+c))^p)^n,x, algorithm="fricas")

[Out] integral(((b*sec(d*x + c))^p*a)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int ((b \sec(dx + c))^p a)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*sec(d*x+c))^p)^n,x, algorithm="giac")

[Out] integrate(((b*sec(d*x + c))^p*a)^n, x)

maple [F] time = 1.66, size = 0, normalized size = 0.00

$$\int (a(b \sec(dx + c))^p)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*(b*sec(d*x+c))^p)^n,x)

[Out] int((a*(b*sec(d*x+c))^p)^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int ((b \sec(dx + c))^p a)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*sec(d*x+c))^p)^n,x, algorithm="maxima")

[Out] integrate(((b*sec(d*x + c))^p*a)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(a \left(\frac{b}{\cos(c + dx)} \right)^p \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*(b/cos(c + d*x))^p)^n,x)

[Out] int((a*(b/cos(c + d*x))^p)^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a(b \sec(c + dx))^p)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*(b*sec(d*x+c))**p)**n,x)

[Out] Integral((a*(b*sec(c + d*x))**p)**n, x)

3.70 $\int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=97

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{7/2}}{7b^3d} + \frac{10 \sin(c + dx)(b \sec(c + dx))^{3/2}}{21bd} + \frac{10 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

[Out] $10/21*(b*\sec(d*x+c))^{(3/2)*\sin(d*x+c)/b/d+2/7*(b*\sec(d*x+c))^{(7/2)*\sin(d*x+c)/b^3/d+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{7/2}}{7b^3d} + \frac{10 \sin(c + dx)(b \sec(c + dx))^{3/2}}{21bd} + \frac{10 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^4*Sqrt[b*Sec[c + d*x]], x]`

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*d) + (10*(b*\text{Sec}[c + d*x])^{(3/2)*\text{Sin}[c + d*x]}/(21*b*d) + (2*(b*\text{Sec}[c + d*x])^{(7/2)*\text{Sin}[c + d*x]}/(7*b^3*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \sec^4(c + dx) \sqrt{b \sec(c + dx)} dx &= \frac{\int (b \sec(c + dx))^{9/2} dx}{b^4} \\
 &= \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^3 d} + \frac{5 \int (b \sec(c + dx))^{5/2} dx}{7b^2} \\
 &= \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21bd} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^3 d} + \frac{5}{21} \int \sqrt{b \sec(c + dx)} dx \\
 &= \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21bd} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^3 d} + \frac{1}{21} (5\sqrt{b \sec(c + dx)}) \\
 &= \frac{10\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21bd}
 \end{aligned}$$

Mathematica [A] time = 0.22, size = 69, normalized size = 0.71

$$\frac{\sec^2(c + dx) \sqrt{b \sec(c + dx)} \left(5 \sin(2(c + dx)) + 6 \tan(c + dx) + 10 \cos^2(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{21d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^4*Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (Sec[c + d*x]^2*Sqrt[b*Sec[c + d*x]]*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)
```

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \sec(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^4, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^4, x)

maple [C] time = 1.20, size = 152, normalized size = 1.57

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(5i(\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF}\left(\frac{i}{2}, \frac{\cos(dx + c)}{1 + \cos(dx + c)}\right) \right)}{21d \sin(dx + c)^3 \cos(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-5*cos(d*x+c)^3+5*cos(d*x+c)^2-3*cos(d*x+c)+3)*(b/cos(d*x+c))^(1/2)/sin(d*x+c)^3/cos(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c+dx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^4,x)

[Out] int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^4, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} \sec^4(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**4*(b*sec(d*x+c))**(1/2), x)
```

```
[Out] Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**4, x)
```

3.71 $\int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=95

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{5/2}}{5b^2d} + \frac{6 \sin(c + dx)\sqrt{b \sec(c + dx)}}{5d} - \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

[Out] $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b^2/d-6/5*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{5/2}}{5b^2d} + \frac{6 \sin(c + dx)\sqrt{b \sec(c + dx)}}{5d} - \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3*Sqrt[b*Sec[c + d*x]],x]

[Out] $(-6*b*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (6*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(5*d) + (2*(b*\text{Sec}[c + d*x])^{(5/2)}*\text{Sin}[c + d*x])/(5*b^2*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \sec^3(c + dx) \sqrt{b \sec(c + dx)} dx &= \frac{\int (b \sec(c + dx))^{7/2} dx}{b^3} \\
 &= \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^2d} + \frac{3 \int (b \sec(c + dx))^{3/2} dx}{5b} \\
 &= \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^2d} - \frac{1}{5}(3b) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
 &= \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^2d} - \frac{(3b) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{5\sqrt{\cos(c + dx)}} \\
 &= -\frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^2d}
 \end{aligned}$$

Mathematica [A] time = 0.17, size = 69, normalized size = 0.73

$$\frac{\sec^2(c + dx) \sqrt{b \sec(c + dx)} \left(7 \sin(c + dx) + 3 \sin(3(c + dx)) - 12 \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10d}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3*Sqrt[b*Sec[c + d*x]], x]

[Out] (Sec[c + d*x]^2*Sqrt[b*Sec[c + d*x]]*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)])/(10*d)

fricas [F] time = 0.90, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^3, x)

maple [C] time = 0.92, size = 356, normalized size = 3.75

$$\frac{2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(3i(\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF} \left(\right. \right.}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2),x)

[Out]
$$-2/5/d*(1+\cos(d*x+c))^2*(-1+\cos(d*x+c))^2*(3*I*\cos(d*x+c)^3*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)-3*I*\cos(d*x+c)^3*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)+3*I*\cos(d*x+c)^2*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)-3*I*\cos(d*x+c)^2*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)+3*\cos(d*x+c)^3-2*\cos(d*x+c)^2-1)*(b/\cos(d*x+c))^(1/2)/\cos(d*x+c)^2/\sin(d*x+c)^5$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c+dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^3, x)`

[Out] `int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^3, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} \sec^3(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**3*(b*sec(d*x+c))**(1/2), x)`

[Out] `Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**3, x)`

3.72 $\int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=69

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{3/2}}{3bd} + \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

[Out] $2/3*(b*\sec(d*x+c))^{3/2}*\sin(d*x+c)/b/d+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{1/2}/d$

Rubi [A] time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{3/2}}{3bd} + \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^2*Sqrt[b*Sec[c + d*x]], x]`

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*(b*\text{Sec}[c + d*x])^{3/2}*\text{Sin}[c + d*x])/(3*b*d)$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx &= \frac{\int (b \sec(c + dx))^{5/2} dx}{b^2} \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3bd} + \frac{1}{3} \int \sqrt{b \sec(c + dx)} dx \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3bd} + \frac{1}{3} \left(\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3bd} \end{aligned}$$

Mathematica [A] time = 0.09, size = 51, normalized size = 0.74

$$\frac{2(b \sec(c + dx))^{3/2} \left(\sin(c + dx) + \cos^{\frac{3}{2}}(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3bd}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^2*Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (2*(b*Sec[c + d*x])^(3/2)*(Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + Sin[c + d*x]))/(3*b*d)
```

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^2, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^2, x)

maple [C] time = 0.79, size = 130, normalized size = 1.88

$$\frac{2\sqrt{\frac{b}{\cos(dx+c)}}(-1+\cos(dx+c))\left(i\sin(dx+c)\cos(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)},i\right)\sqrt{\frac{1}{1+\cos(dx+c)}}\right)}{3d\sin(dx+c)^3\cos(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2),x)

[Out] -2/3/d*(b/cos(d*x+c))^(1/2)*(-1+cos(d*x+c))*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*(1+cos(d*x+c))^2/sin(d*x+c)^3/cos(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx+c)} \sec(dx+c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c+dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^2,x)

[Out] int((b/cos(c + d*x))^(1/2)/cos(c + d*x)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c+dx)} \sec^2(c+dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(1/2), x)
```

```
[Out] Integral(sqrt(b*sec(c + d*x))*sec(c + d*x)**2, x)
```

3.73 $\int \sec(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=63

$$\frac{2 \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $-2*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]*Sqrt[b*Sec[c + d*x]], x]`

[Out] $(-2*b*\text{EllipticE}[(c + d*x)/2, 2])/((d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \sec(c + dx) \sqrt{b \sec(c + dx)} dx &= \frac{\int (b \sec(c + dx))^{3/2} dx}{b} \\ &= \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} - b \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} - \frac{b \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= -\frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} \end{aligned}$$

Mathematica [A] time = 0.05, size = 47, normalized size = 0.75

$$\frac{2\sqrt{b \sec(c + dx)} \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]*Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (2*Sqrt[b*Sec[c + d*x]]*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) +
Sin[c + d*x]))/d
```

fricas [F] time = 0.99, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c), x)

maple [C] time = 0.92, size = 314, normalized size = 4.98

$$2\sqrt{\frac{b}{\cos(dx+c)}} (1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x)

[Out] $-2/d*(b/\cos(d*x+c))^{1/2}*(1+\cos(d*x+c))^2*(-1+\cos(d*x+c))^2*(I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}-I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+\cos(d*x+c)-1)/\sin(d*x+c)^5$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\frac{b}{\cos(c+dx)}}}{\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/cos(c + d*x),x)

[Out] int((b/cos(c + d*x))^(1/2)/cos(c + d*x), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*sec(c + d*x))*sec(c + d*x), x)

3.74 $\int \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=38

$$\frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3771, 2641}

$$\frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[c + d*x]], x]

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/d$

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{b \sec(c + dx)} dx &= \left(\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}\right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 38, normalized size = 1.00

$$\frac{2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\sec(c+dx)}}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[c + d*x]], x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b\sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b\sec(dx+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.81, size = 98, normalized size = 2.58

$$\frac{2i\sqrt{\frac{b}{\cos(dx+c)}}(-1+\cos(dx+c))\sqrt{\frac{1}{1+\cos(dx+c)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right)(1+\cos(dx+c))^2}{d\sin(dx+c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(1/2), x)

[Out] -2*I/d*(b/cos(d*x+c))^(1/2)*(-1+cos(d*x+c))*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1+cos(d*x+c))^2/sin(d*x+c)^2

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c)), x)

mupad [B] time = 0.23, size = 35, normalized size = 0.92

$$\frac{2 \sqrt{\cos(c + dx)} \sqrt{\frac{b}{\cos(c+dx)}} F\left(\frac{c}{2} + \frac{dx}{2} \middle| 2\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2),x)

[Out] (2*cos(c + d*x)^(1/2)*(b/cos(c + d*x))^(1/2)*ellipticF(c/2 + (d*x)/2, 2))/d

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*sec(c + d*x)), x)

3.75 $\int \cos(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=39

$$\frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $2*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3771, 2639}

$$\frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]*Sqrt[b*Sec[c + d*x]],x]`

[Out] `(2*b*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])`

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned} \int \cos(c + dx) \sqrt{b \sec(c + dx)} dx &= b \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{b \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 39, normalized size = 1.00

$$\frac{2bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*Sqrt[b*Sec[c + d*x]], x]

[Out] (2*b*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c), x)

maple [C] time = 0.87, size = 303, normalized size = 7.77

$$2 \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - i \cos(dx + c) \sin(dx + c) \right) E$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x)`

[Out] $2/d*(I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*EllipticF(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}-I*\cos(d*x+c)*\sin(d*x+c)*EllipticE(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\sin(d*x+c)*EllipticF(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*EllipticE(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-\cos(d*x+c)^2+\cos(d*x+c))*(b/\cos(d*x+c))^{1/2}/\sin(d*x+c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*sec(d*x + c))*cos(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \cos(c + dx) \sqrt{\frac{b}{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)*(b/cos(c + d*x))^(1/2),x)`

[Out] `int(cos(c + d*x)*(b/cos(c + d*x))^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/2),x)`

[Out] `Integral(sqrt(b*sec(c + d*x))*cos(c + d*x), x)`

3.76 $\int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=67

$$\frac{2b \sin(c + dx)}{3d \sqrt{b \sec(c + dx)}} + \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

[Out] $2/3*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.05, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b \sin(c + dx)}{3d \sqrt{b \sec(c + dx)}} + \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^2*Sqrt[b*Sec[c + d*x]],x]`

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*b*\text{Sin}[c + d*x])/(3*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3769

`Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx) \sqrt{b \sec(c + dx)} dx &= b^2 \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\ &= \frac{2b \sin(c + dx)}{3d \sqrt{b \sec(c + dx)}} + \frac{1}{3} \int \sqrt{b \sec(c + dx)} dx \\ &= \frac{2b \sin(c + dx)}{3d \sqrt{b \sec(c + dx)}} + \frac{1}{3} \left(\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx)}{3d \sqrt{b \sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 51, normalized size = 0.76

$$\frac{\sqrt{b \sec(c + dx)} \left(\sin(2(c + dx)) + 2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^2*Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (Sqrt[b*Sec[c + d*x]]*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin
[2*(c + d*x)]))/(3*d)
```

fricas [F] time = 0.75, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^2, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^2, x)

maple [C] time = 0.89, size = 123, normalized size = 1.84

$$\frac{2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \operatorname{EllipticF} \left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i \right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} - (\cos^2(dx + c)) + c \right)}{3d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x)

[Out] -2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)^2+cos(d*x+c))*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(1/2)/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^2 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} \cos^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Integral(sqrt(b*sec(c + d*x))*cos(c + d*x)**2, x)
```

3.77 $\int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=70

$$\frac{2b^2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $2/5*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(3/2)}+6/5*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^3*Sqrt[b*Sec[c + d*x]],x]`

[Out] $(6*b*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\cos[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^2*\sin[c + d*x])/(5*d*(b*\text{Sec}[c + d*x])^{(3/2)})$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3769

`Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned} \int \cos^3(c + dx) \sqrt{b \sec(c + dx)} dx &= b^3 \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\ &= \frac{2b^2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{1}{5}(3b) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{2b^2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{(3b) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{6bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^2 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.06, size = 57, normalized size = 0.81

$$\frac{\sqrt{b \sec(c + dx)} \left(\sin(c + dx) + \sin(3(c + dx)) + 12\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10d}$$

Antiderivative was successfully verified.

[In] `Integrate[Cos[c + d*x]^3*Sqrt[b*Sec[c + d*x]], x]`

[Out] `(Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*d)`

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \cos(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")`

[Out] `integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^3, x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^3, x)

maple [C] time = 0.96, size = 315, normalized size = 4.50

$$2\sqrt{\frac{b}{\cos(dx+c)}} \left(3i \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - 3i \cos(dx+c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x)

[Out] 2/5/d*(b/cos(d*x+c))^(1/2)*(3*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-3*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+3*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)^4-2*cos(d*x+c)^2+3*cos(d*x+c))/sin(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx+c)} \cos(dx+c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c+dx)^3 \sqrt{\frac{b}{\cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**3*(b*sec(d*x+c))**(1/2),x)

[Out] Timed out

3.78 $\int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=95

$$\frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{10\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

[Out] $2/7*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/2)}+10/21*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{10\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4*Sqrt[b*Sec[c + d*x]], x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*d) + (2*b^3*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(5/2)}) + (10*b*\text{Sin}[c + d*x])/(21*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d^n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \cos^4(c + dx) \sqrt{b \sec(c + dx)} dx &= b^4 \int \frac{1}{(b \sec(c + dx))^{7/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{1}{7} (5b^2) \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{5}{21} \int \sqrt{b \sec(c + dx)} dx \\
 &= \frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{1}{21} (5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \\
 &= \frac{10\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b^3 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{1}{21}
 \end{aligned}$$

Mathematica [A] time = 0.09, size = 63, normalized size = 0.66

$$\frac{\sqrt{b \sec(c + dx)} \left(26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) + 40 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{84d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4*Sqrt[b*Sec[c + d*x]],x]

[Out] (Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*d)

fricas [F] time = 1.15, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \cos(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^4, x)

maple [C] time = 1.00, size = 145, normalized size = 1.53

$$\frac{2(-1 + \cos(dx + c)) \left(5i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} - 3(\cos^4(dx + c)) \right)}{21d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x)

[Out] -2/21/d*(-1+cos(d*x+c))*(5*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*cos(d*x+c)^4+3*cos(d*x+c)^3-5*cos(d*x+c)^2+5*cos(d*x+c))*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(1/2)/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^4 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c + dx)} \cos^4(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**4*(b*sec(d*x+c))**(1/2), x)

[Out] Integral(sqrt(b*sec(c + d*x))*cos(c + d*x)**4, x)

3.79 $\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=98

$$\frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $2/9*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/2)}+14/45*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(3/2)}+14/15*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^5*Sqrt[b*Sec[c + d*x]], x]

[Out] $(14*b*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(9*d*(b*\text{Sec}[c + d*x])^{(7/2)}) + (14*b^2*\text{Sin}[c + d*x])/(45*d*(b*\text{Sec}[c + d*x])^{(3/2)})$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \cos^5(c + dx) \sqrt{b \sec(c + dx)} dx &= b^5 \int \frac{1}{(b \sec(c + dx))^{9/2}} dx \\
&= \frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{1}{9} (7b^3) \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
&= \frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{1}{15} (7b) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
&= \frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{(7b) \int \sqrt{\cos(c + dx)} dx}{15 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
&= \frac{14bE\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^4 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^2 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.25, size = 71, normalized size = 0.72

$$\frac{\sqrt{b \sec(c + dx)} \left((33 \sin(c + dx) + 5 \sin(3(c + dx))) \cos^2(c + dx) + 84 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{90d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^5*Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (Sqrt[b*Sec[c + d*x]]*(84*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]^2*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)])))/(90*d)
```

fricas [F] time = 0.80, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} \cos(dx + c)^5, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^5, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^5, x)

maple [C] time = 0.94, size = 325, normalized size = 3.32

$$2\sqrt{\frac{b}{\cos(dx+c)}} \left(21i \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - 21i \cos(dx+c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x)

[Out] 2/45/d*(b/cos(d*x+c))^(1/2)*(21*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-21*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-5*cos(d*x+c)^6+21*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-21*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-2*cos(d*x+c)^4-14*cos(d*x+c)^2+21*cos(d*x+c))/sin(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \cos(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(d*x + c))*cos(d*x + c)^5, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^5 \sqrt{\frac{b}{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**5*(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

3.80 $\int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=95

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{7/2}}{7b^2d} + \frac{10 \sin(c + dx)(b \sec(c + dx))^{3/2}}{21d} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

[Out] $10/21*(b*\sec(d*x+c))^{(3/2)*\sin(d*x+c)/d+2/7*(b*\sec(d*x+c))^{(7/2)*\sin(d*x+c)}/b^2/d+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{7/2}}{7b^2d} + \frac{10 \sin(c + dx)(b \sec(c + dx))^{3/2}}{21d} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^3*(b*Sec[c + d*x])^(3/2), x]`

[Out] `(10*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(21*d) + (10*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(21*d) + (2*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*b^2*d)`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sec^3(c + dx)(b \sec(c + dx))^{3/2} dx &= \frac{\int (b \sec(c + dx))^{9/2} dx}{b^3} \\ &= \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^2d} + \frac{5 \int (b \sec(c + dx))^{5/2} dx}{7b} \\ &= \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^2d} + \frac{1}{21}(5b) \\ &= \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^2d} + \frac{1}{21}(5b) \\ &= \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} \end{aligned}$$

Mathematica [A] time = 0.19, size = 64, normalized size = 0.67

$$\frac{(b \sec(c + dx))^{5/2} \left(5 \sin(2(c + dx)) + 6 \tan(c + dx) + 10 \cos^2(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{21bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3*(b*Sec[c + d*x])^(3/2), x]

[Out] ((b*Sec[c + d*x])^(5/2)*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*b*d)

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \sec(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b*sec(d*x + c)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^3, x)

maple [C] time = 0.77, size = 152, normalized size = 1.60

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(5i(\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF}\left(\frac{i}{\sqrt{1 + \cos(dx + c)}}\right) \right)}{21d \sin(dx + c)^3 \cos(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-5*cos(d*x+c)^3+5*cos(d*x+c)^2-3*cos(d*x+c)+3)*(b/cos(d*x+c))^(3/2)/sin(d*x+c)^3/cos(d*x+c)^2

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec^3(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\cos(c+dx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^3,x)

[Out] int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{3}{2}} \sec^3(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**3*(b*sec(d*x+c))**(3/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x)**3, x)
```

3.81 $\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=98

$$-\frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)(b \sec(c + dx))^{5/2}}{5bd} + \frac{6b \sin(c + dx) \sqrt{b \sec(c + dx)}}{5d}$$

[Out] $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b/d-6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*b*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$-\frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)(b \sec(c + dx))^{5/2}}{5bd} + \frac{6b \sin(c + dx) \sqrt{b \sec(c + dx)}}{5d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(3/2), x]`

[Out] $(-6*b^2*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (6*b*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(5*d) + (2*(b*\text{Sec}[c + d*x])^{(5/2)}*\text{Sin}[c + d*x])/(5*b*d)$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx &= \frac{\int (b \sec(c + dx))^{7/2} dx}{b^2} \\
 &= \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5bd} + \frac{3}{5} \int (b \sec(c + dx))^{3/2} dx \\
 &= \frac{6b\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5bd} - \frac{1}{5} (3b^2) \int \sec^2(c + dx) dx \\
 &= \frac{6b\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5bd} - \frac{(3b^2)}{5\sqrt{\cos(c + dx)}} \\
 &= -\frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{6b\sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5bd}
 \end{aligned}$$

Mathematica [A] time = 0.18, size = 64, normalized size = 0.65

$$\frac{(b \sec(c + dx))^{5/2} \left(7 \sin(c + dx) + 3 \sin(3(c + dx)) - 12 \cos^{5/2}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10bd}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] ((b*Sec[c + d*x])^(5/2)*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] +
7*Sin[c + d*x] + 3*Sin[3*(c + d*x)])/(10*b*d)
```

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b*sec(d*x + c)^3, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^2, x)

maple [C] time = 0.76, size = 356, normalized size = 3.63

$$2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(3i (\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticE} \left(\frac{i}{\sqrt{1 + \cos(dx + c)}} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2),x)

[Out] 2/5/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)+3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*cos(d*x+c)^3+2*cos(d*x+c)^2+1)*(b/cos(d*x+c))^(3/2)/sin(d*x+c)^5/cos(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\cos(c+dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^2, x)`

[Out] `int((b/cos(c + d*x))^(3/2)/cos(c + d*x)^2, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{3}{2}} \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x)**2, x)`

3.82 $\int \sec(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=67

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} + \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

[Out] $2/3*(b*\sec(d*x+c))^{3/2}*\sin(d*x+c)/d+2/3*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^{3/2}}{3d} + \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]*(b*Sec[c + d*x])^(3/2), x]`

[Out] $(2*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*(b*\text{Sec}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(3*d)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sec(c + dx)(b \sec(c + dx))^{3/2} dx &= \frac{\int (b \sec(c + dx))^{5/2} dx}{b} \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} + \frac{1}{3}b \int \sqrt{b \sec(c + dx)} dx \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} + \frac{1}{3} \left(b \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2b \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} \end{aligned}$$

Mathematica [A] time = 0.08, size = 49, normalized size = 0.73

$$\frac{2b \sqrt{b \sec(c + dx)} \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(3/2), x]

[Out] (2*b*Sqrt[b*Sec[c + d*x]]*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d)

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c), x)

maple [C] time = 0.66, size = 122, normalized size = 1.82

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \right)}{3d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x)

[Out] -2/3/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*(b/cos(d*x+c))^(3/2)/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)/cos(c + d*x),x)

[Out] int((b/cos(c + d*x))^(3/2)/cos(c + d*x), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{3}{2}} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)*(b*sec(d*x+c))**(3/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**(3/2)*sec(c + d*x), x)
```

3.83 $\int (b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=66

$$\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $-2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2*b*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.03, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3771, 2639}

$$\frac{2b \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c + d*x])^{(3/2)}, x]$

[Out] $(-2*b^2*\text{EllipticE}[(c + d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3768

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[(b^2*(n-2))/(n-1), \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 3771

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n-1)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (b \sec(c + dx))^{3/2} dx &= \frac{2b\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} - b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
&= \frac{2b\sqrt{b \sec(c + dx)} \sin(c + dx)}{d} - \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
&= -\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b\sqrt{b \sec(c + dx)} \sin(c + dx)}{d}
\end{aligned}$$

Mathematica [A] time = 0.03, size = 48, normalized size = 0.73

$$\frac{2b\sqrt{b \sec(c + dx)} \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2),x]

[Out] (2*b*Sqrt[b*Sec[c + d*x]]*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/d

fricas [F] time = 0.80, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.84, size = 320, normalized size = 4.85

$$2(1 + \cos(dx + c))^2(-1 + \cos(dx + c))^2 \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(3/2), x)

[Out] $-2/d*(1+\cos(d*x+c))^{1/2}*(-1+\cos(d*x+c))^{1/2}*(I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}-I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+\cos(d*x+c)-1)*\cos(d*x+c)*(b/\cos(d*x+c))^{3/2}/\sin(d*x+c)^5$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2), x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2), x)

[Out] int((b/cos(c + d*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(3/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**(3/2), x)
```

3.84 $\int \cos(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=39

$$\frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d}$$

[Out] $2*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.03, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3771, 2641}

$$\frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]*(b*Sec[c + d*x])^(3/2), x]`

[Out] `(2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned} \int \cos(c + dx)(b \sec(c + dx))^{3/2} dx &= b \int \sqrt{b \sec(c + dx)} dx \\ &= (b\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 39, normalized size = 1.00

$$\frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(3/2), x]

[Out] (2*b*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \cos(dx + c) \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b*cos(d*x + c)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c), x)

maple [C] time = 0.75, size = 98, normalized size = 2.51

$$\frac{2i \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}} \left(\frac{\cos(dx+c)}{1+\cos(dx+c)}\right)^{\frac{3}{2}} (-1 + \cos(dx + c)) \text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) (1 + \cos(dx + c))^3 \sqrt{\frac{1}{1+\cos(dx+c)}}}{d \sin(dx + c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)*(b*sec(d*x+c))^(3/2),x)`

[Out] `-2*I/d*(b/cos(d*x+c))^(3/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(3/2)*(-1+cos(d*x+c))*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1+cos(d*x+c))^3*(1/(1+cos(d*x+c)))^(1/2)/sin(d*x+c)^2`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \cos(c + dx) \left(\frac{b}{\cos(c + dx)} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)*(b/cos(c + d*x))^(3/2),x)`

[Out] `int(cos(c + d*x)*(b/cos(c + d*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{3}{2}} \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))**(3/2),x)`

[Out] `Integral((b*sec(c + d*x))**(3/2)*cos(c + d*x), x)`

3.85 $\int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=41

$$\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3771, 2639}

$$\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^(3/2), x]

[Out] (2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx)(b \sec(c + dx))^{3/2} dx &= b^2 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{b^2 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(3/2), x]

[Out] (2*b^2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \cos(dx + c)^2 \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b*cos(d*x + c)^2*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^2, x)

maple [C] time = 0.77, size = 309, normalized size = 7.54

$$2 \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - i \cos(dx + c) \sin(dx + c) \operatorname{EllipticE}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2), x)`

[Out] $2/d*(I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}-I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-\cos(d*x+c)^2+\cos(d*x+c))*\cos(d*x+c)*(b/\cos(d*x+c))^{3/2}/\sin(d*x+c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(3/2), x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 \left(\frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2), x)`

[Out] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(3/2), x)`

[Out] Timed out

3.86 $\int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=70

$$\frac{2b^2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

[Out] $2/3*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+2/3*b*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.05, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b^2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^3*(b*Sec[c + d*x])^(3/2),x]`

[Out] $(2*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*b^2*\text{Sin}[c + d*x])/(3*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3769

`Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \cos^3(c + dx)(b \sec(c + dx))^{3/2} dx &= b^3 \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\ &= \frac{2b^2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{1}{3}b \int \sqrt{b \sec(c + dx)} dx \\ &= \frac{2b^2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{1}{3} \left(b\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \right) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b^2 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 52, normalized size = 0.74

$$\frac{b\sqrt{b \sec(c + dx)} \left(\sin(2(c + dx)) + 2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3*(b*Sec[c + d*x])^(3/2), x]

[Out] (b*Sqrt[b*Sec[c + d*x]]*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*d)

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \cos(dx + c)^3 \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b*cos(d*x + c)^3*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^3, x)

maple [C] time = 0.82, size = 129, normalized size = 1.84

$$\frac{2(1 + \cos(dx + c))^2 \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}} (-1 + \cos(dx + c)) \cos(dx + c) \left(-i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1+\cos(dx+c)}{1-\cos(dx+c)}}\right)}{3d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2),x)

[Out] 2/3/d*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(3/2)*(-1+cos(d*x+c))*cos(d*x+c)*(-I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c))))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+cos(d*x+c)^2-cos(d*x+c))/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^3 \left(\frac{b}{\cos(c + dx)}\right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**3*(b*sec(d*x+c))**(3/2),x)

[Out] Timed out

3.87 $\int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=72

$$\frac{2b^3 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

[Out] $2/5*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+6/5*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

Rubi [A] time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^3 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^4*(b*Sec[c + d*x])^(3/2), x]`

[Out] $(6*b^2*\text{EllipticE}[(c + d*x)/2, 2])/((5*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^3*\text{Sin}[c + d*x]))/(5*d*(b*\text{Sec}[c + d*x])^(3/2))$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3769

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3771


```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \cos^4(c + dx)(b \sec(c + dx))^{3/2} dx &= b^4 \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\ &= \frac{2b^3 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{1}{5} (3b^2) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{2b^3 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{(3b^2) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{6b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^3 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 58, normalized size = 0.81

$$\frac{b\sqrt{b \sec(c + dx)} \left(\sin(c + dx) + \sin(3(c + dx)) + 12\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^4*(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] (b*Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] +
Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*d)
```

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \cos(dx + c)^4 \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b*cos(d*x + c)^4*sec(d*x + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^4, x)

maple [C] time = 0.84, size = 319, normalized size = 4.43

$$2 \left(-3i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} + 3i \cos(dx + c) \sin(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x)

[Out]
$$-2/5/d*(-3*I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}+3*I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-3*I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+3*I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+\cos(d*x+c)^4+2*\cos(d*x+c)^2-3*\cos(d*x+c))*\cos(d*x+c)*(b/\cos(d*x+c))^{3/2}/\sin(d*x+c)$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^4 \left(\frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**4*(b*sec(d*x+c))**(3/2), x)

[Out] Timed out

3.88 $\int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=98

$$\frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^2 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

[Out] $2/7*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^(5/2)+10/21*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^(1/2)+10/21*b*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/d$

Rubi [A] time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^2 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^5*(b*Sec[c + d*x])^(3/2), x]

[Out] $(10*b*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(2*1*d) + (2*b^4*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^(5/2)) + (10*b^2*\text{Sin}[c + d*x])/(21*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \cos^5(c + dx)(b \sec(c + dx))^{3/2} dx &= b^5 \int \frac{1}{(b \sec(c + dx))^{7/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{1}{7} (5b^3) \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\
 &= \frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^2 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{1}{21} (5b) \int \sqrt{b \sec(c + dx)} dx \\
 &= \frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^2 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{1}{21} (5b\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \\
 &= \frac{10b\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b^4 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}}
 \end{aligned}$$

Mathematica [A] time = 0.08, size = 64, normalized size = 0.65

$$\frac{b\sqrt{b \sec(c + dx)} \left(26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) + 40\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{84d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^5*(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] (b*Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] +
26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*d)
```

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \cos(dx + c)^5 \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b*cos(d*x + c)^5*sec(d*x + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^5, x)

maple [C] time = 0.94, size = 151, normalized size = 1.54

$$\frac{2(1 + \cos(dx + c))^2 \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}} (-1 + \cos(dx + c)) \cos(dx + c) \left(-5i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1+\cos(dx+c)}{1-\cos(dx+c)}}\right)}{21d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2),x)

[Out] 2/21/d*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(3/2)*(-1+cos(d*x+c))*cos(d*x+c)*(-5*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+3*cos(d*x+c)^4-3*cos(d*x+c)^3+5*cos(d*x+c)^2-5*cos(d*x+c))/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^5, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^5 \left(\frac{b}{\cos(c + dx)}\right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2),x)

[Out] int(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**5*(b*sec(d*x+c))**(3/2), x)

[Out] Timed out

3.89 $\int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=100

$$\frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $2/9*b^5*\sin(d*x+c)/d/(b*\sec(d*x+c))^(7/2)+14/45*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(3/2)+14/15*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^(1/2))/d/\cos(d*x+c)^(1/2)/(b*\sec(d*x+c))^(1/2)$

Rubi [A] time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^6*(b*Sec[c + d*x])^(3/2),x]`

[Out] $(14*b^2*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^5*\text{Sin}[c + d*x])/(9*d*(b*\text{Sec}[c + d*x])^(7/2)) + (14*b^3*\text{Sin}[c + d*x])/(45*d*(b*\text{Sec}[c + d*x])^(3/2))$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3769

`Int[(csc[(c_.) + (d_.)*(x_)*(b_.)]^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3771


```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \cos^6(c + dx)(b \sec(c + dx))^{3/2} dx &= b^6 \int \frac{1}{(b \sec(c + dx))^{9/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{1}{9} (7b^4) \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{1}{15} (7b^2) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{(7b^2) \int \sqrt{\cos(c + dx)} dx}{15\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 &= \frac{14b^2 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^5 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^3 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] time = 0.15, size = 72, normalized size = 0.72

$$\frac{b\sqrt{b \sec(c + dx)} \left((33 \sin(c + dx) + 5 \sin(3(c + dx))) \cos^2(c + dx) + 84\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{90d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^6*(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] (b*Sqrt[b*Sec[c + d*x]]*(84*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] +
Cos[c + d*x]^2*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)])))/(90*d)
```

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b \cos(dx + c)^6 \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b*cos(d*x + c)^6*sec(d*x + c), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^6, x)

maple [C] time = 0.95, size = 331, normalized size = 3.31

$$2 \left(21i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - 21i \cos(dx + c) \sin(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x)

[Out] 2/45/d*(21*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-21*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-5*cos(d*x+c)^6+21*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-21*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-2*cos(d*x+c)^4-14*cos(d*x+c)^2+21*cos(d*x+c))*cos(d*x+c)*(b/cos(d*x+c))^(3/2)/sin(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \cos(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(3/2)*cos(d*x + c)^6, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^6 \left(\frac{b}{\cos(c + dx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2), x)
```

```
[Out] int(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**6*(b*sec(d*x+c))**(3/2), x)
```

```
[Out] Timed out
```

3.90 $\int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=98

$$\frac{10b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\sec(c+dx)}}{21d} + \frac{2\sin(c+dx)(b\sec(c+dx))^{7/2}}{7bd} + \frac{10b\sin(c+dx)(b\sec(c+dx))^{5/2}}{21d}$$

[Out] $10/21*b*(b*\sec(d*x+c))^{(3/2)}*\sin(d*x+c)/d+2/7*(b*\sec(d*x+c))^{(7/2)}*\sin(d*x+c)/b/d+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{10b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\sec(c+dx)}}{21d} + \frac{2\sin(c+dx)(b\sec(c+dx))^{7/2}}{7bd} + \frac{10b\sin(c+dx)(b\sec(c+dx))^{5/2}}{21d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(5/2),x]

[Out] $(10*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*d) + (10*b*(b*\text{Sec}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(21*d) + (2*(b*\text{Sec}[c + d*x])^{(7/2)}*\text{Sin}[c + d*x])/(7*b*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \sec^2(c + dx)(b \sec(c + dx))^{5/2} dx &= \frac{\int (b \sec(c + dx))^{9/2} dx}{b^2} \\
 &= \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7bd} + \frac{5}{7} \int (b \sec(c + dx))^{5/2} dx \\
 &= \frac{10b(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7bd} + \frac{1}{21} \int (b \sec(c + dx))^{5/2} dx \\
 &= \frac{10b(b \sec(c + dx))^{3/2} \sin(c + dx)}{21d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7bd} + \frac{1}{21} \int (b \sec(c + dx))^{5/2} dx \\
 &= \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{10b(b \sec(c + dx))^{3/2}}{21d}
 \end{aligned}$$

Mathematica [A] time = 0.19, size = 61, normalized size = 0.62

$$\frac{(b \sec(c + dx))^{5/2} \left(5 \sin(2(c + dx)) + 6 \tan(c + dx) + 10 \cos^2(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{21d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] ((b*Sec[c + d*x])^(5/2)*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] +
5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d)
```

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \sec(dx + c)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b^2*sec(d*x + c)^4, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^2, x)

maple [C] time = 0.80, size = 152, normalized size = 1.55

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(5i(\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, I\right) - 5\cos(dx + c)^3 + 5\cos(dx + c)^2 - 3\cos(dx + c) + 3 \right) (b/\cos(dx + c))^{5/2}}{21d \sin(dx + c)^3 \cos(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-5*cos(d*x+c)^3+5*cos(d*x+c)^2-3*cos(d*x+c)+3)*(b/cos(d*x+c))^(5/2)/sin(d*x+c)^3/cos(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^2, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}}{\cos(c+dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)/cos(c + d*x)^2,x)

[Out] int((b/cos(c + d*x))^(5/2)/cos(c + d*x)^2, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{5}{2}} \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(5/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**(5/2)*sec(c + d*x)**2, x)
```

3.91 $\int \sec(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=97

$$-\frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{6b^2 \sin(c + dx)\sqrt{b \sec(c + dx)}}{5d} + \frac{2 \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d}$$

[Out] $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/d-6/5*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*b^2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3768, 3771, 2639}

$$\frac{6b^2 \sin(c + dx)\sqrt{b \sec(c + dx)}}{5d} - \frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)(b \sec(c + dx))^{5/2}}{5d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]*(b*Sec[c + d*x])^(5/2),x]

[Out] $(-6*b^3*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (6*b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(5*d) + (2*(b*\text{Sec}[c + d*x])^{(5/2)}*\text{Sin}[c + d*x])/(5*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)*(b_.)]^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771


```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^(n)*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \sec(c + dx)(b \sec(c + dx))^{5/2} dx &= \frac{\int (b \sec(c + dx))^{7/2} dx}{b} \\
 &= \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d} + \frac{1}{5}(3b) \int (b \sec(c + dx))^{3/2} dx \\
 &= \frac{6b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d} - \frac{1}{5} (3b^3) \\
 &= \frac{6b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d} - \frac{(3b^3)}{5\sqrt{\cos(c + dx)}} \\
 &= -\frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}} + \frac{6b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d}
 \end{aligned}$$

Mathematica [A] time = 0.18, size = 61, normalized size = 0.63

$$\frac{(b \sec(c + dx))^{5/2} \left(7 \sin(c + dx) + 3 \sin(3(c + dx)) - 12 \cos^2(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] ((b*Sec[c + d*x])^(5/2)*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] +
7*Sin[c + d*x] + 3*Sin[3*(c + d*x)]))/(10*d)
```

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b^2*sec(d*x + c)^3, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c), x)

maple [C] time = 0.82, size = 348, normalized size = 3.59

$$2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(3i \left(\cos^3(dx + c) \right) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticE} \left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, I \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x)

[Out] 2/5/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)+3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*cos(d*x+c)^3+2*cos(d*x+c)^2+1)*(b/cos(d*x+c))^(5/2)/sin(d*x+c)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)} \right)^{5/2}}{\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(5/2)/cos(c + d*x), x)`

[Out] `int((b/cos(c + d*x))^(5/2)/cos(c + d*x), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{5}{2}} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)*(b*sec(d*x+c))**(5/2), x)`

[Out] `Integral((b*sec(c + d*x))**(5/2)*sec(c + d*x), x)`

3.92 $\int (b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=70

$$\frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d}$$

[Out] $2/3*b*(b*\sec(d*x+c))^{3/2}*\sin(d*x+c)/d+2/3*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3771, 2641}

$$\frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b \sin(c + dx) (b \sec(c + dx))^{3/2}}{3d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c + d*x])^{5/2}, x]$

[Out] $(2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*b*(b*\text{Sec}[c + d*x])^{3/2}*\text{Sin}[c + d*x])/(3*d)$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3768

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[(b^2*(n-2))/(n-1), \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3771

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{n-1}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned} \int (b \sec(c + dx))^{5/2} dx &= \frac{2b(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} + \frac{1}{3} b^2 \int \sqrt{b \sec(c + dx)} dx \\ &= \frac{2b(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} + \frac{1}{3} (b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} \\ &= \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b(b \sec(c + dx))^{3/2} \sin(c + dx)}{3d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 51, normalized size = 0.73

$$\frac{2b^2 \sqrt{b \sec(c + dx)} \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2), x]

[Out] (2*b^2*Sqrt[b*Sec[c + d*x]]*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*d)

fricas [F] time = 0.71, size = 0, normalized size = 0.00

$$\text{integral}(\sqrt{b \sec(dx + c)} b^2 \sec(dx + c)^2, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^2*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.79, size = 128, normalized size = 1.83

$$\frac{2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - \cos(dx + c) \right)}{3d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^(5/2),x)`

[Out] `-2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*cos(d*x+c)*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(5/2)/sin(d*x+c)^3`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(5/2),x)`

[Out] `int((b/cos(c + d*x))^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**(5/2),x)`

[Out] `Integral((b*sec(c + d*x))**(5/2), x)`

3.93 $\int \cos(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=68

$$\frac{2b^2 \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $-2*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2*b^2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2b^2 \sin(c + dx) \sqrt{b \sec(c + dx)}}{d} - \frac{2b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]*(b*Sec[c + d*x])^(5/2), x]`

[Out] $(-2*b^3*\text{EllipticE}[(c + d*x)/2, 2])/((d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2639

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned}
 \int \cos(c + dx)(b \sec(c + dx))^{5/2} dx &= b \int (b \sec(c + dx))^{3/2} dx \\
 &= \frac{2b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d} - b^3 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
 &= \frac{2b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d} - \frac{b^3 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 &= -\frac{2b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d}
 \end{aligned}$$

Mathematica [A] time = 0.02, size = 50, normalized size = 0.74

$$\frac{2b^2 \sqrt{b \sec(c + dx)} \left(\sin(c + dx) - \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(5/2), x]

[Out] (2*b^2*Sqrt[b*Sec[c + d*x]]*(-(Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]) + Sin[c + d*x]))/d

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c) \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c), x)

maple [C] time = 0.96, size = 324, normalized size = 4.76

$$2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(i \cos(dx + c) \sin(dx + c) \operatorname{EllipticE}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{1}{1 + \cos(dx + c)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x)

[Out] 2/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*cos(d*x+c)*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*cos(d*x+c)^2*(b/cos(d*x+c))^(5/2)/sin(d*x+c)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx) \left(\frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)*(b/cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)*(b/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))**(5/2),x)

[Out] Timed out

3.94 $\int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=41

$$\frac{2b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\sec(c+dx)}}{d}$$

[Out] $2*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.04, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3771, 2641}

$$\frac{2b^2\sqrt{\cos(c+dx)}F\left(\frac{1}{2}(c+dx)\middle|2\right)\sqrt{b\sec(c+dx)}}{d}$$

Antiderivative was successfully verified.

[In] `Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^(5/2), x]`

[Out] `(2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d`

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2641

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]`

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned} \int \cos^2(c + dx)(b \sec(c + dx))^{5/2} dx &= b^2 \int \sqrt{b \sec(c + dx)} dx \\ &= (b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx \\ &= \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(5/2), x]

[Out] (2*b^2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/d

fricas [F] time = 0.92, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c)^2 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)^2*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^2, x)

maple [C] time = 0.81, size = 98, normalized size = 2.39

$$\frac{2i \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \left(\frac{\cos(dx+c)}{1+\cos(dx+c)}\right)^{\frac{5}{2}} (-1 + \cos(dx + c)) \text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) (1 + \cos(dx + c))^5 \left(\frac{1}{1+\cos(dx+c)}\right)^{\frac{3}{2}}}{d \sin(dx + c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2),x)`

[Out] `-2*I/d*(b/cos(d*x+c))^(5/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(5/2)*(-1+cos(d*x+c))*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1+cos(d*x+c))^5*(1/(1+cos(d*x+c)))^(3/2)/sin(d*x+c)^2`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 \left(\frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(5/2),x)`

[Out] Timed out

3.95 $\int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=41

$$\frac{2b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

[Out] $2*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3771, 2639}

$$\frac{2b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3*(b*Sec[c + d*x])^(5/2), x]

[Out] (2*b^3*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \cos^3(c + dx)(b \sec(c + dx))^{5/2} dx &= b^3 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{b^3 \int \sqrt{\cos(c + dx)} dx}{\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{2b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 38, normalized size = 0.93

$$\frac{2 \cos^{\frac{5}{2}}(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right) (b \sec(c + dx))^{5/2}}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3*(b*Sec[c + d*x])^(5/2), x]

[Out] (2*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2]*(b*Sec[c + d*x])^(5/2))/d

fricas [F] time = 0.88, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c)^3 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)^3*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^3, x)

maple [C] time = 0.82, size = 311, normalized size = 7.59

$$2 \left(-i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} + i \cos(dx + c) \sin(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2),x)`

[Out]
$$-2/d*(-I*\cos(d*x+c)*\sin(d*x+c)*\text{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\cos(d*x+c)*\sin(d*x+c)*\text{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*\text{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\sin(d*x+c)*\text{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+\cos(d*x+c)^2-\cos(d*x+c))*\cos(d*x+c)^2*(b/\cos(d*x+c))^{5/2}/\sin(d*x+c)$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^3, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^3 \left(\frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2),x)`

[Out] `int(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**3*(b*sec(d*x+c))**(5/2),x)`

[Out] Timed out

3.96 $\int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=72

$$\frac{2b^3 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

[Out] $2/3*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^(1/2)+2/3*b^2*(\cos(1/2*d*x+1/2*c)^2)^(1/2)/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^(1/2))*\cos(d*x+c)^(1/2)*(b*\sec(d*x+c))^(1/2)/d$

Rubi [A] time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b^3 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4*(b*Sec[c + d*x])^(5/2),x]

[Out] $(2*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*d) + (2*b^3*\text{Sin}[c + d*x])/(3*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \cos^4(c + dx)(b \sec(c + dx))^{5/2} dx &= b^4 \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\ &= \frac{2b^3 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{1}{3}b^2 \int \sqrt{b \sec(c + dx)} dx \\ &= \frac{2b^3 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} + \frac{1}{3} \left(b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)} \right) \int \frac{1}{\sqrt{\cos(c + dx)}} \\ &= \frac{2b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3d} + \frac{2b^3 \sin(c + dx)}{3d\sqrt{b \sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 54, normalized size = 0.75

$$\frac{b^2 \sqrt{b \sec(c + dx)} \left(\sin(2(c + dx)) + 2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4*(b*Sec[c + d*x])^(5/2), x]

[Out] (b^2*Sqrt[b*Sec[c + d*x]]*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*d)

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c)^4 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)^4*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^4, x)

maple [C] time = 0.82, size = 131, normalized size = 1.82

$$\frac{2(1 + \cos(dx + c))^2 \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} (-1 + \cos(dx + c)) (\cos^2(dx + c)) \left(-i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right)\right)}{3d \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x)

[Out] 2/3/d*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(5/2)*(-1+cos(d*x+c))*cos(d*x+c)^2*(-I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+cos(d*x+c)^2-cos(d*x+c))/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^4, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^4 \left(\frac{b}{\cos(c + dx)}\right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**4*(b*sec(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

3.97 $\int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=72

$$\frac{2b^4 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

[Out] $2/5*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(3/2)}+6/5*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^4 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^5*(b*\text{Sec}[c + d*x])^{(5/2)}, x]$

[Out] $(6*b^3*\text{EllipticE}[(c + d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^4*\text{Sin}[c + d*x])/(5*d*(b*\text{Sec}[c + d*x])^{(3/2)})$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3769

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_*)]*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned} \int \cos^5(c + dx)(b \sec(c + dx))^{5/2} dx &= b^5 \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\ &= \frac{2b^4 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{1}{5} (3b^3) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{2b^4 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{(3b^3) \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{6b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^4 \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 60, normalized size = 0.83

$$\frac{b^2 \sqrt{b \sec(c + dx)} \left(\sin(c + dx) + \sin(3(c + dx)) + 12 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10d}$$

Antiderivative was successfully verified.

[In] `Integrate[Cos[c + d*x]^5*(b*Sec[c + d*x])^(5/2), x]`

[Out] `(b^2*Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*d)`

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c)^5 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")`

[Out] `integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)^5*sec(d*x + c)^2, x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^5, x)

maple [C] time = 0.94, size = 321, normalized size = 4.46

$$2 \left(\frac{b}{\cos(dx+c)} \right)^{\frac{5}{2}} \left(\cos^2(dx+c) \right) \left(-3i \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) \sqrt{\frac{1}{1+\cos(dx+c)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x)

[Out] $-2/5/d*(b/\cos(d*x+c))^{5/2}*\cos(d*x+c)^2*(-3*I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}+3*I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-3*I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+3*I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+\cos(d*x+c)^4+2*\cos(d*x+c)^2-3*\cos(d*x+c))/\sin(d*x+c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{\frac{5}{2}} \cos(dx+c)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^5*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^5, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c+dx)^5 \left(\frac{b}{\cos(c+dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**5*(b*sec(d*x+c))**(5/2), x)

[Out] Timed out

3.98 $\int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=100

$$\frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^3 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

[Out] $2/7*b^5*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/2)}+10/21*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+10/21*b^2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^3 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^6*(b*Sec[c + d*x])^(5/2), x]

[Out] $(10*b^2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*d) + (2*b^5*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(5/2)}) + (10*b^3*\text{Sin}[c + d*x])/(21*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d^n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \cos^6(c + dx)(b \sec(c + dx))^{5/2} dx &= b^6 \int \frac{1}{(b \sec(c + dx))^{7/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{1}{7} (5b^4) \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\
 &= \frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^3 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{1}{21} (5b^2) \int \sqrt{b \sec(c + dx)} dx \\
 &= \frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10b^3 \sin(c + dx)}{21d\sqrt{b \sec(c + dx)}} + \frac{1}{21} (5b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \\
 &= \frac{10b^2 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21d} + \frac{2b^5 \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \dots
 \end{aligned}$$

Mathematica [A] time = 0.08, size = 66, normalized size = 0.66

$$\frac{b^2 \sqrt{b \sec(c + dx)} \left(26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) + 40 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{84d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^6*(b*Sec[c + d*x])^(5/2), x]

[Out] (b^2*Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*d)

fricas [F] time = 1.39, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c)^6 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)^6*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^6, x)

maple [C] time = 0.97, size = 153, normalized size = 1.53

$$2(1 + \cos(dx + c))^2 \left(\frac{b}{\cos(dx+c)} \right)^{\frac{5}{2}} (-1 + \cos(dx + c)) (\cos^2(dx + c)) \left(5i \sin(dx + c) \operatorname{EllipticF} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) \right)$$

21d sin(dx +

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(5/2)*(-1+cos(d*x+c))*cos(d*x+c)^2*(5*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*cos(d*x+c)^4+3*cos(d*x+c)^3-5*cos(d*x+c)^2+5*cos(d*x+c))/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^6*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^6, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^6 \left(\frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2),x)

[Out] int(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**6*(b*sec(d*x+c))**(5/2),x)

[Out] Timed out

3.99 $\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=100

$$\frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{14b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

[Out] $2/9*b^6*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/2)}+14/45*b^4*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(3/2)}+14/15*b^3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{14b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)}\sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^7*(b*\text{Sec}[c + d*x])^{(5/2)}, x]$

[Out] $(14*b^3*\text{EllipticE}[(c + d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (2*b^6*\text{Sin}[c + d*x])/(9*d*(b*\text{Sec}[c + d*x])^{(7/2)}) + (14*b^4*\text{Sin}[c + d*x])/(45*d*(b*\text{Sec}[c + d*x])^{(3/2)})$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticE}[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3769

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_*)]*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^{2*n}), \text{Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \cos^7(c + dx)(b \sec(c + dx))^{5/2} dx &= b^7 \int \frac{1}{(b \sec(c + dx))^{9/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{1}{9} (7b^5) \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{1}{15} (7b^3) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
&= \frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{(7b^3) \int \sqrt{\cos(c + dx)} dx}{15\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
&= \frac{14b^3 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^6 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b^4 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.18, size = 74, normalized size = 0.74

$$\frac{b^2 \sqrt{b \sec(c + dx)} \left((33 \sin(c + dx) + 5 \sin(3(c + dx))) \cos^2(c + dx) + 84 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{90d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^7*(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] (b^2*Sqrt[b*Sec[c + d*x]]*(84*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2]
+ Cos[c + d*x]^2*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)])))/(90*d)
```

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^2 \cos(dx + c)^7 \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*b^2*cos(d*x + c)^7*sec(d*x + c)^2, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^7 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^7, x)

maple [C] time = 1.03, size = 333, normalized size = 3.33

$$\frac{2 \left(\frac{b}{\cos(dx+c)} \right)^{\frac{5}{2}} \left(\cos^2(dx+c) \right) \left(5 \left(\cos^6(dx+c) \right) + 21i \cos(dx+c) \sin(dx+c) \operatorname{EllipticE} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) \sqrt{\frac{1+\cos(dx+c)}{1-\cos(dx+c)}} \right)}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2),x)

[Out] $-2/45/d*(b/\cos(d*x+c))^{5/2}*\cos(d*x+c)^2*(5*\cos(d*x+c)^6+21*I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2})*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-21*I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}+21*I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-21*I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+2*\cos(d*x+c)^4+14*\cos(d*x+c)^2-21*\cos(d*x+c))/\sin(d*x+c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \cos(dx + c)^7 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^7*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(5/2)*cos(d*x + c)^7, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^7 \left(\frac{b}{\cos(c + dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2), x)
```

```
[Out] int(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**7*(b*sec(d*x+c))**(5/2), x)
```

```
[Out] Timed out
```


3.100 $\int (b \sec(c + dx))^{7/2} dx$

Optimal. Leaf size=98

$$-\frac{6b^4 E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{6b^3 \sin(c+dx)\sqrt{b\sec(c+dx)}}{5d} + \frac{2b \sin(c+dx)(b\sec(c+dx))^{5/2}}{5d}$$

[Out] $2/5*b*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/d-6/5*b^4*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*b^3*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.05, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3768, 3771, 2639}

$$\frac{6b^3 \sin(c+dx)\sqrt{b\sec(c+dx)}}{5d} - \frac{6b^4 E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{2b \sin(c+dx)(b\sec(c+dx))^{5/2}}{5d}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(7/2), x]

[Out] $(-6*b^4*\text{EllipticE}[(c+d*x)/2, 2])/((5*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (6*b^3*\text{Sqrt}[b*\text{Sec}[c+d*x]]*\text{Sin}[c+d*x])/(5*d) + (2*b*(b*\text{Sec}[c+d*x])^{(5/2)}*\text{Sin}[c+d*x])/(5*d)$

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int (b \sec(c + dx))^{7/2} dx &= \frac{2b(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d} + \frac{1}{5} (3b^2) \int (b \sec(c + dx))^{3/2} dx \\
&= \frac{6b^3 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2b(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d} - \frac{1}{5} (3b^4) \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
&= \frac{6b^3 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2b(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d} - \frac{(3b^4) \int \sqrt{\cos(c + dx)}}{5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} dx \\
&= -\frac{6b^4 E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{6b^3 \sqrt{b \sec(c + dx)} \sin(c + dx)}{5d} + \frac{2b(b \sec(c + dx))^{5/2} \sin(c + dx)}{5d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 62, normalized size = 0.63

$$\frac{b(b \sec(c + dx))^{5/2} \left(7 \sin(c + dx) + 3 \sin(3(c + dx)) - 12 \cos^2(c + dx) E\left(\frac{1}{2}(c + dx) \middle| 2\right)\right)}{10d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(7/2),x]

[Out] (b*(b*Sec[c + d*x])^(5/2)*(-12*Cos[c + d*x]^(5/2)*EllipticE[(c + d*x)/2, 2] + 7*Sin[c + d*x] + 3*Sin[3*(c + d*x)])/(10*d)

fricas [F] time = 0.80, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{b \sec(dx + c)} b^3 \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(7/2),x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*b^3*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(7/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(7/2), x)

maple [C] time = 0.92, size = 354, normalized size = 3.61

$$2(-1 + \cos(dx + c))^2 \left(3i \left(\cos^3(dx + c) \right) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticE}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) - 3i \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(7/2),x)

[Out] $\frac{2}{5}d(-1 + \cos(dx + c))^2(3I\cos(dx + c)^3\sin(dx + c)(\frac{1}{1 + \cos(dx + c)})^{1/2}(\frac{\cos(dx + c)}{1 + \cos(dx + c)})^{1/2}\operatorname{EllipticE}(I\frac{-1 + \cos(dx + c)}{\sin(dx + c)}, I) - 3I\cos(dx + c)^3\sin(dx + c)(\frac{1}{1 + \cos(dx + c)})^{1/2}(\frac{\cos(dx + c)}{1 + \cos(dx + c)})^{1/2}\operatorname{EllipticF}(I\frac{-1 + \cos(dx + c)}{\sin(dx + c)}, I) + 3I\cos(dx + c)^2\sin(dx + c)(\frac{1}{1 + \cos(dx + c)})^{1/2}(\frac{\cos(dx + c)}{1 + \cos(dx + c)})^{1/2}\operatorname{EllipticE}(I\frac{-1 + \cos(dx + c)}{\sin(dx + c)}, I) - 3I\cos(dx + c)^2\sin(dx + c)(\frac{1}{1 + \cos(dx + c)})^{1/2}(\frac{\cos(dx + c)}{1 + \cos(dx + c)})^{1/2}\operatorname{EllipticF}(I\frac{-1 + \cos(dx + c)}{\sin(dx + c)}, I) - 3\cos(dx + c)^3 + 2\cos(dx + c)^2 + 1)\cos(dx + c)(1 + \cos(dx + c))^{5/2}(b/\cos(dx + c))^{7/2}/\sin(dx + c)^5$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(7/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(7/2),x)

[Out] int((b/cos(c + d*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(7/2),x)
```

```
[Out] Timed out
```

$$3.101 \quad \int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{7/2}}{7b^4d} + \frac{10 \sin(c+dx)(b \sec(c+dx))^{3/2}}{21b^2d} + \frac{10 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21bd}$$

[Out] 10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^2/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^4/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b/d

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{7/2}}{7b^4d} + \frac{10 \sin(c+dx)(b \sec(c+dx))^{3/2}}{21b^2d} + \frac{10 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21bd}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^5/Sqrt[b*Sec[c + d*x]],x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(21*b*d) + (10*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(21*b^2*d) + (2*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*b^4*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sec^5(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\int (b \sec(c+dx))^{9/2} dx}{b^5} \\
&= \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^4 d} + \frac{5 \int (b \sec(c+dx))^{5/2} dx}{7b^3} \\
&= \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^2 d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^4 d} + \frac{5 \int \sqrt{b \sec(c+dx)} dx}{21b} \\
&= \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^2 d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^4 d} + \frac{(5\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)})}{21b} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21bd} + \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^2 d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^4 d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 69, normalized size = 0.69

$$\frac{\sec^3(c+dx) \left(5 \sin(2(c+dx)) + 6 \tan(c+dx) + 10 \cos^{\frac{5}{2}}(c+dx) F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{21d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^5/Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (Sec[c + d*x]^3*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d*Sqrt[b*Sec[c + d*x]])
```

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^4}{b}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^4/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^5}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^5/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.99, size = 152, normalized size = 1.52

$$\frac{2(-1 + \cos(dx + c)) \left(5i \left(\cos^3(dx + c) \right) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) - 5 \right)}{21d \cos(dx + c)^4 \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2), x)

[Out] -2/21/d*(-1+cos(d*x+c))*(5*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)-5*cos(d*x+c)^3+5*cos(d*x+c)^2-3*cos(d*x+c)+3)*(1+cos(d*x+c))^2/cos(d*x+c)^4/(b/cos(d*x+c))^(1/2)/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^5}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(1/2), x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^5/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^5 \sqrt{\frac{b}{\cos(c + dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2)),x)
```

```
[Out] int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(1/2)), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**5/(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Integral(sec(c + d*x)**5/sqrt(b*sec(c + d*x)), x)
```


$$3.102 \quad \int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=97

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{5/2}}{5b^3d} + \frac{6 \sin(c+dx)\sqrt{b \sec(c+dx)}}{5bd} - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] 2/5*(b*sec(d*x+c))^(5/2)*sin(d*x+c)/b^3/d-6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+6/5*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/b/d

Rubi [A] time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{5/2}}{5b^3d} + \frac{6 \sin(c+dx)\sqrt{b \sec(c+dx)}}{5bd} - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^4/Sqrt[b*Sec[c + d*x]], x]

[Out] (-6*EllipticE[(c + d*x)/2, 2])/(5*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (6*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(5*b*d) + (2*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(5*b^3*d)

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\int (b \sec(c+dx))^{7/2} dx}{b^4} \\
&= \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^3d} + \frac{3 \int (b \sec(c+dx))^{3/2} dx}{5b^2} \\
&= \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5bd} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^3d} - \frac{3}{5} \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \\
&= \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5bd} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^3d} - \frac{3 \int \sqrt{\cos(c+dx)} dx}{5\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\
&= -\frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5d\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5bd} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^3d}
\end{aligned}$$

Mathematica [A] time = 0.24, size = 61, normalized size = 0.63

$$\frac{2 \tan(c+dx) \left(\sec^2(c+dx) + 3 \right) - \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{\sqrt{\cos(c+dx)}}}{5d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^4/Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] ((-6*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*(3 + Sec[c + d*x]^2)
*Tan[c + d*x])/(5*d*Sqrt[b*Sec[c + d*x]])
```

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^3}{b}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^3/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/sqrt(b*sec(d*x + c)), x)

maple [C] time = 1.05, size = 356, normalized size = 3.67

$$\frac{2(-1 + \cos(dx + c))^2 \left(3i \left(\cos^3(dx + c) \right) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) - 3 \right)}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2),x)

[Out]
$$-2/5/d*(-1+\cos(d*x+c))^2*(3*I*\cos(d*x+c)^3*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)-3*I*\cos(d*x+c)^3*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)+3*I*\cos(d*x+c)^2*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)-3*I*\cos(d*x+c)^2*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^(1/2)*(\cos(d*x+c)/(1+\cos(d*x+c)))^(1/2)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)+3*\cos(d*x+c)^3-2*\cos(d*x+c)^2-1*(1+\cos(d*x+c))^2/\sin(d*x+c)^5/\cos(d*x+c)^3/(b/\cos(d*x+c))^(1/2)$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^4/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^4 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2)), x)

[Out] int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**4/(b*sec(d*x+c))**(1/2), x)

[Out] Integral(sec(c + d*x)**4/sqrt(b*sec(c + d*x)), x)

$$3.103 \quad \int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{3/2}}{3b^2d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3bd}$$

[Out] $2/3*(b*\sec(d*x+c))^{3/2}*\sin(d*x+c)/b^2/d+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b/d$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{3/2}}{3b^2d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3bd}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/Sqrt[b*Sec[c + d*x]],x]

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*b*d) + (2*(b*\text{Sec}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(3*b^2*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c + dx)}{\sqrt{b \sec(c + dx)}} dx &= \frac{\int (b \sec(c + dx))^{5/2} dx}{b^3} \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3b^2 d} + \frac{\int \sqrt{b \sec(c + dx)} dx}{3b} \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3b^2 d} + \frac{(\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b} \\ &= \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3bd} + \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3b^2 d} \end{aligned}$$

Mathematica [A] time = 0.08, size = 51, normalized size = 0.71

$$\frac{2\sqrt{b \sec(c + dx)} \left(\tan(c + dx) + \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/Sqrt[b*Sec[c + d*x]], x]

[Out] (2*Sqrt[b*Sec[c + d*x]]*(Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan[c + d*x]))/(3*b*d)

fricas [F] time = 1.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \sec(dx + c)^2}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^2/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.89, size = 130, normalized size = 1.81

$$\frac{2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - \cos(dx + c) \right)}{3d \sin(dx + c)^3 \cos(dx + c)^2 \sqrt{\frac{b}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x)

[Out] -2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c))))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*(1+cos(d*x+c))^2/sin(d*x+c)^3/cos(d*x+c)^2/(b/cos(d*x+c))^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^3 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2)),x)

[Out] int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**3/(b*sec(d*x+c))**(1/2), x)
```

```
[Out] Integral(sec(c + d*x)**3/sqrt(b*sec(c + d*x)), x)
```


$$3.104 \quad \int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=65

$$\frac{2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{bd} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

[Out] $-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b/d$

Rubi [A] time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{bd} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/Sqrt[b*Sec[c + d*x]], x]

[Out] $(-2*\text{EllipticE}[(c+d*x)/2, 2])/(d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (2*\text{Sqrt}[b*\text{Sec}[c+d*x]]*\text{Sin}[c+d*x])/(b*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\int (b \sec(c+dx))^{3/2} dx}{b^2} \\ &= \frac{2\sqrt{b \sec(c+dx)} \sin(c+dx)}{bd} - \int \frac{1}{\sqrt{b \sec(c+dx)}} dx \\ &= \frac{2\sqrt{b \sec(c+dx)} \sin(c+dx)}{bd} - \frac{\int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\ &= -\frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2\sqrt{b \sec(c+dx)} \sin(c+dx)}{bd} \end{aligned}$$

Mathematica [A] time = 0.08, size = 48, normalized size = 0.74

$$\frac{2 \tan(c+dx) - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{\sqrt{\cos(c+dx)}}}{d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^2/Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] ((-2*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*Tan[c + d*x])/(d*Sqr
t[b*Sec[c + d*x]])
```

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)/b, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.96, size = 319, normalized size = 4.91

$$\frac{2(1 + \cos(dx+c))^2(-1 + \cos(dx+c))^2 \left(i \cos(dx+c) \sin(dx+c) \operatorname{EllipticE}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} \sqrt{\frac{1}{1-\cos(dx+c)}} \right)}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x)

[Out] 2/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*cos(d*x+c)*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*(b/cos(d*x+c))^(1/2)/b/sin(d*x+c)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c+dx)^2 \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2)), x)`

[Out] `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(1/2), x)`

[Out] `Integral(sec(c + d*x)**2/sqrt(b*sec(c + d*x)), x)`

$$3.105 \quad \int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=41

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{bd}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b/d$

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3771, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{bd}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/Sqrt[b*Sec[c + d*x]], x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\int \sqrt{b \sec(c+dx)} dx}{b} \\ &= \frac{(\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{bd} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/Sqrt[b*Sec[c + d*x]], x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b*d)

fricas [F] time = 0.61, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)}}{b}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.85, size = 98, normalized size = 2.39

$$\frac{2i \left(\frac{1}{1+\cos(dx+c)} \right)^{\frac{3}{2}} (-1 + \cos(dx+c)) \operatorname{EllipticF} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) (1 + \cos(dx+c))^2}{d \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \sin(dx+c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)/(b*sec(d*x+c))^(1/2), x)`

[Out] `-2*I/d*(1/(1+cos(d*x+c)))^(3/2)*(-1+cos(d*x+c))*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1+cos(d*x+c))^2/(b/cos(d*x+c))^(1/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)/sin(d*x+c)^2`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/2), x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)/sqrt(b*sec(d*x + c)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c+dx) \sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/2)), x)`

[Out] `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)/(b*sec(d*x+c))**(1/2), x)`

[Out] `Integral(sec(c + d*x)/sqrt(b*sec(c + d*x)), x)`

$$3.106 \quad \int \frac{1}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=38

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3771, 2639}

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[b*Sec[c + d*x]],x]

[Out] (2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \sec(c+dx)}} dx &= \frac{\int \sqrt{\cos(c+dx)} dx}{\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \\ &= \frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 38, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[b*Sec[c + d*x]], x]

[Out] (2*EllipticE[(c + d*x)/2, 2])/(d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b\sec(dx+c)}}{b\sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b\sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(1/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.90, size = 306, normalized size = 8.05

$$2\left(i\sin(dx+c)\cos(dx+c)\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\text{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right)\sqrt{\frac{1}{1+\cos(dx+c)}} - i\cos(dx+c)\sin(dx+c)\right)E$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(d*x+c))^(1/2), x)

[Out] 2/d*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)-I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)*(cos

$(d*x+c)/(1+\cos(d*x+c))^{1/2}+I*\sin(d*x+c)*\text{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*\text{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-\cos(d*x+c)^2+\cos(d*x+c))*(b/\cos(d*x+c))^{1/2}/\sin(d*x+c)/b$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b/cos(c + d*x))^(1/2),x)

[Out] int(1/(b/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))**(1/2),x)

[Out] Integral(1/sqrt(b*sec(c + d*x)), x)

$$3.107 \quad \int \frac{\cos(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=69

$$\frac{2 \sin(c+dx)}{3d\sqrt{b \sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3bd}$$

[Out] $2/3*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b/d$

Rubi [A] time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2 \sin(c+dx)}{3d\sqrt{b \sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3bd}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/Sqrt[b*Sec[c + d*x]],x]

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*b*d) + (2*\text{Sin}[c + d*x])/(3*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d^n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= b \int \frac{1}{(b \sec(c+dx))^{3/2}} dx \\ &= \frac{2 \sin(c+dx)}{3d \sqrt{b \sec(c+dx)}} + \frac{\int \sqrt{b \sec(c+dx)} dx}{3b} \\ &= \frac{2 \sin(c+dx)}{3d \sqrt{b \sec(c+dx)}} + \frac{(\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b} \\ &= \frac{2 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3bd} + \frac{2 \sin(c+dx)}{3d \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.08, size = 60, normalized size = 0.87

$$\frac{b \sec^2(c+dx) \left(\sin(2(c+dx)) + 2 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{3d(b \sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]/Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (b*Sec[c + d*x]^2*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*d*(b*Sec[c + d*x])^(3/2))
```

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx+c)} \cos(dx+c)}{b \sec(dx+c)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)/(b*sec(d*x + c)), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/sqrt(b*sec(d*x + c)), x)

maple [C] time = 1.02, size = 126, normalized size = 1.83

$$\frac{2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \operatorname{EllipticF} \left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i \right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} - (\cos^2(dx + c)) + c \right)}{3d \sin(dx + c)^3 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x)

[Out] -2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)^2+cos(d*x+c))*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(1/2)/sin(d*x+c)^3/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)}{\sqrt{\frac{b}{\cos(c + dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b/cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)/(b/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)/(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Integral(cos(c + d*x)/sqrt(b*sec(c + d*x)), x)
```

$$3.108 \quad \int \frac{\cos^2(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=67

$$\frac{2b \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] $2/5*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(3/2)}+6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/Sqrt[b*Sec[c + d*x]], x]

[Out] $(6*\text{EllipticE}[(c+d*x)/2, 2])/(5*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (2*b*\text{Sin}[c+d*x])/(5*d*(b*\text{Sec}[c+d*x])^{(3/2)})$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx &= b^2 \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\ &= \frac{2b \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{3}{5} \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\ &= \frac{2b \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} + \frac{3 \int \sqrt{\cos(c + dx)} dx}{5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5d\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b \sin(c + dx)}{5d(b \sec(c + dx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 60, normalized size = 0.90

$$\frac{\sqrt{b \sec(c + dx)} \left(\sin(c + dx) + \sin(3(c + dx)) + 12\sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10bd}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^2/Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*b*d)
```

fricas [F] time = 0.55, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \cos(dx + c)^2}{b \sec(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^2/(b*sec(d*x + c)), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/sqrt(b*sec(d*x + c)), x)

maple [C] time = 1.17, size = 318, normalized size = 4.75

$$2 \left(3i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - 3i \cos(dx + c) \sin(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x)

[Out] 2/5/d*(3*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-3*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+3*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)^4-2*cos(d*x+c)^2+3*cos(d*x+c))*(b/cos(d*x+c))^(1/2)/sin(d*x+c)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^2}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/2),x)

[Out] int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(1/2), x)

[Out] Integral(cos(c + d*x)**2/sqrt(b*sec(c + d*x)), x)

$$3.109 \quad \int \frac{\cos^3(c+dx)}{\sqrt{b} \sec(c+dx)} dx$$

Optimal. Leaf size=97

$$\frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21d\sqrt{b} \sec(c+dx)} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b} \sec(c+dx)}{21bd}$$

[Out] $2/7*b^2*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/2)}+10/21*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b/d$

Rubi [A] time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21d\sqrt{b} \sec(c+dx)} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b} \sec(c+dx)}{21bd}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3/Sqrt[b*Sec[c + d*x]], x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*b*d) + (2*b^2*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(5/2)}) + (10*\text{Sin}[c + d*x])/(21*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= b^3 \int \frac{1}{(b \sec(c+dx))^{7/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{1}{7}(5b) \int \frac{1}{(b \sec(c+dx))^{3/2}} dx \\
&= \frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21d\sqrt{b \sec(c+dx)}} + \frac{5 \int \sqrt{b \sec(c+dx)} dx}{21b} \\
&= \frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21d\sqrt{b \sec(c+dx)}} + \frac{(5\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b} \\
&= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21bd} + \frac{2b^2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21d\sqrt{b \sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 66, normalized size = 0.68

$$\frac{\sqrt{b \sec(c+dx)} \left(26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) + 40 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{84bd}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^3/Sqrt[b*Sec[c + d*x]], x]
```

```
[Out] (Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26
*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b*d)
```

fricas [F] time = 0.77, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx+c)} \cos(dx+c)^3}{b \sec(dx+c)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")
```

[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^3/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^3/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.98, size = 148, normalized size = 1.53

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(5i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \right)}{21d \sin(dx + c)^3 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*cos(d*x+c)^4+3*cos(d*x+c)^3-5*cos(d*x+c)^2+5*cos(d*x+c))*(b/cos(d*x+c))^(1/2)/sin(d*x+c)^3/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^3}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^3/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^3}{\sqrt{\frac{b}{\cos(c + dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(c + d*x)^3/(b/cos(c + d*x))^(1/2),x)
```

```
[Out] int(cos(c + d*x)^3/(b/cos(c + d*x))^(1/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)**3/(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

$$3.110 \quad \int \frac{\cos^4(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=95

$$\frac{2b^3 \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14b \sin(c+dx)}{45d(b \sec(c+dx))^{3/2}} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] $2/9*b^3*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/2)}+14/45*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(3/2)}+14/15*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^3 \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14b \sin(c+dx)}{45d(b \sec(c+dx))^{3/2}} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^4/Sqrt[b*Sec[c + d*x]], x]

[Out] $(14*\text{EllipticE}[(c+d*x)/2, 2])/(15*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (2*b^3*\text{Sin}[c+d*x])/(9*d*(b*\text{Sec}[c+d*x])^{(7/2)}) + (14*b*\text{Sin}[c+d*x])/(45*d*(b*\text{Sec}[c+d*x])^{(3/2)})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1)/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\cos^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx &= b^4 \int \frac{1}{(b \sec(c + dx))^{9/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{1}{9} (7b^2) \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{7}{15} \int \frac{1}{\sqrt{b \sec(c + dx)}} dx \\
 &= \frac{2b^3 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{7 \int \sqrt{\cos(c + dx)} dx}{15 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 &= \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^3 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14b \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] time = 0.20, size = 70, normalized size = 0.74

$$\frac{4(33 \sin(c + dx) + 5 \sin(3(c + dx))) \cos(c + dx) + \frac{336E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{\sqrt{\cos(c + dx)}}}{360d \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^4/Sqrt[b*Sec[c + d*x]], x]

[Out] ((336*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 4*Cos[c + d*x]*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)]))/(360*d*Sqrt[b*Sec[c + d*x]])

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \cos(dx + c)^4}{b \sec(dx + c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^4/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^4}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^4/sqrt(b*sec(d*x + c)), x)

maple [C] time = 0.99, size = 328, normalized size = 3.45

$$2 \left(21i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - 21i \cos(dx + c) \sin(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2),x)

[Out] 2/45/d*(21*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-21*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-5*cos(d*x+c)^6+21*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-21*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-2*cos(d*x+c)^4-14*cos(d*x+c)^2+21*cos(d*x+c))*(b/cos(d*x+c))^(1/2)/sin(d*x+c)/b

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^4}{\sqrt{b \sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^4/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^4/sqrt(b*sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^4}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^4/(b/cos(c + d*x))^(1/2), x)`

[Out] `int(cos(c + d*x)^4/(b/cos(c + d*x))^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^4(c + dx)}{\sqrt{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**4/(b*sec(d*x+c))**(1/2), x)`

[Out] `Integral(cos(c + d*x)**4/sqrt(b*sec(c + d*x)), x)`

$$3.111 \quad \int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{7/2}}{7b^5d} + \frac{10 \sin(c+dx)(b \sec(c+dx))^{3/2}}{21b^3d} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^2d}$$

[Out] 10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^3/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^5/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{7/2}}{7b^5d} + \frac{10 \sin(c+dx)(b \sec(c+dx))^{3/2}}{21b^3d} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^2d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^6/(b*Sec[c + d*x])^(3/2),x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(21*b^2*d) + (10*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(21*b^3*d) + (2*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*b^5*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> -Simp[(b*Csc[c + d*x])*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{3/2}} dx &= \frac{\int (b \sec(c + dx))^{9/2} dx}{b^6} \\ &= \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^5 d} + \frac{5 \int (b \sec(c + dx))^{5/2} dx}{7b^4} \\ &= \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21b^3 d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^5 d} + \frac{5 \int \sqrt{b \sec(c + dx)}}{21b^2} \\ &= \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21b^3 d} + \frac{2(b \sec(c + dx))^{7/2} \sin(c + dx)}{7b^5 d} + \frac{(5\sqrt{\cos(c + dx)} \sqrt{b})}{21b^2} \\ &= \frac{10\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21b^2 d} + \frac{10(b \sec(c + dx))^{3/2} \sin(c + dx)}{21b^3 d} + \frac{2}{21b^2} \end{aligned}$$

Mathematica [A] time = 0.17, size = 69, normalized size = 0.69

$$\frac{\sec^4(c + dx) \left(5 \sin(2(c + dx)) + 6 \tan(c + dx) + 10 \cos^{\frac{5}{2}}(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{21d(b \sec(c + dx))^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^6/(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] (Sec[c + d*x]^4*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] + 5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*d*(b*Sec[c + d*x])^(3/2))
```

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \sec(dx + c)^4}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^4/b^2, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^6}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.89, size = 152, normalized size = 1.52

$$\frac{2(1 + \cos(dx+c))^2(-1 + \cos(dx+c)) \left(5i(\cos^3(dx+c)) \sin(dx+c) \sqrt{\frac{1}{1+\cos(dx+c)}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i}{\sqrt{1+\cos(dx+c)}}\right) \right)}{21d \sin(dx+c)^3 \cos(dx+c)^5 \left(\frac{b}{\cos(dx+c)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x)

[Out] $-2/21/d*(1+\cos(d*x+c))^2*(-1+\cos(d*x+c))*(5*I*\cos(d*x+c)^3*\sin(d*x+c)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c),I)-5*\cos(d*x+c)^3+5*\cos(d*x+c)^2-3*\cos(d*x+c)+3)/\sin(d*x+c)^3/\cos(d*x+c)^5/(b/\cos(d*x+c))^{3/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^6}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^6 \left(\frac{b}{\cos(c+dx)} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2)), x)`

[Out] `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(3/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**6/(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral(sec(c + d*x)**6/(b*sec(c + d*x))**(3/2), x)`

$$3.112 \quad \int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{5/2}}{5b^4d} + \frac{6 \sin(c+dx)\sqrt{b \sec(c+dx)}}{5b^2d} - \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b^4/d-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/b/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b^2/d$

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{5/2}}{5b^4d} + \frac{6 \sin(c+dx)\sqrt{b \sec(c+dx)}}{5b^2d} - \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^5/(b*Sec[c + d*x])^(3/2), x]

[Out] $(-6*\text{EllipticE}[(c+d*x)/2, 2])/(5*b*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (6*\text{Sqrt}[b*\text{Sec}[c+d*x]]*\text{Sin}[c+d*x])/(5*b^2*d) + (2*(b*\text{Sec}[c+d*x])^{(5/2)}*\text{Sin}[c+d*x])/(5*b^4*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{3/2}} dx &= \frac{\int (b \sec(c+dx))^{7/2} dx}{b^5} \\
&= \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^4d} + \frac{3 \int (b \sec(c+dx))^{3/2} dx}{5b^3} \\
&= \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5b^2d} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^4d} - \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b} \\
&= \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5b^2d} + \frac{2(b \sec(c+dx))^{5/2} \sin(c+dx)}{5b^4d} - \frac{3 \int \sqrt{\cos(c+dx)}}{5b\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} \\
&= -\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{6\sqrt{b \sec(c+dx)} \sin(c+dx)}{5b^2d} + \frac{2(b \sec(c+dx))^{5/2}}{5b^4d}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 64, normalized size = 0.64

$$\frac{2 \tan(c+dx) \left(\sec^2(c+dx) + 3 \right) - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{\sqrt{\cos(c+dx)}}}{5bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^5/(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] ((-6*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*(3 + Sec[c + d*x]^2)
*Tan[c + d*x])/(5*b*d*Sqrt[b*Sec[c + d*x]])
```

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^3}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```


[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^3/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^5}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.96, size = 356, normalized size = 3.56

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c))^2 \left(3i(\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF} \left(\right. \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x)

[Out] -2/5/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)+3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)+3*cos(d*x+c)^3-2*cos(d*x+c)^2-1)/sin(d*x+c)^5/cos(d*x+c)^4/(b/cos(d*x+c))^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^5}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^5 \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2)), x)

[Out] int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(c + dx)}{(b \sec(c + dx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**5/(b*sec(d*x+c))**(3/2), x)

[Out] Integral(sec(c + d*x)**5/(b*sec(c + d*x))**(3/2), x)

$$3.113 \quad \int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{3/2}}{3b^3d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^2d}$$

[Out] $2/3*(b*\sec(d*x+c))^{(3/2)}*\sin(d*x+c)/b^3/d+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^2/d$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{3/2}}{3b^3d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^2d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^4/(b*Sec[c + d*x])^(3/2), x]

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*b^2*d) + (2*(b*\text{Sec}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(3*b^3*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{3/2}} dx &= \frac{\int (b \sec(c + dx))^{5/2} dx}{b^4} \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3b^3 d} + \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} \\ &= \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3b^3 d} + \frac{(\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} \\ &= \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3b^2 d} + \frac{2(b \sec(c + dx))^{3/2} \sin(c + dx)}{3b^3 d} \end{aligned}$$

Mathematica [A] time = 0.09, size = 56, normalized size = 0.78

$$\frac{2 \sec^3(c + dx) \left(\sin(c + dx) + \cos^{\frac{3}{2}}(c + dx) F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d(b \sec(c + dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^4/(b*Sec[c + d*x])^(3/2), x]

[Out] (2*Sec[c + d*x]^3*(Cos[c + d*x]^(3/2)*EllipticF[(c + d*x)/2, 2] + Sin[c + d*x]))/(3*d*(b*Sec[c + d*x])^(3/2))

fricas [F] time = 0.59, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \sec(dx + c)^2}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^2/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^4}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.86, size = 125, normalized size = 1.74

$$\frac{2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - \cos(dx + c) \right)}{3db^3 \sin(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2),x)

[Out] -2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*(1+cos(d*x+c))^2*(b/cos(d*x+c))^(3/2)/b^3/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^4}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^4 \left(\frac{b}{\cos(c+dx)} \right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2)),x)

[Out] int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(3/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**4/(b*sec(d*x+c))**(3/2), x)
```

```
[Out] Integral(sec(c + d*x)**4/(b*sec(c + d*x))**(3/2), x)
```

$$3.114 \quad \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=68

$$\frac{2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{b^2 d} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

[Out] $-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/b/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b^2/d$

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{b^2 d} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/(b*Sec[c + d*x])^(3/2), x]

[Out] $(-2*\text{EllipticE}[(c+d*x)/2, 2])/ (b*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (2*\text{Sqrt}[b*\text{Sec}[c+d*x]]*\text{Sin}[c+d*x])/ (b^2*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> -Simp[(b*Csc[c + d*x])*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx &= \frac{\int (b \sec(c + dx))^{3/2} dx}{b^3} \\ &= \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{b^2 d} - \frac{\int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{b} \\ &= \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{b^2 d} - \frac{\int \sqrt{\cos(c + dx)} dx}{b\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= -\frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{bd\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{b^2 d} \end{aligned}$$

Mathematica [A] time = 0.06, size = 51, normalized size = 0.75

$$\frac{2 \tan(c + dx) - \frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{\sqrt{\cos(c + dx)}}}{bd\sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^3/(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] ((-2*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*Tan[c + d*x])/(b*d*S
qrt[b*Sec[c + d*x]])
```

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \sec(dx + c)}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)/b^2, x)
```


giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 1.02, size = 324, normalized size = 4.76

$$2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(i \cos(dx + c) \sin(dx + c) \operatorname{EllipticE}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{1}{1 - \cos(dx + c)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x)

[Out] 2/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*cos(d*x+c)*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)/cos(d*x+c)^2/sin(d*x+c)^5/(b/cos(d*x+c))^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^3 \left(\frac{b}{\cos(c + dx)} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2)), x)`

[Out] `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(3/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**3/(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral(sec(c + d*x)**3/(b*sec(c + d*x))**(3/2), x)`

$$3.115 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=41

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^2 d}$$

[Out] 2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3771, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^2 d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^2*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx &= \frac{\int \sqrt{b \sec(c+dx)} dx}{b^2} \\ &= \frac{(\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^2} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^2 d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^2 d}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^2*d)

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)}}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.74, size = 98, normalized size = 2.39

$$\frac{2i \left(\frac{1}{1+\cos(dx+c)} \right)^{\frac{5}{2}} (-1 + \cos(dx+c)) \operatorname{EllipticF} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) (1 + \cos(dx+c))^2}{d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{3}{2}} \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{\frac{3}{2}} \sin(dx+c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2), x)`

[Out] `-2*I/d*(1/(1+cos(d*x+c)))^(5/2)*(-1+cos(d*x+c))*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1+cos(d*x+c))^2/(b/cos(d*x+c))^(3/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(3/2)/sin(d*x+c)^2`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c+dx)^2 \left(\frac{b}{\cos(c+dx)} \right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2)), x)`

[Out] `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(3/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(3/2), x)`

$$3.116 \quad \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=41

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{bd\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/b/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3771, 2639}

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{bd\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Sec[c + d*x])^(3/2), x]

[Out] (2*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{3/2}} dx &= \frac{\int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b} \\ &= \frac{\int \sqrt{\cos(c+dx)} dx}{b \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\ &= \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 41, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{bd \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(3/2), x]

[Out] (2*EllipticE[(c + d*x)/2, 2])/(b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)}}{b^2 \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/(b^2*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.78, size = 311, normalized size = 7.59

$$2 \left(-i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} + i \cos(dx + c) \sin(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)/(b*sec(d*x+c))^(3/2), x)`

[Out] $-2/d * (-I * \cos(d*x+c) * \sin(d*x+c) * \operatorname{EllipticF}(I * (-1 + \cos(d*x+c)) / \sin(d*x+c), I) * (1 / (1 + \cos(d*x+c)))^{1/2} * (\cos(d*x+c) / (1 + \cos(d*x+c)))^{1/2} + I * \cos(d*x+c) * \sin(d*x+c) * \operatorname{EllipticE}(I * (-1 + \cos(d*x+c)) / \sin(d*x+c), I) * (1 / (1 + \cos(d*x+c)))^{1/2} * (\cos(d*x+c) / (1 + \cos(d*x+c)))^{1/2} - I * \sin(d*x+c) * \operatorname{EllipticF}(I * (-1 + \cos(d*x+c)) / \sin(d*x+c), I) * (1 / (1 + \cos(d*x+c)))^{1/2} * (\cos(d*x+c) / (1 + \cos(d*x+c)))^{1/2} + I * \sin(d*x+c) * \operatorname{EllipticE}(I * (-1 + \cos(d*x+c)) / \sin(d*x+c), I) * (1 / (1 + \cos(d*x+c)))^{1/2} * (\cos(d*x+c) / (1 + \cos(d*x+c)))^{1/2} + \cos(d*x+c)^2 - \cos(d*x+c) / \cos(d*x+c)^2 / (b / \cos(d*x+c))^{3/2} / \sin(d*x+c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)/(b*sec(d*x+c))^(3/2), x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)/(b*sec(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) \left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(3/2)), x)`

[Out] `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(3/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)/(b*sec(d*x+c))**(3/2),x)
```

```
[Out] Integral(sec(c + d*x)/(b*sec(c + d*x))**(3/2), x)
```

$$3.117 \quad \int \frac{1}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=72

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^2d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}$$

[Out] 2/3*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^2/d

Rubi [A] time = 0.03, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3769, 3771, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^2d} + \frac{2 \sin(c+dx)}{3bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(-3/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^2*d) + (2*Sin[c + d*x])/(3*b*d*Sqrt[b*Sec[c + d*x]])

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(b \sec(c + dx))^{3/2}} dx &= \frac{2 \sin(c + dx)}{3bd\sqrt{b \sec(c + dx)}} + \frac{\int \sqrt{b \sec(c + dx)} dx}{3b^2} \\
&= \frac{2 \sin(c + dx)}{3bd\sqrt{b \sec(c + dx)}} + \frac{(\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{3b^2} \\
&= \frac{2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{3b^2 d} + \frac{2 \sin(c + dx)}{3bd\sqrt{b \sec(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 59, normalized size = 0.82

$$\frac{\sec^2(c + dx) \left(\sin(2(c + dx)) + 2\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{3d(b \sec(c + dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(-3/2), x]

[Out] (Sec[c + d*x]^2*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*d*(b*Sec[c + d*x])^(3/2))

fricas [F] time = 0.84, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)}}{b^2 \sec(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/(b^2*sec(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(-3/2), x)

maple [C] time = 0.79, size = 131, normalized size = 1.82

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} - \left(\frac{b}{\cos(dx + c)} \right)^{\frac{3}{2}} \right)}{3d \sin(dx + c)^3 \cos(dx + c)^2 \left(\frac{b}{\cos(dx + c)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(d*x+c))^(3/2), x)

[Out] -2/3/d*(1+cos(d*x+c))²*(-1+cos(d*x+c))*(I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)²+cos(d*x+c))/sin(d*x+c)³/cos(d*x+c)²/(b/cos(d*x+c))^(3/2))

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(3/2), x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(-3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b/cos(c + d*x))^(3/2), x)

[Out] int(1/(b/cos(c + d*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(d*x+c))**(3/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**(-3/2), x)
```

$$3.118 \quad \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=69

$$\frac{2 \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] 2/5*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2 \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} + \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Sec[c + d*x])^(3/2),x]

[Out] (6*EllipticE[(c + d*x)/2, 2])/(5*b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*d*(b*Sec[c + d*x])^(3/2))

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{3/2}} dx &= b \int \frac{1}{(b \sec(c+dx))^{5/2}} dx \\ &= \frac{2 \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} + \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b} \\ &= \frac{2 \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} + \frac{3 \int \sqrt{\cos(c+dx)} dx}{5b \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\ &= \frac{6E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{5bd \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{5d(b \sec(c+dx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 60, normalized size = 0.87

$$\frac{\sqrt{b \sec(c+dx)} \left(\sin(c+dx) + \sin(3(c+dx)) + 12 \sqrt{\cos(c+dx)} E\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{10b^2d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(3/2), x]
```

```
[Out] (Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*b^2*d)
```

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)} \cos(dx+c)}{b^2 \sec(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)/(b^2*sec(d*x + c)^2), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(3/2), x)

maple [C] time = 0.95, size = 323, normalized size = 4.68

$$\frac{6i \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}}}{5} - \frac{6i \cos(dx+c) \sin(dx+c) \operatorname{EllipticE}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}}}{5} \sqrt{\frac{1}{1+\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x)

[Out] 2/5/d*(3*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-3*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+3*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)^4-2*cos(d*x+c)^2+3*cos(d*x+c))/sin(d*x+c)/cos(d*x+c)^2/(b/cos(d*x+c))^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c+dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b/cos(c + d*x))^(3/2),x)

[Out] `int(cos(c + d*x)/(b/cos(c + d*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral(cos(c + d*x)/(b*sec(c + d*x))**(3/2), x)`

$$3.119 \quad \int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=98

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^2d} + \frac{10 \sin(c+dx)}{21bd\sqrt{b \sec(c+dx)}} + \frac{2b \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}}$$

[Out] $2/7*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/2)}+10/21*\sin(d*x+c)/b/d/(b*\sec(d*x+c))^{(1/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^2/d$

Rubi [A] time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2641}

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^2d} + \frac{10 \sin(c+dx)}{21bd\sqrt{b \sec(c+dx)}} + \frac{2b \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(3/2), x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*b^2*d) + (2*b*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(5/2)}) + (10*\text{Sin}[c + d*x])/(21*b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{3/2}} dx &= b^2 \int \frac{1}{(b \sec(c + dx))^{7/2}} dx \\
 &= \frac{2b \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{5}{7} \int \frac{1}{(b \sec(c + dx))^{3/2}} dx \\
 &= \frac{2b \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21bd\sqrt{b \sec(c + dx)}} + \frac{5 \int \sqrt{b \sec(c + dx)} dx}{21b^2} \\
 &= \frac{2b \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21bd\sqrt{b \sec(c + dx)}} + \frac{(5\sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21b^2} \\
 &= \frac{10\sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21b^2d} + \frac{2b \sin(c + dx)}{7d(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21bd\sqrt{b \sec(c + dx)}}
 \end{aligned}$$

Mathematica [A] time = 0.10, size = 66, normalized size = 0.67

$$\frac{\sqrt{b \sec(c + dx)} \left(26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) + 40 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{84b^2d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(3/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b^2*d)

fricas [F] time = 0.82, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx + c)} \cos(dx + c)^2}{b^2 \sec(dx + c)^2}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] $\text{integral}(\sqrt{b \sec(dx + c)} \cos(dx + c)^2 / (b^2 \sec(dx + c)^2), x)$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\cos(dx+c)^2/(b \sec(dx+c))^{3/2}, x, \text{algorithm}="giac")$

[Out] $\text{integrate}(\cos(dx + c)^2 / (b \sec(dx + c))^{3/2}, x)$

maple [C] time = 0.90, size = 153, normalized size = 1.56

$$\frac{2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c)) \left(5i \sin(dx + c) \text{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} - \right)}{21d \sin(dx + c)^3 \cos(dx + c)^2 \left(\frac{b}{\cos(dx + c)}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(\cos(dx+c)^2/(b \sec(dx+c))^{3/2}, x)$

[Out] $-2/21/d*(1+\cos(dx+c))^{2*(-1+\cos(dx+c))}*(5*I*\sin(dx+c)*\text{EllipticF}(I*(-1+\cos(dx+c))/\sin(dx+c), I)*(1/(1+\cos(dx+c)))^{1/2}*(\cos(dx+c)/(1+\cos(dx+c)))^{1/2}-3*\cos(dx+c)^4+3*\cos(dx+c)^3-5*\cos(dx+c)^2+5*\cos(dx+c))/\sin(dx+c)^3/\cos(dx+c)^2/(b/\cos(dx+c))^{3/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\cos(dx+c)^2/(b \sec(dx+c))^{3/2}, x, \text{algorithm}="maxima")$

[Out] $\text{integrate}(\cos(dx + c)^2 / (b \sec(dx + c))^{3/2}, x)$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^2}{\left(\frac{b}{\cos(c + dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2/(b/cos(c + d*x))^(3/2), x)`

[Out] `int(cos(c + d*x)^2/(b/cos(c + d*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(3/2), x)`

$$3.120 \quad \int \frac{\cos^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=97

$$\frac{2b^2 \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14 \sin(c+dx)}{45d(b \sec(c+dx))^{3/2}} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] 2/9*b^2*sin(d*x+c)/d/(b*sec(d*x+c))^(7/2)+14/45*sin(d*x+c)/d/(b*sec(d*x+c))^(3/2)+14/15*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{2b^2 \sin(c+dx)}{9d(b \sec(c+dx))^{7/2}} + \frac{14 \sin(c+dx)}{45d(b \sec(c+dx))^{3/2}} + \frac{14E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{15bd\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^3/(b*Sec[c + d*x])^(3/2), x]

[Out] (14*EllipticE[(c + d*x)/2, 2])/(15*b*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*b^2*Sin[c + d*x])/(9*d*(b*Sec[c + d*x])^(7/2)) + (14*Sin[c + d*x])/(45*d*(b*Sec[c + d*x])^(3/2))

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned}
 \int \frac{\cos^3(c + dx)}{(b \sec(c + dx))^{3/2}} dx &= b^3 \int \frac{1}{(b \sec(c + dx))^{9/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{1}{9}(7b) \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{7 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{15b} \\
 &= \frac{2b^2 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}} + \frac{7 \int \sqrt{\cos(c + dx)} dx}{15b \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
 &= \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15bd \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b^2 \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14 \sin(c + dx)}{45d(b \sec(c + dx))^{3/2}}
 \end{aligned}$$

Mathematica [A] time = 0.08, size = 73, normalized size = 0.75

$$\frac{84E\left(\frac{1}{2}(c + dx) \middle| 2\right) + (33 \sin(c + dx) + 5 \sin(3(c + dx))) \cos^{\frac{3}{2}}(c + dx)}{90d \cos^{\frac{3}{2}}(c + dx) (b \sec(c + dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Integrate[Cos[c + d*x]^3/(b*Sec[c + d*x])^(3/2), x]`

[Out] `(84*EllipticE[(c + d*x)/2, 2] + Cos[c + d*x]^(3/2)*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)]))/(90*d*Cos[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2))`

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \cos(dx + c)^3}{b^2 \sec(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")`

[Out] `integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^3/(b^2*sec(d*x + c)^2), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^3}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2), x, algorithm="giac")`

[Out] `integrate(cos(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)`

maple [C] time = 0.98, size = 333, normalized size = 3.43

$$\frac{14i \sin(dx+c) \cos(dx+c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}}}{15} - \frac{14i \cos(dx+c) \sin(dx+c) \operatorname{EllipticE}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}}}{15}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2), x)`

[Out] `2/45/d*(21*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)-21*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-5*cos(d*x+c)^6+21*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-21*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-2*cos(d*x+c)^4-14*cos(d*x+c)^2+21*cos(d*x+c))/sin(d*x+c)/cos(d*x+c)^2/(b/cos(d*x+c))^(3/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^3}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3/(b*sec(d*x+c))^(3/2), x, algorithm="maxima")`

[Out] `integrate(cos(d*x + c)^3/(b*sec(d*x + c))^(3/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^3}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^3/(b/cos(c + d*x))^(3/2), x)

[Out] int(cos(c + d*x)^3/(b/cos(c + d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**3/(b*sec(d*x+c))**(3/2), x)

[Out] Timed out

$$3.121 \quad \int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{7/2}}{7b^6d} + \frac{10 \sin(c+dx)(b \sec(c+dx))^{3/2}}{21b^4d} + \frac{10 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^3d}$$

[Out] 10/21*(b*sec(d*x+c))^(3/2)*sin(d*x+c)/b^4/d+2/7*(b*sec(d*x+c))^(7/2)*sin(d*x+c)/b^6/d+10/21*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{7/2}}{7b^6d} + \frac{10 \sin(c+dx)(b \sec(c+dx))^{3/2}}{21b^4d} + \frac{10 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^3d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^7/(b*Sec[c + d*x])^(5/2), x]

[Out] (10*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(21*b^3*d) + (10*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(21*b^4*d) + (2*(b*Sec[c + d*x])^(7/2)*Sin[c + d*x])/(7*b^6*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec^7(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\int (b \sec(c+dx))^{9/2} dx}{b^7} \\ &= \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^6d} + \frac{5 \int (b \sec(c+dx))^{5/2} dx}{7b^5} \\ &= \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^4d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^6d} + \frac{5 \int \sqrt{b \sec(c+dx)} dx}{21b^3} \\ &= \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^4d} + \frac{2(b \sec(c+dx))^{7/2} \sin(c+dx)}{7b^6d} + \frac{(5\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)})}{21b^3} \\ &= \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^3d} + \frac{10(b \sec(c+dx))^{3/2} \sin(c+dx)}{21b^4d} + \end{aligned}$$

Mathematica [A] time = 0.14, size = 64, normalized size = 0.64

$$\frac{(b \sec(c+dx))^{5/2} \left(5 \sin(2(c+dx)) + 6 \tan(c+dx) + 10 \cos^{\frac{5}{2}}(c+dx) F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{21b^5d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^7/(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] ((b*Sec[c + d*x])^(5/2)*(10*Cos[c + d*x]^(5/2)*EllipticF[(c + d*x)/2, 2] +
5*Sin[2*(c + d*x)] + 6*Tan[c + d*x]))/(21*b^5*d)
```

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^4}{b^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^4/b^3, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^7}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^7/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.90, size = 152, normalized size = 1.52

$$\frac{2(1+\cos(dx+c))^2(-1+\cos(dx+c))\left(5i(\cos^3(dx+c))\sin(dx+c)\sqrt{\frac{1}{1+\cos(dx+c)}}\sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}}\operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\cos(dx+c)}\right)\right)}{21d\sin(dx+c)^3\cos(dx+c)^6\left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-5*cos(d*x+c)^3+5*cos(d*x+c)^2-3*cos(d*x+c)+3)/sin(d*x+c)^3/cos(d*x+c)^6/(b/cos(d*x+c))^(5/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^7}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^7/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^7/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^7\left(\frac{b}{\cos(c+dx)}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2)),x)`

[Out] `int(1/(cos(c + d*x)^7*(b/cos(c + d*x))^(5/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^7(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**7/(b*sec(d*x+c))**(5/2),x)`

[Out] `Integral(sec(c + d*x)**7/(b*sec(c + d*x))**(5/2), x)`

$$3.122 \quad \int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=100

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{5/2}}{5b^5d} + \frac{6 \sin(c+dx)\sqrt{b \sec(c+dx)}}{5b^3d} - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5b^2d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] $2/5*(b*\sec(d*x+c))^{(5/2)}*\sin(d*x+c)/b^5/d-6/5*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+6/5*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{5/2}}{5b^5d} + \frac{6 \sin(c+dx)\sqrt{b \sec(c+dx)}}{5b^3d} - \frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5b^2d\sqrt{\cos(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^6/(b*Sec[c + d*x])^(5/2), x]

[Out] $(-6*\text{EllipticE}[(c+d*x)/2, 2])/(5*b^2*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (6*\text{Sqrt}[b*\text{Sec}[c+d*x]]*\text{Sin}[c+d*x])/(5*b^3*d) + (2*(b*\text{Sec}[c+d*x])^{(5/2)}*\text{Sin}[c+d*x])/(5*b^5*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c-Pi/2+d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := -Simp[(b*Csc[c+d*x]*(b*Csc[c+d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c+d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\sec^6(c + dx)}{(b \sec(c + dx))^{5/2}} dx &= \frac{\int (b \sec(c + dx))^{7/2} dx}{b^6} \\
 &= \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^5 d} + \frac{3 \int (b \sec(c + dx))^{3/2} dx}{5b^4} \\
 &= \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5b^3 d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^5 d} - \frac{3 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{5b^2} \\
 &= \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5b^3 d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^5 d} - \frac{3 \int \sqrt{\cos(c + dx)}}{5b^2 \sqrt{\cos(c + dx)} \sqrt{b}} \\
 &= -\frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{6\sqrt{b \sec(c + dx)} \sin(c + dx)}{5b^3 d} + \frac{2(b \sec(c + dx))^{5/2} \sin(c + dx)}{5b^5 d}
 \end{aligned}$$

Mathematica [A] time = 0.12, size = 64, normalized size = 0.64

$$\frac{2 \tan(c + dx) (\sec^2(c + dx) + 3) - \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{\sqrt{\cos(c + dx)}}}{5b^2 d \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^6/(b*Sec[c + d*x])^(5/2), x]

[Out] ((-6*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*(3 + Sec[c + d*x]^2)*Tan[c + d*x])/(5*b^2*d*Sqrt[b*Sec[c + d*x]])

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \sec(dx + c)^3}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^3/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^6}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 1.03, size = 351, normalized size = 3.51

$$\frac{2(1 + \cos(dx + c))^2 (-1 + \cos(dx + c))^2 \left(3i (\cos^3(dx + c)) \sin(dx + c) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \operatorname{EllipticF}\left(\frac{i(\cos(dx + c) - 1)}{\sqrt{1 + \cos(dx + c)}}\right), I\right)}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x)

[Out] -2/5/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^3*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)+3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)-3*I*cos(d*x+c)^2*sin(d*x+c)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)+3*cos(d*x+c)^3-2*cos(d*x+c)^2-1)*(b/cos(d*x+c))^(5/2)/b^5/sin(d*x+c)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^6}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^6/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^6/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^6 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2)), x)`

[Out] `int(1/(cos(c + d*x)^6*(b/cos(c + d*x))^(5/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^6(c+dx)}{(b \sec(c+dx))^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**6/(b*sec(d*x+c))**(5/2), x)`

[Out] `Integral(sec(c + d*x)**6/(b*sec(c + d*x))**(5/2), x)`

$$3.123 \quad \int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{3/2}}{3b^4d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^3d}$$

[Out] $2/3*(b*\sec(d*x+c))^{(3/2)}*\sin(d*x+c)/b^4/d+2/3*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2641}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^{3/2}}{3b^4d} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^3d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^5/(b*Sec[c + d*x])^(5/2),x]

[Out] $(2*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(3*b^3*d) + (2*(b*\text{Sec}[c + d*x])^{(3/2)}*\text{Sin}[c + d*x])/(3*b^4*d)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec^5(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\int (b \sec(c+dx))^{5/2} dx}{b^5} \\ &= \frac{2(b \sec(c+dx))^{3/2} \sin(c+dx)}{3b^4d} + \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^3} \\ &= \frac{2(b \sec(c+dx))^{3/2} \sin(c+dx)}{3b^4d} + \frac{(\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^3} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^3d} + \frac{2(b \sec(c+dx))^{3/2} \sin(c+dx)}{3b^4d} \end{aligned}$$

Mathematica [A] time = 0.07, size = 51, normalized size = 0.71

$$\frac{2\sqrt{b \sec(c+dx)} \left(\tan(c+dx) + \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{3b^3d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^5/(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] (2*sqrt[b*Sec[c + d*x]]*(sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Tan
[c + d*x]))/(3*b^3*d)
```

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)} \sec(dx+c)^2}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)^2/b^3, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^5}{(b \sec(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.88, size = 130, normalized size = 1.81

$$\frac{2(-1 + \cos(dx + c)) \left(i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - \cos(dx + c) \right)}{3d \sin(dx + c)^3 \cos(dx + c)^4 \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2),x)

[Out] -2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)*(1+cos(d*x+c))^2/sin(d*x+c)^3/cos(d*x+c)^4/(b/cos(d*x+c))^(5/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^5}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^5/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^5/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c + dx)^5 \left(\frac{b}{\cos(c+dx)}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2)),x)

[Out] int(1/(cos(c + d*x)^5*(b/cos(c + d*x))^(5/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**5/(b*sec(d*x+c))**(5/2), x)

[Out] Integral(sec(c + d*x)**5/(b*sec(c + d*x))**(5/2), x)

$$3.124 \quad \int \frac{\sec^4(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=68

$$\frac{2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{b^3 d} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

[Out] $-2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3768, 3771, 2639}

$$\frac{2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{b^3 d} - \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^4/(b*Sec[c + d*x])^(5/2), x]

[Out] $(-2*\text{EllipticE}[(c+d*x)/2, 2])/(b^2*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (2*\text{Sqrt}[b*\text{Sec}[c+d*x]]*\text{Sin}[c+d*x])/(b^3*d)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n-1))/(d*(n-1)), x] + Dist[(b^2*(n-2))/(n-1), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{5/2}} dx &= \frac{\int (b \sec(c + dx))^{3/2} dx}{b^4} \\ &= \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{b^3 d} - \frac{\int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{b^2} \\ &= \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{b^3 d} - \frac{\int \sqrt{\cos(c + dx)} dx}{b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\ &= -\frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2\sqrt{b \sec(c + dx)} \sin(c + dx)}{b^3 d} \end{aligned}$$

Mathematica [A] time = 0.05, size = 51, normalized size = 0.75

$$\frac{2 \tan(c + dx) - \frac{2E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{\sqrt{\cos(c + dx)}}}{b^2 d \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]^4/(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] ((-2*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 2*Tan[c + d*x])/(b^2*d
*Sqrt[b*Sec[c + d*x]])
```

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \sec(dx + c)}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))*sec(d*x + c)/b^3, x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 1.06, size = 324, normalized size = 4.76

$$2(1 + \cos(dx+c))^2(-1 + \cos(dx+c))^2 \left(i \cos(dx+c) \sin(dx+c) \operatorname{EllipticE}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} \sqrt{\frac{c}{1+\cos(dx+c)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x)

[Out] 2/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))^2*(I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*cos(d*x+c)*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)+1)/cos(d*x+c)^3/(b/cos(d*x+c))^(5/2)/sin(d*x+c)^5

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^4}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^4/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^4/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx)^4 \left(\frac{b}{\cos(c+dx)}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2)),x)`

[Out] `int(1/(cos(c + d*x)^4*(b/cos(c + d*x))^(5/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**4/(b*sec(d*x+c))**(5/2),x)`

[Out] `Integral(sec(c + d*x)**4/(b*sec(c + d*x))**(5/2), x)`

$$3.125 \quad \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=41

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^3 d}$$

[Out] 2*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3771, 2641}

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^3 d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^3/(b*Sec[c + d*x])^(5/2),x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^3*d)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\int \sqrt{b \sec(c+dx)} dx}{b^3} \\ &= \frac{(\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{b^3} \\ &= \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^3 d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 41, normalized size = 1.00

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{b^3 d}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^3/(b*Sec[c + d*x])^(5/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(b^3*d)

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)}}{b^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/b^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \sec(dx+c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.76, size = 98, normalized size = 2.39

$$\frac{2i \left(\frac{1}{1+\cos(dx+c)} \right)^{\frac{7}{2}} (-1 + \cos(dx+c)) \operatorname{EllipticF} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) (1 + \cos(dx+c))^2}{d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{5}{2}} \left(\frac{\cos(dx+c)}{1+\cos(dx+c)} \right)^{\frac{5}{2}} \sin(dx+c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2), x)`

[Out] `-2*I/d*(1/(1+cos(d*x+c)))^(7/2)*(-1+cos(d*x+c))*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c), I)*(1+cos(d*x+c))^2/(b/cos(d*x+c))^(5/2)/(cos(d*x+c)/(1+cos(d*x+c)))^(5/2)/sin(d*x+c)^2`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^3}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^3/(b*sec(d*x+c))^(5/2), x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^3/(b*sec(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c+dx)^3 \left(\frac{b}{\cos(c+dx)} \right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2)), x)`

[Out] `int(1/(cos(c + d*x)^3*(b/cos(c + d*x))^(5/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**3/(b*sec(d*x+c))**(5/2), x)`

[Out] `Integral(sec(c + d*x)**3/(b*sec(c + d*x))**(5/2), x)`

$$3.126 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=41

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

[Out] $2*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c), 2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3771, 2639}

$$\frac{2E\left(\frac{1}{2}(c+dx)\middle|2\right)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(5/2), x]

[Out] (2*EllipticE[(c + d*x)/2, 2])/(b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3771

Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{b^2} \\ &= \frac{\int \sqrt{\cos(c+dx)} dx}{b^2 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \\ &= \frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{b^2 d \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 38, normalized size = 0.93

$$\frac{2E\left(\frac{1}{2}(c+dx) \middle| 2\right)}{d \cos^{\frac{5}{2}}(c+dx) (b \sec(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(5/2), x]

[Out] (2*EllipticE[(c + d*x)/2, 2])/(d*Cos[c + d*x]^(5/2)*(b*Sec[c + d*x])^(5/2))

fricas [F] time = 0.50, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)}}{b^3 \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/(b^3*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.75, size = 311, normalized size = 7.59

$$2i \sin(dx + c) \cos(dx + c) \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} - 2i \cos(dx + c) \sin(dx + c) E$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2), x)`

[Out] $2/d*(I*\sin(d*x+c)*\cos(d*x+c)*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}-I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}+I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I)*(1/(1+\cos(d*x+c)))^{1/2}*(\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2}-\cos(d*x+c)^2+\cos(d*x+c))/\cos(d*x+c)^3/(b/\cos(d*x+c))^{5/2}/\sin(d*x+c)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, algorithm="maxima")`

[Out] `integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2)), x)`

[Out] `int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(5/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(5/2), x)
```

```
[Out] Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(5/2), x)
```


$$3.127 \quad \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^3d} + \frac{2 \sin(c+dx)}{3b^2d\sqrt{b \sec(c+dx)}}$$

[Out] 2/3*sin(d*x+c)/b^2/d/(b*sec(d*x+c))^(1/2)+2/3*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticF(sin(1/2*d*x+1/2*c),2^(1/2))*cos(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/b^3/d

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3769, 3771, 2641}

$$\frac{2 \sin(c+dx)}{3b^2d\sqrt{b \sec(c+dx)}} + \frac{2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^3d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Sec[c + d*x])^(5/2), x]

[Out] (2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2]*Sqrt[b*Sec[c + d*x]])/(3*b^3*d) + (2*Sin[c + d*x])/(3*b^2*d*Sqrt[b*Sec[c + d*x]])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d^n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{b} \\ &= \frac{2 \sin(c+dx)}{3b^2 d \sqrt{b \sec(c+dx)}} + \frac{\int \sqrt{b \sec(c+dx)} dx}{3b^3} \\ &= \frac{2 \sin(c+dx)}{3b^2 d \sqrt{b \sec(c+dx)}} + \frac{(\sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{3b^3} \\ &= \frac{2 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{3b^3 d} + \frac{2 \sin(c+dx)}{3b^2 d \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 62, normalized size = 0.86

$$\frac{\sec^2(c+dx) \left(\sin(2(c+dx)) + 2\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{3bd(b \sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] (Sec[c + d*x]^2*(2*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + Sin[2*(c + d*x)]))/(3*b*d*(b*Sec[c + d*x])^(3/2))
```

fricas [F] time = 0.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx+c)}}{b^3 \sec(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(d*x + c))/(b^3*sec(d*x + c)^2), x)
```

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.74, size = 131, normalized size = 1.82

$$\frac{2(-1 + \cos(dx+c)) \left(i \sin(dx+c) \operatorname{EllipticF}\left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i\right) \sqrt{\frac{1}{1+\cos(dx+c)}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - (\cos^2(dx+c)) + c \right)}{3d \sin(dx+c)^3 \cos(dx+c)^3 \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x)

[Out] -2/3/d*(-1+cos(d*x+c))*(I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-cos(d*x+c)^2+cos(d*x+c))*(1+cos(d*x+c))^2/sin(d*x+c)^3/cos(d*x+c)^3/(b/cos(d*x+c))^(5/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\cos(c+dx) \left(\frac{b}{\cos(c+dx)}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(5/2)),x)`

[Out] `int(1/(cos(c + d*x)*(b/cos(c + d*x))^(5/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)/(b*sec(d*x+c))**(5/2),x)`

[Out] `Integral(sec(c + d*x)/(b*sec(c + d*x))**(5/2), x)`

$$3.128 \quad \int \frac{1}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5b^2d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{2\sin(c+dx)}{5bd(b\sec(c+dx))^{3/2}}$$

[Out] 2/5*sin(d*x+c)/b/d/(b*sec(d*x+c))^(3/2)+6/5*(cos(1/2*d*x+1/2*c)^2)^(1/2)/cos(1/2*d*x+1/2*c)*EllipticE(sin(1/2*d*x+1/2*c),2^(1/2))/b^2/d/cos(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.03, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3769, 3771, 2639}

$$\frac{6E\left(\frac{1}{2}(c+dx)\middle|2\right)}{5b^2d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{2\sin(c+dx)}{5bd(b\sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(-5/2), x]

[Out] (6*EllipticE[(c + d*x)/2, 2])/(5*b^2*d*Sqrt[Cos[c + d*x]]*Sqrt[b*Sec[c + d*x]]) + (2*Sin[c + d*x])/(5*b*d*(b*Sec[c + d*x])^(3/2))

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d^n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(b \sec(c + dx))^{5/2}} dx &= \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} + \frac{3 \int \frac{1}{\sqrt{b \sec(c+dx)}} dx}{5b^2} \\
&= \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}} + \frac{3 \int \sqrt{\cos(c + dx)} dx}{5b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
&= \frac{6E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{5b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2 \sin(c + dx)}{5bd(b \sec(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 60, normalized size = 0.83

$$\frac{\sqrt{b \sec(c + dx)} \left(\sin(c + dx) + \sin(3(c + dx)) + 12 \sqrt{\cos(c + dx)} E\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{10b^3 d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(-5/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*(12*Sqrt[Cos[c + d*x]]*EllipticE[(c + d*x)/2, 2] + Sin[c + d*x] + Sin[3*(c + d*x)]))/(10*b^3*d)

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx + c)}}{b^3 \sec(dx + c)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/(b^3*sec(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(-5/2), x)

maple [C] time = 0.83, size = 321, normalized size = 4.46

$$2 \left(3i \cos(dx + c) \sin(dx + c) \operatorname{EllipticE} \left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i \right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} - 3i \sin(dx + c) \cos(dx + c) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(d*x+c))^(5/2), x)

[Out]
$$-2/5/d*(3*I*\cos(d*x+c)*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I) * (1/(1+\cos(d*x+c)))^{1/2} * (\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2} - 3*I*\sin(d*x+c)*\cos(d*x+c) * (\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2} * \operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I) * (1/(1+\cos(d*x+c)))^{1/2} + 3*I*\sin(d*x+c)*\operatorname{EllipticE}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I) * (1/(1+\cos(d*x+c)))^{1/2} * (\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2} - 3*I*\sin(d*x+c)*\operatorname{EllipticF}(I*(-1+\cos(d*x+c))/\sin(d*x+c), I) * (1/(1+\cos(d*x+c)))^{1/2} * (\cos(d*x+c)/(1+\cos(d*x+c)))^{1/2} + \cos(d*x+c)^4 + 2*\cos(d*x+c)^2 - 3*\cos(d*x+c))/\cos(d*x+c)^3/\sin(d*x+c)/(b/\cos(d*x+c))^{5/2}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(5/2), x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(-5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b/cos(c + d*x))^(5/2), x)

[Out] int(1/(b/cos(c + d*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))**(5/2),x)

[Out] Integral((b*sec(c + d*x))**(-5/2), x)

$$3.129 \quad \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=97

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^3d} + \frac{10 \sin(c+dx)}{21b^2d\sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}}$$

[Out] $2/7*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/2)}+10/21*\sin(d*x+c)/b^2/d/(b*\sec(d*x+c))^{(1/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^3/d$

Rubi [A] time = 0.06, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3769, 3771, 2641}

$$\frac{10 \sin(c+dx)}{21b^2d\sqrt{b \sec(c+dx)}} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^3d} + \frac{2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]/(b*Sec[c + d*x])^(5/2),x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*b^3*d) + (2*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(5/2)}) + (10*\text{Sin}[c + d*x])/(21*b^2*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] :=> Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= b \int \frac{1}{(b \sec(c+dx))^{7/2}} dx \\
&= \frac{2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{5 \int \frac{1}{(b \sec(c+dx))^{3/2}} dx}{7b} \\
&= \frac{2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21b^2 d \sqrt{b \sec(c+dx)}} + \frac{5 \int \sqrt{b \sec(c+dx)} dx}{21b^3} \\
&= \frac{2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21b^2 d \sqrt{b \sec(c+dx)}} + \frac{(5 \sqrt{\cos(c+dx)} \sqrt{b \sec(c+dx)}) \int \frac{1}{\sqrt{\cos(c+dx)}} dx}{21b^3} \\
&= \frac{10 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^3 d} + \frac{2 \sin(c+dx)}{7d(b \sec(c+dx))^{5/2}} + \frac{10 \sin(c+dx)}{21b^2 d \sqrt{b \sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 66, normalized size = 0.68

$$\frac{\sqrt{b \sec(c+dx)} \left(26 \sin(2(c+dx)) + 3 \sin(4(c+dx)) + 40 \sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \right)}{84b^3 d}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] (Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26
*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b^3*d)
```

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{b \sec(dx+c)} \cos(dx+c)}{b^3 \sec(dx+c)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)/(b^3*sec(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.84, size = 153, normalized size = 1.58

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(5i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \right)}{21d \sin(dx + c)^3 \cos(dx + c)^3 \left(\frac{b}{\cos(dx + c)} \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*cos(d*x+c)^4+3*cos(d*x+c)^3-5*cos(d*x+c)^2+5*cos(d*x+c))/sin(d*x+c)^3/cos(d*x+c)^3/(b/cos(d*x+c))^(5/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)}{\left(\frac{b}{\cos(c + dx)}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)/(b/cos(c + d*x))^(5/2), x)`

[Out] `int(cos(c + d*x)/(b/cos(c + d*x))^(5/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)/(b*sec(d*x+c))**(5/2), x)`

[Out] `Integral(cos(c + d*x)/(b*sec(c + d*x))**(5/2), x)`

$$3.130 \quad \int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=98

$$\frac{14E\left(\frac{1}{2}(c+dx)\middle|2\right)}{15b^2d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{14\sin(c+dx)}{45bd(b\sec(c+dx))^{3/2}} + \frac{2b\sin(c+dx)}{9d(b\sec(c+dx))^{7/2}}$$

[Out] $2/9*b*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/2)}+14/45*\sin(d*x+c)/b/d/(b*\sec(d*x+c))^{(3/2)}+14/15*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticE}(\sin(1/2*d*x+1/2*c),2^{(1/2)})/b^2/d/\cos(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3769, 3771, 2639}

$$\frac{14E\left(\frac{1}{2}(c+dx)\middle|2\right)}{15b^2d\sqrt{\cos(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{14\sin(c+dx)}{45bd(b\sec(c+dx))^{3/2}} + \frac{2b\sin(c+dx)}{9d(b\sec(c+dx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(5/2), x]

[Out] $(14*\text{EllipticE}[(c+d*x)/2, 2])/((15*b^2*d*\text{Sqrt}[\text{Cos}[c+d*x]]*\text{Sqrt}[b*\text{Sec}[c+d*x]]) + (2*b*\text{Sin}[c+d*x])/(9*d*(b*\text{Sec}[c+d*x])^{(7/2)}) + (14*\text{Sin}[c+d*x])/(45*b*d*(b*\text{Sec}[c+d*x])^{(3/2)})$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2639

Int[Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_)+(d_)*(x_)])*(b_)^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n+1))/(b*d*n), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

```
Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] :=> Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx &= b^2 \int \frac{1}{(b \sec(c + dx))^{9/2}} dx \\
&= \frac{2b \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{7}{9} \int \frac{1}{(b \sec(c + dx))^{5/2}} dx \\
&= \frac{2b \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14 \sin(c + dx)}{45bd(b \sec(c + dx))^{3/2}} + \frac{7 \int \frac{1}{\sqrt{b \sec(c + dx)}} dx}{15b^2} \\
&= \frac{2b \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14 \sin(c + dx)}{45bd(b \sec(c + dx))^{3/2}} + \frac{7 \int \sqrt{\cos(c + dx)} dx}{15b^2 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} \\
&= \frac{14E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{15b^2 d \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}} + \frac{2b \sin(c + dx)}{9d(b \sec(c + dx))^{7/2}} + \frac{14 \sin(c + dx)}{45bd(b \sec(c + dx))^{3/2}}
\end{aligned}$$

Mathematica [A] time = 0.13, size = 73, normalized size = 0.74

$$\frac{4(33 \sin(c + dx) + 5 \sin(3(c + dx))) \cos(c + dx) + \frac{336E\left(\frac{1}{2}(c + dx) \middle| 2\right)}{\sqrt{\cos(c + dx)}}}{360b^2 d \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(5/2), x]
```

```
[Out] ((336*EllipticE[(c + d*x)/2, 2])/Sqrt[Cos[c + d*x]] + 4*Cos[c + d*x]*(33*Sin[c + d*x] + 5*Sin[3*(c + d*x)]))/(360*b^2*d*Sqrt[b*Sec[c + d*x]])
```

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)} \cos(dx + c)^2}{b^3 \sec(dx + c)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")
```

[Out] integral(sqrt(b*sec(d*x + c))*cos(d*x + c)^2/(b^3*sec(d*x + c)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(5/2), x)

maple [C] time = 0.94, size = 333, normalized size = 3.40

$$\frac{2 \left(5 \left(\cos^6(dx+c) \right) + 21i \cos(dx+c) \sin(dx+c) \operatorname{EllipticE} \left(\frac{i(-1+\cos(dx+c))}{\sin(dx+c)}, i \right) \sqrt{\frac{1}{1+\cos(dx+c)}} \sqrt{\frac{\cos(dx+c)}{1+\cos(dx+c)}} - 21i \right)}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*sec(d*x+c))^(5/2),x)

[Out] -2/45/d*(5*cos(d*x+c)^6+21*I*cos(d*x+c)*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-21*I*sin(d*x+c)*cos(d*x+c)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)+21*I*sin(d*x+c)*EllipticE(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-21*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)+2*cos(d*x+c)^4+14*cos(d*x+c)^2-21*cos(d*x+c))/cos(d*x+c)^3/(b/cos(d*x+c))^(5/2)/sin(d*x+c)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\cos(c + dx)^2}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2/(b/cos(c + d*x))^(5/2), x)

[Out] int(cos(c + d*x)^2/(b/cos(c + d*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(5/2), x)

[Out] Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(5/2), x)

$$3.131 \quad \int \frac{1}{(b \sec(c+dx))^{7/2}} dx$$

Optimal. Leaf size=100

$$\frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^4d} + \frac{10 \sin(c+dx)}{21b^3d\sqrt{b \sec(c+dx)}} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}}$$

[Out] $2/7*\sin(d*x+c)/b/d/(b*\sec(d*x+c))^{(5/2)}+10/21*\sin(d*x+c)/b^3/d/(b*\sec(d*x+c))^{(1/2)}+10/21*(\cos(1/2*d*x+1/2*c)^2)^{(1/2)}/\cos(1/2*d*x+1/2*c)*\text{EllipticF}(\sin(1/2*d*x+1/2*c),2^{(1/2)})*\cos(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/b^4/d$

Rubi [A] time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3769, 3771, 2641}

$$\frac{10 \sin(c+dx)}{21b^3d\sqrt{b \sec(c+dx)}} + \frac{10\sqrt{\cos(c+dx)} F\left(\frac{1}{2}(c+dx) \middle| 2\right) \sqrt{b \sec(c+dx)}}{21b^4d} + \frac{2 \sin(c+dx)}{7bd(b \sec(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(-7/2), x]

[Out] $(10*\text{Sqrt}[\text{Cos}[c + d*x]]*\text{EllipticF}[(c + d*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(21*b^4*d) + (2*\text{Sin}[c + d*x])/(7*b*d*(b*\text{Sec}[c + d*x])^{(5/2)}) + (10*\text{Sin}[c + d*x])/(21*b^3*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 2641

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rule 3769

Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Csc[c + d*x])^(n + 1))/(b*d*n), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3771

Int[(csc[(c_) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(b \sec(c + dx))^{7/2}} dx &= \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} + \frac{5 \int \frac{1}{(b \sec(c + dx))^{3/2}} dx}{7b^2} \\
&= \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21b^3 d \sqrt{b \sec(c + dx)}} + \frac{5 \int \sqrt{b \sec(c + dx)} dx}{21b^4} \\
&= \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21b^3 d \sqrt{b \sec(c + dx)}} + \frac{(5 \sqrt{\cos(c + dx)} \sqrt{b \sec(c + dx)}) \int \frac{1}{\sqrt{\cos(c + dx)}} dx}{21b^4} \\
&= \frac{10 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \sqrt{b \sec(c + dx)}}{21b^4 d} + \frac{2 \sin(c + dx)}{7bd(b \sec(c + dx))^{5/2}} + \frac{10 \sin(c + dx)}{21b^3 d \sqrt{b \sec(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 66, normalized size = 0.66

$$\frac{\sqrt{b \sec(c + dx)} \left(26 \sin(2(c + dx)) + 3 \sin(4(c + dx)) + 40 \sqrt{\cos(c + dx)} F\left(\frac{1}{2}(c + dx) \middle| 2\right) \right)}{84b^4 d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(-7/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*(40*Sqrt[Cos[c + d*x]]*EllipticF[(c + d*x)/2, 2] + 26*Sin[2*(c + d*x)] + 3*Sin[4*(c + d*x)]))/(84*b^4*d)

fricas [F] time = 0.88, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{b \sec(dx + c)}}{b^4 \sec(dx + c)^4}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(7/2), x, algorithm="fricas")

[Out] integral(sqrt(b*sec(d*x + c))/(b^4*sec(d*x + c)^4), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(7/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(-7/2), x)

maple [C] time = 0.90, size = 153, normalized size = 1.53

$$\frac{2(1 + \cos(dx + c))^2(-1 + \cos(dx + c)) \left(5i \sin(dx + c) \operatorname{EllipticF}\left(\frac{i(-1 + \cos(dx + c))}{\sin(dx + c)}, i\right) \sqrt{\frac{1}{1 + \cos(dx + c)}} \sqrt{\frac{\cos(dx + c)}{1 + \cos(dx + c)}} \right)}{21d \left(\frac{b}{\cos(dx + c)}\right)^{\frac{7}{2}} \cos(dx + c)^4 \sin(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(d*x+c))^(7/2),x)

[Out] -2/21/d*(1+cos(d*x+c))^2*(-1+cos(d*x+c))*(5*I*sin(d*x+c)*EllipticF(I*(-1+cos(d*x+c))/sin(d*x+c),I)*(1/(1+cos(d*x+c)))^(1/2)*(cos(d*x+c)/(1+cos(d*x+c)))^(1/2)-3*cos(d*x+c)^4+3*cos(d*x+c)^3-5*cos(d*x+c)^2+5*cos(d*x+c))/(b/cos(d*x+c))^(7/2)/cos(d*x+c)^4/sin(d*x+c)^3

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(7/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(-7/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{b}{\cos(c + dx)}\right)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b/cos(c + d*x))^(7/2),x)

[Out] int(1/(b/cos(c + d*x))^(7/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(c + dx))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(d*x+c))**(7/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**(-7/2), x)
```

3.132 $\int \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=107

$$\frac{\sin(c + dx) \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{4d} + \frac{3 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{8d} + \frac{3 \sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{8d \sqrt{\sec(c + dx)}}$$

[Out] 3/8*sec(d*x+c)^(3/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+1/4*sec(d*x+c)^(7/2)*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d+3/8*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{\sin(c + dx) \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{4d} + \frac{3 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{8d} + \frac{3 \sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{8d \sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(9/2)*Sqrt[b*Sec[c + d*x]], x]

[Out] (3*ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]]/(8*d*Sqrt[Sec[c + d*x]]) + (3*Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(8*d) + (Sec[c + d*x]^(7/2)*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(4*d)

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{9}{2}}(c+dx)\sqrt{b\sec(c+dx)} dx &= \frac{\sqrt{b\sec(c+dx)} \int \sec^5(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{\sec^{\frac{7}{2}}(c+dx)\sqrt{b\sec(c+dx)} \sin(c+dx)}{4d} + \frac{(3\sqrt{b\sec(c+dx)}) \int \sec^3(c+dx) dx}{4\sqrt{\sec(c+dx)}} \\
&= \frac{3\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)} \sin(c+dx)}{8d} + \frac{\sec^{\frac{7}{2}}(c+dx)\sqrt{b\sec(c+dx)} \sin(c+dx)}{4d} \\
&= \frac{3 \tanh^{-1}(\sin(c+dx))\sqrt{b\sec(c+dx)}}{8d\sqrt{\sec(c+dx)}} + \frac{3\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)} \sin(c+dx)}{8d}
\end{aligned}$$

Mathematica [A] time = 0.15, size = 64, normalized size = 0.60

$$\frac{\sqrt{b\sec(c+dx)} \left(3 \tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx) (2 \sec^2(c+dx) + 3) \right)}{8d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(9/2)*Sqrt[b*Sec[c + d*x]], x]

[Out] (Sqrt[b*Sec[c + d*x]]*(3*ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*(3 + 2*Sec[c + d*x]^2)*Tan[c + d*x]))/(8*d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.67, size = 229, normalized size = 2.14

$$\left[\frac{3\sqrt{b} \cos(dx+c)^3 \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2(3 \cos(dx+c)^2 + 2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{16 d \cos(dx+c)^3}, - \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(9/2)*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/16*(3*sqrt(b)*cos(d*x + c)^3*log(-(b*cos(d*x + c)^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*(3*cos(d*x + c)^2 + 2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/

$d*\cos(dx + c)^3$, $-1/8*(3*\sqrt{-b}*\arctan(\sqrt{-b}*\sqrt{b/\cos(dx + c)})*\sqrt{\cos(dx + c)}*\sin(dx + c)/b)*\cos(dx + c)^3 - (3*\cos(dx + c)^2 + 2)*\sqrt{b/\cos(dx + c)}*\sin(dx + c)/\sqrt{\cos(dx + c)})/(d*\cos(dx + c)^3]$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sec(dx + c)^{\frac{9}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(dx+c)^(9/2)*(b*sec(dx+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(dx + c))*sec(dx + c)^(9/2), x)

maple [A] time = 1.20, size = 131, normalized size = 1.22

$$\frac{\left(3 \left(\cos^4(dx + c)\right) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - 3 \left(\cos^4(dx + c)\right) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) + 3 \left(\cos^2(dx + c)\right) \sin(dx+c)}{8d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(dx+c)^(9/2)*(b*sec(dx+c))^(1/2),x)

[Out] $1/8/d*(3*\cos(dx+c)^4*\ln(-(-\sin(dx+c)-1+\cos(dx+c))/\sin(dx+c))-3*\cos(dx+c)^4*\ln(-(-1+\cos(dx+c)+\sin(dx+c))/\sin(dx+c))+3*\cos(dx+c)^2*\sin(dx+c)+2*\sin(dx+c))*\cos(dx+c)*(1/\cos(dx+c))^(9/2)*(b/\cos(dx+c))^(1/2)$

maxima [B] time = 1.06, size = 1656, normalized size = 15.48

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(dx+c)^(9/2)*(b*sec(dx+c))^(1/2),x, algorithm="maxima")

[Out] $-1/16*(12*(\sin(8*d*x + 8*c) + 4*\sin(6*d*x + 6*c) + 6*\sin(4*d*x + 4*c) + 4*\sin(2*d*x + 2*c))*\cos(7/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 44*(\sin(8*d*x + 8*c) + 4*\sin(6*d*x + 6*c) + 6*\sin(4*d*x + 4*c) + 4*\sin(2*d*x + 2*c))*\cos(5/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 44*(\sin(8*d*x + 8*c) + 4*\sin(6*d*x + 6*c) + 6*\sin(4*d*x + 4*c) + 4*\sin(2*d*x + 2*c))*\cos(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 12*(\sin(8*d*x + 8*c) + 4*\sin(6*d*x + 6*c) + 6*\sin(4*d*x + 4*c) + 4*\sin(2*d*x + 2*c))*\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 3*(2*(4*\cos(6*d*x + 6*c) + 6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\cos(8*d*x + 8*c) + \cos(8*d*x + 8*c)^2 + 8*(6*\cos(4*d*x + 4*c) + 4*\cos(2*d*x + 2*c) + 1)*\cos(6*d*x + 6*c) + 16*c$

```

os(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4
*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x
+ 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3
*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6
*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*s
in(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + 3
*(2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*cos(
8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2
*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2*d*x + 2*c)
+ 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x + 2*c)^2 + 4*(
2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin(8*d*x + 8
*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*sin
(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 + 48*sin(4*d*
x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x + 2*c) + 1)
*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan
2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c)))) + 1) - 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) +
6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(7/2*arctan2(sin(2*d*x + 2*
c), cos(2*d*x + 2*c))) - 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(
4*d*x + 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(5/2*arctan2(sin(2*d*x + 2*c), co
s(2*d*x + 2*c))) + 44*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x
+ 4*c) + 4*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*
x + 2*c))) + 12*(cos(8*d*x + 8*c) + 4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c)
+ 4*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*
c))))*sqrt(b)/((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x +
2*c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4
*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos
(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x
+ 2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))
*sin(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d
*x + 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2
+ 48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d
*x + 2*c) + 1)*d)

```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{\frac{b}{\cos(c + dx)}} \left(\frac{1}{\cos(c + dx)} \right)^{9/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(9/2),x)


```
[Out] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(9/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**(9/2)*(b*sec(d*x+c))**(1/2), x)
```

```
[Out] Timed out
```

3.133 $\int \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=70

$$\frac{\sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{\sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

[Out] $1/3*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)^3*(b*\sec(d*x+c))^{(1/2)}/d+\sin(d*x+c)*\sec(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{\sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{\sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^(7/2)*Sqrt[b*Sec[c + d*x]],x]`

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d + (\text{Sec}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d)$

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 3767

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{7}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx &= \frac{\sqrt{b \sec(c+dx)} \int \sec^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= -\frac{\sqrt{b \sec(c+dx)} \operatorname{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d \sqrt{\sec(c+dx)}} \\
&= \frac{\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d} + \frac{\sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{3d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 45, normalized size = 0.64

$$\frac{\left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx)\right) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(7/2)*Sqrt[b*Sec[c + d*x]],x]

[Out] (Sqrt[b*Sec[c + d*x]]*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.63, size = 43, normalized size = 0.61

$$\frac{(2 \cos(dx+c)^2 + 1) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3d \cos(dx+c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*(2*cos(d*x + c)^2 + 1)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(5/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx+c)} \sec(dx+c)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(7/2), x)

maple [A] time = 1.07, size = 52, normalized size = 0.74

$$\frac{\left(2\left(\cos^2(dx+c)\right)+1\right)\left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}}\sqrt{\frac{b}{\cos(dx+c)}}\cos(dx+c)\sin(dx+c)}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2), x)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*(1/cos(d*x+c))^(7/2)*(b/cos(d*x+c))^(1/2)*cos(d*x+c)*sin(d*x+c)

maxima [B] time = 0.83, size = 294, normalized size = 4.20

$$\frac{4((3\cos(2dx+2c) + 1)\sin(6dx+6c) + 3(3\cos(2dx+2c) + 1)\sin(4dx+4c) - 3\cos(6dx+6c)\sin(2dx+2c) - 9\cos(4dx+4c)\sin(2dx+2c))\sqrt{b}/((2(3\cos(4dx+4c) + 3\cos(2dx+2c) + 1)\cos(6dx+6c) + \cos(6dx+6c)^2 + 6(3\cos(2dx+2c) + 1)\cos(4dx+4c) + 9\cos(4dx+4c)^2 + 9\cos(2dx+2c)^2 + 6(\sin(4dx+4c) + \sin(2dx+2c))\sin(6dx+6c) + \sin(6dx+6c)^2 + 9\sin(4dx+4c)^2 + 18\sin(4dx+4c)\sin(2dx+2c) + 9\sin(2dx+2c)^2 + 6\cos(2dx+2c) + 1)d)}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(1/2), x, algorithm="maxima")

[Out] 4/3*((3*cos(2*d*x + 2*c) + 1)*sin(6*d*x + 6*c) + 3*(3*cos(2*d*x + 2*c) + 1)*sin(4*d*x + 4*c) - 3*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) - 9*cos(4*d*x + 4*c)*sin(2*d*x + 2*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 2.55, size = 126, normalized size = 1.80

$$\frac{2\cos(c+dx)\sqrt{\frac{b}{\cos(c+dx)}}\sqrt{\frac{1}{\cos(c+dx)}}(4\sin(c+dx)+5\sin(3c+3dx)+\sin(5c+5dx)+\cos(c+dx)10i+4\sin(c+dx)+\cos(3c+3dx)*5i+\cos(5c+5dx)*1i+5\sin(3c+3dx)+\sin(5c+5dx))}{3d(10\cos(c+dx)+5\cos(3c+3dx)+\cos(5c+5dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(7/2), x)

[Out] (2*cos(c + d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*(cos(c + d*x)*10i + 4*sin(c + d*x) + cos(3*c + 3*d*x)*5i + cos(5*c + 5*d*x)*1i + 5*sin(3

```
*c + 3*d*x) + sin(5*c + 5*d*x)))/(3*d*(10*cos(c + d*x) + 5*cos(3*c + 3*d*x)
+ cos(5*c + 5*d*x)))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**(7/2)*(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

3.134 $\int \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=72

$$\frac{\sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{2d} + \frac{\sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

[Out] $1/2 * \sec(d*x+c)^{(3/2)} * \sin(d*x+c) * (b * \sec(d*x+c))^{(1/2)} / d + 1/2 * \operatorname{arctanh}(\sin(d*x+c)) * (b * \sec(d*x+c))^{(1/2)} / d / \sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{\sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{2d} + \frac{\sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]],x]`

[Out] $(\operatorname{ArcTanh}[\sin[c + d*x]] * \operatorname{Sqrt}[b * \operatorname{Sec}[c + d*x]]) / (2 * d * \operatorname{Sqrt}[\operatorname{Sec}[c + d*x]]) + (\operatorname{Sec}[c + d*x]^{(3/2)} * \operatorname{Sqrt}[b * \operatorname{Sec}[c + d*x]] * \sin[c + d*x]) / (2 * d)$

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3770

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx &= \frac{\sqrt{b \sec(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{\sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{2d} + \frac{\sqrt{b \sec(c+dx)} \int \sec(c+dx) dx}{2\sqrt{\sec(c+dx)}} \\
&= \frac{\tanh^{-1}(\sin(c+dx)) \sqrt{b \sec(c+dx)}}{2d\sqrt{\sec(c+dx)}} + \frac{\sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{2d}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 50, normalized size = 0.69

$$\frac{\sqrt{b \sec(c+dx)} \left(\tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx) \right)}{2d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]],x]

[Out] (Sqrt[b*Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.77, size = 199, normalized size = 2.76

$$\left[\frac{\sqrt{b} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4d \cos(dx+c)}, -\sqrt{-b} \arctan\left(\frac{\sqrt{-b}}{\dots} \right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c)^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx+c)} \sec(dx+c)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(5/2), x)

maple [A] time = 1.07, size = 114, normalized size = 1.58

$$\frac{\left((\cos^2(dx+c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx+c)) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx+c) \right) \cos(dx+c)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c)*cos(d*x+c)*(1/cos(d*x+c))^(5/2)*(b/cos(d*x+c))^(1/2)

maxima [B] time = 0.97, size = 661, normalized size = 9.18

$$\frac{\left(4(\sin(4dx+4c) + 2\sin(2dx+2c)) \cos\left(\frac{3}{2} \arctan(\sin(2dx+2c), \cos(2dx+2c))\right) - 4(\sin(4dx+4c) + \right)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))

*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{\frac{b}{\cos(c + dx)}} \left(\frac{1}{\cos(c + dx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(5/2), x)

[Out] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(5/2)*(b*sec(d*x+c))**(1/2), x)

[Out] Timed out

$$3.135 \quad \int \sec^2(c + dx) \sqrt{b \sec(c + dx)} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{\sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(3/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]], x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m + 1/2)}*b^{(n - 1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m + n)}, x], x] /; \text{FreeQ}\{a, b, m\}, x \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n + 1/2, 0] \ \&\& \ \text{IntegerQ}[m + n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2 - 1)}, x], x], x, \text{Cot}[c + d*x]], x] /; \text{FreeQ}\{c, d\}, x \ \&\& \ \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} dx &= \frac{\sqrt{b \sec(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{\sec(c+dx)}} \\ &= -\frac{\sqrt{b \sec(c+dx)} \operatorname{Subst}\left(\int 1 dx, x, -\tan(c+dx)\right)}{d \sqrt{\sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sin(c+dx) \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]],x]

[Out] (Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d

fricas [A] time = 0.81, size = 30, normalized size = 0.94

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx+c)} \sec(dx+c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sec(d*x + c)^(3/2), x)

maple [A] time = 1.08, size = 39, normalized size = 1.22

$$\frac{\cos(dx+c)\sin(dx+c)\left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}}\sqrt{\frac{b}{\cos(dx+c)}}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x)`

[Out] `1/d*cos(d*x+c)*sin(d*x+c)*(1/cos(d*x+c))^(3/2)*(b/cos(d*x+c))^(1/2)`

maxima [A] time = 1.05, size = 54, normalized size = 1.69

$$\frac{2\sqrt{b}\sin(2dx+2c)}{(\cos(2dx+2c)^2+\sin(2dx+2c)^2+2\cos(2dx+2c)+1)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `2*sqrt(b)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)`

mupad [B] time = 0.29, size = 46, normalized size = 1.44

$$\frac{(\cos(dx) - \sin(dx)1i)(\sin(c) + \cos(c)1i)\sqrt{\frac{b}{\cos(c+dx)}}\sqrt{\frac{1}{\cos(c+dx)}}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c+d*x))^(1/2)*(1/cos(c+d*x))^(3/2),x)`

[Out] `((cos(d*x) - sin(d*x)*1i)*(cos(c)*1i + sin(c))*(b/cos(c+d*x))^(1/2)*(1/cos(c+d*x))^(3/2))/d`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c+dx)} \sec^{\frac{3}{2}}(c+dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**(1/2),x)`

[Out] `Integral(sqrt(b*sec(c+d*x))*sec(c+d*x)**(3/2),x)`

3.136 $\int \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} dx$

Optimal. Leaf size=33

$$\frac{\sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{d\sqrt{\sec(c + dx)}}$$

[Out] arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{d\sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]], x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} dx &= \frac{\sqrt{b \sec(c + dx)} \int \sec(c + dx) dx}{\sqrt{\sec(c + dx)}} \\ &= \frac{\tanh^{-1}(\sin(c + dx))\sqrt{b \sec(c + dx)}}{d\sqrt{\sec(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 33, normalized size = 1.00

$$\frac{\sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{d\sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]], x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.69, size = 111, normalized size = 3.36

$$\left[\frac{\sqrt{b} \log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2d}, \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/2*sqrt(b)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/d, -sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(dx + c)} \sqrt{\sec(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))*sqrt(sec(d*x + c)), x)

maple [A] time = 0.98, size = 52, normalized size = 1.58

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) \sqrt{\frac{1}{\cos(dx+c)}} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2), x)

[Out] -2/d*arctanh((-1+cos(d*x+c))/sin(d*x+c))*(1/cos(d*x+c))^(1/2)*(b/cos(d*x+c))^(1/2)*cos(d*x+c)

maxima [B] time = 1.02, size = 65, normalized size = 1.97

$$\frac{\sqrt{b} \left(\log \left(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1 \right) - \log \left(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1 \right) \right)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 1/2*sqrt(b)*(log(cos(d*x + c)^2 + sin(d*x + c)^2 + 2*sin(d*x + c) + 1) - log(cos(d*x + c)^2 + sin(d*x + c)^2 - 2*sin(d*x + c) + 1))/d

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2),x)

[Out] int((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(c+dx)} \sqrt{\sec(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(1/2)*(b*sec(d*x+c))**(1/2),x)

[Out] Integral(sqrt(b*sec(c + d*x))*sqrt(sec(c + d*x)), x)

$$3.137 \quad \int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx$$

Optimal. Leaf size=24

$$\frac{x\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

[Out] $x*(b*\sec(d*x+c))^{(1/2)}/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[c + d*x]]/Sqrt[Sec[c + d*x]], x]`

[Out] `(x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} dx &= \frac{\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} \int 1 dx \\ &= \frac{x\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 24, normalized size = 1.00

$$\frac{x\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[c + d*x]]/Sqrt[Sec[c + d*x]], x]

[Out] (x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]

fricas [A] time = 0.60, size = 98, normalized size = 4.08

$$\left[\frac{\sqrt{-b} \log\left(-2\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b\right)}{2d}, \frac{\sqrt{b} \arctan\left(\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2), x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/d, sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(dx+c)}}{\sqrt{\sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))/sqrt(sec(d*x + c)), x)

maple [A] time = 1.03, size = 32, normalized size = 1.33

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} (dx+c)}{d \sqrt{\frac{1}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2), x)

[Out] 1/d*(b/cos(d*x+c))^(1/2)/(1/cos(d*x+c))^(1/2)*(d*x+c)

maxima [A] time = 0.50, size = 26, normalized size = 1.08

$$\frac{2\sqrt{b} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")

[Out] 2*sqrt(b)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

mupad [B] time = 0.20, size = 24, normalized size = 1.00

$$\frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{\sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(1/2),x)

[Out] (x*(b/cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2)

sympy [A] time = 0.58, size = 5, normalized size = 0.21

$$\sqrt{b}x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(1/2),x)

[Out] sqrt(b)*x

$$3.138 \quad \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}}$$

[Out] $\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[c + d*x]]/\text{Sec}[c + d*x]^{(3/2)}, x]$

[Out] $(\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[\text{Sec}[c + d*x]])$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/\text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x\} \&\& \text{!IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{3}{2}}(c+dx)} dx &= \frac{\sqrt{b \sec(c+dx)} \int \cos(c+dx) dx}{\sqrt{\sec(c+dx)}} \\ &= \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{d\sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 32, normalized size = 1.00

$$\frac{\sin(c + dx)\sqrt{b \sec(c + dx)}}{d\sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(3/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.67, size = 30, normalized size = 0.94

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2), x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/d

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(dx + c)}}{\sec(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2), x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))/sec(d*x + c)^(3/2), x)

maple [A] time = 1.16, size = 41, normalized size = 1.28

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{d \left(\frac{1}{\cos(dx+c)} \right)^{\frac{3}{2}} \cos(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2), x)

[Out] 1/d*(b/cos(d*x+c))^(1/2)*sin(d*x+c)/(1/cos(d*x+c))^(3/2)/cos(d*x+c)

maxima [A] time = 0.63, size = 13, normalized size = 0.41

$$\frac{\sqrt{b} \sin(dx + c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")

[Out] sqrt(b)*sin(d*x + c)/d

mupad [B] time = 0.22, size = 32, normalized size = 1.00

$$\frac{\sin(c + dx) \sqrt{\frac{b}{\cos(c+dx)}}}{d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(3/2),x)

[Out] (sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(d*(1/cos(c + d*x))^(1/2))

sympy [A] time = 13.39, size = 36, normalized size = 1.12

$$\begin{cases} \frac{\sqrt{b} \tan(c+dx)}{d \sec(c+dx)} & \text{for } d \neq 0 \\ \frac{x \sqrt{b \sec(c)}}{\sec^{\frac{3}{2}}(c)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(3/2),x)

[Out] Piecewise((sqrt(b)*tan(c + d*x)/(d*sec(c + d*x)), Ne(d, 0)), (x*sqrt(b*sec(c))/sec(c)**(3/2), True))

$$3.139 \quad \int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{5}{2}}(c+dx)} dx$$

Optimal. Leaf size=63

$$\frac{x\sqrt{b \sec(c+dx)}}{2\sqrt{\sec(c+dx)}} + \frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{2d \sec^{\frac{3}{2}}(c+dx)}$$

[Out] $1/2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(3/2)}+1/2*x*(b*\sec(d*x+c))^{(1/2)}/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{x\sqrt{b \sec(c+dx)}}{2\sqrt{\sec(c+dx)}} + \frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{2d \sec^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(5/2), x]

[Out] $(x*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(2*\text{Sqrt}[\text{Sec}[c + d*x]]) + (\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(2*d*\text{Sec}[c + d*x]^{(3/2)})$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n - 1)]/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{b \sec(c+dx)}}{\sec^2(c+dx)^{\frac{5}{2}}} dx &= \frac{\sqrt{b \sec(c+dx)} \int \cos^2(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{2d \sec^{\frac{3}{2}}(c+dx)} + \frac{\sqrt{b \sec(c+dx)} \int 1 dx}{2\sqrt{\sec(c+dx)}} \\
&= \frac{x\sqrt{b \sec(c+dx)}}{2\sqrt{\sec(c+dx)}} + \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{2d \sec^{\frac{3}{2}}(c+dx)}
\end{aligned}$$

Mathematica [A] time = 0.07, size = 45, normalized size = 0.71

$$\frac{(2(c+dx) + \sin(2(c+dx)))\sqrt{b \sec(c+dx)}}{4d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(5/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.78, size = 158, normalized size = 2.51

$$\left[\frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + \sqrt{-b} \log\left(-2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)\right)}{4d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(-b)*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(dx+c)}}{\sec(dx+c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))/sec(d*x + c)^(5/2), x)

maple [A] time = 1.30, size = 54, normalized size = 0.86

$$\frac{(\cos(dx+c)\sin(dx+c)+dx+c)\sqrt{\frac{b}{\cos(dx+c)}}}{2d\cos(dx+c)^2\left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*(b/cos(d*x+c))^(1/2)/cos(d*x+c)^2/(1/cos(d*x+c))^(5/2)

maxima [A] time = 0.80, size = 25, normalized size = 0.40

$$\frac{(2dx+2c+\sin(2dx+2c))\sqrt{b}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))*sqrt(b)/d

mupad [B] time = 0.44, size = 41, normalized size = 0.65

$$\frac{(\sin(2c+2dx)+2dx)\sqrt{\frac{b}{\cos(c+dx)}}}{4d\sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(5/2),x)

[Out] ((sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*d*(1/cos(c + d*x))^(1/2))

sympy [A] time = 110.95, size = 82, normalized size = 1.30

$$\begin{cases} \frac{\sqrt{b} x \tan^2(c+dx)}{2 \sec^2(c+dx)} + \frac{\sqrt{b} x}{2 \sec^2(c+dx)} + \frac{\sqrt{b} \tan(c+dx)}{2d \sec^2(c+dx)} & \text{for } d \neq 0 \\ \frac{x \sqrt{b \sec(c)}}{\sec^{\frac{5}{2}}(c)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(5/2),x)

[Out] Piecewise((sqrt(b)*x*tan(c + d*x)**2/(2*sec(c + d*x)**2) + sqrt(b)*x/(2*sec(c + d*x)**2) + sqrt(b)*tan(c + d*x)/(2*d*sec(c + d*x)**2), Ne(d, 0)), (x*sqrt(b*sec(c))/sec(c)**(5/2), True))

$$3.140 \quad \int \frac{\sqrt{b \sec(c+dx)}}{7 \sec^2(c+dx)} dx$$

Optimal. Leaf size=70

$$\frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{b \sec(c+dx)}}{3d\sqrt{\sec(c+dx)}}$$

[Out] $\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(1/2)}-1/3*\sin(d*x+c)^3*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{b \sec(c+dx)}}{3d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(7/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]]) - (Sqrt[b*Sec[c + d*x]]*Sin[c + d*x]^3)/(3*d*Sqrt[Sec[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{7}{2}}(c+dx)} dx &= \frac{\sqrt{b \sec(c+dx)} \int \cos^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= -\frac{\sqrt{b \sec(c+dx)} \operatorname{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{d\sqrt{\sec(c+dx)}} \\
&= \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{d\sqrt{\sec(c+dx)}} - \frac{\sqrt{b \sec(c+dx)} \sin^3(c+dx)}{3d\sqrt{\sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.12, size = 45, normalized size = 0.64

$$\frac{\sin(c+dx)(\cos(2(c+dx))+5)\sqrt{b \sec(c+dx)}}{6d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(7/2), x]

[Out] ((5 + Cos[2*(c + d*x)])*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(6*d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.88, size = 48, normalized size = 0.69

$$\frac{(\cos(dx+c)^3 + 2 \cos(dx+c)) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3d\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2), x, algorithm="fricas")

[Out] 1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(dx+c)}}{\sec(dx+c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2), x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))/sec(d*x + c)^(7/2), x)

maple [A] time = 1.39, size = 52, normalized size = 0.74

$$\frac{(2 + \cos^2(dx + c)) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx + c)}{3d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}} \cos(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2), x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*(b/cos(d*x+c))^(1/2)*sin(d*x+c)/(1/cos(d*x+c))^(7/2)/cos(d*x+c)^3

maxima [A] time = 1.05, size = 42, normalized size = 0.60

$$\frac{\sqrt{b} \left(\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan\left(\frac{\sin(3dx + 3c)}{\cos(3dx + 3c)}\right)\right) \right)}{12d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(7/2), x, algorithm="maxima")

[Out] 1/12*sqrt(b)*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/d

mupad [B] time = 0.52, size = 45, normalized size = 0.64

$$\frac{(9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(7/2), x)

[Out] ((9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(7/2),x)
```

```
[Out] Timed out
```

$$3.141 \quad \int \frac{\sqrt{b \sec(c+dx)}}{9 \sec^2(c+dx)} dx$$

Optimal. Leaf size=98

$$\frac{3x\sqrt{b \sec(c+dx)}}{8\sqrt{\sec(c+dx)}} + \frac{3 \sin(c+dx)\sqrt{b \sec(c+dx)}}{8d \sec^{\frac{3}{2}}(c+dx)} + \frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{4d \sec^{\frac{7}{2}}(c+dx)}$$

[Out] 1/4*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(7/2)+3/8*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(3/2)+3/8*x*(b*sec(d*x+c))^(1/2)/sec(d*x+c)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3x\sqrt{b \sec(c+dx)}}{8\sqrt{\sec(c+dx)}} + \frac{3 \sin(c+dx)\sqrt{b \sec(c+dx)}}{8d \sec^{\frac{3}{2}}(c+dx)} + \frac{\sin(c+dx)\sqrt{b \sec(c+dx)}}{4d \sec^{\frac{7}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(9/2), x]

[Out] (3*x*Sqrt[b*Sec[c + d*x]])/(8*Sqrt[Sec[c + d*x]]) + (Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(4*d*Sec[c + d*x]^(7/2)) + (3*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(8*d*Sec[c + d*x]^(3/2))

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_.))^(m_.)*((b_.)*(v_.))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIn[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIn[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{b \sec(c+dx)}}{\sec^{\frac{9}{2}}(c+dx)} dx &= \frac{\sqrt{b \sec(c+dx)} \int \cos^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{4d \sec^{\frac{7}{2}}(c+dx)} + \frac{(3\sqrt{b \sec(c+dx)}) \int \cos^2(c+dx) dx}{4\sqrt{\sec(c+dx)}} \\
&= \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{4d \sec^{\frac{7}{2}}(c+dx)} + \frac{3\sqrt{b \sec(c+dx)} \sin(c+dx)}{8d \sec^{\frac{3}{2}}(c+dx)} + \frac{(3\sqrt{b \sec(c+dx)}) \int 1 dx}{8\sqrt{\sec(c+dx)}} \\
&= \frac{3x\sqrt{b \sec(c+dx)}}{8\sqrt{\sec(c+dx)}} + \frac{\sqrt{b \sec(c+dx)} \sin(c+dx)}{4d \sec^{\frac{7}{2}}(c+dx)} + \frac{3\sqrt{b \sec(c+dx)} \sin(c+dx)}{8d \sec^{\frac{3}{2}}(c+dx)}
\end{aligned}$$

Mathematica [A] time = 0.12, size = 55, normalized size = 0.56

$$\frac{(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx)))\sqrt{b \sec(c+dx)}}{32d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[c + d*x]]/Sec[c + d*x]^(9/2), x]

[Out] (Sqrt[b*Sec[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])) / (32*d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.68, size = 202, normalized size = 2.06

$$\left[\frac{2(2 \cos(dx+c)^4 + 3 \cos(dx+c)^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}} + 3 \sqrt{-b} \log \left(-2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c) \right) \right] / 16d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2), x, algorithm="fricas")

[Out] [1/16*(2*(2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) + 3*sqrt(-b)*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/8*((2*cos(d

$(x + c)^4 + 3 \cos(dx + c)^2 \sqrt{b/\cos(dx + c)} \sin(dx + c) / \sqrt{\cos(dx + c)} + 3 \sqrt{b} \arctan(\sqrt{b/\cos(dx + c)} \sin(dx + c) / (\sqrt{b} \sqrt{\cos(dx + c)})) / d]$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(dx + c)}}{\sec(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(d*x + c))/sec(d*x + c)^(9/2), x)

maple [A] time = 1.36, size = 74, normalized size = 0.76

$$\frac{(2(\cos^3(dx + c)) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c) \sqrt{\frac{b}{\cos(dx + c)}}}{8d \left(\frac{1}{\cos(dx + c)}\right)^{\frac{9}{2}} \cos(dx + c)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2),x)

[Out] 1/8/d*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)*(b/cos(d*x+c))^(1/2)/(1/cos(d*x+c))^(9/2)/cos(d*x+c)^4

maxima [A] time = 0.78, size = 49, normalized size = 0.50

$$\frac{\left(12 dx + 12 c + \sin(4 dx + 4 c) + 8 \sin\left(\frac{1}{2} \arctan\left(\frac{\sin(4 dx + 4 c)}{\cos(4 dx + 4 c)}\right)\right)\right) \sqrt{b}}{32 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/2)/sec(d*x+c)^(9/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

mupad [B] time = 0.71, size = 52, normalized size = 0.53

$$\frac{\sqrt{\frac{b}{\cos(c + dx)}} (8 \sin(2c + 2dx) + \sin(4c + 4dx) + 12dx)}{32d \sqrt{\frac{1}{\cos(c + dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b/cos(c + d*x))^(1/2)/(1/cos(c + d*x))^(9/2),x)
```

```
[Out] ((b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(32*d*(1/cos(c + d*x))^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(1/2)/sec(d*x+c)**(9/2),x)
```

```
[Out] Timed out
```

3.142 $\int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=110

$$\frac{b \sin(c + dx) \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{4d} + \frac{3b \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{8d} + \frac{3b \sqrt{b \sec(c + dx)} \tanh^{-1}(\sqrt{\sec(c + dx)})}{8d \sqrt{\sec(c + dx)}}$$

[Out] $3/8*b*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d+1/4*b*\sec(d*x+c)^{(7/2)}*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d+3/8*b*\arctanh(\sin(d*x+c))*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{b \sin(c + dx) \sec^{\frac{7}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{4d} + \frac{3b \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{8d} + \frac{3b \sqrt{b \sec(c + dx)} \tanh^{-1}(\sqrt{\sec(c + dx)})}{8d \sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(3/2), x]

[Out] $(3*b*\text{ArcTanh}[\text{Sin}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(8*d*\text{Sqrt}[\text{Sec}[c + d*x]]) + (3*b*\text{Sec}[c + d*x]^{(3/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(8*d) + (b*\text{Sec}[c + d*x]^{(7/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(4*d)$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := -Simp[(b*Csc[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx &= \frac{(b\sqrt{b \sec(c+dx)}) \int \sec^5(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{b \sec^{\frac{7}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{4d} + \frac{(3b\sqrt{b \sec(c+dx)}) \int \sec^3(c+dx) dx}{4\sqrt{\sec(c+dx)}} \\
&= \frac{3b \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{8d} + \frac{b \sec^{\frac{7}{2}}(c+dx) \sqrt{b \sec(c+dx)}}{4d} \\
&= \frac{3b \tanh^{-1}(\sin(c+dx)) \sqrt{b \sec(c+dx)}}{8d \sqrt{\sec(c+dx)}} + \frac{3b \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)}}{8d}
\end{aligned}$$

Mathematica [A] time = 0.12, size = 64, normalized size = 0.58

$$\frac{(b \sec(c+dx))^{3/2} (3 \tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx) (2 \sec^2(c+dx) + 3))}{8d \sec^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(3/2), x]

[Out] ((b*Sec[c + d*x])^(3/2)*(3*ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*(3 + 2*Sec[c + d*x]^2)*Tan[c + d*x]))/(8*d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.82, size = 236, normalized size = 2.15

$$\left[\frac{3 b^{\frac{3}{2}} \cos(dx+c)^3 \log\left(-\frac{b \cos(dx+c)^2 - 2 \sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2(3b \cos(dx+c)^2 + 2b) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{16 d \cos(dx+c)^3}, \dots \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [1/16*(3*b^(3/2)*cos(d*x + c)^3*log(-(b*cos(d*x + c)^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*(3*b*cos(d*x + c)^2 + 2*b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c))

))/((d*cos(d*x + c)^3), -1/8*(3*sqrt(-b)*b*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c)^3 - (3*b*cos(d*x + c)^2 + 2*b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)^3)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(7/2), x)

maple [A] time = 0.94, size = 131, normalized size = 1.19

$$\frac{3 \left(\cos^4(dx + c) \right) \ln \left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)} \right) - 3 \left(\cos^4(dx + c) \right) \ln \left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)} \right) + 3 \left(\cos^2(dx + c) \right) \sin(dx + c)}{8d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x)

[Out] 1/8/d*(3*cos(d*x+c)^4*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-3*cos(d*x+c)^4*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))+3*cos(d*x+c)^2*sin(d*x+c)+2*sin(d*x+c))*cos(d*x+c)*(1/cos(d*x+c))^(7/2)*(b/cos(d*x+c))^(3/2)

maxima [B] time = 1.18, size = 1742, normalized size = 15.84

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] -1/16*(12*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 44*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 44*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 12*(b*sin(8*d*x + 8*c) + 4*b*sin(6*d*x + 6*c) + 6*b*sin(4*d*x + 4*c) + 4*b*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 3*(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x + 4*c)^2

```

+ 16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x + 6*c)^2
+ 36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b*s
in(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*co
s(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c) + 4*b*cos(2*
d*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) + b)*cos(4*d*x
+ 4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*b*sin(4*d*x + 4
*c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(4*d*x + 4*c) + 2
*b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b*log(cos(1/2*arctan2(sin(2*d*x +
2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x +
2*c))))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + 3*
(b*cos(8*d*x + 8*c)^2 + 16*b*cos(6*d*x + 6*c)^2 + 36*b*cos(4*d*x + 4*c)^2 +
16*b*cos(2*d*x + 2*c)^2 + b*sin(8*d*x + 8*c)^2 + 16*b*sin(6*d*x + 6*c)^2 +
36*b*sin(4*d*x + 4*c)^2 + 48*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*b*si
n(2*d*x + 2*c)^2 + 2*(4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos
(2*d*x + 2*c) + b)*cos(8*d*x + 8*c) + 8*(6*b*cos(4*d*x + 4*c) + 4*b*cos(2*d
*x + 2*c) + b)*cos(6*d*x + 6*c) + 12*(4*b*cos(2*d*x + 2*c) + b)*cos(4*d*x +
4*c) + 8*b*cos(2*d*x + 2*c) + 4*(2*b*sin(6*d*x + 6*c) + 3*b*sin(4*d*x + 4*
c) + 2*b*sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 16*(3*b*sin(4*d*x + 4*c) + 2*
b*sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + b*log(cos(1/2*arctan2(sin(2*d*x + 2
*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 12*
(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*cos
(2*d*x + 2*c) + b)*sin(7/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4
4*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b*c
os(2*d*x + 2*c) + b)*sin(5/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) +
44*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4*b
*cos(2*d*x + 2*c) + b)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))
+ 12*(b*cos(8*d*x + 8*c) + 4*b*cos(6*d*x + 6*c) + 6*b*cos(4*d*x + 4*c) + 4
*b*cos(2*d*x + 2*c) + b)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
))) * sqrt(b) / ((2*(4*cos(6*d*x + 6*c) + 6*cos(4*d*x + 4*c) + 4*cos(2*d*x + 2*
c) + 1)*cos(8*d*x + 8*c) + cos(8*d*x + 8*c)^2 + 8*(6*cos(4*d*x + 4*c) + 4*c
os(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 16*cos(6*d*x + 6*c)^2 + 12*(4*cos(2
*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 36*cos(4*d*x + 4*c)^2 + 16*cos(2*d*x +
2*c)^2 + 4*(2*sin(6*d*x + 6*c) + 3*sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*s
in(8*d*x + 8*c) + sin(8*d*x + 8*c)^2 + 16*(3*sin(4*d*x + 4*c) + 2*sin(2*d*x
+ 2*c))*sin(6*d*x + 6*c) + 16*sin(6*d*x + 6*c)^2 + 36*sin(4*d*x + 4*c)^2 +
48*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 16*sin(2*d*x + 2*c)^2 + 8*cos(2*d*x
+ 2*c) + 1)*d)

```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{3/2} \left(\frac{1}{\cos(c + dx)} \right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(7/2), x)
```

```
[Out] int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(7/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**(7/2)*(b*sec(d*x+c))**(3/2), x)
```

```
[Out] Timed out
```

$$3.143 \quad \int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^{3/2} dx$$

Optimal. Leaf size=72

$$\frac{b \sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{b \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

[Out] $1/3*b*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)^3*(b*\sec(d*x+c))^{(1/2)}/d+b*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{b \sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{b \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2), x]`

[Out] `(b*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d + (b*Sec[c + d*x])^(5/2)*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x]^3/(3*d)`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 3767

`Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]`

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx &= \frac{(b\sqrt{b \sec(c+dx)}) \int \sec^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= -\frac{(b\sqrt{b \sec(c+dx)}) \operatorname{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d\sqrt{\sec(c+dx)}} \\
&= \frac{b\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d} + \frac{b \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}}{3d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 45, normalized size = 0.62

$$\frac{\left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx)\right) (b \sec(c+dx))^{3/2}}{d \sec^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2), x]

[Out] ((b*Sec[c + d*x])^(3/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.82, size = 44, normalized size = 0.61

$$\frac{(2b \cos(dx+c)^2 + b) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3d \cos(dx+c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] 1/3*(2*b*cos(d*x + c)^2 + b)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(5/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{\frac{3}{2}} \sec(dx+c)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(5/2), x)

maple [A] time = 0.88, size = 52, normalized size = 0.72

$$\frac{(2(\cos^2(dx+c)) + 1) \left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}} \cos(dx+c) \sin(dx+c)}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2), x)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*(1/cos(d*x+c))^(5/2)*(b/cos(d*x+c))^(3/2)*cos(d*x+c)*sin(d*x+c)

maxima [B] time = 1.00, size = 299, normalized size = 4.15

$$\frac{4(3b \cos(6dx + 6c) \sin(2dx + 2c) + 9b \cos(4dx + 4c) \sin(2dx + 2c) - (3b \cos(2dx + 2c) + b) \sin(6dx + 6c) - 3(3b \cos(2dx + 2c) + b) \sin(4dx + 4c)) \sqrt{b}}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1) \cos(4dx + 4c) + 9 \cos(4dx + 4c)^2 + 9 \cos(2dx + 2c)^2 + 6(\sin(4dx + 4c) + \sin(2dx + 2c)) \sin(6dx + 6c) + \sin(6dx + 6c)^2 + 9 \sin(4dx + 4c)^2 + 18 \sin(4dx + 4c) \sin(2dx + 2c) + 9 \sin(2dx + 2c)^2 + 6 \cos(2dx + 2c) + 1) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^(3/2), x, algorithm="maxima")

[Out] -4/3*(3*b*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b*cos(2*d*x + 2*c) + b)*sin(6*d*x + 6*c) - 3*(3*b*cos(2*d*x + 2*c) + b)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 1.36, size = 127, normalized size = 1.76

$$\frac{2b \cos(c + dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} (4 \sin(c + dx) + 5 \sin(3c + 3dx) + \sin(5c + 5dx) + \cos(c + dx) \cos(3c + 3dx) + \cos(5c + 5dx))}{3d (10 \cos(c + dx) + 5 \cos(3c + 3dx) + \cos(5c + 5dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(5/2), x)

[Out] (2*b*cos(c + d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*(cos(c + d*x)*10i + 4*sin(c + d*x) + cos(3*c + 3*d*x)*5i + cos(5*c + 5*d*x)*1i + 5*sin(3*c + 3*d*x) + sin(5*c + 5*d*x)))/(3*d*(10*cos(c + d*x) + 5*cos(3*c + 3*d*x) + cos(5*c + 5*d*x)))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(5/2)*(b*sec(d*x+c))**(3/2),x)

[Out] Timed out

3.144 $\int \sec^2(c + dx)(b \sec(c + dx))^{3/2} dx$

Optimal. Leaf size=74

$$\frac{b \sin(c + dx) \sec^2(c + dx) \sqrt{b \sec(c + dx)}}{2d} + \frac{b \sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

[Out] $1/2*b*\sec(d*x+c)^(3/2)*\sin(d*x+c)*(b*\sec(d*x+c))^(1/2)/d+1/2*b*\operatorname{arctanh}(\sin(d*x+c))*(b*\sec(d*x+c))^(1/2)/d/\sec(d*x+c)^(1/2)$

Rubi [A] time = 0.02, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{b \sin(c + dx) \sec^2(c + dx) \sqrt{b \sec(c + dx)}}{2d} + \frac{b \sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sec}[c + d*x]^{(3/2)}*(b*\operatorname{Sec}[c + d*x])^{(3/2)}, x]$

[Out] $(b*\operatorname{ArcTanh}[\operatorname{Sin}[c + d*x]]*\operatorname{Sqrt}[b*\operatorname{Sec}[c + d*x]])/(2*d*\operatorname{Sqrt}[\operatorname{Sec}[c + d*x]]) + (b*\operatorname{Sec}[c + d*x]^{(3/2)}*\operatorname{Sqrt}[b*\operatorname{Sec}[c + d*x]]*\operatorname{Sin}[c + d*x])/(2*d)$

Rule 17

$\operatorname{Int}[(u_*)*((a_*)*(v_))^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\operatorname{Sqrt}[b*v])/ \operatorname{Sqrt}[a*v], \operatorname{Int}[u*v^{(m+n)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, m\}, x \&\& \text{!IntegerQ}[m] \&\& \operatorname{IGtQ}[n+1/2, 0] \&\& \operatorname{IntegerQ}[m+n]$

Rule 3768

$\operatorname{Int}[(\operatorname{csc}[(c_*) + (d_*)*(x_*)])*(b_*)^{(n_*)}, x_Symbol] \rightarrow -\operatorname{Simp}[(b*\operatorname{Cos}[c + d*x])*(b*\operatorname{Csc}[c + d*x])^{(n-1)})/(d*(n-1)), x] + \operatorname{Dist}[(b^2*(n-2))/(n-1), \operatorname{Int}[(b*\operatorname{Csc}[c + d*x])^{(n-2)}, x], x] /;$ $\operatorname{FreeQ}\{b, c, d\}, x \&\& \operatorname{GtQ}[n, 1] \&\& \operatorname{IntegerQ}[2*n]$

Rule 3770

$\operatorname{Int}[\operatorname{csc}[(c_*) + (d_*)*(x_*)], x_Symbol] \rightarrow -\operatorname{Simp}[\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]]/d, x] /;$ $\operatorname{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2} dx &= \frac{(b\sqrt{b \sec(c+dx)}) \int \sec^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{b \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{2d} + \frac{(b\sqrt{b \sec(c+dx)}) \int \sec(c+dx) dx}{2\sqrt{\sec(c+dx)}} \\
&= \frac{b \tanh^{-1}(\sin(c+dx)) \sqrt{b \sec(c+dx)}}{2d\sqrt{\sec(c+dx)}} + \frac{b \sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{2d}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 50, normalized size = 0.68

$$\frac{(b \sec(c+dx))^{3/2} (\tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx))}{2d \sec^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2), x]

[Out] ((b*Sec[c + d*x])^(3/2)*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.85, size = 202, normalized size = 2.73

$$\left[\frac{b^{\frac{3}{2}} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2b \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4d \cos(dx+c)}, \sqrt{-b} b \arctan\left(\frac{\sqrt{-b}}{\dots}\right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [1/4*(b^(3/2)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*b*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)), -1/2*(sqrt(-b)*b*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b*cos(d*x + c) - b*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(3/2), x)

maple [A] time = 0.90, size = 114, normalized size = 1.54

$$\frac{\left((\cos^2(dx + c)) \ln\left(-\frac{-1 + \cos(dx + c) + \sin(dx + c)}{\sin(dx + c)}\right) - (\cos^2(dx + c)) \ln\left(-\frac{-\sin(dx + c) - 1 + \cos(dx + c)}{\sin(dx + c)}\right) - \sin(dx + c) \right) \cos(dx + c)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*cos(d*x+c)*(1/cos(d*x+c))^(3/2)*(b/cos(d*x+c))^(3/2)

maxima [B] time = 0.92, size = 691, normalized size = 9.34

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] -1/4*(4*(b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b*sin(4*d*x + 4*c) + 2*b*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b*cos(4*d*x + 4*c))^2 + 4*b*cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) + (b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2*d*x + 2*c) + b)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))) + 1) - 4*(b*cos(4*d*x + 4*c) + 2*b*cos(2*d*x + 2*c) + b)*sin(3/2*ar

```
ctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b*cos(4*d*x + 4*c) + 2*b*cos(2*d*x + 2*c) + b)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) * sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{3/2} \left(\frac{1}{\cos(c + dx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(3/2), x)
```

```
[Out] int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(3/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**(3/2), x)
```

```
[Out] Timed out
```

$$3.145 \quad \int \sqrt{\sec(c + dx)} (b \sec(c + dx))^{3/2} dx$$

Optimal. Leaf size=33

$$\frac{b \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

[Out] b*sin(d*x+c)*sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(1/2)/d

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{b \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2), x]

[Out] (b*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/d

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned}
\int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{3/2} dx &= \frac{(b\sqrt{b \sec(c+dx)}) \int \sec^2(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= -\frac{(b\sqrt{b \sec(c+dx)}) \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{d\sqrt{\sec(c+dx)}} \\
&= \frac{b\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 0.97

$$\frac{\sin(c+dx)(b \sec(c+dx))^{3/2}}{d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2), x]

[Out] ((b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])

fricas [A] time = 0.66, size = 31, normalized size = 0.94

$$\frac{b\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{d\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] b*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{\frac{3}{2}} \sqrt{\sec(dx+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)*sqrt(sec(d*x + c)), x)

maple [A] time = 1.01, size = 39, normalized size = 1.18

$$\frac{\sqrt{\frac{1}{\cos(dx+c)}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}} \cos(dx+c) \sin(dx+c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2), x)`

[Out] `1/d*(1/cos(d*x+c))^(1/2)*(b/cos(d*x+c))^(3/2)*cos(d*x+c)*sin(d*x+c)`

maxima [A] time = 0.73, size = 54, normalized size = 1.64

$$\frac{2 b^{\frac{3}{2}} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c)^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(3/2), x, algorithm="maxima")`

[Out] `2*b^(3/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c)^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d)`

mupad [B] time = 0.22, size = 47, normalized size = 1.42

$$\frac{b (\cos(dx) - \sin(dx) 1i) (\sin(c) + \cos(c) 1i) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}}}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(1/2), x)`

[Out] `(b*(cos(d*x) - sin(d*x)*1i)*(cos(c)*1i + sin(c))*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/d`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{3}{2}} \sqrt{\sec(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(1/2)*(b*sec(d*x+c))**(3/2), x)`

[Out] `Integral((b*sec(c + d*x))**(3/2)*sqrt(sec(c + d*x)), x)`

$$3.146 \quad \int \frac{(b \sec(c+dx))^{3/2}}{\sqrt{\sec(c+dx)}} dx$$

Optimal. Leaf size=34

$$\frac{b\sqrt{b \sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\sec(c+dx)}}$$

[Out] b*arctanh(sin(d*x+c))*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{b\sqrt{b \sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(3/2)/Sqrt[Sec[c + d*x]],x]

[Out] (b*ArcTanh[Sin[c + d*x]]*Sqrt[b*Sec[c + d*x]])/(d*Sqrt[Sec[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c+dx))^{3/2}}{\sqrt{\sec(c+dx)}} dx &= \frac{(b\sqrt{b \sec(c+dx)}) \int \sec(c+dx) dx}{\sqrt{\sec(c+dx)}} \\ &= \frac{b \tanh^{-1}(\sin(c+dx))\sqrt{b \sec(c+dx)}}{d\sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 33, normalized size = 0.97

$$\frac{(b \sec(c + dx))^{3/2} \tanh^{-1}(\sin(c + dx))}{d \sec^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2)/Sqrt[Sec[c + d*x]], x]

[Out] (ArcTanh[Sin[c + d*x]]*(b*Sec[c + d*x])^(3/2))/(d*Sec[c + d*x]^(3/2))

fricas [A] time = 1.10, size = 112, normalized size = 3.29

$$\left[\frac{b^2 \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2d}, \frac{\sqrt{-b} b \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2), x, algorithm="fricas")

[Out] [1/2*b^(3/2)*log(-(b*cos(d*x + c)^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/d, -sqrt(-b)*b*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{3/2}}{\sqrt{\sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)/sqrt(sec(d*x + c)), x)

maple [A] time = 1.28, size = 52, normalized size = 1.53

$$\frac{2 \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right) \cos(dx+c) \left(\frac{b}{\cos(dx+c)}\right)^{3/2}}{d \sqrt{\frac{1}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x)`

[Out] $-2/d*\operatorname{arctanh}((-1+\cos(dx+c))/\sin(dx+c))*\cos(dx+c)*(b/\cos(dx+c))^{3/2}/(1/\cos(dx+c))^{1/2}$

maxima [B] time = 0.78, size = 68, normalized size = 2.00

$$\frac{(b \log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - b \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1))}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

[Out] $1/2*(b*\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2*\sin(dx+c) + 1) - b*\log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2*\sin(dx+c) + 1))*\operatorname{sqrt}(b)/d$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}}{\sqrt{\frac{1}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c+d*x))^(3/2)/(1/cos(c+d*x))^(1/2),x)`

[Out] `int((b/cos(c+d*x))^(3/2)/(1/cos(c+d*x))^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(c+dx))^{\frac{3}{2}}}{\sqrt{\sec(c+dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(1/2),x)`

[Out] `Integral((b*sec(c+d*x))**(3/2)/sqrt(sec(c+d*x)),x)`

$$3.147 \quad \int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=25

$$\frac{bx\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

[Out] $b*x*(b*\sec(d*x+c))^{(1/2)}/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{bx\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c+d*x])^{(3/2)}/\text{Sec}[c+d*x]^{(3/2)},x]$

[Out] $(b*x*\text{Sqrt}[b*\text{Sec}[c+d*x]])/\text{Sqrt}[\text{Sec}[c+d*x]]$

Rule 8

$\text{Int}[a_, x_Symbol] := \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)*((b_.)*(v_))^{(n_)}], x_Symbol] := \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& !\text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx &= \frac{(b\sqrt{b \sec(c+dx)})}{\sqrt{\sec(c+dx)}} \int 1 dx \\ &= \frac{bx\sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 24, normalized size = 0.96

$$\frac{x(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(3/2), x]

[Out] (x*(b*Sec[c + d*x])^(3/2))/Sec[c + d*x]^(3/2)

fricas [A] time = 0.79, size = 99, normalized size = 3.96

$$\left[\frac{\sqrt{-b} b \log \left(-2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{2d}, \frac{b^{\frac{3}{2}} \arctan \left(\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}} \right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2), x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*b*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/d, b^(3/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx+c))^{\frac{3}{2}}}{\sec(dx+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(3/2), x)

maple [A] time = 0.98, size = 32, normalized size = 1.28

$$\frac{\left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}(dx+c)}{d\left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2), x)

[Out] 1/d*(b/cos(d*x+c))^(3/2)/(1/cos(d*x+c))^(3/2)*(d*x+c)

maxima [A] time = 0.77, size = 26, normalized size = 1.04

$$\frac{2 b^{\frac{3}{2}} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")

[Out] 2*b^(3/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

mupad [B] time = 0.20, size = 25, normalized size = 1.00

$$\frac{b x \sqrt{\frac{b}{\cos(c+d x)}}}{\sqrt{\frac{1}{\cos(c+d x)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(3/2),x)

[Out] (b*x*(b/cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2)

sympy [A] time = 18.08, size = 5, normalized size = 0.20

$$b^{\frac{3}{2}}x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(3/2),x)

[Out] b**(3/2)*x

$$3.148 \quad \int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=33

$$\frac{b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

[Out] b*sin(d*x+c)*(b*sec(d*x+c))^(1/2)/d/sec(d*x+c)^(1/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(5/2), x]

[Out] (b*Sqrt[b*Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[Sec[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2637

Int[sin[Pi/2 + (c_.) + (d_.)*(x_.)], x_Symbol] := Simp[Sin[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx &= \frac{(b \sqrt{b \sec(c+dx)}) \int \cos(c+dx) dx}{\sqrt{\sec(c+dx)}} \\ &= \frac{b \sqrt{b \sec(c+dx)} \sin(c+dx)}{d \sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 32, normalized size = 0.97

$$\frac{\sin(c + dx)(b \sec(c + dx))^{3/2}}{d \sec^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(5/2), x]

[Out] ((b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.65, size = 31, normalized size = 0.94

$$\frac{b \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2), x, algorithm="fricas")

[Out] b*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/d

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{3}{2}}}{\sec(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(5/2), x)

maple [A] time = 1.10, size = 41, normalized size = 1.24

$$\frac{\sin(dx + c) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}{d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}} \cos(dx + c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2), x)

[Out] 1/d*sin(d*x+c)*(b/cos(d*x+c))^(3/2)/(1/cos(d*x+c))^(5/2)/cos(d*x+c)

maxima [A] time = 0.69, size = 13, normalized size = 0.39

$$\frac{b^{\frac{3}{2}} \sin(dx + c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")

[Out] b^(3/2)*sin(d*x + c)/d

mupad [B] time = 0.25, size = 33, normalized size = 1.00

$$\frac{b \sin(c + dx) \sqrt{\frac{b}{\cos(c+dx)}}}{d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(5/2),x)

[Out] (b*sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(5/2),x)

[Out] Timed out

$$3.149 \quad \int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=65

$$\frac{bx\sqrt{b \sec(c+dx)}}{2\sqrt{\sec(c+dx)}} + \frac{b \sin(c+dx)\sqrt{b \sec(c+dx)}}{2d \sec^{\frac{3}{2}}(c+dx)}$$

[Out] $1/2*b*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(3/2)}+1/2*b*x*(b*\sec(d*x+c))^{(1/2)}/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{bx\sqrt{b \sec(c+dx)}}{2\sqrt{\sec(c+dx)}} + \frac{b \sin(c+dx)\sqrt{b \sec(c+dx)}}{2d \sec^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c+d*x])^{(3/2)}/\text{Sec}[c+d*x]^{(7/2)},x]$

[Out] $(b*x*\text{Sqrt}[b*\text{Sec}[c+d*x]])/(2*\text{Sqrt}[\text{Sec}[c+d*x]]) + (b*\text{Sqrt}[b*\text{Sec}[c+d*x]]* \text{Sin}[c+d*x])/(2*d*\text{Sec}[c+d*x]^{(3/2)})$

Rule 8

$\text{Int}[a_, x_Symbol] \text{ :> Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m)}*((b_.)*(v_))^{(n)}, x_Symbol] \text{ :> Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] \text{ /; FreeQ}[\{a, b, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n+1/2, 0] \ \&\& \ \text{IntegerQ}[m+n]$

Rule 2635

$\text{Int}[(b_.)*\sin[(c_.)+(d_.)*(x_)]^{(n)}, x_Symbol] \text{ :> -Simp}[(b*\text{Cos}[c+d*x])* (b*\text{Sin}[c+d*x])^{(n-1)})/(d*n), x] + \text{Dist}[(b^{2*(n-1)})/n, \text{Int}[(b*\text{Sin}[c+d*x])^{(n-2)}, x], x] \text{ /; FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
\int \frac{(b \sec(c + dx))^{3/2}}{\sec^7(c + dx)} dx &= \frac{(b\sqrt{b \sec(c + dx)}) \int \cos^2(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
&= \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{2d \sec^3(c + dx)} + \frac{(b\sqrt{b \sec(c + dx)}) \int 1 dx}{2\sqrt{\sec(c + dx)}} \\
&= \frac{bx\sqrt{b \sec(c + dx)}}{2\sqrt{\sec(c + dx)}} + \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{2d \sec^3(c + dx)}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 45, normalized size = 0.69

$$\frac{(2(c + dx) + \sin(2(c + dx)))(b \sec(c + dx))^{3/2}}{4d \sec^3(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(7/2), x]

[Out] ((b*Sec[c + d*x])^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.85, size = 161, normalized size = 2.48

$$\left[\frac{2b\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + \sqrt{-b} b \log\left(-2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)\right)}{4d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2), x, algorithm="fricas")

[Out] [1/4*(2*b*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(-b)*b*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/2*(b*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + b^(3/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{3}{2}}}{\sec(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(7/2), x)

maple [A] time = 1.19, size = 54, normalized size = 0.83

$$\frac{(\cos(dx + c) \sin(dx + c) + dx + c) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}{2d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}} \cos(dx + c)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*(b/cos(d*x+c))^(3/2)/(1/cos(d*x+c))^(7/2)/cos(d*x+c)^2

maxima [A] time = 0.67, size = 28, normalized size = 0.43

$$\frac{(2(dx + c)b + b \sin(2dx + 2c))\sqrt{b}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(7/2),x, algorithm="maxima")

[Out] 1/4*(2*(d*x + c)*b + b*sin(2*d*x + 2*c))*sqrt(b)/d

mupad [B] time = 0.43, size = 42, normalized size = 0.65

$$\frac{b (\sin(2c + 2dx) + 2dx) \sqrt{\frac{b}{\cos(c+dx)}}}{4d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(7/2),x)

```
[Out] (b*(sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*d*(1/cos(c + d*x))^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(7/2), x)
```

```
[Out] Timed out
```

$$3.150 \quad \int \frac{(b \sec(c+dx))^{3/2}}{9 \sec^2(c+dx)} dx$$

Optimal. Leaf size=72

$$\frac{b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}} - \frac{b \sin^3(c+dx) \sqrt{b \sec(c+dx)}}{3d \sqrt{\sec(c+dx)}}$$

[Out] $b \sin(d*x+c) * (b \sec(d*x+c))^{(1/2)} / d / \sec(d*x+c)^{(1/2)} - 1/3 * b \sin(d*x+c)^3 * (b \sec(d*x+c))^{(1/2)} / d / \sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{b \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}} - \frac{b \sin^3(c+dx) \sqrt{b \sec(c+dx)}}{3d \sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b \text{Sec}[c + d*x])^{(3/2)} / \text{Sec}[c + d*x]^{(9/2)}, x]$

[Out] $(b \text{Sqrt}[b \text{Sec}[c + d*x]] * \text{Sin}[c + d*x]) / (d \text{Sqrt}[\text{Sec}[c + d*x]]) - (b \text{Sqrt}[b \text{Sec}[c + d*x]] * \text{Sin}[c + d*x]^3) / (3 * d \text{Sqrt}[\text{Sec}[c + d*x]])$

Rule 17

$\text{Int}[(u_.) * ((a_.) * (v_.))^{(m_.)} * ((b_.) * (v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)} * b^{(n-1/2)} * \text{Sqrt}[b*v]) / \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

$\text{Int}[\sin[(c_.) + (d_.) * (x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{((n-1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(b \sec(c + dx))^{3/2}}{\sec^2(c + dx)} dx &= \frac{(b\sqrt{b \sec(c + dx)}) \int \cos^3(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
&= -\frac{(b\sqrt{b \sec(c + dx)}) \text{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d\sqrt{\sec(c + dx)}} \\
&= \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{d\sqrt{\sec(c + dx)}} - \frac{b\sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d\sqrt{\sec(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 45, normalized size = 0.62

$$\frac{\sin(c + dx)(\cos(2(c + dx)) + 5)(b \sec(c + dx))^{3/2}}{6d \sec^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(9/2), x]

[Out] ((5 + Cos[2*(c + d*x)])*(b*Sec[c + d*x])^(3/2)*Sin[c + d*x])/(6*d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.44, size = 51, normalized size = 0.71

$$\frac{(b \cos(dx + c)^3 + 2b \cos(dx + c)) \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)}{3d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2), x, algorithm="fricas")

[Out] 1/3*(b*cos(d*x + c)^3 + 2*b*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{3/2}}{\sec^2(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(9/2), x)

maple [A] time = 1.10, size = 52, normalized size = 0.72

$$\frac{(2 + \cos^2(dx + c)) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}} \sin(dx + c)}{3d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{9}{2}} \cos(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2), x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*(b/cos(d*x+c))^(3/2)*sin(d*x+c)/(1/cos(d*x+c))^(9/2)/cos(d*x+c)^3

maxima [A] time = 0.74, size = 45, normalized size = 0.62

$$\frac{\left(b \sin(3dx + 3c) + 9b \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)\right) \sqrt{b}}{12d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(9/2), x, algorithm="maxima")

[Out] 1/12*(b*sin(3*d*x + 3*c) + 9*b*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d

mupad [B] time = 0.44, size = 46, normalized size = 0.64

$$\frac{b (9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(9/2), x)

[Out] (b*(9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(9/2),x)
```

```
[Out] Timed out
```

$$3.151 \quad \int \frac{(b \sec(c+dx))^{3/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=101

$$\frac{3bx\sqrt{b \sec(c+dx)}}{8\sqrt{\sec(c+dx)}} + \frac{3b \sin(c+dx)\sqrt{b \sec(c+dx)}}{8d \sec^{\frac{3}{2}}(c+dx)} + \frac{b \sin(c+dx)\sqrt{b \sec(c+dx)}}{4d \sec^{\frac{7}{2}}(c+dx)}$$

[Out] $1/4*b*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(7/2)}+3/8*b*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(3/2)}+3/8*b*x*(b*\sec(d*x+c))^{(1/2)}/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{3bx\sqrt{b \sec(c+dx)}}{8\sqrt{\sec(c+dx)}} + \frac{3b \sin(c+dx)\sqrt{b \sec(c+dx)}}{8d \sec^{\frac{3}{2}}(c+dx)} + \frac{b \sin(c+dx)\sqrt{b \sec(c+dx)}}{4d \sec^{\frac{7}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(11/2), x]

[Out] $(3*b*x*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(8*\text{Sqrt}[\text{Sec}[c + d*x]]) + (b*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(4*d*\text{Sec}[c + d*x]^{(7/2)}) + (3*b*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(8*d*\text{Sec}[c + d*x]^{(3/2)})$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*SIN[c + d*x])^(n - 1)]/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{(b \sec(c + dx))^{3/2}}{\sec^{11/2}(c + dx)} dx &= \frac{(b\sqrt{b \sec(c + dx)}) \int \cos^4(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
&= \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{4d \sec^{7/2}(c + dx)} + \frac{(3b\sqrt{b \sec(c + dx)}) \int \cos^2(c + dx) dx}{4\sqrt{\sec(c + dx)}} \\
&= \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{4d \sec^{7/2}(c + dx)} + \frac{3b\sqrt{b \sec(c + dx)} \sin(c + dx)}{8d \sec^{3/2}(c + dx)} + \frac{(3b\sqrt{b \sec(c + dx)}) \int 1 dx}{8\sqrt{\sec(c + dx)}} \\
&= \frac{3bx\sqrt{b \sec(c + dx)}}{8\sqrt{\sec(c + dx)}} + \frac{b\sqrt{b \sec(c + dx)} \sin(c + dx)}{4d \sec^{7/2}(c + dx)} + \frac{3b\sqrt{b \sec(c + dx)} \sin(c + dx)}{8d \sec^{3/2}(c + dx)}
\end{aligned}$$

Mathematica [A] time = 0.16, size = 55, normalized size = 0.54

$$\frac{(12(c + dx) + 8 \sin(2(c + dx)) + \sin(4(c + dx)))(b \sec(c + dx))^{3/2}}{32d \sec^{3/2}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(3/2)/Sec[c + d*x]^(11/2),x]

[Out] ((b*Sec[c + d*x])^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])))/(32*d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.73, size = 207, normalized size = 2.05

$$\left[\frac{3\sqrt{-b}b \log\left(-2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b\right) + \frac{2(2b \cos(dx+c)^4 + 3b \cos(dx+c)^2)}{\sqrt{\cos(dx+c)}}}{16d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x, algorithm="fricas")

[Out] [1/16*(3*sqrt(-b)*b*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b) + 2*(2*b*cos(d*x + c)^4 + 3*b*cos(d

$x + c)^2 \sqrt{b/\cos(dx + c)} \sin(dx + c) / \sqrt{\cos(dx + c)}) / d, 1/8 * (3 * b^{3/2} * \arctan(\sqrt{b/\cos(dx + c)} \sin(dx + c) / (\sqrt{b} \sqrt{\cos(dx + c)}))) + (2 * b * \cos(dx + c)^4 + 3 * b * \cos(dx + c)^2 * \sqrt{b/\cos(dx + c)} \sin(dx + c) / \sqrt{\cos(dx + c)}) / d]$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{3}{2}}}{\sec(dx + c)^{\frac{11}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(3/2)/sec(d*x + c)^(11/2), x)

maple [A] time = 1.19, size = 74, normalized size = 0.73

$$\frac{\left(2 \left(\cos^3(dx + c)\right) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c\right) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}{8d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{11}{2}} \cos(dx + c)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x)

[Out] 1/8/d*(2*cos(d*x+c)^3*sin(d*x+c)+3*cos(d*x+c)*sin(d*x+c)+3*d*x+3*c)*(b/cos(d*x+c))^(3/2)/(1/cos(d*x+c))^(11/2)/cos(d*x+c)^4

maxima [A] time = 1.02, size = 53, normalized size = 0.52

$$\frac{\left(12(dx + c)b + b \sin(4dx + 4c) + 8b \sin\left(\frac{1}{2} \arctan\left(\frac{\sin(4dx + 4c)}{\cos(4dx + 4c)}\right)\right)\right) \sqrt{b}}{32d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(3/2)/sec(d*x+c)^(11/2),x, algorithm="maxima")

[Out] 1/32*(12*(d*x + c)*b + b*sin(4*d*x + 4*c) + 8*b*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))*sqrt(b)/d

mupad [B] time = 0.59, size = 53, normalized size = 0.52

$$\frac{b \sqrt{\frac{b}{\cos(c+dx)}} (8 \sin(2c + 2dx) + \sin(4c + 4dx) + 12dx)}{32d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b/cos(c + d*x))^(3/2)/(1/cos(c + d*x))^(11/2),x)
```

```
[Out] (b*(b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))  
/(32*d*(1/cos(c + d*x))^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(3/2)/sec(d*x+c)**(11/2),x)
```

```
[Out] Timed out
```

$$3.152 \quad \int \sec^{\frac{7}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx$$

Optimal. Leaf size=116

$$\frac{b^2 \sin^5(c + dx) \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{5d} + \frac{2b^2 \sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{b^2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d}$$

[Out] $2/3*b^2*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)^3*(b*\sec(d*x+c))^{(1/2)}/d+1/5*b^2*\sec(d*x+c)^{(9/2)}*\sin(d*x+c)^5*(b*\sec(d*x+c))^{(1/2)}/d+b^2*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.02, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{b^2 \sin^5(c + dx) \sec^{\frac{9}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{5d} + \frac{2b^2 \sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{b^2 \sin(c + dx) \sqrt{\sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(5/2), x]

[Out] $(b^2*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d + (2*b^2*\text{Sec}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d) + (b^2*\text{Sec}[c + d*x]^{(9/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^5)/(5*d)$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{7}{2}}(c+dx)(b \sec(c+dx))^{5/2} dx &= \frac{(b^2 \sqrt{b \sec(c+dx)}) \int \sec^6(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= -\frac{(b^2 \sqrt{b \sec(c+dx)}) \text{Subst}\left(\int (1+2x^2+x^4) dx, x, -\tan(c+dx)\right)}{d\sqrt{\sec(c+dx)}} \\
&= \frac{b^2 \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d} + \frac{2b^2 \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}}{3d}
\end{aligned}$$

Mathematica [A] time = 0.23, size = 57, normalized size = 0.49

$$\frac{\left(\frac{1}{5} \tan^5(c+dx) + \frac{2}{3} \tan^3(c+dx) + \tan(c+dx)\right) (b \sec(c+dx))^{5/2}}{d \sec^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(7/2)*(b*Sec[c + d*x])^(5/2), x]

[Out] ((b*Sec[c + d*x])^(5/2)*(Tan[c + d*x] + (2*Tan[c + d*x]^3)/3 + Tan[c + d*x]^5/5))/(d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.55, size = 63, normalized size = 0.54

$$\frac{(8b^2 \cos(dx+c)^4 + 4b^2 \cos(dx+c)^2 + 3b^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{15d \cos(dx+c)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] 1/15*(8*b^2*cos(d*x + c)^4 + 4*b^2*cos(d*x + c)^2 + 3*b^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(9/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{\frac{5}{2}} \sec(dx+c)^{\frac{7}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^(7/2), x)

maple [A] time = 0.94, size = 62, normalized size = 0.53

$$\frac{\left(8 \left(\cos^4(dx + c)\right) + 4 \left(\cos^2(dx + c)\right) + 3\right) \sin(dx + c) \cos(dx + c) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}}}{15d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2), x)

[Out] 1/15/d*(8*cos(d*x+c)^4+4*cos(d*x+c)^2+3)*sin(d*x+c)*cos(d*x+c)*(b/cos(d*x+c))^(5/2)*(1/cos(d*x+c))^(7/2)

maxima [B] time = 1.18, size = 705, normalized size = 6.08

$$15 \left(2 \left(5 \cos(8dx + 8c) + 10 \cos(6dx + 6c) + 10 \cos(4dx + 4c) + 5 \cos(2dx + 2c) + 1 \right) \cos(10dx + 10c) - \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)*(b*sec(d*x+c))^(5/2), x, algorithm="maxima")

[Out] -16/15*(5*(2*b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*cos(10*d*x + 10*c) + 25*(2*b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*cos(8*d*x + 8*c) + 50*(2*b^2*sin(4*d*x + 4*c) + b^2*sin(2*d*x + 2*c))*cos(6*d*x + 6*c) - (10*b^2*cos(4*d*x + 4*c) + 5*b^2*cos(2*d*x + 2*c) + b^2)*sin(10*d*x + 10*c) - 5*(10*b^2*cos(4*d*x + 4*c) + 5*b^2*cos(2*d*x + 2*c) + b^2)*sin(8*d*x + 8*c) - 10*(10*b^2*cos(4*d*x + 4*c) + 5*b^2*cos(2*d*x + 2*c) + b^2)*sin(6*d*x + 6*c))*sqrt(b)/((2*(5*cos(8*d*x + 8*c) + 10*cos(6*d*x + 6*c) + 10*cos(4*d*x + 4*c) + 5*cos(2*d*x + 2*c) + 1)*cos(10*d*x + 10*c) + cos(10*d*x + 10*c)^2 + 10*(10*cos(6*d*x + 6*c) + 10*cos(4*d*x + 4*c) + 5*cos(2*d*x + 2*c) + 1)*cos(8*d*x + 8*c) + 25*cos(8*d*x + 8*c)^2 + 20*(10*cos(4*d*x + 4*c) + 5*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + 100*cos(6*d*x + 6*c)^2 + 20*(5*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 100*cos(4*d*x + 4*c)^2 + 25*cos(2*d*x + 2*c)^2 + 10*(sin(8*d*x + 8*c) + 2*sin(6*d*x + 6*c) + 2*sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(10*d*x + 10*c) + sin(10*d*x + 10*c)^2 + 50*(2*sin(6*d*x + 6*c) + 2*sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(8*d*x + 8*c) + 25*sin(8*d*x + 8*c)^2 + 100*(2*sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + 100*sin(6*d*x + 6*c)^2 + 100*sin(4*d*x + 4*c)^2 + 100*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 25*sin(2*d*x + 2*c)^2 + 10*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 4.70, size = 205, normalized size = 1.77

$$\sqrt{\frac{b}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}} \left(\frac{b^2 \sqrt{\frac{1}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}} 8i}{15d} + \frac{b^2 \sqrt{\frac{1}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}} (-2 \sin(2c + 2dx)^2 + \sin(4c + 4dx) 1i + 1) 16i}{3d} + \frac{b^2 \sqrt{\frac{1}{2 \sin\left(\frac{c}{2} + \frac{dx}{2}\right)^2 - 1}}}{16 \left(\sin(c + dx)^2 - 1\right)^2} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(7/2), x)`

[Out] `-((-b/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*((b^2*(-1/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*8i)/(15*d) + (b^2*(-1/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*(sin(4*c + 4*d*x)*1i - 2*sin(2*c + 2*d*x)^2 + 1)*16i)/(3*d) + (b^2*(-1/(2*sin(c/2 + (d*x)/2)^2 - 1))^(1/2)*(sin(2*c + 2*d*x)*1i - 2*sin(c + d*x)^2 + 1)*8i)/(3*d))*(sin(5*c + 5*d*x)*1i + 2*sin((5*c)/2 + (5*d*x)/2)^2 - 1))/(16*(sin(c + d*x)^2 - 1)^2)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(7/2)*(b*sec(d*x+c))**(5/2), x)`

[Out] Timed out

$$3.153 \quad \int \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^{5/2} dx$$

Optimal. Leaf size=76

$$\frac{b^2 \sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{b^2 \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

[Out] $1/3*b^2*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)^3*(b*\sec(d*x+c))^{(1/2)}/d+b^2*\sin(d*x+c)*\sec(d*x+c)^{(1/2)}*(b*\sec(d*x+c))^{(1/2)}/d$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3767}

$$\frac{b^2 \sin^3(c + dx) \sec^{\frac{5}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{3d} + \frac{b^2 \sin(c + dx) \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)}}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(3/2)}*(b*\text{Sec}[c + d*x])^{(5/2)}, x]$

[Out] $(b^2*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/d + (b^2*\text{Sec}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d)$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1 + x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c + d*x]], x] /;$ FreeQ[{c, d}, x] && IGtQ[n/2, 0]

Rubi steps

$$\begin{aligned}
\int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2} dx &= \frac{(b^2 \sqrt{b \sec(c+dx)}) \int \sec^4(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= -\frac{(b^2 \sqrt{b \sec(c+dx)}) \operatorname{Subst}\left(\int (1+x^2) dx, x, -\tan(c+dx)\right)}{d \sqrt{\sec(c+dx)}} \\
&= \frac{b^2 \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)} \sin(c+dx)}{d} + \frac{b^2 \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}}{3d}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 45, normalized size = 0.59

$$\frac{\left(\frac{1}{3} \tan^3(c+dx) + \tan(c+dx)\right) (b \sec(c+dx))^{5/2}}{d \sec^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2), x]

[Out] ((b*Sec[c + d*x])^(5/2)*(Tan[c + d*x] + Tan[c + d*x]^3/3))/(d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.53, size = 48, normalized size = 0.63

$$\frac{(2b^2 \cos(dx+c)^2 + b^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3d \cos(dx+c)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] 1/3*(2*b^2*cos(d*x + c)^2 + b^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*cos(d*x + c)^(5/2))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{\frac{5}{2}} \sec(dx+c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*sec(d*x + c)^(3/2), x)

maple [A] time = 0.95, size = 52, normalized size = 0.68

$$\frac{(2(\cos^2(dx+c)) + 1) \cos(dx+c) \sin(dx+c) \left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}{3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2), x)

[Out] 1/3/d*(2*cos(d*x+c)^2+1)*cos(d*x+c)*sin(d*x+c)*(1/cos(d*x+c))^(3/2)*(b/cos(d*x+c))^(5/2)

maxima [B] time = 0.67, size = 311, normalized size = 4.09

$$\frac{4(3b^2 \cos(6dx + 6c))}{3(2(3 \cos(4dx + 4c) + 3 \cos(2dx + 2c) + 1) \cos(6dx + 6c) + \cos(6dx + 6c)^2 + 6(3 \cos(2dx + 2c) + 1))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^(5/2), x, algorithm="maxima")

[Out] -4/3*(3*b^2*cos(6*d*x + 6*c)*sin(2*d*x + 2*c) + 9*b^2*cos(4*d*x + 4*c)*sin(2*d*x + 2*c) - (3*b^2*cos(2*d*x + 2*c) + b^2)*sin(6*d*x + 6*c) - 3*(3*b^2*cos(2*d*x + 2*c) + b^2)*sin(4*d*x + 4*c))*sqrt(b)/((2*(3*cos(4*d*x + 4*c) + 3*cos(2*d*x + 2*c) + 1)*cos(6*d*x + 6*c) + cos(6*d*x + 6*c)^2 + 6*(3*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + 9*cos(4*d*x + 4*c)^2 + 9*cos(2*d*x + 2*c)^2 + 6*(sin(4*d*x + 4*c) + sin(2*d*x + 2*c))*sin(6*d*x + 6*c) + sin(6*d*x + 6*c)^2 + 9*sin(4*d*x + 4*c)^2 + 18*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 9*sin(2*d*x + 2*c)^2 + 6*cos(2*d*x + 2*c) + 1)*d)

mupad [B] time = 1.37, size = 129, normalized size = 1.70

$$\frac{2b^2 \cos(c + dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} (4 \sin(c + dx) + 5 \sin(3c + 3dx) + \sin(5c + 5dx) + \cos(c + dx))}{3d (10 \cos(c + dx) + 5 \cos(3c + 3dx) + \cos(5c + 5dx))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(3/2), x)

[Out] (2*b^2*cos(c + d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*(cos(c + d*x)*10i + 4*sin(c + d*x) + cos(3*c + 3*d*x)*5i + cos(5*c + 5*d*x)*1i + 5*sin(3*c + 3*d*x) + sin(5*c + 5*d*x)))/(3*d*(10*cos(c + d*x) + 5*cos(3*c + 3*d*x) + cos(5*c + 5*d*x)))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**(5/2), x)

[Out] Timed out

3.154 $\int \sqrt{\sec(c + dx)} (b \sec(c + dx))^{5/2} dx$

Optimal. Leaf size=78

$$\frac{b^2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{2d} + \frac{b^2 \sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

[Out] $1/2*b^2*\sec(d*x+c)^{(3/2)}*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d+1/2*b^2*\operatorname{arctanh}(\sin(d*x+c))*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{b^2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx) \sqrt{b \sec(c + dx)}}{2d} + \frac{b^2 \sqrt{b \sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{2d \sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2), x]

[Out] $(b^2*\operatorname{ArcTanh}[\sin[c + d*x]]*\operatorname{Sqrt}[b*\operatorname{Sec}[c + d*x]])/(2*d*\operatorname{Sqrt}[\operatorname{Sec}[c + d*x]]) + (b^2*\operatorname{Sec}[c + d*x]^{(3/2)}*\operatorname{Sqrt}[b*\operatorname{Sec}[c + d*x]]*\sin[c + d*x])/(2*d)$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] :> Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] :> -Simp[(b*Csc[c + d*x])*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sqrt{\sec(c+dx)} (b \sec(c+dx))^{5/2} dx &= \frac{(b^2 \sqrt{b \sec(c+dx)}) \int \sec^3(c+dx) dx}{\sqrt{\sec(c+dx)}} \\
&= \frac{b^2 \sec^2(c+dx) \sqrt{b \sec(c+dx)} \sin(c+dx)}{2d} + \frac{(b^2 \sqrt{b \sec(c+dx)}) \int \sec(c+dx) dx}{2\sqrt{\sec(c+dx)}} \\
&= \frac{b^2 \tanh^{-1}(\sin(c+dx)) \sqrt{b \sec(c+dx)}}{2d \sqrt{\sec(c+dx)}} + \frac{b^2 \sec^2(c+dx) \sqrt{b \sec(c+dx)}}{2d}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 50, normalized size = 0.64

$$\frac{(b \sec(c+dx))^{5/2} (\tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx))}{2d \sec^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2), x]

[Out] ((b*Sec[c + d*x])^(5/2)*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.60, size = 208, normalized size = 2.67

$$\left[\frac{b^{5/2} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right) + \frac{2b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4d \cos(dx+c)}, -\frac{\sqrt{-b} b^2 \arctan\left(\frac{\sqrt{-b} \sin(dx+c)}{\cos(dx+c)}\right)}{4d \cos(dx+c)} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [1/4*(b^(5/2)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c)), -1/2*(sqrt(-b)*b^2*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(d*cos(d*x + c))]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{5}{2}} \sqrt{\sec(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)*sqrt(sec(d*x + c)), x)

maple [A] time = 0.96, size = 114, normalized size = 1.46

$$\frac{\left((\cos^2(dx + c)) \ln\left(-\frac{-1 + \cos(dx + c) + \sin(dx + c)}{\sin(dx + c)}\right) - (\cos^2(dx + c)) \ln\left(-\frac{-\sin(dx + c) - 1 + \cos(dx + c)}{\sin(dx + c)}\right) - \sin(dx + c) \right) \cos(dx + c)}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*cos(d*x+c)*(1/cos(d*x+c))^(1/2)*(b/cos(d*x+c))^(5/2)

maxima [B] time = 0.94, size = 747, normalized size = 9.58

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)*(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] -1/4*(4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(b^2*sin(4*d*x + 4*c) + 2*b^2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))), cos(2*d*x + 2*c))) + 1) + (b^2*cos(4*d*x + 4*c)^2 + 4*b^2*cos(2*d*x + 2*c)^2 + b^2*sin(4*d*x + 4*c)^2 + 4*b^2*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b^2*sin(2*d*x + 2*c)^2 + 4*b^2*cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*cos(2*d*x + 2*c) + b^2)*cos(4*d*x + 4*c))*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(b^2*cos(4*d*x

+ 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 4*(b^2*cos(4*d*x + 4*c) + 2*b^2*cos(2*d*x + 2*c) + b^2)*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))*sqrt(b)/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*d)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{5/2} \sqrt{\frac{1}{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(1/2), x)

[Out] int((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(1/2)*(b*sec(d*x+c))**(5/2), x)

[Out] Timed out

$$3.155 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sqrt{\sec(c+dx)}} dx$$

Optimal. Leaf size=35

$$\frac{b^2 \sin(c+dx) \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}{d}$$

[Out] $b^2 \sin(dx+c) \sec(dx+c)^{(1/2)} (b \sec(dx+c))^{(1/2)} / d$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{b^2 \sin(c+dx) \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}{d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b \cdot \text{Sec}[c + d \cdot x])^{(5/2)} / \text{Sqrt}[\text{Sec}[c + d \cdot x]], x]$

[Out] $(b^2 \cdot \text{Sqrt}[\text{Sec}[c + d \cdot x]] \cdot \text{Sqrt}[b \cdot \text{Sec}[c + d \cdot x]] \cdot \text{Sin}[c + d \cdot x]) / d$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a \cdot x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.) \cdot ((a_.) \cdot (v_.))^{(m_.)} \cdot ((b_.) \cdot (v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)} \cdot b^{(n-1/2)} \cdot \text{Sqrt}[b \cdot v]) / \text{Sqrt}[a \cdot v], \text{Int}[u \cdot v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\amp; \text{!IntegerQ}[m] \&\amp; \text{IGtQ}[n+1/2, 0] \&\amp; \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.) \cdot (x_.)]^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d \cdot x]], x] /; \text{FreeQ}[\{c, d\}, x] \&\amp; \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c + dx))^{5/2}}{\sqrt{\sec(c + dx)}} dx &= \frac{(b^2 \sqrt{b \sec(c + dx)}) \int \sec^2(c + dx) dx}{\sqrt{\sec(c + dx)}} \\ &= -\frac{(b^2 \sqrt{b \sec(c + dx)}) \text{Subst}(\int 1 dx, x, -\tan(c + dx))}{d \sqrt{\sec(c + dx)}} \\ &= \frac{b^2 \sqrt{\sec(c + dx)} \sqrt{b \sec(c + dx)} \sin(c + dx)}{d} \end{aligned}$$

Mathematica [A] time = 0.03, size = 32, normalized size = 0.91

$$\frac{\sin(c + dx)(b \sec(c + dx))^{5/2}}{d \sec^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2)/Sqrt[Sec[c + d*x]],x]

[Out] ((b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(d*Sec[c + d*x]^(3/2))

fricas [A] time = 0.72, size = 33, normalized size = 0.94

$$\frac{b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="fricas")

[Out] b^2*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{5}{2}}}{\sqrt{\sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)/sqrt(sec(d*x + c)), x)

maple [A] time = 1.09, size = 39, normalized size = 1.11

$$\frac{\left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \cos(dx+c) \sin(dx+c)}{d \sqrt{\frac{1}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x)`

[Out] `1/d*(b/cos(d*x+c))^(5/2)*cos(d*x+c)*sin(d*x+c)/(1/cos(d*x+c))^(1/2)`

maxima [A] time = 1.20, size = 54, normalized size = 1.54

$$\frac{2 b^{\frac{5}{2}} \sin(2 dx + 2 c)}{(\cos(2 dx + 2 c)^2 + \sin(2 dx + 2 c)^2 + 2 \cos(2 dx + 2 c) + 1) d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(1/2),x, algorithm="maxima")`

[Out] `2*b^(5/2)*sin(2*d*x + 2*c)/((cos(2*d*x + 2*c))^2 + sin(2*d*x + 2*c)^2 + 2*cos(2*d*x + 2*c) + 1)*d`

mupad [B] time = 0.72, size = 66, normalized size = 1.89

$$\frac{b^2 \sqrt{\frac{b}{\cos(c+dx)}} (\cos(2c + 2dx) \operatorname{li} + \sin(2c + 2dx) + 1i)}{d (\cos(2c + 2dx) + 1) \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(1/2),x)`

[Out] `(b^2*(b/cos(c + d*x))^(1/2)*(cos(2*c + 2*d*x)*1i + sin(2*c + 2*d*x) + 1i))/(d*(cos(2*c + 2*d*x) + 1)*(1/cos(c + d*x))^(1/2))`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(1/2),x)`

[Out] Timed out

$$3.156 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=36

$$\frac{b^2 \sqrt{b \sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{d \sqrt{\sec(c+dx)}}$$

[Out] $b^2 \operatorname{arctanh}(\sin(dx+c)) \cdot (b \sec(dx+c))^{1/2} / d / \sec(dx+c)^{1/2}$

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{b^2 \sqrt{b \sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{d \sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b \cdot \text{Sec}[c + d \cdot x])^{5/2} / \text{Sec}[c + d \cdot x]^{3/2}, x]$

[Out] $(b^2 \cdot \text{ArcTanh}[\text{Sin}[c + d \cdot x]] \cdot \text{Sqrt}[b \cdot \text{Sec}[c + d \cdot x]]) / (d \cdot \text{Sqrt}[\text{Sec}[c + d \cdot x]])$

Rule 17

$\text{Int}[(u \cdot (a \cdot v))^m \cdot (b \cdot v)^n, x_Symbol] \rightarrow \text{Dist}[(a^{m+1/2} \cdot b^{n-1/2} \cdot \text{Sqrt}[b \cdot v]) / \text{Sqrt}[a \cdot v], \text{Int}[u \cdot v^{m+n}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

$\text{Int}[\text{csc}[(c \cdot x) + (d \cdot x)], x_Symbol] \rightarrow -\text{Simp}[\text{ArcTanh}[\text{Cos}[c + d \cdot x]] / d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx &= \frac{(b^2 \sqrt{b \sec(c+dx)}) \int \sec(c+dx) dx}{\sqrt{\sec(c+dx)}} \\ &= \frac{b^2 \tanh^{-1}(\sin(c+dx)) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 33, normalized size = 0.92

$$\frac{(b \sec(c + dx))^{5/2} \tanh^{-1}(\sin(c + dx))}{d \sec^{\frac{5}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*(b*Sec[c + d*x])^(5/2))/(d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.62, size = 114, normalized size = 3.17

$$\left[\frac{b^{\frac{5}{2}} \log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2d}, \frac{\sqrt{-b} b^2 \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2), x, algorithm="fricas")

[Out] [1/2*b^(5/2)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/d, -sqrt(-b)*b^2*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{5}{2}}}{\sec(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(3/2), x)

maple [A] time = 0.96, size = 52, normalized size = 1.44

$$\frac{2 \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \cos(dx+c) \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right)}{d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x)`

[Out] $-2/d*(b/\cos(d*x+c))^{5/2}*\cos(d*x+c)*\operatorname{arctanh}((-1+\cos(d*x+c))/\sin(d*x+c))/(1/\cos(d*x+c))^{3/2}$

maxima [B] time = 1.04, size = 72, normalized size = 2.00

$$\frac{(b^2 \log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2 \sin(dx+c) + 1) - b^2 \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2 \sin(dx+c) + 1))}{2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(3/2),x, algorithm="maxima")`

[Out] $1/2*(b^2*\log(\cos(d*x+c)^2 + \sin(d*x+c)^2 + 2*\sin(d*x+c) + 1) - b^2*\log(\cos(d*x+c)^2 + \sin(d*x+c)^2 - 2*\sin(d*x+c) + 1))*\sqrt{b}/d$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c+d*x))^(5/2)/(1/cos(c+d*x))^(3/2),x)`

[Out] `int((b/cos(c+d*x))^(5/2)/(1/cos(c+d*x))^(3/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(3/2),x)`

[Out] Timed out

$$3.157 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=27

$$\frac{b^2 x \sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

[Out] $b^2 x (b \sec(d x + c))^{1/2} / \sec(d x + c)^{1/2}$

Rubi [A] time = 0.00, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{b^2 x \sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(5/2),x]

[Out] (b^2*x*Sqrt[b*Sec[c + d*x]])/Sqrt[Sec[c + d*x]]

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx &= \frac{(b^2 \sqrt{b \sec(c+dx)}) \int 1 dx}{\sqrt{\sec(c+dx)}} \\ &= \frac{b^2 x \sqrt{b \sec(c+dx)}}{\sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 24, normalized size = 0.89

$$\frac{x(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(5/2), x]

[Out] (x*(b*Sec[c + d*x])^(5/2))/Sec[c + d*x]^(5/2)

fricas [A] time = 0.85, size = 101, normalized size = 3.74

$$\left[\frac{\sqrt{-b} b^2 \log\left(-2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b\right)}{2d}, \frac{b^{\frac{5}{2}} \arctan\left(\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}}\right)}{d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2), x, algorithm="fricas")

[Out] [1/2*sqrt(-b)*b^2*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/d, b^(5/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{5}{2}}}{\sec(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(5/2), x)

maple [A] time = 1.02, size = 32, normalized size = 1.19

$$\frac{\left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} (dx+c)}{d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2), x)

[Out] 1/d*(b/cos(d*x+c))^(5/2)/(1/cos(d*x+c))^(5/2)*(d*x+c)

maxima [A] time = 0.76, size = 26, normalized size = 0.96

$$\frac{2 b^{\frac{5}{2}} \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(5/2),x, algorithm="maxima")

[Out] 2*b^(5/2)*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/d

mupad [B] time = 0.12, size = 27, normalized size = 1.00

$$\frac{b^2 x \sqrt{\frac{b}{\cos(c+dx)}}}{\sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(5/2),x)

[Out] (b^2*x*(b/cos(c + d*x))^(1/2))/(1/cos(c + d*x))^(1/2)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(5/2),x)

[Out] Timed out

$$3.158 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=35

$$\frac{b^2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

[Out] $b^2 \sin(d*x+c) * (b * \sec(d*x+c))^{(1/2)} / d / \sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{b^2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b * \text{Sec}[c + d*x])^{(5/2)} / \text{Sec}[c + d*x]^{(7/2)}, x]$

[Out] $(b^2 * \text{Sqrt}[b * \text{Sec}[c + d*x]] * \text{Sin}[c + d*x]) / (d * \text{Sqrt}[\text{Sec}[c + d*x]])$

Rule 17

$\text{Int}[(u_.) * ((a_.) * (v_.))^{(m_.)} * ((b_.) * (v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)} * b^{(n-1/2)} * \text{Sqrt}[b*v]) / \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n+1/2, 0] && IntegerQ[m+n]

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.) * (x_.)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x] / d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx &= \frac{(b^2 \sqrt{b \sec(c+dx)}) \int \cos(c+dx) dx}{\sqrt{\sec(c+dx)}} \\ &= \frac{b^2 \sqrt{b \sec(c+dx)} \sin(c+dx)}{d \sqrt{\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 32, normalized size = 0.91

$$\frac{\sin(c + dx)(b \sec(c + dx))^{5/2}}{d \sec^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(7/2), x]

[Out] ((b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.75, size = 33, normalized size = 0.94

$$\frac{b^2 \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2), x, algorithm="fricas")

[Out] b^2*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/d

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{5}{2}}}{\sec(dx + c)^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(7/2), x)

maple [A] time = 1.13, size = 41, normalized size = 1.17

$$\frac{\left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \sin(dx+c)}{d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}} \cos(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2), x)

[Out] 1/d*(b/cos(d*x+c))^(5/2)*sin(d*x+c)/(1/cos(d*x+c))^(7/2)/cos(d*x+c)

maxima [A] time = 0.99, size = 13, normalized size = 0.37

$$\frac{b^{\frac{5}{2}} \sin(dx + c)}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(7/2),x, algorithm="maxima")

[Out] b^(5/2)*sin(d*x + c)/d

mupad [B] time = 0.33, size = 35, normalized size = 1.00

$$\frac{b^2 \sin(c + dx) \sqrt{\frac{b}{\cos(c+dx)}}}{d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(7/2),x)

[Out] (b^2*sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(7/2),x)

[Out] Timed out

$$3.159 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=69

$$\frac{b^2 x \sqrt{b \sec(c+dx)}}{2 \sqrt{\sec(c+dx)}} + \frac{b^2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{2d \sec^3(c+dx)}$$

[Out] $1/2*b^2*\sin(d*x+c)*(b*\sec(d*x+c))^{(1/2)}/d/\sec(d*x+c)^{(3/2)}+1/2*b^2*x*(b*\sec(d*x+c))^{(1/2)}/\sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{b^2 x \sqrt{b \sec(c+dx)}}{2 \sqrt{\sec(c+dx)}} + \frac{b^2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{2d \sec^3(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(9/2), x]

[Out] $(b^2*x*\text{Sqrt}[b*\text{Sec}[c + d*x]])/(2*\text{Sqrt}[\text{Sec}[c + d*x]]) + (b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(2*d*\text{Sec}[c + d*x]^{(3/2)})$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{(b \sec(c + dx))^{5/2}}{\sec^2(c + dx)} dx &= \frac{(b^2 \sqrt{b \sec(c + dx)}) \int \cos^2(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
&= \frac{b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{2d \sec^{\frac{3}{2}}(c + dx)} + \frac{(b^2 \sqrt{b \sec(c + dx)}) \int 1 dx}{2\sqrt{\sec(c + dx)}} \\
&= \frac{b^2 x \sqrt{b \sec(c + dx)}}{2\sqrt{\sec(c + dx)}} + \frac{b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{2d \sec^{\frac{3}{2}}(c + dx)}
\end{aligned}$$

Mathematica [A] time = 0.11, size = 45, normalized size = 0.65

$$\frac{(2(c + dx) + \sin(2(c + dx)))(b \sec(c + dx))^{5/2}}{4d \sec^{\frac{5}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(9/2), x]

[Out] ((b*Sec[c + d*x])^(5/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.83, size = 167, normalized size = 2.42

$$\left[\frac{2b^2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + \sqrt{-b} b^2 \log\left(-2\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)\right)}{4d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2), x, algorithm="fricas")

[Out] [1/4*(2*b^2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(-b)*b^2*log(-2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/d, 1/2*(b^2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + b^(5/2)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/d]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{\frac{5}{2}}}{\sec(dx + c)^{\frac{9}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(9/2), x)

maple [A] time = 0.98, size = 54, normalized size = 0.78

$$\frac{(\cos(dx + c) \sin(dx + c) + dx + c) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}{2d \cos(dx + c)^2 \left(\frac{1}{\cos(dx+c)}\right)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*(b/cos(d*x+c))^(5/2)/cos(d*x+c)^2/(1/cos(d*x+c))^(9/2)

maxima [A] time = 0.69, size = 32, normalized size = 0.46

$$\frac{(2(dx + c)b^2 + b^2 \sin(2dx + 2c))\sqrt{b}}{4d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(9/2),x, algorithm="maxima")

[Out] 1/4*(2*(d*x + c)*b^2 + b^2*sin(2*d*x + 2*c))*sqrt(b)/d

mupad [B] time = 0.37, size = 44, normalized size = 0.64

$$\frac{b^2 (\sin(2c + 2dx) + 2dx) \sqrt{\frac{b}{\cos(c+dx)}}}{4d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(9/2),x)

```
[Out] (b^2*(sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*d*(1/cos(c + d*x))^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(9/2), x)
```

```
[Out] Timed out
```

$$3.160 \quad \int \frac{(b \sec(c+dx))^{5/2}}{\sec^2(c+dx)} dx$$

Optimal. Leaf size=76

$$\frac{b^2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}} - \frac{b^2 \sin^3(c+dx) \sqrt{b \sec(c+dx)}}{3d \sqrt{\sec(c+dx)}}$$

[Out] $b^2 \sin(d*x+c) * (b * \sec(d*x+c))^{(1/2)} / d / \sec(d*x+c)^{(1/2)} - 1/3 * b^2 * \sin(d*x+c)^3 * (b * \sec(d*x+c))^{(1/2)} / d / \sec(d*x+c)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2633}

$$\frac{b^2 \sin(c+dx) \sqrt{b \sec(c+dx)}}{d \sqrt{\sec(c+dx)}} - \frac{b^2 \sin^3(c+dx) \sqrt{b \sec(c+dx)}}{3d \sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(11/2), x]

[Out] $(b^2 * \text{Sqrt}[b * \text{Sec}[c + d*x]] * \text{Sin}[c + d*x]) / (d * \text{Sqrt}[\text{Sec}[c + d*x]]) - (b^2 * \text{Sqrt}[b * \text{Sec}[c + d*x]] * \text{Sin}[c + d*x]^3) / (3 * d * \text{Sqrt}[\text{Sec}[c + d*x]])$

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2633

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := -Dist[d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned}
\int \frac{(b \sec(c + dx))^{5/2}}{\sec^{11/2}(c + dx)} dx &= \frac{(b^2 \sqrt{b \sec(c + dx)}) \int \cos^3(c + dx) dx}{\sqrt{\sec(c + dx)}} \\
&= -\frac{(b^2 \sqrt{b \sec(c + dx)}) \operatorname{Subst}\left(\int (1 - x^2) dx, x, -\sin(c + dx)\right)}{d \sqrt{\sec(c + dx)}} \\
&= \frac{b^2 \sqrt{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sec(c + dx)}} - \frac{b^2 \sqrt{b \sec(c + dx)} \sin^3(c + dx)}{3d \sqrt{\sec(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.17, size = 45, normalized size = 0.59

$$\frac{\sin(c + dx)(\cos(2(c + dx)) + 5)(b \sec(c + dx))^{5/2}}{6d \sec^2(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(5/2)/Sec[c + d*x]^(11/2), x]

[Out] ((5 + Cos[2*(c + d*x)])*(b*Sec[c + d*x])^(5/2)*Sin[c + d*x])/(6*d*Sec[c + d*x]^(5/2))

fricas [A] time = 0.74, size = 55, normalized size = 0.72

$$\frac{(b^2 \cos(dx + c)^3 + 2b^2 \cos(dx + c)) \sqrt{\frac{b}{\cos(dx + c)}} \sin(dx + c)}{3d \sqrt{\cos(dx + c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2), x, algorithm="fricas")

[Out] 1/3*(b^2*cos(d*x + c)^3 + 2*b^2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^{5/2}}{\sec^{11/2}(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(5/2)/sec(d*x + c)^(11/2), x)

maple [A] time = 0.97, size = 52, normalized size = 0.68

$$\frac{(2 + \cos^2(dx + c)) \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \sin(dx + c)}{3d \left(\frac{1}{\cos(dx+c)}\right)^{\frac{11}{2}} \cos(dx + c)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2), x)

[Out] 1/3/d*(2+cos(d*x+c)^2)*(b/cos(d*x+c))^(5/2)*sin(d*x+c)/(1/cos(d*x+c))^(11/2)/cos(d*x+c)^3

maxima [A] time = 0.88, size = 49, normalized size = 0.64

$$\frac{\left(b^2 \sin(3dx + 3c) + 9b^2 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)\right) \sqrt{b}}{12d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(5/2)/sec(d*x+c)^(11/2), x, algorithm="maxima")

[Out] 1/12*(b^2*sin(3*d*x + 3*c) + 9*b^2*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))*sqrt(b)/d

mupad [B] time = 0.50, size = 48, normalized size = 0.63

$$\frac{b^2 (9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(5/2)/(1/cos(c + d*x))^(11/2), x)

[Out] (b^2*(9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**(5/2)/sec(d*x+c)**(11/2),x)
```

```
[Out] Timed out
```

$$3.161 \quad \int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b} \sec(c+dx)} dx$$

Optimal. Leaf size=72

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{2d\sqrt{b} \sec(c+dx)} + \frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d\sqrt{b} \sec(c+dx)}$$

[Out] $1/2*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*\sec(d*x+c)^{(1/2)}/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{2d\sqrt{b} \sec(c+dx)} + \frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{2d\sqrt{b} \sec(c+dx)}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^(7/2)/Sqrt[b*Sec[c + d*x]], x]`

[Out] $(\operatorname{ArcTanh}[\sin[c + d*x]]*\sqrt{\sec[c + d*x]})/(2*d*\sqrt{b*\sec[c + d*x]}) + (\sec[c + d*x]^{(5/2)}*\sin[c + d*x])/(2*d*\sqrt{b*\sec[c + d*x]})$

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3770

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{\sec^{\frac{7}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec^3(c+dx) dx}{\sqrt{b \sec(c+dx)}} \\
&= \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2d\sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{2\sqrt{b \sec(c+dx)}} \\
&= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\sec(c+dx)}}{2d\sqrt{b \sec(c+dx)}} + \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2d\sqrt{b \sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 50, normalized size = 0.69

$$\frac{\sqrt{\sec(c+dx)} \left(\tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx) \right)}{2d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(7/2)/Sqrt[b*Sec[c + d*x]], x]

[Out] (Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.72, size = 205, normalized size = 2.85

$$\left[\frac{\sqrt{b} \cos(dx+c) \log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4bd \cos(dx+c)}, -\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\cos(dx+c)}}{\sin(dx+c)} \right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b*d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b*d*cos(d*x + c))]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{7}{2}}}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(7/2)/sqrt(b*sec(d*x + c)), x)

maple [A] time = 0.87, size = 114, normalized size = 1.58

$$\frac{\left((\cos^2(dx+c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx+c)) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx+c) \right) \left(\frac{1}{\cos(dx+c)} \right)}{2d \sqrt{\frac{b}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x)

[Out] $-1/2/d*(\cos(d*x+c)^2*\ln(-(-1+\cos(d*x+c)+\sin(d*x+c))/\sin(d*x+c))-\cos(d*x+c)^2*\ln(-(-\sin(d*x+c)-1+\cos(d*x+c))/\sin(d*x+c))-\sin(d*x+c))*(1/\cos(d*x+c))^(7/2)*\cos(d*x+c)/(b/\cos(d*x+c))^(1/2)$

maxima [B] time = 1.08, size = 661, normalized size = 9.18

$$4(\sin(4dx+4c)+2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan\left(\frac{\sin(2dx+2c)}{\cos(2dx+2c)}\right)\right)-4(\sin(4dx+4c)+$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] $-1/4*(4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - 4*(\sin(4*d*x + 4*c) + 2*\sin(2*d*x + 2*c))*\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) - (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d*x + 4*c)^2 + 4*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + 4*\sin(2*d*x + 2*c)^2 + 4*\cos(2*d*x + 2*c) + 1)*\log(\cos(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 + 2*\sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) + (2*(2*\cos(2*d*x + 2*c) + 1)*\cos(4*d*x + 4*c) + \cos(4*d*x + 4*c)^2 + 4*\cos(2*d*x + 2*c)^2 + \sin(4*d$

```
*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 +
4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/
2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) +
2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
)) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c)))))/((2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c)
+ cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4
*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) +
1)*sqrt(b)*d)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{7/2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(1/2), x)

[Out] int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(7/2)/(b*sec(d*x+c))**(1/2), x)

[Out] Timed out

$$3.162 \quad \int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b} \sec(c+dx)} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d\sqrt{b} \sec(c+dx)}$$

[Out] $\sec(d*x+c)^{(3/2)}*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d\sqrt{b} \sec(c+dx)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(5/2)}/\text{Sqrt}[b*\text{Sec}[c + d*x]], x]$

[Out] $(\text{Sec}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n+1/2, 0] \ \&\& \ \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\sec^{\frac{5}{2}}(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec^2(c+dx) dx}{\sqrt{b \sec(c+dx)}} \\ &= -\frac{\sqrt{\sec(c+dx)} \operatorname{Subst}\left(\int 1 dx, x, -\tan(c+dx)\right)}{d\sqrt{b \sec(c+dx)}} \\ &= \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{d\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 32, normalized size = 1.00

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(5/2)/Sqrt[b*Sec[c + d*x]], x]

[Out] (Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.44, size = 33, normalized size = 1.03

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{bd\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{5}{2}}}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(5/2)/sqrt(b*sec(d*x + c)), x)

maple [A] time = 0.82, size = 39, normalized size = 1.22

$$\frac{\left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}} \cos(dx+c) \sin(dx+c)}{d \sqrt{\frac{b}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x)`

[Out] `1/d*(1/cos(d*x+c))^(5/2)*cos(d*x+c)*sin(d*x+c)/(b/cos(d*x+c))^(1/2)`

maxima [B] time = 0.96, size = 59, normalized size = 1.84

$$\frac{2\sqrt{b}\sin(2dx+2c)}{(b\cos(2dx+2c)^2 + b\sin(2dx+2c)^2 + 2b\cos(2dx+2c) + b)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `2*sqrt(b)*sin(2*d*x + 2*c)/((b*cos(2*d*x + 2*c)^2 + b*sin(2*d*x + 2*c)^2 + 2*b*cos(2*d*x + 2*c) + b)*d)`

mupad [B] time = 0.27, size = 51, normalized size = 1.59

$$\frac{(\cos(dx) - \sin(dx) 1i) (\cos(c) - \sin(c) 1i) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} 1i}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(c + d*x))^(5/2)/(b/cos(c + d*x))^(1/2),x)`

[Out] `((cos(d*x) - sin(d*x)*1i)*(cos(c) - sin(c)*1i)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*1i)/(b*d)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(1/2),x)`

[Out] Timed out

$$3.163 \quad \int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=33

$$\frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{b \sec(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/d/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(3/2)/Sqrt[b*Sec[c + d*x]], x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(d*Sqrt[b*Sec[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{\sqrt{b \sec(c+dx)}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{\sqrt{b \sec(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 33, normalized size = 1.00

$$\frac{\sqrt{\sec(c + dx)} \tanh^{-1}(\sin(c + dx))}{d\sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)/Sqrt[b*Sec[c + d*x]], x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.64, size = 114, normalized size = 3.45

$$\left[\frac{\log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2\sqrt{b}d}, \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{bd} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/(sqrt(b)*d), -sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/(b*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{3}{2}}}{\sqrt{b \sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(3/2)/sqrt(b*sec(d*x + c)), x)

maple [A] time = 0.83, size = 52, normalized size = 1.58

$$\frac{2 \cos(dx+c) \left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}} \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right)}{d\sqrt{\frac{b}{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x)`

[Out] $-2/d*\cos(d*x+c)*(1/\cos(d*x+c))^{3/2}*\operatorname{arctanh}((-1+\cos(d*x+c))/\sin(d*x+c))/(b/\cos(d*x+c))^{1/2}$

maxima [B] time = 1.07, size = 65, normalized size = 1.97

$$\frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2\sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2\sin(dx+c) + 1)}{2\sqrt{b}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] $1/2*(\log(\cos(d*x+c)^2 + \sin(d*x+c)^2 + 2*\sin(d*x+c) + 1) - \log(\cos(d*x+c)^2 + \sin(d*x+c)^2 - 2*\sin(d*x+c) + 1))/(\operatorname{sqrt}(b)*d)$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}}{\sqrt{\frac{b}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(c+d*x))^(3/2)/(b/cos(c+d*x))^(1/2),x)`

[Out] `int((1/cos(c+d*x))^(3/2)/(b/cos(c+d*x))^(1/2),x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^{\frac{3}{2}}(c+dx)}{\sqrt{b}\sec(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(1/2),x)`

[Out] `Integral(sec(c+d*x)**(3/2)/sqrt(b*sec(c+d*x)),x)`

$$3.164 \quad \int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=24

$$\frac{x\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}}$$

[Out] $x*\sec(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[Sec[c + d*x]]/Sqrt[b*Sec[c + d*x]], x]`

[Out] `(x*Sqrt[Sec[c + d*x]])/Sqrt[b*Sec[c + d*x]]`

Rule 8

`Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} dx &= \frac{\sqrt{\sec(c+dx)} \int 1 dx}{\sqrt{b \sec(c+dx)}} \\ &= \frac{x\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.01, size = 24, normalized size = 1.00

$$\frac{x\sqrt{\sec(c+dx)}}{\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]/Sqrt[b*Sec[c + d*x]], x]

[Out] (x*Sqrt[Sec[c + d*x]])/Sqrt[b*Sec[c + d*x]]

fricas [A] time = 0.83, size = 101, normalized size = 4.21

$$\left[\frac{\sqrt{-b} \log\left(2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}}\cos(dx+c)^{\frac{3}{2}}\sin(dx+c)+2b\cos(dx+c)^2-b\right)}{2bd}, \frac{\arctan\left(\frac{\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{\sqrt{b}\sqrt{\cos(dx+c)}}\right)}{\sqrt{b}d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/(b*d), arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/(sqrt(b)*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\sec(dx+c)}}{\sqrt{b\sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(sec(d*x + c))/sqrt(b*sec(d*x + c)), x)

maple [A] time = 0.75, size = 32, normalized size = 1.33

$$\frac{\sqrt{\frac{1}{\cos(dx+c)}}(dx+c)}{d\sqrt{\frac{b}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2), x)

[Out] 1/d*(1/cos(d*x+c))^(1/2)/(b/cos(d*x+c))^(1/2)*(d*x+c)

maxima [A] time = 0.52, size = 26, normalized size = 1.08

$$\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{\sqrt{b}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(sqrt(b)*d)

mupad [B] time = 0.30, size = 27, normalized size = 1.12

$$\frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{b \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(1/2)/(b/cos(c + d*x))^(1/2),x)

[Out] (x*(b/cos(c + d*x))^(1/2))/(b*(1/cos(c + d*x))^(1/2))

sympy [A] time = 9.46, size = 5, normalized size = 0.21

$$\frac{x}{\sqrt{b}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(1/2),x)

[Out] x/sqrt(b)

$$3.165 \quad \int \frac{1}{\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=32

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 2637}

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]]), x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m-1/2)}*b^{(n+1/2)}*\text{Sqrt}[a*v])/ \text{Sqrt}[b*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x \&\& \text{IntegerQ}[m] \&\& \text{ILtQ}[n-1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos(c+dx) dx}{\sqrt{b \sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{d\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 32, normalized size = 1.00

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]]),x]

[Out] (Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.79, size = 33, normalized size = 1.03

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b*d)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \sec(dx+c)} \sqrt{\sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*sec(d*x + c))*sqrt(sec(d*x + c))), x)

maple [A] time = 0.94, size = 41, normalized size = 1.28

$$\frac{\sin(dx+c)}{d \sqrt{\frac{1}{\cos(dx+c)}} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x)

[Out] 1/d*sin(d*x+c)/(1/cos(d*x+c))^(1/2)/(b/cos(d*x+c))^(1/2)/cos(d*x+c)

maxima [A] time = 0.94, size = 13, normalized size = 0.41

$$\frac{\sin(dx+c)}{\sqrt{b}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(sqrt(b)*d)

mupad [B] time = 0.31, size = 35, normalized size = 1.09

$$\frac{\sin(c + dx) \sqrt{\frac{b}{\cos(c+dx)}}}{bd \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)),x)

[Out] (sin(c + d*x)*(b/cos(c + d*x))^(1/2))/(b*d*(1/cos(c + d*x))^(1/2))

sympy [A] time = 19.62, size = 36, normalized size = 1.12

$$\begin{cases} \frac{\tan(c+dx)}{\sqrt{b}d \sec(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{\sqrt{b \sec(c)} \sqrt{\sec(c)}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(1/2),x)

[Out] Piecewise((tan(c + d*x)/(sqrt(b)*d*sec(c + d*x)), Ne(d, 0)), (x/(sqrt(b*sec(c))*sqrt(sec(c))), True))

$$3.166 \quad \int \frac{1}{\sec^{\frac{3}{2}}(c+dx) \sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=63

$$\frac{x\sqrt{\sec(c+dx)}}{2\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2d\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)/d/\sec(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}+1/2*x*\sec(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 2635, 8}

$$\frac{x\sqrt{\sec(c+dx)}}{2\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2d\sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]]),x]

[Out] (x*Sqrt[Sec[c + d*x]])/(2*Sqrt[b*Sec[c + d*x]]) + Sin[c + d*x]/(2*d*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)]/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx = \frac{\sqrt{\sec(c+dx)} \int \cos^2(c+dx) dx}{\sqrt{b\sec(c+dx)}}$$

$$= \frac{\sin(c+dx)}{2d\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \int 1 dx}{2\sqrt{b\sec(c+dx)}}$$

$$= \frac{x\sqrt{\sec(c+dx)}}{2\sqrt{b\sec(c+dx)}} + \frac{\sin(c+dx)}{2d\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}}$$

Mathematica [A] time = 0.06, size = 45, normalized size = 0.71

$$\frac{(2(c+dx) + \sin(2(c+dx)))\sqrt{\sec(c+dx)}}{4d\sqrt{b\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sec[c + d*x]^(3/2)*Sqrt[b*Sec[c + d*x]]), x]

[Out] (Sqrt[Sec[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.61, size = 165, normalized size = 2.62

$$\frac{2\sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) - \sqrt{-b} \log\left(2\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)\right)}{4bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) - sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b*d), 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b\sec(dx+c)} \sec(dx+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/(sqrt(b*sec(d*x + c))*sec(d*x + c)^(3/2)), x)`

maple [A] time = 0.94, size = 54, normalized size = 0.86

$$\frac{\cos(dx+c)\sin(dx+c)+dx+c}{2d\cos(dx+c)^2\left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}}\sqrt{\frac{b}{\cos(dx+c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x)`

[Out] `1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)/cos(d*x+c)^2/(1/cos(d*x+c))^(3/2)/(b/cos(d*x+c))^(1/2)`

maxima [A] time = 0.95, size = 25, normalized size = 0.40

$$\frac{2dx+2c+\sin(2dx+2c)}{4\sqrt{bd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(1/2),x, algorithm="maxima")`

[Out] `1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(sqrt(b)*d)`

mapad [B] time = 0.43, size = 44, normalized size = 0.70

$$\frac{(\sin(2c+2dx)+2dx)\sqrt{\frac{b}{\cos(c+dx)}}}{4bd\sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((b/cos(c+d*x))^(1/2)*(1/cos(c+d*x))^(3/2)),x)`

[Out] `((sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*b*d*(1/cos(c + d*x))^(1/2))`

sympy [A] time = 30.23, size = 82, normalized size = 1.30

$$\begin{cases} \frac{x \tan^2(c+dx)}{2\sqrt{b} \sec^2(c+dx)} + \frac{x}{2\sqrt{b} \sec^2(c+dx)} + \frac{\tan(c+dx)}{2\sqrt{b} d \sec^2(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{\sqrt{b \sec(c)} \sec^{\frac{3}{2}}(c)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(1/2),x)`

[Out] `Piecewise((x*tan(c + d*x)**2/(2*sqrt(b)*sec(c + d*x)**2) + x/(2*sqrt(b)*sec(c + d*x)**2) + tan(c + d*x)/(2*sqrt(b)*d*sec(c + d*x)**2), Ne(d, 0)), (x/(sqrt(b*sec(c))*sec(c)**(3/2)), True))`

$$3.167 \quad \int \frac{1}{\sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}} dx$$

Optimal. Leaf size=70

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\sec(c+dx)}}{3d\sqrt{b \sec(c+dx)}}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/d/(b*\sec(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\sec(d*x+c)^{(1/2)}/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 2633}

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{d\sqrt{b \sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\sec(c+dx)}}{3d\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(\text{Sec}[c + d*x]^{(5/2)}*\text{Sqrt}[b*\text{Sec}[c + d*x]]), x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(d*\text{Sqrt}[b*\text{Sec}[c + d*x]]) - (\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m - 1/2)}*b^{(n + 1/2)}*\text{Sqrt}[a*v])/ \text{Sqrt}[b*v], \text{Int}[u*v^{(m + n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2633

$\text{Int}[\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{((n - 1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos^3(c+dx) dx}{\sqrt{b\sec(c+dx)}} \\ &= -\frac{\sqrt{\sec(c+dx)} \text{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{d\sqrt{b\sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{d\sqrt{b\sec(c+dx)}} - \frac{\sqrt{\sec(c+dx)} \sin^3(c+dx)}{3d\sqrt{b\sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 45, normalized size = 0.64

$$\frac{\sin(c+dx)(\cos(2(c+dx))+5)\sqrt{\sec(c+dx)}}{6d\sqrt{b\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]]),x]

[Out] ((5 + Cos[2*(c + d*x)])*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(6*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.67, size = 51, normalized size = 0.73

$$\frac{(\cos(dx+c)^3 + 2\cos(dx+c))\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{3bd\sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="fricas")

[Out] 1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b\sec(dx+c)} \sec(dx+c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*sec(d*x + c))*sec(d*x + c)^(5/2)), x)

maple [A] time = 1.05, size = 52, normalized size = 0.74

$$\frac{\sin(dx + c) \left(2 + \cos^2(dx + c)\right)}{3d \cos(dx + c)^3 \left(\frac{1}{\cos(dx + c)}\right)^{\frac{5}{2}} \sqrt{\frac{b}{\cos(dx + c)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2), x)

[Out] 1/3/d*sin(d*x+c)*(2+cos(d*x+c)^2)/cos(d*x+c)^3/(1/cos(d*x+c))^(5/2)/(b/cos(d*x+c))^(1/2)

maxima [A] time = 0.67, size = 42, normalized size = 0.60

$$\frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan\left(\frac{\sin(3dx + 3c)}{\cos(3dx + 3c)}\right)\right)}{12 \sqrt{bd}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2), x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(sqrt(b)*d)

mupad [B] time = 0.52, size = 48, normalized size = 0.69

$$\frac{(9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12bd \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(5/2)), x)

[Out] ((9*sin(c + d*x) + sin(3*c + 3*d*x))*b/cos(c + d*x))^(1/2)/(12*b*d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(1/2),x)
```

```
[Out] Timed out
```

$$3.168 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{2bd\sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{2bd\sqrt{b \sec(c+dx)}}$$

[Out] $1/2*\sec(d*x+c)^{(5/2)}*\sin(d*x+c)/b/d/(b*\sec(d*x+c))^{(1/2)}+1/2*\operatorname{arctanh}(\sin(d*x+c))*\sec(d*x+c)^{(1/2)}/b/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{2bd\sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{2bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(9/2)}/(b*\text{Sec}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{ArcTanh}[\text{Sin}[c + d*x]]*\text{Sqrt}[\text{Sec}[c + d*x]])/(2*b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + (\text{Sec}[c + d*x]^{(5/2)}*\text{Sin}[c + d*x])/(2*b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 17

$\text{Int}[(u_*)*((a_*)*(v_))^{(m_)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3768

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_)]*(b_*)^{(n_)}), x_Symbol] \rightarrow -\text{Simp}[(b*\text{Cos}[c + d*x])*(b*\text{Csc}[c + d*x])^{(n-1)})/(d*(n-1)), x] + \text{Dist}[(b^{2*(n-2)})/(n-1), \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3770

$\text{Int}[\text{csc}[(c_*) + (d_*)*(x_)], x_Symbol] \rightarrow -\text{Simp}[\text{ArcTanh}[\text{Cos}[c + d*x]]/d, x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec^3(c+dx) dx}{b\sqrt{b} \sec(c+dx)} \\
&= \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2bd\sqrt{b} \sec(c+dx)} + \frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{2b\sqrt{b} \sec(c+dx)} \\
&= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\sec(c+dx)}}{2bd\sqrt{b} \sec(c+dx)} + \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2bd\sqrt{b} \sec(c+dx)}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 50, normalized size = 0.64

$$\frac{\sec^{\frac{3}{2}}(c+dx) \left(\tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx) \right)}{2d(b \sec(c+dx))^{\frac{3}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(9/2)/(b*Sec[c + d*x])^(3/2), x]

[Out] (Sec[c + d*x]^(3/2)*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.95, size = 205, normalized size = 2.63

$$\left[\frac{\sqrt{b} \cos(dx+c) \log\left(-\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4b^2d \cos(dx+c)}, \sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\cos(dx+c)}}{\sqrt{\cos(dx+c)}} \right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^2*d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^2*d*cos(d*x + c))]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{9}{2}}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(9/2)/(b*sec(d*x + c))^(3/2), x)

maple [A] time = 0.82, size = 114, normalized size = 1.46

$$\frac{\left((\cos^2(dx+c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx+c)) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx+c) \right) \left(\frac{1}{\cos(dx+c)} \right)}{2d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c)*(1/cos(d*x+c))^(9/2)*cos(d*x+c)/(b/cos(d*x+c))^(3/2)

maxima [B] time = 0.77, size = 670, normalized size = 8.59

$$4(\sin(4dx+4c)+2\sin(2dx+2c))\cos\left(\frac{3}{2}\arctan(\sin(2dx+2c),\cos(2dx+2c))\right)-4(\sin(4dx+4c)+$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c)

```

+ 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d
*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 +
4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2
*c)))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 - 2*sin(1/
2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) - 4*(cos(4*d*x + 4*c) +
2*cos(2*d*x + 2*c) + 1)*sin(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)
)) + 4*(cos(4*d*x + 4*c) + 2*cos(2*d*x + 2*c) + 1)*sin(1/2*arctan2(sin(2*d*
x + 2*c), cos(2*d*x + 2*c)))))/((b*cos(4*d*x + 4*c)^2 + 4*b*cos(2*d*x + 2*c)
^2 + b*sin(4*d*x + 4*c)^2 + 4*b*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*b*sin
(2*d*x + 2*c)^2 + 2*(2*b*cos(2*d*x + 2*c) + b)*cos(4*d*x + 4*c) + 4*b*cos(2
*d*x + 2*c) + b)*sqrt(b)*d)

```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{9/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(9/2)/(b/cos(c + d*x))^(3/2), x)

[Out] int((1/cos(c + d*x))^(9/2)/(b/cos(c + d*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(9/2)/(b*sec(d*x+c))**(3/2), x)

[Out] Timed out

$$3.169 \quad \int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{bd\sqrt{b \sec(c+dx)}}$$

[Out] $\sec(d*x+c)^{(3/2)}*\sin(d*x+c)/b/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(7/2)}/(b*\text{Sec}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{Sec}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{!IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \&\& \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\sec^{\frac{7}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{3}{2}}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec^2(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \text{Subst}(\int 1 dx, x, -\tan(c+dx))}{bd\sqrt{b \sec(c+dx)}} \\ &= \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{bd\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{d(b \sec(c+dx))^{\frac{3}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(3/2), x]

[Out] (Sec[c + d*x]^(5/2)*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.68, size = 33, normalized size = 0.94

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{b^2 d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{7}{2}}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(7/2)/(b*sec(d*x + c))^(3/2), x)

maple [A] time = 0.82, size = 39, normalized size = 1.11

$$\frac{\left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}} \cos(dx+c) \sin(dx+c)}{d \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2),x)`

[Out] `1/d*(1/cos(d*x+c))^(7/2)*cos(d*x+c)*sin(d*x+c)/(b/cos(d*x+c))^(3/2)`

maxima [B] time = 0.85, size = 67, normalized size = 1.91

$$\frac{2\sqrt{b} \sin(2dx + 2c)}{(b^2 \cos(2dx + 2c)^2 + b^2 \sin(2dx + 2c)^2 + 2b^2 \cos(2dx + 2c) + b^2)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] `2*sqrt(b)*sin(2*d*x + 2*c)/((b^2*cos(2*d*x + 2*c)^2 + b^2*sin(2*d*x + 2*c)^2 + 2*b^2*cos(2*d*x + 2*c) + b^2)*d)`

mupad [B] time = 0.27, size = 51, normalized size = 1.46

$$\frac{(\cos(dx) - \sin(dx) 1i) (\cos(c) - \sin(c) 1i) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} 1i}{b^2 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(c + d*x))^(7/2)/(b/cos(c + d*x))^(3/2),x)`

[Out] `((cos(d*x) - sin(d*x)*1i)*(cos(c) - sin(c)*1i)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*1i)/(b^2*d)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(7/2)/(b*sec(d*x+c))**(3/2),x)`

[Out] Timed out

$$3.170 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=36

$$\frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{bd\sqrt{b \sec(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/b/d/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(b*d*Sqrt[b*Sec[c + d*x]])

Rule 17

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{3/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx))\sqrt{\sec(c+dx)}}{bd\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 33, normalized size = 0.92

$$\frac{\sec^2(c + dx) \tanh^{-1}(\sin(c + dx))}{d(b \sec(c + dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(3/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sec[c + d*x]^(3/2))/(d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.71, size = 114, normalized size = 3.17

$$\left[\frac{\log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2b^{\frac{3}{2}}d}, \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{b^2d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/(b^(3/2)*d), -sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/(b^2*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^{\frac{5}{2}}}{(b \sec(dx + c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(5/2)/(b*sec(d*x + c))^(3/2), x)

maple [A] time = 0.75, size = 52, normalized size = 1.44

$$\frac{2 \left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}} \cos(dx+c) \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right)}{d \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x)`

[Out] $-2/d*(1/\cos(d*x+c))^{5/2}*\cos(d*x+c)*\operatorname{arctanh}((-1+\cos(d*x+c))/\sin(d*x+c))/(b/\cos(d*x+c))^{3/2}$

maxima [B] time = 0.87, size = 65, normalized size = 1.81

$$\frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2\sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2\sin(dx+c) + 1)}{2b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")`

[Out] $1/2*(\log(\cos(d*x+c)^2 + \sin(d*x+c)^2 + 2*\sin(d*x+c) + 1) - \log(\cos(d*x+c)^2 + \sin(d*x+c)^2 - 2*\sin(d*x+c) + 1))/(b^{3/2}*d)$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{5/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(c+d*x))^(5/2)/(b/cos(c+d*x))^(3/2),x)`

[Out] `int((1/cos(c+d*x))^(5/2)/(b/cos(c+d*x))^(3/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(3/2),x)`

[Out] Timed out

$$3.171 \quad \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=27

$$\frac{x\sqrt{\sec(c+dx)}}{b\sqrt{b \sec(c+dx)}}$$

[Out] $x \sec(d*x+c)^{(1/2)}/b/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{\sec(c+dx)}}{b\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(3/2)}/(b*\text{Sec}[c + d*x])^{(3/2)}, x]$

[Out] $(x*\text{Sqrt}[\text{Sec}[c + d*x]])/(b*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] := \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)*((b_.)*(v_))^{(n_)}], x_Symbol] := \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{3/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int 1 dx}{b\sqrt{b \sec(c+dx)}} \\ &= \frac{x\sqrt{\sec(c+dx)}}{b\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 24, normalized size = 0.89

$$\frac{x \sec^3(c+dx)}{(b \sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)/(b*Sec[c + d*x])^(3/2), x]

[Out] (x*Sec[c + d*x]^(3/2))/(b*Sec[c + d*x])^(3/2)

fricas [A] time = 1.00, size = 101, normalized size = 3.74

$$\left[\frac{\sqrt{-b} \log \left(2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{2b^2d}, \frac{\arctan \left(\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}} \right)}{b^{\frac{3}{2}}d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/(b^2*d), arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/(b^(3/2)*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{3}{2}}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(3/2)/(b*sec(d*x + c))^(3/2), x)

maple [A] time = 0.76, size = 32, normalized size = 1.19

$$\frac{\left(\frac{1}{\cos(dx+c)} \right)^{\frac{3}{2}} (dx+c)}{d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2), x)

[Out] 1/d*(1/cos(d*x+c))^(3/2)/(b/cos(d*x+c))^(3/2)*(d*x+c)

maxima [A] time = 0.71, size = 26, normalized size = 0.96

$$\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(b^(3/2)*d)

mupad [B] time = 0.25, size = 27, normalized size = 1.00

$$\frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{b^2 \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(3/2)/(b/cos(c + d*x))^(3/2),x)

[Out] (x*(b/cos(c + d*x))^(1/2))/(b^2*(1/cos(c + d*x))^(1/2))

sympy [A] time = 26.54, size = 5, normalized size = 0.19

$$\frac{x}{b^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(3/2),x)

[Out] x/b**(3/2)

$$3.172 \quad \int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{bd\sqrt{b \sec(c+dx)}}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/b/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[\text{Sec}[c + d*x]]/(b*\text{Sec}[c + d*x])^{(3/2)}, x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x \&\& \text{IntegerQ}[m] \&\& \text{IGtQ}[n+1/2, 0] \&\& \text{IntegerQ}[m+n]$

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{3/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{bd\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{d(b \sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(3/2), x]

[Out] (Sec[c + d*x]^(3/2)*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.73, size = 33, normalized size = 0.94

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b^2 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2), x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^2*d)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\sec(dx+c)}}{(b \sec(dx+c))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2), x, algorithm="giac")

[Out] integrate(sqrt(sec(d*x + c))/(b*sec(d*x + c))^(3/2), x)

maple [A] time = 0.86, size = 41, normalized size = 1.17

$$\frac{\sin(dx+c) \sqrt{\frac{1}{\cos(dx+c)}}}{d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{3}{2}} \cos(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2), x)

[Out] 1/d*sin(d*x+c)*(1/cos(d*x+c))^(1/2)/(b/cos(d*x+c))^(3/2)/cos(d*x+c)

maxima [A] time = 1.04, size = 13, normalized size = 0.37

$$\frac{\sin(dx+c)}{b^{\frac{3}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(b^(3/2)*d)

mupad [B] time = 0.42, size = 39, normalized size = 1.11

$$\frac{\sin(2c + 2dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}}}{2b^2 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(1/2)/(b/cos(c + d*x))^(3/2),x)

[Out] (sin(2*c + 2*d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/(2*b^2*d)

sympy [A] time = 15.08, size = 36, normalized size = 1.03

$$\begin{cases} \frac{\tan(c+dx)}{b^2 d \sec(c+dx)} & \text{for } d \neq 0 \\ \frac{x\sqrt{\sec(c)}}{(b \sec(c))^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(3/2),x)

[Out] Piecewise((tan(c + d*x)/(b**(3/2)*d*sec(c + d*x)), Ne(d, 0)), (x*sqrt(sec(c))/(b*sec(c))**(3/2), True))

$$3.173 \quad \int \frac{1}{\sqrt{\sec(c+dx)} (b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=69

$$\frac{x\sqrt{\sec(c+dx)}}{2b\sqrt{b\sec(c+dx)}} + \frac{\sin(c+dx)}{2bd\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}}$$

[Out] 1/2*sin(d*x+c)/b/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+1/2*x*sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 2635, 8}

$$\frac{x\sqrt{\sec(c+dx)}}{2b\sqrt{b\sec(c+dx)}} + \frac{\sin(c+dx)}{2bd\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2)),x]

[Out] (x*Sqrt[Sec[c + d*x]])/(2*b*Sqrt[b*Sec[c + d*x]]) + Sin[c + d*x]/(2*b*d*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1))/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\sec(c+dx)}(b \sec(c+dx))^{3/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos^2(c+dx) dx}{b\sqrt{b \sec(c+dx)}} \\ &= \frac{\sin(c+dx)}{2bd\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \int 1 dx}{2b\sqrt{b \sec(c+dx)}} \\ &= \frac{x\sqrt{\sec(c+dx)}}{2b\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2bd\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 45, normalized size = 0.65

$$\frac{(2(c+dx) + \sin(2(c+dx))) \sec^{\frac{3}{2}}(c+dx)}{4d(b \sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(3/2)),x]

[Out] (Sec[c + d*x]^(3/2)*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.63, size = 165, normalized size = 2.39

$$\left[\frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) - \sqrt{-b} \log\left(2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)\right)}{4b^2d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) - sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^2*d), 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b^2*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx+c))^{\frac{3}{2}} \sqrt{\sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(d*x + c))^(3/2)*sqrt(sec(d*x + c))), x)

maple [A] time = 0.98, size = 54, normalized size = 0.78

$$\frac{\cos(dx+c)\sin(dx+c)+dx+c}{2d\cos(dx+c)^2\sqrt{\frac{1}{\cos(dx+c)}}\left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)/cos(d*x+c)^2/(1/cos(d*x+c))^(1/2)/(b/cos(d*x+c))^(3/2)

maxima [A] time = 0.88, size = 25, normalized size = 0.36

$$\frac{2dx+2c+\sin(2dx+2c)}{4b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(3/2),x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(3/2)*d)

mupad [B] time = 0.37, size = 44, normalized size = 0.64

$$\frac{(\sin(2c+2dx)+2dx)\sqrt{\frac{b}{\cos(c+dx)}}}{4b^2d\sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(1/2)),x)

[Out] ((sin(2*c + 2*d*x) + 2*d*x)*(b/cos(c + d*x))^(1/2))/(4*b^2*d*(1/cos(c + d*x))^(1/2))

sympy [A] time = 29.99, size = 82, normalized size = 1.19

$$\begin{cases} \frac{x \tan^2(c+dx)}{2b^{\frac{3}{2}} \sec^2(c+dx)} + \frac{x}{2b^{\frac{3}{2}} \sec^2(c+dx)} + \frac{\tan(c+dx)}{2b^{\frac{3}{2}} d \sec^2(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{(b \sec(c))^{\frac{3}{2}} \sqrt{\sec(c)}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(3/2),x)

[Out] Piecewise((x*tan(c + d*x)**2/(2*b**(3/2)*sec(c + d*x)**2) + x/(2*b**(3/2)*sec(c + d*x)**2) + tan(c + d*x)/(2*b**(3/2)*d*sec(c + d*x)**2), Ne(d, 0)), (x/((b*sec(c))**(3/2)*sqrt(sec(c))), True))

$$3.174 \quad \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=76

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{bd\sqrt{b \sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\sec(c+dx)}}{3bd\sqrt{b \sec(c+dx)}}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/b/d/(b*\sec(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\sec(d*x+c)^{(1/2)}/b/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 2633}

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{bd\sqrt{b \sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\sec(c+dx)}}{3bd\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(\text{Sec}[c + d*x]^{(3/2)}*(b*\text{Sec}[c + d*x])^{(3/2)}), x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]]) - (\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*b*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m-1/2)}*b^{(n+1/2)}*\text{Sqrt}[a*v])/ \text{Sqrt}[b*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2633

$\text{Int}[\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1-x^2)^{((n-1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{\frac{3}{2}}} dx = \frac{\sqrt{\sec(c+dx)} \int \cos^3(c+dx) dx}{b\sqrt{b \sec(c+dx)}}$$

$$= -\frac{\sqrt{\sec(c+dx)} \text{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{bd\sqrt{b \sec(c+dx)}}$$

$$= \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{bd\sqrt{b \sec(c+dx)}} - \frac{\sqrt{\sec(c+dx)} \sin^3(c+dx)}{3bd\sqrt{b \sec(c+dx)}}$$

Mathematica [A] time = 0.10, size = 45, normalized size = 0.59

$$\frac{\sin(c+dx)(\cos(2(c+dx))+5)\sec^{\frac{3}{2}}(c+dx)}{6d(b \sec(c+dx))^{\frac{3}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(3/2)),x]

[Out] ((5 + Cos[2*(c + d*x)])*Sec[c + d*x]^(3/2)*Sin[c + d*x])/(6*d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.71, size = 51, normalized size = 0.67

$$\frac{(\cos(dx+c)^3 + 2 \cos(dx+c)) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3 b^2 d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")

[Out] 1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^2*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx+c))^{\frac{3}{2}} \sec(dx+c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(d*x + c))^(3/2)*sec(d*x + c)^(3/2)), x)

maple [A] time = 1.02, size = 52, normalized size = 0.68

$$\frac{\sin(dx + c) (2 + \cos^2(dx + c))}{3d \cos(dx + c)^3 \left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2), x)

[Out] 1/3/d*sin(d*x+c)*(2+cos(d*x+c)^2)/cos(d*x+c)^3/(1/cos(d*x+c))^(3/2)/(b/cos(d*x+c))^(3/2)

maxima [A] time = 0.72, size = 42, normalized size = 0.55

$$\frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)}{12 b^{\frac{3}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(3/2), x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(3/2)*d)

mupad [B] time = 0.33, size = 48, normalized size = 0.63

$$\frac{(9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12 b^2 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(3/2)), x)

[Out] ((9*sin(c + d*x) + sin(3*c + 3*d*x))*(b/cos(c + d*x))^(1/2))/(12*b^2*d*(1/cos(c + d*x))^(1/2))

sympy [A] time = 121.59, size = 65, normalized size = 0.86

$$\begin{cases} \frac{2 \tan^3(c+dx)}{3 b^{\frac{3}{2}} d \sec^3(c+dx)} + \frac{\tan(c+dx)}{b^{\frac{3}{2}} d \sec^3(c+dx)} & \text{for } d \neq 0 \\ \frac{x}{(b \sec(c))^{\frac{3}{2}} \sec^{\frac{3}{2}}(c)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(3/2),x)
```

```
[Out] Piecewise((2*tan(c + d*x)**3/(3*b**(3/2)*d*sec(c + d*x)**3) + tan(c + d*x)/  
(b**(3/2)*d*sec(c + d*x)**3), Ne(d, 0)), (x/((b*sec(c))**(3/2)*sec(c)**(3/2  
)), True))
```

$$3.175 \quad \int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^{3/2}} dx$$

Optimal. Leaf size=107

$$\frac{3x\sqrt{\sec(c+dx)}}{8b\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{4bd \sec^{\frac{5}{2}}(c+dx)\sqrt{b \sec(c+dx)}} + \frac{3 \sin(c+dx)}{8bd\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/b/d/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2)+3/8*sin(d*x+c)/b/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+3/8*x*sec(d*x+c)^(1/2)/b/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 2635, 8}

$$\frac{3x\sqrt{\sec(c+dx)}}{8b\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{4bd \sec^{\frac{5}{2}}(c+dx)\sqrt{b \sec(c+dx)}} + \frac{3 \sin(c+dx)}{8bd\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2)), x]

[Out] (3*x*Sqrt[Sec[c + d*x]])/(8*b*Sqrt[b*Sec[c + d*x]]) + Sin[c + d*x]/(4*b*d*Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]]) + (3*Sin[c + d*x])/(8*b*d*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^{\frac{5}{2}}(c+dx)(b\sec(c+dx))^{3/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos^4(c+dx) dx}{b\sqrt{b\sec(c+dx)}} \\
&= \frac{\sin(c+dx)}{4bd \sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} + \frac{(3\sqrt{\sec(c+dx)}) \int \cos^2(c+dx) dx}{4b\sqrt{b\sec(c+dx)}} \\
&= \frac{\sin(c+dx)}{4bd \sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} + \frac{3\sin(c+dx)}{8bd\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} + \frac{(3\sqrt{\sec(c+dx)}) \int \cos^2(c+dx) dx}{8bd\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}} \\
&= \frac{3x\sqrt{\sec(c+dx)}}{8b\sqrt{b\sec(c+dx)}} + \frac{\sin(c+dx)}{4bd \sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} + \frac{3\sin(c+dx)}{8bd\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.12, size = 55, normalized size = 0.51

$$\frac{(12(c+dx) + 8\sin(2(c+dx)) + \sin(4(c+dx))) \sec^{\frac{3}{2}}(c+dx)}{32d(b\sec(c+dx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^(3/2)),x]

[Out] (Sec[c + d*x]^(3/2)*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])) / (32*d*(b*Sec[c + d*x])^(3/2))

fricas [A] time = 0.72, size = 208, normalized size = 1.94

$$\left[\frac{2(2\cos(dx+c)^4 + 3\cos(dx+c)^2)\sqrt{\frac{b}{\cos(dx+c)}}\sin(dx+c)}{\sqrt{\cos(dx+c)}} - 3\sqrt{-b}\log\left(2\sqrt{-b}\sqrt{\frac{b}{\cos(dx+c)}}\cos(dx+c)^{\frac{3}{2}}\sin(dx+c) + 2b\cos(dx+c)\right) \right] / 16b^2d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(3/2),x, algorithm="fricas")

[Out] [1/16*(2*(2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) - 3*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^2*d), 1/8*((2*

$\cos(dx + c)^4 + 3\cos(dx + c)^2 \sqrt{b/\cos(dx + c)} \sin(dx + c) / \sqrt{\cos(dx + c)} + 3\sqrt{b} \arctan(\sqrt{b/\cos(dx + c)} \sin(dx + c) / (\sqrt{b} \sqrt{\cos(dx + c)})) / (b^2 d)$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{3}{2}} \sec(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(dx+c)^(5/2)/(b*sec(dx+c))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(dx + c))^(3/2)*sec(dx + c)^(5/2)), x)

maple [A] time = 1.03, size = 74, normalized size = 0.69

$$\frac{2(\cos^3(dx + c)) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c}{8d \cos(dx + c)^4 \left(\frac{1}{\cos(dx+c)}\right)^{\frac{5}{2}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(dx+c)^(5/2)/(b*sec(dx+c))^(3/2),x)

[Out] 1/8/d*(2*cos(dx+c)^3*sin(dx+c)+3*cos(dx+c)*sin(dx+c)+3*dx+3*c)/cos(dx+c)^4/(1/cos(dx+c))^(5/2)/(b/cos(dx+c))^(3/2)

maxima [A] time = 0.97, size = 49, normalized size = 0.46

$$\frac{12dx + 12c + \sin(4dx + 4c) + 8 \sin\left(\frac{1}{2} \arctan\left(\frac{\sin(4dx + 4c)}{\cos(4dx + 4c)}\right)\right)}{32b^{\frac{3}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(dx+c)^(5/2)/(b*sec(dx+c))^(3/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(3/2)*d)

mupad [B] time = 0.62, size = 55, normalized size = 0.51

$$\frac{\sqrt{\frac{b}{\cos(c+dx)}} (8 \sin(2c + 2dx) + \sin(4c + 4dx) + 12dx)}{32b^2 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((b/cos(c + d*x))^(3/2)*(1/cos(c + d*x))^(5/2)),x)
```

```
[Out] ((b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(
32*b^2*d*(1/cos(c + d*x))^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(3/2),x)
```

```
[Out] Timed out
```

$$3.176 \quad \int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=78

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{2b^2 d \sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{2b^2 d \sqrt{b \sec(c+dx)}}$$

[Out] $1/2 * \sec(d*x+c)^{(5/2)} * \sin(d*x+c) / b^2/d / (b * \sec(d*x+c))^{(1/2)} + 1/2 * \operatorname{arctanh}(\sin(d*x+c)) * \sec(d*x+c)^{(1/2)} / b^2/d / (b * \sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3768, 3770}

$$\frac{\sin(c+dx) \sec^{\frac{5}{2}}(c+dx)}{2b^2 d \sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{2b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sec[c + d*x]^(11/2)/(b*Sec[c + d*x])^(5/2), x]`

[Out] `(ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(2*b^2*d*Sqrt[b*Sec[c + d*x]]) + (Sec[c + d*x]^(5/2)*Sin[c + d*x])/(2*b^2*d*Sqrt[b*Sec[c + d*x]])`

Rule 17

`Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]`

Rule 3768

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3770

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{\sec^{\frac{11}{2}}(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec^3(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
&= \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2b^2 d \sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{2b^2 \sqrt{b \sec(c+dx)}} \\
&= \frac{\tanh^{-1}(\sin(c+dx)) \sqrt{\sec(c+dx)}}{2b^2 d \sqrt{b \sec(c+dx)}} + \frac{\sec^{\frac{5}{2}}(c+dx) \sin(c+dx)}{2b^2 d \sqrt{b \sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 53, normalized size = 0.68

$$\frac{\sqrt{\sec(c+dx)} \left(\tanh^{-1}(\sin(c+dx)) + \tan(c+dx) \sec(c+dx) \right)}{2b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(11/2)/(b*Sec[c + d*x])^(5/2), x]

[Out] (Sqrt[Sec[c + d*x]]*(ArcTanh[Sin[c + d*x]] + Sec[c + d*x]*Tan[c + d*x]))/(2*b^2*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.94, size = 205, normalized size = 2.63

$$\left[\frac{\sqrt{b} \cos(dx+c) \log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2} \right) + \frac{2\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}}}{4b^3 d \cos(dx+c)}, -\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\cos(dx+c)}}{\sin(dx+c)} \right) \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [1/4*(sqrt(b)*cos(d*x + c)*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2) + 2*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^3*d*cos(d*x + c)), -1/2*(sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)*cos(d*x + c) - sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)))/(b^3*d*cos(d*x + c))]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{11}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(11/2)/(b*sec(d*x + c))^(5/2), x)

maple [A] time = 0.82, size = 114, normalized size = 1.46

$$\frac{\left((\cos^2(dx+c)) \ln\left(-\frac{-1+\cos(dx+c)+\sin(dx+c)}{\sin(dx+c)}\right) - (\cos^2(dx+c)) \ln\left(-\frac{-\sin(dx+c)-1+\cos(dx+c)}{\sin(dx+c)}\right) - \sin(dx+c) \right) \left(\frac{1}{\cos(dx+c)} \right)}{2d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x)

[Out] -1/2/d*(cos(d*x+c)^2*ln(-(-1+cos(d*x+c)+sin(d*x+c))/sin(d*x+c))-cos(d*x+c)^2*ln(-(-sin(d*x+c)-1+cos(d*x+c))/sin(d*x+c))-sin(d*x+c))*(1/cos(d*x+c))^(11/2)*cos(d*x+c)/(b/cos(d*x+c))^(5/2)

maxima [B] time = 0.84, size = 688, normalized size = 8.82

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(11/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] -1/4*(4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(3/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - 4*(sin(4*d*x + 4*c) + 2*sin(2*d*x + 2*c))*cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) - (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))^2 + sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c)))^2 + 2*sin(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))) + 1) + (2*(2*cos(2*d*x + 2*c) + 1)*cos(4*d*x + 4*c) + cos(4*d*x + 4*c)^2 + 4*cos(2*d*x + 2*c)^2 + sin(4*d*x + 4*c)^2 + 4*sin(4*d*x + 4*c)*sin(2*d*x + 2*c) + 4*sin(2*d*x + 2*c)^2 + 4*cos(2*d*x + 2*c) + 1)*log(cos(1/2*arctan2(sin(2*d*x + 2*c), cos(2*d*x + 2*c))))

$$\begin{aligned} & *c)))^2 + \sin(1/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c)))^2 - 2*\sin(1/ \\ & 2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c))) + 1) - 4*(\cos(4*d*x + 4*c) + \\ & 2*\cos(2*d*x + 2*c) + 1)*\sin(3/2*\arctan2(\sin(2*d*x + 2*c), \cos(2*d*x + 2*c) \\ &)) + 4*(\cos(4*d*x + 4*c) + 2*\cos(2*d*x + 2*c) + 1)*\sin(1/2*\arctan2(\sin(2*d* \\ & x + 2*c), \cos(2*d*x + 2*c)))))/((b^2*\cos(4*d*x + 4*c)^2 + 4*b^2*\cos(2*d*x + \\ & 2*c)^2 + b^2*\sin(4*d*x + 4*c)^2 + 4*b^2*\sin(4*d*x + 4*c)*\sin(2*d*x + 2*c) + \\ & 4*b^2*\sin(2*d*x + 2*c)^2 + 4*b^2*\cos(2*d*x + 2*c) + b^2 + 2*(2*b^2*\cos(2*d \\ & *x + 2*c) + b^2)*\cos(4*d*x + 4*c))*\sqrt{b}*d) \end{aligned}$$

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{11/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(11/2)/(b/cos(c + d*x))^(5/2), x)

[Out] int((1/cos(c + d*x))^(11/2)/(b/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(11/2)/(b*sec(d*x+c))**(5/2), x)

[Out] Timed out

$$3.177 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}$$

[Out] $\sec(d*x+c)^{(3/2)}*\sin(d*x+c)/b^2/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 3767, 8}

$$\frac{\sin(c+dx) \sec^{\frac{3}{2}}(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(9/2)}/(b*\text{Sec}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Sec}[c + d*x]^{(3/2)}*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] \rightarrow \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n+1/2, 0] \ \&\& \ \text{IntegerQ}[m+n]$

Rule 3767

$\text{Int}[\text{csc}[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{ExpandIntegrand}[(1+x^2)^{(n/2-1)}, x], x], x, \text{Cot}[c+d*x]], x] /; \text{FreeQ}[\{c, d\}, x] \ \&\& \ \text{IGtQ}[n/2, 0]$

Rubi steps

$$\begin{aligned} \int \frac{\sec^{\frac{9}{2}}(c+dx)}{(b \sec(c+dx))^{\frac{5}{2}}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec^2(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\ &= -\frac{\sqrt{\sec(c+dx)} \operatorname{Subst}\left(\int 1 dx, x, -\tan(c+dx)\right)}{b^2 d \sqrt{b \sec(c+dx)}} \\ &= \frac{\sec^{\frac{3}{2}}(c+dx) \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 32, normalized size = 0.91

$$\frac{\sin(c+dx) \sec^{\frac{7}{2}}(c+dx)}{d(b \sec(c+dx))^{\frac{5}{2}}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(9/2)/(b*Sec[c + d*x])^(5/2), x]

[Out] (Sec[c + d*x]^(7/2)*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(5/2))

fricas [A] time = 0.55, size = 33, normalized size = 0.94

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{b^3 d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{9}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(9/2)/(b*sec(d*x + c))^(5/2), x)

maple [A] time = 0.83, size = 39, normalized size = 1.11

$$\frac{\left(\frac{1}{\cos(dx+c)}\right)^{\frac{9}{2}} \cos(dx+c) \sin(dx+c)}{d \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2),x)`

[Out] `1/d*(1/cos(d*x+c))^(9/2)*cos(d*x+c)*sin(d*x+c)/(b/cos(d*x+c))^(5/2)`

maxima [B] time = 0.72, size = 67, normalized size = 1.91

$$\frac{2\sqrt{b} \sin(2dx + 2c)}{(b^3 \cos(2dx + 2c)^2 + b^3 \sin(2dx + 2c)^2 + 2b^3 \cos(2dx + 2c) + b^3)d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(9/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] `2*sqrt(b)*sin(2*d*x + 2*c)/((b^3*cos(2*d*x + 2*c)^2 + b^3*sin(2*d*x + 2*c)^2 + 2*b^3*cos(2*d*x + 2*c) + b^3)*d)`

mupad [B] time = 0.26, size = 51, normalized size = 1.46

$$\frac{(\cos(dx) - \sin(dx) 1i) (\cos(c) - \sin(c) 1i) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}} 1i}{b^3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(c + d*x))^(9/2)/(b/cos(c + d*x))^(5/2),x)`

[Out] `((cos(d*x) - sin(d*x)*1i)*(cos(c) - sin(c)*1i)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2)*1i)/(b^3*d)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(9/2)/(b*sec(d*x+c))**(5/2),x)`

[Out] Timed out

$$3.178 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=36

$$\frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{b^2 d \sqrt{b \sec(c+dx)}}$$

[Out] arctanh(sin(d*x+c))*sec(d*x+c)^(1/2)/b^2/d/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.01, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 3770}

$$\frac{\sqrt{\sec(c+dx)} \tanh^{-1}(\sin(c+dx))}{b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sqrt[Sec[c + d*x]])/(b^2*d*Sqrt[b*Sec[c + d*x]])

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \sec(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\ &= \frac{\tanh^{-1}(\sin(c+dx)) \sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 33, normalized size = 0.92

$$\frac{\sec^2(c + dx) \tanh^{-1}(\sin(c + dx))}{d(b \sec(c + dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(7/2)/(b*Sec[c + d*x])^(5/2), x]

[Out] (ArcTanh[Sin[c + d*x]]*Sec[c + d*x]^(5/2))/(d*(b*Sec[c + d*x])^(5/2))

fricas [A] time = 0.97, size = 114, normalized size = 3.17

$$\left[\frac{\log\left(\frac{b \cos(dx+c)^2 - 2\sqrt{b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c) - 2b}{\cos(dx+c)^2}\right)}{2b^{\frac{5}{2}}d}, \frac{\sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b}\right)}{b^3d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [1/2*log(-(b*cos(d*x + c))^2 - 2*sqrt(b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c) - 2*b)/cos(d*x + c)^2)/(b^(5/2)*d), -sqrt(-b)*arctan(sqrt(-b)*sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/b)/(b^3*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^{\frac{7}{2}}}{(b \sec(dx + c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(7/2)/(b*sec(d*x + c))^(5/2), x)

maple [A] time = 0.78, size = 52, normalized size = 1.44

$$\frac{2 \cos(dx + c) \left(\frac{1}{\cos(dx+c)}\right)^{\frac{7}{2}} \operatorname{arctanh}\left(\frac{-1+\cos(dx+c)}{\sin(dx+c)}\right)}{d \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2),x)`

[Out] $-2/d*\cos(d*x+c)*(1/\cos(d*x+c))^{7/2}*\operatorname{arctanh}((-1+\cos(d*x+c))/\sin(d*x+c))/(b/\cos(d*x+c))^{5/2}$

maxima [B] time = 0.90, size = 65, normalized size = 1.81

$$\frac{\log(\cos(dx+c)^2 + \sin(dx+c)^2 + 2\sin(dx+c) + 1) - \log(\cos(dx+c)^2 + \sin(dx+c)^2 - 2\sin(dx+c) + 1)}{2b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^(7/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")`

[Out] $1/2*(\log(\cos(d*x+c)^2 + \sin(d*x+c)^2 + 2*\sin(d*x+c) + 1) - \log(\cos(d*x+c)^2 + \sin(d*x+c)^2 - 2*\sin(d*x+c) + 1))/(b^{5/2}*d)$

mupad [F] time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^{7/2}}{\left(\frac{b}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1/cos(c+d*x))^(7/2)/(b/cos(c+d*x))^(5/2),x)`

[Out] `int((1/cos(c+d*x))^(7/2)/(b/cos(c+d*x))^(5/2),x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**(7/2)/(b*sec(d*x+c))^(5/2),x)`

[Out] Timed out

$$3.179 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=27

$$\frac{x\sqrt{\sec(c+dx)}}{b^2\sqrt{b \sec(c+dx)}}$$

[Out] $x*\sec(d*x+c)^{(1/2)}/b^2/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.00, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 8}

$$\frac{x\sqrt{\sec(c+dx)}}{b^2\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(5/2)}/(b*\text{Sec}[c + d*x])^{(5/2)}, x]$

[Out] $(x*\text{Sqrt}[\text{Sec}[c + d*x]])/(b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 8

$\text{Int}[a_, x_Symbol] := \text{Simp}[a*x, x] /; \text{FreeQ}[a, x]$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)*((b_.)*(v_))^{(n_)}], x_Symbol] := \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /; \text{FreeQ}[\{a, b, m\}, x] \&\amp; \text{IntegerQ}[m] \&\amp; \text{IGtQ}[n+1/2, 0] \&\amp; \text{IntegerQ}[m+n]$

Rubi steps

$$\begin{aligned} \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int 1 dx}{b^2\sqrt{b \sec(c+dx)}} \\ &= \frac{x\sqrt{\sec(c+dx)}}{b^2\sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.02, size = 24, normalized size = 0.89

$$\frac{x \sec^2(c+dx)}{(b \sec(c+dx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(5/2)/(b*Sec[c + d*x])^(5/2), x]

[Out] (x*Sec[c + d*x]^(5/2))/(b*Sec[c + d*x])^(5/2)

fricas [A] time = 0.95, size = 101, normalized size = 3.74

$$\left[\frac{\sqrt{-b} \log \left(2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)^2 - b \right)}{2b^3d}, \frac{\arctan \left(\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{b} \sqrt{\cos(dx+c)}} \right)}{b^{\frac{5}{2}}d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [-1/2*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b)/(b^3*d), arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c))))/(b^(5/2)*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{5}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(5/2)/(b*sec(d*x + c))^(5/2), x)

maple [A] time = 0.81, size = 32, normalized size = 1.19

$$\frac{\left(\frac{1}{\cos(dx+c)} \right)^{\frac{5}{2}} (dx+c)}{d \left(\frac{b}{\cos(dx+c)} \right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2), x)

[Out] 1/d*(1/cos(d*x+c))^(5/2)/(b/cos(d*x+c))^(5/2)*(d*x+c)

maxima [A] time = 0.82, size = 26, normalized size = 0.96

$$\frac{2 \arctan\left(\frac{\sin(dx+c)}{\cos(dx+c)+1}\right)}{b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 2*arctan(sin(d*x + c)/(cos(d*x + c) + 1))/(b^(5/2)*d)

mupad [B] time = 0.32, size = 27, normalized size = 1.00

$$\frac{x \sqrt{\frac{b}{\cos(c+dx)}}}{b^3 \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(5/2)/(b/cos(c + d*x))^(5/2),x)

[Out] (x*(b/cos(c + d*x))^(1/2))/(b^3*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(5/2)/(b*sec(d*x+c))**(5/2),x)

[Out] Timed out

$$3.180 \quad \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=35

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/b^2/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {17, 2637}

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^{(3/2)}/(b*\text{Sec}[c + d*x])^{(5/2)}, x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 17

$\text{Int}[(u_.)*((a_.)*(v_.))^{(m_.)}*((b_.)*(v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m+1/2)}*b^{(n-1/2)}*\text{Sqrt}[b*v])/ \text{Sqrt}[a*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ $\text{FreeQ}\{a, b, m\}, x \ \&\& \ !\text{IntegerQ}[m] \ \&\& \ \text{IGtQ}[n+1/2, 0] \ \&\& \ \text{IntegerQ}[m+n]$

Rule 2637

$\text{Int}[\sin[\text{Pi}/2 + (c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[\text{Sin}[c + d*x]/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \int \frac{\sec^3(c+dx)}{(b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.03, size = 35, normalized size = 1.00

$$\frac{\sin(c + dx)\sqrt{\sec(c + dx)}}{b^2 d \sqrt{b \sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)/(b*Sec[c + d*x])^(5/2), x]

[Out] (Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(b^2*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.76, size = 33, normalized size = 0.94

$$\frac{\sqrt{\frac{b}{\cos(dx+c)}} \sqrt{\cos(dx+c)} \sin(dx+c)}{b^3 d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] sqrt(b/cos(d*x + c))*sqrt(cos(d*x + c))*sin(d*x + c)/(b^3*d)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^{\frac{3}{2}}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^(3/2)/(b*sec(d*x + c))^(5/2), x)

maple [A] time = 0.84, size = 41, normalized size = 1.17

$$\frac{\sin(dx+c) \left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}}}{d \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}} \cos(dx+c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2), x)

[Out] 1/d*sin(d*x+c)*(1/cos(d*x+c))^(3/2)/(b/cos(d*x+c))^(5/2)/cos(d*x+c)

maxima [A] time = 0.90, size = 13, normalized size = 0.37

$$\frac{\sin(dx + c)}{b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] sin(d*x + c)/(b^(5/2)*d)

mupad [B] time = 0.39, size = 39, normalized size = 1.11

$$\frac{\sin(2c + 2dx) \sqrt{\frac{b}{\cos(c+dx)}} \sqrt{\frac{1}{\cos(c+dx)}}}{2b^3d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^(3/2)/(b/cos(c + d*x))^(5/2),x)

[Out] (sin(2*c + 2*d*x)*(b/cos(c + d*x))^(1/2)*(1/cos(c + d*x))^(1/2))/(2*b^3*d)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(5/2),x)

[Out] Timed out

$$3.181 \quad \int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=69

$$\frac{x\sqrt{\sec(c+dx)}}{2b^2\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2b^2d\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}}$$

[Out] $1/2*\sin(d*x+c)/b^2/d/\sec(d*x+c)^{(1/2)}/(b*\sec(d*x+c))^{(1/2)+1/2*x*\sec(d*x+c)^{(1/2)}/b^2/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.01, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {17, 2635, 8}

$$\frac{x\sqrt{\sec(c+dx)}}{2b^2\sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2b^2d\sqrt{\sec(c+dx)}\sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(5/2), x]

[Out] $(x*\text{Sqrt}[\text{Sec}[c + d*x]])/(2*b^2*\text{Sqrt}[b*\text{Sec}[c + d*x]]) + \text{Sin}[c + d*x]/(2*b^2*d*\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 17

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(a^(m + 1/2)*b^(n - 1/2)*Sqrt[b*v])/Sqrt[a*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && IGtQ[n + 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{\sec(c+dx)}}{(b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos^2(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
&= \frac{\sin(c+dx)}{2b^2 d \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}} + \frac{\sqrt{\sec(c+dx)} \int 1 dx}{2b^2 \sqrt{b \sec(c+dx)}} \\
&= \frac{x \sqrt{\sec(c+dx)}}{2b^2 \sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{2b^2 d \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 48, normalized size = 0.70

$$\frac{(2(c+dx) + \sin(2(c+dx)))\sqrt{\sec(c+dx)}}{4b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]/(b*Sec[c + d*x])^(5/2), x]

[Out] (Sqrt[Sec[c + d*x]]*(2*(c + d*x) + Sin[2*(c + d*x)]))/(4*b^2*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.93, size = 165, normalized size = 2.39

$$\left[\frac{2 \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) - \sqrt{-b} \log\left(2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c)\right)}{4b^3 d} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2), x, algorithm="fricas")

[Out] [1/4*(2*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) - sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^3*d), 1/2*(sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + sqrt(b)*arctan(sqrt(b/cos(d*x + c))*sin(d*x + c)/(sqrt(b)*sqrt(cos(d*x + c)))))/(b^3*d)]

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\sec(dx+c)}}{(b \sec(dx+c))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(sec(d*x + c))/(b*sec(d*x + c))^(5/2), x)

maple [A] time = 0.91, size = 54, normalized size = 0.78

$$\frac{(\cos(dx+c)\sin(dx+c)+dx+c)\sqrt{\frac{1}{\cos(dx+c)}}}{2d\cos(dx+c)^2\left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x)

[Out] 1/2/d*(cos(d*x+c)*sin(d*x+c)+d*x+c)*(1/cos(d*x+c))^(1/2)/cos(d*x+c)^2/(b/cos(d*x+c))^(5/2)

maxima [A] time = 0.96, size = 25, normalized size = 0.36

$$\frac{2dx+2c+\sin(2dx+2c)}{4b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="maxima")

[Out] 1/4*(2*d*x + 2*c + sin(2*d*x + 2*c))/(b^(5/2)*d)

mupad [B] time = 0.63, size = 64, normalized size = 0.93

$$\frac{\sqrt{\frac{b}{\cos(c+dx)}}(\sin(c+dx)+\sin(3c+3dx)+4dx\cos(c+dx))}{8b^3d\cos(c+dx)\sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c+d*x))^(1/2)/(b/cos(c+d*x))^(5/2),x)

[Out] ((b/cos(c+d*x))^(1/2)*(sin(c+d*x)+sin(3*c+3*d*x)+4*d*x*cos(c+d*x)))/(8*b^3*d*cos(c+d*x)*(1/cos(c+d*x))^(1/2))

sympy [A] time = 110.56, size = 82, normalized size = 1.19

$$\begin{cases} \frac{x \tan^2(c+dx)}{2b^2 \sec^2(c+dx)} + \frac{x}{2b^2 \sec^2(c+dx)} + \frac{\tan(c+dx)}{2b^2 d \sec^2(c+dx)} & \text{for } d \neq 0 \\ \frac{x \sqrt{\sec(c)}}{(b \sec(c))^{\frac{5}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(5/2), x)

[Out] Piecewise((x*tan(c + d*x)**2/(2*b**(5/2)*sec(c + d*x)**2) + x/(2*b**(5/2)*sec(c + d*x)**2) + tan(c + d*x)/(2*b**(5/2)*d*sec(c + d*x)**2), Ne(d, 0)), (x*sqrt(sec(c))/(b*sec(c))**(5/2), True))

$$3.182 \quad \int \frac{1}{\sqrt{\sec(c+dx)} (b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=76

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\sec(c+dx)}}{3b^2 d \sqrt{b \sec(c+dx)}}$$

[Out] $\sin(d*x+c)*\sec(d*x+c)^{(1/2)}/b^2/d/(b*\sec(d*x+c))^{(1/2)}-1/3*\sin(d*x+c)^3*\sec(d*x+c)^{(1/2)}/b^2/d/(b*\sec(d*x+c))^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {18, 2633}

$$\frac{\sin(c+dx)\sqrt{\sec(c+dx)}}{b^2 d \sqrt{b \sec(c+dx)}} - \frac{\sin^3(c+dx)\sqrt{\sec(c+dx)}}{3b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(\text{Sqrt}[\text{Sec}[c + d*x]]*(b*\text{Sec}[c + d*x])^{(5/2)}), x]$

[Out] $(\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x])/(b^2*d*\text{Sqrt}[b*\text{Sec}[c + d*x]]) - (\text{Sqrt}[\text{Sec}[c + d*x]]*\text{Sin}[c + d*x]^3)/(3*b^2*d*\text{Sqrt}[b*\text{Sec}[c + d*x]])$

Rule 18

$\text{Int}[(u_.)*((a_.)*(v_))^{(m_)}*((b_.)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a^{(m-1/2)}*b^{(n+1/2)}*\text{Sqrt}[a*v])/ \text{Sqrt}[b*v], \text{Int}[u*v^{(m+n)}, x], x] /;$ FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2633

$\text{Int}[\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow -\text{Dist}[d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1-x^2)^{((n-1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] /;$ FreeQ[{c, d}, x] && IGtQ[(n-1)/2, 0]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{\sec(c+dx)} (b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos^3(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\ &= -\frac{\sqrt{\sec(c+dx)} \operatorname{Subst}\left(\int (1-x^2) dx, x, -\sin(c+dx)\right)}{b^2 d \sqrt{b \sec(c+dx)}} \\ &= \frac{\sqrt{\sec(c+dx)} \sin(c+dx)}{b^2 d \sqrt{b \sec(c+dx)}} - \frac{\sqrt{\sec(c+dx)} \sin^3(c+dx)}{3b^2 d \sqrt{b \sec(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 48, normalized size = 0.63

$$\frac{\sin(c+dx)(\cos(2(c+dx))+5)\sqrt{\sec(c+dx)}}{6b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^(5/2)),x]

[Out] ((5 + Cos[2*(c + d*x)])*Sqrt[Sec[c + d*x]]*Sin[c + d*x])/(6*b^2*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.71, size = 51, normalized size = 0.67

$$\frac{(\cos(dx+c)^3 + 2 \cos(dx+c)) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{3b^3 d \sqrt{\cos(dx+c)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")

[Out] 1/3*(cos(d*x + c)^3 + 2*cos(d*x + c))*sqrt(b/cos(d*x + c))*sin(d*x + c)/(b^3*d*sqrt(cos(d*x + c)))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx+c))^{5/2} \sqrt{\sec(dx+c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(d*x + c))^(5/2)*sqrt(sec(d*x + c))), x)

maple [A] time = 1.13, size = 52, normalized size = 0.68

$$\frac{\sin(dx + c) (2 + \cos^2(dx + c))}{3d \cos(dx + c)^3 \sqrt{\frac{1}{\cos(dx+c)}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2), x)

[Out] 1/3/d*sin(d*x+c)*(2+cos(d*x+c)^2)/cos(d*x+c)^3/(1/cos(d*x+c))^(1/2)/(b/cos(d*x+c))^(5/2)

maxima [A] time = 1.04, size = 42, normalized size = 0.55

$$\frac{\sin(3dx + 3c) + 9 \sin\left(\frac{1}{3} \arctan(\sin(3dx + 3c), \cos(3dx + 3c))\right)}{12 b^{\frac{5}{2}} d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(5/2), x, algorithm="maxima")

[Out] 1/12*(sin(3*d*x + 3*c) + 9*sin(1/3*arctan2(sin(3*d*x + 3*c), cos(3*d*x + 3*c))))/(b^(5/2)*d)

mupad [B] time = 0.42, size = 48, normalized size = 0.63

$$\frac{(9 \sin(c + dx) + \sin(3c + 3dx)) \sqrt{\frac{b}{\cos(c+dx)}}}{12 b^3 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(1/2)), x)

[Out] ((9*sin(c + d*x) + sin(3*c + 3*d*x))*b/cos(c + d*x))^(1/2)/(12*b^3*d*(1/cos(c + d*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(d*x+c)**(1/2)/(b*sec(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

$$3.183 \quad \int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx$$

Optimal. Leaf size=107

$$\frac{3x\sqrt{\sec(c+dx)}}{8b^2\sqrt{b\sec(c+dx)}} + \frac{\sin(c+dx)}{4b^2d\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} + \frac{3\sin(c+dx)}{8b^2d\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}}$$

[Out] 1/4*sin(d*x+c)/b^2/d/sec(d*x+c)^(5/2)/(b*sec(d*x+c))^(1/2)+3/8*sin(d*x+c)/b^2/d/sec(d*x+c)^(1/2)/(b*sec(d*x+c))^(1/2)+3/8*x*sec(d*x+c)^(1/2)/b^2/(b*sec(d*x+c))^(1/2)

Rubi [A] time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {18, 2635, 8}

$$\frac{3x\sqrt{\sec(c+dx)}}{8b^2\sqrt{b\sec(c+dx)}} + \frac{\sin(c+dx)}{4b^2d\sec^{\frac{5}{2}}(c+dx)\sqrt{b\sec(c+dx)}} + \frac{3\sin(c+dx)}{8b^2d\sqrt{\sec(c+dx)}\sqrt{b\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2)), x]

[Out] (3*x*Sqrt[Sec[c + d*x]])/(8*b^2*Sqrt[b*Sec[c + d*x]]) + Sin[c + d*x]/(4*b^2*d*Sec[c + d*x]^(5/2)*Sqrt[b*Sec[c + d*x]]) + (3*Sin[c + d*x])/(8*b^2*d*Sqrt[Sec[c + d*x]]*Sqrt[b*Sec[c + d*x]])

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 18

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(a^(m - 1/2)*b^(n + 1/2)*Sqrt[a*v])/Sqrt[b*v], Int[u*v^(m + n), x], x] /; FreeQ[{a, b, m}, x] && !IntegerQ[m] && ILtQ[n - 1/2, 0] && IntegerQ[m + n]

Rule 2635

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := -Simp[(b*Cos[c + d*x])*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[(b^2*(n - 1))/n, Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^{5/2}} dx &= \frac{\sqrt{\sec(c+dx)} \int \cos^4(c+dx) dx}{b^2 \sqrt{b \sec(c+dx)}} \\
&= \frac{\sin(c+dx)}{4b^2 d \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}} + \frac{(3\sqrt{\sec(c+dx)}) \int \cos^2(c+dx) dx}{4b^2 \sqrt{b \sec(c+dx)}} \\
&= \frac{\sin(c+dx)}{4b^2 d \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \sqrt{\sec(c+dx)} \sqrt{b \sec(c+dx)}} + \dots \\
&= \frac{3x \sqrt{\sec(c+dx)}}{8b^2 \sqrt{b \sec(c+dx)}} + \frac{\sin(c+dx)}{4b^2 d \sec^{\frac{5}{2}}(c+dx) \sqrt{b \sec(c+dx)}} + \frac{3 \sin(c+dx)}{8b^2 d \sqrt{\sec(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 58, normalized size = 0.54

$$\frac{(12(c+dx) + 8 \sin(2(c+dx)) + \sin(4(c+dx))) \sqrt{\sec(c+dx)}}{32b^2 d \sqrt{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^(5/2)),x]

[Out] (Sqrt[Sec[c + d*x]]*(12*(c + d*x) + 8*Sin[2*(c + d*x)] + Sin[4*(c + d*x)])) / (32*b^2*d*Sqrt[b*Sec[c + d*x]])

fricas [A] time = 0.68, size = 208, normalized size = 1.94

$$\left[\frac{2(2 \cos(dx+c)^4 + 3 \cos(dx+c)^2) \sqrt{\frac{b}{\cos(dx+c)}} \sin(dx+c)}{\sqrt{\cos(dx+c)}} - 3 \sqrt{-b} \log \left(2 \sqrt{-b} \sqrt{\frac{b}{\cos(dx+c)}} \cos(dx+c)^{\frac{3}{2}} \sin(dx+c) + 2b \cos(dx+c) \right) \right] / 16b^3d$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(d*x+c)^(3/2)/(b*sec(d*x+c))^(5/2),x, algorithm="fricas")

[Out] [1/16*(2*(2*cos(d*x + c)^4 + 3*cos(d*x + c)^2)*sqrt(b/cos(d*x + c))*sin(d*x + c)/sqrt(cos(d*x + c)) - 3*sqrt(-b)*log(2*sqrt(-b)*sqrt(b/cos(d*x + c))*cos(d*x + c)^(3/2)*sin(d*x + c) + 2*b*cos(d*x + c)^2 - b))/(b^3*d), 1/8*((2*

$\cos(dx + c)^4 + 3\cos(dx + c)^2 \sqrt{b/\cos(dx + c)} \sin(dx + c) / \sqrt{\cos(dx + c)} + 3\sqrt{b} \arctan(\sqrt{b/\cos(dx + c)} \sin(dx + c) / (\sqrt{b} \sqrt{\cos(dx + c)})) / (b^3 d)$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{5}{2}} \sec(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(dx+c)^(3/2)/(b*sec(dx+c))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(dx + c))^(5/2)*sec(dx + c)^(3/2)), x)

maple [A] time = 1.12, size = 74, normalized size = 0.69

$$\frac{2(\cos^3(dx + c)) \sin(dx + c) + 3 \cos(dx + c) \sin(dx + c) + 3dx + 3c}{8d \cos(dx + c)^4 \left(\frac{1}{\cos(dx+c)}\right)^{\frac{3}{2}} \left(\frac{b}{\cos(dx+c)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sec(dx+c)^(3/2)/(b*sec(dx+c))^(5/2),x)

[Out] 1/8/d*(2*cos(dx+c)^3*sin(dx+c)+3*cos(dx+c)*sin(dx+c)+3*d*x+3*c)/cos(dx+c)^4/(1/cos(dx+c))^(3/2)/(b/cos(dx+c))^(5/2)

maxima [A] time = 1.14, size = 49, normalized size = 0.46

$$\frac{12dx + 12c + \sin(4dx + 4c) + 8 \sin\left(\frac{1}{2} \arctan\left(\frac{\sin(4dx + 4c)}{\cos(4dx + 4c)}\right)\right)}{32b^{\frac{5}{2}}d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sec(dx+c)^(3/2)/(b*sec(dx+c))^(5/2),x, algorithm="maxima")

[Out] 1/32*(12*d*x + 12*c + sin(4*d*x + 4*c) + 8*sin(1/2*arctan2(sin(4*d*x + 4*c), cos(4*d*x + 4*c))))/(b^(5/2)*d)

mupad [B] time = 0.46, size = 55, normalized size = 0.51

$$\frac{\sqrt{\frac{b}{\cos(c+dx)}} (8 \sin(2c + 2dx) + \sin(4c + 4dx) + 12dx)}{32b^3 d \sqrt{\frac{1}{\cos(c+dx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((b/cos(c + d*x))^(5/2)*(1/cos(c + d*x))^(3/2)),x)
```

```
[Out] ((b/cos(c + d*x))^(1/2)*(8*sin(2*c + 2*d*x) + sin(4*c + 4*d*x) + 12*d*x))/(
32*b^3*d*(1/cos(c + d*x))^(1/2))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sec(d*x+c)**(3/2)/(b*sec(d*x+c))**(5/2),x)
```

```
[Out] Timed out
```

3.184 $\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{4/3} {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4bd\sqrt{\sin^2(c + dx)}}$$

[Out] 3/4*hypergeom([-2/3, 1/2], [1/3], cos(d*x+c)^2)*(b*sec(d*x+c))^(4/3)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{4/3} {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4bd\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sin[c + d*x])/(4*b*d*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \sec^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx &= \frac{\int (b \sec(c + dx))^{7/3} dx}{b^2} \\
&= \frac{\left(\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{7/3}} dx}{b^2} \\
&= \frac{3 {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sin(c + dx)}{4bd \sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.07, size = 60, normalized size = 1.03

$$\frac{3 \sqrt{-\tan^2(c + dx)} \csc(c + dx) (b \sec(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \sec^2(c + dx)\right)}{7bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Csc[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(7*b*d)

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^2, x)

maple [F] time = 0.52, size = 0, normalized size = 0.00

$$\int (\sec^2(dx + c)) (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3), x)`

[Out] `int(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(1/3), x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}}{\cos(c+dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(1/3)/cos(c + d*x)^2, x)`

[Out] `int((b/cos(c + d*x))^(1/3)/cos(c + d*x)^2, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sec(c + dx)} \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(1/3), x)`

[Out] `Integral((b*sec(c + d*x))**(1/3)*sec(c + d*x)**2, x)`

3.185 $\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

Optimal. Leaf size=53

$$\frac{3 \sin(c + dx) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

[Out] 3*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c + dx) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]*(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \sec(c + dx) \sqrt[3]{b \sec(c + dx)} dx &= \frac{\int (b \sec(c + dx))^{4/3} dx}{b} \\
&= \frac{\left(\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{4/3}} dx}{b} \\
&= \frac{3 {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{d \sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 60, normalized size = 1.13

$$\frac{3 \sqrt{-\tan^2(c + dx)} \cot(c + dx) (b \sec(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sec^2(c + dx)\right)}{4bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(1/3),x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(4*b*d)

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \sec(dx + c)\right)^{\frac{1}{3}} \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c), x)

maple [F] time = 0.43, size = 0, normalized size = 0.00

$$\int \sec(dx + c) (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

[Out] `int(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)*(b*sec(d*x+c))^(1/3),x,algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}}{\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(1/3)/cos(c + d*x),x)`

[Out] `int((b/cos(c + d*x))^(1/3)/cos(c + d*x), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sec(c + dx)} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)*(b*sec(d*x+c))**(1/3),x)`

[Out] `Integral((b*sec(c + d*x))**(1/3)*sec(c + d*x), x)`

3.186 $\int \sqrt[3]{b \sec(c + dx)} dx$

Optimal. Leaf size=56

$$\frac{3b \sin(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{2/3}}$$

[Out] $-3/2*b*hypergeom([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(2/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.02, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$\frac{3b \sin(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*b*Hypergeometric2F1[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(2*d*(b*\text{Sec}[c + d*x])^{(2/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

$\text{Int}[(b* \sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n - 1)}*((\text{Sin}[c + d*x]/b)^{(n - 1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[n]$

Rubi steps

$$\int \sqrt[3]{b \sec(c + dx)} dx = \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\sqrt[3]{\frac{\cos(c + dx)}{b}}} dx$$

$$= -\frac{3 \cos(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{2d \sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.04, size = 55, normalized size = 0.98

$$\frac{3 \sqrt{-\tan^2(c + dx)} \cot(c + dx) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \sec^2(c + dx)\right)}{d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(1/3),x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sqrt[-Tan[c + d*x]^2])/d

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{\frac{1}{3}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/3),x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.54, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^(1/3),x)`

[Out] `int((b*sec(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(1/3),x)`

[Out] `int((b/cos(c + d*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sec(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**(1/3),x)`

[Out] `Integral((b*sec(c + d*x))**(1/3), x)`

3.187 $\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{5/3}}$$

[Out] $-3/5*b^2*\text{hypergeom}([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{5/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]*(b*\text{Sec}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*b^2*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(5*d*(b*\text{Sec}[c + d*x])^{(5/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

$\text{Int}[(u_.)*(v_.)^{(m_.)}*((b_.)*(v_))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \cos(c + dx) \sqrt[3]{b \sec(c + dx)} dx &= b \int \frac{1}{(b \sec(c + dx))^{2/3}} dx \\
&= \left(b \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{2/3} dx \\
&= -\frac{3 \cos^2(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{5d \sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 58, normalized size = 1.00

$$-\frac{3b \sqrt{-\tan^2(c + dx)} \cot(c + dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sec^2(c + dx)\right)}{2d(b \sec(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(1/3), x]

[Out] (-3*b*Cot[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(2*d*(b*Sec[c + d*x])^(2/3))

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{1/3} \cos(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{1/3} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c), x)

maple [F] time = 0.58, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

[Out] `int(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))^(1/3),x,algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx) \left(\frac{b}{\cos(c + dx)} \right)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)*(b/cos(c + d*x))^(1/3),x)`

[Out] `int(cos(c + d*x)*(b/cos(c + d*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sec(c + dx)} \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))**(1/3),x)`

[Out] `Integral((b*sec(c + d*x))**(1/3)*cos(c + d*x), x)`

3.188 $\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

Optimal. Leaf size=58

$$\frac{3b^3 \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right)}{8d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{8/3}}$$

[Out] $-3/8*b^3*hypergeom([1/2, 4/3], [7/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{\wedge}(8/3)/(\sin(d*x+c)^2)^{\wedge}(1/2)$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3b^3 \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right)}{8d \sqrt{\sin^2(c + dx)} (b \sec(c + dx))^{8/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^2*(b*\text{Sec}[c + d*x])^{\wedge}(1/3), x]$

[Out] $(-3*b^3*Hypergeometric2F1[1/2, 4/3, 7/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(8*d*(b*\text{Sec}[c + d*x])^{\wedge}(8/3)*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

$\text{Int}[(u_*)*(v_*)^{\wedge}(m_*)*((b_*)*(v_*))^{\wedge}(n_), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{\wedge}(m+n), x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{\wedge}(n_), x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{\wedge}(n+1)*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_*)]*(b_*))^{\wedge}(n_), x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{\wedge}(n-1)*((\text{Sin}[c + d*x]/b)^{\wedge}(n-1)*\text{Int}[1/(\text{Sin}[c + d*x]/b)^{\wedge}n, x]), x] /; \text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \cos^2(c + dx) \sqrt[3]{b \sec(c + dx)} dx &= b^2 \int \frac{1}{(b \sec(c + dx))^{5/3}} dx \\
&= \left(b^2 \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{5/3} dx \\
&= -\frac{3 \cos^3(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{8d \sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.15, size = 59, normalized size = 1.02

$$\frac{3 \sin(2(c + dx)) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(-\frac{5}{6}, \frac{1}{2}; \frac{1}{6}; \sec^2(c + dx)\right)}{10d \sqrt{-\tan^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Hypergeometric2F1[-5/6, 1/2, 1/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sin[2*(c + d*x)])/(10*d*Sqrt[-Tan[c + d*x]^2])

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c)^2, x)

maple [F] time = 1.06, size = 0, normalized size = 0.00

$$\int (\cos^2(dx + c)) (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3), x)`

[Out] `int(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(1/3), x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(1/3)*cos(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 \left(\frac{b}{\cos(c + dx)} \right)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3), x)`

[Out] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sec(c + dx)} \cos^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(1/3), x)`

[Out] `Integral((b*sec(c + d*x))**(1/3)*cos(c + d*x)**2, x)`

$$3.189 \quad \int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{7/3} {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c + dx)\right)}{7bd\sqrt{\sin^2(c + dx)}}$$

[Out] 3/7*hypergeom([-7/6, 1/2], [-1/6], cos(d*x+c)^2)*(b*sec(d*x+c))^(7/3)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{7/3} {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c + dx)\right)}{7bd\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^(4/3), x]

[Out] (3*Hypergeometric2F1[-7/6, 1/2, -1/6, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(7/3))*Sin[c + d*x]/(7*b*d*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \sec^2(c + dx)(b \sec(c + dx))^{4/3} dx &= \frac{\int (b \sec(c + dx))^{10/3} dx}{b^2} \\
&= \frac{\left(\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{10/3}} dx}{b^2} \\
&= \frac{3b {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \cos^2(c + dx)\right) \sec(c + dx) \sqrt[3]{b \sec(c + dx)} \tan(c + dx)}{7d \sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 60, normalized size = 1.03

$$\frac{3\sqrt{-\tan^2(c + dx)} \csc(c + dx)(b \sec(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{3}, \frac{8}{3}; \sec^2(c + dx)\right)}{10bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^(4/3), x]

[Out] (3*Csc[c + d*x]*Hypergeometric2F1[1/2, 5/3, 8/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(7/3)*Sqrt[-Tan[c + d*x]^2])/(10*b*d)

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \sec(dx + c)\right)^{\frac{1}{3}} b \sec(dx + c)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^2, x)

maple [F] time = 0.44, size = 0, normalized size = 0.00

$$\int (\sec^2(dx + c)) (b \sec(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3),x)`

[Out] `int(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}}{\cos(c+dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(4/3)/cos(c + d*x)^2,x)`

[Out] `int((b/cos(c + d*x))^(4/3)/cos(c + d*x)^2, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{4}{3}} \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**(4/3),x)`

[Out] `Integral((b*sec(c + d*x))**(4/3)*sec(c + d*x)**2, x)`

3.190 $\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx$

Optimal. Leaf size=55

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{4/3} {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)}}$$

[Out] 3/4*hypergeom([-2/3, 1/2], [1/3], cos(d*x+c)^2)*(b*sec(d*x+c))^(4/3)*sin(d*x+c)/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{4/3} {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right)}{4d\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]*(b*Sec[c + d*x])^(4/3), x]

[Out] (3*Hypergeometric2F1[-2/3, 1/2, 1/3, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sin[c + d*x])/(4*d*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \sec(c + dx)(b \sec(c + dx))^{4/3} dx &= \frac{\int (b \sec(c + dx))^{7/3} dx}{b} \\
&= \frac{\left(\sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{7/3}} dx}{b} \\
&= \frac{3 {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \cos^2(c + dx)\right) (b \sec(c + dx))^{4/3} \sin(c + dx)}{4d\sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 60, normalized size = 1.09

$$\frac{3\sqrt{-\tan^2(c + dx)} \cot(c + dx)(b \sec(c + dx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \sec^2(c + dx)\right)}{7bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^(4/3), x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(7/3)*Sqrt[-Tan[c + d*x]^2])/(7*b*d)

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{1/3} b \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{4/3} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c), x)

maple [F] time = 0.43, size = 0, normalized size = 0.00

$$\int \sec(dx + c) (b \sec(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x)`

[Out] `int(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^{4/3}}{\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(4/3)/cos(c + d*x), x)`

[Out] `int((b/cos(c + d*x))^(4/3)/cos(c + d*x), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{4}{3}} \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)*(b*sec(d*x+c))**(4/3), x)`

[Out] `Integral((b*sec(c + d*x))**(4/3)*sec(c + d*x), x)`

3.191 $\int (b \sec(c + dx))^{4/3} dx$

Optimal. Leaf size=54

$$\frac{3b \sin(c + dx) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

[Out] 3*b*hypergeom([-1/6, 1/2], [5/6], cos(d*x+c)^2)*(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$\frac{3b \sin(c + dx) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right)}{d \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(4/3), x]

[Out] (3*b*Hypergeometric2F1[-1/6, 1/2, 5/6, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*Sqrt[Sin[c + d*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int (b \sec(c + dx))^{4/3} dx = \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{4/3}} dx$$

$$= \frac{3b {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{d\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.01, size = 57, normalized size = 1.06

$$\frac{3\sqrt{-\tan^2(c + dx)} \cot(c + dx) (b \sec(c + dx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sec^2(c + dx)\right)}{4d}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(4/3), x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(4*d)

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{1/3} b \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{4/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3), x)

maple [F] time = 0.44, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{4/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^(4/3),x)`

[Out] `int((b*sec(d*x+c))^(4/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(4/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(4/3),x)`

[Out] `int((b/cos(c + d*x))^(4/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**(4/3),x)`

[Out] `Integral((b*sec(c + d*x))**(4/3), x)`

3.192 $\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}(b \sec(c + dx))^{2/3}}$$

[Out] $-3/2*b^2*hypergeom([1/3, 1/2], [4/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(2/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3b^2 \sin(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right)}{2d\sqrt{\sin^2(c + dx)}(b \sec(c + dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]*(b*Sec[c + d*x])^(4/3), x]

[Out] $(-3*b^2*Hypergeometric2F1[1/3, 1/2, 4/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(2*d*(b*\text{Sec}[c + d*x])^{(2/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \cos(c + dx)(b \sec(c + dx))^{4/3} dx &= b \int \sqrt[3]{b \sec(c + dx)} dx \\
&= \left(b \sqrt[3]{\frac{\cos(c + dx)}{b}} \sqrt[3]{b \sec(c + dx)} \right) \int \frac{1}{\sqrt[3]{\frac{\cos(c + dx)}{b}}} dx \\
&= \frac{3b \cos(c + dx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \cos^2(c + dx)\right) \sqrt[3]{b \sec(c + dx)} \sin(c + dx)}{2d \sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.01, size = 56, normalized size = 0.97

$$\frac{3b \sqrt{-\tan^2(c + dx)} \cot(c + dx) \sqrt[3]{b \sec(c + dx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \sec^2(c + dx)\right)}{d}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^(4/3), x]

[Out] (3*b*Cot[c + d*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(1/3)*Sqrt[-Tan[c + d*x]^2])/d

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \sec(dx + c)\right)^{\frac{1}{3}} b \cos(dx + c) \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*b*cos(d*x + c)*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c), x)

maple [F] time = 0.64, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \sec(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)*(b*sec(d*x+c))^(4/3), x)`

[Out] `int(cos(d*x+c)*(b*sec(d*x+c))^(4/3), x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))^(4/3), x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx) \left(\frac{b}{\cos(c + dx)} \right)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)*(b/cos(c + d*x))^(4/3), x)`

[Out] `int(cos(c + d*x)*(b/cos(c + d*x))^(4/3), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))**(4/3), x)`

[Out] Timed out

3.193 $\int \cos^2(c + dx)(b \sec(c + dx))^{4/3} dx$

Optimal. Leaf size=58

$$-\frac{3b^3 \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5d\sqrt{\sin^2(c + dx)}(b \sec(c + dx))^{5/3}}$$

[Out] $-3/5*b^3*\text{hypergeom}([1/2, 5/6], [11/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(5/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$-\frac{3b^3 \sin(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c + dx)\right)}{5d\sqrt{\sin^2(c + dx)}(b \sec(c + dx))^{5/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^2*(b*\text{Sec}[c + d*x])^{(4/3)}, x]$

[Out] $(-3*b^3*\text{Hypergeometric2F1}[1/2, 5/6, 11/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(5*d*(b*\text{Sec}[c + d*x])^{(5/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ $\text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_)]*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /;$ $\text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \cos^2(c+dx)(b \sec(c+dx))^{4/3} dx &= b^2 \int \frac{1}{(b \sec(c+dx))^{2/3}} dx \\
&= \left(b^2 \sqrt[3]{\frac{\cos(c+dx)}{b}} \sqrt[3]{b \sec(c+dx)} \right) \int \left(\frac{\cos(c+dx)}{b} \right)^{2/3} dx \\
&= -\frac{3b \cos^2(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \cos^2(c+dx)\right) \sqrt[3]{b \sec(c+dx)} \sin(c+dx)}{5d\sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.01, size = 60, normalized size = 1.03

$$-\frac{3b^2 \sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sec^2(c+dx)\right)}{2d(b \sec(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^(4/3), x]

[Out] (-3*b^2*Cot[c + d*x]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(2*d*(b*Sec[c + d*x])^(2/3))

fricas [F] time = 0.65, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx+c))^{1/3} b \cos(dx+c)^2 \sec(dx+c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*b*cos(d*x + c)^2*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{4/3} \cos(dx+c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c)^2, x)

maple [F] time = 1.15, size = 0, normalized size = 0.00

$$\int (\cos^2(dx + c)) (b \sec(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3),x)`

[Out] `int(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(4/3)*cos(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(c + dx)^2 \left(\frac{b}{\cos(c + dx)} \right)^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3),x)`

[Out] `int(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**(4/3),x)`

[Out] Timed out

$$3.194 \quad \int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \sin(c+dx)(b \sec(c+dx))^{2/3} {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx)\right)}{2bd\sqrt{\sin^2(c+dx)}}$$

[Out] 3/2*hypergeom([-1/3, 1/2], [2/3], cos(d*x+c)^2)*(b*sec(d*x+c))^(2/3)*sin(d*x+c)/b/d/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c+dx)(b \sec(c+dx))^{2/3} {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx)\right)}{2bd\sqrt{\sin^2(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Hypergeometric2F1[-1/3, 1/2, 2/3, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sin[c + d*x])/(2*b*d*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/((Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx &= \frac{\int (b \sec(c+dx))^{5/3} dx}{b^2} \\
&= \frac{\left(\left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b} \right)^{5/3}} dx}{b^2} \\
&= \frac{3 {}_2F_1 \left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \cos^2(c+dx) \right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{2bd \sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 60, normalized size = 1.03

$$\frac{3\sqrt{-\tan^2(c+dx)} \csc(c+dx) (b \sec(c+dx))^{2/3} {}_2F_1 \left(\frac{1}{2}, \frac{5}{6}, \frac{11}{6}; \sec^2(c+dx) \right)}{5bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Csc[c + d*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(5*b*d)

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \sec(dx+c))^{\frac{2}{3}} \sec(dx+c)}{b}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.37, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)

[Out] int(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3)),x)

[Out] int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(1/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(1/3),x)

[Out] Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(1/3), x)

$$3.195 \quad \int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

Optimal. Leaf size=53

$$\frac{3 \sin(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{d \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

[Out] -3*hypergeom([1/6, 1/2], [7/6], cos(d*x+c)^2)*sin(d*x+c)/d/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{d \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Sec[c + d*x])^(1/3), x]

[Out] (-3*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sin[c + d*x])/(d*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{\sec(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx &= \frac{\int (b \sec(c+dx))^{2/3} dx}{b} \\
&= \frac{\left(\left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b} \right)^{2/3}} dx}{b} \\
&= \frac{3 \cos(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{bd \sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 60, normalized size = 1.13

$$\frac{3 \sqrt{-\tan^2(c+dx)} \cot(c+dx) (b \sec(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sec^2(c+dx)\right)}{2bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(1/3), x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(2*b*d)

fricas [F] time = 0.57, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \sec(dx+c))^{2/3}}{b}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)/b, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.44, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*sec(d*x+c))^(1/3), x)

[Out] int(sec(d*x+c)/(b*sec(d*x+c))^(1/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(1/3), x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) \left(\frac{b}{\cos(c+dx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/3)), x)

[Out] int(1/(cos(c + d*x)*(b/cos(c + d*x))^(1/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))**(1/3), x)

[Out] Integral(sec(c + d*x)/(b*sec(c + d*x))**(1/3), x)

$$3.196 \quad \int \frac{1}{\sqrt[3]{b \sec(c+dx)}} dx$$

Optimal. Leaf size=56

$$-\frac{3b \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{4/3}}$$

[Out] $-3/4*b*hypergeom([1/2, 2/3], [5/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(4/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$-\frac{3b \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^(-1/3), x]

[Out] $(-3*b*Hypergeometric2F1[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(4*d*(b*\text{Sec}[c + d*x])^{(4/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \frac{1}{\sqrt[3]{b \sec(c+dx)}} dx = \left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \int \sqrt[3]{\frac{\cos(c+dx)}{b}} dx$$

$$= -\frac{3 \cos^2(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{4bd\sqrt{\sin^2(c+dx)}}$$

Mathematica [A] time = 0.05, size = 55, normalized size = 0.98

$$-\frac{3\sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sec^2(c+dx)\right)}{d\sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(-1/3), x]

[Out] (-3*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(d*(b*Sec[c + d*x])^(1/3))

fricas [F] time = 0.62, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}}}{b \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.45, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*sec(d*x+c))^(1/3),x)`

[Out] `int(1/(b*sec(d*x+c))^(1/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(1/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b/cos(c + d*x))^(1/3),x)`

[Out] `int(1/(b/cos(c + d*x))^(1/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(d*x+c))**(1/3),x)`

[Out] `Integral((b*sec(c + d*x))**(1/3), x)`

$$3.197 \quad \int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{7/3}}$$

[Out] $-3/7*b^2*\text{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]/(b*\text{Sec}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*b^2*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(7/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \&\& \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n, x\} \&\& \text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_)]*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}\{b, c, d, n, x\} \&\& \text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \frac{\cos(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx &= b \int \frac{1}{(b \sec(c+dx))^{4/3}} dx \\
&= \left(b \left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \left(\frac{\cos(c+dx)}{b} \right)^{4/3} dx \\
&= -\frac{3 \cos^3(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{7bd\sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 58, normalized size = 1.00

$$-\frac{3b\sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \sec^2(c+dx)\right)}{4d(b \sec(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(1/3), x]

[Out] (-3*b*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(4*d*(b*Sec[c + d*x])^(4/3))

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}} \cos(dx+c)}{b \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*cos(d*x + c)/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/3), x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.45, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*sec(d*x+c))^(1/3), x)

[Out] int(cos(d*x+c)/(b*sec(d*x+c))^(1/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(1/3), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b/cos(c + d*x))^(1/3), x)

[Out] int(cos(c + d*x)/(b/cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))**(1/3), x)

[Out] Integral(cos(c + d*x)/(b*sec(c + d*x))**(1/3), x)

$$3.198 \quad \int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

Optimal. Leaf size=58

$$\frac{3b^3 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{10/3}}$$

[Out] $-3/10*b^3*\text{hypergeom}([1/2, 5/3], [8/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(10/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3b^3 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{10/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^2/(b*\text{Sec}[c + d*x])^{(1/3)}, x]$

[Out] $(-3*b^3*\text{Hypergeometric2F1}[1/2, 5/3, 8/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(10*d*(b*\text{Sec}[c + d*x])^{(10/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_)]*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/((\text{Sin}[c + d*x]/b)^n, x)], x] /; \text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx &= b^2 \int \frac{1}{(b \sec(c+dx))^{7/3}} dx \\
&= \left(b^2 \left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \left(\frac{\cos(c+dx)}{b} \right)^{7/3} dx \\
&= -\frac{3 \cos^4(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{10bd\sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.07, size = 60, normalized size = 1.03

$$-\frac{3b^2\sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \sec^2(c+dx)\right)}{7d(b \sec(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(1/3), x]

[Out] (-3*b^2*Cot[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(7*d*(b*Sec[c + d*x])^(7/3))

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}} \cos(dx+c)^2}{b \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*cos(d*x + c)^2/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.93, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)

[Out] int(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{\left(\frac{b}{\cos(c+dx)}\right)^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/3),x)

[Out] int(cos(c + d*x)^2/(b/cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(1/3),x)

[Out] Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(1/3), x)

$$3.199 \quad \int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$$

Optimal. Leaf size=56

$$\frac{3 \sin(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

[Out] -3*hypergeom([1/6, 1/2], [7/6], cos(d*x+c)^2)*sin(d*x+c)/b/d/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right)}{bd \sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2/(b*Sec[c + d*x])^(4/3), x]

[Out] (-3*Hypergeometric2F1[1/6, 1/2, 7/6, Cos[c + d*x]^2]*Sin[c + d*x])/(b*d*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{\sec^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx &= \frac{\int (b \sec(c+dx))^{2/3} dx}{b^2} \\
&= \frac{\left(\left(\frac{\cos(c+dx)}{b}\right)^{2/3} (b \sec(c+dx))^{2/3}\right) \int \frac{1}{\left(\frac{\cos(c+dx)}{b}\right)^{2/3}} dx}{b^2} \\
&= \frac{3 \cos(c+dx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{b^2 d \sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.01, size = 60, normalized size = 1.07

$$\frac{3\sqrt{-\tan^2(c+dx)} \cot(c+dx) (b \sec(c+dx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sec^2(c+dx)\right)}{2b^2 d}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2/(b*Sec[c + d*x])^(4/3), x]

[Out] (3*Cot[c + d*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(2*b^2*d)

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}}}{b^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)/b^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^2}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)

maple [F] time = 0.44, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3), x)

[Out] int(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^2}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2/(b*sec(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx)^2 \left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3)), x)

[Out] int(1/(cos(c + d*x)^2*(b/cos(c + d*x))^(4/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(c + dx)}{(b \sec(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**2/(b*sec(d*x+c))**(4/3), x)

[Out] Integral(sec(c + d*x)**2/(b*sec(c + d*x))**(4/3), x)

$$3.200 \quad \int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx$$

Optimal. Leaf size=55

$$\frac{3 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{4/3}}$$

[Out] $-3/4 \cdot \text{hypergeom}\left(\left[\frac{1}{2}, \frac{2}{3}\right], \left[\frac{5}{3}\right], \cos(d*x+c)^2\right) \cdot \sin(d*x+c) / d / (b \cdot \sec(d*x+c))^{4/3} / (\sin(d*x+c)^2)^{1/2}$

Rubi [A] time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right)}{4d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]/(b*Sec[c + d*x])^(4/3), x]

[Out] $(-3 \cdot \text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Cos}[c + d*x]^2] \cdot \text{Sin}[c + d*x]) / (4 \cdot d \cdot (b \cdot \text{Sec}[c + d*x])^{4/3} \cdot \text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/((Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{\sec(c+dx)}{(b \sec(c+dx))^{4/3}} dx &= \frac{\int \frac{1}{\sqrt[3]{b \sec(c+dx)}} dx}{b} \\
&= \frac{\left(\left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \sqrt[3]{\frac{\cos(c+dx)}{b}} dx}{b} \\
&= -\frac{3 \cos^2(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{4b^2 d \sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.01, size = 58, normalized size = 1.05

$$-\frac{3\sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sec^2(c+dx)\right)}{bd \sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]/(b*Sec[c + d*x])^(4/3), x]

[Out] (-3*Cot[c + d*x]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(b*d*(b*Sec[c + d*x])^(1/3))

fricas [F] time = 1.02, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{(b \sec(dx+c))^{2/3}}{b^2 \sec(dx+c)}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)/(b^2*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)}{(b \sec(dx+c))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(4/3), x)

maple [F] time = 0.43, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)/(b*sec(d*x+c))^(4/3), x)

[Out] int(sec(d*x+c)/(b*sec(d*x+c))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate(sec(d*x + c)/(b*sec(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(c + dx) \left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(c + d*x)*(b/cos(c + d*x))^(4/3)), x)

[Out] int(1/(cos(c + d*x)*(b/cos(c + d*x))^(4/3)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(c + dx)}{(b \sec(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)/(b*sec(d*x+c))**(4/3), x)

[Out] Integral(sec(c + d*x)/(b*sec(c + d*x))**(4/3), x)

$$3.201 \quad \int \frac{1}{(b \sec(c+dx))^{4/3}} dx$$

Optimal. Leaf size=56

$$\frac{3b \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{7/3}}$$

[Out] $-3/7*b*\text{hypergeom}([1/2, 7/6], [13/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(7/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3772, 2643}

$$\frac{3b \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c+dx)\right)}{7d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[c + d*x])^{(-4/3)}, x]$

[Out] $(-3*b*\text{Hypergeometric2F1}[1/2, 7/6, 13/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(7*d*(b*\text{Sec}[c + d*x])^{(7/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 2643

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n + 1)}*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n - 1)}*((\text{Sin}[c + d*x]/b)^{(n - 1)}*\text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int \frac{1}{(b \sec(c + dx))^{4/3}} dx = \left(\frac{\cos(c + dx)}{b} \right)^{2/3} (b \sec(c + dx))^{2/3} \int \left(\frac{\cos(c + dx)}{b} \right)^{4/3} dx$$

$$= -\frac{3 \cos^3(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \cos^2(c + dx)\right) (b \sec(c + dx))^{2/3} \sin(c + dx)}{7b^2 d \sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.00, size = 57, normalized size = 1.02

$$\frac{3\sqrt{-\tan^2(c + dx)} \cot(c + dx) {}_2F_1\left(-\frac{2}{3}, \frac{1}{2}; \frac{1}{3}; \sec^2(c + dx)\right)}{4d(b \sec(c + dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^(-4/3), x]

[Out] (-3*Cot[c + d*x]*Hypergeometric2F1[-2/3, 1/2, 1/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(4*d*(b*Sec[c + d*x])^(4/3))

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx + c))^{2/3}}{b^2 \sec(dx + c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)/(b^2*sec(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3), x)

maple [F] time = 0.38, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*sec(d*x+c))^(4/3),x)`

[Out] `int(1/(b*sec(d*x+c))^(4/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(d*x+c))^(4/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(4/3), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b/cos(c + d*x))^(4/3),x)`

[Out] `int(1/(b/cos(c + d*x))^(4/3), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(d*x+c))**(4/3),x)`

[Out] `Integral((b*sec(c + d*x))**(4/3), x)`

$$3.202 \quad \int \frac{\cos(c+dx)}{(b \sec(c+dx))^{4/3}} dx$$

Optimal. Leaf size=58

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{10/3}}$$

[Out] $-3/10*b^2*\text{hypergeom}([1/2, 5/3], [8/3], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(10/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{3b^2 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c+dx)\right)}{10d\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{10/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]/(b*\text{Sec}[c + d*x])^{(4/3)}, x]$

[Out] $(-3*b^2*\text{Hypergeometric2F1}[1/2, 5/3, 8/3, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(10*d*(b*\text{Sec}[c + d*x])^{(10/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n+1)}*\text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2])/(b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n, x\} \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_*) + (d_*)*(x_)]*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)}*((\text{Sin}[c + d*x]/b)^{(n-1)}*\text{Int}[1/((\text{Sin}[c + d*x]/b)^n, x)], x] /; \text{FreeQ}\{b, c, d, n, x\} \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \frac{\cos(c+dx)}{(b \sec(c+dx))^{4/3}} dx &= b \int \frac{1}{(b \sec(c+dx))^{7/3}} dx \\
&= \left(b \left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \left(\frac{\cos(c+dx)}{b} \right)^{7/3} dx \\
&= -\frac{3 \cos^4(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{5}{3}; \frac{8}{3}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{10b^2 d \sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 58, normalized size = 1.00

$$-\frac{3b\sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{7}{6}, \frac{1}{2}; -\frac{1}{6}; \sec^2(c+dx)\right)}{7d(b \sec(c+dx))^{7/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]/(b*Sec[c + d*x])^(4/3), x]

[Out] (-3*b*Cot[c + d*x]*Hypergeometric2F1[-7/6, 1/2, -1/6, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(7*d*(b*Sec[c + d*x])^(7/3))

fricas [F] time = 0.78, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}} \cos(dx+c)}{b^2 \sec(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*cos(d*x + c)/(b^2*sec(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(4/3), x)

maple [F] time = 0.45, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)/(b*sec(d*x+c))^(4/3), x)

[Out] int(cos(d*x+c)/(b*sec(d*x+c))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)/(b*sec(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)/(b/cos(c + d*x))^(4/3), x)

[Out] int(cos(c + d*x)/(b/cos(c + d*x))^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(c + dx)}{(b \sec(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)/(b*sec(d*x+c))**(4/3), x)

[Out] Integral(cos(c + d*x)/(b*sec(c + d*x))**(4/3), x)

$$3.203 \quad \int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx$$

Optimal. Leaf size=58

$$\frac{3b^3 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c+dx)\right)}{13d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{13/3}}$$

[Out] $-3/13*b^3*\text{hypergeom}([1/2, 13/6], [19/6], \cos(d*x+c)^2)*\sin(d*x+c)/d/(b*\sec(d*x+c))^{(13/3)}/(\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3772, 2643}

$$\frac{3b^3 \sin(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c+dx)\right)}{13d \sqrt{\sin^2(c+dx)} (b \sec(c+dx))^{13/3}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2/(b*Sec[c + d*x])^(4/3), x]

[Out] $(-3*b^3*\text{Hypergeometric2F1}[1/2, 13/6, 19/6, \text{Cos}[c + d*x]^2]*\text{Sin}[c + d*x])/(13*d*(b*\text{Sec}[c + d*x])^{(13/3)}*\text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^2(c+dx)}{(b \sec(c+dx))^{4/3}} dx &= b^2 \int \frac{1}{(b \sec(c+dx))^{10/3}} dx \\
&= \left(b^2 \left(\frac{\cos(c+dx)}{b} \right)^{2/3} (b \sec(c+dx))^{2/3} \right) \int \left(\frac{\cos(c+dx)}{b} \right)^{10/3} dx \\
&= -\frac{3 \cos^5(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{13}{6}; \frac{19}{6}; \cos^2(c+dx)\right) (b \sec(c+dx))^{2/3} \sin(c+dx)}{13b^2 d \sqrt{\sin^2(c+dx)}}
\end{aligned}$$

Mathematica [A] time = 0.09, size = 60, normalized size = 1.03

$$-\frac{3b^2 \sqrt{-\tan^2(c+dx)} \cot(c+dx) {}_2F_1\left(-\frac{5}{3}, \frac{1}{2}; -\frac{2}{3}; \sec^2(c+dx)\right)}{10d(b \sec(c+dx))^{10/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2/(b*Sec[c + d*x])^(4/3), x]

[Out] (-3*b^2*Cot[c + d*x]*Hypergeometric2F1[-5/3, 1/2, -2/3, Sec[c + d*x]^2]*Sqrt[-Tan[c + d*x]^2])/(10*d*(b*Sec[c + d*x])^(10/3))

fricas [F] time = 0.87, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{2/3} \cos(dx+c)^2}{b^2 \sec(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*cos(d*x + c)^2/(b^2*sec(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx+c)^2}{(b \sec(dx+c))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3), x, algorithm="giac")

[Out] integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)

maple [F] time = 1.03, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3), x)

[Out] int(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos(dx + c)^2}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2/(b*sec(d*x+c))^(4/3), x, algorithm="maxima")

[Out] integrate(cos(d*x + c)^2/(b*sec(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(c + dx)^2}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(c + d*x)^2/(b/cos(c + d*x))^(4/3), x)

[Out] int(cos(c + d*x)^2/(b/cos(c + d*x))^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(c + dx)}{(b \sec(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)**2/(b*sec(d*x+c))**(4/3), x)

[Out] Integral(cos(c + d*x)**2/(b*sec(c + d*x))**(4/3), x)

3.204 $\int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx$

Optimal. Leaf size=81

$$\frac{3b \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \sec^m(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(-3m - 1); \frac{1}{6}(5 - 3m); \cos^2(c + dx)\right)}{d(3m + 1)\sqrt{\sin^2(c + dx)}}$$

[Out] 3*b*hypergeom([1/2, -1/6-1/2*m], [5/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^m*(b*sec(d*x+c))^(1/3)*sin(d*x+c)/d/(1+3*m)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{3b \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \sec^m(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(-3m - 1); \frac{1}{6}(5 - 3m); \cos^2(c + dx)\right)}{d(3m + 1)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^(4/3), x]

[Out] (3*b*Hypergeometric2F1[1/2, (-1 - 3*m)/6, (5 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^m*(b*Sec[c + d*x])^(1/3)*Sin[c + d*x])/(d*(1 + 3*m)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^m(c + dx)(b \sec(c + dx))^{4/3} dx &= \frac{(b \sqrt[3]{b \sec(c + dx)}) \int \sec^{\frac{4}{3}+m}(c + dx) dx}{\sqrt[3]{\sec(c + dx)}} \\ &= \left(b \cos^{\frac{1}{3}+m}(c + dx) \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} \right) \int \cos^{-\frac{4}{3}-m}(c + dx) dx \\ &= \frac{3b {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(-1 - 3m); \frac{1}{6}(5 - 3m); \cos^2(c + dx)\right) \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)}}{d(1 + 3m)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 83, normalized size = 1.02

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx)(b \sec(c + dx))^{4/3} \sec^{m-1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m + \frac{4}{3}\right); \frac{1}{2}\left(m + \frac{10}{3}\right); \sec^2(c + dx)\right)}{d\left(m + \frac{4}{3}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^(4/3), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (4/3 + m)/2, (10/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(4/3)*Sqrt[-Tan[c + d*x]^2])/(d*(4/3 + m))

fricas [F] time = 0.83, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^{\frac{1}{3}} b \sec(dx + c)^m \sec(dx + c), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*b*sec(d*x + c)^m*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^m, x)

maple [F] time = 0.83, size = 0, normalized size = 0.00

$$\int (\sec^m(dx + c)) (b \sec(dx + c))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x)

[Out] int(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{4}{3}} \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(4/3)*sec(d*x + c)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{4}{3}} \left(\frac{1}{\cos(c + dx)} \right)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(4/3)*(1/cos(c + d*x))^m,x)

[Out] int((b/cos(c + d*x))^(4/3)*(1/cos(c + d*x))^m, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**m*(b*sec(d*x+c))**(4/3),x)

[Out] Timed out

3.205 $\int \sec^m(c + dx)(b \sec(c + dx))^{2/3} dx$

Optimal. Leaf size=82

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{2/3} \sec^{m-1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(1 - 3m); \frac{1}{6}(7 - 3m); \cos^2(c + dx)\right)}{d(1 - 3m)\sqrt{\sin^2(c + dx)}}$$

[Out] -3*hypergeom([1/2, 1/6-1/2*m], [7/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*(b*sec(d*x+c))^(2/3)*sin(d*x+c)/d/(1-3*m)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{3 \sin(c + dx)(b \sec(c + dx))^{2/3} \sec^{m-1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(1 - 3m); \frac{1}{6}(7 - 3m); \cos^2(c + dx)\right)}{d(1 - 3m)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^(2/3), x]

[Out] (-3*Hypergeometric2F1[1/2, (1 - 3*m)/6, (7 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(2/3)*Sin[c + d*x])/(d*(1 - 3*m)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m + n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^m(c+dx)(b \sec(c+dx))^{2/3} dx &= \frac{(b \sec(c+dx))^{2/3} \int \sec^{\frac{2}{3}+m}(c+dx) dx}{\sec^{\frac{2}{3}}(c+dx)} \\ &= \left(\cos^{\frac{2}{3}+m}(c+dx) \sec^m(c+dx)(b \sec(c+dx))^{2/3} \right) \int \cos^{-\frac{2}{3}-m}(c+dx) dx \\ &= -\frac{{}_3F_2\left(\frac{1}{2}, \frac{1}{6}(1-3m); \frac{1}{6}(7-3m); \cos^2(c+dx)\right) \sec^{-1+m}(c+dx)(b \sec(c+dx))^{2/3}}{d(1-3m)\sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.14, size = 83, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx)(b \sec(c+dx))^{2/3} \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m+\frac{2}{3}\right); \frac{1}{2}\left(m+\frac{8}{3}\right); \sec^2(c+dx)\right)}{d\left(m+\frac{2}{3}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^(2/3), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (2/3 + m)/2, (8/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(2/3)*Sqrt[-Tan[c + d*x]^2])/(d*(2/3 + m))

fricas [F] time = 0.54, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \sec(dx+c)\right)^{\frac{2}{3}} \sec(dx+c)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^{\frac{2}{3}} \sec(dx+c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x, algorithm="giac")`

[Out] `integrate((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m, x)`

maple [F] time = 0.73, size = 0, normalized size = 0.00

$$\int (\sec^m(dx + c)) (b \sec(dx + c))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x)`

[Out] `int(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{2}{3}} \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(2/3),x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{2}{3}} \left(\frac{1}{\cos(c + dx)} \right)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^(2/3)*(1/cos(c + d*x))^m,x)`

[Out] `int((b/cos(c + d*x))^(2/3)*(1/cos(c + d*x))^m, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^{\frac{2}{3}} \sec^m(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**m*(b*sec(d*x+c))**(2/3),x)`

[Out] `Integral((b*sec(c + d*x))**(2/3)*sec(c + d*x)**m, x)`

3.206 $\int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx$

Optimal. Leaf size=82

$$\frac{3 \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \sec^{m-1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(2 - 3m); \frac{1}{6}(8 - 3m); \cos^2(c + dx)\right)}{d(2 - 3m) \sqrt{\sin^2(c + dx)}}$$

[Out] $-3 \text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{3} - \frac{1}{2}m\right], \left[\frac{4}{3} - \frac{1}{2}m\right], \cos(d*x+c)^2\right) * \sec(d*x+c)^{-1+m} * (b * \sec(d*x+c))^{1/3} * \sin(d*x+c) / d / (2 - 3m) / (\sin(d*x+c)^2)^{1/2}$

Rubi [A] time = 0.04, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{3 \sin(c + dx) \sqrt[3]{b \sec(c + dx)} \sec^{m-1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(2 - 3m); \frac{1}{6}(8 - 3m); \cos^2(c + dx)\right)}{d(2 - 3m) \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[c + d*x]^m * (b * \text{Sec}[c + d*x])^{1/3}, x]$

[Out] $(-3 * \text{Hypergeometric2F1}[1/2, (2 - 3*m)/6, (8 - 3*m)/6, \text{Cos}[c + d*x]^2] * \text{Sec}[c + d*x]^{-1 + m} * (b * \text{Sec}[c + d*x])^{1/3} * \text{Sin}[c + d*x]) / (d * (2 - 3*m) * \text{Sqrt}[\text{Sin}[c + d*x]^2])$

Rule 20

$\text{Int}[(u_.) * ((a_.) * (v_.))^{(m_.)} * ((b_.) * (v_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b^{\text{IntPart}[n]} * (b*v)^{\text{FracPart}[n]}) / (a^{\text{IntPart}[n]} * (a*v)^{\text{FracPart}[n]}) , \text{Int}[u * (a*v)^{(m+n)}, x], x] /;$ FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2643

$\text{Int}[(b_.) * \sin[(c_.) + (d_.) * (x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x] * (b * \text{Sin}[c + d*x])^{(n+1)} * \text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2]) / (b * d * (n+1) * \text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /;$ FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.) * (x_.)] * (b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b * \text{Csc}[c + d*x])^{(n-1)} * ((\text{Sin}[c + d*x] / b)^{(n-1)} * \text{Int}[1 / (\text{Sin}[c + d*x] / b)^n, x]), x] /;$ Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} dx &= \frac{\sqrt[3]{b \sec(c + dx)} \int \sec^{\frac{1}{3}+m}(c + dx) dx}{\sqrt[3]{\sec(c + dx)}} \\ &= \left(\cos^{\frac{1}{3}+m}(c + dx) \sec^m(c + dx) \sqrt[3]{b \sec(c + dx)} \right) \int \cos^{-\frac{1}{3}-m}(c + dx) dx \\ &= -\frac{{}_3F_2\left(\frac{1}{2}, \frac{1}{6}(2-3m), \frac{1}{6}(8-3m); \cos^2(c + dx)\right) \sec^{-1+m}(c + dx) \sqrt[3]{b \sec(c + dx)}}{d(2-3m)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.14, size = 83, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx) \sqrt[3]{b \sec(c + dx)} \sec^{m-1}(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m + \frac{1}{3}\right); \frac{1}{2}\left(m + \frac{7}{3}\right); \sec^2(c + dx)\right)}{d\left(m + \frac{1}{3}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^(1/3), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (1/3 + m)/2, (7/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^(1/3)*Sqrt[-Tan[c + d*x]^2])/(d*(1/3 + m))

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \sec(dx + c)\right)^{\frac{1}{3}} \sec(dx + c)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m, x)

maple [F] time = 0.73, size = 0, normalized size = 0.00

$$\int (\sec^m(dx + c)) (b \sec(dx + c))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x)

[Out] int(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^{\frac{1}{3}} \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^{\frac{1}{3}} \left(\frac{1}{\cos(c + dx)} \right)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^(1/3)*(1/cos(c + d*x))^m,x)

[Out] int((b/cos(c + d*x))^(1/3)*(1/cos(c + d*x))^m, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sec(c + dx)} \sec^m(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**m*(b*sec(d*x+c))**(1/3),x)

[Out] Integral((b*sec(c + d*x))**(1/3)*sec(c + d*x)**m, x)

$$3.207 \quad \int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx$$

Optimal. Leaf size=82

$$\frac{3 \sin(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(4-3m); \frac{1}{6}(10-3m); \cos^2(c+dx)\right)}{d(4-3m)\sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

[Out] -3*hypergeom([1/2, 2/3-1/2*m], [5/3-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*sin(d*x+c)/d/(4-3*m)/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{3 \sin(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(4-3m); \frac{1}{6}(10-3m); \cos^2(c+dx)\right)}{d(4-3m)\sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^m/(b*Sec[c + d*x])^(1/3), x]

[Out] (-3*Hypergeometric2F1[1/2, (4 - 3*m)/6, (10 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sin[c + d*x]/(d*(4 - 3*m)*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^m(c+dx)}{\sqrt[3]{b \sec(c+dx)}} dx &= \frac{\sqrt[3]{\sec(c+dx)} \int \sec^{-\frac{1}{3}+m}(c+dx) dx}{\sqrt[3]{b \sec(c+dx)}} \\ &= \frac{\left(\cos^{\frac{2}{3}+m}(c+dx) \sec^{1+m}(c+dx)\right) \int \cos^{\frac{1}{3}-m}(c+dx) dx}{\sqrt[3]{b \sec(c+dx)}} \\ &= -\frac{3 {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(4-3m); \frac{1}{6}(10-3m); \cos^2(c+dx)\right) \sec^{-1+m}(c+dx) \sin(c+dx)}{d(4-3m) \sqrt[3]{b \sec(c+dx)} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.13, size = 83, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m-\frac{1}{3}\right); \frac{1}{2}\left(m+\frac{5}{3}\right); \sec^2(c+dx)\right)}{d\left(m-\frac{1}{3}\right) \sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m/(b*Sec[c + d*x])^(1/3), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (-1/3 + m)/2, (5/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sqrt[-Tan[c + d*x]^2])/(d*(-1/3 + m)*(b*Sec[c + d*x])^(1/3))

fricas [F] time = 0.75, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}} \sec(dx+c)^m}{b \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^m}{(b \sec(dx+c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(1/3), x)

maple [F] time = 0.68, size = 0, normalized size = 0.00

$$\int \frac{\sec^m(dx + c)}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x)

[Out] int(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(1/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^m}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(1/3),x)

[Out] int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(1/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^m(c + dx)}{\sqrt[3]{b \sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**m/(b*sec(d*x+c))**(1/3),x)

[Out] Integral(sec(c + d*x)**m/(b*sec(c + d*x))**(1/3), x)

$$3.208 \quad \int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx$$

Optimal. Leaf size=82

$$\frac{3 \sin(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(5-3m); \frac{1}{6}(11-3m); \cos^2(c+dx)\right)}{d(5-3m)\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{2/3}}$$

[Out] -3*hypergeom([1/2, 5/6-1/2*m], [11/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*sin(d*x+c)/d/(5-3*m)/(b*sec(d*x+c))^(2/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{3 \sin(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(5-3m); \frac{1}{6}(11-3m); \cos^2(c+dx)\right)}{d(5-3m)\sqrt{\sin^2(c+dx)}(b \sec(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^m/(b*Sec[c + d*x])^(2/3), x]

[Out] (-3*Hypergeometric2F1[1/2, (5 - 3*m)/6, (11 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sin[c + d*x])/(d*(5 - 3*m)*(b*Sec[c + d*x])^(2/3)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{2/3}} dx &= \frac{\sec^{\frac{2}{3}}(c+dx) \int \sec^{-\frac{2}{3}+m}(c+dx) dx}{(b \sec(c+dx))^{2/3}} \\ &= \frac{\left(\cos^{\frac{1}{3}+m}(c+dx) \sec^{1+m}(c+dx)\right) \int \cos^{\frac{2}{3}-m}(c+dx) dx}{(b \sec(c+dx))^{2/3}} \\ &= \frac{3 {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(5-3m); \frac{1}{6}(11-3m); \cos^2(c+dx)\right) \sec^{-1+m}(c+dx) \sin(c+dx)}{d(5-3m)(b \sec(c+dx))^{2/3} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.14, size = 83, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m-\frac{2}{3}\right); \frac{1}{2}\left(m+\frac{4}{3}\right); \sec^2(c+dx)\right)}{d\left(m-\frac{2}{3}\right)(b \sec(c+dx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m/(b*Sec[c + d*x])^(2/3), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (-2/3 + m)/2, (4/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sqrt[-Tan[c + d*x]^2])/(d*(-2/3 + m)*(b*Sec[c + d*x])^(2/3))

fricas [F] time = 0.74, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{1}{3}} \sec(dx+c)^m}{b \sec(dx+c)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(1/3)*sec(d*x + c)^m/(b*sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^m}{(b \sec(dx+c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(2/3), x)

maple [F] time = 0.58, size = 0, normalized size = 0.00

$$\int \frac{\sec^m(dx + c)}{(b \sec(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x)

[Out] int(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(2/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(2/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^m}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(2/3),x)

[Out] int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(2/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**m/(b*sec(d*x+c))**(2/3),x)
```

```
[Out] Integral(sec(c + d*x)**m/(b*sec(c + d*x))**(2/3), x)
```

$$3.209 \quad \int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx$$

Optimal. Leaf size=85

$$\frac{3 \sin(c+dx) \sec^{m-2}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(7-3m); \frac{1}{6}(13-3m); \cos^2(c+dx)\right)}{bd(7-3m)\sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

[Out] -3*hypergeom([1/2, 7/6-1/2*m], [13/6-1/2*m], cos(d*x+c)^2)*sec(d*x+c)^(-2+m)*sin(d*x+c)/b/d/(7-3*m)/(b*sec(d*x+c))^(1/3)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{3 \sin(c+dx) \sec^{m-2}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(7-3m); \frac{1}{6}(13-3m); \cos^2(c+dx)\right)}{bd(7-3m)\sqrt{\sin^2(c+dx)} \sqrt[3]{b \sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^m/(b*Sec[c + d*x])^(4/3), x]

[Out] (-3*Hypergeometric2F1[1/2, (7 - 3*m)/6, (13 - 3*m)/6, Cos[c + d*x]^2]*Sec[c + d*x]^(-2 + m)*Sin[c + d*x])/(b*d*(7 - 3*m)*(b*Sec[c + d*x])^(1/3)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{\sec^m(c+dx)}{(b \sec(c+dx))^{4/3}} dx &= \frac{\sqrt[3]{\sec(c+dx)} \int \sec^{-\frac{4}{3}+m}(c+dx) dx}{b \sqrt[3]{b \sec(c+dx)}} \\ &= \frac{\left(\cos^{\frac{2}{3}+m}(c+dx) \sec^{1+m}(c+dx)\right) \int \cos^{\frac{4}{3}-m}(c+dx) dx}{b \sqrt[3]{b \sec(c+dx)}} \\ &= -\frac{3 {}_2F_1\left(\frac{1}{2}, \frac{1}{6}(7-3m); \frac{1}{6}(13-3m); \cos^2(c+dx)\right) \sec^{-2+m}(c+dx) \sin(c+dx)}{bd(7-3m) \sqrt[3]{b \sec(c+dx)} \sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.18, size = 83, normalized size = 0.98

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx) \sec^{m-1}(c+dx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(m-\frac{4}{3}\right); \frac{1}{2}\left(m+\frac{2}{3}\right); \sec^2(c+dx)\right)}{d\left(m-\frac{4}{3}\right)(b \sec(c+dx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m/(b*Sec[c + d*x])^(4/3), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (-4/3 + m)/2, (2/3 + m)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*Sqrt[-Tan[c + d*x]^2])/(d*(-4/3 + m)*(b*Sec[c + d*x])^(4/3))

fricas [F] time = 0.73, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx+c))^{\frac{2}{3}} \sec(dx+c)^m}{b^2 \sec(dx+c)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^(2/3)*sec(d*x + c)^m/(b^2*sec(d*x + c)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx+c)^m}{(b \sec(dx+c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x, algorithm="giac")

[Out] integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(4/3), x)

maple [F] time = 0.64, size = 0, normalized size = 0.00

$$\int \frac{\sec^m(dx + c)}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x)

[Out] int(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(dx + c)^m}{(b \sec(dx + c))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m/(b*sec(d*x+c))^(4/3),x, algorithm="maxima")

[Out] integrate(sec(d*x + c)^m/(b*sec(d*x + c))^(4/3), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1}{\cos(c+dx)}\right)^m}{\left(\frac{b}{\cos(c+dx)}\right)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(4/3),x)

[Out] int((1/cos(c + d*x))^m/(b/cos(c + d*x))^(4/3), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^m(c + dx)}{(b \sec(c + dx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(d*x+c)**m/(b*sec(d*x+c))**(4/3),x)
```

```
[Out] Integral(sec(c + d*x)**m/(b*sec(c + d*x))**(4/3), x)
```

3.210 $\int \sec^m(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=89

$$\frac{\sin(c + dx) \sec^{m-1}(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-m - n + 1); \frac{1}{2}(-m - n + 3); \cos^2(c + dx)\right)}{d(-m - n + 1)\sqrt{\sin^2(c + dx)}}$$

[Out] -hypergeom([1/2, 1/2-1/2*m-1/2*n], [3/2-1/2*m-1/2*n], cos(d*x+c)^2)*sec(d*x+c)^(-1+m)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(1-m-n)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {20, 3772, 2643}

$$\frac{\sin(c + dx) \sec^{m-1}(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-m - n + 1); \frac{1}{2}(-m - n + 3); \cos^2(c + dx)\right)}{d(-m - n + 1)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^m*(b*Sec[c + d*x])^n,x]

[Out] -((Hypergeometric2F1[1/2, (1 - m - n)/2, (3 - m - n)/2, Cos[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(1 - m - n)*Sqrt[Sin[c + d*x]^2]))

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m + n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^m(c + dx)(b \sec(c + dx))^n dx &= (\sec^{-n}(c + dx)(b \sec(c + dx))^n) \int \sec^{m+n}(c + dx) dx \\ &= (\cos^{m+n}(c + dx) \sec^m(c + dx)(b \sec(c + dx))^n) \int \cos^{-m-n}(c + dx) dx \\ &= -\frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 - m - n); \frac{1}{2}(3 - m - n); \cos^2(c + dx)\right) \sec^{-1+m}(c + dx)(b \sec(c + dx))^n}{d(1 - m - n)\sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.07, size = 76, normalized size = 0.85

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx) \sec^{m-1}(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{m+n}{2}; \frac{1}{2}(m+n+2); \sec^2(c + dx)\right)}{d(m+n)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^m*(b*Sec[c + d*x])^n,x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (m + n)/2, (2 + m + n)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(-1 + m)*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(m + n))

fricas [F] time = 0.70, size = 0, normalized size = 0.00

$$\text{integral}((b \sec(dx + c))^n \sec(dx + c)^m, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*sec(d*x + c)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^m, x)

maple [F] time = 2.42, size = 0, normalized size = 0.00

$$\int (\sec^m(dx + c))(b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^m*(b*sec(d*x+c))^n,x)

[Out] int(sec(d*x+c)^m*(b*sec(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^m*(b*sec(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)}\right)^n \left(\frac{1}{\cos(c + dx)}\right)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^m,x)

[Out] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^m, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \sec^m(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**m*(b*sec(d*x+c))**n,x)

[Out] Integral((b*sec(c + d*x))**n*sec(c + d*x)**m, x)

3.211 $\int \sec^2(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=72

$$\frac{\sin(c + dx)(b \sec(c + dx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-n - 1); \frac{1-n}{2}; \cos^2(c + dx)\right)}{bd(n + 1)\sqrt{\sin^2(c + dx)}}$$

[Out] hypergeom([1/2, -1/2-1/2*n], [1/2-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^(1+n)*sin(d*x+c)/b/d/(1+n)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{\sin(c + dx)(b \sec(c + dx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-n - 1); \frac{1-n}{2}; \cos^2(c + dx)\right)}{bd(n + 1)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^2*(b*Sec[c + d*x])^n,x]

[Out] (Hypergeometric2F1[1/2, (-1 - n)/2, (1 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(1 + n)*Sin[c + d*x])/(b*d*(1 + n)*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \sec^2(c + dx)(b \sec(c + dx))^n dx &= \frac{\int (b \sec(c + dx))^{2+n} dx}{b^2} \\
&= \frac{\left(\frac{\cos(c+dx)}{b}\right)^n (b \sec(c + dx))^n \int \left(\frac{\cos(c+dx)}{b}\right)^{-2-n} dx}{b^2} \\
&= \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{2}(-1 - n); \frac{1-n}{2}; \cos^2(c + dx)\right) (b \sec(c + dx))^{1+n} \sin(c + dx)}{bd(1 + n)\sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 71, normalized size = 0.99

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx) \sec(c + dx) (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+2}{2}; \frac{n+4}{2}; \sec^2(c + dx)\right)}{d(n + 2)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^2*(b*Sec[c + d*x])^n,x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (2 + n)/2, (4 + n)/2, Sec[c + d*x]^2]*Sec[c + d*x]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(2 + n))

fricas [F] time = 0.58, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^n \sec(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*sec(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^2, x)

maple [F] time = 1.47, size = 0, normalized size = 0.00

$$\int (\sec^2(dx + c)) (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(d*x+c)^2*(b*sec(d*x+c))^n,x)`

[Out] `int(sec(d*x+c)^2*(b*sec(d*x+c))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^n*sec(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\cos(c+dx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^n/cos(c + d*x)^2,x)`

[Out] `int((b/cos(c + d*x))^n/cos(c + d*x)^2, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \sec^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(d*x+c)**2*(b*sec(d*x+c))**n,x)`

[Out] `Integral((b*sec(c + d*x))**n*sec(c + d*x)**2, x)`

3.212 $\int \sec(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=61

$$\frac{\sin(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, -\frac{n}{2}; \frac{2-n}{2}; \cos^2(c + dx)\right)}{dn\sqrt{\sin^2(c + dx)}}$$

[Out] hypergeom([1/2, -1/2*n], [1-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/n/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {16, 3772, 2643}

$$\frac{\sin(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, -\frac{n}{2}; \frac{2-n}{2}; \cos^2(c + dx)\right)}{dn\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]*(b*Sec[c + d*x])^n,x]

[Out] (Hypergeometric2F1[1/2, -n/2, (2 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*n*Sqrt[Sin[c + d*x]^2])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \sec(c + dx)(b \sec(c + dx))^n dx &= \frac{\int (b \sec(c + dx))^{1+n} dx}{b} \\
&= \frac{\left(\left(\frac{\cos(c+dx)}{b}\right)^n (b \sec(c + dx))^n\right) \int \left(\frac{\cos(c+dx)}{b}\right)^{-1-n} dx}{b} \\
&= \frac{{}_2F_1\left(\frac{1}{2}, -\frac{n}{2}; \frac{2-n}{2}; \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{dn\sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.04, size = 65, normalized size = 1.07

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \sec^2(c + dx)\right)}{d(n + 1)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]*(b*Sec[c + d*x])^n,x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(1 + n))

fricas [F] time = 0.75, size = 0, normalized size = 0.00

$$\text{integral}((b \sec(dx + c))^n \sec(dx + c), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*sec(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c), x)

maple [F] time = 1.69, size = 0, normalized size = 0.00

$$\int \sec(dx + c) (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)*(b*sec(d*x+c))^n,x)

[Out] int(sec(d*x+c)*(b*sec(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\cos(c+dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n/cos(c + d*x),x)

[Out] int((b/cos(c + d*x))^n/cos(c + d*x), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \sec(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)*(b*sec(d*x+c))**n,x)

[Out] Integral((b*sec(c + d*x))**n*sec(c + d*x), x)

3.213 $\int (b \sec(c + dx))^n dx$

Optimal. Leaf size=73

$$\frac{b \sin(c + dx)(b \sec(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(c + dx)\right)}{d(1-n)\sqrt{\sin^2(c + dx)}}$$

[Out] -b*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^(n-1)*sin(d*x+c)/d/(1-n)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{b \sin(c + dx)(b \sec(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(c + dx)\right)}{d(1-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^n,x]

[Out] -((b*Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[c + d*x]^2]*(b*Sec[c + d*x])^(n - 1)*Sin[c + d*x])/(d*(1 - n)*Sqrt[Sin[c + d*x]^2]))

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int (b \sec(c + dx))^n dx = \left(\frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \int \left(\frac{\cos(c + dx)}{b} \right)^{-n} dx$$

$$= - \frac{\cos(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(1-n)\sqrt{\sin^2(c + dx)}}$$

Mathematica [A] time = 0.04, size = 61, normalized size = 0.84

$$\frac{\sqrt{-\tan^2(c + dx)} \cot(c + dx) (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{n+2}{2}; \sec^2(c + dx)\right)}{dn}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^n,x]

[Out] (Cot[c + d*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*sqrt[-Tan[c + d*x]^2])/(d*n)

fricas [F] time = 0.76, size = 0, normalized size = 0.00

$$\text{integral}((b \sec(dx + c))^n, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n, x)

maple [F] time = 1.07, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(d*x+c))^n,x)`

[Out] `int((b*sec(d*x+c))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))^n,x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^n, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(c + d*x))^n,x)`

[Out] `int((b/cos(c + d*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(d*x+c))**n,x)`

[Out] `Integral((b*sec(c + d*x))**n, x)`

3.214 $\int \cos(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=75

$$\frac{b^2 \sin(c + dx)(b \sec(c + dx))^{n-2} {}_2F_1\left(\frac{1}{2}, \frac{2-n}{2}; \frac{4-n}{2}; \cos^2(c + dx)\right)}{d(2-n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-b^2 \text{hypergeom}\left(\left[\frac{1}{2}, 1-1/2*n\right], \left[2-1/2*n\right], \cos(d*x+c)^2\right) * (b*\sec(d*x+c))^{(-2+n)} * \sin(d*x+c) / d / (2-n) / (\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {16, 3772, 2643}

$$\frac{b^2 \sin(c + dx)(b \sec(c + dx))^{n-2} {}_2F_1\left(\frac{1}{2}, \frac{2-n}{2}; \frac{4-n}{2}; \cos^2(c + dx)\right)}{d(2-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x] * (b*\text{Sec}[c + d*x])^n, x]$

[Out] $-((b^2*\text{Hypergeometric2F1}[1/2, (2-n)/2, (4-n)/2, \text{Cos}[c + d*x]^2] * (b*\text{Sec}[c + d*x])^{(-2+n)} * \text{Sin}[c + d*x]) / (d*(2-n)*\text{Sqrt}[\text{Sin}[c + d*x]^2]))$

Rule 16

$\text{Int}[(u_.) * (v_.)^{(m_.)} * ((b_.) * (v_.)^{(n_.)}), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u * (b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_.) * \sin[(c_.) + (d_.) * (x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x] * (b*\text{Sin}[c + d*x])^{(n+1)} * \text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2]) / (b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.) * (x_.)] * (b_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)} * ((\text{Sin}[c + d*x]/b)^{(n-1)} * \text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \cos(c + dx)(b \sec(c + dx))^n dx &= b \int (b \sec(c + dx))^{-1+n} dx \\
&= \left(b \left(\frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{1-n} dx \\
&= -\frac{\cos^2(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{2-n}{2}; \frac{4-n}{2}; \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(2-n)\sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.10, size = 68, normalized size = 0.91

$$\frac{b\sqrt{-\tan^2(c + dx)} \cot(c + dx)(b \sec(c + dx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{n-1}{2}; \frac{n+1}{2}; \sec^2(c + dx)\right)}{d(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]*(b*Sec[c + d*x])^n,x]

[Out] (b*Cot[c + d*x]*Hypergeometric2F1[1/2, (-1 + n)/2, (1 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^(-1 + n)*Sqrt[-Tan[c + d*x]^2])/(d*(-1 + n))

fricas [F] time = 0.86, size = 0, normalized size = 0.00

$$\text{integral}((b \sec(dx + c))^n \cos(dx + c), x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*cos(d*x + c), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*cos(d*x + c), x)

maple [F] time = 2.38, size = 0, normalized size = 0.00

$$\int \cos(dx + c) (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)*(b*sec(d*x+c))^n,x)`

[Out] `int(cos(d*x+c)*(b*sec(d*x+c))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \cos(dx + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))^n,x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^n*cos(d*x + c), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx) \left(\frac{b}{\cos(c + dx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)*(b/cos(c + d*x))^n,x)`

[Out] `int(cos(c + d*x)*(b/cos(c + d*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \cos(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)*(b*sec(d*x+c))**n,x)`

[Out] `Integral((b*sec(c + d*x))**n*cos(c + d*x), x)`

3.215 $\int \cos^2(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=75

$$\frac{b^3 \sin(c + dx)(b \sec(c + dx))^{n-3} {}_2F_1\left(\frac{1}{2}, \frac{3-n}{2}; \frac{5-n}{2}; \cos^2(c + dx)\right)}{d(3-n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-b^3 \text{hypergeom}([1/2, 3/2-1/2*n], [5/2-1/2*n], \cos(d*x+c)^2) * (b*\sec(d*x+c))^{(-3+n)} * \sin(d*x+c) / d / (3-n) / (\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{b^3 \sin(c + dx)(b \sec(c + dx))^{n-3} {}_2F_1\left(\frac{1}{2}, \frac{3-n}{2}; \frac{5-n}{2}; \cos^2(c + dx)\right)}{d(3-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[c + d*x]^2*(b*Sec[c + d*x])^n,x]

[Out] $-((b^3 \text{Hypergeometric2F1}[1/2, (3-n)/2, (5-n)/2, \text{Cos}[c + d*x]^2] * (b*\text{Sec}[c + d*x])^{(-3+n)} * \text{Sin}[c + d*x]) / (d*(3-n)*\text{Sqrt}[\text{Sin}[c + d*x]^2]))$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*SIN[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2]) / (b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \cos^2(c + dx)(b \sec(c + dx))^n dx &= b^2 \int (b \sec(c + dx))^{-2+n} dx \\
&= \left(b^2 \left(\frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{2-n} dx \\
&= -\frac{\cos^3(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{3-n}{2}; \frac{5-n}{2}; \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(3-n)\sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.08, size = 71, normalized size = 0.95

$$\frac{\cos^2(c + dx)\sqrt{-\tan^2(c + dx)} \cot(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n-2}{2}; \frac{n}{2}; \sec^2(c + dx)\right)}{d(n-2)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^2*(b*Sec[c + d*x])^n,x]

[Out] (Cos[c + d*x]^2*Cot[c + d*x]*Hypergeometric2F1[1/2, (-2 + n)/2, n/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(-2 + n))

fricas [F] time = 0.90, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx + c))^n \cos(dx + c)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*cos(d*x + c)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*cos(d*x + c)^2, x)

maple [F] time = 4.02, size = 0, normalized size = 0.00

$$\int (\cos^2(dx + c))(b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^2*(b*sec(d*x+c))^n,x)`

[Out] `int(cos(d*x+c)^2*(b*sec(d*x+c))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \cos(dx + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^2*(b*sec(d*x+c))^n,x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^n*cos(d*x + c)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^2 \left(\frac{b}{\cos(c + dx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^2*(b/cos(c + d*x))^n,x)`

[Out] `int(cos(c + d*x)^2*(b/cos(c + d*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \cos^2(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**2*(b*sec(d*x+c))**n,x)`

[Out] `Integral((b*sec(c + d*x))**n*cos(c + d*x)**2, x)`

3.216 $\int \cos^3(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=75

$$\frac{b^4 \sin(c + dx)(b \sec(c + dx))^{n-4} {}_2F_1\left(\frac{1}{2}, \frac{4-n}{2}; \frac{6-n}{2}; \cos^2(c + dx)\right)}{d(4-n)\sqrt{\sin^2(c + dx)}}$$

[Out] $-b^4 \text{hypergeom}\left(\left[\frac{1}{2}, 2-1/2*n\right], \left[3-1/2*n\right], \cos(d*x+c)^2\right) * (b*\sec(d*x+c))^{(-4+n)} * \sin(d*x+c) / d / (4-n) / (\sin(d*x+c)^2)^{(1/2)}$

Rubi [A] time = 0.06, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3772, 2643}

$$\frac{b^4 \sin(c + dx)(b \sec(c + dx))^{n-4} {}_2F_1\left(\frac{1}{2}, \frac{4-n}{2}; \frac{6-n}{2}; \cos^2(c + dx)\right)}{d(4-n)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[c + d*x]^3 * (b*\text{Sec}[c + d*x])^n, x]$

[Out] $-((b^4*\text{Hypergeometric2F1}[1/2, (4-n)/2, (6-n)/2, \text{Cos}[c + d*x]^2] * (b*\text{Sec}[c + d*x])^{(-4+n)} * \text{Sin}[c + d*x]) / (d*(4-n)*\text{Sqrt}[\text{Sin}[c + d*x]^2]))$

Rule 16

$\text{Int}[(u_.) * (v_.)^{(m_.)} * ((b_.) * (v_.)^{(n_.)}), x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u * (b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2643

$\text{Int}[(b_.) * \sin[(c_.) + (d_.) * (x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(\text{Cos}[c + d*x] * (b*\text{Sin}[c + d*x])^{(n+1)} * \text{Hypergeometric2F1}[1/2, (n+1)/2, (n+3)/2, \text{Sin}[c + d*x]^2]) / (b*d*(n+1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]), x] /; \text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[2*n]$

Rule 3772

$\text{Int}[(\text{csc}[(c_.) + (d_.) * (x_.)] * (b_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(b*\text{Csc}[c + d*x])^{(n-1)} * ((\text{Sin}[c + d*x]/b)^{(n-1)} * \text{Int}[1/(\text{Sin}[c + d*x]/b)^n, x]), x] /; \text{FreeQ}\{b, c, d, n\}, x \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned}
\int \cos^3(c + dx)(b \sec(c + dx))^n dx &= b^3 \int (b \sec(c + dx))^{-3+n} dx \\
&= \left(b^3 \left(\frac{\cos(c + dx)}{b} \right)^n (b \sec(c + dx))^n \right) \int \left(\frac{\cos(c + dx)}{b} \right)^{3-n} dx \\
&= -\frac{\cos^4(c + dx) {}_2F_1\left(\frac{1}{2}, \frac{4-n}{2}; \frac{6-n}{2}; \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(4-n)\sqrt{\sin^2(c + dx)}}
\end{aligned}$$

Mathematica [A] time = 0.11, size = 73, normalized size = 0.97

$$\frac{\cos^3(c + dx)\sqrt{-\tan^2(c + dx)} \cot(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{n-3}{2}; \frac{n-1}{2}; \sec^2(c + dx)\right)}{d(n-3)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[c + d*x]^3*(b*Sec[c + d*x])^n,x]

[Out] (Cos[c + d*x]^3*Cot[c + d*x]*Hypergeometric2F1[1/2, (-3 + n)/2, (-1 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*sqrt[-Tan[c + d*x]^2])/(d*(-3 + n))

fricas [F] time = 0.93, size = 0, normalized size = 0.00

$$\text{integral}((b \sec(dx + c))^n \cos(dx + c)^3, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*cos(d*x + c)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(d*x+c)^3*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*cos(d*x + c)^3, x)

maple [F] time = 6.42, size = 0, normalized size = 0.00

$$\int (\cos^3(dx + c)) (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(d*x+c)^3*(b*sec(d*x+c))^n,x)`

[Out] `int(cos(d*x+c)^3*(b*sec(d*x+c))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \cos(dx + c)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)^3*(b*sec(d*x+c))^n,x, algorithm="maxima")`

[Out] `integrate((b*sec(d*x + c))^n*cos(d*x + c)^3, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(c + dx)^3 \left(\frac{b}{\cos(c + dx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(c + d*x)^3*(b/cos(c + d*x))^n,x)`

[Out] `int(cos(c + d*x)^3*(b/cos(c + d*x))^n, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(d*x+c)**3*(b*sec(d*x+c))**n,x)`

[Out] Timed out

3.217 $\int \sec^{\frac{5}{2}}(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=80

$$\frac{2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-2n - 3); \frac{1}{4}(1 - 2n); \cos^2(c + dx)\right)}{d(2n + 3)\sqrt{\sin^2(c + dx)}}$$

[Out] 2*hypergeom([1/2, -3/4-1/2*n], [1/4-1/2*n], cos(d*x+c)^2)*sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(3+2*n)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{2 \sin(c + dx) \sec^{\frac{3}{2}}(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-2n - 3); \frac{1}{4}(1 - 2n); \cos^2(c + dx)\right)}{d(2n + 3)\sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^n,x]

[Out] (2*Hypergeometric2F1[1/2, (-3 - 2*n)/4, (1 - 2*n)/4, Cos[c + d*x]^2]*Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(3 + 2*n)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_)*sin[(c_)+(d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^{\frac{5}{2}}(c+dx)(b \sec(c+dx))^n dx &= (\sec^{-n}(c+dx)(b \sec(c+dx))^n) \int \sec^{\frac{5}{2}+n}(c+dx) dx \\ &= \left(\cos^{\frac{1}{2}+n}(c+dx) \sqrt{\sec(c+dx)} (b \sec(c+dx))^n \right) \int \cos^{-\frac{5}{2}-n}(c+dx) dx \\ &= \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-3-2n); \frac{1}{4}(1-2n); \cos^2(c+dx)\right) \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^n}{d(3+2n)\sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 81, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx) \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n+\frac{5}{2}\right); \frac{1}{2}\left(n+\frac{9}{2}\right); \sec^2(c+dx)\right)}{d\left(n+\frac{5}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(5/2)*(b*Sec[c + d*x])^n,x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (5/2 + n)/2, (9/2 + n)/2, Sec[c + d*x]^2]*Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(5/2 + n))

fricas [F] time = 0.95, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx+c))^n \sec(dx+c)^{\frac{5}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*sec(d*x + c)^(5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^n \sec(dx+c)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^(5/2), x)

maple [F] time = 0.88, size = 0, normalized size = 0.00

$$\int \left(\sec^{\frac{5}{2}}(dx + c) \right) (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x)

[Out] int(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(5/2)*(b*sec(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^n \left(\frac{1}{\cos(c + dx)} \right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(5/2),x)

[Out] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(5/2)*(b*sec(d*x+c))**n,x)

[Out] Timed out

3.218 $\int \sec^2(c + dx)(b \sec(c + dx))^n dx$

Optimal. Leaf size=80

$$\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)} (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-2n - 1); \frac{1}{4}(3 - 2n); \cos^2(c + dx)\right)}{d(2n + 1) \sqrt{\sin^2(c + dx)}}$$

[Out] 2*hypergeom([1/2, -1/4-1/2*n], [3/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)*sec(d*x+c)^(1/2)/d/(1+2*n)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{2 \sin(c + dx) \sqrt{\sec(c + dx)} (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-2n - 1); \frac{1}{4}(3 - 2n); \cos^2(c + dx)\right)}{d(2n + 1) \sqrt{\sin^2(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n,x]

[Out] (2*Hypergeometric2F1[1/2, (-1 - 2*n)/4, (3 - 2*n)/4, Cos[c + d*x]^2]*Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(1 + 2*n)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/((Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^{\frac{3}{2}}(c+dx)(b \sec(c+dx))^n dx &= (\sec^{-n}(c+dx)(b \sec(c+dx))^n) \int \sec^{\frac{3}{2}+n}(c+dx) dx \\ &= \left(\cos^{\frac{1}{2}+n}(c+dx) \sqrt{\sec(c+dx)} (b \sec(c+dx))^n \right) \int \cos^{-\frac{3}{2}-n}(c+dx) dx \\ &= \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{4}(-1-2n); \frac{1}{4}(3-2n); \cos^2(c+dx)\right) \sqrt{\sec(c+dx)} (b \sec(c+dx))^n}{d(1+2n)\sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.09, size = 81, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx) \sqrt{\sec(c+dx)} (b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n+\frac{3}{2}\right); \frac{1}{2}\left(n+\frac{7}{2}\right); \sec^2(c+dx)\right)}{d\left(n+\frac{3}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[c + d*x]^(3/2)*(b*Sec[c + d*x])^n,x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (3/2 + n)/2, (7/2 + n)/2, Sec[c + d*x]^2]*Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(3/2 + n))

fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left((b \sec(dx+c))^n \sec(dx+c)^{\frac{3}{2}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*sec(d*x + c)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^n \sec(dx+c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^(3/2), x)

maple [F] time = 1.01, size = 0, normalized size = 0.00

$$\int \left(\sec^{\frac{3}{2}}(dx + c) \right) (b \sec(dx + c))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x)

[Out] int(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sec(dx + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)^(3/2)*(b*sec(d*x+c))^n,x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n*sec(d*x + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^n \left(\frac{1}{\cos(c + dx)} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(3/2),x)

[Out] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \sec^{\frac{3}{2}}(c + dx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(d*x+c)**(3/2)*(b*sec(d*x+c))**n,x)

[Out] Integral((b*sec(c + d*x))**n*sec(c + d*x)**(3/2), x)

3.219 $\int \sqrt{\sec(c + dx)} (b \sec(c + dx))^n dx$

Optimal. Leaf size=80

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(1 - 2n); \frac{1}{4}(5 - 2n); \cos^2(c + dx)\right)}{d(1 - 2n)\sqrt{\sin^2(c + dx)}\sqrt{\sec(c + dx)}}$$

[Out] -2*hypergeom([1/2, 1/4-1/2*n], [5/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(1-2*n)/sec(d*x+c)^(1/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{2 \sin(c + dx)(b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(1 - 2n); \frac{1}{4}(5 - 2n); \cos^2(c + dx)\right)}{d(1 - 2n)\sqrt{\sin^2(c + dx)}\sqrt{\sec(c + dx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n,x]

[Out] (-2*Hypergeometric2F1[1/2, (1 - 2*n)/4, (5 - 2*n)/4, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x]/(d*(1 - 2*n)*Sqrt[Sec[c + d*x]]*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m + n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2643

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sqrt{\sec(c+dx)} (b \sec(c+dx))^n dx &= (\sec^{-n}(c+dx)(b \sec(c+dx))^n) \int \sec^{\frac{1}{2}+n}(c+dx) dx \\ &= \left(\cos^{\frac{1}{2}+n}(c+dx) \sqrt{\sec(c+dx)} (b \sec(c+dx))^n \right) \int \cos^{-\frac{1}{2}-n}(c+dx) dx \\ &= -\frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{4}(1-2n); \frac{1}{4}(5-2n); \cos^2(c+dx)\right) (b \sec(c+dx))^n \sin(c+dx)}{d(1-2n)\sqrt{\sec(c+dx)}\sqrt{\sin^2(c+dx)}} \end{aligned}$$

Mathematica [A] time = 0.10, size = 81, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c+dx)} \csc(c+dx) (b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n+\frac{1}{2}\right); \frac{1}{2}\left(n+\frac{5}{2}\right); \sec^2(c+dx)\right)}{d\left(n+\frac{1}{2}\right)\sqrt{\sec(c+dx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sec[c + d*x]]*(b*Sec[c + d*x])^n,x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (1/2 + n)/2, (5/2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(1/2 + n)*Sqrt[Sec[c + d*x]])

fricas [F] time = 0.89, size = 0, normalized size = 0.00

$$\text{integral}((b \sec(dx+c))^n \sqrt{\sec(dx+c)}, x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n*sqrt(sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx+c))^n \sqrt{\sec(dx+c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n*sqrt(sec(d*x + c)), x)

maple [F] time = 0.89, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n (\sqrt{\sec(dx + c)}) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x)

[Out] int((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(dx + c))^n \sqrt{\sec(dx + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n*sec(d*x+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n*sqrt(sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(c + dx)} \right)^n \sqrt{\frac{1}{\cos(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(1/2),x)

[Out] int((b/cos(c + d*x))^n*(1/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(c + dx))^n \sqrt{\sec(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**n*sec(d*x+c)**(1/2),x)

[Out] Integral((b*sec(c + d*x))**n*sqrt(sec(c + d*x)), x)

$$3.220 \quad \int \frac{(b \sec(c+dx))^n}{\sqrt{\sec(c+dx)}} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3-2n); \frac{1}{4}(7-2n); \cos^2(c+dx)\right)}{d(3-2n)\sqrt{\sin^2(c+dx)} \sec^{\frac{3}{2}}(c+dx)}$$

[Out] -2*hypergeom([1/2, 3/4-1/2*n], [7/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*si
n(d*x+c)/d/(3-2*n)/sec(d*x+c)^(3/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00,
number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} =$
0.143, Rules used = {20, 3772, 2643}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3-2n); \frac{1}{4}(7-2n); \cos^2(c+dx)\right)}{d(3-2n)\sqrt{\sin^2(c+dx)} \sec^{\frac{3}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^n/Sqrt[Sec[c + d*x]], x]

[Out] (-2*Hypergeometric2F1[1/2, (3 - 2*n)/4, (7 - 2*n)/4, Cos[c + d*x]^2]*(b*Sec
[c + d*x])^n*Sin[c + d*x])/(d*(3 - 2*n)*Sec[c + d*x]^(3/2)*Sqrt[Sin[c + d*x
]^2))

Rule 20

Int[(u_)*((a_)*(v_))^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart
[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m + n
) , x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !
IntegerQ[m + n]

Rule 2643

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(
b*Sin[c + d*x])^(n + 1)*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c
+ d*x]^2])/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x]
&& !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(b*Csc[c + d*x]
)^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx &= (\sec^{-n}(c + dx)(b \sec(c + dx))^n) \int \sec^{-\frac{1}{2}+n}(c + dx) dx \\ &= \left(\cos^{\frac{1}{2}+n}(c + dx) \sqrt{\sec(c + dx)} (b \sec(c + dx))^n \right) \int \cos^{\frac{1}{2}-n}(c + dx) dx \\ &= -\frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{4}(3 - 2n); \frac{1}{4}(7 - 2n); \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(3 - 2n) \sec^{\frac{3}{2}}(c + dx) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.11, size = 81, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx) (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n - \frac{1}{2}\right); \frac{1}{2}\left(n + \frac{3}{2}\right); \sec^2(c + dx)\right)}{d\left(n - \frac{1}{2}\right) \sec^{\frac{3}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^n/Sqrt[Sec[c + d*x]], x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (-1/2 + n)/2, (3/2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(-1/2 + n)*Sec[c + d*x]^(3/2))

fricas [F] time = 1.03, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(1/2), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n/sqrt(sec(d*x + c)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n/sqrt(sec(d*x + c)), x)

maple [F] time = 0.88, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x)

[Out] int((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sqrt{\sec(dx + c)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(1/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n/sqrt(sec(d*x + c)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\sqrt{\frac{1}{\cos(c+dx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(1/2),x)

[Out] int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(c + dx))^n}{\sqrt{\sec(c + dx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))**n/sec(d*x+c)**(1/2),x)

[Out] Integral((b*sec(c + d*x))**n/sqrt(sec(c + d*x)), x)

$$3.221 \quad \int \frac{(b \sec(c+dx))^n}{\sec^{\frac{3}{2}}(c+dx)} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5-2n); \frac{1}{4}(9-2n); \cos^2(c+dx)\right)}{d(5-2n)\sqrt{\sin^2(c+dx)} \sec^{\frac{5}{2}}(c+dx)}$$

[Out] -2*hypergeom([1/2, 5/4-1/2*n], [9/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(5-2*n)/sec(d*x+c)^(5/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5-2n); \frac{1}{4}(9-2n); \cos^2(c+dx)\right)}{d(5-2n)\sqrt{\sin^2(c+dx)} \sec^{\frac{5}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^n/Sec[c + d*x]^(3/2), x]

[Out] (-2*Hypergeometric2F1[1/2, (5 - 2*n)/4, (9 - 2*n)/4, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(5 - 2*n)*Sec[c + d*x]^(5/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_.), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx &= (\sec^{-n}(c + dx)(b \sec(c + dx))^n) \int \sec^{-\frac{3}{2}+n}(c + dx) dx \\ &= \left(\cos^{\frac{1}{2}+n}(c + dx) \sqrt{\sec(c + dx)} (b \sec(c + dx))^n \right) \int \cos^{\frac{3}{2}-n}(c + dx) dx \\ &= \frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{4}(5 - 2n); \frac{1}{4}(9 - 2n); \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(5 - 2n) \sec^{\frac{5}{2}}(c + dx) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.14, size = 81, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx) (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n - \frac{3}{2}\right); \frac{1}{2}\left(n + \frac{1}{2}\right); \sec^2(c + dx)\right)}{d\left(n - \frac{3}{2}\right) \sec^{\frac{5}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^n/Sec[c + d*x]^(3/2), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (-3/2 + n)/2, (1/2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*sqrt[-Tan[c + d*x]^2])/(d*(-3/2 + n)*Sec[c + d*x]^(5/2))

fricas [F] time = 0.75, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{3}{2}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(3/2), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n/sec(d*x + c)^(3/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n/sec(d*x + c)^(3/2), x)

maple [F] time = 0.97, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x)

[Out] int((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n/sec(d*x + c)^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\left(\frac{1}{\cos(c+dx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(3/2),x)

[Out] int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{3}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c)**n/sec(d*x+c)**(3/2),x)
```

```
[Out] Integral((b*sec(c + d*x)**n/sec(c + d*x)**(3/2), x)
```

$$3.222 \quad \int \frac{(b \sec(c+dx))^n}{\sec^{\frac{5}{2}}(c+dx)} dx$$

Optimal. Leaf size=80

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(7-2n); \frac{1}{4}(11-2n); \cos^2(c+dx)\right)}{d(7-2n)\sqrt{\sin^2(c+dx)} \sec^{\frac{7}{2}}(c+dx)}$$

[Out] -2*hypergeom([1/2, 7/4-1/2*n], [11/4-1/2*n], cos(d*x+c)^2)*(b*sec(d*x+c))^n*sin(d*x+c)/d/(7-2*n)/sec(d*x+c)^(7/2)/(sin(d*x+c)^2)^(1/2)

Rubi [A] time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {20, 3772, 2643}

$$\frac{2 \sin(c+dx)(b \sec(c+dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{4}(7-2n); \frac{1}{4}(11-2n); \cos^2(c+dx)\right)}{d(7-2n)\sqrt{\sin^2(c+dx)} \sec^{\frac{7}{2}}(c+dx)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[c + d*x])^n/Sec[c + d*x]^(5/2), x]

[Out] (-2*Hypergeometric2F1[1/2, (7 - 2*n)/4, (11 - 2*n)/4, Cos[c + d*x]^2]*(b*Sec[c + d*x])^n*Sin[c + d*x])/(d*(7 - 2*n)*Sec[c + d*x]^(7/2)*Sqrt[Sin[c + d*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_)*((b_.)*(v_))^(n_), x_Symbol] := Dist[(b^IntPart[n]*(b*v)^FracPart[n])/(a^IntPart[n]*(a*v)^FracPart[n]), Int[u*(a*v)^(m+n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m+n]

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(Cos[c + d*x]*(b*Sin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; Fr

eeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx &= (\sec^{-n}(c + dx)(b \sec(c + dx))^n) \int \sec^{-\frac{5}{2}+n}(c + dx) dx \\ &= \left(\cos^{\frac{1}{2}+n}(c + dx) \sqrt{\sec(c + dx)} (b \sec(c + dx))^n \right) \int \cos^{\frac{5}{2}-n}(c + dx) dx \\ &= -\frac{{}_2F_1\left(\frac{1}{2}, \frac{1}{4}(7-2n); \frac{1}{4}(11-2n); \cos^2(c + dx)\right) (b \sec(c + dx))^n \sin(c + dx)}{d(7-2n) \sec^{\frac{7}{2}}(c + dx) \sqrt{\sin^2(c + dx)}} \end{aligned}$$

Mathematica [A] time = 0.17, size = 81, normalized size = 1.01

$$\frac{\sqrt{-\tan^2(c + dx)} \csc(c + dx) (b \sec(c + dx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\left(n - \frac{5}{2}\right); \frac{1}{2}\left(n - \frac{1}{2}\right); \sec^2(c + dx)\right)}{d\left(n - \frac{5}{2}\right) \sec^{\frac{7}{2}}(c + dx)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[c + d*x])^n/Sec[c + d*x]^(5/2), x]

[Out] (Csc[c + d*x]*Hypergeometric2F1[1/2, (-5/2 + n)/2, (-1/2 + n)/2, Sec[c + d*x]^2]*(b*Sec[c + d*x])^n*Sqrt[-Tan[c + d*x]^2])/(d*(-5/2 + n)*Sec[c + d*x]^(7/2))

fricas [F] time = 1.00, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{5}{2}}}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(5/2), x, algorithm="fricas")

[Out] integral((b*sec(d*x + c))^n/sec(d*x + c)^(5/2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(d*x + c))^n/sec(d*x + c)^(5/2), x)

maple [F] time = 0.99, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x)

[Out] int((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(dx + c))^n}{\sec(dx + c)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(d*x+c))^n/sec(d*x+c)^(5/2),x, algorithm="maxima")

[Out] integrate((b*sec(d*x + c))^n/sec(d*x + c)^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(c+dx)}\right)^n}{\left(\frac{1}{\cos(c+dx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(5/2),x)

[Out] int((b/cos(c + d*x))^n/(1/cos(c + d*x))^(5/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(c + dx))^n}{\sec^{\frac{5}{2}}(c + dx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(d*x+c))**n/sec(d*x+c)**(5/2),x)
```

```
[Out] Integral((b*sec(c + d*x))**n/sec(c + d*x)**(5/2), x)
```

3.223 $\int (d \sec(a + bx))^{7/2} \sin(a + bx) dx$

Optimal. Leaf size=20

$$\frac{2d(d \sec(a + bx))^{5/2}}{5b}$$

[Out] $2/5*d*(d*\sec(b*x+a))^(5/2)/b$

Rubi [A] time = 0.04, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2622, 30}

$$\frac{2d(d \sec(a + bx))^{5/2}}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Sec}[a + b*x])^(7/2)*\text{Sin}[a + b*x], x]$

[Out] $(2*d*(d*\text{Sec}[a + b*x])^(5/2))/(5*b)$

Rule 30

$\text{Int}[(x_)^(m_.), x_Symbol] \rightarrow \text{Simp}[x^(m + 1)/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2622

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*\sec[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2)], x], x, a*\text{Sec}[e + f*x]], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n + 1)/2] \ \&\& \ !(\text{IntegerQ}[(m + 1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (d \sec(a + bx))^{7/2} \sin(a + bx) dx &= \frac{d \text{Subst}\left(\int x^{3/2} dx, x, d \sec(a + bx)\right)}{b} \\ &= \frac{2d(d \sec(a + bx))^{5/2}}{5b} \end{aligned}$$

Mathematica [A] time = 0.06, size = 20, normalized size = 1.00

$$\frac{2d(d \sec(a + bx))^{5/2}}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(7/2)*Sin[a + b*x],x]

[Out] (2*d*(d*Sec[a + b*x])^(5/2))/(5*b)

fricas [A] time = 0.88, size = 28, normalized size = 1.40

$$\frac{2d^3 \sqrt{\frac{d}{\cos(bx+a)}}}{5b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(7/2)*sin(b*x+a),x, algorithm="fricas")

[Out] 2/5*d^3*sqrt(d/cos(b*x + a))/(b*cos(b*x + a)^2)

giac [B] time = 1.65, size = 154, normalized size = 7.70

$$\frac{4 \left(5 \left(\sqrt{-d} \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - \sqrt{-d \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + d} \right)^4 d - 10 \left(\sqrt{-d} \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - \sqrt{-d \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + d} \right)^5 \right)}{5 \left(\sqrt{-d} \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - \sqrt{-d \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + d} - \sqrt{-d} \right)^5} b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(7/2)*sin(b*x+a),x, algorithm="giac")

[Out] -4/5*(5*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))^4*d - 10*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))^2*d^2 + d^3)*d^3*sgn(cos(b*x + a))/((sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d) - sqrt(-d))^5*b)

maple [A] time = 0.14, size = 17, normalized size = 0.85

$$\frac{2d(d \sec(bx+a))^{\frac{5}{2}}}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(7/2)*sin(b*x+a),x)

[Out] 2/5*d*(d*sec(b*x+a))^(5/2)/b

maxima [A] time = 0.72, size = 23, normalized size = 1.15

$$\frac{2 \left(\frac{d}{\cos(bx+a)} \right)^{\frac{7}{2}} \cos(bx+a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(7/2)*sin(b*x+a),x, algorithm="maxima")

[Out] 2/5*(d/cos(b*x + a))^(7/2)*cos(b*x + a)/b

mupad [B] time = 1.59, size = 77, normalized size = 3.85

$$\frac{8d^3 \sqrt{\frac{d}{\cos(a+bx)}} (4 \cos(2a + 2bx) + \cos(4a + 4bx) + 3)}{5b (15 \cos(2a + 2bx) + 6 \cos(4a + 4bx) + \cos(6a + 6bx) + 10)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d/cos(a + b*x))^(7/2),x)

[Out] (8*d^3*(d/cos(a + b*x))^(1/2)*(4*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 3))/
(5*b*(15*cos(2*a + 2*b*x) + 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) + 10))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))**(7/2)*sin(b*x+a),x)

[Out] Timed out

3.224 $\int (d \sec(a + bx))^{5/2} \sin(a + bx) dx$

Optimal. Leaf size=20

$$\frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

[Out] $2/3*d*(d*\sec(b*x+a))^(3/2)/b$

Rubi [A] time = 0.04, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2622, 30}

$$\frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Sec}[a + b*x])^(5/2)*\text{Sin}[a + b*x], x]$

[Out] $(2*d*(d*\text{Sec}[a + b*x])^(3/2))/(3*b)$

Rule 30

$\text{Int}[(x_)^(m_), x_Symbol] \rightarrow \text{Simp}[x^(m + 1)/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2622

$\text{Int}[\text{csc}[(e_) + (f_)*(x_)]^(n_)*((a_)*\text{sec}[(e_) + (f_)*(x_)])^(m_), x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n + 1)/2] \ \&\& \ !(\text{IntegerQ}[(m + 1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (d \sec(a + bx))^{5/2} \sin(a + bx) dx &= \frac{d \text{Subst}\left(\int \sqrt{x} dx, x, d \sec(a + bx)\right)}{b} \\ &= \frac{2d(d \sec(a + bx))^{3/2}}{3b} \end{aligned}$$

Mathematica [A] time = 0.04, size = 20, normalized size = 1.00

$$\frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x],x]

[Out] (2*d*(d*Sec[a + b*x])^(3/2))/(3*b)

fricas [A] time = 0.76, size = 28, normalized size = 1.40

$$\frac{2d^2\sqrt{\frac{d}{\cos(bx+a)}}}{3b\cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a),x, algorithm="fricas")

[Out] 2/3*d^2*sqrt(d/cos(b*x + a))/(b*cos(b*x + a))

giac [B] time = 1.59, size = 111, normalized size = 5.55

$$\frac{4\left(3\left(\sqrt{-d}\tan\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-\sqrt{-d\tan\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4+d}\right)^2d-d^2\right)d^2\operatorname{sgn}(\cos(bx+a))}{3\left(\sqrt{-d}\tan\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-\sqrt{-d\tan\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4+d}-\sqrt{-d}\right)^3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a),x, algorithm="giac")

[Out] 4/3*(3*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))^2*d - d^2)*d^2*sgn(cos(b*x + a))/((sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d) - sqrt(-d))^3*b)

maple [A] time = 0.12, size = 17, normalized size = 0.85

$$\frac{2d(d\sec(bx+a))^{\frac{3}{2}}}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(5/2)*sin(b*x+a),x)

[Out] 2/3*d*(d*sec(b*x+a))^(3/2)/b

maxima [A] time = 0.39, size = 23, normalized size = 1.15

$$\frac{2 \left(\frac{d}{\cos(bx+a)} \right)^{\frac{5}{2}} \cos(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a),x, algorithm="maxima")

[Out] 2/3*(d/cos(b*x + a))^(5/2)*cos(b*x + a)/b

mupad [B] time = 0.25, size = 39, normalized size = 1.95

$$\frac{4d^2 \cos(a+bx) \sqrt{\frac{d}{\cos(a+bx)}}}{3b(\cos(2a+2bx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d/cos(a + b*x))^(5/2),x)

[Out] (4*d^2*cos(a + b*x)*(d/cos(a + b*x))^(1/2))/(3*b*(cos(2*a + 2*b*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))**(5/2)*sin(b*x+a),x)

[Out] Timed out

3.225 $\int (d \sec(a + bx))^{3/2} \sin(a + bx) dx$

Optimal. Leaf size=18

$$\frac{2d\sqrt{d \sec(a + bx)}}{b}$$

[Out] 2*d*(d*sec(b*x+a))^(1/2)/b

Rubi [A] time = 0.04, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2622, 30}

$$\frac{2d\sqrt{d \sec(a + bx)}}{b}$$

Antiderivative was successfully verified.

[In] Int[(d*Sec[a + b*x])^(3/2)*Sin[a + b*x],x]

[Out] (2*d*Sqrt[d*Sec[a + b*x]])/b

Rule 30

Int[(x_)^(m_.), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2622

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int (d \sec(a + bx))^{3/2} \sin(a + bx) dx &= \frac{d \operatorname{Subst}\left(\int \frac{1}{\sqrt{x}} dx, x, d \sec(a + bx)\right)}{b} \\ &= \frac{2d\sqrt{d \sec(a + bx)}}{b} \end{aligned}$$

Mathematica [A] time = 0.03, size = 18, normalized size = 1.00

$$\frac{2d\sqrt{d \sec(a + bx)}}{b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(3/2)*Sin[a + b*x],x]

[Out] (2*d*Sqrt[d*Sec[a + b*x]])/b

fricas [A] time = 0.96, size = 18, normalized size = 1.00

$$\frac{2d\sqrt{\frac{d}{\cos(bx+a)}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="fricas")

[Out] 2*d*sqrt(d/cos(b*x + a))/b

giac [B] time = 1.49, size = 62, normalized size = 3.44

$$-\frac{4d^2\operatorname{sgn}(\cos(bx+a))}{\left(\sqrt{-d}\tan\left(\frac{1}{2}bx+\frac{1}{2}a\right)^2-\sqrt{-d}\tan\left(\frac{1}{2}bx+\frac{1}{2}a\right)^4+d-\sqrt{-d}\right)b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="giac")

[Out] -4*d^2*sgn(cos(b*x + a))/((sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d) - sqrt(-d))*b)

maple [A] time = 0.12, size = 17, normalized size = 0.94

$$\frac{2d\sqrt{d}\sec(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(3/2)*sin(b*x+a),x)

[Out] 2*d*(d*sec(b*x+a))^(1/2)/b

maxima [A] time = 0.61, size = 23, normalized size = 1.28

$$\frac{2\left(\frac{d}{\cos(bx+a)}\right)^{\frac{3}{2}}\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="maxima")

[Out] 2*(d/cos(b*x + a))^(3/2)*cos(b*x + a)/b

mupad [B] time = 0.10, size = 18, normalized size = 1.00

$$\frac{2d \sqrt{\frac{d}{\cos(a+bx)}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d/cos(a + b*x))^(3/2),x)

[Out] (2*d*(d/cos(a + b*x))^(1/2))/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \sec(a + bx))^{\frac{3}{2}} \sin(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))**(3/2)*sin(b*x+a),x)

[Out] Integral((d*sec(a + b*x))**(3/2)*sin(a + b*x), x)

3.226 $\int \sqrt{d \sec(a + bx)} \sin(a + bx) dx$

Optimal. Leaf size=18

$$-\frac{2d}{b\sqrt{d \sec(a + bx)}}$$

[Out] $-2*d/b/(d*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.03, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2622, 30}

$$-\frac{2d}{b\sqrt{d \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[d*\text{Sec}[a + b*x]]*\text{Sin}[a + b*x], x]$

[Out] $(-2*d)/(b*\text{Sqrt}[d*\text{Sec}[a + b*x]])$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] := \text{Simp}[x^{(m + 1)}/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2622

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^{(n_.)}*((a_.)*\text{sec}[(e_.) + (f_.)*(x_)])^{(m_.)}, x_Symbol] := \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m + n - 1)}/(-1 + x^2/a^2)^{((n + 1)/2)}, x], x, a*\text{Sec}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n + 1)/2] \ \&\& \ !(\text{IntegerQ}[(m + 1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int \sqrt{d \sec(a + bx)} \sin(a + bx) dx &= \frac{d \text{Subst}\left(\int \frac{1}{x^{3/2}} dx, x, d \sec(a + bx)\right)}{b} \\ &= -\frac{2d}{b\sqrt{d \sec(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.04, size = 18, normalized size = 1.00

$$-\frac{2d}{b\sqrt{d \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Sec[a + b*x]]*Sin[a + b*x],x]

[Out] (-2*d)/(b*Sqrt[d*Sec[a + b*x]])

fricas [A] time = 0.64, size = 23, normalized size = 1.28

$$-\frac{2\sqrt{\frac{d}{\cos(bx+a)}}\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="fricas")

[Out] -2*sqrt(d/cos(b*x + a))*cos(b*x + a)/b

giac [A] time = 0.32, size = 22, normalized size = 1.22

$$-\frac{2\sqrt{d\cos(bx+a)}\operatorname{sgn}(\cos(bx+a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="giac")

[Out] -2*sqrt(d*cos(b*x + a))*sgn(cos(b*x + a))/b

maple [A] time = 0.19, size = 17, normalized size = 0.94

$$-\frac{2d}{b\sqrt{d}\sec(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(1/2)*sin(b*x+a),x)

[Out] -2*d/b/(d*sec(b*x+a))^(1/2)

maxima [A] time = 0.47, size = 23, normalized size = 1.28

$$-\frac{2\sqrt{\frac{d}{\cos(bx+a)}}\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="maxima")

[Out] -2*sqrt(d/cos(b*x + a))*cos(b*x + a)/b

mupad [B] time = 0.23, size = 23, normalized size = 1.28

$$-\frac{2 \cos(a + b x) \sqrt{\frac{d}{\cos(a + b x)}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d/cos(a + b*x))^(1/2),x)

[Out] -(2*cos(a + b*x)*(d/cos(a + b*x))^(1/2))/b

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \sec(a + b x)} \sin(a + b x) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))**(1/2)*sin(b*x+a),x)

[Out] Integral(sqrt(d*sec(a + b*x))*sin(a + b*x), x)

$$3.227 \quad \int \frac{\sin(a+bx)}{\sqrt{d \sec(a+bx)}} dx$$

Optimal. Leaf size=20

$$-\frac{2d}{3b(d \sec(a + bx))^{3/2}}$$

[Out] $-2/3*d/b/(d*\sec(b*x+a))^{(3/2)}$

Rubi [A] time = 0.03, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2622, 30}

$$-\frac{2d}{3b(d \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]/Sqrt[d*Sec[a + b*x]],x]

[Out] $(-2*d)/(3*b*(d*Sec[a + b*x])^{(3/2)})$

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2622

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx &= \frac{d \operatorname{Subst}\left(\int \frac{1}{x^{5/2}} dx, x, d \sec(a + bx)\right)}{b} \\ &= -\frac{2d}{3b(d \sec(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A] time = 0.05, size = 20, normalized size = 1.00

$$-\frac{2d}{3b(d \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]/Sqrt[d*Sec[a + b*x]], x]

[Out] (-2*d)/(3*b*(d*Sec[a + b*x])^(3/2))

fricas [A] time = 0.73, size = 28, normalized size = 1.40

$$-\frac{2 \sqrt{\frac{d}{\cos(bx+a)}} \cos(bx+a)^2}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*sec(b*x+a))^(1/2), x, algorithm="fricas")

[Out] -2/3*sqrt(d/cos(b*x + a))*cos(b*x + a)^2/(b*d)

giac [B] time = 1.76, size = 35, normalized size = 1.75

$$-\frac{2 \sqrt{d \cos(bx+a)} |b| \cos(bx+a) \operatorname{sgn}(b) \operatorname{sgn}(\cos(bx+a))}{3b^2d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*sec(b*x+a))^(1/2), x, algorithm="giac")

[Out] -2/3*sqrt(d*cos(b*x + a))*abs(b)*cos(b*x + a)*sgn(b)*sgn(cos(b*x + a))/(b^2*d)

maple [A] time = 0.15, size = 17, normalized size = 0.85

$$-\frac{2d}{3b(d \sec(bx + a))^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)/(d*sec(b*x+a))^(1/2), x)

[Out] -2/3*d/b/(d*sec(b*x+a))^(3/2)

maxima [A] time = 0.38, size = 23, normalized size = 1.15

$$-\frac{2 \cos(bx + a)}{3b \sqrt{\frac{d}{\cos(bx+a)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] -2/3*cos(b*x + a)/(b*sqrt(d/cos(b*x + a)))

mupad [B] time = 0.24, size = 28, normalized size = 1.40

$$-\frac{2 \cos(a + bx)^2 \sqrt{\frac{d}{\cos(a+bx)}}}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/(d/cos(a + b*x))^(1/2),x)

[Out] -(2*cos(a + b*x)^2*(d/cos(a + b*x))^(1/2))/(3*b*d)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin(a + bx)}{\sqrt{d \sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*sec(b*x+a))**(1/2),x)

[Out] Integral(sin(a + b*x)/sqrt(d*sec(a + b*x)), x)

3.228 $\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx$

Optimal. Leaf size=41

$$\frac{2d^3}{b\sqrt{d \sec(a + bx)}} + \frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

[Out] $2/3*d*(d*\sec(b*x+a))^(3/2)/b+2*d^3/b/(d*\sec(b*x+a))^(1/2)$

Rubi [A] time = 0.05, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2622, 14}

$$\frac{2d^3}{b\sqrt{d \sec(a + bx)}} + \frac{2d(d \sec(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] Int[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x]^3,x]

[Out] (2*d^3)/(b*Sqrt[d*Sec[a + b*x]]) + (2*d*(d*Sec[a + b*x])^(3/2))/(3*b)

Rule 14

Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2622

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
\int (d \sec(a + bx))^{5/2} \sin^3(a + bx) dx &= \frac{d^3 \operatorname{Subst} \left(\int \frac{-1 + \frac{x^2}{d^2}}{x^{3/2}} dx, x, d \sec(a + bx) \right)}{b} \\
&= \frac{d^3 \operatorname{Subst} \left(\int \left(-\frac{1}{x^{3/2}} + \frac{\sqrt{x}}{d^2} \right) dx, x, d \sec(a + bx) \right)}{b} \\
&= \frac{2d^3}{b\sqrt{d \sec(a + bx)}} + \frac{2d(d \sec(a + bx))^{3/2}}{3b}
\end{aligned}$$

Mathematica [A] time = 0.22, size = 32, normalized size = 0.78

$$\frac{d(3 \cos(2(a + bx)) + 5)(d \sec(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(5/2)*Sin[a + b*x]^3,x]

[Out] (d*(5 + 3*Cos[2*(a + b*x)])*(d*Sec[a + b*x])^(3/2))/(3*b)

fricas [A] time = 0.80, size = 42, normalized size = 1.02

$$\frac{2 \left(3 d^2 \cos(bx + a)^2 + d^2 \right) \sqrt{\frac{d}{\cos(bx+a)}}}{3 b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x, algorithm="fricas")

[Out] 2/3*(3*d^2*cos(b*x + a)^2 + d^2)*sqrt(d/cos(b*x + a))/(b*cos(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \sec(bx + a))^{5/2} \sin(bx + a)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x, algorithm="giac")

[Out] integrate((d*sec(b*x + a))^(5/2)*sin(b*x + a)^3, x)

maple [B] time = 0.91, size = 357, normalized size = 8.71

$$(-1 + \cos(bx + a)) \left(12 (\cos^3(bx + a)) \sqrt{-\frac{\cos(bx+a)}{(\cos(bx+a)+1)^2}} + 12 (\cos^2(bx + a)) \sqrt{-\frac{\cos(bx+a)}{(\cos(bx+a)+1)^2}} + 3 \ln \left(-\frac{2(\cos^2(bx+a) + \cos(bx+a) + 1)}{\cos(bx+a) + 1} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x)

[Out]
$$-1/6/b*(-1+\cos(b*x+a))*(12*\cos(b*x+a)^3*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}+12*\cos(b*x+a)^2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}+3*\ln(-2*(2*\cos(b*x+a))^2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}-\cos(b*x+a)^2+2*\cos(b*x+a)-2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}-1)/\sin(b*x+a)^2*\cos(b*x+a)^2-3*\ln(-(2*\cos(b*x+a))^2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}-\cos(b*x+a)^2+2*\cos(b*x+a)-2*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}-1)/\sin(b*x+a)^2*\cos(b*x+a)^2+4*\cos(b*x+a)*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}+4*(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}*\cos(b*x+a)*(d/\cos(b*x+a))^{(5/2)}/(-\cos(b*x+a)/(\cos(b*x+a)+1)^2)^{(1/2)}/\sin(b*x+a)^2$$

maxima [A] time = 0.64, size = 36, normalized size = 0.88

$$\frac{2 \left(\frac{3d^2}{\sqrt{\frac{d}{\cos(bx+a)}}} + \left(\frac{d}{\cos(bx+a)} \right)^{\frac{3}{2}} \right) d}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*sin(b*x+a)^3,x, algorithm="maxima")

[Out]
$$2/3*(3*d^2/\sqrt{d/\cos(b*x+a)} + (d/\cos(b*x+a))^{(3/2)})*d/b$$

mupad [B] time = 0.59, size = 50, normalized size = 1.22

$$\frac{d^2 \sqrt{\frac{d}{\cos(a+bx)}} \left(\frac{13 \cos(a+bx)}{3} + \cos(3a + 3bx) \right)}{b (\cos(2a + 2bx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3*(d/cos(a + b*x))^(5/2),x)

```
[Out] (d^2*(d/cos(a + b*x))^(1/2)*((13*cos(a + b*x))/3 + cos(3*a + 3*b*x)))/(b*(c  
os(2*a + 2*b*x) + 1))
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*sec(b*x+a))**(5/2)*sin(b*x+a)**3,x)
```

```
[Out] Timed out
```

3.229 $\int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx$

Optimal. Leaf size=43

$$\frac{2d(d \sec(a + bx))^{7/2}}{7b} - \frac{2d^3(d \sec(a + bx))^{3/2}}{3b}$$

[Out] $-2/3*d^3*(d*\sec(b*x+a))^{(3/2)}/b+2/7*d*(d*\sec(b*x+a))^{(7/2)}/b$

Rubi [A] time = 0.05, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2622, 14}

$$\frac{2d(d \sec(a + bx))^{7/2}}{7b} - \frac{2d^3(d \sec(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Sec}[a + b*x])^{(9/2)}*\text{Sin}[a + b*x]^3, x]$

[Out] $(-2*d^3*(d*\text{Sec}[a + b*x])^{(3/2)})/(3*b) + (2*d*(d*\text{Sec}[a + b*x])^{(7/2)})/(7*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{InverseFunctionQ}[v]$

Rule 2622

$\text{Int}[\text{csc}[(e_.) + (f_)*(x_)]^{(n_)}*((a_)*\sec[(e_.) + (f_)*(x_)])^{(m_)}, x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1+x^2/a^2)^{(n+1)/2}], x], x, a*\text{Sec}[e+f*x], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n+1)/2] \ \&\& \ !(\text{IntegerQ}[(m+1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (d \sec(a + bx))^{9/2} \sin^3(a + bx) dx &= \frac{d^3 \text{Subst}\left(\int \sqrt{x} \left(-1 + \frac{x^2}{d^2}\right) dx, x, d \sec(a + bx)\right)}{b} \\ &= \frac{d^3 \text{Subst}\left(\int \left(-\sqrt{x} + \frac{x^{5/2}}{d^2}\right) dx, x, d \sec(a + bx)\right)}{b} \\ &= -\frac{2d^3(d \sec(a + bx))^{3/2}}{3b} + \frac{2d(d \sec(a + bx))^{7/2}}{7b} \end{aligned}$$

Mathematica [A] time = 0.11, size = 42, normalized size = 0.98

$$\frac{d^4(7 \cos(2(a + bx)) + 1) \sec^3(a + bx) \sqrt{d} \sec(a + bx)}{21b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(9/2)*Sin[a + b*x]^3,x]

[Out] -1/21*(d^4*(1 + 7*Cos[2*(a + b*x)])*Sec[a + b*x]^3*Sqrt[d*Sec[a + b*x]])/b

fricas [A] time = 0.91, size = 44, normalized size = 1.02

$$\frac{2(7d^4 \cos(bx + a)^2 - 3d^4) \sqrt{\frac{d}{\cos(bx+a)}}}{21b \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x, algorithm="fricas")

[Out] -2/21*(7*d^4*cos(b*x + a)^2 - 3*d^4)*sqrt(d/cos(b*x + a))/(b*cos(b*x + a)^3)

giac [B] time = 1.83, size = 257, normalized size = 5.98

$$\frac{16 \left(21 \left(\sqrt{-d} \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - \sqrt{-d \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + d} \right)^5 \sqrt{-d} d + 7 \left(\sqrt{-d} \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - \sqrt{-d \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^4 + d} \right)^5 \right)}{21^5 b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x, algorithm="giac")

[Out] 16/21*(21*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))^5*sqrt(-d)*d + 7*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))^4*d^2 - 28*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))^3*sqrt(-d)*d^2 + 7*(sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d))*sqrt(-d)*d^3 + d^4)*d^4*sgn(cos(b*x + a))/((sqrt(-d)*tan(1/2*b*x + 1/2*a)^2 - sqrt(-d*tan(1/2*b*x + 1/2*a)^4 + d) - sqrt(-d))^7*b)

maple [A] time = 0.68, size = 36, normalized size = 0.84

$$\frac{2(7(\cos^2(bx + a)) - 3) \cos(bx + a) \left(\frac{d}{\cos(bx+a)}\right)^{\frac{9}{2}}}{21b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x)`

[Out] $-2/21/b*(7*\cos(b*x+a)^2-3)*\cos(b*x+a)*(d/\cos(b*x+a))^{9/2}$

maxima [A] time = 0.34, size = 38, normalized size = 0.88

$$\frac{2 \left(7 d^2 \left(\frac{d}{\cos(bx+a)} \right)^{\frac{3}{2}} - 3 \left(\frac{d}{\cos(bx+a)} \right)^{\frac{7}{2}} \right) d}{21 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*sec(b*x+a))^(9/2)*sin(b*x+a)^3,x, algorithm="maxima")`

[Out] $-2/21*(7*d^2*(d/\cos(b*x + a))^{3/2} - 3*(d/\cos(b*x + a))^{7/2})*d/b$

mupad [B] time = 4.34, size = 95, normalized size = 2.21

$$\frac{4 d^4 e^{a 1 i + b x 1 i} \sqrt{\frac{d}{\frac{e^{-a 1 i - b x 1 i}}{2} + \frac{e^{a 1 i + b x 1 i}}{2}}} (2 e^{a 2 i + b x 2 i} + 7 e^{a 4 i + b x 4 i} + 7)}{21 b (e^{a 2 i + b x 2 i} + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^3*(d/cos(a + b*x))^(9/2),x)`

[Out] $-(4*d^4*\exp(a*1i + b*x*1i)*(d/(\exp(- a*1i - b*x*1i)/2 + \exp(a*1i + b*x*1i)/2))^{1/2}*(2*\exp(a*2i + b*x*2i) + 7*\exp(a*4i + b*x*4i) + 7))/(21*b*(\exp(a*2i + b*x*2i) + 1)^3)$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*sec(b*x+a))**(9/2)*sin(b*x+a)**3,x)`

[Out] Timed out

3.230 $\int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx$

Optimal. Leaf size=128

$$\frac{4d^4 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{7b} - \frac{4cd^3 (d \csc(a + bx))^{3/2}}{7b \sqrt{c \sec(a + bx)}} - \frac{2cd (d \csc(a + bx))^{7/2}}{7b \sqrt{c \sec(a + bx)}}$$

[Out] $-4/7 * c * d^3 * (d * \csc(b * x + a))^{(3/2)} / b / (c * \sec(b * x + a))^{(1/2)} - 2/7 * c * d * (d * \csc(b * x + a))^{(7/2)} / b / (c * \sec(b * x + a))^{(1/2)} - 4/7 * d^4 * (\sin(a + 1/4 * \pi + b * x)^2)^{(1/2)} / \sin(a + 1/4 * \pi + b * x) * \text{EllipticF}(\cos(a + 1/4 * \pi + b * x), 2^{(1/2)}) * (d * \csc(b * x + a))^{(1/2)} * (c * \sec(b * x + a))^{(1/2)} * \sin(2 * b * x + 2 * a)^{(1/2)} / b$

Rubi [A] time = 0.20, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2625, 2630, 2573, 2641}

$$-\frac{4cd^3 (d \csc(a + bx))^{3/2}}{7b \sqrt{c \sec(a + bx)}} + \frac{4d^4 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{7b} - \frac{2cd (d \csc(a + bx))^{7/2}}{7b \sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d * \text{Csc}[a + b * x])^{(9/2)} * \text{Sqrt}[c * \text{Sec}[a + b * x]], x]$

[Out] $(-4 * c * d^3 * (d * \text{Csc}[a + b * x])^{(3/2)}) / (7 * b * \text{Sqrt}[c * \text{Sec}[a + b * x]]) - (2 * c * d * (d * \text{Csc}[a + b * x])^{(7/2)}) / (7 * b * \text{Sqrt}[c * \text{Sec}[a + b * x]]) + (4 * d^4 * \text{Sqrt}[d * \text{Csc}[a + b * x]] * \text{EllipticF}[a - \pi/4 + b * x, 2] * \text{Sqrt}[c * \text{Sec}[a + b * x]] * \text{Sqrt}[\sin[2 * a + 2 * b * x]]) / (7 * b)$

Rule 2573

$\text{Int}[1 / (\text{Sqrt}[\cos[(e_.) + (f_.) * (x_.)] * (b_.)] * \text{Sqrt}[(a_.) * \sin[(e_.) + (f_.) * (x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\sin[2 * e + 2 * f * x]] / (\text{Sqrt}[a * \sin[e + f * x]] * \text{Sqrt}[b * \cos[e + f * x]]), \text{Int}[1 / \text{Sqrt}[\sin[2 * e + 2 * f * x]], x], x] /;$ FreeQ[{a, b, e, f}, x]

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.) * (x_.)] * (a_.))^{(m)} * ((b_.) * \sec[(e_.) + (f_.) * (x_.)])^{(n)}], x_Symbol] \rightarrow -\text{Simp}[(a * b * (a * \csc[e + f * x])^{(m - 1)} * (b * \sec[e + f * x])^{(n - 1)}) / (f * (m - 1)), x] + \text{Dist}[(a^{2 * (m + n - 2)}) / (m - 1), \text{Int}[(a * \csc[e + f * x])^{(m - 2)} * (b * \sec[e + f * x])^n, x], x] /;$ FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2 * m, 2 * n] && !GtQ[n, m]

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2641

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int (d \csc(a + bx))^{9/2} \sqrt{c \sec(a + bx)} dx &= -\frac{2cd(d \csc(a + bx))^{7/2}}{7b\sqrt{c \sec(a + bx)}} + \frac{1}{7} (6d^2) \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx \\ &= -\frac{4cd^3(d \csc(a + bx))^{3/2}}{7b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b\sqrt{c \sec(a + bx)}} + \frac{1}{7} (4d^4) \int \sqrt{d \csc(a + bx)} dx \\ &= -\frac{4cd^3(d \csc(a + bx))^{3/2}}{7b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b\sqrt{c \sec(a + bx)}} + \frac{1}{7} (4d^4 \sqrt{c \cos(a + bx)}) \\ &= -\frac{4cd^3(d \csc(a + bx))^{3/2}}{7b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b\sqrt{c \sec(a + bx)}} + \frac{1}{7} (4d^4 \sqrt{d \csc(a + bx)}) \\ &= -\frac{4cd^3(d \csc(a + bx))^{3/2}}{7b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b\sqrt{c \sec(a + bx)}} + \frac{4d^4 \sqrt{d \csc(a + bx)} F}{7b \left(\csc^2(a + bx) - 2 \right)} \end{aligned}$$

Mathematica [C] time = 1.51, size = 122, normalized size = 0.95

$$\frac{2d^4 \cos(2(a + bx)) \cot(a + bx) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} \left((\cos(2(a + bx)) - 2) \csc^4(a + bx) - 2(-\cot^2(a + bx)) \right)}{7b \left(\csc^2(a + bx) - 2 \right)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Csc[a + b*x])^(9/2)*Sqrt[c*Sec[a + b*x]],x]
```

```
[Out] (2*d^4*Cos[2*(a + b*x)]*Cot[a + b*x]*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*((-2 + Cos[2*(a + b*x)])*Csc[a + b*x]^4 - 2*(-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2]*Sec[a + b*x]^2))/(7*b*(-2 + Csc[a + b*x]^2))
```

fricas [F] time = 0.92, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)} d^4 \csc(bx+a)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d^4*csc(b*x + a)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx+a))^{\frac{9}{2}} \sqrt{c \sec(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(9/2)*sqrt(c*sec(b*x + a)), x)

maple [B] time = 1.28, size = 534, normalized size = 4.17

$$\left(4 \sin(bx+a) \left(\cos^3(bx+a)\right) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \text{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x)

[Out]
$$\begin{aligned} & -1/7/b*(4*\sin(b*x+a)*\cos(b*x+a)^3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2*2^{(1/2)})) \\ & +4*\sin(b*x+a)*\cos(b*x+a)^2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2*2^{(1/2)}) \\ & -4*\sin(b*x+a)*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2*2^{(1/2)}) \\ & -4*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2*2^{(1/2)}) \\ & -2*\cos(b*x+a)^3*2^{(1/2)}+3*\cos(b*x+a)*2^{(1/2)}*(d/\sin(b*x+a))^{(9/2)}*(c/\cos(b*x+a))^{(1/2)}*\sin(b*x+a)*2^{(1/2)} \end{aligned}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc (bx + a))^{\frac{9}{2}} \sqrt{c \sec (bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(9/2)*sqrt(c*sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{\frac{c}{\cos (a + b x)}} \left(\frac{d}{\sin (a + b x)} \right)^{9/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(9/2),x)

[Out] int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(9/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(9/2)*(c*sec(b*x+a))**(1/2),x)

[Out] Timed out

3.231 $\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx$

Optimal. Leaf size=69

$$\frac{8cd^3\sqrt{d \csc(a + bx)}}{5b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b\sqrt{c \sec(a + bx)}}$$

[Out] $-2/5*c*d*(d*\csc(b*x+a))^{(5/2)}/b/(c*\sec(b*x+a))^{(1/2)}-8/5*c*d^3*(d*\csc(b*x+a))^{(1/2)}/b/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2625, 2619}

$$\frac{8cd^3\sqrt{d \csc(a + bx)}}{5b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(7/2)*Sqrt[c*Sec[a + b*x]], x]

[Out] $(-8*c*d^3*Sqrt[d*Csc[a + b*x]])/(5*b*Sqrt[c*Sec[a + b*x]]) - (2*c*d*(d*Csc[a + b*x])^{(5/2)})/(5*b*Sqrt[c*Sec[a + b*x]])$

Rule 2619

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]

Rule 2625

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Dist[(a^2*(m + n - 2))/(m - 1), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rubi steps

$$\int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx = -\frac{2cd(d \csc(a + bx))^{5/2}}{5b\sqrt{c \sec(a + bx)}} + \frac{1}{5} (4d^2) \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx$$

$$= -\frac{8cd^3 \sqrt{d \csc(a + bx)}}{5b\sqrt{c \sec(a + bx)}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b\sqrt{c \sec(a + bx)}}$$

Mathematica [A] time = 0.12, size = 56, normalized size = 0.81

$$\frac{2d^3 \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} (4 \cos(a + bx) + \cot(a + bx) \csc(a + bx))}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(7/2)*Sqrt[c*Sec[a + b*x]],x]

[Out] (-2*d^3*Sqrt[d*Csc[a + b*x]]*(4*Cos[a + b*x] + Cot[a + b*x]*Csc[a + b*x])*Sqrt[c*Sec[a + b*x]])/(5*b)

fricas [A] time = 0.89, size = 67, normalized size = 0.97

$$\frac{2(4d^3 \cos(bx + a)^3 - 5d^3 \cos(bx + a)) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{5(b \cos(bx + a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] -2/5*(4*d^3*cos(b*x + a)^3 - 5*d^3*cos(b*x + a))*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*cos(b*x + a)^2 - b)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{7/2} \sqrt{c \sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(7/2)*sqrt(c*sec(b*x + a)), x)

maple [A] time = 1.19, size = 54, normalized size = 0.78

$$\frac{2(4(\cos^2(bx + a)) - 5) \cos(bx + a) \left(\frac{d}{\sin(bx+a)}\right)^{7/2} \sqrt{\frac{c}{\cos(bx+a)}} \sin(bx + a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x)`

[Out] $2/5/b*(4*\cos(b*x+a)^2-5)*\cos(b*x+a)*(d/\sin(b*x+a))^{7/2}*(c/\cos(b*x+a))^{1/2}*\sin(b*x+a)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{7}{2}} \sqrt{c \sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate((d*csc(b*x + a))^(7/2)*sqrt(c*sec(b*x + a)), x)`

mupad [B] time = 1.34, size = 85, normalized size = 1.23

$$\frac{4 d^3 \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (3 \cos(a+bx) - 4 \cos(3a+3bx) + \cos(5a+5bx))}{5b (\cos(4a+4bx) - 4 \cos(2a+2bx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(7/2),x)`

[Out] $-(4*d^3*(c/\cos(a + b*x))^{1/2}*(d/\sin(a + b*x))^{1/2}*(3*\cos(a + b*x) - 4*\cos(3*a + 3*b*x) + \cos(5*a + 5*b*x)))/(5*b*(\cos(4*a + 4*b*x) - 4*\cos(2*a + 2*b*x) + 3))$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))**(7/2)*(c*sec(b*x+a))**(1/2),x)`

[Out] Timed out

3.232 $\int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx$

Optimal. Leaf size=93

$$\frac{2d^2 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} - \frac{2cd(d \csc(a + bx))^{3/2}}{3b \sqrt{c \sec(a + bx)}}$$

[Out] $-2/3*c*d*(d*\csc(b*x+a))^{(3/2)}/b/(c*\sec(b*x+a))^{(1/2)}-2/3*d^2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b$

Rubi [A] time = 0.14, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2625, 2630, 2573, 2641}

$$\frac{2d^2 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} - \frac{2cd(d \csc(a + bx))^{3/2}}{3b \sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(5/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]], x]$

[Out] $(-2*c*d*(d*\text{Csc}[a + b*x])^{(3/2)})/(3*b*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (2*d^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(3*b)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(m-1)), x] + \text{Dist}[(a^2*(m+n-2))/(m-1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{IntegersQ}[2*m, 2*n] \ \&\& \ !\text{GtQ}[n, m]$

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a*\text{Csc}[e + f*x])^m*(b*\text{Sec}[e + f*x])^n*(a*\text{Sin}[e + f*x])$

```
)^m*(b*cos[e + f*x])^n, Int[1/((a*sin[e + f*x])^m*(b*cos[e + f*x])^n), x],
x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
]
```

Rule 2641

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c -
Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx &= -\frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} + \frac{1}{3} (2d^2) \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx \\
 &= -\frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} + \frac{1}{3} (2d^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}) \\
 &= -\frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} + \frac{1}{3} (2d^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}) \\
 &= -\frac{2cd(d \csc(a + bx))^{3/2}}{3b\sqrt{c \sec(a + bx)}} + \frac{2d^2 \sqrt{d \csc(a + bx)} F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a + bx)}}{3b}
 \end{aligned}$$

Mathematica [C] time = 0.89, size = 109, normalized size = 1.17

$$\frac{d(\cos(a + bx) + \cos(3(a + bx))) \sec^2(a + bx) \sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2} \left((-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; c \sec^2(a + bx)\right) \right)}{3b(\csc^2(a + bx) - 2)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]],x]
```

```
[Out] -1/3*(d*(Cos[a + b*x] + Cos[3*(a + b*x)])*(d*Csc[a + b*x])^(3/2)*(Cot[a + b*x]^2 + (-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2]))*Sec[a + b*x]^2*Sqrt[c*Sec[a + b*x]]/(b*(-2 + Csc[a + b*x]^2))
```

fricas [F] time = 0.84, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} d^2 \csc(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")
```

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d^2*csc(b*x + a)^2, x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc (bx + a))^{\frac{5}{2}} \sqrt{c \sec (bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a)), x)

maple [B] time = 1.23, size = 281, normalized size = 3.02

$$\left(2 \sin (bx + a) \cos (bx + a) \sqrt{\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \sqrt{\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \sqrt{\frac{-1 + \cos (bx + a)}{\sin (bx + a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1 - \cos (bx + a)}{\sin (bx + a)}}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x)

[Out] 1/3/b*(2*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-cos(b*x+a)*2^(1/2))*(d/sin(b*x+a))^(5/2)*(c/cos(b*x+a))^(1/2)*sin(b*x+a)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc (bx + a))^{\frac{5}{2}} \sqrt{c \sec (bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{\frac{c}{\cos (a + bx)}} \left(\frac{d}{\sin (a + bx)} \right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2), x)
```

```
[Out] int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(b*x+a))**(5/2)*(c*sec(b*x+a))**(1/2), x)
```

```
[Out] Timed out
```


$$3.233 \quad \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx$$

Optimal. Leaf size=31

$$\frac{2cd\sqrt{d \csc(a + bx)}}{b\sqrt{c \sec(a + bx)}}$$

[Out] $-2*c*d*(d*\csc(b*x+a))^{(1/2)}/b/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2619}

$$\frac{2cd\sqrt{d \csc(a + bx)}}{b\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(3/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]], x]$

[Out] $(-2*c*d*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(b*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 2619

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m + n - 2, 0] \&\& \text{NeQ}[n, 1]$

Rubi steps

$$\int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx = -\frac{2cd\sqrt{d \csc(a + bx)}}{b\sqrt{c \sec(a + bx)}}$$

Mathematica [A] time = 0.07, size = 31, normalized size = 1.00

$$\frac{2cd\sqrt{d \csc(a + bx)}}{b\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(d*\text{Csc}[a + b*x])^{(3/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]], x]$

[Out] $(-2*c*d*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(b*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

fricas [A] time = 0.78, size = 36, normalized size = 1.16

$$\frac{2d \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] -2*d*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)/b

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a)), x)

maple [A] time = 1.22, size = 42, normalized size = 1.35

$$\frac{2 \left(\frac{d}{\sin(bx+a)} \right)^{\frac{3}{2}} \sqrt{\frac{c}{\cos(bx+a)}} \cos(bx+a) \sin(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x)

[Out] -2/b*(d/sin(b*x+a))^(3/2)*(c/cos(b*x+a))^(1/2)*cos(b*x+a)*sin(b*x+a)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a)), x)

mupad [B] time = 0.37, size = 36, normalized size = 1.16

$$\frac{2d \cos(a + bx) \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(3/2),x)
```

```
[Out] -(2*d*cos(a + b*x)*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2))/b
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(b*x+a))**(3/2)*(c*sec(b*x+a))**(1/2),x)
```

```
[Out] Timed out
```

3.234 $\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx$

Optimal. Leaf size=53

$$\frac{\sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b}$$

[Out] $-(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticF}(\cos(a+1/4*\text{Pi}+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b$

Rubi [A] time = 0.09, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2630, 2573, 2641}

$$\frac{\sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]],x]

[Out] (Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/b

Rule 2573

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} dx &= \left(\sqrt{c \cos(a+bx)} \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)} \right) \int \frac{1}{\sqrt{c \cos(a+bx)}} dx \\ &= \left(\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)} \right) \int \frac{1}{\sqrt{\sin(2a+2bx)}} dx \\ &= \frac{\sqrt{d \csc(a+bx)} F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)}}{b} \end{aligned}$$

Mathematica [C] time = 0.67, size = 68, normalized size = 1.28

$$\frac{\tan^3(a+bx) \left(-\cot^2(a+bx)\right)^{7/4} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a+bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]],x]

[Out] ((-Cot[a + b*x]^2)^(7/4)*Sqrt[d*Csc[a + b*x]]*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2]*Sqrt[c*Sec[a + b*x]]*Tan[a + b*x]^3)/b

fricas [F] time = 0.80, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a)), x)

maple [B] time = 1.42, size = 155, normalized size = 2.92

$$\frac{\sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \left(\sin^2(bx+a)\right) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \text{EllipticF}\left(\right)}{b(-1+\cos(bx+a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x)`

[Out] $-1/b*(c/\cos(b*x+a))^{1/2}*(d/\sin(b*x+a))^{1/2}*\sin(b*x+a)^2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})/(-1+\cos(b*x+a))*2^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a)), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \sqrt{\frac{c}{\cos(a + bx)}} \sqrt{\frac{d}{\sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2),x)`

[Out] `int((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))**(1/2)*(c*sec(b*x+a))**(1/2),x)`

[Out] `Integral(sqrt(c*sec(a + b*x))*sqrt(d*csc(a + b*x)), x)`

$$3.235 \quad \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx$$

Optimal. Leaf size=270

$$\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{\sqrt{c \sec(a+bx)}}{2\sqrt{2}}$$

[Out] $1/2 * \arctan(-1 + 2^{1/2} * \tan(b*x+a)^{1/2}) * (c * \sec(b*x+a))^{1/2} / b * 2^{1/2} / (d * c \sec(b*x+a))^{1/2} / \tan(b*x+a)^{1/2} + 1/2 * \arctan(1 + 2^{1/2} * \tan(b*x+a)^{1/2}) * (c * \sec(b*x+a))^{1/2} / b * 2^{1/2} / (d * c \sec(b*x+a))^{1/2} / \tan(b*x+a)^{1/2} + 1/4 * \ln(1 - 2^{1/2} * \tan(b*x+a)^{1/2} + \tan(b*x+a)) * (c * \sec(b*x+a))^{1/2} / b * 2^{1/2} / (d * c \sec(b*x+a))^{1/2} / \tan(b*x+a)^{1/2} - 1/4 * \ln(1 + 2^{1/2} * \tan(b*x+a)^{1/2} + \tan(b*x+a)) * (c * \sec(b*x+a))^{1/2} / b * 2^{1/2} / (d * c \sec(b*x+a))^{1/2} / \tan(b*x+a)^{1/2}$

Rubi [A] time = 0.14, antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {2629, 3476, 329, 297, 1162, 617, 204, 1165, 628}

$$\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{\sqrt{c \sec(a+bx)}}{2\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sec[a + b*x]]/Sqrt[d*Csc[a + b*x]],x]

[Out] $-((\text{ArcTan}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b*x]]] * \text{Sqrt}[c * \text{Sec}[a + b*x]]) / (\text{Sqrt}[2] * b * \text{Sqrt}[d * \text{Csc}[a + b*x]] * \text{Sqrt}[\text{Tan}[a + b*x]])) + (\text{ArcTan}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b*x]]] * \text{Sqrt}[c * \text{Sec}[a + b*x]]) / (\text{Sqrt}[2] * b * \text{Sqrt}[d * \text{Csc}[a + b*x]] * \text{Sqrt}[\text{Tan}[a + b*x]]) + (\text{Log}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]] * \text{Sqrt}[c * \text{Sec}[a + b*x]]) / (2 * \text{Sqrt}[2] * b * \text{Sqrt}[d * \text{Csc}[a + b*x]] * \text{Sqrt}[\text{Tan}[a + b*x]]) - (\text{Log}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]] * \text{Sqrt}[c * \text{Sec}[a + b*x]]) / (2 * \text{Sqrt}[2] * b * \text{Sqrt}[d * \text{Csc}[a + b*x]] * \text{Sqrt}[\text{Tan}[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 297

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a,

b}], x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 329

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^n]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 617

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 628

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1162

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1165

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && ! IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx &= \frac{\sqrt{c \sec(a + bx)} \int \sqrt{\tan(a + bx)} dx}{\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\
 &= \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{\sqrt{x}}{1+x^2} dx, x, \tan(a + bx)\right)}{b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\
 &= \frac{(2\sqrt{c \sec(a + bx)}) \operatorname{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\
 &= -\frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} + \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\
 &= \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a + bx)}\right)}{2b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} + \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a + bx)}\right)}{2b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\
 &= \frac{\log\left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2} b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{\log\left(1 + \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2} b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\
 &= -\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2} b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} + \frac{\tan^{-1}\left(1 + \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2} b\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}
 \end{aligned}$$

Mathematica [A] time = 1.37, size = 171, normalized size = 0.63

$$\frac{\cot(a + bx)\sqrt{c \sec(a + bx)} \left(\log\left(\sqrt{\cot^2(a + bx)} - \sqrt{2} \sqrt[4]{\cot^2(a + bx)} + 1\right) - \log\left(\sqrt{\cot^2(a + bx)} + \sqrt{2} \sqrt[4]{\cot^2(a + bx)} + 1\right) \right)}{2\sqrt{2} b\sqrt[4]{\cot^2(a + bx)} \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sec[a + b*x]]/Sqrt[d*Csc[a + b*x]], x]

[Out] (Cot[a + b*x]*(2*ArcTan[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] - 2*ArcTan[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] + Log[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] +

$\text{Sqrt}[\text{Cot}[a + b*x]^2] - \text{Log}[1 + \text{Sqrt}[2]*(\text{Cot}[a + b*x]^2)^{(1/4)} + \text{Sqrt}[\text{Cot}[a + b*x]^2]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]/(2*\text{Sqrt}[2]*b*(\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Sqrt}[d*\text{Csc}[a + b*x]])$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(bx + a)}}{\sqrt{d \csc(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sec(b*x + a))/sqrt(d*csc(b*x + a)), x)

maple [C] time = 1.00, size = 275, normalized size = 1.02

$$\sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \left(i \text{EllipticPi} \left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x)

[Out] $-1/2/b*(c/\cos(b*x+a))^{(1/2)}*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*(I*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2-1/2*I, 1/2*2^{(1/2)})-I*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2+1/2*I, 1/2*2^{(1/2)})-\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2-1/2*I, 1/2*2^{(1/2)})-\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2+1/2*I, 1/2*2^{(1/2)})))*\sin(b*x+a)/(d/\sin(b*x+a))^{(1/2)/(-1+\cos(b*x+a))*2^{(1/2)}}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(bx + a)}}{\sqrt{d \csc(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*sec(b*x + a))/sqrt(d*csc(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{c}{\cos(a+bx)}}}{\sqrt{\frac{d}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(1/2),x)

[Out] int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(1/2)/(d*csc(b*x+a))**(1/2),x)

[Out] Integral(sqrt(c*sec(a + b*x))/sqrt(d*csc(a + b*x)), x)

$$3.236 \quad \int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{3/2}} dx$$

Optimal. Leaf size=93

$$\frac{\sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{2bd^2} - \frac{c}{bd \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

[Out] $-c/b/d/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-1/2*(\sin(a+1/4*\text{Pi}+b*x)^{2})^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticF}(\cos(a+1/4*\text{Pi}+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b/d^2$

Rubi [A] time = 0.15, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2627, 2630, 2573, 2641}

$$\frac{\sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{2bd^2} - \frac{c}{bd \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[c*\text{Sec}[a+b*x]]/(d*\text{Csc}[a+b*x])^{(3/2)}, x]$

[Out] $-(c/(b*d*\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{Sqrt}[c*\text{Sec}[a+b*x]])) + (\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{EllipticF}[a-\text{Pi}/4+b*x, 2]*\text{Sqrt}[c*\text{Sec}[a+b*x]]*\text{Sqrt}[\text{Sin}[2*a+2*b*x]])/(2*b*d^2)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /;$ FreeQ[{a, b, e, f}, x]

Rule 2627

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(b*(a*\text{Csc}[e + f*x])^{(m+1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(a*f*(m+n)), x] + \text{Dist}[(m+1)/(a^2*(m+n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m+2)}*(b*\text{Sec}[e + f*x])^n, x], x] /;$ FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m+n, 0] && IntegersQ[2*m, 2*n]

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a*\text{Csc}[e + f*x])^m*(b*\text{Sec}[e + f*x])^n*(a*\text{Sin}[e + f*x])$

$\int (b \cos[e + f x])^n \int \frac{1}{(a \sin[e + f x])^m (b \cos[e + f x])^n} dx, x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{IntegerQ}[m - 1/2] \ \&\& \ \text{IntegerQ}[n - 1/2]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2 * \text{EllipticF}[(1 * (c - \text{Pi}/2 + d * x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{3/2}} dx &= -\frac{c}{bd \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} + \frac{\int \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)} dx}{2d^2} \\ &= -\frac{c}{bd \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} + \frac{(\sqrt{c} \cos(a + bx) \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)})}{2d^2} \\ &= -\frac{c}{bd \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} + \frac{(\sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)})}{2d^2} \\ &= -\frac{c}{bd \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} + \frac{\sqrt{d} \csc(a + bx) F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a + bx)}}{2bd^2} \end{aligned}$$

Mathematica [C] time = 0.69, size = 80, normalized size = 0.86

$$\frac{(c \sec(a + bx))^{3/2} \left((-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right) + \cos(2(a + bx)) + 1 \right)}{2bcd \sqrt{d} \csc(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(3/2), x]

[Out] $-1/2 * ((1 + \cos[2 * (a + b * x)]) + (-\cot[a + b * x]^2)^{(3/4)} * \text{Hypergeometric2F1}[1/2, 3/4, 3/2, \csc[a + b * x]^2]) * (c * \sec[a + b * x])^{(3/2)} / (b * c * d * \text{Sqrt}[d * \csc[a + b * x]])$

fricas [F] time = 1.42, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d} \csc(bx + a) \sqrt{c \sec(bx + a)}}{d^2 \csc(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2),x, algorithm="fricas")
 [Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(d^2*csc(b*x + a)^2), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2),x, algorithm="giac")
 [Out] integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(3/2), x)
maple [A] time = 1.02, size = 186, normalized size = 2.00

$$\frac{\left(\sin(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{\sqrt{2}}{2}\right) \right)}{2b(-1 + \cos(bx + a)) \left(\frac{d}{\sin(bx + a)}\right)^{\frac{3}{2}} \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2),x)
 [Out] -1/2/b*(sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*cos(b*x+a)^2-cos(b*x+a)*2^(1/2))*(c/cos(b*x+a))^(1/2)/(-1+cos(b*x+a))/(d/sin(b*x+a))^(3/2)/sin(b*x+a)*2^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(3/2),x, algorithm="maxima")
 [Out] integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(3/2), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{c}{\cos(a+bx)}}}{\left(\frac{d}{\sin(a+bx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(3/2), x)`

[Out] `int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(a + bx)}}{(d \csc(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))**(1/2)/(d*csc(b*x+a))**(3/2), x)`

[Out] `Integral(sqrt(c*sec(a + b*x))/(d*csc(a + b*x))**(3/2), x)`

$$3.237 \quad \int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx$$

Optimal. Leaf size=322

$$\frac{3 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3 \tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3\sqrt{c \sec(a+bx)}}{8\sqrt{2} bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

[Out] $-1/2*c/b/d/(d*csc(b*x+a))^{(3/2)}/(c*sec(b*x+a))^{(1/2)}+3/8*arctan(-1+2^{(1/2)*tan(b*x+a)^{(1/2)}}*(c*sec(b*x+a))^{(1/2)}/b/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/tan(b*x+a)^{(1/2)}+3/8*arctan(1+2^{(1/2)*tan(b*x+a)^{(1/2)}}*(c*sec(b*x+a))^{(1/2)}/b/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/tan(b*x+a)^{(1/2)}+3/16*\ln(1-2^{(1/2)*tan(b*x+a)^{(1/2)}+tan(b*x+a)}*(c*sec(b*x+a))^{(1/2)}/b/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/tan(b*x+a)^{(1/2)}-3/16*\ln(1+2^{(1/2)*tan(b*x+a)^{(1/2)}+tan(b*x+a)}*(c*sec(b*x+a))^{(1/2)}/b/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/tan(b*x+a)^{(1/2)}$

Rubi [A] time = 0.21, antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2627, 2629, 3476, 329, 297, 1162, 617, 204, 1165, 628}

$$\frac{3 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3 \tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3\sqrt{c \sec(a+bx)}}{8\sqrt{2} bd^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(5/2), x]

[Out] $-c/(2*b*d*(d*Csc[a + b*x])^{(3/2)}*Sqrt[c*Sec[a + b*x]]) - (3*ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[c*Sec[a + b*x]])/(4*Sqrt[2]*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (3*ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[c*Sec[a + b*x]])/(4*Sqrt[2]*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (3*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(8*Sqrt[2]*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) - (3*Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(8*Sqrt[2]*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 297

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4)

), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 329

Int[((c_.)*(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 617

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 628

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1162

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1165

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2627

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))/(a*f*(m + n)), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] &

& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{c \sec(a+bx)}}{(d \csc(a+bx))^{5/2}} dx &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} + \frac{3 \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d \csc(a+bx)}} dx}{4d^2} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} + \frac{(3\sqrt{c \sec(a+bx)}) \int \sqrt{\tan(a+bx)} dx}{4d^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} + \frac{(3\sqrt{c \sec(a+bx)}) \text{Subst}\left(\int \frac{\sqrt{x}}{1+x^2} dx, x, \tan(a+bx)\right)}{4bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} + \frac{(3\sqrt{c \sec(a+bx)}) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{2bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} - \frac{(3\sqrt{c \sec(a+bx)}) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{4bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} + \frac{(3\sqrt{c \sec(a+bx)}) \text{Subst}\left(\int \frac{1}{1-\sqrt{2}xx^2} dx, x, \sqrt{\tan(a+bx)}\right)}{8bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} + \frac{3 \log\left(1 - \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{c \sec(a+bx)}}{8\sqrt{2} bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}} \\
 &= -\frac{c}{2bd(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} - \frac{3 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bd^2 \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}
 \end{aligned}$$

Mathematica [A] time = 1.82, size = 222, normalized size = 0.69

$$c\sqrt{d \csc(a + bx)} \left(-4\sqrt[4]{\cot^2(a + bx)} + 4 \cos(2(a + bx))\sqrt[4]{\cot^2(a + bx)} + 3\sqrt{2} \log \left(\sqrt{\cot^2(a + bx)} - \sqrt{2} \sqrt[4]{\cot^2(a + bx)} \right) \right)$$

16

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sec[a + b*x]]/(d*Csc[a + b*x])^(5/2), x]

[Out] (c*Sqrt[d*Csc[a + b*x]]*(6*Sqrt[2]*ArcTan[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4)]) - 6*Sqrt[2]*ArcTan[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] - 4*(Cot[a + b*x]^2)^(1/4) + 4*Cos[2*(a + b*x)]*(Cot[a + b*x]^2)^(1/4) + 3*Sqrt[2]*Log[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4) + Sqrt[Cot[a + b*x]^2]] - 3*Sqrt[2]*Log[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4) + Sqrt[Cot[a + b*x]^2]]))/(16*b*d^3*(Cot[a + b*x]^2)^(1/4)*Sqrt[c*Sec[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(5/2), x)

maple [C] time = 1.10, size = 514, normalized size = 1.60

$$\left(3i \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} + \frac{i}{2} \frac{\sqrt{2}}{2} \right) - \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2),x)

[Out] 1/8/b*(3*I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))-3*I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))+3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))+3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-2*2^(1/2)*cos(b*x+a)^2+2*cos(b*x+a)*2^(1/2))*(c/cos(b*x+a))^(1/2)/(-1+cos(b*x+a))/(d/sin(b*x+a))^(5/2)/sin(b*x+a)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sec(bx + a)}}{(d \csc(bx + a))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(1/2)/(d*csc(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*sec(b*x + a))/(d*csc(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{c}{\cos(a+bx)}}}{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(5/2),x)

[Out] int((c/cos(a + b*x))^(1/2)/(d/sin(a + b*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(1/2)/(d*csc(b*x+a))**(5/2),x)

[Out] Timed out

3.238 $\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx$

Optimal. Leaf size=104

$$\frac{64cd^5\sqrt{c\sec(a+bx)}}{21b\sqrt{d\csc(a+bx)}} - \frac{16cd^3\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{3/2}}{21b} - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{7/2}}{7b}$$

[Out] $-16/21*c*d^3*(d*\csc(b*x+a))^{(3/2)}*(c*\sec(b*x+a))^{(1/2)}/b-2/7*c*d*(d*\csc(b*x+a))^{(7/2)}*(c*\sec(b*x+a))^{(1/2)}/b+64/21*c*d^5*(c*\sec(b*x+a))^{(1/2)}/b/(d*\csc(b*x+a))^{(1/2)}$

Rubi [A] time = 0.16, antiderivative size = 104, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2625, 2619}

$$\frac{64cd^5\sqrt{c\sec(a+bx)}}{21b\sqrt{d\csc(a+bx)}} - \frac{16cd^3\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{3/2}}{21b} - \frac{2cd\sqrt{c\sec(a+bx)}(d\csc(a+bx))^{7/2}}{7b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(9/2)}*(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(64*c*d^5*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(21*b*\text{Sqrt}[d*\text{Csc}[a + b*x]]) - (16*c*d^3*(d*\text{Csc}[a + b*x])^{(3/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(21*b) - (2*c*d*(d*\text{Csc}[a + b*x])^{(7/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(7*b)$

Rule 2619

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(n-1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m + n - 2, 0] \&\& \text{NeQ}[n, 1]$

Rule 2625

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(m-1)), x] + \text{Dist}[(a^2*(m+n-2))/(m-1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{IntegersQ}[2*m, 2*n] \&\& !\text{GtQ}[n, m]$

Rubi steps

$$\begin{aligned} \int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{3/2} dx &= -\frac{2cd(d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)}}{7b} + \frac{1}{7} (8d^2) \int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx \\ &= -\frac{16cd^3(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}}{21b} - \frac{2cd(d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)}}{7b} \\ &= \frac{64cd^5 \sqrt{c \sec(a + bx)}}{21b \sqrt{d \csc(a + bx)}} - \frac{16cd^3(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}}{21b} - \frac{2cd(d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)}}{7b} \end{aligned}$$

Mathematica [A] time = 0.29, size = 57, normalized size = 0.55

$$\frac{2cd^5 (3 \csc^4(a + bx) + 8 \csc^2(a + bx) - 32) \sqrt{c \sec(a + bx)}}{21b \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(9/2)*(c*Sec[a + b*x])^(3/2), x]

[Out] (-2*c*d^5*(-32 + 8*Csc[a + b*x]^2 + 3*Csc[a + b*x]^4)*Sqrt[c*Sec[a + b*x]])/(21*b*Sqrt[d*Csc[a + b*x]])

fricas [A] time = 1.51, size = 85, normalized size = 0.82

$$\frac{2(32cd^4 \cos(bx + a)^4 - 56cd^4 \cos(bx + a)^2 + 21cd^4) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{21(b \cos(bx + a)^2 - b) \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2), x, algorithm="fricas")

[Out] -2/21*(32*c*d^4*cos(b*x + a)^4 - 56*c*d^4*cos(b*x + a)^2 + 21*c*d^4)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/((b*cos(b*x + a)^2 - b)*sin(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{9}{2}} (c \sec(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2), x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(3/2), x)

maple [A] time = 1.08, size = 64, normalized size = 0.62

$$\frac{2 \left(32 \left(\cos^4 (bx + a) \right) - 56 \left(\cos^2 (bx + a) \right) + 21 \right) \cos (bx + a) \left(\frac{d}{\sin (bx + a)} \right)^{\frac{9}{2}} \left(\frac{c}{\cos (bx + a)} \right)^{\frac{3}{2}} \sin (bx + a)}{21b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2),x)

[Out] 2/21/b*(32*cos(b*x+a)^4-56*cos(b*x+a)^2+21)*cos(b*x+a)*(d/sin(b*x+a))^(9/2)*(c/cos(b*x+a))^(3/2)*sin(b*x+a)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc (bx + a))^{\frac{9}{2}} (c \sec (bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(3/2), x)

mupad [B] time = 2.17, size = 110, normalized size = 1.06

$$\frac{16 c d^4 \sqrt{\frac{c}{\cos (a+b x)}} \sqrt{\frac{d}{\sin (a+b x)}} (41 \sin (a+b x)-29 \sin (3 a+3 b x)+12 \sin (5 a+5 b x)-2 \sin (7 a+7 b x))}{21 b (15 \cos (2 a+2 b x)-6 \cos (4 a+4 b x)+\cos (6 a+6 b x)-10)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(9/2),x)

[Out] -(16*c*d^4*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(41*sin(a + b*x) - 29*sin(3*a + 3*b*x) + 12*sin(5*a + 5*b*x) - 2*sin(7*a + 7*b*x)))/(21*b*(15*cos(2*a + 2*b*x) - 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) - 10))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(9/2)*(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

3.239 $\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx$

Optimal. Leaf size=166

$$\frac{24c^2d^4E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{5b\sqrt{\sin(2a + 2bx)}\sqrt{c \sec(a + bx)}\sqrt{d \csc(a + bx)}} + \frac{24cd^5\sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3\sqrt{c \sec(a + bx)}\sqrt{d \csc(a + bx)}}{5b}$$

[Out] $24/5*c*d^5*(c*\sec(b*x+a))^{(1/2)}/b/(d*\csc(b*x+a))^{(3/2)} - 2/5*c*d*(d*\csc(b*x+a))^{(5/2)}*(c*\sec(b*x+a))^{(1/2)}/b - 12/5*c*d^3*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}/b + 24/5*c^2*d^4*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})/b/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.27, antiderivative size = 166, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2625, 2626, 2630, 2572, 2639}

$$\frac{24c^2d^4E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{5b\sqrt{\sin(2a + 2bx)}\sqrt{c \sec(a + bx)}\sqrt{d \csc(a + bx)}} + \frac{24cd^5\sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3\sqrt{c \sec(a + bx)}\sqrt{d \csc(a + bx)}}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(7/2)}*(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(24*c*d^5*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(5*b*(d*\text{Csc}[a + b*x])^{(3/2)}) - (12*c*d^3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(5*b) - (2*c*d*(d*\text{Csc}[a + b*x])^{(5/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(5*b) - (24*c^2*d^4*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2])/ (5*b*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2572

$\text{Int}[\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]], x_Symbol] \rightarrow \text{Dist}[(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]])/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], \text{Int}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}[\{a, b, e, f\}, x]$

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(m-1)), x] + \text{Dist}[(a^{2*(m+n-2)})/(m-1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{IntegersQ}[2*m, 2*n] \&\& !\text{GtQ}[n, m]$

Rule 2626


```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2639

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
 \int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2} dx &= -\frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b} + \frac{1}{5} (6d^2) \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx \\
 &= -\frac{12cd^3 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5b} - \frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b} \\
 &= \frac{24cd^5 \sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5b} - \frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b} \\
 &= \frac{24cd^5 \sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5b} - \frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b} \\
 &= \frac{24cd^5 \sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5b} - \frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b} \\
 &= \frac{24cd^5 \sqrt{c \sec(a + bx)}}{5b(d \csc(a + bx))^{3/2}} - \frac{12cd^3 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5b} - \frac{2cd(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}}{5b}
 \end{aligned}$$

Mathematica [C] time = 1.14, size = 114, normalized size = 0.69

$$\frac{2cd^3 \tan^2(a + bx) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} \left(12 \cos^2(a + bx) \sqrt[4]{-\cot^2(a + bx)} {}_2F_1 \left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx) \right) \right)}{5b}$$

$\cos(b*x+a)/\sin(b*x+a)^{(1/2)} * \text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)}) - 24*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)} * \text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)}) + 12*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)} * \text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)}) - 12*\cos(b*x+a)^3*2^{(1/2)} - 24*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)} * \text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)}) + 12*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)} * ((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)} * \text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)}) + 6*2^{(1/2)}*\cos(b*x+a)^2 + 12*\cos(b*x+a)*2^{(1/2)} - 5*2^{(1/2)}*\cos(b*x+a)*(d/\sin(b*x+a))^{(7/2)}*(c/\cos(b*x+a))^{(3/2)}*\sin(b*x+a)*2^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{3/2} \left(\frac{d}{\sin(a + bx)} \right)^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(7/2),x)

[Out] int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(7/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(7/2)*(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

3.240 $\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx$

Optimal. Leaf size=69

$$\frac{8cd^3\sqrt{c \sec(a + bx)}}{3b\sqrt{d \csc(a + bx)}} - \frac{2cd\sqrt{c \sec(a + bx)}(d \csc(a + bx))^{3/2}}{3b}$$

[Out] $-2/3*c*d*(d*\csc(b*x+a))^{(3/2)}*(c*\sec(b*x+a))^{(1/2)}/b+8/3*c*d^3*(c*\sec(b*x+a))^{(1/2)}/b/(d*\csc(b*x+a))^{(1/2)}$

Rubi [A] time = 0.10, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2625, 2619}

$$\frac{8cd^3\sqrt{c \sec(a + bx)}}{3b\sqrt{d \csc(a + bx)}} - \frac{2cd\sqrt{c \sec(a + bx)}(d \csc(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(5/2)}*(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(8*c*d^3*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(3*b*\text{Sqrt}[d*\text{Csc}[a + b*x]]) - (2*c*d*(d*\text{Csc}[a + b*x])^{(3/2)}*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(3*b)$

Rule 2619

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \ \&\amp; \ \text{EqQ}[m + n - 2, 0] \ \&\amp; \ \text{NeQ}[n, 1]$

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[(a^2*(m + n - 2))/(m - 1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^{(n)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \ \&\amp; \ \text{GtQ}[m, 1] \ \&\amp; \ \text{IntegersQ}[2*m, 2*n] \ \&\amp; \ !\text{GtQ}[n, m]$

Rubi steps

$$\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2} dx = -\frac{2cd(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}}{3b} + \frac{1}{3} (4d^2) \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx$$

$$= \frac{8cd^3 \sqrt{c \sec(a + bx)}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}}{3b}$$

Mathematica [A] time = 0.14, size = 45, normalized size = 0.65

$$\frac{2cd^3 (\csc^2(a + bx) - 4) \sqrt{c \sec(a + bx)}}{3b \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2),x]

[Out] (-2*c*d^3*(-4 + Csc[a + b*x]^2)*Sqrt[c*Sec[a + b*x]])/(3*b*Sqrt[d*Csc[a + b*x]])

fricas [A] time = 0.96, size = 58, normalized size = 0.84

$$\frac{2(4cd^2 \cos(bx + a)^2 - 3cd^2) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{3b \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] -2/3*(4*c*d^2*cos(b*x + a)^2 - 3*c*d^2)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*sin(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2), x)

maple [A] time = 1.04, size = 54, normalized size = 0.78

$$\frac{2(4(\cos^2(bx + a)) - 3) \cos(bx + a) \left(\frac{d}{\sin(bx+a)}\right)^{\frac{5}{2}} \left(\frac{c}{\cos(bx+a)}\right)^{\frac{3}{2}} \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x)`

[Out] $-2/3/b*(4*\cos(b*x+a)^2-3)*\cos(b*x+a)*(d/\sin(b*x+a))^{5/2}*(c/\cos(b*x+a))^{3/2}*\sin(b*x+a)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")`

[Out] `integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2), x)`

mupad [B] time = 0.76, size = 61, normalized size = 0.88

$$\frac{2 c d^2 (2 \sin(a + b x) - \sin(3 a + 3 b x)) \sqrt{\frac{c}{\cos(a+b x)}} \sqrt{\frac{d}{\sin(a+b x)}}}{3 b \sin(a + b x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(5/2),x)`

[Out] $(2*c*d^2*(2*\sin(a + b*x) - \sin(3*a + 3*b*x))*(c/\cos(a + b*x))^{1/2}*(d/\sin(a + b*x))^{1/2})/(3*b*\sin(a + b*x)^2)$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))**(5/2)*(c*sec(b*x+a))**(3/2),x)`

[Out] Timed out

3.241 $\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx$

Optimal. Leaf size=125

$$\frac{4c^2 d^2 E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{b \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} + \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b}$$

[Out] $4*c*d^3*(c*\sec(b*x+a))^{(1/2)}/b/(d*\csc(b*x+a))^{(3/2)}-2*c*d*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}/b+4*c^2*d^2*(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticE}(\cos(a+1/4*\text{Pi}+b*x),2^{(1/2)})/b/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.20, antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2625, 2626, 2630, 2572, 2639}

$$\frac{4c^2 d^2 E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{b \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} + \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(3/2)}*(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(4*c*d^3*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(b*(d*\text{Csc}[a + b*x])^{(3/2)}) - (2*c*d*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]])/b - (4*c^2*d^2*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2])/(b*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2572

$\text{Int}[\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]]$, x_Symbol] $\rightarrow \text{Dist}[(\text{Sqrt}[a*\sin[e + f*x]]*\text{Sqrt}[b*\cos[e + f*x]])/\text{Sqrt}[\sin[2*e + 2*f*x]]$, $\text{Int}[\text{Sqrt}[\sin[2*e + 2*f*x]]$, x], x] /; $\text{FreeQ}[\{a, b, e, f\}, x]$

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}$, x_Symbol] $\rightarrow -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(m-1))$, x] + $\text{Dist}[(a^2*(m+n-2))/(m-1)$, $\text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^n$, x], x] /; $\text{FreeQ}[\{a, b, e, f, n\}, x]$ && $\text{GtQ}[m, 1]$ && $\text{IntegersQ}[2*m, 2*n]$ && ! $\text{GtQ}[n, m]$

Rule 2626

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}$, x_Symbol] $\rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)}$

1))/ (f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_.)])^n, x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx &= -\frac{2cd\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{b} + (2d^2) \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx \\
 &= \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{b} - (4c^2d^2) \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx \\
 &= \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{b} - \frac{(4c^2d^2) \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx}{\sqrt{d \csc(a + bx)}} \\
 &= \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{b} - \frac{(4c^2d^2) \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx}{\sqrt{d \csc(a + bx)}} \\
 &= \frac{4cd^3 \sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2cd\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{b} - \frac{(4c^2d^2) \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx}{b\sqrt{d \csc(a + bx)}}
 \end{aligned}$$

Mathematica [C] time = 0.52, size = 99, normalized size = 0.79

$$\frac{2cd \tan^2(a + bx) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)} \left(2 \cos^2(a + bx) \sqrt{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) + c \right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2), x]

[Out] $(-2*c*d*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(\text{Cos}[2*(a + b*x)]*\text{Cot}[a + b*x]^2 + 2*\text{Cos}[a + b*x]^2*(-\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Hypergeometric2F1}[-1/2, 1/4, 1/2, \text{Csc}[a + b*x]^2])*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Tan}[a + b*x]^2)/b$

fricas [F] time = 1.02, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)} cd \csc(bx+a) \sec(bx+a), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")`

[Out] `integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c*d*csc(b*x + a)*sec(b*x + a), x)`

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx+a))^{\frac{3}{2}} (c \sec(bx+a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")`

[Out] `integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2), x)`

maple [B] time = 1.15, size = 498, normalized size = 3.98

$$\left(-4 \cos(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \text{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x)`

[Out] $-1/b*(-4*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})+2*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})-4*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})+2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})+2*\cos(b*x+a)$

$*2^{(1/2)}-2^{(1/2)})*\cos(b*x+a)*(d/\sin(b*x+a))^{(3/2)}*(c/\cos(b*x+a))^{(3/2)}*\sin(b*x+a)*2^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{3/2} \left(\frac{d}{\sin(a + bx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2),x)

[Out] int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(3/2)*(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

$$3.242 \quad \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx$$

Optimal. Leaf size=31

$$\frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

[Out] $2*c*d*(c*\sec(b*x+a))^{(1/2)}/b/(d*\csc(b*x+a))^{(1/2)}$

Rubi [A] time = 0.05, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2619}

$$\frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(2*c*d*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(b*\text{Sqrt}[d*\text{Csc}[a + b*x]])$

Rule 2619

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m + n - 2, 0] \&\& \text{NeQ}[n, 1]$

Rubi steps

$$\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx = \frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

Mathematica [A] time = 0.07, size = 31, normalized size = 1.00

$$\frac{2cd\sqrt{c \sec(a + bx)}}{b\sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(2*c*d*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(b*\text{Sqrt}[d*\text{Csc}[a + b*x]])$

fricas [A] time = 0.57, size = 36, normalized size = 1.16

$$\frac{2c\sqrt{\frac{c}{\cos(bx+a)}}\sqrt{\frac{d}{\sin(bx+a)}}\sin(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] 2*c*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*sin(b*x + a)/b

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(bx+a)} (c \sec(bx+a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2), x)

maple [A] time = 1.15, size = 42, normalized size = 1.35

$$\frac{2\sqrt{\frac{d}{\sin(bx+a)}}\left(\frac{c}{\cos(bx+a)}\right)^{\frac{3}{2}}\cos(bx+a)\sin(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x)

[Out] 2/b*(d/sin(b*x+a))^(1/2)*(c/cos(b*x+a))^(3/2)*cos(b*x+a)*sin(b*x+a)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(bx+a)} (c \sec(bx+a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2), x)

mupad [B] time = 0.30, size = 36, normalized size = 1.16

$$\frac{2c\sin(a+bx)\sqrt{\frac{c}{\cos(a+bx)}}\sqrt{\frac{d}{\sin(a+bx)}}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(1/2),x)
```

```
[Out] (2*c*sin(a + b*x)*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2))/b
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(b*x+a))**(1/2)*(c*sec(b*x+a))**(3/2),x)
```

```
[Out] Timed out
```

$$3.243 \quad \int \frac{(c \sec(a+bx))^{3/2}}{\sqrt{d} \csc(a+bx)} dx$$

Optimal. Leaf size=89

$$\frac{2cd\sqrt{c \sec(a+bx)}}{b(d \csc(a+bx))^{3/2}} - \frac{2c^2 E\left(a+bx - \frac{\pi}{4} \middle| 2\right)}{b\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

[Out] $2*c*d*(c*\sec(b*x+a))^{(1/2)}/b/(d*\csc(b*x+a))^{(3/2)}+2*c^2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x),2^{(1/2)})/b/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.14, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2626, 2630, 2572, 2639}

$$\frac{2cd\sqrt{c \sec(a+bx)}}{b(d \csc(a+bx))^{3/2}} - \frac{2c^2 E\left(a+bx - \frac{\pi}{4} \middle| 2\right)}{b\sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(3/2)/Sqrt[d*Csc[a + b*x]], x]

[Out] $(2*c*d*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(b*(d*\text{Csc}[a + b*x])^{(3/2)}) - (2*c^2*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2])/(b*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]], x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[2*e + 2*f*x]], Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2626

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])

)^m*(b*cos[e + f*x])ⁿ, Int[1/((a*sin[e + f*x])^m*(b*cos[e + f*x])ⁿ), x],
 x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P
 i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(c \sec(a + bx))^{3/2}}{\sqrt{d \csc(a + bx)}} dx &= \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - (2c^2) \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx \\ &= \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{(2c^2) \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)}} \\ &= \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{(2c^2) \int \sqrt{\sin(2a + 2bx)} dx}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \\ &= \frac{2cd\sqrt{c \sec(a + bx)}}{b(d \csc(a + bx))^{3/2}} - \frac{2c^2 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{b\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] time = 0.38, size = 66, normalized size = 0.74

$$\frac{2cd\sqrt{c \sec(a + bx)} \left(\sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) - 1 \right)}{b(d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(3/2)/Sqrt[d*Csc[a + b*x]], x]

[Out] (-2*c*d*(-1 + (-Cot[a + b*x]^2)^(1/4))*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]]/(b*(d*Csc[a + b*x])^(3/2))

fricas [F] time = 0.60, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} c \sec(bx + a)}{d \csc(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c*sec(b*x + a)/(d*csc(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{\sqrt{d} \csc(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(3/2)/sqrt(d*csc(b*x + a)), x)

maple [B] time = 1.09, size = 497, normalized size = 5.58

$$\left(\cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{\sqrt{2}}{2}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x)

[Out] -1/b*(cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-2*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-2*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+cos(b*x+a)*2^(1/2)-2^(1/2))*cos(b*x+a)*(c/cos(b*x+a))^(3/2)/(d/sin(b*x+a))^(1/2)/sin(b*x+a)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{\sqrt{d} \csc(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(3/2)/sqrt(d*csc(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}}{\sqrt{\frac{d}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(1/2),x)

[Out] int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(3/2)/(d*csc(b*x+a))**(1/2),x)

[Out] Timed out

$$3.244 \quad \int \frac{(c \sec(a+bx))^{3/2}}{(d \csc(a+bx))^{3/2}} dx$$

Optimal. Leaf size=327

$$\frac{c^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} b d^2 \sqrt{c \sec(a+bx)}} - \frac{c^2 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d}}{\sqrt{2} b d^2 \sqrt{c \sec(a+bx)}}$$

[Out] 2*c*(c*sec(b*x+a))^(1/2)/b/d/(d*csc(b*x+a))^(1/2)-1/2*c^2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/2*c^2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/4*c^2*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/4*c^2*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b/d^2*2^(1/2)/(c*sec(b*x+a))^(1/2)

Rubi [A] time = 0.22, antiderivative size = 327, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2624, 2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{c^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} b d^2 \sqrt{c \sec(a+bx)}} - \frac{c^2 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d}}{\sqrt{2} b d^2 \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(3/2), x]

[Out] (2*c*Sqrt[c*Sec[a + b*x]])/(b*d*Sqrt[d*Csc[a + b*x]]) + (c^2*ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])/(Sqrt[2]*b*d^2*Sqrt[c*Sec[a + b*x]]) - (c^2*ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])/(Sqrt[2]*b*d^2*Sqrt[c*Sec[a + b*x]]) + (c^2*Sqrt[d*Csc[a + b*x]]*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[Tan[a + b*x]])/(2*Sqrt[2]*b*d^2*Sqrt[c*Sec[a + b*x]]) - (c^2*Sqrt[d*Csc[a + b*x]]*Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[Tan[a + b*x]])/(2*Sqrt[2]*b*d^2*Sqrt[c*Sec[a + b*x]])

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

```
Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_)*(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2624

```
Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n
_), x_Symbol] := Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))
/(f*a*(n - 1)), x] + Dist[(b^2*(m + 1))/(a^2*(n - 1)), Int[(a*Csc[e + f*x])
```

$(m + 2) \cdot (b \cdot \sec[e + f \cdot x])^{n - 2}$, x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[n, 1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{3/2}} dx &= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} - \frac{c^2 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{d^2} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} - \frac{(c^2\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{d^2\sqrt{c \sec(a + bx)}} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} - \frac{(c^2\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1}{\sqrt{x}(1+x^2)} dx, x, \tan(a + bx) \right)}{bd^2\sqrt{c \sec(a + bx)}} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} - \frac{(2c^2\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, \sqrt{\tan(a + bx)} \right)}{bd^2\sqrt{c \sec(a + bx)}} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} - \frac{(c^2\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)} \right)}{bd^2\sqrt{c \sec(a + bx)}} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} - \frac{(c^2\sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a + bx)} \right)}{2bd^2\sqrt{c \sec(a + bx)}} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} + \frac{c^2\sqrt{d \csc(a + bx)} \log \left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx) \right) \sqrt{\tan(a + bx)}}{2\sqrt{2} bd^2\sqrt{c \sec(a + bx)}} \\
&= \frac{2c\sqrt{c \sec(a + bx)}}{bd\sqrt{d \csc(a + bx)}} + \frac{c^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a + bx)} \right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{\sqrt{2} bd^2\sqrt{c \sec(a + bx)}}
\end{aligned}$$

Mathematica [C] time = 0.25, size = 64, normalized size = 0.20

$$\frac{2c\sqrt{c \sec(a + bx)} \left(\cot^2(a + bx) {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a + bx) \right) + 3 \right)}{3bd\sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(3/2), x]

[Out] (2*c*(3 + Cot[a + b*x]^2*Hypergeometric2F1[3/4, 1, 7/4, -Cot[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(3*b*d*Sqrt[d*Csc[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{(d \csc(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(3/2), x)

maple [C] time = 1.06, size = 646, normalized size = 1.98

$$\left(i \operatorname{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} - \frac{i}{2} \frac{\sqrt{2}}{2} \right) \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \sin \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x)

[Out] 1/2/b*(I*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)-I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)+EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)+((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)-2*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2*cos(b*x+a)*2^(1/2)-2*2^(1/2))*cos(b*x+a)*(c/cos(b*x+a))^(3/2)/(-1+cos(b*x+a))/(d/sin(b*x+a))^(3/2)/sin(b*x+a)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{(d \csc(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(3/2),x)

[Out] int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(3/2)/(d*csc(b*x+a))**(3/2),x)

[Out] Timed out

$$3.245 \quad \int \frac{(c \sec(a+bx))^{3/2}}{(d \csc(a+bx))^{5/2}} dx$$

Optimal. Leaf size=94

$$\frac{2c\sqrt{c \sec(a+bx)}}{bd(d \csc(a+bx))^{3/2}} - \frac{3c^2 E\left(a+bx - \frac{\pi}{4} \middle| 2\right)}{bd^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

[Out] $2*c*(c*\sec(b*x+a))^{(1/2)}/b/d/(d*\csc(b*x+a))^{(3/2)}+3*c^2*(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticE}(\cos(a+1/4*\text{Pi}+b*x),2^{(1/2)})/b/d^2/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.15, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2624, 2630, 2572, 2639}

$$\frac{2c\sqrt{c \sec(a+bx)}}{bd(d \csc(a+bx))^{3/2}} - \frac{3c^2 E\left(a+bx - \frac{\pi}{4} \middle| 2\right)}{bd^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(5/2), x]

[Out] $(2*c*\text{Sqrt}[c*\text{Sec}[a + b*x]])/(b*d*(d*\text{Csc}[a + b*x])^{(3/2)}) - (3*c^2*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2])/(b*d^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]], x_Symbol] := Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[2*e + 2*f*x]], Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2624

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))/(f*a*(n - 1)), x] + Dist[(b^2*(m + 1))/(a^2*(n - 1)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[n, 1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])

$)^m (b \cos[e + f x])^n, \text{Int}[1/((a \sin[e + f x])^m (b \cos[e + f x])^n), x],$
 $x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x\} \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$
 $]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2 * \text{EllipticE}[(1 * (c - P$
 $i/2 + d * x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{(c \sec(a + bx))^{3/2}}{(d \csc(a + bx))^{5/2}} dx &= \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{(3c^2) \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx}{d^2} \\ &= \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{(3c^2) \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{d^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)}} \\ &= \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{(3c^2) \int \sqrt{\sin(2a + 2bx)} dx}{d^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \\ &= \frac{2c\sqrt{c \sec(a + bx)}}{bd(d \csc(a + bx))^{3/2}} - \frac{3c^2 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{bd^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] time = 0.42, size = 69, normalized size = 0.73

$$\frac{c\sqrt{c \sec(a + bx)} \left(3\sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) - 2\right)}{bd(d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(3/2)/(d*Csc[a + b*x])^(5/2), x]

[Out] -((c*(-2 + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(b*d*(d*Csc[a + b*x])^(3/2)))

fricas [F] time = 0.82, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} c \sec(bx + a)}{d^3 \csc(bx + a)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c*sec(b*x + a)/(d^3*csc(b*x + a)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{(d \csc(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(5/2), x)

maple [B] time = 1.01, size = 512, normalized size = 5.45

$$\left(6 \cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \operatorname{EllipticE}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{\sqrt{2}}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x)

[Out] 1/2/b*(6*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-3*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+6*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*cos(b*x+a)^2-3*cos(b*x+a)*2^(1/2)+2*2^(1/2))*cos(b*x+a)*(c/cos(b*x+a))^(3/2)/sin(b*x+a)^3/(d/sin(b*x+a))^(5/2)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{3}{2}}}{(d \csc(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sec(b*x+a))^(3/2)/(d*csc(b*x+a))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((c*sec(b*x + a))^(3/2)/(d*csc(b*x + a))^(5/2), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(5/2),x)
```

```
[Out] int((c/cos(a + b*x))^(3/2)/(d/sin(a + b*x))^(5/2), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sec(b*x+a))**(3/2)/(d*csc(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```

3.246 $\int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx$

Optimal. Leaf size=166

$$\frac{40c^2d^4\sqrt{\sin(2a+2bx)}F\left(a+bx-\frac{\pi}{4}\middle|2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{21b} + \frac{40cd^5(c\sec(a+bx))^{3/2}}{21b\sqrt{d\csc(a+bx)}} - \frac{20cd^3(c\sec(a+bx))^{5/2}}{21b\sqrt{d\csc(a+bx)}}$$

[Out] $-20/21*c*d^3*(d*\csc(b*x+a))^{(3/2)}*(c*\sec(b*x+a))^{(3/2)}/b-2/7*c*d*(d*\csc(b*x+a))^{(7/2)}*(c*\sec(b*x+a))^{(3/2)}/b+40/21*c*d^5*(c*\sec(b*x+a))^{(3/2)}/b/(d*\csc(b*x+a))^{(1/2)}-40/21*c^2*d^4*(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticF}(\cos(a+1/4*\text{Pi}+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b$

Rubi [A] time = 0.26, antiderivative size = 166, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2625, 2626, 2630, 2573, 2641}

$$\frac{40c^2d^4\sqrt{\sin(2a+2bx)}F\left(a+bx-\frac{\pi}{4}\middle|2\right)\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}}{21b} + \frac{40cd^5(c\sec(a+bx))^{3/2}}{21b\sqrt{d\csc(a+bx)}} - \frac{20cd^3(c\sec(a+bx))^{5/2}}{21b\sqrt{d\csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(9/2)}*(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(40*c*d^5*(c*\text{Sec}[a + b*x])^{(3/2)})/(21*b*\text{Sqrt}[d*\text{Csc}[a + b*x]]) - (20*c*d^3*(d*\text{Csc}[a + b*x])^{(3/2)}*(c*\text{Sec}[a + b*x])^{(3/2)})/(21*b) - (2*c*d*(d*\text{Csc}[a + b*x])^{(7/2)}*(c*\text{Sec}[a + b*x])^{(3/2)})/(7*b) + (40*c^2*d^4*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(21*b)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]])], x_Symbol] :> \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f, x\}$

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n)}], x_Symbol] :> -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(m-1)), x] + \text{Dist}[(a^2*(m+n-2))/(m-1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n, x\} \&\& \text{GtQ}[m, 1] \&\& \text{IntegersQ}[2*m, 2*n] \&\& !\text{GtQ}[n, m]$

Rule 2626

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2641

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx &= -\frac{2cd(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2}}{7b} + \frac{1}{7} (10d^2) \int (d \csc(a + bx))^{9/2} (c \sec(a + bx))^{5/2} dx \\ &= -\frac{20cd^3(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{21b} - \frac{2cd(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{3/2}}{7b} \\ &= \frac{40cd^5(c \sec(a + bx))^{3/2}}{21b\sqrt{d \csc(a + bx)}} - \frac{20cd^3(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{21b} \\ &= \frac{40cd^5(c \sec(a + bx))^{3/2}}{21b\sqrt{d \csc(a + bx)}} - \frac{20cd^3(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{21b} \\ &= \frac{40cd^5(c \sec(a + bx))^{3/2}}{21b\sqrt{d \csc(a + bx)}} - \frac{20cd^3(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{21b} \\ &= \frac{40cd^5(c \sec(a + bx))^{3/2}}{21b\sqrt{d \csc(a + bx)}} - \frac{20cd^3(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{21b} \end{aligned}$$

Mathematica [C] time = 1.45, size = 92, normalized size = 0.55

$$\frac{2cd^5(c \sec(a + bx))^{3/2} \left(20(-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right) + \cot^2(a + bx) (3 \csc^2(a + bx) + 13) \right)}{21b\sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(9/2)*(c*Sec[a + b*x])^(5/2), x]

[Out] $(-2*c*d^5*(-7 + \cot[a + b*x])^2*(13 + 3*\csc[a + b*x]^2) + 20*(-\cot[a + b*x]^2)^{(3/4)}*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \csc[a + b*x]^2])*(c*\sec[a + b*x])^{(3/2)}/(21*b*\sqrt{d*\csc[a + b*x]})$

fricas [F] time = 0.64, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} c^2 d^4 \csc(bx + a)^4 \sec(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c^2*d^4*csc(b*x + a)^4*sec(b*x + a)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{9}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(5/2), x)

maple [B] time = 1.19, size = 555, normalized size = 3.34

$$\frac{\left(40 \left(\cos^4(bx + a)\right) \sin(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \text{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{1}{2}\right) + 40 \sin(bx + a) \cos(bx + a)^3 \left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \text{EllipticF}\left(\left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)}, \frac{1}{2}\right) - 40 \sin(bx + a) \cos(bx + a)^2 \left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \text{EllipticF}\left(\left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)}, \frac{1}{2}\right) - 40 \sin(bx + a) \cos(bx + a) \left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \text{EllipticF}\left(\left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)}, \frac{1}{2}\right) - 40 \sin(bx + a) \left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \left(\frac{-1 + \cos(bx + a)}{\sin(bx + a)}\right)^{(1/2)} \text{EllipticF}\left(\left(\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}\right)^{(1/2)}, \frac{1}{2}\right)}{\sin(bx + a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2), x)

[Out] $-1/21/b*(40*\cos(b*x+a)^4*\sin(b*x+a)*\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\text{EllipticF}\left(\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}, 1/2\right)+40*\sin(b*x+a)*\cos(b*x+a)^3*\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\text{EllipticF}\left(\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}, 1/2\right)-40*\sin(b*x+a)*\cos(b*x+a)^2*\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\text{EllipticF}\left(\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}, 1/2\right)-40*\sin(b*x+a)*\cos(b*x+a)*\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\text{EllipticF}\left(\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}, 1/2\right)-40*\sin(b*x+a)*\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}*\text{EllipticF}\left(\left(\frac{1-\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{(1/2)}, 1/2\right)$

$$\begin{aligned}
 & -1 + \cos(b*x+a) + \sin(b*x+a) / \sin(b*x+a) \wedge (1/2) * ((-1 + \cos(b*x+a)) / \sin(b*x+a)) \wedge (1/2) \\
 & * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a)) \wedge (1/2), 1/2 * 2 \wedge (1/2)) - 40 * \\
 & \sin(b*x+a) * \cos(b*x+a) * ((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a)) \wedge (1/2) * ((-1 + \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a)) \wedge (1/2) * \\
 & \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a)) \wedge (1/2), 1/2 * 2 \wedge (1/2)) - 20 * \cos(b*x+a) \wedge 4 * 2 \wedge (1/2) \\
 & + 30 * 2 \wedge (1/2) * \cos(b*x+a) \wedge 2 - 7 * 2 \wedge (1/2) * \cos(b*x+a) * (d / \sin(b*x+a)) \wedge (9/2) * (c / \cos(b*x+a)) \wedge (5/2) * \sin(b*x+a) * 2 \wedge (1/2)
 \end{aligned}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{9}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(9/2)*(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{5/2} \left(\frac{d}{\sin(a + bx)} \right)^{9/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(9/2),x)

[Out] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(9/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(9/2)*(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

3.247 $\int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx$

Optimal. Leaf size=106

$$-\frac{64c^3d^3\sqrt{d \csc(a + bx)}}{15b\sqrt{c \sec(a + bx)}} + \frac{16cd^3(c \sec(a + bx))^{3/2}\sqrt{d \csc(a + bx)}}{15b} - \frac{2cd(c \sec(a + bx))^{3/2}(d \csc(a + bx))^{5/2}}{5b}$$

[Out] $-2/5*c*d*(d*\csc(b*x+a))^{(5/2)}*(c*\sec(b*x+a))^{(3/2)}/b+16/15*c*d^3*(c*\sec(b*x+a))^{(3/2)}*(d*\csc(b*x+a))^{(1/2)}/b-64/15*c^3*d^3*(d*\csc(b*x+a))^{(1/2)}/b/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.16, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2625, 2626, 2619}

$$-\frac{64c^3d^3\sqrt{d \csc(a + bx)}}{15b\sqrt{c \sec(a + bx)}} + \frac{16cd^3(c \sec(a + bx))^{3/2}\sqrt{d \csc(a + bx)}}{15b} - \frac{2cd(c \sec(a + bx))^{3/2}(d \csc(a + bx))^{5/2}}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(7/2)}*(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(-64*c^3*d^3*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(15*b*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (16*c*d^3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(3/2)})/(15*b) - (2*c*d*(d*\text{Csc}[a + b*x])^{(5/2)}*(c*\text{Sec}[a + b*x])^{(3/2)})/(5*b)$

Rule 2619

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] /;$ FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[(a^2*(m + n - 2))/(m - 1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /;$ FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2626

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[(a^2*(m + n - 2))/(m - 1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /;$

1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned} \int (d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2} dx &= -\frac{2cd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}}{5b} + \frac{1}{5} (8d^2) \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx \\ &= \frac{16cd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{15b} - \frac{2cd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}}{5b} \\ &= -\frac{64c^3 d^3 \sqrt{d \csc(a + bx)}}{15b \sqrt{c \sec(a + bx)}} + \frac{16cd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{15b} \end{aligned}$$

Mathematica [A] time = 0.19, size = 57, normalized size = 0.54

$$-\frac{2cd^3 (c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)} (32 \cos^2(a + bx) + 3 \cot^2(a + bx) - 5)}{15b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2), x]

[Out] (-2*c*d^3*(-5 + 32*Cos[a + b*x]^2 + 3*Cot[a + b*x]^2)*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2))/(15*b)

fricas [A] time = 0.97, size = 89, normalized size = 0.84

$$-\frac{2 \left(32 c^2 d^3 \cos(bx + a)^4 - 40 c^2 d^3 \cos(bx + a)^2 + 5 c^2 d^3 \right) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{15 (b \cos(bx + a)^3 - b \cos(bx + a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2), x, algorithm="fricas")

[Out] -2/15*(32*c^2*d^3*cos(b*x + a)^4 - 40*c^2*d^3*cos(b*x + a)^2 + 5*c^2*d^3)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*cos(b*x + a)^3 - b*cos(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{7/2} (c \sec(bx + a))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2), x)

maple [A] time = 1.12, size = 64, normalized size = 0.60

$$\frac{2 \left(32 \left(\cos^4 (bx + a) \right) - 40 \left(\cos^2 (bx + a) \right) + 5 \right) \cos (bx + a) \left(\frac{d}{\sin (bx+a)} \right)^{\frac{7}{2}} \left(\frac{c}{\cos (bx+a)} \right)^{\frac{5}{2}} \sin (bx + a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x)

[Out] 2/15/b*(32*cos(b*x+a)^4-40*cos(b*x+a)^2+5)*cos(b*x+a)*(d/sin(b*x+a))^(7/2)*(c/cos(b*x+a))^(5/2)*sin(b*x+a)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc (bx + a))^{\frac{7}{2}} (c \sec (bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2), x)

mupad [B] time = 2.32, size = 112, normalized size = 1.06

$$\frac{16 c^2 d^3 \sqrt{\frac{c}{\cos (a+b x)}} \sqrt{\frac{d}{\sin (a+b x)}} (5 \cos (a+b x)-3 \cos (3 a+3 b x)-4 \cos (5 a+5 b x)+2 \cos (7 a+7 b x))}{15 b (\cos (2 a+2 b x)+2 \cos (4 a+4 b x)-\cos (6 a+6 b x)-2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(7/2),x)

[Out] (16*c^2*d^3*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(5*cos(a + b*x) - 3*cos(3*a + 3*b*x) - 4*cos(5*a + 5*b*x) + 2*cos(7*a + 7*b*x)))/(15*b*(cos(2*a + 2*b*x) + 2*cos(4*a + 4*b*x) - cos(6*a + 6*b*x) - 2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(b*x+a))**(7/2)*(c*sec(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```

3.248 $\int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx$

Optimal. Leaf size=131

$$\frac{4c^2 d^2 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} + \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd (c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}}$$

[Out] $-2/3 * c * d * (d * \csc(b * x + a))^{3/2} * (c * \sec(b * x + a))^{3/2} / b + 4/3 * c * d^3 * (c * \sec(b * x + a))^{3/2} / b / (d * \csc(b * x + a))^{1/2} - 4/3 * c^2 * d^2 * (\sin(a + 1/4 * \pi + b * x)^2)^{1/2} / \sin(a + 1/4 * \pi + b * x) * \text{EllipticF}(\cos(a + 1/4 * \pi + b * x), 2^{1/2}) * (d * \csc(b * x + a))^{1/2} * (c * \sec(b * x + a))^{1/2} * \sin(2 * b * x + 2 * a)^{1/2} / b$

Rubi [A] time = 0.20, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2625, 2626, 2630, 2573, 2641}

$$\frac{4c^2 d^2 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} + \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}} - \frac{2cd (c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d * \text{Csc}[a + b * x])^{5/2} * (c * \text{Sec}[a + b * x])^{5/2}, x]$

[Out] $(4 * c * d^3 * (c * \text{Sec}[a + b * x])^{3/2}) / (3 * b * \text{Sqrt}[d * \text{Csc}[a + b * x]]) - (2 * c * d * (d * \text{Csc}[a + b * x])^{3/2} * (c * \text{Sec}[a + b * x])^{3/2}) / (3 * b) + (4 * c^2 * d^2 * \text{Sqrt}[d * \text{Csc}[a + b * x]] * \text{EllipticF}[a - \pi/4 + b * x, 2] * \text{Sqrt}[c * \text{Sec}[a + b * x]] * \text{Sqrt}[\sin[2 * a + 2 * b * x]]) / (3 * b)$

Rule 2573

$\text{Int}[1 / (\text{Sqrt}[\cos[(e_.) + (f_.) * (x_.)] * (b_.)] * \text{Sqrt}[(a_.) * \sin[(e_.) + (f_.) * (x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\sin[2 * e + 2 * f * x]] / (\text{Sqrt}[a * \sin[e + f * x]] * \text{Sqrt}[b * \cos[e + f * x]]), \text{Int}[1 / \text{Sqrt}[\sin[2 * e + 2 * f * x]], x], x] /;$ FreeQ[{a, b, e, f}, x]

Rule 2625

$\text{Int}[(\csc[(e_.) + (f_.) * (x_.)] * (a_.))^{(m_.)} * ((b_.) * \sec[(e_.) + (f_.) * (x_.)])^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(a * b * (a * \csc[e + f * x])^{(m - 1)} * (b * \sec[e + f * x])^{(n - 1)}) / (f * (m - 1)), x] + \text{Dist}[(a^2 * (m + n - 2)) / (m - 1), \text{Int}[(a * \csc[e + f * x])^{(m - 2)} * (b * \sec[e + f * x])^n, x], x] /;$ FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2 * m, 2 * n] && !GtQ[n, m]

Rule 2626

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] + Dist[(b^2*(m + n - 2))/(n - 1), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2641

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int (d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2} dx &= -\frac{2cd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{3b} + (2d^2) \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx \\ &= \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} - \frac{2cd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{3b} + \frac{1}{3} \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx \\ &= \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} - \frac{2cd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{3b} + \frac{1}{3} \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx \\ &= \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} - \frac{2cd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{3b} + \frac{1}{3} \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2} dx \\ &= \frac{4cd^3 (c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} - \frac{2cd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}{3b} + \frac{4c^2}{3} \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2} dx \end{aligned}$$

Mathematica [C] time = 0.69, size = 87, normalized size = 0.66

$$\frac{2c^3 d \tan^2(a + bx) (d \csc(a + bx))^{3/2} \left(2 \left(-\cot^2(a + bx) \right)^{3/4} {}_2F_1 \left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx) \right) + \cot^2(a + bx) - 1 \right)}{3b\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2),x]

[Out] $(-2*c^3*d*(d*Csc[a + b*x])^{3/2}*(-1 + Cot[a + b*x]^2 + 2*(-Cot[a + b*x]^2)^{3/4}*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*Tan[a + b*x]^2)/(3*b*Sqrt[c*Sec[a + b*x]])$

fricas [F] time = 0.67, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} c^2 d^2 \csc(bx + a)^2 \sec(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c^2*d^2*csc(b*x + a)^2*sec(b*x + a)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2), x)

maple [B] time = 1.16, size = 300, normalized size = 2.29

$$\left(4 \sin(bx + a) \left(\cos^2(bx + a)\right) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \text{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a)}{\sin(bx + a)}}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x)

[Out] $\frac{1}{3} b^* (4 * \sin(b*x+a) * \cos(b*x+a)^2 * ((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2} * ((-1 + \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2} * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2}, 1/2 * 2^{1/2}) + 4 * \sin(b*x+a) * \cos(b*x+a) * ((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2} * ((-1 + \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2} * ((-1 + \cos(b*x+a)) / \sin(b*x+a))^{1/2} * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2}, 1/2 * 2^{1/2}) - 2 * 2^{1/2} * \cos(b*x+a)^2 * 2^{1/2} * \cos(b*x+a) * (d / \sin(b*x+a))^{5/2} * (c / \cos(b*x+a))^{5/2} * \sin(b*x+a) * 2^{1/2})$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)} \right)^{5/2} \left(\frac{d}{\sin(a + bx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2),x)

[Out] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(5/2)*(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

$$3.249 \quad \int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx$$

Optimal. Leaf size=69

$$\frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} - \frac{8c^3 d \sqrt{d \csc(a + bx)}}{3b \sqrt{c \sec(a + bx)}}$$

[Out] $2/3*c*d*(c*\sec(b*x+a))^{3/2}*(d*\csc(b*x+a))^{1/2}/b-8/3*c^3*d*(d*\csc(b*x+a))^{1/2}/b/(c*\sec(b*x+a))^{1/2}$

Rubi [A] time = 0.10, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2626, 2619}

$$\frac{2cd(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b} - \frac{8c^3 d \sqrt{d \csc(a + bx)}}{3b \sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{3/2}*(c*\text{Sec}[a + b*x])^{5/2}, x]$

[Out] $(-8*c^3*d*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(3*b*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (2*c*d*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{3/2})/(3*b)$

Rule 2619

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m + n - 2, 0] \&\& \text{NeQ}[n, 1]$

Rule 2626

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*(b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] + \text{Dist}[(b^2*(m + n - 2))/(n - 1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, m\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rubi steps

$$\int (d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2} dx = \frac{2cd\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{3b} + \frac{1}{3} (4c^2) \int (d \csc(a + bx))^{3/2} \sqrt{d \csc(a + bx)} dx$$

$$= -\frac{8c^3 d \sqrt{d \csc(a + bx)}}{3b \sqrt{c \sec(a + bx)}} + \frac{2cd\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{3/2}}{3b}$$

Mathematica [A] time = 0.21, size = 45, normalized size = 0.65

$$-\frac{2cd(2 \cos(2(a + bx)) + 1)(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2),x]

[Out] (-2*c*d*(1 + 2*Cos[2*(a + b*x)])*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2))/(3*b)

fricas [A] time = 0.84, size = 58, normalized size = 0.84

$$-\frac{2(4c^2d \cos(bx + a)^2 - c^2d) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{3b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] -2/3*(4*c^2*d*cos(b*x + a)^2 - c^2*d)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*cos(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2), x)

maple [A] time = 1.04, size = 54, normalized size = 0.78

$$-\frac{2(4(\cos^2(bx + a)) - 1) \cos(bx + a) \left(\frac{d}{\sin(bx+a)}\right)^{\frac{3}{2}} \left(\frac{c}{\cos(bx+a)}\right)^{\frac{5}{2}} \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x)`

[Out] $-2/3/b*(4*\cos(b*x+a)^2-1)*\cos(b*x+a)*(d/\sin(b*x+a))^{3/2}*(c/\cos(b*x+a))^{5/2}*\sin(b*x+a)$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(3/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")`

[Out] `integrate((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2), x)`

mupad [B] time = 0.83, size = 64, normalized size = 0.93

$$\frac{4c^2 d \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (2 \cos(a+bx) + \cos(3a+3bx))}{3b (\cos(2a+2bx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(3/2),x)`

[Out] $-(4*c^2*d*(c/\cos(a + b*x))^{1/2}*(d/\sin(a + b*x))^{1/2}*(2*\cos(a + b*x) + \cos(3*a + 3*b*x)))/(3*b*(\cos(2*a + 2*b*x) + 1))$

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))**(3/2)*(c*sec(b*x+a))**(5/2),x)`

[Out] Timed out

3.250 $\int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx$

Optimal. Leaf size=93

$$\frac{2c^2 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} + \frac{2cd(c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}}$$

[Out] $2/3*c*d*(c*\sec(b*x+a))^{(3/2)}/b/(d*\csc(b*x+a))^{(1/2)}-2/3*c^2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b$

Rubi [A] time = 0.14, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2626, 2630, 2573, 2641}

$$\frac{2c^2 \sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}{3b} + \frac{2cd(c \sec(a + bx))^{3/2}}{3b \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(2*c*d*(c*\text{Sec}[a + b*x])^{(3/2)})/(3*b*\text{Sqrt}[d*\text{Csc}[a + b*x]]) + (2*c^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(3*b)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2626

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(n-1)), x] + \text{Dist}[(b^2*(m+n-2))/(n-1), \text{Int}[(a*\text{Csc}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n-2)}, x], x] /; \text{FreeQ}\{a, b, e, f, m\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a*\text{Csc}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(n-1)}], x]$

)^m*(b*cos[e + f*x])^n, Int[1/((a*sin[e + f*x])^m*(b*cos[e + f*x])^n), x],
 x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c -
 Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2} dx &= \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} + \frac{1}{3} (2c^2) \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx \\ &= \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} + \frac{1}{3} (2c^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \\ &= \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} + \frac{1}{3} (2c^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)} \\ &= \frac{2cd(c \sec(a + bx))^{3/2}}{3b\sqrt{d \csc(a + bx)}} + \frac{2c^2 \sqrt{d \csc(a + bx)} F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a + bx)}}{3b} \end{aligned}$$

Mathematica [C] time = 0.60, size = 68, normalized size = 0.73

$$\frac{2cd(c \sec(a + bx))^{3/2} \left((-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right) - 1 \right)}{3b\sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2),x]

[Out] (-2*c*d*(-1 + (-Cot[a + b*x]^2)^(3/4))*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*(c*Sec[a + b*x])^(3/2)/(3*b*Sqrt[d*Csc[a + b*x]])

fricas [F] time = 0.51, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} c^2 \sec(bx + a)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c^2*sec(b*x + a)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(bx + a)} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2), x)

maple [A] time = 1.25, size = 188, normalized size = 2.02

$$\frac{\left(2 \sin(bx + a) \cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a)}{\sin(bx + a)}}\right)\right)}{3b(-1 + \cos(bx + a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2),x)

[Out] -1/3/b*(2*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2))*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-cos(b*x+a)*2^(1/2)+2^(1/2))*cos(b*x+a)*(d/sin(b*x+a))^(1/2)*(c/cos(b*x+a))^(5/2)*sin(b*x+a)/(-1+cos(b*x+a))*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(bx + a)} (c \sec(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(a + bx)}\right)^{5/2} \sqrt{\frac{d}{\sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2),x)
```

```
[Out] int((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2), x)
```

```
sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(b*x+a))**(1/2)*(c*sec(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```

$$3.251 \quad \int \frac{(c \sec(a+bx))^{5/2}}{\sqrt{d} \csc(a+bx)} dx$$

Optimal. Leaf size=33

$$\frac{2cd(c \sec(a+bx))^{3/2}}{3b(d \csc(a+bx))^{3/2}}$$

[Out] $2/3*c*d*(c*\sec(b*x+a))^(3/2)/b/(d*\csc(b*x+a))^(3/2)$

Rubi [A] time = 0.05, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2619}

$$\frac{2cd(c \sec(a+bx))^{3/2}}{3b(d \csc(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(5/2)/Sqrt[d*Csc[a + b*x]],x]

[Out] (2*c*d*(c*Sec[a + b*x])^(3/2))/(3*b*(d*Csc[a + b*x])^(3/2))

Rule 2619

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]

Rubi steps

$$\int \frac{(c \sec(a+bx))^{5/2}}{\sqrt{d} \csc(a+bx)} dx = \frac{2cd(c \sec(a+bx))^{3/2}}{3b(d \csc(a+bx))^{3/2}}$$

Mathematica [A] time = 0.11, size = 33, normalized size = 1.00

$$\frac{2cd(c \sec(a+bx))^{3/2}}{3b(d \csc(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(5/2)/Sqrt[d*Csc[a + b*x]],x]

[Out] (2*c*d*(c*Sec[a + b*x])^(3/2))/(3*b*(d*Csc[a + b*x])^(3/2))

fricas [B] time = 0.78, size = 58, normalized size = 1.76

$$\frac{2 \left(c^2 \cos(bx + a)^2 - c^2 \right) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{3bd \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x, algorithm="fricas")

[Out] -2/3*(c^2*cos(b*x + a)^2 - c^2)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*d*cos(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{5}{2}}}{\sqrt{d} \csc(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(5/2)/sqrt(d*csc(b*x + a)), x)

maple [A] time = 0.99, size = 42, normalized size = 1.27

$$\frac{2 \left(\frac{c}{\cos(bx+a)} \right)^{\frac{5}{2}} \cos(bx + a) \sin(bx + a)}{3b \sqrt{\frac{d}{\sin(bx+a)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x)

[Out] 2/3/b*(c/cos(b*x+a))^(5/2)*cos(b*x+a)*sin(b*x+a)/(d/sin(b*x+a))^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{5}{2}}}{\sqrt{d} \csc(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(5/2)/sqrt(d*csc(b*x + a)), x)

mupad [B] time = 0.86, size = 66, normalized size = 2.00

$$\frac{c^2 \sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}} (\cos(a+bx) - \cos(3a+3bx))}{3bd(\cos(2a+2bx)+1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(1/2),x)

[Out] (c^2*(c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)*(cos(a + b*x) - cos(3*a + 3*b*x)))/(3*b*d*(cos(2*a + 2*b*x) + 1))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(5/2)/(d*csc(b*x+a))**(1/2),x)

[Out] Timed out

$$3.252 \quad \int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{3/2}} dx$$

Optimal. Leaf size=98

$$\frac{2c(c \sec(a+bx))^{3/2}}{3bd\sqrt{d \csc(a+bx)}} - \frac{c^2\sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3bd^2}$$

[Out] $2/3*c*(c*\sec(b*x+a))^(3/2)/b/d/(d*\csc(b*x+a))^(1/2)+1/3*c^2*(\sin(a+1/4*\pi+b*x)^2)^(1/2)/\sin(a+1/4*\pi+b*x)*\text{EllipticF}(\cos(a+1/4*\pi+b*x), 2^(1/2))*(d*\csc(b*x+a))^(1/2)*(c*\sec(b*x+a))^(1/2)*\sin(2*b*x+2*a)^(1/2)/b/d^2$

Rubi [A] time = 0.15, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2624, 2630, 2573, 2641}

$$\frac{2c(c \sec(a+bx))^{3/2}}{3bd\sqrt{d \csc(a+bx)}} - \frac{c^2\sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3bd^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sec}[a + b*x])^(5/2)/(d*\text{Csc}[a + b*x])^(3/2), x]$

[Out] $(2*c*(c*\text{Sec}[a + b*x])^(3/2))/(3*b*d*\text{Sqrt}[d*\text{Csc}[a + b*x]]) - (c^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{EllipticF}[a - \pi/4 + b*x, 2]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(3*b*d^2)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f, x\}$

Rule 2624

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] \rightarrow \text{Simp}[(b*(a*\text{Csc}[e + f*x])^(m+1)*(b*\text{Sec}[e + f*x])^(n-1))/(f*a*(n-1)), x] + \text{Dist}[(b^2*(m+1))/(a^2*(n-1)), \text{Int}[(a*\text{Csc}[e + f*x])^(m+2)*(b*\text{Sec}[e + f*x])^(n-2), x], x] /; \text{FreeQ}\{a, b, e, f, x\} \&\& \text{GtQ}[n, 1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] \rightarrow \text{Dist}[(a*\text{Csc}[e + f*x])^m*(b*\text{Sec}[e + f*x])^n*(a*\text{Sin}[e + f*x])$

)^m*(b*cos[e + f*x])ⁿ, Int[1/((a*sin[e + f*x])^m*(b*cos[e + f*x])ⁿ), x],
 x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c -
 Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{3/2}} dx &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d} \csc(a + bx)} - \frac{c^2 \int \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)} dx}{3d^2} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d} \csc(a + bx)} - \frac{(c^2 \sqrt{c} \cos(a + bx) \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)} \sqrt{d} \sin(a + bx))}{3d^2} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d} \csc(a + bx)} - \frac{(c^2 \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)})}{3d^2} \int \frac{1}{\sqrt{\sin(2a + 2bx)}} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd\sqrt{d} \csc(a + bx)} - \frac{c^2 \sqrt{d} \csc(a + bx) F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3bd^2} \end{aligned}$$

Mathematica [C] time = 0.53, size = 70, normalized size = 0.71

$$\frac{c(c \sec(a + bx))^{3/2} \left((-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right) + 2 \right)}{3bd\sqrt{d} \csc(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(3/2), x]

[Out] (c*(2 + (-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*(c*Sec[a + b*x])^(3/2))/(3*b*d*Sqrt[d*Csc[a + b*x]])

fricas [F] time = 0.69, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d} \csc(bx + a) \sqrt{c \sec(bx + a)} c^2 \sec(bx + a)^2}{d^2 \csc(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*c^2*sec(b*x + a)^2/(d^2*csc(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec (bx + a))^{\frac{5}{2}}}{(d \csc (bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(3/2), x)

maple [A] time = 1.07, size = 190, normalized size = 1.94

$$\frac{\left(\sin (bx + a) \cos (bx + a) \sqrt{\frac{1-\cos (bx+a)+\sin (bx+a)}{\sin (bx+a)}} \sqrt{\frac{-1+\cos (bx+a)+\sin (bx+a)}{\sin (bx+a)}} \sqrt{\frac{-1+\cos (bx+a)}{\sin (bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos (bx+a)+\sin (bx+a)}{\sin (bx+a)}}\right)\right)}{3b(-1+\cos (bx+a))\left(\frac{d}{\sin (bx+a)}\right)^{\frac{3}{2}} \sin (bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x)

[Out] 1/3/b*(sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+cos(b*x+a)*2^(1/2)-2^(1/2))*cos(b*x+a)*(c/cos(b*x+a))^(5/2)/(-1+cos(b*x+a))/(d/sin(b*x+a))^(3/2)/sin(b*x+a)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec (bx + a))^{\frac{5}{2}}}{(d \csc (bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}}{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(3/2), x)

[Out] int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))**(5/2)/(d*csc(b*x+a))**(3/2), x)

[Out] Timed out

$$3.253 \quad \int \frac{(c \sec(a+bx))^{5/2}}{(d \csc(a+bx))^{5/2}} dx$$

Optimal. Leaf size=329

$$\frac{c^2 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c^2 \tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c^2 \sqrt{c \sec(a+bx)}}{2 \sqrt{d \csc(a+bx)}}$$

[Out] $2/3 * c * (c * \sec(b * x + a))^{3/2} / b / d / (d * \csc(b * x + a))^{3/2} - 1/2 * c^2 * \arctan(-1 + 2^{(1/2)} * \tan(b * x + a)^{(1/2)}) * (c * \sec(b * x + a))^{1/2} / b / d^2 * 2^{(1/2)} / (d * \csc(b * x + a))^{1/2} / \tan(b * x + a)^{(1/2)} - 1/2 * c^2 * \arctan(1 + 2^{(1/2)} * \tan(b * x + a)^{(1/2)}) * (c * \sec(b * x + a))^{1/2} / b / d^2 * 2^{(1/2)} / (d * \csc(b * x + a))^{1/2} / \tan(b * x + a)^{(1/2)} - 1/4 * c^2 * \ln(1 - 2^{(1/2)} * \tan(b * x + a)^{(1/2)} + \tan(b * x + a)) * (c * \sec(b * x + a))^{1/2} / b / d^2 * 2^{(1/2)} / (d * \csc(b * x + a))^{1/2} / \tan(b * x + a)^{(1/2)} + 1/4 * c^2 * \ln(1 + 2^{(1/2)} * \tan(b * x + a)^{(1/2)} + \tan(b * x + a)) * (c * \sec(b * x + a))^{1/2} / b / d^2 * 2^{(1/2)} / (d * \csc(b * x + a))^{1/2} / \tan(b * x + a)^{(1/2)}$

Rubi [A] time = 0.22, antiderivative size = 329, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2624, 2629, 3476, 329, 297, 1162, 617, 204, 1165, 628}

$$\frac{c^2 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c^2 \tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{\sqrt{2} b d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{c^2 \sqrt{c \sec(a+bx)}}{2 \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(5/2), x]

[Out] $(2 * c * (c * \text{Sec}[a + b * x])^{3/2}) / (3 * b * d * (d * \text{Csc}[a + b * x])^{3/2}) + (c^2 * \text{ArcTan}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b * x]]] * \text{Sqrt}[c * \text{Sec}[a + b * x]]) / (\text{Sqrt}[2] * b * d^2 * \text{Sqrt}[d * \text{Csc}[a + b * x]] * \text{Sqrt}[\text{Tan}[a + b * x]]) - (c^2 * \text{ArcTan}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b * x]]] * \text{Sqrt}[c * \text{Sec}[a + b * x]]) / (\text{Sqrt}[2] * b * d^2 * \text{Sqrt}[d * \text{Csc}[a + b * x]] * \text{Sqrt}[\text{Tan}[a + b * x]]) - (c^2 * \text{Log}[1 - \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b * x]] + \text{Tan}[a + b * x]] * \text{Sqrt}[c * \text{Sec}[a + b * x]]) / (2 * \text{Sqrt}[2] * b * d^2 * \text{Sqrt}[d * \text{Csc}[a + b * x]] * \text{Sqrt}[\text{Tan}[a + b * x]]) + (c^2 * \text{Log}[1 + \text{Sqrt}[2] * \text{Sqrt}[\text{Tan}[a + b * x]] + \text{Tan}[a + b * x]] * \text{Sqrt}[c * \text{Sec}[a + b * x]]) / (2 * \text{Sqrt}[2] * b * d^2 * \text{Sqrt}[d * \text{Csc}[a + b * x]] * \text{Sqrt}[\text{Tan}[a + b * x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 297

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2624

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))
/(f*a*(n - 1)), x] + Dist[(b^2*(m + 1))/(a^2*(n - 1)), Int[(a*Csc[e + f*x])
```

$^{(m+2)}(b \sec[e + f x])^{(n-2)}, x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegersQ}[2m, 2n]$

Rule 2629

$\text{Int}[(\text{csc}[e] + (f)(x)](a))^{(m)}((b)\sec[e + f x])^{(n)}, x_Symbol] \rightarrow \text{Dist}[(a \text{Csc}[e + f x])^m (b \sec[e + f x])^n / \text{Tan}[e + f x]^n, \text{Int}[\text{Tan}[e + f x]^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \ \&\& \ !\text{IntegerQ}[n] \ \&\& \ \text{EqQ}[m + n, 0]$

Rule 3476

$\text{Int}[(b)\tan[(c) + (d)(x)]^{(n)}, x_Symbol] \rightarrow \text{Dist}[b/d, \text{Subst}[\text{Int}[x^n / (b^2 + x^2), x], x, b \text{Tan}[c + d x]], x] /; \text{FreeQ}[\{b, c, d, n\}, x] \ \&\& \ !\text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned} \int \frac{(c \sec(a + bx))^{5/2}}{(d \csc(a + bx))^{5/2}} dx &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{c^2 \int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d \csc(a + bx)}} dx}{d^2} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{(c^2 \sqrt{c \sec(a + bx)}) \int \sqrt{\tan(a + bx)} dx}{d^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{(c^2 \sqrt{c \sec(a + bx)}) \text{Subst}\left(\int \frac{\sqrt{x}}{1+x^2} dx, x, \tan(a + bx)\right)}{bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{(2c^2 \sqrt{c \sec(a + bx)}) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} + \frac{(c^2 \sqrt{c \sec(a + bx)}) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{(c^2 \sqrt{c \sec(a + bx)}) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{\tan(a + bx)}\right)}{2bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{(c^2 \sqrt{c \sec(a + bx)}) \text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a + bx)}\right)}{2bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} - \frac{c^2 \log\left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c \sec(a + bx)}}{2\sqrt{2} bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} + \\ &= \frac{2c(c \sec(a + bx))^{3/2}}{3bd(d \csc(a + bx))^{3/2}} + \frac{c^2 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2} bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} - \frac{c^2 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{\sqrt{2} bd^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}} \end{aligned}$$

Mathematica [A] time = 1.80, size = 280, normalized size = 0.85

$$c(c \sec(a + bx))^{3/2} \left(8 \cos(a + bx) \sqrt[4]{\cot^2(a + bx)} \cot(a + bx) - 8 \sqrt[4]{\cot^2(a + bx)} \csc(a + bx) + 3\sqrt{2} \cos(a + bx) \right)$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sec[a + b*x])^(5/2)/(d*Csc[a + b*x])^(5/2),x]

[Out]
$$\begin{aligned} & -1/12*(c*(6*\text{Sqrt}[2]*\text{ArcTan}[1 - \text{Sqrt}[2]*(\text{Cot}[a + b*x]^2)^{(1/4)}]*\text{Cos}[a + b*x] \\ & * \text{Cot}[a + b*x] - 6*\text{Sqrt}[2]*\text{ArcTan}[1 + \text{Sqrt}[2]*(\text{Cot}[a + b*x]^2)^{(1/4)}]*\text{Cos}[a \\ & + b*x]*\text{Cot}[a + b*x] + 8*\text{Cos}[a + b*x]*\text{Cot}[a + b*x]*(\text{Cot}[a + b*x]^2)^{(1/4)} - \\ & 8*(\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Csc}[a + b*x] + 3*\text{Sqrt}[2]*\text{Cos}[a + b*x]*\text{Cot}[a + b*x] \\ & * \text{Log}[1 - \text{Sqrt}[2]*(\text{Cot}[a + b*x]^2)^{(1/4)} + \text{Sqrt}[\text{Cot}[a + b*x]^2]] - 3*\text{Sqrt}[2] \\ & * \text{Cos}[a + b*x]*\text{Cot}[a + b*x]*\text{Log}[1 + \text{Sqrt}[2]*(\text{Cot}[a + b*x]^2)^{(1/4)} + \text{Sqrt}[\text{Co} \\ & \text{t}[a + b*x]^2])*(c*\text{Sec}[a + b*x])^{(3/2)})/(b*d^2*(\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Sqrt}[\\ & d*\text{Csc}[a + b*x]]) \end{aligned}$$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{5}{2}}}{(d \csc(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(5/2), x)

maple [C] time = 1.19, size = 536, normalized size = 1.63

$$\left(3i \cos(bx + a) \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \text{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \dots \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2), x)`

[Out] $\frac{1}{6} \frac{1}{b} \left(3 I \cos(bx+a) \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} - \frac{1}{2} I, \frac{1}{2} \sqrt{2} \right)^{1/2} - 3 I \cos(bx+a) \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} + \frac{1}{2} I, \frac{1}{2} \sqrt{2} \right)^{1/2} - 3 \cos(bx+a) \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} - \frac{1}{2} I, \frac{1}{2} \sqrt{2} \right)^{1/2} - 3 \cos(bx+a) \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} + \frac{1}{2} I, \frac{1}{2} \sqrt{2} \right)^{1/2} + 2 \cos(bx+a) \sqrt{2} - 2 \sqrt{2} \right) \cos(bx+a) \left(\frac{c}{\cos(bx+a)} \right)^{5/2} / (-1+\cos(bx+a)) / (d/\sin(bx+a))^{5/2} / \sin(bx+a) \sqrt{2} \right)^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sec(bx + a))^{\frac{5}{2}}}{(d \csc(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sec(b*x+a))^(5/2)/(d*csc(b*x+a))^(5/2), x, algorithm="maxima")`

[Out] `integrate((c*sec(b*x + a))^(5/2)/(d*csc(b*x + a))^(5/2), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{c}{\cos(a+bx)} \right)^{5/2}}{\left(\frac{d}{\sin(a+bx)} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(5/2), x)`

[Out] `int((c/cos(a + b*x))^(5/2)/(d/sin(a + b*x))^(5/2), x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sec(b*x+a))**(5/2)/(d*csc(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```

$$3.254 \quad \int \frac{(d \csc(a+bx))^{9/2}}{\sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=69

$$\frac{8cd^3(d \csc(a+bx))^{3/2}}{21b(c \sec(a+bx))^{3/2}} - \frac{2cd(d \csc(a+bx))^{7/2}}{7b(c \sec(a+bx))^{3/2}}$$

[Out] $-8/21*c*d^3*(d*\csc(b*x+a))^{3/2}/b/(c*\sec(b*x+a))^{3/2}-2/7*c*d*(d*\csc(b*x+a))^{7/2}/b/(c*\sec(b*x+a))^{3/2}$

Rubi [A] time = 0.10, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2625, 2619}

$$\frac{8cd^3(d \csc(a+bx))^{3/2}}{21b(c \sec(a+bx))^{3/2}} - \frac{2cd(d \csc(a+bx))^{7/2}}{7b(c \sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(9/2)/Sqrt[c*Sec[a + b*x]], x]

[Out] $(-8*c*d^3*(d*Csc[a + b*x])^{3/2})/(21*b*(c*Sec[a + b*x])^{3/2}) - (2*c*d*(d*Csc[a + b*x])^{7/2})/(7*b*(c*Sec[a + b*x])^{3/2})$

Rule 2619

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]

Rule 2625

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Dist[(a^2*(m + n - 2))/(m - 1), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rubi steps

$$\int \frac{(d \csc(a + bx))^{9/2}}{\sqrt{c \sec(a + bx)}} dx = -\frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{3/2}} + \frac{1}{7} (4d^2) \int \frac{(d \csc(a + bx))^{5/2}}{\sqrt{c \sec(a + bx)}} dx$$

$$= -\frac{8cd^3(d \csc(a + bx))^{3/2}}{21b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{7/2}}{7b(c \sec(a + bx))^{3/2}}$$

Mathematica [A] time = 0.27, size = 45, normalized size = 0.65

$$\frac{2cd(2 \cos(2(a + bx)) - 5)(d \csc(a + bx))^{7/2}}{21b(c \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(9/2)/Sqrt[c*Sec[a + b*x]], x]

[Out] (2*c*d*(-5 + 2*Cos[2*(a + b*x)])*(d*Csc[a + b*x])^(7/2))/(21*b*(c*Sec[a + b*x])^(3/2))

fricas [A] time = 0.59, size = 79, normalized size = 1.14

$$-\frac{2(4d^4 \cos(bx + a)^4 - 7d^4 \cos(bx + a)^2) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{21(bc \cos(bx + a)^2 - bc) \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2), x, algorithm="fricas")

[Out] -2/21*(4*d^4*cos(b*x + a)^4 - 7*d^4*cos(b*x + a)^2)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/((b*c*cos(b*x + a)^2 - b*c)*sin(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{9/2}}{\sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2), x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(9/2)/sqrt(c*sec(b*x + a)), x)

maple [A] time = 1.21, size = 54, normalized size = 0.78

$$\frac{2 \left(4 \left(\cos^2 (bx + a) \right) - 7 \right) \left(\frac{d}{\sin(bx+a)} \right)^{\frac{9}{2}} \cos (bx + a) \sin (bx + a)}{21b \sqrt{\frac{c}{\cos(bx+a)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2),x)`

[Out] `2/21/b*(4*cos(b*x+a)^2-7)*(d/sin(b*x+a))^(9/2)*cos(b*x+a)*sin(b*x+a)/(c/cos(b*x+a))^(1/2)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc (bx + a))^{\frac{9}{2}}}{\sqrt{c \sec (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate((d*csc(b*x + a))^(9/2)/sqrt(c*sec(b*x + a)), x)`

mupad [B] time = 1.83, size = 99, normalized size = 1.43

$$\frac{8 d^4 \sqrt{\frac{d}{\sin(a+bx)}} (11 \sin (2 a + 2 b x) - 7 \sin (4 a + 4 b x) + \sin (6 a + 6 b x))}{21 b \sqrt{\frac{c}{\cos(a+bx)}} (15 \cos (2 a + 2 b x) - 6 \cos (4 a + 4 b x) + \cos (6 a + 6 b x) - 10)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(1/2),x)`

[Out] `(8*d^4*(d/sin(a + b*x))^(1/2)*(11*sin(2*a + 2*b*x) - 7*sin(4*a + 4*b*x) + sin(6*a + 6*b*x)))/(21*b*(c/cos(a + b*x))^(1/2)*(15*cos(2*a + 2*b*x) - 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) - 10))`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))**(9/2)/(c*sec(b*x+a))**(1/2),x)`

[Out] Timed out

$$3.255 \quad \int \frac{(d \csc(a+bx))^{7/2}}{\sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=128

$$\frac{4d^4 E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{5b\sqrt{\sin(2a+2bx)}\sqrt{c \sec(a+bx)}\sqrt{d \csc(a+bx)}} - \frac{4cd^3\sqrt{d \csc(a+bx)}}{5b(c \sec(a+bx))^{3/2}} - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

[Out] $-2/5*c*d*(d*\csc(b*x+a))^{(5/2)}/b/(c*\sec(b*x+a))^{(3/2)}-4/5*c*d^3*(d*\csc(b*x+a))^{(1/2)}/b/(c*\sec(b*x+a))^{(3/2)}+4/5*d^4*(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticE}(\cos(a+1/4*\text{Pi}+b*x),2^{(1/2)})/b/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.19, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2625, 2630, 2572, 2639}

$$\frac{4cd^3\sqrt{d \csc(a+bx)}}{5b(c \sec(a+bx))^{3/2}} - \frac{4d^4 E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{5b\sqrt{\sin(2a+2bx)}\sqrt{c \sec(a+bx)}\sqrt{d \csc(a+bx)}} - \frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(7/2)/Sqrt[c*Sec[a + b*x]],x]

[Out] $(-4*c*d^3*\text{Sqrt}[d*\text{Csc}[a+b*x]])/(5*b*(c*\text{Sec}[a+b*x])^{(3/2)}) - (2*c*d*(d*\text{Csc}[a+b*x])^{(5/2)})/(5*b*(c*\text{Sec}[a+b*x])^{(3/2)}) - (4*d^4*\text{EllipticE}[a-\text{Pi}/4+b*x,2])/(5*b*\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{Sqrt}[c*\text{Sec}[a+b*x]]*\text{Sqrt}[\text{Sin}[2*a+2*b*x]])$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[2*e + 2*f*x]], Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2625

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> -Simp[(a*b*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n-1))/(f*(m-1)), x] + Dist[(a^2*(m+n-2))/(m-1), Int[(a*Csc[e + f*x])^(m-2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2639

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{7/2}}{\sqrt{c \sec(a + bx)}} dx &= -\frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} + \frac{1}{5} (2d^2) \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx \\ &= -\frac{4cd^3 \sqrt{d \csc(a + bx)}}{5b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} - \frac{1}{5} (4d^4) \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx \\ &= -\frac{4cd^3 \sqrt{d \csc(a + bx)}}{5b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} - \frac{(4d^4) \int \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}}{5\sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx \\ &= -\frac{4cd^3 \sqrt{d \csc(a + bx)}}{5b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} - \frac{(4d^4) \int \sqrt{\sin(2a + 2bx)} dx}{5\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \\ &= -\frac{4cd^3 \sqrt{d \csc(a + bx)}}{5b(c \sec(a + bx))^{3/2}} - \frac{2cd(d \csc(a + bx))^{5/2}}{5b(c \sec(a + bx))^{3/2}} - \frac{4d^4 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{5b\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] time = 1.08, size = 104, normalized size = 0.81

$$\frac{2d^2 \tan^2(a + bx)(d \csc(a + bx))^{3/2} \left(\sin(2(a + bx)) \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) - (\cos(2(a + bx))) \right)}{5b\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Csc[a + b*x])^(7/2)/Sqrt[c*Sec[a + b*x]], x]
```

```
[Out] (-2*d^2*(d*Csc[a + b*x])^(3/2)*(-((-2 + Cos[2*(a + b*x)])*Cot[a + b*x]^3) + (-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2]*Sin[2*(a + b*x)]*Tan[a + b*x]^2)/(5*b*Sqrt[c*Sec[a + b*x]])
```


fricas [F] time = 0.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d} \csc(bx+a) \sqrt{c \sec(bx+a)} d^3 \csc(bx+a)^3}{c \sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d^3*csc(b*x + a)^3/(c*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx+a))^{\frac{7}{2}}}{\sqrt{c \sec(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(7/2)/sqrt(c*sec(b*x + a)), x)

maple [B] time = 1.23, size = 976, normalized size = 7.62

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x)

[Out]
$$\begin{aligned} & -1/5/b*(4*\cos(b*x+a)^3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*E \\ & \text{llipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})-2*\cos(b*x+a)^3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*E \\ & \text{llipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})+4*\cos(b*x+a)^2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*E \\ & \text{llipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})-2*\cos(b*x+a)^2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*E \\ & \text{llipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})-4*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*E \\ & \text{llipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})+2*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2} \end{aligned}$$

$\sin(b*x+a)/\sin(b*x+a)^{(1/2)*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)*\text{EllipticF}((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2))}-2*\cos(b*x+a)^3*2^{(1/2)}-4*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*\text{EllipticE}((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2))}+2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*\text{EllipticF}((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2))}+2^{(1/2)*\cos(b*x+a)^2+2*\cos(b*x+a)*2^{(1/2))}*(d/\sin(b*x+a))^{(7/2)*\sin(b*x+a)/(c/\cos(b*x+a))^{(1/2)}/\cos(b*x+a)*2^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{7}{2}}}{\sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(7/2)/sqrt(c*sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{7/2}}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(1/2),x)

[Out] int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(1/2),x)

[Out] Timed out

$$3.256 \quad \int \frac{(d \csc(a+bx))^{5/2}}{\sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=33

$$-\frac{2cd(d \csc(a+bx))^{3/2}}{3b(c \sec(a+bx))^{3/2}}$$

[Out] $-2/3*c*d*(d*\csc(b*x+a))^{3/2}/b/(c*\sec(b*x+a))^{3/2}$

Rubi [A] time = 0.05, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2619}

$$-\frac{2cd(d \csc(a+bx))^{3/2}}{3b(c \sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(5/2)/Sqrt[c*Sec[a + b*x]],x]

[Out] $(-2*c*d*(d*Csc[a + b*x])^{3/2})/(3*b*(c*Sec[a + b*x])^{3/2})$

Rule 2619

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]

Rubi steps

$$\int \frac{(d \csc(a+bx))^{5/2}}{\sqrt{c \sec(a+bx)}} dx = -\frac{2cd(d \csc(a+bx))^{3/2}}{3b(c \sec(a+bx))^{3/2}}$$

Mathematica [A] time = 0.12, size = 33, normalized size = 1.00

$$-\frac{2cd(d \csc(a+bx))^{3/2}}{3b(c \sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(5/2)/Sqrt[c*Sec[a + b*x]],x]

[Out] $(-2*c*d*(d*Csc[a + b*x])^{3/2})/(3*b*(c*Sec[a + b*x])^{3/2})$

fricas [A] time = 0.77, size = 51, normalized size = 1.55

$$\frac{2 d^2 \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx+a)^2}{3bc \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] -2/3*d^2*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^2/(b*c*sin(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{5}{2}}}{\sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(5/2)/sqrt(c*sec(b*x + a)), x)

maple [A] time = 1.16, size = 42, normalized size = 1.27

$$\frac{2 \left(\frac{d}{\sin(bx+a)} \right)^{\frac{5}{2}} \cos(bx+a) \sin(bx+a)}{3b \sqrt{\frac{c}{\cos(bx+a)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x)

[Out] -2/3/b*(d/sin(b*x+a))^(5/2)*cos(b*x+a)*sin(b*x+a)/(c/cos(b*x+a))^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{5}{2}}}{\sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(5/2)/sqrt(c*sec(b*x + a)), x)

mupad [B] time = 0.80, size = 49, normalized size = 1.48

$$-\frac{d^2 \sin(2a + 2bx) \sqrt{\frac{d}{\sin(a+bx)}}}{3b \sin(a + bx)^2 \sqrt{\frac{c}{\cos(a+bx)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(1/2),x)

[Out] -(d^2*sin(2*a + 2*b*x)*(d/sin(a + b*x))^(1/2))/(3*b*sin(a + b*x)^2*(c/cos(a + b*x))^(1/2))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(1/2),x)

[Out] Timed out

$$3.257 \quad \int \frac{(d \csc(a+bx))^{3/2}}{\sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=89

$$-\frac{2d^2 E\left(a+bx-\frac{\pi}{4} \middle| 2\right)}{b\sqrt{\sin(2a+2bx)}\sqrt{c \sec(a+bx)}\sqrt{d \csc(a+bx)}} - \frac{2cd\sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}}$$

[Out] $-2*c*d*(d*csc(b*x+a))^{(1/2)}/b/(c*sec(b*x+a))^{(3/2)+2*d^2*(\sin(a+1/4*Pi+b*x))^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*EllipticE(\cos(a+1/4*Pi+b*x),2^{(1/2)})/b/(d*csc(b*x+a))^{(1/2)}/(c*sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.14, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2625, 2630, 2572, 2639}

$$-\frac{2d^2 E\left(a+bx-\frac{\pi}{4} \middle| 2\right)}{b\sqrt{\sin(2a+2bx)}\sqrt{c \sec(a+bx)}\sqrt{d \csc(a+bx)}} - \frac{2cd\sqrt{d \csc(a+bx)}}{b(c \sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(3/2)/Sqrt[c*Sec[a + b*x]], x]

[Out] $(-2*c*d*Sqrt[d*Csc[a + b*x]])/(b*(c*Sec[a + b*x])^{(3/2)}) - (2*d^2*EllipticE[a - Pi/4 + b*x, 2])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[\sin[2*a + 2*b*x]])$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[\sin[2*e + 2*f*x]]], Int[Sqrt[\sin[2*e + 2*f*x]]], x, x] /; FreeQ[{a, b, e, f}, x]

Rule 2625

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> -Simp[(a*b*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n-1))/(f*(m-1)), x] + Dist[(a^2*(m+n-2))/(m-1), Int[(a*Csc[e + f*x])^(m-2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])

$\int (b \cos(e + f x))^n \int \frac{1}{(a \sin(e + f x))^m (b \cos(e + f x))^n} dx, x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \} \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin(c + d x)], x_Symbol] \rightarrow \text{Simp}[(2 \text{EllipticE}[(1/2)(c - P i/2 + d x)], x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx &= -\frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} - (2d^2) \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx \\ &= -\frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} - \frac{(2d^2) \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{\sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)}} \\ &= -\frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} - \frac{(2d^2) \int \sqrt{\sin(2a + 2bx)} dx}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \\ &= -\frac{2cd\sqrt{d \csc(a + bx)}}{b(c \sec(a + bx))^{3/2}} - \frac{2d^2 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{b\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] time = 0.48, size = 80, normalized size = 0.90

$$\frac{2d^2 \tan(a + bx) \left(\sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) + \cot^2(a + bx) \right)}{b\sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(3/2)/Sqrt[c*Sec[a + b*x]],x]

[Out] (-2*d^2*(Cot[a + b*x]^2 + (-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Tan[a + b*x])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])

fricas [F] time = 0.68, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} d \csc(bx + a)}{c \sec(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")
 [Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d*csc(b*x + a)/(c*sec(b*x + a)), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc (bx + a))^{\frac{3}{2}}}{\sqrt{c \sec (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")
 [Out] integrate((d*csc(b*x + a))^(3/2)/sqrt(c*sec(b*x + a)), x)
maple [B] time = 1.20, size = 494, normalized size = 5.55

$$\left(2 \cos (bx + a) \sqrt{\frac{1-\cos (bx+a)+\sin (bx+a)}{\sin (bx+a)}} \sqrt{\frac{-1+\cos (bx+a)+\sin (bx+a)}{\sin (bx+a)}} \sqrt{\frac{-1+\cos (bx+a)}{\sin (bx+a)}} \text{EllipticE}\left(\sqrt{\frac{1-\cos (bx+a)+\sin (bx+a)}{\sin (bx+a)}}, \frac{\sqrt{2}}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x)
 [Out] 1/b*(2*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-cos(b*x+a)*2^(1/2)*(d/sin(b*x+a))^(3/2)*sin(b*x+a)/(c/cos(b*x+a))^(1/2)/cos(b*x+a)*2^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc (bx + a))^{\frac{3}{2}}}{\sqrt{c \sec (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(3/2)/sqrt(c*sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(1/2),x)

[Out] int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(1/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(1/2),x)

[Out] Timed out

$$3.258 \quad \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=270

$$\frac{\sqrt{\tan(a+bx)} \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)}}{\sqrt{2} b \sqrt{c \sec(a+bx)}} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} b \sqrt{c \sec(a+bx)}}$$

[Out] 1/2*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/2*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)-1/4*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)+1/4*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(d*csc(b*x+a))^(1/2)*tan(b*x+a)^(1/2)/b*2^(1/2)/(c*sec(b*x+a))^(1/2)

Rubi [A] time = 0.13, antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{\sqrt{\tan(a+bx)} \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)}}{\sqrt{2} b \sqrt{c \sec(a+bx)}} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} b \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Csc[a + b*x]]/Sqrt[c*Sec[a + b*x]],x]

[Out] -((ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])/(Sqrt[2]*b*Sqrt[c*Sec[a + b*x]])) + (ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])/(Sqrt[2]*b*Sqrt[c*Sec[a + b*x]]) - (Sqrt[d*Csc[a + b*x]]*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[Tan[a + b*x]])/(2*Sqrt[2]*b*Sqrt[c*Sec[a + b*x]]) + (Sqrt[d*Csc[a + b*x]]*Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[Tan[a + b*x]])/(2*Sqrt[2]*b*Sqrt[c*Sec[a + b*x]])

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}

, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 329

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n))/c^n)^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 617

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 628

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1162

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1165

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] :=> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && ! IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx &= \frac{(\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{\sqrt{c \sec(a+bx)}} \\
 &= \frac{(\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \operatorname{Subst}\left(\int \frac{1}{\sqrt{x(1+x^2)}} dx, x, \tan(a+bx)\right)}{b\sqrt{c \sec(a+bx)}} \\
 &= \frac{(2\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \operatorname{Subst}\left(\int \frac{1}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{b\sqrt{c \sec(a+bx)}} \\
 &= \frac{(\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \operatorname{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a+bx)}\right)}{b\sqrt{c \sec(a+bx)}} + \frac{(\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)})}{\sqrt{c \sec(a+bx)}} \\
 &= \frac{(\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \operatorname{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a+bx)}\right)}{2b\sqrt{c \sec(a+bx)}} + \frac{(\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)})}{\sqrt{c \sec(a+bx)}} \\
 &= -\frac{\sqrt{d \csc(a+bx)} \log\left(1 - \sqrt{2} \sqrt{\tan(a+bx)} + \tan(a+bx)\right) \sqrt{\tan(a+bx)}}{2\sqrt{2} b\sqrt{c \sec(a+bx)}} + \frac{\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{\sqrt{c \sec(a+bx)}} \\
 &= -\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{\sqrt{2} b\sqrt{c \sec(a+bx)}} + \frac{\tan^{-1}\left(1 + \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}}{\sqrt{2} b\sqrt{c \sec(a+bx)}}
 \end{aligned}$$

Mathematica [C] time = 0.13, size = 55, normalized size = 0.20

$$\frac{2 \cot(a+bx) \sqrt{d \csc(a+bx)} {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a+bx)\right)}{3b\sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[a + b*x]]/Sqrt[c*Sec[a + b*x]],x]

[Out] (-2*Cot[a + b*x]*Sqrt[d*Csc[a + b*x]]*Hypergeometric2F1[3/4, 1, 7/4, -Cot[a + b*x]^2])/(3*b*Sqrt[c*Sec[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(bx+a)}}{\sqrt{c \sec(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(b*x + a))/sqrt(c*sec(b*x + a)), x)

maple [C] time = 1.24, size = 316, normalized size = 1.17

$$\sqrt{\frac{d}{\sin(bx+a)}} \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \left(i \operatorname{EllipticPi} \left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x)

[Out] $-1/2/b*(d/\sin(b*x+a))^{1/2}*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*(I*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-I*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2}))-2*\operatorname{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})+\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})+\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2}))*\sin(b*x+a)^2/(c/\cos(b*x+a))^{1/2}/\cos(b*x+a)/(-1+\cos(b*x+a))*2^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(bx+a)}}{\sqrt{c \sec(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(b*x + a))/sqrt(c*sec(b*x + a)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{d}{\sin(a+bx)}}}{\sqrt{\frac{c}{\cos(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(1/2),x)

[Out] int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(a + bx)}}{\sqrt{c \sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(1/2),x)

[Out] Integral(sqrt(d*csc(a + b*x))/sqrt(c*sec(a + b*x)), x)

$$3.259 \quad \int \frac{1}{\sqrt{d} \csc(a+bx) \sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=53

$$\frac{E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{b\sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d} \csc(a + bx)}$$

[Out] $-(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticE}(\cos(a+1/4*\text{Pi}+b*x), 2^{(1/2)})/b/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.09, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2630, 2572, 2639}

$$\frac{E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{b\sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d} \csc(a + bx)}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]),x]

[Out] EllipticE[a - Pi/4 + b*x, 2]/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[2*e + 2*f*x]], Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]] , x_Symbol] :> Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx = \frac{\int \sqrt{c \cos(a+bx)} \sqrt{d \sin(a+bx)} dx}{\sqrt{c \cos(a+bx)} \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{d \sin(a+bx)}}$$

$$= \frac{\int \sqrt{\sin(2a+2bx)} dx}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)}}$$

$$= \frac{E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{b \sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)} \sqrt{\sin(2a+2bx)}}$$

Mathematica [C] time = 0.22, size = 66, normalized size = 1.25

$$\frac{\tan(a+bx) \sqrt[4]{-\cot^2(a+bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a+bx)\right)}{b \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]),x]

[Out] ((-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2]*Tan[a + b*x])/(b*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])

fricas [F] time = 0.86, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)}}{cd \csc(bx+a) \sec(bx+a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c*d*csc(b*x + a)*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d \csc(bx+a)} \sqrt{c \sec(bx+a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))), x)

maple [B] time = 1.01, size = 509, normalized size = 9.60

$$\left(2 \cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \text{EllipticE} \left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2), x)

[Out] $-1/2/b*(2*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})-\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})+2*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})-((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})+2^{1/2}*\cos(b*x+a)^2-\cos(b*x+a)*2^{1/2})/\cos(b*x+a)/\sin(b*x+a)/(d/\sin(b*x+a))^{1/2}/(c/\cos(b*x+a))^{1/2}*2^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2), x, algorithm="maxima")

[Out] integrate(1/(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}} \sqrt{\frac{d}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)),x)`

[Out] `int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(1/2)), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(1/2),x)`

[Out] `Integral(1/(sqrt(c*sec(a + b*x))*sqrt(d*csc(a + b*x))), x)`

$$3.260 \quad \int \frac{1}{(d \csc(a+bx))^{3/2} \sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=322

$$\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{4\sqrt{2} bd^2 \sqrt{c \sec(a+bx)}} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{4\sqrt{2} bd^2 \sqrt{c \sec(a+bx)}}$$

[Out] $-1/2*c/b/d/(c*\sec(b*x+a))^{(3/2)}/(d*\csc(b*x+a))^{(1/2)}+1/8*\arctan(-1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+1/8*\arctan(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-1/16*\ln(1-2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+1/16*\ln(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.20, antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2627, 2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{4\sqrt{2} bd^2 \sqrt{c \sec(a+bx)}} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{4\sqrt{2} bd^2 \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]),x]

[Out] $-c/(2*b*d*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^{(3/2)}) - (\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*Csc[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(4*\text{Sqrt}[2]*b*d^2*\text{Sqrt}[c*Sec[a + b*x]]) + (\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*Csc[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(4*\text{Sqrt}[2]*b*d^2*\text{Sqrt}[c*Sec[a + b*x]]) - (\text{Sqrt}[d*Csc[a + b*x]]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b*d^2*\text{Sqrt}[c*Sec[a + b*x]]) + (\text{Sqrt}[d*Csc[a + b*x]]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b*d^2*\text{Sqrt}[c*Sec[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),

$x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& (\text{GtQ}[a/b, 0] \mid\mid (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 329

$\text{Int}[\{(c_.)*(x_)^m\}*((a_) + (b_.)*(x_)^n)^p, x_Symbol] \rightarrow \text{With}[\{k = \text{Denominator}[m]\}, \text{Dist}[k/c, \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(a + (b*x^{k*n})/c^n)^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 617

$\text{Int}[\{(a_) + (b_.)*(x_) + (c_.)*(x_)^2\}^{-1}, x_Symbol] \rightarrow \text{With}[\{q = 1 - 4*S\text{implify}[(a*c)/b^2]\}, \text{Dist}[-2/b, \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + (2*c*x)/b], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \mid\mid \text{!RationalQ}[b^2 - 4*a*c])] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 628

$\text{Int}[\{(d_) + (e_.)*(x_)\}/\{(a_) + (b_.)*(x_) + (c_.)*(x_)^2\}, x_Symbol] \rightarrow \text{Simp}[(d*\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]])/b, x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 1162

$\text{Int}[\{(d_) + (e_.)*(x_)^2\}/\{(a_) + (c_.)*(x_)^4\}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[(2*d)/e, 2]\}, \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

Rule 1165

$\text{Int}[\{(d_) + (e_.)*(x_)^2\}/\{(a_) + (c_.)*(x_)^4\}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-(2*d)/e, 2]\}, \text{Dist}[e/(2*c*q), \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

Rule 2627

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*\text{sec}[(e_.) + (f_.)*(x_)])^n, x_Symbol] \rightarrow \text{Simp}[(b*(a*\text{Csc}[e + f*x])^{m+1}*(b*\text{Sec}[e + f*x])^{n-1})/(a*f*(m+n)), x] + \text{Dist}[(m+1)/(a^2*(m+n)), \text{Int}[(a*\text{Csc}[e + f*x])^{m+2}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&$

& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} dx &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} + \frac{\int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4d^2} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} + \frac{(\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)})}{4d^2 \sqrt{c \sec(a + bx)}} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} + \frac{(\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)})}{4bd^2} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} + \frac{(\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)})}{2bd^2} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} + \frac{(\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)})}{4bd^2} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} + \frac{(\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)})}{8bd^2} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} - \frac{\sqrt{d} \csc(a + bx) \log\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right)}{8\sqrt{2} bd^2} \\
 &= -\frac{c}{2bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} - \frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right)}{4\sqrt{2} bd^2 \sqrt{c}}
 \end{aligned}$$

Mathematica [C] time = 0.26, size = 66, normalized size = 0.20

$$\frac{\cot(a + bx) \left(\csc^2(a + bx) {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a + bx)\right) + 3 \right)}{6b\sqrt{c \sec(a + bx)} (d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]),x]

[Out] -1/6*(Cot[a + b*x]*(3 + Csc[a + b*x]^2*Hypergeometric2F1[3/4, 1, 7/4, -Cot[a + b*x]^2]))/(b*(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a))), x)

maple [C] time = 0.91, size = 658, normalized size = 2.04

$$\left(i \operatorname{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} - \frac{i}{2}, \frac{\sqrt{2}}{2} \right) \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \sin \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x)

```
[Out] -1/8/b*(I*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)-I*(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)+EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)+((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)-2*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2*cos(b*x+a)^3*2^(1/2)-2*2^(1/2)*cos(b*x+a)^2)/(-1+cos(b*x+a))/(d/sin(b*x+a))^(3/2)/(c/cos(b*x+a))^(1/2)/cos(b*x+a)/sin(b*x+a)*2^(1/2)
```

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} \sqrt{c \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/((d*csc(b*x + a))^(3/2)*sqrt(c*sec(b*x + a))), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}} \left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(3/2)),x)
```

```
[Out] int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(3/2)), x)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sec(a + bx)} (d \csc(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(1/2), x)
```

```
[Out] Integral(1/(sqrt(c*sec(a + b*x))*(d*csc(a + b*x))**(3/2)), x)
```


$$3.261 \quad \int \frac{1}{(d \csc(a+bx))^{5/2} \sqrt{c \sec(a+bx)}} dx$$

Optimal. Leaf size=95

$$\frac{E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{2bd^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} - \frac{c}{3bd(c\sec(a+bx))^{3/2}(d\csc(a+bx))^{3/2}}$$

[Out] $-1/3*c/b/d/(d*csc(b*x+a))^{(3/2)}/(c*sec(b*x+a))^{(3/2)}-1/2*(\sin(a+1/4*Pi+b*x))^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*EllipticE(\cos(a+1/4*Pi+b*x),2^{(1/2)})/b/d^2/(d*csc(b*x+a))^{(1/2)}/(c*sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.14, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2627, 2630, 2572, 2639}

$$\frac{E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{2bd^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} - \frac{c}{3bd(c\sec(a+bx))^{3/2}(d\csc(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]]),x]

[Out] $-c/(3*b*d*(d*Csc[a + b*x])^{(3/2)}*(c*Sec[a + b*x])^{(3/2)}) + EllipticE[a - Pi/4 + b*x, 2]/(2*b*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]*Sqrt[\sin[2*a + 2*b*x]])$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[\sin[2*e + 2*f*x]], Int[Sqrt[\sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2627

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))/(a*f*(m + n)), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])

)^m*(b*cos[e + f*x])^n, Int[1/((a*sin[e + f*x])^m*(b*cos[e + f*x])^n), x],
 x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P
 i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)}} dx &= -\frac{c}{3bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} + \frac{\int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{2d^2} \\ &= -\frac{c}{3bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} + \frac{\int \sqrt{c \cos(a + bx)}}{2d^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)}} \\ &= -\frac{c}{3bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} + \frac{\int \sqrt{\sin(2a + 2bx)}}{2d^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} \\ &= -\frac{c}{3bd(d \csc(a + bx))^{3/2}(c \sec(a + bx))^{3/2}} + \frac{E\left(a - \frac{\pi}{4} + bx\right)}{2bd^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} \end{aligned}$$

Mathematica [C] time = 0.36, size = 84, normalized size = 0.88

$$\frac{\tan(a + bx) \left(-3 \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) + \cos(2(a + bx)) + 1 \right)}{6bd^2 \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(5/2)*Sqrt[c*Sec[a + b*x]]),x]

[Out] -1/6*((1 + Cos[2*(a + b*x)] - 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-
 1/2, 1/4, 1/2, Csc[a + b*x]^2])*Tan[a + b*x])/(b*d^2*Sqrt[d*Csc[a + b*x]]*S
 qrt[c*Sec[a + b*x]])

fricas [F] time = 1.35, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)}}{cd^3 \csc(bx + a)^3 \sec(bx + a)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c*d^3*csc(b*x + a)^3*sec(b*x + a)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc (bx + a))^{\frac{5}{2}} \sqrt{c \sec (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a))), x)

maple [B] time = 1.01, size = 523, normalized size = 5.51

$$\left(2 \left(\cos^4 (bx + a) \right) \sqrt{2} + 3 \cos (bx + a) \sqrt{\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \sqrt{\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \sqrt{\frac{-1 + \cos (bx + a)}{\sin (bx + a)}} \right) \text{EllipticF} \left(\sqrt{\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x)

[Out] 1/12/b*(2*cos(b*x+a)^4*2^(1/2)+3*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))-6*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))+3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))-6*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))-5*2^(1/2)*cos(b*x+a)^2+3*cos(b*x+a)*2^(1/2))/cos(b*x+a)/sin(b*x+a)^3/(d/sin(b*x+a))^(5/2)/(c/cos(b*x+a))^(1/2)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc (bx + a))^{\frac{5}{2}} \sqrt{c \sec (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(1/((d*csc(b*x + a))^(5/2)*sqrt(c*sec(b*x + a))), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{\frac{c}{\cos(a+bx)}} \left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2)),x)

[Out] int(1/((c/cos(a + b*x))^(1/2)*(d/sin(a + b*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(1/2),x)

[Out] Timed out

$$3.262 \quad \int \frac{(d \csc(a+bx))^{11/2}}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=110

$$\frac{8d^5 \sqrt{d \csc(a+bx)}}{45bc \sqrt{c \sec(a+bx)}} + \frac{2d^3 (d \csc(a+bx))^{5/2}}{45bc \sqrt{c \sec(a+bx)}} - \frac{2d (d \csc(a+bx))^{9/2}}{9bc \sqrt{c \sec(a+bx)}}$$

[Out] $2/45*d^3*(d*\csc(b*x+a))^{(5/2)}/b/c/(c*\sec(b*x+a))^{(1/2)}-2/9*d*(d*\csc(b*x+a))^{(9/2)}/b/c/(c*\sec(b*x+a))^{(1/2)}+8/45*d^5*(d*\csc(b*x+a))^{(1/2)}/b/c/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.15, antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2623, 2625, 2619}

$$\frac{8d^5 \sqrt{d \csc(a+bx)}}{45bc \sqrt{c \sec(a+bx)}} + \frac{2d^3 (d \csc(a+bx))^{5/2}}{45bc \sqrt{c \sec(a+bx)}} - \frac{2d (d \csc(a+bx))^{9/2}}{9bc \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(11/2)/(c*Sec[a + b*x])^(3/2),x]

[Out] $(8*d^5*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(45*b*c*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (2*d^3*(d*\text{Csc}[a + b*x])^{(5/2)})/(45*b*c*\text{Sqrt}[c*\text{Sec}[a + b*x]]) - (2*d*(d*\text{Csc}[a + b*x])^{(9/2)})/(9*b*c*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 2619

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(n - 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n - 2, 0] && NeQ[n, 1]

Rule 2623

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(f*b*(m - 1)), x] + Dist[(a^2*(n + 1))/(b^2*(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2625

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n -

1))/ (f*(m - 1)), x] + Dist[(a^2*(m + n - 2))/(m - 1), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])ⁿ, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{11/2}}{(c \sec(a + bx))^{3/2}} dx &= -\frac{2d(d \csc(a + bx))^{9/2}}{9bc\sqrt{c \sec(a + bx)}} - \frac{d^2 \int (d \csc(a + bx))^{7/2} \sqrt{c \sec(a + bx)} dx}{9c^2} \\ &= \frac{2d^3(d \csc(a + bx))^{5/2}}{45bc\sqrt{c \sec(a + bx)}} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc\sqrt{c \sec(a + bx)}} - \frac{(4d^4) \int (d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)} dx}{45c^2} \\ &= \frac{8d^5 \sqrt{d \csc(a + bx)}}{45bc\sqrt{c \sec(a + bx)}} + \frac{2d^3(d \csc(a + bx))^{5/2}}{45bc\sqrt{c \sec(a + bx)}} - \frac{2d(d \csc(a + bx))^{9/2}}{9bc\sqrt{c \sec(a + bx)}} \end{aligned}$$

Mathematica [A] time = 0.29, size = 57, normalized size = 0.52

$$\frac{2d^3(2 \cos(2(a + bx)) - 7) \cot^2(a + bx)(d \csc(a + bx))^{5/2}}{45bc\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(11/2)/(c*Sec[a + b*x])^(3/2), x]

[Out] (2*d³*(-7 + 2*Cos[2*(a + b*x)])*Cot[a + b*x]²*(d*Csc[a + b*x])^(5/2))/(45*b*c*Sqrt[c*Sec[a + b*x]])

fricas [A] time = 2.42, size = 88, normalized size = 0.80

$$\frac{2 \left(4d^5 \cos(bx + a)^5 - 9d^5 \cos(bx + a)^3 \right) \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}}}{45 \left(bc^2 \cos(bx + a)^4 - 2bc^2 \cos(bx + a)^2 + bc^2 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2), x, algorithm="fricas")

[Out] 2/45*(4*d⁵*cos(b*x + a)⁵ - 9*d⁵*cos(b*x + a)³)*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))/(b*c²*cos(b*x + a)⁴ - 2*b*c²*cos(b*x + a)² + b*c²)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{11}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(11/2)/(c*sec(b*x + a))^(3/2), x)

maple [A] time = 1.10, size = 54, normalized size = 0.49

$$\frac{2 \left(4 \left(\cos^2 (bx + a) \right) - 9 \right) \left(\frac{d}{\sin (bx+a)} \right)^{\frac{11}{2}} \cos (bx + a) \sin (bx + a)}{45 b \left(\frac{c}{\cos (bx+a)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x)

[Out] 2/45/b*(4*cos(b*x+a)^2-9)*(d/sin(b*x+a))^(11/2)*cos(b*x+a)*sin(b*x+a)/(c/cos(b*x+a))^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc (bx + a))^{\frac{11}{2}}}{(c \sec (bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(11/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(11/2)/(c*sec(b*x + a))^(3/2), x)

mupad [B] time = 3.69, size = 125, normalized size = 1.14

$$\frac{8 d^5 \sqrt{\frac{d}{\sin (a+b x)}} (9 \cos (2 a+2 b x)+14 \cos (4 a+4 b x)-9 \cos (6 a+6 b x)+\cos (8 a+8 b x)-15)}{45 b c \sqrt{\frac{c}{\cos (a+b x)}} (28 \cos (4 a+4 b x)-56 \cos (2 a+2 b x)-8 \cos (6 a+6 b x)+\cos (8 a+8 b x)+35)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(11/2)/(c/cos(a + b*x))^(3/2),x)

[Out] (8*d^5*(d/sin(a + b*x))^(1/2)*(9*cos(2*a + 2*b*x) + 14*cos(4*a + 4*b*x) - 9*cos(6*a + 6*b*x) + cos(8*a + 8*b*x) - 15))/(45*b*c*(c/cos(a + b*x))^(1/2)*(28*cos(4*a + 4*b*x) - 56*cos(2*a + 2*b*x) - 8*cos(6*a + 6*b*x) + cos(8*a + 8*b*x) + 35))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(11/2)/(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

$$3.263 \quad \int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=135

$$\frac{2d^4 \sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{21bc^2} + \frac{2d^3 (d \csc(a+bx))^{3/2}}{21bc \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{7/2}}{7bc \sqrt{c \sec(a+bx)}}$$

[Out] $2/21*d^3*(d*csc(b*x+a))^(3/2)/b/c/(c*sec(b*x+a))^(1/2)-2/7*d*(d*csc(b*x+a))^(7/2)/b/c/(c*sec(b*x+a))^(1/2)+2/21*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/c^2$

Rubi [A] time = 0.20, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2623, 2625, 2630, 2573, 2641}

$$\frac{2d^4 \sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{21bc^2} + \frac{2d^3 (d \csc(a+bx))^{3/2}}{21bc \sqrt{c \sec(a+bx)}} - \frac{2d(d \csc(a+bx))^{7/2}}{7bc \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(9/2)/(c*Sec[a + b*x])^(3/2),x]

[Out] $(2*d^3*(d*Csc[a + b*x])^(3/2))/(21*b*c*Sqrt[c*Sec[a + b*x]]) - (2*d*(d*Csc[a + b*x])^(7/2))/(7*b*c*Sqrt[c*Sec[a + b*x]]) - (2*d^4*Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(21*b*c^2)$

Rule 2573

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x, x] /; FreeQ[{a, b, e, f}, x]

Rule 2623

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(f*b*(m - 1)), x] + Dist[(a^2*(n + 1))/(b^2*(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2625

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := -Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Dist[(a^2*(m + n - 2))/(m - 1), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2641

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{9/2}}{(c \sec(a + bx))^{3/2}} dx &= -\frac{2d(d \csc(a + bx))^{7/2}}{7bc\sqrt{c} \sec(a + bx)} - \frac{d^2 \int (d \csc(a + bx))^{5/2} \sqrt{c \sec(a + bx)} dx}{7c^2} \\ &= \frac{2d^3(d \csc(a + bx))^{3/2}}{21bc\sqrt{c} \sec(a + bx)} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc\sqrt{c} \sec(a + bx)} - \frac{(2d^4) \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{21c^2} \\ &= \frac{2d^3(d \csc(a + bx))^{3/2}}{21bc\sqrt{c} \sec(a + bx)} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc\sqrt{c} \sec(a + bx)} - \frac{(2d^4 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)})}{21c^2} \\ &= \frac{2d^3(d \csc(a + bx))^{3/2}}{21bc\sqrt{c} \sec(a + bx)} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc\sqrt{c} \sec(a + bx)} - \frac{(2d^4 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(a + bx)})}{21c^2} \\ &= \frac{2d^3(d \csc(a + bx))^{3/2}}{21bc\sqrt{c} \sec(a + bx)} - \frac{2d(d \csc(a + bx))^{7/2}}{7bc\sqrt{c} \sec(a + bx)} - \frac{2d^4 \sqrt{d \csc(a + bx)} F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c}}{21bc^2} \end{aligned}$$

Mathematica [C] time = 1.46, size = 119, normalized size = 0.88

$$\frac{d^3 \cos(2(a + bx))(d \csc(a + bx))^{3/2} \left((\cos(2(a + bx)) + 5) \csc^4(a + bx) - 2(-\cot^2(a + bx))^{3/4} \sec^2(a + bx) {}_2F_1\left(\frac{1}{2}, 1; \frac{3}{2}; -\cot^2(a + bx)\right) \right)}{21bc \left(\csc^2(a + bx) - 2 \right) \sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(9/2)/(c*Sec[a + b*x])^(3/2), x]

[Out]
$$-1/21*(d^3*\cos[2*(a + b*x)]*(d*Csc[a + b*x])^(3/2)*((5 + \cos[2*(a + b*x)])*Csc[a + b*x]^4 - 2*(-\cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2]*Sec[a + b*x]^2))/(b*c*(-2 + Csc[a + b*x]^2)*\sqrt{c*Sec[a + b*x]})$$

fricas [F] time = 1.16, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} d^4 \csc(bx + a)^4}{c^2 \sec(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2), x, algorithm="fricas")

[Out]
$$\text{integral}(\sqrt{d \csc(bx + a)} * \sqrt{c \sec(bx + a)} * d^4 * \csc(bx + a)^4 / (c^2 * \sec(bx + a)^2), x)$$

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{9}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2), x, algorithm="giac")

[Out]
$$\text{integrate}((d \csc(bx + a))^{9/2} / (c \sec(bx + a))^{3/2}, x)$$

maple [B] time = 1.23, size = 542, normalized size = 4.01

$$\left(2 \sin(bx + a) (\cos^3(bx + a)) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \text{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a)}{\sin(bx + a)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2), x)

[Out]
$$1/21/b*(2*\sin(b*x+a)*\cos(b*x+a)^3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^(1/2)*((-1+\cos(b*x+a))/\sin(b*x+a))$$

$a)^{(1/2)} * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)}, 1/2 * 2^{(1/2)}) + 2 * \sin(b*x+a) * \cos(b*x+a)^2 * ((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)} * ((-1 + \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)} * ((-1 + \cos(b*x+a)) / \sin(b*x+a))^{(1/2)} * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)}, 1/2 * 2^{(1/2)}) - 2 * \sin(b*x+a) * \cos(b*x+a) * ((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)} * ((-1 + \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)} * ((-1 + \cos(b*x+a)) / \sin(b*x+a))^{(1/2)} * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)}, 1/2 * 2^{(1/2)}) - 2 * \sin(b*x+a) * ((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)} * ((-1 + \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)} * ((-1 + \cos(b*x+a)) / \sin(b*x+a))^{(1/2)} * \text{EllipticF}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{(1/2)}, 1/2 * 2^{(1/2)}) - \cos(b*x+a)^3 * 2^{(1/2)} - 2 * \cos(b*x+a) * 2^{(1/2)} * (d / \sin(b*x+a))^{(9/2)} * \sin(b*x+a) / (c / \cos(b*x+a))^{(3/2)} / \cos(b*x+a)^2 * 2^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{9}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{9/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(3/2),x)

[Out] int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(9/2)/(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

$$3.264 \quad \int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=33

$$-\frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{5/2}}$$

[Out] $-2/5*c*d*(d*\csc(b*x+a))^{(5/2)}/b/(c*\sec(b*x+a))^{(5/2)}$

Rubi [A] time = 0.05, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2619}

$$-\frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(7/2)}/(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(-2*c*d*(d*\text{Csc}[a + b*x])^{(5/2)})/(5*b*(c*\text{Sec}[a + b*x])^{(5/2)})$

Rule 2619

$\text{Int}[(\csc[(e_.) + (f_.)*(x_)]*(a_.))^{(m_)}*((b_.)*\sec[(e_.) + (f_.)*(x_)])^{(n_)}], x_Symbol] :> \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(n-1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \ \&\& \ \text{EqQ}[m + n - 2, 0] \ \&\& \ \text{NeQ}[n, 1]$

Rubi steps

$$\int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{3/2}} dx = -\frac{2cd(d \csc(a+bx))^{5/2}}{5b(c \sec(a+bx))^{5/2}}$$

Mathematica [A] time = 0.14, size = 45, normalized size = 1.36

$$-\frac{2d^3 \cot^2(a+bx) \sqrt{d \csc(a+bx)}}{5bc \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(d*\text{Csc}[a + b*x])^{(7/2)}/(c*\text{Sec}[a + b*x])^{(3/2)}, x]$

[Out] $(-2*d^3*\text{Cot}[a + b*x]^2*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(5*b*c*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

fricas [B] time = 0.86, size = 59, normalized size = 1.79

$$\frac{2d^3 \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx+a)^3}{5(bc^2 \cos(bx+a)^2 - bc^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] 2/5*d^3*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^3/(b*c^2*cos(b*x + a)^2 - b*c^2)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{7}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(3/2), x)

maple [A] time = 1.10, size = 42, normalized size = 1.27

$$\frac{2 \left(\frac{d}{\sin(bx+a)} \right)^{\frac{7}{2}} \cos(bx+a) \sin(bx+a)}{5b \left(\frac{c}{\cos(bx+a)} \right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x)

[Out] -2/5/b*(d/sin(b*x+a))^(7/2)*cos(b*x+a)*sin(b*x+a)/(c/cos(b*x+a))^(3/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{7}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(3/2), x)

mupad [B] time = 1.29, size = 70, normalized size = 2.12

$$\frac{2 d^3 (\cos(4 a + 4 b x) - 1) \sqrt{\frac{d}{\sin(a+b x)}}}{5 b c \sqrt{\frac{c}{\cos(a+b x)}} (\cos(4 a + 4 b x) - 4 \cos(2 a + 2 b x) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(3/2),x)

[Out] (2*d^3*(cos(4*a + 4*b*x) - 1)*(d/sin(a + b*x))^(1/2))/(5*b*c*(c/cos(a + b*x))^(1/2)*(cos(4*a + 4*b*x) - 4*cos(2*a + 2*b*x) + 3))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

$$3.265 \quad \int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=98

$$\frac{d^2 \sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3bc^2} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc \sqrt{c \sec(a+bx)}}$$

[Out] $-2/3*d*(d*\csc(b*x+a))^{(3/2)}/b/c/(c*\sec(b*x+a))^{(1/2)}+1/3*d^2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x),2^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*(c*\sec(b*x+a))^{(1/2)}*\sin(2*b*x+2*a)^{(1/2)}/b/c^2$

Rubi [A] time = 0.15, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2623, 2630, 2573, 2641}

$$\frac{d^2 \sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{3bc^2} - \frac{2d(d \csc(a+bx))^{3/2}}{3bc \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a+b*x])^{(5/2)}/(c*\text{Sec}[a+b*x])^{(3/2)},x]$

[Out] $(-2*d*(d*\text{Csc}[a+b*x])^{(3/2)})/(3*b*c*\text{Sqrt}[c*\text{Sec}[a+b*x]]) - (d^2*\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sec}[a+b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(3*b*c^2)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /;$ FreeQ[{a, b, e, f}, x]

Rule 2623

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_)}, x_Symbol] \rightarrow -\text{Simp}[(a*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n+1)})/(f*b*(m-1)), x] + \text{Dist}[(a^2*(n+1))/(b^2*(m-1)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^{(n+2)}, x], x] /;$ FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(a*\text{Csc}[e + f*x])^m*(b*\text{Sec}[e + f*x])^n*(a*\text{Sin}[e + f*x])$

$\int (b \cos[e + f x])^n \int \frac{1}{(a \sin[e + f x])^m (b \cos[e + f x])^n} dx, x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{IntegerQ}[m - 1/2] \ \&\& \ \text{IntegerQ}[n - 1/2]$

Rule 2641

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2*\text{EllipticF}[(1*(c - \text{Pi}/2 + d*x))/2, 2])/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{3/2}} dx &= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc\sqrt{c \sec(a + bx)}} - \frac{d^2 \int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{3c^2} \\ &= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc\sqrt{c \sec(a + bx)}} - \frac{(d^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)})}{3c^2} \\ &= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc\sqrt{c \sec(a + bx)}} - \frac{(d^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)})}{3c^2} \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx \\ &= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc\sqrt{c \sec(a + bx)}} - \frac{d^2 \sqrt{d \csc(a + bx)} F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}}{3bc^2} \end{aligned}$$

Mathematica [C] time = 0.83, size = 105, normalized size = 1.07

$$\frac{d \cos(2(a + bx)) \sec^3(a + bx) (d \csc(a + bx))^{3/2} \left(2 \cot^2(a + bx) - (-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right)\right)}{3b \left(\csc^2(a + bx) - 2\right) (c \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(5/2)/(c*Sec[a + b*x])^(3/2), x]

[Out] $-1/3*(d*\text{Cos}[2*(a + b*x)]*(d*\text{Csc}[a + b*x])^{3/2}*(2*\text{Cot}[a + b*x]^2 - (-\text{Cot}[a + b*x]^2)^{3/4}*\text{Hypergeometric2F1}[1/2, 3/4, 3/2, \text{Csc}[a + b*x]^2])*\text{Sec}[a + b*x]^3)/(b*(-2 + \text{Csc}[a + b*x]^2)*(c*\text{Sec}[a + b*x])^{3/2})$

fricas [F] time = 0.72, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} d^2 \csc(bx + a)^2}{c^2 \sec(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d^2*csc(b*x + a)^2/(c^2*sec(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{5}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(3/2), x)

maple [B] time = 1.19, size = 286, normalized size = 2.92

$$\frac{\left(\sin(bx + a) \cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}\right) \right)}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x)

[Out] -1/3/b*(sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+cos(b*x+a)*2^(1/2))*d/sin(b*x+a))^(5/2)*sin(b*x+a)/(c/cos(b*x+a))^(3/2)/cos(b*x+a)^2*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{5}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(3/2), x)

[Out] int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(3/2), x)

[Out] Timed out

$$3.266 \quad \int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=327

$$\frac{d^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{c \sec(a+bx)}}{\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{d^2 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{c \sec(a+bx)}}{\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{d^2 \sqrt{c \sec(a+bx)}}{2\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

[Out] $-2*d*(d*csc(b*x+a))^{(1/2)}/b/c/(c*sec(b*x+a))^{(1/2)}-1/2*d^2*arctan(-1+2^{(1/2)})*\tan(b*x+a)^{(1/2)}*(c*sec(b*x+a))^{(1/2)}/b/c^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}-1/2*d^2*arctan(1+2^{(1/2)})*\tan(b*x+a)^{(1/2)}*(c*sec(b*x+a))^{(1/2)}/b/c^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}-1/4*d^2*\ln(1-2^{(1/2)})*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(c*sec(b*x+a))^{(1/2)}/b/c^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}+1/4*d^2*\ln(1+2^{(1/2)})*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(c*sec(b*x+a))^{(1/2)}/b/c^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}$

Rubi [A] time = 0.22, antiderivative size = 327, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2623, 2629, 3476, 329, 297, 1162, 617, 204, 1165, 628}

$$\frac{d^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{c \sec(a+bx)}}{\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{d^2 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{c \sec(a+bx)}}{\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} - \frac{d^2 \sqrt{c \sec(a+bx)}}{2\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(3/2), x]

[Out] $(-2*d*Sqrt[d*Csc[a + b*x]])/(b*c*Sqrt[c*Sec[a + b*x]]) + (d^2*ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[c*Sec[a + b*x]])/(Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) - (d^2*ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[c*Sec[a + b*x]])/(Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) - (d^2*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(2*Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (d^2*Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(2*Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :-> -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 297

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2623

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)
)/(f*b*(m - 1)), x] + Dist[(a^2*(n + 1))/(b^2*(m - 1)), Int[(a*Csc[e + f*x]
```

)^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_)])^n), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n]/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^n), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{3/2}} dx &= \frac{2d\sqrt{d} \csc(a + bx)}{bc\sqrt{c} \sec(a + bx)} - \frac{d^2 \int \frac{\sqrt{c \sec(a+bx)}}{\sqrt{d} \csc(a+bx)} dx}{c^2} \\
&= \frac{2d\sqrt{d} \csc(a + bx)}{bc\sqrt{c} \sec(a + bx)} - \frac{(d^2\sqrt{c} \sec(a + bx)) \int \sqrt{\tan(a + bx)} dx}{c^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
&= \frac{2d\sqrt{d} \csc(a + bx)}{bc\sqrt{c} \sec(a + bx)} - \frac{(d^2\sqrt{c} \sec(a + bx)) \text{Subst}\left(\int \frac{\sqrt{x}}{1+x^2} dx, x, \tan(a + bx)\right)}{bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
&= \frac{2d\sqrt{d} \csc(a + bx)}{bc\sqrt{c} \sec(a + bx)} - \frac{(2d^2\sqrt{c} \sec(a + bx)) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
&= \frac{2d\sqrt{d} \csc(a + bx)}{bc\sqrt{c} \sec(a + bx)} + \frac{(d^2\sqrt{c} \sec(a + bx)) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)}\right)}{bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} - \frac{(d^2\sqrt{c} \sec(a + bx)) \text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a + bx)}\right)}{2bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
&= \frac{2d\sqrt{d} \csc(a + bx)}{bc\sqrt{c} \sec(a + bx)} - \frac{d^2 \log\left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right) \sqrt{c} \sec(a + bx)}{2\sqrt{2} bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} + \frac{d^2 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c} \sec(a + bx)}{\sqrt{2} bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} - \frac{d^2 \tan^{-1}\left(1 + \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c} \sec(a + bx)}{\sqrt{2} bc^2\sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}}
\end{aligned}$$

Mathematica [A] time = 0.67, size = 199, normalized size = 0.61

$$\frac{d\sqrt{d} \csc(a + bx) \left(8\sqrt[4]{\cot^2(a + bx)} + \sqrt{2} \log\left(\sqrt{\cot^2(a + bx)} - \sqrt{2} \sqrt[4]{\cot^2(a + bx)} + 1\right) - \sqrt{2} \log\left(\sqrt{\cot^2(a + bx)} + \sqrt{2} \sqrt[4]{\cot^2(a + bx)} + 1\right) \right)}{4bc\sqrt[4]{\cot^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(3/2), x]

[Out] -1/4*(d*Sqrt[d*Csc[a + b*x]]*(2*Sqrt[2]*ArcTan[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] - 2*Sqrt[2]*ArcTan[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] + 8*(Cot[a + b*x]^2)^(1/4) + Sqrt[2]*Log[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4) + Sqrt[Cot[a + b*x]^2]] - Sqrt[2]*Log[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4) + Sqrt[Cot[a + b*x]^2]])))/(b*c*(Cot[a + b*x]^2)^(1/4)*Sqrt[c*Sec[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

$+a))^{1/2} * ((-1 + \cos(b*x+a)) / \sin(b*x+a))^{1/2} * \text{EllipticPi}(((1 - \cos(b*x+a) + \sin(b*x+a)) / \sin(b*x+a))^{1/2}, 1/2 + 1/2*I, 1/2 * 2^{1/2}) + 2 * \cos(b*x+a) * 2^{1/2}) * (d / \sin(b*x+a))^{3/2} * \sin(b*x+a) / (c / \cos(b*x+a))^{3/2} / \cos(b*x+a)^{2 * 2^{1/2}}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{3}{2}}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(3/2), x)

[Out] int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(3/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(3/2), x)

[Out] Timed out

$$3.267 \quad \int \frac{\sqrt{d} \csc(a+bx)}{(c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=92

$$\frac{\sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{2bc^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

[Out] d/b/c/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(1/2)-1/2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*(d*csc(b*x+a))^(1/2)*(c*sec(b*x+a))^(1/2)*sin(2*b*x+2*a)^(1/2)/b/c^2

Rubi [A] time = 0.14, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2628, 2630, 2573, 2641}

$$\frac{\sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \middle| 2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{2bc^2} + \frac{d}{bc \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(3/2), x]

[Out] d/(b*c*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]]) + (Sqrt[d*Csc[a + b*x]]*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[c*Sec[a + b*x]]*Sqrt[Sin[2*a + 2*b*x]])/(2*b*c^2)

Rule 2573

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2628

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2630

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])

$\int \frac{1}{(a \sin[e + f x])^m (b \cos[e + f x])^n} dx$,
 Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x],
 x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2641

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{d \csc(a + bx)}}{(c \sec(a + bx))^{3/2}} dx &= \frac{d}{bc \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{\int \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} dx}{2c^2} \\ &= \frac{d}{bc \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{(\sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)})}{2c^2} \\ &= \frac{d}{bc \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{(\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)})}{2c^2} \\ &= \frac{d}{bc \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} + \frac{\sqrt{d \csc(a + bx)} F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sec(a + bx)}}{2bc^2} \end{aligned}$$

Mathematica [C] time = 0.66, size = 84, normalized size = 0.91

$$\frac{d \sec^3(a + bx) \left(-(-\cot^2(a + bx))^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right) + \cos(2(a + bx)) + 1 \right)}{2b(c \sec(a + bx))^{3/2} \sqrt{d \csc(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(3/2), x]

[Out] (d*(1 + Cos[2*(a + b*x)] - (-Cot[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])*Sec[a + b*x]^3)/(2*b*Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2))

fricas [F] time = 1.21, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)}}{c^2 \sec(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")
 [Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^2*sec(b*x + a)^2), x)
giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")
 [Out] integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(3/2), x)
maple [A] time = 1.22, size = 193, normalized size = 2.10

$$\frac{\left(-\sin(bx+a)\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}\operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{\sqrt{2}}{2}\right)\right)}{2b(-1+\cos(bx+a))\left(\frac{c}{\cos(bx+a)}\right)^{\frac{3}{2}}\cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x)
 [Out] 1/2/b*(-sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*cos(b*x+a)^2-cos(b*x+a)*2^(1/2))*(d/sin(b*x+a))^(1/2)*sin(b*x+a)/(-1+cos(b*x+a))/(c/cos(b*x+a))^(3/2)/cos(b*x+a)^2*2^(1/2)
maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")
 [Out] integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(3/2), x)
mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{d}{\sin(a+bx)}}}{\left(\frac{c}{\cos(a+bx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(3/2), x)`

[Out] `int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(3/2), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(a + bx)} \frac{1}{3}}{(c \sec(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(3/2), x)`

[Out] `Integral(sqrt(d*csc(a + b*x))/(c*sec(a + b*x))**(3/2), x)`

$$3.268 \quad \int \frac{1}{\sqrt{d} \csc(a+bx) (c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=322

$$\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)} + \frac{\sqrt{c \sec(a+bx)}}{8\sqrt{2} bc^2}$$

[Out] 1/2*d/b/c/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(1/2)+1/8*arctan(-1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/8*arctan(1+2^(1/2)*tan(b*x+a)^(1/2))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)+1/16*ln(1-2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)-1/16*ln(1+2^(1/2)*tan(b*x+a)^(1/2)+tan(b*x+a))*(c*sec(b*x+a))^(1/2)/b/c^2*2^(1/2)/(d*csc(b*x+a))^(1/2)/tan(b*x+a)^(1/2)

Rubi [A] time = 0.21, antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2628, 2629, 3476, 329, 297, 1162, 617, 204, 1165, 628}

$$\frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)} + \frac{\tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{c \sec(a+bx)}}{4\sqrt{2} bc^2 \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)} + \frac{\sqrt{c \sec(a+bx)}}{8\sqrt{2} bc^2}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)), x]

[Out] d/(2*b*c*(d*Csc[a + b*x])^(3/2)*Sqrt[c*Sec[a + b*x]]) - (ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[c*Sec[a + b*x]])/(4*Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]]]*Sqrt[c*Sec[a + b*x]])/(4*Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(8*Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) - (Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(8*Sqrt[2]*b*c^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 297

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4)

), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 329

Int[((c_.)*(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 617

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 628

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1162

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1165

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2628

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^(m)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1]

&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{3/2}} dx &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} + \frac{\int \frac{\sqrt{c \sec(a + bx)}}{\sqrt{d} \csc(a + bx)} dx}{4c^2} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} + \frac{\sqrt{c \sec(a + bx)} \int \sqrt{\tan(a + bx)}}{4c^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} + \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{\sqrt{x}}{1+x^2} dx\right)}{4bc^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} + \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{x^2}{1+x^4} dx\right)}{2bc^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} - \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{1-x^2}{1+x^4} dx\right)}{4bc^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} + \frac{\sqrt{c \sec(a + bx)} \operatorname{Subst}\left(\int \frac{1}{1-\sqrt{2}x} dx\right)}{8bc^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} + \frac{\log\left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx)\right)}{8\sqrt{2} bc^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}} \\
 &= \frac{d}{2bc(d \csc(a + bx))^{3/2} \sqrt{c \sec(a + bx)}} - \frac{\tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a + bx)}\right) \sqrt{c \sec(a + bx)}}{4\sqrt{2} bc^2 \sqrt{d} \csc(a + bx) \sqrt{\tan(a + bx)}}
 \end{aligned}$$

Mathematica [A] time = 2.02, size = 223, normalized size = 0.69

$$\sqrt{d \csc(a + bx)} \left(4 \sqrt[4]{\cot^2(a + bx)} - 4 \cos(2(a + bx)) \sqrt[4]{\cot^2(a + bx)} + \sqrt{2} \log \left(\sqrt{\cot^2(a + bx)} - \sqrt{2} \sqrt[4]{\cot^2(a + bx)} \right) \right)$$

16bca

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(3/2)),x]

[Out] (Sqrt[d*Csc[a + b*x]]*(2*Sqrt[2]*ArcTan[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] - 2*Sqrt[2]*ArcTan[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] + 4*(Cot[a + b*x]^2)^(1/4) - 4*Cos[2*(a + b*x)]*(Cot[a + b*x]^2)^(1/4) + Sqrt[2]*Log[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4) + Sqrt[Cot[a + b*x]^2]] - Sqrt[2]*Log[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4) + Sqrt[Cot[a + b*x]^2]]))/(16*b*c*d*(Cot[a + b*x]^2)^(1/4)*Sqrt[c*Sec[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d \csc(bx + a)} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2)), x)

maple [C] time = 0.99, size = 520, normalized size = 1.61

$$\left(i \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} - \frac{i}{2}, \frac{\sqrt{2}}{2} \right) - \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x)

[Out]
$$-1/8/b*(I*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2}))*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}-I*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2}))-((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2}))-((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2}))-2*2^{1/2}*cos(b*x+a)^2+2*cos(b*x+a)*2^{1/2})*sin(b*x+a)/(-1+\cos(b*x+a))/cos(b*x+a)^2/(d/\sin(b*x+a))^{1/2}/(c/\cos(b*x+a))^{3/2}*2^{1/2}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d} \csc(bx+a) (c \sec(bx+a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2} \sqrt{\frac{d}{\sin(a+bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(1/2)),x)

[Out] int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(1/2)), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sec(a + bx))^{\frac{3}{2}} \sqrt{d} \csc(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(3/2),x)
```

```
[Out] Integral(1/((c*sec(a + b*x))**(3/2)*sqrt(d*csc(a + b*x))), x)
```

$$3.269 \quad \int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=135

$$\frac{\sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4}\middle|2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{12bc^2d^2} - \frac{c}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}} + \frac{1}{6bcd\sqrt{c \sec(a+bx)}}$$

[Out] $-1/3*c/b/d/(c*\sec(b*x+a))^{5/2}/(d*\csc(b*x+a))^{1/2}+1/6/b/c/d/(d*\csc(b*x+a))^{1/2}/(c*\sec(b*x+a))^{1/2}-1/12*(\sin(a+1/4*Pi+b*x))^2)^{1/2}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x),2^{1/2})*(d*\csc(b*x+a))^{1/2}*(c*\sec(b*x+a))^{1/2}*\sin(2*b*x+2*a)^{1/2}/b/c^2/d^2$

Rubi [A] time = 0.21, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2627, 2628, 2630, 2573, 2641}

$$\frac{\sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4}\middle|2\right) \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}}{12bc^2d^2} - \frac{c}{3bd(c \sec(a+bx))^{5/2} \sqrt{d \csc(a+bx)}} + \frac{1}{6bcd\sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/((d*\text{Csc}[a+b*x])^{3/2}*(c*\text{Sec}[a+b*x])^{3/2}),x]$

[Out] $-c/(3*b*d*\text{Sqrt}[d*\text{Csc}[a+b*x]]*(c*\text{Sec}[a+b*x])^{5/2})+1/(6*b*c*d*\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{Sqrt}[c*\text{Sec}[a+b*x]])+(\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{EllipticF}[a-Pi/4+b*x,2]*\text{Sqrt}[c*\text{Sec}[a+b*x]]*\text{Sqrt}[\text{Sin}[2*a+2*b*x]])/(12*b*c^2*d^2)$

Rule 2573

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.)+(f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.)+(f_.)*(x_.)])],x_Symbol] :> \text{Dist}[\text{Sqrt}[\text{Sin}[2*e+2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e+f*x]]*\text{Sqrt}[b*\text{Cos}[e+f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e+2*f*x]],x],x] /; \text{FreeQ}\{a,b,e,f\},x]$

Rule 2627

$\text{Int}[(\csc[(e_.)+(f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.)+(f_.)*(x_.)])^{(n_.)},x_Symbol] :> \text{Simp}[(b*(a*\text{Csc}[e+f*x])^{(m+1)}*(b*\text{Sec}[e+f*x])^{(n-1)})/(a*f*(m+n)),x] + \text{Dist}[(m+1)/(a^2*(m+n)), \text{Int}[(a*\text{Csc}[e+f*x])^{(m+2)}*(b*\text{Sec}[e+f*x])^n,x],x] /; \text{FreeQ}\{a,b,e,f,n\},x] \&\& \text{LtQ}[m,-1] \&\& \text{NeQ}[m+n,0] \&\& \text{IntegersQ}[2*m,2*n]$

Rule 2628

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^(m*(b*Sec[e + f*x])^(n + 2)), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2641

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticF[(1*(c - Pi/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} dx &= -\frac{c}{3bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{5/2}} + \frac{\int \frac{\sqrt{d} \csc(a + bx)}{(c \sec(a + bx))^{3/2}} dx}{6d^2} \\ &= -\frac{c}{3bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{5/2}} + \frac{1}{6bcd\sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} \\ &= -\frac{c}{3bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{5/2}} + \frac{1}{6bcd\sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} \\ &= -\frac{c}{3bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{5/2}} + \frac{1}{6bcd\sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} \\ &= -\frac{c}{3bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{5/2}} + \frac{1}{6bcd\sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} \end{aligned}$$

Mathematica [C] time = 0.57, size = 89, normalized size = 0.66

$$\frac{\csc^2(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \csc^2(a + bx)\right)}{\sqrt[4]{-\cot^2(a + bx)}} - 2 \cos(2(a + bx))}{12bcd\sqrt{c \sec(a + bx)} \sqrt{d} \csc(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2)),x]

[Out] (-2*Cos[2*(a + b*x)] + (Csc[a + b*x]^2*Hypergeometric2F1[1/2, 3/4, 3/2, Csc[a + b*x]^2])/(-Cot[a + b*x]^2)^(1/4))/(12*b*c*d*Sqrt[d*Csc[a + b*x]]*Sqrt[c*Sec[a + b*x]])

fricas [F] time = 0.98, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d} \csc(bx + a) \sqrt{c} \sec(bx + a)}{c^2 d^2 \csc(bx + a)^2 \sec(bx + a)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^2*d^2*csc(b*x + a)^2*sec(b*x + a)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2)), x)

maple [A] time = 0.88, size = 220, normalized size = 1.63

$$\frac{\left(2 \left(\cos^4(bx + a)\right) \sqrt{2} + \sin(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \text{EllipticF}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{\pi}{4}\right)\right)}{12b(-1 + \cos(bx + a)) \left(\frac{d}{\sin(bx + a)}\right)^{\frac{3}{2}} \left(\frac{c}{\cos(bx + a)}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x)

[Out] -1/12/b*(2*cos(b*x+a)^4*2^(1/2)+sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1

$/2*2^{(1/2)}-2*\cos(b*x+a)^3*2^{(1/2)}-2^{(1/2)}*\cos(b*x+a)^2+\cos(b*x+a)*2^{(1/2)}$
 $/(-1+\cos(b*x+a))/(d/\sin(b*x+a))^{(3/2)}/(c/\cos(b*x+a))^{(3/2)}/\cos(b*x+a)^2/\sin$
 $(b*x+a)*2^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2} \left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2)),x)

[Out] int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(3/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a)**(3/2)/(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

$$3.270 \quad \int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=371

$$\frac{3 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{c \sec(a+bx)}}{32\sqrt{2} bc^2 d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{c \sec(a+bx)}}{32\sqrt{2} bc^2 d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3\sqrt{c \sec(a+bx)}}{64\sqrt{2} bc^2 d^2}$$

[Out] $-1/4*c/b/d/(d*csc(b*x+a))^{(3/2)}/(c*sec(b*x+a))^{(5/2)}+3/16/b/c/d/(d*csc(b*x+a))^{(3/2)}/(c*sec(b*x+a))^{(1/2)}+3/64*arctan(-1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(c*sec(b*x+a))^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}+3/64*arctan(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(c*sec(b*x+a))^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}+3/128*\ln(1-2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(c*sec(b*x+a))^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}-3/128*\ln(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(c*sec(b*x+a))^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(d*csc(b*x+a))^{(1/2)}/\tan(b*x+a)^{(1/2)}$

Rubi [A] time = 0.30, antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {2627, 2628, 2629, 3476, 329, 297, 1162, 617, 204, 1165, 628}

$$\frac{3 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{c \sec(a+bx)}}{32\sqrt{2} bc^2 d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{c \sec(a+bx)}}{32\sqrt{2} bc^2 d^2 \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{3\sqrt{c \sec(a+bx)}}{64\sqrt{2} bc^2 d^2}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2)),x]

[Out] $-c/(4*b*d*(d*Csc[a + b*x])^{(3/2)}*(c*Sec[a + b*x])^{(5/2)}) + 3/(16*b*c*d*(d*Csc[a + b*x])^{(3/2)}*Sqrt[c*Sec[a + b*x]]) - (3*ArcTan[1 - Sqrt[2]*Sqrt[Tan[a + b*x]])*Sqrt[c*Sec[a + b*x]])/(32*Sqrt[2]*b*c^2*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (3*ArcTan[1 + Sqrt[2]*Sqrt[Tan[a + b*x]])*Sqrt[c*Sec[a + b*x]])/(32*Sqrt[2]*b*c^2*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) + (3*Log[1 - Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(64*Sqrt[2]*b*c^2*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]]) - (3*Log[1 + Sqrt[2]*Sqrt[Tan[a + b*x]] + Tan[a + b*x]]*Sqrt[c*Sec[a + b*x]])/(64*Sqrt[2]*b*c^2*d^2*Sqrt[d*Csc[a + b*x]]*Sqrt[Tan[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 297

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2627

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
```

```

_.), x_Symbol] := Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1)
)/(a*f*(m + n)), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m +
2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] &
& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

```

Rule 2628

```

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1
))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(
b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1]
&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

```

Rule 2629

```

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^
n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ
[n] && EqQ[m + n, 0]

```

Rule 3476

```

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]

```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{3/2}} dx &= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3 \int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^3} dx}{8d^2} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}} \\
&= -\frac{c}{4bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} + \frac{3}{16bcd(d \csc(a + bx))^{3/2} \sqrt{d}}
\end{aligned}$$

Mathematica [A] time = 2.28, size = 246, normalized size = 0.66

$$\sqrt{d \csc(a + bx)} \left(8 \sqrt[4]{\cot^2(a + bx)} - 12 \cos(2(a + bx)) \sqrt[4]{\cot^2(a + bx)} + 4 \cos(4(a + bx)) \sqrt[4]{\cot^2(a + bx)} + 3\sqrt{2} \log \right)$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(3/2)),x]

[Out] (Sqrt[d*Csc[a + b*x]]*(6*Sqrt[2]*ArcTan[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] - 6*Sqrt[2]*ArcTan[1 + Sqrt[2]*(Cot[a + b*x]^2)^(1/4)] + 8*(Cot[a + b*x]^2)^(1/4) - 12*Cos[2*(a + b*x)]*(Cot[a + b*x]^2)^(1/4) + 4*Cos[4*(a + b*x)]*(Cot[a + b*x]^2)^(1/4) + 3*Sqrt[2]*Log[1 - Sqrt[2]*(Cot[a + b*x]^2)^(1/4) +

$\text{Sqrt}[\text{Cot}[a + b*x]^2] - 3*\text{Sqrt}[2]*\text{Log}[1 + \text{Sqrt}[2]*(\text{Cot}[a + b*x]^2)^{(1/4)} + \text{Sqrt}[\text{Cot}[a + b*x]^2]]/(128*b*c*d^3*(\text{Cot}[a + b*x]^2)^{(1/4)}*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2)), x)

maple [C] time = 0.85, size = 548, normalized size = 1.48

$$\frac{\left(8 \left(\cos^4(bx + a)\right) \sqrt{2} + 3i \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \text{EllipticPi}\left(\sqrt{\frac{1 - \cos(bx + a)}{\sin(bx + a)}}\right)\right)}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x)

[Out] $-1/64/b*(8*\cos(b*x+a)^4*2^{(1/2)}+3*I*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}-3*I*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})-8*\cos(b*x+a)^3*2^{(1/2)}-3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})$

/2))-3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))-6*2^(1/2)*cos(b*x+a)^2+6*cos(b*x+a)*2^(1/2))/(-1+cos(b*x+a))/cos(b*x+a)^2/sin(b*x+a)/(d/sin(b*x+a))^(5/2)/(c/cos(b*x+a))^(3/2)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(3/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{3/2} \left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(5/2)),x)

[Out] int(1/((c/cos(a + b*x))^(3/2)*(d/sin(a + b*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(3/2),x)

[Out] Timed out

$$3.271 \quad \int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=33

$$\frac{2cd(d \csc(a+bx))^{7/2}}{7b(c \sec(a+bx))^{7/2}}$$

[Out] $-2/7*c*d*(d*\csc(b*x+a))^{(7/2)}/b/(c*\sec(b*x+a))^{(7/2)}$

Rubi [A] time = 0.05, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2619}

$$\frac{2cd(d \csc(a+bx))^{7/2}}{7b(c \sec(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(9/2)}/(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(-2*c*d*(d*\text{Csc}[a + b*x])^{(7/2)})/(7*b*(c*\text{Sec}[a + b*x])^{(7/2)})$

Rule 2619

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a*b*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)})/(f*(n-1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \ \&\& \ \text{EqQ}[m + n - 2, 0] \ \&\& \ \text{NeQ}[n, 1]$

Rubi steps

$$\int \frac{(d \csc(a+bx))^{9/2}}{(c \sec(a+bx))^{5/2}} dx = -\frac{2cd(d \csc(a+bx))^{7/2}}{7b(c \sec(a+bx))^{7/2}}$$

Mathematica [A] time = 0.16, size = 45, normalized size = 1.36

$$-\frac{2d^4 \cot^3(a+bx) \sqrt{d \csc(a+bx)}}{7bc^2 \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(d*\text{Csc}[a + b*x])^{(9/2)}/(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(-2*d^4*\text{Cot}[a + b*x]^3*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(7*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

fricas [B] time = 4.02, size = 67, normalized size = 2.03

$$\frac{2d^4 \sqrt{\frac{c}{\cos(bx+a)}} \sqrt{\frac{d}{\sin(bx+a)}} \cos(bx+a)^4}{7(b^3 \cos(bx+a)^2 - bc^3) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] 2/7*d^4*sqrt(c/cos(b*x + a))*sqrt(d/sin(b*x + a))*cos(b*x + a)^4/((b*c^3*cos(b*x + a)^2 - b*c^3)*sin(b*x + a))

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{9}{2}}}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(5/2), x)

maple [A] time = 1.11, size = 42, normalized size = 1.27

$$\frac{2 \left(\frac{d}{\sin(bx+a)} \right)^{\frac{9}{2}} \cos(bx+a) \sin(bx+a)}{7b \left(\frac{c}{\cos(bx+a)} \right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x)

[Out] -2/7/b*(d/sin(b*x+a))^(9/2)*cos(b*x+a)*sin(b*x+a)/(c/cos(b*x+a))^(5/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{9}{2}}}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(9/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(9/2)/(c*sec(b*x + a))^(5/2), x)

mupad [B] time = 1.76, size = 93, normalized size = 2.82

$$\frac{2 d^4 \sqrt{\frac{d}{\sin(a+b x)}} (3 \sin (2 a+2 b x)-\sin (6 a+6 b x))}{7 b c^2 \sqrt{\frac{c}{\cos(a+b x)}} (15 \cos (2 a+2 b x)-6 \cos (4 a+4 b x)+\cos (6 a+6 b x)-10)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(9/2)/(c/cos(a + b*x))^(5/2),x)

[Out] (2*d^4*(d/sin(a + b*x))^(1/2)*(3*sin(2*a + 2*b*x) - sin(6*a + 6*b*x)))/(7*b*c^2*(c/cos(a + b*x))^(1/2)*(15*cos(2*a + 2*b*x) - 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) - 10))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(9/2)/(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

$$3.272 \quad \int \frac{(d \csc(a+bx))^{7/2}}{(c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=135

$$\frac{6d^4 E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{5bc^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{6d^3 \sqrt{d \csc(a+bx)}}{5bc(c \sec(a+bx))^{3/2}} - \frac{2d(d \csc(a+bx))^{5/2}}{5bc(c \sec(a+bx))^{3/2}}$$

[Out] $-2/5*d*(d*\csc(b*x+a))^{(5/2)}/b/c/(c*\sec(b*x+a))^{(3/2)}+6/5*d^3*(d*\csc(b*x+a))^{(1/2)}/b/c/(c*\sec(b*x+a))^{(3/2)}-6/5*d^4*(\sin(a+1/4*\pi+b*x)^2)^{(1/2)}/\sin(a+1/4*\pi+b*x)*\text{EllipticE}(\cos(a+1/4*\pi+b*x),2^{(1/2)})/b/c^2/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.20, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2623, 2625, 2630, 2572, 2639}

$$\frac{6d^4 E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{5bc^2 \sqrt{\sin(2a+2bx)} \sqrt{c \sec(a+bx)} \sqrt{d \csc(a+bx)}} + \frac{6d^3 \sqrt{d \csc(a+bx)}}{5bc(c \sec(a+bx))^{3/2}} - \frac{2d(d \csc(a+bx))^{5/2}}{5bc(c \sec(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[a + b*x])^(7/2)/(c*Sec[a + b*x])^(5/2), x]

[Out] $(6*d^3*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(5*b*c*(c*\text{Sec}[a + b*x])^{(3/2)}) - (2*d*(d*\text{Csc}[a + b*x])^{(5/2)})/(5*b*c*(c*\text{Sec}[a + b*x])^{(3/2)}) + (6*d^4*\text{EllipticE}[a - \pi/4 + b*x, 2])/(5*b*c^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\sin[2*a + 2*b*x]])$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]] , x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[2*e + 2*f*x]], Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2623

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(f*b*(m - 1)), x] + Dist[(a^2*(n + 1))/(b^2*(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2625

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := -Simp[(a*b*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n - 1))/(f*(m - 1)), x] + Dist[(a^2*(m + n - 2))/(m - 1), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2639

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{7/2}}{(c \sec(a + bx))^{5/2}} dx &= -\frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} - \frac{(3d^2) \int \frac{(d \csc(a + bx))^{3/2}}{\sqrt{c \sec(a + bx)}} dx}{5c^2} \\ &= \frac{6d^3 \sqrt{d \csc(a + bx)}}{5bc(c \sec(a + bx))^{3/2}} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} + \frac{(6d^4) \int \frac{1}{\sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} dx}{5c^2} \\ &= \frac{6d^3 \sqrt{d \csc(a + bx)}}{5bc(c \sec(a + bx))^{3/2}} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} + \frac{(6d^4) \int \sqrt{c \cos(a + bx)} \sqrt{d \sec(a + bx)}}{5c^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)}} \\ &= \frac{6d^3 \sqrt{d \csc(a + bx)}}{5bc(c \sec(a + bx))^{3/2}} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} + \frac{(6d^4) \int \sqrt{\sin(2a + 2bx)} dx}{5c^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \\ &= \frac{6d^3 \sqrt{d \csc(a + bx)}}{5bc(c \sec(a + bx))^{3/2}} - \frac{2d(d \csc(a + bx))^{5/2}}{5bc(c \sec(a + bx))^{3/2}} + \frac{6d^4 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{5bc^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] time = 1.77, size = 101, normalized size = 0.75

$$\frac{d^5 \sqrt{c \sec(a + bx)} \left(6 \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) + (1 - 3 \cos(2(a + bx))) \cot^2(a + bx) \csc^2(a + bx) \right)}{5bc^3 (d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(7/2)/(c*Sec[a + b*x])^(5/2),x]

[Out] (d^5*((1 - 3*Cos[2*(a + b*x)])*Cot[a + b*x]^2*Csc[a + b*x]^2 + 6*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(5*b*c^3*(d*Csc[a + b*x])^(3/2))

fricas [F] time = 1.03, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d \csc(bx + a)} \sqrt{c \sec(bx + a)} d^3 \csc(bx + a)^3}{c^3 \sec(bx + a)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))*d^3*csc(b*x + a)^3/(c^3*sec(b*x + a)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{7}{2}}}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(5/2), x)

maple [B] time = 1.21, size = 977, normalized size = 7.24

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x)

[Out] 1/5/b*(6*cos(b*x+a)^3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-3*cos(b*x+a)^3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+6*cos(b*x+a)^2*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-3*cos(b*x+a)^2*((1-cos(b*x+a)+sin(b*x+a))

$$\frac{1}{\sin(bx+a)^{1/2}} * ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} * \text{EllipticF}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2*2^{1/2}) - 6*\cos(bx+a)*((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} * \text{EllipticE}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2*2^{1/2}) + 3*\cos(bx+a)*((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} * \text{EllipticF}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2*2^{1/2}) - 6*((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} * \text{EllipticE}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2*2^{1/2}) + 3*((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} * ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} * \text{EllipticF}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2*2^{1/2}) - 3*\cos(bx+a)^3*2^{1/2} - 2^{1/2}*\cos(bx+a)^2 + 3*\cos(bx+a)*2^{1/2} * (d/\sin(bx+a))^{7/2} * \sin(bx+a) / (c/\cos(bx+a))^{5/2} / \cos(bx+a)^3*2^{1/2}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx+a))^{\frac{7}{2}}}{(c \sec(bx+a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(7/2)/(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{7/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(5/2), x)

[Out] int((d/sin(a + b*x))^(7/2)/(c/cos(a + b*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(5/2), x)

[Out] Timed out

$$3.273 \quad \int \frac{(d \csc(a+bx))^{5/2}}{(c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=329

$$\frac{d^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} bc^2 \sqrt{c \sec(a+bx)}} - \frac{d^2 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} bc^2 \sqrt{c \sec(a+bx)}}$$

[Out] $-2/3*d*(d*\csc(b*x+a))^{(3/2)}/b/c/(c*\sec(b*x+a))^{(3/2)}-1/2*d^2*\arctan(-1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-1/2*d^2*\arctan(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+1/4*d^2*\ln(1-2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-1/4*d^2*\ln(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.21, antiderivative size = 329, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2623, 2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{d^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} bc^2 \sqrt{c \sec(a+bx)}} - \frac{d^2 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{\sqrt{2} bc^2 \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(5/2)}/(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(-2*d*(d*\text{Csc}[a + b*x])^{(3/2)})/(3*b*c*(c*\text{Sec}[a + b*x])^{(3/2)}) + (d^2*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) - (d^2*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (d^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(2*\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) - (d^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(2*\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 204

$\text{Int}[(a_0 + (b_0*x)^2)^{-1}, x_Symbol] \rightarrow -\text{Simp}[\text{ArcTan}[(\text{Rt}[-b, 2]*x)/\text{Rt}[-a, 2]]/(\text{Rt}[-a, 2]*\text{Rt}[-b, 2]), x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 211

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2623

```
Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)
)/(f*b*(m - 1)), x] + Dist[(a^2*(n + 1))/(b^2*(m - 1)), Int[(a*Csc[e + f*x]
```

)^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
\int \frac{(d \csc(a + bx))^{5/2}}{(c \sec(a + bx))^{5/2}} dx &= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} - \frac{d^2 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{c^2} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} - \frac{(d^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{c^2 \sqrt{c \sec(a + bx)}} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} - \frac{(d^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1}{\sqrt{x}(1+x^2)} dx, x, \tan(a + bx) \right)}{bc^2 \sqrt{c \sec(a + bx)}} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} - \frac{(2d^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, \sqrt{\tan(a + bx)} \right)}{bc^2 \sqrt{c \sec(a + bx)}} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} - \frac{(d^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \sqrt{\tan(a + bx)} \right)}{bc^2 \sqrt{c \sec(a + bx)}} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} - \frac{(d^2 \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}) \text{Subst} \left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \sqrt{\tan(a + bx)} \right)}{2bc^2 \sqrt{c \sec(a + bx)}} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} + \frac{d^2 \sqrt{d \csc(a + bx)} \log \left(1 - \sqrt{2} \sqrt{\tan(a + bx)} + \tan(a + bx) \right) \sqrt{\tan(a + bx)}}{2\sqrt{2} bc^2 \sqrt{c \sec(a + bx)}} \\
&= -\frac{2d(d \csc(a + bx))^{3/2}}{3bc(c \sec(a + bx))^{3/2}} + \frac{d^2 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a + bx)} \right) \sqrt{d \csc(a + bx)} \sqrt{\tan(a + bx)}}{\sqrt{2} bc^2 \sqrt{c \sec(a + bx)}}
\end{aligned}$$

Mathematica [C] time = 0.20, size = 55, normalized size = 0.17

$$\frac{2d(d \csc(a + bx))^{3/2} \left({}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a + bx) \right) - 1 \right)}{3bc(c \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(5/2)/(c*Sec[a + b*x])^(5/2),x]

[Out] (2*d*(d*Csc[a + b*x])^(3/2)*(-1 + Hypergeometric2F1[3/4, 1, 7/4, -Cot[a + b*x]^2]))/(3*b*c*(c*Sec[a + b*x])^(3/2))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{5}{2}}}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

[Out] `integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(5/2), x)`

maple [C] time = 1.23, size = 1239, normalized size = 3.77

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x)`

[Out] `-1/6/b*(3*I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*cos(b*x+a)*sin(b*x+a)-3*I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*cos(b*x+a)*sin(b*x+a)+3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*cos(b*x+a)*sin(b*x+a)+3*I*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))+3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*cos(b*x+a)*sin(b*x+a)-3*I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)-6*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/`

$2*2^{(1/2)}+3*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)*\sin(b*x+a)+3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})*\sin(b*x+a)-6*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)*EllipticF(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2*2^{(1/2)})+2*2^{(1/2)*\cos(b*x+a)^2*(d/\sin(b*x+a))^{(5/2)*\sin(b*x+a)/\cos(b*x+a)^3/(c/\cos(b*x+a))^{(5/2)*2^{(1/2)}}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc(bx + a))^{\frac{5}{2}}}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(5/2)/(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{5/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(5/2), x)

[Out] int((d/sin(a + b*x))^(5/2)/(c/cos(a + b*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(5/2), x)

[Out] Timed out

$$3.274 \quad \int \frac{(d \csc(a+bx))^{3/2}}{(c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=94

$$\frac{3d^2 E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{bc^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc (c \sec(a + bx))^{3/2}}$$

[Out] $-2*d*(d*\csc(b*x+a))^{(1/2)}/b/c/(c*\sec(b*x+a))^{(3/2)}+3*d^2*(\sin(a+1/4*\pi+b*x)^2)^{(1/2)}/\sin(a+1/4*\pi+b*x)*\text{EllipticE}(\cos(a+1/4*\pi+b*x),2^{(1/2)})/b/c^2/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.15, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2623, 2630, 2572, 2639}

$$\frac{3d^2 E\left(a + bx - \frac{\pi}{4} \middle| 2\right)}{bc^2 \sqrt{\sin(2a + 2bx)} \sqrt{c \sec(a + bx)} \sqrt{d \csc(a + bx)}} - \frac{2d \sqrt{d \csc(a + bx)}}{bc (c \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[a + b*x])^{(3/2)}/(c*\text{Sec}[a + b*x])^{(5/2)}, x]$

[Out] $(-2*d*\text{Sqrt}[d*\text{Csc}[a + b*x]])/(b*c*(c*\text{Sec}[a + b*x])^{(3/2)}) - (3*d^2*\text{EllipticE}[a - \pi/4 + b*x, 2])/(b*c^2*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[c*\text{Sec}[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2572

$\text{Int}[\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]]$, x_Symbol] $\rightarrow \text{Dist}[(\text{Sqrt}[a*\sin[e + f*x]]*\text{Sqrt}[b*\cos[e + f*x]])/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]$, $\text{Int}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]$, x], x] /; $\text{FreeQ}\{a, b, e, f\}, x]$

Rule 2623

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}$, x_Symbol] $\rightarrow -\text{Simp}[(a*(a*\csc[e + f*x])^{(m-1)}*(b*\sec[e + f*x])^{(n+1)})/(f*b*(m-1))$, x] + $\text{Dist}[(a^{2*(n+1)})/(b^{2*(m-1)})$, $\text{Int}[(a*\csc[e + f*x])^{(m-2)}*(b*\sec[e + f*x])^{(n+2)}$, x], x] /; $\text{FreeQ}\{a, b, e, f\}, x]$ && $\text{GtQ}[m, 1]$ && $\text{LtQ}[n, -1]$ && $\text{IntegersQ}[2*m, 2*n]$

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}$, x_Symbol] $\rightarrow \text{Dist}[(a*\csc[e + f*x])^m*(b*\sec[e + f*x])^n*(a*\sin[e + f*x])$

)^m*(b*cos[e + f*x])^n, Int[1/((a*sin[e + f*x])^m*(b*cos[e + f*x])^n), x],
 x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
]

Rule 2639

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P
 i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(d \csc(a + bx))^{3/2}}{(c \sec(a + bx))^{5/2}} dx &= -\frac{2d\sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} - \frac{(3d^2) \int \frac{1}{\sqrt{d \csc(a+bx)} \sqrt{c \sec(a+bx)}} dx}{c^2} \\ &= -\frac{2d\sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} - \frac{(3d^2) \int \sqrt{c \cos(a + bx)} \sqrt{d \sin(a + bx)} dx}{c^2 \sqrt{c \cos(a + bx)} \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{d \sin(a + bx)}} \\ &= -\frac{2d\sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} - \frac{(3d^2) \int \sqrt{\sin(2a + 2bx)} dx}{c^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \\ &= -\frac{2d\sqrt{d \csc(a + bx)}}{bc(c \sec(a + bx))^{3/2}} - \frac{3d^2 E\left(a - \frac{\pi}{4} + bx \mid 2\right)}{bc^2 \sqrt{d \csc(a + bx)} \sqrt{c \sec(a + bx)} \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] time = 0.70, size = 80, normalized size = 0.85

$$\frac{d^3 \sqrt{c \sec(a + bx)} \left(3 \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) + 2 \cot^2(a + bx) \right)}{bc^3 (d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[a + b*x])^(3/2)/(c*Sec[a + b*x])^(5/2),x]

[Out] -((d^3*(2*Cot[a + b*x]^2 + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2,
 , 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(b*c^3*(d*Csc[a + b*x])^(
 (3/2))))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc (bx + a))^{\frac{3}{2}}}{(c \sec (bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(5/2), x)

maple [B] time = 1.17, size = 507, normalized size = 5.39

$$\left(6 \cos (bx + a) \sqrt{\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \sqrt{\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}} \sqrt{\frac{-1 + \cos (bx + a)}{\sin (bx + a)}} \operatorname{EllipticE} \left(\sqrt{\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)}}, \frac{\sqrt{2}}{2} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x)

[Out] $\frac{1}{2} b (6 \cos (bx + a) \left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \operatorname{EllipticE} \left(\left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}}, \frac{1}{2} \sqrt{2} \right) - 3 \cos (bx + a) \left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \operatorname{EllipticF} \left(\left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}}, \frac{1}{2} \sqrt{2} \right) + 6 \left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \operatorname{EllipticE} \left(\left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}}, \frac{1}{2} \sqrt{2} \right) - 3 \left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \left(\frac{-1 + \cos (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}} \operatorname{EllipticF} \left(\left(\frac{1 - \cos (bx + a) + \sin (bx + a)}{\sin (bx + a)} \right)^{\frac{1}{2}}, \frac{1}{2} \sqrt{2} \right) + 2^{\frac{1}{2}} \cos (bx + a)^2 - 3 \cos (bx + a) \sqrt{2} \right) \left(\frac{d}{\sin (bx + a)} \right)^{\frac{3}{2}} \sin (bx + a) / \cos (bx + a)^3 / (c / \cos (bx + a))^{\frac{5}{2}} \sqrt{2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \csc (bx + a))^{\frac{3}{2}}}{(c \sec (bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((d*csc(b*x + a))^(3/2)/(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(a+bx)}\right)^{3/2}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(5/2),x)

[Out] int((d/sin(a + b*x))^(3/2)/(c/cos(a + b*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

$$3.275 \quad \int \frac{\sqrt{d} \csc(a+bx)}{(c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=322

$$\frac{3 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)}{4\sqrt{2} bc^2 \sqrt{c} \sec(a+bx)} + \frac{3 \tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)}{4\sqrt{2} bc^2 \sqrt{c} \sec(a+bx)}$$

[Out] $1/2*d/b/c/(c*\sec(b*x+a))^{(3/2)}/(d*\csc(b*x+a))^{(1/2)}+3/8*\arctan(-1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+3/8*\arctan(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-3/16*\ln(1-2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+3/16*\ln(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.20, antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2628, 2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{3 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)}{4\sqrt{2} bc^2 \sqrt{c} \sec(a+bx)} + \frac{3 \tan^{-1}\left(\sqrt{2} \sqrt{\tan(a+bx)} + 1\right) \sqrt{\tan(a+bx)} \sqrt{d} \csc(a+bx)}{4\sqrt{2} bc^2 \sqrt{c} \sec(a+bx)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(5/2), x]

[Out] $d/(2*b*c*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(3/2)}) - (3*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(4*\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (3*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(4*\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) - (3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b*c^2*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),

$x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& (\text{GtQ}[a/b, 0] \mid\mid (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 329

$\text{Int}[(c_*)(x_)^m*((a_) + (b_*)(x_)^n)^p], x_Symbol] :> \text{With}[\{k = \text{Denominator}[m]\}, \text{Dist}[k/c, \text{Subst}[\text{Int}[x^{k*(m+1)-1}*(a + (b*x^{k*n}))/c^n]^p, x], x, (c*x)^{1/k}], x]] /; \text{FreeQ}[\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 617

$\text{Int}[(a_) + (b_*)(x_) + (c_*)(x_)^2]^{-1}, x_Symbol] :> \text{With}[\{q = 1 - 4*\text{Simplify}[(a*c)/b^2]\}, \text{Dist}[-2/b, \text{Subst}[\text{Int}[1/(q - x^2)], x], x, 1 + (2*c*x)/b], x] /; \text{RationalQ}[q] \&\& (\text{EqQ}[q^2, 1] \mid\mid \text{!RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 628

$\text{Int}[(d_) + (e_*)(x_)] / [(a_) + (b_*)(x_) + (c_*)(x_)^2], x_Symbol] :> \text{Simp}[(d*\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]])/b, x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 1162

$\text{Int}[(d_) + (e_*)(x_)^2] / [(a_) + (c_*)(x_)^4], x_Symbol] :> \text{With}[\{q = \text{Rt}[(2*d)/e, 2]\}, \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

Rule 1165

$\text{Int}[(d_) + (e_*)(x_)^2] / [(a_) + (c_*)(x_)^4], x_Symbol] :> \text{With}[\{q = \text{Rt}[-(2*d)/e, 2]\}, \text{Dist}[e/(2*c*q), \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

Rule 2628

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.)^m*((b_.)*\text{sec}[(e_.) + (f_.)*(x_)])^n), x_Symbol] :> -\text{Simp}[(a*(a*\text{Csc}[e + f*x])^{m-1}*(b*\text{Sec}[e + f*x])^{n+1})/(b*f*(m+n)), x] + \text{Dist}[(n+1)/(b^2*(m+n)), \text{Int}[(a*\text{Csc}[e + f*x])^m*(b*\text{Sec}[e + f*x])^{n+2}], x], x] /; \text{FreeQ}[\{a, b, e, f, m\}, x] \&\& \text{LtQ}[n, -1]$

&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2629

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]

Rule 3476

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{d \csc(a+bx)}}{(c \sec(a+bx))^{5/2}} dx &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} + \frac{3 \int \frac{\sqrt{d \csc(a+bx)}}{\sqrt{c \sec(a+bx)}} dx}{4c^2} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} + \frac{(3\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \int \frac{1}{\sqrt{\tan(a+bx)}} dx}{4c^2 \sqrt{c \sec(a+bx)}} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} + \frac{(3\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \text{Subst}\left(\int \frac{1}{\sqrt{x}} dx, \sqrt{\tan(a+bx)}\right)}{4bc^2 \sqrt{c \sec(a+bx)}} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} + \frac{(3\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \text{Subst}\left(\int \frac{1}{1+x} dx, \sqrt{\tan(a+bx)}\right)}{2bc^2 \sqrt{c \sec(a+bx)}} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} + \frac{(3\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \text{Subst}\left(\int \frac{1-x}{1+x} dx, \sqrt{\tan(a+bx)}\right)}{4bc^2 \sqrt{c \sec(a+bx)}} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} + \frac{(3\sqrt{d \csc(a+bx)} \sqrt{\tan(a+bx)}) \text{Subst}\left(\int \frac{1}{1-\sqrt{x}} dx, \sqrt{\tan(a+bx)}\right)}{8bc^2 \sqrt{c \sec(a+bx)}} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} - \frac{3\sqrt{d \csc(a+bx)} \log\left(1 - \sqrt{2} \sqrt{\tan(a+bx)} + \sqrt{2} \sqrt{\tan(a+bx)}\right)}{8\sqrt{2} bc^2 \sqrt{c \sec(a+bx)}} \\
 &= \frac{d}{2bc\sqrt{d \csc(a+bx)} (c \sec(a+bx))^{3/2}} - \frac{3 \tan^{-1}\left(1 - \sqrt{2} \sqrt{\tan(a+bx)}\right) \sqrt{d \csc(a+bx)}}{4\sqrt{2} bc^2 \sqrt{c \sec(a+bx)}}
 \end{aligned}$$

Mathematica [C] time = 0.23, size = 70, normalized size = 0.22

$$\frac{\cot(a + bx)\sqrt{d \csc(a + bx)} \left({}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a + bx)\right) + \cos(2(a + bx)) - 1 \right)}{4bc^2\sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[a + b*x]]/(c*Sec[a + b*x])^(5/2), x]

[Out] -1/4*(Cot[a + b*x]*Sqrt[d*Csc[a + b*x]]*(-1 + Cos[2*(a + b*x)] + 2*Hypergeometric2F1[3/4, 1, 7/4, -Cot[a + b*x]^2]))/(b*c^2*Sqrt[c*Sec[a + b*x]])

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2), x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(bx + a)}}{(c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2), x, algorithm="giac")

[Out] integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(5/2), x)

maple [C] time = 1.20, size = 658, normalized size = 2.04

$$\left(3i \operatorname{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} - \frac{i}{2} \frac{\sqrt{2}}{2} \right) \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \right) \operatorname{si}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2), x)

[Out] -1/8/b*(3*I*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*E

$$\text{EllipticPi}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}, 1/2-1/2I, 1/2\sqrt{2}\right)^{1/2} - 3I\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2} \text{EllipticPi}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}, 1/2+1/2I, 1/2\sqrt{2}\right) \sin(bx+a) + 3\text{EllipticPi}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}, 1/2-1/2I, 1/2\sqrt{2}\right) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2} \sin(bx+a) - 6\sin(bx+a) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2} \text{EllipticF}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}, 1/2\sqrt{2}\right) + 3\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2} \left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2} \text{EllipticPi}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}, 1/2+1/2I, 1/2\sqrt{2}\right) \sin(bx+a) - 2\cos(bx+a)^3\sqrt{2} + 2\sqrt{2}\cos(bx+a)^2 \left(\frac{d}{\sin(bx+a)}\right)^{1/2} \sin(bx+a) / \left(\frac{-1+\cos(bx+a)}{\cos(bx+a)}\right)^3 / \left(\frac{c}{\cos(bx+a)}\right)^{5/2} \sqrt{2}$$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \csc(bx+a)}}{(c \sec(bx+a))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(b*x + a))/(c*sec(b*x + a))^(5/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{d}{\sin(a+bx)}}}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(5/2),x)

[Out] int((d/sin(a + b*x))^(1/2)/(c/cos(a + b*x))^(5/2), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

$$3.276 \quad \int \frac{1}{\sqrt{d} \csc(a+bx) (c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=95

$$\frac{E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{2bc^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} + \frac{d}{3bc(c\sec(a+bx))^{3/2}(d\csc(a+bx))^{3/2}}$$

[Out] $1/3*d/b/c/(d*\csc(b*x+a))^{(3/2)}/(c*\sec(b*x+a))^{(3/2)}-1/2*(\sin(a+1/4*\text{Pi}+b*x))^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticE}(\cos(a+1/4*\text{Pi}+b*x),2^{(1/2)})/b/c^2/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.14, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2628, 2630, 2572, 2639}

$$\frac{E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{2bc^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} + \frac{d}{3bc(c\sec(a+bx))^{3/2}(d\csc(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(\text{Sqrt}[d*\text{Csc}[a+b*x]]*(c*\text{Sec}[a+b*x])^{(5/2)}),x]$

[Out] $d/(3*b*c*(d*\text{Csc}[a+b*x])^{(3/2)}*(c*\text{Sec}[a+b*x])^{(3/2)}) + \text{EllipticE}[a - \text{Pi}/4 + b*x, 2]/(2*b*c^2*\text{Sqrt}[d*\text{Csc}[a+b*x]]*\text{Sqrt}[c*\text{Sec}[a+b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2572

$\text{Int}[\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]] , x_Symbol] \rightarrow \text{Dist}[(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]])/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], \text{Int}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2628

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow -\text{Simp}[(a*(a*\text{Csc}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n+1)})/(b*f*(m+n)), x] + \text{Dist}[(n+1)/(b^2*(m+n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f, m\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{NeQ}[m+n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2630

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a*\text{Csc}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(n)}], x]$

$)^m (b \cos[e + f x])^n$, $\text{Int}[1/((a \sin[e + f x])^m (b \cos[e + f x])^n), x]$,
 $x] /;$ $\text{FreeQ}\{a, b, e, f, m, n\}, x$ && $\text{IntegerQ}[m - 1/2]$ && $\text{IntegerQ}[n - 1/2]$
 $]$

Rule 2639

$\text{Int}[\text{Sqrt}[\sin[(c \cdot) + (d \cdot)(x \cdot)]]], x_Symbol] :> \text{Simp}[(2 \cdot \text{EllipticE}[(1 \cdot (c - P$
 $i/2 + d \cdot x))/2, 2])/d, x] /;$ $\text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{5/2}} dx &= \frac{d}{3bc(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} + \frac{\int \frac{1}{\sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} dx}{2c^2} \\ &= \frac{d}{3bc(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} + \frac{\int \sqrt{c \cos(a + bx)}}{2c^2 \sqrt{c \cos(a + bx)} \sqrt{d} \csc(a + bx)} \\ &= \frac{d}{3bc(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} + \frac{\int \sqrt{\sin(2a + 2bx)}}{2c^2 \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} \\ &= \frac{d}{3bc(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}} + \frac{E\left(a - \frac{\pi}{4} + bx\right)}{2bc^2 \sqrt{d} \csc(a + bx) \sqrt{c \sec(a + bx)}} \end{aligned}$$

Mathematica [C] time = 0.36, size = 79, normalized size = 0.83

$$\frac{d \sqrt{c \sec(a + bx)} \left(3 \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx)\right) + \cos(2(a + bx)) + 1 \right)}{6bc^3 (d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[d*Csc[a + b*x]]*(c*Sec[a + b*x])^(5/2)),x]

[Out] (d*(1 + Cos[2*(a + b*x)] + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(6*b*c^3*(d*Csc[a + b*x])^(3/2))

fricas [F] time = 2.40, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{d} \csc(bx + a) \sqrt{c \sec(bx + a)}}{c^3 d \csc(bx + a) \sec(bx + a)^3}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^3*d*csc(b*x + a)*sec(b*x + a)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d \csc(bx + a)} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2)), x)

maple [B] time = 0.98, size = 522, normalized size = 5.49

$$\left(2 \left(\cos^4(bx + a) \right) \sqrt{2} - 3 \cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \right) \text{EllipticF} \left(\sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x)

[Out] -1/12/b*(2*cos(b*x+a)^4*2^(1/2)-3*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+6*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+6*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2^(1/2)*cos(b*x+a)^2-3*cos(b*x+a)*2^(1/2))/(d/sin(b*x+a))^(1/2)/(c/cos(b*x+a))^(5/2)/cos(b*x+a)^3/sin(b*x+a)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d \csc(bx + a)} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*csc(b*x+a))^(1/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(1/(sqrt(d*csc(b*x + a))*(c*sec(b*x + a))^(5/2)), x)
```

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \sqrt{\frac{d}{\sin(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2)),x)
```

```
[Out] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(1/2)), x)
```

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*csc(b*x+a))**(1/2)/(c*sec(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```

$$3.277 \quad \int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=371

$$\frac{3 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{32 \sqrt{2} bc^2 d^2 \sqrt{c \sec(a+bx)}} + \frac{3 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d}}{32 \sqrt{2} bc^2 d^2 \sqrt{c \sec(a+bx)}}$$

[Out] $-1/4*c/b/d/(c*\sec(b*x+a))^{(7/2)}/(d*\csc(b*x+a))^{(1/2)}+1/16/b/c/d/(c*\sec(b*x+a))^{(3/2)}/(d*\csc(b*x+a))^{(1/2)}+3/64*\arctan(-1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+3/64*\arctan(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-3/128*\ln(1-2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+3/128*\ln(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^2*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.29, antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {2627, 2628, 2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{3 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{32 \sqrt{2} bc^2 d^2 \sqrt{c \sec(a+bx)}} + \frac{3 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d}}{32 \sqrt{2} bc^2 d^2 \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2)),x]

[Out] $-c/(4*b*d*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(7/2)}) + 1/(16*b*c*d*\text{Sqrt}[d*\text{Csc}[a + b*x]]*(c*\text{Sec}[a + b*x])^{(3/2)}) - (3*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(32*\text{Sqrt}[2]*b*c^2*d^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (3*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(32*\text{Sqrt}[2]*b*c^2*d^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) - (3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(64*\text{Sqrt}[2]*b*c^2*d^2*\text{Sqrt}[c*\text{Sec}[a + b*x]]) + (3*\text{Sqrt}[d*\text{Csc}[a + b*x]]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(64*\text{Sqrt}[2]*b*c^2*d^2*\text{Sqrt}[c*\text{Sec}[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

```
Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2627

```
Int[(csc[(e_) + (f_)*(x_)])*(a_)^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n
```

```

_.), x_Symbol] := Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1)
)/(a*f*(m + n)), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m +
2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] &
& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

```

Rule 2628

```

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1
))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(
b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1]
&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

```

Rule 2629

```

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n
_), x_Symbol] := Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^
n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ
[n] && EqQ[m + n, 0]

```

Rule 3476

```

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[b/d, Subst[Int[
x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !
IntegerQ[n]

```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} dx &= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{\int \frac{\sqrt{d} \csc(a+bx)}{(c \sec(a+bx))^{5/2}} dx}{8d^2} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} \\
&= -\frac{c}{4bd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}} + \frac{1}{16bcd\sqrt{d} \csc(a + bx) (c \sec(a + bx))^{7/2}}
\end{aligned}$$

Mathematica [C] time = 0.37, size = 81, normalized size = 0.22

$$-\frac{\csc^3(a + bx) \left(2 {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a + bx) \right) + \cos(2(a + bx)) - \cos(4(a + bx)) \right)}{32bc(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(5/2)),x]

[Out] -1/32*(Csc[a + b*x]^3*(Cos[2*(a + b*x)] - Cos[4*(a + b*x)] + 2*Hypergeometric2F1[3/4, 1, 7/4, -Cot[a + b*x]^2]))/(b*c*(d*Csc[a + b*x])^(3/2)*(c*Sec[a + b*x])^(3/2))

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2)), x)

maple [C] time = 0.86, size = 686, normalized size = 1.85

$$\frac{\left(3i \operatorname{EllipticPi} \left(\sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2} - \frac{i}{2}, \frac{\sqrt{2}}{2} \right) \sqrt{\frac{1 - \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a) + \sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1 + \cos(bx+a)}{\sin(bx+a)}} \right) \operatorname{si}}{\dots}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x)

[Out] -1/64/b*(3*I*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-3*I*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)+8*cos(b*x+a)^5*2^(1/2)-8*cos(b*x+a)^4*2^(1/2)-6*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+3*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)+3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos

$(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})*\sin(b*x+a)-2*\cos(b*x+a)^3*2^{(1/2)}+2*2^{(1/2)*\cos(b*x+a)^2)/(-1+\cos(b*x+a))}/(d/\sin(b*x+a))^{(3/2)/(c/\cos(b*x+a))^{(5/2)/\cos(b*x+a)^3/\sin(b*x+a)*2^{(1/2)}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{3}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(3/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(1/((d*csc(b*x + a))^(3/2)*(c*sec(b*x + a))^(5/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \left(\frac{d}{\sin(a+bx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(3/2)),x)

[Out] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(3/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))**(3/2)/(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

$$3.278 \quad \int \frac{1}{(d \csc(a+bx))^{5/2} (c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=135

$$\frac{3E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{20bc^2d^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} - \frac{c}{5bd(c\sec(a+bx))^{7/2}(d\csc(a+bx))^{3/2}} + \frac{1}{10bcd(c\sec(a+bx))^{5/2}(d\csc(a+bx))^{3/2}}$$

[Out] $-1/5*c/b/d/(d*\csc(b*x+a))^{(3/2)}/(c*\sec(b*x+a))^{(7/2)}+1/10/b/c/d/(d*\csc(b*x+a))^{(3/2)}/(c*\sec(b*x+a))^{(3/2)}-3/20*(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticE}(\cos(a+1/4*\text{Pi}+b*x),2^{(1/2)})/b/c^2/d^2/(d*\csc(b*x+a))^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A] time = 0.21, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2627, 2628, 2630, 2572, 2639}

$$\frac{3E\left(a+bx-\frac{\pi}{4}\middle|2\right)}{20bc^2d^2\sqrt{\sin(2a+2bx)}\sqrt{c\sec(a+bx)}\sqrt{d\csc(a+bx)}} - \frac{c}{5bd(c\sec(a+bx))^{7/2}(d\csc(a+bx))^{3/2}} + \frac{1}{10bcd(c\sec(a+bx))^{5/2}(d\csc(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2)),x]

[Out] $-c/(5*b*d*(d*Csc[a + b*x])^{(3/2)}*(c*Sec[a + b*x])^{(7/2)}) + 1/(10*b*c*d*(d*Csc[a + b*x])^{(3/2)}*(c*Sec[a + b*x])^{(3/2)}) + (3*EllipticE[a - \text{Pi}/4 + b*x, 2])/((20*b*c^2*d^2*\text{Sqrt}[d*Csc[a + b*x]]*\text{Sqrt}[c*Sec[a + b*x]]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]]))$

Rule 2572

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]] , x_Symbol] :> Dist[(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[2*e + 2*f*x]], Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2627

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))/(a*f*(m + n)), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2628

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2630

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Dist[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2639

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2*EllipticE[(1*(c - P i/2 + d*x))/2, 2])/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{5/2}} dx &= -\frac{c}{5bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{7/2}} + \frac{3 \int \frac{1}{\sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}}}{10d^2} \\ &= -\frac{c}{5bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{7/2}} + \frac{1}{10bcd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} \\ &= -\frac{c}{5bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{7/2}} + \frac{1}{10bcd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} \\ &= -\frac{c}{5bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{7/2}} + \frac{1}{10bcd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} \\ &= -\frac{c}{5bd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{7/2}} + \frac{1}{10bcd(d \csc(a + bx))^{3/2} (c \sec(a + bx))^{5/2}} \end{aligned}$$

Mathematica [C] time = 0.65, size = 90, normalized size = 0.67

$$\frac{\sqrt{c \sec(a + bx)} \left(3 \sqrt[4]{-\cot^2(a + bx)} {}_2F_1 \left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \csc^2(a + bx) \right) - 2 \cos^2(a + bx) \cos(2(a + bx)) \right)}{20bc^3d(d \csc(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(5/2)*(c*Sec[a + b*x])^(5/2)),x]

[Out] ((-2*Cos[a + b*x]^2*Cos[2*(a + b*x)] + 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[-1/2, 1/4, 1/2, Csc[a + b*x]^2])*Sqrt[c*Sec[a + b*x]])/(20*b*c^3*d*(d*Csc[a + b*x])^(3/2))

fricas [F] time = 1.01, size = 0, normalized size = 0.00

$$\text{integral} \left(\frac{\sqrt{d} \csc(bx + a) \sqrt{c} \sec(bx + a)}{c^3 d^3 \csc(bx + a)^3 \sec(bx + a)^3}, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(d*csc(b*x + a))*sqrt(c*sec(b*x + a))/(c^3*d^3*csc(b*x + a)^3*sec(b*x + a)^3), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2)), x)

maple [B] time = 0.87, size = 536, normalized size = 3.97

$$\left(4 \left(\cos^6(bx + a) \right) \sqrt{2} - 6 \left(\cos^4(bx + a) \right) \sqrt{2} - 6 \cos(bx + a) \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x)

[Out] 1/40/b*(4*cos(b*x+a)^6*2^(1/2)-6*cos(b*x+a)^4*2^(1/2)-6*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+3*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b

$x+a)/\sin(b*x+a))^{1/2}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})-6*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticE}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})+3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2})-2^{1/2}*\cos(b*x+a)^2+3*\cos(b*x+a)*2^{1/2}))/\cos(b*x+a)^3/\sin(b*x+a)^3/(d/\sin(b*x+a))^{5/2}/(c/\cos(b*x+a))^{5/2}*2^{1/2}$

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{5}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(5/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(1/((d*csc(b*x + a))^(5/2)*(c*sec(b*x + a))^(5/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \left(\frac{d}{\sin(a+bx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2)),x)

[Out] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(5/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))**(5/2)/(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

$$3.279 \quad \int \frac{1}{(d \csc(a+bx))^{7/2} (c \sec(a+bx))^{5/2}} dx$$

Optimal. Leaf size=406

$$\frac{5 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{128 \sqrt{2} bc^2 d^4 \sqrt{c \sec(a+bx)}} + \frac{5 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d}}{128 \sqrt{2} bc^2 d^4 \sqrt{c \sec(a+bx)}}$$

[Out] $-1/6*c/b/d/(d*\csc(b*x+a))^{(5/2)}/(c*\sec(b*x+a))^{(7/2)}-5/48*c/b/d^3/(c*\sec(b*x+a))^{(7/2)}/(d*\csc(b*x+a))^{(1/2)}+5/192/b/c/d^3/(c*\sec(b*x+a))^{(3/2)}/(d*\csc(b*x+a))^{(1/2)}+5/256*\arctan(-1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^4*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+5/256*\arctan(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)})*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^4*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}-5/512*\ln(1-2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^4*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}+5/512*\ln(1+2^{(1/2)}*\tan(b*x+a)^{(1/2)}+\tan(b*x+a))*(d*\csc(b*x+a))^{(1/2)}*\tan(b*x+a)^{(1/2)}/b/c^2/d^4*2^{(1/2)}/(c*\sec(b*x+a))^{(1/2)}$

Rubi [A] time = 0.39, antiderivative size = 406, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 11, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.440$, Rules used = {2627, 2628, 2629, 3476, 329, 211, 1165, 628, 1162, 617, 204}

$$\frac{5 \tan^{-1} \left(1 - \sqrt{2} \sqrt{\tan(a+bx)} \right) \sqrt{\tan(a+bx)} \sqrt{d \csc(a+bx)}}{128 \sqrt{2} bc^2 d^4 \sqrt{c \sec(a+bx)}} + \frac{5 \tan^{-1} \left(\sqrt{2} \sqrt{\tan(a+bx)} + 1 \right) \sqrt{\tan(a+bx)} \sqrt{d}}{128 \sqrt{2} bc^2 d^4 \sqrt{c \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2)),x]

[Out] $-c/(6*b*d*(d*Csc[a + b*x])^{(5/2)}*(c*Sec[a + b*x])^{(7/2)}) - (5*c)/(48*b*d^3*\text{Sqrt}[d*Csc[a + b*x]]*(c*Sec[a + b*x])^{(7/2)}) + 5/(192*b*c*d^3*\text{Sqrt}[d*Csc[a + b*x]]*(c*Sec[a + b*x])^{(3/2)}) - (5*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*Csc[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(128*\text{Sqrt}[2]*b*c^2*d^4*\text{Sqrt}[c*Sec[a + b*x]]) + (5*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]]]*\text{Sqrt}[d*Csc[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(128*\text{Sqrt}[2]*b*c^2*d^4*\text{Sqrt}[c*Sec[a + b*x]]) - (5*\text{Sqrt}[d*Csc[a + b*x]]*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(256*\text{Sqrt}[2]*b*c^2*d^4*\text{Sqrt}[c*Sec[a + b*x]]) + (5*\text{Sqrt}[d*Csc[a + b*x]]*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{Tan}[a + b*x]] + \text{Tan}[a + b*x]]*\text{Sqrt}[\text{Tan}[a + b*x]])/(256*\text{Sqrt}[2]*b*c^2*d^4*\text{Sqrt}[c*Sec[a + b*x]])$

Rule 204

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :-> -Simp[ArcTan[(Rt[-b, 2]*x)/Rt[-a, 2]]/(Rt[-a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[

a, 0] || LtQ[b, 0])

Rule 211

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 329

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + (b*x^(k*n)))/c^
n]^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 617

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[(a*c)/b^2]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + (2*c*x)/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 628

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[(d*Log[RemoveContent[a + b*x + c*x^2, x]])/b, x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1162

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(2*d)/e, 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1165

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
(-2*d)/e, 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2627

```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(b*(a*Csc[e + f*x])^(m + 1)*(b*Sec[e + f*x])^(n - 1))/(a*f*(m + n)), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] & NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2628

```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> -Simp[(a*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))/(b*f*(m + n)), x] + Dist[(n + 1)/(b^2*(m + n)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2629

```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Dist[((a*Csc[e + f*x])^m*(b*Sec[e + f*x])^n)/Tan[e + f*x]^n, Int[Tan[e + f*x]^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[n] && EqQ[m + n, 0]
```

Rule 3476

```
Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] :> Dist[b/d, Subst[Int[x^n/(b^2 + x^2), x], x, b*Tan[c + d*x]], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{1}{(d \csc(a + bx))^{7/2} (c \sec(a + bx))^{5/2}} dx &= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} + \frac{5 \int \frac{1}{(d \csc(a+bx))^{3/2} (c \sec(a+bx))^{5/2}} dx}{12d^2} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}} \\
&= -\frac{c}{6bd(d \csc(a + bx))^{5/2} (c \sec(a + bx))^{7/2}} - \frac{5c}{48bd^3 \sqrt{d \csc(a + bx)} (c \sec(a + bx))^{5/2}}
\end{aligned}$$

Mathematica [C] time = 0.40, size = 95, normalized size = 0.23

$$\frac{\cot(a + bx) \sqrt{d \csc(a + bx)} \left(10 {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; -\cot^2(a + bx) \right) + 3 \cos(2(a + bx)) - 9 \cos(4(a + bx)) + 2 \cos(6(a + bx)) \right)}{384bc^2d^4 \sqrt{c \sec(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Csc[a + b*x])^(7/2)*(c*Sec[a + b*x])^(5/2)),x]

[Out] $-1/384*(\cot[a + b*x]*\sqrt{d*\csc[a + b*x]}*(4 + 3*\cos[2*(a + b*x)] - 9*\cos[4*(a + b*x)] + 2*\cos[6*(a + b*x)] + 10*\text{Hypergeometric2F1}[3/4, 1, 7/4, -\cot[a + b*x]^2]))/(b*c^2*d^4*\sqrt{c*\sec[a + b*x]})$

fricas [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="fricas")`

[Out] Timed out

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="giac")`

[Out] `integrate(1/((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2)), x)`

maple [C] time = 0.89, size = 712, normalized size = 1.75

$$\frac{\left(64\sqrt{2} \left(\cos^7(bx + a)\right) - 64 \left(\cos^6(bx + a)\right) \sqrt{2} + 15i \sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}}\right)}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x)`

[Out] $1/768/b*(64*2^{(1/2)}*\cos(b*x+a)^7-64*\cos(b*x+a)^6*2^{(1/2)}+15*I*\sin(b*x+a)*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}-15*I*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})-104*\cos(b*x+a)^5*2^{(1/2)}+104*\cos(b*x+a)^4*2^{(1/2)}-15*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}$

2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(b*x+a)-15*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)+30*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+10*cos(b*x+a)^3*2^(1/2)-10*2^(1/2)*cos(b*x+a)^2/(-1+cos(b*x+a))/cos(b*x+a)^3/sin(b*x+a)^3/(d/sin(b*x+a))^(7/2)/(c/cos(b*x+a))^(5/2)*2^(1/2)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(bx + a))^{\frac{7}{2}} (c \sec(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))^(7/2)/(c*sec(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(1/((d*csc(b*x + a))^(7/2)*(c*sec(b*x + a))^(5/2)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{c}{\cos(a+bx)}\right)^{5/2} \left(\frac{d}{\sin(a+bx)}\right)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(7/2)),x)

[Out] int(1/((c/cos(a + b*x))^(5/2)*(d/sin(a + b*x))^(7/2)), x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(b*x+a))**(7/2)/(c*sec(b*x+a))**(5/2),x)

[Out] Timed out

3.280 $\int \csc^n(e + fx) \sec^m(e + fx) dx$

Optimal. Leaf size=81

$$\frac{\cos^2(e + fx)^{\frac{m+1}{2}} \sec^{m+1}(e + fx) \csc^{n-1}(e + fx) {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] (cos(f*x+e)^2)^(1/2+1/2*m)*csc(f*x+e)^(-1+n)*hypergeom([1/2-1/2*n, 1/2+1/2*m], [3/2-1/2*n], sin(f*x+e)^2)*sec(f*x+e)^(1+m)/f/(1-n)

Rubi [A] time = 0.07, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2631, 2577}

$$\frac{\cos^2(e + fx)^{\frac{m+1}{2}} \sec^{m+1}(e + fx) \csc^{n-1}(e + fx) {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^n*Sec[e + f*x]^m,x]

[Out] ((Cos[e + f*x]^2)^((1 + m)/2)*Csc[e + f*x]^(-1 + n)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x]^(1 + m))/(f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2)]/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int \csc^n(e + fx) \sec^m(e + fx) dx = \left(\cos^{1+m}(e + fx) \csc^{-1+n}(e + fx) \sec^{1+m}(e + fx) \sin^{-1+n}(e + fx) \right) \int \cos^{-m}(e + fx) dx$$

$$= \frac{\cos^2(e + fx)^{\frac{1+m}{2}} \csc^{-1+n}(e + fx) {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sec^{1+m}(e + fx)}{f(1-n)}$$

Mathematica [C] time = 2.28, size = 278, normalized size = 3.43

$$(n-3) \sec^m(e + fx) \csc^{n-1}(e + fx)$$

$$f(n-1) \left((n-3) F_1\left(\frac{1}{2} - \frac{n}{2}; m, -m - n + 1; \frac{3}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 2 \tan^2\left(\frac{1}{2}(e + fx)\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[Csc[e + f*x]^n*Sec[e + f*x]^m,x]

[Out] -(((-3 + n)*AppellF1[1/2 - n/2, m, 1 - m - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Csc[e + f*x]^(-1 + n)*Sec[e + f*x]^m)/(f*(-1 + n))*((-3 + n)*AppellF1[1/2 - n/2, m, 1 - m - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*((-1 + m + n)*AppellF1[3/2 - n/2, m, 2 - m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + m*AppellF1[3/2 - n/2, 1 + m, 1 - m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))

fricas [F] time = 5.82, size = 0, normalized size = 0.00

$$\text{integral}\left(\csc(fx + e)^n \sec(fx + e)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^n*sec(f*x+e)^m,x, algorithm="fricas")

[Out] integral(csc(f*x + e)^n*sec(f*x + e)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \csc(fx + e)^n \sec(fx + e)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^n*sec(f*x+e)^m,x, algorithm="giac")

[Out] integrate(csc(f*x + e)^n*sec(f*x + e)^m, x)

maple [F] time = 2.01, size = 0, normalized size = 0.00

$$\int (\csc^n(fx + e)) (\sec^m(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^n*sec(f*x+e)^m,x)

[Out] int(csc(f*x+e)^n*sec(f*x+e)^m,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \csc(fx + e)^n \sec(fx + e)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^n*sec(f*x+e)^m,x, algorithm="maxima")

[Out] integrate(csc(f*x + e)^n*sec(f*x + e)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{1}{\cos(e + fx)} \right)^m \left(\frac{1}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1/cos(e + f*x))^m*(1/sin(e + f*x))^n,x)

[Out] int((1/cos(e + f*x))^m*(1/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \csc^n(e + fx) \sec^m(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**n*sec(f*x+e)**m,x)

[Out] Integral(csc(e + f*x)**n*sec(e + f*x)**m, x)

3.281 $\int \csc^n(e + fx)(a \sec(e + fx))^m dx$

Optimal. Leaf size=86

$$\frac{\cos^2(e + fx)^{\frac{m+1}{2}} \csc^{n-1}(e + fx)(a \sec(e + fx))^{m+1} {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{af(1-n)}$$

[Out] $(\cos(f*x+e)^2)^{(1/2+1/2*m)} * \csc(f*x+e)^{(-1+n)} * \text{hypergeom}([1/2-1/2*n, 1/2+1/2*m], [3/2-1/2*n], \sin(f*x+e)^2) * (a*\sec(f*x+e))^{(1+m)} / a/f/(1-n)$

Rubi [A] time = 0.09, antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{\cos^2(e + fx)^{\frac{m+1}{2}} \csc^{n-1}(e + fx)(a \sec(e + fx))^{m+1} {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{af(1-n)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^n * (a*\text{Sec}[e + f*x])^m, x]$

[Out] $((\text{Cos}[e + f*x]^2)^{((1 + m)/2)} * \text{Csc}[e + f*x]^{(-1 + n)} * \text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 - n)/2, \text{Sin}[e + f*x]^2] * (a*\text{Sec}[e + f*x])^{(1 + m)}) / (a*f*(1 - n))$

Rule 2577

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)} * ((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(b^{(2*\text{IntPart}[(n - 1)/2] + 1)} * (b*\text{Cos}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])} * (a*\text{Sin}[e + f*x])^{(m + 1)} * \text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Sin}[e + f*x]^2]) / (a*f*(m + 1) * (\text{Cos}[e + f*x]^2)^{\text{FracPart}[(n - 1)/2])}, x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x]$

Rule 2631

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)} * ((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(a^{2*(a*\text{Csc}[e + f*x])^{(m - 1)} * (b*\text{Sec}[e + f*x])^{(n + 1)} * (a*\text{Sin}[e + f*x])^{(m - 1)} * (b*\text{Cos}[e + f*x])^{(n + 1)}) / b^2, \text{Int}[1/((a*\text{Sin}[e + f*x])^m * (b*\text{Cos}[e + f*x])^n), x], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{!SimplerQ}[-m, -n]$

Rubi steps

$$\int \csc^n(e + fx)(a \sec(e + fx))^m dx = \frac{\left((a \cos(e + fx))^{1+m} \csc^{-1+n}(e + fx)(a \sec(e + fx))^{1+m} \sin^{-1+n}(e + fx)\right) \int (a \sec(e + fx))^m dx}{a^2}$$

$$= \frac{\cos^2(e + fx)^{\frac{1+m}{2}} \csc^{-1+n}(e + fx) {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) (a \sec(e + fx))^m}{af(1-n)}$$

Mathematica [C] time = 0.59, size = 280, normalized size = 3.26

$$(n-3) \csc^{n-1}(e + fx)(a \sec(e + fx))^m$$

$$\frac{f(n-1) \left((n-3) F_1\left(\frac{1}{2} - \frac{n}{2}; m, -m - n + 1; \frac{3}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 2 \tan^2\left(\frac{1}{2}(e + fx)\right) \right) (a \sec(e + fx))^m}{f(n-1)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Csc[e + f*x]^n*(a*Sec[e + f*x])^m,x]

[Out] -(((-3 + n)*AppellF1[1/2 - n/2, m, 1 - m - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Csc[e + f*x]^(-1 + n)*(a*Sec[e + f*x])^m)/(f*(-1 + n))*((-3 + n)*AppellF1[1/2 - n/2, m, 1 - m - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*((-1 + m + n)*AppellF1[3/2 - n/2, m, 2 - m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + m*AppellF1[3/2 - n/2, 1 + m, 1 - m - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))

fricas [F] time = 1.70, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(a \sec(fx + e)\right)^m \csc(fx + e)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^n*(a*sec(f*x+e))^m,x, algorithm="fricas")

[Out] integral((a*sec(f*x + e))^m*csc(f*x + e)^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \left(a \sec(fx + e)\right)^m \csc(fx + e)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^n*(a*sec(f*x+e))^m,x, algorithm="giac")

[Out] integrate((a*sec(f*x + e))^m*csc(f*x + e)^n, x)

maple [F] time = 1.84, size = 0, normalized size = 0.00

$$\int (\csc^n(fx + e)) (a \sec(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^n*(a*sec(f*x+e))^m,x)

[Out] int(csc(f*x+e)^n*(a*sec(f*x+e))^m,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec(fx + e))^m \csc(fx + e)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^n*(a*sec(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((a*sec(f*x + e))^m*csc(f*x + e)^n, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{a}{\cos(e + fx)} \right)^m \left(\frac{1}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(e + f*x))^m*(1/sin(e + f*x))^n,x)

[Out] int((a/cos(e + f*x))^m*(1/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec(e + fx))^m \csc^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**n*(a*sec(f*x+e))**m,x)

[Out] Integral((a*sec(e + f*x))**m*csc(e + f*x)**n, x)

3.282 $\int (b \csc(e + fx))^n \sec^m(e + fx) dx$

Optimal. Leaf size=84

$$\frac{b \cos^2(e + fx)^{\frac{m+1}{2}} \sec^{m+1}(e + fx) (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] b*(cos(f*x+e)^2)^(1/2+1/2*m)*(b*csc(f*x+e))^(1-n)*hypergeom([1/2-1/2*n, 1/2+1/2*m], [3/2-1/2*n], sin(f*x+e)^2)*sec(f*x+e)^(1+m)/f/(1-n)

Rubi [A] time = 0.09, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{b \cos^2(e + fx)^{\frac{m+1}{2}} \sec^{m+1}(e + fx) (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^m,x]

[Out] (b*(Cos[e + f*x]^2)^((1 + m)/2)*(b*Csc[e + f*x])^(1 - n)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2]*Sec[e + f*x]^(1 + m))/(f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1))*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx = \left(b^2 \cos^{1+m}(e + fx) (b \csc(e + fx))^{-1+n} \sec^{1+m}(e + fx) (b \sin(e + fx))^{-1+n} \right) \\ = \frac{b \cos^2(e + fx)^{\frac{1+m}{2}} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sec^{1+m}(e + fx)}{f(1-n)}$$

Mathematica [C] time = 0.32, size = 281, normalized size = 3.35

$$\frac{b(n-3) \sec^m(e + fx) (b \csc(e + fx))^n}{f(n-1) \left((n-3) F_1\left(\frac{1-n}{2}; m, -m-n+1; \frac{3-n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 2 \tan^2\left(\frac{1}{2}(e + fx)\right) \left((m+n) \right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^m,x]

[Out] -((b*(-3 + n)*AppellF1[(1 - n)/2, m, 1 - m - n, (3 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(b*Csc[e + f*x])^(-1 + n)*Sec[e + f*x]^m)/(f*(-1 + n)*((-3 + n)*AppellF1[(1 - n)/2, m, 1 - m - n, (3 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*((-1 + m + n)*AppellF1[(3 - n)/2, m, 2 - m - n, (5 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + m*AppellF1[(3 - n)/2, 1 + m, 1 - m - n, (5 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))

fricas [F] time = 3.48, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \sec(fx + e)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^m,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e)^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^m,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^m, x)

maple [F] time = 1.88, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (\sec^m(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n*sec(f*x+e)^m,x)

[Out] int((b*csc(f*x+e))^n*sec(f*x+e)^m,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec^m(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^m,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\sin(e + fx)} \right)^n \left(\frac{1}{\cos(e + fx)} \right)^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/sin(e + f*x))^n*(1/cos(e + f*x))^m,x)

[Out] int((b/sin(e + f*x))^n*(1/cos(e + f*x))^m, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \sec^m(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))**n*sec(f*x+e)**m,x)

[Out] Integral((b*csc(e + f*x))**n*sec(e + f*x)**m, x)

3.283 $\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx$

Optimal. Leaf size=89

$$\frac{b \cos^2(e + fx)^{\frac{m+1}{2}} (a \sec(e + fx))^{m+1} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{af(1-n)}$$

[Out] b*(cos(f*x+e)^2)^(1/2+1/2*m)*(b*csc(f*x+e))^(1-n)*hypergeom([1/2-1/2*n, 1/2+1/2*m], [3/2-1/2*n], sin(f*x+e)^2)*(a*sec(f*x+e))^(1+m)/a/f/(1-n)

Rubi [A] time = 0.09, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2631, 2577}

$$\frac{b \cos^2(e + fx)^{\frac{m+1}{2}} (a \sec(e + fx))^{m+1} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{m+1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{af(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*(a*Sec[e + f*x])^m,x]

[Out] (b*(Cos[e + f*x]^2)^(1+m/2)*(b*Csc[e + f*x])^(1-n)*Hypergeometric2F1[(1+m)/2, (1-n)/2, (3-n)/2, Sin[e + f*x]^2]*(a*Sec[e + f*x])^(1+m))/(a*f*(1-n))

Rule 2577

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[(b^(2*IntPart[(n-1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n-1)/2])*(a*Sin[e + f*x])^(m+1)*Hypergeometric2F1[(1+m)/2, (1-n)/2, (3+m)/2, Sin[e + f*x]^2])/(a*f*(m+1)*(Cos[e + f*x]^2)^FracPart[(n-1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_) + (f_)*(x_)]*(a_.))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n+1))*(a*Sin[e + f*x])^(m-1)*(b*Cos[e + f*x])^(n+1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int (b \csc(e + fx))^n (a \sec(e + fx))^m dx = \frac{(b^2 (a \cos(e + fx))^{1+m} (b \csc(e + fx))^{-1+n} (a \sec(e + fx))^{1+m} (b \sin(e + fx))^n)}{a^2}$$

$$= \frac{b \cos^2(e + fx)^{\frac{1+m}{2}} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) (a \sec(e + fx))^m}{af(1-n)}$$

Mathematica [C] time = 0.19, size = 283, normalized size = 3.18

$$\frac{b(n-3)(a \sec(e + fx))^m (b \csc(e + fx))^n}{f(n-1) \left((n-3) F_1\left(\frac{1-n}{2}; m, -m-n+1; \frac{3-n}{2}; \tan^2\left(\frac{1}{2}(e + fx)\right), -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 2 \tan^2\left(\frac{1}{2}(e + fx)\right) \left((m+n) \right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Csc[e + f*x])^n*(a*Sec[e + f*x])^m,x]

[Out] -((b*(-3 + n)*AppellF1[(1 - n)/2, m, 1 - m - n, (3 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(b*Csc[e + f*x])^(-1 + n)*(a*Sec[e + f*x])^m)/(f*(-1 + n)*((-3 + n)*AppellF1[(1 - n)/2, m, 1 - m - n, (3 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - 2*((-1 + m + n)*AppellF1[(3 - n)/2, m, 2 - m - n, (5 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + m*AppellF1[(3 - n)/2, 1 + m, 1 - m - n, (5 - n)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))

fricas [F] time = 1.39, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \left(a \sec(fx + e)\right)^m, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*(a*sec(f*x + e))^m, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*(a*sec(f*x + e))^m, x)

maple [F] time = 1.87, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x)

[Out] int((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (a \sec(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(a*sec(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n*(a*sec(f*x + e))^m, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{a}{\cos(e + fx)} \right)^m \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a/cos(e + f*x))^m*(b/sin(e + f*x))^n,x)

[Out] int((a/cos(e + f*x))^m*(b/sin(e + f*x))^n, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sec(e + fx))^m (b \csc(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))**n*(a*sec(f*x+e))**m,x)

[Out] Integral((a*sec(e + f*x))**m*(b*csc(e + f*x))**n, x)

3.284 $\int (b \csc(e + fx))^n \sec^5(e + fx) dx$

Optimal. Leaf size=48

$$\frac{(b \csc(e + fx))^{n+5} {}_2F_1\left(3, \frac{n+5}{2}; \frac{n+7}{2}; \csc^2(e + fx)\right)}{b^5 f(n+5)}$$

[Out] (b*csc(f*x+e))^(5+n)*hypergeom([3, 5/2+1/2*n], [7/2+1/2*n], csc(f*x+e)^2)/b^5/f/(5+n)

Rubi [A] time = 0.05, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2621, 364}

$$\frac{(b \csc(e + fx))^{n+5} {}_2F_1\left(3, \frac{n+5}{2}; \frac{n+7}{2}; \csc^2(e + fx)\right)}{b^5 f(n+5)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^5,x]

[Out] ((b*Csc[e + f*x])^(5 + n)*Hypergeometric2F1[3, (5 + n)/2, (7 + n)/2, Csc[e + f*x]^2])/(b^5*f*(5 + n))

Rule 364

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a^p*(c*x)^(m + 1)*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, -((b*x^n)/a)])/ (c*(m + 1)), x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\int (b \csc(e + fx))^n \sec^5(e + fx) dx = - \frac{\text{Subst} \left(\int \frac{x^{4+n}}{\left(-1 + \frac{x^2}{b^2}\right)^3} dx, x, b \csc(e + fx) \right)}{b^5 f}$$

$$= \frac{(b \csc(e + fx))^{5+n} {}_2F_1 \left(3, \frac{5+n}{2}; \frac{7+n}{2}; \csc^2(e + fx) \right)}{b^5 f(5+n)}$$

Mathematica [A] time = 0.04, size = 51, normalized size = 1.06

$$\frac{b(b \csc(e + fx))^{n-1} {}_2F_1 \left(3, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx) \right)}{f(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^5,x]

[Out] -((b*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[3, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(-1 + n)))

fricas [F] time = 1.18, size = 0, normalized size = 0.00

$$\text{integral} \left((b \csc(fx + e))^n \sec(fx + e)^5, x \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^5,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e)^5, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^5,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^5, x)

maple [F] time = 1.24, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (\sec^5(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n*sec(f*x+e)^5,x)

[Out] int((b*csc(f*x+e))^n*sec(f*x+e)^5,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec^5(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^5,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^5, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/sin(e + f*x))^n/cos(e + f*x)^5,x)

[Out] int((b/sin(e + f*x))^n/cos(e + f*x)^5, x)

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))**n*sec(f*x+e)**5,x)

[Out] Timed out

3.285 $\int (b \csc(e + fx))^n \sec^3(e + fx) dx$

Optimal. Leaf size=49

$$-\frac{(b \csc(e + fx))^{n+3} {}_2F_1\left(2, \frac{n+3}{2}; \frac{n+5}{2}; \csc^2(e + fx)\right)}{b^3 f(n+3)}$$

[Out] $-(b*\csc(f*x+e))^{(3+n)}*\text{hypergeom}([2, 3/2+1/2*n], [5/2+1/2*n], \csc(f*x+e)^2)/b^3/f/(3+n)$

Rubi [A] time = 0.05, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2621, 364}

$$-\frac{(b \csc(e + fx))^{n+3} {}_2F_1\left(2, \frac{n+3}{2}; \frac{n+5}{2}; \csc^2(e + fx)\right)}{b^3 f(n+3)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Csc}[e + f*x])^n*\text{Sec}[e + f*x]^3, x]$

[Out] $-\left(\left((b*\text{Csc}[e + f*x])^{(3 + n)}*\text{Hypergeometric2F1}[2, (3 + n)/2, (5 + n)/2, \text{Csc}[e + f*x]^2]\right)/(b^3*f*(3 + n))\right)$

Rule 364

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_) + (b_*)*(x_)^{(n_)})^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(a^p*(c*x)^{(m+1)}*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, -((b*x^n)/a)])/(c*(m+1)), x] /;$ $\text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 2621

$\text{Int}[(\csc[(e_*) + (f_*)*(x_)]*(a_*)^{(m_*)}\sec[(e_*) + (f_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow -\text{Dist}[(f*a^n)^{-1}, \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1+x^2/a^2)^{((n+1)/2)}, x], x, a*\text{Csc}[e + f*x]], x] /;$ $\text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n+1)/2] \ \&\& \ !(\text{IntegerQ}[(m+1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx = -\frac{\text{Subst}\left(\int \frac{x^{2+n}}{\left(-1+\frac{x^2}{b^2}\right)^2} dx, x, b \csc(e + fx)\right)}{b^3 f}$$

$$= -\frac{(b \csc(e + fx))^{3+n} {}_2F_1\left(2, \frac{3+n}{2}; \frac{5+n}{2}; \csc^2(e + fx)\right)}{b^3 f(3+n)}$$

Mathematica [A] time = 0.04, size = 51, normalized size = 1.04

$$\frac{b(b \csc(e + fx))^{n-1} {}_2F_1\left(2, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^3,x]

[Out] -((b*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(-1 + n)))

fricas [F] time = 1.63, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \sec(fx + e)^3, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^3,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e)^3, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^3,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^3, x)

maple [F] time = 1.20, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (\sec^3(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n*sec(f*x+e)^3,x)

[Out] int((b*csc(f*x+e))^n*sec(f*x+e)^3,x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec^3(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^3,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^3, x)

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/sin(e + f*x))^n/cos(e + f*x)^3,x)

[Out] int((b/sin(e + f*x))^n/cos(e + f*x)^3, x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \sec^3(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))**n*sec(f*x+e)**3,x)

[Out] Integral((b*csc(e + f*x))**n*sec(e + f*x)**3, x)

3.286 $\int (b \csc(e + fx))^n \sec(e + fx) dx$

Optimal. Leaf size=48

$$\frac{(b \csc(e + fx))^{n+1} {}_2F_1\left(1, \frac{n+1}{2}; \frac{n+3}{2}; \csc^2(e + fx)\right)}{bf(n+1)}$$

[Out] (b*csc(f*x+e))^(1+n)*hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], csc(f*x+e)^2)/b/f/(1+n)

Rubi [A] time = 0.04, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2621, 364}

$$\frac{(b \csc(e + fx))^{n+1} {}_2F_1\left(1, \frac{n+1}{2}; \frac{n+3}{2}; \csc^2(e + fx)\right)}{bf(n+1)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*Sec[e + f*x],x]

[Out] ((b*Csc[e + f*x])^(1 + n)*Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Csc[e + f*x]^2])/(b*f*(1 + n))

Rule 364

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[(a^p*(c*x)^(m + 1)*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, -((b*x^n)/a)])/ (c*(m + 1)), x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\int (b \csc(e + fx))^n \sec(e + fx) dx = -\frac{\text{Subst}\left(\int \frac{x^n}{-1+\frac{x^2}{b^2}} dx, x, b \csc(e + fx)\right)}{bf}$$

$$= \frac{(b \csc(e + fx))^{1+n} {}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \csc^2(e + fx)\right)}{bf(1+n)}$$

Mathematica [A] time = 0.03, size = 51, normalized size = 1.06

$$\frac{b(b \csc(e + fx))^{n-1} {}_2F_1\left(1, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x], x]

[Out] -((b*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[1, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/(f*(-1 + n)))

fricas [F] time = 1.66, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \sec(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e), x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e), x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e), x)

maple [F] time = 1.16, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n*sec(f*x+e),x)`

[Out] `int((b*csc(f*x+e))^n*sec(f*x+e),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*sec(f*x+e),x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*sec(f*x + e), x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/sin(e + f*x))^n/cos(e + f*x),x)`

[Out] `int((b/sin(e + f*x))^n/cos(e + f*x), x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \sec(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))**n*sec(f*x+e),x)`

[Out] `Integral((b*csc(e + f*x))**n*sec(e + f*x), x)`

$$3.287 \quad \int \cos(e + fx)(b \csc(e + fx))^n dx$$

Optimal. Leaf size=24

$$\frac{b(b \csc(e + fx))^{n-1}}{f(1-n)}$$

[Out] b*(b*csc(f*x+e))⁽⁻¹⁺ⁿ⁾/f/(1-n)

Rubi [A] time = 0.03, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2621, 30}

$$\frac{b(b \csc(e + fx))^{n-1}}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]*(b*Csc[e + f*x])ⁿ, x]

[Out] (b*(b*Csc[e + f*x])^(-1 + n))/(f*(1 - n))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2621

Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*sec[(e_) + (f_)*(x_)]^(n_), x_Symbol] := -Dist[(f*aⁿ)⁽⁻¹⁾, Subst[Int[x^(m + n - 1)/(-1 + x²/a²)^{(n + 1)/2}, x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(e + fx)(b \csc(e + fx))^n dx &= -\frac{b \text{Subst}\left(\int x^{-2+n} dx, x, b \csc(e + fx)\right)}{f} \\ &= \frac{b(b \csc(e + fx))^{-1+n}}{f(1-n)} \end{aligned}$$

Mathematica [A] time = 0.02, size = 23, normalized size = 0.96

$$\frac{b(b \csc(e + fx))^{n-1}}{f(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]*(b*Csc[e + f*x])^n,x]

[Out] -((b*(b*Csc[e + f*x])^(-1 + n))/(f*(-1 + n)))

fricas [A] time = 0.86, size = 29, normalized size = 1.21

$$\frac{\left(\frac{b}{\sin(fx+e)}\right)^n \sin(fx+e)}{fn-f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] -(b/sin(f*x + e))^n*sin(f*x + e)/(f*n - f)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e), x)

maple [B] time = 0.30, size = 66, normalized size = 2.75

$$\frac{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right) e^{n \ln\left(\frac{b\left(1+\tan^2\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}{2 \tan\left(\frac{e}{2} + \frac{fx}{2}\right)}\right)}}{f(-1+n)\left(1+\tan^2\left(\frac{e}{2} + \frac{fx}{2}\right)\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)*(b*csc(f*x+e))^n,x)

[Out] -2/f/(-1+n)*tan(1/2*e+1/2*f*x)*exp(n*ln(1/2*b*(1+tan(1/2*e+1/2*f*x)^2)/tan(1/2*e+1/2*f*x)))/(1+tan(1/2*e+1/2*f*x)^2)

maxima [A] time = 0.71, size = 29, normalized size = 1.21

$$\frac{b^n \sin(fx+e)^{-n} \sin(fx+e)}{f(n-1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x, algorithm="maxima")

[Out] $-b^n \sin(fx + e)^{-n} \sin(fx + e) / (f(n - 1))$

mupad [B] time = 0.32, size = 28, normalized size = 1.17

$$-\frac{\sin(e + fx) \left(\frac{b}{\sin(e + fx)} \right)^n}{f(n - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)*(b/sin(e + f*x))^n,x)

[Out] $-(\sin(e + fx) * (b/\sin(e + fx))^n) / (f * (n - 1))$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \cos(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)*(b*csc(f*x+e))^n,x)

[Out] Integral((b*csc(e + f*x))^n*cos(e + f*x), x)

3.288 $\int \cos^3(e + fx)(b \csc(e + fx))^n dx$

Optimal. Leaf size=52

$$\frac{b(b \csc(e + fx))^{n-1}}{f(1-n)} - \frac{b^3(b \csc(e + fx))^{n-3}}{f(3-n)}$$

[Out] $-b^3*(b*\csc(f*x+e))^{(-3+n)}/f/(3-n)+b*(b*\csc(f*x+e))^{(-1+n)}/f/(1-n)$

Rubi [A] time = 0.05, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2621, 14}

$$\frac{b(b \csc(e + fx))^{n-1}}{f(1-n)} - \frac{b^3(b \csc(e + fx))^{n-3}}{f(3-n)}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^3*(b*Csc[e + f*x])^n,x]

[Out] $-((b^3*(b*Csc[e + f*x])^{(-3 + n)})/(f*(3 - n))) + (b*(b*Csc[e + f*x])^{(-1 + n)})/(f*(1 - n))$

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2621

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.)^(m_))*sec[(e_.) + (f_.)*(x_)]^(n_), x_Symbol] := -Dist[(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
\int \cos^3(e + fx)(b \csc(e + fx))^n dx &= -\frac{b^3 \operatorname{Subst}\left(\int x^{-4+n} \left(-1 + \frac{x^2}{b^2}\right) dx, x, b \csc(e + fx)\right)}{f} \\
&= -\frac{b^3 \operatorname{Subst}\left(\int \left(-x^{-4+n} + \frac{x^{-2+n}}{b^2}\right) dx, x, b \csc(e + fx)\right)}{f} \\
&= -\frac{b^3(b \csc(e + fx))^{-3+n}}{f(3-n)} + \frac{b(b \csc(e + fx))^{-1+n}}{f(1-n)}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 45, normalized size = 0.87

$$-\frac{b((n-1)\cos(2(e+fx)) + n-5)(b \csc(e+fx))^{n-1}}{2f(n-3)(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^3*(b*Csc[e + f*x])^n,x]

[Out] -1/2*(b*(-5 + n + (-1 + n)*Cos[2*(e + f*x)])*(b*Csc[e + f*x])^(-1 + n))/(f*(-3 + n)*(-1 + n))

fricas [A] time = 0.88, size = 49, normalized size = 0.94

$$-\frac{\left((n-1)\cos(fx+e)^2 - 2\right)\left(\frac{b}{\sin(fx+e)}\right)^n \sin(fx+e)}{fn^2 - 4fn + 3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^3*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] -((n-1)*cos(f*x+e)^2 - 2)*(b/sin(f*x+e))^n*sin(f*x+e)/(f*n^2 - 4*f*n + 3*f)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^3*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x+e))^n*cos(f*x+e)^3, x)

maple [F] time = 3.86, size = 0, normalized size = 0.00

$$\int (\cos^3(fx + e)) (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^3*(b*csc(f*x+e))^n,x)

[Out] int(cos(f*x+e)^3*(b*csc(f*x+e))^n,x)

maxima [A] time = 0.71, size = 58, normalized size = 1.12

$$\frac{\frac{b^n \sin(fx+e)^{-n} \sin(fx+e)^3}{n-3} - \frac{b^n \sin(fx+e)^{-n} \sin(fx+e)}{n-1}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^3*(b*csc(f*x+e))^n,x, algorithm="maxima")

[Out] (b^n*sin(f*x + e)^(-n)*sin(f*x + e)^3/(n - 3) - b^n*sin(f*x + e)^(-n)*sin(f*x + e)/(n - 1))/f

mupad [B] time = 0.63, size = 66, normalized size = 1.27

$$\frac{\left(\frac{b}{\sin(e+fx)}\right)^n (9 \sin(e+fx) + \sin(3e+3fx) - n \sin(e+fx) - n \sin(3e+3fx))}{4f(n^2 - 4n + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^3*(b/sin(e + f*x))^n,x)

[Out] ((b/sin(e + f*x))^n*(9*sin(e + f*x) + sin(3*e + 3*f*x) - n*sin(e + f*x) - n*sin(3*e + 3*f*x)))/(4*f*(n^2 - 4*n + 3))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**3*(b*csc(f*x+e))**n,x)

[Out] Timed out

3.289 $\int \cos^5(e + fx)(b \csc(e + fx))^n dx$

Optimal. Leaf size=78

$$\frac{b^5(b \csc(e + fx))^{n-5}}{f(5-n)} - \frac{2b^3(b \csc(e + fx))^{n-3}}{f(3-n)} + \frac{b(b \csc(e + fx))^{n-1}}{f(1-n)}$$

[Out] $b^5*(b*\csc(f*x+e))^{(-5+n)}/f/(5-n)-2*b^3*(b*\csc(f*x+e))^{(-3+n)}/f/(3-n)+b*(b*\csc(f*x+e))^{(-1+n)}/f/(1-n)$

Rubi [A] time = 0.06, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2621, 270}

$$\frac{b^5(b \csc(e + fx))^{n-5}}{f(5-n)} - \frac{2b^3(b \csc(e + fx))^{n-3}}{f(3-n)} + \frac{b(b \csc(e + fx))^{n-1}}{f(1-n)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[e + f*x]^5*(b*\text{Csc}[e + f*x])^n, x]$

[Out] $(b^5*(b*\text{Csc}[e + f*x])^{(-5 + n)})/(f*(5 - n)) - (2*b^3*(b*\text{Csc}[e + f*x])^{(-3 + n)})/(f*(3 - n)) + (b*(b*\text{Csc}[e + f*x])^{(-1 + n)})/(f*(1 - n))$

Rule 270

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_*) + (b_*)*(x_)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2621

$\text{Int}[(\csc[(e_*) + (f_*)*(x_*)]*(a_*)^{(m_*)}\sec[(e_*) + (f_*)*(x_*)]^{(n_*)}), x_Symbol] \rightarrow -\text{Dist}[(f*a^n)^{-1}, \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1+x^2/a^2)^{((n+1)/2)}, x], x, a*\csc[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n+1)/2] \ \&\& \ !(\text{IntegerQ}[(m+1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned}
\int \cos^5(e + fx)(b \csc(e + fx))^n dx &= -\frac{b^5 \text{Subst}\left(\int x^{-6+n} \left(-1 + \frac{x^2}{b^2}\right)^2 dx, x, b \csc(e + fx)\right)}{f} \\
&= -\frac{b^5 \text{Subst}\left(\int \left(x^{-6+n} - \frac{2x^{-4+n}}{b^2} + \frac{x^{-2+n}}{b^4}\right) dx, x, b \csc(e + fx)\right)}{f} \\
&= \frac{b^5 (b \csc(e + fx))^{-5+n}}{f(5-n)} - \frac{2b^3 (b \csc(e + fx))^{-3+n}}{f(3-n)} + \frac{b (b \csc(e + fx))^{-1+n}}{f(1-n)}
\end{aligned}$$

Mathematica [A] time = 0.56, size = 81, normalized size = 1.04

$$\frac{\sin^5(e + fx) \left((n^2 - 8n + 15) \csc^4(e + fx) - 2(n^2 - 6n + 5) \csc^2(e + fx) + n^2 - 4n + 3 \right) (b \csc(e + fx))^n}{f(n-5)(n-3)(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^5*(b*Csc[e + f*x])^n,x]

[Out] -(((b*Csc[e + f*x])^n*(3 - 4*n + n^2 - 2*(5 - 6*n + n^2)*Csc[e + f*x]^2 + (15 - 8*n + n^2)*Csc[e + f*x]^4)*Sin[e + f*x]^5)/(f*(-5 + n)*(-3 + n)*(-1 + n)))

fricas [A] time = 1.08, size = 73, normalized size = 0.94

$$\frac{\left((n^2 - 4n + 3) \cos(fx + e)^4 - 4(n-1) \cos(fx + e)^2 + 8 \right) \left(\frac{b}{\sin(fx+e)} \right)^n \sin(fx + e)}{fn^3 - 9fn^2 + 23fn - 15f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^5*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] -((n^2 - 4*n + 3)*cos(f*x + e)^4 - 4*(n - 1)*cos(f*x + e)^2 + 8)*(b/sin(f*x + e))^n*sin(f*x + e)/(f*n^3 - 9*f*n^2 + 23*f*n - 15*f)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^5 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^5*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e)^5, x)

maple [F] time = 4.43, size = 0, normalized size = 0.00

$$\int (\cos^5(fx + e))(b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^5*(b*csc(f*x+e))^n,x)

[Out] int(cos(f*x+e)^5*(b*csc(f*x+e))^n,x)

maxima [A] time = 0.69, size = 86, normalized size = 1.10

$$-\frac{\frac{b^n \sin(fx+e)^{-n} \sin(fx+e)^5}{n-5} - \frac{2b^n \sin(fx+e)^{-n} \sin(fx+e)^3}{n-3} + \frac{b^n \sin(fx+e)^{-n} \sin(fx+e)}{n-1}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^5*(b*csc(f*x+e))^n,x, algorithm="maxima")

[Out] -(b^n*sin(f*x + e)^(-n)*sin(f*x + e)^5/(n - 5) - 2*b^n*sin(f*x + e)^(-n)*sin(f*x + e)^3/(n - 3) + b^n*sin(f*x + e)^(-n)*sin(f*x + e)/(n - 1))/f

mupad [B] time = 1.41, size = 134, normalized size = 1.72

$$\frac{\left(\frac{b}{\sin(e+fx)}\right)^n (150 \sin(e+fx) + 25 \sin(3e+3fx) + 3 \sin(5e+5fx) + 3n^2 \sin(3e+3fx) + n^2 \sin(5e+5fx) - 24n \sin(e+fx) - 28n \sin(3e+3fx) - 4n \sin(5e+5fx) + 2n^2 \sin(e+fx))}{16f(n^3 - 9n^2 + 23n - 15)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^5*(b/sin(e + f*x))^n,x)

[Out] -((b/sin(e + f*x))^n*(150*sin(e + f*x) + 25*sin(3*e + 3*f*x) + 3*sin(5*e + 5*f*x) + 3*n^2*sin(3*e + 3*f*x) + n^2*sin(5*e + 5*f*x) - 24*n*sin(e + f*x) - 28*n*sin(3*e + 3*f*x) - 4*n*sin(5*e + 5*f*x) + 2*n^2*sin(e + f*x)))/(16*f*(23*n - 9*n^2 + n^3 - 15))

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**5*(b*csc(f*x+e))**n,x)

[Out] Timed out

3.290 $\int (b \csc(e + fx))^n \sec^6(e + fx) dx$

Optimal. Leaf size=72

$$\frac{b\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{7}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] b*(b*csc(f*x+e))⁽⁻¹⁺ⁿ⁾*hypergeom([7/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)²) *sec(f*x+e)*(cos(f*x+e)²)^(1/2)/f/(1-n)

Rubi [A] time = 0.08, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{b\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{7}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])ⁿ*Sec[e + f*x]⁶,x]

[Out] (b*Sqrt[Cos[e + f*x]²]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[7/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]²]*Sec[e + f*x])/(f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^{(2*IntPart[(n - 1)/2] + 1)}*(b*Cos[e + f*x])^{(2*FracPart[(n - 1)/2])}*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]²])/(a*f*(m + 1)*(Cos[e + f*x]²)^{FracPart[(n - 1)/2]}), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a²*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1))/b², Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])ⁿ), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int (b \csc(e + fx))^n \sec^6(e + fx) dx = \left(b^2 (b \csc(e + fx))^{-1+n} (b \sin(e + fx))^{-1+n} \right) \int \sec^6(e + fx) (b \sin(e + fx))^{-n} dx$$

$$= \frac{b \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{7}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sec(e + fx)}{f(1-n)}$$

Mathematica [A] time = 0.61, size = 77, normalized size = 1.07

$$\frac{\tan(e + fx) \sec^2(e + fx)^{-n/2} (b \csc(e + fx))^n {}_2F_1\left(-\frac{n}{2} - 2, \frac{1}{2} - \frac{n}{2}; \frac{3}{2} - \frac{n}{2}; -\tan^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^6,x]

[Out] ((b*Csc[e + f*x])^n*Hypergeometric2F1[-2 - n/2, 1/2 - n/2, 3/2 - n/2, -Tan[e + f*x]^2]*Tan[e + f*x])/(f*(1 - n)*(Sec[e + f*x]^2)^(n/2))

fricas [F] time = 1.06, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \sec(fx + e)^6, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^6,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e)^6, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^6,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^6, x)

maple [F] time = 1.28, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (\sec^6(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n*sec(f*x+e)^6,x)`

[Out] `int((b*csc(f*x+e))^n*sec(f*x+e)^6,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^6 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*sec(f*x+e)^6,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*sec(f*x + e)^6, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/sin(e + f*x))^n/cos(e + f*x)^6,x)`

[Out] `int((b/sin(e + f*x))^n/cos(e + f*x)^6, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))**n*sec(f*x+e)**6,x)`

[Out] Timed out

3.291 $\int (b \csc(e + fx))^n \sec^4(e + fx) dx$

Optimal. Leaf size=72

$$\frac{b\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] b*(b*csc(f*x+e))^(−1+n)*hypergeom([5/2, 1/2−1/2*n], [3/2−1/2*n], sin(f*x+e)^2)*sec(f*x+e)*(cos(f*x+e)^2)^(1/2)/f/(1−n)

Rubi [A] time = 0.08, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{b\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*Sec[e + f*x]^4,x]

[Out] (b*Sqrt[Cos[e + f*x]^2]*(b*Csc[e + f*x])^(−1 + n)*Hypergeometric2F1[5/2, (1 − n)/2, (3 − n)/2, Sin[e + f*x]^2]*Sec[e + f*x])/(f*(1 − n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n − 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n − 1)/2])*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 − n)/2, (3 + m)/2, Sin[e + f*x]^2)]/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n − 1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m − 1)*(b*Sec[e + f*x])^(n + 1))*(a*Sin[e + f*x])^(m − 1)*(b*Cos[e + f*x])^(n + 1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x]^n), x), x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[−m, −n]

Rubi steps

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx = \left(b^2 (b \csc(e + fx))^{-1+n} (b \sin(e + fx))^{-1+n} \right) \int \sec^4(e + fx) (b \sin(e + fx))^{-n} dx$$

$$= \frac{b \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sec(e + fx)}{f(1-n)}$$

Mathematica [A] time = 0.53, size = 77, normalized size = 1.07

$$\frac{\tan(e + fx) \sec^2(e + fx)^{-n/2} (b \csc(e + fx))^n {}_2F_1\left(-\frac{n}{2} - 1, \frac{1}{2} - \frac{n}{2}; \frac{3}{2} - \frac{n}{2}; -\tan^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^4,x]

[Out] ((b*Csc[e + f*x])^n*Hypergeometric2F1[-1 - n/2, 1/2 - n/2, 3/2 - n/2, -Tan[e + f*x]^2]*Tan[e + f*x])/(f*(1 - n)*(Sec[e + f*x]^2)^(n/2))

fricas [F] time = 1.11, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \sec(fx + e)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^4,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^4,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^4, x)

maple [F] time = 1.22, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (\sec^4(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n*sec(f*x+e)^4,x)`

[Out] `int((b*csc(f*x+e))^n*sec(f*x+e)^4,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*sec(f*x+e)^4,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*sec(f*x + e)^4, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/sin(e + f*x))^n/cos(e + f*x)^4,x)`

[Out] `int((b/sin(e + f*x))^n/cos(e + f*x)^4, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \sec^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))**n*sec(f*x+e)**4,x)`

[Out] `Integral((b*csc(e + f*x))**n*sec(e + f*x)**4, x)`

3.292 $\int (b \csc(e + fx))^n \sec^2(e + fx) dx$

Optimal. Leaf size=72

$$\frac{b\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

[Out] b*(b*csc(f*x+e))⁽⁻¹⁺ⁿ⁾*hypergeom([3/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)²)
*sec(f*x+e)*(cos(f*x+e)²)^(1/2)/f/(1-n)

Rubi [A] time = 0.08, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{b\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])ⁿ*Sec[e + f*x]²,x]

[Out] (b*Sqrt[Cos[e + f*x]²]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[3/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]²]*Sec[e + f*x])/ (f*(1 - n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^{(2*IntPart[(n - 1)/2] + 1)}*(b*Cos[e + f*x])^{(2*FracPart[(n - 1)/2])}*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]²])/(a*f*(m + 1)*(Cos[e + f*x]²)^{FracPart[(n - 1)/2]}), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a²*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1))/b², Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])ⁿ), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx = \left(b^2 (b \csc(e + fx))^{-1+n} (b \sin(e + fx))^{-1+n} \right) \int \sec^2(e + fx) (b \sin(e + fx))^{-n} dx$$

$$= \frac{b \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sec(e + fx)}{f(1-n)}$$

Mathematica [A] time = 0.47, size = 75, normalized size = 1.04

$$\frac{\tan(e + fx) \sec^2(e + fx)^{-n/2} (b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2} - \frac{n}{2}, -\frac{n}{2}; \frac{3}{2} - \frac{n}{2}; -\tan^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sec[e + f*x]^2,x]

[Out] ((b*Csc[e + f*x])^n*Hypergeometric2F1[1/2 - n/2, -1/2*n, 3/2 - n/2, -Tan[e + f*x]^2]*Tan[e + f*x])/(f*(1 - n)*(Sec[e + f*x]^2)^(n/2))

fricas [F] time = 0.86, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \sec(fx + e)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^2,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*sec(f*x + e)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec^2(fx + e) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*sec(f*x+e)^2,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*sec(f*x + e)^2, x)

maple [F] time = 1.12, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (\sec^2(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n*sec(f*x+e)^2,x)`

[Out] `int((b*csc(f*x+e))^n*sec(f*x+e)^2,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sec(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*sec(f*x+e)^2,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*sec(f*x + e)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\cos(e+fx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/sin(e + f*x))^n/cos(e + f*x)^2,x)`

[Out] `int((b/sin(e + f*x))^n/cos(e + f*x)^2, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \sec^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))**n*sec(f*x+e)**2,x)`

[Out] `Integral((b*csc(e + f*x))**n*sec(e + f*x)**2, x)`

3.293 $\int (b \csc(e + fx))^n dx$

Optimal. Leaf size=72

$$\frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

[Out] b*cos(f*x+e)*(b*csc(f*x+e))^(n-1)*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)/(cos(f*x+e)^2)^(1/2)

Rubi [A] time = 0.03, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3772, 2643}

$$\frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n,x]

[Out] (b*cos[e + f*x]*(b*csc[e + f*x])^(n-1)*Hypergeometric2F1[1/2, (1-n)/2, (3-n)/2, Sin[e + f*x]^2])/(f*(1-n)*Sqrt[Cos[e + f*x]^2])

Rule 2643

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(Cos[c + d*x]*(b*Ssin[c + d*x])^(n+1)*Hypergeometric2F1[1/2, (n+1)/2, (n+3)/2, Sin[c + d*x]^2])/(b*d*(n+1)*Sqrt[Cos[c + d*x]^2]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3772

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n-1)*((Sin[c + d*x]/b)^(n-1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\int (b \csc(e + fx))^n dx = (b \csc(e + fx))^n \left(\frac{\sin(e + fx)}{b} \right)^n \int \left(\frac{\sin(e + fx)}{b} \right)^{-n} dx$$

$$= \frac{\cos(e + fx)(b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sin(e + fx)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

Mathematica [A] time = 0.11, size = 65, normalized size = 0.90

$$\frac{\sin(e + fx) \cos(e + fx) \sin^2(e + fx)^{\frac{n-1}{2}} (b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{3}{2}; \cos^2(e + fx)\right)}{f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n,x]

[Out] -((Cos[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[1/2, (1 + n)/2, 3/2, Cos[e + f*x]^2]*Sin[e + f*x]*(Sin[e + f*x]^2)^((-1 + n)/2))/f)

fricas [F] time = 1.17, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n, x)

maple [F] time = 1.51, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n,x)`

[Out] `int((b*csc(f*x+e))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/sin(e + f*x))^n,x)`

[Out] `int((b/sin(e + f*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))**n,x)`

[Out] `Integral((b*csc(e + f*x))**n, x)`

3.294 $\int \cos^2(e + fx)(b \csc(e + fx))^n dx$

Optimal. Leaf size=72

$$\frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(-\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

[Out] b*cos(f*x+e)*(b*csc(f*x+e))⁽⁻¹⁺ⁿ⁾*hypergeom([-1/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)²)/f/(1-n)/(cos(f*x+e)²)^(1/2)

Rubi [A] time = 0.08, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(-\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^2*(b*Csc[e + f*x])^n,x]

[Out] (b*Cos[e + f*x]*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[-1/2, (1 - n)/2, (3 - n)/2, Sin[e + f*x]^2])/f*(1 - n)*Sqrt[Cos[e + f*x]^2]

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^{(2*FracPart[(n - 1)/2])}*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2])/(a*f*(m + 1)*(Cos[e + f*x]^2)^{FracPart[(n - 1)/2]}), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])ⁿ), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int \cos^2(e + fx)(b \csc(e + fx))^n dx = \left(b^2 (b \csc(e + fx))^{-1+n} (b \sin(e + fx))^{-1+n} \right) \int \cos^2(e + fx)(b \sin(e + fx))^{-n} dx$$

$$= \frac{b \cos(e + fx)(b \csc(e + fx))^{-1+n} {}_2F_1\left(-\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

Mathematica [B] time = 0.48, size = 165, normalized size = 2.29

$$\frac{2 \tan\left(\frac{1}{2}(e + fx)\right) \sec^2\left(\frac{1}{2}(e + fx)\right)^{-n} (b \csc(e + fx))^n \left({}_2F_1\left(1 - n, \frac{1}{2} - \frac{n}{2}; \frac{3}{2} - \frac{n}{2}; -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 4 {}_2F_1\left(2 - n, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) \right)}{f(n-1)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^2*(b*Csc[e + f*x])^n,x]

[Out] (-2*(b*Csc[e + f*x])^n*(Hypergeometric2F1[1 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2] - 4*Hypergeometric2F1[2 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2] + 4*Hypergeometric2F1[3 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2])/(f*(-1 + n)*(Sec[(e + f*x)/2]^2)^n)

fricas [F] time = 2.14, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \cos(fx + e)^2, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*cos(f*x + e)^2, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e)^2, x)

maple [F] time = 2.68, size = 0, normalized size = 0.00

$$\int (\cos^2(fx + e))(b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(f*x+e)^2*(b*csc(f*x+e))^n,x)`

[Out] `int(cos(f*x+e)^2*(b*csc(f*x+e))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(f*x+e)^2*(b*csc(f*x+e))^n,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*cos(f*x + e)^2, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(e + fx)^2 \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(e + f*x)^2*(b/sin(e + f*x))^n,x)`

[Out] `int(cos(e + f*x)^2*(b/sin(e + f*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \cos^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(f*x+e)**2*(b*csc(f*x+e))**n,x)`

[Out] `Integral((b*csc(e + f*x))**n*cos(e + f*x)**2, x)`

3.295 $\int \cos^4(e + fx)(b \csc(e + fx))^n dx$

Optimal. Leaf size=72

$$\frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(-\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

[Out] b*cos(f*x+e)*(b*csc(f*x+e))^(1-n)*hypergeom([-3/2, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)/f/(1-n)/(cos(f*x+e)^2)^(1/2)

Rubi [A] time = 0.08, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2631, 2577}

$$\frac{b \cos(e + fx)(b \csc(e + fx))^{n-1} {}_2F_1\left(-\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^4*(b*Csc[e + f*x])^n,x]

[Out] (b*Cos[e + f*x]*(b*Csc[e + f*x])^(1-n)*Hypergeometric2F1[-3/2, (1-n)/2, (3-n)/2, Sin[e + f*x]^2])/(f*(1-n)*Sqrt[Cos[e + f*x]^2])

Rule 2577

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[(b^(2*IntPart[(n-1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n-1)/2])*(a*Sin[e + f*x])^(m+1)*Hypergeometric2F1[(1+m)/2, (1-n)/2, (3+m)/2, Sin[e + f*x]^2])/(a*f*(m+1)*(Cos[e + f*x]^2)^FracPart[(n-1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n+1))*(a*Sin[e + f*x])^(m-1)*(b*Cos[e + f*x])^(n+1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int \cos^4(e + fx)(b \csc(e + fx))^n dx = \left(b^2(b \csc(e + fx))^{-1+n}(b \sin(e + fx))^{-1+n} \right) \int \cos^4(e + fx)(b \sin(e + fx))^{-n} dx$$

$$= \frac{b \cos(e + fx)(b \csc(e + fx))^{-1+n} {}_2F_1\left(-\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{f(1-n)\sqrt{\cos^2(e + fx)}}$$

Mathematica [B] time = 0.62, size = 246, normalized size = 3.42

$$\frac{2 \tan\left(\frac{1}{2}(e + fx)\right) \sec^2\left(\frac{1}{2}(e + fx)\right)^{-n} (b \csc(e + fx))^n \left({}_2F_1\left(1 - n, \frac{1}{2} - \frac{n}{2}; \frac{3}{2} - \frac{n}{2}; -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) - 8 \left({}_2F_1\left(2 - n, \frac{1}{2} - \frac{n}{2}, \frac{3}{2} - \frac{n}{2}, -\tan^2\left(\frac{1}{2}(e + fx)\right)\right)\right)}{\dots}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^4*(b*Csc[e + f*x])^n,x]

[Out] (-2*(b*Csc[e + f*x])^n*(Hypergeometric2F1[1 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2] - 8*(Hypergeometric2F1[2 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2] - 3*Hypergeometric2F1[3 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2] + 4*Hypergeometric2F1[4 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2] - 2*Hypergeometric2F1[5 - n, 1/2 - n/2, 3/2 - n/2, -Tan[(e + f*x)/2]^2]))*Tan[(e + f*x)/2])/(f*(-1 + n)*(Sec[(e + f*x)/2]^2)^n)

fricas [F] time = 1.32, size = 0, normalized size = 0.00

$$\text{integral}\left(\left(b \csc(fx + e)\right)^n \cos(fx + e)^4, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4*(b*csc(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*cos(f*x + e)^4, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4*(b*csc(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*cos(f*x + e)^4, x)

maple [F] time = 3.06, size = 0, normalized size = 0.00

$$\int (\cos^4(fx + e))(b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(f*x+e)^4*(b*csc(f*x+e))^n,x)`

[Out] `int(cos(f*x+e)^4*(b*csc(f*x+e))^n,x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \cos(fx + e)^4 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(f*x+e)^4*(b*csc(f*x+e))^n,x, algorithm="maxima")`

[Out] `integrate((b*csc(f*x + e))^n*cos(f*x + e)^4, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(e + fx)^4 \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(e + f*x)^4*(b/sin(e + f*x))^n,x)`

[Out] `int(cos(e + f*x)^4*(b/sin(e + f*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \cos^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(f*x+e)**4*(b*csc(f*x+e))**n,x)`

[Out] `Integral((b*csc(e + f*x))**n*cos(e + f*x)**4, x)`

3.296 $\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx$

Optimal. Leaf size=81

$$\frac{b \cos^2(e + fx)^{5/4} (c \sec(e + fx))^{5/2} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{5}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{cf(1-n)}$$

[Out] b*(cos(f*x+e)^2)^(5/4)*(b*csc(f*x+e))^(n-1)*hypergeom([5/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(c*sec(f*x+e))^(5/2)/c/f/(1-n)

Rubi [A] time = 0.11, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2631, 2577}

$$\frac{b \cos^2(e + fx)^{5/4} (c \sec(e + fx))^{5/2} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{5}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{cf(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*(c*Sec[e + f*x])^(3/2), x]

[Out] (b*(Cos[e + f*x]^2)^(5/4)*(b*Csc[e + f*x])^(n-1)*Hypergeometric2F1[5/4, (1-n)/2, (3-n)/2, Sin[e + f*x]^2]*(c*Sec[e + f*x])^(5/2))/(c*f*(1-n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[(b^(2*IntPart[(n-1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n-1)/2])*(a*Sin[e + f*x])^(m+1)*Hypergeometric2F1[(1+m)/2, (1-n)/2, (3+m)/2, Sin[e + f*x]^2])/(a*f*(m+1)*(Cos[e + f*x]^2)^FracPart[(n-1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_.)])^n, x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n+1))*(a*Sin[e + f*x])^(m-1)*(b*Cos[e + f*x])^(n+1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int (b \csc(e + fx))^n (c \sec(e + fx))^{3/2} dx = \frac{(b^2 (c \cos(e + fx))^{5/2} (b \csc(e + fx))^{-1+n} (c \sec(e + fx))^{5/2} (b \sin(e + fx)))}{c^2}$$

$$= \frac{b \cos^2(e + fx)^{5/4} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{5}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) (c \sec(e + fx))^{3/2}}{cf(1-n)}$$

Mathematica [A] time = 1.89, size = 92, normalized size = 1.14

$$\frac{2 \cot(e + fx) (c \sec(e + fx))^{3/2} \left(-\tan^2(e + fx)\right)^{\frac{n+1}{2}} (b \csc(e + fx))^n {}_2F_1\left(\frac{n+1}{2}, \frac{1}{4}(2n+3); \frac{1}{4}(2n+7); \sec^2(e + fx)\right)}{f(2n+3)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*(c*Sec[e + f*x])^(3/2),x]

[Out] (2*Cot[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[(1 + n)/2, (3 + 2*n)/4, (7 + 2*n)/4, Sec[e + f*x]^2]*(c*Sec[e + f*x])^(3/2)*(-Tan[e + f*x]^2)^(1 + n)/2)/(f*(3 + 2*n))

fricas [F] time = 1.62, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \sec(fx + e)} (b \csc(fx + e))^n c \sec(fx + e), x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n*c*sec(f*x + e), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(fx + e))^{\frac{3}{2}} (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((c*sec(f*x + e))^(3/2)*(b*csc(f*x + e))^n, x)

maple [F] time = 0.89, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (c \sec(fx + e))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x)`

[Out] `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sec(fx + e))^{\frac{3}{2}} (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate((c*sec(f*x + e))^(3/2)*(b*csc(f*x + e))^n, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{c}{\cos(e + fx)} \right)^{3/2} \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(e + f*x))^(3/2)*(b/sin(e + f*x))^n,x)`

[Out] `int((c/cos(e + f*x))^(3/2)*(b/sin(e + f*x))^n, x)`

sympy [F(-1)] time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))**n*(c*sec(f*x+e))**(3/2),x)`

[Out] Timed out

3.297 $\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$

Optimal. Leaf size=81

$$\frac{b \cos^2(e + fx)^{3/4} (c \sec(e + fx))^{3/2} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{3}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{cf(1-n)}$$

[Out] b*(cos(f*x+e)^2)^(3/4)*(b*csc(f*x+e))^(n-1)*hypergeom([3/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(c*sec(f*x+e))^(3/2)/c/f/(1-n)

Rubi [A] time = 0.10, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2631, 2577}

$$\frac{b \cos^2(e + fx)^{3/4} (c \sec(e + fx))^{3/2} (b \csc(e + fx))^{n-1} {}_2F_1\left(\frac{3}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{cf(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n*Sqrt[c*Sec[e + f*x]], x]

[Out] (b*(Cos[e + f*x]^2)^(3/4)*(b*Csc[e + f*x])^(n-1)*Hypergeometric2F1[3/4, (1-n)/2, (3-n)/2, Sin[e + f*x]^2]*(c*Sec[e + f*x])^(3/2))/(c*f*(1-n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(b^(2*IntPart[(n-1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n-1)/2])*(a*Sin[e + f*x])^(m+1)*Hypergeometric2F1[(1+m)/2, (1-n)/2, (3+m)/2, Sin[e + f*x]^2)]/(a*f*(m+1)*(Cos[e + f*x]^2)^FracPart[(n-1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n+1))*(a*Sin[e + f*x])^(m-1)*(b*Cos[e + f*x])^(n+1)]/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx = \frac{(b^2(c \cos(e + fx))^{3/2}(b \csc(e + fx))^{-1+n}(c \sec(e + fx))^{3/2}(b \sin(e + fx))^{-1})}{c^2}$$

$$= \frac{b \cos^2(e + fx)^{3/4}(b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{3}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) (c \sec(e + fx))}{cf(1-n)}$$

Mathematica [A] time = 1.72, size = 90, normalized size = 1.11

$$\frac{2 \cot(e + fx) \sqrt{c \sec(e + fx)} (-\tan^2(e + fx))^{\frac{n+1}{2}} (b \csc(e + fx))^n {}_2F_1\left(\frac{n+1}{2}, \frac{1}{4}(2n+1); \frac{1}{4}(2n+5); \sec^2(e + fx)\right)}{2fn + f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*Sqrt[c*Sec[e + f*x]], x]

[Out] (2*Cot[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[(1 + n)/2, (1 + 2*n)/4, (5 + 2*n)/4, Sec[e + f*x]^2]*Sqrt[c*Sec[e + f*x]]*(-Tan[e + f*x]^2)^(1 + n)/2)/(f + 2*f*n)

fricas [F] time = 0.88, size = 0, normalized size = 0.00

$$\text{integral}\left(\sqrt{c \sec(fx + e)} (b \csc(fx + e))^n, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2), x, algorithm="fricas")

[Out] integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n, x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sec(fx + e)} (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n, x)

maple [F] time = 0.92, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n \sqrt{c \sec(fx + e)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x)`

[Out] `int((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x)`

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sec(fx + e)} (b \csc(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n, x)`

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{\frac{c}{\cos(e + fx)}} \left(\frac{b}{\sin(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c/cos(e + f*x))^(1/2)*(b/sin(e + f*x))^n,x)`

[Out] `int((c/cos(e + f*x))^(1/2)*(b/sin(e + f*x))^n, x)`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int (b \csc(e + fx))^n \sqrt{c \sec(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*csc(f*x+e))^n*(c*sec(f*x+e))^(1/2),x)`

[Out] `Integral((b*csc(e + f*x))^n*sqrt(c*sec(e + f*x)), x)`

$$3.298 \quad \int \frac{(b \csc(e+fx))^n}{\sqrt{c \sec(e+fx)}} dx$$

Optimal. Leaf size=81

$$\frac{b^4 \sqrt{\cos^2(e+fx)} \sqrt{c \sec(e+fx)} (b \csc(e+fx))^{n-1} {}_2F_1\left(\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e+fx)\right)}{cf(1-n)}$$

[Out] b*(cos(f*x+e)^2)^(1/4)*(b*csc(f*x+e))^(1-n)*hypergeom([1/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)^2)*(c*sec(f*x+e))^(1/2)/c/f/(1-n)

Rubi [A] time = 0.10, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2631, 2577}

$$\frac{b^4 \sqrt{\cos^2(e+fx)} \sqrt{c \sec(e+fx)} (b \csc(e+fx))^{n-1} {}_2F_1\left(\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e+fx)\right)}{cf(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])^n/Sqrt[c*Sec[e + f*x]], x]

[Out] (b*(Cos[e + f*x]^2)^(1/4)*(b*Csc[e + f*x])^(1-n)*Hypergeometric2F1[1/4, (1-n)/2, (3-n)/2, Sin[e + f*x]^2]*Sqrt[c*Sec[e + f*x]])/(c*f*(1-n))

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[(b^(2*IntPart[(n-1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n-1)/2])*(a*Sin[e + f*x])^(m+1)*Hypergeometric2F1[(1+m)/2, (1-n)/2, (3+m)/2, Sin[e + f*x]^2])/(a*f*(m+1)*(Cos[e + f*x]^2)^FracPart[(n-1)/2]), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_.)])^n, x_Symbol] :> Dist[(a^2*(a*Csc[e + f*x])^(m-1)*(b*Sec[e + f*x])^(n+1)*(a*Sin[e + f*x])^(m-1)*(b*Cos[e + f*x])^(n+1))/b^2, Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx = \frac{(b^2 \sqrt{c \cos(e + fx)} (b \csc(e + fx))^{-1+n} \sqrt{c \sec(e + fx)} (b \sin(e + fx))^{-1+n}) \int \sqrt{c \cos(e + fx)}}{c^2}$$

$$= \frac{b^4 \sqrt{\cos^2(e + fx)} (b \csc(e + fx))^{-1+n} {}_2F_1\left(\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right) \sqrt{c \sec(e + fx)}}{c f(1-n)}$$

Mathematica [C] time = 3.18, size = 326, normalized size = 4.02

$$\frac{4(n-3) \sin\left(\frac{1}{2}(e+fx)\right)}{f(n-1) \sqrt{c \sec(e+fx)} \left(2(3-2n) \sin^2\left(\frac{1}{2}(e+fx)\right) F_1\left(\frac{3}{2} - \frac{n}{2}; -\frac{1}{2}, \frac{5}{2} - n; \frac{5}{2} - \frac{n}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Csc[e + f*x])^n/Sqrt[c*Sec[e + f*x]],x]

[Out] (-4*(-3 + n)*AppellF1[1/2 - n/2, -1/2, 3/2 - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*(b*Csc[e + f*x])^n*Sin[(e + f*x)/2])/((f*(-1 + n)*Sqrt[c*Sec[e + f*x]]*(2*(-3 + n)*AppellF1[1/2 - n/2, -1/2, 3/2 - n, 3/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^2 - AppellF1[3/2 - n/2, 1/2, 3/2 - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(-1 + Cos[e + f*x]) + 2*(3 - 2*n)*AppellF1[3/2 - n/2, -1/2, 5/2 - n, 5/2 - n/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sin[(e + f*x)/2]^2))

fricas [F] time = 1.05, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{c \sec(fx + e)} (b \csc(fx + e))^n}{c \sec(fx + e)}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n/(c*sec(f*x + e)), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n/sqrt(c*sec(f*x + e)), x)

maple [F] time = 0.79, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x)

[Out] int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(fx + e))^n}{\sqrt{c \sec(fx + e)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n/sqrt(c*sec(f*x + e)), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\sqrt{\frac{c}{\cos(e+fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(1/2),x)

[Out] int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(1/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(e + fx))^n}{\sqrt{c \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*csc(f*x+e)**n/(c*sec(f*x+e))**(1/2),x)
```

```
[Out] Integral((b*csc(e + f*x)**n/sqrt(c*sec(e + f*x)), x)
```

$$3.299 \quad \int \frac{(b \csc(e+fx))^n}{(c \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=81

$$\frac{b(b \csc(e+fx))^{n-1} {}_2F_1\left(-\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e+fx)\right)}{cf(1-n)\sqrt[4]{\cos^2(e+fx)}\sqrt{c \sec(e+fx)}}$$

[Out] b*(b*csc(f*x+e))⁽⁻¹⁺ⁿ⁾*hypergeom([-1/4, 1/2-1/2*n], [3/2-1/2*n], sin(f*x+e)²)/c/f/(1-n)/(cos(f*x+e)²)^(1/4)/(c*sec(f*x+e))^(1/2)

Rubi [A] time = 0.11, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2631, 2577}

$$\frac{b(b \csc(e+fx))^{n-1} {}_2F_1\left(-\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e+fx)\right)}{cf(1-n)\sqrt[4]{\cos^2(e+fx)}\sqrt{c \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Csc[e + f*x])ⁿ/(c*Sec[e + f*x])^(3/2), x]

[Out] (b*(b*Csc[e + f*x])^(-1 + n)*Hypergeometric2F1[-1/4, (1 - n)/2, (3 - n)/2, Sin[e + f*x]²]/(c*f*(1 - n)*(Cos[e + f*x]²)^(1/4)*Sqrt[c*Sec[e + f*x]])

Rule 2577

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(b^{(2*IntPart[(n - 1)/2] + 1)}*(b*Cos[e + f*x])^{(2*FracPart[(n - 1)/2])}*(a*Sin[e + f*x])^(m + 1)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]²]/(a*f*(m + 1)*(Cos[e + f*x]²)^{FracPart[(n - 1)/2]}), x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2631

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] :> Dist[(a²*(a*Csc[e + f*x])^(m - 1)*(b*Sec[e + f*x])^(n + 1)*(a*Sin[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1))/b², Int[1/((a*Sin[e + f*x])^m*(b*Cos[e + f*x])ⁿ), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !SimplerQ[-m, -n]

Rubi steps

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{3/2}} dx = \frac{(b^2(b \csc(e + fx))^{-1+n}(b \sin(e + fx))^{-1+n}) \int (c \cos(e + fx))^{3/2} (b \sin(e + fx))^{-n} dx}{c^2 \sqrt{c \cos(e + fx)} \sqrt{c \sec(e + fx)}}$$

$$= \frac{b(b \csc(e + fx))^{-1+n} {}_2F_1\left(-\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \sin^2(e + fx)\right)}{cf(1-n)\sqrt[4]{\cos^2(e + fx)} \sqrt{c \sec(e + fx)}}$$

Mathematica [A] time = 1.02, size = 115, normalized size = 1.42

$$\frac{2 \cos(2(e + fx)) \cot(e + fx) \sqrt{c \sec(e + fx)} \left(-\tan^2(e + fx)\right)^{\frac{n+1}{2}} (b \csc(e + fx))^n {}_2F_1\left(\frac{n+1}{2}, \frac{1}{4}(2n-3); \frac{1}{4}(2n+1)\right)}{c^2 f(2n-3) (\sec^2(e + fx) - 2)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n/(c*Sec[e + f*x])^(3/2),x]

[Out] (-2*Cos[2*(e + f*x)]*Cot[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[(1 + n)/2, (-3 + 2*n)/4, (1 + 2*n)/4, Sec[e + f*x]^2]*Sqrt[c*Sec[e + f*x]]*(-Tan[e + f*x]^2)^((1 + n)/2))/(c^2*f*(-3 + 2*n)*(-2 + Sec[e + f*x]^2))

fricas [F] time = 1.20, size = 0, normalized size = 0.00

$$\text{integral}\left(\frac{\sqrt{c \sec(fx + e)} (b \csc(fx + e))^n}{c^2 \sec(fx + e)^2}, x\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sec(f*x + e))*(b*csc(f*x + e))^n/(c^2*sec(f*x + e)^2), x)

giac [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(fx + e))^n}{(c \sec(fx + e))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n/(c*sec(f*x + e))^(3/2), x)

maple [F] time = 0.73, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(fx + e))^n}{(c \sec(fx + e))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2), x)

[Out] int((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2), x)

maxima [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(fx + e))^n}{(c \sec(fx + e))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n/(c*sec(f*x+e))^(3/2), x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n/(c*sec(f*x + e))^(3/2), x)

mupad [F] time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\sin(e+fx)}\right)^n}{\left(\frac{c}{\cos(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(3/2), x)

[Out] int((b/sin(e + f*x))^n/(c/cos(e + f*x))^(3/2), x)

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \csc(e + fx))^n}{(c \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*csc(f*x+e)**n/(c*sec(f*x+e))**(3/2),x)
```

```
[Out] Integral((b*csc(e + f*x)**n/(c*sec(e + f*x))**(3/2), x)
```


Chapter 4

Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.0.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] :=
  If[ExpnType[result]<=ExpnType[optimal],
```

```

If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],
  If[LeafCount[result]<=2*LeafCount[optimal],
    "A",
    "B"],
  "C"],
If[FreeQ[result,Integrate] && FreeQ[result,Int],
  "C",
"F"]]

```

```
(* ::Text:: *)
```

```
(*The following summarizes the type number assigned an *)
```

```
(*expression based on the functions it involves*)
```

```
(*1 = rational function*)
```

```
(*2 = algebraic function*)
```

```
(*3 = elementary function*)
```

```
(*4 = special function*)
```

```
(*5 = hyperpergeometric function*)
```

```
(*6 = appell function*)
```

```
(*7 = rootsum function*)
```

```
(*8 = integrate function*)
```

```
(*9 = unknown function*)
```

```
ExpnType[expn_] :=
```

```
  If[AtomQ[expn],
```

```
    1,
```

```
  If[ListQ[expn],
```

```
    Max[Map[ExpnType,expn]],
```

```
  If[Head[expn]===Power,
```

```
    If[IntegerQ[expn[[2]],
```

```
      ExpnType[expn[[1]],
```

```
    If[Head[expn[[2]]===Rational,
```

```
      If[IntegerQ[expn[[1]] || Head[expn[[1]]===Rational,
```

```
        1,
```

```
        Max[ExpnType[expn[[1]],2]],
```

```
      Max[ExpnType[expn[[1]],ExpnType[expn[[2]],3]],
```

```
  If[Head[expn]===Plus || Head[expn]===Times,
```

```
    Max[ExpnType[First[expn],ExpnType[Rest[expn]]],
```

```
  If[ElementaryFunctionQ[Head[expn]],
```

```
    Max[3,ExpnType[expn[[1]]],
```

```
  If[SpecialFunctionQ[Head[expn]],
```

```
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
```

```
  If[HypergeometricFunctionQ[Head[expn]],
```

```
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
```

```
  If[AppellFunctionQ[Head[expn]],
```

```
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],
```



```

If[Head[expn]===RootSum,
  Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
If[Head[expn]===Integrate || Head[expn]===Int,
  Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
9]]]]]]]]]]

ElementaryFunctionQ[func_] :=
MemberQ[{
  Exp,Log,
  Sin,Cos,Tan,Cot,Sec,Csc,
  ArcSin,ArcCos,ArcTan,ArcCot,ArcSec,ArcCsc,
  Sinh,Cosh,Tanh,Coth,Sech,Csch,
  ArcSinh,ArcCosh,ArcTanh,ArcCoth,ArcSech,ArcCsch
},func]

SpecialFunctionQ[func_] :=
MemberQ[{
  Erf, Erfc, Erfi,
  FresnelS, FresnelC,
  ExpIntegralE, ExpIntegralEi, LogIntegral,
  SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
  Gamma, LogGamma, PolyGamma,
  Zeta, PolyLog, ProductLog,
  EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]

AppellFunctionQ[func_] :=
MemberQ[{AppellF1},func]

```

4.0.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019  Added debug flag, added 'dilog' to special functions
#                      see problem 156, file Apostol_Problems

GradeAntiderivative := proc(result,optimal)
local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
    debug:=false;

    leaf_count_result:=leafcount(result);
    #do NOT call ExpnType() if leaf size is too large. Recursion problem
    if leaf_count_result > 500000 then
        return "B";
    fi;

    leaf_count_optimal:=leafcount(optimal);

    ExpnType_result:=ExpnType(result);
    ExpnType_optimal:=ExpnType(optimal);

    if debug then
        print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
ExpnType_optimal);
    fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#   is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
#   antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F";
end if;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;

```

```

if is_contains_complex(result) then
  if is_contains_complex(optimal) then
    if debug then
      print("both result and optimal complex");
    fi;
    #both result and optimal complex
    if leaf_count_result<=2*leaf_count_optimal then
      return "A";
    else
      return "B";
    end if
  else #result contains complex but optimal is not
    if debug then
      print("result contains complex but optimal is not");
    fi;
    return "C";
  end if
else # result do not contain complex
  # this assumes optimal do not as well
  if debug then
    print("result do not contain complex, this assumes optimal do not
as well");
  fi;
  if leaf_count_result<=2*leaf_count_optimal then
    if debug then
      print("leaf_count_result<=2*leaf_count_optimal");
    fi;
    return "A";
  else
    if debug then
      print("leaf_count_result>2*leaf_count_optimal");
    fi;
    return "B";
  end if
end if
else #ExpnType(result) > ExpnType(optimal)
  if debug then
    print("ExpnType(result) > ExpnType(optimal)");
  fi;
  return "C";
end if

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false

```

```

#
#Nasser 032417
is_contains_complex:= proc(expression)
  return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+`') or type(expn,'`*`') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  end if
end proc:

```

```

elif HypergeometricFunctionQ(op(0,expn)) then
  max(5,apply(max,map(ExpnType,[op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
  max(6,apply(max,map(ExpnType,[op(expn)])))
elif op(0,expn)='int' then
  max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product. rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

```

```
#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:
```

4.0.3 Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#           Port of original Maple grading function by
#           Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#           added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                    asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                    asinh,acosh,atanh,acoth,asech,acsch
                    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]
```

```

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'`^`')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)
))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,
ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+`') or type
(expn,'`*`')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))

```

```

elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
expn))
    m1 = max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif is_appell_function(expn.func):
    m1 = max(map(expnType, list(expn.args)))
    return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
elif isinstance(expn,RootSum):
    m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,
Apply[List,expn]],7]],
    return max(7,m1)
elif str(expn).find("Integral") != -1:
    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        return "F"

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
    else:
        return "C"

```


4.0.4 SageMath grading function

```

#Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:

```

```

        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    debug=False
    if debug: print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
sinh_integral'
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func ," is special_function")
        else:
            print ("func ", func ," is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U
']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in
sagemath

def is_atom(expn):

    debug=False
    if debug: print ("Enter is_atom")

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-type-in-maple/
    try:
        if expn.parent() is SR:

```

```

        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
    return False

except AttributeError as error:
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #instance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(instance(expn.args[0],
Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
    elif expn.operator() == operator.pow: #instance(expn,Pow)
        if type(expn.operands()[1])==Integer: #instance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #instance(expn.args[1],
Rational)
            if type(expn.operands()[0])==Rational: #instance(expn.args[0],
Rational)
                return 1
            else:
                return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.
args[0]))
        else:
            return max(3,expnType(expn.operands()[0]),expnType(expn.operands()
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
    elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
instance(expn,Add) or instance(expn,Mul)
        m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
        m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))

```

```

    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.
func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #
is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
#is checked before calling the grading function that is passed.
#but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(
expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

```

```
#main function
```

```
def grade_antiderivative(result,optimal):
```

```

    if debug: print ("Enter grade_antiderivative for sagemath")

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    if debug: print ("leaf_count_result=", leaf_count_result, "
leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",
expnType_optimal)

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex

```

```
        if leaf_count_result <= 2*leaf_count_optimal:
            return "A"
        else:
            return "B"
    else: #result contains complex but optimal is not
        return "C"
else: # result do not contain complex, this assumes optimal do not as
well
    if leaf_count_result <= 2*leaf_count_optimal:
        return "A"
    else:
        return "B"
else:
    return "C"
```