

Computer algebra independent integration tests

4-Trig-functions/4.6-Cosecant/4.6.1.4-d-cot- \wedge n-a+b-csc- \wedge m

Nasser M. Abbasi

July 17, 2021 Compiled on July 17, 2021 at 9:45pm

Contents

1	Introduction	5
1.1	Listing of CAS systems tested	5
1.2	Results	6
1.3	Performance	10
1.4	list of integrals that has no closed form antiderivative	11
1.5	list of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	12
1.7	Timing	13
1.8	Verification	13
1.9	Important notes about some of the results	13
1.9.1	Important note about Maxima results	13
1.9.2	Important note about FriCAS and Giac/XCAS results	14
1.9.3	Important note about finding leaf size of antiderivative	14
1.9.4	Important note about Mupad results	15
1.10	Design of the test system	16
2	detailed summary tables of results	17
2.1	List of integrals sorted by grade for each CAS	17
2.1.1	Rubi	17
2.1.2	Mathematica	17
2.1.3	Maple	17
2.1.4	Maxima	18

2.1.5	FriCAS	18
2.1.6	Sympy	18
2.1.7	Giac	18
2.1.8	Mupad	18
2.2	Detailed conclusion table per each integral for all CAS systems	19
2.3	Detailed conclusion table specific for Rubi results	24
3	Listing of integrals	27
3.1	$\int \frac{\tan^4(x)}{a+a \csc(x)} dx$	27
3.2	$\int \frac{\tan^3(x)}{a+a \csc(x)} dx$	31
3.3	$\int \frac{\tan^2(x)}{a+a \csc(x)} dx$	35
3.4	$\int \frac{\tan(x)}{a+a \csc(x)} dx$	39
3.5	$\int \frac{\cot(x)}{a+a \csc(x)} dx$	42
3.6	$\int \frac{\cot^2(x)}{a+a \csc(x)} dx$	45
3.7	$\int \frac{\cot^3(x)}{a+a \csc(x)} dx$	48
3.8	$\int \frac{\cot^4(x)}{a+a \csc(x)} dx$	51
3.9	$\int \frac{\cot^5(x)}{a+a \csc(x)} dx$	55
3.10	$\int \frac{\cot^6(x)}{a+a \csc(x)} dx$	58
3.11	$\int \frac{\cot^7(x)}{a+a \csc(x)} dx$	62
3.12	$\int \frac{\tan^5(x)}{a+b \csc(x)} dx$	66
3.13	$\int \frac{\tan^3(x)}{a+b \csc(x)} dx$	70
3.14	$\int \frac{\tan(x)}{a+b \csc(x)} dx$	74
3.15	$\int \frac{\cot(x)}{a+b \csc(x)} dx$	77
3.16	$\int \frac{\cot^3(x)}{a+b \csc(x)} dx$	80
3.17	$\int \frac{\cot^5(x)}{a+b \csc(x)} dx$	83
3.18	$\int \frac{\cot^7(x)}{a+b \csc(x)} dx$	87
3.19	$\int \frac{\tan^4(x)}{a+b \csc(x)} dx$	91
3.20	$\int \frac{\tan^2(x)}{a+b \csc(x)} dx$	99
3.21	$\int \frac{\cot^2(x)}{a+b \csc(x)} dx$	105
3.22	$\int \frac{\cot^4(x)}{a+b \csc(x)} dx$	110

3.23	$\int \frac{\cot^6(x)}{a+b \csc(x)} dx$	116
4	Listing of Grading functions	125
4.0.1	Mathematica and Rubi grading function	125
4.0.2	Maple grading function	127
4.0.3	Sympy grading function	132
4.0.4	SageMath grading function	135

Chapter 1

Introduction

This report gives the result of running the computer algebra independent integration problems. The listing of the problems are maintained by and can be downloaded from <https://rulebasedintegration.org>

The number of integrals in this report is [23]. This is test number [131].

1.1 Listing of CAS systems tested

The following systems were tested at this time.

1. Mathematica 12.3 (64 bit) on windows 10.
2. Rubi 4.16.1 in Mathematica 12.1 on windows 10.
3. Maple 2021.1 (64 bit) on windows 10.
4. Maxima 5.44 on Linux. (via sageMath 9.3)
5. Fricas 1.3.7 on Linux (via sageMath 9.3)
6. Giac/Xcas 1.7 on Linux. (via sageMath 9.3)
7. Sympy 1.8 under Python 3.8.8 using Anaconda distribution on Ubuntu.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 under windows 10 (64 bit)

Maxima, Fricas and Giac/Xcas were called from inside SageMath. This was done using SageMath integrate command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly using Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in the table below reflects the above.

System	solved	Failed
Rubi	% 100.00 (23)	% 0.00 (0)
Mathematica	% 100.00 (23)	% 0.00 (0)
Maple	% 100.00 (23)	% 0.00 (0)
Maxima	% 78.26 (18)	% 21.74 (5)
Fricas	% 100.00 (23)	% 0.00 (0)
Sympy	% 0.00 (0)	% 100.00 (23)
Giac	% 100.00 (23)	% 0.00 (0)
Mupad	% 100.00 (23)	% 0.00 (0)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	86.96	13.04	0.00	0.00
Maple	78.26	21.74	0.00	0.00
Maxima	60.87	17.39	0.00	21.74
Fricas	73.91	26.09	0.00	0.00
Sympy	0.00	0.00	0.00	100.00
Giac	86.96	13.04	0.00	0.00
Mupad	0.00	100.00	0.00	0.00

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure F .

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned F(-1).

The third is due to an exception generated. Assigned F(-2). This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	0	0.00 %	0.00 %	0.00 %
Maxima	5	0.00 %	0.00 %	100.00 %
Fricas	0	0.00 %	0.00 %	0.00 %
Sympy	23	100.00 %	0.00 %	0.00 %
Giac	0	0.00 %	0.00 %	0.00 %
Mupad	0	0.00 %	0.00 %	0.00 %

Table 1.4: Time and leaf size performance for each CAS

1.3 Performance

The table below summarizes the performance of each CAS system in terms of CPU time and leaf size of results.


System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.12	73.00	1.03	58.00	1.00
Mathematica	0.23	89.26	1.30	71.00	1.04
Maple	0.40	100.04	1.42	64.00	1.32
Maxima	0.35	79.56	1.45	54.00	1.12
Fricas	0.75	166.35	1.92	70.00	1.27
Sympy	0.00	0.00	0.00	0.00	0.00
Giac	0.46	91.52	1.23	65.00	1.23
Mupad	1.00	689.09	6.59	113.00	2.33

Table 1.5: Time and leaf size performance for each CAS


The following are bar charts for the normalized leafsize and time used columns from the above table.

Normalized mean size of antiderivative

Lower is better

**Mean time used (seconds)**

Lower is better



1.4 list of integrals that has no closed form antiderivative

{}

1.5 list of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of _int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call has completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out is not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica. Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative produced was correct.

Verification phase has 3 minutes time out. An integral whose result was not verified could still be correct. Further investigation is needed on those integrals which failed verifications. Such integrals are marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since these integrals are run in a batch mode, using an automated script, and by using `sagemath` (`SageMath` uses `Maxima`), then any integral where `Maxima` needs an interactive response from the user to answer a question during evaluation of the integral in order to complete the integration, will fail and is counted as failed.

The exception raised is `ValueError`. Therefore `Maxima` result below is lower than what could result if `Maxima` was run directly and each question `Maxima` asks was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate if the error is due to the interactive question being asked or not.

Maxima integrate was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath loading of Maxima abs_integrate was found to cause some problem. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS and Giac/X-CAS results

There are few integrals which failed due to SageMath not able to translate the result back to SageMath syntax and not because these CAS system were not able to do the integrations.

These will fail with error `Exception raised: NotImplementedError`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals.

Hopefully the next version of SageMath will have complete translation of FriCAS and Xcas syntax and I will re-run all the tests again when this happens.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi and Maple, the buildin system function `LeafSize` is used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special buildin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy and Giac antiderivatives is determined using the following function, thanks to user slelievre at <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/>

ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which is called directly from Python, the following code is used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative, Maple was used to determine the leaf size of Mupad output by post processing.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine,'cos(x)*sin(x)')
the_variable = evalin(symengine,'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



Chapter 2

detailed summary tables of results

2.1 List of integrals sorted by grade for each CAS

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

B grade: { 1, 8, 10 }

C grade: { }

F grade: { }

2.1.3 Maple

A grade: { 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 }

B grade: { 1, 8, 10, 22, 23 }

C grade: { }

F grade: { }

2.1.4 Maxima

A grade: { 2, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18 }

B grade: { 1, 3, 8, 10 }

C grade: { }

F grade: { 19, 20, 21, 22, 23 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 19, 20, 21 }

B grade: { 8, 10, 17, 18, 22, 23 }

C grade: { }

F grade: { }

2.1.6 Sympy

A grade: { }

B grade: { }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 6, 7, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23 }

B grade: { 8, 10, 22 }

C grade: { }

F grade: { }

2.1.8 MuPad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23 }

C grade: { }

F grade: { }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$

Problem 1	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	B	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	111	102	194	58	0	80	83
normalized size	1	1.00	2.02	1.85	3.53	1.05	0.00	1.45	1.51
time (sec)	N/A	0.103	0.164	0.485	0.419	1.405	0.000	0.615	0.479
Problem 2	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	50	55	58	71	0	52	134
normalized size	1	1.00	0.74	0.81	0.85	1.04	0.00	0.76	1.97
time (sec)	N/A	0.070	0.135	0.451	0.320	0.519	0.000	0.468	0.435
Problem 3	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	F	A	B	
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	62	64	94	38	0	49	51
normalized size	1	1.00	1.59	1.64	2.41	0.97	0.00	1.26	1.31
time (sec)	N/A	0.079	0.108	0.448	0.423	0.728	0.000	0.312	0.291

Problem 4	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	30	33	31	35	0	34	62
normalized size	1	1.00	0.75	0.82	0.78	0.88	0.00	0.85	1.55
time (sec)	N/A	0.049	0.054	0.411	0.322	0.534	0.000	0.524	0.339

Problem 5	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	9	9	9	19	9	9	0	9	25
normalized size	1	1.00	1.00	2.11	1.00	1.00	0.00	1.00	2.78
time (sec)	N/A	0.022	0.009	0.324	0.318	0.700	0.000	0.343	0.294

Problem 6	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	30	21	30	25	0	17	45
normalized size	1	1.00	2.00	1.40	2.00	1.67	0.00	1.13	3.00
time (sec)	N/A	0.040	0.042	0.377	0.426	0.490	0.000	0.277	0.282

Problem 7	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	11	16	18	19	0	14	36
normalized size	1	1.00	0.69	1.00	1.12	1.19	0.00	0.88	2.25
time (sec)	N/A	0.042	0.020	0.313	0.322	0.604	0.000	0.567	0.304

Problem 8	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	B	F	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	90	64	86	64	0	65	85
normalized size	1	1.00	2.90	2.06	2.77	2.06	0.00	2.10	2.74
time (sec)	N/A	0.060	0.046	0.398	0.419	0.481	0.000	0.676	0.298

Problem 9	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	26	35	29	45	0	29	80
normalized size	1	1.00	0.72	0.97	0.81	1.25	0.00	0.81	2.22
time (sec)	N/A	0.049	0.028	0.385	0.314	0.722	0.000	0.354	0.297

Problem 10	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	163	108	134	104	0	109	123
normalized size	1	1.00	3.33	2.20	2.73	2.12	0.00	2.22	2.51
time (sec)	N/A	0.081	0.052	0.378	0.418	0.581	0.000	0.333	0.333

Problem 11	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	39	55	42	70	0	41	113
normalized size	1	1.00	0.67	0.95	0.72	1.21	0.00	0.71	1.95
time (sec)	N/A	0.058	0.051	0.379	0.316	0.632	0.000	0.535	0.431

Problem 12	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	178	178	172	216	251	260	0	252	445
normalized size	1	1.00	0.97	1.21	1.41	1.46	0.00	1.42	2.50
time (sec)	N/A	0.272	0.787	0.426	0.332	1.423	0.000	0.415	2.300

Problem 13	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	115	117	120	144	0	139	172
normalized size	1	1.00	1.02	1.04	1.06	1.27	0.00	1.23	1.52
time (sec)	N/A	0.167	0.488	0.506	0.320	1.868	0.000	0.226	1.137

Problem 14	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	56	60	50	58	0	53	80
normalized size	1	1.00	0.85	0.91	0.76	0.88	0.00	0.80	1.21
time (sec)	N/A	0.091	0.061	0.536	0.317	1.249	0.000	0.473	0.442

Problem 15	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	11	21	11	11	0	12	55
normalized size	1	1.00	0.58	1.11	0.58	0.58	0.00	0.63	2.89
time (sec)	N/A	0.024	0.008	0.325	0.314	0.549	0.000	0.420	0.418

Problem 16	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	39	44	42	47	0	44	75
normalized size	1	1.00	1.03	1.16	1.11	1.24	0.00	1.16	1.97
time (sec)	N/A	0.063	0.049	0.381	0.311	0.477	0.000	0.653	0.459

Problem 17	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	F	A	B	
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	85	100	84	151	0	90	157
normalized size	1	1.00	1.18	1.39	1.17	2.10	0.00	1.25	2.18
time (sec)	N/A	0.083	0.135	0.320	0.311	0.511	0.000	0.388	0.567

Problem 18	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	F	A	B	
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	132	181	149	327	0	155	284
normalized size	1	1.00	1.08	1.48	1.22	2.68	0.00	1.27	2.33
time (sec)	N/A	0.123	0.282	0.332	0.316	0.559	0.000	0.408	0.870

Problem 19	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	205	194	210	0	483	0	216	4468
normalized size	1	1.29	1.22	1.32	0.00	3.04	0.00	1.36	28.10
time (sec)	N/A	0.324	1.072	0.421	0.000	0.576	0.000	0.587	7.377

Problem 20	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	112	87	106	0	296	0	103	2341
normalized size	1	1.33	1.04	1.26	0.00	3.52	0.00	1.23	27.87
time (sec)	N/A	0.187	0.267	0.515	0.000	0.471	0.000	0.364	2.258

Problem 21	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	71	105	0	206	0	80	697
normalized size	1	1.00	1.16	1.72	0.00	3.38	0.00	1.31	11.43
time (sec)	N/A	0.176	0.064	0.352	0.000	0.595	0.000	0.314	0.526

Problem 22	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-2)	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	158	206	0	453	0	163	2163
normalized size	1	1.00	1.74	2.26	0.00	4.98	0.00	1.79	23.77
time (sec)	N/A	0.289	0.413	0.329	0.000	0.632	0.000	0.380	1.190

Problem 23	Optimal	Rubi	Mathematica	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-2)	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	186	186	312	363	0	852	0	299	4075
normalized size	1	1.00	1.68	1.95	0.00	4.58	0.00	1.61	21.91
time (sec)	N/A	0.278	0.922	0.340	0.000	0.954	0.000	1.014	1.586

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [19] had the largest ratio of [.6923]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	5	3	1.00	13	0.231
2	A	3	2	1.00	13	0.154
3	A	4	3	1.00	13	0.231
4	A	3	2	1.00	11	0.182
5	A	2	2	1.00	11	0.182
6	A	3	2	1.00	13	0.154
7	A	3	2	1.00	13	0.154
8	A	4	3	1.00	13	0.231
9	A	3	2	1.00	13	0.154
10	A	5	3	1.00	13	0.231
11	A	3	2	1.00	13	0.154
12	A	3	2	1.00	13	0.154
13	A	3	2	1.00	13	0.154
14	A	3	2	1.00	11	0.182
15	A	4	4	1.00	11	0.364

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
16	A	3	2	1.00	13	0.154
17	A	3	2	1.00	13	0.154
18	A	3	2	1.00	13	0.154
19	A	16	9	1.29	13	0.692
20	A	10	9	1.33	13	0.692
21	A	8	8	1.00	13	0.615
22	A	7	7	1.00	13	0.538
23	A	16	9	1.00	13	0.692

Chapter 3

Listing of integrals

3.1 $\int \frac{\tan^4(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=55

$$\frac{x}{a} - \frac{\tan^5(x)(1 - \csc(x))}{5a} + \frac{\tan^3(x)(5 - 4 \csc(x))}{15a} - \frac{\tan(x)(15 - 8 \csc(x))}{15a}$$

[Out] $x/a - 1/15*(15-8*csc(x))*tan(x)/a + 1/15*(5-4*csc(x))*tan(x)^3/a - 1/5*(1-csc(x))*tan(x)^5/a$

Rubi [A] time = 0.10, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.231, Rules used = {3888, 3882, 8}

$$\frac{x}{a} - \frac{\tan^5(x)(1 - \csc(x))}{5a} + \frac{\tan^3(x)(5 - 4 \csc(x))}{15a} - \frac{\tan(x)(15 - 8 \csc(x))}{15a}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]^4/(a + a*Csc[x]), x]

[Out] $x/a - ((15 - 8*Csc[x])*Tan[x])/(15*a) + ((5 - 4*Csc[x])*Tan[x]^3)/(15*a) - ((1 - Csc[x])*Tan[x]^5)/(5*a)$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 3882

```
Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^(m_)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> -Simp[((e*Cot[c + d*x])^(m + 1)*(a + b*Csc[c + d*x]))/(d *e*(m + 1)), x] - Dist[1/(e^2*(m + 1)), Int[(e*Cot[c + d*x])^(m + 2)*(a*(m + 1) + b*(m + 2)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e}, x] && LtQ[m, -1]
```

Rule 3888

```
Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^(m_)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] :> Dist[a^(2*n)/e^(2*n), Int[(e*Cot[c + d*x])^(m + 2*n)/(-a + b*Csc[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[a^2 - b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\tan^4(x)}{a + a \csc(x)} dx &= \frac{\int (-a + a \csc(x)) \tan^6(x) dx}{a^2} \\ &= -\frac{(1 - \csc(x)) \tan^5(x)}{5a} + \frac{\int (5a - 4a \csc(x)) \tan^4(x) dx}{5a^2} \\ &= \frac{(5 - 4 \csc(x)) \tan^3(x)}{15a} - \frac{(1 - \csc(x)) \tan^5(x)}{5a} + \frac{\int (-15a + 8a \csc(x)) \tan^2(x) dx}{15a^2} \\ &= -\frac{(15 - 8 \csc(x)) \tan(x)}{15a} + \frac{(5 - 4 \csc(x)) \tan^3(x)}{15a} - \frac{(1 - \csc(x)) \tan^5(x)}{5a} + \frac{\int 15a dx}{15a^2} \\ &= \frac{x}{a} - \frac{(15 - 8 \csc(x)) \tan(x)}{15a} + \frac{(5 - 4 \csc(x)) \tan^3(x)}{15a} - \frac{(1 - \csc(x)) \tan^5(x)}{5a} \end{aligned}$$

Mathematica [B] time = 0.16, size = 111, normalized size = 2.02

$$\frac{-64 \sin(x) + 240x \sin(2x) - 178 \sin(2x) - 128 \sin(3x) + 120x \sin(4x) - 89 \sin(4x) + 6(120x - 89) \cos(x) + 128 \cos(2x) - 960a (\cos(\frac{x}{2}) - \sin(\frac{x}{2}))^3 (\sin(\frac{x}{2}) + \cos(\frac{x}{2}))^5}{960a (\cos(\frac{x}{2}) - \sin(\frac{x}{2}))^3 (\sin(\frac{x}{2}) + \cos(\frac{x}{2}))^5}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]^4/(a + a*Csc[x]), x]`

[Out] `(200 + 6*(-89 + 120*x)*Cos[x] + 128*Cos[2*x] - 178*Cos[3*x] + 240*x*Cos[3*x] + 184*Cos[4*x] - 64*Sin[x] - 178*Sin[2*x] + 240*x*Sin[2*x] - 128*Sin[3*x] - 89*Sin[4*x] + 120*x*Sin[4*x])/((960*a*(Cos[x/2] - Sin[x/2]))^3*(Cos[x/2] + Sin[x/2])^5)`

fricas [A] time = 1.41, size = 58, normalized size = 1.05

$$\frac{15x \cos(x)^3 + 23 \cos(x)^4 - 19 \cos(x)^2 + (15x \cos(x)^3 - 8 \cos(x)^2 + 1) \sin(x) + 4}{15(a \cos(x)^3 \sin(x) + a \cos(x)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^4/(a+a*csc(x)),x, algorithm="fricas")`

[Out] $\frac{1}{15}*(15*x*\cos(x)^3 + 23*\cos(x)^4 - 19*\cos(x)^2 + (15*x*\cos(x)^3 - 8*\cos(x)^2 + 1)*\sin(x) + 4)/(a*\cos(x)^3*\sin(x) + a*\cos(x)^3)$

giac [A] time = 0.62, size = 80, normalized size = 1.45

$$\frac{x}{a} + \frac{15 \tan\left(\frac{1}{2}x\right)^2 - 36 \tan\left(\frac{1}{2}x\right) + 17}{24 a \left(\tan\left(\frac{1}{2}x\right) - 1\right)^3} + \frac{55 \tan\left(\frac{1}{2}x\right)^4 + 260 \tan\left(\frac{1}{2}x\right)^3 + 450 \tan\left(\frac{1}{2}x\right)^2 + 300 \tan\left(\frac{1}{2}x\right) + 71}{40 a \left(\tan\left(\frac{1}{2}x\right) + 1\right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^4/(a+a*csc(x)),x, algorithm="giac")`

[Out] $x/a + \frac{1}{24}*(15*\tan(1/2*x)^2 - 36*\tan(1/2*x) + 17)/(a*(\tan(1/2*x) - 1)^3) + \frac{1}{40}*(55*\tan(1/2*x)^4 + 260*\tan(1/2*x)^3 + 450*\tan(1/2*x)^2 + 300*\tan(1/2*x) + 71)/(a*(\tan(1/2*x) + 1)^5)$

maple [B] time = 0.48, size = 102, normalized size = 1.85

$$-\frac{1}{6a \left(\tan\left(\frac{x}{2}\right) - 1\right)^3} - \frac{1}{4a \left(\tan\left(\frac{x}{2}\right) - 1\right)^2} + \frac{5}{8a \left(\tan\left(\frac{x}{2}\right) - 1\right)} + \frac{2}{5a \left(\tan\left(\frac{x}{2}\right) + 1\right)^5} - \frac{1}{a \left(\tan\left(\frac{x}{2}\right) + 1\right)^4} + \frac{1}{a \left(\tan\left(\frac{x}{2}\right) + 1\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^4/(a+a*csc(x)),x)`

[Out] $-\frac{1}{6}/a/(\tan(1/2*x)-1)^3 - \frac{1}{4}/a/(\tan(1/2*x)-1)^2 + \frac{5}{8}/a/(\tan(1/2*x)-1) + \frac{2}{5}/a/(\tan(1/2*x)+1)^5 - \frac{1}{a}/(\tan(1/2*x)+1)^4 + \frac{1}{a}/(\tan(1/2*x)+1)^2 + \frac{11}{8}/a/(\tan(1/2*x)+1) + 2/a*\arctan(\tan(1/2*x))$

maxima [B] time = 0.42, size = 194, normalized size = 3.53

$$\frac{2 \left(\frac{\sin(x)}{\cos(x)+1} - \frac{46 \sin(x)^2}{(\cos(x)+1)^2} - \frac{13 \sin(x)^3}{(\cos(x)+1)^3} + \frac{100 \sin(x)^4}{(\cos(x)+1)^4} + \frac{35 \sin(x)^5}{(\cos(x)+1)^5} - \frac{30 \sin(x)^6}{(\cos(x)+1)^6} - \frac{15 \sin(x)^7}{(\cos(x)+1)^7} + 8 \right)}{15 \left(a + \frac{2 a \sin(x)}{\cos(x)+1} - \frac{2 a \sin(x)^2}{(\cos(x)+1)^2} - \frac{6 a \sin(x)^3}{(\cos(x)+1)^3} + \frac{6 a \sin(x)^5}{(\cos(x)+1)^5} + \frac{2 a \sin(x)^6}{(\cos(x)+1)^6} - \frac{2 a \sin(x)^7}{(\cos(x)+1)^7} - \frac{a \sin(x)^8}{(\cos(x)+1)^8} \right)} + \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^4/(a+a*csc(x)),x, algorithm="maxima")`

[Out]
$$\frac{2}{15} \frac{\sin(x)}{\cos(x) + 1} - \frac{46 \sin(x)^2}{\cos(x) + 1} - \frac{13 \sin(x)^3}{\cos(x) + 1} + \frac{100 \sin(x)^4}{\cos(x) + 1} + \frac{35 \sin(x)^5}{\cos(x) + 1} - \frac{30 \sin(x)^6}{\cos(x) + 1} + \frac{15 \sin(x)^7}{\cos(x) + 1} - \frac{8}{(a + 2a \sin(x))(\cos(x) + 1)} - \frac{2a \sin(x)^2}{\cos(x) + 1} - \frac{6a \sin(x)^3}{\cos(x) + 1} + \frac{6a \sin(x)^5}{\cos(x) + 1} + \frac{2a \sin(x)^6}{\cos(x) + 1} - \frac{2a \sin(x)^7}{\cos(x) + 1} - \frac{a \sin(x)^8}{\cos(x) + 1} + \frac{2 \arctan(\sin(x))}{\cos(x) + 1})/a$$

mupad [B] time = 0.48, size = 83, normalized size = 1.51

$$\frac{x}{a} - \frac{-2 \tan\left(\frac{x}{2}\right)^7 - 4 \tan\left(\frac{x}{2}\right)^6 + \frac{14 \tan\left(\frac{x}{2}\right)^5}{3} + \frac{40 \tan\left(\frac{x}{2}\right)^4}{3} - \frac{26 \tan\left(\frac{x}{2}\right)^3}{15} - \frac{92 \tan\left(\frac{x}{2}\right)^2}{15} + \frac{2 \tan\left(\frac{x}{2}\right)}{15} + \frac{16}{15}}{a \left(\tan\left(\frac{x}{2}\right) - 1\right)^3 \left(\tan\left(\frac{x}{2}\right) + 1\right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^4/(a + a/sin(x)),x)`

[Out]
$$\frac{x}{a} - \frac{(2 \tan(x/2))/15 - (92 \tan(x/2)^2)/15 - (26 \tan(x/2)^3)/15 + (40 \tan(x/2)^4)/3 + (14 \tan(x/2)^5)/3 - 4 \tan(x/2)^6 - 2 \tan(x/2)^7 + 16/15}{(a * (\tan(x/2) - 1)^3 * (\tan(x/2) + 1)^5)}$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\tan^4(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)**4/(a+a*csc(x)),x)`

[Out] `Integral(tan(x)**4/(csc(x) + 1), x)/a`

3.2 $\int \frac{\tan^3(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=68

$$\frac{1}{8a(1 - \sin(x))} + \frac{3}{4a(\sin(x) + 1)} - \frac{1}{8a(\sin(x) + 1)^2} + \frac{5 \log(1 - \sin(x))}{16a} + \frac{11 \log(\sin(x) + 1)}{16a}$$

[Out] $5/16*\ln(1-\sin(x))/a+11/16*\ln(1+\sin(x))/a+1/8/a/(1-\sin(x))-1/8/a/(1+\sin(x))^2+3/4/a/(1+\sin(x))$

Rubi [A] time = 0.07, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3879, 88}

$$\frac{1}{8a(1 - \sin(x))} + \frac{3}{4a(\sin(x) + 1)} - \frac{1}{8a(\sin(x) + 1)^2} + \frac{5 \log(1 - \sin(x))}{16a} + \frac{11 \log(\sin(x) + 1)}{16a}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]^3/(a + a*Csc[x]), x]

[Out] $(5*\Log[1 - \Sin[x]])/(16*a) + (11*\Log[1 + \Sin[x]])/(16*a) + 1/(8*a*(1 - \Sin[x])) - 1/(8*a*(1 + \Sin[x])^2) + 3/(4*a*(1 + \Sin[x]))$

Rule 88

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))
```

Rule 3879

```
Int[cot[(c_.) + (d_.)*(x_)]^(m_.)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_), x_Symbol] :> Dist[1/(a^(m - n - 1)*b^n*d), Subst[Int[((a - b*x)^(m - 1)/2)*(a + b*x)^((m - 1)/2 + n))/x^(m + n), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[(m - 1)/2] && EqQ[a^2 - b^2, 0] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan^3(x)}{a + a \csc(x)} dx &= a^4 \text{Subst} \left(\int \frac{x^4}{(a - ax)^2(a + ax)^3} dx, x, \sin(x) \right) \\
&= a^4 \text{Subst} \left(\int \left(\frac{1}{8a^5(-1+x)^2} + \frac{5}{16a^5(-1+x)} + \frac{1}{4a^5(1+x)^3} - \frac{3}{4a^5(1+x)^2} + \frac{11}{16a^5(1+x)} \right) dx \right. \\
&\quad \left. = \frac{5 \log(1 - \sin(x))}{16a} + \frac{11 \log(1 + \sin(x))}{16a} + \frac{1}{8a(1 - \sin(x))} - \frac{1}{8a(1 + \sin(x))^2} + \frac{3}{4a(1 + \sin(x))} \right)
\end{aligned}$$

Mathematica [A] time = 0.14, size = 50, normalized size = 0.74

$$\frac{\frac{2(5 \sin^2(x) - 3 \sin(x) - 6)}{(\sin(x) - 1)(\sin(x) + 1)^2} + 5 \log(1 - \sin(x)) + 11 \log(\sin(x) + 1)}{16a}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]^3/(a + a*Csc[x]),x]`

[Out] $\frac{(5 \log[1 - \sin[x]] + 11 \log[1 + \sin[x]] + (2(-6 - 3 \sin[x] + 5 \sin[x]^2))/((-1 + \sin[x])*(1 + \sin[x])^2))/(16*a)}$

fricas [A] time = 0.52, size = 71, normalized size = 1.04

$$\frac{10 \cos(x)^2 + 11 (\cos(x)^2 \sin(x) + \cos(x)^2) \log(\sin(x) + 1) + 5 (\cos(x)^2 \sin(x) + \cos(x)^2) \log(-\sin(x) + 1) + 6 \sin(x) + 2}{16 (\cos(x)^2 \sin(x) + \cos(x)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^3/(a+a*csc(x)),x, algorithm="fricas")`

[Out] $\frac{1}{16}*(10*\cos(x)^2 + 11*(\cos(x)^2*\sin(x) + \cos(x)^2)*\log(\sin(x) + 1) + 5*(\cos(x)^2*\sin(x) + \cos(x)^2)*\log(-\sin(x) + 1) + 6*\sin(x) + 2)/(a*\cos(x)^2*\sin(x) + a*\cos(x)^2)$

giac [A] time = 0.47, size = 52, normalized size = 0.76

$$\frac{11 \log(\sin(x) + 1)}{16a} + \frac{5 \log(-\sin(x) + 1)}{16a} + \frac{5 \sin(x)^2 - 3 \sin(x) - 6}{8a(\sin(x) + 1)^2(\sin(x) - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^3/(a+a*csc(x)),x, algorithm="giac")`

[Out] $\frac{11}{16} \ln(\sin(x) + 1)/a + \frac{5}{16} \ln(-\sin(x) + 1)/a + \frac{1}{8} (5 \sin(x)^2 - 3 \sin(x) - 6)/(a (\sin(x) + 1)^2 (\sin(x) - 1))$

maple [A] time = 0.45, size = 55, normalized size = 0.81

$$-\frac{1}{8a(-1+\sin(x))} + \frac{5 \ln(-1+\sin(x))}{16a} - \frac{1}{8a(1+\sin(x))^2} + \frac{3}{4a(1+\sin(x))} + \frac{11 \ln(1+\sin(x))}{16a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \tan(x)^3/(a+a*\csc(x)), x)$

[Out] $-\frac{1}{8} \ln(-1+\sin(x))/a + \frac{5}{16} \ln(-1+\sin(x))/a - \frac{1}{8} \ln(1+\sin(x))^2/a + \frac{3}{4} \ln(1+\sin(x))/a + \frac{11}{16} \ln(1+\sin(x))/a$

maxima [A] time = 0.32, size = 58, normalized size = 0.85

$$\frac{5 \sin(x)^2 - 3 \sin(x) - 6}{8(a \sin(x)^3 + a \sin(x)^2 - a \sin(x) - a)} + \frac{11 \log(\sin(x) + 1)}{16a} + \frac{5 \log(\sin(x) - 1)}{16a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\tan(x)^3/(a+a*\csc(x)), x, \text{algorithm}=\text{"maxima"})$

[Out] $\frac{1}{8} (5 \sin(x)^2 - 3 \sin(x) - 6)/(a \sin(x)^3 + a \sin(x)^2 - a \sin(x) - a) + \frac{11}{16} \ln(\sin(x) + 1)/a + \frac{5}{16} \ln(\sin(x) - 1)/a$

mupad [B] time = 0.43, size = 134, normalized size = 1.97

$$\frac{5 \ln(\tan(\frac{x}{2}) - 1)}{8a} + \frac{11 \ln(\tan(\frac{x}{2}) + 1)}{8a} - \frac{\ln(\tan(\frac{x}{2})^2 + 1)}{a} + \frac{-\frac{3 \tan(\frac{x}{2})^5}{4} + \frac{\tan(\frac{x}{2})^4}{2} + \frac{9 \tan(\frac{x}{2})^3}{2} + \frac{\tan(\frac{x}{2})}{4}}{a \tan(\frac{x}{2})^6 + 2a \tan(\frac{x}{2})^5 - a \tan(\frac{x}{2})^4 - 4a \tan(\frac{x}{2})^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \tan(x)^3/(a + a/\sin(x)), x)$

[Out] $(5 \ln(\tan(x/2) - 1))/(8*a) + (11 \ln(\tan(x/2) + 1))/(8*a) - \ln(\tan(x/2)^2 + 1)/a + (\tan(x/2)^2/2 - (3 \tan(x/2))/4 + (9 \tan(x/2)^3)/2 + \tan(x/2)^4/2 - (3 \tan(x/2)^5)/4)/(a + 2*a \tan(x/2) - a \tan(x/2)^2 - 4*a \tan(x/2)^3 - a \tan(x/2)^4 + 2*a \tan(x/2)^5 + a \tan(x/2)^6)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\tan^3(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tan(x)**3/(a+a*csc(x)),x)  
[Out] Integral(tan(x)**3/(csc(x) + 1), x)/a
```

3.3 $\int \frac{\tan^2(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=39

$$-\frac{x}{a} - \frac{\tan^3(x)(1 - \csc(x))}{3a} + \frac{\tan(x)(3 - 2 \csc(x))}{3a}$$

[Out] $-\frac{x}{a} + \frac{(3 - 2 \csc(x)) \tan(x)}{3a} - \frac{(1 - \csc(x)) \tan^3(x)}{3a}$

Rubi [A] time = 0.08, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.231, Rules used = {3888, 3882, 8}

$$-\frac{x}{a} - \frac{\tan^3(x)(1 - \csc(x))}{3a} + \frac{\tan(x)(3 - 2 \csc(x))}{3a}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]^2/(a + a*Csc[x]), x]

[Out] $-\frac{x}{a} + \frac{(3 - 2 \csc(x)) \tan(x)}{3a} - \frac{(1 - \csc(x)) \tan^3(x)}{3a}$

Rule 8

Int[a_, x_Symbol] :> Simplify[a*x, x] /; FreeQ[a, x]

Rule 3882

```
Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^m_*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)), x_Symbol] :> -Simplify[((e*Cot[c + d*x])^(m + 1)*(a + b*Csc[c + d*x]))/(d*e*(m + 1)), x] - Dist[1/(e^(2*(m + 1))), Int[(e*Cot[c + d*x])^(m + 2)*(a*(m + 1) + b*(m + 2)*Csc[c + d*x]), x], x] /; FreeQ[{a, b, c, d, e}, x] && LtQ[m, -1]
```

Rule 3888

```
Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^m_*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.))^(n_), x_Symbol] :> Dist[a^(2*n)/e^(2*n), Int[(e*Cot[c + d*x])^(m + 2*n)/(-a + b*Csc[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[a^2 - b^2, 0] && ILtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan^2(x)}{a + a \csc(x)} dx &= \frac{\int (-a + a \csc(x)) \tan^4(x) dx}{a^2} \\
&= -\frac{(1 - \csc(x)) \tan^3(x)}{3a} + \frac{\int (3a - 2a \csc(x)) \tan^2(x) dx}{3a^2} \\
&= \frac{(3 - 2 \csc(x)) \tan(x)}{3a} - \frac{(1 - \csc(x)) \tan^3(x)}{3a} + \frac{\int -3a dx}{3a^2} \\
&= -\frac{x}{a} + \frac{(3 - 2 \csc(x)) \tan(x)}{3a} - \frac{(1 - \csc(x)) \tan^3(x)}{3a}
\end{aligned}$$

Mathematica [A] time = 0.11, size = 62, normalized size = 1.59

$$-\frac{-2 \sin(x) + 4 \cos(2x) + (6x - 5)(\sin(x) + 1) \cos(x)}{6a \left(\cos\left(\frac{x}{2}\right) - \sin\left(\frac{x}{2}\right)\right) \left(\sin\left(\frac{x}{2}\right) + \cos\left(\frac{x}{2}\right)\right)^3}$$

Antiderivative was successfully verified.

[In] Integrate[Tan[x]^2/(a + a*Csc[x]), x]

[Out] $\frac{-1/6*(4*\cos(2*x) - 2*\sin(x) + (-5 + 6*x)*\cos(x)*(1 + \sin(x)))/(a*(\cos(x/2) - \sin(x/2))*(\cos(x/2) + \sin(x/2))^3)}$

fricas [A] time = 0.73, size = 38, normalized size = 0.97

$$-\frac{3x \cos(x) + 4 \cos(x)^2 + (3x \cos(x) - 1) \sin(x) - 2}{3(a \cos(x) \sin(x) + a \cos(x))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tan(x)^2/(a+a*csc(x)), x, algorithm="fricas")

[Out] $\frac{-1/3*(3*x*\cos(x) + 4*\cos(x)^2 + (3*x*\cos(x) - 1)*\sin(x) - 2)/(a*\cos(x)*\sin(x) + a*\cos(x))}$

giac [A] time = 0.31, size = 49, normalized size = 1.26

$$-\frac{x}{a} - \frac{1}{2a \left(\tan\left(\frac{1}{2}x\right) - 1\right)} - \frac{9 \tan\left(\frac{1}{2}x\right)^2 + 24 \tan\left(\frac{1}{2}x\right) + 11}{6a \left(\tan\left(\frac{1}{2}x\right) + 1\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^2/(a+a*csc(x)),x, algorithm="giac")`

[Out]
$$-\frac{x}{a} - \frac{1}{2} \cdot \frac{1}{a \cdot (\tan(1/2 \cdot x) - 1)} - \frac{1}{6} \cdot \frac{9 \cdot \tan(1/2 \cdot x)^2 + 24 \cdot \tan(1/2 \cdot x) + 11}{a \cdot (\tan(1/2 \cdot x) + 1)^3}$$

maple [A] time = 0.45, size = 64, normalized size = 1.64

$$-\frac{1}{2a \left(\tan\left(\frac{x}{2}\right)-1\right)} + \frac{2}{3a \left(\tan\left(\frac{x}{2}\right)+1\right)^3} - \frac{1}{a \left(\tan\left(\frac{x}{2}\right)+1\right)^2} - \frac{3}{2a \left(\tan\left(\frac{x}{2}\right)+1\right)} - \frac{2 \arctan\left(\tan\left(\frac{x}{2}\right)\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^2/(a+a*csc(x)),x)`

[Out]
$$-\frac{1}{2} \cdot \frac{a}{\tan(1/2 \cdot x) - 1} + \frac{2}{3} \cdot \frac{a}{(\tan(1/2 \cdot x) + 1)^3} - \frac{1}{a} \cdot \frac{1}{(\tan(1/2 \cdot x) + 1)^2} - \frac{3}{2} \cdot \frac{a}{(\tan(1/2 \cdot x) + 1)} - 2 \cdot \frac{a \cdot \arctan(\tan(1/2 \cdot x))}{a}$$

maxima [B] time = 0.42, size = 94, normalized size = 2.41

$$-\frac{2 \left(\frac{\sin(x)}{\cos(x)+1} - \frac{6 \sin(x)^2}{(\cos(x)+1)^2} - \frac{3 \sin(x)^3}{(\cos(x)+1)^3} + 2\right)}{3 \left(a + \frac{2 a \sin(x)}{\cos(x)+1} - \frac{2 a \sin(x)^3}{(\cos(x)+1)^3} - \frac{a \sin(x)^4}{(\cos(x)+1)^4}\right)} - \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^2/(a+a*csc(x)),x, algorithm="maxima")`

[Out]
$$-\frac{2}{3} \cdot \frac{\sin(x)}{\cos(x) + 1} - \frac{6 \sin(x)^2}{(\cos(x) + 1)^2} - \frac{3 \sin(x)^3}{(\cos(x) + 1)^3} + \frac{2}{a + 2 \cdot \frac{\sin(x)}{\cos(x) + 1} - 2 \cdot \frac{\sin(x)^3}{(\cos(x) + 1)^3} - \frac{\sin(x)^4}{(\cos(x) + 1)^4}} - 2 \cdot \frac{\arctan(\sin(x)/(\cos(x) + 1))}{a}$$

mupad [B] time = 0.29, size = 51, normalized size = 1.31

$$\frac{-2 \tan\left(\frac{x}{2}\right)^3 - 4 \tan\left(\frac{x}{2}\right)^2 + \frac{2 \tan\left(\frac{x}{2}\right)}{3} + \frac{4}{3}}{a \left(\tan\left(\frac{x}{2}\right)-1\right) \left(\tan\left(\frac{x}{2}\right)+1\right)^3} - \frac{x}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^2/(a + a/sin(x)),x)`

[Out]
$$-\frac{((2 \cdot \tan(x/2))/3 - 4 \cdot \tan(x/2)^2 - 2 \cdot \tan(x/2)^3 + 4/3)/(a \cdot (\tan(x/2) - 1) \cdot (\tan(x/2) + 1)^3) - x/a}{a}$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\tan^2(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tan(x)**2/(a+a*csc(x)),x)  
[Out] Integral(tan(x)**2/(csc(x) + 1), x)/a
```

3.4 $\int \frac{\tan(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=40

$$-\frac{1}{2a(\sin(x) + 1)} - \frac{\log(1 - \sin(x))}{4a} - \frac{3 \log(\sin(x) + 1)}{4a}$$

[Out] $-1/4*\ln(1-\sin(x))/a-3/4*\ln(1+\sin(x))/a-1/2/a/(1+\sin(x))$

Rubi [A] time = 0.05, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.182, Rules used = {3879, 88}

$$-\frac{1}{2a(\sin(x) + 1)} - \frac{\log(1 - \sin(x))}{4a} - \frac{3 \log(\sin(x) + 1)}{4a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\tan[x]/(a + a*\csc[x]), x]$

[Out] $-\text{Log}[1 - \sin[x]]/(4*a) - (3*\text{Log}[1 + \sin[x]])/(4*a) - 1/(2*a*(1 + \sin[x]))$

Rule 88

```
Int[((a_.) + (b_.*(x_))^^(m_.*((c_.) + (d_.*(x_))^^(n_.*((e_.) + (f_.*(x_))^^(p_.), x_Symbol) :> Int[ExpandIntegrand[(a + b*x)^^m*(c + d*x)^^n*(e + f*x)^^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))
```

Rule 3879

```
Int[cot[(c_.) + (d_.*(x_))^^(m_.*((c_.) + (d_.*(x_))))*(b_.) + (a_.))^^(n_.), x_Symbol] :> Dist[1/(a^(m - n - 1)*b^n*d), Subst[Int[((a - b*x)^((m - 1)/2)*(a + b*x)^((m - 1)/2 + n))/x^(m + n), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[(m - 1)/2] && EqQ[a^2 - b^2, 0] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan(x)}{a + a \csc(x)} dx &= a^2 \operatorname{Subst} \left(\int \frac{x^2}{(a - ax)(a + ax)^2} dx, x, \sin(x) \right) \\
&= a^2 \operatorname{Subst} \left(\int \left(-\frac{1}{4a^3(-1+x)} + \frac{1}{2a^3(1+x)^2} - \frac{3}{4a^3(1+x)} \right) dx, x, \sin(x) \right) \\
&= -\frac{\log(1 - \sin(x))}{4a} - \frac{3 \log(1 + \sin(x))}{4a} - \frac{1}{2a(1 + \sin(x))}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 30, normalized size = 0.75

$$-\frac{\frac{2}{\sin(x)+1} + \log(1 - \sin(x)) + 3 \log(\sin(x) + 1)}{4a}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]/(a + a*Csc[x]),x]`

[Out] $-1/4*(\text{Log}[1 - \text{Sin}[x]] + 3*\text{Log}[1 + \text{Sin}[x]] + 2/(1 + \text{Sin}[x]))/a$

fricas [A] time = 0.53, size = 35, normalized size = 0.88

$$-\frac{3 (\sin(x) + 1) \log(\sin(x) + 1) + (\sin(x) + 1) \log(-\sin(x) + 1) + 2}{4 (a \sin(x) + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+a*csc(x)),x, algorithm="fricas")`

[Out] $-1/4*(3*(\sin(x) + 1)*\log(\sin(x) + 1) + (\sin(x) + 1)*\log(-\sin(x) + 1) + 2)/(a*\sin(x) + a)$

giac [A] time = 0.52, size = 34, normalized size = 0.85

$$-\frac{3 \log(\sin(x) + 1)}{4 a} - \frac{\log(-\sin(x) + 1)}{4 a} - \frac{1}{2 a (\sin(x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+a*csc(x)),x, algorithm="giac")`

[Out] $-3/4*\log(\sin(x) + 1)/a - 1/4*\log(-\sin(x) + 1)/a - 1/2/(a*(\sin(x) + 1))$

maple [A] time = 0.41, size = 33, normalized size = 0.82

$$-\frac{\ln(-1 + \sin(x))}{4a} - \frac{1}{2a(1 + \sin(x))} - \frac{3 \ln(1 + \sin(x))}{4a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)/(a+a*csc(x)),x)`

[Out] $-1/4/a*\ln(-1+\sin(x))-1/2/a/(1+\sin(x))-3/4*\ln(1+\sin(x))/a$

maxima [A] time = 0.32, size = 31, normalized size = 0.78

$$-\frac{3 \log (\sin (x)+1)}{4 a}-\frac{\log (\sin (x)-1)}{4 a}-\frac{1}{2 (a \sin (x)+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $-3/4*\log(\sin(x) + 1)/a - 1/4*\log(\sin(x) - 1)/a - 1/2/(a*\sin(x) + a)$

mupad [B] time = 0.34, size = 62, normalized size = 1.55

$$\frac{\tan \left(\frac{x}{2}\right)}{a \tan \left(\frac{x}{2}\right)^2+2 a \tan \left(\frac{x}{2}\right)+a}-\frac{3 \ln \left(\tan \left(\frac{x}{2}\right)+1\right)}{2 a}-\frac{\ln \left(\tan \left(\frac{x}{2}\right)-1\right)}{2 a}+\frac{\ln \left(\tan \left(\frac{x}{2}\right)^2+1\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)/(a + a/sin(x)),x)`

[Out] $\tan(x/2)/(a + 2*a*tan(x/2) + a*tan(x/2)^2) - (3*log(tan(x/2) + 1))/(2*a) - \log(tan(x/2) - 1)/(2*a) + \log(tan(x/2)^2 + 1)/a$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\tan (x)}{\csc (x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+a*csc(x)),x)`

[Out] `Integral(tan(x)/(csc(x) + 1), x)/a`

3.5 $\int \frac{\cot(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=9

$$\frac{\log(\sin(x) + 1)}{a}$$

[Out] $\ln(1+\sin(x))/a$

Rubi [A] time = 0.02, antiderivative size = 9, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {3879, 31}

$$\frac{\log(\sin(x) + 1)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[x]/(a + a*\text{Csc}[x]), x]$

[Out] $\text{Log}[1 + \text{Sin}[x]]/a$

Rule 31

$\text{Int}[(a_1 + b_1*x_1)^{-1}, x_{\text{Symbol}}] \Rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]/b, x] /; \text{FreeQ}[\{a, b\}, x]$

Rule 3879

$\text{Int}[\cot[(c_1 + d_1*x_1)^{(m_1 - n_1)*d_1}]*\csc[(c_1 + d_1*x_1)*(b_1 + a_1)^{(n_1 - 1)/2}*(a + b*x)^{((m - 1)/2 + n_1)}/x^{(m + n_1)}, x], \text{Sin}[c + d*x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{IntegerQ}[n]$

Rubi steps

$$\begin{aligned} \int \frac{\cot(x)}{a + a \csc(x)} dx &= \text{Subst}\left(\int \frac{1}{a + ax} dx, x, \sin(x)\right) \\ &= \frac{\log(1 + \sin(x))}{a} \end{aligned}$$

Mathematica [A] time = 0.01, size = 9, normalized size = 1.00

$$\frac{\log(\sin(x) + 1)}{a}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cot[x]/(a + a*Csc[x]),x]
[Out] Log[1 + Sin[x]]/a
fricas [A] time = 0.70, size = 9, normalized size = 1.00
```

$$\frac{\log(\sin(x) + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cot(x)/(a+a*csc(x)),x, algorithm="fricas")
[Out] log(sin(x) + 1)/a
giac [A] time = 0.34, size = 9, normalized size = 1.00
```

$$\frac{\log(\sin(x) + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cot(x)/(a+a*csc(x)),x, algorithm="giac")
[Out] log(sin(x) + 1)/a
maple [A] time = 0.32, size = 19, normalized size = 2.11
```

$$-\frac{\ln(\csc(x))}{a} + \frac{\ln(1 + \csc(x))}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cot(x)/(a+a*csc(x)),x)
[Out] -1/a*ln(csc(x))+1/a*ln(1+csc(x))
maxima [A] time = 0.32, size = 9, normalized size = 1.00
```

$$\frac{\log(\sin(x) + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cot(x)/(a+a*csc(x)),x, algorithm="maxima")
[Out] log(sin(x) + 1)/a
```

mupad [B] time = 0.29, size = 25, normalized size = 2.78

$$\frac{2 \ln \left(\tan \left(\frac{x}{2}\right)+1\right)-\ln \left(\tan \left(\frac{x}{2}\right)^2+1\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)/(a + a/sin(x)),x)`

[Out] `(2*log(tan(x/2) + 1) - log(tan(x/2)^2 + 1))/a`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)/(a+a*csc(x)),x)`

[Out] `Integral(cot(x)/(csc(x) + 1), x)/a`

3.6 $\int \frac{\cot^2(x)}{a + a \csc(x)} dx$

Optimal. Leaf size=15

$$-\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{a}$$

[Out] $-\frac{x}{a} - \text{arctanh}(\cos(x))/a$

Rubi [A] time = 0.04, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3888, 3770}

$$-\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{a}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^2/(a + a*Csc[x]), x]

[Out] $-(x/a) - \text{ArcTanh}[\cos(x)]/a$

Rule 3770

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rule 3888

Int[(cot[(c_.) + (d_.)*(x_)]*(e_.))^m_]*csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_.)^n_, x_Symbol] :> Dist[a^(2*n)/e^(2*n), Int[(e*Cot[c + d*x])^(m + 2*n)/(-a + b*Csc[c + d*x])^n, x], x] /; FreeQ[{a, b, c, d, e, m}, x] && EqQ[a^2 - b^2, 0] && ILtQ[n, 0]

Rubi steps

$$\begin{aligned} \int \frac{\cot^2(x)}{a + a \csc(x)} dx &= \frac{\int (-a + a \csc(x)) dx}{a^2} \\ &= -\frac{x}{a} + \frac{\int \csc(x) dx}{a} \\ &= -\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{a} \end{aligned}$$

Mathematica [A] time = 0.04, size = 30, normalized size = 2.00

$$-\frac{x}{a} + \frac{\log(\sin(\frac{x}{2}))}{a} - \frac{\log(\cos(\frac{x}{2}))}{a}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^2/(a + a*Csc[x]),x]`

[Out] $-(x/a) - \text{Log}[\text{Cos}[x/2]]/a + \text{Log}[\text{Sin}[x/2]]/a$

fricas [A] time = 0.49, size = 25, normalized size = 1.67

$$-\frac{2x + \log\left(\frac{1}{2}\cos(x) + \frac{1}{2}\right) - \log\left(-\frac{1}{2}\cos(x) + \frac{1}{2}\right)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^2/(a+a*csc(x)),x, algorithm="fricas")`

[Out] $-1/2*(2*x + \log(1/2*\cos(x) + 1/2) - \log(-1/2*\cos(x) + 1/2))/a$

giac [A] time = 0.28, size = 17, normalized size = 1.13

$$-\frac{x}{a} + \frac{\log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^2/(a+a*csc(x)),x, algorithm="giac")`

[Out] $-x/a + \log(\text{abs}(\tan(1/2*x)))/a$

maple [A] time = 0.38, size = 21, normalized size = 1.40

$$\frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{a} - \frac{2\arctan\left(\tan\left(\frac{x}{2}\right)\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^2/(a+a*csc(x)),x)`

[Out] $1/a*\ln(\tan(1/2*x))-2/a*\arctan(\tan(1/2*x))$

maxima [A] time = 0.43, size = 30, normalized size = 2.00

$$-\frac{2\arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} + \frac{\log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^2/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $-2\arctan(\sin(x)/(\cos(x) + 1))/a + \log(\sin(x)/(\cos(x) + 1))/a$

mupad [B] time = 0.28, size = 45, normalized size = 3.00

$$\frac{2 \operatorname{atan}\left(\frac{4}{4 \tan\left(\frac{x}{2}\right)+4}-\frac{4 \tan\left(\frac{x}{2}\right)}{4 \tan\left(\frac{x}{2}\right)+4}\right)}{a}+\frac{\ln \left(\tan \left(\frac{x}{2}\right)\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^2/(a + a/sin(x)),x)`

[Out] $(2 \operatorname{atan}(4/(4 \tan(x/2) + 4) - (4 \tan(x/2))/(4 \tan(x/2) + 4)))/a + \log(\tan(x/2))/a$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot^2(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)**2/(a+a*csc(x)),x)`

[Out] `Integral(cot(x)**2/(csc(x) + 1), x)/a`

3.7 $\int \frac{\cot^3(x)}{a + a \csc(x)} dx$

Optimal. Leaf size=16

$$-\frac{\csc(x)}{a} - \frac{\log(\sin(x))}{a}$$

[Out] $-\csc(x)/a - \ln(\sin(x))/a$

Rubi [A] time = 0.04, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3879, 43}

$$-\frac{\csc(x)}{a} - \frac{\log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^3/(a + a*Csc[x]), x]

[Out] $-(\Csc[x]/a) - \Log[\Sin[x]]/a$

Rule 43

```
Int[((a_) + (b_)*(x_))^m_*((c_) + (d_)*(x_))^n_, x_Symbol] :> Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 3879

```
Int[cot[(c_) + (d_)*(x_)]^m_*csc[(c_) + (d_)*(x_)]*(b_)^(n_), x_Symbol] :> Dist[1/(a^(m - n - 1)*b^n*d), Subst[Int[((a - b*x)^(m - 1)/2)*(a + b*x)^(n)/x^(m + n), x], x, Sin[c + d*x]], x] /; Fre
eQ[{a, b, c, d}, x] && IntegerQ[(m - 1)/2] && EqQ[a^2 - b^2, 0] && IntegerQ
[n]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cot^3(x)}{a + a \csc(x)} dx &= \frac{\text{Subst}\left(\int \frac{a - ax}{x^2} dx, x, \sin(x)\right)}{a^2} \\
&= \frac{\text{Subst}\left(\int \left(\frac{a}{x^2} - \frac{a}{x}\right) dx, x, \sin(x)\right)}{a^2} \\
&= -\frac{\csc(x)}{a} - \frac{\log(\sin(x))}{a}
\end{aligned}$$

Mathematica [A] time = 0.02, size = 11, normalized size = 0.69

$$-\frac{\csc(x) + \log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[x]^3/(a + a*Csc[x]), x]

[Out] $-(\csc(x) + \log(\sin(x)))/a$

fricas [A] time = 0.60, size = 19, normalized size = 1.19

$$-\frac{\log\left(\frac{1}{2} \sin(x)\right) \sin(x) + 1}{a \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^3/(a+a*csc(x)), x, algorithm="fricas")

[Out] $-(\log(1/2*\sin(x))*\sin(x) + 1)/(a*\sin(x))$

giac [A] time = 0.57, size = 14, normalized size = 0.88

$$-\frac{\frac{1}{\sin(x)} + \log(|\sin(x)|)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^3/(a+a*csc(x)), x, algorithm="giac")

[Out] $-(1/\sin(x) + \log(\sin(x)))/a$

maple [A] time = 0.31, size = 16, normalized size = 1.00

$$-\frac{\csc(x)}{a} + \frac{\ln(\csc(x))}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \cot(x)^3 / (a + a * \csc(x)) dx$

[Out] $-\csc(x)/a + 1/a * \ln(\csc(x))$

maxima [A] time = 0.32, size = 18, normalized size = 1.12

$$-\frac{\log(\sin(x))}{a} - \frac{1}{a \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\cot(x)^3 / (a + a * \csc(x)), x, \text{algorithm}=\text{"maxima"})$

[Out] $-\log(\sin(x))/a - 1/(a * \sin(x))$

mupad [B] time = 0.30, size = 36, normalized size = 2.25

$$-\frac{\frac{\tan(\frac{x}{2})}{2} - \ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + \ln\left(\tan\left(\frac{x}{2}\right)\right) + \frac{1}{2\tan(\frac{x}{2})}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \cot(x)^3 / (a + a/\sin(x)) dx$

[Out] $-(\tan(x/2)/2 - \log(\tan(x/2)^2 + 1) + \log(\tan(x/2)) + 1/(2*\tan(x/2)))/a$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot^3(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\cot(x)^{**3} / (a + a * \csc(x)), x)$

[Out] $\text{Integral}(\cot(x)^{**3} / (\csc(x) + 1), x)/a$

$$3.8 \quad \int \frac{\cot^4(x)}{a+a \csc(x)} dx$$

Optimal. Leaf size=31

$$\frac{x}{a} + \frac{\tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x)(2 - \csc(x))}{2a}$$

[Out] $x/a + 1/2*\text{arctanh}(\cos(x))/a + 1/2*\cot(x)*(2 - \csc(x))/a$

Rubi [A] time = 0.06, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.231, Rules used = {3888, 3881, 3770}

$$\frac{x}{a} + \frac{\tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x)(2 - \csc(x))}{2a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\cot[x]^4/(a + a*\csc[x]), x]$

[Out] $x/a + \text{ArcTanh}[\cos[x]]/(2*a) + (\cot[x]*(2 - \csc[x]))/(2*a)$

Rule 3770

$\text{Int}[\csc[(c_.) + (d_.)*(x_.)], x_{\text{Symbol}}] \rightarrow -\text{Simp}[\text{ArcTanh}[\cos[c + d*x]]/d, x]$
 $/; \text{FreeQ}[\{c, d\}, x]$

Rule 3881

$\text{Int}[(\cot[(c_.) + (d_.)*(x_.)]*(e_.)^{(m_.)})*(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_)), x_{\text{Symbol}}] \rightarrow -\text{Simp}[(e*(e*\cot[c + d*x])^{(m - 1)}*(a*m + b*(m - 1)*\csc[c + d*x]))/(d*m*(m - 1)), x] - \text{Dist}[e^{2/m}, \text{Int}[(e*\cot[c + d*x])^{(m - 2)}*(a*m + b*(m - 1)*\csc[c + d*x]), x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{GtQ}[m, 1]$

Rule 3888

$\text{Int}[(\cot[(c_.) + (d_.)*(x_.)]*(e_.)^{(m_.)})*(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.)^{(n_.)}), x_{\text{Symbol}}] \rightarrow \text{Dist}[a^{(2*n)}/e^{(2*n)}, \text{Int}[(e*\cot[c + d*x])^{(m + 2*n)}/(-a + b*\csc[c + d*x])^n, x], x] /; \text{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \text{EqQ}[a^2 - b^2, 0] \&& \text{ILtQ}[n, 0]$

Rubi steps

$$\begin{aligned}
\int \frac{\cot^4(x)}{a + a \csc(x)} dx &= \frac{\int \cot^2(x)(-a + a \csc(x)) dx}{a^2} \\
&= \frac{\cot(x)(2 - \csc(x))}{2a} - \frac{\int (-2a + a \csc(x)) dx}{2a^2} \\
&= \frac{x}{a} + \frac{\cot(x)(2 - \csc(x))}{2a} - \frac{\int \csc(x) dx}{2a} \\
&= \frac{x}{a} + \frac{\tanh^{-1}(\cos(x))}{2a} + \frac{\cot(x)(2 - \csc(x))}{2a}
\end{aligned}$$

Mathematica [B] time = 0.05, size = 90, normalized size = 2.90

$$\frac{x}{a} - \frac{\tan\left(\frac{x}{2}\right)}{2a} + \frac{\cot\left(\frac{x}{2}\right)}{2a} - \frac{\csc^2\left(\frac{x}{2}\right)}{8a} + \frac{\sec^2\left(\frac{x}{2}\right)}{8a} - \frac{\log\left(\sin\left(\frac{x}{2}\right)\right)}{2a} + \frac{\log\left(\cos\left(\frac{x}{2}\right)\right)}{2a}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[x]^4/(a + a*Csc[x]),x]

[Out] $x/a + \text{Cot}[x/2]/(2*a) - \text{Csc}[x/2]^2/(8*a) + \text{Log}[\text{Cos}[x/2]]/(2*a) - \text{Log}[\text{Sin}[x/2]]/(2*a) + \text{Sec}[x/2]^2/(8*a) - \text{Tan}[x/2]/(2*a)$

fricas [B] time = 0.48, size = 64, normalized size = 2.06

$$\frac{4 x \cos(x)^2 + (\cos(x)^2 - 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - (\cos(x)^2 - 1) \log\left(-\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 4 \cos(x) \sin(x) - 4 x + 2 \cos(x)}{4(a \cos(x)^2 - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^4/(a+a*csc(x)),x, algorithm="fricas")

[Out] $1/4*(4*x*cos(x)^2 + (cos(x)^2 - 1)*log(1/2*cos(x) + 1/2) - (cos(x)^2 - 1)*log(-1/2*cos(x) + 1/2) - 4*cos(x)*sin(x) - 4*x + 2*cos(x))/(a*cos(x)^2 - a)$

giac [B] time = 0.68, size = 65, normalized size = 2.10

$$\frac{x}{a} - \frac{\log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{2a} + \frac{a \tan\left(\frac{1}{2}x\right)^2 - 4a \tan\left(\frac{1}{2}x\right)}{8a^2} + \frac{6 \tan\left(\frac{1}{2}x\right)^2 + 4 \tan\left(\frac{1}{2}x\right) - 1}{8a \tan\left(\frac{1}{2}x\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^4/(a+a*csc(x)),x, algorithm="giac")`

[Out] $x/a - \frac{1}{2} \log(\tan(1/2*x))/a + \frac{1}{8} (a \tan(1/2*x)^2 - 4*a \tan(1/2*x))/a^2 + \frac{1}{8} (6 \tan(1/2*x)^2 + 4 \tan(1/2*x) - 1)/(a \tan(1/2*x)^2)$

maple [B] time = 0.40, size = 64, normalized size = 2.06

$$\frac{\tan^2\left(\frac{x}{2}\right)}{8a} - \frac{\tan\left(\frac{x}{2}\right)}{2a} - \frac{1}{8a \tan\left(\frac{x}{2}\right)^2} + \frac{1}{2a \tan\left(\frac{x}{2}\right)} - \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{2a} + \frac{2 \arctan\left(\tan\left(\frac{x}{2}\right)\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^4/(a+a*csc(x)),x)`

[Out] $\frac{1}{8} \frac{a \tan(1/2*x)^2 - 1/2 \tan(1/2*x) - 1/8 \ln(\tan(1/2*x)) + 1/2 \arctan(\tan(1/2*x))}{a \tan(1/2*x)^2}$

maxima [B] time = 0.42, size = 86, normalized size = 2.77

$$-\frac{\frac{4 \sin(x)}{\cos(x)+1} - \frac{\sin(x)^2}{(\cos(x)+1)^2}}{8a} + \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} - \frac{\log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{2a} + \frac{\left(\frac{4 \sin(x)}{\cos(x)+1} - 1\right)(\cos(x) + 1)^2}{8a \sin(x)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^4/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $\frac{-1/8*(4*\sin(x)/(\cos(x) + 1) - \sin(x)^2/(\cos(x) + 1)^2)/a + 2*\arctan(\sin(x)/(\cos(x) + 1))/a - 1/2*\log(\sin(x)/(\cos(x) + 1))/a + 1/8*(4*\sin(x)/(\cos(x) + 1) - 1)*(\cos(x) + 1)^2/(a*\sin(x)^2)}$

mupad [B] time = 0.30, size = 85, normalized size = 2.74

$$\frac{\tan\left(\frac{x}{2}\right)^2}{8a} - \frac{2 \operatorname{atan}\left(\frac{4}{4 \tan\left(\frac{x}{2}\right)+2} - \frac{2 \tan\left(\frac{x}{2}\right)}{4 \tan\left(\frac{x}{2}\right)+2}\right)}{a} - \frac{\tan\left(\frac{x}{2}\right)}{2a} - \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{2a} + \frac{2 \tan\left(\frac{x}{2}\right) - \frac{1}{2}}{4a \tan\left(\frac{x}{2}\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^4/(a + a/sin(x)),x)`

[Out] $\frac{\tan(x/2)^2/(8*a) - (2*\operatorname{atan}(4/(4*\tan(x/2) + 2) - (2*\tan(x/2))/(4*\tan(x/2) + 2))/a - \tan(x/2)/(2*a) - \log(\tan(x/2))/(2*a) + (2*\tan(x/2) - 1/2)/(4*a*\tan(x/2)^2)}$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot^4(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)**4/(a+a*csc(x)),x)
[Out] Integral(cot(x)**4/(csc(x) + 1), x)/a

3.9 $\int \frac{\cot^5(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=36

$$-\frac{\csc^3(x)}{3a} + \frac{\csc^2(x)}{2a} + \frac{\csc(x)}{a} + \frac{\log(\sin(x))}{a}$$

[Out] $\csc(x)/a + 1/2*\csc(x)^2/a - 1/3*\csc(x)^3/a + \ln(\sin(x))/a$

Rubi [A] time = 0.05, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3879, 75}

$$-\frac{\csc^3(x)}{3a} + \frac{\csc^2(x)}{2a} + \frac{\csc(x)}{a} + \frac{\log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[x]^5/(a + a*\text{Csc}[x]), x]$

[Out] $\text{Csc}[x]/a + \text{Csc}[x]^{2/(2*a)} - \text{Csc}[x]^{3/(3*a)} + \text{Log}[\text{Sin}[x]]/a$

Rule 75

```
Int[((d_)*(x_))^(n_)*((a_) + (b_)*(x_))*((e_) + (f_)*(x_))^(p_), x_Sym
bol] :> Int[ExpandIntegrand[(a + b*x)*(d*x)^n*(e + f*x)^p, x], x]; FreeQ[
{a, b, d, e, f, n}, x] && IGtQ[p, 0] && EqQ[b*e + a*f, 0] && !(ILtQ[n + p
+ 2, 0] && GtQ[n + 2*p, 0])
```

Rule 3879

```
Int[cot[(c_) + (d_)*(x_)]^(m_)*csc[(c_) + (d_)*(x_)]*(b_)^(n
_), x_Symbol] :> Dist[1/(a^(m - n - 1)*b^n*d), Subst[Int[((a - b*x)^((
m - 1)/2)*(a + b*x)^((m - 1)/2 + n))/x^(m + n), x], x, Sin[c + d*x]], x]; Fre
eQ[{a, b, c, d}, x] && IntegerQ[(m - 1)/2] && EqQ[a^2 - b^2, 0] && IntegerQ
[n]
```

Rubi steps

$$\begin{aligned} \int \frac{\cot^5(x)}{a + a \csc(x)} dx &= \frac{\text{Subst}\left(\int \frac{(a-ax)^2(a+ax)}{x^4} dx, x, \sin(x)\right)}{a^4} \\ &= \frac{\text{Subst}\left(\int \left(\frac{a^3}{x^4} - \frac{a^3}{x^3} - \frac{a^3}{x^2} + \frac{a^3}{x}\right) dx, x, \sin(x)\right)}{a^4} \\ &= \frac{\csc(x)}{a} + \frac{\csc^2(x)}{2a} - \frac{\csc^3(x)}{3a} + \frac{\log(\sin(x))}{a} \end{aligned}$$

Mathematica [A] time = 0.03, size = 26, normalized size = 0.72

$$\frac{-\frac{1}{3} \csc^3(x) + \frac{\csc^2(x)}{2} + \csc(x) + \log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[x]^5/(a + a*Csc[x]), x]

[Out] (Csc[x] + Csc[x]^2/2 - Csc[x]^3/3 + Log[Sin[x]])/a

fricas [A] time = 0.72, size = 45, normalized size = 1.25

$$\frac{6 (\cos(x)^2 - 1) \log\left(\frac{1}{2} \sin(x)\right) \sin(x) + 6 \cos(x)^2 - 3 \sin(x) - 4}{6 (a \cos(x)^2 - a) \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^5/(a+a*csc(x)), x, algorithm="fricas")

[Out] $\frac{1}{6} \cdot (6 \cdot (\cos(x)^2 - 1) \cdot \log(1/2 \cdot \sin(x)) \cdot \sin(x) + 6 \cdot \cos(x)^2 - 3 \cdot \sin(x) - 4) / ((a \cdot \cos(x)^2 - a) \cdot \sin(x))$

giac [A] time = 0.35, size = 29, normalized size = 0.81

$$\frac{\frac{6 \sin(x)^2 + 3 \sin(x) - 2}{\sin(x)^3} + 6 \log(|\sin(x)|)}{6 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^5/(a+a*csc(x)), x, algorithm="giac")

[Out] $\frac{1}{6} \cdot ((6 \cdot \sin(x)^2 + 3 \cdot \sin(x) - 2) / \sin(x)^3 + 6 \cdot \log(\text{abs}(\sin(x)))) / a$

maple [A] time = 0.38, size = 35, normalized size = 0.97

$$\frac{1}{2a \sin(x)^2} - \frac{1}{3a \sin(x)^3} + \frac{\ln(\sin(x))}{a} + \frac{1}{a \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^5/(a+a*csc(x)),x)`

[Out] $\frac{1}{2}/a/\sin(x)^2 - \frac{1}{3}/a/\sin(x)^3 + \ln(\sin(x))/a + 1/a/\sin(x)$

maxima [A] time = 0.31, size = 29, normalized size = 0.81

$$\frac{\log(\sin(x))}{a} + \frac{6 \sin(x)^2 + 3 \sin(x) - 2}{6 a \sin(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^5/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $\log(\sin(x))/a + \frac{1}{6}*(6*\sin(x)^2 + 3*\sin(x) - 2)/(a*\sin(x)^3)$

mupad [B] time = 0.30, size = 80, normalized size = 2.22

$$\frac{3 \tan\left(\frac{x}{2}\right)}{8 a} - \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} + \frac{\tan\left(\frac{x}{2}\right)^2}{8 a} - \frac{\tan\left(\frac{x}{2}\right)^3}{24 a} + \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{a} + \frac{3 \tan\left(\frac{x}{2}\right)^2 + \tan\left(\frac{x}{2}\right) - \frac{1}{3}}{8 a \tan\left(\frac{x}{2}\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^5/(a + a/sin(x)),x)`

[Out] $\frac{(3*\tan(x/2))/(8*a) - \log(\tan(x/2)^2 + 1)/a + \tan(x/2)^2/(8*a) - \tan(x/2)^3/(24*a) + \log(\tan(x/2))/a + (\tan(x/2) + 3*\tan(x/2)^2 - 1/3)/(8*a*\tan(x/2)^3)}$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot^5(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)**5/(a+a*csc(x)),x)`

[Out] `Integral(cot(x)**5/(csc(x) + 1), x)/a`

3.10 $\int \frac{\cot^6(x)}{a+a \csc(x)} dx$

Optimal. Leaf size=49

$$-\frac{x}{a} - \frac{3 \tanh^{-1}(\cos(x))}{8a} + \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a}$$

[Out] $-\frac{x}{a} - \frac{3 \operatorname{arctanh}(\cos(x))}{8a} + \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a}$

Rubi [A] time = 0.08, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.231, Rules used = {3888, 3881, 3770}

$$-\frac{x}{a} - \frac{3 \tanh^{-1}(\cos(x))}{8a} + \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\cot[x]^6/(a + a \csc[x]), x]$

[Out] $-\frac{(x/a) - (3 \operatorname{ArcTanh}[\cos[x]])/(8a) + (\cot[x]^3(4 - 3 \csc[x]))/(12a) - (\cot[x](8 - 3 \csc[x]))/(8a)}$

Rule 3770

$\operatorname{Int}[\csc[(c_.) + (d_.)*(x_.)], x_{\text{Symbol}}] \rightarrow -\operatorname{Simp}[\operatorname{ArcTanh}[\cos[c + d*x]]/d, x]$
/; FreeQ[{c, d}, x]

Rule 3881

$\operatorname{Int}[(\cot[(c_.) + (d_.)*(x_.)]*(\operatorname{e}^{(m_*)}))^{(m_*)}*(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_)), x_{\text{Symbol}}] \rightarrow -\operatorname{Simp}[(\operatorname{e}^{(e_*)}\operatorname{Cot}[c + d*x])^{(m - 1)}*(a*m + b*(m - 1)*\csc[c + d*x])/((d*m*(m - 1)), x) - \operatorname{Dist}[\operatorname{e}^{2/m}, \operatorname{Int}[(\operatorname{e}^{(e_*)}\operatorname{Cot}[c + d*x])^{(m - 2)}*(a*m + b*(m - 1)*\csc[c + d*x]), x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e\}, x] \&& \operatorname{GtQ}[m, 1]$

Rule 3888

$\operatorname{Int}[(\cot[(c_.) + (d_.)*(x_.)]*(\operatorname{e}^{(m_*)}))^{(m_*)}*(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_))^{(n_*)}, x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[a^{(2*n)}/\operatorname{e}^{(2*n)}, \operatorname{Int}[(\operatorname{e}^{(e_*)}\operatorname{Cot}[c + d*x])^{(m + 2*n)}/(-a + b*\csc[c + d*x])^n, x], x] /; \operatorname{FreeQ}[\{a, b, c, d, e, m\}, x] \&& \operatorname{EqQ}[a^2 - b^2, 0] \&& \operatorname{ILtQ}[n, 0]$

Rubi steps

$$\begin{aligned}
\int \frac{\cot^6(x)}{a + a \csc(x)} dx &= \frac{\int \cot^4(x)(-a + a \csc(x)) dx}{a^2} \\
&= \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\int \cot^2(x)(-4a + 3a \csc(x)) dx}{4a^2} \\
&= \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a} + \frac{\int (-8a + 3a \csc(x)) dx}{8a^2} \\
&= -\frac{x}{a} + \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a} + \frac{3 \int \csc(x) dx}{8a} \\
&= -\frac{x}{a} - \frac{3 \tanh^{-1}(\cos(x))}{8a} + \frac{\cot^3(x)(4 - 3 \csc(x))}{12a} - \frac{\cot(x)(8 - 3 \csc(x))}{8a}
\end{aligned}$$

Mathematica [B] time = 0.05, size = 163, normalized size = 3.33

$$-\frac{x}{a} + \frac{2 \tan\left(\frac{x}{2}\right)}{3a} - \frac{2 \cot\left(\frac{x}{2}\right)}{3a} - \frac{\csc^4\left(\frac{x}{2}\right)}{64a} + \frac{5 \csc^2\left(\frac{x}{2}\right)}{32a} + \frac{\sec^4\left(\frac{x}{2}\right)}{64a} - \frac{5 \sec^2\left(\frac{x}{2}\right)}{32a} + \frac{3 \log\left(\sin\left(\frac{x}{2}\right)\right)}{8a} - \frac{3 \log\left(\cos\left(\frac{x}{2}\right)\right)}{8a} + \frac{\cot\left(\frac{x}{2}\right)}{2}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^6/(a + a*Csc[x]), x]`

[Out] $-\frac{(x/a) - (2*\text{Cot}[x/2])/(3*a) + (5*\text{Csc}[x/2]^2)/(32*a) + (\text{Cot}[x/2]*\text{Csc}[x/2]^2)/(24*a) - \text{Csc}[x/2]^4/(64*a) - (3*\text{Log}[\text{Cos}[x/2]])/(8*a) + (3*\text{Log}[\text{Sin}[x/2]])/(8*a) - (5*\text{Sec}[x/2]^2)/(32*a) + \text{Sec}[x/2]^4/(64*a) + (2*\text{Tan}[x/2])/(3*a) - (\text{Se}c[x/2]^2*\text{Tan}[x/2])/(24*a)}$

fricas [B] time = 0.58, size = 104, normalized size = 2.12

$$\frac{48 x \cos(x)^4 - 96 x \cos(x)^2 + 30 \cos(x)^3 + 9 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log\left(\frac{1}{2} \cos(x) + \frac{1}{2}\right) - 9 (\cos(x)^4 - 2 \cos(x)^2 + 1)}{48 (\cos(x)^4 - 2 \cos(x)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^6/(a+a*csc(x)), x, algorithm="fricas")`

[Out] $-1/48*(48*x*\cos(x)^4 - 96*x*\cos(x)^2 + 30*\cos(x)^3 + 9*(\cos(x)^4 - 2*\cos(x)^2 + 1)*\log(1/2*\cos(x) + 1/2) - 9*(\cos(x)^4 - 2*\cos(x)^2 + 1)*\log(-1/2*\cos(x) + 1/2) - 16*(4*\cos(x)^3 - 3*\cos(x))*\sin(x) + 48*x - 18*\cos(x))/(a*\cos(x)^4 - 2*a*\cos(x)^2 + a)$

giac [B] time = 0.33, size = 109, normalized size = 2.22

$$-\frac{x}{a} + \frac{3 \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{8a} + \frac{3a^3 \tan\left(\frac{1}{2}x\right)^4 - 8a^3 \tan\left(\frac{1}{2}x\right)^3 - 24a^3 \tan\left(\frac{1}{2}x\right)^2 + 120a^3 \tan\left(\frac{1}{2}x\right) - 150 \tan\left(\frac{1}{2}x\right)^4 + 192a^4}{192a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^6/(a+a*csc(x)),x, algorithm="giac")`

[Out] $-x/a + 3/8*\log(\text{abs}(\tan(1/2*x)))/a + 1/192*(3*a^3*tan(1/2*x)^4 - 8*a^3*tan(1/2*x)^3 - 24*a^3*tan(1/2*x)^2 + 120*a^3*tan(1/2*x))/a^4 - 1/192*(150*tan(1/2*x)^4 + 120*tan(1/2*x)^3 - 24*tan(1/2*x)^2 - 8*tan(1/2*x) + 3)/(a*tan(1/2*x)^4)$

maple [B] time = 0.38, size = 108, normalized size = 2.20

$$\frac{\tan^4\left(\frac{x}{2}\right)}{64a} - \frac{\tan^3\left(\frac{x}{2}\right)}{24a} - \frac{\tan^2\left(\frac{x}{2}\right)}{8a} + \frac{5\tan\left(\frac{x}{2}\right)}{8a} - \frac{1}{64a\tan\left(\frac{x}{2}\right)^4} + \frac{1}{24a\tan\left(\frac{x}{2}\right)^3} + \frac{1}{8a\tan\left(\frac{x}{2}\right)^2} - \frac{5}{8a\tan\left(\frac{x}{2}\right)} + \frac{3\ln\left(\tan\left(\frac{x}{2}\right)\right)}{8a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^6/(a+a*csc(x)),x)`

[Out] $1/64/a*\tan(1/2*x)^4 - 1/24/a*\tan(1/2*x)^3 - 1/8/a*\tan(1/2*x)^2 + 5/8/a*\tan(1/2*x) - 1/64/a/\tan(1/2*x)^4 + 1/24/a/\tan(1/2*x)^3 + 1/8/a/\tan(1/2*x)^2 - 5/8/a/\tan(1/2*x) + 3/8/a*\ln(\tan(1/2*x)) - 2/a*\arctan(\tan(1/2*x))$

maxima [B] time = 0.42, size = 134, normalized size = 2.73

$$\frac{120 \sin(x)}{\cos(x)+1} - \frac{24 \sin(x)^2}{(\cos(x)+1)^2} - \frac{8 \sin(x)^3}{(\cos(x)+1)^3} + \frac{3 \sin(x)^4}{(\cos(x)+1)^4} - \frac{2 \arctan\left(\frac{\sin(x)}{\cos(x)+1}\right)}{a} + \frac{3 \log\left(\frac{\sin(x)}{\cos(x)+1}\right)}{8a} + \frac{\left(\frac{8 \sin(x)}{\cos(x)+1} + \frac{24 \sin(x)^2}{(\cos(x)+1)^2} - \frac{120 \sin(x)}{\cos(x)+1}\right)}{192a \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^6/(a+a*csc(x)),x, algorithm="maxima")`

[Out] $1/192*(120*\sin(x)/(\cos(x) + 1) - 24*\sin(x)^2/(\cos(x) + 1)^2 - 8*\sin(x)^3/(\cos(x) + 1)^3 + 3*\sin(x)^4/(\cos(x) + 1)^4)/a - 2*\arctan(\sin(x)/(\cos(x) + 1))/a + 3/8*\log(\sin(x)/(\cos(x) + 1))/a + 1/192*(8*\sin(x)/(\cos(x) + 1) + 24*\sin(x)^2/(\cos(x) + 1)^2 - 120*\sin(x)^3/(\cos(x) + 1)^3 - 3)*(\cos(x) + 1)^4/(a*\sin(x)^4)$

mupad [B] time = 0.33, size = 123, normalized size = 2.51

$$\frac{5 \tan\left(\frac{x}{2}\right)}{8 a} + \frac{2 \operatorname{atan}\left(\frac{4}{4 \tan\left(\frac{x}{2}\right) + \frac{3}{2}} - \frac{3 \tan\left(\frac{x}{2}\right)}{2 \left(4 \tan\left(\frac{x}{2}\right) + \frac{3}{2}\right)}\right)}{a} - \frac{\tan\left(\frac{x}{2}\right)^2}{8 a} - \frac{\tan\left(\frac{x}{2}\right)^3}{24 a} + \frac{\tan\left(\frac{x}{2}\right)^4}{64 a} + \frac{3 \ln\left(\tan\left(\frac{x}{2}\right)\right)}{8 a} + \frac{-10 \tan\left(\frac{x}{2}\right)^3 + 2}{16 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^6/(a + a/sin(x)),x)`

[Out] $\frac{(5*\tan(x/2))/(8*a) + (2*\operatorname{atan}(4/(4*\tan(x/2) + 3/2) - (3*\tan(x/2))/(2*(4*\tan(x/2) + 3/2))))/a - \tan(x/2)^2/(8*a) - \tan(x/2)^3/(24*a) + \tan(x/2)^4/(64*a) + (3*\log(\tan(x/2)))/(8*a) + ((2*\tan(x/2))/3 + 2*\tan(x/2)^2 - 10*\tan(x/2)^3 - 1/4)/(16*a*\tan(x/2)^4)}$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot^6(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)**6/(a+a*csc(x)),x)`

[Out] `Integral(cot(x)**6/(csc(x) + 1), x)/a`

3.11 $\int \frac{\cot^7(x)}{a + a \csc(x)} dx$

Optimal. Leaf size=58

$$-\frac{\csc^5(x)}{5a} + \frac{\csc^4(x)}{4a} + \frac{2\csc^3(x)}{3a} - \frac{\csc^2(x)}{a} - \frac{\csc(x)}{a} - \frac{\log(\sin(x))}{a}$$

[Out] $-\csc(x)/a - \csc(x)^2/a + 2/3*\csc(x)^3/a + 1/4*\csc(x)^4/a - 1/5*\csc(x)^5/a - \ln(\sin(x))/a$

Rubi [A] time = 0.06, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3879, 88}

$$-\frac{\csc^5(x)}{5a} + \frac{\csc^4(x)}{4a} + \frac{2\csc^3(x)}{3a} - \frac{\csc^2(x)}{a} - \frac{\csc(x)}{a} - \frac{\log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^7/(a + a*Csc[x]), x]

[Out] $-(\Csc[x]/a) - \Csc[x]^2/a + (2*\Csc[x]^3)/(3*a) + \Csc[x]^4/(4*a) - \Csc[x]^5/(5*a) - \Log[\Sin[x]]/a$

Rule 88

```
Int[((a_.) + (b_.*(x_))^m_.*((c_.) + (d_.*(x_))^n_.*((e_.) + (f_.*(x_))^p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))
```

Rule 3879

```
Int[cot[(c_.) + (d_.*(x_))^m_.*(csc[(c_.) + (d_.*(x_))]*(b_.) + (a_))^n_, x_Symbol] :> Dist[1/(a^(m - n - 1)*b^n*d), Subst[Int[((a - b*x)^(m - 1)/2)*(a + b*x)^(n/2 + 1))/x^(m + n), x], x, Sin[c + d*x]], x] /; FreeQ[{a, b, c, d}, x] && IntegerQ[(m - 1)/2] && EqQ[a^2 - b^2, 0] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \int \frac{\cot^7(x)}{a + a \csc(x)} dx &= \frac{\text{Subst}\left(\int \frac{(a-ax)^3(a+ax)^2}{x^6} dx, x, \sin(x)\right)}{a^6} \\ &= \frac{\text{Subst}\left(\int \left(\frac{a^5}{x^6} - \frac{a^5}{x^5} - \frac{2a^5}{x^4} + \frac{2a^5}{x^3} + \frac{a^5}{x^2} - \frac{a^5}{x}\right) dx, x, \sin(x)\right)}{a^6} \\ &= -\frac{\csc(x)}{a} - \frac{\csc^2(x)}{a} + \frac{2 \csc^3(x)}{3a} + \frac{\csc^4(x)}{4a} - \frac{\csc^5(x)}{5a} - \frac{\log(\sin(x))}{a} \end{aligned}$$

Mathematica [A] time = 0.05, size = 39, normalized size = 0.67

$$-\frac{\frac{\csc^5(x)}{5} - \frac{\csc^4(x)}{4} - \frac{2 \csc^3(x)}{3} + \csc^2(x) + \csc(x) + \log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^7/(a + a*Csc[x]), x]`

[Out] $-\left((\text{Csc}[x] + \text{Csc}[x]^2 - (2 \text{Csc}[x]^3)/3 - \text{Csc}[x]^4/4 + \text{Csc}[x]^5/5 + \text{Log}[\text{Sin}[x]])/a\right)$

fricas [A] time = 0.63, size = 70, normalized size = 1.21

$$\frac{60 \cos(x)^4 + 60 (\cos(x)^4 - 2 \cos(x)^2 + 1) \log\left(\frac{1}{2} \sin(x)\right) \sin(x) - 80 \cos(x)^2 - 15 (4 \cos(x)^2 - 3) \sin(x) + 32}{60 (a \cos(x)^4 - 2 a \cos(x)^2 + a) \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^7/(a+a*csc(x)), x, algorithm="fricas")`

[Out] $\frac{-1/60*(60*\cos(x)^4 + 60*(\cos(x)^4 - 2*\cos(x)^2 + 1)*\log(1/2*\sin(x))*\sin(x) - 80*\cos(x)^2 - 15*(4*\cos(x)^2 - 3)*\sin(x) + 32)/((a*\cos(x)^4 - 2*a*\cos(x)^2 + a)*\sin(x))}{60 a}$

giac [A] time = 0.54, size = 41, normalized size = 0.71

$$-\frac{\frac{60 \sin(x)^4 + 60 \sin(x)^3 - 40 \sin(x)^2 - 15 \sin(x) + 12}{\sin(x)^5} + 60 \log(|\sin(x)|)}{60 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^7/(a+a*csc(x)), x, algorithm="giac")`

[Out] $-1/60*((60*\sin(x)^4 + 60*\sin(x)^3 - 40*\sin(x)^2 - 15*\sin(x) + 12)/\sin(x)^5 + 60*\log(\sin(x)))/a$

maple [A] time = 0.38, size = 55, normalized size = 0.95

$$-\frac{1}{a \sin(x)^2} + \frac{2}{3a \sin(x)^3} + \frac{1}{4a \sin(x)^4} - \frac{\ln(\sin(x))}{a} - \frac{1}{5a \sin(x)^5} - \frac{1}{a \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \cot(x)^7/(a+a*csc(x)) dx$

[Out] $-1/a/\sin(x)^2 + 2/3/a/\sin(x)^3 + 1/4/a/\sin(x)^4 - \ln(\sin(x))/a - 1/5/a/\sin(x)^5 - 1/a/\sin(x)$

maxima [A] time = 0.32, size = 42, normalized size = 0.72

$$-\frac{\log(\sin(x))}{a} - \frac{60 \sin(x)^4 + 60 \sin(x)^3 - 40 \sin(x)^2 - 15 \sin(x) + 12}{60 a \sin(x)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(\cot(x)^7/(a+a*csc(x)), x, \text{algorithm}=\text{"maxima"})$

[Out] $-\log(\sin(x))/a - 1/60*(60*\sin(x)^4 + 60*\sin(x)^3 - 40*\sin(x)^2 - 15*\sin(x) + 12)/(a*\sin(x)^5)$

mupad [B] time = 0.43, size = 113, normalized size = 1.95

$$\frac{180 \tan\left(\frac{x}{2}\right)^3 - 960 \tan\left(\frac{x}{2}\right)^5 \ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right) - 50 \tan\left(\frac{x}{2}\right)^2 - 15 \tan\left(\frac{x}{2}\right) + 300 \tan\left(\frac{x}{2}\right)^4 + 300 \tan\left(\frac{x}{2}\right)^6 + 180 \tan\left(\frac{x}{2}\right)^8}{960 a \tan\left(\frac{x}{2}\right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \cot(x)^7/(a + a/\sin(x)) dx$

[Out] $-(180*\tan(x/2)^3 - 960*\tan(x/2)^5*\log(\tan(x/2)^2 + 1) - 50*\tan(x/2)^2 - 15*\tan(x/2) + 300*\tan(x/2)^4 + 300*\tan(x/2)^6 + 180*\tan(x/2)^7 - 50*\tan(x/2)^8 - 15*\tan(x/2)^9 + 6*\tan(x/2)^{10} + 960*\tan(x/2)^5*\log(\tan(x/2)) + 6)/(960*a*\tan(x/2)^5)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\cot^7(x)}{\csc(x)+1} dx}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cot(x)**7/(a+a*csc(x)),x)
[Out] Integral(cot(x)**7/(csc(x) + 1), x)/a
```

3.12 $\int \frac{\tan^5(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=178

$$-\frac{(8a^2 + 21ab + 15b^2) \log(1 - \csc(x))}{16(a + b)^3} - \frac{(8a^2 - 21ab + 15b^2) \log(\csc(x) + 1)}{16(a - b)^3} + \frac{b^6 \log(a + b \csc(x))}{a (a^2 - b^2)^3} + \frac{5a + 7b}{16(a + b)^2 (1 - \csc(x))^2}$$

[Out] $1/16/(a+b)/(1-\csc(x))^2 + 1/16*(5*a+7*b)/(a+b)^2/(1-\csc(x)) + 1/16/(a-b)/(1+\csc(x))^2 + 1/16*(5*a-7*b)/(a-b)^2/(1+\csc(x)) - 1/16*(8*a^2+21*a*b+15*b^2)*\ln(1-\csc(x))/(a+b)^3 - 1/16*(8*a^2-21*a*b+15*b^2)*\ln(1+\csc(x))/(a-b)^3 + b^6*\ln(a+b*\csc(x))/a/(a^2-b^2)^3 - \ln(\sin(x))/a$

Rubi [A] time = 0.27, antiderivative size = 178, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3885, 894}

$$\frac{b^6 \log(a + b \csc(x))}{a (a^2 - b^2)^3} - \frac{(8a^2 + 21ab + 15b^2) \log(1 - \csc(x))}{16(a + b)^3} - \frac{(8a^2 - 21ab + 15b^2) \log(\csc(x) + 1)}{16(a - b)^3} + \frac{5a + 7b}{16(a + b)^2 (1 - \csc(x))^2}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]^5/(a + b*Csc[x]), x]

[Out] $1/(16*(a + b)*(1 - \csc(x))^2) + (5*a + 7*b)/(16*(a + b)^2*(1 - \csc(x))) + 1/(16*(a - b)*(1 + \csc(x))^2) + (5*a - 7*b)/(16*(a - b)^2*(1 + \csc(x))) - ((8*a^2 + 21*a*b + 15*b^2)*\text{Log}[1 - \csc(x)])/(16*(a + b)^3) - ((8*a^2 - 21*a*b + 15*b^2)*\text{Log}[1 + \csc(x)])/(16*(a - b)^3) + (b^6*\text{Log}[a + b*\csc(x)])/(a*(a^2 - b^2)^3) - \text{Log}[\sin(x)]/a$

Rule 894

```
Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (c_)*(x_)^2)^p_, x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && (EqQ[p, 1] && IntegersQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))
```

Rule 3885

```
Int[cot[(c_) + (d_)*(x_)]^(m_)*csc[(c_) + (d_)*(x_)]*(b_)^(n_), x_Symbol] :> -Dist[(-1)^(m - 1)/2]/(d*b^(m - 1)), Subst[Int[((b^2 - x^2)^(m - 1)/2)*(a + x)^n/x, x], x, b*Csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned}
 \int \frac{\tan^5(x)}{a + b \csc(x)} dx &= b^6 \text{Subst} \left(\int \frac{1}{x(a+x)(b^2-x^2)^3} dx, x, b \csc(x) \right) \\
 &= b^6 \text{Subst} \left(\int \left(\frac{1}{8b^4(a+b)(b-x)^3} + \frac{5a+7b}{16b^5(a+b)^2(b-x)^2} + \frac{8a^2+21ab+15b^2}{16b^6(a+b)^3(b-x)} + \frac{1}{ab^6x} + \frac{1}{a} \right) dx, x, b \csc(x) \right) \\
 &= \frac{1}{16(a+b)(1-\csc(x))^2} + \frac{5a+7b}{16(a+b)^2(1-\csc(x))} + \frac{1}{16(a-b)(1+\csc(x))^2} + \frac{5a-7b}{16(a-b)^2(1+\csc(x))} + \dots
 \end{aligned}$$

Mathematica [A] time = 0.79, size = 172, normalized size = 0.97

$$\frac{\csc(x)(a \sin(x) + b) \left(-\frac{(8a^2+21ab+15b^2) \log(1-\sin(x))}{(a+b)^3} - \frac{(8a^2-21ab+15b^2) \log(\sin(x)+1)}{(a-b)^3} + \frac{16b^6 \log(a \sin(x)+b)}{a(a-b)^3(a+b)^3} + \frac{7a+9b}{(a+b)^2(\sin(x)-1)} \right)}{16(a+b \csc(x))}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]^5/(a + b*Csc[x]), x]`

[Out] $\frac{(\csc(x)*(b + a*\sin(x))*(-(8*a^2 + 21*a*b + 15*b^2)*\log[1 - \sin(x)]/(a + b)^3 - (8*a^2 - 21*a*b + 15*b^2)*\log[1 + \sin(x)]/(a - b)^3 + (16*b^6*\log[b + a*\sin(x)])/(a*(a - b)^3*(a + b)^3) + 1/((a + b)*(-1 + \sin(x))^2) + (7*a + 9*b)/((a + b)^2*(-1 + \sin(x))) + 1/((a - b)*(1 + \sin(x))^2) + (-7*a + 9*b)/((a - b)^2*(1 + \sin(x))))/(16*(a + b*\csc(x)))$

fricas [A] time = 1.42, size = 260, normalized size = 1.46

$$16 b^6 \cos(x)^4 \log(a \sin(x) + b) + 4 a^6 - 8 a^4 b^2 + 4 a^2 b^4 - (8 a^6 + 3 a^5 b - 24 a^4 b^2 - 10 a^3 b^3 + 24 a^2 b^4 + 15 a b^5) \cos(x)^2 + 24 a^5 b^3 - 48 a^4 b^5 + 12 a^3 b^7 - 12 a^2 b^9 + 3 a b^{11} - b^{13}) \sin(x)^4$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^5/(a+b*csc(x)), x, algorithm="fricas")`

[Out] $\frac{1}{16} (16 b^6 \cos(x)^4 \log(a \sin(x) + b) + 4 a^6 - 8 a^4 b^2 + 4 a^2 b^4 - (8 a^6 + 3 a^5 b - 24 a^4 b^2 - 10 a^3 b^3 + 24 a^2 b^4 + 15 a b^5) \cos(x)^4 \log(\sin(x) + 1) - (8 a^6 - 3 a^5 b - 24 a^4 b^2 + 10 a^3 b^3 + 24 a^2 b^4 - 15 a b^5) \cos(x)^4 \log(-\sin(x) + 1) - 8 (2 a^6 - 5 a^4 b^2 + 3 a^2 b^4) \cos(x)^2 - 2 (2 a^5 b - 4 a^3 b^3 + 2 a^2 b^5 - (5 a^5 b - 14 a^3 b^3 + 9 a^2 b^5) \cos(x)^2) \sin(x)) / ((a^7 - 3 a^5 b^2 + 3 a^3 b^4 - a^2 b^6) \cos(x)^4)$

giac [A] time = 0.42, size = 252, normalized size = 1.42

$$\frac{b^6 \log(|a \sin(x) + b|)}{a^7 - 3a^5b^2 + 3a^3b^4 - ab^6} - \frac{(8a^2 - 21ab + 15b^2) \log(\sin(x) + 1)}{16(a^3 - 3a^2b + 3ab^2 - b^3)} - \frac{(8a^2 + 21ab + 15b^2) \log(-\sin(x) + 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} - \frac{6a^5}{}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tan(x)^5/(a+b*csc(x)),x, algorithm="giac")

[Out] $b^6 \log(\text{abs}(a \sin(x) + b)) / (a^7 - 3a^5b^2 + 3a^3b^4 - a*b^6) - 1/16*(8*a^2 - 21*a*b + 15*b^2)*\log(\sin(x) + 1) / (a^3 - 3*a^2*b + 3*a*b^2 - b^3) - 1/16*(8*a^2 + 21*a*b + 15*b^2)*\log(-\sin(x) + 1) / (a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 1/8*(6*a^5 - 16*a^3*b^2 + 10*a*b^4 + (5*a^4*b - 14*a^2*b^3 + 9*b^5)*sin(x)^3 - 4*(2*a^5 - 5*a^3*b^2 + 3*a*b^4)*sin(x)^2 - (3*a^4*b - 10*a^2*b^3 + 7*b^5)*sin(x)) / ((a + b)^3*(a - b)^3*(\sin(x) + 1)^2*(\sin(x) - 1)^2)$

maple [A] time = 0.43, size = 216, normalized size = 1.21

$$\frac{b^6 \ln(b + a \sin(x))}{(a + b)^3 (a - b)^3 a} + \frac{1}{2(8a + 8b)(-1 + \sin(x))^2} + \frac{7a}{16(a + b)^2 (-1 + \sin(x))} + \frac{9b}{16(a + b)^2 (-1 + \sin(x))} - \frac{\ln(-1 + \sin(x))}{2(a + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(tan(x)^5/(a+b*csc(x)),x)

[Out] $b^6 / (a+b)^3 (a-b)^3 / a * \ln(b+a \sin(x)) + 1/2 / (8*a+8*b) / (-1+\sin(x))^2 + 7/16 / (a+b)^2 / (-1+\sin(x)) * a + 9/16 / (a+b)^2 / (-1+\sin(x)) * b - 1/2 / (a+b)^3 * \ln(-1+\sin(x)) * a^2 - 2/1/16 / (a+b)^3 * \ln(-1+\sin(x)) * a * b - 15/16 / (a+b)^3 * \ln(-1+\sin(x)) * b^2 + 1/2 / (8*a-8*b) / (1+\sin(x))^2 - 7/16 / (a-b)^2 / (1+\sin(x)) * a + 9/16 / (a-b)^2 / (1+\sin(x)) * b - 1/2 / (a-b)^3 * \ln(1+\sin(x)) * a^2 + 21/16 / (a-b)^3 * \ln(1+\sin(x)) * a * b - 15/16 / (a-b)^3 * \ln(1+\sin(x)) * b^2$

maxima [A] time = 0.33, size = 251, normalized size = 1.41

$$\frac{b^6 \log(a \sin(x) + b)}{a^7 - 3a^5b^2 + 3a^3b^4 - ab^6} - \frac{(8a^2 - 21ab + 15b^2) \log(\sin(x) + 1)}{16(a^3 - 3a^2b + 3ab^2 - b^3)} - \frac{(8a^2 + 21ab + 15b^2) \log(\sin(x) - 1)}{16(a^3 + 3a^2b + 3ab^2 + b^3)} - \frac{5a^2b}{8(a^3 - 3a^2b + 3ab^2 - b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tan(x)^5/(a+b*csc(x)),x, algorithm="maxima")

[Out] $b^6 \log(a \sin(x) + b) / (a^7 - 3a^5b^2 + 3a^3b^4 - a*b^6) - 1/16*(8*a^2 - 21*a*b + 15*b^2)*\log(\sin(x) + 1) / (a^3 - 3*a^2*b + 3*a*b^2 - b^3) - 1/16*(8*a^2 + 21*a*b + 15*b^2)*\log(\sin(x) - 1) / (a^3 + 3*a^2*b + 3*a*b^2 + b^3) - 1/8*((5*a^2*b - 9*b^3)*\sin(x)^3 + 6*a^3 - 10*a*b^2 - 4*(2*a^3 - 3*a*b^2)*\sin(x)^2 - 4*(2*a^3 - 3*a*b^2)*\sin(x)) / ((a + b)^3*(a - b)^3*(\sin(x) + 1)^2*(\sin(x) - 1)^2)$

$$(x)^2 - (3*a^2*b - 7*b^3)*\sin(x))/((a^4 - 2*a^2*b^2 + b^4)*\sin(x)^4 + a^4 - 2*a^2*b^2 + b^4 - 2*(a^4 - 2*a^2*b^2 + b^4)*\sin(x)^2)$$

mupad [B] time = 2.30, size = 445, normalized size = 2.50

$$\frac{\frac{2 \tan\left(\frac{x}{2}\right)^2 (2 a b^2 - a^3)}{a^4 - 2 a^2 b^2 + b^4} - \frac{4 \tan\left(\frac{x}{2}\right)^4 (3 a b^2 - 2 a^3)}{a^4 - 2 a^2 b^2 + b^4} + \frac{2 \tan\left(\frac{x}{2}\right)^6 (2 a b^2 - a^3)}{a^4 - 2 a^2 b^2 + b^4} + \frac{\tan\left(\frac{x}{2}\right)^7 (3 a^2 b - 7 b^3)}{4 (a^4 - 2 a^2 b^2 + b^4)} - \frac{\tan\left(\frac{x}{2}\right)^3 (11 a^2 b - 15 b^3)}{4 (a^4 - 2 a^2 b^2 + b^4)} - \frac{\tan\left(\frac{x}{2}\right)^5 (11 a^2 b - 15 b^3)}{4 (a^4 - 2 a^2 b^2 + b^4)}}{\tan\left(\frac{x}{2}\right)^8 - 4 \tan\left(\frac{x}{2}\right)^6 + 6 \tan\left(\frac{x}{2}\right)^4 - 4 \tan\left(\frac{x}{2}\right)^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^5/(a + b/sin(x)), x)`

[Out] $((2*\tan(x/2)^2*(2*a*b^2 - a^3))/(a^4 + b^4 - 2*a^2*b^2) - (4*\tan(x/2)^4*(3*a*b^2 - 2*a^3))/(a^4 + b^4 - 2*a^2*b^2) + (2*\tan(x/2)^6*(2*a*b^2 - a^3))/(a^4 + b^4 - 2*a^2*b^2) + (\tan(x/2)^7*(3*a^2*b - 7*b^3))/(4*(a^4 + b^4 - 2*a^2*b^2)) - (\tan(x/2)^3*(11*a^2*b - 15*b^3))/(4*(a^4 + b^4 - 2*a^2*b^2)) - (\tan(x/2)^5*(11*a^2*b - 15*b^3))/(4*(a^4 + b^4 - 2*a^2*b^2)) + (b*tan(x/2)*(3*a^2 - 7*b^2))/(4*(a^4 + b^4 - 2*a^2*b^2)))/(6*tan(x/2)^4 - 4*tan(x/2)^2 - 4*tan(x/2)^6 + tan(x/2)^8 + 1) - log(tan(x/2) - 1)*((5*b)/(8*(a + b)^2) + 1/(a + b) + b^2/(4*(a + b)^3)) - log(tan(x/2) + 1)*(b^2/(4*(a - b)^3) - (5*b)/(8*(a - b)^2) + 1/(a - b)) + log(tan(x/2)^2 + 1)/a - (b^6*log(b + 2*a*tan(x/2) + b*tan(x/2)^2))/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tan^5(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)**5/(a+b*csc(x)), x)`

[Out] `Integral(tan(x)**5/(a + b*csc(x)), x)`

3.13 $\int \frac{\tan^3(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=113

$$\frac{b^4 \log(a + b \csc(x))}{a (a^2 - b^2)^2} - \frac{1}{4(a + b)(1 - \csc(x))} - \frac{1}{4(a - b)(\csc(x) + 1)} + \frac{(2a + 3b) \log(1 - \csc(x))}{4(a + b)^2} + \frac{(2a - 3b) \log(\csc(x))}{4(a - b)^2}$$

[Out] $-1/4/(a+b)/(1-\csc(x))-1/4/(a-b)/(1+\csc(x))+1/4*(2*a+3*b)*\ln(1-\csc(x))/(a+b)$
 $^2+1/4*(2*a-3*b)*\ln(1+\csc(x))/(a-b)^2+b^4*4*\ln(a+b*\csc(x))/a/(a^2-b^2)^2+2+\ln(s$
 $\ln(x))/a$

Rubi [A] time = 0.17, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3885, 894}

$$\frac{b^4 \log(a + b \csc(x))}{a (a^2 - b^2)^2} - \frac{1}{4(a + b)(1 - \csc(x))} - \frac{1}{4(a - b)(\csc(x) + 1)} + \frac{(2a + 3b) \log(1 - \csc(x))}{4(a + b)^2} + \frac{(2a - 3b) \log(\csc(x))}{4(a - b)^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\tan[x]^3/(a + b*\csc[x]), x]$

[Out] $-1/(4*(a + b)*(1 - \csc[x])) - 1/(4*(a - b)*(1 + \csc[x])) + ((2*a + 3*b)*\log[1 - \csc[x]])/(4*(a + b)^2) + ((2*a - 3*b)*\log[1 + \csc[x]])/(4*(a - b)^2) + (b^4*\log[a + b*\csc[x]])/(a*(a^2 - b^2)^2) + \log[\sin[x]]/a$

Rule 894

```
Int[((d_.) + (e_)*(x_))^m_*((f_)*(x_))^n_*((a_)*(c_)*(x_)^2)^p_, x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x)^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && (EqQ[p, 1] && IntegersQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))
```

Rule 3885

```
Int[cot[(c_)*(x_)^m_]*csc[(c_)*(x_)]*(b_)*(a_))^(n_), x_Symbol] :> -Dist[(-1)^((m - 1)/2)/(d*b^(m - 1)), Subst[Int[((b^2 - x^2)^(m - 1)/2)*(a + x)^n)/x, x], x, b*csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan^3(x)}{a + b \csc(x)} dx &= - \left(b^4 \text{Subst} \left(\int \frac{1}{x(a+x)(b^2-x^2)^2} dx, x, b \csc(x) \right) \right) \\
&= - \left(b^4 \text{Subst} \left(\int \left(\frac{1}{4b^3(a+b)(b-x)^2} + \frac{2a+3b}{4b^4(a+b)^2(b-x)} + \frac{1}{ab^4x} - \frac{1}{a(a-b)^2(a+b)^2(a+x)} \right) dx, x, b \csc(x) \right) \right) \\
&= - \frac{1}{4(a+b)(1-\csc(x))} - \frac{1}{4(a-b)(1+\csc(x))} + \frac{(2a+3b)\log(1-\csc(x))}{4(a+b)^2} + \frac{(2a-3b)\log(1+\csc(x))}{4(a-b)^2}
\end{aligned}$$

Mathematica [A] time = 0.49, size = 115, normalized size = 1.02

$$\frac{\csc(x)(a \sin(x) + b) \left(\frac{4b^4 \log(a \sin(x) + b)}{a(a-b)^2(a+b)^2} - \frac{1}{(a+b)(\sin(x)-1)} + \frac{1}{(a-b)(\sin(x)+1)} + \frac{(2a+3b) \log(1-\sin(x))}{(a+b)^2} + \frac{(2a-3b) \log(\sin(x)+1)}{(a-b)^2} \right)}{4(a+b \csc(x))}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]^3/(a + b*Csc[x]), x]`

[Out] `(Csc[x]*(b + a*Sin[x])*(((2*a + 3*b)*Log[1 - Sin[x]])/(a + b)^2 + ((2*a - 3*b)*Log[1 + Sin[x]])/(a - b)^2 + (4*b^4*Log[b + a*Sin[x]])/(a*(a - b)^2*(a + b)^2) - 1/((a + b)*(-1 + Sin[x])) + 1/((a - b)*(1 + Sin[x]))))/((4*(a + b)*Csc[x]))`

fricas [A] time = 1.87, size = 144, normalized size = 1.27

$$\frac{4 b^4 \cos(x)^2 \log(a \sin(x) + b) + 2 a^4 - 2 a^2 b^2 + (2 a^4 + a^3 b - 4 a^2 b^2 - 3 a b^3) \cos(x)^2 \log(\sin(x) + 1) + (2 a^4 - a^3 b - 4 a^2 b^2 - 3 a b^3) \cos(x)^2 \log(-\sin(x) + 1) - 2*(a^3 b - a b^3) \sin(x)) / ((a^5 - 2 a^3 b^2 + a b^4) \cos(x)^2)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^3/(a+b*csc(x)), x, algorithm="fricas")`

[Out] `1/4*(4*b^4*cos(x)^2*log(a*sin(x) + b) + 2*a^4 - 2*a^2*b^2 + (2*a^4 + a^3*b - 4*a^2*b^2 - 3*a*b^3)*cos(x)^2*log(sin(x) + 1) + (2*a^4 - a^3*b - 4*a^2*b^2 + 3*a*b^3)*cos(x)^2*log(-sin(x) + 1) - 2*(a^3*b - a*b^3)*sin(x)) / ((a^5 - 2*a^3*b^2 + a*b^4)*cos(x)^2)`

giac [A] time = 0.23, size = 139, normalized size = 1.23

$$\frac{b^4 \log(|a \sin(x) + b|)}{a^5 - 2 a^3 b^2 + a b^4} + \frac{(2 a - 3 b) \log(\sin(x) + 1)}{4 (a^2 - 2 a b + b^2)} + \frac{(2 a + 3 b) \log(-\sin(x) + 1)}{4 (a^2 + 2 a b + b^2)} - \frac{a^3 - a b^2 - (a^2 b - b^3) \sin(x)}{2 (a + b)^2 (a - b)^2 (\sin(x) + 1) (\cos(x))^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^3/(a+b*csc(x)),x, algorithm="giac")`

[Out] $b^4 \log(\text{abs}(a \sin(x) + b)) / (a^5 - 2a^3b^2 + a^2b^4) + 1/4*(2a - 3b)*\log(\sin(x) + 1) / (a^2 - 2a*b + b^2) + 1/4*(2a + 3b)*\log(-\sin(x) + 1) / (a^2 + 2a*b + b^2) - 1/2*(a^3 - a^2b^2 - (a^2b - b^3)*\sin(x)) / ((a + b)^2*(a - b)^2) * (\sin(x) + 1)*(\sin(x) - 1)$

maple [A] time = 0.51, size = 117, normalized size = 1.04

$$\frac{b^4 \ln(b + a \sin(x))}{(a + b)^2 (a - b)^2 a} - \frac{1}{(4a + 4b)(-1 + \sin(x))} + \frac{a \ln(-1 + \sin(x))}{2(a + b)^2} + \frac{3 \ln(-1 + \sin(x)) b}{4(a + b)^2} + \frac{1}{(4a - 4b)(1 + \sin(x))} - \frac{3 \ln(-1 + \sin(x)) b}{4(a - b)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^3/(a+b*csc(x)),x)`

[Out] $b^4 / (a+b)^2 / (a-b)^2 / a * \ln(b+a \sin(x)) - 1 / (4a+4b) / (-1+\sin(x)) + 1/2*a / (a+b)^2 * \ln(-1+\sin(x)) + 3/4 / (a+b)^2 * \ln(-1+\sin(x)) * b + 1 / (4a-4b) / (1+\sin(x)) - 3/4 / (a-b)^2 * \ln(1+\sin(x)) * b + 1/2*a * \ln(1+\sin(x)) / (a-b)^2$

maxima [A] time = 0.32, size = 120, normalized size = 1.06

$$\frac{b^4 \log(a \sin(x) + b)}{a^5 - 2a^3b^2 + ab^4} + \frac{(2a - 3b) \log(\sin(x) + 1)}{4(a^2 - 2ab + b^2)} + \frac{(2a + 3b) \log(\sin(x) - 1)}{4(a^2 + 2ab + b^2)} + \frac{b \sin(x) - a}{2((a^2 - b^2) \sin(x)^2 - a^2 + b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^3/(a+b*csc(x)),x, algorithm="maxima")`

[Out] $b^4 \log(a \sin(x) + b) / (a^5 - 2a^3b^2 + a^2b^4) + 1/4*(2a - 3b)*\log(\sin(x) + 1) / (a^2 - 2a*b + b^2) + 1/4*(2a + 3b)*\log(\sin(x) - 1) / (a^2 + 2a*b + b^2) + 1/2*(b*\sin(x) - a) / ((a^2 - b^2)*\sin(x)^2 - a^2 + b^2)$

mupad [B] time = 1.14, size = 172, normalized size = 1.52

$$\frac{\ln\left(\tan\left(\frac{x}{2}\right) + 1\right) (2a - 3b)}{2(a - b)^2} - \frac{\frac{b \tan\left(\frac{x}{2}\right)^3}{a^2 - b^2} - \frac{2a \tan\left(\frac{x}{2}\right)^2}{a^2 - b^2} + \frac{b \tan\left(\frac{x}{2}\right)}{a^2 - b^2}}{\tan\left(\frac{x}{2}\right)^4 - 2 \tan\left(\frac{x}{2}\right)^2 + 1} - \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} + \frac{\ln\left(\tan\left(\frac{x}{2}\right) - 1\right) (2a + 3b)}{2(a + b)^2} + \frac{b^4 \ln(-1 + \sin(x))}{4(a - b)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^3/(a + b/sin(x)),x)`

[Out] $(\log(\tan(x/2) + 1)*(2a - 3b)) / (2*(a - b)^2) - ((b*\tan(x/2)^3) / (a^2 - b^2)) - (2*a*\tan(x/2)^2) / (a^2 - b^2) + (b*\tan(x/2)) / (a^2 - b^2) / (\tan(x/2)^4 - 2)$

```
*tan(x/2)^2 + 1) - log(tan(x/2)^2 + 1)/a + (log(tan(x/2) - 1)*(2*a + 3*b))/  
(2*(a + b)^2) + (b^4*log(b + 2*a*tan(x/2) + b*tan(x/2)^2))/(a*(a^2 - b^2)^2  
)
```

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tan^3(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tan(x)**3/(a+b*csc(x)),x)  
[Out] Integral(tan(x)**3/(a + b*csc(x)), x)
```

3.14 $\int \frac{\tan(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=66

$$\frac{b^2 \log(a + b \csc(x))}{a (a^2 - b^2)} - \frac{\log(1 - \csc(x))}{2(a + b)} - \frac{\log(\csc(x) + 1)}{2(a - b)} - \frac{\log(\sin(x))}{a}$$

[Out] $-1/2 \ln(1 - \csc(x))/(a + b) - 1/2 \ln(1 + \csc(x))/(a - b) + b^2 \ln(a + b \csc(x))/a / (a^2 - b^2) - \ln(\sin(x))/a$

Rubi [A] time = 0.09, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.182, Rules used = {3885, 894}

$$\frac{b^2 \log(a + b \csc(x))}{a (a^2 - b^2)} - \frac{\log(1 - \csc(x))}{2(a + b)} - \frac{\log(\csc(x) + 1)}{2(a - b)} - \frac{\log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]/(a + b*Csc[x]), x]

[Out] $-\text{Log}[1 - \csc(x)]/(2*(a + b)) - \text{Log}[1 + \csc(x)]/(2*(a - b)) + (b^2 \text{Log}[a + b * \csc(x])]/(a*(a^2 - b^2)) - \text{Log}[\sin(x)]/a$

Rule 894

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] :> Int[ExpandIntegrand[((d + e*x)^m*(f + g*x)^n*(a + c*x^2)^p, x), x]; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && ((EqQ[p, 1] && IntegersQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))
```

Rule 3885

```
Int[cot[(c_.) + (d_.)*(x_)]^(m_)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n_), x_Symbol] :> -Dist[(-1)^(m - 1)/2]/(d*b^(m - 1)), Subst[Int[((b^2 - x^2)^(m - 1)/2)*(a + x)^n/x, x], x, b*Csc[c + d*x]], x]; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan(x)}{a + b \csc(x)} dx &= b^2 \operatorname{Subst} \left(\int \frac{1}{x(a+x)(b^2-x^2)} dx, x, b \csc(x) \right) \\
&= b^2 \operatorname{Subst} \left(\int \left(\frac{1}{2b^2(a+b)(b-x)} + \frac{1}{ab^2x} + \frac{1}{a(a-b)(a+b)(a+x)} - \frac{1}{2(a-b)b^2(b+x)} \right) dx, x, b \csc(x) \right) \\
&= -\frac{\log(1-\csc(x))}{2(a+b)} - \frac{\log(1+\csc(x))}{2(a-b)} + \frac{b^2 \log(a+b \csc(x))}{a(a^2-b^2)} - \frac{\log(\sin(x))}{a}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 56, normalized size = 0.85

$$-\frac{-2b^2 \log(a \sin(x) + b) + a(a-b) \log(1 - \sin(x)) + a(a+b) \log(\sin(x) + 1)}{2a(a-b)(a+b)}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]/(a + b*Csc[x]), x]`

[Out] $\frac{-1/2*(a*(a - b)*\operatorname{Log}[1 - \operatorname{Sin}[x]] + a*(a + b)*\operatorname{Log}[1 + \operatorname{Sin}[x]] - 2*b^2*\operatorname{Log}[b + a*\operatorname{Sin}[x]])/(a*(a - b)*(a + b))}{}$

fricas [A] time = 1.25, size = 58, normalized size = 0.88

$$\frac{2b^2 \log(a \sin(x) + b) - (a^2 + ab) \log(\sin(x) + 1) - (a^2 - ab) \log(-\sin(x) + 1)}{2(a^3 - ab^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+b*csc(x)), x, algorithm="fricas")`

[Out] $\frac{1/2*(2*b^2*\operatorname{log}(a*\operatorname{sin}(x) + b) - (a^2 + a*b)*\operatorname{log}(\operatorname{sin}(x) + 1) - (a^2 - a*b)*\operatorname{log}(-\operatorname{sin}(x) + 1))/(a^3 - a*b^2)}{}$

giac [A] time = 0.47, size = 53, normalized size = 0.80

$$\frac{b^2 \log(|a \sin(x) + b|)}{a^3 - ab^2} - \frac{\log(\sin(x) + 1)}{2(a - b)} - \frac{\log(-\sin(x) + 1)}{2(a + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+b*csc(x)), x, algorithm="giac")`

[Out] $\frac{b^2*\operatorname{log}(\operatorname{abs}(a*\operatorname{sin}(x) + b))/(a^3 - a*b^2) - 1/2*\operatorname{log}(\operatorname{sin}(x) + 1)/(a - b) - 1/2*\operatorname{log}(-\operatorname{sin}(x) + 1)/(a + b)}{}$

maple [A] time = 0.54, size = 60, normalized size = 0.91

$$\frac{b^2 \ln(b + a \sin(x))}{(a + b)(a - b)a} - \frac{\ln(-1 + \sin(x))}{2a + 2b} - \frac{\ln(1 + \sin(x))}{2a - 2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)/(a+b*csc(x)),x)`

[Out] $b^2/(a+b)/(a-b)/a*\ln(b+a*\sin(x))-1/(2*a+2*b)*\ln(-1+\sin(x))-1/(2*a-2*b)*\ln(1+\sin(x))$

maxima [A] time = 0.32, size = 50, normalized size = 0.76

$$\frac{b^2 \log(a \sin(x) + b)}{a^3 - ab^2} - \frac{\log(\sin(x) + 1)}{2(a - b)} - \frac{\log(\sin(x) - 1)}{2(a + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+b*csc(x)),x, algorithm="maxima")`

[Out] $b^2*\log(a*\sin(x) + b)/(a^3 - a*b^2) - 1/2*\log(\sin(x) + 1)/(a - b) - 1/2*\log(\sin(x) - 1)/(a + b)$

mupad [B] time = 0.44, size = 80, normalized size = 1.21

$$\frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{\ln\left(\tan\left(\frac{x}{2}\right) + 1\right)}{a - b} - \frac{\ln\left(\tan\left(\frac{x}{2}\right) - 1\right)}{a + b} + \frac{b^2 \ln\left(b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b\right)}{a (a^2 - b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)/(a + b/sin(x)),x)`

[Out] $\log(\tan(x/2)^2 + 1)/a - \log(\tan(x/2) + 1)/(a - b) - \log(\tan(x/2) - 1)/(a + b) + (b^2*\log(b + 2*a*tan(x/2) + b*tan(x/2)^2))/(a*(a^2 - b^2))$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tan(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)/(a+b*csc(x)),x)`

[Out] `Integral(tan(x)/(a + b*csc(x)), x)`

3.15 $\int \frac{\cot(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=19

$$\frac{\log(a + b \csc(x))}{a} + \frac{\log(\sin(x))}{a}$$

[Out] $\ln(a+b*\csc(x))/a+\ln(\sin(x))/a$

Rubi [A] time = 0.02, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.364, Rules used = {3885, 36, 29, 31}

$$\frac{\log(a + b \csc(x))}{a} + \frac{\log(\sin(x))}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[x]/(a + b*\text{Csc}[x]), x]$

[Out] $\text{Log}[a + b*\text{Csc}[x]]/a + \text{Log}[\text{Sin}[x]]/a$

Rule 29

$\text{Int}[(x_.)^{-1}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[x], x]$

Rule 31

$\text{Int}[((a_.) + (b_.)*(x_.))^{-1}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}[\{a, b\}, x]$

Rule 36

$\text{Int}[1/(((a_.) + (b_.)*(x_.))*(c_.) + (d_.)*(x_.))), x_Symbol] \rightarrow \text{Dist}[b/(b*c - a*d), \text{Int}[1/(a + b*x), x], x] - \text{Dist}[d/(b*c - a*d), \text{Int}[1/(c + d*x), x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&& \text{NeQ}[b*c - a*d, 0]$

Rule 3885

$\text{Int}[\cot[(c_.) + (d_.)*(x_.)]^{(m_.)}*(\csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^{(n_.)}, x_Symbol] \rightarrow -\text{Dist}[(-1)^{(m - 1)/2}/(d*b^{(m - 1)}), \text{Subst}[\text{Int}[(b^2 - x^2)^{(m - 1)/2}*(a + x)^n/x, x], x, b*\text{Csc}[c + d*x]], x] /; \text{FreeQ}[\{a, b, c, d, n\}, x] \&& \text{IntegerQ}[(m - 1)/2] \&& \text{NeQ}[a^2 - b^2, 0]$

Rubi steps

$$\begin{aligned}
\int \frac{\cot(x)}{a + b \csc(x)} dx &= -\text{Subst}\left(\int \frac{1}{x(a+x)} dx, x, b \csc(x)\right) \\
&= -\frac{\text{Subst}\left(\int \frac{1}{x} dx, x, b \csc(x)\right)}{a} + \frac{\text{Subst}\left(\int \frac{1}{a+x} dx, x, b \csc(x)\right)}{a} \\
&= \frac{\log(a + b \csc(x))}{a} + \frac{\log(\sin(x))}{a}
\end{aligned}$$

Mathematica [A] time = 0.01, size = 11, normalized size = 0.58

$$\frac{\log(a \sin(x) + b)}{a}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]/(a + b*Csc[x]), x]`

[Out] `Log[b + a*Sin[x]]/a`

fricas [A] time = 0.55, size = 11, normalized size = 0.58

$$\frac{\log(a \sin(x) + b)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)/(a+b*csc(x)), x, algorithm="fricas")`

[Out] `log(a*sin(x) + b)/a`

giac [A] time = 0.42, size = 12, normalized size = 0.63

$$\frac{\log(|a \sin(x) + b|)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)/(a+b*csc(x)), x, algorithm="giac")`

[Out] `log(abs(a*sin(x) + b))/a`

maple [A] time = 0.32, size = 21, normalized size = 1.11

$$\frac{\ln(a + b \csc(x))}{a} - \frac{\ln(\csc(x))}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)/(a+b*csc(x)),x)`

[Out] `ln(a+b*csc(x))/a-1/a*ln(csc(x))`

maxima [A] time = 0.31, size = 11, normalized size = 0.58

$$\frac{\log(a \sin(x) + b)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)/(a+b*csc(x)),x, algorithm="maxima")`

[Out] `log(a*sin(x) + b)/a`

mupad [B] time = 0.42, size = 55, normalized size = 2.89

$$\frac{2 \operatorname{atanh}\left(\frac{a \left(2 b^3 \sin(x)+\frac{5 a b^2}{2}-a^3-\frac{a b^2 \cos(2 x)}{2}\right)}{\left(-a^2+\sin(x) a b+2 b^2\right)^2}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)/(a + b/sin(x)),x)`

[Out] `(2*atanh((a*(2*b^3*sin(x) + (5*a*b^2)/2 - a^3 - (a*b^2*cos(2*x))/2))/(2*b^2 - a^2 + a*b*sin(x))^2))/a`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)/(a+b*csc(x)),x)`

[Out] `Integral(cot(x)/(a + b*csc(x)), x)`

3.16 $\int \frac{\cot^3(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=38

$$-\frac{\left(1 - \frac{a^2}{b^2}\right) \log(a + b \csc(x))}{a} - \frac{\log(\sin(x))}{a} - \frac{\csc(x)}{b}$$

[Out] $-\csc(x)/b - (1-a^2/b^2)*\ln(a+b*\csc(x))/a - \ln(\sin(x))/a$

Rubi [A] time = 0.06, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3885, 894}

$$-\frac{\left(1 - \frac{a^2}{b^2}\right) \log(a + b \csc(x))}{a} - \frac{\log(\sin(x))}{a} - \frac{\csc(x)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^3/(a + b*Csc[x]), x]

[Out] $-(\Csc[x]/b) - ((1 - a^2/b^2)*\Log[a + b*\Csc[x]])/a - \Log[\Sin[x]]/a$

Rule 894

```
Int[((d_.) + (e_.)*(x_))^m_*((f_.) + (g_.)*(x_))^n_*((a_) + (c_.)*(x_)^2)^p_, x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x)^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && ((EqQ[p, 1] && IntegersQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))
```

Rule 3885

```
Int[cot[(c_.) + (d_.)*(x_)]^m_*((csc[(c_.) + (d_.)*(x_.)]*(b_.) + (a_.))^n_2)^((m - 1)/2)/(d*b^(m - 1)), Subst[Int[((b^2 - x^2)^((m - 1)/2)*(a + x)^n)/x, x], x, b*Csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cot^3(x)}{a + b \csc(x)} dx &= \frac{\text{Subst}\left(\int \frac{b^2 - x^2}{x(a+x)} dx, x, b \csc(x)\right)}{b^2} \\
&= \frac{\text{Subst}\left(\int \left(-1 + \frac{b^2}{ax} + \frac{a^2 - b^2}{a(a+x)}\right) dx, x, b \csc(x)\right)}{b^2} \\
&= -\frac{\csc(x)}{b} - \frac{\left(1 - \frac{a^2}{b^2}\right) \log(a + b \csc(x))}{a} - \frac{\log(\sin(x))}{a}
\end{aligned}$$

Mathematica [A] time = 0.05, size = 39, normalized size = 1.03

$$\frac{(a^2 - b^2) \log(a \sin(x) + b) + a^2(-\log(\sin(x))) - ab \csc(x)}{ab^2}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^3/(a + b*Csc[x]), x]`

[Out] `(-(a*b*Csc[x]) - a^2*Log[Sin[x]] + (a^2 - b^2)*Log[b + a*Sin[x]])/(a*b^2)`

fricas [A] time = 0.48, size = 47, normalized size = 1.24

$$-\frac{a^2 \log\left(-\frac{1}{2} \sin(x)\right) \sin(x) - (a^2 - b^2) \log(a \sin(x) + b) \sin(x) + ab}{ab^2 \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^3/(a+b*csc(x)), x, algorithm="fricas")`

[Out] `-(a^2*log(-1/2*sin(x))*sin(x) - (a^2 - b^2)*log(a*sin(x) + b)*sin(x) + a*b)/(a*b^2*sin(x))`

giac [A] time = 0.65, size = 44, normalized size = 1.16

$$-\frac{a \log(|\sin(x)|)}{b^2} + \frac{(a^2 - b^2) \log(|a \sin(x) + b|)}{ab^2} - \frac{1}{b \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^3/(a+b*csc(x)), x, algorithm="giac")`

[Out] `-a*log(abs(sin(x)))/b^2 + (a^2 - b^2)*log(abs(a*sin(x) + b))/(a*b^2) - 1/(b*sin(x))`

maple [A] time = 0.38, size = 44, normalized size = 1.16

$$\frac{a \ln(b + a \sin(x))}{b^2} - \frac{\ln(b + a \sin(x))}{a} - \frac{1}{b \sin(x)} - \frac{a \ln(\sin(x))}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^3/(a+b*csc(x)),x)`

[Out] `1/b^2*a*ln(b+a*sin(x))-1/a*ln(b+a*sin(x))-1/b/sin(x)-a/b^2*ln(sin(x))`

maxima [A] time = 0.31, size = 42, normalized size = 1.11

$$-\frac{a \log(\sin(x))}{b^2} + \frac{(a^2 - b^2) \log(a \sin(x) + b)}{ab^2} - \frac{1}{b \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^3/(a+b*csc(x)),x, algorithm="maxima")`

[Out] `-a*log(sin(x))/b^2 + (a^2 - b^2)*log(a*sin(x) + b)/(a*b^2) - 1/(b*sin(x))`

mupad [B] time = 0.46, size = 75, normalized size = 1.97

$$\ln\left(b \tan\left(\frac{x}{2}\right)^2 + 2a \tan\left(\frac{x}{2}\right) + b\right) \left(\frac{a}{b^2} - \frac{1}{a}\right) - \frac{\tan\left(\frac{x}{2}\right)}{2b} + \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{1}{2b \tan\left(\frac{x}{2}\right)} - \frac{a \ln\left(\tan\left(\frac{x}{2}\right)\right)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^3/(a + b/sin(x)),x)`

[Out] `log(b + 2*a*tan(x/2) + b*tan(x/2)^2)*(a/b^2 - 1/a) - tan(x/2)/(2*b) + log(tan(x/2)^2 + 1)/a - 1/(2*b*tan(x/2)) - (a*log(tan(x/2)))/b^2`

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot^3(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)**3/(a+b*csc(x)),x)`

[Out] `Integral(cot(x)**3/(a + b*csc(x)), x)`

3.17 $\int \frac{\cot^5(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=72

$$\frac{(a^2 - b^2)^2 \log(a + b \csc(x))}{ab^4} - \frac{(a^2 - 2b^2) \csc(x)}{b^3} + \frac{a \csc^2(x)}{2b^2} + \frac{\log(\sin(x))}{a} - \frac{\csc^3(x)}{3b}$$

[Out] $-(a^2 - 2b^2) * \csc(x) / b^3 + 1/2 * a * \csc(x)^2 / b^2 - 1/3 * \csc(x)^3 / b + (a^2 - b^2)^2 / 2 * \ln(a + b * \csc(x)) / a + b^4 * \ln(\sin(x)) / a$

Rubi [A] time = 0.08, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3885, 894}

$$-\frac{(a^2 - 2b^2) \csc(x)}{b^3} + \frac{(a^2 - b^2)^2 \log(a + b \csc(x))}{ab^4} + \frac{a \csc^2(x)}{2b^2} + \frac{\log(\sin(x))}{a} - \frac{\csc^3(x)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^5/(a + b*Csc[x]), x]

[Out] $-(((a^2 - 2b^2) * \csc(x)) / b^3) + (a * \csc(x)^2) / (2 * b^2) - \csc(x)^3 / (3 * b) + ((a^2 - b^2)^2 * \log[a + b * \csc(x)]) / (a * b^4) + \log[\sin(x)] / a$

Rule 894

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x)^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && ((EqQ[p, 1] && IntegersQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))
```

Rule 3885

```
Int[cot[(c_.) + (d_.)*(x_)]^(m_)*csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_), x_Symbol] :> -Dist[(-1)^((m - 1)/2)/(d*b^(m - 1)), Subst[Int[((b^2 - x^2)^(m - 1)/2)*(a + x)^n)/x, x], x, b*Csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cot^5(x)}{a + b \csc(x)} dx &= -\frac{\text{Subst}\left(\int \frac{(b^2-x^2)^2}{x(a+x)} dx, x, b \csc(x)\right)}{b^4} \\
&= -\frac{\text{Subst}\left(\int \left(a^2\left(1-\frac{2b^2}{a^2}\right) + \frac{b^4}{ax} - ax + x^2 - \frac{(a^2-b^2)^2}{a(a+x)}\right) dx, x, b \csc(x)\right)}{b^4} \\
&= -\frac{(a^2-2b^2)\csc(x)}{b^3} + \frac{a\csc^2(x)}{2b^2} - \frac{\csc^3(x)}{3b} + \frac{(a^2-b^2)^2 \log(a+b \csc(x))}{ab^4} + \frac{\log(\sin(x))}{a}
\end{aligned}$$

Mathematica [A] time = 0.14, size = 85, normalized size = 1.18

$$\frac{3a^2b^2 \csc^2(x) - 6ab(a^2 - 2b^2)\csc(x) - 6a^2(a^2 - 2b^2)\log(\sin(x)) + 6(a^2 - b^2)^2 \log(a \sin(x) + b) - 2ab^3 \csc^3(x)}{6ab^4}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^5/(a + b*Csc[x]), x]`

[Out] $\frac{(-6*a*b*(a^2 - 2*b^2)*Csc[x] + 3*a^2*b^2*Csc[x]^2 - 2*a*b^3*Csc[x]^3 - 6*a^2*(a^2 - 2*b^2)*Log[Sin[x]] + 6*(a^2 - b^2)^2*Log[b + a*Sin[x]])}{(6*a*b^4)}$

fricas [B] time = 0.51, size = 151, normalized size = 2.10

$$\frac{3 a^2 b^2 \sin(x) - 6 a^3 b + 10 a b^3 + 6 (a^3 b - 2 a b^3) \cos(x)^2 + 6 (a^4 - 2 a^2 b^2 + b^4 - (a^4 - 2 a^2 b^2 + b^4) \cos(x)^2) \log(a \sin(x))}{6 (a b^4 \cos(x)^2 - a b^4) \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^5/(a+b*csc(x)), x, algorithm="fricas")`

[Out] $\frac{-1/6*(3*a^2*b^2*sin(x) - 6*a^3*b + 10*a*b^3 + 6*(a^3*b - 2*a*b^3)*cos(x)^2 + 6*(a^4 - 2*a^2*b^2 + b^4 - (a^4 - 2*a^2*b^2 + b^4)*cos(x)^2)*log(a*sin(x)) + b)*sin(x) - 6*(a^4 - 2*a^2*b^2 - (a^4 - 2*a^2*b^2)*cos(x)^2)*log(1/2*sin(x))*sin(x))/((a*b^4*cos(x)^2 - a*b^4)*sin(x))}{(a*b^4*cos(x)^2 - a*b^4)}$

giac [A] time = 0.39, size = 90, normalized size = 1.25

$$-\frac{(a^3 - 2 a b^2) \log(|\sin(x)|)}{b^4} + \frac{(a^4 - 2 a^2 b^2 + b^4) \log(|a \sin(x) + b|)}{a b^4} + \frac{3 a b^2 \sin(x) - 2 b^3 - 6 (a^2 b - 2 b^3) \sin(x)^2}{6 b^4 \sin(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^5/(a+b*csc(x)),x, algorithm="giac")`

[Out] $-\frac{(a^3 - 2a^2b^2)\ln(\sin(x))}{b^4} + \frac{(a^4 - 2a^2b^2 + b^4)\ln(\sin(x))}{b^4} + \frac{1}{6}(3a^2b^2\sin(x) - 2b^3 - 6(a^2b^2 - 2b^3)\sin(x))\ln(\sin(x))$

maple [A] time = 0.32, size = 100, normalized size = 1.39

$$\frac{a^3 \ln(b + a \sin(x))}{b^4} - \frac{2a \ln(b + a \sin(x))}{b^2} + \frac{\ln(b + a \sin(x))}{a} - \frac{1}{3b \sin(x)^3} - \frac{a^2}{b^3 \sin(x)} + \frac{2}{b \sin(x)} + \frac{a}{2b^2 \sin(x)^2} - \frac{a^3 \ln(b + a \sin(x))}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^5/(a+b*csc(x)),x)`

[Out] $\frac{1}{b^4}a^3 \ln(b + a \sin(x)) - \frac{2}{b^2}a^2 \ln(b + a \sin(x)) + \frac{1}{a}a \ln(b + a \sin(x)) - \frac{1}{3b} \sin(x)^3 - \frac{1}{b^3} \sin(x)^2 + \frac{2}{b} \sin(x) + \frac{2}{b^2} \sin(x)^2$

maxima [A] time = 0.31, size = 84, normalized size = 1.17

$$-\frac{(a^3 - 2ab^2)\log(\sin(x))}{b^4} + \frac{(a^4 - 2a^2b^2 + b^4)\log(a \sin(x) + b)}{ab^4} + \frac{3ab \sin(x) - 6(a^2 - 2b^2)\sin(x)^2 - 2b^2}{6b^3 \sin(x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^5/(a+b*csc(x)),x, algorithm="maxima")`

[Out] $-\frac{(a^3 - 2a^2b^2)\log(\sin(x))}{b^4} + \frac{(a^4 - 2a^2b^2 + b^4)\log(a \sin(x) + b)}{b^4} + \frac{1}{6}(3a^2b^2\sin(x) - 6(a^2b^2 - 2b^3)\sin(x)^2 - 2b^3)\ln(\sin(x))$

mupad [B] time = 0.57, size = 157, normalized size = 2.18

$$\tan\left(\frac{x}{2}\right) \left(\frac{7}{8b} - \frac{a^2}{2b^3} \right) - \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{\tan\left(\frac{x}{2}\right)^3}{24b} + \frac{a \tan\left(\frac{x}{2}\right)^2}{8b^2} - \frac{\tan\left(\frac{x}{2}\right)^2 (4a^2 - 7b^2) + \frac{b^2}{3} - ab \tan\left(\frac{x}{2}\right)}{8b^3 \tan\left(\frac{x}{2}\right)^3} + \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^5/(a + b/sin(x)),x)`

[Out] $\tan(x/2) * (7/(8*b) - a^2/(2*b^3)) - \log(\tan(x/2)^2 + 1)/a - \tan(x/2)^3/(24*b) + (a*tan(x/2)^2)/(8*b^2) - (\tan(x/2)^2*(4*a^2 - 7*b^2) + b^2/3 - a*b*tan(x/2)^2)/(8*b^3*tan(x/2)^3)$

$x/2))/(8*b^3*tan(x/2)^3) + (\log(tan(x/2))*(2*a*b^2 - a^3))/b^4 + (\log(b + 2*a*tan(x/2) + b*tan(x/2)^2)*(a^2 - b^2)^2)/(a*b^4)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot^5(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)**5/(a+b*csc(x)),x)
[Out] Integral(cot(x)**5/(a + b*csc(x)), x)

$$3.18 \quad \int \frac{\cot^7(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=122

$$\frac{(a^2 - b^2)^3 \log(a + b \csc(x))}{ab^6} + \frac{a(a^2 - 3b^2) \csc^2(x)}{2b^4} - \frac{(a^2 - 3b^2) \csc^3(x)}{3b^3} - \frac{(a^4 - 3a^2b^2 + 3b^4) \csc(x)}{b^5} + \frac{a \csc^4(x)}{4b^2} -$$

[Out] $-(a^4 - 3a^2b^2 + 3b^4) \csc(x)/b^5 + 1/2*a*(a^2 - 3b^2)*\csc(x)^2/b^4 - 1/3*(a^2 - 3b^2)*\csc(x)^3/b^3 + 1/4*a*\csc(x)^4/b^2 - 1/5*\csc(x)^5/b + (a^2 - b^2)^3 \ln(a + b \csc(x))/a - b^6 \ln(\sin(x))/a$

Rubi [A] time = 0.12, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.154, Rules used = {3885, 894}

$$-\frac{(a^2 - 3b^2) \csc^3(x)}{3b^3} + \frac{a(a^2 - 3b^2) \csc^2(x)}{2b^4} - \frac{(-3a^2b^2 + a^4 + 3b^4) \csc(x)}{b^5} + \frac{(a^2 - b^2)^3 \log(a + b \csc(x))}{ab^6} + \frac{a \csc^4(x)}{4b^2}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^7/(a + b*Csc[x]), x]

[Out] $-((a^4 - 3a^2b^2 + 3b^4)*\Csc[x])/b^5 + (a*(a^2 - 3b^2)*\Csc[x]^2)/(2*b^4) - ((a^2 - 3b^2)*\Csc[x]^3)/(3*b^3) + (a*\Csc[x]^4)/(4*b^2) - \Csc[x]^5/(5*b) + ((a^2 - b^2)^3 \Log[a + b \Csc[x]])/(a*b^6) - \Log[\Sin[x]]/a$

Rule 894

```
Int[((d_.) + (e_)*(x_))^m_*((f_.) + (g_)*(x_))^n_*((a_) + (c_)*(x_)^2)^p_, x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*(f + g*x)^n*(a + c*x)^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p] && ((EqQ[p, 1] && IntegerQ[m, n]) || (ILtQ[m, 0] && ILtQ[n, 0]))
```

Rule 3885

```
Int[cot[(c_.) + (d_)*(x_)]^m_*((c_.) + (d_)*(x_))*((b_.) + (a_))^n_, x_Symbol] :> -Dist[(-1)^((m - 1)/2)/(d*b^(m - 1)), Subst[Int[((b^2 - x^2)^((m - 1)/2)*(a + x)^n)/x, x], x, b*Csc[c + d*x]], x] /; FreeQ[{a, b, c, d, n}, x] && IntegerQ[(m - 1)/2] && NeQ[a^2 - b^2, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{\cot^7(x)}{a + b \csc(x)} dx &= \frac{\text{Subst}\left(\int \frac{(b^2-x^2)^3}{x(a+x)} dx, x, b \csc(x)\right)}{b^6} \\ &= \frac{\text{Subst}\left(\int \left(-a^4\left(1 + \frac{3b^2(-a^2+b^2)}{a^4}\right) + \frac{b^6}{ax} + a(a^2-3b^2)x - (a^2-3b^2)x^2 + ax^3 - x^4 + \frac{(a^2-b^2)^3}{a(a+x)}\right) dx, x, b \csc(x)\right)}{b^6} \\ &= -\frac{(a^4-3a^2b^2+3b^4)\csc(x)}{b^5} + \frac{a(a^2-3b^2)\csc^2(x)}{2b^4} - \frac{(a^2-3b^2)\csc^3(x)}{3b^3} + \frac{a\csc^4(x)}{4b^2} - \frac{\csc^5(x)}{5b} \end{aligned}$$

Mathematica [A] time = 0.28, size = 132, normalized size = 1.08

$$\frac{30ab^2(a^2-3b^2)\csc^2(x) + \frac{60(a^2-b^2)^3 \log(a \sin(x)+b)}{a} - 20b^3(a^2-3b^2)\csc^3(x) - 60b(a^4-3a^2b^2+3b^4)\csc(x) - 60a(a^4-3a^2b^2+3b^4)\csc(x) - 60a(a^4-3a^2b^2+3b^4)\csc(x)}{60b^6}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^7/(a + b*Csc[x]), x]`

[Out] $(-60*b*(a^4 - 3*a^2*b^2 + 3*b^4)*Csc[x] + 30*a*b^2*(a^2 - 3*b^2)*Csc[x]^2 - 20*b^3*(a^2 - 3*b^2)*Csc[x]^3 + 15*a*b^4*Csc[x]^4 - 12*b^5*Csc[x]^5 - 60*a*(a^4 - 3*a^2*b^2 + 3*b^4)*Log[Sin[x]] + (60*(a^2 - b^2)^3*Log[b + a*Sin[x]])/a)/(60*b^6)$

fricas [B] time = 0.56, size = 327, normalized size = 2.68

$$60a^5b - 160a^3b^3 + 132ab^5 + 60(a^5b - 3a^3b^3 + 3ab^5)\cos(x)^4 - 20(6a^5b - 17a^3b^3 + 15ab^5)\cos(x)^2 - 60(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\cos(x)^4 - 20(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\cos(x)^2 + 60(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\cos(x)^4 - 20(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\cos(x)^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^7/(a+b*csc(x)), x, algorithm="fricas")`

[Out] $-1/60*(60*a^5*b - 160*a^3*b^3 + 132*a*b^5 + 60*(a^5*b - 3*a^3*b^3 + 3*a*b^5)*cos(x)^4 - 20*(6*a^5*b - 17*a^3*b^3 + 15*a*b^5)*cos(x)^2 - 60*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cos(x)^4 - 20*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cos(x)^2 + 60*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cos(x)^4 - 20*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*cos(x)^2)*log(a*sin(x) + b)*sin(x) + 60*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 + (a^6 - 3*a^4*b^2 + 3*a^2*b^4)*cos(x)^4 - 20*(a^6 - 3*a^4*b^2 + 3*a^2*b^4)*cos(x)^2)*log(-1/2*sin(x))*sin(x) - 15*(2*a^4*b^2 - 5*a^2*b^4 - 2*(a^4*b^2 - 3*a^2*b^4)*cos(x)^2)*sin(x))/((a*b^6*cos(x)^4 - 2*a*b^6*cos(x)^2 + a*b^6)*sin(x))$

giac [A] time = 0.41, size = 155, normalized size = 1.27

$$-\frac{(a^5 - 3a^3b^2 + 3ab^4)\log(|\sin(x)|)}{b^6} + \frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\log(|a\sin(x) + b|)}{ab^6} + \frac{15ab^4\sin(x) - 12b^5 - 60}{1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^7/(a+b*csc(x)),x, algorithm="giac")

[Out] $-(a^5 - 3a^3b^2 + 3ab^4)\log(\sin(x))/b^6 + (a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\log(a\sin(x) + b)/(a*b^6) + 1/60*(15*a*b^4*\sin(x) - 12*b^5 - 60*(a^4*b - 3a^2*b^3 + 3b^5)*\sin(x)^4 + 30*(a^3*b^2 - 3*a*b^4)*\sin(x)^3 - 20*(a^2*b^3 - 3*b^5)*\sin(x)^2)/(b^6*\sin(x)^5)$

maple [A] time = 0.33, size = 181, normalized size = 1.48

$$\frac{a^5 \ln(b + a \sin(x))}{b^6} - \frac{3a^3 \ln(b + a \sin(x))}{b^4} + \frac{3a \ln(b + a \sin(x))}{b^2} - \frac{\ln(b + a \sin(x))}{a} - \frac{1}{5b \sin(x)^5} - \frac{a^2}{3b^3 \sin(x)^3} + \frac{1}{b \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cot(x)^7/(a+b*csc(x)),x)

[Out] $1/b^6*a^5*\ln(b+a*\sin(x))-3/b^4*a^3*\ln(b+a*\sin(x))+3/b^2*a*\ln(b+a*\sin(x))-1/a*\ln(b+a*\sin(x))-1/5/b/\sin(x)^5-1/3/b^3/\sin(x)^3*a^2+1/b/\sin(x)^3-1/b^5/\sin(x)*a^4+3/b^3/\sin(x)*a^2-3/b/\sin(x)+1/4*a/b^2/\sin(x)^4+1/2/b^4*a^3/\sin(x)^2-3/2*a/b^2/\sin(x)^2-1/b^6*a^5*\ln(\sin(x))+3/b^4*a^3*\ln(\sin(x))-3*a/b^2*\ln(\sin(x))$

maxima [A] time = 0.32, size = 149, normalized size = 1.22

$$-\frac{(a^5 - 3a^3b^2 + 3ab^4)\log(\sin(x))}{b^6} + \frac{(a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\log(a\sin(x) + b)}{ab^6} + \frac{15ab^3\sin(x) - 60(a^4 - 3a^2b^2 + 3b^4)\sin(x)^4 - 12b^4\sin(x)^3 + 30(a^3b^2 - 3a^2b^3)\sin(x)^2 - 20(a^2b^2 - 3b^4)\sin(x)^2}{(b^5\sin(x)^5)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^7/(a+b*csc(x)),x, algorithm="maxima")

[Out] $-(a^5 - 3a^3b^2 + 3ab^4)\log(\sin(x))/b^6 + (a^6 - 3a^4b^2 + 3a^2b^4 - b^6)\log(a\sin(x) + b)/(a*b^6) + 1/60*(15*a*b^4*\sin(x) - 60*(a^4 - 3a^2b^2 + 3b^4)*\sin(x)^4 - 12*b^4\sin(x)^3 + 30*(a^3b^2 - 3a^2b^3)*\sin(x)^2 - 20*(a^2b^2 - 3b^4)*\sin(x)^2)/(b^5*\sin(x)^5)$

mupad [B] time = 0.87, size = 284, normalized size = 2.33

$$\tan\left(\frac{x}{2}\right)^3 \left(\frac{3}{32b} - \frac{a^2}{24b^3} \right) - \frac{19\tan\left(\frac{x}{2}\right)}{16b} + \frac{\ln\left(\tan\left(\frac{x}{2}\right)^2 + 1\right)}{a} - \frac{\tan\left(\frac{x}{2}\right)^5}{160b} - \tan\left(\frac{x}{2}\right)^2 \left(\frac{a}{32b^2} + \frac{a\left(\frac{9}{32b} - \frac{a^2}{8b^3}\right)}{b} \right) + \frac{11a^2\tan\left(\frac{x}{2}\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^7/(a + b/sin(x)),x)`

[Out] $\tan(x/2)^3*(3/(32*b) - a^2/(24*b^3)) - (19*tan(x/2))/(16*b) + \log(\tan(x/2)^2 + 1)/a - \tan(x/2)^5/(160*b) - \tan(x/2)^2*(a/(32*b^2) + (a*(9/(32*b) - a^2/(8*b^3))/b) + (11*a^2*tan(x/2))/(8*b^3) + (a*tan(x/2)^4)/(64*b^2) - (a^4*tan(x/2))/(2*b^5) - (\log(\tan(x/2))*(3*a*b^4 + a^5 - 3*a^3*b^2))/b^6 - (\cot(x/2)^5*(\tan(x/2)^3*(10*a*b^3 - 4*a^3*b) - \tan(x/2)^2*(3*b^4 - (4*a^2*b^2)/3) + \tan(x/2)^4*(16*a^4 + 38*b^4 - 44*a^2*b^2) + b^4/5 - (a*b^3*tan(x/2))/2))/(32*b^5) + (\log(b + 2*a*tan(x/2) + b*tan(x/2)^2)*(a^2 - b^2)^3)/(a*b^6)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot^7(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)**7/(a+b*csc(x)),x)`

[Out] `Integral(cot(x)**7/(a + b*csc(x)), x)`

$$3.19 \quad \int \frac{\tan^4(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=159

$$\frac{a \tan^3(x)}{3(a^2 - b^2)} + \frac{ab^2 \tan(x)}{(a^2 - b^2)^2} - \frac{a \tan(x)}{a^2 - b^2} - \frac{b \sec^3(x)}{3(a^2 - b^2)} + \frac{b \sec(x)}{a^2 - b^2} + \frac{2b^5 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a(a^2-b^2)^{5/2}} - \frac{b^3 \sec(x)}{(a^2-b^2)^2} + \frac{x}{a}$$

[Out] $x/a+2*b^5*arctanh((a+b*tan(1/2*x))/(a^2-b^2)^(1/2))/a/(a^2-b^2)^(5/2)-b^3*s\ec(x)/(a^2-b^2)^2+b*\sec(x)/(a^2-b^2)-1/3*b*\sec(x)^3/(a^2-b^2)+a*b^2*tan(x)/(a^2-b^2)^2-a*tan(x)/(a^2-b^2)+1/3*a*tan(x)^3/(a^2-b^2)$

Rubi [A] time = 0.32, antiderivative size = 205, normalized size of antiderivative = 1.29, number of steps used = 16, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.692, Rules used = {3898, 2902, 2606, 3473, 8, 2735, 2660, 618, 206}

$$\frac{b^4 x}{a(a^2 - b^2)^2} - \frac{ab^2 x}{(a^2 - b^2)^2} + \frac{ax}{a^2 - b^2} + \frac{a \tan^3(x)}{3(a^2 - b^2)} + \frac{ab^2 \tan(x)}{(a^2 - b^2)^2} - \frac{a \tan(x)}{a^2 - b^2} - \frac{b \sec^3(x)}{3(a^2 - b^2)} - \frac{b^3 \sec(x)}{(a^2 - b^2)^2} + \frac{b \sec(x)}{a^2 - b^2} + \frac{2b^5 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a(a^2-b^2)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]^4/(a + b*Csc[x]), x]

[Out] $-((a*b^2*x)/(a^2 - b^2)^2) + (b^4*x)/(a*(a^2 - b^2)^2) + (a*x)/(a^2 - b^2) + (2*b^5*ArcTanh[(a + b*Tan[x/2])/Sqrt[a^2 - b^2]])/(a*(a^2 - b^2)^(5/2)) - (b^3*Sec[x])/(a^2 - b^2)^2 + (b*Sec[x])/(a^2 - b^2) - (b*Sec[x]^3)/(3*(a^2 - b^2)) + (a*b^2*Tan[x])/(a^2 - b^2)^2 - (a*Tan[x])/(a^2 - b^2) + (a*Tan[x]^3)/(3*(a^2 - b^2))$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 206

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 618

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},

$x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 2606

```
Int[((a_)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.*tan[(e_.) + (f_.*(x_))])^(n_.), x_Symbol] :> Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^(n - 1)/2], x], x, Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Rule 2660

```
Int[((a_) + (b_.*sin[(c_.) + (d_.*(x_))])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*x^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2735

```
Int[((a_) + (b_.*sin[(e_.) + (f_.*(x_))])/((c_.) + (d_.*sin[(e_.) + (f_.*(x_))]), x_Symbol] :> Simp[(b*x)/d, x] - Dist[(b*c - a*d)/d, Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

Rule 2902

```
Int[((cos[(e_.) + (f_.*(x_))]*(g_.*(p_))*((d_.*sin[(e_.) + (f_.*(x_))])^(n_))/((a_) + (b_.*sin[(e_.) + (f_.*(x_))]), x_Symbol] :> Dist[(a*d^2)/(a^2 - b^2), Int[(g*Cos[e + f*x])^p*(d*Sin[e + f*x])^(n - 2), x], x] + (-Dist[(b*d)/(a^2 - b^2), Int[(g*Cos[e + f*x])^p*(d*Sin[e + f*x])^(n - 1), x], x] - Dist[(a^2*d^2)/(g^2*(a^2 - b^2)), Int[((g*Cos[e + f*x])^(p + 2)*(d*Sin[e + f*x])^(n - 2))/(a + b*Sin[e + f*x]), x], x]) /; FreeQ[{a, b, d, e, f, g}, x] && NeQ[a^2 - b^2, 0] && IntegersQ[2*n, 2*p] && LtQ[p, -1] && GtQ[n, 1]
```

Rule 3473

```
Int[((b_.*tan[(c_.) + (d_.*(x_))])^(n_), x_Symbol] :> Simp[(b*(b*Tan[c + d*x])^(n - 1))/(d*(n - 1)), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3898

```
Int[cot[(c_.) + (d_.*(x_))^(m_.)*(csc[(c_.) + (d_.*(x_))]*(b_.) + (a_.)^(n_)), x_Symbol] :> Int[(Cos[c + d*x]^m*(b + a*Sin[c + d*x])^n)/Sin[c + d*x]^(m + n), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && IntegerQ[n] && IntegerQ[m] && (IntegerQ[m/2] || LeQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan^4(x)}{a + b \csc(x)} dx &= \int \frac{\sin(x) \tan^4(x)}{b + a \sin(x)} dx \\
&= \frac{a \int \tan^4(x) dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan^3(x) dx}{a^2 - b^2} + \frac{b^2 \int \frac{\sin(x) \tan^2(x)}{b+a \sin(x)} dx}{a^2 - b^2} \\
&= \frac{a \tan^3(x)}{3(a^2 - b^2)} + \frac{(ab^2) \int \tan^2(x) dx}{(a^2 - b^2)^2} - \frac{b^3 \int \sec(x) \tan(x) dx}{(a^2 - b^2)^2} + \frac{b^4 \int \frac{\sin(x)}{b+a \sin(x)} dx}{(a^2 - b^2)^2} - \frac{a \int \tan^2(x) dx}{a^2 - b^2} \\
&= \frac{b^4 x}{a(a^2 - b^2)^2} + \frac{b \sec(x)}{a^2 - b^2} - \frac{b \sec^3(x)}{3(a^2 - b^2)} + \frac{ab^2 \tan(x)}{(a^2 - b^2)^2} - \frac{a \tan(x)}{a^2 - b^2} + \frac{a \tan^3(x)}{3(a^2 - b^2)} - \frac{(ab^2) \int 1 dx}{(a^2 - b^2)^2} \\
&= -\frac{ab^2 x}{(a^2 - b^2)^2} + \frac{b^4 x}{a(a^2 - b^2)^2} + \frac{ax}{a^2 - b^2} - \frac{b^3 \sec(x)}{(a^2 - b^2)^2} + \frac{b \sec(x)}{a^2 - b^2} - \frac{b \sec^3(x)}{3(a^2 - b^2)} + \frac{ab^2 \tan(x)}{(a^2 - b^2)^2} - \\
&= -\frac{ab^2 x}{(a^2 - b^2)^2} + \frac{b^4 x}{a(a^2 - b^2)^2} + \frac{ax}{a^2 - b^2} - \frac{b^3 \sec(x)}{(a^2 - b^2)^2} + \frac{b \sec(x)}{a^2 - b^2} - \frac{b \sec^3(x)}{3(a^2 - b^2)} + \frac{ab^2 \tan(x)}{(a^2 - b^2)^2} - \\
&= -\frac{ab^2 x}{(a^2 - b^2)^2} + \frac{b^4 x}{a(a^2 - b^2)^2} + \frac{ax}{a^2 - b^2} + \frac{2b^5 \tanh^{-1}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{a(a^2-b^2)^{5/2}} - \frac{b^3 \sec(x)}{(a^2 - b^2)^2} + \frac{b \sec(x)}{a^2 - b^2} -
\end{aligned}$$

Mathematica [A] time = 1.07, size = 194, normalized size = 1.22

$$\csc(x)(a \sin(x) + b) \left(\frac{\sec^3(x) (-4a^4 \sin(3x) + 3a^4 x \cos(3x) + 2a^3 b + 3a^2 b^2 \sin(x) + 7a^2 b^2 \sin(3x) - 6a^2 b^2 x \cos(3x) + 6ab(a^2 - 2b^2) \cos(2x) + 9x(a^2 - b^2) \cos(x))}{(a-b)^2(a+b)^2} \right) + 12a(a + b \csc(x))$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]^4/(a + b*Csc[x]), x]`

[Out] `(Csc[x]*(b + a*Sin[x])*((-24*b^5*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]])/(-a^2 + b^2)^(5/2) + (Sec[x]^3*(2*a^3*b - 8*a*b^3 + 9*(a^2 - b^2)^2*x*Cos[x] + 6*a*b*(a^2 - 2*b^2)*Cos[2*x] + 3*a^4*x*Cos[3*x] - 6*a^2*b^2*x*Cos[3*x]))/((a - b)^2*(a + b)^2))`

$$+ 3*b^4*x*Cos[3*x] + 3*a^2*b^2*Sin[x] - 4*a^4*Sin[3*x] + 7*a^2*b^2*Sin[3*x]))/((a - b)^2*(a + b)^2))/((12*a*(a + b)*Csc[x]))$$

fricas [A] time = 0.58, size = 483, normalized size = 3.04

$$\left[\frac{3 \sqrt{a^2 - b^2} b^5 \cos(x)^3 \log\left(\frac{(a^2 - b^2) \cos(x)^2 + 2 ab \sin(x) + a^2 + b^2 + 2 (b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2 ab \sin(x) - a^2 - b^2}\right) - 2 a^5 b + 4 a^3 b^3 - 2 a b^5 + 6}{6 (a^7 -$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tan(x)^4/(a+b*csc(x)),x, algorithm="fricas")

$$\begin{aligned} \text{[Out]} \quad & [1/6*(3*sqrt(a^2 - b^2)*b^5*cos(x)^3*log(((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) - 2*a^5*b + 4*a^3*b^3 - 2*a*b^5 + 6*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*x*cos(x)^3 + 6*(a^5*b - 3*a^3*b^3 + 2*a*b^5)*cos(x)^2 + 2*(a^6 - 2*a^4*b^2 + a^2*b^4 - (4*a^6 - 11*a^4*b^2 + 7*a^2*b^4)*cos(x)^2)*sin(x))/((a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6)*cos(x)^3), 1/3*(3*sqrt(-a^2 + b^2)*b^5*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/(a^2 - b^2)*cos(x)^3 - a^5*b + 2*a^3*b^3 - a*b^5 + 3*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*x*cos(x)^3 + 3*(a^5*b - 3*a^3*b^3 + 2*a*b^5)*cos(x)^2 + (a^6 - 2*a^4*b^2 + a^2*b^4 - (4*a^6 - 11*a^4*b^2 + 7*a^2*b^4)*cos(x)^2)*sin(x))/((a^7 - 3*a^5*b^2 + 3*a^3*b^4 - a*b^6)*cos(x)^3)] \end{aligned}$$

giac [A] time = 0.59, size = 216, normalized size = 1.36

$$\begin{aligned} & -\frac{2 \left(\pi \left\lfloor \frac{x}{2\pi} + \frac{1}{2} \right\rfloor \operatorname{sgn}(b) + \arctan\left(\frac{b \tan\left(\frac{1}{2}x\right) + a}{\sqrt{-a^2 + b^2}}\right) \right) b^5}{(a^5 - 2 a^3 b^2 + a b^4) \sqrt{-a^2 + b^2}} + \frac{x}{a} + \frac{2 \left(3 a^3 \tan\left(\frac{1}{2}x\right)^5 - 6 a b^2 \tan\left(\frac{1}{2}x\right)^5 + 3 b^3 \tan\left(\frac{1}{2}x\right)^4 - 10 a^2 b^2 \tan\left(\frac{1}{2}x\right)^3 + 15 a b^4 \tan\left(\frac{1}{2}x\right)^2 - 10 a^5 \tan\left(\frac{1}{2}x\right) + 15 a^3 b^2 \tan\left(\frac{1}{2}x\right) - 15 a^2 b^3 \tan\left(\frac{1}{2}x\right) + 15 a b^5 \tan\left(\frac{1}{2}x\right) - 15 b^6 \tan\left(\frac{1}{2}x\right) + 10 a^7 + 10 a^5 b^2 + 10 a^3 b^4 + 10 a b^6 + b^8 \right)}{6 a^7} \end{aligned}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(tan(x)^4/(a+b*csc(x)),x, algorithm="giac")

$$\begin{aligned} \text{[Out]} \quad & -2*(\pi*\operatorname{floor}(1/2*x/\pi + 1/2)*\operatorname{sgn}(b) + \arctan((b*\tan(1/2*x) + a)/\sqrt{-a^2 + b^2})*b^5)/((a^5 - 2*a^3*b^2 + a*b^4)*\sqrt{-a^2 + b^2}) + x/a + 2/3*(3*a^3 * \tan(1/2*x)^5 - 6*a*b^2*\tan(1/2*x)^5 + 3*b^3*\tan(1/2*x)^4 - 10*a^3*\tan(1/2*x)^3 + 16*a*b^2*\tan(1/2*x)^3 + 6*a^2*b*\tan(1/2*x)^2 - 12*b^3*\tan(1/2*x)^2 + 3*a^3*\tan(1/2*x) - 6*a*b^2*\tan(1/2*x) - 2*a^2*b + 5*b^3)/((a^4 - 2*a^2*b^2 + b^4)*(tan(1/2*x)^2 - 1)^3) \end{aligned}$$

maple [A] time = 0.42, size = 210, normalized size = 1.32

$$-\frac{2b^5 \arctan\left(\frac{2 \tan\left(\frac{x}{2}\right) b+2 a}{2 \sqrt{-a^2+b^2}}\right)}{(a-b)^2 (a+b)^2 a \sqrt{-a^2+b^2}} - \frac{64}{3 \left(\tan\left(\frac{x}{2}\right)-1\right)^3 (64 a+64 b)} - \frac{32}{(64 a+64 b) \left(\tan\left(\frac{x}{2}\right)-1\right)^2} + \frac{a}{(a+b)^2 \left(\tan\left(\frac{x}{2}\right)-1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^4/(a+b*csc(x)),x)`

[Out] `-2/(a-b)^2/(a+b)^2*b^5/a/(-a^2+b^2)^(1/2)*arctan(1/2*(2*tan(1/2*x)*b+2*a)/(-a^2+b^2)^(1/2))-64/3/(\tan(1/2*x)-1)^3/(64*a+64*b)-32/(64*a+64*b)/(\tan(1/2*x)-1)^2+1/(a+b)^2/(\tan(1/2*x)-1)*a+3/2/(a+b)^2/(\tan(1/2*x)-1)*b-64/3/(\tan(1/2*x)+1)^3/(64*a-64*b)+32/(64*a-64*b)/(\tan(1/2*x)+1)^2-3/2/(a-b)^2/(\tan(1/2*x)+1)*b+1/(a-b)^2/(\tan(1/2*x)+1)*a+2/a*arctan(\tan(1/2*x))`

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^4/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details)Is $4*a^2-4*b^2$ positive or negative?

mupad [B] time = 7.38, size = 4468, normalized size = 28.10

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^4/(a + b/sin(x)),x)`

[Out] `((4*tan(x/2)^2*(a^2*b - 2*b^3))/(a^4 + b^4 - 2*a^2*b^2) - (2*(2*a^2*b - 5*b^3))/(3*(a^4 + b^4 - 2*a^2*b^2)) + (2*b^3*tan(x/2)^4)/(a^4 + b^4 - 2*a^2*b^2) - (4*a*tan(x/2)^3*(5*a^2 - 8*b^2))/(3*(a^4 + b^4 - 2*a^2*b^2)) + (2*a*tan(x/2)*(a^2 - 2*b^2))/(a^4 + b^4 - 2*a^2*b^2) + (2*a*tan(x/2)^5*(a^2 - 2*b^2))/(a^4 + b^4 - 2*a^2*b^2))/(3*tan(x/2)^2 - 3*tan(x/2)^4 + tan(x/2)^6 - 1) + (2*atan((320*a*b^21*tan(x/2)))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) + (64*a^21*b*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15)`

$$\begin{aligned}
& 15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3 \\
& 520*a^17*b^5 - 704*a^19*b^3) - (2560*a^3*b^19*tan(x/2))/(320*a*b^21 + 64*a^ \\
& 21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29 \\
& 504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19* \\
& b^3) + (9280*a^5*b^17*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9 \\
& 280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^ \\
& 13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) - (20160*a^7*b^15*t \\
& an(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^ \\
& 7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 \\
& + 3520*a^17*b^5 - 704*a^19*b^3) + (29184*a^9*b^13*tan(x/2))/(320*a*b^21 + \\
& 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 \\
& - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704* \\
& a^19*b^3) - (29504*a^11*b^11*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b \\
& ^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 2 \\
& 1120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) + (21120*a^1 \\
& 3*b^9*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 2 \\
& 0160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a \\
& ^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) - (10560*a^15*b^7*tan(x/2))/(320*a* \\
& b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a \\
& ^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 \\
& - 704*a^19*b^3) + (3520*a^17*b^5*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560* \\
& a^3*b^19 + 9280*a^5*b^17 - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^1 \\
& 1 + 21120*a^13*b^9 - 10560*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3) - (704* \\
& a^19*b^3*tan(x/2))/(320*a*b^21 + 64*a^21*b - 2560*a^3*b^19 + 9280*a^5*b^17 \\
& - 20160*a^7*b^15 + 29184*a^9*b^13 - 29504*a^11*b^11 + 21120*a^13*b^9 - 1056 \\
& 0*a^15*b^7 + 3520*a^17*b^5 - 704*a^19*b^3))/a + (b^5*atan(((b^5*((a + b)^5 \\
& *(a - b)^5)^(1/2)*(tan(x/2)*(64*a^22*b - 64*b^23 + 576*a^2*b^21 - 2560*a^4* \\
& b^19 + 7360*a^6*b^17 - 14880*a^8*b^15 + 21696*a^10*b^13 - 22880*a^12*b^11 + \\
& 17280*a^14*b^9 - 9120*a^16*b^7 + 3200*a^18*b^5 - 672*a^20*b^3) + 32*a*b^22 \\
& - 320*a^3*b^20 + 1440*a^5*b^18 - 3840*a^7*b^16 + 6720*a^9*b^14 - 8064*a^11* \\
& b^12 + 6720*a^13*b^10 - 3840*a^15*b^8 + 1440*a^17*b^6 - 320*a^19*b^4 + 32* \\
& a^21*b^2 + (b^5*((a + b)^5*(a - b)^5)^(1/2)*(32*a^23*b + 96*a^3*b^21 - 864* \\
& a^5*b^19 + 3488*a^7*b^17 - 8320*a^9*b^15 + 12992*a^11*b^13 - 13888*a^13*b^1 \\
& 1 + 10304*a^15*b^9 - 5248*a^17*b^7 + 1760*a^19*b^5 - 352*a^21*b^3 + tan(x/2) \\
&)*(64*a^2*b^22 - 512*a^4*b^20 + 1792*a^6*b^18 - 3584*a^8*b^16 + 4480*a^10*b \\
& ^14 - 3584*a^12*b^12 + 1792*a^14*b^10 - 512*a^16*b^8 + 64*a^18*b^6) + (b^5* \\
& ((a + b)^5*(a - b)^5)^(1/2)*(tan(x/2)*(96*a^24*b - 64*a^2*b^23 + 736*a^4*b^ \\
& 21 - 3840*a^6*b^19 + 12000*a^8*b^17 - 24960*a^10*b^15 + 36288*a^12*b^13 - 3 \\
& 7632*a^14*b^11 + 27840*a^16*b^9 - 14400*a^18*b^7 + 4960*a^20*b^5 - 1024*a^2* \\
& 2*b^3) + 32*a^3*b^22 - 320*a^5*b^20 + 1440*a^7*b^18 - 3840*a^9*b^16 + 6720* \\
& a^11*b^14 - 8064*a^13*b^12 + 6720*a^15*b^10 - 3840*a^17*b^8 + 1440*a^19*b^6 \\
& - 320*a^21*b^4 + 32*a^23*b^2)))/(a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10* \\
& a^7*b^4 + 5*a^9*b^2)*i)/(a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + \\
& 5*a^9*b^2) + (b^5*((a + b)^5*(a - b)^5)^(1/2)*(tan(x/2)*(64*a^22*b - 64*b^
\end{aligned}$$

$$\begin{aligned}
& 23 + 576*a^2*b^21 - 2560*a^4*b^19 + 7360*a^6*b^17 - 14880*a^8*b^15 + 21696*a^10*b^13 - 22880*a^12*b^11 + 17280*a^14*b^9 - 9120*a^16*b^7 + 3200*a^18*b^5 - 672*a^20*b^3) + 32*a*b^22 - 320*a^3*b^20 + 1440*a^5*b^18 - 3840*a^7*b^16 + 6720*a^9*b^14 - 8064*a^11*b^12 + 6720*a^13*b^10 - 3840*a^15*b^8 + 1440*a^17*b^6 - 320*a^19*b^4 + 32*a^21*b^2 - (b^5*((a + b)^5*(a - b)^5))^(1/2)*(32*a^23*b + 96*a^3*b^21 - 864*a^5*b^19 + 3488*a^7*b^17 - 8320*a^9*b^15 + 12992*a^11*b^13 - 13888*a^13*b^11 + 10304*a^15*b^9 - 5248*a^17*b^7 + 1760*a^19*b^5 - 352*a^21*b^3 + \tan(x/2)*(64*a^2*b^22 - 512*a^4*b^20 + 1792*a^6*b^18 - 3584*a^8*b^16 + 4480*a^10*b^14 - 3584*a^12*b^12 + 1792*a^14*b^10 - 512*a^16*b^8 + 64*a^18*b^6) - (b^5*((a + b)^5*(a - b)^5))^(1/2)*(\tan(x/2)*(96*a^24*b - 64*a^2*b^23 + 736*a^4*b^21 - 3840*a^6*b^19 + 12000*a^8*b^17 - 24960*a^10*b^15 + 36288*a^12*b^13 - 37632*a^14*b^11 + 27840*a^16*b^9 - 14400*a^18*b^7 + 4960*a^20*b^5 - 1024*a^22*b^3) + 32*a^3*b^22 - 320*a^5*b^20 + 1440*a^7*b^18 - 3840*a^9*b^16 + 6720*a^11*b^14 - 8064*a^13*b^12 + 6720*a^15*b^10 - 3840*a^17*b^8 + 1440*a^19*b^6 - 320*a^21*b^4 + 32*a^23*b^2))/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2)))^11i)/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2))/((2*tan(x/2)*(64*b^22 - 512*a^2*b^20 + 1792*a^4*b^18 - 3584*a^6*b^16 + 4480*a^8*b^14 - 3584*a^10*b^12 + 1792*a^12*b^10 - 512*a^14*b^8 + 64*a^16*b^6) - 128*a*b^21 + 832*a^3*b^19 - 2304*a^5*b^17 + 3520*a^7*b^15 - 3200*a^9*b^13 + 1728*a^11*b^11 - 512*a^13*b^9 + 64*a^15*b^7 + (b^5*((a + b)^5*(a - b)^5))^(1/2)*(\tan(x/2)*(64*a^22*b - 64*b^23 + 576*a^2*b^21 - 2560*a^4*b^19 + 7360*a^6*b^17 - 14880*a^8*b^15 + 21696*a^10*b^13 - 22880*a^12*b^11 + 17280*a^14*b^9 - 9120*a^16*b^7 + 3200*a^18*b^5 - 672*a^20*b^3) + 32*a*b^22 - 320*a^3*b^20 + 1440*a^5*b^18 - 3840*a^7*b^16 + 6720*a^9*b^14 - 8064*a^11*b^12 + 6720*a^13*b^10 - 3840*a^15*b^8 + 1440*a^17*b^6 - 320*a^19*b^4 + 32*a^21*b^2 + (b^5*((a + b)^5*(a - b)^5))^(1/2)*(32*a^23*b + 96*a^3*b^21 - 864*a^5*b^19 + 3488*a^7*b^17 - 8320*a^9*b^15 + 12992*a^11*b^13 - 13888*a^13*b^11 + 10304*a^15*b^9 - 5248*a^17*b^7 + 1760*a^19*b^5 - 352*a^21*b^3 + \tan(x/2)*(64*a^2*b^22 - 512*a^4*b^20 + 1792*a^6*b^18 - 3584*a^8*b^16 + 4480*a^10*b^14 - 3584*a^12*b^12 + 1792*a^14*b^10 - 512*a^16*b^8 + 64*a^18*b^6) + (b^5*((a + b)^5*(a - b)^5))^(1/2)*(\tan(x/2)*(96*a^24*b - 64*a^2*b^23 + 736*a^4*b^21 - 3840*a^6*b^19 + 12000*a^8*b^17 - 24960*a^10*b^15 + 36288*a^12*b^13 - 37632*a^14*b^11 + 27840*a^16*b^9 - 14400*a^18*b^7 + 4960*a^20*b^5 - 1024*a^22*b^3) + 32*a^3*b^22 - 320*a^5*b^20 + 1440*a^7*b^18 - 3840*a^9*b^16 + 6720*a^11*b^14 - 8064*a^13*b^12 + 6720*a^15*b^10 - 3840*a^17*b^8 + 1440*a^19*b^6 - 320*a^21*b^4 + 32*a^23*b^2))/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2)))^10i)/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2))/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2)))^10i)/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2) - (b^5*((a + b)^5*(a - b)^5))^(1/2)*(t an(x/2)*(64*a^22*b - 64*b^23 + 576*a^2*b^21 - 2560*a^4*b^19 + 7360*a^6*b^17 - 14880*a^8*b^15 + 21696*a^10*b^13 - 22880*a^12*b^11 + 17280*a^14*b^9 - 9120*a^16*b^7 + 3200*a^18*b^5 - 672*a^20*b^3) + 32*a*b^22 - 320*a^3*b^20 + 1440*a^5*b^18 - 3840*a^7*b^16 + 6720*a^9*b^14 - 8064*a^11*b^12 + 6720*a^13*b^10 - 3840*a^15*b^8 + 1440*a^17*b^6 - 320*a^19*b^4 + 32*a^21*b^2 - (b^5*((a + b)^5*(a - b)^5))^(1/2)*(\tan(x/2)*(64*a^22*b - 64*b^23 + 576*a^2*b^21 - 2560*a^4*b^19 + 7360*a^6*b^17 - 14880*a^8*b^15 + 21696*a^10*b^13 - 22880*a^12*b^11 + 17280*a^14*b^9 - 9120*a^16*b^7 + 3200*a^18*b^5 - 672*a^20*b^3) + 32*a*b^22 - 320*a^3*b^20 + 1440*a^5*b^18 - 3840*a^7*b^16 + 6720*a^9*b^14 - 8064*a^11*b^12 + 6720*a^13*b^10 - 3840*a^15*b^8 + 1440*a^17*b^6 - 320*a^19*b^4 + 32*a^21*b^2))/((a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2)))^10i)
\end{aligned}$$

$$\begin{aligned}
& + b)^5 * (a - b)^5 * (1/2) * (32*a^23*b + 96*a^3*b^21 - 864*a^5*b^19 + 3488*a^7*b^17 - 8320*a^9*b^15 + 12992*a^11*b^13 - 13888*a^13*b^11 + 10304*a^15*b^9 - 5248*a^17*b^7 + 1760*a^19*b^5 - 352*a^21*b^3 + \tan(x/2)*(64*a^2*b^22 - 512*a^4*b^20 + 1792*a^6*b^18 - 3584*a^8*b^16 + 4480*a^10*b^14 - 3584*a^12*b^12 + 1792*a^14*b^10 - 512*a^16*b^8 + 64*a^18*b^6) - (b^5*((a + b)^5*(a - b)^5)^{(1/2)} * (\tan(x/2)*(96*a^24*b - 64*a^2*b^23 + 736*a^4*b^21 - 3840*a^6*b^19 + 12000*a^8*b^17 - 24960*a^10*b^15 + 36288*a^12*b^13 - 37632*a^14*b^11 + 27840*a^16*b^9 - 14400*a^18*b^7 + 4960*a^20*b^5 - 1024*a^22*b^3) + 32*a^3*b^22 - 320*a^5*b^20 + 1440*a^7*b^18 - 3840*a^9*b^16 + 6720*a^11*b^14 - 8064*a^13*b^12 + 6720*a^15*b^10 - 3840*a^17*b^8 + 1440*a^19*b^6 - 320*a^21*b^4 + 32*a^23*b^2)) / (a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2)) / (a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2))) * ((a + b)^5 * (a - b)^5)^{(1/2)} * 2i) / (a*b^10 - a^11 - 5*a^3*b^8 + 10*a^5*b^6 - 10*a^7*b^4 + 5*a^9*b^2)
\end{aligned}$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tan^4(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tan(x)**4/(a+b*csc(x)),x)
[Out] Integral(tan(x)**4/(a + b*csc(x)), x)
```

$$3.20 \quad \int \frac{\tan^2(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=84

$$\frac{a \tan(x)}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} + \frac{2b^3 \tanh^{-1} \left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}} \right)}{a (a^2 - b^2)^{3/2}} - \frac{x}{a}$$

[Out] $-\frac{x}{a+2b^3 \operatorname{arctanh}((a+b \tan(1/2*x))/(a^2-b^2)^{(1/2)})/a/(a^2-b^2)^{(3/2)}-b*\sec(x)/(a^2-b^2)+a*\tan(x)/(a^2-b^2)}$

Rubi [A] time = 0.19, antiderivative size = 112, normalized size of antiderivative = 1.33, number of steps used = 10, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.692, Rules used = {3898, 2902, 2606, 8, 3473, 2735, 2660, 618, 206}

$$\frac{b^2 x}{a (a^2 - b^2)} - \frac{ax}{a^2 - b^2} + \frac{a \tan(x)}{a^2 - b^2} - \frac{b \sec(x)}{a^2 - b^2} + \frac{2b^3 \tanh^{-1} \left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}} \right)}{a (a^2 - b^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Tan[x]^2/(a + b*Csc[x]), x]

[Out] $-\frac{((a*x)/(a^2 - b^2)) + (b^2*x)/(a*(a^2 - b^2)) + (2*b^3*\operatorname{ArcTanh}[(a + b*\operatorname{Tan}[x/2])/Sqrt[a^2 - b^2]])/(a*(a^2 - b^2)^{(3/2)}) - (b*\operatorname{Sec}[x])/(a^2 - b^2) + (a*\operatorname{Tan}[x])/(a^2 - b^2)}$

Rule 8

Int[a_, x_Symbol] :> Simp[a*x, x] /; FreeQ[a, x]

Rule 206

Int[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 618

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^{-1}, x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 2606

```
Int[((a_)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.*tan[(e_.) + (f_.*(x_))])^(n_.), x_Symbol] :> Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^(n - 1)/2], x], x, Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Rule 2660

```
Int[((a_) + (b_.*sin[(c_.) + (d_.*(x_))])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2735

```
Int[((a_) + (b_.*sin[(e_.) + (f_.*(x_))])/((c_.) + (d_.*sin[(e_.) + (f_.*(x_))]), x_Symbol] :> Simp[(b*x)/d, x] - Dist[(b*c - a*d)/d, Int[1/(c + d*Sin[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c - a*d, 0]
```

Rule 2902

```
Int[((cos[(e_.) + (f_.*(x_))]*(g_.*(p_*((d_.*sin[(e_.) + (f_.*(x_))])^(n_))))/((a_) + (b_.*sin[(e_.) + (f_.*(x_))]), x_Symbol] :> Dist[(a*d^2)/(a^2 - b^2), Int[(g*Cos[e + f*x])^p*(d*Sin[e + f*x])^(n - 2), x], x] + (-Dist[(b*d)/(a^2 - b^2), Int[(g*Cos[e + f*x])^p*(d*Sin[e + f*x])^(n - 1), x], x] - Dist[(a^2*d^2)/(g^2*(a^2 - b^2)), Int[((g*Cos[e + f*x])^(p + 2)*(d*Sin[e + f*x])^(n - 2))/(a + b*Sin[e + f*x]), x], x]) /; FreeQ[{a, b, d, e, f, g}, x] && NeQ[a^2 - b^2, 0] && IntegersQ[2*n, 2*p] && LtQ[p, -1] && GtQ[n, 1]
```

Rule 3473

```
Int[((b_.*tan[(c_.) + (d_.*(x_))])^(n_), x_Symbol] :> Simp[(b*(b*Tan[c + d*x])^(n - 1))/(d*(n - 1)), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]
```

Rule 3898

```
Int[cot[(c_.) + (d_.*(x_))^(m_.)*(csc[(c_.) + (d_.*(x_))]*(b_.) + (a_.)^(n_)), x_Symbol] :> Int[(Cos[c + d*x]^m*(b + a*Sin[c + d*x])^n)/Sin[c + d*x]^(m + n), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && IntegerQ[n] && IntegerQ[m] && (IntegerQ[m/2] || LeQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int \frac{\tan^2(x)}{a + b \csc(x)} dx &= \int \frac{\sin(x) \tan^2(x)}{b + a \sin(x)} dx \\
&= \frac{a \int \tan^2(x) dx}{a^2 - b^2} - \frac{b \int \sec(x) \tan(x) dx}{a^2 - b^2} + \frac{b^2 \int \frac{\sin(x)}{b+a \sin(x)} dx}{a^2 - b^2} \\
&= \frac{b^2 x}{a(a^2 - b^2)} + \frac{a \tan(x)}{a^2 - b^2} - \frac{a \int 1 dx}{a^2 - b^2} - \frac{b \operatorname{Subst}(\int 1 dx, x, \sec(x))}{a^2 - b^2} - \frac{b^3 \int \frac{1}{b+a \sin(x)} dx}{a(a^2 - b^2)} \\
&= -\frac{ax}{a^2 - b^2} + \frac{b^2 x}{a(a^2 - b^2)} - \frac{b \sec(x)}{a^2 - b^2} + \frac{a \tan(x)}{a^2 - b^2} - \frac{(2b^3) \operatorname{Subst}\left(\int \frac{1}{b+2ax+bx^2} dx, x, \tan\left(\frac{x}{2}\right)\right)}{a(a^2 - b^2)} \\
&= -\frac{ax}{a^2 - b^2} + \frac{b^2 x}{a(a^2 - b^2)} - \frac{b \sec(x)}{a^2 - b^2} + \frac{a \tan(x)}{a^2 - b^2} + \frac{(4b^3) \operatorname{Subst}\left(\int \frac{1}{4(a^2-b^2)-x^2} dx, x, 2a + 2b \tan\left(\frac{x}{2}\right)\right)}{a(a^2 - b^2)} \\
&= -\frac{ax}{a^2 - b^2} + \frac{b^2 x}{a(a^2 - b^2)} + \frac{2b^3 \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a(a^2 - b^2)^{3/2}} - \frac{b \sec(x)}{a^2 - b^2} + \frac{a \tan(x)}{a^2 - b^2}
\end{aligned}$$

Mathematica [A] time = 0.27, size = 87, normalized size = 1.04

$$\frac{\sqrt{b^2 - a^2} \left(a^2(-x) + a^2 \tan(x) - ab \sec(x) + b^2 x\right) - 2b^3 \tan^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}}\right)}{a(b^2 - a^2)^{3/2}}$$

Antiderivative was successfully verified.

[In] `Integrate[Tan[x]^2/(a + b*Csc[x]), x]`

[Out] $-\left((-2*b^3*\operatorname{ArcTan}\left((a + b*Tan[x/2])/Sqrt[-a^2 + b^2]\right))/Sqrt[-a^2 + b^2]\right) + Sqrt[-a^2 + b^2]*(-(a^2*x) + b^2*x - a*b*Sec[x] + a^2*Tan[x]))/(a*(-a^2 + b^2)^(3/2))$

fricas [A] time = 0.47, size = 296, normalized size = 3.52

$$\frac{\sqrt{a^2 - b^2} b^3 \cos(x) \log\left(-\frac{(a^2 - 2 b^2) \cos(x)^2 + 2 a b \sin(x) + a^2 + b^2 - 2 (b \cos(x) \sin(x) + a \cos(x)) \sqrt{a^2 - b^2}}{a^2 \cos(x)^2 - 2 a b \sin(x) - a^2 - b^2}\right) + 2 a^3 b - 2 a b^3 + 2 (a^4 - 2 a^2 b^2 + b^4) \cos(x)}{2 (a^5 - 2 a^3 b^2 + a b^4) \cos(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^2/(a+b*csc(x)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [-1/2 * (\sqrt{a^2 - b^2}) * b^3 * \cos(x) * \log(-((a^2 - 2*b^2) * \cos(x)^2 + 2*a*b*\sin(x) + a^2 + b^2 - 2*(b*\cos(x)*\sin(x) + a*\cos(x)) * \sqrt{a^2 - b^2})) / (a^2 * \cos(x)^2 - 2*a*b*\sin(x) - a^2 - b^2)) + 2*a^3*b - 2*a*b^3 + 2*(a^4 - 2*a^2*b^2 + b^4) * x * \cos(x) - 2*(a^4 - a^2*b^2) * \sin(x)) / ((a^5 - 2*a^3*b^2 + a*b^4) * \cos(x)), \\ & (\sqrt{-a^2 + b^2}) * b^3 * \arctan(-\sqrt{-a^2 + b^2}) * (b * \sin(x) + a) / ((a^2 - b^2) * \cos(x))) * \cos(x) - a^3*b + a*b^3 - (a^4 - 2*a^2*b^2 + b^4) * x * \cos(x) + (a^4 - a^2*b^2) * \sin(x)) / ((a^5 - 2*a^3*b^2 + a*b^4) * \cos(x))] \end{aligned}$$

giac [A] time = 0.36, size = 103, normalized size = 1.23

$$-\frac{2 \left(\pi \left[\frac{x}{2\pi} + \frac{1}{2} \right] \operatorname{sgn}(b) + \arctan \left(\frac{b \tan \left(\frac{1}{2} x \right) + a}{\sqrt{-a^2 + b^2}} \right) \right) b^3}{(a^3 - ab^2) \sqrt{-a^2 + b^2}} - \frac{x}{a} - \frac{2 \left(a \tan \left(\frac{1}{2} x \right) - b \right)}{(a^2 - b^2) \left(\tan \left(\frac{1}{2} x \right)^2 - 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^2/(a+b*csc(x)),x, algorithm="giac")`

[Out]
$$\begin{aligned} & -2 * (\pi * \operatorname{floor}(1/2*x/\pi + 1/2) * \operatorname{sgn}(b) + \arctan(b * \tan(1/2*x) + a) / \sqrt{-a^2 + b^2})) * b^3 / ((a^3 - a*b^2) * \sqrt{-a^2 + b^2}) - x/a - 2 * (a * \tan(1/2*x) - b) / ((a^2 - b^2) * (\tan(1/2*x)^2 - 1)) \end{aligned}$$

maple [A] time = 0.52, size = 106, normalized size = 1.26

$$-\frac{16}{(16a - 16b) \left(\tan \left(\frac{x}{2} \right) + 1 \right)} - \frac{2b^3 \arctan \left(\frac{2 \tan \left(\frac{x}{2} \right) b + 2a}{2\sqrt{-a^2 + b^2}} \right)}{(a - b)(a + b)a\sqrt{-a^2 + b^2}} - \frac{16}{(16a + 16b) \left(\tan \left(\frac{x}{2} \right) - 1 \right)} - \frac{2 \arctan \left(\tan \left(\frac{x}{2} \right) \right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(tan(x)^2/(a+b*csc(x)),x)`

[Out]
$$\begin{aligned} & -16 / (16*a - 16*b) / (\tan(1/2*x) + 1) - 2 / (a - b) / (a + b) * b^3 / a / (-a^2 + b^2)^{(1/2)} * \arctan(1/2 * (2 * \tan(1/2*x) * b + 2*a) / (-a^2 + b^2)^{(1/2)}) - 16 / (16*a + 16*b) / (\tan(1/2*x) - 1) - 2 / a * \arctan(\tan(1/2*x)) \end{aligned}$$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(tan(x)^2/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details) Is $4*a^2-4*b^2$ positive or negative?

mupad [B] time = 2.26, size = 2341, normalized size = 27.87

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int \tan(x)^2 / (a + b/\sin(x)) \, dx$

[Out]
$$\begin{aligned} & ((2*b)/(a^2 - b^2) - (2*a*tan(x/2))/(a^2 - b^2))/(tan(x/2)^2 - 1) + (2*atan((64*a^11*b*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) - (192*a*b^11*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) + (768*a^3*b^9*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) - (1216*a^5*b^7*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) + (960*a^7*b^5*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3) - (384*a^9*b^3*tan(x/2))/(192*a*b^11 - 64*a^11*b - 768*a^3*b^9 + 1216*a^5*b^7 - 960*a^7*b^5 + 384*a^9*b^3))/a + (b^3*atan(((b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(64*a^12*b + 64*b^13 - 320*a^2*b^11 + 736*a^4*b^9 - 992*a^6*b^7 + 800*a^8*b^5 - 352*a^10*b^3) - 32*a*b^12 + 160*a^3*b^10 - 320*a^5*b^8 + 320*a^7*b^6 - 160*a^9*b^4 + 32*a^11*b^2 + (b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(64*a^2*b^12 - 256*a^4*b^10 + 384*a^6*b^8 - 256*a^8*b^6 + 64*a^10*b^4) - 32*a^13*b + 64*a^3*b^11 - 288*a^5*b^9 + 512*a^7*b^7 - 448*a^9*b^5 + 192*a^11*b^3 + (b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(96*a^14*b + 64*a^2*b^13 - 416*a^4*b^11 + 1120*a^6*b^9 - 1600*a^8*b^7 + 1280*a^10*b^5 - 544*a^12*b^3) - 32*a^3*b^12 + 160*a^5*b^10 - 320*a^7*b^8 + 320*a^9*b^6 - 160*a^11*b^4 + 32*a^13*b^2))/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)))/((a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)*1i)/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2) + (b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(64*a^12*b + 64*b^13 - 320*a^2*b^11 + 736*a^4*b^9 - 992*a^6*b^7 + 800*a^8*b^5 - 352*a^10*b^3) - 32*a*b^12 + 160*a^3*b^10 - 320*a^5*b^8 + 320*a^7*b^6 - 160*a^9*b^4 + 32*a^11*b^2 - (b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(64*a^2*b^12 - 256*a^4*b^10 + 384*a^6*b^8 - 256*a^8*b^6 + 64*a^10*b^4) - 32*a^13*b + 64*a^3*b^11 - 288*a^5*b^9 + 512*a^7*b^7 - 448*a^9*b^5 + 192*a^11*b^3 - (b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(96*a^14*b + 64*a^2*b^13 - 416*a^4*b^11 + 1120*a^6*b^9 - 1600*a^8*b^7 + 1280*a^10*b^5 - 544*a^12*b^3) - 32*a^3*b^12 + 160*a^5*b^10 - 320*a^7*b^8 + 320*a^9*b^6 - 160*a^11*b^4 + 32*a^13*b^2))/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)))/((a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)*1i)/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2))/((2*tan(x/2)*(64*b^12 - 256*a^2*b^10 + 384*a^4*b^8 - 256*a^6*b^6 + 64*a^8*b^4) - 64*a*b^11 + 192*a^3*b^9 - 192*a^5*b^7 + 64*a^7*b^5 + (b^3*((a + b)^3*(a - b)^3)^{(1/2)}*(tan(x/2)*(64*a^12*b + 64*b^13 - 416*a^4*b^11 + 1120*a^6*b^9 - 1600*a^8*b^7 + 1280*a^10*b^5 - 544*a^12*b^3) - 32*a^3*b^12 + 160*a^5*b^10 - 320*a^7*b^8 + 320*a^9*b^6 - 160*a^11*b^4 + 32*a^13*b^2)))/((a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)*1i)/(a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)) \end{aligned}$$

$$\begin{aligned}
& *b^{13} - 320*a^2*b^{11} + 736*a^4*b^9 - 992*a^6*b^7 + 800*a^8*b^5 - 352*a^{10}*b^3 \\
& - 32*a*b^{12} + 160*a^3*b^{10} - 320*a^5*b^8 + 320*a^7*b^6 - 160*a^9*b^4 + \\
& 32*a^{11}*b^2 + (b^3*((a+b)^3*(a-b)^3))^{(1/2)}*(\tan(x/2)*(64*a^2*b^{12} - 256 \\
& *a^4*b^{10} + 384*a^6*b^8 - 256*a^8*b^6 + 64*a^{10}*b^4) - 32*a^{13}*b + 64*a^3*b^11 \\
& - 288*a^5*b^9 + 512*a^7*b^7 - 448*a^9*b^5 + 192*a^{11}*b^3 + (b^3*((a+b)^3*(a-b)^3))^{(1/2)}*(\tan(x/2)*(96*a^{14}*b + 64*a^2*b^{13} - 416*a^4*b^{11} + 11 \\
& 20*a^6*b^9 - 1600*a^8*b^7 + 1280*a^{10}*b^5 - 544*a^{12}*b^3) - 32*a^3*b^{12} + 1 \\
& 60*a^5*b^{10} - 320*a^7*b^8 + 320*a^9*b^6 - 160*a^{11}*b^4 + 32*a^{13}*b^2)) / (a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)) \\
& / (a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2) - (b^3*((a+b)^3*(a-b)^3))^{(1/2)}*(\tan(x/2)*(64*a^{12}*b + 64*b^{13} - 320*a^2*b^{11} + 736*a^4*b^9 - 992*a^6*b^7 + \\
& 800*a^8*b^5 - 352*a^{10}*b^3) - 32*a*b^{12} + 160*a^3*b^{10} - 320*a^5*b^8 + 320*a^7*b^6 - 160*a^9*b^4 + 32*a^{11}*b^2 - (b^3*((a+b)^3*(a-b)^3))^{(1/2)}*(\tan(x/2)*(64*a^2*b^{12} - 256*a^4*b^{10} + 384*a^6*b^8 - 256*a^8*b^6 + 64*a^{10}*b^4) - 32*a^{13}*b + 64*a^3*b^{11} - 288*a^5*b^9 + 512*a^7*b^7 - 448*a^9*b^5 + 192 \\
& *a^{11}*b^3 - (b^3*((a+b)^3*(a-b)^3))^{(1/2)}*(\tan(x/2)*(96*a^{14}*b + 64*a^2*b^{13} - 416*a^4*b^{11} + 1120*a^6*b^9 - 1600*a^8*b^7 + 1280*a^{10}*b^5 - 544*a^{12}*b^3) - 32*a^3*b^{12} + 160*a^5*b^{10} - 320*a^7*b^8 + 320*a^9*b^6 - 160*a^{11}*b^4 + 32*a^{13}*b^2)) / (a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)) / (a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)) * ((a+b)^3 \\
& *(a-b)^3)^{(1/2)*2i} / (a*b^6 - a^7 - 3*a^3*b^4 + 3*a^5*b^2)
\end{aligned}$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\tan^2(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(tan(x)**2/(a+b*csc(x)),x)
[Out] Integral(tan(x)**2/(a + b*csc(x)), x)
```

3.21 $\int \frac{\cot^2(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=61

$$\frac{2\sqrt{a^2 - b^2} \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab} - \frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b}$$

[Out] $-\frac{x}{a} - \frac{\text{arctanh}(\cos(x))}{b} + \frac{2 \text{arctanh}\left(\frac{(a+b) \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab} - \frac{\tanh^{-1}(\cos(x))}{b}$

Rubi [A] time = 0.18, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.615, Rules used = {3894, 4051, 3770, 3919, 3831, 2660, 618, 206}

$$\frac{2\sqrt{a^2 - b^2} \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab} - \frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[x]^2/(a + b \text{Csc}[x]), x]$

[Out] $-\frac{x}{a} - \frac{\text{ArcTanh}[\cos(x)]}{b} + \frac{(2 \sqrt{a^2 - b^2}) \text{ArcTanh}\left(\frac{a + b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{a b}$

Rule 206

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 618

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2660

```
Int[((a_) + (b_)*sin[(c_) + (d_)*(x_)])^(-1), x_Symbol] :> With[{e = FreFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]]
```

Rule 3770

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rule 3831

```
Int[csc[(e_.) + (f_.*(x_)]/(csc[(e_.) + (f_.*(x_)]*(b_.) + (a_)), x_Symbo
l] :> Dist[1/b, Int[1/(1 + (a*Sin[e + f*x])/b), x], x] /; FreeQ[{a, b, e, f
}, x] && NeQ[a^2 - b^2, 0]
```

Rule 3894

```
Int[cot[(c_.) + (d_.*(x_)]^2*(csc[(c_.) + (d_.*(x_)]*(b_.) + (a_))^(n_),
x_Symbol] :> Int[(-1 + Csc[c + d*x]^2)*(a + b*Csc[c + d*x])^n, x] /; FreeQ[
{a, b, c, d, n}, x] && NeQ[a^2 - b^2, 0]
```

Rule 3919

```
Int[((csc[(e_.) + (f_.*(x_)]*(d_.) + (c_))/((csc[(e_.) + (f_.*(x_)]*(b_.) +
(a_)), x_Symbol] :> Simplify[(c*x)/a, x] - Dist[(b*c - a*d)/a, Int[Csc[e + f*x]
]/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && NeQ[b*c -
a*d, 0]
```

Rule 4051

```
Int[((A_.) + csc[(e_.) + (f_.*(x_)]^2*(C_))/((csc[(e_.) + (f_.*(x_)]*(b_.
) + (a_)), x_Symbol] :> Dist[C/b, Int[Csc[e + f*x], x], x] + Dist[1/b, Int[
(A*b - a*C*Csc[e + f*x])/(a + b*Csc[e + f*x]), x], x] /; FreeQ[{a, b, e, f,
A, C}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cot^2(x)}{a + b \csc(x)} dx &= \int \frac{-1 + \csc^2(x)}{a + b \csc(x)} dx \\
&= \frac{\int \csc(x) dx}{b} + \frac{\int \frac{-b-a \csc(x)}{a+b \csc(x)} dx}{b} \\
&= -\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b} - \left(\frac{a}{b} - \frac{b}{a} \right) \int \frac{\csc(x)}{a + b \csc(x)} dx \\
&= -\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b} - \frac{\left(\frac{a}{b} - \frac{b}{a} \right) \int \frac{1}{1 + \frac{a \sin(x)}{b}} dx}{b} \\
&= -\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b} - \frac{\left(2 \left(\frac{a}{b} - \frac{b}{a} \right) \right) \text{Subst} \left(\int \frac{1}{1 + \frac{2ax}{b} + x^2} dx, x, \tan \left(\frac{x}{2} \right) \right)}{b} \\
&= -\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b} + \frac{\left(4 \left(\frac{a}{b} - \frac{b}{a} \right) \right) \text{Subst} \left(\int \frac{1}{-4 \left(1 - \frac{a^2}{b^2} \right) - x^2} dx, x, \frac{2a}{b} + 2 \tan \left(\frac{x}{2} \right) \right)}{b} \\
&= -\frac{x}{a} - \frac{\tanh^{-1}(\cos(x))}{b} + \frac{2 \sqrt{a^2 - b^2} \tanh^{-1} \left(\frac{b \left(\frac{a}{b} + \tan \left(\frac{x}{2} \right) \right)}{\sqrt{a^2 - b^2}} \right)}{ab}
\end{aligned}$$

Mathematica [A] time = 0.06, size = 71, normalized size = 1.16

$$\frac{2 \sqrt{b^2 - a^2} \tan^{-1} \left(\frac{a+b \tan \left(\frac{x}{2} \right)}{\sqrt{b^2 - a^2}} \right) + a \log \left(\sin \left(\frac{x}{2} \right) \right) - a \log \left(\cos \left(\frac{x}{2} \right) \right) - bx}{ab}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^2/(a + b*Csc[x]), x]`

[Out] $(-\frac{(b*x) + 2*\sqrt{-a^2 + b^2}*\text{ArcTan}[(a + b*Tan[x/2])/\sqrt{-a^2 + b^2}]}{a*\text{Log}[\text{Cos}[x/2]] + a*\text{Log}[\text{Sin}[x/2]])/(a*b)$

fricas [A] time = 0.59, size = 206, normalized size = 3.38

$$\frac{2 b x + a \log \left(\frac{1}{2} \cos(x) + \frac{1}{2} \right) - a \log \left(-\frac{1}{2} \cos(x) + \frac{1}{2} \right) - \sqrt{a^2 - b^2} \log \left(\frac{(a^2 - 2 b^2) \cos(x)^2 + 2 a b \sin(x) + a^2 + b^2 + 2 (b \cos(x) \sin(x) - a b)}{a^2 \cos(x)^2 - 2 a b \sin(x) - a^2 - b^2} \right)}{2 a b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^2/(a+b*csc(x)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [-1/2*(2*b*x + a*log(1/2*cos(x) + 1/2) - a*log(-1/2*cos(x) + 1/2) - \sqrt{a^2 - b^2}*\log((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2))/(a*b), \\ & -1/2*(2*b*x + a*log(1/2*cos(x) + 1/2) - a*log(-1/2*cos(x) + 1/2) - 2*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a))/((a^2 - b^2)*cos(x)))/(a*b)] \end{aligned}$$

giac [A] time = 0.31, size = 80, normalized size = 1.31

$$-\frac{x}{a} + \frac{\log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{b} - \frac{2\left(\pi\left[\frac{x}{2\pi} + \frac{1}{2}\right]\operatorname{sgn}(b) + \arctan\left(\frac{b\tan\left(\frac{1}{2}x\right)+a}{\sqrt{-a^2+b^2}}\right)\right)(a^2 - b^2)}{\sqrt{-a^2 + b^2}ab}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^2/(a+b*csc(x)),x, algorithm="giac")`

[Out]
$$-x/a + \log(\operatorname{abs}(\tan(1/2*x)))/b - 2*(\pi*\operatorname{floor}(1/2*x/\pi + 1/2)*\operatorname{sgn}(b) + \arctan((b*\tan(1/2*x) + a)/\sqrt{-a^2 + b^2})*(a^2 - b^2)/(\sqrt{-a^2 + b^2}*a*b))$$

maple [A] time = 0.35, size = 105, normalized size = 1.72

$$-\frac{2a\arctan\left(\frac{2\tan\left(\frac{x}{2}\right)b+2a}{2\sqrt{-a^2+b^2}}\right)}{b\sqrt{-a^2+b^2}} + \frac{2b\arctan\left(\frac{2\tan\left(\frac{x}{2}\right)b+2a}{2\sqrt{-a^2+b^2}}\right)}{a\sqrt{-a^2+b^2}} + \frac{\ln\left(\tan\left(\frac{x}{2}\right)\right)}{b} - \frac{2\arctan\left(\tan\left(\frac{x}{2}\right)\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^2/(a+b*csc(x)),x)`

[Out]
$$\begin{aligned} & -2/b*a/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*tan(1/2*x)*b+2*a)/(-a^2+b^2)^{(1/2)})+2*b/a/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*tan(1/2*x)*b+2*a)/(-a^2+b^2)^{(1/2)})+1/b*\ln(\tan(1/2*x))-2/a*\arctan(\tan(1/2*x)) \end{aligned}$$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^2/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details) Is $4*a^2-4*b^2$ positive or negative?

mupad [B] time = 0.53, size = 697, normalized size = 11.43

$$\frac{2 \operatorname{atan}\left(\frac{64 b^3}{-64 a^3-64 \tan\left(\frac{x}{2}\right) a^2 b+64 a b^2+64 \tan\left(\frac{x}{2}\right) b^3}+\frac{64 a^3 \tan\left(\frac{x}{2}\right)}{-64 a^3-64 \tan\left(\frac{x}{2}\right) a^2 b+64 a b^2+64 \tan\left(\frac{x}{2}\right) b^3}-\frac{64 a^2 b}{-64 a^3-64 \tan\left(\frac{x}{2}\right) a^2 b+64 a b^2+64 \tan\left(\frac{x}{2}\right) b^3}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^2/(a + b/sin(x)),x)`

[Out] $(2*\operatorname{atan}((64*b^3)/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2)) + (64*a^3*tan(x/2))/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2))) - (64*a^2*b)/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2)) - (64*a*b^2*tan(x/2))/(64*b^3*tan(x/2) + 64*a*b^2 - 64*a^3 - 64*a^2*b*tan(x/2)))/a + \log(\tan(x/2))/b - (2*\operatorname{atanh}((512*a^4*(a^2 - b^2)^(1/2))/(256*a*b^4 - 64*b^5*tan(x/2) + 512*a^5 - 768*a^3*b^2 + 832*a^2*b^3*tan(x/2) + (1024*a^6*tan(x/2))/b - 1792*a^4*b*tan(x/2)) - (512*a^2*(a^2 - b^2)^(1/2))/(256*a*b^2 - 64*b^3*tan(x/2) - 768*a^3 + (512*a^5)/b^2 - (1792*a^4*tan(x/2))/b + (1024*a^6*tan(x/2))/b^3 + 832*a^2*b*tan(x/2)) + (64*b^2*(a^2 - b^2)^(1/2))/(256*a*b^2 - 64*b^3*tan(x/2) - 768*a^3 + (512*a^5)/b^2 - (1792*a^4*tan(x/2))/b + (1024*a^6*tan(x/2))/b^3 + 832*a^2*b*tan(x/2)) - (1280*a^3*tan(x/2)*(a^2 - b^2)^(1/2))/(256*a*b^3 - 64*b^4*tan(x/2) - 1792*a^4*tan(x/2) - 768*a^3*b + (512*a^5)/b + 832*a^2*b^2*tan(x/2) + (1024*a^6*tan(x/2))/b^2 + (1024*a^5*tan(x/2)*(a^2 - b^2)^(1/2))/(1024*a^6*tan(x/2) - 64*b^6*tan(x/2) + 256*a*b^5 + 512*a^5*b - 768*a^3*b^3 + 832*a^2*b^4*tan(x/2) - 1792*a^4*b^2*tan(x/2) + (320*a*b*tan(x/2)*(a^2 - b^2)^(1/2))/(256*a*b^2 - 64*b^3*tan(x/2) - 768*a^3 + (512*a^5)/b^2 - (1792*a^4*tan(x/2))/b + (1024*a^6*tan(x/2))/b^3 + 832*a^2*b*tan(x/2)))*(a^2 - b^2)^(1/2))/(a*b)$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot^2(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)**2/(a+b*csc(x)),x)`

[Out] `Integral(cot(x)**2/(a + b*csc(x)), x)`

3.22 $\int \frac{\cot^4(x)}{a+b \csc(x)} dx$

Optimal. Leaf size=91

$$-\frac{(2a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{2(a^2 - b^2)^{3/2} \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab^3} + \frac{a \cot(x)}{b^2} + \frac{x}{a} - \frac{\cot(x) \csc(x)}{2b}$$

[Out] $x/a - 1/2*(2*a^2 - 3*b^2)*\text{arctanh}(\cos(x))/b^3 + 2*(a^2 - b^2)^{3/2}*\text{arctanh}((a+b*\tan(1/2*x))/(a^2 - b^2)^{1/2})/a/b^3 + a*\cot(x)/b^2 - 1/2*\cot(x)*\csc(x)/b$

Rubi [A] time = 0.29, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.538, Rules used = {3898, 2893, 3057, 2660, 618, 206, 3770}

$$-\frac{(2a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{2(a^2 - b^2)^{3/2} \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab^3} + \frac{a \cot(x)}{b^2} + \frac{x}{a} - \frac{\cot(x) \csc(x)}{2b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\cot[x]^4/(a + b*\csc[x]), x]$

[Out] $x/a - ((2*a^2 - 3*b^2)*\text{ArcTanh}[\text{Cos}[x]])/(2*b^3) + (2*(a^2 - b^2)^{3/2}*\text{ArcTanh}[(a + b*\text{Tan}[x/2])/Sqrt[a^2 - b^2]])/(a*b^3) + (a*\cot[x])/b^2 - (\cot[x]*\csc[x])/(2*b)$

Rule 206

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1*ArcTanh[(Rt[-b, 2]*x)/Rt[a, 2]])/(Rt[a, 2]*Rt[-b, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 618

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 2660

```
Int[((a_) + (b_)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[
```

$a^2 - b^2, 0]$

Rule 2893

```
Int[cos[(e_.) + (f_.)*(x_)]^4*((d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((a_) +
(b_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Simp[(Cos[e + f*x]*(a + b
*Sin[e + f*x])^(m + 1)*(d*Sin[e + f*x])^(n + 1))/(a*d*f*(n + 1)), x] + (-Di
st[1/(a^2*d^2*(n + 1)*(n + 2)), Int[(a + b*Sin[e + f*x])^m*(d*Sin[e + f*x])
^(n + 2)*Simp[a^2*n*(n + 2) - b^2*(m + n + 2)*(m + n + 3) + a*b*m*Sin[e + f
*x] - (a^2*(n + 1)*(n + 2) - b^2*(m + n + 2)*(m + n + 4))*Sin[e + f*x]^2, x
], x], x] - Simp[(b*(m + n + 2)*Cos[e + f*x]*(a + b*Sin[e + f*x])^(m + 1)*(d
*Sin[e + f*x])^(n + 2))/(a^2*d^2*f*(n + 1)*(n + 2)), x]) /; FreeQ[{a, b, d
, e, f, m}, x] && NeQ[a^2 - b^2, 0] && (IGtQ[m, 0] || IntegersQ[2*m, 2*n])
&& !m < -1 && LtQ[n, -1] && (LtQ[n, -2] || EqQ[m + n + 4, 0])
```

Rule 3057

```
Int[((A_.) + (B_.)*sin[(e_.) + (f_.)*(x_)] + (C_.)*sin[(e_.) + (f_.)*(x_)]^
2)/(((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])*((c_) + (d_.)*sin[(e_.) + (f_.)
*(x_)])), x_Symbol] :> Simp[(C*x)/(b*d), x] + (Dist[(A*b^2 - a*b*B + a^2*C)
/(b*(b*c - a*d)), Int[1/(a + b*Sin[e + f*x]), x], x] - Dist[(c^2*C - B*c*d
+ A*d^2)/(d*(b*c - a*d)), Int[1/(c + d*Sin[e + f*x]), x], x]) /; FreeQ[{a,
b, c, d, e, f, A, B, C}, x] && NeQ[b*c - a*d, 0] && NeQ[a^2 - b^2, 0] && Ne
Q[c^2 - d^2, 0]
```

Rule 3770

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x]
/; FreeQ[{c, d}, x]
```

Rule 3898

```
Int[cot[(c_.) + (d_.)*(x_)]^(m_)*(csc[(c_.) + (d_.)*(x_)]*(b_.) + (a_))^(n
_), x_Symbol] :> Int[(Cos[c + d*x]^m*(b + a*Sin[c + d*x])^n)/Sin[c + d*x]^
m + n), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && IntegerQ[n] &&
IntegerQ[m] && (IntegerQ[m/2] || LeQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int \frac{\cot^4(x)}{a+b \csc(x)} dx &= \int \frac{\cos(x) \cot^3(x)}{b+a \sin(x)} dx \\
&= \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} - \frac{\int \frac{\csc(x)(-2a^2+3b^2-ab \sin(x)-2b^2 \sin^2(x))}{b+a \sin(x)} dx}{2b^2} \\
&= \frac{x}{a} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} - \frac{(a^2-b^2)^2 \int \frac{1}{b+a \sin(x)} dx}{ab^3} - \frac{(-2a^2+3b^2) \int \csc(x) dx}{2b^3} \\
&= \frac{x}{a} - \frac{(2a^2-3b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} - \frac{(2(a^2-b^2)^2) \text{Subst}\left(\int \frac{1}{b+2ax+2b^2 \sin^2(x)} dx, x, \cos(x)\right)}{ab^3} \\
&= \frac{x}{a} - \frac{(2a^2-3b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b} + \frac{(4(a^2-b^2)^2) \text{Subst}\left(\int \frac{1}{4(a^2-b^2) \sin^2(x)} dx, x, \cos(x)\right)}{ab^3} \\
&= \frac{x}{a} - \frac{(2a^2-3b^2) \tanh^{-1}(\cos(x))}{2b^3} + \frac{2(a^2-b^2)^{3/2} \tanh^{-1}\left(\frac{a+b \tan(\frac{x}{2})}{\sqrt{a^2-b^2}}\right)}{ab^3} + \frac{a \cot(x)}{b^2} - \frac{\cot(x) \csc(x)}{2b}
\end{aligned}$$

Mathematica [A] time = 0.41, size = 158, normalized size = 1.74

$$\frac{8a^3 \log \left(\sin \left(\frac{x}{2}\right)\right)-8a^3 \log \left(\cos \left(\frac{x}{2}\right)\right)-16 \left(b^2-a^2\right)^{3/2} \tan ^{-1}\left(\frac{a+b \tan \left(\frac{x}{2}\right)}{\sqrt{b^2-a^2}}\right)-4 a^2 b \tan \left(\frac{x}{2}\right)+4 a^2 b \cot \left(\frac{x}{2}\right)-a b^2 \csc ^2\left(\frac{x}{2}\right)}{8 a b^3}$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^4/(a + b*Csc[x]), x]`

[Out]
$$\begin{aligned}
&(8 b^3 x - 16 (-a^2 + b^2)^{(3/2)} \operatorname{ArcTan}[(a + b \operatorname{Tan}[x/2])/\operatorname{Sqrt}[-a^2 + b^2]] \\
&+ 4 a^2 b^2 \operatorname{Cot}[x/2] - a b^2 \operatorname{Csc}[x/2]^2 - 8 a^3 \operatorname{Log}[\operatorname{Cos}[x/2]] + 12 a b^2 \operatorname{Log}[\operatorname{Cos}[x/2]] \\
&+ 8 a^3 \operatorname{Log}[\operatorname{Sin}[x/2]] - 12 a b^2 \operatorname{Log}[\operatorname{Sin}[x/2]] + a b^2 \operatorname{Sec}[x/2]^2 \\
&- 4 a^2 b \operatorname{Tan}[x/2])/(8 a b^3)
\end{aligned}$$

fricas [B] time = 0.63, size = 453, normalized size = 4.98

$$\frac{4 b^3 x \cos(x)^2 - 4 a^2 b \cos(x) \sin(x) - 4 b^3 x + 2 a b^2 \cos(x) - 2 ((a^2 - b^2) \cos(x)^2 - a^2 + b^2) \sqrt{a^2 - b^2} \log\left(-\frac{(a^2 - b^2) \cos(x)}{\sqrt{a^2 - b^2}}\right)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^4/(a+b*csc(x)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [1/4*(4*b^3*x*cos(x)^2 - 4*a^2*b*cos(x)*sin(x) - 4*b^3*x + 2*a*b^2*cos(x) - \\ & 2*((a^2 - b^2)*cos(x)^2 - a^2 + b^2)*sqrt(a^2 - b^2)*log(-(a^2 - 2*b^2)*c \\ & os(x)^2 + 2*a*b*sin(x) + a^2 + b^2 - 2*(b*cos(x)*sin(x) + a*cos(x))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) + (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(1/2*cos(x) + 1/2) - (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(-1/2*cos(x) + 1/2))/(a*b^3*cos(x)^2 - a*b^3), 1/4*(4*b^3*x*cos(x)^2 - 4*a^2*b*cos(x)*sin(x) - 4*b^3*x + 2*a*b^2*cos(x) + 4*((a^2 - b^2)*cos(x)^2 - a^2 + b^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/(a^2 - b^2)*cos(x))) + (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(1/2*cos(x) + 1/2) - (2*a^3 - 3*a*b^2 - (2*a^3 - 3*a*b^2)*cos(x)^2)*log(-1/2*cos(x) + 1/2))/(a*b^3*cos(x)^2 - a*b^3)] \end{aligned}$$

giac [B] time = 0.38, size = 163, normalized size = 1.79

$$\frac{x}{a} + \frac{b \tan\left(\frac{1}{2}x\right)^2 - 4a \tan\left(\frac{1}{2}x\right)}{8b^2} + \frac{(2a^2 - 3b^2) \log\left(\left|\tan\left(\frac{1}{2}x\right)\right|\right)}{2b^3} - \frac{2(a^4 - 2a^2b^2 + b^4) \left(\pi \left[\frac{x}{2\pi} + \frac{1}{2}\right] \operatorname{sgn}(b) + \arctan\left(\frac{x}{2\pi} + \frac{1}{2}\right)\right)}{\sqrt{-a^2 + b^2} ab^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^4/(a+b*csc(x)),x, algorithm="giac")`

[Out]
$$\begin{aligned} & x/a + 1/8*(b*tan(1/2*x)^2 - 4*a*tan(1/2*x))/b^2 + 1/2*(2*a^2 - 3*b^2)*log(a \\ & bs(tan(1/2*x))/b^3 - 2*(a^4 - 2*a^2*b^2 + b^4)*(pi*floor(1/2*x/pi + 1/2)*s \\ & gn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))/(sqrt(-a^2 + b^2)*a*b^ \\ & 3) - 1/8*(12*a^2*tan(1/2*x)^2 - 18*b^2*tan(1/2*x)^2 - 4*a*b*tan(1/2*x) + b^ \\ & 2)/(b^3*tan(1/2*x)^2) \end{aligned}$$

maple [B] time = 0.33, size = 206, normalized size = 2.26

$$\frac{\tan^2\left(\frac{x}{2}\right)}{8b} - \frac{a \tan\left(\frac{x}{2}\right)}{2b^2} - \frac{2a^3 \arctan\left(\frac{2 \tan\left(\frac{x}{2}\right)b+2a}{2\sqrt{-a^2+b^2}}\right)}{b^3 \sqrt{-a^2+b^2}} + \frac{4a \arctan\left(\frac{2 \tan\left(\frac{x}{2}\right)b+2a}{2\sqrt{-a^2+b^2}}\right)}{b \sqrt{-a^2+b^2}} - \frac{2b \arctan\left(\frac{2 \tan\left(\frac{x}{2}\right)b+2a}{2\sqrt{-a^2+b^2}}\right)}{a \sqrt{-a^2+b^2}} - \frac{1}{8b \tan\left(\frac{x}{2}\right)^2} +$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^4/(a+b*csc(x)),x)`

[Out]
$$\begin{aligned} & 1/8/b*tan(1/2*x)^2 - 1/2/b^2*a*tan(1/2*x) - 2/b^3*a^3/(-a^2+b^2)^(1/2)*arctan(1 \\ & /2*(2*tan(1/2*x)*b+2*a)/(-a^2+b^2)^(1/2)) + 4/b*a/(-a^2+b^2)^(1/2)*arctan(1/2 \\ & *(2*tan(1/2*x)*b+2*a)/(-a^2+b^2)^(1/2)) - 2*b/a/(-a^2+b^2)^(1/2)*arctan(1/2*(2 \\ & *tan(1/2*x)*b+2*a)/(-a^2+b^2)^(1/2)) - 1/8/b/tan(1/2*x)^2 + 1/b^3*ln(tan(1/2*x) \\ &)*a^2 - 3/2/b*ln(tan(1/2*x)) + 1/2*a/b^2/tan(1/2*x) + 2/a*arctan(tan(1/2*x)) \end{aligned}$$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)^4/(a+b*csc(x)),x, algorithm="maxima")

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details)Is $4*a^2-4*b^2$ positive or negative?

mupad [B] time = 1.19, size = 2163, normalized size = 23.77

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cot(x)^4/(a + b/sin(x)),x)

[Out]
$$\frac{(a^3 \log(\sin(x/2)/\cos(x/2)))/(a*b^3 - a*b^3*\cos(2*x)) - (2*b^3*\atan((2*b^3*\cos(x/2) + 2*a^3*\sin(x/2) - 3*a*b^2*\sin(x/2))/(2*b^3*\sin(x/2) - 2*a^3*\cos(x/2) + 3*a*b^2*\cos(x/2))))/(a*b^3 - a*b^3*\cos(2*x)) - (\atan((32*a^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} - 14*b^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2}) - 14*b^12*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2}) + 19*a*b^5*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} + 16*a^5*b*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} + 13*a*b^{11}*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2}) - 36*a^3*b^3*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} - 24*a^3*b^9*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} + 8*a^5*b^7*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} + 2*a^7*b^5*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} + 63*a^2*b^4*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} - 82*a^4*b^2*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} + 72*a^2*b^{10}*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} - 106*a^4*b^8*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} + 56*a^6*b^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} - 11*a^8*b^4*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2} + 2*a^{10}*b^2*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2})/(a^{15}*\sin(x/2)*32i + a^2*b^{13}*\cos(x/2)*27i - a^4*b^{11}*\cos(x/2)*165i + a^6*b^9*\cos(x/2)*390i - a^8*b^7*\cos(x/2)*469i + a^{10}*b^5*\cos(x/2)*309i - a^{12}*b^3*\cos(x/2)*108i + a^3*b^{12}*\sin(x/2)*108i - a^5*b^{10}*\sin(x/2)*482i + a^7*b^8*\sin(x/2)*982i - a^9*b^6*\sin(x/2)*1080i + a^{11}*b^4*\sin(x/2)*670i - a^{13}*b^2*\sin(x/2)*224i + a^{14}*b*\cos(x/2)*16i - a*b^{14}*\sin(x/2)*6i)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{1/2}*(2i)/(a*b^3 - a*b^3*\cos(2*x)) - (a^3*\cos(2*x)*\log(\sin(x/2)/\cos(x/2)))/(a*b^3 - a*b^3*\cos(2*x)) + (\atan((32*a^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2} - 14*b^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2}) - 14*b^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2}) - 14*b^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{3/2})$$

$$\begin{aligned}
& 3*a^2*b^4 - 3*a^4*b^2)^{(3/2)} - 14*b^{12}*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 19*a*b^5*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(3/2)} \\
& + 16*a^5*b*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 13*a*b^{11} \\
& * \cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} - 36*a^3*b^3*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(3/2)} - 24*a^3*b^9*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 8*a^5*b^7*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 2*a^7*b^5*\cos(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 63*a^2*b^4*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(3/2)} - 82*a^4*b^2*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(3/2)} + 72*a^2*b^{10}*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} - 106*a^4*b^8*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 56*a^6*b^6*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} - 11*a^8*b^4*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)} + 2*a^{10}*b^2*\sin(x/2)*(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)}/(a^{15}*\sin(x/2)*32i + a^2*b^{13}*\cos(x/2)*27i - a^4*b^{11}*\cos(x/2)*165i + a^6*b^9*\cos(x/2)*390i - a^8*b^7*\cos(x/2)*469i + a^{10}*b^5*\cos(x/2)*309i - a^{12}*b^3*\cos(x/2)*108i + a^3*b^{12}*\sin(x/2)*108i - a^5*b^{10}*\sin(x/2)*482i + a^7*b^8*\sin(x/2)*982i - a^9*b^6*\sin(x/2)*1080i + a^{11}*b^4*\sin(x/2)*670i - a^{13}*b^2*\sin(x/2)*224i + a^{14}*b*\cos(x/2)*16i - a*b^{14}*\sin(x/2)*6i)*\cos(2*x) \\
& *(a^6 - b^6 + 3*a^2*b^4 - 3*a^4*b^2)^{(1/2)*2i}/(a*b^3 - a*b^3*\cos(2*x)) - (3*a*b^2*\log(\sin(x/2)/\cos(x/2)))/(2*(a*b^3 - a*b^3*\cos(2*x))) + (2*b^3*\cos(2*x)*\atan((2*b^3*\cos(x/2) + 2*a^3*\sin(x/2) - 3*a*b^2*\sin(x/2))/(2*b^3*\sin(x/2) - 2*a^3*\cos(x/2) + 3*a*b^2*\cos(x/2)))/(a*b^3 - a*b^3*\cos(2*x)) - (a*b^2*\cos(x))/(a*b^3 - a*b^3*\cos(2*x)) + (a^2*b*\sin(2*x))/(a*b^3 - a*b^3*\cos(2*x)) + (3*a*b^2*\cos(2*x)*\log(\sin(x/2)/\cos(x/2)))/(2*(a*b^3 - a*b^3*\cos(2*x)))
\end{aligned}$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot^4(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cot(x)**4/(a+b*csc(x)),x)
[Out] Integral(cot(x)**4/(a + b*csc(x)), x)
```

$$3.23 \quad \int \frac{\cot^6(x)}{a+b \csc(x)} dx$$

Optimal. Leaf size=186

$$\frac{2(a^2 - b^2)^{5/2} \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab^5} + \frac{a(a^2 - 3b^2) \cot(x)}{b^4} - \frac{(a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} - \frac{(a^2 - 3b^2) \cot(x) \csc(x)}{2b^3} - \frac{(a^4 - 2a^2b^2 + b^4) \tanh^{-1}(\cos(x))}{2b^5}$$

[Out] $-\frac{x}{a} - \frac{3}{8} \operatorname{arctanh}(\cos(x)) / b - \frac{1}{2} (a^2 - 3b^2) \operatorname{arctanh}(\cos(x)) / b^3 - \frac{(a^4 - 3a^2b^2 + b^4) \operatorname{arctanh}((a+b \tan(1/2x)) / (a^2 - b^2)^{1/2})}{a/b^5} + \frac{a(b^2 + a^2 - 3b^2) \cot(x)}{b^4} - \frac{(a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} - \frac{(a^2 - 3b^2) \cot(x) \csc(x)}{2b^3} - \frac{(a^4 - 2a^2b^2 + b^4) \tanh^{-1}(\cos(x))}{2b^5}$

Rubi [A] time = 0.28, antiderivative size = 186, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 9, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.692$, Rules used = {3898, 2897, 3770, 3767, 8, 3768, 2660, 618, 206}

$$\frac{a(a^2 - 3b^2) \cot(x)}{b^4} - \frac{(a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} - \frac{(-3a^2b^2 + a^4 + 3b^4) \tanh^{-1}(\cos(x))}{b^5} + \frac{2(a^2 - b^2)^{5/2} \tanh^{-1}\left(\frac{a+b \tan\left(\frac{x}{2}\right)}{\sqrt{a^2-b^2}}\right)}{ab^5}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\cot[x]^6 / (a + b \csc[x]), x]$

[Out] $-\frac{x}{a} - \frac{3 \operatorname{ArcTanh}[\cos(x)]}{8b} - \frac{(a^2 - 3b^2) \operatorname{ArcTanh}[\cos(x)]}{2b^3} - \frac{(a^4 - 3a^2b^2 + 3b^4) \operatorname{ArcTanh}[\cos(x)]}{b^5} + \frac{(2(a^2 - b^2)^{5/2} \operatorname{ArcTanh}[(a + b \tan(x/2)) / \sqrt{a^2 - b^2}])}{a \cdot b^5} + \frac{(a \cot(x))}{b^2} + \frac{(a^2 - 3b^2) \cot(x)}{b^4} + \frac{(a \cot(x))^3}{3b^2} - \frac{(3 \cot(x) \csc(x))}{8b} - \frac{(a^2 - 3b^2) \cot(x) \csc(x)}{2b^3} - \frac{(\cot(x) \csc(x))^3}{4b}$

Rule 8

$\operatorname{Int}[a_, x_Symbol] :> \operatorname{Simp}[a_*x, x] /; \operatorname{FreeQ}[a, x]$

Rule 206

$\operatorname{Int}[(a_+ + b_-)(x_-)^2, x_Symbol] :> \operatorname{Simp}[(1 \operatorname{ArcTanh}[(\operatorname{Rt}[-b, 2] x) / \operatorname{Rt}[a, 2]]) / (\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2]), x] /; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{NegQ}[a/b] \& (\operatorname{GtQ}[a, 0] \|\operatorname{LtQ}[b, 0])$

Rule 618

$\operatorname{Int}[(a_+ + b_-)(x_-) + (c_-)(x_-)^2, x_Symbol] :> \operatorname{Dist}[-2, \operatorname{Subst}[\operatorname{Int}[1 / \operatorname{Simp}[b^2 - 4a_*c - x^2, x], x], x, b + 2c_*x], x] /; \operatorname{FreeQ}[\{a, b, c\},$

$x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 2660

```
Int[((a_) + (b_.)*sin[(c_.) + (d_.)*(x_)])^(-1), x_Symbol] :> With[{e = FreeFactors[Tan[(c + d*x)/2], x]}, Dist[(2*e)/d, Subst[Int[1/(a + 2*b*e*x + a*e^2*x^2), x], x, Tan[(c + d*x)/2]/e], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0]
```

Rule 2897

```
Int[cos[(e_.) + (f_.)*(x_)]^(p_)*((d_.)*sin[(e_.) + (f_.)*(x_)])^(n_)*((a_) + (b_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Int[ExpandTrig[(d*sin[e + f*x])^n*(a + b*sin[e + f*x])^m*(1 - sin[e + f*x]^2)^(p/2), x], x] /; FreeQ[{a, b, d, e, f}, x] && NeQ[a^2 - b^2, 0] && IntegersQ[m, 2*n, p/2] && (LtQ[m, -1] || (EqQ[m, -1] && GtQ[p, 0]))
```

Rule 3767

```
Int[csc[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> -Dist[d^(-1), Subst[Int[ExpandIntegrand[(1 + x^2)^(n/2 - 1), x], x, Cot[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[n/2, 0]
```

Rule 3768

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> -Simp[(b*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1))/(d*(n - 1)), x] + Dist[(b^2*(n - 2))/(n - 1), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]
```

Rule 3770

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> -Simp[ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rule 3898

```
Int[cot[(c_.) + (d_.)*(x_)]^(m_)*csc[(c_.) + (d_.)*(x_)]*(b_.)^(n_), x_Symbol] :> Int[(Cos[c + d*x]^m*(b + a*Sin[c + d*x])^n)/Sin[c + d*x]^(m + n), x] /; FreeQ[{a, b, c, d}, x] && NeQ[a^2 - b^2, 0] && IntegerQ[n] && IntegerQ[m] && (IntegerQ[m/2] || LeQ[m, 1])
```

Rubi steps

$$\begin{aligned}
\int \frac{\cot^6(x)}{a + b \csc(x)} dx &= \int \frac{\cos(x) \cot^5(x)}{b + a \sin(x)} dx \\
&= \int \left(-\frac{1}{a} + \frac{(a^4 - 3a^2b^2 + 3b^4) \csc(x)}{b^5} + \frac{(-a^3 + 3ab^2) \csc^2(x)}{b^4} + \frac{(a^2 - 3b^2) \csc^3(x)}{b^3} - \frac{a \csc^4(x)}{b^2} \right. \\
&\quad \left. - \frac{x}{a} - \frac{a \int \csc^4(x) dx}{b^2} + \frac{\int \csc^5(x) dx}{b} - \frac{(a(a^2 - 3b^2)) \int \csc^2(x) dx}{b^4} + \frac{(a^2 - 3b^2) \int \csc^3(x) dx}{b^3} \right. \\
&\quad \left. - \frac{x}{a} - \frac{(a^4 - 3a^2b^2 + 3b^4) \tanh^{-1}(\cos(x))}{b^5} - \frac{(a^2 - 3b^2) \cot(x) \csc(x)}{2b^3} - \frac{\cot(x) \csc^3(x)}{4b} + \frac{a \operatorname{Sub}(x, \csc(x), \csc^2(x))}{b^2} \right. \\
&\quad \left. - \frac{x}{a} - \frac{(a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} - \frac{(a^4 - 3a^2b^2 + 3b^4) \tanh^{-1}(\cos(x))}{b^5} + \frac{a \cot(x)}{b^2} + \frac{a(a^2 - 3b^2) \operatorname{Sub}(x, \csc(x), \csc^2(x))}{b^4} \right. \\
&\quad \left. - \frac{x}{a} - \frac{3 \tanh^{-1}(\cos(x))}{8b} - \frac{(a^2 - 3b^2) \tanh^{-1}(\cos(x))}{2b^3} - \frac{(a^4 - 3a^2b^2 + 3b^4) \tanh^{-1}(\cos(x))}{b^5} + \frac{a \operatorname{Sub}(x, \csc(x), \csc^2(x))}{b^6} \right) +
\end{aligned}$$

Mathematica [A] time = 0.92, size = 312, normalized size = 1.68

$$192a^5 \log \left(\sin \left(\frac{x}{2} \right) \right) - 192a^5 \log \left(\cos \left(\frac{x}{2} \right) \right) - 96a^4b \tan \left(\frac{x}{2} \right) - 24a^3b^2 \csc^2 \left(\frac{x}{2} \right) + 24a^3b^2 \sec^2 \left(\frac{x}{2} \right) - 480a^3b^2 \log \left(\sin \left(\frac{x}{2} \right) \right) + 480a^3b^2 \log \left(\cos \left(\frac{x}{2} \right) \right)$$

Antiderivative was successfully verified.

[In] `Integrate[Cot[x]^6/(a + b*Csc[x]), x]`

[Out]
$$\begin{aligned}
& (-192*b^5*x + 384*(-a^2 + b^2)^(5/2)*ArcTan[(a + b*Tan[x/2])/Sqrt[-a^2 + b^2]] + 32*a^2*b*(3*a^2 - 7*b^2)*Cot[x/2] - 24*a^3*b^2*Csc[x/2]^2 + 54*a*b^4*Csc[x/2]^2 - 3*a*b^4*Csc[x/2]^4 - 192*a^5*Log[Cos[x/2]] + 480*a^3*b^2*Log[Cos[x/2]] - 360*a*b^4*Log[Cos[x/2]] + 192*a^5*Log[Sin[x/2]] - 480*a^3*b^2*Log[Sin[x/2]] + 360*a*b^4*Log[Sin[x/2]] + 24*a^3*b^2*Sec[x/2]^2 - 54*a*b^4*Sec[x/2]^2 + 3*a*b^4*Sec[x/2]^4 - 64*a^2*b^3*Csc[x]^3*Sin[x/2]^4 + 4*a^2*b^3*Csc[x/2]^4*Sin[x] - 96*a^4*b*Tan[x/2] + 224*a^2*b^3*Tan[x/2])/(192*a*b^5)
\end{aligned}$$

fricas [B] time = 0.95, size = 852, normalized size = 4.58

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^6/(a+b*csc(x)),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [-1/48*(48*b^5*x*cos(x)^4 - 96*b^5*x*cos(x)^2 + 48*b^5*x - 6*(4*a^3*b^2 - 9*a*b^4)*cos(x)^3 - 24*((a^4 - 2*a^2*b^2 + b^4)*cos(x)^4 + a^4 - 2*a^2*b^2 + b^4 - 2*(a^4 - 2*a^2*b^2 + b^4)*cos(x)^2)*sqrt(a^2 - b^2)*log((a^2 - 2*b^2)*cos(x)^2 + 2*a*b*sin(x) + a^2 + b^2 + 2*(b*cos(x)*sin(x) + a*cos(x)))*sqrt(a^2 - b^2))/(a^2*cos(x)^2 - 2*a*b*sin(x) - a^2 - b^2)) + 6*(4*a^3*b^2 - 7*a*b^4)*cos(x) + 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(1/2*cos(x) + 1/2) - 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(-1/2*cos(x) + 1/2) + 16*((3*a^4*b - 7*a^2*b^3)*cos(x)^3 - 3*(a^4*b - 2*a^2*b^3)*cos(x)*sin(x))/(a*b^5*cos(x)^4 - 2*a*b^5*cos(x)^2 + a*b^5), -1/48*(48*b^5*x*cos(x)^4 - 96*b^5*x*cos(x)^2 + 48*b^5*x - 6*(4*a^3*b^2 - 9*a*b^4)*cos(x)^3 - 48*((a^4 - 2*a^2*b^2 + b^4)*cos(x)^4 + a^4 - 2*a^2*b^2 + b^4 - 2*(a^4 - 2*a^2*b^2 + b^4)*cos(x)^2)*sqrt(-a^2 + b^2)*arctan(-sqrt(-a^2 + b^2)*(b*sin(x) + a)/((a^2 - b^2)*cos(x))) + 6*(4*a^3*b^2 - 7*a*b^4)*cos(x) + 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(1/2*cos(x) + 1/2) - 3*(8*a^5 - 20*a^3*b^2 + 15*a*b^4 + (8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^4 - 2*(8*a^5 - 20*a^3*b^2 + 15*a*b^4)*cos(x)^2)*log(-1/2*cos(x) + 1/2) + 16*((3*a^4*b - 7*a^2*b^3)*cos(x)^3 - 3*(a^4*b - 2*a^2*b^3)*cos(x))*sin(x))/(a*b^5*cos(x)^4 - 2*a*b^5*cos(x)^2 + a*b^5)] \end{aligned}$$

giac [A] time = 1.01, size = 299, normalized size = 1.61

$$-\frac{x}{a} + \frac{3b^3 \tan\left(\frac{1}{2}x\right)^4 - 8ab^2 \tan\left(\frac{1}{2}x\right)^3 + 24a^2b \tan\left(\frac{1}{2}x\right)^2 - 48b^3 \tan\left(\frac{1}{2}x\right)^2 - 96a^3 \tan\left(\frac{1}{2}x\right) + 216ab^2 \tan\left(\frac{1}{2}x\right)}{192b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^6/(a+b*csc(x)),x, algorithm="giac")`

[Out]
$$\begin{aligned} & -x/a + 1/192*(3*b^3*tan(1/2*x)^4 - 8*a*b^2*tan(1/2*x)^3 + 24*a^2*b*tan(1/2*x)^2 - 48*b^3*tan(1/2*x)^2 - 96*a^3*tan(1/2*x) + 216*a*b^2*tan(1/2*x))/b^4 \\ & + 1/8*(8*a^4 - 20*a^2*b^2 + 15*b^4)*log(abs(tan(1/2*x)))/b^5 - 2*(a^6 - 3*a^4*b^2 + 3*a^2*b^4 - b^6)*(pi*floor(1/2*x/pi + 1/2)*sgn(b) + arctan((b*tan(1/2*x) + a)/sqrt(-a^2 + b^2)))/(sqrt(-a^2 + b^2)*a*b^5) - 1/192*(400*a^4*tan(1/2*x)^4 - 1000*a^2*b^2*tan(1/2*x)^4 + 750*b^4*tan(1/2*x)^4 - 96*a^3*b*tan(1/2*x)^3 + 216*a*b^3*tan(1/2*x)^3 + 24*a^2*b^2*tan(1/2*x)^2 - 48*b^4*tan(1/2*x)^2 - 8*a*b^3*tan(1/2*x) + 3*b^4)/(b^5*tan(1/2*x)^4) \end{aligned}$$

maple [B] time = 0.34, size = 363, normalized size = 1.95

$$\frac{\tan^4\left(\frac{x}{2}\right)}{64b} - \frac{\left(\tan^3\left(\frac{x}{2}\right)\right)a}{24b^2} + \frac{a^2\left(\tan^2\left(\frac{x}{2}\right)\right)}{8b^3} - \frac{\tan^2\left(\frac{x}{2}\right)}{4b} - \frac{a^3\tan\left(\frac{x}{2}\right)}{2b^4} + \frac{9a\tan\left(\frac{x}{2}\right)}{8b^2} - \frac{2a^5\arctan\left(\frac{2\tan\left(\frac{x}{2}\right)b+2a}{2\sqrt{-a^2+b^2}}\right)}{b^5\sqrt{-a^2+b^2}} + \frac{6a^3\arctan\left(\frac{2\tan\left(\frac{x}{2}\right)b+2a}{b^3\sqrt{-a^2+b^2}}\right)}{b^3\sqrt{-a^2+b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^6/(a+b*csc(x)),x)`

[Out]
$$\begin{aligned} & 1/64/b*\tan(1/2*x)^4 - 1/24/b^2*\tan(1/2*x)^3*a + 1/8/b^3*a^2*\tan(1/2*x)^2 - 1/4/b*\tan(1/2*x)^2 - 1/2/b^4*a^3*\tan(1/2*x) + 9/8/b^2*a*\tan(1/2*x) - 2*a^5/b^5/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*\tan(1/2*x)*b+2*a)/(-a^2+b^2)^{(1/2)}) + 6/b^3*a^3/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*\tan(1/2*x)*b+2*a)/(-a^2+b^2)^{(1/2)}) - 6/b*a/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*\tan(1/2*x)*b+2*a)/(-a^2+b^2)^{(1/2)}) + 2*b/a/(-a^2+b^2)^{(1/2)}*\arctan(1/2*(2*\tan(1/2*x)*b+2*a)/(-a^2+b^2)^{(1/2)}) - 1/64/b/\tan(1/2*x)^4 - 1/8/b^3/\tan(1/2*x)^2*a^2 + 1/4/b/\tan(1/2*x)^2 + 1/b^5*\ln(\tan(1/2*x))*a^4 - 5/2/b^3*\ln(\tan(1/2*x))*a^2 + 15/8/b*\ln(\tan(1/2*x)) + 1/24*a/b^2/\tan(1/2*x)^3 + 1/2*a^3/b^4/\tan(1/2*x) - 9/8*a/b^2/\tan(1/2*x) - 2/a*\arctan(\tan(1/2*x)) \end{aligned}$$

maxima [F(-2)] time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cot(x)^6/(a+b*csc(x)),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*a^2-4*b^2>0)', see `assume?` for more details) Is $4*a^2-4*b^2$ positive or negative?

mupad [B] time = 1.59, size = 4075, normalized size = 21.91

result too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cot(x)^6/(a + b/sin(x)),x)`

[Out]
$$\begin{aligned} & \tan(x/2)*(a/(8*b^2) + (2*a*(1/(2*b) - a^2/(4*b^3)))/b) - \tan(x/2)^2*(1/(4*b) - a^2/(8*b^3)) + \tan(x/2)^4/(64*b) - (2*\arctan(((4*(24*a*b^16 + 58*a^3*b^14 - 345*a^5*b^12 + 543*a^7*b^10 - 440*a^9*b^8 + 184*a^11*b^6 - 32*a^13*b^4))/b^12 + (((4*(24*a^3*b^16 - 32*a^5*b^14))/b^12 - (4*\tan(x/2)*(16*a^2*b^17 - 136*a^4*b^15 + 128*a^6*b^13))/b^12)*11)/a - (4*(93*a^3*b^15 - 240*a^5$$

$$\begin{aligned}
 & *b^{13} + 208*a^7*b^{11} - 64*a^9*b^9)) / b^{12} + (4*tan(x/2)*(62*a^2*b^{16} - 335*a^4*b^{14} + 604*a^6*b^{12} - 456*a^8*b^{10} + 128*a^{10}*b^8)) / b^{12}) * i) / a - (4*tan(x/2)*(16*b^{17} - 180*a^2*b^{15} + 252*a^4*b^{13} - 148*a^6*b^{11} + 100*a^8*b^9 - 40*a^{10}*b^7 + 8*a^{12}*b^5)) / b^{12}) * i) / a - (4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12} + (4*tan(x/2)*(8*a^{16} + 62*b^{16} - 410*a^2*b^{14} + 929*a^4*b^{12} - 1096*a^6*b^{10} + 873*a^8*b^8 - 550*a^10*b^6 + 255*a^12*b^4 - 68*a^{14}*b^2)) / b^{12}) / a - ((4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12} + ((4*(24*a*b^{16} + 58*a^3*b^{14} - 345*a^5*b^{12} + 543*a^7*b^{10} - 440*a^9*b^8 + 184*a^{11}*b^6 - 32*a^{13}*b^4)) / b^{12} + ((4*(93*a^3*b^{15} - 240*a^5*b^{13} + 208*a^7*b^11 - 64*a^9*b^9)) / b^{12} + ((4*(24*a^3*b^{16} - 32*a^5*b^{14})) / b^{12} - (4*tan(x/2)*(16*a^2*b^{17} - 136*a^4*b^{15} + 128*a^6*b^{13})) / b^{12}) * i) / a - (4*tan(x/2)*(62*a^2*b^{16} - 335*a^4*b^{14} + 604*a^6*b^{12} - 456*a^8*b^{10} + 128*a^{10}*b^8)) / b^{12}) * i) / a - (4*tan(x/2)*(16*b^{17} - 180*a^2*b^{15} + 252*a^4*b^{13} - 148*a^6*b^{11} + 100*a^8*b^9 - 40*a^{10}*b^7 + 8*a^{12}*b^5)) / b^{12}) * i) / a - (4*tan(x/2)*(8*a^{16} + 62*b^{16} - 410*a^2*b^{14} + 929*a^4*b^{12} - 1096*a^6*b^{10} + 873*a^8*b^8 - 550*a^10*b^6 + 255*a^12*b^4 - 68*a^{14}*b^2)) / b^{12}) / a) / (((((4*(24*a*b^{16} + 58*a^3*b^{14} - 345*a^5*b^{12} + 543*a^7*b^{10} - 440*a^9*b^8 + 184*a^{11}*b^6 - 32*a^{13}*b^4)) / b^{12} + (((4*(24*a^3*b^{16} - 32*a^5*b^{14})) / b^{12} - (4*tan(x/2)*(16*a^2*b^{17} - 136*a^4*b^{15} + 128*a^6*b^{13})) / b^{12}) * i) / a - (4*(93*a^3*b^{15} - 240*a^5*b^{13} + 208*a^7*b^{11} - 64*a^9*b^9)) / b^{12} + (4*tan(x/2)*(62*a^2*b^{16} - 335*a^4*b^{14} + 604*a^6*b^{12} - 456*a^8*b^{10} + 128*a^{10}*b^8)) / b^{12}) * i) / a - (4*tan(x/2)*(16*b^{17} - 180*a^2*b^{15} + 252*a^4*b^{13} - 148*a^6*b^{11} + 100*a^8*b^9 - 40*a^{10}*b^7 + 8*a^{12}*b^5)) / b^{12}) * i) / a - (4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12} + ((4*(24*a*b^{16} + 58*a^3*b^{14} - 345*a^5*b^{12} + 543*a^7*b^{10} - 440*a^9*b^8 + 184*a^{11}*b^6 - 32*a^{13}*b^4)) / b^{12} + (((4*(93*a^3*b^{15} - 240*a^5*b^{13} + 208*a^7*b^{11} - 64*a^9*b^9)) / b^{12} + ((4*(24*a^3*b^{16} - 32*a^5*b^{14})) / b^{12} - (4*tan(x/2)*(16*a^2*b^{17} - 136*a^4*b^{15} + 128*a^6*b^{13})) / b^{12}) * i) / a - (4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12} + ((4*(24*a^3*b^{16} - 32*a^5*b^{14})) / b^{12} - (4*tan(x/2)*(16*a^2*b^{17} - 136*a^4*b^{15} + 128*a^6*b^{13})) / b^{12}) * i) / a - (4*tan(x/2)*(62*a^2*b^{16} - 335*a^4*b^{14} + 604*a^6*b^{12} - 456*a^8*b^{10} + 128*a^{10}*b^8)) / b^{12}) * i) / a - (4*tan(x/2)*(16*b^{17} - 180*a^2*b^{15} + 252*a^4*b^{13} - 148*a^6*b^{11} + 100*a^8*b^9 - 40*a^{10}*b^7 + 8*a^{12}*b^5)) / b^{12}) * i) / a - (4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12} + ((4*(24*a*b^{16} + 58*a^3*b^{14} - 345*a^5*b^{12} + 543*a^7*b^{10} - 440*a^9*b^8 + 184*a^{11}*b^6 - 32*a^{13}*b^4)) / b^{12} + (((4*(93*a^3*b^{15} - 240*a^5*b^{13} + 208*a^7*b^{11} - 64*a^9*b^9)) / b^{12} + ((4*(24*a^3*b^{16} - 32*a^5*b^{14})) / b^{12} - (4*tan(x/2)*(16*a^2*b^{17} - 136*a^4*b^{15} + 128*a^6*b^{13})) / b^{12}) * i) / a - (4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12} + ((4*(24*a^3*b^{16} - 32*a^5*b^{14})) / b^{12} - (4*tan(x/2)*(16*a^2*b^{17} - 136*a^4*b^{15} + 128*a^6*b^{13})) / b^{12}) * i) / a - (4*tan(x/2)*(8*a^{16} + 62*b^{16} - 410*a^2*b^{14} + 929*a^4*b^{12} - 1096*a^6*b^{10} + 873*a^8*b^8 - 550*a^10*b^6 + 255*a^12*b^4 - 68*a^{14}*b^2)) / b^{12}) / a + (8*(15*a*b^{14} + 8*a^{15} - 20*a^3*b^{12} - 97*a^5*b^{10} + 305*a^7*b^8 - 366*a^9*b^6 + 223*a^{11}*b^4 - 68*a^{13}*b^2)) / b^{12} - (8*tan(x/2)*(4*a^2*b^{13} - 4*b^{15} + 12*a^4*b^{11} - 20*a^6*b^9 + 8*a^8*b^7)) / b^{12})) / a - (a*tan(x/2)^3) / (24*b^2) - (tan(x/2)^3*(18*a^2*b^2 - 8*a^3) + tan(x/2)^2*(2*a^2*b - 4*b^3) + b^3/4 - (2*a^2*b^2*tan(x/2)) / 3) / (16*b^4*tan(x/2)^4) + (log(tan(x/2)) * (a^4 + (15*b^4) / 8 - (5*a^2*b^2) / 2)) / b^5 - (atan(((a + b)^5 * (a - b)^5)^{(1/2)} * ((4*tan(x/2) * (8*a^16 + 62*b^16 - 410*a^2*b^14 + 929*a^4*b^12 - 1096*a^6*b^10 + 873*a^8*b^8 - 550*a^10*b^6 + 255*a^12*b^4 - 68*a^14*b^2)) / b^12) - (4*(53*a*b^{15} - 120*a^3*b^{13} + 48*a^5*b^{11} + 56*a^7*b^9 - 48*a^9*b^7 + 8*a^{11}*b^5)) / b^{12}) / a) / 2
 \end{aligned}$$

$$\begin{aligned} & \text{---} \\ & ((4*(24*a^3*b^16 - 32*a^5*b^14))/b^12 - (4*tan(x/2)*(16*a^2*b^17 - 136*a^4*b^15 + 128*a^6*b^13))/b^12)*((a + b)^5*(a - b)^5)^{(1/2)}/(a*b^5)) \\ & ((a*b^5))/((a*b^5)))/((a*b^5)))*((a + b)^5*(a - b)^5)^{(1/2)*2i}/(a*b^5) \end{aligned}$$

sympy [F] time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cot^6(x)}{a + b \csc(x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cot(x)**6/(a+b*csc(x)),x)
[Out] Integral(cot(x)**6/(a + b*csc(x)), x)

Chapter 4

Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.0.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)

GradeAntiderivative[result_,optimal_] :=
  If[ExpnType[result]<=ExpnType[optimal],
```

```

If[FreeQ[result,Complex] || Not[FreeQ[optimal,Complex]],

  If[LeafCount[result]<=2*LeafCount[optimal],

    "A",
    "B"],
   "C"],

If[FreeQ[result,Integrate] && FreeQ[result,Int],

  "C",
  "F"]]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)
(*9 = unknown function*)

ExpnType[expn_] :=
  If[AtomQ[expn],
   1,
   If[ListQ[expn],
    Max[Map[ExpnType, expn]],
    If[Head[expn]==Power,
     If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
      If[Head[expn[[2]]]==Rational,
       If[IntegerQ[expn[[1]]] || Head[expn[[1]]]==Rational,
        1,
        Max[ExpnType[expn[[1]]],2]],
      Max[ExpnType[expn[[1]]],ExpnType[expn[[2]]],3]]],
     If[Head[expn]==Plus || Head[expn]==Times,
      Max[ExpnType[First[expn]],ExpnType[Rest[expn]]],
      If[ElementaryFunctionQ[Head[expn]],
       Max[3,ExpnType[expn[[1]]]],
       If[SpecialFunctionQ[Head[expn]],
        Apply[Max,Append[Map[ExpnType,Apply[List,expn]],4]],
        If[HypergeometricFunctionQ[Head[expn]],
         Apply[Max,Append[Map[ExpnType,Apply[List,expn]],5]],
         If[AppellFunctionQ[Head[expn]],
          Apply[Max,Append[Map[ExpnType,Apply[List,expn]],6]],


```

```

If [Head[expn]==RootSum,
  Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
  If[Head[expn]==Integrate || Head[expn]==Int,
    Apply[Max,Append[Map[ExpnType,Apply[List,expn]],8]],
  9]]]]]]]]]
}

ElementaryFunctionQ[func_] :=
MemberQ[{  

  Exp, Log,  

  Sin, Cos, Tan, Cot, Sec, Csc,  

  ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  

  Sinh, Cosh, Tanh, Coth, Sech, Csch,  

  ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsch
}, func]

SpecialFunctionQ[func_] :=
MemberQ[{  

  Erf, Erfc, Erfi,  

  FresnelS, FresnelC,  

  ExpIntegralE, ExpIntegralEi, LogIntegral,  

  SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,  

  Gamma, LogGamma, PolyGamma,  

  Zeta, PolyLog, ProductLog,  

  EllipticF, EllipticE, EllipticPi
}, func]

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]

```

4.0.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#               if leaf size is "too large". Set at 500,000

```

```

#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
# see problem 156, file Apostol_Problems

GradeAntiderivative := proc(result,optimal)
local leaf_count_result, leaf_count_optimal,ExpnType_result,ExpnType_optimal,
      debug:=false;

leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
    return "B";
fi;

leaf_count_optimal:=leafcount(optimal);

ExpnType_result:=ExpnType(result);
ExpnType_optimal:=ExpnType(optimal);

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",
ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
#   "F" if the result fails to integrate an expression that
#       is integrable
#   "C" if result involves higher level functions than necessary
#   "B" if result is more than twice the size of the optimal
#       antiderivative
#   "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F";
end if;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;

```

```

if is_contains_complex(result) then
    if is_contains_complex(optimal) then
        if debug then
            print("both result and optimal complex");
        fi;
        #both result and optimal complex
        if leaf_count_result<=2*leaf_count_optimal then
            return "A";
        else
            return "B";
        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C";
    end if
else # result do not contain complex
    # this assumes optimal do not as well
    if debug then
        print("result do not contain complex, this assumes optimal do not
as well");
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B";
    end if
    end if
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C";
end if

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false

```

```

#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc;

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
# 6 = appell function
# 7 = rootsum function
# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
    if type(expn,'atomic') then
        1
    elif type(expn,'list') then
        apply(max,map(ExpnType,expn))
    elif type(expn,'sqrt') then
        if type(op(1,expn),'rational') then
            1
        else
            max(2,ExpnType(op(1,expn)))
        end if
    elif type(expn,'`^') then
        if type(op(2,expn),'integer') then
            ExpnType(op(1,expn))
        elif type(op(2,expn),'rational') then
            if type(op(1,expn),'rational') then
                1
            else
                max(2,ExpnType(op(1,expn)))
            end if
        else
            max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
        end if
    elif type(expn,'`+`) or type(expn,'`*`) then
        max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif ElementaryFunctionQ(op(0,expn)) then
        max(3,ExpnType(op(1,expn)))
    elif SpecialFunctionQ(op(0,expn)) then
        max(4,apply(max,map(ExpnType,[op(expn)])))
    end if
end proc;

```

```

elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
9
end if
end proc:

ElementaryFunctionQ := proc(func)
member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,
    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
if nops(u)=2 then
    op(2,u)
else
    apply(op(0,u),op(2..nops(u),u))
end if
end proc:

```

```
#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
    MmaTranslator[Mma][LeafCount](u);
end proc:
```

4.0.3 Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                   asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                   asinh,acosh,atanh,acoth,asech,acsch
                  ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                  ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]
```

```

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

    except AttributeError as error:
        return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'````')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        )
    else:
        return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,
ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+``') or type
(expn,'`*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))

```

```

    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,
        expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,
        Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:
        m1 = max(map(expnType, list(expn.args)))
        return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

#main function
def grade_antiderivative(result,optimal):

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        return "F"

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    return "A"
                else:
                    return "B"
            else: #result contains complex but optimal is not
                return "C"
        else: # result do not contain complex, this assumes optimal do not as
        well
            if leaf_count_result <= 2*leaf_count_optimal:
                return "A"
            else:
                return "B"
        else:
            return "C"
    else:
        return "C"

```

4.0.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#                  Albert Rich to use with Sagemath. This is used to
#                  grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#                  'arctan2','floor','abs','log_integral'

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow:    #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    debug=False
    m = func.name() in ['exp','log','ln',
                        'sin','cos','tan','cot','sec','csc',
                        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
                        'sinh','cosh','tanh','coth','sech','csch',
                        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
                        'arctan2','floor','abs'
                        ]
    if debug:
        if m:
```

```

        print ("func ", func , " is elementary_function")
    else:
        print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    debug=False
    if debug: print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
                       'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','
sinh_integral'
                       'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
                       'polylog','lambert_w','elliptic_f','elliptic_e',
                       'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U
    ']

def is_appell_function(func):
    return func.name() in ['hypergeometric']    #[appellf1] can't find this in
sagemath

def is_atom(expn):

    debug=False
    if debug: print ("Enter is_atom")

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-
    sagemath-equivalent-to-atomic-type-in-maple/
    try:
        if expn.parent() is SR:

```

```

        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(),"base_ring") and hasattr(expn.parent(),"gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().
gens()
    return False

except AttributeError as error:
    return False

def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list:  #isinstance(expn,list):
        return max(map(expnType, expn))  #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],
Rational)):
            return 1
        else:
            return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.
args[0]))
    elif expn.operator() == operator.pow:  #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer:  #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0])  #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational:  #isinstance(expn.args[1],
Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],
Rational)
                return 1
            else:
                return max(2,expnType(expn.operands()[0]))  #max(2,expnType(expn.
args[0]))
        else:
            return max(3,expnType(expn.operands()[0]),expnType(expn.operands()
[1])) #max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1]))
    elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #
isinstance(expn,Add) or isinstance(expn,Mul)
        m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
        m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))

```

```

        return max(m1,m2)  #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.operator()):  #is_elementary_function(expn.
        func)
        return max(3,expnType(expn.operands()[0]))
    elif is_special_function(expn.operator()): #is_special_function(expn.func)
        m1 = max(map(expnType, expn.operands()))           #max(map(expnType, list(
        expn.args)))
        return max(4,m1)  #max(4,m1)
    elif is_hypergeometric_function(expn.operator()): #
        is_hypergeometric_function(expn.func)
        m1 = max(map(expnType, expn.operands()))           #max(map(expnType, list(
        expn.args)))
        return max(5,m1)  #max(5,m1)
    elif is_appell_function(expn.operator()):
        m1 = max(map(expnType, expn.operands()))           #max(map(expnType, list(
        expn.args)))
        return max(6,m1)  #max(6,m1)
    elif str(expn).find("Integral") != -1: #this will never happen, since it
        #is checked before calling the grading function that is passed.
        #but kept it here.
        m1 = max(map(expnType, expn.operands()))           #max(map(expnType, list(
        expn.args)))
        return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
    else:
        return 9

#main function
def grade_antiderivative(result,optimal):

    if debug: print ("Enter grade_antiderivative for sageMath")

    leaf_count_result  = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    if debug: print ("leaf_count_result=", leaf_count_result, "
    leaf_count_optimal=",leaf_count_optimal)

    expnType_result  = expnType(result)
    expnType_optimal = expnType(optimal)

    if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",",
    expnType_optimal)

    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex

```

```
if leaf_count_result <= 2*leaf_count_optimal:  
    return "A"  
else:  
    return "B"  
else: #result contains complex but optimal is not  
    return "C"  
else: # result do not contain complex, this assumes optimal do not as  
well  
    if leaf_count_result <= 2*leaf_count_optimal:  
        return "A"  
    else:  
        return "B"  
else:  
    return "C"
```