

Computer algebra independent integration tests

Summer 2022 edition

1-Algebraic-functions/1.2-Trinomial-products/1.2.2-Quartic/44-

1.2.2.7-P-x-d+e-x²-^q-a+b-x²+c-x⁴-^p

Nasser M. Abbasi

September 28, 2022

Compiled on September 28, 2022 at 1:24am

Contents

1	Introduction	3
2	detailed summary tables of results	19
3	Listing of integrals	35
4	Appendix	259

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	list of integrals that has no closed form antiderivative	11
1.5	List of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	13
1.7	Timing	13
1.8	Verification	14
1.9	Important notes about some of the results	14
1.10	Design of the test system	17

This report gives the result of running the computer algebra independent integration test. The download section in the appendix contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [42]. This is test number [44].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.1 (June 29, 2022) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.0.1 on windows 10.
3. Maple 2022.1 (June 1, 2022) on windows 10.
4. Maxima 5.46 (April 13, 2022) using Lisp SBCL 2.1.11.debian on Linux via sagemath 9.6.
5. Fricas 1.3.8 (June 21, 2022) based on sbcl 2.1.11.debian on Linux via sagemath 9.6.
6. Giac/Xcas 1.9.0-13 (July 3, 2022) on Linux via sagemath 9.6.
7. Sympy 1.10.1 (March 20, 2022) Using Python 3.10.4 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly from Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (42)	0.00 (0)
Mathematica	95.24 (40)	4.76 (2)
Maple	95.24 (40)	4.76 (2)
Fricas	21.43 (9)	78.57 (33)
Sympy	14.29 (6)	85.71 (36)
Giac	11.90 (5)	88.10 (37)
Mupad	2.38 (1)	97.62 (41)
Maxima	0.00 (0)	100.00 (42)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Maple	23.81	33.33	38.10	4.76
Mathematica	16.67	0.00	78.57	4.76
Fricas	14.29	7.14	0.00	78.57
Giac	9.52	2.38	0.00	88.10
Mupad	N/A	2.38	0.00	97.62
Maxima	0.00	0.00	0.00	100.00
Sympy	0.00	0.00	14.29	85.71

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and

Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	2	100.00 %	0.00 %	0.00 %
Maple	2	100.00 %	0.00 %	0.00 %
Fricas	33	15.15 %	33.33 %	51.52 %
Giac	37	91.89 %	2.70 %	5.41 %
Maxima	42	85.71 %	0.00 %	14.29 %
Sympy	36	83.33 %	16.67 %	0.00 %
Mupad	41	100.00 %	0.00 %	0.00 %

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.



Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.53	576.24	1.08	396.50	1.00
Mathematica	8.85	422.63	0.82	259.50	0.66
Maple	0.16	1063.33	1.72	450.00	1.33
Maxima	0.00	0.00	0.00	0.00	0.00
Fricas	0.85	617.44	2.59	669.00	2.70
Sympy	2.89	186.00	0.53	167.00	0.51
Giac	5.04	359.00	1.32	360.00	1.17
Mupad	1.85	397.00	3.75	397.00	3.75

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used from the above table.



1.4 list of integrals that has no closed form antiderivative

{

1.5 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for `Rubi` and `Mathematica`.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
```

```
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified.

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,..}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	20
2.2	Detailed conclusion table per each integral for all CAS systems	23
2.3	Detailed conclusion table specific for Rubi results	32

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	21
2.1.2	Mathematica	21
2.1.3	Maple	21
2.1.4	Maxima	21
2.1.5	FriCAS	22
2.1.6	Sympy	22
2.1.7	Giac	22
2.1.8	Mupad	22

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 35, 37, 38, 39, 40, 41, 42 }

B grade: { }

C grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 34, 36 }

F grade: { 15, 33 }

2.1.3 Maple

A grade: { 20, 21, 30, 31, 37, 38, 39, 40, 41, 42 }

B grade: { 17, 18, 19, 22, 23, 24, 25, 26, 27, 28, 29, 34, 35, 36 }

C grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 16, 32 }

F grade: { 15, 33 }

2.1.4 Maxima

A grade: { }

B grade: { }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 }

2.1.5 FriCAS

A grade: { 37, 38, 39, 40, 41, 42 }

B grade: { 34, 35, 36 }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33 }

2.1.6 Sympy

A grade: { }

B grade: { }

C grade: { 1, 2, 3, 4, 10, 11 }

F grade: { 5, 6, 7, 8, 9, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42 }

2.1.7 Giac

A grade: { 34, 37, 38, 41 }

B grade: { 42 }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 39, 40 }

2.1.8 Mupad

A grade: { }

B grade: { 34 }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 36, 37, 38, 39, 40, 41, 42 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column N.S. in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, **Mathematica** was abbrevi-

	Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
viated to MMA.	grade	A	A	C	C	F	F(-2)	C	F	F
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	453	878	217	533	0	0	364	0	-1
	N.S.	1	1.94	0.48	1.18	0.00	0.00	0.80	0.00	-0.00
	time (sec)	N/A	0.340	10.195	0.184	0.000	0.000	3.888	0.000	0.000

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	367	706	159	403	0	0	262	0	-1
N.S.	1	1.92	0.43	1.10	0.00	0.00	0.71	0.00	-0.00
time (sec)	N/A	0.253	10.116	0.146	0.000	0.000	2.877	0.000	0.000

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	277	432	121	269	0	0	167	0	-1
N.S.	1	1.56	0.44	0.97	0.00	0.00	0.60	0.00	-0.00
time (sec)	N/A	0.143	10.071	0.140	0.000	0.000	1.977	0.000	0.000

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	226	226	77	169	0	0	78	0	-1
N.S.	1	1.00	0.34	0.75	0.00	0.00	0.35	0.00	-0.00
time (sec)	N/A	0.046	10.037	0.123	0.000	0.000	0.927	0.000	0.000

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	369	369	138	192	0	0	0	0	-1
N.S.	1	1.00	0.37	0.52	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.244	10.225	0.154	0.000	0.000	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	641	641	297	679	0	0	0	0	-1
N.S.	1	1.00	0.46	1.06	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.763	10.582	0.161	0.000	0.000	0.000	0.000	0.000

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	875	875	453	1591	0	0	0	0	-1
N.S.	1	1.00	0.52	1.82	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.173	11.126	0.158	0.000	0.000	0.000	0.000	0.000

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	912	912	222	588	0	0	0	0	-1
N.S.	1	1.00	0.24	0.64	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.450	10.201	0.195	0.000	0.000	0.000	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	694	694	166	458	0	0	0	0	-1
N.S.	1	1.00	0.24	0.66	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.338	10.132	0.162	0.000	0.000	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	395	395	126	320	0	0	167	0	-1
N.S.	1	1.00	0.32	0.81	0.00	0.00	0.42	0.00	-0.00
time (sec)	N/A	0.211	10.087	0.139	0.000	0.000	4.912	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	C	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	262	262	99	217	0	0	78	0	-1
N.S.	1	1.00	0.38	0.83	0.00	0.00	0.30	0.00	-0.00
time (sec)	N/A	0.066	10.038	0.138	0.000	0.000	2.778	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	732	732	432	564	0	0	0	0	-1
N.S.	1	1.00	0.59	0.77	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.531	10.535	0.171	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1494	1494	427	1384	0	0	0	0	-1
N.S.	1	1.00	0.29	0.93	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.362	10.851	0.169	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	2452	2452	630	2326	0	0	0	0	-1
N.S.	1	1.00	0.26	0.95	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	2.619	11.491	0.173	0.000	0.000	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-1)	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	169	169	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.137	0.404	0.048	0.000	0.000	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	94	81	0	0	0	0	-1
N.S.	1	1.00	1.96	1.69	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.018	10.097	0.161	0.000	0.000	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	755	755	901	1708	0	0	0	0	-1
N.S.	1	1.00	1.19	2.26	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.853	14.311	0.133	0.000	0.000	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	528	528	674	1201	0	0	0	0	-1
N.S.	1	1.00	1.28	2.27	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.465	12.739	0.183	0.000	0.000	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	368	368	521	759	0	0	0	0	-1
N.S.	1	1.00	1.42	2.06	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.164	11.311	0.039	0.000	0.000	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	283	283	302	362	0	0	0	0	-1
N.S.	1	1.00	1.07	1.28	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.049	10.142	0.018	0.000	0.000	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	436	436	298	359	0	0	0	0	-1
N.S.	1	1.00	0.68	0.82	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.269	10.340	0.158	0.000	0.000	0.000	0.000	0.000

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	782	782	1853	1495	0	0	0	0	-1
N.S.	1	1.00	2.37	1.91	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.959	12.001	0.153	0.000	0.000	0.000	0.000	0.000

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1125	1125	781	4476	0	0	0	0	-1
N.S.	1	1.00	0.69	3.98	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	2.270	13.309	0.146	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	859	859	1058	2445	0	0	0	0	-1
N.S.	1	1.00	1.23	2.85	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.890	14.740	0.164	0.000	0.000	0.000	0.000	0.000

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	628	633	766	1891	0	0	0	0	-1
N.S.	1	1.01	1.22	3.01	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.376	12.978	0.162	0.000	0.000	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	481	481	597	1390	0	0	0	0	-1
N.S.	1	1.00	1.24	2.89	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.238	11.694	0.039	0.000	0.000	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	398	398	497	931	0	0	0	0	-1
N.S.	1	1.00	1.25	2.34	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.136	10.903	0.037	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	867	1045	1736	3241	0	0	0	0	-1
N.S.	1	1.21	2.00	3.74	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.966	13.291	0.145	0.000	0.000	0.000	0.000	0.000

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	1301	2112	1116	8276	0	0	0	0	-1
N.S.	1	1.62	0.86	6.36	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	2.209	14.939	0.204	0.000	0.000	0.000	0.000	0.000

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-1)	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	273	273	310	365	0	0	0	0	-1
N.S.	1	1.00	1.14	1.34	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.100	10.334	0.164	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	271	271	312	369	0	0	0	0	-1
N.S.	1	1.00	1.15	1.36	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.123	10.299	0.158	0.000	0.000	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	74	93	0	0	0	0	-1
N.S.	1	1.00	0.70	0.88	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.066	10.137	0.136	0.000	0.000	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-1)	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	218	218	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.230	0.429	0.036	0.000	0.000	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	81	296	0	279	0	80	397
N.S.	1	1.00	0.76	2.79	0.00	2.63	0.00	0.75	3.75
time (sec)	N/A	0.090	0.090	0.373	0.000	0.407	0.000	4.106	1.845

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	145	145	139	222	0	603	0	0	-1
N.S.	1	1.00	0.96	1.53	0.00	4.16	0.00	0.00	-0.01
time (sec)	N/A	0.141	0.540	0.132	0.000	1.798	0.000	0.000	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	239	239	151	784	0	669	0	0	-1
N.S.	1	1.00	0.63	3.28	0.00	2.80	0.00	0.00	-0.00
time (sec)	N/A	0.120	0.555	0.159	0.000	1.702	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	309	309	212	442	0	468	0	360	-1
N.S.	1	1.00	0.69	1.43	0.00	1.51	0.00	1.17	-0.00
time (sec)	N/A	0.413	0.568	0.213	0.000	0.397	0.000	3.622	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	283	283	166	373	0	360	0	265	-1
N.S.	1	1.00	0.59	1.32	0.00	1.27	0.00	0.94	-0.00
time (sec)	N/A	0.230	0.414	0.188	0.000	0.380	0.000	3.712	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	286	286	174	251	0	771	0	0	-1
N.S.	1	1.00	0.61	0.88	0.00	2.70	0.00	0.00	-0.00
time (sec)	N/A	0.497	0.489	0.203	0.000	0.838	0.000	0.000	0.000

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	294	169	288	0	767	0	0	-1
N.S.	1	1.00	0.57	0.98	0.00	2.61	0.00	0.00	-0.00
time (sec)	N/A	0.536	0.625	0.219	0.000	0.708	0.000	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	288	288	173	330	0	797	0	370	-1
N.S.	1	1.00	0.60	1.15	0.00	2.77	0.00	1.28	-0.00
time (sec)	N/A	0.542	0.612	0.202	0.000	0.740	0.000	7.319	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F(-1)	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	294	294	186	412	0	843	0	720	-1
N.S.	1	1.00	0.63	1.40	0.00	2.87	0.00	2.45	-0.00
time (sec)	N/A	0.511	0.768	0.202	0.000	0.700	0.000	6.430	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [30] had the largest ratio of [41]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	15	5	1.94	28	0.179
2	A	12	5	1.92	28	0.179
3	A	8	5	1.56	26	0.192
4	A	3	3	1.00	19	0.158
5	A	3	3	1.00	28	0.107
6	A	6	6	1.00	28	0.214
7	A	7	6	1.00	28	0.214
8	A	12	7	1.00	28	0.250
9	A	10	6	1.00	28	0.214
10	A	7	5	1.00	26	0.192
11	A	4	4	1.00	19	0.210
12	A	9	7	1.00	28	0.250
13	A	15	10	1.00	28	0.357
14	A	22	11	1.00	28	0.393
15	A	6	3	1.00	26	0.115
16	A	2	2	1.00	27	0.074
17	A	6	4	1.00	33	0.121
18	A	5	4	1.00	33	0.121
19	A	4	4	1.00	31	0.129
20	A	3	3	1.00	24	0.125
21	A	3	3	1.00	33	0.091
22	A	6	6	1.00	33	0.182
23	A	7	6	1.00	33	0.182
24	A	5	5	1.00	33	0.152
25	A	4	4	1.01	33	0.121

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
26	A	4	4	1.00	31	0.129
27	A	4	4	1.00	24	0.167
28	A	9	7	1.21	33	0.212
29	A	15	10	1.62	33	0.303
30	A	1	1	1.00	41	0.024
31	A	1	1	1.00	41	0.024
32	A	4	4	1.00	31	0.129
33	A	6	3	1.00	31	0.097
34	A	11	7	1.00	28	0.250
35	A	4	3	1.00	30	0.100
36	A	1	1	1.00	30	0.033
37	A	6	6	1.00	38	0.158
38	A	6	6	1.00	37	0.162
39	A	8	7	1.00	40	0.175
40	A	8	7	1.00	40	0.175
41	A	8	7	1.00	40	0.175
42	A	8	7	1.00	40	0.175

Chapter 3

Listing of integrals

Local contents

3.1	$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+cx^4}} dx$	36
3.2	$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$	42
3.3	$\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$	47
3.4	$\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$	52
3.5	$\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$	56
3.6	$\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+cx^4}} dx$	60
3.7	$\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+cx^4}} dx$	65
3.8	$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$	72
3.9	$\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$	78
3.10	$\int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$	83
3.11	$\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$	88
3.12	$\int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx$	92
3.13	$\int \frac{A+Bx^2}{(d+ex^2)^2(a+cx^4)^{3/2}} dx$	98
3.14	$\int \frac{A+Bx^2}{(d+ex^2)^3(a+cx^4)^{3/2}} dx$	106
3.15	$\int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$	115
3.16	$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$	118
3.17	$\int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$	121
3.18	$\int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+bx^2+cx^4}} dx$	127

3.19	$\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$	133
3.20	$\int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$	138
3.21	$\int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$	142
3.22	$\int \frac{A+Bx^2}{(d+ex^2)^2\sqrt{a+bx^2+cx^4}} dx$	146
3.23	$\int \frac{A+Bx^2}{(d+ex^2)^3\sqrt{a+bx^2+cx^4}} dx$	153
3.24	$\int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$	160
3.25	$\int \frac{(A+Bx^2)(d+ex^2)^2}{(a+bx^2+cx^4)^{3/2}} dx$	167
3.26	$\int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$	173
3.27	$\int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$	178
3.28	$\int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$	183
3.29	$\int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$	191
3.30	$\int \frac{\sqrt{a} + \sqrt{c} x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$	198
3.31	$\int \frac{1 + \sqrt{\frac{c}{a}} x^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$	202
3.32	$\int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$	206
3.33	$\int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$	210
3.34	$\int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$	213
3.35	$\int \frac{\sqrt{a+bx^2+cx^4}}{ad-cdx^4} dx$	218
3.36	$\int \frac{\sqrt{a+bx^2-cx^4}}{ad+cdx^4} dx$	222
3.37	$\int x\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4} dx$	226
3.38	$\int \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4} dx$	231
3.39	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x} dx$	236
3.40	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx$	241
3.41	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx$	247
3.42	$\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx$	253


```
*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(42*c^(9/4)*Sqrt[a
+ c*x^4]) - (a^(3/4)*d*e*(B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^
4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]
)/(2*c^(5/4)*Sqrt[a + c*x^4]) - (3*a^(5/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqr
t[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c
^(1/4)*x)/a^(1/4)], 1/2]]/(10*c^(7/4)*Sqrt[a + c*x^4]) + (a^(1/4)*d^2*(B*d
+ 3*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2
]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(2*c^(3/4)*Sqrt[a + c*x^4]
)
```

Rule 226

```
Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2]]/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 311

```
Int[(x_)^2/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, D
ist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a +
b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 327

```
Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 1210

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2]]/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

Rule 1735

```
Int[(Px)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + cx^4}} dx &= \int \left(\frac{Ad^3}{\sqrt{a + cx^4}} + \frac{d^2(Bd + 3Ae)x^2}{\sqrt{a + cx^4}} + \frac{3de(Bd + Ae)x^4}{\sqrt{a + cx^4}} + \frac{e^2(3Bd + Ae)x^6}{\sqrt{a + cx^4}} + \dots \right) \\
&= (Ad^3) \int \frac{1}{\sqrt{a + cx^4}} dx + (Be^3) \int \frac{x^8}{\sqrt{a + cx^4}} dx + (3de(Bd + Ae)) \int \frac{x^4}{\sqrt{a + cx^4}} dx + \dots \\
&= \frac{de(Bd + Ae)x\sqrt{a + cx^4}}{c} + \frac{e^2(3Bd + Ae)x^3\sqrt{a + cx^4}}{5c} + \frac{Be^3x^5\sqrt{a + cx^4}}{7c} + \dots \\
&= -\frac{5aBe^3x\sqrt{a + cx^4}}{21c^2} + \frac{de(Bd + Ae)x\sqrt{a + cx^4}}{c} + \frac{e^2(3Bd + Ae)x^3\sqrt{a + cx^4}}{5c} + \dots \\
&= -\frac{5aBe^3x\sqrt{a + cx^4}}{21c^2} + \frac{de(Bd + Ae)x\sqrt{a + cx^4}}{c} + \frac{e^2(3Bd + Ae)x^3\sqrt{a + cx^4}}{5c} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.19, size = 217, normalized size = 0.48

$$\frac{ex(a + cx^4)(-25aBe^2 + 21Ace(5d + ex^2) + 3Bc(35d^2 + 21dex^2 + 5e^2x^4)) + 5(21Acd(cx^2 - ae^2) + aBe(-21cd^2 + 5ae^2))x\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; \frac{3}{2}; -\frac{cx^4}{a}\right) + 7c(5Bcd^3 + 15Acd^2e - 9aBde^2 - 3aAe^3)x^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{7}{4}; -\frac{cx^4}{a}\right)}{105c^2\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + c*x^4], x]

[Out] (e*x*(a + c*x^4)*(-25*a*B*e^2 + 21*A*c*e*(5*d + e*x^2) + 3*B*c*(35*d^2 + 21*d*e*x^2 + 5*e^2*x^4)) + 5*(21*A*c*d*(c*d^2 - a*e^2) + a*B*e*(-21*c*d^2 + 5*a*e^2))*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 7*c*(5*B*c*d^3 + 15*A*c*d^2*e - 9*a*B*d*e^2 - 3*a*A*e^3)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)]/(105*c^2*Sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.18, size = 533, normalized size = 1.18

method	result
--------	--------

elliptic	$\frac{B e^3 x^5 \sqrt{c x^4 + a}}{7c} + \frac{(A e^3 + 3B d e^2) x^3 \sqrt{c x^4 + a}}{5c} + \frac{(3A d e^2 + 3B d^2 e - \frac{5a B e^3}{7c}) x \sqrt{c x^4 + a}}{3c} + \frac{\left(A d^3 - \frac{a(3A d e^2 + 3B d^2 e - \frac{5a B e^3}{7c})}{3c} \right)}{3c^2}$
risch	$\frac{e x (15 B e^2 x^4 c + 21 A c e^2 x^2 + 63 B c d e x^2 + 105 c d e A - 25 B a e^2 + 105 B c d^2) \sqrt{c x^4 + a}}{105 c^2} - \frac{i(63 A a c e^3 - 315 A c^2 d^2 e + 189 B a c d e^2 - 105 B c^2 d^3)}{105 c^2}$
default	$B e^3 \left(\frac{x^5 \sqrt{c x^4 + a}}{7c} - \frac{5 a x \sqrt{c x^4 + a}}{21 c^2} + \frac{5 a^2 \sqrt{1 - \frac{i \sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i \sqrt{c} x^2}{\sqrt{a}}}}{21 c^2} \frac{\text{EllipticF}\left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{c x^4 + a}} \right) + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$B e^3 \left(\frac{1}{7} \frac{x^5 \sqrt{c x^4 + a}}{c^2} - \frac{5}{21} \frac{a x \sqrt{c x^4 + a}}{c^2} + \frac{5 a^2 \sqrt{1 - \frac{i \sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i \sqrt{c} x^2}{\sqrt{a}}}}{21 c^2} \frac{\text{EllipticF}\left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{c x^4 + a}} \right) + \dots$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)^3/sqrt(c*x^4+a),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)*(x^2*e + d)^3/sqrt(c*x^4 + a), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)
```

Sympy [C] Result contains complex when optimal does not.

time = 3.89, size = 364, normalized size = 0.80

$$\frac{Ae^{\frac{1}{2}\pi i} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{3Ad^2e^{\frac{3}{2}\pi i} {}_2F_1\left(\frac{3}{2}, \frac{3}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{3Ade^{\frac{5}{2}\pi i} {}_2F_1\left(\frac{5}{2}, \frac{5}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{Ae^{\frac{7}{2}\pi i} {}_2F_1\left(\frac{7}{2}, \frac{7}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{Bd^3e^{\frac{1}{2}\pi i} {}_2F_1\left(\frac{1}{2}, \frac{3}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{3Bd^2e^{\frac{3}{2}\pi i} {}_2F_1\left(\frac{3}{2}, \frac{3}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{3Bde^{\frac{5}{2}\pi i} {}_2F_1\left(\frac{5}{2}, \frac{5}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{Be^{\frac{7}{2}\pi i} {}_2F_1\left(\frac{7}{2}, \frac{7}{2}\right) \sqrt{\frac{a-dx}{a}}}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+a)**(1/2),x)
```

```
[Out] A*d**3*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + 3*A*d**2*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + 3*A*d*e**2*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + A*e**3*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4)) + B*d**3*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + 3*B*d**2*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + 3*B*d*e**2*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4)) + B*e**3*x**9*gamma(9/4)*hyper((1/2, 9/4), (13/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(13/4))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/sqrt(c*x^4 + a), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(1/2), x)
```

```
[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(1/2), x)
```

$$3.2 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{\sqrt{a+cx^4}} dx$$

Optimal. Leaf size=367

$$\frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} + \frac{(5Bcd^2 + 10Acde - 3aBe^2)x\sqrt{a + cx^4}}{5c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(5Bcd^2 + 10Acde - 3aBe^2)}{5c^{3/2}(\sqrt{a} + \sqrt{c}x^2)}$$

```
[Out] 1/3*e*(A*e+2*B*d)*x*(c*x^4+a)^(1/2)/c+1/5*B*e^2*x^3*(c*x^4+a)^(1/2)/c+1/5*(
10*A*c*d*e-3*B*a*e^2+5*B*c*d^2)*x*(c*x^4+a)^(1/2)/c^(3/2)/(a^(1/2)+x^2*c^(1
/2))-1/5*a^(1/4)*(10*A*c*d*e-3*B*a*e^2+5*B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a
^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c
^(1/4)*x/a^(1/4))),1/2)*2^(1/2)*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x
^2*c^(1/2)))^(1/2)/c^(7/4)/(c*x^4+a)^(1/2)+1/30*(cos(2*arctan(c^(1/4)*x/a
^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c
^(1/4)*x/a^(1/4))),1/2)*2^(1/2)*(15*A*c^(3/2)*d^2-9*a^(3/2)*B*e^2+15*c*d*(2
*A*e+B*d)*a^(1/2)-5*a*e*(A*e+2*B*d)*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+
a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(7/4)/(c*x^4+a)^(1/2)
```

Rubi [A]

time = 0.25, antiderivative size = 706, normalized size of antiderivative = 1.92, number of steps used = 12, number of rules used = 5, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.179$,

Rules used = {1735, 226, 311, 1210, 327}

1735: Int[(A + B*x^2)*(d + e*x^2)^2/Sqrt[a + c*x^4], x] -> (e*(2*B*d + A*e)*x*Sqrt[a + c*x^4])/(3*c) + (B*e^2*x^3*Sqrt[a + c*x^4])/(5*c) - (3*a*B*e^2*x*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) + (d*(B*d + 2*A*e)*x*Sqrt[a + c*x^4])/(Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) + (3*a^(5/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(5*c^(7/4)*Sqrt[a + c*x^4]) - (a^(1/4)*d*(B*d + 2*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(3/4)*Sqrt[a + c*x^4]) + (A*d^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]) - (3*a^(5/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(10*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*e*(2*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(5/4)*Sqrt[a + c*x^4]) + (a^(1/4)*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2))

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)*(d + e*x^2)^2/Sqrt[a + c*x^4], x]

```
[Out] (e*(2*B*d + A*e)*x*Sqrt[a + c*x^4])/(3*c) + (B*e^2*x^3*Sqrt[a + c*x^4])/(5*c) - (3*a*B*e^2*x*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) + (d*(B*d + 2*A*e)*x*Sqrt[a + c*x^4])/(Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) + (3*a^(5/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(5*c^(7/4)*Sqrt[a + c*x^4]) - (a^(1/4)*d*(B*d + 2*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(3/4)*Sqrt[a + c*x^4]) + (A*d^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(1/4)*Sqrt[a + c*x^4]) - (3*a^(5/4)*B*e^2*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(10*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*e*(2*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)]^2*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(5/4)*Sqrt[a + c*x^4]) + (a^(1/4)*Sqrt[a + c*x^4])/(5*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2))
```

)*d*(B*d + 2*A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]/(2*c^(3/4)*Sqrt[a + c*x^4])

Rule 226

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 311

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 1210

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 1735

Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + cx^4}} dx &= \int \left(\frac{Ad^2}{\sqrt{a + cx^4}} + \frac{d(Bd + 2Ae)x^2}{\sqrt{a + cx^4}} + \frac{e(2Bd + Ae)x^4}{\sqrt{a + cx^4}} + \frac{Be^2x^6}{\sqrt{a + cx^4}} \right) dx \\
&= (Ad^2) \int \frac{1}{\sqrt{a + cx^4}} dx + (Be^2) \int \frac{x^6}{\sqrt{a + cx^4}} dx + (e(2Bd + Ae)) \int \frac{x^4}{\sqrt{a + cx^4}} dx \\
&= \frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} + \frac{Ad^2(\sqrt{a} + \sqrt{c}x^2)}{2\sqrt[4]{a}} \sqrt{\frac{a + cx^4}{a}} \\
&= \frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} + \frac{d(Bd + 2Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{Ad^2}{2\sqrt[4]{a}} \\
&= \frac{e(2Bd + Ae)x\sqrt{a + cx^4}}{3c} + \frac{Be^2x^3\sqrt{a + cx^4}}{5c} - \frac{3aBe^2x\sqrt{a + cx^4}}{5c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} + \frac{d(Bd + 2Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.12, size = 159, normalized size = 0.43

$$\frac{ex(10Bd + 5Ae + 3Be^2)(a + cx^4) - 5(-3Acd^2 + 2ABde + aAe^2)x\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a}\right) + (5Bcd^2 + 10Acde - 3aBe^2)x^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{cx^4}{a}\right)}{15c\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + c*x^4], x]

[Out] (e*x*(10*B*d + 5*A*e + 3*B*e*x^2)*(a + c*x^4) - 5*(-3*A*c*d^2 + 2*a*B*d*e + a*A*e^2)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + (5*B*c*d^2 + 10*A*c*d*e - 3*a*B*e^2)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)])/(15*c*Sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.15, size = 403, normalized size = 1.10

method	result
elliptic	$ \frac{Be^2x^3\sqrt{cx^4 + a}}{5c} + \frac{(Ae^2 + 2Bde)x\sqrt{cx^4 + a}}{3c} + \frac{\left(Ad^2 - \frac{a(Ae^2 + 2Bde)}{3c}\right)\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \text{Elliptic} $

risch	$\frac{ex(3Be^2x^2+5Ae+10Bd)\sqrt{cx^4+a}}{15c} - \frac{i(30cdeA-9Ba^2e^2+15Bcd^2)\sqrt{a}\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{c}}$
default	$B e^2 \left(\frac{x^3\sqrt{cx^4+a}}{5c} - \frac{3ia^{\frac{3}{2}}\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right),i\right)-\text{EllipticE}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\right),i\right)}{5c^{\frac{3}{2}}\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $B e^2 \left(\frac{1}{5} \frac{x^3 \sqrt{c x^4 + a}}{c} - \frac{3}{5} \frac{I a^{3/2}}{c^{3/2}} \frac{\sqrt{1 - \frac{i \sqrt{c} x^2}{\sqrt{a}}}}{\sqrt{1 + \frac{i \sqrt{c} x^2}{\sqrt{a}}}} \left(\text{EllipticF} \left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}} \right), I \right) - \text{EllipticE} \left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}} \right), I \right) + (A e^2 + 2 B d e) \frac{1}{3} \frac{x \sqrt{c x^4 + a}}{c} - \frac{1}{3} \frac{a}{c} \frac{\sqrt{1 - \frac{i \sqrt{c} x^2}{\sqrt{a}}}}{\sqrt{1 + \frac{i \sqrt{c} x^2}{\sqrt{a}}}} \left(\text{EllipticF} \left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}} \right), I \right) + I (2 A d e + B d^2) \frac{a^{1/2}}{\sqrt{1 - \frac{i \sqrt{c} x^2}{\sqrt{a}}}} \frac{\sqrt{1 + \frac{i \sqrt{c} x^2}{\sqrt{a}}}}{\sqrt{c x^4 + a}} \left(\text{EllipticF} \left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}} \right), I \right) - \text{EllipticE} \left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}} \right), I \right) + A d^2 \frac{1}{\sqrt{1 - \frac{i \sqrt{c} x^2}{\sqrt{a}}}} \frac{\sqrt{1 + \frac{i \sqrt{c} x^2}{\sqrt{a}}}}{\sqrt{c x^4 + a}} \left(\text{EllipticF} \left(x \sqrt{\frac{i \sqrt{c}}{\sqrt{a}}} \right), I \right) \right)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)^2/sqrt(c*x^4+a),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)*(x^2*e + d)^2/sqrt(c*x^4 + a), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [C] Result contains complex when optimal does not.
time = 2.88, size = 262, normalized size = 0.71

$$\frac{Ad^2x\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{Ade^2x^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{2\sqrt{a}\Gamma\left(\frac{1}{4}\right)} + \frac{Ae^2x^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{Bd^2x^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{1}{4}\right)} + \frac{Bde^2x^5\Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{2\sqrt{a}\Gamma\left(\frac{3}{4}\right)} + \frac{Be^2x^7\Gamma\left(\frac{7}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| \frac{cx^4e^{i\pi}}{a}\right)}{4\sqrt{a}\Gamma\left(\frac{11}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+a)**(1/2), x)

[Out] A*d**2*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + A*d*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(2*sqrt(a)*gamma(7/4)) + A*e**2*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4)) + B*d**2*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*d*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(2*sqrt(a)*gamma(9/4)) + B*e**2*x**7*gamma(7/4)*hyper((1/2, 7/4), (11/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(11/4))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(1/2), x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^2/sqrt(c*x^4 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(1/2), x)

[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(1/2), x)

3.3 $\int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+cx^4}} dx$

Optimal. Leaf size=277

$$\frac{Bex\sqrt{a+cx^4}}{3c} + \frac{(Bd+ Ae)x\sqrt{a+cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(Bd+ Ae)(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}}{c^{3/4}\sqrt{a+cx^4}} E\left(2 \tan^{-1}\left(\frac{\sqrt{a+cx^4}}{\sqrt{a} + \sqrt{c}x^2}\right)\right)$$

[Out] $1/3*B*e*x*(c*x^4+a)^{(1/2)}/c+(A*e+B*d)*x*(c*x^4+a)^{(1/2)}/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-a^{(1/4)}*(A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}+1/6*a^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*((3*A*c*d-B*a*e)/a^{(1/2)}+3*(A*e+B*d)*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(5/4)}/(c*x^4+a)^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 432, normalized size of antiderivative = 1.56, number of steps used = 8, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {1735, 226, 311, 1210, 327}

$$\frac{a^{3/4}Be(\sqrt{a} + \sqrt{c}x) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{6c^{3/4}\sqrt{a+cx^4}} + \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{c}x) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} (Ae + Bd) F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{2c^{3/4}\sqrt{a+cx^4}} - \frac{\sqrt[4]{a}(\sqrt{a} + \sqrt{c}x) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} (Ae + Bd) E\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{c^{3/4}\sqrt{a+cx^4}} + \frac{Ad(\sqrt{a} + \sqrt{c}x) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\right)}{2\sqrt[4]{a}\sqrt{c}\sqrt{a+cx^4}} + \frac{x\sqrt{a+cx^4}(Ae + Bd)}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} + \frac{Bex\sqrt{a+cx^4}}{3c}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(A + B*x^2)*(d + e*x^2)/\text{Sqrt}[a + c*x^4], x]$

[Out] $(B*e*x*\text{Sqrt}[a + c*x^4])/(3*c) + ((B*d + A*e)*x*\text{Sqrt}[a + c*x^4])/(\text{Sqrt}[c]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (a^{(1/4)}*(B*d + A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((c^{(3/4)}*\text{Sqrt}[a + c*x^4]) + (A*d*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((2*a^{(1/4)}*c^{(1/4)}*\text{Sqrt}[a + c*x^4]) - (a^{(3/4)}*B*e*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2]))/(6*c^{(5/4)}*\text{Sqrt}[a + c*x^4]) + (a^{(1/4)}*(B*d + A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(2*c^{(3/4)}*\text{Sqrt}[a + c*x^4])$

Rule 226

$\text{Int}[1/\text{Sqrt}[(a_) + (b_.)*(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*\text{Sqrt}[a + b*x^4]))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$

Rule 311

$\text{Int}[(x_)^2/\text{Sqrt}[(a_) + (b_.)(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[b/a, 2]\}, \text{Dist}[1/q, \text{Int}[1/\text{Sqrt}[a + b*x^4], x], x] - \text{Dist}[1/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^4], x], x] \text{ /; FreeQ}\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$

Rule 327

$\text{Int}[((c_.)(x_)^m)*((a_) + (b_.)(x_)^n)^p, x_Symbol] \text{ :> Simp}[c^{(n-1)}*(c*x)^{m-n+1}*((a + b*x^n)^{p+1}/(b*(m+n*p+1))), x] - \text{Dist}[a*c^n*((m-n+1)/(b*(m+n*p+1))), \text{Int}[(c*x)^{m-n}*(a + b*x^n)^p, x], x] \text{ /; FreeQ}\{a, b, c, p\}, x] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, n-1] \ \&\& \ \text{NeQ}[m+n*p+1, 0] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 1210

$\text{Int}[(d_) + (e_.)(x_)^2]/\text{Sqrt}[(a_) + (c_.)(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*\text{Sqrt}[a + c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] \text{ /; EqQ}[e + d*q^2, 0]] \text{ /; FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$

Rule 1735

$\text{Int}[(Px_)*((d_) + (e_.)(x_)^2)^q*(a_) + (c_.)(x_)^4)^p, x_Symbol] \text{ :> Int}[\text{ExpandIntegrand}[1/\text{Sqrt}[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^{p+1/2}], x] \text{ /; FreeQ}\{a, c, d, e\}, x] \ \&\& \ \text{PolyQ}[Px, x^2] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{IntegerQ}[p + 1/2] \ \&\& \ \text{IntegerQ}[q]$

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + cx^4}} dx &= \int \left(\frac{Ad}{\sqrt{a + cx^4}} + \frac{(Bd + Ae)x^2}{\sqrt{a + cx^4}} + \frac{Bex^4}{\sqrt{a + cx^4}} \right) dx \\ &= (Ad) \int \frac{1}{\sqrt{a + cx^4}} dx + (Be) \int \frac{x^4}{\sqrt{a + cx^4}} dx + (Bd + Ae) \int \frac{x^2}{\sqrt{a + cx^4}} dx \\ &= \frac{Bex\sqrt{a + cx^4}}{3c} + \frac{Ad(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\right)}{2\sqrt[4]{a}\sqrt[4]{c}\sqrt{a + cx^4}} \\ &= \frac{Bex\sqrt{a + cx^4}}{3c} + \frac{(Bd + Ae)x\sqrt{a + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(Bd + Ae)(\sqrt{a} + \sqrt{c}x^2)}{c^3} \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.07, size = 121, normalized size = 0.44

$$\frac{Bex(a + cx^4) + (3Acd - aBe)x\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a}\right) + c(Bd + Ae)x^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{2}, \frac{3}{4}, \frac{7}{4}, -\frac{cx^4}{a}\right)}{3c\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/Sqrt[a + c*x^4], x]

[Out] (B*e*x*(a + c*x^4) + (3*A*c*d - a*B*e)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -(c*x^4)/a] + c*(B*d + A*e)*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/2, 3/4, 7/4, -(c*x^4)/a])/(3*c*Sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.14, size = 269, normalized size = 0.97

method	result
elliptic	$\frac{Bex\sqrt{cx^4+a}}{3c} + \frac{(Ad - \frac{aBe}{3c})\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} + \frac{i(Ae+Bd)\sqrt{a}\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{cx^4+a}}$
default	$Be\left(\frac{x\sqrt{cx^4+a}}{3c} - \frac{a\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{3c\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}\right) + \frac{i(Ae+Bd)\sqrt{a}\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{cx^4+a}}$
risch	$\frac{Bex\sqrt{cx^4+a}}{3c} + \frac{i(3Ace+3Bcd)\sqrt{a}\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - \text{EllipticE}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)\right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{c}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2), x, method=_RETURNVERBOSE)

[Out] B*e*(1/3*x*(c*x^4+a)^(1/2)/c-1/3*a/c/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2), I)+I*(A*e+B*d)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2), I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2), I))+A*d/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2), I)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="maxima")``[Out] integrate((B*x^2 + A)*(x^2*e + d)/sqrt(c*x^4 + a), x)`**Fricas [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="fricas")``[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)`**Sympy [C] Result contains complex when optimal does not.**

time = 1.98, size = 167, normalized size = 0.60

$$\frac{A dx \Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{5}{4}\right)} + \frac{A e x^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)} + \frac{B dx^3 \Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)} + \frac{B e x^5 \Gamma\left(\frac{5}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{5}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{9}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+a)**(1/2),x)`

```
[Out] A*d*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + A*e*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*d*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4)) + B*e*x**5*gamma(5/4)*hyper((1/2, 5/4), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(9/4))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="giac")`

[Out] integrate((B*x^2 + A)*(x^2*e + d)/sqrt(c*x^4 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(1/2), x)

[Out] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(1/2), x)

3.4 $\int \frac{A+Bx^2}{\sqrt{a+cx^4}} dx$

Optimal. Leaf size=226

$$\frac{Bx\sqrt{a+cx^4}}{\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)} - \frac{\sqrt[4]{a}B(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}E\left(2\tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{c^{3/4}\sqrt{a+cx^4}} + \frac{\sqrt[4]{a}\left(B+\frac{A\sqrt{c}}{\sqrt{a}}\right)}{\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)}$$

[Out] $B*x*(c*x^4+a)^{(1/2)}/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-a^{(1/4)}*B*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}+1/2*a^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(B+A*c^{(1/2)}/a^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {1212, 226, 1210}

$$\frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}\left(\frac{A\sqrt{c}}{\sqrt{a}}+B\right)F\left(2\text{ArcTan}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{2c^{3/4}\sqrt{a+cx^4}} - \frac{\sqrt[4]{a}B(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}E\left(2\text{ArcTan}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{2}\right)}{c^{3/4}\sqrt{a+cx^4}} + \frac{Bx\sqrt{a+cx^4}}{\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(A + B*x^2)/\text{Sqrt}[a + c*x^4], x]$

[Out] $(B*x*\text{Sqrt}[a + c*x^4])/(\text{Sqrt}[c]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (a^{(1/4)}*B*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (c^{(3/4)}*\text{Sqrt}[a + c*x^4]) + (a^{(1/4)}*(B + (A*\text{Sqrt}[c])/ \text{Sqrt}[a])* (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (2*c^{(3/4)}*\text{Sqrt}[a + c*x^4])$

Rule 226

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*\text{Sqrt}[a + b*x^4]))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[b/a]$

Rule 1210

$\text{Int}[((d_) + (e_)*(x_)^2)/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*$

```
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0]] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

Rule 1212

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0]] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rubi steps

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4}} dx = \left(A + \frac{\sqrt{a} B}{\sqrt{c}} \right) \int \frac{1}{\sqrt{a + cx^4}} dx - \frac{(\sqrt{a} B) \int \frac{1 - \frac{\sqrt{c} x^2}{\sqrt{a}}}{\sqrt{a + cx^4}} dx}{\sqrt{c}}$$

$$= \frac{Bx\sqrt{a + cx^4}}{\sqrt{c} (\sqrt{a} + \sqrt{c} x^2)} - \frac{{}^4\sqrt{a} B (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} E\left(2 \tan^{-1}\left(\frac{{}^4\sqrt{c} x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right)}{c^{3/4} \sqrt{a + cx^4}}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.04, size = 77, normalized size = 0.34

$$\frac{\sqrt{1 + \frac{cx^4}{a}} \left(3Ax {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) + Bx^3 {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{7}{4}; -\frac{cx^4}{a}\right) \right)}{3\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(A + B*x^2)/Sqrt[a + c*x^4], x]
```

```
[Out] (Sqrt[1 + (c*x^4)/a]*(3*A*x*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)]
+ B*x^3*Hypergeometric2F1[1/2, 3/4, 7/4, -((c*x^4)/a)]))/(3*Sqrt[a + c*x^4]
)
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 169, normalized size = 0.75

method	result
--------	--------

default	$\frac{iB\sqrt{a} \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} \left(\text{EllipticF}\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - \text{EllipticE}\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) \right) + A\sqrt{1 - \frac{i\sqrt{c}}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} \sqrt{c}}$
elliptic	$\frac{iB\sqrt{a} \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} \left(\text{EllipticF}\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - \text{EllipticE}\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) \right) + A\sqrt{1 - \frac{i\sqrt{c}}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a} \sqrt{c}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $I*B*a^{1/2}/(I/a^{1/2}*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}*(1+I/a^{1/2}*c^{1/2}*x^2)^{1/2}/(c*x^4+a)^{1/2}/c^{1/2}*(\text{EllipticF}(x*(I/a^{1/2})*c^{1/2}),I)-\text{EllipticE}(x*(I/a^{1/2})*c^{1/2}),I)+A/(I/a^{1/2}*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}*(1+I/a^{1/2}*c^{1/2}*x^2)^{1/2}/(c*x^4+a)^{1/2}*\text{EllipticF}(x*(I/a^{1/2})*c^{1/2}),I)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)/sqrt(c*x^4 + a), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Symbolic function `elliptic_ec` takes exactly 1 arguments (2 given)

Sympy [C] Result contains complex when optimal does not.

time = 0.93, size = 78, normalized size = 0.35

$$\frac{Ax\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\frac{1}{4}, \frac{1}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{5}{4}\right)} + \frac{Bx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\frac{1}{2}, \frac{3}{4} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4\sqrt{a} \Gamma\left(\frac{7}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(c*x**4+a)**(1/2),x)

[Out] A*x*gamma(1/4)*hyper((1/4, 1/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(5/4)) + B*x**3*gamma(3/4)*hyper((1/2, 3/4), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*sqrt(a)*gamma(7/4))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/sqrt(c*x^4 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{\sqrt{c x^4 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/(a + c*x^4)^(1/2),x)

[Out] int((A + B*x^2)/(a + c*x^4)^(1/2), x)

$$3.5 \quad \int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+cx^4}} dx$$

Optimal. Leaf size=369

$$\frac{(Bd - Ae) \tan^{-1} \left(\frac{\sqrt{cd^2 + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + cx^4}} \right) (\sqrt{a} B - A\sqrt{c}) (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} F\left(2 \tan^{-1} \left(\frac{\sqrt{cd^2 + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + cx^4}} \right), \frac{1}{2} \right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 + ae^2} \cdot 2\sqrt[4]{a} \sqrt[4]{c} (\sqrt{c} d - \sqrt{a} e) \sqrt{a + cx^4}}$$

[Out] $-1/2*(-A*e+B*d)*\arctan(x*(a*e^2+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+a)^{(1/2)})/d^{(1/2)}/e^{(1/2)}/(a*e^2+c*d^2)^{(1/2)}-1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)})))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*2^{(1/2)})*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}+1/4*a^{(3/4)}*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)})))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)}, 1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(1/4)}/d/e/(-a*e^2+c*d^2)/(c*x^4+a)^{(1/2)}$

Rubi [A]

time = 0.24, antiderivative size = 369, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {1723, 226, 1721}

$$\frac{a^{3/4}(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} \left(\frac{\sqrt{c}d + e}{\sqrt{a}} \right)^2 (Bd - Ae) \text{III} \left(-\frac{(\sqrt{c}d - \sqrt{a}e)^2}{\sqrt{a} \sqrt{c}de}; 2 \text{ArcTan} \left(\frac{\sqrt{c}x}{\sqrt{a}} \right) \right) \frac{1}{2} - \frac{(Bd - Ae) \text{ArcTan} \left(\frac{\sqrt{a} \sqrt{c} x^2 + cd^2}{\sqrt{d} \sqrt{e} \sqrt{a + cx^4}} \right)}{2\sqrt{d} \sqrt{e} \sqrt{ae^2 + cd^2}} - \frac{(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} (\sqrt{a} B - A\sqrt{c}) F \left(2 \text{ArcTan} \left(\frac{\sqrt{c}x}{\sqrt{a}} \right) \right) \frac{1}{2}}{2\sqrt[4]{a} \sqrt[4]{c} \sqrt{a + cx^4} (\sqrt{c}d - \sqrt{a}e)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x]

[Out] $-1/2*((B*d - A*e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])])/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[c*d^2 + a*e^2]) - ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])* (\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((2*a^{(1/4)}*c^{(1/4)}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*\text{Sqrt}[a + c*x^4]) + (a^{(3/4)}*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2))/(4*c^{(1/4)}*d*e*(c*d^2 - a*e^2)*\text{Sqrt}[a + c*x^4])$

Rule 226

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] :> With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4])]*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 1721

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e)
) + a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2]), x]
+ Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*((a + c*x^4)/(a*(A + B*x^2)^2))]/(
4*d*e*A*q*Sqrt[a + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)],
2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e
^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1723

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e
+ d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x],
x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2
- a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rubi steps

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + cx^4}} dx = -\frac{(\sqrt{a}B - A\sqrt{c}) \int \frac{1}{\sqrt{a + cx^4}} dx}{\sqrt{c}d - \sqrt{a}e} + \frac{(\sqrt{a}(Bd - Ae)) \int \frac{1 + \frac{\sqrt{c}x^2}{\sqrt{a}}}{(d + ex^2)\sqrt{a + cx^4}} dx}{\sqrt{c}d - \sqrt{a}e}$$

$$= -\frac{(Bd - Ae) \tan^{-1}\left(\frac{\sqrt{cd^2 + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a + cx^4}}\right)}{2\sqrt{d}\sqrt{e}\sqrt{cd^2 + ae^2}} - \frac{(\sqrt{a}B - A\sqrt{c})(\sqrt{a} + \sqrt{c}x^2)}{2\sqrt{a}\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.22, size = 138, normalized size = 0.37

$$\frac{i\sqrt{1 + \frac{cx^4}{a}} \left(BdF \left(i \sinh^{-1} \left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x \right) \middle| -1 \right) + (-Bd + Ae) \Pi \left(-\frac{i\sqrt{a}e}{\sqrt{c}d}; i \sinh^{-1} \left(\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} x \right) \middle| -1 \right) \right)}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} de\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x]

[Out] $((-I)\sqrt{1 + (c*x^4)/a}*(B*d*EllipticF[I*ArcSinh[\sqrt{(I*\sqrt{c})}/\sqrt{a}]]*x, -1) + (-B*d) + A*e)*EllipticPi[((-I)\sqrt{a}*e)/(\sqrt{c}*d), I*ArcSinh[\sqrt{(I*\sqrt{c})}/\sqrt{a}]]*x, -1))/(\sqrt{(I*\sqrt{c})}/\sqrt{a})*d*e*\sqrt{a + c*x^4})$

Maple [C] Result contains complex when optimal does not.

time = 0.15, size = 192, normalized size = 0.52

method	result
default	$\frac{B\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{e\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} + \frac{(Ae - Bd)\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\text{EllipticPi}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{ed\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}}$
elliptic	$\frac{B\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{e\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\text{EllipticPi}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{d\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $B/e/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}, I) + (A*e - B*d)/e/d/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}*(1-I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}*(1+I/a^{(1/2)*c^{(1/2)}}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticPi(x*(I/a^{(1/2)*c^{(1/2)}})^{(1/2)}, I*a^{(1/2)}/c^{(1/2)}*e/d, (-I/a^{(1/2)*c^{(1/2)}})^{(1/2)}/(I/a^{(1/2)*c^{(1/2)}})^{(1/2)})$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(x^2*e + d)), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+a)**(1/2),x)

[Out] Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(x^2*e + d)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a} (ex^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)),x)

[Out] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)), x)

$$\frac{\sqrt[4]{x}/\sqrt[4]{a}, 1/2]}{(2a^{1/4}d(\sqrt{c}d - \sqrt{a}e)\sqrt{a + cx^4}) + ((\sqrt{c}d + \sqrt{a}e)(Bcd^3 - 3Acd^2e - aBde^2 - aAe^3)(\sqrt{a} + \sqrt{c}x^2)\sqrt{(a + cx^4)/(\sqrt{a} + \sqrt{c}x^2)^2})\text{EllipticPi}[-1/4(\sqrt{c}d - \sqrt{a}e)^2/(\sqrt{a}\sqrt{c}de), 2\text{ArcTan}[(c^{1/4}x)/\sqrt[4]{a}], 1/2)]/(8a^{1/4}c^{1/4}d^2e(\sqrt{c}d - \sqrt{a}e)(cd^2 + ae^2)\sqrt{a + cx^4})}$$

Rule 226

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 1210

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]
```

Rule 1711

```
Int[((P4x)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + c*x^4]/(2*d*(q + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), Int[((d + e*x^2)^(q + 1)/Sqrt[a + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*c*d^2*(q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[c*d^2 + a*e^2, 0] && ILtQ[q, -1]
```

Rule 1721

```
Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*((a + c*x^4)/(a*(A + B*x^2)^2))]/(4*d*e*A*q*Sqrt[a + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1723

```
Int[((A_.) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q
```

))/((c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]

Rule 1729

Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :>
 With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2]/((d + e*x^2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]

Rubi steps

$$\begin{aligned} \int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{\int \frac{-2Acd^2 - aBde - aAe^2 - 2cd(Bd - Ae)x^2 - ce(Bd - Ae)x^4}{(d + ex^2)\sqrt{a + cx^4}} dx}{2d(cd^2 + ae^2)} \\ &= -\frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{\int \frac{-\sqrt{a}c^{3/2}de(Bd - Ae) + ce(-2Acd^2 - aBde - aAe^2) + (-2c^2d}{(d + ex^2)\sqrt{a + cx^4}}}{2cde(cd^2 + ae^2)}}{2d(cd^2 + ae^2)(d + ex^2)} \\ &= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{\sqrt{a}\sqrt{c}(Bd - Ae)}{2cde(cd^2 + ae^2)} \\ &= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + cx^4}}{2d(cd^2 + ae^2)(d + ex^2)} - \frac{(Bcd^3 - 3Acd^2e)}{2cde(cd^2 + ae^2)} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.58, size = 297, normalized size = 0.46

$$\frac{\frac{de(-Bd + Ae)x(a + cx^4)}{(cd^2 + ae^2)(d + ex^2)} - \frac{i\sqrt{1 + \frac{cx^4}{a}} \left(i\sqrt{a}\sqrt{c}de(Bd - Ae)E\left(i\sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{a}}z\right)\right) - 1 \right) + \sqrt{c}d(\sqrt{c}d - i\sqrt{a}e)(Bd - Ae)F\left(i\sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{a}}z\right)\right) - 1 \right) + (-Bcd^3 + 3Acd^2e + aBde^2 + aAe^3)\Pi\left(\frac{i\sqrt{a}e}{\sqrt{c}d}; i\sinh^{-1}\left(\sqrt{\frac{i\sqrt{c}}{a}}z\right)\right) - 1 \right)}{\sqrt{\frac{i\sqrt{c}}{a}}(cd^2e + ae^2)}}{2d^2\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + c*x^4]), x]

```
[Out] ((d*e*(-(B*d) + A*e)*x*(a + c*x^4))/((c*d^2 + a*e^2)*(d + e*x^2)) - (I*Sqrt
[1 + (c*x^4)/a]*(I*Sqrt[a]*Sqrt[c]*d*e*(B*d - A*e)*EllipticE[I*ArcSinh[Sqrt
[(I*Sqrt[c])/Sqrt[a]]*x], -1] + Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(B*d -
A*e)*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (-B*c*d^3) +
3*A*c*d^2*e + a*B*d*e^2 + a*A*e^3)*EllipticPi[((-I)*Sqrt[a]*e)/(Sqrt[c]*d),
I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1]))/(Sqrt[(I*Sqrt[c])/Sqrt[a]]*(
c*d^2*e + a*e^3)))/(2*d^2*Sqrt[a + c*x^4])
```

Maple [C] Result contains complex when optimal does not.

time = 0.16, size = 679, normalized size = 1.06

method	result
default	$\frac{B \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}} \operatorname{EllipticPi}\left(x \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, \frac{i\sqrt{a} e}{\sqrt{c} d}, \sqrt{\frac{-i\sqrt{c}}{\sqrt{a}}}\right) + (Ae - Bd) \frac{e^2 x \sqrt{c x^4 + a}}{2d(a e^2 + c d^2)(e x^2 + d)}}{ed \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{c x^4 + a}}$
elliptic	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] B/e/d/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)
)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2)
,I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)
)+(A*e-B*d)/e*(1/2*e^2*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)/(e*x^2+d)-1/2*c/(a
*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/
a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(
1/2),I)-1/2*I*c^(1/2)*e/d/(a*e^2+c*d^2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*
(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(
1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*I*c^(1/2)*e/d/(a*e^2+c*d^
2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a
^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticE(x*(I/a^(1/2)*c^(1/2))^(
1/2),I)+1/2/d^2/(a*e^2+c*d^2)*e^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^
(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi
(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/
2)/(I/a^(1/2)*c^(1/2))^(1/2))*a+3/2/(a*e^2+c*d^2)/(I/a^(1/2)*c^(1/2))^(1/2)
*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(
1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1
/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*c)
```


Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(x^2*e + d)^2), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+a)**(1/2),x)

[Out] Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)**2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(x^2*e + d)^2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + a} (ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^2),x)

[Out] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^2), x)

$$3.7 \quad \int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+cx^4}} dx$$

Optimal. Leaf size=875

$$\frac{\sqrt{c} (5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3) x \sqrt{a+cx^4}}{8d^2 (cd^2 + ae^2)^2 (\sqrt{a} + \sqrt{c} x^2)} - \frac{e(Bd - Ae)x \sqrt{a+cx^4}}{4d (cd^2 + ae^2) (d + ex^2)^2} - \frac{e(5Bcd^3 - 9Acd^2e - aBde^2 - 3aAe^3)}{8d^2 (cd^2 + ae^2)}$$

[Out] $1/16*(3*A*e*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)-B*(-a^2*d*e^4-10*a*c*d^3*e^2+3*c^2*d^5))*\arctan(x*(a*e^2+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+a)^{(1/2)})/d^{(5/2)}/(a*e^2+c*d^2)^{(5/2)}/e^{(1/2)}-1/4*e*(-A*e+B*d)*x*(c*x^4+a)^{(1/2)}/d/(a*e^2+c*d^2)/(e*x^2+d)^2-1/8*e*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^3)*x*(c*x^4+a)^{(1/2)}/d^2/(a*e^2+c*d^2)^2/(e*x^2+d)+1/8*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^3)*x*c^{(1/2)}*(c*x^4+a)^{(1/2)}/d^2/(a*e^2+c*d^2)^2/(a^{(1/2)}+x^2*c^{(1/2)})-1/8*a^{(1/4)}*c^{(1/4)}*(-3*A*a*e^3-9*A*c*d^2*e-B*a*d*e^2+5*B*c*d^3)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/d^2/(a*e^2+c*d^2)^2/(c*x^4+a)^{(1/2)}-1/32*(3*A*e*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)-B*(-a^2*d*e^4-10*a*c*d^3*e^2+3*c^2*d^5))*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*2^{(1/2)})*(e*a^{(1/2)}+d*c^{(1/2)}))*((a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^3/e/(a*e^2+c*d^2)^2/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}+1/8*c^{(1/4)}*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(4*A*c*d^2+a*e*(3*A*e+B*d)+d*(-A*e+B*d))*a^{(1/2)}*c^{(1/2)}*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/d^2/(a*e^2+c*d^2)/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}$

Rubi [A]

time = 1.17, antiderivative size = 875, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1711, 1729, 1210, 1723, 226, 1721}

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^3*sqrt[a + c*x^4]),x]

[Out] $(\text{sqrt}[c]*(5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*x*\text{sqrt}[a + c*x^4])/((8*d^2*(c*d^2 + a*e^2)^2*(\text{sqrt}[a] + \text{sqrt}[c]*x^2)) - (e*(B*d - A*e)*x*\text{sqrt}[a + c*x^4]))/(4*d*(c*d^2 + a*e^2)*(d + e*x^2)^2) - (e*(5*B*c*d^3 - 9*A*c*d$

$$\begin{aligned} &^2e - a*B*d*e^2 - 3*a*A*e^3)*x*\text{Sqrt}[a + c*x^4]/(8*d^2*(c*d^2 + a*e^2)^2*(\\ &d + e*x^2)) + ((3*A*e*(5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4) - B*(3*c^2*d^5 \\ &- 10*a*c*d^3*e^2 - a^2*d*e^4))*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt} \\ &[e]*\text{Sqrt}[a + c*x^4])]/(16*d^(5/2)*\text{Sqrt}[e]*(c*d^2 + a*e^2)^(5/2)) - (a^(1/4) \\ &)*c^(1/4)*(5*B*c*d^3 - 9*A*c*d^2*e - a*B*d*e^2 - 3*a*A*e^3)*(\text{Sqrt}[a] + \text{Sqrt} \\ &[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c \\ &(1/4)*x)/a^(1/4)], 1/2]/(8*d^2*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) + (c^(1/ \\ &4)*(4*A*c*d^2 + \text{Sqrt}[a]*\text{Sqrt}[c]*d*(B*d - A*e) + a*e*(B*d + 3*A*e))*(\text{Sqrt}[a] \\ &+ \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{Arc} \\ &\text{Tan}[(c^(1/4)*x)/a^(1/4)], 1/2]/(8*a^(1/4)*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d \\ &^2 + a*e^2)*\text{Sqrt}[a + c*x^4]) - ((\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(3*A*e*(5*c^2*d^4 + \\ &2*a*c*d^2*e^2 + a^2*e^4) - B*(3*c^2*d^5 - 10*a*c*d^3*e^2 - a^2*d*e^4))*(\text{S} \\ &\text{qrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi} \\ &[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^(1/4)*x) \\ &/a^(1/4)], 1/2]/(32*a^(1/4)*c^(1/4)*d^3*e*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + \\ &a*e^2)^2*\text{Sqrt}[a + c*x^4]) \end{aligned}$$

Rule 226

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]
```

Rule 1210

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*
EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

Rule 1711

```
Int[((P4x)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol
] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]
}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + c*x^4]/(
2*d*(q + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), I
nt[((d + e*x^2)^(q + 1)/Sqrt[a + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2
*q + 3) + 2*c*d^2*(q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(C
*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x]] /; FreeQ[{a, c, d, e}, x]
&& PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[c*d^2 + a*e^2, 0] && ILt
Q[q, -1]
```

Rule 1721

```
Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4])
, x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e
```

```
) + a*(e/d), 2]*(x/Sqrt[a + c*x^4])/(2*d*e*Rt[c*(d/e) + a*(e/d), 2])), x]
+ Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*((a + c*x^4)/(a*(A + B*x^2)^2))]/(
4*d*e*A*q*Sqrt[a + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)],
2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e
^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1723

```
Int[((A_.) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]
), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q
))/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e
+ d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x],
x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2
- a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1729

```
Int[(P4x_)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] :=
With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff
[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Dist
[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2
)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2, 2]
&& NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rubi steps

$\frac{1}{2}x^2)^{1/2}/(cx^4+a)^{1/2}*\text{EllipticPi}(x*(I/a^{1/2}*c^{1/2})^{1/2}, I*a^{1/2}/c^{1/2}*e/d, (-I/a^{1/2}*c^{1/2})^{1/2}/(I/a^{1/2}*c^{1/2})^{1/2})*a+3/2/(a*e^2+c*d^2)/(I/a^{1/2}*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}*(1+I/a^{1/2}*c^{1/2}*x^2)^{1/2}/(cx^4+a)^{1/2}*\text{EllipticPi}(x*(I/a^{1/2}*c^{1/2})^{1/2}, I*a^{1/2}/c^{1/2}*e/d, (-I/a^{1/2}*c^{1/2})^{1/2}/(I/a^{1/2}*c^{1/2})^{1/2})*c)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(x^2*e + d)^3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*x^4 + a)*(B*x^2 + A)/(c*d^3*x^4 + a*d^3 + (c*x^10 + a*x^6)*e^3 + 3*(c*d*x^8 + a*d*x^4)*e^2 + 3*(c*d^2*x^6 + a*d^2*x^2)*e), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{\sqrt{a + cx^4} (d + ex^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+a)**(1/2),x)

[Out] Integral((A + B*x**2)/(sqrt(a + c*x**4)*(d + e*x**2)**3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + a)*(x^2*e + d)^3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{\sqrt{c x^4 + a} (e x^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^3), x)

[Out] int((A + B*x^2)/((a + c*x^4)^(1/2)*(d + e*x^2)^3), x)

$$3.8 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=912

$$\frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2)}{2ac^2\sqrt{a+cx^4}} + \frac{Be^3x\sqrt{a+cx^4}}{3c^2} + \frac{e^2(3Bd^2 - aAe^2)}{c^3}$$

[Out] $\frac{1}{2}x(Acd(-3ae^2+cd^2)-aBe(-ae^2+3cd^2)+c(-Aae^3+3Acd^2e-3Bde^2+Bcd^3)x^2)/a/c^2/(cx^4+a)^{(1/2)}+1/3B^3e^3x^3/(cx^4+a)^{(1/2)}/c^2+e^2(Ae+3Bd)x^2/(cx^4+a)^{(1/2)}/c^3/(a^{(1/2)}+x^2c^{(1/2)})-1/2(-Aae^3+3Acd^2e-3Bde^2+Bcd^3)x^2/(cx^4+a)^{(1/2)}/a/c^3/(a^{(1/2)}+x^2c^{(1/2)})-a^{(1/4)}e^2(Ae+3Bd)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))\text{EllipticE}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2,2^{(1/2)})(a^{(1/2)}+x^2c^{(1/2)})(cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)})^2)^{(1/2)}/c^{(7/4)}/(cx^4+a)^{(1/2)}+1/2(-Aae^3+3Acd^2e-3Bde^2+Bcd^3)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))\text{EllipticE}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2,2^{(1/2)})(a^{(1/2)}+x^2c^{(1/2)})(cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(7/4)}/(cx^4+a)^{(1/2)}-1/6a^{(3/4)}B^3e^3(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2,2^{(1/2)})(a^{(1/2)}+x^2c^{(1/2)})(cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)})^2)^{(1/2)}/c^{(9/4)}/(cx^4+a)^{(1/2)}+1/2a^{(1/4)}e^2(Ae+3Bd)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2,2^{(1/2)})(a^{(1/2)}+x^2c^{(1/2)})(cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)})^2)^{(1/2)}/c^{(7/4)}/(cx^4+a)^{(1/2)}+1/2e^2(3Acd^2e-Bae^2+3Bcd^2)(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2,2^{(1/2)})(a^{(1/2)}+x^2c^{(1/2)})(cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(9/4)}/(cx^4+a)^{(1/2)}+1/4(\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))^2)^{(1/2)}/\cos(2\arctan(c^{(1/4)}x/a^{(1/4)}))\text{EllipticF}(\sin(2\arctan(c^{(1/4)}x/a^{(1/4)})),1/2,2^{(1/2)})(A^2cd^3+a^2B^3e^3-3Acd^2e(Ae+Bd)-c^{(3/2)}d^2(3Ae+Bd)a^{(1/2)}+a^{(3/2)}e^2(Ae+3Bd)c^{(1/2)})(a^{(1/2)}+x^2c^{(1/2)})(cx^4+a)/(a^{(1/2)}+x^2c^{(1/2)})^2)^{(1/2)}/a^{(5/4)}/c^{(9/4)}/(cx^4+a)^{(1/2)}$

Rubi [A]

time = 0.45, antiderivative size = 912, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 7, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1735, 1193, 1212, 226, 1210, 311, 327}

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x]

[Out] (x*(A*c*d*(c*d^2 - 3*a*e^2) - a*B*e*(3*c*d^2 - a*e^2) + c*(B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*x^2))/(2*a*c^2*Sqrt[a + c*x^4]) + (B*e^3*x*Sqrt[a + c*x^4])/(3*c^2) + (e^2*(3*B*d + A*e)*x*Sqrt[a + c*x^4])/(c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - ((B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*Sqrt[a + c*x^4])/(2*a*c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - (a^(1/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(c^(7/4)*Sqrt[a + c*x^4]) + ((B*c*d^3 + 3*A*c*d^2*e - 3*a*B*d*e^2 - a*A*e^3)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(7/4)*Sqrt[a + c*x^4]) - (a^(3/4)*B*e^3*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(6*c^(9/4)*Sqrt[a + c*x^4]) + (a^(1/4)*e^2*(3*B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*c^(7/4)*Sqrt[a + c*x^4]) + (e*(3*B*c*d^2 + 3*A*c*d*e - a*B*e^2)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(9/4)*Sqrt[a + c*x^4]) + ((A*c^2*d^3 + a^2*B*e^3 - 3*a*c*d*e*(B*d + A*e) + a^(3/2)*Sqrt[c]*e^2*(3*B*d + A*e) - Sqrt[a]*c^(3/2)*d^2*(B*d + 3*A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*c^(9/4)*Sqrt[a + c*x^4])

Rule 226

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 311

Int[(x_)^2/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 2]}, Dist[1/q, Int[1/Sqrt[a + b*x^4], x], x] - Dist[1/q, Int[(1 - q*x^2)/Sqrt[a + b*x^4], x], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 1193

Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[(-x)*(d + e*x^2)*((a + c*x^4)^(p + 1)/(4*a*(p + 1))), x] + Dist[1/(4*a*(p + 1))

```
), Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /
; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ
[2*p]
```

Rule 1210

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

Rule 1212

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rule 1735

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)}{c^2(a + cx^4)^{3/2}} \right. \\
&= \frac{\int \frac{Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3)x^2}{(a + cx^4)^{3/2}} dx}{c^2} + \frac{(Be^3) \int \frac{x^4}{\sqrt{a + cx^4}}}{c} \\
&= \frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3))}{2ac^2\sqrt{a + cx^4}} \\
&= \frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3))}{2ac^2\sqrt{a + cx^4}} \\
&= \frac{x(Acd(cd^2 - 3ae^2) - aBe(3cd^2 - ae^2) + c(Bcd^3 + 3Acd^2e - 3aBde^2 - aAe^3))}{2ac^2\sqrt{a + cx^4}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.20, size = 222, normalized size = 0.24

$$\frac{3Acd^3 + ae^2(-3d + 2ex^2) + aBez(5ae^2 + c(-9d^2 + 18dex^2 + 2e^2x^4)) + (aBe(9cd^2 - 5ae^2) + 3Acd(cd^2 + 3ae^2))x\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; -\frac{cx^4}{a}\right) + 2c(Bcd^3 + 3Acd^2e - 9aBde^2 - 3aAe^3)x^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{3}{4}, \frac{7}{4}; -\frac{cx^4}{a}\right)}{6ac^2\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x]

[Out] (3*A*c*x*(c*d^3 + a*e^2*(-3*d + 2*e*x^2)) + a*B*e*x*(5*a*e^2 + c*(-9*d^2 + 18*d*e*x^2 + 2*e^2*x^4)) + (a*B*e*(9*c*d^2 - 5*a*e^2) + 3*A*c*d*(c*d^2 + 3*a*e^2))*x*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*c*(B*c*d^3 + 3*A*c*d^2*e - 9*a*B*d*e^2 - 3*a*A*e^3)*x^3*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c^2*sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.20, size = 588, normalized size = 0.64 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2), x, method=_RETURNVERBOSE)

[Out] B*e^3*(1/2/c^2*x*a/((x^4+a/c)*c)^(1/2)+1/3/c^2*x*(c*x^4+a)^(1/2)-5/6*a/c^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2))

$$\frac{1}{2}x^2)^{1/2}/(cx^4+a)^{1/2}*\text{EllipticF}(x*(I/a^{1/2}*c^{1/2})^{1/2},I)+(A$$

$$*e^3+3*B*d*e^2)*(-1/2/c*x^3/((x^4+a/c)*c)^{1/2}+3/2*I/c^{3/2}*a^{1/2}/(I/a^{1/2}$$

$$*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}*(1+I/a^{1/2}*c^{1/2}*x$$

$$^2)^{1/2}/(c*x^4+a)^{1/2}*(\text{EllipticF}(x*(I/a^{1/2}*c^{1/2})^{1/2},I)-\text{Ellipti}$$

$$cE(x*(I/a^{1/2}*c^{1/2})^{1/2},I)))+(3*A*d*e^2+3*B*d^2*e)*(-1/2/c*x/((x^4+a$$

$$/c)*c)^{1/2}+1/2/c/(I/a^{1/2}*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}$$

$$*(1+I/a^{1/2}*c^{1/2}*x^2)^{1/2}/(c*x^4+a)^{1/2}*\text{EllipticF}(x*(I/a^{1/2}*c^{1/2}$$

$$)^{1/2},I)+(3*A*d^2*e+B*d^3)*(1/2*x^3/a/((x^4+a/c)*c)^{1/2}-1/2*I/a^{1/2}$$

$$/I/a^{1/2}*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}*(1+I/a^{1/2}$$

$$*c^{1/2}*x^2)^{1/2}/(c*x^4+a)^{1/2}/c^{1/2}*(\text{EllipticF}(x*(I/a^{1/2}*c^{1/2}$$

$$)^{1/2},I)-\text{EllipticE}(x*(I/a^{1/2}*c^{1/2})^{1/2},I))+A*d^3*(1/2*x/a/((x^4+$$

$$a/c)*c)^{1/2}+1/2/a/(I/a^{1/2}*c^{1/2})^{1/2}*(1-I/a^{1/2}*c^{1/2}*x^2)^{1/2}$$

$$*(1+I/a^{1/2}*c^{1/2}*x^2)^{1/2}/(c*x^4+a)^{1/2}*\text{EllipticF}(x*(I/a^{1/2}*c$$

$$^{1/2})^{1/2},I))$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/(c*x^4 + a)^(3/2), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**3/(a + c*x**4)**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/(c*x^4 + a)^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(B x^2 + A) (e x^2 + d)^3}{(c x^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + c*x^4)^(3/2), x)

$$3.9 \quad \int \frac{(A+Bx^2)(d+ex^2)^2}{(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=694

$$\frac{x(ACd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a+cx^4}} + \frac{Be^2x\sqrt{a+cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{(Bcd^2 + 2Acde - aBe^2)x}{2ac^{3/2}(\sqrt{a} + \sqrt{c}x^2)}$$

```
[Out] 1/2*x*(A*c*d^2-2*a*B*d*e-a*A*e^2+(2*A*c*d*e-B*a*e^2+B*c*d^2)*x^2)/a/c/(c*x^
4+a)^(1/2)+B*e^2*x*(c*x^4+a)^(1/2)/c^(3/2)/(a^(1/2)+x^2*c^(1/2))-1/2*(2*A*c
*d*e-B*a*e^2+B*c*d^2)*x*(c*x^4+a)^(1/2)/a/c^(3/2)/(a^(1/2)+x^2*c^(1/2))-a^(
1/4)*B*e^2*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*
x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2
)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(7/4)/(c*x^4+a)^(
1/2)+1/2*(2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(
1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(
1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))
^2)^(1/2)/a^(3/4)/c^(7/4)/(c*x^4+a)^(1/2)+1/2*a^(1/4)*B*e^2*(cos(2*arctan(c^
(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2
*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(
a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(7/4)/(c*x^4+a)^(1/2)+1/2*e*(A*e+2*B*d)*(co
s(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*El
lipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))
*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(5/4)/(c*x^4+a)^(1/2)-
1/4*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/
4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c
^(1/2))*(B*c*d^2+2*A*c*d*e-a*B*e^2-(-A*a*e^2+A*c*d^2-2*B*a*d*e)*c^(1/2)/a^(
1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(7/4)/(c*x^4+a)^(
1/2)
```

Rubi [A]

time = 0.34, antiderivative size = 694, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 6, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {1735, 1193, 1212, 226, 1210, 311}

(1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2) - (1/2)*sqrt(a+cx^4)/sqrt(a+cx^4)^(3/2)

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x]

[Out] (x*(A*c*d^2 - 2*a*B*d*e - a*A*e^2 + (B*c*d^2 + 2*A*c*d*e - a*B*e^2)*x^2))/(2*a*c*Sqrt[a + c*x^4]) + (B*e^2*x*Sqrt[a + c*x^4])/(c^(3/2)*(Sqrt[a] + Sqrt[c]*x^2)) - ((B*c*d^2 + 2*A*c*d*e - a*B*e^2)*x*Sqrt[a + c*x^4])/(2*a*c^(3/2)

$$\begin{aligned} &)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2) - (a^{(1/4)}*B*e^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a \\ & + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)} \\ &], 1/2])/((c^{(7/4)}*\text{Sqrt}[a + c*x^4]) + ((B*c*d^2 + 2*A*c*d*e - a*B*e^2)*(\text{Sqrt} \\ & [a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2* \\ & \text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(2*a^{(3/4)}*c^{(7/4)}*\text{Sqrt}[a + c*x^4]) + (a \\ & ^{(1/4)}*B*e^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^ \\ & 2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(2*c^{(7/4)}*\text{Sqrt}[a + c* \\ & x^4]) + (e*(2*B*d + A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] \\ & + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(2*a^{(1/4)} \\ & *c^{(5/4)}*\text{Sqrt}[a + c*x^4]) - ((B*c*d^2 + 2*A*c*d*e - a*B*e^2 - (\text{Sqrt}[c]*(A*c \\ & *d^2 - 2*a*B*d*e - a*A*e^2))/\text{Sqrt}[a])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x \\ & ^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2 \\ &])/(4*a^{(3/4)}*c^{(7/4)}*\text{Sqrt}[a + c*x^4]) \end{aligned}$$

Rule 226

$$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*\text{Sqrt}[a + b*x^4]))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[b/a]$$

Rule 311

$$\text{Int}[(x_)^2/\text{Sqrt}[(a_) + (b_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[b/a, 2]\}, \text{Dist}[1/q, \text{Int}[1/\text{Sqrt}[a + b*x^4], x], x] - \text{Dist}[1/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^4], x], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[b/a]$$

Rule 1193

$$\text{Int}[(d_) + (e_)*(x_)^2]*((a_) + (c_)*(x_)^4)^{(p_)}, x_Symbol] := \text{Simp}[(-x)*(d + e*x^2)*((a + c*x^4)^{(p + 1)}/(4*a*(p + 1))), x] + \text{Dist}[1/(4*a*(p + 1)), \text{Int}[\text{Simp}[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{NeQ}[c*d^2 + a*e^2, 0] \&\& \text{LtQ}[p, -1] \&\& \text{IntegerQ}[2*p]$$

Rule 1210

$$\text{Int}[(d_) + (e_)*(x_)^2]/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*\text{Sqrt}[a + c*x^4]))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{PosQ}[c/a]$$

Rule 1212

$$\text{Int}[(d_) + (e_)*(x_)^2]/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] := \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + c*x^4], x], x] /; \text{NeQ}[e + d*q, 0] /; \text{FreeQ}[\{a, c,$$

$d, e\}, x] \&\& \text{PosQ}[c/a]$

Rule 1735

`Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]`
`:> Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p`
`+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a`
`*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]`

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2}{c(a + cx^4)^{3/2}} + \frac{e(2Bd + Ae)}{c\sqrt{a + cx^4}} \right) dx \\ &= \frac{\int \frac{Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2}{(a + cx^4)^{3/2}} dx}{c} + \frac{(Be^2) \int \frac{x^2}{\sqrt{a + cx^4}} dx}{c} + \frac{e(2Bd + Ae) \int \frac{1}{\sqrt{a + cx^4}} dx}{c} \\ &= \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a + cx^4}} + \frac{e(2Bd + Ae) \int \frac{1}{\sqrt{a + cx^4}} dx}{c} \\ &= \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be^2 x \sqrt{a + cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{cx^4})} \\ &= \frac{x(Acd^2 - 2aBde - aAe^2 + (Bcd^2 + 2Acde - aBe^2)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be^2 x \sqrt{a + cx^4}}{c^{3/2}(\sqrt{a} + \sqrt{cx^4})} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.13, size = 166, normalized size = 0.24

$$\frac{3A(cd^2 - ae^2)x + 6aBex(-d + ex^2) + 3(Acd^2 + 2aBde + aAe^2)x\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) + 2(Bcd^2 + 2Acde - 3aBe^2)x^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{7}{4}; -\frac{cx^4}{a}\right)}{6ac\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] `Integrate[((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x]`

[Out] `(3*A*(c*d^2 - a*e^2)*x + 6*a*B*e*x*(-d + e*x^2) + 3*(A*c*d^2 + 2*a*B*d*e + a*A*e^2)*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -(c*x^4)/a`

)] + 2*(B*c*d^2 + 2*A*c*d*e - 3*a*B*e^2)*x^3*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c*sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.16, size = 458, normalized size = 0.66

method	result
elliptic	$-\frac{2c\left(-\frac{(2cdeA - Ba e^2 + Bc d^2)x^3}{4c^2a} + \frac{(Aa e^2 - Ac d^2 + 2adeB)x}{4c^2a}\right)}{\sqrt{\left(x^4 + \frac{a}{c}\right)c}} + \frac{\left(\frac{e(Ae + 2Bd)}{c} - \frac{Aa e^2 - Ac d^2 + 2adeB}{2ac}\right) \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}}$
default	$B e^2 \left(-\frac{x^3}{2c\sqrt{\left(x^4 + \frac{a}{c}\right)c}} + \frac{3i\sqrt{a} \sqrt{1 - \frac{i\sqrt{c} x^2}{\sqrt{a}}} \sqrt{1 + \frac{i\sqrt{c} x^2}{\sqrt{a}}}}{2c^{\frac{3}{2}} \sqrt{\frac{i\sqrt{c}}{\sqrt{a}}} \sqrt{cx^4 + a}} \left(\text{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - \text{EllipticE}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)

[Out] B*e^2*(-1/2/c*x^3/((x^4+a/c)*c)^(1/2)+3/2*I/c^(3/2)*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)))+(A*e^2+2*B*d*e)*(-1/2/c*x/((x^4+a/c)*c)^(1/2)+1/2/c/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I))+(2*A*d*e+B*d^2)*(1/2*x^3/a/((x^4+a/c)*c)^(1/2)-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)))+A*d^2*(1/2*x/a/((x^4+a/c)*c)^(1/2)+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^2/(c*x^4 + a)^(3/2), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Symbolic function `elliptic_ec` takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+a)**(3/2),x)`

[Out] `Integral((A + B*x**2)*(d + e*x**2)**2/(a + c*x**4)**(3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="giac")`

[Out] `integrate((B*x^2 + A)*(x^2*e + d)^2/(c*x^4 + a)^(3/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + a)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2),x)`

[Out] `int(((A + B*x^2)*(d + e*x^2)^2)/(a + c*x^4)^(3/2), x)`

$$3.10 \quad \int \frac{(A+Bx^2)(d+ex^2)}{(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=395

$$\frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a+cx^4}} - \frac{(Bd + Ae)x\sqrt{a+cx^4}}{2a\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} + \frac{(Bd + Ae)(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}} E$$

[Out] 1/2*x*(A*c*d-a*B*e+c*(A*e+B*d)*x^2)/a/c/(c*x^4+a)^(1/2)-1/2*(A*e+B*d)*x*(c*x^4+a)^(1/2)/a/c^(1/2)/(a^(1/2)+x^2*c^(1/2))+1/2*(A*e+B*d)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(3/4)/(c*x^4+a)^(1/2)+1/2*B*e*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/c^(5/4)/(c*x^4+a)^(1/2)+1/4*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(5/4)/c^(5/4)/(c*x^4+a)^(1/2)

Rubi [A]

time = 0.21, antiderivative size = 395, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {1735, 1193, 1212, 226, 1210}

$$\frac{(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right) (-\sqrt{a}\sqrt{c}(Ae+Bd) - aBe + Ad) + (\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} (Ae+Bd) E\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right) - \frac{x\sqrt{a+cx^4}(Ae+Bd)}{2a\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} + \frac{x(-aBe+cx^2(Ae+Bd) + Ad)}{2ac\sqrt{a+cx^4}} + \frac{Be(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right) \middle| \frac{1}{2}\right)}{2\sqrt{a}c^{3/4}\sqrt{a+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x]

[Out] (x*(A*c*d - a*B*e + c*(B*d + A*e)*x^2))/(2*a*c*Sqrt[a + c*x^4]) - ((B*d + A*e)*x*Sqrt[a + c*x^4])/(2*a*Sqrt[c]*(Sqrt[a] + Sqrt[c]*x^2)) + ((B*d + A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(3/4)*c^(3/4)*Sqrt[a + c*x^4]) + (B*e*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(2*a^(1/4)*c^(5/4)*Sqrt[a + c*x^4]) + ((A*c*d - a*B*e - Sqrt[a]*Sqrt[c]*(B*d + A*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2])/(4*a^(5/4)*c^(5/4)*Sqrt[a + c*x^4])

Rule 226

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^4]))*

EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 1193

Int[((d_) + (e_)*(x_)^2)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Simp[(-x)*(d + e*x^2)*((a + c*x^4)^(p + 1)/(4*a*(p + 1))), x] + Dist[1/(4*a*(p + 1)), Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2*p]

Rule 1210

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] :> With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2)^2))/(q*Sqrt[a + c*x^4])*EllipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 1212

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] :> With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c, d, e}, x] && PosQ[c/a]

Rule 1735

Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)}{(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd - aBe + c(Bd + Ae)x^2}{c(a + cx^4)^{3/2}} + \frac{Be}{c\sqrt{a + cx^4}} \right) dx \\
&= \frac{\int \frac{Acd - aBe + c(Bd + Ae)x^2}{(a + cx^4)^{3/2}} dx}{c} + \frac{(Be) \int \frac{1}{\sqrt{a + cx^4}} dx}{c} \\
&= \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}\right)}{2^4\sqrt{a}c^{5/4}\sqrt{a + cx^4}} \\
&= \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a + cx^4}} + \frac{Be(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}\right)}{2^4\sqrt{a}c^{5/4}\sqrt{a + cx^4}} \\
&= \frac{x(Acd - aBe + c(Bd + Ae)x^2)}{2ac\sqrt{a + cx^4}} - \frac{(Bd + Ae)x\sqrt{a + cx^4}}{2a\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} + \frac{(Bd + Ae)(\sqrt{a} + \sqrt{c}x^2)}{2a\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.09, size = 126, normalized size = 0.32

$$\frac{3(Acd - aBe)x + 3(Acd + aBe)x\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}; \frac{5}{4}; -\frac{cx^4}{a}\right) + 2c(Bd + Ae)x^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{7}{4}; -\frac{cx^4}{a}\right)}{6ac\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x]

[Out] (3*(A*c*d - a*B*e)*x + 3*(A*c*d + a*B*e)*x*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*c*(B*d + A*e)*x^3*sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)]/(6*a*c*sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.14, size = 320, normalized size = 0.81

method	result
elliptic	$ -\frac{2c\left(-\frac{(Ae+Bd)x^3}{4ca} - \frac{(Acd-aBe)x}{4ac^2}\right)}{\sqrt{\left(x^4 + \frac{a}{c}\right)c}} + \frac{\left(\frac{Be}{c} + \frac{Acd-aBe}{2ac}\right)\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right) - i(A $

default	$Be \left(-\frac{x}{2c\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2c\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \right) + (Ae + Bd) \left(-\frac{x}{2c\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2c\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \right)$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $B*e*(-1/2/c*x/((x^4+a/c)*c)^(1/2)+1/2/c/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*\operatorname{EllipticF}(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+(A*e+B*d)*(1/2*x^3/a/((x^4+a/c)*c)^(1/2)-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(\operatorname{EllipticF}(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-\operatorname{EllipticE}(x*(I/a^(1/2)*c^(1/2))^(1/2),I))+A*d*(1/2*x/a/((x^4+a/c)*c)^(1/2)+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*\operatorname{EllipticF}(x*(I/a^(1/2)*c^(1/2))^(1/2),I))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)*(x^2*e + d)/(c*x^4 + a)^(3/2), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Symbolic function `elliptic_ec` takes exactly 1 arguments (2 given)

Sympy [C] Result contains complex when optimal does not.

time = 4.91, size = 167, normalized size = 0.42

$$\frac{Adx\Gamma(\frac{1}{4}) {}_2F_1\left(\frac{1}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma(\frac{5}{4})} + \frac{Aex^3\Gamma(\frac{3}{4}) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma(\frac{7}{4})} + \frac{Bdx^3\Gamma(\frac{3}{4}) {}_2F_1\left(\frac{3}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma(\frac{7}{4})} + \frac{Bex^5\Gamma(\frac{5}{4}) {}_2F_1\left(\frac{5}{4}, \frac{3}{2} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma(\frac{9}{4})}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+a)**(3/2),x)
```

```
[Out] A*d*x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**
(3/2)*gamma(5/4)) + A*e*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*ex
p_polar(I*pi)/a)/(4*a**(3/2)*gamma(7/4)) + B*d*x**3*gamma(3/4)*hyper((3/4,
3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(7/4)) + B*e*x**5*
gamma(5/4)*hyper((5/4, 3/2), (9/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*
gamma(9/4))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)*(x^2*e + d)/(c*x^4 + a)^(3/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2),x)
```

```
[Out] int(((A + B*x^2)*(d + e*x^2))/(a + c*x^4)^(3/2), x)
```


3.11 $\int \frac{A+Bx^2}{(a+cx^4)^{3/2}} dx$

Optimal. Leaf size=262

$$\frac{x(A+Bx^2)}{2a\sqrt{a+cx^4}} - \frac{Bx\sqrt{a+cx^4}}{2a\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)} + \frac{B(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2\tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right)\middle|\frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}}$$

[Out] $1/2*x*(B*x^2+A)/a/(c*x^4+a)^{(1/2)}-1/2*B*x*(c*x^4+a)^{(1/2)}/a/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})+1/2*B*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(3/4)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}-1/4*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/a^{(5/4)}/c^{(3/4)}/(c*x^4+a)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 262, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {1193, 1212, 226, 1210}

$$-\frac{(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}(\sqrt{a}B-A\sqrt{c})F\left(2\text{ArcTan}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right)\middle|\frac{1}{2}\right)}{4a^{5/4}c^{3/4}\sqrt{a+cx^4}} + \frac{B(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}E\left(2\text{ArcTan}\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}\right)\middle|\frac{1}{2}\right)}{2a^{3/4}c^{3/4}\sqrt{a+cx^4}} + \frac{x(A+Bx^2)}{2a\sqrt{a+cx^4}} - \frac{Bx\sqrt{a+cx^4}}{2a\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/(a + c*x^4)^(3/2), x]

[Out] $(x*(A+B*x^2))/(2*a*\text{Sqrt}[a+c*x^4]) - (B*x*\text{Sqrt}[a+c*x^4])/(2*a*\text{Sqrt}[c]*(\text{Sqrt}[a]+\text{Sqrt}[c]*x^2)) + (B*(\text{Sqrt}[a]+\text{Sqrt}[c]*x^2)*\text{Sqrt}[(a+c*x^4)/(\text{Sqrt}[a]+\text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(2*a^{(3/4)}*c^{(3/4)}*\text{Sqrt}[a+c*x^4]) - ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])*(\text{Sqrt}[a]+\text{Sqrt}[c]*x^2)*\text{Sqrt}[(a+c*x^4)/(\text{Sqrt}[a]+\text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/(4*a^{(5/4)}*c^{(3/4)}*\text{Sqrt}[a+c*x^4])$

Rule 226

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] :> With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

Rule 1193

Int[((d_) + (e_)*(x_)^2)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Simp[(-x)*(d + e*x^2)*((a + c*x^4)^(p + 1)/(4*a*(p + 1))), x] + Dist[1/(4*a*(p + 1))

```
), Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /
; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ
[2*p]
```

Rule 1210

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
  (1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*E
  llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
  }, x] && PosQ[c/a]
```

Rule 1212

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
  Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
  nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
  d, e}, x] && PosQ[c/a]
```

Rubi steps

$$\begin{aligned} \int \frac{A + Bx^2}{(a + cx^4)^{3/2}} dx &= \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} - \frac{\int \frac{-A+Bx^2}{\sqrt{a + cx^4}} dx}{2a} \\ &= \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} + \frac{\left(A - \frac{\sqrt{a} B}{\sqrt{c}}\right) \int \frac{1}{\sqrt{a + cx^4}} dx}{2a} + \frac{B \int \frac{1 - \frac{\sqrt{c} x^2}{\sqrt{a}}}{\sqrt{a + cx^4}} dx}{2\sqrt{a} \sqrt{c}} \\ &= \frac{x(A + Bx^2)}{2a\sqrt{a + cx^4}} - \frac{Bx\sqrt{a + cx^4}}{2a\sqrt{c} (\sqrt{a} + \sqrt{c} x^2)} + \frac{B(\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} E\left(2 \operatorname{arctan}\left(\frac{\sqrt{c} x}{\sqrt{a}}\right)\right)}{2a^{3/4} c^{3/4} \sqrt{a + cx^4}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.04, size = 99, normalized size = 0.38

$$\frac{3Ax + 3Ax\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{1}{4}, \frac{1}{2}, \frac{5}{4}, -\frac{cx^4}{a}\right) + 2Bx^3\sqrt{1 + \frac{cx^4}{a}} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}, \frac{7}{4}, -\frac{cx^4}{a}\right)}{6a\sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/(a + c*x^4)^(3/2),x]

[Out] (3*A*x + 3*A*x*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[1/4, 1/2, 5/4, -((c*x^4)/a)] + 2*B*x^3*Sqrt[1 + (c*x^4)/a]*Hypergeometric2F1[3/4, 3/2, 7/4, -((c*x^4)/a)])/(6*a*Sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.

time = 0.14, size = 217, normalized size = 0.83

method	result
elliptic	$-\frac{2c\left(-\frac{Bx^3}{4ac}-\frac{Ax}{4ac}\right)}{\sqrt{\left(x^4+\frac{a}{c}\right)c}} + \frac{A\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}} - \frac{iB\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}}$
default	$B\left(\frac{x^3}{2a\sqrt{\left(x^4+\frac{a}{c}\right)c}} - \frac{i\sqrt{1-\frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1+\frac{i\sqrt{c}x^2}{\sqrt{a}}}\left(\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)-\operatorname{EllipticE}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}},i\right)\right)}{2\sqrt{a}\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4+a}\sqrt{c}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)

[Out] B*(1/2*x^3/a/((x^4+a/c)*c)^(1/2)-1/2*I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)/c^(1/2)*(EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)-EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I))+A*(1/2*x/a/((x^4+a/c)*c)^(1/2)+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(c*x^4 + a)^(3/2), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [C] Result contains complex when optimal does not.

time = 2.78, size = 78, normalized size = 0.30

$$\frac{Ax\Gamma\left(\frac{1}{4}\right) {}_2F_1\left(\begin{matrix} \frac{1}{4}, \frac{3}{2} \\ \frac{5}{4} \end{matrix} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{5}{4}\right)} + \frac{Bx^3\Gamma\left(\frac{3}{4}\right) {}_2F_1\left(\begin{matrix} \frac{3}{4}, \frac{3}{2} \\ \frac{7}{4} \end{matrix} \middle| \frac{cx^4 e^{i\pi}}{a}\right)}{4a^{\frac{3}{2}}\Gamma\left(\frac{7}{4}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(c*x**4+a)**(3/2),x)

[Out] A*x*gamma(1/4)*hyper((1/4, 3/2), (5/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(5/4)) + B*x**3*gamma(3/4)*hyper((3/4, 3/2), (7/4,), c*x**4*exp_polar(I*pi)/a)/(4*a**(3/2)*gamma(7/4))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(c*x^4 + a)^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/(a + c*x^4)^(3/2),x)

[Out] int((A + B*x^2)/(a + c*x^4)^(3/2), x)

$$3.12 \quad \int \frac{A+Bx^2}{(d+ex^2)(a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=732

$$\frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a+cx^4}} - \frac{\sqrt{c}(Bd - Ae)x\sqrt{a+cx^4}}{2a(cd^2 + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e^{3/2}(Bd - Ae)\tan^{-1}\left(\frac{\sqrt{cd^2 + ae^2}x}{\sqrt{d}\sqrt{e}\sqrt{a+cx^4}}\right)}{2\sqrt{d}(cd^2 + ae^2)^{3/2}}$$

[Out] $-1/2*e^{(3/2)}*(-A*e+B*d)*\arctan(x*(a*e^2+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+a)^{(1/2)})/(a*e^2+c*d^2)^{(3/2)}/d^{(1/2)}+1/2*x*(A*c*d+a*B*e+c*(-A*e+B*d)*x^2)/a/(a*e^2+c*d^2)/(c*x^4+a)^{(1/2)}-1/2*(-A*e+B*d)*x*c^{(1/2)}*(c*x^4+a)^{(1/2)}/a/(a*e^2+c*d^2)/(a^{(1/2)}+x^2*c^{(1/2)})+1/2*c^{(1/4)}*(-A*e+B*d)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(a*e^2+c*d^2)/(c*x^4+a)^{(1/2)}-1/2*c^{(1/4)}*e*(-A*e+B*d)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/(a*e^2+c*d^2)/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+a)^{(1/2)}+1/4*a^{(3/4)}*e*(-A*e+B*d)*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(1/4)}/d/(-a^2*e^4+c^2*d^4)/(c*x^4+a)^{(1/2)}+1/4*(cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(A*c*d+a*B*e-(-A*e+B*d)*a^{(1/2)}*c^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(5/4)}/c^{(1/4)}/(a*e^2+c*d^2)/(c*x^4+a)^{(1/2)}$

Rubi [A]

time = 0.53, antiderivative size = 732, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1735, 1193, 1212, 226, 1210, 1231, 1721}

$$\frac{(\sqrt{c} + \sqrt{c}x^2) \sqrt{\frac{cd^2 + ae^2}{(c^2d^2 + ae^2)^2}} \left(\frac{2ax \arctan\left(\frac{cx}{d}\right)}{\sqrt{c}} \right) - \sqrt{c} \sqrt{cd^2 + ae^2} (Bd - Ae) + aBe + Ad}{2a \sqrt{c} \sqrt{cd^2 + ae^2} (ae^2 + cd^2)} - \frac{e^{3/2} (\sqrt{c} + \sqrt{c}x^2) \sqrt{\frac{cd^2 + ae^2}{(c^2d^2 + ae^2)^2}} (Bd - Ae) \left(\frac{2ax \arctan\left(\frac{cx}{d}\right)}{\sqrt{c}} \right)}{2a \sqrt{c} \sqrt{cd^2 + ae^2} (ae^2 + cd^2)} - \frac{e^{3/2} (\sqrt{c} + \sqrt{c}x^2) \sqrt{\frac{cd^2 + ae^2}{(c^2d^2 + ae^2)^2}} (Bd - Ae) \left(\frac{2ax \arctan\left(\frac{cx}{d}\right)}{\sqrt{c}} \right)}{2a \sqrt{c} \sqrt{cd^2 + ae^2} (ae^2 + cd^2)} - \frac{e^{3/2} (\sqrt{c} + \sqrt{c}x^2) \sqrt{\frac{cd^2 + ae^2}{(c^2d^2 + ae^2)^2}} (Bd - Ae) \left(\frac{2ax \arctan\left(\frac{cx}{d}\right)}{\sqrt{c}} \right)}{2a \sqrt{c} \sqrt{cd^2 + ae^2} (ae^2 + cd^2)} - \frac{e^{3/2} (\sqrt{c} + \sqrt{c}x^2) \sqrt{\frac{cd^2 + ae^2}{(c^2d^2 + ae^2)^2}} (Bd - Ae) \left(\frac{2ax \arctan\left(\frac{cx}{d}\right)}{\sqrt{c}} \right)}{2a \sqrt{c} \sqrt{cd^2 + ae^2} (ae^2 + cd^2)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*(a + c*x^4)^(3/2)),x]

[Out] $(x*(A*c*d + a*B*e + c*(B*d - A*e)*x^2))/(2*a*(c*d^2 + a*e^2)*\text{Sqrt}[a + c*x^4]) - (\text{Sqrt}[c]*(B*d - A*e)*x*\text{Sqrt}[a + c*x^4])/(2*a*(c*d^2 + a*e^2)*(\text{Sqrt}[a + \text{Sqrt}[c]*x^2]) - (e^{(3/2)}*(B*d - A*e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])])/(2*\text{Sqrt}[d]*(c*d^2 + a*e^2)^{(3/2)}) + (c^{(1/4)}$

```

*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^
2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(2*a^(3/4)*(c*d^2 + a*
e^2)*Sqrt[a + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt
[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1
/4)], 1/2]]/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 + a*e^2)*Sqrt[a + c*x
^4]) + ((A*c*d + a*B*e - Sqrt[a]*Sqrt[c]*(B*d - A*e))*(Sqrt[a] + Sqrt[c]*x^
2)*Sqrt[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*
x)/a^(1/4)], 1/2]]/(4*a^(5/4)*c^(1/4)*(c*d^2 + a*e^2)*Sqrt[a + c*x^4]) + (a
^(3/4)*e*((Sqrt[c]*d)/Sqrt[a] + e)^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqr
t[(a + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[c]*d - Sqrt
[a]*e)^2/(Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], 1/2]]/(4*c^(
1/4)*d*(c^2*d^4 - a^2*e^4)*Sqrt[a + c*x^4])

```

Rule 226

```

Int[1/Sqrt[(a_) + (b_)*(x_)^4], x_Symbol] := With[{q = Rt[b/a, 4]}, Simp[(
1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2]]/(2*q*Sqrt[a + b*x^4]))*
EllipticF[2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, b}, x] && PosQ[b/a]

```

Rule 1193

```

Int[((d_) + (e_)*(x_)^2)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[(-x
)*(d + e*x^2)*((a + c*x^4)^(p + 1)/(4*a*(p + 1))), x] + Dist[1/(4*a*(p + 1
)), Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /
; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ
[2*p]

```

Rule 1210

```

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2]]/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]

```

Rule 1212

```

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (c_)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]

```

Rule 1231

```

Int[1/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] := With[
{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4]
, x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^

```

2)*Sqrt[a + c*x^4]), x], x]] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]

Rule 1721

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :> With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2]))], x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*((a + c*x^4)/(a*(A + B*x^2)^2))]/(4*d*e*A*q*Sqrt[a + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rule 1735

Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]

Rubi steps

$$\begin{aligned}
 \int \frac{A + Bx^2}{(d + ex^2)(a + cx^4)^{3/2}} dx &= \int \left(\frac{Acd + aBe + c(Bd - Ae)x^2}{(cd^2 + ae^2)(a + cx^4)^{3/2}} + \frac{e(-Bd + Ae)}{(cd^2 + ae^2)(d + ex^2)\sqrt{a + cx^4}} \right) dx \\
 &= \frac{\int \frac{Acd + aBe + c(Bd - Ae)x^2}{(a + cx^4)^{3/2}} dx}{cd^2 + ae^2} - \frac{(e(Bd - Ae)) \int \frac{1}{(d + ex^2)\sqrt{a + cx^4}} dx}{cd^2 + ae^2} \\
 &= \frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a + cx^4}} - \frac{\int \frac{-Acd - aBe + c(Bd - Ae)x^2}{\sqrt{a + cx^4}} dx}{2a(cd^2 + ae^2)} - \frac{(\sqrt{c} e(Bd - Ae))}{(\sqrt{c} d - Ae)} \\
 &= \frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a + cx^4}} - \frac{e^{3/2}(Bd - Ae) \tan^{-1} \left(\frac{\sqrt{cd^2 + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + cx^4}} \right)}{2\sqrt{d} (cd^2 + ae^2)^{3/2}} \\
 &= \frac{x(Acd + aBe + c(Bd - Ae)x^2)}{2a(cd^2 + ae^2)\sqrt{a + cx^4}} - \frac{\sqrt{c} (Bd - Ae)x\sqrt{a + cx^4}}{2a(cd^2 + ae^2)(\sqrt{a} + \sqrt{c} x^2)} - \frac{e^{3/2}(Bd - Ae)}{(\sqrt{c} d - Ae)}
 \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.53, size = 432, normalized size = 0.59

$$\frac{A\sqrt{\frac{c}{a}}\operatorname{arctan}\left(\frac{\sqrt{\frac{c}{a}}x+B\sqrt{\frac{c}{a}}}{\sqrt{\frac{c}{a}}}\right)+B\sqrt{\frac{c}{a}}\operatorname{arctan}\left(\frac{\sqrt{\frac{c}{a}}x+B\sqrt{\frac{c}{a}}}{\sqrt{\frac{c}{a}}}\right)-A\sqrt{\frac{c}{a}}\operatorname{arctan}\left(\frac{\sqrt{\frac{c}{a}}x+B\sqrt{\frac{c}{a}}}{\sqrt{\frac{c}{a}}}\right)-\sqrt{a}\sqrt{c}d(B-A)\sqrt{1+\frac{cx^2}{a}}E\left(\operatorname{arcsinh}\left(\sqrt{\frac{c}{a}}x\right)\right)+(\sqrt{a}B-A\sqrt{c})d(\sqrt{c}d-i\sqrt{a}c)\sqrt{1+\frac{cx^2}{a}}F\left(\operatorname{arcsinh}\left(\sqrt{\frac{c}{a}}x\right)\right)+2aBd\sqrt{1+\frac{cx^2}{a}}\Pi\left(\frac{-\sqrt{a}}{\sqrt{c}};\operatorname{arcsinh}\left(\sqrt{\frac{c}{a}}x\right)\right)-2aAe^2\sqrt{1+\frac{cx^2}{a}}\Pi\left(\frac{-\sqrt{a}}{\sqrt{c}};\operatorname{arcsinh}\left(\sqrt{\frac{c}{a}}x\right)\right)}{2a\sqrt{\frac{c}{a}}d(ad^2+ae^2)\sqrt{a+cx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(A + B*x^2)/((d + e*x^2)*(a + c*x^4)^(3/2)),x]
```

```
[Out] (A*Sqrt[(I*Sqrt[c])/Sqrt[a]]*c*d^2*x + a*B*Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*e*x
+ B*Sqrt[(I*Sqrt[c])/Sqrt[a]]*c*d^2*x^3 - A*Sqrt[(I*Sqrt[c])/Sqrt[a]]*c*d*e
*x^3 - Sqrt[a]*Sqrt[c]*d*(B*d - A*e)*Sqrt[1 + (c*x^4)/a]*EllipticE[I*ArcSin
h[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + (Sqrt[a]*B - I*A*Sqrt[c])*d*(Sqrt[c]*
d - I*Sqrt[a]*e)*Sqrt[1 + (c*x^4)/a]*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/S
qrt[a]]*x], -1] + (2*I)*a*B*d*e*Sqrt[1 + (c*x^4)/a]*EllipticPi[(-I)*Sqrt[a
]*e]/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] - (2*I)*a*A*e
^2*Sqrt[1 + (c*x^4)/a]*EllipticPi[(-I)*Sqrt[a]*e]/(Sqrt[c]*d), I*ArcSinh[S
qrt[(I*Sqrt[c])/Sqrt[a]]*x], -1)]/(2*a*Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*(c*d^2 +
a*e^2)*Sqrt[a + c*x^4])
```

Maple [C] Result contains complex when optimal does not.
time = 0.17, size = 564, normalized size = 0.77

method	result
default	$B \left(\frac{x}{2a\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)}{2a\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} \right) \frac{(Ae - Bd)}{e} + \frac{2c \left(\frac{ex^3}{4a(ae^2 + cd^2)} - \frac{1}{\sqrt{(x^4 + \frac{a}{c})c}} \right)}{\sqrt{(x^4 + \frac{a}{c})c}}$
elliptic	$-\frac{2c \left(\frac{(Ae - Bd)x^3}{4a(ae^2 + cd^2)} - \frac{(Acd + aBe)x}{4a(ae^2 + cd^2)c} \right)}{\sqrt{(x^4 + \frac{a}{c})c}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}\sqrt{1 + \frac{i\sqrt{c}x^2}{\sqrt{a}}}\operatorname{EllipticF}\left(x\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}, i\right)Acd}{2a(ae^2 + cd^2)\sqrt{\frac{i\sqrt{c}}{\sqrt{a}}}\sqrt{cx^4 + a}} + \frac{\sqrt{1 - \frac{i\sqrt{c}x^2}{\sqrt{a}}}}{\sqrt{(x^4 + \frac{a}{c})c}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] B/e*(1/2*x/a/((x^4+a/c)*c)^(1/2)+1/2/a/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/
2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*Ellip
ticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+(A*e-B*d)/e*(-2*c*(1/4/a*e/(a*e^2+c*d^
```


$$2) * x^{3-1/4} * d/a / (a * e^2 + c * d^2) * x / ((x^4 + a/c) * c)^{1/2} + 1/2 * c * d/a / (a * e^2 + c * d^2) / (I/a^{1/2} * c^{1/2})^{1/2} * (1 - I/a^{1/2} * c^{1/2} * x^2)^{1/2} * (1 + I/a^{1/2} * c^{1/2} * x^2)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticF}(x * (I/a^{1/2} * c^{1/2})^{1/2}, I) + 1/2 * I/a^{1/2} * c^{1/2} * e / (a * e^2 + c * d^2) / (I/a^{1/2} * c^{1/2})^{1/2} * (1 - I/a^{1/2} * c^{1/2} * x^2)^{1/2} * (1 + I/a^{1/2} * c^{1/2} * x^2)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticE}(x * (I/a^{1/2} * c^{1/2})^{1/2}, I) - 1/2 * I/a^{1/2} * c^{1/2} * e / (a * e^2 + c * d^2) / (I/a^{1/2} * c^{1/2})^{1/2} * (1 - I/a^{1/2} * c^{1/2} * x^2)^{1/2} * (1 + I/a^{1/2} * c^{1/2} * x^2)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticE}(x * (I/a^{1/2} * c^{1/2})^{1/2}, I) + 1/(a * e^2 + c * d^2) * e^2/d / (I/a^{1/2} * c^{1/2})^{1/2} * (1 - I/a^{1/2} * c^{1/2} * x^2)^{1/2} * (1 + I/a^{1/2} * c^{1/2} * x^2)^{1/2} / (c * x^4 + a)^{1/2} * \text{EllipticPi}(x * (I/a^{1/2} * c^{1/2})^{1/2}, I * a^{1/2}/c^{1/2} * e/d, (-I/a^{1/2} * c^{1/2})^{1/2} / (I/a^{1/2} * c^{1/2})^{1/2})^{1/2})$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(x^2*e + d)), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(a + cx^4)^{\frac{3}{2}} (d + ex^2)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+a)**(3/2),x)

[Out] Integral((A + B*x**2)/((a + c*x**4)**(3/2)*(d + e*x**2)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(x^2*e + d)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{(c x^4 + a)^{3/2} (e x^2 + d)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)),x)

[Out] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)), x)

$$\arctan(c^{(1/4)}*x/a^{(1/4)})*EllipticF(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*2^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(B*c*d^2-2*A*c*d*e-a*B*e^2-(-A*a*e^2+A*c*d^2+2*B*a*d*e)*c^{(1/2)}/a^{(1/2)})*((c*x^4+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(a*e^2+c*d^2)^2/(c*x^4+a)^{(1/2)}$$

Rubi [A]

time = 1.36, antiderivative size = 1494, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 10, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {1735, 1193, 1212, 226, 1210, 1238, 1729, 1723, 1721, 1231}

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^2*(a + c*x^4)^(3/2)),x]

[Out] $(c*x*(A*c*d^2 + 2*a*B*d*e - a*A*e^2 + (B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x^2))/((2*a*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) + (\text{Sqrt}[c]*e^2*(B*d - A*e)*x*\text{Sqrt}[a + c*x^4]))/(2*d*(c*d^2 + a*e^2)^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (\text{Sqrt}[c]*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*x*\text{Sqrt}[a + c*x^4])/((2*a*(c*d^2 + a*e^2)^2*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (e^3*(B*d - A*e)*x*\text{Sqrt}[a + c*x^4]))/(2*d*(c*d^2 + a*e^2)^2*(d + e*x^2)) - (e^{(3/2)}*(B*d - A*e)*(3*c*d^2 + a*e^2)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])])/(4*d^{(3/2)}*(c*d^2 + a*e^2)^{(5/2)}) - (e^{(3/2)}*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])])/(2*\text{Sqrt}[d]*(c*d^2 + a*e^2)^{(5/2)}) - (a^{(1/4)}*c^{(1/4)}*e^2*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((2*d*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) + (c^{(1/4)}*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((2*a^{(3/4)}*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) - (c^{(1/4)}*e*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((2*a^{(1/4)}*d*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)*\text{Sqrt}[a + c*x^4]) - (c^{(1/4)}*e*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((2*a^{(1/4)}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) - (c^{(1/4)}*(B*c*d^2 - 2*A*c*d*e - a*B*e^2 - (\text{Sqrt}[c]*(A*c*d^2 + 2*a*B*d*e - a*A*e^2))/\text{Sqrt}[a])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((4*a^{(3/4)}*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) + (e*(\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(B*d - A*e)*(3*c*d^2 + a*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/((8*a^{(1/4)}*c^{(1/4)}*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) + (a^{(3/4)}*e*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d -$

$\text{Sqrt}[a]*e^{2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e)}, 2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2)/(\text{Sqrt}[a]*e^{2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e)}, 2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2))/(\text{Sqrt}[a]*e^{2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e)}, 2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2))/(\text{Sqrt}[a]*e^{2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e)}, 2*\text{ArcTan}[(c^{1/4}*x)/a^{1/4}], 1/2))$

Rule 226

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[b/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[(a + b*x^4)/(a*(1 + q^2*x^2)^2])/(2*q*\text{Sqrt}[a + b*x^4]))* \text{EllipticF}[2*\text{ArcTan}[q*x], 1/2], x]] \text{ /; FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[b/a]$

Rule 1193

$\text{Int}[(d_) + (e_)*(x_)^2]*((a_) + (c_)*(x_)^4)^{(p_)}, x_Symbol] \text{ :> Simp}[(-x)*(d + e*x^2)*((a + c*x^4)^{(p + 1)/(4*a*(p + 1))}), x] + \text{Dist}[1/(4*a*(p + 1)), \text{Int}[\text{Simp}[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^{(p + 1)}, x], x] \text{ /; FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntegerQ}[2*p]$

Rule 1210

$\text{Int}[(d_) + (e_)*(x_)^2]/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[(a + c*x^4)/(a*(1 + q^2*x^2)^2])/(q*\text{Sqrt}[a + c*x^4]))* \text{EllipticE}[2*\text{ArcTan}[q*x], 1/2], x] \text{ /; EqQ}[e + d*q^2, 0] \text{ /; FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$

Rule 1212

$\text{Int}[(d_) + (e_)*(x_)^2]/\text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + c*x^4], x], x] \text{ /; NeQ}[e + d*q, 0] \text{ /; FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{PosQ}[c/a]$

Rule 1231

$\text{Int}[1/(((d_) + (e_)*(x_)^2)*\text{Sqrt}[(a_) + (c_)*(x_)^4]), x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(c*d + a*e*q)/(c*d^2 - a*e^2), \text{Int}[1/\text{Sqrt}[a + c*x^4], x], x] - \text{Dist}[(a*e*(e + d*q))/(c*d^2 - a*e^2), \text{Int}[(1 + q*x^2)/((d + e*x^2)*\text{Sqrt}[a + c*x^4]), x], x]] \text{ /; FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{NeQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{NeQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1238

$\text{Int}[(d_) + (e_)*(x_)^2]^{(q_)} / \text{Sqrt}[(a_) + (c_)*(x_)^4], x_Symbol] \text{ :> Simp} [(-e^2)*x*(d + e*x^2)^{(q + 1)}*(\text{Sqrt}[a + c*x^4]/(2*d*(q + 1)*(c*d^2 + a*e^2))), x] + \text{Dist}[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), \text{Int}[(d + e*x^2)^{(q + 1)}/\text{Sqrt}[a + c*x^4]*\text{Simp}[a*e^2*(2*q + 3) + 2*c*d^2*(q + 1) - 2*e*c*d*(q + 1)*x^2 + c*e^2*(2*q + 5)*x^4, x], x], x] \text{ /; FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{ILtQ}[q, -1]$

Rule 1721

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2]))], x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*((a + c*x^4)/(a*(A + B*x^2)^2))]/(4*d*e*A*q*Sqrt[a + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1723

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x], x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1729

```
Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2]/((d + e*x^2)*Sqrt[a + c*x^4]), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rule 1735

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^2 (a + cx^4)^{3/2}} dx &= \int \left(\frac{c(Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{(cd^2 + ae^2)^2 (a + cx^4)^{3/2}} + \frac{e}{(cd^2 + ae^2)} \right) dx \\
&= \frac{c \int \frac{Ac d^2 + 2a B d e - a A e^2 + (B c d^2 - 2 A c d e - a B e^2) x^2}{(a + c x^4)^{3/2}} dx}{(cd^2 + ae^2)^2} - \frac{(e(Bd - Ae)) \int \frac{1}{(d+ex^2)^2 \sqrt{a+cx^4}} dx}{cd^2 + ae^2} \\
&= \frac{cx(Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a(cd^2 + ae^2)^2 \sqrt{a + cx^4}} - \frac{e^3(Bd - Ae)x}{2d(cd^2 + ae^2)^2} \\
&= \frac{cx(Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a(cd^2 + ae^2)^2 \sqrt{a + cx^4}} - \frac{e^3(Bd - Ae)x}{2d(cd^2 + ae^2)^2} \\
&= \frac{cx(Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a(cd^2 + ae^2)^2 \sqrt{a + cx^4}} + \frac{\sqrt{c} e^2 (Bd - Ae)}{2d(cd^2 + ae^2)^2} \\
&= \frac{cx(Acd^2 + 2aBde - aAe^2 + (Bcd^2 - 2Acde - aBe^2)x^2)}{2a(cd^2 + ae^2)^2 \sqrt{a + cx^4}} + \frac{\sqrt{c} e^2 (Bd - Ae)}{2d(cd^2 + ae^2)^2}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.85, size = 427, normalized size = 0.29

$$\frac{\sqrt{\frac{15c}{2a}} \operatorname{ArcCosh}\left(\frac{Bd - Ae}{\sqrt{a + cx^4}}\right) \operatorname{EllipticE}\left[\operatorname{ArcSinh}\left(\frac{\sqrt{a + cx^4}}{\sqrt{a + cx^4}}\right), -1\right] + \frac{2a}{\sqrt{a + cx^4}} \left(-\sqrt{c} \sqrt{d - Bd^2 + 2Acde - aBe^2} \operatorname{EllipticE}\left[\operatorname{ArcSinh}\left(\frac{\sqrt{a + cx^4}}{\sqrt{a + cx^4}}\right), -1\right] + \left(\sqrt{c} d \sqrt{d - Bd^2 + 2Acde - aBe^2} + i \sqrt{c} \sqrt{d(Bd - Ae) + a(2Bd - Ae)} \right) \operatorname{EllipticF}\left[\operatorname{ArcSinh}\left(\frac{\sqrt{a + cx^4}}{\sqrt{a + cx^4}}\right), -1\right] + a(-5Bd^2 + 7Acde + aBe^2) \operatorname{EllipticPi}\left[\frac{-\frac{2a}{\sqrt{a + cx^4}}}{\sqrt{a + cx^4}}, \operatorname{ArcSinh}\left(\frac{\sqrt{a + cx^4}}{\sqrt{a + cx^4}}\right), -1\right] \right)}{2a \sqrt{\frac{15c}{2a}} (cd^2 + ae^2)^2 (d + ex^2) \sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*(a + c*x^4)^(3/2)),x]

[Out] (Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*(a*e^3*(-(B*d) + A*e)*x*(a + c*x^4) + c*d*x*(d + e*x^2))*(-(a*A*e^2) + B*c*d^2*x^2 + A*c*d*(d - 2*e*x^2) + a*B*e*(2*d - e*x^2)) - (d + e*x^2)*Sqrt[1 + (c*x^4)/a]*(-(Sqrt[a]*Sqrt[c]*d*(-(B*c*d^3) + 2*A*c*d^2*e + 2*a*B*d*e^2 - a*A*e^3)*EllipticE[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1]) + I*(Sqrt[c]*d*(Sqrt[c]*d - I*Sqrt[a]*e)*(A*c*d^2 + I*Sqrt[a]*Sqrt[c]*d*(B*d - A*e) + a*e*(2*B*d - A*e))*EllipticF[I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1] + a*e*(-5*B*c*d^3 + 7*A*c*d^2*e + a*B*d*e^2 + a*A*e^3)*EllipticPi[(-I)*Sqrt[a]*e/(Sqrt[c]*d), I*ArcSinh[Sqrt[(I*Sqrt[c])/Sqrt[a]]*x], -1)))/(2*a*Sqrt[(I*Sqrt[c])/Sqrt[a]]*(c*d^3 + a*d*e^2)^2*(d + e*x^2)*Sqrt[a + c*x^4])

Maple [C] Result contains complex when optimal does not.
time = 0.17, size = 1384, normalized size = 0.93

method	result	size
default	Expression too large to display	1384
elliptic	Expression too large to display	1664

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & B/e*(-2*c*(1/4/a*e/(a*e^2+c*d^2)*x^3-1/4*d/a/(a*e^2+c*d^2)*x)/((x^4+a/c)*c) \\ & ^{(1/2)}+1/2*c*d/a/(a*e^2+c*d^2)/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)} \\ & ^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(\\ & I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)+1/2*I/a^{(1/2)}*c^{(1/2)}*e/(a*e^2+c*d^2)/(I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)} \\ & /((c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)-1/2*I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)}*e/(a*e^2+c*d^2)/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)} \\ & *(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticE(x*(I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)})^{(1/2)},I)+1/(a*e^2+c*d^2)*e^2/d/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I \\ & /a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)} \\ & *EllipticPi(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I*a^{(1/2)}/c^{(1/2)}*e/d,(-I/a^{(1/2)}*c^{(1/2)} \\ & ^{(1/2)})^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)})+(A*e-B*d)/e*(1/2*e^4/(a*e^2+c*d^2) \\ & ^2/d*x*(c*x^4+a)^{(1/2)}/(e*x^2+d)-2*c*(1/2/a*c*d*e/(a*e^2+c*d^2)^2*x^3+1/4/ \\ & a*(a*e^2-c*d^2)/(a*e^2+c*d^2)^2*x)/((x^4+a/c)*c)^{(1/2)}-1/(I/a^{(1/2)}*c^{(1/2)} \\ & ^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c* \\ & x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)*e^2*c/(a*e^2+c*d^2)^2 \\ & +1/2/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticF(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I \\ & ^{(1/2)}*c^2/a/(a*e^2+c*d^2)^2*d^2-1/2*I*a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(1/2)} \\ & ^{(1/2)}*e^3/d/(a*e^2+c*d^2)^2*EllipticF(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)+1/2*I* \\ & a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(1/2)}*e^3/d/(a*e^2+c*d^2)^2*Elliptic \\ & E(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)+I/a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I \\ & /a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)} \\ & ^{(1/2)}*c^{(3/2)}*d*e/(a*e^2+c*d^2)^2*EllipticF(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)-I/a^{(1/2)} \\ & /((I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(3/2)}*d*e/(a*e^2+c*d^2)^2*EllipticE(x \\ & *(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I)+1/2*e^4/d^2/(a*e^2+c*d^2)^2/(I/a^{(1/2)}*c^{(1/2)} \\ & ^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c \\ & *x^4+a)^{(1/2)}*EllipticPi(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)},I*a^{(1/2)}/c^{(1/2)}*e/d, \\ & (-I/a^{(1/2)}*c^{(1/2)})^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)})*a+7/2*e^2/(a*e^2+c*d^2) \\ & ^2/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)} \\ & ^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*EllipticPi(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, \end{aligned}$$

$I \cdot a^{(1/2)} / c^{(1/2)} \cdot e / d, (-I / a^{(1/2)} \cdot c^{(1/2)})^{(1/2)} / (I / a^{(1/2)} \cdot c^{(1/2)})^{(1/2)} \cdot c)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(x^2*e + d)^2), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(a + cx^4)^{\frac{3}{2}} (d + ex^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+a)**(3/2),x)

[Out] Integral((A + B*x**2)/((a + c*x**4)**(3/2)*(d + e*x**2)**2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(x^2*e + d)^2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(cx^4 + a)^{3/2} (ex^2 + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^2), x)
```

```
[Out] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^2), x)
```

$$3.14 \quad \int \frac{A+Bx^2}{(d+ex^2)^3 (a+cx^4)^{3/2}} dx$$

Optimal. Leaf size=2452

result too large to display

```
[Out] -1/4*e^(3/2)*(a*e^2+3*c*d^2)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(3/2)/(a*e^2+c*d^2)^(7/2)-3/16*e^(3/2)*(-A*e+B*d)*(a^2*e^4+2*a*c*d^2*e^2+5*c^2*d^4)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/d^(5/2)/(a*e^2+c*d^2)^(7/2)-1/2*c*e^(3/2)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*arctan(x*(a*e^2+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+a)^(1/2))/(a*e^2+c*d^2)^(7/2)/d^(1/2)+1/2*c*x*(A*c*d*(-3*a*e^2+c*d^2)+a*B*e*(-a*e^2+3*c*d^2)+c*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*x^2)/a/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/4*e^3*(-A*e+B*d)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^2/(e*x^2+d)^2-3/8*e^3*(-A*e+B*d)*(a*e^2+3*c*d^2)*x*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^3/(e*x^2+d)-1/2*e^3*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^3/(e*x^2+d)-1/2*c^(3/2)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*x*(c*x^4+a)^(1/2)/a/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))+3/8*e^2*(-A*e+B*d)*(a*e^2+3*c*d^2)*x*c^(1/2)*(c*x^4+a)^(1/2)/d^2/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))+1/2*e^2*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*x*c^(1/2)*(c*x^4+a)^(1/2)/d/(a*e^2+c*d^2)^3/(a^(1/2)+x^2*c^(1/2))-3/8*a^(1/4)*c^(1/4)*e^2*(-A*e+B*d)*(a*e^2+3*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d^2/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/2*a^(1/4)*c^(1/4)*e^2*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/d/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)+1/2*c^(5/4)*(A*a*e^3-3*A*c*d^2*e-3*B*a*d*e^2+B*c*d^3)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(3/4)/(a*e^2+c*d^2)^3/(c*x^4+a)^(1/2)-1/2*c^(1/4)*e*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*2^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)/a^(1/4)/(a*e^2+c*d^2)^3/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+a)^(1/2)+1/8*e*(a*e^2+3*c*d^2)*(-2*A*c*d*e-B*a*e^2+B*c*d^2)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^2/d/e/a^(1/2)/c^(1/2),1/2*2^(1/2))*(e*a^(1/2)+d*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+a)/(a^(1/2)+x^2*c^(1/2)))^2)^(1/2)
```

$$\begin{aligned} & /a^{1/4}/c^{1/4}/d^2/(a^2e^2+cd^2)^{3/2}/(-ea^{1/2}+dc^{1/2})/(cx^4+a)^{1/2} \\ & +3/32e*(-Ae+Bd)*(a^2e^4+2a^2cd^2e^2+5c^2d^4)*(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticPi}(\sin(2\arctan(c^{1/4}x/a^{1/4})), -1/4*(-ea^{1/2}+dc^{1/2})^2/d/e/a^{1/2}/c^{1/2}, 1/2*2^{1/2})*(ea^{1/2}+dc^{1/2})*(a^{1/2}+x^2c^{1/2})*((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^2)^{1/2}/a^{1/4}/c^{1/4}/d^3/(a^2e^2+cd^2)^{3/2}/(-ea^{1/2}+dc^{1/2})/(cx^4+a)^{1/2}+1/4c^{3/4}*(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticF}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2*2^{1/2})*(A^2c^2d^3-a^2B^2e^3+3A^2cd^2e^2-c^{3/2}d^2*(-3A^2e+B^2d)*a^{1/2}+a^{3/2}e^2*(-Ae+3Bd)*c^{1/2})*(a^{1/2}+x^2c^{1/2})*((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^2)^{1/2}/a^{5/4}/(a^2e^2+cd^2)^{3/2}/(cx^4+a)^{1/2}+1/4a^{3/4}c^{3/4}e*(A^2a^2e^3-3A^2cd^2e^2-3B^2a^2d^2e^2+B^2cd^3)*(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticPi}(\sin(2\arctan(c^{1/4}x/a^{1/4})), -1/4*(-ea^{1/2}+dc^{1/2})^2/d/e/a^{1/2}/c^{1/2}, 1/2*2^{1/2})*(a^{1/2}+x^2c^{1/2})*(e+dc^{1/2}/a^{1/2})^2*((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^2)^{1/2}/d/(-a^2e^2+cd^2)/(a^2e^2+cd^2)^{3/2}/(cx^4+a)^{1/2}-1/8c^{1/4}e*(-Ae+Bd)*(\cos(2\arctan(c^{1/4}x/a^{1/4}))^2)^{1/2}/\cos(2\arctan(c^{1/4}x/a^{1/4})) * \text{EllipticF}(\sin(2\arctan(c^{1/4}x/a^{1/4})), 1/2*2^{1/2})*(a^{1/2}+x^2c^{1/2})*(4c^2d^2+3a^2e^2-d^2ea^{1/2})*c^{1/2})*((cx^4+a)/(a^{1/2}+x^2c^{1/2}))^2)^{1/2}/a^{1/4}/d^2/(a^2e^2+cd^2)^2/(-ea^{1/2}+dc^{1/2})/(cx^4+a)^{1/2} \end{aligned}$$
Rubi [A]

time = 2.62, antiderivative size = 2452, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 11, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.393$, Rules used = {1735, 1193, 1212, 226, 1210, 1238, 1711, 1729, 1723, 1721, 1231}

Too large to display

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^3*(a + c*x^4)^(3/2)), x]

[Out] $(cx*(Acd*(cd^2 - 3ae^2) + aB^2e*(3cd^2 - ae^2) + c*(Bcd^3 - 3A^2cd^2e - 3a^2Bd^2e^2 + aA^2e^3)*x^2))/(2a*(cd^2 + ae^2)^3\sqrt{a + cx^4}) + (3\sqrt{c}e^2*(Bd - Ae)*(3cd^2 + ae^2)*x\sqrt{a + cx^4})/(8d^2*(cd^2 + ae^2)^3*(\sqrt{a} + \sqrt{c}x^2)) + (\sqrt{c}e^2*(Bcd^2 - 2A^2cd^2e - aB^2e^2)*x\sqrt{a + cx^4})/(2d*(cd^2 + ae^2)^3*(\sqrt{a} + \sqrt{c}x^2)) - (c^{3/2}*(Bcd^3 - 3A^2cd^2e - 3a^2Bd^2e^2 + aA^2e^3)*x\sqrt{a + cx^4})/(2a*(cd^2 + ae^2)^3*(\sqrt{a} + \sqrt{c}x^2)) - (e^3*(Bd - Ae)*x\sqrt{a + cx^4})/(4d*(cd^2 + ae^2)^2*(d + e*x^2)^2) - (3e^3*(Bd - Ae)*(3cd^2 + ae^2)*x\sqrt{a + cx^4})/(8d^2*(cd^2 + ae^2)^3*(d + e*x^2)) - (e^3*(Bcd^2 - 2A^2cd^2e - aB^2e^2)*x\sqrt{a + cx^4})/(2d*(cd^2 + ae^2)^3*(d + e*x^2)) - (e^{3/2}*(3cd^2 + ae^2)*(Bcd^2 - 2A^2cd^2e - aB^2e^2)*\text{ArcTan}[(\sqrt{cd^2 + ae^2}x)/(\sqrt{d}\sqrt{e}\sqrt{a + cx^4})])/(4d^{3/2}*(cd^2 + ae^2)^{7/2}) - (ce^{3/2}*(Bcd^3 - 3A^2cd^2e -$

$$\begin{aligned}
& 3*a*B*d*e^2 + a*A*e^3)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])]/(2*\text{Sqrt}[d]*(c*d^2 + a*e^2)^{(7/2)}) - (3*e^{(3/2)}*(B*d - A*e)* \\
& (5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4)*\text{ArcTan}[(\text{Sqrt}[c*d^2 + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + c*x^4])]/(16*d^{(5/2)}*(c*d^2 + a*e^2)^{(7/2)}) - (3*a^{(1/4)}*c^{(1/4)}*e^2*(B*d - A*e)*(3*c*d^2 + a*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (8*d^2*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) - (a^{(1/4)}*c^{(1/4)}*e^2*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (2*d*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) + (c^{(5/4)}*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (2*a^{(3/4)}*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) - (c^{(1/4)}*e*(B*d - A*e)*(4*c*d^2 - \text{Sqrt}[a]*\text{Sqrt}[c]*d*e + 3*a*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (8*a^{(1/4)}*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) - (c^{(1/4)}*e*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (2*a^{(1/4)}*d*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^2*\text{Sqrt}[a + c*x^4]) - (c^{(5/4)}*e*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (2*a^{(1/4)}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) + (c^{(3/4)}*(A*c^2*d^3 - a^2*B*e^3 - \text{Sqrt}[a]*c^{(3/2)}*d^2*(B*d - 3*A*e) + 3*a*c*d*e*(B*d - A*e) + a^{(3/2)}*\text{Sqrt}[c]*e^2*(3*B*d - A*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (4*a^{(5/4)}*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) + (e*(\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(3*c*d^2 + a*e^2)*(B*c*d^2 - 2*A*c*d*e - a*B*e^2)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (8*a^{(1/4)}*c^{(1/4)}*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) + (a^{(3/4)}*c^{(3/4)}*e*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(B*c*d^3 - 3*A*c*d^2*e - 3*a*B*d*e^2 + a*A*e^3)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (4*d*(c*d^2 - a*e^2)*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4]) + (3*e*(\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(B*d - A*e)*(5*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], 1/2])/ (32*a^{(1/4)}*c^{(1/4)}*d^3*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 + a*e^2)^3*\text{Sqrt}[a + c*x^4])
\end{aligned}$$

Rule 226

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^4], x_Symbol] :-> With[{q = Rt[b/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^4]))*EllipticF[2*ArcTan[q*x], 1/2], x]] /; FreeQ[{a, b}, x] && PosQ[b/a]

```

Rule 1193

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[(-x
)*(d + e*x^2)*((a + c*x^4)^(p + 1)/(4*a*(p + 1))), x] + Dist[1/(4*a*(p + 1)
), Int[Simp[d*(4*p + 5) + e*(4*p + 7)*x^2, x]*(a + c*x^4)^(p + 1), x], x] /
; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && IntegerQ
[2*p]
```

Rule 1210

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*
(1 + q^2*x^2)*(Sqrt[(a + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + c*x^4]))*E
llipticE[2*ArcTan[q*x], 1/2], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, c, d, e
}, x] && PosQ[c/a]
```

Rule 1212

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := With[{q =
Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + c*x^4], x], x] - Dist[e/q, I
nt[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, c,
d, e}, x] && PosQ[c/a]
```

Rule 1231

```
Int[1/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (c_.)*(x_)^4]), x_Symbol] := With[
{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4]
, x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^
2)*Sqrt[a + c*x^4]), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2
, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rule 1238

```
Int[((d_) + (e_.)*(x_)^2)^(q_)/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol] := Simp
[(-e^2)*x*(d + e*x^2)^(q + 1)*(Sqrt[a + c*x^4]/(2*d*(q + 1)*(c*d^2 + a*e^2)
)), x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), Int[((d + e*x^2)^(q + 1)/Sqr
t[a + c*x^4])*Simp[a*e^2*(2*q + 3) + 2*c*d^2*(q + 1) - 2*e*c*d*(q + 1)*x^2
+ c*e^2*(2*q + 5)*x^4, x], x] /; FreeQ[{a, c, d, e}, x] && ILtQ[q, -1]
```

Rule 1711

```
Int[((P4x_)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (c_.)*(x_)^4], x_Symbol
] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]
}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + c*x^4]/(
2*d*(q + 1)*(c*d^2 + a*e^2))), x] + Dist[1/(2*d*(q + 1)*(c*d^2 + a*e^2)), I
```

```

nt[((d + e*x^2)^(q + 1)/Sqrt[a + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2
*q + 3) + 2*c*d^2*(q + 1)) + 2*d*(B*c*d - A*c*e + a*C*e)*(q + 1)*x^2 + c*(C
*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x], x] /; FreeQ[{a, c, d, e}, x]
&& PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[c*d^2 + a*e^2, 0] && ILt
Q[q, -1]

```

Rule 1721

```

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[c*(d/e)
+ a*(e/d), 2]*(x/Sqrt[a + c*x^4])]/(2*d*e*Rt[c*(d/e) + a*(e/d), 2])), x]
+ Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*((a + c*x^4)/(a*(A + B*x^2)^2))]/(
4*d*e*A*q*Sqrt[a + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)],
2*ArcTan[q*x], 1/2], x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e
^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

```

Rule 1723

```

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q)
)/(c*d^2 - a*e^2), Int[1/Sqrt[a + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e
+ d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + c*x^4]), x],
x] /; FreeQ[{a, c, d, e, A, B}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2
- a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]

```

Rule 1729

```

Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (c_)*(x_)^4]), x_Symbol] :=
With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff
[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + c*x^4], x], x] + Dist
[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2)
*Sqrt[a + c*x^4]), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[P4x, x^2, 2]
&& NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]

```

Rule 1735

```

Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol]
:= Int[ExpandIntegrand[1/Sqrt[a + c*x^4], Px*(d + e*x^2)^q*(a + c*x^4)^(p
+ 1/2), x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a
*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]

```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^3 (a + cx^4)^{3/2}} dx &= \int \left(\frac{c(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{(cd^2 + ae^2)^3 (a + cx^4)^{3/2}} \right. \\
&= \frac{c \int \frac{Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3)x^2}{(a + cx^4)^{3/2}} dx}{(cd^2 + ae^2)^3} - \frac{(e(Bd - Ae))}{(cd^2 + ae^2)^3} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}} \\
&= \frac{cx(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 11.49, size = 630, normalized size = 0.26

$$\frac{\sqrt{c} \sqrt{a} \sqrt{a + cx^4} \left(\frac{c(Acd(cd^2 - 3ae^2) + aBe(3cd^2 - ae^2) + c(Bcd^3 - 3Acd^2e - 3aBde^2 + aAe^3))}{(cd^2 + ae^2)^3} - \frac{(e(Bd - Ae))}{(cd^2 + ae^2)^3} \right)}{2a(cd^2 + ae^2)^3 \sqrt{a + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^3*(a + c*x^4)^(3/2)), x]

[Out] (Sqrt[(I*Sqrt[c])/Sqrt[a]]*d*x*(-2*a*d*e^3*(B*d - A*e)*(c*d^2 + a*e^2)*(a + c*x^4) + a*e^3*(-13*B*c*d^3 + 17*A*c*d^2*e + a*B*d*e^2 + 3*a*A*e^3)*(d + e*x^2)*(a + c*x^4) + 4*c*d^2*(d + e*x^2)^2*(B*(-a^2*e^3) + c^2*d^3*x^2 + 3*a*c*d*e*(d - e*x^2)) + A*c*(c*d^2*(d - 3*e*x^2) + a*e^2*(-3*d + e*x^2)))) - (d + e*x^2)^2*Sqrt[1 + (c*x^4)/a]*(Sqrt[a]*Sqrt[c]*d*(3*A*e*(-4*c^2*d^4 +

$$7*a*c*d^2*e^2 + a^2*e^4) + B*(4*c^2*d^5 - 25*a*c*d^3*e^2 + a^2*d*e^4)*\text{EllipticE}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1] + I*(\text{Sqrt}[c]*d*(\text{Sqrt}[c]*d - I*\text{Sqrt}[a]*e)*(4*A*c^2*d^4 + (4*I)*\text{Sqrt}[a]*c^{(3/2)}*d^3*(B*d - 2*A*e) + 19*a*c*d^2*e*(B*d - A*e) - (2*I)*a^{(3/2)}*\text{Sqrt}[c]*d*e^2*(3*B*d - A*e) - a^2*e^3*(B*d + 3*A*e))*\text{EllipticF}[I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1] + a*e*(3*A*e*(21*c^2*d^4 + 2*a*c*d^2*e^2 + a^2*e^4) + B*(-35*c^2*d^5 + 26*a*c*d^3*e^2 + a^2*d*e^4))*\text{EllipticPi}[((-I)*\text{Sqrt}[a]*e)/(\text{Sqrt}[c]*d), I*\text{ArcSinh}[\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*x], -1)))/(8*a*\text{Sqrt}[(I*\text{Sqrt}[c])/ \text{Sqrt}[a]]*(c*d^3 + a*d*e^2)^{3/2}*(d + e*x^2)^2*\text{Sqrt}[a + c*x^4])$$

Maple [C] Result contains complex when optimal does not.

time = 0.17, size = 2326, normalized size = 0.95

method	result	size
default	Expression too large to display	2326
elliptic	Expression too large to display	2596

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2), x, method=_RETURNVERBOSE)`

[Out]
$$\frac{(Ae-Bd)/e*(1/4*e^4/(a*e^2+c*d^2)^2/d*x*(c*x^4+a)^{(1/2)}/(e*x^2+d)^2+1/8*e^4*(3*a*e^2+17*c*d^2)/d^2/(a*e^2+c*d^2)^3*x*(c*x^4+a)^{(1/2)}/(e*x^2+d)-2*c*(-1/4*c*e*(a*e^2-3*c*d^2)/a/(a*e^2+c*d^2)^3*x^3+1/4*c*d*(3*a*e^2-c*d^2)/a/(a*e^2+c*d^2)^3*x)/((x^4+a/c)*c)^{(1/2)}-1/8/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*\text{EllipticF}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)*e^4*c/d/(a*e^2+c*d^2)^3*a-27/8/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*\text{EllipticF}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)*e^2*c^2*d/(a*e^2+c*d^2)^3+1/2/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*\text{EllipticF}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)*c^3*d^3/a/(a*e^2+c*d^2)^3+3/2*I/a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(5/2)}*e/(a*e^2+c*d^2)^3*d^2*\text{EllipticE}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)-3/2*I/a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(5/2)}*e/(a*e^2+c*d^2)^3*d^2*\text{EllipticE}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)+3/8*I*a^{(3/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(1/2)}*e^5/d^2/(a*e^2+c*d^2)^3*\text{EllipticE}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)-3/8*I*a^{(3/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(1/2)}*e^5/d^2/(a*e^2+c*d^2)^3*\text{EllipticF}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)-21/8*I*a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(3/2)}*e^3/(a*e^2+c*d^2)^3*\text{EllipticF}(x*(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}, I)+21/8*I*a^{(1/2)}/(I/a^{(1/2)}*c^{(1/2)})^{(1/2)}*(1-I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}*(1+I/a^{(1/2)}*c^{(1/2)}*x^2)^{(1/2)}/(c*x^4+a)^{(1/2)}*c^{(1/2)}$$

$$\begin{aligned} & 3/2)*e^3/(a*e^2+c*d^2)^3*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+3/8*e^6/d \\ & ^3/(a*e^2+c*d^2)^3/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2) \\ & *(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c \\ & ^{(1/2)})^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c \\ & ^{(1/2)})^(1/2))*a^2+3/4*e^4/(a*e^2+c*d^2)^3/d/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I \\ & /a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2) \\ & *EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c \\ & ^{(1/2)})^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*a*c+63/8*e^2*d/(a*e^2+c*d^2)^3/(I/ \\ & a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2) \\ & *x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2),I*a^(1/2) \\ &)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2))*c^2)+B/ \\ & e*(1/2*e^4/(a*e^2+c*d^2)^2/d*x*(c*x^4+a)^(1/2)/(e*x^2+d)-2*c*(1/2/a*c*d*e/(\\ & a*e^2+c*d^2)^2*x^3+1/4/a*(a*e^2-c*d^2)/(a*e^2+c*d^2)^2*x)/((x^4+a/c)*c)^(1/ \\ & 2)-1/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2) \\ & *c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a^(1/2)*c^(1/2))^(1/2),I \\ &)*e^2*c/(a*e^2+c*d^2)^2+1/2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)* \\ & x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticF(x*(I/a \\ & ^{(1/2)*c^(1/2)})^(1/2),I)*c^2/a/(a*e^2+c*d^2)^2*d^2-1/2*I*a^(1/2)/(I/a^(1/2) \\ & *c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(\\ & 1/2)/(c*x^4+a)^(1/2)*c^(1/2)*e^3/d/(a*e^2+c*d^2)^2*EllipticF(x*(I/a^(1/2)*c \\ & ^{(1/2)})^(1/2),I)+1/2*I*a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/ \\ & 2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*c^(1/2)*e^3/d \\ & /(a*e^2+c*d^2)^2*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+I/a^(1/2)/(I/a^(1 \\ & /2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2 \\ &)^(1/2)/(c*x^4+a)^(1/2)*c^(3/2)*d*e/(a*e^2+c*d^2)^2*EllipticF(x*(I/a^(1/2)* \\ & c^(1/2))^(1/2),I)-I/a^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)* \\ & x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*c^(3/2)*d*e/(a*e \\ & ^2+c*d^2)^2*EllipticE(x*(I/a^(1/2)*c^(1/2))^(1/2),I)+1/2*e^4/d^2/(a*e^2+c*d \\ & ^2)^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)*x^2)^(1/2)*(1+I/a^(1/2) \\ &)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/a^(1/2)*c^(1/2))^(1/2) \\ & ,I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/a^(1/2)*c^(1/2))^(1/2) \\ &)*a+7/2*e^2/(a*e^2+c*d^2)^2/(I/a^(1/2)*c^(1/2))^(1/2)*(1-I/a^(1/2)*c^(1/2)* \\ & x^2)^(1/2)*(1+I/a^(1/2)*c^(1/2)*x^2)^(1/2)/(c*x^4+a)^(1/2)*EllipticPi(x*(I/ \\ & a^(1/2)*c^(1/2))^(1/2),I*a^(1/2)/c^(1/2)*e/d,(-I/a^(1/2)*c^(1/2))^(1/2)/(I/ \\ & a^(1/2)*c^(1/2))^(1/2))*c) \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(x^2*e + d)^3), x)

Fricas [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+a)**(3/2),x)

[Out] Timed out

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + a)^(3/2)*(x^2*e + d)^3), x)

Mupad [F]
time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{(c x^4 + a)^{3/2} (e x^2 + d)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^3),x)

[Out] int((A + B*x^2)/((a + c*x^4)^(3/2)*(d + e*x^2)^3), x)

$$3.15 \quad \int \frac{(A+Bx^2)(d+ex^2)^q}{a+cx^4} dx$$

Optimal. Leaf size=169

$$\frac{\left(A - \frac{\sqrt{-a}B}{\sqrt{c}}\right) x(d+ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d}\right)}{2a} + \frac{\left(A + \frac{\sqrt{-a}B}{\sqrt{c}}\right) x(d+ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; \frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d}\right)}{2a}$$

[Out] 1/2*x*(e*x^2+d)^q*AppellF1(1/2,-q,1,3/2,-e*x^2/d,-x^2*c^(1/2)/(-a)^(1/2))*(A-B*(-a)^(1/2)/c^(1/2))/a/((1+e*x^2/d)^q)+1/2*x*(e*x^2+d)^q*AppellF1(1/2,1,-q,3/2,x^2*c^(1/2)/(-a)^(1/2),-e*x^2/d)*(A+B*(-a)^(1/2)/c^(1/2))/a/((1+e*x^2/d)^q)

Rubi [A]

time = 0.14, antiderivative size = 169, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {1707, 441, 440}

$$\frac{x\left(A - \frac{\sqrt{-a}B}{\sqrt{c}}\right)(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d}\right)}{2a} + \frac{x\left(\frac{\sqrt{-a}B}{\sqrt{c}} + A\right)(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; \frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d}\right)}{2a}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]

[Out] ((A - (Sqrt[-a]*B)/Sqrt[c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, -(Sqrt[c]*x^2)/Sqrt[-a], -(e*x^2)/d])/(2*a*(1 + (e*x^2)/d)^q) + ((A + (Sqrt[-a]*B)/Sqrt[c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (Sqrt[c]*x^2)/Sqrt[-a], -(e*x^2)/d])/(2*a*(1 + (e*x^2)/d)^q)

Rule 440

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x]
;/; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1]
&& (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])
```

Rule 441

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol]
:> Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]),
Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x]
;/; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])
```

Rule 1707

```
Int[(Px_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol]
:> Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + c*x^4)^p, x], x]
;/; FreeQ[{a
```

, c, d, e, q}, x] && PolyQ[Px, x^2] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[p]

Rubi steps

$$\begin{aligned}
 \int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx &= \int \left(-\frac{(\sqrt{-a} B + A\sqrt{c})(d + ex^2)^q}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} - \sqrt{c}x^2)} + \frac{(\sqrt{-a} B - A\sqrt{c})(d + ex^2)^q}{2\sqrt{-a}\sqrt{c}(\sqrt{-a} + \sqrt{c}x^2)} \right) dx \\
 &= -\left(\frac{1}{2} \left(\frac{A}{\sqrt{-a}} + \frac{B}{\sqrt{c}} \right) \int \frac{(d + ex^2)^q}{\sqrt{-a} - \sqrt{c}x^2} dx \right) + \frac{1}{2} \left(\frac{aA}{(-a)^{3/2}} + \frac{B}{\sqrt{c}} \right) \int \frac{(d + ex^2)^q}{\sqrt{-a} + \sqrt{c}x^2} dx \\
 &= -\left(\frac{1}{2} \left(\left(\frac{A}{\sqrt{-a}} + \frac{B}{\sqrt{c}} \right) (d + ex^2)^q \left(1 + \frac{ex^2}{d} \right)^{-q} \right) \int \frac{\left(1 + \frac{ex^2}{d} \right)^q}{\sqrt{-a} - \sqrt{c}x^2} dx \right) \\
 &= \frac{\left(A - \frac{\sqrt{-a} B}{\sqrt{c}} \right) x (d + ex^2)^q \left(1 + \frac{ex^2}{d} \right)^{-q} F_1 \left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{\sqrt{c}x^2}{\sqrt{-a}}, -\frac{ex^2}{d} \right)}{2a}
 \end{aligned}$$

Mathematica [F]

time = 0.40, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + cx^4} dx$$

Verification is not applicable to the result.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]

[Out] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x]

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a), x)

[Out] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^q/(c*x^4 + a), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="fricas")

[Out] integral((B*x^2 + A)*(x^2*e + d)^q/(c*x^4 + a), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**q/(c*x**4+a),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+a),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^q/(c*x^4 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^q)/(a + c*x^4), x)

$$3.16 \quad \int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx$$

Optimal. Leaf size=48

$$\frac{\sqrt{2}(2+x^2)E(\tan^{-1}(x)|\frac{1}{2})}{\sqrt{\frac{2+x^2}{1+x^2}}\sqrt{2+3x^2+x^4}}$$

[Out] $(x^2+2)*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*\text{EllipticE}(x/(x^2+1)^{(1/2)},1/2*2^{(1/2)})*2^{(1/2)}/((x^2+2)/(x^2+1))^{(1/2)}/(x^4+3*x^2+2)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 27, $\frac{\text{number of rules}}{\text{integrand size}} = 0.074$, Rules used = {1470, 422}

$$\frac{\sqrt{2}(x^2+2)E(\text{ArcTan}(x)|\frac{1}{2})}{\sqrt{\frac{x^2+2}{x^2+1}}\sqrt{x^4+3x^2+2}}$$

Antiderivative was successfully verified.

[In] Int[(2 + x^2)/((1 + x^2)*Sqrt[2 + 3*x^2 + x^4]),x]

[Out] (Sqrt[2]*(2 + x^2)*EllipticE[ArcTan[x], 1/2])/((Sqrt[(2 + x^2)/(1 + x^2)]*Sqrt[2 + 3*x^2 + x^4])

Rule 422

Int[Sqrt[(a_) + (b_)*(x_)^2]/((c_) + (d_)*(x_)^2)^(3/2), x_Symbol] := Simp[(Sqrt[a + b*x^2]/(c*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[c*((a + b*x^2)/(a*(c + d*x^2))])))*EllipticE[ArcTan[Rt[d/c, 2]*x], 1 - b*(c/(a*d))], x] /; FreeQ[{a, b, c, d}, x] && PosQ[b/a] && PosQ[d/c]

Rule 1470

Int[((d_) + (e_)*(x_)^(n_))^(q_)*((f_) + (g_)*(x_)^(n_))^(r_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Dist[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d + e*x^n)^FracPart[p]*(a/d + (c*x^n)/e)^FracPart[p]), Int[(d + e*x^n)^(p + q)*(f + g*x^n)^r*(a/d + (c/e)*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q, r}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p]

Rubi steps

$$\int \frac{2+x^2}{(1+x^2)\sqrt{2+3x^2+x^4}} dx = \frac{\left(\sqrt{1+x^2}\sqrt{2+x^2}\right) \int \frac{\sqrt{2+x^2}}{(1+x^2)^{3/2}} dx}{\sqrt{2+3x^2+x^4}}$$

$$= \frac{\sqrt{2}(2+x^2)E(\tan^{-1}(x)|_{\frac{1}{2}})}{\sqrt{\frac{2+x^2}{1+x^2}}\sqrt{2+3x^2+x^4}}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.10, size = 94, normalized size = 1.96

$$\frac{2x + x^3 + i\sqrt{1+x^2}\sqrt{2+x^2}E\left(i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right) - i\sqrt{1+x^2}\sqrt{2+x^2}F\left(i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right)}{\sqrt{2+3x^2+x^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(2 + x^2)/((1 + x^2)*Sqrt[2 + 3*x^2 + x^4]), x]

[Out] (2*x + x^3 + I*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticE[I*ArcSinh[x/Sqrt[2]]], 2) - I*Sqrt[1 + x^2]*Sqrt[2 + x^2]*EllipticF[I*ArcSinh[x/Sqrt[2]], 2])/Sqrt[2 + 3*x^2 + x^4]

Maple [C] Result contains complex when optimal does not.

time = 0.16, size = 81, normalized size = 1.69

method	result	si
risch	$\frac{x(x^2+2)}{\sqrt{x^4+3x^2+2}} - \frac{i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\left(\text{EllipticF}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right) - \text{EllipticE}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right)\right)}{2\sqrt{x^4+3x^2+2}}$	8
default	$\frac{(x^2+2)x}{\sqrt{(x^2+2)(x^2+1)}} - \frac{i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\left(\text{EllipticF}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right) - \text{EllipticE}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right)\right)}{2\sqrt{x^4+3x^2+2}}$	8
elliptic	$\frac{(x^2+2)x}{\sqrt{(x^2+2)(x^2+1)}} - \frac{i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\left(\text{EllipticF}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right) - \text{EllipticE}\left(\frac{i\sqrt{2}x}{2}, \sqrt{2}\right)\right)}{2\sqrt{x^4+3x^2+2}}$	8

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2), x, method=_RETURNVERBOSE)

[Out] (x^2+2)*x/((x^2+2)*(x^2+1))^(1/2)-1/2*I*2^(1/2)*(2*x^2+4)^(1/2)*(x^2+1)^(1/2)/(x^4+3*x^2+2)^(1/2)*(EllipticF(1/2*I*2^(1/2)*x, 2^(1/2))-EllipticE(1/2*I*2^(1/2)*x, 2^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((x^2 + 2)/(sqrt(x^4 + 3*x^2 + 2)*(x^2 + 1)), x)
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly
1 arguments (2 given)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 + 2}{\sqrt{(x^2 + 1)(x^2 + 2)} (x^2 + 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((x**2+2)/(x**2+1)/(x**4+3*x**2+2)**(1/2),x)
```

```
[Out] Integral((x**2 + 2)/(sqrt((x**2 + 1)*(x**2 + 2))*(x**2 + 1)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((x^2+2)/(x^2+1)/(x^4+3*x^2+2)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((x^2 + 2)/(sqrt(x^4 + 3*x^2 + 2)*(x^2 + 1)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^2 + 2}{(x^2 + 1) \sqrt{x^4 + 3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^2 + 2)/((x^2 + 1)*(3*x^2 + x^4 + 2)^(1/2)),x)
```

```
[Out] int((x^2 + 2)/((x^2 + 1)*(3*x^2 + x^4 + 2)^(1/2)), x)
```

$$3.17 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=755

$$\frac{e(7Ace(15cd - 4be) + B(105c^2d^2 + 24b^2e^2 - ce(84bd + 25ae)))x\sqrt{a+bx^2+cx^4}}{105c^3} + \frac{e^2(21Bcd - 6bBe + 7Ac)}{35c^2}$$

[Out] 1/105*e*(7*A*c*e*(-4*b*e+15*c*d)+B*(105*c^2*d^2+24*b^2*e^2-c*e*(25*a*e+84*b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/c^3+1/35*e^2*(7*A*c*e-6*B*b*e+21*B*c*d)*x^3*(c*x^4+b*x^2+a)^(1/2)/c^2+1/7*B*e^3*x^5*(c*x^4+b*x^2+a)^(1/2)/c+1/105*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))*x*(c*x^4+b*x^2+a)^(1/2)/c^(7/2)/(a^(1/2)+x^2*c^(1/2))-1/105*a^(1/4)*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(15/4)/(c*x^4+b*x^2+a)^(1/2)+1/210*a^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*(7*A*c*e*(45*c^2*d^2+8*b^2*e^2-3*c*e*(3*a*e+10*b*d))+B*(105*c^3*d^3-48*b^3*e^3-21*c^2*d*e*(9*a*e+10*b*d)+8*b*c*e^2*(13*a*e+21*b*d)))+(7*A*c*(4*a*b*e^3-15*a*c*d*e^2+15*c^2*d^3)-a*B*e*(105*c^2*d^2+24*b^2*e^2-c*e*(25*a*e+84*b*d)))*c^(1/2)/a^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/c^(15/4)/(c*x^4+b*x^2+a)^(1/2)

Rubi [A]

time = 0.85, antiderivative size = 755, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.121$,

Rules used = {1693, 1211, 1117, 1209}

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] (e*(7*A*c*e*(15*c*d - 4*b*e) + B*(105*c^2*d^2 + 24*b^2*e^2 - c*e*(84*b*d + 25*a*e)))*x*Sqrt[a + b*x^2 + c*x^4]/(105*c^3) + (e^2*(21*B*c*d - 6*b*B*e + 7*A*c*e)*x^3*Sqrt[a + b*x^2 + c*x^4]/(35*c^2) + (B*e^3*x^5*Sqrt[a + b*x^2 + c*x^4])/(7*c) + ((7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a*e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2*(21*b*d + 13*a*e)))*x*Sqrt[a + b*x^2 + c*x^4]/(105*c^(7/2)*(Sqrt[a] + Sqr

$t[c]*x^2) - (a^{1/4}*(7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a*e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2*(21*b*d + 13*a*e)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^{1/4}*x)/a^{1/4}], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(105*c^{15/4}*Sqrt[a + b*x^2 + c*x^4]) + (a^{1/4}*(7*A*c*e*(45*c^2*d^2 + 8*b^2*e^2 - 3*c*e*(10*b*d + 3*a*e)) + B*(105*c^3*d^3 - 48*b^3*e^3 - 21*c^2*d*e*(10*b*d + 9*a*e) + 8*b*c*e^2*(21*b*d + 13*a*e)) + (Sqrt[c]*(7*A*c*(15*c^2*d^3 - 15*a*c*d*e^2 + 4*a*b*e^3) - a*B*e*(105*c^2*d^2 + 24*b^2*e^2 - c*e*(84*b*d + 25*a*e))))/Sqrt[a])*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^{1/4}*x)/a^{1/4}], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(210*c^{15/4}*Sqrt[a + b*x^2 + c*x^4])$

Rule 1117

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x]] \text{ /; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1209

$\text{Int}[(d_) + (e_)*(x_)^2/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] \text{ /; EqQ}[e + d*q^2, 0] \text{ /; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1211

$\text{Int}[(d_) + (e_)*(x_)^2/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] \text{ /; NeQ}[e + d*q, 0] \text{ /; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1693

$\text{Int}[(Pq_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^{(p_)}, x_Symbol] \text{ :> With}[\{q = \text{Expon}[Pq, x^2], e = \text{Coeff}[Pq, x^2, \text{Expon}[Pq, x^2]]\}, \text{Simp}[e*x^{(2*q - 3)}*((a + b*x^2 + c*x^4)^{(p + 1)}/(c*(2*q + 4*p + 1))), x] + \text{Dist}[1/(c*(2*q + 4*p + 1)), \text{Int}[(a + b*x^2 + c*x^4)^p*\text{ExpandToSum}[c*(2*q + 4*p + 1)*Pq - a*e*(2*q - 3)*x^{(2*q - 4)} - b*e*(2*q + 2*p - 1)*x^{(2*q - 2)} - c*e*(2*q + 4*p + 1)*x^{(2*q)}, x], x], x]] \text{ /; FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{PolyQ}[Pq, x^2] \ \&\& \ \text{Expon}[Pq, x^2] > 1 \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{!LtQ}[p, -1]$

$b^2 - 4ac] * d - 2ae)) * \text{Sqrt}[(b + \text{Sqrt}[b^2 - 4ac] + 2cx^2)/(b + \text{Sqrt}[b^2 - 4ac])] * \text{Sqrt}[(2b - 2\text{Sqrt}[b^2 - 4ac] + 4cx^2)/(b - \text{Sqrt}[b^2 - 4ac])] * \text{EllipticF}[\text{I} * \text{ArcSinh}[\text{Sqrt}[2] * \text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4ac])] * x], (b + \text{Sqrt}[b^2 - 4ac])/(b - \text{Sqrt}[b^2 - 4ac])]/(420c^4 \text{Sqrt}[c/(b + \text{Sqrt}[b^2 - 4ac])]) * \text{Sqrt}[a + bx^2 + cx^4])$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1707 vs. $2(739) = 1478$.

time = 0.13, size = 1708, normalized size = 2.26

method	result
elliptic	$\frac{Be^3x^5\sqrt{cx^4+bx^2+a}}{7c} + \frac{(Ae^3+3Bde^2-\frac{6bBe^3}{7c})x^3\sqrt{cx^4+bx^2+a}}{5c} + \frac{\left(3Ade^2+3Bd^2e-\frac{4b(Ae^3+3Bde^2-\frac{6bBe^3}{7c})}{5c}\right)}{3}$
risch	Expression too large to display
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $B e^3 \frac{1}{7} \frac{1}{c} x^5 (c x^4 + b x^2 + a)^{1/2} - \frac{6}{35} \frac{b}{c^2} x^3 (c x^4 + b x^2 + a)^{1/2} + \frac{1}{3} \frac{(-5/7 a/c + 24/35 b^2/c^2)}{c x} (c x^4 + b x^2 + a)^{1/2} - \frac{1}{12} \frac{(-5/7 a/c + 24/35 b^2/c^2)}{c a^2} (c x^4 + b x^2 + a)^{1/2} / ((-b + (-4ac + b^2)^{1/2})/a)^{1/2} * (4 - 2 * (-b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} * (4 + 2 * (b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} * \text{EllipticF}(1/2 x^2 (1/2) * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2})/a/c)^{1/2}) - 1/2 * (18/35 a * b/c^2 - 2/3 * (-5/7 a/c + 24/35 b^2/c^2)/c * b) * a^2 (1/2) / ((-b + (-4ac + b^2)^{1/2})/a)^{1/2} * (4 - 2 * (-b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} * (4 + 2 * (b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) * (\text{EllipticF}(1/2 x^2 (1/2) * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2})/a/c)^{1/2})) + (A e^3 + 3 B d e^2) * (1/5) \frac{1}{c} x^3 (c x^4 + b x^2 + a)^{1/2} - \frac{4}{15} \frac{b}{c^2} x (c x^4 + b x^2 + a)^{1/2} + \frac{1}{15} \frac{a * b}{c^2} x^2 (1/2) / ((-b + (-4ac + b^2)^{1/2})/a)^{1/2} * (4 - 2 * (-b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} * (4 + 2 * (b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} * \text{EllipticF}(1/2 x^2 (1/2) * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2})/a/c)^{1/2}) - 1/2 * (-3/5 a/c + 8/15 b^2/c^2) * a^2 (1/2) / ((-b + (-4ac + b^2)^{1/2})/a)^{1/2} * (4 - 2 * (-b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} * (4 + 2 * (b + (-4ac + b^2)^{1/2})/a x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) * (\text{EllipticF}(1/2 x^2 (1/2) * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2})/a/c)^{1/2}) - \text{EllipticE}(1/2 x^2 (1/2) * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2})/a/c)^{1/2}))$

$$\begin{aligned}
& -4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}))+(3*A*d*e^2+3*B*d^2*e)*(1/3*x*(c \\
& *x^4+b*x^2+a)^{(1/2)}/c-1/12*a/c^2)^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4 \\
& -2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x^2)^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}+1/3*b/c*a^2)^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(EllipticF(1/2*x^2)^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}-EllipticE(1/2*x^2)^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})) \\
& -1/2*(3*A*d^2*e+B*d^3)*a^2)^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(EllipticF(1/2*x^2)^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}-EllipticE(1/2*x^2)^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+1/4*A*d^3*x^2)^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x^2)^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}))
\end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/sqrt(c*x^4 + b*x^2 + a), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**3/sqrt(a + b*x**2 + c*x**4), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/sqrt(c*x^4 + b*x^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^3}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(1/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(1/2), x)

$\text{Sqrt}[c]*x^2)^2*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]/(30*c^{(11/4)}*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1117

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2))/(2*q*\text{Sqrt}[a + b*x^2 + c*x^4))*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2 - b*(q^2/(4*c))], x]] \text{ ; FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1209

$\text{Int}(((d_) + (e_)*(x_)^2)/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol) \text{ :> With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2))/(q*\text{Sqrt}[a + b*x^2 + c*x^4))*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2 - b*(q^2/(4*c))], x] \text{ ; EqQ}[e + d*q^2, 0] \text{ ; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1211

$\text{Int}(((d_) + (e_)*(x_)^2)/\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol) \text{ :> With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] \text{ ; NeQ}[e + d*q, 0] \text{ ; FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1693

$\text{Int}[(Pq_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^{(p_)}, x_Symbol] \text{ :> With}[\{q = \text{Expon}[Pq, x^2], e = \text{Coeff}[Pq, x^2, \text{Expon}[Pq, x^2]]\}, \text{Simp}[e*x^{(2*q - 3)}*((a + b*x^2 + c*x^4)^{(p + 1)}/(c*(2*q + 4*p + 1))), x] + \text{Dist}[1/(c*(2*q + 4*p + 1)), \text{Int}[(a + b*x^2 + c*x^4)^p*\text{ExpandToSum}[c*(2*q + 4*p + 1)*Pq - a*e*(2*q - 3)*x^{(2*q - 4)} - b*e*(2*q + 2*p - 1)*x^{(2*q - 2)} - c*e*(2*q + 4*p + 1)*x^{(2*q)}, x], x], x]] \text{ ; FreeQ}[\{a, b, c, p\}, x] \ \&\& \ \text{PolyQ}[Pq, x^2] \ \&\& \ \text{Expon}[Pq, x^2] > 1 \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{!LtQ}[p, -1]$

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx &= \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{\int \frac{5Acd^2 + (5Bcd^2 + 10Acde - 3aBe^2)x^2 + e(10Bcd - 4bBe + 5Ace)x^4}{\sqrt{a + bx^2 + cx^4}} dx}{5c} \\
&= \frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{\int \frac{-2}{\sqrt{a + bx^2 + cx^4}} dx}{5c} \\
&= \frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} - \frac{(\sqrt{a + bx^2 + cx^4})^{-2}}{5c} \\
&= \frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{(10A - 2e^2)}{5c} \sqrt{a + bx^2 + cx^4}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 12.74, size = 674, normalized size = 1.28

$$\frac{e(10Bcd - 4bBe + 5Ace)x\sqrt{a + bx^2 + cx^4}}{15c^2} + \frac{Be^2x^3\sqrt{a + bx^2 + cx^4}}{5c} + \frac{(10A - 2e^2)\sqrt{a + bx^2 + cx^4}}{5c}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] (4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4)*(5*A*c*e + B*(10*c*d - 4*b*e + 3*c*e*x^2)) + I*(-b + Sqrt[b^2 - 4*a*c])*(10*A*c*e*(3*c*d - b*e) + B*(15*c^2*d^2 + 8*b^2*e^2 - c*e*(20*b*d + 9*a*e)))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - I*(-8*b^3*B*e^2 + 2*b^2*e*(10*B*c*d + 5*A*c*e + 4*B*Sqrt[b^2 - 4*a*c]*e) - b*c*(15*B*c*d^2 + B*e*(20*Sqrt[b^2 - 4*a*c]*d - 17*a*e) + 10*A*e*(3*c*d + Sqrt[b^2 - 4*a*c]*e)) + c*(B*(-9*a*Sqrt[b^2 - 4*a*c]*e^2 + 5*c*d*(3*Sqrt[b^2 - 4*a*c]*d - 4*a*e) + 10*A*c*(3*c*d^2 + e*(3*Sqrt[b^2 - 4*a*c]*d - a*e))))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])/(60*c^3*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1200 vs. 2(516) = 1032.

time = 0.18, size = 1201, normalized size = 2.27

method	result
elliptic	$\frac{B e^2 x^3 \sqrt{c x^4 + b x^2 + a}}{5c} + \frac{(A e^2 + 2Bde - \frac{4bB e^2}{5c}) x \sqrt{c x^4 + b x^2 + a}}{3c} + \frac{\left(A d^2 - \frac{a(A e^2 + 2Bde - \frac{4bB e^2}{5c})}{3c} \right) \sqrt{2} \sqrt{\dots}}{\dots}$
risch	$\frac{ex(3Be^2x^2c + 5Ace - 4Bbe + 10Bcd) \sqrt{cx^4 + bx^2 + a}}{15c^2} - \frac{(10Abc e^2 - 30c^2 deA + 9Bac e^2 - 8B b^2 e^2 + 20Bbcde - 15B c^2 d^2) a \sqrt{2} \sqrt{\dots}}{\dots}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $B e^2 \frac{1}{5} c x^3 (c x^4 + b x^2 + a)^{1/2} - \frac{4}{15} \frac{b}{c^2} x (c x^4 + b x^2 + a)^{1/2} + \frac{1}{15} \frac{a b}{c^2} \frac{2^{1/2}}{(-b + (-4 a^* c + b^2)^{1/2}) / a^{1/2}} \frac{(4 - 2(-b + (-4 a^* c + b^2)^{1/2})^{1/2}) / a^* x^2)^{1/2}}{(4 + 2(b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2}} \frac{(c x^4 + b x^2 + a)^{1/2} * \text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}) - 1/2 * (-3/5 * a / c + 8/15 * b^2 / c^2) * a^* 2^{1/2}}{((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2}} \frac{(4 + 2(b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2}}{(c x^4 + b x^2 + a)^{1/2}} \frac{(b + (-4 a^* c + b^2)^{1/2}) * (\text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}) - \text{EllipticE}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}))}{(A e^2 + 2 B d e) * (1/3 * x * (c x^4 + b x^2 + a)^{1/2} / c - 1/12 * a / c^2)^{1/2}} \frac{((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2} * (4 + 2(b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2}}{(c x^4 + b x^2 + a)^{1/2} * \text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}) + 1/3 * b / c * a^* 2^{1/2}} \frac{((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2} * (4 + 2(b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2}}{(c x^4 + b x^2 + a)^{1/2}} \frac{(b + (-4 a^* c + b^2)^{1/2}) * (\text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}) - \text{EllipticE}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}))}{-1/2 * (2 * A * d * e + B * d^2) * a^* 2^{1/2}} \frac{((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2} * (4 + 2(b + (-4 a^* c + b^2)^{1/2}) / a^* x^2)^{1/2}}{(c x^4 + b x^2 + a)^{1/2}} \frac{(b + (-4 a^* c + b^2)^{1/2}) * (\text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}) - \text{EllipticE}(1/2 * x^2^{1/2} * ((-b + (-4 a^* c + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4 a^* c + b^2)^{1/2}) / a / c)^{1/2}))}{\dots}$

$$a^{1/2}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{1/2}))/a/c)^{1/2}))+1/4*A*d^2*2^{1/2}/$$

$$((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2}*(4-2*(-b+(-4*a*c+b^2)^{1/2}))/a*x^2)^{1/2}$$

$$*(4+2*(b+(-4*a*c+b^2)^{1/2}))/a*x^2)^{1/2}/(c*x^4+b*x^2+a)^{1/2}*EllipticF(1$$

$$/2*x*2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{1/2}))/a/c)^{1/2})$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^2/sqrt(c*x^4 + b*x^2 + a), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**2/sqrt(a + b*x**2 + c*x**4), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^2/sqrt(c*x^4 + b*x^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(1/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(1/2), x)

$$3.19 \quad \int \frac{(A+Bx^2)(d+ex^2)}{\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=368

$$\frac{Bex\sqrt{a+bx^2+cx^4}}{3c} + \frac{(3Bcd - 2bBe + 3Ace)x\sqrt{a+bx^2+cx^4}}{3c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(3Bcd - 2bBe + 3Ace)(\sqrt{a} + \sqrt{c}x^2)}{3c^{3/2}(\sqrt{a} + \sqrt{c}x^2)}$$

[Out] $\frac{1}{3}B^*e^*x^*(c^*x^4+b^*x^2+a)^{(1/2)}/c+1/3*(3^*A^*c^*e-2^*B^*b^*e+3^*B^*c^*d)^*x^*(c^*x^4+b^*x^2+a)^{(1/2)}/c^{(3/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-1/3*a^{(1/4)}*(3^*A^*c^*e-2^*B^*b^*e+3^*B^*c^*d)^*(\cos(2^*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2^*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2^*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c^*x^4+b^*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(c^*x^4+b^*x^2+a)^{(1/2)}+1/6*a^{(1/4)}*(\cos(2^*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2^*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2^*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(3^*B^*c^*d-2^*b^*B^*e+3^*A^*c^*e+(3^*A^*c^*d-B^*a^*e)^*c^{(1/2)}/a^{(1/2)})*((c^*x^4+b^*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)}))^2)^{(1/2)}/c^{(7/4)}/(c^*x^4+b^*x^2+a)^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 368, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {1693, 1211, 1117, 1209}

$$\frac{\sqrt{a}(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\middle| \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) \left(\frac{\sqrt{c}(3Bcd - 2bBe + 3Ace)}{\sqrt{a}} - 2bBe + 3Bcd\right) \sqrt[4]{a}(\sqrt{a} + \sqrt{c}x^2)\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} E\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\middle| \left(2 - \frac{b}{\sqrt{a}\sqrt{c}}\right)\right) (3Ace - 2bBe + 3Bcd) + \frac{x\sqrt{a+bx^2+cx^4}(3Ace - 2bBe + 3Bcd)}{3c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} + \frac{Bex\sqrt{a+bx^2+cx^4}}{3c}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2))/Sqrt[a + b*x^2 + c*x^4], x]

[Out] $(B^*e^*x^*\text{Sqrt}[a + b^*x^2 + c^*x^4])/(3^*c) + ((3^*B^*c^*d - 2^*b^*B^*e + 3^*A^*c^*e)^*x^*\text{Sqrt}[a + b^*x^2 + c^*x^4])/(3^*c^{(3/2)}*(\text{Sqrt}[a] + \text{Sqrt}[c]^*x^2)) - (a^{(1/4)}*(3^*B^*c^*d - 2^*b^*B^*e + 3^*A^*c^*e)^*(\text{Sqrt}[a] + \text{Sqrt}[c]^*x^2)^*\text{Sqrt}[(a + b^*x^2 + c^*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]^*x^2)^2]*\text{EllipticE}[2^*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]^*\text{Sqrt}[c]))/4])/3^*c^{(7/4)}*\text{Sqrt}[a + b^*x^2 + c^*x^4] + (a^{(1/4)}*(3^*B^*c^*d - 2^*b^*B^*e + 3^*A^*c^*e + (\text{Sqrt}[c]^*(3^*A^*c^*d - a^*B^*e))/\text{Sqrt}[a])^*(\text{Sqrt}[a] + \text{Sqrt}[c]^*x^2)^*\text{Sqrt}[(a + b^*x^2 + c^*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]^*x^2)^2]*\text{EllipticF}[2^*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]^*\text{Sqrt}[c]))/4])/6^*c^{(7/4)}*\text{Sqrt}[a + b^*x^2 + c^*x^4])$

Rule 1117

Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))]

], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1209

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4])*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1211

Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1693

Int[(Pq_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{q = Expon[Pq, x^2], e = Coeff[Pq, x^2, Expon[Pq, x^2]]}, Simp[e*x^(2*q - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(2*q + 4*p + 1))), x] + Dist[1/(c*(2*q + 4*p + 1)), Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*q + 4*p + 1)*Pq - a*e*(2*q - 3)*x^(2*q - 4) - b*e*(2*q + 2*p - 1)*x^(2*q - 2) - c*e*(2*q + 4*p + 1)*x^(2*q), x], x], x] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx &= \frac{Bex\sqrt{a + bx^2 + cx^4}}{3c} + \frac{\int \frac{3Ac d - aBe + (3Bcd - 2bBe + 3Ace)x^2}{\sqrt{a + bx^2 + cx^4}} dx}{3c} \\ &= \frac{Bex\sqrt{a + bx^2 + cx^4}}{3c} - \frac{(\sqrt{a}(3Bcd - 2bBe + 3Ace)) \int \frac{1 - \frac{\sqrt{c}x^2}{\sqrt{a}}}{\sqrt{a + bx^2 + cx^4}} dx}{3c^{3/2}} \\ &= \frac{Bex\sqrt{a + bx^2 + cx^4}}{3c} + \frac{(3Bcd - 2bBe + 3Ace)x\sqrt{a + bx^2 + cx^4}}{3c^{3/2}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a}(3B}{3c^{3/2}} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.
 time = 11.31, size = 521, normalized size = 1.42

$$\frac{4Bc\sqrt{\frac{c}{3+2\sqrt{b^2-4ac}}}\arcsin\left(\frac{bx+cx^2}{\sqrt{a+bx^2+cx^4}}\right)-\left(-b+\sqrt{b^2-4ac}\right)\left(-3Bcd+2Bc-3Ad\right)\sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{3+2\sqrt{b^2-4ac}}}\sqrt{\frac{2b-2\sqrt{b^2-4ac}+4cx^2}{3-2\sqrt{b^2-4ac}}}}{3b^2\sqrt{c+2\sqrt{b^2-4ac}}\sqrt{a+bx^2+cx^4}}E\left(\operatorname{arcsinh}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\right)\right)+\left(-2BBe-\left(4Ad+3B\sqrt{b^2-4ac}d-3aBc+3A\sqrt{b^2-4ac}c\right)+b\left(3Bcd+3Aa+2B\sqrt{b^2-4ac}c\right)\right)\sqrt{\frac{b+\sqrt{b^2-4ac}+2cx^2}{3+2\sqrt{b^2-4ac}}}\sqrt{\frac{2b-2\sqrt{b^2-4ac}+4cx^2}{3-2\sqrt{b^2-4ac}}}}{3+2\sqrt{b^2-4ac}}F\left(\operatorname{arcsinh}\left(\sqrt{2}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\right)\right)\sqrt{\frac{b+\sqrt{b^2-4ac}}{3+2\sqrt{b^2-4ac}}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/Sqrt[a + b*x^2 + c*x^4],x]

[Out] (4*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*e*x*(a + b*x^2 + c*x^4) - I*(-b + Sqrt[b^2 - 4*a*c])*(-3*B*c*d + 2*b*B*e - 3*A*c*e)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + I*(-2*b^2*B*e - c*(6*A*c*d + 3*B*Sqrt[b^2 - 4*a*c]*d - 2*a*B*e + 3*A*Sqrt[b^2 - 4*a*c]*e) + b*(3*B*c*d + 3*A*c*e + 2*B*Sqrt[b^2 - 4*a*c]*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(12*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 758 vs. 2(360) = 720.

time = 0.04, size = 759, normalized size = 2.06

method	result
elliptic	$\frac{Bex\sqrt{cx^4 + bx^2 + a}}{3c} + \frac{(Ad - \frac{aBe}{3c})\sqrt{2}\sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}}\sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}}}{4\sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}\sqrt{c}}$
risch	$\frac{Bex\sqrt{cx^4 + bx^2 + a}}{3c} + \frac{(3Ace - 2Bbe + 3Bcd)a\sqrt{2}\sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}}\sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}}}{4\sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}\sqrt{c}}$

default	Be	$\frac{x\sqrt{cx^4 + bx^2 + a}}{3c} - \frac{a\sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}}}{12c \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}}$	Elliptic
---------	----	---	----------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$B*e*(1/3*x*(c*x^4+b*x^2+a)^{(1/2)}/c-1/12*a/c*2^{(1/2)}/((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2))^{(1/2)})/a/c)^{(1/2)}+1/3*b/c*a*2^{(1/2)}/((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2))^{(1/2)}*(EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2))^{(1/2)})/a/c)^{(1/2)}-EllipticE(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2))^{(1/2)})/a/c)^{(1/2)}))-1/2*(A*e+B*d)*a*2^{(1/2)}/((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2))^{(1/2)}*(EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2))^{(1/2)})/a/c)^{(1/2)}-EllipticE(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2))^{(1/2)})/a/c)^{(1/2)}))+1/4*A*d*2^{(1/2)}/((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2))^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2))^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2))^{(1/2)})/a/c)^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)*(x^2*e + d)/sqrt(c*x^4 + b*x^2 + a), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)
```

Sympy [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{(A + Bx^2)(d + ex^2)}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)
```

```
[Out] Integral((A + B*x**2)*(d + e*x**2)/sqrt(a + b*x**2 + c*x**4), x)
```

Giac [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)*(x^2*e + d)/sqrt(c*x^4 + b*x^2 + a), x)
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(1/2),x)
```

```
[Out] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(1/2), x)
```

$$3.20 \quad \int \frac{A+Bx^2}{\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=283

$$\frac{Bx\sqrt{a+bx^2+cx^4}}{\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)} - \frac{\sqrt[4]{a} B(\sqrt{a}+\sqrt{c}x^2) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right) \middle| \frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{c^{3/4}\sqrt{a+bx^2+cx^4}}$$

[Out] $B*x*(c*x^4+b*x^2+a)^{(1/2)}/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})-a^{(1/4)}*B*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})^{(1/2)}/c^{(3/4)}/(c*x^4+b*x^2+a)^{(1/2)}+1/2*a^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*(B+A*c^{(1/2)}/a^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^{(1/2)}/c^{(3/4)}/(c*x^4+b*x^2+a)^{(1/2)})$

Rubi [A]

time = 0.05, antiderivative size = 283, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {1211, 1117, 1209}

$$\frac{\sqrt[4]{a}(\sqrt{a}+\sqrt{c}x^2)\left(\frac{\sqrt[4]{c}x}{\sqrt{a}}+B\right)\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}F\left(2\text{ArcTan}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{2c^{3/4}\sqrt{a+bx^2+cx^4}} - \frac{\sqrt[4]{a}B(\sqrt{a}+\sqrt{c}x^2)\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a}+\sqrt{c}x^2)^2}}E\left(2\text{ArcTan}\left(\frac{\sqrt[4]{c}x}{\sqrt[4]{a}}\right)\middle|\frac{1}{4}\left(2-\frac{b}{\sqrt{a}\sqrt{c}}\right)\right)}{c^{3/4}\sqrt{a+bx^2+cx^4}} + \frac{Bx\sqrt{a+bx^2+cx^4}}{\sqrt{c}(\sqrt{a}+\sqrt{c}x^2)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/Sqrt[a + b*x^2 + c*x^4], x]

[Out] $(B*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(\text{Sqrt}[c]*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) - (a^{(1/4)}*B*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/ (c^{(3/4)}*\text{Sqrt}[a + b*x^2 + c*x^4]) + (a^{(1/4)}*(B + (A*\text{Sqrt}[c])/ \text{Sqrt}[a])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*c^{(3/4)}*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1117

Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] :> With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1209

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol]
:> With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1211

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol]
:> With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rubi steps

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx = \left(A + \frac{\sqrt{a} B}{\sqrt{c}} \right) \int \frac{1}{\sqrt{a + bx^2 + cx^4}} dx - \frac{(\sqrt{a} B) \int \frac{1 - \frac{\sqrt{c} x^2}{\sqrt{a}}}{\sqrt{a + bx^2 + cx^4}} dx}{\sqrt{c}}$$

$$= \frac{Bx\sqrt{a + bx^2 + cx^4}}{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)} - \frac{\sqrt[4]{a} B(\sqrt{a} + \sqrt{c}x^2) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}} E\left(2 \tan^{-1}\left(\frac{\sqrt[4]{c}}{\sqrt[4]{a}}\right)\right)}{c^{3/4}\sqrt{a + bx^2 + cx^4}}$$

Mathematica [C] Result contains complex when optimal does not.
time = 10.14, size = 302, normalized size = 1.07

$$i \frac{\sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \left(B(-b + \sqrt{b^2 - 4ac}) E\left(i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x\right) \Big|_{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}}\right) + (bB - 2Ac - B\sqrt{b^2 - 4ac}) F\left(i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x\right) \Big|_{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}}\right) \right)}{2\sqrt{2} c \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(A + B*x^2)/Sqrt[a + b*x^2 + c*x^4], x]
```

```
[Out] ((I/2)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(B*(-b + Sqrt[b^2 - 4*a*c])*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (b*B - 2*A*c - B*Sqrt[b^2 - 4*a*c])*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])/(Sqrt[2]*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*Sqrt[a + b*x^2 + c*x^4])
```

Maple [A]

time = 0.02, size = 362, normalized size = 1.28

method	result
default	$Ba\sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} \left(\text{EllipticF} \left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac}}{2a}}}{2\sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}} \right) \right)$
elliptic	$Ba\sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} \left(\text{EllipticF} \left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac}}{2a}}}{2\sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}}} \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2*B*a^2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)/(b+(-4*a*c+b^2)^{(1/2)})*(\text{EllipticF}(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-\text{EllipticE}(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+1/4*A*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)}*\text{EllipticF}(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly
1 arguments (2 given)
```

Sympy [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{A + Bx^2}{\sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x**2+A)/(c*x**4+b*x**2+a)**(1/2),x)
```

```
[Out] Integral((A + B*x**2)/sqrt(a + b*x**2 + c*x**4), x)
```

Giac [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((B*x^2 + A)/sqrt(c*x^4 + b*x^2 + a), x)
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{Bx^2 + A}{\sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(1/2),x)
```

```
[Out] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(1/2), x)
```

$$3.21 \quad \int \frac{A+Bx^2}{(d+ex^2)\sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=436

$$\frac{(Bd - Ae) \tan^{-1} \left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}} \right) (\sqrt{a} B - A\sqrt{c}) (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} F \left(\frac{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}}{2^4 \sqrt{a} \sqrt{c} (\sqrt{c} d - \sqrt{a} e) \sqrt{a + bx^2 + cx^4}} \right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}}$$

[Out] $-1/2*(-A*e+B*d)*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})/d^{(1/2)}/e^{(1/2)}/(a*e^2-b*d*e+c*d^2)^{(1/2)}-1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/4*a^{(3/4)}*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(1/4)}/d/e/(-a*e^2+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A]

time = 0.27, antiderivative size = 436, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {1722, 1117, 1720}

$$\frac{a^{3/4}(\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a}} + c \right)^2 (Bd - Ae) \Pi \left(\frac{\sqrt{cd^2 - bde + ae^2}}{4\sqrt{a} \sqrt{c} de}; 2 \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a}} \right) \middle| 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right) - \frac{(Bd - Ae) \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a} \sqrt{c} \sqrt{a + bx^2 + cx^4}} \right) (\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} F \left(2 \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a}} \right) \middle| 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right)}{4\sqrt{c} de \sqrt{a + bx^2 + cx^4} (cd^2 - ae^2)} - \frac{(Bd - Ae) \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a} \sqrt{c} \sqrt{a + bx^2 + cx^4}} \right) (\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} F \left(2 \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a}} \right) \middle| 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right)}{2^4 \sqrt{a} \sqrt{c} \sqrt{a^2 - bde + cd^2}} - \frac{(Bd - Ae) \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a} \sqrt{c} \sqrt{a + bx^2 + cx^4}} \right) (\sqrt{a} + \sqrt{c} x^2) (\sqrt{a} B - A\sqrt{c}) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} F \left(2 \text{ArcTan} \left(\frac{\sqrt{cd^2 - bde + ae^2}}{\sqrt{a}} \right) \middle| 2 - \frac{b}{\sqrt{a} \sqrt{c}} \right)}{2^4 \sqrt{a} \sqrt{c} \sqrt{a + bx^2 + cx^4} (\sqrt{c} d - \sqrt{a} e)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] $-1/2*((B*d - A*e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])]/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[c*d^2 - b*d*e + a*e^2]) - ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[a + b*x^2 + c*x^4]/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2)*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/2*a^{(1/4)}*c^{(1/4)}*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*\text{Sqrt}[a + b*x^2 + c*x^4] + (a^{(3/4)}*((\text{Sqrt}[c]*d)/\text{Sqrt}[a] + e)^2*(B*d - A*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[a + b*x^2 + c*x^4]/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2)*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/4*c^{(1/4)}*d*e*(c*d^2 - a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1117

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x]] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1720

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-B*d - A*e)*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/(4*a*B))], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1722

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/(d + e*x^2)*Sqrt[a + b*x^2 + c*x^4], x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rubi steps

$$\int \frac{A + Bx^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx = -\frac{(\sqrt{a} B - A\sqrt{c}) \int \frac{1}{\sqrt{a + bx^2 + cx^4}} dx}{\sqrt{c} d - \sqrt{a} e} + \frac{(\sqrt{a} (Bd - Ae)) \int \frac{1}{(d+ex^2)\sqrt{a + bx^2 + cx^4}} dx}{\sqrt{c} d - \sqrt{a} e}$$

$$= -\frac{(Bd - Ae) \tan^{-1} \left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}} \right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} - \frac{(\sqrt{a} B - A\sqrt{c}) (\sqrt{a + bx^2 + cx^4})}{\sqrt{c} d - \sqrt{a} e}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.34, size = 298, normalized size = 0.68

$$\frac{i \sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \left(\text{BdF} \left(i \sinh^{-1} \left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \Big|_{b - \sqrt{b^2 - 4ac}}^{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}} \right) + (-Bd + Ae) \Pi \left(\frac{(b + \sqrt{b^2 - 4ac})^e}{2cd}; i \sinh^{-1} \left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \Big|_{b - \sqrt{b^2 - 4ac}}^{\frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}} \right) \right)}{\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} de \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-I)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(B*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + (-B*d + A*e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])))/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*Sqrt[a + b*x^2 + c*x^4])

Maple [A]

time = 0.16, size = 359, normalized size = 0.82

method	result
default	$B\sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} \operatorname{EllipticF} \left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{2a}}}{4e \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}} \sqrt{cx^4 + bx^2 + a}} \right)$
elliptic	$B\sqrt{2} \sqrt{4 + \frac{2bx^2}{a} - \frac{2x^2\sqrt{-4ac + b^2}}{a}} \sqrt{4 + \frac{2bx^2}{a} + \frac{2x^2\sqrt{-4ac + b^2}}{a}} \operatorname{EllipticF} \left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac}}{2a}}}{4e \sqrt{-\frac{b}{a} + \frac{\sqrt{-4ac + b^2}}{a}} \sqrt{cx^4 + bx^2 + a}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/4*B/e*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2)+(A*e-B*d)/e/d*2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),-2/(-b+(-4*a*c+b^2)^(1/2))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d) \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)

$$3.22 \quad \int \frac{A+Bx^2}{(d+ex^2)^2 \sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=782

$$\frac{\sqrt{c} (Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d (cd^2 - bde + ae^2) (\sqrt{a} + \sqrt{c} x^2)} - \frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d (cd^2 - bde + ae^2) (d + ex^2)} - \frac{(B(cd^3 - ade^2) - Ae(3cd^2 - e(2bd - e^2d)))}{4d^{3/2}\sqrt{e} (cd^2 - bde + ae^2)}$$

```
[Out] -1/4*(B*(-a*d*e^2+c*d^3)-A*e*(3*c*d^2-e*(-a*e+2*b*d)))*arctan(x*(a*e^2-b*d*
e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))/d^(3/2)/(a*e^2-b*d*e+
c*d^2)^(3/2)/e^(1/2)-1/2*e*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^(1/2)/d/(a*e^2-b*d*
e+c*d^2)/(e*x^2+d)+1/2*(-A*e+B*d)*x*c^(1/2)*(c*x^4+b*x^2+a)^(1/2)/d/(a*e^2-
b*d*e+c*d^2)/(a^(1/2)+x^2*c^(1/2))-1/2*a^(1/4)*c^(1/4)*(-A*e+B*d)*(cos(2*ar
ctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*Elliptic
E(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)
)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/d/(a*e^2-b*d
*e+c*d^2)/(c*x^4+b*x^2+a)^(1/2)+1/2*A*c^(1/4)*(cos(2*arctan(c^(1/4)*x/a^(1/
4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/
4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x
^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(1/4)/d/(-e*a^(1/2)+d*c^(1/2))
/(c*x^4+b*x^2+a)^(1/2)+1/8*(B*(-a*d*e^2+c*d^3)-A*e*(3*c*d^2-e*(-a*e+2*b*d))
)*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)
))*EllipticPi(sin(2*arctan(c^(1/4)*x/a^(1/4))),-1/4*(-e*a^(1/2)+d*c^(1/2))^
2/d/e/a^(1/2)/c^(1/2),1/2*(2-b/a^(1/2)/c^(1/2))^(1/2))*(e*a^(1/2)+d*c^(1/2)
)*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(
1/4)/c^(1/4)/d^2/e/(a*e^2-b*d*e+c*d^2)/(-e*a^(1/2)+d*c^(1/2))/(c*x^4+b*x^2+
a)^(1/2)
```

Rubi [A]

time = 0.96, antiderivative size = 782, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {1710, 1728, 1209, 1722, 1117, 1720}

$$\frac{\sqrt{c} (Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d (cd^2 - bde + ae^2) (\sqrt{a} + \sqrt{c} x^2)} - \frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d (cd^2 - bde + ae^2) (d + ex^2)} - \frac{(B(cd^3 - ade^2) - Ae(3cd^2 - e(2bd - e^2d)))}{4d^{3/2}\sqrt{e} (cd^2 - bde + ae^2)}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + b*x^2 + c*x^4]),x]

```
[Out] (Sqrt[c]*(B*d - A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)
*(Sqrt[a] + Sqrt[c]*x^2)) - (e*(B*d - A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(2*d
*(c*d^2 - b*d*e + a*e^2)*(d + e*x^2)) - ((B*(c*d^3 - a*d*e^2) - A*e*(3*c*d^
2 - e*(2*b*d - a*e)))*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[
e]*Sqrt[a + b*x^2 + c*x^4]])/(4*d^(3/2)*Sqrt[e]*(c*d^2 - b*d*e + a*e^2)^(3
```

/2)) - (a^(1/4)*c^(1/4)*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*d*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4]) + (A*c^(1/4)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(1/4)*d*(Sqrt[c]*d - Sqrt[a]*e)*Sqrt[a + b*x^2 + c*x^4]) + ((Sqrt[c]*d + Sqrt[a]*e)*(B*(c*d^3 - a*d*e^2) - A*e*(3*c*d^2 - e*(2*b*d - a*e)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[c]*d - Sqrt[a]*e)^2/(Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(8*a^(1/4)*c^(1/4)*d^2*e*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4])

Rule 1117

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1209

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2])]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1710

```
Int[((P4x_)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + b*x^2 + c*x^4]/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)), Int[((d + e*x^2)^(q + 1)/Sqrt[a + b*x^2 + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*d*(c*d - b*e)*(q + 1)) - 2*((B*d - A*e)*(b*e*(q + 2) - c*d*(q + 1)) - C*d*(b*d + a*e*(q + 1)))*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && ILtQ[q, -1]
```

Rule 1720

```
Int[((A_) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[
```

```
-b + c*(d/e) + a*(e/d), 2]))], x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(
(a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2))]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4])
)*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/
(4*a*B))], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] &&
NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ
[c*A^2 - a*B^2, 0]
```

Rule 1722

```
Int[((A_.) + (B_.)*(x_)^2)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2
+ (c_.)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q)
- a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] +
Dist[a*(B*d - A*e)*(e + d*q)/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2
)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && N
eQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0
] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1728

```
Int[(P4x_)/(((d_) + (e_.)*(x_)^2)*Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x
, 2], C = Coeff[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + b*x^2
+ c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a
*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b,
c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*
d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && !GtQ[b^2 - 4*a*c,
0]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} - \frac{\int \frac{-aBde - A(2cd^2 - e(2bd - ae)) - 2cd(Bd - Ae)x^2}{(d + ex^2)\sqrt{a + bx^2 + cx^4}}}{2d(cd^2 - bde + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} - \frac{\int \frac{-\sqrt{a}c^{3/2}de(Bd - Ae) + ce(-aBde - A(2cd^2 - e(2bd - ae)))}{(d + ex^2)\sqrt{a + bx^2 + cx^4}}}{2d(cd^2 - bde + ae^2)} \\
&= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)} \\
&= \frac{\sqrt{c}(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(\sqrt{a} + \sqrt{c}x^2)} - \frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{2d(cd^2 - bde + ae^2)(d + ex^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.
time = 12.00, size = 1853, normalized size = 2.37

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-1/8*I)*((-4*I)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*d*e^2*(B*d - A*e)*x*(a + b*x^2 + c*x^4) + Sqrt[2]*B*(b - Sqrt[b^2 - 4*a*c])*d^2*e*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*(EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + Sqrt[2]*A*(-b + Sqrt[b^2 - 4*a*c])*d*e^2*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*(EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + 2*Sqrt[2]*B*c*d^3*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - 2*Sqrt[2]*A*c*d^2*e*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])])*(d + e*x^2)*Ellip

```

ticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 -
4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - 2*Sqrt[2]*B*c*d^3*Sqrt[(b + Sqrt[b^2 - 4
*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2
- 4*a*c])]*(d + e*x^2)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*Ar
cSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(
b - Sqrt[b^2 - 4*a*c])] + 6*Sqrt[2]*A*c*d^2*e*Sqrt[(b + Sqrt[b^2 - 4*a*c] +
2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c
])]*(d + e*x^2)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[S
qrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqr
t[b^2 - 4*a*c])] - 4*Sqrt[2]*A*b*d*e^2*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^
2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d
+ e*x^2)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*
Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 -
4*a*c])] + 2*Sqrt[2]*a*B*d*e^2*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b +
Sqrt[b^2 - 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d + e*x^2
)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/
(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c
])] + 2*Sqrt[2]*a*A*e^3*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2
- 4*a*c])]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*(d + e*x^2)*Ellipti
cPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt
[b^2 - 4*a*c])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])]/(Sqr
t[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*(c*d^3 + d*e*(-b*d) + a*e))*(d + e*x^2)*S
qrt[a + b*x^2 + c*x^4])

```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1494 vs. $2(745) = 1490$.

time = 0.15, size = 1495, normalized size = 1.91

method	result	size
default	Expression too large to display	1495
elliptic	Expression too large to display	2301

Verification of antiderivative is not currently implemented for this CAS.

```

[In] int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2), x, method=_RETURNVERBOSE)
[Out] B/e/d*2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*
(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2
)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a
)^(1/2), -2/(-b+(-4*a*c+b^2)^(1/2))*a*e/d, (-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1
/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)+(A*e-B*d)/e*(1/2*e^2*x*(c*x^
4+b*x^2+a)^(1/2)/d/(a*e^2-b*d*e+c*d^2)/(e*x^2+d)-1/8*c/(a*e^2-b*d*e+c*d^2)*
2^(1/2)/(-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^
2)^(1/2))^(1/2)*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4+b*x^2
+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2), 1/2*(-4
+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/4*c*e/d/(a*e^2-b*d*e+c*d^2)*a*2^(

```

$$\begin{aligned} & 1/2)/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/4*c*e/d/(a*e^2-b*d*e+c*d^2)*a^2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticE(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})+1/2/(a*e^2-b*d*e+c*d^2)/d^2*e^2*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, -2/(-b+(-4*a*c+b^2)^{(1/2)})*a*e/d, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)})*a-1/(a*e^2-b*d*e+c*d^2)/d*e*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, -2/(-b+(-4*a*c+b^2)^{(1/2)})*a*e/d, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)})*b+3/2/(a*e^2-b*d*e+c*d^2)*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, -2/(-b+(-4*a*c+b^2)^{(1/2)})*a*e/d, (-1/2*(b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)})*c \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)^2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*x^4 + b*x^2 + a)*(B*x^2 + A)/(c*d^2*x^4 + b*d^2*x^2 + a*d^2 + (c*x^8 + b*x^6 + a*x^4)*e^2 + 2*(c*d*x^6 + b*d*x^4 + a*d*x^2)*e), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)/((d + e*x**2)**2*sqrt(a + b*x**2 + c*x**4)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)^2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d)^2 \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(1/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(1/2)), x)

$$3.23 \quad \int \frac{A+Bx^2}{(d+ex^2)^3 \sqrt{a+bx^2+cx^4}} dx$$

Optimal. Leaf size=1125

$$\frac{\sqrt{c} (3Ae(3cd^2 - e(2bd - ae)) - Bd(5cd^2 - e(2bd + ae))) x \sqrt{a + bx^2 + cx^4}}{8d^2 (cd^2 - bde + ae^2)^2 (\sqrt{a} + \sqrt{c} x^2)} - \frac{e(Bd - Ae)x \sqrt{a + bx^2 + cx^4}}{4d (cd^2 - bde + ae^2) (d + ex^2)}$$

[Out] $-1/16*(B*d*(3*c^2*d^4-10*a*c*d^2*e^2+a*e^3*(-a*e+4*b*d))-A*e*(15*c^2*d^4-2*c*d^2*e*(-3*a*e+10*b*d)+e^2*(3*a^2*e^2-8*a*b*d*e+8*b^2*d^2))$
 $*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/d^{(5/2)}/(a*e^2-b*d*e+c*d^2)^{(5/2)}/e^{(1/2)}-1/4*e*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)/(e*x^2+d)^2+1/8*e*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d))$
 $*x*(c*x^4+b*x^2+a)^{(1/2)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(e*x^2+d)-1/8*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d))$
 $*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(a^{(1/2)}+x^2*c^{(1/2)})+1/8*a^{(1/4)}*c^{(1/4)}*(3*A*e*(3*c*d^2-e*(-a*e+2*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d))$
 $*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))$
 $*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(c*x^4+b*x^2+a)^{(1/2)}+1/32*(B*d*(3*c^2*d^4-10*a*c*d^2*e^2+a*e^3*(-a*e+4*b*d))-A*e*(15*c^2*d^4-2*c*d^2*e*(-3*a*e+10*b*d)+e^2*(3*a^2*e^2-8*a*b*d*e+8*b^2*d^2))$
 $*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))$
 $*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), -1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^3/e/(a*e^2-b*d*e+c*d^2)^2/(e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/8*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))$
 $*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*(a*e*(3*A*e+B*d)+4*A*d*(-b*e+c*d)+d*(-A*e+B*d)*a^{(1/2)}*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/d^2/(a*e^2-b*d*e+c*d^2)/(e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A]

time = 2.27, antiderivative size = 1125, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {1710, 1728, 1209, 1722, 1117, 1720}

Antiderivative was successfully verified.

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + b*x^2 + c*x^4]),x]

[Out]
$$-1/8*(\text{Sqrt}[c]*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e)))*x*\text{Sqrt}[a + b*x^2 + c*x^4]/(d^2*(c*d^2 - b*d*e + a*e^2)^2*(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)) - (e*(B*d - A*e)*x*\text{Sqrt}[a + b*x^2 + c*x^4]/(4*d*(c*d^2 - b*d*e + a*e^2)*(d + e*x^2)^2) + (e*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e)))*x*\text{Sqrt}[a + b*x^2 + c*x^4]/(8*d^2*(c*d^2 - b*d*e + a*e^2)^2*(d + e*x^2)) - ((B*(3*c^2*d^5 - 10*a*c*d^3*e^2 + a*d*e^3*(4*b*d - a*e)) - A*e*(15*c^2*d^4 - 2*c*d^2*e*(10*b*d - 3*a*e) + e^2*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2)))*\text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4]))/(16*d^(5/2)*\text{Sqrt}[e]*(c*d^2 - b*d*e + a*e^2)^(5/2)) + (a^(1/4)*c^(1/4)*(3*A*e*(3*c*d^2 - e*(2*b*d - a*e)) - B*(5*c*d^3 - d*e*(2*b*d + a*e)))*(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(8*d^2*(c*d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4]) + (c^(1/4)*(\text{Sqrt}[a]*\text{Sqrt}[c]*d*(B*d - A*e) + a*e*(B*d + 3*A*e) + 4*A*d*(c*d - b*e))*(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(8*a^(1/4)*d^2*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + ((\text{Sqrt}[c]*d + \text{Sqrt}[a]*e)*(B*(3*c^2*d^5 - 10*a*c*d^3*e^2 + a*d*e^3*(4*b*d - a*e)) - A*e*(15*c^2*d^4 - 2*c*d^2*e*(10*b*d - 3*a*e) + e^2*(8*b^2*d^2 - 8*a*b*d*e + 3*a^2*e^2)))*(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a + \text{Sqrt}[c]*x^2)^2]*\text{EllipticPi}[-1/4*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)^2/(\text{Sqrt}[a]*\text{Sqrt}[c]*d*e), 2*\text{ArcTan}[(c^(1/4)*x)/a^(1/4)], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(32*a^(1/4)*c^(1/4)*d^3*e*(\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(c*d^2 - b*d*e + a*e^2)^2*\text{Sqrt}[a + b*x^2 + c*x^4])$$

Rule 1117

Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1209

Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]

Rule 1710

Int[((P4x)*((d_) + (e_.)*(x_)^2)^(q_))/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = C

```

oeff[P4x, x, 4]}, Simp[(-(C*d^2 - B*d*e + A*e^2))*x*(d + e*x^2)^(q + 1)*(Sqrt[a + b*x^2 + c*x^4]/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/(2*d*(q + 1)*(c*d^2 - b*d*e + a*e^2)), Int[((d + e*x^2)^(q + 1)/Sqrt[a + b*x^2 + c*x^4])*Simp[a*d*(C*d - B*e) + A*(a*e^2*(2*q + 3) + 2*d*(c*d - b*e)*(q + 1)) - 2*((B*d - A*e)*(b*e*(q + 2) - c*d*(q + 1)) - C*d*(b*d + a*e*(q + 1)))*x^2 + c*(C*d^2 - B*d*e + A*e^2)*(2*q + 5)*x^4, x], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2] && LeQ[Expon[P4x, x], 4] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && ILtQ[q, -1]

```

Rule 1720

```

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/(4*a*B))], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

```

Rule 1722

```

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q) - a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] + Dist[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/(d + e*x^2)*Sqrt[a + b*x^2 + c*x^4], x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]

```

Rule 1728

```

Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x, 2], C = Coeff[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && !GtQ[b^2 - 4*a*c, 0]

```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx &= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{4d(cd^2 - bde + ae^2)(d + ex^2)^2} - \frac{\int \frac{-4Acd^2 - aBde + Ae(4bd - 3ae) - 2(Bd - Ae)}{(d + ex^2)^2 \sqrt{a + bx^2 + cx^4}}}{4d(cd^2 - bde + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{4d(cd^2 - bde + ae^2)(d + ex^2)^2} + \frac{e(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))}{8d^2(cd^2 - bde + ae^2)} \\
&= -\frac{e(Bd - Ae)x\sqrt{a + bx^2 + cx^4}}{4d(cd^2 - bde + ae^2)(d + ex^2)^2} + \frac{e(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))}{8d^2(cd^2 - bde + ae^2)} \\
&= -\frac{\sqrt{c}(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))x\sqrt{a + bx^2}}{8d^2(cd^2 - bde + ae^2)^2(\sqrt{a} + \sqrt{c}x^2)} \\
&= -\frac{\sqrt{c}(3Ae(3cd^2 - e(2bd - ae)) - B(5cd^3 - de(2bd + ae)))x\sqrt{a + bx^2}}{8d^2(cd^2 - bde + ae^2)^2(\sqrt{a} + \sqrt{c}x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.
time = 13.31, size = 781, normalized size = 0.69

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)^3*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-4*d*e^2*x*(a + b*x^2 + c*x^4)*(2*d*(B*d - A*e)*(c*d^2 + e*(-(b*d) + a*e)) + (-3*A*e*(3*c*d^2 + e*(-2*b*d + a*e)) + B*(5*c*d^3 - d*e*(2*b*d + a*e)))*(d + e*x^2))/(d + e*x^2)^2 - (I*Sqrt[2]*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*((-b + Sqrt[b^2 - 4*a*c])*d*e*(3*A*e*(3*c*d^2 + e*(-2*b*d + a*e)) + B*(-5*c*d^3 + d*e*(2*b*d + a*e)))*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + d*(B*d*(6*c^2*d^3 + c*d*e*(-5*b*d + 5*Sqrt[b^2 - 4*a*c]*d - 6*a*e) - (-b + Sqrt[b^2 - 4*a*c])*e^2*(2*b*d + a*e)) - A*e*(14*c^2*d^3 - 3*(-b + Sqrt[b^2 - 4*a*c])*e^2*(2*b*d - a*e) + c*d*e*(-17*b*d + 9*Sqrt[b^2 - 4*a*c]*d + 2*a*e)))*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + 2*(B*(-3*c^2*d^5 + 10*a*c*d^3*e^2 + a*d*e^3*(-4*b*d + a*e)) + A*e*(15*c^2*d^4 + 2*c*d^2*e*(-10*b*d + 3*a*e) + e^2

$$\frac{(8b^2d^2 - 8abde + 3a^2e^2) \operatorname{EllipticPi}\left[\frac{(b + \sqrt{b^2 - 4ac})e}{2cd}, I \operatorname{ArcSinh}\left[\frac{\sqrt{2}\sqrt{c/(b + \sqrt{b^2 - 4ac})}}{x}\right], \frac{(b + \sqrt{b^2 - 4ac})}{(b - \sqrt{b^2 - 4ac})}\right]}{\sqrt{c/(b + \sqrt{b^2 - 4ac})}} / (32d^3e(c d^2 + e(-bd) + ae))^2 \sqrt{a + bx^2 + cx^4}$$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 4475 vs. $2(1080) = 2160$.

time = 0.15, size = 4476, normalized size = 3.98

method	result	size
default	Expression too large to display	4476
elliptic	Expression too large to display	5671

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{(Ae - Bd) \sqrt{1/4e^2/d/(ae^2 - bde + cd^2)} x (cx^4 + bx^2 + a)^{1/2} / (e^2x^2 + d)^2 + 3/8e^2(ae^2 - 2bde + 3cd^2) / (ae^2 - bde + cd^2)^2 / d^2 x (cx^4 + bx^2 + a)^{1/2} / (e^2x^2 + d) - 3/8e^2c / (ae^2 - bde + cd^2)^2 / da^{1/2} / (-b/a + 1/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a - 2x^2/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a + 2x^2/a (-4ac + b^2)^{1/2})^{1/2} / (cx^4 + bx^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) \operatorname{EllipticF}(1/2, x^2)^{1/2} * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2b(b + (-4ac + b^2)^{1/2})/a/c)^{1/2} * b + 3/8e^2c / (ae^2 - bde + cd^2)^2 / da^{1/2} / (-b/a + 1/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a - 2x^2/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a + 2x^2/a (-4ac + b^2)^{1/2})^{1/2} / (cx^4 + bx^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) \operatorname{EllipticE}(1/2, x^2)^{1/2} * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2b(b + (-4ac + b^2)^{1/2})/a/c)^{1/2} * b - 1/32c / (ae^2 - bde + cd^2)^2 / d^2 (1/2) / (-b/a + 1/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a - 2x^2/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a + 2x^2/a (-4ac + b^2)^{1/2})^{1/2} / (cx^4 + bx^2 + a)^{1/2} \operatorname{EllipticF}(1/2, x^2)^{1/2} * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2b(b + (-4ac + b^2)^{1/2})/a/c)^{1/2} * ae^2 + 3/8 / (ae^2 - bde + cd^2)^2 / d^3 e^4 x^2 (1/2) / (-b/a + 1/a (-4ac + b^2)^{1/2})^{1/2} (1 + 1/2bx^2/a - 1/2x^2/a (-4ac + b^2)^{1/2})^{1/2} (1 + 1/2bx^2/a + 1/2x^2/a (-4ac + b^2)^{1/2})^{1/2} / (cx^4 + bx^2 + a)^{1/2} \operatorname{EllipticPi}(1/2, x^2)^{1/2} * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, -2 / (-b + (-4ac + b^2)^{1/2}) * ae/d, (-1/2 * (b + (-4ac + b^2)^{1/2})/a)^{1/2} * 2^{1/2} / ((-b + (-4ac + b^2)^{1/2})/a)^{1/2} * a^2 + 1 / (ae^2 - bde + cd^2)^2 / d^2 e^2 x^2 (1/2) / (-b/a + 1/a (-4ac + b^2)^{1/2})^{1/2} (1 + 1/2bx^2/a - 1/2x^2/a (-4ac + b^2)^{1/2})^{1/2} (1 + 1/2bx^2/a + 1/2x^2/a (-4ac + b^2)^{1/2})^{1/2} / (cx^4 + bx^2 + a)^{1/2} \operatorname{EllipticPi}(1/2, x^2)^{1/2} * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, -2 / (-b + (-4ac + b^2)^{1/2}) * ae/d, (-1/2 * (b + (-4ac + b^2)^{1/2})/a)^{1/2} * 2^{1/2} / ((-b + (-4ac + b^2)^{1/2})/a)^{1/2} * b^2 + 9/16e^2c^2 / (ae^2 - bde + cd^2)^2 * a^2 (1/2) / (-b/a + 1/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a - 2x^2/a (-4ac + b^2)^{1/2})^{1/2} (4 + 2bx^2/a + 2x^2/a (-4ac + b^2)^{1/2})^{1/2} / (cx^4 + bx^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) \operatorname{EllipticF}(1/2, x^2)^{1/2} * ((-b + (-4ac + b^2)^{1/2})/a)^{1/2}, 1/2 * (-4 + 2b(b + (-4ac + b^2)^{1/2})/a/c)^{1/2} * b$$

$$\begin{aligned}
& *c+b^2)^{1/2})/a/c)^{1/2})-3/16*e^3*c/(a*e^2-b*d*e+c*d^2)^2/d^2*a^2*x^{1/2}) \\
& /(-b/a+1/a*(-4*a*c+b^2)^{1/2})^{1/2}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{1/2}) \\
&)^{1/2}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}/(c*x^4+b*x^2+a)^{1/2} \\
& /b+(-4*a*c+b^2)^{1/2})*EllipticE(1/2*x^2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/ \\
& a)^{1/2},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{1/2}))/a/c)^{1/2})-9/16*e*c^2/(a*e^2-b \\
& *d*e+c*d^2)^2*a^2^{1/2}/(-b/a+1/a*(-4*a*c+b^2)^{1/2})^{1/2}*(4+2*b*x^2/a-2* \\
& x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2} \\
& /c*x^4+b*x^2+a)^{1/2}/b+(-4*a*c+b^2)^{1/2})*EllipticE(1/2*x^2^{1/2}*((\\
& -b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{1/2}))/a/c)^{1/2} \\
&)-5/2/(a*e^2-b*d*e+c*d^2)^2*e^2^{1/2}/(-b/a+1/a*(-4*a*c+b^2)^{1/2})^{1/2} \\
& *(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}*(1+1/2*b*x^2/a+1/2*x^2/ \\
& a*(-4*a*c+b^2)^{1/2})^{1/2}/c*x^4+b*x^2+a)^{1/2}*EllipticPi(1/2*x^2^{1/2}* \\
& ((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},-2/(-b+(-4*a*c+b^2)^{1/2}))*a*e/d,(-1/2*(b \\
& +(-4*a*c+b^2)^{1/2}))/a)^{1/2}*2^{1/2}/((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2})*b* \\
& c+3/16*e^3*c/(a*e^2-b*d*e+c*d^2)^2/d^2*a^2*x^{1/2})/(-b/a+1/a*(-4*a*c+b^2)^{1/2} \\
&)^{1/2}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}*(4+2*b*x^2/a+2*x \\
& ^2/a*(-4*a*c+b^2)^{1/2})^{1/2}/c*x^4+b*x^2+a)^{1/2}/b+(-4*a*c+b^2)^{1/2} \\
&)*EllipticF(1/2*x^2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},1/2*(-4+2*b*(b+ \\
& -4*a*c+b^2)^{1/2}))/a/c)^{1/2})+15/8/(a*e^2-b*d*e+c*d^2)^2*d^2^{1/2}/(-b/a+1 \\
& /a*(-4*a*c+b^2)^{1/2})^{1/2}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2} \\
& *(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}/c*x^4+b*x^2+a)^{1/2} \\
&)*EllipticPi(1/2*x^2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},-2/(-b+(-4*a* \\
& c+b^2)^{1/2}))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^{1/2}))/a)^{1/2}*2^{1/2}/((-b+(-4* \\
& a*c+b^2)^{1/2}))/a)^{1/2})*c^2+1/8*c/(a*e^2-b*d*e+c*d^2)^2*2^{1/2}/(-b/a+1/a \\
& *(-4*a*c+b^2)^{1/2})^{1/2}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}*(\\
& 4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}/c*x^4+b*x^2+a)^{1/2}*Ellipti \\
& cF(1/2*x^2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},1/2*(-4+2*b*(b+(-4*a*c+b \\
& ^2)^{1/2}))/a/c)^{1/2})*e*b-1/(a*e^2-b*d*e+c*d^2)^2/d^2*e^3*2^{1/2}/(-b/a+1/ \\
& a*(-4*a*c+b^2)^{1/2})^{1/2}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2} \\
& *(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}/c*x^4+b*x^2+a)^{1/2} \\
&)*EllipticPi(1/2*x^2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},-2/(-b+(-4*a*c \\
& +b^2)^{1/2}))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^{1/2}))/a)^{1/2}*2^{1/2}/((-b+(-4*a \\
& *c+b^2)^{1/2}))/a)^{1/2})*a*b+3/4/(a*e^2-b*d*e+c*d^2)^2/d*e^2*2^{1/2}/(-b/a+ \\
& 1/a*(-4*a*c+b^2)^{1/2})^{1/2}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2} \\
& *(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{1/2})^{1/2}/c*x^4+b*x^2+a)^{1/2} \\
&)*EllipticPi(1/2*x^2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},-2/(-b+(-4*a \\
& *c+b^2)^{1/2}))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^{1/2}))/a)^{1/2}*2^{1/2}/((-b+(-4 \\
& *a*c+b^2)^{1/2}))/a)^{1/2})*a*c-7/32*c^2/(a*e^2-...
\end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)^3), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(d + ex^2)^3 \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**3/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((A + B*x**2)/((d + e*x**2)**3*sqrt(a + b*x**2 + c*x**4)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^3/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)^3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{Bx^2 + A}{(ex^2 + d)^3 \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)^3*(a + b*x^2 + c*x^4)^(1/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)^3*(a + b*x^2 + c*x^4)^(1/2)), x)

$$3.24 \quad \int \frac{(A+Bx^2)(d+ex^2)^3}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=859

$$\frac{x(Ac(b^2cd^3 - 2acd(cd^2 - 3ae^2) - abe(3cd^2 + ae^2)) + aB(ab^2e^3 + 2ace(3cd^2 - ae^2) - bcd(cd^2 + 3ae^2)) - (a^2c^2(b^2 - 4ac)\sqrt{a +$$

```
[Out] x*(A*c*(b^2*c*d^3-2*a*c*d*(-3*a*e^2+c*d^2)-a*b*e*(a*e^2+3*c*d^2))+a*B*(a*b^2*e^3+2*a*c*e*(-a*e^2+3*c*d^2)-b*c*d*(3*a*e^2+c*d^2))-(a*B*(-b*e+2*c*d)*(c^2*d^2+b^2*e^2-c*e*(3*a*e+b*d))+A*c*(a*b^2*e^3+2*a*c*e*(-a*e^2+3*c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*x^2)/a/c^2/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)+1/3*B*e^3*x*(c*x^4+b*x^2+a)^(1/2)/c^2+1/3*(a*B*(6*c^3*d^3-8*b^3*e^3-9*c^2*d*e*(6*a*e+b*d)+b*c*e^2*(29*a*e+18*b*d))+3*A*c*(2*a*b^2*e^3+6*a*c*e*(-a*e^2+c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*x*(c*x^4+b*x^2+a)^(1/2)/a/c^(5/2)/(-4*a*c+b^2)/(a^(1/2)+x^2*c^(1/2))-1/3*(a*B*(6*c^3*d^3-8*b^3*e^3-9*c^2*d*e*(6*a*e+b*d)+b*c*e^2*(29*a*e+18*b*d))+3*A*c*(2*a*b^2*e^3+6*a*c*e*(-a*e^2+c*d^2)-b*c*d*(3*a*e^2+c*d^2)))*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticE(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2)))^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(11/4)/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^(1/2)-1/6*(cos(2*arctan(c^(1/4)*x/a^(1/4)))^2)^(1/2)/cos(2*arctan(c^(1/4)*x/a^(1/4)))*EllipticF(sin(2*arctan(c^(1/4)*x/a^(1/4))),1/2*(2-b/a^(1/2)/c^(1/2)))^(1/2))*(3*A*c^3*d^3-5*a^2*B*c*e^3+a*e*(-2*b*e+3*c*d)*(3*A*c*e-4*B*b*e+3*B*c*d)-3*c^(5/2)*d^2*(3*A*e+B*d)*a^(1/2)+3*a^(3/2)*e^2*(3*A*c*e-4*B*b*e+9*B*c*d)*c^(1/2))*(a^(1/2)+x^2*c^(1/2))*((c*x^4+b*x^2+a)/(a^(1/2)+x^2*c^(1/2)))^(1/2)/a^(3/4)/c^(11/4)/(b-2*a^(1/2)*c^(1/2))/(c*x^4+b*x^2+a)^(1/2)
```

Rubi [A]

time = 0.89, antiderivative size = 859, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.152$, Rules used = {1692, 1693, 1211, 1117, 1209}

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x]

```
[Out] (x*(A*c*(b^2*c*d^3 - 2*a*c*d*(c*d^2 - 3*a*e^2) - a*b*e*(3*c*d^2 + a*e^2)) + a*B*(a*b^2*e^3 + 2*a*c*e*(3*c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)) - (a*B*(2*c*d - b*e)*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e)) + A*c*(a*b^2*e^3 + 2*a*c*e*(3*c*d^2 - a*e^2) - b*c*d*(c*d^2 + 3*a*e^2)))*x^2)/(a*c^2*(b^2 - 4*a*c)*Sqrt[a + b*x^2 + c*x^4]) + (B*e^3*x*Sqrt[a + b*x^2 + c*x^4])/(3*c^2)
```

$$\begin{aligned}
& + ((a*B*(6*c^3*d^3 - 8*b^3*e^3 - 9*c^2*d*e*(b*d + 6*a*e) + b*c*e^2*(18*b*d \\
& + 29*a*e)) + 3*A*c*(2*a*b^2*e^3 + 6*a*c*e*(c*d^2 - a*e^2) - b*c*d*(c*d^2 + \\
& 3*a*e^2)))*x*\text{Sqrt}[a + b*x^2 + c*x^4]/(3*a*c^{(5/2)}*(b^2 - 4*a*c)*(\text{Sqrt}[a \\
& + \text{Sqrt}[c]*x^2)) - ((a*B*(6*c^3*d^3 - 8*b^3*e^3 - 9*c^2*d*e*(b*d + 6*a*e) + \\
& b*c*e^2*(18*b*d + 29*a*e)) + 3*A*c*(2*a*b^2*e^3 + 6*a*c*e*(c*d^2 - a*e^2) - \\
& b*c*d*(c*d^2 + 3*a*e^2)))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4) \\
& /(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b \\
& /(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]/(3*a^{(3/4)}*c^{(11/4)}*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + \\
& c*x^4]) - ((3*A*c^3*d^3 - 5*a^2*B*c*e^3 - 3*\text{Sqrt}[a]*c^{(5/2)}*d^2*(B*d + 3*A* \\
& e) + a*e*(3*c*d - 2*b*e))*(3*B*c*d - 4*b*B*e + 3*A*c*e) + 3*a^{(3/2)}*\text{Sqrt}[c]* \\
& e^2*(9*B*c*d - 4*b*B*e + 3*A*c*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 \\
& + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}] \\
& , (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4]/(6*a^{(3/4)}*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*c^{(11/4)} \\
& *\text{Sqrt}[a + b*x^2 + c*x^4])
\end{aligned}$$

Rule 1117

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1209

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1211

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1692

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 2]}, Simp[x*(a + b*x^2 + c*x^4)^(p + 1)*((a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Pq, a
```

```

+ b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p
+ 7)*(b*d - 2*a*e)*x^2, x], x], x]] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x^
2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]

```

Rule 1693

```

Int[(Pq_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := With[{q =
Expon[Pq, x^2], e = Coeff[Pq, x^2, Expon[Pq, x^2]]}, Simp[e*x^(2*q - 3)*((
a + b*x^2 + c*x^4)^(p + 1)/(c*(2*q + 4*p + 1))), x] + Dist[1/(c*(2*q + 4*p
+ 1)), Int[(a + b*x^2 + c*x^4)^p*ExpandToSum[c*(2*q + 4*p + 1)*Pq - a*e*(2*
q - 3)*x^(2*q - 4) - b*e*(2*q + 2*p - 1)*x^(2*q - 2) - c*e*(2*q + 4*p + 1)*
x^(2*q), x], x], x]] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x^2] && Expon[P
q, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && !LtQ[p, -1]

```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{3/2}} dx &= \frac{x(Ac(b^2cd^3 - 2acd(cd^2 - 3ae^2) - abe(3cd^2 + ae^2)) + aB(ab^2e^3 + 2ace(3cd^2 \\
&= \frac{x(Ac(b^2cd^3 - 2acd(cd^2 - 3ae^2) - abe(3cd^2 + ae^2)) + aB(ab^2e^3 + 2ace(3cd^2 \\
&= \frac{x(Ac(b^2cd^3 - 2acd(cd^2 - 3ae^2) - abe(3cd^2 + ae^2)) + aB(ab^2e^3 + 2ace(3cd^2 \\
&= \frac{x(Ac(b^2cd^3 - 2acd(cd^2 - 3ae^2) - abe(3cd^2 + ae^2)) + aB(ab^2e^3 + 2ace(3cd^2
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.
time = 14.74, size = 1058, normalized size = 1.23

Antiderivative was successfully verified.

```
[In] Integrate[((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x]
```

```
[Out] (-4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])*x*(3*A*c*(b^2*(c*d^3 - a*e^3*x^2) + b
*(-a^2*e^3) + c^2*d^3*x^2 - 3*a*c*d*e*(d - e*x^2)) + 2*a*c*(a*e^2*(3*d + e
```

```

*x^2) - c*d^2*(d + 3*e*x^2))) + a*B*(4*b^3*e^3*x^2 + b^2*e^2*(4*a*e - 9*c*d
*x^2 + c*e*x^4) - b*c*(3*c*d^2*(d - 3*e*x^2) + a*e^2*(9*d + 13*e*x^2)) - 2*
c*(5*a^2*e^3 + 3*c^2*d^3*x^2 + a*c*e*(-9*d^2 - 9*d*e*x^2 + 2*e^2*x^4))) +
I*(-b + Sqrt[b^2 - 4*a*c])*(a*B*(-6*c^3*d^3 + 8*b^3*e^3 + 9*c^2*d*e*(b*d +
6*a*e) - b*c*e^2*(18*b*d + 29*a*e)) + 3*A*c*(-2*a*b^2*e^3 + 6*a*c*e*(-(c*d^
2) + a*e^2) + b*c*d*(c*d^2 + 3*a*e^2)))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x
^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b
- Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*
a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) - I*(a*B*(8*b^3
*(-b + Sqrt[b^2 - 4*a*c])*e^3 - 6*c^3*d^2*(Sqrt[b^2 - 4*a*c]*d - 6*a*e) + b
*c*e^2*(18*b^2*d - 18*b*Sqrt[b^2 - 4*a*c]*d + 37*a*b*e - 29*a*Sqrt[b^2 - 4*
a*c]*e) + c^2*e*(-9*b^2*d^2 + 2*a*e*(27*Sqrt[b^2 - 4*a*c]*d - 10*a*e) + 9*b
*d*(Sqrt[b^2 - 4*a*c]*d - 8*a*e))) + 3*A*c*(2*a*b^3*e^3 - b^2*(c^2*d^3 + 3*
a*c*d*e^2 + 2*a*Sqrt[b^2 - 4*a*c]*e^3) + b*c*(c*Sqrt[b^2 - 4*a*c]*d^3 + a*e
^2*(3*Sqrt[b^2 - 4*a*c]*d - 8*a*e)) + 2*a*c*(2*c^2*d^3 + 3*a*Sqrt[b^2 - 4*a
*c]*e^3 - 3*c*d*e*(Sqrt[b^2 - 4*a*c]*d - 2*a*e)))*Sqrt[(b + Sqrt[b^2 - 4*a
*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4
*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sq
rt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))]/(12
*a*c^3*(-b^2 + 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^
4])

```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 2444 vs. 2(841) = 1682.

time = 0.16, size = 2445, normalized size = 2.85

method	result
elliptic	$2c \left(\frac{(2Aa^2c^2e^3 - Aab^2ce^3 + 3Aabc^2de^2 - 6Aac^3d^2e + Abc^3d^3 - 3Ba^2bce^3 + 6Ba^2c^2de^2 + Bab^3e^3 - 3Bab^2cde^2 + 3Babc^2d^2e - 2Ba^3d^3)x^3}{2c^3a(4ac - b^2)} \right) \sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right)}$
default	Expression too large to display
risch	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2), x, method=_RETURNVERBOSE)

[Out] B*e^3*(-2*c*(-1/2*b/c^3*(3*a*c-b^2)/(4*a*c-b^2))*x^3-1/2*(2*a*c-b^2)*a/c^3/(4*a*c-b^2)*x)/((x^4+b/c*x^2+a/c)*c)^(1/2)+1/3/c^2*x*(c*x^4+b*x^2+a)^(1/2)+1/4*(-a*(2*a*c-b^2)/(4*a*c-b^2)/c^2-1/3*a/c^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x^2^(1/2)*((-b+(-4

$$\begin{aligned}
& *a*c+b^2)^{(1/2)}/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}-1/ \\
& 2*(-5/3*b/c^2-b*(3*a*c-b^2)/c^2/(4*a*c-b^2))*a^2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(\text{Elliptic} \\
& \text{cF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-\text{EllipticE}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+(A*e^3+3*B*d*e^2)*(- \\
& 2*c*(1/2/c^2*(2*a*c-b^2)/(4*a*c-b^2)*x^3-1/2*a*b/c^2/(4*a*c-b^2)*x)/((x^4+b \\
& /c*x^2+a/c)*c)^{(1/2)}-1/4*a*b/c/(4*a*c-b^2)*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*\text{EllipticF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/2*(\\
& 1/c+(2*a*c-b^2)/c/(4*a*c-b^2))*a^2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(\text{EllipticF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-\text{EllipticE}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+ \\
& 2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+(3*A*d*e^2+3*B*d^2*e)*(-2*c*(1/2*b \\
& /c/(4*a*c-b^2)*x^3+a/c/(4*a*c-b^2)*x)/((x^4+b/c*x^2+a/c)*c)^{(1/2)}+1/2*a/(4* \\
& a*c-b^2)*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*\text{EllipticF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b \\
& *(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}-1/2*b/(4*a*c-b^2)*a^2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)}) \\
& *(\text{EllipticF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+ \\
& (-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-\text{EllipticE}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+(3*A*d^2*e+ \\
& B*d^3)*(-2*c*(-1/(4*a*c-b^2)*x^3-1/2*b/(4*a*c-b^2)/c*x)/((x^4+b/c*x^2+a/c)* \\
& c)^{(1/2)}-1/4*b/(4*a*c-b^2)*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*\text{EllipticF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})+c/(4*a*c-b^2)*a^2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a \\
& *c+b^2)^{(1/2))*(\text{EllipticF}(1/2*x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1 \\
& /2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-\text{EllipticE}(1/2*x^2^{(1/2)}*((-b+ \\
& (-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)} \\
&))+A*d^3*(-2*c*(1/2*a*b/(4*a*c-b^2)*x^3-1/2*(2*a*c-b^2)/a/(4*a*c-b^2)/c*x)/ \\
& ((x^4+b/c*x^2+a/c)*c)^{(1/2)}+1/4*(1/a-(2*a*c-b^2)/a/(4*a*c-b^2))*2^{(1/2)/((-b+ \\
& (-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4 \\
& +2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*\text{EllipticF}(1/2* \\
& x^2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}, 1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/2*b*c/(4*a*c-b^2)*2^{(1/2)/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*
\end{aligned}$$

$$x^2)^{1/2}/(c*x^4+b*x^2+a)^{1/2}/(b+(-4*a*c+b^2)^{1/2})*(EllipticF(1/2*x*2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{1/2}))/a/c)^{1/2})-EllipticE(1/2*x*2^{1/2}*((-b+(-4*a*c+b^2)^{1/2}))/a)^{1/2},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{1/2}))/a/c)^{1/2}))$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/(c*x^4 + b*x^2 + a)^(3/2), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^3}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**3/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**3/(a + b*x**2 + c*x**4)**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^3/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^3/(c*x^4 + b*x^2 + a)^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(B x^2 + A) (e x^2 + d)^3}{(c x^4 + b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^3)/(a + b*x^2 + c*x^4)^(3/2), x)

$$a + b*x^2 + c*x^4) - ((A*c^2*d^2 + 3*a^{(3/2)}*B*Sqrt[c]*e^2 - Sqrt[a]*c^{(3/2)}*d*(B*d + 2*A*e) + a*e*(2*B*c*d - 2*b*B*e + A*c*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(Sqrt[a]*Sqrt[c]))/4]/(2*a^{(3/4)}*(b - 2*Sqrt[a]*Sqrt[c])*c^{(7/4)}*Sqrt[a + b*x^2 + c*x^4])$$

Rule 1117

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1209

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1211

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1692

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 2]}, Simp[x*(a + b*x^2 + c*x^4)^(p + 1)*((a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)/(2*a*(p + 1)*(b^2 - 4*a*c))), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Pq, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{3/2}} dx &= -\frac{x \left(c \left(\frac{aB(bcd^2 - 4acde + abe^2)}{c} - A(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) \right) - (Ac(bcd^2 - 4acde + abe^2)) \right)}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x \left(c \left(\frac{aB(bcd^2 - 4acde + abe^2)}{c} - A(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) \right) - (Ac(bcd^2 - 4acde + abe^2)) \right)}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x \left(c \left(\frac{aB(bcd^2 - 4acde + abe^2)}{c} - A(b^2d^2 - 2abde - 2a(cd^2 - ae^2)) \right) - (Ac(bcd^2 - 4acde + abe^2)) \right)}{ac(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 12.98, size = 766, normalized size = 1.22

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] (-4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])] * x * (- (a*B*(a*b*e^2 + 2*c^2*d^2*x^2 + b^2*e^2*x^2 + b*c*d*(d - 2*e*x^2) - 2*a*c*e*(2*d + e*x^2))) + A*c*(b^2*d^2 + 2*a^2*e^2 + b*c*d^2*x^2 + a*b*e*(-2*d + e*x^2) - 2*a*c*d*(d + 2*e*x^2))) - I*(-b + Sqrt[b^2 - 4*a*c]) * (- (A*c*(b*c*d^2 - 4*a*c*d*e + a*b*e^2)) + 2*a*B*(c^2*d^2 + b^2*e^2 - c*e*(b*d + 3*a*e))) * Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])] * EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]] * x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + I*(2*a*B*(b^2*(-b + Sqrt[b^2 - 4*a*c])*e^2 + c^2*d*(Sqrt[b^2 - 4*a*c]*d - 4*a*e) + c*e*(b^2*d - b*Sqrt[b^2 - 4*a*c]*d + 4*a*b*e - 3*a*Sqrt[b^2 - 4*a*c]*e)) + A*c*(b^2*(c*d^2 + a*e^2) - b*Sqrt[b^2 - 4*a*c]*(c*d^2 + a*e^2) - 4*a*c*(c*d^2 - Sqrt[b^2 - 4*a*c]*d*e + a*e^2))) * Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])] * EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]] * x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]/(4*a*c^2*(-b^2 + 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])] * Sqrt[a + b*x^2 + c*x^4])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1890 vs. 2(618) = 1236.

time = 0.16, size = 1891, normalized size = 3.01

method	result
elliptic	$-\frac{2c \left(\frac{Aabc e^2 - 4de c^2 aA + Ab c^2 d^2 + 2B a^2 c e^2 - Ba b^2 e^2 + 2Babcde - 2Ba c^2 d^2}{2c^2 a(4ac - b^2)} x^3 + \frac{(2A a^2 c e^2 - 2Aabcde - 2Aa c^2 d^2 + A b^2 c d^2 - B a^2 b e^2 + 4de a^2}{2c^2 (4ac - b^2) a} \right)}{\sqrt{\left(x^4 + \frac{bx^2}{c} + \frac{a}{c}\right) c}}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & B e^2 (-2c (1/2/c^2 (2ac - b^2) / (4ac - b^2)) x^3 - 1/2 ab/c^2 / (4ac - b^2) * x) \\ & / ((x^4 + b/c x^2 + a/c) * c)^{1/2} - 1/4 ab/c / (4ac - b^2) * 2^{1/2} / ((-b + (-4ac + b^2) \\ &)^{1/2}) / a^{1/2} * (4 - 2(-b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} * (4 + 2(b + (-4ac + \\ & + b^2)^{1/2}) / a * x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} * \text{EllipticF}(1/2 * x^2^{1/2} * ((- \\ & b + (-4ac + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2} \\ &) - 1/2 * (1/c + (2ac - b^2) / c / (4ac - b^2)) * a * 2^{1/2} / ((-b + (-4ac + b^2)^{1/2}) / a \\ &)^{1/2} * (4 - 2(-b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} * (4 + 2(b + (-4ac + b^2)^{1/2} \\ &) / a * x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) * (\text{EllipticF}(1/2 \\ & * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2} \\ &) / a / c)^{1/2}) - \text{EllipticE}(1/2 * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2}, \\ & 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2})) + (A e^2 + 2 * B * d * e) * (-2c (1/2 \\ & * b/c / (4ac - b^2) * x^3 + a/c / (4ac - b^2) * x) / ((x^4 + b/c x^2 + a/c) * c)^{1/2} + 1/2 * a / (\\ & 4ac - b^2) * 2^{1/2} / ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4ac + b^2)^{1/2} \\ &) / a * x^2)^{1/2} * (4 + 2(b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} / (c x^4 + b x^2 + a \\ &)^{1/2} * \text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 \\ & * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2}) - 1/2 * b / (4ac - b^2) * a * 2^{1/2} / ((-b + (-4 \\ & ac + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} * (4 + 2(b + \\ & (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2} \\ &) * (\text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (\\ & b + (-4ac + b^2)^{1/2}) / a / c)^{1/2}) - \text{EllipticE}(1/2 * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2} \\ &) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2})) + (2 * A * d * e + \\ & B * d^2) * (-2c (-1 / (4ac - b^2) * x^3 - 1/2 * b / (4ac - b^2) / c * x) / ((x^4 + b/c x^2 + a/c) * \\ & c)^{1/2} - 1/4 * b / (4ac - b^2) * 2^{1/2} / ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * (4 - 2(- \\ & b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} * (4 + 2(b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} \\ &) / (c x^4 + b x^2 + a)^{1/2} * \text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a) \\ &)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2}) + c / (4ac - b^2) * a * 2^{1/2} / \\ & ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2} * (4 - 2(-b + (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} * (4 + 2(b + \\ & (-4ac + b^2)^{1/2}) / a * x^2)^{1/2} / (c x^4 + b x^2 + a)^{1/2} / (b + (-4ac + b^2)^{1/2}) * (\text{EllipticF}(1/2 * x^2^{1/2} * ((-b + (-4ac + b^2)^{1/2}) / a)^{1/2}, 1 \\ & / 2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2}) - \text{EllipticE}(1/2 * x^2^{1/2} * ((-b + \\ & (-4ac + b^2)^{1/2}) / a)^{1/2}, 1/2 * (-4 + 2 * b * (b + (-4ac + b^2)^{1/2}) / a / c)^{1/2})) \end{aligned}$$

))+A*d^2*(-2*c*(1/2/a*b/(4*a*c-b^2)*x^3-1/2*(2*a*c-b^2)/a/(4*a*c-b^2)/c*x)/((x^4+b/c*x^2+a/c)*c)^(1/2)+1/4*(1/a-(2*a*c-b^2)/a/(4*a*c-b^2))*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-1/2*b*c/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^2/(c*x^4 + b*x^2 + a)^(3/2), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^2}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**2/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)**2/(a + b*x**2 + c*x**4)**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((B*x^2+A)*(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")``[Out] integrate((B*x^2 + A)*(x^2*e + d)^2/(c*x^4 + b*x^2 + a)^(3/2), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^2}{(cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2),x)``[Out] int(((A + B*x^2)*(d + e*x^2)^2)/(a + b*x^2 + c*x^4)^(3/2), x)`

$$3.26 \quad \int \frac{(A+Bx^2)(d+ex^2)}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=481

$$\frac{x(aB(bd-2ae) - A(b^2d-2acd-abe) - (Ac(bd-2ae) - aB(2cd-be))x^2)}{a(b^2-4ac)\sqrt{a+bx^2+cx^4}} - \frac{(Ac(bd-2ae) - aB(2cd-be))}{a\sqrt{c}(b^2-4ac)}$$

[Out] $-x*(a*B*(-2*a*e+b*d)-A*(-a*b*e-2*a*c*d+b^2*d)-(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*x^2)/a/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}-(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*x*(c*x^4+b*x^2+a)^{(1/2)}/a/(-4*a*c+b^2)/c^{(1/2)}/(a^{(1/2)}+x^2*c^{(1/2)})+(A*c*(-2*a*e+b*d)-a*B*(-b*e+2*c*d))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^{(1/2)})^2/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(3/4)}/(-4*a*c+b^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^{(1/2)})^2/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(B*a^{(1/2)}-A*c^{(1/2)})*(-e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(3/4)}/(b-2*a^{(1/2)}*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A]

time = 0.24, antiderivative size = 481, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {1692, 1211, 1117, 1209}

$$\frac{(\sqrt{c} + \sqrt{c}x) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{c} + \sqrt{c}x)^2}} F\left(2\text{ArcTan}\left(\frac{\sqrt{c}}{\sqrt{a}}\right) \middle| 2 - \frac{a}{\sqrt{a}\sqrt{c}}\right) (A(bd-2ae) - aB(2cd-be))}{a^{3/4}c^{3/4}(b^2-4ac)\sqrt{a+bx^2+cx^4}} + \frac{(\sqrt{c} + \sqrt{c}x) (\sqrt{c}B - A\sqrt{c}) \sqrt{\frac{a+bx^2+cx^4}{(\sqrt{c} + \sqrt{c}x)^2}} (\sqrt{c}d - \sqrt{c}e) F\left(2\text{ArcTan}\left(\frac{\sqrt{c}}{\sqrt{a}}\right) \middle| 2 - \frac{a}{\sqrt{a}\sqrt{c}}\right)}{2a^{3/4}c^{3/4}(b^2-4ac)\sqrt{a+bx^2+cx^4}} + \frac{x\sqrt{a+bx^2+cx^4} (Ac(bd-2ae) - aB(2cd-be))}{a\sqrt{c}(b^2-4ac)\sqrt{a+bx^2+cx^4}} - \frac{x(-A(-abe-2cd+e^2d) - (2^2(Ac(bd-2ae) - aB(2cd-be))) + aB(bd-2ae))}{a(b^2-4ac)\sqrt{a+bx^2+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] $-((x*(a*B*(b*d-2*a*e) - A*(b^2*d-2*a*c*d - a*b*e) - (A*c*(b*d-2*a*e) - a*B*(2*c*d-b*e))*x^2)/(a*(b^2-4*a*c)*\text{Sqrt}[a+b*x^2+c*x^4])) - ((A*c*(b*d-2*a*e) - a*B*(2*c*d-b*e))*x*\text{Sqrt}[a+b*x^2+c*x^4])/(a*\text{Sqrt}[c]*(b^2-4*a*c)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + ((A*c*(b*d-2*a*e) - a*B*(2*c*d-b*e))*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a+b*x^2+c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2-b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^{(3/4)}*c^{(3/4)}*(b^2-4*a*c)*\text{Sqrt}[a+b*x^2+c*x^4]) + ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])* (\text{Sqrt}[c]*d - \text{Sqrt}[a]*e)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a+b*x^2+c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2-b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{(3/4)}*(b-2*\text{Sqrt}[a]*\text{Sqrt}[c])*c^{(3/4)})*\text{Sqrt}[a+b*x^2+c*x^4])$

Rule 1117

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1209

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1211

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1692

```
Int[(Pq_)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := With[{d = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 0], e = Coeff[PolynomialRemainder[Pq, a + b*x^2 + c*x^4, x], x, 2]}, Simp[x*(a + b*x^2 + c*x^4)^(p + 1)*((a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[(a + b*x^2 + c*x^4)^(p + 1)*ExpandToSum[2*a*(p + 1)*(b^2 - 4*a*c)*PolynomialQuotient[Pq, a + b*x^2 + c*x^4, x] + b^2*d*(2*p + 3) - 2*a*c*d*(4*p + 5) - a*b*e + c*(4*p + 7)*(b*d - 2*a*e)*x^2, x], x], x] /; FreeQ[{a, b, c}, x] && PolyQ[Pq, x^2] && Expon[Pq, x^2] > 1 && NeQ[b^2 - 4*a*c, 0] && LtQ[p, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{3/2}} dx &= -\frac{x(aB(bd - 2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x(aB(bd - 2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x(aB(bd - 2ae) - A(b^2d - 2acd - abe) - (Ac(bd - 2ae) - aB(2cd - be))x^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 11.69, size = 597, normalized size = 1.24

$$\frac{\sqrt{\frac{a^2(b^2 - 4ac)^2 + 4a^2c^2}{(b^2 - 4ac)^2 + 4c^2}} \operatorname{arctanh}\left(\frac{\sqrt{\frac{a^2(b^2 - 4ac)^2 + 4a^2c^2}}}{b^2 - 4ac}\right) - (Ac(bd - 2ae) - aB(2cd - be)) \operatorname{arctanh}\left(\frac{\sqrt{\frac{a^2(b^2 - 4ac)^2 + 4a^2c^2}}}{b^2 - 4ac}\right) + \frac{2a^2c^2}{b^2 - 4ac} \operatorname{arctanh}\left(\frac{\sqrt{\frac{a^2(b^2 - 4ac)^2 + 4a^2c^2}}}{b^2 - 4ac}\right) + \frac{2a^2c^2}{b^2 - 4ac} \operatorname{arctanh}\left(\frac{\sqrt{\frac{a^2(b^2 - 4ac)^2 + 4a^2c^2}}}{b^2 - 4ac}\right)}{a^2(b^2 - 4ac)^2 \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] (4*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x*(a*B*(-2*a*e + 2*c*d*x^2 + b*(d - e*x^2)) + A*(-(b^2*d) + b*(a*e - c*d*x^2) + 2*a*c*(d + e*x^2))) + I*(-b + Sqrt[b^2 - 4*a*c])*(A*c*(b*d - 2*a*e) + a*B*(-2*c*d + b*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]) - I*(A*c*(-(b^2*d) + 4*a*c*d + b*Sqrt[b^2 - 4*a*c]*d - 2*a*Sqrt[b^2 - 4*a*c]*e) + a*B*(b*(-b + Sqrt[b^2 - 4*a*c])*e + c*(-2*Sqrt[b^2 - 4*a*c]*d + 4*a*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))/(4*a*c*(-b^2 + 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])])*Sqrt[a + b*x^2 + c*x^4])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1389 vs. 2(471) = 942.

time = 0.04, size = 1390, normalized size = 2.89

method	result
--------	--------

elliptic	$-\frac{2c\left(-\frac{(2aceA-Abcd-Babe+2acdB)x^3}{2ca(4ac-b^2)}-\frac{(Aabe+2acdA-Ab^2d-2e a^2B+Babd)x}{2ca(4ac-b^2)}\right)}{\sqrt{\left(x^4+\frac{bx^2}{c}+\frac{a}{c}\right)c}}+\frac{\left(\frac{Be}{c}+\frac{Acd-aBe}{ac}-\frac{Aabe+2acdA-Ab^2d-2e a^2B+Babd}{a(4ac-b^2)}\right)}{c}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$B*e*(-2*c*(1/2*b/c/(4*a*c-b^2)*x^3+a/c/(4*a*c-b^2)*x)/((x^4+b/c*x^2+a/c)*c)^{(1/2)}+1/2*a/(4*a*c-b^2)*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/2*b/(4*a*c-b^2)*a*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+A*d*(-2*c*(-1/(4*a*c-b^2)*x^3-1/2*b/(4*a*c-b^2)/c*x)/((x^4+b/c*x^2+a/c)*c)^{(1/2)}-1/4*b/(4*a*c-b^2)*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})+c/(4*a*c-b^2)*a*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+A*d*(-2*c*(1/2*a*b/(4*a*c-b^2)*x^3-1/2*(2*a*c-b^2)/a/(4*a*c-b^2)/c*x)/((x^4+b/c*x^2+a/c)*c)^{(1/2)}+1/4*(1/a-(2*a*c-b^2)/a/(4*a*c-b^2))*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-1/2*b*c/(4*a*c-b^2)*2^{(1/2)}/((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)}*(4-2*(-b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}*(4+2*(b+(-4*a*c+b^2)^{(1/2)})/a*x^2)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*(EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})-EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}))$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)/(c*x^4 + b*x^2 + a)^(3/2), x)

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)*(d + e*x**2)/(a + b*x**2 + c*x**4)**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)/(c*x^4 + b*x^2 + a)^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)}{(cx^4 + bx^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2),x)

[Out] int(((A + B*x^2)*(d + e*x^2))/(a + b*x^2 + c*x^4)^(3/2), x)

$$3.27 \quad \int \frac{A+Bx^2}{(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=398

$$\frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a+bx^2+cx^4}} - \frac{(Ab - 2aB)\sqrt{c}x\sqrt{a+bx^2+cx^4}}{a(b^2 - 4ac)(\sqrt{a} + \sqrt{c}x^2)} + \frac{(Ab - 2aB)^4\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)}{a^2(b^2 - 4ac)^2(\sqrt{a} + \sqrt{c}x^2)}$$

[Out] $x*(A*b^2 - a*b*B - 2*a*A*c + (A*b - 2*a*B)*c*x^2)/a/(-4*a*c + b^2)/(c*x^4 + b*x^2 + a)^{(1/2)} - (A*b - 2*a*B)*x*c^{(1/2)}*(c*x^4 + b*x^2 + a)^{(1/2)}/a/(-4*a*c + b^2)/(a^{(1/2)} + x^2*c^{(1/2)}) + (A*b - 2*a*B)*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*(2 - b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)} + x^2*c^{(1/2)})*((c*x^4 + b*x^2 + a)/(a^{(1/2)} + x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(-4*a*c + b^2)/(c*x^4 + b*x^2 + a)^{(1/2)} + 1/2*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})), 1/2*(2 - b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(B*a^{(1/2)} - A*c^{(1/2)})*(a^{(1/2)} + x^2*c^{(1/2)})*((c*x^4 + b*x^2 + a)/(a^{(1/2)} + x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/c^{(1/4)}/(b - 2*a^{(1/2)}*c^{(1/2)})/(c*x^4 + b*x^2 + a)^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 398, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {1192, 1211, 1117, 1209}

$$\frac{\sqrt{c}(\sqrt{a} + \sqrt{c}x^2)(Ab - 2aB)\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}E\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\middle|1 - \frac{b}{\sqrt{a}\sqrt{c}}\right)}{a^{3/4}(b^2 - 4ac)\sqrt{a+bx^2+cx^4}} + \frac{(\sqrt{a} + \sqrt{c}x^2)(\sqrt{a}B - A\sqrt{c})\sqrt{\frac{a+bx^2+cx^4}{(\sqrt{a} + \sqrt{c}x^2)^2}}F\left(2\text{ArcTan}\left(\frac{\sqrt{c}x}{\sqrt{a}}\right)\middle|1 - \frac{b}{\sqrt{a}\sqrt{c}}\right)}{2a^{3/4}\sqrt{c}(b - 2\sqrt{a}\sqrt{c})\sqrt{a+bx^2+cx^4}} - \frac{\sqrt{c}x(Ab - 2aB)\sqrt{a+bx^2+cx^4}}{a(b^2 - 4ac)(\sqrt{a} + \sqrt{c}x^2)} + \frac{x(cx^2(Ab - 2aB) - 2aAc - abB + Ab^2)}{a(b^2 - 4ac)\sqrt{a+bx^2+cx^4}}$$

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] $(x*(A*b^2 - a*b*B - 2*a*A*c + (A*b - 2*a*B)*c*x^2))/(a*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) - ((A*b - 2*a*B)*\text{Sqrt}[c]*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*(b^2 - 4*a*c)*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)) + ((A*b - 2*a*B)*c^{(1/4)}*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticE}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(a^{(3/4)}*(b^2 - 4*a*c)*\text{Sqrt}[a + b*x^2 + c*x^4]) + ((\text{Sqrt}[a]*B - A*\text{Sqrt}[c])*(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]*x^2)^2]*\text{EllipticF}[2*\text{ArcTan}[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(\text{Sqrt}[a]*\text{Sqrt}[c]))/4])/(2*a^{(3/4)}*(b - 2*\text{Sqrt}[a]*\text{Sqrt}[c])*c^{(1/4)}*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1117

Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]]/

$(2*q*\text{Sqrt}[a + b*x^2 + c*x^4])*\text{EllipticF}[2*\text{ArcTan}[q*x], 1/2 - b*(q^2/(4*c))]$
 $], x]] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1192

$\text{Int}[\{(d_)+(e_)*(x_)^2*\{(a_)+(b_)*(x_)^2+(c_)*(x_)^4\}^{(p_)}, x_Symbol]$
 $:= \text{Simp}[x*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)*\{(a + b*x^2 + c*x^4\}^{(p + 1)}/(2*a*(p + 1)*(b^2 - 4*a*c)), x] + \text{Dist}[1/(2*a*(p + 1)*(b^2 - 4*a*c)), \text{Int}[\text{Simp}[(2*p + 3)*d*b^2 - a*b*e - 2*a*c*d*(4*p + 5) + (4*p + 7)*(d*b - 2*a*e)*c*x^2, x]*(a + b*x^2 + c*x^4)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{NeQ}[c*d^2 - b*d*e + a*e^2, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntegerQ}[2*p]$

Rule 1209

$\text{Int}[\{(d_)+(e_)*(x_)^2/\text{Sqrt}[(a_)+(b_)*(x_)^2+(c_)*(x_)^4], x_Symbol]$
 $:= \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d)*x*(\text{Sqrt}[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + \text{Simp}[d*(1 + q^2*x^2)*(\text{Sqrt}[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2)^2)]/(q*\text{Sqrt}[a + b*x^2 + c*x^4])*\text{EllipticE}[2*\text{ArcTan}[q*x], 1/2 - b*(q^2/(4*c))], x] /; \text{EqQ}[e + d*q^2, 0] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rule 1211

$\text{Int}[\{(d_)+(e_)*(x_)^2/\text{Sqrt}[(a_)+(b_)*(x_)^2+(c_)*(x_)^4], x_Symbol]$
 $:= \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d*q)/q, \text{Int}[1/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] - \text{Dist}[e/q, \text{Int}[(1 - q*x^2)/\text{Sqrt}[a + b*x^2 + c*x^4], x], x] /; \text{NeQ}[e + d*q, 0] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{PosQ}[c/a]$

Rubi steps

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{3/2}} dx = \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{\int \frac{-a(bB - 2Ac) + (Ab - 2aB)cx^2}{\sqrt{a + bx^2 + cx^4}} dx}{a(b^2 - 4ac)}$$

$$= \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} + \frac{((Ab - 2aB)\sqrt{c}) \int \frac{1 - \sqrt{c}x^2}{\sqrt{a + bx^2 + cx^4}}}{\sqrt{a}(b^2 - 4ac)}$$

$$= \frac{x(Ab^2 - abB - 2aAc + (Ab - 2aB)cx^2)}{a(b^2 - 4ac)\sqrt{a + bx^2 + cx^4}} - \frac{(Ab - 2aB)\sqrt{c}x\sqrt{a + bx^2 + cx^4}}{a(b^2 - 4ac)(\sqrt{a} + \sqrt{c}x^2)} +$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.90, size = 497, normalized size = 1.25

$$4\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\frac{a(b^2+2ac^2)-A(b^2-2ac+bc^2)+i(AB-2aB)(-b+\sqrt{b^2-4ac})}{b+\sqrt{b^2-4ac}}\sqrt{\frac{b+\sqrt{b^2-4ac}+2ac^2}{b+\sqrt{b^2-4ac}}}\sqrt{\frac{2b-2\sqrt{b^2-4ac}+4ac^2}{b-\sqrt{b^2-4ac}}}\frac{E\left(\operatorname{arcsinh}\left(\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right)\right)}{4a(b^2-4ac)}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\sqrt{a+bx^2+cx^4}}{-2aB\sqrt{b^2-4ac}+A(-b^2+4ac+b\sqrt{b^2-4ac})}\sqrt{\frac{b+\sqrt{b^2-4ac}+2ac^2}{b+\sqrt{b^2-4ac}}}\sqrt{\frac{2b-2\sqrt{b^2-4ac}+4ac^2}{b-\sqrt{b^2-4ac}}}\frac{F\left(\operatorname{arcsinh}\left(\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\frac{b+\sqrt{b^2-4ac}}{b-\sqrt{b^2-4ac}}\right)\right)}{4a(b^2-4ac)}\sqrt{\frac{c}{b+\sqrt{b^2-4ac}}}\sqrt{a+bx^2+cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x]

[Out] -1/4*(4*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x*(a*B*(b + 2*c*x^2) - A*(b^2 - 2*a*c + b*c*x^2)) + I*(A*b - 2*a*B)*(-b + Sqrt[b^2 - 4*a*c])*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - I*(-2*a*B*Sqrt[b^2 - 4*a*c] + A*(-b^2 + 4*a*c + b*Sqrt[b^2 - 4*a*c]))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x, (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])])/(a*(b^2 - 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[a + b*x^2 + c*x^4])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 930 vs. 2(392) = 784.

time = 0.04, size = 931, normalized size = 2.34

method	result
elliptic	$-\frac{2c\left(\frac{(Ab-2aB)x^3}{2a(4ac-b^2)}-\frac{(2acA-Ab^2+abB)x}{2ac(4ac-b^2)}\right)}{\sqrt{\left(x^4+\frac{bx^2}{c}+\frac{a}{c}\right)c}}+\frac{\left(\frac{A}{a}-\frac{2acA-Ab^2+abB}{(4ac-b^2)a}\right)\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4+\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}}{4\sqrt{-b+\sqrt{-4ac+b^2}}}$
default	$B\left(-\frac{2c\left(-\frac{x^3}{4ac-b^2}-\frac{bx}{2(4ac-b^2)c}\right)}{\sqrt{\left(x^4+\frac{bx^2}{c}+\frac{a}{c}\right)c}}-\frac{b\sqrt{2}\sqrt{4-\frac{2(-b+\sqrt{-4ac+b^2})x^2}{a}}\sqrt{4+\frac{2(b+\sqrt{-4ac+b^2})x^2}{a}}}{4(4ac-b^2)\sqrt{\frac{-b+\sqrt{-4ac+b^2}}{a}}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2), x, method=_RETURNVERBOSE)

[Out] B*(-2*c*(-1/(4*a*c-b^2)*x^3-1/2*b/(4*a*c-b^2)/c*x)/((x^4+b*c*x^2+a/c)*c)^(1/2)-1/4*b/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-

```

4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*
x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2
),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+c/(4*a*c-b^2)*a*2^(1/2)/((
-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(
4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^
2)^(1/2))*(EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-
4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a
*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))))+A*
(-2*c*(1/2/a*b/(4*a*c-b^2)*x^3-1/2*(2*a*c-b^2)/a/(4*a*c-b^2)/c*x)/((x^4+b/c
*x^2+a/c)*c)^(1/2)+1/4*(1/a-(2*a*c-b^2)/a/(4*a*c-b^2))*2^(1/2)/((-b+(-4*a*c
+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4
*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2
)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(
1/2))-1/2*b*c/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(
-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2
)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*x*2^(1/2)*((-
b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2
))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b
+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))))

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((B*x^2 + A)/(c*x^4 + b*x^2 + a)^(3/2), x)
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly
1 arguments (2 given)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{A + Bx^2}{(a + bx^2 + cx^4)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Integral((A + B*x**2)/(a + b*x**2 + c*x**4)**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/(c*x^4 + b*x^2 + a)^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{(c x^4 + b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2),x)

[Out] int((A + B*x^2)/(a + b*x^2 + c*x^4)^(3/2), x)

$$3.28 \quad \int \frac{A+Bx^2}{(d+ex^2)(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=867

$$\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be) - A(bcd - b^2e + 2ace))x^2) + \sqrt{c}(aB(aBcd - b^2e + 2ace))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}} + \frac{\sqrt{c}(aB(aBcd - b^2e + 2ace))}{a}$$

[Out] $-1/2*e^{(3/2)}*(-A*e+B*d)*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}/(a*e^2-b*d*e+c*d^2)^{(3/2)}/d^{(1/2)}-x*(a*b*c*(-A*e+B*d)-(-2*a*c+b^2)*(-A*b*e+A*c*d+B*a*e)+c*(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*x^2)/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(a^{(1/2)}+x^2*c^{(1/2)})-c^{(1/4)}*(a*B*(-b*e+2*c*d)-A*(2*a*c*e-b^2*e+b*c*d))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)}))*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/4*a^{(3/4)}*e*(-A*e+B*d)*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(a^{(1/2)}+x^2*c^{(1/2)})*(e+d*c^{(1/2)}/a^{(1/2)})^2*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/c^{(1/4)}/d/(-a*e^2+c*d^2)/(a*e^2-b*d*e+c*d^2)/(c*x^4+b*x^2+a)^{(1/2)}+1/2*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)}*(B*a^{(1/2)}-A*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)}))*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(-e*a^{(1/2)}+d*c^{(1/2)})/(b-2*a^{(1/2)}*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A]

time = 0.97, antiderivative size = 1045, normalized size of antiderivative = 1.21, number of steps used = 9, number of rules used = 7, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.212$, Rules used = {1734, 1192, 1211, 1117, 1209, 1230, 1720}

Antiderivative was successfully verified.

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)), x]

[Out] $-((x*(a*b*c*(B*d - A*e) - (b^2 - 2*a*c)*(A*c*d - A*b*e + a*B*e) + c*(a*B*(2*c*d - b*e) - A*(b*c*d - b^2*e + 2*a*c*e))*x^2))/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4]) + (\text{Sqrt}[c]*(a*B*(2*c*d - b*e) - A*(b*c*d - b^2*e + 2*a*c*e))*x*\text{Sqrt}[a + b*x^2 + c*x^4])/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*\text{Sqrt}[a + b*x^2 + c*x^4])$

$$\begin{aligned} &^2 - b*d*e + a*e^2)*(Sqrt[a] + Sqrt[c]*x^2)) - (e^{(3/2)}*(B*d - A*e)*ArcTan[\\ &(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])) \\ &/((2*Sqrt[d]*(c*d^2 - b*d*e + a*e^2)^{(3/2)}) - (c^{(1/4)}*(a*B*(2*c*d - b*e) - \\ &A*(b*c*d - b^2*e + 2*a*c*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^ \\ &4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^{(1/4)}*x)/a^{(1/4)}], (2 - \\ &b/(Sqrt[a]*Sqrt[c]))/4])/(a^{(3/4)}*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)*Sqr \\ &rt[a + b*x^2 + c*x^4]) - (c^{(1/4)}*e*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqr \\ &t[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^{(1/4)} \\ &)*x)/a^{(1/4)}], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^{(1/4)}*(Sqrt[c]*d - Sqrt[a \\ &]*e)*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4]) - (c^{(1/4)}*(a*B*e - S \\ &qrt[a]*Sqrt[c]*(B*d - A*e) + A*(c*d - b*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a \\ &+ b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^{(1/4)}*x) \\ &/a^{(1/4)}], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^{(3/4)}*(b - 2*Sqrt[a]*Sqrt[c]) \\ &*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4]) + (a^{(3/4)}*e*((Sqrt[c]*d) \\ &/Sqrt[a] + e)^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4) \\ &)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[c]*d - Sqrt[a]*e)^2/(Sqr \\ &t[a]*Sqrt[c]*d*e), 2*ArcTan[(c^{(1/4)}*x)/a^{(1/4)}], (2 - b/(Sqrt[a]*Sqrt[c])) \\ &/4])/(4*c^{(1/4)}*d*(c*d^2 - a*e^2)*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + \\ &c*x^4]) \end{aligned}$$

Rule 1117

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(2*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticF[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1192

```
Int[((d_) + (e_.)*(x_)^2)*((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4)^(p_), x_Symbol] := Simp[x*(a*b*e - d*(b^2 - 2*a*c) - c*(b*d - 2*a*e)*x^2)*((a + b*x^2 + c*x^4)^(p + 1)/(2*a*(p + 1)*(b^2 - 4*a*c))), x] + Dist[1/(2*a*(p + 1)*(b^2 - 4*a*c)), Int[Simp[(2*p + 3)*d*b^2 - a*b*e - 2*a*c*d*(4*p + 5) + (4*p + 7)*(d*b - 2*a*e)*c*x^2, x]*(a + b*x^2 + c*x^4)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[p, -1] && IntegerQ[2*p]
```

Rule 1209

```
Int[((d_) + (e_.)*(x_)^2)/Sqrt[(a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4], x_Symbol] := With[{q = Rt[c/a, 4]}, Simp[(-d)*x*(Sqrt[a + b*x^2 + c*x^4]/(a*(1 + q^2*x^2))), x] + Simp[d*(1 + q^2*x^2)*(Sqrt[(a + b*x^2 + c*x^4)/(a*(1 + q^2*x^2)^2)]/(q*Sqrt[a + b*x^2 + c*x^4]))*EllipticE[2*ArcTan[q*x], 1/2 - b*(q^2/(4*c))], x] /; EqQ[e + d*q^2, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1211

```
Int[((d_) + (e_)*(x_)^2)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol]
:> With[{q = Rt[c/a, 2]}, Dist[(e + d*q)/q, Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[e/q, Int[(1 - q*x^2)/Sqrt[a + b*x^2 + c*x^4], x], x] /; NeQ[e + d*q, 0] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && PosQ[c/a]
```

Rule 1230

```
Int[1/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol]
:> With[{q = Rt[c/a, 2]}, Dist[(c*d + a*e*q)/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[(a*e*(e + d*q))/(c*d^2 - a*e^2), Int[(1 + q*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a]
```

Rule 1720

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol]
:> With[{q = Rt[B/A, 2]}, Simp[(-(B*d - A*e))*(ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/(4*a*B))], x] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]
```

Rule 1734

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol]
:> Int[ExpandIntegrand[1/Sqrt[a + b*x^2 + c*x^4], Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e}, x] && PolyQ[Px, x^2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\begin{aligned}
\int \frac{A + Bx^2}{(d + ex^2)(a + bx^2 + cx^4)^{3/2}} dx &= \int \left(\frac{Acd - Abe + aBe + c(Bd - Ae)x^2}{(cd^2 - bde + ae^2)(a + bx^2 + cx^4)^{3/2}} + \frac{e(-Bd + Ae)}{(cd^2 - bde + ae^2)(d + ex^2)} \right) dx \\
&= \frac{\int \frac{Acd - Abe + aBe + c(Bd - Ae)x^2}{(a + bx^2 + cx^4)^{3/2}} dx}{cd^2 - bde + ae^2} - \frac{(e(Bd - Ae)) \int \frac{1}{(d + ex^2)\sqrt{a + bx^2 + cx^4}} dx}{cd^2 - bde + ae^2} \\
&= -\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be)))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be)))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}} \\
&= -\frac{x(abc(Bd - Ae) - (b^2 - 2ac)(Acd - Abe + aBe) + c(aB(2cd - be)))}{a(b^2 - 4ac)(cd^2 - bde + ae^2)\sqrt{a + bx^2 + cx^4}}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.
time = 13.29, size = 1736, normalized size = 2.00

Antiderivative was successfully verified.

[In] Integrate[(A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)),x]

[Out] (-4*A*b^2*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 4*a*b*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 8*a*A*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x + 4*A*b^3*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 4*a*b^2*B*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 12*a*A*b*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x + 8*a^2*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x - 4*A*b*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x^3 + 8*a*B*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d^2*x^3 + 4*A*b^2*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - 4*a*b*B*c*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - 8*a*A*c^2*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*e*x^3 - I*(-b + Sqrt[b^2 - 4*a*c])*d*(a*B*(2*c*d - b*e) + A*(-(b*c*d) + b^2*e - 2*a*c*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + I*d*(a*B*(b*(b - Sqrt[b^2 - 4*a*c])*e + 2*c*(Sqrt[b^2 - 4*a*c]*d - 2*a*e)) + A*(-(b^3*e) + b*c*(-(Sqrt[b^2 - 4

```

*a*c]*d) + 4*a*e) + b^2*(c*d + Sqrt[b^2 - 4*a*c]*e) - 2*a*c*(2*c*d + Sqrt[b
^2 - 4*a*c]*e))*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a
*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*E
llipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt[b^
2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - (2*I)*a*b^2*B*d*e*Sqrt[(b + Sqrt[b^2
- 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4*a*
c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*
e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + Sqrt
[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (8*I)*a^2*B*c*d*e*Sqrt[(b + Sqrt[
b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2 - 4
*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 - 4*a*c
])e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b + S
qrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] + (2*I)*a*A*b^2*e^2*Sqrt[(b + Sq
rt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b^2
- 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 - 4*
a*c])e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x], (b
+ Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - (8*I)*a^2*A*c*e^2*Sqrt[(b +
Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c])]*Sqrt[(2*b - 2*Sqrt[b
^2 - 4*a*c] + 4*c*x^2)/(b - Sqrt[b^2 - 4*a*c])]*EllipticPi[((b + Sqrt[b^2 -
4*a*c])e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*x],
(b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]/(4*a*(-b^2 + 4*a*c)*Sqrt[
c/(b + Sqrt[b^2 - 4*a*c])]*d*(c*d^2 + e*(-(b*d) + a*e))*Sqrt[a + b*x^2 + c*
x^4])

```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 3240 vs. $2(832) = 1664$.

time = 0.14, size = 3241, normalized size = 3.74

method	result	size
default	Expression too large to display	3241
elliptic	Expression too large to display	4597

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] B/e*(-2*c*(1/2/a*b/(4*a*c-b^2)*x^3-1/2*(2*a*c-b^2)/a/(4*a*c-b^2)/c*x)/((x^4
+b/c*x^2+a/c)*c)^(1/2)+1/4*(1/a-(2*a*c-b^2)/a/(4*a*c-b^2))*2^(1/2)/((-b+(-4
*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b
+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(
1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a
/c)^(1/2))-1/2*b*c/(4*a*c-b^2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4
-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(
1/2)/(c*x^4+b*x^2+a)^(1/2)/(b+(-4*a*c+b^2)^(1/2))*(EllipticF(1/2*x*2^(1/2)
)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(
1/2))-EllipticE(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*
```

$$\begin{aligned}
& b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})))+(A*e-B*d)/e*(-2*c*(1/2*(2*a*c*e-b^2* \\
& e+b*c*d)/a/(4*a*c-b^2)/(a*e^2-b*d*e+c*d^2)*x^3+1/2*(3*a*b*c*e-2*a*c^2*d-b^3 \\
& *e+b^2*c*d)/a/(4*a*c-b^2)/(a*e^2-b*d*e+c*d^2)/c*x)/((x^4+b*c*x^2+a/c)*c)^{(1 \\
& /2)}-1/4*2^{(1/2)}*(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(- \\
& 4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x \\
& ^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)} \\
& ,1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})/a/(a*e^2-b*d*e+c*d^2)*e*b+1 \\
& /4*2^{(1/2)}*(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c \\
& +b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b* \\
& x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2* \\
& (-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})/a/(a*e^2-b*d*e+c*d^2)*c*d+3/4*2^{(\\
& 1/2)}*(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2) \\
& ^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a \\
&)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2 \\
& *b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})/(4*a*c-b^2)/(a*e^2-b*d*e+c*d^2)*b*c*e \\
& -1/2*2^{(1/2)}*(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a \\
& *c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+ \\
& b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/ \\
& 2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})/(4*a*c-b^2)/(a*e^2-b*d*e+c*d^2 \\
&)*c^2*d-1/4*2^{(1/2)}*(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/ \\
& a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/ \\
& (c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(\\
& 1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})/a/(4*a*c-b^2)/(a*e^2-b* \\
& d*e+c*d^2)*b^3*e+1/4*2^{(1/2)}*(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2 \\
& /a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2) \\
&)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1 \\
& /2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})/a/(4*a*c-b^2)/ \\
& (a*e^2-b*d*e+c*d^2)*b^2*c*d-c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^{(1/2)}/(-b \\
& /a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(\\
& 1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(\\
& b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(\\
& 1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})*a*e+1/2*c/(a*e^2-b*d*e+ \\
& c*d^2)/(4*a*c-b^2)*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a \\
& -2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2) \\
&)^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticF(1/2*x*2^{(1/2) \\
& }*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(\\
& 1/2)})*b^2*e-1/2*c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^{(1/2)}/(-b/a+1/a*(-4* \\
& a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b \\
& *x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)}/(b+(-4*a*c+b \\
& ^2)^{(1/2)})*EllipticF(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(- \\
& 4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})*b*d+c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c \\
& -b^2)*2^{(1/2)}/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a-2*x^2/a*(-4* \\
& a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}/(c*x^4 \\
& +b*x^2+a)^{(1/2)}/(b+(-4*a*c+b^2)^{(1/2)})*EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c \\
& +b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})*a*e-1/
\end{aligned}$$

$$2*c/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^{(1/2)/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)/(b+(-4*a*c+b^2)^{(1/2)})}*EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)}*b^2*e+1/2*c^2/(a*e^2-b*d*e+c*d^2)/(4*a*c-b^2)*2^{(1/2)/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}}*(4+2*b*x^2/a-2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(4+2*b*x^2/a+2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)/(b+(-4*a*c+b^2)^{(1/2)})}*EllipticE(1/2*x*2^{(1/2)}*((-b+(-4*a*c+b^2)^{(1/2)})/a)^{(1/2)},1/2*(-4+2*b*(b+(-4*a*c+b^2)^{(1/2)})/a/c)^{(1/2)})*b*d+e^2/(a*e^2-b*d*e+c*d^2)/d*2^{(1/2)/(-b/a+1/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)}*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^{(1/2)})^{(1/2)/(c*x^4+b*x^2+a)^{(1/2)}*EllipticPi(1/2*x...$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(x^2*e + d)), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(x^2*e + d)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{(e x^2 + d) (c x^4 + b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(3/2)), x)

$$3.29 \quad \int \frac{A+Bx^2}{(d+ex^2)^2(a+bx^2+cx^4)^{3/2}} dx$$

Optimal. Leaf size=1301

$$\frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae))) - c(aB(a(b^2 - 4ac)(cd^2 - bde + ae^2)^2 \sqrt{a + b$$

[Out] $\frac{1}{4}e^{3/2}*(A*e*(7*c*d^2-e*(-a*e+4*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*\arctan(x*(a*e^2-b*d*e+c*d^2)^{(1/2)}/d^{(1/2)}/e^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})/d^{(3/2)}/(a*e^2-b*d*e+c*d^2)^{(5/2)}+x*(a*b*c*(A*e*(-b*e+2*c*d)-B*(-a*e^2+c*d^2))+(-2*a*c+b^2)*(a*B*e*(-b*e+2*c*d)+A*(c^2*d^2+b^2*e^2-c*e*(a*e+2*b*d)))-c*(a*B*(2*c^2*d^2+b^2*e^2-2*c*e*(a*e+b*d))+A*(2*b^2*c*d*e-4*a*c^2*d*e-b^3*e^2-b*c*(-3*a*e^2+c*d^2)))*x^2)/a/(-4*a*c+b^2)/(a*e^2-b*d*e+c*d^2)^2/(c*x^4+b*x^2+a)^{(1/2)}-1/2*e^3*(-A*e+B*d)*x*(c*x^4+b*x^2+a)^{(1/2)}/d/(a*e^2-b*d*e+c*d^2)^2/(e*x^2+d)+1/2*(a*B*d*(-4*c^2*d^2-3*b^2*e^2+4*c*e*(2*a*e+b*d))+A*(2*b^3*d*e^2+2*b*c*d*(-3*a*e^2+c*d^2)-4*a*c*e*(a*e^2-2*c*d^2)+b^2*(a*e^3-4*c*d^2*e)))*x*c^{(1/2)}*(c*x^4+b*x^2+a)^{(1/2)}/a/(4*a*c-b^2)/d/(c*d^2+e*(a*e-b*d))^2/(a^{(1/2)}+x^2*c^{(1/2)})-1/2*c^{(1/4)}*(a*B*d*(4*c^2*d^2+3*b^2*e^2-4*c*e*(2*a*e+b*d))-A*(2*b^3*d*e^2+2*b*c*d*(-3*a*e^2+c*d^2)-4*a*c*e*(a*e^2-2*c*d^2)+b^2*(a*e^3-4*c*d^2*e)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticE}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/(-4*a*c+b^2)/d/(c*d^2+e*(a*e-b*d))^2/(c*x^4+b*x^2+a)^{(1/2)}-1/8*e*(A*e*(7*c*d^2-e*(-a*e+4*b*d))-B*d*(5*c*d^2-e*(a*e+2*b*d)))*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticPi}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),-1/4*(-e*a^{(1/2)}+d*c^{(1/2)})^2/d/e/a^{(1/2)}/c^{(1/2)},1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*(e*a^{(1/2)}+d*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(1/4)}/c^{(1/4)}/d^2/(a*e^2-b*d*e+c*d^2)^2/(-e*a^{(1/2)}+d*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}+1/2*c^{(1/4)}*(\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))^2)^{(1/2)}/\cos(2*\arctan(c^{(1/4)}*x/a^{(1/4)}))*\text{EllipticF}(\sin(2*\arctan(c^{(1/4)}*x/a^{(1/4)})),1/2*(2-b/a^{(1/2)}/c^{(1/2)})^{(1/2)})*((-A*e+B*d)*(-b*e+c*d)*a^{(1/2)}+a*e*(-2*A*e+B*d)*c^{(1/2)}+A*d*(b*e-c*d)*c^{(1/2)})*(a^{(1/2)}+x^2*c^{(1/2)})*((c*x^4+b*x^2+a)/(a^{(1/2)}+x^2*c^{(1/2)})^2)^{(1/2)}/a^{(3/4)}/d/(-c*d^2+e*(-a*e+b*d))/(e*a^{(1/2)}-d*c^{(1/2)})/(b-2*a^{(1/2)}*c^{(1/2)})/(c*x^4+b*x^2+a)^{(1/2)}$

Rubi [A]

time = 2.21, antiderivative size = 2112, normalized size of antiderivative = 1.62, number of steps used = 15, number of rules used = 10, integrand size = 33, $\frac{\text{number of rules}}{\text{integrand size}} = 0.303$, Rules used = {1734, 1192, 1211, 1117, 1209, 1237, 1728, 1722, 1720, 1230}

Antiderivative was successfully verified.

[In] Int[(A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x]

[Out] (x*(a*b*c*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2)) + (b^2 - 2*a*c)*(a*B*e*(2*c*d - b*e) + A*(c^2*d^2 + b^2*e^2 - c*e*(2*b*d + a*e))) - c*(a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*x^2)/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]) + (Sqrt[c]*e^2*(B*d - A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*(Sqrt[a] + Sqrt[c]*x^2)) + (Sqrt[c]*(a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*x*Sqrt[a + b*x^2 + c*x^4])/(a*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)^2*(Sqrt[a] + Sqrt[c]*x^2)) - (e^3*(B*d - A*e)*x*Sqrt[a + b*x^2 + c*x^4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*(d + e*x^2)) - (e^(3/2)*(B*d - A*e)*(3*c*d^2 - e*(2*b*d - a*e))*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(4*d^(3/2)*(c*d^2 - b*d*e + a*e^2)^(5/2)) + (e^(3/2)*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2))*ArcTan[(Sqrt[c*d^2 - b*d*e + a*e^2]*x)/(Sqrt[d]*Sqrt[e]*Sqrt[a + b*x^2 + c*x^4])])/(2*Sqrt[d]*(c*d^2 - b*d*e + a*e^2)^(5/2)) - (a^(1/4)*c^(1/4)*e^2*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*d*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]) - (c^(1/4)*(a*B*(2*c^2*d^2 + b^2*e^2 - 2*c*e*(b*d + a*e)) + A*(2*b^2*c*d*e - 4*a*c^2*d*e - b^3*e^2 - b*c*(c*d^2 - 3*a*e^2)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticE[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(a^(3/4)*(b^2 - 4*a*c)*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]) - (c^(1/4)*e*(B*d - A*e)*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(1/4)*d*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 - b*d*e + a*e^2)*Sqrt[a + b*x^2 + c*x^4]) + (c^(1/4)*e*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(1/4)*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]) - (c^(1/4)*(a^(3/2)*B*Sqrt[c]*e^2 + A*(c*d - b*e)^2 + a*e*(2*B*c*d - b*B*e - A*c*e) - Sqrt[a]*Sqrt[c]*(B*c*d^2 - A*e*(2*c*d - b*e)))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticF[2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(2*a^(3/4)*(b - 2*Sqrt[a]*Sqrt[c])*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]) + (e*(Sqrt[c]*d + Sqrt[a]*e)*(B*d - A*e)*(3*c*d^2 - e*(2*b*d - a*e))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[c]*d - Sqrt[a]*e)^2/(Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(8*a^(1/4)*c^(1/4)*d^2*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4]) - (a^(3/4)*e*((Sqrt[c]*d)/Sqrt[a] + e)^2*(A*e*(2*c*d - b*e) - B*(c*d^2 - a*e^2))*(Sqrt[a] + Sqrt[c]*x^2)*Sqrt[(a + b*x^2 + c*x^4)/(Sqrt[a] + Sqrt[c]*x^2)^2]*EllipticPi[-1/4*(Sqrt[c]*d - Sqrt[a]*e)^2/(Sqrt[a]*Sqrt[c]*d*e), 2*ArcTan[(c^(1/4)*x)/a^(1/4)], (2 - b/(Sqrt[a]*Sqrt[c]))/4])/(8*a^(1/4)*c^(1/4)*d^2*(Sqrt[c]*d - Sqrt[a]*e)*(c*d^2 - b*d*e + a*e^2)^2*Sqrt[a + b*x^2 + c*x^4])

$a] * e)^2 / (\text{Sqrt}[a] * \text{Sqrt}[c] * d * e), 2 * \text{ArcTan}[(c^{1/4} * x) / a^{1/4}], (2 - b / (\text{Sqrt}[a] * \text{Sqrt}[c])) / 4] / (4 * c^{1/4} * d * (c * d^2 - a * e^2) * (c * d^2 - b * d * e + a * e^2)^2 * \text{Sqrt}[a + b * x^2 + c * x^4])$

Rule 1117

$\text{Int}[1 / \text{Sqrt}[(a_) + (b_) * (x_)^2 + (c_) * (x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(1 + q^2 * x^2) * (\text{Sqrt}[a + b * x^2 + c * x^4] / (a * (1 + q^2 * x^2)^2))] / (2 * q * \text{Sqrt}[a + b * x^2 + c * x^4))] * \text{EllipticF}[2 * \text{ArcTan}[q * x], 1/2 - b * (q^2 / (4 * c))] , x] /; \text{FreeQ}[\{a, b, c\}, x] \&\& \text{NeQ}[b^2 - 4 * a * c, 0] \&\& \text{PosQ}[c/a]$

Rule 1192

$\text{Int}[(d_) + (e_) * (x_)^2 * ((a_) + (b_) * (x_)^2 + (c_) * (x_)^4)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[x * (a * b * e - d * (b^2 - 2 * a * c) - c * (b * d - 2 * a * e) * x^2) * ((a + b * x^2 + c * x^4)^{(p + 1)} / (2 * a * (p + 1) * (b^2 - 4 * a * c))), x] + \text{Dist}[1 / (2 * a * (p + 1) * (b^2 - 4 * a * c)), \text{Int}[\text{Simp}[(2 * p + 3) * d * b^2 - a * b * e - 2 * a * c * d * (4 * p + 5) + (4 * p + 7) * (d * b - 2 * a * e) * c * x^2, x] * (a + b * x^2 + c * x^4)^{(p + 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4 * a * c, 0] \&\& \text{NeQ}[c * d^2 - b * d * e + a * e^2, 0] \&\& \text{LtQ}[p, -1] \&\& \text{IntegerQ}[2 * p]$

Rule 1209

$\text{Int}[(d_) + (e_) * (x_)^2 / \text{Sqrt}[(a_) + (b_) * (x_)^2 + (c_) * (x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 4]\}, \text{Simp}[(-d) * x * (\text{Sqrt}[a + b * x^2 + c * x^4] / (a * (1 + q^2 * x^2))), x] + \text{Simp}[d * (1 + q^2 * x^2) * (\text{Sqrt}[a + b * x^2 + c * x^4] / (a * (1 + q^2 * x^2)^2))] / (q * \text{Sqrt}[a + b * x^2 + c * x^4))] * \text{EllipticE}[2 * \text{ArcTan}[q * x], 1/2 - b * (q^2 / (4 * c))] , x] /; \text{EqQ}[e + d * q^2, 0] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4 * a * c, 0] \&\& \text{PosQ}[c/a]$

Rule 1211

$\text{Int}[(d_) + (e_) * (x_)^2 / \text{Sqrt}[(a_) + (b_) * (x_)^2 + (c_) * (x_)^4], x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(e + d * q) / q, \text{Int}[1 / \text{Sqrt}[a + b * x^2 + c * x^4], x], x] - \text{Dist}[e / q, \text{Int}[(1 - q * x^2) / \text{Sqrt}[a + b * x^2 + c * x^4], x], x] /; \text{NeQ}[e + d * q, 0] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4 * a * c, 0] \&\& \text{PosQ}[c/a]$

Rule 1230

$\text{Int}[1 / (((d_) + (e_) * (x_)^2) * \text{Sqrt}[(a_) + (b_) * (x_)^2 + (c_) * (x_)^4]), x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[c/a, 2]\}, \text{Dist}[(c * d + a * e * q) / (c * d^2 - a * e^2), \text{Int}[1 / \text{Sqrt}[a + b * x^2 + c * x^4], x], x] - \text{Dist}[(a * e * (e + d * q)) / (c * d^2 - a * e^2), \text{Int}[(1 + q * x^2) / ((d + e * x^2) * \text{Sqrt}[a + b * x^2 + c * x^4]), x], x]] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&\& \text{NeQ}[b^2 - 4 * a * c, 0] \&\& \text{NeQ}[c * d^2 - b * d * e + a * e^2, 0] \&\& \text{NeQ}[c * d^2 - a * e^2, 0] \&\& \text{PosQ}[c/a]$

Rule 1237

```
Int[((d_) + (e_)*(x_)^2)^(q_)/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_
Symbol] := Simp[(-e^2)*x*(d + e*x^2)^(q + 1)*(Sqrt[a + b*x^2 + c*x^4]/(2*d*
(q + 1)*(c*d^2 - b*d*e + a*e^2))), x] + Dist[1/(2*d*(q + 1)*(c*d^2 - b*d*e
+ a*e^2)), Int[((d + e*x^2)^(q + 1)/Sqrt[a + b*x^2 + c*x^4])*Simp[a*e^2*(2*
q + 3) + 2*d*(c*d - b*e)*(q + 1) - 2*e*(c*d*(q + 1) - b*e*(q + 2))*x^2 + c*
e^2*(2*q + 5)*x^4, x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a
*c, 0] && ILtQ[q, -1]
```

Rule 1720

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 +
(c_)*(x_)^4]), x_Symbol] := With[{q = Rt[B/A, 2]}, Simp[(-B*d - A*e)*(A
rcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[
-b + c*(d/e) + a*(e/d), 2])), x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(
a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2))]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4])
)*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/
(4*a*B))], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] &&
NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ
[c*A^2 - a*B^2, 0]
```

Rule 1722

```
Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 +
(c_)*(x_)^4]), x_Symbol] := With[{q = Rt[c/a, 2]}, Dist[(A*(c*d + a*e*q)
- a*B*(e + d*q))/(c*d^2 - a*e^2), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] +
Dist[a*(B*d - A*e)*((e + d*q)/(c*d^2 - a*e^2)), Int[(1 + q*x^2)/((d + e*x^2)
)*Sqrt[a + b*x^2 + c*x^4], x], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && N
eQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0]
&& PosQ[c/a] && NeQ[c*A^2 - a*B^2, 0]
```

Rule 1728

```
Int[(P4x_)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4])
, x_Symbol] := With[{q = Rt[c/a, 2], A = Coeff[P4x, x, 0], B = Coeff[P4x, x
, 2], C = Coeff[P4x, x, 4]}, Dist[-C/(e*q), Int[(1 - q*x^2)/Sqrt[a + b*x^2
+ c*x^4], x], x] + Dist[1/(c*e), Int[(A*c*e + a*C*d*q + (B*c*e - C*(c*d - a
*e*q))*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x]] /; FreeQ[{a, b,
c, d, e}, x] && PolyQ[P4x, x^2, 2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*
d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && !GtQ[b^2 - 4*a*c,
0]
```

Rule 1734

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(
p_), x_Symbol] := Int[ExpandIntegrand[1/Sqrt[a + b*x^2 + c*x^4], Px*(d + e
```

```
*x^2)^q*(a + b*x^2 + c*x^4)^(p + 1/2), x], x] /; FreeQ[{a, b, c, d, e}, x]
&& PolyQ[Px, x^2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] &
& IntegerQ[p + 1/2] && IntegerQ[q]
```

Rubi steps

$$\int \frac{A + Bx^2}{(d + ex^2)^2 (a + bx^2 + cx^4)^{3/2}} dx = \int \left(\frac{aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be))}{(cd^2 - bde + ae^2)^2 (a + bx^2 + cx^4)^{3/2}} \right. \\ \left. + \frac{aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be)) - B(cd^2 - ae^2)x^2}{(a + bx^2 + cx^4)^{3/2}} \right) dx \quad (e(Bc^2 - Bcd + Ae^2) - Bcd + Ae^2) \\ = \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be)))}{(cd^2 - bde + ae^2)^2} - \frac{(e(Bc^2 - Bcd + Ae^2) - Bcd + Ae^2)}{(cd^2 - bde + ae^2)^2} \\ = \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be)))}{(cd^2 - bde + ae^2)^2} - \frac{(e(Bc^2 - Bcd + Ae^2) - Bcd + Ae^2)}{(cd^2 - bde + ae^2)^2} \\ = \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be)))}{(cd^2 - bde + ae^2)^2} - \frac{(e(Bc^2 - Bcd + Ae^2) - Bcd + Ae^2)}{(cd^2 - bde + ae^2)^2} \\ = \frac{x(abc(Ae(2cd - be) - B(cd^2 - ae^2)) + (b^2 - 2ac)(aBe(2cd - be) + A(c^2d^2 + b^2e^2 - ce(2bd + ae)) - c(Ae(2cd - be)))}{(cd^2 - bde + ae^2)^2} - \frac{(e(Bc^2 - Bcd + Ae^2) - Bcd + Ae^2)}{(cd^2 - bde + ae^2)^2}$$

Mathematica [C] Result contains complex when optimal does not.

time = 14.94, size = 1116, normalized size = 0.86

Antiderivative was successfully verified.

```
[In] Integrate[(A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x]
```

```
[Out] (4*Sqrt[c/(b + Sqrt[b^2 - 4*a*c])]*d*(-(a*(b^2 - 4*a*c)*e^3*(B*d - A*e)*x*(a + b*x^2 + c*x^4)) + 2*d*x*(d + e*x^2)*(a*B*(-(b^3*e^2) + b^2*c*e*(2*d - e*x^2) + b*c*(3*a*e^2 - c*d*(d - 2*e*x^2)) - 2*c^2*(c*d^2*x^2 + a*e*(2*d - e
```

```

*x^2))) + A*(b^4*e^2 + b^3*c*e*(-2*d + e*x^2) + 2*a*c^2*(a*e^2 - c*d*(d - 2
*e*x^2)) + b^2*c*(-4*a*e^2 + c*d*(d - 2*e*x^2)) + b*c^2*(c*d^2*x^2 - 3*a*e*
(-2*d + e*x^2)))) - I*Sqrt[2]*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b +
Sqrt[b^2 - 4*a*c]]*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(d + e*x^2)
*((-b + Sqrt[b^2 - 4*a*c])*d*(a*B*d*(-4*c^2*d^2 - 3*b^2*e^2 + 4*c*e*(b*d +
2*a*e)) + A*(2*b^3*d*e^2 + 2*b*c*d*(c*d^2 - 3*a*e^2) - 4*a*c*e*(-2*c*d^2 +
a*e^2) + b^2*(-4*c*d^2*e + a*e^3)))*EllipticE[I*ArcSinh[Sqrt[2]*Sqrt[c/(b +
Sqrt[b^2 - 4*a*c]])*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] -
d*(a*B*d*(3*b^2*(b - Sqrt[b^2 - 4*a*c])*e^2 - 4*c^2*d*(Sqrt[b^2 - 4*a*c]*d
- 6*a*e) + 2*c*(-3*b + 2*Sqrt[b^2 - 4*a*c])*e*(b*d + 2*a*e)) + A*(-2*b^4*d
*e^2 + b^2*(-2*c^2*d^3 + a*Sqrt[b^2 - 4*a*c])*e^3 - 4*c*d*e*(Sqrt[b^2 - 4*a*
c]*d - 3*a*e)) - 4*a*c*(-2*c^2*d^3 + a*Sqrt[b^2 - 4*a*c])*e^3 - 2*c*d*e*(Sqr
t[b^2 - 4*a*c]*d - 2*a*e)) + b^3*e*(4*c*d^2 + e*(2*Sqrt[b^2 - 4*a*c]*d - a*
e)) + 2*b*c*(c*d^2*(Sqrt[b^2 - 4*a*c]*d - 8*a*e) + a*e^2*(-3*Sqrt[b^2 - 4*a
*c]*d + 2*a*e)))*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]
)]]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])] - 2*a*(-b^2 + 4*a*c
)*e*(A*e*(7*c*d^2 + e*(-4*b*d + a*e)) + B*(-5*c*d^3 + d*e*(2*b*d + a*e)))*E
llipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b
+ Sqrt[b^2 - 4*a*c]])*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])]
)/(8*a*(b^2 - 4*a*c)*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])*(c*d^3 + d*e*(-(b*d) +
a*e))^2*(d + e*x^2)*Sqrt[a + b*x^2 + c*x^4])

```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 8275 vs. $2(1252) = 2504$.

time = 0.20, size = 8276, normalized size = 6.36

method	result	size
default	Expression too large to display	8276
elliptic	Expression too large to display	9725

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] result too large to display
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(x^2*e + d)^2), x)
```

Fricas [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)/(e*x**2+d)**2/(c*x**4+b*x**2+a)**(3/2),x)

[Out] Timed out

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)/(e*x^2+d)^2/(c*x^4+b*x^2+a)^(3/2),x, algorithm="giac")

[Out] integrate((B*x^2 + A)/((c*x^4 + b*x^2 + a)^(3/2)*(x^2*e + d)^2), x)

Mupad [F]
time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{B x^2 + A}{(e x^2 + d)^2 (c x^4 + b x^2 + a)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)),x)

[Out] int((A + B*x^2)/((d + e*x^2)^2*(a + b*x^2 + c*x^4)^(3/2)), x)

$$3.30 \quad \int \frac{\sqrt{a} + \sqrt{c} x^2}{(d+ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Optimal. Leaf size=273

$$\frac{(\sqrt{c} d - \sqrt{a} e) \tan^{-1} \left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}} \right) (\sqrt{c} d + \sqrt{a} e) (\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}}}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} + \frac{\sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}}}{4\sqrt{a} \sqrt{c}}$$

[Out] $-1/2 \arctan(x(ae^2 - bde + cd^2)^{1/2}/d^{1/2}/e^{1/2}/(cx^4 + bx^2 + a)^{1/2}) * (-ea^{1/2} + dc^{1/2})/d^{1/2}/e^{1/2}/(ae^2 - bde + cd^2)^{1/2} + 1/4 * (\cos(2 \arctan(c^{1/4} x/a^{1/4}))^2)^{1/2} / \cos(2 \arctan(c^{1/4} x/a^{1/4})) * \text{EllipticPi}(\sin(2 \arctan(c^{1/4} x/a^{1/4})), -1/4 * (-ea^{1/2} + dc^{1/2})^2/d/e/a^{1/2}/c^{1/2}, 1/2 * (2 - b/a^{1/2}/c^{1/2})^{1/2}) * (ea^{1/2} + dc^{1/2}) * (a^{1/2} + x^2 c^{1/2}) * ((cx^4 + bx^2 + a)/(a^{1/2} + x^2 c^{1/2}))^{1/2} / a^{1/4} / c^{1/4} / d / e / (cx^4 + bx^2 + a)^{1/2}$

Rubi [A]

time = 0.10, antiderivative size = 273, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 41, $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$, Rules used = {1720}

$$\frac{(\sqrt{a} + \sqrt{c} x^2) \sqrt{\frac{a + bx^2 + cx^4}{(\sqrt{a} + \sqrt{c} x^2)^2}} (\sqrt{a} e + \sqrt{c} d) \Pi \left(-\frac{(\sqrt{c} d - \sqrt{a} e)^2}{4\sqrt{a} \sqrt{c} de}; 2 \text{ArcTan} \left(\frac{\sqrt{c} x}{\sqrt{a}} \right) \middle| \frac{1}{4} \left(2 - \frac{b}{\sqrt{a} \sqrt{c}} \right) \right)}{4\sqrt{a} \sqrt{c} de \sqrt{a + bx^2 + cx^4}} - \frac{(\sqrt{c} d - \sqrt{a} e) \text{ArcTan} \left(\frac{x \sqrt{ae^2 - bde + cd^2}}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}} \right)}{2\sqrt{d} \sqrt{e} \sqrt{ae^2 - bde + cd^2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(\text{Sqrt}[a] + \text{Sqrt}[c]x^2)/((d + ex^2)*\text{Sqrt}[a + bx^2 + cx^4]), x]$

[Out] $-1/2 * ((\text{Sqrt}[c]d - \text{Sqrt}[a]e) * \text{ArcTan}[(\text{Sqrt}[c]d^2 - bde + ae^2)x]/(\text{Sqrt}[d] * \text{Sqrt}[e] * \text{Sqrt}[a + bx^2 + cx^4])) / (\text{Sqrt}[d] * \text{Sqrt}[e] * \text{Sqrt}[c]d^2 - bde + ae^2) + ((\text{Sqrt}[c]d + \text{Sqrt}[a]e) * (\text{Sqrt}[a] + \text{Sqrt}[c]x^2) * \text{Sqrt}[(a + bx^2 + cx^4)/(\text{Sqrt}[a] + \text{Sqrt}[c]x^2)^2]) * \text{EllipticPi}[-1/4 * (\text{Sqrt}[c]d - \text{Sqrt}[a]e)^2 / (\text{Sqrt}[a] * \text{Sqrt}[c]de), 2 * \text{ArcTan}[(c^{1/4}x)/a^{1/4}], (2 - b/(\text{Sqrt}[a] * \text{Sqrt}[c]))/4]) / (4 * a^{1/4} * c^{1/4} * d * e * \text{Sqrt}[a + bx^2 + cx^4])$

Rule 1720

$\text{Int}[(A + Bx)(x^2)/((d + ex^2)*\text{Sqrt}[(a + bx^2 + cx^4)], x_Symbol] :> \text{With}[\{q = \text{Rt}[B/A, 2]\}, \text{Simp}[(-Bd - Ae) * (\text{ArcTan}[\text{Rt}[-b + c(d/e) + a(e/d), 2] * (x/\text{Sqrt}[a + bx^2 + cx^4])]) / (2d * e * \text{Rt}[-b + c(d/e) + a(e/d), 2]))], x] + \text{Simp}[(Bd + Ae) * (A + Bx^2) * (\text{Sqrt}[A^2 * (a + bx^2 + cx^4)/(a(A + Bx^2)^2)]) / (4d * e * A * q * \text{Sqrt}[a + bx^2 + cx^4]) * \text{EllipticPi}[\text{Cancel}[-(Bd - Ae)^2 / (4d * e * A * B)], 2 * \text{ArcTan}[q * x], 1/2 - b/(A/(4 * a * B))], x] /; \text{FreeQ}\{a, b, c, d, e, A, B\}, x] \&\& \text{NeQ}[b^2 - 4 * a * c, 0] \&\&$

NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rubi steps

$$\int \frac{\sqrt{a} + \sqrt{c} x^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx = -\frac{(\sqrt{c} d - \sqrt{a} e) \tan^{-1} \left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}} \right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} + \frac{(\sqrt{c} d + \sqrt{a} e)}{\dots}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.33, size = 310, normalized size = 1.14

$$\frac{i \sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \left(\sqrt{c} d F \left(i \sinh^{-1} \left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right) + (-\sqrt{c} d + \sqrt{a} e) \Pi \left(\frac{(b + \sqrt{b^2 - 4ac})^e}{2cd}; i \sinh^{-1} \left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x \right) \middle| \frac{b + \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}} \right) \right)}{\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} de \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[a] + Sqrt[c]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-1)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(Sqrt[c]*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + ((-Sqrt[c]*d) + Sqrt[a]*e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))]/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])*d*e*Sqrt[a + b*x^2 + c*x^4])

Maple [A]

time = 0.16, size = 365, normalized size = 1.34

method	result
default	$\frac{\sqrt{c} \sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} \operatorname{EllipticF} \left(\frac{x \sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{2a}}}{\dots} \right)}{4e \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}} \sqrt{cx^4 + bx^2 + a}}$

elliptic	$\frac{\sqrt{a(cx^4 + bx^2 + a)} \sqrt{(cx^4 + bx^2 + a)c} (\sqrt{a} + x^2\sqrt{c})}{c\sqrt{2} \sqrt{4 + \frac{2bx^2}{a} - \frac{2x^2\sqrt{-4ac + b^2}}{a}}}$
----------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNV
ERBOSE)`

[Out] $\frac{1}{4}c^{1/2}/e^{2^{1/2}}/((-b+(-4ac+b^2)^{1/2})/a)^{1/2}*(4-2*(-b+(-4ac+b^2)^{1/2})/ax^2)^{1/2}*(4+2*(b+(-4ac+b^2)^{1/2})/ax^2)^{1/2}/(cx^4+bx^2+a)^{1/2}*\text{EllipticF}(1/2*x^2^{1/2}*((-b+(-4ac+b^2)^{1/2})/a)^{1/2}, 1/2*(-4+2*b*(b+(-4ac+b^2)^{1/2})/a/c)^{1/2})+1/e*(ea^{1/2}-d*c^{1/2})/d^{2^{1/2}}/(-b/a+1/a*(-4ac+b^2)^{1/2})^{1/2}*(1+1/2*b*x^2/a-1/2*x^2/a*(-4ac+b^2)^{1/2})^{1/2}*(1+1/2*b*x^2/a+1/2*x^2/a*(-4ac+b^2)^{1/2})^{1/2}/(cx^4+bx^2+a)^{1/2}*\text{EllipticPi}(1/2*x^2^{1/2}*((-b+(-4ac+b^2)^{1/2})/a)^{1/2}, -2/(-b+(-4ac+b^2)^{1/2})*a/e/d, (-1/2*(b+(-4ac+b^2)^{1/2})/a)^{1/2}*2^{1/2}/((-b+(-4ac+b^2)^{1/2})/a)^{1/2})$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")`

[Out] `integrate((sqrt(c)*x^2 + sqrt(a))/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")`

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a} + \sqrt{c} x^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**(1/2)+x**2*c**(1/2))/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)

[Out] Integral((sqrt(a) + sqrt(c)*x**2)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^(1/2)+x^2*c^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{a} + \sqrt{c} x^2}{(ex^2 + d) \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^(1/2) + c^(1/2)*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)),x)

[Out] int((a^(1/2) + c^(1/2)*x^2)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)

$$3.31 \quad \int \frac{1 + \sqrt{\frac{c}{a}} x^2}{(d+ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Optimal. Leaf size=271

$$\frac{\left(\sqrt{\frac{c}{a}} d - e\right) \tan^{-1}\left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}}\right) + \left(\sqrt{\frac{c}{a}} d + e\right) \left(1 + \sqrt{\frac{c}{a}} x^2\right) \sqrt{\frac{a + bx^2 + cx^4}{a \left(1 + \sqrt{\frac{c}{a}} x^2\right)^2}} \Pi}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} + \frac{4\sqrt{\frac{c}{a}} de \sqrt{a}}$$

[Out] $-1/2*\arctan(x*(a*e^2-b*d*e+c*d^2)^(1/2)/d^(1/2)/e^(1/2)/(c*x^4+b*x^2+a)^(1/2))*(-e+d*(c/a)^(1/2))/d^(1/2)/e^(1/2)/(a*e^2-b*d*e+c*d^2)^(1/2)+1/4*(\cos(2*\arctan((c/a)^(1/4)*x))^2)^(1/2)/\cos(2*\arctan((c/a)^(1/4)*x))*\text{EllipticPi}(\sin(2*\arctan((c/a)^(1/4)*x)), -1/4*(-e+d*(c/a)^(1/2))^2/d/e/(c/a)^(1/2), 1/2*(2-b*(c/a)^(1/2)/c)^(1/2))*(e+d*(c/a)^(1/2))*(1+x^2*(c/a)^(1/2))*((c*x^4+b*x^2+a)/a/(1+x^2*(c/a)^(1/2))^2)^(1/2)/(c/a)^(1/4)/d/e/(c*x^4+b*x^2+a)^(1/2)$

Rubi [A]

time = 0.12, antiderivative size = 271, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 41, $\frac{\text{number of rules}}{\text{integrand size}} = 0.024$, Rules used = {1720}

$$\frac{\left(x^2 \sqrt{\frac{c}{a}} + 1\right) \sqrt{\frac{a + bx^2 + cx^4}{a \left(x^2 \sqrt{\frac{c}{a}} + 1\right)^2}} \left(d \sqrt{\frac{c}{a}} + e\right) \Pi\left(-\frac{\left(\sqrt{\frac{c}{a}} d - e\right)^2}{4 \sqrt{\frac{c}{a}} de}; 2 \text{ArcTan}\left(\sqrt{\frac{c}{a}} x\right) \Big|_{\frac{1}{4}} \left(2 - \frac{b \sqrt{\frac{c}{a}}}{c}\right)\right)}{4de \sqrt{\frac{c}{a}} \sqrt{a + bx^2 + cx^4}} - \frac{\left(d \sqrt{\frac{c}{a}} - e\right) \text{ArcTan}\left(\frac{x \sqrt{ae^2 - bde + cd^2}}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}}\right)}{2\sqrt{d} \sqrt{e} \sqrt{ae^2 - bde + cd^2}}$$

Antiderivative was successfully verified.

[In] Int[(1 + Sqrt[c/a]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x]

[Out] $-1/2*((\text{Sqrt}[c/a]*d - e)*\text{ArcTan}[(\text{Sqrt}[c*d^2 - b*d*e + a*e^2]*x)/(\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[a + b*x^2 + c*x^4])]) / (\text{Sqrt}[d]*\text{Sqrt}[e]*\text{Sqrt}[c*d^2 - b*d*e + a*e^2]) + ((\text{Sqrt}[c/a]*d + e)*(1 + \text{Sqrt}[c/a]*x^2)*\text{Sqrt}[(a + b*x^2 + c*x^4)/(a*(1 + \text{Sqrt}[c/a]*x^2)^2)])*\text{EllipticPi}[-1/4*(\text{Sqrt}[c/a]*d - e)^2/(\text{Sqrt}[c/a]*d*e), 2*\text{ArcTan}[(c/a)^(1/4)*x], (2 - (b*\text{Sqrt}[c/a])/c)/4]) / (4*(c/a)^(1/4)*d*e*\text{Sqrt}[a + b*x^2 + c*x^4])$

Rule 1720

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] :> With[{q = Rt[B/A, 2]}, Simp[(- (B*d - A*e)) * (ArcTan[Rt[-b + c*(d/e) + a*(e/d), 2]*(x/Sqrt[a + b*x^2 + c*x^4])]/(2*d*e*Rt[

$-b + c*(d/e) + a*(e/d), 2]))$, x] + Simp[(B*d + A*e)*(A + B*x^2)*(Sqrt[A^2*(a + b*x^2 + c*x^4)/(a*(A + B*x^2)^2)]/(4*d*e*A*q*Sqrt[a + b*x^2 + c*x^4]))*EllipticPi[Cancel[-(B*d - A*e)^2/(4*d*e*A*B)], 2*ArcTan[q*x], 1/2 - b*(A/(4*a*B))], x]] /; FreeQ[{a, b, c, d, e, A, B}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && PosQ[c/a] && EqQ[c*A^2 - a*B^2, 0]

Rubi steps

$$\int \frac{1 + \sqrt{\frac{c}{a}} x^2}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx = -\frac{\left(\sqrt{\frac{c}{a}} d - e\right) \tan^{-1}\left(\frac{\sqrt{cd^2 - bde + ae^2} x}{\sqrt{d} \sqrt{e} \sqrt{a + bx^2 + cx^4}}\right)}{2\sqrt{d} \sqrt{e} \sqrt{cd^2 - bde + ae^2}} + \frac{\left(\sqrt{\frac{c}{a}} d + e\right) \left(1 - \dots\right)}{\dots}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.30, size = 312, normalized size = 1.15

$$\frac{i \sqrt{\frac{b + \sqrt{b^2 - 4ac} + 2cx^2}{b + \sqrt{b^2 - 4ac}}} \sqrt{1 + \frac{2cx^2}{b - \sqrt{b^2 - 4ac}}} \left(\sqrt{\frac{c}{a}} d F\left(i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x\right) \middle| \frac{\pm \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right) + \left(-\sqrt{\frac{c}{a}} d + e\right) \Pi\left(\frac{(b + \sqrt{b^2 - 4ac})^e}{2cd}; i \sinh^{-1}\left(\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} x\right) \middle| \frac{\pm \sqrt{b^2 - 4ac}}{b - \sqrt{b^2 - 4ac}}\right) \right)}{\sqrt{2} \sqrt{\frac{c}{b + \sqrt{b^2 - 4ac}}} de \sqrt{a + bx^2 + cx^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(1 + Sqrt[c/a]*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]),x]

[Out] ((-1)*Sqrt[(b + Sqrt[b^2 - 4*a*c] + 2*c*x^2)/(b + Sqrt[b^2 - 4*a*c]])*Sqrt[1 + (2*c*x^2)/(b - Sqrt[b^2 - 4*a*c]])*(Sqrt[c/a]*d*EllipticF[I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c])) + ((-Sqrt[c/a]*d) + e)*EllipticPi[((b + Sqrt[b^2 - 4*a*c])*e)/(2*c*d), I*ArcSinh[Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])]*x], (b + Sqrt[b^2 - 4*a*c])/(b - Sqrt[b^2 - 4*a*c]))]/(Sqrt[2]*Sqrt[c/(b + Sqrt[b^2 - 4*a*c]])*d*e*Sqrt[a + b*x^2 + c*x^4])

Maple [A]

time = 0.16, size = 369, normalized size = 1.36

method	result
--------	--------

default	$\sqrt{\frac{c}{a}} \sqrt{2} \sqrt{4 - \frac{2(-b + \sqrt{-4ac + b^2})x^2}{a}} \sqrt{4 + \frac{2(b + \sqrt{-4ac + b^2})x^2}{a}} \operatorname{EllipticF} \left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac}}{2a}}}{\sqrt{c x^4 + b x^2 + a}} \right)$
elliptic	$\sqrt{\frac{(c x^4 + b x^2 + a)c}{a}} a \left(1 + x^2 \sqrt{\frac{c}{a}} \right) \left(\frac{\sqrt{2} \sqrt{1 + \frac{b x^2}{2a} - \frac{x^2 \sqrt{-4ac + b^2}}{2a}} \sqrt{1 + \frac{b x^2}{2a} + \frac{x^2 \sqrt{-4ac + b^2}}{2a}} \operatorname{EllipticF} \left(\frac{x\sqrt{2} \sqrt{\frac{-b + \sqrt{-4ac}}{2a}}}{\sqrt{c x^4 + b x^2 + a}} \right)}{4e \sqrt{\frac{-b + \sqrt{-4ac + b^2}}{a}} \sqrt{c x^4 + b x^2 + a}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/4*(c/a)^(1/2)/e*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*(4-2*(-b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)*(4+2*(b+(-4*a*c+b^2)^(1/2))/a*x^2)^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticF(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),1/2*(-4+2*b*(b+(-4*a*c+b^2)^(1/2))/a/c)^(1/2))+1/e*(e-d*(c/a)^(1/2))/d*2^(1/2)/((-b/a+1/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a-1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)*(1+1/2*b*x^2/a+1/2*x^2/a*(-4*a*c+b^2)^(1/2))^(1/2)/(c*x^4+b*x^2+a)^(1/2)*EllipticPi(1/2*x*2^(1/2)*((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2),-2/((-b+(-4*a*c+b^2)^(1/2))*a*e/d,(-1/2*(b+(-4*a*c+b^2)^(1/2))/a)^(1/2)*2^(1/2)/((-b+(-4*a*c+b^2)^(1/2))/a)^(1/2))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((x^2*sqrt(c/a) + 1)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)), x)
```

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="fricas")
```

```
[Out] Timed out
```

Sympy [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

$$\int \frac{x^2 \sqrt{\frac{c}{a} + 1}}{(d + ex^2) \sqrt{a + bx^2 + cx^4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+x**2*(c/a)**(1/2))/(e*x**2+d)/(c*x**4+b*x**2+a)**(1/2),x)
```

```
[Out] Integral((x**2*sqrt(c/a) + 1)/((d + e*x**2)*sqrt(a + b*x**2 + c*x**4)), x)
```

Giac [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+x^2*(c/a)^(1/2))/(e*x^2+d)/(c*x^4+b*x^2+a)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((x^2*sqrt(c/a) + 1)/(sqrt(c*x^4 + b*x^2 + a)*(x^2*e + d)), x)
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{x^2 \sqrt{\frac{c}{a} + 1}}{(ex^2 + d) \sqrt{cx^4 + bx^2 + a}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^2*(c/a)^(1/2) + 1)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)),x)
```

```
[Out] int((x^2*(c/a)^(1/2) + 1)/((d + e*x^2)*(a + b*x^2 + c*x^4)^(1/2)), x)
```

$$3.32 \quad \int \frac{946+315x^2}{(7+5x^2)\sqrt{2+3x^2+x^4}} dx$$

Optimal. Leaf size=106

$$\frac{631(1+x^2)\sqrt{\frac{2+x^2}{1+x^2}}F(\tan^{-1}(x)|\frac{1}{2})}{2\sqrt{2}\sqrt{2+3x^2+x^4}} - \frac{2525(2+x^2)\Pi(\frac{2}{7};\tan^{-1}(x)|\frac{1}{2})}{14\sqrt{2}\sqrt{\frac{2+x^2}{1+x^2}}\sqrt{2+3x^2+x^4}}$$

[Out] $-2525/28*(x^2+2)*(1/(x^2+1))^{(1/2)}*(x^2+1)^{(1/2)}*\text{EllipticPi}(x/(x^2+1)^{(1/2)}, 2/7, 1/2*2^{(1/2)})*2^{(1/2)}/((x^2+2)/(x^2+1))^{(1/2)}/(x^4+3*x^2+2)^{(1/2)}+631/4*(x^2+1)^{(3/2)}*(1/(x^2+1))^{(1/2)}*\text{EllipticF}(x/(x^2+1)^{(1/2)}, 1/2*2^{(1/2)})*((x^2+2)/(x^2+1))^{(1/2)}*2^{(1/2)}/(x^4+3*x^2+2)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.129$, Rules used = {1718, 1113, 1470, 553}

$$\frac{631(x^2+1)\sqrt{\frac{x^2+2}{x^2+1}}F(\text{ArcTan}(x)|\frac{1}{2})}{2\sqrt{2}\sqrt{x^4+3x^2+2}} - \frac{2525(x^2+2)\Pi(\frac{2}{7};\text{ArcTan}(x)|\frac{1}{2})}{14\sqrt{2}\sqrt{\frac{x^2+2}{x^2+1}}\sqrt{x^4+3x^2+2}}$$

Antiderivative was successfully verified.

[In] Int[(946 + 315*x^2)/((7 + 5*x^2)*Sqrt[2 + 3*x^2 + x^4]),x]

[Out] $(631*(1+x^2)*\text{Sqrt}[(2+x^2)/(1+x^2)]*\text{EllipticF}[\text{ArcTan}[x], 1/2])/(2*\text{Sqrt}[2]*\text{Sqrt}[2+3*x^2+x^4]) - (2525*(2+x^2)*\text{EllipticPi}[2/7, \text{ArcTan}[x], 1/2])/(14*\text{Sqrt}[2]*\text{Sqrt}[(2+x^2)/(1+x^2)]*\text{Sqrt}[2+3*x^2+x^4])$

Rule 553

Int[Sqrt[(c_) + (d_)*(x_)^2]/(((a_) + (b_)*(x_)^2)*Sqrt[(e_) + (f_)*(x_)^2]), x_Symbol] :> Simp[c*(Sqrt[e + f*x^2]/(a*e*Rt[d/c, 2]*Sqrt[c + d*x^2]*Sqrt[c*((e + f*x^2)/(e*(c + d*x^2))])))*EllipticPi[1 - b*(c/(a*d)), ArcTan[Rt[d/c, 2]*x], 1 - c*(f/(d*e))], x] /; FreeQ[{a, b, c, d, e, f}, x] && PosQ[d/c]

Rule 1113

Int[1/Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4], x_Symbol] :> With[{q = Rt[b^2 - 4*a*c, 2]}, Simp[(2*a + (b + q)*x^2)*(Sqrt[(2*a + (b - q)*x^2)/(2*a + (b + q)*x^2)]/(2*a*Rt[(b + q)/(2*a), 2]*Sqrt[a + b*x^2 + c*x^4])))*EllipticF[ArcTan[Rt[(b + q)/(2*a), 2]*x], 2*(q/(b + q))], x] /; PosQ[(b + q)/a] &&

!(PosQ[(b - q)/a] && SimplerSqrtQ[(b - q)/(2*a), (b + q)/(2*a)])] /; FreeQ[{a, b, c}, x] && GtQ[b^2 - 4*a*c, 0]

Rule 1470

Int[((d_) + (e_)*(x_)^(n_))^(q_)*((f_) + (g_)*(x_)^(n_))^(r_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Dist[(a + b*x^n + c*x^(2*n))^FracPart[p]/((d + e*x^n)^FracPart[p]*(a/d + (c*x^n)/e)^FracPart[p]), Int[(d + e*x^n)^(p + q)*(f + g*x^n)^r*(a/d + (c/e)*x^n)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p, q, r}, x] && EqQ[n2, 2*n] && NeQ[b^2 - 4*a*c, 0] && EqQ[c*d^2 - b*d*e + a*e^2, 0] && !IntegerQ[p]

Rule 1718

Int[((A_) + (B_)*(x_)^2)/(((d_) + (e_)*(x_)^2)*Sqrt[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]), x_Symbol] := With[{q = Sqrt[b^2 - 4*a*c]}, Dist[(2*a*B - A*(b + q))/(2*a*e - d*(b + q)), Int[1/Sqrt[a + b*x^2 + c*x^4], x], x] - Dist[(B*d - A*e)/(2*a*e - d*(b + q)), Int[(2*a + (b + q)*x^2)/((d + e*x^2)*Sqrt[a + b*x^2 + c*x^4]), x], x] /; RationalQ[q] /; FreeQ[{a, b, c, d, e, A, B}, x] && GtQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NeQ[c*A^2 - b*A*B + a*B^2, 0]

Rubi steps

$$\begin{aligned} \int \frac{946 + 315x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx &= \frac{631}{2} \int \frac{1}{\sqrt{2 + 3x^2 + x^4}} dx - \frac{2525}{8} \int \frac{4 + 4x^2}{(7 + 5x^2)\sqrt{2 + 3x^2 + x^4}} dx \\ &= \frac{631(1 + x^2) \sqrt{\frac{2 + x^2}{1 + x^2}} F(\tan^{-1}(x) | \frac{1}{2})}{2\sqrt{2} \sqrt{2 + 3x^2 + x^4}} - \frac{\left(2525 \sqrt{\frac{1}{2} + \frac{x^2}{4}} \sqrt{4 + 4x^2}\right) f}{8\sqrt{2 + 3x^2 + x^4}} \\ &= \frac{631(1 + x^2) \sqrt{\frac{2 + x^2}{1 + x^2}} F(\tan^{-1}(x) | \frac{1}{2})}{2\sqrt{2} \sqrt{2 + 3x^2 + x^4}} - \frac{2525(2 + x^2) \Pi\left(\frac{2}{7}; \tan^{-1}(x) | \frac{1}{2}\right)}{14\sqrt{2} \sqrt{\frac{2 + x^2}{1 + x^2}} \sqrt{2 + 3x^2 + x^4}} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 10.14, size = 74, normalized size = 0.70

$$\frac{i\sqrt{1+x^2}\sqrt{2+x^2}\left(441F\left(i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right)+505\Pi\left(\frac{10}{7};i\sinh^{-1}\left(\frac{x}{\sqrt{2}}\right)\middle|2\right)\right)}{7\sqrt{2+3x^2+x^4}}$$

Antiderivative was successfully verified.

[In] Integrate[(946 + 315*x^2)/((7 + 5*x^2)*Sqrt[2 + 3*x^2 + x^4]),x]

[Out] $\left(\frac{-1}{7}i\right)\sqrt{1+x^2}\sqrt{2+x^2}\left(441\text{EllipticF}\left[\frac{i\sqrt{2}x}{\sqrt{2}},2\right]+505\text{EllipticPi}\left[\frac{10}{7},\frac{i\sqrt{2}x}{\sqrt{2}},2\right]\right)\sqrt{2+3x^2+x^4}$

Maple [C] Result contains complex when optimal does not.

time = 0.14, size = 93, normalized size = 0.88

method	result
default	$-\frac{63i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\text{EllipticF}\left(\frac{i\sqrt{2}x}{\sqrt{2}},\sqrt{2}\right)}{2\sqrt{x^4+3x^2+2}}-\frac{505i\sqrt{2}\sqrt{1+\frac{x^2}{2}}\sqrt{x^2+1}\text{EllipticPi}\left(\frac{i\sqrt{2}x}{\sqrt{2}},\frac{10}{7},2\right)}{7\sqrt{x^4+3x^2+2}}$
elliptic	$-\frac{63i\sqrt{2}\sqrt{2x^2+4}\sqrt{x^2+1}\text{EllipticF}\left(\frac{i\sqrt{2}x}{\sqrt{2}},\sqrt{2}\right)}{2\sqrt{x^4+3x^2+2}}-\frac{505i\sqrt{2}\sqrt{1+\frac{x^2}{2}}\sqrt{x^2+1}\text{EllipticPi}\left(\frac{i\sqrt{2}x}{\sqrt{2}},\frac{10}{7},2\right)}{7\sqrt{x^4+3x^2+2}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x,method=_RETURNVERBOSE)

[Out] $-63/2i2^{(1/2)}(2x^2+4)^{(1/2)}(x^2+1)^{(1/2)}/(x^4+3x^2+2)^{(1/2)}\text{EllipticF}\left(\frac{1}{2}i2^{(1/2)}x,2^{(1/2)}\right)-505/7i2^{(1/2)}(1+1/2x^2)^{(1/2)}(x^2+1)^{(1/2)}/(x^4+3x^2+2)^{(1/2)}\text{EllipticPi}\left(\frac{1}{2}i2^{(1/2)}x,10/7,2^{(1/2)}\right)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="maxima")

[Out] integrate((315*x^2 + 946)/(sqrt(x^4 + 3*x^2 + 2)*(5*x^2 + 7)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(x^4 + 3*x^2 + 2)*(315*x^2 + 946)/(5*x^6 + 22*x^4 + 31*x^2 + 14), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{315x^2 + 946}{\sqrt{(x^2 + 1)(x^2 + 2)}(5x^2 + 7)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((315*x**2+946)/(5*x**2+7)/(x**4+3*x**2+2)**(1/2),x)

[Out] Integral((315*x**2 + 946)/(sqrt((x**2 + 1)*(x**2 + 2))*(5*x**2 + 7)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((315*x^2+946)/(5*x^2+7)/(x^4+3*x^2+2)^(1/2),x, algorithm="giac")

[Out] integrate((315*x^2 + 946)/(sqrt(x^4 + 3*x^2 + 2)*(5*x^2 + 7)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{315x^2 + 946}{(5x^2 + 7)\sqrt{x^4 + 3x^2 + 2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((315*x^2 + 946)/((5*x^2 + 7)*(3*x^2 + x^4 + 2)^(1/2)),x)

[Out] int((315*x^2 + 946)/((5*x^2 + 7)*(3*x^2 + x^4 + 2)^(1/2)), x)

$$3.33 \quad \int \frac{(A+Bx^2)(d+ex^2)^q}{a+bx^2+cx^4} dx$$

Optimal. Leaf size=218

$$\frac{\left(B - \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) x(d+ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b-\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{b - \sqrt{b^2-4ac}} + \frac{\left(B + \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) x(d+ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b+\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{\sqrt{b^2-4ac} + b}$$

[Out] $x*(e*x^2+d)^q*AppellF1(1/2,1,-q,3/2,-2*c*x^2/(b-(-4*a*c+b^2)^(1/2)),-e*x^2/d)*(B+(2*A*c-B*b)/(-4*a*c+b^2)^(1/2))/((1+e*x^2/d)^q)/(b-(-4*a*c+b^2)^(1/2))+x*(e*x^2+d)^q*AppellF1(1/2,1,-q,3/2,-2*c*x^2/(b+(-4*a*c+b^2)^(1/2)),-e*x^2/d)*(B+(-2*A*c+B*b)/(-4*a*c+b^2)^(1/2))/((1+e*x^2/d)^q)/(b+(-4*a*c+b^2)^(1/2))$

Rubi [A]

time = 0.23, antiderivative size = 218, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 31, $\frac{\text{number of rules}}{\text{integrand size}} = 0.097$, Rules used = {1706, 441, 440}

$$\frac{x(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} \left(B - \frac{bB-2Ac}{\sqrt{b^2-4ac}}\right) F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b-\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{b - \sqrt{b^2-4ac}} + \frac{x(d+ex^2)^q \left(\frac{ex^2}{d} + 1\right)^{-q} \left(\frac{bB-2Ac}{\sqrt{b^2-4ac}} + B\right) F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b+\sqrt{b^2-4ac}}, -\frac{ex^2}{d}\right)}{\sqrt{b^2-4ac} + b}$$

Antiderivative was successfully verified.

[In] Int[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]

[Out] $((B - (b*B - 2*A*c)/\text{Sqrt}[b^2 - 4*a*c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (-2*c*x^2)/(b - \text{Sqrt}[b^2 - 4*a*c]), -(e*x^2/d)])/((b - \text{Sqrt}[b^2 - 4*a*c])*(1 + (e*x^2/d)^q) + ((B + (b*B - 2*A*c)/\text{Sqrt}[b^2 - 4*a*c])*x*(d + e*x^2)^q*AppellF1[1/2, 1, -q, 3/2, (-2*c*x^2)/(b + \text{Sqrt}[b^2 - 4*a*c]), -(e*x^2/d)])/((b + \text{Sqrt}[b^2 - 4*a*c])*(1 + (e*x^2/d)^q))$

Rule 440

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Simp[a^p*c^q*x*AppellF1[1/n, -p, -q, 1 + 1/n, (-b)*(x^n/a), (-d)*(x^n/c)], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && (IntegerQ[p] || GtQ[a, 0]) && (IntegerQ[q] || GtQ[c, 0])

Rule 441

Int[((a_) + (b_.)*(x_)^(n_))^(p_)*((c_) + (d_.)*(x_)^(n_))^(q_), x_Symbol] :> Dist[a^IntPart[p]*((a + b*x^n)^FracPart[p]/(1 + b*(x^n/a))^FracPart[p]), Int[(1 + b*(x^n/a))^p*(c + d*x^n)^q, x], x] /; FreeQ[{a, b, c, d, n, p, q}, x] && NeQ[b*c - a*d, 0] && NeQ[n, -1] && !(IntegerQ[p] || GtQ[a, 0])

Rule 1706

```
Int[(Px_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] :> Int[ExpandIntegrand[Px*(d + e*x^2)^q*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, e, q}, x] && PolyQ[Px, x^2] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && IntegerQ[p]
```

Rubi steps

$$\begin{aligned} \int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx &= \int \left(\frac{\left(B + \frac{-bB+2Ac}{\sqrt{b^2-4ac}}\right)(d + ex^2)^q}{b - \sqrt{b^2-4ac} + 2cx^2} + \frac{\left(B - \frac{-bB+2Ac}{\sqrt{b^2-4ac}}\right)(d + ex^2)^q}{b + \sqrt{b^2-4ac} + 2cx^2} \right) dx \\ &= \left(B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right) \int \frac{(d + ex^2)^q}{b - \sqrt{b^2 - 4ac} + 2cx^2} dx + \left(B + \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right) \int \frac{(d + ex^2)^q}{b + \sqrt{b^2 - 4ac} + 2cx^2} dx \\ &= \left(\left(B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right) (d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} \right) \int \frac{\left(1 + \frac{ex^2}{d}\right)^q}{b - \sqrt{b^2 - 4ac} + 2cx^2} dx \\ &= \frac{\left(B - \frac{bB - 2Ac}{\sqrt{b^2 - 4ac}}\right) x(d + ex^2)^q \left(1 + \frac{ex^2}{d}\right)^{-q} F_1\left(\frac{1}{2}; 1, -q; \frac{3}{2}; -\frac{2cx^2}{b - \sqrt{b^2 - 4ac}}\right)}{b - \sqrt{b^2 - 4ac}} \end{aligned}$$

Mathematica [F]

time = 0.43, size = 0, normalized size = 0.00

$$\int \frac{(A + Bx^2)(d + ex^2)^q}{a + bx^2 + cx^4} dx$$

Verification is not applicable to the result.

[In] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]

[Out] Integrate[((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x]

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a), x)

[Out] int((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="maxima")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^q/(c*x^4 + b*x^2 + a), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="fricas")

[Out] integral((B*x^2 + A)*(x^2*e + d)^q/(c*x^4 + b*x^2 + a), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x**2+A)*(e*x**2+d)**q/(c*x**4+b*x**2+a),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((B*x^2+A)*(e*x^2+d)^q/(c*x^4+b*x^2+a),x, algorithm="giac")

[Out] integrate((B*x^2 + A)*(x^2*e + d)^q/(c*x^4 + b*x^2 + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(Bx^2 + A)(ex^2 + d)^q}{cx^4 + bx^2 + a} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4),x)

[Out] int(((A + B*x^2)*(d + e*x^2)^q)/(a + b*x^2 + c*x^4), x)

$$3.34 \quad \int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx$$

Optimal. Leaf size=106

$$-\frac{1}{2} \tan^{-1}(\sqrt{3} - 2\sqrt{1+x^2}) + \frac{1}{2} \tan^{-1}(\sqrt{3} + 2\sqrt{1+x^2}) + \frac{1}{4}\sqrt{3} \log(2+x^2 - \sqrt{3}\sqrt{1+x^2}) - \frac{1}{4}\sqrt{3} \log$$

[Out] 1/2*arctan(-3^(1/2)+2*(x^2+1)^(1/2))+1/2*arctan(3^(1/2)+2*(x^2+1)^(1/2))+1/4*ln(2+x^2-3^(1/2)*(x^2+1)^(1/2))*3^(1/2)-1/4*ln(2+x^2+3^(1/2)*(x^2+1)^(1/2))*3^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 7, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {1699, 840, 1183, 648, 632, 210, 642}

$$-\frac{1}{2}\text{ArcTan}(\sqrt{3} - 2\sqrt{x^2+1}) + \frac{1}{2}\text{ArcTan}(2\sqrt{x^2+1} + \sqrt{3}) + \frac{1}{4}\sqrt{3} \log(x^2 - \sqrt{3}\sqrt{x^2+1} + 2) - \frac{1}{4}\sqrt{3} \log(x^2 + \sqrt{3}\sqrt{x^2+1} + 2)$$

Antiderivative was successfully verified.

[In] Int[(x*(1 + 2*x^2))/(Sqrt[1 + x^2]*(1 + x^2 + x^4)),x]

[Out] -1/2*ArcTan[Sqrt[3] - 2*Sqrt[1 + x^2]] + ArcTan[Sqrt[3] + 2*Sqrt[1 + x^2]]/2 + (Sqrt[3]*Log[2 + x^2 - Sqrt[3]*Sqrt[1 + x^2]])/4 - (Sqrt[3]*Log[2 + x^2 + Sqrt[3]*Sqrt[1 + x^2]])/4

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 632

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In

`t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]`

Rule 840

`Int[((f_) + (g_)*(x_))/(Sqrt[(d_) + (e_)*(x_)]*((a_) + (b_)*(x_) + (c_)*(x_)^2)), x_Symbol] := Dist[2, Subst[Int[(e*f - d*g + g*x^2)/(c*d^2 - b*d*e + a*e^2 - (2*c*d - b*e)*x^2 + c*x^4), x], x, Sqrt[d + e*x]], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0]`

Rule 1183

`Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]`

Rule 1699

`Int[(Px_)*(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Dist[1/2, Subst[Int[(Px /. x -> Sqrt[x])*(d + e*x)^q*(a + b*x + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, b, c, d, e, p, q}, x] && PolyQ[Px, x^2]`

Rubi steps

$$\begin{aligned}
 \int \frac{x(1+2x^2)}{\sqrt{1+x^2}(1+x^2+x^4)} dx &= \frac{1}{2} \text{Subst} \left(\int \frac{1+2x}{\sqrt{1+x}(1+x+x^2)} dx, x, x^2 \right) \\
 &= \text{Subst} \left(\int \frac{-1+2x^2}{1-x^2+x^4} dx, x, \sqrt{1+x^2} \right) \\
 &= \frac{\text{Subst} \left(\int \frac{-\sqrt{3}+3x}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right)}{2\sqrt{3}} + \frac{\text{Subst} \left(\int \frac{-\sqrt{3}-3x}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right)}{2\sqrt{3}} \\
 &= \frac{1}{4} \text{Subst} \left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right) + \frac{1}{4} \text{Subst} \left(\int \frac{1}{1+\sqrt{3}x+x^2} dx, x, \sqrt{1+x^2} \right) \\
 &= \frac{1}{4} \sqrt{3} \log \left(2+x^2-\sqrt{3}\sqrt{1+x^2} \right) - \frac{1}{4} \sqrt{3} \log \left(2+x^2+\sqrt{3}\sqrt{1+x^2} \right) \\
 &= -\frac{1}{2} \tan^{-1} \left(\sqrt{3}-2\sqrt{1+x^2} \right) + \frac{1}{2} \tan^{-1} \left(\sqrt{3}+2\sqrt{1+x^2} \right) + \frac{1}{4} \sqrt{3} \log \left(\frac{2+x^2-\sqrt{3}\sqrt{1+x^2}}{2+x^2+\sqrt{3}\sqrt{1+x^2}} \right)
 \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.09, size = 81, normalized size = 0.76

$$\frac{1}{2}(1 - i\sqrt{3}) \tan^{-1} \left(\frac{1}{2}(1 - i\sqrt{3}) \sqrt{1+x^2} \right) + \frac{1}{2}(1 + i\sqrt{3}) \tan^{-1} \left(\frac{1}{2}(1 + i\sqrt{3}) \sqrt{1+x^2} \right)$$

Antiderivative was successfully verified.

[In] Integrate[(x*(1 + 2*x^2))/(Sqrt[1 + x^2]*(1 + x^2 + x^4)),x]

[Out] ((1 - I*Sqrt[3])*ArcTan[((1 - I*Sqrt[3])*Sqrt[1 + x^2])/2])/2 + ((1 + I*Sqrt[3])*ArcTan[((1 + I*Sqrt[3])*Sqrt[1 + x^2])/2])/2

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 295 vs. 2(80) = 160.

time = 0.37, size = 296, normalized size = 2.79

method	result
default	$\frac{\sqrt{2} \sqrt{\frac{2(1+x)^2}{(1-x)^2} + 2} \left(\sqrt{3} \operatorname{arctanh} \left(\frac{\sqrt{\frac{2(1+x)^2}{(1-x)^2} + 2} \sqrt{3}}{2} \right) + \operatorname{arctan} \left(\frac{\sqrt{\frac{2(1+x)^2}{(1-x)^2} + 2} (1+x)}{\left(\frac{(1+x)^2}{(1-x)^2} + 1 \right) (1-x)} \right) \right)}{4 \sqrt{\frac{(1+x)^2}{(1-x)^2} + 1} \left(1 + \frac{1+x}{1-x} \right)}$
trager	$-4 \ln \left(\frac{16 \operatorname{RootOf}(16_Z^4 - 4_Z^2 + 1)^5 x^2 - 16 \operatorname{RootOf}(16_Z^4 - 4_Z^2 + 1)^3 x^2 - 12 \operatorname{RootOf}(16_Z^4 - 4_Z^2 + 1)^3 + 4 \operatorname{RootOf}(16_Z^4 - 4_Z^2 + 1)}{(4 \operatorname{RootOf}(16_Z^4 - 4_Z^2 + 1)^2 x - x + 1) (4 \operatorname{RootOf}(16_Z^4 - 4_Z^2 + 1)^2 x - x + 1)} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/4*2^(1/2)*(2*(1+x)^2/(1-x)^2+2)^(1/2)*(3^(1/2)*arctanh(1/2*(2*(1+x)^2/(1-x)^2+2)^(1/2)*3^(1/2))+arctan(1/((1+x)^2/(1-x)^2+1)*(2*(1+x)^2/(1-x)^2+2)^(1/2)*(1+x)/(1-x)))/(((1+x)^2/(1-x)^2+1)/(1+(1+x)/(1-x)))^(1/2)/(1+(1+x)/(1-x))-1/4*2^(1/2)*(2*(-1+x)^2/(-1-x)^2+2)^(1/2)*(3^(1/2)*arctanh(1/2*(2*(-1+x)^2/(-1-x)^2+2)^(1/2)*3^(1/2))+arctan(1/((-1+x)^2/(-1-x)^2+1)*(2*(-1+x)^2/(-1-x)^2+2)^(1/2)*(-1+x)/(-1-x)))/((-1+x)^2/(-1-x)^2+1)/(1+(-1+x)/(-1-x)))^(1/2)/(1+(-1+x)/(-1-x))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate((2*x^2 + 1)*x/((x^4 + x^2 + 1)*sqrt(x^2 + 1)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 279 vs. 2(80) = 160.

time = 0.41, size = 279, normalized size = 2.63

$$\frac{1}{4}\sqrt{3}\log(32x^4+80x^2+32)\sqrt{x^2+1}-16(2x^3+\sqrt{3})\sqrt{x^2+1}+\frac{1}{4}\sqrt{3}\log(32x^4+80x^2+32)\sqrt{x^2+1}-\frac{1}{4}\sqrt{3}\log(32x^4+80x^2+32)\sqrt{x^2+1}-\arctan\left(\frac{\sqrt{2x^4+5x^2+2}\sqrt{x^2+1}}{x+\sqrt{x^2+1}}\right)+\sqrt{3}-2\sqrt{x^2+1}-\arctan\left(\frac{\sqrt{2x^4+5x^2+2}\sqrt{x^2+1}}{x+\sqrt{x^2+1}}\right)-\sqrt{3}-2\sqrt{x^2+1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="fricas")

[Out] 1/4*sqrt(3)*log(32*x^4 + 80*x^2 + 32*sqrt(3)*(x^3 + x) - 16*(2*x^3 + sqrt(3))*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 32) - 1/4*sqrt(3)*log(32*x^4 + 80*x^2 - 32*sqrt(3)*(x^3 + x) - 16*(2*x^3 - sqrt(3))*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 32) - arctan(2*sqrt(2*x^4 + 5*x^2 + 2*sqrt(3)*(x^3 + x) - (2*x^3 + sqrt(3))*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 2)*(x + sqrt(x^2 + 1)) + sqrt(3) - 2*sqrt(x^2 + 1)) - arctan(2*sqrt(2*x^4 + 5*x^2 - 2*sqrt(3)*(x^3 + x) - (2*x^3 - sqrt(3))*(2*x^2 + 1) + 4*x)*sqrt(x^2 + 1) + 2)*(x + sqrt(x^2 + 1)) - sqrt(3) - 2*sqrt(x^2 + 1))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x(2x^2 + 1)}{\sqrt{x^2 + 1} (x^2 - x + 1) (x^2 + x + 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x**2+1)/(x**4+x**2+1)/(x**2+1)**(1/2),x)

[Out] Integral(x*(2*x**2 + 1)/(sqrt(x**2 + 1)*(x**2 - x + 1)*(x**2 + x + 1)), x)

Giac [A]

time = 4.11, size = 80, normalized size = 0.75

$$-\frac{1}{4}\sqrt{3}\log\left(x^2+\sqrt{3}\sqrt{x^2+1}+2\right)+\frac{1}{4}\sqrt{3}\log\left(x^2-\sqrt{3}\sqrt{x^2+1}+2\right)+\frac{1}{2}\arctan\left(\sqrt{3}+2\sqrt{x^2+1}\right)+\frac{1}{2}\arctan\left(-\sqrt{3}+2\sqrt{x^2+1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x*(2*x^2+1)/(x^4+x^2+1)/(x^2+1)^(1/2),x, algorithm="giac")

[Out] -1/4*sqrt(3)*log(x^2 + sqrt(3)*sqrt(x^2 + 1) + 2) + 1/4*sqrt(3)*log(x^2 - sqrt(3)*sqrt(x^2 + 1) + 2) + 1/2*arctan(sqrt(3) + 2*sqrt(x^2 + 1)) + 1/2*arctan(-sqrt(3) + 2*sqrt(x^2 + 1))

Mupad [B]

time = 1.85, size = 397, normalized size = 3.75

$$\frac{\ln\left(\frac{(x-1)\sqrt{3x^2+3}\sqrt{x^2+1}+\sqrt{3x^2+3}}{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}\right)}{\sqrt{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}}-\frac{\ln\left(\frac{(x+1)\sqrt{3x^2+3}\sqrt{x^2+1}+\sqrt{3x^2+3}}{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}\right)}{\sqrt{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}}-\frac{\ln\left(\frac{(x-1)\sqrt{3x^2+3}\sqrt{x^2+1}+\sqrt{3x^2+3}}{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}\right)}{\sqrt{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}}-\frac{\ln\left(\frac{(x+1)\sqrt{3x^2+3}\sqrt{x^2+1}+\sqrt{3x^2+3}}{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}\right)}{\sqrt{\left(\frac{1}{2}\sqrt{3x^2+3}\right)^2+\sqrt{3x^2+3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((x*(2*x^2 + 1))/((x^2 + 1)^{(1/2)}*(x^2 + x^4 + 1)),x)$

[Out] $((\log(x - (3^{(1/2)}*1i)/2 - 1/2) - \log(x/2 + (3^{(1/2)}/2 + 1i/2)*(x^2 + 1)^{(1/2)} + (3^{(1/2)}*x*1i)/2 + 1))*((3^{(1/2)}*1i)/2 + 2*((3^{(1/2)}*1i)/2 + 1/2)^3 + 1/2))/((((3^{(1/2)}*1i)/2 + 1/2)^2 + 1)^{(1/2)}*(3^{(1/2)}*1i + 4*((3^{(1/2)}*1i)/2 + 1/2)^3 + 1)) + ((\log(x - (3^{(1/2)}*1i)/2 + 1/2) - \log((3^{(1/2)}/2 - 1i/2)*(x^2 + 1)^{(1/2)} - x/2 + (3^{(1/2)}*x*1i)/2 + 1))*((3^{(1/2)}*1i)/2 + 2*((3^{(1/2)}*1i)/2 - 1/2)^3 - 1/2))/((((3^{(1/2)}*1i)/2 - 1/2)^2 + 1)^{(1/2)}*(3^{(1/2)}*1i + 4*((3^{(1/2)}*1i)/2 - 1/2)^3 - 1)) + ((\log(x + (3^{(1/2)}*1i)/2 - 1/2) - \log(x/2 + (3^{(1/2)}/2 - 1i/2)*(x^2 + 1)^{(1/2)} - (3^{(1/2)}*x*1i)/2 + 1))*((3^{(1/2)}*1i)/2 + 2*((3^{(1/2)}*1i)/2 - 1/2)^3 - 1/2))/((((3^{(1/2)}*1i)/2 - 1/2)^2 + 1)^{(1/2)}*(3^{(1/2)}*1i + 4*((3^{(1/2)}*1i)/2 - 1/2)^3 - 1)) + ((\log(x + (3^{(1/2)}*1i)/2 + 1/2) - \log((3^{(1/2)}/2 + 1i/2)*(x^2 + 1)^{(1/2)} - x/2 - (3^{(1/2)}*x*1i)/2 + 1))*((3^{(1/2)}*1i)/2 + 2*((3^{(1/2)}*1i)/2 + 1/2)^3 + 1/2))/((((3^{(1/2)}*1i)/2 + 1/2)^2 + 1)^{(1/2)}*(3^{(1/2)}*1i + 4*((3^{(1/2)}*1i)/2 + 1/2)^3 + 1))$

$$3.35 \quad \int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cd x^4} dx$$

Optimal. Leaf size=145

$$-\frac{\sqrt{b - 2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{\sqrt{b - 2\sqrt{a}\sqrt{c}} x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} + \frac{\sqrt{b + 2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{\sqrt{b + 2\sqrt{a}\sqrt{c}} x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

[Out] $-1/4*\operatorname{arctanh}(x*(b-2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})*(b-2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/d/a^{(1/2)}/c^{(1/2)}+1/4*\operatorname{arctanh}(x*(b+2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/(c*x^4+b*x^2+a)^{(1/2)})*(b+2*a^{(1/2)}*c^{(1/2)})^{(1/2)}/d/a^{(1/2)}/c^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 145, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2096, 1107, 214}

$$\frac{\sqrt{2\sqrt{a}\sqrt{c} + b} \tanh^{-1}\left(\frac{x\sqrt{2\sqrt{a}\sqrt{c} + b}}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} - \frac{\sqrt{b - 2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{x\sqrt{b - 2\sqrt{a}\sqrt{c}}}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sqrt}[a + b*x^2 + c*x^4]/(a*d - c*d*x^4), x]$

[Out] $-1/4*(\operatorname{Sqrt}[b - 2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b - 2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]]*x)/\operatorname{Sqrt}[a + b*x^2 + c*x^4]])/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]*d) + (\operatorname{Sqrt}[b + 2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[b + 2*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]]*x)/\operatorname{Sqrt}[a + b*x^2 + c*x^4]])/(4*\operatorname{Sqrt}[a]*\operatorname{Sqrt}[c]*d)$

Rule 214

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b]$

Rule 1107

$\operatorname{Int}[(a_ + (b_)*(x_)^2 + (c_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Rt}[b^2 - 4*a*c, 2]\}, \operatorname{Dist}[c/q, \operatorname{Int}[1/(b/2 - q/2 + c*x^2), x], x] - \operatorname{Dist}[c/q, \operatorname{Int}[1/(b/2 + q/2 + c*x^2), x], x]] /; \operatorname{FreeQ}\{a, b, c\}, x \ \&\& \operatorname{NeQ}[b^2 - 4*a*c, 0] \ \&\& \operatorname{PosQ}[b^2 - 4*a*c]$

Rule 2096

```
Int[Sqrt[v_]/((d_) + (e_)*(x_)^4), x_Symbol] := With[{a = Coeff[v, x, 0],
b = Coeff[v, x, 2], c = Coeff[v, x, 4]}, Dist[a/d, Subst[Int[1/(1 - 2*b*x^2
+ (b^2 - 4*a*c)*x^4), x], x, x/Sqrt[v]], x] /; EqQ[c*d + a*e, 0] && PosQ[a
*c]] /; FreeQ[{d, e}, x] && PolyQ[v, x^2, 2]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{a + bx^2 + cx^4}}{ad - cdx^4} dx &= \frac{\text{Subst}\left(\int \frac{1}{1-2bx^2+(b^2-4ac)x^4} dx, x, \frac{x}{\sqrt{a + bx^2 + cx^4}}\right)}{d} \\ &= \frac{(b^2 - 4ac) \text{Subst}\left(\int \frac{1}{-b-2\sqrt{a}\sqrt{c} + (b^2-4ac)x^2} dx, x, \frac{x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} - \frac{(b^2 - 4ac) \text{Subst}\left(\int \frac{1}{-b+2\sqrt{a}\sqrt{c} + (b^2-4ac)x^2} dx, x, \frac{x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} \\ &= -\frac{\sqrt{b - 2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{\sqrt{b - 2\sqrt{a}\sqrt{c}} x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} + \frac{\sqrt{b + 2\sqrt{a}\sqrt{c}} \tanh^{-1}\left(\frac{\sqrt{b + 2\sqrt{a}\sqrt{c}} x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d} \end{aligned}$$

Mathematica [A]

time = 0.54, size = 139, normalized size = 0.96

$$\frac{-\sqrt{-b - 2\sqrt{a}\sqrt{c}} \tan^{-1}\left(\frac{\sqrt{-b - 2\sqrt{a}\sqrt{c}} x}{\sqrt{a + bx^2 + cx^4}}\right) + \sqrt{-b + 2\sqrt{a}\sqrt{c}} \tan^{-1}\left(\frac{\sqrt{-b + 2\sqrt{a}\sqrt{c}} x}{\sqrt{a + bx^2 + cx^4}}\right)}{4\sqrt{a}\sqrt{c}d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2 + c*x^4]/(a*d - c*d*x^4), x]

[Out] $(-\text{Sqrt}[-b - 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*\text{ArcTan}[(\text{Sqrt}[-b - 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*x)/\text{Sqrt}[a + b*x^2 + c*x^4]]) + \text{Sqrt}[-b + 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*\text{ArcTan}[(\text{Sqrt}[-b + 2*\text{Sqrt}[a]*\text{Sqrt}[c]]*x)/\text{Sqrt}[a + b*x^2 + c*x^4]])/(4*\text{Sqrt}[a]*\text{Sqrt}[c]*d)$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 221 vs. $2(105) = 210$.

time = 0.13, size = 222, normalized size = 1.53

method	result
--------	--------

+ a)*(a*d^2*sqrt(1/(a*c*d^4)) + x^2) - (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) + a*d*x)*sqrt((2*a*c*d^2*sqrt(1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 - a) + 1/8*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(c*x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) - x^2) + (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) - a*d*x)*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2)))/(c*x^4 - a) - 1/8*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2))*log(-(sqrt(c*x^4 + b*x^2 + a)*(a*d^2*sqrt(1/(a*c*d^4)) - x^2) - (a*c*d^3*x^3*sqrt(1/(a*c*d^4)) - a*d*x)*sqrt(-(2*a*c*d^2*sqrt(1/(a*c*d^4)) - b)/(a*c*d^2)))/(c*x^4 - a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a + bx^2 + cx^4}}{-a + cx^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*x**4+b*x**2+a)**(1/2)/(-c*d*x**4+a*d),x)

[Out] -Integral(sqrt(a + b*x**2 + c*x**4)/(-a + c*x**4), x)/d

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*x^4+b*x^2+a)^(1/2)/(-c*d*x^4+a*d),x, algorithm="giac")

[Out] integrate(-sqrt(c*x^4 + b*x^2 + a)/(c*d*x^4 - a*d), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{cx^4 + bx^2 + a}}{ad - cd x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2 + c*x^4)^(1/2)/(a*d - c*d*x^4),x)

[Out] int((a + b*x^2 + c*x^4)^(1/2)/(a*d - c*d*x^4), x)

$$3.36 \quad \int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cd x^4} dx$$

Optimal. Leaf size=239

$$\frac{\sqrt{b + \sqrt{b^2 + 4ac}} \tan^{-1} \left(\frac{\sqrt{b + \sqrt{b^2 + 4ac}} x (b - \sqrt{b^2 + 4ac} - 2cx^2)}{2\sqrt{2} \sqrt{a} \sqrt{c} \sqrt{a + bx^2 - cx^4}} \right)}{2\sqrt{2} \sqrt{a} \sqrt{c} d} + \frac{\sqrt{-b + \sqrt{b^2 + 4ac}} \tanh^{-1} \left(\frac{\sqrt{-b + \sqrt{b^2 + 4ac}} x (b + \sqrt{b^2 + 4ac} - 2cx^2)}{2\sqrt{2} \sqrt{a} \sqrt{c} \sqrt{a + bx^2 - cx^4}} \right)}{2\sqrt{2} \sqrt{a} \sqrt{c} d}$$

[Out] $1/4 \cdot \arctanh(1/4 * x * (b - 2 * c * x^2 + (4 * a * c + b^2)^{1/2}) * (-b + (4 * a * c + b^2)^{1/2}))^{1/2} * 2^{1/2} / a^{1/2} / c^{1/2} / (-c * x^4 + b * x^2 + a)^{1/2} * (-b + (4 * a * c + b^2)^{1/2})^{1/2} / d * 2^{1/2} / a^{1/2} / c^{1/2} - 1/4 \cdot \arctan(1/4 * x * (b - 2 * c * x^2 - (4 * a * c + b^2)^{1/2}) * (b + (4 * a * c + b^2)^{1/2}))^{1/2} * 2^{1/2} / a^{1/2} / c^{1/2} / (-c * x^4 + b * x^2 + a)^{1/2} / (b + (4 * a * c + b^2)^{1/2})^{1/2} / d * 2^{1/2} / a^{1/2} / c^{1/2}$

Rubi [A]

time = 0.12, antiderivative size = 239, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 30, $\frac{\text{number of rules}}{\text{integrand size}} = 0.033$, Rules used = {2097}

$$\frac{\sqrt{\sqrt{4ac + b^2} - b} \tanh^{-1} \left(\frac{x \sqrt{\sqrt{4ac + b^2} - b} (\sqrt{4ac + b^2} + b - 2cx^2)}{2\sqrt{2} \sqrt{a} \sqrt{c} \sqrt{a + bx^2 - cx^4}} \right)}{2\sqrt{2} \sqrt{a} \sqrt{c} d} - \frac{\sqrt{\sqrt{4ac + b^2} + b} \text{ArcTan} \left(\frac{x \sqrt{\sqrt{4ac + b^2} + b} (-\sqrt{4ac + b^2} + b - 2cx^2)}{2\sqrt{2} \sqrt{a} \sqrt{c} \sqrt{a + bx^2 - cx^4}} \right)}{2\sqrt{2} \sqrt{a} \sqrt{c} d}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[a + b*x^2 - c*x^4]/(a*d + c*d*x^4), x]$

[Out] $-1/2 * (\text{Sqrt}[b + \text{Sqrt}[b^2 + 4*a*c]] * \text{ArcTan}[(\text{Sqrt}[b + \text{Sqrt}[b^2 + 4*a*c]] * x * (b - \text{Sqrt}[b^2 + 4*a*c] - 2*c*x^2)) / (2*\text{Sqrt}[2]*\text{Sqrt}[a]*\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2 - c*x^4])]) / (\text{Sqrt}[2]*\text{Sqrt}[a]*\text{Sqrt}[c]*d) + (\text{Sqrt}[-b + \text{Sqrt}[b^2 + 4*a*c]] * \text{ArcTanh}[(\text{Sqrt}[-b + \text{Sqrt}[b^2 + 4*a*c]] * x * (b + \text{Sqrt}[b^2 + 4*a*c] - 2*c*x^2)) / (2*\text{Sqrt}[2]*\text{Sqrt}[a]*\text{Sqrt}[c]*\text{Sqrt}[a + b*x^2 - c*x^4])]) / (2*\text{Sqrt}[2]*\text{Sqrt}[a]*\text{Sqrt}[c]*d)$

Rule 2097

$\text{Int}[\text{Sqrt}[(a_) + (b_)*(x_)^2 + (c_)*(x_)^4]/((d_) + (e_)*(x_)^4), x_Symbol] \rightarrow \text{With}\{q = \text{Sqrt}[b^2 - 4*a*c]\}, \text{Simp}[(-a)*(\text{Sqrt}[b + q]/(2*\text{Sqrt}[2]*\text{Rt}[(-a)*c, 2]*d))*\text{ArcTan}[\text{Sqrt}[b + q]*x*((b - q + 2*c*x^2)/(2*\text{Sqrt}[2]*\text{Rt}[(-a)*c, 2]*\text{Sqrt}[a + b*x^2 + c*x^4])], x] + \text{Simp}[a*(\text{Sqrt}[-b + q]/(2*\text{Sqrt}[2]*\text{Rt}[(-a)*c, 2]*d))*\text{ArcTanh}[\text{Sqrt}[-b + q]*x*((b + q + 2*c*x^2)/(2*\text{Sqrt}[2]*\text{Rt}[(-a)*c, 2]*\text{Sqrt}[a + b*x^2 + c*x^4])], x]] /; \text{FreeQ}\{a, b, c, d, e\}, x] \&\& \text{EqQ}[c*d + a*e, 0] \&\& \text{NegQ}[a*c]$

Rubi steps

$$\int \frac{\sqrt{a + bx^2 - cx^4}}{ad + cd x^4} dx = -\frac{\sqrt{b + \sqrt{b^2 + 4ac}} \tan^{-1} \left(\frac{\sqrt{b + \sqrt{b^2 + 4ac}} x (b - \sqrt{b^2 + 4ac} - 2cx^2)}{2\sqrt{2} \sqrt{a} \sqrt{c} \sqrt{a + bx^2 - cx^4}} \right)}{2\sqrt{2} \sqrt{a} \sqrt{c} d} + \frac{\sqrt{-b + \dots}}{\dots}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.55, size = 151, normalized size = 0.63

$$\frac{i \left(\sqrt{-b - 2i\sqrt{a} \sqrt{c}} \tan^{-1} \left(\frac{\sqrt{-b - 2i\sqrt{a} \sqrt{c}} x}{\sqrt{a + bx^2 - cx^4}} \right) - \sqrt{-b + 2i\sqrt{a} \sqrt{c}} \tan^{-1} \left(\frac{\sqrt{-b + 2i\sqrt{a} \sqrt{c}} x}{\sqrt{a + bx^2 - cx^4}} \right) \right)}{4\sqrt{a} \sqrt{c} d}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a + b*x^2 - c*x^4]/(a*d + c*d*x^4), x]

[Out] ((I/4)*(Sqrt[-b - (2*I)*Sqrt[a]*Sqrt[c]]*ArcTan[(Sqrt[-b - (2*I)*Sqrt[a]*Sqrt[c]]*x)/Sqrt[a + b*x^2 - c*x^4]] - Sqrt[-b + (2*I)*Sqrt[a]*Sqrt[c]]*ArcTan[(Sqrt[-b + (2*I)*Sqrt[a]*Sqrt[c]]*x)/Sqrt[a + b*x^2 - c*x^4]])/(Sqrt[a]*Sqrt[c]*d)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 783 vs. 2(183) = 366.

time = 0.16, size = 784, normalized size = 3.28

method	result
elliptic	$\frac{\left(\sqrt{b + \sqrt{4ac + b^2}} \operatorname{b\,ln} \left(\frac{-cx^4 + bx^2 + a}{x^2} - \frac{\sqrt{-cx^4 + bx^2 + a} \sqrt{2} \sqrt{b + \sqrt{4ac + b^2}} + \sqrt{4ac + b^2}}{16dac} \right) \right)}{16dac} + \dots$
default	$\frac{\left(\sqrt{b + \sqrt{4ac + b^2}} \operatorname{b\,ln} \left(\frac{-cx^4 + bx^2 + a}{x^2} - \frac{\sqrt{-cx^4 + bx^2 + a} \sqrt{2} \sqrt{b + \sqrt{4ac + b^2}} + \sqrt{4ac + b^2}}{16ac} \right) \right)}{16ac} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d), x, method=_RETURNVERBOSE)

*sqrt(-1/(a*c*d^4)) + b)/(a*c*d^2))*log((sqrt(-c*x^4 + b*x^2 + a)*a*d^2*sqrt(-1/(a*c*d^4)) - sqrt(-c*x^4 + b*x^2 + a)*x^2 + (a*c*d^3*x^3*sqrt(-1/(a*c*d^4)) + a*d*x)*sqrt(-(2*a*c*d^2*sqrt(-1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 + a)) + 1/8*sqrt(-(2*a*c*d^2*sqrt(-1/(a*c*d^4)) + b)/(a*c*d^2))*log((sqrt(-c*x^4 + b*x^2 + a)*a*d^2*sqrt(-1/(a*c*d^4)) - sqrt(-c*x^4 + b*x^2 + a)*x^2 - (a*c*d^3*x^3*sqrt(-1/(a*c*d^4)) + a*d*x)*sqrt(-(2*a*c*d^2*sqrt(-1/(a*c*d^4)) + b)/(a*c*d^2)))/(c*x^4 + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{\sqrt{a + bx^2 - cx^4}}{a + cx^4} dx}{d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c*x**4+b*x**2+a)**(1/2)/(c*d*x**4+a*d), x)

[Out] Integral(sqrt(a + b*x**2 - c*x**4)/(a + c*x**4), x)/d

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((-c*x^4+b*x^2+a)^(1/2)/(c*d*x^4+a*d), x, algorithm="giac")

[Out] integrate(sqrt(-c*x^4 + b*x^2 + a)/(c*d*x^4 + a*d), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{-cx^4 + bx^2 + a}}{cdx^4 + ad} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a + b*x^2 - c*x^4)^(1/2)/(a*d + c*d*x^4), x)

[Out] int((a + b*x^2 - c*x^4)^(1/2)/(a*d + c*d*x^4), x)

3.37 $\int x \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx$

Optimal. Leaf size=309

$$\frac{e(12bcd - 16ad^2 - 7be^2)(e + 2dx)\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{128d^4(a + bx^2)} + \frac{bx^2(c + ex + dx^2)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{5d(a + bx^2)}$$

[Out] $\frac{1}{5}bx^2(d^2x^2+ex+c)^{3/2}((bx^2+a)^2)^{1/2}/d/(bx^2+a)-\frac{1}{240}(42b^2d^2ex-80a^2d^2+32b^2cd-35b^2e^2)(d^2x^2+ex+c)^{3/2}((bx^2+a)^2)^{1/2}/d^3/(bx^2+a)+\frac{1}{256}e(4cd-e^2)(-16a^2d^2+12b^2cd-7b^2e^2)\operatorname{arctanh}\left(\frac{1}{2}(2dx+e)/d\right)/(d^2x^2+ex+c)^{1/2}((bx^2+a)^2)^{1/2}/d^{9/2}/(bx^2+a)+\frac{1}{128}e(-16a^2d^2+12b^2cd-7b^2e^2)(2dx+e)(d^2x^2+ex+c)^{1/2}((bx^2+a)^2)^{1/2}/d^4/(bx^2+a)$

Rubi [A]

time = 0.41, antiderivative size = 309, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 38, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {6876, 1667, 793, 626, 635, 212}

$$\frac{e\sqrt{a^2+2abx^2+b^2x^4}(4cd-e^2)(-16ad^2+12bcd-7be^2)\operatorname{tanh}^{-1}\left(\frac{2dx+e}{\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{256d^{9/2}(a+bx^2)} + \frac{e\sqrt{a^2+2abx^2+b^2x^4}(2dx+e)\sqrt{c+dx^2+ex}(-16ad^2+12bcd-7be^2)}{128d^4(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4}(c+dx^2+ex)^{3/2}(-80ad^2+32bcd+42bdex-35be^2)}{240d^3(a+bx^2)} + \frac{bx^2\sqrt{a^2+2abx^2+b^2x^4}(c+dx^2+ex)^{3/2}}{5d(a+bx^2)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[x\operatorname{Sqrt}[c + ex + dx^2]*\operatorname{Sqrt}[a^2 + 2a*bx^2 + b^2*x^4], x]$

[Out] $(e*(12*b*c*d - 16*a*d^2 - 7*b*e^2)*(e + 2*d*x)*\operatorname{Sqrt}[c + ex + dx^2]*\operatorname{Sqrt}[a^2 + 2*a*bx^2 + b^2*x^4])/(128*d^4*(a + b*x^2)) + (b*x^2*(c + ex + dx^2)^{3/2}*\operatorname{Sqrt}[a^2 + 2*a*bx^2 + b^2*x^4])/(5*d*(a + b*x^2)) - ((32*b*c*d - 80*a*d^2 - 35*b*e^2 + 42*b*d*e*x)*(c + ex + dx^2)^{3/2}*\operatorname{Sqrt}[a^2 + 2*a*bx^2 + b^2*x^4])/(240*d^3*(a + b*x^2)) + (e*(4*c*d - e^2)*(12*b*c*d - 16*a*d^2 - 7*b*e^2)*\operatorname{Sqrt}[a^2 + 2*a*bx^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + ex + dx^2])])/(256*d^{9/2}*(a + b*x^2))$

Rule 212

$\operatorname{Int}[(a_0 + (b_0)*(x_0)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 626

$\operatorname{Int}[(a_0 + (b_0)*(x_0) + (c_0)*(x_0)^2)^{p_0}, x_Symbol] \rightarrow \operatorname{Simp}[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - \operatorname{Dist}[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), \operatorname{Int}[(a + b*x + c*x^2)^{p-1}, x], x] /;$ FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]

Rule 635

```
Int[1/Sqrt[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 793

```
Int[((d_) + (e_)*(x_))*((f_) + (g_)*(x_))*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[(-(b*e*g*(p + 2) - c*(e*f + d*g)*(2*p + 3) - 2*c*e*g*(p + 1)*x))*((a + b*x + c*x^2)^(p + 1)/(2*c^2*(p + 1)*(2*p + 3))), x] + Dist[(b^2*e*g*(p + 2) - 2*a*c*e*g + c*(2*c*d*f - b*(e*f + d*g))*(2*p + 3))/(2*c^2*(2*p + 3)), Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, p}, x] && NeQ[b^2 - 4*a*c, 0] && !LeQ[p, -1]
```

Rule 1667

```
Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)^(m + q - 1)*((a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1))), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))
```

Rule 6876

```
Int[(u_)*((a_) + (b_)*(x_)^(n_) + (c_)*(x_)^(n2_))^(p_), x_Symbol] := Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

3.38 $\int \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4} dx$

Optimal. Leaf size=283

$$\frac{(4bcd - 16ad^2 - 5be^2)(e + 2dx)\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{64d^3(a + bx^2)} - \frac{5be(c + ex + dx^2)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{24d^2(a + bx^2)}$$

[Out] $-5/24*b*e*(d*x^2+e*x+c)^{(3/2)*((b*x^2+a)^2)^{(1/2)}/d^2/(b*x^2+a)+1/4*b*x*(d*x^2+e*x+c)^{(3/2)*((b*x^2+a)^2)^{(1/2)}/d/(b*x^2+a)-1/128*(4*c*d-e^2)*(-16*a*d^2+4*b*c*d-5*b*e^2)*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/d^{(7/2)/(b*x^2+a)-1/64*(-16*a*d^2+4*b*c*d-5*b*e^2)*(2*d*x+e)*(d*x^2+e*x+c)^{(1/2)*((b*x^2+a)^2)^{(1/2)}/d^3/(b*x^2+a)}$

Rubi [A]

time = 0.23, antiderivative size = 283, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 37, $\frac{\text{number of rules}}{\text{integrand size}} = 0.162$, Rules used = {6876, 1675, 654, 626, 635, 212}

$$\frac{5be\sqrt{a^2+2abx^2+b^2x^4}(c+dx+ex)^{3/2}}{24d^3(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4}(4cd-e^2)(-16ad^2+4bcd-5be^2)\operatorname{tanh}^{-1}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx+ex}}\right)}{128d^{7/2}(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4}(2dx+e)\sqrt{c+dx^2+ex}(-16ad^2+4bcd-5be^2)}{64d^3(a+bx^2)} + \frac{bx\sqrt{a^2+2abx^2+b^2x^4}(c+dx+ex)^{3/2}}{4d(a+bx^2)}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]`

[Out] $-1/64*((4*b*c*d - 16*a*d^2 - 5*b*e^2)*(e + 2*d*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(d^3*(a + b*x^2)) - (5*b*e*(c + e*x + d*x^2)^{(3/2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(24*d^2*(a + b*x^2)) + (b*x*(c + e*x + d*x^2)^{(3/2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(4*d*(a + b*x^2)) - ((4*c*d - e^2)*(4*b*c*d - 16*a*d^2 - 5*b*e^2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(128*d^{(7/2)*(a + b*x^2)})$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 626

`Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(b + 2*c*x)*((a + b*x + c*x^2)^p/(2*c*(2*p + 1))), x] - Dist[p*((b^2 - 4*a*c)/(2*c*(2*p + 1))), Int[(a + b*x + c*x^2)^(p - 1), x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[p, 0] && IntegerQ[4*p]`

Rule 635


```
Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 654

```
Int[((d_.) + (e_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*((a + b*x + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Dist[(2*c*d - b*e)/(2*c), Int[(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, p}, x] && NeQ[2*c*d - b*e, 0] && NeQ[p, -1]
```

Rule 1675

```
Int[(Pq_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], e = Coeff[Pq, x, Expon[Pq, x]]}, Simp[e*x^(q - 1)*((a + b*x + c*x^2)^(p + 1)/(c*(q + 2*p + 1))), x] + Dist[1/(c*(q + 2*p + 1)), Int[(a + b*x + c*x^2)^p*ExpandToSum[c*(q + 2*p + 1)*Pq - a*e*(q - 1)*x^(q - 2) - b*e*(q + p)*x^(q - 1) - c*e*(q + 2*p + 1)*x^q, x], x], x] /; FreeQ[{a, b, c, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && !LeQ[p, -1]
```

Rule 6876

```
Int[(u_)*((a_) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^p_, x_Symbol] := Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int (2ab+2b^2x^2) \sqrt{c+ex+dx^2} dx}{2ab+2b^2x^2} \\
&= \frac{bx(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{4d(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4}}{4d} \\
&= -\frac{5be(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{24d^2(a+bx^2)} + \frac{bx(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{4d} \\
&= -\frac{(4bcd-16ad^2-5be^2)(e+2dx)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{64d^3(a+bx^2)} \\
&= -\frac{(4bcd-16ad^2-5be^2)(e+2dx)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{64d^3(a+bx^2)} \\
&= -\frac{(4bcd-16ad^2-5be^2)(e+2dx)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{64d^3(a+bx^2)}
\end{aligned}$$

Mathematica [A]

time = 0.41, size = 166, normalized size = 0.59

$$\frac{\sqrt{(a+bx^2)^2} \left(2\sqrt{d} \sqrt{c+x(e+dx)} (48ad^2(e+2dx) + b(15e^3 - 10de^2x + 8d^2e^2x^2 + 48d^3x^3 + 4cd(-13e + 6dx))) + 3(4cd - e^2) (4bcd - 16ad^2 - 5be^2) \log(e + 2dx - 2\sqrt{d} \sqrt{c+x(e+dx)}) \right)}{384d^{7/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4], x]

[Out] (Sqrt[(a + b*x^2)^2]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(48*a*d^2*(e + 2*d*x) + b*(15*e^3 - 10*d*e^2*x + 8*d^2*e*x^2 + 48*d^3*x^3 + 4*c*d*(-13*e + 6*d*x))) + 3*(4*c*d - e^2)*(4*b*c*d - 16*a*d^2 - 5*b*e^2)*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]])/(384*d^(7/2)*(a + b*x^2))

Maple [A]

time = 0.19, size = 373, normalized size = 1.32

method	result
risch	$ \frac{(48x^3d^3b+8be^2d^2+96ad^3x+24bcd^2x-10bd^2e^2x+48ad^2e-52bcde+15e^3b)\sqrt{dx^2+ex+c}\sqrt{(bx^2+a)^2}}{192d^3(bx^2+a)} + \left(\frac{\ln\left(\frac{e+dx}{\sqrt{d}}\right)}{\sqrt{d}} \right) $

default	$\sqrt{(bx^2 + a)^2} \left(96(dx^2 + ex + c)^{\frac{3}{2}} d^{\frac{7}{2}} bx - 80(dx^2 + ex + c)^{\frac{3}{2}} d^{\frac{5}{2}} be + 192d^{\frac{9}{2}} \sqrt{dx^2 + ex + c} \operatorname{arctan} \frac{ax - 48\sqrt{dx^2 + ex + c}}{d} \right)$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{384} \left((bx^2 + a)^2 \sqrt{dx^2 + ex + c} \left(96(dx^2 + ex + c)^{\frac{3}{2}} d^{\frac{7}{2}} bx - 80(dx^2 + ex + c)^{\frac{3}{2}} d^{\frac{5}{2}} be + 192d^{\frac{9}{2}} \sqrt{dx^2 + ex + c} \operatorname{arctan} \frac{ax - 48\sqrt{dx^2 + ex + c}}{d} \right) \right)$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested a dditional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c*d-%e^2>0)', see 'assume?' for more det

Fricas [A]

time = 0.38, size = 360, normalized size = 1.27

$$\frac{3(16b^2d^2 - 64acd + 5b^2e - 8(3bd - 2a^2d)\sqrt{d} \log(8d^2x^2 + 8dxe - 4\sqrt{d^2 + 2x^2}(2d + e)\sqrt{d + 4cd + e^2}) + 4(8b^2d^2 - 10b^2cd + 15bd^2 + 24(bd^2 + 4ad^2)e + 4(2b^2d^2 - 13bd^2 + 12ad^2)\sqrt{d^2 + 2x^2})) - 3(16b^2d^2 - 64acd + 5b^2e - 8(3bd - 2a^2d)\sqrt{d} \arctan(\frac{\sqrt{d^2 + 2x^2}\sqrt{d + 4cd + e^2}}{2d + e}) + 2(8b^2d^2 - 10b^2cd + 15bd^2 + 24(bd^2 + 4ad^2)e + 4(2b^2d^2 - 13bd^2 + 12ad^2)\sqrt{d^2 + 2x^2}))}{384d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="fricas")`

[Out] $\frac{1}{768} \left(3(16b^2d^2 - 64acd + 5b^2e - 8(3bd - 2a^2d)\sqrt{d} \log(8d^2x^2 + 8dxe - 4\sqrt{d^2 + 2x^2}(2d + e)\sqrt{d + 4cd + e^2}) + 4(8b^2d^2 - 10b^2cd + 15bd^2 + 24(bd^2 + 4ad^2)e + 4(2b^2d^2 - 13bd^2 + 12ad^2)\sqrt{d^2 + 2x^2})) - 3(16b^2d^2 - 64acd + 5b^2e - 8(3bd - 2a^2d)\sqrt{d} \arctan(\frac{\sqrt{d^2 + 2x^2}\sqrt{d + 4cd + e^2}}{2d + e}) + 2(8b^2d^2 - 10b^2cd + 15bd^2 + 24(bd^2 + 4ad^2)e + 4(2b^2d^2 - 13bd^2 + 12ad^2)\sqrt{d^2 + 2x^2})) \right) \sqrt{d} \sqrt{-d} \operatorname{arctan} \left(\frac{1}{2} \sqrt{d^2 + 2x^2} \sqrt{d + 4cd + e^2} \right) \sqrt{d + 4cd + e^2}$

$(-d)/(d^2*x^2 + d*x*e + c*d)) + 2*(48*b*d^4*x^3 - 10*b*d^2*x*e^2 + 15*b*d*e^3 + 24*(b*c*d^3 + 4*a*d^4)*x + 4*(2*b*d^3*x^2 - 13*b*c*d^2 + 12*a*d^3)*e)*\sqrt{d*x^2 + x*e + c})/d^4]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2),x)

[Out] Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2), x)

Giac [A]

time = 3.71, size = 265, normalized size = 0.94

$$\frac{1}{192} \sqrt{dx^2 + ex + c} \left(z \left(4 \left(6 \operatorname{sgn}(bx^2 + a) + \frac{b \operatorname{sgn}(bx^2 + a)}{d} \right) x + \frac{12bc \operatorname{sgn}(bx^2 + a) + 48ae \operatorname{sgn}(bx^2 + a) - 5bd^2 \operatorname{sgn}(bx^2 + a)}{d^2} \right) x - \frac{52bd \operatorname{sgn}(bx^2 + a) - 48ae^2 \operatorname{sgn}(bx^2 + a) - 15b^2 \operatorname{sgn}(bx^2 + a)}{d^2} \right) + \frac{(16b^2d^2 \operatorname{sgn}(bx^2 + a) - 64ae^2 \operatorname{sgn}(bx^2 + a) - 24bd^2 \operatorname{sgn}(bx^2 + a) + 16ae^2 \operatorname{sgn}(bx^2 + a) + 5b^2 \operatorname{sgn}(bx^2 + a)) \log \left(\frac{-2(\sqrt{dx^2 + ex + c})\sqrt{d} - z}{128d} \right)}{128d^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2),x, algorithm="giac")

[Out] $\frac{1}{192} \sqrt{dx^2 + ex + c} \left(2 \left(4 \left(6 \operatorname{sgn}(bx^2 + a) + \frac{b \operatorname{sgn}(bx^2 + a)}{d} \right) x + \frac{12bc \operatorname{sgn}(bx^2 + a) + 48ae \operatorname{sgn}(bx^2 + a) - 5bd^2 \operatorname{sgn}(bx^2 + a)}{d^2} \right) x - \frac{52bd \operatorname{sgn}(bx^2 + a) - 48ae^2 \operatorname{sgn}(bx^2 + a) - 15b^2 \operatorname{sgn}(bx^2 + a)}{d^2} \right) + \frac{1}{128} \left(16b^2d^2 \operatorname{sgn}(bx^2 + a) - 64a^2c \operatorname{sgn}(bx^2 + a) - 24b^2c \operatorname{sgn}(bx^2 + a) + 16ae^2 \operatorname{sgn}(bx^2 + a) + 5b^2e^4 \operatorname{sgn}(bx^2 + a) \right) \log \left(\frac{-2(\sqrt{dx^2 + ex + c})\sqrt{d} - z}{128d} \right) \right)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2),x)

[Out] int(((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2), x)

$$3.39 \quad \int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x} dx$$

Optimal. Leaf size=286

$$\frac{(8ad^2 - be^2 - 2bdex) \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{8d^2(a + bx^2)} + \frac{b(c + ex + dx^2)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{3d(a + bx^2)} + \frac{e(8a^2d - b^2e)}{8d^2(a + bx^2)}$$

[Out] $1/3*b*(d*x^2+e*x+c)^{(3/2)*((b*x^2+a)^2)^{(1/2)}/d/(b*x^2+a)+1/16*e*(8*a*d^2-b^2e)/(4*c*d-e^2)*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/d^{(5/2)/(b*x^2+a)-a*\operatorname{arctanh}(1/2*(e*x+2*c)/c^{(1/2)/(d*x^2+e*x+c)^{(1/2)})}*c^{(1/2)*((b*x^2+a)^2)^{(1/2)/(b*x^2+a)+1/8*(-2*b*d*e*x+8*a*d^2-b^2e)*}(d*x^2+e*x+c)^{(1/2)*((b*x^2+a)^2)^{(1/2)}/d^2/(b*x^2+a)}$

Rubi [A]

time = 0.50, antiderivative size = 286, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6876, 1667, 828, 857, 635, 212, 738}

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \sqrt{c + dx^2 + ex} (8ad^2 - 2bdex - be^2)}{8d^2(a + bx^2)} + \frac{e\sqrt{a^2 + 2abx^2 + b^2x^4} (8ad^2 - b(4cd - e^2)) \operatorname{tanh}^{-1}\left(\frac{2dxe}{\sqrt{d}\sqrt{c + dx^2 + ex}}\right)}{16d^{5/2}(a + bx^2)} + \frac{b\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2}}{3d(a + bx^2)} - \frac{a\sqrt{c}\sqrt{a^2 + 2abx^2 + b^2x^4} \operatorname{tanh}^{-1}\left(\frac{2cex}{\sqrt{c}\sqrt{c + dx^2 + ex}}\right)}{a + bx^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/x, x]$

[Out] $((8*a*d^2 - b*e^2 - 2*b*d*e*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(8*d^2*(a + b*x^2)) + (b*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(3*d*(a + b*x^2)) + (e*(8*a*d^2 - b*(4*c*d - e^2))*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(16*d^{(5/2)}*(a + b*x^2)) - (a*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(2*c + e*x)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(a + b*x^2)$

Rule 212

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

$\operatorname{Int}[1/\operatorname{Sqrt}[(a + (b \cdot x) + (c \cdot x)^2), x_Symbol] \rightarrow \operatorname{Dist}[2, \operatorname{Subst}[\operatorname{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\operatorname{Sqrt}[a + b*x + c*x^2]], x] /;$ FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738

```
Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol]
:> Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 828

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Simp[(d + e*x)^(m + 1)*(c*e*f*(m + 2*p + 2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m + 2*p + 1)*x)*((a + b*x + c*x^2)^p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2))), x] - Dist[p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p - 1)*Simp[c*e*f*(b*d - 2*a*e)*(m + 2*p + 2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

Rule 857

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1667

```
Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:> With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)^(m + q - 1)*((a + b*x + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1))), x] + Dist[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + b*x + c*x^2)^p*ExpandToSum[c*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)^(q - 2)*(b*d*e*(p + 1) + a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - e*(2*c*d - b*e)*(m + q + p)*x), x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, b, c, d, e, m, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !(IGtQ[m, 0] && RationalQ[a, b, c, d, e] && (IntegerQ[p] || ILtQ[p + 1/2, 0]))
```

Rule 6876

```
Int[(u_)*((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^p, x_Symbol]
:> Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ
```

$[b^2 - 4ac, 0]$ && IntegerQ[p - 1/2]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2)\sqrt{c+ex+dx^2}}{x} dx}{2ab+2b^2x^2} \\
 &= \frac{b(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3d(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4}}{\sqrt{c+ex+dx^2}} \\
 &= \frac{(8ad^2-be^2-2bdex) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4}}{\sqrt{c+ex+dx^2}} \\
 &= \frac{(8ad^2-be^2-2bdex) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4}}{\sqrt{c+ex+dx^2}} \\
 &= \frac{(8ad^2-be^2-2bdex) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4}}{\sqrt{c+ex+dx^2}} \\
 &= \frac{(8ad^2-be^2-2bdex) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8d^2(a+bx^2)} + \frac{\sqrt{a^2+2abx^2+b^2x^4}}{\sqrt{c+ex+dx^2}}
 \end{aligned}$$

Mathematica [A]

time = 0.49, size = 174, normalized size = 0.61

$$\frac{\sqrt{(a+bx^2)^2} \left(2\sqrt{d} \sqrt{c+x(e+dx)} (2Ad^2 + b(8cd - 3e^2 + 2dex + 8d^2x^2)) + 96a\sqrt{c} d^{5/2} \tanh^{-1} \left(\frac{\sqrt{d}x - \sqrt{c+x(e+dx)}}{\sqrt{c}} \right) - 3e(8ad^2 + b(-4cd + e^2)) \log(e + 2dx - 2\sqrt{d} \sqrt{c+x(e+dx)}) \right)}{48d^{5/2}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x,x]

[Out] (Sqrt[(a + b*x^2)^2]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(24*a*d^2 + b*(8*c*d - 3*e^2 + 2*d*e*x + 8*d^2*x^2)) + 96*a*Sqrt[c]*d^(5/2)*ArcTanh[(Sqrt[d]*x - Sqrt[c + x*(e + d*x)])/Sqrt[c]] - 3*e*(8*a*d^2 + b*(-4*c*d + e^2))*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]])/(48*d^(5/2)*(a + b*x^2))

Maple [A]

time = 0.20, size = 251, normalized size = 0.88

method	result
--------	--------

default	$\frac{\sqrt{(bx^2 + a)^2} \left(16(dx^2 + ex + c)^{\frac{3}{2}} d^{\frac{5}{2}} b - 12\sqrt{dx^2 + ex + c} d^{\frac{5}{2}} be - 48d^{\frac{7}{2}} \sqrt{c} \ln \left(\frac{2c + ex + 2\sqrt{c} \sqrt{dx^2 + ex + c}}{x} \right) \right)}{a}$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x,method=_RETURNVERBOSE)
```

```
[Out] 1/48*((b*x^2+a)^2)^(1/2)*(16*(d*x^2+e*x+c)^(3/2)*d^(5/2)*b-12*(d*x^2+e*x+c)^(1/2)*d^(5/2)*b*e*x-48*d^(7/2)*c^(1/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a+48*(d*x^2+e*x+c)^(1/2)*d^(7/2)*a-6*(d*x^2+e*x+c)^(1/2)*d^(3/2)*b*e^2+24*d^3*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*a*e-1/2*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c*d^2*e+3*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*d*e^3)/(b*x^2+a)/d^(7/2)
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c*d-%e^2>0)', see 'assume?' for more details)
```

Fricas [A]

time = 0.84, size = 771, normalized size = 2.70

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="fricas")
```

```
[Out] [1/96*(48*a*sqrt(c)*d^3*log((4*c*d*x^2 + x^2*e^2 + 8*c*x*e - 4*sqrt(d*x^2 + x*e + c))*(x*e + 2*c)*sqrt(c) + 8*c^2)/x^2) + 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*x*e + 4*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(8*b*d^3*x^2 + 2*b*d^2*x*e + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + x*e + c))/d^3, 1/48*(24*a*sqrt(c)*d^3*log((4*c*d*x^2 + x^2*e^2 + 8*c*x*e - 4*sqrt(d*x^2 + x*e + c))*(x*e + 2*c)*sqrt(c) + 8*c^2)/x^2) - 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*x*e + c*d)) + 2*(8*b*d^3*x^2
```


$$2 + 2*b*d^2*x*e + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + x*e + c))/d^3, 1/96*(96*a*sqrt(-c)*d^3*arctan(1/2*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(-c)/(c*d*x^2 + c*x*e + c^2)) + 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(d)*log(8*d^2*x^2 + 8*d*x*e + 4*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(8*b*d^3*x^2 + 2*b*d^2*x*e + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + x*e + c))/d^3, 1/48*(48*a*sqrt(-c)*d^3*arctan(1/2*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(-c)/(c*d*x^2 + c*x*e + c^2)) - 3*(b*e^3 - 4*(b*c*d - 2*a*d^2)*e)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*x*e + c*d)) + 2*(8*b*d^3*x^2 + 2*b*d^2*x*e + 8*b*c*d^2 + 24*a*d^3 - 3*b*d*e^2)*sqrt(d*x^2 + x*e + c))/d^3]$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x,x)

[Out] Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2)/x, x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, integration of abs or sign assumes constant sign by intervals (correct if the argument is real):Check [abs(sa

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x,x)

[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x, x)

$$3.40 \quad \int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x^2} dx$$

Optimal. Leaf size=294

$$\frac{((bc + 4ad)e + 2d(bc + 2ad)x)\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{4cd(a + bx^2)} - \frac{a(c + ex + dx^2)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{cx(a + bx^2)}$$

[Out] $-a*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/c/x/(b*x^2+a)+1/8*(8*a*d^2+4*b*c*d-b*e^2)*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)})/(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/d^{(3/2)}/(b*x^2+a)-1/2*a*e*\operatorname{arctanh}(1/2*(e*x+2*c)/c^{(1/2)})/(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/(b*x^2+a)/c^{(1/2)}+1/4*((4*a*d+b*c)*e+2*d*(2*a*d+b*c)*x)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c/d/(b*x^2+a)$

Rubi [A]

time = 0.54, antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6876, 1664, 828, 857, 635, 212, 738}

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} (8ad^2 + 4bcd - be^2) \operatorname{tanh}^{-1}\left(\frac{2dx+e}{\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{8d^{3/2}(a+bx^2)} - \frac{a\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2}}{cx(a+bx^2)} + \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \sqrt{c + dx^2 + ex} (e(4ad + bc) + 2dx(2ad + bc))}{4cd(a+bx^2)} - \frac{ae\sqrt{a^2 + 2abx^2 + b^2x^4} \operatorname{tanh}^{-1}\left(\frac{2dx+e}{\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{2\sqrt{c}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] `Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^2,x]`

[Out] $((b*c + 4*a*d)*e + 2*d*(b*c + 2*a*d)*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]/(4*c*d*(a + b*x^2)) - (a*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]/(c*x*(a + b*x^2)) + ((4*b*c*d + 8*a*d^2 - b*e^2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(8*d^{(3/2)}*(a + b*x^2)) - (a*e*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(2*c + e*x)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(2*\operatorname{Sqrt}[c]*(a + b*x^2))$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 635

`Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]`

Rule 738

```
Int[1/(((d_.) + (e_.)*(x_.))*Sqrt[(a_.) + (b_.)*(x_.) + (c_.)*(x_)^2]), x_Symbol]
:= Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 828

```
Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Simp[(d + e*x)^(m + 1)*(c*e*f*(m + 2*p + 2) - g*(c*d + 2*c*d*p - b*e*p) + g*c*e*(m + 2*p + 1)*x)*((a + b*x + c*x^2)^p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2))), x] - Dist[p/(c*e^2*(m + 2*p + 1)*(m + 2*p + 2)), Int[(d + e*x)^m*(a + b*x + c*x^2)^(p - 1)*Simp[c*e*f*(b*d - 2*a*e)*(m + 2*p + 2) + g*(a*e*(b*e - 2*c*d*m + b*e*m) + b*d*(b*e*p - c*d - 2*c*d*p)) + (c*e*f*(2*c*d - b*e)*(m + 2*p + 2) + g*(b^2*e^2*(p + m + 1) - 2*c^2*d^2*(1 + 2*p) - c*e*(b*d*(m - 2*p) + 2*a*e*(m + 2*p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && (IntegerQ[p] || !RationalQ[m] || (GeQ[m, -1] && LtQ[m, 0])) && !ILtQ[m + 2*p, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

Rule 857

```
Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1664

```
Int[(Pq_)*((d_.) + (e_.)*(x_.))^(m_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= With[{Q = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Q + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]
```

Rule 6876

```
Int[(u_)*((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^(p_.), x_Symbol]
:= Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^2} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2) \sqrt{c+ex+dx^2}}{x^2} dx}{2ab+2b^2x^2} \\
&= -\frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{cx(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4}}{a+bx^2} \\
&= \frac{((bc+4ad)e+2d(bc+2ad)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} \\
&= \frac{((bc+4ad)e+2d(bc+2ad)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} \\
&= \frac{((bc+4ad)e+2d(bc+2ad)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)} \\
&= \frac{((bc+4ad)e+2d(bc+2ad)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cd(a+bx^2)}
\end{aligned}$$

Mathematica [A]

time = 0.63, size = 169, normalized size = 0.57

$$\frac{\sqrt{(a+bx^2)^2} \left(8ad^{3/2}ex \tanh^{-1} \left(\frac{\sqrt{d}x - \sqrt{c+x(e+dx)}}{\sqrt{c}} \right) + \sqrt{c} \left(2\sqrt{d} \sqrt{c+x(e+dx)} (-4ad+bx(e+2dx)) + (-8ad^2+b(-4cd+e^2))x \log \left(d(e+2dx-2\sqrt{d} \sqrt{c+x(e+dx)}) \right) \right) \right)}{8\sqrt{c} d^{3/2}x(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^2,x]

```
[Out] (Sqrt[(a + b*x^2)^2]*(8*a*d^(3/2)*e*x*ArcTanh[(Sqrt[d]*x - Sqrt[c + x*(e + d*x)])/Sqrt[c]] + Sqrt[c]*(2*Sqrt[d]*Sqrt[c + x*(e + d*x)]*(-4*a*d + b*x*(e + 2*d*x)) + (-8*a*d^2 + b*(-4*c*d + e^2))*x*Log[d*(e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)])])))/(8*Sqrt[c]*d^(3/2)*x*(a + b*x^2))
```

Maple [A]

time = 0.22, size = 288, normalized size = 0.98

method	result
--------	--------

risch	$-\frac{a\sqrt{dx^2+ex+c}\sqrt{(bx^2+a)^2}}{x(bx^2+a)} + \left(\frac{bx\sqrt{dx^2+ex+c}}{2} + \frac{be\sqrt{dx^2+ex+c}}{4d} - \frac{be^2 \ln\left(\frac{\frac{e}{2}+dx}{\sqrt{d}} + \sqrt{dx^2+ex+c}\right)}{8d^{\frac{3}{2}}}\right)$
default	$-\frac{\sqrt{(bx^2+a)^2}\left(-8\sqrt{dx^2+ex+c}d^{\frac{7}{2}}ax^2-4\sqrt{dx^2+ex+c}d^{\frac{5}{2}}bcx^2+4d^{\frac{5}{2}}\sqrt{c}\ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right)\right)}{x(bx^2+a)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

[Out]
$$-1/8*((b*x^2+a)^2)^(1/2)*(-8*(d*x^2+e*x+c)^(1/2)*d^(7/2)*a*x^2-4*(d*x^2+e*x+c)^(1/2)*d^(5/2)*b*c*x^2+4*d^(5/2)*c^(1/2)*\ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a*e*x+8*(d*x^2+e*x+c)^(3/2)*d^(5/2)*a-8*(d*x^2+e*x+c)^(1/2)*d^(5/2)*a*e*x-2*(d*x^2+e*x+c)^(1/2)*d^(3/2)*b*c*e*x-8*\ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*a*c*d^3*x-4*\ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c^2*d^2*x+\ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c*d*e^2*x)/(b*x^2+a)/c/x/d^(5/2)$$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation may help (example of legal syntax is 'assume(4*c*d-%e^2>0)', see 'assume?' for more details)

Fricas [A]

time = 0.71, size = 767, normalized size = 2.61

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="fricas")`

[Out]
$$[1/16*(4*a*\sqrt{c}*d^2*x*e*\log((4*c*d*x^2+x^2*e^2+8*c*x*e-4*\sqrt{d*x^2+x*e+c})*(x*e+2*c))*\sqrt{c}+8*c^2)/x^2)+(b*c*x*e^2-4*(b*c^2*d+)$$

```

2*a*c*d^2)*x)*sqrt(d)*log(8*d^2*x^2 + 8*d*x*e - 4*sqrt(d*x^2 + x*e + c)*(2*
d*x + e)*sqrt(d) + 4*c*d + e^2) + 4*(2*b*c*d^2*x^2 + b*c*d*x*e - 4*a*c*d^2)
*sqrt(d*x^2 + x*e + c))/(c*d^2*x), 1/8*(2*a*sqrt(c)*d^2*x*e*log((4*c*d*x^2
+ x^2*e^2 + 8*c*x*e - 4*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(c) + 8*c^2)/
x^2) + (b*c*x*e^2 - 4*(b*c^2*d + 2*a*c*d^2)*x)*sqrt(-d)*arctan(1/2*sqrt(d*x
^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*x*e + c*d)) + 2*(2*b*c*d^2*
x^2 + b*c*d*x*e - 4*a*c*d^2)*sqrt(d*x^2 + x*e + c))/(c*d^2*x), 1/16*(8*a*sq
rt(-c)*d^2*x*arctan(1/2*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(-c)/(c*d*x^2
+ c*x*e + c^2))*e + (b*c*x*e^2 - 4*(b*c^2*d + 2*a*c*d^2)*x)*sqrt(d)*log(8*
d^2*x^2 + 8*d*x*e - 4*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e
^2) + 4*(2*b*c*d^2*x^2 + b*c*d*x*e - 4*a*c*d^2)*sqrt(d*x^2 + x*e + c))/(c*d
^2*x), 1/8*(4*a*sqrt(-c)*d^2*x*arctan(1/2*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)
*sqrt(-c)/(c*d*x^2 + c*x*e + c^2))*e + (b*c*x*e^2 - 4*(b*c^2*d + 2*a*c*d^2)
*x)*sqrt(-d)*arctan(1/2*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2
+ d*x*e + c*d)) + 2*(2*b*c*d^2*x^2 + b*c*d*x*e - 4*a*c*d^2)*sqrt(d*x^2 + x
*e + c))/(c*d^2*x)]

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**2,x)
```

```
[Out] Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2)/x**2, x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^2,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);OUTPUT:Warning, integration of abs or sign a
ssumes constant sign by intervals (correct if the argument is real):Check [
abs(sa
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^2,x)
```

```
[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^2, x)
```

$$3.41 \quad \int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x^3} dx$$

Optimal. Leaf size=288

$$\frac{(ae + 2(2bc + ad)x)\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{4cx(a + bx^2)} - \frac{a(c + ex + dx^2)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{2cx^2(a + bx^2)} + \frac{be\sqrt{a^2 + 2abx^2 + b^2x^4}}{2cx^2(a + bx^2)}$$

[Out] $-1/2*a*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/c/x^2/(b*x^2+a)-1/8*(4*a*c*d-a*e^2+8*b*c^2)*\operatorname{arctanh}(1/2*(e*x+2*c)/c^{(1/2)})/(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c^{(3/2)}/(b*x^2+a)+1/2*b*e*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)})/(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/(b*x^2+a)/d^{(1/2)}+1/4*(a*e+2*(a*d+2*b*c))*x*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c/x/(b*x^2+a)$

Rubi [A]

time = 0.54, antiderivative size = 288, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6876, 1664, 826, 857, 635, 212, 738}

$$-\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} (4acd - ae^2 + 8bc^2) \operatorname{tanh}^{-1}\left(\frac{2e+ex}{2\sqrt{c+dx^2+ex}}\right)}{8c^{3/2}(a+bx^2)} - \frac{a\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2}}{2cx^2(a+bx^2)} + \frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \sqrt{c + dx^2 + ex} (2x(ad + 2bc) + ae)}{4cx(a+bx^2)} + \frac{be\sqrt{a^2 + 2abx^2 + b^2x^4} \operatorname{tanh}^{-1}\left(\frac{2d+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{2\sqrt{d}(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Int[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^3,x]

[Out] $((a*e + 2*(2*b*c + a*d)*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(4*c*x*(a + b*x^2)) - (a*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(2*c*x^2*(a + b*x^2)) + (b*e*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(2*\operatorname{Sqrt}[d]*(a + b*x^2)) - ((8*b*c^2 + 4*a*c*d - a*e^2)*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(2*c + e*x)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(8*c^{(3/2)}*(a + b*x^2))$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 635

Int[1/Sqrt[(a_) + (b_.)*(x_) + (c_.)*(x_)^2], x_Symbol] := Dist[2, Subst[Int[1/(4*c - x^2), x], x, (b + 2*c*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 738


```
Int[1/(((d_.) + (e_.)*(x_.))*Sqrt[(a_.) + (b_.)*(x_.) + (c_.)*(x_)^2]), x_Symbol]
:= Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 826

```
Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Simp[(d + e*x)^(m + 1)*(e*f*(m + 2*p + 2) - d*g*(2*p + 1) + e*g*(m + 1)*x)*((a + b*x + c*x^2)^p/(e^2*(m + 1)*(m + 2*p + 2))), x]
+ Dist[p/(e^2*(m + 1)*(m + 2*p + 2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p - 1)*Simp[g*(b*d + 2*a*e + 2*a*e*m + 2*b*d*p) - f*b*e*(m + 2*p + 2) + (g*(2*c*d + b*e + b*e*m + 4*c*d*p) - 2*c*e*f*(m + 2*p + 2))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, m}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && RationalQ[p] && p > 0 && (LtQ[m, -1] || EqQ[p, 1] || (IntegerQ[p] && !RationalQ[m])) && NeQ[m, -1] && !ILtQ[m + 2*p + 1, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])
```

Rule 857

```
Int[((d_.) + (e_.)*(x_.))^(m_.)*((f_.) + (g_.)*(x_.))*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1664

```
Int[(Pq_)*((d_.) + (e_.)*(x_.))^(m_.)*((a_.) + (b_.)*(x_.) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= With[{Q = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Q + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]
```

Rule 6876

```
Int[(u_)*((a_.) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^(p_.), x_Symbol]
:= Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^3} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4}}{2ab+2b^2x^2} \int \frac{(2ab+2b^2x^2)\sqrt{c+ex+dx^2}}{x^3} dx \\
&= -\frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4}}{a+bx^2} \\
&= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \\
&= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \\
&= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)} \\
&= \frac{(ae+2(2bc+ad)x)\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{4cx(a+bx^2)} - \frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{2cx^2(a+bx^2)}
\end{aligned}$$

Mathematica [A]

time = 0.61, size = 173, normalized size = 0.60

$$\frac{\sqrt{(a+bx^2)^2} \left(\sqrt{d} (8bc^2 + 4acd - ae^2) x^2 \tanh^{-1} \left(\frac{-\sqrt{d}x + \sqrt{c+x(e+dx)}}{\sqrt{c}} \right) + \sqrt{c} \left(\sqrt{d} (2ac + aex - 4bcx^2) \sqrt{c+x(e+dx)} + 2bce x^2 \log(e + 2dx - 2\sqrt{d} \sqrt{c+x(e+dx)}) \right) \right)}{4c^{3/2} \sqrt{d} x^2 (a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^3,x]

[Out] $-\frac{1}{4} * (\text{Sqrt}[(a + b*x^2)^2] * (\text{Sqrt}[d] * (8*b*c^2 + 4*a*c*d - a*e^2) * x^2 * \text{ArcTanh}[-(\text{Sqrt}[d]*x) + \text{Sqrt}[c + x*(e + d*x)]] / \text{Sqrt}[c]) + \text{Sqrt}[c] * (\text{Sqrt}[d] * (2*a*c + a*e*x - 4*b*c*x^2) * \text{Sqrt}[c + x*(e + d*x)] + 2*b*c*e*x^2 * \text{Log}[e + 2*d*x - 2*\text{Sqrt}[d] * \text{Sqrt}[c + x*(e + d*x)]])) / (c^{3/2} * \text{Sqrt}[d] * x^2 * (a + b*x^2))$

Maple [A]

time = 0.20, size = 330, normalized size = 1.15

method	result
risch	$ -\frac{a\sqrt{dx^2+ex+c} (ex+2c) \sqrt{(bx^2+a)^2}}{4x^2c(bx^2+a)} + \left(b\sqrt{dx^2+ex+c} + \frac{eb \ln\left(\frac{\frac{e}{2}+dx}{\sqrt{d}} + \sqrt{dx^2+ex+c}\right)}{2\sqrt{d}} \right) \ln\left(\frac{2c+e}{\dots}\right) $

default	$-\frac{\sqrt{(bx^2+a)^2} \left(4d^{\frac{5}{2}}c^{\frac{3}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) ax^2+8d^{\frac{3}{2}}c^{\frac{5}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) \right)}{b}$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x,method=_RETURNVERBOSE)
[Out] -1/8*((b*x^2+a)^(1/2))*(4*d^(5/2)*c^(3/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a*x^2+8*d^(3/2)*c^(5/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*b*x^2+2*(d*x^2+e*x+c)^(1/2)*d^(5/2)*a*e*x^3-4*(d*x^2+e*x+c)^(1/2)*d^(5/2)*a*c*x^2-d^(3/2)*c^(1/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a*e^2*x^2-2*(d*x^2+e*x+c)^(3/2)*d^(3/2)*a*e*x+2*(d*x^2+e*x+c)^(1/2)*d^(3/2)*a*e^2*x^2-8*(d*x^2+e*x+c)^(1/2)*d^(3/2)*b*c^2*x^2+4*(d*x^2+e*x+c)^(3/2)*d^(3/2)*a*c-4*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*d*b*c^2*e*x^2)/d^(3/2)/x^2/c^2/(b*x^2+a)
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="maxima")
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c*d-%e^2>0)', see 'assume?' for more det
```

Fricas [A]

time = 0.74, size = 797, normalized size = 2.77

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="fricas")
[Out] [1/16*(4*b*c^2*sqrt(d)*x^2*e*log(8*d^2*x^2 + 8*d*x*e + 4*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + (a*d*x^2*e^2 - 4*(2*b*c^2*d + a*c*d^2)*x^2)*sqrt(c)*log((4*c*d*x^2 + x^2*e^2 + 8*c*x*e + 4*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(c) + 8*c^2)/x^2) + 4*(4*b*c^2*d*x^2 - a*c*d*x*e - 2*a*c^2*d)*sqrt(d*x^2 + x*e + c))/(c^2*d*x^2), -1/16*(8*b*c^2*sqrt(-d)*x^2*arctan(1/2*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*x*e + c*d))*
```

$$e - (a*d*x^2*e^2 - 4*(2*b*c^2*d + a*c*d^2)*x^2)*\sqrt{c}*\log((4*c*d*x^2 + x^2*e^2 + 8*c*x*e + 4*\sqrt{d*x^2 + x*e + c})*(x*e + 2*c)*\sqrt{c} + 8*c^2)/x^2) - 4*(4*b*c^2*d*x^2 - a*c*d*x*e - 2*a*c^2*d)*\sqrt{d*x^2 + x*e + c})/(c^2*d*x^2), 1/8*(2*b*c^2*\sqrt{d}*x^2*e*\log(8*d^2*x^2 + 8*d*x*e + 4*\sqrt{d*x^2 + x*e + c})*(2*d*x + e)*\sqrt{d} + 4*c*d + e^2) - (a*d*x^2*e^2 - 4*(2*b*c^2*d + a*c*d^2)*x^2)*\sqrt{-c}*\arctan(1/2*\sqrt{d*x^2 + x*e + c}*(x*e + 2*c)*\sqrt{-c})/(c*d*x^2 + c*x*e + c^2)) + 2*(4*b*c^2*d*x^2 - a*c*d*x*e - 2*a*c^2*d)*\sqrt{d*x^2 + x*e + c})/(c^2*d*x^2), -1/8*(4*b*c^2*\sqrt{-d}*x^2*\arctan(1/2*\sqrt{d*x^2 + x*e + c}*(2*d*x + e)*\sqrt{-d})/(d^2*x^2 + d*x*e + c*d))*e + (a*d*x^2*e^2 - 4*(2*b*c^2*d + a*c*d^2)*x^2)*\sqrt{-c}*\arctan(1/2*\sqrt{d*x^2 + x*e + c}*(x*e + 2*c)*\sqrt{-c})/(c*d*x^2 + c*x*e + c^2)) - 2*(4*b*c^2*d*x^2 - a*c*d*x*e - 2*a*c^2*d)*\sqrt{d*x^2 + x*e + c})/(c^2*d*x^2)]$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c + dx^2 + ex} \sqrt{(a + bx^2)^2}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**3,x)

[Out] Integral(sqrt(c + d*x**2 + e*x)*sqrt((a + b*x**2)**2)/x**3, x)

Giac [A]

time = 7.32, size = 370, normalized size = 1.28

$$\frac{\log\left(\frac{-2(\sqrt{d}x - \sqrt{d^2 + ex + c})(d - \sqrt{d}x)\operatorname{sgn}(bx^2 + a)}{2\sqrt{d}} + \sqrt{d^2 + ex + c}\operatorname{sgn}(bx^2 + a)\right) + \frac{(b\sqrt{d}\operatorname{sgn}(bx^2 + a) + 4\operatorname{arctan}(bx^2 + a) - a\sqrt{d}\operatorname{sgn}(bx^2 + a))\operatorname{arctan}\left(\frac{-\sqrt{d}x - \sqrt{d^2 + ex + c}}{2\sqrt{d}}\right) + 4(\sqrt{d}x - \sqrt{d^2 + ex + c})\operatorname{arctan}(bx^2 + a) + 8(\sqrt{d}x - \sqrt{d^2 + ex + c})\operatorname{arctan}(bx^2 + a) + 4(\sqrt{d}x - \sqrt{d^2 + ex + c})\operatorname{arctan}(bx^2 + a) + (\sqrt{d}x - \sqrt{d^2 + ex + c})\operatorname{arctan}(bx^2 + a) + (\sqrt{d}x - \sqrt{d^2 + ex + c})\operatorname{arctan}(bx^2 + a)}{4(\sqrt{d}x - \sqrt{d^2 + ex + c})}}{4(\sqrt{d}x - \sqrt{d^2 + ex + c})}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^3,x, algorithm="giac")

[Out] $-1/2*b*e*\log(\operatorname{abs}(-2*(\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})*d - \sqrt{d}*e))*\operatorname{sgn}(b*x^2 + a)/\sqrt{d} + \sqrt{d*x^2 + x*e + c}*b*\operatorname{sgn}(b*x^2 + a) + 1/4*(8*b*c^2*\operatorname{sgn}(b*x^2 + a) + 4*a*c*d*\operatorname{sgn}(b*x^2 + a) - a*e^2*\operatorname{sgn}(b*x^2 + a))*\operatorname{arctan}(-(\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})/\sqrt{-c})/(\sqrt{-c}*c) + 1/4*(4*(\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})^3*a*c*d*\operatorname{sgn}(b*x^2 + a) + 8*(\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})^2*a*c*\sqrt{d}*e*\operatorname{sgn}(b*x^2 + a) + 4*(\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})*a*c^2*d*\operatorname{sgn}(b*x^2 + a) + (\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})^3*a*e^2*\operatorname{sgn}(b*x^2 + a) + (\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})*a*c*e^2*\operatorname{sgn}(b*x^2 + a))/((\sqrt{d}*x - \sqrt{d*x^2 + x*e + c})^2 - c)^2*c)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^3,x)
```

```
[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^3, x)
```

$$3.42 \quad \int \frac{\sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{x^4} dx$$

Optimal. Leaf size=294

$$\frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c + ex + dx^2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{8c^2x^2(a + bx^2)} - \frac{a(c + ex + dx^2)^{3/2} \sqrt{a^2 + 2abx^2 + b^2x^4}}{3cx^3(a + bx^2)} + \dots$$

[Out] $-1/3*a*(d*x^2+e*x+c)^{(3/2)}*((b*x^2+a)^2)^{(1/2)}/c/x^3/(b*x^2+a)-1/16*e*(8*b*c^2-a*(4*c*d-e^2))*\operatorname{arctanh}(1/2*(e*x+2*c)/c^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*((b*x^2+a)^2)^{(1/2)}/c^{(5/2)}/(b*x^2+a)+b*\operatorname{arctanh}(1/2*(2*d*x+e)/d^{(1/2)}/(d*x^2+e*x+c)^{(1/2)})*d^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/(b*x^2+a)+1/8*(2*a*c*e-(-a*e^2+8*b*c^2)*x)*(d*x^2+e*x+c)^{(1/2)}*((b*x^2+a)^2)^{(1/2)}/c^2/x^2/(b*x^2+a)$

Rubi [A]

time = 0.51, antiderivative size = 294, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 40, $\frac{\text{number of rules}}{\text{integrand size}} = 0.175$, Rules used = {6876, 1664, 824, 857, 635, 212, 738}

$$\frac{\sqrt{a^2 + 2abx^2 + b^2x^4} \sqrt{c + dx^2 + ex} (2ace - x(8bc^2 - ae^2))}{8c^2x^2(a + bx^2)} - \frac{e\sqrt{a^2 + 2abx^2 + b^2x^4} (8bc^2 - a(4cd - e^2)) \operatorname{tanh}^{-1}\left(\frac{2c+ex}{2\sqrt{c}\sqrt{c+dx^2+ex}}\right)}{16c^{5/2}(a + bx^2)} + \frac{b\sqrt{d}\sqrt{a^2 + 2abx^2 + b^2x^4} \operatorname{tanh}^{-1}\left(\frac{2dx+e}{2\sqrt{d}\sqrt{c+dx^2+ex}}\right)}{a + bx^2} - \frac{a\sqrt{a^2 + 2abx^2 + b^2x^4} (c + dx^2 + ex)^{3/2}}{3cx^3(a + bx^2)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/x^4, x]$

[Out] $((2*a*c*e - (8*b*c^2 - a*e^2)*x)*\operatorname{Sqrt}[c + e*x + d*x^2]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(8*c^2*x^2*(a + b*x^2)) - (a*(c + e*x + d*x^2)^{(3/2)}*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4])/(3*c*x^3*(a + b*x^2)) + (b*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(e + 2*d*x)/(2*\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(a + b*x^2) - (e*(8*b*c^2 - a*(4*c*d - e^2))*\operatorname{Sqrt}[a^2 + 2*a*b*x^2 + b^2*x^4]*\operatorname{ArcTanh}[(2*c + e*x)/(2*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[c + e*x + d*x^2])])/(16*c^{(5/2)}*(a + b*x^2))$

Rule 212

$\operatorname{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 635

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_) + (b_)*(x_) + (c_)*(x_)^2], x_Symbol] \rightarrow \operatorname{Dist}[2, \operatorname{Subst}[\operatorname{Int}[1/(4*c - x^2), x], x, (b + 2*c*x)/\operatorname{Sqrt}[a + b*x + c*x^2]], x] /; \operatorname{FreeQ}\{a, b, c\}, x \ \&\& \operatorname{NeQ}[b^2 - 4*a*c, 0]$

Rule 738

```
Int[1/(((d_.) + (e_.)*(x_))*Sqrt[(a_.) + (b_.)*(x_) + (c_.)*(x_)^2]), x_Symbol]
:= Dist[-2, Subst[Int[1/(4*c*d^2 - 4*b*d*e + 4*a*e^2 - x^2), x], x, (2*a*e - b*d - (2*c*d - b*e)*x)/Sqrt[a + b*x + c*x^2]], x] /; FreeQ[{a, b, c, d, e}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[2*c*d - b*e, 0]
```

Rule 824

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= Simp[(-(d + e*x)^(m + 1))*((a + b*x + c*x^2)^p/(e^2*(m + 1)*(m + 2)*(c*d^2 - b*d*e + a*e^2)))*((d*g - e*f*(m + 2))*(c*d^2 - b*d*e + a*e^2) - d*p*(2*c*d - b*e)*(e*f - d*g) - e*(g*(m + 1)*(c*d^2 - b*d*e + a*e^2) + p*(2*c*d - b*e)*(e*f - d*g))*x], x] - Dist[p/(e^2*(m + 1)*(m + 2)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 2)*(a + b*x + c*x^2)^(p - 1)*Simp[2*a*c*e*(e*f - d*g)*(m + 2) + b^2*e*(d*g*(p + 1) - e*f*(m + p + 2)) + b*(a*e^2*g*(m + 1) - c*d*(d*g*(2*p + 1) - e*f*(m + 2*p + 2)) - c*(2*c*d*(d*g*(2*p + 1) - e*f*(m + 2*p + 2)) - e*(2*a*e*g*(m + 1) - b*(d*g*(m - 2*p) + e*f*(m + 2*p + 2)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && GtQ[p, 0] && LtQ[m, -2] && LtQ[m + 2*p, 0] && !ILtQ[m + 2*p + 3, 0]
```

Rule 857

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= Dist[g/e, Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p, x], x] + Dist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + b*x + c*x^2)^p, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, p}, x]
&& NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 1664

```
Int[(Pq_)*((d_.) + (e_.)*(x_))^(m_)*((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol]
:= With[{Q = PolynomialQuotient[Pq, d + e*x, x], R = PolynomialRemainder[Pq, d + e*x, x]}, Simp[(e*R*(d + e*x)^(m + 1)*(a + b*x + c*x^2)^(p + 1))/((m + 1)*(c*d^2 - b*d*e + a*e^2)), x] + Dist[1/((m + 1)*(c*d^2 - b*d*e + a*e^2)), Int[(d + e*x)^(m + 1)*(a + b*x + c*x^2)^p*ExpandToSum[(m + 1)*(c*d^2 - b*d*e + a*e^2)*Q + c*d*R*(m + 1) - b*e*R*(m + p + 2) - c*e*R*(m + 2*p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, p}, x] && PolyQ[Pq, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && LtQ[m, -1]
```

Rule 6876

```
Int[(u_)*((a_) + (b_.)*(x_)^(n_.) + (c_.)*(x_)^(n2_.))^(p_), x_Symbol]
:= Dist[Sqrt[a + b*x^n + c*x^(2*n)]/((4*c)^(p - 1/2)*(b + 2*c*x^n)), Int[u*(b + 2*c*x^n)^(2*p), x], x] /; FreeQ[{a, b, c, n, p}, x] && EqQ[n2, 2*n] && EqQ[b^2 - 4*a*c, 0] && IntegerQ[p - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{x^4} dx &= \frac{\sqrt{a^2+2abx^2+b^2x^4} \int \frac{(2ab+2b^2x^2) \sqrt{c+ex+dx^2}}{x^4} dx}{2ab+2b^2x^2} \\
&= -\frac{a(c+ex+dx^2)^{3/2} \sqrt{a^2+2abx^2+b^2x^4}}{3cx^3(a+bx^2)} - \frac{\sqrt{a^2+2abx^2+b^2x^4}}{a+bx^2} \\
&= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} \\
&= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} \\
&= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)} \\
&= \frac{(2ace - (8bc^2 - ae^2)x) \sqrt{c+ex+dx^2} \sqrt{a^2+2abx^2+b^2x^4}}{8c^2x^2(a+bx^2)}
\end{aligned}$$

Mathematica [A]

time = 0.77, size = 186, normalized size = 0.63

$$\frac{\sqrt{(a+bx^2)^2} \left(-3e(8bc^2 + a(-4cd + e^2))x^3 \tanh^{-1} \left(\frac{-\sqrt{d}x + \sqrt{c+x(dx)}}{\sqrt{c}} \right) - \sqrt{c} \left(\sqrt{c+x(dx)} (24bc^2x^2 + a(8c^2 - 3e^2x^2 + 2cx(e+4dx))) + 24bc^2\sqrt{d}x^3 \log(e+2dx - 2\sqrt{d}\sqrt{c+x(dx)}) \right) \right)}{24c^{5/2}x^3(a+bx^2)}$$

Antiderivative was successfully verified.

[In] Integrate[(Sqrt[c + e*x + d*x^2]*Sqrt[a^2 + 2*a*b*x^2 + b^2*x^4])/x^4,x]

```
[Out] (Sqrt[(a + b*x^2)^2]*(-3*e*(8*b*c^2 + a*(-4*c*d + e^2))*x^3*ArcTanh[(-(Sqrt[d]*x) + Sqrt[c + x*(e + d*x)])/Sqrt[c]] - Sqrt[c]*(Sqrt[c + x*(e + d*x)]*(24*b*c^2*x^2 + a*(8*c^2 - 3*e^2*x^2 + 2*c*x*(e + 4*d*x))) + 24*b*c^2*Sqrt[d]*x^3*Log[e + 2*d*x - 2*Sqrt[d]*Sqrt[c + x*(e + d*x)]]))/(24*c^(5/2)*x^3*(a + b*x^2))
```

Maple [A]

time = 0.20, size = 412, normalized size = 1.40

method	result
--------	--------

risch	$-\frac{\sqrt{dx^2+ex+c} (8acd^2x^2-3ae^2x^2+24b^2c^2x^2+2acex+8c^2a)\sqrt{(bx^2+a)^2}}{24x^3c^2(bx^2+a)} + \frac{\left(b\sqrt{d} \ln\left(\frac{\frac{x}{2}+dx}{\sqrt{d}}+\sqrt{dx^2+ex+c}\right) \right)}{1}$
default	$-\frac{\sqrt{(bx^2+a)^2} \left(-12d^{\frac{5}{2}}c^{\frac{3}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) ae^3x^3+24d^{\frac{3}{2}}c^{\frac{5}{2}} \ln\left(\frac{2c+ex+2\sqrt{c}\sqrt{dx^2+ex+c}}{x}\right) \right)}{1}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x,method=_RETURNVERBOSE)
[Out] -1/48*((b*x^2+a)^2)^(1/2)*(-12*d^(5/2)*c^(3/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a*e*x^3+24*d^(3/2)*c^(5/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*b*e*x^3-6*(d*x^2+e*x+c)^(1/2)*d^(5/2)*a*e^2*x^4-48*(d*x^2+e*x+c)^(1/2)*d^(5/2)*b*c^2*x^4+12*(d*x^2+e*x+c)^(1/2)*d^(5/2)*a*c*e*x^3+3*d^(3/2)*c^(1/2)*ln((2*c+e*x+2*c^(1/2)*(d*x^2+e*x+c)^(1/2))/x)*a*e^3*x^3+6*(d*x^2+e*x+c)^(3/2)*d^(3/2)*a*e^2*x^2+48*(d*x^2+e*x+c)^(3/2)*d^(3/2)*b*c^2*x^2-6*(d*x^2+e*x+c)^(1/2)*d^(3/2)*a*e^3*x^3-48*(d*x^2+e*x+c)^(1/2)*d^(3/2)*b*c^2*e*x^3-12*(d*x^2+e*x+c)^(3/2)*d^(3/2)*a*c*e*x-48*ln(1/2*(2*(d*x^2+e*x+c)^(1/2)*d^(1/2)+2*d*x+e)/d^(1/2))*b*c^3*d^2*x^3+16*(d*x^2+e*x+c)^(3/2)*d^(3/2)*a*c^2/d^(3/2)/x^3/c^3/(b*x^2+a)
```

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="maxima")
```

```
[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(4*c*d-%e^2>0)', see 'assume?' for more det
```

Fricas [A]

time = 0.70, size = 843, normalized size = 2.87

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="fricas")

[Out] [1/96*(48*b*c^3*sqrt(d)*x^3*log(8*d^2*x^2 + 8*d*x*e + 4*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 3*(a*x^3*e^3 + 4*(2*b*c^2 - a*c*d)*x^3*e)*sqrt(c)*log((4*c*d*x^2 + x^2*e^2 + 8*c*x*e - 4*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(c) + 8*c^2)/x^2) + 4*(3*a*c*x^2*e^2 - 2*a*c^2*x*e - 8*a*c^3 - 8*(3*b*c^3 + a*c^2*d)*x^2)*sqrt(d*x^2 + x*e + c))/(c^3*x^3), -1/96*(96*b*c^3*sqrt(-d)*x^3*arctan(1/2*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*x*e + c*d)) - 3*(a*x^3*e^3 + 4*(2*b*c^2 - a*c*d)*x^3*e)*sqrt(c)*log((4*c*d*x^2 + x^2*e^2 + 8*c*x*e - 4*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(c) + 8*c^2)/x^2) - 4*(3*a*c*x^2*e^2 - 2*a*c^2*x*e - 8*a*c^3 - 8*(3*b*c^3 + a*c^2*d)*x^2)*sqrt(d*x^2 + x*e + c))/(c^3*x^3), 1/48*(24*b*c^3*sqrt(d)*x^3*log(8*d^2*x^2 + 8*d*x*e + 4*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(d) + 4*c*d + e^2) + 3*(a*x^3*e^3 + 4*(2*b*c^2 - a*c*d)*x^3*e)*sqrt(-c)*arctan(1/2*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(-c)/(c*d*x^2 + c*x*e + c^2)) + 2*(3*a*c*x^2*e^2 - 2*a*c^2*x*e - 8*a*c^3 - 8*(3*b*c^3 + a*c^2*d)*x^2)*sqrt(d*x^2 + x*e + c))/(c^3*x^3), -1/48*(48*b*c^3*sqrt(-d)*x^3*arctan(1/2*sqrt(d*x^2 + x*e + c)*(2*d*x + e)*sqrt(-d)/(d^2*x^2 + d*x*e + c*d)) - 3*(a*x^3*e^3 + 4*(2*b*c^2 - a*c*d)*x^3*e)*sqrt(-c)*arctan(1/2*sqrt(d*x^2 + x*e + c)*(x*e + 2*c)*sqrt(-c)/(c*d*x^2 + c*x*e + c^2)) - 2*(3*a*c*x^2*e^2 - 2*a*c^2*x*e - 8*a*c^3 - 8*(3*b*c^3 + a*c^2*d)*x^2)*sqrt(d*x^2 + x*e + c))/(c^3*x^3)]

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x**2+e*x+c)**(1/2)*((b*x**2+a)**2)**(1/2)/x**4,x)

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 720 vs. 2(230) = 460.

time = 6.43, size = 720, normalized size = 2.45

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*x^2+e*x+c)^(1/2)*((b*x^2+a)^2)^(1/2)/x^4,x, algorithm="giac")

[Out] -b*sqrt(d)*log(abs(2*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*sqrt(d) + e))*sgn(b*x^2 + a) + 1/8*(8*b*c^2*e*sgn(b*x^2 + a) - 4*a*c*d*e*sgn(b*x^2 + a) + a*e^3*sgn(b*x^2 + a))*arctan(-(sqrt(d)*x - sqrt(d*x^2 + x*e + c))/sqrt(-c))/(sqrt(-c)*c^2) + 1/24*(24*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^5*b*c^2*sqrt(d)

```

*e*sgn(b*x^2 + a) + 12*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^5*a*c*d^(3/2)*e*
sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^4*b*c^3*d*sgn(b*x^2
+ a) + 48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^4*a*c^2*d^2*sgn(b*x^2 + a) -
48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^3*b*c^3*sqrt(d)*e*sgn(b*x^2 + a) +
48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^3*a*c^2*d^(3/2)*e*sgn(b*x^2 + a) - 9
6*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^2*b*c^4*d*sgn(b*x^2 + a) - 3*(sqrt(d)
*x - sqrt(d*x^2 + x*e + c))^5*a*sqrt(d)*e^3*sgn(b*x^2 + a) + 24*(sqrt(d)*x
- sqrt(d*x^2 + x*e + c))*b*c^4*sqrt(d)*e*sgn(b*x^2 + a) + 36*(sqrt(d)*x - s
qrt(d*x^2 + x*e + c))*a*c^3*d^(3/2)*e*sgn(b*x^2 + a) + 48*b*c^5*d*sgn(b*x^2
+ a) + 16*a*c^4*d^2*sgn(b*x^2 + a) + 48*(sqrt(d)*x - sqrt(d*x^2 + x*e + c)
)^2*a*c^2*d*e^2*sgn(b*x^2 + a) + 8*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))^3*a*
c*sqrt(d)*e^3*sgn(b*x^2 + a) + 3*(sqrt(d)*x - sqrt(d*x^2 + x*e + c))*a*c^2*
sqrt(d)*e^3*sgn(b*x^2 + a))/(((sqrt(d)*x - sqrt(d*x^2 + x*e + c))^2 - c)^3*
c^2*sqrt(d))

```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(bx^2 + a)^2} \sqrt{dx^2 + ex + c}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^4,x)

[Out] int((((a + b*x^2)^2)^(1/2)*(c + e*x + d*x^2)^(1/2))/x^4, x)

Chapter 4

Appendix

Local contents

4.1	Download section	260
4.2	Listing of Grading functions	260

4.1 Download section

The following zip files contain the raw integrals used in this test.

Mathematica format Mathematica_syntax.zip

Maple and Mupad format Maple_syntax.zip

Sympy format SYMPY_syntax.zip

Sage math format SAGE_syntax.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*           is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*           antiderivative*)
(* "A" if result can be considered optimal*)
```

```

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A","none"}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","none"}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)

```



```

ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]

```

4.2.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
#do NOT call ExpnType() if leaf size is too large. Recursion problem
if leaf_count_result > 500000 then
      return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

```

```

    if debug then
        print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
    fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#   is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
#   antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A","";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of r
                    convert(leaf_count_result,string)," vs. $2 (" ,
                    convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_co

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex

```

```

    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A","";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string),")=",convert(2*leaf_cou

    fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                  convert(ExpnType_result,string)," vs. order ",
                  convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function

```

```

# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,

```

```

    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

```



```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

4.2.4 SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fricas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or instan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```