

Computer algebra independent integration tests

Summer 2022 edition

4-Trig-functions/4.1-Sine/65-4.1.0-a-sin-^m-b-trg-ⁿ

Nasser M. Abbasi

September 27, 2022

Compiled on September 27, 2022 at 2:13am

Contents

1	Introduction	3
2	detailed summary tables of results	19
3	Listing of integrals	151
4	Appendix	2113

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	list of integrals that has no closed form antiderivative	11
1.5	List of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	13
1.7	Timing	13
1.8	Verification	14
1.9	Important notes about some of the results	14
1.10	Design of the test system	17

This report gives the result of running the computer algebra independent integration test. The download section in the appendix contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [538]. This is test number [65].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.1 (June 29, 2022) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.0.1 on windows 10.
3. Maple 2022.1 (June 1, 2022) on windows 10.
4. Maxima 5.46 (April 13, 2022) using Lisp SBCL 2.1.11.debian on Linux via sagemath 9.6.
5. Fricas 1.3.8 (June 21, 2022) based on sbcl 2.1.11.debian on Linux via sagemath 9.6.
6. Giac/Xcas 1.9.0-13 (July 3, 2022) on Linux via sagemath 9.6.
7. Sympy 1.10.1 (March 20, 2022) Using Python 3.10.4 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly from Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (538)	0.00 (0)
Mathematica	100.00 (538)	0.00 (0)
Maple	82.34 (443)	17.66 (95)
Fricas	78.07 (420)	21.93 (118)
Mupad	46.10 (248)	53.90 (290)
Maxima	45.17 (243)	54.83 (295)
Giac	39.59 (213)	60.41 (325)
Sympy	18.96 (102)	81.04 (436)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	99.44	0.00	0.56	0.00
Mathematica	76.77	2.97	20.26	0.00
Maple	45.35	19.33	17.66	17.66
Maxima	44.05	1.12	0.00	54.83
Fricas	39.03	14.13	24.91	21.93
Giac	29.00	10.59	0.00	60.41
Sympy	8.55	10.41	0.00	81.04
Mupad	N/A	46.10	0.00	53.90

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and

Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	0	0.00 %	0.00 %	0.00 %
Maple	95	100.00 %	0.00 %	0.00 %
Fricas	118	83.90 %	11.02 %	5.08 %
Giac	325	98.77 %	1.23 %	0.00 %
Maxima	295	99.66 %	0.34 %	0.00 %
Sympy	436	53.67 %	27.06 %	19.27 %
Mupad	290	100.00 %	0.00 %	0.00 %

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.



Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.05	78.71	1.00	68.00	1.00
Mathematica	0.85	107.44	1.57	57.00	0.88
Maple	0.21	236.12	2.86	108.00	1.70
Maxima	0.34	51.16	1.04	40.00	0.96
Fricas	0.93	150.09	1.67	74.00	1.28
Sympy	2.99	238.61	5.58	66.00	1.68
Giac	4.20	75.99	1.62	55.00	1.15
Mupad	1.10	52.19	1.14	38.50	0.92

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used from the above table.



1.4 list of integrals that has no closed form antiderivative

{

1.5 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {486, 488, 489, 490, 491, 496, 497, 498, 499, 501, 502}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for `Rubi` and `Mathematica`.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
```

```
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified.

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,..}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	20
2.2	Detailed conclusion table per each integral for all CAS systems	27
2.3	Detailed conclusion table specific for Rubi results	135

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	21
2.1.2	Mathematica	21
2.1.3	Maple	22
2.1.4	Maxima	23
2.1.5	FriCAS	23
2.1.6	Sympy	24
2.1.7	Giac	25
2.1.8	Mupad	26

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538 }

B grade: { }

C grade: { 35, 36, 37 }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 38, 39, 40, 41, 42, 43, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 139, 141, 143, 145, 146, 147, 148, 149, 151, 152, 154, 156, 157, 158, 159, 160, 161, 163, 164, 165, 167, 169, 171, 172, 173, 174, 175, 177, 179, 180, 181, 182, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 204, 205, 206, 207, 208, 209, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 244, 252, 253, 254, 255, 256, 264, 265, 266, 273, 274, 275, 284, 285, 286, 288, 297, 298, 299, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349,

350, 351, 352, 353, 354, 355, 356, 357, 358, 361, 362, 363, 364, 365, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 483, 484, 485, 487, 492, 493, 494, 495, 500, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538 }

B grade: { 44, 87, 88, 126, 150, 153, 155, 176, 178, 183, 210, 221, 359, 360, 366, 496 }

C grade: { 35, 36, 37, 138, 140, 142, 144, 162, 166, 168, 170, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 243, 245, 246, 247, 248, 249, 250, 251, 257, 258, 259, 260, 261, 262, 263, 267, 268, 269, 270, 271, 272, 276, 277, 278, 279, 280, 281, 282, 283, 287, 289, 290, 291, 292, 293, 294, 295, 296, 300, 301, 302, 303, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 477, 478, 479, 480, 481, 482, 486, 488, 489, 490, 491, 497, 498, 499, 501, 502 }

F grade: { }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 75, 77, 80, 81, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 111, 113, 115, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 197, 200, 204, 206, 211, 212, 213, 214, 215, 216, 217, 232, 234, 239, 240, 252, 253, 254, 255, 256, 264, 265, 266, 267, 268, 269, 270, 273, 274, 275, 284, 285, 286, 291, 292, 294, 295, 297, 298, 299, 342, 357, 374, 388, 400, 410, 411, 412, 413, 424, 425, 426, 427, 437, 438, 439, 440, 453, 454, 455, 456, 457, 469, 470, 471, 476, 477, 478, 479 }

B grade: { 10, 74, 76, 78, 79, 82, 110, 112, 114, 116, 163, 196, 198, 199, 201, 202, 203, 205, 207, 208, 209, 210, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 233, 235, 236, 237, 238, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 257, 258, 259, 260, 261, 276, 277, 278, 279, 280, 281, 288, 293, 371, 372, 373, 375, 376, 377, 385, 386, 387, 389, 390, 397, 398, 399, 401, 402, 403, 414, 415, 416, 428, 429, 430, 441, 442, 443, 458, 459, 460, 461, 462, 463, 464, 465, 468, 480, 481, 482, 494 }

C grade: { 262, 263, 271, 272, 282, 283, 287, 289, 290, 296, 300, 301, 302, 303, 341, 356, 378, 379, 380, 381, 382, 383, 384, 391, 392, 393, 394, 395, 396, 404, 405, 406, 407, 408, 409, 417, 418, 419, 420, 421, 422, 423, 431, 432, 433, 434, 435, 436, 444, 445, 446, 447, 448, 449, 450, 451, 452, 466, 467, 472, 473, 474, 475, 493, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537 }

F grade: { 33, 34, 35, 36, 37, 38, 39, 40, 41, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335,

336, 337, 338, 339, 340, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 538 }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 76, 77, 78, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 112, 114, 115, 116, 118, 119, 120, 121, 122, 123, 124, 125, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 204, 205, 206, 207, 208, 209, 210, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 340, 341, 342, 355, 356, 357, 371, 372, 373, 374, 375, 376, 377, 385, 386, 387, 388, 389, 390, 397, 398, 399, 400, 401, 402, 403, 410, 411, 412, 413, 414, 415, 416, 424, 425, 426, 427, 428, 429, 430, 437, 438, 439, 440, 441, 442, 443, 492, 493, 494 }

B grade: { 75, 79, 111, 113, 117, 126 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 378, 379, 380, 381, 382, 383, 384, 391, 392, 393, 394, 395, 396, 404, 405, 406, 407, 408, 409, 417, 418, 419, 420, 421, 422, 423, 431, 432, 433, 434, 435, 436, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 55, 56, 57, 58, 59, 60, 61, 62, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 84, 85, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 105, 106, 107, 108, 109, 110, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 141, 142, 143, 144, 145, 146, 147, 148, 149, 151, 155, 157, 158, 159, 160, 161, 163, 165, 167, 169, 170, 171, 172, 173, 174, 175, 183, 187, 188, 189, 190, 191, 192, 193, 204, 205, 206, 207, 208, 209, 210, 221, 222, 223, 225, 231, 242, 251, 252, 253, 254, 255, 256, 264, 265, 266, 273, 274, 275, 284, 285, 286, 288, 297, 298, 299, 324, 327, 328, 329, 332, 333, 334, 335, 340, 341, 342, 355, 356, 357, 371, 372, 373, 374, 385, 386, 387,

388, 397, 398, 399, 400, 410, 411, 412, 413, 424, 425, 426, 427, 437, 438, 439, 440, 453, 454, 455, 468, 469, 470, 471, 492, 493, 494 }

B grade: { 53, 54, 63, 83, 86, 104, 111, 126, 127, 128, 140, 150, 152, 153, 154, 156, 162, 164, 166, 168, 176, 177, 178, 179, 180, 181, 182, 184, 185, 186, 224, 226, 227, 228, 229, 230, 243, 244, 245, 246, 247, 248, 249, 250, 262, 263, 271, 272, 282, 283, 287, 289, 290, 296, 300, 301, 302, 303, 375, 376, 377, 389, 390, 401, 402, 403, 414, 415, 416, 428, 429, 430, 441, 442, 443, 476 }

C grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 269, 270, 293, 294, 295, 378, 379, 380, 381, 382, 383, 384, 391, 392, 393, 394, 395, 396, 404, 405, 406, 407, 408, 409, 417, 418, 419, 420, 421, 422, 423, 431, 432, 433, 434, 435, 436, 444, 445, 446, 447, 448, 449, 458, 459, 460, 480, 481, 482, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537 }

F grade: { 33, 34, 35, 36, 37, 38, 39, 40, 41, 257, 258, 259, 260, 261, 267, 268, 276, 277, 278, 279, 280, 281, 291, 292, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 325, 326, 330, 331, 336, 337, 338, 339, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 450, 451, 452, 456, 457, 461, 462, 463, 464, 465, 466, 467, 472, 473, 474, 475, 477, 478, 479, 483, 484, 485, 486, 487, 488, 489, 490, 491, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 538 }

2.1.6 Sympy

A grade: { 1, 3, 5, 7, 42, 43, 44, 49, 50, 51, 52, 66, 68, 70, 80, 81, 82, 83, 98, 99, 100, 101, 103, 104, 149, 151, 157, 158, 159, 160, 165, 175, 176, 178, 179, 186, 187, 188, 204, 205, 206, 207, 208, 252, 254, 255 }

B grade: { 2, 4, 6, 8, 58, 59, 60, 61, 62, 67, 69, 89, 90, 91, 92, 102, 120, 121, 122, 123, 124, 125, 133, 134, 135, 136, 137, 138, 139, 145, 146, 147, 148, 150, 161, 162, 163, 164, 171, 172, 173, 174, 177, 185, 189, 190, 191, 192, 253, 256, 340, 341, 342, 355, 356, 357 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 45, 46, 47, 48, 53, 54, 55, 56, 57, 63, 64, 65, 71, 72, 73, 74, 75, 76, 77, 78, 79, 84, 85, 86, 87, 88, 93, 94, 95, 96, 97, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 126, 127, 128, 129, 130, 131, 132, 140, 141, 142, 143, 144, 152, 153, 154, 155, 156, 166, 167, 168, 169, 170, 180, 181, 182, 183, 184, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409,

410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538 }

2.1.7 Giac

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 100, 101, 102, 103, 104, 110, 118, 123, 125, 133, 134, 135, 136, 137, 139, 140, 141, 142, 143, 144, 149, 151, 157, 158, 159, 160, 161, 163, 164, 165, 166, 167, 168, 169, 170, 177, 178, 179, 185, 186, 188, 189, 204, 205, 226, 227, 247, 248, 252, 253, 254, 255, 256, 342, 357, 371, 372, 373, 374, 375, 376, 377, 385, 386, 387, 388, 389, 390, 397, 398, 399, 401, 402, 403, 410, 411, 412, 414, 415, 416, 424, 425, 426, 428, 429, 430, 437, 438, 439, 441, 442, 443 }

B grade: { 98, 99, 105, 106, 107, 108, 109, 111, 112, 113, 114, 115, 116, 117, 119, 120, 121, 122, 124, 126, 127, 128, 129, 130, 131, 132, 138, 145, 146, 147, 148, 150, 152, 153, 154, 155, 156, 162, 171, 172, 173, 174, 175, 176, 180, 181, 182, 183, 184, 340, 341, 355, 356, 400, 413, 427, 440 }

C grade: { }

F grade: { 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 187, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 249, 250, 251, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 378, 379, 380, 381, 382, 383, 384, 391, 392, 393, 394, 395, 396, 404, 405, 406, 407, 408, 409, 417, 418, 419, 420, 421, 422, 423, 431, 432, 433, 434, 435, 436, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 511, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 28, 29, 39, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 207, 208, 209, 210, 221, 252, 253, 254, 255, 256, 264, 265, 266, 273, 274, 275, 284, 285, 286, 287, 288, 289, 290, 297, 298, 299, 300, 301, 302, 303, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 340, 341, 342, 355, 356, 357, 374, 381, 388, 399, 400, 413, 427, 440, 453, 454, 455, 468, 469, 470, 471, 476, 492, 493, 494, 511 }

C grade: { }

F grade: { 25, 26, 27, 30, 31, 32, 33, 34, 35, 36, 37, 38, 40, 41, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 257, 258, 259, 260, 261, 262, 263, 267, 268, 269, 270, 271, 272, 276, 277, 278, 279, 280, 281, 282, 283, 291, 292, 293, 294, 295, 296, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 337, 338, 339, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 375, 376, 377, 378, 379, 380, 382, 383, 384, 385, 386, 387, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 472, 473, 474, 475, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, 505, 506, 507, 508, 509, 510, 512, 513, 514, 515, 516, 517, 518, 519, 520, 521, 522, 523, 524, 525, 526, 527, 528, 529, 530, 531, 532, 533, 534, 535, 536, 537, 538 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column **N.S.** in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, **Mathematica** was abbrevi-

	Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
viated to MMA .	grade	A	A	A	A	A	A	A	A	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	11	11	22	12	11	11	14	11	11
	N.S.	1	1.00	2.00	1.09	1.00	1.00	1.27	1.00	1.00
	time (sec)	N/A	0.003	0.008	0.067	0.327	0.377	0.044	4.759	0.020

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	23	27	24	23	46	18	18
N.S.	1	1.00	0.92	1.08	0.96	0.92	1.84	0.72	0.72
time (sec)	N/A	0.006	0.021	0.037	0.287	0.406	0.075	5.378	0.430

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	29	22	22	22	37	25	24
N.S.	1	1.00	1.07	0.81	0.81	0.81	1.37	0.93	0.89
time (sec)	N/A	0.007	0.010	0.093	0.283	0.387	0.103	5.553	0.353

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	33	38	33	36	95	32	50
N.S.	1	1.00	0.72	0.83	0.72	0.78	2.07	0.70	1.09
time (sec)	N/A	0.013	0.031	0.089	0.287	0.392	0.171	4.853	0.470

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	44	32	34	34	60	38	32
N.S.	1	1.00	1.05	0.76	0.81	0.81	1.43	0.90	0.76
time (sec)	N/A	0.009	0.011	0.102	0.287	0.385	0.249	5.247	0.358

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	45	48	48	47	139	46	43
N.S.	1	1.00	0.67	0.72	0.72	0.70	2.07	0.69	0.64
time (sec)	N/A	0.021	0.032	0.109	0.274	0.388	0.405	3.910	0.578

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	59	42	44	44	80	50	43
N.S.	1	1.00	1.09	0.78	0.81	0.81	1.48	0.93	0.80
time (sec)	N/A	0.011	0.009	0.112	0.333	0.374	0.607	4.342	0.375

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	55	58	59	56	184	60	90
N.S.	1	1.00	0.62	0.66	0.67	0.64	2.09	0.68	1.02
time (sec)	N/A	0.034	0.040	0.135	0.316	0.428	0.898	4.689	1.505

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	45	84	0	73	0	0	34
N.S.	1	1.00	0.75	1.40	0.00	1.22	0.00	0.00	0.57
time (sec)	N/A	0.018	0.053	0.105	0.000	0.147	0.000	0.000	0.483

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	35	118	0	68	0	0	34
N.S.	1	1.00	0.85	2.88	0.00	1.66	0.00	0.00	0.83
time (sec)	N/A	0.014	0.032	0.060	0.000	0.101	0.000	0.000	0.414

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	33	72	0	60	0	0	34
N.S.	1	1.00	0.80	1.76	0.00	1.46	0.00	0.00	0.83
time (sec)	N/A	0.012	0.031	0.057	0.000	0.110	0.000	0.000	0.404

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	77	0	55	0	0	15
N.S.	1	1.00	1.11	4.05	0.00	2.89	0.00	0.00	0.79
time (sec)	N/A	0.006	0.020	0.106	0.000	0.098	0.000	0.000	0.366

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	21	57	0	47	0	0	15
N.S.	1	1.00	1.11	3.00	0.00	2.47	0.00	0.00	0.79
time (sec)	N/A	0.006	0.021	0.052	0.000	0.099	0.000	0.000	0.387

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	32	110	0	81	0	0	34
N.S.	1	1.00	0.86	2.97	0.00	2.19	0.00	0.00	0.92
time (sec)	N/A	0.011	0.042	0.051	0.000	0.099	0.000	0.000	0.473

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	33	72	0	97	0	0	34
N.S.	1	1.00	0.80	1.76	0.00	2.37	0.00	0.00	0.83
time (sec)	N/A	0.012	0.039	0.051	0.000	0.092	0.000	0.000	0.585

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	51	132	0	132	0	0	34
N.S.	1	1.00	0.85	2.20	0.00	2.20	0.00	0.00	0.57
time (sec)	N/A	0.017	0.039	0.060	0.000	0.103	0.000	0.000	0.560

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	55	104	0	87	0	0	42
N.S.	1	1.00	0.79	1.49	0.00	1.24	0.00	0.00	0.60
time (sec)	N/A	0.020	0.083	0.059	0.000	0.103	0.000	0.000	0.488

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	44	142	0	80	0	0	42
N.S.	1	1.00	0.94	3.02	0.00	1.70	0.00	0.00	0.89
time (sec)	N/A	0.013	0.065	0.056	0.000	0.123	0.000	0.000	0.450

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	40	88	0	72	0	0	42
N.S.	1	1.00	0.85	1.87	0.00	1.53	0.00	0.00	0.89
time (sec)	N/A	0.013	0.029	0.047	0.000	0.107	0.000	0.000	0.440

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	24	91	0	63	0	0	18
N.S.	1	1.00	1.14	4.33	0.00	3.00	0.00	0.00	0.86
time (sec)	N/A	0.007	0.013	0.094	0.000	0.117	0.000	0.000	0.368

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	24	69	0	55	0	0	18
N.S.	1	1.00	1.14	3.29	0.00	2.62	0.00	0.00	0.86
time (sec)	N/A	0.006	0.016	0.065	0.000	0.092	0.000	0.000	0.377

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	39	132	0	99	0	0	42
N.S.	1	1.00	0.91	3.07	0.00	2.30	0.00	0.00	0.98
time (sec)	N/A	0.013	0.058	0.055	0.000	0.110	0.000	0.000	0.511

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	47	47	43	88	0	115	0	0	42
N.S.	1	1.00	0.91	1.87	0.00	2.45	0.00	0.00	0.89
time (sec)	N/A	0.012	0.079	0.057	0.000	0.094	0.000	0.000	0.613

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	55	160	0	158	0	0	42
N.S.	1	1.00	0.79	2.29	0.00	2.26	0.00	0.00	0.60
time (sec)	N/A	0.021	0.210	0.063	0.000	0.104	0.000	0.000	0.594

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	80	108	0	105	0	0	-1
N.S.	1	1.00	0.78	1.05	0.00	1.02	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.111	0.063	0.000	0.102	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	66	152	0	101	0	0	-1
N.S.	1	1.00	0.88	2.03	0.00	1.35	0.00	0.00	-0.01
time (sec)	N/A	0.023	0.082	0.059	0.000	0.102	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	62	97	0	81	0	0	-1
N.S.	1	1.00	0.83	1.29	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.023	0.041	0.067	0.000	0.111	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	98	0	67	0	0	36
N.S.	1	1.00	0.98	2.28	0.00	1.56	0.00	0.00	0.84
time (sec)	N/A	0.013	0.019	0.138	0.000	0.115	0.000	0.000	0.395

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	74	0	62	0	0	36
N.S.	1	1.00	0.98	1.72	0.00	1.44	0.00	0.00	0.84
time (sec)	N/A	0.013	0.023	0.066	0.000	0.092	0.000	0.000	0.485

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	54	141	0	108	0	0	-1
N.S.	1	1.00	0.74	1.93	0.00	1.48	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.038	0.060	0.000	0.096	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	55	105	0	127	0	0	-1
N.S.	1	1.00	0.71	1.36	0.00	1.65	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.058	0.076	0.000	0.100	0.000	0.000	0.000

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	105	68	168	0	170	0	0	-1
N.S.	1	1.00	0.65	1.60	0.00	1.62	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.117	0.063	0.000	0.119	0.000	0.000	0.000

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.011	0.042	0.016	0.000	0.000	0.000	0.000	0.000

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.012	0.027	0.015	0.000	0.000	0.000	0.000	0.000

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	C	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	517	58	55	0	0	0	0	0	-1
N.S.	1	0.11	0.11	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.011	0.026	0.016	0.000	0.000	0.000	0.000	0.000

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	C	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	252	58	55	0	0	0	0	0	-1
N.S.	1	0.23	0.22	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.011	0.031	0.014	0.000	0.000	0.000	0.000	0.000

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	C	C	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	271	56	53	0	0	0	0	0	-1
N.S.	1	0.21	0.20	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.011	0.031	0.014	0.000	0.000	0.000	0.000	0.000

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	56	56	53	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.011	0.031	0.016	0.000	0.000	0.000	0.000	0.000

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	0	0	0	0	0	54
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	0.86
time (sec)	N/A	0.011	0.034	0.020	0.000	0.000	0.000	0.000	0.783

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	63	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.012	0.030	0.020	0.000	0.000	0.000	0.000	0.000

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	76	0	0	0	0	0	-1
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.061	0.032	0.000	0.000	0.000	0.000	0.000

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	22	24	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	1.47	1.60	0.87
time (sec)	N/A	0.014	0.003	0.045	0.288	0.405	0.152	7.055	0.056

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	22	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	1.47	0.87	0.87
time (sec)	N/A	0.012	0.003	0.030	0.283	0.370	0.095	4.688	0.037

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	37	14	13	13	19	13	28
N.S.	1	1.00	2.47	0.93	0.87	0.87	1.27	0.87	1.87
time (sec)	N/A	0.007	0.011	0.033	0.286	0.391	0.062	3.890	0.455

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	12	12	12	12	18	14	0	18	16
N.S.	1	1.00	1.00	1.00	1.50	1.17	0.00	1.50	1.33
time (sec)	N/A	0.003	0.006	0.020	0.298	0.466	0.000	4.385	0.506

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	10	10	10	11	12	12	0	12	20
N.S.	1	1.00	1.00	1.10	1.20	1.20	0.00	1.20	2.00
time (sec)	N/A	0.007	0.006	0.023	0.288	0.344	0.000	4.783	0.515

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	17	13	0	13	13
N.S.	1	1.00	1.00	0.93	1.13	0.87	0.00	0.87	0.87
time (sec)	N/A	0.015	0.009	0.036	0.285	0.356	0.000	3.840	0.382

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	13	0	13	13
N.S.	1	1.00	1.00	0.93	0.87	0.87	0.00	0.87	0.87
time (sec)	N/A	0.012	0.007	0.036	0.290	0.344	0.000	4.053	0.448

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	47	60	46	53	88	54	45
N.S.	1	1.00	0.77	0.98	0.75	0.87	1.44	0.89	0.74
time (sec)	N/A	0.027	0.119	0.161	0.303	0.380	1.255	4.095	0.398

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	50	36	43	66	54	36
N.S.	1	1.00	0.80	1.09	0.78	0.93	1.43	1.17	0.78
time (sec)	N/A	0.025	0.072	0.128	0.295	0.359	0.586	3.571	0.399

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	27	40	26	33	44	26	26
N.S.	1	1.00	0.87	1.29	0.84	1.06	1.42	0.84	0.84
time (sec)	N/A	0.022	0.047	0.096	0.295	0.379	0.380	4.456	0.359

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	21	20	13	13
N.S.	1	1.00	1.00	0.93	0.87	1.40	1.33	0.87	0.87
time (sec)	N/A	0.012	0.003	0.026	0.293	0.374	0.244	2.548	0.023

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	23	19	18	31	0	18	14
N.S.	1	1.00	1.64	1.36	1.29	2.21	0.00	1.29	1.00
time (sec)	N/A	0.006	0.007	0.034	0.509	0.398	0.000	2.927	0.375

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	13	29	0	13	13
N.S.	1	1.00	1.00	1.47	0.87	1.93	0.00	0.87	0.87
time (sec)	N/A	0.020	0.005	0.042	0.296	0.360	0.000	3.366	0.381

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	56	42	26	39	0	26	25
N.S.	1	1.00	1.81	1.35	0.84	1.26	0.00	0.84	0.81
time (sec)	N/A	0.021	0.034	0.064	0.283	0.357	0.000	2.849	0.387

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	77	60	36	51	0	36	35
N.S.	1	1.00	1.67	1.30	0.78	1.11	0.00	0.78	0.76
time (sec)	N/A	0.024	0.033	0.078	0.296	0.416	0.000	3.676	0.407

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	98	78	46	61	0	46	45
N.S.	1	1.00	1.61	1.28	0.75	1.00	0.00	0.75	0.74
time (sec)	N/A	0.026	0.029	0.044	0.293	0.402	0.000	3.384	0.405

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	52	64	48	57	189	60	89
N.S.	1	1.00	0.59	0.73	0.55	0.65	2.15	0.68	1.01
time (sec)	N/A	0.044	0.110	0.153	0.302	0.380	0.873	4.064	1.517

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	40	54	37	47	136	46	43
N.S.	1	1.00	0.60	0.81	0.55	0.70	2.03	0.69	0.64
time (sec)	N/A	0.037	0.066	0.124	0.344	0.372	0.401	6.040	0.558

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	23	43	24	36	92	18	50
N.S.	1	1.00	0.50	0.93	0.52	0.78	2.00	0.39	1.09
time (sec)	N/A	0.026	0.027	0.037	0.289	0.374	0.166	4.528	0.460

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	23	27	24	23	46	18	18
N.S.	1	1.00	0.92	1.08	0.96	0.92	1.84	0.72	0.72
time (sec)	N/A	0.007	0.020	0.000	0.319	0.375	0.074	4.074	0.002

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	28	34	36	3160	36	27
N.S.	1	1.00	1.00	1.22	1.48	1.57	137.39	1.57	1.17
time (sec)	N/A	0.010	0.010	0.034	0.286	0.395	26.090	3.782	0.457

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	34	48	46	61	0	48	69
N.S.	1	1.00	1.00	1.41	1.35	1.79	0.00	1.41	2.03
time (sec)	N/A	0.016	0.013	0.046	0.313	0.408	0.000	3.665	1.225

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	55	66	65	71	0	82	125
N.S.	1	1.00	1.00	1.20	1.18	1.29	0.00	1.49	2.27
time (sec)	N/A	0.031	0.034	0.069	0.297	0.418	0.000	4.252	6.539

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	76	84	91	84	0	73	177
N.S.	1	1.00	1.00	1.11	1.20	1.11	0.00	0.96	2.33
time (sec)	N/A	0.041	0.050	0.081	0.284	0.395	0.000	3.683	7.325

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	48	34	36	26	44	27	25
N.S.	1	1.00	1.55	1.10	1.16	0.84	1.42	0.87	0.81
time (sec)	N/A	0.022	0.092	0.105	0.287	0.371	0.856	3.632	0.373

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	27	34	26	26	46	27	26
N.S.	1	1.00	0.87	1.10	0.84	0.84	1.48	0.87	0.84
time (sec)	N/A	0.022	0.069	0.086	0.290	0.346	0.576	3.350	0.378

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	35	34	26	26	44	26	37
N.S.	1	1.00	1.13	1.10	0.84	0.84	1.42	0.84	1.19
time (sec)	N/A	0.021	0.014	0.065	0.281	0.396	0.377	3.908	0.511

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	27	34	26	26	46	27	26
N.S.	1	1.00	0.87	1.10	0.84	0.84	1.48	0.87	0.84
time (sec)	N/A	0.022	0.047	0.049	0.303	0.364	0.246	5.592	0.344

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	24	20	13	13
N.S.	1	1.00	1.00	0.93	0.87	1.60	1.33	0.87	0.87
time (sec)	N/A	0.012	0.002	0.036	0.288	0.373	0.151	4.469	0.368

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	25	25	25	25	0	29	25
N.S.	1	1.00	0.89	0.89	0.89	0.89	0.00	1.04	0.89
time (sec)	N/A	0.015	0.015	0.037	0.299	0.429	0.000	4.643	0.429

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	21	40	19	22	0	23	20
N.S.	1	1.00	1.00	1.90	0.90	1.05	0.00	1.10	0.95
time (sec)	N/A	0.014	0.020	0.039	0.291	0.357	0.000	5.702	0.486

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	25	23	31	34	0	42	27
N.S.	1	1.00	0.93	0.85	1.15	1.26	0.00	1.56	1.00
time (sec)	N/A	0.010	0.020	0.040	0.285	0.404	0.000	4.817	0.383

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	60	25	25	0	25	23
N.S.	1	1.00	1.00	2.22	0.93	0.93	0.00	0.93	0.85
time (sec)	N/A	0.014	0.021	0.046	0.303	0.388	0.000	4.579	0.448

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	39	25	0	25	13
N.S.	1	1.00	1.00	1.47	2.60	1.67	0.00	1.67	0.87
time (sec)	N/A	0.019	0.004	0.060	0.294	0.348	0.000	5.143	0.383

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	78	25	25	0	25	25
N.S.	1	1.00	1.00	2.52	0.81	0.81	0.00	0.81	0.81
time (sec)	N/A	0.021	0.038	0.062	0.292	0.357	0.000	3.994	0.540

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	28	42	49	25	0	25	25
N.S.	1	1.00	0.90	1.35	1.58	0.81	0.00	0.81	0.81
time (sec)	N/A	0.025	0.027	0.071	0.301	0.358	0.000	4.812	0.398

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	31	96	25	25	0	25	25
N.S.	1	1.00	1.00	3.10	0.81	0.81	0.00	0.81	0.81
time (sec)	N/A	0.023	0.025	0.088	0.297	0.351	0.000	4.819	0.631

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	28	60	59	25	0	25	35
N.S.	1	1.00	0.90	1.94	1.90	0.81	0.00	0.81	1.13
time (sec)	N/A	0.022	0.032	0.092	0.287	0.349	0.000	3.923	0.418

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	47	78	46	63	88	82	45
N.S.	1	1.00	0.77	1.28	0.75	1.03	1.44	1.34	0.74
time (sec)	N/A	0.027	0.152	0.127	0.281	0.386	3.632	3.385	0.379

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	68	36	53	66	68	36
N.S.	1	1.00	0.80	1.48	0.78	1.15	1.43	1.48	0.78
time (sec)	N/A	0.025	0.086	0.129	0.284	0.382	1.245	3.971	0.383

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	27	58	26	41	44	26	26
N.S.	1	1.00	0.87	1.87	0.84	1.32	1.42	0.84	0.84
time (sec)	N/A	0.022	0.054	0.089	0.328	0.370	0.576	3.965	0.032

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	31	20	13	13
N.S.	1	1.00	1.00	0.93	0.87	2.07	1.33	0.87	0.87
time (sec)	N/A	0.015	0.002	0.042	0.287	0.340	0.233	3.730	0.359

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	31	54	41	42	0	41	38
N.S.	1	1.00	0.78	1.35	1.02	1.05	0.00	1.02	0.95
time (sec)	N/A	0.027	0.089	0.052	0.520	0.347	0.000	3.569	0.479

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	38	28	29	46	0	29	24
N.S.	1	1.00	1.36	1.00	1.04	1.64	0.00	1.04	0.86
time (sec)	N/A	0.011	0.008	0.050	0.528	0.375	0.000	5.027	0.401

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	13	39	0	13	13
N.S.	1	1.00	1.00	1.47	0.87	2.60	0.00	0.87	0.87
time (sec)	N/A	0.019	0.006	0.063	0.294	0.350	0.000	3.649	0.411

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	77	42	26	49	0	26	25
N.S.	1	1.00	2.48	1.35	0.84	1.58	0.00	0.84	0.81
time (sec)	N/A	0.022	0.031	0.078	0.296	0.355	0.000	3.942	0.398

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	98	60	36	61	0	36	35
N.S.	1	1.00	2.13	1.30	0.78	1.33	0.00	0.78	0.76
time (sec)	N/A	0.024	0.031	0.046	0.302	0.365	0.000	4.682	0.428

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	111	62	82	48	66	231	74	109
N.S.	1	1.00	0.56	0.74	0.43	0.59	2.08	0.67	0.98
time (sec)	N/A	0.067	0.139	0.123	0.286	0.401	1.764	3.878	1.936

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	33	72	33	56	189	32	90
N.S.	1	1.00	0.37	0.80	0.37	0.62	2.10	0.36	1.00
time (sec)	N/A	0.059	0.035	0.099	0.295	0.389	1.665	3.428	1.504

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	40	61	37	46	136	46	43
N.S.	1	1.00	0.58	0.88	0.54	0.67	1.97	0.67	0.62
time (sec)	N/A	0.048	0.051	0.066	0.334	0.359	0.569	3.256	0.524

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	33	38	33	36	95	32	50
N.S.	1	1.00	0.72	0.83	0.72	0.78	2.07	0.70	1.09
time (sec)	N/A	0.013	0.015	0.000	0.292	0.333	0.169	3.051	0.424

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	38	46	48	0	48	53
N.S.	1	1.00	1.00	1.00	1.21	1.26	0.00	1.26	1.39
time (sec)	N/A	0.018	0.013	0.044	0.278	0.373	0.000	3.288	0.581

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	40	58	56	74	0	58	98
N.S.	1	1.00	0.82	1.18	1.14	1.51	0.00	1.18	2.00
time (sec)	N/A	0.021	0.071	0.047	0.289	0.389	0.000	4.582	3.969

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	45	76	71	74	0	63	126
N.S.	1	1.00	0.82	1.38	1.29	1.35	0.00	1.15	2.29
time (sec)	N/A	0.035	0.092	0.073	0.294	0.383	0.000	4.796	6.585

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	99	94	91	84	0	73	177
N.S.	1	1.00	1.27	1.21	1.17	1.08	0.00	0.94	2.27
time (sec)	N/A	0.053	0.022	0.083	0.283	0.384	0.000	4.212	7.380

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	64	112	111	94	0	107	229
N.S.	1	1.00	0.65	1.13	1.12	0.95	0.00	1.08	2.31
time (sec)	N/A	0.063	0.224	0.109	0.286	0.386	0.000	4.523	7.442

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	68	52	46	36	65	85	35
N.S.	1	1.00	1.48	1.13	1.00	0.78	1.41	1.85	0.76
time (sec)	N/A	0.028	0.269	0.187	0.294	0.366	3.368	4.235	0.409

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	52	36	36	68	82	36
N.S.	1	1.00	0.80	1.13	0.78	0.78	1.48	1.78	0.78
time (sec)	N/A	0.028	0.197	0.124	0.282	0.382	2.473	2.966	0.409

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	50	52	36	36	65	43	36
N.S.	1	1.00	1.09	1.13	0.78	0.78	1.41	0.93	0.78
time (sec)	N/A	0.028	0.021	0.106	0.301	0.362	2.819	3.724	0.477

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	52	36	36	68	68	36
N.S.	1	1.00	0.80	1.13	0.78	0.78	1.48	1.48	0.78
time (sec)	N/A	0.024	0.109	0.122	0.282	0.374	1.246	2.785	0.376

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	48	52	26	36	65	26	26
N.S.	1	1.00	1.55	1.68	0.84	1.16	2.10	0.84	0.84
time (sec)	N/A	0.021	0.087	0.102	0.284	0.427	0.848	3.427	0.383

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	52	36	36	68	54	36
N.S.	1	1.00	0.80	1.13	0.78	0.78	1.48	1.17	0.78
time (sec)	N/A	0.026	0.070	0.073	0.281	0.367	0.581	3.893	0.370

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	34	20	13	13
N.S.	1	1.00	1.00	0.93	0.87	2.27	1.33	0.87	0.87
time (sec)	N/A	0.012	0.003	0.057	0.289	0.388	0.353	3.484	0.047

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	35	35	37	35	0	226	53
N.S.	1	1.00	0.88	0.88	0.92	0.88	0.00	5.65	1.32
time (sec)	N/A	0.018	0.027	0.050	0.281	0.371	0.000	2.846	0.537

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	39	50	32	33	0	99	31
N.S.	1	1.00	1.05	1.35	0.86	0.89	0.00	2.68	0.84
time (sec)	N/A	0.023	0.027	0.045	0.286	0.362	0.000	4.665	0.495

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	33	51	41	54	0	182	37
N.S.	1	1.00	0.77	1.19	0.95	1.26	0.00	4.23	0.86
time (sec)	N/A	0.025	0.031	0.058	0.305	0.392	0.000	6.070	0.404

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	70	35	35	0	100	35
N.S.	1	1.00	1.00	1.84	0.92	0.92	0.00	2.63	0.92
time (sec)	N/A	0.017	0.024	0.063	0.293	0.389	0.000	5.552	0.535

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	37	35	54	44	0	226	38
N.S.	1	1.00	0.86	0.81	1.26	1.02	0.00	5.26	0.88
time (sec)	N/A	0.016	0.038	0.056	0.287	0.376	0.000	4.405	0.397

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	41	88	35	35	0	72	35
N.S.	1	1.00	1.00	2.15	0.85	0.85	0.00	1.76	0.85
time (sec)	N/A	0.016	0.024	0.071	0.290	0.351	0.000	4.815	0.542

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-2)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	59	35	0	48	13
N.S.	1	1.00	1.00	1.47	3.93	2.33	0.00	3.20	0.87
time (sec)	N/A	0.019	0.006	0.069	0.286	0.334	0.000	4.686	0.422

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	46	106	35	35	0	116	35
N.S.	1	1.00	1.00	2.30	0.76	0.76	0.00	2.52	0.76
time (sec)	N/A	0.025	0.026	0.083	0.296	0.344	0.000	5.303	0.594

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	38	42	69	35	0	93	25
N.S.	1	1.00	1.23	1.35	2.23	1.13	0.00	3.00	0.81
time (sec)	N/A	0.022	0.042	0.102	0.282	0.376	0.000	5.252	0.416

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	46	124	35	35	0	160	35
N.S.	1	1.00	1.00	2.70	0.76	0.76	0.00	3.48	0.76
time (sec)	N/A	0.027	0.034	0.051	0.288	0.380	0.000	3.545	0.767

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	38	60	79	35	0	139	35
N.S.	1	1.00	0.83	1.30	1.72	0.76	0.00	3.02	0.76
time (sec)	N/A	0.029	0.041	0.055	0.299	0.346	0.000	5.273	0.442

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	46	142	35	35	0	204	35
N.S.	1	1.00	1.00	3.09	0.76	0.76	0.00	4.43	0.76
time (sec)	N/A	0.028	0.026	0.073	0.287	0.372	0.000	5.323	1.022

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	38	78	89	35	0	183	45
N.S.	1	1.00	0.83	1.70	1.93	0.76	0.00	3.98	0.98
time (sec)	N/A	0.030	0.092	0.056	0.291	0.350	0.000	4.233	0.420

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	52	68	66	84	0	68	147
N.S.	1	1.00	0.79	1.03	1.00	1.27	0.00	1.03	2.23
time (sec)	N/A	0.029	0.140	0.079	0.283	0.388	0.000	4.081	7.242

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	50	96	45	45	0	144	50
N.S.	1	1.00	1.00	1.92	0.90	0.90	0.00	2.88	1.00
time (sec)	N/A	0.019	0.031	0.086	0.274	0.386	0.000	3.382	0.540

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	75	48	56	60	1085	145	88
N.S.	1	1.00	1.42	0.91	1.06	1.13	20.47	2.74	1.66
time (sec)	N/A	0.020	0.026	0.064	0.282	0.378	2.395	3.492	5.370

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	33	35	35	1086	170	66
N.S.	1	1.00	1.00	0.82	0.88	0.88	27.15	4.25	1.65
time (sec)	N/A	0.019	0.012	0.043	0.305	0.382	1.519	3.690	0.504

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	60	38	46	50	473	101	62
N.S.	1	1.00	1.58	1.00	1.21	1.32	12.45	2.66	1.63
time (sec)	N/A	0.020	0.022	0.046	0.280	0.381	0.972	3.307	1.685

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	23	25	25	369	25	35
N.S.	1	1.00	1.00	0.85	0.93	0.93	13.67	0.93	1.30
time (sec)	N/A	0.015	0.012	0.033	0.290	0.385	0.667	3.997	0.414

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	42	28	34	38	92	57	35
N.S.	1	1.00	1.83	1.22	1.48	1.65	4.00	2.48	1.52
time (sec)	N/A	0.010	0.014	0.045	0.294	0.378	0.444	4.247	0.462

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	19	12	11	13	17	12	26
N.S.	1	1.00	1.73	1.09	1.00	1.18	1.55	1.09	2.36
time (sec)	N/A	0.003	0.010	0.020	0.306	0.384	0.193	4.657	0.406

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	B	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	31	12	28	30	0	56	11
N.S.	1	1.00	2.82	1.09	2.55	2.73	0.00	5.09	1.00
time (sec)	N/A	0.007	0.017	0.050	0.289	0.358	0.000	4.112	0.386

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	42	30	36	52	0	55	23
N.S.	1	1.00	1.83	1.30	1.57	2.26	0.00	2.39	1.00
time (sec)	N/A	0.015	0.022	0.052	0.296	0.339	0.000	3.863	0.084

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	36	23	40	56	0	124	35
N.S.	1	1.00	1.33	0.85	1.48	2.07	0.00	4.59	1.30
time (sec)	N/A	0.015	0.025	0.051	0.298	0.377	0.000	3.949	0.096

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	57	40	50	67	0	101	33
N.S.	1	1.00	1.50	1.05	1.32	1.76	0.00	2.66	0.87
time (sec)	N/A	0.019	0.021	0.068	0.305	0.403	0.000	4.143	0.394

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	46	33	65	67	0	170	46
N.S.	1	1.00	1.18	0.85	1.67	1.72	0.00	4.36	1.18
time (sec)	N/A	0.020	0.068	0.066	0.285	0.353	0.000	3.222	0.409

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	72	50	60	77	0	145	45
N.S.	1	1.00	1.36	0.94	1.13	1.45	0.00	2.74	0.85
time (sec)	N/A	0.020	0.020	0.085	0.291	0.377	0.000	3.240	0.403

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	56	43	85	77	0	214	56
N.S.	1	1.00	0.98	0.75	1.49	1.35	0.00	3.75	0.98
time (sec)	N/A	0.022	0.102	0.097	0.291	0.393	0.000	3.576	0.400

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	50	62	42	43	82	42	43
N.S.	1	1.00	1.00	1.24	0.84	0.86	1.64	0.84	0.86
time (sec)	N/A	0.026	0.020	0.062	0.284	0.345	0.990	3.365	0.475

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	41	66	63	52	119	55	47
N.S.	1	1.00	0.67	1.08	1.03	0.85	1.95	0.90	0.77
time (sec)	N/A	0.033	0.103	0.042	0.535	0.405	0.745	2.923	0.589

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	52	32	33	61	32	35
N.S.	1	1.00	1.00	1.37	0.84	0.87	1.61	0.84	0.92
time (sec)	N/A	0.025	0.013	0.047	0.285	0.356	0.549	3.491	0.448

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	31	56	43	40	75	43	43
N.S.	1	1.00	0.78	1.40	1.08	1.00	1.88	1.08	1.08
time (sec)	N/A	0.025	0.100	0.031	0.505	0.363	0.441	2.953	0.568

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	23	42	20	22	39	20	23
N.S.	1	1.00	1.00	1.83	0.87	0.96	1.70	0.87	1.00
time (sec)	N/A	0.015	0.011	0.030	0.278	0.346	0.357	2.788	0.446

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	29	21	18	29	29	35	15
N.S.	1	1.00	1.93	1.40	1.20	1.93	1.93	2.33	1.00
time (sec)	N/A	0.006	0.012	0.032	0.506	0.357	0.311	3.000	0.411

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	11	11	11	14	13	13	20	13	13
N.S.	1	1.00	1.00	1.27	1.18	1.18	1.82	1.18	1.18
time (sec)	N/A	0.007	0.007	0.032	0.284	0.363	0.311	2.742	0.419

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	27	30	36	50	0	38	22
N.S.	1	1.00	1.17	1.30	1.57	2.17	0.00	1.65	0.96
time (sec)	N/A	0.015	0.012	0.038	0.305	0.400	0.000	7.083	0.021

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	13	31	22	33	0	16	14
N.S.	1	1.00	0.59	1.41	1.00	1.50	0.00	0.73	0.64
time (sec)	N/A	0.024	0.015	0.089	0.289	0.340	0.000	3.171	0.393

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	27	50	61	85	0	63	48
N.S.	1	1.00	0.55	1.02	1.24	1.73	0.00	1.29	0.98
time (sec)	N/A	0.031	0.011	0.054	0.281	0.362	0.000	4.049	0.429

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	46	50	32	43	0	32	33
N.S.	1	1.00	1.21	1.32	0.84	1.13	0.00	0.84	0.87
time (sec)	N/A	0.025	0.028	0.061	0.282	0.348	0.000	3.971	0.412

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	27	68	79	95	0	73	67
N.S.	1	1.00	0.39	0.97	1.13	1.36	0.00	1.04	0.96
time (sec)	N/A	0.032	0.014	0.073	0.277	0.365	0.000	4.270	0.094

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	45	63	45	71	1484	231	85
N.S.	1	1.00	0.78	1.09	0.78	1.22	25.59	3.98	1.47
time (sec)	N/A	0.032	0.075	0.050	0.281	0.381	5.625	4.378	0.694

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	103	70	66	93	719	163	129
N.S.	1	1.00	1.56	1.06	1.00	1.41	10.89	2.47	1.95
time (sec)	N/A	0.031	0.030	0.063	0.283	0.377	2.528	3.961	1.346

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	35	53	35	61	614	187	62
N.S.	1	1.00	0.81	1.23	0.81	1.42	14.28	4.35	1.44
time (sec)	N/A	0.027	0.045	0.041	0.292	0.361	1.503	3.595	0.462

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	86	60	56	83	241	140	77
N.S.	1	1.00	1.76	1.22	1.14	1.69	4.92	2.86	1.57
time (sec)	N/A	0.020	0.023	0.040	0.288	0.413	0.940	4.216	0.539

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	34	25	23	41	42	36	36
N.S.	1	1.00	1.21	0.89	0.82	1.46	1.50	1.29	1.29
time (sec)	N/A	0.012	0.071	0.034	0.311	0.358	0.352	4.439	0.429

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	34	34	75	50	46	72	58	93	48
N.S.	1	1.00	2.21	1.47	1.35	2.12	1.71	2.74	1.41
time (sec)	N/A	0.016	0.023	0.034	0.284	0.371	0.533	3.166	0.449

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	18	24	13	13
N.S.	1	1.00	1.00	0.93	0.87	1.20	1.60	0.87	0.87
time (sec)	N/A	0.013	0.010	0.017	0.284	0.332	0.346	4.892	0.392

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	34	23	36	65	0	119	34
N.S.	1	1.00	1.26	0.85	1.33	2.41	0.00	4.41	1.26
time (sec)	N/A	0.015	0.030	0.059	0.289	0.363	0.000	4.342	0.050

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	143	52	61	96	0	140	49
N.S.	1	1.00	2.92	1.06	1.24	1.96	0.00	2.86	1.00
time (sec)	N/A	0.030	0.183	0.066	0.289	0.377	0.000	3.676	0.034

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	61	43	64	102	0	188	39
N.S.	1	1.00	1.42	1.00	1.49	2.37	0.00	4.37	0.91
time (sec)	N/A	0.027	0.017	0.059	0.289	0.368	0.000	4.074	0.379

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	205	70	73	112	0	163	60
N.S.	1	1.00	3.11	1.06	1.11	1.70	0.00	2.47	0.91
time (sec)	N/A	0.029	0.303	0.076	0.294	0.346	0.000	3.541	0.458

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	56	61	82	112	0	232	74
N.S.	1	1.00	0.97	1.05	1.41	1.93	0.00	4.00	1.28
time (sec)	N/A	0.030	0.162	0.079	0.308	0.355	0.000	3.902	0.438

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	68	90	56	68	105	56	55
N.S.	1	1.00	1.00	1.32	0.82	1.00	1.54	0.82	0.81
time (sec)	N/A	0.032	0.026	0.112	0.288	0.359	1.857	3.434	0.525

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	53	94	75	89	141	68	56
N.S.	1	1.00	0.66	1.18	0.94	1.11	1.76	0.85	0.70
time (sec)	N/A	0.038	0.205	0.110	0.514	0.360	1.378	3.724	1.583

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	80	44	56	82	41	45
N.S.	1	1.00	1.00	1.51	0.83	1.06	1.55	0.77	0.85
time (sec)	N/A	0.029	0.021	0.046	0.294	0.355	1.028	4.904	0.479

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	57	57	43	84	55	79	97	55	46
N.S.	1	1.00	0.75	1.47	0.96	1.39	1.70	0.96	0.81
time (sec)	N/A	0.027	0.132	0.038	0.499	0.400	0.786	6.315	0.725

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	37	68	35	48	63	35	32
N.S.	1	1.00	1.00	1.84	0.95	1.30	1.70	0.95	0.86
time (sec)	N/A	0.018	0.015	0.040	0.320	0.348	0.611	5.413	0.450

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	33	26	34	69	48	62	24
N.S.	1	1.00	1.22	0.96	1.26	2.56	1.78	2.30	0.89
time (sec)	N/A	0.012	0.008	0.029	0.518	0.375	0.496	4.343	0.451

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	60	25	38	42	25	22
N.S.	1	1.00	1.00	2.31	0.96	1.46	1.62	0.96	0.85
time (sec)	N/A	0.014	0.012	0.032	0.295	0.345	0.495	4.476	0.434

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	13	34	71	13	13
N.S.	1	1.00	1.00	1.47	0.87	2.27	4.73	0.87	0.87
time (sec)	N/A	0.019	0.008	0.030	0.272	0.349	0.780	4.396	0.383

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	26	24	13	13
N.S.	1	1.00	1.00	0.93	0.87	1.73	1.60	0.87	0.87
time (sec)	N/A	0.013	0.008	0.022	0.280	0.334	0.417	3.996	0.426

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	31	40	50	94	0	52	32
N.S.	1	1.00	0.82	1.05	1.32	2.47	0.00	1.37	0.84
time (sec)	N/A	0.018	0.011	0.061	0.277	0.368	0.000	3.665	0.025

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	45	50	35	54	0	35	36
N.S.	1	1.00	1.22	1.35	0.95	1.46	0.00	0.95	0.97
time (sec)	N/A	0.025	0.029	0.049	0.279	0.341	0.000	4.182	0.418

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	31	68	73	130	0	72	61
N.S.	1	1.00	0.47	1.03	1.11	1.97	0.00	1.09	0.92
time (sec)	N/A	0.029	0.011	0.062	0.272	0.355	0.000	3.640	0.382

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	43	68	44	66	0	31	45
N.S.	1	1.00	0.81	1.28	0.83	1.25	0.00	0.58	0.85
time (sec)	N/A	0.027	0.017	0.062	0.297	0.362	0.000	3.292	0.461

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	31	86	91	140	0	85	79
N.S.	1	1.00	0.35	0.97	1.02	1.57	0.00	0.96	0.89
time (sec)	N/A	0.034	0.014	0.093	0.283	0.378	0.000	4.784	0.448

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	55	89	56	100	1664	277	92
N.S.	1	1.00	0.80	1.29	0.81	1.45	24.12	4.01	1.33
time (sec)	N/A	0.033	0.075	0.062	0.288	0.389	10.636	3.282	1.420

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	141	98	89	132	869	209	157
N.S.	1	1.00	1.58	1.10	1.00	1.48	9.76	2.35	1.76
time (sec)	N/A	0.035	0.033	0.072	0.280	0.407	5.733	3.348	0.549

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	47	81	49	90	733	232	74
N.S.	1	1.00	0.81	1.40	0.84	1.55	12.64	4.00	1.28
time (sec)	N/A	0.029	0.105	0.056	0.277	0.382	4.857	4.295	0.642

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	123	88	79	122	330	164	105
N.S.	1	1.00	1.76	1.26	1.13	1.74	4.71	2.34	1.50
time (sec)	N/A	0.024	0.026	0.049	0.289	0.368	1.991	5.310	0.581

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	46	33	38	70	61	164	52
N.S.	1	1.00	1.10	0.79	0.90	1.67	1.45	3.90	1.24
time (sec)	N/A	0.016	0.081	0.039	0.279	0.405	0.614	4.897	0.422

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	113	78	71	112	92	139	78
N.S.	1	1.00	2.05	1.42	1.29	2.04	1.67	2.53	1.42
time (sec)	N/A	0.031	0.026	0.050	0.287	0.354	1.145	4.508	0.488

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	22	25	39	44	25	25
N.S.	1	1.00	1.00	1.47	1.67	2.60	2.93	1.67	1.67
time (sec)	N/A	0.020	0.005	0.033	0.282	0.327	0.610	4.322	0.394

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	113	68	65	111	58	98	48
N.S.	1	1.00	2.05	1.24	1.18	2.02	1.05	1.78	0.87
time (sec)	N/A	0.031	0.029	0.046	0.284	0.369	1.062	5.566	0.466

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	14	13	27	24	13	23
N.S.	1	1.00	1.00	0.93	0.87	1.80	1.60	0.87	1.53
time (sec)	N/A	0.013	0.008	0.033	0.272	0.357	0.540	3.061	0.409

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	44	33	51	105	0	165	79
N.S.	1	1.00	1.10	0.82	1.28	2.62	0.00	4.12	1.98
time (sec)	N/A	0.019	0.075	0.050	0.274	0.374	0.000	3.491	0.442

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	129	70	79	132	0	163	66
N.S.	1	1.00	1.84	1.00	1.13	1.89	0.00	2.33	0.94
time (sec)	N/A	0.030	2.999	0.063	0.284	0.373	0.000	4.062	0.390

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	54	61	74	138	0	232	82
N.S.	1	1.00	0.93	1.05	1.28	2.38	0.00	4.00	1.41
time (sec)	N/A	0.029	0.244	0.081	0.274	0.356	0.000	3.943	0.469

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	268	88	91	148	0	209	78
N.S.	1	1.00	3.01	0.99	1.02	1.66	0.00	2.35	0.88
time (sec)	N/A	0.033	0.361	0.087	0.294	0.352	0.000	3.791	0.071

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	91	79	92	148	0	278	64
N.S.	1	1.00	1.32	1.14	1.33	2.14	0.00	4.03	0.93
time (sec)	N/A	0.033	0.023	0.052	0.291	0.373	0.000	3.911	0.415

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	27	22	14	33	29	14	19
N.S.	1	1.00	1.59	1.29	0.82	1.94	1.71	0.82	1.12
time (sec)	N/A	0.019	0.017	0.037	0.276	0.348	0.009	3.247	0.084

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	17	17	17	22	14	30	15	14	13
N.S.	1	1.00	1.00	1.29	0.82	1.76	0.88	0.82	0.76
time (sec)	N/A	0.018	0.008	0.046	0.280	0.354	0.029	3.207	0.439

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	19	18	24	37	0	18
N.S.	1	1.00	1.00	0.86	0.82	1.09	1.68	0.00	0.82
time (sec)	N/A	0.018	0.024	0.032	0.286	0.367	4.732	0.000	0.128

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	19	18	21	37	21	18
N.S.	1	1.00	1.00	0.86	0.82	0.95	1.68	0.95	0.82
time (sec)	N/A	0.015	0.012	0.020	0.281	0.372	0.708	5.939	0.423

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	18	18	36	18	18
N.S.	1	1.00	1.00	0.95	0.90	0.90	1.80	0.90	0.90
time (sec)	N/A	0.018	0.010	0.026	0.288	0.362	0.699	4.618	0.525

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	19	18	26	34	0	37
N.S.	1	1.00	1.00	0.95	0.90	1.30	1.70	0.00	1.85
time (sec)	N/A	0.018	0.017	0.025	0.288	0.362	1.224	0.000	0.183

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	19	18	26	36	0	53
N.S.	1	1.00	1.00	0.86	0.82	1.18	1.64	0.00	2.41
time (sec)	N/A	0.019	0.019	0.022	0.274	0.352	5.055	0.000	0.807

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	19	18	26	36	0	65
N.S.	1	1.00	1.00	0.86	0.82	1.18	1.64	0.00	2.95
time (sec)	N/A	0.019	0.028	0.037	0.274	0.343	46.042	0.000	6.709

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	19	18	26	0	0	65
N.S.	1	1.00	1.00	0.86	0.82	1.18	0.00	0.00	2.95
time (sec)	N/A	0.018	0.042	0.020	0.291	0.346	0.000	0.000	4.036

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	60	249	0	119	0	0	-1
N.S.	1	1.00	0.48	1.98	0.00	0.94	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.104	0.119	0.000	0.127	0.000	0.000	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	60	236	0	107	0	0	-1
N.S.	1	1.00	0.48	1.87	0.00	0.85	0.00	0.00	-0.01
time (sec)	N/A	0.076	0.096	0.114	0.000	0.106	0.000	0.000	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	57	223	0	106	0	0	-1
N.S.	1	1.00	0.58	2.28	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.037	0.121	0.000	0.105	0.000	0.000	0.000

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	57	208	0	90	0	0	-1
N.S.	1	1.00	0.58	2.12	0.00	0.92	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.049	0.099	0.000	0.116	0.000	0.000	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	58	194	0	87	0	0	-1
N.S.	1	1.00	0.84	2.81	0.00	1.26	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.064	0.104	0.000	0.122	0.000	0.000	0.000

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	58	188	0	78	0	0	-1
N.S.	1	1.00	0.84	2.72	0.00	1.13	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.072	0.147	0.000	0.126	0.000	0.000	0.000

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	60	168	0	105	0	0	-1
N.S.	1	1.00	0.88	2.47	0.00	1.54	0.00	0.00	-0.01
time (sec)	N/A	0.044	0.061	0.129	0.000	0.102	0.000	0.000	0.000

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	60	242	0	103	0	0	-1
N.S.	1	1.00	0.83	3.36	0.00	1.43	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.060	0.118	0.000	0.098	0.000	0.000	0.000

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	59	365	0	120	0	0	-1
N.S.	1	1.00	0.59	3.65	0.00	1.20	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.050	0.198	0.000	0.095	0.000	0.000	0.000

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	59	396	0	114	0	0	-1
N.S.	1	1.00	0.59	3.96	0.00	1.14	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.044	0.144	0.000	0.102	0.000	0.000	0.000

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	57	63	36	34	73	53	-1
N.S.	1	1.00	1.27	1.40	0.80	0.76	1.62	1.18	-0.02
time (sec)	N/A	0.028	0.222	0.077	0.289	0.366	1.200	5.656	0.000

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	57	92	36	28	71	46	-1
N.S.	1	1.00	1.33	2.14	0.84	0.65	1.65	1.07	-0.02
time (sec)	N/A	0.029	0.127	0.065	0.288	0.368	1.195	5.617	0.000

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	46	70	35	36	70	0	-1
N.S.	1	1.00	1.07	1.63	0.81	0.84	1.63	0.00	-0.02
time (sec)	N/A	0.034	0.052	0.211	0.295	0.365	2.427	0.000	0.000

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	48	85	34	38	71	0	66
N.S.	1	1.00	1.12	1.98	0.79	0.88	1.65	0.00	1.53
time (sec)	N/A	0.037	0.074	0.207	0.315	0.364	5.296	0.000	1.064

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	70	98	37	38	71	0	93
N.S.	1	1.00	1.63	2.28	0.86	0.88	1.65	0.00	2.16
time (sec)	N/A	0.034	0.185	0.214	0.287	0.387	40.731	0.000	3.666

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	70	111	37	38	0	0	93
N.S.	1	1.00	1.56	2.47	0.82	0.84	0.00	0.00	2.07
time (sec)	N/A	0.034	0.217	0.373	0.283	0.353	0.000	0.000	3.780

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	94	124	37	38	0	0	279
N.S.	1	1.00	2.09	2.76	0.82	0.84	0.00	0.00	6.20
time (sec)	N/A	0.033	0.372	0.465	0.290	0.353	0.000	0.000	5.330

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	156	156	57	275	0	133	0	0	-1
N.S.	1	1.00	0.37	1.76	0.00	0.85	0.00	0.00	-0.01
time (sec)	N/A	0.112	0.098	0.188	0.000	0.133	0.000	0.000	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	156	156	57	262	0	121	0	0	-1
N.S.	1	1.00	0.37	1.68	0.00	0.78	0.00	0.00	-0.01
time (sec)	N/A	0.107	0.068	0.206	0.000	0.149	0.000	0.000	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	65	249	0	120	0	0	-1
N.S.	1	1.00	0.51	1.95	0.00	0.94	0.00	0.00	-0.01
time (sec)	N/A	0.092	0.050	0.177	0.000	0.117	0.000	0.000	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	65	255	0	102	0	0	-1
N.S.	1	1.00	0.51	1.99	0.00	0.80	0.00	0.00	-0.01
time (sec)	N/A	0.091	0.079	0.209	0.000	0.132	0.000	0.000	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	58	221	0	100	0	0	-1
N.S.	1	1.00	0.59	2.23	0.00	1.01	0.00	0.00	-0.01
time (sec)	N/A	0.068	0.050	0.223	0.000	0.099	0.000	0.000	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	58	208	0	88	0	0	-1
N.S.	1	1.00	0.59	2.10	0.00	0.89	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.052	0.166	0.000	0.104	0.000	0.000	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	60	213	0	115	0	0	-1
N.S.	1	1.00	0.60	2.13	0.00	1.15	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.046	0.168	0.000	0.114	0.000	0.000	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	60	286	0	113	0	0	-1
N.S.	1	1.00	0.59	2.80	0.00	1.11	0.00	0.00	-0.01
time (sec)	N/A	0.075	0.050	0.206	0.000	0.101	0.000	0.000	0.000

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	65	366	0	120	0	0	-1
N.S.	1	1.00	0.64	3.59	0.00	1.18	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.043	0.319	0.000	0.105	0.000	0.000	0.000

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	65	398	0	114	0	0	-1
N.S.	1	1.00	0.64	3.90	0.00	1.12	0.00	0.00	-0.01
time (sec)	N/A	0.077	0.048	0.290	0.000	0.100	0.000	0.000	0.000

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	B	A	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	111	103	36	44	0	0	35
N.S.	1	1.00	2.13	1.98	0.69	0.85	0.00	0.00	0.67
time (sec)	N/A	0.024	0.192	0.182	0.300	0.363	0.000	0.000	0.636

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	83	332	98	313	0	0	-1
N.S.	1	1.00	0.83	3.32	0.98	3.13	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.161	0.418	0.505	0.480	0.000	0.000	0.000

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	80	293	98	299	0	0	-1
N.S.	1	1.00	0.81	2.96	0.99	3.02	0.00	0.00	-0.01
time (sec)	N/A	0.048	0.152	0.361	0.518	0.474	0.000	0.000	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	68	254	83	281	0	0	-1
N.S.	1	1.00	0.87	3.26	1.06	3.60	0.00	0.00	-0.01
time (sec)	N/A	0.044	0.102	0.365	0.494	0.491	0.000	0.000	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	65	211	83	259	0	0	-1
N.S.	1	1.00	0.84	2.74	1.08	3.36	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.065	0.443	0.505	0.467	0.000	0.000	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	51	183	67	238	0	48	-1
N.S.	1	1.00	0.88	3.16	1.16	4.10	0.00	0.83	-0.02
time (sec)	N/A	0.038	0.048	0.375	0.509	0.404	0.000	4.354	0.000

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	50	182	68	246	0	52	-1
N.S.	1	1.00	0.85	3.08	1.15	4.17	0.00	0.88	-0.02
time (sec)	N/A	0.035	0.032	0.433	0.514	0.404	0.000	4.147	0.000

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	66	422	79	309	0	0	-1
N.S.	1	1.00	0.85	5.41	1.01	3.96	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.059	0.684	0.793	0.417	0.000	0.000	0.000

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	70	624	80	318	0	0	-1
N.S.	1	1.00	0.86	7.70	0.99	3.93	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.094	0.654	0.600	0.416	0.000	0.000	0.000

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	81	882	100	342	0	0	-1
N.S.	1	1.00	0.81	8.82	1.00	3.42	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.134	0.683	0.562	0.430	0.000	0.000	0.000

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	87	1082	102	342	0	0	-1
N.S.	1	1.00	0.84	10.50	0.99	3.32	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.115	0.875	0.555	0.432	0.000	0.000	0.000

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	89	242	0	122	0	0	-1
N.S.	1	1.00	0.72	1.95	0.00	0.98	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.267	0.642	0.000	0.124	0.000	0.000	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	74	229	0	121	0	0	-1
N.S.	1	1.00	0.77	2.39	0.00	1.26	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.186	0.753	0.000	0.112	0.000	0.000	0.000

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	73	216	0	109	0	0	-1
N.S.	1	1.00	0.76	2.25	0.00	1.14	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.172	0.573	0.000	0.106	0.000	0.000	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	58	203	0	105	0	0	-1
N.S.	1	1.00	0.88	3.08	0.00	1.59	0.00	0.00	-0.02
time (sec)	N/A	0.046	0.108	0.744	0.000	0.121	0.000	0.000	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	56	190	0	91	0	0	-1
N.S.	1	1.00	0.85	2.88	0.00	1.38	0.00	0.00	-0.02
time (sec)	N/A	0.044	0.075	0.720	0.000	0.098	0.000	0.000	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	56	203	0	102	0	0	-1
N.S.	1	1.00	0.86	3.12	0.00	1.57	0.00	0.00	-0.02
time (sec)	N/A	0.041	0.063	0.819	0.000	0.105	0.000	0.000	0.000

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	47	188	0	93	0	0	-1
N.S.	1	1.00	0.73	2.94	0.00	1.45	0.00	0.00	-0.02
time (sec)	N/A	0.041	0.065	0.688	0.000	0.107	0.000	0.000	0.000

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	65	209	0	131	0	0	-1
N.S.	1	1.00	0.69	2.22	0.00	1.39	0.00	0.00	-0.01
time (sec)	N/A	0.061	0.128	0.862	0.000	0.119	0.000	0.000	0.000

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	62	190	0	129	0	0	-1
N.S.	1	1.00	0.63	1.94	0.00	1.32	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.110	0.858	0.000	0.123	0.000	0.000	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	82	408	0	145	0	0	-1
N.S.	1	1.00	0.65	3.24	0.00	1.15	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.131	0.984	0.000	0.126	0.000	0.000	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	137	407	133	419	0	0	-1
N.S.	1	1.00	1.01	3.01	0.99	3.10	0.00	0.00	-0.01
time (sec)	N/A	0.063	1.446	0.822	0.505	0.489	0.000	0.000	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	78	371	118	405	0	0	-1
N.S.	1	1.00	0.69	3.28	1.04	3.58	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.460	0.885	0.503	0.472	0.000	0.000	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	118	310	118	393	0	0	-1
N.S.	1	1.00	1.04	2.74	1.04	3.48	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.842	0.872	0.499	0.509	0.000	0.000	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	65	286	103	380	0	0	-1
N.S.	1	1.00	0.71	3.14	1.13	4.18	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.213	0.694	0.501	0.404	0.000	0.000	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	76	280	103	347	0	0	-1
N.S.	1	1.00	0.84	3.08	1.13	3.81	0.00	0.00	-0.01
time (sec)	N/A	0.049	0.139	0.621	0.523	0.445	0.000	0.000	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	62	275	102	340	0	95	-1
N.S.	1	1.00	0.67	2.96	1.10	3.66	0.00	1.02	-0.01
time (sec)	N/A	0.044	0.187	0.723	0.528	0.431	0.000	5.222	0.000

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	69	283	103	334	0	91	-1
N.S.	1	1.00	0.74	3.04	1.11	3.59	0.00	0.98	-0.01
time (sec)	N/A	0.046	0.175	0.642	0.530	0.433	0.000	5.103	0.000

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	91	705	117	406	0	0	-1
N.S.	1	1.00	0.79	6.13	1.02	3.53	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.186	1.039	0.518	0.461	0.000	0.000	0.000

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	B	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	92	905	117	418	0	0	-1
N.S.	1	1.00	0.80	7.87	1.02	3.63	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.267	1.187	0.501	0.429	0.000	0.000	0.000

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	A	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	137	102	1165	134	438	0	0	-1
N.S.	1	1.00	0.74	8.50	0.98	3.20	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.337	1.342	0.489	0.450	0.000	0.000	0.000

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	22	22	22	19	18	21	37	21	18
N.S.	1	1.00	1.00	0.86	0.82	0.95	1.68	0.95	0.82
time (sec)	N/A	0.016	0.015	0.027	0.274	0.351	2.195	3.973	0.097

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	18	14	13	14	170	13	25
N.S.	1	1.00	0.86	0.67	0.62	0.67	8.10	0.62	1.19
time (sec)	N/A	0.016	0.011	0.131	0.319	0.362	4.696	5.884	0.474

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	18	14	13	20	24	13	25
N.S.	1	1.00	0.86	0.67	0.62	0.95	1.14	0.62	1.19
time (sec)	N/A	0.016	0.011	0.135	0.282	0.379	6.431	6.221	0.470

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	21	21	18	14	13	20	24	13	25
N.S.	1	1.00	0.86	0.67	0.62	0.95	1.14	0.62	1.19
time (sec)	N/A	0.017	0.010	0.129	0.282	0.349	44.486	6.476	0.447

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	16	14	13	12	323	13	25
N.S.	1	1.00	0.84	0.74	0.68	0.63	17.00	0.68	1.32
time (sec)	N/A	0.016	0.008	0.148	0.279	0.377	5.307	5.746	0.438

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	70	532	0	0	0	0	-1
N.S.	1	1.00	0.53	4.03	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.109	0.074	1.595	0.000	0.000	0.000	0.000	0.000

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	70	518	0	0	0	0	-1
N.S.	1	1.00	0.74	5.45	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.064	0.151	0.000	0.000	0.000	0.000	0.000

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	67	505	0	0	0	0	-1
N.S.	1	1.00	1.26	9.53	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.045	0.147	0.000	0.000	0.000	0.000	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	70	493	0	0	0	0	-1
N.S.	1	1.00	0.75	5.30	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.078	1.231	0.000	0.000	0.000	0.000	0.000

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	134	70	528	0	0	0	0	-1
N.S.	1	1.00	0.52	3.94	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.118	0.107	0.160	0.000	0.000	0.000	0.000	0.000

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	320	320	70	514	0	2205	0	0	-1
N.S.	1	1.00	0.22	1.61	0.00	6.89	0.00	0.00	-0.00
time (sec)	N/A	0.184	0.082	0.102	0.000	28.536	0.000	0.000	0.000

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	280	280	67	271	0	2003	0	0	-1
N.S.	1	1.00	0.24	0.97	0.00	7.15	0.00	0.00	-0.00
time (sec)	N/A	0.129	0.042	0.105	0.000	29.973	0.000	0.000	0.000

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	37	38	0	42	0	0	50
N.S.	1	1.00	1.00	1.03	0.00	1.14	0.00	0.00	1.35
time (sec)	N/A	0.035	0.059	0.115	0.000	0.359	0.000	0.000	0.987

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	57	50	0	54	0	0	95
N.S.	1	1.00	0.76	0.67	0.00	0.72	0.00	0.00	1.27
time (sec)	N/A	0.072	0.170	0.133	0.000	0.392	0.000	0.000	2.063

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	112	67	60	0	64	0	0	216
N.S.	1	1.00	0.60	0.54	0.00	0.57	0.00	0.00	1.93
time (sec)	N/A	0.115	0.162	0.134	0.000	0.459	0.000	0.000	6.208

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	71	216	0	0	0	0	-1
N.S.	1	1.00	0.54	1.65	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.117	0.086	0.138	0.000	0.000	0.000	0.000	0.000

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	67	182	0	0	0	0	-1
N.S.	1	1.00	0.72	1.96	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.058	0.132	0.000	0.000	0.000	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	67	186	0	106	0	0	-1
N.S.	1	1.00	0.68	1.90	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.076	0.115	0.118	0.000	0.099	0.000	0.000	0.000

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	133	70	215	0	119	0	0	-1
N.S.	1	1.00	0.53	1.62	0.00	0.89	0.00	0.00	-0.01
time (sec)	N/A	0.114	0.109	0.127	0.000	0.114	0.000	0.000	0.000

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	320	320	67	654	0	1868	0	0	-1
N.S.	1	1.00	0.21	2.04	0.00	5.84	0.00	0.00	-0.00
time (sec)	N/A	0.183	0.051	0.099	0.000	17.013	0.000	0.000	0.000

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	313	313	67	642	0	1865	0	0	-1
N.S.	1	1.00	0.21	2.05	0.00	5.96	0.00	0.00	-0.00
time (sec)	N/A	0.177	0.109	0.118	0.000	16.749	0.000	0.000	0.000

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	40	38	0	50	0	0	64
N.S.	1	1.00	1.08	1.03	0.00	1.35	0.00	0.00	1.73
time (sec)	N/A	0.038	0.083	0.098	0.000	0.401	0.000	0.000	1.505

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	57	50	0	61	0	0	207
N.S.	1	1.00	0.54	0.47	0.00	0.58	0.00	0.00	1.95
time (sec)	N/A	0.113	0.211	0.099	0.000	0.448	0.000	0.000	6.162

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	141	141	67	60	0	73	0	0	193
N.S.	1	1.00	0.48	0.43	0.00	0.52	0.00	0.00	1.37
time (sec)	N/A	0.157	0.212	0.123	0.000	0.487	0.000	0.000	6.674

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	166	166	72	545	0	0	0	0	-1
N.S.	1	1.00	0.43	3.28	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.149	0.120	0.146	0.000	0.000	0.000	0.000	0.000

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	70	532	0	0	0	0	-1
N.S.	1	1.00	0.53	4.06	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.115	0.126	0.126	0.000	0.000	0.000	0.000	0.000

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	67	519	0	0	0	0	-1
N.S.	1	1.00	0.71	5.46	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.067	0.141	0.000	0.000	0.000	0.000	0.000

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	67	508	0	0	0	0	-1
N.S.	1	1.00	0.71	5.40	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.078	0.087	0.109	0.000	0.000	0.000	0.000	0.000

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	133	133	70	529	0	0	0	0	-1
N.S.	1	1.00	0.53	3.98	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.111	0.123	0.133	0.000	0.000	0.000	0.000	0.000

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	168	168	72	544	0	0	0	0	-1
N.S.	1	1.00	0.43	3.24	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.153	0.125	0.115	0.000	0.000	0.000	0.000	0.000

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	320	320	67	510	0	2074	0	0	-1
N.S.	1	1.00	0.21	1.59	0.00	6.48	0.00	0.00	-0.00
time (sec)	N/A	0.174	0.083	0.112	0.000	29.811	0.000	0.000	0.000

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	315	315	67	532	0	2268	0	0	-1
N.S.	1	1.00	0.21	1.69	0.00	7.20	0.00	0.00	-0.00
time (sec)	N/A	0.181	0.090	0.109	0.000	30.679	0.000	0.000	0.000

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	40	38	0	60	0	0	89
N.S.	1	1.00	1.08	1.03	0.00	1.62	0.00	0.00	2.41
time (sec)	N/A	0.040	0.096	0.098	0.000	0.416	0.000	0.000	1.778

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	57	50	0	74	0	0	176
N.S.	1	1.00	0.54	0.47	0.00	0.70	0.00	0.00	1.66
time (sec)	N/A	0.114	0.219	0.098	0.000	0.483	0.000	0.000	6.335

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	141	141	67	60	0	87	0	0	207
N.S.	1	1.00	0.48	0.43	0.00	0.62	0.00	0.00	1.47
time (sec)	N/A	0.152	0.325	0.274	0.000	0.620	0.000	0.000	6.430

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	226	226	57	692	0	1282	0	0	44
N.S.	1	1.00	0.25	3.06	0.00	5.67	0.00	0.00	0.19
time (sec)	N/A	0.104	0.041	0.835	0.000	14.974	0.000	0.000	2.048

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	33	0	16	0	0	53
N.S.	1	1.00	1.00	2.06	0.00	1.00	0.00	0.00	3.31
time (sec)	N/A	0.016	0.015	0.132	0.000	0.352	0.000	0.000	0.862

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	38	166	0	447	0	0	25
N.S.	1	1.00	0.31	1.36	0.00	3.66	0.00	0.00	0.20
time (sec)	N/A	0.055	0.010	0.106	0.000	0.669	0.000	0.000	0.721

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	143	38	2595	0	457	0	0	25
N.S.	1	1.00	0.27	18.15	0.00	3.20	0.00	0.00	0.17
time (sec)	N/A	0.074	0.010	0.119	0.000	0.657	0.000	0.000	0.809

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	70	216	0	0	0	0	-1
N.S.	1	1.00	0.53	1.64	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.114	0.079	0.195	0.000	0.000	0.000	0.000	0.000

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	68	189	0	0	0	0	-1
N.S.	1	1.00	0.74	2.05	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.077	0.138	0.000	0.000	0.000	0.000	0.000

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	65	151	0	60	0	0	-1
N.S.	1	1.00	1.23	2.85	0.00	1.13	0.00	0.00	-0.02
time (sec)	N/A	0.036	0.046	0.104	0.000	0.092	0.000	0.000	0.000

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	65	185	0	106	0	0	-1
N.S.	1	1.00	0.67	1.91	0.00	1.09	0.00	0.00	-0.01
time (sec)	N/A	0.075	0.075	0.102	0.000	0.100	0.000	0.000	0.000

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	134	70	213	0	120	0	0	-1
N.S.	1	1.00	0.52	1.59	0.00	0.90	0.00	0.00	-0.01
time (sec)	N/A	0.113	0.090	0.129	0.000	0.129	0.000	0.000	0.000

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	280	280	65	312	0	1697	0	0	-1
N.S.	1	1.00	0.23	1.11	0.00	6.06	0.00	0.00	-0.00
time (sec)	N/A	0.119	0.042	0.116	0.000	15.441	0.000	0.000	0.000

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	36	38	0	39	0	0	31
N.S.	1	1.00	1.03	1.09	0.00	1.11	0.00	0.00	0.89
time (sec)	N/A	0.036	0.041	0.086	0.000	0.443	0.000	0.000	0.815

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	52	50	0	51	0	0	77
N.S.	1	1.00	0.69	0.67	0.00	0.68	0.00	0.00	1.03
time (sec)	N/A	0.074	0.112	0.098	0.000	0.401	0.000	0.000	1.493

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	112	112	67	60	0	61	0	0	123
N.S.	1	1.00	0.60	0.54	0.00	0.54	0.00	0.00	1.10
time (sec)	N/A	0.111	0.163	0.115	0.000	0.406	0.000	0.000	3.826

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	174	55	292	0	1185	0	0	44
N.S.	1	1.00	0.32	1.68	0.00	6.81	0.00	0.00	0.25
time (sec)	N/A	0.061	0.019	0.112	0.000	14.637	0.000	0.000	1.599

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	199	199	55	937	0	1545	0	0	44
N.S.	1	1.00	0.28	4.71	0.00	7.76	0.00	0.00	0.22
time (sec)	N/A	0.078	0.026	0.293	0.000	27.631	0.000	0.000	1.625

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F(-2)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	201	57	1261	0	1321	0	0	44
N.S.	1	1.00	0.28	6.27	0.00	6.57	0.00	0.00	0.22
time (sec)	N/A	0.082	0.026	0.104	0.000	15.076	0.000	0.000	1.901

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	B	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	226	226	57	1934	0	1670	0	0	44
N.S.	1	1.00	0.25	8.56	0.00	7.39	0.00	0.00	0.19
time (sec)	N/A	0.099	0.031	0.117	0.000	27.371	0.000	0.000	1.875

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.043	0.081	0.000	0.000	0.000	0.000	0.000

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.027	0.034	0.055	0.000	0.000	0.000	0.000	0.000

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.011	0.026	0.002	0.000	0.000	0.000	0.000	0.000

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.030	0.040	0.000	0.000	0.000	0.000	0.000

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.029	0.046	0.000	0.000	0.000	0.000	0.000

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.040	0.080	0.000	0.000	0.000	0.000	0.000

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.039	0.058	0.000	0.000	0.000	0.000	0.000

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.011	0.032	0.003	0.000	0.000	0.000	0.000	0.000

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.038	0.045	0.000	0.000	0.000	0.000	0.000

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.036	0.055	0.000	0.000	0.000	0.000	0.000

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.038	0.075	0.000	0.000	0.000	0.000	0.000

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.034	0.047	0.000	0.000	0.000	0.000	0.000

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.011	0.027	0.002	0.000	0.000	0.000	0.000	0.000

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.032	0.039	0.000	0.000	0.000	0.000	0.000

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.030	0.041	0.000	0.000	0.000	0.000	0.000

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.035	0.064	0.000	0.000	0.000	0.000	0.000

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.033	0.037	0.000	0.000	0.000	0.000	0.000

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.012	0.026	0.001	0.000	0.000	0.000	0.000	0.000

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.032	0.041	0.000	0.000	0.000	0.000	0.000

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F(-1)	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	55	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	0.028	0.043	0.000	0.000	0.000	0.000	0.000

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	57	0	0	144	0	0	44
N.S.	1	1.00	0.45	0.00	0.00	1.12	0.00	0.00	0.34
time (sec)	N/A	0.102	0.030	0.020	0.000	0.419	0.000	0.000	1.237

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	224	224	57	0	0	0	0	0	44
N.S.	1	1.00	0.25	0.00	0.00	0.00	0.00	0.00	0.20
time (sec)	N/A	0.221	0.033	0.019	0.000	0.000	0.000	0.000	1.049

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	249	249	57	0	0	0	0	0	44
N.S.	1	1.00	0.23	0.00	0.00	0.00	0.00	0.00	0.18
time (sec)	N/A	0.241	0.040	0.018	0.000	0.000	0.000	0.000	1.640

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	57	0	0	197	0	0	44
N.S.	1	1.00	0.37	0.00	0.00	1.27	0.00	0.00	0.28
time (sec)	N/A	0.119	0.039	0.027	0.000	0.527	0.000	0.000	1.139

Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	57	0	0	195	0	0	44
N.S.	1	1.00	0.37	0.00	0.00	1.26	0.00	0.00	0.28
time (sec)	N/A	0.079	0.038	0.026	0.000	0.447	0.000	0.000	1.644

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	57	0	0	152	0	0	44
N.S.	1	1.00	0.45	0.00	0.00	1.19	0.00	0.00	0.34
time (sec)	N/A	0.054	0.020	0.021	0.000	0.527	0.000	0.000	1.485

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	225	225	55	0	0	0	0	0	44
N.S.	1	1.00	0.24	0.00	0.00	0.00	0.00	0.00	0.20
time (sec)	N/A	0.202	0.020	0.020	0.000	0.000	0.000	0.000	1.118

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F(-1)	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	250	55	0	0	0	0	0	44
N.S.	1	1.00	0.22	0.00	0.00	0.00	0.00	0.00	0.18
time (sec)	N/A	0.230	0.024	0.018	0.000	0.000	0.000	0.000	1.695

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	57	0	0	189	0	0	44
N.S.	1	1.00	0.37	0.00	0.00	1.22	0.00	0.00	0.28
time (sec)	N/A	0.080	0.023	0.020	0.000	0.733	0.000	0.000	1.200

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-1)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	155	155	57	0	0	219	0	0	44
N.S.	1	1.00	0.37	0.00	0.00	1.41	0.00	0.00	0.28
time (sec)	N/A	0.079	0.025	0.018	0.000	0.607	0.000	0.000	1.914

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	0	0	18	0	0	10
N.S.	1	1.00	1.00	0.00	0.00	1.12	0.00	0.00	0.62
time (sec)	N/A	0.015	0.009	0.014	0.000	0.611	0.000	0.000	0.826

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	0	0	10	0	0	94
N.S.	1	1.00	1.00	0.00	0.00	0.62	0.00	0.00	5.88
time (sec)	N/A	0.015	0.014	0.018	0.000	0.393	0.000	0.000	0.822

Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	79	0	0	0	0	0	71
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	0.89
time (sec)	N/A	0.026	0.075	0.041	0.000	0.000	0.000	0.000	2.346

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	82	0	0	0	0	0	-1
N.S.	1	1.00	0.96	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.085	0.031	0.000	0.000	0.000	0.000	0.000

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	85	0	0	0	0	0	-1
N.S.	1	1.00	1.02	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.053	0.034	0.000	0.000	0.000	0.000	0.000

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	85	0	0	0	0	0	-1
N.S.	1	1.00	0.97	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.070	0.032	0.000	0.000	0.000	0.000	0.000

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	55	0	77	70	2040	248	132
N.S.	1	1.00	0.74	0.00	1.04	0.95	27.57	3.35	1.78
time (sec)	N/A	0.047	0.231	0.091	0.287	0.413	5.686	5.652	1.612

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	48	1318	53	46	525	118	62
N.S.	1	1.00	0.96	26.36	1.06	0.92	10.50	2.36	1.24
time (sec)	N/A	0.035	0.065	1.273	0.353	0.416	3.183	3.177	0.879

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	24	24	25	25	24	24	56	24	25
N.S.	1	1.00	1.04	1.04	1.00	1.00	2.33	1.00	1.04
time (sec)	N/A	0.015	0.007	1.111	0.318	0.407	0.456	3.240	0.605

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	-1
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.016	0.053	0.000	0.000	0.000	0.000	0.000

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	51	0	0	0	0	0	-1
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.033	0.019	0.037	0.000	0.000	0.000	0.000	0.000

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	63	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.041	0.086	0.000	0.000	0.000	0.000	0.000

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	63	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.036	0.065	0.000	0.000	0.000	0.000	0.000

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	63	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.011	0.033	0.008	0.000	0.000	0.000	0.000	0.000

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	63	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.034	0.042	0.000	0.000	0.000	0.000	0.000

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	63	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.032	0.034	0.000	0.000	0.000	0.000	0.000

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	78	0	0	0	0	0	-1
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.100	0.040	0.000	0.000	0.000	0.000	0.000

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	75	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.032	0.050	0.039	0.000	0.000	0.000	0.000	0.000

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	75	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.055	0.034	0.000	0.000	0.000	0.000	0.000

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	78	0	0	0	0	0	-1
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.067	0.035	0.000	0.000	0.000	0.000	0.000

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	78	0	0	0	0	0	-1
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.061	0.045	0.000	0.000	0.000	0.000	0.000

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	83	0	78	84	2451	249	132
N.S.	1	1.00	1.09	0.00	1.03	1.11	32.25	3.28	1.74
time (sec)	N/A	0.047	0.216	0.140	0.287	0.430	5.873	3.042	1.544

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	50	1076	52	50	688	117	65
N.S.	1	1.00	1.00	21.52	1.04	1.00	13.76	2.34	1.30
time (sec)	N/A	0.040	0.092	1.172	0.288	0.379	2.672	2.688	0.912

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	26	26	25	25	60	25	26
N.S.	1	1.00	1.04	1.04	1.00	1.00	2.40	1.00	1.04
time (sec)	N/A	0.016	0.007	0.931	0.299	0.444	0.827	3.767	0.175

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	52	0	0	0	0	0	-1
N.S.	1	1.00	1.06	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.037	0.093	0.066	0.000	0.000	0.000	0.000	0.000

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	154	0	0	0	0	0	-1
N.S.	1	1.00	3.14	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.033	1.852	0.074	0.000	0.000	0.000	0.000	0.000

Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	244	0	0	0	0	0	-1
N.S.	1	1.00	4.98	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.032	2.850	0.069	0.000	0.000	0.000	0.000	0.000

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	68	0	0	0	0	0	-1
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.093	0.293	0.000	0.000	0.000	0.000	0.000

Problem 362	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	68	0	0	0	0	0	-1
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.066	0.374	0.000	0.000	0.000	0.000	0.000

Problem 363	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	64	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.012	0.035	0.014	0.000	0.000	0.000	0.000	0.000

Problem 364	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	80	0	0	0	0	0	-1
N.S.	1	1.00	1.16	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.160	0.060	0.000	0.000	0.000	0.000	0.000

Problem 365	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	82	0	0	0	0	0	-1
N.S.	1	1.00	1.19	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.160	0.072	0.000	0.000	0.000	0.000	0.000

Problem 366	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	158	0	0	0	0	0	-1
N.S.	1	1.00	2.08	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.296	0.091	0.000	0.000	0.000	0.000	0.000

Problem 367	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	76	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.110	0.075	0.000	0.000	0.000	0.000	0.000

Problem 368	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	82	0	0	0	0	0	-1
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.032	0.080	0.079	0.000	0.000	0.000	0.000	0.000

Problem 369	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	82	0	0	0	0	0	-1
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.091	0.059	0.000	0.000	0.000	0.000	0.000

Problem 370	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	79	0	0	0	0	0	-1
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.103	0.053	0.000	0.000	0.000	0.000	0.000

Problem 371	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	58	517	67	61	0	108	-1
N.S.	1	1.00	0.68	6.08	0.79	0.72	0.00	1.27	-0.01
time (sec)	N/A	0.040	0.239	2.740	0.280	0.397	0.000	2.663	0.000

Problem 372	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	48	507	53	50	0	83	-1
N.S.	1	1.00	0.76	8.05	0.84	0.79	0.00	1.32	-0.02
time (sec)	N/A	0.036	0.151	0.262	0.274	0.390	0.000	3.395	0.000

Problem 373	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	36	497	37	37	0	57	-1
N.S.	1	1.00	0.88	12.12	0.90	0.90	0.00	1.39	-0.02
time (sec)	N/A	0.030	0.123	0.294	0.294	0.364	0.000	4.769	0.000

Problem 374	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	25	25	0	24	23
N.S.	1	1.00	1.00	0.94	1.39	1.39	0.00	1.33	1.28
time (sec)	N/A	0.020	0.031	0.112	0.287	0.399	0.000	5.881	0.204

Problem 375	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	58	58	73	169	75	265	0	64	-1
N.S.	1	1.00	1.26	2.91	1.29	4.57	0.00	1.10	-0.02
time (sec)	N/A	0.032	0.263	0.200	0.492	0.436	0.000	4.290	0.000

Problem 376	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	95	603	111	382	0	103	-1
N.S.	1	1.00	1.02	6.48	1.19	4.11	0.00	1.11	-0.01
time (sec)	N/A	0.054	0.450	0.269	0.484	0.481	0.000	4.651	0.000

Problem 377	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	107	1089	145	474	0	134	-1
N.S.	1	1.00	0.87	8.85	1.18	3.85	0.00	1.09	-0.01
time (sec)	N/A	0.057	0.630	0.267	0.539	0.505	0.000	4.767	0.000

Problem 378	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	73	165	0	115	0	0	-1
N.S.	1	1.00	0.59	1.34	0.00	0.93	0.00	0.00	-0.01
time (sec)	N/A	0.099	0.109	0.445	0.000	0.121	0.000	0.000	0.000

Problem 379	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	61	143	0	102	0	0	-1
N.S.	1	1.00	0.64	1.51	0.00	1.07	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.078	0.283	0.000	0.134	0.000	0.000	0.000

Problem 380	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	51	123	0	90	0	0	-1
N.S.	1	1.00	0.76	1.84	0.00	1.34	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.076	0.238	0.000	0.119	0.000	0.000	0.000

Problem 381	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	98	0	61	0	0	35
N.S.	1	1.00	1.00	2.58	0.00	1.61	0.00	0.00	0.92
time (sec)	N/A	0.015	0.031	0.280	0.000	0.088	0.000	0.000	0.563

Problem 382	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	47	184	0	107	0	0	-1
N.S.	1	1.00	0.76	2.97	0.00	1.73	0.00	0.00	-0.02
time (sec)	N/A	0.041	0.086	0.240	0.000	0.109	0.000	0.000	0.000

Problem 383	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	63	335	0	161	0	0	-1
N.S.	1	1.00	0.66	3.53	0.00	1.69	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.169	0.408	0.000	0.139	0.000	0.000	0.000

Problem 384	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	73	485	0	204	0	0	-1
N.S.	1	1.00	0.59	3.94	0.00	1.66	0.00	0.00	-0.01
time (sec)	N/A	0.099	0.318	0.290	0.000	0.120	0.000	0.000	0.000

Problem 385	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	52	969	76	58	0	106	-1
N.S.	1	1.00	0.63	11.67	0.92	0.70	0.00	1.28	-0.01
time (sec)	N/A	0.043	0.103	1.633	0.309	0.382	0.000	4.002	0.000

Problem 386	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	42	959	58	46	0	81	-1
N.S.	1	1.00	0.67	15.22	0.92	0.73	0.00	1.29	-0.02
time (sec)	N/A	0.039	0.069	0.253	0.275	0.384	0.000	4.008	0.000

Problem 387	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	30	949	39	33	0	50	-1
N.S.	1	1.00	0.73	23.15	0.95	0.80	0.00	1.22	-0.02
time (sec)	N/A	0.034	0.049	0.235	0.287	0.402	0.000	5.936	0.000

Problem 388	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	18	17	25	19	0	27	18
N.S.	1	1.00	1.00	0.94	1.39	1.06	0.00	1.50	1.00
time (sec)	N/A	0.024	0.023	0.046	0.285	0.380	0.000	4.590	0.478

Problem 389	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	85	235	91	298	0	79	-1
N.S.	1	1.00	1.10	3.05	1.18	3.87	0.00	1.03	-0.01
time (sec)	N/A	0.037	0.665	0.251	0.505	0.418	0.000	5.186	0.000

Problem 390	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	97	644	129	416	0	134	-1
N.S.	1	1.00	0.86	5.70	1.14	3.68	0.00	1.19	-0.01
time (sec)	N/A	0.055	1.628	0.227	0.690	0.479	0.000	3.773	0.000

Problem 391	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	70	330	0	116	0	0	-1
N.S.	1	1.00	0.55	2.58	0.00	0.91	0.00	0.00	-0.01
time (sec)	N/A	0.108	0.109	0.303	0.000	0.121	0.000	0.000	0.000

Problem 392	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	60	320	0	105	0	0	-1
N.S.	1	1.00	0.61	3.27	0.00	1.07	0.00	0.00	-0.01
time (sec)	N/A	0.077	0.089	0.252	0.000	0.107	0.000	0.000	0.000

Problem 393	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	48	310	0	91	0	0	-1
N.S.	1	1.00	0.73	4.70	0.00	1.38	0.00	0.00	-0.02
time (sec)	N/A	0.046	0.062	0.217	0.000	0.104	0.000	0.000	0.000

Problem 394	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	66	66	48	320	0	90	0	0	-1
N.S.	1	1.00	0.73	4.85	0.00	1.36	0.00	0.00	-0.02
time (sec)	N/A	0.026	0.049	0.262	0.000	0.104	0.000	0.000	0.000

Problem 395	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	57	322	0	122	0	0	-1
N.S.	1	1.00	0.63	3.58	0.00	1.36	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.112	0.227	0.000	0.104	0.000	0.000	0.000

Problem 396	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	77	622	0	180	0	0	-1
N.S.	1	1.00	0.62	5.02	0.00	1.45	0.00	0.00	-0.01
time (sec)	N/A	0.090	0.149	0.237	0.000	0.118	0.000	0.000	0.000

Problem 397	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	52	532	70	75	0	108	-1
N.S.	1	1.00	0.61	6.26	0.82	0.88	0.00	1.27	-0.01
time (sec)	N/A	0.043	0.313	0.234	0.294	0.402	0.000	5.238	0.000

Problem 398	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	42	522	55	61	0	80	-1
N.S.	1	1.00	0.67	8.29	0.87	0.97	0.00	1.27	-0.02
time (sec)	N/A	0.039	0.254	0.258	0.302	0.417	0.000	3.548	0.000

Problem 399	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F(-1)	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	32	357	38	45	0	53	50
N.S.	1	1.00	0.78	8.71	0.93	1.10	0.00	1.29	1.22
time (sec)	N/A	0.034	0.150	0.220	0.285	0.382	0.000	4.445	0.908

Problem 400	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	25	30	0	36	39
N.S.	1	1.00	1.00	0.85	1.25	1.50	0.00	1.80	1.95
time (sec)	N/A	0.024	0.036	0.044	0.284	0.413	0.000	4.578	0.624

Problem 401	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	87	237	91	354	0	91	-1
N.S.	1	1.00	1.12	3.04	1.17	4.54	0.00	1.17	-0.01
time (sec)	N/A	0.038	0.139	0.218	0.512	0.464	0.000	4.311	0.000

Problem 402	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	109	699	129	482	0	128	-1
N.S.	1	1.00	0.96	6.19	1.14	4.27	0.00	1.13	-0.01
time (sec)	N/A	0.057	1.360	0.325	0.491	0.509	0.000	4.333	0.000

Problem 403	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	143	119	1161	163	584	0	159	-1
N.S.	1	1.00	0.83	8.12	1.14	4.08	0.00	1.11	-0.01
time (sec)	N/A	0.067	0.930	0.204	0.502	0.515	0.000	3.529	0.000

Problem 404	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	74	168	0	140	0	0	-1
N.S.	1	1.00	0.57	1.29	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.101	0.165	0.312	0.000	0.180	0.000	0.000	0.000

Problem 405	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	64	144	0	124	0	0	-1
N.S.	1	1.00	0.64	1.44	0.00	1.24	0.00	0.00	-0.01
time (sec)	N/A	0.073	0.105	0.255	0.000	0.117	0.000	0.000	0.000

Problem 406	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	52	126	0	110	0	0	-1
N.S.	1	1.00	0.74	1.80	0.00	1.57	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.100	0.224	0.000	0.109	0.000	0.000	0.000

Problem 407	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	51	128	0	110	0	0	-1
N.S.	1	1.00	0.73	1.83	0.00	1.57	0.00	0.00	-0.01
time (sec)	N/A	0.025	0.066	0.203	0.000	0.122	0.000	0.000	0.000

Problem 408	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	67	202	0	143	0	0	-1
N.S.	1	1.00	0.68	2.06	0.00	1.46	0.00	0.00	-0.01
time (sec)	N/A	0.071	0.142	0.342	0.000	0.110	0.000	0.000	0.000

Problem 409	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	79	352	0	209	0	0	-1
N.S.	1	1.00	0.64	2.86	0.00	1.70	0.00	0.00	-0.01
time (sec)	N/A	0.106	0.303	0.284	0.000	0.113	0.000	0.000	0.000

Problem 410	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	52	56	67	66	0	117	-1
N.S.	1	1.00	0.60	0.64	0.77	0.76	0.00	1.34	-0.01
time (sec)	N/A	0.038	0.161	0.288	0.284	0.395	0.000	4.156	0.000

Problem 411	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	42	46	53	55	0	92	-1
N.S.	1	1.00	0.65	0.71	0.82	0.85	0.00	1.42	-0.02
time (sec)	N/A	0.033	0.119	0.211	0.269	0.387	0.000	3.659	0.000

Problem 412	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	32	36	39	44	0	67	-1
N.S.	1	1.00	0.74	0.84	0.91	1.02	0.00	1.56	-0.02
time (sec)	N/A	0.030	0.081	0.187	0.306	0.384	0.000	4.049	0.000

Problem 413	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	25	30	0	36	28
N.S.	1	1.00	1.00	0.85	1.25	1.50	0.00	1.80	1.40
time (sec)	N/A	0.021	0.039	0.051	0.339	0.385	0.000	3.971	0.544

Problem 414	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	73	161	76	271	0	59	-1
N.S.	1	1.00	1.24	2.73	1.29	4.59	0.00	1.00	-0.02
time (sec)	N/A	0.028	0.081	0.211	0.500	0.438	0.000	4.131	0.000

Problem 415	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	93	425	110	389	0	110	-1
N.S.	1	1.00	1.00	4.57	1.18	4.18	0.00	1.18	-0.01
time (sec)	N/A	0.048	0.855	0.248	0.519	0.455	0.000	4.313	0.000

Problem 416	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	107	729	145	486	0	143	-1
N.S.	1	1.00	0.87	5.93	1.18	3.95	0.00	1.16	-0.01
time (sec)	N/A	0.056	1.620	0.248	0.554	0.462	0.000	4.456	0.000

Problem 417	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	73	338	0	126	0	0	-1
N.S.	1	1.00	0.59	2.75	0.00	1.02	0.00	0.00	-0.01
time (sec)	N/A	0.103	0.328	0.514	0.000	0.123	0.000	0.000	0.000

Problem 418	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	63	328	0	115	0	0	-1
N.S.	1	1.00	0.66	3.45	0.00	1.21	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.243	0.230	0.000	0.124	0.000	0.000	0.000

Problem 419	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	60	318	0	101	0	0	-1
N.S.	1	1.00	0.90	4.75	0.00	1.51	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.105	0.274	0.000	0.119	0.000	0.000	0.000

Problem 420	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	306	0	70	0	0	-1
N.S.	1	1.00	1.00	8.05	0.00	1.84	0.00	0.00	-0.03
time (sec)	N/A	0.016	0.030	0.438	0.000	0.108	0.000	0.000	0.000

Problem 421	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	48	316	0	118	0	0	-1
N.S.	1	1.00	0.76	5.02	0.00	1.87	0.00	0.00	-0.02
time (sec)	N/A	0.044	0.117	0.231	0.000	0.113	0.000	0.000	0.000

Problem 422	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	74	618	0	171	0	0	-1
N.S.	1	1.00	0.78	6.51	0.00	1.80	0.00	0.00	-0.01
time (sec)	N/A	0.072	0.147	0.268	0.000	0.130	0.000	0.000	0.000

Problem 423	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	86	918	0	216	0	0	-1
N.S.	1	1.00	0.70	7.46	0.00	1.76	0.00	0.00	-0.01
time (sec)	N/A	0.102	0.106	0.281	0.000	0.116	0.000	0.000	0.000

Problem 424	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	52	56	67	66	0	119	-1
N.S.	1	1.00	0.60	0.64	0.77	0.76	0.00	1.37	-0.01
time (sec)	N/A	0.044	0.302	0.257	0.291	0.428	0.000	4.747	0.000

Problem 425	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	42	46	53	55	0	94	-1
N.S.	1	1.00	0.65	0.71	0.82	0.85	0.00	1.45	-0.02
time (sec)	N/A	0.041	0.180	0.201	0.285	0.409	0.000	3.890	0.000

Problem 426	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	32	36	39	44	0	69	-1
N.S.	1	1.00	0.74	0.84	0.91	1.02	0.00	1.60	-0.02
time (sec)	N/A	0.033	0.126	0.168	0.289	0.410	0.000	2.811	0.000

Problem 427	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	25	30	0	38	28
N.S.	1	1.00	1.00	0.85	1.25	1.50	0.00	1.90	1.40
time (sec)	N/A	0.024	0.042	0.046	0.285	0.393	0.000	3.700	0.567

Problem 428	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	89	221	93	338	0	77	-1
N.S.	1	1.00	1.14	2.83	1.19	4.33	0.00	0.99	-0.01
time (sec)	N/A	0.038	1.376	0.211	0.506	0.500	0.000	3.492	0.000

Problem 429	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	98	426	111	392	0	98	-1
N.S.	1	1.00	1.05	4.58	1.19	4.22	0.00	1.05	-0.01
time (sec)	N/A	0.054	0.337	0.216	0.489	0.535	0.000	3.502	0.000

Problem 430	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	109	729	144	490	0	135	-1
N.S.	1	1.00	0.89	5.93	1.17	3.98	0.00	1.10	-0.01
time (sec)	N/A	0.061	0.466	0.240	0.496	0.475	0.000	3.620	0.000

Problem 431	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	81	173	0	118	0	0	-1
N.S.	1	1.00	0.64	1.37	0.00	0.94	0.00	0.00	-0.01
time (sec)	N/A	0.094	0.121	0.282	0.000	0.119	0.000	0.000	0.000

Problem 432	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	71	153	0	107	0	0	-1
N.S.	1	1.00	0.72	1.56	0.00	1.09	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.100	0.229	0.000	0.151	0.000	0.000	0.000

Problem 433	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	59	131	0	94	0	0	-1
N.S.	1	1.00	0.82	1.82	0.00	1.31	0.00	0.00	-0.01
time (sec)	N/A	0.027	0.049	0.193	0.000	0.106	0.000	0.000	0.000

Problem 434	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	58	191	0	110	0	0	-1
N.S.	1	1.00	0.85	2.81	0.00	1.62	0.00	0.00	-0.01
time (sec)	N/A	0.049	0.077	0.199	0.000	0.138	0.000	0.000	0.000

Problem 435	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	62	343	0	161	0	0	-1
N.S.	1	1.00	0.61	3.36	0.00	1.58	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.167	0.222	0.000	0.122	0.000	0.000	0.000

Problem 436	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	74	493	0	214	0	0	-1
N.S.	1	1.00	0.56	3.73	0.00	1.62	0.00	0.00	-0.01
time (sec)	N/A	0.104	0.238	0.259	0.000	0.109	0.000	0.000	0.000

Problem 437	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-2)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	62	56	67	66	0	119	-1
N.S.	1	1.00	0.71	0.64	0.77	0.76	0.00	1.37	-0.01
time (sec)	N/A	0.045	0.263	0.270	0.287	0.412	0.000	5.316	0.000

Problem 438	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	52	46	53	55	0	94	-1
N.S.	1	1.00	0.80	0.71	0.82	0.85	0.00	1.45	-0.02
time (sec)	N/A	0.039	0.155	0.205	0.296	0.372	0.000	4.335	0.000

Problem 439	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F(-1)	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	36	39	44	0	69	-1
N.S.	1	1.00	0.98	0.84	0.91	1.02	0.00	1.60	-0.02
time (sec)	N/A	0.035	0.113	0.180	0.285	0.397	0.000	4.728	0.000

Problem 440	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	20	17	25	30	0	38	28
N.S.	1	1.00	1.00	0.85	1.25	1.50	0.00	1.90	1.40
time (sec)	N/A	0.026	0.061	0.046	0.277	0.414	0.000	4.259	0.569

Problem 441	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	90	377	94	343	0	86	-1
N.S.	1	1.00	1.11	4.65	1.16	4.23	0.00	1.06	-0.01
time (sec)	N/A	0.039	0.156	0.214	0.505	0.490	0.000	4.641	0.000

Problem 442	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	98	437	111	398	0	107	-1
N.S.	1	1.00	1.05	4.70	1.19	4.28	0.00	1.15	-0.01
time (sec)	N/A	0.053	1.612	0.220	0.485	0.470	0.000	4.380	0.000

Problem 443	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	110	737	143	494	0	139	-1
N.S.	1	1.00	0.89	5.99	1.16	4.02	0.00	1.13	-0.01
time (sec)	N/A	0.061	1.893	0.237	0.537	0.436	0.000	3.775	0.000

Problem 444	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	126	83	343	0	126	0	0	-1
N.S.	1	1.00	0.66	2.72	0.00	1.00	0.00	0.00	-0.01
time (sec)	N/A	0.095	0.363	0.237	0.000	0.122	0.000	0.000	0.000

Problem 445	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	66	333	0	115	0	0	-1
N.S.	1	1.00	0.67	3.40	0.00	1.17	0.00	0.00	-0.01
time (sec)	N/A	0.061	0.289	0.207	0.000	0.125	0.000	0.000	0.000

Problem 446	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	60	321	0	102	0	0	-1
N.S.	1	1.00	0.83	4.46	0.00	1.42	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.068	0.250	0.000	0.147	0.000	0.000	0.000

Problem 447	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	51	313	0	118	0	0	-1
N.S.	1	1.00	0.75	4.60	0.00	1.74	0.00	0.00	-0.01
time (sec)	N/A	0.047	0.132	0.230	0.000	0.117	0.000	0.000	0.000

Problem 448	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	79	623	0	175	0	0	-1
N.S.	1	1.00	0.77	6.11	0.00	1.72	0.00	0.00	-0.01
time (sec)	N/A	0.078	0.187	0.251	0.000	0.105	0.000	0.000	0.000

Problem 449	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	87	923	0	220	0	0	-1
N.S.	1	1.00	0.66	6.99	0.00	1.67	0.00	0.00	-0.01
time (sec)	N/A	0.111	0.137	0.277	0.000	0.111	0.000	0.000	0.000

Problem 450	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	449	449	169	538	0	0	0	0	-1
N.S.	1	1.00	0.38	1.20	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.310	1.386	1.395	0.000	0.000	0.000	0.000	0.000

Problem 451	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	414	414	157	512	0	0	0	0	-1
N.S.	1	1.00	0.38	1.24	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.237	0.759	0.274	0.000	0.000	0.000	0.000	0.000

Problem 452	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	376	376	122	273	0	0	0	0	-1
N.S.	1	1.00	0.32	0.73	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.173	0.444	0.283	0.000	0.000	0.000	0.000	0.000

Problem 453	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	33	33	37	40	0	48	0	0	36
N.S.	1	1.00	1.12	1.21	0.00	1.45	0.00	0.00	1.09
time (sec)	N/A	0.033	0.061	0.234	0.000	0.434	0.000	0.000	0.873

Problem 454	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	52	52	0	79	0	0	83
N.S.	1	1.00	0.73	0.73	0.00	1.11	0.00	0.00	1.17
time (sec)	N/A	0.070	0.142	0.239	0.000	0.506	0.000	0.000	1.805

Problem 455	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	106	106	65	62	0	104	0	0	169
N.S.	1	1.00	0.61	0.58	0.00	0.98	0.00	0.00	1.59
time (sec)	N/A	0.106	0.154	0.258	0.000	0.500	0.000	0.000	5.695

Problem 456	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	90	212	0	0	0	0	-1
N.S.	1	1.00	0.70	1.66	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.136	15.701	0.363	0.000	0.000	0.000	0.000	0.000

Problem 457	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	66	184	0	0	0	0	-1
N.S.	1	1.00	0.73	2.02	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.104	10.923	0.287	0.000	0.000	0.000	0.000	0.000

Problem 458	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	66	153	0	61	0	0	-1
N.S.	1	1.00	1.25	2.89	0.00	1.15	0.00	0.00	-0.02
time (sec)	N/A	0.066	0.278	0.251	0.000	0.157	0.000	0.000	0.000

Problem 459	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	75	278	0	133	0	0	-1
N.S.	1	1.00	0.79	2.93	0.00	1.40	0.00	0.00	-0.01
time (sec)	N/A	0.107	10.333	0.239	0.000	0.122	0.000	0.000	0.000

Problem 460	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	111	532	0	185	0	0	-1
N.S.	1	1.00	0.85	4.09	0.00	1.42	0.00	0.00	-0.01
time (sec)	N/A	0.142	10.760	0.264	0.000	0.124	0.000	0.000	0.000

Problem 461	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	86	524	0	0	0	0	-1
N.S.	1	1.00	0.75	4.56	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.109	10.391	0.373	0.000	0.000	0.000	0.000	0.000

Problem 462	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	85	85	74	511	0	0	0	0	-1
N.S.	1	1.00	0.87	6.01	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.081	10.223	0.286	0.000	0.000	0.000	0.000	0.000

Problem 463	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	60	497	0	0	0	0	-1
N.S.	1	1.00	1.18	9.75	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.056	10.807	0.289	0.000	0.000	0.000	0.000	0.000

Problem 464	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	63	482	0	0	0	0	-1
N.S.	1	1.00	0.78	5.95	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.084	10.192	0.207	0.000	0.000	0.000	0.000	0.000

Problem 465	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	F(-2)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	82	1030	0	0	0	0	-1
N.S.	1	1.00	0.71	8.96	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.109	10.353	0.229	0.000	0.000	0.000	0.000	0.000

Problem 466	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	363	363	145	648	0	0	0	0	-1
N.S.	1	1.00	0.40	1.79	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.185	0.993	0.198	0.000	0.000	0.000	0.000	0.000

Problem 467	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	328	328	113	304	0	0	0	0	-1
N.S.	1	1.00	0.34	0.93	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.136	0.519	0.233	0.000	0.000	0.000	0.000	0.000

Problem 468	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	70	0	52	0	0	57
N.S.	1	1.00	1.00	2.33	0.00	1.73	0.00	0.00	1.90
time (sec)	N/A	0.025	0.073	0.178	0.000	0.411	0.000	0.000	1.250

Problem 469	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	42	82	0	78	0	0	103
N.S.	1	1.00	0.69	1.34	0.00	1.28	0.00	0.00	1.69
time (sec)	N/A	0.053	0.098	0.194	0.000	0.417	0.000	0.000	2.662

Problem 470	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	52	92	0	103	0	0	163
N.S.	1	1.00	0.57	1.01	0.00	1.13	0.00	0.00	1.79
time (sec)	N/A	0.085	0.143	0.220	0.000	0.454	0.000	0.000	6.239

Problem 471	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	62	102	0	126	0	0	192
N.S.	1	1.00	0.51	0.84	0.00	1.04	0.00	0.00	1.59
time (sec)	N/A	0.109	0.187	0.267	0.000	0.432	0.000	0.000	6.414

Problem 472	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	490	490	176	572	0	0	0	0	-1
N.S.	1	1.00	0.36	1.17	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.369	0.977	0.290	0.000	0.000	0.000	0.000	0.000

Problem 473	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	453	453	165	546	0	0	0	0	-1
N.S.	1	1.00	0.36	1.21	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.298	0.815	0.247	0.000	0.000	0.000	0.000	0.000

Problem 474	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	418	418	151	516	0	0	0	0	-1
N.S.	1	1.00	0.36	1.23	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.231	0.632	0.224	0.000	0.000	0.000	0.000	0.000

Problem 475	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F(-1)	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	411	411	144	957	0	0	0	0	-1
N.S.	1	1.00	0.35	2.33	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.244	0.576	0.218	0.000	0.000	0.000	0.000	0.000

Problem 476	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	45	40	0	73	0	0	84
N.S.	1	1.00	1.29	1.14	0.00	2.09	0.00	0.00	2.40
time (sec)	N/A	0.037	0.085	0.181	0.000	0.415	0.000	0.000	1.862

Problem 477	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	172	172	103	246	0	0	0	0	-1
N.S.	1	1.00	0.60	1.43	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.186	10.626	0.253	0.000	0.000	0.000	0.000	0.000

Problem 478	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	135	135	87	218	0	0	0	0	-1
N.S.	1	1.00	0.64	1.61	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.147	0.312	0.263	0.000	0.000	0.000	0.000	0.000

Problem 479	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	84	190	0	0	0	0	-1
N.S.	1	1.00	0.89	2.02	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.107	10.414	0.267	0.000	0.000	0.000	0.000	0.000

Problem 480	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	78	284	0	139	0	0	-1
N.S.	1	1.00	0.78	2.84	0.00	1.39	0.00	0.00	-0.01
time (sec)	N/A	0.110	10.376	0.201	0.000	0.116	0.000	0.000	0.000

Problem 481	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	137	119	539	0	190	0	0	-1
N.S.	1	1.00	0.87	3.93	0.00	1.39	0.00	0.00	-0.01
time (sec)	N/A	0.145	10.635	0.230	0.000	0.116	0.000	0.000	0.000

Problem 482	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	174	174	131	793	0	246	0	0	-1
N.S.	1	1.00	0.75	4.56	0.00	1.41	0.00	0.00	-0.01
time (sec)	N/A	0.197	10.878	0.260	0.000	0.125	0.000	0.000	0.000

Problem 483	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	96	0	0	0	0	0	-1
N.S.	1	1.00	1.28	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	9.753	0.142	0.000	0.000	0.000	0.000	0.000

Problem 484	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	96	0	0	0	0	0	-1
N.S.	1	1.00	1.28	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	5.860	0.119	0.000	0.000	0.000	0.000	0.000

Problem 485	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	106	0	0	0	0	0	-1
N.S.	1	1.00	1.38	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.064	5.692	0.120	0.000	0.000	0.000	0.000	0.000

Problem 486	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	289	0	0	0	0	0	-1
N.S.	1	1.00	3.75	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.068	18.673	0.115	0.000	0.000	0.000	0.000	0.000

Problem 487	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	116	0	0	0	0	0	-1
N.S.	1	1.00	1.51	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.079	17.302	0.108	0.000	0.000	0.000	0.000	0.000

Problem 488	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	285	0	0	0	0	0	-1
N.S.	1	1.00	3.31	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.064	1.011	0.096	0.000	0.000	0.000	0.000	0.000

Problem 489	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	287	0	0	0	0	0	-1
N.S.	1	1.00	3.22	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.198	0.082	0.000	0.000	0.000	0.000	0.000

Problem 490	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	287	0	0	0	0	0	-1
N.S.	1	1.00	3.22	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.064	0.193	0.076	0.000	0.000	0.000	0.000	0.000

Problem 491	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	289	0	0	0	0	0	-1
N.S.	1	1.00	3.14	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.066	0.194	0.060	0.000	0.000	0.000	0.000	0.000

Problem 492	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	80	80	80	0	91	89	0	0	134
N.S.	1	1.00	1.00	0.00	1.14	1.11	0.00	0.00	1.68
time (sec)	N/A	0.050	0.273	0.228	0.319	0.380	0.000	0.000	1.601

Problem 493	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	A	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	47	1732	63	56	0	0	67
N.S.	1	1.00	0.90	33.31	1.21	1.08	0.00	0.00	1.29
time (sec)	N/A	0.036	0.114	1.218	0.282	0.353	0.000	0.000	0.935

Problem 494	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	22	120	30	30	0	0	27
N.S.	1	1.00	0.88	4.80	1.20	1.20	0.00	0.00	1.08
time (sec)	N/A	0.022	0.017	0.784	0.313	0.370	0.000	0.000	0.195

Problem 495	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	92	0	0	0	0	0	-1
N.S.	1	1.00	1.88	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.241	0.050	0.000	0.000	0.000	0.000	0.000

Problem 496	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	201	0	0	0	0	0	-1
N.S.	1	1.00	4.19	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.038	3.090	0.051	0.000	0.000	0.000	0.000	0.000

Problem 497	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F(-2)	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	8327	0	0	0	0	0	-1
N.S.	1	1.00	114.07	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.061	25.273	0.263	0.000	0.000	0.000	0.000	0.000

Problem 498	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	6192	0	0	0	0	0	-1
N.S.	1	1.00	84.82	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	22.008	0.237	0.000	0.000	0.000	0.000	0.000

Problem 499	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	4143	0	0	0	0	0	-1
N.S.	1	1.00	56.75	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	18.011	0.172	0.000	0.000	0.000	0.000	0.000

Problem 500	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	61	0	0	0	0	0	-1
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.048	0.009	0.000	0.000	0.000	0.000	0.000

Problem 501	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	2638	0	0	0	0	0	-1
N.S.	1	1.00	36.14	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	13.744	0.057	0.000	0.000	0.000	0.000	0.000

Problem 502	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	3833	0	0	0	0	0	-1
N.S.	1	1.00	52.51	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	15.361	0.055	0.000	0.000	0.000	0.000	0.000

Problem 503	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	104	0	0	0	0	0	-1
N.S.	1	1.00	1.37	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.076	42.084	0.079	0.000	0.000	0.000	0.000	0.000

Problem 504	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	75	0	0	0	0	0	-1
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.067	10.126	0.076	0.000	0.000	0.000	0.000	0.000

Problem 505	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	72	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.073	10.111	0.058	0.000	0.000	0.000	0.000	0.000

Problem 506	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	78	78	73	0	0	0	0	0	-1
N.S.	1	1.00	0.94	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.086	10.109	0.057	0.000	0.000	0.000	0.000	0.000

Problem 507	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	67	214	0	103	0	0	-1
N.S.	1	1.00	0.67	2.14	0.00	1.03	0.00	0.00	-0.01
time (sec)	N/A	0.053	0.135	1.502	0.000	0.101	0.000	0.000	0.000

Problem 508	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	62	538	0	100	0	0	-1
N.S.	1	1.00	0.83	7.17	0.00	1.33	0.00	0.00	-0.01
time (sec)	N/A	0.041	0.086	0.142	0.000	0.119	0.000	0.000	0.000

Problem 509	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	55	187	0	89	0	0	-1
N.S.	1	1.00	0.76	2.60	0.00	1.24	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.058	0.207	0.000	0.115	0.000	0.000	0.000

Problem 510	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	43	513	0	63	0	0	-1
N.S.	1	1.00	0.98	11.66	0.00	1.43	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.036	0.205	0.000	0.098	0.000	0.000	0.000

Problem 511	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	162	0	59	0	0	63
N.S.	1	1.00	0.98	3.77	0.00	1.37	0.00	0.00	1.47
time (sec)	N/A	0.016	0.029	0.115	0.000	0.099	0.000	0.000	0.619

Problem 512	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	68	68	57	514	0	86	0	0	-1
N.S.	1	1.00	0.84	7.56	0.00	1.26	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.096	0.147	0.000	0.091	0.000	0.000	0.000

Problem 513	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	55	319	0	105	0	0	-1
N.S.	1	1.00	0.74	4.31	0.00	1.42	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.072	0.118	0.000	0.097	0.000	0.000	0.000

Problem 514	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	100	100	68	1055	0	138	0	0	-1
N.S.	1	1.00	0.68	10.55	0.00	1.38	0.00	0.00	-0.01
time (sec)	N/A	0.042	0.108	0.138	0.000	0.116	0.000	0.000	0.000

Problem 515	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	68	216	0	107	0	0	-1
N.S.	1	1.00	0.66	2.10	0.00	1.04	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.084	0.119	0.000	0.127	0.000	0.000	0.000

Problem 516	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	62	544	0	105	0	0	-1
N.S.	1	1.00	0.81	7.06	0.00	1.36	0.00	0.00	-0.01
time (sec)	N/A	0.042	0.134	0.119	0.000	0.100	0.000	0.000	0.000

Problem 517	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	56	189	0	92	0	0	-1
N.S.	1	1.00	0.75	2.52	0.00	1.23	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.043	0.113	0.000	0.120	0.000	0.000	0.000

Problem 518	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	45	531	0	65	0	0	-1
N.S.	1	1.00	0.98	11.54	0.00	1.41	0.00	0.00	-0.02
time (sec)	N/A	0.030	0.020	0.131	0.000	0.102	0.000	0.000	0.000

Problem 519	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	44	44	43	162	0	61	0	0	-1
N.S.	1	1.00	0.98	3.68	0.00	1.39	0.00	0.00	-0.02
time (sec)	N/A	0.022	0.015	0.132	0.000	0.098	0.000	0.000	0.000

Problem 520	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	54	520	0	89	0	0	-1
N.S.	1	1.00	0.76	7.32	0.00	1.25	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.016	0.099	0.000	0.094	0.000	0.000	0.000

Problem 521	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	58	319	0	108	0	0	-1
N.S.	1	1.00	0.81	4.43	0.00	1.50	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.129	0.118	0.000	0.095	0.000	0.000	0.000

Problem 522	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	68	1055	0	148	0	0	-1
N.S.	1	1.00	0.66	10.24	0.00	1.44	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.182	0.149	0.000	0.095	0.000	0.000	0.000

Problem 523	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	70	208	0	106	0	0	-1
N.S.	1	1.00	0.69	2.04	0.00	1.04	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.068	0.122	0.000	0.107	0.000	0.000	0.000

Problem 524	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	57	546	0	103	0	0	-1
N.S.	1	1.00	0.79	7.58	0.00	1.43	0.00	0.00	-0.01
time (sec)	N/A	0.039	0.115	0.141	0.000	0.098	0.000	0.000	0.000

Problem 525	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	64	181	0	92	0	0	-1
N.S.	1	1.00	0.86	2.45	0.00	1.24	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.056	0.135	0.000	0.178	0.000	0.000	0.000

Problem 526	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	42	533	0	66	0	0	-1
N.S.	1	1.00	0.98	12.40	0.00	1.53	0.00	0.00	-0.02
time (sec)	N/A	0.015	0.014	0.118	0.000	0.125	0.000	0.000	0.000

Problem 527	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	45	165	0	62	0	0	-1
N.S.	1	1.00	0.98	3.59	0.00	1.35	0.00	0.00	-0.02
time (sec)	N/A	0.017	0.014	0.187	0.000	0.157	0.000	0.000	0.000

Problem 528	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	70	70	52	522	0	89	0	0	-1
N.S.	1	1.00	0.74	7.46	0.00	1.27	0.00	0.00	-0.01
time (sec)	N/A	0.031	0.067	0.126	0.000	0.099	0.000	0.000	0.000

Problem 529	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	60	319	0	108	0	0	-1
N.S.	1	1.00	0.78	4.14	0.00	1.40	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.046	0.118	0.000	0.093	0.000	0.000	0.000

Problem 530	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	103	103	70	216	0	106	0	0	-1
N.S.	1	1.00	0.68	2.10	0.00	1.03	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.069	0.115	0.000	0.125	0.000	0.000	0.000

Problem 531	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	74	74	60	547	0	103	0	0	-1
N.S.	1	1.00	0.81	7.39	0.00	1.39	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.027	0.134	0.000	0.118	0.000	0.000	0.000

Problem 532	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	63	189	0	92	0	0	-1
N.S.	1	1.00	0.82	2.45	0.00	1.19	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.015	0.101	0.000	0.102	0.000	0.000	0.000

Problem 533	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	45	533	0	66	0	0	-1
N.S.	1	1.00	0.98	11.59	0.00	1.43	0.00	0.00	-0.02
time (sec)	N/A	0.016	0.021	0.116	0.000	0.116	0.000	0.000	0.000

Problem 534	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	45	165	0	62	0	0	-1
N.S.	1	1.00	0.98	3.59	0.00	1.35	0.00	0.00	-0.02
time (sec)	N/A	0.017	0.015	0.122	0.000	0.136	0.000	0.000	0.000

Problem 535	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	55	510	0	89	0	0	-1
N.S.	1	1.00	0.75	6.99	0.00	1.22	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.037	0.113	0.000	0.119	0.000	0.000	0.000

Problem 536	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	60	319	0	108	0	0	-1
N.S.	1	1.00	0.78	4.14	0.00	1.40	0.00	0.00	-0.01
time (sec)	N/A	0.033	0.050	0.125	0.000	0.099	0.000	0.000	0.000

Problem 537	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	C	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	105	105	73	1054	0	144	0	0	-1
N.S.	1	1.00	0.70	10.04	0.00	1.37	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.085	0.125	0.000	0.093	0.000	0.000	0.000

Problem 538	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	102	0	0	0	0	0	-1
N.S.	1	1.00	1.17	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.048	7.148	0.066	0.000	0.000	0.000	0.000	0.000

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [257] had the largest ratio of [25]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	1	1	1.00	6	0.167
2	A	2	2	1.00	8	0.250
3	A	2	1	1.00	8	0.125
4	A	3	2	1.00	8	0.250
5	A	2	1	1.00	8	0.125
6	A	4	2	1.00	8	0.250
7	A	2	1	1.00	8	0.125
8	A	5	2	1.00	8	0.250
9	A	3	2	1.00	8	0.250
10	A	2	2	1.00	8	0.250
11	A	2	2	1.00	8	0.250
12	A	1	1	1.00	8	0.125
13	A	1	1	1.00	8	0.125
14	A	2	2	1.00	8	0.250
15	A	2	2	1.00	8	0.250
16	A	3	2	1.00	8	0.250
17	A	3	2	1.00	10	0.200
18	A	2	2	1.00	10	0.200
19	A	2	2	1.00	10	0.200
20	A	1	1	1.00	10	0.100
21	A	1	1	1.00	10	0.100
22	A	2	2	1.00	10	0.200
23	A	2	2	1.00	10	0.200
24	A	3	2	1.00	10	0.200
25	A	4	3	1.00	12	0.250

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
26	A	3	3	1.00	12	0.250
27	A	3	3	1.00	12	0.250
28	A	2	2	1.00	12	0.167
29	A	2	2	1.00	12	0.167
30	A	3	3	1.00	12	0.250
31	A	3	3	1.00	12	0.250
32	A	4	3	1.00	12	0.250
33	A	1	1	1.00	12	0.083
34	A	1	1	1.00	12	0.083
35	C	1	1	0.11	12	0.083
36	C	1	1	0.23	12	0.083
37	C	1	1	0.21	12	0.083
38	A	1	1	1.00	12	0.083
39	A	1	1	1.00	8	0.125
40	A	1	1	1.00	10	0.100
41	A	2	2	1.00	21	0.095
42	A	2	2	1.00	15	0.133
43	A	2	2	1.00	15	0.133
44	A	2	2	1.00	13	0.154
45	A	1	1	1.00	6	0.167
46	A	2	2	1.00	13	0.154
47	A	2	2	1.00	15	0.133
48	A	2	2	1.00	15	0.133
49	A	3	2	1.00	17	0.118
50	A	3	2	1.00	17	0.118
51	A	3	2	1.00	17	0.118
52	A	2	2	1.00	15	0.133
53	A	2	2	1.00	8	0.250
54	A	2	2	1.00	17	0.118
55	A	3	2	1.00	17	0.118
56	A	3	2	1.00	17	0.118
57	A	3	2	1.00	17	0.118
58	A	5	3	1.00	17	0.176
59	A	4	3	1.00	17	0.176
60	A	3	3	1.00	17	0.176

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
61	A	2	2	1.00	8	0.250
62	A	3	3	1.00	13	0.231
63	A	2	2	1.00	15	0.133
64	A	3	3	1.00	17	0.176
65	A	4	3	1.00	17	0.176
66	A	3	2	1.00	17	0.118
67	A	3	2	1.00	17	0.118
68	A	3	2	1.00	17	0.118
69	A	3	2	1.00	17	0.118
70	A	2	2	1.00	15	0.133
71	A	3	2	1.00	15	0.133
72	A	3	2	1.00	15	0.133
73	A	2	2	1.00	8	0.250
74	A	2	1	1.00	15	0.067
75	A	2	2	1.00	17	0.118
76	A	3	2	1.00	17	0.118
77	A	3	2	1.00	17	0.118
78	A	3	2	1.00	17	0.118
79	A	3	2	1.00	17	0.118
80	A	3	2	1.00	17	0.118
81	A	3	2	1.00	17	0.118
82	A	3	2	1.00	17	0.118
83	A	2	2	1.00	15	0.133
84	A	4	4	1.00	17	0.235
85	A	3	2	1.00	8	0.250
86	A	2	2	1.00	17	0.118
87	A	3	2	1.00	17	0.118
88	A	3	2	1.00	17	0.118
89	A	6	3	1.00	17	0.176
90	A	5	3	1.00	17	0.176
91	A	4	3	1.00	17	0.176
92	A	3	2	1.00	8	0.250
93	A	4	3	1.00	15	0.200
94	A	4	4	1.00	15	0.267
95	A	3	2	1.00	15	0.133

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
96	A	4	3	1.00	17	0.176
97	A	5	3	1.00	17	0.176
98	A	4	3	1.00	17	0.176
99	A	3	2	1.00	17	0.118
100	A	4	3	1.00	17	0.176
101	A	3	2	1.00	17	0.118
102	A	3	2	1.00	17	0.118
103	A	3	2	1.00	17	0.118
104	A	2	2	1.00	15	0.133
105	A	4	3	1.00	15	0.200
106	A	3	2	1.00	17	0.118
107	A	4	3	1.00	17	0.176
108	A	3	2	1.00	15	0.133
109	A	3	2	1.00	8	0.250
110	A	3	2	1.00	15	0.133
111	A	2	2	1.00	17	0.118
112	A	3	2	1.00	17	0.118
113	A	3	2	1.00	17	0.118
114	A	3	2	1.00	17	0.118
115	A	4	3	1.00	17	0.176
116	A	3	2	1.00	17	0.118
117	A	4	3	1.00	17	0.176
118	A	5	4	1.00	17	0.235
119	A	3	2	1.00	15	0.133
120	A	4	3	1.00	15	0.200
121	A	4	3	1.00	15	0.200
122	A	4	3	1.00	15	0.200
123	A	3	2	1.00	15	0.133
124	A	3	3	1.00	13	0.231
125	A	1	1	1.00	6	0.167
126	A	2	2	1.00	13	0.154
127	A	3	3	1.00	15	0.200
128	A	3	2	1.00	15	0.133
129	A	4	3	1.00	15	0.200
130	A	4	3	1.00	15	0.200

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
131	A	4	3	1.00	15	0.200
132	A	4	3	1.00	15	0.200
133	A	3	2	1.00	17	0.118
134	A	5	4	1.00	17	0.235
135	A	3	2	1.00	17	0.118
136	A	4	4	1.00	17	0.235
137	A	3	2	1.00	15	0.133
138	A	2	2	1.00	8	0.250
139	A	2	2	1.00	13	0.154
140	A	3	3	1.00	15	0.200
141	A	3	2	1.00	17	0.118
142	A	4	4	1.00	17	0.235
143	A	3	2	1.00	17	0.118
144	A	5	4	1.00	17	0.235
145	A	4	3	1.00	17	0.176
146	A	5	4	1.00	17	0.235
147	A	4	3	1.00	17	0.176
148	A	4	4	1.00	15	0.267
149	A	2	2	1.00	8	0.250
150	A	2	2	1.00	15	0.133
151	A	2	2	1.00	15	0.133
152	A	3	2	1.00	15	0.133
153	A	4	4	1.00	17	0.235
154	A	4	3	1.00	17	0.176
155	A	5	4	1.00	17	0.235
156	A	4	3	1.00	17	0.176
157	A	3	2	1.00	17	0.118
158	A	6	4	1.00	17	0.235
159	A	3	2	1.00	17	0.118
160	A	5	4	1.00	17	0.235
161	A	3	2	1.00	15	0.133
162	A	3	2	1.00	8	0.250
163	A	2	1	1.00	15	0.067
164	A	2	2	1.00	17	0.118
165	A	2	2	1.00	15	0.133

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
166	A	4	3	1.00	15	0.200
167	A	3	2	1.00	17	0.118
168	A	5	4	1.00	17	0.235
169	A	3	2	1.00	17	0.118
170	A	6	4	1.00	17	0.235
171	A	4	3	1.00	17	0.176
172	A	6	4	1.00	17	0.235
173	A	4	3	1.00	17	0.176
174	A	5	4	1.00	15	0.267
175	A	3	2	1.00	8	0.250
176	A	3	2	1.00	15	0.133
177	A	2	2	1.00	17	0.118
178	A	3	3	1.00	17	0.176
179	A	2	2	1.00	15	0.133
180	A	4	3	1.00	15	0.200
181	A	5	4	1.00	17	0.235
182	A	4	3	1.00	17	0.176
183	A	6	4	1.00	17	0.235
184	A	4	3	1.00	17	0.176
185	A	3	2	1.00	9	0.222
186	A	3	2	1.00	9	0.222
187	A	2	2	1.00	19	0.105
188	A	2	2	1.00	19	0.105
189	A	2	2	1.00	19	0.105
190	A	2	2	1.00	19	0.105
191	A	2	2	1.00	19	0.105
192	A	2	2	1.00	19	0.105
193	A	2	2	1.00	19	0.105
194	A	5	4	1.00	21	0.190
195	A	5	4	1.00	21	0.190
196	A	4	4	1.00	21	0.190
197	A	4	4	1.00	21	0.190
198	A	3	3	1.00	21	0.143
199	A	3	3	1.00	21	0.143
200	A	3	3	1.00	21	0.143

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
201	A	3	3	1.00	21	0.143
202	A	4	4	1.00	21	0.190
203	A	4	4	1.00	21	0.190
204	A	3	2	1.00	21	0.095
205	A	3	2	1.00	21	0.095
206	A	3	2	1.00	21	0.095
207	A	3	2	1.00	21	0.095
208	A	3	2	1.00	21	0.095
209	A	3	2	1.00	21	0.095
210	A	3	2	1.00	21	0.095
211	A	6	4	1.00	21	0.190
212	A	6	4	1.00	21	0.190
213	A	5	4	1.00	21	0.190
214	A	5	4	1.00	21	0.190
215	A	4	3	1.00	21	0.143
216	A	4	3	1.00	21	0.143
217	A	4	4	1.00	21	0.190
218	A	4	4	1.00	21	0.190
219	A	4	3	1.00	21	0.143
220	A	4	3	1.00	21	0.143
221	A	3	2	1.00	19	0.105
222	A	7	6	1.00	19	0.316
223	A	7	6	1.00	19	0.316
224	A	6	6	1.00	19	0.316
225	A	6	6	1.00	19	0.316
226	A	5	5	1.00	19	0.263
227	A	5	5	1.00	19	0.263
228	A	6	6	1.00	19	0.316
229	A	6	6	1.00	19	0.316
230	A	7	6	1.00	19	0.316
231	A	7	6	1.00	19	0.316
232	A	5	4	1.00	21	0.190
233	A	4	4	1.00	21	0.190
234	A	4	4	1.00	21	0.190
235	A	3	3	1.00	21	0.143

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
236	A	3	3	1.00	21	0.143
237	A	3	3	1.00	21	0.143
238	A	3	3	1.00	21	0.143
239	A	4	4	1.00	21	0.190
240	A	4	4	1.00	21	0.190
241	A	5	4	1.00	21	0.190
242	A	8	7	1.00	21	0.333
243	A	7	7	1.00	21	0.333
244	A	7	7	1.00	21	0.333
245	A	6	6	1.00	21	0.286
246	A	6	6	1.00	21	0.286
247	A	6	6	1.00	21	0.286
248	A	6	6	1.00	21	0.286
249	A	7	7	1.00	21	0.333
250	A	7	7	1.00	21	0.333
251	A	8	7	1.00	21	0.333
252	A	2	2	1.00	19	0.105
253	A	3	2	1.00	11	0.182
254	A	3	2	1.00	11	0.182
255	A	3	2	1.00	11	0.182
256	A	3	2	1.00	11	0.182
257	A	4	3	1.00	25	0.120
258	A	3	3	1.00	25	0.120
259	A	2	2	1.00	25	0.080
260	A	3	3	1.00	25	0.120
261	A	4	3	1.00	25	0.120
262	A	11	8	1.00	25	0.320
263	A	10	7	1.00	25	0.280
264	A	1	1	1.00	25	0.040
265	A	2	2	1.00	25	0.080
266	A	3	2	1.00	25	0.080
267	A	4	4	1.00	25	0.160
268	A	3	3	1.00	25	0.120
269	A	3	3	1.00	25	0.120
270	A	4	4	1.00	25	0.160

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
271	A	11	8	1.00	25	0.320
272	A	11	8	1.00	25	0.320
273	A	1	1	1.00	25	0.040
274	A	3	3	1.00	25	0.120
275	A	4	3	1.00	25	0.120
276	A	5	4	1.00	25	0.160
277	A	4	4	1.00	25	0.160
278	A	3	3	1.00	25	0.120
279	A	3	3	1.00	25	0.120
280	A	4	4	1.00	25	0.160
281	A	5	4	1.00	25	0.160
282	A	11	8	1.00	25	0.320
283	A	11	8	1.00	25	0.320
284	A	1	1	1.00	25	0.040
285	A	3	3	1.00	25	0.120
286	A	4	3	1.00	25	0.120
287	A	12	8	1.00	21	0.381
288	A	1	1	1.00	13	0.077
289	A	10	7	1.00	13	0.538
290	A	11	8	1.00	13	0.615
291	A	4	3	1.00	25	0.120
292	A	3	3	1.00	25	0.120
293	A	2	2	1.00	25	0.080
294	A	3	3	1.00	25	0.120
295	A	4	3	1.00	25	0.120
296	A	10	7	1.00	25	0.280
297	A	1	1	1.00	25	0.040
298	A	2	2	1.00	25	0.080
299	A	3	2	1.00	25	0.080
300	A	10	7	1.00	21	0.333
301	A	11	8	1.00	21	0.381
302	A	11	8	1.00	21	0.381
303	A	12	8	1.00	21	0.381
304	A	1	1	1.00	21	0.048
305	A	1	1	1.00	21	0.048

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
306	A	1	1	1.00	12	0.083
307	A	1	1	1.00	21	0.048
308	A	1	1	1.00	21	0.048
309	A	1	1	1.00	21	0.048
310	A	1	1	1.00	21	0.048
311	A	1	1	1.00	12	0.083
312	A	1	1	1.00	21	0.048
313	A	1	1	1.00	21	0.048
314	A	1	1	1.00	21	0.048
315	A	1	1	1.00	21	0.048
316	A	1	1	1.00	12	0.083
317	A	1	1	1.00	21	0.048
318	A	1	1	1.00	21	0.048
319	A	1	1	1.00	21	0.048
320	A	1	1	1.00	21	0.048
321	A	1	1	1.00	12	0.083
322	A	1	1	1.00	21	0.048
323	A	1	1	1.00	21	0.048
324	A	8	8	1.00	21	0.381
325	A	11	7	1.00	21	0.333
326	A	12	8	1.00	21	0.381
327	A	9	9	1.00	21	0.429
328	A	9	9	1.00	21	0.429
329	A	8	8	1.00	21	0.381
330	A	11	7	1.00	21	0.333
331	A	12	8	1.00	21	0.381
332	A	9	9	1.00	21	0.429
333	A	9	9	1.00	21	0.429
334	A	1	1	1.00	13	0.077
335	A	1	1	1.00	13	0.077
336	A	1	1	1.00	17	0.059
337	A	1	1	1.00	19	0.053
338	A	1	1	1.00	19	0.053
339	A	1	1	1.00	21	0.048
340	A	3	2	1.00	19	0.105

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
341	A	3	2	1.00	19	0.105
342	A	2	2	1.00	17	0.118
343	A	2	2	1.00	17	0.118
344	A	2	2	1.00	19	0.105
345	A	1	1	1.00	19	0.053
346	A	1	1	1.00	19	0.053
347	A	1	1	1.00	10	0.100
348	A	1	1	1.00	19	0.053
349	A	1	1	1.00	19	0.053
350	A	1	1	1.00	23	0.043
351	A	1	1	1.00	23	0.043
352	A	1	1	1.00	23	0.043
353	A	1	1	1.00	23	0.043
354	A	1	1	1.00	23	0.043
355	A	3	2	1.00	19	0.105
356	A	3	2	1.00	19	0.105
357	A	2	2	1.00	17	0.118
358	A	2	2	1.00	17	0.118
359	A	2	2	1.00	19	0.105
360	A	2	2	1.00	19	0.105
361	A	1	1	1.00	19	0.053
362	A	1	1	1.00	19	0.053
363	A	1	1	1.00	10	0.100
364	A	1	1	1.00	19	0.053
365	A	1	1	1.00	19	0.053
366	A	1	1	1.00	23	0.043
367	A	1	1	1.00	23	0.043
368	A	1	1	1.00	23	0.043
369	A	1	1	1.00	23	0.043
370	A	1	1	1.00	23	0.043
371	A	3	2	1.00	21	0.095
372	A	3	2	1.00	21	0.095
373	A	3	2	1.00	21	0.095
374	A	2	2	1.00	19	0.105
375	A	5	5	1.00	19	0.263

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
376	A	6	6	1.00	21	0.286
377	A	7	6	1.00	21	0.286
378	A	5	3	1.00	21	0.143
379	A	4	3	1.00	21	0.143
380	A	3	3	1.00	21	0.143
381	A	2	2	1.00	12	0.167
382	A	3	3	1.00	21	0.143
383	A	4	3	1.00	21	0.143
384	A	5	3	1.00	21	0.143
385	A	3	2	1.00	21	0.095
386	A	3	2	1.00	21	0.095
387	A	3	2	1.00	21	0.095
388	A	2	2	1.00	19	0.105
389	A	6	6	1.00	19	0.316
390	A	7	7	1.00	21	0.333
391	A	5	4	1.00	21	0.190
392	A	4	4	1.00	21	0.190
393	A	3	3	1.00	21	0.143
394	A	3	3	1.00	12	0.250
395	A	4	4	1.00	21	0.190
396	A	5	4	1.00	21	0.190
397	A	3	2	1.00	21	0.095
398	A	3	2	1.00	21	0.095
399	A	3	2	1.00	21	0.095
400	A	2	2	1.00	19	0.105
401	A	6	6	1.00	19	0.316
402	A	7	7	1.00	21	0.333
403	A	8	7	1.00	21	0.333
404	A	5	4	1.00	21	0.190
405	A	4	4	1.00	21	0.190
406	A	3	3	1.00	21	0.143
407	A	3	3	1.00	12	0.250
408	A	4	4	1.00	21	0.190
409	A	5	4	1.00	21	0.190
410	A	3	2	1.00	21	0.095

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
411	A	3	2	1.00	21	0.095
412	A	3	2	1.00	21	0.095
413	A	2	2	1.00	19	0.105
414	A	5	5	1.00	19	0.263
415	A	6	6	1.00	21	0.286
416	A	7	6	1.00	21	0.286
417	A	5	3	1.00	21	0.143
418	A	4	3	1.00	21	0.143
419	A	3	3	1.00	21	0.143
420	A	2	2	1.00	12	0.167
421	A	3	3	1.00	21	0.143
422	A	4	3	1.00	21	0.143
423	A	5	3	1.00	21	0.143
424	A	3	2	1.00	21	0.095
425	A	3	2	1.00	21	0.095
426	A	3	2	1.00	21	0.095
427	A	2	2	1.00	19	0.105
428	A	6	6	1.00	19	0.316
429	A	6	6	1.00	21	0.286
430	A	7	7	1.00	21	0.333
431	A	5	4	1.00	21	0.190
432	A	4	4	1.00	21	0.190
433	A	3	3	1.00	12	0.250
434	A	3	3	1.00	21	0.143
435	A	4	4	1.00	21	0.190
436	A	5	4	1.00	21	0.190
437	A	3	2	1.00	21	0.095
438	A	3	2	1.00	21	0.095
439	A	3	2	1.00	21	0.095
440	A	2	2	1.00	19	0.105
441	A	6	6	1.00	19	0.316
442	A	6	6	1.00	21	0.286
443	A	7	7	1.00	21	0.333
444	A	5	4	1.00	21	0.190
445	A	4	4	1.00	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
446	A	3	3	1.00	12	0.250
447	A	3	3	1.00	21	0.143
448	A	4	4	1.00	21	0.190
449	A	5	4	1.00	21	0.190
450	A	13	9	1.00	25	0.360
451	A	12	9	1.00	25	0.360
452	A	11	8	1.00	25	0.320
453	A	1	1	1.00	25	0.040
454	A	2	2	1.00	25	0.080
455	A	3	2	1.00	25	0.080
456	A	5	4	1.00	25	0.160
457	A	4	4	1.00	25	0.160
458	A	3	3	1.00	25	0.120
459	A	4	4	1.00	25	0.160
460	A	5	4	1.00	25	0.160
461	A	5	4	1.00	23	0.174
462	A	4	4	1.00	23	0.174
463	A	3	3	1.00	23	0.130
464	A	4	4	1.00	23	0.174
465	A	5	4	1.00	23	0.174
466	A	12	9	1.00	23	0.391
467	A	11	8	1.00	23	0.348
468	A	1	1	1.00	23	0.043
469	A	2	2	1.00	23	0.087
470	A	3	2	1.00	23	0.087
471	A	4	2	1.00	23	0.087
472	A	14	10	1.00	25	0.400
473	A	13	10	1.00	25	0.400
474	A	12	9	1.00	25	0.360
475	A	12	9	1.00	25	0.360
476	A	1	1	1.00	25	0.040
477	A	6	5	1.00	25	0.200
478	A	5	5	1.00	25	0.200
479	A	4	4	1.00	25	0.160
480	A	4	4	1.00	25	0.160

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
481	A	5	5	1.00	25	0.200
482	A	6	5	1.00	25	0.200
483	A	2	2	1.00	23	0.087
484	A	2	2	1.00	23	0.087
485	A	2	2	1.00	23	0.087
486	A	2	2	1.00	23	0.087
487	A	2	2	1.00	23	0.087
488	A	2	2	1.00	17	0.118
489	A	2	2	1.00	19	0.105
490	A	2	2	1.00	19	0.105
491	A	2	2	1.00	21	0.095
492	A	3	2	1.00	19	0.105
493	A	3	2	1.00	19	0.105
494	A	2	2	1.00	17	0.118
495	A	2	2	1.00	17	0.118
496	A	2	2	1.00	19	0.105
497	A	2	2	1.00	19	0.105
498	A	2	2	1.00	19	0.105
499	A	2	2	1.00	19	0.105
500	A	2	2	1.00	10	0.200
501	A	2	2	1.00	19	0.105
502	A	2	2	1.00	19	0.105
503	A	2	2	1.00	23	0.087
504	A	2	2	1.00	23	0.087
505	A	2	2	1.00	23	0.087
506	A	2	2	1.00	23	0.087
507	A	5	4	1.00	21	0.190
508	A	4	4	1.00	21	0.190
509	A	4	4	1.00	21	0.190
510	A	3	3	1.00	19	0.158
511	A	2	2	1.00	12	0.167
512	A	4	4	1.00	19	0.210
513	A	4	4	1.00	21	0.190
514	A	5	4	1.00	21	0.190
515	A	5	4	1.00	21	0.190

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
516	A	4	4	1.00	21	0.190
517	A	4	4	1.00	21	0.190
518	A	3	3	1.00	21	0.143
519	A	3	3	1.00	19	0.158
520	A	3	3	1.00	12	0.250
521	A	4	4	1.00	19	0.210
522	A	5	4	1.00	21	0.190
523	A	5	4	1.00	21	0.190
524	A	4	4	1.00	21	0.190
525	A	4	4	1.00	19	0.210
526	A	2	2	1.00	12	0.167
527	A	3	3	1.00	19	0.158
528	A	4	4	1.00	21	0.190
529	A	4	4	1.00	21	0.190
530	A	5	4	1.00	21	0.190
531	A	4	4	1.00	19	0.210
532	A	3	3	1.00	12	0.250
533	A	3	3	1.00	19	0.158
534	A	3	3	1.00	21	0.143
535	A	4	4	1.00	21	0.190
536	A	4	4	1.00	21	0.190
537	A	5	4	1.00	21	0.190
538	A	2	2	1.00	21	0.095

Chapter 3

Listing of integrals

Local contents

3.1	$\int \sin(a + bx) dx$	152
3.2	$\int \sin^2(a + bx) dx$	155
3.3	$\int \sin^3(a + bx) dx$	158
3.4	$\int \sin^4(a + bx) dx$	161
3.5	$\int \sin^5(a + bx) dx$	164
3.6	$\int \sin^6(a + bx) dx$	167
3.7	$\int \sin^7(a + bx) dx$	170
3.8	$\int \sin^8(a + bx) dx$	173
3.9	$\int \sin^{\frac{7}{2}}(bx) dx$	177
3.10	$\int \sin^{\frac{5}{2}}(bx) dx$	180
3.11	$\int \sin^{\frac{3}{2}}(bx) dx$	183
3.12	$\int \sqrt{\sin(bx)} dx$	186
3.13	$\int \frac{1}{\sqrt{\sin(bx)}} dx$	189
3.14	$\int \frac{1}{\sin^{\frac{3}{2}}(bx)} dx$	192
3.15	$\int \frac{1}{\sin^{\frac{5}{2}}(bx)} dx$	195
3.16	$\int \frac{1}{\sin^{\frac{7}{2}}(bx)} dx$	198
3.17	$\int \sin^{\frac{7}{2}}(a + bx) dx$	201
3.18	$\int \sin^{\frac{5}{2}}(a + bx) dx$	204
3.19	$\int \sin^{\frac{3}{2}}(a + bx) dx$	207
3.20	$\int \sqrt{\sin(a + bx)} dx$	210
3.21	$\int \frac{1}{\sqrt{\sin(a + bx)}} dx$	213
3.22	$\int \frac{1}{\sin^{\frac{3}{2}}(a+bx)} dx$	216
3.23	$\int \frac{1}{\sin^{\frac{5}{2}}(a+bx)} dx$	219
3.24	$\int \frac{1}{\sin^{\frac{7}{2}}(a+bx)} dx$	222

3.25	$\int (c \sin(a + bx))^{7/2} dx$	226
3.26	$\int (c \sin(a + bx))^{5/2} dx$	229
3.27	$\int (c \sin(a + bx))^{3/2} dx$	232
3.28	$\int \sqrt{c \sin(a + bx)} dx$	235
3.29	$\int \frac{1}{\sqrt{c \sin(a + bx)}} dx$	238
3.30	$\int \frac{1}{(c \sin(a + bx))^{3/2}} dx$	241
3.31	$\int \frac{1}{(c \sin(a + bx))^{5/2}} dx$	244
3.32	$\int \frac{1}{(c \sin(a + bx))^{7/2}} dx$	247
3.33	$\int (c \sin(a + bx))^{4/3} dx$	251
3.34	$\int (c \sin(a + bx))^{2/3} dx$	254
3.35	$\int \sqrt[3]{c \sin(a + bx)} dx$	257
3.36	$\int \frac{1}{\sqrt[3]{c \sin(a + bx)}} dx$	260
3.37	$\int \frac{1}{(c \sin(a + bx))^{2/3}} dx$	263
3.38	$\int \frac{1}{(c \sin(a + bx))^{4/3}} dx$	266
3.39	$\int \sin^n(a + bx) dx$	269
3.40	$\int (c \sin(a + bx))^n dx$	272
3.41	$\int (a \sin(e + fx))^m (b \sin(e + fx))^n dx$	275
3.42	$\int \cos^3(a + bx) \sin(a + bx) dx$	278
3.43	$\int \cos^2(a + bx) \sin(a + bx) dx$	281
3.44	$\int \cos(a + bx) \sin(a + bx) dx$	284
3.45	$\int \tan(a + bx) dx$	287
3.46	$\int \sec(a + bx) \tan(a + bx) dx$	290
3.47	$\int \sec^2(a + bx) \tan(a + bx) dx$	293
3.48	$\int \sec^3(a + bx) \tan(a + bx) dx$	296
3.49	$\int \cos^7(a + bx) \sin^2(a + bx) dx$	299
3.50	$\int \cos^5(a + bx) \sin^2(a + bx) dx$	302
3.51	$\int \cos^3(a + bx) \sin^2(a + bx) dx$	305
3.52	$\int \cos(a + bx) \sin^2(a + bx) dx$	308
3.53	$\int \tan^2(a + bx) dx$	311
3.54	$\int \sec^2(a + bx) \tan^2(a + bx) dx$	314
3.55	$\int \sec^4(a + bx) \tan^2(a + bx) dx$	317
3.56	$\int \sec^6(a + bx) \tan^2(a + bx) dx$	320
3.57	$\int \sec^8(a + bx) \tan^2(a + bx) dx$	323
3.58	$\int \cos^6(a + bx) \sin^2(a + bx) dx$	326
3.59	$\int \cos^4(a + bx) \sin^2(a + bx) dx$	330
3.60	$\int \cos^2(a + bx) \sin^2(a + bx) dx$	334
3.61	$\int \sin^2(a + bx) dx$	337
3.62	$\int \sin(a + bx) \tan(a + bx) dx$	340
3.63	$\int \sec(a + bx) \tan^2(a + bx) dx$	345
3.64	$\int \sec^3(a + bx) \tan^2(a + bx) dx$	348
3.65	$\int \sec^5(a + bx) \tan^2(a + bx) dx$	352
3.66	$\int \cos^5(a + bx) \sin^3(a + bx) dx$	356

3.67	$\int \cos^4(a + bx) \sin^3(a + bx) dx$	359
3.68	$\int \cos^3(a + bx) \sin^3(a + bx) dx$	362
3.69	$\int \cos^2(a + bx) \sin^3(a + bx) dx$	365
3.70	$\int \cos(a + bx) \sin^3(a + bx) dx$	368
3.71	$\int \sin^2(a + bx) \tan(a + bx) dx$	371
3.72	$\int \sin(a + bx) \tan^2(a + bx) dx$	374
3.73	$\int \tan^3(a + bx) dx$	377
3.74	$\int \sec(a + bx) \tan^3(a + bx) dx$	380
3.75	$\int \sec^2(a + bx) \tan^3(a + bx) dx$	383
3.76	$\int \sec^3(a + bx) \tan^3(a + bx) dx$	386
3.77	$\int \sec^4(a + bx) \tan^3(a + bx) dx$	389
3.78	$\int \sec^5(a + bx) \tan^3(a + bx) dx$	392
3.79	$\int \sec^6(a + bx) \tan^3(a + bx) dx$	395
3.80	$\int \cos^7(a + bx) \sin^4(a + bx) dx$	398
3.81	$\int \cos^5(a + bx) \sin^4(a + bx) dx$	401
3.82	$\int \cos^3(a + bx) \sin^4(a + bx) dx$	404
3.83	$\int \cos(a + bx) \sin^4(a + bx) dx$	407
3.84	$\int \sin^2(a + bx) \tan^2(a + bx) dx$	410
3.85	$\int \tan^4(a + bx) dx$	414
3.86	$\int \sec^2(a + bx) \tan^4(a + bx) dx$	417
3.87	$\int \sec^4(a + bx) \tan^4(a + bx) dx$	420
3.88	$\int \sec^6(a + bx) \tan^4(a + bx) dx$	423
3.89	$\int \cos^6(a + bx) \sin^4(a + bx) dx$	426
3.90	$\int \cos^4(a + bx) \sin^4(a + bx) dx$	430
3.91	$\int \cos^2(a + bx) \sin^4(a + bx) dx$	434
3.92	$\int \sin^4(a + bx) dx$	438
3.93	$\int \sin^3(a + bx) \tan(a + bx) dx$	441
3.94	$\int \sin(a + bx) \tan^3(a + bx) dx$	444
3.95	$\int \sec(a + bx) \tan^4(a + bx) dx$	448
3.96	$\int \sec^3(a + bx) \tan^4(a + bx) dx$	451
3.97	$\int \sec^5(a + bx) \tan^4(a + bx) dx$	455
3.98	$\int \cos^7(a + bx) \sin^5(a + bx) dx$	459
3.99	$\int \cos^6(a + bx) \sin^5(a + bx) dx$	463
3.100	$\int \cos^5(a + bx) \sin^5(a + bx) dx$	466
3.101	$\int \cos^4(a + bx) \sin^5(a + bx) dx$	469
3.102	$\int \cos^3(a + bx) \sin^5(a + bx) dx$	472
3.103	$\int \cos^2(a + bx) \sin^5(a + bx) dx$	475
3.104	$\int \cos(a + bx) \sin^5(a + bx) dx$	478
3.105	$\int \sin^4(a + bx) \tan(a + bx) dx$	481
3.106	$\int \sin^3(a + bx) \tan^2(a + bx) dx$	485
3.107	$\int \sin^2(a + bx) \tan^3(a + bx) dx$	488
3.108	$\int \sin(a + bx) \tan^4(a + bx) dx$	492
3.109	$\int \tan^5(a + bx) dx$	495
3.110	$\int \sec(a + bx) \tan^5(a + bx) dx$	498
3.111	$\int \sec^2(a + bx) \tan^5(a + bx) dx$	501

3.112	$\int \sec^3(a + bx) \tan^5(a + bx) dx$	504
3.113	$\int \sec^4(a + bx) \tan^5(a + bx) dx$	507
3.114	$\int \sec^5(a + bx) \tan^5(a + bx) dx$	510
3.115	$\int \sec^6(a + bx) \tan^5(a + bx) dx$	513
3.116	$\int \sec^7(a + bx) \tan^5(a + bx) dx$	517
3.117	$\int \sec^8(a + bx) \tan^5(a + bx) dx$	520
3.118	$\int \sin^3(a + bx) \tan^3(a + bx) dx$	524
3.119	$\int \sin(a + bx) \tan^6(a + bx) dx$	528
3.120	$\int \cos^5(a + bx) \cot(a + bx) dx$	531
3.121	$\int \cos^4(a + bx) \cot(a + bx) dx$	535
3.122	$\int \cos^3(a + bx) \cot(a + bx) dx$	539
3.123	$\int \cos^2(a + bx) \cot(a + bx) dx$	543
3.124	$\int \cos(a + bx) \cot(a + bx) dx$	546
3.125	$\int \cot(a + bx) dx$	550
3.126	$\int \csc(a + bx) \sec(a + bx) dx$	553
3.127	$\int \csc(a + bx) \sec^2(a + bx) dx$	556
3.128	$\int \csc(a + bx) \sec^3(a + bx) dx$	559
3.129	$\int \csc(a + bx) \sec^4(a + bx) dx$	562
3.130	$\int \csc(a + bx) \sec^5(a + bx) dx$	566
3.131	$\int \csc(a + bx) \sec^6(a + bx) dx$	570
3.132	$\int \csc(a + bx) \sec^7(a + bx) dx$	574
3.133	$\int \cos^5(a + bx) \cot^2(a + bx) dx$	578
3.134	$\int \cos^4(a + bx) \cot^2(a + bx) dx$	581
3.135	$\int \cos^3(a + bx) \cot^2(a + bx) dx$	585
3.136	$\int \cos^2(a + bx) \cot^2(a + bx) dx$	588
3.137	$\int \cos(a + bx) \cot^2(a + bx) dx$	592
3.138	$\int \cot^2(a + bx) dx$	595
3.139	$\int \cot(a + bx) \csc(a + bx) dx$	598
3.140	$\int \csc^2(a + bx) \sec(a + bx) dx$	601
3.141	$\int \csc^2(a + bx) \sec^2(a + bx) dx$	604
3.142	$\int \csc^2(a + bx) \sec^3(a + bx) dx$	607
3.143	$\int \csc^2(a + bx) \sec^4(a + bx) dx$	611
3.144	$\int \csc^2(a + bx) \sec^5(a + bx) dx$	614
3.145	$\int \cos^4(a + bx) \cot^3(a + bx) dx$	618
3.146	$\int \cos^3(a + bx) \cot^3(a + bx) dx$	623
3.147	$\int \cos^2(a + bx) \cot^3(a + bx) dx$	627
3.148	$\int \cos(a + bx) \cot^3(a + bx) dx$	631
3.149	$\int \cot^3(a + bx) dx$	635
3.150	$\int \cot^2(a + bx) \csc(a + bx) dx$	638
3.151	$\int \cot(a + bx) \csc^2(a + bx) dx$	641
3.152	$\int \csc^3(a + bx) \sec(a + bx) dx$	644
3.153	$\int \csc^3(a + bx) \sec^2(a + bx) dx$	647
3.154	$\int \csc^3(a + bx) \sec^3(a + bx) dx$	651
3.155	$\int \csc^3(a + bx) \sec^4(a + bx) dx$	655
3.156	$\int \csc^3(a + bx) \sec^5(a + bx) dx$	659

3.157	$\int \cos^5(a + bx) \cot^4(a + bx) dx$	663
3.158	$\int \cos^4(a + bx) \cot^4(a + bx) dx$	666
3.159	$\int \cos^3(a + bx) \cot^4(a + bx) dx$	670
3.160	$\int \cos^2(a + bx) \cot^4(a + bx) dx$	673
3.161	$\int \cos(a + bx) \cot^4(a + bx) dx$	677
3.162	$\int \cot^4(a + bx) dx$	680
3.163	$\int \cot^3(a + bx) \csc(a + bx) dx$	683
3.164	$\int \cot^2(a + bx) \csc^2(a + bx) dx$	686
3.165	$\int \cot(a + bx) \csc^3(a + bx) dx$	689
3.166	$\int \csc^4(a + bx) \sec(a + bx) dx$	692
3.167	$\int \csc^4(a + bx) \sec^2(a + bx) dx$	696
3.168	$\int \csc^4(a + bx) \sec^3(a + bx) dx$	699
3.169	$\int \csc^4(a + bx) \sec^4(a + bx) dx$	703
3.170	$\int \csc^4(a + bx) \sec^5(a + bx) dx$	706
3.171	$\int \cos^4(a + bx) \cot^5(a + bx) dx$	710
3.172	$\int \cos^3(a + bx) \cot^5(a + bx) dx$	715
3.173	$\int \cos^2(a + bx) \cot^5(a + bx) dx$	720
3.174	$\int \cos(a + bx) \cot^5(a + bx) dx$	724
3.175	$\int \cot^5(a + bx) dx$	728
3.176	$\int \cot^4(a + bx) \csc(a + bx) dx$	731
3.177	$\int \cot^3(a + bx) \csc^2(a + bx) dx$	735
3.178	$\int \cot^2(a + bx) \csc^3(a + bx) dx$	738
3.179	$\int \cot(a + bx) \csc^4(a + bx) dx$	742
3.180	$\int \csc^5(a + bx) \sec(a + bx) dx$	745
3.181	$\int \csc^5(a + bx) \sec^2(a + bx) dx$	749
3.182	$\int \csc^5(a + bx) \sec^3(a + bx) dx$	753
3.183	$\int \csc^5(a + bx) \sec^4(a + bx) dx$	757
3.184	$\int \csc^5(a + bx) \sec^5(a + bx) dx$	761
3.185	$\int \cot^2(x) \csc^4(x) dx$	765
3.186	$\int \cot^3(x) \csc^4(x) dx$	768
3.187	$\int (d \cos(a + bx))^{3/2} \sin(a + bx) dx$	771
3.188	$\int \sqrt{d \cos(a + bx)} \sin(a + bx) dx$	774
3.189	$\int \frac{\sin(a+bx)}{\sqrt{d \cos(a + bx)}} dx$	777
3.190	$\int \frac{\sin(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	780
3.191	$\int \frac{\sin(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	783
3.192	$\int \frac{\sin(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	786
3.193	$\int \frac{\sin(a+bx)}{(d \cos(a+bx))^{9/2}} dx$	789
3.194	$\int (d \cos(a + bx))^{9/2} \sin^2(a + bx) dx$	792
3.195	$\int (d \cos(a + bx))^{7/2} \sin^2(a + bx) dx$	796
3.196	$\int (d \cos(a + bx))^{5/2} \sin^2(a + bx) dx$	800
3.197	$\int (d \cos(a + bx))^{3/2} \sin^2(a + bx) dx$	804
3.198	$\int \sqrt{d \cos(a + bx)} \sin^2(a + bx) dx$	808

3.199	$\int \frac{\sin^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx$	811
3.200	$\int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	815
3.201	$\int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	818
3.202	$\int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	822
3.203	$\int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{9/2}} dx$	826
3.204	$\int \sqrt{d \cos(a+bx)} \sin^3(a+bx) dx$	830
3.205	$\int \frac{\sin^3(a+bx)}{\sqrt{d \cos(a+bx)}} dx$	833
3.206	$\int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	836
3.207	$\int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	839
3.208	$\int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	842
3.209	$\int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{9/2}} dx$	845
3.210	$\int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{11/2}} dx$	848
3.211	$\int (d \cos(a+bx))^{9/2} \sin^4(a+bx) dx$	851
3.212	$\int (d \cos(a+bx))^{7/2} \sin^4(a+bx) dx$	855
3.213	$\int (d \cos(a+bx))^{5/2} \sin^4(a+bx) dx$	859
3.214	$\int (d \cos(a+bx))^{3/2} \sin^4(a+bx) dx$	863
3.215	$\int \sqrt{d \cos(a+bx)} \sin^4(a+bx) dx$	867
3.216	$\int \frac{\sin^4(a+bx)}{\sqrt{d \cos(a+bx)}} dx$	870
3.217	$\int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	874
3.218	$\int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	878
3.219	$\int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	882
3.220	$\int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{9/2}} dx$	886
3.221	$\int \cos^{\frac{3}{2}}(a+bx) \sin^5(a+bx) dx$	890
3.222	$\int (d \cos(a+bx))^{9/2} \csc(a+bx) dx$	893
3.223	$\int (d \cos(a+bx))^{7/2} \csc(a+bx) dx$	898
3.224	$\int (d \cos(a+bx))^{5/2} \csc(a+bx) dx$	902
3.225	$\int (d \cos(a+bx))^{3/2} \csc(a+bx) dx$	906
3.226	$\int \sqrt{d \cos(a+bx)} \csc(a+bx) dx$	910
3.227	$\int \frac{\csc(a+bx)}{\sqrt{d \cos(a+bx)}} dx$	914
3.228	$\int \frac{\csc(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	918
3.229	$\int \frac{\csc(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	923
3.230	$\int \frac{\csc(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	928
3.231	$\int \frac{\csc(a+bx)}{(d \cos(a+bx))^{9/2}} dx$	933
3.232	$\int (d \cos(a+bx))^{11/2} \csc^2(a+bx) dx$	938
3.233	$\int (d \cos(a+bx))^{9/2} \csc^2(a+bx) dx$	942

3.234	$\int (d \cos(a + bx))^{7/2} \csc^2(a + bx) dx$	946
3.235	$\int (d \cos(a + bx))^{5/2} \csc^2(a + bx) dx$	950
3.236	$\int (d \cos(a + bx))^{3/2} \csc^2(a + bx) dx$	954
3.237	$\int \sqrt{d \cos(a + bx)} \csc^2(a + bx) dx$	958
3.238	$\int \frac{\csc^2(a+bx)}{\sqrt{d \cos(a + bx)}} dx$	961
3.239	$\int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	965
3.240	$\int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	969
3.241	$\int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	973
3.242	$\int (d \cos(a + bx))^{11/2} \csc^3(a + bx) dx$	977
3.243	$\int (d \cos(a + bx))^{9/2} \csc^3(a + bx) dx$	982
3.244	$\int (d \cos(a + bx))^{7/2} \csc^3(a + bx) dx$	987
3.245	$\int (d \cos(a + bx))^{5/2} \csc^3(a + bx) dx$	992
3.246	$\int (d \cos(a + bx))^{3/2} \csc^3(a + bx) dx$	997
3.247	$\int \sqrt{d \cos(a + bx)} \csc^3(a + bx) dx$	1001
3.248	$\int \frac{\csc^3(a+bx)}{\sqrt{d \cos(a + bx)}} dx$	1006
3.249	$\int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{3/2}} dx$	1011
3.250	$\int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{5/2}} dx$	1017
3.251	$\int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{7/2}} dx$	1023
3.252	$\int \sqrt[5]{d \cos(a + bx)} \sin(a + bx) dx$	1029
3.253	$\int \cos^3(x) \sqrt{\sin(x)} dx$	1032
3.254	$\int \cos^3(x) \sin^{3/2}(x) dx$	1035
3.255	$\int \cos^3(x) \sin^{5/2}(x) dx$	1038
3.256	$\int \frac{\cos^3(x)}{\sqrt{\sin(x)}} dx$	1041
3.257	$\int (d \cos(a + bx))^{9/2} \sqrt{c \sin(a + bx)} dx$	1044
3.258	$\int (d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)} dx$	1048
3.259	$\int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx$	1052
3.260	$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a+bx))^{3/2}} dx$	1055
3.261	$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a+bx))^{7/2}} dx$	1059
3.262	$\int (d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)} dx$	1063
3.263	$\int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx$	1069
3.264	$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a+bx))^{5/2}} dx$	1075
3.265	$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a+bx))^{9/2}} dx$	1078
3.266	$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a+bx))^{13/2}} dx$	1081
3.267	$\int (d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2} dx$	1084

3.268	$\int \frac{(c \sin(a+bx))^{3/2}}{\sqrt{d \cos(a+bx)}} dx$	1088
3.269	$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{5/2}} dx$	1091
3.270	$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{9/2}} dx$	1095
3.271	$\int \sqrt{d \cos(a+bx)} (c \sin(a+bx))^{3/2} dx$	1099
3.272	$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{3/2}} dx$	1105
3.273	$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{7/2}} dx$	1111
3.274	$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{11/2}} dx$	1114
3.275	$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{15/2}} dx$	1118
3.276	$\int (d \cos(a+bx))^{9/2} (c \sin(a+bx))^{5/2} dx$	1122
3.277	$\int (d \cos(a+bx))^{5/2} (c \sin(a+bx))^{5/2} dx$	1126
3.278	$\int \sqrt{d \cos(a+bx)} (c \sin(a+bx))^{5/2} dx$	1130
3.279	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{3/2}} dx$	1134
3.280	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{7/2}} dx$	1138
3.281	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{11/2}} dx$	1142
3.282	$\int \frac{(c \sin(a+bx))^{5/2}}{\sqrt{d \cos(a+bx)}} dx$	1146
3.283	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{5/2}} dx$	1152
3.284	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{9/2}} dx$	1158
3.285	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{13/2}} dx$	1161
3.286	$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{17/2}} dx$	1165
3.287	$\int \frac{\sin^{\frac{7}{2}}(a+bx)}{\cos^{\frac{7}{2}}(a+bx)} dx$	1169
3.288	$\int \frac{\sin^{\frac{3}{2}}(x)}{\cos^{\frac{7}{2}}(x)} dx$	1175
3.289	$\int \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} dx$	1178
3.290	$\int \frac{\sin^{\frac{5}{2}}(x)}{\sqrt{\cos(x)}} dx$	1183
3.291	$\int \frac{(d \cos(a+bx))^{7/2}}{\sqrt{c \sin(a+bx)}} dx$	1189
3.292	$\int \frac{(d \cos(a+bx))^{3/2}}{\sqrt{c \sin(a+bx)}} dx$	1193
3.293	$\int \frac{1}{\sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}} dx$	1196
3.294	$\int \frac{1}{(d \cos(a+bx))^{5/2} \sqrt{c \sin(a+bx)}} dx$	1199
3.295	$\int \frac{1}{(d \cos(a+bx))^{9/2} \sqrt{c \sin(a+bx)}} dx$	1203
3.296	$\int \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} dx$	1207

3.297	$\int \frac{1}{(d \cos(a+bx))^{3/2} \sqrt{c \sin(a+bx)}} dx$	1213
3.298	$\int \frac{1}{(d \cos(a+bx))^{7/2} \sqrt{c \sin(a+bx)}} dx$	1216
3.299	$\int \frac{1}{(d \cos(a+bx))^{11/2} \sqrt{c \sin(a+bx)}} dx$	1219
3.300	$\int \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} dx$	1222
3.301	$\int \frac{\cos^{\frac{3}{2}}(a+bx)}{\sin^{\frac{3}{2}}(a+bx)} dx$	1228
3.302	$\int \frac{\cos^{\frac{5}{2}}(a+bx)}{\sin^{\frac{5}{2}}(a+bx)} dx$	1234
3.303	$\int \frac{\cos^{\frac{7}{2}}(a+bx)}{\sin^{\frac{7}{2}}(a+bx)} dx$	1240
3.304	$\int \cos^4(e+fx) \sqrt[3]{b \sin(e+fx)} dx$	1247
3.305	$\int \cos^2(e+fx) \sqrt[3]{b \sin(e+fx)} dx$	1250
3.306	$\int \sqrt[3]{b \sin(e+fx)} dx$	1253
3.307	$\int \sec^2(e+fx) \sqrt[3]{b \sin(e+fx)} dx$	1256
3.308	$\int \sec^4(e+fx) \sqrt[3]{b \sin(e+fx)} dx$	1259
3.309	$\int \cos^4(e+fx) (b \sin(e+fx))^{5/3} dx$	1262
3.310	$\int \cos^2(e+fx) (b \sin(e+fx))^{5/3} dx$	1265
3.311	$\int (b \sin(e+fx))^{5/3} dx$	1268
3.312	$\int \sec^2(e+fx) (b \sin(e+fx))^{5/3} dx$	1271
3.313	$\int \sec^4(e+fx) (b \sin(e+fx))^{5/3} dx$	1274
3.314	$\int \frac{\cos^4(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$	1277
3.315	$\int \frac{\cos^2(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$	1280
3.316	$\int \frac{1}{\sqrt[3]{b \sin(e+fx)}} dx$	1283
3.317	$\int \frac{\sec^2(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$	1286
3.318	$\int \frac{\sec^4(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$	1289
3.319	$\int \frac{\cos^4(e+fx)}{(b \sin(e+fx))^{5/3}} dx$	1292
3.320	$\int \frac{\cos^2(e+fx)}{(b \sin(e+fx))^{5/3}} dx$	1295
3.321	$\int \frac{1}{(b \sin(e+fx))^{5/3}} dx$	1298
3.322	$\int \frac{\sec^2(e+fx)}{(b \sin(e+fx))^{5/3}} dx$	1301
3.323	$\int \frac{\sec^4(e+fx)}{(b \sin(e+fx))^{5/3}} dx$	1304
3.324	$\int \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} dx$	1307
3.325	$\int \frac{\sin^{\frac{3}{2}}(a+bx)}{\cos^{\frac{3}{2}}(a+bx)} dx$	1312
3.326	$\int \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)} dx$	1316

3.327	$\int \frac{\sin^{\frac{5}{3}}(a+bx)}{\cos^{\frac{5}{3}}(a+bx)} dx$	1321
3.328	$\int \frac{\sin^{\frac{7}{3}}(a+bx)}{\cos^{\frac{7}{3}}(a+bx)} dx$	1326
3.329	$\int \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} dx$	1331
3.330	$\int \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} dx$	1336
3.331	$\int \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} dx$	1340
3.332	$\int \frac{\cos^{\frac{5}{3}}(a+bx)}{\sin^{\frac{5}{3}}(a+bx)} dx$	1345
3.333	$\int \frac{\cos^{\frac{7}{3}}(a+bx)}{\sin^{\frac{7}{3}}(a+bx)} dx$	1350
3.334	$\int \frac{\cos^{\frac{2}{3}}(x)}{\sin^{\frac{2}{3}}(x)} dx$	1355
3.335	$\int \frac{\sin^{\frac{2}{3}}(x)}{\cos^{\frac{2}{3}}(x)} dx$	1358
3.336	$\int \cos^n(e+fx) \sin^m(e+fx) dx$	1361
3.337	$\int (d \cos(e+fx))^n \sin^m(e+fx) dx$	1364
3.338	$\int \cos^n(e+fx) (b \sin(e+fx))^m dx$	1367
3.339	$\int (d \cos(e+fx))^n (b \sin(e+fx))^m dx$	1370
3.340	$\int \cos^5(a+bx) (c \sin(a+bx))^m dx$	1373
3.341	$\int \cos^3(a+bx) (c \sin(a+bx))^m dx$	1377
3.342	$\int \cos(a+bx) (c \sin(a+bx))^m dx$	1381
3.343	$\int \sec(a+bx) (c \sin(a+bx))^m dx$	1384
3.344	$\int \sec^3(a+bx) (c \sin(a+bx))^m dx$	1387
3.345	$\int \cos^4(a+bx) (c \sin(a+bx))^m dx$	1390
3.346	$\int \cos^2(a+bx) (c \sin(a+bx))^m dx$	1393
3.347	$\int (c \sin(a+bx))^m dx$	1396
3.348	$\int \sec^2(a+bx) (c \sin(a+bx))^m dx$	1399
3.349	$\int \sec^4(a+bx) (c \sin(a+bx))^m dx$	1402
3.350	$\int (d \cos(a+bx))^{3/2} (c \sin(a+bx))^m dx$	1405
3.351	$\int \sqrt{d \cos(a+bx)} (c \sin(a+bx))^m dx$	1408
3.352	$\int \frac{(c \sin(a+bx))^m}{\sqrt{d \cos(a+bx)}} dx$	1411
3.353	$\int \frac{(c \sin(a+bx))^m}{(d \cos(a+bx))^{3/2}} dx$	1414
3.354	$\int \frac{(c \sin(a+bx))^m}{(d \cos(a+bx))^{5/2}} dx$	1417
3.355	$\int (d \cos(a+bx))^n \sin^5(a+bx) dx$	1420
3.356	$\int (d \cos(a+bx))^n \sin^3(a+bx) dx$	1425
3.357	$\int (d \cos(a+bx))^n \sin(a+bx) dx$	1429
3.358	$\int (d \cos(a+bx))^n \csc(a+bx) dx$	1432
3.359	$\int (d \cos(a+bx))^n \csc^3(a+bx) dx$	1435
3.360	$\int (d \cos(a+bx))^n \csc^5(a+bx) dx$	1438
3.361	$\int (d \cos(a+bx))^n \sin^4(a+bx) dx$	1441
3.362	$\int (d \cos(a+bx))^n \sin^2(a+bx) dx$	1444

3.363	$\int (d \cos(a + bx))^n dx$	1447
3.364	$\int (d \cos(a + bx))^n \csc^2(a + bx) dx$	1450
3.365	$\int (d \cos(a + bx))^n \csc^4(a + bx) dx$	1453
3.366	$\int (d \cos(a + bx))^n (c \sin(a + bx))^{5/2} dx$	1456
3.367	$\int (d \cos(a + bx))^n (c \sin(a + bx))^{3/2} dx$	1459
3.368	$\int (d \cos(a + bx))^n \sqrt{c \sin(a + bx)} dx$	1462
3.369	$\int \frac{(d \cos(a + bx))^n}{\sqrt{c \sin(a + bx)}} dx$	1465
3.370	$\int \frac{(d \cos(a + bx))^n}{(c \sin(a + bx))^{3/2}} dx$	1468
3.371	$\int \sqrt{b \sec(e + fx)} \sin^7(e + fx) dx$	1471
3.372	$\int \sqrt{b \sec(e + fx)} \sin^5(e + fx) dx$	1475
3.373	$\int \sqrt{b \sec(e + fx)} \sin^3(e + fx) dx$	1479
3.374	$\int \sqrt{b \sec(e + fx)} \sin(e + fx) dx$	1482
3.375	$\int \csc(e + fx) \sqrt{b \sec(e + fx)} dx$	1485
3.376	$\int \csc^3(e + fx) \sqrt{b \sec(e + fx)} dx$	1489
3.377	$\int \csc^5(e + fx) \sqrt{b \sec(e + fx)} dx$	1494
3.378	$\int \sqrt{b \sec(e + fx)} \sin^6(e + fx) dx$	1499
3.379	$\int \sqrt{b \sec(e + fx)} \sin^4(e + fx) dx$	1503
3.380	$\int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx$	1506
3.381	$\int \sqrt{b \sec(e + fx)} dx$	1509
3.382	$\int \csc^2(e + fx) \sqrt{b \sec(e + fx)} dx$	1512
3.383	$\int \csc^4(e + fx) \sqrt{b \sec(e + fx)} dx$	1515
3.384	$\int \csc^6(e + fx) \sqrt{b \sec(e + fx)} dx$	1519
3.385	$\int (b \sec(e + fx))^{3/2} \sin^7(e + fx) dx$	1523
3.386	$\int (b \sec(e + fx))^{3/2} \sin^5(e + fx) dx$	1527
3.387	$\int (b \sec(e + fx))^{3/2} \sin^3(e + fx) dx$	1531
3.388	$\int (b \sec(e + fx))^{3/2} \sin(e + fx) dx$	1535
3.389	$\int \csc(e + fx) (b \sec(e + fx))^{3/2} dx$	1538
3.390	$\int \csc^3(e + fx) (b \sec(e + fx))^{3/2} dx$	1543
3.391	$\int (b \sec(e + fx))^{3/2} \sin^6(e + fx) dx$	1548
3.392	$\int (b \sec(e + fx))^{3/2} \sin^4(e + fx) dx$	1552
3.393	$\int (b \sec(e + fx))^{3/2} \sin^2(e + fx) dx$	1556
3.394	$\int (b \sec(e + fx))^{3/2} dx$	1559
3.395	$\int \csc^2(e + fx) (b \sec(e + fx))^{3/2} dx$	1562
3.396	$\int \csc^4(e + fx) (b \sec(e + fx))^{3/2} dx$	1566
3.397	$\int (b \sec(e + fx))^{5/2} \sin^7(e + fx) dx$	1570
3.398	$\int (b \sec(e + fx))^{5/2} \sin^5(e + fx) dx$	1574
3.399	$\int (b \sec(e + fx))^{5/2} \sin^3(e + fx) dx$	1578
3.400	$\int (b \sec(e + fx))^{5/2} \sin(e + fx) dx$	1581
3.401	$\int \csc(e + fx) (b \sec(e + fx))^{5/2} dx$	1584
3.402	$\int \csc^3(e + fx) (b \sec(e + fx))^{5/2} dx$	1589
3.403	$\int \csc^5(e + fx) (b \sec(e + fx))^{5/2} dx$	1595
3.404	$\int (b \sec(e + fx))^{5/2} \sin^6(e + fx) dx$	1601

3.405	$\int (b \sec(e + fx))^{5/2} \sin^4(e + fx) dx$	1605
3.406	$\int (b \sec(e + fx))^{5/2} \sin^2(e + fx) dx$	1609
3.407	$\int (b \sec(e + fx))^{5/2} dx$	1612
3.408	$\int \csc^2(e + fx)(b \sec(e + fx))^{5/2} dx$	1615
3.409	$\int \csc^4(e + fx)(b \sec(e + fx))^{5/2} dx$	1619
3.410	$\int \frac{\sin^7(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1623
3.411	$\int \frac{\sin^5(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1626
3.412	$\int \frac{\sin^3(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1630
3.413	$\int \frac{\sin(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1633
3.414	$\int \frac{\csc(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1636
3.415	$\int \frac{\csc^3(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1640
3.416	$\int \frac{\csc^5(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1645
3.417	$\int \frac{\sin^6(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1651
3.418	$\int \frac{\sin^4(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1655
3.419	$\int \frac{\sin^2(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1659
3.420	$\int \frac{1}{\sqrt{b \sec(e + fx)}} dx$	1662
3.421	$\int \frac{\csc^2(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1665
3.422	$\int \frac{\csc^4(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1669
3.423	$\int \frac{\csc^6(e+fx)}{\sqrt{b \sec(e + fx)}} dx$	1673
3.424	$\int \frac{\sin^7(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1677
3.425	$\int \frac{\sin^5(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1680
3.426	$\int \frac{\sin^3(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1683
3.427	$\int \frac{\sin(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1686
3.428	$\int \frac{\csc(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1689
3.429	$\int \frac{\csc^3(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1694
3.430	$\int \frac{\csc^5(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1699
3.431	$\int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1705
3.432	$\int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1709
3.433	$\int \frac{1}{(b \sec(e+fx))^{3/2}} dx$	1713
3.434	$\int \frac{\csc^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1716

3.435	$\int \frac{\csc^4(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1719
3.436	$\int \frac{\csc^6(e+fx)}{(b \sec(e+fx))^{3/2}} dx$	1723
3.437	$\int \frac{\sin^7(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1727
3.438	$\int \frac{\sin^5(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1730
3.439	$\int \frac{\sin^3(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1733
3.440	$\int \frac{\sin(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1736
3.441	$\int \frac{\csc(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1739
3.442	$\int \frac{\csc^3(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1744
3.443	$\int \frac{\csc^5(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1749
3.444	$\int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1755
3.445	$\int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1759
3.446	$\int \frac{1}{(b \sec(e+fx))^{5/2}} dx$	1763
3.447	$\int \frac{\csc^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1767
3.448	$\int \frac{\csc^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1771
3.449	$\int \frac{\csc^6(e+fx)}{(b \sec(e+fx))^{5/2}} dx$	1775
3.450	$\int \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{9/2} dx$	1779
3.451	$\int \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{5/2} dx$	1784
3.452	$\int \sqrt{b \sec(e+fx)} \sqrt{a \sin(e+fx)} dx$	1789
3.453	$\int \frac{\sqrt{b \sec(e+fx)}}{(a \sin(e+fx))^{3/2}} dx$	1794
3.454	$\int \frac{\sqrt{b \sec(e+fx)}}{(a \sin(e+fx))^{7/2}} dx$	1797
3.455	$\int \frac{\sqrt{b \sec(e+fx)}}{(a \sin(e+fx))^{11/2}} dx$	1800
3.456	$\int \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{7/2} dx$	1804
3.457	$\int \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{3/2} dx$	1808
3.458	$\int \frac{\sqrt{b \sec(e+fx)}}{\sqrt{a \sin(e+fx)}} dx$	1812
3.459	$\int \frac{\sqrt{b \sec(e+fx)}}{(a \sin(e+fx))^{5/2}} dx$	1815
3.460	$\int \frac{\sqrt{b \sec(e+fx)}}{(a \sin(e+fx))^{9/2}} dx$	1819
3.461	$\int \frac{\sin^{9/2}(e+fx)}{\sqrt{b \sec(e+fx)}} dx$	1823
3.462	$\int \frac{\sin^{5/2}(e+fx)}{\sqrt{b \sec(e+fx)}} dx$	1827
3.463	$\int \frac{\sqrt{\sin(e+fx)}}{\sqrt{b \sec(e+fx)}} dx$	1831
3.464	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{3/2}(e+fx)} dx$	1835

3.465	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{7}{2}}(e+fx)} dx$	1839
3.466	$\int \frac{\sin^{\frac{3}{2}}(e+fx)}{\sqrt{b \sec(e+fx)}} dx$	1843
3.467	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sqrt{\sin(e+fx)}} dx$	1848
3.468	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{5}{2}}(e+fx)} dx$	1853
3.469	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{9}{2}}(e+fx)} dx$	1856
3.470	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{13}{2}}(e+fx)} dx$	1859
3.471	$\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{17}{2}}(e+fx)} dx$	1862
3.472	$\int \frac{(a \sin(e+fx))^{9/2}}{(b \sec(e+fx))^{3/2}} dx$	1866
3.473	$\int \frac{(a \sin(e+fx))^{5/2}}{(b \sec(e+fx))^{3/2}} dx$	1872
3.474	$\int \frac{\sqrt{a \sin(e+fx)}}{(b \sec(e+fx))^{3/2}} dx$	1878
3.475	$\int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{3/2}} dx$	1883
3.476	$\int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{7/2}} dx$	1889
3.477	$\int \frac{(a \sin(e+fx))^{7/2}}{(b \sec(e+fx))^{3/2}} dx$	1892
3.478	$\int \frac{(a \sin(e+fx))^{3/2}}{(b \sec(e+fx))^{3/2}} dx$	1896
3.479	$\int \frac{1}{(b \sec(e+fx))^{3/2} \sqrt{a \sin(e+fx)}} dx$	1900
3.480	$\int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{5/2}} dx$	1904
3.481	$\int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{9/2}} dx$	1908
3.482	$\int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{13/2}} dx$	1912
3.483	$\int (d \sec(a+bx))^{5/2} (c \sin(a+bx))^m dx$	1917
3.484	$\int (d \sec(a+bx))^{3/2} (c \sin(a+bx))^m dx$	1920
3.485	$\int \sqrt{d \sec(a+bx)} (c \sin(a+bx))^m dx$	1923
3.486	$\int \frac{(c \sin(a+bx))^m}{\sqrt{d \sec(a+bx)}} dx$	1926
3.487	$\int \frac{(c \sin(a+bx))^m}{(d \sec(a+bx))^{3/2}} dx$	1929
3.488	$\int \sec^n(e+fx) \sin^m(e+fx) dx$	1932
3.489	$\int \sec^n(e+fx) (a \sin(e+fx))^m dx$	1935
3.490	$\int (b \sec(e+fx))^n \sin^m(e+fx) dx$	1938
3.491	$\int (b \sec(e+fx))^n (a \sin(e+fx))^m dx$	1941
3.492	$\int (b \sec(e+fx))^n \sin^5(e+fx) dx$	1944
3.493	$\int (b \sec(e+fx))^n \sin^3(e+fx) dx$	1947
3.494	$\int (b \sec(e+fx))^n \sin(e+fx) dx$	1951
3.495	$\int \csc(e+fx) (b \sec(e+fx))^n dx$	1954
3.496	$\int \csc^3(e+fx) (b \sec(e+fx))^n dx$	1957
3.497	$\int (b \sec(e+fx))^n \sin^6(e+fx) dx$	1960
3.498	$\int (b \sec(e+fx))^n \sin^4(e+fx) dx$	1963

3.499	$\int (b \sec(e + fx))^n \sin^2(e + fx) dx$	1966
3.500	$\int (b \sec(e + fx))^n dx$	1970
3.501	$\int \csc^2(e + fx)(b \sec(e + fx))^n dx$	1973
3.502	$\int \csc^4(e + fx)(b \sec(e + fx))^n dx$	1977
3.503	$\int (b \sec(a + bx))^n (c \sin(a + bx))^{3/2} dx$	1981
3.504	$\int (b \sec(a + bx))^n \sqrt{c \sin(a + bx)} dx$	1984
3.505	$\int \frac{(b \sec(a + bx))^n}{\sqrt{c \sin(a + bx)}} dx$	1987
3.506	$\int \frac{(b \sec(a + bx))^n}{(c \sin(a + bx))^{3/2}} dx$	1990
3.507	$\int \sqrt{d \csc(e + fx)} \sin^4(e + fx) dx$	1993
3.508	$\int \sqrt{d \csc(e + fx)} \sin^3(e + fx) dx$	1997
3.509	$\int \sqrt{d \csc(e + fx)} \sin^2(e + fx) dx$	2001
3.510	$\int \sqrt{d \csc(e + fx)} \sin(e + fx) dx$	2005
3.511	$\int \sqrt{d \csc(e + fx)} dx$	2009
3.512	$\int \csc(e + fx) \sqrt{d \csc(e + fx)} dx$	2012
3.513	$\int \csc^2(e + fx) \sqrt{d \csc(e + fx)} dx$	2016
3.514	$\int \csc^3(e + fx) \sqrt{d \csc(e + fx)} dx$	2020
3.515	$\int (d \csc(e + fx))^{3/2} \sin^5(e + fx) dx$	2024
3.516	$\int (d \csc(e + fx))^{3/2} \sin^4(e + fx) dx$	2028
3.517	$\int (d \csc(e + fx))^{3/2} \sin^3(e + fx) dx$	2032
3.518	$\int (d \csc(e + fx))^{3/2} \sin^2(e + fx) dx$	2035
3.519	$\int (d \csc(e + fx))^{3/2} \sin(e + fx) dx$	2039
3.520	$\int (d \csc(e + fx))^{3/2} dx$	2042
3.521	$\int \csc(e + fx)(d \csc(e + fx))^{3/2} dx$	2046
3.522	$\int \csc^2(e + fx)(d \csc(e + fx))^{3/2} dx$	2050
3.523	$\int \frac{\sin^3(e + fx)}{\sqrt{d \csc(e + fx)}} dx$	2054
3.524	$\int \frac{\sin^2(e + fx)}{\sqrt{d \csc(e + fx)}} dx$	2058
3.525	$\int \frac{\sin(e + fx)}{\sqrt{d \csc(e + fx)}} dx$	2062
3.526	$\int \frac{1}{\sqrt{d \csc(e + fx)}} dx$	2066
3.527	$\int \frac{\csc(e + fx)}{\sqrt{d \csc(e + fx)}} dx$	2069
3.528	$\int \frac{\csc^2(e + fx)}{\sqrt{d \csc(e + fx)}} dx$	2072
3.529	$\int \frac{\csc^3(e + fx)}{\sqrt{d \csc(e + fx)}} dx$	2076
3.530	$\int \frac{\sin^2(e + fx)}{(d \csc(e + fx))^{3/2}} dx$	2080
3.531	$\int \frac{\sin(e + fx)}{(d \csc(e + fx))^{3/2}} dx$	2084
3.532	$\int \frac{1}{(d \csc(e + fx))^{3/2}} dx$	2088
3.533	$\int \frac{\csc(e + fx)}{(d \csc(e + fx))^{3/2}} dx$	2091
3.534	$\int \frac{\csc^2(e + fx)}{(d \csc(e + fx))^{3/2}} dx$	2095

3.535	$\int \frac{\csc^3(e+fx)}{(d \csc(e+fx))^{3/2}} dx$	2098
3.536	$\int \frac{\csc^4(e+fx)}{(d \csc(e+fx))^{3/2}} dx$	2102
3.537	$\int \frac{\csc^5(e+fx)}{(d \csc(e+fx))^{3/2}} dx$	2106
3.538	$\int (b \csc(e+fx))^n (a \sin(e+fx))^m dx$	2110

3.1 $\int \sin(a + bx) dx$

Optimal. Leaf size=11

$$-\frac{\cos(a + bx)}{b}$$

[Out] $-\cos(b*x+a)/b$

Rubi [A]

time = 0.00, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2718}

$$-\frac{\cos(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x],x]

[Out] $-(\text{Cos}[a + b*x])/b$

Rule 2718

Int[sin[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Cos[c + d*x]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \sin(a + bx) dx = -\frac{\cos(a + bx)}{b}$$

Mathematica [A]

time = 0.01, size = 22, normalized size = 2.00

$$-\frac{\cos(a) \cos(bx)}{b} + \frac{\sin(a) \sin(bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x],x]

[Out] $-\left(\text{Cos}[a] * \text{Cos}[b*x]\right)/b + \left(\text{Sin}[a] * \text{Sin}[b*x]\right)/b$

Maple [A]

time = 0.07, size = 12, normalized size = 1.09

method	result	size
derivativedivides	$-\frac{\cos(bx+a)}{b}$	12
default	$-\frac{\cos(bx+a)}{b}$	12
risch	$-\frac{\cos(bx+a)}{b}$	12
norman	$\frac{2\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}$	32
meijerg	$\frac{\sin(a)\sin(bx)}{b} + \frac{\cos(a)\sqrt{\pi}\left(\frac{1}{\sqrt{\pi}} - \frac{\cos(bx)}{\sqrt{\pi}}\right)}{b}$	34

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a),x,method=_RETURNVERBOSE)`

[Out] $-\cos(b*x+a)/b$

Maxima [A]

time = 0.33, size = 11, normalized size = 1.00

$$-\frac{\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a),x, algorithm="maxima")`

[Out] $-\cos(b*x + a)/b$

Fricas [A]

time = 0.38, size = 11, normalized size = 1.00

$$-\frac{\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a),x, algorithm="fricas")`

[Out] $-\cos(b*x + a)/b$

Sympy [A]

time = 0.04, size = 14, normalized size = 1.27

$$\begin{cases} -\frac{\cos(a+bx)}{b} & \text{for } b \neq 0 \\ x \sin(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a),x)`

[Out] `Piecewise((-cos(a + b*x)/b, Ne(b, 0)), (x*sin(a), True))`

Giac [A]

time = 4.76, size = 11, normalized size = 1.00

$$-\frac{\cos(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a),x, algorithm="giac")`

[Out] `-cos(b*x + a)/b`

Mupad [B]

time = 0.02, size = 11, normalized size = 1.00

$$-\frac{\cos(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x),x)`

[Out] `-cos(a + b*x)/b`

3.2 $\int \sin^2(a + bx) dx$

Optimal. Leaf size=25

$$\frac{x}{2} - \frac{\cos(a + bx) \sin(a + bx)}{2b}$$

[Out] 1/2*x-1/2*cos(b*x+a)*sin(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$\frac{x}{2} - \frac{\sin(a + bx) \cos(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2,x]

[Out] x/2 - (Cos[a + b*x]*Sin[a + b*x])/(2*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sin^2(a + bx) dx &= -\frac{\cos(a + bx) \sin(a + bx)}{2b} + \frac{\int 1 dx}{2} \\ &= \frac{x}{2} - \frac{\cos(a + bx) \sin(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 23, normalized size = 0.92

$$-\frac{-2(a + bx) + \sin(2(a + bx))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^2,x]

[Out] $-1/4*(-2*(a + b*x) + \text{Sin}[2*(a + b*x)])/b$

Maple [A]

time = 0.04, size = 27, normalized size = 1.08

method	result	size
risch	$\frac{x}{2} - \frac{\sin(2bx+2a)}{4b}$	19
derivativedivides	$-\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx+a}{2}$ b	27
default	$-\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx+a}{2}$ b	27
norman	$\frac{\tan^3\left(\frac{bx+a}{2}\right)}{b} + x\left(\tan^2\left(\frac{bx+a}{2}\right)\right) + \frac{x}{2} - \frac{\tan\left(\frac{bx+a}{2}\right)}{b} + \frac{x\left(\tan^4\left(\frac{bx+a}{2}\right)\right)}{2}$ $(1+\tan^2\left(\frac{bx+a}{2}\right))^2$	77

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] $1/b*(-1/2*\cos(b*x+a)*\sin(b*x+a)+1/2*b*x+1/2*a)$

Maxima [A]

time = 0.29, size = 24, normalized size = 0.96

$$\frac{2bx + 2a - \sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2,x, algorithm="maxima")

[Out] $1/4*(2*b*x + 2*a - \sin(2*b*x + 2*a))/b$

Fricas [A]

time = 0.41, size = 23, normalized size = 0.92

$$\frac{bx - \cos(bx + a)\sin(bx + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2,x, algorithm="fricas")

[Out] $1/2*(b*x - \cos(b*x + a)*\sin(b*x + a))/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(19) = 38$.

time = 0.07, size = 46, normalized size = 1.84

$$\begin{cases} \frac{x \sin^2(a+bx)}{2} + \frac{x \cos^2(a+bx)}{2} - \frac{\sin(a+bx)\cos(a+bx)}{2b} & \text{for } b \neq 0 \\ x \sin^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**2,x)

[Out] Piecewise((x*sin(a + b*x)**2/2 + x*cos(a + b*x)**2/2 - sin(a + b*x)*cos(a + b*x)/(2*b), Ne(b, 0)), (x*sin(a)**2, True))

Giac [A]

time = 5.38, size = 18, normalized size = 0.72

$$\frac{1}{2}x - \frac{\sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/2*x - 1/4*sin(2*b*x + 2*a)/b

Mupad [B]

time = 0.43, size = 18, normalized size = 0.72

$$\frac{x}{2} - \frac{\sin(2a + 2bx)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2,x)

[Out] x/2 - sin(2*a + 2*b*x)/(4*b)

3.3 $\int \sin^3(a + bx) dx$

Optimal. Leaf size=27

$$-\frac{\cos(a + bx)}{b} + \frac{\cos^3(a + bx)}{3b}$$

[Out] $-\cos(b*x+a)/b+1/3*\cos(b*x+a)^3/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2713}

$$\frac{\cos^3(a + bx)}{3b} - \frac{\cos(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[a + b*x]^3, x]$

[Out] $-(\text{Cos}[a + b*x]/b) + \text{Cos}[a + b*x]^3/(3*b)$

Rule 2713

$\text{Int}[\sin[(c_.) + (d_.)*(x_)]^{(n_.)}, x_Symbol] \text{ :> } \text{Dist}[-d^{(-1)}, \text{Subst}[\text{Int}[\text{Expand}[(1 - x^2)^{((n - 1)/2)}, x], x], x, \text{Cos}[c + d*x]], x] \text{ /; } \text{FreeQ}\{c, d\}, x \text{ \&\& } \text{IGtQ}[(n - 1)/2, 0]$

Rubi steps

$$\begin{aligned} \int \sin^3(a + bx) dx &= -\frac{\text{Subst}(\int (1 - x^2) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\cos(a + bx)}{b} + \frac{\cos^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 29, normalized size = 1.07

$$-\frac{3 \cos(a + bx)}{4b} + \frac{\cos(3(a + bx))}{12b}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Sin}[a + b*x]^3, x]$

[Out] $(-3*\text{Cos}[a + b*x])/(4*b) + \text{Cos}[3*(a + b*x)]/(12*b)$

Maple [A]

time = 0.09, size = 22, normalized size = 0.81

method	result	size
derivativedivides	$-\frac{(2+\sin^2(bx+a))\cos(bx+a)}{3b}$	22
default	$-\frac{(2+\sin^2(bx+a))\cos(bx+a)}{3b}$	22
risch	$-\frac{3\cos(bx+a)}{4b} + \frac{\cos(3bx+3a)}{12b}$	27
norman	$-\frac{4\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{4}{3b}$ $\frac{1}{\left(1+\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3}$	39

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
[Out] -1/3/b*(2+sin(b*x+a)^2)*cos(b*x+a)
```

Maxima [A]

time = 0.28, size = 22, normalized size = 0.81

$$\frac{\cos(bx+a)^3 - 3\cos(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^3,x, algorithm="maxima")
```

```
[Out] 1/3*(cos(b*x + a)^3 - 3*cos(b*x + a))/b
```

Fricas [A]

time = 0.39, size = 22, normalized size = 0.81

$$\frac{\cos(bx+a)^3 - 3\cos(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^3,x, algorithm="fricas")
```

```
[Out] 1/3*(cos(b*x + a)^3 - 3*cos(b*x + a))/b
```

Sympy [A]

time = 0.10, size = 37, normalized size = 1.37

$$\begin{cases} -\frac{\sin^2(a+bx)\cos(a+bx)}{b} - \frac{2\cos^3(a+bx)}{3b} & \text{for } b \neq 0 \\ x \sin^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**3,x)

[Out] Piecewise((-sin(a + b*x)**2*cos(a + b*x)/b - 2*cos(a + b*x)**3/(3*b), Ne(b, 0)), (x*sin(a)**3, True))

Giac [A]

time = 5.55, size = 25, normalized size = 0.93

$$\frac{\cos (bx+a)^3}{3 b}-\frac{\cos (bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/3*cos(b*x + a)^3/b - cos(b*x + a)/b

Mupad [B]

time = 0.35, size = 24, normalized size = 0.89

$$-\frac{3 \cos (a+b x)-\cos (a+b x)^3}{3 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3,x)

[Out] -(3*cos(a + b*x) - cos(a + b*x)^3)/(3*b)

3.4 $\int \sin^4(a + bx) dx$

Optimal. Leaf size=46

$$\frac{3x}{8} - \frac{3 \cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos(a + bx) \sin^3(a + bx)}{4b}$$

[Out] $3/8*x - 3/8*\cos(b*x+a)*\sin(b*x+a)/b - 1/4*\cos(b*x+a)*\sin(b*x+a)^3/b$

Rubi [A]

time = 0.01, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$-\frac{\sin^3(a + bx) \cos(a + bx)}{4b} - \frac{3 \sin(a + bx) \cos(a + bx)}{8b} + \frac{3x}{8}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^4,x]

[Out] $(3*x)/8 - (3*\text{Cos}[a + b*x]*\text{Sin}[a + b*x])/(8*b) - (\text{Cos}[a + b*x]*\text{Sin}[a + b*x]^3)/(4*b)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sin^4(a + bx) dx &= -\frac{\cos(a + bx) \sin^3(a + bx)}{4b} + \frac{3}{4} \int \sin^2(a + bx) dx \\ &= -\frac{3 \cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos(a + bx) \sin^3(a + bx)}{4b} + \frac{3 \int 1 dx}{8} \\ &= \frac{3x}{8} - \frac{3 \cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos(a + bx) \sin^3(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 33, normalized size = 0.72

$$\frac{12(a + bx) - 8 \sin(2(a + bx)) + \sin(4(a + bx))}{32b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^4,x]**[Out]** (12*(a + b*x) - 8*Sin[2*(a + b*x)] + Sin[4*(a + b*x)])/(32*b)**Maple [A]**

time = 0.09, size = 38, normalized size = 0.83

method	result
risch	$\frac{3x}{8} + \frac{\sin(4bx+4a)}{32b} - \frac{\sin(2bx+2a)}{4b}$
derivativedivides	$-\frac{\left(\sin^3(bx+a) + \frac{3\sin(bx+a)}{2}\right) \cos(bx+a)}{4b} + \frac{3bx}{8} + \frac{3a}{8}$
default	$-\frac{\left(\sin^3(bx+a) + \frac{3\sin(bx+a)}{2}\right) \cos(bx+a)}{4b} + \frac{3bx}{8} + \frac{3a}{8}$
norman	$\frac{\frac{3x}{8} - \frac{3 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{4b} - \frac{11 \left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{11 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{3 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{3x \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2} + \frac{9x \left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4} + \frac{3x}{4}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^4,x,method=_RETURNVERBOSE)**[Out]** 1/b*(-1/4*(sin(b*x+a)^3+3/2*sin(b*x+a))*cos(b*x+a)+3/8*b*x+3/8*a)**Maxima [A]**

time = 0.29, size = 33, normalized size = 0.72

$$\frac{12bx + 12a + \sin(4bx + 4a) - 8 \sin(2bx + 2a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4,x, algorithm="maxima")**[Out]** 1/32*(12*b*x + 12*a + sin(4*b*x + 4*a) - 8*sin(2*b*x + 2*a))/b**Fricas [A]**

time = 0.39, size = 36, normalized size = 0.78

$$\frac{3bx + (2 \cos(bx + a))^3 - 5 \cos(bx + a) \sin(bx + a)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/8*(3*b*x + (2*cos(b*x + a)^3 - 5*cos(b*x + a))*sin(b*x + a))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(41) = 82$.

time = 0.17, size = 95, normalized size = 2.07

$$\begin{cases} \frac{3x \sin^4(a+bx)}{8} + \frac{3x \sin^2(a+bx) \cos^2(a+bx)}{4} + \frac{3x \cos^4(a+bx)}{8} - \frac{5 \sin^3(a+bx) \cos(a+bx)}{8b} - \frac{3 \sin(a+bx) \cos^3(a+bx)}{8b} & \text{for } b \neq 0 \\ x \sin^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**4,x)

[Out] Piecewise(((3*x*sin(a + b*x)**4/8 + 3*x*sin(a + b*x)**2*cos(a + b*x)**2/4 + 3*x*cos(a + b*x)**4/8 - 5*sin(a + b*x)**3*cos(a + b*x)/(8*b) - 3*sin(a + b*x)*cos(a + b*x)**3/(8*b), Ne(b, 0)), (x*sin(a)**4, True))

Giac [A]

time = 4.85, size = 32, normalized size = 0.70

$$\frac{3}{8}x + \frac{\sin(4bx + 4a)}{32b} - \frac{\sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4,x, algorithm="giac")

[Out] 3/8*x + 1/32*sin(4*b*x + 4*a)/b - 1/4*sin(2*b*x + 2*a)/b

Mupad [B]

time = 0.47, size = 50, normalized size = 1.09

$$\frac{3x}{8} - \frac{\frac{5 \tan(a+bx)^3}{8} + \frac{3 \tan(a+bx)}{8}}{b (\tan(a+bx)^4 + 2 \tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4,x)

[Out] (3*x)/8 - ((3*tan(a + b*x))/8 + (5*tan(a + b*x)^3)/8)/(b*(2*tan(a + b*x)^2 + tan(a + b*x)^4 + 1))

3.5 $\int \sin^5(a + bx) dx$

Optimal. Leaf size=42

$$-\frac{\cos(a + bx)}{b} + \frac{2 \cos^3(a + bx)}{3b} - \frac{\cos^5(a + bx)}{5b}$$

[Out] $-\cos(b*x+a)/b+2/3*\cos(b*x+a)^3/b-1/5*\cos(b*x+a)^5/b$

Rubi [A]

time = 0.01, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2713}

$$-\frac{\cos^5(a + bx)}{5b} + \frac{2 \cos^3(a + bx)}{3b} - \frac{\cos(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^5,x]

[Out] $-(\text{Cos}[a + b*x]/b) + (2*\text{Cos}[a + b*x]^3)/(3*b) - \text{Cos}[a + b*x]^5/(5*b)$

Rule 2713

Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] :> Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]

Rubi steps

$$\begin{aligned} \int \sin^5(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 - 2x^2 + x^4) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos(a + bx)}{b} + \frac{2 \cos^3(a + bx)}{3b} - \frac{\cos^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 44, normalized size = 1.05

$$-\frac{5 \cos(a + bx)}{8b} + \frac{5 \cos(3(a + bx))}{48b} - \frac{\cos(5(a + bx))}{80b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^5,x]

[Out] $(-5*\text{Cos}[a + b*x])/(8*b) + (5*\text{Cos}[3*(a + b*x)])/(48*b) - \text{Cos}[5*(a + b*x)]/(80*b)$

Maple [A]

time = 0.10, size = 32, normalized size = 0.76

method	result	size
derivativedivides	$-\frac{\left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{5b}$	32
default	$-\frac{\left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{5b}$	32
risch	$-\frac{5 \cos(bx+a)}{8b} - \frac{\cos(5bx+5a)}{80b} + \frac{5 \cos(3bx+3a)}{48b}$	41
norman	$\frac{-\frac{16}{15b} - \frac{16(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{32(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^5}$	55

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

[Out] $-1/5/b*(8/3+\sin(b*x+a)^4+4/3*\sin(b*x+a)^2)*\cos(b*x+a)$

Maxima [A]

time = 0.29, size = 34, normalized size = 0.81

$$\frac{3 \cos(bx+a)^5 - 10 \cos(bx+a)^3 + 15 \cos(bx+a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^5,x, algorithm="maxima")`

[Out] $-1/15*(3*\cos(b*x + a)^5 - 10*\cos(b*x + a)^3 + 15*\cos(b*x + a))/b$

Fricas [A]

time = 0.39, size = 34, normalized size = 0.81

$$\frac{3 \cos(bx+a)^5 - 10 \cos(bx+a)^3 + 15 \cos(bx+a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $-1/15*(3*\cos(b*x + a)^5 - 10*\cos(b*x + a)^3 + 15*\cos(b*x + a))/b$

Sympy [A]

time = 0.25, size = 60, normalized size = 1.43

$$\begin{cases} -\frac{\sin^4(a+bx) \cos(a+bx)}{b} - \frac{4 \sin^2(a+bx) \cos^3(a+bx)}{3b} - \frac{8 \cos^5(a+bx)}{15b} & \text{for } b \neq 0 \\ x \sin^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**5,x)

[Out] Piecewise((-sin(a + b*x)**4*cos(a + b*x)/b - 4*sin(a + b*x)**2*cos(a + b*x)**3/(3*b) - 8*cos(a + b*x)**5/(15*b), Ne(b, 0)), (x*sin(a)**5, True))

Giac [A]

time = 5.25, size = 38, normalized size = 0.90

$$-\frac{\cos(bx+a)^5}{5b} + \frac{2\cos(bx+a)^3}{3b} - \frac{\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^5,x, algorithm="giac")

[Out] -1/5*cos(b*x + a)^5/b + 2/3*cos(b*x + a)^3/b - cos(b*x + a)/b

Mupad [B]

time = 0.36, size = 32, normalized size = 0.76

$$-\frac{\frac{\cos(a+bx)^5}{5} - \frac{2\cos(a+bx)^3}{3} + \cos(a+bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^5,x)

[Out] -(cos(a + b*x) - (2*cos(a + b*x)^3)/3 + cos(a + b*x)^5/5)/b

3.6 $\int \sin^6(a + bx) dx$

Optimal. Leaf size=67

$$\frac{5x}{16} - \frac{5 \cos(a + bx) \sin(a + bx)}{16b} - \frac{5 \cos(a + bx) \sin^3(a + bx)}{24b} - \frac{\cos(a + bx) \sin^5(a + bx)}{6b}$$

[Out] 5/16*x-5/16*cos(b*x+a)*sin(b*x+a)/b-5/24*cos(b*x+a)*sin(b*x+a)^3/b-1/6*cos(b*x+a)*sin(b*x+a)^5/b

Rubi [A]

time = 0.02, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$-\frac{\sin^5(a + bx) \cos(a + bx)}{6b} - \frac{5 \sin^3(a + bx) \cos(a + bx)}{24b} - \frac{5 \sin(a + bx) \cos(a + bx)}{16b} + \frac{5x}{16}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^6,x]

[Out] (5*x)/16 - (5*Cos[a + b*x]*Sin[a + b*x])/(16*b) - (5*Cos[a + b*x]*Sin[a + b*x]^3)/(24*b) - (Cos[a + b*x]*Sin[a + b*x]^5)/(6*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sine[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sin^6(a + bx) dx &= -\frac{\cos(a + bx) \sin^5(a + bx)}{6b} + \frac{5}{6} \int \sin^4(a + bx) dx \\ &= -\frac{5 \cos(a + bx) \sin^3(a + bx)}{24b} - \frac{\cos(a + bx) \sin^5(a + bx)}{6b} + \frac{5}{8} \int \sin^2(a + bx) dx \\ &= -\frac{5 \cos(a + bx) \sin(a + bx)}{16b} - \frac{5 \cos(a + bx) \sin^3(a + bx)}{24b} - \frac{\cos(a + bx) \sin^5(a + bx)}{6b} + \frac{5x}{16} \\ &= \frac{5x}{16} - \frac{5 \cos(a + bx) \sin(a + bx)}{16b} - \frac{5 \cos(a + bx) \sin^3(a + bx)}{24b} - \frac{\cos(a + bx) \sin^5(a + bx)}{6b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 45, normalized size = 0.67

$$\frac{60a + 60bx - 45 \sin(2(a + bx)) + 9 \sin(4(a + bx)) - \sin(6(a + bx))}{192b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^6,x]`

```
[Out] (60*a + 60*b*x - 45*Sin[2*(a + b*x)] + 9*Sin[4*(a + b*x)] - Sin[6*(a + b*x)])/(192*b)
```

Maple [A]

time = 0.11, size = 48, normalized size = 0.72

method	result
risch	$\frac{5x}{16} - \frac{\sin(6bx+6a)}{192b} + \frac{3 \sin(4bx+4a)}{64b} - \frac{15 \sin(2bx+2a)}{64b}$
derivativedivides	$-\frac{\left(\sin^5(bx+a) + \frac{5(\sin^3(bx+a))}{4} + \frac{15 \sin(bx+a)}{8}\right) \cos(bx+a)}{6b} + \frac{5bx}{16} + \frac{5a}{16}$
default	$-\frac{\left(\sin^5(bx+a) + \frac{5(\sin^3(bx+a))}{4} + \frac{15 \sin(bx+a)}{8}\right) \cos(bx+a)}{6b} + \frac{5bx}{16} + \frac{5a}{16}$
norman	$\frac{5x}{16} - \frac{5 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{85 \left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} - \frac{33 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{33 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{85 \left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} + \frac{5 \left(\tan^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} + \frac{15}{(1+\tan^2)}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^6,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/6*(sin(b*x+a)^5+5/4*sin(b*x+a)^3+15/8*sin(b*x+a))*cos(b*x+a)+5/16*b*x+5/16*a)
```

Maxima [A]

time = 0.27, size = 48, normalized size = 0.72

$$\frac{4 \sin(2bx + 2a)^3 + 60bx + 60a + 9 \sin(4bx + 4a) - 48 \sin(2bx + 2a)}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^6,x, algorithm="maxima")`

```
[Out] 1/192*(4*sin(2*b*x + 2*a)^3 + 60*b*x + 60*a + 9*sin(4*b*x + 4*a) - 48*sin(2*b*x + 2*a))/b
```

Fricas [A]

time = 0.39, size = 47, normalized size = 0.70

$$\frac{15bx - (8 \cos(bx + a))^5 - 26 \cos(bx + a)^3 + 33 \cos(bx + a) \sin(bx + a)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^6,x, algorithm="fricas")

[Out] 1/48*(15*b*x - (8*cos(b*x + a)^5 - 26*cos(b*x + a)^3 + 33*cos(b*x + a))*sin(b*x + a))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 139 vs. 2(61) = 122.

time = 0.40, size = 139, normalized size = 2.07

$$\begin{cases} \frac{5x \sin^6(a+bx)}{16} + \frac{15x \sin^4(a+bx) \cos^2(a+bx)}{16} + \frac{15x \sin^2(a+bx) \cos^4(a+bx)}{16} + \frac{5x \cos^6(a+bx)}{16} - \frac{11 \sin^5(a+bx) \cos(a+bx)}{16b} - \frac{5 \sin^3(a+bx) \cos^3(a+bx)}{6b} - \frac{5 \sin(a+bx) \cos^5(a+bx)}{16b} & \text{for } b \neq 0 \\ x \sin^6(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**6,x)

[Out] Piecewise(((5*x*sin(a + b*x)**6/16 + 15*x*sin(a + b*x)**4*cos(a + b*x)**2/16 + 15*x*sin(a + b*x)**2*cos(a + b*x)**4/16 + 5*x*cos(a + b*x)**6/16 - 11*sin(a + b*x)**5*cos(a + b*x)/(16*b) - 5*sin(a + b*x)**3*cos(a + b*x)**3/(6*b) - 5*sin(a + b*x)*cos(a + b*x)**5/(16*b), Ne(b, 0)), (x*sin(a)**6, True))

Giac [A]

time = 3.91, size = 46, normalized size = 0.69

$$\frac{5}{16}x - \frac{\sin(6bx + 6a)}{192b} + \frac{3 \sin(4bx + 4a)}{64b} - \frac{15 \sin(2bx + 2a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^6,x, algorithm="giac")

[Out] 5/16*x - 1/192*sin(6*b*x + 6*a)/b + 3/64*sin(4*b*x + 4*a)/b - 15/64*sin(2*b*x + 2*a)/b

Mupad [B]

time = 0.58, size = 43, normalized size = 0.64

$$\frac{5x}{16} - \frac{\frac{15 \sin(2a+2bx)}{64} - \frac{3 \sin(4a+4bx)}{64} + \frac{\sin(6a+6bx)}{192}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^6,x)

[Out] (5*x)/16 - ((15*sin(2*a + 2*b*x))/64 - (3*sin(4*a + 4*b*x))/64 + sin(6*a + 6*b*x)/192)/b

3.7 $\int \sin^7(a + bx) dx$

Optimal. Leaf size=54

$$-\frac{\cos(a + bx)}{b} + \frac{\cos^3(a + bx)}{b} - \frac{3 \cos^5(a + bx)}{5b} + \frac{\cos^7(a + bx)}{7b}$$

[Out] $-\cos(b*x+a)/b+\cos(b*x+a)^3/b-3/5*\cos(b*x+a)^5/b+1/7*\cos(b*x+a)^7/b$

Rubi [A]

time = 0.01, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$,

Rules used = {2713}

$$\frac{\cos^7(a + bx)}{7b} - \frac{3 \cos^5(a + bx)}{5b} + \frac{\cos^3(a + bx)}{b} - \frac{\cos(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Sin[a + b*x]^7,x]`

[Out] $-(\text{Cos}[a + b*x]/b) + \text{Cos}[a + b*x]^3/b - (3*\text{Cos}[a + b*x]^5)/(5*b) + \text{Cos}[a + b*x]^7/(7*b)$

Rule 2713

`Int[sin[(c_.) + (d_.)*(x_)]^(n_), x_Symbol] := Dist[-d^(-1), Subst[Int[Expand[(1 - x^2)^((n - 1)/2), x], x], x, Cos[c + d*x]], x] /; FreeQ[{c, d}, x] && IGtQ[(n - 1)/2, 0]`

Rubi steps

$$\begin{aligned} \int \sin^7(a + bx) dx &= -\frac{\text{Subst}\left(\int (1 - 3x^2 + 3x^4 - x^6) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos(a + bx)}{b} + \frac{\cos^3(a + bx)}{b} - \frac{3 \cos^5(a + bx)}{5b} + \frac{\cos^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 59, normalized size = 1.09

$$-\frac{35 \cos(a + bx)}{64b} + \frac{7 \cos(3(a + bx))}{64b} - \frac{7 \cos(5(a + bx))}{320b} + \frac{\cos(7(a + bx))}{448b}$$

Antiderivative was successfully verified.

[In] `Integrate[Sin[a + b*x]^7,x]`

[Out] $(-35 \cos[a + b*x]) / (64*b) + (7 \cos[3*(a + b*x)]) / (64*b) - (7 \cos[5*(a + b*x)]) / (320*b) + \cos[7*(a + b*x)] / (448*b)$

Maple [A]

time = 0.11, size = 42, normalized size = 0.78

method	result	size
derivativedivides	$-\frac{\left(\frac{16}{5} + \sin^6(bx+a) + \frac{6(\sin^4(bx+a))}{5} + \frac{8(\sin^2(bx+a))}{5}\right) \cos(bx+a)}{7b}$	42
default	$-\frac{\left(\frac{16}{5} + \sin^6(bx+a) + \frac{6(\sin^4(bx+a))}{5} + \frac{8(\sin^2(bx+a))}{5}\right) \cos(bx+a)}{7b}$	42
risch	$-\frac{35 \cos(bx+a)}{64b} + \frac{\cos(7bx+7a)}{448b} - \frac{7 \cos(5bx+5a)}{320b} + \frac{7 \cos(3bx+3a)}{64b}$	55
norman	$\frac{\frac{32(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{32}{35b} - \frac{32(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{96(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{5b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^7}$	71

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^7,x,method=_RETURNVERBOSE)`

[Out] $-1/7/b*(16/5+\sin(b*x+a)^6+6/5*\sin(b*x+a)^4+8/5*\sin(b*x+a)^2)*\cos(b*x+a)$

Maxima [A]

time = 0.33, size = 44, normalized size = 0.81

$$\frac{5 \cos(bx+a)^7 - 21 \cos(bx+a)^5 + 35 \cos(bx+a)^3 - 35 \cos(bx+a)}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^7,x, algorithm="maxima")`

[Out] $1/35*(5*\cos(b*x + a)^7 - 21*\cos(b*x + a)^5 + 35*\cos(b*x + a)^3 - 35*\cos(b*x + a))/b$

Fricas [A]

time = 0.37, size = 44, normalized size = 0.81

$$\frac{5 \cos(bx+a)^7 - 21 \cos(bx+a)^5 + 35 \cos(bx+a)^3 - 35 \cos(bx+a)}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^7,x, algorithm="fricas")`

[Out] $1/35*(5*\cos(b*x + a)^7 - 21*\cos(b*x + a)^5 + 35*\cos(b*x + a)^3 - 35*\cos(b*x + a))/b$

Sympy [A]

time = 0.61, size = 80, normalized size = 1.48

$$\begin{cases} -\frac{\sin^6(a+bx)\cos(a+bx)}{b} - \frac{2\sin^4(a+bx)\cos^3(a+bx)}{b} - \frac{8\sin^2(a+bx)\cos^5(a+bx)}{5b} - \frac{16\cos^7(a+bx)}{35b} & \text{for } b \neq 0 \\ x \sin^7(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**7,x)

[Out] Piecewise((-sin(a + b*x)**6*cos(a + b*x)/b - 2*sin(a + b*x)**4*cos(a + b*x)**3/b - 8*sin(a + b*x)**2*cos(a + b*x)**5/(5*b) - 16*cos(a + b*x)**7/(35*b), Ne(b, 0)), (x*sin(a)**7, True))

Giac [A]

time = 4.34, size = 50, normalized size = 0.93

$$\frac{\cos(bx+a)^7}{7b} - \frac{3\cos(bx+a)^5}{5b} + \frac{\cos(bx+a)^3}{b} - \frac{\cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^7,x, algorithm="giac")

[Out] 1/7*cos(b*x + a)^7/b - 3/5*cos(b*x + a)^5/b + cos(b*x + a)^3/b - cos(b*x + a)/b

Mupad [B]

time = 0.38, size = 43, normalized size = 0.80

$$\frac{\cos(a+bx)(5\cos(a+bx)^6 - 21\cos(a+bx)^4 + 35\cos(a+bx)^2 - 35)}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^7,x)

[Out] (cos(a + b*x)*(35*cos(a + b*x)^2 - 21*cos(a + b*x)^4 + 5*cos(a + b*x)^6 - 35))/(35*b)

3.8 $\int \sin^8(a + bx) dx$

Optimal. Leaf size=88

$$\frac{35x}{128} - \frac{35 \cos(a + bx) \sin(a + bx)}{128b} - \frac{35 \cos(a + bx) \sin^3(a + bx)}{192b} - \frac{7 \cos(a + bx) \sin^5(a + bx)}{48b} - \frac{\cos(a + bx) \sin^7(a + bx)}{8b}$$

[Out] 35/128*x-35/128*cos(b*x+a)*sin(b*x+a)/b-35/192*cos(b*x+a)*sin(b*x+a)^3/b-7/48*cos(b*x+a)*sin(b*x+a)^5/b-1/8*cos(b*x+a)*sin(b*x+a)^7/b

Rubi [A]

time = 0.03, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$,

Rules used = {2715, 8}

$$-\frac{\sin^7(a + bx) \cos(a + bx)}{8b} - \frac{7 \sin^5(a + bx) \cos(a + bx)}{48b} - \frac{35 \sin^3(a + bx) \cos(a + bx)}{192b} - \frac{35 \sin(a + bx) \cos(a + bx)}{128b} + \frac{35x}{128}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^8,x]

[Out] (35*x)/128 - (35*Cos[a + b*x]*Sin[a + b*x])/(128*b) - (35*Cos[a + b*x]*Sin[a + b*x]^3)/(192*b) - (7*Cos[a + b*x]*Sin[a + b*x]^5)/(48*b) - (Cos[a + b*x]*Sin[a + b*x]^7)/(8*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \sin^8(a+bx) dx &= -\frac{\cos(a+bx)\sin^7(a+bx)}{8b} + \frac{7}{8} \int \sin^6(a+bx) dx \\
&= -\frac{7\cos(a+bx)\sin^5(a+bx)}{48b} - \frac{\cos(a+bx)\sin^7(a+bx)}{8b} + \frac{35}{48} \int \sin^4(a+bx) dx \\
&= -\frac{35\cos(a+bx)\sin^3(a+bx)}{192b} - \frac{7\cos(a+bx)\sin^5(a+bx)}{48b} - \frac{\cos(a+bx)\sin^7(a+bx)}{8b} \\
&= -\frac{35\cos(a+bx)\sin(a+bx)}{128b} - \frac{35\cos(a+bx)\sin^3(a+bx)}{192b} - \frac{7\cos(a+bx)\sin^5(a+bx)}{48b} \\
&= \frac{35x}{128} - \frac{35\cos(a+bx)\sin(a+bx)}{128b} - \frac{35\cos(a+bx)\sin^3(a+bx)}{192b} - \frac{7\cos(a+bx)\sin^5(a+bx)}{48b}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.62

$$\frac{840a + 840bx - 672\sin(2(a+bx)) + 168\sin(4(a+bx)) - 32\sin(6(a+bx)) + 3\sin(8(a+bx))}{3072b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^8, x]`

```
[Out] (840*a + 840*b*x - 672*Sin[2*(a + b*x)] + 168*Sin[4*(a + b*x)] - 32*Sin[6*(a + b*x)] + 3*Sin[8*(a + b*x)])/(3072*b)
```

Maple [A]

time = 0.14, size = 58, normalized size = 0.66

method	result
derivativedivides	$-\frac{\left(\sin^7(bx+a) + \frac{7(\sin^5(bx+a))}{6} + \frac{35(\sin^3(bx+a))}{24} + \frac{35\sin(bx+a)}{16}\right)\cos(bx+a)}{8b} + \frac{35bx}{128} + \frac{35a}{128}$
default	$-\frac{\left(\sin^7(bx+a) + \frac{7(\sin^5(bx+a))}{6} + \frac{35(\sin^3(bx+a))}{24} + \frac{35\sin(bx+a)}{16}\right)\cos(bx+a)}{8b} + \frac{35bx}{128} + \frac{35a}{128}$
risch	$\frac{35x}{128} + \frac{\sin(8bx+8a)}{1024b} - \frac{\sin(6bx+6a)}{96b} + \frac{7\sin(4bx+4a)}{128b} - \frac{7\sin(2bx+2a)}{32b}$
norman	$\frac{35x}{128} - \frac{35\tan\left(\frac{bx+a}{2}\right)}{64b} - \frac{805\left(\tan^3\left(\frac{bx+a}{2}\right)\right)}{192b} - \frac{2681\left(\tan^5\left(\frac{bx+a}{2}\right)\right)}{192b} - \frac{5053\left(\tan^7\left(\frac{bx+a}{2}\right)\right)}{192b} + \frac{5053\left(\tan^9\left(\frac{bx+a}{2}\right)\right)}{192b} + \frac{2681\left(\tan^{11}\left(\frac{bx+a}{2}\right)\right)}{192b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^8, x, method=_RETURNVERBOSE)`

[Out] $1/b*(-1/8*(\sin(b*x+a)^7+7/6*\sin(b*x+a)^5+35/24*\sin(b*x+a)^3+35/16*\sin(b*x+a))*\cos(b*x+a)+35/128*b*x+35/128*a)$

Maxima [A]

time = 0.32, size = 59, normalized size = 0.67

$$\frac{128 \sin(2bx + 2a)^3 + 840bx + 840a + 3 \sin(8bx + 8a) + 168 \sin(4bx + 4a) - 768 \sin(2bx + 2a)}{3072b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^8,x, algorithm="maxima")`

[Out] $1/3072*(128*\sin(2*b*x + 2*a)^3 + 840*b*x + 840*a + 3*\sin(8*b*x + 8*a) + 168*\sin(4*b*x + 4*a) - 768*\sin(2*b*x + 2*a))/b$

Fricas [A]

time = 0.43, size = 56, normalized size = 0.64

$$\frac{105bx + (48 \cos(bx + a))^7 - 200 \cos(bx + a)^5 + 326 \cos(bx + a)^3 - 279 \cos(bx + a) \sin(bx + a)}{384b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^8,x, algorithm="fricas")`

[Out] $1/384*(105*b*x + (48*\cos(b*x + a))^7 - 200*\cos(b*x + a)^5 + 326*\cos(b*x + a)^3 - 279*\cos(b*x + a))*\sin(b*x + a)/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 184 vs. $2(82) = 164$.

time = 0.90, size = 184, normalized size = 2.09

$$\begin{cases} \frac{35x \sin^6(a+bx)}{128} + \frac{35x \sin^6(a+bx) \cos^2(a+bx)}{32} + \frac{105x \sin^4(a+bx) \cos^4(a+bx)}{64} + \frac{35x \sin^2(a+bx) \cos^6(a+bx)}{32} + \frac{35x \cos^8(a+bx)}{128} - \frac{93 \sin^7(a+bx) \cos(a+bx)}{128b} - \frac{511 \sin^5(a+bx) \cos^3(a+bx)}{384b} - \frac{385 \sin^3(a+bx) \cos^5(a+bx)}{384b} - \frac{35 \sin(a+bx) \cos^7(a+bx)}{128b} & \text{for } b \neq 0 \\ x \sin^8(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**8,x)`

[Out] `Piecewise((35*x*sin(a + b*x)**8/128 + 35*x*sin(a + b*x)**6*cos(a + b*x)**2/32 + 105*x*sin(a + b*x)**4*cos(a + b*x)**4/64 + 35*x*sin(a + b*x)**2*cos(a + b*x)**6/32 + 35*x*cos(a + b*x)**8/128 - 93*sin(a + b*x)**7*cos(a + b*x)/(128*b) - 511*sin(a + b*x)**5*cos(a + b*x)**3/(384*b) - 385*sin(a + b*x)**3*cos(a + b*x)**5/(384*b) - 35*sin(a + b*x)*cos(a + b*x)**7/(128*b), Ne(b, 0)), (x*sin(a)**8, True))`

Giac [A]

time = 4.69, size = 60, normalized size = 0.68

$$\frac{35}{128}x + \frac{\sin(8bx + 8a)}{1024b} - \frac{\sin(6bx + 6a)}{96b} + \frac{7 \sin(4bx + 4a)}{128b} - \frac{7 \sin(2bx + 2a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^8,x, algorithm="giac")

[Out] $35/128*x + 1/1024*\sin(8*b*x + 8*a)/b - 1/96*\sin(6*b*x + 6*a)/b + 7/128*\sin(4*b*x + 4*a)/b - 7/32*\sin(2*b*x + 2*a)/b$

Mupad [B]

time = 1.50, size = 90, normalized size = 1.02

$$\frac{35x}{128} - \frac{\frac{93 \tan(a+bx)^7}{128} + \frac{511 \tan(a+bx)^5}{384} + \frac{385 \tan(a+bx)^3}{384} + \frac{35 \tan(a+bx)}{128}}{b (\tan(a+bx)^8 + 4 \tan(a+bx)^6 + 6 \tan(a+bx)^4 + 4 \tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^8,x)

[Out] $(35*x)/128 - ((35*\tan(a + b*x))/128 + (385*\tan(a + b*x)^3)/384 + (511*\tan(a + b*x)^5)/384 + (93*\tan(a + b*x)^7)/128)/(b*(4*\tan(a + b*x)^2 + 6*\tan(a + b*x)^4 + 4*\tan(a + b*x)^6 + \tan(a + b*x)^8 + 1))$

3.9 $\int \sin^{\frac{7}{2}}(bx) dx$

Optimal. Leaf size=60

$$-\frac{10F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{21b} - \frac{10 \cos(bx) \sqrt{\sin(bx)}}{21b} - \frac{2 \cos(bx) \sin^{\frac{5}{2}}(bx)}{7b}$$

[Out] $-10/21*(\sin(1/4*\text{Pi}+1/2*b*x)^2)^{(1/2)}/\sin(1/4*\text{Pi}+1/2*b*x)*\text{EllipticF}(\cos(1/4*\text{Pi}+1/2*b*x), 2^{(1/2)})/b-2/7*\cos(b*x)*\sin(b*x)^{(5/2)}/b-10/21*\cos(b*x)*\sin(b*x)^{(1/2)}/b$

Rubi [A]

time = 0.02, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2720}

$$-\frac{10F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{21b} - \frac{2 \sin^{\frac{5}{2}}(bx) \cos(bx)}{7b} - \frac{10 \sqrt{\sin(bx)} \cos(bx)}{21b}$$

Antiderivative was successfully verified.

[In] Int[Sin[b*x]^(7/2), x]

[Out] $(-10*\text{EllipticF}[\text{Pi}/4 - (b*x)/2, 2])/(21*b) - (10*\text{Cos}[b*x]*\text{Sqrt}[\text{Sin}[b*x]])/(21*b) - (2*\text{Cos}[b*x]*\text{Sin}[b*x]^{(5/2)})/(7*b)$

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sine[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sine[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sin^{\frac{7}{2}}(bx) dx &= -\frac{2 \cos(bx) \sin^{\frac{5}{2}}(bx)}{7b} + \frac{5}{7} \int \sin^{\frac{3}{2}}(bx) dx \\ &= -\frac{10 \cos(bx) \sqrt{\sin(bx)}}{21b} - \frac{2 \cos(bx) \sin^{\frac{5}{2}}(bx)}{7b} + \frac{5}{21} \int \frac{1}{\sqrt{\sin(bx)}} dx \\ &= -\frac{10F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{21b} - \frac{10 \cos(bx) \sqrt{\sin(bx)}}{21b} - \frac{2 \cos(bx) \sin^{\frac{5}{2}}(bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 45, normalized size = 0.75

$$\frac{-20F\left(\frac{1}{4}(\pi - 2bx) \mid 2\right) + (-23 \cos(bx) + 3 \cos(3bx)) \sqrt{\sin(bx)}}{42b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[b*x]^(7/2),x]`

```
[Out] (-20*EllipticF[(Pi - 2*b*x)/4, 2] + (-23*Cos[b*x] + 3*Cos[3*b*x])*Sqrt[Sin[
b*x]])/(42*b)
```

Maple [A]

time = 0.10, size = 84, normalized size = 1.40

method	result
default	$\frac{\frac{2 \sin(bx) (\cos^4(bx))}{7} + \frac{5 \sqrt{\sin(bx) + 1} \sqrt{-2 \sin(bx) + 2} \sqrt{-\sin(bx)} \operatorname{EllipticF}\left(\sqrt{\sin(bx) + 1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx) \sqrt{\sin(bx)} b} - \frac{16(\cos(bx))^{5/2}}{21 b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x)^(7/2),x,method=_RETURNVERBOSE)`

```
[Out] (2/7*sin(b*x)*cos(b*x)^4+5/21*(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin
(b*x))^(1/2)*EllipticF((sin(b*x)+1)^(1/2),1/2*2^(1/2))-16/21*cos(b*x)^2*si
n(b*x))/cos(b*x)/sin(b*x)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x)^(7/2),x, algorithm="maxima")``[Out] integrate(sin(b*x)^(7/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.15, size = 73, normalized size = 1.22

$$\frac{5 \sqrt{2} \sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx) + i \sin(bx)) + 5 \sqrt{2} \sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx) - i \sin(bx)) + 2 (3 \cos(bx)^3 - 8 \cos(bx)) \sqrt{\sin(bx)}}{21 b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x)^(7/2),x, algorithm="fricas")`

[Out] $\frac{1}{21} \cdot (5\sqrt{2}\sqrt{-1}\text{weierstrassPInverse}(4, 0, \cos(bx) + I\sin(bx)) + 5\sqrt{2}\sqrt{1}\text{weierstrassPInverse}(4, 0, \cos(bx) - I\sin(bx)) + 2 \cdot (3\cos(bx)^3 - 8\cos(bx))\sqrt{\sin(bx)})/b$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x)**(7/2), x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3060 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x)^(7/2), x, algorithm="giac")`

[Out] `integrate(sin(b*x)^(7/2), x)`

Mupad [B]

time = 0.48, size = 34, normalized size = 0.57

$$-\frac{\cos(bx) \sin(bx)^{9/2} {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; \frac{3}{2}; \cos(bx)^2\right)}{b (\sin(bx)^2)^{9/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x)^(7/2), x)`

[Out] $-(\cos(bx) \cdot \sin(bx)^{(9/2)} \cdot \text{hypergeom}([-5/4, 1/2], 3/2, \cos(bx)^2)) / (b \cdot (\sin(bx)^2)^{(9/4)})$

3.10 $\int \sin^{\frac{5}{2}}(bx) dx$

Optimal. Leaf size=41

$$-\frac{6E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{5b} - \frac{2 \cos(bx) \sin^{\frac{3}{2}}(bx)}{5b}$$

[Out] $-6/5*(\sin(1/4*\text{Pi}+1/2*b*x)^2)^{(1/2)}/\sin(1/4*\text{Pi}+1/2*b*x)*\text{EllipticE}(\cos(1/4*\text{Pi}+1/2*b*x), 2^{(1/2)})/b-2/5*\cos(b*x)*\sin(b*x)^{(3/2)}/b$

Rubi [A]

time = 0.01, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2719}

$$-\frac{6E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{5b} - \frac{2 \sin^{\frac{3}{2}}(bx) \cos(bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sin[b*x]^(5/2), x]

[Out] $(-6*\text{EllipticE}[\text{Pi}/4 - (b*x)/2, 2])/(5*b) - (2*\text{Cos}[b*x]*\text{Sin}[b*x]^{(3/2)})/(5*b)$

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*SIN[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sin^{\frac{5}{2}}(bx) dx &= -\frac{2 \cos(bx) \sin^{\frac{3}{2}}(bx)}{5b} + \frac{3}{5} \int \sqrt{\sin(bx)} dx \\ &= -\frac{6E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{5b} - \frac{2 \cos(bx) \sin^{\frac{3}{2}}(bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 35, normalized size = 0.85

$$\frac{2\left(3E\left(\frac{1}{4}(\pi - 2bx) \mid 2\right) + \cos(bx) \sin^{\frac{3}{2}}(bx)\right)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[b*x]^(5/2), x]``[Out] (-2*(3*EllipticE[(Pi - 2*b*x)/4, 2] + Cos[b*x]*Sin[b*x]^(3/2)))/(5*b)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 117 vs. 2(58) = 116.

time = 0.06, size = 118, normalized size = 2.88

method	result
default	$\frac{\frac{2(\sin^4(bx))}{5} - \frac{2(\sin^2(bx))}{5} - \frac{6\sqrt{\sin(bx)+1}\sqrt{-2\sin(bx)+2}\sqrt{-\sin(bx)}}{5} \text{EllipticE}\left(\sqrt{\sin(bx)+1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx)\sqrt{\sin(bx)}b} +$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x)^(5/2), x, method=_RETURNVERBOSE)`

```
[Out] (2/5*sin(b*x)^4-2/5*sin(b*x)^2-6/5*(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)
*(-sin(b*x))^(1/2)*EllipticE((sin(b*x)+1)^(1/2), 1/2*2^(1/2))+3/5*(sin(b*x)+
1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*EllipticF((sin(b*x)+1)^(1/
2), 1/2*2^(1/2)))/cos(b*x)/sin(b*x)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x)^(5/2), x, algorithm="maxima")``[Out] integrate(sin(b*x)^(5/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 68, normalized size = 1.66

$$\frac{-2\cos(bx)\sin(bx)^{\frac{3}{2}} - 3i\sqrt{2}\sqrt{-i}\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx) + i\sin(bx))) + 3i\sqrt{2}\sqrt{i}\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx) - i\sin(bx)))}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x)^(5/2),x, algorithm="fricas")

[Out] $-1/5*(2*\cos(b*x)*\sin(b*x)^{(3/2)} - 3*I*\sqrt{2}*\sqrt{-I}*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x) + I*\sin(b*x))) + 3*I*\sqrt{2}*\sqrt{I}*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x) - I*\sin(b*x))))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^{\frac{5}{2}}(bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x)**(5/2),x)

[Out] Integral(sin(b*x)**(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x)^(5/2),x, algorithm="giac")

[Out] integrate(sin(b*x)^(5/2), x)

Mupad [B]

time = 0.41, size = 34, normalized size = 0.83

$$\frac{\cos(bx) \sin(bx)^{7/2} {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{3}{2}; \cos(bx)^2\right)}{b (\sin(bx)^2)^{7/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x)^(5/2),x)

[Out] $-(\cos(b*x)*\sin(b*x)^{(7/2)}*\text{hypergeom}([-3/4, 1/2], 3/2, \cos(b*x)^2))/(b*(\sin(b*x)^2)^{(7/4)})$

3.11 $\int \sin^{\frac{3}{2}}(bx) dx$

Optimal. Leaf size=41

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{3b} - \frac{2 \cos(bx) \sqrt{\sin(bx)}}{3b}$$

[Out] $-2/3*(\sin(1/4*\text{Pi}+1/2*b*x)^2)^{(1/2)}/\sin(1/4*\text{Pi}+1/2*b*x)*\text{EllipticF}(\cos(1/4*\text{Pi}+1/2*b*x),2^{(1/2)})/b-2/3*\cos(b*x)*\sin(b*x)^{(1/2)}/b$

Rubi [A]

time = 0.01, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2720}

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{3b} - \frac{2 \sqrt{\sin(bx)} \cos(bx)}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[b*x]^{(3/2)}, x]$

[Out] $(-2*\text{EllipticF}[\text{Pi}/4 - (b*x)/2, 2])/(3*b) - (2*\text{Cos}[b*x]*\text{Sqrt}[\text{Sin}[b*x]])/(3*b)$

Rule 2715

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n-1)})/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sin^{\frac{3}{2}}(bx) dx &= -\frac{2 \cos(bx) \sqrt{\sin(bx)}}{3b} + \frac{1}{3} \int \frac{1}{\sqrt{\sin(bx)}} dx \\ &= -\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{3b} - \frac{2 \cos(bx) \sqrt{\sin(bx)}}{3b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 33, normalized size = 0.80

$$\frac{2 \left(F\left(\frac{1}{4}(\pi - 2bx) \mid 2\right) + \cos(bx) \sqrt{\sin(bx)} \right)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[b*x]^(3/2), x]``[Out] (-2*(EllipticF[(Pi - 2*b*x)/4, 2] + Cos[b*x]*Sqrt[Sin[b*x]]))/(3*b)`**Maple [A]**

time = 0.06, size = 72, normalized size = 1.76

method	result	size
default	$\frac{\sqrt{\sin(bx) + 1} \sqrt{-2 \sin(bx) + 2} \sqrt{-\sin(bx)} \operatorname{EllipticF}\left(\sqrt{\sin(bx) + 1}, \frac{\sqrt{2}}{2}\right) - 2 \frac{(\cos^2(bx) \sin(bx))^{3/2}}{3}}{\cos(bx) \sqrt{\sin(bx)} b}$	72

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x)^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] (1/3*(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*EllipticF((
sin(b*x)+1)^(1/2), 1/2*2^(1/2))-2/3*cos(b*x)^2*sin(b*x))/cos(b*x)/sin(b*x)^(
1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x)^(3/2), x, algorithm="maxima")``[Out] integrate(sin(b*x)^(3/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 60, normalized size = 1.46

$$\frac{\sqrt{2} \sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx) + i \sin(bx)) + \sqrt{2} \sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx) - i \sin(bx)) - 2 \cos(bx) \sqrt{\sin(bx)}}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x)^(3/2), x, algorithm="fricas")`

[Out] $\frac{1}{3}(\sqrt{2})\sqrt{-1}\text{weierstrassPInverse}(4, 0, \cos(bx) + I\sin(bx)) + \sqrt{2}\sqrt{I}\text{weierstrassPInverse}(4, 0, \cos(bx) - I\sin(bx)) - 2\cos(bx)\sqrt{\sin(bx)}/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^{\frac{3}{2}}(bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x)**(3/2),x)`

[Out] `Integral(sin(b*x)**(3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x)^(3/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x)^(3/2), x)`

Mupad [B]

time = 0.40, size = 34, normalized size = 0.83

$$-\frac{\cos(bx) \sin(bx)^{5/2} {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{2}; \cos(bx)^2\right)}{b (\sin(bx)^2)^{5/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x)^(3/2),x)`

[Out] `-(cos(b*x)*sin(b*x)^(5/2)*hypergeom([-1/4, 1/2], 3/2, cos(b*x)^2))/(b*(sin(b*x)^2)^(5/4))`

3.12 $\int \sqrt{\sin(bx)} dx$

Optimal. Leaf size=19

$$-\frac{2E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

[Out] $-2*(\sin(1/4*\text{Pi}+1/2*b*x)^2)^{(1/2)}/\sin(1/4*\text{Pi}+1/2*b*x)*\text{EllipticE}(\cos(1/4*\text{Pi}+1/2*b*x), 2^{(1/2)})/b$

Rubi [A]

time = 0.01, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2719}

$$-\frac{2E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sin[b*x]], x]

[Out] $(-2*\text{EllipticE}[\text{Pi}/4 - (b*x)/2, 2])/b$

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \sqrt{\sin(bx)} dx = -\frac{2E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

Mathematica [A]

time = 0.02, size = 21, normalized size = 1.11

$$-\frac{2E\left(\frac{1}{2}\left(\frac{\pi}{2} - bx\right) \mid 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sin[b*x]], x]

[Out] $(-2*\text{EllipticE}[(\text{Pi}/2 - b*x)/2, 2])/b$

Maple [A]

time = 0.11, size = 77, normalized size = 4.05

method	result
default	$-\frac{\sqrt{\sin(bx)+1} \sqrt{-2\sin(bx)+2} \sqrt{-\sin(bx)} \left(2 \operatorname{EllipticE}\left(\sqrt{\sin(bx)+1}, \frac{\sqrt{2}}{2}\right) - \operatorname{EllipticF}\left(\sqrt{\sin(bx)+1}, \frac{\sqrt{2}}{2}\right) \right)}{\cos(bx) \sqrt{\sin(bx)} b}$
risch	$-\frac{i\sqrt{2} \sqrt{-i(e^{2ibx}-1)e^{-ibx}}}{b} + i \left(\frac{2i(i-ie^{2ibx})}{\sqrt{e^{ibx}(i-ie^{2ibx})}} - \frac{\sqrt{e^{ibx}+1} \sqrt{-2e^{ibx}+2} \sqrt{-e^{ibx}} \left(-2 \operatorname{EllipticE}\left(\sqrt{e^{ibx}+1}, \frac{\sqrt{2}}{2}\right) - \operatorname{EllipticF}\left(\sqrt{e^{ibx}+1}, \frac{\sqrt{2}}{2}\right) \right)}{\sqrt{-ie^{3ibx}} \cos(bx) \sqrt{\sin(bx)} b} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*(2*EllipticE((sin(b*x)+1)^(1/2),1/2*2^(1/2))-EllipticF((sin(b*x)+1)^(1/2),1/2*2^(1/2)))/cos(b*x)/sin(b*x)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(sin(b*x)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 55, normalized size = 2.89

$$\frac{i\sqrt{2}\sqrt{-i}\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx)+i\sin(bx))) - i\sqrt{2}\sqrt{i}\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx)-i\sin(bx)))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x)^(1/2),x, algorithm="fricas")
```

```
[Out] (I*sqrt(2)*sqrt(-I)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x) + I*sin(b*x))) - I*sqrt(2)*sqrt(I)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x) - I*sin(b*x))))/b
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\sin(bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x)**(1/2),x)

[Out] Integral(sqrt(sin(b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(sin(b*x)), x)

Mupad [B]

time = 0.37, size = 15, normalized size = 0.79

$$-\frac{2 E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x)^(1/2),x)

[Out] -(2*ellipticE(pi/4 - (b*x)/2, 2))/b

$$3.13 \quad \int \frac{1}{\sqrt{\sin(bx)}} dx$$

Optimal. Leaf size=19

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

[Out] $-2*(\sin(1/4*\text{Pi}+1/2*b*x)^2)^{(1/2)}/\sin(1/4*\text{Pi}+1/2*b*x)*\text{EllipticF}(\cos(1/4*\text{Pi}+1/2*b*x),2^{(1/2)})/b$

Rubi [A]

time = 0.01, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2720}

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[Sin[b*x]],x]

[Out] $(-2*\text{EllipticF}[\text{Pi}/4 - (b*x)/2, 2])/b$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\sqrt{\sin(bx)}} dx = -\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

Mathematica [A]

time = 0.02, size = 21, normalized size = 1.11

$$-\frac{2F\left(\frac{1}{2}\left(\frac{\pi}{2} - bx\right) \mid 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[Sin[b*x]],x]

[Out] $(-2*\text{EllipticF}[(\text{Pi}/2 - b*x)/2, 2])/b$

Maple [A]

time = 0.05, size = 57, normalized size = 3.00

method	result	size
default	$\frac{\sqrt{\sin(bx)+1} \sqrt{-2\sin(bx)+2} \sqrt{-\sin(bx)} \operatorname{EllipticF}\left(\sqrt{\sin(bx)+1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx) \sqrt{\sin(bx)} b}$	57

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/sin(b*x)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] (sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*EllipticF((sin(b*x)+1)^(1/2),1/2*2^(1/2))/cos(b*x)/sin(b*x)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(b*x)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/sqrt(sin(b*x)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 47, normalized size = 2.47

$$\frac{\sqrt{2} \sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx) + i \sin(bx)) + \sqrt{2} \sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx) - i \sin(bx))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(b*x)^(1/2),x, algorithm="fricas")
```

```
[Out] (sqrt(2)*sqrt(-I)*weierstrassPInverse(4, 0, cos(b*x) + I*sin(b*x)) + sqrt(2)*sqrt(I)*weierstrassPInverse(4, 0, cos(b*x) - I*sin(b*x)))/b
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\sin(bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(b*x)**(1/2),x)
```

```
[Out] Integral(1/sqrt(sin(b*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(sin(b*x)), x)

Mupad [B]

time = 0.39, size = 15, normalized size = 0.79

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sin(b*x)^(1/2),x)

[Out] -(2*ellipticF(pi/4 - (b*x)/2, 2))/b

$$3.14 \quad \int \frac{1}{\sin^{\frac{3}{2}}(bx)} dx$$

Optimal. Leaf size=37

$$\frac{2E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b} - \frac{2 \cos(bx)}{b\sqrt{\sin(bx)}}$$

[Out] 2*(sin(1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/4*Pi+1/2*b*x)*EllipticE(cos(1/4*Pi+1/2*b*x),2^(1/2))/b-2*cos(b*x)/b/sin(b*x)^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2719}

$$\frac{2E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b} - \frac{2 \cos(bx)}{b\sqrt{\sin(bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[b*x]^(-3/2),x]

[Out] (2*EllipticE[Pi/4 - (b*x)/2, 2])/b - (2*Cos[b*x])/(b*Sqrt[Sin[b*x]])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sin^{\frac{3}{2}}(bx)} dx &= -\frac{2 \cos(bx)}{b\sqrt{\sin(bx)}} - \int \sqrt{\sin(bx)} dx \\ &= \frac{2E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{b} - \frac{2 \cos(bx)}{b\sqrt{\sin(bx)}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 32, normalized size = 0.86

$$\frac{2 \left(E\left(\frac{1}{4}(\pi - 2bx) \mid 2\right) - \frac{\cos(bx)}{\sqrt{\sin(bx)}} \right)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[b*x]^(-3/2), x]``[Out] (2*(EllipticE[(Pi - 2*b*x)/4, 2] - Cos[b*x]/Sqrt[Sin[b*x]]))/b`**Maple [A]**

time = 0.05, size = 110, normalized size = 2.97

method	result
default	$\frac{2 \sqrt{\sin(bx) + 1} \sqrt{-2 \sin(bx) + 2} \sqrt{-\sin(bx)} \operatorname{EllipticE}\left(\sqrt{\sin(bx) + 1}, \frac{\sqrt{2}}{2}\right) - \sqrt{\sin(bx) + 1} \sqrt{-\sin(bx)}}{\cos(bx) \sqrt{\sin(bx)} b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(b*x)^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] (2*(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*EllipticE((sin(b*x)+1)^(1/2), 1/2*2^(1/2))-sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*EllipticF((sin(b*x)+1)^(1/2), 1/2*2^(1/2))-2*cos(b*x)^2)/cos(b*x)/sin(b*x)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x)^(3/2), x, algorithm="maxima")``[Out] integrate(sin(b*x)^(-3/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 81, normalized size = 2.19

$$\frac{-i \sqrt{2} \sqrt{-i} \sin(bx) \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(bx) + i \sin(bx))) + i \sqrt{2} \sqrt{i} \sin(bx) \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(bx) - i \sin(bx))) - 2 \cos(bx) \sqrt{\sin(bx)}}{b \sin(bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x)^(3/2), x, algorithm="fricas")`

[Out] $(-I\sqrt{2}\sqrt{-I}\sin(bx)\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx) + I\sin(bx))) + I\sqrt{2}\sqrt{I}\sin(bx)\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx) - I\sin(bx))) - 2\cos(bx)\sqrt{\sin(bx)})/(b\sin(bx))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sin^{\frac{3}{2}}(bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(b*x)**(3/2),x)`

[Out] `Integral(sin(b*x)**(-3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(b*x)^(3/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x)^(-3/2), x)`

Mupad [B]

time = 0.47, size = 34, normalized size = 0.92

$$-\frac{\cos(bx) (\sin(bx)^2)^{1/4} {}_2F_1\left(\frac{1}{2}, \frac{5}{4}; \frac{3}{2}; \cos(bx)^2\right)}{b \sqrt{\sin(bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sin(b*x)^(3/2),x)`

[Out] `-(cos(b*x)*(sin(b*x)^2)^(1/4)*hypergeom([1/2, 5/4], 3/2, cos(b*x)^2))/(b*sin(b*x)^(1/2))`

3.15

$$\int \frac{1}{\sin^{\frac{5}{2}}(bx)} dx$$

Optimal. Leaf size=41

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{3b} - \frac{2 \cos(bx)}{3b \sin^{\frac{3}{2}}(bx)}$$

[Out] $-2/3*(\sin(1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/4*Pi+1/2*b*x)*\text{EllipticF}(\cos(1/4*Pi+1/2*b*x), 2^{(1/2)})/b-2/3*\cos(b*x)/b/\sin(b*x)^{(3/2)}$

Rubi [A]

time = 0.01, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2720}

$$-\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{3b} - \frac{2 \cos(bx)}{3b \sin^{\frac{3}{2}}(bx)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[b*x]^{(-5/2)}, x]$

[Out] $(-2*\text{EllipticF}[\text{Pi}/4 - (b*x)/2, 2])/(3*b) - (2*\text{Cos}[b*x])/(3*b*\text{Sin}[b*x]^{(3/2)})$

Rule 2716

$\text{Int}[(b_* \sin[(c_*) + (d_*)*(x_*)])^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1))), x] + \text{Dist}[(n + 2)/(b^2*(n + 1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n + 2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sin^{\frac{5}{2}}(bx)} dx &= -\frac{2 \cos(bx)}{3b \sin^{\frac{3}{2}}(bx)} + \frac{1}{3} \int \frac{1}{\sqrt{\sin(bx)}} dx \\ &= -\frac{2F\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{3b} - \frac{2 \cos(bx)}{3b \sin^{\frac{3}{2}}(bx)} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 33, normalized size = 0.80

$$\frac{2 \left(F\left(\frac{1}{4}(\pi - 2bx) \mid 2\right) + \frac{\cos(bx)}{\sin^{\frac{3}{2}}(bx)} \right)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[b*x]^(-5/2), x]``[Out] (-2*(EllipticF[(Pi - 2*b*x)/4, 2] + Cos[b*x]/Sin[b*x]^(3/2)))/(3*b)`**Maple [A]**

time = 0.05, size = 72, normalized size = 1.76

method	result	s
default	$\frac{\sqrt{\sin(bx) + 1} \sqrt{-2 \sin(bx) + 2} \sqrt{-\sin(bx)} \operatorname{EllipticF}\left(\sqrt{\sin(bx) + 1}, \frac{\sqrt{2}}{2}\right) \sin(bx) - 2(\cos^2(bx))}{3 \sin(bx)^{\frac{3}{2}} \cos(bx)b}$	7

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(b*x)^(5/2), x, method=_RETURNVERBOSE)`
`[Out] 1/3/sin(b*x)^(3/2)*((sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*EllipticF((sin(b*x)+1)^(1/2), 1/2*2^(1/2))*sin(b*x)-2*cos(b*x)^2)/cos(b*x)/b`
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x)^(5/2), x, algorithm="maxima")``[Out] integrate(sin(b*x)^(-5/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 97, normalized size = 2.37

$$\frac{\sqrt{-i} \left(\sqrt{2} \cos(bx)^2 - \sqrt{2} \right) \operatorname{weierstrassPInverse}(4, 0, \cos(bx) + i \sin(bx)) + \sqrt{i} \left(\sqrt{2} \cos(bx)^2 - \sqrt{2} \right) \operatorname{weierstrassPInverse}(4, 0, \cos(bx) - i \sin(bx)) + 2 \cos(bx) \sqrt{\sin(bx)}}{3 (b \cos(bx)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x)^(5/2), x, algorithm="fricas")`

[Out] $\frac{1}{3}(\sqrt{-1})(\sqrt{2}\cos(bx)^2 - \sqrt{2})\text{weierstrassPInverse}(4, 0, \cos(bx) + I\sin(bx)) + \sqrt{I}(\sqrt{2}\cos(bx)^2 - \sqrt{2})\text{weierstrassPInverse}(4, 0, \cos(bx) - I\sin(bx)) + 2\cos(bx)\sqrt{\sin(bx)}/(b\cos(bx)^2 - b)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sin^{\frac{5}{2}}(bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(b*x)**(5/2), x)`

[Out] `Integral(sin(b*x)**(-5/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(b*x)^(5/2), x, algorithm="giac")`

[Out] `integrate(sin(b*x)^(-5/2), x)`

Mupad [B]

time = 0.59, size = 34, normalized size = 0.83

$$\frac{\cos(bx) (\sin(bx)^2)^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{3}{2}; \cos(bx)^2\right)}{b \sin(bx)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sin(b*x)^(5/2), x)`

[Out] `-(cos(b*x)*(sin(b*x)^2)^(3/4)*hypergeom([1/2, 7/4], 3/2, cos(b*x)^2))/(b*sin(b*x)^(3/2))`

3.16 $\int \frac{1}{\sin^2(bx)} dx$

Optimal. Leaf size=60

$$\frac{6E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{5b} - \frac{2 \cos(bx)}{5b \sin^{\frac{5}{2}}(bx)} - \frac{6 \cos(bx)}{5b \sqrt{\sin(bx)}}$$

[Out] 6/5*(sin(1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/4*Pi+1/2*b*x)*EllipticE(cos(1/4*Pi+1/2*b*x),2^(1/2))/b-2/5*cos(b*x)/b/sin(b*x)^(5/2)-6/5*cos(b*x)/b/sin(b*x)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2719}

$$\frac{6E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{5b} - \frac{2 \cos(bx)}{5b \sin^{\frac{5}{2}}(bx)} - \frac{6 \cos(bx)}{5b \sqrt{\sin(bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[b*x]^(-7/2),x]

[Out] (6*EllipticE[Pi/4 - (b*x)/2, 2])/(5*b) - (2*Cos[b*x])/(5*b*Sin[b*x]^(5/2)) - (6*Cos[b*x])/(5*b*Sqrt[Sin[b*x]])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sin^{\frac{7}{2}}(bx)} dx &= -\frac{2 \cos(bx)}{5b \sin^{\frac{5}{2}}(bx)} + \frac{3}{5} \int \frac{1}{\sin^{\frac{3}{2}}(bx)} dx \\
&= -\frac{2 \cos(bx)}{5b \sin^{\frac{5}{2}}(bx)} - \frac{6 \cos(bx)}{5b \sqrt{\sin(bx)}} - \frac{3}{5} \int \sqrt{\sin(bx)} dx \\
&= \frac{6E\left(\frac{\pi}{4} - \frac{bx}{2} \mid 2\right)}{5b} - \frac{2 \cos(bx)}{5b \sin^{\frac{5}{2}}(bx)} - \frac{6 \cos(bx)}{5b \sqrt{\sin(bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 51, normalized size = 0.85

$$\frac{-7 \cos(bx) + 3 \cos(3bx) + 12E\left(\frac{1}{4}(\pi - 2bx) \mid 2\right) \sin^{\frac{5}{2}}(bx)}{10b \sin^{\frac{5}{2}}(bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[b*x]^(-7/2), x]`

```
[Out] (-7*Cos[b*x] + 3*Cos[3*b*x] + 12*EllipticE[(Pi - 2*b*x)/4, 2]*Sin[b*x]^(5/2)) / (10*b*Sin[b*x]^(5/2))
```

Maple [A]

time = 0.06, size = 132, normalized size = 2.20

method	result
default	$ \frac{6 \sqrt{\sin(bx) + 1} \sqrt{-2 \sin(bx) + 2} \sqrt{-\sin(bx)} (\sin^2(bx)) \operatorname{EllipticE}\left(\sqrt{\sin(bx) + 1}, \frac{\sqrt{2}}{2}\right) - 3 \sqrt{\sin(bx)}}{5 \sin(bx)^{\frac{5}{2}}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(b*x)^(7/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/5/sin(b*x)^(5/2)*(6*(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*sin(b*x)^2*EllipticE((sin(b*x)+1)^(1/2), 1/2*2^(1/2))-3*(sin(b*x)+1)^(1/2)*(-2*sin(b*x)+2)^(1/2)*(-sin(b*x))^(1/2)*sin(b*x)^2*EllipticF((sin(b*x)+1)^(1/2), 1/2*2^(1/2))+6*sin(b*x)^4-4*sin(b*x)^2-2)/cos(b*x)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x)^(7/2),x, algorithm="maxima")

[Out] integrate(sin(b*x)^(-7/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 132, normalized size = 2.20

$$\frac{3\sqrt{-1}(i\sqrt{2}\cos(bx)^2 - i\sqrt{2})\sin(bx)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx) + i\sin(bx))) + 3\sqrt{1}(-i\sqrt{2}\cos(bx)^2 + i\sqrt{2})\sin(bx)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx) - i\sin(bx))) + 2(3\cos(bx)^3 - 4\cos(bx))\sqrt{\sin(bx)}}{5(b\cos(bx)^2 - b)\sin(bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x)^(7/2),x, algorithm="fricas")

[Out]
$$-1/5*(3*\sqrt{-1}*(I*\sqrt{2}*\cos(b*x)^2 - I*\sqrt{2})*\sin(b*x)*\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(b*x) + I*\sin(b*x))) + 3*\sqrt{1}*(-I*\sqrt{2}*\cos(b*x)^2 + I*\sqrt{2})*\sin(b*x)*\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(b*x) - I*\sin(b*x))) + 2*(3*\cos(b*x)^3 - 4*\cos(b*x))*\sqrt{\sin(b*x)})/(b*\cos(b*x)^2 - b)*\sin(b*x)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sin^{7/2}(bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x)**(7/2),x)

[Out] Integral(sin(b*x)**(-7/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x)^(7/2),x, algorithm="giac")

[Out] integrate(sin(b*x)^(-7/2), x)

Mupad [B]

time = 0.56, size = 34, normalized size = 0.57

$$-\frac{\cos(bx)(\sin(bx)^2)^{5/4} {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{3}{2}; \cos(bx)^2\right)}{b\sin(bx)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sin(b*x)^(7/2),x)

[Out]
$$-(\cos(b*x)*(\sin(b*x)^2)^{(5/4)}*\operatorname{hypergeom}([1/2, 9/4], 3/2, \cos(b*x)^2))/(b*\sin(b*x)^{(5/2)})$$

3.17 $\int \sin^{\frac{7}{2}}(a + bx) dx$

Optimal. Leaf size=70

$$\frac{10F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{21b} - \frac{10 \cos(a + bx) \sqrt{\sin(a + bx)}}{21b} - \frac{2 \cos(a + bx) \sin^{\frac{5}{2}}(a + bx)}{7b}$$

[Out] -10/21*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticF(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))/b-2/7*cos(b*x+a)*sin(b*x+a)^(5/2)/b-10/21*cos(b*x+a)*sin(b*x+a)^(1/2)/b

Rubi [A]

time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2715, 2720}

$$\frac{10F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{21b} - \frac{2 \sin^{\frac{5}{2}}(a + bx) \cos(a + bx)}{7b} - \frac{10 \sqrt{\sin(a + bx)} \cos(a + bx)}{21b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(7/2), x]

[Out] (10*EllipticF[(a - Pi/2 + b*x)/2, 2])/(21*b) - (10*Cos[a + b*x]*Sqrt[Sin[a + b*x]])/(21*b) - (2*Cos[a + b*x]*Sin[a + b*x]^(5/2))/(7*b)

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Sint[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sint[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sin^{\frac{7}{2}}(a + bx) dx &= -\frac{2 \cos(a + bx) \sin^{\frac{5}{2}}(a + bx)}{7b} + \frac{5}{7} \int \sin^{\frac{3}{2}}(a + bx) dx \\ &= -\frac{10 \cos(a + bx) \sqrt{\sin(a + bx)}}{21b} - \frac{2 \cos(a + bx) \sin^{\frac{5}{2}}(a + bx)}{7b} + \frac{5}{21} \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{10F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{21b} - \frac{10 \cos(a + bx) \sqrt{\sin(a + bx)}}{21b} - \frac{2 \cos(a + bx) \sin^{\frac{5}{2}}(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 55, normalized size = 0.79

$$\frac{-20F\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) + (-23 \cos(a + bx) + 3 \cos(3(a + bx)))\sqrt{\sin(a + bx)}}{42b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^(7/2),x]`

```
[Out] (-20*EllipticF[(-2*a + Pi - 2*b*x)/4, 2] + (-23*Cos[a + b*x] + 3*Cos[3*(a + b*x)])*Sqrt[Sin[a + b*x]])/(42*b)
```

Maple [A]

time = 0.06, size = 104, normalized size = 1.49

method	result
default	$\frac{\frac{2 \sin(bx+a) (\cos^4(bx+a))}{7} + \frac{5 \sqrt{\sin(bx+a)+1} \sqrt{-2 \sin(bx+a)+2} \sqrt{-\sin(bx+a)} \operatorname{EllipticF}\left(\sqrt{\sin(bx+a)}\right)}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^(7/2),x,method=_RETURNVERBOSE)`

```
[Out] (2/7*sin(b*x+a)*cos(b*x+a)^4+5/21*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-16/21*cos(b*x+a)^2*sin(b*x+a))/cos(b*x+a)/sin(b*x+a)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(7/2),x, algorithm="maxima")``[Out] integrate(sin(b*x + a)^(7/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 87, normalized size = 1.24

$$\frac{5 \sqrt{2} \sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + 5 \sqrt{2} \sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)) + 2(3 \cos(bx+a)^3 - 8 \cos(bx+a)) \sqrt{\sin(bx+a)}}{21b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(7/2),x, algorithm="fricas")`

[Out] $\frac{1}{21} \cdot (5\sqrt{2}\sqrt{-1} \operatorname{weierstrassPInverse}(4, 0, \cos(bx + a) + I\sin(bx + a)) + 5\sqrt{2}\sqrt{I} \operatorname{weierstrassPInverse}(4, 0, \cos(bx + a) - I\sin(bx + a)) + 2 \cdot (3\cos(bx + a)^3 - 8\cos(bx + a)) \cdot \sqrt{\sin(bx + a)}) / b$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(7/2), x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3060 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(7/2), x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^(7/2), x)`

Mupad [B]

time = 0.49, size = 42, normalized size = 0.60

$$-\frac{\cos(ax + bx) \sin(ax + bx)^{9/2} {}_2F_1\left(-\frac{5}{4}, \frac{1}{2}; \frac{3}{2}; \cos(ax + bx)^2\right)}{b (\sin(ax + bx)^2)^{9/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^(7/2), x)`

[Out] $-(\cos(a + bx) \sin(a + bx)^{9/2} \operatorname{hypergeom}([-5/4, 1/2], 3/2, \cos(a + bx)^2)) / (b (\sin(a + bx)^2)^{9/4})$

3.18 $\int \sin^{\frac{5}{2}}(a + bx) dx$

Optimal. Leaf size=47

$$\frac{6E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{5b} - \frac{2 \cos(a + bx) \sin^{\frac{3}{2}}(a + bx)}{5b}$$

[Out] $-6/5*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})/b-2/5*\cos(b*x+a)*\sin(b*x+a)^{(3/2)}/b$

Rubi [A]

time = 0.01, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2715, 2719}

$$\frac{6E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{5b} - \frac{2 \sin^{\frac{3}{2}}(a + bx) \cos(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(5/2), x]

[Out] $(6*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2])/(5*b) - (2*\text{Cos}[a + b*x]*\text{Sin}[a + b*x]^{(3/2)})/(5*b)$

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sin^{\frac{5}{2}}(a + bx) dx &= -\frac{2 \cos(a + bx) \sin^{\frac{3}{2}}(a + bx)}{5b} + \frac{3}{5} \int \sqrt{\sin(a + bx)} dx \\ &= \frac{6E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{5b} - \frac{2 \cos(a + bx) \sin^{\frac{3}{2}}(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 44, normalized size = 0.94

$$\frac{6E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) + \sqrt{\sin(a + bx)} \sin(2(a + bx))}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^(5/2), x]``[Out] -1/5*(6*EllipticE[(-2*a + Pi - 2*b*x)/4, 2] + Sqrt[Sin[a + b*x]]*Sin[2*(a + b*x)])/b`**Maple [A]**

time = 0.06, size = 142, normalized size = 3.02

method	result
default	$\frac{\frac{2(\sin^4(bx+a))}{5} - \frac{2(\sin^2(bx+a))}{5} - \frac{6\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}}{5} \text{EllipticE}\left(\sqrt{\sin(bx+a)}\right)}{\cos(bx+a)}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^(5/2), x, method=_RETURNVERBOSE)`

```
[Out] (2/5*sin(b*x+a)^4-2/5*sin(b*x+a)^2-6/5*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticE((sin(b*x+a)+1)^(1/2),1/2*2^(1/2))+3/5*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2),1/2*2^(1/2)))/cos(b*x+a)/sin(b*x+a)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(5/2), x, algorithm="maxima")``[Out] integrate(sin(b*x + a)^(5/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 80, normalized size = 1.70

$$\frac{2 \cos(bx+a) \sin(bx+a)^{\frac{3}{2}} - 3i \sqrt{2} \sqrt{-i} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a))) + 3i \sqrt{2} \sqrt{i} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)))}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(5/2), x, algorithm="fricas")`

[Out] $-1/5*(2*\cos(b*x + a)*\sin(b*x + a)^{(3/2)} - 3*I*\sqrt{2}*\sqrt{-I}*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x + a) + I*\sin(b*x + a))) + 3*I*\sqrt{2}*\sqrt{I}*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x + a) - I*\sin(b*x + a))))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^{\frac{5}{2}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(5/2),x)`

[Out] `Integral(sin(a + b*x)**(5/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(5/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^(5/2), x)`

Mupad [B]

time = 0.45, size = 42, normalized size = 0.89

$$-\frac{\cos(a + bx) \sin(a + bx)^{7/2} {}_2F_1\left(-\frac{3}{4}, \frac{1}{2}; \frac{3}{2}; \cos(a + bx)^2\right)}{b (\sin(a + bx)^2)^{7/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^(5/2),x)`

[Out] `-(cos(a + b*x)*sin(a + b*x)^(7/2)*hypergeom([-3/4, 1/2], 3/2, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(7/4))`

3.19 $\int \sin^{\frac{3}{2}}(a + bx) dx$

Optimal. Leaf size=47

$$\frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{3b} - \frac{2 \cos(a + bx) \sqrt{\sin(a + bx)}}{3b}$$

[Out] $-2/3*(\sin(1/2*a+1/4*\pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*\pi+1/2*b*x)*\text{Elliptic F}(\cos(1/2*a+1/4*\pi+1/2*b*x), 2^{(1/2)})/b-2/3*\cos(b*x+a)*\sin(b*x+a)^{(1/2)}/b$

Rubi [A]

time = 0.01, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2715, 2720}

$$\frac{2F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{3b} - \frac{2 \sqrt{\sin(a + bx)} \cos(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(3/2), x]

[Out] $(2*\text{EllipticF}[(a - \pi/2 + b*x)/2, 2])/(3*b) - (2*\text{Cos}[a + b*x]*\text{Sqrt}[\text{Sin}[a + b*x]])/(3*b)$

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sin^{\frac{3}{2}}(a + bx) dx &= -\frac{2 \cos(a + bx) \sqrt{\sin(a + bx)}}{3b} + \frac{1}{3} \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{3b} - \frac{2 \cos(a + bx) \sqrt{\sin(a + bx)}}{3b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 40, normalized size = 0.85

$$\frac{2\left(F\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) + \cos(a + bx)\sqrt{\sin(a + bx)}\right)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^(3/2), x]`

```
[Out] (-2*(EllipticF[(-2*a + Pi - 2*b*x)/4, 2] + Cos[a + b*x]*Sqrt[Sin[a + b*x]]
)/(3*b)
```

Maple [A]

time = 0.05, size = 88, normalized size = 1.87

method	result
default	$\frac{\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \operatorname{EllipticF}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] (1/3*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*Ellip
ticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-2/3*cos(b*x+a)^2*sin(b*x+a))/cos(b*x
+a)/sin(b*x+a)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(3/2), x, algorithm="maxima")``[Out] integrate(sin(b*x + a)^(3/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 72, normalized size = 1.53

$$\frac{\sqrt{2} \sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + \sqrt{2} \sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)) - 2 \cos(bx+a) \sqrt{\sin(bx+a)}}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(3/2), x, algorithm="fricas")`

[Out] $\frac{1}{3}(\sqrt{2}\sqrt{-1}\text{weierstrassPInverse}(4, 0, \cos(bx + a) + I\sin(bx + a)) + \sqrt{2}\sqrt{1}\text{weierstrassPInverse}(4, 0, \cos(bx + a) - I\sin(bx + a)) - 2\cos(bx + a)\sqrt{\sin(bx + a)})/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^{\frac{3}{2}}(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(3/2),x)`

[Out] `Integral(sin(a + b*x)**(3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(3/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^(3/2), x)`

Mupad [B]

time = 0.44, size = 42, normalized size = 0.89

$$\frac{\cos(a + bx) \sin(a + bx)^{5/2} {}_2F_1\left(-\frac{1}{4}, \frac{1}{2}; \frac{3}{2}; \cos(a + bx)^2\right)}{b (\sin(a + bx)^2)^{5/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^(3/2),x)`

[Out] `-(cos(a + b*x)*sin(a + b*x)^(5/2)*hypergeom([-1/4, 1/2], 3/2, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(5/4))`

3.20 $\int \sqrt{\sin(a + bx)} dx$

Optimal. Leaf size=21

$$\frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{b}$$

[Out] $-2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})/b$

Rubi [A]

time = 0.01, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2719}

$$\frac{2E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sin[a + b*x]],x]

[Out] (2*EllipticE[(a - Pi/2 + b*x)/2, 2])/b

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \sqrt{\sin(a + bx)} dx = \frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{b}$$

Mathematica [A]

time = 0.01, size = 24, normalized size = 1.14

$$-\frac{2E\left(\frac{1}{2}\left(-a + \frac{\pi}{2} - bx\right) \middle| 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sin[a + b*x]],x]

[Out] (-2*EllipticE[(-a + Pi/2 - b*x)/2, 2])/b

Maple [A]

time = 0.09, size = 91, normalized size = 4.33

method	result
default	$\frac{\sqrt{\sin (bx+a)+1} \sqrt{-2 \sin (bx+a)+2} \sqrt{-\sin (bx+a)} \left({}_2\text{EllipticE}\left(\sqrt{\sin (bx+a)+1}, \frac{\sqrt{2}}{2}\right)\right)}{\cos (bx+a) \sqrt{\sin (bx+a)} b}$
risch	$-\frac{i \sqrt{2} \sqrt{-i\left(e^{2i(bx+a)}-1\right)} e^{-i(bx+a)}}{b} + i \left(\frac{2i\left(i-e^{2i(bx+a)}\right)}{\sqrt{e^{i(bx+a)}\left(i-i e^{2i(bx+a)}\right)}} - \frac{\sqrt{e^{i(bx+a)}+1} \sqrt{-2 e^{i(bx+a)}+1}}{\sqrt{e^{i(bx+a)}\left(i-i e^{2i(bx+a)}\right)}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*(2*EllipticE((sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-EllipticF((sin(b*x+a)+1)^(1/2),1/2*2^(1/2)))/cos(b*x+a)/sin(b*x+a)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(sin(b*x + a)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 63, normalized size = 3.00

$$\frac{i \sqrt{2} \sqrt{-i} \text{weierstrassZeta}(4,0, \text{weierstrassPInverse}(4,0, \cos (bx+a)+i \sin (bx+a))) - i \sqrt{2} \sqrt{i} \text{weierstrassZeta}(4,0, \text{weierstrassPInverse}(4,0, \cos (bx+a)-i \sin (bx+a)))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^(1/2),x, algorithm="fricas")
```

```
[Out] (I*sqrt(2)*sqrt(-I)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*sqrt(I)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a))))/b
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\sin (a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**(1/2),x)

[Out] Integral(sqrt(sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(sin(b*x + a)), x)

Mupad [B]

time = 0.37, size = 18, normalized size = 0.86

$$\frac{2 E\left(\frac{a}{2} - \frac{\pi}{4} + \frac{bx}{2} \mid 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^(1/2),x)

[Out] (2*ellipticE(a/2 - pi/4 + (b*x)/2, 2))/b

$$3.21 \quad \int \frac{1}{\sqrt{\sin(a + bx)}} dx$$

Optimal. Leaf size=21

$$\frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right)}{b}$$

[Out] $-2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})/b$

Rubi [A]

time = 0.01, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2720}

$$\frac{2F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \mid 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[Sin[a + b*x]],x]

[Out] (2*EllipticF[(a - Pi/2 + b*x)/2, 2])/b

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{1}{\sqrt{\sin(a + bx)}} dx = \frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right)}{b}$$

Mathematica [A]

time = 0.02, size = 24, normalized size = 1.14

$$-\frac{2F\left(\frac{1}{2}\left(-a + \frac{\pi}{2} - bx\right) \mid 2\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[1/Sqrt[Sin[a + b*x]],x]

[Out] (-2*EllipticF[(-a + Pi/2 - b*x)/2, 2])/b

Maple [A]

time = 0.06, size = 69, normalized size = 3.29

method	result
default	$\frac{\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \operatorname{EllipticF}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx+a) \sqrt{\sin(bx+a)} b}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/sin(b*x+a)^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] (sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF(
(sin(b*x+a)+1)^(1/2),1/2*2^(1/2))/cos(b*x+a)/sin(b*x+a)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(b*x+a)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/sqrt(sin(b*x + a)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 55, normalized size = 2.62

$$\frac{\sqrt{2} \sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + \sqrt{2} \sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(b*x+a)^(1/2),x, algorithm="fricas")
```

```
[Out] (sqrt(2)*sqrt(-I)*weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a))
+ sqrt(2)*sqrt(I)*weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)))
/b
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\sin(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(b*x+a)**(1/2),x)
```

[Out] Integral(1/sqrt(sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(sin(b*x + a)), x)

Mupad [B]

time = 0.38, size = 18, normalized size = 0.86

$$-\frac{2F\left(\frac{\pi}{4} - \frac{a}{2} - \frac{bx}{2} \mid 2\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sin(a + b*x)^(1/2),x)

[Out] -(2*ellipticF(pi/4 - a/2 - (b*x)/2, 2))/b

$$3.22 \quad \int \frac{1}{\sin^{\frac{3}{2}}(a+bx)} dx$$

Optimal. Leaf size=43

$$-\frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{b} - \frac{2\cos(a+bx)}{b\sqrt{\sin(a+bx)}}$$

[Out] $2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})/b-2*\cos(b*x+a)/b/\sin(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2716, 2719}

$$-\frac{2E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{b} - \frac{2\cos(a+bx)}{b\sqrt{\sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(-3/2), x]

[Out] $(-2*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2])/b - (2*\text{Cos}[a + b*x])/(b*\text{Sqrt}[\text{Sin}[a + b*x]])$

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sin^{\frac{3}{2}}(a+bx)} dx &= -\frac{2\cos(a+bx)}{b\sqrt{\sin(a+bx)}} - \int \sqrt{\sin(a+bx)} dx \\ &= -\frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{b} - \frac{2\cos(a+bx)}{b\sqrt{\sin(a+bx)}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 39, normalized size = 0.91

$$\frac{2 \left(E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) - \frac{\cos(a+bx)}{\sqrt{\sin(a+bx)}} \right)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^(-3/2), x]``[Out] (2*(EllipticE[(-2*a + Pi - 2*b*x)/4, 2] - Cos[a + b*x]/Sqrt[Sin[a + b*x]])) /b`**Maple [A]**

time = 0.06, size = 132, normalized size = 3.07

method	result
default	$\frac{2\sqrt{\sin(bx+a)+1}\sqrt{-2\sin(bx+a)+2}\sqrt{-\sin(bx+a)}\operatorname{EllipticE}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) - \cos(bx+a)\sqrt{\sin(bx+a)}}{\cos(bx+a)\sqrt{\sin(bx+a)}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(b*x+a)^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] (2*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticE((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-2*cos(b*x+a)^2)/cos(b*x+a)/sin(b*x+a)^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x+a)^(3/2), x, algorithm="maxima")``[Out] integrate(sin(b*x + a)^(-3/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 99, normalized size = 2.30

$$\frac{-i\sqrt{2}\sqrt{-i}\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\cos(bx+a)+i\sin(bx+a))+i\sqrt{2}\sqrt{i}\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\cos(bx+a)-i\sin(bx+a))-2\cos(bx+a)\sqrt{\sin(bx+a)}}{b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)^(3/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*sqrt(-I)*sin(b*x + a)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a))) + I*sqrt(2)*sqrt(I)*sin(b*x + a)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a))) - 2*cos(b*x + a)*sqrt(sin(b*x + a)))/(b*sin(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sin^{\frac{3}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)**(3/2),x)

[Out] Integral(sin(a + b*x)**(-3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)^(3/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^(-3/2), x)

Mupad [B]

time = 0.51, size = 42, normalized size = 0.98

$$-\frac{\cos(a + bx) (\sin(a + bx)^2)^{1/4} {}_2F_1\left(\frac{1}{2}, \frac{5}{4}; \frac{3}{2}; \cos(a + bx)^2\right)}{b \sqrt{\sin(a + bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sin(a + b*x)^(3/2),x)

[Out] -(cos(a + b*x)*(sin(a + b*x)^2)^(1/4)*hypergeom([1/2, 5/4], 3/2, cos(a + b*x)^2))/(b*sin(a + b*x)^(1/2))

$$3.23 \quad \int \frac{1}{\sin^{\frac{5}{2}}(a+bx)} dx$$

Optimal. Leaf size=47

$$\frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{3b} - \frac{2 \cos(a + bx)}{3b \sin^{\frac{3}{2}}(a + bx)}$$

[Out] -2/3*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticF(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))/b-2/3*cos(b*x+a)/b/sin(b*x+a)^(3/2)

Rubi [A]

time = 0.01, antiderivative size = 47, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2716, 2720}

$$\frac{2F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{3b} - \frac{2 \cos(a + bx)}{3b \sin^{\frac{3}{2}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(-5/2),x]

[Out] (2*EllipticF[(a - Pi/2 + b*x)/2, 2])/(3*b) - (2*Cos[a + b*x])/(3*b*Sin[a + b*x]^(3/2))

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sin^{\frac{5}{2}}(a + bx)} dx &= -\frac{2 \cos(a + bx)}{3b \sin^{\frac{3}{2}}(a + bx)} + \frac{1}{3} \int \frac{1}{\sqrt{\sin(a + bx)}} dx \\ &= \frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{3b} - \frac{2 \cos(a + bx)}{3b \sin^{\frac{3}{2}}(a + bx)} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 43, normalized size = 0.91

$$\frac{2 \left(F\left(\frac{1}{4}(2a - \pi + 2bx) \mid 2\right) - \frac{\cos(a+bx)}{\sin^{\frac{3}{2}}(a+bx)} \right)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^(-5/2), x]``[Out] (2*(EllipticF[(2*a - Pi + 2*b*x)/4, 2] - Cos[a + b*x]/Sin[a + b*x]^(3/2)))/(3*b)`**Maple [A]**

time = 0.06, size = 88, normalized size = 1.87

method	result
default	$\frac{\sqrt{\sin(bx+a)+1} \sqrt{-2\sin(bx+a)+2} \sqrt{-\sin(bx+a)} \operatorname{EllipticF}\left(\sqrt{\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right) \sin(bx+a)}{3 \sin(bx+a)^{\frac{3}{2}} \cos(bx+a)b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(b*x+a)^(5/2), x, method=_RETURNVERBOSE)``[Out] 1/3/sin(b*x+a)^(3/2)*((sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))*sin(b*x+a)-2*cos(b*x+a)^2)/cos(b*x+a)/b`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x+a)^(5/2), x, algorithm="maxima")``[Out] integrate(sin(b*x + a)^(-5/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 115, normalized size = 2.45

$$\frac{\sqrt{-i} \left(\sqrt{2} \cos(bx+a)^2 - \sqrt{2} \right) \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + \sqrt{i} \left(\sqrt{2} \cos(bx+a)^2 - \sqrt{2} \right) \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)) + 2 \cos(bx+a) \sqrt{\sin(bx+a)}}{3 (b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(b*x+a)^(5/2), x, algorithm="fricas")`

[Out] $\frac{1}{3}(\sqrt{-1}(\sqrt{2}\cos(bx + a)^2 - \sqrt{2}))\text{weierstrassPInverse}(4, 0, \cos(bx + a) + I\sin(bx + a)) + \sqrt{I}(\sqrt{2}\cos(bx + a)^2 - \sqrt{2})\text{weierstrassPInverse}(4, 0, \cos(bx + a) - I\sin(bx + a)) + 2\cos(bx + a)\sqrt{\sin(bx + a)}}{(b\cos(bx + a)^2 - b)}$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sin^{\frac{5}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(b*x+a)**(5/2),x)`

[Out] `Integral(sin(a + b*x)**(-5/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(b*x+a)^(5/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^(-5/2), x)`

Mupad [B]

time = 0.61, size = 42, normalized size = 0.89

$$\frac{\cos(a + bx) (\sin(a + bx)^2)^{3/4} {}_2F_1\left(\frac{1}{2}, \frac{7}{4}; \frac{3}{2}; \cos(a + bx)^2\right)}{b \sin(a + bx)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sin(a + b*x)^(5/2),x)`

[Out] `-(cos(a + b*x)*(sin(a + b*x)^2)^(3/4)*hypergeom([1/2, 7/4], 3/2, cos(a + b*x)^2))/(b*sin(a + b*x)^(3/2))`

$$3.24 \quad \int \frac{1}{\sin^2(a+bx)} dx$$

Optimal. Leaf size=70

$$-\frac{6E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right)}{5b} - \frac{2 \cos(a + bx)}{5b \sin^{\frac{5}{2}}(a + bx)} - \frac{6 \cos(a + bx)}{5b \sqrt{\sin(a + bx)}}$$

[Out] 6/5*(sin(1/2*a+1/4*Pi+1/2*b*x)^2)^(1/2)/sin(1/2*a+1/4*Pi+1/2*b*x)*EllipticE(cos(1/2*a+1/4*Pi+1/2*b*x),2^(1/2))/b-2/5*cos(b*x+a)/b/sin(b*x+a)^(5/2)-6/5*cos(b*x+a)/b/sin(b*x+a)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2716, 2719}

$$-\frac{6E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{5b} - \frac{2 \cos(a + bx)}{5b \sin^{\frac{5}{2}}(a + bx)} - \frac{6 \cos(a + bx)}{5b \sqrt{\sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(-7/2),x]

[Out] (-6*EllipticE[(a - Pi/2 + b*x)/2, 2])/(5*b) - (2*Cos[a + b*x])/(5*b*Sin[a + b*x]^(5/2)) - (6*Cos[a + b*x])/(5*b*Sqrt[Sin[a + b*x]])

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sin^{\frac{7}{2}}(a+bx)} dx &= -\frac{2 \cos(a+bx)}{5b \sin^{\frac{5}{2}}(a+bx)} + \frac{3}{5} \int \frac{1}{\sin^{\frac{3}{2}}(a+bx)} dx \\
&= -\frac{2 \cos(a+bx)}{5b \sin^{\frac{5}{2}}(a+bx)} - \frac{6 \cos(a+bx)}{5b \sqrt{\sin(a+bx)}} - \frac{3}{5} \int \sqrt{\sin(a+bx)} dx \\
&= -\frac{6E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right)}{5b} - \frac{2 \cos(a+bx)}{5b \sin^{\frac{5}{2}}(a+bx)} - \frac{6 \cos(a+bx)}{5b \sqrt{\sin(a+bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.21, size = 55, normalized size = 0.79

$$\frac{2 \left(3E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) - \frac{\cos(a+bx)(1+3\sin^2(a+bx))}{\sin^{\frac{5}{2}}(a+bx)} \right)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^(-7/2), x]`

```
[Out] (2*(3*EllipticE[(-2*a + Pi - 2*b*x)/4, 2] - (Cos[a + b*x]*(1 + 3*Sin[a + b*x]^2))/Sin[a + b*x]^(5/2)))/(5*b)
```

Maple [A]

time = 0.06, size = 160, normalized size = 2.29

method	result
default	$ \frac{6 \sqrt{\sin(bx+a)+1} \sqrt{-2 \sin(bx+a)+2} \sqrt{-\sin(bx+a)} (\sin^2(bx+a)) \operatorname{EllipticE}\left(\sqrt{\sin(bx+a)+1}\right)}{5b} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(b*x+a)^(7/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/5/sin(b*x+a)^(5/2)*(6*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*sin(b*x+a)^2*EllipticE((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-3*(sin(b*x+a)+1)^(1/2)*(-2*sin(b*x+a)+2)^(1/2)*(-sin(b*x+a))^(1/2)*sin(b*x+a)^2*EllipticF((sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))+6*sin(b*x+a)^4-4*sin(b*x+a)^2-2)/cos(b*x+a)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)^(7/2),x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^(-7/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 158, normalized size = 2.26

$$\frac{3\sqrt{-1}\left(i\sqrt{2}\cos(bx+a)^2-i\sqrt{2}\right)\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a)))+3\sqrt{1}\left(-i\sqrt{2}\cos(bx+a)^2+i\sqrt{2}\right)\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a)))+2\left(3\cos(bx+a)^3-4\cos(bx+a)\right)\sqrt{\sin(bx+a)}}{5\left(b\cos(bx+a)^2-b\right)\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)^(7/2),x, algorithm="fricas")

[Out] $-1/5*(3*\sqrt{-1}*(I*\sqrt{2}*\cos(b*x + a)^2 - I*\sqrt{2})*\sin(b*x + a)*\operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(b*x + a) + I*\sin(b*x + a))) + 3*\sqrt{1}*(-I*\sqrt{2}*\cos(b*x + a)^2 + I*\sqrt{2})*\sin(b*x + a)*\operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(b*x + a) - I*\sin(b*x + a))) + 2*(3*\cos(b*x + a)^3 - 4*\cos(b*x + a))*\sqrt{\sin(b*x + a)})/((b*\cos(b*x + a)^2 - b)*\sin(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sin^{7/2}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)**(7/2),x)

[Out] Integral(sin(a + b*x)**(-7/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(b*x+a)^(7/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^(-7/2), x)

Mupad [B]

time = 0.59, size = 42, normalized size = 0.60

$$\frac{\cos(a + bx) (\sin(a + bx)^2)^{5/4} {}_2F_1\left(\frac{1}{2}, \frac{9}{4}; \frac{3}{2}; \cos(a + bx)^2\right)}{b \sin(a + bx)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/sin(a + b*x)^(7/2),x)
```

```
[Out] -(cos(a + b*x)*(sin(a + b*x)^2)^(5/4)*hypergeom([1/2, 9/4], 3/2, cos(a + b*x)^2))/(b*sin(a + b*x)^(5/2))
```


3.25 $\int (c \sin(a + bx))^{7/2} dx$

Optimal. Leaf size=103

$$\frac{10c^4 F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a + bx)}}{21b \sqrt{c \sin(a + bx)}} - \frac{10c^3 \cos(a + bx) \sqrt{c \sin(a + bx)}}{21b} - \frac{2c \cos(a + bx) (c \sin(a + bx))^5}{7b}$$

[Out] $-2/7*c*\cos(b*x+a)*(c*\sin(b*x+a))^{(5/2)}/b-10/21*c^4*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\sin(b*x+a)^{(1/2)}/b/(c*\sin(b*x+a))^{(1/2)}-10/21*c^3*\cos(b*x+a)*(c*\sin(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.04, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2720}

$$\frac{10c^4 \sqrt{\sin(a + bx)} F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{21b \sqrt{c \sin(a + bx)}} - \frac{10c^3 \cos(a + bx) \sqrt{c \sin(a + bx)}}{21b} - \frac{2c \cos(a + bx) (c \sin(a + bx))^{5/2}}{7b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(7/2)}, x]$

[Out] $(10*c^4*\text{EllipticF}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[\text{Sin}[a + b*x]])/(21*b*\text{Sqrt}[c*\text{Sin}[a + b*x]]) - (10*c^3*\text{Cos}[a + b*x]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(21*b) - (2*c*\text{Cos}[a + b*x]*(c*\text{Sin}[a + b*x])^{(5/2)})/(7*b)$

Rule 2715

$\text{Int}[(d_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d, x\} \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d, x\}$

Rule 2721

$\text{Int}[(d_*)*\sin[(c_*) + (d_*)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d, x\} \&\& \text{LtQ}[-1, n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
\int (c \sin(a + bx))^{7/2} dx &= -\frac{2c \cos(a + bx)(c \sin(a + bx))^{5/2}}{7b} + \frac{1}{7}(5c^2) \int (c \sin(a + bx))^{3/2} dx \\
&= -\frac{10c^3 \cos(a + bx) \sqrt{c \sin(a + bx)}}{21b} - \frac{2c \cos(a + bx)(c \sin(a + bx))^{5/2}}{7b} + \frac{1}{21}(5c^4) \int (c \sin(a + bx))^{1/2} dx \\
&= -\frac{10c^3 \cos(a + bx) \sqrt{c \sin(a + bx)}}{21b} - \frac{2c \cos(a + bx)(c \sin(a + bx))^{5/2}}{7b} + \frac{(5c^4 \sqrt{\sin(a + bx)})}{21} \\
&= \frac{10c^4 F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{\sin(a + bx)}}{21b \sqrt{c \sin(a + bx)}} - \frac{10c^3 \cos(a + bx) \sqrt{c \sin(a + bx)}}{21b} - \frac{2c \cos(a + bx)(c \sin(a + bx))^{5/2}}{7b}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 80, normalized size = 0.78

$$\frac{c^3 \left(-20F\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) + (-23 \cos(a + bx) + 3 \cos(3(a + bx))) \sqrt{\sin(a + bx)} \right) \sqrt{c \sin(a + bx)}}{42b \sqrt{\sin(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(c*Sin[a + b*x])^(7/2), x]`

```
[Out] (c^3*(-20*EllipticF[(-2*a + Pi - 2*b*x)/4, 2] + (-23*Cos[a + b*x] + 3*Cos[3*(a + b*x)])*Sqrt[Sin[a + b*x]])*Sqrt[c*Sin[a + b*x]])/(42*b*Sqrt[Sin[a + b*x]])
```

Maple [A]

time = 0.06, size = 108, normalized size = 1.05

method	result
default	$ -\frac{c^4 \left(-6(\sin^5(bx+a)) + 5 \sqrt{-\sin(bx+a) + 1} \sqrt{2 \sin(bx+a) + 2} \left(\sqrt{\sin(bx+a)} \right) \text{EllipticF} \left(\sqrt{-\sin(bx+a)} \right) \right)}{21 \cos(bx+a) \sqrt{c \sin(bx+a)} b} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(b*x+a))^(7/2), x, method=_RETURNVERBOSE)`

```
[Out] -1/21*c^4*(-6*sin(b*x+a)^5+5*(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*EllipticF((-sin(b*x+a)+1)^(1/2), 1/2*2^(1/2))-4*sin(b*x+a)^3+10*sin(b*x+a))/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(7/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(7/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 105, normalized size = 1.02

$$\frac{5\sqrt{2}\sqrt{-ic^2}\text{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a))+5\sqrt{2}\sqrt{ic^2}\text{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a))+2(3c^3\cos(bx+a)^3-8c^3\cos(bx+a))\sqrt{c\sin(bx+a)}}{21b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(7/2),x, algorithm="fricas")

[Out] 1/21*(5*sqrt(2)*sqrt(-I*c)*c^3*weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a)) + 5*sqrt(2)*sqrt(I*c)*c^3*weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*(3*c^3*cos(b*x + a)^3 - 8*c^3*cos(b*x + a))*sqrt(c*sin(b*x + a)))/b

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(7/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3877 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(7/2),x)

[Out] int((c*sin(a + b*x))^(7/2), x)

3.26 $\int (c \sin(a + bx))^{5/2} dx$

Optimal. Leaf size=75

$$\frac{6c^2 E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{c \sin(a + bx)}}{5b \sqrt{\sin(a + bx)}} - \frac{2c \cos(a + bx) (c \sin(a + bx))^{3/2}}{5b}$$

[Out] $-2/5*c*\cos(b*x+a)*(c*\sin(b*x+a))^{(3/2)}/b-6/5*c^2*(\sin(1/2*a+1/4*Pi+1/2*b*x))^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*(c*\sin(b*x+a))^{(1/2)}/b/\sin(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2719}

$$\frac{6c^2 E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{c \sin(a + bx)}}{5b \sqrt{\sin(a + bx)}} - \frac{2c \cos(a + bx) (c \sin(a + bx))^{3/2}}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(5/2)}, x]$

[Out] $(6*c^2*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(5*b*\text{Sqrt}[\text{Sin}[a + b*x]]) - (2*c*\text{Cos}[a + b*x]*(c*\text{Sin}[a + b*x])^{(3/2)})/(5*b)$

Rule 2715

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d, x\}$

Rule 2721

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^{(n-1)}/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
\int (c \sin(a + bx))^{5/2} dx &= -\frac{2c \cos(a + bx)(c \sin(a + bx))^{3/2}}{5b} + \frac{1}{5}(3c^2) \int \sqrt{c \sin(a + bx)} dx \\
&= -\frac{2c \cos(a + bx)(c \sin(a + bx))^{3/2}}{5b} + \frac{(3c^2 \sqrt{c \sin(a + bx)}) \int \sqrt{\sin(a + bx)} dx}{5 \sqrt{\sin(a + bx)}} \\
&= \frac{6c^2 E\left(\frac{1}{2}(a - \frac{\pi}{2} + bx) \mid 2\right) \sqrt{c \sin(a + bx)}}{5b \sqrt{\sin(a + bx)}} - \frac{2c \cos(a + bx)(c \sin(a + bx))^{3/2}}{5b}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 66, normalized size = 0.88

$$\frac{(c \sin(a + bx))^{5/2} \left(6E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) + \sqrt{\sin(a + bx)} \sin(2(a + bx))\right)}{5b \sin^{\frac{5}{2}}(a + bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[(c*Sin[a + b*x])^(5/2),x]`

```
[Out] -1/5*((c*Sin[a + b*x])^(5/2)*(6*EllipticE[(-2*a + Pi - 2*b*x)/4, 2] + Sqrt[
Sin[a + b*x]]*Sin[2*(a + b*x)]))/(b*Sin[a + b*x]^(5/2))
```

Maple [A]

time = 0.06, size = 152, normalized size = 2.03

method	result
default	$ -\frac{c^3 \left(6 \sqrt{-\sin(bx + a) + 1} \sqrt{2 \sin(bx + a) + 2} \left(\sqrt{\sin(bx + a)}\right) \text{EllipticE}\left(\sqrt{-\sin(bx + a) + 1}, \frac{\sqrt{2}}{2}\right)\right)}{b} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/5*c^3*(6*(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*E
llipticE((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-3*(-sin(b*x+a)+1)^(1/2)*(2*sin(
b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*EllipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2
)))-2*sin(b*x+a)^4+2*sin(b*x+a)^2)/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(5/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 101, normalized size = 1.35

$$\frac{-2\sqrt{c\sin(bx+a)}c^2\cos(bx+a)\sin(bx+a) - 3i\sqrt{2}\sqrt{-ic}c^2\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a))) + 3i\sqrt{2}\sqrt{ic}c^2\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a)))}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2),x, algorithm="fricas")

[Out]
$$-1/5*(2*\text{sqrt}(c*\sin(b*x + a))*c^2*\cos(b*x + a)*\sin(b*x + a) - 3*I*\text{sqrt}(2)*\text{sqrt}(-I*c)*c^2*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x + a) + I*\sin(b*x + a))) + 3*I*\text{sqrt}(2)*\text{sqrt}(I*c)*c^2*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(b*x + a) - I*\sin(b*x + a))))/b$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(5/2),x)

[Out] Integral((c*sin(a + b*x))**(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(5/2),x)

[Out] int((c*sin(a + b*x))^(5/2), x)

3.27 $\int (c \sin(a + bx))^{3/2} dx$

Optimal. Leaf size=75

$$\frac{2c^2 F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a + bx)}}{3b \sqrt{c \sin(a + bx)}} - \frac{2c \cos(a + bx) \sqrt{c \sin(a + bx)}}{3b}$$

[Out] $-2/3*c^2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\sin(b*x+a)^{(1/2)}/b/(c*\sin(b*x+a))^{(1/2)}-2/3*c*\cos(b*x+a)*(c*\sin(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.02, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 2721, 2720}

$$\frac{2c^2 \sqrt{\sin(a + bx)} F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \middle| 2\right)}{3b \sqrt{c \sin(a + bx)}} - \frac{2c \cos(a + bx) \sqrt{c \sin(a + bx)}}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(3/2)}, x]$

[Out] $(2*c^2*\text{EllipticF}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[\text{Sin}[a + b*x]])/(3*b*\text{Sqrt}[c*\text{Sin}[a + b*x]]) - (2*c*\text{Cos}[a + b*x]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(3*b)$

Rule 2715

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 2721

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int (c \sin(a + bx))^{3/2} dx &= -\frac{2c \cos(a + bx) \sqrt{c \sin(a + bx)}}{3b} + \frac{1}{3} c^2 \int \frac{1}{\sqrt{c \sin(a + bx)}} dx \\
&= -\frac{2c \cos(a + bx) \sqrt{c \sin(a + bx)}}{3b} + \frac{\left(c^2 \sqrt{\sin(a + bx)}\right) \int \frac{1}{\sqrt{\sin(a + bx)}} dx}{3 \sqrt{c \sin(a + bx)}} \\
&= \frac{2c^2 F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{\sin(a + bx)}}{3b \sqrt{c \sin(a + bx)}} - \frac{2c \cos(a + bx) \sqrt{c \sin(a + bx)}}{3b}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 62, normalized size = 0.83

$$-\frac{2\left(F\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) + \cos(a + bx) \sqrt{\sin(a + bx)}\right) (c \sin(a + bx))^{3/2}}{3b \sin^{\frac{3}{2}}(a + bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[(c*Sin[a + b*x])^(3/2),x]`

```
[Out] (-2*(EllipticF[(-2*a + Pi - 2*b*x)/4, 2] + Cos[a + b*x]*Sqrt[Sin[a + b*x]])
*(c*Sin[a + b*x])^(3/2))/(3*b*Sin[a + b*x]^(3/2))
```

Maple [A]

time = 0.07, size = 97, normalized size = 1.29

method	result
default	$ -\frac{c^2 \left(\sqrt{-\sin(bx + a) + 1} \sqrt{2 \sin(bx + a) + 2} \left(\sqrt{\sin(bx + a)} \right) \text{EllipticF} \left(\sqrt{-\sin(bx + a) + 1}, \frac{\sqrt{2}}{2} \right) \right)}{3 \cos(bx + a) \sqrt{c \sin(bx + a)} b} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/3*c^2*((-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*EllipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-2*sin(b*x+a)^3+2*sin(b*x+a))/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.11, size = 81, normalized size = 1.08

$$\frac{\sqrt{2} \sqrt{-i c} \operatorname{cweierstrassPInverse}(4, 0, \cos(bx + a) + i \sin(bx + a)) + \sqrt{2} \sqrt{i c} \operatorname{cweierstrassPInverse}(4, 0, \cos(bx + a) - i \sin(bx + a)) - 2 \sqrt{c \sin(bx + a)} c \cos(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2),x, algorithm="fricas")

[Out] 1/3*(sqrt(2)*sqrt(-I*c)*cweierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a)) + sqrt(2)*sqrt(I*c)*cweierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)) - 2*sqrt(c*sin(b*x + a))*c*cos(b*x + a))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(3/2),x)

[Out] Integral((c*sin(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(3/2),x)

[Out] int((c*sin(a + b*x))^(3/2), x)

3.28 $\int \sqrt{c \sin(a + bx)} dx$

Optimal. Leaf size=43

$$\frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{c \sin(a + bx)}}{b \sqrt{\sin(a + bx)}}$$

[Out] $-2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*(c*\sin(b*x+a))^{(1/2)}/b/\sin(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {2721, 2719}

$$\frac{2E\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \mid 2\right) \sqrt{c \sin(a + bx)}}{b \sqrt{\sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sin[a + b*x]],x]

[Out] $(2*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(b*\text{Sqrt}[\text{Sin}[a + b*x]])$

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sqrt{c \sin(a + bx)} dx &= \frac{\sqrt{c \sin(a + bx)} \int \sqrt{\sin(a + bx)} dx}{\sqrt{\sin(a + bx)}} \\ &= \frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{c \sin(a + bx)}}{b \sqrt{\sin(a + bx)}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 0.98

$$\frac{2E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sqrt{c \sin(a + bx)}}{b \sqrt{\sin(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[c*Sin[a + b*x]],x]``[Out] (-2*EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[c*Sin[a + b*x]])/(b*Sqrt[Sin[a + b*x]])`**Maple [A]**

time = 0.14, size = 98, normalized size = 2.28

method	result
default	$\frac{c \sqrt{-\sin(bx+a)+1} \sqrt{2 \sin(bx+a)+2} \left(\sqrt{\sin(bx+a)}\right) \left(2 \operatorname{EllipticE}\left(\sqrt{-\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)\right)}{\cos(bx+a) \sqrt{c \sin(bx+a)} b}$
risch	$-\frac{i \sqrt{2} \sqrt{-ic(e^{2i(bx+a)} - 1) e^{-i(bx+a)}}}{b} + i \frac{\sqrt{e^{i(bx+a)} + 1} \sqrt{-2e^{i(bx+a)}}}{c \sqrt{e^{i(bx+a)} (-ic e^{2i(bx+a)} + ic)}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)``[Out] -c*(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*(2*EllipticE((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-EllipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2)))/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^(1/2),x, algorithm="maxima")``[Out] integrate(sqrt(c*sin(b*x + a)), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 67, normalized size = 1.56

$$\frac{i \sqrt{2} \sqrt{-ic} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a))) - i \sqrt{2} \sqrt{ic} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] (I*sqrt(2)*sqrt(-I*c)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*sqrt(I*c)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a))))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(1/2),x)

[Out] Integral(sqrt(c*sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a)), x)

Mupad [B]

time = 0.40, size = 36, normalized size = 0.84

$$\frac{2 \sqrt{c \sin(a + bx)} E\left(\frac{a}{2} - \frac{\pi}{4} + \frac{bx}{2} \middle| 2\right)}{b \sqrt{\sin(a + bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/2),x)

[Out] (2*(c*sin(a + b*x))^(1/2)*ellipticE(a/2 - pi/4 + (b*x)/2, 2))/(b*sin(a + b*x)^(1/2))

$$3.29 \quad \int \frac{1}{\sqrt{c \sin(a + bx)}} dx$$

Optimal. Leaf size=43

$$\frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{\sin(a + bx)}}{b \sqrt{c \sin(a + bx)}}$$

[Out] $-2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\sin(b*x+a)^{(1/2)}/b/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$,

Rules used = {2721, 2720}

$$\frac{2 \sqrt{\sin(a + bx)} F\left(\frac{1}{2}\left(a + bx - \frac{\pi}{2}\right) \mid 2\right)}{b \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[c*Sin[a + b*x]],x]

[Out] $(2*\text{EllipticF}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[\text{Sin}[a + b*x]])/(b*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{c \sin(a + bx)}} dx &= \frac{\sqrt{\sin(a + bx)} \int \frac{1}{\sqrt{\sin(a + bx)}} dx}{\sqrt{c \sin(a + bx)}} \\ &= \frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{\sin(a + bx)}}{b \sqrt{c \sin(a + bx)}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 0.98

$$-\frac{2F\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sqrt{\sin(a + bx)}}{b \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/Sqrt[c*Sin[a + b*x]],x]``[Out] (-2*EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]])/(b*Sqrt[c*Sin[a + b*x]])`**Maple [A]**

time = 0.07, size = 74, normalized size = 1.72

method	result
default	$-\frac{\sqrt{-\sin(bx+a)+1} \sqrt{2\sin(bx+a)+2} \left(\sqrt{\sin(bx+a)}\right) \text{EllipticF}\left(\sqrt{-\sin(bx+a)+1}, \frac{\sqrt{2}}{2}\right)}{\cos(bx+a) \sqrt{c \sin(bx+a)} b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)``[Out] -(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*EllipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")``[Out] integrate(1/sqrt(c*sin(b*x + a)), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 62, normalized size = 1.44

$$\frac{\sqrt{2} \sqrt{-ic} \text{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + \sqrt{2} \sqrt{ic} \text{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a))}{bc}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")`

[Out] $(\sqrt{2}*\sqrt{-I*c}*weierstrassPInverse(4, 0, \cos(b*x + a) + I*\sin(b*x + a) + \sqrt{2}*\sqrt{I*c}*weierstrassPInverse(4, 0, \cos(b*x + a) - I*\sin(b*x + a)))/(b*c)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(c*sin(b*x+a))**(1/2),x)`

[Out] `Integral(1/sqrt(c*sin(a + b*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(c*sin(b*x+a))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/sqrt(c*sin(b*x + a)), x)`

Mupad [B]

time = 0.48, size = 36, normalized size = 0.84

$$-\frac{2 \sqrt{\sin(a + bx)} F\left(\frac{\pi}{4} - \frac{a}{2} - \frac{bx}{2} \mid 2\right)}{b \sqrt{c \sin(a + bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(c*sin(a + b*x))^(1/2),x)`

[Out] `-(2*sin(a + b*x)^(1/2)*ellipticF(pi/4 - a/2 - (b*x)/2, 2))/(b*(c*sin(a + b*x))^(1/2))`

$$3.30 \quad \int \frac{1}{(c \sin(a+bx))^{3/2}} dx$$

Optimal. Leaf size=73

$$-\frac{2 \cos(a+bx)}{bc \sqrt{c \sin(a+bx)}} - \frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{c \sin(a+bx)}}{bc^2 \sqrt{\sin(a+bx)}}$$

[Out] $-2*\cos(b*x+a)/b/c/(c*\sin(b*x+a))^{(1/2)}+2*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x),2^{(1/2)})*(c*\sin(b*x+a))^{(1/2)}/b/c^2/\sin(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2719}

$$-\frac{2E\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \mid 2\right) \sqrt{c \sin(a+bx)}}{bc^2 \sqrt{\sin(a+bx)}} - \frac{2 \cos(a+bx)}{bc \sqrt{c \sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] `Int[(c*SIn[a + b*x])^(-3/2),x]`

[Out] $(-2*\text{Cos}[a + b*x])/(b*c*\text{Sqrt}[c*\text{Sin}[a + b*x]]) - (2*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(b*c^2*\text{Sqrt}[\text{Sin}[a + b*x]])$

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*SIn[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*SIn[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*SIn[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(c \sin(a + bx))^{3/2}} dx &= -\frac{2 \cos(a + bx)}{bc \sqrt{c \sin(a + bx)}} - \frac{\int \sqrt{c \sin(a + bx)} dx}{c^2} \\
 &= -\frac{2 \cos(a + bx)}{bc \sqrt{c \sin(a + bx)}} - \frac{\sqrt{c \sin(a + bx)} \int \sqrt{\sin(a + bx)} dx}{c^2 \sqrt{\sin(a + bx)}} \\
 &= -\frac{2 \cos(a + bx)}{bc \sqrt{c \sin(a + bx)}} - \frac{2E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{c \sin(a + bx)}}{bc^2 \sqrt{\sin(a + bx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 54, normalized size = 0.74

$$-\frac{2\left(\cos(a + bx) - E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sqrt{\sin(a + bx)}\right)}{bc \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(c*Sin[a + b*x])^(-3/2),x]`

```
[Out] (-2*(Cos[a + b*x] - EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sqrt[Sin[a + b*x]]
)/(b*c*Sqrt[c*Sin[a + b*x]])
```

Maple [A]

time = 0.06, size = 141, normalized size = 1.93

method	result
default	$ \frac{2 \sqrt{-\sin(bx + a) + 1} \sqrt{2 \sin(bx + a) + 2} \left(\sqrt{\sin(bx + a)} \operatorname{EllipticE}\left(\sqrt{-\sin(bx + a) + 1}, \frac{\sqrt{2}}{2}\right) - \sqrt{\sin(bx + a)} \operatorname{EllipticF}\left(\sqrt{-\sin(bx + a) + 1}, \frac{\sqrt{2}}{2}\right)\right)}{c \cos(bx + a) \sqrt{c \sin(bx + a)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(c*sin(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/c*(2*(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*EllipticE((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(1/2)*EllipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-2*cos(b*x+a)^2/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(-3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 108, normalized size = 1.48

$$\frac{-i\sqrt{2}\sqrt{-ic}\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a)))+i\sqrt{2}\sqrt{ic}\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a)))-2\sqrt{c\sin(bx+a)}\cos(bx+a)}{bc^2\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(3/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*sqrt(-I*c)*sin(b*x + a)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a))) + I*sqrt(2)*sqrt(I*c)*sin(b*x + a)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a))) - 2*sqrt(c*sin(b*x + a))*cos(b*x + a))/(b*c^2*sin(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))**(3/2),x)

[Out] Integral((c*sin(a + b*x))**(-3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \sin(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(a + b*x))^(3/2),x)

[Out] int(1/(c*sin(a + b*x))^(3/2), x)

3.31 $\int \frac{1}{(c \sin(a+bx))^{5/2}} dx$

Optimal. Leaf size=77

$$-\frac{2 \cos(a+bx)}{3bc(c \sin(a+bx))^{3/2}} + \frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a+bx)}}{3bc^2 \sqrt{c \sin(a+bx)}}$$

[Out] $-2/3*\cos(b*x+a)/b/c/(c*\sin(b*x+a))^{(3/2)}-2/3*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticF}(\cos(1/2*a+1/4*Pi+1/2*b*x), 2^{(1/2)})*\sin(b*x+a)^{(1/2)}/b/c^2/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2720}

$$\frac{2 \sqrt{\sin(a+bx)} F\left(\frac{1}{2}\left(a+bx - \frac{\pi}{2}\right) \middle| 2\right)}{3bc^2 \sqrt{c \sin(a+bx)}} - \frac{2 \cos(a+bx)}{3bc(c \sin(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(-5/2)}, x]$

[Out] $(-2*\text{Cos}[a + b*x])/(3*b*c*(c*\text{Sin}[a + b*x])^{(3/2)}) + (2*\text{EllipticF}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[\text{Sin}[a + b*x]])/(3*b*c^2*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2716

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)), x] + \text{Dist}[(n + 2)/(b^2*(n + 1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n + 2)}, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 2721

$\text{Int}[(b_*)*\sin[(c_*) + (d_*)(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^n/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /;$ FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(c \sin(a + bx))^{5/2}} dx &= -\frac{2 \cos(a + bx)}{3bc(c \sin(a + bx))^{3/2}} + \frac{\int \frac{1}{\sqrt{c \sin(a + bx)}} dx}{3c^2} \\
&= -\frac{2 \cos(a + bx)}{3bc(c \sin(a + bx))^{3/2}} + \frac{\sqrt{\sin(a + bx)} \int \frac{1}{\sqrt{\sin(a + bx)}} dx}{3c^2 \sqrt{c \sin(a + bx)}} \\
&= -\frac{2 \cos(a + bx)}{3bc(c \sin(a + bx))^{3/2}} + \frac{2F\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{\sin(a + bx)}}{3bc^2 \sqrt{c \sin(a + bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 55, normalized size = 0.71

$$-\frac{2\left(\cos(a + bx) + F\left(\frac{1}{4}(-2a + \pi - 2bx) \middle| 2\right) \sin^{\frac{3}{2}}(a + bx)\right)}{3bc(c \sin(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[(c*Sin[a + b*x])^(-5/2),x]`

```
[Out] (-2*(Cos[a + b*x] + EllipticF[(-2*a + Pi - 2*b*x)/4, 2]*Sin[a + b*x]^(3/2))
)/(3*b*c*(c*Sin[a + b*x])^(3/2))
```

Maple [A]

time = 0.08, size = 105, normalized size = 1.36

method	result
default	$-\frac{\sqrt{-\sin(bx + a) + 1} \sqrt{2 \sin(bx + a) + 2} \left(\sin^{\frac{5}{2}}(bx + a)\right) \operatorname{EllipticF}\left(\sqrt{-\sin(bx + a) + 1}, \frac{\sqrt{2}}{2}\right) - 2(\sin^3)}{3c^2 \sin(bx + a)^2 \cos(bx + a) \sqrt{c \sin(bx + a)} b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(c*sin(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/3/c^2*((-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(5/2)*Ell
ipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-2*sin(b*x+a)^3+2*sin(b*x+a))/sin(
b*x+a)^2/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(-5/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 127, normalized size = 1.65

$$\frac{(\sqrt{2} \cos(bx+a)^2 - \sqrt{2})\sqrt{-i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) + i \sin(bx+a)) + (\sqrt{2} \cos(bx+a)^2 - \sqrt{2})\sqrt{i} \operatorname{weierstrassPInverse}(4, 0, \cos(bx+a) - i \sin(bx+a)) + 2\sqrt{c \sin(bx+a)} \cos(bx+a)}{3(bc^2 \cos(bx+a)^2 - bc^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(5/2),x, algorithm="fricas")

[Out] 1/3*((sqrt(2)*cos(b*x + a)^2 - sqrt(2))*sqrt(-I*c)*weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a)) + (sqrt(2)*cos(b*x + a)^2 - sqrt(2))*sqrt(I*c)*weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*sqrt(c*sin(b*x + a))*cos(b*x + a)/(b*c^3*cos(b*x + a)^2 - b*c^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(a + bx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(5/2),x)

[Out] Integral((c*sin(a + b*x))^(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(-5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \sin(a + bx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(a + b*x))^(5/2),x)

[Out] int(1/(c*sin(a + b*x))^(5/2), x)

3.32 $\int \frac{1}{(c \sin(a+bx))^{7/2}} dx$

Optimal. Leaf size=105

$$-\frac{2 \cos(a+bx)}{5bc(c \sin(a+bx))^{5/2}} - \frac{6 \cos(a+bx)}{5bc^3 \sqrt{c \sin(a+bx)}} - \frac{6E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \middle| 2\right) \sqrt{c \sin(a+bx)}}{5bc^4 \sqrt{\sin(a+bx)}}$$

[Out] $-2/5*\cos(b*x+a)/b/c/(c*\sin(b*x+a))^{(5/2)}-6/5*\cos(b*x+a)/b/c^3/(c*\sin(b*x+a))^{(1/2)}+6/5*(\sin(1/2*a+1/4*Pi+1/2*b*x)^2)^{(1/2)}/\sin(1/2*a+1/4*Pi+1/2*b*x)*\text{EllipticE}(\cos(1/2*a+1/4*Pi+1/2*b*x),2^{(1/2)})*(c*\sin(b*x+a))^{(1/2)}/b/c^4/\sin(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2716, 2721, 2719}

$$-\frac{6E\left(\frac{1}{2}\left(a+bx-\frac{\pi}{2}\right) \middle| 2\right) \sqrt{c \sin(a+bx)}}{5bc^4 \sqrt{\sin(a+bx)}} - \frac{6 \cos(a+bx)}{5bc^3 \sqrt{c \sin(a+bx)}} - \frac{2 \cos(a+bx)}{5bc(c \sin(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[(c*Sin[a + b*x])^(-7/2),x]`

[Out] $(-2*\text{Cos}[a + b*x])/(5*b*c*(c*\text{Sin}[a + b*x])^{(5/2)}) - (6*\text{Cos}[a + b*x])/(5*b*c^3*\text{Sqrt}[c*\text{Sin}[a + b*x]]) - (6*\text{EllipticE}[(a - \text{Pi}/2 + b*x)/2, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(5*b*c^4*\text{Sqrt}[\text{Sin}[a + b*x]])$

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \int \frac{1}{(c \sin(a + bx))^{7/2}} dx &= -\frac{2 \cos(a + bx)}{5bc(c \sin(a + bx))^{5/2}} + \frac{3 \int \frac{1}{(c \sin(a + bx))^{3/2}} dx}{5c^2} \\
 &= -\frac{2 \cos(a + bx)}{5bc(c \sin(a + bx))^{5/2}} - \frac{6 \cos(a + bx)}{5bc^3 \sqrt{c \sin(a + bx)}} - \frac{3 \int \sqrt{c \sin(a + bx)} dx}{5c^4} \\
 &= -\frac{2 \cos(a + bx)}{5bc(c \sin(a + bx))^{5/2}} - \frac{6 \cos(a + bx)}{5bc^3 \sqrt{c \sin(a + bx)}} - \frac{\left(3 \sqrt{c \sin(a + bx)}\right) \int \sqrt{\sin(a + bx)}}{5c^4 \sqrt{\sin(a + bx)}} \\
 &= -\frac{2 \cos(a + bx)}{5bc(c \sin(a + bx))^{5/2}} - \frac{6 \cos(a + bx)}{5bc^3 \sqrt{c \sin(a + bx)}} - \frac{6E\left(\frac{1}{2}\left(a - \frac{\pi}{2} + bx\right) \mid 2\right) \sqrt{c \sin(a + bx)}}{5bc^4 \sqrt{\sin(a + bx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.12, size = 68, normalized size = 0.65

$$\frac{2 \left(\cot(a + bx) - 3E\left(\frac{1}{4}(-2a + \pi - 2bx) \mid 2\right) \sin^{\frac{3}{2}}(a + bx) + \frac{3}{2} \sin(2(a + bx)) \right)}{5bc^2(c \sin(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(-7/2),x]

[Out] (-2*(Cot[a + b*x] - 3*EllipticE[(-2*a + Pi - 2*b*x)/4, 2]*Sin[a + b*x]^(3/2) + (3*Sin[2*(a + b*x)]/2))/(5*b*c^2*(c*Sin[a + b*x])^(3/2))

Maple [A]

time = 0.06, size = 168, normalized size = 1.60

method	result
default	$ \frac{6 \sqrt{-\sin(bx + a) + 1} \sqrt{2 \sin(bx + a) + 2} \left(\sin^{\frac{7}{2}}(bx + a)\right) \text{EllipticE}\left(\sqrt{-\sin(bx + a) + 1}, \frac{\sqrt{2}}{2}\right) - 3 \sqrt{-\sin(bx + a) + 1}}{5c^3 \sin(bx + a)} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(b*x+a))^(7/2),x,method=_RETURNVERBOSE)

[Out] 1/5/c^3*(6*(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(7/2)*EllipticE((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))-3*(-sin(b*x+a)+1)^(1/2)*(2*sin(b*x+a)+2)^(1/2)*sin(b*x+a)^(7/2)*EllipticF((-sin(b*x+a)+1)^(1/2),1/2*2^(1/2))+6*sin(b*x+a)^5-4*sin(b*x+a)^3-2*sin(b*x+a))/sin(b*x+a)^3/cos(b*x+a)/(c*sin(b*x+a))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(c*sin(b*x+a))^(7/2),x, algorithm="maxima")``[Out] integrate((c*sin(b*x + a))^(-7/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 170, normalized size = 1.62

$$\frac{3(i\sqrt{2}\cos(bx+a)^2 - i\sqrt{2})\sqrt{-i}c\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx+a)+i\sin(bx+a))) + 3(-i\sqrt{2}\cos(bx+a)^2 + i\sqrt{2})\sqrt{i}c\sin(bx+a)\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(bx+a)-i\sin(bx+a))) + 2(3\cos(bx+a)^2 - 4\cos(bx+a))\sqrt{c\sin(bx+a)}}{5(bc^4\cos(bx+a)^2 - bc^4)\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(c*sin(b*x+a))^(7/2),x, algorithm="fricas")`

```
[Out] -1/5*(3*(I*sqrt(2)*cos(b*x + a)^2 - I*sqrt(2))*sqrt(-I*c)*sin(b*x + a)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) + I*sin(b*x + a))) + 3*(-I*sqrt(2)*cos(b*x + a)^2 + I*sqrt(2))*sqrt(I*c)*sin(b*x + a)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(b*x + a) - I*sin(b*x + a))) + 2*(3*cos(b*x + a)^3 - 4*cos(b*x + a))*sqrt(c*sin(b*x + a)))/((b*c^4*cos(b*x + a)^2 - b*c^4)*sin(b*x + a))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(a + bx))^{\frac{7}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(c*sin(b*x+a))**(7/2),x)``[Out] Integral((c*sin(a + b*x))**(-7/2), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(c*sin(b*x+a))^(7/2),x, algorithm="giac")``[Out] integrate((c*sin(b*x + a))^(-7/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(c \sin(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(a + b*x))^(7/2),x)

[Out] int(1/(c*sin(a + b*x))^(7/2), x)

3.33 $\int (c \sin(a + bx))^{4/3} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \sin^2(a + bx)\right) (c \sin(a + bx))^{7/3}}{7bc \sqrt{\cos^2(a + bx)}}$$

[Out] 3/7*cos(b*x+a)*hypergeom([1/2, 7/6], [13/6], sin(b*x+a)^2)*(c*sin(b*x+a))^(7/3)/b/c/(cos(b*x+a)^2)^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(a + bx)(c \sin(a + bx))^{7/3} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \sin^2(a + bx)\right)}{7bc \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(4/3), x]

[Out] (3*Cos[a + b*x]*Hypergeometric2F1[1/2, 7/6, 13/6, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(7/3))/(7*b*c*Sqrt[Cos[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (c \sin(a + bx))^{4/3} dx = \frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \sin^2(a + bx)\right) (c \sin(a + bx))^{7/3}}{7bc \sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{7}{6}; \frac{13}{6}; \sin^2(a + bx)\right) (c \sin(a + bx))^{4/3} \tan(a + bx)}{7b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(4/3),x]

[Out] (3*sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/2, 7/6, 13/6, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(4/3)*Tan[a + b*x])/(7*b)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int (c \sin(bx + a))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(4/3),x)

[Out] int((c*sin(b*x+a))^(4/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(4/3),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(4/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(4/3),x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^(1/3)*c*sin(b*x + a), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^{\frac{4}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(4/3),x)

[Out] Integral((c*sin(a + b*x))**(4/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(4/3),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(4/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int (c \sin(a + b x))^{4/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(4/3),x)

[Out] int((c*sin(a + b*x))^(4/3), x)

3.34 $\int (c \sin(a + bx))^{2/3} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \sin^2(a + bx)\right) (c \sin(a + bx))^{5/3}}{5bc \sqrt{\cos^2(a + bx)}}$$

[Out] 3/5*cos(b*x+a)*hypergeom([1/2, 5/6],[11/6],sin(b*x+a)^2)*(c*sin(b*x+a))^(5/3)/b/c/(cos(b*x+a)^2)^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(a + bx)(c \sin(a + bx))^{5/3} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \sin^2(a + bx)\right)}{5bc \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*SIn[a + b*x])^(2/3),x]

[Out] (3*Cos[a + b*x]*Hypergeometric2F1[1/2, 5/6, 11/6, Sin[a + b*x]^2]*(c*SIn[a + b*x])^(5/3))/(5*b*c*Sqrt[Cos[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*SIn[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (c \sin(a + bx))^{2/3} dx = \frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \sin^2(a + bx)\right) (c \sin(a + bx))^{5/3}}{5bc \sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{5}{6}; \frac{11}{6}; \sin^2(a + bx)\right) (c \sin(a + bx))^{2/3} \tan(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(2/3),x]

[Out] (3*Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/2, 5/6, 11/6, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(2/3)*Tan[a + b*x])/(5*b)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int (c \sin(bx + a))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(2/3),x)

[Out] int((c*sin(b*x+a))^(2/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(2/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(2/3),x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^(2/3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(2/3),x)

[Out] Integral((c*sin(a + b*x))**(2/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^(2/3),x, algorithm="giac")``[Out] integrate((c*sin(b*x + a))^(2/3), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int (c \sin(a + bx))^{2/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(a + b*x))^(2/3),x)``[Out] int((c*sin(a + b*x))^(2/3), x)`

3.35 $\int \sqrt[3]{c \sin(a + bx)} dx$

Optimal. Leaf size=517

$$\sqrt[3]{\frac{3}{2}(3 - i\sqrt{3})} \sqrt[3]{c} E \left(\sin^{-1} \left(\frac{\sqrt{2} \sqrt{1 - \frac{(c \sin(a + bx))^{2/3}}{c^{2/3}}}}{\sqrt{3 + i\sqrt{3}}} \right) \middle| \frac{3i - \sqrt{3}}{3i + \sqrt{3}} \right) \sec(a + bx) \sqrt{1 - \frac{(c \sin(a + bx))^{2/3}}{c^{2/3}}}$$

b

[Out] $\frac{3}{4} c^{1/3} \text{EllipticF}\left(2^{1/2} \sqrt{1 - (c \sin(bx+a))^{2/3}/c^{2/3}}\right)^{1/2} / (3 - I \cdot 3^{1/2})^{1/2}, ((3I + 3^{1/2}) / (3I - 3^{1/2}))^{1/2} \sec(bx+a) (1 - I \cdot 3^{1/2}) \cdot (1 - (c \sin(bx+a))^{2/3}/c^{2/3})^{1/2} \cdot ((I - 3^{1/2}) / (3I - 3^{1/2}) + 2 \cdot (c \sin(bx+a))^{2/3}/c^{2/3} / (3I + 3^{1/2}))^{1/2} \cdot (3 - I \cdot 3^{1/2})^{1/2} \cdot (2 \cdot (c \sin(bx+a))^{2/3}/c^{2/3} / (3 - I \cdot 3^{1/2}) + (3^{1/2} + I) / (3I + 3^{1/2}))^{1/2} / b \cdot 2^{1/2} - 3/2 \cdot c^{1/3} \cdot \text{EllipticE}\left(2^{1/2} \sqrt{1 - (c \sin(bx+a))^{2/3}/c^{2/3}}\right)^{1/2} / (3 + I \cdot 3^{1/2})^{1/2}, ((3I - 3^{1/2}) / (3I + 3^{1/2}))^{1/2} \sec(bx+a) (1 - (c \sin(bx+a))^{2/3}/c^{2/3})^{1/2} \cdot ((I - 3^{1/2}) / (3I - 3^{1/2}) + 2 \cdot (c \sin(bx+a))^{2/3}/c^{2/3} / (3I + 3^{1/2}))^{1/2} \cdot (18 - 6I \cdot 3^{1/2})^{1/2} \cdot (2 \cdot (c \sin(bx+a))^{2/3}/c^{2/3} / (3 - I \cdot 3^{1/2}) + (3^{1/2} + I) / (3I + 3^{1/2}))^{1/2} / b$

Rubi [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 0.11, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(a + bx) (c \sin(a + bx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(a + bx)\right)}{4bc \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(1/3),x]

[Out] (3*Cos[a + b*x]*Hypergeometric2F1[1/2, 2/3, 5/3, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(4/3))/(4*b*c*Sqrt[Cos[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \sqrt[3]{c \sin(a + bx)} dx = \frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(a + bx)\right) (c \sin(a + bx))^{4/3}}{4bc \sqrt{\cos^2(a + bx)}}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.03, size = 55, normalized size = 0.11

$$\frac{3 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(a + bx)\right) \sqrt[3]{c \sin(a + bx)} \tan(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(1/3),x]

[Out] (3*sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/2, 2/3, 5/3, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1/3)*Tan[a + b*x])/(4*b)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int (c \sin(bx + a))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(1/3),x)

[Out] int((c*sin(b*x+a))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/3),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/3),x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^(1/3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{c \sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(1/3),x)

[Out] Integral((c*sin(a + b*x))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/3),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (c \sin(a + bx))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/3),x)

[Out] int((c*sin(a + b*x))^(1/3), x)

$$3.36 \quad \int \frac{1}{\sqrt[3]{c \sin(a + bx)}} dx$$

Optimal. Leaf size=252

$$3\sqrt{3 - i\sqrt{3}} F\left(\sin^{-1}\left(\frac{\sqrt{2}\sqrt{1 - \frac{(c \sin(a + bx))^{2/3}}{c^{2/3}}}}{\sqrt{3 - i\sqrt{3}}}\right) \middle| \frac{3i + \sqrt{3}}{3i - \sqrt{3}}\right) \sec(a + bx) \sqrt{1 - \frac{(c \sin(a + bx))^{2/3}}{c^{2/3}}}$$

$$\sqrt{2} b \sqrt[3]{c}$$

[Out] $-3/2 * \text{EllipticF}(2^{(1/2)} * (1 - (c * \sin(b * x + a))^{(2/3)} / c^{(2/3)})^{(1/2)} / (3 - I * 3^{(1/2)})^{(1/2)}, ((3 * I + 3^{(1/2)}) / (3 * I - 3^{(1/2)}))^{(1/2)}) * \sec(b * x + a) * (1 - (c * \sin(b * x + a))^{(2/3)} / c^{(2/3)})^{(1/2)} * ((1 - 3^{(1/2)}) / (3 * I - 3^{(1/2)}) + 2 * (c * \sin(b * x + a))^{(2/3)} / c^{(2/3)}) / (3 + I * 3^{(1/2)})^{(1/2)} * (3 - I * 3^{(1/2)})^{(1/2)} * (2 * (c * \sin(b * x + a))^{(2/3)} / c^{(2/3)} / (3 - I * 3^{(1/2)}) + (3^{(1/2)} + I) / (3 * I + 3^{(1/2)}))^{(1/2)} / b / c^{(1/3)} * 2^{(1/2)}$

Rubi [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 0.23, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(a + bx) (c \sin(a + bx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(a + bx)\right)}{2bc \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c * \text{Sin}[a + b * x])^{(-1/3)}, x]$

[Out] $(3 * \text{Cos}[a + b * x] * \text{Hypergeometric2F1}[1/3, 1/2, 4/3, \text{Sin}[a + b * x]^2] * (c * \text{Sin}[a + b * x])^{(2/3)}) / (2 * b * c * \text{Sqrt}[\text{Cos}[a + b * x]^2])$

Rule 2722

$\text{Int}[(b * \sin(c + d * x) + (d * x))^{(n)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d * x] * ((b * \text{Sin}[c + d * x])^{(n + 1)} / (b * d * (n + 1) * \text{Sqrt}[\text{Cos}[c + d * x]^2])) * \text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d * x]^2, x] /;$ $\text{FreeQ}\{b, c, d, n\}, x$ && $! \text{IntegerQ}[2 * n]$

Rubi steps

$$\int \frac{1}{\sqrt[3]{c \sin(a + bx)}} dx = \frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(a + bx)\right) (c \sin(a + bx))^{2/3}}{2bc \sqrt{\cos^2(a + bx)}}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.03, size = 55, normalized size = 0.22

$$\frac{3\sqrt{\cos^2(a+bx)} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(a+bx)\right) \tan(a+bx)}{2b^3\sqrt[3]{c\sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(-1/3), x]

[Out] (3*Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/3, 1/2, 4/3, Sin[a + b*x]^2]*Tan[a + b*x])/(2*b*(c*Sin[a + b*x])^(1/3))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(bx + a))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(b*x+a))^(1/3), x)

[Out] int(1/(c*sin(b*x+a))^(1/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(1/3), x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(1/3), x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^(2/3)/(c*sin(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(c*sin(b*x+a))**(1/3),x)`

[Out] `Integral((c*sin(a + b*x))**(-1/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(c*sin(b*x+a))^(1/3),x, algorithm="giac")`

[Out] `integrate((c*sin(b*x + a))^(1/3), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(c \sin(a + bx))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(c*sin(a + b*x))^(1/3),x)`

[Out] `int(1/(c*sin(a + b*x))^(1/3), x)`

$$3.37 \quad \int \frac{1}{(c \sin(a+bx))^{2/3}} dx$$

Optimal. Leaf size=271

$$3^{3/4} F\left(\cos^{-1}\left(\frac{c^{2/3} - (1 - \sqrt{3})(c \sin(a+bx))^{2/3}}{c^{2/3} - (1 + \sqrt{3})(c \sin(a+bx))^{2/3}}\right) \middle| \frac{1}{4}(2 + \sqrt{3})\right) \sec(a+bx) \sqrt[3]{c \sin(a+bx)} (c^{2/3} - (c \sin(a+bx))^{2/3})$$

$$2bc^{5/3} \sqrt{\frac{(c \sin(a+bx))^{2/3} (c^{2/3} - (c \sin(a+bx))^{2/3})}{(c^{2/3} - (1 + \sqrt{3})(c \sin(a+bx))^{2/3})^2}}$$

[Out] $1/2*3^{(3/4)}*((c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1-3^{(1/2))))^{2/(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1+3^{(1/2))))^{2/(1/2)/(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1-3^{(1/2))})*(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1+3^{(1/2)))}}*EllipticF((1-(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1-3^{(1/2))))^{2/(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1+3^{(1/2)))})^{2/(1/2)},1/4*6^{(1/2)}+1/4*2^{(1/2)})*\sec(b*x+a)*(c*\sin(b*x+a))^{(1/3)}*(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)})*(c^{(4/3)}*(1+(c*\sin(b*x+a))^{(2/3)}/c^{(2/3)}+(c*\sin(b*x+a))^{(4/3)}/c^{(4/3)))/(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1+3^{(1/2))))^{2/(1/2)}/b/c^{(5/3)}/(-(c*\sin(b*x+a))^{(2/3)}*(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)))/(c^{(2/3)}-(c*\sin(b*x+a))^{(2/3)}*(1+3^{(1/2))))^{2/(1/2)}$

Rubi [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.01, antiderivative size = 56, normalized size of antiderivative = 0.21, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$,

Rules used = {2722}

$$\frac{3 \cos(a+bx) \sqrt[3]{c \sin(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}, \frac{7}{6}; \sin^2(a+bx)\right)}{bc \sqrt{\cos^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(-2/3),x]

[Out] (3*Cos[a + b*x]*Hypergeometric2F1[1/6, 1/2, 7/6, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1/3))/(b*c*Sqrt[Cos[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{(c \sin(a + bx))^{2/3}} dx = \frac{3 \cos(a + bx) {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \sin^2(a + bx)\right) \sqrt[3]{c \sin(a + bx)}}{bc \sqrt{\cos^2(a + bx)}}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.03, size = 53, normalized size = 0.20

$$\frac{3 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{2}; \frac{7}{6}; \sin^2(a + bx)\right) \tan(a + bx)}{b(c \sin(a + bx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(-2/3),x]

[Out] (3*sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/6, 1/2, 7/6, Sin[a + b*x]^2]*Tan[a + b*x])/(b*(c*Sin[a + b*x])^(2/3))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(bx + a))^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(b*x+a))^(2/3),x)

[Out] int(1/(c*sin(b*x+a))^(2/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(2/3),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(2/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(2/3),x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^(1/3)/(c*sin(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(a + bx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))**(2/3),x)

[Out] Integral((c*sin(a + b*x))**(-2/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(2/3),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(2/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(c \sin(a + bx))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(a + b*x))^(2/3),x)

[Out] int(1/(c*sin(a + b*x))^(2/3), x)

$$3.38 \quad \int \frac{1}{(c \sin(a+bx))^{4/3}} dx$$

Optimal. Leaf size=56

$$-\frac{3 \cos(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sin^2(a+bx)\right)}{bc \sqrt{\cos^2(a+bx)} \sqrt[3]{c \sin(a+bx)}}$$

[Out] $-3*\cos(b*x+a)*\text{hypergeom}([-1/6, 1/2], [5/6], \sin(b*x+a)^2)/b/c/(c*\sin(b*x+a))^{(1/3)}/(\cos(b*x+a)^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 56, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$-\frac{3 \cos(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sin^2(a+bx)\right)}{bc \sqrt{\cos^2(a+bx)} \sqrt[3]{c \sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(-4/3)}, x]$

[Out] $(-3*\text{Cos}[a + b*x]*\text{Hypergeometric2F1}[-1/6, 1/2, 5/6, \text{Sin}[a + b*x]^2])/(b*c*\text{Sqrt}[\text{Cos}[a + b*x]^2]*(c*\text{Sin}[a + b*x])^{(1/3)})$

Rule 2722

$\text{Int}[(b*.)*\sin[(c*.) + (d*.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n + 1)}/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2]))*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2], x] /; \text{FreeQ}\{b, c, d, n\}, x \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\int \frac{1}{(c \sin(a+bx))^{4/3}} dx = -\frac{3 \cos(a+bx) {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sin^2(a+bx)\right)}{bc \sqrt{\cos^2(a+bx)} \sqrt[3]{c \sin(a+bx)}}$$

Mathematica [A]

time = 0.03, size = 53, normalized size = 0.95

$$-\frac{3 \sqrt{\cos^2(a+bx)} {}_2F_1\left(-\frac{1}{6}, \frac{1}{2}; \frac{5}{6}; \sin^2(a+bx)\right) \tan(a+bx)}{b(c \sin(a+bx))^{4/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(-4/3),x]

[Out] (-3*Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[-1/6, 1/2, 5/6, Sin[a + b*x]^2]*Tan[a + b*x])/(b*(c*Sin[a + b*x])^(4/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(bx + a))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(b*x+a))^(4/3),x)

[Out] int(1/(c*sin(b*x+a))^(4/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(4/3),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(-4/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(4/3),x, algorithm="fricas")

[Out] integral(-(c*sin(b*x + a))^(2/3)/(c^2*cos(b*x + a)^2 - c^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(c \sin(a + bx))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))**(4/3),x)

[Out] Integral((c*sin(a + b*x))**(-4/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(c*sin(b*x+a))^(4/3),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(-4/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(c \sin(a + bx))^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(c*sin(a + b*x))^(4/3),x)

[Out] int(1/(c*sin(a + b*x))^(4/3), x)

3.39 $\int \sin^n(a + bx) dx$

Optimal. Leaf size=63

$$\frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \sin^2(a + bx)\right) \sin^{1+n}(a + bx)}{b(1+n) \sqrt{\cos^2(a + bx)}}$$

[Out] cos(b*x+a)*hypergeom([1/2, 1/2+1/2*n], [3/2+1/2*n], sin(b*x+a)^2)*sin(b*x+a)^(1+n)/b/(1+n)/(cos(b*x+a)^2)^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {2722}

$$\frac{\cos(a + bx) \sin^{n+1}(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \sin^2(a + bx)\right)}{b(n+1) \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^n,x]

[Out] (Cos[a + b*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Sin[a + b*x]^2]*Sin[a + b*x]^(1 + n))/(b*(1 + n)*Sqrt[Cos[a + b*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \sin^n(a + bx) dx = \frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \sin^2(a + bx)\right) \sin^{1+n}(a + bx)}{b(1+n) \sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.03, size = 63, normalized size = 1.00

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \sin^2(a + bx)\right) \sec(a + bx) \sin^{1+n}(a + bx)}{b(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^n,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Sin[a + b*x]^2]*Sec[a + b*x]*Sin[a + b*x]^(1 + n))/(b*(1 + n))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \sin^n (bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^n,x)

[Out] int(sin(b*x+a)^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^n,x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^n,x, algorithm="fricas")

[Out] integral(sin(b*x + a)^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^n (a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**n,x)

[Out] Integral(sin(a + b*x)**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^n,x, algorithm="giac")

[Out] integrate(sin(b*x + a)^n, x)

Mupad [B]

time = 0.78, size = 54, normalized size = 0.86

$$\frac{\cos(a + bx) \sin(a + bx)^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{1}{2} - \frac{n}{2}; \frac{3}{2}; \cos(a + bx)^2\right)}{b (\sin(a + bx)^2)^{\frac{n}{2} + \frac{1}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^n,x)

[Out] -(cos(a + b*x)*sin(a + b*x)^(n + 1)*hypergeom([1/2, 1/2 - n/2], 3/2, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(n/2 + 1/2))

3.40 $\int (c \sin(a + bx))^n dx$

Optimal. Leaf size=68

$$\frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+n}}{bc(1+n) \sqrt{\cos^2(a + bx)}}$$

[Out] $\cos(b*x+a)*\text{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \sin(b*x+a)^2)*(c*\sin(b*x+a))^{(1+n)}/b/c/(1+n)/(\cos(b*x+a)^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\frac{\cos(a + bx)(c \sin(a + bx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \sin^2(a + bx)\right)}{bc(n+1) \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^n, x]$

[Out] $(\text{Cos}[a + b*x]*\text{Hypergeometric2F1}[1/2, (1 + n)/2, (3 + n)/2, \text{Sin}[a + b*x]^2]*(c*\text{Sin}[a + b*x])^{(1 + n)})/(b*c*(1 + n)*\text{Sqrt}[\text{Cos}[a + b*x]^2])$

Rule 2722

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n + 1)})/(b*d*(n + 1)*\text{Sqrt}[\text{Cos}[c + d*x]^2])]*\text{Hypergeometric2F1}[1/2, (n + 1)/2, (n + 3)/2, \text{Sin}[c + d*x]^2], x] /; \text{FreeQ}\{b, c, d, n\}, x] \&\& \text{!IntegerQ}[2*n]$

Rubi steps

$$\int (c \sin(a + bx))^n dx = \frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+n}}{bc(1+n) \sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.03, size = 63, normalized size = 0.93

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^n \tan(a + bx)}{b(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^n,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^n*Tan[a + b*x])/(b*(1 + n))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int (c \sin(bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^n,x)

[Out] int((c*sin(b*x+a))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^n,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^n,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^n,x)

[Out] Integral((c*sin(a + b*x))^n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^n,x, algorithm="giac")``[Out] integrate((c*sin(b*x + a))^n, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(a + b*x))^n,x)``[Out] int((c*sin(a + b*x))^n, x)`

3.41 $\int (a \sin(e + fx))^m (b \sin(e + fx))^n dx$

Optimal. Leaf size=81

$$\frac{\cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m + n); \frac{1}{2}(3 + m + n); \sin^2(e + fx)\right) (a \sin(e + fx))^{1+m} (b \sin(e + fx))^n}{af(1 + m + n) \sqrt{\cos^2(e + fx)}}$$

[Out] cos(f*x+e)*hypergeom([1/2, 1/2+1/2*m+1/2*n], [3/2+1/2*m+1/2*n], sin(f*x+e)^2) *(a*sin(f*x+e))^(1+m)*(b*sin(f*x+e))^n/a/f/(1+m+n)/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {20, 2722}

$$\frac{\cos(e + fx)(a \sin(e + fx))^{m+1} (b \sin(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m + n + 1); \frac{1}{2}(m + n + 3); \sin^2(e + fx)\right)}{af(m + n + 1) \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sin[e + f*x])^m*(b*Sin[e + f*x])^n,x]

[Out] (Cos[e + f*x]*Hypergeometric2F1[1/2, (1 + m + n)/2, (3 + m + n)/2, Sin[e + f*x]^2]*(a*Sin[e + f*x])^(1 + m)*(b*Sin[e + f*x])^n)/(a*f*(1 + m + n)*Sqrt[Cos[e + f*x]^2])

Rule 20

Int[(u_.)*((a_.)*(v_))^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[b^IntPart[n]*((b*v)^FracPart[n]/(a^IntPart[n]*(a*v)^FracPart[n])), Int[u*(a*v)^(m + n), x], x] /; FreeQ[{a, b, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && !IntegerQ[m + n]

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int (a \sin(e + fx))^m (b \sin(e + fx))^n dx &= ((a \sin(e + fx))^{-n} (b \sin(e + fx))^n) \int (a \sin(e + fx))^{m+n} dx \\ &= \frac{\cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m + n); \frac{1}{2}(3 + m + n); \sin^2(e + fx)\right) (a \sin(e + fx))^{1+m} (b \sin(e + fx))^n}{af(1 + m + n) \sqrt{\cos^2(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 76, normalized size = 0.94

$$\frac{\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m + n); \frac{1}{2}(3 + m + n); \sin^2(e + fx)\right) (a \sin(e + fx))^m (b \sin(e + fx))^n \tan(e + fx)}{f(1 + m + n)}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sin[e + f*x])^m*(b*Sin[e + f*x])^n,x]

[Out] (Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/2, (1 + m + n)/2, (3 + m + n)/2, Sin[e + f*x]^2]*(a*Sin[e + f*x])^m*(b*Sin[e + f*x])^n*Tan[e + f*x])/(f*(1 + m + n))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int (a \sin(fx + e))^m (b \sin(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^m*(b*sin(f*x+e))^n,x)

[Out] int((a*sin(f*x+e))^m*(b*sin(f*x+e))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^m*(b*sin(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((a*sin(f*x + e))^m*(b*sin(f*x + e))^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^m*(b*sin(f*x+e))^n,x, algorithm="fricas")

[Out] integral((a*sin(f*x + e))^m*(b*sin(f*x + e))^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sin(e + fx))^m (b \sin(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))**m*(b*sin(f*x+e))**n,x)

[Out] Integral((a*sin(e + f*x))**m*(b*sin(e + f*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^m*(b*sin(f*x+e))^n,x, algorithm="giac")

[Out] integrate((a*sin(f*x + e))^m*(b*sin(f*x + e))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \sin(e + f x))^m (b \sin(e + f x))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^m*(b*sin(e + f*x))^n,x)

[Out] int((a*sin(e + f*x))^m*(b*sin(e + f*x))^n, x)

3.42 $\int \cos^3(a + bx) \sin(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\cos^4(a + bx)}{4b}$$

[Out] -1/4*cos(b*x+a)^4/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2645, 30}

$$-\frac{\cos^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*Sin[a + b*x],x]

[Out] -1/4*Cos[a + b*x]^4/b

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \sin(a + bx) dx &= -\frac{\text{Subst}\left(\int x^3 dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$-\frac{\cos^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^3*Sin[a + b*x],x]

[Out] -1/4*Cos[a + b*x]^4/b

Maple [A]

time = 0.04, size = 14, normalized size = 0.93

method	result	size
derivativedivides	$-\frac{\cos^4(bx+a)}{4b}$	14
default	$-\frac{\cos^4(bx+a)}{4b}$	14
risch	$-\frac{\cos(4bx+4a)}{32b} - \frac{\cos(2bx+2a)}{8b}$	30
norman	$\frac{2\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{2\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^4}$	50

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] -1/4*cos(b*x+a)^4/b

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*sin(b*x+a),x, algorithm="maxima")

[Out] -1/4*cos(b*x + a)^4/b

Fricas [A]

time = 0.41, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*sin(b*x+a),x, algorithm="fricas")

[Out] -1/4*cos(b*x + a)^4/b

Sympy [A]

time = 0.15, size = 22, normalized size = 1.47

$$\begin{cases} -\frac{\cos^4(a+bx)}{4b} & \text{for } b \neq 0 \\ x \sin(a) \cos^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)**3*sin(b*x+a),x)``[Out] Piecewise((-cos(a + b*x)**4/(4*b), Ne(b, 0)), (x*sin(a)*cos(a)**3, True))`**Giac [A]**

time = 7.05, size = 24, normalized size = 1.60

$$-\frac{\sin(bx+a)^4 - 2\sin(bx+a)^2}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^3*sin(b*x+a),x, algorithm="giac")``[Out] -1/4*(sin(b*x + a)^4 - 2*sin(b*x + a)^2)/b`**Mupad [B]**

time = 0.06, size = 13, normalized size = 0.87

$$-\frac{\cos(a+bx)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)^3*sin(a + b*x),x)``[Out] -cos(a + b*x)^4/(4*b)`

3.43 $\int \cos^2(a + bx) \sin(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\cos^3(a + bx)}{3b}$$

[Out] -1/3*cos(b*x+a)^3/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2645, 30}

$$-\frac{\cos^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Sin[a + b*x],x]

[Out] -1/3*Cos[a + b*x]^3/b

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2645

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) \sin(a + bx) dx &= -\frac{\text{Subst}(\int x^2 dx, x, \cos(a + bx))}{b} \\ &= -\frac{\cos^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$-\frac{\cos^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^2*Sin[a + b*x],x]

[Out] -1/3*Cos[a + b*x]^3/b

Maple [A]

time = 0.03, size = 14, normalized size = 0.93

method	result	size
derivativedivides	$-\frac{\cos^3(bx+a)}{3b}$	14
default	$-\frac{\cos^3(bx+a)}{3b}$	14
risch	$-\frac{\cos(bx+a)}{4b} - \frac{\cos(3bx+3a)}{12b}$	27
norman	$-\frac{2(\tan^4(\frac{bx}{2} + \frac{a}{2})) - \frac{2}{3b}}{(1+\tan^2(\frac{bx}{2} + \frac{a}{2}))^3}$	39

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] -1/3*cos(b*x+a)^3/b

Maxima [A]

time = 0.28, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a),x, algorithm="maxima")

[Out] -1/3*cos(b*x + a)^3/b

Fricas [A]

time = 0.37, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a),x, algorithm="fricas")

[Out] -1/3*cos(b*x + a)^3/b

Sympy [A]

time = 0.09, size = 22, normalized size = 1.47

$$\begin{cases} -\frac{\cos^3(a+bx)}{3b} & \text{for } b \neq 0 \\ x \sin(a) \cos^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2*sin(b*x+a),x)

[Out] Piecewise((-cos(a + b*x)**3/(3*b), Ne(b, 0)), (x*sin(a)*cos(a)**2, True))

Giac [A]

time = 4.69, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a),x, algorithm="giac")

[Out] -1/3*cos(b*x + a)^3/b

Mupad [B]

time = 0.04, size = 13, normalized size = 0.87

$$-\frac{\cos(a+bx)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2*sin(a + b*x),x)

[Out] -cos(a + b*x)^3/(3*b)

3.44 $\int \cos(a + bx) \sin(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sin^2(a + bx)}{2b}$$

[Out] 1/2*sin(b*x+a)^2/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2644, 30}

$$\frac{\sin^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Sin[a + b*x],x]

[Out] Sin[a + b*x]^2/(2*b)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \sin(a + bx) dx &= \frac{\text{Subst}(\int x dx, x, \sin(a + bx))}{b} \\ &= \frac{\sin^2(a + bx)}{2b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 37 vs. 2(15) = 30.

time = 0.01, size = 37, normalized size = 2.47

$$\frac{1}{2} \left(-\frac{\cos(2a) \cos(2bx)}{2b} + \frac{\sin(2a) \sin(2bx)}{2b} \right)$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Sin[a + b*x],x]

[Out] $(-1/2*(\text{Cos}[2*a]*\text{Cos}[2*b*x])/b + (\text{Sin}[2*a]*\text{Sin}[2*b*x])/(2*b))/2$

Maple [A]

time = 0.03, size = 14, normalized size = 0.93

method	result	size
derivativedivides	$\frac{\sin^2(bx+a)}{2b}$	14
default	$\frac{\sin^2(bx+a)}{2b}$	14
risch	$-\frac{\cos(2bx+2a)}{4b}$	15
norman	$\frac{2\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b\left(1+\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)^2}$	32

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] $1/2*\sin(b*x+a)^2/b$

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^2}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a),x, algorithm="maxima")

[Out] $-1/2*\cos(b*x + a)^2/b$

Fricas [A]

time = 0.39, size = 13, normalized size = 0.87

$$-\frac{\cos(bx+a)^2}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a),x, algorithm="fricas")

[Out] $-1/2*\cos(b*x + a)^2/b$

Sympy [A]

time = 0.06, size = 19, normalized size = 1.27

$$\begin{cases} \frac{\sin^2(a+bx)}{2b} & \text{for } b \neq 0 \\ x \sin(a) \cos(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)*sin(b*x+a),x)
```

```
[Out] Piecewise((sin(a + b*x)**2/(2*b), Ne(b, 0)), (x*sin(a)*cos(a), True))
```

Giac [A]

time = 3.89, size = 13, normalized size = 0.87

$$\frac{\sin(bx + a)^2}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)*sin(b*x+a),x, algorithm="giac")
```

```
[Out] 1/2*sin(b*x + a)^2/b
```

Mupad [B]

time = 0.45, size = 28, normalized size = 1.87

$$\begin{cases} \frac{x \sin(2a)}{2} & \text{if } b = 0 \\ -\frac{\cos(2a+2bx)}{4b} & \text{if } b \neq 0 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)*sin(a + b*x),x)
```

```
[Out] piecewise(b == 0, (x*sin(2*a))/2, b ~= 0, -cos(2*a + 2*b*x)/(4*b))
```

3.45 $\int \tan(a + bx) dx$

Optimal. Leaf size=12

$$-\frac{\log(\cos(a + bx))}{b}$$

[Out] $-\ln(\cos(b*x+a))/b$

Rubi [A]

time = 0.00, antiderivative size = 12, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3556}

$$-\frac{\log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Tan[a + b*x], x]

[Out] $-(\text{Log}[\text{Cos}[a + b*x]])/b$

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \tan(a + bx) dx = -\frac{\log(\cos(a + bx))}{b}$$

Mathematica [A]

time = 0.01, size = 12, normalized size = 1.00

$$-\frac{\log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Tan[a + b*x], x]

[Out] $-(\text{Log}[\text{Cos}[a + b*x]])/b$

Maple [A]

time = 0.02, size = 12, normalized size = 1.00

method	result	size
derivativedivides	$\frac{\ln(\sec(bx+a))}{b}$	12
default	$\frac{\ln(\sec(bx+a))}{b}$	12
norman	$\frac{\ln(1+\tan^2(bx+a))}{2b}$	17
risch	$ix + \frac{2ia}{b} - \frac{\ln(e^{2i(bx+a)}+1)}{b}$	30

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)*sin(b*x+a),x,method=_RETURNVERBOSE)`

[Out] `1/b*ln(sec(b*x+a))`

Maxima [A]

time = 0.30, size = 18, normalized size = 1.50

$$-\frac{\log(-\sin(bx+a)^2+1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)*sin(b*x+a),x, algorithm="maxima")`

[Out] `-1/2*log(-sin(b*x + a)^2 + 1)/b`

Fricas [A]

time = 0.47, size = 14, normalized size = 1.17

$$-\frac{\log(-\cos(bx+a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)*sin(b*x+a),x, algorithm="fricas")`

[Out] `-log(-cos(b*x + a))/b`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin(a+bx) \sec(a+bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)*sin(b*x+a),x)`

[Out] `Integral(sin(a + b*x)*sec(a + b*x), x)`

Giac [A]

time = 4.38, size = 18, normalized size = 1.50

$$-\frac{\log\left(\frac{|\cos(bx+a)|}{|b|}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a),x, algorithm="giac")

[Out] -log(abs(cos(b*x + a))/abs(b))/b

Mupad [B]

time = 0.51, size = 16, normalized size = 1.33

$$\frac{\ln(\tan(a + bx)^2 + 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/cos(a + b*x),x)

[Out] log(tan(a + b*x)^2 + 1)/(2*b)

3.46 $\int \sec(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=10

$$\frac{\sec(a + bx)}{b}$$

[Out] sec(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 10, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2686, 8}

$$\frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]*Tan[a + b*x],x]

[Out] Sec[a + b*x]/b

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \sec(a + bx) \tan(a + bx) dx &= \frac{\text{Subst}(\int 1 dx, x, \sec(a + bx))}{b} \\ &= \frac{\sec(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 10, normalized size = 1.00

$$\frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]*Tan[a + b*x],x]

[Out] Sec[a + b*x]/b

Maple [A]

time = 0.02, size = 11, normalized size = 1.10

method	result	size
derivativdivides	$\frac{\sec(bx+a)}{b}$	11
default	$\frac{\sec(bx+a)}{b}$	11
norman	$-\frac{2}{b\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)}$	21
risch	$\frac{2e^{i(bx+a)}}{b(e^{2i(bx+a)}+1)}$	28

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] sec(b*x+a)/b

Maxima [A]

time = 0.29, size = 12, normalized size = 1.20

$$\frac{1}{b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a),x, algorithm="maxima")

[Out] 1/(b*cos(b*x + a))

Fricas [A]

time = 0.34, size = 12, normalized size = 1.20

$$\frac{1}{b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a),x, algorithm="fricas")

[Out] 1/(b*cos(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin(a + bx) \sec^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**2*sin(b*x+a),x)`

[Out] `Integral(sin(a + b*x)*sec(a + b*x)**2, x)`

Giac [A]

time = 4.78, size = 12, normalized size = 1.20

$$\frac{1}{b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2*sin(b*x+a),x, algorithm="giac")`

[Out] `1/(b*cos(b*x + a))`

Mupad [B]

time = 0.52, size = 20, normalized size = 2.00

$$-\frac{2}{b \left(\tan \left(\frac{a}{2} + \frac{bx}{2} \right)^2 - 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)/cos(a + b*x)^2,x)`

[Out] `-2/(b*(tan(a/2 + (b*x)/2)^2 - 1))`

3.47 $\int \sec^2(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sec^2(a + bx)}{2b}$$

[Out] 1/2*sec(b*x+a)^2/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2686, 30}

$$\frac{\sec^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2*Tan[a + b*x],x]

[Out] Sec[a + b*x]^2/(2*b)

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \sec^2(a + bx) \tan(a + bx) dx &= \frac{\text{Subst}(\int x dx, x, \sec(a + bx))}{b} \\ &= \frac{\sec^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$\frac{\sec^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2*Tan[a + b*x],x]

[Out] Sec[a + b*x]^2/(2*b)

Maple [A]

time = 0.04, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$\frac{\sec^2(bx+a)}{2b}$	14
default	$\frac{\sec^2(bx+a)}{2b}$	14
risch	$\frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}+1)^2}$	28
norman	$\frac{2\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)^2}$	32

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^3*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/2*sec(b*x+a)^2/b

Maxima [A]

time = 0.29, size = 17, normalized size = 1.13

$$-\frac{1}{2(\sin(bx+a)^2-1)b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a),x, algorithm="maxima")

[Out] -1/2/((sin(b*x + a)^2 - 1)*b)

Fricas [A]

time = 0.36, size = 13, normalized size = 0.87

$$\frac{1}{2b\cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a),x, algorithm="fricas")

[Out] 1/2/(b*cos(b*x + a)^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin(a + bx) \sec^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3*sin(b*x+a),x)

[Out] Integral(sin(a + b*x)*sec(a + b*x)**3, x)

Giac [A]

time = 3.84, size = 13, normalized size = 0.87

$$\frac{1}{2b \cos(bx + a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a),x, algorithm="giac")

[Out] 1/2/(b*cos(b*x + a)^2)

Mupad [B]

time = 0.38, size = 13, normalized size = 0.87

$$\frac{\tan(a + bx)^2}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/cos(a + b*x)^3,x)

[Out] tan(a + b*x)^2/(2*b)

3.48 $\int \sec^3(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sec^3(a + bx)}{3b}$$

[Out] 1/3*sec(b*x+a)^3/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2686, 30}

$$\frac{\sec^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^3*Tan[a + b*x],x]

[Out] Sec[a + b*x]^3/(3*b)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2686

Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \sec^3(a + bx) \tan(a + bx) dx &= \frac{\text{Subst}\left(\int x^2 dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\sec^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$\frac{\sec^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^3*Tan[a + b*x],x]

[Out] Sec[a + b*x]^3/(3*b)

Maple [A]

time = 0.04, size = 14, normalized size = 0.93

method	result	size
derivativedivides	$\frac{\sec^3(bx+a)}{3b}$	14
default	$\frac{\sec^3(bx+a)}{3b}$	14
risch	$\frac{8e^{3i(bx+a)}}{3b(e^{2i(bx+a)}+1)^3}$	28
norman	$\frac{-\frac{2(\tan^4(\frac{bx}{2}+\frac{a}{2}))}{b}-\frac{2}{3b}}{(\tan^2(\frac{bx}{2}+\frac{a}{2})-1)^3}$	39

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^4*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] 1/3*sec(b*x+a)^3/b

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$\frac{1}{3b \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a),x, algorithm="maxima")

[Out] 1/3/(b*cos(b*x + a)^3)

Fricas [A]

time = 0.34, size = 13, normalized size = 0.87

$$\frac{1}{3b \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a),x, algorithm="fricas")

[Out] 1/3/(b*cos(b*x + a)^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin(a + bx) \sec^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4*sin(b*x+a),x)

[Out] Integral(sin(a + b*x)*sec(a + b*x)**4, x)

Giac [A]

time = 4.05, size = 13, normalized size = 0.87

$$\frac{1}{3 b \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a),x, algorithm="giac")

[Out] 1/3/(b*cos(b*x + a)^3)

Mupad [B]

time = 0.45, size = 13, normalized size = 0.87

$$\frac{1}{3 b \cos(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/cos(a + b*x)^4,x)

[Out] 1/(3*b*cos(a + b*x)^3)

3.49 $\int \cos^7(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=61

$$\frac{\sin^3(a + bx)}{3b} - \frac{3 \sin^5(a + bx)}{5b} + \frac{3 \sin^7(a + bx)}{7b} - \frac{\sin^9(a + bx)}{9b}$$

[Out] $1/3*\sin(b*x+a)^3/b-3/5*\sin(b*x+a)^5/b+3/7*\sin(b*x+a)^7/b-1/9*\sin(b*x+a)^9/b$

Rubi [A]

time = 0.03, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$,

Rules used = {2644, 276}

$$-\frac{\sin^9(a + bx)}{9b} + \frac{3 \sin^7(a + bx)}{7b} - \frac{3 \sin^5(a + bx)}{5b} + \frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^7*Sin[a + b*x]^2,x]`

[Out] $\text{Sin}[a + b*x]^3/(3*b) - (3*\text{Sin}[a + b*x]^5)/(5*b) + (3*\text{Sin}[a + b*x]^7)/(7*b) - \text{Sin}[a + b*x]^9/(9*b)$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^(m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2644

`Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned} \int \cos^7(a + bx) \sin^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(1 - x^2)^3 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 - 3x^4 + 3x^6 - x^8) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^3(a + bx)}{3b} - \frac{3 \sin^5(a + bx)}{5b} + \frac{3 \sin^7(a + bx)}{7b} - \frac{\sin^9(a + bx)}{9b} \end{aligned}$$

Mathematica [A]

time = 0.12, size = 47, normalized size = 0.77

$$\frac{(1606 + 1389 \cos(2(a + bx)) + 330 \cos(4(a + bx)) + 35 \cos(6(a + bx))) \sin^3(a + bx)}{10080b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^7*Sin[a + b*x]^2,x]`

```
[Out] ((1606 + 1389*Cos[2*(a + b*x)] + 330*Cos[4*(a + b*x)] + 35*Cos[6*(a + b*x)])
)*Sin[a + b*x]^3/(10080*b)
```

Maple [A]

time = 0.16, size = 60, normalized size = 0.98

method	result
risch	$\frac{7 \sin(bx+a)}{128b} - \frac{\sin(9bx+9a)}{2304b} - \frac{5 \sin(7bx+7a)}{1792b} - \frac{\sin(5bx+5a)}{160b}$
derivativedivides	$-\frac{\sin(bx+a)(\cos^8(bx+a))}{9} + \frac{\left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{b \cdot 63}$
default	$-\frac{\sin(bx+a)(\cos^8(bx+a))}{9} + \frac{\left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{b \cdot 63}$
norman	$\frac{8 \left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} - \frac{16 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5b} + \frac{632 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{35b} - \frac{2848 \left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{315b} + \frac{632 \left(\tan^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{35b} - \frac{16 \left(\tan^{13}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5b}$ $\frac{\hspace{15em}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^9}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^7*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/9*sin(b*x+a)*cos(b*x+a)^8+1/63*(16/5+cos(b*x+a)^6+6/5*cos(b*x+a)^4+
8/5*cos(b*x+a)^2)*sin(b*x+a)
```

Maxima [A]

time = 0.30, size = 46, normalized size = 0.75

$$\frac{35 \sin(bx + a)^9 - 135 \sin(bx + a)^7 + 189 \sin(bx + a)^5 - 105 \sin(bx + a)^3}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7*sin(b*x+a)^2,x, algorithm="maxima")`

```
[Out] -1/315*(35*sin(b*x + a)^9 - 135*sin(b*x + a)^7 + 189*sin(b*x + a)^5 - 105*s
in(b*x + a)^3)/b
```

Fricas [A]

time = 0.38, size = 53, normalized size = 0.87

$$\frac{(35 \cos(bx + a)^8 - 5 \cos(bx + a)^6 - 6 \cos(bx + a)^4 - 8 \cos(bx + a)^2 - 16) \sin(bx + a)}{315 b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7*sin(b*x+a)^2,x, algorithm="fricas")``[Out] -1/315*(35*cos(b*x + a)^8 - 5*cos(b*x + a)^6 - 6*cos(b*x + a)^4 - 8*cos(b*x + a)^2 - 16)*sin(b*x + a)/b`**Sympy [A]**

time = 1.25, size = 88, normalized size = 1.44

$$\begin{cases} \frac{16 \sin^9(a+bx)}{315b} + \frac{8 \sin^7(a+bx) \cos^2(a+bx)}{35b} + \frac{2 \sin^5(a+bx) \cos^4(a+bx)}{5b} + \frac{\sin^3(a+bx) \cos^6(a+bx)}{3b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos^7(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)**7*sin(b*x+a)**2,x)``[Out] Piecewise((16*sin(a + b*x)**9/(315*b) + 8*sin(a + b*x)**7*cos(a + b*x)**2/(35*b) + 2*sin(a + b*x)**5*cos(a + b*x)**4/(5*b) + sin(a + b*x)**3*cos(a + b*x)**6/(3*b), Ne(b, 0)), (x*sin(a)**2*cos(a)**7, True))`**Giac [A]**

time = 4.09, size = 54, normalized size = 0.89

$$-\frac{\sin(9bx + 9a)}{2304b} - \frac{5 \sin(7bx + 7a)}{1792b} - \frac{\sin(5bx + 5a)}{160b} + \frac{7 \sin(bx + a)}{128b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7*sin(b*x+a)^2,x, algorithm="giac")``[Out] -1/2304*sin(9*b*x + 9*a)/b - 5/1792*sin(7*b*x + 7*a)/b - 1/160*sin(5*b*x + 5*a)/b + 7/128*sin(b*x + a)/b`**Mupad [B]**

time = 0.40, size = 45, normalized size = 0.74

$$\frac{-\frac{\sin(a+bx)^9}{9} + \frac{3 \sin(a+bx)^7}{7} - \frac{3 \sin(a+bx)^5}{5} + \frac{\sin(a+bx)^3}{3}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)^7*sin(a + b*x)^2,x)``[Out] (sin(a + b*x)^3/3 - (3*sin(a + b*x)^5)/5 + (3*sin(a + b*x)^7)/7 - sin(a + b*x)^9/9)/b`

3.50 $\int \cos^5(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sin^3(a + bx)}{3b} - \frac{2 \sin^5(a + bx)}{5b} + \frac{\sin^7(a + bx)}{7b}$$

[Out] 1/3*sin(b*x+a)^3/b-2/5*sin(b*x+a)^5/b+1/7*sin(b*x+a)^7/b

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 276}

$$\frac{\sin^7(a + bx)}{7b} - \frac{2 \sin^5(a + bx)}{5b} + \frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^5*Sin[a + b*x]^2,x]

[Out] Sin[a + b*x]^3/(3*b) - (2*Sin[a + b*x]^5)/(5*b) + Sin[a + b*x]^7/(7*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx) \sin^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(1 - x^2)^2 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 - 2x^4 + x^6) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^3(a + bx)}{3b} - \frac{2 \sin^5(a + bx)}{5b} + \frac{\sin^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 37, normalized size = 0.80

$$\frac{(157 + 108 \cos(2(a + bx)) + 15 \cos(4(a + bx))) \sin^3(a + bx)}{840b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^5*Sin[a + b*x]^2,x]``[Out] ((157 + 108*Cos[2*(a + b*x)] + 15*Cos[4*(a + b*x)])*Sin[a + b*x]^3)/(840*b)`**Maple [A]**

time = 0.13, size = 50, normalized size = 1.09

method	result	size
derivativedivides	$\frac{-\frac{\sin(bx+a)(\cos^6(bx+a))}{7} + \frac{\left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{35b}}{b}$	50
default	$\frac{-\frac{\sin(bx+a)(\cos^6(bx+a))}{7} + \frac{\left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{35}}{b}$	50
risch	$\frac{5 \sin(bx+a)}{64b} - \frac{\sin(7bx+7a)}{448b} - \frac{3 \sin(5bx+5a)}{320b} - \frac{\sin(3bx+3a)}{192b}$	55
norman	$\frac{\frac{8(\tan^3(\frac{bx+a}{2}))}{3b} - \frac{32(\tan^5(\frac{bx+a}{2}))}{15b} + \frac{304(\tan^7(\frac{bx+a}{2}))}{35b} - \frac{32(\tan^9(\frac{bx+a}{2}))}{15b} + \frac{8(\tan^{11}(\frac{bx+a}{2}))}{3b}}{(1 + \tan^2(\frac{bx+a}{2}))^7}}$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5*sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/7*sin(b*x+a)*cos(b*x+a)^6+1/35*(8/3+cos(b*x+a)^4+4/3*cos(b*x+a)^2)*sin(b*x+a)`**Maxima [A]**

time = 0.29, size = 36, normalized size = 0.78

$$\frac{15 \sin(bx + a)^7 - 42 \sin(bx + a)^5 + 35 \sin(bx + a)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5*sin(b*x+a)^2,x, algorithm="maxima")``[Out] 1/105*(15*sin(b*x + a)^7 - 42*sin(b*x + a)^5 + 35*sin(b*x + a)^3)/b`**Fricas [A]**

time = 0.36, size = 43, normalized size = 0.93

$$\frac{(15 \cos(bx + a)^6 - 3 \cos(bx + a)^4 - 4 \cos(bx + a)^2 - 8) \sin(bx + a)}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^2,x, algorithm="fricas")

[Out] $-1/105*(15*\cos(b*x + a)^6 - 3*\cos(b*x + a)^4 - 4*\cos(b*x + a)^2 - 8)*\sin(b*x + a)/b$

Sympy [A]

time = 0.59, size = 66, normalized size = 1.43

$$\begin{cases} \frac{8 \sin^7(a+bx)}{105b} + \frac{4 \sin^5(a+bx) \cos^2(a+bx)}{15b} + \frac{\sin^3(a+bx) \cos^4(a+bx)}{3b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**5*sin(b*x+a)**2,x)

[Out] Piecewise((8*sin(a + b*x)**7/(105*b) + 4*sin(a + b*x)**5*cos(a + b*x)**2/(15*b) + sin(a + b*x)**3*cos(a + b*x)**4/(3*b), Ne(b, 0)), (x*sin(a)**2*cos(a)**5, True))

Giac [A]

time = 3.57, size = 54, normalized size = 1.17

$$-\frac{\sin(7bx + 7a)}{448b} - \frac{3 \sin(5bx + 5a)}{320b} - \frac{\sin(3bx + 3a)}{192b} + \frac{5 \sin(bx + a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^2,x, algorithm="giac")

[Out] $-1/448*\sin(7*b*x + 7*a)/b - 3/320*\sin(5*b*x + 5*a)/b - 1/192*\sin(3*b*x + 3*a)/b + 5/64*\sin(b*x + a)/b$

Mupad [B]

time = 0.40, size = 36, normalized size = 0.78

$$\frac{15 \sin(a + bx)^7 - 42 \sin(a + bx)^5 + 35 \sin(a + bx)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5*sin(a + b*x)^2,x)

[Out] $(35*\sin(a + b*x)^3 - 42*\sin(a + b*x)^5 + 15*\sin(a + b*x)^7)/(105*b)$

3.51 $\int \cos^3(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\sin^3(a + bx)}{3b} - \frac{\sin^5(a + bx)}{5b}$$

[Out] 1/3*sin(b*x+a)^3/b-1/5*sin(b*x+a)^5/b

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 14}

$$\frac{\sin^3(a + bx)}{3b} - \frac{\sin^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*Sin[a + b*x]^2,x]

[Out] Sin[a + b*x]^3/(3*b) - Sin[a + b*x]^5/(5*b)

Rule 14

```
Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2644

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \sin^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(1 - x^2) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 - x^4) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^3(a + bx)}{3b} - \frac{\sin^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 27, normalized size = 0.87

$$\frac{(7 + 3 \cos(2(a + bx))) \sin^3(a + bx)}{30b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^3*Sin[a + b*x]^2,x]

[Out] ((7 + 3*Cos[2*(a + b*x)])*Sin[a + b*x]^3)/(30*b)

Maple [A]

time = 0.10, size = 40, normalized size = 1.29

method	result	size
derivativedivides	$-\frac{\sin(bx+a)\cos^4(bx+a)}{5} + \frac{(2+\cos^2(bx+a))\sin(bx+a)}{15}$	40
default	$-\frac{\sin(bx+a)\cos^4(bx+a)}{5} + \frac{(2+\cos^2(bx+a))\sin(bx+a)}{15}$	40
risch	$\frac{\sin(bx+a)}{8b} - \frac{\sin(5bx+5a)}{80b} - \frac{\sin(3bx+3a)}{48b}$	41
norman	$\frac{8\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 16\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 8\left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} \frac{1}{\left(1+\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^5}$	66

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3*sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/5*sin(b*x+a)*cos(b*x+a)^4+1/15*(2+cos(b*x+a)^2)*sin(b*x+a))

Maxima [A]

time = 0.30, size = 26, normalized size = 0.84

$$-\frac{3 \sin (bx + a)^5 - 5 \sin (bx + a)^3}{15 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*sin(b*x+a)^2,x, algorithm="maxima")

[Out] -1/15*(3*sin(b*x + a)^5 - 5*sin(b*x + a)^3)/b

Fricas [A]

time = 0.38, size = 33, normalized size = 1.06

$$-\frac{(3 \cos (bx + a)^4 - \cos (bx + a)^2 - 2) \sin (bx + a)}{15 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/15*(3*cos(b*x + a)^4 - cos(b*x + a)^2 - 2)*sin(b*x + a)/b

Sympy [A]

time = 0.38, size = 44, normalized size = 1.42

$$\begin{cases} \frac{2 \sin^5(a+bx)}{15b} + \frac{\sin^3(a+bx) \cos^2(a+bx)}{3b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3*sin(b*x+a)**2,x)

[Out] Piecewise((2*sin(a + b*x)**5/(15*b) + sin(a + b*x)**3*cos(a + b*x)**2/(3*b), Ne(b, 0)), (x*sin(a)**2*cos(a)**3, True))

Giac [A]

time = 4.46, size = 26, normalized size = 0.84

$$\frac{3 \sin(bx + a)^5 - 5 \sin(bx + a)^3}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*sin(b*x+a)^2,x, algorithm="giac")

[Out] -1/15*(3*sin(b*x + a)^5 - 5*sin(b*x + a)^3)/b

Mupad [B]

time = 0.36, size = 26, normalized size = 0.84

$$\frac{5 \sin(a + bx)^3 - 3 \sin(a + bx)^5}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3*sin(a + b*x)^2,x)

[Out] (5*sin(a + b*x)^3 - 3*sin(a + b*x)^5)/(15*b)

3.52 $\int \cos(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sin^3(a + bx)}{3b}$$

[Out] 1/3*sin(b*x+a)^3/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2644, 30}

$$\frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Sin[a + b*x]^2,x]

[Out] Sin[a + b*x]^3/(3*b)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \sin^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$\frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Sin[a + b*x]^2,x]

[Out] Sin[a + b*x]^3/(3*b)

Maple [A]

time = 0.03, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$\frac{\sin^3(bx+a)}{3b}$	14
default	$\frac{\sin^3(bx+a)}{3b}$	14
risch	$\frac{\sin(bx+a)}{4b} - \frac{\sin(3bx+3a)}{12b}$	27
norman	$\frac{8\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3}$	32

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/3*sin(b*x+a)^3/b

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$\frac{\sin(bx+a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a)^2,x, algorithm="maxima")

[Out] 1/3*sin(b*x + a)^3/b

Fricas [A]

time = 0.37, size = 21, normalized size = 1.40

$$-\frac{(\cos(bx+a)^2 - 1)\sin(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/3*(cos(b*x + a)^2 - 1)*sin(b*x + a)/b

Sympy [A]

time = 0.24, size = 20, normalized size = 1.33

$$\begin{cases} \frac{\sin^3(a+bx)}{3b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)**2,x)``[Out] Piecewise((sin(a + b*x)**3/(3*b), Ne(b, 0)), (x*sin(a)**2*cos(a), True))`**Giac [A]**

time = 2.55, size = 13, normalized size = 0.87

$$\frac{\sin(bx + a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)^2,x, algorithm="giac")``[Out] 1/3*sin(b*x + a)^3/b`**Mupad [B]**

time = 0.02, size = 13, normalized size = 0.87

$$\frac{\sin(a + bx)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)*sin(a + b*x)^2,x)``[Out] sin(a + b*x)^3/(3*b)`

3.53 $\int \tan^2(a + bx) dx$

Optimal. Leaf size=14

$$-x + \frac{\tan(a + bx)}{b}$$

[Out] $-x + \tan(b*x+a)/b$

Rubi [A]

time = 0.01, antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$\frac{\tan(a + bx)}{b} - x$$

Antiderivative was successfully verified.

[In] Int[Tan[a + b*x]^2,x]

[Out] -x + Tan[a + b*x]/b

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \tan^2(a + bx) dx &= \frac{\tan(a + bx)}{b} - \int 1 dx \\ &= -x + \frac{\tan(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 23, normalized size = 1.64

$$-\frac{\tan^{-1}(\tan(a + bx))}{b} + \frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Tan[a + b*x]^2,x]

[Out] -(ArcTan[Tan[a + b*x]]/b) + Tan[a + b*x]/b

Maple [A]

time = 0.03, size = 19, normalized size = 1.36

method	result	size
derivativedivides	$\frac{\tan(bx+a)-bx-a}{b}$	19
default	$\frac{\tan(bx+a)-bx-a}{b}$	19
risch	$-x + \frac{2i}{b(e^{2i(bx+a)}+1)}$	24
norman	$\frac{x - \frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} - x \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1}$	47

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2*sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(tan(b*x+a)-b*x-a)

Maxima [A]

time = 0.51, size = 18, normalized size = 1.29

$$-\frac{bx + a - \tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^2,x, algorithm="maxima")

[Out] -(b*x + a - tan(b*x + a))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(14) = 28.

time = 0.40, size = 31, normalized size = 2.21

$$-\frac{bx \cos(bx + a) - \sin(bx + a)}{b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^2,x, algorithm="fricas")

[Out] -(b*x*cos(b*x + a) - sin(b*x + a))/(b*cos(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^2(a + bx) \sec^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**2*sin(b*x+a)**2,x)`

[Out] `Integral(sin(a + b*x)**2*sec(a + b*x)**2, x)`

Giac [A]

time = 2.93, size = 18, normalized size = 1.29

$$-\frac{bx + a - \tan(bx + a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2*sin(b*x+a)^2,x, algorithm="giac")`

[Out] `-(b*x + a - tan(b*x + a))/b`

Mupad [B]

time = 0.38, size = 14, normalized size = 1.00

$$\frac{\tan(a + bx)}{b} - x$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^2/cos(a + b*x)^2,x)`

[Out] `tan(a + b*x)/b - x`

3.54 $\int \sec^2(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\tan^3(a + bx)}{3b}$$

[Out] 1/3*tan(b*x+a)^3/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 30}

$$\frac{\tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2*Tan[a + b*x]^2,x]

[Out] Tan[a + b*x]^3/(3*b)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2687

Int[sec[(e_) + (f_)*(x_)]^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^2(a + bx) \tan^2(a + bx) dx &= \frac{\text{Subst}(\int x^2 dx, x, \tan(a + bx))}{b} \\ &= \frac{\tan^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$\frac{\tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2*Tan[a + b*x]^2,x]

[Out] Tan[a + b*x]^3/(3*b)

Maple [A]

time = 0.04, size = 22, normalized size = 1.47

method	result	size
derivativedivides	$\frac{\sin^3(bx+a)}{3b \cos(bx+a)^3}$	22
default	$\frac{\sin^3(bx+a)}{3b \cos(bx+a)^3}$	22
norman	$-\frac{8 \left(\tan^3 \left(\frac{bx+a}{2} \right) \right)}{3b \left(\tan^2 \left(\frac{bx+a}{2} \right) - 1 \right)^3}$	32
risch	$-\frac{2i(3e^{4i(bx+a)}+1)}{3b(e^{2i(bx+a)}+1)^3}$	33

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^4*sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/3/b*sin(b*x+a)^3/cos(b*x+a)^3

Maxima [A]

time = 0.30, size = 13, normalized size = 0.87

$$\frac{\tan(bx+a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^2,x, algorithm="maxima")

[Out] 1/3*tan(b*x + a)^3/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(13) = 26.

time = 0.36, size = 29, normalized size = 1.93

$$-\frac{(\cos(bx+a)^2 - 1) \sin(bx+a)}{3b \cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/3*(cos(b*x + a)^2 - 1)*sin(b*x + a)/(b*cos(b*x + a)^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^2(a + bx) \sec^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4*sin(b*x+a)**2,x)

[Out] Integral(sin(a + b*x)**2*sec(a + b*x)**4, x)

Giac [A]

time = 3.37, size = 13, normalized size = 0.87

$$\frac{\tan(bx + a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/3*tan(b*x + a)^3/b

Mupad [B]

time = 0.38, size = 13, normalized size = 0.87

$$\frac{\tan(a + bx)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/cos(a + b*x)^4,x)

[Out] tan(a + b*x)^3/(3*b)

3.55 $\int \sec^4(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\tan^3(a + bx)}{3b} + \frac{\tan^5(a + bx)}{5b}$$

[Out] 1/3*tan(b*x+a)^3/b+1/5*tan(b*x+a)^5/b

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 14}

$$\frac{\tan^5(a + bx)}{5b} + \frac{\tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^4*Tan[a + b*x]^2,x]

[Out] Tan[a + b*x]^3/(3*b) + Tan[a + b*x]^5/(5*b)

Rule 14

```
Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2687

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])
```

Rubi steps

$$\begin{aligned} \int \sec^4(a + bx) \tan^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(1 + x^2) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 + x^4) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^3(a + bx)}{3b} + \frac{\tan^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 56, normalized size = 1.81

$$-\frac{2 \tan(a + bx)}{15b} - \frac{\sec^2(a + bx) \tan(a + bx)}{15b} + \frac{\sec^4(a + bx) \tan(a + bx)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^4*Tan[a + b*x]^2,x]``[Out] (-2*Tan[a + b*x])/(15*b) - (Sec[a + b*x]^2*Tan[a + b*x])/(15*b) + (Sec[a + b*x]^4*Tan[a + b*x])/(5*b)`**Maple [A]**

time = 0.06, size = 42, normalized size = 1.35

method	result	size
derivativedivides	$\frac{\frac{\sin^3(bx+a)}{5 \cos(bx+a)^5} + \frac{2(\sin^3(bx+a))}{15 \cos(bx+a)^3}}{b}$	42
default	$\frac{\frac{\sin^3(bx+a)}{5 \cos(bx+a)^5} + \frac{2(\sin^3(bx+a))}{15 \cos(bx+a)^3}}{b}$	42
risch	$-\frac{4i(15 e^{6i(bx+a)} - 5 e^{4i(bx+a)} + 5 e^{2i(bx+a)} + 1)}{15b(e^{2i(bx+a)} + 1)^5}$	55
norman	$\frac{-\frac{8(\tan^3(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{16(\tan^5(\frac{bx}{2} + \frac{a}{2}))}{15b} - \frac{8(\tan^7(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^5}$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^6*sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/5*sin(b*x+a)^3/cos(b*x+a)^5+2/15*sin(b*x+a)^3/cos(b*x+a)^3)`**Maxima [A]**

time = 0.28, size = 26, normalized size = 0.84

$$\frac{3 \tan(bx + a)^5 + 5 \tan(bx + a)^3}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6*sin(b*x+a)^2,x, algorithm="maxima")``[Out] 1/15*(3*tan(b*x + a)^5 + 5*tan(b*x + a)^3)/b`**Fricas [A]**

time = 0.36, size = 39, normalized size = 1.26

$$-\frac{(2 \cos(bx + a)^4 + \cos(bx + a)^2 - 3) \sin(bx + a)}{15b \cos(bx + a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/15*(2*cos(b*x + a)^4 + cos(b*x + a)^2 - 3)*sin(b*x + a)/(b*cos(b*x + a)^5)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^2(a + bx) \sec^6(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**6*sin(b*x+a)**2,x)

[Out] Integral(sin(a + b*x)**2*sec(a + b*x)**6, x)

Giac [A]

time = 2.85, size = 26, normalized size = 0.84

$$\frac{3 \tan(bx + a)^5 + 5 \tan(bx + a)^3}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/15*(3*tan(b*x + a)^5 + 5*tan(b*x + a)^3)/b

Mupad [B]

time = 0.39, size = 25, normalized size = 0.81

$$\frac{\tan(a + bx)^3 (3 \tan(a + bx)^2 + 5)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/cos(a + b*x)^6,x)

[Out] (tan(a + b*x)^3*(3*tan(a + b*x)^2 + 5))/(15*b)

3.56 $\int \sec^6(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\tan^3(a + bx)}{3b} + \frac{2 \tan^5(a + bx)}{5b} + \frac{\tan^7(a + bx)}{7b}$$

[Out] 1/3*tan(b*x+a)^3/b+2/5*tan(b*x+a)^5/b+1/7*tan(b*x+a)^7/b

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 276}

$$\frac{\tan^7(a + bx)}{7b} + \frac{2 \tan^5(a + bx)}{5b} + \frac{\tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^6*Tan[a + b*x]^2,x]

[Out] Tan[a + b*x]^3/(3*b) + (2*Tan[a + b*x]^5)/(5*b) + Tan[a + b*x]^7/(7*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2687

Int[sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^6(a + bx) \tan^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(1 + x^2)^2 dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 + 2x^4 + x^6) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^3(a + bx)}{3b} + \frac{2 \tan^5(a + bx)}{5b} + \frac{\tan^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 77, normalized size = 1.67

$$-\frac{8 \tan(a+bx)}{105b} - \frac{4 \sec^2(a+bx) \tan(a+bx)}{105b} - \frac{\sec^4(a+bx) \tan(a+bx)}{35b} + \frac{\sec^6(a+bx) \tan(a+bx)}{7b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^6*Tan[a + b*x]^2,x]`

`[Out] (-8*Tan[a + b*x])/(105*b) - (4*Sec[a + b*x]^2*Tan[a + b*x])/(105*b) - (Sec[a + b*x]^4*Tan[a + b*x])/(35*b) + (Sec[a + b*x]^6*Tan[a + b*x])/(7*b)`

Maple [A]

time = 0.08, size = 60, normalized size = 1.30

method	result	size
derivativedivides	$\frac{\frac{\sin^3(bx+a)}{7 \cos(bx+a)^7} + \frac{4(\sin^3(bx+a))}{35 \cos(bx+a)^5} + \frac{8(\sin^3(bx+a))}{105 \cos(bx+a)^3}}{b}$	60
default	$\frac{\frac{\sin^3(bx+a)}{7 \cos(bx+a)^7} + \frac{4(\sin^3(bx+a))}{35 \cos(bx+a)^5} + \frac{8(\sin^3(bx+a))}{105 \cos(bx+a)^3}}{b}$	60
risch	$-\frac{16i(70e^{8i(bx+a)} - 35e^{6i(bx+a)} + 21e^{4i(bx+a)} + 7e^{2i(bx+a)} + 1)}{105b(e^{2i(bx+a)} + 1)^7}$	66
norman	$\frac{\frac{8(\tan^3(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{32(\tan^5(\frac{bx}{2} + \frac{a}{2}))}{15b} - \frac{304(\tan^7(\frac{bx}{2} + \frac{a}{2}))}{35b} - \frac{32(\tan^9(\frac{bx}{2} + \frac{a}{2}))}{15b} - \frac{8(\tan^{11}(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^7}$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^8*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

`[Out] 1/b*(1/7*sin(b*x+a)^3/cos(b*x+a)^7+4/35*sin(b*x+a)^3/cos(b*x+a)^5+8/105*sin(b*x+a)^3/cos(b*x+a)^3)`

Maxima [A]

time = 0.30, size = 36, normalized size = 0.78

$$\frac{15 \tan(bx+a)^7 + 42 \tan(bx+a)^5 + 35 \tan(bx+a)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^8*sin(b*x+a)^2,x, algorithm="maxima")`

`[Out] 1/105*(15*tan(b*x + a)^7 + 42*tan(b*x + a)^5 + 35*tan(b*x + a)^3)/b`

Fricas [A]

time = 0.42, size = 51, normalized size = 1.11

$$-\frac{(8 \cos(bx+a)^6 + 4 \cos(bx+a)^4 + 3 \cos(bx+a)^2 - 15) \sin(bx+a)}{105b \cos(bx+a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^8*sin(b*x+a)^2,x, algorithm="fricas")`

[Out] $-1/105*(8*\cos(b*x + a)^6 + 4*\cos(b*x + a)^4 + 3*\cos(b*x + a)^2 - 15)*\sin(b*x + a)/(b*\cos(b*x + a)^7)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**8*sin(b*x+a)**2,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [A]

time = 3.68, size = 36, normalized size = 0.78

$$\frac{15 \tan(bx + a)^7 + 42 \tan(bx + a)^5 + 35 \tan(bx + a)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^8*sin(b*x+a)^2,x, algorithm="giac")`

[Out] $1/105*(15*\tan(b*x + a)^7 + 42*\tan(b*x + a)^5 + 35*\tan(b*x + a)^3)/b$

Mupad [B]

time = 0.41, size = 35, normalized size = 0.76

$$\frac{\tan(a + bx)^3 (15 \tan(a + bx)^4 + 42 \tan(a + bx)^2 + 35)}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^2/cos(a + b*x)^8,x)`

[Out] $(\tan(a + b*x)^3*(42*\tan(a + b*x)^2 + 15*\tan(a + b*x)^4 + 35))/(105*b)$

3.57 $\int \sec^8(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=61

$$\frac{\tan^3(a + bx)}{3b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{3 \tan^7(a + bx)}{7b} + \frac{\tan^9(a + bx)}{9b}$$

[Out] $1/3*\tan(b*x+a)^3/b+3/5*\tan(b*x+a)^5/b+3/7*\tan(b*x+a)^7/b+1/9*\tan(b*x+a)^9/b$

Rubi [A]

time = 0.03, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$,

Rules used = {2687, 276}

$$\frac{\tan^9(a + bx)}{9b} + \frac{3 \tan^7(a + bx)}{7b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{\tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] `Int[Sec[a + b*x]^8*Tan[a + b*x]^2,x]`

[Out] `Tan[a + b*x]^3/(3*b) + (3*Tan[a + b*x]^5)/(5*b) + (3*Tan[a + b*x]^7)/(7*b) + Tan[a + b*x]^9/(9*b)`

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2687

`Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])`

Rubi steps

$$\begin{aligned} \int \sec^8(a + bx) \tan^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(1 + x^2)^3 dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 + 3x^4 + 3x^6 + x^8) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^3(a + bx)}{3b} + \frac{3 \tan^5(a + bx)}{5b} + \frac{3 \tan^7(a + bx)}{7b} + \frac{\tan^9(a + bx)}{9b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 98, normalized size = 1.61

$$\frac{16 \tan(a+bx)}{315b} - \frac{8 \sec^2(a+bx) \tan(a+bx)}{315b} - \frac{2 \sec^4(a+bx) \tan(a+bx)}{105b} - \frac{\sec^6(a+bx) \tan(a+bx)}{63b} + \frac{\sec^8(a+bx) \tan(a+bx)}{9b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^8*Tan[a + b*x]^2,x]`

`[Out] (-16*Tan[a + b*x])/(315*b) - (8*Sec[a + b*x]^2*Tan[a + b*x])/(315*b) - (2*Sec[a + b*x]^4*Tan[a + b*x])/(105*b) - (Sec[a + b*x]^6*Tan[a + b*x])/(63*b) + (Sec[a + b*x]^8*Tan[a + b*x])/(9*b)`

Maple [A]

time = 0.04, size = 78, normalized size = 1.28

method	result	size
risch	$\frac{32i(315 e^{10i(bx+a)} - 189 e^{8i(bx+a)} + 84 e^{6i(bx+a)} + 36 e^{4i(bx+a)} + 9 e^{2i(bx+a)} + 1)}{315b(e^{2i(bx+a)} + 1)^9}$	77
derivativedivides	$\frac{\frac{\sin^3(bx+a)}{9 \cos(bx+a)^9} + \frac{2(\sin^3(bx+a))}{21 \cos(bx+a)^7} + \frac{8(\sin^3(bx+a))}{105 \cos(bx+a)^5} + \frac{16(\sin^3(bx+a))}{315 \cos(bx+a)^3}}{b}$	78
default	$\frac{\frac{\sin^3(bx+a)}{9 \cos(bx+a)^9} + \frac{2(\sin^3(bx+a))}{21 \cos(bx+a)^7} + \frac{8(\sin^3(bx+a))}{105 \cos(bx+a)^5} + \frac{16(\sin^3(bx+a))}{315 \cos(bx+a)^3}}{b}$	78

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^10*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

`[Out] 1/b*(1/9*sin(b*x+a)^3/cos(b*x+a)^9+2/21*sin(b*x+a)^3/cos(b*x+a)^7+8/105*sin(b*x+a)^3/cos(b*x+a)^5+16/315*sin(b*x+a)^3/cos(b*x+a)^3)`

Maxima [A]

time = 0.29, size = 46, normalized size = 0.75

$$\frac{35 \tan(bx+a)^9 + 135 \tan(bx+a)^7 + 189 \tan(bx+a)^5 + 105 \tan(bx+a)^3}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^10*sin(b*x+a)^2,x, algorithm="maxima")`

`[Out] 1/315*(35*tan(b*x + a)^9 + 135*tan(b*x + a)^7 + 189*tan(b*x + a)^5 + 105*tan(b*x + a)^3)/b`

Fricas [A]

time = 0.40, size = 61, normalized size = 1.00

$$\frac{(16 \cos(bx+a)^8 + 8 \cos(bx+a)^6 + 6 \cos(bx+a)^4 + 5 \cos(bx+a)^2 - 35) \sin(bx+a)}{315b \cos(bx+a)^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^10*sin(b*x+a)^2,x, algorithm="fricas")

[Out] $-1/315*(16*\cos(b*x + a)^8 + 8*\cos(b*x + a)^6 + 6*\cos(b*x + a)^4 + 5*\cos(b*x + a)^2 - 35)*\sin(b*x + a)/(b*\cos(b*x + a)^9)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**10*sin(b*x+a)**2,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 8569 deep

Giac [A]

time = 3.38, size = 46, normalized size = 0.75

$$\frac{35 \tan (bx + a)^9 + 135 \tan (bx + a)^7 + 189 \tan (bx + a)^5 + 105 \tan (bx + a)^3}{315 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^10*sin(b*x+a)^2,x, algorithm="giac")

[Out] $1/315*(35*\tan(b*x + a)^9 + 135*\tan(b*x + a)^7 + 189*\tan(b*x + a)^5 + 105*\tan(b*x + a)^3)/b$

Mupad [B]

time = 0.41, size = 45, normalized size = 0.74

$$\frac{\frac{\tan(a+bx)^9}{9} + \frac{3 \tan(a+bx)^7}{7} + \frac{3 \tan(a+bx)^5}{5} + \frac{\tan(a+bx)^3}{3}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/cos(a + b*x)^10,x)

[Out] $(\tan(a + b*x)^3/3 + (3*\tan(a + b*x)^5)/5 + (3*\tan(a + b*x)^7)/7 + \tan(a + b*x)^9/9)/b$

3.58 $\int \cos^6(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=88

$$\frac{5x}{128} + \frac{5 \cos(a + bx) \sin(a + bx)}{128b} + \frac{5 \cos^3(a + bx) \sin(a + bx)}{192b} + \frac{\cos^5(a + bx) \sin(a + bx)}{48b} - \frac{\cos^7(a + bx) \sin(a + bx)}{8b}$$

[Out] 5/128*x+5/128*cos(b*x+a)*sin(b*x+a)/b+5/192*cos(b*x+a)^3*sin(b*x+a)/b+1/48*cos(b*x+a)^5*sin(b*x+a)/b-1/8*cos(b*x+a)^7*sin(b*x+a)/b

Rubi [A]

time = 0.04, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2648, 2715, 8}

$$-\frac{\sin(a + bx) \cos^7(a + bx)}{8b} + \frac{\sin(a + bx) \cos^5(a + bx)}{48b} + \frac{5 \sin(a + bx) \cos^3(a + bx)}{192b} + \frac{5 \sin(a + bx) \cos(a + bx)}{128b} + \frac{5x}{128}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^6*Sin[a + b*x]^2,x]

[Out] (5*x)/128 + (5*Cos[a + b*x]*Sin[a + b*x])/(128*b) + (5*Cos[a + b*x]^3*Sin[a + b*x])/(192*b) + (Cos[a + b*x]^5*Sin[a + b*x])/(48*b) - (Cos[a + b*x]^7*Sin[a + b*x])/(8*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2648

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^n*((a_)*sin[(e_) + (f_)*(x_)])^m, x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^n, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \cos^6(a+bx) \sin^2(a+bx) dx &= -\frac{\cos^7(a+bx) \sin(a+bx)}{8b} + \frac{1}{8} \int \cos^6(a+bx) dx \\
&= \frac{\cos^5(a+bx) \sin(a+bx)}{48b} - \frac{\cos^7(a+bx) \sin(a+bx)}{8b} + \frac{5}{48} \int \cos^4(a+bx) dx \\
&= \frac{5 \cos^3(a+bx) \sin(a+bx)}{192b} + \frac{\cos^5(a+bx) \sin(a+bx)}{48b} - \frac{\cos^7(a+bx) \sin(a+bx)}{8b} \\
&= \frac{5 \cos(a+bx) \sin(a+bx)}{128b} + \frac{5 \cos^3(a+bx) \sin(a+bx)}{192b} + \frac{\cos^5(a+bx) \sin(a+bx)}{48b} \\
&= \frac{5x}{128} + \frac{5 \cos(a+bx) \sin(a+bx)}{128b} + \frac{5 \cos^3(a+bx) \sin(a+bx)}{192b} + \frac{\cos^5(a+bx) \sin(a+bx)}{48b}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 52, normalized size = 0.59

$$\frac{120bx + 48 \sin(2(a+bx)) - 24 \sin(4(a+bx)) - 16 \sin(6(a+bx)) - 3 \sin(8(a+bx))}{3072b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^6*Sin[a + b*x]^2,x]`

```
[Out] (120*b*x + 48*Sin[2*(a + b*x)] - 24*Sin[4*(a + b*x)] - 16*Sin[6*(a + b*x)] - 3*Sin[8*(a + b*x)])/(3072*b)
```

Maple [A]

time = 0.15, size = 64, normalized size = 0.73

method	result
risch	$\frac{5x}{128} - \frac{\sin(8bx+8a)}{1024b} - \frac{\sin(6bx+6a)}{192b} - \frac{\sin(4bx+4a)}{128b} + \frac{\sin(2bx+2a)}{64b}$
derivativedivides	$-\frac{(\cos^7(bx+a) \sin(bx+a))}{8} + \frac{\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15 \cos(bx+a)}{8}\right) \sin(bx+a)}{48} + \frac{5bx}{128} + \frac{5a}{128}$
default	$-\frac{(\cos^7(bx+a) \sin(bx+a))}{8} + \frac{\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15 \cos(bx+a)}{8}\right) \sin(bx+a)}{48} + \frac{5bx}{128} + \frac{5a}{128}$
norman	$\frac{5x}{128} - \frac{5 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{64b} + \frac{397 \left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{192b} - \frac{895 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{192b} + \frac{1765 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{192b} - \frac{1765 \left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{192b} + \frac{895 \left(\tan^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{192b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^6*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/8*\cos(b*x+a)^7*\sin(b*x+a)+1/48*(\cos(b*x+a)^5+5/4*\cos(b*x+a)^3+15/8*\cos(b*x+a))*\sin(b*x+a)+5/128*b*x+5/128*a)$

Maxima [A]

time = 0.30, size = 48, normalized size = 0.55

$$\frac{64 \sin(2bx + 2a)^3 + 120bx + 120a - 3 \sin(8bx + 8a) - 24 \sin(4bx + 4a)}{3072b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6*sin(b*x+a)^2,x, algorithm="maxima")`

[Out] $1/3072*(64*\sin(2*b*x + 2*a)^3 + 120*b*x + 120*a - 3*\sin(8*b*x + 8*a) - 24*\sin(4*b*x + 4*a))/b$

Fricas [A]

time = 0.38, size = 57, normalized size = 0.65

$$\frac{15bx - (48 \cos(bx + a)^7 - 8 \cos(bx + a)^5 - 10 \cos(bx + a)^3 - 15 \cos(bx + a)) \sin(bx + a)}{384b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6*sin(b*x+a)^2,x, algorithm="fricas")`

[Out] $1/384*(15*b*x - (48*\cos(b*x + a)^7 - 8*\cos(b*x + a)^5 - 10*\cos(b*x + a)^3 - 15*\cos(b*x + a))*\sin(b*x + a))/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(80) = 160$.

time = 0.87, size = 189, normalized size = 2.15

$$\begin{cases} \frac{5x \sin^6(a+bx)}{128} + \frac{5x \sin^6(a+bx) \cos^2(a+bx)}{32} + \frac{15x \sin^4(a+bx) \cos^4(a+bx)}{64} + \frac{5x \sin^2(a+bx) \cos^6(a+bx)}{32} + \frac{5x \cos^8(a+bx)}{128} + \frac{5 \sin^7(a+bx) \cos(a+bx)}{128b} + \frac{55 \sin^5(a+bx) \cos^3(a+bx)}{384b} + \frac{73 \sin^3(a+bx) \cos^5(a+bx)}{384b} - \frac{5 \sin(a+bx) \cos^7(a+bx)}{128b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos^6(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**6*sin(b*x+a)**2,x)`

[Out] `Piecewise((5*x*sin(a + b*x)**8/128 + 5*x*sin(a + b*x)**6*cos(a + b*x)**2/32 + 15*x*sin(a + b*x)**4*cos(a + b*x)**4/64 + 5*x*sin(a + b*x)**2*cos(a + b*x)**6/32 + 5*x*cos(a + b*x)**8/128 + 5*sin(a + b*x)**7*cos(a + b*x)/(128*b) + 55*sin(a + b*x)**5*cos(a + b*x)**3/(384*b) + 73*sin(a + b*x)**3*cos(a + b*x)**5/(384*b) - 5*sin(a + b*x)*cos(a + b*x)**7/(128*b), Ne(b, 0)), (x*sin(a)**2*cos(a)**6, True))`

Giac [A]

time = 4.06, size = 60, normalized size = 0.68

$$\frac{5}{128}x - \frac{\sin(8bx + 8a)}{1024b} - \frac{\sin(6bx + 6a)}{192b} - \frac{\sin(4bx + 4a)}{128b} + \frac{\sin(2bx + 2a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^2,x, algorithm="giac")

[Out] 5/128*x - 1/1024*sin(8*b*x + 8*a)/b - 1/192*sin(6*b*x + 6*a)/b - 1/128*sin(4*b*x + 4*a)/b + 1/64*sin(2*b*x + 2*a)/b

Mupad [B]

time = 1.52, size = 89, normalized size = 1.01

$$\frac{5x}{128} + \frac{\frac{5 \tan(a+bx)^7}{128} + \frac{55 \tan(a+bx)^5}{384} + \frac{73 \tan(a+bx)^3}{384} - \frac{5 \tan(a+bx)}{128}}{b (\tan(a+bx)^8 + 4 \tan(a+bx)^6 + 6 \tan(a+bx)^4 + 4 \tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6*sin(a + b*x)^2,x)

[Out] (5*x)/128 + ((73*tan(a + b*x)^3)/384 - (5*tan(a + b*x))/128 + (55*tan(a + b*x)^5)/384 + (5*tan(a + b*x)^7)/128)/(b*(4*tan(a + b*x)^2 + 6*tan(a + b*x)^4 + 4*tan(a + b*x)^6 + tan(a + b*x)^8 + 1))

3.59 $\int \cos^4(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=67

$$\frac{x}{16} + \frac{\cos(a + bx) \sin(a + bx)}{16b} + \frac{\cos^3(a + bx) \sin(a + bx)}{24b} - \frac{\cos^5(a + bx) \sin(a + bx)}{6b}$$

[Out] 1/16*x+1/16*cos(b*x+a)*sin(b*x+a)/b+1/24*cos(b*x+a)^3*sin(b*x+a)/b-1/6*cos(b*x+a)^5*sin(b*x+a)/b

Rubi [A]

time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2648, 2715, 8}

$$-\frac{\sin(a + bx) \cos^5(a + bx)}{6b} + \frac{\sin(a + bx) \cos^3(a + bx)}{24b} + \frac{\sin(a + bx) \cos(a + bx)}{16b} + \frac{x}{16}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Sin[a + b*x]^2,x]

[Out] x/16 + (Cos[a + b*x]*Sin[a + b*x])/(16*b) + (Cos[a + b*x]^3*Sin[a + b*x])/(24*b) - (Cos[a + b*x]^5*Sin[a + b*x])/(6*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m_, x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \cos^4(a+bx) \sin^2(a+bx) dx &= -\frac{\cos^5(a+bx) \sin(a+bx)}{6b} + \frac{1}{6} \int \cos^4(a+bx) dx \\
&= \frac{\cos^3(a+bx) \sin(a+bx)}{24b} - \frac{\cos^5(a+bx) \sin(a+bx)}{6b} + \frac{1}{8} \int \cos^2(a+bx) dx \\
&= \frac{\cos(a+bx) \sin(a+bx)}{16b} + \frac{\cos^3(a+bx) \sin(a+bx)}{24b} - \frac{\cos^5(a+bx) \sin(a+bx)}{6b} \\
&= \frac{x}{16} + \frac{\cos(a+bx) \sin(a+bx)}{16b} + \frac{\cos^3(a+bx) \sin(a+bx)}{24b} - \frac{\cos^5(a+bx) \sin(a+bx)}{6b}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 40, normalized size = 0.60

$$-\frac{-12bx - 3 \sin(2(a+bx)) + 3 \sin(4(a+bx)) + \sin(6(a+bx))}{192b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^4*Sin[a + b*x]^2,x]``[Out] -1/192*(-12*b*x - 3*Sin[2*(a + b*x)] + 3*Sin[4*(a + b*x)] + Sin[6*(a + b*x)])/b`**Maple [A]**

time = 0.12, size = 54, normalized size = 0.81

method	result
risch	$\frac{x}{16} - \frac{\sin(6bx+6a)}{192b} - \frac{\sin(4bx+4a)}{64b} + \frac{\sin(2bx+2a)}{64b}$
derivativedivides	$-\frac{(\cos^5(bx+a) \sin(bx+a))}{6} + \frac{(\cos^3(bx+a) + \frac{3 \cos(bx+a)}{2}) \sin(bx+a)}{24} + \frac{bx+a}{16}$
default	$-\frac{(\cos^5(bx+a) \sin(bx+a))}{6} + \frac{(\cos^3(bx+a) + \frac{3 \cos(bx+a)}{2}) \sin(bx+a)}{24} + \frac{bx+a}{16}$
norman	$\frac{x}{16} - \frac{\tan(\frac{bx}{2} + \frac{a}{2})}{8b} + \frac{47(\tan^3(\frac{bx}{2} + \frac{a}{2}))}{24b} - \frac{13(\tan^5(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{13(\tan^7(\frac{bx}{2} + \frac{a}{2}))}{4b} - \frac{47(\tan^9(\frac{bx}{2} + \frac{a}{2}))}{24b} + \frac{\tan^{11}(\frac{bx}{2} + \frac{a}{2})}{8b} + \frac{3x(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{8(1+\tan^2(\frac{bx}{2} + \frac{a}{2}))}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^4*sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/6*cos(b*x+a)^5*sin(b*x+a)+1/24*(cos(b*x+a)^3+3/2*cos(b*x+a))*sin(b*x+a)+1/16*b*x+1/16*a)`

Maxima [A]

time = 0.34, size = 37, normalized size = 0.55

$$\frac{4 \sin(2bx + 2a)^3 + 12bx + 12a - 3 \sin(4bx + 4a)}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^2,x, algorithm="maxima")**[Out]** 1/192*(4*sin(2*b*x + 2*a)^3 + 12*b*x + 12*a - 3*sin(4*b*x + 4*a))/b**Fricas [A]**

time = 0.37, size = 47, normalized size = 0.70

$$\frac{3bx - (8 \cos(bx + a)^5 - 2 \cos(bx + a)^3 - 3 \cos(bx + a)) \sin(bx + a)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^2,x, algorithm="fricas")**[Out]** 1/48*(3*b*x - (8*cos(b*x + a)^5 - 2*cos(b*x + a)^3 - 3*cos(b*x + a))*sin(b*x + a))/b**Sympy [B]** Leaf count of result is larger than twice the leaf count of optimal. 136 vs. 2(56) = 112.

time = 0.40, size = 136, normalized size = 2.03

$$\begin{cases} \frac{x \sin^6(a+bx)}{16} + \frac{3x \sin^4(a+bx) \cos^2(a+bx)}{16} + \frac{3x \sin^2(a+bx) \cos^4(a+bx)}{16} + \frac{x \cos^6(a+bx)}{16} + \frac{\sin^5(a+bx) \cos(a+bx)}{16b} + \frac{\sin^3(a+bx) \cos^3(a+bx)}{6b} - \frac{\sin(a+bx) \cos^5(a+bx)}{16b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**4*sin(b*x+a)**2,x)**[Out]** Piecewise((x*sin(a + b*x)**6/16 + 3*x*sin(a + b*x)**4*cos(a + b*x)**2/16 + 3*x*sin(a + b*x)**2*cos(a + b*x)**4/16 + x*cos(a + b*x)**6/16 + sin(a + b*x)**5*cos(a + b*x)/(16*b) + sin(a + b*x)**3*cos(a + b*x)**3/(6*b) - sin(a + b*x)*cos(a + b*x)**5/(16*b), Ne(b, 0)), (x*sin(a)**2*cos(a)**4, True))**Giac [A]**

time = 6.04, size = 46, normalized size = 0.69

$$\frac{1}{16} x - \frac{\sin(6bx + 6a)}{192b} - \frac{\sin(4bx + 4a)}{64b} + \frac{\sin(2bx + 2a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^2,x, algorithm="giac")

[Out] $\frac{1}{16}x - \frac{1}{192}\frac{\sin(6bx + 6a)}{b} - \frac{1}{64}\frac{\sin(4bx + 4a)}{b} + \frac{1}{64}\frac{\sin(2bx + 2a)}{b}$

Mupad [B]

time = 0.56, size = 43, normalized size = 0.64

$$\frac{x}{16} - \frac{\frac{\sin(4a+4bx)}{64} - \frac{\sin(2a+2bx)}{64} + \frac{\sin(6a+6bx)}{192}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^4*sin(a + b*x)^2,x)`

[Out] $x/16 - (\sin(4a + 4bx)/64 - \sin(2a + 2bx)/64 + \sin(6a + 6bx)/192)/b$

3.60 $\int \cos^2(a + bx) \sin^2(a + bx) dx$

Optimal. Leaf size=46

$$\frac{x}{8} + \frac{\cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos^3(a + bx) \sin(a + bx)}{4b}$$

[Out] 1/8*x+1/8*cos(b*x+a)*sin(b*x+a)/b-1/4*cos(b*x+a)^3*sin(b*x+a)/b

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2648, 2715, 8}

$$-\frac{\sin(a + bx) \cos^3(a + bx)}{4b} + \frac{\sin(a + bx) \cos(a + bx)}{8b} + \frac{x}{8}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Sin[a + b*x]^2,x]

[Out] x/8 + (Cos[a + b*x]*Sin[a + b*x])/(8*b) - (Cos[a + b*x]^3*Sin[a + b*x])/(4*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2648

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^n_)*((a_.)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^n_, x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \cos^2(a+bx) \sin^2(a+bx) dx &= -\frac{\cos^3(a+bx) \sin(a+bx)}{4b} + \frac{1}{4} \int \cos^2(a+bx) dx \\
&= \frac{\cos(a+bx) \sin(a+bx)}{8b} - \frac{\cos^3(a+bx) \sin(a+bx)}{4b} + \frac{\int 1 dx}{8} \\
&= \frac{x}{8} + \frac{\cos(a+bx) \sin(a+bx)}{8b} - \frac{\cos^3(a+bx) \sin(a+bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 23, normalized size = 0.50

$$-\frac{-4(a+bx) + \sin(4(a+bx))}{32b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^2*Sin[a + b*x]^2,x]``[Out] -1/32*(-4*(a + b*x) + Sin[4*(a + b*x)])/b`**Maple [A]**

time = 0.04, size = 43, normalized size = 0.93

method	result
risch	$\frac{x}{8} - \frac{\sin(4bx+4a)}{32b}$
derivativedivides	$-\frac{(\cos^3(bx+a) \sin(bx+a))}{4} + \frac{\cos(bx+a) \sin(bx+a)}{8} + \frac{bx}{8} + \frac{a}{8}$ b
default	$-\frac{(\cos^3(bx+a) \sin(bx+a))}{4} + \frac{\cos(bx+a) \sin(bx+a)}{8} + \frac{bx}{8} + \frac{a}{8}$ b
norman	$\frac{x}{8} - \frac{\tan(\frac{bx}{2} + \frac{a}{2})}{4b} + \frac{7(\tan^3(\frac{bx}{2} + \frac{a}{2}))}{4b} - \frac{7(\tan^5(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{\tan^7(\frac{bx}{2} + \frac{a}{2})}{4b} + \frac{x(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{2} + \frac{3x(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{4} + \frac{x(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{2}$ $(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^4$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^2*sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/4*cos(b*x+a)^3*sin(b*x+a)+1/8*cos(b*x+a)*sin(b*x+a)+1/8*b*x+1/8*a)`**Maxima [A]**

time = 0.29, size = 24, normalized size = 0.52

$$\frac{4bx + 4a - \sin(4bx + 4a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^2,x, algorithm="maxima")

[Out] 1/32*(4*b*x + 4*a - sin(4*b*x + 4*a))/b

Fricas [A]

time = 0.37, size = 36, normalized size = 0.78

$$\frac{bx - (2 \cos(bx + a))^3 - \cos(bx + a) \sin(bx + a)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^2,x, algorithm="fricas")

[Out] 1/8*(b*x - (2*cos(b*x + a))^3 - cos(b*x + a))*sin(b*x + a)/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(37) = 74.

time = 0.17, size = 92, normalized size = 2.00

$$\begin{cases} \frac{x \sin^4(a+bx)}{8} + \frac{x \sin^2(a+bx) \cos^2(a+bx)}{4} + \frac{x \cos^4(a+bx)}{8} + \frac{\sin^3(a+bx) \cos(a+bx)}{8b} - \frac{\sin(a+bx) \cos^3(a+bx)}{8b} & \text{for } b \neq 0 \\ x \sin^2(a) \cos^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2*sin(b*x+a)**2,x)

[Out] Piecewise((x*sin(a + b*x)**4/8 + x*sin(a + b*x)**2*cos(a + b*x)**2/4 + x*cos(a + b*x)**4/8 + sin(a + b*x)**3*cos(a + b*x)/(8*b) - sin(a + b*x)*cos(a + b*x)**3/(8*b), Ne(b, 0)), (x*sin(a)**2*cos(a)**2, True))

Giac [A]

time = 4.53, size = 18, normalized size = 0.39

$$\frac{1}{8}x - \frac{\sin(4bx + 4a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/8*x - 1/32*sin(4*b*x + 4*a)/b

Mupad [B]

time = 0.46, size = 50, normalized size = 1.09

$$\frac{x}{8} - \frac{\frac{\tan(a+bx)}{8} - \frac{\tan(a+bx)^3}{8}}{b(\tan(a+bx)^4 + 2\tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2*sin(a + b*x)^2,x)

[Out] x/8 - (tan(a + b*x)/8 - tan(a + b*x)^3/8)/(b*(2*tan(a + b*x)^2 + tan(a + b*x)^4 + 1))

3.61 $\int \sin^2(a + bx) dx$

Optimal. Leaf size=25

$$\frac{x}{2} - \frac{\cos(a + bx) \sin(a + bx)}{2b}$$

[Out] 1/2*x-1/2*cos(b*x+a)*sin(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$\frac{x}{2} - \frac{\sin(a + bx) \cos(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2,x]

[Out] x/2 - (Cos[a + b*x]*Sin[a + b*x])/(2*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sin^2(a + bx) dx &= -\frac{\cos(a + bx) \sin(a + bx)}{2b} + \frac{\int 1 dx}{2} \\ &= \frac{x}{2} - \frac{\cos(a + bx) \sin(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 23, normalized size = 0.92

$$-\frac{-2(a + bx) + \sin(2(a + bx))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^2,x]

[Out] $-1/4*(-2*(a + b*x) + \text{Sin}[2*(a + b*x)])/b$

Maple [A]

time = 0.00, size = 27, normalized size = 1.08

method	result	size
risch	$\frac{x}{2} - \frac{\sin(2bx+2a)}{4b}$	19
derivativdivides	$-\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx+a}{2}$ b	27
default	$-\frac{\cos(bx+a)\sin(bx+a)}{2} + \frac{bx+a}{2}$ b	27
norman	$\frac{\tan^3\left(\frac{bx+a}{2}\right)}{b} + x\left(\tan^2\left(\frac{bx+a}{2}\right)\right) + \frac{x}{2} - \frac{\tan\left(\frac{bx+a}{2}\right)}{b} + \frac{x\left(\tan^4\left(\frac{bx+a}{2}\right)\right)}{2}$ $(1+\tan^2\left(\frac{bx+a}{2}\right))^2$	77

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] $1/b*(-1/2*\cos(b*x+a)*\sin(b*x+a)+1/2*b*x+1/2*a)$

Maxima [A]

time = 0.32, size = 24, normalized size = 0.96

$$\frac{2bx + 2a - \sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2,x, algorithm="maxima")

[Out] $1/4*(2*b*x + 2*a - \sin(2*b*x + 2*a))/b$

Fricas [A]

time = 0.38, size = 23, normalized size = 0.92

$$\frac{bx - \cos(bx + a)\sin(bx + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2,x, algorithm="fricas")

[Out] $1/2*(b*x - \cos(b*x + a)*\sin(b*x + a))/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(19) = 38$.

time = 0.07, size = 46, normalized size = 1.84

$$\begin{cases} \frac{x \sin^2(a+bx)}{2} + \frac{x \cos^2(a+bx)}{2} - \frac{\sin(a+bx)\cos(a+bx)}{2b} & \text{for } b \neq 0 \\ x \sin^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**2,x)

[Out] Piecewise((x*sin(a + b*x)**2/2 + x*cos(a + b*x)**2/2 - sin(a + b*x)*cos(a + b*x)/(2*b), Ne(b, 0)), (x*sin(a)**2, True))

Giac [A]

time = 4.07, size = 18, normalized size = 0.72

$$\frac{1}{2}x - \frac{\sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/2*x - 1/4*sin(2*b*x + 2*a)/b

Mupad [B]

time = 0.00, size = 18, normalized size = 0.72

$$\frac{x}{2} - \frac{\sin(2a + 2bx)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2,x)

[Out] x/2 - sin(2*a + 2*b*x)/(4*b)

3.62 $\int \sin(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=23

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\sin(a + bx)}{b}$$

[Out] arctanh(sin(b*x+a))/b-sin(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2672, 327, 212}

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]*Tan[a + b*x],x]

[Out] ArcTanh[Sin[a + b*x]]/b - Sin[a + b*x]/b

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2672

Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]

Rubi steps

$$\begin{aligned} \int \sin(a + bx) \tan(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \sin(a + bx)\right)}{b} \\ &= -\frac{\sin(a + bx)}{b} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\sin(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 23, normalized size = 1.00

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]*Tan[a + b*x],x]``[Out] ArcTanh[Sin[a + b*x]]/b - Sin[a + b*x]/b`**Maple [A]**

time = 0.03, size = 28, normalized size = 1.22

method	result	size
derivativedivides	$\frac{-\sin(bx+a) + \ln(\sec(bx+a) + \tan(bx+a))}{b}$	28
default	$\frac{-\sin(bx+a) + \ln(\sec(bx+a) + \tan(bx+a))}{b}$	28
norman	$-\frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b}$	64
risch	$\frac{ie^{i(bx+a)}}{2b} - \frac{ie^{-i(bx+a)}}{2b} - \frac{\ln(e^{i(bx+a)} - i)}{b} + \frac{\ln(e^{i(bx+a)} + i)}{b}$	67

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(-sin(b*x+a)+ln(sec(b*x+a)+tan(b*x+a)))`**Maxima [A]**

time = 0.29, size = 34, normalized size = 1.48

$$\frac{\log(\sin(bx + a) + 1) - \log(\sin(bx + a) - 1) - 2 \sin(bx + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

Giac [A]

time = 3.78, size = 36, normalized size = 1.57

$$\frac{\log(|\sin(bx + a) + 1|) - \log(|\sin(bx + a) - 1|) - 2 \sin(bx + a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/2*(log(abs(sin(b*x + a) + 1)) - log(abs(sin(b*x + a) - 1)) - 2*sin(b*x + a))/b

Mupad [B]

time = 0.46, size = 27, normalized size = 1.17

$$\frac{2 \operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{b} - \frac{\sin(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/cos(a + b*x),x)

[Out] (2*atanh(tan(a/2 + (b*x)/2)))/b - sin(a + b*x)/b

3.63 $\int \sec(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=34

$$-\frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\sec(a + bx) \tan(a + bx)}{2b}$$

[Out] $-1/2*\operatorname{arctanh}(\sin(b*x+a))/b+1/2*\sec(b*x+a)*\tan(b*x+a)/b$

Rubi [A]

time = 0.02, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2691, 3855}

$$\frac{\tan(a + bx) \sec(a + bx)}{2b} - \frac{\tanh^{-1}(\sin(a + bx))}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Sec[a + b*x]*Tan[a + b*x]^2,x]`

[Out] $-1/2*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]]/b + (\operatorname{Sec}[a + b*x]*\operatorname{Tan}[a + b*x])/(2*b)$

Rule 2691

```
Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]
```

Rule 3855

```
Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \sec(a + bx) \tan^2(a + bx) dx &= \frac{\sec(a + bx) \tan(a + bx)}{2b} - \frac{1}{2} \int \sec(a + bx) dx \\ &= -\frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\sec(a + bx) \tan(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 34, normalized size = 1.00

$$-\frac{\tanh^{-1}(\sin(a + bx))}{2b} + \frac{\sec(a + bx) \tan(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]*Tan[a + b*x]^2,x]

[Out] -1/2*ArcTanh[Sin[a + b*x]]/b + (Sec[a + b*x]*Tan[a + b*x])/(2*b)

Maple [A]

time = 0.05, size = 48, normalized size = 1.41

method	result	size
derivativedivides	$\frac{\frac{\sin^3(bx+a)}{2 \cos(bx+a)^2} + \frac{\sin(bx+a)}{2} - \frac{\ln(\sec(bx+a)+\tan(bx+a))}{2}}{b}$	48
default	$\frac{\frac{\sin^3(bx+a)}{2 \cos(bx+a)^2} + \frac{\sin(bx+a)}{2} - \frac{\ln(\sec(bx+a)+\tan(bx+a))}{2}}{b}$	48
risch	$-\frac{i(e^{3i(bx+a)}-e^{i(bx+a)})}{b(e^{2i(bx+a)}+1)^2} + \frac{\ln(e^{i(bx+a)}-i)}{2b} - \frac{\ln(e^{i(bx+a)}+i)}{2b}$	78
norman	$\frac{\frac{\tan(\frac{bx}{2}+\frac{a}{2})}{b} + \frac{\tan^3(\frac{bx}{2}+\frac{a}{2})}{b}}{(\tan^2(\frac{bx}{2}+\frac{a}{2})-1)^2} + \frac{\ln(\tan(\frac{bx}{2}+\frac{a}{2})-1)}{2b} - \frac{\ln(\tan(\frac{bx}{2}+\frac{a}{2})+1)}{2b}$	81

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^3*sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(1/2*sin(b*x+a)^3/cos(b*x+a)^2+1/2*sin(b*x+a)-1/2*ln(sec(b*x+a)+tan(b*x+a)))

Maxima [A]

time = 0.31, size = 46, normalized size = 1.35

$$-\frac{2 \sin(bx+a)}{\sin(bx+a)^2-1} + \log(\sin(bx+a)+1) - \log(\sin(bx+a)-1)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^2,x, algorithm="maxima")

[Out] -1/4*(2*sin(b*x + a)/(sin(b*x + a)^2 - 1) + log(sin(b*x + a) + 1) - log(sin(b*x + a) - 1))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 61 vs. 2(30) = 60.

time = 0.41, size = 61, normalized size = 1.79

$$\frac{\cos(bx+a)^2 \log(\sin(bx+a)+1) - \cos(bx+a)^2 \log(-\sin(bx+a)+1) - 2 \sin(bx+a)}{4b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^2,x, algorithm="fricas")

[Out] $-1/4*(\cos(b*x + a)^2*\log(\sin(b*x + a) + 1) - \cos(b*x + a)^2*\log(-\sin(b*x + a) + 1) - 2*\sin(b*x + a))/(b*\cos(b*x + a)^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^2(a + bx) \sec^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3*sin(b*x+a)**2,x)

[Out] Integral(sin(a + b*x)**2*sec(a + b*x)**3, x)

Giac [A]

time = 3.67, size = 48, normalized size = 1.41

$$\frac{\frac{2 \sin(bx+a)}{\sin(bx+a)^2-1} + \log(|\sin(bx+a)+1|) - \log(|\sin(bx+a)-1|)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^2,x, algorithm="giac")

[Out] $-1/4*(2*\sin(b*x + a)/(\sin(b*x + a)^2 - 1) + \log(\text{abs}(\sin(b*x + a) + 1)) - \log(\text{abs}(\sin(b*x + a) - 1)))/b$

Mupad [B]

time = 1.22, size = 69, normalized size = 2.03

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 - 2 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1 \right)} - \frac{\text{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/cos(a + b*x)^3,x)

[Out] $(\tan(a/2 + (b*x)/2) + \tan(a/2 + (b*x)/2)^3)/(b*(\tan(a/2 + (b*x)/2)^4 - 2*\tan(a/2 + (b*x)/2)^2 + 1) - \text{atanh}(\tan(a/2 + (b*x)/2))/b$

3.64 $\int \sec^3(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=55

$$-\frac{\tanh^{-1}(\sin(a + bx))}{8b} - \frac{\sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec^3(a + bx) \tan(a + bx)}{4b}$$

[Out] $-1/8*\operatorname{arctanh}(\sin(b*x+a))/b-1/8*\sec(b*x+a)*\tan(b*x+a)/b+1/4*\sec(b*x+a)^3*\tan(b*x+a)/b$

Rubi [A]

time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2691, 3853, 3855}

$$-\frac{\tanh^{-1}(\sin(a + bx))}{8b} + \frac{\tan(a + bx) \sec^3(a + bx)}{4b} - \frac{\tan(a + bx) \sec(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sec}[a + b*x]^3*\operatorname{Tan}[a + b*x]^2, x]$

[Out] $-1/8*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]]/b - (\operatorname{Sec}[a + b*x]*\operatorname{Tan}[a + b*x])/(8*b) + (\operatorname{Sec}[a + b*x]^3*\operatorname{Tan}[a + b*x])/(4*b)$

Rule 2691

$\operatorname{Int}[(a_*)*\sec[(e_*) + (f_*)(x_)]^{(m_*)}*((b_*)*\tan[(e_*) + (f_*)(x_)]^{(n_*)}, x_Symbol] :> \operatorname{Simp}[b*(a*\operatorname{Sec}[e + f*x])^m*((b*\operatorname{Tan}[e + f*x])^{(n-1)})/(f*(m+n-1)), x] - \operatorname{Dist}[b^2*((n-1)/(m+n-1)), \operatorname{Int}[(a*\operatorname{Sec}[e + f*x])^m*(b*\operatorname{Tan}[e + f*x])^{(n-2)}, x], x] /; \operatorname{FreeQ}\{a, b, e, f, m\}, x] \&\& \operatorname{GtQ}[n, 1] \&\& \operatorname{NeQ}[m+n-1, 0] \&\& \operatorname{IntegersQ}[2*m, 2*n]$

Rule 3853

$\operatorname{Int}[(\operatorname{csc}[c_*) + (d_*)(x_)]*(b_*)^{(n_*)}, x_Symbol] :> \operatorname{Simp}[(-b)*\operatorname{Cos}[c + d*x]*(b*\operatorname{Csc}[c + d*x])^{(n-1)})/(d*(n-1)), x] + \operatorname{Dist}[b^2*((n-2)/(n-1)), \operatorname{Int}[(b*\operatorname{Csc}[c + d*x])^{(n-2)}, x], x] /; \operatorname{FreeQ}\{b, c, d\}, x] \&\& \operatorname{GtQ}[n, 1] \&\& \operatorname{IntegerQ}[2*n]$

Rule 3855

$\operatorname{Int}[\operatorname{csc}[c_*) + (d_*)(x_)], x_Symbol] :> \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]]/d, x] /; \operatorname{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned}
\int \sec^3(a+bx) \tan^2(a+bx) dx &= \frac{\sec^3(a+bx) \tan(a+bx)}{4b} - \frac{1}{4} \int \sec^3(a+bx) dx \\
&= -\frac{\sec(a+bx) \tan(a+bx)}{8b} + \frac{\sec^3(a+bx) \tan(a+bx)}{4b} - \frac{1}{8} \int \sec(a+bx) dx \\
&= -\frac{\tanh^{-1}(\sin(a+bx))}{8b} - \frac{\sec(a+bx) \tan(a+bx)}{8b} + \frac{\sec^3(a+bx) \tan(a+bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 1.00

$$-\frac{\tanh^{-1}(\sin(a+bx))}{8b} - \frac{\sec(a+bx) \tan(a+bx)}{8b} + \frac{\sec^3(a+bx) \tan(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^3*Tan[a + b*x]^2,x]``[Out] -1/8*ArcTanh[Sin[a + b*x]]/b - (Sec[a + b*x]*Tan[a + b*x])/(8*b) + (Sec[a + b*x]^3*Tan[a + b*x])/(4*b)`**Maple [A]**

time = 0.07, size = 66, normalized size = 1.20

method	result	size
derivativedivides	$\frac{\frac{\sin^3(bx+a)}{4 \cos(bx+a)^4} + \frac{\sin^3(bx+a)}{8 \cos(bx+a)^2} + \frac{\sin(bx+a)}{8} - \frac{\ln(\sec(bx+a) + \tan(bx+a))}{8}}{b}$	66
default	$\frac{\frac{\sin^3(bx+a)}{4 \cos(bx+a)^4} + \frac{\sin^3(bx+a)}{8 \cos(bx+a)^2} + \frac{\sin(bx+a)}{8} - \frac{\ln(\sec(bx+a) + \tan(bx+a))}{8}}{b}$	66
risch	$\frac{i(e^{7i(bx+a)} - 7e^{5i(bx+a)} + 7e^{3i(bx+a)} - e^{i(bx+a)})}{4b(e^{2i(bx+a)} + 1)^4} + \frac{\ln(e^{i(bx+a)} - i)}{8b} - \frac{\ln(e^{i(bx+a)} + i)}{8b}$	100
norman	$\frac{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{4b} + \frac{7\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{7\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)}{4b} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{8b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{8b}$	110

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^5*sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/4*sin(b*x+a)^3/cos(b*x+a)^4+1/8*sin(b*x+a)^3/cos(b*x+a)^2+1/8*sin(b*x+a)-1/8*ln(sec(b*x+a)+tan(b*x+a)))`**Maxima [A]**

time = 0.30, size = 65, normalized size = 1.18

$$\frac{2\left(\sin(bx+a)^3 + \sin(bx+a)\right)}{\sin(bx+a)^4 - 2\sin(bx+a)^2 + 1} - \log(\sin(bx+a) + 1) + \log(\sin(bx+a) - 1)$$

16b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^2,x, algorithm="maxima")

[Out] 1/16*(2*(sin(b*x + a)^3 + sin(b*x + a))/(sin(b*x + a)^4 - 2*sin(b*x + a)^2 + 1) - log(sin(b*x + a) + 1) + log(sin(b*x + a) - 1))/b

Fricas [A]

time = 0.42, size = 71, normalized size = 1.29

$$\frac{\cos(bx+a)^4 \log(\sin(bx+a)+1) - \cos(bx+a)^4 \log(-\sin(bx+a)+1) + 2(\cos(bx+a)^2 - 2)\sin(bx+a)}{16b \cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/16*(cos(b*x + a)^4*log(sin(b*x + a) + 1) - cos(b*x + a)^4*log(-sin(b*x + a) + 1) + 2*(cos(b*x + a)^2 - 2)*sin(b*x + a))/(b*cos(b*x + a)^4)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^2(a + bx) \sec^5(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**5*sin(b*x+a)**2,x)

[Out] Integral(sin(a + b*x)**2*sec(a + b*x)**5, x)

Giac [A]

time = 4.25, size = 82, normalized size = 1.49

$$\frac{4 \left(\frac{1}{\sin(bx+a)} + \sin(bx+a) \right) - \log \left(\left| \frac{1}{\sin(bx+a)} + \sin(bx+a) + 2 \right| \right) + \log \left(\left| \frac{1}{\sin(bx+a)} + \sin(bx+a) - 2 \right| \right)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/32*(4*(1/sin(b*x + a) + sin(b*x + a))/((1/sin(b*x + a) + sin(b*x + a))^2 - 4) - log(abs(1/sin(b*x + a) + sin(b*x + a) + 2)) + log(abs(1/sin(b*x + a) + sin(b*x + a) - 2)))/b

Mupad [B]

time = 6.54, size = 125, normalized size = 2.27

$$b \left(\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^7}{4} + \frac{7 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5}{4} + \frac{7 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3}{4} + \frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{4} - \frac{\operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{4b} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(a + b*x)^2/cos(a + b*x)^5,x)
```

```
[Out] (tan(a/2 + (b*x)/2)/4 + (7*tan(a/2 + (b*x)/2)^3)/4 + (7*tan(a/2 + (b*x)/2)^5)/4 + tan(a/2 + (b*x)/2)^7/4)/(b*(6*tan(a/2 + (b*x)/2)^4 - 4*tan(a/2 + (b*x)/2)^2 - 4*tan(a/2 + (b*x)/2)^6 + tan(a/2 + (b*x)/2)^8 + 1)) - atanh(tan(a/2 + (b*x)/2))/(4*b)
```

3.65 $\int \sec^5(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=76

$$\frac{\tanh^{-1}(\sin(a + bx))}{16b} - \frac{\sec(a + bx) \tan(a + bx)}{16b} - \frac{\sec^3(a + bx) \tan(a + bx)}{24b} + \frac{\sec^5(a + bx) \tan(a + bx)}{6b}$$

[Out] -1/16*arctanh(sin(b*x+a))/b-1/16*sec(b*x+a)*tan(b*x+a)/b-1/24*sec(b*x+a)^3*tan(b*x+a)/b+1/6*sec(b*x+a)^5*tan(b*x+a)/b

Rubi [A]

time = 0.04, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2691, 3853, 3855}

$$\frac{\tanh^{-1}(\sin(a + bx))}{16b} + \frac{\tan(a + bx) \sec^5(a + bx)}{6b} - \frac{\tan(a + bx) \sec^3(a + bx)}{24b} - \frac{\tan(a + bx) \sec(a + bx)}{16b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^5*Tan[a + b*x]^2,x]

[Out] -1/16*ArcTanh[Sin[a + b*x]]/b - (Sec[a + b*x]*Tan[a + b*x])/(16*b) - (Sec[a + b*x]^3*Tan[a + b*x])/(24*b) + (Sec[a + b*x]^5*Tan[a + b*x])/(6*b)

Rule 2691

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sec^5(a+bx) \tan^2(a+bx) dx &= \frac{\sec^5(a+bx) \tan(a+bx)}{6b} - \frac{1}{6} \int \sec^5(a+bx) dx \\
&= -\frac{\sec^3(a+bx) \tan(a+bx)}{24b} + \frac{\sec^5(a+bx) \tan(a+bx)}{6b} - \frac{1}{8} \int \sec^3(a+bx) dx \\
&= -\frac{\sec(a+bx) \tan(a+bx)}{16b} - \frac{\sec^3(a+bx) \tan(a+bx)}{24b} + \frac{\sec^5(a+bx) \tan(a+bx)}{6b} \\
&= -\frac{\tanh^{-1}(\sin(a+bx))}{16b} - \frac{\sec(a+bx) \tan(a+bx)}{16b} - \frac{\sec^3(a+bx) \tan(a+bx)}{24b}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 76, normalized size = 1.00

$$-\frac{\tanh^{-1}(\sin(a+bx))}{16b} - \frac{\sec(a+bx) \tan(a+bx)}{16b} - \frac{\sec^3(a+bx) \tan(a+bx)}{24b} + \frac{\sec^5(a+bx) \tan(a+bx)}{6b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^5*Tan[a + b*x]^2,x]`

```
[Out] -1/16*ArcTanh[Sin[a + b*x]]/b - (Sec[a + b*x]*Tan[a + b*x])/(16*b) - (Sec[a + b*x]^3*Tan[a + b*x])/(24*b) + (Sec[a + b*x]^5*Tan[a + b*x])/(6*b)
```

Maple [A]

time = 0.08, size = 84, normalized size = 1.11

method	result
derivativedivides	$\frac{\frac{\sin^3(bx+a)}{6 \cos(bx+a)^6} + \frac{\sin^3(bx+a)}{8 \cos(bx+a)^4} + \frac{\sin^3(bx+a)}{16 \cos(bx+a)^2} + \frac{\sin(bx+a)}{16} - \frac{\ln(\sec(bx+a)+\tan(bx+a))}{16}}{b}$
default	$\frac{\frac{\sin^3(bx+a)}{6 \cos(bx+a)^6} + \frac{\sin^3(bx+a)}{8 \cos(bx+a)^4} + \frac{\sin^3(bx+a)}{16 \cos(bx+a)^2} + \frac{\sin(bx+a)}{16} - \frac{\ln(\sec(bx+a)+\tan(bx+a))}{16}}{b}$
risch	$\frac{i(3e^{11i(bx+a)}+17e^{9i(bx+a)}-114e^{7i(bx+a)}+114e^{5i(bx+a)}-17e^{3i(bx+a)}-3e^{i(bx+a)})}{24b(e^{2i(bx+a)}+1)^6} + \frac{\ln(e^{i(bx+a)}-i)}{16b} - \frac{\ln(e^{i(bx+a)}+i)}{16b}$
norman	$\frac{\tan\left(\frac{bx}{2}+\frac{a}{2}\right)}{8b} + \frac{47\left(\tan^3\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{24b} + \frac{13\left(\tan^5\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{4b} + \frac{13\left(\tan^7\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{4b} + \frac{47\left(\tan^9\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{24b} + \frac{\tan^{11}\left(\frac{bx}{2}+\frac{a}{2}\right)}{8b} + \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{16b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^7*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/6*sin(b*x+a)^3/cos(b*x+a)^6+1/8*sin(b*x+a)^3/cos(b*x+a)^4+1/16*sin(b*x+a)^3/cos(b*x+a)^2+1/16*sin(b*x+a)-1/16*ln(sec(b*x+a)+tan(b*x+a)))
```


Maxima [A]

time = 0.28, size = 91, normalized size = 1.20

$$\frac{2 \left(3 \sin(bx+a)^5 - 8 \sin(bx+a)^3 - 3 \sin(bx+a) \right)}{\sin(bx+a)^6 - 3 \sin(bx+a)^4 + 3 \sin(bx+a)^2 - 1} - 3 \log(\sin(bx+a) + 1) + 3 \log(\sin(bx+a) - 1)$$

$$96 b$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^2,x, algorithm="maxima")`

```
[Out] 1/96*(2*(3*sin(b*x + a)^5 - 8*sin(b*x + a)^3 - 3*sin(b*x + a))/(sin(b*x + a)^6 - 3*sin(b*x + a)^4 + 3*sin(b*x + a)^2 - 1) - 3*log(sin(b*x + a) + 1) + 3*log(sin(b*x + a) - 1))/b
```

Fricas [A]

time = 0.40, size = 84, normalized size = 1.11

$$\frac{3 \cos(bx+a)^6 \log(\sin(bx+a) + 1) - 3 \cos(bx+a)^6 \log(-\sin(bx+a) + 1) + 2(3 \cos(bx+a)^4 + 2 \cos(bx+a)^2 - 8) \sin(bx+a)}{96 b \cos(bx+a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^2,x, algorithm="fricas")`

```
[Out] -1/96*(3*cos(b*x + a)^6*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^6*log(-sin(b*x + a) + 1) + 2*(3*cos(b*x + a)^4 + 2*cos(b*x + a)^2 - 8)*sin(b*x + a))/(b*cos(b*x + a)^6)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**7*sin(b*x+a)**2,x)`

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep
```

Giac [A]

time = 3.68, size = 73, normalized size = 0.96

$$\frac{2 \left(3 \sin(bx+a)^5 - 8 \sin(bx+a)^3 - 3 \sin(bx+a) \right)}{(\sin(bx+a)^2 - 1)^3} - 3 \log(|\sin(bx+a) + 1|) + 3 \log(|\sin(bx+a) - 1|)$$

$$96 b$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^2,x, algorithm="giac")`

[Out] $\frac{1}{96} \cdot (2 \cdot (3 \cdot \sin(bx + a)^5 - 8 \cdot \sin(bx + a)^3 - 3 \cdot \sin(bx + a)) / (\sin(bx + a)^2 - 1)^3 - 3 \cdot \log(\text{abs}(\sin(bx + a) + 1)) + 3 \cdot \log(\text{abs}(\sin(bx + a) - 1))) / b$

Mupad [B]

time = 7.33, size = 177, normalized size = 2.33

$$\frac{\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{11}}{8} + \frac{47 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^9}{24} + \frac{13 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^7}{4} + \frac{13 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5}{4} + \frac{47 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3}{24} + \frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{8}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{12} - 6 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{10} + 15 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 - 20 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + 15 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 - 6 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1 \right)} - \frac{\text{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^2/cos(a + b*x)^7,x)`

[Out] $\frac{(\tan(a/2 + (b*x)/2)/8 + (47*\tan(a/2 + (b*x)/2)^3)/24 + (13*\tan(a/2 + (b*x)/2)^5)/4 + (13*\tan(a/2 + (b*x)/2)^7)/4 + (47*\tan(a/2 + (b*x)/2)^9)/24 + \tan(a/2 + (b*x)/2)^{11}/8) / (b*(15*\tan(a/2 + (b*x)/2)^4 - 6*\tan(a/2 + (b*x)/2)^2 - 20*\tan(a/2 + (b*x)/2)^6 + 15*\tan(a/2 + (b*x)/2)^8 - 6*\tan(a/2 + (b*x)/2)^{10} + \tan(a/2 + (b*x)/2)^{12} + 1)) - \text{atanh}(\tan(a/2 + (b*x)/2)) / (8*b)}$

3.66 $\int \cos^5(a + bx) \sin^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\cos^6(a + bx)}{6b} + \frac{\cos^8(a + bx)}{8b}$$

[Out] $-1/6*\cos(b*x+a)^6/b+1/8*\cos(b*x+a)^8/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 14}

$$\frac{\cos^8(a + bx)}{8b} - \frac{\cos^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^5*\text{Sin}[a + b*x]^3, x]$

[Out] $-1/6*\text{Cos}[a + b*x]^6/b + \text{Cos}[a + b*x]^8/(8*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m, x\} \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{InverseFunctionQ}[v]]$

Rule 2645

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_.)}*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{(n-1)/2}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m, x\} \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx) \sin^3(a + bx) dx &= -\frac{\text{Subst}(\int x^5(1 - x^2) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\text{Subst}(\int (x^5 - x^7) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\cos^6(a + bx)}{6b} + \frac{\cos^8(a + bx)}{8b} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 48, normalized size = 1.55

$$\frac{-72 \cos(2(a + bx)) - 12 \cos(4(a + bx)) + 8 \cos(6(a + bx)) + 3 \cos(8(a + bx))}{3072b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^5*Sin[a + b*x]^3,x]``[Out] (-72*Cos[2*(a + b*x)] - 12*Cos[4*(a + b*x)] + 8*Cos[6*(a + b*x)] + 3*Cos[8*(a + b*x)])/(3072*b)`**Maple [A]**

time = 0.10, size = 34, normalized size = 1.10

method	result	size
derivativedivides	$\frac{-\frac{(\cos^6(bx+a))(\sin^2(bx+a))}{8} - \frac{(\cos^6(bx+a))}{24}}{b}$	34
default	$\frac{-\frac{(\cos^6(bx+a))(\sin^2(bx+a))}{8} - \frac{(\cos^6(bx+a))}{24}}{b}$	34
risch	$\frac{\cos(8bx+8a)}{1024b} + \frac{\cos(6bx+6a)}{384b} - \frac{\cos(4bx+4a)}{256b} - \frac{3 \cos(2bx+2a)}{128b}$	58
norman	$\frac{4\left(\tan^4\left(\frac{bx+a}{2}\right)\right)}{b} + \frac{4\left(\tan^{12}\left(\frac{bx+a}{2}\right)\right)}{b} - \frac{16\left(\tan^6\left(\frac{bx+a}{2}\right)\right)}{3b} - \frac{16\left(\tan^{10}\left(\frac{bx+a}{2}\right)\right)}{3b} + \frac{40\left(\tan^8\left(\frac{bx+a}{2}\right)\right)}{3b}$ $\left(1 + \tan^2\left(\frac{bx+a}{2}\right)\right)^8$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5*sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/8*cos(b*x+a)^6*sin(b*x+a)^2-1/24*cos(b*x+a)^6)`**Maxima [A]**

time = 0.29, size = 36, normalized size = 1.16

$$\frac{3 \sin(bx + a)^8 - 8 \sin(bx + a)^6 + 6 \sin(bx + a)^4}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5*sin(b*x+a)^3,x, algorithm="maxima")``[Out] 1/24*(3*sin(b*x + a)^8 - 8*sin(b*x + a)^6 + 6*sin(b*x + a)^4)/b`**Fricas [A]**

time = 0.37, size = 26, normalized size = 0.84

$$\frac{3 \cos(bx + a)^8 - 4 \cos(bx + a)^6}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^5*sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $1/24*(3*\cos(b*x + a)^8 - 4*\cos(b*x + a)^6)/b$

Sympy [A]

time = 0.86, size = 44, normalized size = 1.42

$$\begin{cases} -\frac{\sin^2(a+bx)\cos^6(a+bx)}{6b} - \frac{\cos^8(a+bx)}{24b} & \text{for } b \neq 0 \\ x \sin^3(a) \cos^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**5*sin(b*x+a)**3,x)`

[Out] `Piecewise((-sin(a + b*x)**2*cos(a + b*x)**6/(6*b) - cos(a + b*x)**8/(24*b), Ne(b, 0)), (x*sin(a)**3*cos(a)**5, True))`

Giac [A]

time = 3.63, size = 27, normalized size = 0.87

$$\frac{\cos(bx + a)^8}{8b} - \frac{\cos(bx + a)^6}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^5*sin(b*x+a)^3,x, algorithm="giac")`

[Out] $1/8*\cos(b*x + a)^8/b - 1/6*\cos(b*x + a)^6/b$

Mupad [B]

time = 0.37, size = 25, normalized size = 0.81

$$\frac{\cos(a + bx)^6 (3 \cos(a + bx)^2 - 4)}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^5*sin(a + b*x)^3,x)`

[Out] $(\cos(a + b*x)^6*(3*\cos(a + b*x)^2 - 4))/(24*b)$

3.67 $\int \cos^4(a + bx) \sin^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\cos^5(a + bx)}{5b} + \frac{\cos^7(a + bx)}{7b}$$

[Out] $-1/5*\cos(b*x+a)^5/b+1/7*\cos(b*x+a)^7/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 14}

$$\frac{\cos^7(a + bx)}{7b} - \frac{\cos^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^4*\text{Sin}[a + b*x]^3, x]$

[Out] $-1/5*\text{Cos}[a + b*x]^5/b + \text{Cos}[a + b*x]^7/(7*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x] \&\& \text{SumQ}[u] \&\& !\text{LinearQ}[u, x] \&\& !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b\}, x] \&\& \text{InverseFunctionQ}[v]$

Rule 2645

$\text{Int}[(\cos[(e_.) + (f_)*(x_)]*(a_))^{(m_)}*\sin[(e_.) + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{((n - 1)/2)}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n - 1)/2] \&\& !(IntegerQ[(m - 1)/2] \&\& \text{GtQ}[m, 0] \&\& \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int \cos^4(a + bx) \sin^3(a + bx) dx &= -\frac{\text{Subst}(\int x^4(1 - x^2) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\text{Subst}(\int (x^4 - x^6) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\cos^5(a + bx)}{5b} + \frac{\cos^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 27, normalized size = 0.87

$$\frac{\cos^5(a + bx)(-9 + 5 \cos(2(a + bx)))}{70b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4*Sin[a + b*x]^3,x]

[Out] (Cos[a + b*x]^5*(-9 + 5*Cos[2*(a + b*x)]))/(70*b)

Maple [A]

time = 0.09, size = 34, normalized size = 1.10

method	result	size
derivativedivides	$\frac{\frac{(\cos^5(bx+a))(\sin^2(bx+a))}{7} - \frac{2(\cos^5(bx+a))}{35}}{b}$	34
default	$\frac{\frac{(\cos^5(bx+a))(\sin^2(bx+a))}{7} - \frac{2(\cos^5(bx+a))}{35}}{b}$	34
risch	$-\frac{3 \cos(bx+a)}{64b} + \frac{\cos(7bx+7a)}{448b} + \frac{\cos(5bx+5a)}{320b} - \frac{\cos(3bx+3a)}{64b}$	55
norman	$\frac{-\frac{4(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{8(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{4(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{4}{35b} - \frac{4(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{5b} + \frac{8(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{5b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^7}$	103

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/7*cos(b*x+a)^5*sin(b*x+a)^2-2/35*cos(b*x+a)^5)

Maxima [A]

time = 0.29, size = 26, normalized size = 0.84

$$\frac{5 \cos(bx + a)^7 - 7 \cos(bx + a)^5}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^3,x, algorithm="maxima")

[Out] 1/35*(5*cos(b*x + a)^7 - 7*cos(b*x + a)^5)/b

Fricas [A]

time = 0.35, size = 26, normalized size = 0.84

$$\frac{5 \cos(bx + a)^7 - 7 \cos(bx + a)^5}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/35*(5*cos(b*x + a)^7 - 7*cos(b*x + a)^5)/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(22) = 44$.

time = 0.58, size = 46, normalized size = 1.48

$$\begin{cases} -\frac{\sin^2(a+bx)\cos^5(a+bx)}{5b} - \frac{2\cos^7(a+bx)}{35b} & \text{for } b \neq 0 \\ x \sin^3(a) \cos^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**4*sin(b*x+a)**3,x)

[Out] Piecewise((-sin(a + b*x)**2*cos(a + b*x)**5/(5*b) - 2*cos(a + b*x)**7/(35*b), Ne(b, 0)), (x*sin(a)**3*cos(a)**4, True))

Giac [A]

time = 3.35, size = 27, normalized size = 0.87

$$\frac{\cos(bx + a)^7}{7b} - \frac{\cos(bx + a)^5}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/7*cos(b*x + a)^7/b - 1/5*cos(b*x + a)^5/b

Mupad [B]

time = 0.38, size = 26, normalized size = 0.84

$$-\frac{7\cos(a + bx)^5 - 5\cos(a + bx)^7}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4*sin(a + b*x)^3,x)

[Out] -(7*cos(a + b*x)^5 - 5*cos(a + b*x)^7)/(35*b)

3.68 $\int \cos^3(a + bx) \sin^3(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\sin^4(a + bx)}{4b} - \frac{\sin^6(a + bx)}{6b}$$

[Out] $1/4*\sin(b*x+a)^4/b-1/6*\sin(b*x+a)^6/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 14}

$$\frac{\sin^4(a + bx)}{4b} - \frac{\sin^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^3*Sin[a + b*x]^3,x]`

[Out] `Sin[a + b*x]^4/(4*b) - Sin[a + b*x]^6/(6*b)`

Rule 14

```
Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2644

```
Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \sin^3(a + bx) dx &= \frac{\text{Subst}\left(\int x^3(1 - x^2) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^3 - x^5) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^4(a + bx)}{4b} - \frac{\sin^6(a + bx)}{6b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 35, normalized size = 1.13

$$\frac{1}{8} \left(-\frac{3 \cos(2(a + bx))}{8b} + \frac{\cos(6(a + bx))}{24b} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*Sin[a + b*x]^3,x]``[Out] ((-3*Cos[2*(a + b*x)])/(8*b) + Cos[6*(a + b*x)]/(24*b))/8`**Maple [A]**

time = 0.06, size = 34, normalized size = 1.10

method	result	size
risch	$\frac{\cos(6bx+6a)}{192b} - \frac{3 \cos(2bx+2a)}{64b}$	30
derivativedivides	$\frac{-(\cos^4(bx+a))(\sin^2(bx+a)) - (\cos^4(bx+a))}{6b}$	34
default	$\frac{-(\cos^4(bx+a))(\sin^2(bx+a)) - (\cos^4(bx+a))}{6b}$	34
norman	$\frac{4(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{4(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{8(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b}$ $\frac{1}{(1+\tan^2(\frac{bx}{2} + \frac{a}{2}))^6}$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^3*sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/6*cos(b*x+a)^4*sin(b*x+a)^2-1/12*cos(b*x+a)^4)`**Maxima [A]**

time = 0.28, size = 26, normalized size = 0.84

$$\frac{2 \sin(bx + a)^6 - 3 \sin(bx + a)^4}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^3*sin(b*x+a)^3,x, algorithm="maxima")``[Out] -1/12*(2*sin(b*x + a)^6 - 3*sin(b*x + a)^4)/b`**Fricas [A]**

time = 0.40, size = 26, normalized size = 0.84

$$\frac{2 \cos(bx + a)^6 - 3 \cos(bx + a)^4}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3*sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $1/12*(2*\cos(b*x + a)^6 - 3*\cos(b*x + a)^4)/b$

Sympy [A]

time = 0.38, size = 44, normalized size = 1.42

$$\begin{cases} -\frac{\sin^2(a+bx)\cos^4(a+bx)}{4b} - \frac{\cos^6(a+bx)}{12b} & \text{for } b \neq 0 \\ x \sin^3(a) \cos^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**3*sin(b*x+a)**3,x)`

[Out] `Piecewise((-sin(a + b*x)**2*cos(a + b*x)**4/(4*b) - cos(a + b*x)**6/(12*b), Ne(b, 0)), (x*sin(a)**3*cos(a)**3, True))`

Giac [A]

time = 3.91, size = 26, normalized size = 0.84

$$-\frac{2 \sin (bx + a)^6 - 3 \sin (bx + a)^4}{12 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3*sin(b*x+a)^3,x, algorithm="giac")`

[Out] $-1/12*(2*\sin(b*x + a)^6 - 3*\sin(b*x + a)^4)/b$

Mupad [B]

time = 0.51, size = 37, normalized size = 1.19

$$\frac{\cos(a + bx)^4 (\cos(a + bx)^2 - 1)}{4b} - \frac{\cos(a + bx)^6}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^3*sin(a + b*x)^3,x)`

[Out] $(\cos(a + b*x)^4*(\cos(a + b*x)^2 - 1))/(4*b) - \cos(a + b*x)^6/(12*b)$

3.69 $\int \cos^2(a + bx) \sin^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\cos^3(a + bx)}{3b} + \frac{\cos^5(a + bx)}{5b}$$

[Out] $-1/3*\cos(b*x+a)^3/b+1/5*\cos(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 14}

$$\frac{\cos^5(a + bx)}{5b} - \frac{\cos^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^2*\text{Sin}[a + b*x]^3, x]$

[Out] $-1/3*\text{Cos}[a + b*x]^3/b + \text{Cos}[a + b*x]^5/(5*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_.)}*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}), x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{((n - 1)/2)}, x], x, a*\text{Cos}[e + f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) \sin^3(a + bx) dx &= -\frac{\text{Subst}(\int x^2(1 - x^2) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\text{Subst}(\int (x^2 - x^4) dx, x, \cos(a + bx))}{b} \\ &= -\frac{\cos^3(a + bx)}{3b} + \frac{\cos^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 27, normalized size = 0.87

$$\frac{\cos^3(a + bx)(-7 + 3 \cos(2(a + bx)))}{30b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^2*Sin[a + b*x]^3,x]

[Out] (Cos[a + b*x]^3*(-7 + 3*Cos[2*(a + b*x)]))/(30*b)

Maple [A]

time = 0.05, size = 34, normalized size = 1.10

method	result	size
derivativedivides	$-\frac{(\cos^3(bx+a))(\sin^2(bx+a))}{5} - \frac{2(\cos^3(bx+a))}{15}$	34
default	$-\frac{(\cos^3(bx+a))(\sin^2(bx+a))}{5} - \frac{2(\cos^3(bx+a))}{15}$	34
risch	$-\frac{\cos(bx+a)}{8b} + \frac{\cos(5bx+5a)}{80b} - \frac{\cos(3bx+3a)}{48b}$	41
norman	$-\frac{\frac{4}{15b} - \frac{4(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{4(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{4(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^5}$	71

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/5*cos(b*x+a)^3*sin(b*x+a)^2-2/15*cos(b*x+a)^3)

Maxima [A]

time = 0.30, size = 26, normalized size = 0.84

$$\frac{3 \cos(bx + a)^5 - 5 \cos(bx + a)^3}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^3,x, algorithm="maxima")

[Out] 1/15*(3*cos(b*x + a)^5 - 5*cos(b*x + a)^3)/b

Fricas [A]

time = 0.36, size = 26, normalized size = 0.84

$$\frac{3 \cos(bx + a)^5 - 5 \cos(bx + a)^3}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/15*(3*cos(b*x + a)^5 - 5*cos(b*x + a)^3)/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 46 vs. $2(22) = 44$.

time = 0.25, size = 46, normalized size = 1.48

$$\begin{cases} -\frac{\sin^2(a+bx)\cos^3(a+bx)}{3b} - \frac{2\cos^5(a+bx)}{15b} & \text{for } b \neq 0 \\ x \sin^3(a) \cos^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2*sin(b*x+a)**3,x)

[Out] Piecewise((-sin(a + b*x)**2*cos(a + b*x)**3/(3*b) - 2*cos(a + b*x)**5/(15*b), Ne(b, 0)), (x*sin(a)**3*cos(a)**2, True))

Giac [A]

time = 5.59, size = 27, normalized size = 0.87

$$\frac{\cos(bx + a)^5}{5b} - \frac{\cos(bx + a)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/5*cos(b*x + a)^5/b - 1/3*cos(b*x + a)^3/b

Mupad [B]

time = 0.34, size = 26, normalized size = 0.84

$$-\frac{5\cos(a + bx)^3 - 3\cos(a + bx)^5}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2*sin(a + b*x)^3,x)

[Out] -(5*cos(a + b*x)^3 - 3*cos(a + b*x)^5)/(15*b)

3.70 $\int \cos(a + bx) \sin^3(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sin^4(a + bx)}{4b}$$

[Out] 1/4*sin(b*x+a)^4/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2644, 30}

$$\frac{\sin^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Sin[a + b*x]^3,x]

[Out] Sin[a + b*x]^4/(4*b)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \sin^3(a + bx) dx &= \frac{\text{Subst}\left(\int x^3 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$\frac{\sin^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Sin[a + b*x]^3,x]

[Out] Sin[a + b*x]^4/(4*b)

Maple [A]

time = 0.04, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$\frac{\sin^4(bx+a)}{4b}$	14
default	$\frac{\sin^4(bx+a)}{4b}$	14
risch	$\frac{\cos(4bx+4a)}{32b} - \frac{\cos(2bx+2a)}{8b}$	30
norman	$\frac{4\left(\tan^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b\left(1+\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)^4}$	32

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/4*sin(b*x+a)^4/b

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$\frac{\sin(bx+a)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a)^3,x, algorithm="maxima")

[Out] 1/4*sin(b*x + a)^4/b

Fricas [A]

time = 0.37, size = 24, normalized size = 1.60

$$\frac{\cos(bx+a)^4 - 2\cos(bx+a)^2}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/4*(cos(b*x + a)^4 - 2*cos(b*x + a)^2)/b

Sympy [A]

time = 0.15, size = 20, normalized size = 1.33

$$\begin{cases} \frac{\sin^4(a+bx)}{4b} & \text{for } b \neq 0 \\ x \sin^3(a) \cos(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)**3,x)``[Out] Piecewise((sin(a + b*x)**4/(4*b), Ne(b, 0)), (x*sin(a)**3*cos(a), True))`**Giac [A]**

time = 4.47, size = 13, normalized size = 0.87

$$\frac{\sin(bx + a)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)^3,x, algorithm="giac")``[Out] 1/4*sin(b*x + a)^4/b`**Mupad [B]**

time = 0.37, size = 13, normalized size = 0.87

$$\frac{\sin(a + bx)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)*sin(a + b*x)^3,x)``[Out] sin(a + b*x)^4/(4*b)`

3.71 $\int \sin^2(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=28

$$\frac{\cos^2(a + bx)}{2b} - \frac{\log(\cos(a + bx))}{b}$$

[Out] 1/2*cos(b*x+a)^2/b-ln(cos(b*x+a))/b

Rubi [A]

time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 14}

$$\frac{\cos^2(a + bx)}{2b} - \frac{\log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2*Tan[a + b*x],x]

[Out] Cos[a + b*x]^2/(2*b) - Log[Cos[a + b*x]]/b

Rule 14

```
Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2670

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \sin^2(a + bx) \tan(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x} - x\right) dx, x, \cos(a + bx)\right)}{b} \\ &= \frac{\cos^2(a + bx)}{2b} - \frac{\log(\cos(a + bx))}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 25, normalized size = 0.89

$$-\frac{\frac{1}{2} \cos^2(a + bx) + \log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^2*Tan[a + b*x],x]
```

```
[Out] -((-1/2*Cos[a + b*x]^2 + Log[Cos[a + b*x]])/b)
```

Maple [A]

time = 0.04, size = 25, normalized size = 0.89

method	result	size
derivativedivides	$-\frac{\frac{\sin^2(bx+a)}{2} - \ln(\cos(bx+a))}{b}$	25
default	$-\frac{\frac{\sin^2(bx+a)}{2} - \ln(\cos(bx+a))}{b}$	25
risch	$ix + \frac{e^{2i(bx+a)}}{8b} + \frac{e^{-2i(bx+a)}}{8b} + \frac{2ia}{b} - \frac{\ln(e^{2i(bx+a)}+1)}{b}$	58
norman	$-\frac{2\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b\left(1+\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)^2} + \frac{\ln\left(1+\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{b}$	85

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(b*x+a)*sin(b*x+a)^3,x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(-1/2*sin(b*x+a)^2-ln(cos(b*x+a)))
```

Maxima [A]

time = 0.30, size = 25, normalized size = 0.89

$$-\frac{\sin^2(bx + a) + \log(\sin^2(bx + a) - 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)*sin(b*x+a)^3,x, algorithm="maxima")
```

```
[Out] -1/2*(sin(b*x + a)^2 + log(sin(b*x + a)^2 - 1))/b
```

Fricas [A]

time = 0.43, size = 25, normalized size = 0.89

$$\frac{\cos^2(bx + a) - 2 \log(-\cos(bx + a))}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/2*(cos(b*x + a)^2 - 2*log(-cos(b*x + a)))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^3(a + bx) \sec(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)**3,x)

[Out] Integral(sin(a + b*x)**3*sec(a + b*x), x)

Giac [A]

time = 4.64, size = 29, normalized size = 1.04

$$\frac{\cos(bx + a)^2 - \log\left(\frac{\cos(bx+a)^2}{b^2}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/2*(cos(b*x + a)^2 - log(cos(b*x + a)^2/b^2))/b

Mupad [B]

time = 0.43, size = 25, normalized size = 0.89

$$\frac{\cos(a + bx)^2 + \ln(\tan(a + bx)^2 + 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/cos(a + b*x),x)

[Out] (log(tan(a + b*x)^2 + 1) + cos(a + b*x)^2)/(2*b)

3.72 $\int \sin(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=21

$$\frac{\cos(a + bx)}{b} + \frac{\sec(a + bx)}{b}$$

[Out] $\cos(b*x+a)/b+\sec(b*x+a)/b$

Rubi [A]

time = 0.01, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 14}

$$\frac{\cos(a + bx)}{b} + \frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[a + b*x]*\text{Tan}[a + b*x]^2, x]$

[Out] $\text{Cos}[a + b*x]/b + \text{Sec}[a + b*x]/b$

Rule 14

$\text{Int}[(u_)*((c_.)*(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_.)*(v_)) /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{InverseFunctionQ}[v]$

Rule 2670

$\text{Int}[\text{sin}[(e_.) + (f_.)*(x_)]^{(m_.)}*\text{tan}[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[-f^{(-1)}, \text{Subst}[\text{Int}[(1 - x^2)^{(m+n-1)/2}/x^n, x], x, \text{Cos}[e + f*x]], x] /; \text{FreeQ}\{e, f\}, x \ \&\& \ \text{IntegersQ}[m, n, (m+n-1)/2]$

Rubi steps

$$\begin{aligned} \int \sin(a + bx) \tan^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^2} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-1 + \frac{1}{x^2}\right) dx, x, \cos(a + bx)\right)}{b} \\ &= \frac{\cos(a + bx)}{b} + \frac{\sec(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 21, normalized size = 1.00

$$\frac{\cos(a + bx)}{b} + \frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]*Tan[a + b*x]^2,x]

[Out] Cos[a + b*x]/b + Sec[a + b*x]/b

Maple [A]

time = 0.04, size = 40, normalized size = 1.90

method	result	size
norman	$-\frac{4}{b\left(1+\tan^2\left(\frac{bx+a}{2}\right)\right)\left(\tan^2\left(\frac{bx+a}{2}\right)-1\right)}$	36
derivativedivides	$\frac{\frac{\sin^4(bx+a)}{\cos(bx+a)} + (2+\sin^2(bx+a)) \cos(bx+a)}{b}$	40
default	$\frac{\frac{\sin^4(bx+a)}{\cos(bx+a)} + (2+\sin^2(bx+a)) \cos(bx+a)}{b}$	40
risch	$\frac{e^{i(bx+a)}}{2b} + \frac{e^{-i(bx+a)}}{2b} + \frac{2e^{i(bx+a)}}{b(e^{2i(bx+a)}+1)}$	57

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/b*(sin(b*x+a)^4/cos(b*x+a)+(2+sin(b*x+a)^2)*cos(b*x+a))

Maxima [A]

time = 0.29, size = 19, normalized size = 0.90

$$\frac{\frac{1}{\cos(bx+a)} + \cos(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^3,x, algorithm="maxima")

[Out] (1/cos(b*x + a) + cos(b*x + a))/b

Fricas [A]

time = 0.36, size = 22, normalized size = 1.05

$$\frac{\cos(bx+a)^2 + 1}{b \cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^3,x, algorithm="fricas")

[Out] (cos(b*x + a)^2 + 1)/(b*cos(b*x + a))

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: HeuristicGCDFailed

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2*sin(b*x+a)**3,x)

[Out] Exception raised: HeuristicGCDFailed >> no luck

Giac [A]

time = 5.70, size = 23, normalized size = 1.10

$$\frac{\cos(bx + a)}{b} + \frac{1}{b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^3,x, algorithm="giac")

[Out] cos(b*x + a)/b + 1/(b*cos(b*x + a))

Mupad [B]

time = 0.49, size = 20, normalized size = 0.95

$$-\frac{4}{b \left(\tan \left(\frac{a}{2} + \frac{bx}{2} \right)^4 - 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/cos(a + b*x)^2,x)

[Out] -4/(b*(tan(a/2 + (b*x)/2)^4 - 1))

3.73 $\int \tan^3(a + bx) dx$

Optimal. Leaf size=27

$$\frac{\log(\cos(a + bx))}{b} + \frac{\tan^2(a + bx)}{2b}$$

[Out] $\ln(\cos(b*x+a))/b+1/2*\tan(b*x+a)^2/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$\frac{\tan^2(a + bx)}{2b} + \frac{\log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tan}[a + b*x]^3, x]$

[Out] $\text{Log}[\text{Cos}[a + b*x]]/b + \text{Tan}[a + b*x]^2/(2*b)$

Rule 3554

$\text{Int}[(b_*)*\tan[(c_*) + (d_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)})/(d*(n - 1))], x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] /;$ $\text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1]$

Rule 3556

$\text{Int}[\tan[(c_*) + (d_*)(x_)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \int \tan^3(a + bx) dx &= \frac{\tan^2(a + bx)}{2b} - \int \tan(a + bx) dx \\ &= \frac{\log(\cos(a + bx))}{b} + \frac{\tan^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 25, normalized size = 0.93

$$\frac{2 \log(\cos(a + bx)) + \tan^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Tan[a + b*x]^3,x]

[Out] (2*Log[Cos[a + b*x]] + Tan[a + b*x]^2)/(2*b)

Maple [A]

time = 0.04, size = 23, normalized size = 0.85

method	result	size
derivativedivides	$\frac{\frac{(\tan^2(bx+a))}{2} + \ln(\cos(bx+a))}{b}$	23
default	$\frac{(\tan^2(bx+a))}{2} + \ln(\cos(bx+a))}{b}$	23
risch	$-ix - \frac{2ia}{b} + \frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}+1)^2} + \frac{\ln(e^{2i(bx+a)}+1)}{b}$	56
norman	$\frac{2\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^2} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b} - \frac{\ln\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	84

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^3*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/b*(1/2*tan(b*x+a)^2+ln(cos(b*x+a)))

Maxima [A]

time = 0.28, size = 31, normalized size = 1.15

$$-\frac{\frac{1}{\sin(bx+a)^2-1} - \log(\sin(bx+a)^2-1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^3,x, algorithm="maxima")

[Out] -1/2*(1/(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2 - 1))/b

Fricas [A]

time = 0.40, size = 34, normalized size = 1.26

$$\frac{2 \cos(bx+a)^2 \log(-\cos(bx+a)) + 1}{2b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/2*(2*cos(b*x + a)^2*log(-cos(b*x + a)) + 1)/(b*cos(b*x + a)^2)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: HeuristicGCDFailed

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3*sin(b*x+a)**3,x)

[Out] Exception raised: HeuristicGCDFailed >> no luck

Giac [A]

time = 4.82, size = 42, normalized size = 1.56

$$\frac{\log\left(\frac{\cos(bx+a)^2}{b^2}\right)}{2b} - \frac{\cos(bx+a)^2 - 1}{2b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/2*log(cos(b*x + a)^2/b^2)/b - 1/2*(cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^2)

Mupad [B]

time = 0.38, size = 27, normalized size = 1.00

$$-\frac{\ln(\tan(a+bx)^2+1) - \tan(a+bx)^2}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/cos(a + b*x)^3,x)

[Out] -(log(tan(a + b*x)^2 + 1) - tan(a + b*x)^2)/(2*b)

3.74 $\int \sec(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=27

$$-\frac{\sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b}$$

[Out] `-sec(b*x+a)/b+1/3*sec(b*x+a)^3/b`

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {2686}

$$\frac{\sec^3(a + bx)}{3b} - \frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Sec[a + b*x]*Tan[a + b*x]^3,x]`

[Out] `-(Sec[a + b*x]/b) + Sec[a + b*x]^3/(3*b)`

Rule 2686

```
Int[((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Rubi steps

$$\begin{aligned} \int \sec(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}\left(\int (-1 + x^2) dx, x, \sec(a + bx)\right)}{b} \\ &= -\frac{\sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 27, normalized size = 1.00

$$-\frac{\sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] `Integrate[Sec[a + b*x]*Tan[a + b*x]^3,x]`

[Out] $-(\text{Sec}[a + b*x]/b) + \text{Sec}[a + b*x]^3/(3*b)$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(25) = 50$.

time = 0.05, size = 60, normalized size = 2.22

method	result	size
norman	$-\frac{4\left(\tan^2\left(\frac{bx+a}{2}\right)\right)}{b} + \frac{4}{3b}$ $\frac{1}{\left(\tan^2\left(\frac{bx+a}{2}\right)-1\right)^3}$	39
risch	$-\frac{2\left(3e^{5i(bx+a)}+2e^{3i(bx+a)}+3e^{i(bx+a)}\right)}{3b\left(e^{2i(bx+a)}+1\right)^3}$	53
derivativedivides	$\frac{\frac{\sin^4(bx+a)}{3\cos(bx+a)^3} - \frac{\sin^4(bx+a)}{3\cos(bx+a)} - \frac{(2+\sin^2(bx+a))\cos(bx+a)}{3}}{b}$	60
default	$\frac{\frac{\sin^4(bx+a)}{3\cos(bx+a)^3} - \frac{\sin^4(bx+a)}{3\cos(bx+a)} - \frac{(2+\sin^2(bx+a))\cos(bx+a)}{3}}{b}$	60

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^4*sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $1/b*(1/3*\sin(b*x+a)^4/\cos(b*x+a)^3-1/3*\sin(b*x+a)^4/\cos(b*x+a)-1/3*(2+\sin(b*x+a)^2)*\cos(b*x+a))$

Maxima [A]

time = 0.30, size = 25, normalized size = 0.93

$$-\frac{3\cos(bx+a)^2-1}{3b\cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4*sin(b*x+a)^3,x, algorithm="maxima")`

[Out] $-1/3*(3*\cos(b*x + a)^2 - 1)/(b*\cos(b*x + a)^3)$

Fricas [A]

time = 0.39, size = 25, normalized size = 0.93

$$-\frac{3\cos(bx+a)^2-1}{3b\cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4*sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $-1/3*(3*\cos(b*x + a)^2 - 1)/(b*\cos(b*x + a)^3)$

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4*sin(b*x+a)**3,x)

[Out] Timed out

Giac [A]

time = 4.58, size = 25, normalized size = 0.93

$$-\frac{3 \cos (bx+a)^2-1}{3 b \cos (bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^3,x, algorithm="giac")

[Out] -1/3*(3*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^3)

Mupad [B]

time = 0.45, size = 23, normalized size = 0.85

$$-\frac{\cos (a+b x)^2-\frac{1}{3}}{b \cos (a+b x)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/cos(a + b*x)^4,x)

[Out] -(cos(a + b*x)^2 - 1/3)/(b*cos(a + b*x)^3)

3.75 $\int \sec^2(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\tan^4(a + bx)}{4b}$$

[Out] 1/4*tan(b*x+a)^4/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 30}

$$\frac{\tan^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2*Tan[a + b*x]^3,x]

[Out] Tan[a + b*x]^4/(4*b)

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2687

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^2(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}\left(\int x^3 dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$\frac{\tan^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2*Tan[a + b*x]^3,x]

[Out] Tan[a + b*x]^4/(4*b)

Maple [A]

time = 0.06, size = 22, normalized size = 1.47

method	result	size
derivativedivides	$\frac{\sin^4(bx+a)}{4b \cos(bx+a)^4}$	22
default	$\frac{\sin^4(bx+a)}{4b \cos(bx+a)^4}$	22
norman	$\frac{4\left(\tan^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)^4}$	32
risch	$-\frac{2(e^{6i(bx+a)}+e^{2i(bx+a)})}{b(e^{2i(bx+a)}+1)^4}$	38

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^5*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] 1/4/b*sin(b*x+a)^4/cos(b*x+a)^4

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(13) = 26.

time = 0.29, size = 39, normalized size = 2.60

$$\frac{2 \sin (bx+a)^2-1}{4\left(\sin (bx+a)^4-2 \sin (bx+a)^2+1\right) b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^3,x, algorithm="maxima")

[Out] 1/4*(2*sin(b*x + a)^2 - 1)/((sin(b*x + a)^4 - 2*sin(b*x + a)^2 + 1)*b)

Fricas [A]

time = 0.35, size = 25, normalized size = 1.67

$$-\frac{2 \cos (bx+a)^2-1}{4 b \cos (bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^3,x, algorithm="fricas")

[Out] -1/4*(2*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^4)

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**5*sin(b*x+a)**3,x)

[Out] Timed out

Giac [A]

time = 5.14, size = 25, normalized size = 1.67

$$-\frac{2 \cos (bx + a)^2 - 1}{4 b \cos (bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^3,x, algorithm="giac")

[Out] -1/4*(2*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^4)

Mupad [B]

time = 0.38, size = 13, normalized size = 0.87

$$\frac{\tan(a + bx)^4}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/cos(a + b*x)^5,x)

[Out] tan(a + b*x)^4/(4*b)

3.76 $\int \sec^3(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\sec^3(a + bx)}{3b} + \frac{\sec^5(a + bx)}{5b}$$

[Out] $-1/3*\sec(b*x+a)^3/b+1/5*\sec(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 14}

$$\frac{\sec^5(a + bx)}{5b} - \frac{\sec^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^3*\text{Tan}[a + b*x]^3, x]$

[Out] $-1/3*\text{Sec}[a + b*x]^3/b + \text{Sec}[a + b*x]^5/(5*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{InverseFunctionQ}[v]$

Rule 2686

$\text{Int}[(a_)*\sec[(e_)+(f_)*(x_)]^{(m_)}*((b_)*\tan[(e_)+(f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a/f, \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \text{Sec}[e+f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n+1])$

Rubi steps

$$\begin{aligned} \int \sec^3(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}(\int x^2(-1 + x^2) dx, x, \sec(a + bx))}{b} \\ &= \frac{\text{Subst}(\int (-x^2 + x^4) dx, x, \sec(a + bx))}{b} \\ &= -\frac{\sec^3(a + bx)}{3b} + \frac{\sec^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 31, normalized size = 1.00

$$-\frac{\sec^3(a+bx)}{3b} + \frac{\sec^5(a+bx)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^3*Tan[a + b*x]^3,x]``[Out] -1/3*Sec[a + b*x]^3/b + Sec[a + b*x]^5/(5*b)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 77 vs. 2(27) = 54.

time = 0.06, size = 78, normalized size = 2.52

method	result	size
risch	$-\frac{8(5e^{7i(bx+a)} - 2e^{5i(bx+a)} + 5e^{3i(bx+a)})}{15b(e^{2i(bx+a)} + 1)^5}$	53
norman	$\frac{\frac{4}{15b} - \frac{4(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{4(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{4(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^5}$	71
derivativedivides	$\frac{\frac{\sin^4(bx+a)}{5 \cos(bx+a)^5} + \frac{\sin^4(bx+a)}{15 \cos(bx+a)^3} - \frac{\sin^4(bx+a)}{15 \cos(bx+a)} - \frac{(2 + \sin^2(bx+a)) \cos(bx+a)}{15}}{b}$	78
default	$\frac{\frac{\sin^4(bx+a)}{5 \cos(bx+a)^5} + \frac{\sin^4(bx+a)}{15 \cos(bx+a)^3} - \frac{\sin^4(bx+a)}{15 \cos(bx+a)} - \frac{(2 + \sin^2(bx+a)) \cos(bx+a)}{15}}{b}$	78

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^6*sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/5*sin(b*x+a)^4/cos(b*x+a)^5+1/15*sin(b*x+a)^4/cos(b*x+a)^3-1/15*sin(b*x+a)^4/cos(b*x+a)-1/15*(2+sin(b*x+a)^2)*cos(b*x+a))`**Maxima [A]**

time = 0.29, size = 25, normalized size = 0.81

$$-\frac{5 \cos(bx+a)^2 - 3}{15 b \cos(bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6*sin(b*x+a)^3,x, algorithm="maxima")``[Out] -1/15*(5*cos(b*x + a)^2 - 3)/(b*cos(b*x + a)^5)`**Fricas [A]**

time = 0.36, size = 25, normalized size = 0.81

$$-\frac{5 \cos(bx+a)^2 - 3}{15 b \cos(bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^6*sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $-1/15*(5*\cos(b*x + a)^2 - 3)/(b*\cos(b*x + a)^5)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**6*sin(b*x+a)**3,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [A]

time = 3.99, size = 25, normalized size = 0.81

$$-\frac{5 \cos (bx+a)^2-3}{15 b \cos (bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^6*sin(b*x+a)^3,x, algorithm="giac")`

[Out] $-1/15*(5*\cos(b*x + a)^2 - 3)/(b*\cos(b*x + a)^5)$

Mupad [B]

time = 0.54, size = 25, normalized size = 0.81

$$-\frac{\frac{\cos(a+bx)^2}{3} - \frac{1}{5}}{b \cos (a+bx)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^3/cos(a + b*x)^6,x)`

[Out] $-(\cos(a + b*x)^{2/3} - 1/5)/(b*\cos(a + b*x)^5)$

3.77 $\int \sec^4(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\sec^4(a + bx)}{4b} + \frac{\sec^6(a + bx)}{6b}$$

[Out] $-1/4*\sec(b*x+a)^4/b+1/6*\sec(b*x+a)^6/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 14}

$$\frac{\sec^6(a + bx)}{6b} - \frac{\sec^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^4*\text{Tan}[a + b*x]^3,x]$

[Out] $-1/4*\text{Sec}[a + b*x]^4/b + \text{Sec}[a + b*x]^6/(6*b)$

Rule 14

$\text{Int}[(u_*)((c_*)(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)] /; FreeQ[{a, b}, x] && InverseFunctionQ[v]

Rule 2686

$\text{Int}[(a_)*\sec[(e_.) + (f_.)*(x_)]^{(m_.)}*((b_)*\tan[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[a/f, \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \text{Sec}[e + f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])

Rubi steps

$$\begin{aligned} \int \sec^4(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}(\int x^3(-1 + x^2) dx, x, \sec(a + bx))}{b} \\ &= \frac{\text{Subst}(\int (-x^3 + x^5) dx, x, \sec(a + bx))}{b} \\ &= -\frac{\sec^4(a + bx)}{4b} + \frac{\sec^6(a + bx)}{6b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 28, normalized size = 0.90

$$\frac{3 \sec^4(a + bx) - 2 \sec^6(a + bx)}{12b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^4*Tan[a + b*x]^3,x]``[Out] -1/12*(3*Sec[a + b*x]^4 - 2*Sec[a + b*x]^6)/b`**Maple [A]**

time = 0.07, size = 42, normalized size = 1.35

method	result	size
derivativedivides	$\frac{\frac{\sin^4(bx+a)}{6 \cos(bx+a)^6} + \frac{\sin^4(bx+a)}{12 \cos(bx+a)^4}}{b}$	42
default	$\frac{\frac{\sin^4(bx+a)}{6 \cos(bx+a)^6} + \frac{\sin^4(bx+a)}{12 \cos(bx+a)^4}}{b}$	42
risch	$-\frac{4(3e^{8i(bx+a)} - 2e^{6i(bx+a)} + 3e^{4i(bx+a)})}{3b(e^{2i(bx+a)} + 1)^6}$	53
norman	$\frac{\frac{4(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{4(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{8(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^6}$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^7*sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/6*sin(b*x+a)^4/cos(b*x+a)^6+1/12*sin(b*x+a)^4/cos(b*x+a)^4)`**Maxima [A]**

time = 0.30, size = 49, normalized size = 1.58

$$-\frac{3 \sin(bx + a)^2 - 1}{12 (\sin(bx + a)^6 - 3 \sin(bx + a)^4 + 3 \sin(bx + a)^2 - 1)b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^3,x, algorithm="maxima")``[Out] -1/12*(3*sin(b*x + a)^2 - 1)/((sin(b*x + a)^6 - 3*sin(b*x + a)^4 + 3*sin(b*x + a)^2 - 1)*b)`**Fricas [A]**

time = 0.36, size = 25, normalized size = 0.81

$$-\frac{3 \cos(bx + a)^2 - 2}{12b \cos(bx + a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^7*sin(b*x+a)^3,x, algorithm="fricas")`

[Out] `-1/12*(3*cos(b*x + a)^2 - 2)/(b*cos(b*x + a)^6)`

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**7*sin(b*x+a)**3,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [A]

time = 4.81, size = 25, normalized size = 0.81

$$\frac{3 \cos (bx + a)^2 - 2}{12 b \cos (bx + a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^7*sin(b*x+a)^3,x, algorithm="giac")`

[Out] `-1/12*(3*cos(b*x + a)^2 - 2)/(b*cos(b*x + a)^6)`

Mupad [B]

time = 0.40, size = 25, normalized size = 0.81

$$\frac{\tan(a + bx)^4 (2 \tan(a + bx)^2 + 3)}{12 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^3/cos(a + b*x)^7,x)`

[Out] `(tan(a + b*x)^4*(2*tan(a + b*x)^2 + 3))/(12*b)`

3.78 $\int \sec^5(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\sec^5(a + bx)}{5b} + \frac{\sec^7(a + bx)}{7b}$$

[Out] $-1/5*\sec(b*x+a)^5/b+1/7*\sec(b*x+a)^7/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 14}

$$\frac{\sec^7(a + bx)}{7b} - \frac{\sec^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^5*\text{Tan}[a + b*x]^3, x]$

[Out] $-1/5*\text{Sec}[a + b*x]^5/b + \text{Sec}[a + b*x]^7/(7*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m, x\} \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{InverseFunctionQ}[v]]$

Rule 2686

$\text{Int}[(a_)*\sec[(e_)+(f_)*(x_)]^{(m_)}*((b_)*\tan[(e_)+(f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[a/f, \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{(n-1)/2}], x], x, \text{Sec}[e+f*x]], x] /; \text{FreeQ}\{a, e, f, m, x\} \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n+1])$

Rubi steps

$$\begin{aligned} \int \sec^5(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}(\int x^4(-1 + x^2) dx, x, \sec(a + bx))}{b} \\ &= \frac{\text{Subst}(\int (-x^4 + x^6) dx, x, \sec(a + bx))}{b} \\ &= -\frac{\sec^5(a + bx)}{5b} + \frac{\sec^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 31, normalized size = 1.00

$$-\frac{\sec^5(a+bx)}{5b} + \frac{\sec^7(a+bx)}{7b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^5*Tan[a + b*x]^3,x]``[Out] -1/5*Sec[a + b*x]^5/b + Sec[a + b*x]^7/(7*b)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(27) = 54$.

time = 0.09, size = 96, normalized size = 3.10

method	result	size
risch	$-\frac{32(7e^{9i(bx+a)} - 6e^{7i(bx+a)} + 7e^{5i(bx+a)})}{35b(e^{2i(bx+a)} + 1)^7}$	53
derivativedivides	$\frac{\frac{\sin^4(bx+a)}{7 \cos(bx+a)^7} + \frac{3(\sin^4(bx+a))}{35 \cos(bx+a)^5} + \frac{\sin^4(bx+a)}{35 \cos(bx+a)^3} - \frac{\sin^4(bx+a)}{35 \cos(bx+a)} - \frac{(2+\sin^2(bx+a)) \cos(bx+a)}{35}}{b}$	96
default	$\frac{\frac{\sin^4(bx+a)}{7 \cos(bx+a)^7} + \frac{3(\sin^4(bx+a))}{35 \cos(bx+a)^5} + \frac{\sin^4(bx+a)}{35 \cos(bx+a)^3} - \frac{\sin^4(bx+a)}{35 \cos(bx+a)} - \frac{(2+\sin^2(bx+a)) \cos(bx+a)}{35}}{b}$	96
norman	$\frac{-\frac{4(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{8(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{4}{35b} - \frac{4(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{4(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{8(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{5b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^7}$	103

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^8*sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/7*sin(b*x+a)^4/cos(b*x+a)^7+3/35*sin(b*x+a)^4/cos(b*x+a)^5+1/35*sin(b*x+a)^4/cos(b*x+a)^3-1/35*sin(b*x+a)^4/cos(b*x+a)-1/35*(2+sin(b*x+a)^2)*cos(b*x+a))`**Maxima [A]**

time = 0.30, size = 25, normalized size = 0.81

$$-\frac{7 \cos(bx+a)^2 - 5}{35 b \cos(bx+a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^8*sin(b*x+a)^3,x, algorithm="maxima")``[Out] -1/35*(7*cos(b*x + a)^2 - 5)/(b*cos(b*x + a)^7)`

Fricas [A]

time = 0.35, size = 25, normalized size = 0.81

$$-\frac{7 \cos (bx+a)^2-5}{35 b \cos (bx+a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^8*sin(b*x+a)^3,x, algorithm="fricas")``[Out] -1/35*(7*cos(b*x + a)^2 - 5)/(b*cos(b*x + a)^7)`**Sympy [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**8*sin(b*x+a)**3,x)``[Out] Exception raised: SystemError >> excessive stack use: stack is 6189 deep`**Giac [A]**

time = 4.82, size = 25, normalized size = 0.81

$$-\frac{7 \cos (bx+a)^2-5}{35 b \cos (bx+a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^8*sin(b*x+a)^3,x, algorithm="giac")``[Out] -1/35*(7*cos(b*x + a)^2 - 5)/(b*cos(b*x + a)^7)`**Mupad [B]**

time = 0.63, size = 25, normalized size = 0.81

$$-\frac{7 \cos (a+bx)^2-5}{35 b \cos (a+bx)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^3/cos(a + b*x)^8,x)``[Out] -(7*cos(a + b*x)^2 - 5)/(35*b*cos(a + b*x)^7)`

3.79 $\int \sec^6(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=31

$$-\frac{\sec^6(a + bx)}{6b} + \frac{\sec^8(a + bx)}{8b}$$

[Out] $-1/6*\sec(b*x+a)^6/b+1/8*\sec(b*x+a)^8/b$

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 14}

$$\frac{\sec^8(a + bx)}{8b} - \frac{\sec^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]^6*\text{Tan}[a + b*x]^3,x]$

[Out] $-1/6*\text{Sec}[a + b*x]^6/b + \text{Sec}[a + b*x]^8/(8*b)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2686

$\text{Int}[(a_)*\sec[(e_)+(f_)*(x_)]^{(m_)}*((b_)*\tan[(e_)+(f_)*(x_)]^{(n_)}), x_Symbol] \rightarrow \text{Dist}[a/f, \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \text{Sec}[e+f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])

Rubi steps

$$\begin{aligned} \int \sec^6(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}(\int x^5(-1 + x^2) dx, x, \sec(a + bx))}{b} \\ &= \frac{\text{Subst}(\int (-x^5 + x^7) dx, x, \sec(a + bx))}{b} \\ &= -\frac{\sec^6(a + bx)}{6b} + \frac{\sec^8(a + bx)}{8b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 28, normalized size = 0.90

$$\frac{4 \sec^6(a + bx) - 3 \sec^8(a + bx)}{24b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^6*Tan[a + b*x]^3,x]**[Out]** -1/24*(4*Sec[a + b*x]^6 - 3*Sec[a + b*x]^8)/b**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(27) = 54.

time = 0.09, size = 60, normalized size = 1.94

method	result	size
risch	$-\frac{32(e^{10i(bx+a)} - e^{8i(bx+a)} + e^{6i(bx+a)})}{3b(e^{2i(bx+a)} + 1)^8}$	49
derivativedivides	$\frac{\frac{\sin^4(bx+a)}{8 \cos(bx+a)^8} + \frac{\sin^4(bx+a)}{12 \cos(bx+a)^6} + \frac{\sin^4(bx+a)}{24 \cos(bx+a)^4}}{b}$	60
default	$\frac{\frac{\sin^4(bx+a)}{8 \cos(bx+a)^8} + \frac{\sin^4(bx+a)}{12 \cos(bx+a)^6} + \frac{\sin^4(bx+a)}{24 \cos(bx+a)^4}}{b}$	60
norman	$\frac{\frac{4(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{4(\tan^{12}(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{16(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{16(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{40(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^8}$	98

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^9*sin(b*x+a)^3,x,method=_RETURNVERBOSE)**[Out]** 1/b*(1/8*sin(b*x+a)^4/cos(b*x+a)^8+1/12*sin(b*x+a)^4/cos(b*x+a)^6+1/24*sin(b*x+a)^4/cos(b*x+a)^4)**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(27) = 54.

time = 0.29, size = 59, normalized size = 1.90

$$\frac{4 \sin^2(bx + a) - 1}{24 (\sin^8(bx + a) - 4 \sin^6(bx + a) + 6 \sin^4(bx + a) - 4 \sin^2(bx + a) + 1)b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^9*sin(b*x+a)^3,x, algorithm="maxima")**[Out]** 1/24*(4*sin(b*x + a)^2 - 1)/((sin(b*x + a)^8 - 4*sin(b*x + a)^6 + 6*sin(b*x + a)^4 - 4*sin(b*x + a)^2 + 1)*b)

Fricas [A]

time = 0.35, size = 25, normalized size = 0.81

$$\frac{4 \cos (bx + a)^2 - 3}{24 b \cos (bx + a)^8}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^9*sin(b*x+a)^3,x, algorithm="fricas")``[Out] -1/24*(4*cos(b*x + a)^2 - 3)/(b*cos(b*x + a)^8)`**Sympy [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**9*sin(b*x+a)**3,x)``[Out] Exception raised: SystemError >> excessive stack use: stack is 8569 deep`**Giac [A]**

time = 3.92, size = 25, normalized size = 0.81

$$\frac{4 \cos (bx + a)^2 - 3}{24 b \cos (bx + a)^8}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^9*sin(b*x+a)^3,x, algorithm="giac")``[Out] -1/24*(4*cos(b*x + a)^2 - 3)/(b*cos(b*x + a)^8)`**Mupad [B]**

time = 0.42, size = 35, normalized size = 1.13

$$\frac{\tan(a + bx)^4 (3 \tan(a + bx)^4 + 8 \tan(a + bx)^2 + 6)}{24 b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^3/cos(a + b*x)^9,x)``[Out] (tan(a + b*x)^4*(8*tan(a + b*x)^2 + 3*tan(a + b*x)^4 + 6))/(24*b)`

3.80 $\int \cos^7(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=61

$$\frac{\sin^5(a + bx)}{5b} - \frac{3 \sin^7(a + bx)}{7b} + \frac{\sin^9(a + bx)}{3b} - \frac{\sin^{11}(a + bx)}{11b}$$

[Out] $1/5*\sin(b*x+a)^5/b-3/7*\sin(b*x+a)^7/b+1/3*\sin(b*x+a)^9/b-1/11*\sin(b*x+a)^{11}/b$

Rubi [A]

time = 0.03, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 276}

$$-\frac{\sin^{11}(a + bx)}{11b} + \frac{\sin^9(a + bx)}{3b} - \frac{3 \sin^7(a + bx)}{7b} + \frac{\sin^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^7*Sin[a + b*x]^4,x]

[Out] Sin[a + b*x]^5/(5*b) - (3*Sin[a + b*x]^7)/(7*b) + Sin[a + b*x]^9/(3*b) - Sin[a + b*x]^11/(11*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^7(a + bx) \sin^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4(1 - x^2)^3 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^4 - 3x^6 + 3x^8 - x^{10}) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^5(a + bx)}{5b} - \frac{3 \sin^7(a + bx)}{7b} + \frac{\sin^9(a + bx)}{3b} - \frac{\sin^{11}(a + bx)}{11b} \end{aligned}$$

Mathematica [A]

time = 0.15, size = 47, normalized size = 0.77

$$\frac{(3042 + 3335 \cos(2(a + bx)) + 910 \cos(4(a + bx)) + 105 \cos(6(a + bx))) \sin^5(a + bx)}{36960b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^7*Sin[a + b*x]^4,x]`

```
[Out] ((3042 + 3335*Cos[2*(a + b*x)] + 910*Cos[4*(a + b*x)] + 105*Cos[6*(a + b*x)]
)*Sin[a + b*x]^5)/(36960*b)
```

Maple [A]

time = 0.13, size = 78, normalized size = 1.28

method	result	size
derivativedivides	$\frac{-\frac{(\sin^3(bx+a))(\cos^8(bx+a))}{11} - \frac{\sin(bx+a)(\cos^8(bx+a))}{33} + \frac{\left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{231}}{b}$	78
default	$\frac{-\frac{(\sin^3(bx+a))(\cos^8(bx+a))}{11} - \frac{\sin(bx+a)(\cos^8(bx+a))}{33} + \frac{\left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{231}}{b}$	78
risch	$\frac{7 \sin(bx+a)}{512b} + \frac{\sin(11bx+11a)}{11264b} + \frac{\sin(9bx+9a)}{3072b} - \frac{\sin(7bx+7a)}{7168b} - \frac{11 \sin(5bx+5a)}{5120b} - \frac{\sin(3bx+3a)}{512b}$	83

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^7*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/11*sin(b*x+a)^3*cos(b*x+a)^8-1/33*sin(b*x+a)*cos(b*x+a)^8+1/231*(16
/5+cos(b*x+a)^6+6/5*cos(b*x+a)^4+8/5*cos(b*x+a)^2)*sin(b*x+a)
```

Maxima [A]

time = 0.28, size = 46, normalized size = 0.75

$$\frac{105 \sin(bx + a)^{11} - 385 \sin(bx + a)^9 + 495 \sin(bx + a)^7 - 231 \sin(bx + a)^5}{1155b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7*sin(b*x+a)^4,x, algorithm="maxima")`

```
[Out] -1/1155*(105*sin(b*x + a)^11 - 385*sin(b*x + a)^9 + 495*sin(b*x + a)^7 - 23
1*sin(b*x + a)^5)/b
```

Fricas [A]

time = 0.39, size = 63, normalized size = 1.03

$$\frac{(105 \cos(bx + a)^{10} - 140 \cos(bx + a)^8 + 5 \cos(bx + a)^6 + 6 \cos(bx + a)^4 + 8 \cos(bx + a)^2 + 16) \sin(bx + a)}{1155b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7*sin(b*x+a)^4,x, algorithm="fricas")

[Out] $1/1155*(105*\cos(b*x + a)^{10} - 140*\cos(b*x + a)^8 + 5*\cos(b*x + a)^6 + 6*\cos(b*x + a)^4 + 8*\cos(b*x + a)^2 + 16)*\sin(b*x + a)/b$

Sympy [A]

time = 3.63, size = 88, normalized size = 1.44

$$\begin{cases} \frac{16 \sin^{11}(a+bx)}{1155b} + \frac{8 \sin^9(a+bx) \cos^2(a+bx)}{105b} + \frac{6 \sin^7(a+bx) \cos^4(a+bx)}{35b} + \frac{\sin^5(a+bx) \cos^6(a+bx)}{5b} & \text{for } b \neq 0 \\ x \sin^4(a) \cos^7(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**7*sin(b*x+a)**4,x)

[Out] Piecewise((16*sin(a + b*x)**11/(1155*b) + 8*sin(a + b*x)**9*cos(a + b*x)**2/(105*b) + 6*sin(a + b*x)**7*cos(a + b*x)**4/(35*b) + sin(a + b*x)**5*cos(a + b*x)**6/(5*b), Ne(b, 0)), (x*sin(a)**4*cos(a)**7, True))

Giac [A]

time = 3.38, size = 82, normalized size = 1.34

$$\frac{\sin(11bx + 11a)}{11264b} + \frac{\sin(9bx + 9a)}{3072b} - \frac{\sin(7bx + 7a)}{7168b} - \frac{11 \sin(5bx + 5a)}{5120b} - \frac{\sin(3bx + 3a)}{512b} + \frac{7 \sin(bx + a)}{512b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7*sin(b*x+a)^4,x, algorithm="giac")

[Out] $1/11264*\sin(11*b*x + 11*a)/b + 1/3072*\sin(9*b*x + 9*a)/b - 1/7168*\sin(7*b*x + 7*a)/b - 11/5120*\sin(5*b*x + 5*a)/b - 1/512*\sin(3*b*x + 3*a)/b + 7/512*\sin(b*x + a)/b$

Mupad [B]

time = 0.38, size = 45, normalized size = 0.74

$$\frac{-\frac{\sin(a+bx)^{11}}{11} + \frac{\sin(a+bx)^9}{3} - \frac{3 \sin(a+bx)^7}{7} + \frac{\sin(a+bx)^5}{5}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^7*sin(a + b*x)^4,x)

[Out] $(\sin(a + b*x)^5/5 - (3*\sin(a + b*x)^7)/7 + \sin(a + b*x)^9/3 - \sin(a + b*x)^{11}/11)/b$

3.81 $\int \cos^5(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sin^5(a + bx)}{5b} - \frac{2 \sin^7(a + bx)}{7b} + \frac{\sin^9(a + bx)}{9b}$$

[Out] 1/5*sin(b*x+a)^5/b-2/7*sin(b*x+a)^7/b+1/9*sin(b*x+a)^9/b

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 276}

$$\frac{\sin^9(a + bx)}{9b} - \frac{2 \sin^7(a + bx)}{7b} + \frac{\sin^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^5*Sin[a + b*x]^4,x]

[Out] Sin[a + b*x]^5/(5*b) - (2*Sin[a + b*x]^7)/(7*b) + Sin[a + b*x]^9/(9*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx) \sin^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4(1 - x^2)^2 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^4 - 2x^6 + x^8) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^5(a + bx)}{5b} - \frac{2 \sin^7(a + bx)}{7b} + \frac{\sin^9(a + bx)}{9b} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 37, normalized size = 0.80

$$\frac{(249 + 220 \cos(2(a + bx))) + 35 \cos(4(a + bx))) \sin^5(a + bx)}{2520b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^5*Sin[a + b*x]^4,x]**[Out]** ((249 + 220*Cos[2*(a + b*x)] + 35*Cos[4*(a + b*x)])*Sin[a + b*x]^5)/(2520*b)**Maple [A]**

time = 0.13, size = 68, normalized size = 1.48

method	result	size
derivativedivides	$\frac{-\frac{(\sin^3(bx+a))(\cos^6(bx+a))}{9} - \frac{\sin(bx+a)(\cos^6(bx+a))}{21} + \frac{\left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{105}}{b}$	68
default	$\frac{-\frac{(\sin^3(bx+a))(\cos^6(bx+a))}{9} - \frac{\sin(bx+a)(\cos^6(bx+a))}{21} + \frac{\left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{105}}{b}$	68
risch	$\frac{3 \sin(bx+a)}{128b} + \frac{\sin(9bx+9a)}{2304b} + \frac{\sin(7bx+7a)}{1792b} - \frac{\sin(5bx+5a)}{320b} - \frac{\sin(3bx+3a)}{192b}$	69
norman	$\frac{32 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5b} - \frac{384 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{35b} + \frac{6976 \left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{315b} - \frac{384 \left(\tan^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{35b} + \frac{32 \left(\tan^{13}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^9}$	98

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^5*sin(b*x+a)^4,x,method=_RETURNVERBOSE)**[Out]** 1/b*(-1/9*sin(b*x+a)^3*cos(b*x+a)^6-1/21*sin(b*x+a)*cos(b*x+a)^6+1/105*(8/3+cos(b*x+a)^4+4/3*cos(b*x+a)^2)*sin(b*x+a))**Maxima [A]**

time = 0.28, size = 36, normalized size = 0.78

$$\frac{35 \sin(bx + a)^9 - 90 \sin(bx + a)^7 + 63 \sin(bx + a)^5}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^4,x, algorithm="maxima")**[Out]** 1/315*(35*sin(b*x + a)^9 - 90*sin(b*x + a)^7 + 63*sin(b*x + a)^5)/b**Fricas [A]**

time = 0.38, size = 53, normalized size = 1.15

$$\frac{(35 \cos(bx + a)^8 - 50 \cos(bx + a)^6 + 3 \cos(bx + a)^4 + 4 \cos(bx + a)^2 + 8) \sin(bx + a)}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/315*(35*cos(b*x + a)^8 - 50*cos(b*x + a)^6 + 3*cos(b*x + a)^4 + 4*cos(b*x + a)^2 + 8)*sin(b*x + a)/b

Sympy [A]

time = 1.25, size = 66, normalized size = 1.43

$$\begin{cases} \frac{8 \sin^9(a+bx)}{315b} + \frac{4 \sin^7(a+bx) \cos^2(a+bx)}{35b} + \frac{\sin^5(a+bx) \cos^4(a+bx)}{5b} & \text{for } b \neq 0 \\ x \sin^4(a) \cos^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**5*sin(b*x+a)**4,x)

[Out] Piecewise((8*sin(a + b*x)**9/(315*b) + 4*sin(a + b*x)**7*cos(a + b*x)**2/(35*b) + sin(a + b*x)**5*cos(a + b*x)**4/(5*b), Ne(b, 0)), (x*sin(a)**4*cos(a)**5, True))

Giac [A]

time = 3.97, size = 68, normalized size = 1.48

$$\frac{\sin(9bx + 9a)}{2304b} + \frac{\sin(7bx + 7a)}{1792b} - \frac{\sin(5bx + 5a)}{320b} - \frac{\sin(3bx + 3a)}{192b} + \frac{3 \sin(bx + a)}{128b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/2304*sin(9*b*x + 9*a)/b + 1/1792*sin(7*b*x + 7*a)/b - 1/320*sin(5*b*x + 5*a)/b - 1/192*sin(3*b*x + 3*a)/b + 3/128*sin(b*x + a)/b

Mupad [B]

time = 0.38, size = 36, normalized size = 0.78

$$\frac{35 \sin(a + bx)^9 - 90 \sin(a + bx)^7 + 63 \sin(a + bx)^5}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5*sin(a + b*x)^4,x)

[Out] (63*sin(a + b*x)^5 - 90*sin(a + b*x)^7 + 35*sin(a + b*x)^9)/(315*b)

3.82 $\int \cos^3(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b}$$

[Out] 1/5*sin(b*x+a)^5/b-1/7*sin(b*x+a)^7/b

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 14}

$$\frac{\sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*Sin[a + b*x]^4,x]

[Out] Sin[a + b*x]^5/(5*b) - Sin[a + b*x]^7/(7*b)

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \sin^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4(1 - x^2) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^4 - x^6) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^5(a + bx)}{5b} - \frac{\sin^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 27, normalized size = 0.87

$$\frac{(9 + 5 \cos(2(a + bx))) \sin^5(a + bx)}{70b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*Sin[a + b*x]^4,x]``[Out] ((9 + 5*Cos[2*(a + b*x)])*Sin[a + b*x]^5)/(70*b)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 57 vs. 2(27) = 54.

time = 0.09, size = 58, normalized size = 1.87

method	result	size
risch	$\frac{3 \sin(bx+a)}{64b} + \frac{\sin(7bx+7a)}{448b} - \frac{\sin(5bx+5a)}{320b} - \frac{\sin(3bx+3a)}{64b}$	55
derivativedivides	$-\frac{(\cos^4(bx+a))(\sin^3(bx+a))}{7} - \frac{3 \sin(bx+a)(\cos^4(bx+a))}{35b} + \frac{(2+\cos^2(bx+a)) \sin(bx+a)}{35}$	58
default	$-\frac{(\cos^4(bx+a))(\sin^3(bx+a))}{7} - \frac{3 \sin(bx+a)(\cos^4(bx+a))}{35b} + \frac{(2+\cos^2(bx+a)) \sin(bx+a)}{35}$	58
norman	$\frac{32(\tan^5(\frac{bx+a}{2}))}{5b} - \frac{192(\tan^7(\frac{bx+a}{2}))}{35b} + \frac{32(\tan^9(\frac{bx+a}{2}))}{5b}$ $\frac{1}{(1+\tan^2(\frac{bx+a}{2}))^7}$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^3*sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/7*cos(b*x+a)^4*sin(b*x+a)^3-3/35*sin(b*x+a)*cos(b*x+a)^4+1/35*(2*cos(b*x+a)^2)*sin(b*x+a))`**Maxima [A]**

time = 0.33, size = 26, normalized size = 0.84

$$\frac{5 \sin(bx + a)^7 - 7 \sin(bx + a)^5}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^3*sin(b*x+a)^4,x, algorithm="maxima")``[Out] -1/35*(5*sin(b*x + a)^7 - 7*sin(b*x + a)^5)/b`**Fricas [A]**

time = 0.37, size = 41, normalized size = 1.32

$$\frac{(5 \cos(bx + a)^6 - 8 \cos(bx + a)^4 + \cos(bx + a)^2 + 2) \sin(bx + a)}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3*sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $1/35*(5*\cos(b*x + a)^6 - 8*\cos(b*x + a)^4 + \cos(b*x + a)^2 + 2)*\sin(b*x + a)/b$

Sympy [A]

time = 0.58, size = 44, normalized size = 1.42

$$\begin{cases} \frac{2 \sin^7(a+bx)}{35b} + \frac{\sin^5(a+bx) \cos^2(a+bx)}{5b} & \text{for } b \neq 0 \\ x \sin^4(a) \cos^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**3*sin(b*x+a)**4,x)`

[Out] `Piecewise((2*sin(a + b*x)**7/(35*b) + sin(a + b*x)**5*cos(a + b*x)**2/(5*b), Ne(b, 0)), (x*sin(a)**4*cos(a)**3, True))`

Giac [A]

time = 3.96, size = 26, normalized size = 0.84

$$\frac{5 \sin(bx + a)^7 - 7 \sin(bx + a)^5}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3*sin(b*x+a)^4,x, algorithm="giac")`

[Out] $-1/35*(5*\sin(b*x + a)^7 - 7*\sin(b*x + a)^5)/b$

Mupad [B]

time = 0.03, size = 26, normalized size = 0.84

$$\frac{7 \sin(a + bx)^5 - 5 \sin(a + bx)^7}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^3*sin(a + b*x)^4,x)`

[Out] $(7*\sin(a + b*x)^5 - 5*\sin(a + b*x)^7)/(35*b)$

3.83 $\int \cos(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sin^5(a + bx)}{5b}$$

[Out] 1/5*sin(b*x+a)^5/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2644, 30}

$$\frac{\sin^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Sin[a + b*x]^4,x]

[Out] Sin[a + b*x]^5/(5*b)

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \sin^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$\frac{\sin^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Sin[a + b*x]^4,x]

[Out] Sin[a + b*x]^5/(5*b)

Maple [A]

time = 0.04, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$\frac{\sin^5(bx+a)}{5b}$	14
default	$\frac{\sin^5(bx+a)}{5b}$	14
norman	$\frac{32\left(\tan^5\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{5b\left(1+\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)^5}$	32
risch	$\frac{\sin(bx+a)}{8b} + \frac{\sin(5bx+5a)}{80b} - \frac{\sin(3bx+3a)}{16b}$	41

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] 1/5*sin(b*x+a)^5/b

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$\frac{\sin^5(bx+a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a)^4,x, algorithm="maxima")

[Out] 1/5*sin(b*x + a)^5/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 31 vs. 2(13) = 26.

time = 0.34, size = 31, normalized size = 2.07

$$\frac{(\cos(bx+a)^4 - 2\cos(bx+a)^2 + 1)\sin(bx+a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/5*(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*sin(b*x + a)/b

Sympy [A]

time = 0.23, size = 20, normalized size = 1.33

$$\begin{cases} \frac{\sin^5(a+bx)}{5b} & \text{for } b \neq 0 \\ x \sin^4(a) \cos(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)**4,x)``[Out] Piecewise((sin(a + b*x)**5/(5*b), Ne(b, 0)), (x*sin(a)**4*cos(a), True))`**Giac [A]**

time = 3.73, size = 13, normalized size = 0.87

$$\frac{\sin(bx + a)^5}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)^4,x, algorithm="giac")``[Out] 1/5*sin(b*x + a)^5/b`**Mupad [B]**

time = 0.36, size = 13, normalized size = 0.87

$$\frac{\sin(a + bx)^5}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)*sin(a + b*x)^4,x)``[Out] sin(a + b*x)^5/(5*b)`

3.84 $\int \sin^2(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=40

$$-\frac{3x}{2} + \frac{3 \tan(a + bx)}{2b} - \frac{\sin^2(a + bx) \tan(a + bx)}{2b}$$

[Out] $-3/2*x+3/2*\tan(b*x+a)/b-1/2*\sin(b*x+a)^2*\tan(b*x+a)/b$

Rubi [A]

time = 0.03, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2671, 294, 327, 209}

$$\frac{3 \tan(a + bx)}{2b} - \frac{\sin^2(a + bx) \tan(a + bx)}{2b} - \frac{3x}{2}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2*Tan[a + b*x]^2,x]

[Out] $(-3*x)/2 + (3*\tan[a + b*x])/(2*b) - (\sin[a + b*x]^2*\tan[a + b*x])/(2*b)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2671

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[In

```
t[(ff*x)^(m + n)/(b^2 + ff^2*x^2)^(m/2 + 1), x], x, b*(Tan[e + f*x]/ff)], x
]] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2]
```

Rubi steps

$$\begin{aligned} \int \sin^2(a + bx) \tan^2(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^4}{(1+x^2)^2} dx, x, \tan(a + bx)\right)}{b} \\ &= -\frac{\sin^2(a + bx) \tan(a + bx)}{2b} + \frac{3\text{Subst}\left(\int \frac{x^2}{1+x^2} dx, x, \tan(a + bx)\right)}{2b} \\ &= \frac{3 \tan(a + bx)}{2b} - \frac{\sin^2(a + bx) \tan(a + bx)}{2b} - \frac{3\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \tan(a + bx)\right)}{2b} \\ &= -\frac{3x}{2} + \frac{3 \tan(a + bx)}{2b} - \frac{\sin^2(a + bx) \tan(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 31, normalized size = 0.78

$$\frac{-6(a + bx) + \sin(2(a + bx)) + 4 \tan(a + bx)}{4b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^2*Tan[a + b*x]^2,x]
```

```
[Out] (-6*(a + b*x) + Sin[2*(a + b*x)] + 4*Tan[a + b*x])/(4*b)
```

Maple [A]

time = 0.05, size = 54, normalized size = 1.35

method	result	size
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{\cos(bx+a)} + \left(\sin^3(bx+a) + \frac{3 \sin(bx+a)}{2}\right) \cos(bx+a) - \frac{3bx}{2} - \frac{3a}{2}}{b}$	54
default	$\frac{\frac{\sin^5(bx+a)}{\cos(bx+a)} + \left(\sin^3(bx+a) + \frac{3 \sin(bx+a)}{2}\right) \cos(bx+a) - \frac{3bx}{2} - \frac{3a}{2}}{b}$	54
risch	$-\frac{3x}{2} - \frac{ie^{2i(bx+a)}}{8b} + \frac{ie^{-2i(bx+a)}}{8b} + \frac{2i}{b(e^{2i(bx+a)}+1)}$	54
norman	$\frac{\frac{3x}{2} - \frac{3 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} - \frac{2(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right))}{b} - \frac{3(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right))}{b} + \frac{3x(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right))}{2} - \frac{3x(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right))}{2} - \frac{3x(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right))}{2}}{(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right))^2 (\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1)}$	124

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(b*x+a)^2*sin(b*x+a)^4,x,method=_RETURNVERBOSE)
```

[Out] $1/b*(\sin(b*x+a)^5/\cos(b*x+a)+(\sin(b*x+a)^3+3/2*\sin(b*x+a))*\cos(b*x+a)-3/2*b*x-3/2*a)$

Maxima [A]

time = 0.52, size = 41, normalized size = 1.02

$$-\frac{3bx + 3a - \frac{\tan(bx+a)}{\tan(bx+a)^2+1} - 2 \tan(bx+a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2*sin(b*x+a)^4,x, algorithm="maxima")`

[Out] $-1/2*(3*b*x + 3*a - \tan(b*x + a)/(\tan(b*x + a)^2 + 1) - 2*\tan(b*x + a))/b$

Fricas [A]

time = 0.35, size = 42, normalized size = 1.05

$$-\frac{3bx \cos(bx+a) - (\cos(bx+a)^2 + 2) \sin(bx+a)}{2b \cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2*sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $-1/2*(3*b*x*\cos(b*x + a) - (\cos(b*x + a)^2 + 2)*\sin(b*x + a))/(b*\cos(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^4(a + bx) \sec^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**2*sin(b*x+a)**4,x)`

[Out] `Integral(sin(a + b*x)**4*sec(a + b*x)**2, x)`

Giac [A]

time = 3.57, size = 41, normalized size = 1.02

$$-\frac{3bx + 3a - \frac{\tan(bx+a)}{\tan(bx+a)^2+1} - 2 \tan(bx+a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2*sin(b*x+a)^4,x, algorithm="giac")`

[Out] $-1/2*(3*b*x + 3*a - \tan(b*x + a)/(\tan(b*x + a)^2 + 1) - 2*\tan(b*x + a))/b$

Mupad [B]

time = 0.48, size = 38, normalized size = 0.95

$$\frac{\frac{\cos(a+bx) \sin(a+bx)}{2} + \frac{\sin(a+bx)}{\cos(a+bx)}}{b} - \frac{3x}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^4/cos(a + b*x)^2,x)`

[Out] `((cos(a + b*x)*sin(a + b*x))/2 + sin(a + b*x)/cos(a + b*x))/b - (3*x)/2`

3.85 $\int \tan^4(a + bx) dx$

Optimal. Leaf size=28

$$x - \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b}$$

[Out] $x - \tan(b*x+a)/b + 1/3*\tan(b*x+a)^3/b$

Rubi [A]

time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$\frac{\tan^3(a + bx)}{3b} - \frac{\tan(a + bx)}{b} + x$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Tan}[a + b*x]^4, x]$

[Out] $x - \text{Tan}[a + b*x]/b + \text{Tan}[a + b*x]^3/(3*b)$

Rule 8

$\text{Int}[a_, x_Symbol] \text{ :> Simp}[a*x, x] \text{ /; FreeQ}[a, x]$

Rule 3554

$\text{Int}[(b_.*\text{tan}[(c_.) + (d_.)*(x_)])^{(n_)}, x_Symbol] \text{ :> Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)/(d*(n - 1))}), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] \text{ /; FreeQ}\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1]$

Rubi steps

$$\begin{aligned} \int \tan^4(a + bx) dx &= \frac{\tan^3(a + bx)}{3b} - \int \tan^2(a + bx) dx \\ &= -\frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b} + \int 1 dx \\ &= x - \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 38, normalized size = 1.36

$$\frac{\tan^{-1}(\tan(a + bx))}{b} - \frac{\tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Tan[a + b*x]^4, x]

[Out] ArcTan[Tan[a + b*x]]/b - Tan[a + b*x]/b + Tan[a + b*x]^3/(3*b)

Maple [A]

time = 0.05, size = 28, normalized size = 1.00

method	result	size
derivativedivides	$\frac{\frac{(\tan^3(bx+a))}{3} - \tan(bx+a) + bx+a}{b}$	2
default	$\frac{\frac{(\tan^3(bx+a))}{3} - \tan(bx+a) + bx+a}{b}$	2
risch	$x - \frac{4i(3e^{4i(bx+a)} + 3e^{2i(bx+a)} + 2)}{3b(e^{2i(bx+a)} + 1)^3}$	4
norman	$\frac{x\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - x + \frac{2\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} - \frac{20\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} + \frac{2\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + 3x\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 3x\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^3}$	1

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^4*sin(b*x+a)^4, x, method=_RETURNVERBOSE)

[Out] 1/b*(1/3*tan(b*x+a)^3-tan(b*x+a)+b*x+a)

Maxima [A]

time = 0.53, size = 29, normalized size = 1.04

$$\frac{\tan(bx+a)^3 + 3bx + 3a - 3\tan(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^4, x, algorithm="maxima")

[Out] 1/3*(tan(b*x + a)^3 + 3*b*x + 3*a - 3*tan(b*x + a))/b

Fricas [A]

time = 0.38, size = 46, normalized size = 1.64

$$\frac{3bx \cos(bx+a)^3 - (4 \cos(bx+a)^2 - 1) \sin(bx+a)}{3b \cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^4, x, algorithm="fricas")

[Out] 1/3*(3*b*x*cos(b*x + a)^3 - (4*cos(b*x + a)^2 - 1)*sin(b*x + a))/(b*cos(b*x + a)^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^4(a + bx) \sec^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4*sin(b*x+a)**4,x)

[Out] Integral(sin(a + b*x)**4*sec(a + b*x)**4, x)

Giac [A]

time = 5.03, size = 29, normalized size = 1.04

$$\frac{\tan(bx + a)^3 + 3bx + 3a - 3 \tan(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/3*(tan(b*x + a)^3 + 3*b*x + 3*a - 3*tan(b*x + a))/b

Mupad [B]

time = 0.40, size = 24, normalized size = 0.86

$$x - \frac{\tan(a + bx) - \frac{\tan(a+bx)^3}{3}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/cos(a + b*x)^4,x)

[Out] x - (tan(a + b*x) - tan(a + b*x)^3/3)/b

3.86 $\int \sec^2(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\tan^5(a + bx)}{5b}$$

[Out] 1/5*tan(b*x+a)^5/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 30}

$$\frac{\tan^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2*Tan[a + b*x]^4,x]

[Out] Tan[a + b*x]^5/(5*b)

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2687

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^2(a + bx) \tan^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4 dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$\frac{\tan^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2*Tan[a + b*x]^4,x]

[Out] Tan[a + b*x]^5/(5*b)

Maple [A]

time = 0.06, size = 22, normalized size = 1.47

method	result	size
derivativedivides	$\frac{\sin^5(bx+a)}{5b \cos(bx+a)^5}$	22
default	$\frac{\sin^5(bx+a)}{5b \cos(bx+a)^5}$	22
norman	$-\frac{32 \left(\tan^5 \left(\frac{bx+a}{2} \right) \right)}{5b \left(\tan^2 \left(\frac{bx+a}{2} \right) - 1 \right)^5}$	32
risch	$\frac{2i(5e^{8i(bx+a)} + 10e^{4i(bx+a)} + 1)}{5b(e^{2i(bx+a)} + 1)^5}$	44

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^6*sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] 1/5/b*sin(b*x+a)^5/cos(b*x+a)^5

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$\frac{\tan(bx+a)^5}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^4,x, algorithm="maxima")

[Out] 1/5*tan(b*x + a)^5/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(13) = 26.

time = 0.35, size = 39, normalized size = 2.60

$$\frac{(\cos(bx+a)^4 - 2\cos(bx+a)^2 + 1)\sin(bx+a)}{5b\cos(bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/5*(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*sin(b*x + a)/(b*cos(b*x + a)^5)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**6*sin(b*x+a)**4,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [A]

time = 3.65, size = 13, normalized size = 0.87

$$\frac{\tan(bx + a)^5}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/5*tan(b*x + a)^5/b

Mupad [B]

time = 0.41, size = 13, normalized size = 0.87

$$\frac{\tan(a + bx)^5}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/cos(a + b*x)^6,x)

[Out] tan(a + b*x)^5/(5*b)

3.87 $\int \sec^4(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\tan^5(a + bx)}{5b} + \frac{\tan^7(a + bx)}{7b}$$

[Out] 1/5*tan(b*x+a)^5/b+1/7*tan(b*x+a)^7/b

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 14}

$$\frac{\tan^7(a + bx)}{7b} + \frac{\tan^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^4*Tan[a + b*x]^4,x]

[Out] Tan[a + b*x]^5/(5*b) + Tan[a + b*x]^7/(7*b)

Rule 14

```
Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2687

```
Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])
```

Rubi steps

$$\begin{aligned} \int \sec^4(a + bx) \tan^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4(1 + x^2) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^4 + x^6) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^5(a + bx)}{5b} + \frac{\tan^7(a + bx)}{7b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 77 vs. $2(31) = 62$.

time = 0.03, size = 77, normalized size = 2.48

$$\frac{2 \tan(a + bx)}{35b} + \frac{\sec^2(a + bx) \tan(a + bx)}{35b} - \frac{8 \sec^4(a + bx) \tan(a + bx)}{35b} + \frac{\sec^6(a + bx) \tan(a + bx)}{7b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^4*Tan[a + b*x]^4,x]

[Out] (2*Tan[a + b*x])/(35*b) + (Sec[a + b*x]^2*Tan[a + b*x])/(35*b) - (8*Sec[a + b*x]^4*Tan[a + b*x])/(35*b) + (Sec[a + b*x]^6*Tan[a + b*x])/(7*b)

Maple [A]

time = 0.08, size = 42, normalized size = 1.35

method	result	size
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{7 \cos(bx+a)^7} + \frac{2(\sin^5(bx+a))}{35 \cos(bx+a)^5}}{b}$	42
default	$\frac{\frac{\sin^5(bx+a)}{7 \cos(bx+a)^7} + \frac{2(\sin^5(bx+a))}{35 \cos(bx+a)^5}}{b}$	42
norman	$\frac{-\frac{32(\tan^5(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{192(\tan^7(\frac{bx}{2} + \frac{a}{2}))}{35b} - \frac{32(\tan^9(\frac{bx}{2} + \frac{a}{2}))}{5b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^7}$	66
risch	$\frac{4i(35e^{10i(bx+a)} - 35e^{8i(bx+a)} + 70e^{6i(bx+a)} - 14e^{4i(bx+a)} + 7e^{2i(bx+a)} + 1)}{35b(e^{2i(bx+a)} + 1)^7}$	77

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^8*sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] 1/b*(1/7*sin(b*x+a)^5/cos(b*x+a)^7+2/35*sin(b*x+a)^5/cos(b*x+a)^5)

Maxima [A]

time = 0.30, size = 26, normalized size = 0.84

$$\frac{5 \tan(bx + a)^7 + 7 \tan(bx + a)^5}{35b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^8*sin(b*x+a)^4,x, algorithm="maxima")

[Out] 1/35*(5*tan(b*x + a)^7 + 7*tan(b*x + a)^5)/b

Fricas [A]

time = 0.35, size = 49, normalized size = 1.58

$$\frac{(2 \cos(bx + a)^6 + \cos(bx + a)^4 - 8 \cos(bx + a)^2 + 5) \sin(bx + a)}{35b \cos(bx + a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^8*sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $1/35*(2*\cos(b*x + a)^6 + \cos(b*x + a)^4 - 8*\cos(b*x + a)^2 + 5)*\sin(b*x + a)/(b*\cos(b*x + a)^7)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**8*sin(b*x+a)**4,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 8569 deep

Giac [A]

time = 3.94, size = 26, normalized size = 0.84

$$\frac{5 \tan (bx + a)^7 + 7 \tan (bx + a)^5}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^8*sin(b*x+a)^4,x, algorithm="giac")`

[Out] $1/35*(5*\tan(b*x + a)^7 + 7*\tan(b*x + a)^5)/b$

Mupad [B]

time = 0.40, size = 25, normalized size = 0.81

$$\frac{\tan(a + bx)^5 (5 \tan(a + bx)^2 + 7)}{35 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^4/cos(a + b*x)^8,x)`

[Out] $(\tan(a + b*x)^5*(5*\tan(a + b*x)^2 + 7))/(35*b)$

3.88 $\int \sec^6(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\tan^5(a + bx)}{5b} + \frac{2 \tan^7(a + bx)}{7b} + \frac{\tan^9(a + bx)}{9b}$$

[Out] 1/5*tan(b*x+a)^5/b+2/7*tan(b*x+a)^7/b+1/9*tan(b*x+a)^9/b

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 276}

$$\frac{\tan^9(a + bx)}{9b} + \frac{2 \tan^7(a + bx)}{7b} + \frac{\tan^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^6*Tan[a + b*x]^4,x]

[Out] Tan[a + b*x]^5/(5*b) + (2*Tan[a + b*x]^7)/(7*b) + Tan[a + b*x]^9/(9*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2687

Int[sec[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^6(a + bx) \tan^4(a + bx) dx &= \frac{\text{Subst}\left(\int x^4(1 + x^2)^2 dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^4 + 2x^6 + x^8) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^5(a + bx)}{5b} + \frac{2 \tan^7(a + bx)}{7b} + \frac{\tan^9(a + bx)}{9b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 98 vs. 2(46) = 92.

time = 0.03, size = 98, normalized size = 2.13

$$\frac{8 \tan(a+bx)}{315b} + \frac{4 \sec^2(a+bx) \tan(a+bx)}{315b} + \frac{\sec^4(a+bx) \tan(a+bx)}{105b} - \frac{10 \sec^6(a+bx) \tan(a+bx)}{63b} + \frac{\sec^8(a+bx) \tan(a+bx)}{9b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^6*Tan[a + b*x]^4,x]

[Out] (8*Tan[a + b*x])/(315*b) + (4*Sec[a + b*x]^2*Tan[a + b*x])/(315*b) + (Sec[a + b*x]^4*Tan[a + b*x])/(105*b) - (10*Sec[a + b*x]^6*Tan[a + b*x])/(63*b) + (Sec[a + b*x]^8*Tan[a + b*x])/(9*b)

Maple [A]

time = 0.05, size = 60, normalized size = 1.30

method	result	size
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{9 \cos(bx+a)^9} + \frac{4(\sin^5(bx+a))}{63 \cos(bx+a)^7} + \frac{8(\sin^5(bx+a))}{315 \cos(bx+a)^5}}{b}$	60
default	$\frac{\frac{\sin^5(bx+a)}{9 \cos(bx+a)^9} + \frac{4(\sin^5(bx+a))}{63 \cos(bx+a)^7} + \frac{8(\sin^5(bx+a))}{315 \cos(bx+a)^5}}{b}$	60
risch	$\frac{16i(210e^{12i(bx+a)} - 315e^{10i(bx+a)} + 441e^{8i(bx+a)} - 126e^{6i(bx+a)} + 36e^{4i(bx+a)} + 9e^{2i(bx+a)} + 1)}{315b(e^{2i(bx+a)} + 1)^9}$	88

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^10*sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] 1/b*(1/9*sin(b*x+a)^5/cos(b*x+a)^9+4/63*sin(b*x+a)^5/cos(b*x+a)^7+8/315*sin(b*x+a)^5/cos(b*x+a)^5)

Maxima [A]

time = 0.30, size = 36, normalized size = 0.78

$$\frac{35 \tan(bx+a)^9 + 90 \tan(bx+a)^7 + 63 \tan(bx+a)^5}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^10*sin(b*x+a)^4,x, algorithm="maxima")

[Out] 1/315*(35*tan(b*x + a)^9 + 90*tan(b*x + a)^7 + 63*tan(b*x + a)^5)/b

Fricas [A]

time = 0.36, size = 61, normalized size = 1.33

$$\frac{(8 \cos(bx+a)^8 + 4 \cos(bx+a)^6 + 3 \cos(bx+a)^4 - 50 \cos(bx+a)^2 + 35) \sin(bx+a)}{315b \cos(bx+a)^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^10*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/315*(8*cos(b*x + a)^8 + 4*cos(b*x + a)^6 + 3*cos(b*x + a)^4 - 50*cos(b*x + a)^2 + 35)*sin(b*x + a)/(b*cos(b*x + a)^9)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**10*sin(b*x+a)**4,x)

[Out] Timed out

Giac [A]

time = 4.68, size = 36, normalized size = 0.78

$$\frac{35 \tan(bx + a)^9 + 90 \tan(bx + a)^7 + 63 \tan(bx + a)^5}{315 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^10*sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/315*(35*tan(b*x + a)^9 + 90*tan(b*x + a)^7 + 63*tan(b*x + a)^5)/b

Mupad [B]

time = 0.43, size = 35, normalized size = 0.76

$$\frac{\tan(a + bx)^5 (35 \tan(a + bx)^4 + 90 \tan(a + bx)^2 + 63)}{315 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/cos(a + b*x)^10,x)

[Out] (tan(a + b*x)^5*(90*tan(a + b*x)^2 + 35*tan(a + b*x)^4 + 63))/(315*b)

3.89 $\int \cos^6(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=111

$$\frac{3x}{256} + \frac{3 \cos(a + bx) \sin(a + bx)}{256b} + \frac{\cos^3(a + bx) \sin(a + bx)}{128b} + \frac{\cos^5(a + bx) \sin(a + bx)}{160b} - \frac{3 \cos^7(a + bx) \sin(a + bx)}{80b}$$

[Out] 3/256*x+3/256*cos(b*x+a)*sin(b*x+a)/b+1/128*cos(b*x+a)^3*sin(b*x+a)/b+1/160*cos(b*x+a)^5*sin(b*x+a)/b-3/80*cos(b*x+a)^7*sin(b*x+a)/b-1/10*cos(b*x+a)^7*sin(b*x+a)^3/b

Rubi [A]

time = 0.07, antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2648, 2715, 8}

$$-\frac{\sin^3(a + bx) \cos^7(a + bx)}{10b} - \frac{3 \sin(a + bx) \cos^7(a + bx)}{80b} + \frac{\sin(a + bx) \cos^5(a + bx)}{160b} + \frac{\sin(a + bx) \cos^3(a + bx)}{128b} + \frac{3 \sin(a + bx) \cos(a + bx)}{256b} + \frac{3x}{256}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^6*Sin[a + b*x]^4,x]

[Out] (3*x)/256 + (3*Cos[a + b*x]*Sin[a + b*x])/(256*b) + (Cos[a + b*x]^3*Sin[a + b*x])/(128*b) + (Cos[a + b*x]^5*Sin[a + b*x])/(160*b) - (3*Cos[a + b*x]^7*Sin[a + b*x])/(80*b) - (Cos[a + b*x]^7*Sin[a + b*x]^3)/(10*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2648

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^n]*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2715

Int[((b_)*sin[(c_) + (d_)*(x_)])^n], x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \cos^6(a+bx) \sin^4(a+bx) dx &= -\frac{\cos^7(a+bx) \sin^3(a+bx)}{10b} + \frac{3}{10} \int \cos^6(a+bx) \sin^2(a+bx) dx \\
&= -\frac{3 \cos^7(a+bx) \sin(a+bx)}{80b} - \frac{\cos^7(a+bx) \sin^3(a+bx)}{10b} + \frac{3}{80} \int \cos^6(a+bx) dx \\
&= \frac{\cos^5(a+bx) \sin(a+bx)}{160b} - \frac{3 \cos^7(a+bx) \sin(a+bx)}{80b} - \frac{\cos^7(a+bx) \sin^3(a+bx)}{10b} \\
&= \frac{\cos^3(a+bx) \sin(a+bx)}{128b} + \frac{\cos^5(a+bx) \sin(a+bx)}{160b} - \frac{3 \cos^7(a+bx) \sin(a+bx)}{80b} \\
&= \frac{3 \cos(a+bx) \sin(a+bx)}{256b} + \frac{\cos^3(a+bx) \sin(a+bx)}{128b} + \frac{\cos^5(a+bx) \sin(a+bx)}{160b} \\
&= \frac{3x}{256} + \frac{3 \cos(a+bx) \sin(a+bx)}{256b} + \frac{\cos^3(a+bx) \sin(a+bx)}{128b} + \frac{\cos^5(a+bx)}{160b}
\end{aligned}$$

Mathematica [A]

time = 0.14, size = 62, normalized size = 0.56

$$\frac{120bx + 20 \sin(2(a+bx)) - 40 \sin(4(a+bx)) - 10 \sin(6(a+bx)) + 5 \sin(8(a+bx)) + 2 \sin(10(a+bx))}{10240b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^6*Sin[a + b*x]^4,x]`

```
[Out] (120*b*x + 20*Sin[2*(a + b*x)] - 40*Sin[4*(a + b*x)] - 10*Sin[6*(a + b*x)]
+ 5*Sin[8*(a + b*x)] + 2*Sin[10*(a + b*x)])/(10240*b)
```

Maple [A]

time = 0.12, size = 82, normalized size = 0.74

method	result
risch	$\frac{3x}{256} + \frac{\sin(10bx+10a)}{5120b} + \frac{\sin(8bx+8a)}{2048b} - \frac{\sin(6bx+6a)}{1024b} - \frac{\sin(4bx+4a)}{256b} + \frac{\sin(2bx+2a)}{512b}$
derivativedivides	$-\frac{(\cos^7(bx+a))(\sin^3(bx+a))}{10} - \frac{3(\cos^7(bx+a))\sin(bx+a)}{80} + \frac{\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15\cos(bx+a)}{8}\right)\sin(bx+a)}{160} + \frac{3bx}{256} + \frac{3a}{256}$
default	$-\frac{(\cos^7(bx+a))(\sin^3(bx+a))}{10} - \frac{3(\cos^7(bx+a))\sin(bx+a)}{80} + \frac{\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15\cos(bx+a)}{8}\right)\sin(bx+a)}{160} + \frac{3bx}{256} + \frac{3a}{256}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^6*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/10*cos(b*x+a)^7*sin(b*x+a)^3-3/80*cos(b*x+a)^7*sin(b*x+a)+1/160*(co
s(b*x+a)^5+5/4*cos(b*x+a)^3+15/8*cos(b*x+a))*sin(b*x+a)+3/256*b*x+3/256*a)
```

Maxima [A]

time = 0.29, size = 48, normalized size = 0.43

$$\frac{32 \sin(2bx + 2a)^5 + 120bx + 120a + 5 \sin(8bx + 8a) - 40 \sin(4bx + 4a)}{10240b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^4,x, algorithm="maxima")

[Out] 1/10240*(32*sin(2*b*x + 2*a)^5 + 120*b*x + 120*a + 5*sin(8*b*x + 8*a) - 40*sin(4*b*x + 4*a))/b

Fricas [A]

time = 0.40, size = 66, normalized size = 0.59

$$\frac{15bx + (128 \cos(bx + a)^9 - 176 \cos(bx + a)^7 + 8 \cos(bx + a)^5 + 10 \cos(bx + a)^3 + 15 \cos(bx + a)) \sin(bx + a)}{1280b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/1280*(15*b*x + (128*cos(b*x + a)^9 - 176*cos(b*x + a)^7 + 8*cos(b*x + a)^5 + 10*cos(b*x + a)^3 + 15*cos(b*x + a))*sin(b*x + a))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 231 vs. $2(100) = 200$.

time = 1.76, size = 231, normalized size = 2.08

$$\begin{cases} \frac{3x \sin^{10}(a+bx)}{256} + \frac{15x \sin^8(a+bx) \cos^2(a+bx)}{256} + \frac{15x \sin^6(a+bx) \cos^4(a+bx)}{128} + \frac{15x \sin^4(a+bx) \cos^6(a+bx)}{128} + \frac{15x \sin^2(a+bx) \cos^8(a+bx)}{256} + \frac{3x \cos^{10}(a+bx)}{256} + \frac{3 \sin^8(a+bx) \cos(a+bx)}{2560} + \frac{7 \sin^7(a+bx) \cos^2(a+bx)}{1280} + \frac{\sin^6(a+bx) \cos^3(a+bx)}{100} - \frac{7 \sin^5(a+bx) \cos^4(a+bx)}{1280} - \frac{3 \sin^4(a+bx) \cos^5(a+bx)}{2560} & \text{for } b \neq 0 \\ x \sin^4(a) \cos^6(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**6*sin(b*x+a)**4,x)

[Out] Piecewise((3*x*sin(a + b*x)**10/256 + 15*x*sin(a + b*x)**8*cos(a + b*x)**2/256 + 15*x*sin(a + b*x)**6*cos(a + b*x)**4/128 + 15*x*sin(a + b*x)**4*cos(a + b*x)**6/128 + 15*x*sin(a + b*x)**2*cos(a + b*x)**8/256 + 3*x*cos(a + b*x)**10/256 + 3*sin(a + b*x)**9*cos(a + b*x)/(256*b) + 7*sin(a + b*x)**7*cos(a + b*x)**3/(128*b) + sin(a + b*x)**5*cos(a + b*x)**5/(10*b) - 7*sin(a + b*x)**3*cos(a + b*x)**7/(128*b) - 3*sin(a + b*x)*cos(a + b*x)**9/(256*b), Ne(b, 0)), (x*sin(a)**4*cos(a)**6, True))

Giac [A]

time = 3.88, size = 74, normalized size = 0.67

$$\frac{3}{256}x + \frac{\sin(10bx + 10a)}{5120b} + \frac{\sin(8bx + 8a)}{2048b} - \frac{\sin(6bx + 6a)}{1024b} - \frac{\sin(4bx + 4a)}{256b} + \frac{\sin(2bx + 2a)}{512b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^4,x, algorithm="giac")

[Out] $\frac{3}{256}x + \frac{1}{5120}\sin(10bx + 10a)/b + \frac{1}{2048}\sin(8bx + 8a)/b - \frac{1}{1024}\sin(6bx + 6a)/b - \frac{1}{256}\sin(4bx + 4a)/b + \frac{1}{512}\sin(2bx + 2a)/b$

Mupad [B]

time = 1.94, size = 109, normalized size = 0.98

$$\frac{3x}{256} + \frac{\frac{3\tan(a+bx)^9}{256} + \frac{7\tan(a+bx)^7}{128} + \frac{\tan(a+bx)^5}{10} - \frac{7\tan(a+bx)^3}{128} - \frac{3\tan(a+bx)}{256}}{b(\tan(a+bx)^{10} + 5\tan(a+bx)^8 + 10\tan(a+bx)^6 + 10\tan(a+bx)^4 + 5\tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6*sin(a + b*x)^4,x)

[Out] $\frac{(3x)/256 + (\tan(a + bx))^5/10 - (7\tan(a + bx)^3)/128 - (3\tan(a + bx))/256 + (7\tan(a + bx)^7)/128 + (3\tan(a + bx)^9)/256}{(b(5\tan(a + bx)^2 + 10\tan(a + bx)^4 + 10\tan(a + bx)^6 + 5\tan(a + bx)^8 + \tan(a + bx)^{10} + 1))}$

3.90 $\int \cos^4(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=90

$$\frac{3x}{128} + \frac{3 \cos(a + bx) \sin(a + bx)}{128b} + \frac{\cos^3(a + bx) \sin(a + bx)}{64b} - \frac{\cos^5(a + bx) \sin(a + bx)}{16b} - \frac{\cos^5(a + bx) \sin^3(a + bx)}{8b}$$

[Out] 3/128*x+3/128*cos(b*x+a)*sin(b*x+a)/b+1/64*cos(b*x+a)^3*sin(b*x+a)/b-1/16*cos(b*x+a)^5*sin(b*x+a)/b-1/8*cos(b*x+a)^5*sin(b*x+a)^3/b

Rubi [A]

time = 0.06, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2648, 2715, 8}

$$-\frac{\sin^3(a + bx) \cos^5(a + bx)}{8b} - \frac{\sin(a + bx) \cos^5(a + bx)}{16b} + \frac{\sin(a + bx) \cos^3(a + bx)}{64b} + \frac{3 \sin(a + bx) \cos(a + bx)}{128b} + \frac{3x}{128}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Sin[a + b*x]^4,x]

[Out] (3*x)/128 + (3*Cos[a + b*x]*Sin[a + b*x])/(128*b) + (Cos[a + b*x]^3*Sin[a + b*x])/(64*b) - (Cos[a + b*x]^5*Sin[a + b*x])/(16*b) - (Cos[a + b*x]^5*Sin[a + b*x]^3)/(8*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n]*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^n], x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Sin[c + d*x])^(n - 1)/(d*n)), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \cos^4(a+bx) \sin^4(a+bx) dx &= -\frac{\cos^5(a+bx) \sin^3(a+bx)}{8b} + \frac{3}{8} \int \cos^4(a+bx) \sin^2(a+bx) dx \\
&= -\frac{\cos^5(a+bx) \sin(a+bx)}{16b} - \frac{\cos^5(a+bx) \sin^3(a+bx)}{8b} + \frac{1}{16} \int \cos^4(a+bx) dx \\
&= \frac{\cos^3(a+bx) \sin(a+bx)}{64b} - \frac{\cos^5(a+bx) \sin(a+bx)}{16b} - \frac{\cos^5(a+bx) \sin^3(a+bx)}{8b} \\
&= \frac{3 \cos(a+bx) \sin(a+bx)}{128b} + \frac{\cos^3(a+bx) \sin(a+bx)}{64b} - \frac{\cos^5(a+bx) \sin(a+bx)}{16b} \\
&= \frac{3x}{128} + \frac{3 \cos(a+bx) \sin(a+bx)}{128b} + \frac{\cos^3(a+bx) \sin(a+bx)}{64b} - \frac{\cos^5(a+bx) \sin(a+bx)}{16b}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 33, normalized size = 0.37

$$\frac{24(a+bx) - 8 \sin(4(a+bx)) + \sin(8(a+bx))}{1024b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^4*Sin[a + b*x]^4,x]``[Out] (24*(a + b*x) - 8*Sin[4*(a + b*x)] + Sin[8*(a + b*x)])/(1024*b)`**Maple [A]**

time = 0.10, size = 72, normalized size = 0.80

method	result
risch	$\frac{3x}{128} + \frac{\sin(8bx+8a)}{1024b} - \frac{\sin(4bx+4a)}{128b}$
derivativedivides	$-\frac{(\cos^5(bx+a))(\sin^3(bx+a))}{8} - \frac{(\cos^5(bx+a))\sin(bx+a)}{16} + \frac{(\cos^3(bx+a) + \frac{3\cos(bx+a)}{2})\sin(bx+a)}{64} + \frac{3bx}{128} + \frac{3a}{128}$
default	$-\frac{(\cos^5(bx+a))(\sin^3(bx+a))}{8} - \frac{(\cos^5(bx+a))\sin(bx+a)}{16} + \frac{(\cos^3(bx+a) + \frac{3\cos(bx+a)}{2})\sin(bx+a)}{64} + \frac{3bx}{128} + \frac{3a}{128}$
norman	$\frac{3x}{128} - \frac{3 \tan(\frac{bx+a}{2})}{64b} - \frac{23(\tan^3(\frac{bx+a}{2}))}{64b} + \frac{333(\tan^5(\frac{bx+a}{2}))}{64b} - \frac{671(\tan^7(\frac{bx+a}{2}))}{64b} + \frac{671(\tan^9(\frac{bx+a}{2}))}{64b} - \frac{333(\tan^{11}(\frac{bx+a}{2}))}{64b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^4*sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/8*cos(b*x+a)^5*sin(b*x+a)^3-1/16*cos(b*x+a)^5*sin(b*x+a)+1/64*(cos(b*x+a)^3+3/2*cos(b*x+a))*sin(b*x+a)+3/128*b*x+3/128*a)`

Maxima [A]

time = 0.30, size = 33, normalized size = 0.37

$$\frac{24bx + 24a + \sin(8bx + 8a) - 8\sin(4bx + 4a)}{1024b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^4*sin(b*x+a)^4,x, algorithm="maxima")``[Out] 1/1024*(24*b*x + 24*a + sin(8*b*x + 8*a) - 8*sin(4*b*x + 4*a))/b`**Fricas [A]**

time = 0.39, size = 56, normalized size = 0.62

$$\frac{3bx + (16\cos(bx + a))^7 - 24\cos(bx + a)^5 + 2\cos(bx + a)^3 + 3\cos(bx + a)\sin(bx + a)}{128b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^4*sin(b*x+a)^4,x, algorithm="fricas")``[Out] 1/128*(3*b*x + (16*cos(b*x + a))^7 - 24*cos(b*x + a)^5 + 2*cos(b*x + a)^3 + 3*cos(b*x + a))*sin(b*x + a)/b`**Sympy [B]** Leaf count of result is larger than twice the leaf count of optimal. 189 vs. $2(80) = 160$.

time = 1.67, size = 189, normalized size = 2.10

$$\begin{cases} \frac{3x\sin^6(a+bx)}{128} + \frac{3x\sin^6(a+bx)\cos^2(a+bx)}{32} + \frac{9x\sin^4(a+bx)\cos^4(a+bx)}{64} + \frac{3x\sin^2(a+bx)\cos^6(a+bx)}{32} + \frac{3x\cos^8(a+bx)}{128} + \frac{3\sin^7(a+bx)\cos(a+bx)}{128b} + \frac{11\sin^5(a+bx)\cos^3(a+bx)}{128b} - \frac{11\sin^3(a+bx)\cos^5(a+bx)}{128b} - \frac{3\sin(a+bx)\cos^7(a+bx)}{128b} & \text{for } b \neq 0 \\ x\sin^4(a)\cos^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)**4*sin(b*x+a)**4,x)`

```
[Out] Piecewise((3*x*sin(a + b*x)**8/128 + 3*x*sin(a + b*x)**6*cos(a + b*x)**2/32
+ 9*x*sin(a + b*x)**4*cos(a + b*x)**4/64 + 3*x*sin(a + b*x)**2*cos(a + b*x)
)**6/32 + 3*x*cos(a + b*x)**8/128 + 3*sin(a + b*x)**7*cos(a + b*x)/(128*b)
+ 11*sin(a + b*x)**5*cos(a + b*x)**3/(128*b) - 11*sin(a + b*x)**3*cos(a + b
*x)**5/(128*b) - 3*sin(a + b*x)*cos(a + b*x)**7/(128*b), Ne(b, 0)), (x*sin(
a)**4*cos(a)**4, True))
```

Giac [A]

time = 3.43, size = 32, normalized size = 0.36

$$\frac{3}{128}x + \frac{\sin(8bx + 8a)}{1024b} - \frac{\sin(4bx + 4a)}{128b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*sin(b*x+a)^4,x, algorithm="giac")

[Out] 3/128*x + 1/1024*sin(8*b*x + 8*a)/b - 1/128*sin(4*b*x + 4*a)/b

Mupad [B]

time = 1.50, size = 90, normalized size = 1.00

$$\frac{3x}{128} - \frac{-\frac{3 \tan(a+bx)^7}{128} - \frac{11 \tan(a+bx)^5}{128} + \frac{11 \tan(a+bx)^3}{128} + \frac{3 \tan(a+bx)}{128}}{b (\tan(a+bx)^8 + 4 \tan(a+bx)^6 + 6 \tan(a+bx)^4 + 4 \tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4*sin(a + b*x)^4,x)

[Out] (3*x)/128 - ((3*tan(a + b*x))/128 + (11*tan(a + b*x)^3)/128 - (11*tan(a + b*x)^5)/128 - (3*tan(a + b*x)^7)/128)/(b*(4*tan(a + b*x)^2 + 6*tan(a + b*x)^4 + 4*tan(a + b*x)^6 + tan(a + b*x)^8 + 1))

3.91 $\int \cos^2(a + bx) \sin^4(a + bx) dx$

Optimal. Leaf size=69

$$\frac{x}{16} + \frac{\cos(a + bx) \sin(a + bx)}{16b} - \frac{\cos^3(a + bx) \sin(a + bx)}{8b} - \frac{\cos^3(a + bx) \sin^3(a + bx)}{6b}$$

[Out] 1/16*x+1/16*cos(b*x+a)*sin(b*x+a)/b-1/8*cos(b*x+a)^3*sin(b*x+a)/b-1/6*cos(b*x+a)^3*sin(b*x+a)^3/b

Rubi [A]

time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$,

Rules used = {2648, 2715, 8}

$$-\frac{\sin^3(a + bx) \cos^3(a + bx)}{6b} - \frac{\sin(a + bx) \cos^3(a + bx)}{8b} + \frac{\sin(a + bx) \cos(a + bx)}{16b} + \frac{x}{16}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Sin[a + b*x]^4,x]

[Out] x/16 + (Cos[a + b*x]*Sin[a + b*x])/(16*b) - (Cos[a + b*x]^3*Sin[a + b*x])/(8*b) - (Cos[a + b*x]^3*Sin[a + b*x]^3)/(6*b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Sin[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*Sin[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \cos^2(a+bx) \sin^4(a+bx) dx &= -\frac{\cos^3(a+bx) \sin^3(a+bx)}{6b} + \frac{1}{2} \int \cos^2(a+bx) \sin^2(a+bx) dx \\
&= -\frac{\cos^3(a+bx) \sin(a+bx)}{8b} - \frac{\cos^3(a+bx) \sin^3(a+bx)}{6b} + \frac{1}{8} \int \cos^2(a+bx) dx \\
&= \frac{\cos(a+bx) \sin(a+bx)}{16b} - \frac{\cos^3(a+bx) \sin(a+bx)}{8b} - \frac{\cos^3(a+bx) \sin^3(a+bx)}{6b} \\
&= \frac{x}{16} + \frac{\cos(a+bx) \sin(a+bx)}{16b} - \frac{\cos^3(a+bx) \sin(a+bx)}{8b} - \frac{\cos^3(a+bx) \sin^3(a+bx)}{6b}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 40, normalized size = 0.58

$$\frac{12bx - 3 \sin(2(a+bx)) - 3 \sin(4(a+bx)) + \sin(6(a+bx))}{192b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^2*Sin[a + b*x]^4,x]``[Out] (12*b*x - 3*Sin[2*(a + b*x)] - 3*Sin[4*(a + b*x)] + Sin[6*(a + b*x)])/(192*b)`**Maple [A]**

time = 0.07, size = 61, normalized size = 0.88

method	result
risch	$\frac{x}{16} + \frac{\sin(6bx+6a)}{192b} - \frac{\sin(4bx+4a)}{64b} - \frac{\sin(2bx+2a)}{64b}$
derivativedivides	$-\frac{(\cos^3(bx+a))(\sin^3(bx+a))}{6} - \frac{(\cos^3(bx+a))\sin(bx+a)}{8} + \frac{\cos(bx+a)\sin(bx+a)}{16} + \frac{bx}{16} + \frac{a}{16}$
default	$-\frac{(\cos^3(bx+a))(\sin^3(bx+a))}{6} - \frac{(\cos^3(bx+a))\sin(bx+a)}{8} + \frac{\cos(bx+a)\sin(bx+a)}{16} + \frac{bx}{16} + \frac{a}{16}$
norman	$\frac{x}{16} - \frac{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{17\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} + \frac{19\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} - \frac{19\left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{17\left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} + \frac{\tan^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} + \frac{3x\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8} + \frac{3a\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^2*sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/6*cos(b*x+a)^3*sin(b*x+a)^3-1/8*cos(b*x+a)^3*sin(b*x+a)+1/16*cos(b*x+a)*sin(b*x+a)+1/16*b*x+1/16*a)`**Maxima [A]**

time = 0.33, size = 37, normalized size = 0.54

$$-\frac{4 \sin(2bx + 2a)^3 - 12bx - 12a + 3 \sin(4bx + 4a)}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^4,x, algorithm="maxima")

[Out] -1/192*(4*sin(2*b*x + 2*a)^3 - 12*b*x - 12*a + 3*sin(4*b*x + 4*a))/b

Fricas [A]

time = 0.36, size = 46, normalized size = 0.67

$$\frac{3bx + (8 \cos(bx + a)^5 - 14 \cos(bx + a)^3 + 3 \cos(bx + a)) \sin(bx + a)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/48*(3*b*x + (8*cos(b*x + a)^5 - 14*cos(b*x + a)^3 + 3*cos(b*x + a))*sin(b*x + a))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 136 vs. 2(58) = 116.

time = 0.57, size = 136, normalized size = 1.97

$$\begin{cases} \frac{x \sin^6(a+bx)}{16} + \frac{3x \sin^4(a+bx) \cos^2(a+bx)}{16} + \frac{3x \sin^2(a+bx) \cos^4(a+bx)}{16} + \frac{x \cos^6(a+bx)}{16} + \frac{\sin^5(a+bx) \cos(a+bx)}{16b} - \frac{\sin^3(a+bx) \cos^3(a+bx)}{6b} - \frac{\sin(a+bx) \cos^5(a+bx)}{16b} & \text{for } b \neq 0 \\ x \sin^4(a) \cos^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2*sin(b*x+a)**4,x)

[Out] Piecewise((x*sin(a + b*x)**6/16 + 3*x*sin(a + b*x)**4*cos(a + b*x)**2/16 + 3*x*sin(a + b*x)**2*cos(a + b*x)**4/16 + x*cos(a + b*x)**6/16 + sin(a + b*x)**5*cos(a + b*x)/(16*b) - sin(a + b*x)**3*cos(a + b*x)**3/(6*b) - sin(a + b*x)*cos(a + b*x)**5/(16*b), Ne(b, 0)), (x*sin(a)**4*cos(a)**2, True))

Giac [A]

time = 3.26, size = 46, normalized size = 0.67

$$\frac{1}{16}x + \frac{\sin(6bx + 6a)}{192b} - \frac{\sin(4bx + 4a)}{64b} - \frac{\sin(2bx + 2a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/16*x + 1/192*sin(6*b*x + 6*a)/b - 1/64*sin(4*b*x + 4*a)/b - 1/64*sin(2*b*x + 2*a)/b

Mupad [B]

time = 0.52, size = 43, normalized size = 0.62

$$\frac{x}{16} - \frac{\sin(2a+2bx)}{64} + \frac{\sin(4a+4bx)}{64} - \frac{\sin(6a+6bx)}{192}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^2*sin(a + b*x)^4,x)
```

```
[Out] x/16 - (sin(2*a + 2*b*x)/64 + sin(4*a + 4*b*x)/64 - sin(6*a + 6*b*x)/192)/b
```

3.92 $\int \sin^4(a + bx) dx$

Optimal. Leaf size=46

$$\frac{3x}{8} - \frac{3 \cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos(a + bx) \sin^3(a + bx)}{4b}$$

[Out] $3/8*x - 3/8*\cos(b*x+a)*\sin(b*x+a)/b - 1/4*\cos(b*x+a)*\sin(b*x+a)^3/b$

Rubi [A]

time = 0.01, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {2715, 8}

$$-\frac{\sin^3(a + bx) \cos(a + bx)}{4b} - \frac{3 \sin(a + bx) \cos(a + bx)}{8b} + \frac{3x}{8}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^4,x]

[Out] $(3*x)/8 - (3*\text{Cos}[a + b*x]*\text{Sin}[a + b*x])/(8*b) - (\text{Cos}[a + b*x]*\text{Sin}[a + b*x]^3)/(4*b)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2715

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*SIN[c + d*x])^(n - 1)/(d*n), x] + Dist[b^2*((n - 1)/n), Int[(b*SIN[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \sin^4(a + bx) dx &= -\frac{\cos(a + bx) \sin^3(a + bx)}{4b} + \frac{3}{4} \int \sin^2(a + bx) dx \\ &= -\frac{3 \cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos(a + bx) \sin^3(a + bx)}{4b} + \frac{3 \int 1 dx}{8} \\ &= \frac{3x}{8} - \frac{3 \cos(a + bx) \sin(a + bx)}{8b} - \frac{\cos(a + bx) \sin^3(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 33, normalized size = 0.72

$$\frac{12(a + bx) - 8 \sin(2(a + bx)) + \sin(4(a + bx))}{32b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^4,x]``[Out] (12*(a + b*x) - 8*Sin[2*(a + b*x)] + Sin[4*(a + b*x)])/(32*b)`**Maple [A]**

time = 0.00, size = 38, normalized size = 0.83

method	result
risch	$\frac{3x}{8} + \frac{\sin(4bx+4a)}{32b} - \frac{\sin(2bx+2a)}{4b}$
derivativdivides	$-\frac{\left(\sin^3(bx+a) + \frac{3\sin(bx+a)}{2}\right)\cos(bx+a)}{4} + \frac{3bx}{8} + \frac{3a}{8}$
default	$-\frac{\left(\sin^3(bx+a) + \frac{3\sin(bx+a)}{2}\right)\cos(bx+a)}{4} + \frac{3bx}{8} + \frac{3a}{8}$
norman	$\frac{\frac{3x}{8} - \frac{3\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{4b} - \frac{11\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{11\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{3\left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{3x\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2} + \frac{9x\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4} + \frac{3x\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/4*(sin(b*x+a)^3+3/2*sin(b*x+a))*cos(b*x+a)+3/8*b*x+3/8*a)`**Maxima [A]**

time = 0.29, size = 33, normalized size = 0.72

$$\frac{12bx + 12a + \sin(4bx + 4a) - 8 \sin(2bx + 2a)}{32b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^4,x, algorithm="maxima")``[Out] 1/32*(12*b*x + 12*a + sin(4*b*x + 4*a) - 8*sin(2*b*x + 2*a))/b`**Fricas [A]**

time = 0.33, size = 36, normalized size = 0.78

$$\frac{3bx + (2 \cos(bx + a))^3 - 5 \cos(bx + a) \sin(bx + a)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4,x, algorithm="fricas")

[Out] $1/8*(3*b*x + (2*\cos(b*x + a))^3 - 5*\cos(b*x + a))*\sin(b*x + a)/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(41) = 82$.

time = 0.17, size = 95, normalized size = 2.07

$$\begin{cases} \frac{3x \sin^4(a+bx)}{8} + \frac{3x \sin^2(a+bx) \cos^2(a+bx)}{4} + \frac{3x \cos^4(a+bx)}{8} - \frac{5 \sin^3(a+bx) \cos(a+bx)}{8b} - \frac{3 \sin(a+bx) \cos^3(a+bx)}{8b} & \text{for } b \neq 0 \\ x \sin^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**4,x)

[Out] Piecewise((3*x*sin(a + b*x)**4/8 + 3*x*sin(a + b*x)**2*cos(a + b*x)**2/4 + 3*x*cos(a + b*x)**4/8 - 5*sin(a + b*x)**3*cos(a + b*x)/(8*b) - 3*sin(a + b*x)*cos(a + b*x)**3/(8*b), Ne(b, 0)), (x*sin(a)**4, True))

Giac [A]

time = 3.05, size = 32, normalized size = 0.70

$$\frac{3}{8}x + \frac{\sin(4bx + 4a)}{32b} - \frac{\sin(2bx + 2a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4,x, algorithm="giac")

[Out] $3/8*x + 1/32*\sin(4*b*x + 4*a)/b - 1/4*\sin(2*b*x + 2*a)/b$

Mupad [B]

time = 0.42, size = 50, normalized size = 1.09

$$\frac{3x}{8} - \frac{\frac{5 \tan(a+bx)^3}{8} + \frac{3 \tan(a+bx)}{8}}{b (\tan(a+bx)^4 + 2 \tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4,x)

[Out] $(3*x)/8 - ((3*\tan(a + b*x))/8 + (5*\tan(a + b*x)^3)/8)/(b*(2*\tan(a + b*x)^2 + \tan(a + b*x)^4 + 1))$

3.93 $\int \sin^3(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=38

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

[Out] arctanh(sin(b*x+a))/b-sin(b*x+a)/b-1/3*sin(b*x+a)^3/b

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2672, 308, 212}

$$-\frac{\sin^3(a + bx)}{3b} - \frac{\sin(a + bx)}{b} + \frac{\tanh^{-1}(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3*Tan[a + b*x],x]

[Out] ArcTanh[Sin[a + b*x]]/b - Sin[a + b*x]/b - Sin[a + b*x]^3/(3*b)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 308

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2672

Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]

Rubi steps

$$\begin{aligned}
\int \sin^3(a+bx) \tan(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{x^4}{1-x^2} dx, x, \sin(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(-1 - x^2 + \frac{1}{1-x^2}\right) dx, x, \sin(a+bx)\right)}{b} \\
&= -\frac{\sin(a+bx)}{b} - \frac{\sin^3(a+bx)}{3b} + \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sin(a+bx)\right)}{b} \\
&= \frac{\tanh^{-1}(\sin(a+bx))}{b} - \frac{\sin(a+bx)}{b} - \frac{\sin^3(a+bx)}{3b}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 38, normalized size = 1.00

$$\frac{\tanh^{-1}(\sin(a+bx))}{b} - \frac{\sin(a+bx)}{b} - \frac{\sin^3(a+bx)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^3*Tan[a + b*x],x]``[Out] ArcTanh[Sin[a + b*x]]/b - Sin[a + b*x]/b - Sin[a + b*x]^3/(3*b)`**Maple [A]**

time = 0.04, size = 38, normalized size = 1.00

method	result	size
derivativedivides	$\frac{-\frac{(\sin^3(bx+a))}{3} - \sin(bx+a) + \ln(\sec(bx+a) + \tan(bx+a))}{b}$	38
default	$\frac{-\frac{(\sin^3(bx+a))}{3} - \sin(bx+a) + \ln(\sec(bx+a) + \tan(bx+a))}{b}$	38
risch	$\frac{5ie^{i(bx+a)}}{8b} - \frac{5ie^{-i(bx+a)}}{8b} + \frac{\ln(e^{i(bx+a)}+i)}{b} - \frac{\ln(e^{i(bx+a)}-i)}{b} + \frac{\sin(3bx+3a)}{12b}$	81
norman	$\frac{-\frac{2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} - \frac{20(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right))}{3b} - \frac{2(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right))}{b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b}$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/3*sin(b*x+a)^3-sin(b*x+a)+ln(sec(b*x+a)+tan(b*x+a)))`**Maxima [A]**

time = 0.28, size = 46, normalized size = 1.21

$$\frac{2 \sin(bx+a)^3 - 3 \log(\sin(bx+a) + 1) + 3 \log(\sin(bx+a) - 1) + 6 \sin(bx+a)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^4,x, algorithm="maxima")

[Out] $-1/6*(2*\sin(b*x + a)^3 - 3*\log(\sin(b*x + a) + 1) + 3*\log(\sin(b*x + a) - 1) + 6*\sin(b*x + a))/b$

Fricas [A]

time = 0.37, size = 48, normalized size = 1.26

$$\frac{2(\cos(bx+a)^2 - 4)\sin(bx+a) + 3\log(\sin(bx+a) + 1) - 3\log(-\sin(bx+a) + 1)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^4,x, algorithm="fricas")

[Out] $1/6*(2*(\cos(b*x + a)^2 - 4)*\sin(b*x + a) + 3*\log(\sin(b*x + a) + 1) - 3*\log(-\sin(b*x + a) + 1))/b$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)**4,x)

[Out] Timed out

Giac [A]

time = 3.29, size = 48, normalized size = 1.26

$$\frac{2\sin(bx+a)^3 - 3\log(|\sin(bx+a) + 1|) + 3\log(|\sin(bx+a) - 1|) + 6\sin(bx+a)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^4,x, algorithm="giac")

[Out] $-1/6*(2*\sin(b*x + a)^3 - 3*\log(\text{abs}(\sin(b*x + a) + 1)) + 3*\log(\text{abs}(\sin(b*x + a) - 1)) + 6*\sin(b*x + a))/b$

Mupad [B]

time = 0.58, size = 53, normalized size = 1.39

$$\frac{2 \operatorname{atanh}\left(\frac{\sin\left(\frac{a}{2} + \frac{bx}{2}\right)}{\cos\left(\frac{a}{2} + \frac{bx}{2}\right)}\right)}{b} - \frac{5 \sin(a + bx)}{4b} + \frac{\sin(3a + 3bx)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/cos(a + b*x),x)

[Out] $(2*\operatorname{atanh}(\sin(a/2 + (b*x)/2)/\cos(a/2 + (b*x)/2)))/b - (5*\sin(a + b*x))/(4*b) + \sin(3*a + 3*b*x)/(12*b)$

3.94 $\int \sin(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=49

$$-\frac{3 \tanh^{-1}(\sin(a + bx))}{2b} + \frac{3 \sin(a + bx)}{2b} + \frac{\sin(a + bx) \tan^2(a + bx)}{2b}$$

[Out] $-3/2*\operatorname{arctanh}(\sin(b*x+a))/b+3/2*\sin(b*x+a)/b+1/2*\sin(b*x+a)*\tan(b*x+a)^2/b$

Rubi [A]

time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2672, 294, 327, 212}

$$\frac{3 \sin(a + bx)}{2b} + \frac{\sin(a + bx) \tan^2(a + bx)}{2b} - \frac{3 \tanh^{-1}(\sin(a + bx))}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]*Tan[a + b*x]^3,x]

[Out] $(-3*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]])/(2*b) + (3*\operatorname{Sin}[a + b*x])/(2*b) + (\operatorname{Sin}[a + b*x]*\operatorname{Tan}[a + b*x]^2)/(2*b)$

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2672

```
Int[((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(
ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rubi steps

$$\begin{aligned} \int \sin(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^4}{(1-x^2)^2} dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin(a + bx) \tan^2(a + bx)}{2b} - \frac{3 \text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \sin(a + bx)\right)}{2b} \\ &= \frac{3 \sin(a + bx)}{2b} + \frac{\sin(a + bx) \tan^2(a + bx)}{2b} - \frac{3 \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sin(a + bx)\right)}{2b} \\ &= -\frac{3 \tanh^{-1}(\sin(a + bx))}{2b} + \frac{3 \sin(a + bx)}{2b} + \frac{\sin(a + bx) \tan^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 40, normalized size = 0.82

$$\frac{-3 \tanh^{-1}(\sin(a + bx)) + (2 + \cos(2(a + bx))) \sec(a + bx) \tan(a + bx)}{2b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]*Tan[a + b*x]^3,x]
```

```
[Out] (-3*ArcTanh[Sin[a + b*x]] + (2 + Cos[2*(a + b*x)])*Sec[a + b*x]*Tan[a + b*x
])/ (2*b)
```

Maple [A]

time = 0.05, size = 58, normalized size = 1.18

method	result	size
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{2 \cos(bx+a)^2} + \frac{(\sin^3(bx+a))}{2} + \frac{3 \sin(bx+a)}{2} - \frac{3 \ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$	58
default	$\frac{\frac{\sin^5(bx+a)}{2 \cos(bx+a)^2} + \frac{(\sin^3(bx+a))}{2} + \frac{3 \sin(bx+a)}{2} - \frac{3 \ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$	58
risch	$-\frac{i e^{i(bx+a)}}{2b} + \frac{i e^{-i(bx+a)}}{2b} - \frac{i(e^{3i(bx+a)} - e^{i(bx+a)})}{b(e^{2i(bx+a)} + 1)^2} + \frac{3 \ln(e^{i(bx+a)} - i)}{2b} - \frac{3 \ln(e^{i(bx+a)} + i)}{2b}$	108

norman	$\frac{\frac{3 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} - \frac{2\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{3\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^2} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{2b} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{2b}$	114
--------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^3*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] `1/b*(1/2*sin(b*x+a)^5/cos(b*x+a)^2+1/2*sin(b*x+a)^3+3/2*sin(b*x+a)-3/2*ln(sec(b*x+a)+tan(b*x+a)))`

Maxima [A]

time = 0.29, size = 56, normalized size = 1.14

$$\frac{\frac{2 \sin(bx+a)}{\sin(bx+a)^2-1} + 3 \log(\sin(bx+a)+1) - 3 \log(\sin(bx+a)-1) - 4 \sin(bx+a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3*sin(b*x+a)^4,x, algorithm="maxima")`

[Out] `-1/4*(2*sin(b*x + a)/(sin(b*x + a)^2 - 1) + 3*log(sin(b*x + a) + 1) - 3*log(sin(b*x + a) - 1) - 4*sin(b*x + a))/b`

Fricas [A]

time = 0.39, size = 74, normalized size = 1.51

$$\frac{3 \cos(bx+a)^2 \log(\sin(bx+a)+1) - 3 \cos(bx+a)^2 \log(-\sin(bx+a)+1) - 2(2 \cos(bx+a)^2 + 1) \sin(bx+a)}{4b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3*sin(b*x+a)^4,x, algorithm="fricas")`

[Out] `-1/4*(3*cos(b*x + a)^2*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^2*log(-sin(b*x + a) + 1) - 2*(2*cos(b*x + a)^2 + 1)*sin(b*x + a))/(b*cos(b*x + a)^2)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^4(a + bx) \sec^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**3*sin(b*x+a)**4,x)`

[Out] `Integral(sin(a + b*x)**4*sec(a + b*x)**3, x)`

Giac [A]

time = 4.58, size = 58, normalized size = 1.18

$$\frac{\frac{2 \sin(bx+a)}{\sin(bx+a)^2-1} + 3 \log(|\sin(bx+a) + 1|) - 3 \log(|\sin(bx+a) - 1|) - 4 \sin(bx+a)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3*sin(b*x+a)^4,x, algorithm="giac")`

`[Out] -1/4*(2*sin(b*x + a)/(sin(b*x + a)^2 - 1) + 3*log(abs(sin(b*x + a) + 1)) - 3*log(abs(sin(b*x + a) - 1)) - 4*sin(b*x + a))/b`

Mupad [B]

time = 3.97, size = 98, normalized size = 2.00

$$\frac{3 \operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{b} - \frac{3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5 - 2 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3 + 3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{b \left(-\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 - 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^4/cos(a + b*x)^3,x)`

`[Out] - (3*atanh(tan(a/2 + (b*x)/2)))/b - (3*tan(a/2 + (b*x)/2) - 2*tan(a/2 + (b*x)/2)^3 + 3*tan(a/2 + (b*x)/2)^5)/(b*(tan(a/2 + (b*x)/2)^2 + tan(a/2 + (b*x)/2)^4 - tan(a/2 + (b*x)/2)^6 - 1))`

3.95 $\int \sec(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=55

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{8b} - \frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec(a + bx) \tan^3(a + bx)}{4b}$$

[Out] $3/8*\operatorname{arctanh}(\sin(b*x+a))/b-3/8*\sec(b*x+a)*\tan(b*x+a)/b+1/4*\sec(b*x+a)*\tan(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2691, 3855}

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{8b} + \frac{\tan^3(a + bx) \sec(a + bx)}{4b} - \frac{3 \tan(a + bx) \sec(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] `Int[Sec[a + b*x]*Tan[a + b*x]^4,x]`

[Out] $(3*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]])/(8*b) - (3*\operatorname{Sec}[a + b*x]*\operatorname{Tan}[a + b*x])/(8*b) + (\operatorname{Sec}[a + b*x]*\operatorname{Tan}[a + b*x]^3)/(4*b)$

Rule 2691

`Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]`

Rule 3855

`Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \int \sec(a + bx) \tan^4(a + bx) dx &= \frac{\sec(a + bx) \tan^3(a + bx)}{4b} - \frac{3}{4} \int \sec(a + bx) \tan^2(a + bx) dx \\ &= -\frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec(a + bx) \tan^3(a + bx)}{4b} + \frac{3}{8} \int \sec(a + bx) dx \\ &= \frac{3 \tanh^{-1}(\sin(a + bx))}{8b} - \frac{3 \sec(a + bx) \tan(a + bx)}{8b} + \frac{\sec(a + bx) \tan^3(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 45, normalized size = 0.82

$$\frac{6 \tanh^{-1}(\sin(a + bx)) - (1 + 5 \cos(2(a + bx))) \sec^3(a + bx) \tan(a + bx)}{16b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]*Tan[a + b*x]^4, x]``[Out] (6*ArcTanh[Sin[a + b*x]] - (1 + 5*Cos[2*(a + b*x)])*Sec[a + b*x]^3*Tan[a + b*x])/(16*b)`**Maple [A]**

time = 0.07, size = 76, normalized size = 1.38

method	result
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{4 \cos(bx+a)^4} - \frac{\sin^5(bx+a)}{8 \cos(bx+a)^2} - \frac{(\sin^3(bx+a))}{8} - \frac{3 \sin(bx+a)}{8} + \frac{3 \ln(\sec(bx+a)+\tan(bx+a))}{8}}{b}$
default	$\frac{\frac{\sin^5(bx+a)}{4 \cos(bx+a)^4} - \frac{\sin^5(bx+a)}{8 \cos(bx+a)^2} - \frac{(\sin^3(bx+a))}{8} - \frac{3 \sin(bx+a)}{8} + \frac{3 \ln(\sec(bx+a)+\tan(bx+a))}{8}}{b}$
risch	$\frac{i(5 e^{7i(bx+a)} - 3 e^{5i(bx+a)} + 3 e^{3i(bx+a)} - 5 e^{i(bx+a)})}{4b(e^{2i(bx+a)} + 1)^4} + \frac{3 \ln(e^{i(bx+a)} + i)}{8b} - \frac{3 \ln(e^{i(bx+a)} - i)}{8b}$
norman	$-\frac{3 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{4b} + \frac{11 \left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} + \frac{11 \left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} - \frac{3 \left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{8b} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^5*sin(b*x+a)^4, x, method=_RETURNVERBOSE)``[Out] 1/b*(1/4*sin(b*x+a)^5/cos(b*x+a)^4-1/8*sin(b*x+a)^5/cos(b*x+a)^2-1/8*sin(b*x+a)^3-3/8*sin(b*x+a)+3/8*ln(sec(b*x+a)+tan(b*x+a)))`**Maxima [A]**

time = 0.29, size = 71, normalized size = 1.29

$$\frac{2 \left(5 \sin(bx+a)^3 - 3 \sin(bx+a) \right)}{\sin(bx+a)^4 - 2 \sin(bx+a)^2 + 1} + 3 \log(\sin(bx+a) + 1) - 3 \log(\sin(bx+a) - 1)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5*sin(b*x+a)^4, x, algorithm="maxima")``[Out] 1/16*(2*(5*sin(b*x + a)^3 - 3*sin(b*x + a))/(sin(b*x + a)^4 - 2*sin(b*x + a)^2 + 1) + 3*log(sin(b*x + a) + 1) - 3*log(sin(b*x + a) - 1))/b`

Fricas [A]

time = 0.38, size = 74, normalized size = 1.35

$$\frac{3 \cos(bx+a)^4 \log(\sin(bx+a)+1) - 3 \cos(bx+a)^4 \log(-\sin(bx+a)+1) - 2(5 \cos(bx+a)^2 - 2) \sin(bx+a)}{16 b \cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5*sin(b*x+a)^4,x, algorithm="fricas")`

```
[Out] 1/16*(3*cos(b*x + a)^4*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^4*log(-sin(b*x + a) + 1) - 2*(5*cos(b*x + a)^2 - 2)*sin(b*x + a))/(b*cos(b*x + a)^4)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**5*sin(b*x+a)**4,x)`

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep
```

Giac [A]

time = 4.80, size = 63, normalized size = 1.15

$$\frac{2(5 \sin(bx+a)^3 - 3 \sin(bx+a))}{(\sin(bx+a)^2 - 1)^2} + 3 \log(|\sin(bx+a)+1|) - 3 \log(|\sin(bx+a)-1|)$$

$$16b$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5*sin(b*x+a)^4,x, algorithm="giac")`

```
[Out] 1/16*(2*(5*sin(b*x + a)^3 - 3*sin(b*x + a))/(sin(b*x + a)^2 - 1)^2 + 3*log(abs(sin(b*x + a) + 1)) - 3*log(abs(sin(b*x + a) - 1)))/b
```

Mupad [B]

time = 6.59, size = 126, normalized size = 2.29

$$\frac{3 \operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{4b} - \frac{\frac{3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^7}{4} - \frac{11 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5}{4} - \frac{11 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3}{4} + \frac{3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{4}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 - 4 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + 6 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 - 4 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^4/cos(a + b*x)^5,x)`

```
[Out] (3*atanh(tan(a/2 + (b*x)/2)))/(4*b) - ((3*tan(a/2 + (b*x)/2))/4 - (11*tan(a/2 + (b*x)/2)^3)/4 - (11*tan(a/2 + (b*x)/2)^5)/4 + (3*tan(a/2 + (b*x)/2)^7)/4)/(b*(6*tan(a/2 + (b*x)/2)^4 - 4*tan(a/2 + (b*x)/2)^2 - 4*tan(a/2 + (b*x)/2)^6 + tan(a/2 + (b*x)/2)^8 + 1))
```

3.96 $\int \sec^3(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=78

$$\frac{\tanh^{-1}(\sin(a + bx))}{16b} + \frac{\sec(a + bx) \tan(a + bx)}{16b} - \frac{\sec^3(a + bx) \tan(a + bx)}{8b} + \frac{\sec^3(a + bx) \tan^3(a + bx)}{6b}$$

[Out] 1/16*arctanh(sin(b*x+a))/b+1/16*sec(b*x+a)*tan(b*x+a)/b-1/8*sec(b*x+a)^3*tan(b*x+a)/b+1/6*sec(b*x+a)^3*tan(b*x+a)^3/b

Rubi [A]

time = 0.05, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2691, 3853, 3855}

$$\frac{\tanh^{-1}(\sin(a + bx))}{16b} + \frac{\tan^3(a + bx) \sec^3(a + bx)}{6b} - \frac{\tan(a + bx) \sec^3(a + bx)}{8b} + \frac{\tan(a + bx) \sec(a + bx)}{16b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^3*Tan[a + b*x]^4,x]

[Out] ArcTanh[Sin[a + b*x]]/(16*b) + (Sec[a + b*x]*Tan[a + b*x])/(16*b) - (Sec[a + b*x]^3*Tan[a + b*x])/(8*b) + (Sec[a + b*x]^3*Tan[a + b*x]^3)/(6*b)

Rule 2691

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sec^3(a+bx) \tan^4(a+bx) dx &= \frac{\sec^3(a+bx) \tan^3(a+bx)}{6b} - \frac{1}{2} \int \sec^3(a+bx) \tan^2(a+bx) dx \\
&= -\frac{\sec^3(a+bx) \tan(a+bx)}{8b} + \frac{\sec^3(a+bx) \tan^3(a+bx)}{6b} + \frac{1}{8} \int \sec^3(a+bx) dx \\
&= \frac{\sec(a+bx) \tan(a+bx)}{16b} - \frac{\sec^3(a+bx) \tan(a+bx)}{8b} + \frac{\sec^3(a+bx) \tan^3(a+bx)}{6b} \\
&= \frac{\tanh^{-1}(\sin(a+bx))}{16b} + \frac{\sec(a+bx) \tan(a+bx)}{16b} - \frac{\sec^3(a+bx) \tan(a+bx)}{8b}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 99, normalized size = 1.27

$$\frac{\tanh^{-1}(\sin(a+bx))}{16b} + \frac{\sec(a+bx) \tan(a+bx)}{16b} + \frac{\sec^3(a+bx) \tan(a+bx)}{24b} - \frac{\sec^5(a+bx) \tan(a+bx)}{6b} + \frac{\sec^3(a+bx) \tan^3(a+bx)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^3*Tan[a + b*x]^4,x]`

```
[Out] ArcTanh[Sin[a + b*x]]/(16*b) + (Sec[a + b*x]*Tan[a + b*x])/(16*b) + (Sec[a + b*x]^3*Tan[a + b*x])/(24*b) - (Sec[a + b*x]^5*Tan[a + b*x])/(6*b) + (Sec[a + b*x]^3*Tan[a + b*x]^3)/(3*b)
```

Maple [A]

time = 0.08, size = 94, normalized size = 1.21

method	result
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{6 \cos(bx+a)^6} + \frac{\sin^5(bx+a)}{24 \cos(bx+a)^4} - \frac{\sin^5(bx+a)}{48 \cos(bx+a)^2} - \frac{(\sin^3(bx+a))}{48} - \frac{\sin(bx+a)}{16} + \frac{\ln(\sec(bx+a)+\tan(bx+a))}{16}}{b}$
default	$\frac{\frac{\sin^5(bx+a)}{6 \cos(bx+a)^6} + \frac{\sin^5(bx+a)}{24 \cos(bx+a)^4} - \frac{\sin^5(bx+a)}{48 \cos(bx+a)^2} - \frac{(\sin^3(bx+a))}{48} - \frac{\sin(bx+a)}{16} + \frac{\ln(\sec(bx+a)+\tan(bx+a))}{16}}{b}$
risch	$-\frac{i(3e^{11i(bx+a)} - 47e^{9i(bx+a)} + 78e^{7i(bx+a)} - 78e^{5i(bx+a)} + 47e^{3i(bx+a)} - 3e^{i(bx+a)})}{24b(e^{2i(bx+a)} + 1)^6} + \frac{\ln(e^{i(bx+a)} + i)}{16b} - \frac{\ln(e^{i(bx+a)} - i)}{16b}$
norman	$-\frac{\tan\left(\frac{bx+a}{2}\right)}{8b} + \frac{17\left(\tan^3\left(\frac{bx+a}{2}\right)\right)}{24b} + \frac{19\left(\tan^5\left(\frac{bx+a}{2}\right)\right)}{4b} + \frac{19\left(\tan^7\left(\frac{bx+a}{2}\right)\right)}{4b} + \frac{17\left(\tan^9\left(\frac{bx+a}{2}\right)\right)}{24b} - \frac{\tan^{11}\left(\frac{bx+a}{2}\right)}{8b} - \frac{\ln\left(\tan^2\left(\frac{bx+a}{2}\right) - 1\right)}{6}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^7*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/6*sin(b*x+a)^5/cos(b*x+a)^6+1/24*sin(b*x+a)^5/cos(b*x+a)^4-1/48*sin(b*x+a)^5/cos(b*x+a)^2-1/48*sin(b*x+a)^3-1/16*sin(b*x+a)+1/16*ln(sec(b*x+a)+tan(b*x+a)))
```

Maxima [A]

time = 0.28, size = 91, normalized size = 1.17

$$\frac{2 \left(3 \sin(bx+a)^5 + 8 \sin(bx+a)^3 - 3 \sin(bx+a) \right)}{\sin(bx+a)^6 - 3 \sin(bx+a)^4 + 3 \sin(bx+a)^2 - 1} - 3 \log(\sin(bx+a) + 1) + 3 \log(\sin(bx+a) - 1)$$

$$96 b$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^4,x, algorithm="maxima")`

```
[Out] -1/96*(2*(3*sin(b*x + a)^5 + 8*sin(b*x + a)^3 - 3*sin(b*x + a))/(sin(b*x + a)^6 - 3*sin(b*x + a)^4 + 3*sin(b*x + a)^2 - 1) - 3*log(sin(b*x + a) + 1) + 3*log(sin(b*x + a) - 1))/b
```

Fricas [A]

time = 0.38, size = 84, normalized size = 1.08

$$\frac{3 \cos(bx+a)^6 \log(\sin(bx+a) + 1) - 3 \cos(bx+a)^6 \log(-\sin(bx+a) + 1) + 2(3 \cos(bx+a)^4 - 14 \cos(bx+a)^2 + 8) \sin(bx+a)}{96 b \cos(bx+a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^4,x, algorithm="fricas")`

```
[Out] 1/96*(3*cos(b*x + a)^6*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^6*log(-sin(b*x + a) + 1) + 2*(3*cos(b*x + a)^4 - 14*cos(b*x + a)^2 + 8)*sin(b*x + a))/(b*cos(b*x + a)^6)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**7*sin(b*x+a)**4,x)`

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 6189 deep
```

Giac [A]

time = 4.21, size = 73, normalized size = 0.94

$$\frac{2 \left(3 \sin(bx+a)^5 + 8 \sin(bx+a)^3 - 3 \sin(bx+a) \right)}{(\sin(bx+a)^2 - 1)^3} - 3 \log(|\sin(bx+a) + 1|) + 3 \log(|\sin(bx+a) - 1|)$$

$$96 b$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^7*sin(b*x+a)^4,x, algorithm="giac")`

[Out] $-1/96*(2*(3*\sin(b*x + a)^5 + 8*\sin(b*x + a)^3 - 3*\sin(b*x + a))/(\sin(b*x + a)^2 - 1)^3 - 3*\log(\text{abs}(\sin(b*x + a) + 1)) + 3*\log(\text{abs}(\sin(b*x + a) - 1)))/b$

Mupad [B]

time = 7.38, size = 177, normalized size = 2.27

$$\frac{\operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b} + \frac{-\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{11}}{8} + \frac{17\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^9}{24} + \frac{19\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^7}{4} + \frac{19\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5}{4} + \frac{17\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3}{24} - \frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{8}}{b\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{12} - 6\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{10} + 15\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 - 20\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + 15\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 - 6\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{int}(\sin(a + b*x)^4/\cos(a + b*x)^7, x)$

[Out] $\operatorname{atanh}(\tan(a/2 + (b*x)/2))/(8*b) + ((17*\tan(a/2 + (b*x)/2)^3)/24 - \tan(a/2 + (b*x)/2)/8 + (19*\tan(a/2 + (b*x)/2)^5)/4 + (19*\tan(a/2 + (b*x)/2)^7)/4 + (17*\tan(a/2 + (b*x)/2)^9)/24 - \tan(a/2 + (b*x)/2)^{11}/8)/(b*(15*\tan(a/2 + (b*x)/2)^4 - 6*\tan(a/2 + (b*x)/2)^2 - 20*\tan(a/2 + (b*x)/2)^6 + 15*\tan(a/2 + (b*x)/2)^8 - 6*\tan(a/2 + (b*x)/2)^{10} + \tan(a/2 + (b*x)/2)^{12} + 1))$

3.97 $\int \sec^5(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=99

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{128b} + \frac{3 \sec(a + bx) \tan(a + bx)}{128b} + \frac{\sec^3(a + bx) \tan(a + bx)}{64b} - \frac{\sec^5(a + bx) \tan(a + bx)}{16b} + \frac{\sec^5(a + bx)}{16b}$$

[Out] 3/128*arctanh(sin(b*x+a))/b+3/128*sec(b*x+a)*tan(b*x+a)/b+1/64*sec(b*x+a)^3*tan(b*x+a)/b-1/16*sec(b*x+a)^5*tan(b*x+a)/b+1/8*sec(b*x+a)^5*tan(b*x+a)^3/b

Rubi [A]

time = 0.06, antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2691, 3853, 3855}

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{128b} + \frac{\tan^3(a + bx) \sec^5(a + bx)}{8b} - \frac{\tan(a + bx) \sec^5(a + bx)}{16b} + \frac{\tan(a + bx) \sec^3(a + bx)}{64b} + \frac{3 \tan(a + bx) \sec(a + bx)}{128b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^5*Tan[a + b*x]^4,x]

[Out] (3*ArcTanh[Sin[a + b*x]])/(128*b) + (3*Sec[a + b*x]*Tan[a + b*x])/(128*b) + (Sec[a + b*x]^3*Tan[a + b*x])/(64*b) - (Sec[a + b*x]^5*Tan[a + b*x])/(16*b) + (Sec[a + b*x]^5*Tan[a + b*x]^3)/(8*b)

Rule 2691

Int[((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_.)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \sec^5(a+bx) \tan^4(a+bx) dx &= \frac{\sec^5(a+bx) \tan^3(a+bx)}{8b} - \frac{3}{8} \int \sec^5(a+bx) \tan^2(a+bx) dx \\
&= -\frac{\sec^5(a+bx) \tan(a+bx)}{16b} + \frac{\sec^5(a+bx) \tan^3(a+bx)}{8b} + \frac{1}{16} \int \sec^5(a+bx) \tan^2(a+bx) dx \\
&= \frac{\sec^3(a+bx) \tan(a+bx)}{64b} - \frac{\sec^5(a+bx) \tan(a+bx)}{16b} + \frac{\sec^5(a+bx) \tan^3(a+bx)}{8b} \\
&= \frac{3 \sec(a+bx) \tan(a+bx)}{128b} + \frac{\sec^3(a+bx) \tan(a+bx)}{64b} - \frac{\sec^5(a+bx) \tan(a+bx)}{16b} \\
&= \frac{3 \tanh^{-1}(\sin(a+bx))}{128b} + \frac{3 \sec(a+bx) \tan(a+bx)}{128b} + \frac{\sec^3(a+bx) \tan(a+bx)}{64b}
\end{aligned}$$

Mathematica [A]

time = 0.22, size = 64, normalized size = 0.65

$$\frac{96 \tanh^{-1}(\sin(a+bx)) + (182 - 307 \cos(2(a+bx)) + 26 \cos(4(a+bx)) + 3 \cos(6(a+bx))) \sec^7(a+bx) \tan(a+bx)}{4096b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^5*Tan[a + b*x]^4,x]`

```
[Out] (96*ArcTanh[Sin[a + b*x]] + (182 - 307*Cos[2*(a + b*x)] + 26*Cos[4*(a + b*x)] + 3*Cos[6*(a + b*x)])*Sec[a + b*x]^7*Tan[a + b*x])/(4096*b)
```

Maple [A]

time = 0.11, size = 112, normalized size = 1.13

method	result
derivativedivides	$\frac{\frac{\sin^5(bx+a)}{8 \cos(bx+a)^8} + \frac{\sin^5(bx+a)}{16 \cos(bx+a)^6} + \frac{\sin^5(bx+a)}{64 \cos(bx+a)^4} - \frac{\sin^5(bx+a)}{128 \cos(bx+a)^2} - \frac{(\sin^3(bx+a))}{128} - \frac{3 \sin(bx+a)}{128} + \frac{3 \ln(\sec(bx+a) + \tan(bx+a))}{128}}{b}$
default	$\frac{\frac{\sin^5(bx+a)}{8 \cos(bx+a)^8} + \frac{\sin^5(bx+a)}{16 \cos(bx+a)^6} + \frac{\sin^5(bx+a)}{64 \cos(bx+a)^4} - \frac{\sin^5(bx+a)}{128 \cos(bx+a)^2} - \frac{(\sin^3(bx+a))}{128} - \frac{3 \sin(bx+a)}{128} + \frac{3 \ln(\sec(bx+a) + \tan(bx+a))}{128}}{b}$
risch	$-\frac{i(3e^{15i(bx+a)} + 23e^{13i(bx+a)} - 333e^{11i(bx+a)} + 671e^{9i(bx+a)} - 671e^{7i(bx+a)} + 333e^{5i(bx+a)} - 23e^{3i(bx+a)} - 3e^{i(bx+a)})}{64b(e^{2i(bx+a)} + 1)^8}$
norman	$-\frac{\frac{3 \tan\left(\frac{bx+a}{2}\right)}{64b} + \frac{23(\tan^3\left(\frac{bx+a}{2}\right))}{64b} + \frac{333(\tan^5\left(\frac{bx+a}{2}\right))}{64b} + \frac{671(\tan^7\left(\frac{bx+a}{2}\right))}{64b} + \frac{671(\tan^9\left(\frac{bx+a}{2}\right))}{64b} + \frac{333(\tan^{11}\left(\frac{bx+a}{2}\right))}{64b}}{(\tan^2\left(\frac{bx+a}{2}\right) - 1)^8}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^9*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/8*sin(b*x+a)^5/cos(b*x+a)^8+1/16*sin(b*x+a)^5/cos(b*x+a)^6+1/64*sin(b*x+a)^5/cos(b*x+a)^4-1/128*sin(b*x+a)^5/cos(b*x+a)^2-1/128*sin(b*x+a)^3-3/128*sin(b*x+a)+3/128*ln(sec(b*x+a)+tan(b*x+a)))
```

Maxima [A]

time = 0.29, size = 111, normalized size = 1.12

$$\frac{2 \left(3 \sin(bx+a)^7 - 11 \sin(bx+a)^5 - 11 \sin(bx+a)^3 + 3 \sin(bx+a) \right)}{\sin(bx+a)^8 - 4 \sin(bx+a)^6 + 6 \sin(bx+a)^4 - 4 \sin(bx+a)^2 + 1} - 3 \log(\sin(bx+a) + 1) + 3 \log(\sin(bx+a) - 1)$$

$$256b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^9*sin(b*x+a)^4,x, algorithm="maxima")

[Out] -1/256*(2*(3*sin(b*x + a)^7 - 11*sin(b*x + a)^5 - 11*sin(b*x + a)^3 + 3*sin(b*x + a))/(sin(b*x + a)^8 - 4*sin(b*x + a)^6 + 6*sin(b*x + a)^4 - 4*sin(b*x + a)^2 + 1) - 3*log(sin(b*x + a) + 1) + 3*log(sin(b*x + a) - 1))/b

Fricas [A]

time = 0.39, size = 94, normalized size = 0.95

$$\frac{3 \cos(bx+a)^8 \log(\sin(bx+a) + 1) - 3 \cos(bx+a)^8 \log(-\sin(bx+a) + 1) + 2(3 \cos(bx+a)^6 + 2 \cos(bx+a)^4 - 24 \cos(bx+a)^2 + 16) \sin(bx+a)}{256b \cos(bx+a)^8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^9*sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/256*(3*cos(b*x + a)^8*log(sin(b*x + a) + 1) - 3*cos(b*x + a)^8*log(-sin(b*x + a) + 1) + 2*(3*cos(b*x + a)^6 + 2*cos(b*x + a)^4 - 24*cos(b*x + a)^2 + 16)*sin(b*x + a))/(b*cos(b*x + a)^8)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**9*sin(b*x+a)**4,x)**[Out]** Timed out**Giac [A]**

time = 4.52, size = 107, normalized size = 1.08

$$\frac{4 \left(3 \left(\frac{1}{\sin(bx+a)} + \sin(bx+a) \right)^3 - \frac{20}{\sin(bx+a)} - 20 \sin(bx+a) \right)}{\left(\left(\frac{1}{\sin(bx+a)} + \sin(bx+a) \right)^2 - 4 \right)^2} - 3 \log \left(\left| \frac{1}{\sin(bx+a)} + \sin(bx+a) + 2 \right| \right) + 3 \log \left(\left| \frac{1}{\sin(bx+a)} + \sin(bx+a) - 2 \right| \right)$$

$$512b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^9*sin(b*x+a)^4,x, algorithm="giac")

[Out] $-1/512*(4*(3*(1/\sin(b*x + a) + \sin(b*x + a))^3 - 20/\sin(b*x + a) - 20*\sin(b*x + a))/((1/\sin(b*x + a) + \sin(b*x + a))^2 - 4)^2 - 3*\log(\text{abs}(1/\sin(b*x + a) + \sin(b*x + a) + 2)) + 3*\log(\text{abs}(1/\sin(b*x + a) + \sin(b*x + a) - 2)))/b$

Mupad [B]

time = 7.44, size = 229, normalized size = 2.31

$$\frac{3 \operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{64b} + \frac{-\frac{3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{15}}{64} + \frac{23 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{13}}{64} + \frac{333 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{11}}{64} + \frac{671 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^9}{64} + \frac{671 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^7}{64} + \frac{333 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5}{64} + \frac{23 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3}{64} - \frac{3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{64}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{16} - 8 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{14} + 28 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{12} - 56 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{10} + 70 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 - 56 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + 28 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 - 8 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(\sin(a + b*x)^4/\cos(a + b*x)^9, x)$

[Out] $(3*\operatorname{atanh}(\tan(a/2 + (b*x)/2)))/(64*b) + ((23*\tan(a/2 + (b*x)/2)^3)/64 - (3*\tan(a/2 + (b*x)/2))/64 + (333*\tan(a/2 + (b*x)/2)^5)/64 + (671*\tan(a/2 + (b*x)/2)^7)/64 + (671*\tan(a/2 + (b*x)/2)^9)/64 + (333*\tan(a/2 + (b*x)/2)^{11})/64 + (23*\tan(a/2 + (b*x)/2)^{13})/64 - (3*\tan(a/2 + (b*x)/2)^{15})/64)/(b*(28*\tan(a/2 + (b*x)/2)^4 - 8*\tan(a/2 + (b*x)/2)^2 - 56*\tan(a/2 + (b*x)/2)^6 + 70*\tan(a/2 + (b*x)/2)^8 - 56*\tan(a/2 + (b*x)/2)^{10} + 28*\tan(a/2 + (b*x)/2)^{12} - 8*\tan(a/2 + (b*x)/2)^{14} + \tan(a/2 + (b*x)/2)^{16} + 1))$

3.98 $\int \cos^7(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=46

$$-\frac{\cos^8(a + bx)}{8b} + \frac{\cos^{10}(a + bx)}{5b} - \frac{\cos^{12}(a + bx)}{12b}$$

[Out] $-1/8*\cos(b*x+a)^8/b+1/5*\cos(b*x+a)^{10}/b-1/12*\cos(b*x+a)^{12}/b$

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2645, 272, 45}

$$-\frac{\cos^{12}(a + bx)}{12b} + \frac{\cos^{10}(a + bx)}{5b} - \frac{\cos^8(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^7*Sin[a + b*x]^5,x]

[Out] $-1/8*\text{Cos}[a + b*x]^8/b + \text{Cos}[a + b*x]^10/(5*b) - \text{Cos}[a + b*x]^12/(12*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2645

Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned}
\int \cos^7(a+bx) \sin^5(a+bx) dx &= -\frac{\text{Subst}\left(\int x^7(1-x^2)^2 dx, x, \cos(a+bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int (1-x)^2 x^3 dx, x, \cos^2(a+bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int (x^3 - 2x^4 + x^5) dx, x, \cos^2(a+bx)\right)}{2b} \\
&= -\frac{\cos^8(a+bx)}{8b} + \frac{\cos^{10}(a+bx)}{5b} - \frac{\cos^{12}(a+bx)}{12b}
\end{aligned}$$

Mathematica [A]

time = 0.27, size = 68, normalized size = 1.48

$$-\frac{600 \cos(2(a+bx)) + 75 \cos(4(a+bx)) - 100 \cos(6(a+bx)) - 30 \cos(8(a+bx)) + 12 \cos(10(a+bx)) + 5 \cos(12(a+bx))}{122880b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^7*Sin[a + b*x]^5,x]`

```
[Out] -1/122880*(600*Cos[2*(a + b*x)] + 75*Cos[4*(a + b*x)] - 100*Cos[6*(a + b*x)]
- 30*Cos[8*(a + b*x)] + 12*Cos[10*(a + b*x)] + 5*Cos[12*(a + b*x)]/b
```

Maple [A]

time = 0.19, size = 52, normalized size = 1.13

method	result	size
derivativedivides	$-\frac{\frac{(\sin^4(bx+a))(\cos^8(bx+a))}{12} - \frac{(\sin^2(bx+a))(\cos^8(bx+a))}{30} - \frac{(\cos^8(bx+a))}{120}}{b}$	52
default	$-\frac{\frac{(\sin^4(bx+a))(\cos^8(bx+a))}{12} - \frac{(\sin^2(bx+a))(\cos^8(bx+a))}{30} - \frac{(\cos^8(bx+a))}{120}}{b}$	52
risch	$-\frac{\cos(12bx+12a)}{24576b} - \frac{\cos(10bx+10a)}{10240b} + \frac{\cos(8bx+8a)}{4096b} + \frac{5 \cos(6bx+6a)}{6144b} - \frac{5 \cos(4bx+4a)}{8192b} - \frac{5 \cos(2bx+2a)}{1024b}$	86

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^7*sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/12*sin(b*x+a)^4*cos(b*x+a)^8-1/30*sin(b*x+a)^2*cos(b*x+a)^8-1/120*cos(b*x+a)^8)
```

Maxima [A]

time = 0.29, size = 46, normalized size = 1.00

$$-\frac{10 \sin(bx+a)^{12} - 36 \sin(bx+a)^{10} + 45 \sin(bx+a)^8 - 20 \sin(bx+a)^6}{120b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7*sin(b*x+a)^5,x, algorithm="maxima")

[Out] $-1/120*(10*\sin(b*x + a)^{12} - 36*\sin(b*x + a)^{10} + 45*\sin(b*x + a)^8 - 20*\sin(b*x + a)^6)/b$

Fricas [A]

time = 0.37, size = 36, normalized size = 0.78

$$-\frac{10 \cos (bx+a)^{12}-24 \cos (bx+a)^{10}+15 \cos (bx+a)^8}{120 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7*sin(b*x+a)^5,x, algorithm="fricas")

[Out] $-1/120*(10*\cos(b*x + a)^{12} - 24*\cos(b*x + a)^{10} + 15*\cos(b*x + a)^8)/b$

Sympy [A]

time = 3.37, size = 65, normalized size = 1.41

$$\begin{cases} -\frac{\sin^4(a+bx)\cos^8(a+bx)}{8b} - \frac{\sin^2(a+bx)\cos^{10}(a+bx)}{20b} - \frac{\cos^{12}(a+bx)}{120b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos^7(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**7*sin(b*x+a)**5,x)

[Out] Piecewise((-sin(a + b*x)**4*cos(a + b*x)**8/(8*b) - sin(a + b*x)**2*cos(a + b*x)**10/(20*b) - cos(a + b*x)**12/(120*b), Ne(b, 0)), (x*sin(a)**5*cos(a)**7, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 85 vs. $2(40) = 80$.
time = 4.23, size = 85, normalized size = 1.85

$$-\frac{\cos(12bx+12a)}{24576b} - \frac{\cos(10bx+10a)}{10240b} + \frac{\cos(8bx+8a)}{4096b} + \frac{5\cos(6bx+6a)}{6144b} - \frac{5\cos(4bx+4a)}{8192b} - \frac{5\cos(2bx+2a)}{1024b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7*sin(b*x+a)^5,x, algorithm="giac")

[Out] $-1/24576*\cos(12*b*x + 12*a)/b - 1/10240*\cos(10*b*x + 10*a)/b + 1/4096*\cos(8*b*x + 8*a)/b + 5/6144*\cos(6*b*x + 6*a)/b - 5/8192*\cos(4*b*x + 4*a)/b - 5/1024*\cos(2*b*x + 2*a)/b$

Mupad [B]

time = 0.41, size = 35, normalized size = 0.76

$$\frac{\cos(a+bx)^8(10\cos(a+bx)^4 - 24\cos(a+bx)^2 + 15)}{120b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^7*sin(a + b*x)^5,x)
```

```
[Out] -(cos(a + b*x)^8*(10*cos(a + b*x)^4 - 24*cos(a + b*x)^2 + 15))/(120*b)
```

3.99 $\int \cos^6(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=46

$$-\frac{\cos^7(a + bx)}{7b} + \frac{2 \cos^9(a + bx)}{9b} - \frac{\cos^{11}(a + bx)}{11b}$$

[Out] $-1/7*\cos(b*x+a)^7/b+2/9*\cos(b*x+a)^9/b-1/11*\cos(b*x+a)^{11}/b$

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 276}

$$-\frac{\cos^{11}(a + bx)}{11b} + \frac{2 \cos^9(a + bx)}{9b} - \frac{\cos^7(a + bx)}{7b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^6*Sin[a + b*x]^5,x]`

[Out] $-1/7*\text{Cos}[a + b*x]^7/b + (2*\text{Cos}[a + b*x]^9)/(9*b) - \text{Cos}[a + b*x]^{11}/(11*b)$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2645

`Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

Rubi steps

$$\begin{aligned} \int \cos^6(a + bx) \sin^5(a + bx) dx &= -\frac{\text{Subst}\left(\int x^6(1 - x^2)^2 dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int (x^6 - 2x^8 + x^{10}) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos^7(a + bx)}{7b} + \frac{2 \cos^9(a + bx)}{9b} - \frac{\cos^{11}(a + bx)}{11b} \end{aligned}$$

Mathematica [A]

time = 0.20, size = 37, normalized size = 0.80

$$\frac{\cos^7(a + bx)(-365 + 364 \cos(2(a + bx)) - 63 \cos(4(a + bx)))}{5544b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^6*Sin[a + b*x]^5,x]

[Out] (Cos[a + b*x]^7*(-365 + 364*Cos[2*(a + b*x)] - 63*Cos[4*(a + b*x)]))/(5544*b)

Maple [A]

time = 0.12, size = 52, normalized size = 1.13

method	result	size
derivativedivides	$\frac{\frac{(\cos^7(bx+a))(\sin^4(bx+a))}{11} - \frac{4(\cos^7(bx+a))(\sin^2(bx+a))}{99} - \frac{8(\cos^7(bx+a))}{693}}{b}$	52
default	$\frac{\frac{(\cos^7(bx+a))(\sin^4(bx+a))}{11} - \frac{4(\cos^7(bx+a))(\sin^2(bx+a))}{99} - \frac{8(\cos^7(bx+a))}{693}}{b}$	52
risch	$-\frac{5 \cos(bx+a)}{512b} - \frac{\cos(11bx+11a)}{11264b} - \frac{\cos(9bx+9a)}{9216b} + \frac{5 \cos(7bx+7a)}{7168b} + \frac{\cos(5bx+5a)}{1024b} - \frac{5 \cos(3bx+3a)}{1536b}$	83

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^6*sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/11*cos(b*x+a)^7*sin(b*x+a)^4-4/99*cos(b*x+a)^7*sin(b*x+a)^2-8/693*cos(b*x+a)^7)

Maxima [A]

time = 0.28, size = 36, normalized size = 0.78

$$-\frac{63 \cos(bx + a)^{11} - 154 \cos(bx + a)^9 + 99 \cos(bx + a)^7}{693b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/693*(63*cos(b*x + a)^11 - 154*cos(b*x + a)^9 + 99*cos(b*x + a)^7)/b

Fricas [A]

time = 0.38, size = 36, normalized size = 0.78

$$-\frac{63 \cos(bx + a)^{11} - 154 \cos(bx + a)^9 + 99 \cos(bx + a)^7}{693b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^5,x, algorithm="fricas")

[Out] $-1/693*(63*\cos(b*x + a)^{11} - 154*\cos(b*x + a)^9 + 99*\cos(b*x + a)^7)/b$

Sympy [A]

time = 2.47, size = 68, normalized size = 1.48

$$\begin{cases} -\frac{\sin^4(a+bx)\cos^7(a+bx)}{7b} - \frac{4\sin^2(a+bx)\cos^9(a+bx)}{63b} - \frac{8\cos^{11}(a+bx)}{693b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos^6(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**6*sin(b*x+a)**5,x)

[Out] Piecewise((-sin(a + b*x)**4*cos(a + b*x)**7/(7*b) - 4*sin(a + b*x)**2*cos(a + b*x)**9/(63*b) - 8*cos(a + b*x)**11/(693*b), Ne(b, 0)), (x*sin(a)**5*cos(a)**6, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(40) = 80.

time = 2.97, size = 82, normalized size = 1.78

$$-\frac{\cos(11bx + 11a)}{11264b} - \frac{\cos(9bx + 9a)}{9216b} + \frac{5\cos(7bx + 7a)}{7168b} + \frac{\cos(5bx + 5a)}{1024b} - \frac{5\cos(3bx + 3a)}{1536b} - \frac{5\cos(bx + a)}{512b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6*sin(b*x+a)^5,x, algorithm="giac")

[Out] $-1/11264*\cos(11*b*x + 11*a)/b - 1/9216*\cos(9*b*x + 9*a)/b + 5/7168*\cos(7*b*x + 7*a)/b + 1/1024*\cos(5*b*x + 5*a)/b - 5/1536*\cos(3*b*x + 3*a)/b - 5/512*\cos(b*x + a)/b$

Mupad [B]

time = 0.41, size = 36, normalized size = 0.78

$$-\frac{63\cos(a+bx)^{11} - 154\cos(a+bx)^9 + 99\cos(a+bx)^7}{693b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6*sin(a + b*x)^5,x)

[Out] $-(99*\cos(a + b*x)^7 - 154*\cos(a + b*x)^9 + 63*\cos(a + b*x)^{11})/(693*b)$

3.100 $\int \cos^5(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sin^6(a + bx)}{6b} - \frac{\sin^8(a + bx)}{4b} + \frac{\sin^{10}(a + bx)}{10b}$$

[Out] 1/6*sin(b*x+a)^6/b-1/4*sin(b*x+a)^8/b+1/10*sin(b*x+a)^10/b

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2644, 272, 45}

$$\frac{\sin^{10}(a + bx)}{10b} - \frac{\sin^8(a + bx)}{4b} + \frac{\sin^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^5*Sin[a + b*x]^5,x]

[Out] Sin[a + b*x]^6/(6*b) - Sin[a + b*x]^8/(4*b) + Sin[a + b*x]^10/(10*b)

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
\int \cos^5(a+bx) \sin^5(a+bx) dx &= \frac{\text{Subst}\left(\int x^5(1-x^2)^2 dx, x, \sin(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int (1-x)^2 x^2 dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int (x^2 - 2x^3 + x^4) dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{\sin^6(a+bx)}{6b} - \frac{\sin^8(a+bx)}{4b} + \frac{\sin^{10}(a+bx)}{10b}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 50, normalized size = 1.09

$$\frac{1}{32} \left(-\frac{5 \cos(2(a+bx))}{16b} + \frac{5 \cos(6(a+bx))}{96b} - \frac{\cos(10(a+bx))}{160b} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^5*Sin[a + b*x]^5,x]``[Out] ((-5*Cos[2*(a + b*x)])/(16*b) + (5*Cos[6*(a + b*x)])/(96*b) - Cos[10*(a + b*x)]/(160*b))/32`**Maple [A]**

time = 0.11, size = 52, normalized size = 1.13

method	result	size
risch	$-\frac{\cos(10bx+10a)}{5120b} + \frac{5 \cos(6bx+6a)}{3072b} - \frac{5 \cos(2bx+2a)}{512b}$	44
derivativedivides	$-\frac{(\cos^6(bx+a))(\sin^4(bx+a))}{10} - \frac{(\cos^6(bx+a))(\sin^2(bx+a))}{20} - \frac{(\cos^6(bx+a))}{60}$	52
default	$-\frac{(\cos^6(bx+a))(\sin^4(bx+a))}{10} - \frac{(\cos^6(bx+a))(\sin^2(bx+a))}{20} - \frac{(\cos^6(bx+a))}{60}$	52

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/10*cos(b*x+a)^6*sin(b*x+a)^4-1/20*cos(b*x+a)^6*sin(b*x+a)^2-1/60*cos(b*x+a)^6)`**Maxima [A]**

time = 0.30, size = 36, normalized size = 0.78

$$\frac{6 \sin(bx+a)^{10} - 15 \sin(bx+a)^8 + 10 \sin(bx+a)^6}{60b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^5,x, algorithm="maxima")

[Out] 1/60*(6*sin(b*x + a)^10 - 15*sin(b*x + a)^8 + 10*sin(b*x + a)^6)/b

Fricas [A]

time = 0.36, size = 36, normalized size = 0.78

$$-\frac{6 \cos (b x+a)^{10}-15 \cos (b x+a)^{8}+10 \cos (b x+a)^{6}}{60 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^5,x, algorithm="fricas")

[Out] -1/60*(6*cos(b*x + a)^10 - 15*cos(b*x + a)^8 + 10*cos(b*x + a)^6)/b

Sympy [A]

time = 2.82, size = 65, normalized size = 1.41

$$\begin{cases} -\frac{\sin^4(a+bx)\cos^6(a+bx)}{6b} - \frac{\sin^2(a+bx)\cos^8(a+bx)}{12b} - \frac{\cos^{10}(a+bx)}{60b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos^5(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**5*sin(b*x+a)**5,x)

[Out] Piecewise((-sin(a + b*x)**4*cos(a + b*x)**6/(6*b) - sin(a + b*x)**2*cos(a + b*x)**8/(12*b) - cos(a + b*x)**10/(60*b), Ne(b, 0)), (x*sin(a)**5*cos(a)**5, True))

Giac [A]

time = 3.72, size = 43, normalized size = 0.93

$$-\frac{\cos (10 b x+10 a)}{5120 b}+\frac{5 \cos (6 b x+6 a)}{3072 b}-\frac{5 \cos (2 b x+2 a)}{512 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*sin(b*x+a)^5,x, algorithm="giac")

[Out] -1/5120*cos(10*b*x + 10*a)/b + 5/3072*cos(6*b*x + 6*a)/b - 5/512*cos(2*b*x + 2*a)/b

Mupad [B]

time = 0.48, size = 36, normalized size = 0.78

$$-\frac{\frac{\cos (a+b x)^{10}}{10}-\frac{\cos (a+b x)^{8}}{4}+\frac{\cos (a+b x)^{6}}{6}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5*sin(a + b*x)^5,x)

[Out] -(cos(a + b*x)^6/6 - cos(a + b*x)^8/4 + cos(a + b*x)^10/10)/b

3.101 $\int \cos^4(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=46

$$-\frac{\cos^5(a + bx)}{5b} + \frac{2 \cos^7(a + bx)}{7b} - \frac{\cos^9(a + bx)}{9b}$$

[Out] $-1/5*\cos(b*x+a)^5/b+2/7*\cos(b*x+a)^7/b-1/9*\cos(b*x+a)^9/b$

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 276}

$$-\frac{\cos^9(a + bx)}{9b} + \frac{2 \cos^7(a + bx)}{7b} - \frac{\cos^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^4*\text{Sin}[a + b*x]^5, x]$

[Out] $-1/5*\text{Cos}[a + b*x]^5/b + (2*\text{Cos}[a + b*x]^7)/(7*b) - \text{Cos}[a + b*x]^9/(9*b)$

Rule 276

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{Exp andIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0]$

Rule 2645

$\text{Int}[(\cos[(e_*) + (f_*)(x_)]*(a_*)^{(m_*)}*\sin[(e_*) + (f_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{(n-1)/2}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n-1)/2] \&\& !(\text{IntegerQ}[(m-1)/2] \&\& \text{GtQ}[m, 0] \&\& \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int \cos^4(a + bx) \sin^5(a + bx) dx &= -\frac{\text{Subst}\left(\int x^4(1 - x^2)^2 dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int (x^4 - 2x^6 + x^8) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos^5(a + bx)}{5b} + \frac{2 \cos^7(a + bx)}{7b} - \frac{\cos^9(a + bx)}{9b} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 37, normalized size = 0.80

$$\frac{\cos^5(a + bx)(-249 + 220 \cos(2(a + bx)) - 35 \cos(4(a + bx)))}{2520b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^4*Sin[a + b*x]^5,x]`

`[Out] (Cos[a + b*x]^5*(-249 + 220*Cos[2*(a + b*x)] - 35*Cos[4*(a + b*x)]))/(2520*b)`

Maple [A]

time = 0.12, size = 52, normalized size = 1.13

method	result
derivativdivides	$-\frac{(\cos^5(bx+a))(\sin^4(bx+a))}{9} - \frac{4(\cos^5(bx+a))(\sin^2(bx+a))}{63} - \frac{8(\cos^5(bx+a))}{315}$
default	$-\frac{(\cos^5(bx+a))(\sin^4(bx+a))}{9} - \frac{4(\cos^5(bx+a))(\sin^2(bx+a))}{63} - \frac{8(\cos^5(bx+a))}{315}$
risch	$-\frac{3 \cos(bx+a)}{128b} - \frac{\cos(9bx+9a)}{2304b} + \frac{\cos(7bx+7a)}{1792b} + \frac{\cos(5bx+5a)}{320b} - \frac{\cos(3bx+3a)}{192b}$
norman	$\frac{-\frac{16}{315b} - \frac{112(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{32(\tan^{12}(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{16(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{32(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{16(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{35b} - \frac{64(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{35b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^9}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^4*sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

`[Out] 1/b*(-1/9*cos(b*x+a)^5*sin(b*x+a)^4-4/63*cos(b*x+a)^5*sin(b*x+a)^2-8/315*cos(b*x+a)^5)`

Maxima [A]

time = 0.28, size = 36, normalized size = 0.78

$$-\frac{35 \cos(bx + a)^9 - 90 \cos(bx + a)^7 + 63 \cos(bx + a)^5}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^4*sin(b*x+a)^5,x, algorithm="maxima")`

`[Out] -1/315*(35*cos(b*x + a)^9 - 90*cos(b*x + a)^7 + 63*cos(b*x + a)^5)/b`

Fricas [A]

time = 0.37, size = 36, normalized size = 0.78

$$-\frac{35 \cos(bx + a)^9 - 90 \cos(bx + a)^7 + 63 \cos(bx + a)^5}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4*sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $-1/315*(35*\cos(b*x + a)^9 - 90*\cos(b*x + a)^7 + 63*\cos(b*x + a)^5)/b$

Sympy [A]

time = 1.25, size = 68, normalized size = 1.48

$$\begin{cases} -\frac{\sin^4(a+bx)\cos^5(a+bx)}{5b} - \frac{4\sin^2(a+bx)\cos^7(a+bx)}{35b} - \frac{8\cos^9(a+bx)}{315b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos^4(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**4*sin(b*x+a)**5,x)`

[Out] `Piecewise((-sin(a + b*x)**4*cos(a + b*x)**5/(5*b) - 4*sin(a + b*x)**2*cos(a + b*x)**7/(35*b) - 8*cos(a + b*x)**9/(315*b), Ne(b, 0)), (x*sin(a)**5*cos(a)**4, True))`

Giac [A]

time = 2.78, size = 68, normalized size = 1.48

$$-\frac{\cos(9bx + 9a)}{2304b} + \frac{\cos(7bx + 7a)}{1792b} + \frac{\cos(5bx + 5a)}{320b} - \frac{\cos(3bx + 3a)}{192b} - \frac{3\cos(bx + a)}{128b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $-1/2304*\cos(9*b*x + 9*a)/b + 1/1792*\cos(7*b*x + 7*a)/b + 1/320*\cos(5*b*x + 5*a)/b - 1/192*\cos(3*b*x + 3*a)/b - 3/128*\cos(b*x + a)/b$

Mupad [B]

time = 0.38, size = 36, normalized size = 0.78

$$-\frac{35\cos(a+bx)^9 - 90\cos(a+bx)^7 + 63\cos(a+bx)^5}{315b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^4*sin(a + b*x)^5,x)`

[Out] $-(63*\cos(a + b*x)^5 - 90*\cos(a + b*x)^7 + 35*\cos(a + b*x)^9)/(315*b)$

3.102 $\int \cos^3(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\sin^6(a + bx)}{6b} - \frac{\sin^8(a + bx)}{8b}$$

[Out] 1/6*sin(b*x+a)^6/b-1/8*sin(b*x+a)^8/b

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 14}

$$\frac{\sin^6(a + bx)}{6b} - \frac{\sin^8(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*Sin[a + b*x]^5,x]

[Out] Sin[a + b*x]^6/(6*b) - Sin[a + b*x]^8/(8*b)

Rule 14

```
Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2644

```
Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \sin^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^5(1 - x^2) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^5 - x^7) dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^6(a + bx)}{6b} - \frac{\sin^8(a + bx)}{8b} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 48, normalized size = 1.55

$$\frac{-72 \cos(2(a + bx)) + 12 \cos(4(a + bx)) + 8 \cos(6(a + bx)) - 3 \cos(8(a + bx))}{3072b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*Sin[a + b*x]^5,x]``[Out] (-72*Cos[2*(a + b*x)] + 12*Cos[4*(a + b*x)] + 8*Cos[6*(a + b*x)] - 3*Cos[8*(a + b*x)])/(3072*b)`**Maple [A]**

time = 0.10, size = 52, normalized size = 1.68

method	result	size
derivativedivides	$\frac{-\frac{(\cos^4(bx+a))(\sin^4(bx+a))}{8} - \frac{(\cos^4(bx+a))(\sin^2(bx+a))}{12} - \frac{(\cos^4(bx+a))}{24}}{b}$	52
default	$\frac{-\frac{(\cos^4(bx+a))(\sin^4(bx+a))}{8} - \frac{(\cos^4(bx+a))(\sin^2(bx+a))}{12} - \frac{(\cos^4(bx+a))}{24}}{b}$	52
risch	$-\frac{\cos(8bx+8a)}{1024b} + \frac{\cos(6bx+6a)}{384b} + \frac{\cos(4bx+4a)}{256b} - \frac{3 \cos(2bx+2a)}{128b}$	58
norman	$\frac{\frac{32(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{32(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{32(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^8}$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^3*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/8*cos(b*x+a)^4*sin(b*x+a)^4-1/12*cos(b*x+a)^4*sin(b*x+a)^2-1/24*cos(b*x+a)^4)`**Maxima [A]**

time = 0.28, size = 26, normalized size = 0.84

$$\frac{3 \sin(bx + a)^8 - 4 \sin(bx + a)^6}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^3*sin(b*x+a)^5,x, algorithm="maxima")``[Out] -1/24*(3*sin(b*x + a)^8 - 4*sin(b*x + a)^6)/b`**Fricas [A]**

time = 0.43, size = 36, normalized size = 1.16

$$\frac{3 \cos(bx + a)^8 - 8 \cos(bx + a)^6 + 6 \cos(bx + a)^4}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3*sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $-1/24*(3*\cos(b*x + a)^8 - 8*\cos(b*x + a)^6 + 6*\cos(b*x + a)^4)/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(22) = 44$.

time = 0.85, size = 65, normalized size = 2.10

$$\begin{cases} -\frac{\sin^4(a+bx)\cos^4(a+bx)}{4b} - \frac{\sin^2(a+bx)\cos^6(a+bx)}{6b} - \frac{\cos^8(a+bx)}{24b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**3*sin(b*x+a)**5,x)`

[Out] `Piecewise((-sin(a + b*x)**4*cos(a + b*x)**4/(4*b) - sin(a + b*x)**2*cos(a + b*x)**6/(6*b) - cos(a + b*x)**8/(24*b), Ne(b, 0)), (x*sin(a)**5*cos(a)**3, True))`

Giac [A]

time = 3.43, size = 26, normalized size = 0.84

$$-\frac{3 \sin (bx + a)^8 - 4 \sin (bx + a)^6}{24 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $-1/24*(3*\sin(b*x + a)^8 - 4*\sin(b*x + a)^6)/b$

Mupad [B]

time = 0.38, size = 26, normalized size = 0.84

$$\frac{4 \sin (a + bx)^6 - 3 \sin (a + bx)^8}{24 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^3*sin(a + b*x)^5,x)`

[Out] $(4*\sin(a + b*x)^6 - 3*\sin(a + b*x)^8)/(24*b)$

3.103 $\int \cos^2(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=46

$$-\frac{\cos^3(a + bx)}{3b} + \frac{2 \cos^5(a + bx)}{5b} - \frac{\cos^7(a + bx)}{7b}$$

[Out] $-1/3*\cos(b*x+a)^3/b+2/5*\cos(b*x+a)^5/b-1/7*\cos(b*x+a)^7/b$

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 276}

$$-\frac{\cos^7(a + bx)}{7b} + \frac{2 \cos^5(a + bx)}{5b} - \frac{\cos^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^2*Sin[a + b*x]^5,x]`

[Out] $-1/3*\text{Cos}[a + b*x]^3/b + (2*\text{Cos}[a + b*x]^5)/(5*b) - \text{Cos}[a + b*x]^7/(7*b)$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2645

`Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) \sin^5(a + bx) dx &= -\frac{\text{Subst}\left(\int x^2(1 - x^2)^2 dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int (x^2 - 2x^4 + x^6) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos^3(a + bx)}{3b} + \frac{2 \cos^5(a + bx)}{5b} - \frac{\cos^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 37, normalized size = 0.80

$$\frac{\cos^3(a + bx)(-157 + 108 \cos(2(a + bx)) - 15 \cos(4(a + bx)))}{840b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^2*Sin[a + b*x]^5,x]`

```
[Out] (Cos[a + b*x]^3*(-157 + 108*Cos[2*(a + b*x)] - 15*Cos[4*(a + b*x)])/(840*b)
```

Maple [A]

time = 0.07, size = 52, normalized size = 1.13

method	result	size
derivativedivides	$\frac{-\frac{(\cos^3(bx+a))(\sin^4(bx+a))}{7} - \frac{4(\cos^3(bx+a))(\sin^2(bx+a))}{35} - \frac{8(\cos^3(bx+a))}{105}}{b}$	52
default	$\frac{-\frac{(\cos^3(bx+a))(\sin^4(bx+a))}{7} - \frac{4(\cos^3(bx+a))(\sin^2(bx+a))}{35} - \frac{8(\cos^3(bx+a))}{105}}{b}$	52
risch	$-\frac{5 \cos(bx+a)}{64b} - \frac{\cos(7bx+7a)}{448b} + \frac{3 \cos(5bx+5a)}{320b} - \frac{\cos(3bx+3a)}{192b}$	55
norman	$\frac{-\frac{16}{105b} - \frac{32(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{16(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{15b} - \frac{16(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{5b} + \frac{16(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^7}$	87

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^2*sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/7*cos(b*x+a)^3*sin(b*x+a)^4-4/35*cos(b*x+a)^3*sin(b*x+a)^2-8/105*cos(b*x+a)^3)
```

Maxima [A]

time = 0.28, size = 36, normalized size = 0.78

$$\frac{15 \cos(bx + a)^7 - 42 \cos(bx + a)^5 + 35 \cos(bx + a)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^2*sin(b*x+a)^5,x, algorithm="maxima")`

```
[Out] -1/105*(15*cos(b*x + a)^7 - 42*cos(b*x + a)^5 + 35*cos(b*x + a)^3)/b
```

Fricas [A]

time = 0.37, size = 36, normalized size = 0.78

$$\frac{15 \cos(bx + a)^7 - 42 \cos(bx + a)^5 + 35 \cos(bx + a)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^5,x, algorithm="fricas")

[Out] -1/105*(15*cos(b*x + a)^7 - 42*cos(b*x + a)^5 + 35*cos(b*x + a)^3)/b

Sympy [A]

time = 0.58, size = 68, normalized size = 1.48

$$\begin{cases} -\frac{\sin^4(a+bx)\cos^3(a+bx)}{3b} - \frac{4\sin^2(a+bx)\cos^5(a+bx)}{15b} - \frac{8\cos^7(a+bx)}{105b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos^2(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2*sin(b*x+a)**5,x)

[Out] Piecewise((-sin(a + b*x)**4*cos(a + b*x)**3/(3*b) - 4*sin(a + b*x)**2*cos(a + b*x)**5/(15*b) - 8*cos(a + b*x)**7/(105*b), Ne(b, 0)), (x*sin(a)**5*cos(a)**2, True))

Giac [A]

time = 3.89, size = 54, normalized size = 1.17

$$-\frac{\cos(7bx + 7a)}{448b} + \frac{3\cos(5bx + 5a)}{320b} - \frac{\cos(3bx + 3a)}{192b} - \frac{5\cos(bx + a)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*sin(b*x+a)^5,x, algorithm="giac")

[Out] -1/448*cos(7*b*x + 7*a)/b + 3/320*cos(5*b*x + 5*a)/b - 1/192*cos(3*b*x + 3*a)/b - 5/64*cos(b*x + a)/b

Mupad [B]

time = 0.37, size = 36, normalized size = 0.78

$$-\frac{15\cos(a+bx)^7 - 42\cos(a+bx)^5 + 35\cos(a+bx)^3}{105b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2*sin(a + b*x)^5,x)

[Out] -(35*cos(a + b*x)^3 - 42*cos(a + b*x)^5 + 15*cos(a + b*x)^7)/(105*b)

3.104 $\int \cos(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\sin^6(a + bx)}{6b}$$

[Out] 1/6*sin(b*x+a)^6/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2644, 30}

$$\frac{\sin^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Sin[a + b*x]^5,x]

[Out] Sin[a + b*x]^6/(6*b)

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_) + (f_)*(x_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \sin^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^5 dx, x, \sin(a + bx)\right)}{b} \\ &= \frac{\sin^6(a + bx)}{6b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$\frac{\sin^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x]*Sin[a + b*x]^5,x]
```

```
[Out] Sin[a + b*x]^6/(6*b)
```

Maple [A]

time = 0.06, size = 14, normalized size = 0.93

method	result	size
derivativedivides	$\frac{\sin^6(bx+a)}{6b}$	14
default	$\frac{\sin^6(bx+a)}{6b}$	14
norman	$\frac{32 \left(\tan^6 \left(\frac{bx+a}{2} \right) \right)}{3b \left(1 + \tan^2 \left(\frac{bx+a}{2} \right) \right)^6}$	32
risch	$-\frac{\cos(6bx+6a)}{192b} + \frac{\cos(4bx+4a)}{32b} - \frac{5 \cos(2bx+2a)}{64b}$	44

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(b*x+a)*sin(b*x+a)^5,x,method=_RETURNVERBOSE)
```

```
[Out] 1/6*sin(b*x+a)^6/b
```

Maxima [A]

time = 0.29, size = 13, normalized size = 0.87

$$\frac{\sin(bx+a)^6}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)*sin(b*x+a)^5,x, algorithm="maxima")
```

```
[Out] 1/6*sin(b*x + a)^6/b
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(13) = 26.

time = 0.39, size = 34, normalized size = 2.27

$$-\frac{\cos(bx+a)^6 - 3 \cos(bx+a)^4 + 3 \cos(bx+a)^2}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)*sin(b*x+a)^5,x, algorithm="fricas")
```

```
[Out] -1/6*(cos(b*x + a)^6 - 3*cos(b*x + a)^4 + 3*cos(b*x + a)^2)/b
```

Sympy [A]

time = 0.35, size = 20, normalized size = 1.33

$$\begin{cases} \frac{\sin^6(a+bx)}{6b} & \text{for } b \neq 0 \\ x \sin^5(a) \cos(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)**5,x)``[Out] Piecewise((sin(a + b*x)**6/(6*b), Ne(b, 0)), (x*sin(a)**5*cos(a), True))`**Giac [A]**

time = 3.48, size = 13, normalized size = 0.87

$$\frac{\sin(bx + a)^6}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)*sin(b*x+a)^5,x, algorithm="giac")``[Out] 1/6*sin(b*x + a)^6/b`**Mupad [B]**

time = 0.05, size = 13, normalized size = 0.87

$$\frac{\sin(a + bx)^6}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)*sin(a + b*x)^5,x)``[Out] sin(a + b*x)^6/(6*b)`

3.105 $\int \sin^4(a + bx) \tan(a + bx) dx$

Optimal. Leaf size=40

$$\frac{\cos^2(a + bx)}{b} - \frac{\cos^4(a + bx)}{4b} - \frac{\log(\cos(a + bx))}{b}$$

[Out] $\cos(b*x+a)^2/b-1/4*\cos(b*x+a)^4/b-\ln(\cos(b*x+a))/b$

Rubi [A]

time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2670, 272, 45}

$$-\frac{\cos^4(a + bx)}{4b} + \frac{\cos^2(a + bx)}{b} - \frac{\log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Sin[a + b*x]^4*Tan[a + b*x],x]`

[Out] `Cos[a + b*x]^2/b - Cos[a + b*x]^4/(4*b) - Log[Cos[a + b*x]]/b`

Rule 45

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LtQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 272

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Rule 2670

`Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]`

Rubi steps

$$\begin{aligned}
\int \sin^4(a+bx) \tan(a+bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x} dx, x, \cos(a+bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{(1-x)^2}{x} dx, x, \cos^2(a+bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \left(-2 + \frac{1}{x} + x\right) dx, x, \cos^2(a+bx)\right)}{2b} \\
&= \frac{\cos^2(a+bx)}{b} - \frac{\cos^4(a+bx)}{4b} - \frac{\log(\cos(a+bx))}{b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 35, normalized size = 0.88

$$-\frac{-\cos^2(a+bx) + \frac{1}{4}\cos^4(a+bx) + \log(\cos(a+bx))}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^4*Tan[a + b*x], x]``[Out] -((-Cos[a + b*x]^2 + Cos[a + b*x]^4/4 + Log[Cos[a + b*x]])/b)`**Maple [A]**

time = 0.05, size = 35, normalized size = 0.88

method	result
derivativedivides	$\frac{-\frac{(\sin^4(bx+a))}{4} - \frac{(\sin^2(bx+a))}{2} - \ln(\cos(bx+a))}{b}$
default	$\frac{-\frac{(\sin^4(bx+a))}{4} - \frac{(\sin^2(bx+a))}{2} - \ln(\cos(bx+a))}{b}$
risch	$ix + \frac{3e^{2i(bx+a)}}{16b} + \frac{3e^{-2i(bx+a)}}{16b} + \frac{2ia}{b} - \frac{\ln(e^{2i(bx+a)}+1)}{b} - \frac{\cos(4bx+4a)}{32b}$
norman	$\frac{-\frac{2(\tan^2(\frac{bx+a}{2}))}{b} - \frac{2(\tan^6(\frac{bx+a}{2}))}{b} - \frac{8(\tan^4(\frac{bx+a}{2}))}{b}}{(1+\tan^2(\frac{bx+a}{2}))^4} + \frac{\ln(1+\tan^2(\frac{bx+a}{2}))}{b} - \frac{\ln(\tan(\frac{bx+a}{2})-1)}{b} - \frac{\ln(\tan(\frac{bx+a}{2}))}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/4*sin(b*x+a)^4-1/2*sin(b*x+a)^2-ln(cos(b*x+a)))`**Maxima [A]**

time = 0.28, size = 37, normalized size = 0.92

$$-\frac{\sin^4(bx+a) + 2\sin^2(bx+a) + 2\log(\sin^2(bx+a) - 1)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/4*(sin(b*x + a)^4 + 2*sin(b*x + a)^2 + 2*log(sin(b*x + a)^2 - 1))/b

Fricas [A]

time = 0.37, size = 35, normalized size = 0.88

$$\frac{\cos(bx+a)^4 - 4\cos(bx+a)^2 + 4\log(-\cos(bx+a))}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^5,x, algorithm="fricas")

[Out] -1/4*(cos(b*x + a)^4 - 4*cos(b*x + a)^2 + 4*log(-cos(b*x + a)))/b

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)**5,x)

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 226 vs. 2(38) = 76.

time = 2.85, size = 226, normalized size = 5.65

$$\frac{3 \left(\frac{\cos(bx+a)+1}{\cos(bx+a)-1} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} \right)^2 - \frac{20(\cos(bx+a)+1)}{\cos(bx+a)-1} - \frac{20(\cos(bx+a)-1)}{\cos(bx+a)+1} + 44}{\left(\frac{\cos(bx+a)+1}{\cos(bx+a)-1} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 2 \right)^2} - 2 \log \left(\left| -\frac{\cos(bx+a)+1}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 2 \right| \right) + 2 \log \left(\left| -\frac{\cos(bx+a)+1}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 2 \right| \right)$$

4b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*sin(b*x+a)^5,x, algorithm="giac")

[Out] -1/4*((3*((cos(b*x + a) + 1)/(cos(b*x + a) - 1) + (cos(b*x + a) - 1)/(cos(b*x + a) + 1))^2 - 20*(cos(b*x + a) + 1)/(cos(b*x + a) - 1) - 20*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 44)/((cos(b*x + a) + 1)/(cos(b*x + a) - 1) + (cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 2)^2 - 2*log(abs(-(cos(b*x + a) + 1)/(cos(b*x + a) - 1) - (cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 2)) + 2*log(abs(-(cos(b*x + a) + 1)/(cos(b*x + a) - 1) - (cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 2)))/b

Mupad [B]

time = 0.54, size = 53, normalized size = 1.32

$$\frac{\ln(\tan(a+bx)^2 + 1)}{2b} + \frac{\tan(a+bx)^2 + \frac{3}{4}}{b(\tan(a+bx)^4 + 2\tan(a+bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(a + b*x)^5/cos(a + b*x),x)
```

```
[Out] log(tan(a + b*x)^2 + 1)/(2*b) + (tan(a + b*x)^2 + 3/4)/(b*(2*tan(a + b*x)^2  
+ tan(a + b*x)^4 + 1))
```

3.106 $\int \sin^3(a + bx) \tan^2(a + bx) dx$

Optimal. Leaf size=37

$$\frac{2 \cos(a + bx)}{b} - \frac{\cos^3(a + bx)}{3b} + \frac{\sec(a + bx)}{b}$$

[Out] 2*cos(b*x+a)/b-1/3*cos(b*x+a)^3/b+sec(b*x+a)/b

Rubi [A]

time = 0.02, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2670, 276}

$$-\frac{\cos^3(a + bx)}{3b} + \frac{2 \cos(a + bx)}{b} + \frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3*Tan[a + b*x]^2,x]

[Out] (2*Cos[a + b*x])/b - Cos[a + b*x]^3/(3*b) + Sec[a + b*x]/b

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned} \int \sin^3(a + bx) \tan^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x^2} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-2 + \frac{1}{x^2} + x^2\right) dx, x, \cos(a + bx)\right)}{b} \\ &= \frac{2 \cos(a + bx)}{b} - \frac{\cos^3(a + bx)}{3b} + \frac{\sec(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 39, normalized size = 1.05

$$\frac{7 \cos(a + bx)}{4b} - \frac{\cos(3(a + bx))}{12b} + \frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^3*Tan[a + b*x]^2,x]``[Out] (7*Cos[a + b*x])/(4*b) - Cos[3*(a + b*x)]/(12*b) + Sec[a + b*x]/b`**Maple [A]**

time = 0.04, size = 50, normalized size = 1.35

method	result	size
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{\cos(bx+a)} + \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{b}$	50
default	$\frac{\frac{\sin^6(bx+a)}{\cos(bx+a)} + \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{b}$	50
norman	$\frac{-\frac{16}{3b} - \frac{32(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{3b}}{\left(1 + \tan^2(\frac{bx}{2} + \frac{a}{2})\right)^3 \left(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1\right)}$	54
risch	$\frac{7e^{i(bx+a)}}{8b} + \frac{7e^{-i(bx+a)}}{8b} + \frac{2e^{i(bx+a)}}{b(e^{2i(bx+a)}+1)} - \frac{\cos(3bx+3a)}{12b}$	71

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^2*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(sin(b*x+a)^6/cos(b*x+a)+(8/3+sin(b*x+a)^4+4/3*sin(b*x+a)^2)*cos(b*x+a)`
`)`**Maxima [A]**

time = 0.29, size = 32, normalized size = 0.86

$$\frac{\cos(bx + a)^3 - \frac{3}{\cos(bx+a)} - 6 \cos(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^2*sin(b*x+a)^5,x, algorithm="maxima")``[Out] -1/3*(cos(b*x + a)^3 - 3/cos(b*x + a) - 6*cos(b*x + a))/b`**Fricas [A]**

time = 0.36, size = 33, normalized size = 0.89

$$\frac{\cos(bx + a)^4 - 6 \cos(bx + a)^2 - 3}{3b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^5,x, algorithm="fricas")

[Out] -1/3*(cos(b*x + a)^4 - 6*cos(b*x + a)^2 - 3)/(b*cos(b*x + a))

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: HeuristicGCDFailed

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2*sin(b*x+a)**5,x)

[Out] Exception raised: HeuristicGCDFailed >> no luck

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 99 vs. 2(35) = 70.
time = 4.66, size = 99, normalized size = 2.68

$$\frac{2 \left(\frac{3}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1} + \frac{\frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 5}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right)^3} \right)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*sin(b*x+a)^5,x, algorithm="giac")

[Out] 2/3*(3/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1) + (12*(cos(b*x + a) - 1) / (cos(b*x + a) + 1) - 3*(cos(b*x + a) - 1)^2 / (cos(b*x + a) + 1)^2 - 5) / ((cos(b*x + a) - 1) / (cos(b*x + a) + 1) - 1)^3) / b

Mupad [B]

time = 0.50, size = 31, normalized size = 0.84

$$-\frac{(\cos(a + bx) + 1)^3 (\cos(a + bx) - 3)}{3b \cos(a + bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^5/cos(a + b*x)^2,x)

[Out] -((cos(a + b*x) + 1)^3*(cos(a + b*x) - 3))/(3*b*cos(a + b*x))

3.107 $\int \sin^2(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=43

$$-\frac{\cos^2(a + bx)}{2b} + \frac{2 \log(\cos(a + bx))}{b} + \frac{\sec^2(a + bx)}{2b}$$

[Out] $-1/2*\cos(b*x+a)^2/b+2*\ln(\cos(b*x+a))/b+1/2*\sec(b*x+a)^2/b$

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2670, 272, 45}

$$-\frac{\cos^2(a + bx)}{2b} + \frac{\sec^2(a + bx)}{2b} + \frac{2 \log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2*Tan[a + b*x]^3,x]

[Out] $-1/2*\cos[a + b*x]^2/b + (2*\log[\cos[a + b*x]])/b + \sec[a + b*x]^2/(2*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \sin^2(a + bx) \tan^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x^3} dx, x, \cos(a + bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{(1-x)^2}{x^2} dx, x, \cos^2(a + bx)\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^2} - \frac{2}{x}\right) dx, x, \cos^2(a + bx)\right)}{2b} \\
&= -\frac{\cos^2(a + bx)}{2b} + \frac{2 \log(\cos(a + bx))}{b} + \frac{\sec^2(a + bx)}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 33, normalized size = 0.77

$$\frac{4 \log(\cos(a + bx)) + \sec^2(a + bx) + \sin^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^2*Tan[a + b*x]^3,x]``[Out] (4*Log[Cos[a + b*x]] + Sec[a + b*x]^2 + Sin[a + b*x]^2)/(2*b)`**Maple [A]**

time = 0.06, size = 51, normalized size = 1.19

method	result
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{2 \cos(bx+a)^2} + \frac{(\sin^4(bx+a))}{2} + \sin^2(bx+a) + 2 \ln(\cos(bx+a))}{b}$
default	$\frac{\frac{\sin^6(bx+a)}{2 \cos(bx+a)^2} + \frac{(\sin^4(bx+a))}{2} + \sin^2(bx+a) + 2 \ln(\cos(bx+a))}{b}$
risch	$-2ix - \frac{e^{2i(bx+a)}}{8b} - \frac{e^{-2i(bx+a)}}{8b} - \frac{4ia}{b} + \frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}+1)^2} + \frac{2 \ln(e^{2i(bx+a)}+1)}{b}$
norman	$\frac{\frac{4(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{4(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^2 (\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^2} + \frac{2 \ln(\tan(\frac{bx}{2} + \frac{a}{2}) - 1)}{b} + \frac{2 \ln(\tan(\frac{bx}{2} + \frac{a}{2}) + 1)}{b} - \frac{2 \ln(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^3*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/2*sin(b*x+a)^6/cos(b*x+a)^2+1/2*sin(b*x+a)^4+sin(b*x+a)^2+2*ln(cos(b*x+a)))`

Maxima [A]

time = 0.31, size = 41, normalized size = 0.95

$$\frac{\sin (bx+a)^2 - \frac{1}{\sin (bx+a)^2-1} + 2 \log (\sin (bx+a)^2 - 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3*sin(b*x+a)^5,x, algorithm="maxima")``[Out] 1/2*(sin(b*x + a)^2 - 1/(sin(b*x + a)^2 - 1) + 2*log(sin(b*x + a)^2 - 1))/b`**Fricas [A]**

time = 0.39, size = 54, normalized size = 1.26

$$-\frac{2 \cos (bx+a)^4 - 8 \cos (bx+a)^2 \log (-\cos (bx+a)) - \cos (bx+a)^2 - 2}{4 b \cos (bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3*sin(b*x+a)^5,x, algorithm="fricas")``[Out] -1/4*(2*cos(b*x + a)^4 - 8*cos(b*x + a)^2*log(-cos(b*x + a)) - cos(b*x + a)^2 - 2)/(b*cos(b*x + a)^2)`**Sympy [F(-1)]** Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**3*sin(b*x+a)**5,x)``[Out] Timed out`**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(39) = 78.

time = 6.07, size = 182, normalized size = 4.23

$$\frac{4 \left(\frac{\cos (bx+a)+1}{\cos (bx+a)-1} + \frac{\cos (bx+a)-1}{\cos (bx+a)+1} \right)}{\left(\frac{\cos (bx+a)+1}{\cos (bx+a)-1} + \frac{\cos (bx+a)-1}{\cos (bx+a)+1} \right)^2 - 4} + \log \left(\left| -\frac{\cos (bx+a)+1}{\cos (bx+a)-1} - \frac{\cos (bx+a)-1}{\cos (bx+a)+1} + 2 \right| \right) - \log \left(\left| -\frac{\cos (bx+a)+1}{\cos (bx+a)-1} - \frac{\cos (bx+a)-1}{\cos (bx+a)+1} - 2 \right| \right)$$

b

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3*sin(b*x+a)^5,x, algorithm="giac")``[Out] -(4*((cos(b*x + a) + 1)/(cos(b*x + a) - 1) + (cos(b*x + a) - 1)/(cos(b*x + a) + 1)))/(((cos(b*x + a) + 1)/(cos(b*x + a) - 1) + (cos(b*x + a) - 1)/(cos(`

$b*x + a) + 1))^2 - 4) + \log(\text{abs}(-(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 2)) - \log(\text{abs}(-(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 2)))/b$

Mupad [B]

time = 0.40, size = 37, normalized size = 0.86

$$-\frac{\ln(\tan(a + bx)^2 + 1) + \frac{\cos(a + bx)^2}{2} - \frac{\tan(a + bx)^2}{2}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^3,x)`

[Out] `-(log(tan(a + b*x)^2 + 1) + cos(a + b*x)^2/2 - tan(a + b*x)^2/2)/b`

3.108 $\int \sin(a + bx) \tan^4(a + bx) dx$

Optimal. Leaf size=38

$$-\frac{\cos(a + bx)}{b} - \frac{2 \sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b}$$

[Out] $-\cos(b*x+a)/b-2*\sec(b*x+a)/b+1/3*\sec(b*x+a)^3/b$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 276}

$$-\frac{\cos(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b} - \frac{2 \sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[a + b*x]*\text{Tan}[a + b*x]^4, x]$

[Out] $-(\text{Cos}[a + b*x])/b - (2*\text{Sec}[a + b*x])/b + \text{Sec}[a + b*x]^3/(3*b)$

Rule 276

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{Exp andIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \&\& \text{IGtQ}[p, 0]$

Rule 2670

$\text{Int}[\text{sin}[(e_*) + (f_*)(x_)]^{(m_*)}*\text{tan}[(e_*) + (f_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-f^{(-1)}, \text{Subst}[\text{Int}[(1 - x^2)^{(m + n - 1)/2}/x^n, x], x, \text{Cos}[e + f*x]], x] /; \text{FreeQ}\{e, f, x\} \&\& \text{IntegersQ}[m, n, (m + n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \sin(a + bx) \tan^4(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x^4} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^4} - \frac{2}{x^2}\right) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos(a + bx)}{b} - \frac{2 \sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 38, normalized size = 1.00

$$-\frac{\cos(a+bx)}{b} - \frac{2\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]*Tan[a + b*x]^4,x]
```

```
[Out] -(Cos[a + b*x]/b) - (2*Sec[a + b*x])/b + Sec[a + b*x]^3/(3*b)
```

Maple [A]

time = 0.06, size = 70, normalized size = 1.84

method	result	size
norman	$\frac{\frac{16}{3b} - \frac{32(\tan^2(\frac{bx+a}{2}))}{3b}}{(1+\tan^2(\frac{bx+a}{2}))(\tan^2(\frac{bx+a}{2})-1)^3}$	54
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{3\cos(bx+a)^3} - \frac{\sin^6(bx+a)}{\cos(bx+a)} - \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right)\cos(bx+a)}{b}$	70
default	$\frac{\frac{\sin^6(bx+a)}{3\cos(bx+a)^3} - \frac{\sin^6(bx+a)}{\cos(bx+a)} - \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right)\cos(bx+a)}{b}$	70
risch	$-\frac{e^{i(bx+a)}}{2b} - \frac{e^{-i(bx+a)}}{2b} - \frac{4(3e^{5i(bx+a)} + 4e^{3i(bx+a)} + 3e^{i(bx+a)})}{3b(e^{2i(bx+a)} + 1)^3}$	82

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(b*x+a)^4*sin(b*x+a)^5,x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(1/3*sin(b*x+a)^6/cos(b*x+a)^3-sin(b*x+a)^6/cos(b*x+a)-(8/3+sin(b*x+a)^4+4/3*sin(b*x+a)^2)*cos(b*x+a))
```

Maxima [A]

time = 0.29, size = 35, normalized size = 0.92

$$-\frac{\frac{6\cos(bx+a)^2-1}{\cos(bx+a)^3} + 3\cos(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)^4*sin(b*x+a)^5,x, algorithm="maxima")
```

```
[Out] -1/3*((6*cos(b*x + a)^2 - 1)/cos(b*x + a)^3 + 3*cos(b*x + a))/b
```

Fricas [A]

time = 0.39, size = 35, normalized size = 0.92

$$-\frac{3\cos(bx+a)^4 + 6\cos(bx+a)^2 - 1}{3b\cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4*sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $-1/3*(3*\cos(b*x + a)^4 + 6*\cos(b*x + a)^2 - 1)/(b*\cos(b*x + a)^3)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**4*sin(b*x+a)**5,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 100 vs. 2(36) = 72.

time = 5.55, size = 100, normalized size = 2.63

$$\frac{2 \left(\frac{3}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1} - \frac{\frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 5}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right)^3} \right)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $2/3*(3/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1) - (12*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 3*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 5)/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)^3)/b$

Mupad [B]

time = 0.53, size = 35, normalized size = 0.92

$$\frac{3 \cos(a + bx)^4 + 6 \cos(a + bx)^2 - 1}{3b \cos(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^4,x)`

[Out] $-(6*\cos(a + b*x)^2 + 3*\cos(a + b*x)^4 - 1)/(3*b*\cos(a + b*x)^3)$

3.109 $\int \tan^5(a + bx) dx$

Optimal. Leaf size=43

$$-\frac{\log(\cos(a + bx))}{b} - \frac{\tan^2(a + bx)}{2b} + \frac{\tan^4(a + bx)}{4b}$$

[Out] $-\ln(\cos(b*x+a))/b-1/2*\tan(b*x+a)^2/b+1/4*\tan(b*x+a)^4/b$

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$\frac{\tan^4(a + bx)}{4b} - \frac{\tan^2(a + bx)}{2b} - \frac{\log(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Tan[a + b*x]^5,x]

[Out] $-(\text{Log}[\text{Cos}[a + b*x]]/b) - \text{Tan}[a + b*x]^2/(2*b) + \text{Tan}[a + b*x]^4/(4*b)$

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \tan^5(a + bx) dx &= \frac{\tan^4(a + bx)}{4b} - \int \tan^3(a + bx) dx \\ &= -\frac{\tan^2(a + bx)}{2b} + \frac{\tan^4(a + bx)}{4b} + \int \tan(a + bx) dx \\ &= -\frac{\log(\cos(a + bx))}{b} - \frac{\tan^2(a + bx)}{2b} + \frac{\tan^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 37, normalized size = 0.86

$$-\frac{4 \log(\cos(a + bx)) + 2 \tan^2(a + bx) - \tan^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Tan[a + b*x]^5,x]

[Out] $-1/4*(4*\text{Log}[\text{Cos}[a + b*x]] + 2*\text{Tan}[a + b*x]^2 - \text{Tan}[a + b*x]^4)/b$

Maple [A]

time = 0.06, size = 35, normalized size = 0.81

method	result
derivativedivides	$\frac{(\tan^4(bx+a))}{4} - \frac{(\tan^2(bx+a))}{b} - \ln(\cos(bx+a))$
default	$\frac{(\tan^4(bx+a))}{4} - \frac{(\tan^2(bx+a))}{b} - \ln(\cos(bx+a))$
risch	$ix + \frac{2ia}{b} - \frac{4(e^{6i(bx+a)} + e^{4i(bx+a)} + e^{2i(bx+a)})}{b(e^{2i(bx+a)} + 1)^4} - \frac{\ln(e^{2i(bx+a)} + 1)}{b}$
norman	$\frac{-\frac{2(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{2(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{8(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^4} + \frac{\ln(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}) - 1)}{b} - \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^5*sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] $1/b*(1/4*\tan(b*x+a)^4 - 1/2*\tan(b*x+a)^2 - \ln(\cos(b*x+a)))$

Maxima [A]

time = 0.29, size = 54, normalized size = 1.26

$$\frac{\frac{4 \sin(bx+a)^2 - 3}{\sin(bx+a)^4 - 2 \sin(bx+a)^2 + 1} - 2 \log(\sin(bx+a)^2 - 1)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^5,x, algorithm="maxima")

[Out] $1/4*((4*\sin(b*x + a)^2 - 3)/(\sin(b*x + a)^4 - 2*\sin(b*x + a)^2 + 1) - 2*\log(\sin(b*x + a)^2 - 1))/b$

Fricas [A]

time = 0.38, size = 44, normalized size = 1.02

$$\frac{4 \cos(bx+a)^4 \log(-\cos(bx+a)) + 4 \cos(bx+a)^2 - 1}{4b \cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5*sin(b*x+a)^5,x, algorithm="fricas")

[Out] $-1/4*(4*\cos(b*x + a)^4*\log(-\cos(b*x + a)) + 4*\cos(b*x + a)^2 - 1)/(b*\cos(b*x + a)^4)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**5*sin(b*x+a)**5,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 226 vs. 2(39) = 78.

time = 4.41, size = 226, normalized size = 5.26

$$\frac{3 \left(\frac{\cos(bx+a)+1}{\cos(bx+a)-1} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} \right)^2 + \frac{20(\cos(bx+a)+1)}{\cos(bx+a)-1} + \frac{20(\cos(bx+a)-1)}{\cos(bx+a)+1} + 44}{\left(\frac{\cos(bx+a)+1}{\cos(bx+a)-1} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 2 \right)^2} + 2 \log \left(\left| \frac{\cos(bx+a)+1}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 2 \right| \right) - 2 \log \left(\left| \frac{\cos(bx+a)+1}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 2 \right| \right)$$

4b

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^5*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $1/4*((3*((\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) + (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1)))^2 + 20*(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) + 20*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 44)/((\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) + (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 2)^2 + 2*\log(\text{abs}(-(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 2)) - 2*\log(\text{abs}(-(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 2)))/b$

Mupad [B]

time = 0.40, size = 38, normalized size = 0.88

$$\frac{\frac{\ln(\tan(a+bx)^2+1)}{2} - \frac{\tan(a+bx)^2}{2} + \frac{\tan(a+bx)^4}{4}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^5,x)`

[Out] $(\log(\tan(a + b*x)^2 + 1)/2 - \tan(a + b*x)^2/2 + \tan(a + b*x)^4/4)/b$

3.110 $\int \sec(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=41

$$\frac{\sec(a + bx)}{b} - \frac{2 \sec^3(a + bx)}{3b} + \frac{\sec^5(a + bx)}{5b}$$

[Out] $\sec(b*x+a)/b-2/3*\sec(b*x+a)^3/b+1/5*\sec(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2686, 200}

$$\frac{\sec^5(a + bx)}{5b} - \frac{2 \sec^3(a + bx)}{3b} + \frac{\sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[a + b*x]*\text{Tan}[a + b*x]^5, x]$

[Out] $\text{Sec}[a + b*x]/b - (2*\text{Sec}[a + b*x]^3)/(3*b) + \text{Sec}[a + b*x]^5/(5*b)$

Rule 200

$\text{Int}[(a_ + (b_)*(x_)^{(n_)})^{(p_)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2686

$\text{Int}[(a_)*\sec[(e_ + (f_)*(x_))]^{(m_)}*((b_)*\tan[(e_ + (f_)*(x_))]^{(n_)}), x_Symbol] :> \text{Dist}[a/f, \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1 + x^2)^{(n-1)/2}], x], x, \text{Sec}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m, x\} \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[m/2] \ \&\& \ \text{LtQ}[0, m, n + 1])$

Rubi steps

$$\begin{aligned} \int \sec(a + bx) \tan^5(a + bx) dx &= \frac{\text{Subst}\left(\int (-1 + x^2)^2 dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (1 - 2x^2 + x^4) dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\sec(a + bx)}{b} - \frac{2 \sec^3(a + bx)}{3b} + \frac{\sec^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 41, normalized size = 1.00

$$\frac{\sec(a + bx)}{b} - \frac{2 \sec^3(a + bx)}{3b} + \frac{\sec^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]*Tan[a + b*x]^5,x]**[Out]** Sec[a + b*x]/b - (2*Sec[a + b*x]^3)/(3*b) + Sec[a + b*x]^5/(5*b)**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 87 vs. $2(37) = 74$.

time = 0.07, size = 88, normalized size = 2.15

method	result	size
norman	$\frac{-\frac{16}{15b} + \frac{16(\tan^2(\frac{bx+a}{2}))}{3b} - \frac{32(\tan^4(\frac{bx+a}{2}))}{3b}}{(\tan^2(\frac{bx+a}{2}) - 1)^5}$	55
risch	$\frac{2e^{9i(bx+a)} + \frac{8e^{7i(bx+a)}}{3} + \frac{116e^{5i(bx+a)}}{15} + \frac{8e^{3i(bx+a)}}{3} + 2e^{i(bx+a)}}{b(e^{2i(bx+a)} + 1)^5}$	75
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{5 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{15 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{5 \cos(bx+a)} + \frac{\left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{5}}{b}$	88
default	$\frac{\frac{\sin^6(bx+a)}{5 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{15 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{5 \cos(bx+a)} + \frac{\left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{5}}{b}$	88

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^6*sin(b*x+a)^5,x,method=_RETURNVERBOSE)**[Out]** 1/b*(1/5*sin(b*x+a)^6/cos(b*x+a)^5-1/15*sin(b*x+a)^6/cos(b*x+a)^3+1/5*sin(b*x+a)^6/cos(b*x+a)+1/5*(8/3+sin(b*x+a)^4+4/3*sin(b*x+a)^2)*cos(b*x+a))**Maxima [A]**

time = 0.29, size = 35, normalized size = 0.85

$$\frac{15 \cos(bx + a)^4 - 10 \cos(bx + a)^2 + 3}{15 b \cos(bx + a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^5,x, algorithm="maxima")**[Out]** 1/15*(15*cos(b*x + a)^4 - 10*cos(b*x + a)^2 + 3)/(b*cos(b*x + a)^5)

Fricas [A]

time = 0.35, size = 35, normalized size = 0.85

$$\frac{15 \cos (bx+a)^4 - 10 \cos (bx+a)^2 + 3}{15 b \cos (bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6*sin(b*x+a)^5,x, algorithm="fricas")``[Out] 1/15*(15*cos(b*x + a)^4 - 10*cos(b*x + a)^2 + 3)/(b*cos(b*x + a)^5)`**Sympy [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**6*sin(b*x+a)**5,x)``[Out] Exception raised: SystemError >> excessive stack use: stack is 6189 deep`**Giac [A]**

time = 4.82, size = 72, normalized size = 1.76

$$\frac{16 \left(\frac{5(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{10(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 1 \right)}{15 b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6*sin(b*x+a)^5,x, algorithm="giac")``[Out] 16/15*(5*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 10*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 1)/(b*((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^5)`**Mupad [B]**

time = 0.54, size = 35, normalized size = 0.85

$$\frac{15 \cos (a+bx)^4 - 10 \cos (a+bx)^2 + 3}{15 b \cos (a+bx)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^5/cos(a + b*x)^6,x)``[Out] (15*cos(a + b*x)^4 - 10*cos(a + b*x)^2 + 3)/(15*b*cos(a + b*x)^5)`

3.111 $\int \sec^2(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=15

$$\frac{\tan^6(a + bx)}{6b}$$

[Out] 1/6*tan(b*x+a)^6/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 30}

$$\frac{\tan^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2*Tan[a + b*x]^5,x]

[Out] Tan[a + b*x]^6/(6*b)

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2687

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^2(a + bx) \tan^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^5 dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^6(a + bx)}{6b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$\frac{\tan^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2*Tan[a + b*x]^5,x]

[Out] Tan[a + b*x]^6/(6*b)

Maple [A]

time = 0.07, size = 22, normalized size = 1.47

method	result	size
derivativedivides	$\frac{\sin^6(bx+a)}{6b \cos(bx+a)^6}$	22
default	$\frac{\sin^6(bx+a)}{6b \cos(bx+a)^6}$	22
norman	$\frac{32 \left(\tan^6 \left(\frac{bx+a}{2} \right) \right)}{3b \left(\tan^2 \left(\frac{bx+a}{2} \right) - 1 \right)^6}$	32
risch	$\frac{2e^{10i(bx+a)} + 20e^{6i(bx+a)} + 2e^{2i(bx+a)}}{b(e^{2i(bx+a)} + 1)^6}$	53

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^7*sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/6/b*sin(b*x+a)^6/cos(b*x+a)^6

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(13) = 26.

time = 0.29, size = 59, normalized size = 3.93

$$-\frac{3 \sin (bx+a)^4 - 3 \sin (bx+a)^2 + 1}{6 \left(\sin (bx+a)^6 - 3 \sin (bx+a)^4 + 3 \sin (bx+a)^2 - 1 \right) b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7*sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/6*(3*sin(b*x + a)^4 - 3*sin(b*x + a)^2 + 1)/((sin(b*x + a)^6 - 3*sin(b*x + a)^4 + 3*sin(b*x + a)^2 - 1)*b)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 35 vs. 2(13) = 26.

time = 0.33, size = 35, normalized size = 2.33

$$\frac{3 \cos (bx+a)^4 - 3 \cos (bx+a)^2 + 1}{6 b \cos (bx+a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7*sin(b*x+a)^5,x, algorithm="fricas")

[Out] $1/6*(3*\cos(b*x + a)^4 - 3*\cos(b*x + a)^2 + 1)/(b*\cos(b*x + a)^6)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**7*sin(b*x+a)**5,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 8569 deep

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(13) = 26$.

time = 4.69, size = 48, normalized size = 3.20

$$-\frac{32(\cos(bx+a)-1)^3}{3b\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}+1\right)^6(\cos(bx+a)+1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^7*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $-32/3*(\cos(b*x + a) - 1)^3/(b*((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)^6 * (\cos(b*x + a) + 1)^3)$

Mupad [B]

time = 0.42, size = 13, normalized size = 0.87

$$\frac{\tan(a + bx)^6}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^7,x)`

[Out] $\tan(a + b*x)^6/(6*b)$

3.112 $\int \sec^3(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sec^3(a + bx)}{3b} - \frac{2 \sec^5(a + bx)}{5b} + \frac{\sec^7(a + bx)}{7b}$$

[Out] $1/3*\sec(b*x+a)^3/b-2/5*\sec(b*x+a)^5/b+1/7*\sec(b*x+a)^7/b$

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 276}

$$\frac{\sec^7(a + bx)}{7b} - \frac{2 \sec^5(a + bx)}{5b} + \frac{\sec^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] `Int[Sec[a + b*x]^3*Tan[a + b*x]^5,x]`

[Out] `Sec[a + b*x]^3/(3*b) - (2*Sec[a + b*x]^5)/(5*b) + Sec[a + b*x]^7/(7*b)`

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2686

`Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^(n - 1)/2], x], x, Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])`

Rubi steps

$$\begin{aligned} \int \sec^3(a + bx) \tan^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^2(-1 + x^2)^2 dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^2 - 2x^4 + x^6) dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\sec^3(a + bx)}{3b} - \frac{2 \sec^5(a + bx)}{5b} + \frac{\sec^7(a + bx)}{7b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 46, normalized size = 1.00

$$\frac{\sec^3(a + bx)}{3b} - \frac{2\sec^5(a + bx)}{5b} + \frac{\sec^7(a + bx)}{7b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^3*Tan[a + b*x]^5,x]``[Out] Sec[a + b*x]^3/(3*b) - (2*Sec[a + b*x]^5)/(5*b) + Sec[a + b*x]^7/(7*b)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(40) = 80.

time = 0.08, size = 106, normalized size = 2.30

method	result	size
risch	$\frac{8e^{11i(bx+a)} - 32e^{9i(bx+a)} + 304e^{7i(bx+a)} - 32e^{5i(bx+a)} + 8e^{3i(bx+a)}}{3 \cdot 15 \cdot 35 \cdot 15 \cdot 3} \cdot \frac{1}{b(e^{2i(bx+a)} + 1)^7}$	75
norman	$-\frac{16}{105b} - \frac{16(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{16(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{15b} - \frac{16(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{32(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{3b}$ $\frac{1}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^7}$	87
derivativdivides	$\frac{\frac{\sin^6(bx+a)}{7 \cos(bx+a)^7} + \frac{\sin^6(bx+a)}{35 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{105 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{35 \cos(bx+a)} + \frac{\left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{35}}{b}$	106
default	$\frac{\frac{\sin^6(bx+a)}{7 \cos(bx+a)^7} + \frac{\sin^6(bx+a)}{35 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{105 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{35 \cos(bx+a)} + \frac{\left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3}\right) \cos(bx+a)}{35}}{b}$	106

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^8*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/7*sin(b*x+a)^6/cos(b*x+a)^7+1/35*sin(b*x+a)^6/cos(b*x+a)^5-1/105*sin(b*x+a)^6/cos(b*x+a)^3+1/35*sin(b*x+a)^6/cos(b*x+a)+1/35*(8/3+sin(b*x+a)^4+4/3*sint(b*x+a)^2)*cos(b*x+a))`**Maxima [A]**

time = 0.30, size = 35, normalized size = 0.76

$$\frac{35 \cos(bx + a)^4 - 42 \cos(bx + a)^2 + 15}{105 b \cos(bx + a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^8*sin(b*x+a)^5,x, algorithm="maxima")``[Out] 1/105*(35*cos(b*x + a)^4 - 42*cos(b*x + a)^2 + 15)/(b*cos(b*x + a)^7)`

Fricas [A]

time = 0.34, size = 35, normalized size = 0.76

$$\frac{35 \cos (bx+a)^4 - 42 \cos (bx+a)^2 + 15}{105 b \cos (bx+a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^8*sin(b*x+a)^5,x, algorithm="fricas")**[Out]** 1/105*(35*cos(b*x + a)^4 - 42*cos(b*x + a)^2 + 15)/(b*cos(b*x + a)^7)**Sympy [F(-1)]** Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**8*sin(b*x+a)**5,x)**[Out]** Timed out**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 116 vs. 2(40) = 80.

time = 5.30, size = 116, normalized size = 2.52

$$\frac{16 \left(\frac{7(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{21(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - \frac{35(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{70(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + 1 \right)}{105 b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^8*sin(b*x+a)^5,x, algorithm="giac")**[Out]** 16/105*(7*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 21*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 35*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 + 70*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 + 1)/(b*((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^7)**Mupad [B]**

time = 0.59, size = 35, normalized size = 0.76

$$\frac{35 \cos (a+bx)^4 - 42 \cos (a+bx)^2 + 15}{105 b \cos (a+bx)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^5/cos(a + b*x)^8,x)**[Out]** (35*cos(a + b*x)^4 - 42*cos(a + b*x)^2 + 15)/(105*b*cos(a + b*x)^7)

3.113 $\int \sec^4(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=31

$$\frac{\tan^6(a + bx)}{6b} + \frac{\tan^8(a + bx)}{8b}$$

[Out] 1/6*tan(b*x+a)^6/b+1/8*tan(b*x+a)^8/b

Rubi [A]

time = 0.02, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 14}

$$\frac{\tan^8(a + bx)}{8b} + \frac{\tan^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^4*Tan[a + b*x]^5,x]

[Out] Tan[a + b*x]^6/(6*b) + Tan[a + b*x]^8/(8*b)

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2687

Int[sec[(e_) + (f_)*(x_)]^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \sec^4(a + bx) \tan^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^5(1 + x^2) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^5 + x^7) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\tan^6(a + bx)}{6b} + \frac{\tan^8(a + bx)}{8b} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 38, normalized size = 1.23

$$\frac{6 \sec^4(a + bx) - 8 \sec^6(a + bx) + 3 \sec^8(a + bx)}{24b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^4*Tan[a + b*x]^5,x]``[Out] (6*Sec[a + b*x]^4 - 8*Sec[a + b*x]^6 + 3*Sec[a + b*x]^8)/(24*b)`**Maple [A]**

time = 0.10, size = 42, normalized size = 1.35

method	result	size
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{8 \cos(bx+a)^8} + \frac{\sin^6(bx+a)}{24 \cos(bx+a)^6}}{b}$	42
default	$\frac{\frac{\sin^6(bx+a)}{8 \cos(bx+a)^8} + \frac{\sin^6(bx+a)}{24 \cos(bx+a)^6}}{b}$	42
norman	$\frac{\frac{32 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right) \right)}{3b} + \frac{32 \left(\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right) \right)}{3b} + \frac{32 \left(\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right) \right)}{3b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1 \right)^8}$	66
risch	$\frac{4 e^{12i(bx+a)} - \frac{16 e^{10i(bx+a)}}{3} + \frac{40 e^{8i(bx+a)}}{3} - \frac{16 e^{6i(bx+a)}}{3} + 4 e^{4i(bx+a)}}{b(e^{2i(bx+a)} + 1)^8}$	75

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^9*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/8*sin(b*x+a)^6/cos(b*x+a)^8+1/24*sin(b*x+a)^6/cos(b*x+a)^6)`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(27) = 54$.

time = 0.28, size = 69, normalized size = 2.23

$$\frac{6 \sin(bx + a)^4 - 4 \sin(bx + a)^2 + 1}{24 (\sin(bx + a)^8 - 4 \sin(bx + a)^6 + 6 \sin(bx + a)^4 - 4 \sin(bx + a)^2 + 1)b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^9*sin(b*x+a)^5,x, algorithm="maxima")``[Out] 1/24*(6*sin(b*x + a)^4 - 4*sin(b*x + a)^2 + 1)/((sin(b*x + a)^8 - 4*sin(b*x + a)^6 + 6*sin(b*x + a)^4 - 4*sin(b*x + a)^2 + 1)*b)`**Fricas [A]**

time = 0.38, size = 35, normalized size = 1.13

$$\frac{6 \cos(bx + a)^4 - 8 \cos(bx + a)^2 + 3}{24 b \cos(bx + a)^8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^9*sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $1/24*(6*\cos(b*x + a)^4 - 8*\cos(b*x + a)^2 + 3)/(b*\cos(b*x + a)^8)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**9*sin(b*x+a)**5,x)`

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 93 vs. $2(27) = 54$.

time = 5.25, size = 93, normalized size = 3.00

$$-\frac{32 \left(\frac{(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + \frac{(\cos(bx+a)-1)^5}{(\cos(bx+a)+1)^5} \right)}{3b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^9*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $-32/3*((\cos(b*x + a) - 1)^3/(\cos(b*x + a) + 1)^3 - (\cos(b*x + a) - 1)^4/(\cos(b*x + a) + 1)^4 + (\cos(b*x + a) - 1)^5/(\cos(b*x + a) + 1)^5)/(b*((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)^8)$

Mupad [B]

time = 0.42, size = 25, normalized size = 0.81

$$\frac{\tan(a + bx)^6 (3 \tan(a + bx)^2 + 4)}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^9,x)`

[Out] $(\tan(a + b*x)^6*(3*\tan(a + b*x)^2 + 4))/(24*b)$

3.114 $\int \sec^5(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sec^5(a + bx)}{5b} - \frac{2 \sec^7(a + bx)}{7b} + \frac{\sec^9(a + bx)}{9b}$$

[Out] 1/5*sec(b*x+a)^5/b-2/7*sec(b*x+a)^7/b+1/9*sec(b*x+a)^9/b

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 276}

$$\frac{\sec^9(a + bx)}{9b} - \frac{2 \sec^7(a + bx)}{7b} + \frac{\sec^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^5*Tan[a + b*x]^5,x]

[Out] Sec[a + b*x]^5/(5*b) - (2*Sec[a + b*x]^7)/(7*b) + Sec[a + b*x]^9/(9*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^(n - 1)/2], x], x, Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \sec^5(a + bx) \tan^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^4(-1 + x^2)^2 dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^4 - 2x^6 + x^8) dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\sec^5(a + bx)}{5b} - \frac{2 \sec^7(a + bx)}{7b} + \frac{\sec^9(a + bx)}{9b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 46, normalized size = 1.00

$$\frac{\sec^5(a + bx)}{5b} - \frac{2 \sec^7(a + bx)}{7b} + \frac{\sec^9(a + bx)}{9b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^5*Tan[a + b*x]^5,x]``[Out] Sec[a + b*x]^5/(5*b) - (2*Sec[a + b*x]^7)/(7*b) + Sec[a + b*x]^9/(9*b)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 123 vs. 2(40) = 80.

time = 0.05, size = 124, normalized size = 2.70

method	result
risch	$\frac{32 e^{13i(bx+a)} - 384 e^{11i(bx+a)} + 6976 e^{9i(bx+a)} - 384 e^{7i(bx+a)} + 32 e^{5i(bx+a)}}{5 b (e^{2i(bx+a)} + 1)^9}$
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{9 \cos(bx+a)^9} + \frac{\sin^6(bx+a)}{21 \cos(bx+a)^7} + \frac{\sin^6(bx+a)}{105 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{315 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{105 \cos(bx+a)} + \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3} \right) \cos(bx+a)}{b}$
default	$\frac{\frac{\sin^6(bx+a)}{9 \cos(bx+a)^9} + \frac{\sin^6(bx+a)}{21 \cos(bx+a)^7} + \frac{\sin^6(bx+a)}{105 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{315 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{105 \cos(bx+a)} + \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{3} \right) \cos(bx+a)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^10*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/9*sin(b*x+a)^6/cos(b*x+a)^9+1/21*sin(b*x+a)^6/cos(b*x+a)^7+1/105*sin(b*x+a)^6/cos(b*x+a)^5-1/315*sin(b*x+a)^6/cos(b*x+a)^3+1/105*sin(b*x+a)^6/cos(b*x+a)+1/105*(8/3+sin(b*x+a)^4+4/3*sin(b*x+a)^2)*cos(b*x+a))`**Maxima [A]**

time = 0.29, size = 35, normalized size = 0.76

$$\frac{63 \cos(bx + a)^4 - 90 \cos(bx + a)^2 + 35}{315 b \cos(bx + a)^9}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^10*sin(b*x+a)^5,x, algorithm="maxima")``[Out] 1/315*(63*cos(b*x + a)^4 - 90*cos(b*x + a)^2 + 35)/(b*cos(b*x + a)^9)`**Fricas [A]**

time = 0.38, size = 35, normalized size = 0.76

$$\frac{63 \cos(bx + a)^4 - 90 \cos(bx + a)^2 + 35}{315 b \cos(bx + a)^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^10*sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $1/315*(63*\cos(b*x + a)^4 - 90*\cos(b*x + a)^2 + 35)/(b*\cos(b*x + a)^9)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**10*sin(b*x+a)**5,x)`

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(40) = 80.

time = 3.55, size = 160, normalized size = 3.48

$$\frac{16 \left(\frac{9(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{36(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - \frac{126(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{441(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - \frac{315(\cos(bx+a)-1)^5}{(\cos(bx+a)+1)^5} + \frac{210(\cos(bx+a)-1)^6}{(\cos(bx+a)+1)^6} + 1 \right)}{315 b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^10*sin(b*x+a)^5,x, algorithm="giac")`

[Out] $16/315*(9*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 36*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 - 126*(\cos(b*x + a) - 1)^3/(\cos(b*x + a) + 1)^3 + 441*(\cos(b*x + a) - 1)^4/(\cos(b*x + a) + 1)^4 - 315*(\cos(b*x + a) - 1)^5/(\cos(b*x + a) + 1)^5 + 210*(\cos(b*x + a) - 1)^6/(\cos(b*x + a) + 1)^6 + 1)/(b*((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)^9)$

Mupad [B]

time = 0.77, size = 35, normalized size = 0.76

$$\frac{63 \cos(a + b x)^4 - 90 \cos(a + b x)^2 + 35}{315 b \cos(a + b x)^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^10,x)`

[Out] $(63*\cos(a + b*x)^4 - 90*\cos(a + b*x)^2 + 35)/(315*b*\cos(a + b*x)^9)$

3.115 $\int \sec^6(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sec^6(a + bx)}{6b} - \frac{\sec^8(a + bx)}{4b} + \frac{\sec^{10}(a + bx)}{10b}$$

[Out] 1/6*sec(b*x+a)^6/b-1/4*sec(b*x+a)^8/b+1/10*sec(b*x+a)^10/b

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2686, 272, 45}

$$\frac{\sec^{10}(a + bx)}{10b} - \frac{\sec^8(a + bx)}{4b} + \frac{\sec^6(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^6*Tan[a + b*x]^5,x]

[Out] Sec[a + b*x]^6/(6*b) - Sec[a + b*x]^8/(4*b) + Sec[a + b*x]^10/(10*b)

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 2686

```
Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(
n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2)
, x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2]
&& !(IntegerQ[m/2] && LtQ[0, m, n + 1])
```

Rubi steps

$$\begin{aligned}
\int \sec^6(a+bx) \tan^5(a+bx) dx &= \frac{\text{Subst}\left(\int x^5(-1+x^2)^2 dx, x, \sec(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int (-1+x)^2 x^2 dx, x, \sec^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int (x^2-2x^3+x^4) dx, x, \sec^2(a+bx)\right)}{2b} \\
&= \frac{\sec^6(a+bx)}{6b} - \frac{\sec^8(a+bx)}{4b} + \frac{\sec^{10}(a+bx)}{10b}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 38, normalized size = 0.83

$$\frac{10 \sec^6(a+bx) - 15 \sec^8(a+bx) + 6 \sec^{10}(a+bx)}{60b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^6*Tan[a + b*x]^5, x]``[Out] (10*Sec[a + b*x]^6 - 15*Sec[a + b*x]^8 + 6*Sec[a + b*x]^10)/(60*b)`**Maple [A]**

time = 0.06, size = 60, normalized size = 1.30

method	result	size
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{10 \cos(bx+a)^{10}} + \frac{\sin^6(bx+a)}{20 \cos(bx+a)^8} + \frac{\sin^6(bx+a)}{60 \cos(bx+a)^6}}{b}$	60
default	$\frac{\frac{\sin^6(bx+a)}{10 \cos(bx+a)^{10}} + \frac{\sin^6(bx+a)}{20 \cos(bx+a)^8} + \frac{\sin^6(bx+a)}{60 \cos(bx+a)^6}}{b}$	60
risch	$\frac{\frac{32 e^{14i(bx+a)}}{3} - \frac{64 e^{12i(bx+a)}}{3} + \frac{192 e^{10i(bx+a)}}{5} - \frac{64 e^{8i(bx+a)}}{3} + \frac{32 e^{6i(bx+a)}}{3}}{b(e^{2i(bx+a)}+1)^{10}}$	75

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^11*sin(b*x+a)^5, x, method=_RETURNVERBOSE)``[Out] 1/b*(1/10*sin(b*x+a)^6/cos(b*x+a)^10+1/20*sin(b*x+a)^6/cos(b*x+a)^8+1/60*sin(b*x+a)^6/cos(b*x+a)^6)`**Maxima [A]**

time = 0.30, size = 79, normalized size = 1.72

$$\frac{10 \sin(bx+a)^4 - 5 \sin(bx+a)^2 + 1}{60 (\sin(bx+a)^{10} - 5 \sin(bx+a)^8 + 10 \sin(bx+a)^6 - 10 \sin(bx+a)^4 + 5 \sin(bx+a)^2 - 1) b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^11*sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/60*(10*sin(b*x + a)^4 - 5*sin(b*x + a)^2 + 1)/((sin(b*x + a)^10 - 5*sin(b*x + a)^8 + 10*sin(b*x + a)^6 - 10*sin(b*x + a)^4 + 5*sin(b*x + a)^2 - 1)*b)

Fricas [A]

time = 0.35, size = 35, normalized size = 0.76

$$\frac{10 \cos (b x+a)^4-15 \cos (b x+a)^2+6}{60 b \cos (b x+a)^{10}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^11*sin(b*x+a)^5,x, algorithm="fricas")

[Out] 1/60*(10*cos(b*x + a)^4 - 15*cos(b*x + a)^2 + 6)/(b*cos(b*x + a)^10)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**11*sin(b*x+a)**5,x)

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 139 vs. 2(40) = 80.

time = 5.27, size = 139, normalized size = 3.02

$$\frac{32 \left(\frac{5(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{10(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + \frac{18(\cos(bx+a)-1)^5}{(\cos(bx+a)+1)^5} - \frac{10(\cos(bx+a)-1)^6}{(\cos(bx+a)+1)^6} + \frac{5(\cos(bx+a)-1)^7}{(\cos(bx+a)+1)^7} \right)}{15 b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^{10}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^11*sin(b*x+a)^5,x, algorithm="giac")

[Out] -32/15*(5*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 - 10*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 + 18*(cos(b*x + a) - 1)^5/(cos(b*x + a) + 1)^5 - 10*(cos(b*x + a) - 1)^6/(cos(b*x + a) + 1)^6 + 5*(cos(b*x + a) - 1)^7/(cos(b*x + a) + 1)^7)/(b*((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^10)

Mupad [B]

time = 0.44, size = 35, normalized size = 0.76

$$\frac{\tan(a + b x)^6 (6 \tan(a + b x)^4 + 15 \tan(a + b x)^2 + 10)}{60 b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(a + b*x)^5/cos(a + b*x)^11,x)
```

```
[Out] (tan(a + b*x)^6*(15*tan(a + b*x)^2 + 6*tan(a + b*x)^4 + 10))/(60*b)
```

3.116 $\int \sec^7(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sec^7(a + bx)}{7b} - \frac{2 \sec^9(a + bx)}{9b} + \frac{\sec^{11}(a + bx)}{11b}$$

[Out] $1/7*\sec(b*x+a)^7/b-2/9*\sec(b*x+a)^9/b+1/11*\sec(b*x+a)^{11}/b$

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2686, 276}

$$\frac{\sec^{11}(a + bx)}{11b} - \frac{2 \sec^9(a + bx)}{9b} + \frac{\sec^7(a + bx)}{7b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^7*Tan[a + b*x]^5,x]

[Out] Sec[a + b*x]^7/(7*b) - (2*Sec[a + b*x]^9)/(9*b) + Sec[a + b*x]^11/(11*b)

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \sec^7(a + bx) \tan^5(a + bx) dx &= \frac{\text{Subst}\left(\int x^6(-1 + x^2)^2 dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int (x^6 - 2x^8 + x^{10}) dx, x, \sec(a + bx)\right)}{b} \\ &= \frac{\sec^7(a + bx)}{7b} - \frac{2 \sec^9(a + bx)}{9b} + \frac{\sec^{11}(a + bx)}{11b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 46, normalized size = 1.00

$$\frac{\sec^7(a + bx)}{7b} - \frac{2\sec^9(a + bx)}{9b} + \frac{\sec^{11}(a + bx)}{11b}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^7*Tan[a + b*x]^5,x]**[Out]** Sec[a + b*x]^7/(7*b) - (2*Sec[a + b*x]^9)/(9*b) + Sec[a + b*x]^11/(11*b)**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(40) = 80.

time = 0.07, size = 142, normalized size = 3.09

method	result
risch	$\frac{128 e^{15i(bx+a)} - 2560 e^{13i(bx+a)} + 47360 e^{11i(bx+a)} - 2560 e^{9i(bx+a)} + 128 e^{7i(bx+a)}}{7 \cdot 63 \cdot 63 \cdot 63 \cdot 7} \cdot b(e^{2i(bx+a)} + 1)^{11}$
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{11 \cos(bx+a)^{11}} + \frac{5(\sin^6(bx+a))}{99 \cos(bx+a)^9} + \frac{5(\sin^6(bx+a))}{231 \cos(bx+a)^7} + \frac{\sin^6(bx+a)}{231 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{693 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{231 \cos(bx+a)} + \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{231}\right)}{b}$
default	$\frac{\frac{\sin^6(bx+a)}{11 \cos(bx+a)^{11}} + \frac{5(\sin^6(bx+a))}{99 \cos(bx+a)^9} + \frac{5(\sin^6(bx+a))}{231 \cos(bx+a)^7} + \frac{\sin^6(bx+a)}{231 \cos(bx+a)^5} - \frac{\sin^6(bx+a)}{693 \cos(bx+a)^3} + \frac{\sin^6(bx+a)}{231 \cos(bx+a)} + \left(\frac{8}{3} + \sin^4(bx+a) + \frac{4(\sin^2(bx+a))}{231}\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^12*sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/b*(1/11*sin(b*x+a)^6/cos(b*x+a)^11+5/99*sin(b*x+a)^6/cos(b*x+a)^9+5/231*sin(b*x+a)^6/cos(b*x+a)^7+1/231*sin(b*x+a)^6/cos(b*x+a)^5-1/693*sin(b*x+a)^6/cos(b*x+a)^3+1/231*sin(b*x+a)^6/cos(b*x+a)+1/231*(8/3+sin(b*x+a)^4+4/3*sin(b*x+a)^2)*cos(b*x+a))

Maxima [A]

time = 0.29, size = 35, normalized size = 0.76

$$\frac{99 \cos(bx + a)^4 - 154 \cos(bx + a)^2 + 63}{693 b \cos(bx + a)^{11}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^12*sin(b*x+a)^5,x, algorithm="maxima")**[Out]** 1/693*(99*cos(b*x + a)^4 - 154*cos(b*x + a)^2 + 63)/(b*cos(b*x + a)^11)

Fricas [A]

time = 0.37, size = 35, normalized size = 0.76

$$\frac{99 \cos (bx+a)^4 - 154 \cos (bx+a)^2 + 63}{693 b \cos (bx+a)^{11}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^12*sin(b*x+a)^5,x, algorithm="fricas")

[Out] 1/693*(99*cos(b*x + a)^4 - 154*cos(b*x + a)^2 + 63)/(b*cos(b*x + a)^11)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**12*sin(b*x+a)**5,x)

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 204 vs. 2(40) = 80.

time = 5.32, size = 204, normalized size = 4.43

$$\frac{16 \left(\frac{11 \cos(bx+a)-1}{\cos(bx+a)+1} + \frac{55 (\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - \frac{297 (\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{1485 (\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - \frac{2079 (\cos(bx+a)-1)^5}{(\cos(bx+a)+1)^5} + \frac{2541 (\cos(bx+a)-1)^6}{(\cos(bx+a)+1)^6} - \frac{1155 (\cos(bx+a)-1)^7}{(\cos(bx+a)+1)^7} + \frac{462 (\cos(bx+a)-1)^8}{(\cos(bx+a)+1)^8} + 1 \right)}{693 b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^{11}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^12*sin(b*x+a)^5,x, algorithm="giac")

[Out] 16/693*(11*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 55*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 297*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 + 1485*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 - 2079*(cos(b*x + a) - 1)^5/(cos(b*x + a) + 1)^5 + 2541*(cos(b*x + a) - 1)^6/(cos(b*x + a) + 1)^6 - 1155*(cos(b*x + a) - 1)^7/(cos(b*x + a) + 1)^7 + 462*(cos(b*x + a) - 1)^8/(cos(b*x + a) + 1)^8 + 1)/(b*((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^11)

Mupad [B]

time = 1.02, size = 35, normalized size = 0.76

$$\frac{99 \cos (a+bx)^4 - 154 \cos (a+bx)^2 + 63}{693 b \cos (a+bx)^{11}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^5/cos(a + b*x)^12,x)

[Out] (99*cos(a + b*x)^4 - 154*cos(a + b*x)^2 + 63)/(693*b*cos(a + b*x)^11)

3.117 $\int \sec^8(a + bx) \tan^5(a + bx) dx$

Optimal. Leaf size=46

$$\frac{\sec^8(a + bx)}{8b} - \frac{\sec^{10}(a + bx)}{5b} + \frac{\sec^{12}(a + bx)}{12b}$$

[Out] 1/8*sec(b*x+a)^8/b-1/5*sec(b*x+a)^10/b+1/12*sec(b*x+a)^12/b

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2686, 272, 45}

$$\frac{\sec^{12}(a + bx)}{12b} - \frac{\sec^{10}(a + bx)}{5b} + \frac{\sec^8(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^8*Tan[a + b*x]^5,x]

[Out] Sec[a + b*x]^8/(8*b) - Sec[a + b*x]^10/(5*b) + Sec[a + b*x]^12/(12*b)

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned}
\int \sec^8(a+bx) \tan^5(a+bx) dx &= \frac{\text{Subst}\left(\int x^7(-1+x^2)^2 dx, x, \sec(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int (-1+x)^2 x^3 dx, x, \sec^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int (x^3 - 2x^4 + x^5) dx, x, \sec^2(a+bx)\right)}{2b} \\
&= \frac{\sec^8(a+bx)}{8b} - \frac{\sec^{10}(a+bx)}{5b} + \frac{\sec^{12}(a+bx)}{12b}
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 38, normalized size = 0.83

$$\frac{15 \sec^8(a+bx) - 24 \sec^{10}(a+bx) + 10 \sec^{12}(a+bx)}{120b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sec[a + b*x]^8*Tan[a + b*x]^5,x]``[Out] (15*Sec[a + b*x]^8 - 24*Sec[a + b*x]^10 + 10*Sec[a + b*x]^12)/(120*b)`**Maple [A]**

time = 0.06, size = 78, normalized size = 1.70

method	result	size
risch	$\frac{32 e^{16i(bx+a)} - \frac{384 e^{14i(bx+a)}}{5} + \frac{1856 e^{12i(bx+a)}}{15} - \frac{384 e^{10i(bx+a)}}{5} + 32 e^{8i(bx+a)}}{b(e^{2i(bx+a)}+1)^{12}}$	75
derivativedivides	$\frac{\frac{\sin^6(bx+a)}{12 \cos(bx+a)^{12}} + \frac{\sin^6(bx+a)}{20 \cos(bx+a)^{10}} + \frac{\sin^6(bx+a)}{40 \cos(bx+a)^8} + \frac{\sin^6(bx+a)}{120 \cos(bx+a)^6}}{b}$	78
default	$\frac{\frac{\sin^6(bx+a)}{12 \cos(bx+a)^{12}} + \frac{\sin^6(bx+a)}{20 \cos(bx+a)^{10}} + \frac{\sin^6(bx+a)}{40 \cos(bx+a)^8} + \frac{\sin^6(bx+a)}{120 \cos(bx+a)^6}}{b}$	78

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^13*sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/12*sin(b*x+a)^6/cos(b*x+a)^12+1/20*sin(b*x+a)^6/cos(b*x+a)^10+1/40*sin(b*x+a)^6/cos(b*x+a)^8+1/120*sin(b*x+a)^6/cos(b*x+a)^6)`**Maxima [B]** Leaf count of result is larger than twice the leaf count of optimal. 89 vs. 2(40) = 80.

time = 0.29, size = 89, normalized size = 1.93

$$\frac{15 \sin(bx+a)^4 - 6 \sin(bx+a)^2 + 1}{120 (\sin(bx+a)^{12} - 6 \sin(bx+a)^{10} + 15 \sin(bx+a)^8 - 20 \sin(bx+a)^6 + 15 \sin(bx+a)^4 - 6 \sin(bx+a)^2 + 1)b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^13*sin(b*x+a)^5,x, algorithm="maxima")

[Out] 1/120*(15*sin(b*x + a)^4 - 6*sin(b*x + a)^2 + 1)/((sin(b*x + a)^12 - 6*sin(b*x + a)^10 + 15*sin(b*x + a)^8 - 20*sin(b*x + a)^6 + 15*sin(b*x + a)^4 - 6*sin(b*x + a)^2 + 1)*b)

Fricas [A]

time = 0.35, size = 35, normalized size = 0.76

$$\frac{15 \cos (bx+a)^4 - 24 \cos (bx+a)^2 + 10}{120 b \cos (bx+a)^{12}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^13*sin(b*x+a)^5,x, algorithm="fricas")

[Out] 1/120*(15*cos(b*x + a)^4 - 24*cos(b*x + a)^2 + 10)/(b*cos(b*x + a)^12)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**13*sin(b*x+a)**5,x)

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 183 vs. 2(40) = 80.

time = 4.23, size = 183, normalized size = 3.98

$$\frac{32 \left(\frac{5(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{15(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + \frac{39(\cos(bx+a)-1)^5}{(\cos(bx+a)+1)^5} - \frac{42(\cos(bx+a)-1)^6}{(\cos(bx+a)+1)^6} + \frac{39(\cos(bx+a)-1)^7}{(\cos(bx+a)+1)^7} - \frac{15(\cos(bx+a)-1)^8}{(\cos(bx+a)+1)^8} + \frac{5(\cos(bx+a)-1)^9}{(\cos(bx+a)+1)^9} \right)}{15 b \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^{12}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^13*sin(b*x+a)^5,x, algorithm="giac")

[Out] -32/15*(5*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 - 15*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 + 39*(cos(b*x + a) - 1)^5/(cos(b*x + a) + 1)^5 - 42*(cos(b*x + a) - 1)^6/(cos(b*x + a) + 1)^6 + 39*(cos(b*x + a) - 1)^7/(cos(b*x + a) + 1)^7 - 15*(cos(b*x + a) - 1)^8/(cos(b*x + a) + 1)^8 + 5*(cos(b*x + a) - 1)^9/(cos(b*x + a) + 1)^9)/(b*((cos(b*x + a) - 1)/(cos(b*x + a) + 1))^12)

Mupad [B]

time = 0.42, size = 45, normalized size = 0.98

$$\frac{\frac{\tan(a+bx)^{12}}{12} + \frac{3 \tan(a+bx)^{10}}{10} + \frac{3 \tan(a+bx)^8}{8} + \frac{\tan(a+bx)^6}{6}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5/cos(a + b*x)^13,x)`

[Out] `(tan(a + b*x)^6/6 + (3*tan(a + b*x)^8)/8 + (3*tan(a + b*x)^10)/10 + tan(a + b*x)^12/12)/b`

3.118 $\int \sin^3(a + bx) \tan^3(a + bx) dx$

Optimal. Leaf size=66

$$-\frac{5 \tanh^{-1}(\sin(a + bx))}{2b} + \frac{5 \sin(a + bx)}{2b} + \frac{5 \sin^3(a + bx)}{6b} + \frac{\sin^3(a + bx) \tan^2(a + bx)}{2b}$$

[Out] $-5/2*\operatorname{arctanh}(\sin(b*x+a))/b+5/2*\sin(b*x+a)/b+5/6*\sin(b*x+a)^3/b+1/2*\sin(b*x+a)^3*\tan(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2672, 294, 308, 212}

$$\frac{5 \sin^3(a + bx)}{6b} + \frac{5 \sin(a + bx)}{2b} + \frac{\sin^3(a + bx) \tan^2(a + bx)}{2b} - \frac{5 \tanh^{-1}(\sin(a + bx))}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3*Tan[a + b*x]^3,x]

[Out] $(-5*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]])/(2*b) + (5*\operatorname{Sin}[a + b*x])/(2*b) + (5*\operatorname{Sin}[a + b*x]^3)/(6*b) + (\operatorname{Sin}[a + b*x]^3*\operatorname{Tan}[a + b*x]^2)/(2*b)$

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 294

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 308

Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2672

Int[((a_)*sin[(e_) + (f_)*(x_)])^(m_)*tan[(e_) + (f_)*(x_)]^(n_), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(

```
ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x]
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \sin^3(a + bx) \tan^3(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^6}{(1-x^2)^2} dx, x, \sin(a + bx)\right)}{b} \\
&= \frac{\sin^3(a + bx) \tan^2(a + bx)}{2b} - \frac{5\text{Subst}\left(\int \frac{x^4}{1-x^2} dx, x, \sin(a + bx)\right)}{2b} \\
&= \frac{\sin^3(a + bx) \tan^2(a + bx)}{2b} - \frac{5\text{Subst}\left(\int \left(-1 - x^2 + \frac{1}{1-x^2}\right) dx, x, \sin(a + bx)\right)}{2b} \\
&= \frac{5 \sin(a + bx)}{2b} + \frac{5 \sin^3(a + bx)}{6b} + \frac{\sin^3(a + bx) \tan^2(a + bx)}{2b} - \frac{5\text{Subst}\left(\int \frac{1}{1-x^2}\right)}{2b} \\
&= -\frac{5 \tanh^{-1}(\sin(a + bx))}{2b} + \frac{5 \sin(a + bx)}{2b} + \frac{5 \sin^3(a + bx)}{6b} + \frac{\sin^3(a + bx) \tan^2(a + bx)}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.14, size = 52, normalized size = 0.79

$$\frac{-60 \tanh^{-1}(\sin(a + bx)) + (37 + 24 \cos(2(a + bx)) - \cos(4(a + bx))) \sec(a + bx) \tan(a + bx)}{24b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^3*Tan[a + b*x]^3,x]
```

```
[Out] (-60*ArcTanh[Sin[a + b*x]] + (37 + 24*Cos[2*(a + b*x)] - Cos[4*(a + b*x)])*
Sec[a + b*x]*Tan[a + b*x])/(24*b)
```

Maple [A]

time = 0.08, size = 68, normalized size = 1.03

method	result
derivativedivides	$\frac{\frac{\sin^7(bx+a)}{2 \cos(bx+a)^2} + \frac{(\sin^5(bx+a))}{2} + \frac{5(\sin^3(bx+a))}{6} + \frac{5 \sin(bx+a)}{2} - \frac{5 \ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$
default	$\frac{\frac{\sin^7(bx+a)}{2 \cos(bx+a)^2} + \frac{(\sin^5(bx+a))}{2} + \frac{5(\sin^3(bx+a))}{6} + \frac{5 \sin(bx+a)}{2} - \frac{5 \ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$
risch	$\frac{ie^{3i(bx+a)}}{24b} - \frac{9ie^{i(bx+a)}}{8b} + \frac{9ie^{-i(bx+a)}}{8b} - \frac{ie^{-3i(bx+a)}}{24b} - \frac{i(e^{3i(bx+a)} - e^{i(bx+a)})}{b(e^{2i(bx+a)} + 1)^2} - \frac{5 \ln(e^{i(bx+a)} + i)}{2b} + \frac{5 \ln(e^{i(bx+a)} - i)}{2b}$

norman	$\frac{\frac{5 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{b} + \frac{20\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} - \frac{22\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} + \frac{20\left(\tan^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} + \frac{5\left(\tan^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^2} + \frac{5 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{2b}$
--------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^3*sin(b*x+a)^6,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b} \left(\frac{1}{2} \sin(bx+a)^7 / \cos(bx+a)^2 + \frac{1}{2} \sin(bx+a)^5 + \frac{5}{6} \sin(bx+a)^3 + \frac{5}{2} \sin(bx+a) - \frac{5}{2} \ln(\sec(bx+a) + \tan(bx+a)) \right)$

Maxima [A]

time = 0.28, size = 66, normalized size = 1.00

$$\frac{4 \sin(bx+a)^3 - \frac{6 \sin(bx+a)}{\sin(bx+a)^2 - 1} - 15 \log(\sin(bx+a) + 1) + 15 \log(\sin(bx+a) - 1) + 24 \sin(bx+a)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3*sin(b*x+a)^6,x, algorithm="maxima")`

[Out] $\frac{1}{12} \left(4 \sin(bx+a)^3 - 6 \sin(bx+a) / (\sin(bx+a)^2 - 1) - 15 \log(\sin(bx+a) + 1) + 15 \log(\sin(bx+a) - 1) + 24 \sin(bx+a) \right) / b$

Fricas [A]

time = 0.39, size = 84, normalized size = 1.27

$$\frac{15 \cos(bx+a)^2 \log(\sin(bx+a) + 1) - 15 \cos(bx+a)^2 \log(-\sin(bx+a) + 1) + 2(2 \cos(bx+a)^4 - 14 \cos(bx+a)^2 - 3) \sin(bx+a)}{12b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3*sin(b*x+a)^6,x, algorithm="fricas")`

[Out] $\frac{-1}{12} \left(15 \cos(bx+a)^2 \log(\sin(bx+a) + 1) - 15 \cos(bx+a)^2 \log(-\sin(bx+a) + 1) + 2(2 \cos(bx+a)^4 - 14 \cos(bx+a)^2 - 3) \sin(bx+a) \right) / (b \cos(bx+a)^2)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**3*sin(b*x+a)**6,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [A]

time = 4.08, size = 68, normalized size = 1.03

$$\frac{4 \sin(bx+a)^3 - \frac{6 \sin(bx+a)}{\sin(bx+a)^2 - 1} - 15 \log(|\sin(bx+a) + 1|) + 15 \log(|\sin(bx+a) - 1|) + 24 \sin(bx+a)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*sin(b*x+a)^6,x, algorithm="giac")

[Out] $\frac{1}{12}*(4*\sin(b*x + a)^3 - 6*\sin(b*x + a)/(\sin(b*x + a)^2 - 1) - 15*\log(\text{abs}(\sin(b*x + a) + 1)) + 15*\log(\text{abs}(\sin(b*x + a) - 1)) + 24*\sin(b*x + a))/b$

Mupad [B]

time = 7.24, size = 147, normalized size = 2.23

$$\frac{5 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^9 + \frac{20 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^7}{3} - \frac{22 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^5}{3} + \frac{20 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^3}{3} + 5 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{10} + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 - 2 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 - 2 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1 \right)} - \frac{5 \operatorname{atanh}\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^6/cos(a + b*x)^3,x)

[Out] $\frac{(5*\tan(a/2 + (b*x)/2) + (20*\tan(a/2 + (b*x)/2)^3)/3 - (22*\tan(a/2 + (b*x)/2)^5)/3 + (20*\tan(a/2 + (b*x)/2)^7)/3 + 5*\tan(a/2 + (b*x)/2)^9)/(b*(\tan(a/2 + (b*x)/2)^2 - 2*\tan(a/2 + (b*x)/2)^4 - 2*\tan(a/2 + (b*x)/2)^6 + \tan(a/2 + (b*x)/2)^8 + \tan(a/2 + (b*x)/2)^{10} + 1)) - (5*\operatorname{atanh}(\tan(a/2 + (b*x)/2)))}{b}$

3.119 $\int \sin(a + bx) \tan^6(a + bx) dx$

Optimal. Leaf size=50

$$\frac{\cos(a + bx)}{b} + \frac{3 \sec(a + bx)}{b} - \frac{\sec^3(a + bx)}{b} + \frac{\sec^5(a + bx)}{5b}$$

[Out] $\cos(b*x+a)/b+3*\sec(b*x+a)/b-\sec(b*x+a)^3/b+1/5*\sec(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 276}

$$\frac{\cos(a + bx)}{b} + \frac{\sec^5(a + bx)}{5b} - \frac{\sec^3(a + bx)}{b} + \frac{3 \sec(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[a + b*x]*\text{Tan}[a + b*x]^6, x]$

[Out] $\text{Cos}[a + b*x]/b + (3*\text{Sec}[a + b*x])/b - \text{Sec}[a + b*x]^3/b + \text{Sec}[a + b*x]^5/(5*b)$

Rule 276

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_) + (b_*)*(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2670

$\text{Int}[\text{sin}[(e_*) + (f_*)*(x_)]^{(m_*)}*\text{tan}[(e_*) + (f_*)*(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-f^{(-1)}, \text{Subst}[\text{Int}[(1 - x^2)^{(m+n-1)/2}/x^n, x], x, \text{Cos}[e + f*x]], x] /; \text{FreeQ}\{e, f, x\} \ \&\& \ \text{IntegersQ}[m, n, (m+n-1)/2]$

Rubi steps

$$\begin{aligned} \int \sin(a + bx) \tan^6(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^3}{x^6} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-1 + \frac{1}{x^6} - \frac{3}{x^4} + \frac{3}{x^2}\right) dx, x, \cos(a + bx)\right)}{b} \\ &= \frac{\cos(a + bx)}{b} + \frac{3 \sec(a + bx)}{b} - \frac{\sec^3(a + bx)}{b} + \frac{\sec^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 50, normalized size = 1.00

$$\frac{\cos(a + bx)}{b} + \frac{3 \sec(a + bx)}{b} - \frac{\sec^3(a + bx)}{b} + \frac{\sec^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]*Tan[a + b*x]^6,x]``[Out] Cos[a + b*x]/b + (3*Sec[a + b*x])/b - Sec[a + b*x]^3/b + Sec[a + b*x]^5/(5*b)`**Maple [A]**

time = 0.09, size = 96, normalized size = 1.92

method	result	size
norman	$\frac{-\frac{32}{5b} + \frac{128 \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 32 \left(\tan^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{5b \left(1 + \tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) - 1 \right)^5}$	70
derivativedivides	$\frac{\frac{\sin^8(bx+a)}{5 \cos(bx+a)^5} - \frac{\sin^8(bx+a)}{5 \cos(bx+a)^3} + \frac{\sin^8(bx+a)}{\cos(bx+a)} + \left(\frac{16}{5} + \sin^6(bx+a) + \frac{6 \left(\sin^4(bx+a) \right)}{5} + \frac{8 \left(\sin^2(bx+a) \right)}{5} \right) \cos(bx+a)}{b}$	96
default	$\frac{\frac{\sin^8(bx+a)}{5 \cos(bx+a)^5} - \frac{\sin^8(bx+a)}{5 \cos(bx+a)^3} + \frac{\sin^8(bx+a)}{\cos(bx+a)} + \left(\frac{16}{5} + \sin^6(bx+a) + \frac{6 \left(\sin^4(bx+a) \right)}{5} + \frac{8 \left(\sin^2(bx+a) \right)}{5} \right) \cos(bx+a)}{b}$	96
risch	$\frac{e^{i(bx+a)}}{2b} + \frac{e^{-i(bx+a)}}{2b} + \frac{6e^{9i(bx+a)} + 16e^{7i(bx+a)} + \frac{132e^{5i(bx+a)}}{5} + 16e^{3i(bx+a)} + 6e^{i(bx+a)}}{b(e^{2i(bx+a)} + 1)^5}$	104

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^6*sin(b*x+a)^7,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/5*sin(b*x+a)^8/cos(b*x+a)^5-1/5*sin(b*x+a)^8/cos(b*x+a)^3+sin(b*x+a)^8/cos(b*x+a)+(16/5+sin(b*x+a)^6+6/5*sin(b*x+a)^4+8/5*sin(b*x+a)^2)*cos(b*x+a))`**Maxima [A]**

time = 0.27, size = 45, normalized size = 0.90

$$\frac{\frac{15 \cos(bx+a)^4 - 5 \cos(bx+a)^2 + 1}{\cos(bx+a)^5} + 5 \cos(bx+a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6*sin(b*x+a)^7,x, algorithm="maxima")``[Out] 1/5*((15*cos(b*x + a)^4 - 5*cos(b*x + a)^2 + 1)/cos(b*x + a)^5 + 5*cos(b*x + a))/b`

Fricas [A]

time = 0.39, size = 45, normalized size = 0.90

$$\frac{5 \cos(bx + a)^6 + 15 \cos(bx + a)^4 - 5 \cos(bx + a)^2 + 1}{5 b \cos(bx + a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^7,x, algorithm="fricas")**[Out]** 1/5*(5*cos(b*x + a)^6 + 15*cos(b*x + a)^4 - 5*cos(b*x + a)^2 + 1)/(b*cos(b*x + a)^5)**Sympy [F(-1)]** Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**6*sin(b*x+a)**7,x)**[Out]** Timed out**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(48) = 96.

time = 3.38, size = 144, normalized size = 2.88

$$\frac{2 \left(\frac{5}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1}-1} - \frac{50 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 80 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^2 + 30 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^3 + \frac{5 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^4 + 11}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right)^5} + 11 \right)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6*sin(b*x+a)^7,x, algorithm="giac")**[Out]** -2/5*(5/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1) - (50*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 80*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 30*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 + 5*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 + 11)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^5)/b**Mupad [B]**

time = 0.54, size = 50, normalized size = 1.00

$$\frac{\cos(a + bx)}{b} + \frac{3}{b \cos(a + bx)} - \frac{1}{b \cos(a + bx)^3} + \frac{1}{5 b \cos(a + bx)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^7/cos(a + b*x)^6,x)**[Out]** cos(a + b*x)/b + 3/(b*cos(a + b*x)) - 1/(b*cos(a + b*x)^3) + 1/(5*b*cos(a + b*x)^5)

3.120 $\int \cos^5(a + bx) \cot(a + bx) dx$

Optimal. Leaf size=53

$$-\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\cos(a + bx)}{b} + \frac{\cos^3(a + bx)}{3b} + \frac{\cos^5(a + bx)}{5b}$$

[Out] $-\operatorname{arctanh}(\cos(b*x+a))/b + \cos(b*x+a)/b + 1/3*\cos(b*x+a)^3/b + 1/5*\cos(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2672, 308, 212}

$$\frac{\cos^5(a + bx)}{5b} + \frac{\cos^3(a + bx)}{3b} + \frac{\cos(a + bx)}{b} - \frac{\tanh^{-1}(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^5*Cot[a + b*x],x]`

[Out] $-(\operatorname{ArcTanh}[\cos[a + b*x]]/b) + \cos[a + b*x]/b + \cos[a + b*x]^3/(3*b) + \cos[a + b*x]^5/(5*b)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 308

`Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]`

Rule 2672

`Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]`

Rubi steps

$$\begin{aligned}
\int \cos^5(a+bx) \cot(a+bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{1-x^2} dx, x, \cos(a+bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int (-1-x^2-x^4+\frac{1}{1-x^2}) dx, x, \cos(a+bx)\right)}{b} \\
&= \frac{\cos(a+bx)}{b} + \frac{\cos^3(a+bx)}{3b} + \frac{\cos^5(a+bx)}{5b} - \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(a+bx)\right)}{b} \\
&= -\frac{\tanh^{-1}(\cos(a+bx))}{b} + \frac{\cos(a+bx)}{b} + \frac{\cos^3(a+bx)}{3b} + \frac{\cos^5(a+bx)}{5b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 75, normalized size = 1.42

$$\frac{11 \cos(a+bx)}{8b} + \frac{7 \cos(3(a+bx))}{48b} + \frac{\cos(5(a+bx))}{80b} - \frac{\log(\cos(\frac{1}{2}(a+bx)))}{b} + \frac{\log(\sin(\frac{1}{2}(a+bx)))}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^5*Cot[a + b*x], x]`

```
[Out] (11*Cos[a + b*x])/(8*b) + (7*Cos[3*(a + b*x)])/(48*b) + Cos[5*(a + b*x)]/(80*b) - Log[Cos[(a + b*x)/2]]/b + Log[Sin[(a + b*x)/2]]/b
```

Maple [A]

time = 0.06, size = 48, normalized size = 0.91

method	result	size
derivativedivides	$\frac{\frac{\cos^5(bx+a)}{5} + \frac{\cos^3(bx+a)}{3} + \cos(bx+a) + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	48
default	$\frac{\frac{\cos^5(bx+a)}{5} + \frac{\cos^3(bx+a)}{3} + \cos(bx+a) + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	48
risch	$\frac{11 e^{i(bx+a)}}{16b} + \frac{11 e^{-i(bx+a)}}{16b} - \frac{\ln(e^{i(bx+a)}+1)}{b} + \frac{\ln(e^{i(bx+a)}-1)}{b} + \frac{\cos(5bx+5a)}{80b} + \frac{7 \cos(3bx+3a)}{48b}$	91
norman	$\frac{\frac{6(\tan^8(\frac{bx+a}{2}))}{b} + \frac{46}{15b} + \frac{12(\tan^6(\frac{bx+a}{2}))}{b} + \frac{28(\tan^2(\frac{bx+a}{2}))}{3b} + \frac{56(\tan^4(\frac{bx+a}{2}))}{3b}}{(1+\tan^2(\frac{bx+a}{2}))^5} + \frac{\ln(\tan(\frac{bx+a}{2}))}{b}$	102

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^6/sin(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/5*cos(b*x+a)^5+1/3*cos(b*x+a)^3+cos(b*x+a)+ln(csc(b*x+a)-cot(b*x+a)))
```

Maxima [A]

time = 0.28, size = 56, normalized size = 1.06

$$\frac{6 \cos (bx+a)^5+10 \cos (bx+a)^3+30 \cos (bx+a)-15 \log (\cos (bx+a)+1)+15 \log (\cos (bx+a)-1)}{30 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a),x, algorithm="maxima")

[Out] 1/30*(6*cos(b*x + a)^5 + 10*cos(b*x + a)^3 + 30*cos(b*x + a) - 15*log(cos(b*x + a) + 1) + 15*log(cos(b*x + a) - 1))/b

Fricas [A]

time = 0.38, size = 60, normalized size = 1.13

$$\frac{6 \cos (bx+a)^5+10 \cos (bx+a)^3+30 \cos (bx+a)-15 \log \left(\frac{1}{2} \cos (bx+a)+\frac{1}{2}\right)+15 \log \left(-\frac{1}{2} \cos (bx+a)+\frac{1}{2}\right)}{30 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a),x, algorithm="fricas")

[Out] 1/30*(6*cos(b*x + a)^5 + 10*cos(b*x + a)^3 + 30*cos(b*x + a) - 15*log(1/2*cos(b*x + a) + 1/2) + 15*log(-1/2*cos(b*x + a) + 1/2))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 1085 vs. 2(41) = 82.

time = 2.40, size = 1085, normalized size = 20.47

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**6/sin(b*x+a),x)

[Out] Piecewise(((15*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**10/(15*b*tan(a/2 + b*x/2)**10 + 75*b*tan(a/2 + b*x/2)**8 + 150*b*tan(a/2 + b*x/2)**6 + 150*b*tan(a/2 + b*x/2)**4 + 75*b*tan(a/2 + b*x/2)**2 + 15*b) + 75*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(15*b*tan(a/2 + b*x/2)**10 + 75*b*tan(a/2 + b*x/2)**8 + 150*b*tan(a/2 + b*x/2)**6 + 150*b*tan(a/2 + b*x/2)**4 + 75*b*tan(a/2 + b*x/2)**2 + 15*b) + 150*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(15*b*tan(a/2 + b*x/2)**10 + 75*b*tan(a/2 + b*x/2)**8 + 150*b*tan(a/2 + b*x/2)**6 + 150*b*tan(a/2 + b*x/2)**4 + 75*b*tan(a/2 + b*x/2)**2 + 15*b) + 150*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(15*b*tan(a/2 + b*x/2)**10 + 75*b*tan(a/2 + b*x/2)**8 + 150*b*tan(a/2 + b*x/2)**6 + 150*b*tan(a/2 + b*x/2)**4 + 75*b*tan(a/2 + b*x/2)**2 + 15*b) + 75*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(15*b*tan(a/2 + b*x/2)**10 + 75*b*tan(a/2 + b*x/2)**8 + 150*b*tan(a/2 + b*x/2)**6 + 150*b*tan(a/2 + b*x/2)**4 + 75*b*tan(a/2 + b*x/2)**2 + 15*b) + 1

$5 \cdot \log(\tan(a/2 + b \cdot x/2)) / (15 \cdot b \cdot \tan(a/2 + b \cdot x/2)^{10} + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^8 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^6 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^4 + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^2 + 15 \cdot b) + 90 \cdot \tan(a/2 + b \cdot x/2)^8 / (15 \cdot b \cdot \tan(a/2 + b \cdot x/2)^{10} + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^8 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^6 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^4 + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^2 + 15 \cdot b) + 180 \cdot \tan(a/2 + b \cdot x/2)^6 / (15 \cdot b \cdot \tan(a/2 + b \cdot x/2)^{10} + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^8 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^6 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^4 + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^2 + 15 \cdot b) + 280 \cdot \tan(a/2 + b \cdot x/2)^4 / (15 \cdot b \cdot \tan(a/2 + b \cdot x/2)^{10} + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^8 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^6 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^4 + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^2 + 15 \cdot b) + 140 \cdot \tan(a/2 + b \cdot x/2)^2 / (15 \cdot b \cdot \tan(a/2 + b \cdot x/2)^{10} + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^8 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^6 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^4 + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^2 + 15 \cdot b) + 46 / (15 \cdot b \cdot \tan(a/2 + b \cdot x/2)^{10} + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^8 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^6 + 150 \cdot b \cdot \tan(a/2 + b \cdot x/2)^4 + 75 \cdot b \cdot \tan(a/2 + b \cdot x/2)^2 + 15 \cdot b), \text{Ne}(b, 0)), (x \cdot \cos(a))^6 / \sin(a), \text{True})$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 145 vs. 2(49) = 98.

time = 3.49, size = 145, normalized size = 2.74

$$\frac{4 \left(\frac{70(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{140(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{90(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{45(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - 23 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right)^5} + 15 \log \left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right)$$

30 b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a),x, algorithm="giac")

[Out] $\frac{1}{30} \cdot (4 \cdot (70 \cdot (\cos(b \cdot x + a) - 1) / (\cos(b \cdot x + a) + 1) - 140 \cdot (\cos(b \cdot x + a) - 1)^2 / (\cos(b \cdot x + a) + 1)^2 + 90 \cdot (\cos(b \cdot x + a) - 1)^3 / (\cos(b \cdot x + a) + 1)^3 - 45 \cdot (\cos(b \cdot x + a) - 1)^4 / (\cos(b \cdot x + a) + 1)^4 - 23) / ((\cos(b \cdot x + a) - 1) / (\cos(b \cdot x + a) + 1) - 1)^5 + 15 \cdot \log(\text{abs}(-\cos(b \cdot x + a) + 1) / \text{abs}(\cos(b \cdot x + a) + 1))) / b$

Mupad [B]

time = 5.37, size = 88, normalized size = 1.66

$$\frac{\ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{b} + \frac{6 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 + 12 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + \frac{56 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{3} + \frac{28 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{3} + \frac{46}{15}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1 \right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6/sin(a + b*x),x)

[Out] $\log(\tan(a/2 + (b \cdot x)/2)) / b + ((28 \cdot \tan(a/2 + (b \cdot x)/2)^2) / 3 + (56 \cdot \tan(a/2 + (b \cdot x)/2)^4) / 3 + 12 \cdot \tan(a/2 + (b \cdot x)/2)^6 + 6 \cdot \tan(a/2 + (b \cdot x)/2)^8 + 46 / 15) / (b \cdot (\tan(a/2 + (b \cdot x)/2)^2 + 1)^5)$

3.121 $\int \cos^4(a + bx) \cot(a + bx) dx$

Optimal. Leaf size=40

$$\frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{b} + \frac{\sin^4(a + bx)}{4b}$$

[Out] $\ln(\sin(b*x+a))/b - \sin(b*x+a)^2/b + 1/4*\sin(b*x+a)^4/b$

Rubi [A]

time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2670, 272, 45}

$$\frac{\sin^4(a + bx)}{4b} - \frac{\sin^2(a + bx)}{b} + \frac{\log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^4*Cot[a + b*x],x]`

[Out] `Log[Sin[a + b*x]]/b - Sin[a + b*x]^2/b + Sin[a + b*x]^4/(4*b)`

Rule 45

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int
[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n},
x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && Le
Q[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 2670

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol]
:= Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*
x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \cos^4(a + bx) \cot(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x} dx, x, -\sin(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1-x)^2}{x} dx, x, \sin^2(a + bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(-2 + \frac{1}{x} + x\right) dx, x, \sin^2(a + bx)\right)}{2b} \\
&= \frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{b} + \frac{\sin^4(a + bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 40, normalized size = 1.00

$$\frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{b} + \frac{\sin^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^4*Cot[a + b*x],x]``[Out] Log[Sin[a + b*x]]/b - Sin[a + b*x]^2/b + Sin[a + b*x]^4/(4*b)`**Maple [A]**

time = 0.04, size = 33, normalized size = 0.82

method	result	size
derivativedivides	$\frac{(\cos^4(bx+a))}{4} + \frac{(\cos^2(bx+a))}{2} + \ln(\sin(bx+a))$	33
default	$\frac{(\cos^4(bx+a))}{4} + \frac{(\cos^2(bx+a))}{2} + \ln(\sin(bx+a))$	33
risch	$-ix + \frac{3e^{2i(bx+a)}}{16b} + \frac{3e^{-2i(bx+a)}}{16b} - \frac{2ia}{b} + \frac{\ln(e^{2i(bx+a)}-1)}{b} + \frac{\cos(4bx+4a)}{32b}$	71
norman	$\frac{-\frac{4(\tan^2(\frac{bx+a}{2}))}{b} - \frac{4(\tan^6(\frac{bx+a}{2}))}{b} - \frac{4(\tan^4(\frac{bx+a}{2}))}{b}}{(1+\tan^2(\frac{bx+a}{2}))^4} + \frac{\ln(\tan(\frac{bx+a}{2}))}{b} - \frac{\ln(1+\tan^2(\frac{bx+a}{2}))}{b}$	100

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(1/4*cos(b*x+a)^4+1/2*cos(b*x+a)^2+ln(sin(b*x+a)))`**Maxima [A]**

time = 0.31, size = 35, normalized size = 0.88

$$\frac{\sin(bx + a)^4 - 4 \sin(bx + a)^2 + 2 \log(\sin(bx + a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a),x, algorithm="maxima")

[Out] 1/4*(sin(b*x + a)^4 - 4*sin(b*x + a)^2 + 2*log(sin(b*x + a)^2))/b

Fricas [A]

time = 0.38, size = 35, normalized size = 0.88

$$\frac{\cos(bx + a)^4 + 2 \cos(bx + a)^2 + 4 \log\left(\frac{1}{2} \sin(bx + a)\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a),x, algorithm="fricas")

[Out] 1/4*(cos(b*x + a)^4 + 2*cos(b*x + a)^2 + 4*log(1/2*sin(b*x + a)))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 1086 vs. 2(31) = 62.

time = 1.52, size = 1086, normalized size = 27.15

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**5/sin(b*x+a),x)

[Out] Piecewise((-log(tan(a/2 + b*x/2)**2 + 1))*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2)**2 + 1))*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 6*log(tan(a/2 + b*x/2)**2 + 1))*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2)**2 + 1))*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2)**2 + 1)/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 4*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 6*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 4*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b)

+ b*x/2)**2 + b) - 4*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b), Ne(b, 0)), (x*cos(a)**5/sin(a), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 170 vs. 2(38) = 76.

time = 3.69, size = 170, normalized size = 4.25

$$\frac{\frac{52(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{102(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{52(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{25(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - 25}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^4} - 6 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) + 12 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right|\right)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a),x, algorithm="giac")

[Out] -1/12*((52*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 102*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 52*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 - 25*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 - 25)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)^4 - 6*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)) + 12*log(abs(-cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1))/b

Mupad [B]

time = 0.50, size = 66, normalized size = 1.65

$$\frac{\ln(\tan(a + bx))}{b} - \frac{\ln(\tan(a + bx)^2 + 1)}{2b} + \frac{\frac{\tan(a+bx)^2}{2} + \frac{3}{4}}{b(\tan(a + bx)^4 + 2\tan(a + bx)^2 + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5/sin(a + b*x),x)

[Out] log(tan(a + b*x))/b - log(tan(a + b*x)^2 + 1)/(2*b) + (tan(a + b*x)^2/2 + 3/4)/(b*(2*tan(a + b*x)^2 + tan(a + b*x)^4 + 1))

3.122 $\int \cos^3(a + bx) \cot(a + bx) dx$

Optimal. Leaf size=38

$$-\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\cos(a + bx)}{b} + \frac{\cos^3(a + bx)}{3b}$$

[Out] $-\text{arctanh}(\cos(b*x+a))/b + \cos(b*x+a)/b + 1/3*\cos(b*x+a)^3/b$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2672, 308, 212}

$$\frac{\cos^3(a + bx)}{3b} + \frac{\cos(a + bx)}{b} - \frac{\tanh^{-1}(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^3*\text{Cot}[a + b*x], x]$

[Out] $-(\text{ArcTanh}[\text{Cos}[a + b*x]]/b) + \text{Cos}[a + b*x]/b + \text{Cos}[a + b*x]^3/(3*b)$

Rule 212

$\text{Int}[(a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 308

$\text{Int}[(x_)^{(m_)} / ((a_) + (b_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{Int}[\text{PolynomialDivide}[x^m, a + b*x^n, x], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{GtQ}[m, 2*n - 1]$

Rule 2672

$\text{Int}[(a_)*\sin[(e_) + (f_)*(x_)]^{(m_)}*\tan[(e_) + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{With}\{\text{ff} = \text{FreeFactors}[\text{Sin}[e + f*x], x]\}, \text{Dist}[\text{ff}/f, \text{Subst}[\text{Int}[(\text{ff}*x)^{(m+n)} / (a^2 - \text{ff}^2*x^2)^{(n+1)/2}, x], x, a*(\text{Sin}[e + f*x]/\text{ff})], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n+1)/2]$

Rubi steps

$$\begin{aligned}
\int \cos^3(a+bx) \cot(a+bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^4}{1-x^2} dx, x, \cos(a+bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \left(-1-x^2+\frac{1}{1-x^2}\right) dx, x, \cos(a+bx)\right)}{b} \\
&= \frac{\cos(a+bx)}{b} + \frac{\cos^3(a+bx)}{3b} - \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(a+bx)\right)}{b} \\
&= -\frac{\tanh^{-1}(\cos(a+bx))}{b} + \frac{\cos(a+bx)}{b} + \frac{\cos^3(a+bx)}{3b}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 60, normalized size = 1.58

$$\frac{5 \cos(a+bx)}{4b} + \frac{\cos(3(a+bx))}{12b} - \frac{\log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*Cot[a + b*x], x]`

```
[Out] (5*Cos[a + b*x])/(4*b) + Cos[3*(a + b*x)]/(12*b) - Log[Cos[(a + b*x)/2]]/b
+ Log[Sin[(a + b*x)/2]]/b
```

Maple [A]

time = 0.05, size = 38, normalized size = 1.00

method	result	size
derivativedivides	$\frac{(\cos^3(bx+a))}{3} + \cos(bx+a) + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	38
default	$\frac{(\cos^3(bx+a))}{3} + \cos(bx+a) + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	38
norman	$\frac{4\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 4\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \frac{8}{3b} + \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	70
risch	$\frac{5e^{i(bx+a)}}{8b} + \frac{5e^{-i(bx+a)}}{8b} - \frac{\ln(e^{i(bx+a)}+1)}{b} + \frac{\ln(e^{i(bx+a)}-1)}{b} + \frac{\cos(3bx+3a)}{12b}$	77

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^4/sin(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/3*cos(b*x+a)^3+cos(b*x+a)+ln(csc(b*x+a)-cot(b*x+a)))
```

Maxima [A]

time = 0.28, size = 46, normalized size = 1.21

$$\frac{2 \cos(bx+a)^3 + 6 \cos(bx+a) - 3 \log(\cos(bx+a)+1) + 3 \log(\cos(bx+a)-1)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a),x, algorithm="maxima")

[Out] 1/6*(2*cos(b*x + a)^3 + 6*cos(b*x + a) - 3*log(cos(b*x + a) + 1) + 3*log(cos(b*x + a) - 1))/b

Fricas [A]

time = 0.38, size = 50, normalized size = 1.32

$$\frac{2 \cos (b x+a)^3+6 \cos (b x+a)-3 \log \left(\frac{1}{2} \cos (b x+a)+\frac{1}{2}\right)+3 \log \left(-\frac{1}{2} \cos (b x+a)+\frac{1}{2}\right)}{6 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a),x, algorithm="fricas")

[Out] 1/6*(2*cos(b*x + a)^3 + 6*cos(b*x + a) - 3*log(1/2*cos(b*x + a) + 1/2) + 3*log(-1/2*cos(b*x + a) + 1/2))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 473 vs. 2(29) = 58.

time = 0.97, size = 473, normalized size = 12.45

$$\left\{ \begin{array}{l} \frac{3 \log (\cos (x+\frac{a}{b})) \cos ^2(x+\frac{a}{b})}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} + \frac{9 \log (\cos (x+\frac{a}{b})) \cos ^2(x+\frac{a}{b})}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} + \frac{9 \log (\cos (x+\frac{a}{b})) \cos ^2(x+\frac{a}{b})}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} + \frac{3 \log (\cos (x+\frac{a}{b}))}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} + \frac{12 \cos ^2(x+\frac{a}{b})}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} + \frac{12 \cos ^2(x+\frac{a}{b})}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} + \frac{3}{20 \cos ^2(x+\frac{a}{b})+20 \cos (x+\frac{a}{b})+10} \end{array} \right. \text{ for } b \neq 0$$

otherwise

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**4/sin(b*x+a),x)

[Out] Piecewise(((3*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b) + 9*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b) + 9*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b) + 3*log(tan(a/2 + b*x/2))/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b) + 12*tan(a/2 + b*x/2)**4/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b) + 12*tan(a/2 + b*x/2)**2/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b) + 8/(3*b*tan(a/2 + b*x/2)**6 + 9*b*tan(a/2 + b*x/2)**4 + 9*b*tan(a/2 + b*x/2)**2 + 3*b), Ne(b, 0)), (x*cos(a)**4/sin(a), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 101 vs. 2(36) = 72.

time = 3.31, size = 101, normalized size = 2.66

$$\frac{8 \left(\frac{3(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 2 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right)^3} + 3 \log \left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right)$$

6 b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a),x, algorithm="giac")

[Out] $\frac{1}{6} * (8 * (3 * (\cos(b*x + a) - 1) / (\cos(b*x + a) + 1) - 3 * (\cos(b*x + a) - 1)^2 / (\cos(b*x + a) + 1)^2 - 2) / ((\cos(b*x + a) - 1) / (\cos(b*x + a) + 1) - 1)^3 + 3 * \log(\frac{-\cos(b*x + a) + 1}{\cos(b*x + a) + 1})) / b$

Mupad [B]

time = 1.68, size = 62, normalized size = 1.63

$$\frac{\ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{b} + \frac{4\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 + 4\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + \frac{8}{3}}{b\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 + 1\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4/sin(a + b*x),x)

[Out] $\log(\tan(a/2 + (b*x)/2))/b + (4*\tan(a/2 + (b*x)/2)^2 + 4*\tan(a/2 + (b*x)/2)^4 + 8/3)/(b*(\tan(a/2 + (b*x)/2)^2 + 1)^3)$

3.123 $\int \cos^2(a + bx) \cot(a + bx) dx$

Optimal. Leaf size=27

$$\frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{2b}$$

[Out] ln(sin(b*x+a))/b-1/2*sin(b*x+a)^2/b

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 14}

$$\frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Cot[a + b*x],x]

[Out] Log[Sin[a + b*x]]/b - Sin[a + b*x]^2/(2*b)

Rule 14

```
Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2670

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) \cot(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{1-x^2}{x} dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(\frac{1}{x} - x\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 27, normalized size = 1.00

$$\frac{\log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^2*Cot[a + b*x],x]``[Out] Log[Sin[a + b*x]]/b - Sin[a + b*x]^2/(2*b)`**Maple [A]**

time = 0.03, size = 23, normalized size = 0.85

method	result	size
derivativedivides	$\frac{\frac{\cos^2(bx+a)}{2} + \ln(\sin(bx+a))}{b}$	23
default	$\frac{\frac{\cos^2(bx+a)}{2} + \ln(\sin(bx+a))}{b}$	23
risch	$-ix + \frac{e^{2i(bx+a)}}{8b} + \frac{e^{-2i(bx+a)}}{8b} - \frac{2ia}{b} + \frac{\ln(e^{2i(bx+a)}-1)}{b}$	57
norman	$-\frac{2\left(\tan^2\left(\frac{bx+a}{2}\right)\right)}{b\left(1+\tan^2\left(\frac{bx+a}{2}\right)\right)^2} + \frac{\ln\left(\tan\left(\frac{bx+a}{2}\right)\right)}{b} - \frac{\ln\left(1+\tan^2\left(\frac{bx+a}{2}\right)\right)}{b}$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^3/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(1/2*cos(b*x+a)^2+ln(sin(b*x+a)))`**Maxima [A]**

time = 0.29, size = 25, normalized size = 0.93

$$-\frac{\sin(bx+a)^2 - \log(\sin(bx+a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^3/sin(b*x+a),x, algorithm="maxima")``[Out] -1/2*(sin(b*x + a)^2 - log(sin(b*x + a)^2))/b`**Fricas [A]**

time = 0.38, size = 25, normalized size = 0.93

$$\frac{\cos(bx+a)^2 + 2 \log\left(\frac{1}{2} \sin(bx+a)\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a),x, algorithm="fricas")

[Out] 1/2*(cos(b*x + a)^2 + 2*log(1/2*sin(b*x + a)))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 369 vs. $2(20) = 40$.

time = 0.67, size = 369, normalized size = 13.67

$$\left\{ \begin{array}{l} \frac{-\log(\tan^2(\frac{a}{2} + \frac{bx}{2}) + 1) \tan^2(\frac{a}{2} + \frac{bx}{2})}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} - \frac{2 \log(\tan^2(\frac{a}{2} + \frac{bx}{2}) + 1) \tan^2(\frac{a}{2} + \frac{bx}{2})}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} - \frac{\log(\tan^2(\frac{a}{2} + \frac{bx}{2}) + 1)}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} + \frac{\log(\tan(\frac{a}{2} + \frac{bx}{2})) \tan^2(\frac{a}{2} + \frac{bx}{2})}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} + \frac{2 \log(\tan(\frac{a}{2} + \frac{bx}{2})) \tan^2(\frac{a}{2} + \frac{bx}{2})}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} + \frac{\log(\tan(\frac{a}{2} + \frac{bx}{2}))}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} - \frac{2 \tan^2(\frac{a}{2} + \frac{bx}{2})}{b \tan^2(\frac{a}{2} + \frac{bx}{2}) + 2b \tan^2(\frac{a}{2} + \frac{bx}{2}) + b} \text{ for } b \neq 0 \\ \frac{x \cos^3(a)}{\sin(a)} \text{ otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3/sin(b*x+a),x)

[Out] Piecewise((-log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - 2*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2)**2 + 1)/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) + 2*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - 2*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b), Ne(b, 0)), (x*cos(a)**3/sin(a), True))

Giac [A]

time = 4.00, size = 25, normalized size = 0.93

$$\frac{\sin(bx + a)^2 - \log(\sin(bx + a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a),x, algorithm="giac")

[Out] -1/2*(sin(b*x + a)^2 - log(sin(b*x + a)^2))/b

Mupad [B]

time = 0.41, size = 35, normalized size = 1.30

$$\frac{\frac{\cos(a+bx)^2}{2} - \frac{\ln(\tan(a+bx)^2+1)}{2}}{b} + \ln(\tan(a+bx))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3/sin(a + b*x),x)

[Out] (log(tan(a + b*x)) - log(tan(a + b*x)^2 + 1)/2 + cos(a + b*x)^2/2)/b

3.124 $\int \cos(a + bx) \cot(a + bx) dx$

Optimal. Leaf size=23

$$-\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\cos(a + bx)}{b}$$

[Out] -arctanh(cos(b*x+a))/b+cos(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {2672, 327, 212}

$$\frac{\cos(a + bx)}{b} - \frac{\tanh^{-1}(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Cot[a + b*x],x]

[Out] -(ArcTanh[Cos[a + b*x]]/b) + Cos[a + b*x]/b

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2672

Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \cot(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \cos(a + bx)\right)}{b} \\ &= \frac{\cos(a + bx)}{b} - \frac{\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\cos(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 42, normalized size = 1.83

$$\frac{\cos(a + bx)}{b} - \frac{\log\left(\cos\left(\frac{1}{2}(a + bx)\right)\right)}{b} + \frac{\log\left(\sin\left(\frac{1}{2}(a + bx)\right)\right)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]*Cot[a + b*x],x]``[Out] Cos[a + b*x]/b - Log[Cos[(a + b*x)/2]]/b + Log[Sin[(a + b*x)/2]]/b`**Maple [A]**

time = 0.04, size = 28, normalized size = 1.22

method	result	size
derivativedivides	$\frac{\cos(bx+a)+\ln(\csc(bx+a)-\cot(bx+a))}{b}$	28
default	$\frac{\cos(bx+a)+\ln(\csc(bx+a)-\cot(bx+a))}{b}$	28
norman	$-\frac{2\left(\tan^2\left(\frac{bx+a}{2}\right)\right)}{b\left(1+\tan^2\left(\frac{bx+a}{2}\right)\right)} + \frac{\ln\left(\tan\left(\frac{bx+a}{2}\right)\right)}{b}$	47
risch	$\frac{e^{i(bx+a)}}{2b} + \frac{e^{-i(bx+a)}}{2b} + \frac{\ln(e^{i(bx+a)}-1)}{b} - \frac{\ln(e^{i(bx+a)}+1)}{b}$	63

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^2/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(cos(b*x+a)+ln(csc(b*x+a)-cot(b*x+a)))`**Maxima [A]**

time = 0.29, size = 34, normalized size = 1.48

$$\frac{2 \cos(bx + a) - \log(\cos(bx + a) + 1) + \log(\cos(bx + a) - 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a),x, algorithm="maxima")

[Out] 1/2*(2*cos(b*x + a) - log(cos(b*x + a) + 1) + log(cos(b*x + a) - 1))/b

Fricas [A]

time = 0.38, size = 38, normalized size = 1.65

$$\frac{2 \cos (b x+a)-\log \left(\frac{1}{2} \cos (b x+a)+\frac{1}{2}\right)+\log \left(-\frac{1}{2} \cos (b x+a)+\frac{1}{2}\right)}{2 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a),x, algorithm="fricas")

[Out] 1/2*(2*cos(b*x + a) - log(1/2*cos(b*x + a) + 1/2) + log(-1/2*cos(b*x + a) + 1/2))/b

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 92 vs. 2(17) = 34.

time = 0.44, size = 92, normalized size = 4.00

$$\begin{cases} \frac{\log \left(\tan \left(\frac{a}{2}+\frac{b x}{2}\right)\right) \tan ^2\left(\frac{a}{2}+\frac{b x}{2}\right)}{b \tan ^2\left(\frac{a}{2}+\frac{b x}{2}\right)+b}+\frac{\log \left(\tan \left(\frac{a}{2}+\frac{b x}{2}\right)\right)}{b \tan ^2\left(\frac{a}{2}+\frac{b x}{2}\right)+b}+\frac{2}{b \tan ^2\left(\frac{a}{2}+\frac{b x}{2}\right)+b} & \text { for } b \neq 0 \\ \frac{x \cos ^2(a)}{\sin (a)} & \text { otherwise } \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2/sin(b*x+a),x)

[Out] Piecewise((log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))/(b*tan(a/2 + b*x/2)**2 + b) + 2/(b*tan(a/2 + b*x/2)**2 + b), Ne(b, 0)), (x*cos(a)**2/sin(a), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 57 vs. 2(23) = 46.

time = 4.25, size = 57, normalized size = 2.48

$$\frac{\frac{4}{\frac{\cos (b x+a)-1}{\cos (b x+a)+1}-1}-\log \left(\frac{|-\cos (b x+a)+1|}{|\cos (b x+a)+1|}\right)}{2 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a),x, algorithm="giac")

[Out] -1/2*(4/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1) - log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.46, size = 35, normalized size = 1.52

$$\frac{2}{b \left(\tan \left(\frac{a}{2}+\frac{b x}{2}\right)^2+1\right)}+\frac{\ln \left(\tan \left(\frac{a}{2}+\frac{b x}{2}\right)\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^2/sin(a + b*x),x)
```

```
[Out] 2/(b*(tan(a/2 + (b*x)/2)^2 + 1)) + log(tan(a/2 + (b*x)/2))/b
```

3.125 $\int \cot(a + bx) dx$

Optimal. Leaf size=11

$$\frac{\log(\sin(a + bx))}{b}$$

[Out] $\ln(\sin(b*x+a))/b$

Rubi [A]

time = 0.00, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3556}

$$\frac{\log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[a + b*x], x]$

[Out] $\text{Log}[\text{Sin}[a + b*x]]/b$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\int \cot(a + bx) dx = \frac{\log(\sin(a + bx))}{b}$$

Mathematica [A]

time = 0.01, size = 19, normalized size = 1.73

$$\frac{\log(\cos(a + bx)) + \log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Cot}[a + b*x], x]$

[Out] $(\text{Log}[\text{Cos}[a + b*x]] + \text{Log}[\text{Tan}[a + b*x]])/b$

Maple [A]

time = 0.02, size = 12, normalized size = 1.09

method	result	size
derivativedivides	$\frac{\ln(\sin(bx+a))}{b}$	12
default	$\frac{\ln(\sin(bx+a))}{b}$	12
risch	$-ix - \frac{2ia}{b} + \frac{\ln(e^{2i(bx+a)}-1)}{b}$	29
norman	$\frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{\ln\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)/sin(b*x+a),x,method=_RETURNVERBOSE)`

[Out] $\ln(\sin(b*x+a))/b$

Maxima [A]

time = 0.31, size = 11, normalized size = 1.00

$$\frac{\log(\sin(bx+a))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)/sin(b*x+a),x, algorithm="maxima")`

[Out] $\log(\sin(b*x + a))/b$

Fricas [A]

time = 0.38, size = 13, normalized size = 1.18

$$\frac{\log\left(\frac{1}{2} \sin(bx+a)\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)/sin(b*x+a),x, algorithm="fricas")`

[Out] $\log(1/2*\sin(b*x + a))/b$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 17 vs. $2(8) = 16$.

time = 0.19, size = 17, normalized size = 1.55

$$\begin{cases} \frac{\log(\sin(a+bx))}{b} & \text{for } b \neq 0 \\ \frac{x \cos(a)}{\sin(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a),x)

[Out] Piecewise((log(sin(a + b*x))/b, Ne(b, 0)), (x*cos(a)/sin(a), True))

Giac [A]

time = 4.66, size = 12, normalized size = 1.09

$$\frac{\log(|\sin(bx + a)|)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a),x, algorithm="giac")

[Out] log(abs(sin(b*x + a)))/b

Mupad [B]

time = 0.41, size = 26, normalized size = 2.36

$$-\frac{\ln(\tan(a + bx)^2 + 1) - 2 \ln(\tan(a + bx))}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)/sin(a + b*x),x)

[Out] -(log(tan(a + b*x)^2 + 1) - 2*log(tan(a + b*x)))/(2*b)

3.126 $\int \csc(a + bx) \sec(a + bx) dx$

Optimal. Leaf size=11

$$\frac{\log(\tan(a + bx))}{b}$$

[Out] ln(tan(b*x+a))/b

Rubi [A]

time = 0.01, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2700, 29}

$$\frac{\log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]*Sec[a + b*x],x]

[Out] Log[Tan[a + b*x]]/b

Rule 29

Int[(x_)^(-1), x_Symbol] :> Simp[Log[x], x]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned} \int \csc(a + bx) \sec(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{1}{x} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\log(\tan(a + bx))}{b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 31 vs. 2(11) = 22.

time = 0.02, size = 31, normalized size = 2.82

$$2 \left(-\frac{\log(\cos(a + bx))}{2b} + \frac{\log(\sin(a + bx))}{2b} \right)$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[a + b*x]*Sec[a + b*x],x]
```

```
[Out] 2*(-1/2*Log[Cos[a + b*x]]/b + Log[Sin[a + b*x]]/(2*b))
```

Maple [A]

time = 0.05, size = 12, normalized size = 1.09

method	result	size
derivativedivides	$\frac{\ln(\tan(bx+a))}{b}$	12
default	$\frac{\ln(\tan(bx+a))}{b}$	12
risch	$\frac{\ln(e^{2i(bx+a)}-1)}{b} - \frac{\ln(e^{2i(bx+a)}+1)}{b}$	35
norman	$\frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{b}$	50

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(b*x+a)/sin(b*x+a),x,method=_RETURNVERBOSE)
```

```
[Out] ln(tan(b*x+a))/b
```

Maxima [B] Leaf count of result is larger than twice the leaf count of optimal. 28 vs. 2(11) = 22.

time = 0.29, size = 28, normalized size = 2.55

$$-\frac{\log(\sin(bx+a)^2-1) - \log(\sin(bx+a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)/sin(b*x+a),x, algorithm="maxima")
```

```
[Out] -1/2*(log(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2))/b
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 30 vs. 2(11) = 22.

time = 0.36, size = 30, normalized size = 2.73

$$-\frac{\log(\cos(bx+a)^2) - \log\left(-\frac{1}{4}\cos(bx+a)^2 + \frac{1}{4}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)/sin(b*x+a),x, algorithm="fricas")
```

```
[Out] -1/2*(log(cos(b*x + a)^2) - log(-1/4*cos(b*x + a)^2 + 1/4))/b
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(a + bx)}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a),x)**[Out]** Integral(sec(a + b*x)/sin(a + b*x), x)**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(11) = 22.

time = 4.11, size = 56, normalized size = 5.09

$$\frac{\log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) - 2 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a),x, algorithm="giac")**[Out]** 1/2*(log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)) - 2*log(abs(-(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)))/b**Mupad [B]**

time = 0.39, size = 11, normalized size = 1.00

$$\frac{\ln(\tan(a + bx))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)*sin(a + b*x)),x)**[Out]** log(tan(a + b*x))/b

3.127 $\int \csc(a + bx) \sec^2(a + bx) dx$

Optimal. Leaf size=23

$$-\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\sec(a + bx)}{b}$$

[Out] -arctanh(cos(b*x+a))/b+sec(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2702, 327, 213}

$$\frac{\sec(a + bx)}{b} - \frac{\tanh^{-1}(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]*Sec[a + b*x]^2,x]

[Out] -(ArcTanh[Cos[a + b*x]]/b) + Sec[a + b*x]/b

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \csc(a+bx) \sec^2(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{x^2}{-1+x^2} dx, x, \sec(a+bx)\right)}{b} \\ &= \frac{\sec(a+bx)}{b} + \frac{\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sec(a+bx)\right)}{b} \\ &= -\frac{\tanh^{-1}(\cos(a+bx))}{b} + \frac{\sec(a+bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 1.83

$$-\frac{\log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\sec(a+bx)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]*Sec[a + b*x]^2,x]``[Out] -(Log[Cos[(a + b*x)/2]]/b) + Log[Sin[(a + b*x)/2]]/b + Sec[a + b*x]/b`**Maple [A]**

time = 0.05, size = 30, normalized size = 1.30

method	result	size
derivativedivides	$\frac{\frac{1}{\cos(bx+a)} + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	30
default	$\frac{\frac{1}{\cos(bx+a)} + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	30
norman	$-\frac{2}{b\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	36
risch	$\frac{2e^{i(bx+a)}}{b(e^{2i(bx+a)}+1)} - \frac{\ln(e^{i(bx+a)}+1)}{b} + \frac{\ln(e^{i(bx+a)}-1)}{b}$	62

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^2/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(1/cos(b*x+a)+ln(csc(b*x+a)-cot(b*x+a)))`**Maxima [A]**

time = 0.30, size = 36, normalized size = 1.57

$$\frac{\frac{2}{\cos(bx+a)} - \log(\cos(bx+a) + 1) + \log(\cos(bx+a) - 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a),x, algorithm="maxima")`

[Out] $1/2*(2/\cos(b*x + a) - \log(\cos(b*x + a) + 1) + \log(\cos(b*x + a) - 1))/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(23) = 46$.

time = 0.34, size = 52, normalized size = 2.26

$$\frac{\cos(bx + a) \log\left(\frac{1}{2} \cos(bx + a) + \frac{1}{2}\right) - \cos(bx + a) \log\left(-\frac{1}{2} \cos(bx + a) + \frac{1}{2}\right) - 2}{2b \cos(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a),x, algorithm="fricas")`

[Out] $-1/2*(\cos(b*x + a)*\log(1/2*\cos(b*x + a) + 1/2) - \cos(b*x + a)*\log(-1/2*\cos(b*x + a) + 1/2) - 2)/(b*\cos(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(a + bx)}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**2/sin(b*x+a),x)`

[Out] `Integral(sec(a + b*x)**2/sin(a + b*x), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 55 vs. $2(23) = 46$.

time = 3.86, size = 55, normalized size = 2.39

$$\frac{\frac{4}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1}+1} + \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a),x, algorithm="giac")`

[Out] $1/2*(4/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1) + \log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)))/b$

Mupad [B]

time = 0.08, size = 23, normalized size = 1.00

$$-\frac{\text{atanh}(\cos(a + bx)) - \frac{1}{\cos(a + bx)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)^2*sin(a + b*x)),x)`

[Out] $-(\text{atanh}(\cos(a + b*x)) - 1/\cos(a + b*x))/b$

3.128 $\int \csc(a + bx) \sec^3(a + bx) dx$

Optimal. Leaf size=27

$$\frac{\log(\tan(a + bx))}{b} + \frac{\tan^2(a + bx)}{2b}$$

[Out] $\ln(\tan(b*x+a))/b+1/2*\tan(b*x+a)^2/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2700, 14}

$$\frac{\tan^2(a + bx)}{2b} + \frac{\log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[a + b*x]*\text{Sec}[a + b*x]^3, x]$

[Out] $\text{Log}[\text{Tan}[a + b*x]]/b + \text{Tan}[a + b*x]^2/(2*b)$

Rule 14

$\text{Int}[(u_*)((c_.)*(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /;$ FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2700

$\text{Int}[\csc[(e_.) + (f_.)*(x_)]^{(m_.)}*\sec[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[1/f, \text{Subst}[\text{Int}[(1 + x^2)^{(m + n)/2 - 1}/x^m, x], x, \text{Tan}[e + f*x]], x] /;$ FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned} \int \csc(a + bx) \sec^3(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{1+x^2}{x} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(\frac{1}{x} + x\right) dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\log(\tan(a + bx))}{b} + \frac{\tan^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 36, normalized size = 1.33

$$\frac{-2 \log(\cos(a + bx)) - 2 \log(\sin(a + bx)) - \sec^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]*Sec[a + b*x]^3,x]``[Out] -1/2*(2*Log[Cos[a + b*x]] - 2*Log[Sin[a + b*x]] - Sec[a + b*x]^2)/b`**Maple [A]**

time = 0.05, size = 23, normalized size = 0.85

method	result	size
derivativedivides	$\frac{\frac{1}{2 \cos(bx+a)^2} + \ln(\tan(bx+a))}{b}$	23
default	$\frac{\frac{1}{2 \cos(bx+a)^2} + \ln(\tan(bx+a))}{b}$	23
risch	$\frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}+1)^2} + \frac{\ln(e^{2i(bx+a)}-1)}{b} - \frac{\ln(e^{2i(bx+a)}+1)}{b}$	62
norman	$\frac{2\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b\left(\tan^2\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)^2} + \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2}+\frac{a}{2}\right)+1\right)}{b}$	81

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^3/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(1/2/cos(b*x+a)^2+ln(tan(b*x+a)))`**Maxima [A]**

time = 0.30, size = 40, normalized size = 1.48

$$\frac{\frac{1}{\sin(bx+a)^2-1} + \log(\sin(bx+a)^2-1) - \log(\sin(bx+a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3/sin(b*x+a),x, algorithm="maxima")``[Out] -1/2*(1/(sin(b*x + a)^2 - 1) + log(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2))/b`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 56 vs. 2(25) = 50.

time = 0.38, size = 56, normalized size = 2.07

$$\frac{-\cos(bx+a)^2 \log(\cos(bx+a)^2) - \cos(bx+a)^2 \log\left(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}\right) - 1}{2b \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a),x, algorithm="fricas")

[Out] $-1/2*(\cos(b*x + a)^2*\log(\cos(b*x + a)^2) - \cos(b*x + a)^2*\log(-1/4*\cos(b*x + a)^2 + 1/4) - 1)/(b*\cos(b*x + a)^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(a + bx)}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3/sin(b*x+a),x)

[Out] Integral(sec(a + b*x)**3/sin(a + b*x), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(25) = 50.

time = 3.95, size = 124, normalized size = 4.59

$$\frac{\frac{2(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 3}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right)^2} + \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) - 2 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a),x, algorithm="giac")

[Out] $1/2*((2*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 3*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 3)/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)^2 + \log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)) - 2*\log(\text{abs}(-(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)))/b$

Mupad [B]

time = 0.10, size = 35, normalized size = 1.30

$$\frac{\frac{\ln(\sin(a+bx)^2)}{2} - \ln(\cos(a+bx)) + \frac{1}{2\cos(a+bx)^2}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^3*sin(a + b*x)),x)

[Out] $(\log(\sin(a + b*x)^2)/2 - \log(\cos(a + b*x)) + 1/(2*\cos(a + b*x)^2))/b$

3.129 $\int \csc(a + bx) \sec^4(a + bx) dx$

Optimal. Leaf size=38

$$-\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b}$$

[Out] $-\text{arctanh}(\cos(b*x+a))/b + \sec(b*x+a)/b + 1/3*\sec(b*x+a)^3/b$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2702, 308, 213}

$$\frac{\sec^3(a + bx)}{3b} + \frac{\sec(a + bx)}{b} - \frac{\tanh^{-1}(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]*Sec[a + b*x]^4,x]`

[Out] $-(\text{ArcTanh}[\text{Cos}[a + b*x]]/b) + \text{Sec}[a + b*x]/b + \text{Sec}[a + b*x]^3/(3*b)$

Rule 213

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rule 308

`Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned}
\int \csc(a+bx) \sec^4(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, \sec(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(1+x^2 + \frac{1}{-1+x^2}\right) dx, x, \sec(a+bx)\right)}{b} \\
&= \frac{\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b} + \frac{\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sec(a+bx)\right)}{b} \\
&= -\frac{\tanh^{-1}(\cos(a+bx))}{b} + \frac{\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 57, normalized size = 1.50

$$-\frac{\log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]*Sec[a + b*x]^4, x]``[Out] -(Log[Cos[(a + b*x)/2]]/b) + Log[Sin[(a + b*x)/2]]/b + Sec[a + b*x]/b + Sec[a + b*x]^3/(3*b)`**Maple [A]**

time = 0.07, size = 40, normalized size = 1.05

method	result	size
derivativedivides	$\frac{\frac{1}{3 \cos(bx+a)^3} + \frac{1}{\cos(bx+a)} + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	40
default	$\frac{\frac{1}{3 \cos(bx+a)^3} + \frac{1}{\cos(bx+a)} + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	40
norman	$\frac{\frac{4 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{8}{3b} - \frac{4 \left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^3} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	70
risch	$\frac{2e^{5i(bx+a)} + \frac{20e^{3i(bx+a)}}{3} + 2e^{i(bx+a)}}{b(e^{2i(bx+a)} + 1)^3} + \frac{\ln(e^{i(bx+a)} - 1)}{b} - \frac{\ln(e^{i(bx+a)} + 1)}{b}$	87

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^4/sin(b*x+a), x, method=_RETURNVERBOSE)``[Out] 1/b*(1/3/cos(b*x+a)^3+1/cos(b*x+a)+ln(csc(b*x+a)-cot(b*x+a)))`

Maxima [A]

time = 0.31, size = 50, normalized size = 1.32

$$\frac{2 \left(\frac{3 \cos(bx+a)^2+1}{\cos(bx+a)^3} - 3 \log(\cos(bx+a)+1) + 3 \log(\cos(bx+a)-1) \right)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^4/sin(b*x+a),x, algorithm="maxima")``[Out] 1/6*(2*(3*cos(b*x + a)^2 + 1)/cos(b*x + a)^3 - 3*log(cos(b*x + a) + 1) + 3*log(cos(b*x + a) - 1))/b`**Fricas [A]**

time = 0.40, size = 67, normalized size = 1.76

$$\frac{3 \cos(bx+a)^3 \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 3 \cos(bx+a)^3 \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 6 \cos(bx+a)^2 - 2}{6b \cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^4/sin(b*x+a),x, algorithm="fricas")``[Out] -1/6*(3*cos(b*x + a)^3*log(1/2*cos(b*x + a) + 1/2) - 3*cos(b*x + a)^3*log(-1/2*cos(b*x + a) + 1/2) - 6*cos(b*x + a)^2 - 2)/(b*cos(b*x + a)^3)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(a+bx)}{\sin(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**4/sin(b*x+a),x)``[Out] Integral(sec(a + b*x)**4/sin(a + b*x), x)`**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 101 vs. 2(36) = 72.

time = 4.14, size = 101, normalized size = 2.66

$$\frac{8 \left(\frac{3(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 2 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^3} + 3 \log \left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right)$$

$6b$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^4/sin(b*x+a),x, algorithm="giac")`

[Out] $\frac{1}{6} \cdot \frac{8 \cdot (3 \cdot (\cos(bx + a) - 1) / (\cos(bx + a) + 1) + 3 \cdot (\cos(bx + a) - 1)^2 / (\cos(bx + a) + 1)^2 + 2) / ((\cos(bx + a) - 1) / (\cos(bx + a) + 1) + 1)^3 + 3 \cdot \log(\frac{-\cos(bx + a) + 1}{\cos(bx + a) + 1})}{b}$

Mupad [B]

time = 0.39, size = 33, normalized size = 0.87

$$\frac{\operatorname{atanh}(\cos(a + bx)) - \frac{\cos(a + bx)^2 + \frac{1}{3}}{\cos(a + bx)^3}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)^4*sin(a + b*x)),x)`

[Out] $-(\operatorname{atanh}(\cos(a + b*x)) - (\cos(a + b*x)^2 + 1/3)/\cos(a + b*x)^3)/b$

3.130 $\int \csc(a + bx) \sec^5(a + bx) dx$

Optimal. Leaf size=39

$$\frac{\log(\tan(a + bx))}{b} + \frac{\tan^2(a + bx)}{b} + \frac{\tan^4(a + bx)}{4b}$$

[Out] $\ln(\tan(b*x+a))/b + \tan(b*x+a)^2/b + 1/4*\tan(b*x+a)^4/b$

Rubi [A]

time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2700, 272, 45}

$$\frac{\tan^4(a + bx)}{4b} + \frac{\tan^2(a + bx)}{b} + \frac{\log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[a + b*x]*\text{Sec}[a + b*x]^5, x]$

[Out] $\text{Log}[\text{Tan}[a + b*x]]/b + \text{Tan}[a + b*x]^2/b + \text{Tan}[a + b*x]^4/(4*b)$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ $\text{FreeQ}\{a, b, c, d, n, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{IGtQ}[m, 0] \ \&\& \ (\ !\text{IntegerQ}[n] \ || \ (\text{EqQ}[c, 0] \ \&\& \ \text{LeQ}[7*m + 4*n + 4, 0]) \ || \ \text{LtQ}[9*m + 5*(n + 1), 0] \ || \ \text{GtQ}[m + n + 2, 0])$

Rule 272

$\text{Int}[(x_.)^(m_.)*((a_.) + (b_.)*(x_.)^(n_.))^(p_.), x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m + 1)/n) - 1})*(a + b*x)^p, x], x, x^n], x] /;$ $\text{FreeQ}\{a, b, m, n, p, x\} \ \&\& \ \text{IntegerQ}[\text{Simplify}[(m + 1)/n]]$

Rule 2700

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_.)]^(m_.)*\text{sec}[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol] \rightarrow \text{Dist}[1/f, \text{Subst}[\text{Int}[(1 + x^2)^{((m + n)/2 - 1)}/x^m, x], x, \text{Tan}[e + f*x]], x] /;$ $\text{FreeQ}\{e, f, x\} \ \&\& \ \text{IntegersQ}[m, n, (m + n)/2]$

Rubi steps

$$\begin{aligned}
\int \csc(a+bx) \sec^5(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{x} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x)^2}{x} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(2 + \frac{1}{x} + x\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\log(\tan(a+bx))}{b} + \frac{\tan^2(a+bx)}{b} + \frac{\tan^4(a+bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 46, normalized size = 1.18

$$\frac{4 \log(\cos(a+bx)) - 4 \log(\sin(a+bx)) - 2 \sec^2(a+bx) - \sec^4(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]*Sec[a + b*x]^5,x]``[Out] -1/4*(4*Log[Cos[a + b*x]] - 4*Log[Sin[a + b*x]] - 2*Sec[a + b*x]^2 - Sec[a + b*x]^4)/b`**Maple [A]**

time = 0.07, size = 33, normalized size = 0.85

method	result
derivativedivides	$\frac{\frac{1}{4 \cos^4(bx+a)} + \frac{1}{2 \cos^2(bx+a)^2} + \ln(\tan(bx+a))}{b}$
default	$\frac{\frac{1}{4 \cos^4(bx+a)} + \frac{1}{2 \cos^2(bx+a)^2} + \ln(\tan(bx+a))}{b}$
risch	$\frac{2e^{6i(bx+a)} + 8e^{4i(bx+a)} + 2e^{2i(bx+a)}}{b(e^{2i(bx+a)} + 1)^4} - \frac{\ln(e^{2i(bx+a)} + 1)}{b} + \frac{\ln(e^{2i(bx+a)} - 1)}{b}$
norman	$\frac{\frac{2}{3b} + \frac{4(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{4(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{2(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^4} + \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}) - 1)}{b} - \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^5/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(1/4/cos(b*x+a)^4+1/2/cos(b*x+a)^2+ln(tan(b*x+a)))`

Maxima [A]

time = 0.28, size = 65, normalized size = 1.67

$$\frac{\frac{2 \sin(bx+a)^2 - 3}{\sin(bx+a)^4 - 2 \sin(bx+a)^2 + 1} + 2 \log(\sin(bx+a)^2 - 1) - 2 \log(\sin(bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5/sin(b*x+a),x, algorithm="maxima")`

`[Out] -1/4*((2*sin(b*x + a)^2 - 3)/(sin(b*x + a)^4 - 2*sin(b*x + a)^2 + 1) + 2*log(sin(b*x + a)^2 - 1) - 2*log(sin(b*x + a)^2))/b`

Fricas [A]

time = 0.35, size = 67, normalized size = 1.72

$$\frac{2 \cos(bx+a)^4 \log(\cos(bx+a)^2) - 2 \cos(bx+a)^4 \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) - 2 \cos(bx+a)^2 - 1}{4b \cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5/sin(b*x+a),x, algorithm="fricas")`

`[Out] -1/4*(2*cos(b*x + a)^4*log(cos(b*x + a)^2) - 2*cos(b*x + a)^4*log(-1/4*cos(b*x + a)^2 + 1/4) - 2*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^4)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(a + bx)}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**5/sin(b*x+a),x)``[Out] Integral(sec(a + b*x)**5/sin(a + b*x), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 170 vs. 2(37) = 74.

time = 3.22, size = 170, normalized size = 4.36

$$\frac{\frac{52 \cos(bx+a)-1}{\cos(bx+a)+1} + \frac{102 (\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{52 (\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{25 (\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + 25}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^4} + 6 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) - 12 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5/sin(b*x+a),x, algorithm="giac")`

[Out] $\frac{1}{12} \left(\frac{52(\cos(bx + a) - 1)}{\cos(bx + a) + 1} + 102(\cos(bx + a) - 1)^2 / (\cos(bx + a) + 1)^2 + 52(\cos(bx + a) - 1)^3 / (\cos(bx + a) + 1)^3 + 25(\cos(bx + a) - 1)^4 / (\cos(bx + a) + 1)^4 + 25 \right) / \left(\frac{\cos(bx + a) - 1}{\cos(bx + a) + 1} + 1 \right)^4 + 6 \log(\text{abs}(-\cos(bx + a) + 1) / \text{abs}(\cos(bx + a) + 1)) - 12 \log(\text{abs}(-(\cos(bx + a) - 1) / (\cos(bx + a) + 1) - 1)) / b$

Mupad [B]

time = 0.41, size = 46, normalized size = 1.18

$$\frac{\frac{\ln(\sin(a+bx)^2)}{2} - \ln(\cos(a+bx)) + \frac{\frac{\cos(a+bx)^2}{2} + \frac{1}{4}}{\cos(a+bx)^4}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)^5*sin(a + b*x)),x)`

[Out] $(\log(\sin(a + b*x)^2)/2 - \log(\cos(a + b*x)) + (\cos(a + b*x)^2/2 + 1/4)/\cos(a + b*x)^4)/b$

3.131 $\int \csc(a + bx) \sec^6(a + bx) dx$

Optimal. Leaf size=53

$$-\frac{\tanh^{-1}(\cos(a + bx))}{b} + \frac{\sec(a + bx)}{b} + \frac{\sec^3(a + bx)}{3b} + \frac{\sec^5(a + bx)}{5b}$$

[Out] $-\text{arctanh}(\cos(b*x+a))/b + \sec(b*x+a)/b + 1/3*\sec(b*x+a)^3/b + 1/5*\sec(b*x+a)^5/b$

Rubi [A]

time = 0.02, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2702, 308, 213}

$$\frac{\sec^5(a + bx)}{5b} + \frac{\sec^3(a + bx)}{3b} + \frac{\sec(a + bx)}{b} - \frac{\tanh^{-1}(\cos(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]*Sec[a + b*x]^6,x]`

[Out] $-(\text{ArcTanh}[\text{Cos}[a + b*x]])/b + \text{Sec}[a + b*x]/b + \text{Sec}[a + b*x]^3/(3*b) + \text{Sec}[a + b*x]^5/(5*b)$

Rule 213

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rule 308

`Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_)), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned}
\int \csc(a+bx) \sec^6(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{x^6}{-1+x^2} dx, x, \sec(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \left(1+x^2+x^4+\frac{1}{-1+x^2}\right) dx, x, \sec(a+bx)\right)}{b} \\
&= \frac{\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b} + \frac{\sec^5(a+bx)}{5b} + \frac{\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sec(a+bx)\right)}{b} \\
&= -\frac{\tanh^{-1}(\cos(a+bx))}{b} + \frac{\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b} + \frac{\sec^5(a+bx)}{5b}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 72, normalized size = 1.36

$$-\frac{\log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{b} + \frac{\sec(a+bx)}{b} + \frac{\sec^3(a+bx)}{3b} + \frac{\sec^5(a+bx)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]*Sec[a + b*x]^6, x]`

```
[Out] -(Log[Cos[(a + b*x)/2]]/b) + Log[Sin[(a + b*x)/2]]/b + Sec[a + b*x]/b + Sec[a + b*x]^3/(3*b) + Sec[a + b*x]^5/(5*b)
```

Maple [A]

time = 0.08, size = 50, normalized size = 0.94

method	result	size
derivativedivides	$\frac{\frac{1}{5 \cos(bx+a)^5} + \frac{1}{3 \cos(bx+a)^3} + \frac{1}{\cos(bx+a)} + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	50
default	$\frac{\frac{1}{5 \cos(bx+a)^5} + \frac{1}{3 \cos(bx+a)^3} + \frac{1}{\cos(bx+a)} + \ln(\csc(bx+a) - \cot(bx+a))}{b}$	50
norman	$\frac{-\frac{46}{15b} + \frac{12 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{6 \left(\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{28 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b} - \frac{56 \left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{3b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^5} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	102
risch	$\frac{2 e^{9i(bx+a)} + 32 e^{7i(bx+a)} + 356 e^{5i(bx+a)} + 32 e^{3i(bx+a)} + 2 e^{i(bx+a)}}{b(e^{2i(bx+a)} + 1)^5} - \frac{\ln(e^{i(bx+a)} + 1)}{b} + \frac{\ln(e^{i(bx+a)} - 1)}{b}$	109

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^6/sin(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/5/cos(b*x+a)^5+1/3/cos(b*x+a)^3+1/cos(b*x+a)+ln(csc(b*x+a)-cot(b*x+a)))
```

Maxima [A]

time = 0.29, size = 60, normalized size = 1.13

$$\frac{2 \left(15 \cos(bx+a)^4 + 5 \cos(bx+a)^2 + 3 \right)}{\cos(bx+a)^5} - 15 \log(\cos(bx+a) + 1) + 15 \log(\cos(bx+a) - 1)}{30b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6/sin(b*x+a),x, algorithm="maxima")`

`[Out] 1/30*(2*(15*cos(b*x + a)^4 + 5*cos(b*x + a)^2 + 3)/cos(b*x + a)^5 - 15*log(cos(b*x + a) + 1) + 15*log(cos(b*x + a) - 1))/b`

Fricas [A]

time = 0.38, size = 77, normalized size = 1.45

$$\frac{15 \cos(bx+a)^5 \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 15 \cos(bx+a)^5 \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 30 \cos(bx+a)^4 - 10 \cos(bx+a)^2 - 6}{30b \cos(bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^6/sin(b*x+a),x, algorithm="fricas")`

`[Out] -1/30*(15*cos(b*x + a)^5*log(1/2*cos(b*x + a) + 1/2) - 15*cos(b*x + a)^5*log(-1/2*cos(b*x + a) + 1/2) - 30*cos(b*x + a)^4 - 10*cos(b*x + a)^2 - 6)/(b*cos(b*x + a)^5)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^6(a + bx)}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**6/sin(b*x+a),x)`

`[Out] Integral(sec(a + b*x)**6/sin(a + b*x), x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 145 vs. 2(49) = 98.

time = 3.24, size = 145, normalized size = 2.74

$$\frac{4 \left(\frac{70(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{140(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{90(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{45(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + 23 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^5} + 15 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)}{30b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^6/sin(b*x+a),x, algorithm="giac")

[Out] $\frac{1}{30} \cdot \left(4 \cdot \frac{70 \cdot (\cos(bx + a) - 1)}{(\cos(bx + a) + 1)} + 140 \cdot (\cos(bx + a) - 1)^2 / (\cos(bx + a) + 1)^2 + 90 \cdot (\cos(bx + a) - 1)^3 / (\cos(bx + a) + 1)^3 + 45 \cdot (\cos(bx + a) - 1)^4 / (\cos(bx + a) + 1)^4 + 23 \right) / \left((\cos(bx + a) - 1) / (\cos(bx + a) + 1) + 1 \right)^5 + 15 \cdot \log(\text{abs}(-\cos(bx + a) + 1) / \text{abs}(\cos(bx + a) + 1)) / b$

Mupad [B]

time = 0.40, size = 45, normalized size = 0.85

$$\frac{\cos(a + bx)^4 + \frac{\cos(a+bx)^2}{3} + \frac{1}{5}}{b \cos(a + bx)^5} - \frac{\text{atanh}(\cos(a + bx))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^6*sin(a + b*x)),x)

[Out] $(\cos(a + bx)^2/3 + \cos(a + bx)^4 + 1/5) / (b \cdot \cos(a + bx)^5) - \text{atanh}(\cos(a + bx)) / b$

3.132 $\int \csc(a + bx) \sec^7(a + bx) dx$

Optimal. Leaf size=57

$$\frac{\log(\tan(a + bx))}{b} + \frac{3 \tan^2(a + bx)}{2b} + \frac{3 \tan^4(a + bx)}{4b} + \frac{\tan^6(a + bx)}{6b}$$

[Out] $\ln(\tan(b*x+a))/b+3/2*\tan(b*x+a)^2/b+3/4*\tan(b*x+a)^4/b+1/6*\tan(b*x+a)^6/b$

Rubi [A]

time = 0.02, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2700, 272, 45}

$$\frac{\tan^6(a + bx)}{6b} + \frac{3 \tan^4(a + bx)}{4b} + \frac{3 \tan^2(a + bx)}{2b} + \frac{\log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]*Sec[a + b*x]^7,x]`

[Out] `Log[Tan[a + b*x]]/b + (3*Tan[a + b*x]^2)/(2*b) + (3*Tan[a + b*x]^4)/(4*b) + Tan[a + b*x]^6/(6*b)`

Rule 45

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])`

Rule 272

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]`

Rule 2700

`Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]`

Rubi steps

$$\begin{aligned}
\int \csc(a+bx) \sec^7(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^3}{x} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x)^3}{x} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(3 + \frac{1}{x} + 3x + x^2\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\log(\tan(a+bx))}{b} + \frac{3 \tan^2(a+bx)}{2b} + \frac{3 \tan^4(a+bx)}{4b} + \frac{\tan^6(a+bx)}{6b}
\end{aligned}$$

Mathematica [A]

time = 0.10, size = 56, normalized size = 0.98

$$\frac{12 \log(\cos(a+bx)) - 12 \log(\sin(a+bx)) - 6 \sec^2(a+bx) - 3 \sec^4(a+bx) - 2 \sec^6(a+bx)}{12b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]*Sec[a + b*x]^7,x]``[Out] -1/12*(12*Log[Cos[a + b*x]] - 12*Log[Sin[a + b*x]] - 6*Sec[a + b*x]^2 - 3*Sec[a + b*x]^4 - 2*Sec[a + b*x]^6)/b`**Maple [A]**

time = 0.10, size = 43, normalized size = 0.75

method	result
derivativedivides	$\frac{\frac{1}{6 \cos(bx+a)^6} + \frac{1}{4 \cos(bx+a)^4} + \frac{1}{2 \cos(bx+a)^2} + \ln(\tan(bx+a))}{b}$
default	$\frac{\frac{1}{6 \cos(bx+a)^6} + \frac{1}{4 \cos(bx+a)^4} + \frac{1}{2 \cos(bx+a)^2} + \ln(\tan(bx+a))}{b}$
risch	$\frac{2 e^{10i(bx+a)} + 12 e^{8i(bx+a)} + \frac{92 e^{6i(bx+a)}}{3} + 12 e^{4i(bx+a)} + 2 e^{2i(bx+a)}}{b(e^{2i(bx+a)}+1)^6} + \frac{\ln(e^{2i(bx+a)}-1)}{b} - \frac{\ln(e^{2i(bx+a)}+1)}{b}$
norman	$\frac{\frac{6(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{6(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{12(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{12(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{68(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{3b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2})-1)^6} + \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{b}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^7/sin(b*x+a),x,method=_RETURNVERBOSE)``[Out] 1/b*(1/6/cos(b*x+a)^6+1/4/cos(b*x+a)^4+1/2/cos(b*x+a)^2+ln(tan(b*x+a)))`

Maxima [A]

time = 0.29, size = 85, normalized size = 1.49

$$\frac{6 \sin(bx+a)^4 - 15 \sin(bx+a)^2 + 11}{\sin(bx+a)^6 - 3 \sin(bx+a)^4 + 3 \sin(bx+a)^2 - 1} + 6 \log(\sin(bx+a)^2 - 1) - 6 \log(\sin(bx+a)^2)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7/sin(b*x+a),x, algorithm="maxima")

[Out] -1/12*((6*sin(b*x + a)^4 - 15*sin(b*x + a)^2 + 11)/(sin(b*x + a)^6 - 3*sin(b*x + a)^4 + 3*sin(b*x + a)^2 - 1) + 6*log(sin(b*x + a)^2 - 1) - 6*log(sin(b*x + a)^2))/b

Fricas [A]

time = 0.39, size = 77, normalized size = 1.35

$$\frac{6 \cos(bx+a)^6 \log(\cos(bx+a)^2) - 6 \cos(bx+a)^6 \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) - 6 \cos(bx+a)^4 - 3 \cos(bx+a)^2 - 2}{12b \cos(bx+a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7/sin(b*x+a),x, algorithm="fricas")

[Out] -1/12*(6*cos(b*x + a)^6*log(cos(b*x + a)^2) - 6*cos(b*x + a)^6*log(-1/4*cos(b*x + a)^2 + 1/4) - 6*cos(b*x + a)^4 - 3*cos(b*x + a)^2 - 2)/(b*cos(b*x + a)^6)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^7(a + bx)}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**7/sin(b*x+a),x)**[Out]** Integral(sec(a + b*x)**7/sin(a + b*x), x)**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 214 vs. 2(51) = 102.

time = 3.58, size = 214, normalized size = 3.75

$$\frac{\frac{522(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{1485(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{1580(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{1485(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + \frac{522(\cos(bx+a)-1)^5}{(\cos(bx+a)+1)^5} + \frac{147(\cos(bx+a)-1)^6}{(\cos(bx+a)+1)^6} + 147}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^6} + 30 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) - 60 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}$$

60b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^7/sin(b*x+a),x, algorithm="giac")

[Out] $\frac{1}{60} * \left(\frac{522 * (\cos(b*x + a) - 1)}{(\cos(b*x + a) + 1)} + 1485 * (\cos(b*x + a) - 1)^2 / (\cos(b*x + a) + 1)^2 + 1580 * (\cos(b*x + a) - 1)^3 / (\cos(b*x + a) + 1)^3 + 1485 * (\cos(b*x + a) - 1)^4 / (\cos(b*x + a) + 1)^4 + 522 * (\cos(b*x + a) - 1)^5 / (\cos(b*x + a) + 1)^5 + 147 * (\cos(b*x + a) - 1)^6 / (\cos(b*x + a) + 1)^6 + 147 / \left(\frac{(\cos(b*x + a) - 1)}{(\cos(b*x + a) + 1)} + 1 \right)^6 + 30 * \log(\text{abs}(-\cos(b*x + a) + 1) / \text{abs}(\cos(b*x + a) + 1)) - 60 * \log(\text{abs}(-(\cos(b*x + a) - 1) / (\cos(b*x + a) + 1) - 1)) \right) / b$

Mupad [B]

time = 0.40, size = 56, normalized size = 0.98

$$\frac{\frac{\ln(\sin(a+bx)^2)}{2} - \ln(\cos(a+bx)) + \frac{\frac{\cos(a+bx)^4}{2} + \frac{\cos(a+bx)^2}{4} + \frac{1}{6}}{\cos(a+bx)^6}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)^7*sin(a + b*x)),x)`

[Out] $(\log(\sin(a + b*x)^2)/2 - \log(\cos(a + b*x)) + (\cos(a + b*x)^2/4 + \cos(a + b*x)^4/2 + 1/6)/\cos(a + b*x)^6)/b$

3.133 $\int \cos^5(a + bx) \cot^2(a + bx) dx$

Optimal. Leaf size=50

$$-\frac{\csc(a + bx)}{b} - \frac{3 \sin(a + bx)}{b} + \frac{\sin^3(a + bx)}{b} - \frac{\sin^5(a + bx)}{5b}$$

[Out] $-\csc(b*x+a)/b-3*\sin(b*x+a)/b+\sin(b*x+a)^3/b-1/5*\sin(b*x+a)^5/b$

Rubi [A]

time = 0.03, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2670, 276}

$$-\frac{\sin^5(a + bx)}{5b} + \frac{\sin^3(a + bx)}{b} - \frac{3 \sin(a + bx)}{b} - \frac{\csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^5*\text{Cot}[a + b*x]^2, x]$

[Out] $-(\text{Csc}[a + b*x]/b) - (3*\text{Sin}[a + b*x])/b + \text{Sin}[a + b*x]^3/b - \text{Sin}[a + b*x]^5/(5*b)$

Rule 276

$\text{Int}[(c_*)(x_*)^{(m_*)}((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2670

$\text{Int}[\sin[(e_*) + (f_*)(x_*)]^{(m_*)}*\tan[(e_*) + (f_*)(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-f^{-1}, \text{Subst}[\text{Int}[(1 - x^2)^{(m+n-1)/2}/x^n, x], x, \text{Cos}[e + f*x]], x] /; \text{FreeQ}\{e, f, x\} \ \&\& \ \text{IntegersQ}[m, n, (m+n-1)/2]$

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx) \cot^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^3}{x^2} dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-3 + \frac{1}{x^2} + 3x^2 - x^4\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\csc(a + bx)}{b} - \frac{3 \sin(a + bx)}{b} + \frac{\sin^3(a + bx)}{b} - \frac{\sin^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 50, normalized size = 1.00

$$-\frac{\csc(a+bx)}{b} - \frac{3\sin(a+bx)}{b} + \frac{\sin^3(a+bx)}{b} - \frac{\sin^5(a+bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^5*Cot[a + b*x]^2,x]**[Out]** -(Csc[a + b*x]/b) - (3*Sin[a + b*x])/b + Sin[a + b*x]^3/b - Sin[a + b*x]^5/(5*b)**Maple [A]**

time = 0.06, size = 62, normalized size = 1.24

method	result
derivativedivides	$\frac{-\frac{\cos^8(bx+a)}{\sin(bx+a)} - \left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)}{b}$
default	$-\frac{\cos^8(bx+a)}{\sin(bx+a)} - \left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5}\right) \sin(bx+a)$
risch	$\frac{19ie^{i(bx+a)}}{16b} - \frac{19ie^{-i(bx+a)}}{16b} - \frac{2ie^{i(bx+a)}}{b(e^{2i(bx+a)}-1)} - \frac{\sin(5bx+5a)}{80b} - \frac{3\sin(3bx+3a)}{16b}$
norman	$\frac{-\frac{1}{2b} - \frac{9(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{47(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{2b} - \frac{182(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{5b} - \frac{47(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{2b} - \frac{9(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\tan^{12}(\frac{bx}{2} + \frac{a}{2})}{2b}}{(1+\tan^2(\frac{bx}{2} + \frac{a}{2}))^5 \tan(\frac{bx}{2} + \frac{a}{2})}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^7/sin(b*x+a)^2,x,method=_RETURNVERBOSE)**[Out]** 1/b*(-1/sin(b*x+a)*cos(b*x+a)^8-(16/5+cos(b*x+a)^6+6/5*cos(b*x+a)^4+8/5*cos(b*x+a)^2)*sin(b*x+a)**Maxima [A]**

time = 0.28, size = 42, normalized size = 0.84

$$\frac{\sin(bx+a)^5 - 5\sin(bx+a)^3 + \frac{5}{\sin(bx+a)} + 15\sin(bx+a)}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^2,x, algorithm="maxima")**[Out]** -1/5*(sin(b*x + a)^5 - 5*sin(b*x + a)^3 + 5/sin(b*x + a) + 15*sin(b*x + a))/b

Fricas [A]

time = 0.34, size = 43, normalized size = 0.86

$$\frac{\cos (bx+a)^6+2 \cos (bx+a)^4+8 \cos (bx+a)^2-16}{5 b \sin (bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^2,x, algorithm="fricas")**[Out]** 1/5*(cos(b*x + a)^6 + 2*cos(b*x + a)^4 + 8*cos(b*x + a)^2 - 16)/(b*sin(b*x + a))**Sympy [B]** Leaf count of result is larger than twice the leaf count of optimal. 82 vs. 2(39) = 78.

time = 0.99, size = 82, normalized size = 1.64

$$\left\{ \begin{array}{ll} -\frac{16 \sin ^5(a+b x)}{5 b}-\frac{8 \sin ^3(a+b x) \cos ^2(a+b x)}{b}-\frac{6 \sin (a+b x) \cos ^4(a+b x)}{b}-\frac{\cos ^6(a+b x)}{b \sin (a+b x)} & \text { for } b \neq 0 \\ \frac{x \cos ^7(a)}{\sin ^2(a)} & \text { otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**7/sin(b*x+a)**2,x)**[Out]** Piecewise((-16*sin(a + b*x)**5/(5*b) - 8*sin(a + b*x)**3*cos(a + b*x)**2/b - 6*sin(a + b*x)*cos(a + b*x)**4/b - cos(a + b*x)**6/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)**7/sin(a)**2, True))**Giac [A]**

time = 3.36, size = 42, normalized size = 0.84

$$\frac{\sin (bx+a)^5-5 \sin (bx+a)^3+\frac{5}{\sin (bx+a)}+15 \sin (bx+a)}{5 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^2,x, algorithm="giac")**[Out]** -1/5*(sin(b*x + a)^5 - 5*sin(b*x + a)^3 + 5/sin(b*x + a) + 15*sin(b*x + a))/b**Mupad [B]**

time = 0.48, size = 43, normalized size = 0.86

$$\frac{\sin (a+b x)^6-5 \sin (a+b x)^4+15 \sin (a+b x)^2+5}{5 b \sin (a+b x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^7/sin(a + b*x)^2,x)**[Out]** -(15*sin(a + b*x)^2 - 5*sin(a + b*x)^4 + sin(a + b*x)^6 + 5)/(5*b*sin(a + b*x))

3.134 $\int \cos^4(a + bx) \cot^2(a + bx) dx$

Optimal. Leaf size=61

$$-\frac{15x}{8} - \frac{15 \cot(a + bx)}{8b} + \frac{5 \cos^2(a + bx) \cot(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot(a + bx)}{4b}$$

[Out] $-15/8*x-15/8*\cot(b*x+a)/b+5/8*\cos(b*x+a)^2*\cot(b*x+a)/b+1/4*\cos(b*x+a)^4*\cot(b*x+a)/b$

Rubi [A]

time = 0.03, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2671, 294, 327, 209}

$$-\frac{15 \cot(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot(a + bx)}{4b} + \frac{5 \cos^2(a + bx) \cot(a + bx)}{8b} - \frac{15x}{8}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Cot[a + b*x]^2,x]

[Out] $(-15*x)/8 - (15*\cot[a + b*x])/(8*b) + (5*\cos[a + b*x]^2*\cot[a + b*x])/(8*b) + (\cos[a + b*x]^4*\cot[a + b*x])/(4*b)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2671

```
Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol]
:> With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[Int[(ff*x)^(m + n)/(b^2 + ff^2*x^2)^(m/2 + 1), x], x, b*(Tan[e + f*x]/ff)], x]]
/; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2]
```

Rubi steps

$$\begin{aligned} \int \cos^4(a + bx) \cot^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{(1+x^2)^3} dx, x, \cot(a + bx)\right)}{b} \\ &= \frac{\cos^4(a + bx) \cot(a + bx)}{4b} - \frac{5\text{Subst}\left(\int \frac{x^4}{(1+x^2)^2} dx, x, \cot(a + bx)\right)}{4b} \\ &= \frac{5\cos^2(a + bx) \cot(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot(a + bx)}{4b} - \frac{15\text{Subst}\left(\int \frac{x^2}{1+x^2} dx, x, \cot(a + bx)\right)}{4b} \\ &= -\frac{15\cot(a + bx)}{8b} + \frac{5\cos^2(a + bx) \cot(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot(a + bx)}{4b} \\ &= -\frac{15x}{8} - \frac{15\cot(a + bx)}{8b} + \frac{5\cos^2(a + bx) \cot(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 41, normalized size = 0.67

$$-\frac{60a + 60bx + 32\cot(a + bx) + 16\sin(2(a + bx)) + \sin(4(a + bx))}{32b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4*Cot[a + b*x]^2,x]

[Out] -1/32*(60*a + 60*b*x + 32*Cot[a + b*x] + 16*Sin[2*(a + b*x)] + Sin[4*(a + b*x)])/b

Maple [A]

time = 0.04, size = 66, normalized size = 1.08

method	result
derivativedivides	$\frac{-\frac{\cos^7(bx+a)}{\sin(bx+a)} - \left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15\cos(bx+a)}{8}\right) \sin(bx+a) - \frac{15bx}{8} - \frac{15a}{8}}{b}$
default	$\frac{-\frac{\cos^7(bx+a)}{\sin(bx+a)} - \left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15\cos(bx+a)}{8}\right) \sin(bx+a) - \frac{15bx}{8} - \frac{15a}{8}}{b}$

risch	$-\frac{15x}{8} + \frac{ie^{2i(bx+a)}}{4b} - \frac{ie^{-2i(bx+a)}}{4b} - \frac{2i}{b(e^{2i(bx+a)}-1)} - \frac{\sin(4bx+4a)}{32b}$
norman	$\frac{-\frac{1}{2b} - \frac{15(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{4b} - \frac{5(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{5(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{15(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{\tan^{10}(\frac{bx}{2} + \frac{a}{2})}{2b} - \frac{15x \tan(\frac{bx}{2} + \frac{a}{2})}{8} - \frac{15x(\tan(\frac{bx}{2} + \frac{a}{2}))^2}{8}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^4 \tan(\frac{bx}{2} + \frac{a}{2})}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^6/sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/\sin(b*x+a)*\cos(b*x+a)^7-(\cos(b*x+a))^5+5/4*\cos(b*x+a)^3+15/8*\cos(b*x+a))*\sin(b*x+a)-15/8*b*x-15/8*a)$

Maxima [A]

time = 0.53, size = 63, normalized size = 1.03

$$\frac{15bx + 15a + \frac{15 \tan(bx+a)^4 + 25 \tan(bx+a)^2 + 8}{\tan(bx+a)^5 + 2 \tan(bx+a)^3 + \tan(bx+a)}}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^2,x, algorithm="maxima")`

[Out] $-1/8*(15*b*x + 15*a + (15*\tan(b*x + a)^4 + 25*\tan(b*x + a)^2 + 8)/(\tan(b*x + a)^5 + 2*\tan(b*x + a)^3 + \tan(b*x + a)))/b$

Fricas [A]

time = 0.41, size = 52, normalized size = 0.85

$$\frac{2 \cos(bx+a)^5 + 5 \cos(bx+a)^3 - 15bx \sin(bx+a) - 15 \cos(bx+a)}{8b \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^2,x, algorithm="fricas")`

[Out] $1/8*(2*\cos(b*x + a)^5 + 5*\cos(b*x + a)^3 - 15*b*x*\sin(b*x + a) - 15*\cos(b*x + a))/(b*\sin(b*x + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(54) = 108.

time = 0.75, size = 119, normalized size = 1.95

$$\begin{cases} -\frac{15x \sin^4(a+bx)}{8} - \frac{15x \sin^2(a+bx) \cos^2(a+bx)}{4} - \frac{15x \cos^4(a+bx)}{8} - \frac{15 \sin^3(a+bx) \cos(a+bx)}{8b} - \frac{25 \sin(a+bx) \cos^3(a+bx)}{8b} - \frac{\cos^5(a+bx)}{b \sin(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^6(a)}{\sin^2(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**6/sin(b*x+a)**2,x)`

[Out] Piecewise((-15*x*sin(a + b*x)**4/8 - 15*x*sin(a + b*x)**2*cos(a + b*x)**2/4 - 15*x*cos(a + b*x)**4/8 - 15*sin(a + b*x)**3*cos(a + b*x)/(8*b) - 25*sin(a + b*x)*cos(a + b*x)**3/(8*b) - cos(a + b*x)**5/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)**6/sin(a)**2, True))

Giac [A]

time = 2.92, size = 55, normalized size = 0.90

$$-\frac{15bx + 15a + \frac{7 \tan(bx+a)^3 + 9 \tan(bx+a)}{(\tan(bx+a)^2 + 1)^2} + \frac{8}{\tan(bx+a)}}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a)^2,x, algorithm="giac")

[Out] -1/8*(15*b*x + 15*a + (7*tan(b*x + a)^3 + 9*tan(b*x + a))/(tan(b*x + a)^2 + 1)^2 + 8/tan(b*x + a))/b

Mupad [B]

time = 0.59, size = 47, normalized size = 0.77

$$-\frac{15x}{8} - \frac{\cos(a + bx)^4 \left(\frac{15 \tan(a+bx)^4}{8} + \frac{25 \tan(a+bx)^2}{8} + 1 \right)}{b \tan(a + bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6/sin(a + b*x)^2,x)

[Out] - (15*x)/8 - (cos(a + b*x)^4*((25*tan(a + b*x)^2)/8 + (15*tan(a + b*x)^4)/8 + 1))/(b*tan(a + b*x))

3.135 $\int \cos^3(a + bx) \cot^2(a + bx) dx$

Optimal. Leaf size=38

$$-\frac{\csc(a + bx)}{b} - \frac{2 \sin(a + bx)}{b} + \frac{\sin^3(a + bx)}{3b}$$

[Out] $-\csc(b*x+a)/b-2*\sin(b*x+a)/b+1/3*\sin(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2670, 276}

$$\frac{\sin^3(a + bx)}{3b} - \frac{2 \sin(a + bx)}{b} - \frac{\csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[a + b*x]^3*\text{Cot}[a + b*x]^2, x]$

[Out] $-(\text{Csc}[a + b*x]/b) - (2*\text{Sin}[a + b*x])/b + \text{Sin}[a + b*x]^3/(3*b)$

Rule 276

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{Exp andIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /;$ FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2670

$\text{Int}[\sin[(e_*) + (f_*)(x_)]^{(m_*)}*\tan[(e_*) + (f_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-f^{(-1)}, \text{Subst}[\text{Int}[(1 - x^2)^{(m + n - 1)/2}/x^n, x], x, \text{Cos}[e + f*x]], x] /;$ FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \cot^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x^2} dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-2 + \frac{1}{x^2} + x^2\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\csc(a + bx)}{b} - \frac{2 \sin(a + bx)}{b} + \frac{\sin^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 38, normalized size = 1.00

$$-\frac{\csc(a+bx)}{b} - \frac{2\sin(a+bx)}{b} + \frac{\sin^3(a+bx)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*Cot[a + b*x]^2,x]``[Out] -(Csc[a + b*x]/b) - (2*Sin[a + b*x])/b + Sin[a + b*x]^3/(3*b)`**Maple [A]**

time = 0.05, size = 52, normalized size = 1.37

method	result	size
derivativedivides	$\frac{-\frac{\cos^6(bx+a)}{\sin(bx+a)} - \left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{b}$	52
default	$\frac{-\frac{\cos^6(bx+a)}{\sin(bx+a)} - \left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{b}$	52
risch	$\frac{7ie^{i(bx+a)}}{8b} - \frac{7ie^{-i(bx+a)}}{8b} - \frac{2ie^{i(bx+a)}}{b(e^{2i(bx+a)}-1)} - \frac{\sin(3bx+3a)}{12b}$	74
norman	$\frac{-\frac{1}{2b} - \frac{6(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{25(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{3b} - \frac{6(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\tan^8(\frac{bx}{2} + \frac{a}{2})}{2b}}{(1+\tan^2(\frac{bx}{2} + \frac{a}{2}))^3 \tan(\frac{bx}{2} + \frac{a}{2})}$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5/sin(b*x+a)^2,x,method=_RETURNVERBOSE)``[Out] 1/b*(-cos(b*x+a)^6/sin(b*x+a)-(8/3+cos(b*x+a)^4+4/3*cos(b*x+a)^2)*sin(b*x+a))`**Maxima [A]**

time = 0.28, size = 32, normalized size = 0.84

$$\frac{\sin(bx+a)^3 - \frac{3}{\sin(bx+a)} - 6\sin(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5/sin(b*x+a)^2,x, algorithm="maxima")``[Out] 1/3*(sin(b*x + a)^3 - 3/sin(b*x + a) - 6*sin(b*x + a))/b`**Fricas [A]**

time = 0.36, size = 33, normalized size = 0.87

$$\frac{\cos(bx+a)^4 + 4\cos(bx+a)^2 - 8}{3b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^5/sin(b*x+a)^2,x, algorithm="fricas")`

[Out] `1/3*(cos(b*x + a)^4 + 4*cos(b*x + a)^2 - 8)/(b*sin(b*x + a))`

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 61 vs. $2(29) = 58$.

time = 0.55, size = 61, normalized size = 1.61

$$\begin{cases} -\frac{8\sin^3(a+bx)}{3b} - \frac{4\sin(a+bx)\cos^2(a+bx)}{b} - \frac{\cos^4(a+bx)}{b\sin(a+bx)} & \text{for } b \neq 0 \\ \frac{x\cos^5(a)}{\sin^2(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**5/sin(b*x+a)**2,x)`

[Out] `Piecewise((-8*sin(a + b*x)**3/(3*b) - 4*sin(a + b*x)*cos(a + b*x)**2/b - cos(a + b*x)**4/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)**5/sin(a)**2, True))`

Giac [A]

time = 3.49, size = 32, normalized size = 0.84

$$\frac{\sin(bx + a)^3 - \frac{3}{\sin(bx+a)} - 6\sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^5/sin(b*x+a)^2,x, algorithm="giac")`

[Out] `1/3*(sin(b*x + a)^3 - 3/sin(b*x + a) - 6*sin(b*x + a))/b`

Mupad [B]

time = 0.45, size = 35, normalized size = 0.92

$$-\frac{\sin(a + bx)^4 + 6\sin(a + bx)^2 + 3}{3b\sin(a + bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^5/sin(a + b*x)^2,x)`

[Out] `-(6*sin(a + b*x)^2 - sin(a + b*x)^4 + 3)/(3*b*sin(a + b*x))`

3.136 $\int \cos^2(a + bx) \cot^2(a + bx) dx$

Optimal. Leaf size=40

$$-\frac{3x}{2} - \frac{3 \cot(a + bx)}{2b} + \frac{\cos^2(a + bx) \cot(a + bx)}{2b}$$

[Out] $-3/2*x-3/2*\cot(b*x+a)/b+1/2*\cos(b*x+a)^2*\cot(b*x+a)/b$

Rubi [A]

time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2671, 294, 327, 209}

$$-\frac{3 \cot(a + bx)}{2b} + \frac{\cos^2(a + bx) \cot(a + bx)}{2b} - \frac{3x}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Cot[a + b*x]^2,x]

[Out] $(-3*x)/2 - (3*\cot[a + b*x])/(2*b) + (\cos[a + b*x]^2*\cot[a + b*x])/(2*b)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2671

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[In

```
t[(ff*x)^(m + n)/(b^2 + ff^2*x^2)^(m/2 + 1), x], x, b*(Tan[e + f*x]/ff)], x
]] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2]
```

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) \cot^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^4}{(1+x^2)^2} dx, x, \cot(a + bx)\right)}{b} \\ &= \frac{\cos^2(a + bx) \cot(a + bx)}{2b} - \frac{3\text{Subst}\left(\int \frac{x^2}{1+x^2} dx, x, \cot(a + bx)\right)}{2b} \\ &= -\frac{3 \cot(a + bx)}{2b} + \frac{\cos^2(a + bx) \cot(a + bx)}{2b} + \frac{3\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(a + bx)\right)}{2b} \\ &= -\frac{3x}{2} - \frac{3 \cot(a + bx)}{2b} + \frac{\cos^2(a + bx) \cot(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 31, normalized size = 0.78

$$-\frac{6(a + bx) + 4 \cot(a + bx) + \sin(2(a + bx))}{4b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x]^2*Cot[a + b*x]^2,x]
```

```
[Out] -1/4*(6*(a + b*x) + 4*Cot[a + b*x] + Sin[2*(a + b*x)])/b
```

Maple [A]

time = 0.03, size = 56, normalized size = 1.40

method	result	size
risch	$-\frac{3x}{2} + \frac{ie^{2i(bx+a)}}{8b} - \frac{ie^{-2i(bx+a)}}{8b} - \frac{2i}{b(e^{2i(bx+a)}-1)}$	54
derivativedivides	$-\frac{\cos^5(bx+a)}{\sin(bx+a)} - \left(\cos^3(bx+a) + \frac{3\cos(bx+a)}{2}\right) \sin(bx+a) - \frac{3bx}{2} - \frac{3a}{2}$ b	56
default	$-\frac{\cos^5(bx+a)}{\sin(bx+a)} - \left(\cos^3(bx+a) + \frac{3\cos(bx+a)}{2}\right) \sin(bx+a) - \frac{3bx}{2} - \frac{3a}{2}$ b	56
norman	$-\frac{1}{2b} - \frac{3\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b} + \frac{3\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b} + \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{2b} - \frac{3x \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{2} - 3x\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \frac{3x\left(\tan^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2}$ $(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right))^2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)$	122

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(b*x+a)^4/sin(b*x+a)^2,x,method=_RETURNVERBOSE)
```

[Out] $1/b*(-\cos(b*x+a)^5/\sin(b*x+a)-(\cos(b*x+a)^3+3/2*\cos(b*x+a))*\sin(b*x+a)-3/2*b*x-3/2*a)$

Maxima [A]

time = 0.51, size = 43, normalized size = 1.08

$$-\frac{3bx + 3a + \frac{3 \tan(bx+a)^2+2}{\tan(bx+a)^3+\tan(bx+a)}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4/sin(b*x+a)^2,x, algorithm="maxima")`

[Out] $-1/2*(3*b*x + 3*a + (3*\tan(b*x + a)^2 + 2)/(\tan(b*x + a)^3 + \tan(b*x + a)))/b$

Fricas [A]

time = 0.36, size = 40, normalized size = 1.00

$$\frac{\cos(bx+a)^3 - 3bx \sin(bx+a) - 3 \cos(bx+a)}{2b \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4/sin(b*x+a)^2,x, algorithm="fricas")`

[Out] $1/2*(\cos(b*x + a)^3 - 3*b*x*\sin(b*x + a) - 3*\cos(b*x + a))/(b*\sin(b*x + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 75 vs. $2(34) = 68$.

time = 0.44, size = 75, normalized size = 1.88

$$\begin{cases} -\frac{3x \sin^2(a+bx)}{2} - \frac{3x \cos^2(a+bx)}{2} - \frac{3 \sin(a+bx) \cos(a+bx)}{2b} - \frac{\cos^3(a+bx)}{b \sin(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^4(a)}{\sin^2(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**4/sin(b*x+a)**2,x)`

[Out] `Piecewise((-3*x*sin(a + b*x)**2/2 - 3*x*cos(a + b*x)**2/2 - 3*sin(a + b*x)*cos(a + b*x)/(2*b) - cos(a + b*x)**3/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)**4/sin(a)**2, True))`

Giac [A]

time = 2.95, size = 43, normalized size = 1.08

$$-\frac{3bx + 3a + \frac{3 \tan(bx+a)^2+2}{\tan(bx+a)^3+\tan(bx+a)}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^2,x, algorithm="giac")

[Out]
$$-1/2*(3*b*x + 3*a + (3*\tan(b*x + a)^2 + 2)/(\tan(b*x + a)^3 + \tan(b*x + a))) / b$$

Mupad [B]

time = 0.57, size = 43, normalized size = 1.08

$$\frac{9 \cos(a + b x) - \cos(3 a + 3 b x) + 12 b x \sin(a + b x)}{8 b \sin(a + b x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4/sin(a + b*x)^2,x)

[Out]
$$-(9*\cos(a + b*x) - \cos(3*a + 3*b*x) + 12*b*x*\sin(a + b*x))/(8*b*\sin(a + b*x))$$

3.137 $\int \cos(a + bx) \cot^2(a + bx) dx$

Optimal. Leaf size=23

$$-\frac{\csc(a + bx)}{b} - \frac{\sin(a + bx)}{b}$$

[Out] -csc(b*x+a)/b-sin(b*x+a)/b

Rubi [A]

time = 0.02, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 14}

$$-\frac{\sin(a + bx)}{b} - \frac{\csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Cot[a + b*x]^2,x]

[Out] -(Csc[a + b*x]/b) - Sin[a + b*x]/b

Rule 14

Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \cot^2(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^2} dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(-1 + \frac{1}{x^2}\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\csc(a + bx)}{b} - \frac{\sin(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 23, normalized size = 1.00

$$-\frac{\csc(a + bx)}{b} - \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Cot[a + b*x]^2,x]

[Out] -(Csc[a + b*x]/b) - Sin[a + b*x]/b

Maple [A]

time = 0.03, size = 42, normalized size = 1.83

method	result	size
derivativedivides	$\frac{-\frac{\cos^4(bx+a)}{\sin(bx+a)} - (2+\cos^2(bx+a)) \sin(bx+a)}{b}$	42
default	$\frac{-\frac{\cos^4(bx+a)}{\sin(bx+a)} - (2+\cos^2(bx+a)) \sin(bx+a)}{b}$	42
risch	$\frac{ie^{i(bx+a)}}{2b} - \frac{ie^{-i(bx+a)}}{2b} - \frac{2ie^{i(bx+a)}}{b(e^{2i(bx+a)}-1)}$	60
norman	$\frac{-\frac{1}{2b} - \frac{3(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\tan^4(\frac{bx}{2} + \frac{a}{2})}{2b}}{(1+\tan^2(\frac{bx}{2} + \frac{a}{2})) \tan(\frac{bx}{2} + \frac{a}{2})}$	66

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3/sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(-cos(b*x+a)^4/sin(b*x+a)-(2+cos(b*x+a)^2)*sin(b*x+a))

Maxima [A]

time = 0.28, size = 20, normalized size = 0.87

$$\frac{\frac{1}{\sin(bx+a)} + \sin(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^2,x, algorithm="maxima")

[Out] -(1/sin(b*x + a) + sin(b*x + a))/b

Fricas [A]

time = 0.35, size = 22, normalized size = 0.96

$$\frac{\cos(bx+a)^2 - 2}{b \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^2,x, algorithm="fricas")

[Out] (cos(b*x + a)^2 - 2)/(b*sin(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 39 vs. $2(17) = 34$.

time = 0.36, size = 39, normalized size = 1.70

$$\begin{cases} -\frac{2 \sin(a+bx)}{b} - \frac{\cos^2(a+bx)}{b \sin(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^3(a)}{\sin^2(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3/sin(b*x+a)**2,x)

[Out] Piecewise((-2*sin(a + b*x)/b - cos(a + b*x)**2/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)**3/sin(a)**2, True))

Giac [A]

time = 2.79, size = 20, normalized size = 0.87

$$-\frac{\frac{1}{\sin(bx+a)} + \sin(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^2,x, algorithm="giac")

[Out] -(1/sin(b*x + a) + sin(b*x + a))/b

Mupad [B]

time = 0.45, size = 23, normalized size = 1.00

$$-\frac{\sin(a+bx)^2+1}{b \sin(a+bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3/sin(a + b*x)^2,x)

[Out] -(sin(a + b*x)^2 + 1)/(b*sin(a + b*x))

3.138 $\int \cot^2(a + bx) dx$

Optimal. Leaf size=15

$$-x - \frac{\cot(a + bx)}{b}$$

[Out] $-x - \cot(b*x+a)/b$

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$-\frac{\cot(a + bx)}{b} - x$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^2,x]

[Out] $-x - \cot[a + b*x]/b$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_.)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \cot^2(a + bx) dx &= -\frac{\cot(a + bx)}{b} - \int 1 dx \\ &= -x - \frac{\cot(a + bx)}{b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 29, normalized size = 1.93

$$-\frac{\cot(a + bx) {}_2F_1\left(-\frac{1}{2}, 1; \frac{1}{2}; -\tan^2(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^2,x]

[Out] -((Cot[a + b*x]*Hypergeometric2F1[-1/2, 1, 1/2, -Tan[a + b*x]^2])/b)

Maple [A]

time = 0.03, size = 21, normalized size = 1.40

method	result	size
derivativedivides	$\frac{-\cot(bx+a)-bx-a}{b}$	21
default	$\frac{-\cot(bx+a)-bx-a}{b}$	21
risch	$-x - \frac{2i}{b(e^{2i(bx+a)}-1)}$	24
norman	$\frac{-\frac{1}{2b} + \frac{\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)}{2b} - x \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)}$	47

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2/sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(-cot(b*x+a)-b*x-a)

Maxima [A]

time = 0.51, size = 18, normalized size = 1.20

$$-\frac{bx + a + \frac{1}{\tan(bx+a)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^2,x, algorithm="maxima")

[Out] -(b*x + a + 1/tan(b*x + a))/b

Fricas [A]

time = 0.36, size = 29, normalized size = 1.93

$$-\frac{bx \sin(bx + a) + \cos(bx + a)}{b \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^2,x, algorithm="fricas")

[Out] -(b*x*sin(b*x + a) + cos(b*x + a))/(b*sin(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 29 vs. $2(10) = 20$.

time = 0.31, size = 29, normalized size = 1.93

$$\begin{cases} -x - \frac{\cos(a+bx)}{b \sin(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^2(a)}{\sin^2(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2/sin(b*x+a)**2,x)

[Out] Piecewise((-x - cos(a + b*x)/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)**2/sin(a)**2, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(15) = 30$.
time = 3.00, size = 35, normalized size = 2.33

$$-\frac{2bx + 2a + \frac{1}{\tan(\frac{1}{2}bx + \frac{1}{2}a)} - \tan(\frac{1}{2}bx + \frac{1}{2}a)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^2,x, algorithm="giac")

[Out] -1/2*(2*b*x + 2*a + 1/tan(1/2*b*x + 1/2*a) - tan(1/2*b*x + 1/2*a))/b

Mupad [B]

time = 0.41, size = 15, normalized size = 1.00

$$-x - \frac{\cot(a + bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2/sin(a + b*x)^2,x)

[Out] - x - cot(a + b*x)/b

3.139 $\int \cot(a + bx) \csc(a + bx) dx$

Optimal. Leaf size=11

$$-\frac{\csc(a + bx)}{b}$$

[Out] -csc(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 11, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {2686, 8}

$$-\frac{\csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]*Csc[a + b*x],x]

[Out] -(Csc[a + b*x]/b)

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \cot(a + bx) \csc(a + bx) dx &= -\frac{\text{Subst}(\int 1 dx, x, \csc(a + bx))}{b} \\ &= -\frac{\csc(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 11, normalized size = 1.00

$$-\frac{\csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]*Csc[a + b*x],x]

[Out] -(Csc[a + b*x]/b)

Maple [A]

time = 0.03, size = 14, normalized size = 1.27

method	result	size
derivativdivides	$-\frac{1}{\sin(bx+a)b}$	14
default	$-\frac{1}{\sin(bx+a)b}$	14
risch	$-\frac{2ie^{i(bx+a)}}{b(e^{2i(bx+a)}-1)}$	29
norman	$-\frac{\frac{1}{2b} - \frac{\tan^2\left(\frac{bx+a}{2}\right)}{2b}}{\tan\left(\frac{bx+a}{2}\right)}$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)/sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] -1/sin(b*x+a)/b

Maxima [A]

time = 0.28, size = 13, normalized size = 1.18

$$-\frac{1}{b \sin (bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^2,x, algorithm="maxima")

[Out] -1/(b*sin(b*x + a))

Fricas [A]

time = 0.36, size = 13, normalized size = 1.18

$$-\frac{1}{b \sin (bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/(b*sin(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 20 vs. $2(8) = 16$.

time = 0.31, size = 20, normalized size = 1.82

$$\begin{cases} -\frac{1}{b \sin(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos(a)}{\sin^2(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)**2,x)

[Out] Piecewise((-1/(b*sin(a + b*x)), Ne(b, 0)), (x*cos(a)/sin(a)**2, True))

Giac [A]

time = 2.74, size = 13, normalized size = 1.18

$$-\frac{1}{b \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^2,x, algorithm="giac")

[Out] -1/(b*sin(b*x + a))

Mupad [B]

time = 0.42, size = 13, normalized size = 1.18

$$-\frac{1}{b \sin(a + bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)/sin(a + b*x)^2,x)

[Out] -1/(b*sin(a + b*x))

3.140 $\int \csc^2(a + bx) \sec(a + bx) dx$

Optimal. Leaf size=23

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\csc(a + bx)}{b}$$

[Out] arctanh(sin(b*x+a))/b-csc(b*x+a)/b

Rubi [A]

time = 0.01, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2701, 327, 213}

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^2*Sec[a + b*x],x]

[Out] ArcTanh[Sin[a + b*x]]/b - Csc[a + b*x]/b

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \csc^2(a + bx) \sec(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^2}{-1+x^2} dx, x, \csc(a + bx)\right)}{b} \\ &= -\frac{\csc(a + bx)}{b} - \frac{\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\csc(a + bx)}{b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 27, normalized size = 1.17

$$-\frac{\csc(a + bx) {}_2F_1\left(-\frac{1}{2}, 1; \frac{1}{2}; \sin^2(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2*Sec[a + b*x],x]

[Out] -((Csc[a + b*x]*Hypergeometric2F1[-1/2, 1, 1/2, Sin[a + b*x]^2])/b)

Maple [A]

time = 0.04, size = 30, normalized size = 1.30

method	result	size
derivativedivides	$-\frac{1}{\sin(bx+a)} + \frac{\ln(\sec(bx+a) + \tan(bx+a))}{b}$	30
default	$-\frac{1}{\sin(bx+a)} + \frac{\ln(\sec(bx+a) + \tan(bx+a))}{b}$	30
risch	$-\frac{2ie^{i(bx+a)}}{b(e^{2i(bx+a)}-1)} + \frac{\ln(e^{i(bx+a)}+i)}{b} - \frac{\ln(e^{i(bx+a)}-i)}{b}$	65
norman	$-\frac{1}{2b} - \frac{\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)}{2b} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b}$	69

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)/sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/sin(b*x+a)+ln(sec(b*x+a)+tan(b*x+a)))

Maxima [A]

time = 0.30, size = 36, normalized size = 1.57

$$-\frac{\frac{2}{\sin(bx+a)} - \log(\sin(bx+a) + 1) + \log(\sin(bx+a) - 1)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^2,x, algorithm="maxima")

[Out] -1/2*(2/sin(b*x + a) - log(sin(b*x + a) + 1) + log(sin(b*x + a) - 1))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(23) = 46.

time = 0.40, size = 50, normalized size = 2.17

$$\frac{\log(\sin(bx + a) + 1) \sin(bx + a) - \log(-\sin(bx + a) + 1) \sin(bx + a) - 2}{2b \sin(bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^2,x, algorithm="fricas")

[Out] 1/2*(log(sin(b*x + a) + 1)*sin(b*x + a) - log(-sin(b*x + a) + 1)*sin(b*x + a) - 2)/(b*sin(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(a + bx)}{\sin^2(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)**2,x)

[Out] Integral(sec(a + b*x)/sin(a + b*x)**2, x)

Giac [A]

time = 7.08, size = 38, normalized size = 1.65

$$\frac{\frac{2}{\sin(bx+a)} - \log(|\sin(bx + a) + 1|) + \log(|\sin(bx + a) - 1|)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^2,x, algorithm="giac")

[Out] -1/2*(2/sin(b*x + a) - log(abs(sin(b*x + a) + 1)) + log(abs(sin(b*x + a) - 1)))/b

Mupad [B]

time = 0.02, size = 22, normalized size = 0.96

$$\frac{\operatorname{atanh}(\sin(a + bx)) - \frac{1}{\sin(a + bx)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)*sin(a + b*x)^2),x)

[Out] (atanh(sin(a + b*x)) - 1/sin(a + b*x))/b

3.141 $\int \csc^2(a + bx) \sec^2(a + bx) dx$

Optimal. Leaf size=22

$$-\frac{\cot(a + bx)}{b} + \frac{\tan(a + bx)}{b}$$

[Out] $-\cot(b*x+a)/b+\tan(b*x+a)/b$

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2700, 14}

$$\frac{\tan(a + bx)}{b} - \frac{\cot(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^2*Sec[a + b*x]^2,x]`

[Out] $-(\text{Cot}[a + b*x]/b) + \text{Tan}[a + b*x]/b$

Rule 14

`Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]`

Rule 2700

`Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]`

Rubi steps

$$\begin{aligned} \int \csc^2(a + bx) \sec^2(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{1+x^2}{x^2} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^2}\right) dx, x, \tan(a + bx)\right)}{b} \\ &= -\frac{\cot(a + bx)}{b} + \frac{\tan(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 13, normalized size = 0.59

$$-\frac{2 \cot(2(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2*Sec[a + b*x]^2,x]

[Out] (-2*Cot[2*(a + b*x)])/b

Maple [A]

time = 0.09, size = 31, normalized size = 1.41

method	result	size
derivativedivides	$\frac{1}{\sin(bx+a)\cos(bx+a)} - 2\cot(bx+a)$ b	31
default	$\frac{1}{\sin(bx+a)\cos(bx+a)} - 2\cot(bx+a)$ b	31
risch	$-\frac{4i}{b(e^{2i(bx+a)}+1)(e^{2i(bx+a)}-1)}$	33
norman	$\frac{1}{2b} - \frac{3(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{\tan^4(\frac{bx}{2} + \frac{a}{2})}{2b}$ $\frac{1}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)\tan(\frac{bx}{2} + \frac{a}{2})}$	66

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2/sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/b*(1/sin(b*x+a)/cos(b*x+a)-2*cot(b*x+a))

Maxima [A]

time = 0.29, size = 22, normalized size = 1.00

$$-\frac{\frac{1}{\tan(bx+a)} - \tan(bx+a)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^2,x, algorithm="maxima")

[Out] -(1/tan(b*x + a) - tan(b*x + a))/b

Fricas [A]

time = 0.34, size = 33, normalized size = 1.50

$$-\frac{2\cos(bx+a)^2 - 1}{b\cos(bx+a)\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^2,x, algorithm="fricas")

[Out] -(2*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)*sin(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(a + bx)}{\sin^2(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2/sin(b*x+a)**2,x)

[Out] Integral(sec(a + b*x)**2/sin(a + b*x)**2, x)

Giac [A]

time = 3.17, size = 16, normalized size = 0.73

$$-\frac{2}{b \tan(2bx + 2a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^2,x, algorithm="giac")

[Out] -2/(b*tan(2*b*x + 2*a))

Mupad [B]

time = 0.39, size = 14, normalized size = 0.64

$$-\frac{2 \cot(2a + 2bx)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^2*sin(a + b*x)^2),x)

[Out] -(2*cot(2*a + 2*b*x))/b

3.142 $\int \csc^2(a + bx) \sec^3(a + bx) dx$

Optimal. Leaf size=49

$$\frac{3 \tanh^{-1}(\sin(a + bx))}{2b} - \frac{3 \csc(a + bx)}{2b} + \frac{\csc(a + bx) \sec^2(a + bx)}{2b}$$

[Out] 3/2*arctanh(sin(b*x+a))/b-3/2*csc(b*x+a)/b+1/2*csc(b*x+a)*sec(b*x+a)^2/b

Rubi [A]

time = 0.03, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2701, 294, 327, 213}

$$-\frac{3 \csc(a + bx)}{2b} + \frac{3 \tanh^{-1}(\sin(a + bx))}{2b} + \frac{\csc(a + bx) \sec^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^2*Sec[a + b*x]^3,x]

[Out] (3*ArcTanh[Sin[a + b*x]])/(2*b) - (3*Csc[a + b*x])/(2*b) + (Csc[a + b*x]*Sec[a + b*x]^2)/(2*b)

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1)*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

```
Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol]
:> Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^2(a + bx) \sec^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^4}{(-1+x^2)^2} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\csc(a + bx) \sec^2(a + bx)}{2b} - \frac{3\text{Subst}\left(\int \frac{x^2}{-1+x^2} dx, x, \csc(a + bx)\right)}{2b} \\ &= -\frac{3 \csc(a + bx)}{2b} + \frac{\csc(a + bx) \sec^2(a + bx)}{2b} - \frac{3\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \csc(a + bx)\right)}{2b} \\ &= \frac{3 \tanh^{-1}(\sin(a + bx))}{2b} - \frac{3 \csc(a + bx)}{2b} + \frac{\csc(a + bx) \sec^2(a + bx)}{2b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 27, normalized size = 0.55

$$-\frac{\csc(a + bx) {}_2F_1\left(-\frac{1}{2}, 2; \frac{1}{2}; \sin^2(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2*Sec[a + b*x]^3,x]

[Out] -((Csc[a + b*x]*Hypergeometric2F1[-1/2, 2, 1/2, Sin[a + b*x]^2])/b)

Maple [A]

time = 0.05, size = 50, normalized size = 1.02

method	result	size
derivativedivides	$\frac{1}{2 \sin(bx+a) \cos(bx+a)^2} - \frac{3}{2 \sin(bx+a)} + \frac{3 \ln(\sec(bx+a) + \tan(bx+a))}{2}$	50
default	$\frac{1}{2 \sin(bx+a) \cos(bx+a)^2} - \frac{3}{2 \sin(bx+a)} + \frac{3 \ln(\sec(bx+a) + \tan(bx+a))}{2}$	50
risch	$-\frac{i(3e^{5i(bx+a)} + 2e^{3i(bx+a)} + 3e^{i(bx+a)})}{b(e^{2i(bx+a)} + 1)^2(e^{2i(bx+a)} - 1)} - \frac{3 \ln(e^{i(bx+a)} - i)}{2b} + \frac{3 \ln(e^{i(bx+a)} + i)}{2b}$	104
norman	$-\frac{1}{2b} + \frac{3\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b} + \frac{3\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b} - \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{2b} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{2b} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{2b}$	117

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^3/sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] `1/b*(1/2/sin(b*x+a)/cos(b*x+a)^2-3/2/sin(b*x+a)+3/2*ln(sec(b*x+a)+tan(b*x+a)))`

Maxima [A]

time = 0.28, size = 61, normalized size = 1.24

$$\frac{2(3\sin(bx+a)^2-2)}{\sin(bx+a)^3-\sin(bx+a)} - 3\log(\sin(bx+a)+1) + 3\log(\sin(bx+a)-1)$$

$$4b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3/sin(b*x+a)^2,x, algorithm="maxima")`

[Out] `-1/4*(2*(3*sin(b*x + a)^2 - 2)/(sin(b*x + a)^3 - sin(b*x + a)) - 3*log(sin(b*x + a) + 1) + 3*log(sin(b*x + a) - 1))/b`

Fricas [A]

time = 0.36, size = 85, normalized size = 1.73

$$\frac{3\cos(bx+a)^2\log(\sin(bx+a)+1)\sin(bx+a) - 3\cos(bx+a)^2\log(-\sin(bx+a)+1)\sin(bx+a) - 6\cos(bx+a)^2 + 2}{4b\cos(bx+a)^2\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3/sin(b*x+a)^2,x, algorithm="fricas")`

[Out] `1/4*(3*cos(b*x + a)^2*log(sin(b*x + a) + 1)*sin(b*x + a) - 3*cos(b*x + a)^2*log(-sin(b*x + a) + 1)*sin(b*x + a) - 6*cos(b*x + a)^2 + 2)/(b*cos(b*x + a)^2*sin(b*x + a))`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(a+bx)}{\sin^2(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**3/sin(b*x+a)**2,x)`

[Out] `Integral(sec(a + b*x)**3/sin(a + b*x)**2, x)`

Giac [A]

time = 4.05, size = 63, normalized size = 1.29

$$\frac{2(3\sin(bx+a)^2-2)}{\sin(bx+a)^3-\sin(bx+a)} - 3\log(|\sin(bx+a)+1|) + 3\log(|\sin(bx+a)-1|)$$

$$4b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a)^2,x, algorithm="giac")

[Out] $-1/4*(2*(3*\sin(b*x + a)^2 - 2)/(\sin(b*x + a)^3 - \sin(b*x + a)) - 3*\log(\text{abs}(\sin(b*x + a) + 1)) + 3*\log(\text{abs}(\sin(b*x + a) - 1)))/b$

Mupad [B]

time = 0.43, size = 48, normalized size = 0.98

$$\frac{3 \operatorname{atanh}(\sin(a + bx))}{2b} + \frac{\frac{3 \sin(a+bx)^2}{2} - 1}{b (\sin(a + bx) - \sin(a + bx)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^3*sin(a + b*x)^2),x)

[Out] $(3*\operatorname{atanh}(\sin(a + b*x)))/(2*b) + ((3*\sin(a + b*x)^2)/2 - 1)/(b*(\sin(a + b*x) - \sin(a + b*x)^3))$

3.143 $\int \csc^2(a + bx) \sec^4(a + bx) dx$

Optimal. Leaf size=38

$$-\frac{\cot(a + bx)}{b} + \frac{2 \tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b}$$

[Out] $-\cot(b*x+a)/b+2*\tan(b*x+a)/b+1/3*\tan(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2700, 276}

$$\frac{\tan^3(a + bx)}{3b} + \frac{2 \tan(a + bx)}{b} - \frac{\cot(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[a + b*x]^2*\text{Sec}[a + b*x]^4, x]$

[Out] $-(\text{Cot}[a + b*x]/b) + (2*\text{Tan}[a + b*x])/b + \text{Tan}[a + b*x]^3/(3*b)$

Rule 276

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_)})^{(p_*)}, x_Symbol] :> \text{Int}[\text{Exp andIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0]$

Rule 2700

$\text{Int}[\text{csc}[(e_*) + (f_*)(x_)]^{(m_*)}\text{sec}[(e_*) + (f_*)(x_)]^{(n_*)}, x_Symbol] :> \text{Dist}[1/f, \text{Subst}[\text{Int}[(1 + x^2)^{(m+n)/2 - 1}/x^m, x], x, \text{Tan}[e + f*x]], x] /; \text{FreeQ}\{e, f\}, x] \&\& \text{IntegersQ}[m, n, (m+n)/2]$

Rubi steps

$$\begin{aligned} \int \csc^2(a + bx) \sec^4(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{x^2} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(2 + \frac{1}{x^2} + x^2\right) dx, x, \tan(a + bx)\right)}{b} \\ &= -\frac{\cot(a + bx)}{b} + \frac{2 \tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 46, normalized size = 1.21

$$-\frac{\cot(a+bx)}{b} + \frac{5 \tan(a+bx)}{3b} + \frac{\sec^2(a+bx) \tan(a+bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2*Sec[a + b*x]^4,x]**[Out]** -(Cot[a + b*x]/b) + (5*Tan[a + b*x])/(3*b) + (Sec[a + b*x]^2*Tan[a + b*x])/(3*b)**Maple [A]**

time = 0.06, size = 50, normalized size = 1.32

method	result	size
risch	$-\frac{16i(2e^{2i(bx+a)}+1)}{3b(e^{2i(bx+a)}+1)^3(e^{2i(bx+a)}-1)}$	46
derivativedivides	$\frac{\frac{1}{3 \sin(bx+a) \cos(bx+a)^3} + \frac{4}{3 \sin(bx+a) \cos(bx+a)} - \frac{8 \cot(bx+a)}{3}}{b}$	50
default	$\frac{\frac{1}{3 \sin(bx+a) \cos(bx+a)^3} + \frac{4}{3 \sin(bx+a) \cos(bx+a)} - \frac{8 \cot(bx+a)}{3}}{b}$	50
norman	$\frac{\frac{1}{2b} - \frac{6(\tan^2(\frac{bx+a}{2}))}{b} + \frac{25(\tan^4(\frac{bx+a}{2}))}{3b} - \frac{6(\tan^6(\frac{bx+a}{2}))}{b} + \frac{\tan^8(\frac{bx+a}{2})}{2b}}{(\tan^2(\frac{bx+a}{2})-1)^3 \tan(\frac{bx+a}{2})}$	98

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^4/sin(b*x+a)^2,x,method=_RETURNVERBOSE)**[Out]** 1/b*(1/3/sin(b*x+a)/cos(b*x+a)^3+4/3/sin(b*x+a)/cos(b*x+a)-8/3*cot(b*x+a))**Maxima [A]**

time = 0.28, size = 32, normalized size = 0.84

$$\frac{\tan(bx+a)^3 - \frac{3}{\tan(bx+a)} + 6 \tan(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^2,x, algorithm="maxima")**[Out]** 1/3*(tan(b*x + a)^3 - 3/tan(b*x + a) + 6*tan(b*x + a))/b**Fricas [A]**

time = 0.35, size = 43, normalized size = 1.13

$$-\frac{8 \cos(bx+a)^4 - 4 \cos(bx+a)^2 - 1}{3b \cos(bx+a)^3 \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^2,x, algorithm="fricas")

[Out] -1/3*(8*cos(b*x + a)^4 - 4*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^3*sin(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(a + bx)}{\sin^2(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4/sin(b*x+a)**2,x)

[Out] Integral(sec(a + b*x)**4/sin(a + b*x)**2, x)

Giac [A]

time = 3.97, size = 32, normalized size = 0.84

$$\frac{\tan(bx + a)^3 - \frac{3}{\tan(bx+a)} + 6 \tan(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^2,x, algorithm="giac")

[Out] 1/3*(tan(b*x + a)^3 - 3/tan(b*x + a) + 6*tan(b*x + a))/b

Mupad [B]

time = 0.41, size = 33, normalized size = 0.87

$$\frac{\tan(a + bx)^4 + 6 \tan(a + bx)^2 - 3}{3b \tan(a + bx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^4*sin(a + b*x)^2),x)

[Out] (6*tan(a + b*x)^2 + tan(a + b*x)^4 - 3)/(3*b*tan(a + b*x))

3.144 $\int \csc^2(a + bx) \sec^5(a + bx) dx$

Optimal. Leaf size=70

$$\frac{15 \tanh^{-1}(\sin(a + bx))}{8b} - \frac{15 \csc(a + bx)}{8b} + \frac{5 \csc(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc(a + bx) \sec^4(a + bx)}{4b}$$

[Out] 15/8*arctanh(sin(b*x+a))/b-15/8*csc(b*x+a)/b+5/8*csc(b*x+a)*sec(b*x+a)^2/b+1/4*csc(b*x+a)*sec(b*x+a)^4/b

Rubi [A]

time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2701, 294, 327, 213}

$$-\frac{15 \csc(a + bx)}{8b} + \frac{15 \tanh^{-1}(\sin(a + bx))}{8b} + \frac{\csc(a + bx) \sec^4(a + bx)}{4b} + \frac{5 \csc(a + bx) \sec^2(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^2*Sec[a + b*x]^5,x]

[Out] (15*ArcTanh[Sin[a + b*x]])/(8*b) - (15*Csc[a + b*x])/(8*b) + (5*Csc[a + b*x]*Sec[a + b*x]^2)/(8*b) + (Csc[a + b*x]*Sec[a + b*x]^4)/(4*b)

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && ! LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2701

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol]
:> Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^2(a + bx) \sec^5(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{(-1+x^2)^3} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\csc(a + bx) \sec^4(a + bx)}{4b} - \frac{5 \text{Subst}\left(\int \frac{x^4}{(-1+x^2)^2} dx, x, \csc(a + bx)\right)}{4b} \\ &= \frac{5 \csc(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc(a + bx) \sec^4(a + bx)}{4b} - \frac{15 \text{Subst}\left(\int \frac{x^2}{-1+x^2} dx, x, \csc(a + bx)\right)}{8b} \\ &= -\frac{15 \csc(a + bx)}{8b} + \frac{5 \csc(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc(a + bx) \sec^4(a + bx)}{4b} \\ &= \frac{15 \tanh^{-1}(\sin(a + bx))}{8b} - \frac{15 \csc(a + bx)}{8b} + \frac{5 \csc(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc(a + bx) \sec^4(a + bx)}{4b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 27, normalized size = 0.39

$$-\frac{\csc(a + bx) {}_2F_1\left(-\frac{1}{2}, 3; \frac{1}{2}; \sin^2(a + bx)\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2*Sec[a + b*x]^5,x]

[Out] -((Csc[a + b*x]*Hypergeometric2F1[-1/2, 3, 1/2, Sin[a + b*x]^2])/b)

Maple [A]

time = 0.07, size = 68, normalized size = 0.97

method	result
derivativedivides	$\frac{\frac{1}{4 \sin(bx+a) \cos(bx+a)^4} + \frac{5}{8 \sin(bx+a) \cos(bx+a)^2} - \frac{15}{8 \sin(bx+a)} + \frac{15 \ln(\sec(bx+a) + \tan(bx+a))}{8}}{b}$
default	$\frac{\frac{1}{4 \sin(bx+a) \cos(bx+a)^4} + \frac{5}{8 \sin(bx+a) \cos(bx+a)^2} - \frac{15}{8 \sin(bx+a)} + \frac{15 \ln(\sec(bx+a) + \tan(bx+a))}{8}}{b}$
risch	$-\frac{i(15 e^{9i(bx+a)} + 40 e^{7i(bx+a)} + 18 e^{5i(bx+a)} + 40 e^{3i(bx+a)} + 15 e^{i(bx+a)})}{4b(e^{2i(bx+a)} + 1)^4(e^{2i(bx+a)} - 1)} - \frac{15 \ln(e^{i(bx+a)} - i)}{8b} + \frac{15 \ln(e^{i(bx+a)} + i)}{8b}$

norman	$\frac{-\frac{1}{2b} + \frac{15 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \frac{5 \left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \frac{5 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \frac{15 \left(\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \frac{\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)}{2b}}{4b} - \frac{15 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^4 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)}$
--------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^5/sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] `1/b*(1/4/sin(b*x+a)/cos(b*x+a)^4+5/8/sin(b*x+a)/cos(b*x+a)^2-15/8/sin(b*x+a)+15/8*ln(sec(b*x+a)+tan(b*x+a)))`

Maxima [A]

time = 0.28, size = 79, normalized size = 1.13

$$\frac{2 \left(15 \sin(bx+a)^4 - 25 \sin(bx+a)^2 + 8 \right)}{\sin(bx+a)^5 - 2 \sin(bx+a)^3 + \sin(bx+a)} - 15 \log(\sin(bx+a) + 1) + 15 \log(\sin(bx+a) - 1)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^5/sin(b*x+a)^2,x, algorithm="maxima")`

[Out] `-1/16*(2*(15*sin(b*x + a)^4 - 25*sin(b*x + a)^2 + 8)/(sin(b*x + a)^5 - 2*sin(b*x + a)^3 + sin(b*x + a)) - 15*log(sin(b*x + a) + 1) + 15*log(sin(b*x + a) - 1))/b`

Fricas [A]

time = 0.37, size = 95, normalized size = 1.36

$$\frac{15 \cos(bx+a)^4 \log(\sin(bx+a) + 1) \sin(bx+a) - 15 \cos(bx+a)^4 \log(-\sin(bx+a) + 1) \sin(bx+a) - 30 \cos(bx+a)^4 + 10 \cos(bx+a)^2 + 4}{16b \cos(bx+a)^4 \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^5/sin(b*x+a)^2,x, algorithm="fricas")`

[Out] `1/16*(15*cos(b*x + a)^4*log(sin(b*x + a) + 1)*sin(b*x + a) - 15*cos(b*x + a)^4*log(-sin(b*x + a) + 1)*sin(b*x + a) - 30*cos(b*x + a)^4 + 10*cos(b*x + a)^2 + 4)/(b*cos(b*x + a)^4*sin(b*x + a))`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(a + bx)}{\sin^2(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**5/sin(b*x+a)**2,x)`

[Out] `Integral(sec(a + b*x)**5/sin(a + b*x)**2, x)`

Giac [A]

time = 4.27, size = 73, normalized size = 1.04

$$\frac{2 \left(7 \sin(bx+a)^3 - 9 \sin(bx+a) \right)}{\left(\sin(bx+a)^2 - 1 \right)^2} + \frac{16}{\sin(bx+a)} - 15 \log(|\sin(bx+a) + 1|) + 15 \log(|\sin(bx+a) - 1|)$$

$$16b$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^5/sin(b*x+a)^2,x, algorithm="giac")`

```
[Out] -1/16*(2*(7*sin(b*x + a)^3 - 9*sin(b*x + a))/(sin(b*x + a)^2 - 1)^2 + 16/sin(b*x + a) - 15*log(abs(sin(b*x + a) + 1)) + 15*log(abs(sin(b*x + a) - 1)))/b
```

Mupad [B]

time = 0.09, size = 67, normalized size = 0.96

$$\frac{15 \operatorname{atanh}(\sin(a + bx))}{8b} - \frac{\frac{15 \sin(a+bx)^4}{8} - \frac{25 \sin(a+bx)^2}{8} + 1}{b (\sin(a + bx)^5 - 2 \sin(a + bx)^3 + \sin(a + bx))}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(cos(a + b*x)^5*sin(a + b*x)^2),x)`

```
[Out] (15*atanh(sin(a + b*x)))/(8*b) - ((15*sin(a + b*x)^4)/8 - (25*sin(a + b*x)^2)/8 + 1)/(b*(sin(a + b*x) - 2*sin(a + b*x)^3 + sin(a + b*x)^5))
```


3.145 $\int \cos^4(a + bx) \cot^3(a + bx) dx$

Optimal. Leaf size=58

$$-\frac{\csc^2(a + bx)}{2b} - \frac{3 \log(\sin(a + bx))}{b} + \frac{3 \sin^2(a + bx)}{2b} - \frac{\sin^4(a + bx)}{4b}$$

[Out] $-1/2*\csc(b*x+a)^2/b-3*\ln(\sin(b*x+a))/b+3/2*\sin(b*x+a)^2/b-1/4*\sin(b*x+a)^4/b$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2670, 272, 45}

$$-\frac{\sin^4(a + bx)}{4b} + \frac{3 \sin^2(a + bx)}{2b} - \frac{\csc^2(a + bx)}{2b} - \frac{3 \log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Cot[a + b*x]^3,x]

[Out] $-1/2*\text{Csc}[a + b*x]^2/b - (3*\text{Log}[\text{Sin}[a + b*x]])/b + (3*\text{Sin}[a + b*x]^2)/(2*b) - \text{Sin}[a + b*x]^4/(4*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \cos^4(a+bx) \cot^3(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^3}{x^3} dx, x, -\sin(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1-x)^3}{x^2} dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(3 + \frac{1}{x^2} - \frac{3}{x} - x\right) dx, x, \sin^2(a+bx)\right)}{2b} \\
&= -\frac{\csc^2(a+bx)}{2b} - \frac{3 \log(\sin(a+bx))}{b} + \frac{3 \sin^2(a+bx)}{2b} - \frac{\sin^4(a+bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 45, normalized size = 0.78

$$-\frac{2 \csc^2(a+bx) + 12 \log(\sin(a+bx)) - 6 \sin^2(a+bx) + \sin^4(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^4*Cot[a + b*x]^3,x]``[Out] -1/4*(2*Csc[a + b*x]^2 + 12*Log[Sin[a + b*x]] - 6*Sin[a + b*x]^2 + Sin[a + b*x]^4)/b`**Maple [A]**

time = 0.05, size = 63, normalized size = 1.09

method	result
derivativedivides	$\frac{-\frac{\cos^8(bx+a)}{2 \sin(bx+a)^2} - \frac{\cos^6(bx+a)}{2} - \frac{3(\cos^4(bx+a))}{4} - \frac{3(\cos^2(bx+a))}{2} - 3 \ln(\sin(bx+a))}{b}$
default	$\frac{-\frac{\cos^8(bx+a)}{2 \sin(bx+a)^2} - \frac{\cos^6(bx+a)}{2} - \frac{3(\cos^4(bx+a))}{4} - \frac{3(\cos^2(bx+a))}{2} - 3 \ln(\sin(bx+a))}{b}$
risch	$3ix - \frac{5e^{2i(bx+a)}}{16b} - \frac{5e^{-2i(bx+a)}}{16b} + \frac{6ia}{b} + \frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}-1)^2} - \frac{3 \ln(e^{2i(bx+a)}-1)}{b} - \frac{\cos(4bx+4a)}{32b}$
norman	$\frac{-\frac{1}{8b} - \frac{\tan^{12}\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} + \frac{10\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{57\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} + \frac{57\left(\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^4 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{3 \ln\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^7/sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/2/sin(b*x+a)^2*cos(b*x+a)^8-1/2*cos(b*x+a)^6-3/4*cos(b*x+a)^4-3/2*cos(b*x+a)^2-3*ln(sin(b*x+a)))`

Maxima [A]

time = 0.28, size = 45, normalized size = 0.78

$$\frac{\sin (bx+a)^4 - 6 \sin (bx+a)^2 + \frac{2}{\sin (bx+a)^2} + 6 \log (\sin (bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^3,x, algorithm="maxima")

[Out] -1/4*(sin(b*x + a)^4 - 6*sin(b*x + a)^2 + 2/sin(b*x + a)^2 + 6*log(sin(b*x + a)^2))/b

Fricas [A]

time = 0.38, size = 71, normalized size = 1.22

$$\frac{8 \cos (bx+a)^6 + 24 \cos (bx+a)^4 - 51 \cos (bx+a)^2 + 96 (\cos (bx+a)^2 - 1) \log \left(\frac{1}{2} \sin (bx+a)\right) + 3}{32 (b \cos (bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^3,x, algorithm="fricas")

[Out] -1/32*(8*cos(b*x + a)^6 + 24*cos(b*x + a)^4 - 51*cos(b*x + a)^2 + 96*(cos(b*x + a)^2 - 1)*log(1/2*sin(b*x + a)) + 3)/(b*cos(b*x + a)^2 - b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 1484 vs. 2(48) = 96.

time = 5.62, size = 1484, normalized size = 25.59

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**7/sin(b*x+a)**3,x)

[Out] Piecewise((24*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**10/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 96*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**8/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 144*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**6/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 96*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 24*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 3

```

2*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 24*log(tan(a/2 + b*x/2
)))*tan(a/2 + b*x/2)**10/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**
8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x
/2)**2) - 96*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(8*b*tan(a/2 + b*x/2
)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2
+ b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 144*log(tan(a/2 + b*x/2))*tan(a/2
+ b*x/2)**6/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan
(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 96
*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**10 + 32*b
*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4
+ 8*b*tan(a/2 + b*x/2)**2) - 24*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(
8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 + b*x/2)
**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - tan(a/2 + b*x/2
)**12/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*tan(a/2 +
b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 57*tan(a
/2 + b*x/2)**8/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 + 48*b*
tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) +
80*tan(a/2 + b*x/2)**6/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**
8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x
/2)**2) + 57*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 +
b*x/2)**8 + 48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(
a/2 + b*x/2)**2) - 1/(8*b*tan(a/2 + b*x/2)**10 + 32*b*tan(a/2 + b*x/2)**8 +
48*b*tan(a/2 + b*x/2)**6 + 32*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)
**2), Ne(b, 0)), (x*cos(a)**7/sin(a)**3, True))

```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 231 vs. 2(52) = 104.

time = 4.38, size = 231, normalized size = 3.98

$$\frac{\frac{12(\cos(bx+a)-1)+1}{\cos(bx+a)+1}(\cos(bx+a)+1)}{\cos(bx+a)-1} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + \frac{2\left(\frac{76(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{118(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{76(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{25(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - 25\right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^4} - 12 \log\left(\frac{-\cos(bx+a)+1}{\cos(bx+a)+1}\right) + 24 \log\left(\left|\frac{-\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right|\right)$$

8b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/8*((12*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)*(cos(b*x + a) + 1)/(cos(b*x + a) - 1) + (cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 2*(76*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 118*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 76*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 - 25*(cos(b*x + a) - 1)^4/(cos(b*x + a) + 1)^4 - 25)/(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)^4 - 12*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)) + 24*log(abs(-(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)))/b

Mupad [B]

time = 0.69, size = 85, normalized size = 1.47

$$\frac{3 \ln(\tan(a + bx)^2 + 1)}{2b} - \frac{3 \ln(\tan(a + bx))}{b} - \frac{\frac{3 \tan(a+bx)^4}{2} + \frac{9 \tan(a+bx)^2}{4} + \frac{1}{2}}{b (\tan(a + bx)^6 + 2 \tan(a + bx)^4 + \tan(a + bx)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^7/sin(a + b*x)^3,x)

[Out] (3*log(tan(a + b*x)^2 + 1))/(2*b) - (3*log(tan(a + b*x)))/b - ((9*tan(a + b*x)^2)/4 + (3*tan(a + b*x)^4)/2 + 1/2)/(b*(tan(a + b*x)^2 + 2*tan(a + b*x)^4 + tan(a + b*x)^6))

3.146 $\int \cos^3(a + bx) \cot^3(a + bx) dx$

Optimal. Leaf size=66

$$\frac{5 \tanh^{-1}(\cos(a + bx))}{2b} - \frac{5 \cos(a + bx)}{2b} - \frac{5 \cos^3(a + bx)}{6b} - \frac{\cos^3(a + bx) \cot^2(a + bx)}{2b}$$

[Out] $5/2*\operatorname{arctanh}(\cos(b*x+a))/b-5/2*\cos(b*x+a)/b-5/6*\cos(b*x+a)^3/b-1/2*\cos(b*x+a)^3*\cot(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2672, 294, 308, 212}

$$-\frac{5 \cos^3(a + bx)}{6b} - \frac{5 \cos(a + bx)}{2b} - \frac{\cos^3(a + bx) \cot^2(a + bx)}{2b} + \frac{5 \tanh^{-1}(\cos(a + bx))}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^3*Cot[a + b*x]^3,x]`

[Out] $(5*\operatorname{ArcTanh}[\cos[a + b*x]])/(2*b) - (5*\cos[a + b*x])/(2*b) - (5*\cos[a + b*x]^3)/(6*b) - (\cos[a + b*x]^3*\cot[a + b*x]^2)/(2*b)$

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 294

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !ILtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 308

`Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]`

Rule 2672

`Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(`

```
ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x]
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \cot^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{(1-x^2)^2} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos^3(a + bx) \cot^2(a + bx)}{2b} + \frac{5\text{Subst}\left(\int \frac{x^4}{1-x^2} dx, x, \cos(a + bx)\right)}{2b} \\ &= -\frac{\cos^3(a + bx) \cot^2(a + bx)}{2b} + \frac{5\text{Subst}\left(\int \left(-1 - x^2 + \frac{1}{1-x^2}\right) dx, x, \cos(a + bx)\right)}{2b} \\ &= -\frac{5 \cos(a + bx)}{2b} - \frac{5 \cos^3(a + bx)}{6b} - \frac{\cos^3(a + bx) \cot^2(a + bx)}{2b} + \frac{5\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(a + bx)\right)}{2b} \\ &= \frac{5 \tanh^{-1}(\cos(a + bx))}{2b} - \frac{5 \cos(a + bx)}{2b} - \frac{5 \cos^3(a + bx)}{6b} - \frac{\cos^3(a + bx) \cot^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 103, normalized size = 1.56

$$-\frac{9 \cos(a + bx)}{4b} - \frac{\cos(3(a + bx))}{12b} - \frac{\csc^2\left(\frac{1}{2}(a + bx)\right)}{8b} + \frac{5 \log\left(\cos\left(\frac{1}{2}(a + bx)\right)\right)}{2b} - \frac{5 \log\left(\sin\left(\frac{1}{2}(a + bx)\right)\right)}{2b} + \frac{\sec^2\left(\frac{1}{2}(a + bx)\right)}{8b}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[a + b*x]^3*Cot[a + b*x]^3,x]
```

```
[Out] (-9*Cos[a + b*x])/(4*b) - Cos[3*(a + b*x)]/(12*b) - Csc[(a + b*x)/2]^2/(8*b)
) + (5*Log[Cos[(a + b*x)/2]])/(2*b) - (5*Log[Sin[(a + b*x)/2]])/(2*b) + Sec
[(a + b*x)/2]^2/(8*b)
```

Maple [A]

time = 0.06, size = 70, normalized size = 1.06

method	result
derivativedivides	$-\frac{\frac{\cos^7(bx+a)}{2 \sin(bx+a)^2} - \frac{\cos^5(bx+a)}{2} - \frac{5(\cos^3(bx+a))}{6} - \frac{5 \cos(bx+a)}{2} - \frac{5 \ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$
default	$-\frac{\frac{\cos^7(bx+a)}{2 \sin(bx+a)^2} - \frac{\cos^5(bx+a)}{2} - \frac{5(\cos^3(bx+a))}{6} - \frac{5 \cos(bx+a)}{2} - \frac{5 \ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$
norman	$-\frac{1}{8b} + \frac{\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{75\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} - \frac{65\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{12b} - \frac{55\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} - \frac{5 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b} \right. \\ \left. \frac{1}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^3 \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)^2}$

risch	$-\frac{e^{3i(bx+a)}}{24b} - \frac{9e^{i(bx+a)}}{8b} - \frac{9e^{-i(bx+a)}}{8b} - \frac{e^{-3i(bx+a)}}{24b} + \frac{e^{3i(bx+a)} + e^{i(bx+a)}}{b(e^{2i(bx+a)} - 1)^2} - \frac{5\ln(e^{i(bx+a)} - 1)}{2b} + \frac{5\ln(e^{i(bx+a)} + 1)}{2b}$
-------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^6/sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/2*\cos(b*x+a)^7/\sin(b*x+a)^2-1/2*\cos(b*x+a)^5-5/6*\cos(b*x+a)^3-5/2*\cos(b*x+a)-5/2*\ln(\csc(b*x+a)-\cot(b*x+a)))$

Maxima [A]

time = 0.28, size = 66, normalized size = 1.00

$$\frac{4 \cos(bx+a)^3 - \frac{6 \cos(bx+a)}{\cos(bx+a)^2-1} + 24 \cos(bx+a) - 15 \log(\cos(bx+a) + 1) + 15 \log(\cos(bx+a) - 1)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^3,x, algorithm="maxima")`

[Out] $-1/12*(4*\cos(b*x+a)^3 - 6*\cos(b*x+a)/(\cos(b*x+a)^2 - 1) + 24*\cos(b*x+a) - 15*\log(\cos(b*x+a) + 1) + 15*\log(\cos(b*x+a) - 1))/b$

Fricas [A]

time = 0.38, size = 93, normalized size = 1.41

$$\frac{4 \cos(bx+a)^5 + 20 \cos(bx+a)^3 - 15 (\cos(bx+a)^2 - 1) \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 15 (\cos(bx+a)^2 - 1) \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 30 \cos(bx+a)}{12(b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $-1/12*(4*\cos(b*x+a)^5 + 20*\cos(b*x+a)^3 - 15*(\cos(b*x+a)^2 - 1)*\log(1/2*\cos(b*x+a) + 1/2) + 15*(\cos(b*x+a)^2 - 1)*\log(-1/2*\cos(b*x+a) + 1/2) - 30*\cos(b*x+a))/(b*\cos(b*x+a)^2 - b)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 719 vs. 2(58) = 116.

time = 2.53, size = 719, normalized size = 10.89

--

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**6/sin(b*x+a)**3,x)`

[Out] `Piecewise((-60*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(24*b*tan(a/2 + b*x/2)**8 + 72*b*tan(a/2 + b*x/2)**6 + 72*b*tan(a/2 + b*x/2)**4 + 24*b*tan(a/2 + b*x/2)**2) - 180*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(24*b*tan(a/2 + b*x/2)**8 + 72*b*tan(a/2 + b*x/2)**6 + 72*b*tan(a/2 + b*x/2)**4 + 24*b*tan(a/2 + b*x/2)**2))`

$2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) - 180*\log(\tan(a/2 + b*x/2))*\tan(a/2 + b*x/2)**4/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) - 60*\log(\tan(a/2 + b*x/2))*\tan(a/2 + b*x/2)**2/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) + 3*\tan(a/2 + b*x/2)**10/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) - 165*\tan(a/2 + b*x/2)**6/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) - 225*\tan(a/2 + b*x/2)**4/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) - 130*\tan(a/2 + b*x/2)**2/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2) - 3/(24*b*\tan(a/2 + b*x/2)**8 + 72*b*\tan(a/2 + b*x/2)**6 + 72*b*\tan(a/2 + b*x/2)**4 + 24*b*\tan(a/2 + b*x/2)**2), Ne(b, 0)), (x*cos(a)**6/sin(a)**3, True))$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(58) = 116.

time = 3.96, size = 163, normalized size = 2.47

$$\frac{3 \left(\frac{10(\cos(bx+a)-1)}{\cos(bx+a)+1} + 1 \right) (\cos(bx+a)+1)}{\cos(bx+a)-1} - \frac{3(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{16 \left(\frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{9(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 7 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right)^3} - 30 \log \left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right)}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a)^3,x, algorithm="giac")

[Out] $1/24*(3*(10*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)*(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - 3*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 16*(12*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 9*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 - 7)/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)^3 - 30*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)))/b$

Mupad [B]

time = 1.35, size = 129, normalized size = 1.95

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8b} - \frac{5 \ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{2b} - \frac{\frac{49 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6}{8} + \frac{67 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{8} + \frac{121 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{24} + \frac{1}{8}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 + 3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + 3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6/sin(a + b*x)^3,x)

[Out] $\tan(a/2 + (b*x)/2)^2/(8*b) - (5*\log(\tan(a/2 + (b*x)/2)))/(2*b) - ((121*\tan(a/2 + (b*x)/2)^2)/24 + (67*\tan(a/2 + (b*x)/2)^4)/8 + (49*\tan(a/2 + (b*x)/2)^6)/8 + 1/8)/(b*(\tan(a/2 + (b*x)/2)^2 + 3*\tan(a/2 + (b*x)/2)^4 + 3*\tan(a/2 + (b*x)/2)^6 + \tan(a/2 + (b*x)/2)^8))$

3.147 $\int \cos^2(a + bx) \cot^3(a + bx) dx$

Optimal. Leaf size=43

$$-\frac{\csc^2(a + bx)}{2b} - \frac{2 \log(\sin(a + bx))}{b} + \frac{\sin^2(a + bx)}{2b}$$

[Out] $-1/2*\csc(b*x+a)^2/b-2*\ln(\sin(b*x+a))/b+1/2*\sin(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2670, 272, 45}

$$\frac{\sin^2(a + bx)}{2b} - \frac{\csc^2(a + bx)}{2b} - \frac{2 \log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Cot[a + b*x]^3,x]

[Out] $-1/2*\text{Csc}[a + b*x]^2/b - (2*\text{Log}[\text{Sin}[a + b*x]])/b + \text{Sin}[a + b*x]^2/(2*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \cos^2(a + bx) \cot^3(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x^3} dx, x, -\sin(a + bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1-x)^2}{x^2} dx, x, \sin^2(a + bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^2} - \frac{2}{x}\right) dx, x, \sin^2(a + bx)\right)}{2b} \\
&= -\frac{\csc^2(a + bx)}{2b} - \frac{2 \log(\sin(a + bx))}{b} + \frac{\sin^2(a + bx)}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 35, normalized size = 0.81

$$-\frac{\csc^2(a + bx) + 4 \log(\sin(a + bx)) - \sin^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^2*Cot[a + b*x]^3,x]``[Out] -1/2*(Csc[a + b*x]^2 + 4*Log[Sin[a + b*x]] - Sin[a + b*x]^2)/b`**Maple [A]**

time = 0.04, size = 53, normalized size = 1.23

method	result	size
derivativedivides	$\frac{-\frac{\cos^6(bx+a)}{2 \sin(bx+a)^2} - \frac{(\cos^4(bx+a))}{2} - (\cos^2(bx+a)) - 2 \ln(\sin(bx+a))}{b}$	53
default	$\frac{-\frac{\cos^6(bx+a)}{2 \sin(bx+a)^2} - \frac{(\cos^4(bx+a))}{2} - (\cos^2(bx+a)) - 2 \ln(\sin(bx+a))}{b}$	53
risch	$2ix - \frac{e^{2i(bx+a)}}{8b} - \frac{e^{-2i(bx+a)}}{8b} + \frac{4ia}{b} + \frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}-1)^2} - \frac{2 \ln(e^{2i(bx+a)}-1)}{b}$	85
norman	$\frac{-\frac{1}{8b} - \frac{\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} + \frac{9\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} - \frac{2 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{2 \ln\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$	101

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5/sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/2*cos(b*x+a)^6/sin(b*x+a)^2-1/2*cos(b*x+a)^4-cos(b*x+a)^2-2*ln(sin(b*x+a)))`

Maxima [A]

time = 0.29, size = 35, normalized size = 0.81

$$\frac{\sin (b x+a)^2-\frac{1}{\sin (b x+a)^2}-2 \log (\sin (b x+a)^2)}{2 b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5/sin(b*x+a)^3,x, algorithm="maxima")``[Out] 1/2*(sin(b*x + a)^2 - 1/sin(b*x + a)^2 - 2*log(sin(b*x + a)^2))/b`**Fricas [A]**

time = 0.36, size = 61, normalized size = 1.42

$$\frac{2 \cos (b x+a)^4-3 \cos (b x+a)^2+8(\cos (b x+a)^2-1) \log \left(\frac{1}{2} \sin (b x+a)\right)-1}{4\left(b \cos (b x+a)^2-b\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5/sin(b*x+a)^3,x, algorithm="fricas")``[Out] -1/4*(2*cos(b*x + a)^4 - 3*cos(b*x + a)^2 + 8*(cos(b*x + a)^2 - 1)*log(1/2* sin(b*x + a)) - 1)/(b*cos(b*x + a)^2 - b)`**Sympy [B]** Leaf count of result is larger than twice the leaf count of optimal. 614 vs. $2(34) = 68$.

time = 1.50, size = 614, normalized size = 14.28

$$\frac{\cos (a+b x)}{\sin (a+b x)^2}+\frac{\cos (a+b x)}{\sin (a+b x)^4}+\frac{\cos (a+b x)}{\sin (a+b x)^6}+\frac{\cos (a+b x)}{\sin (a+b x)^8}+\frac{\cos (a+b x)}{\sin (a+b x)^{10}}+\frac{\cos (a+b x)}{\sin (a+b x)^{12}}+\frac{\cos (a+b x)}{\sin (a+b x)^{14}}+\frac{\cos (a+b x)}{\sin (a+b x)^{16}}+\frac{\cos (a+b x)}{\sin (a+b x)^{18}}+\frac{\cos (a+b x)}{\sin (a+b x)^{20}}+\frac{\cos (a+b x)}{\sin (a+b x)^{22}}+\frac{\cos (a+b x)}{\sin (a+b x)^{24}}+\frac{\cos (a+b x)}{\sin (a+b x)^{26}}+\frac{\cos (a+b x)}{\sin (a+b x)^{28}}+\frac{\cos (a+b x)}{\sin (a+b x)^{30}}+\frac{\cos (a+b x)}{\sin (a+b x)^{32}}+\frac{\cos (a+b x)}{\sin (a+b x)^{34}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)**5/sin(b*x+a)**3,x)`
`[Out] Piecewise((16*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**6/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 32*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 16*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 16*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 32*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 16*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - tan(a/2 + b*x/2)**8/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 18*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a`

/2 + b*x/2)**2) - 1/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2), Ne(b, 0)), (x*cos(a)**5/sin(a)**3, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(39) = 78.

time = 3.59, size = 187, normalized size = 4.35

$$\frac{\frac{8 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right) (\cos(bx+a)+1)}{\cos(bx+a)-1} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + \frac{8 \left(\frac{4 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - \frac{3 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} \right)^2 - 3 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right)^2} \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right)^2} - 8 \log \left(\frac{|\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right) + 16 \log \left(\left| -\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right| \right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/8*((8*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)*(cos(b*x + a) + 1)/(cos(b*x + a) - 1) + (cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 8*(4*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 3*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 3)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)^2 - 8*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)) + 16*log(abs(-(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1))))/b

Mupad [B]

time = 0.46, size = 62, normalized size = 1.44

$$\frac{\ln(\tan(a + bx)^2 + 1)}{b} - \frac{2 \ln(\tan(a + bx))}{b} - \frac{\tan(a + bx)^2 + \frac{1}{2}}{b(\tan(a + bx)^4 + \tan(a + bx)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5/sin(a + b*x)^3,x)

[Out] log(tan(a + b*x)^2 + 1)/b - (2*log(tan(a + b*x)))/b - (tan(a + b*x)^2 + 1/2)/(b*(tan(a + b*x)^2 + tan(a + b*x)^4))

3.148 $\int \cos(a + bx) \cot^3(a + bx) dx$

Optimal. Leaf size=49

$$\frac{3 \tanh^{-1}(\cos(a + bx))}{2b} - \frac{3 \cos(a + bx)}{2b} - \frac{\cos(a + bx) \cot^2(a + bx)}{2b}$$

[Out] 3/2*arctanh(cos(b*x+a))/b-3/2*cos(b*x+a)/b-1/2*cos(b*x+a)*cot(b*x+a)^2/b

Rubi [A]

time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2672, 294, 327, 212}

$$-\frac{3 \cos(a + bx)}{2b} - \frac{\cos(a + bx) \cot^2(a + bx)}{2b} + \frac{3 \tanh^{-1}(\cos(a + bx))}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Cot[a + b*x]^3,x]

[Out] (3*ArcTanh[Cos[a + b*x]])/(2*b) - (3*Cos[a + b*x])/(2*b) - (Cos[a + b*x]*Cot[a + b*x]^2)/(2*b)

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !ILtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 327

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2672

```
Int[((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(
ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x]
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \cot^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^4}{(1-x^2)^2} dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\cos(a + bx) \cot^2(a + bx)}{2b} + \frac{3\text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \cos(a + bx)\right)}{2b} \\ &= -\frac{3 \cos(a + bx)}{2b} - \frac{\cos(a + bx) \cot^2(a + bx)}{2b} + \frac{3\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(a + bx)\right)}{2b} \\ &= \frac{3 \tanh^{-1}(\cos(a + bx))}{2b} - \frac{3 \cos(a + bx)}{2b} - \frac{\cos(a + bx) \cot^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 86, normalized size = 1.76

$$-\frac{\cos(a + bx)}{b} - \frac{\csc^2\left(\frac{1}{2}(a + bx)\right)}{8b} + \frac{3 \log\left(\cos\left(\frac{1}{2}(a + bx)\right)\right)}{2b} - \frac{3 \log\left(\sin\left(\frac{1}{2}(a + bx)\right)\right)}{2b} + \frac{\sec^2\left(\frac{1}{2}(a + bx)\right)}{8b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Cot[a + b*x]^3,x]

[Out] -(Cos[a + b*x]/b) - Csc[(a + b*x)/2]^2/(8*b) + (3*Log[Cos[(a + b*x)/2]])/(2*b) - (3*Log[Sin[(a + b*x)/2]])/(2*b) + Sec[(a + b*x)/2]^2/(8*b)

Maple [A]

time = 0.04, size = 60, normalized size = 1.22

method	result	size
derivativedivides	$\frac{-\frac{\cos^5(bx+a)}{2 \sin(bx+a)^2} - \frac{(\cos^3(bx+a))}{2} - \frac{3 \cos(bx+a)}{2} - \frac{3 \ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$	60
default	$\frac{-\frac{\cos^5(bx+a)}{2 \sin(bx+a)^2} - \frac{(\cos^3(bx+a))}{2} - \frac{3 \cos(bx+a)}{2} - \frac{3 \ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$	60
norman	$\frac{-\frac{1}{8b} + \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{9\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b}}{\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b}$	82

risch	$-\frac{e^{i(bx+a)}}{2b} - \frac{e^{-i(bx+a)}}{2b} + \frac{e^{3i(bx+a)} + e^{i(bx+a)}}{b(e^{2i(bx+a)} - 1)^2} - \frac{3 \ln(e^{i(bx+a)} - 1)}{2b} + \frac{3 \ln(e^{i(bx+a)} + 1)}{2b}$	100
-------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^4/sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $1/b * (-1/2 * \cos(b*x+a)^5 / \sin(b*x+a)^2 - 1/2 * \cos(b*x+a)^3 - 3/2 * \cos(b*x+a) - 3/2 * \ln(\csc(b*x+a) - \cot(b*x+a)))$

Maxima [A]

time = 0.29, size = 56, normalized size = 1.14

$$\frac{\frac{2 \cos(bx+a)}{\cos(bx+a)^2 - 1} - 4 \cos(bx+a) + 3 \log(\cos(bx+a) + 1) - 3 \log(\cos(bx+a) - 1)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4/sin(b*x+a)^3,x, algorithm="maxima")`

[Out] $1/4 * (2 * \cos(b*x + a) / (\cos(b*x + a)^2 - 1) - 4 * \cos(b*x + a) + 3 * \log(\cos(b*x + a) + 1) - 3 * \log(\cos(b*x + a) - 1)) / b$

Fricas [A]

time = 0.41, size = 83, normalized size = 1.69

$$\frac{4 \cos(bx+a)^3 - 3(\cos(bx+a)^2 - 1) \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 3(\cos(bx+a)^2 - 1) \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 6 \cos(bx+a)}{4(b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4/sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $-1/4 * (4 * \cos(b*x + a)^3 - 3 * (\cos(b*x + a)^2 - 1) * \log(1/2 * \cos(b*x + a) + 1/2) + 3 * (\cos(b*x + a)^2 - 1) * \log(-1/2 * \cos(b*x + a) + 1/2) - 6 * \cos(b*x + a)) / (b * \cos(b*x + a)^2 - b)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 241 vs. 2(42) = 84.

time = 0.94, size = 241, normalized size = 4.92

$$\begin{cases} -\frac{12 \log\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right) \tan^4\left(\frac{a}{2} + \frac{bx}{2}\right) - 12 \log\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right) \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right) + \frac{\tan^6\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b \tan^4\left(\frac{a}{2} + \frac{bx}{2}\right) + 8b \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)} - \frac{18 \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b \tan^4\left(\frac{a}{2} + \frac{bx}{2}\right) + 8b \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)} - \frac{1}{8b \tan^4\left(\frac{a}{2} + \frac{bx}{2}\right) + 8b \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)} & \text{for } b \neq 0 \\ \frac{x \cos^4(a)}{\sin^3(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**4/sin(b*x+a)**3,x)`

[Out] $\text{Piecewise}((-12 * \log(\tan(a/2 + b*x/2)) * \tan(a/2 + b*x/2)**4 / (8*b*\tan(a/2 + b*x/2)**4 + 8*b*\tan(a/2 + b*x/2)**2) - 12 * \log(\tan(a/2 + b*x/2)) * \tan(a/2 + b*x/2)$

2)**2/(8*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + tan(a/2 + b*x/2)
)**6/(8*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 18*tan(a/2 + b*x
 /2)**2/(8*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 1/(8*b*tan(a/2
 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2), Ne(b, 0)), (x*cos(a)**4/sin(a)**3,
 True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(43) = 86.

time = 4.22, size = 140, normalized size = 2.86

$$\frac{\frac{14(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 1}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2}} + \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 6 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)$$

$$8b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^3,x, algorithm="giac")

[Out] -1/8*((14*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 3*(cos(b*x + a) - 1)^2/(c
 os(b*x + a) + 1)^2 - 1)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - (cos(b*x +
 a) - 1)^2/(cos(b*x + a) + 1)^2) + (cos(b*x + a) - 1)/(cos(b*x + a) + 1) +
 6*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.54, size = 77, normalized size = 1.57

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8b} - \frac{3 \ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{2b} - \frac{\frac{17 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8} + \frac{1}{8}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4/sin(a + b*x)^3,x)

[Out] tan(a/2 + (b*x)/2)^2/(8*b) - (3*log(tan(a/2 + (b*x)/2)))/(2*b) - ((17*tan(a
 /2 + (b*x)/2)^2)/8 + 1/8)/(b*(tan(a/2 + (b*x)/2)^2 + tan(a/2 + (b*x)/2)^4))

3.149 $\int \cot^3(a + bx) dx$

Optimal. Leaf size=28

$$-\frac{\cot^2(a + bx)}{2b} - \frac{\log(\sin(a + bx))}{b}$$

[Out] $-1/2*\cot(b*x+a)^2/b-\ln(\sin(b*x+a))/b$

Rubi [A]

time = 0.01, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$-\frac{\cot^2(a + bx)}{2b} - \frac{\log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[a + b*x]^3, x]$

[Out] $-1/2*\text{Cot}[a + b*x]^2/b - \text{Log}[\text{Sin}[a + b*x]]/b$

Rule 3554

$\text{Int}[(b_.)*\tan[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*((b*\text{Tan}[c + d*x])^{(n - 1)}/(d*(n - 1))), x] - \text{Dist}[b^2, \text{Int}[(b*\text{Tan}[c + d*x])^{(n - 2)}, x], x] /;$ $\text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1]$

Rule 3556

$\text{Int}[\tan[(c_.) + (d_.)*(x_.)], x_Symbol] \rightarrow \text{Simp}[-\text{Log}[\text{RemoveContent}[\text{Cos}[c + d*x], x]]/d, x] /;$ $\text{FreeQ}\{c, d\}, x$

Rubi steps

$$\begin{aligned} \int \cot^3(a + bx) dx &= -\frac{\cot^2(a + bx)}{2b} - \int \cot(a + bx) dx \\ &= -\frac{\cot^2(a + bx)}{2b} - \frac{\log(\sin(a + bx))}{b} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 34, normalized size = 1.21

$$-\frac{\cot^2(a + bx) + 2 \log(\cos(a + bx)) + 2 \log(\tan(a + bx))}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^3,x]

[Out] $-1/2*(\text{Cot}[a + b*x]^2 + 2*\text{Log}[\text{Cos}[a + b*x]] + 2*\text{Log}[\text{Tan}[a + b*x]])/b$

Maple [A]

time = 0.03, size = 25, normalized size = 0.89

method	result	size
derivativedivides	$-\frac{(\cot^2(bx+a)) - \ln(\sin(bx+a))}{2b}$	25
default	$-\frac{(\cot^2(bx+a)) - \ln(\sin(bx+a))}{2b}$	25
risch	$ix + \frac{2ia}{b} + \frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}-1)^2} - \frac{\ln(e^{2i(bx+a)}-1)}{b}$	57
norman	$-\frac{1}{8b} - \frac{\tan^4(\frac{bx+a}{2})}{8b} + \frac{\ln(1+\tan^2(\frac{bx+a}{2}))}{b} - \frac{\ln(\tan(\frac{bx+a}{2}))}{b}$	69

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3/sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] $1/b*(-1/2*\cot(b*x+a)^2 - \ln(\sin(b*x+a)))$

Maxima [A]

time = 0.31, size = 23, normalized size = 0.82

$$-\frac{1}{\sin(bx+a)^2} + \log(\sin(bx+a)^2)$$

$$2b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^3,x, algorithm="maxima")

[Out] $-1/2*(1/\sin(b*x + a)^2 + \log(\sin(b*x + a)^2))/b$

Fricas [A]

time = 0.36, size = 41, normalized size = 1.46

$$\frac{2(\cos(bx+a)^2 - 1)\log(\frac{1}{2}\sin(bx+a)) - 1}{2(b\cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^3,x, algorithm="fricas")

[Out] $-1/2*(2*(\cos(b*x + a)^2 - 1)*\log(1/2*\sin(b*x + a)) - 1)/(b*\cos(b*x + a)^2 - b)$

Sympy [A]

time = 0.35, size = 42, normalized size = 1.50

$$\begin{cases} -\frac{\log(\sin(a+bx))}{b} - \frac{\cos^2(a+bx)}{2b\sin^2(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^3(a)}{\sin^3(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3/sin(b*x+a)**3,x)**[Out]** Piecewise((-log(sin(a + b*x))/b - cos(a + b*x)**2/(2*b*sin(a + b*x)**2), Ne(b, 0)), (x*cos(a)**3/sin(a)**3, True))**Giac [A]**

time = 4.44, size = 36, normalized size = 1.29

$$\frac{\frac{\sin(bx+a)^2-1}{\sin(bx+a)^2} - \log(\sin(bx+a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^3,x, algorithm="giac")**[Out]** 1/2*((sin(b*x + a)^2 - 1)/sin(b*x + a)^2 - log(sin(b*x + a)^2))/b**Mupad [B]**

time = 0.43, size = 36, normalized size = 1.29

$$\frac{\cot(a+bx)^2 - \ln(\tan(a+bx)^2 + 1) + 2 \ln(\tan(a+bx))}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3/sin(a + b*x)^3,x)**[Out]** -(2*log(tan(a + b*x)) - log(tan(a + b*x)^2 + 1) + cot(a + b*x)^2)/(2*b)

3.150 $\int \cot^2(a + bx) \csc(a + bx) dx$

Optimal. Leaf size=34

$$\frac{\tanh^{-1}(\cos(a + bx))}{2b} - \frac{\cot(a + bx) \csc(a + bx)}{2b}$$

[Out] 1/2*arctanh(cos(b*x+a))/b-1/2*cot(b*x+a)*csc(b*x+a)/b

Rubi [A]

time = 0.02, antiderivative size = 34, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2691, 3855}

$$\frac{\tanh^{-1}(\cos(a + bx))}{2b} - \frac{\cot(a + bx) \csc(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^2*Csc[a + b*x],x]

[Out] ArcTanh[Cos[a + b*x]]/(2*b) - (Cot[a + b*x]*Csc[a + b*x])/(2*b)

Rule 2691

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegersQ[2*m, 2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] := Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \cot^2(a + bx) \csc(a + bx) dx &= -\frac{\cot(a + bx) \csc(a + bx)}{2b} - \frac{1}{2} \int \csc(a + bx) dx \\ &= \frac{\tanh^{-1}(\cos(a + bx))}{2b} - \frac{\cot(a + bx) \csc(a + bx)}{2b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 75 vs. 2(34) = 68.

time = 0.02, size = 75, normalized size = 2.21

$$-\frac{\csc^2\left(\frac{1}{2}(a+bx)\right)}{8b} + \frac{\log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{2b} - \frac{\log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{2b} + \frac{\sec^2\left(\frac{1}{2}(a+bx)\right)}{8b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^2*Csc[a + b*x], x]

[Out] -1/8*Csc[(a + b*x)/2]^2/b + Log[Cos[(a + b*x)/2]]/(2*b) - Log[Sin[(a + b*x)/2]]/(2*b) + Sec[(a + b*x)/2]^2/(8*b)

Maple [A]

time = 0.03, size = 50, normalized size = 1.47

method	result	size
derivativedivides	$\frac{-\frac{\cos^3(bx+a)}{2\sin(bx+a)^2} - \frac{\cos(bx+a)}{2} - \frac{\ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$	50
default	$\frac{-\frac{\cos^3(bx+a)}{2\sin(bx+a)^2} - \frac{\cos(bx+a)}{2} - \frac{\ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$	50
norman	$\frac{-\frac{1}{8b} + \frac{\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b}$	51
risch	$\frac{e^{3i(bx+a)} + e^{i(bx+a)}}{b(e^{2i(bx+a)} - 1)^2} - \frac{\ln(e^{i(bx+a)} - 1)}{2b} + \frac{\ln(e^{i(bx+a)} + 1)}{2b}$	72

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2/sin(b*x+a)^3, x, method=_RETURNVERBOSE)

[Out] 1/b*(-1/2*cos(b*x+a)^3/sin(b*x+a)^2-1/2*cos(b*x+a)-1/2*ln(csc(b*x+a)-cot(b*x+a)))

Maxima [A]

time = 0.28, size = 46, normalized size = 1.35

$$\frac{\frac{2\cos(bx+a)}{\cos(bx+a)^2-1} + \log(\cos(bx+a)+1) - \log(\cos(bx+a)-1)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^3, x, algorithm="maxima")

[Out] 1/4*(2*cos(b*x + a)/(cos(b*x + a)^2 - 1) + log(cos(b*x + a) + 1) - log(cos(b*x + a) - 1))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(30) = 60.

time = 0.37, size = 72, normalized size = 2.12

$$\frac{(\cos(bx+a)^2-1)\log\left(\frac{1}{2}\cos(bx+a)+\frac{1}{2}\right) - (\cos(bx+a)^2-1)\log\left(-\frac{1}{2}\cos(bx+a)+\frac{1}{2}\right) + 2\cos(bx+a)}{4(b\cos(bx+a)^2-b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^2/sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $\frac{1}{4} * ((\cos(b*x + a)^2 - 1) * \log(1/2 * \cos(b*x + a) + 1/2) - (\cos(b*x + a)^2 - 1) * \log(-1/2 * \cos(b*x + a) + 1/2) + 2 * \cos(b*x + a)) / (b * \cos(b*x + a)^2 - b)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(27) = 54$.

time = 0.53, size = 58, normalized size = 1.71

$$\begin{cases} -\frac{\log\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{2b} + \frac{\tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b} - \frac{1}{8b \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)} & \text{for } b \neq 0 \\ \frac{x \cos^2(a)}{\sin^3(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**2/sin(b*x+a)**3,x)`

[Out] `Piecewise((-log(tan(a/2 + b*x/2))/(2*b) + tan(a/2 + b*x/2)**2/(8*b) - 1/(8*b*tan(a/2 + b*x/2)**2), Ne(b, 0)), (x*cos(a)**2/sin(a)**3, True))`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 93 vs. $2(30) = 60$.

time = 3.17, size = 93, normalized size = 2.74

$$\frac{\frac{\left(\frac{2(\cos(bx+a)-1)}{\cos(bx+a)+1} + 1\right)(\cos(bx+a)+1)}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 2 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^2/sin(b*x+a)^3,x, algorithm="giac")`

[Out] $\frac{1}{8} * ((2 * (\cos(b*x + a) - 1) / (\cos(b*x + a) + 1) + 1) * (\cos(b*x + a) + 1) / (\cos(b*x + a) - 1) - (\cos(b*x + a) - 1) / (\cos(b*x + a) + 1) - 2 * \log(\text{abs}(-\cos(b*x + a) + 1) / \text{abs}(\cos(b*x + a) + 1))) / b$

Mupad [B]

time = 0.45, size = 48, normalized size = 1.41

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8b} - \frac{1}{8b \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2} - \frac{\ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^2/sin(a + b*x)^3,x)`

[Out] $\tan(a/2 + (b*x)/2)^2/(8*b) - 1/(8*b*\tan(a/2 + (b*x)/2)^2) - \log(\tan(a/2 + (b*x)/2))/(2*b)$

3.151 $\int \cot(a + bx) \csc^2(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\csc^2(a + bx)}{2b}$$

[Out] -1/2*csc(b*x+a)^2/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2686, 30}

$$-\frac{\csc^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]*Csc[a + b*x]^2,x]

[Out] -1/2*Csc[a + b*x]^2/b

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \cot(a + bx) \csc^2(a + bx) dx &= -\frac{\text{Subst}(\int x dx, x, \csc(a + bx))}{b} \\ &= -\frac{\csc^2(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$-\frac{\csc^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]*Csc[a + b*x]^2,x]

[Out] $-1/2*\text{Csc}[a + b*x]^2/b$

Maple [A]

time = 0.02, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$-\frac{1}{2\sin(bx+a)^2b}$	14
default	$-\frac{1}{2\sin(bx+a)^2b}$	14
risch	$\frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}-1)^2}$	28
norman	$-\frac{\frac{1}{8b} - \frac{\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2}$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)/sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] $-1/2/\sin(b*x+a)^2/b$

Maxima [A]

time = 0.28, size = 13, normalized size = 0.87

$$-\frac{1}{2b\sin(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^3,x, algorithm="maxima")

[Out] $-1/2/(b*\sin(b*x + a)^2)$

Fricas [A]

time = 0.33, size = 18, normalized size = 1.20

$$\frac{1}{2(b\cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^3,x, algorithm="fricas")

[Out] $1/2/(b*\cos(b*x + a)^2 - b)$

Sympy [A]

time = 0.35, size = 24, normalized size = 1.60

$$\begin{cases} -\frac{1}{2b\sin^2(a+bx)} & \text{for } b \neq 0 \\ \frac{x\cos(a)}{\sin^3(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)/sin(b*x+a)**3,x)``[Out] Piecewise((-1/(2*b*sin(a + b*x)**2), Ne(b, 0)), (x*cos(a)/sin(a)**3, True))`**Giac [A]**

time = 4.89, size = 13, normalized size = 0.87

$$-\frac{1}{2b\sin(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)/sin(b*x+a)^3,x, algorithm="giac")``[Out] -1/2/(b*sin(b*x + a)^2)`**Mupad [B]**

time = 0.39, size = 13, normalized size = 0.87

$$-\frac{1}{2b\sin(a+bx)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)/sin(a + b*x)^3,x)``[Out] -1/(2*b*sin(a + b*x)^2)`

3.152 $\int \csc^3(a + bx) \sec(a + bx) dx$

Optimal. Leaf size=27

$$-\frac{\cot^2(a + bx)}{2b} + \frac{\log(\tan(a + bx))}{b}$$

[Out] $-1/2*\cot(b*x+a)^2/b+\ln(\tan(b*x+a))/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2700, 14}

$$\frac{\log(\tan(a + bx))}{b} - \frac{\cot^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[a + b*x]^3*\text{Sec}[a + b*x], x]$

[Out] $-1/2*\text{Cot}[a + b*x]^2/b + \text{Log}[\text{Tan}[a + b*x]]/b$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{InverseFunctionQ}[v]$

Rule 2700

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^{(m_.)}*\text{sec}[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[1/f, \text{Subst}[\text{Int}[(1 + x^2)^{(m+n)/2 - 1}/x^m, x], x, \text{Tan}[e + f*x]], x] /; \text{FreeQ}\{e, f\}, x \ \&\& \ \text{IntegersQ}[m, n, (m+n)/2]$

Rubi steps

$$\begin{aligned} \int \csc^3(a + bx) \sec(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{1+x^2}{x^3} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(\frac{1}{x^3} + \frac{1}{x}\right) dx, x, \tan(a + bx)\right)}{b} \\ &= -\frac{\cot^2(a + bx)}{2b} + \frac{\log(\tan(a + bx))}{b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 34, normalized size = 1.26

$$\frac{\csc^2(a + bx) + 2 \log(\cos(a + bx)) - 2 \log(\sin(a + bx))}{2b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^3*Sec[a + b*x],x]``[Out] -1/2*(Csc[a + b*x]^2 + 2*Log[Cos[a + b*x]] - 2*Log[Sin[a + b*x]])/b`**Maple [A]**

time = 0.06, size = 23, normalized size = 0.85

method	result	size
derivativedivides	$-\frac{\frac{1}{2 \sin^2(bx+a)} + \ln(\tan(bx+a))}{b}$	23
default	$-\frac{\frac{1}{2 \sin^2(bx+a)} + \ln(\tan(bx+a))}{b}$	23
risch	$\frac{2e^{2i(bx+a)}}{b(e^{2i(bx+a)}-1)^2} - \frac{\ln(e^{2i(bx+a)}+1)}{b} + \frac{\ln(e^{2i(bx+a)}-1)}{b}$	62
norman	$-\frac{\frac{1}{8b} - \frac{\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b}$	84

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)/sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/2/sin(b*x+a)^2+ln(tan(b*x+a)))`**Maxima [A]**

time = 0.29, size = 36, normalized size = 1.33

$$\frac{\frac{1}{\sin^2(bx+a)} + \log(\sin(bx+a)^2 - 1) - \log(\sin(bx+a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)/sin(b*x+a)^3,x, algorithm="maxima")``[Out] -1/2*(1/sin(b*x + a)^2 + log(sin(b*x + a)^2 - 1) - log(sin(b*x + a)^2))/b`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(25) = 50$.

time = 0.36, size = 65, normalized size = 2.41

$$\frac{(\cos(bx+a)^2 - 1) \log(\cos(bx+a)^2) - (\cos(bx+a)^2 - 1) \log\left(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}\right) - 1}{2(b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^3,x, algorithm="fricas")

[Out] $-1/2*((\cos(b*x + a)^2 - 1)*\log(\cos(b*x + a)^2) - (\cos(b*x + a)^2 - 1)*\log(-1/4*\cos(b*x + a)^2 + 1/4) - 1)/(b*\cos(b*x + a)^2 - b)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(a + bx)}{\sin^3(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)**3,x)

[Out] Integral(sec(a + b*x)/sin(a + b*x)**3, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(25) = 50.

time = 4.34, size = 119, normalized size = 4.41

$$\frac{\left(\frac{4(\cos(bx+a)-1)-1}{\cos(bx+a)+1}\right)(\cos(bx+a)+1) - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 4 \log\left(\frac{|\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) + 8 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^3,x, algorithm="giac")

[Out] $-1/8*((4*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)*(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 4*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)) + 8*\log(\text{abs}(-(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)))/b$

Mupad [B]

time = 0.05, size = 34, normalized size = 1.26

$$\frac{\ln(\cos(a + bx)) - \frac{\ln(\sin(a + bx)^2)}{2} + \frac{1}{2\sin(a + bx)^2}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)*sin(a + b*x)^3),x)

[Out] $-(\log(\cos(a + b*x)) - \log(\sin(a + b*x)^2)/2 + 1/(2*\sin(a + b*x)^2))/b$

3.153 $\int \csc^3(a + bx) \sec^2(a + bx) dx$

Optimal. Leaf size=49

$$-\frac{3 \tanh^{-1}(\cos(a + bx))}{2b} + \frac{3 \sec(a + bx)}{2b} - \frac{\csc^2(a + bx) \sec(a + bx)}{2b}$$

[Out] $-3/2*\operatorname{arctanh}(\cos(b*x+a))/b+3/2*\sec(b*x+a)/b-1/2*\csc(b*x+a)^2*\sec(b*x+a)/b$

Rubi [A]

time = 0.03, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2702, 294, 327, 213}

$$\frac{3 \sec(a + bx)}{2b} - \frac{3 \tanh^{-1}(\cos(a + bx))}{2b} - \frac{\csc^2(a + bx) \sec(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^3*Sec[a + b*x]^2,x]`

[Out] $(-3*\operatorname{ArcTanh}[\operatorname{Cos}[a + b*x]])/(2*b) + (3*\operatorname{Sec}[a + b*x])/(2*b) - (\operatorname{Csc}[a + b*x]^2*\operatorname{Sec}[a + b*x])/(2*b)$

Rule 213

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rule 294

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 327

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol]
:> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^3(a + bx) \sec^2(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^4}{(-1+x^2)^2} dx, x, \sec(a + bx)\right)}{b} \\ &= -\frac{\csc^2(a + bx) \sec(a + bx)}{2b} + \frac{3\text{Subst}\left(\int \frac{x^2}{-1+x^2} dx, x, \sec(a + bx)\right)}{2b} \\ &= \frac{3 \sec(a + bx)}{2b} - \frac{\csc^2(a + bx) \sec(a + bx)}{2b} + \frac{3\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sec(a + bx)\right)}{2b} \\ &= -\frac{3 \tanh^{-1}(\cos(a + bx))}{2b} + \frac{3 \sec(a + bx)}{2b} - \frac{\csc^2(a + bx) \sec(a + bx)}{2b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 143 vs. 2(49) = 98.

time = 0.18, size = 143, normalized size = 2.92

$$\frac{\csc^4(a + bx) (2 - 6 \cos(2(a + bx)) + 2 \cos(3(a + bx)) + 3 \cos(3(a + bx))) \log(\cos(\frac{1}{2}(a + bx))) - 3 \cos(3(a + bx)) \log(\sin(\frac{1}{2}(a + bx))) + \cos(a + bx) (-2 - 3 \log(\cos(\frac{1}{2}(a + bx))) + 3 \log(\sin(\frac{1}{2}(a + bx))))}{2b (\csc^2(\frac{1}{2}(a + bx)) - \sec^2(\frac{1}{2}(a + bx)))}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^3*Sec[a + b*x]^2,x]

[Out] (Csc[a + b*x]^4*(2 - 6*Cos[2*(a + b*x)] + 2*Cos[3*(a + b*x)] + 3*Cos[3*(a + b*x)]*Log[Cos[(a + b*x)/2]] - 3*Cos[3*(a + b*x)]*Log[Sin[(a + b*x)/2]] + Cos[a + b*x]*(-2 - 3*Log[Cos[(a + b*x)/2]] + 3*Log[Sin[(a + b*x)/2]]))/ (2*b*(Csc[(a + b*x)/2]^2 - Sec[(a + b*x)/2]^2))

Maple [A]

time = 0.07, size = 52, normalized size = 1.06

method	result	size
derivativedivides	$-\frac{1}{2 \sin(bx+a)^2 \cos(bx+a)} + \frac{3}{2 \cos(bx+a)} + \frac{3 \ln(\csc(bx+a) - \cot(bx+a))}{2b}$	52
default	$-\frac{1}{2 \sin(bx+a)^2 \cos(bx+a)} + \frac{3}{2 \cos(bx+a)} + \frac{3 \ln(\csc(bx+a) - \cot(bx+a))}{2b}$	52
norman	$\frac{\frac{1}{8b} + \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{9\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right) \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b}$	82

risch	$\frac{3e^{5i(bx+a)} - 2e^{3i(bx+a)} + 3e^{i(bx+a)}}{b(e^{2i(bx+a)} - 1)^2(e^{2i(bx+a)} + 1)} - \frac{3\ln(e^{i(bx+a)} + 1)}{2b} + \frac{3\ln(e^{i(bx+a)} - 1)}{2b}$	100
-------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^2/sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/2/\sin(b*x+a)^2/\cos(b*x+a)+3/2/\cos(b*x+a)+3/2*\ln(\csc(b*x+a)-\cot(b*x+a)))$

Maxima [A]

time = 0.29, size = 61, normalized size = 1.24

$$\frac{2(3\cos(bx+a)^2-2)}{\cos(bx+a)^3-\cos(bx+a)} - 3\log(\cos(bx+a)+1) + 3\log(\cos(bx+a)-1)$$

$4b$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a)^3,x, algorithm="maxima")`

[Out] $1/4*(2*(3*\cos(b*x+a)^2-2)/(\cos(b*x+a)^3-\cos(b*x+a))-3*\log(\cos(b*x+a)+1)+3*\log(\cos(b*x+a)-1))/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 96 vs. $2(43) = 86$.

time = 0.38, size = 96, normalized size = 1.96

$$\frac{6\cos(bx+a)^2-3(\cos(bx+a)^3-\cos(bx+a))\log(\frac{1}{2}\cos(bx+a)+\frac{1}{2})+3(\cos(bx+a)^3-\cos(bx+a))\log(-\frac{1}{2}\cos(bx+a)+\frac{1}{2})-4}{4(b\cos(bx+a)^3-b\cos(bx+a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $1/4*(6*\cos(b*x+a)^2-3*(\cos(b*x+a)^3-\cos(b*x+a))*\log(1/2*\cos(b*x+a)+1/2)+3*(\cos(b*x+a)^3-\cos(b*x+a))*\log(-1/2*\cos(b*x+a)+1/2)-4)/(b*\cos(b*x+a)^3-b*\cos(b*x+a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(a+bx)}{\sin^3(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**2/sin(b*x+a)**3,x)`

[Out] `Integral(sec(a+b*x)**2/sin(a+b*x)**3, x)`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 140 vs. 2(43) = 86.

time = 3.68, size = 140, normalized size = 2.86

$$\frac{\frac{14(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 1}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2}} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 6 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)$$

$8b$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^3,x, algorithm="giac")

[Out] 1/8*((14*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 3*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 1)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + (cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2) - (cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 6*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.03, size = 49, normalized size = 1.00

$$-\frac{3 \operatorname{atanh}(\cos(a + bx))}{2b} - \frac{\frac{3 \cos(a+bx)^2}{2} - 1}{b (\cos(a + bx) - \cos(a + bx)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^2*sin(a + b*x)^3),x)

[Out] - (3*atanh(cos(a + b*x)))/(2*b) - ((3*cos(a + b*x)^2)/2 - 1)/(b*(cos(a + b*x) - cos(a + b*x)^3))

3.154 $\int \csc^3(a + bx) \sec^3(a + bx) dx$

Optimal. Leaf size=43

$$-\frac{\cot^2(a + bx)}{2b} + \frac{2 \log(\tan(a + bx))}{b} + \frac{\tan^2(a + bx)}{2b}$$

[Out] $-1/2*\cot(b*x+a)^2/b+2*\ln(\tan(b*x+a))/b+1/2*\tan(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2700, 272, 45}

$$\frac{\tan^2(a + bx)}{2b} - \frac{\cot^2(a + bx)}{2b} + \frac{2 \log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^3*Sec[a + b*x]^3,x]

[Out] $-1/2*\cot[a + b*x]^2/b + (2*\log[\tan[a + b*x]])/b + \tan[a + b*x]^2/(2*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LtQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned}
\int \csc^3(a+bx) \sec^3(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{x^3} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x)^2}{x^2} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^2} + \frac{2}{x}\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= -\frac{\cot^2(a+bx)}{2b} + \frac{2 \log(\tan(a+bx))}{b} + \frac{\tan^2(a+bx)}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 61, normalized size = 1.42

$$8 \left(-\frac{\csc^2(a+bx)}{16b} - \frac{\log(\cos(a+bx))}{4b} + \frac{\log(\sin(a+bx))}{4b} + \frac{\sec^2(a+bx)}{16b} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^3*Sec[a + b*x]^3,x]``[Out] 8*(-1/16*Csc[a + b*x]^2/b - Log[Cos[a + b*x]]/(4*b) + Log[Sin[a + b*x]]/(4*b) + Sec[a + b*x]^2/(16*b))`**Maple [A]**

time = 0.06, size = 43, normalized size = 1.00

method	result
derivativedivides	$\frac{1}{2 \sin^2(bx+a) \cos^2(bx+a)} - \frac{1}{\sin^2(bx+a)} + 2 \ln(\tan(bx+a))$
default	$\frac{1}{2 \sin^2(bx+a) \cos^2(bx+a)} - \frac{1}{\sin^2(bx+a)} + 2 \ln(\tan(bx+a))$
risch	$\frac{4 e^{6i(bx+a)} + 4 e^{2i(bx+a)}}{b(e^{2i(bx+a)} + 1)^2 (e^{2i(bx+a)} - 1)^2} + \frac{2 \ln(e^{2i(bx+a)} - 1)}{b} - \frac{2 \ln(e^{2i(bx+a)} + 1)}{b}$
norman	$\frac{-\frac{1}{8b} - \frac{\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} + \frac{9\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{4b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^2 \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)} + \frac{2 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{2 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b} - \frac{2 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^3/sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/2/sin(b*x+a)^2/cos(b*x+a)^2-1/sin(b*x+a)^2+2*ln(tan(b*x+a)))`

Maxima [A]

time = 0.29, size = 64, normalized size = 1.49

$$\frac{\frac{2 \sin(bx+a)^2 - 1}{\sin(bx+a)^4 - \sin(bx+a)^2} + 2 \log(\sin(bx+a)^2 - 1) - 2 \log(\sin(bx+a)^2)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a)^3,x, algorithm="maxima")

[Out] -1/2*((2*sin(b*x + a)^2 - 1)/(sin(b*x + a)^4 - sin(b*x + a)^2) + 2*log(sin(b*x + a)^2 - 1) - 2*log(sin(b*x + a)^2))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 102 vs. 2(39) = 78.

time = 0.37, size = 102, normalized size = 2.37

$$\frac{2 \cos(bx+a)^2 - 2(\cos(bx+a)^4 - \cos(bx+a)^2) \log(\cos(bx+a)^2) + 2(\cos(bx+a)^4 - \cos(bx+a)^2) \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) - 1}{2(b \cos(bx+a)^4 - b \cos(bx+a)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/2*(2*cos(b*x + a)^2 - 2*(cos(b*x + a)^4 - cos(b*x + a)^2)*log(cos(b*x + a)^2) + 2*(cos(b*x + a)^4 - cos(b*x + a)^2)*log(-1/4*cos(b*x + a)^2 + 1/4) - 1)/(b*cos(b*x + a)^4 - b*cos(b*x + a)^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(a + bx)}{\sin^3(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3/sin(b*x+a)**3,x)

[Out] Integral(sec(a + b*x)**3/sin(a + b*x)**3, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(39) = 78.

time = 4.07, size = 188, normalized size = 4.37

$$\frac{\frac{8 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - \frac{8 \left(\frac{4 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + \frac{3 \frac{\cos(bx+a)-1}{\cos(bx+a)+1}^2 + 3}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^2} \right)}{8b} - 8 \log\left(\frac{|\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) + 16 \log\left(\left| -\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right| \right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a)^3,x, algorithm="giac")

```
[Out] -1/8*((8*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)*(cos(b*x + a) + 1)/(cos
(b*x + a) - 1) - (cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 8*(4*(cos(b*x + a)
- 1)/(cos(b*x + a) + 1) + 3*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 3)
/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^2 - 8*log(abs(-cos(b*x + a) +
1)/abs(cos(b*x + a) + 1)) + 16*log(abs(-(cos(b*x + a) - 1)/(cos(b*x + a) +
1) - 1)))/b
```

Mupad [B]

time = 0.38, size = 39, normalized size = 0.91

$$\frac{\tan(a + bx)^2}{2b} - \frac{1}{2b \tan(a + bx)^2} + \frac{2 \ln(\tan(a + bx))}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(a + b*x)^3*sin(a + b*x)^3),x)
```

```
[Out] tan(a + b*x)^2/(2*b) - 1/(2*b*tan(a + b*x)^2) + (2*log(tan(a + b*x)))/b
```

3.155 $\int \csc^3(a + bx) \sec^4(a + bx) dx$

Optimal. Leaf size=66

$$-\frac{5 \tanh^{-1}(\cos(a + bx))}{2b} + \frac{5 \sec(a + bx)}{2b} + \frac{5 \sec^3(a + bx)}{6b} - \frac{\csc^2(a + bx) \sec^3(a + bx)}{2b}$$

[Out] $-5/2*\operatorname{arctanh}(\cos(b*x+a))/b+5/2*\sec(b*x+a)/b+5/6*\sec(b*x+a)^3/b-1/2*\csc(b*x+a)^2*\sec(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2702, 294, 308, 213}

$$\frac{5 \sec^3(a + bx)}{6b} + \frac{5 \sec(a + bx)}{2b} - \frac{5 \tanh^{-1}(\cos(a + bx))}{2b} - \frac{\csc^2(a + bx) \sec^3(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^3*Sec[a + b*x]^4,x]`

[Out] $(-5*\operatorname{ArcTanh}[\operatorname{Cos}[a + b*x]])/(2*b) + (5*\operatorname{Sec}[a + b*x])/(2*b) + (5*\operatorname{Sec}[a + b*x]^3)/(6*b) - (\operatorname{Csc}[a + b*x]^2*\operatorname{Sec}[a + b*x]^3)/(2*b)$

Rule 213

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rule 294

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 308

`Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2`

), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
 \int \csc^3(a + bx) \sec^4(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^6}{(-1+x^2)^2} dx, x, \sec(a + bx)\right)}{b} \\
 &= -\frac{\csc^2(a + bx) \sec^3(a + bx)}{2b} + \frac{5 \text{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, \sec(a + bx)\right)}{2b} \\
 &= -\frac{\csc^2(a + bx) \sec^3(a + bx)}{2b} + \frac{5 \text{Subst}\left(\int \left(1 + x^2 + \frac{1}{-1+x^2}\right) dx, x, \sec(a + bx)\right)}{2b} \\
 &= \frac{5 \sec(a + bx)}{2b} + \frac{5 \sec^3(a + bx)}{6b} - \frac{\csc^2(a + bx) \sec^3(a + bx)}{2b} + \frac{5 \text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \sec(a + bx)\right)}{2b} \\
 &= -\frac{5 \tanh^{-1}(\cos(a + bx))}{2b} + \frac{5 \sec(a + bx)}{2b} + \frac{5 \sec^3(a + bx)}{6b} - \frac{\csc^2(a + bx)}{2b}
 \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 205 vs. 2(66) = 132.

time = 0.30, size = 205, normalized size = 3.11

$\frac{2 \cos^2(a + bx) (22 - 40 \cos(2(a + bx)) + 13 \cos(3(a + bx)) - 30 \cos(4(a + bx)) + 13 \cos(5(a + bx)) + 15 \cos(3(a + bx)) \log(\cos(\frac{1}{2}(a + bx))) + 15 \cos(5(a + bx)) \log(\cos(\frac{1}{2}(a + bx))) - 15 \cos(3(a + bx)) \log(\sin(\frac{1}{2}(a + bx))) - 15 \cos(5(a + bx)) \log(\sin(\frac{1}{2}(a + bx))) + \cos(a + bx) (-26 - 30 \log(\cos(\frac{1}{2}(a + bx))) + 30 \log(\sin(\frac{1}{2}(a + bx))))}{3b (\cos^2(\frac{1}{2}(a + bx)) - \sec^2(\frac{1}{2}(a + bx)))^2}$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^3*Sec[a + b*x]^4,x]

[Out] (2*Csc[a + b*x]^8*(22 - 40*Cos[2*(a + b*x)] + 13*Cos[3*(a + b*x)] - 30*Cos[4*(a + b*x)] + 13*Cos[5*(a + b*x)] + 15*Cos[3*(a + b*x)]*Log[Cos[(a + b*x)/2]] + 15*Cos[5*(a + b*x)]*Log[Cos[(a + b*x)/2]] - 15*Cos[3*(a + b*x)]*Log[Sin[(a + b*x)/2]] - 15*Cos[5*(a + b*x)]*Log[Sin[(a + b*x)/2]] + Cos[a + b*x]*(-26 - 30*Log[Cos[(a + b*x)/2]] + 30*Log[Sin[(a + b*x)/2]]))/ (3*b*(Csc[(a + b*x)/2]^2 - Sec[(a + b*x)/2]^2)^3)

Maple [A]

time = 0.08, size = 70, normalized size = 1.06

method	result	size
derivativedivides	$\frac{\frac{1}{3 \sin(bx+a)^2 \cos(bx+a)^3} - \frac{5}{6 \sin(bx+a)^2 \cos(bx+a)} + \frac{5}{2 \cos(bx+a)} + \frac{5 \ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$	70
default	$\frac{\frac{1}{3 \sin(bx+a)^2 \cos(bx+a)^3} - \frac{5}{6 \sin(bx+a)^2 \cos(bx+a)} + \frac{5}{2 \cos(bx+a)} + \frac{5 \ln(\csc(bx+a) - \cot(bx+a))}{2}}{b}$	70

norman	$\frac{\frac{1}{8b} + \frac{\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} + \frac{75\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} - \frac{65\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{12b} - \frac{55\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^3 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^2} + \frac{5 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b}$	114
risch	$\frac{15 e^{9i(bx+a)} + 20 e^{7i(bx+a)} - 22 e^{5i(bx+a)} + 20 e^{3i(bx+a)} + 15 e^{i(bx+a)}}{3b(e^{2i(bx+a)} + 1)^3 (e^{2i(bx+a)} - 1)^2} + \frac{5 \ln(e^{i(bx+a)} - 1)}{2b} - \frac{5 \ln(e^{i(bx+a)} + 1)}{2b}$	123

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^4/sin(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b} * \left(\frac{1}{3} \sin(b*x+a)^2 / \cos(b*x+a)^3 - \frac{5}{6} \sin(b*x+a)^2 / \cos(b*x+a) + \frac{5}{2} \cos(b*x+a) + \frac{5}{2} \ln(\csc(b*x+a) - \cot(b*x+a)) \right)$

Maxima [A]

time = 0.29, size = 73, normalized size = 1.11

$$\frac{2 \left(15 \cos(bx+a)^4 - 10 \cos(bx+a)^2 - 2 \right)}{\cos(bx+a)^5 - \cos(bx+a)^3} - \frac{15 \log(\cos(bx+a) + 1) + 15 \log(\cos(bx+a) - 1)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4/sin(b*x+a)^3,x, algorithm="maxima")`

[Out] $\frac{1}{12} * \left(2 * (15 * \cos(b*x + a)^4 - 10 * \cos(b*x + a)^2 - 2) / (\cos(b*x + a)^5 - \cos(b*x + a)^3) - 15 * \log(\cos(b*x + a) + 1) + 15 * \log(\cos(b*x + a) - 1) \right) / b$

Fricas [A]

time = 0.35, size = 112, normalized size = 1.70

$$\frac{30 \cos(bx+a)^4 - 20 \cos(bx+a)^2 - 15 (\cos(bx+a)^5 - \cos(bx+a)^3) \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 15 (\cos(bx+a)^5 - \cos(bx+a)^3) \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 4}{12 (b \cos(bx+a)^5 - b \cos(bx+a)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^4/sin(b*x+a)^3,x, algorithm="fricas")`

[Out] $\frac{1}{12} * \left(30 * \cos(b*x + a)^4 - 20 * \cos(b*x + a)^2 - 15 * (\cos(b*x + a)^5 - \cos(b*x + a)^3) * \log\left(\frac{1}{2} * \cos(b*x + a) + \frac{1}{2}\right) + 15 * (\cos(b*x + a)^5 - \cos(b*x + a)^3) * \log\left(-\frac{1}{2} * \cos(b*x + a) + \frac{1}{2}\right) - 4 \right) / (b * \cos(b*x + a)^5 - b * \cos(b*x + a)^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(a + bx)}{\sin^3(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**4/sin(b*x+a)**3,x)`

[Out] Integral(sec(a + b*x)**4/sin(a + b*x)**3, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(58) = 116.

time = 3.54, size = 163, normalized size = 2.47

$$\frac{3 \left(\frac{10(\cos(bx+a)-1)}{\cos(bx+a)+1} - 1 \right) (\cos(bx+a)+1)}{\cos(bx+a)-1} + \frac{3(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{16 \left(\frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{9(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 7 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1 \right)^3} - 30 \log \left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right)$$

$$24b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^3,x, algorithm="giac")

[Out] -1/24*(3*(10*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)*(cos(b*x + a) + 1)/(cos(b*x + a) - 1) + 3*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 16*(12*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 9*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 7)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^3 - 30*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.46, size = 60, normalized size = 0.91

$$\frac{-\frac{5 \cos(a+bx)^4}{2} + \frac{5 \cos(a+bx)^2}{3} + \frac{1}{3}}{b (\cos(a+bx)^3 - \cos(a+bx)^5)} - \frac{5 \operatorname{atanh}(\cos(a+bx))}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^4*sin(a + b*x)^3),x)

[Out] ((5*cos(a + b*x)^2)/3 - (5*cos(a + b*x)^4)/2 + 1/3)/(b*(cos(a + b*x)^3 - cos(a + b*x)^5)) - (5*atanh(cos(a + b*x)))/(2*b)

3.156 $\int \csc^3(a + bx) \sec^5(a + bx) dx$

Optimal. Leaf size=58

$$-\frac{\cot^2(a + bx)}{2b} + \frac{3 \log(\tan(a + bx))}{b} + \frac{3 \tan^2(a + bx)}{2b} + \frac{\tan^4(a + bx)}{4b}$$

[Out] $-1/2*\cot(b*x+a)^2/b+3*\ln(\tan(b*x+a))/b+3/2*\tan(b*x+a)^2/b+1/4*\tan(b*x+a)^4/b$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2700, 272, 45}

$$\frac{\tan^4(a + bx)}{4b} + \frac{3 \tan^2(a + bx)}{2b} - \frac{\cot^2(a + bx)}{2b} + \frac{3 \log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^3*Sec[a + b*x]^5,x]

[Out] $-1/2*\cot[a + b*x]^2/b + (3*\log[\tan[a + b*x]])/b + (3*\tan[a + b*x]^2)/(2*b) + \tan[a + b*x]^4/(4*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned}
\int \csc^3(a+bx) \sec^5(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^3}{x^3} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x)^3}{x^2} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(3 + \frac{1}{x^2} + \frac{3}{x} + x\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= -\frac{\cot^2(a+bx)}{2b} + \frac{3 \log(\tan(a+bx))}{b} + \frac{3 \tan^2(a+bx)}{2b} + \frac{\tan^4(a+bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.16, size = 56, normalized size = 0.97

$$\frac{2 \csc^2(a+bx) + 12 \log(\cos(a+bx)) - 12 \log(\sin(a+bx)) - 4 \sec^2(a+bx) - \sec^4(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^3*Sec[a + b*x]^5,x]``[Out] -1/4*(2*Csc[a + b*x]^2 + 12*Log[Cos[a + b*x]] - 12*Log[Sin[a + b*x]] - 4*Sec[a + b*x]^2 - Sec[a + b*x]^4)/b`**Maple [A]**

time = 0.08, size = 61, normalized size = 1.05

method	result
derivativedivides	$\frac{\frac{1}{4 \sin^2(bx+a) \cos(bx+a)^4} + \frac{3}{4 \sin^2(bx+a)^2 \cos(bx+a)^2} - \frac{3}{2 \sin^2(bx+a)^2} + 3 \ln(\tan(bx+a))}{b}$
default	$\frac{\frac{1}{4 \sin^2(bx+a) \cos(bx+a)^4} + \frac{3}{4 \sin^2(bx+a)^2 \cos(bx+a)^2} - \frac{3}{2 \sin^2(bx+a)^2} + 3 \ln(\tan(bx+a))}{b}$
risch	$\frac{6 e^{10i(bx+a)} + 12 e^{8i(bx+a)} - 4 e^{6i(bx+a)} + 12 e^{4i(bx+a)} + 6 e^{2i(bx+a)}}{b(e^{2i(bx+a)} + 1)^4 (e^{2i(bx+a)} - 1)^2} + \frac{3 \ln(e^{2i(bx+a)} - 1)}{b} - \frac{3 \ln(e^{2i(bx+a)} + 1)}{b}$
norman	$\frac{-\frac{1}{8b} - \frac{\tan^{12}\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{10 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} + \frac{57 \left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} + \frac{57 \left(\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^4 \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^5/sin(b*x+a)^3,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/4/sin(b*x+a)^2/cos(b*x+a)^4+3/4/sin(b*x+a)^2/cos(b*x+a)^2-3/2/sin(b*x+a)^2+3*ln(tan(b*x+a)))`

Maxima [A]

time = 0.31, size = 82, normalized size = 1.41

$$\frac{\frac{6 \sin(bx+a)^4 - 9 \sin(bx+a)^2 + 2}{\sin(bx+a)^6 - 2 \sin(bx+a)^4 + \sin(bx+a)^2} + 6 \log(\sin(bx+a)^2 - 1) - 6 \log(\sin(bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^3,x, algorithm="maxima")

[Out] -1/4*((6*sin(b*x + a)^4 - 9*sin(b*x + a)^2 + 2)/(sin(b*x + a)^6 - 2*sin(b*x + a)^4 + sin(b*x + a)^2) + 6*log(sin(b*x + a)^2 - 1) - 6*log(sin(b*x + a)^2))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 112 vs. 2(52) = 104.

time = 0.36, size = 112, normalized size = 1.93

$$\frac{6 \cos(bx+a)^4 - 3 \cos(bx+a)^2 - 6(\cos(bx+a)^6 - \cos(bx+a)^4) \log(\cos(bx+a)^2) + 6(\cos(bx+a)^6 - \cos(bx+a)^4) \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) - 1}{4(b \cos(bx+a)^6 - b \cos(bx+a)^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^3,x, algorithm="fricas")

[Out] 1/4*(6*cos(b*x + a)^4 - 3*cos(b*x + a)^2 - 6*(cos(b*x + a)^6 - cos(b*x + a)^4)*log(cos(b*x + a)^2) + 6*(cos(b*x + a)^6 - cos(b*x + a)^4)*log(-1/4*cos(b*x + a)^2 + 1/4) - 1)/(b*cos(b*x + a)^6 - b*cos(b*x + a)^4)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(a + bx)}{\sin^3(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**5/sin(b*x+a)**3,x)

[Out] Integral(sec(a + b*x)**5/sin(a + b*x)**3, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 232 vs. 2(52) = 104.

time = 3.90, size = 232, normalized size = 4.00

$$\frac{\frac{12(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}-1)(\cos(bx+a)+1)}{\cos(bx+a)-1} - \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - \frac{2\left(\frac{76(\cos(bx+a)-1)+118(\cos(bx+a)-1)^2}{\cos(bx+a)+1} + \frac{76(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^2} + \frac{25(\cos(bx+a)-1)^4+25}{(\cos(bx+a)+1)^4}\right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^4} - 12 \log\left(\frac{|\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) + 24 \log\left(\frac{|\cos(bx+a)-1|}{|\cos(bx+a)+1|} - 1\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^3,x, algorithm="giac")

[Out]
$$-1/8*((12*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)*(\cos(b*x + a) + 1)/(\cos(b*x + a) - 1) - (\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 2*(76*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 118*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 76*(\cos(b*x + a) - 1)^3/(\cos(b*x + a) + 1)^3 + 25*(\cos(b*x + a) - 1)^4/(\cos(b*x + a) + 1)^4 + 25)/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)^4 - 12*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)) + 24*\log(\text{abs}(-(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)))/b$$

Mupad [B]

time = 0.44, size = 74, normalized size = 1.28

$$\frac{3 \ln(\sin(a + bx)^2)}{2b} - \frac{3 \ln(\cos(a + bx))}{b} + \frac{-\frac{3 \cos(a+bx)^4}{2} + \frac{3 \cos(a+bx)^2}{4} + \frac{1}{4}}{b (\cos(a + bx)^4 - \cos(a + bx)^6)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^5*sin(a + b*x)^3),x)

[Out]
$$(3*\log(\sin(a + b*x)^2))/(2*b) - (3*\log(\cos(a + b*x)))/b + ((3*\cos(a + b*x)^2)/4 - (3*\cos(a + b*x)^4)/2 + 1/4)/(b*(\cos(a + b*x)^4 - \cos(a + b*x)^6))$$

3.157 $\int \cos^5(a + bx) \cot^4(a + bx) dx$

Optimal. Leaf size=68

$$\frac{4 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{6 \sin(a + bx)}{b} - \frac{4 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b}$$

[Out] $4*\csc(b*x+a)/b-1/3*\csc(b*x+a)^3/b+6*\sin(b*x+a)/b-4/3*\sin(b*x+a)^3/b+1/5*\sin(b*x+a)^5/b$

Rubi [A]

time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2670, 276}

$$\frac{\sin^5(a + bx)}{5b} - \frac{4 \sin^3(a + bx)}{3b} + \frac{6 \sin(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{4 \csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^5*Cot[a + b*x]^4,x]`

[Out] $(4*\text{Csc}[a + b*x])/b - \text{Csc}[a + b*x]^3/(3*b) + (6*\text{Sin}[a + b*x])/b - (4*\text{Sin}[a + b*x]^3)/(3*b) + \text{Sin}[a + b*x]^5/(5*b)$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2670

`Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]`

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx) \cot^4(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^4}{x^4} dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(6 + \frac{1}{x^4} - \frac{4}{x^2} - 4x^2 + x^4\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{4 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{6 \sin(a + bx)}{b} - \frac{4 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 68, normalized size = 1.00

$$\frac{4 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{6 \sin(a + bx)}{b} - \frac{4 \sin^3(a + bx)}{3b} + \frac{\sin^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^5*Cot[a + b*x]^4,x]`
`[Out] (4*Csc[a + b*x])/b - Csc[a + b*x]^3/(3*b) + (6*Sin[a + b*x])/b - (4*Sin[a + b*x]^3)/(3*b) + Sin[a + b*x]^5/(5*b)`
Maple [A]

time = 0.11, size = 90, normalized size = 1.32

method	result
derivativedivides	$\frac{-\frac{\cos^{10}(bx+a)}{3 \sin(bx+a)^3} + \frac{7(\cos^{10}(bx+a))}{3 \sin(bx+a)} + \frac{7\left(\frac{128}{35} + \cos^8(bx+a) + \frac{8(\cos^6(bx+a))}{7} + \frac{48(\cos^4(bx+a))}{35} + \frac{64(\cos^2(bx+a))}{35}\right) \sin(bx+a)}{3}}{b}$
default	$\frac{-\frac{\cos^{10}(bx+a)}{3 \sin(bx+a)^3} + \frac{7(\cos^{10}(bx+a))}{3 \sin(bx+a)} + \frac{7\left(\frac{128}{35} + \cos^8(bx+a) + \frac{8(\cos^6(bx+a))}{7} + \frac{48(\cos^4(bx+a))}{35} + \frac{64(\cos^2(bx+a))}{35}\right) \sin(bx+a)}{3}}{b}$
risch	$-\frac{ie^{5i(bx+a)}}{160b} - \frac{13ie^{3i(bx+a)}}{96b} - \frac{41ie^{i(bx+a)}}{16b} + \frac{41ie^{-i(bx+a)}}{16b} + \frac{13ie^{-3i(bx+a)}}{96b} + \frac{ie^{-5i(bx+a)}}{160b} + \frac{8i(3e^{5i(bx+a)} - 3e^{3i(bx+a)} - 3e^{i(bx+a)} + 3e^{-i(bx+a)} - 3e^{-3i(bx+a)} + 3e^{-5i(bx+a)} - 3e^{-7i(bx+a)} + 3e^{-9i(bx+a)} - 3e^{-11i(bx+a)} + 3e^{-13i(bx+a)} - 3e^{-15i(bx+a)} + 3e^{-17i(bx+a)} - 3e^{-19i(bx+a)} + 3e^{-21i(bx+a)} - 3e^{-23i(bx+a)} + 3e^{-25i(bx+a)} - 3e^{-27i(bx+a)} + 3e^{-29i(bx+a)} - 3e^{-31i(bx+a)} + 3e^{-33i(bx+a)} - 3e^{-35i(bx+a)})}{3b}$
norman	$\frac{-\frac{1}{24b} + \frac{5(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{3b} + \frac{137(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{6b} + \frac{65(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{1883(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{20b} + \frac{65(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{b} + \frac{137(\tan^{12}(\frac{bx}{2} + \frac{a}{2}))}{6b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^5 \tan(\frac{bx}{2} + \frac{a}{2})^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^9/sin(b*x+a)^4,x,method=_RETURNVERBOSE)`
`[Out] 1/b*(-1/3/sin(b*x+a)^3*cos(b*x+a)^10+7/3/sin(b*x+a)*cos(b*x+a)^10+7/3*(128/35+cos(b*x+a)^8+8/7*cos(b*x+a)^6+48/35*cos(b*x+a)^4+64/35*cos(b*x+a)^2)*sin(b*x+a)`
Maxima [A]

time = 0.29, size = 56, normalized size = 0.82

$$\frac{3 \sin(bx + a)^5 - 20 \sin(bx + a)^3 + \frac{5(12 \sin(bx + a)^2 - 1)}{\sin(bx + a)^3} + 90 \sin(bx + a)}{15b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^9/sin(b*x+a)^4,x, algorithm="maxima")`
`[Out] 1/15*(3*sin(b*x + a)^5 - 20*sin(b*x + a)^3 + 5*(12*sin(b*x + a)^2 - 1)/sin(b*x + a)^3 + 90*sin(b*x + a))/b`

Fricas [A]

time = 0.36, size = 68, normalized size = 1.00

$$\frac{3 \cos (bx+a)^8 + 8 \cos (bx+a)^6 + 48 \cos (bx+a)^4 - 192 \cos (bx+a)^2 + 128}{15 (b \cos (bx+a))^2 - b} \sin (bx+a)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^9/sin(b*x+a)^4,x, algorithm="fricas")``[Out] -1/15*(3*cos(b*x + a)^8 + 8*cos(b*x + a)^6 + 48*cos(b*x + a)^4 - 192*cos(b*x + a)^2 + 128)/((b*cos(b*x + a)^2 - b)*sin(b*x + a))`**Sympy [A]**

time = 1.86, size = 105, normalized size = 1.54

$$\begin{cases} \frac{128 \sin^5(a+bx)}{15b} + \frac{64 \sin^3(a+bx) \cos^2(a+bx)}{3b} + \frac{16 \sin(a+bx) \cos^4(a+bx)}{b} + \frac{8 \cos^6(a+bx)}{3b \sin(a+bx)} - \frac{\cos^8(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^9(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)**9/sin(b*x+a)**4,x)``[Out] Piecewise(((128*sin(a + b*x)**5/(15*b) + 64*sin(a + b*x)**3*cos(a + b*x)**2/(3*b) + 16*sin(a + b*x)*cos(a + b*x)**4/b + 8*cos(a + b*x)**6/(3*b*sin(a + b*x)) - cos(a + b*x)**8/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**9/sin(a)**4, True))`**Giac [A]**

time = 3.43, size = 56, normalized size = 0.82

$$\frac{3 \sin (bx+a)^5 - 20 \sin (bx+a)^3 + \frac{5(12 \sin (bx+a)^2 - 1)}{\sin (bx+a)^3} + 90 \sin (bx+a)}{15 b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^9/sin(b*x+a)^4,x, algorithm="giac")``[Out] 1/15*(3*sin(b*x + a)^5 - 20*sin(b*x + a)^3 + 5*(12*sin(b*x + a)^2 - 1)/sin(b*x + a)^3 + 90*sin(b*x + a))/b`**Mupad [B]**

time = 0.52, size = 55, normalized size = 0.81

$$\frac{3 \sin (a+bx)^8 - 20 \sin (a+bx)^6 + 90 \sin (a+bx)^4 + 60 \sin (a+bx)^2 - 5}{15 b \sin (a+bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)^9/sin(a + b*x)^4,x)``[Out] (60*sin(a + b*x)^2 + 90*sin(a + b*x)^4 - 20*sin(a + b*x)^6 + 3*sin(a + b*x)^8 - 5)/(15*b*sin(a + b*x)^3)`

3.158 $\int \cos^4(a + bx) \cot^4(a + bx) dx$

Optimal. Leaf size=80

$$\frac{35x}{8} + \frac{35 \cot(a + bx)}{8b} - \frac{35 \cot^3(a + bx)}{24b} + \frac{7 \cos^2(a + bx) \cot^3(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot^3(a + bx)}{4b}$$

[Out] 35/8*x+35/8*cot(b*x+a)/b-35/24*cot(b*x+a)^3/b+7/8*cos(b*x+a)^2*cot(b*x+a)^3/b+1/4*cos(b*x+a)^4*cot(b*x+a)^3/b

Rubi [A]

time = 0.04, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2671, 294, 308, 209}

$$-\frac{35 \cot^3(a + bx)}{24b} + \frac{35 \cot(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot^3(a + bx)}{4b} + \frac{7 \cos^2(a + bx) \cot^3(a + bx)}{8b} + \frac{35x}{8}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Cot[a + b*x]^4,x]

[Out] (35*x)/8 + (35*Cot[a + b*x])/(8*b) - (35*Cot[a + b*x]^3)/(24*b) + (7*Cos[a + b*x]^2*Cot[a + b*x]^3)/(8*b) + (Cos[a + b*x]^4*Cot[a + b*x]^3)/(4*b)

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 308

Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2671

Int[sin[(e_) + (f_)*(x_)]^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[In

`t[(ff*x)^(m + n)/(b^2 + ff^2*x^2)^(m/2 + 1), x], x, b*(Tan[e + f*x]/ff)], x
]] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned}
 \int \cos^4(a + bx) \cot^4(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^8}{(1+x^2)^3} dx, x, \cot(a + bx)\right)}{b} \\
 &= \frac{\cos^4(a + bx) \cot^3(a + bx)}{4b} - \frac{7\text{Subst}\left(\int \frac{x^6}{(1+x^2)^2} dx, x, \cot(a + bx)\right)}{4b} \\
 &= \frac{7 \cos^2(a + bx) \cot^3(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot^3(a + bx)}{4b} - \frac{35\text{Subst}\left(\int \frac{x^4}{1+x^2} dx, x, \cot(a + bx)\right)}{4b} \\
 &= \frac{7 \cos^2(a + bx) \cot^3(a + bx)}{8b} + \frac{\cos^4(a + bx) \cot^3(a + bx)}{4b} - \frac{35\text{Subst}\left(\int (-1 + x^2) dx, x, \cot(a + bx)\right)}{4b} \\
 &= \frac{35 \cot(a + bx)}{8b} - \frac{35 \cot^3(a + bx)}{24b} + \frac{7 \cos^2(a + bx) \cot^3(a + bx)}{8b} + \frac{\cos^4(a + bx)}{8b} \\
 &= \frac{35x}{8} + \frac{35 \cot(a + bx)}{8b} - \frac{35 \cot^3(a + bx)}{24b} + \frac{7 \cos^2(a + bx) \cot^3(a + bx)}{8b} + \frac{\cos^4(a + bx)}{8b}
 \end{aligned}$$

Mathematica [A]

time = 0.21, size = 53, normalized size = 0.66

$$\frac{420(a + bx) - 32 \cot(a + bx) (-10 + \csc^2(a + bx)) + 72 \sin(2(a + bx)) + 3 \sin(4(a + bx))}{96b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4*Cot[a + b*x]^4,x]

[Out] (420*(a + b*x) - 32*Cot[a + b*x]*(-10 + Csc[a + b*x]^2) + 72*Sin[2*(a + b*x)] + 3*Sin[4*(a + b*x)])/(96*b)

Maple [A]

time = 0.11, size = 94, normalized size = 1.18

method	result
derivativedivides	$ \frac{-\frac{\cos^9(bx+a)}{3 \sin(bx+a)^3} + \frac{2(\cos^9(bx+a))}{\sin(bx+a)} + 2\left(\cos^7(bx+a) + \frac{7(\cos^5(bx+a))}{6} + \frac{35(\cos^3(bx+a))}{24} + \frac{35 \cos(bx+a)}{16}\right) \sin(bx+a) + \frac{35bx}{8} + \frac{35a}{8}}{b} $
default	$ \frac{-\frac{\cos^9(bx+a)}{3 \sin(bx+a)^3} + \frac{2(\cos^9(bx+a))}{\sin(bx+a)} + 2\left(\cos^7(bx+a) + \frac{7(\cos^5(bx+a))}{6} + \frac{35(\cos^3(bx+a))}{24} + \frac{35 \cos(bx+a)}{16}\right) \sin(bx+a) + \frac{35bx}{8} + \frac{35a}{8}}{b} $

risch	$\frac{35x}{8} - \frac{ie^{4i(bx+a)}}{64b} - \frac{3ie^{2i(bx+a)}}{8b} + \frac{3ie^{-2i(bx+a)}}{8b} + \frac{ie^{-4i(bx+a)}}{64b} + \frac{4i(6e^{4i(bx+a)} - 9e^{2i(bx+a)} + 5)}{3b(e^{2i(bx+a)} - 1)^3}$
norman	$-\frac{1}{24b} + \frac{35(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{24b} + \frac{63(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{8b} + \frac{35(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{8b} - \frac{35(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{8b} - \frac{63(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{8b} - \frac{35(\tan^{12}(\frac{bx}{2} + \frac{a}{2}))}{24b}$

$$\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^8/sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/3/\sin(b*x+a)^3*\cos(b*x+a)^9+2/\sin(b*x+a)*\cos(b*x+a)^9+2*(\cos(b*x+a)^7+7/6*\cos(b*x+a)^5+35/24*\cos(b*x+a)^3+35/16*\cos(b*x+a))*\sin(b*x+a)+35/8*b*x+35/8*a)$

Maxima [A]

time = 0.51, size = 75, normalized size = 0.94

$$\frac{105bx + 105a + \frac{105 \tan(bx+a)^6 + 175 \tan(bx+a)^4 + 56 \tan(bx+a)^2 - 8}{\tan(bx+a)^7 + 2 \tan(bx+a)^5 + \tan(bx+a)^3}}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8/sin(b*x+a)^4,x, algorithm="maxima")`

[Out] $1/24*(105*b*x + 105*a + (105*\tan(b*x + a)^6 + 175*\tan(b*x + a)^4 + 56*\tan(b*x + a)^2 - 8)/(\tan(b*x + a)^7 + 2*\tan(b*x + a)^5 + \tan(b*x + a)^3)/b$

Fricas [A]

time = 0.36, size = 89, normalized size = 1.11

$$\frac{6 \cos(bx+a)^7 + 21 \cos(bx+a)^5 - 140 \cos(bx+a)^3 - 105 (bx \cos(bx+a)^2 - bx) \sin(bx+a) + 105 \cos(bx+a)}{24 (b \cos(bx+a)^2 - b) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8/sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $-1/24*(6*\cos(b*x + a)^7 + 21*\cos(b*x + a)^5 - 140*\cos(b*x + a)^3 - 105*(b*x*\cos(b*x + a)^2 - b*x)*\sin(b*x + a) + 105*\cos(b*x + a))/((b*\cos(b*x + a)^2 - b)*\sin(b*x + a))$

Sympy [A]

time = 1.38, size = 141, normalized size = 1.76

$$\begin{cases} \frac{35x \sin^4(a+bx)}{8} + \frac{35x \sin^2(a+bx) \cos^2(a+bx)}{4} + \frac{35x \cos^4(a+bx)}{8} + \frac{35 \sin^3(a+bx) \cos(a+bx)}{8b} + \frac{175 \sin(a+bx) \cos^3(a+bx)}{24b} + \frac{7 \cos^5(a+bx)}{3b \sin(a+bx)} - \frac{\cos^7(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^8(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**8/sin(b*x+a)**4,x)

[Out] Piecewise((35*x*sin(a + b*x)**4/8 + 35*x**2*cos(a + b*x)**2/4 + 35*x*cos(a + b*x)**4/8 + 35*sin(a + b*x)**3*cos(a + b*x)/(8*b) + 175*sin(a + b*x)*cos(a + b*x)**3/(24*b) + 7*cos(a + b*x)**5/(3*b*sin(a + b*x)) - cos(a + b*x)**7/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**8/sin(a)**4, True))

Giac [A]

time = 3.72, size = 68, normalized size = 0.85

$$\frac{105bx + 105a + \frac{3(11\tan(bx+a)^3 + 13\tan(bx+a))}{(\tan(bx+a)^2 + 1)^2} + \frac{8(9\tan(bx+a)^2 - 1)}{\tan(bx+a)^3}}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^8/sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/24*(105*b*x + 105*a + 3*(11*tan(b*x + a)^3 + 13*tan(b*x + a))/(tan(b*x + a)^2 + 1)^2 + 8*(9*tan(b*x + a)^2 - 1)/tan(b*x + a)^3)/b

Mupad [B]

time = 1.58, size = 56, normalized size = 0.70

$$\frac{35x}{8} + \frac{\cos(a + bx)^4 \left(\frac{35\tan(a+bx)^6}{8} + \frac{175\tan(a+bx)^4}{24} + \frac{7\tan(a+bx)^2}{3} - \frac{1}{3} \right)}{b \tan(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^8/sin(a + b*x)^4,x)

[Out] (35*x)/8 + (cos(a + b*x)^4*((7*tan(a + b*x)^2)/3 + (175*tan(a + b*x)^4)/24 + (35*tan(a + b*x)^6)/8 - 1/3))/(b*tan(a + b*x)^3)

3.159 $\int \cos^3(a + bx) \cot^4(a + bx) dx$

Optimal. Leaf size=53

$$\frac{3 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{3 \sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

[Out] $3*\csc(b*x+a)/b-1/3*\csc(b*x+a)^3/b+3*\sin(b*x+a)/b-1/3*\sin(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2670, 276}

$$-\frac{\sin^3(a + bx)}{3b} + \frac{3 \sin(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{3 \csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*Cot[a + b*x]^4,x]

[Out] $(3*\text{Csc}[a + b*x])/b - \text{Csc}[a + b*x]^3/(3*b) + (3*\text{Sin}[a + b*x])/b - \text{Sin}[a + b*x]^3/(3*b)$

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx) \cot^4(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^3}{x^4} dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(3 + \frac{1}{x^4} - \frac{3}{x^2} - x^2\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{3 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{3 \sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 53, normalized size = 1.00

$$\frac{3 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{3 \sin(a + bx)}{b} - \frac{\sin^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*Cot[a + b*x]^4,x]``[Out] (3*Csc[a + b*x])/b - Csc[a + b*x]^3/(3*b) + (3*Sin[a + b*x])/b - Sin[a + b*x]^3/(3*b)`**Maple [A]**

time = 0.05, size = 80, normalized size = 1.51

method	result
derivativedivides	$\frac{-\frac{\cos^8(bx+a)}{3 \sin(bx+a)^3} + \frac{5(\cos^8(bx+a))}{3 \sin(bx+a)} + \frac{5 \left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5} \right) \sin(bx+a)}{3b}}$
default	$\frac{-\frac{\cos^8(bx+a)}{3 \sin(bx+a)^3} + \frac{5(\cos^8(bx+a))}{3 \sin(bx+a)} + \frac{5 \left(\frac{16}{5} + \cos^6(bx+a) + \frac{6(\cos^4(bx+a))}{5} + \frac{8(\cos^2(bx+a))}{5} \right) \sin(bx+a)}{3b}}$
risch	$-\frac{ie^{3i(bx+a)}}{24b} - \frac{11ie^{i(bx+a)}}{8b} + \frac{11ie^{-i(bx+a)}}{8b} + \frac{ie^{-3i(bx+a)}}{24b} + \frac{2i(9e^{5i(bx+a)} - 14e^{3i(bx+a)} + 9e^{i(bx+a)})}{3b(e^{2i(bx+a)} - 1)^3}$
norman	$\frac{-\frac{1}{24b} + \frac{5(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{91(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{8b} + \frac{35(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{2b} + \frac{91(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{8b} + \frac{5(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{4b} - \frac{\tan^{12}(\frac{bx}{2} + \frac{a}{2})}{24b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^3 \tan(\frac{bx}{2} + \frac{a}{2})^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^7/sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/3/sin(b*x+a)^3*cos(b*x+a)^8+5/3/sin(b*x+a)*cos(b*x+a)^8+5/3*(16/5+cos(b*x+a)^6+6/5*cos(b*x+a)^4+8/5*cos(b*x+a)^2)*sin(b*x+a)`**Maxima [A]**

time = 0.29, size = 44, normalized size = 0.83

$$\frac{\sin(bx + a)^3 - \frac{9 \sin(bx+a)^2 - 1}{\sin(bx+a)^3} - 9 \sin(bx + a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7/sin(b*x+a)^4,x, algorithm="maxima")``[Out] -1/3*(sin(b*x + a)^3 - (9*sin(b*x + a)^2 - 1)/sin(b*x + a)^3 - 9*sin(b*x + a))/b`

Fricas [A]

time = 0.35, size = 56, normalized size = 1.06

$$\frac{\cos(bx+a)^6 + 6 \cos(bx+a)^4 - 24 \cos(bx+a)^2 + 16}{3(b \cos(bx+a)^2 - b) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7/sin(b*x+a)^4,x, algorithm="fricas")``[Out] -1/3*(cos(b*x + a)^6 + 6*cos(b*x + a)^4 - 24*cos(b*x + a)^2 + 16)/((b*cos(b*x + a)^2 - b)*sin(b*x + a))`**Sympy [A]**

time = 1.03, size = 82, normalized size = 1.55

$$\begin{cases} \frac{16 \sin^3(a+bx)}{3b} + \frac{8 \sin(a+bx) \cos^2(a+bx)}{b} + \frac{2 \cos^4(a+bx)}{b \sin(a+bx)} - \frac{\cos^6(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^7(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)**7/sin(b*x+a)**4,x)``[Out] Piecewise((16*sin(a + b*x)**3/(3*b) + 8*sin(a + b*x)*cos(a + b*x)**2/b + 2*cos(a + b*x)**4/(b*sin(a + b*x)) - cos(a + b*x)**6/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**7/sin(a)**4, True))`**Giac [A]**

time = 4.90, size = 41, normalized size = 0.77

$$\frac{\left(\frac{1}{\sin(bx+a)} + \sin(bx+a)\right)^3 - \frac{12}{\sin(bx+a)} - 12 \sin(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^7/sin(b*x+a)^4,x, algorithm="giac")``[Out] -1/3*((1/sin(b*x + a) + sin(b*x + a))^3 - 12/sin(b*x + a) - 12*sin(b*x + a))/b`**Mupad [B]**

time = 0.48, size = 45, normalized size = 0.85

$$\frac{-\sin(a+bx)^6 + 9 \sin(a+bx)^4 + 9 \sin(a+bx)^2 - 1}{3b \sin(a+bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)^7/sin(a + b*x)^4,x)``[Out] (9*sin(a + b*x)^2 + 9*sin(a + b*x)^4 - sin(a + b*x)^6 - 1)/(3*b*sin(a + b*x)^3)`

3.160 $\int \cos^2(a + bx) \cot^4(a + bx) dx$

Optimal. Leaf size=57

$$\frac{5x}{2} + \frac{5 \cot(a + bx)}{2b} - \frac{5 \cot^3(a + bx)}{6b} + \frac{\cos^2(a + bx) \cot^3(a + bx)}{2b}$$

[Out] $5/2*x+5/2*\cot(b*x+a)/b-5/6*\cot(b*x+a)^3/b+1/2*\cos(b*x+a)^2*\cot(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 57, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2671, 294, 308, 209}

$$-\frac{5 \cot^3(a + bx)}{6b} + \frac{5 \cot(a + bx)}{2b} + \frac{\cos^2(a + bx) \cot^3(a + bx)}{2b} + \frac{5x}{2}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Cot[a + b*x]^4,x]

[Out] $(5*x)/2 + (5*\text{Cot}[a + b*x])/(2*b) - (5*\text{Cot}[a + b*x]^3)/(6*b) + (\text{Cos}[a + b*x]^2*\text{Cot}[a + b*x]^3)/(2*b)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 308

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2671

Int[sin[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := With[{ff = FreeFactors[Tan[e + f*x], x]}, Dist[b*(ff/f), Subst[In

`t[(ff*x)^(m + n)/(b^2 + ff^2*x^2)^(m/2 + 1), x], x, b*(Tan[e + f*x]/ff)], x
]] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2]`

Rubi steps

$$\begin{aligned} \int \cos^2(a + bx) \cot^4(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{(1+x^2)^2} dx, x, \cot(a + bx)\right)}{b} \\ &= \frac{\cos^2(a + bx) \cot^3(a + bx)}{2b} - \frac{5\text{Subst}\left(\int \frac{x^4}{1+x^2} dx, x, \cot(a + bx)\right)}{2b} \\ &= \frac{\cos^2(a + bx) \cot^3(a + bx)}{2b} - \frac{5\text{Subst}\left(\int \left(-1 + x^2 + \frac{1}{1+x^2}\right) dx, x, \cot(a + bx)\right)}{2b} \\ &= \frac{5 \cot(a + bx)}{2b} - \frac{5 \cot^3(a + bx)}{6b} + \frac{\cos^2(a + bx) \cot^3(a + bx)}{2b} - \frac{5\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \cot(a + bx)\right)}{2b} \\ &= \frac{5x}{2} + \frac{5 \cot(a + bx)}{2b} - \frac{5 \cot^3(a + bx)}{6b} + \frac{\cos^2(a + bx) \cot^3(a + bx)}{2b} \end{aligned}$$

Mathematica [A]

time = 0.13, size = 43, normalized size = 0.75

$$\frac{30(a + bx) - 4 \cot(a + bx) (-7 + \csc^2(a + bx)) + 3 \sin(2(a + bx))}{12b}$$

Antiderivative was successfully verified.

[In] `Integrate[Cos[a + b*x]^2*Cot[a + b*x]^4, x]`

[Out] `(30*(a + b*x) - 4*Cot[a + b*x]*(-7 + Csc[a + b*x]^2) + 3*Sin[2*(a + b*x)])/(12*b)`

Maple [A]

time = 0.04, size = 84, normalized size = 1.47

method	result
risch	$\frac{5x}{2} - \frac{ie^{2i(bx+a)}}{8b} + \frac{ie^{-2i(bx+a)}}{8b} + \frac{2i(9e^{4i(bx+a)} - 12e^{2i(bx+a)} + 7)}{3b(e^{2i(bx+a)} - 1)^3}$
derivativedivides	$-\frac{\cos^7(bx+a)}{3 \sin(bx+a)^3} + \frac{4(\cos^7(bx+a))}{3 \sin(bx+a)} + \frac{4\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15 \cos(bx+a)}{8}\right) \sin(bx+a)}{3} + \frac{5bx + 5a}{2}$
default	$-\frac{\cos^7(bx+a)}{3 \sin(bx+a)^3} + \frac{4(\cos^7(bx+a))}{3 \sin(bx+a)} + \frac{4\left(\cos^5(bx+a) + \frac{5(\cos^3(bx+a))}{4} + \frac{15 \cos(bx+a)}{8}\right) \sin(bx+a)}{3} + \frac{5bx + 5a}{2}$

norman	$\frac{-\frac{1}{24b} + \frac{25(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{24b} + \frac{25(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{12b} - \frac{25(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{12b} - \frac{25(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{24b} + \frac{\tan^{10}(\frac{bx}{2} + \frac{a}{2})}{24b} + \frac{5x(\tan^3(\frac{bx}{2} + \frac{a}{2}))}{2}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))^2 \tan(\frac{bx}{2} + \frac{a}{2})^3} + 5$
--------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^6/sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] $1/b * (-1/3 * \cos(b*x+a)^7 / \sin(b*x+a)^3 + 4/3 * \cos(b*x+a)^7 / \sin(b*x+a) + 5/4 * \cos(b*x+a)^3 + 15/8 * \cos(b*x+a)) * \sin(b*x+a) + 5/2 * b*x + 5/2 * a$

Maxima [A]

time = 0.50, size = 55, normalized size = 0.96

$$\frac{15bx + 15a + \frac{15 \tan(bx+a)^4 + 10 \tan(bx+a)^2 - 2}{\tan(bx+a)^5 + \tan(bx+a)^3}}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^4,x, algorithm="maxima")`

[Out] $1/6 * (15 * b * x + 15 * a + (15 * \tan(b * x + a)^4 + 10 * \tan(b * x + a)^2 - 2) / (\tan(b * x + a)^5 + \tan(b * x + a)^3)) / b$

Fricas [A]

time = 0.40, size = 79, normalized size = 1.39

$$\frac{3 \cos(bx+a)^5 - 20 \cos(bx+a)^3 - 15 (bx \cos(bx+a)^2 - bx) \sin(bx+a) + 15 \cos(bx+a)}{6 (b \cos(bx+a)^2 - b) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $-1/6 * (3 * \cos(b * x + a)^5 - 20 * \cos(b * x + a)^3 - 15 * (b * x * \cos(b * x + a)^2 - b * x) * \sin(b * x + a) + 15 * \cos(b * x + a)) / ((b * \cos(b * x + a)^2 - b) * \sin(b * x + a))$

Sympy [A]

time = 0.79, size = 97, normalized size = 1.70

$$\begin{cases} \frac{5x \sin^2(a+bx)}{2} + \frac{5x \cos^2(a+bx)}{2} + \frac{5 \sin(a+bx) \cos(a+bx)}{2b} + \frac{5 \cos^3(a+bx)}{3b \sin(a+bx)} - \frac{\cos^5(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^6(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**6/sin(b*x+a)**4,x)`

[Out] Piecewise((5*x*sin(a + b*x)**2/2 + 5*x*cos(a + b*x)**2/2 + 5*sin(a + b*x)*cos(a + b*x)/(2*b) + 5*cos(a + b*x)**3/(3*b*sin(a + b*x)) - cos(a + b*x)**5/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**6/sin(a)**4, True))

Giac [A]

time = 6.31, size = 55, normalized size = 0.96

$$\frac{15bx + 15a + \frac{3 \tan(bx+a)}{\tan(bx+a)^2+1} + \frac{2(6 \tan(bx+a)^2-1)}{\tan(bx+a)^3}}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/6*(15*b*x + 15*a + 3*tan(b*x + a)/(tan(b*x + a)^2 + 1) + 2*(6*tan(b*x + a)^2 - 1)/tan(b*x + a)^3)/b

Mupad [B]

time = 0.72, size = 46, normalized size = 0.81

$$\frac{5x}{2} + \frac{\cos(a + bx)^2 \left(\frac{5 \tan(a+bx)^4}{2} + \frac{5 \tan(a+bx)^2}{3} - \frac{1}{3} \right)}{b \tan(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6/sin(a + b*x)^4,x)

[Out] (5*x)/2 + (cos(a + b*x)^2*((5*tan(a + b*x)^2)/3 + (5*tan(a + b*x)^4)/2 - 1/3))/(b*tan(a + b*x)^3)

3.161 $\int \cos(a + bx) \cot^4(a + bx) dx$

Optimal. Leaf size=37

$$\frac{2 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b}$$

[Out] 2*csc(b*x+a)/b-1/3*csc(b*x+a)^3/b+sin(b*x+a)/b

Rubi [A]

time = 0.02, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2670, 276}

$$\frac{\sin(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{2 \csc(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*Cot[a + b*x]^4,x]

[Out] (2*Csc[a + b*x])/b - Csc[a + b*x]^3/(3*b) + Sin[a + b*x]/b

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned} \int \cos(a + bx) \cot^4(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{(1-x^2)^2}{x^4} dx, x, -\sin(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^4} - \frac{2}{x^2}\right) dx, x, -\sin(a + bx)\right)}{b} \\ &= \frac{2 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 37, normalized size = 1.00

$$\frac{2 \csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} + \frac{\sin(a + bx)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]*Cot[a + b*x]^4,x]``[Out] (2*Csc[a + b*x])/b - Csc[a + b*x]^3/(3*b) + Sin[a + b*x]/b`**Maple [A]**

time = 0.04, size = 68, normalized size = 1.84

method	result	size
derivativedivides	$\frac{-\frac{\cos^6(bx+a)}{3 \sin(bx+a)^3} + \frac{\cos^6(bx+a)}{\sin(bx+a)} + \left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{b}$	68
default	$\frac{-\frac{\cos^6(bx+a)}{3 \sin(bx+a)^3} + \frac{\cos^6(bx+a)}{\sin(bx+a)} + \left(\frac{8}{3} + \cos^4(bx+a) + \frac{4(\cos^2(bx+a))}{3}\right) \sin(bx+a)}{b}$	68
risch	$-\frac{ie^{i(bx+a)}}{2b} + \frac{ie^{-i(bx+a)}}{2b} + \frac{4i(3e^{5i(bx+a)} - 4e^{3i(bx+a)} + 3e^{i(bx+a)})}{3b(e^{2i(bx+a)} - 1)^3}$	85
norman	$\frac{-\frac{1}{24b} + \frac{5(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{6b} + \frac{15(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{5(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{6b} - \frac{\tan^8(\frac{bx}{2} + \frac{a}{2})}{24b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2})) \tan(\frac{bx}{2} + \frac{a}{2})^3}$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5/sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/3*cos(b*x+a)^6/sin(b*x+a)^3+cos(b*x+a)^6/sin(b*x+a)+(8/3+cos(b*x+a)^4+4/3*cos(b*x+a)^2)*sin(b*x+a)`**Maxima [A]**

time = 0.32, size = 35, normalized size = 0.95

$$\frac{\frac{6 \sin(bx+a)^2 - 1}{\sin(bx+a)^3} + 3 \sin(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5/sin(b*x+a)^4,x, algorithm="maxima")``[Out] 1/3*((6*sin(b*x + a)^2 - 1)/sin(b*x + a)^3 + 3*sin(b*x + a))/b`**Fricas [A]**

time = 0.35, size = 48, normalized size = 1.30

$$\frac{3 \cos(bx+a)^4 - 12 \cos(bx+a)^2 + 8}{3(b \cos(bx+a)^2 - b) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a)^4,x, algorithm="fricas")

[Out] -1/3*(3*cos(b*x + a)^4 - 12*cos(b*x + a)^2 + 8)/((b*cos(b*x + a)^2 - b)*sin(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 63 vs. 2(29) = 58.

time = 0.61, size = 63, normalized size = 1.70

$$\begin{cases} \frac{8 \sin(a+bx)}{3b} + \frac{4 \cos^2(a+bx)}{3b \sin(a+bx)} - \frac{\cos^4(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^5(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**5/sin(b*x+a)**4,x)

[Out] Piecewise((8*sin(a + b*x)/(3*b) + 4*cos(a + b*x)**2/(3*b*sin(a + b*x)) - cos(a + b*x)**4/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**5/sin(a)**4, True))

Giac [A]

time = 5.41, size = 35, normalized size = 0.95

$$\frac{\frac{6 \sin(bx+a)^2-1}{\sin(bx+a)^3} + 3 \sin(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/3*((6*sin(b*x + a)^2 - 1)/sin(b*x + a)^3 + 3*sin(b*x + a))/b

Mupad [B]

time = 0.45, size = 32, normalized size = 0.86

$$\frac{\sin(a + bx)^4 + 2 \sin(a + bx)^2 - \frac{1}{3}}{b \sin(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5/sin(a + b*x)^4,x)

[Out] (2*sin(a + b*x)^2 + sin(a + b*x)^4 - 1/3)/(b*sin(a + b*x)^3)

3.162 $\int \cot^4(a + bx) dx$

Optimal. Leaf size=27

$$x + \frac{\cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b}$$

[Out] $x + \cot(b*x+a)/b - 1/3*\cot(b*x+a)^3/b$

Rubi [A]

time = 0.01, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 8}

$$-\frac{\cot^3(a + bx)}{3b} + \frac{\cot(a + bx)}{b} + x$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^4,x]

[Out] $x + \text{Cot}[a + b*x]/b - \text{Cot}[a + b*x]^3/(3*b)$

Rule 8

Int[a_, x_Symbol] := Simp[a*x, x] /; FreeQ[a, x]

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \cot^4(a + bx) dx &= -\frac{\cot^3(a + bx)}{3b} - \int \cot^2(a + bx) dx \\ &= \frac{\cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} + \int 1 dx \\ &= x + \frac{\cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 33, normalized size = 1.22

$$-\frac{\cot^3(a + bx) {}_2F_1\left(-\frac{3}{2}, 1; -\frac{1}{2}; -\tan^2(a + bx)\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^4,x]

[Out] -1/3*(Cot[a + b*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, -Tan[a + b*x]^2])/b

Maple [A]

time = 0.03, size = 26, normalized size = 0.96

method	result	size
derivativedivides	$-\frac{(\cot^3(bx+a))}{3} + \frac{\cot(bx+a)+bx+a}{b}$	26
default	$-\frac{(\cot^3(bx+a))}{3} + \frac{\cot(bx+a)+bx+a}{b}$	26
risch	$x + \frac{4i(3e^{4i(bx+a)} - 3e^{2i(bx+a)} + 2)}{3b(e^{2i(bx+a)} - 1)^3}$	46
norman	$\frac{x\left(\tan^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \frac{1}{24b} + \frac{5\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} - \frac{5\left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{8b} + \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{24b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^3}$	80

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4/sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/3*cot(b*x+a)^3+cot(b*x+a)+b*x+a)

Maxima [A]

time = 0.52, size = 34, normalized size = 1.26

$$\frac{3bx + 3a + \frac{3 \tan(bx+a)^2 - 1}{\tan(bx+a)^3}}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^4,x, algorithm="maxima")

[Out] 1/3*(3*b*x + 3*a + (3*tan(b*x + a)^2 - 1)/tan(b*x + a)^3)/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 69 vs. 2(25) = 50.

time = 0.38, size = 69, normalized size = 2.56

$$\frac{4 \cos(bx+a)^3 + 3(bx \cos(bx+a)^2 - bx) \sin(bx+a) - 3 \cos(bx+a)}{3(b \cos(bx+a)^2 - b) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^4,x, algorithm="fricas")

[Out] $\frac{1}{3} \cdot (4 \cdot \cos(b \cdot x + a)^3 + 3 \cdot (b \cdot x \cdot \cos(b \cdot x + a)^2 - b \cdot x) \cdot \sin(b \cdot x + a) - 3 \cdot \cos(b \cdot x + a)) / ((b \cdot \cos(b \cdot x + a)^2 - b) \cdot \sin(b \cdot x + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 48 vs. $2(20) = 40$.

time = 0.50, size = 48, normalized size = 1.78

$$\begin{cases} x + \frac{\cos(a+bx)}{b \sin(a+bx)} - \frac{\cos^3(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^4(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**4/sin(b*x+a)**4,x)`

[Out] `Piecewise((x + cos(a + b*x)/(b*sin(a + b*x)) - cos(a + b*x)**3/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**4/sin(a)**4, True))`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 62 vs. $2(25) = 50$.
time = 4.34, size = 62, normalized size = 2.30

$$\frac{\tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3 + 24bx + 24a + \frac{15 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^2 - 1}{\tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)^3} - 15 \tan\left(\frac{1}{2}bx + \frac{1}{2}a\right)}{24b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^4/sin(b*x+a)^4,x, algorithm="giac")`

[Out] $\frac{1}{24} \cdot (\tan(1/2 \cdot b \cdot x + 1/2 \cdot a)^3 + 24 \cdot b \cdot x + 24 \cdot a + (15 \cdot \tan(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 - 1) / \tan(1/2 \cdot b \cdot x + 1/2 \cdot a)^3 - 15 \cdot \tan(1/2 \cdot b \cdot x + 1/2 \cdot a)) / b$

Mupad [B]

time = 0.45, size = 24, normalized size = 0.89

$$x + \frac{\tan(a + bx)^2 - \frac{1}{3}}{b \tan(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^4/sin(a + b*x)^4,x)`

[Out] $x + (\tan(a + b \cdot x)^2 - 1/3) / (b \cdot \tan(a + b \cdot x)^3)$

3.163 $\int \cot^3(a + bx) \csc(a + bx) dx$

Optimal. Leaf size=26

$$\frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b}$$

[Out] $\csc(b*x+a)/b-1/3*\csc(b*x+a)^3/b$

Rubi [A]

time = 0.01, antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 1, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.067$, Rules used = {2686}

$$\frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cot}[a + b*x]^3*\text{Csc}[a + b*x], x]$

[Out] $\text{Csc}[a + b*x]/b - \text{Csc}[a + b*x]^3/(3*b)$

Rule 2686

$\text{Int}[(a_*)*\text{sec}[(e_*) + (f_*)*(x_*)]^{(m_*)}*((b_*)*\text{tan}[(e_*) + (f_*)*(x_*)]^{(n_*)}, x_Symbol] :> \text{Dist}[a/f, \text{Subst}[\text{Int}[(a*x)^{(m-1)}*(-1+x^2)^{((n-1)/2)}, x], x, \text{Sec}[e+f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])

Rubi steps

$$\begin{aligned} \int \cot^3(a + bx) \csc(a + bx) dx &= -\frac{\text{Subst}\left(\int (-1 + x^2) dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 26, normalized size = 1.00

$$\frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[\text{Cot}[a + b*x]^3*\text{Csc}[a + b*x], x]$

[Out] $\text{Csc}[a + b*x]/b - \text{Csc}[a + b*x]^3/(3*b)$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 59 vs. $2(24) = 48$.

time = 0.03, size = 60, normalized size = 2.31

method	result	size
risch	$\frac{2i(3e^{5i(bx+a)} - 2e^{3i(bx+a)} + 3e^{i(bx+a)})}{3b(e^{2i(bx+a)} - 1)^3}$	54
derivativedivides	$-\frac{\frac{\cos^4(bx+a)}{3\sin(bx+a)^3} + \frac{\cos^4(bx+a)}{3\sin(bx+a)} + \frac{(2+\cos^2(bx+a))\sin(bx+a)}{3}}{b}$	60
default	$-\frac{\frac{\cos^4(bx+a)}{3\sin(bx+a)^3} + \frac{\cos^4(bx+a)}{3\sin(bx+a)} + \frac{(2+\cos^2(bx+a))\sin(bx+a)}{3}}{b}$	60
norman	$-\frac{\frac{1}{24b} + \frac{3(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{8b}}{\tan(\frac{bx}{2} + \frac{a}{2})^3} + \frac{3(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{8b} - \frac{\tan^6(\frac{bx}{2} + \frac{a}{2})}{24b}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^3/sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/3*\cos(b*x+a)^4/\sin(b*x+a)^3+1/3*\cos(b*x+a)^4/\sin(b*x+a)+1/3*(2+\cos(b*x+a)^2)*\sin(b*x+a))$

Maxima [A]

time = 0.30, size = 25, normalized size = 0.96

$$\frac{3 \sin (bx + a)^2 - 1}{3 b \sin (bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3/sin(b*x+a)^4,x, algorithm="maxima")`

[Out] $1/3*(3*\sin(b*x + a)^2 - 1)/(b*\sin(b*x + a)^3)$

Fricas [A]

time = 0.35, size = 38, normalized size = 1.46

$$\frac{3 \cos (bx + a)^2 - 2}{3 (b \cos (bx + a)^2 - b) \sin (bx + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^3/sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $1/3*(3*\cos(b*x + a)^2 - 2)/((b*\cos(b*x + a)^2 - b)*\sin(b*x + a))$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 42 vs. $2(19) = 38$.

time = 0.50, size = 42, normalized size = 1.62

$$\begin{cases} \frac{2}{3b \sin(a+bx)} - \frac{\cos^2(a+bx)}{3b \sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^3(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3/sin(b*x+a)**4,x)

[Out] Piecewise((2/(3*b*sin(a + b*x)) - cos(a + b*x)**2/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)**3/sin(a)**4, True))

Giac [A]

time = 4.48, size = 25, normalized size = 0.96

$$\frac{3 \sin(bx + a)^2 - 1}{3b \sin(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^4,x, algorithm="giac")

[Out] 1/3*(3*sin(b*x + a)^2 - 1)/(b*sin(b*x + a)^3)

Mupad [B]

time = 0.43, size = 22, normalized size = 0.85

$$\frac{\sin(a + bx)^2 - \frac{1}{3}}{b \sin(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3/sin(a + b*x)^4,x)

[Out] (sin(a + b*x)^2 - 1/3)/(b*sin(a + b*x)^3)

3.164 $\int \cot^2(a + bx) \csc^2(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\cot^3(a + bx)}{3b}$$

[Out] -1/3*cot(b*x+a)^3/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 30}

$$-\frac{\cot^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^2*Csc[a + b*x]^2,x]

[Out] -1/3*Cot[a + b*x]^3/b

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2687

Int[sec[(e_) + (f_)*(x_)]^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \cot^2(a + bx) \csc^2(a + bx) dx &= \frac{\text{Subst}\left(\int x^2 dx, x, -\cot(a + bx)\right)}{b} \\ &= -\frac{\cot^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$-\frac{\cot^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^2*Csc[a + b*x]^2,x]

[Out] -1/3*Cot[a + b*x]^3/b

Maple [A]

time = 0.03, size = 22, normalized size = 1.47

method	result	size
derivativdivides	$-\frac{\cos^3(bx+a)}{3\sin(bx+a)^3b}$	22
default	$-\frac{\cos^3(bx+a)}{3\sin(bx+a)^3b}$	22
risch	$\frac{2i(3e^{4i(bx+a)}+1)}{3b(e^{2i(bx+a)}-1)^3}$	33
norman	$-\frac{\frac{1}{24b} + \frac{\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} + \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{24b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^3}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2/sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] -1/3*cos(b*x+a)^3/sin(b*x+a)^3/b

Maxima [A]

time = 0.27, size = 13, normalized size = 0.87

$$-\frac{1}{3b \tan(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^4,x, algorithm="maxima")

[Out] -1/3/(b*tan(b*x + a)^3)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(13) = 26.

time = 0.35, size = 34, normalized size = 2.27

$$\frac{\cos(bx+a)^3}{3(b \cos(bx+a)^2 - b) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/3*cos(b*x + a)^3/((b*cos(b*x + a)^2 - b)*sin(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 71 vs. $2(12) = 24$.

time = 0.78, size = 71, normalized size = 4.73

$$\begin{cases} \frac{\tan^3\left(\frac{a}{2} + \frac{bx}{2}\right)}{24b} - \frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b} + \frac{1}{8b \tan\left(\frac{a}{2} + \frac{bx}{2}\right)} - \frac{1}{24b \tan^3\left(\frac{a}{2} + \frac{bx}{2}\right)} & \text{for } b \neq 0 \\ \frac{x \cos^2(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2/sin(b*x+a)**4,x)

[Out] Piecewise((tan(a/2 + b*x/2)**3/(24*b) - tan(a/2 + b*x/2)/(8*b) + 1/(8*b*tan(a/2 + b*x/2)) - 1/(24*b*tan(a/2 + b*x/2)**3), Ne(b, 0)), (x*cos(a)**2/sin(a)**4, True))

Giac [A]

time = 4.40, size = 13, normalized size = 0.87

$$-\frac{1}{3b \tan(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^4,x, algorithm="giac")

[Out] -1/3/(b*tan(b*x + a)^3)

Mupad [B]

time = 0.38, size = 13, normalized size = 0.87

$$-\frac{\cot(a + bx)^3}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2/sin(a + b*x)^4,x)

[Out] -cot(a + b*x)^3/(3*b)

3.165 $\int \cot(a + bx) \csc^3(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\csc^3(a + bx)}{3b}$$

[Out] -1/3*csc(b*x+a)^3/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2686, 30}

$$-\frac{\csc^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]*Csc[a + b*x]^3,x]

[Out] -1/3*Csc[a + b*x]^3/b

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2686

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \cot(a + bx) \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int x^2 dx, x, \csc(a + bx)\right)}{b} \\ &= -\frac{\csc^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$-\frac{\csc^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]*Csc[a + b*x]^3,x]

[Out] -1/3*Csc[a + b*x]^3/b

Maple [A]

time = 0.02, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$-\frac{1}{3 \sin(bx+a)^3 b}$	14
default	$-\frac{1}{3 \sin(bx+a)^3 b}$	14
risch	$\frac{8ie^{3i(bx+a)}}{3b(e^{2i(bx+a)}-1)^3}$	29
norman	$-\frac{\frac{1}{24b} - \frac{\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)}{8b} - \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{24b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^3}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)/sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] -1/3/sin(b*x+a)^3/b

Maxima [A]

time = 0.28, size = 13, normalized size = 0.87

$$-\frac{1}{3 b \sin (b x + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^4,x, algorithm="maxima")

[Out] -1/3/(b*sin(b*x + a)^3)

Fricas [A]

time = 0.33, size = 26, normalized size = 1.73

$$\frac{1}{3 (b \cos (b x + a)^2 - b) \sin (b x + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^4,x, algorithm="fricas")

[Out] 1/3/((b*cos(b*x + a)^2 - b)*sin(b*x + a))

Sympy [A]

time = 0.42, size = 24, normalized size = 1.60

$$\begin{cases} -\frac{1}{3b\sin^3(a+bx)} & \text{for } b \neq 0 \\ \frac{x\cos(a)}{\sin^4(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)/sin(b*x+a)**4,x)
```

```
[Out] Piecewise((-1/(3*b*sin(a + b*x)**3), Ne(b, 0)), (x*cos(a)/sin(a)**4, True))
```

Giac [A]

time = 4.00, size = 13, normalized size = 0.87

$$-\frac{1}{3b\sin(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)/sin(b*x+a)^4,x, algorithm="giac")
```

```
[Out] -1/3/(b*sin(b*x + a)^3)
```

Mupad [B]

time = 0.43, size = 13, normalized size = 0.87

$$-\frac{1}{3b\sin(a+bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)/sin(a + b*x)^4,x)
```

```
[Out] -1/(3*b*sin(a + b*x)^3)
```

3.166 $\int \csc^4(a + bx) \sec(a + bx) dx$

Optimal. Leaf size=38

$$\frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b}$$

[Out] arctanh(sin(b*x+a))/b-csc(b*x+a)/b-1/3*csc(b*x+a)^3/b

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2701, 308, 213}

$$-\frac{\csc^3(a + bx)}{3b} - \frac{\csc(a + bx)}{b} + \frac{\tanh^{-1}(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^4*Sec[a + b*x],x]

[Out] ArcTanh[Sin[a + b*x]]/b - Csc[a + b*x]/b - Csc[a + b*x]^3/(3*b)

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 308

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2701

Int[(csc[(e_.) + (f_.)*(x_)])*(a_.)^(m_)*sec[(e_.) + (f_.)*(x_)^(n_.)], x_Symbol] := Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
\int \csc^4(a + bx) \sec(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, \csc(a + bx)\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \left(1 + x^2 + \frac{1}{-1+x^2}\right) dx, x, \csc(a + bx)\right)}{b} \\
&= -\frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b} - \frac{\text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \csc(a + bx)\right)}{b} \\
&= \frac{\tanh^{-1}(\sin(a + bx))}{b} - \frac{\csc(a + bx)}{b} - \frac{\csc^3(a + bx)}{3b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 31, normalized size = 0.82

$$-\frac{\csc^3(a + bx) {}_2F_1\left(-\frac{3}{2}, 1; -\frac{1}{2}; \sin^2(a + bx)\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^4*Sec[a + b*x],x]

[Out] -1/3*(Csc[a + b*x]^3*Hypergeometric2F1[-3/2, 1, -1/2, Sin[a + b*x]^2])/b

Maple [A]

time = 0.06, size = 40, normalized size = 1.05

method	result	size
derivativedivides	$-\frac{\frac{1}{3 \sin(bx+a)^3} - \frac{1}{\sin(bx+a)} + \ln(\sec(bx+a) + \tan(bx+a))}{b}$	40
default	$-\frac{\frac{1}{3 \sin(bx+a)^3} - \frac{1}{\sin(bx+a)} + \ln(\sec(bx+a) + \tan(bx+a))}{b}$	40
risch	$-\frac{2i(3e^{5i(bx+a)} - 10e^{3i(bx+a)} + 3e^{i(bx+a)})}{3b(e^{2i(bx+a)} - 1)^3} - \frac{\ln(e^{i(bx+a)} - i)}{b} + \frac{\ln(e^{i(bx+a)} + i)}{b}$	90
norman	$-\frac{\frac{1}{24b} - \frac{5(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{8b} - \frac{5(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{8b} - \frac{\tan^6(\frac{bx}{2} + \frac{a}{2})}{24b}}{\tan(\frac{bx}{2} + \frac{a}{2})^3} + \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}) + 1)}{b} - \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}) - 1)}{b}$	101

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)/sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/3/sin(b*x+a)^3-1/sin(b*x+a)+ln(sec(b*x+a)+tan(b*x+a)))

Maxima [A]

time = 0.28, size = 50, normalized size = 1.32

$$-\frac{2\left(3 \sin(bx+a)^2 + 1\right)}{\sin(bx+a)^3} - 3 \log(\sin(bx+a) + 1) + 3 \log(\sin(bx+a) - 1)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^4,x, algorithm="maxima")

[Out] $-1/6*(2*(3*\sin(b*x + a)^2 + 1)/\sin(b*x + a)^3 - 3*\log(\sin(b*x + a) + 1) + 3*\log(\sin(b*x + a) - 1))/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 94 vs. $2(36) = 72$.

time = 0.37, size = 94, normalized size = 2.47

$$\frac{3(\cos(bx+a)^2 - 1)\log(\sin(bx+a) + 1)\sin(bx+a) - 3(\cos(bx+a)^2 - 1)\log(-\sin(bx+a) + 1)\sin(bx+a) - 6\cos(bx+a)^2 + 8}{6(b\cos(bx+a)^2 - b)\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^4,x, algorithm="fricas")

[Out] $1/6*(3*(\cos(b*x + a)^2 - 1)*\log(\sin(b*x + a) + 1)*\sin(b*x + a) - 3*(\cos(b*x + a)^2 - 1)*\log(-\sin(b*x + a) + 1)*\sin(b*x + a) - 6*\cos(b*x + a)^2 + 8)/((b*\cos(b*x + a)^2 - b)*\sin(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(a + bx)}{\sin^4(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)**4,x)

[Out] Integral(sec(a + b*x)/sin(a + b*x)**4, x)

Giac [A]

time = 3.67, size = 52, normalized size = 1.37

$$\frac{2\left(\frac{3\sin(bx+a)^2+1}{\sin(bx+a)^3}\right) - 3\log(|\sin(bx+a) + 1|) + 3\log(|\sin(bx+a) - 1|)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^4,x, algorithm="giac")

[Out] $-1/6*(2*(3*\sin(b*x + a)^2 + 1)/\sin(b*x + a)^3 - 3*\log(\text{abs}(\sin(b*x + a) + 1)) + 3*\log(\text{abs}(\sin(b*x + a) - 1)))/b$

Mupad [B]

time = 0.02, size = 32, normalized size = 0.84

$$\frac{\operatorname{atanh}(\sin(a + bx)) - \frac{\sin(a+bx)^2 + \frac{1}{3}}{\sin(a+bx)^3}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(cos(a + b*x)*sin(a + b*x)^4),x)
```

```
[Out] (atanh(sin(a + b*x)) - (sin(a + b*x)^2 + 1/3)/sin(a + b*x)^3)/b
```

3.167 $\int \csc^4(a + bx) \sec^2(a + bx) dx$

Optimal. Leaf size=37

$$-\frac{2 \cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} + \frac{\tan(a + bx)}{b}$$

[Out] $-2*\cot(b*x+a)/b-1/3*\cot(b*x+a)^3/b+\tan(b*x+a)/b$

Rubi [A]

time = 0.02, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2700, 276}

$$\frac{\tan(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} - \frac{2 \cot(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[a + b*x]^4*\text{Sec}[a + b*x]^2,x]$

[Out] $(-2*\text{Cot}[a + b*x])/b - \text{Cot}[a + b*x]^3/(3*b) + \text{Tan}[a + b*x]/b$

Rule 276

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{Exp andIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \&\& \text{IGtQ}[p, 0]$

Rule 2700

$\text{Int}[\text{csc}[(e_*) + (f_*)(x_)]^{(m_*)}*\text{sec}[(e_*) + (f_*)(x_)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/f, \text{Subst}[\text{Int}[(1 + x^2)^{(m+n)/2 - 1}/x^m, x], x, \text{Tan}[e + f*x]], x] /; \text{FreeQ}\{e, f, x\} \&\& \text{IntegersQ}[m, n, (m+n)/2]$

Rubi steps

$$\begin{aligned} \int \csc^4(a + bx) \sec^2(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{x^4} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^4} + \frac{2}{x^2}\right) dx, x, \tan(a + bx)\right)}{b} \\ &= -\frac{2 \cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} + \frac{\tan(a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 45, normalized size = 1.22

$$-\frac{5 \cot(a + bx)}{3b} - \frac{\cot(a + bx) \csc^2(a + bx)}{3b} + \frac{\tan(a + bx)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^4*Sec[a + b*x]^2,x]``[Out] (-5*Cot[a + b*x])/(3*b) - (Cot[a + b*x]*Csc[a + b*x]^2)/(3*b) + Tan[a + b*x]/b`**Maple [A]**

time = 0.05, size = 50, normalized size = 1.35

method	result	size
risch	$\frac{16i(2e^{2i(bx+a)}-1)}{3b(e^{2i(bx+a)}-1)^3(e^{2i(bx+a)}+1)}$	46
derivativedivides	$\frac{-\frac{1}{3 \sin(bx+a)^3 \cos(bx+a)} + \frac{4}{3 \sin(bx+a) \cos(bx+a)} - \frac{8 \cot(bx+a)}{3}}{b}$	50
default	$-\frac{1}{3 \sin(bx+a)^3 \cos(bx+a)} + \frac{4}{3 \sin(bx+a) \cos(bx+a)} - \frac{8 \cot(bx+a)}{3}$	50
norman	$\frac{\frac{1}{24b} + \frac{5 \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{6b} - \frac{15 \left(\tan^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{4b} + \frac{5 \left(\tan^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{6b} + \frac{\tan^8 \left(\frac{bx}{2} + \frac{a}{2} \right)}{24b}}{\left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) - 1 \right) \tan \left(\frac{bx}{2} + \frac{a}{2} \right)^3}$	98

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^2/sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/3/sin(b*x+a)^3/cos(b*x+a)+4/3/sin(b*x+a)/cos(b*x+a)-8/3*cot(b*x+a))`**Maxima [A]**

time = 0.28, size = 35, normalized size = 0.95

$$-\frac{\frac{6 \tan(bx+a)^2+1}{\tan(bx+a)^3} - 3 \tan(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^2/sin(b*x+a)^4,x, algorithm="maxima")``[Out] -1/3*((6*tan(b*x + a)^2 + 1)/tan(b*x + a)^3 - 3*tan(b*x + a))/b`**Fricas [A]**

time = 0.34, size = 54, normalized size = 1.46

$$-\frac{8 \cos(bx+a)^4 - 12 \cos(bx+a)^2 + 3}{3(b \cos(bx+a)^3 - b \cos(bx+a)) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^4,x, algorithm="fricas")

[Out] $-1/3*(8*\cos(b*x + a)^4 - 12*\cos(b*x + a)^2 + 3)/((b*\cos(b*x + a))^3 - b*\cos(b*x + a))*\sin(b*x + a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(a + bx)}{\sin^4(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2/sin(b*x+a)**4,x)

[Out] Integral(sec(a + b*x)**2/sin(a + b*x)**4, x)

Giac [A]

time = 4.18, size = 35, normalized size = 0.95

$$-\frac{\frac{6 \tan(bx+a)^2+1}{\tan(bx+a)^3} - 3 \tan(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^4,x, algorithm="giac")

[Out] $-1/3*((6*\tan(b*x + a)^2 + 1)/\tan(b*x + a)^3 - 3*\tan(b*x + a))/b$

Mupad [B]

time = 0.42, size = 36, normalized size = 0.97

$$\frac{\tan(a + bx)}{b} - \frac{2 \tan(a + bx)^2 + \frac{1}{3}}{b \tan(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^2*sin(a + b*x)^4),x)

[Out] $\tan(a + b*x)/b - (2*\tan(a + b*x)^2 + 1/3)/(b*\tan(a + b*x)^3)$

3.168 $\int \csc^4(a + bx) \sec^3(a + bx) dx$

Optimal. Leaf size=66

$$\frac{5 \tanh^{-1}(\sin(a + bx))}{2b} - \frac{5 \csc(a + bx)}{2b} - \frac{5 \csc^3(a + bx)}{6b} + \frac{\csc^3(a + bx) \sec^2(a + bx)}{2b}$$

[Out] $5/2*\operatorname{arctanh}(\sin(b*x+a))/b-5/2*\csc(b*x+a)/b-5/6*\csc(b*x+a)^3/b+1/2*\csc(b*x+a)^3*\sec(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2701, 294, 308, 213}

$$-\frac{5 \csc^3(a + bx)}{6b} - \frac{5 \csc(a + bx)}{2b} + \frac{5 \tanh^{-1}(\sin(a + bx))}{2b} + \frac{\csc^3(a + bx) \sec^2(a + bx)}{2b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[a + b*x]^4*\operatorname{Sec}[a + b*x]^3, x]$

[Out] $(5*\operatorname{ArcTanh}[\operatorname{Sin}[a + b*x]])/(2*b) - (5*\operatorname{Csc}[a + b*x])/(2*b) - (5*\operatorname{Csc}[a + b*x]^3)/(6*b) + (\operatorname{Csc}[a + b*x]^3*\operatorname{Sec}[a + b*x]^2)/(2*b)$

Rule 213

$\operatorname{Int}[(a + (b*x)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[b, 2])^{-1})*\operatorname{ArcTanh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /;$ $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c*x)^m*(a + (b*x)^n)^p, x_Symbol] \rightarrow \operatorname{Simp}[c^{n-1}*(c*x)^{m-n+1}*(a + b*x^n)^{p+1}/(b*n*(p+1)), x] - \operatorname{Dist}[c^{n-1}*(m-n+1)/(b*n*(p+1)), \operatorname{Int}[(c*x)^{m-n}*(a + b*x^n)^{p+1}, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, x\} \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{LtQ}[p, -1] \ \&\& \ \operatorname{GtQ}[m+1, n] \ \&\& \ !\operatorname{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \ \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 308

$\operatorname{Int}[(x)^m/((a + (b*x)^n)), x_Symbol] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[x^m, a + b*x^n, x], x] /;$ $\operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{IGtQ}[m, 0] \ \&\& \ \operatorname{IGtQ}[n, 0] \ \&\& \ \operatorname{GtQ}[m, 2*n-1]$

Rule 2701

$\operatorname{Int}[(\csc[(e*x) + (f*x)]*(a))^m*\sec[(e*x) + (f*x)]^n, x_Symbol] \rightarrow \operatorname{Dist}[-(f*a^n)^{-1}, \operatorname{Subst}[\operatorname{Int}[x^{m+n-1}/(-1 + x^2/a^2)^{(n+1)}, x], x]]$

1)/2), x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \csc^4(a + bx) \sec^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{(-1+x^2)^2} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\csc^3(a + bx) \sec^2(a + bx)}{2b} - \frac{5 \text{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, \csc(a + bx)\right)}{2b} \\ &= \frac{\csc^3(a + bx) \sec^2(a + bx)}{2b} - \frac{5 \text{Subst}\left(\int \left(1 + x^2 + \frac{1}{-1+x^2}\right) dx, x, \csc(a + bx)\right)}{2b} \\ &= -\frac{5 \csc(a + bx)}{2b} - \frac{5 \csc^3(a + bx)}{6b} + \frac{\csc^3(a + bx) \sec^2(a + bx)}{2b} - \frac{5 \text{Subst}\left(\int \frac{1}{-1+x^2} dx, x, \csc(a + bx)\right)}{2b} \\ &= \frac{5 \tanh^{-1}(\sin(a + bx))}{2b} - \frac{5 \csc(a + bx)}{2b} - \frac{5 \csc^3(a + bx)}{6b} + \frac{\csc^3(a + bx) \sec^2(a + bx)}{2b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 31, normalized size = 0.47

$$-\frac{\csc^3(a + bx) {}_2F_1\left(-\frac{3}{2}, 2; -\frac{1}{2}; \sin^2(a + bx)\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^4*Sec[a + b*x]^3,x]

[Out] -1/3*(Csc[a + b*x]^3*Hypergeometric2F1[-3/2, 2, -1/2, Sin[a + b*x]^2])/b

Maple [A]

time = 0.06, size = 68, normalized size = 1.03

method	result
derivativedivides	$-\frac{\frac{1}{3 \sin^3(bx+a) \cos^2(bx+a)} + \frac{5}{6 \sin(bx+a) \cos^2(bx+a)} - \frac{5}{2 \sin(bx+a)} + \frac{5 \ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$
default	$-\frac{\frac{1}{3 \sin^3(bx+a) \cos^2(bx+a)} + \frac{5}{6 \sin(bx+a) \cos^2(bx+a)} - \frac{5}{2 \sin(bx+a)} + \frac{5 \ln(\sec(bx+a) + \tan(bx+a))}{2}}{b}$
risch	$-\frac{i(15 e^{9i(bx+a)} - 20 e^{7i(bx+a)} - 22 e^{5i(bx+a)} - 20 e^{3i(bx+a)} + 15 e^{i(bx+a)})}{3b(e^{2i(bx+a)} - 1)^3(e^{2i(bx+a)} + 1)^2} - \frac{5 \ln(e^{i(bx+a)} - i)}{2b} + \frac{5 \ln(e^{i(bx+a)} + i)}{2b}$
norman	$-\frac{\frac{1}{24b} - \frac{25 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} + \frac{25 \left(\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{12b} + \frac{25 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{12b} - \frac{25 \left(\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24b} - \frac{\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)}{24b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^3} - \frac{5 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^3/sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/3/\sin(b*x+a)^3/\cos(b*x+a)^2+5/6/\sin(b*x+a)/\cos(b*x+a)^2-5/2/\sin(b*x+a)+5/2*\ln(\sec(b*x+a)+\tan(b*x+a)))$

Maxima [A]

time = 0.27, size = 73, normalized size = 1.11

$$\frac{2 \left(15 \sin(bx+a)^4 - 10 \sin(bx+a)^2 - 2 \right)}{\sin(bx+a)^5 - \sin(bx+a)^3} - 15 \log(\sin(bx+a) + 1) + 15 \log(\sin(bx+a) - 1)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3/sin(b*x+a)^4,x, algorithm="maxima")`

[Out] $-1/12*(2*(15*\sin(b*x + a)^4 - 10*\sin(b*x + a)^2 - 2)/(\sin(b*x + a)^5 - \sin(b*x + a)^3) - 15*\log(\sin(b*x + a) + 1) + 15*\log(\sin(b*x + a) - 1))/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 130 vs. 2(58) = 116.

time = 0.35, size = 130, normalized size = 1.97

$$\frac{30 \cos(bx+a)^4 - 15(\cos(bx+a)^4 - \cos(bx+a)^2) \log(\sin(bx+a) + 1) \sin(bx+a) + 15(\cos(bx+a)^4 - \cos(bx+a)^2) \log(-\sin(bx+a) + 1) \sin(bx+a) - 40 \cos(bx+a)^2 + 6}{12(b \cos(bx+a)^4 - b \cos(bx+a)^2) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^3/sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $-1/12*(30*\cos(b*x + a)^4 - 15*(\cos(b*x + a)^4 - \cos(b*x + a)^2)*\log(\sin(b*x + a) + 1)*\sin(b*x + a) + 15*(\cos(b*x + a)^4 - \cos(b*x + a)^2)*\log(-\sin(b*x + a) + 1)*\sin(b*x + a) - 40*\cos(b*x + a)^2 + 6)/((b*\cos(b*x + a)^4 - b*\cos(b*x + a)^2)*\sin(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(a + bx)}{\sin^4(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)**3/sin(b*x+a)**4,x)`

[Out] `Integral(sec(a + b*x)**3/sin(a + b*x)**4, x)`

Giac [A]

time = 3.64, size = 72, normalized size = 1.09

$$\frac{\frac{6 \sin(bx+a)}{\sin(bx+a)^2-1} + \frac{4(6 \sin(bx+a)^2+1)}{\sin(bx+a)^3} - 15 \log(|\sin(bx+a)+1|) + 15 \log(|\sin(bx+a)-1|)}{12b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3/sin(b*x+a)^4,x, algorithm="giac")`

```
[Out] -1/12*(6*sin(b*x + a)/(sin(b*x + a)^2 - 1) + 4*(6*sin(b*x + a)^2 + 1)/sin(b
*x + a)^3 - 15*log(abs(sin(b*x + a) + 1)) + 15*log(abs(sin(b*x + a) - 1)))/
b
```

Mupad [B]

time = 0.38, size = 61, normalized size = 0.92

$$\frac{5 \operatorname{atanh}(\sin(a + bx))}{2b} - \frac{-\frac{5 \sin(a+bx)^4}{2} + \frac{5 \sin(a+bx)^2}{3} + \frac{1}{3}}{b (\sin(a + bx)^3 - \sin(a + bx)^5)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(cos(a + b*x)^3*sin(a + b*x)^4),x)`

```
[Out] (5*atanh(sin(a + b*x)))/(2*b) - ((5*sin(a + b*x)^2)/3 - (5*sin(a + b*x)^4)/
2 + 1/3)/(b*(sin(a + b*x)^3 - sin(a + b*x)^5))
```

3.169 $\int \csc^4(a + bx) \sec^4(a + bx) dx$

Optimal. Leaf size=53

$$-\frac{3 \cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} + \frac{3 \tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b}$$

[Out] $-3*\cot(b*x+a)/b-1/3*\cot(b*x+a)^3/b+3*\tan(b*x+a)/b+1/3*\tan(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2700, 276}

$$\frac{\tan^3(a + bx)}{3b} + \frac{3 \tan(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} - \frac{3 \cot(a + bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^4*Sec[a + b*x]^4,x]

[Out] $(-3*\cot[a + b*x])/b - \cot[a + b*x]^3/(3*b) + (3*\tan[a + b*x])/b + \tan[a + b*x]^3/(3*b)$

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned} \int \csc^4(a + bx) \sec^4(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^3}{x^4} dx, x, \tan(a + bx)\right)}{b} \\ &= \frac{\text{Subst}\left(\int \left(3 + \frac{1}{x^4} + \frac{3}{x^2} + x^2\right) dx, x, \tan(a + bx)\right)}{b} \\ &= -\frac{3 \cot(a + bx)}{b} - \frac{\cot^3(a + bx)}{3b} + \frac{3 \tan(a + bx)}{b} + \frac{\tan^3(a + bx)}{3b} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 43, normalized size = 0.81

$$16 \left(-\frac{\cot(2(a+bx))}{3b} - \frac{\cot(2(a+bx)) \csc^2(2(a+bx))}{6b} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^4*Sec[a + b*x]^4,x]``[Out] 16*(-1/3*Cot[2*(a + b*x)]/b - (Cot[2*(a + b*x)]*Csc[2*(a + b*x)]^2)/(6*b))`**Maple [A]**

time = 0.06, size = 68, normalized size = 1.28

method	result
risch	$\frac{32i(3e^{4i(bx+a)}-1)}{3b(e^{2i(bx+a)}-1)^3(e^{2i(bx+a)}+1)^3}$
derivativedivides	$\frac{\frac{1}{3 \sin(bx+a)^3 \cos(bx+a)^3} - \frac{2}{3 \sin(bx+a)^3 \cos(bx+a)} + \frac{8}{3 \sin(bx+a) \cos(bx+a)} - \frac{16 \cot(bx+a)}{3}}{b}$
default	$\frac{\frac{1}{3 \sin(bx+a)^3 \cos(bx+a)^3} - \frac{2}{3 \sin(bx+a)^3 \cos(bx+a)} + \frac{8}{3 \sin(bx+a) \cos(bx+a)} - \frac{16 \cot(bx+a)}{3}}{b}$
norman	$\frac{\frac{1}{24b} + \frac{5(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{4b} - \frac{91(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{8b} + \frac{35(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{2b} - \frac{91(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{8b} + \frac{5(\tan^{10}(\frac{bx}{2} + \frac{a}{2}))}{4b} + \frac{\tan^{12}(\frac{bx}{2} + \frac{a}{2})}{24b}}{(\tan^2(\frac{bx}{2} + \frac{a}{2}) - 1)^3 \tan(\frac{bx}{2} + \frac{a}{2})^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^4/sin(b*x+a)^4,x,method=_RETURNVERBOSE)``[Out] 1/b*(1/3/sin(b*x+a)^3/cos(b*x+a)^3-2/3/sin(b*x+a)^3/cos(b*x+a)+8/3/sin(b*x+a)/cos(b*x+a)-16/3*cot(b*x+a))`**Maxima [A]**

time = 0.30, size = 44, normalized size = 0.83

$$\frac{\tan(bx+a)^3 - \frac{9 \tan(bx+a)^2 + 1}{\tan(bx+a)^3} + 9 \tan(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^4/sin(b*x+a)^4,x, algorithm="maxima")``[Out] 1/3*(tan(b*x + a)^3 - (9*tan(b*x + a)^2 + 1)/tan(b*x + a)^3 + 9*tan(b*x + a))/b`**Fricas [A]**

time = 0.36, size = 66, normalized size = 1.25

$$\frac{16 \cos(bx+a)^6 - 24 \cos(bx+a)^4 + 6 \cos(bx+a)^2 + 1}{3(b \cos(bx+a)^5 - b \cos(bx+a)^3) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^4,x, algorithm="fricas")

[Out] $-1/3*(16*\cos(b*x + a)^6 - 24*\cos(b*x + a)^4 + 6*\cos(b*x + a)^2 + 1)/((b*\cos(b*x + a)^5 - b*\cos(b*x + a)^3)*\sin(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(a + bx)}{\sin^4(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4/sin(b*x+a)**4,x)

[Out] Integral(sec(a + b*x)**4/sin(a + b*x)**4, x)

Giac [A]

time = 3.29, size = 31, normalized size = 0.58

$$-\frac{8(3 \tan(2bx + 2a)^2 + 1)}{3b \tan(2bx + 2a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^4,x, algorithm="giac")

[Out] $-8/3*(3*\tan(2*b*x + 2*a)^2 + 1)/(b*\tan(2*b*x + 2*a)^3)$

Mupad [B]

time = 0.46, size = 45, normalized size = 0.85

$$-\frac{-\tan(a + bx)^6 - 9 \tan(a + bx)^4 + 9 \tan(a + bx)^2 + 1}{3b \tan(a + bx)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^4*sin(a + b*x)^4),x)

[Out] $-(9*\tan(a + b*x)^2 - 9*\tan(a + b*x)^4 - \tan(a + b*x)^6 + 1)/(3*b*\tan(a + b*x)^3)$

3.170 $\int \csc^4(a + bx) \sec^5(a + bx) dx$

Optimal. Leaf size=89

$$\frac{35 \tanh^{-1}(\sin(a + bx))}{8b} - \frac{35 \csc(a + bx)}{8b} - \frac{35 \csc^3(a + bx)}{24b} + \frac{7 \csc^3(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b}$$

[Out] 35/8*arctanh(sin(b*x+a))/b-35/8*csc(b*x+a)/b-35/24*csc(b*x+a)^3/b+7/8*csc(b*x+a)^3*sec(b*x+a)^2/b+1/4*csc(b*x+a)^3*sec(b*x+a)^4/b

Rubi [A]

time = 0.03, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2701, 294, 308, 213}

$$-\frac{35 \csc^3(a + bx)}{24b} - \frac{35 \csc(a + bx)}{8b} + \frac{35 \tanh^{-1}(\sin(a + bx))}{8b} + \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b} + \frac{7 \csc^3(a + bx) \sec^2(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^4*Sec[a + b*x]^5,x]

[Out] (35*ArcTanh[Sin[a + b*x]])/(8*b) - (35*Csc[a + b*x])/(8*b) - (35*Csc[a + b*x]^3)/(24*b) + (7*Csc[a + b*x]^3*Sec[a + b*x]^2)/(8*b) + (Csc[a + b*x]^3*Sec[a + b*x]^4)/(4*b)

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 308

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2701

```
Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*sec[(e_.) + (f_.)*(x_.)]^(n_.), x_Symbol]
:> Dist[-(f*a^n)^(-1), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Csc[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^4(a + bx) \sec^5(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^8}{(-1+x^2)^3} dx, x, \csc(a + bx)\right)}{b} \\ &= \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b} - \frac{7\text{Subst}\left(\int \frac{x^6}{(-1+x^2)^2} dx, x, \csc(a + bx)\right)}{4b} \\ &= \frac{7 \csc^3(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b} - \frac{35\text{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, \csc(a + bx)\right)}{4b} \\ &= \frac{7 \csc^3(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b} - \frac{35\text{Subst}\left(\int (1 + \frac{x^2}{-1+x^2}) dx, x, \csc(a + bx)\right)}{4b} \\ &= -\frac{35 \csc(a + bx)}{8b} - \frac{35 \csc^3(a + bx)}{24b} + \frac{7 \csc^3(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b} \\ &= \frac{35 \tanh^{-1}(\sin(a + bx))}{8b} - \frac{35 \csc(a + bx)}{8b} - \frac{35 \csc^3(a + bx)}{24b} + \frac{7 \csc^3(a + bx) \sec^2(a + bx)}{8b} + \frac{\csc^3(a + bx) \sec^4(a + bx)}{4b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 31, normalized size = 0.35

$$-\frac{\csc^3(a + bx) {}_2F_1\left(-\frac{3}{2}, 3; -\frac{1}{2}; \sin^2(a + bx)\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^4*Sec[a + b*x]^5,x]

[Out] -1/3*(Csc[a + b*x]^3*Hypergeometric2F1[-3/2, 3, -1/2, Sin[a + b*x]^2])/b

Maple [A]

time = 0.09, size = 86, normalized size = 0.97

method	result
derivativedivides	$\frac{\frac{1}{4 \sin^3(bx+a) \cos^4(bx+a)} - \frac{7}{12 \sin^3(bx+a) \cos^2(bx+a)} + \frac{35}{24 \sin(bx+a) \cos^2(bx+a)} - \frac{35}{8 \sin(bx+a)} + \frac{35 \ln(\sec(bx+a) + \tan(bx+a))}{8}}{b}$
default	$\frac{\frac{1}{4 \sin^3(bx+a) \cos^4(bx+a)} - \frac{7}{12 \sin^3(bx+a) \cos^2(bx+a)} + \frac{35}{24 \sin(bx+a) \cos^2(bx+a)} - \frac{35}{8 \sin(bx+a)} + \frac{35 \ln(\sec(bx+a) + \tan(bx+a))}{8}}{b}$

risch	$-\frac{i(105 e^{13i(bx+a)} + 70 e^{11i(bx+a)} - 329 e^{9i(bx+a)} - 204 e^{7i(bx+a)} - 329 e^{5i(bx+a)} + 70 e^{3i(bx+a)} + 105 e^{i(bx+a)})}{12b(e^{2i(bx+a)} + 1)^4 (e^{2i(bx+a)} - 1)^3} - \frac{35 \ln(e^{i(bx+a)} + 1)}{24b}$
norman	$-\frac{1}{24b} - \frac{35 \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{24b} + \frac{63 \left(\tan^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{8b} - \frac{35 \left(\tan^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{8b} - \frac{35 \left(\tan^8 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{8b} + \frac{63 \left(\tan^{10} \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{8b} - \frac{35 \left(\tan^{12} \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{24b} - \frac{35 \ln \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) - 1 \right)^4 \tan \left(\frac{bx}{2} + \frac{a}{2} \right)^3}{\left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) - 1 \right)^4 \tan \left(\frac{bx}{2} + \frac{a}{2} \right)^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^5/sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] $1/b*(1/4/\sin(b*x+a)^3/\cos(b*x+a)^4-7/12/\sin(b*x+a)^3/\cos(b*x+a)^2+35/24/\sin(b*x+a)/\cos(b*x+a)^2-35/8/\sin(b*x+a)+35/8*\ln(\sec(b*x+a)+\tan(b*x+a)))$

Maxima [A]

time = 0.28, size = 91, normalized size = 1.02

$$-\frac{2 \left(105 \sin(bx+a)^6 - 175 \sin(bx+a)^4 + 56 \sin(bx+a)^2 + 8 \right)}{\sin(bx+a)^7 - 2 \sin(bx+a)^5 + \sin(bx+a)^3} - 105 \log(\sin(bx+a) + 1) + 105 \log(\sin(bx+a) - 1)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^5/sin(b*x+a)^4,x, algorithm="maxima")`

[Out] $-1/48*(2*(105*\sin(b*x + a)^6 - 175*\sin(b*x + a)^4 + 56*\sin(b*x + a)^2 + 8)/(\sin(b*x + a)^7 - 2*\sin(b*x + a)^5 + \sin(b*x + a)^3) - 105*\log(\sin(b*x + a) + 1) + 105*\log(\sin(b*x + a) - 1))/b$

Fricas [A]

time = 0.38, size = 140, normalized size = 1.57

$$-\frac{210 \cos(bx+a)^6 - 280 \cos(bx+a)^4 - 105 (\cos(bx+a)^6 - \cos(bx+a)^4) \log(\sin(bx+a) + 1) \sin(bx+a) + 105 (\cos(bx+a)^6 - \cos(bx+a)^4) \log(-\sin(bx+a) + 1) \sin(bx+a) + 42 \cos(bx+a)^2 + 12}{48 (b \cos(bx+a)^6 - b \cos(bx+a)^4) \sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^5/sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $-1/48*(210*\cos(b*x + a)^6 - 280*\cos(b*x + a)^4 - 105*(\cos(b*x + a)^6 - \cos(b*x + a)^4)*\log(\sin(b*x + a) + 1)*\sin(b*x + a) + 105*(\cos(b*x + a)^6 - \cos(b*x + a)^4)*\log(-\sin(b*x + a) + 1)*\sin(b*x + a) + 42*\cos(b*x + a)^2 + 12)/((b*\cos(b*x + a)^6 - b*\cos(b*x + a)^4)*\sin(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(a + bx)}{\sin^4(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**5/sin(b*x+a)**4,x)

[Out] Integral(sec(a + b*x)**5/sin(a + b*x)**4, x)

Giac [A]

time = 4.78, size = 85, normalized size = 0.96

$$\frac{6 \left(\frac{11 \sin(bx+a)^3 - 13 \sin(bx+a)}{(\sin(bx+a)^2 - 1)^2} + \frac{16 (9 \sin(bx+a)^2 + 1)}{\sin(bx+a)^3} - 105 \log(|\sin(bx+a) + 1|) + 105 \log(|\sin(bx+a) - 1|) \right)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^4,x, algorithm="giac")

[Out] -1/48*(6*(11*sin(b*x + a)^3 - 13*sin(b*x + a))/(sin(b*x + a)^2 - 1)^2 + 16*(9*sin(b*x + a)^2 + 1)/sin(b*x + a)^3 - 105*log(abs(sin(b*x + a) + 1)) + 105*log(abs(sin(b*x + a) - 1)))/b

Mupad [B]

time = 0.45, size = 79, normalized size = 0.89

$$\frac{35 \operatorname{atanh}(\sin(a + bx))}{8b} - \frac{\frac{35 \sin(a+bx)^6}{8} - \frac{175 \sin(a+bx)^4}{24} + \frac{7 \sin(a+bx)^2}{3} + \frac{1}{3}}{b (\sin(a + bx)^7 - 2 \sin(a + bx)^5 + \sin(a + bx)^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^5*sin(a + b*x)^4),x)

[Out] (35*atanh(sin(a + b*x)))/(8*b) - ((7*sin(a + b*x)^2)/3 - (175*sin(a + b*x)^4)/24 + (35*sin(a + b*x)^6)/8 + 1/3)/(b*(sin(a + b*x)^3 - 2*sin(a + b*x)^5 + sin(a + b*x)^7))

3.171 $\int \cos^4(a + bx) \cot^5(a + bx) dx$

Optimal. Leaf size=69

$$\frac{2 \csc^2(a + bx)}{b} - \frac{\csc^4(a + bx)}{4b} + \frac{6 \log(\sin(a + bx))}{b} - \frac{2 \sin^2(a + bx)}{b} + \frac{\sin^4(a + bx)}{4b}$$

[Out] 2*csc(b*x+a)^2/b-1/4*csc(b*x+a)^4/b+6*ln(sin(b*x+a))/b-2*sin(b*x+a)^2/b+1/4*sin(b*x+a)^4/b

Rubi [A]

time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$,

Rules used = {2670, 272, 45}

$$\frac{\sin^4(a + bx)}{4b} - \frac{2 \sin^2(a + bx)}{b} - \frac{\csc^4(a + bx)}{4b} + \frac{2 \csc^2(a + bx)}{b} + \frac{6 \log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^4*Cot[a + b*x]^5,x]

[Out] (2*Csc[a + b*x]^2)/b - Csc[a + b*x]^4/(4*b) + (6*Log[Sin[a + b*x]])/b - (2*Sin[a + b*x]^2)/b + Sin[a + b*x]^4/(4*b)

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \cos^4(a+bx) \cot^5(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^4}{x^5} dx, x, -\sin(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1-x^2)^4}{x^3} dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(-4 + \frac{1}{x^3} - \frac{4}{x^2} + \frac{6}{x} + x\right) dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{2 \csc^2(a+bx)}{b} - \frac{\csc^4(a+bx)}{4b} + \frac{6 \log(\sin(a+bx))}{b} - \frac{2 \sin^2(a+bx)}{b} + \frac{\sin^4(a+bx)}{b}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 55, normalized size = 0.80

$$\frac{8 \csc^2(a+bx) - \csc^4(a+bx) + 24 \log(\sin(a+bx)) - 8 \sin^2(a+bx) + \sin^4(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^4*Cot[a + b*x]^5,x]`

```
[Out] (8*Csc[a + b*x]^2 - Csc[a + b*x]^4 + 24*Log[Sin[a + b*x]] - 8*Sin[a + b*x]^2 + Sin[a + b*x]^4)/(4*b)
```

Maple [A]

time = 0.06, size = 89, normalized size = 1.29

method	result
derivativedivides	$\frac{-\frac{\cos^{10}(bx+a)}{4 \sin(bx+a)^4} + \frac{3(\cos^{10}(bx+a))}{4 \sin(bx+a)^2} + \frac{3(\cos^8(bx+a))}{4} + \cos^6(bx+a) + \frac{3(\cos^4(bx+a))}{2} + 3(\cos^2(bx+a)) + 6 \ln(\sin(bx+a))}{b}$
default	$\frac{-\frac{\cos^{10}(bx+a)}{4 \sin(bx+a)^4} + \frac{3(\cos^{10}(bx+a))}{4 \sin(bx+a)^2} + \frac{3(\cos^8(bx+a))}{4} + \cos^6(bx+a) + \frac{3(\cos^4(bx+a))}{2} + 3(\cos^2(bx+a)) + 6 \ln(\sin(bx+a))}{b}$
risch	$-6ix + \frac{e^{4i(bx+a)}}{64b} + \frac{7e^{2i(bx+a)}}{16b} + \frac{7e^{-2i(bx+a)}}{16b} + \frac{e^{-4i(bx+a)}}{64b} - \frac{12ia}{b} - \frac{4(2e^{6i(bx+a)} - 3e^{4i(bx+a)} + 2e^{2i(bx+a)} - 1)}{b(e^{2i(bx+a)} - 1)^4}$
norman	$\frac{-\frac{1}{64b} + \frac{3(\tan^2(\frac{bx+a}{2}))}{8b} + \frac{3(\tan^{14}(\frac{bx+a}{2}))}{8b} - \frac{\tan^{16}(\frac{bx+a}{2})}{64b} - \frac{93(\tan^6(\frac{bx+a}{2}))}{8b} - \frac{93(\tan^{10}(\frac{bx+a}{2}))}{8b} - \frac{591(\tan^8(\frac{bx+a}{2}))}{32b}}{(1 + \tan^2(\frac{bx+a}{2}))^4 \tan(\frac{bx+a}{2})^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^9/sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/4/sin(b*x+a)^4*cos(b*x+a)^10+3/4/sin(b*x+a)^2*cos(b*x+a)^10+3/4*cos(b*x+a)^8+cos(b*x+a)^6+3/2*cos(b*x+a)^4+3*cos(b*x+a)^2+6*ln(sin(b*x+a)))
```

Maxima [A]

time = 0.29, size = 56, normalized size = 0.81

$$\frac{\sin (bx+a)^4 - 8 \sin (bx+a)^2 + \frac{8 \sin (bx+a)^2 - 1}{\sin (bx+a)^4} + 12 \log (\sin (bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^9/sin(b*x+a)^5,x, algorithm="maxima")

[Out] 1/4*(sin(b*x + a)^4 - 8*sin(b*x + a)^2 + (8*sin(b*x + a)^2 - 1)/sin(b*x + a)^4 + 12*log(sin(b*x + a)^2))/b

Fricas [A]

time = 0.39, size = 100, normalized size = 1.45

$$\frac{8 \cos (bx+a)^8 + 32 \cos (bx+a)^6 - 115 \cos (bx+a)^4 + 38 \cos (bx+a)^2 + 192 (\cos (bx+a)^4 - 2 \cos (bx+a)^2 + 1) \log \left(\frac{1}{2} \sin (bx+a)\right) + 29}{32 (b \cos (bx+a)^4 - 2 b \cos (bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^9/sin(b*x+a)^5,x, algorithm="fricas")

[Out] 1/32*(8*cos(b*x + a)^8 + 32*cos(b*x + a)^6 - 115*cos(b*x + a)^4 + 38*cos(b*x + a)^2 + 192*(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*log(1/2*sin(b*x + a)) + 29)/(b*cos(b*x + a)^4 - 2*b*cos(b*x + a)^2 + b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 1664 vs. 2(58) = 116.

time = 10.64, size = 1664, normalized size = 24.12

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**9/sin(b*x+a)**5,x)

[Out] Piecewise((-384*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**12/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 1536*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**10/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 2304*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**8/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 1536*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**6/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 384*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)

```

)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/
2 + b*x/2)**4) + 384*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**12/(64*b*tan(a
/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 +
256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 1536*log(tan(a/2 +
b*x/2))*tan(a/2 + b*x/2)**10/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b
*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*ta
n(a/2 + b*x/2)**4) + 2304*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(64*b*t
an(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**
8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 1536*log(tan(a/
2 + b*x/2))*tan(a/2 + b*x/2)**6/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2
+ b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b
*tan(a/2 + b*x/2)**4) + 384*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(64*b
*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)
**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - tan(a/2 + b*x
/2)**16/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan
(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) +
24*tan(a/2 + b*x/2)**14/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)
**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2
+ b*x/2)**4) - 744*tan(a/2 + b*x/2)**10/(64*b*tan(a/2 + b*x/2)**12 + 256*b
*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)*
**6 + 64*b*tan(a/2 + b*x/2)**4) - 1182*tan(a/2 + b*x/2)**8/(64*b*tan(a/2 + b
*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*x/2)**8 + 256*b*
tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 744*tan(a/2 + b*x/2)**6/(
64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan(a/2 + b*
x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 24*tan(a/
2 + b*x/2)**2/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384
*b*tan(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)*
**4) - 1/(64*b*tan(a/2 + b*x/2)**12 + 256*b*tan(a/2 + b*x/2)**10 + 384*b*tan
(a/2 + b*x/2)**8 + 256*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4), N
e(b, 0)), (x*cos(a)**9/sin(a)**5, True))

```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 277 vs. 2(65) = 130.

time = 3.28, size = 277, normalized size = 4.01

$$\frac{\left(\frac{28(\cos(bx+a)-1) + 288(\cos(bx+a)-1)^2}{(\cos(bx+a)-1)^2} + 1\right)(\cos(bx+a)+1)^2 + \frac{28(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{32\left(\frac{84(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{126(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{84(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{25(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - 25\right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right)^4} - 192 \log\left(\frac{1 - \cos(bx+a)+1}{\cos(bx+a)+1}\right) + 384 \log\left(\frac{1 - \cos(bx+a)-1}{\cos(bx+a)+1} + 1\right)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)^9/sin(b*x+a)^5,x, algorithm="giac")
```

```
[Out] -1/64*((28*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 288*(cos(b*x + a) - 1)^2
/(cos(b*x + a) + 1)^2 + 1)*(cos(b*x + a) + 1)^2/(cos(b*x + a) - 1)^2 + 28*(
cos(b*x + a) - 1)/(cos(b*x + a) + 1) + (cos(b*x + a) - 1)^2/(cos(b*x + a) +
1)^2 + 32*(84*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 126*(cos(b*x + a) -
1)^2/(cos(b*x + a) + 1)^2 + 84*(cos(b*x + a) - 1)^3/(cos(b*x + a) + 1)^3 -
```


$$25*(\cos(b*x + a) - 1)^4/(\cos(b*x + a) + 1)^4 - 25)/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 1)^4 - 192*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)) + 384*\log(\text{abs}(-(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)))/b$$

Mupad [B]

time = 1.42, size = 92, normalized size = 1.33

$$\frac{6 \ln(\tan(a + bx))}{b} - \frac{3 \ln(\tan(a + bx)^2 + 1)}{b} + \frac{3 \tan(a + bx)^6 + \frac{9 \tan(a + bx)^4}{2} + \tan(a + bx)^2 - \frac{1}{4}}{b (\tan(a + bx)^8 + 2 \tan(a + bx)^6 + \tan(a + bx)^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^9/sin(a + b*x)^5,x)

[Out] (6*log(tan(a + b*x)))/b - (3*log(tan(a + b*x)^2 + 1))/b + (tan(a + b*x)^2 + (9*tan(a + b*x)^4)/2 + 3*tan(a + b*x)^6 - 1/4)/(b*(tan(a + b*x)^4 + 2*tan(a + b*x)^6 + tan(a + b*x)^8))

3.172 $\int \cos^3(a + bx) \cot^5(a + bx) dx$

Optimal. Leaf size=89

$$-\frac{35 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{35 \cos(a + bx)}{8b} + \frac{35 \cos^3(a + bx)}{24b} + \frac{7 \cos^3(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos^3(a + bx) \cot^4(a + bx)}{4b}$$

[Out] $-35/8*\operatorname{arctanh}(\cos(b*x+a))/b+35/8*\cos(b*x+a)/b+35/24*\cos(b*x+a)^3/b+7/8*\cos(b*x+a)^3*\cot(b*x+a)^2/b-1/4*\cos(b*x+a)^3*\cot(b*x+a)^4/b$

Rubi [A]

time = 0.04, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2672, 294, 308, 212}

$$\frac{35 \cos^3(a + bx)}{24b} + \frac{35 \cos(a + bx)}{8b} - \frac{\cos^3(a + bx) \cot^4(a + bx)}{4b} + \frac{7 \cos^3(a + bx) \cot^2(a + bx)}{8b} - \frac{35 \tanh^{-1}(\cos(a + bx))}{8b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]^3*\operatorname{Cot}[a + b*x]^5, x]$

[Out] $(-35*\operatorname{ArcTanh}[\operatorname{Cos}[a + b*x]])/(8*b) + (35*\operatorname{Cos}[a + b*x])/(8*b) + (35*\operatorname{Cos}[a + b*x]^3)/(24*b) + (7*\operatorname{Cos}[a + b*x]^3*\operatorname{Cot}[a + b*x]^2)/(8*b) - (\operatorname{Cos}[a + b*x]^3*\operatorname{Cot}[a + b*x]^4)/(4*b)$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[p, -1] \ \&\& \operatorname{GtQ}[m+1, n] \ \&\& \operatorname{!} \operatorname{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 308

$\operatorname{Int}[(x_)^{(m_)}((a_ + (b_)*(x_)^{(n_)}), x_Symbol] \rightarrow \operatorname{Int}[\operatorname{PolynomialDivide}[x^m, a + b*x^n, x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{IGtQ}[m, 0] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{GtQ}[m, 2*n-1]$

Rule 2672

```
Int[((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] :> With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(
ff*x)^(m + n)/(a^2 - ff^2*x^2)^(n + 1)/2], x], x, a*(Sin[e + f*x]/ff)], x]
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \cos^3(a + bx) \cot^5(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^8}{(1-x^2)^3} dx, x, \cos(a + bx)\right)}{b} \\
&= -\frac{\cos^3(a + bx) \cot^4(a + bx)}{4b} + \frac{7\text{Subst}\left(\int \frac{x^6}{(1-x^2)^2} dx, x, \cos(a + bx)\right)}{4b} \\
&= \frac{7\cos^3(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos^3(a + bx) \cot^4(a + bx)}{4b} - \frac{35\text{Subst}\left(\int \frac{x}{1-x^2} dx, x, \cos(a + bx)\right)}{4b} \\
&= \frac{7\cos^3(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos^3(a + bx) \cot^4(a + bx)}{4b} - \frac{35\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \cos(a + bx)\right)}{4b} \\
&= \frac{35\cos(a + bx)}{8b} + \frac{35\cos^3(a + bx)}{24b} + \frac{7\cos^3(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos^3(a + bx) \cot^4(a + bx)}{4b} \\
&= -\frac{35 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{35\cos(a + bx)}{8b} + \frac{35\cos^3(a + bx)}{24b} + \frac{7\cos^3(a + bx) \cot^2(a + bx)}{8b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 141, normalized size = 1.58

$$\frac{13\cos(a + bx)}{4b} + \frac{\cos(3(a + bx))}{12b} + \frac{13\csc^2\left(\frac{1}{2}(a + bx)\right)}{32b} - \frac{\csc^4\left(\frac{1}{2}(a + bx)\right)}{64b} - \frac{35\log\left(\cos\left(\frac{1}{2}(a + bx)\right)\right)}{8b} + \frac{35\log\left(\sin\left(\frac{1}{2}(a + bx)\right)\right)}{8b} - \frac{13\sec^2\left(\frac{1}{2}(a + bx)\right)}{32b} + \frac{\sec^4\left(\frac{1}{2}(a + bx)\right)}{64b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^3*Cot[a + b*x]^5,x]

[Out] (13*Cos[a + b*x])/(4*b) + Cos[3*(a + b*x)]/(12*b) + (13*Csc[(a + b*x)/2]^2)/(32*b) - Csc[(a + b*x)/2]^4/(64*b) - (35*Log[Cos[(a + b*x)/2]])/(8*b) + (35*Log[Sin[(a + b*x)/2]])/(8*b) - (13*Sec[(a + b*x)/2]^2)/(32*b) + Sec[(a + b*x)/2]^4/(64*b)

Maple [A]

time = 0.07, size = 98, normalized size = 1.10

method	result
derivativedivides	$ -\frac{\cos^9(bx+a)}{4\sin(bx+a)^4} + \frac{5(\cos^9(bx+a))}{8\sin(bx+a)^2} + \frac{5(\cos^7(bx+a))}{8} + \frac{7(\cos^5(bx+a))}{8} + \frac{35(\cos^3(bx+a))}{24} + \frac{35\cos(bx+a)}{8} + \frac{35\ln(\csc(bx+a) - \cot(bx+a))}{8} $

default	$\frac{-\frac{\cos^9(bx+a)}{4\sin(bx+a)^4} + \frac{5(\cos^9(bx+a))}{8\sin(bx+a)^2} + \frac{5(\cos^7(bx+a))}{8} + \frac{7(\cos^5(bx+a))}{8} + \frac{35(\cos^3(bx+a))}{24} + \frac{35\cos(bx+a)}{8} + \frac{35\ln(\csc(bx+a)-\cot(bx+a))}{8}}{b}$
norman	$\frac{-\frac{1}{64b} + \frac{21(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{64b} - \frac{21(\tan^{12}(\frac{bx}{2} + \frac{a}{2}))}{64b} + \frac{\tan^{14}(\frac{bx}{2} + \frac{a}{2})}{64b} + \frac{21(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{2b} + \frac{511(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{32b} + \frac{847(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{96b}}{(1+\tan^2(\frac{bx}{2} + \frac{a}{2}))^3 \tan(\frac{bx}{2} + \frac{a}{2})^4}$
risch	$\frac{e^{3i(bx+a)}}{24b} + \frac{13e^{i(bx+a)}}{8b} + \frac{13e^{-i(bx+a)}}{8b} + \frac{e^{-3i(bx+a)}}{24b} - \frac{13e^{7i(bx+a)} - 5e^{5i(bx+a)} - 5e^{3i(bx+a)} + 13e^{i(bx+a)}}{4b(e^{2i(bx+a)} - 1)^4} + 35\ln(\csc(bx+a) - \cot(bx+a))$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^8/sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

[Out] $1/b*(-1/4*\cos(b*x+a)^9/\sin(b*x+a)^4+5/8/\sin(b*x+a)^2*\cos(b*x+a)^9+5/8*\cos(b*x+a)^7+7/8*\cos(b*x+a)^5+35/24*\cos(b*x+a)^3+35/8*\cos(b*x+a)+35/8*\ln(\csc(b*x+a)-\cot(b*x+a)))$

Maxima [A]

time = 0.28, size = 89, normalized size = 1.00

$$\frac{16 \cos(bx+a)^3 - \frac{6(13 \cos(bx+a)^3 - 11 \cos(bx+a))}{\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1} + 144 \cos(bx+a) - 105 \log(\cos(bx+a) + 1) + 105 \log(\cos(bx+a) - 1)}{48b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8/sin(b*x+a)^5,x, algorithm="maxima")`

[Out] $1/48*(16*\cos(b*x+a)^3 - 6*(13*\cos(b*x+a)^3 - 11*\cos(b*x+a))/(\cos(b*x+a)^4 - 2*\cos(b*x+a)^2 + 1) + 144*\cos(b*x+a) - 105*\log(\cos(b*x+a) + 1) + 105*\log(\cos(b*x+a) - 1))/b$

Fricas [A]

time = 0.41, size = 132, normalized size = 1.48

$$\frac{16 \cos(bx+a)^7 + 112 \cos(bx+a)^5 - 350 \cos(bx+a)^3 - 105 (\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log(\frac{1}{2} \cos(bx+a) + \frac{1}{2}) + 105 (\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}) + 210 \cos(bx+a)}{48 (b \cos(bx+a)^4 - 2b \cos(bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^8/sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $1/48*(16*\cos(b*x+a)^7 + 112*\cos(b*x+a)^5 - 350*\cos(b*x+a)^3 - 105*(\cos(b*x+a)^4 - 2*\cos(b*x+a)^2 + 1)*\log(1/2*\cos(b*x+a) + 1/2) + 105*(\cos(b*x+a)^4 - 2*\cos(b*x+a)^2 + 1)*\log(-1/2*\cos(b*x+a) + 1/2) + 210*\cos(b*x+a))/(b*\cos(b*x+a)^4 - 2*b*\cos(b*x+a)^2 + b)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 869 vs. 2(80) = 160.

time = 5.73, size = 869, normalized size = 9.76

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**8/sin(b*x+a)**5,x)

[Out] Piecewise((840*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**10/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 2520*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 2520*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 840*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 3*tan(a/2 + b*x/2)**14/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) - 63*tan(a/2 + b*x/2)**12/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 2016*tan(a/2 + b*x/2)**8/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 3066*tan(a/2 + b*x/2)**6/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 1694*tan(a/2 + b*x/2)**4/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) + 63*tan(a/2 + b*x/2)**2/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4) - 3/(192*b*tan(a/2 + b*x/2)**10 + 576*b*tan(a/2 + b*x/2)**8 + 576*b*tan(a/2 + b*x/2)**6 + 192*b*tan(a/2 + b*x/2)**4), Ne(b, 0)), (x*cos(a)**8/sin(a)**5, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(79) = 158.

time = 3.35, size = 209, normalized size = 2.35

$$\frac{3 \left(\frac{24(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{210(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 1 \right) (\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} - \frac{72(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - \frac{256 \left(\frac{9(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{6(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 5 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1 \right)^3} - 420 \log \left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|} \right)$$

192b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^8/sin(b*x+a)^5,x, algorithm="giac")

[Out] -1/192*(3*(24*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 210*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 1)*(cos(b*x + a) + 1)^2/(cos(b*x + a) - 1)^2 - 72*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 3*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 256*(9*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 6*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 5)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)^3 - 420*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.55, size = 157, normalized size = 1.76

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{64b} - \frac{3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8b} + \frac{35 \ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b} + \frac{\frac{67 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8}{8} + \frac{839 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6}{64} + \frac{1487 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{192} + \frac{21 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{64} - \frac{1}{64}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^{10} + 3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^8 + 3 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^8/sin(a + b*x)^5,x)

[Out] tan(a/2 + (b*x)/2)^4/(64*b) - (3*tan(a/2 + (b*x)/2)^2)/(8*b) + (35*log(tan(a/2 + (b*x)/2)))/(8*b) + ((21*tan(a/2 + (b*x)/2)^2)/64 + (1487*tan(a/2 + (b*x)/2)^4)/192 + (839*tan(a/2 + (b*x)/2)^6)/64 + (67*tan(a/2 + (b*x)/2)^8)/8 - 1/64)/(b*(tan(a/2 + (b*x)/2)^4 + 3*tan(a/2 + (b*x)/2)^6 + 3*tan(a/2 + (b*x)/2)^8 + tan(a/2 + (b*x)/2)^10))

3.173 $\int \cos^2(a + bx) \cot^5(a + bx) dx$

Optimal. Leaf size=58

$$\frac{3 \csc^2(a + bx)}{2b} - \frac{\csc^4(a + bx)}{4b} + \frac{3 \log(\sin(a + bx))}{b} - \frac{\sin^2(a + bx)}{2b}$$

[Out] $3/2*\csc(b*x+a)^2/b-1/4*\csc(b*x+a)^4/b+3*\ln(\sin(b*x+a))/b-1/2*\sin(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2670, 272, 45}

$$-\frac{\sin^2(a + bx)}{2b} - \frac{\csc^4(a + bx)}{4b} + \frac{3 \csc^2(a + bx)}{2b} + \frac{3 \log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*Cot[a + b*x]^5,x]

[Out] $(3*\text{Csc}[a + b*x]^2)/(2*b) - \text{Csc}[a + b*x]^4/(4*b) + (3*\text{Log}[\text{Sin}[a + b*x]])/b - \text{Sin}[a + b*x]^2/(2*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2670

Int[sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-f^(-1), Subst[Int[(1 - x^2)^((m + n - 1)/2)/x^n, x], x, Cos[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \cos^2(a+bx) \cot^5(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1-x^2)^3}{x^5} dx, x, -\sin(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1-x)^3}{x^3} dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(-1 + \frac{1}{x^3} - \frac{3}{x^2} + \frac{3}{x}\right) dx, x, \sin^2(a+bx)\right)}{2b} \\
&= \frac{3 \csc^2(a+bx)}{2b} - \frac{\csc^4(a+bx)}{4b} + \frac{3 \log(\sin(a+bx))}{b} - \frac{\sin^2(a+bx)}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 47, normalized size = 0.81

$$\frac{6 \csc^2(a+bx) - \csc^4(a+bx) + 12 \log(\sin(a+bx)) - 2 \sin^2(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^2*Cot[a + b*x]^5,x]``[Out] (6*Csc[a + b*x]^2 - Csc[a + b*x]^4 + 12*Log[Sin[a + b*x]] - 2*Sin[a + b*x]^2)/(4*b)`**Maple [A]**

time = 0.06, size = 81, normalized size = 1.40

method	result
derivativedivides	$\frac{-\frac{\cos^8(bx+a)}{4 \sin(bx+a)^4} + \frac{\cos^8(bx+a)}{2 \sin(bx+a)^2} + \frac{(\cos^6(bx+a))}{2} + \frac{3(\cos^4(bx+a))}{4} + \frac{3(\cos^2(bx+a))}{2} + 3 \ln(\sin(bx+a))}{b}$
default	$\frac{-\frac{\cos^8(bx+a)}{4 \sin(bx+a)^4} + \frac{\cos^8(bx+a)}{2 \sin(bx+a)^2} + \frac{(\cos^6(bx+a))}{2} + \frac{3(\cos^4(bx+a))}{4} + \frac{3(\cos^2(bx+a))}{2} + 3 \ln(\sin(bx+a))}{b}$
risch	$-3ix + \frac{e^{2i(bx+a)}}{8b} + \frac{e^{-2i(bx+a)}}{8b} - \frac{6ia}{b} - \frac{2(3e^{6i(bx+a)} - 4e^{4i(bx+a)} + 3e^{2i(bx+a)})}{b(e^{2i(bx+a)} - 1)^4} + \frac{3 \ln(e^{2i(bx+a)} - 1)}{b}$
norman	$-\frac{1}{64b} + \frac{9\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{32b} + \frac{9\left(\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{32b} - \frac{\tan^{12}\left(\frac{bx}{2} + \frac{a}{2}\right)}{64b} - \frac{83\left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{32b} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{3 \ln\left(1 + \tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^7/sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/4*cos(b*x+a)^8/sin(b*x+a)^4+1/2/sin(b*x+a)^2*cos(b*x+a)^8+1/2*cos(b*x+a)^6+3/4*cos(b*x+a)^4+3/2*cos(b*x+a)^2+3*ln(sin(b*x+a)))`

Maxima [A]

time = 0.28, size = 49, normalized size = 0.84

$$\frac{2 \sin (bx+a)^2 - \frac{6 \sin (bx+a)^2-1}{\sin (bx+a)^4} - 6 \log (\sin (bx+a)^2)}{4 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/4*(2*sin(b*x + a)^2 - (6*sin(b*x + a)^2 - 1)/sin(b*x + a)^4 - 6*log(sin(b*x + a)^2))/b

Fricas [A]

time = 0.38, size = 90, normalized size = 1.55

$$\frac{2 \cos (bx+a)^6 - 5 \cos (bx+a)^4 - 2 \cos (bx+a)^2 + 12 (\cos (bx+a)^4 - 2 \cos (bx+a)^2 + 1) \log \left(\frac{1}{2} \sin (bx+a)\right) + 4}{4 (b \cos (bx+a)^4 - 2 b \cos (bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^5,x, algorithm="fricas")

[Out] 1/4*(2*cos(b*x + a)^6 - 5*cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 12*(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*log(1/2*sin(b*x + a)) + 4)/(b*cos(b*x + a)^4 - 2*b*cos(b*x + a)^2 + b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 733 vs. 2(48) = 96.

time = 4.86, size = 733, normalized size = 12.64

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**7/sin(b*x+a)**5,x)

[Out] Piecewise((-192*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**8/(64*b*tan(a/2 + b*x/2)**8 + 128*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 384*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**6/(64*b*tan(a/2 + b*x/2)**8 + 128*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 192*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(64*b*tan(a/2 + b*x/2)**8 + 128*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 192*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(64*b*tan(a/2 + b*x/2)**8 + 128*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 384*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(64*b*tan(a/2 + b*x/2)**8 + 128*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 192*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(64*b*tan(a/2 + b*x/2)**8 + 128*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - tan(a/2 + b*

$x/2)^{12}/(64*b*\tan(a/2 + b*x/2)^8 + 128*b*\tan(a/2 + b*x/2)^6 + 64*b*\tan(a/2 + b*x/2)^4) + 18*\tan(a/2 + b*x/2)^{10}/(64*b*\tan(a/2 + b*x/2)^8 + 128*b*\tan(a/2 + b*x/2)^6 + 64*b*\tan(a/2 + b*x/2)^4) - 166*\tan(a/2 + b*x/2)^6/(64*b*\tan(a/2 + b*x/2)^8 + 128*b*\tan(a/2 + b*x/2)^6 + 64*b*\tan(a/2 + b*x/2)^4) + 18*\tan(a/2 + b*x/2)^2/(64*b*\tan(a/2 + b*x/2)^8 + 128*b*\tan(a/2 + b*x/2)^6 + 64*b*\tan(a/2 + b*x/2)^4) - 1/(64*b*\tan(a/2 + b*x/2)^8 + 128*b*\tan(a/2 + b*x/2)^6 + 64*b*\tan(a/2 + b*x/2)^4), Ne(b, 0)), (x*cos(a))^{7/5}, True))$

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 232 vs. 2(52) = 104.

time = 4.29, size = 232, normalized size = 4.00

$$\frac{20 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{18 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - \frac{111 (\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{36 (\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} - \frac{72 (\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + 1}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^2 \left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^2}}{64b} - 96 \log\left(\frac{|\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) + 192 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right|\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^7/sin(b*x+a)^5,x, algorithm="giac")

[Out] $-1/64*(20*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + (\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + (18*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 111*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 36*(\cos(b*x + a) - 1)^3/(\cos(b*x + a) + 1)^3 - 72*(\cos(b*x + a) - 1)^4/(\cos(b*x + a) + 1)^4 + 1)/((\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - (\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2)^2 - 96*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)) + 192*\log(\text{abs}(-(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)))/b$

Mupad [B]

time = 0.64, size = 74, normalized size = 1.28

$$\frac{3 \ln(\tan(a + bx))}{b} - \frac{3 \ln(\tan(a + bx)^2 + 1)}{2b} + \frac{\frac{3 \tan(a+bx)^4}{2} + \frac{3 \tan(a+bx)^2}{4} - \frac{1}{4}}{b (\tan(a + bx)^6 + \tan(a + bx)^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^7/sin(a + b*x)^5,x)

[Out] $(3*\log(\tan(a + b*x)))/b - (3*\log(\tan(a + b*x)^2 + 1))/(2*b) + ((3*\tan(a + b*x)^2)/4 + (3*\tan(a + b*x)^4)/2 - 1/4)/(b*(\tan(a + b*x)^4 + \tan(a + b*x)^6))$

3.174 $\int \cos(a + bx) \cot^5(a + bx) dx$

Optimal. Leaf size=70

$$-\frac{15 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{15 \cos(a + bx)}{8b} + \frac{5 \cos(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos(a + bx) \cot^4(a + bx)}{4b}$$

[Out] $-15/8*\operatorname{arctanh}(\cos(b*x+a))/b+15/8*\cos(b*x+a)/b+5/8*\cos(b*x+a)*\cot(b*x+a)^2/b-1/4*\cos(b*x+a)*\cot(b*x+a)^4/b$

Rubi [A]

time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {2672, 294, 327, 212}

$$\frac{15 \cos(a + bx)}{8b} - \frac{\cos(a + bx) \cot^4(a + bx)}{4b} + \frac{5 \cos(a + bx) \cot^2(a + bx)}{8b} - \frac{15 \tanh^{-1}(\cos(a + bx))}{8b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Cos}[a + b*x]*\operatorname{Cot}[a + b*x]^5, x]$

[Out] $(-15*\operatorname{ArcTanh}[\operatorname{Cos}[a + b*x]])/(8*b) + (15*\operatorname{Cos}[a + b*x])/(8*b) + (5*\operatorname{Cos}[a + b*x]*\operatorname{Cot}[a + b*x]^2)/(8*b) - (\operatorname{Cos}[a + b*x]*\operatorname{Cot}[a + b*x]^4)/(4*b)$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \operatorname{LtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[p, -1] \ \&\& \operatorname{GtQ}[m+1, n] \ \&\& \operatorname{!} \operatorname{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 327

$\operatorname{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*(m+n*p+1))), x] - \operatorname{Dist}[a*c^n*((m-n+1)/(b*(m+n*p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, p\}, x] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{GtQ}[m, n-1] \ \&\& \operatorname{NeQ}[m+n*p+1, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 2672

```
Int[((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.)*tan[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := With[{ff = FreeFactors[Sin[e + f*x], x]}, Dist[ff/f, Subst[Int[(
ff*x)^(m + n)/(a^2 - ff^2*x^2)^((n + 1)/2), x], x, a*(Sin[e + f*x]/ff)], x
] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \cos(a + bx) \cot^5(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^6}{(1-x^2)^3} dx, x, \cos(a + bx)\right)}{b} \\
&= -\frac{\cos(a + bx) \cot^4(a + bx)}{4b} + \frac{5\text{Subst}\left(\int \frac{x^4}{(1-x^2)^2} dx, x, \cos(a + bx)\right)}{4b} \\
&= \frac{5 \cos(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos(a + bx) \cot^4(a + bx)}{4b} - \frac{15\text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \cos(a + bx)\right)}{8b} \\
&= \frac{15 \cos(a + bx)}{8b} + \frac{5 \cos(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos(a + bx) \cot^4(a + bx)}{4b} - \frac{15 \tanh^{-1}(\cos(a + bx))}{8b} \\
&= -\frac{15 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{15 \cos(a + bx)}{8b} + \frac{5 \cos(a + bx) \cot^2(a + bx)}{8b} - \frac{\cos(a + bx) \cot^4(a + bx)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 123, normalized size = 1.76

$$\frac{\cos(a + bx)}{b} + \frac{9 \csc^2\left(\frac{1}{2}(a + bx)\right)}{32b} - \frac{\csc^4\left(\frac{1}{2}(a + bx)\right)}{64b} - \frac{15 \log\left(\cos\left(\frac{1}{2}(a + bx)\right)\right)}{8b} + \frac{15 \log\left(\sin\left(\frac{1}{2}(a + bx)\right)\right)}{8b} - \frac{9 \sec^2\left(\frac{1}{2}(a + bx)\right)}{32b} + \frac{\sec^4\left(\frac{1}{2}(a + bx)\right)}{64b}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*Cot[a + b*x]^5, x]

[Out] Cos[a + b*x]/b + (9*Csc[(a + b*x)/2]^2)/(32*b) - Csc[(a + b*x)/2]^4/(64*b) - (15*Log[Cos[(a + b*x)/2]])/(8*b) + (15*Log[Sin[(a + b*x)/2]])/(8*b) - (9*Sec[(a + b*x)/2]^2)/(32*b) + Sec[(a + b*x)/2]^4/(64*b)

Maple [A]

time = 0.05, size = 88, normalized size = 1.26

method	result
derivativedivides	$ -\frac{\cos^7(bx+a)}{4 \sin(bx+a)^4} + \frac{3(\cos^7(bx+a))}{8 \sin(bx+a)^2} + \frac{3(\cos^5(bx+a))}{8} + \frac{5(\cos^3(bx+a))}{8} + \frac{15 \cos(bx+a)}{8} + \frac{15 \ln(\csc(bx+a) - \cot(bx+a))}{8} $
default	$ -\frac{\cos^7(bx+a)}{4 \sin(bx+a)^4} + \frac{3(\cos^7(bx+a))}{8 \sin(bx+a)^2} + \frac{3(\cos^5(bx+a))}{8} + \frac{5(\cos^3(bx+a))}{8} + \frac{15 \cos(bx+a)}{8} + \frac{15 \ln(\csc(bx+a) - \cot(bx+a))}{8} $

norman	$\frac{-\frac{1}{64b} + \frac{15(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{64b} - \frac{15(\tan^8(\frac{bx}{2} + \frac{a}{2}))}{64b} + \frac{\tan^{10}(\frac{bx}{2} + \frac{a}{2})}{64b} + \frac{5(\tan^4(\frac{bx}{2} + \frac{a}{2}))}{2b}}{(1 + \tan^2(\frac{bx}{2} + \frac{a}{2})) \tan(\frac{bx}{2} + \frac{a}{2})^4} + \frac{15 \ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{8b}$
risch	$\frac{e^{i(bx+a)}}{2b} + \frac{e^{-i(bx+a)}}{2b} - \frac{9e^{7i(bx+a)} - e^{5i(bx+a)} - e^{3i(bx+a)} + 9e^{i(bx+a)}}{4b(e^{2i(bx+a)} - 1)^4} + \frac{15 \ln(e^{i(bx+a)} - 1)}{8b} - \frac{15 \ln(e^{i(bx+a)} + 1)}{8b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^6/sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

[Out] $1/b * (-1/4 * \cos(b*x+a)^7 / \sin(b*x+a)^4 + 3/8 * \cos(b*x+a)^7 / \sin(b*x+a)^2 + 3/8 * \cos(b*x+a)^5 + 5/8 * \cos(b*x+a)^3 + 15/8 * \cos(b*x+a) + 15/8 * \ln(\csc(b*x+a) - \cot(b*x+a)))$

Maxima [A]

time = 0.29, size = 79, normalized size = 1.13

$$\frac{2(9 \cos(bx+a)^3 - 7 \cos(bx+a))}{\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1} - 16 \cos(bx+a) + 15 \log(\cos(bx+a) + 1) - 15 \log(\cos(bx+a) - 1)$$

$$16b$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^5,x, algorithm="maxima")`

[Out] $-1/16 * (2 * (9 * \cos(b*x + a)^3 - 7 * \cos(b*x + a)) / (\cos(b*x + a)^4 - 2 * \cos(b*x + a)^2 + 1) - 16 * \cos(b*x + a) + 15 * \log(\cos(b*x + a) + 1) - 15 * \log(\cos(b*x + a) - 1)) / b$

Fricas [A]

time = 0.37, size = 122, normalized size = 1.74

$$\frac{16 \cos(bx+a)^5 - 50 \cos(bx+a)^3 - 15(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log(\frac{1}{2} \cos(bx+a) + \frac{1}{2}) + 15(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}) + 30 \cos(bx+a)}{16(b \cos(bx+a)^4 - 2b \cos(bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^6/sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $1/16 * (16 * \cos(b*x + a)^5 - 50 * \cos(b*x + a)^3 - 15 * (\cos(b*x + a)^4 - 2 * \cos(b*x + a)^2 + 1) * \log(1/2 * \cos(b*x + a) + 1/2) + 15 * (\cos(b*x + a)^4 - 2 * \cos(b*x + a)^2 + 1) * \log(-1/2 * \cos(b*x + a) + 1/2) + 30 * \cos(b*x + a)) / (b * \cos(b*x + a)^4 - 2 * b * \cos(b*x + a)^2 + b)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 330 vs. $2(63) = 126$.

time = 1.99, size = 330, normalized size = 4.71

$$\left\{ \begin{array}{l} \frac{120 \log(\tan(\frac{x}{2} + \frac{a}{2})) \tan^6(\frac{x}{2} + \frac{a}{2})}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} + \frac{120 \log(\tan(\frac{x}{2} + \frac{a}{2})) \tan^4(\frac{x}{2} + \frac{a}{2})}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} + \frac{\tan^{10}(\frac{x}{2} + \frac{a}{2})}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} - \frac{15 \tan^8(\frac{x}{2} + \frac{a}{2})}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} + \frac{160 \tan^6(\frac{x}{2} + \frac{a}{2})}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} + \frac{15 \tan^2(\frac{x}{2} + \frac{a}{2})}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} - \frac{1}{64b \tan^6(\frac{x}{2} + \frac{a}{2}) + 64b \tan^4(\frac{x}{2} + \frac{a}{2})} \end{array} \right. \text{for } b \neq 0$$

otherwise

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**6/sin(b*x+a)**5,x)

[Out] Piecewise((120*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 120*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + tan(a/2 + b*x/2)**10/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 15*tan(a/2 + b*x/2)**8/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 160*tan(a/2 + b*x/2)**4/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) + 15*tan(a/2 + b*x/2)**2/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4) - 1/(64*b*tan(a/2 + b*x/2)**6 + 64*b*tan(a/2 + b*x/2)**4), Ne(b, 0)), (x*cos(a)**6/sin(a)**5, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 164 vs. 2(62) = 124.

time = 5.31, size = 164, normalized size = 2.34

$$\frac{\left(\frac{16(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{90(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 1\right)(\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} - \frac{16(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{128}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1} - 60 \log\left(\frac{|\cos(bx+a)+1|}{|\cos(bx+a)-1|}\right)$$

64 b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^6/sin(b*x+a)^5,x, algorithm="giac")

[Out] -1/64*((16*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 90*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 1)*(cos(b*x + a) + 1)^2/(cos(b*x + a) - 1)^2 - 16*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - (cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 128/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1) - 60*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.58, size = 105, normalized size = 1.50

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{64b} - \frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{4b} + \frac{15 \ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b} + \frac{\frac{9 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{4} + \frac{15 \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{64} - \frac{1}{64}}{b \left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^6 + \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^6/sin(a + b*x)^5,x)

[Out] tan(a/2 + (b*x)/2)^4/(64*b) - tan(a/2 + (b*x)/2)^2/(4*b) + (15*log(tan(a/2 + (b*x)/2)))/(8*b) + ((15*tan(a/2 + (b*x)/2)^2)/64 + (9*tan(a/2 + (b*x)/2)^4)/4 - 1/64)/(b*(tan(a/2 + (b*x)/2)^4 + tan(a/2 + (b*x)/2)^6))

3.175 $\int \cot^5(a + bx) dx$

Optimal. Leaf size=42

$$\frac{\cot^2(a + bx)}{2b} - \frac{\cot^4(a + bx)}{4b} + \frac{\log(\sin(a + bx))}{b}$$

[Out] $1/2*\cot(b*x+a)^2/b-1/4*\cot(b*x+a)^4/b+\ln(\sin(b*x+a))/b$

Rubi [A]

time = 0.02, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3554, 3556}

$$-\frac{\cot^4(a + bx)}{4b} + \frac{\cot^2(a + bx)}{2b} + \frac{\log(\sin(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^5, x]

[Out] Cot[a + b*x]^2/(2*b) - Cot[a + b*x]^4/(4*b) + Log[Sin[a + b*x]]/b

Rule 3554

Int[((b_.)*tan[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[b*((b*Tan[c + d*x])^(n - 1)/(d*(n - 1))), x] - Dist[b^2, Int[(b*Tan[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1]

Rule 3556

Int[tan[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-Log[RemoveContent[Cos[c + d*x], x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \cot^5(a + bx) dx &= -\frac{\cot^4(a + bx)}{4b} - \int \cot^3(a + bx) dx \\ &= \frac{\cot^2(a + bx)}{2b} - \frac{\cot^4(a + bx)}{4b} + \int \cot(a + bx) dx \\ &= \frac{\cot^2(a + bx)}{2b} - \frac{\cot^4(a + bx)}{4b} + \frac{\log(\sin(a + bx))}{b} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 46, normalized size = 1.10

$$\frac{2 \cot^2(a + bx) - \cot^4(a + bx) + 4 \log(\cos(a + bx)) + 4 \log(\tan(a + bx))}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^5,x]

[Out] (2*Cot[a + b*x]^2 - Cot[a + b*x]^4 + 4*Log[Cos[a + b*x]] + 4*Log[Tan[a + b*x]])/(4*b)

Maple [A]

time = 0.04, size = 33, normalized size = 0.79

method	result	size
derivativedivides	$-\frac{(\cot^4(bx+a))}{4} + \frac{(\cot^2(bx+a))}{2} + \ln(\sin(bx+a))$	33
default	$-\frac{(\cot^4(bx+a))}{4} + \frac{(\cot^2(bx+a))}{2} + \ln(\sin(bx+a))$	33
risch	$-ix - \frac{2ia}{b} - \frac{4(e^{6i(bx+a)} - e^{4i(bx+a)} + e^{2i(bx+a)})}{b(e^{2i(bx+a)} - 1)^4} + \frac{\ln(e^{2i(bx+a)} - 1)}{b}$	77
norman	$-\frac{1}{64b} + \frac{3(\tan^2(\frac{bx}{2} + \frac{a}{2}))}{16b} + \frac{3(\tan^6(\frac{bx}{2} + \frac{a}{2}))}{16b} - \frac{\tan^8(\frac{bx}{2} + \frac{a}{2})}{64b} + \frac{\ln(\tan(\frac{bx}{2} + \frac{a}{2}))}{b} - \frac{\ln(1 + \tan^2(\frac{bx}{2} + \frac{a}{2}))}{b}$	101

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^5/sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/4*cot(b*x+a)^4+1/2*cot(b*x+a)^2+ln(sin(b*x+a)))

Maxima [A]

time = 0.28, size = 38, normalized size = 0.90

$$\frac{\frac{4 \sin(bx+a)^2 - 1}{\sin(bx+a)^4} + 2 \log(\sin(bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a)^5,x, algorithm="maxima")

[Out] 1/4*((4*sin(b*x + a)^2 - 1)/sin(b*x + a)^4 + 2*log(sin(b*x + a)^2))/b

Fricas [A]

time = 0.41, size = 70, normalized size = 1.67

$$\frac{4 \cos(bx+a)^2 - 4(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log\left(\frac{1}{2} \sin(bx+a)\right) - 3}{4(b \cos(bx+a)^4 - 2b \cos(bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5/sin(b*x+a)^5,x, algorithm="fricas")

[Out] $-1/4*(4*\cos(b*x + a)^2 - 4*(\cos(b*x + a)^4 - 2*\cos(b*x + a)^2 + 1)*\log(1/2*\sin(b*x + a)) - 3)/(b*\cos(b*x + a)^4 - 2*b*\cos(b*x + a)^2 + b)$

Sympy [A]

time = 0.61, size = 61, normalized size = 1.45

$$\begin{cases} \frac{\log(\sin(a+bx))}{b} + \frac{\cos^2(a+bx)}{2b\sin^2(a+bx)} - \frac{\cos^4(a+bx)}{4b\sin^4(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^5(a)}{\sin^5(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**5/sin(b*x+a)**5,x)`

[Out] `Piecewise((log(sin(a + b*x))/b + cos(a + b*x)**2/(2*b*sin(a + b*x)**2) - cos(a + b*x)**4/(4*b*sin(a + b*x)**4), Ne(b, 0)), (x*cos(a)**5/sin(a)**5, True))`

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 164 vs. 2(38) = 76.

time = 4.90, size = 164, normalized size = 3.90

$$\frac{\left(\frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{48(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 1\right)(\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} + \frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 32 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right) + 64 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1\right|\right)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^5/sin(b*x+a)^5,x, algorithm="giac")`

[Out] $-1/64*((12*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 48*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 1)*(\cos(b*x + a) + 1)^2/(\cos(b*x + a) - 1)^2 + 12*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + (\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 - 32*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)) + 64*\log(\text{abs}(-(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 1)))/b$

Mupad [B]

time = 0.42, size = 52, normalized size = 1.24

$$\frac{\ln(\tan(a+bx))}{b} - \frac{\ln(\tan(a+bx)^2 + 1)}{2b} + \frac{\frac{\tan(a+bx)^2}{2} - \frac{1}{4}}{b \tan(a+bx)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^5/sin(a + b*x)^5,x)`

[Out] $\log(\tan(a + b*x))/b - \log(\tan(a + b*x)^2 + 1)/(2*b) + (\tan(a + b*x)^2/2 - 1/4)/(b*\tan(a + b*x)^4)$

3.176 $\int \cot^4(a + bx) \csc(a + bx) dx$

Optimal. Leaf size=55

$$-\frac{3 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{3 \cot(a + bx) \csc(a + bx)}{8b} - \frac{\cot^3(a + bx) \csc(a + bx)}{4b}$$

[Out] $-3/8*\operatorname{arctanh}(\cos(b*x+a))/b+3/8*\cot(b*x+a)*\csc(b*x+a)/b-1/4*\cot(b*x+a)^3*\csc(b*x+a)/b$

Rubi [A]

time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$,

Rules used = {2691, 3855}

$$-\frac{3 \tanh^{-1}(\cos(a + bx))}{8b} - \frac{\cot^3(a + bx) \csc(a + bx)}{4b} + \frac{3 \cot(a + bx) \csc(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Cot}[a + b*x]^4*\operatorname{Csc}[a + b*x], x]$

[Out] $(-3*\operatorname{ArcTanh}[\operatorname{Cos}[a + b*x]])/(8*b) + (3*\operatorname{Cot}[a + b*x]*\operatorname{Csc}[a + b*x])/(8*b) - (\operatorname{Cot}[a + b*x]^3*\operatorname{Csc}[a + b*x])/(4*b)$

Rule 2691

$\operatorname{Int}[(a_*)*\sec[(e_*) + (f_*)*(x_*)]^{(m_*)}*((b_*)*\tan[(e_*) + (f_*)*(x_*)])^{(n_*)}, x_Symbol] \rightarrow \operatorname{Simp}[b*(a*\sec[e + f*x])^{m*((b*\tan[e + f*x])^{(n-1)}/(f*(m+n-1)))}, x] - \operatorname{Dist}[b^2*((n-1)/(m+n-1)), \operatorname{Int}[(a*\sec[e + f*x])^{m*(b*\tan[e + f*x])^{(n-2)}}, x], x] /;$ $\operatorname{FreeQ}\{a, b, e, f, m\}, x \ \&\& \operatorname{GtQ}[n, 1] \ \&\& \operatorname{NeQ}[m+n-1, 0] \ \&\& \operatorname{IntegersQ}[2*m, 2*n]$

Rule 3855

$\operatorname{Int}[\csc[(c_*) + (d_*)*(x_*)], x_Symbol] \rightarrow \operatorname{Simp}[-\operatorname{ArcTanh}[\operatorname{Cos}[c + d*x]]/d, x] /;$ $\operatorname{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \cot^4(a + bx) \csc(a + bx) dx &= -\frac{\cot^3(a + bx) \csc(a + bx)}{4b} - \frac{3}{4} \int \cot^2(a + bx) \csc(a + bx) dx \\ &= \frac{3 \cot(a + bx) \csc(a + bx)}{8b} - \frac{\cot^3(a + bx) \csc(a + bx)}{4b} + \frac{3}{8} \int \csc(a + bx) dx \\ &= -\frac{3 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{3 \cot(a + bx) \csc(a + bx)}{8b} - \frac{\cot^3(a + bx) \csc(a + bx)}{4b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 113 vs. 2(55) = 110.

time = 0.03, size = 113, normalized size = 2.05

$$\frac{5 \csc^2\left(\frac{1}{2}(a+bx)\right)}{32b} - \frac{\csc^4\left(\frac{1}{2}(a+bx)\right)}{64b} - \frac{3 \log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{8b} + \frac{3 \log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{8b} - \frac{5 \sec^2\left(\frac{1}{2}(a+bx)\right)}{32b} + \frac{\sec^4\left(\frac{1}{2}(a+bx)\right)}{64b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^4*Csc[a + b*x],x]

[Out] (5*Csc[(a + b*x)/2]^2)/(32*b) - Csc[(a + b*x)/2]^4/(64*b) - (3*Log[Cos[(a + b*x)/2]])/(8*b) + (3*Log[Sin[(a + b*x)/2]])/(8*b) - (5*Sec[(a + b*x)/2]^2)/(32*b) + Sec[(a + b*x)/2]^4/(64*b)

Maple [A]

time = 0.05, size = 78, normalized size = 1.42

method	result	size
derivativedivides	$\frac{-\frac{\cos^5(bx+a)}{4 \sin(bx+a)^4} + \frac{\cos^5(bx+a)}{8 \sin(bx+a)^2} + \frac{\left(\frac{\cos^3(bx+a)}{8} + \frac{3 \cos(bx+a)}{8} + \frac{3 \ln(\csc(bx+a) - \cot(bx+a))}{8}\right)}{b}}{b}$	78
default	$\frac{-\frac{\cos^5(bx+a)}{4 \sin(bx+a)^4} + \frac{\cos^5(bx+a)}{8 \sin(bx+a)^2} + \frac{\left(\frac{\cos^3(bx+a)}{8} + \frac{3 \cos(bx+a)}{8} + \frac{3 \ln(\csc(bx+a) - \cot(bx+a))}{8}\right)}{b}}{b}$	78
norman	$\frac{-\frac{1}{64b} + \frac{\tan^2\left(\frac{bx+a}{2}\right)}{8b} - \frac{\tan^6\left(\frac{bx+a}{2}\right)}{8b} + \frac{\tan^8\left(\frac{bx+a}{2}\right)}{64b}}{\tan\left(\frac{bx+a}{2}\right)^4} + \frac{3 \ln\left(\tan\left(\frac{bx+a}{2}\right)\right)}{8b}$	83
risch	$-\frac{5e^{7i(bx+a)} + 3e^{5i(bx+a)} + 3e^{3i(bx+a)} + 5e^{i(bx+a)}}{4b(e^{2i(bx+a)} - 1)^4} + \frac{3 \ln(e^{i(bx+a)} - 1)}{8b} - \frac{3 \ln(e^{i(bx+a)} + 1)}{8b}$	99

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4/sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/4*cos(b*x+a)^5/sin(b*x+a)^4+1/8*cos(b*x+a)^5/sin(b*x+a)^2+1/8*cos(b*x+a)^3+3/8*cos(b*x+a)+3/8*ln(csc(b*x+a)-cot(b*x+a)))

Maxima [A]

time = 0.29, size = 71, normalized size = 1.29

$$\frac{2\left(5 \cos(bx+a)^3 - 3 \cos(bx+a)\right)}{\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1} + \frac{3 \log(\cos(bx+a) + 1) - 3 \log(\cos(bx+a) - 1)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/16*(2*(5*cos(b*x + a)^3 - 3*cos(b*x + a))/(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1) + 3*log(cos(b*x + a) + 1) - 3*log(cos(b*x + a) - 1))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 112 vs. 2(49) = 98.

time = 0.35, size = 112, normalized size = 2.04

$$\frac{10 \cos(bx+a)^3 + 3(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 3(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) - 6 \cos(bx+a)}{16(b \cos(bx+a)^4 - 2b \cos(bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^5,x, algorithm="fricas")

[Out] $-1/16*(10*\cos(b*x + a)^3 + 3*(\cos(b*x + a)^4 - 2*\cos(b*x + a)^2 + 1)*\log(1/2*\cos(b*x + a) + 1/2) - 3*(\cos(b*x + a)^4 - 2*\cos(b*x + a)^2 + 1)*\log(-1/2*\cos(b*x + a) + 1/2) - 6*\cos(b*x + a))/(b*\cos(b*x + a)^4 - 2*b*\cos(b*x + a)^2 + b)$

Sympy [A]

time = 1.15, size = 92, normalized size = 1.67

$$\begin{cases} \frac{3 \log\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b} + \frac{\tan^4\left(\frac{a}{2} + \frac{bx}{2}\right)}{64b} - \frac{\tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)}{8b} + \frac{1}{8b \tan^2\left(\frac{a}{2} + \frac{bx}{2}\right)} - \frac{1}{64b \tan^4\left(\frac{a}{2} + \frac{bx}{2}\right)} & \text{for } b \neq 0 \\ \frac{x \cos^4(a)}{\sin^5(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**4/sin(b*x+a)**5,x)

[Out] Piecewise(((3*log(tan(a/2 + b*x/2)))/(8*b) + tan(a/2 + b*x/2)**4/(64*b) - tan(a/2 + b*x/2)**2/(8*b) + 1/(8*b*tan(a/2 + b*x/2)**2) - 1/(64*b*tan(a/2 + b*x/2)**4), Ne(b, 0)), (x*cos(a)**4/sin(a)**5, True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 139 vs. 2(49) = 98.

time = 4.51, size = 139, normalized size = 2.53

$$\frac{\left(\frac{8(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{18(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 1\right)(\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} - \frac{8(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 12 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4/sin(b*x+a)^5,x, algorithm="giac")

[Out] $-1/64*((8*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 18*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 1)*(\cos(b*x + a) + 1)^2/(\cos(b*x + a) - 1)^2 - 8*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - (\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 - 12*\log(\text{abs}(-\cos(b*x + a) + 1)/\text{abs}(\cos(b*x + a) + 1)))/b$

Mupad [B]

time = 0.49, size = 78, normalized size = 1.42

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{64b} - \frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8b} + \frac{3 \ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b} + \frac{\cot\left(\frac{a}{2} + \frac{bx}{2}\right)^4 \left(\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^2}{8} - \frac{1}{64}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^4/sin(a + b*x)^5,x)

[Out] tan(a/2 + (b*x)/2)^4/(64*b) - tan(a/2 + (b*x)/2)^2/(8*b) + (3*log(tan(a/2 + (b*x)/2)))/(8*b) + (cot(a/2 + (b*x)/2)^4*(tan(a/2 + (b*x)/2)^2/8 - 1/64))/

b

3.177 $\int \cot^3(a + bx) \csc^2(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\cot^4(a + bx)}{4b}$$

[Out] -1/4*cot(b*x+a)^4/b

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2687, 30}

$$-\frac{\cot^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^3*Csc[a + b*x]^2,x]

[Out] -1/4*Cot[a + b*x]^4/b

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2687

Int[sec[(e_.) + (f_.)*(x_)]^(m_)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])

Rubi steps

$$\begin{aligned} \int \cot^3(a + bx) \csc^2(a + bx) dx &= -\frac{\text{Subst}\left(\int x^3 dx, x, -\cot(a + bx)\right)}{b} \\ &= -\frac{\cot^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.00, size = 15, normalized size = 1.00

$$-\frac{\cot^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^3*Csc[a + b*x]^2,x]

[Out] -1/4*Cot[a + b*x]^4/b

Maple [A]

time = 0.03, size = 22, normalized size = 1.47

method	result	size
derivativdivides	$-\frac{\cos^4(bx+a)}{4 \sin(bx+a)^4 b}$	22
default	$-\frac{\cos^4(bx+a)}{4 \sin(bx+a)^4 b}$	22
risch	$-\frac{2(e^{6i(bx+a)}+e^{2i(bx+a)})}{b(e^{2i(bx+a)}-1)^4}$	38
norman	$-\frac{\frac{1}{64b} + \frac{\tan^2(\frac{bx}{2} + \frac{a}{2})}{16b} + \frac{\tan^6(\frac{bx}{2} + \frac{a}{2})}{16b} - \frac{\tan^8(\frac{bx}{2} + \frac{a}{2})}{64b}}{\tan^4(\frac{bx}{2} + \frac{a}{2})}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^3/sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] -1/4*cos(b*x+a)^4/sin(b*x+a)^4/b

Maxima [A]

time = 0.28, size = 25, normalized size = 1.67

$$\frac{2 \sin(bx + a)^2 - 1}{4 b \sin(bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^5,x, algorithm="maxima")

[Out] 1/4*(2*sin(b*x + a)^2 - 1)/(b*sin(b*x + a)^4)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 39 vs. 2(13) = 26.

time = 0.33, size = 39, normalized size = 2.60

$$-\frac{2 \cos(bx + a)^2 - 1}{4 (b \cos(bx + a)^4 - 2 b \cos(bx + a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^5,x, algorithm="fricas")

[Out] -1/4*(2*cos(b*x + a)^2 - 1)/(b*cos(b*x + a)^4 - 2*b*cos(b*x + a)^2 + b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 44 vs. $2(12) = 24$.

time = 0.61, size = 44, normalized size = 2.93

$$\begin{cases} \frac{1}{4b \sin^2(a+bx)} - \frac{\cos^2(a+bx)}{4b \sin^4(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos^3(a)}{\sin^5(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3/sin(b*x+a)**5,x)

[Out] Piecewise((1/(4*b*sin(a + b*x)**2) - cos(a + b*x)**2/(4*b*sin(a + b*x)**4), Ne(b, 0)), (x*cos(a)**3/sin(a)**5, True))

Giac [A]

time = 4.32, size = 25, normalized size = 1.67

$$\frac{2 \sin(bx + a)^2 - 1}{4 b \sin(bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3/sin(b*x+a)^5,x, algorithm="giac")

[Out] 1/4*(2*sin(b*x + a)^2 - 1)/(b*sin(b*x + a)^4)

Mupad [B]

time = 0.39, size = 25, normalized size = 1.67

$$-\frac{(\sin(a + bx)^2 - 1)^2}{4 b \sin(a + bx)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^3/sin(a + b*x)^5,x)

[Out] -(sin(a + b*x)^2 - 1)^2/(4*b*sin(a + b*x)^4)

3.178 $\int \cot^2(a + bx) \csc^3(a + bx) dx$

Optimal. Leaf size=55

$$\frac{\tanh^{-1}(\cos(a + bx))}{8b} + \frac{\cot(a + bx) \csc(a + bx)}{8b} - \frac{\cot(a + bx) \csc^3(a + bx)}{4b}$$

[Out] 1/8*arctanh(cos(b*x+a))/b+1/8*cot(b*x+a)*csc(b*x+a)/b-1/4*cot(b*x+a)*csc(b*x+a)^3/b

Rubi [A]

time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2691, 3853, 3855}

$$\frac{\tanh^{-1}(\cos(a + bx))}{8b} - \frac{\cot(a + bx) \csc^3(a + bx)}{4b} + \frac{\cot(a + bx) \csc(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]^2*Csc[a + b*x]^3,x]

[Out] ArcTanh[Cos[a + b*x]]/(8*b) + (Cot[a + b*x]*Csc[a + b*x])/(8*b) - (Cot[a + b*x]*Csc[a + b*x]^3)/(4*b)

Rule 2691

Int[((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_.)*((b_.)*tan[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[b*(a*Sec[e + f*x])^m*((b*Tan[e + f*x])^(n - 1)/(f*(m + n - 1))), x] - Dist[b^2*((n - 1)/(m + n - 1)), Int[(a*Sec[e + f*x])^m*(b*Tan[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && NeQ[m + n - 1, 0] && IntegerQ[2*m, 2*n]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n - 1)/(d*(n - 1))), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3855

Int[csc[(c_.) + (d_.)*(x_)], x_Symbol] :> Simp[-ArcTanh[Cos[c + d*x]]/d, x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \cot^2(a+bx) \csc^3(a+bx) dx &= -\frac{\cot(a+bx) \csc^3(a+bx)}{4b} - \frac{1}{4} \int \csc^3(a+bx) dx \\ &= \frac{\cot(a+bx) \csc(a+bx)}{8b} - \frac{\cot(a+bx) \csc^3(a+bx)}{4b} - \frac{1}{8} \int \csc(a+bx) dx \\ &= \frac{\tanh^{-1}(\cos(a+bx))}{8b} + \frac{\cot(a+bx) \csc(a+bx)}{8b} - \frac{\cot(a+bx) \csc^3(a+bx)}{4b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 113 vs. 2(55) = 110.

time = 0.03, size = 113, normalized size = 2.05

$$\frac{\csc^2\left(\frac{1}{2}(a+bx)\right)}{32b} - \frac{\csc^4\left(\frac{1}{2}(a+bx)\right)}{64b} + \frac{\log\left(\cos\left(\frac{1}{2}(a+bx)\right)\right)}{8b} - \frac{\log\left(\sin\left(\frac{1}{2}(a+bx)\right)\right)}{8b} - \frac{\sec^2\left(\frac{1}{2}(a+bx)\right)}{32b} + \frac{\sec^4\left(\frac{1}{2}(a+bx)\right)}{64b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]^2*Csc[a + b*x]^3,x]

[Out] Csc[(a + b*x)/2]^2/(32*b) - Csc[(a + b*x)/2]^4/(64*b) + Log[Cos[(a + b*x)/2]]/(8*b) - Log[Sin[(a + b*x)/2]]/(8*b) - Sec[(a + b*x)/2]^2/(32*b) + Sec[(a + b*x)/2]^4/(64*b)

Maple [A]

time = 0.05, size = 68, normalized size = 1.24

method	result	size
norman	$-\frac{1}{64b} + \frac{\tan^8\left(\frac{bx+a}{2}\right)}{64b} - \frac{\ln\left(\tan\left(\frac{bx+a}{2}\right)\right)}{8b}$	51
derivativedivides	$\frac{-\frac{\cos^3(bx+a)}{4 \sin(bx+a)^4} - \frac{\cos^3(bx+a)}{8 \sin(bx+a)^2} - \frac{\cos(bx+a)}{8} - \frac{\ln(\csc(bx+a) - \cot(bx+a))}{8}}{b}$	68
default	$\frac{-\frac{\cos^3(bx+a)}{4 \sin(bx+a)^4} - \frac{\cos^3(bx+a)}{8 \sin(bx+a)^2} - \frac{\cos(bx+a)}{8} - \frac{\ln(\csc(bx+a) - \cot(bx+a))}{8}}{b}$	68
risch	$-\frac{e^{7i(bx+a)} + 7e^{5i(bx+a)} + 7e^{3i(bx+a)} + e^{i(bx+a)}}{4b(e^{2i(bx+a)} - 1)^4} - \frac{\ln(e^{i(bx+a)} - 1)}{8b} + \frac{\ln(e^{i(bx+a)} + 1)}{8b}$	95

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2/sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/4*cos(b*x+a)^3/sin(b*x+a)^4-1/8*cos(b*x+a)^3/sin(b*x+a)^2-1/8*cos(b*x+a)-1/8*ln(csc(b*x+a)-cot(b*x+a)))

Maxima [A]

time = 0.28, size = 65, normalized size = 1.18

$$\frac{2(\cos(bx+a)^3 + \cos(bx+a))}{\cos(bx+a)^4 - 2\cos(bx+a)^2 + 1} - \log(\cos(bx+a) + 1) + \log(\cos(bx+a) - 1)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^5,x, algorithm="maxima")**[Out]** -1/16*(2*(cos(b*x + a)^3 + cos(b*x + a))/(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1) - log(cos(b*x + a) + 1) + log(cos(b*x + a) - 1))/b**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 111 vs. 2(49) = 98.

time = 0.37, size = 111, normalized size = 2.02

$$\frac{2\cos(bx+a)^3 - (\cos(bx+a)^4 - 2\cos(bx+a)^2 + 1)\log\left(\frac{1}{2}\cos(bx+a) + \frac{1}{2}\right) + (\cos(bx+a)^4 - 2\cos(bx+a)^2 + 1)\log\left(-\frac{1}{2}\cos(bx+a) + \frac{1}{2}\right) + 2\cos(bx+a)}{16(b\cos(bx+a)^4 - 2b\cos(bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^5,x, algorithm="fricas")**[Out]** -1/16*(2*cos(b*x + a)^3 - (cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*log(1/2*cos(b*x + a) + 1/2) + (cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*log(-1/2*cos(b*x + a) + 1/2) + 2*cos(b*x + a))/(b*cos(b*x + a)^4 - 2*b*cos(b*x + a)^2 + b)**Sympy [A]**

time = 1.06, size = 58, normalized size = 1.05

$$\begin{cases} -\frac{\log\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b} + \frac{\tan^4\left(\frac{a}{2} + \frac{bx}{2}\right)}{64b} - \frac{1}{64b\tan^4\left(\frac{a}{2} + \frac{bx}{2}\right)} & \text{for } b \neq 0 \\ \frac{x\cos^2(a)}{\sin^5(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2/sin(b*x+a)**5,x)**[Out]** Piecewise((-log(tan(a/2 + b*x/2))/(8*b) + tan(a/2 + b*x/2)**4/(64*b) - 1/(64*b*tan(a/2 + b*x/2)**4), Ne(b, 0)), (x*cos(a)**2/sin(a)**5, True))**Giac [A]**

time = 5.57, size = 98, normalized size = 1.78

$$\frac{\left(\frac{2(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 1\right)(\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 4\log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2/sin(b*x+a)^5,x, algorithm="giac")

[Out] $\frac{1}{64} * ((2 * (\cos(b*x + a) - 1)^2 / (\cos(b*x + a) + 1)^2 - 1) * (\cos(b*x + a) + 1)^2 / (\cos(b*x + a) - 1)^2 + (\cos(b*x + a) - 1)^2 / (\cos(b*x + a) + 1)^2 - 4 * \log(\text{abs}(-\cos(b*x + a) + 1) / \text{abs}(\cos(b*x + a) + 1))) / b$

Mupad [B]

time = 0.47, size = 48, normalized size = 0.87

$$\frac{\tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4}{64b} - \frac{1}{64b \tan\left(\frac{a}{2} + \frac{bx}{2}\right)^4} - \frac{\ln\left(\tan\left(\frac{a}{2} + \frac{bx}{2}\right)\right)}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2/sin(a + b*x)^5,x)

[Out] $\tan(a/2 + (b*x)/2)^4 / (64*b) - 1 / (64*b*\tan(a/2 + (b*x)/2)^4) - \log(\tan(a/2 + (b*x)/2)) / (8*b)$

3.179 $\int \cot(a + bx) \csc^4(a + bx) dx$

Optimal. Leaf size=15

$$-\frac{\csc^4(a + bx)}{4b}$$

[Out] -1/4*csc(b*x+a)^4/b

Rubi [A]

time = 0.01, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {2686, 30}

$$-\frac{\csc^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cot[a + b*x]*Csc[a + b*x]^4,x]

[Out] -1/4*Csc[a + b*x]^4/b

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2686

Int[((a_)*sec[(e_) + (f_)*(x_)]^(m_))*((b_)*tan[(e_) + (f_)*(x_)]^(n_)), x_Symbol] := Dist[a/f, Subst[Int[(a*x)^(m - 1)*(-1 + x^2)^((n - 1)/2), x], x, Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n + 1])

Rubi steps

$$\begin{aligned} \int \cot(a + bx) \csc^4(a + bx) dx &= -\frac{\text{Subst}\left(\int x^3 dx, x, \csc(a + bx)\right)}{b} \\ &= -\frac{\csc^4(a + bx)}{4b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$-\frac{\csc^4(a + bx)}{4b}$$

Antiderivative was successfully verified.

[In] Integrate[Cot[a + b*x]*Csc[a + b*x]^4,x]

[Out] -1/4*Csc[a + b*x]^4/b

Maple [A]

time = 0.03, size = 14, normalized size = 0.93

method	result	size
derivativdivides	$-\frac{1}{4 \sin(bx+a)^4 b}$	14
default	$-\frac{1}{4 \sin(bx+a)^4 b}$	14
risch	$-\frac{4 e^{4i(bx+a)}}{b(e^{2i(bx+a)}-1)^4}$	28
norman	$-\frac{\frac{1}{64b} - \frac{\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)}{16b} - \frac{\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)}{16b} - \frac{\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)}{64b}}{\tan\left(\frac{bx}{2} + \frac{a}{2}\right)^4}$	67

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)/sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] -1/4/sin(b*x+a)^4/b

Maxima [A]

time = 0.27, size = 13, normalized size = 0.87

$$-\frac{1}{4 b \sin (b x + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/4/(b*sin(b*x + a)^4)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 27 vs. 2(13) = 26.

time = 0.36, size = 27, normalized size = 1.80

$$-\frac{1}{4 (b \cos (b x + a)^4 - 2 b \cos (b x + a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)/sin(b*x+a)^5,x, algorithm="fricas")

[Out] -1/4/(b*cos(b*x + a)^4 - 2*b*cos(b*x + a)^2 + b)

Sympy [A]

time = 0.54, size = 24, normalized size = 1.60

$$\begin{cases} -\frac{1}{4b \sin^4(a+bx)} & \text{for } b \neq 0 \\ \frac{x \cos(a)}{\sin^5(a)} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)/sin(b*x+a)**5,x)
```

```
[Out] Piecewise((-1/(4*b*sin(a + b*x)**4), Ne(b, 0)), (x*cos(a)/sin(a)**5, True))
```

Giac [A]

time = 3.06, size = 13, normalized size = 0.87

$$-\frac{1}{4b \sin(bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)/sin(b*x+a)^5,x, algorithm="giac")
```

```
[Out] -1/4/(b*sin(b*x + a)^4)
```

Mupad [B]

time = 0.41, size = 23, normalized size = 1.53

$$\frac{\cot(a + bx)^2 (\cot(a + bx)^2 + 2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)/sin(a + b*x)^5,x)
```

```
[Out] -(cot(a + b*x)^2*(cot(a + b*x)^2 + 2))/(4*b)
```

3.180 $\int \csc^5(a + bx) \sec(a + bx) dx$

Optimal. Leaf size=40

$$-\frac{\cot^2(a + bx)}{b} - \frac{\cot^4(a + bx)}{4b} + \frac{\log(\tan(a + bx))}{b}$$

[Out] $-\cot(b*x+a)^2/b-1/4*\cot(b*x+a)^4/b+\ln(\tan(b*x+a))/b$

Rubi [A]

time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2700, 272, 45}

$$-\frac{\cot^4(a + bx)}{4b} - \frac{\cot^2(a + bx)}{b} + \frac{\log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^5*Sec[a + b*x],x]

[Out] $-(\text{Cot}[a + b*x]^2/b) - \text{Cot}[a + b*x]^4/(4*b) + \text{Log}[\text{Tan}[a + b*x]]/b$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LtQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^((m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned}
\int \csc^5(a+bx) \sec(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{x^5} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x^2)^2}{x^3} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(\frac{1}{x^3} + \frac{2}{x^2} + \frac{1}{x}\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= -\frac{\cot^2(a+bx)}{b} - \frac{\cot^4(a+bx)}{4b} + \frac{\log(\tan(a+bx))}{b}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 44, normalized size = 1.10

$$-\frac{2 \csc^2(a+bx) + \csc^4(a+bx) + 4 \log(\cos(a+bx)) - 4 \log(\sin(a+bx))}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^5*Sec[a + b*x], x]``[Out] -1/4*(2*Csc[a + b*x]^2 + Csc[a + b*x]^4 + 4*Log[Cos[a + b*x]] - 4*Log[Sin[a + b*x]])/b`**Maple [A]**

time = 0.05, size = 33, normalized size = 0.82

method	result
derivativedivides	$-\frac{\frac{1}{4 \sin^4(bx+a)} - \frac{1}{2 \sin^2(bx+a)^2} + \ln(\tan(bx+a))}{b}$
default	$-\frac{\frac{1}{4 \sin^4(bx+a)} - \frac{1}{2 \sin^2(bx+a)^2} + \ln(\tan(bx+a))}{b}$
risch	$\frac{2e^{6i(bx+a)} - 8e^{4i(bx+a)} + 2e^{2i(bx+a)}}{b(e^{2i(bx+a)} - 1)^4} + \frac{\ln(e^{2i(bx+a)} - 1)}{b} - \frac{\ln(e^{2i(bx+a)} + 1)}{b}$
norman	$-\frac{\frac{1}{64b} - \frac{3 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{16b} - \frac{3 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{16b} - \frac{\tan^8\left(\frac{bx}{2} + \frac{a}{2}\right)}{64b}}{\tan^4\left(\frac{bx}{2} + \frac{a}{2}\right)} + \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)}{b} - \frac{\ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)/sin(b*x+a)^5,x,method=_RETURNVERBOSE)``[Out] 1/b*(-1/4/sin(b*x+a)^4-1/2/sin(b*x+a)^2+ln(tan(b*x+a)))`

Maxima [A]

time = 0.27, size = 51, normalized size = 1.28

$$\frac{\frac{2 \sin(bx+a)^2+1}{\sin(bx+a)^4} + 2 \log(\sin(bx+a)^2 - 1) - 2 \log(\sin(bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^5,x, algorithm="maxima")

[Out] -1/4*((2*sin(b*x + a)^2 + 1)/sin(b*x + a)^4 + 2*log(sin(b*x + a)^2 - 1) - 2*log(sin(b*x + a)^2))/b

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 105 vs. 2(38) = 76.

time = 0.37, size = 105, normalized size = 2.62

$$\frac{2 \cos(bx+a)^2 - 2(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log(\cos(bx+a)^2) + 2(\cos(bx+a)^4 - 2 \cos(bx+a)^2 + 1) \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) - 3}{4(b \cos(bx+a)^4 - 2b \cos(bx+a)^2 + b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^5,x, algorithm="fricas")

[Out] 1/4*(2*cos(b*x + a)^2 - 2*(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*log(cos(b*x + a)^2) + 2*(cos(b*x + a)^4 - 2*cos(b*x + a)^2 + 1)*log(-1/4*cos(b*x + a)^2 + 1/4) - 3)/(b*cos(b*x + a)^4 - 2*b*cos(b*x + a)^2 + b)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec(a + bx)}{\sin^5(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)**5,x)

[Out] Integral(sec(a + b*x)/sin(a + b*x)**5, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 165 vs. 2(38) = 76.

time = 3.49, size = 165, normalized size = 4.12

$$\frac{\left(\frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{48(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 1\right)(\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} + \frac{12(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 32 \log\left(\frac{1-\cos(bx+a)+1}{|\cos(bx+a)+1|}\right) - 64 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)/sin(b*x+a)^5,x, algorithm="giac")

[Out] $\frac{1}{64} * \left(\frac{12 * (\cos(b*x + a) - 1)}{(\cos(b*x + a) + 1)} - 48 * (\cos(b*x + a) - 1)^2 / (\cos(b*x + a) + 1)^2 - 1 \right) * (\cos(b*x + a) + 1)^2 / (\cos(b*x + a) - 1)^2 + 12 * (\cos(b*x + a) - 1) / (\cos(b*x + a) + 1) - (\cos(b*x + a) - 1)^2 / (\cos(b*x + a) + 1)^2 + 32 * \log(\text{abs}(-\cos(b*x + a) + 1) / \text{abs}(\cos(b*x + a) + 1)) - 64 * \log(\text{abs}(-\cos(b*x + a) - 1) / (\cos(b*x + a) + 1 - 1))) / b$

Mupad [B]

time = 0.44, size = 79, normalized size = 1.98

$$\frac{\ln\left(\frac{\cos(2a+2bx)}{2} - \frac{1}{2}\right)}{2b} - \frac{\ln(\cos(a+bx))}{b} - \frac{\frac{\cos(2a+2bx)}{4} - \frac{1}{2}}{b \left(\cos(2a+2bx) - \left(\frac{\cos(2a+2bx)}{2} + \frac{1}{2} \right)^2 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(cos(a + b*x)*sin(a + b*x)^5),x)`

[Out] $\log(\cos(2*a + 2*b*x)/2 - 1/2)/(2*b) - \log(\cos(a + b*x))/b - (\cos(2*a + 2*b*x)/4 - 1/2)/(b*(\cos(2*a + 2*b*x) - (\cos(2*a + 2*b*x)/2 + 1/2)^2))$

3.181 $\int \csc^5(a + bx) \sec^2(a + bx) dx$

Optimal. Leaf size=70

$$-\frac{15 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{15 \sec(a + bx)}{8b} - \frac{5 \csc^2(a + bx) \sec(a + bx)}{8b} - \frac{\csc^4(a + bx) \sec(a + bx)}{4b}$$

[Out] $-15/8*\operatorname{arctanh}(\cos(b*x+a))/b+15/8*\sec(b*x+a)/b-5/8*\csc(b*x+a)^2*\sec(b*x+a)/b-1/4*\csc(b*x+a)^4*\sec(b*x+a)/b$

Rubi [A]

time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2702, 294, 327, 213}

$$\frac{15 \sec(a + bx)}{8b} - \frac{15 \tanh^{-1}(\cos(a + bx))}{8b} - \frac{\csc^4(a + bx) \sec(a + bx)}{4b} - \frac{5 \csc^2(a + bx) \sec(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^5*Sec[a + b*x]^2,x]`

[Out] $(-15*\operatorname{ArcTanh}[\operatorname{Cos}[a + b*x]])/(8*b) + (15*\operatorname{Sec}[a + b*x])/(8*b) - (5*\operatorname{Csc}[a + b*x]^2*\operatorname{Sec}[a + b*x])/(8*b) - (\operatorname{Csc}[a + b*x]^4*\operatorname{Sec}[a + b*x])/(4*b)$

Rule 213

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])`

Rule 294

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 327

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_Symbol]
:> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^5(a + bx) \sec^2(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^6}{(-1+x^2)^3} dx, x, \sec(a + bx)\right)}{b} \\ &= -\frac{\csc^4(a + bx) \sec(a + bx)}{4b} + \frac{5\text{Subst}\left(\int \frac{x^4}{(-1+x^2)^2} dx, x, \sec(a + bx)\right)}{4b} \\ &= -\frac{5 \csc^2(a + bx) \sec(a + bx)}{8b} - \frac{\csc^4(a + bx) \sec(a + bx)}{4b} + \frac{15\text{Subst}\left(\int \frac{x^2}{-1+x^2} dx, x, \sec(a + bx)\right)}{4b} \\ &= \frac{15 \sec(a + bx)}{8b} - \frac{5 \csc^2(a + bx) \sec(a + bx)}{8b} - \frac{\csc^4(a + bx) \sec(a + bx)}{4b} + \frac{15 \tanh^{-1}(\cos(a + bx))}{8b} \\ &= -\frac{15 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{15 \sec(a + bx)}{8b} - \frac{5 \csc^2(a + bx) \sec(a + bx)}{8b} \end{aligned}$$

Mathematica [A]

time = 3.00, size = 129, normalized size = 1.84

$$\frac{14 \csc^2\left(\frac{1}{2}(a + bx)\right) + \csc^4\left(\frac{1}{2}(a + bx)\right) + \frac{\sec^2\left(\frac{1}{2}(a + bx)\right)(78 + \cos(a + bx)(-8(8 + 15 \log(\cos\left(\frac{1}{2}(a + bx)\right)) - 15 \log(\sin\left(\frac{1}{2}(a + bx)\right))) + \sec^4\left(\frac{1}{2}(a + bx)\right) - 14 \tan^2\left(\frac{1}{2}(a + bx)\right))}{-1 + \tan^2\left(\frac{1}{2}(a + bx)\right)}}{64b}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^5*Sec[a + b*x]^2,x]

[Out] -1/64*(14*Csc[(a + b*x)/2]^2 + Csc[(a + b*x)/2]^4 + (Sec[(a + b*x)/2]^2*(78 + Cos[a + b*x]*(-8*(8 + 15*Log[Cos[(a + b*x)/2]] - 15*Log[Sin[(a + b*x)/2]]) + Sec[(a + b*x)/2]^4 - 14*Tan[(a + b*x)/2]^2))/(-1 + Tan[(a + b*x)/2]^2))/b

Maple [A]

time = 0.06, size = 70, normalized size = 1.00

method	result	size
derivativedivides	$-\frac{1}{4 \sin(bx+a)^4 \cos(bx+a)} - \frac{5}{8 \sin(bx+a)^2 \cos(bx+a)} + \frac{15}{8 \cos(bx+a)} + \frac{15 \ln(\csc(bx+a) - \cot(bx+a))}{8}$	70
default	$-\frac{1}{4 \sin(bx+a)^4 \cos(bx+a)} - \frac{5}{8 \sin(bx+a)^2 \cos(bx+a)} + \frac{15}{8 \cos(bx+a)} + \frac{15 \ln(\csc(bx+a) - \cot(bx+a))}{8}$	70

norman	$\frac{\frac{1}{64b} + \frac{15 \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{64b} + \frac{15 \left(\tan^8 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{64b} + \frac{\tan^{10} \left(\frac{bx}{2} + \frac{a}{2} \right)}{64b} - \frac{5 \left(\tan^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{2b}}{\left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) - 1 \right) \tan \left(\frac{bx}{2} + \frac{a}{2} \right)^4} + \frac{15 \ln \left(\tan \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{8b}$	114
risch	$\frac{15 e^{9i(bx+a)} - 40 e^{7i(bx+a)} + 18 e^{5i(bx+a)} - 40 e^{3i(bx+a)} + 15 e^{i(bx+a)}}{4b(e^{2i(bx+a)} - 1)^4 (e^{2i(bx+a)} + 1)} - \frac{15 \ln(e^{i(bx+a)} + 1)}{8b} + \frac{15 \ln(e^{i(bx+a)} - 1)}{8b}$	123

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sec(b*x+a)^2/sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{b} * (-1/4 / \sin(b*x+a)^4 / \cos(b*x+a) - 5/8 / \sin(b*x+a)^2 / \cos(b*x+a) + 15/8 / \cos(b*x+a) + 15/8 * \ln(\csc(b*x+a) - \cot(b*x+a)))$

Maxima [A]

time = 0.28, size = 79, normalized size = 1.13

$$\frac{2 \left(15 \cos(bx+a)^4 - 25 \cos(bx+a)^2 + 8 \right)}{\cos(bx+a)^5 - 2 \cos(bx+a)^3 + \cos(bx+a)} - \frac{15 \log(\cos(bx+a) + 1) + 15 \log(\cos(bx+a) - 1)}{16b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a)^5,x, algorithm="maxima")`

[Out] $\frac{1}{16} * (2 * (15 * \cos(b*x + a)^4 - 25 * \cos(b*x + a)^2 + 8) / (\cos(b*x + a)^5 - 2 * \cos(b*x + a)^3 + \cos(b*x + a)) - 15 * \log(\cos(b*x + a) + 1) + 15 * \log(\cos(b*x + a) - 1)) / b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 132 vs. $2(62) = 124$.

time = 0.37, size = 132, normalized size = 1.89

$$\frac{30 \cos(bx+a)^4 - 50 \cos(bx+a)^2 - 15 (\cos(bx+a)^5 - 2 \cos(bx+a)^3 + \cos(bx+a)) \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 15 (\cos(bx+a)^5 - 2 \cos(bx+a)^3 + \cos(bx+a)) \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 16}{16 (b \cos(bx+a)^5 - 2 b \cos(bx+a)^3 + b \cos(bx+a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)^2/sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $\frac{1}{16} * (30 * \cos(b*x + a)^4 - 50 * \cos(b*x + a)^2 - 15 * (\cos(b*x + a)^5 - 2 * \cos(b*x + a)^3 + \cos(b*x + a)) * \log(1/2 * \cos(b*x + a) + 1/2) + 15 * (\cos(b*x + a)^5 - 2 * \cos(b*x + a)^3 + \cos(b*x + a)) * \log(-1/2 * \cos(b*x + a) + 1/2) + 16) / (b * \cos(b*x + a)^5 - 2 * b * \cos(b*x + a)^3 + b * \cos(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(a + bx)}{\sin^5(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2/sin(b*x+a)**5,x)

[Out] Integral(sec(a + b*x)**2/sin(a + b*x)**5, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(62) = 124.

time = 4.06, size = 163, normalized size = 2.33

$$\frac{\left(\frac{16(\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{90(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 1\right)(\cos(bx+a)+1)^2}{(\cos(bx+a)-1)^2} - \frac{16(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{128}{\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 1} + 60 \log\left(\frac{|-\cos(bx+a)+1|}{|\cos(bx+a)+1|}\right)$$

$64b$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2/sin(b*x+a)^5,x, algorithm="giac")

[Out] 1/64*((16*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 90*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 1)*(cos(b*x + a) + 1)^2/(cos(b*x + a) - 1)^2 - 16*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + (cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 128/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1) + 60*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1))))/b

Mupad [B]

time = 0.39, size = 66, normalized size = 0.94

$$\frac{\frac{15 \cos(a+bx)^4}{8} - \frac{25 \cos(a+bx)^2}{8} + 1}{b (\cos(a+bx)^5 - 2 \cos(a+bx)^3 + \cos(a+bx))} - \frac{15 \operatorname{atanh}(\cos(a+bx))}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^2*sin(a + b*x)^5),x)

[Out] ((15*cos(a + b*x)^4)/8 - (25*cos(a + b*x)^2)/8 + 1)/(b*(cos(a + b*x) - 2*cos(a + b*x)^3 + cos(a + b*x)^5)) - (15*atanh(cos(a + b*x)))/(8*b)

3.182 $\int \csc^5(a + bx) \sec^3(a + bx) dx$

Optimal. Leaf size=58

$$-\frac{3 \cot^2(a + bx)}{2b} - \frac{\cot^4(a + bx)}{4b} + \frac{3 \log(\tan(a + bx))}{b} + \frac{\tan^2(a + bx)}{2b}$$

[Out] $-3/2*\cot(b*x+a)^2/b-1/4*\cot(b*x+a)^4/b+3*\ln(\tan(b*x+a))/b+1/2*\tan(b*x+a)^2/b$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2700, 272, 45}

$$\frac{\tan^2(a + bx)}{2b} - \frac{\cot^4(a + bx)}{4b} - \frac{3 \cot^2(a + bx)}{2b} + \frac{3 \log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^5*Sec[a + b*x]^3,x]

[Out] $(-3*\cot[a + b*x]^2)/(2*b) - \cot[a + b*x]^4/(4*b) + (3*\log[\tan[a + b*x]])/b + \tan[a + b*x]^2/(2*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] :> Dist[1/f, Subst[Int[(1 + x^2)^(m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned}
\int \csc^5(a+bx) \sec^3(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^3}{x^5} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x)^3}{x^3} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(1 + \frac{1}{x^3} + \frac{3}{x^2} + \frac{3}{x}\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= -\frac{3 \cot^2(a+bx)}{2b} - \frac{\cot^4(a+bx)}{4b} + \frac{3 \log(\tan(a+bx))}{b} + \frac{\tan^2(a+bx)}{2b}
\end{aligned}$$

Mathematica [A]

time = 0.24, size = 54, normalized size = 0.93

$$\frac{4 \csc^2(a+bx) + \csc^4(a+bx) + 12 \log(\cos(a+bx)) - 12 \log(\sin(a+bx)) - 2 \sec^2(a+bx)}{4b}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^5*Sec[a + b*x]^3,x]`

```
[Out] -1/4*(4*Csc[a + b*x]^2 + Csc[a + b*x]^4 + 12*Log[Cos[a + b*x]] - 12*Log[Sin[a + b*x]] - 2*Sec[a + b*x]^2)/b
```

Maple [A]

time = 0.08, size = 61, normalized size = 1.05

method	result
derivativedivides	$\frac{-\frac{1}{4 \sin^4(bx+a)} \cos^3(bx+a)^2 + \frac{3}{4 \sin^2(bx+a)} \frac{\cos^3(bx+a)^2}{\cos(bx+a)^2} - \frac{3}{2 \sin^2(bx+a)^2} + 3 \ln(\tan(bx+a))}{b}$
default	$\frac{-\frac{1}{4 \sin^4(bx+a)} \cos^3(bx+a)^2 + \frac{3}{4 \sin^2(bx+a)} \frac{\cos^3(bx+a)^2}{\cos(bx+a)^2} - \frac{3}{2 \sin^2(bx+a)^2} + 3 \ln(\tan(bx+a))}{b}$
risch	$\frac{6 e^{10i(bx+a)} - 12 e^{8i(bx+a)} - 4 e^{6i(bx+a)} - 12 e^{4i(bx+a)} + 6 e^{2i(bx+a)}}{b(e^{2i(bx+a)} - 1)^4 (e^{2i(bx+a)} + 1)^2} - \frac{3 \ln(e^{2i(bx+a)} + 1)}{b} + \frac{3 \ln(e^{2i(bx+a)} - 1)}{b}$
norman	$\frac{-\frac{1}{64b} - \frac{9 \left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{32b} - \frac{9 \left(\tan^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{32b} - \frac{\tan^{12}\left(\frac{bx}{2} + \frac{a}{2}\right)}{64b} + \frac{83 \left(\tan^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{32b}}{\left(\tan^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right)^2 \tan\left(\frac{bx}{2} + \frac{a}{2}\right)^4} + \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b} - \frac{3 \ln\left(\tan\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^3/sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(-1/4/sin(b*x+a)^4/cos(b*x+a)^2+3/4/sin(b*x+a)^2/cos(b*x+a)^2-3/2/sin(b*x+a)^2+3*ln(tan(b*x+a)))
```

Maxima [A]

time = 0.27, size = 74, normalized size = 1.28

$$\frac{6 \sin(bx+a)^4 - 3 \sin(bx+a)^2 - 1}{\sin(bx+a)^6 - \sin(bx+a)^4} + 6 \log(\sin(bx+a)^2 - 1) - 6 \log(\sin(bx+a)^2)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3/sin(b*x+a)^5,x, algorithm="maxima")``[Out] -1/4*((6*sin(b*x + a)^4 - 3*sin(b*x + a)^2 - 1)/(sin(b*x + a)^6 - sin(b*x + a)^4) + 6*log(sin(b*x + a)^2 - 1) - 6*log(sin(b*x + a)^2))/b`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 138 vs. 2(52) = 104.

time = 0.36, size = 138, normalized size = 2.38

$$\frac{6 \cos(bx+a)^4 - 9 \cos(bx+a)^2 - 6(\cos(bx+a)^6 - 2 \cos(bx+a)^4 + \cos(bx+a)^2) \log(\cos(bx+a)^2) + 6(\cos(bx+a)^6 - 2 \cos(bx+a)^4 + \cos(bx+a)^2) \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) + 2}{4(b \cos(bx+a)^6 - 2b \cos(bx+a)^4 + b \cos(bx+a)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)^3/sin(b*x+a)^5,x, algorithm="fricas")``[Out] 1/4*(6*cos(b*x + a)^4 - 9*cos(b*x + a)^2 - 6*(cos(b*x + a)^6 - 2*cos(b*x + a)^4 + cos(b*x + a)^2)*log(cos(b*x + a)^2) + 6*(cos(b*x + a)^6 - 2*cos(b*x + a)^4 + cos(b*x + a)^2)*log(-1/4*cos(b*x + a)^2 + 1/4) + 2)/(b*cos(b*x + a)^6 - 2*b*cos(b*x + a)^4 + b*cos(b*x + a)^2)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^3(a + bx)}{\sin^5(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sec(b*x+a)**3/sin(b*x+a)**5,x)``[Out] Integral(sec(a + b*x)**3/sin(a + b*x)**5, x)`**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 232 vs. 2(52) = 104.

time = 3.94, size = 232, normalized size = 4.00

$$\frac{20 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} - \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{18 \frac{\cos(bx+a)-1}{\cos(bx+a)+1} + 111 \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 36 \frac{(\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + 72 \frac{(\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} - 1}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} + \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2}\right)^2} + 96 \log\left(\frac{1-\cos(bx+a)+1}{|\cos(bx+a)+1|}\right) - 192 \log\left(\left|-\frac{\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right|\right)}{64b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3/sin(b*x+a)^5,x, algorithm="giac")

[Out] $\frac{1}{64} \cdot \frac{20 \cdot (\cos(bx + a) - 1)}{(\cos(bx + a) + 1)} - \frac{(\cos(bx + a) - 1)^2}{(\cos(bx + a) + 1)^2} + \frac{18 \cdot (\cos(bx + a) - 1)}{(\cos(bx + a) + 1)} + \frac{111 \cdot (\cos(bx + a) - 1)^2}{(\cos(bx + a) + 1)^2} + \frac{36 \cdot (\cos(bx + a) - 1)^3}{(\cos(bx + a) + 1)^3} + \frac{72 \cdot (\cos(bx + a) - 1)^4}{(\cos(bx + a) + 1)^4} - \frac{1}{(\cos(bx + a) - 1)} \cdot \frac{(\cos(bx + a) - 1)^2}{(\cos(bx + a) + 1)^2} + \frac{96 \cdot \log\left(\frac{\text{abs}(-\cos(bx + a) + 1)}{\text{abs}(\cos(bx + a) + 1)}\right) - 192 \cdot \log\left(\frac{\text{abs}(-(\cos(bx + a) - 1))}{(\cos(bx + a) + 1) - 1}\right)}{b}$

Mupad [B]

time = 0.47, size = 82, normalized size = 1.41

$$\frac{3 \ln(\sin(a + bx)^2)}{2b} - \frac{3 \ln(\cos(a + bx))}{b} + \frac{\frac{3 \cos(a + bx)^4}{2} - \frac{9 \cos(a + bx)^2}{4} + \frac{1}{2}}{b (\cos(a + bx)^6 - 2 \cos(a + bx)^4 + \cos(a + bx)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^3*sin(a + b*x)^5),x)

[Out] $\frac{3 \cdot \log(\sin(a + bx)^2)}{2b} - \frac{3 \cdot \log(\cos(a + bx))}{b} + \left(\frac{3 \cdot \cos(a + bx)^4}{2} - \frac{9 \cdot \cos(a + bx)^2}{4} + \frac{1}{2} \right) / (b \cdot (\cos(a + bx)^2 - 2 \cdot \cos(a + bx)^4 + \cos(a + bx)^6))$

3.183 $\int \csc^5(a + bx) \sec^4(a + bx) dx$

Optimal. Leaf size=89

$$-\frac{35 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{35 \sec(a + bx)}{8b} + \frac{35 \sec^3(a + bx)}{24b} - \frac{7 \csc^2(a + bx) \sec^3(a + bx)}{8b} - \frac{\csc^4(a + bx) \sec^3(a + bx)}{4b}$$

[Out] $-35/8*\arctanh(\cos(b*x+a))/b+35/8*\sec(b*x+a)/b+35/24*\sec(b*x+a)^3/b-7/8*\csc(b*x+a)^2*\sec(b*x+a)^3/b-1/4*\csc(b*x+a)^4*\sec(b*x+a)^3/b$

Rubi [A]

time = 0.03, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.235$, Rules used = {2702, 294, 308, 213}

$$\frac{35 \sec^3(a + bx)}{24b} + \frac{35 \sec(a + bx)}{8b} - \frac{35 \tanh^{-1}(\cos(a + bx))}{8b} - \frac{\csc^4(a + bx) \sec^3(a + bx)}{4b} - \frac{7 \csc^2(a + bx) \sec^3(a + bx)}{8b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^5*Sec[a + b*x]^4,x]

[Out] $(-35*\text{ArcTanh}[\text{Cos}[a + b*x]])/(8*b) + (35*\text{Sec}[a + b*x])/(8*b) + (35*\text{Sec}[a + b*x]^3)/(24*b) - (7*\text{Csc}[a + b*x]^2*\text{Sec}[a + b*x]^3)/(8*b) - (\text{Csc}[a + b*x]^4*\text{Sec}[a + b*x]^3)/(4*b)$

Rule 213

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[b, 2])^(-1))*ArcTanh[Rt[b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (LtQ[a, 0] || GtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 308

Int[(x_)^(m_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Int[PolynomialDivide[x^m, a + b*x^n, x], x] /; FreeQ[{a, b}, x] && IGtQ[m, 0] && IGtQ[n, 0] && GtQ[m, 2*n - 1]

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_Symbol]
:> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^5(a + bx) \sec^4(a + bx) dx &= \frac{\text{Subst}\left(\int \frac{x^8}{(-1+x^2)^3} dx, x, \sec(a + bx)\right)}{b} \\ &= -\frac{\csc^4(a + bx) \sec^3(a + bx)}{4b} + \frac{7\text{Subst}\left(\int \frac{x^6}{(-1+x^2)^2} dx, x, \sec(a + bx)\right)}{4b} \\ &= -\frac{7 \csc^2(a + bx) \sec^3(a + bx)}{8b} - \frac{\csc^4(a + bx) \sec^3(a + bx)}{4b} + \frac{35\text{Subst}\left(\int \frac{x^4}{-1+x^2} dx, x, \sec(a + bx)\right)}{4b} \\ &= -\frac{7 \csc^2(a + bx) \sec^3(a + bx)}{8b} - \frac{\csc^4(a + bx) \sec^3(a + bx)}{4b} + \frac{35\text{Subst}\left(\int \frac{x^2}{1-x^2} dx, x, \sec(a + bx)\right)}{4b} \\ &= \frac{35 \sec(a + bx)}{8b} + \frac{35 \sec^3(a + bx)}{24b} - \frac{7 \csc^2(a + bx) \sec^3(a + bx)}{8b} - \frac{\csc^4(a + bx)}{4b} \\ &= -\frac{35 \tanh^{-1}(\cos(a + bx))}{8b} + \frac{35 \sec(a + bx)}{8b} + \frac{35 \sec^3(a + bx)}{24b} - \frac{7 \csc^2(a + bx) \sec^3(a + bx)}{8b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 268 vs. 2(89) = 178.

time = 0.36, size = 268, normalized size = 3.01

3987(a + b) (-284 + 658cos(2a + b*x) - 228cos(3a + b*x) + 140cos(4a + b*x) - 76cos(5a + b*x) - 210cos(6a + b*x) + 76cos(7a + b*x) - 315cos(3a + b*x)log(cos((a + b*x)/2)) - 105cos(5a + b*x)log(cos((a + b*x)/2)) + 35cos(a + b*x)(76 + 105log(cos((a + b*x)/2)) - 105log(sin((a + b*x)/2))) + 315cos(3a + b*x)log(sin((a + b*x)/2)) + 105cos(5a + b*x)log(sin((a + b*x)/2)) - 105cos(7a + b*x)log(sin((a + b*x)/2)))/((b*(Csc((a + b*x)/2)^2 - Sec((a + b*x)/2)^2)^3)

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^5*Sec[a + b*x]^4,x]

[Out] -1/24*(Csc[a + b*x]^10*(-204 + 658*Cos[2*(a + b*x)] - 228*Cos[3*(a + b*x)] + 140*Cos[4*(a + b*x)] - 76*Cos[5*(a + b*x)] - 210*Cos[6*(a + b*x)] + 76*Cos[7*(a + b*x)] - 315*Cos[3*(a + b*x)]*Log[Cos[(a + b*x)/2]] - 105*Cos[5*(a + b*x)]*Log[Cos[(a + b*x)/2]] + 105*Cos[7*(a + b*x)]*Log[Cos[(a + b*x)/2]] + 3*Cos[a + b*x]*(76 + 105*Log[Cos[(a + b*x)/2]] - 105*Log[Sin[(a + b*x)/2]]) + 315*Cos[3*(a + b*x)]*Log[Sin[(a + b*x)/2]] + 105*Cos[5*(a + b*x)]*Log[Sin[(a + b*x)/2]] - 105*Cos[7*(a + b*x)]*Log[Sin[(a + b*x)/2]]))/(b*(Csc[(a + b*x)/2]^2 - Sec[(a + b*x)/2]^2)^3)

Maple [A]

time = 0.09, size = 88, normalized size = 0.99

method	result
derivativedivides	$-\frac{1}{4 \sin^4(bx+a) \cos^3(bx+a)} + \frac{7}{12 \sin^2(bx+a) \cos^3(bx+a)} - \frac{35}{24 \sin(bx+a)^2 \cos(bx+a)} + \frac{35}{8 \cos(bx+a)} + \frac{35 \ln(\csc(bx+a) - \cot(bx+a))}{8}$
default	$-\frac{1}{4 \sin^4(bx+a) \cos^3(bx+a)} + \frac{7}{12 \sin^2(bx+a) \cos^3(bx+a)} - \frac{35}{24 \sin(bx+a)^2 \cos(bx+a)} + \frac{35}{8 \cos(bx+a)} + \frac{35 \ln(\csc(bx+a) - \cot(bx+a))}{8}$
risch	$\frac{105 e^{13i(bx+a)} - 70 e^{11i(bx+a)} - 329 e^{9i(bx+a)} + 204 e^{7i(bx+a)} - 329 e^{5i(bx+a)} - 70 e^{3i(bx+a)} + 105 e^{i(bx+a)}}{12b(e^{2i(bx+a)} - 1)^4 (e^{2i(bx+a)} + 1)^3} - \frac{35 \ln(e^{i(bx+a)})}{8b}$
norman	$\frac{\frac{1}{64b} + \frac{21 \left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{64b} + \frac{21 \left(\tan^{12} \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{64b} + \frac{\tan^{14} \left(\frac{bx}{2} + \frac{a}{2} \right)}{64b} - \frac{21 \left(\tan^8 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{2b} + \frac{511 \left(\tan^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{32b} - \frac{847 \left(\tan^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{96b}}{\left(\tan^2 \left(\frac{bx}{2} + \frac{a}{2} \right) - 1 \right)^3 \tan \left(\frac{bx}{2} + \frac{a}{2} \right)^4}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(b*x+a)^4/sin(b*x+a)^5,x,method=_RETURNVERBOSE)
```

```
[Out] 1/b*(-1/4/sin(b*x+a)^4/cos(b*x+a)^3+7/12/sin(b*x+a)^2/cos(b*x+a)^3-35/24/sin(b*x+a)^2/cos(b*x+a)+35/8/cos(b*x+a)+35/8*ln(csc(b*x+a)-cot(b*x+a)))
```

Maxima [A]

time = 0.29, size = 91, normalized size = 1.02

$$\frac{2 \left(105 \cos^6(bx+a) - 175 \cos^4(bx+a) + 56 \cos^2(bx+a) + 8 \right)}{\cos^7(bx+a) - 2 \cos^5(bx+a) + \cos^3(bx+a)} - 105 \log(\cos(bx+a) + 1) + 105 \log(\cos(bx+a) - 1)$$

48b

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)^4/sin(b*x+a)^5,x, algorithm="maxima")
```

```
[Out] 1/48*(2*(105*cos(b*x + a)^6 - 175*cos(b*x + a)^4 + 56*cos(b*x + a)^2 + 8)/(cos(b*x + a)^7 - 2*cos(b*x + a)^5 + cos(b*x + a)^3) - 105*log(cos(b*x + a) + 1) + 105*log(cos(b*x + a) - 1))/b
```

Fricas [A]

time = 0.35, size = 148, normalized size = 1.66

$$\frac{210 \cos^6(bx+a) - 350 \cos^4(bx+a) + 112 \cos^2(bx+a) - 105 (\cos^7(bx+a) - 2 \cos^5(bx+a) + \cos^3(bx+a)) \log\left(\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 105 (\cos^7(bx+a) - 2 \cos^5(bx+a) + \cos^3(bx+a)) \log\left(-\frac{1}{2} \cos(bx+a) + \frac{1}{2}\right) + 16}{48 (b \cos^7(bx+a) - 2b \cos^5(bx+a) + b \cos^3(bx+a))}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(b*x+a)^4/sin(b*x+a)^5,x, algorithm="fricas")
```

```
[Out] 1/48*(210*cos(b*x + a)^6 - 350*cos(b*x + a)^4 + 112*cos(b*x + a)^2 - 105*(cos(b*x + a)^7 - 2*cos(b*x + a)^5 + cos(b*x + a)^3)*log(1/2*cos(b*x + a) + 1/2) + 105*(cos(b*x + a)^7 - 2*cos(b*x + a)^5 + cos(b*x + a)^3)*log(-1/2*cos(b*x + a) + 1/2) + 16)/(b*cos(b*x + a)^7 - 2*b*cos(b*x + a)^5 + b*cos(b*x + a)^3)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(a + bx)}{\sin^5(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4/sin(b*x+a)**5,x)**[Out]** Integral(sec(a + b*x)**4/sin(a + b*x)**5, x)**Giac [B]** Leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(79) = 158.

time = 3.79, size = 209, normalized size = 2.35

$$\frac{3 \left(\frac{24(\cos(bx+a)-1) - 210(\cos(bx+a)-1)^2 - 1}{(\cos(bx+a)+1)^2} (\cos(bx+a)+1)^2 - \frac{72(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{3(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{256 \left(\frac{9(\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{6(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + 5 \right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1} \right)^3} + 420 \log \left(\frac{-\cos(bx+a)+1}{|\cos(bx+a)+1|} \right) \right)}{192b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4/sin(b*x+a)^5,x, algorithm="giac")

[Out] 1/192*(3*(24*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 210*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 - 1)*(cos(b*x + a) + 1)^2/(cos(b*x + a) - 1)^2 - 72*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 3*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 256*(9*(cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 6*(cos(b*x + a) - 1)^2/(cos(b*x + a) + 1)^2 + 5)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)^3 + 420*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)))/b

Mupad [B]

time = 0.07, size = 78, normalized size = 0.88

$$\frac{\frac{35 \cos(a+bx)^6}{8} - \frac{175 \cos(a+bx)^4}{24} + \frac{7 \cos(a+bx)^2}{3} + \frac{1}{3}}{b (\cos(a + bx)^7 - 2 \cos(a + bx)^5 + \cos(a + bx)^3)} - \frac{35 \operatorname{atanh}(\cos(a + bx))}{8b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^4*sin(a + b*x)^5),x)

[Out] ((7*cos(a + b*x)^2)/3 - (175*cos(a + b*x)^4)/24 + (35*cos(a + b*x)^6)/8 + 1/3)/(b*(cos(a + b*x)^3 - 2*cos(a + b*x)^5 + cos(a + b*x)^7)) - (35*atanh(cos(a + b*x)))/(8*b)

3.184 $\int \csc^5(a + bx) \sec^5(a + bx) dx$

Optimal. Leaf size=69

$$-\frac{2 \cot^2(a + bx)}{b} - \frac{\cot^4(a + bx)}{4b} + \frac{6 \log(\tan(a + bx))}{b} + \frac{2 \tan^2(a + bx)}{b} + \frac{\tan^4(a + bx)}{4b}$$

[Out] $-2*\cot(b*x+a)^2/b-1/4*\cot(b*x+a)^4/b+6*\ln(\tan(b*x+a))/b+2*\tan(b*x+a)^2/b+1/4*\tan(b*x+a)^4/b$

Rubi [A]

time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.176$, Rules used = {2700, 272, 45}

$$\frac{\tan^4(a + bx)}{4b} + \frac{2 \tan^2(a + bx)}{b} - \frac{\cot^4(a + bx)}{4b} - \frac{2 \cot^2(a + bx)}{b} + \frac{6 \log(\tan(a + bx))}{b}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^5*Sec[a + b*x]^5,x]

[Out] $(-2*\cot[a + b*x]^2)/b - \cot[a + b*x]^4/(4*b) + (6*\log[\tan[a + b*x]])/b + (2*\tan[a + b*x]^2)/b + \tan[a + b*x]^4/(4*b)$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 2700

Int[csc[(e_.) + (f_.)*(x_)]^(m_.)*sec[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[1/f, Subst[Int[(1 + x^2)^(m + n)/2 - 1)/x^m, x], x, Tan[e + f*x]], x] /; FreeQ[{e, f}, x] && IntegersQ[m, n, (m + n)/2]

Rubi steps

$$\begin{aligned}
\int \csc^5(a+bx) \sec^5(a+bx) dx &= \frac{\text{Subst}\left(\int \frac{(1+x^2)^4}{x^5} dx, x, \tan(a+bx)\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{(1+x)^4}{x^3} dx, x, \tan^2(a+bx)\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \left(4 + \frac{1}{x^3} + \frac{4}{x^2} + \frac{6}{x} + x\right) dx, x, \tan^2(a+bx)\right)}{2b} \\
&= -\frac{2 \cot^2(a+bx)}{b} - \frac{\cot^4(a+bx)}{4b} + \frac{6 \log(\tan(a+bx))}{b} + \frac{2 \tan^2(a+bx)}{b} + \dots
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 91, normalized size = 1.32

$$32 \left(-\frac{3 \csc^2(a+bx)}{64b} - \frac{\csc^4(a+bx)}{128b} - \frac{3 \log(\cos(a+bx))}{16b} + \frac{3 \log(\sin(a+bx))}{16b} + \frac{3 \sec^2(a+bx)}{64b} + \frac{\sec^4(a+bx)}{128b} \right)$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^5*Sec[a + b*x]^5,x]`

```
[Out] 32*((-3*Csc[a + b*x]^2)/(64*b) - Csc[a + b*x]^4/(128*b) - (3*Log[Cos[a + b*x]])/(16*b) + (3*Log[Sin[a + b*x]])/(16*b) + (3*Sec[a + b*x]^2)/(64*b) + Sec[a + b*x]^4/(128*b))
```

Maple [A]

time = 0.05, size = 79, normalized size = 1.14

method	result	size
derivativedivides	$\frac{\frac{1}{4 \sin^4(bx+a)} \cos^4(bx+a) - \frac{1}{2 \sin^4(bx+a)} \cos^2(bx+a) + \frac{3}{2 \sin^2(bx+a)} \cos^2(bx+a) - \frac{3}{\sin^2(bx+a)} + 6 \ln(\tan(bx+a))}{b}$	79
default	$\frac{\frac{1}{4 \sin^4(bx+a)} \cos^4(bx+a) - \frac{1}{2 \sin^4(bx+a)} \cos^2(bx+a) + \frac{3}{2 \sin^2(bx+a)} \cos^2(bx+a) - \frac{3}{\sin^2(bx+a)} + 6 \ln(\tan(bx+a))}{b}$	79
risch	$\frac{12 e^{14i(bx+a)} - 44 e^{10i(bx+a)} - 44 e^{6i(bx+a)} + 12 e^{2i(bx+a)}}{b(e^{2i(bx+a)} - 1)^4 (e^{2i(bx+a)} + 1)^4} + \frac{6 \ln(e^{2i(bx+a)} - 1)}{b} - \frac{6 \ln(e^{2i(bx+a)} + 1)}{b}$	112

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sec(b*x+a)^5/sin(b*x+a)^5,x,method=_RETURNVERBOSE)`

```
[Out] 1/b*(1/4/sin(b*x+a)^4/cos(b*x+a)^4-1/2/sin(b*x+a)^4/cos(b*x+a)^2+3/2/sin(b*x+a)^2/cos(b*x+a)^2-3/sin(b*x+a)^2+6*ln(tan(b*x+a)))
```

Maxima [A]

time = 0.29, size = 92, normalized size = 1.33

$$\frac{12 \sin^6(bx+a) - 18 \sin^4(bx+a) + 4 \sin^2(bx+a) + 1}{\sin^8(bx+a) - 2 \sin^6(bx+a) + \sin^4(bx+a)} + 12 \log(\sin^2(bx+a) - 1) - 12 \log(\sin^2(bx+a))$$

4b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^5,x, algorithm="maxima")

[Out] $-1/4*((12*\sin(b*x + a)^6 - 18*\sin(b*x + a)^4 + 4*\sin(b*x + a)^2 + 1)/(\sin(b*x + a)^8 - 2*\sin(b*x + a)^6 + \sin(b*x + a)^4) + 12*\log(\sin(b*x + a)^2 - 1) - 12*\log(\sin(b*x + a)^2))/b$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 148 vs. 2(65) = 130.

time = 0.37, size = 148, normalized size = 2.14

$$\frac{12 \cos(bx+a)^6 - 18 \cos(bx+a)^4 + 4 \cos(bx+a)^2 - 12 (\cos(bx+a)^8 - 2 \cos(bx+a)^6 + \cos(bx+a)^4) \log(\cos(bx+a)^2) + 12 (\cos(bx+a)^8 - 2 \cos(bx+a)^6 + \cos(bx+a)^4) \log(-\frac{1}{4} \cos(bx+a)^2 + \frac{1}{4}) + 1}{4 (b \cos(bx+a)^8 - 2 b \cos(bx+a)^6 + b \cos(bx+a)^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^5,x, algorithm="fricas")

[Out] $1/4*(12*\cos(b*x + a)^6 - 18*\cos(b*x + a)^4 + 4*\cos(b*x + a)^2 - 12*(\cos(b*x + a)^8 - 2*\cos(b*x + a)^6 + \cos(b*x + a)^4)*\log(\cos(b*x + a)^2) + 12*(\cos(b*x + a)^8 - 2*\cos(b*x + a)^6 + \cos(b*x + a)^4)*\log(-1/4*\cos(b*x + a)^2 + 1/4) + 1)/(b*\cos(b*x + a)^8 - 2*b*\cos(b*x + a)^6 + b*\cos(b*x + a)^4)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^5(a + bx)}{\sin^5(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**5/sin(b*x+a)**5,x)

[Out] Integral(sec(a + b*x)**5/sin(a + b*x)**5, x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 278 vs. 2(65) = 130.

time = 3.91, size = 278, normalized size = 4.03

$$\frac{\left(\frac{28 \cos(bx+a)-11}{\cos(bx+a)+1} - \frac{288 (\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} - 1\right) \cos(bx+a)+1^2}{\cos(bx+a)+1} + \frac{28 (\cos(bx+a)-1)}{\cos(bx+a)+1} - \frac{(\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{32 \left(\frac{24 (\cos(bx+a)-1)}{\cos(bx+a)+1} + \frac{125 (\cos(bx+a)-1)^2}{(\cos(bx+a)+1)^2} + \frac{84 (\cos(bx+a)-1)^3}{(\cos(bx+a)+1)^3} + \frac{25 (\cos(bx+a)-1)^4}{(\cos(bx+a)+1)^4} + 25\right)}{\left(\frac{\cos(bx+a)-1}{\cos(bx+a)+1}\right)^4} + 192 \log\left(\frac{-\cos(bx+a)+1}{\cos(bx+a)+1}\right) - 384 \log\left(\frac{-\cos(bx+a)-1}{\cos(bx+a)+1} - 1\right)}$$

64b

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^5/sin(b*x+a)^5,x, algorithm="giac")

[Out] $1/64*((28*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - 288*(\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 - 1)*(\cos(b*x + a) + 1)^2/(\cos(b*x + a) - 1)^2 + 28*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) - (\cos(b*x + a) - 1)^2/(\cos(b*x + a) + 1)^2 + 32*(84*(\cos(b*x + a) - 1)/(\cos(b*x + a) + 1) + 126*(\cos(b*x + a) - 1$

)²/(cos(b*x + a) + 1)² + 84*(cos(b*x + a) - 1)³/(cos(b*x + a) + 1)³ + 25*(cos(b*x + a) - 1)⁴/(cos(b*x + a) + 1)⁴ + 25)/((cos(b*x + a) - 1)/(cos(b*x + a) + 1) + 1)⁴ + 192*log(abs(-cos(b*x + a) + 1)/abs(cos(b*x + a) + 1)) - 384*log(abs(-(cos(b*x + a) - 1)/(cos(b*x + a) + 1) - 1)))/b

Mupad [B]

time = 0.41, size = 64, normalized size = 0.93

$$\frac{2 \tan(a + bx)^2}{b} + \frac{\tan(a + bx)^4}{4b} + \frac{6 \ln(\tan(a + bx))}{b} - \frac{\cot(a + bx)^4 (2 \tan(a + bx)^2 + \frac{1}{4})}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(a + b*x)^5*sin(a + b*x)^5),x)

[Out] (2*tan(a + b*x)^2)/b + tan(a + b*x)^4/(4*b) + (6*log(tan(a + b*x)))/b - (cot(a + b*x)^4*(2*tan(a + b*x)^2 + 1/4))/b

3.185 $\int \cot^2(x) \csc^4(x) dx$

Optimal. Leaf size=17

$$-\frac{1}{3} \cot^3(x) - \frac{\cot^5(x)}{5}$$

[Out] -1/3*cot(x)^3-1/5*cot(x)^5

Rubi [A]

time = 0.02, antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2687, 14}

$$-\frac{1}{5} \cot^5(x) - \frac{\cot^3(x)}{3}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^2*Csc[x]^4,x]

[Out] -1/3*Cot[x]^3 - Cot[x]^5/5

Rule 14

```
Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2687

```
Int[sec[(e_) + (f_)*(x_)]^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Dist[1/f, Subst[Int[(b*x)^n*(1 + x^2)^(m/2 - 1), x], x, Tan[e + f*x]], x] /; FreeQ[{b, e, f, n}, x] && IntegerQ[m/2] && !(IntegerQ[(n - 1)/2] && LtQ[0, n, m - 1])
```

Rubi steps

$$\begin{aligned} \int \cot^2(x) \csc^4(x) dx &= \text{Subst} \left(\int x^2(1+x^2) dx, x, -\cot(x) \right) \\ &= \text{Subst} \left(\int (x^2 + x^4) dx, x, -\cot(x) \right) \\ &= -\frac{1}{3} \cot^3(x) - \frac{\cot^5(x)}{5} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 27, normalized size = 1.59

$$\frac{2 \cot(x)}{15} + \frac{1}{15} \cot(x) \csc^2(x) - \frac{1}{5} \cot(x) \csc^4(x)$$

Antiderivative was successfully verified.

`[In] Integrate[Cot[x]^2*Csc[x]^4,x]``[Out] (2*Cot[x])/15 + (Cot[x]*Csc[x]^2)/15 - (Cot[x]*Csc[x]^4)/5`**Maple [A]**

time = 0.04, size = 22, normalized size = 1.29

method	result	size
default	$-\frac{\cos^3(x)}{5 \sin(x)^5} - \frac{2(\cos^3(x))}{15 \sin(x)^3}$	22
risch	$-\frac{4i(15e^{6ix} + 5e^{4ix} + 5e^{2ix} - 1)}{15(e^{2ix} - 1)^5}$	36
norman	$-\frac{\frac{1}{160} - \frac{\tan^2(\frac{x}{2})}{96} + \frac{\tan^4(\frac{x}{2})}{16} - \frac{\tan^6(\frac{x}{2})}{96} + \frac{\tan^8(\frac{x}{2})}{160} + \frac{\tan^{10}(\frac{x}{2})}{160}}{\tan(\frac{x}{2})^5}$	50

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(x)^2/sin(x)^6,x,method=_RETURNVERBOSE)``[Out] -1/5/sin(x)^5*cos(x)^3-2/15/sin(x)^3*cos(x)^3`**Maxima [A]**

time = 0.28, size = 14, normalized size = 0.82

$$-\frac{5 \tan(x)^2 + 3}{15 \tan(x)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^2/sin(x)^6,x, algorithm="maxima")``[Out] -1/15*(5*tan(x)^2 + 3)/tan(x)^5`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 33 vs. 2(13) = 26.

time = 0.35, size = 33, normalized size = 1.94

$$\frac{2 \cos(x)^5 - 5 \cos(x)^3}{15 (\cos(x)^4 - 2 \cos(x)^2 + 1) \sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^2/sin(x)^6,x, algorithm="fricas")

[Out] 1/15*(2*cos(x)^5 - 5*cos(x)^3)/((cos(x)^4 - 2*cos(x)^2 + 1)*sin(x))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 29 vs. 2(14) = 28.

time = 0.01, size = 29, normalized size = 1.71

$$\frac{2 \cos(x)}{15 \sin(x)} + \frac{\cos(x)}{15 \sin^3(x)} - \frac{\cos(x)}{5 \sin^5(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)**2/sin(x)**6,x)

[Out] 2*cos(x)/(15*sin(x)) + cos(x)/(15*sin(x)**3) - cos(x)/(5*sin(x)**5)

Giac [A]

time = 3.25, size = 14, normalized size = 0.82

$$-\frac{5 \tan(x)^2 + 3}{15 \tan(x)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^2/sin(x)^6,x, algorithm="giac")

[Out] -1/15*(5*tan(x)^2 + 3)/tan(x)^5

Mupad [B]

time = 0.08, size = 19, normalized size = 1.12

$$-\cos(x)^3 \left(\frac{2}{15 \sin(x)^3} + \frac{1}{5 \sin(x)^5} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^2/sin(x)^6,x)

[Out] -cos(x)^3*(2/(15*sin(x)^3) + 1/(5*sin(x)^5))

3.186 $\int \cot^3(x) \csc^4(x) dx$

Optimal. Leaf size=17

$$\frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6}$$

[Out] 1/4*csc(x)^4-1/6*csc(x)^6

Rubi [A]

time = 0.02, antiderivative size = 17, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 9, $\frac{\text{number of rules}}{\text{integrand size}} = 0.222$, Rules used = {2686, 14}

$$\frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6}$$

Antiderivative was successfully verified.

[In] Int[Cot[x]^3*Csc[x]^4,x]

[Out] Csc[x]^4/4 - Csc[x]^6/6

Rule 14

```
Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2686

```
Int[((a_)*sec[(e_) + (f_)*(x_)])^(m_)*((b_)*tan[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Dist[a/f, Subst[Int[(a*x)^(m-1)*(-1+x^2)^((n-1)/2), x], x, Sec[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[m/2] && LtQ[0, m, n+1])
```

Rubi steps

$$\begin{aligned} \int \cot^3(x) \csc^4(x) dx &= -\text{Subst}\left(\int x^3(-1+x^2) dx, x, \csc(x)\right) \\ &= -\text{Subst}\left(\int (-x^3+x^5) dx, x, \csc(x)\right) \\ &= \frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 17, normalized size = 1.00

$$\frac{\csc^4(x)}{4} - \frac{\csc^6(x)}{6}$$

Antiderivative was successfully verified.

`[In] Integrate[Cot[x]^3*Csc[x]^4,x]``[Out] Csc[x]^4/4 - Csc[x]^6/6`**Maple [A]**

time = 0.05, size = 22, normalized size = 1.29

method	result	size
default	$-\frac{\cos^4(x)}{6 \sin(x)^6} - \frac{\cos^4(x)}{12 \sin(x)^4}$	22
norman	$-\frac{\frac{1}{384} + \frac{3(\tan^4(\frac{x}{2}))}{128} + \frac{3(\tan^8(\frac{x}{2}))}{128} - \frac{(\tan^{12}(\frac{x}{2}))}{384}}{\tan(\frac{x}{2})^6}$	34
risch	$\frac{4e^{8ix} + \frac{8e^{6ix}}{3} + 4e^{4ix}}{(e^{2ix} - 1)^6}$	34

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(x)^3/sin(x)^7,x,method=_RETURNVERBOSE)``[Out] -1/6/sin(x)^6*cos(x)^4-1/12/sin(x)^4*cos(x)^4`**Maxima [A]**

time = 0.28, size = 14, normalized size = 0.82

$$\frac{3 \sin(x)^2 - 2}{12 \sin(x)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3/sin(x)^7,x, algorithm="maxima")``[Out] 1/12*(3*sin(x)^2 - 2)/sin(x)^6`**Fricas [B]** Leaf count of result is larger than twice the leaf count of optimal. 30 vs. $2(13) = 26$.

time = 0.35, size = 30, normalized size = 1.76

$$\frac{3 \cos(x)^2 - 1}{12 (\cos(x)^6 - 3 \cos(x)^4 + 3 \cos(x)^2 - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3/sin(x)^7,x, algorithm="fricas")

[Out] 1/12*(3*cos(x)^2 - 1)/(cos(x)^6 - 3*cos(x)^4 + 3*cos(x)^2 - 1)

Sympy [A]

time = 0.03, size = 15, normalized size = 0.88

$$-\frac{2 - 3 \sin^2(x)}{12 \sin^6(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)**3/sin(x)**7,x)

[Out] -(2 - 3*sin(x)**2)/(12*sin(x)**6)

Giac [A]

time = 3.21, size = 14, normalized size = 0.82

$$\frac{3 \sin(x)^2 - 2}{12 \sin(x)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3/sin(x)^7,x, algorithm="giac")

[Out] 1/12*(3*sin(x)^2 - 2)/sin(x)^6

Mupad [B]

time = 0.44, size = 13, normalized size = 0.76

$$\frac{\frac{\sin(x)^2}{4} - \frac{1}{6}}{\sin(x)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^3/sin(x)^7,x)

[Out] (sin(x)^2/4 - 1/6)/sin(x)^6

3.187 $\int (d \cos(a + bx))^{3/2} \sin(a + bx) dx$

Optimal. Leaf size=22

$$-\frac{2(d \cos(a + bx))^{5/2}}{5bd}$$

[Out] $-2/5*(d*\cos(b*x+a))^{(5/2)}/b/d$

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$-\frac{2(d \cos(a + bx))^{5/2}}{5bd}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x], x]$

[Out] $(-2*(d*\text{Cos}[a + b*x])^{(5/2)})/(5*b*d)$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \text{ :> Simp}[x^{(m + 1)}/(m + 1), x] \text{ /; FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2645

$\text{Int}[(\text{Cos}[(e_.) + (f_.)*(x_)]*(a_.))^{(m_.)}*\text{Sin}[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] \text{ :> Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{((n - 1)/2)}, x], x, a*\text{Cos}[e + f*x]], x] \text{ /; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n - 1)/2] \ \&\& \ !(\text{IntegerQ}[(m - 1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{3/2} \sin(a + bx) dx &= -\frac{\text{Subst}(\int x^{3/2} dx, x, d \cos(a + bx))}{bd} \\ &= -\frac{2(d \cos(a + bx))^{5/2}}{5bd} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 22, normalized size = 1.00

$$-\frac{2(d \cos(a + bx))^{5/2}}{5bd}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(3/2)*sin[a + b*x],x]

[Out] $(-2*(d*\cos[a + b*x])^{(5/2)})/(5*b*d)$

Maple [A]

time = 0.03, size = 19, normalized size = 0.86

method	result	size
derivativeldivides	$-\frac{2(d\cos(bx+a))^{5/2}}{5bd}$	19
default	$-\frac{2(d\cos(bx+a))^{5/2}}{5bd}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(3/2)*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] $-2/5*(d*\cos(b*x+a))^{(5/2)}/b/d$

Maxima [A]

time = 0.29, size = 18, normalized size = 0.82

$$\frac{2(d\cos(bx+a))^{5/2}}{5bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="maxima")

[Out] $-2/5*(d*\cos(b*x + a))^{(5/2)}/(b*d)$

Fricas [A]

time = 0.37, size = 24, normalized size = 1.09

$$\frac{2\sqrt{d\cos(bx+a)}d\cos(bx+a)^2}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="fricas")

[Out] $-2/5*\text{sqrt}(d*\cos(b*x + a))*d*\cos(b*x + a)^2/b$

Sympy [A]

time = 4.73, size = 37, normalized size = 1.68

$$\begin{cases} -\frac{2(d\cos(a+bx))^{3/2}\cos(a+bx)}{5b} & \text{for } b \neq 0 \\ x(d\cos(a))^{3/2}\sin(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(3/2)*sin(b*x+a),x)

[Out] Piecewise((-2*(d*cos(a + b*x))**(3/2)*cos(a + b*x)/(5*b), Ne(b, 0)), (x*(d*cos(a))**(3/2)*sin(a), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*sin(b*x + a), x)

Mupad [B]

time = 0.13, size = 18, normalized size = 0.82

$$\frac{2(d \cos(a + bx))^{5/2}}{5bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d*cos(a + b*x))^(3/2),x)

[Out] -(2*(d*cos(a + b*x))^(5/2))/(5*b*d)

3.188 $\int \sqrt{d \cos(a + bx)} \sin(a + bx) dx$

Optimal. Leaf size=22

$$-\frac{2(d \cos(a + bx))^{3/2}}{3bd}$$

[Out] $-2/3*(d*\cos(b*x+a))^(3/2)/b/d$

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$-\frac{2(d \cos(a + bx))^{3/2}}{3bd}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x],x]`

[Out] $(-2*(d*\text{Cos}[a + b*x])^(3/2))/(3*b*d)$

Rule 30

`Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 2645

`Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} \sin(a + bx) dx &= -\frac{\text{Subst}\left(\int \sqrt{x} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{2(d \cos(a + bx))^{3/2}}{3bd} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 22, normalized size = 1.00

$$-\frac{2(d \cos(a + bx))^{3/2}}{3bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x],x]

[Out] $(-2*(d*\cos[a + b*x])^{(3/2)})/(3*b*d)$

Maple [A]

time = 0.02, size = 19, normalized size = 0.86

method	result	size
derivativedivides	$-\frac{2(d\cos(bx+a))^{\frac{3}{2}}}{3bd}$	19
default	$-\frac{2(d\cos(bx+a))^{\frac{3}{2}}}{3bd}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] $-2/3*(d*\cos(b*x+a))^{(3/2)}/b/d$

Maxima [A]

time = 0.28, size = 18, normalized size = 0.82

$$-\frac{2(d\cos(bx+a))^{\frac{3}{2}}}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="maxima")

[Out] $-2/3*(d*\cos(b*x + a))^{(3/2)}/(b*d)$

Fricas [A]

time = 0.37, size = 21, normalized size = 0.95

$$-\frac{2\sqrt{d\cos(bx+a)}\cos(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="fricas")

[Out] $-2/3*\text{sqrt}(d*\cos(b*x + a))*\cos(b*x + a)/b$

Sympy [A]

time = 0.71, size = 37, normalized size = 1.68

$$\begin{cases} -\frac{2\sqrt{d\cos(a+bx)}\cos(a+bx)}{3b} & \text{for } b \neq 0 \\ x\sqrt{d\cos(a)}\sin(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(1/2)*sin(b*x+a),x)`

[Out] `Piecewise((-2*sqrt(d*cos(a + b*x))*cos(a + b*x)/(3*b), Ne(b, 0)), (x*sqrt(d*cos(a))*sin(a), True))`

Giac [A]

time = 5.94, size = 21, normalized size = 0.95

$$-\frac{2\sqrt{d\cos(bx+a)}\cos(bx+a)}{3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a),x, algorithm="giac")`

[Out] `-2/3*sqrt(d*cos(b*x + a))*cos(b*x + a)/b`

Mupad [B]

time = 0.42, size = 18, normalized size = 0.82

$$-\frac{2(d\cos(a+bx))^{3/2}}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)*(d*cos(a + b*x))^(1/2),x)`

[Out] `-(2*(d*cos(a + b*x))^(3/2))/(3*b*d)`

$$3.189 \quad \int \frac{\sin(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=20

$$-\frac{2\sqrt{d \cos(a+bx)}}{bd}$$

[Out] $-2*(d*\cos(b*x+a))^{(1/2)}/b/d$

Rubi [A]

time = 0.02, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$-\frac{2\sqrt{d \cos(a+bx)}}{bd}$$

Antiderivative was successfully verified.

[In] `Int[Sin[a + b*x]/Sqrt[d*Cos[a + b*x]],x]`

[Out] `(-2*Sqrt[d*Cos[a + b*x]])/(b*d)`

Rule 30

`Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]`

Rule 2645

`Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

Rubi steps

$$\begin{aligned} \int \frac{\sin(a+bx)}{\sqrt{d \cos(a+bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{2\sqrt{d \cos(a+bx)}}{bd} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 20, normalized size = 1.00

$$-\frac{2\sqrt{d\cos(a+bx)}}{bd}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]/Sqrt[d*Cos[a + b*x]],x]

[Out] (-2*Sqrt[d*Cos[a + b*x]])/(b*d)

Maple [A]

time = 0.03, size = 19, normalized size = 0.95

method	result	size
derivativedivides	$-\frac{2\sqrt{d\cos(bx+a)}}{bd}$	19
default	$-\frac{2\sqrt{d\cos(bx+a)}}{bd}$	19
risch	$-\frac{2\cos(bx+a)}{\sqrt{d\cos(bx+a)}b}$	22

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)/(d*cos(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] -2*(d*cos(b*x+a))^(1/2)/b/d

Maxima [A]

time = 0.29, size = 18, normalized size = 0.90

$$-\frac{2\sqrt{d\cos(bx+a)}}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] -2*sqrt(d*cos(b*x + a))/(b*d)

Fricas [A]

time = 0.36, size = 18, normalized size = 0.90

$$-\frac{2\sqrt{d\cos(bx+a)}}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] $-2\sqrt{d\cos(bx + a)}/(bd)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 36 vs. $2(17) = 34$.

time = 0.70, size = 36, normalized size = 1.80

$$\begin{cases} -\frac{2\cos(a+bx)}{b\sqrt{d\cos(a+bx)}} & \text{for } b \neq 0 \\ \frac{x\sin(a)}{\sqrt{d\cos(a)}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)/(d*cos(b*x+a))**(1/2),x)`

[Out] `Piecewise((-2*cos(a + b*x)/(b*sqrt(d*cos(a + b*x))), Ne(b, 0)), (x*sin(a)/sqrt(d*cos(a)), True))`

Giac [A]

time = 4.62, size = 18, normalized size = 0.90

$$-\frac{2\sqrt{d\cos(bx+a)}}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)/(d*cos(b*x+a))^(1/2),x, algorithm="giac")`

[Out] $-2\sqrt{d\cos(bx + a)}/(bd)$

Mupad [B]

time = 0.53, size = 18, normalized size = 0.90

$$-\frac{2\sqrt{d\cos(a+bx)}}{bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)/(d*cos(a + b*x))^(1/2),x)`

[Out] $-(2*(d*cos(a + b*x))^(1/2))/(b*d)$

$$3.190 \quad \int \frac{\sin(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=20

$$\frac{2}{bd\sqrt{d \cos(a+bx)}}$$

[Out] 2/b/d/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$\frac{2}{bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]/(d*Cos[a + b*x])^(3/2), x]

[Out] 2/(b*d*Sqrt[d*Cos[a + b*x]])

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(a+bx)}{(d \cos(a+bx))^{3/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{3/2}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= \frac{2}{bd\sqrt{d \cos(a+bx)}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 20, normalized size = 1.00

$$\frac{2}{bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]/(d*cos[a + b*x])^(3/2),x]

[Out] 2/(b*d*Sqrt[d*cos[a + b*x]])

Maple [A]

time = 0.02, size = 19, normalized size = 0.95

method	result	size
derivativedivides	$\frac{2}{bd\sqrt{d\cos(bx+a)}}$	19
default	$\frac{2}{bd\sqrt{d\cos(bx+a)}}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)/(d*cos(b*x+a))^(3/2),x,method=_RETURNVERBOSE)

[Out] 2/b/d/(d*cos(b*x+a))^(1/2)

Maxima [A]

time = 0.29, size = 18, normalized size = 0.90

$$\frac{2}{\sqrt{d\cos(bx+a)}bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] 2/(sqrt(d*cos(b*x + a))*b*d)

Fricas [A]

time = 0.36, size = 26, normalized size = 1.30

$$\frac{2\sqrt{d\cos(bx+a)}}{bd^2\cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] 2*sqrt(d*cos(b*x + a))/(b*d^2*cos(b*x + a))

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 34 vs. 2(15) = 30.

time = 1.22, size = 34, normalized size = 1.70

$$\begin{cases} \frac{2\cos(a+bx)}{b(d\cos(a+bx))^{\frac{3}{2}}} & \text{for } b \neq 0 \\ \frac{x\sin(a)}{(d\cos(a))^{\frac{3}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)/(d*cos(b*x+a))**(3/2),x)`

[Out] `Piecewise((2*cos(a + b*x)/(b*(d*cos(a + b*x))**(3/2)), Ne(b, 0)), (x*sin(a)/(d*cos(a))**(3/2), True))`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)/(d*cos(b*x+a))^(3/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)/(d*cos(b*x + a))^(3/2), x)`

Mupad [B]

time = 0.18, size = 37, normalized size = 1.85

$$\frac{4 \cos(a + b x) \sqrt{d \cos(a + b x)}}{b d^2 (\cos(2 a + 2 b x) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)/(d*cos(a + b*x))^(3/2),x)`

[Out] `(4*cos(a + b*x)*(d*cos(a + b*x))^(1/2))/(b*d^2*(cos(2*a + 2*b*x) + 1))`

$$3.191 \quad \int \frac{\sin(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=22

$$\frac{2}{3bd(d \cos(a + bx))^{3/2}}$$

[Out] 2/3/b/d/(d*cos(b*x+a))^(3/2)

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$\frac{2}{3bd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]/(d*Cos[a + b*x])^(5/2),x]

[Out] 2/(3*b*d*(d*Cos[a + b*x])^(3/2))

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2645

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(a + bx)}{(d \cos(a + bx))^{5/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{5/2}} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2}{3bd(d \cos(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 22, normalized size = 1.00

$$\frac{2}{3bd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]/(d*Cos[a + b*x])^(5/2), x]

[Out] 2/(3*b*d*(d*Cos[a + b*x])^(3/2))

Maple [A]

time = 0.02, size = 19, normalized size = 0.86

method	result	size
derivativedivides	$\frac{2}{3bd(d \cos(bx+a))^{\frac{3}{2}}}$	19
default	$\frac{2}{3bd(d \cos(bx+a))^{\frac{3}{2}}}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)/(d*cos(b*x+a))^(5/2), x, method=_RETURNVERBOSE)

[Out] 2/3/b/d/(d*cos(b*x+a))^(3/2)

Maxima [A]

time = 0.27, size = 18, normalized size = 0.82

$$\frac{2}{3 (d \cos (bx + a))^{\frac{3}{2}} bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(5/2), x, algorithm="maxima")

[Out] 2/3/((d*cos(b*x + a))^(3/2)*b*d)

Fricas [A]

time = 0.35, size = 26, normalized size = 1.18

$$\frac{2 \sqrt{d \cos (bx + a)}}{3 b d^3 \cos (bx + a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(5/2), x, algorithm="fricas")

[Out] 2/3*sqrt(d*cos(b*x + a))/(b*d^3*cos(b*x + a)^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(17) = 34.

time = 5.06, size = 36, normalized size = 1.64

$$\begin{cases} \frac{2 \cos (a+bx)}{3 b(d \cos (a+bx))^{\frac{5}{2}}} & \text{for } b \neq 0 \\ \frac{x \sin (a)}{(d \cos (a))^{\frac{5}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))**(5/2),x)

[Out] Piecewise((2*cos(a + b*x)/(3*b*(d*cos(a + b*x))**(5/2)), Ne(b, 0)), (x*sin(a)/(d*cos(a))**(5/2), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)/(d*cos(b*x + a))^(5/2), x)

Mupad [B]

time = 0.81, size = 53, normalized size = 2.41

$$\frac{8(\cos(2a + 2bx) + 1)\sqrt{d\cos(a + bx)}}{3bd^3(4\cos(2a + 2bx) + \cos(4a + 4bx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/(d*cos(a + b*x))^(5/2),x)

[Out] (8*(cos(2*a + 2*b*x) + 1)*(d*cos(a + b*x))^(1/2))/(3*b*d^3*(4*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 3))

$$3.192 \quad \int \frac{\sin(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=22

$$\frac{2}{5bd(d \cos(a + bx))^{5/2}}$$

[Out] 2/5/b/d/(d*cos(b*x+a))^(5/2)

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$\frac{2}{5bd(d \cos(a + bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]/(d*Cos[a + b*x])^(7/2), x]

[Out] 2/(5*b*d*(d*Cos[a + b*x])^(5/2))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_.))^(m_.)*sin[(e_) + (f_)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(a + bx)}{(d \cos(a + bx))^{7/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{7/2}} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2}{5bd(d \cos(a + bx))^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 22, normalized size = 1.00

$$\frac{2}{5bd(d \cos(a + bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]/(d*cos[a + b*x])^(7/2),x]

[Out] 2/(5*b*d*(d*cos[a + b*x])^(5/2))

Maple [A]

time = 0.04, size = 19, normalized size = 0.86

method	result	size
derivativedivides	$\frac{2}{5bd(d \cos(bx+a))^{\frac{5}{2}}}$	19
default	$\frac{2}{5bd(d \cos(bx+a))^{\frac{5}{2}}}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)

[Out] 2/5/b/d/(d*cos(b*x+a))^(5/2)

Maxima [A]

time = 0.27, size = 18, normalized size = 0.82

$$\frac{2}{5 (d \cos(bx + a))^{\frac{5}{2}} bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")

[Out] 2/5/((d*cos(b*x + a))^(5/2)*b*d)

Fricas [A]

time = 0.34, size = 26, normalized size = 1.18

$$\frac{2 \sqrt{d \cos(bx + a)}}{5 bd^4 \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")

[Out] 2/5*sqrt(d*cos(b*x + a))/(b*d^4*cos(b*x + a)^3)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(17) = 34.

time = 46.04, size = 36, normalized size = 1.64

$$\begin{cases} \frac{2 \cos(a+bx)}{5b(d \cos(a+bx))^{\frac{7}{2}}} & \text{for } b \neq 0 \\ \frac{x \sin(a)}{(d \cos(a))^{\frac{7}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))**(7/2),x)

[Out] Piecewise((2*cos(a + b*x)/(5*b*(d*cos(a + b*x))**(7/2)), Ne(b, 0)), (x*sin(a)/(d*cos(a))**(7/2), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)/(d*cos(b*x + a))^(7/2), x)

Mupad [B]

time = 6.71, size = 65, normalized size = 2.95

$$\frac{16e^{a3i+b x 3i} \sqrt{d \left(\frac{e^{-a 1i-b x 1i}}{2} + \frac{e^{a 1i+b x 1i}}{2} \right)}}{5bd^4 (e^{a2i+b x 2i} + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/(d*cos(a + b*x))^(7/2),x)

[Out] (16*exp(a*3i + b*x*3i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2))/(5*b*d^4*(exp(a*2i + b*x*2i) + 1)^3)

$$3.193 \quad \int \frac{\sin(a+bx)}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=22

$$\frac{2}{7bd(d \cos(a + bx))^{7/2}}$$

[Out] 2/7/b/d/(d*cos(b*x+a))^(7/2)

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$\frac{2}{7bd(d \cos(a + bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]/(d*Cos[a + b*x])^(9/2),x]

[Out] 2/(7*b*d*(d*Cos[a + b*x])^(7/2))

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2645

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(a + bx)}{(d \cos(a + bx))^{9/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{9/2}} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2}{7bd(d \cos(a + bx))^{7/2}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 22, normalized size = 1.00

$$\frac{2}{7bd(d \cos(a + bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]/(d*cos[a + b*x])^(9/2), x]

[Out] $2/(7*b*d*(d*\cos[a + b*x])^(7/2))$

Maple [A]

time = 0.02, size = 19, normalized size = 0.86

method	result	size
derivativdivides	$\frac{2}{7bd(d\cos(bx+a))^{\frac{7}{2}}}$	19
default	$\frac{2}{7bd(d\cos(bx+a))^{\frac{7}{2}}}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)/(d*cos(b*x+a))^(9/2), x, method=_RETURNVERBOSE)

[Out] $2/7/b/d/(d*\cos(b*x+a))^(7/2)$

Maxima [A]

time = 0.29, size = 18, normalized size = 0.82

$$\frac{2}{7(d\cos(bx+a))^{\frac{7}{2}}bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(9/2), x, algorithm="maxima")

[Out] $2/7/((d*\cos(b*x + a))^(7/2)*b*d)$

Fricas [A]

time = 0.35, size = 26, normalized size = 1.18

$$\frac{2\sqrt{d\cos(bx+a)}}{7bd^5\cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(9/2), x, algorithm="fricas")

[Out] $2/7*\sqrt{d*\cos(b*x + a)}/(b*d^5*\cos(b*x + a)^4)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))**(9/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4847 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)/(d*cos(b*x + a))^(9/2), x)

Mupad [B]

time = 4.04, size = 65, normalized size = 2.95

$$\frac{32 e^{a 4i + b x 4i} \sqrt{d \left(\frac{e^{-a 1i - b x 1i}}{2} + \frac{e^{a 1i + b x 1i}}{2} \right)}}{7 b d^5 (e^{a 2i + b x 2i} + 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)/(d*cos(a + b*x))^(9/2),x)

[Out] (32*exp(a*4i + b*x*4i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2))/(7*b*d^5*(exp(a*2i + b*x*2i) + 1)^4)

3.194 $\int (d \cos(a + bx))^{9/2} \sin^2(a + bx) dx$

Optimal. Leaf size=126

$$\frac{28d^4 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{195b \sqrt{\cos(a + bx)}} + \frac{28d^3 (d \cos(a + bx))^{3/2} \sin(a + bx)}{585b} + \frac{4d (d \cos(a + bx))^{7/2} \sin(a + bx)}{117b}$$

[Out] $28/585*d^3*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)/b+4/117*d*(d*\cos(b*x+a))^{(7/2)}*\sin(b*x+a)/b-2/13*(d*\cos(b*x+a))^{(11/2)}*\sin(b*x+a)/b/d+28/195*d^4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2719}

$$\frac{28d^4 E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{d \cos(a + bx)}}{195b \sqrt{\cos(a + bx)}} + \frac{28d^3 \sin(a + bx) (d \cos(a + bx))^{3/2}}{585b} - \frac{2 \sin(a + bx) (d \cos(a + bx))^{11/2}}{13bd} + \frac{4d \sin(a + bx) (d \cos(a + bx))^{7/2}}{117b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(9/2)}*\text{Sin}[a + b*x]^2, x]$

[Out] $(28*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(195*b*\text{Sqrt}[\text{Cos}[a + b*x]]) + (28*d^3*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x])/(585*b) + (4*d*(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x])/(117*b) - (2*(d*\text{Cos}[a + b*x])^{(11/2)}*\text{Sin}[a + b*x])/(13*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.))^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] :> \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{n*} (a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_)])^{(n_)}], x_Symbol] :> \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n - 1)})/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{9/2} \sin^2(a + bx) dx &= -\frac{2(d \cos(a + bx))^{11/2} \sin(a + bx)}{13bd} + \frac{2}{13} \int (d \cos(a + bx))^{9/2} dx \\
&= \frac{4d(d \cos(a + bx))^{7/2} \sin(a + bx)}{117b} - \frac{2(d \cos(a + bx))^{11/2} \sin(a + bx)}{13bd} + \\
&= \frac{28d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{585b} + \frac{4d(d \cos(a + bx))^{7/2} \sin(a + bx)}{117b} \\
&= \frac{28d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{585b} + \frac{4d(d \cos(a + bx))^{7/2} \sin(a + bx)}{117b} \\
&= \frac{28d^4 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{195b \sqrt{\cos(a + bx)}} + \frac{28d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{585b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.10, size = 60, normalized size = 0.48

$$\frac{d^2(d \cos(a + bx))^{5/2} \sqrt{\cos^2(a + bx)} {}_2F_1\left(-\frac{7}{4}, \frac{3}{2}; \frac{5}{2}; \sin^2(a + bx)\right) \tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(9/2)*Sin[a + b*x]^2,x]
```

```
[Out] (d^2*(d*Cos[a + b*x])^(5/2)*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-7/4,
3/2, 5/2, Sin[a + b*x]^2]*Tan[a + b*x]^3)/(3*b)
```

Maple [A]

time = 0.12, size = 249, normalized size = 1.98

method	result
default	$4 \sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)} d^5 \left(2880 \left(\cos^{15} \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 11520 \left(\cos^{13} \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 19280 \left(\cos^{11} \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 11520 \left(\cos^9 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 2880 \left(\cos^7 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right) \tan^3 \left(\frac{bx}{2} + \frac{a}{2} \right) \sqrt{-d} \left(2 \left(\sin \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(9/2)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{4}{585} \cdot (d \cdot (2 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a))^{-2-1}) \cdot \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{(1/2)} \cdot d^5 \cdot (2880 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^{15} - 11520 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^{13} + 19280 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^{11} - 17520 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^9 + 9284 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^7 - 2808 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^5 + 425 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^3 + 21 \cdot (\sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{(1/2)} \cdot (-2 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 + 1)^{(1/2)} \cdot \text{EllipticE}(\cos(1/2 \cdot b \cdot x + 1/2 \cdot a), 2^{(1/2)}) - 21 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)) / (-d \cdot (2 \cdot \sin(1/2 \cdot b \cdot x + 1/2 \cdot a))^4 - \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{(1/2)} / \sin(1/2 \cdot b \cdot x + 1/2 \cdot a) / (d \cdot (2 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a))^{-2-1})^{(1/2)} / b$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*sin(b*x+a)^2,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(9/2)*sin(b*x + a)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 119, normalized size = 0.94

$$\frac{2 \left(-21i \sqrt{2} d^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a))) + 21i \sqrt{2} d^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a))) + (45 d^4 \cos(bx + a)^5 - 10 d^4 \cos(bx + a)^3 - 14 d^4 \cos(bx + a)) \sqrt{\cos(bx + a)} \sin(bx + a) \right)}{585 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*sin(b*x+a)^2,x, algorithm="fricas")`

[Out]
$$-2/585 \cdot (-21 \cdot I \cdot \sqrt{2}) \cdot d^{(9/2)} \cdot \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b \cdot x + a) + I \cdot \sin(b \cdot x + a))) + 21 \cdot I \cdot \sqrt{2} \cdot d^{(9/2)} \cdot \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b \cdot x + a) - I \cdot \sin(b \cdot x + a))) + (45 \cdot d^4 \cdot \cos(b \cdot x + a)^5 - 10 \cdot d^4 \cdot \cos(b \cdot x + a)^3 - 14 \cdot d^4 \cdot \cos(b \cdot x + a)) \cdot \sqrt{\cos(b \cdot x + a)} \cdot \sin(b \cdot x + a)) / b$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(9/2)*sin(b*x+a)**2,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(9/2)*sin(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(9/2)*sin(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^2 (d \cos(a + bx))^{9/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2*(d*cos(a + b*x))^(9/2),x)

[Out] int(sin(a + b*x)^2*(d*cos(a + b*x))^(9/2), x)

3.195 $\int (d \cos(a + bx))^{7/2} \sin^2(a + bx) dx$

Optimal. Leaf size=126

$$\frac{20d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{231b \sqrt{d \cos(a + bx)}} + \frac{20d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b} + \frac{4d(d \cos(a + bx))^{5/2} \sin(a + bx)}{77b}$$

[Out] $4/77*d*(d*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)/b-2/11*(d*\cos(b*x+a))^{(9/2)}*\sin(b*x+a)/b/d+20/231*d^4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}+20/231*d^3*\sin(b*x+a)*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.08, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2720}

$$\frac{20d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{231b \sqrt{d \cos(a + bx)}} + \frac{20d^3 \sin(a + bx) \sqrt{d \cos(a + bx)}}{231b} - \frac{2 \sin(a + bx) (d \cos(a + bx))^{9/2}}{11bd} + \frac{4d \sin(a + bx) (d \cos(a + bx))^{5/2}}{77b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x]^2, x]$

[Out] $(20*d^4*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(231*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (20*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(231*b) + (4*d*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x])/(77*b) - (2*(d*\text{Cos}[a + b*x])^{(9/2)}*\text{Sin}[a + b*x])/(11*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.))^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^n*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_)])^{(n_)}], x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n - 1)}/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

`Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \int (d \cos(a + bx))^{7/2} \sin^2(a + bx) dx &= -\frac{2(d \cos(a + bx))^{9/2} \sin(a + bx)}{11bd} + \frac{2}{11} \int (d \cos(a + bx))^{7/2} dx \\
 &= \frac{4d(d \cos(a + bx))^{5/2} \sin(a + bx)}{77b} - \frac{2(d \cos(a + bx))^{9/2} \sin(a + bx)}{11bd} + \frac{2}{7} \int (d \cos(a + bx))^{5/2} dx \\
 &= \frac{20d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b} + \frac{4d(d \cos(a + bx))^{5/2} \sin(a + bx)}{77b} - \frac{2}{7} \int (d \cos(a + bx))^{3/2} dx \\
 &= \frac{20d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b} + \frac{4d(d \cos(a + bx))^{5/2} \sin(a + bx)}{77b} - \frac{2}{7} \int (d \cos(a + bx))^{1/2} dx \\
 &= \frac{20d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{231b \sqrt{d \cos(a + bx)}} + \frac{20d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.10, size = 60, normalized size = 0.48

$$\frac{d^2 (d \cos(a + bx))^{3/2} \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{5}{4}, \frac{3}{2}; \frac{5}{2}; \sin^2(a + bx)\right) \tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(7/2)*Sin[a + b*x]^2,x]`

`[Out] (d^2*(d*Cos[a + b*x])^(3/2)*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-5/4, 3/2, 5/2, Sin[a + b*x]^2]*Tan[a + b*x]^3)/(3*b)`

Maple [A]

time = 0.11, size = 236, normalized size = 1.87

method	result
default	$\frac{4 \sqrt{d} \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^4 d^4 \left(672 \cos^{13}\left(\frac{bx}{2} + \frac{a}{2}\right) - 2352 \cos^{11}\left(\frac{bx}{2} + \frac{a}{2}\right) + 3312 \cos^9\left(\frac{bx}{2} + \frac{a}{2}\right) - 1280 \cos^7\left(\frac{bx}{2} + \frac{a}{2}\right) + 256 \cos^5\left(\frac{bx}{2} + \frac{a}{2}\right) - 32 \cos^3\left(\frac{bx}{2} + \frac{a}{2}\right) + 4 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{231 \sqrt{-d} \left(2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(7/2)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{4}{231} \cdot (d \cdot (2 \cos(1/2 \cdot b \cdot x + 1/2 \cdot a))^2 - 1) \cdot \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{(1/2)} \cdot d^4 \cdot (672 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^{13} - 2352 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^{11} + 3312 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^9 - 2400 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^7 + 922 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^5 - 159 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^3 - 5 \cdot (\sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{(1/2)} \cdot (-2 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 + 1)^{(1/2)} \cdot \text{EllipticF}(\cos(1/2 \cdot b \cdot x + 1/2 \cdot a), 2^{(1/2)}) + 5 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)) / (-d \cdot (2 \cdot \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^4 - \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2))^{(1/2)} / \sin(1/2 \cdot b \cdot x + 1/2 \cdot a) / (d \cdot (2 \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a))^2 - 1)^{(1/2)} / b$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*sin(b*x+a)^2,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(7/2)*sin(b*x + a)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 107, normalized size = 0.85

$$\frac{2 \left(5i \sqrt{2} d^{\frac{5}{2}} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) - 5i \sqrt{2} d^{\frac{7}{2}} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)) + (21 d^6 \cos(bx + a)^4 - 6 d^6 \cos(bx + a)^2 - 10 d^6) \sqrt{d \cos(bx + a)} \sin(bx + a) \right)}{231 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*sin(b*x+a)^2,x, algorithm="fricas")`

[Out]
$$-2/231 \cdot (5 \cdot I \cdot \sqrt{2} \cdot d^{(7/2)} \cdot \text{weierstrassPInverse}(-4, 0, \cos(b \cdot x + a) + I \cdot \sin(b \cdot x + a)) - 5 \cdot I \cdot \sqrt{2} \cdot d^{(7/2)} \cdot \text{weierstrassPInverse}(-4, 0, \cos(b \cdot x + a) - I \cdot \sin(b \cdot x + a)) + (21 \cdot d^3 \cdot \cos(b \cdot x + a)^4 - 6 \cdot d^3 \cdot \cos(b \cdot x + a)^2 - 10 \cdot d^3) \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \sin(b \cdot x + a)) / b$$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(7/2)*sin(b*x+a)**2,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 5986 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(7/2)*sin(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(7/2)*sin(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^2 (d \cos(a + bx))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2*(d*cos(a + b*x))^(7/2),x)

[Out] int(sin(a + b*x)^2*(d*cos(a + b*x))^(7/2), x)

3.196 $\int (d \cos(a + bx))^{5/2} \sin^2(a + bx) dx$

Optimal. Leaf size=98

$$\frac{4d^2 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{15b \sqrt{\cos(a + bx)}} + \frac{4d(d \cos(a + bx))^{3/2} \sin(a + bx)}{45b} - \frac{2(d \cos(a + bx))^{7/2} \sin(a + bx)}{9bd}$$

[Out] $4/45*d*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)/b-2/9*(d*\cos(b*x+a))^{(7/2)}*\sin(b*x+a)/b/d+4/15*d^2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2719}

$$\frac{4d^2 E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{15b \sqrt{\cos(a + bx)}} - \frac{2 \sin(a + bx)(d \cos(a + bx))^{7/2}}{9bd} + \frac{4d \sin(a + bx)(d \cos(a + bx))^{3/2}}{45b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x]^2, x]$

[Out] $(4*d^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(15*b*\text{Sqrt}[\text{Cos}[a + b*x]]) + (4*d*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x])/(45*b) - (2*(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x])/(9*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.)^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}), x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n - 1)})/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{5/2} \sin^2(a + bx) dx &= -\frac{2(d \cos(a + bx))^{7/2} \sin(a + bx)}{9bd} + \frac{2}{9} \int (d \cos(a + bx))^{5/2} dx \\ &= \frac{4d(d \cos(a + bx))^{3/2} \sin(a + bx)}{45b} - \frac{2(d \cos(a + bx))^{7/2} \sin(a + bx)}{9bd} + \frac{2}{9} \int (d \cos(a + bx))^{5/2} dx \\ &= \frac{4d(d \cos(a + bx))^{3/2} \sin(a + bx)}{45b} - \frac{2(d \cos(a + bx))^{7/2} \sin(a + bx)}{9bd} + \frac{2}{9} \int (d \cos(a + bx))^{5/2} dx \\ &= \frac{4d^2 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{15b \sqrt{\cos(a + bx)}} + \frac{4d(d \cos(a + bx))^{3/2} \sin(a + bx)}{45b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.04, size = 57, normalized size = 0.58

$$\frac{(d \cos(a + bx))^{5/2} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(-\frac{3}{4}, \frac{3}{2}; \frac{5}{2}; \sin^2(a + bx)\right) \tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(5/2)*Sin[a + b*x]^2,x]
```

```
[Out] ((d*Cos[a + b*x])^(5/2)*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-3/4, 3/2, 5/2, Sin[a + b*x]^2]*Tan[a + b*x]^3)/(3*b)
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 222 vs. 2(110) = 220.

time = 0.12, size = 223, normalized size = 2.28

method	result
default	$\frac{4 \sqrt{d \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{d^3 \left(80 \left(\cos^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 240 \left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 272 \left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 144 \left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 48 \left(\cos^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 8 \left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 8} - 45 \sqrt{-d \left(2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}$

Verification of antiderivative is not currently implemented for this CAS.


```
[In] int((d*cos(b*x+a))^(5/2)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)
[Out] 4/45*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^3*(80*cos(
1/2*b*x+1/2*a)^11-240*cos(1/2*b*x+1/2*a)^9+272*cos(1/2*b*x+1/2*a)^7-144*cos
(1/2*b*x+1/2*a)^5+35*cos(1/2*b*x+1/2*a)^3+3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-
2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))-3*cos
(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/s
in(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*sin(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(5/2)*sin(b*x + a)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 106, normalized size = 1.08

$$\frac{2(-3i\sqrt{2}d^2\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+3i\sqrt{2}d^2\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))+(5d^2\cos(bx+a)^3-2d^2\cos(bx+a))\sqrt{d\cos(bx+a)}\sin(bx+a))}{45b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*sin(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] -2/45*(-3*I*sqrt(2)*d^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4,
0, cos(b*x + a) + I*sin(b*x + a))) + 3*I*sqrt(2)*d^(5/2)*weierstrassZeta(-4
, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) + (5*d^2*co
s(b*x + a)^3 - 2*d^2*cos(b*x + a))*sqrt(d*cos(b*x + a))*sin(b*x + a))/b
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(5/2)*sin(b*x+a)**2,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 4848 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*sin(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(5/2)*sin(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^2 (d \cos(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2*(d*cos(a + b*x))^(5/2),x)

[Out] int(sin(a + b*x)^2*(d*cos(a + b*x))^(5/2), x)

3.197 $\int (d \cos(a + bx))^{3/2} \sin^2(a + bx) dx$

Optimal. Leaf size=98

$$\frac{4d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{21b \sqrt{d \cos(a + bx)}} + \frac{4d \sqrt{d \cos(a + bx)} \sin(a + bx)}{21b} - \frac{2(d \cos(a + bx))^{5/2} \sin(a + bx)}{7bd}$$

[Out] $-2/7*(d*\cos(b*x+a))^{(5/2)*\sin(b*x+a)/b/d+4/21*d^2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)+4/21*d*\sin(b*x+a)*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2720}

$$\frac{4d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{21b \sqrt{d \cos(a + bx)}} - \frac{2 \sin(a + bx) (d \cos(a + bx))^{5/2}}{7bd} + \frac{4d \sin(a + bx) \sqrt{d \cos(a + bx)}}{21b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x]^2, x]$

[Out] $(4*d^2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(21*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (4*d*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(21*b) - (2*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x])/(7*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)]))^{(m_)}, x_Symbol] :> \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^n*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_)])^{(n_)}, x_Symbol] :> \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n - 1)})/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{3/2} \sin^2(a + bx) dx &= -\frac{2(d \cos(a + bx))^{5/2} \sin(a + bx)}{7bd} + \frac{2}{7} \int (d \cos(a + bx))^{3/2} dx \\
&= \frac{4d \sqrt{d \cos(a + bx)} \sin(a + bx)}{21b} - \frac{2(d \cos(a + bx))^{5/2} \sin(a + bx)}{7bd} + \frac{1}{21} \int (d \cos(a + bx))^{3/2} dx \\
&= \frac{4d \sqrt{d \cos(a + bx)} \sin(a + bx)}{21b} - \frac{2(d \cos(a + bx))^{5/2} \sin(a + bx)}{7bd} + \frac{1}{21} \int (d \cos(a + bx))^{3/2} dx \\
&= \frac{4d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{21b \sqrt{d \cos(a + bx)}} + \frac{4d \sqrt{d \cos(a + bx)} \sin(a + bx)}{21b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 57, normalized size = 0.58

$$\frac{(d \cos(a + bx))^{3/2} \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, \frac{3}{2}; \frac{5}{2}; \sin^2(a + bx)\right) \tan^3(a + bx)}{3b}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(3/2)*Sin[a + b*x]^2,x]
```

```
[Out] ((d*Cos[a + b*x])^(3/2)*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, 3/2,
5/2, Sin[a + b*x]^2]*Tan[a + b*x]^3)/(3*b)
```

Maple [A]

time = 0.10, size = 208, normalized size = 2.12

method	result
default	$\frac{4 \sqrt{d} \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d^2 \left(24 \left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 60 \left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 50 \left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 15 \left(\cos^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 3 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{21 \sqrt{-d \left(2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(3/2)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)
[Out] 4/21*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^2*(24*cos(
1/2*b*x+1/2*a)^9-60*cos(1/2*b*x+1/2*a)^7+50*cos(1/2*b*x+1/2*a)^5-15*cos(1/2
*b*x+1/2*a)^3-(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2
)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*
b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*
b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(3/2)*sin(b*x + a)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 90, normalized size = 0.92

$$\frac{2 \left(i \sqrt{2} d^{\frac{3}{2}} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) - i \sqrt{2} d^{\frac{3}{2}} \text{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)) + (3d \cos(bx + a)^2 - 2d) \sqrt{d \cos(bx + a)} \sin(bx + a) \right)}{21 b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] -2/21*(I*sqrt(2)*d^(3/2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*
x + a)) - I*sqrt(2)*d^(3/2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin
(b*x + a)) + (3*d*cos(b*x + a)^2 - 2*d)*sqrt(d*cos(b*x + a))*sin(b*x + a))/
b
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(3/2)*sin(b*x+a)**2,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*sin(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^2 (d \cos(a + bx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2*(d*cos(a + b*x))^(3/2),x)

[Out] int(sin(a + b*x)^2*(d*cos(a + b*x))^(3/2), x)

3.198 $\int \sqrt{d \cos(a + bx)} \sin^2(a + bx) dx$

Optimal. Leaf size=69

$$\frac{4\sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b\sqrt{\cos(a + bx)}} - \frac{2(d \cos(a + bx))^{3/2} \sin(a + bx)}{5bd}$$

[Out] $-2/5*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)/b/d+4/5*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2648, 2721, 2719}

$$\frac{4E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{5b\sqrt{\cos(a + bx)}} - \frac{2 \sin(a + bx)(d \cos(a + bx))^{3/2}}{5bd}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^2,x]`

[Out] `(4*Sqrt[d*Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2])/(5*b*Sqrt[Cos[a + b*x]]) - (2*(d*Cos[a + b*x])^(3/2)*Sin[a + b*x])/(5*b*d)`

Rule 2648

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Ssin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Ssin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*Ssin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
\int \sqrt{d \cos(a + bx)} \sin^2(a + bx) dx &= -\frac{2(d \cos(a + bx))^{3/2} \sin(a + bx)}{5bd} + \frac{2}{5} \int \sqrt{d \cos(a + bx)} dx \\
&= -\frac{2(d \cos(a + bx))^{3/2} \sin(a + bx)}{5bd} + \frac{\left(2\sqrt{d \cos(a + bx)}\right) \int \sqrt{\cos(a + bx)}}{5\sqrt{\cos(a + bx)}} \\
&= \frac{4\sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b\sqrt{\cos(a + bx)}} - \frac{2(d \cos(a + bx))^{3/2} \sin(a + bx)}{5bd}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.06, size = 58, normalized size = 0.84

$$\frac{d^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{3}{2}; \frac{5}{2}; \sin^2(a + bx)\right) \sin^3(a + bx)}{3b\sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^2,x]

[Out] (d*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 3/2, 5/2, Sin[a + b*x]^2]*Sin[a + b*x]^3)/(3*b*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 193 vs. 2(85) = 170.

time = 0.10, size = 194, normalized size = 2.81

method	result
default	$ \frac{4\sqrt{d\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{5\sqrt{-d\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)} d\left(4\left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 8\left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 5\left(\cos^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + \sqrt{\frac{1}{2} - \cos\left(\frac{bx}{2} + \frac{a}{2}\right)}\right) \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{\frac{1}{2} - \cos\left(\frac{bx}{2} + \frac{a}{2}\right)} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*sin(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 4/5*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d*(4*cos(1/2*b*x+1/2*a)^7-8*cos(1/2*b*x+1/2*a)^5+5*cos(1/2*b*x+1/2*a)^3+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))-cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*sin(b*x + a)^2, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.12, size = 87, normalized size = 1.26

$$\frac{2(\sqrt{d \cos(bx+a)} \cos(bx+a) \sin(bx+a) - i\sqrt{2} \sqrt{d} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a))) + i\sqrt{2} \sqrt{d} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a)))}{5b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^2,x, algorithm="fricas")

[Out] $-2/5 * (\sqrt{d \cos(bx+a)} \cos(bx+a) \sin(bx+a) - I \sqrt{2} \sqrt{d} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + I \sin(bx+a))) + I \sqrt{2} \sqrt{d} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - I \sin(bx+a)))) / b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(a + bx)} \sin^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)**2,x)

[Out] Integral(sqrt(d*cos(a + b*x))*sin(a + b*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^2,x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*sin(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^2 \sqrt{d \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2*(d*cos(a + b*x))^(1/2),x)

[Out] int(sin(a + b*x)^2*(d*cos(a + b*x))^(1/2), x)

$$3.199 \quad \int \frac{\sin^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=69

$$\frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3b\sqrt{d \cos(a+bx)}} - \frac{2\sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd}$$

[Out] 4/3*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x), 2^(1/2))*cos(b*x+a)^(1/2)/b/(d*cos(b*x+a))^(1/2)-2/3*sin(b*x+a)*(d*cos(b*x+a))^(1/2)/b/d

Rubi [A]

time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2648, 2721, 2720}

$$\frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3b\sqrt{d \cos(a+bx)}} - \frac{2 \sin(a+bx) \sqrt{d \cos(a+bx)}}{3bd}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2/Sqrt[d*Cos[a + b*x]],x]

[Out] (4*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(3*b*Sqrt[d*Cos[a + b*x]]) - (2*Sqrt[d*Cos[a + b*x]]*Sin[a + b*x])/(3*b*d)

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Ssin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Ssin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*Ssin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\sin^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx &= -\frac{2\sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd} + \frac{2}{3} \int \frac{1}{\sqrt{d \cos(a+bx)}} dx \\ &= -\frac{2\sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd} + \frac{\left(2\sqrt{\cos(a+bx)}\right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{3\sqrt{d \cos(a+bx)}} \\ &= \frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \middle| 2\right)}{3b\sqrt{d \cos(a+bx)}} - \frac{2\sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.07, size = 58, normalized size = 0.84

$$\frac{d \cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{3}{2}; \frac{5}{2}; \sin^2(a+bx)\right) \sin^3(a+bx)}{3b(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^2/Sqrt[d*Cos[a + b*x]], x]

[Out] (d*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, 3/2, 5/2, Sin[a + b*x]^2]*Sin[a + b*x]^3)/(3*b*(d*Cos[a + b*x])^(3/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(85) = 170.

time = 0.15, size = 188, normalized size = 2.72

method	result
default	$\frac{4\sqrt{d} \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \left(2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right) \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}\right)}{3\sqrt{-d} \left(2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right) \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{d}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^2/(d*cos(b*x+a))^(1/2), x, method=_RETURNVERBOSE)

[Out] 4/3*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^4-(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))-sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(b*x + a)^2/sqrt(d*cos(b*x + a)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 78, normalized size = 1.13

$$\frac{2 \left(i \sqrt{2} \sqrt{d} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) + i \sin(bx + a)) - i \sqrt{2} \sqrt{d} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx + a) - i \sin(bx + a)) + \sqrt{d \cos(bx + a)} \sin(bx + a) \right)}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")
```

```
[Out] -2/3*(I*sqrt(2)*sqrt(d)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) - I*sqrt(2)*sqrt(d)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) + sqrt(d*cos(b*x + a))*sin(b*x + a))/(b*d)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^2(a + bx)}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)**2/(d*cos(b*x+a))**(1/2),x)
```

```
[Out] Integral(sin(a + b*x)**2/sqrt(d*cos(a + b*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sin(b*x + a)^2/sqrt(d*cos(b*x + a)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^2}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(a + b*x)^2/(d*cos(a + b*x))^(1/2), x)
```

```
[Out] int(sin(a + b*x)^2/(d*cos(a + b*x))^(1/2), x)
```

$$3.200 \quad \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=68

$$-\frac{4\sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{bd^2 \sqrt{\cos(a+bx)}} + \frac{2 \sin(a+bx)}{bd \sqrt{d \cos(a+bx)}}$$

[Out] $2*\sin(b*x+a)/b/d/(d*\cos(b*x+a))^{(1/2)}-4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/d^{(1/2)}/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2646, 2721, 2719}

$$\frac{2 \sin(a+bx)}{bd \sqrt{d \cos(a+bx)}} - \frac{4E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{d \cos(a+bx)}}{bd^2 \sqrt{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[a + b*x]^2/(d*Cos[a + b*x])^(3/2),x]`

[Out] $(-4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(b*d^2*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(b*d*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2646

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
\int \frac{\sin^2(a+bx)}{(d\cos(a+bx))^{3/2}} dx &= \frac{2\sin(a+bx)}{bd\sqrt{d\cos(a+bx)}} - \frac{2\int\sqrt{d\cos(a+bx)} dx}{d^2} \\
&= \frac{2\sin(a+bx)}{bd\sqrt{d\cos(a+bx)}} - \frac{\left(2\sqrt{d\cos(a+bx)}\right)\int\sqrt{\cos(a+bx)} dx}{d^2\sqrt{\cos(a+bx)}} \\
&= -\frac{4\sqrt{d\cos(a+bx)} E\left(\frac{1}{2}(a+bx)\middle|2\right)}{bd^2\sqrt{\cos(a+bx)}} + \frac{2\sin(a+bx)}{bd\sqrt{d\cos(a+bx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.06, size = 60, normalized size = 0.88

$$\frac{\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{5}{4}, \frac{3}{2}, \frac{5}{2}; \sin^2(a+bx)\right) \sin^3(a+bx)}{3bd\sqrt{d\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^2/(d*Cos[a + b*x])^(3/2), x]

[Out] ((Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[5/4, 3/2, 5/2, Sin[a + b*x]^2]*Sin[a + b*x]^3)/(3*b*d*Sqrt[d*Cos[a + b*x]])

Maple [A]

time = 0.13, size = 168, normalized size = 2.47

method	result
default	$ -\frac{4\sqrt{-2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d\left(\sqrt{2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}\sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}\right) \text{EllipticE}\left(\frac{bx}{2} + \frac{a}{2}\right)}{d\sqrt{-d\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)} \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{d\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^2/(d*cos(b*x+a))^(3/2), x, method=_RETURNVERBOSE)

[Out] -4/d*(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(1/2)*((2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))-sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 105, normalized size = 1.54

$$\frac{2 \left(i \sqrt{2} \sqrt{d} \cos(bx+a) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a))) - i \sqrt{2} \sqrt{d} \cos(bx+a) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a))) - \sqrt{d \cos(bx+a)} \sin(bx+a) \right)}{bd^2 \cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] -2*(I*sqrt(2)*sqrt(d)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - I*sqrt(2)*sqrt(d)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) - sqrt(d*cos(b*x + a))*sin(b*x + a))/(b*d^2*cos(b*x + a))

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**2/(d*cos(b*x+a))**(3/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^2}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/(d*cos(a + b*x))^(3/2),x)

[Out] int(sin(a + b*x)^2/(d*cos(a + b*x))^(3/2), x)

$$3.201 \quad \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=72

$$-\frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}}$$

[Out] 2/3*sin(b*x+a)/b/d/(d*cos(b*x+a))^(3/2)-4/3*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)/b/d^2/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.05, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2646, 2721, 2720}

$$\frac{2 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^2/(d*Cos[a + b*x])^(5/2),x]

[Out] (-4*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(3*b*d^2*Sqrt[d*Cos[a + b*x]]) + (2*Sin[a + b*x])/(3*b*d*(d*Cos[a + b*x])^(3/2))

Rule 2646

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2720

Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx &= \frac{2 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{2 \int \frac{1}{\sqrt{d \cos(a+bx)}} dx}{3d^2} \\
&= \frac{2 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{\left(2 \sqrt{\cos(a+bx)}\right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{3d^2 \sqrt{d \cos(a+bx)}} \\
&= -\frac{4 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}} + \frac{2 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.06, size = 60, normalized size = 0.83

$$\frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{2}, \frac{7}{4}; \frac{5}{2}; \sin^2(a+bx)\right) \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^2/(d*Cos[a + b*x])^(5/2), x]

[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/2, 7/4, 5/2, Sin[a + b*x]^2]*Sin[a + b*x]^3)/(3*b*d*(d*Cos[a + b*x])^(3/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 241 vs. 2(88) = 176.

time = 0.12, size = 242, normalized size = 3.36

method	result
default	$ -\frac{4 \left(2 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \operatorname{EllipticF} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right)}{3d^2 \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \sqrt{-d \left(2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right)}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^2/(d*cos(b*x+a))^(5/2), x, method=_RETURNVERBOSE)

[Out] -4/3*(2*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))*sin(1/2*b*x+1/2*a)^2-(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))-sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/d^2*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(2*cos(1/2*b*x+1/2*a)^2-1)/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(5/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 103, normalized size = 1.43

$$\frac{2 \left(-i \sqrt{2} \sqrt{d} \cos(bx+a)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a)) + i \sqrt{2} \sqrt{d} \cos(bx+a)^2 \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a)) - \sqrt{d \cos(bx+a)} \sin(bx+a) \right)}{3bd^3 \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")
```

```
[Out] -2/3*(-I*sqrt(2)*sqrt(d)*cos(b*x + a)^2*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*sqrt(d)*cos(b*x + a)^2*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) - sqrt(d*cos(b*x + a))*sin(b*x + a))/ (b*d^3*cos(b*x + a)^2)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)**2/(d*cos(b*x+a))**(5/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3880 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(5/2),x, algorithm="giac")
```

```
[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(5/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^2}{(d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(a + b*x)^2/(d*cos(a + b*x))^(5/2),x)
```

```
[Out] int(sin(a + b*x)^2/(d*cos(a + b*x))^(5/2), x)
```

$$3.202 \quad \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=100

$$\frac{4\sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^4 \sqrt{\cos(a+bx)}} + \frac{2 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{4 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}}$$

[Out] $2/5*\sin(b*x+a)/b/d/(d*\cos(b*x+a))^{(5/2)}-4/5*\sin(b*x+a)/b/d^3/(d*\cos(b*x+a))^{(1/2)}+4/5*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/d^4/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2646, 2716, 2721, 2719}

$$\frac{4E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{d \cos(a+bx)}}{5bd^4 \sqrt{\cos(a+bx)}} - \frac{4 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[a + b*x]^2/(d*\text{Cos}[a + b*x])^{(7/2)}, x]$

[Out] $(4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*d^4*\text{Sqrt}[\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(5*b*d*(d*\text{Cos}[a + b*x])^{(5/2)}) - (4*\text{Sin}[a + b*x])/(5*b*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\text{Sin}[e + f*x])^{(m-1)}*((b*\text{Cos}[e + f*x])^{(n+1)})/(b*f*(n+1)), x] + \text{Dist}[a^2*((m-1)/(b^2*(n+1))), \text{Int}[(a*\text{Sin}[e + f*x])^{(m-2)}*(b*\text{Cos}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] \parallel \text{EqQ}[m + n, 0])$

Rule 2716

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_)])^{(n_)}], x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{7/2}} dx &= \frac{2 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{2 \int \frac{1}{(d \cos(a+bx))^{3/2}} dx}{5d^2} \\ &= \frac{2 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{4 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \int \sqrt{d \cos(a+bx)} dx}{5d^4} \\ &= \frac{2 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{4 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{\left(2 \sqrt{d \cos(a+bx)}\right) \int \sqrt{\cos(a+bx)} dx}{5d^4 \sqrt{\cos(a+bx)}} \\ &= \frac{4 \sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^4 \sqrt{\cos(a+bx)}} + \frac{2 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{4 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 59, normalized size = 0.59

$$\frac{\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{3}{2}, \frac{9}{4}; \frac{5}{2}; \sin^2(a+bx)\right) \sin^3(2(a+bx))}{24b(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^2/(d*Cos[a + b*x])^(7/2), x]
```

```
[Out] ((Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[3/2, 9/4, 5/2, Sin[a + b*x]^2]*Sin[2*(a + b*x)]^3)/(24*b*(d*Cos[a + b*x])^(7/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 364 vs. 2(112) = 224.

time = 0.20, size = 365, normalized size = 3.65

method	result
default	$\frac{4 \sqrt{d} \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{24 b (d \cos(a+bx))^{7/2}} \left(4 \operatorname{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \sqrt{2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^2/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

[Out]
$$-4/5*(d*(2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/d^4/\sin(1/2*b*x+1/2*a)^3/(8*\sin(1/2*b*x+1/2*a)^6-12*\sin(1/2*b*x+1/2*a)^4+6*\sin(1/2*b*x+1/2*a)^2-1)*(4*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)})*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\sin(1/2*b*x+1/2*a)^4-8*\sin(1/2*b*x+1/2*a)^6*\cos(1/2*b*x+1/2*a)-4*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)})*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\sin(1/2*b*x+1/2*a)^2+8*\cos(1/2*b*x+1/2*a)*\sin(1/2*b*x+1/2*a)^4+(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)})-\sin(1/2*b*x+1/2*a)^2*\cos(1/2*b*x+1/2*a))*(-2*\sin(1/2*b*x+1/2*a)^4*d+\sin(1/2*b*x+1/2*a)^2*d)^{(1/2)}/(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}/b$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")`

[Out] `integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(7/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 120, normalized size = 1.20

$$\frac{2(-i\sqrt{2}\sqrt{d}\cos(bx+a)^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+i\sqrt{2}\sqrt{d}\cos(bx+a)^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))+\sqrt{d}\cos(bx+a)(2\cos(bx+a)^2-1)\sin(bx+a))}{5bd^4\cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")`

[Out]
$$-2/5*(-I*\sqrt{2}*\sqrt{d}*\cos(b*x + a)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a))) + I*\sqrt{2}*\sqrt{d}*\cos(b*x + a)^3*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a))) + \sqrt{d}*\cos(b*x + a)*(2*\cos(b*x + a)^2 - 1)*\sin(b*x + a))/(b*d^4*\cos(b*x + a)^3)$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**2/(d*cos(b*x+a))**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^2}{(d \cos(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/(d*cos(a + b*x))^(7/2),x)

[Out] int(sin(a + b*x)^2/(d*cos(a + b*x))^(7/2), x)

$$3.203 \quad \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=100

$$-\frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{21bd^4 \sqrt{d \cos(a+bx)}} + \frac{2 \sin(a+bx)}{7bd(d \cos(a+bx))^{7/2}} - \frac{4 \sin(a+bx)}{21bd^3(d \cos(a+bx))^{3/2}}$$

[Out] $2/7*\sin(b*x+a)/b/d/(d*\cos(b*x+a))^{(7/2)}-4/21*\sin(b*x+a)/b/d^3/(d*\cos(b*x+a))^{(3/2)}-4/21*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/d^4/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2646, 2716, 2721, 2720}

$$-\frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{21bd^4 \sqrt{d \cos(a+bx)}} - \frac{4 \sin(a+bx)}{21bd^3(d \cos(a+bx))^{3/2}} + \frac{2 \sin(a+bx)}{7bd(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[a + b*x]^2/(d*\text{Cos}[a + b*x])^{(9/2)}, x]$

[Out] $(-4*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(21*b*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x])/(7*b*d*(d*\text{Cos}[a + b*x])^{(7/2)}) - (4*\text{Sin}[a + b*x])/(21*b*d^3*(d*\text{Cos}[a + b*x])^{(3/2)})$

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.))^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\text{Sin}[e + f*x])^{(m-1)}*((b*\text{Cos}[e + f*x])^{(n+1)})/(b*f*(n+1)), x] + \text{Dist}[a^2*((m-1)/(b^2*(n+1))), \text{Int}[(a*\text{Sin}[e + f*x])^{(m-2)}*(b*\text{Cos}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{IntegersQ}[2*m, 2*n] \ || \ \text{EqQ}[m + n, 0])$

Rule 2716

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n+1)})/(b*d*(n+1)), x] + \text{Dist}[(n+2)/(b^2*(n+1)), \text{Int}[(b*\text{Sin}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{9/2}} dx &= \frac{2 \sin(a+bx)}{7bd(d \cos(a+bx))^{7/2}} - \frac{2 \int \frac{1}{(d \cos(a+bx))^{5/2}} dx}{7d^2} \\ &= \frac{2 \sin(a+bx)}{7bd(d \cos(a+bx))^{7/2}} - \frac{4 \sin(a+bx)}{21bd^3(d \cos(a+bx))^{3/2}} - \frac{2 \int \frac{1}{\sqrt{d \cos(a+bx)}} dx}{21d^4} \\ &= \frac{2 \sin(a+bx)}{7bd(d \cos(a+bx))^{7/2}} - \frac{4 \sin(a+bx)}{21bd^3(d \cos(a+bx))^{3/2}} - \frac{\left(2\sqrt{\cos(a+bx)}\right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{21d^4 \sqrt{d \cos(a+bx)}} \\ &= -\frac{4\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{21bd^4 \sqrt{d \cos(a+bx)}} + \frac{2 \sin(a+bx)}{7bd(d \cos(a+bx))^{7/2}} - \frac{4 \sin(a+bx)}{21bd^3(d \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.04, size = 59, normalized size = 0.59

$$\frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{2}, \frac{11}{4}; \frac{5}{2}; \sin^2(a+bx)\right) \sin^3(2(a+bx))}{24b(d \cos(a+bx))^{9/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^2/(d*Cos[a + b*x])^(9/2), x]
```

```
[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/2, 11/4, 5/2, Sin[a + b*x]^2]*S
in[2*(a + b*x)]^3)/(24*b*(d*Cos[a + b*x])^(9/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 395 vs. 2(112) = 224.

time = 0.14, size = 396, normalized size = 3.96

method	result
default	$4 \left(-8 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \operatorname{EllipticF} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 12 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x+a)^2/(d*cos(b*x+a))^(9/2),x,method=_RETURNVERBOSE)
```

```
[Out] 4/21*(-8*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))*sin(1/2*b*x+1/2*a)^6+12*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))*sin(1/2*b*x+1/2*a)^4-8*sin(1/2*b*x+1/2*a)^6*cos(1/2*b*x+1/2*a)-6*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))*sin(1/2*b*x+1/2*a)^2+8*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^4+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/d^4*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/(2*cos(1/2*b*x+1/2*a)^2-1)^3/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(9/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(9/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 114, normalized size = 1.14

$$\frac{2(-i\sqrt{2}\sqrt{d}\cos(bx+a)^4\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))+i\sqrt{2}\sqrt{d}\cos(bx+a)^4\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))+\sqrt{d}\cos(bx+a)(2\cos(bx+a)^2-3)\sin(bx+a))}{21bd^5\cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")
```

```
[Out] -2/21*(-I*sqrt(2)*sqrt(d)*cos(b*x + a)^4*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + I*sqrt(2)*sqrt(d)*cos(b*x + a)^4*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) + sqrt(d*cos(b*x + a))*(2*cos(b*x + a)^2 - 3)*sin(b*x + a))/(b*d^5*cos(b*x + a)^4)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**2/(d*cos(b*x+a))**(9/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4847 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^2/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^2/(d*cos(b*x + a))^(9/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^2}{(d \cos(a + bx))^{9/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2/(d*cos(a + b*x))^(9/2),x)

[Out] int(sin(a + b*x)^2/(d*cos(a + b*x))^(9/2), x)

3.204 $\int \sqrt{d \cos(a + bx)} \sin^3(a + bx) dx$

Optimal. Leaf size=45

$$-\frac{2(d \cos(a + bx))^{3/2}}{3bd} + \frac{2(d \cos(a + bx))^{7/2}}{7bd^3}$$

[Out] $-2/3*(d*\cos(b*x+a))^(3/2)/b/d+2/7*(d*\cos(b*x+a))^(7/2)/b/d^3$

Rubi [A]

time = 0.03, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2(d \cos(a + bx))^{7/2}}{7bd^3} - \frac{2(d \cos(a + bx))^{3/2}}{3bd}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^3,x]`

[Out] $(-2*(d*\text{Cos}[a + b*x])^(3/2))/(3*b*d) + (2*(d*\text{Cos}[a + b*x])^(7/2))/(7*b*d^3)$

Rule 14

`Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]]`

Rule 2645

`Int[(cos[(e_) + (f_.)*(x_)]*(a_.))^(m_)*sin[(e_) + (f_.)*(x_)]^(n_), x_Symbol] :> Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])]`

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} \sin^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \sqrt{x} \left(1 - \frac{x^2}{d^2}\right) dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\sqrt{x} - \frac{x^{5/2}}{d^2}\right) dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{2(d \cos(a + bx))^{3/2}}{3bd} + \frac{2(d \cos(a + bx))^{7/2}}{7bd^3} \end{aligned}$$

Mathematica [A]

time = 0.22, size = 57, normalized size = 1.27

$$\frac{d \left(16 \cos^2(a + bx) - 16 \sqrt{\cos^2(a + bx)} + 3 \sin^2(2(a + bx)) \right)}{42b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^3,x]

[Out] -1/42*(d*(16*Cos[a + b*x]^2 - 16*(Cos[a + b*x]^2)^(1/4) + 3*Sin[2*(a + b*x)]^2))/(b*Sqrt[d*Cos[a + b*x]])

Maple [A]

time = 0.08, size = 63, normalized size = 1.40

method	result	size
default	$\frac{8 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d \left(6 \left(\sin^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 9 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + \sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) + 1 \right)}{21b}$	63

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] -8/21*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(6*sin(1/2*b*x+1/2*a)^6-9*sin(1/2*b*x+1/2*a)^4+sin(1/2*b*x+1/2*a)^2+1)/b

Maxima [A]

time = 0.29, size = 36, normalized size = 0.80

$$\frac{2 \left(3 (d \cos(bx + a))^{\frac{7}{2}} - 7 (d \cos(bx + a))^{\frac{3}{2}} d^2 \right)}{21 b d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^3,x, algorithm="maxima")

[Out] 2/21*(3*(d*cos(b*x + a))^(7/2) - 7*(d*cos(b*x + a))^(3/2)*d^2)/(b*d^3)

Fricas [A]

time = 0.37, size = 34, normalized size = 0.76

$$\frac{2 \left(3 \cos(bx + a)^3 - 7 \cos(bx + a) \right) \sqrt{d \cos(bx + a)}}{21 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^3,x, algorithm="fricas")

[Out] $2/21*(3*\cos(b*x + a)^3 - 7*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}/b$

Sympy [A]

time = 1.20, size = 73, normalized size = 1.62

$$\begin{cases} -\frac{2\sqrt{d\cos(a+bx)}\sin^2(a+bx)\cos(a+bx)}{3b} - \frac{8\sqrt{d\cos(a+bx)}\cos^3(a+bx)}{21b} & \text{for } b \neq 0 \\ x\sqrt{d\cos(a)}\sin^3(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(1/2)*sin(b*x+a)**3,x)`

[Out] `Piecewise((-2*sqrt(d*cos(a + b*x))*sin(a + b*x)**2*cos(a + b*x)/(3*b) - 8*sqrt(d*cos(a + b*x))*cos(a + b*x)**3/(21*b), Ne(b, 0)), (x*sqrt(d*cos(a))*sin(a)**3, True))`

Giac [A]

time = 5.66, size = 53, normalized size = 1.18

$$\frac{2 \left(3 \sqrt{d \cos(bx + a)} d^3 \cos(bx + a)^3 - 7 \sqrt{d \cos(bx + a)} d^3 \cos(bx + a) \right)}{21 b d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^3,x, algorithm="giac")`

[Out] $2/21*(3*\sqrt{d*\cos(b*x + a)}*d^3*\cos(b*x + a)^3 - 7*\sqrt{d*\cos(b*x + a)}*d^3*\cos(b*x + a))/(b*d^3)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(a + bx)^3 \sqrt{d \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^3*(d*cos(a + b*x))^(1/2),x)`

[Out] `int(sin(a + b*x)^3*(d*cos(a + b*x))^(1/2), x)`

$$3.205 \quad \int \frac{\sin^3(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=43

$$-\frac{2\sqrt{d \cos(a+bx)}}{bd} + \frac{2(d \cos(a+bx))^{5/2}}{5bd^3}$$

[Out] $2/5*(d*\cos(b*x+a))^{(5/2)}/b/d^3-2*(d*\cos(b*x+a))^{(1/2)}/b/d$

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2(d \cos(a+bx))^{5/2}}{5bd^3} - \frac{2\sqrt{d \cos(a+bx)}}{bd}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3/Sqrt[d*Cos[a + b*x]],x]

[Out] $(-2*\text{Sqrt}[d*\text{Cos}[a + b*x]])/(b*d) + (2*(d*\text{Cos}[a + b*x])^{(5/2)})/(5*b*d^3)$

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

Int[(cos[(e_)+(f_)*(x_)]*(a_))^(m_)*sin[(e_)+(f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1-x^2/a^2)^((n-1)/2), x], x, a*Cos[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(a+bx)}{\sqrt{d \cos(a+bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{\sqrt{x}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{\sqrt{x}} - \frac{x^{3/2}}{d^2}\right) dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{2\sqrt{d \cos(a+bx)}}{bd} + \frac{2(d \cos(a+bx))^{5/2}}{5bd^3} \end{aligned}$$

Mathematica [A]

time = 0.13, size = 57, normalized size = 1.33

$$\frac{\cos(a + bx)(-9 + \cos(2(a + bx))) + 8 \cos^2(a + bx)^{3/4} \sec(a + bx)}{5b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^3/Sqrt[d*Cos[a + b*x]], x]`

```
[Out] (Cos[a + b*x]*(-9 + Cos[2*(a + b*x)]) + 8*(Cos[a + b*x]^2)^(3/4)*Sec[a + b*x])/
(5*b*Sqrt[d*Cos[a + b*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 91 vs. 2(37) = 74.

time = 0.06, size = 92, normalized size = 2.14

method	result
default	$\frac{8 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 8 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 8 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{5db}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^3/(d*cos(b*x+a))^(1/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/5*(8*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*sin(1/2*b*x+1/2*a)^4-8*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*sin(1/2*b*x+1/2*a)^2-8*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2))/d/b
```

Maxima [A]

time = 0.29, size = 36, normalized size = 0.84

$$\frac{2 \left(5 \sqrt{d \cos(bx + a)} - \frac{(d \cos(bx + a))^{5/2}}{d^2} \right)}{5bd}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(1/2), x, algorithm="maxima")`

```
[Out] -2/5*(5*sqrt(d*cos(b*x + a)) - (d*cos(b*x + a))^(5/2)/d^2)/(b*d)
```

Fricas [A]

time = 0.37, size = 28, normalized size = 0.65

$$\frac{2 \sqrt{d \cos(bx + a)} (\cos(bx + a)^2 - 5)}{5bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] 2/5*sqrt(d*cos(b*x + a))*(cos(b*x + a)^2 - 5)/(b*d)

Sympy [A]

time = 1.19, size = 71, normalized size = 1.65

$$\begin{cases} \frac{2 \sin^2(a+bx) \cos(a+bx)}{b \sqrt{d \cos(a+bx)}} - \frac{8 \cos^3(a+bx)}{5b \sqrt{d \cos(a+bx)}} & \text{for } b \neq 0 \\ \frac{x \sin^3(a)}{\sqrt{d \cos(a)}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**3/(d*cos(b*x+a))**(1/2),x)

[Out] Piecewise((-2*sin(a + b*x)**2*cos(a + b*x)/(b*sqrt(d*cos(a + b*x))) - 8*cos(a + b*x)**3/(5*b*sqrt(d*cos(a + b*x))), Ne(b, 0)), (x*sin(a)**3/sqrt(d*cos(a)), True))

Giac [A]

time = 5.62, size = 46, normalized size = 1.07

$$\frac{2 \left(\sqrt{d \cos(bx + a)} d^2 \cos(bx + a)^2 - 5 \sqrt{d \cos(bx + a)} d^2 \right)}{5 b d^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] 2/5*(sqrt(d*cos(b*x + a))*d^2*cos(b*x + a)^2 - 5*sqrt(d*cos(b*x + a))*d^2)/(b*d^3)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(a + bx)^3}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/(d*cos(a + b*x))^(1/2),x)

[Out] int(sin(a + b*x)^3/(d*cos(a + b*x))^(1/2), x)

$$3.206 \quad \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=43

$$\frac{2}{bd\sqrt{d \cos(a+bx)}} + \frac{2(d \cos(a+bx))^{3/2}}{3bd^3}$$

[Out] 2/3*(d*cos(b*x+a))^(3/2)/b/d^3+2/b/d/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2(d \cos(a+bx))^{3/2}}{3bd^3} + \frac{2}{bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3/(d*Cos[a + b*x])^(3/2), x]

[Out] 2/(b*d*Sqrt[d*Cos[a + b*x]]) + (2*(d*Cos[a + b*x])^(3/2))/(3*b*d^3)

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

Int[(cos[(e_)+(f_)*(x_)]*(a_))^(m_)*sin[(e_)+(f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1-x^2/a^2)^((n-1)/2), x], x, a*Cos[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{3/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^{3/2}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^{3/2}} - \frac{\sqrt{x}}{d^2}\right) dx, x, d \cos(a+bx)\right)}{bd} \\ &= \frac{2}{bd\sqrt{d \cos(a+bx)}} + \frac{2(d \cos(a+bx))^{3/2}}{3bd^3} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 46, normalized size = 1.07

$$\frac{2\left(-4 + 4\sqrt[4]{\cos^2(a + bx)} + \sin^2(a + bx)\right)}{3bd\sqrt{d\cos(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[a + b*x]^3/(d*Cos[a + b*x])^(3/2), x]``[Out] (-2*(-4 + 4*(Cos[a + b*x]^2)^(1/4) + Sin[a + b*x]^2))/(3*b*d*Sqrt[d*Cos[a + b*x]])`**Maple [A]**

time = 0.21, size = 70, normalized size = 1.63

method	result	size
default	$-\frac{8\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + d\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)}{3d^2\left(2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)b}$	70

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(b*x+a)^3/(d*cos(b*x+a))^(3/2), x, method=_RETURNVERBOSE)``[Out] -8/3/d^2*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2+1)/(2*sin(1/2*b*x+1/2*a)^2-1)/b`**Maxima [A]**

time = 0.29, size = 35, normalized size = 0.81

$$\frac{2\left(\frac{3}{\sqrt{d\cos(bx+a)}} + \frac{(d\cos(bx+a))^{\frac{3}{2}}}{d^2}\right)}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(3/2), x, algorithm="maxima")``[Out] 2/3*(3/sqrt(d*cos(b*x + a)) + (d*cos(b*x + a))^(3/2)/d^2)/(b*d)`**Fricas [A]**

time = 0.36, size = 36, normalized size = 0.84

$$\frac{2\sqrt{d\cos(bx+a)}(\cos(bx+a)^2 + 3)}{3bd^2\cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] 2/3*sqrt(d*cos(b*x + a))*(cos(b*x + a)^2 + 3)/(b*d^2*cos(b*x + a))

Sympy [A]

time = 2.43, size = 70, normalized size = 1.63

$$\begin{cases} \frac{2 \sin^2(a+bx) \cos(a+bx)}{b(d \cos(a+bx))^{\frac{3}{2}}} + \frac{8 \cos^3(a+bx)}{3b(d \cos(a+bx))^{\frac{3}{2}}} & \text{for } b \neq 0 \\ \frac{x \sin^3(a)}{(d \cos(a))^{\frac{3}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**3/(d*cos(b*x+a))**(3/2),x)

[Out] Piecewise((2*sin(a + b*x)**2*cos(a + b*x)/(b*(d*cos(a + b*x))**(3/2)) + 8*cos(a + b*x)**3/(3*b*(d*cos(a + b*x))**(3/2)), Ne(b, 0)), (x*sin(a)**3/(d*cos(a))**(3/2), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^3/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(a + bx)^3}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/(d*cos(a + b*x))^(3/2),x)

[Out] int(sin(a + b*x)^3/(d*cos(a + b*x))^(3/2), x)

$$3.207 \quad \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=43

$$\frac{2}{3bd(d \cos(a+bx))^{3/2}} + \frac{2\sqrt{d \cos(a+bx)}}{bd^3}$$

[Out] 2/3/b/d/(d*cos(b*x+a))^(3/2)+2*(d*cos(b*x+a))^(1/2)/b/d^3

Rubi [A]

time = 0.04, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2\sqrt{d \cos(a+bx)}}{bd^3} + \frac{2}{3bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3/(d*Cos[a + b*x])^(5/2),x]

[Out] 2/(3*b*d*(d*Cos[a + b*x])^(3/2)) + (2*sqrt[d*Cos[a + b*x]])/(b*d^3)

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

Int[(cos[(e_)+(f_)*(x_)]*(a_))^(m_)*sin[(e_)+(f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1-x^2/a^2)^((n-1)/2), x], x, a*Cos[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{5/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^{5/2}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^{5/2}} - \frac{1}{d^2 \sqrt{x}}\right) dx, x, d \cos(a+bx)\right)}{bd} \\ &= \frac{2}{3bd(d \cos(a+bx))^{3/2}} + \frac{2\sqrt{d \cos(a+bx)}}{bd^3} \end{aligned}$$

Mathematica [A]

time = 0.07, size = 48, normalized size = 1.12

$$\frac{2(-4 + 4 \cos^2(a + bx)^{3/4} + 3 \sin^2(a + bx))}{3bd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^3/(d*Cos[a + b*x])^(5/2), x]**[Out]** (-2*(-4 + 4*(Cos[a + b*x]^2)^(3/4) + 3*Sin[a + b*x]^2))/(3*b*d*(d*Cos[a + b*x])^(3/2))**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 84 vs. 2(37) = 74.

time = 0.21, size = 85, normalized size = 1.98

method	result	size
default	$\frac{8 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d \left(3 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 3 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 1 \right)}{3d^3 \left(4 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 4 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 1 \right) b}$	85

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^3/(d*cos(b*x+a))^(5/2), x, method=_RETURNVERBOSE)**[Out]** 8/3/d^3/(4*sin(1/2*b*x+1/2*a)^4-4*sin(1/2*b*x+1/2*a)^2+1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(3*sin(1/2*b*x+1/2*a)^4-3*sin(1/2*b*x+1/2*a)^2+1)/b**Maxima [A]**

time = 0.32, size = 34, normalized size = 0.79

$$\frac{2 \left(\frac{1}{(d \cos(bx+a))^{3/2}} + \frac{3 \sqrt{d \cos(bx+a)}}{d^2} \right)}{3bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(5/2), x, algorithm="maxima")**[Out]** 2/3*(1/(d*cos(b*x + a))^(3/2) + 3*sqrt(d*cos(b*x + a))/d^2)/(b*d)**Fricas [A]**

time = 0.36, size = 38, normalized size = 0.88

$$\frac{2 \sqrt{d \cos(bx+a)} (3 \cos(bx+a)^2 + 1)}{3bd^3 \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")

[Out] 2/3*sqrt(d*cos(b*x + a))*(3*cos(b*x + a)^2 + 1)/(b*d^3*cos(b*x + a)^2)

Sympy [A]

time = 5.30, size = 71, normalized size = 1.65

$$\begin{cases} \frac{2 \sin^2(a+bx) \cos(a+bx)}{3b(d \cos(a+bx))^{\frac{5}{2}}} + \frac{8 \cos^3(a+bx)}{3b(d \cos(a+bx))^{\frac{5}{2}}} & \text{for } b \neq 0 \\ \frac{x \sin^3(a)}{(d \cos(a))^{\frac{5}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**3/(d*cos(b*x+a))**(5/2),x)

[Out] Piecewise((2*sin(a + b*x)**2*cos(a + b*x)/(3*b*(d*cos(a + b*x))**(5/2)) + 8*cos(a + b*x)**3/(3*b*(d*cos(a + b*x))**(5/2)), Ne(b, 0)), (x*sin(a)**3/(d*cos(a))**(5/2), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^3/(d*cos(b*x + a))^(5/2), x)

Mupad [B]

time = 1.06, size = 66, normalized size = 1.53

$$\frac{2 \sqrt{d \cos(a + bx)} (16 \cos(2a + 2bx) + 3 \cos(4a + 4bx) + 13)}{3 b d^3 (4 \cos(2a + 2bx) + \cos(4a + 4bx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/(d*cos(a + b*x))^(5/2),x)

[Out] (2*(d*cos(a + b*x))^(1/2)*(16*cos(2*a + 2*b*x) + 3*cos(4*a + 4*b*x) + 13))/(3*b*d^3*(4*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 3))

$$3.208 \quad \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=43

$$\frac{2}{5bd(d \cos(a+bx))^{5/2}} - \frac{2}{bd^3 \sqrt{d \cos(a+bx)}}$$

[Out] 2/5/b/d/(d*cos(b*x+a))^(5/2)-2/b/d^3/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2}{5bd(d \cos(a+bx))^{5/2}} - \frac{2}{bd^3 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3/(d*Cos[a + b*x])^(7/2), x]

[Out] 2/(5*b*d*(d*Cos[a + b*x])^(5/2)) - 2/(b*d^3*Sqrt[d*Cos[a + b*x]])

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] :> Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n-1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{7/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^{7/2}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^{7/2}} - \frac{1}{d^2 x^{3/2}}\right) dx, x, d \cos(a+bx)\right)}{bd} \\ &= \frac{2}{5bd(d \cos(a+bx))^{5/2}} - \frac{2}{bd^3 \sqrt{d \cos(a+bx)}} \end{aligned}$$

Mathematica [A]

time = 0.18, size = 70, normalized size = 1.63

$$\frac{2\left(5 - 4\sqrt[4]{\cos^2(a + bx)} + 4\left(-1 + \sqrt[4]{\cos^2(a + bx)}\right) \csc^2(a + bx)\right) \tan^2(a + bx)}{5bd^3 \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^3/(d*Cos[a + b*x])^(7/2), x]
```

```
[Out] (2*(5 - 4*(Cos[a + b*x]^2)^(1/4) + 4*(-1 + (Cos[a + b*x]^2)^(1/4))*Csc[a + b*x]^2)*Tan[a + b*x]^2)/(5*b*d^3*Sqrt[d*Cos[a + b*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 97 vs. 2(37) = 74.

time = 0.21, size = 98, normalized size = 2.28

method	result	size
default	$\frac{8\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + d^2\left(5\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 5\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)}{5d^4\left(8\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 12\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 6\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)b}$	98

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x+a)^3/(d*cos(b*x+a))^(7/2), x, method=_RETURNVERBOSE)
```

```
[Out] 8/5/d^4/(8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(5*sin(1/2*b*x+1/2*a)^4-5*sin(1/2*b*x+1/2*a)^2+1)/b
```

Maxima [A]

time = 0.29, size = 37, normalized size = 0.86

$$\frac{2(5d^2 \cos(bx + a)^2 - d^2)}{5(d \cos(bx + a))^{\frac{5}{2}} bd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(7/2), x, algorithm="maxima")
```

```
[Out] -2/5*(5*d^2*cos(b*x + a)^2 - d^2)/((d*cos(b*x + a))^(5/2)*b*d^3)
```

Fricas [A]

time = 0.39, size = 38, normalized size = 0.88

$$\frac{2\sqrt{d \cos(bx + a)} (5 \cos(bx + a)^2 - 1)}{5bd^4 \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")`

[Out] $-2/5\sqrt{d\cos(bx+a)}(5\cos(bx+a)^2-1)/(b*d^4\cos(bx+a)^3)$

Sympy [A]

time = 40.73, size = 71, normalized size = 1.65

$$\begin{cases} \frac{2\sin^2(a+bx)\cos(a+bx)}{5b(d\cos(a+bx))^{\frac{7}{2}}} - \frac{8\cos^3(a+bx)}{5b(d\cos(a+bx))^{\frac{7}{2}}} & \text{for } b \neq 0 \\ \frac{x\sin^3(a)}{(d\cos(a))^{\frac{7}{2}}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**3/(d*cos(b*x+a))**(7/2),x)`

[Out] `Piecewise((2*sin(a + b*x)**2*cos(a + b*x)/(5*b*(d*cos(a + b*x))**(7/2)) - 8*cos(a + b*x)**3/(5*b*(d*cos(a + b*x))**(7/2)), Ne(b, 0)), (x*sin(a)**3/(d*cos(a))**(7/2), True))`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(7/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^3/(d*cos(b*x + a))^(7/2), x)`

Mupad [B]

time = 3.67, size = 93, normalized size = 2.16

$$\frac{4e^{a\operatorname{li}+bx\operatorname{li}}\sqrt{d\left(\frac{e^{-a\operatorname{li}-bx\operatorname{li}}}{2}+\frac{e^{a\operatorname{li}+bx\operatorname{li}}}{2}\right)}(6e^{a2i+bx2i}+5e^{a4i+bx4i}+5)}{5bd^4(e^{a2i+bx2i}+1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^3/(d*cos(a + b*x))^(7/2),x)`

[Out] $-(4\exp(a\operatorname{li}+bx\operatorname{li})*(d(\exp(-a\operatorname{li}-bx\operatorname{li})/2+\exp(a\operatorname{li}+bx\operatorname{li})/2))^{1/2}*(6\exp(a2i+bx2i)+5\exp(a4i+bx4i)+5))/(5b*d^4*(\exp(a2i+bx2i)+1)^3)$

$$3.209 \quad \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=45

$$\frac{2}{7bd(d \cos(a+bx))^{7/2}} - \frac{2}{3bd^3(d \cos(a+bx))^{3/2}}$$

[Out] 2/7/b/d/(d*cos(b*x+a))^(7/2)-2/3/b/d^3/(d*cos(b*x+a))^(3/2)

Rubi [A]

time = 0.03, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2}{7bd(d \cos(a+bx))^{7/2}} - \frac{2}{3bd^3(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3/(d*Cos[a + b*x])^(9/2), x]

[Out] 2/(7*b*d*(d*Cos[a + b*x])^(7/2)) - 2/(3*b*d^3*(d*Cos[a + b*x])^(3/2))

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n-1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{9/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^{9/2}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^{9/2}} - \frac{1}{d^2 x^{5/2}}\right) dx, x, d \cos(a+bx)\right)}{bd} \\ &= \frac{2}{7bd(d \cos(a+bx))^{7/2}} - \frac{2}{3bd^3(d \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.22, size = 70, normalized size = 1.56

$$\frac{2(7 - 4 \cos^2(a + bx)^{3/4} + 4(-1 + \cos^2(a + bx)^{3/4}) \csc^2(a + bx)) \tan^2(a + bx)}{21bd^3(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^3/(d*Cos[a + b*x])^(9/2), x]

[Out] (2*(7 - 4*(Cos[a + b*x]^2)^(3/4) + 4*(-1 + (Cos[a + b*x]^2)^(3/4))*Csc[a + b*x]^2)*Tan[a + b*x]^2)/(21*b*d^3*(d*Cos[a + b*x])^(3/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 110 vs. 2(37) = 74.

time = 0.37, size = 111, normalized size = 2.47

method	result	size
default	$-\frac{8\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + d\left(7\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 7\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)}{21d^5\left(16\left(\sin^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 32\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 24\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 8\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)b}$	111

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^3/(d*cos(b*x+a))^(9/2), x, method=_RETURNVERBOSE)

[Out] -8/21/d^5/(16*sin(1/2*b*x+1/2*a)^8-32*sin(1/2*b*x+1/2*a)^6+24*sin(1/2*b*x+1/2*a)^4-8*sin(1/2*b*x+1/2*a)^2+1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(7*sin(1/2*b*x+1/2*a)^4-7*sin(1/2*b*x+1/2*a)^2+1)/b

Maxima [A]

time = 0.28, size = 37, normalized size = 0.82

$$\frac{2(7d^2 \cos(bx + a)^2 - 3d^2)}{21(d \cos(bx + a))^{7/2} bd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(9/2), x, algorithm="maxima")

[Out] -2/21*(7*d^2*cos(b*x + a)^2 - 3*d^2)/((d*cos(b*x + a))^(7/2)*b*d^3)

Fricas [A]

time = 0.35, size = 38, normalized size = 0.84

$$\frac{2\sqrt{d \cos(bx + a)}(7 \cos(bx + a)^2 - 3)}{21bd^5 \cos(bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")

[Out] -2/21*sqrt(d*cos(b*x + a))*(7*cos(b*x + a)^2 - 3)/(b*d^5*cos(b*x + a)^4)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**3/(d*cos(b*x+a))**(9/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4847 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^3/(d*cos(b*x + a))^(9/2), x)

Mupad [B]

time = 3.78, size = 93, normalized size = 2.07

$$-\frac{8e^{a2i+bx2i} \sqrt{d \left(\frac{e^{-a1i-bx1i}}{2} + \frac{e^{a1i+bx1i}}{2} \right)} (2e^{a2i+bx2i} + 7e^{a4i+bx4i} + 7)}{21bd^5(e^{a2i+bx2i} + 1)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3/(d*cos(a + b*x))^(9/2),x)

[Out] -(8*exp(a*2i + b*x*2i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2)*(2*exp(a*2i + b*x*2i) + 7*exp(a*4i + b*x*4i) + 7))/(21*b*d^5*(exp(a*2i + b*x*2i) + 1)^4)

$$3.210 \quad \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{11/2}} dx$$

Optimal. Leaf size=45

$$\frac{2}{9bd(d \cos(a+bx))^{9/2}} - \frac{2}{5bd^3(d \cos(a+bx))^{5/2}}$$

[Out] 2/9/b/d/(d*cos(b*x+a))^(9/2)-2/5/b/d^3/(d*cos(b*x+a))^(5/2)

Rubi [A]

time = 0.03, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2645, 14}

$$\frac{2}{9bd(d \cos(a+bx))^{9/2}} - \frac{2}{5bd^3(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^3/(d*Cos[a + b*x])^(11/2), x]

[Out] 2/(9*b*d*(d*Cos[a + b*x])^(9/2)) - 2/(5*b*d^3*(d*Cos[a + b*x])^(5/2))

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2645

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n-1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(a+bx)}{(d \cos(a+bx))^{11/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1-x^2}{x^{11/2}} dx, x, d \cos(a+bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(\frac{1}{x^{11/2}} - \frac{1}{d^2 x^{7/2}}\right) dx, x, d \cos(a+bx)\right)}{bd} \\ &= \frac{2}{9bd(d \cos(a+bx))^{9/2}} - \frac{2}{5bd^3(d \cos(a+bx))^{5/2}} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 94 vs. 2(45) = 90.

time = 0.37, size = 94, normalized size = 2.09

$$\frac{2\left(4\sqrt{\cos^2(a+bx)} + \left(9 - 8\sqrt{\cos^2(a+bx)}\right) \csc^2(a+bx) + 4\left(-1 + \sqrt{\cos^2(a+bx)}\right) \csc^4(a+bx)\right) \tan^4(a+bx)}{45bd^5 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^3/(d*Cos[a + b*x])^(11/2), x]

[Out] (2*(4*(Cos[a + b*x]^2)^(1/4) + (9 - 8*(Cos[a + b*x]^2)^(1/4))*Csc[a + b*x]^2 + 4*(-1 + (Cos[a + b*x]^2)^(1/4))*Csc[a + b*x]^4)*Tan[a + b*x]^4)/(45*b*d^5*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 123 vs. 2(37) = 74.

time = 0.46, size = 124, normalized size = 2.76

method	result	size
default	$\frac{8\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + d\left(9\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 9\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)}{45d^6\left(32\left(\sin^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 80\left(\sin^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 80\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 40\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 10\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)b}$	124

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^3/(d*cos(b*x+a))^(11/2), x, method=_RETURNVERBOSE)

[Out] 8/45/d^6/(32*sin(1/2*b*x+1/2*a)^10-80*sin(1/2*b*x+1/2*a)^8+80*sin(1/2*b*x+1/2*a)^6-40*sin(1/2*b*x+1/2*a)^4+10*sin(1/2*b*x+1/2*a)^2-1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(9*sin(1/2*b*x+1/2*a)^4-9*sin(1/2*b*x+1/2*a)^2+1)/b

Maxima [A]

time = 0.29, size = 37, normalized size = 0.82

$$\frac{2(9d^2 \cos(bx+a)^2 - 5d^2)}{45(d \cos(bx+a))^{\frac{9}{2}} bd^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(11/2), x, algorithm="maxima")

[Out] -2/45*(9*d^2*cos(b*x + a)^2 - 5*d^2)/((d*cos(b*x + a))^(9/2)*b*d^3)

Fricas [A]

time = 0.35, size = 38, normalized size = 0.84

$$\frac{2\sqrt{d \cos(bx+a)}(9 \cos(bx+a)^2 - 5)}{45bd^6 \cos(bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(11/2),x, algorithm="fricas")`

[Out] `-2/45*sqrt(d*cos(b*x + a))*(9*cos(b*x + a)^2 - 5)/(b*d^6*cos(b*x + a)^5)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**3/(d*cos(b*x+a))**(11/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^3/(d*cos(b*x+a))^(11/2),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^3/(d*cos(b*x + a))^(11/2), x)`

Mupad [B]

time = 5.33, size = 279, normalized size = 6.20

$$\frac{16 e^{a l i+b x l i} \sqrt{d\left(\frac{e^{-a l i-b x l i}}{2}+\frac{e^{a l i+b x l i}}{2}\right)}}{5 b d^6\left(e^{a 2 i+b x 2 i} l i+l i\right)^2}-\frac{e^{a l i+b x l i} \sqrt{d\left(\frac{e^{-a l i-b x l i}}{2}+\frac{e^{a l i+b x l i}}{2}\right)} 464 i}{45 b d^6\left(e^{a 2 i+b x 2 i} l i+l i\right)^3}-\frac{128 e^{a l i+b x l i} \sqrt{d\left(\frac{e^{-a l i-b x l i}}{2}+\frac{e^{a l i+b x l i}}{2}\right)}}{9 b d^6\left(e^{a 2 i+b x 2 i} l i+l i\right)^4}+\frac{e^{a l i+b x l i} \sqrt{d\left(\frac{e^{-a l i-b x l i}}{2}+\frac{e^{a l i+b x l i}}{2}\right)} 64 i}{9 b d^6\left(e^{a 2 i+b x 2 i} l i+l i\right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^3/(d*cos(a + b*x))^(11/2),x)`

[Out] `(16*exp(a*1i + b*x*1i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2))/(5*b*d^6*(exp(a*2i + b*x*2i)*1i + 1i)^2) - (exp(a*1i + b*x*1i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2)*464i)/(45*b*d^6*(exp(a*2i + b*x*2i)*1i + 1i)^3) - (128*exp(a*1i + b*x*1i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2))/(9*b*d^6*(exp(a*2i + b*x*2i)*1i + 1i)^4) + (exp(a*1i + b*x*1i)*(d*(exp(- a*1i - b*x*1i)/2 + exp(a*1i + b*x*1i)/2))^(1/2)*64i)/(9*b*d^6*(exp(a*2i + b*x*2i)*1i + 1i)^5)`

3.211 $\int (d \cos(a + bx))^{9/2} \sin^4(a + bx) dx$

Optimal. Leaf size=156

$$\frac{56d^4 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{1105b \sqrt{\cos(a + bx)}} + \frac{56d^3 (d \cos(a + bx))^{3/2} \sin(a + bx)}{3315b} + \frac{8d (d \cos(a + bx))^{7/2} \sin(a + bx)}{663b}$$

[Out] $56/3315*d^3*(d*\cos(b*x+a))^{(3/2)*\sin(b*x+a)/b+8/663*d*(d*\cos(b*x+a))^{(7/2)*\sin(b*x+a)/b-12/221*(d*\cos(b*x+a))^{(11/2)*\sin(b*x+a)/b/d-2/17*(d*\cos(b*x+a))^{(11/2)*\sin(b*x+a)^3/b/d+56/1105*d^4*(\cos(1/2*a+1/2*b*x))^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 156, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2719}

$$\frac{56d^4 E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{1105b \sqrt{\cos(a + bx)}} + \frac{56d^3 \sin(a + bx) (d \cos(a + bx))^{3/2}}{3315b} - \frac{2 \sin^3(a + bx) (d \cos(a + bx))^{11/2}}{17bd} - \frac{12 \sin(a + bx) (d \cos(a + bx))^{11/2}}{221bd} + \frac{8d \sin(a + bx) (d \cos(a + bx))^{7/2}}{663b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(9/2)}*\text{Sin}[a + b*x]^4, x]$

[Out] $(56*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(1105*b*\text{Sqrt}[\text{Cos}[a + b*x]]) + (56*d^3*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x])/(3315*b) + (8*d*(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x])/(663*b) - (12*(d*\text{Cos}[a + b*x])^{(11/2)}*\text{Sin}[a + b*x])/(221*b*d) - (2*(d*\text{Cos}[a + b*x])^{(11/2)}*\text{Sin}[a + b*x]^3)/(17*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)/(b*f*(m + n))}), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n - 1)/(d*n)}, x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \int (d \cos(a + bx))^{9/2} \sin^4(a + bx) dx &= -\frac{2(d \cos(a + bx))^{11/2} \sin^3(a + bx)}{17bd} + \frac{6}{17} \int (d \cos(a + bx))^{9/2} \sin^2(a + bx) dx \\
 &= -\frac{12(d \cos(a + bx))^{11/2} \sin(a + bx)}{221bd} - \frac{2(d \cos(a + bx))^{11/2} \sin^3(a + bx)}{17bd} \\
 &= \frac{8d(d \cos(a + bx))^{7/2} \sin(a + bx)}{663b} - \frac{12(d \cos(a + bx))^{11/2} \sin(a + bx)}{221bd} \\
 &= \frac{56d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{3315b} + \frac{8d(d \cos(a + bx))^{7/2} \sin(a + bx)}{663b} \\
 &= \frac{56d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{3315b} + \frac{8d(d \cos(a + bx))^{7/2} \sin(a + bx)}{663b} \\
 &= \frac{56d^4 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{1105b \sqrt{\cos(a + bx)}} + \frac{56d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{3315b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.10, size = 57, normalized size = 0.37

$$\frac{(d \cos(a + bx))^{9/2} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(-\frac{7}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a + bx)\right) \tan^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(9/2)*Sin[a + b*x]^4,x]

[Out] ((d*Cos[a + b*x])^(9/2)*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-7/4, 5/2, 7/2, Sin[a + b*x]^2]*Tan[a + b*x]^5)/(5*b)

Maple [A]

time = 0.19, size = 275, normalized size = 1.76

method	result
default	$\frac{8\sqrt{d}\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^5 d^5 \left(24960\left(\cos^{19}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 124800\left(\cos^{17}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 265440\left(\cos^{15}\right)\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(9/2)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out]
$$-8/3315*(d*(2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*d^5*(24960*\cos(1/2*b*x+1/2*a)^{19}-124800*\cos(1/2*b*x+1/2*a)^{17}+265440*\cos(1/2*b*x+1/2*a)^{15}-312960*\cos(1/2*b*x+1/2*a)^{13}+222520*\cos(1/2*b*x+1/2*a)^{11}-96360*\cos(1/2*b*x+1/2*a)^9+23866*\cos(1/2*b*x+1/2*a)^7-2652*\cos(1/2*b*x+1/2*a)^5-35*\cos(1/2*b*x+1/2*a)^3-21*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(-2*\cos(1/2*b*x+1/2*a)^2+1)^{(1/2)}*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)})+21*\cos(1/2*b*x+1/2*a))/(-d*(2*\sin(1/2*b*x+1/2*a)^4-\sin(1/2*b*x+1/2*a)^2))^{(1/2)}/\sin(1/2*b*x+1/2*a)/(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}/b$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*sin(b*x+a)^4,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(9/2)*sin(b*x + a)^4, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 133, normalized size = 0.85

$\frac{2(-42i\sqrt{2}d^2\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(lr+a)+i\sin(lr+a))) + 42i\sqrt{2}d^2\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(lr+a)-i\sin(lr+a))) - (195d^4\cos(lr+a)^7 - 285d^4\cos(lr+a)^5 + 20d^4\cos(lr+a)^3 + 28d^4\cos(lr+a))\sqrt{d\cos(lr+a)}\sin(lr+a))}{3315}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*sin(b*x+a)^4,x, algorithm="fricas")`

[Out]
$$-2/3315*(-42*I*\sqrt{2}*d^{(9/2)}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(b*x+a)+I*\sin(b*x+a))) + 42*I*\sqrt{2}*d^{(9/2)}*\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(b*x+a)-I*\sin(b*x+a))) - (195*d^4*\cos(b*x+a)^7 - 285*d^4*\cos(b*x+a)^5 + 20*d^4*\cos(b*x+a)^3 + 28*d^4*\cos(b*x+a))*\sqrt{d*\cos(b*x+a)}*\sin(b*x+a))/b$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(9/2)*sin(b*x+a)**4,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*sin(b*x+a)^4,x, algorithm="giac")`

[Out] `integrate((d*cos(b*x + a))^(9/2)*sin(b*x + a)^4, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^4 (d \cos(a + bx))^{9/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^4*(d*cos(a + b*x))^(9/2),x)`

[Out] `int(sin(a + b*x)^4*(d*cos(a + b*x))^(9/2), x)`

3.212 $\int (d \cos(a + bx))^{7/2} \sin^4(a + bx) dx$

Optimal. Leaf size=156

$$\frac{8d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{231b \sqrt{d \cos(a + bx)}} + \frac{8d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b} + \frac{8d(d \cos(a + bx))^{5/2} \sin(a + bx)}{385b} - \frac{4(d \cos(a + bx))^{9/2} \sin(a + bx)}{15bd}$$

[Out] $8/385*d*(d*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)/b-4/55*(d*\cos(b*x+a))^{(9/2)}*\sin(b*x+a)/b/d-2/15*(d*\cos(b*x+a))^{(9/2)}*\sin(b*x+a)^3/b/d+8/231*d^4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}+8/231*d^3*\sin(b*x+a)*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.11, antiderivative size = 156, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2720}

$$\frac{8d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{231b \sqrt{d \cos(a + bx)}} + \frac{8d^3 \sin(a + bx) \sqrt{d \cos(a + bx)}}{231b} - \frac{2 \sin^3(a + bx) (d \cos(a + bx))^{9/2}}{15bd} - \frac{4 \sin(a + bx) (d \cos(a + bx))^{9/2}}{55bd} + \frac{8d \sin(a + bx) (d \cos(a + bx))^{5/2}}{385b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x]^4, x]$

[Out] $(8*d^4*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(231*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (8*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(231*b) + (8*d*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x])/(385*b) - (4*(d*\text{Cos}[a + b*x])^{(9/2)}*\text{Sin}[a + b*x])/(55*b*d) - (2*(d*\text{Cos}[a + b*x])^{(9/2)}*\text{Sin}[a + b*x]^3)/(15*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)/(b*f*(m + n))}), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^n*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n - 1)/(d*n)}, x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \int (d \cos(a + bx))^{7/2} \sin^4(a + bx) dx &= -\frac{2(d \cos(a + bx))^{9/2} \sin^3(a + bx)}{15bd} + \frac{2}{5} \int (d \cos(a + bx))^{7/2} \sin^2(a + bx) dx \\
 &= -\frac{4(d \cos(a + bx))^{9/2} \sin(a + bx)}{55bd} - \frac{2(d \cos(a + bx))^{9/2} \sin^3(a + bx)}{15bd} \\
 &= \frac{8d(d \cos(a + bx))^{5/2} \sin(a + bx)}{385b} - \frac{4(d \cos(a + bx))^{9/2} \sin(a + bx)}{55bd} \\
 &= \frac{8d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b} + \frac{8d(d \cos(a + bx))^{5/2} \sin(a + bx)}{385b} \\
 &= \frac{8d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b} + \frac{8d(d \cos(a + bx))^{5/2} \sin(a + bx)}{385b} \\
 &= \frac{8d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{231b \sqrt{d \cos(a + bx)}} + \frac{8d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{231b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.07, size = 57, normalized size = 0.37

$$\frac{(d \cos(a + bx))^{7/2} \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{5}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a + bx)\right) \tan^5(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(7/2)*Sin[a + b*x]^4,x]

[Out] ((d*Cos[a + b*x])^(7/2)*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-5/4, 5/2, 7/2, Sin[a + b*x]^2]*Tan[a + b*x]^5)/(5*b)

Maple [A]

time = 0.21, size = 262, normalized size = 1.68

method	result
default	$\frac{8\sqrt{d}\left(2\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)^d d^4\left(4928\cos^{17}\left(\frac{bx}{2}+\frac{a}{2}\right)-22176\cos^{15}\left(\frac{bx}{2}+\frac{a}{2}\right)+41216\cos^{13}\left(\frac{bx}{2}+\frac{a}{2}\right)-22176\cos^{11}\left(\frac{bx}{2}+\frac{a}{2}\right)+4928\cos^9\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{1155\sqrt{-}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(7/2)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)`

[Out] $-8/1155*(d*(2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*d^4*(4928*\cos(1/2*b*x+1/2*a)^{17}-22176*\cos(1/2*b*x+1/2*a)^{15}+41216*\cos(1/2*b*x+1/2*a)^{13}-40768*\cos(1/2*b*x+1/2*a)^{11}+22868*\cos(1/2*b*x+1/2*a)^9-6994*\cos(1/2*b*x+1/2*a)^7+926*\cos(1/2*b*x+1/2*a)^5+5*\cos(1/2*b*x+1/2*a)^3+5*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*(-2*\cos(1/2*b*x+1/2*a)^2+1)^{(1/2)}*\text{EllipticF}(\cos(1/2*b*x+1/2*a),2^{(1/2)})-5*\cos(1/2*b*x+1/2*a))/(-d*(2*\sin(1/2*b*x+1/2*a)^4-\sin(1/2*b*x+1/2*a)^2))^{(1/2)}/\sin(1/2*b*x+1/2*a)/(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}/b$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*sin(b*x+a)^4,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(7/2)*sin(b*x + a)^4, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.15, size = 121, normalized size = 0.78

$\frac{2\left(10i\sqrt{2}d^2\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))-10i\sqrt{2}d^2\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))-(77d^3\cos(bx+a)^6-119d^3\cos(bx+a)^4+12d^3\cos(bx+a)^2+20d^3)\sqrt{d\cos(bx+a)}\sin(bx+a)}{1155b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*sin(b*x+a)^4,x, algorithm="fricas")`

[Out] $-2/1155*(10*I*\sqrt{2}*d^{(7/2)}*\text{weierstrassPInverse}(-4,0,\cos(b*x+a)+I*\sin(b*x+a))-10*I*\sqrt{2}*d^{(7/2)}*\text{weierstrassPInverse}(-4,0,\cos(b*x+a)-I*\sin(b*x+a))-(77*d^3*\cos(b*x+a)^6-119*d^3*\cos(b*x+a)^4+12*d^3*\cos(b*x+a)^2+20*d^3)*\sqrt{d*\cos(b*x+a)}*\sin(b*x+a))/b$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(7/2)*sin(b*x+a)**4,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 8856 deep
```

Giac [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(7/2)*sin(b*x+a)^4,x, algorithm="giac")
```

```
[Out] integrate((d*cos(b*x + a))^(7/2)*sin(b*x + a)^4, x)
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.01
```

$$\int \sin(a + bx)^4 (d \cos(a + bx))^{7/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(a + b*x)^4*(d*cos(a + b*x))^(7/2),x)
```

```
[Out] int(sin(a + b*x)^4*(d*cos(a + b*x))^(7/2), x)
```

3.213 $\int (d \cos(a + bx))^{5/2} \sin^4(a + bx) dx$

Optimal. Leaf size=128

$$\frac{8d^2 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{65b \sqrt{\cos(a + bx)}} + \frac{8d(d \cos(a + bx))^{3/2} \sin(a + bx)}{195b} - \frac{4(d \cos(a + bx))^{7/2} \sin(a + bx)}{39bd} - \frac{2(d \cos(a + bx))^{9/2} \sin^3(a + bx)}{13bd^2}$$

[Out] $8/195*d*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)/b-4/39*(d*\cos(b*x+a))^{(7/2)}*\sin(b*x+a)/b/d-2/13*(d*\cos(b*x+a))^{(7/2)}*\sin(b*x+a)^3/b/d+8/65*d^2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.09, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2719}

$$\frac{8d^2 E\left(\frac{1}{2}(a + bx) \middle| 2\right) \sqrt{d \cos(a + bx)}}{65b \sqrt{\cos(a + bx)}} - \frac{2 \sin^3(a + bx) (d \cos(a + bx))^{7/2}}{13bd} - \frac{4 \sin(a + bx) (d \cos(a + bx))^{7/2}}{39bd} + \frac{8d \sin(a + bx) (d \cos(a + bx))^{3/2}}{195b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x]^4, x]$

[Out] $(8*d^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/ (65*b*\text{Sqrt}[\text{Cos}[a + b*x]]) + (8*d*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x])/ (195*b) - (4*(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x])/ (39*b*d) - (2*(d*\text{Cos}[a + b*x])^{(7/2)}*\text{Sin}[a + b*x]^3)/ (13*b*d)$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] := \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] := \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n - 1)})/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{5/2} \sin^4(a + bx) dx &= -\frac{2(d \cos(a + bx))^{7/2} \sin^3(a + bx)}{13bd} + \frac{6}{13} \int (d \cos(a + bx))^{5/2} \sin^2(a + bx) dx \\ &= -\frac{4(d \cos(a + bx))^{7/2} \sin(a + bx)}{39bd} - \frac{2(d \cos(a + bx))^{7/2} \sin^3(a + bx)}{13bd} \\ &= \frac{8d(d \cos(a + bx))^{3/2} \sin(a + bx)}{195b} - \frac{4(d \cos(a + bx))^{7/2} \sin(a + bx)}{39bd} \\ &= \frac{8d(d \cos(a + bx))^{3/2} \sin(a + bx)}{195b} - \frac{4(d \cos(a + bx))^{7/2} \sin(a + bx)}{39bd} \\ &= \frac{8d^2 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{65b \sqrt{\cos(a + bx)}} + \frac{8d(d \cos(a + bx))^{3/2} \sin(a + bx)}{195b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 65, normalized size = 0.51

$$\frac{(d \cos(a + bx))^{5/2} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(-\frac{3}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a + bx)\right) \sin^2(a + bx) \tan^3(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(5/2)*Sin[a + b*x]^4,x]

[Out] ((d*cos[a + b*x])^(5/2)*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-3/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^2*Tan[a + b*x]^3)/(5*b)

Maple [A]

time = 0.18, size = 249, normalized size = 1.95

method	result
default	$\frac{8 \sqrt{d \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d^3 \left(480 \left(\cos^{15}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1920 \left(\cos^{13}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 3040 \left(\cos^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1920 \left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 480 \left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 96 \left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 8 \left(\cos^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 8 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)}{195 \sqrt{-d} \left(2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(5/2)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)
```

```
[Out] -8/195*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^3*(480*cos(1/2*b*x+1/2*a)^15-1920*cos(1/2*b*x+1/2*a)^13+3040*cos(1/2*b*x+1/2*a)^11-2400*cos(1/2*b*x+1/2*a)^9+958*cos(1/2*b*x+1/2*a)^7-156*cos(1/2*b*x+1/2*a)^5-5*cos(1/2*b*x+1/2*a)^3-3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))+3*cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*sin(b*x+a)^4,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(5/2)*sin(b*x + a)^4, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.12, size = 120, normalized size = 0.94

$$\frac{2(-6i\sqrt{2}d^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))) + 6i\sqrt{2}d^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))) - (15d^2\cos(bx+a)^5 - 25d^2\cos(bx+a)^3 + 4d^2\cos(bx+a))\sqrt{d\cos(bx+a)}\sin(bx+a))}{195b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*sin(b*x+a)^4,x, algorithm="fricas")
```

```
[Out] -2/195*(-6*I*sqrt(2)*d^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + 6*I*sqrt(2)*d^(5/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) - (15*d^2*cos(b*x + a)^5 - 25*d^2*cos(b*x + a)^3 + 4*d^2*cos(b*x + a))*sqrt(d*cos(b*x + a))*sin(b*x + a))/b
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(5/2)*sin(b*x+a)**4,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3877 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^(5/2)*sin(b*x+a)^4,x, algorithm="giac")``[Out] integrate((d*cos(b*x + a))^(5/2)*sin(b*x + a)^4, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^4 (d \cos(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^4*(d*cos(a + b*x))^(5/2),x)``[Out] int(sin(a + b*x)^4*(d*cos(a + b*x))^(5/2), x)`

3.214 $\int (d \cos(a + bx))^{3/2} \sin^4(a + bx) dx$

Optimal. Leaf size=128

$$\frac{8d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{77b \sqrt{d \cos(a + bx)}} + \frac{8d \sqrt{d \cos(a + bx)} \sin(a + bx)}{77b} - \frac{12(d \cos(a + bx))^{5/2} \sin(a + bx)}{77bd} - \frac{2(d \cos(a + bx))^{3/2} \sin^3(a + bx)}{77bd}$$

[Out] $-12/77*(d*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)/b/d-2/11*(d*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)^3/b/d+8/77*d^2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}+8/77*d*\sin(b*x+a)*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.09, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2648, 2715, 2721, 2720}

$$\frac{8d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{77b \sqrt{d \cos(a + bx)}} - \frac{2 \sin^3(a + bx) (d \cos(a + bx))^{5/2}}{11bd} - \frac{12 \sin(a + bx) (d \cos(a + bx))^{5/2}}{77bd} + \frac{8d \sin(a + bx) \sqrt{d \cos(a + bx)}}{77b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x]^4, x]$

[Out] $(8*d^2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/((77*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (8*d*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/((77*b) - (12*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x])/((77*b*d) - (2*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x]^3)/(11*b*d))$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)/(b*f*(m + n))}), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2715

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n - 1)/(d*n)}), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Dist[(b*Sin[c + d*x])ⁿ/Sin[c + d*x]ⁿ, Int[Sin[c + d*x]ⁿ, x], x] /; FreeQ[{b, c, d}, x] && LtQ [-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \int (d \cos(a + bx))^{3/2} \sin^4(a + bx) dx &= -\frac{2(d \cos(a + bx))^{5/2} \sin^3(a + bx)}{11bd} + \frac{6}{11} \int (d \cos(a + bx))^{3/2} \sin^2(a + bx) dx \\
 &= -\frac{12(d \cos(a + bx))^{5/2} \sin(a + bx)}{77bd} - \frac{2(d \cos(a + bx))^{5/2} \sin^3(a + bx)}{11bd} \\
 &= \frac{8d \sqrt{d \cos(a + bx)} \sin(a + bx)}{77b} - \frac{12(d \cos(a + bx))^{5/2} \sin(a + bx)}{77bd} \\
 &= \frac{8d \sqrt{d \cos(a + bx)} \sin(a + bx)}{77b} - \frac{12(d \cos(a + bx))^{5/2} \sin(a + bx)}{77bd} \\
 &= \frac{8d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{77b \sqrt{d \cos(a + bx)}} + \frac{8d \sqrt{d \cos(a + bx)} \sin(a + bx)}{77b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.08, size = 65, normalized size = 0.51

$$\frac{(d \cos(a + bx))^{3/2} \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a + bx)\right) \sin^2(a + bx) \tan^3(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(3/2)*Sin[a + b*x]^4,x]

[Out] ((d*Cos[a + b*x])^(3/2)*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^2*Tan[a + b*x]^3)/(5*b)

Maple [A]

time = 0.21, size = 255, normalized size = 1.99

method	result
default	$ \frac{8 \sqrt{d} \left(2 \cos^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)^2 d^2 \left(112 \cos\left(\frac{bx}{2} + \frac{a}{2}\right) \left(\sin^{12}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 280 \left(\sin^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{77 \sqrt{-d} \left(2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(3/2)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)
```

```
[Out] -8/77*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^2*(112*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^12-280*sin(1/2*b*x+1/2*a)^10*cos(1/2*b*x+1/2*a)+228*sin(1/2*b*x+1/2*a)^8*cos(1/2*b*x+1/2*a)-62*sin(1/2*b*x+1/2*a)^6*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))+sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a)^4,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(3/2)*sin(b*x + a)^4, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 102, normalized size = 0.80

$$\frac{2(2i\sqrt{2}d^3\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))-2i\sqrt{2}d^3\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))-(7d\cos(bx+a)^4-13d\cos(bx+a)^2+4d)\sqrt{d\cos(bx+a)}\sin(bx+a))}{77b}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a)^4,x, algorithm="fricas")
```

```
[Out] -2/77*(2*I*sqrt(2)*d^(3/2)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) - 2*I*sqrt(2)*d^(3/2)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) - (7*d*cos(b*x + a)^4 - 13*d*cos(b*x + a)^2 + 4*d)*sqrt(d*cos(b*x + a))*sin(b*x + a))/b
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(3/2)*sin(b*x+a)**4,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3879 deep
```


Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^(3/2)*sin(b*x+a)^4,x, algorithm="giac")``[Out] integrate((d*cos(b*x + a))^(3/2)*sin(b*x + a)^4, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^4 (d \cos(a + bx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^4*(d*cos(a + b*x))^(3/2),x)``[Out] int(sin(a + b*x)^4*(d*cos(a + b*x))^(3/2), x)`

3.215 $\int \sqrt{d \cos(a + bx)} \sin^4(a + bx) dx$

Optimal. Leaf size=99

$$\frac{8\sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{15b\sqrt{\cos(a + bx)}} - \frac{4(d \cos(a + bx))^{3/2} \sin(a + bx)}{15bd} - \frac{2(d \cos(a + bx))^{3/2} \sin^3(a + bx)}{9bd}$$

[Out] $-4/15*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)/b/d-2/9*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)^3/b/d+8/15*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2648, 2721, 2719}

$$-\frac{2 \sin^3(a + bx)(d \cos(a + bx))^{3/2}}{9bd} - \frac{4 \sin(a + bx)(d \cos(a + bx))^{3/2}}{15bd} + \frac{8E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{15b\sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^4,x]

[Out] $(8*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/((15*b*\text{Sqrt}[\text{Cos}[a + b*x]]) - (4*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x]))/(15*b*d) - (2*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x]^3)/(9*b*d)$

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*SIN[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*SIN[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*SIN[c + d*x])^n/SIN[c + d*x]^n, Int[SIN[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \sqrt{d \cos(a + bx)} \sin^4(a + bx) dx &= -\frac{2(d \cos(a + bx))^{3/2} \sin^3(a + bx)}{9bd} + \frac{2}{3} \int \sqrt{d \cos(a + bx)} \sin^2(a + bx) dx \\
&= -\frac{4(d \cos(a + bx))^{3/2} \sin(a + bx)}{15bd} - \frac{2(d \cos(a + bx))^{3/2} \sin^3(a + bx)}{9bd} + \frac{2}{3} \int \sqrt{d \cos(a + bx)} dx \\
&= -\frac{4(d \cos(a + bx))^{3/2} \sin(a + bx)}{15bd} - \frac{2(d \cos(a + bx))^{3/2} \sin^3(a + bx)}{9bd} + \frac{2}{3} \sqrt{d \cos(a + bx)} \\
&= \frac{8\sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \middle| 2\right)}{15b\sqrt{\cos(a + bx)}} - \frac{4(d \cos(a + bx))^{3/2} \sin(a + bx)}{15bd}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 58, normalized size = 0.59

$$\frac{d^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a + bx)\right) \sin^5(a + bx)}{5b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^4,x]

[Out] (d*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^5)/(5*b*Sqrt[d*Cos[a + b*x]])

Maple [A]

time = 0.22, size = 221, normalized size = 2.23

method	result
default	$ \frac{8\sqrt{d\left(2\left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{45\sqrt{-d\left(2\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}} d\left(40\left(\cos^{11}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 120\left(\cos^9\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 118\left(\cos^7\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 36\left(\cos^5\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 5\left(\cos^3\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - \cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right) $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*sin(b*x+a)^4,x,method=_RETURNVERBOSE)

[Out] -8/45*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d*(40*cos(1/2*b*x+1/2*a)^11-120*cos(1/2*b*x+1/2*a)^9+118*cos(1/2*b*x+1/2*a)^7-36*cos(1/2*b*x+1/2*a)^5-5*cos(1/2*b*x+1/2*a)^3-3*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(-2*cos(1/2*b*x+1/2*a)^2+1)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))+3*cos(1/2*b*x+1/2*a))/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2)^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^4,x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*sin(b*x + a)^4, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 100, normalized size = 1.01

$$\frac{2 \left((5 \cos(bx+a)^3 - 11 \cos(bx+a)) \sqrt{d \cos(bx+a)} \sin(bx+a) + 6i \sqrt{2} \sqrt{d} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a))) - 6i \sqrt{2} \sqrt{d} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a))) \right)}{45b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^4,x, algorithm="fricas")

[Out] $\frac{2}{45} * ((5 * \cos(b*x + a))^3 - 11 * \cos(b*x + a)) * \sqrt{d * \cos(b*x + a)} * \sin(b*x + a) + 6 * I * \sqrt{2} * \sqrt{d} * \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I * \sin(b*x + a))) - 6 * I * \sqrt{2} * \sqrt{d} * \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I * \sin(b*x + a))) / b$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)*sin(b*x+a)**4,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*sin(b*x+a)^4,x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*sin(b*x + a)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^4 \sqrt{d \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4*(d*cos(a + b*x))^(1/2),x)

[Out] int(sin(a + b*x)^4*(d*cos(a + b*x))^(1/2), x)

$$3.216 \quad \int \frac{\sin^4(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=99

$$\frac{8\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{7b\sqrt{d \cos(a+bx)}} - \frac{4\sqrt{d \cos(a+bx)} \sin(a+bx)}{7bd} - \frac{2\sqrt{d \cos(a+bx)} \sin^3(a+bx)}{7bd}$$

[Out] 8/7*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)/b/(d*cos(b*x+a))^(1/2)-4/7*sin(b*x+a)*(d*cos(b*x+a))^(1/2)/b/d-2/7*sin(b*x+a)^3*(d*cos(b*x+a))^(1/2)/b/d

Rubi [A]

time = 0.07, antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$,

Rules used = {2648, 2721, 2720}

$$-\frac{2 \sin^3(a+bx) \sqrt{d \cos(a+bx)}}{7bd} - \frac{4 \sin(a+bx) \sqrt{d \cos(a+bx)}}{7bd} + \frac{8 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{7b\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^4/Sqrt[d*Cos[a + b*x]],x]

[Out] (8*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(7*b*sqrt[d*Cos[a + b*x]]) - (4*sqrt[d*Cos[a + b*x]]*Sin[a + b*x])/(7*b*d) - (2*sqrt[d*Cos[a + b*x]]*Sin[a + b*x]^3)/(7*b*d)

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(b*cos[e + f*x])^(n + 1)*((a*sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*cos[e + f*x])^n*(a*sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^4(a+bx)}{\sqrt{d \cos(a+bx)}} dx &= -\frac{2\sqrt{d \cos(a+bx)} \sin^3(a+bx)}{7bd} + \frac{6}{7} \int \frac{\sin^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx \\
&= -\frac{4\sqrt{d \cos(a+bx)} \sin(a+bx)}{7bd} - \frac{2\sqrt{d \cos(a+bx)} \sin^3(a+bx)}{7bd} + \frac{4}{7} \int \frac{1}{\sqrt{d \cos(a+bx)}} dx \\
&= -\frac{4\sqrt{d \cos(a+bx)} \sin(a+bx)}{7bd} - \frac{2\sqrt{d \cos(a+bx)} \sin^3(a+bx)}{7bd} + \frac{(4\sqrt{\cos(a+bx)})}{7\sqrt{d}} \\
&= \frac{8\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{7b\sqrt{d \cos(a+bx)}} - \frac{4\sqrt{d \cos(a+bx)} \sin(a+bx)}{7bd} - \frac{2\sqrt{d \cos(a+bx)} \sin^3(a+bx)}{7bd}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 58, normalized size = 0.59

$$\frac{d \cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a+bx)\right) \sin^5(a+bx)}{5b(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^4/Sqrt[d*Cos[a + b*x]], x]

[Out] (d*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^5)/(5*b*(d*Cos[a + b*x])^(3/2))

Maple [A]

time = 0.17, size = 208, normalized size = 2.10

method	result
default	$ -\frac{8\sqrt{d\left(2\cos^2\left(\frac{bx}{2}+\frac{a}{2}\right)-1\right)\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)}{7\sqrt{-d\left(2\left(\sin^4\left(\frac{bx}{2}+\frac{a}{2}\right)\right)-\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\right)}} \left(4\left(\sin^8\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\cos\left(\frac{bx}{2}+\frac{a}{2}\right)-6\left(\sin^6\left(\frac{bx}{2}+\frac{a}{2}\right)\right)\cos\left(\frac{bx}{2}+\frac{a}{2}\right)+\sqrt{\dots}\right) $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^4/(d*cos(b*x+a))^(1/2), x, method=_RETURNVERBOSE)

[Out] -8/7*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*(4*sin(1/2*b*x+1/2*a)^8*cos(1/2*b*x+1/2*a)-6*sin(1/2*b*x+1/2*a)^6*cos(1/2*b*x+1/2*a)+(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1

$$\frac{\sqrt{2bx+1/2a}, 2^{(1/2)} + \sin(1/2bx+1/2a)^2 \cos(1/2bx+1/2a)}{(-d(2\sin(1/2bx+1/2a)^4 - \sin(1/2bx+1/2a)^2))^{(1/2)} / \sin(1/2bx+1/2a) / (d(2\cos(1/2bx+1/2a)^2 - 1))^{(1/2)}} / b$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^4/sqrt(d*cos(b*x + a)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 88, normalized size = 0.89

$$\frac{2 \left(\sqrt{d \cos(bx+a)} (\cos(bx+a)^2 - 3) \sin(bx+a) - 2i \sqrt{2} \sqrt{d} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a)) + 2i \sqrt{2} \sqrt{d} \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a)) \right)}{7bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] 2/7*(sqrt(d*cos(b*x + a))*(cos(b*x + a)^2 - 3)*sin(b*x + a) - 2*I*sqrt(2)*sqrt(d)*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + 2*I*sqrt(2)*sqrt(d)*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)))/(b*d)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**4/(d*cos(b*x+a))**(1/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3064 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^4/sqrt(d*cos(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^4}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/(d*cos(a + b*x))^(1/2),x)

[Out] int(sin(a + b*x)^4/(d*cos(a + b*x))^(1/2), x)

$$3.217 \quad \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=100

$$-\frac{24\sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^2 \sqrt{\cos(a+bx)}} + \frac{12(d \cos(a+bx))^{3/2} \sin(a+bx)}{5bd^3} + \frac{2 \sin^3(a+bx)}{bd \sqrt{d \cos(a+bx)}}$$

[Out] 12/5*(d*cos(b*x+a))^(3/2)*sin(b*x+a)/b/d^3+2*sin(b*x+a)^3/b/d/(d*cos(b*x+a))^(1/2)-24/5*(cos(1/2*a+1/2*b*x))^2^(1/2)/cos(1/2*a+1/2*b*x)*EllipticE(sin(1/2*a+1/2*b*x),2^(1/2))*(d*cos(b*x+a))^(1/2)/b/d^2/cos(b*x+a)^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2646, 2648, 2721, 2719}

$$\frac{12 \sin(a+bx)(d \cos(a+bx))^{3/2}}{5bd^3} - \frac{24E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{d \cos(a+bx)}}{5bd^2 \sqrt{\cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{bd \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^4/(d*Cos[a + b*x])^(3/2),x]

[Out] (-24*sqrt[d*cos[a + b*x]]*EllipticE[(a + b*x)/2, 2])/(5*b*d^2*sqrt[Cos[a + b*x]]) + (12*(d*cos[a + b*x])^(3/2)*Sin[a + b*x])/(5*b*d^3) + (2*Sin[a + b*x]^3)/(b*d*sqrt[d*cos[a + b*x]])

Rule 2646

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(-a)*(b*cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{3/2}} dx &= \frac{2 \sin^3(a+bx)}{bd \sqrt{d \cos(a+bx)}} - \frac{6 \int \sqrt{d \cos(a+bx)} \sin^2(a+bx) dx}{d^2} \\ &= \frac{12(d \cos(a+bx))^{3/2} \sin(a+bx)}{5bd^3} + \frac{2 \sin^3(a+bx)}{bd \sqrt{d \cos(a+bx)}} - \frac{12 \int \sqrt{d \cos(a+bx)} dx}{5d^2} \\ &= \frac{12(d \cos(a+bx))^{3/2} \sin(a+bx)}{5bd^3} + \frac{2 \sin^3(a+bx)}{bd \sqrt{d \cos(a+bx)}} - \frac{(12 \sqrt{d \cos(a+bx)}) \int dx}{5d^2 \sqrt{\cos(a+bx)}} \\ &= -\frac{24 \sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \middle| 2\right)}{5bd^2 \sqrt{\cos(a+bx)}} + \frac{12(d \cos(a+bx))^{3/2} \sin(a+bx)}{5bd^3} + \frac{2 \sin^3(a+bx)}{bd \sqrt{d \cos(a+bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 60, normalized size = 0.60

$$\frac{\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{5}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a+bx)\right) \sin^5(a+bx)}{5bd \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^4/(d*Cos[a + b*x])^(3/2), x]
```

```
[Out] ((Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[5/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^5)/(5*b*d*Sqrt[d*Cos[a + b*x]])
```

Maple [A]

time = 0.17, size = 213, normalized size = 2.13

method	result
default	$\frac{8 \sqrt{-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d \left(-2 \left(\sin^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 2 \cos \left(\frac{bx}{2} + \frac{a}{2} \right) \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 3 \sqrt{d \cos \left(\frac{bx}{2} + \frac{a}{2} \right)} \right)}{5d \sqrt{-d \left(2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right)}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x+a)^4/(d*cos(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
[Out] -8/5/d*(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(1/2)*(-2*sin(1/2
*b*x+1/2*a)^6*cos(1/2*b*x+1/2*a)+2*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^4+
3*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(c
os(1/2*b*x+1/2*a),2^(1/2))-3*sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/(-d*(
2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/sin(1/2*b*x+1/2*a)/(d*(
2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 115, normalized size = 1.15

$$\frac{2 \left(6i \sqrt{2} \sqrt{d} \cos(bx+a) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) + i \sin(bx+a))) - 6i \sqrt{2} \sqrt{d} \cos(bx+a) \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(bx+a) - i \sin(bx+a))) - \sqrt{d} \cos(bx+a) (\cos(bx+a)^2 + 5) \sin(bx+a) \right)}{5 b^2 \cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")
```

```
[Out] -2/5*(6*I*sqrt(2)*sqrt(d)*cos(b*x + a)*weierstrassZeta(-4, 0, weierstrassPI
nverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) - 6*I*sqrt(2)*sqrt(d)*cos(b*x
+ a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*si
n(b*x + a))) - sqrt(d*cos(b*x + a))*(cos(b*x + a)^2 + 5)*sin(b*x + a))/(b*d
^2*cos(b*x + a))
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)**4/(d*cos(b*x+a))**(3/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 4849 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^4}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/(d*cos(a + b*x))^(3/2),x)

[Out] int(sin(a + b*x)^4/(d*cos(a + b*x))^(3/2), x)

$$3.218 \quad \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=102

$$-\frac{8\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2\sqrt{d \cos(a+bx)}} + \frac{4\sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd^3} + \frac{2 \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}}$$

[Out] 2/3*sin(b*x+a)^3/b/d/(d*cos(b*x+a))^(3/2)-8/3*(cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x),2^(1/2))*cos(b*x+a)^(1/2)/b/d^2/(d*cos(b*x+a))^(1/2)+4/3*sin(b*x+a)*(d*cos(b*x+a))^(1/2)/b/d^3

Rubi [A]

time = 0.08, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2646, 2648, 2721, 2720}

$$\frac{4 \sin(a+bx) \sqrt{d \cos(a+bx)}}{3bd^3} - \frac{8 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2\sqrt{d \cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^4/(d*Cos[a + b*x])^(5/2), x]

[Out] (-8*sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(3*b*d^2*sqrt[d*Cos[a + b*x]]) + (4*sqrt[d*Cos[a + b*x]]*Sin[a + b*x])/(3*b*d^3) + (2*Sin[a + b*x]^3)/(3*b*d*(d*Cos[a + b*x])^(3/2))

Rule 2646

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{5/2}} dx &= \frac{2 \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{2 \int \frac{\sin^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx}{d^2} \\ &= \frac{4 \sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd^3} + \frac{2 \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{4 \int \frac{1}{\sqrt{d \cos(a+bx)}} dx}{3d^2} \\ &= \frac{4 \sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd^3} + \frac{2 \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{(4 \sqrt{\cos(a+bx)}) \int \frac{1}{\sqrt{d \cos(a+bx)}} dx}{3d^2 \sqrt{d \cos(a+bx)}} \\ &= -\frac{8 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}} + \frac{4 \sqrt{d \cos(a+bx)} \sin(a+bx)}{3bd^3} + \frac{2 \sin^3(a+bx)}{3bd(d \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 60, normalized size = 0.59

$$\frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a+bx)\right) \sin^5(a+bx)}{5bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[a + b*x]^4/(d*Cos[a + b*x])^(5/2), x]
```

```
[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[7/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^5)/(5*b*d*(d*Cos[a + b*x])^(3/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(114) = 228.

time = 0.21, size = 286, normalized size = 2.80

method	result
default	$\frac{8 \left(-2 \left(\sin^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 2 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \operatorname{EllipticF} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)^{5/2}}{3d^2 \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right)^{3/2}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(b*x+a)^4/(d*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
[Out] -8/3*(-2*sin(1/2*b*x+1/2*a)^6*cos(1/2*b*x+1/2*a)+2*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2)))*sin(1/2*b*x+1/2*a)^2+2*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^4-(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))-sin(1/2*b*x+1/2*a)^2*cos(1/2*b*x+1/2*a))/d^2*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(2*cos(1/2*b*x+1/2*a)^2-1)/(-d*(2*sin(1/2*b*x+1/2*a)^4-sin(1/2*b*x+1/2*a)^2))^(1/2)/sin(1/2*b*x+1/2*a)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(5/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 113, normalized size = 1.11

$$\frac{2(-2i\sqrt{2}\sqrt{d}\cos(bx+a)^2\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))+2i\sqrt{2}\sqrt{d}\cos(bx+a)^2\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))-\sqrt{d\cos(bx+a)}(\cos(bx+a)^2+1)\sin(bx+a))}{3bd^3\cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")
```

```
[Out] -2/3*(-2*I*sqrt(2)*sqrt(d)*cos(b*x + a)^2*weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a)) + 2*I*sqrt(2)*sqrt(d)*cos(b*x + a)^2*weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a)) - sqrt(d*cos(b*x + a))*(cos(b*x + a)^2 + 1)*sin(b*x + a))/(b*d^3*cos(b*x + a)^2)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(b*x+a)**4/(d*cos(b*x+a))**(5/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^4}{(d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/(d*cos(a + b*x))^(5/2),x)

[Out] int(sin(a + b*x)^4/(d*cos(a + b*x))^(5/2), x)

$$3.219 \quad \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=102

$$\frac{24\sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^4 \sqrt{\cos(a+bx)}} - \frac{12 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{5bd(d \cos(a+bx))^{5/2}}$$

[Out] $2/5*\sin(b*x+a)^3/b/d/(d*\cos(b*x+a))^{(5/2)}-12/5*\sin(b*x+a)/b/d^3/(d*\cos(b*x+a))^{(1/2)}+24/5*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/d^4/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2646, 2721, 2719}

$$\frac{24E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{d \cos(a+bx)}}{5bd^4 \sqrt{\cos(a+bx)}} - \frac{12 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{5bd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[a + b*x]^4/(d*Cos[a + b*x])^(7/2), x]`

[Out] $(24*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*d^4*\text{Sqrt}[\text{Cos}[a + b*x]]) - (12*\text{Sin}[a + b*x])/(5*b*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (2*\text{Sin}[a + b*x]^3)/(5*b*d*(d*\text{Cos}[a + b*x])^{(5/2)})$

Rule 2646

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{7/2}} dx &= \frac{2 \sin^3(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{6 \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{3/2}} dx}{5d^2} \\
 &= -\frac{12 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{5bd(d \cos(a+bx))^{5/2}} + \frac{12 \int \sqrt{d \cos(a+bx)} dx}{5d^4} \\
 &= -\frac{12 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{5bd(d \cos(a+bx))^{5/2}} + \frac{\left(12 \sqrt{d \cos(a+bx)}\right) \int \sqrt{\cos(a+bx)} dx}{5d^4 \sqrt{\cos(a+bx)}} \\
 &= \frac{24 \sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^4 \sqrt{\cos(a+bx)}} - \frac{12 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2 \sin^3(a+bx)}{5bd(d \cos(a+bx))^{5/2}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.04, size = 65, normalized size = 0.64

$$\frac{\cos^3(a+bx) \sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{9}{4}, \frac{5}{2}; \frac{7}{2}; \sin^2(a+bx)\right) \sin^5(a+bx)}{5b(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^4/(d*Cos[a + b*x])^(7/2), x]

[Out] (Cos[a + b*x]^3*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[9/4, 5/2, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^5)/(5*b*(d*Cos[a + b*x])^(7/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 365 vs. 2(114) = 228.

time = 0.32, size = 366, normalized size = 3.59

method	result
default	$ \frac{8 \sqrt{d} \left(2 \cos^2\left(\frac{bx}{2} + \frac{a}{2}\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \left(12 \operatorname{EllipticE}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right), \sqrt{2}\right) \sqrt{2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1}\right)}{5bd^4 \sqrt{\cos(a+bx)}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^4/(d*cos(b*x+a))^(7/2), x, method=_RETURNVERBOSE)

[Out] -8/5*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/d^4/sin(1/2*b*x+1/2*a)^3/(8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)*(12*EllipticE(cos(1/2*b*x+1/2*a), 2^(1/2))*(2*sin(1/2*b*x+1/2*a)

$$\begin{aligned} & ^{-2-1} \wedge (1/2) * (\sin(1/2*b*x+1/2*a)^2)^{\wedge(1/2)} * \sin(1/2*b*x+1/2*a)^4 - 14 * \sin(1/2*b*x+1/2*a)^6 * \cos(1/2*b*x+1/2*a) - 12 * \text{EllipticE}(\cos(1/2*b*x+1/2*a), 2^{\wedge(1/2)}) * (2 * \sin(1/2*b*x+1/2*a)^{-2-1})^{\wedge(1/2)} * (\sin(1/2*b*x+1/2*a)^2)^{\wedge(1/2)} * \sin(1/2*b*x+1/2*a)^2 + 14 * \cos(1/2*b*x+1/2*a) * \sin(1/2*b*x+1/2*a)^4 + 3 * (2 * \sin(1/2*b*x+1/2*a)^{-2-1})^{\wedge(1/2)} * (\sin(1/2*b*x+1/2*a)^2)^{\wedge(1/2)} * \text{EllipticE}(\cos(1/2*b*x+1/2*a), 2^{\wedge(1/2)}) - 3 * \sin(1/2*b*x+1/2*a)^2 * \cos(1/2*b*x+1/2*a) * (-2 * \sin(1/2*b*x+1/2*a)^4 * d + \sin(1/2*b*x+1/2*a)^2 * d)^{\wedge(1/2)} / (d * (2 * \cos(1/2*b*x+1/2*a)^{-2-1})^{\wedge(1/2)}) / b \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(7/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 120, normalized size = 1.18

$$\frac{2(-6i\sqrt{2}\sqrt{d}\cos(bx+a)^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))) + 6i\sqrt{2}\sqrt{d}\cos(bx+a)^3\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))) + \sqrt{d\cos(bx+a)}(7\cos(bx+a)^2-1)\sin(bx+a))}{5bd^4\cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")

[Out]
$$\frac{-2/5 * (-6 * I * \text{sqrt}(2) * \text{sqrt}(d) * \cos(b*x + a)^3 * \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I * \sin(b*x + a))) + 6 * I * \text{sqrt}(2) * \text{sqrt}(d) * \cos(b*x + a)^3 * \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I * \sin(b*x + a))) + \text{sqrt}(d * \cos(b*x + a)) * (7 * \cos(b*x + a)^2 - 1) * \sin(b*x + a)}}{(b * d^4 * \cos(b*x + a)^3)}$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**4/(d*cos(b*x+a))**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^4}{(d \cos(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/(d*cos(a + b*x))^(7/2),x)

[Out] int(sin(a + b*x)^4/(d*cos(a + b*x))^(7/2), x)

$$3.220 \quad \int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=102

$$\frac{8\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{7bd^4\sqrt{d \cos(a+bx)}} - \frac{4 \sin(a+bx)}{7bd^3(d \cos(a+bx))^{3/2}} + \frac{2 \sin^3(a+bx)}{7bd(d \cos(a+bx))^{7/2}}$$

[Out] $-4/7*\sin(b*x+a)/b/d^3/(d*\cos(b*x+a))^{(3/2)}+2/7*\sin(b*x+a)^3/b/d/(d*\cos(b*x+a))^{(7/2)}+8/7*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/d^4/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2646, 2721, 2720}

$$\frac{8\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{7bd^4\sqrt{d \cos(a+bx)}} - \frac{4 \sin(a+bx)}{7bd^3(d \cos(a+bx))^{3/2}} + \frac{2 \sin^3(a+bx)}{7bd(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^4/(d*Cos[a + b*x])^(9/2), x]

[Out] $(8*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(7*b*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]) - (4*\text{Sin}[a + b*x])/(7*b*d^3*(d*\text{Cos}[a + b*x])^{(3/2)}) + (2*\text{Sin}[a + b*x]^3)/(7*b*d*(d*\text{Cos}[a + b*x])^{(7/2)})$

Rule 2646

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^4(a+bx)}{(d \cos(a+bx))^{9/2}} dx &= \frac{2 \sin^3(a+bx)}{7bd(d \cos(a+bx))^{7/2}} - \frac{6 \int \frac{\sin^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx}{7d^2} \\
&= -\frac{4 \sin(a+bx)}{7bd^3(d \cos(a+bx))^{3/2}} + \frac{2 \sin^3(a+bx)}{7bd(d \cos(a+bx))^{7/2}} + \frac{4 \int \frac{1}{\sqrt{d \cos(a+bx)}} dx}{7d^4} \\
&= -\frac{4 \sin(a+bx)}{7bd^3(d \cos(a+bx))^{3/2}} + \frac{2 \sin^3(a+bx)}{7bd(d \cos(a+bx))^{7/2}} + \frac{\left(4 \sqrt{\cos(a+bx)}\right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{7d^4 \sqrt{d \cos(a+bx)}} \\
&= \frac{8 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{7bd^4 \sqrt{d \cos(a+bx)}} - \frac{4 \sin(a+bx)}{7bd^3(d \cos(a+bx))^{3/2}} + \frac{2 \sin^3(a+bx)}{7bd(d \cos(a+bx))^{7/2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 65, normalized size = 0.64

$$\frac{\cos^3(a+bx) \cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{5}{2}, \frac{11}{4}; \frac{7}{2}; \sin^2(a+bx)\right) \sin^5(a+bx)}{5b(d \cos(a+bx))^{9/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^4/(d*Cos[a + b*x])^(9/2), x]

[Out] (Cos[a + b*x]^3*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[5/2, 11/4, 7/2, Sin[a + b*x]^2]*Sin[a + b*x]^5)/(5*b*(d*Cos[a + b*x])^(9/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(114) = 228.

time = 0.29, size = 398, normalized size = 3.90

method	result
default	$8 \left(8 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \sqrt{2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1} \operatorname{EllipticF} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \left(\sin^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 12 \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^4/(d*cos(b*x+a))^(9/2), x, method=_RETURNVERBOSE)

[Out] 8/7*(8*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))*sin(1/2*b*x+1/2*a)^6-12*(sin(1/2*b*x+1/2*a)

$$\begin{aligned} & \left. \right)^2 \right)^{1/2} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{1/2} * \text{EllipticF}(\cos(1/2 * b * x + 1/2 * a), 2^{1/2}) \\ & * \sin(1/2 * b * x + 1/2 * a)^4 - 6 * \sin(1/2 * b * x + 1/2 * a)^6 * \cos(1/2 * b * x + 1/2 * a) + 6 * (\sin \\ & (1/2 * b * x + 1/2 * a)^2)^{1/2} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{1/2} * \text{EllipticF}(\cos(1/2 \\ & * b * x + 1/2 * a), 2^{1/2}) * \sin(1/2 * b * x + 1/2 * a)^2 + 6 * \cos(1/2 * b * x + 1/2 * a) * \sin(1/2 * b * x + \\ & 1/2 * a)^4 - (\sin(1/2 * b * x + 1/2 * a)^2)^{1/2} * (2 * \sin(1/2 * b * x + 1/2 * a)^2 - 1)^{1/2} * \text{Elli} \\ & \text{pticF}(\cos(1/2 * b * x + 1/2 * a), 2^{1/2}) - \sin(1/2 * b * x + 1/2 * a)^2 * \cos(1/2 * b * x + 1/2 * a) / \\ & d^4 * (d * (2 * \cos(1/2 * b * x + 1/2 * a)^2 - 1) * \sin(1/2 * b * x + 1/2 * a)^2)^{1/2} / (2 * \cos(1/2 * b * \\ & x + 1/2 * a)^2 - 1)^3 / (-d * (2 * \sin(1/2 * b * x + 1/2 * a)^4 - \sin(1/2 * b * x + 1/2 * a)^2))^{1/2} / \sin \\ & (1/2 * b * x + 1/2 * a) / (d * (2 * \cos(1/2 * b * x + 1/2 * a)^2 - 1))^{1/2} / b \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(9/2),x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(9/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 114, normalized size = 1.12

$$\frac{2(2i\sqrt{2}\sqrt{d}\cos(bx+a)^4\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))-2i\sqrt{2}\sqrt{d}\cos(bx+a)^4\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))+\sqrt{d\cos(bx+a)}(3\cos(bx+a)^2-1)\sin(bx+a))}{7bd^5\cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")

[Out] $-2/7 * (2 * I * \sqrt{2}) * \sqrt{d} * \cos(b * x + a)^4 * \text{weierstrassPInverse}(-4, 0, \cos(b * x + a) + I * \sin(b * x + a)) - 2 * I * \sqrt{2} * \sqrt{d} * \cos(b * x + a)^4 * \text{weierstrassPInverse}(-4, 0, \cos(b * x + a) - I * \sin(b * x + a)) + \sqrt{d * \cos(b * x + a)} * (3 * \cos(b * x + a)^2 - 1) * \sin(b * x + a) / (b * d^5 * \cos(b * x + a)^4)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**4/(d*cos(b*x+a))**(9/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4847 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^4/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^4/(d*cos(b*x + a))^(9/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(a + bx)^4}{(d \cos(a + bx))^{9/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4/(d*cos(a + b*x))^(9/2),x)

[Out] int(sin(a + b*x)^4/(d*cos(a + b*x))^(9/2), x)

3.221 $\int \cos^{\frac{3}{2}}(a + bx) \sin^5(a + bx) dx$

Optimal. Leaf size=52

$$-\frac{2 \cos^{\frac{5}{2}}(a + bx)}{5b} + \frac{4 \cos^{\frac{9}{2}}(a + bx)}{9b} - \frac{2 \cos^{\frac{13}{2}}(a + bx)}{13b}$$

[Out] $-2/5*\cos(b*x+a)^{(5/2)}/b+4/9*\cos(b*x+a)^{(9/2)}/b-2/13*\cos(b*x+a)^{(13/2)}/b$

Rubi [A]

time = 0.02, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 276}

$$-\frac{2 \cos^{\frac{13}{2}}(a + bx)}{13b} + \frac{4 \cos^{\frac{9}{2}}(a + bx)}{9b} - \frac{2 \cos^{\frac{5}{2}}(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^(3/2)*Sin[a + b*x]^5,x]`

[Out] $(-2*\text{Cos}[a + b*x]^{(5/2)})/(5*b) + (4*\text{Cos}[a + b*x]^{(9/2)})/(9*b) - (2*\text{Cos}[a + b*x]^{(13/2)})/(13*b)$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2645

`Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])`

Rubi steps

$$\begin{aligned} \int \cos^{\frac{3}{2}}(a + bx) \sin^5(a + bx) dx &= -\frac{\text{Subst}\left(\int x^{3/2}(1 - x^2)^2 dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{\text{Subst}\left(\int (x^{3/2} - 2x^{7/2} + x^{11/2}) dx, x, \cos(a + bx)\right)}{b} \\ &= -\frac{2 \cos^{\frac{5}{2}}(a + bx)}{5b} + \frac{4 \cos^{\frac{9}{2}}(a + bx)}{9b} - \frac{2 \cos^{\frac{13}{2}}(a + bx)}{13b} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 111 vs. $2(52) = 104$.

time = 0.19, size = 111, normalized size = 2.13

$$\frac{2\sqrt{\cos(ax+bx)} \left(32 - 32\sqrt[4]{\cos^2(ax+bx)} - 8\sqrt[4]{\cos^2(ax+bx)} \sin^2(ax+bx) - 5\sqrt[4]{\cos^2(ax+bx)} \sin^4(ax+bx) + 45\sqrt[4]{\cos^2(ax+bx)} \sin^6(ax+bx) \right)}{585b\sqrt[4]{\cos^2(ax+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(3/2)*Sin[a + b*x]^5,x]

[Out] (2*Sqrt[Cos[a + b*x]]*(32 - 32*(Cos[a + b*x]^2)^(1/4) - 8*(Cos[a + b*x]^2)^(1/4)*Sin[a + b*x]^2 - 5*(Cos[a + b*x]^2)^(1/4)*Sin[a + b*x]^4 + 45*(Cos[a + b*x]^2)^(1/4)*Sin[a + b*x]^6))/(585*b*(Cos[a + b*x]^2)^(1/4))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 102 vs. $2(40) = 80$.

time = 0.18, size = 103, normalized size = 1.98

method	result
default	$-\frac{32\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1}\left(180\left(\sin^{12}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 540\left(\sin^{10}\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 545\left(\sin^8\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 190\left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 3\left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) + 1\right)}{585b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(3/2)*sin(b*x+a)^5,x,method=_RETURNVERBOSE)

[Out] -32/585*(-2*sin(1/2*b*x+1/2*a)^2+1)^(1/2)*(180*sin(1/2*b*x+1/2*a)^12-540*sin(1/2*b*x+1/2*a)^10+545*sin(1/2*b*x+1/2*a)^8-190*sin(1/2*b*x+1/2*a)^6+3*sin(1/2*b*x+1/2*a)^4+2*sin(1/2*b*x+1/2*a)^2+2)/b

Maxima [A]

time = 0.30, size = 36, normalized size = 0.69

$$\frac{2\left(45\cos^{\frac{13}{2}}(bx+a) - 130\cos^{\frac{9}{2}}(bx+a) + 117\cos^{\frac{5}{2}}(bx+a)\right)}{585b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(3/2)*sin(b*x+a)^5,x, algorithm="maxima")

[Out] -2/585*(45*cos(b*x + a)^(13/2) - 130*cos(b*x + a)^(9/2) + 117*cos(b*x + a)^(5/2))/b

Fricas [A]

time = 0.36, size = 44, normalized size = 0.85

$$\frac{2\left(45\cos^6(bx+a) - 130\cos^4(bx+a) + 117\cos^2(bx+a)\right)\sqrt{\cos(bx+a)}}{585b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(3/2)*sin(b*x+a)^5,x, algorithm="fricas")`

[Out] $-2/585*(45*\cos(b*x + a)^6 - 130*\cos(b*x + a)^4 + 117*\cos(b*x + a)^2)*\sqrt{\cos(b*x + a)}/b$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**(3/2)*sin(b*x+a)**5,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 5988 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(3/2)*sin(b*x+a)^5,x, algorithm="giac")`

[Out] `integrate(cos(b*x + a)^(3/2)*sin(b*x + a)^5, x)`

Mupad [B]

time = 0.64, size = 35, normalized size = 0.67

$$\frac{2 \cos(a + b x)^{5/2} \left(\frac{5 \cos(a + b x)^4}{13} - \frac{10 \cos(a + b x)^2}{9} + 1 \right)}{5 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^(3/2)*sin(a + b*x)^5,x)`

[Out] $-(2*\cos(a + b*x)^(5/2)*((5*\cos(a + b*x)^4)/13 - (10*\cos(a + b*x)^2)/9 + 1))/ (5*b)$

3.222 $\int (d \cos(a + bx))^{9/2} \csc(a + bx) dx$

Optimal. Leaf size=100

$$\frac{d^{9/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{9/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d^3 (d \cos(a + bx))^{3/2}}{3b} + \frac{2d(d \cos(a + bx))^{7/2}}{7b}$$

[Out] $d^{(9/2)}*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-d^{(9/2)}*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+2/3*d^3*(d*\cos(b*x+a))^{(3/2)}/b+2/7*d*(d*\cos(b*x+a))^{(7/2)}/b$

Rubi [A]

time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 327, 335, 304, 209, 212}

$$\frac{d^{9/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{9/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d^3 (d \cos(a + bx))^{3/2}}{3b} + \frac{2d(d \cos(a + bx))^{7/2}}{7b}$$

Antiderivative was successfully verified.

[In] `Int[(d*Cos[a + b*x])^(9/2)*Csc[a + b*x], x]`

[Out] $(d^{(9/2)}*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/b - (d^{(9/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/b + (2*d^3*(d*\operatorname{Cos}[a + b*x])^{(3/2)})/(3*b) + (2*d*(d*\operatorname{Cos}[a + b*x])^{(7/2)})/(7*b)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 304

`Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int (d \cos(ax + b))^{9/2} \csc(ax + b) dx &= -\frac{\text{Subst}\left(\int \frac{x^{9/2}}{1-x^2/d^2} dx, x, d \cos(ax + b)\right)}{bd} \\
&= \frac{2d(d \cos(ax + b))^{7/2}}{7b} - \frac{d \text{Subst}\left(\int \frac{x^{5/2}}{1-x^2/d^2} dx, x, d \cos(ax + b)\right)}{b} \\
&= \frac{2d^3(d \cos(ax + b))^{3/2}}{3b} + \frac{2d(d \cos(ax + b))^{7/2}}{7b} - \frac{d^3 \text{Subst}\left(\int \frac{\sqrt{x}}{1-x^2/d^2} dx, x, d \cos(ax + b)\right)}{b} \\
&= \frac{2d^3(d \cos(ax + b))^{3/2}}{3b} + \frac{2d(d \cos(ax + b))^{7/2}}{7b} - \frac{(2d^3) \text{Subst}\left(\int \frac{x^2}{1-x^4/d^2} dx, x, d \cos(ax + b)\right)}{b} \\
&= \frac{2d^3(d \cos(ax + b))^{3/2}}{3b} + \frac{2d(d \cos(ax + b))^{7/2}}{7b} - \frac{d^5 \text{Subst}\left(\int \frac{1}{d-x^2} dx, x, d \cos(ax + b)\right)}{b} \\
&= \frac{d^{9/2} \tan^{-1}\left(\frac{\sqrt{d \cos(ax + b)}}{\sqrt{d}}\right)}{b} - \frac{d^{9/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(ax + b)}}{\sqrt{d}}\right)}{b} + \dots
\end{aligned}$$

Mathematica [A]

time = 0.16, size = 83, normalized size = 0.83

$$\frac{d^4 \sqrt{d \cos(a + bx)} \left(21 \tan^{-1} \left(\sqrt{\cos(a + bx)} \right) - 21 \tanh^{-1} \left(\sqrt{\cos(a + bx)} \right) + 2 \cos^{\frac{3}{2}}(a + bx) (7 + 3 \cos^2(a + bx)) \right)}{21b \sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(9/2)*Csc[a + b*x], x]`

```
[Out] (d^4*Sqrt[d*Cos[a + b*x]]*(21*ArcTan[Sqrt[Cos[a + b*x]]] - 21*ArcTanh[Sqrt[Cos[a + b*x]]] + 2*Cos[a + b*x]^(3/2)*(7 + 3*Cos[a + b*x]^2)))/(21*b*Sqrt[Cos[a + b*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 331 vs. 2(80) = 160.

time = 0.42, size = 332, normalized size = 3.32

method	result
default	$96 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} \sqrt{-d} d^4 \left(\sin^6 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 21 d^{\frac{9}{2}} \ln \left(\frac{2 \sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} d + d}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 1} \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*cos(b*x+a))^(9/2)*csc(b*x+a), x, method=_RETURNVERBOSE)`

```
[Out] -1/42/(-d)^(1/2)*(96*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)*d^4*sin(1/2*b*x+1/2*a)^6+21*d^(9/2)*ln(2/(cos(1/2*b*x+1/2*a)-1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*(-d)^(1/2)+21*d^(9/2)*ln(2/(cos(1/2*b*x+1/2*a)+1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*(-d)^(1/2)-144*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)*d^4*sin(1/2*b*x+1/2*a)^4+128*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)*d^4*sin(1/2*b*x+1/2*a)^2-40*d^4*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)+42*d^5*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))/b
```

Maxima [A]

time = 0.51, size = 98, normalized size = 0.98

$$\frac{42 d^{\frac{11}{2}} \arctan \left(\frac{\sqrt{d \cos(bx + a)}}{\sqrt{d}} \right) + 21 d^{\frac{11}{2}} \log \left(\frac{\sqrt{d \cos(bx + a)} - \sqrt{d}}{\sqrt{d \cos(bx + a)} + \sqrt{d}} \right) + 12 (d \cos(bx + a))^{\frac{7}{2}} d^2 + 28 (d \cos(bx + a))^{\frac{3}{2}} d^4}{42 bd}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a), x, algorithm="maxima")`

[Out] $1/42*(42*d^{(11/2)}*\arctan(\sqrt{d*\cos(b*x + a)})/\sqrt{d}) + 21*d^{(11/2)}*\log((\sqrt{d*\cos(b*x + a)} - \sqrt{d})/(\sqrt{d*\cos(b*x + a)} + \sqrt{d})) + 12*(d*\cos(b*x + a))^{(7/2)}*d^2 + 28*(d*\cos(b*x + a))^{(3/2)}*d^4/(b*d)$

Fricas [A]

time = 0.48, size = 313, normalized size = 3.13

$$\frac{42\sqrt{-d}d^4\arctan\left(\frac{\sqrt{d\cos(bx+a)}\sqrt{-d}}{\sin(bx+a)}\right) + 21\sqrt{-d}d^4\log\left(\frac{-\sin(bx+a)\sqrt{d\cos(bx+a)}\sqrt{-d}\cos(bx+a)-d\sin(bx+a)}{\sin(bx+a)\sqrt{d\cos(bx+a)}}\right) + 8(3d^4\cos(bx+a)^2 + 7d^4\cos(bx+a))\sqrt{d\cos(bx+a)} - 42d^4\arctan\left(\frac{\sqrt{d\cos(bx+a)}\sqrt{-d}}{\sin(bx+a)}\right) - 21d^4\log\left(\frac{-\sin(bx+a)\sqrt{d\cos(bx+a)}\sqrt{-d}\cos(bx+a)-d\sin(bx+a)}{\sin(bx+a)\sqrt{d\cos(bx+a)}}\right) - 8(3d^4\cos(bx+a)^2 + 7d^4\cos(bx+a))\sqrt{d\cos(bx+a)}}{84}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a),x, algorithm="fricas")`

[Out] $[1/84*(42*\sqrt{-d}*d^4*\arctan(2*\sqrt{d*\cos(b*x + a)})*\sqrt{-d}/(d*\cos(b*x + a) + d)) + 21*\sqrt{-d}*d^4*\log(-(d*\cos(b*x + a))^2 + 4*\sqrt{d*\cos(b*x + a)}*\sqrt{-d}*(\cos(b*x + a) - 1) - 6*d*\cos(b*x + a) + d)/(\cos(b*x + a)^2 + 2*\cos(b*x + a) + 1)) + 8*(3*d^4*\cos(b*x + a)^3 + 7*d^4*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}/b, -1/84*(42*d^{(9/2)}*\arctan(2*\sqrt{d*\cos(b*x + a)})*\sqrt{d}/(d*\cos(b*x + a) - d)) - 21*d^{(9/2)}*\log(-(d*\cos(b*x + a))^2 - 4*\sqrt{d*\cos(b*x + a)}*\sqrt{d}*(\cos(b*x + a) + 1) + 6*d*\cos(b*x + a) + d)/(\cos(b*x + a)^2 - 2*\cos(b*x + a) + 1)) - 8*(3*d^4*\cos(b*x + a)^3 + 7*d^4*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}/b]$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a),x, algorithm="giac")`

[Out] `integrate((d*cos(b*x + a))^(9/2)*csc(b*x + a), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{9/2}}{\sin(a + b x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(9/2)/sin(a + b*x),x)
```

```
[Out] int((d*cos(a + b*x))^(9/2)/sin(a + b*x), x)
```


3.223 $\int (d \cos(a + bx))^{7/2} \csc(a + bx) dx$

Optimal. Leaf size=99

$$\frac{d^{7/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{7/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d^3 \sqrt{d \cos(a + bx)}}{b} + \frac{2d(d \cos(a + bx))^{5/2}}{5b}$$

[Out] $-d^{7/2} \arctan((d \cos(bx+a))^{1/2}/d^{1/2})/b - d^{7/2} \operatorname{arctanh}((d \cos(bx+a))^{1/2}/d^{1/2})/b + 2/5 d^3 (d \cos(bx+a))^{5/2}/b + 2d^3 (d \cos(bx+a))^{1/2}/b$

Rubi [A]

time = 0.05, antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 327, 335, 218, 212, 209}

$$\frac{d^{7/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{7/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d^3 \sqrt{d \cos(a + bx)}}{b} + \frac{2d(d \cos(a + bx))^{5/2}}{5b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d \cos[a + b x])^{7/2} \csc[a + b x], x]$

[Out] $-((d^{7/2} \operatorname{ArcTan}[\sqrt{d \cos[a + b x]}/\sqrt{d}])/b) - (d^{7/2} \operatorname{ArcTanh}[\sqrt{d \cos[a + b x]}/\sqrt{d}])/b + (2d^3 \sqrt{d \cos[a + b x]})/b + (2d^3 (d \cos[a + b x])^{5/2})/(5b)$

Rule 209

$\operatorname{Int}[(a_1 + (b_1)(x_1)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a_1, 2] \operatorname{Rt}[b_1, 2])) \operatorname{ArcTan}[\operatorname{Rt}[b_1, 2](x/\operatorname{Rt}[a_1, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_1 + (b_1)(x_1)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a_1, 2] \operatorname{Rt}[-b_1, 2])) \operatorname{ArcTanh}[\operatorname{Rt}[-b_1, 2](x/\operatorname{Rt}[a_1, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_1 + (b_1)(x_1)^4)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2a), \operatorname{Int}[1/(r - s x^2), x], x] + \operatorname{Dist}[r/(2a), \operatorname{Int}[1/(r + s x^2), x], x] /; \operatorname{FreeQ}\{a, b, x\} \&\& \operatorname{!GtQ}[a/b, 0]$

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{7/2} \csc(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{7/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{bd} \\
&= \frac{2d(d \cos(a + bx))^{5/2}}{5b} - \frac{d \text{Subst}\left(\int \frac{x^{3/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{b} \\
&= \frac{2d^3 \sqrt{d \cos(a + bx)}}{b} + \frac{2d(d \cos(a + bx))^{5/2}}{5b} - \frac{d^3 \text{Subst}\left(\int \frac{1}{\sqrt{x}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{b} \\
&= \frac{2d^3 \sqrt{d \cos(a + bx)}}{b} + \frac{2d(d \cos(a + bx))^{5/2}}{5b} - \frac{(2d^3) \text{Subst}\left(\int \frac{1}{1-\frac{x^4}{d^2}} dx, x, d \cos(a + bx)\right)}{b} \\
&= \frac{2d^3 \sqrt{d \cos(a + bx)}}{b} + \frac{2d(d \cos(a + bx))^{5/2}}{5b} - \frac{d^4 \text{Subst}\left(\int \frac{1}{d-x^2} dx, x, d \cos(a + bx)\right)}{b} \\
&= -\frac{d^{7/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{7/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \dots
\end{aligned}$$

Mathematica [A]

time = 0.15, size = 80, normalized size = 0.81

$$\frac{d^3 \sqrt{d \cos(a + bx)} \left(-5 \tan^{-1} \left(\sqrt{\cos(a + bx)} \right) - 5 \tanh^{-1} \left(\sqrt{\cos(a + bx)} \right) + \sqrt{\cos(a + bx)} (11 + \cos(2(a + bx))) \right)}{5b \sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(7/2)*Csc[a + b*x], x]

[Out] (d^3*sqrt[d*cos[a + b*x]]*(-5*ArcTan[Sqrt[Cos[a + b*x]]] - 5*ArcTanh[Sqrt[Cos[a + b*x]]] + Sqrt[Cos[a + b*x]]*(11 + Cos[2*(a + b*x)])))/(5*b*sqrt[Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 292 vs. 2(81) = 162.

time = 0.36, size = 293, normalized size = 2.96

method	result
default	$- \frac{5d^{\frac{7}{2}} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d + 4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 1} \right)}{\sqrt{-d} - 16 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} \sqrt{d}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(7/2)*csc(b*x+a), x, method=_RETURNVERBOSE)

[Out] -1/10/(-d)^(1/2)*(5*d^(7/2)*ln(2/(cos(1/2*b*x+1/2*a)-1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*(-d)^(1/2)-16*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)*d^3*sin(1/2*b*x+1/2*a)^4+5*d^(7/2)*ln(2/(cos(1/2*b*x+1/2*a)+1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*(-d)^(1/2)+16*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)*d^3*sin(1/2*b*x+1/2*a)^2-24*d^3*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)-10*d^4*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))/b

Maxima [A]

time = 0.52, size = 98, normalized size = 0.99

$$\frac{10 d^{\frac{9}{2}} \arctan \left(\frac{\sqrt{d \cos(bx + a)}}{\sqrt{d}} \right) - 5 d^{\frac{9}{2}} \log \left(\frac{\sqrt{d \cos(bx + a)} - \sqrt{d}}{\sqrt{d \cos(bx + a)} + \sqrt{d}} \right) - 4 (d \cos(bx + a))^{\frac{5}{2}} d^2 - 20 \sqrt{d \cos(bx + a)} d^4}{10bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a), x, algorithm="maxima")

[Out] $-1/10*(10*d^{(9/2)}*\arctan(\sqrt{d*\cos(b*x + a)})/\sqrt{d}) - 5*d^{(9/2)}*\log((\sqrt{d*\cos(b*x + a)} - \sqrt{d})/(\sqrt{d*\cos(b*x + a)} + \sqrt{d})) - 4*(d*\cos(b*x + a))^{(5/2)}*d^2 - 20*\sqrt{d*\cos(b*x + a)}*d^4/(b*d)$

Fricas [A]

time = 0.47, size = 299, normalized size = 3.02

$$\left[\frac{10\sqrt{-d} d^4 \arctan\left(\frac{1/\sqrt{d}\cos(bx+a)\sqrt{-d}}{d\cos(bx+a)}\right) + 5\sqrt{-d} d^4 \log\left(\frac{d\cos(bx+a)\sqrt{-d}\cos(bx+a)-1-d\cos(bx+a)}{d\cos(bx+a)^2\cos(bx+a)}\right) + 8(d^4\cos(bx+a)^2 + 5d^4)\sqrt{d}\cos(bx+a)}{20b} \frac{10d^4 \arctan\left(\frac{1/\sqrt{d}\cos(bx+a)\sqrt{d}}{d\cos(bx+a)}\right) + 5d^4 \log\left(\frac{d\cos(bx+a)\sqrt{d}\cos(bx+a)+1+d\cos(bx+a)}{d\cos(bx+a)^2\cos(bx+a)}\right) + 8(d^4\cos(bx+a)^2 + 5d^4)\sqrt{d}\cos(bx+a)}{20b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a),x, algorithm="fricas")`

[Out] $[1/20*(10*\sqrt{-d}*d^3*\arctan(2*\sqrt{d*\cos(b*x + a)})*\sqrt{-d}/(d*\cos(b*x + a) + d)) + 5*\sqrt{-d}*d^3*\log(-(d*\cos(b*x + a))^2 - 4*\sqrt{d*\cos(b*x + a)}*\sqrt{-d}*(\cos(b*x + a) - 1) - 6*d*\cos(b*x + a) + d)/(\cos(b*x + a)^2 + 2*\cos(b*x + a) + 1)) + 8*(d^3*\cos(b*x + a)^2 + 5*d^3)*\sqrt{d*\cos(b*x + a)})/b, 1/20*(10*d^{(7/2)}*\arctan(2*\sqrt{d*\cos(b*x + a)})*\sqrt{d}/(d*\cos(b*x + a) - d)) + 5*d^{(7/2)}*\log(-(d*\cos(b*x + a))^2 - 4*\sqrt{d*\cos(b*x + a)}*\sqrt{d}*(\cos(b*x + a) + 1) + 6*d*\cos(b*x + a) + d)/(\cos(b*x + a)^2 - 2*\cos(b*x + a) + 1)) + 8*(d^3*\cos(b*x + a)^2 + 5*d^3)*\sqrt{d*\cos(b*x + a)})/b]$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(7/2)*csc(b*x+a),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a),x, algorithm="giac")`

[Out] `integrate((d*cos(b*x + a))^(7/2)*csc(b*x + a), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + bx))^{7/2}}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(a + b*x))^(7/2)/sin(a + b*x),x)`

[Out] `int((d*cos(a + b*x))^(7/2)/sin(a + b*x), x)`

3.224 $\int (d \cos(a + bx))^{5/2} \csc(a + bx) dx$

Optimal. Leaf size=78

$$\frac{d^{5/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{5/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d(d \cos(a + bx))^{3/2}}{3b}$$

[Out] $d^{(5/2)*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-d^{(5/2)*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+2/3*d*(d*\cos(b*x+a))^{(3/2)}/b}$

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 327, 335, 304, 209, 212}

$$\frac{d^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{5/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d(d \cos(a + bx))^{3/2}}{3b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d*\operatorname{Cos}[a + b*x])^{(5/2)}*Csc[a + b*x], x]$

[Out] $(d^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/b - (d^{(5/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/b + (2*d*(d*\operatorname{Cos}[a + b*x])^{(3/2)})/(3*b)}$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}[(x_)^2/((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& !\operatorname{GtQ}[a/b, 0]$

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{5/2} \csc(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{5/2}}{1-x^2/d^2} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2d(d \cos(a + bx))^{3/2}}{3b} - \frac{d \text{Subst}\left(\int \frac{\sqrt{x}}{1-x^2/d^2} dx, x, d \cos(a + bx)\right)}{b} \\ &= \frac{2d(d \cos(a + bx))^{3/2}}{3b} - \frac{(2d) \text{Subst}\left(\int \frac{x^2}{1-x^4/d^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} \\ &= \frac{2d(d \cos(a + bx))^{3/2}}{3b} - \frac{d^3 \text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} + \frac{d^3 S}{b} \\ &= \frac{d^{5/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{5/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2}{b} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 68, normalized size = 0.87

$$\frac{(d \cos(a + bx))^{5/2} \left(3 \tan^{-1}\left(\sqrt{\cos(a + bx)}\right) - 3 \tanh^{-1}\left(\sqrt{\cos(a + bx)}\right) + 2 \cos^{\frac{3}{2}}(a + bx)\right)}{3b \cos^{\frac{5}{2}}(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(5/2)*Csc[a + b*x], x]

[Out] ((d*cos[a + b*x])^(5/2)*(3*ArcTan[Sqrt[Cos[a + b*x]]] - 3*ArcTanh[Sqrt[Cos[a + b*x]]] + 2*cos[a + b*x]^(3/2)))/(3*b*cos[a + b*x]^(5/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(62) = 124.

time = 0.36, size = 254, normalized size = 3.26

method	result
default	$-\frac{3d^{\frac{5}{2}} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right) \sqrt{-d} + 3d^{\frac{5}{2}} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(5/2)*csc(b*x+a), x, method=_RETURNVERBOSE)

[Out]
$$-1/6/(-d)^{(1/2)}*(3*d^{(5/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*(-d)^{(1/2)}+3*d^{(5/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*(-d)^{(1/2)}+8*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}*(-d)^{(1/2)}*d^2*\sin(1/2*b*x+1/2*a)^2-4*d^2*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}*(-d)^{(1/2)}+6*d^3*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-d))/b$$

Maxima [A]

time = 0.49, size = 83, normalized size = 1.06

$$\frac{6 d^{\frac{7}{2}} \arctan \left(\frac{\sqrt{d \cos (b x+a)}}{\sqrt{d}} \right) + 3 d^{\frac{7}{2}} \log \left(\frac{\sqrt{d \cos (b x+a)}-\sqrt{d}}{\sqrt{d \cos (b x+a)}+\sqrt{d}} \right) + 4 (d \cos (b x+a))^{\frac{3}{2}} d^2}{6 b d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a), x, algorithm="maxima")

[Out]
$$1/6*(6*d^{(7/2)}*arctan(sqrt(d*cos(b*x + a))/sqrt(d)) + 3*d^{(7/2)}*log((sqrt(d*cos(b*x + a)) - sqrt(d))/(sqrt(d*cos(b*x + a)) + sqrt(d))) + 4*(d*cos(b*x + a))^{(3/2)}*d^2)/(b*d)$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(62) = 124.

time = 0.49, size = 281, normalized size = 3.60

$$\frac{6 \sqrt{-d} d^2 \arctan \left(\frac{{}_2\sqrt{d \cos (b x+a)} \sqrt{-d}}{d \cos (b x+a)+d} \right) + 8 \sqrt{d \cos (b x+a)} d^2 \cos (b x+a) + 3 \sqrt{-d} d^2 \log \left(\frac{-d \cos (b x+a)+\sqrt{d \cos (b x+a)} \sqrt{-d} \cos (b x+a)-1-d \cos (b x+a)}{\cos (b x+a)+2 \cos (b x+a)+1} \right) - 6 d^2 \arctan \left(\frac{{}_2\sqrt{d \cos (b x+a)} \sqrt{-d}}{d \cos (b x+a)-d} \right) - 8 \sqrt{d \cos (b x+a)} d^2 \cos (b x+a) - 3 d^2 \log \left(\frac{-d \cos (b x+a)-\sqrt{d \cos (b x+a)} \sqrt{-d} \cos (b x+a)+1+d \cos (b x+a)}{\cos (b x+a)-2 \cos (b x+a)+1} \right)}{12 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a),x, algorithm="fricas")

[Out] [1/12*(6*sqrt(-d)*d^2*arctan(2*sqrt(d*cos(b*x + a))*sqrt(-d)/(d*cos(b*x + a) + d)) + 8*sqrt(d*cos(b*x + a))*d^2*cos(b*x + a) + 3*sqrt(-d)*d^2*log(-(d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1) - 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)))/b, -1/12*(6*d^(5/2)*arctan(2*sqrt(d*cos(b*x + a))*sqrt(d)/(d*cos(b*x + a) - d)) - 8*sqrt(d*cos(b*x + a))*d^2*cos(b*x + a) - 3*d^(5/2)*log(-(d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)))/b]

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(5/2)*csc(b*x+a),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(5/2)*csc(b*x + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + bx))^{5/2}}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(5/2)/sin(a + b*x),x)

[Out] int((d*cos(a + b*x))^(5/2)/sin(a + b*x), x)

3.225 $\int (d \cos(a + bx))^{3/2} \csc(a + bx) dx$

Optimal. Leaf size=77

$$-\frac{d^{3/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{3/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d \sqrt{d \cos(a + bx)}}{b}$$

[Out] $-d^{3/2} \arctan((d \cos(bx+a))^{1/2}/d^{1/2})/b - d^{3/2} \operatorname{arctanh}((d \cos(bx+a))^{1/2}/d^{1/2})/b + 2d \sqrt{d \cos(bx+a)}^{1/2}/b$

Rubi [A]

time = 0.04, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 327, 335, 218, 212, 209}

$$-\frac{d^{3/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{3/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \frac{2d \sqrt{d \cos(a + bx)}}{b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d \cos[a + b x])^{3/2} \csc[a + b x], x]$

[Out] $-(d^{3/2} \operatorname{ArcTan}[\operatorname{Sqrt}[d \cos[a + b x]]/\operatorname{Sqrt}[d]])/b - (d^{3/2} \operatorname{ArcTanh}[\operatorname{Sqrt}[d \cos[a + b x]]/\operatorname{Sqrt}[d]])/b + (2d \operatorname{Sqrt}[d \cos[a + b x]])/b$

Rule 209

$\operatorname{Int}[(a + b x)(x^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[b, 2])) \operatorname{ArcTan}[\operatorname{Rt}[b, 2](x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a + b x)(x^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2] \operatorname{Rt}[-b, 2])) \operatorname{ArcTanh}[\operatorname{Rt}[-b, 2](x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a + b x)(x^4)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2a), \operatorname{Int}[1/(r - s x^2), x], x] + \operatorname{Dist}[r/(2a), \operatorname{Int}[1/(r + s x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{3/2} \csc(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{3/2}}{1-x^2} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2d\sqrt{d \cos(a + bx)}}{b} - \frac{d \text{Subst}\left(\int \frac{1}{\sqrt{x}(1-x^2)} dx, x, d \cos(a + bx)\right)}{b} \\ &= \frac{2d\sqrt{d \cos(a + bx)}}{b} - \frac{(2d) \text{Subst}\left(\int \frac{1}{1-x^4} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} \\ &= \frac{2d\sqrt{d \cos(a + bx)}}{b} - \frac{d^2 \text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} - \frac{d^2 \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} \\ &= -\frac{d^{3/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{d^{3/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} + \end{aligned}$$

Mathematica [A]

time = 0.06, size = 65, normalized size = 0.84

$$\frac{\left(-\tan^{-1}\left(\sqrt{\cos(a + bx)}\right) - \tanh^{-1}\left(\sqrt{\cos(a + bx)}\right) + 2\sqrt{\cos(a + bx)}\right) (d \cos(a + bx))^{3/2}}{b \cos^{3/2}(a + bx)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(3/2)*Csc[a + b*x],x]

[Out] ((-ArcTan[Sqrt[Cos[a + b*x]]] - ArcTanh[Sqrt[Cos[a + b*x]]] + 2*Sqrt[Cos[a + b*x]])*(d*Cos[a + b*x])^(3/2))/(b*Cos[a + b*x]^(3/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(63) = 126.

time = 0.44, size = 211, normalized size = 2.74

method	result
default	$-\frac{d^{\frac{3}{2}} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right) \sqrt{-d} + d^{\frac{3}{2}} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 1} \right) \sqrt{-d}}{2bd}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(3/2)*csc(b*x+a),x,method=_RETURNVERBOSE)

[Out] -1/2/(-d)^(1/2)*(d^(3/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*(-d)^(1/2)+d^(3/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*(-d)^(1/2)-4*d*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*(-d)^(1/2)-2*d^2*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))/b

Maxima [A]

time = 0.50, size = 83, normalized size = 1.08

$$\frac{2d^{\frac{5}{2}} \arctan \left(\frac{\sqrt{d} \cos(bx+a)}{\sqrt{d}} \right) - d^{\frac{5}{2}} \log \left(\frac{\sqrt{d} \cos(bx+a) - \sqrt{d}}{\sqrt{d} \cos(bx+a) + \sqrt{d}} \right) - 4 \sqrt{d} \cos(bx+a) d^2}{2bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a),x, algorithm="maxima")

[Out] -1/2*(2*d^(5/2)*arctan(sqrt(d*cos(b*x + a))/sqrt(d)) - d^(5/2)*log((sqrt(d*cos(b*x + a)) - sqrt(d))/(sqrt(d*cos(b*x + a)) + sqrt(d))) - 4*sqrt(d*cos(b*x + a))*d^2)/(b*d)

Fricas [A]

time = 0.47, size = 259, normalized size = 3.36

$$\frac{2\sqrt{-d} d \arctan \left(\frac{{}_2\sqrt{d} \cos(bx+a) \sqrt{-d}}{d \cos(bx+a)} \right) + \sqrt{-d} d \log \left(\frac{d \cos(bx+a)^2 - 4 \sqrt{d} \cos(bx+a) \sqrt{-d} \cos(bx+a) - 1 - d \cos(bx+a)}{\cos(bx+a)^2 - 2 \cos(bx+a) + 1} \right) + 8 \sqrt{d} \cos(bx+a) d}{4b} - \frac{2d^{\frac{5}{2}} \arctan \left(\frac{{}_2\sqrt{d} \cos(bx+a) \sqrt{d}}{d \cos(bx+a)} \right) + d^{\frac{5}{2}} \log \left(\frac{-d \cos(bx+a)^2 - 4 \sqrt{d} \cos(bx+a) \sqrt{d} \cos(bx+a) + 1 + d \cos(bx+a)}{\cos(bx+a)^2 - 2 \cos(bx+a) + 1} \right) + 8 \sqrt{d} \cos(bx+a) d}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(-d)*d*arctan(2*sqrt(d*cos(b*x + a))*sqrt(-d)/(d*cos(b*x + a) + d)) + sqrt(-d)*d*log(-(d*cos(b*x + a))^2 - 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1) - 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*d)/b, 1/4*(2*d^(3/2)*arctan(2*sqrt(d*cos(b*x + a))*sqrt(d)/(d*cos(b*x + a) - d)) + d^(3/2)*log(-(d*cos(b*x + a))^2 - 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*d)/b]

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(3/2)*csc(b*x+a),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{3/2}}{\sin(a + b x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(3/2)/sin(a + b*x),x)

[Out] int((d*cos(a + b*x))^(3/2)/sin(a + b*x), x)

3.226 $\int \sqrt{d \cos(a + bx)} \csc(a + bx) dx$

Optimal. Leaf size=58

$$\frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{\sqrt{d} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b}$$

[Out] $\arctan((d \cos(bx+a))^{1/2}/d^{1/2}) * d^{1/2}/b - \operatorname{arctanh}((d \cos(bx+a))^{1/2}/d^{1/2}) * d^{1/2}/b$

Rubi [A]

time = 0.04, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2645, 335, 304, 209, 212}

$$\frac{\sqrt{d} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{\sqrt{d} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x],x]`

[Out] $(\operatorname{Sqrt}[d] * \operatorname{ArcTan}[\operatorname{Sqrt}[d \cos[a + b*x]]/\operatorname{Sqrt}[d]])/b - (\operatorname{Sqrt}[d] * \operatorname{ArcTanh}[\operatorname{Sqrt}[d \cos[a + b*x]]/\operatorname{Sqrt}[d]])/b$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 304

`Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 335

```
Int[((c_.)*(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} \csc(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{\sqrt{x}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{2\text{Subst}\left(\int \frac{x^2}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd} \\ &= -\frac{d\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} + \frac{d\text{Subst}\left(\int \frac{1}{d+x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} \\ &= \frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} - \frac{\sqrt{d} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 51, normalized size = 0.88

$$\frac{\left(\tan^{-1}\left(\sqrt{\cos(a + bx)}\right) - \tanh^{-1}\left(\sqrt{\cos(a + bx)}\right)\right) \sqrt{d \cos(a + bx)}}{b \sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x], x]
```

```
[Out] ((ArcTan[Sqrt[Cos[a + b*x]]] - ArcTanh[Sqrt[Cos[a + b*x]]])*Sqrt[d*Cos[a +
b*x]])/(b*Sqrt[Cos[a + b*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(46) = 92.

time = 0.38, size = 183, normalized size = 3.16

method	result
default	$\frac{\sqrt{d} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right)}{2\sqrt{-d} b} + \sqrt{-d} + \sqrt{d} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right)}{2\sqrt{-d} b} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(1/2)*csc(b*x+a),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/(-d)^{(1/2)}*(d^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*(-d)^{(1/2)}+d^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*(-d)^{(1/2)}+2*d*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-d))/b$$

Maxima [A]

time = 0.51, size = 67, normalized size = 1.16

$$\frac{2d^{\frac{3}{2}} \arctan \left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}} \right) + d^{\frac{3}{2}} \log \left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}} \right)}{2bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a),x, algorithm="maxima")`

[Out]
$$1/2*(2*d^{(3/2)}*\arctan(\sqrt{d*\cos(b*x+a)})/\sqrt{d}) + d^{(3/2)}*\log((\sqrt{d*\cos(b*x+a)} - \sqrt{d})/(\sqrt{d*\cos(b*x+a)} + \sqrt{d}))/b*d$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 113 vs. 2(46) = 92.

time = 0.40, size = 238, normalized size = 4.10

$$\left[\frac{2\sqrt{-d} \arctan \left(\frac{\sqrt{d \cos(bx+a)} \sqrt{-d} (\cos(bx+a)+1)}{2d \cos(bx+a)} \right) + \sqrt{-d} \log \left(\frac{d \cos(bx+a)^2 + \sqrt{d \cos(bx+a)} \sqrt{-d} (\cos(bx+a)-1) - d \cos(bx+a) + d}{\cos(bx+a)^2 + 2 \cos(bx+a) + 1} \right)}{4b}, \frac{2\sqrt{d} \arctan \left(\frac{\sqrt{d \cos(bx+a)} (\cos(bx+a)-1)}{2\sqrt{d} \cos(bx+a)} \right) + \sqrt{d} \log \left(\frac{d \cos(bx+a)^2 - \sqrt{d \cos(bx+a)} \sqrt{d} (\cos(bx+a)+1) + 6d \cos(bx+a) + d}{\cos(bx+a)^2 - 2 \cos(bx+a) + 1} \right)}{4b} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a),x, algorithm="fricas")`

[Out]
$$[1/4*(2*\sqrt{-d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)})*\sqrt{-d}*(\cos(b*x+a)+1)/(d*\cos(b*x+a)) + \sqrt{-d}*\log((d*\cos(b*x+a)^2 + 4*\sqrt{d*\cos(b*x+a)})*\sqrt{-d}*(\cos(b*x+a)-1) - 6*d*\cos(b*x+a) + d)/(\cos(b*x+a)^2 + 2*\cos(b*x+a) + 1)))/b, 1/4*(2*\sqrt{d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)}*(\cos(b*x+a)-1)/(\sqrt{d}*\cos(b*x+a))) + \sqrt{d}*\log((d*\cos(b*x+a)^2 - 4*\sqrt{d*\cos(b*x+a)})*\sqrt{d}*(\cos(b*x+a)+1) + 6*d*\cos(b*x+a) + d)/(\cos(b*x+a)^2 - 2*\cos(b*x+a) + 1)))/b]$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(a + bx)} \csc(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)*csc(b*x+a),x)

[Out] Integral(sqrt(d*cos(a + b*x))*csc(a + b*x), x)

Giac [A]

time = 4.35, size = 48, normalized size = 0.83

$$\frac{d \left(\frac{\arctan\left(\frac{\sqrt{d \cos(bx + a)}}{\sqrt{-d}}\right)}{\sqrt{-d}} + \frac{\arctan\left(\frac{\sqrt{d \cos(bx + a)}}{\sqrt{d}}\right)}{\sqrt{d}} \right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a),x, algorithm="giac")

[Out] d*(arctan(sqrt(d*cos(b*x + a))/sqrt(-d))/sqrt(-d) + arctan(sqrt(d*cos(b*x + a))/sqrt(d))/sqrt(d))/b

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{d \cos(a + bx)}}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(1/2)/sin(a + b*x),x)

[Out] int((d*cos(a + b*x))^(1/2)/sin(a + b*x), x)

$$3.227 \quad \int \frac{\csc(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=59

$$-\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{b\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{b\sqrt{d}}$$

[Out] $-\arctan((d \cos(bx+a))^{1/2}/d^{1/2})/b/d^{1/2} - \operatorname{arctanh}((d \cos(bx+a))^{1/2}/d^{1/2})/b/d^{1/2}$

Rubi [A]

time = 0.03, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2645, 335, 218, 212, 209}

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{b\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{b\sqrt{d}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]/Sqrt[d*Cos[a + b*x]],x]`

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[d \cos[a + bx]]/\operatorname{Sqrt}[d]]/(b \operatorname{Sqrt}[d])) - \operatorname{ArcTanh}[\operatorname{Sqrt}[d \cos[a + bx]]/\operatorname{Sqrt}[d]]/(b \operatorname{Sqrt}[d])$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
  ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
  Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
  , a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
  !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \frac{\csc(a + bx)}{\sqrt{d \cos(a + bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{x} \left(1 - \frac{x^2}{d^2}\right)} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{2 \text{Subst}\left(\int \frac{1}{1 - \frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \frac{1}{d - x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} - \frac{\text{Subst}\left(\int \frac{1}{d + x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{b} \\ &= -\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b\sqrt{d}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{b\sqrt{d}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 50, normalized size = 0.85

$$-\frac{\left(\tan^{-1}\left(\sqrt{\cos(a + bx)}\right) + \tanh^{-1}\left(\sqrt{\cos(a + bx)}\right)\right) \sqrt{\cos(a + bx)}}{b\sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[a + b*x]/Sqrt[d*Cos[a + b*x]], x]
```

```
[Out] -(((ArcTan[Sqrt[Cos[a + b*x]]] + ArcTanh[Sqrt[Cos[a + b*x]])]*Sqrt[Cos[a +
  b*x]])/(b*Sqrt[d*Cos[a + b*x]]))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 181 vs. $2(47) = 94$.
time = 0.43, size = 182, normalized size = 3.08

method	result
default	$\frac{\ln\left(\frac{2\sqrt{d}\sqrt{-2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)d+d}}{\cos\left(\frac{bx}{2}+\frac{a}{2}\right)+1}\right)\sqrt{-d}+\ln\left(\frac{2\sqrt{d}\sqrt{-2\left(\sin^2\left(\frac{bx}{2}+\frac{a}{2}\right)\right)d+d}}{\cos\left(\frac{bx}{2}+\frac{a}{2}\right)-1}\right)}{2\sqrt{d}\sqrt{-d}b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)/(d*cos(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/d^{(1/2)/(-d)^{(1/2)}*(\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*(-d)^{(1/2)}+\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*(-d)^{(1/2)}-2*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-d))*d^{(1/2)}/b$$

Maxima [A]

time = 0.51, size = 68, normalized size = 1.15

$$\frac{2\sqrt{d}\arctan\left(\frac{\sqrt{d}\cos(bx+a)}{\sqrt{d}}\right)-\sqrt{d}\log\left(\frac{\sqrt{d}\cos(bx+a)-\sqrt{d}}{\sqrt{d}\cos(bx+a)+\sqrt{d}}\right)}{2bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)/(d*cos(b*x+a))^(1/2),x,algorithm="maxima")`

[Out]
$$-1/2*(2*\sqrt{d}*\arctan(\sqrt{d*\cos(b*x+a)}/\sqrt{d})-\sqrt{d}*\log((\sqrt{d*\cos(b*x+a)}-\sqrt{d})/(\sqrt{d*\cos(b*x+a)}+\sqrt{d}))) / (b*d)$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 117 vs. $2(47) = 94$.

time = 0.40, size = 246, normalized size = 4.17

$$\left[\frac{2\sqrt{-d}\arctan\left(\frac{\sqrt{d}\cos(bx+a)\sqrt{-d}\cos(bx+a)+1}{2d\cos(bx+a)}\right)-\sqrt{-d}\log\left(\frac{d\cos(bx+a)^2+\sqrt{d}\cos(bx+a)\sqrt{-d}\cos(bx+a)-1-6d\cos(bx+a)+d}{\cos(bx+a)^2+2\cos(bx+a)+1}\right)}{4bd}, \frac{2\sqrt{d}\arctan\left(\frac{\sqrt{d}\cos(bx+a)\cos(bx+a)-1}{2\sqrt{d}\cos(bx+a)}\right)-\sqrt{d}\log\left(\frac{d\cos(bx+a)^2-4\sqrt{d}\cos(bx+a)\sqrt{d}\cos(bx+a)+1+6d\cos(bx+a)+d}{\cos(bx+a)^2-2\cos(bx+a)+1}\right)}{4bd}\right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)/(d*cos(b*x+a))^(1/2),x,algorithm="fricas")`

[Out]
$$[1/4*(2*\sqrt{-d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)}*\sqrt{-d}*(\cos(b*x+a)+1)/(\sqrt{d*\cos(b*x+a)}))-\sqrt{-d}*\log((d*\cos(b*x+a)^2+4*\sqrt{d*\cos(b*x+a)}))*\sqrt{-d}*(\cos(b*x+a)-1)-6*d*\cos(b*x+a)+d)/(\cos(b*x+a)^2+2*$$

$\cos(b*x + a) + 1)))/(b*d), -1/4*(2*\sqrt{d}*\arctan(1/2*\sqrt{d*\cos(b*x + a)})*(\cos(b*x + a) - 1)/(\sqrt{d}*\cos(b*x + a))) - \sqrt{d}*\log((d*\cos(b*x + a)^2 - 4*\sqrt{d*\cos(b*x + a)}*\sqrt{d}*(\cos(b*x + a) + 1) + 6*d*\cos(b*x + a) + d)/(\cos(b*x + a)^2 - 2*\cos(b*x + a) + 1)))/(b*d)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(a + bx)}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))**(1/2),x)

[Out] Integral(csc(a + b*x)/sqrt(d*cos(a + b*x)), x)

Giac [A]

time = 4.15, size = 52, normalized size = 0.88

$$\frac{d \left(\frac{\arctan\left(\frac{\sqrt{d \cos(bx + a)}}{\sqrt{-d}}\right)}{\sqrt{-d} d} - \frac{\arctan\left(\frac{\sqrt{d \cos(bx + a)}}{\sqrt{d}}\right)}{d^{3/2}} \right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] d*(arctan(sqrt(d*cos(b*x + a))/sqrt(-d))/(sqrt(-d)*d) - arctan(sqrt(d*cos(b*x + a))/sqrt(d))/d^(3/2))/b

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(a + bx) \sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(1/2)),x)

[Out] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(1/2)), x)

$$3.228 \quad \int \frac{\csc(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{3/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{3/2}} + \frac{2}{bd\sqrt{d \cos(a+bx)}}$$

[Out] arctan((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(3/2)-arctanh((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(3/2)+2/b/d/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 331, 335, 304, 209, 212}

$$\frac{\text{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{3/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{3/2}} + \frac{2}{bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]/(d*Cos[a + b*x])^(3/2), x]

[Out] ArcTan[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(b*d^(3/2)) - ArcTanh[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(b*d^(3/2)) + 2/(b*d*Sqrt[d*Cos[a + b*x]])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 331

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))], Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \frac{\csc(a + bx)}{(d \cos(a + bx))^{3/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{3/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2}{bd \sqrt{d \cos(a + bx)}} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{bd^3} \\ &= \frac{2}{bd \sqrt{d \cos(a + bx)}} - \frac{2 \text{Subst}\left(\int \frac{x^2}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd^3} \\ &= \frac{2}{bd \sqrt{d \cos(a + bx)}} - \frac{\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd} + \frac{\text{Subst}\left(\int \frac{1}{d+x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd} \\ &= \frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{bd^{3/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{bd^{3/2}} + \frac{2}{bd \sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 66, normalized size = 0.85

$$\frac{2 + \tan^{-1}\left(\sqrt{\cos(a+bx)}\right) \sqrt{\cos(a+bx)} - \tanh^{-1}\left(\sqrt{\cos(a+bx)}\right) \sqrt{\cos(a+bx)}}{bd\sqrt{d\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]/(d*Cos[a + b*x])^(3/2), x]

[Out] (2 + ArcTan[Sqrt[Cos[a + b*x]])*Sqrt[Cos[a + b*x]] - ArcTanh[Sqrt[Cos[a + b*x]])*Sqrt[Cos[a + b*x]])/(b*d*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 421 vs. 2(64) = 128.

time = 0.68, size = 422, normalized size = 5.41

method	result
default	$-\left(4d^{\frac{5}{2}} \ln\left(\frac{{}_2\sqrt{-d} \sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + d^{-2d}}}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right)}\right)\right) + 2\sqrt{-d} \ln\left(\frac{{}_2\sqrt{d} \sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + d^{-4d}}}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right)^{-1}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)/(d*cos(b*x+a))^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/2*(-(4*d^(5/2)*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)-d))+2*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*d^2+2*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*d^2)*sin(1/2*b*x+1/2*a)^2+2*d^(5/2)*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)-d)-4*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)*d^(3/2)*(-d)^(1/2)+(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*d^2+(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1)*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a))^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*d^2)/d^(7/2)/(-d)^(1/2)/(2*sin(1/2*b*x+1/2*a)^2-1)/b

Maxima [A]

time = 0.79, size = 79, normalized size = 1.01

$$\frac{2 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{\sqrt{d}} + \frac{\log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{\sqrt{d}} + \frac{4}{\sqrt{d \cos(bx+a)}}$$

$2bd$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] $\frac{1}{2} * (2 * \arctan(\sqrt{d \cos(bx + a)} / \sqrt{d}) / \sqrt{d} + \log((\sqrt{d \cos(bx + a)} - \sqrt{d}) / (\sqrt{d \cos(bx + a)} + \sqrt{d}))) / \sqrt{d} + 4 / \sqrt{d \cos(bx + a)}) / (b * d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 148 vs. 2(64) = 128.

time = 0.42, size = 309, normalized size = 3.96

$$\frac{2\sqrt{-d} \arctan\left(\frac{\sqrt{d \cos(bx+a)} \sqrt{-d} \cos(bx+a)}{2d \cos(bx+a)}\right) \cos(bx+a) - \sqrt{-d} \cos(bx+a) \log\left(\frac{d \cos(bx+a) \sqrt{d \cos(bx+a)} \sqrt{-d} \cos(bx+a) - d \cos(bx+a)}{d \cos(bx+a) \sqrt{d \cos(bx+a)} \sqrt{-d} \cos(bx+a)}\right) + 8 \sqrt{d \cos(bx+a)}}{4b^2 \cos(bx+a)} + \frac{2\sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)} \cos(bx+a)}{2\sqrt{d} \cos(bx+a)}\right) \cos(bx+a) + \sqrt{d} \cos(bx+a) \log\left(\frac{d \cos(bx+a) \sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)} + d \cos(bx+a)}{d \cos(bx+a) \sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)}}\right) + 8 \sqrt{d \cos(bx+a)}}{4b^2 \cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] $\left[\frac{1}{4} * (2 * \sqrt{-d} * \arctan(1/2 * \sqrt{d \cos(bx + a)}) * \sqrt{-d} * (\cos(bx + a) + 1) / (d \cos(bx + a))) * \cos(bx + a) - \sqrt{-d} * \cos(bx + a) * \log((d \cos(bx + a))^2 - 4 * \sqrt{d \cos(bx + a)} * \sqrt{-d} * (\cos(bx + a) - 1) - 6 * d * \cos(bx + a) + d) / (\cos(bx + a)^2 + 2 * \cos(bx + a) + 1)) + 8 * \sqrt{d \cos(bx + a)}}{(b * d^2 * \cos(bx + a))}, \frac{1}{4} * (2 * \sqrt{d} * \arctan(1/2 * \sqrt{d \cos(bx + a)}) * (\cos(bx + a) - 1) / (\sqrt{d} * \cos(bx + a))) * \cos(bx + a) + \sqrt{d} * \cos(bx + a) * \log((d \cos(bx + a))^2 - 4 * \sqrt{d \cos(bx + a)} * \sqrt{d} * (\cos(bx + a) + 1) + 6 * d * \cos(bx + a) + d) / (\cos(bx + a)^2 - 2 * \cos(bx + a) + 1)) + 8 * \sqrt{d \cos(bx + a)}}{(b * d^2 * \cos(bx + a))} \right]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(a + bx)}{(d \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(3/2),x)

[Out] Integral(csc(a + b*x)/(d*cos(a + b*x))^(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + b x) (d \cos(a + b x))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(3/2)), x)

[Out] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(3/2)), x)

$$3.229 \quad \int \frac{\csc(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=81

$$-\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{5/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{5/2}} + \frac{2}{3bd(d \cos(a+bx))^{3/2}}$$

[Out] $-\arctan((d \cos(bx+a))^{1/2}/d^{1/2})/b/d^{5/2} - \operatorname{arctanh}((d \cos(bx+a))^{1/2}/d^{1/2})/b/d^{5/2} + 2/3/b/d/(d \cos(bx+a))^{3/2}$

Rubi [A]

time = 0.05, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 331, 335, 218, 212, 209}

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{5/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{5/2}} + \frac{2}{3bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[a + b*x]/(d*\operatorname{Cos}[a + b*x])^{5/2}, x]$

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/(b*d^{5/2})) - \operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/(b*d^{5/2}) + 2/(3*b*d*(d*\operatorname{Cos}[a + b*x])^{3/2})$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& !\operatorname{GtQ}[a/b, 0]$

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \frac{\csc(a + bx)}{(d \cos(a + bx))^{5/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{5/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2}{3bd(d \cos(a + bx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{bd^3} \\ &= \frac{2}{3bd(d \cos(a + bx))^{3/2}} - \frac{2\text{Subst}\left(\int \frac{1}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd^3} \\ &= \frac{2}{3bd(d \cos(a + bx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd^2} - \frac{\text{Subst}\left(\int \frac{1}{d+x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd^2} \\ &= -\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{bd^{5/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{bd^{5/2}} + \frac{2}{3bd(d \cos(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 70, normalized size = 0.86

$$\frac{-2 + 3 \tan^{-1} \left(\sqrt{\cos(a + bx)} \right) \cos^{\frac{3}{2}}(a + bx) + 3 \tanh^{-1} \left(\sqrt{\cos(a + bx)} \right) \cos^{\frac{3}{2}}(a + bx)}{3bd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]/(d*Cos[a + b*x])^(5/2), x]

[Out] -1/3*(-2 + 3*ArcTan[Sqrt[Cos[a + b*x]]]*Cos[a + b*x]^(3/2) + 3*ArcTanh[Sqrt[Cos[a + b*x]]]*Cos[a + b*x]^(3/2))/(b*d*(d*Cos[a + b*x])^(3/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 623 vs. $2(65) = 130$.

time = 0.65, size = 624, normalized size = 7.70

method	result
default	$\frac{\left({}_{24} \ln \left(\frac{{}_2\sqrt{-d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-2d}}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right)} \right) \right) d^{\frac{7}{2}} - 12 \sqrt{-d} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{+4d}}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 1} \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)/(d*cos(b*x+a))^(5/2), x, method=_RETURNVERBOSE)

[Out] 1/6*((24*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d)*d^(7/2)-12*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d)*d^3-12*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d)*d^3)*sin(1/2*b*x+1/2*a)^4+(-24*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d)*d^(7/2)+12*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d)*d^3+12*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d)*d^3)*sin(1/2*b*x+1/2*a)^2+6*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d)*d^(7/2)+4*(-d)^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*d^(5/2)-3*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d)*d^3-3*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2))*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d)*d^3)/(-d)^(1/2)/d^(11/2)/(4*sin(1/2*b*x+1/2*a)^4-4*sin(1/2*b*x+1/2*a)^2+1)/b

Maxima [A]

time = 0.60, size = 80, normalized size = 0.99

$$\frac{6 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{d^{\frac{3}{2}}} - \frac{3 \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{d^{\frac{3}{2}}} - \frac{4}{(d \cos(bx+a))^{\frac{3}{2}}}$$

$6bd$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")

[Out] $-1/6*(6*\arctan(\sqrt{d*\cos(b*x+a)})/\sqrt{d})/d^{3/2} - 3*\log((\sqrt{d*\cos(b*x+a)} - \sqrt{d})/(\sqrt{d*\cos(b*x+a)} + \sqrt{d}))/d^{3/2} - 4/(d*\cos(b*x+a))^{3/2})/(b*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(65) = 130.

time = 0.42, size = 318, normalized size = 3.93

$$\frac{6\sqrt{-d}\arctan\left(\frac{\sqrt{d\cos(bx+a)}\sqrt{-d}\cos(bx+a)}{2d\cos(bx+a)}\right)\cos(bx+a)^2 - 3\sqrt{-d}\cos(bx+a)^2\log\left(\frac{d\cos(bx+a)\sqrt{-d}\cos(bx+a)-d\cos(bx+a)}{\cos(bx+a)^2\cos(bx+a)}\right) + 8\sqrt{d}\cos(bx+a)}{12bd^3\cos(bx+a)^2} - \frac{6\sqrt{d}\arctan\left(\frac{\sqrt{d\cos(bx+a)}\cos(bx+a)}{2\sqrt{d}\cos(bx+a)}\right)\cos(bx+a)^2 - 3\sqrt{d}\cos(bx+a)^2\log\left(\frac{d\cos(bx+a)\sqrt{d}\cos(bx+a)-d\cos(bx+a)}{\cos(bx+a)^2\cos(bx+a)}\right) - 8\sqrt{d}\cos(bx+a)}{12bd^3\cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")

[Out] $[1/12*(6*\sqrt{-d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)})*\sqrt{-d}*(\cos(b*x+a)+1)/(d*\cos(b*x+a)))*\cos(b*x+a)^2 - 3*\sqrt{-d}*\cos(b*x+a)^2*\log((d*\cos(b*x+a)^2 + 4*\sqrt{d*\cos(b*x+a)}*\sqrt{-d}*(\cos(b*x+a)-1) - 6*d*\cos(b*x+a)+d)/(\cos(b*x+a)^2 + 2*\cos(b*x+a)+1)) + 8*\sqrt{d*\cos(b*x+a)})/(b*d^3*\cos(b*x+a)^2), -1/12*(6*\sqrt{d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)})*(\cos(b*x+a)-1)/(\sqrt{d}*\cos(b*x+a)))*\cos(b*x+a)^2 - 3*\sqrt{d}*\cos(b*x+a)^2*\log((d*\cos(b*x+a)^2 - 4*\sqrt{d*\cos(b*x+a)}*\sqrt{d}*(\cos(b*x+a)+1) + 6*d*\cos(b*x+a)+d)/(\cos(b*x+a)^2 - 2*\cos(b*x+a)+1)) - 8*\sqrt{d*\cos(b*x+a)})/(b*d^3*\cos(b*x+a)^2)]$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)/(d*cos(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx) (d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(5/2)),x)

[Out] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(5/2)), x)

$$3.230 \quad \int \frac{\csc(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=100

$$\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{7/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{7/2}} + \frac{2}{5bd(d \cos(a+bx))^{5/2}} + \frac{2}{bd^3 \sqrt{d \cos(a+bx)}}$$

[Out] arctan((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(7/2)-arctanh((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(7/2)+2/5/b/d/(d*cos(b*x+a))^(5/2)+2/b/d^3/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 331, 335, 304, 209, 212}

$$\frac{\text{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{7/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{7/2}} + \frac{2}{bd^3 \sqrt{d \cos(a+bx)}} + \frac{2}{5bd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]/(d*Cos[a + b*x])^(7/2), x]

[Out] ArcTan[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(b*d^(7/2)) - ArcTanh[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(b*d^(7/2)) + 2/(5*b*d*(d*Cos[a + b*x])^(5/2)) + 2/(b*d^3*Sqrt[d*Cos[a + b*x]])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a

/b, 0]

Rule 331

```
Int[((c_.)*(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*(m + n*(p + 1) + 1)/(a*c^n*(m + 1)), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc(a+bx)}{(d \cos(a+bx))^{7/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{7/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{bd} \\
&= \frac{2}{5bd(d \cos(a+bx))^{5/2}} - \frac{\text{Subst}\left(\int \frac{1}{x^{3/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{bd^3} \\
&= \frac{2}{5bd(d \cos(a+bx))^{5/2}} + \frac{2}{bd^3 \sqrt{d \cos(a+bx)}} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a+bx)\right)}{bd^5} \\
&= \frac{2}{5bd(d \cos(a+bx))^{5/2}} + \frac{2}{bd^3 \sqrt{d \cos(a+bx)}} - \frac{2 \text{Subst}\left(\int \frac{x^2}{1-\frac{x^2}{d^2}} dx, x, \sqrt{d \cos(a+bx)}\right)}{bd^5} \\
&= \frac{2}{5bd(d \cos(a+bx))^{5/2}} + \frac{2}{bd^3 \sqrt{d \cos(a+bx)}} - \frac{\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a+bx)}\right)}{bd^3} \\
&= \frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{7/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{7/2}} + \frac{2}{5bd(d \cos(a+bx))^{5/2}}
\end{aligned}$$

Mathematica [A]

time = 0.13, size = 81, normalized size = 0.81

$$\frac{5 \tan^{-1}\left(\sqrt{\cos(a+bx)}\right) \sqrt{\cos(a+bx)} - 5 \tanh^{-1}\left(\sqrt{\cos(a+bx)}\right) \sqrt{\cos(a+bx)} + 2(5 + \sec^2(a+bx))}{5bd^3 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]/(d*Cos[a + b*x])^(7/2), x]`

```
[Out] (5*ArcTan[Sqrt[Cos[a + b*x]]]*Sqrt[Cos[a + b*x]] - 5*ArcTanh[Sqrt[Cos[a + b*x]]]*Sqrt[Cos[a + b*x]] + 2*(5 + Sec[a + b*x]^2))/(5*b*d^3*Sqrt[d*Cos[a + b*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 881 vs. 2(82) = 164.

time = 0.68, size = 882, normalized size = 8.82

method	result
--------	--------

default	$10 \ln \left(\frac{{}_2\sqrt{-d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d^{-2d}}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right)} \right) d^{\frac{9}{2}} - 24 \sqrt{-d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} d^{\frac{7}{2}} + 5 \sqrt{-d} \ln \left(\dots \right)$
---------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{10} \frac{(-d)^{1/2} / d^{15/2} / (8 \sin(1/2 b x + 1/2 a)^6 - 12 \sin(1/2 b x + 1/2 a)^4 + 6 \sin(1/2 b x + 1/2 a)^2 - 1) \cdot (10 \ln(2 / \cos(1/2 b x + 1/2 a)) \cdot ((-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - d)^{9/2} - 24 \cdot (-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} \cdot d^{7/2} + 5 \cdot (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) - 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} + 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4 + 5 \cdot (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) + 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4 - 40 \cdot (2 \cdot \ln(2 / \cos(1/2 b x + 1/2 a)) \cdot ((-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - d)^{9/2} + (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) - 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} + 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4 + (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) + 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4) \cdot \sin(1/2 b x + 1/2 a)^6 + 20 \cdot (6 \cdot \ln(2 / \cos(1/2 b x + 1/2 a)) \cdot ((-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - d)^{9/2} - 4 \cdot (-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} \cdot d^{7/2} + 3 \cdot (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) - 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} + 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4 + 3 \cdot (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) + 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4) \cdot \sin(1/2 b x + 1/2 a)^4 - 10 \cdot (6 \cdot \ln(2 / \cos(1/2 b x + 1/2 a)) \cdot ((-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - d)^{9/2} - 8 \cdot (-d)^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} \cdot d^{7/2} + 3 \cdot (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) - 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} + 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4 + 3 \cdot (-d)^{1/2} \cdot \ln(2 / (\cos(1/2 b x + 1/2 a) + 1)) \cdot (d^{1/2} \cdot (-2 \sin(1/2 b x + 1/2 a)^{2d+d})^{1/2} - 2 \cdot d \cdot \cos(1/2 b x + 1/2 a) - d) \cdot d^4) \cdot \sin(1/2 b x + 1/2 a)^2) / b$$

Maxima [A]

time = 0.56, size = 100, normalized size = 1.00

$$\frac{10 \arctan \left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}} \right)}{d^{\frac{5}{2}}} + \frac{5 \log \left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}} \right)}{d^{\frac{5}{2}}} + \frac{4 (5 d^2 \cos(bx+a)^2 + d^2)}{(d \cos(bx+a))^{\frac{5}{2}} d^2}$$

$10 b d$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")`

[Out]
$$\frac{1}{10} \cdot (10 \cdot \arctan(\sqrt{d \cos(bx+a)}) / \sqrt{d}) / d^{5/2} + 5 \cdot \log((\sqrt{d \cos(bx+a)} - \sqrt{d}) / (\sqrt{d \cos(bx+a)} + \sqrt{d})) / d^{5/2} + 4 \cdot (5 \cdot d^2 \cdot \cos(bx+a)^2 + d^2) / ((d \cdot \cos(bx+a))^{5/2} \cdot d^2) / (b \cdot d)$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 165 vs. 2(82) = 164.

time = 0.43, size = 342, normalized size = 3.42

$$\left[\frac{10\sqrt{d}\arctan\left(\frac{\sqrt{d\cos(bx+a)}\sqrt{d\cos(bx+a)}}{2d\cos(bx+a)}\right)\cos(bx+a)^2 - 5\sqrt{d}\cos(bx+a)^2\log\left(\frac{d\cos(bx+a)-\sqrt{d\cos(bx+a)}\sqrt{d\cos(bx+a)-1}-4d\cos(bx+a)^2}{d\cos(bx+a)^2-2d\cos(bx+a)+1}\right) + 5\sqrt{d}\cos(bx+a)(5\cos(bx+a)^2+1) - 10\sqrt{d}\arctan\left(\frac{\sqrt{d\cos(bx+a)}\sqrt{d\cos(bx+a)}}{2\sqrt{d}\cos(bx+a)}\right)\cos(bx+a)^2 + 5\sqrt{d}\cos(bx+a)^2\log\left(\frac{d\cos(bx+a)-\sqrt{d\cos(bx+a)}\sqrt{d\cos(bx+a)+1}-4d\cos(bx+a)^2}{d\cos(bx+a)^2-2d\cos(bx+a)+1}\right) + 5\sqrt{d}\cos(bx+a)(5\cos(bx+a)^2+1)}{20d^2\cos(bx+a)^3} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")

[Out] [1/20*(10*sqrt(-d)*arctan(1/2*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) + 1)/(d*cos(b*x + a)))*cos(b*x + a)^3 - 5*sqrt(-d)*cos(b*x + a)^3*log((d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1) - 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*(5*cos(b*x + a)^2 + 1))/(b*d^4*cos(b*x + a)^3), 1/20*(10*sqrt(d)*arctan(1/2*sqrt(d*cos(b*x + a))*(cos(b*x + a) - 1)/(sqrt(d)*cos(b*x + a)))*cos(b*x + a)^3 + 5*sqrt(d)*cos(b*x + a)^3*log((d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*(5*cos(b*x + a)^2 + 1))/(b*d^4*cos(b*x + a)^3)]

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)/(d*cos(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx) (d \cos(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(7/2)),x)

[Out] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(7/2)), x)

$$3.231 \quad \int \frac{\csc(a+bx)}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=103

$$\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{9/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{9/2}} + \frac{2}{7bd(d \cos(a+bx))^{7/2}} + \frac{2}{3bd^3(d \cos(a+bx))^{3/2}}$$

[Out] $-\arctan((d \cos(bx+a))^{1/2}/d^{1/2})/b/d^{9/2} - \operatorname{arctanh}((d \cos(bx+a))^{1/2}/d^{1/2})/b/d^{9/2} + 2/7/b/d/(d \cos(bx+a))^{7/2} + 2/3/b/d^3/(d \cos(bx+a))^{3/2}$

Rubi [A]

time = 0.05, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2645, 331, 335, 218, 212, 209}

$$\frac{\operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{9/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{bd^{9/2}} + \frac{2}{3bd^3(d \cos(a+bx))^{3/2}} + \frac{2}{7bd(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]/(d*Cos[a + b*x])^(9/2), x]`

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[d \cos[a + b*x]]/\operatorname{Sqrt}[d]]/(b*d^{9/2})) - \operatorname{ArcTanh}[\operatorname{Sqrt}[d \cos[a + b*x]]/\operatorname{Sqrt}[d]]/(b*d^{9/2}) + 2/(7*b*d*(d \cos[a + b*x])^{7/2}) + 2/(3*b*d^3*(d \cos[a + b*x])^{3/2})$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))], Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \frac{\csc(a + bx)}{(d \cos(a + bx))^{9/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{9/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{bd} \\ &= \frac{2}{7bd(d \cos(a + bx))^{7/2}} - \frac{\text{Subst}\left(\int \frac{1}{x^{5/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{bd^3} \\ &= \frac{2}{7bd(d \cos(a + bx))^{7/2}} + \frac{2}{3bd^3(d \cos(a + bx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{bd^5} \\ &= \frac{2}{7bd(d \cos(a + bx))^{7/2}} + \frac{2}{3bd^3(d \cos(a + bx))^{3/2}} - \frac{2\text{Subst}\left(\int \frac{1}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd^5} \\ &= \frac{2}{7bd(d \cos(a + bx))^{7/2}} + \frac{2}{3bd^3(d \cos(a + bx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{bd^4} \\ &= -\frac{\tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{bd^{9/2}} - \frac{\tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{bd^{9/2}} + \frac{2}{7bd(d \cos(a + bx))^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 87, normalized size = 0.84

$$\frac{-21 \tan^{-1}\left(\sqrt{\cos(a+bx)}\right) \sqrt{\cos(a+bx)} - 21 \tanh^{-1}\left(\sqrt{\cos(a+bx)}\right) \sqrt{\cos(a+bx)} + 14 \sec(a+bx) + 6 \sec^3(a+bx)}{21bd^4 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]/(d*Cos[a + b*x])^(9/2), x]

[Out] (-21*ArcTan[Sqrt[Cos[a + b*x]]]*Sqrt[Cos[a + b*x]] - 21*ArcTanh[Sqrt[Cos[a + b*x]]]*Sqrt[Cos[a + b*x]] + 14*Sec[a + b*x] + 6*Sec[a + b*x]^3)/(21*b*d^4*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1081 vs. 2(83) = 166.

time = 0.88, size = 1082, normalized size = 10.50

method	result	size
default	Expression too large to display	1082

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)/(d*cos(b*x+a))^(9/2), x, method=_RETURNVERBOSE)

[Out] 1/42/(-d)^(1/2)/d^(19/2)/(16*sin(1/2*b*x+1/2*a)^8-32*sin(1/2*b*x+1/2*a)^6+24*sin(1/2*b*x+1/2*a)^4-8*sin(1/2*b*x+1/2*a)^2+1)*(42*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))*d^(11/2)+40*(-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*d^(9/2)-21*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*d^5-21*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*d^5-336*(-2*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))*d^(11/2)+(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*d^5+(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*d^5)*sin(1/2*b*x+1/2*a)^8+672*(-2*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))*d^(11/2)+(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*d^5+(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*d^5)*sin(1/2*b*x+1/2*a)^6+56*(-6*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-d))*d^(11/2)-2*(-d)^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)*d^(9/2)+3*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)-1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+2*d*cos(1/2*b*x+1/2*a)-d))*d^5+3*(-d)^(1/2)*ln(2/(cos(1/2*b*x+1/2*a)+1))*(d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d*cos(1/2*b*x+1/2*a)-d))*d^5)*sin(1/2*b*x+1/2*a)^2-56*(-18*ln(2/cos(1/2*b*x+1/2*a))*((-d)^(1/2)*(-2*

$$\frac{\sin(1/2*b*x+1/2*a)^{2*d+d}*(1/2-d)*d^{(11/2)-2*(-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}*d^{(9/2)+9*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)+2*d*\cos(1/2*b*x+1/2*a)-d))*d^5+9*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)-2*d*\cos(1/2*b*x+1/2*a)-d))*d^5*\sin(1/2*b*x+1/2*a)^4)/b}$$

Maxima [A]

time = 0.55, size = 102, normalized size = 0.99

$$\frac{\frac{42 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{d^{\frac{7}{2}}} - \frac{21 \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{d^{\frac{7}{2}}} - \frac{4(7d^2 \cos(bx+a)^2 + 3d^2)}{(d \cos(bx+a))^{\frac{7}{2}} d^2}}{42bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(9/2),x, algorithm="maxima")

[Out] $-1/42*(42*\arctan(\sqrt{d*\cos(b*x+a)})/\sqrt{d})/d^{(7/2)} - 21*\log((\sqrt{d*\cos(b*x+a)} - \sqrt{d})/(\sqrt{d*\cos(b*x+a)} + \sqrt{d}))/d^{(7/2)} - 4*(7*d^2*\cos(b*x+a)^2 + 3*d^2)/((d*\cos(b*x+a))^{(7/2)}*d^2)/(b*d)$

Fricas [A]

time = 0.43, size = 342, normalized size = 3.32

$$\frac{42 \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) \cos(bx+a)^{-2} \sqrt{d} \cos(bx+a)^2 \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) + 8 \sqrt{d \cos(bx+a)} (7 \cos(bx+a)^2 + 3) - 42 \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) \cos(bx+a)^{-2} \sqrt{d} \cos(bx+a)^2 \log\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 8 \sqrt{d \cos(bx+a)} (7 \cos(bx+a)^2 + 3)}{84 b^2 \cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")

[Out] $[1/84*(42*\sqrt{-d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)})*\sqrt{-d}*(\cos(b*x+a)+1)/(d*\cos(b*x+a)))*\cos(b*x+a)^4 - 21*\sqrt{-d}*\cos(b*x+a)^4*\log((d*\cos(b*x+a)^2 + 4*\sqrt{d*\cos(b*x+a)}*\sqrt{-d}*(\cos(b*x+a)-1) - 6*d*\cos(b*x+a)+d)/(\cos(b*x+a)^2 + 2*\cos(b*x+a)+1)) + 8*\sqrt{d*\cos(b*x+a)}*(7*\cos(b*x+a)^2 + 3)/(b*d^5*\cos(b*x+a)^4), -1/84*(42*\sqrt{d}*\arctan(1/2*\sqrt{d*\cos(b*x+a)}*(\cos(b*x+a)-1)/(\sqrt{d}*\cos(b*x+a)))*\cos(b*x+a)^4 - 21*\sqrt{d}*\cos(b*x+a)^4*\log((d*\cos(b*x+a)^2 - 4*\sqrt{d*\cos(b*x+a)}*\sqrt{d}*(\cos(b*x+a)+1) + 6*d*\cos(b*x+a)+d)/(\cos(b*x+a)^2 - 2*\cos(b*x+a)+1)) - 8*\sqrt{d*\cos(b*x+a)}*(7*\cos(b*x+a)^2 + 3)/(b*d^5*\cos(b*x+a)^4)]$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))**(9/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)/(d*cos(b*x + a))^(9/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx) (d \cos(a + bx))^{9/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(9/2)),x)

[Out] int(1/(sin(a + b*x)*(d*cos(a + b*x))^(9/2)), x)

3.232 $\int (d \cos(a + bx))^{11/2} \csc^2(a + bx) dx$

Optimal. Leaf size=124

$$\frac{d(d \cos(a + bx))^{9/2} \csc(a + bx)}{b} - \frac{15d^6 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{7b \sqrt{d \cos(a + bx)}} - \frac{15d^5 \sqrt{d \cos(a + bx)} \sin(a + bx)}{7b}$$

[Out] $-d*(d*\cos(b*x+a))^{(9/2)}*csc(b*x+a)/b-9/7*d^3*(d*\cos(b*x+a))^{(5/2)}*\sin(b*x+a)/b-15/7*d^6*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}-15/7*d^5*\sin(b*x+a)*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.07, antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2647, 2715, 2721, 2720}

$$\frac{15d^6 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{7b \sqrt{d \cos(a + bx)}} - \frac{15d^5 \sin(a + bx) \sqrt{d \cos(a + bx)}}{7b} - \frac{9d^3 \sin(a + bx) (d \cos(a + bx))^{5/2}}{7b} - \frac{d \csc(a + bx) (d \cos(a + bx))^{9/2}}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(11/2)}*csc[a + b*x]^2, x]$

[Out] $-((d*(d*\text{Cos}[a + b*x])^{(9/2)}*csc[a + b*x])/b) - (15*d^6*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(7*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]) - (15*d^5*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(7*b) - (9*d^3*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sin}[a + b*x])/(7*b)$

Rule 2647

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_)}), x_Symbol] := \text{Simp}[a*(a*\text{Cos}[e + f*x])^{(m-1)}*((b*\text{Sin}[e + f*x])^{(n+1)})/(b*f*(n+1)), x] + \text{Dist}[a^2*((m-1)/(b^2*(n+1))), \text{Int}[(a*\text{Cos}[e + f*x])^{(m-2)}*(b*\text{Sin}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] \parallel \text{EqQ}[m + n, 0])$

Rule 2715

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_.)])^{(n_)}), x_Symbol] := \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n-1)}/(d*n), x] + \text{Dist}[b^2*((n-1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

`Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
 \int (d \cos(a + bx))^{11/2} \csc^2(a + bx) dx &= -\frac{d(d \cos(a + bx))^{9/2} \csc(a + bx)}{b} - \frac{1}{2}(9d^2) \int (d \cos(a + bx))^{7/2} dx \\
 &= -\frac{d(d \cos(a + bx))^{9/2} \csc(a + bx)}{b} - \frac{9d^3 (d \cos(a + bx))^{5/2} \sin(a + bx)}{7b} \\
 &= -\frac{d(d \cos(a + bx))^{9/2} \csc(a + bx)}{b} - \frac{15d^5 \sqrt{d \cos(a + bx)} \sin(a + bx)}{7b} \\
 &= -\frac{d(d \cos(a + bx))^{9/2} \csc(a + bx)}{b} - \frac{15d^5 \sqrt{d \cos(a + bx)} \sin(a + bx)}{7b} \\
 &= -\frac{d(d \cos(a + bx))^{9/2} \csc(a + bx)}{b} - \frac{15d^6 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{7b \sqrt{d \cos(a + bx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.27, size = 89, normalized size = 0.72

$$\frac{d^5 \sqrt{d \cos(a + bx)} \csc(a + bx) \left(\sqrt{\cos(a + bx)} (-45 + 16 \cos(2(a + bx)) + \cos(4(a + bx))) - 60 F\left(\frac{1}{2}(a + bx) \mid 2\right) \sin(a + bx) \right)}{28b \sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(11/2)*Csc[a + b*x]^2,x]`

`[Out] (d^5*Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]*(Sqrt[Cos[a + b*x]]*(-45 + 16*Cos[2*(a + b*x)] + Cos[4*(a + b*x)]) - 60*EllipticF[(a + b*x)/2, 2]*Sin[a + b*x])/(28*b*Sqrt[Cos[a + b*x]])`

Maple [A]

time = 0.64, size = 242, normalized size = 1.95

method	result
--------	--------

default	$-\frac{\sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{d^7 \sin \left(\frac{bx}{2} + \frac{a}{2} \right)} \left(-128 \left(\sin^{12} \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 384 \left(\sin^{10} \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 576 \left(\sin^8 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 14 \left(-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right) d + \dots \right)$
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(11/2)*csc(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out]
$$-1/14*(d*(2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*d^7/(-2*\sin(1/2*b*x+1/2*a)^4*d+\sin(1/2*b*x+1/2*a)^2*d)^{(3/2)}/\cos(1/2*b*x+1/2*a)*\sin(1/2*b*x+1/2*a)*(-128*\sin(1/2*b*x+1/2*a)^{12}+384*\sin(1/2*b*x+1/2*a)^{10}-576*\sin(1/2*b*x+1/2*a)^8+30*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(3/2)}*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\text{EllipticF}(\cos(1/2*b*x+1/2*a),2^{(1/2)})*\cos(1/2*b*x+1/2*a)+512*\sin(1/2*b*x+1/2*a)^6-204*\sin(1/2*b*x+1/2*a)^4+12*\sin(1/2*b*x+1/2*a)^2+7)/(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}/b$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(11/2)*csc(b*x+a)^2,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(11/2)*csc(b*x + a)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 122, normalized size = 0.98

$$\frac{15i\sqrt{2}d^{5/2}\sin(bx+a)\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))-15i\sqrt{2}d^{5/2}\sin(bx+a)\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))+2(2d^6\cos(bx+a)^4+6d^6\cos(bx+a)^2-15d^6)\sqrt{d\cos(bx+a)}}{14b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(11/2)*csc(b*x+a)^2,x, algorithm="fricas")`

[Out]
$$1/14*(15*I*\text{sqrt}(2)*d^{(11/2)}*\sin(b*x + a)*\text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a)) - 15*I*\text{sqrt}(2)*d^{(11/2)}*\sin(b*x + a)*\text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a)) + 2*(2*d^5*\cos(b*x + a)^4 + 6*d^5*\cos(b*x + a)^2 - 15*d^5)*\text{sqrt}(d*\cos(b*x + a)))/(b*\sin(b*x + a))$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(11/2)*csc(b*x+a)**2,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(11/2)*csc(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(11/2)*csc(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{11/2}}{\sin(a + b x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(11/2)/sin(a + b*x)^2,x)

[Out] int((d*cos(a + b*x))^(11/2)/sin(a + b*x)^2, x)

3.233 $\int (d \cos(a + bx))^{9/2} \csc^2(a + bx) dx$

Optimal. Leaf size=96

$$\frac{d(d \cos(a + bx))^{7/2} \csc(a + bx)}{b} - \frac{21d^4 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{5b \sqrt{\cos(a + bx)}} - \frac{7d^3 (d \cos(a + bx))^{3/2} \sin(a + bx)}{5b}$$

[Out] $-d*(d*\cos(b*x+a))^{(7/2)}*csc(b*x+a)/b-7/5*d^3*(d*\cos(b*x+a))^{(3/2)}*\sin(b*x+a)/b-21/5*d^4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2647, 2715, 2721, 2719}

$$\frac{21d^4 E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{5b \sqrt{\cos(a + bx)}} - \frac{7d^3 \sin(a + bx) (d \cos(a + bx))^{3/2}}{5b} - \frac{d \csc(a + bx) (d \cos(a + bx))^{7/2}}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(9/2)}*Csc[a + b*x]^2, x]$

[Out] $-((d*(d*\text{Cos}[a + b*x])^{(7/2)}*Csc[a + b*x])/b) - (21*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*\text{Sqrt}[\text{Cos}[a + b*x]]) - (7*d^3*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sin}[a + b*x])/(5*b)$

Rule 2647

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[a*(a*\text{Cos}[e + f*x])^{(m - 1)}*((b*\text{Sin}[e + f*x])^{(n + 1)})/(b*f*(n + 1)), x] + \text{Dist}[a^2*((m - 1)/(b^2*(n + 1))), \text{Int}[(a*\text{Cos}[e + f*x])^{(m - 2)}*(b*\text{Sin}[e + f*x])^{(n + 2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] \mid\mid \text{EqQ}[m + n, 0])$

Rule 2715

$\text{Int}[(b_.)*\sin[(c_.) + (d_.)*(x_.)]^{(n_.)}, x_Symbol] :> \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Sin}[c + d*x])^{(n - 1)}/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_.)]^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{9/2} \csc^2(a + bx) dx &= -\frac{d(d \cos(a + bx))^{7/2} \csc(a + bx)}{b} - \frac{1}{2}(7d^2) \int (d \cos(a + bx))^{5/2} dx \\ &= -\frac{d(d \cos(a + bx))^{7/2} \csc(a + bx)}{b} - \frac{7d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{5b} \\ &= -\frac{d(d \cos(a + bx))^{7/2} \csc(a + bx)}{b} - \frac{7d^3(d \cos(a + bx))^{3/2} \sin(a + bx)}{5b} \\ &= -\frac{d(d \cos(a + bx))^{7/2} \csc(a + bx)}{b} - \frac{21d^4 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx)\right)}{5b \sqrt{\cos(a + bx)}} \end{aligned}$$

Mathematica [A]

time = 0.19, size = 74, normalized size = 0.77

$$\frac{d^4 \sqrt{d \cos(a + bx)} \left(21E\left(\frac{1}{2}(a + bx) \mid 2\right) + \sqrt{\cos(a + bx)} (5 \cot(a + bx) + \sin(2(a + bx))) \right)}{5b \sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(9/2)*Csc[a + b*x]^2,x]
```

```
[Out] -1/5*(d^4*Sqrt[d*Cos[a + b*x]]*(21*EllipticE[(a + b*x)/2, 2] + Sqrt[Cos[a +
b*x]]*(5*Cot[a + b*x] + Sin[2*(a + b*x)])))/(b*Sqrt[Cos[a + b*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(110) = 220.

time = 0.75, size = 229, normalized size = 2.39

method	result
default	$\frac{\sqrt{d} \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)^{d^6} \sin \left(\frac{bx}{2} + \frac{a}{2} \right) \left(-64 \left(\sin^{10} \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 160 \left(\sin^8 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 42 \left(2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) \right) \right)}{10 \left(-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d \right)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(9/2)*csc(b*x+a)^2,x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{10} \cdot (d \cdot (2 \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 - 1) \cdot \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{1/2} \cdot d^6 / (-2 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^4 \cdot d + \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 \cdot d)^{3/2} / \cos(1/2 \cdot b \cdot x + 1/2 \cdot a) \cdot \sin(1/2 \cdot b \cdot x + 1/2 \cdot a) \cdot (-64 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^{10} + 160 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^8 + 42 \cdot (2 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 - 1)^{3/2} \cdot (\sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2)^{1/2} \cdot \text{EllipticE}(\cos(1/2 \cdot b \cdot x + 1/2 \cdot a), 2^{1/2}) \cdot \cos(1/2 \cdot b \cdot x + 1/2 \cdot a) - 104 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^6 - 4 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^4 + 22 \sin(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 - 5) / (d \cdot (2 \cos(1/2 \cdot b \cdot x + 1/2 \cdot a)^2 - 1))^{1/2} / b$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a)^2,x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(9/2)*csc(b*x + a)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 121, normalized size = 1.26

$\frac{-21i\sqrt{2}d^3\sin(bx+a)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+21i\sqrt{2}d^3\sin(bx+a)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))+2(2d^4\cos(bx+a)^3-7d^4\cos(bx+a))\sqrt{d\cos(bx+a)}}{10b\sin(bx+a)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a)^2,x, algorithm="fricas")`

[Out] $\frac{1}{10} \cdot (-21 \cdot I \cdot \sqrt{2} \cdot d^{9/2} \cdot \sin(b \cdot x + a) \cdot \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b \cdot x + a) + I \cdot \sin(b \cdot x + a))) + 21 \cdot I \cdot \sqrt{2} \cdot d^{9/2} \cdot \sin(b \cdot x + a) \cdot \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b \cdot x + a) - I \cdot \sin(b \cdot x + a))) + 2 \cdot (2 \cdot d^4 \cdot \cos(b \cdot x + a)^3 - 7 \cdot d^4 \cdot \cos(b \cdot x + a)) \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) / (b \cdot \sin(b \cdot x + a))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(9/2)*csc(b*x+a)**2,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(9/2)*csc(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{9/2}}{\sin(a + b x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(9/2)/sin(a + b*x)^2,x)

[Out] int((d*cos(a + b*x))^(9/2)/sin(a + b*x)^2, x)

3.234 $\int (d \cos(a + bx))^{7/2} \csc^2(a + bx) dx$

Optimal. Leaf size=96

$$\frac{d(d \cos(a + bx))^{5/2} \csc(a + bx)}{b} - \frac{5d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{3b \sqrt{d \cos(a + bx)}} - \frac{5d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{3b}$$

[Out] $-d*(d*\cos(b*x+a))^{(5/2)}*csc(b*x+a)/b-5/3*d^4*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*EllipticF(\sin(1/2*a+1/2*b*x),2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}-5/3*d^3*\sin(b*x+a)*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.06, antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2647, 2715, 2721, 2720}

$$\frac{5d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{3b \sqrt{d \cos(a + bx)}} - \frac{5d^3 \sin(a + bx) \sqrt{d \cos(a + bx)}}{3b} - \frac{d \csc(a + bx) (d \cos(a + bx))^{5/2}}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(7/2)}*Csc[a + b*x]^2,x]$

[Out] $-((d*(d*\text{Cos}[a + b*x])^{(5/2)}*Csc[a + b*x])/b) - (5*d^4*\text{Sqrt}[\text{Cos}[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(3*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]) - (5*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sin}[a + b*x])/(3*b)$

Rule 2647

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[a*(a*\text{Cos}[e + f*x])^{(m - 1)}*((b*\text{Sin}[e + f*x])^{(n + 1)})/(b*f*(n + 1)), x] + \text{Dist}[a^2*((m - 1)/(b^2*(n + 1))), \text{Int}[(a*\text{Cos}[e + f*x])^{(m - 2)}*(b*\text{Sin}[e + f*x])^{(n + 2)}, x], x] /; \text{FreeQ}[a, b, e, f], x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] \parallel \text{EqQ}[m + n, 0])$

Rule 2715

$\text{Int}[(b_.*\sin[(c_.) + (d_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Sin}[c + d*x])^{(n - 1)})/(d*n), x] + \text{Dist}[b^2*((n - 1)/n), \text{Int}[(b*\text{Sin}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}[b, c, d], x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[c, d], x]$

Rule 2721

```
Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])
^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ
[-1, n, 1] && IntegerQ[2*n]
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{7/2} \csc^2(a + bx) dx &= -\frac{d(d \cos(a + bx))^{5/2} \csc(a + bx)}{b} - \frac{1}{2}(5d^2) \int (d \cos(a + bx))^{3/2} dx \\ &= -\frac{d(d \cos(a + bx))^{5/2} \csc(a + bx)}{b} - \frac{5d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{3b} \\ &= -\frac{d(d \cos(a + bx))^{5/2} \csc(a + bx)}{b} - \frac{5d^3 \sqrt{d \cos(a + bx)} \sin(a + bx)}{3b} \\ &= -\frac{d(d \cos(a + bx))^{5/2} \csc(a + bx)}{b} - \frac{5d^4 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{3b \sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [A]

time = 0.17, size = 73, normalized size = 0.76

$$\frac{d^3 \sqrt{d \cos(a + bx)} \left(\sqrt{\cos(a + bx)} (-4 + \cos(2(a + bx))) \csc(a + bx) - 5F\left(\frac{1}{2}(a + bx) \mid 2\right) \right)}{3b \sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(7/2)*Csc[a + b*x]^2,x]
```

```
[Out] (d^3*Sqrt[d*Cos[a + b*x]]*(Sqrt[Cos[a + b*x]]*(-4 + Cos[2*(a + b*x)])*Csc[a + b*x] - 5*EllipticF[(a + b*x)/2, 2]))/(3*b*Sqrt[Cos[a + b*x]])
```

Maple [A]

time = 0.57, size = 216, normalized size = 2.25

method	result
default	$\frac{\sqrt{d} \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d^5 \sin \left(\frac{bx}{2} + \frac{a}{2} \right) \left(-32 \left(\sin^8 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) + 10 \left(2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right)^{\frac{3}{2}} \sqrt{\frac{1}{2}} \right)}{6 \left(-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d \right)^{\frac{3}{2}} \cos \left(\frac{bx}{2} + \frac{a}{2} \right)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(7/2)*csc(b*x+a)^2,x,method=_RETURNVERBOSE)
[Out] -1/6*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^5/(-2*sin(
1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(3/2)/cos(1/2*b*x+1/2*a)*sin(1/2
*b*x+1/2*a)*(-32*sin(1/2*b*x+1/2*a)^8+10*(2*sin(1/2*b*x+1/2*a)^2-1)^(3/2)*(
sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))*cos(1/2*b
*x+1/2*a)+64*sin(1/2*b*x+1/2*a)^6-28*sin(1/2*b*x+1/2*a)^4-4*sin(1/2*b*x+1/2
*a)^2+3)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(7/2)*csc(b*x + a)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 109, normalized size = 1.14

$$\frac{5i\sqrt{2}d^{\frac{5}{2}}\sin(bx+a)\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))-5i\sqrt{2}d^{\frac{5}{2}}\sin(bx+a)\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))+2(2d^3\cos(bx+a)^2-5d^3)\sqrt{d\cos(bx+a)}}{6b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] 1/6*(5*I*sqrt(2)*d^(7/2)*sin(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x +
a) + I*sin(b*x + a)) - 5*I*sqrt(2)*d^(7/2)*sin(b*x + a)*weierstrassPInverse
(-4, 0, cos(b*x + a) - I*sin(b*x + a)) + 2*(2*d^3*cos(b*x + a)^2 - 5*d^3)*s
qrt(d*cos(b*x + a)))/(b*sin(b*x + a))
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(7/2)*csc(b*x+a)**2,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(7/2)*csc(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{7/2}}{\sin(a + b x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(7/2)/sin(a + b*x)^2,x)

[Out] int((d*cos(a + b*x))^(7/2)/sin(a + b*x)^2, x)

3.235 $\int (d \cos(a + bx))^{5/2} \csc^2(a + bx) dx$

Optimal. Leaf size=66

$$-\frac{d(d \cos(a + bx))^{3/2} \csc(a + bx)}{b} - \frac{3d^2 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b \sqrt{\cos(a + bx)}}$$

[Out] $-d*(d*\cos(b*x+a))^{3/2}*\csc(b*x+a)/b-3*d^2*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2647, 2721, 2719}

$$-\frac{3d^2 E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{b \sqrt{\cos(a + bx)}} - \frac{d \csc(a + bx) (d \cos(a + bx))^{3/2}}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{5/2}*\text{Csc}[a + b*x]^2, x]$

[Out] $-((d*(d*\text{Cos}[a + b*x])^{3/2}*\text{Csc}[a + b*x])/b) - (3*d^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(b*\text{Sqrt}[\text{Cos}[a + b*x]])$

Rule 2647

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[a*(a*\text{Cos}[e + f*x])^{(m - 1)}*((b*\text{Sin}[e + f*x])^{(n + 1)})/(b*f*(n + 1)), x] + \text{Dist}[a^2*((m - 1)/(b^2*(n + 1))), \text{Int}[(a*\text{Cos}[e + f*x])^{(m - 2)}*(b*\text{Sin}[e + f*x])^{(n + 2)}, x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] \parallel \text{EqQ}[m + n, 0])$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 2721

$\text{Int}[(b*\sin[(c_.) + (d_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Sin}[c + d*x])^{(n)}/\text{Sin}[c + d*x]^n, \text{Int}[\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \&\& \text{LtQ}[-1, n, 1] \&\& \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{5/2} \csc^2(a + bx) dx &= -\frac{d(d \cos(a + bx))^{3/2} \csc(a + bx)}{b} - \frac{1}{2} (3d^2) \int \sqrt{d \cos(a + bx)} dx \\
&= -\frac{d(d \cos(a + bx))^{3/2} \csc(a + bx)}{b} - \frac{\left(3d^2 \sqrt{d \cos(a + bx)}\right) \int \sqrt{\cos(a + bx)} dx}{2\sqrt{\cos(a + bx)}} \\
&= -\frac{d(d \cos(a + bx))^{3/2} \csc(a + bx)}{b} - \frac{3d^2 \sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b\sqrt{\cos(a + bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 58, normalized size = 0.88

$$-\frac{(d \cos(a + bx))^{5/2} \left(\cos^{3/2}(a + bx) \csc(a + bx) + 3E\left(\frac{1}{2}(a + bx) \mid 2\right) \right)}{b \cos^{5/2}(a + bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(5/2)*Csc[a + b*x]^2,x]``[Out] -(((d*Cos[a + b*x])^(5/2)*(Cos[a + b*x]^(3/2)*Csc[a + b*x] + 3*EllipticE[(a + b*x)/2, 2]))/(b*Cos[a + b*x]^(5/2)))`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 202 vs. 2(86) = 172.

time = 0.74, size = 203, normalized size = 3.08

method	result
default	$ \frac{\sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d^4 \sin \left(\frac{bx}{2} + \frac{a}{2} \right) \left(6 \left(2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \right)^{\frac{3}{2}} \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}}{2 \left(-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d \right)^{\frac{3}{2}} \cos \left(\frac{bx}{2} + \frac{a}{2} \right) \sqrt{d} \left(2 \right)} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*cos(b*x+a))^(5/2)*csc(b*x+a)^2,x,method=_RETURNVERBOSE)`

```
[Out] 1/2*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^4/(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(3/2)/cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)*(6*(2*sin(1/2*b*x+1/2*a)^2-1)^(3/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))*cos(1/2*b*x+1/2*a)+8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(5/2)*csc(b*x + a)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 105, normalized size = 1.59

$$\frac{-3i\sqrt{2}d^{\frac{5}{2}}\sin(bx+a)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+3i\sqrt{2}d^{\frac{5}{2}}\sin(bx+a)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))-2\sqrt{d\cos(bx+a)}d^2\cos(bx+a)}{2b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] 1/2*(-3*I*sqrt(2)*d^(5/2)*sin(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + 3*I*sqrt(2)*d^(5/2)*sin(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) - 2*sqrt(d*cos(b*x + a))*d^2*cos(b*x + a)/(b*sin(b*x + a))
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(5/2)*csc(b*x+a)**2,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^2,x, algorithm="giac")
```

```
[Out] integrate((d*cos(b*x + a))^(5/2)*csc(b*x + a)^2, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(d \cos(a + bx))^{5/2}}{\sin(a + bx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(5/2)/sin(a + b*x)^2,x)
```

```
[Out] int((d*cos(a + b*x))^(5/2)/sin(a + b*x)^2, x)
```


3.236 $\int (d \cos(a + bx))^{3/2} \csc^2(a + bx) dx$

Optimal. Leaf size=66

$$-\frac{d\sqrt{d\cos(a+bx)}\csc(a+bx)}{b} - \frac{d^2\sqrt{\cos(a+bx)}F\left(\frac{1}{2}(a+bx)\mid 2\right)}{b\sqrt{d\cos(a+bx)}}$$

[Out] $-d^2(\cos(1/2*a+1/2*b*x))^{1/2}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{1/2})*\cos(b*x+a)^{1/2}/b/(d*\cos(b*x+a))^{1/2}-d*\csc(b*x+a)*(d*\cos(b*x+a))^{1/2}/b$

Rubi [A]

time = 0.04, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2647, 2721, 2720}

$$-\frac{d^2\sqrt{\cos(a+bx)}F\left(\frac{1}{2}(a+bx)\mid 2\right)}{b\sqrt{d\cos(a+bx)}} - \frac{d\csc(a+bx)\sqrt{d\cos(a+bx)}}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{3/2}*\text{Csc}[a + b*x]^2, x]$

[Out] $-((d*\text{Sqrt}[d*\text{Cos}[a + b*x)]*\text{Csc}[a + b*x])/b) - (d^2*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(b*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2647

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[a*(a*\cos[e + f*x])^{(m-1)}*((b*\sin[e + f*x])^{(n+1)})/(b*f*(n+1)), x] + \text{Dist}[a^2*((m-1)/(b^2*(n+1))), \text{Int}[(a*\cos[e + f*x])^{(m-2)}*(b*\sin[e + f*x])^{(n+2)}, x], x] /;$ $\text{FreeQ}\{a, b, e, f, x\} \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ (\text{IntegersQ}[2*m, 2*n] \ || \ \text{EqQ}[m + n, 0])$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ $\text{FreeQ}\{c, d, x\}$

Rule 2721

$\text{Int}[(b*\sin[(c_.) + (d_.)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\sin[c + d*x])^n/\sin[c + d*x]^n, \text{Int}[\sin[c + d*x]^n, x], x] /;$ $\text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{LtQ}[-1, n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{3/2} \csc^2(a + bx) dx &= -\frac{d \sqrt{d \cos(a + bx)} \csc(a + bx)}{b} - \frac{1}{2} d^2 \int \frac{1}{\sqrt{d \cos(a + bx)}} dx \\
&= -\frac{d \sqrt{d \cos(a + bx)} \csc(a + bx)}{b} - \frac{\left(d^2 \sqrt{\cos(a + bx)}\right) \int \frac{1}{\sqrt{\cos(a + bx)}} dx}{2 \sqrt{d \cos(a + bx)}} \\
&= -\frac{d \sqrt{d \cos(a + bx)} \csc(a + bx)}{b} - \frac{d^2 \sqrt{\cos(a + bx)} F\left(\frac{1}{2}(a + bx) \mid 2\right)}{b \sqrt{d \cos(a + bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 56, normalized size = 0.85

$$-\frac{(d \cos(a + bx))^{3/2} \left(\sqrt{\cos(a + bx)} \csc(a + bx) + F\left(\frac{1}{2}(a + bx) \mid 2\right) \right)}{b \cos^{3/2}(a + bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(3/2)*Csc[a + b*x]^2,x]``[Out] -(((d*Cos[a + b*x])^(3/2)*(Sqrt[Cos[a + b*x]]*Csc[a + b*x] + EllipticF[(a + b*x)/2, 2]))/(b*Cos[a + b*x]^(3/2)))`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 189 vs. 2(86) = 172.

time = 0.72, size = 190, normalized size = 2.88

method	result
default	$ -\frac{\sqrt{d} \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d^3 \sin \left(\frac{bx}{2} + \frac{a}{2} \right) \left(2 \left(2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \right)^{\frac{3}{2}} \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \text{EllipticF} \left(\frac{1}{2} \left(\frac{bx}{2} + \frac{a}{2} \right), 2 \right)}{2 \left(-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d \right)^{\frac{3}{2}} \cos \left(\frac{bx}{2} + \frac{a}{2} \right) \sqrt{d} \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right)} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*cos(b*x+a))^(3/2)*csc(b*x+a)^2,x,method=_RETURNVERBOSE)`
`[Out] -1/2*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^3/(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(3/2)/cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)*(2*(2*sin(1/2*b*x+1/2*a)^2-1)^(3/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2))*EllipticF(cos(1/2*b*x+1/2*a),2^(1/2))*cos(1/2*b*x+1/2*a)+4*sin(1/2*b*x+1/2*a)^4-4*sin(1/2*b*x+1/2*a)^2+1)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^2,x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 91, normalized size = 1.38

$$\frac{i\sqrt{2}d^{\frac{3}{2}}\sin(bx+a)\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))-i\sqrt{2}d^{\frac{3}{2}}\sin(bx+a)\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))-2\sqrt{d\cos(bx+a)}d}{2b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^2,x, algorithm="fricas")
```

```
[Out] 1/2*(I*sqrt(2)*d^(3/2)*sin(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x + a)
+ I*sin(b*x + a)) - I*sqrt(2)*d^(3/2)*sin(b*x + a)*weierstrassPInverse(-4,
0, cos(b*x + a) - I*sin(b*x + a)) - 2*sqrt(d*cos(b*x + a))*d)/(b*sin(b*x +
a))
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(3/2)*csc(b*x+a)**2,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 5006 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^2,x, algorithm="giac")
```

```
[Out] integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a)^2, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(d \cos(a + bx))^{3/2}}{\sin(a + bx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(3/2)/sin(a + b*x)^2,x)
```

```
[Out] int((d*cos(a + b*x))^(3/2)/sin(a + b*x)^2, x)
```

3.237 $\int \sqrt{d \cos(a + bx)} \csc^2(a + bx) dx$

Optimal. Leaf size=65

$$-\frac{(d \cos(a + bx))^{3/2} \csc(a + bx)}{bd} - \frac{\sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b\sqrt{\cos(a + bx)}}$$

[Out] $-(d \cos(bx+a))^{3/2} \csc(bx+a)/b/d - (\cos(1/2*a+1/2*b*x))^{1/2} / \cos(1/2*a+1/2*b*x) * \text{EllipticE}(\sin(1/2*a+1/2*b*x), 2^{1/2}) * (d \cos(bx+a))^{1/2} / b / \cos(bx+a)^{1/2}$

Rubi [A]

time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2650, 2721, 2719}

$$-\frac{\csc(a + bx)(d \cos(a + bx))^{3/2}}{bd} - \frac{E\left(\frac{1}{2}(a + bx) \mid 2\right) \sqrt{d \cos(a + bx)}}{b\sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^2,x]`

[Out] $-\left(\frac{(d \cos[a + b*x])^{3/2} \csc[a + b*x]}{b*d}\right) - \left(\frac{\text{Sqrt}[d \cos[a + b*x]] * \text{EllipticE}[(a + b*x)/2, 2]}{b \text{Sqrt}[\cos[a + b*x]]}\right)$

Rule 2650

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_.), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]`

Rubi steps

$$\begin{aligned}
\int \sqrt{d \cos(a + bx)} \csc^2(a + bx) dx &= -\frac{(d \cos(a + bx))^{3/2} \csc(a + bx)}{bd} - \frac{1}{2} \int \sqrt{d \cos(a + bx)} dx \\
&= -\frac{(d \cos(a + bx))^{3/2} \csc(a + bx)}{bd} - \frac{\sqrt{d \cos(a + bx)} \int \sqrt{\cos(a + bx)} dx}{2\sqrt{\cos(a + bx)}} \\
&= -\frac{(d \cos(a + bx))^{3/2} \csc(a + bx)}{bd} - \frac{\sqrt{d \cos(a + bx)} E\left(\frac{1}{2}(a + bx) \mid 2\right)}{b\sqrt{\cos(a + bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 56, normalized size = 0.86

$$-\frac{\sqrt{d \cos(a + bx)} \left(\cos^{\frac{3}{2}}(a + bx) \csc(a + bx) + E\left(\frac{1}{2}(a + bx) \mid 2\right) \right)}{b\sqrt{\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^2,x]

[Out] -((Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^(3/2)*Csc[a + b*x] + EllipticE[(a + b*x)/2, 2]))/(b*Sqrt[Cos[a + b*x]]))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 202 vs. 2(85) = 170.

time = 0.82, size = 203, normalized size = 3.12

method	result
default	$\frac{\sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d^2 \sin \left(\frac{bx}{2} + \frac{a}{2} \right) \left(2 \left(2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right)^{\frac{3}{2}} \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right) \text{EllipticE} \left(\frac{bx+a}{2}, 2 \right)}{2 \left(-2 \left(\sin^4 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d \right)^{\frac{3}{2}} \cos \left(\frac{bx}{2} + \frac{a}{2} \right) \sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right)}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*csc(b*x+a)^2,x,method=_RETURNVERBOSE)

[Out] 1/2*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)*d^2/(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(3/2)/cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)*(2*(2*sin(1/2*b*x+1/2*a)^2-1)^(3/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))*cos(1/2*b*x+1/2*a)+8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^2,x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*csc(b*x + a)^2, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.11, size = 102, normalized size = 1.57

$$\frac{-i\sqrt{2}\sqrt{d}\sin(bx+a)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+i\sqrt{2}\sqrt{d}\sin(bx+a)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))-2\sqrt{d\cos(bx+a)}\cos(bx+a)}{2b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^2,x, algorithm="fricas")

[Out] 1/2*(-I*sqrt(2)*sqrt(d)*sin(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) + I*sin(b*x + a))) + I*sqrt(2)*sqrt(d)*sin(b*x + a)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(b*x + a) - I*sin(b*x + a))) - 2*sqrt(d*cos(b*x + a))*cos(b*x + a)/(b*sin(b*x + a))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(a + bx)} \csc^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)**2,x)

[Out] Integral(sqrt(d*cos(a + b*x))*csc(a + b*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^2,x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*csc(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{d \cos(a + bx)}}{\sin(a + bx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(1/2)/sin(a + b*x)^2,x)

[Out] int((d*cos(a + b*x))^(1/2)/sin(a + b*x)^2, x)

$$3.238 \quad \int \frac{\csc^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=64

$$-\frac{\sqrt{d \cos(a+bx)} \csc(a+bx)}{bd} + \frac{\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{b\sqrt{d \cos(a+bx)}}$$

[Out] (cos(1/2*a+1/2*b*x)^2)^(1/2)/cos(1/2*a+1/2*b*x)*EllipticF(sin(1/2*a+1/2*b*x), 2^(1/2))*cos(b*x+a)^(1/2)/b/(d*cos(b*x+a))^(1/2)-csc(b*x+a)*(d*cos(b*x+a))^(1/2)/b/d

Rubi [A]

time = 0.04, antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$,

Rules used = {2650, 2721, 2720}

$$\frac{\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{b\sqrt{d \cos(a+bx)}} - \frac{\csc(a+bx) \sqrt{d \cos(a+bx)}}{bd}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^2/Sqrt[d*Cos[a + b*x]],x]

[Out] -((Sqrt[d*Cos[a + b*x]]*Csc[a + b*x])/(b*d)) + (Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(b*Sqrt[d*Cos[a + b*x]])

Rule 2650

```
Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol]
:> Simp[(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x]
+ Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m + 2), x], x]
/; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol]
:> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x]
/; FreeQ[{c, d}, x]
```

Rule 2721

```
Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol]
:> Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x]
/; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]
```


Rubi steps

$$\begin{aligned}
\int \frac{\csc^2(a+bx)}{\sqrt{d \cos(a+bx)}} dx &= -\frac{\sqrt{d \cos(a+bx)} \csc(a+bx)}{bd} + \frac{1}{2} \int \frac{1}{\sqrt{d \cos(a+bx)}} dx \\
&= -\frac{\sqrt{d \cos(a+bx)} \csc(a+bx)}{bd} + \frac{\sqrt{\cos(a+bx)} \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{2\sqrt{d \cos(a+bx)}} \\
&= -\frac{\sqrt{d \cos(a+bx)} \csc(a+bx)}{bd} + \frac{\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{b\sqrt{d \cos(a+bx)}}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 47, normalized size = 0.73

$$\frac{-\cot(a+bx) + \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{b\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[a + b*x]^2/Sqrt[d*Cos[a + b*x]], x]``[Out] (-Cot[a + b*x] + Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2])/(b*Sqrt[d*Cos[a + b*x]])`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(84) = 168.

time = 0.69, size = 188, normalized size = 2.94

method	result
default	$\frac{\sqrt{d \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d \sin\left(\frac{bx}{2} + \frac{a}{2}\right) \left(2 \left(2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\right)^{\frac{3}{2}} \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}}}{2 \left(-2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d\right)^{\frac{3}{2}} \cos\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{d \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(b*x+a)^2/(d*cos(b*x+a))^(1/2), x, method=_RETURNVERBOSE)`
`[Out] 1/2*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(3/2)/cos(1/2*b*x+1/2*a)*d*sin(1/2*b*x+1/2*a)*(2*(2*sin(1/2*b*x+1/2*a)^2-1)^(3/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticF(cos(1/2*b*x+1/2*a), 2^(1/2))*cos(1/2*b*x+1/2*a)-4*sin(1/2*b*x+1/2*a)^4+4*sin(1/2*b*x+1/2*a)^2-1)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(b*x + a)^2/sqrt(d*cos(b*x + a)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 93, normalized size = 1.45

$$\frac{-i\sqrt{2}\sqrt{d}\sin(bx+a)\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))+i\sqrt{2}\sqrt{d}\sin(bx+a)\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))-2\sqrt{d\cos(bx+a)}}{2bd\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/2*(-I*sqrt(2)*sqrt(d)*sin(b*x + a)*weierstrassPInverse(-4, 0, cos(b*x + a)
+ I*sin(b*x + a)) + I*sqrt(2)*sqrt(d)*sin(b*x + a)*weierstrassPInverse(-4
, 0, cos(b*x + a) - I*sin(b*x + a)) - 2*sqrt(d*cos(b*x + a)))/(b*d*sin(b*x
+ a))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(a + bx)}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)**2/(d*cos(b*x+a))**(1/2),x)
```

```
[Out] Integral(csc(a + b*x)**2/sqrt(d*cos(a + b*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(csc(b*x + a)^2/sqrt(d*cos(b*x + a)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(a + bx)^2 \sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(1/2)),x)
```

```
[Out] int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(1/2)), x)
```

$$3.239 \quad \int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=94

$$-\frac{\csc(a+bx)}{bd\sqrt{d\cos(a+bx)}} - \frac{3\sqrt{d\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{bd^2\sqrt{\cos(a+bx)}} + \frac{3\sin(a+bx)}{bd\sqrt{d\cos(a+bx)}}$$

[Out] $-\csc(b*x+a)/b/d/(d*\cos(b*x+a))^{(1/2)}+3*\sin(b*x+a)/b/d/(d*\cos(b*x+a))^{(1/2)}-3*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/d^2/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2650, 2716, 2721, 2719}

$$-\frac{3E\left(\frac{1}{2}(a+bx) \mid 2\right)\sqrt{d\cos(a+bx)}}{bd^2\sqrt{\cos(a+bx)}} + \frac{3\sin(a+bx)}{bd\sqrt{d\cos(a+bx)}} - \frac{\csc(a+bx)}{bd\sqrt{d\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^2/(d*Cos[a + b*x])^(3/2),x]`

[Out] $-(\text{Csc}[a + b*x]/(b*d*\text{Sqrt}[d*\text{Cos}[a + b*x]])) - (3*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(b*d^2*\text{Sqrt}[\text{Cos}[a + b*x]]) + (3*\text{Sin}[a + b*x])/(b*d*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2650

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])ⁿ/Sin[c + d*x]ⁿ, Int[Sin[c + d*x]ⁿ, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{3/2}} dx &= -\frac{\csc(a+bx)}{bd\sqrt{d \cos(a+bx)}} + \frac{3}{2} \int \frac{1}{(d \cos(a+bx))^{3/2}} dx \\
 &= -\frac{\csc(a+bx)}{bd\sqrt{d \cos(a+bx)}} + \frac{3 \sin(a+bx)}{bd\sqrt{d \cos(a+bx)}} - \frac{3 \int \sqrt{d \cos(a+bx)} dx}{2d^2} \\
 &= -\frac{\csc(a+bx)}{bd\sqrt{d \cos(a+bx)}} + \frac{3 \sin(a+bx)}{bd\sqrt{d \cos(a+bx)}} - \frac{(3\sqrt{d \cos(a+bx)}) \int \sqrt{\cos(a+bx)}}{2d^2 \sqrt{\cos(a+bx)}} \\
 &= -\frac{\csc(a+bx)}{bd\sqrt{d \cos(a+bx)}} - \frac{3\sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{bd^2 \sqrt{\cos(a+bx)}} + \frac{3 \sin(a+bx)}{bd\sqrt{d \cos(a+bx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.13, size = 65, normalized size = 0.69

$$\frac{-\cos(a+bx) \cot(a+bx) - 3\sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right) + 2 \sin(a+bx)}{bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2/(d*Cos[a + b*x])^(3/2), x]

[Out] (-(Cos[a + b*x]*Cot[a + b*x]) - 3*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + 2*Sin[a + b*x])/(b*d*Sqrt[d*Cos[a + b*x]])

Maple [A]

time = 0.86, size = 209, normalized size = 2.22

method	result
default	$ \frac{\sqrt{d \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} \left(-2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d\right)^{\frac{3}{2}} \left(6 \sqrt{2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)} - 2d^3 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^5 \left(2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)^2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{2d^3 \sin\left(\frac{bx}{2} + \frac{a}{2}\right)^5 \left(2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)^2 \cos\left(\frac{bx}{2} + \frac{a}{2}\right)} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^2/(d*cos(b*x+a))^(3/2), x, method=_RETURNVERBOSE)

[Out] $-1/2*(d*(2*\cos(1/2*b*x+1/2*a)^2-1)*\sin(1/2*b*x+1/2*a)^2)^{(1/2)}/d^3/\sin(1/2*b*x+1/2*a)^5/(2*\sin(1/2*b*x+1/2*a)^2-1)^2/\cos(1/2*b*x+1/2*a)*(-2*\sin(1/2*b*x+1/2*a)^4*d+\sin(1/2*b*x+1/2*a)^2*d)^{(3/2)}*(6*(2*\sin(1/2*b*x+1/2*a)^2-1)^{(1/2)}*(\sin(1/2*b*x+1/2*a)^2)^{(1/2)}*\text{EllipticE}(\cos(1/2*b*x+1/2*a),2^{(1/2)}))*\cos(1/2*b*x+1/2*a)+12*\sin(1/2*b*x+1/2*a)^4-12*\sin(1/2*b*x+1/2*a)^2+1)/(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}/b$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")`

[Out] `integrate(csc(b*x + a)^2/(d*cos(b*x + a))^(3/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 131, normalized size = 1.39

$$\frac{-3i\sqrt{2}\sqrt{d}\cos(bx+a)\sin(bx+a)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+3i\sqrt{2}\sqrt{d}\cos(bx+a)\sin(bx+a)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))-2\sqrt{d\cos(bx+a)}[3\cos(bx+a)^2-2]}{2bd^2\cos(bx+a)\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")`

[Out] $1/2*(-3*I*\sqrt{2}*\sqrt{d}*\cos(b*x + a)*\sin(b*x + a)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) + I*\sin(b*x + a))) + 3*I*\sqrt{2}*\sqrt{d}*\cos(b*x + a)*\sin(b*x + a)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(b*x + a) - I*\sin(b*x + a))) - 2*\sqrt{d*\cos(b*x + a)}*(3*\cos(b*x + a)^2 - 2))/(b*d^2*\cos(b*x + a)*\sin(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(a + bx)}{(d \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**2/(d*cos(b*x+a))**(3/2),x)`

[Out] `Integral(csc(a + b*x)**2/(d*cos(a + b*x))**(3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(3/2),x, algorithm="giac")
```

```
[Out] integrate(csc(b*x + a)^2/(d*cos(b*x + a))^(3/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^2 (d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(3/2)),x)
```

```
[Out] int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(3/2)), x)
```

$$3.240 \quad \int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=98

$$-\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{3/2}} + \frac{5\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}} + \frac{5 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}}$$

[Out] $-\csc(b*x+a)/b/d/(d*\cos(b*x+a))^{(3/2)}+5/3*\sin(b*x+a)/b/d/(d*\cos(b*x+a))^{(3/2)}$
 $+5/3*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticF}(\sin(1/2*a+1/2*b*x), 2^{(1/2)})*\cos(b*x+a)^{(1/2)}/b/d^2/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2650, 2716, 2721, 2720}

$$\frac{5\sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}} + \frac{5 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} - \frac{\csc(a+bx)}{bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^2/(d*Cos[a + b*x])^(5/2), x]`

[Out] $-(\text{Csc}[a + b*x]/(b*d*(d*\text{Cos}[a + b*x])^{(3/2)})) + (5*\text{Sqrt}[\text{Cos}[a + b*x]]*\text{EllipticF}[(a + b*x)/2, 2])/(3*b*d^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (5*\text{Sin}[a + b*x])/(3*b*d*(d*\text{Cos}[a + b*x])^{(3/2)})$

Rule 2650

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rule 2716

`Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 2721

Int[((b_)*sin[(c_.) + (d_)*(x_)])^(n_), x_Symbol] :> Dist[(b*Sin[c + d*x])ⁿ/Sin[c + d*x]ⁿ, Int[Sin[c + d*x]ⁿ, x], x] /; FreeQ[{b, c, d}, x] && LtQ [-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned} \int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{5/2}} dx &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{3/2}} + \frac{5}{2} \int \frac{1}{(d \cos(a+bx))^{5/2}} dx \\ &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{3/2}} + \frac{5 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} + \frac{5 \int \frac{1}{\sqrt{d \cos(a+bx)}} dx}{6d^2} \\ &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{3/2}} + \frac{5 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} + \frac{\left(5 \sqrt{\cos(a+bx)}\right) \int \frac{1}{\sqrt{\cos(a+bx)}} dx}{6d^2 \sqrt{d \cos(a+bx)}} \\ &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{3/2}} + \frac{5 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right)}{3bd^2 \sqrt{d \cos(a+bx)}} + \frac{5 \sin(a+bx)}{3bd(d \cos(a+bx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 62, normalized size = 0.63

$$\frac{-3 \cot(a+bx) + 5 \sqrt{\cos(a+bx)} F\left(\frac{1}{2}(a+bx) \mid 2\right) + 2 \tan(a+bx)}{3bd^2 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2/(d*Cos[a + b*x])^(5/2), x]

[Out] (-3*Cot[a + b*x] + 5*Sqrt[Cos[a + b*x]]*EllipticF[(a + b*x)/2, 2] + 2*Tan[a + b*x])/(3*b*d^2*Sqrt[d*Cos[a + b*x]])

Maple [A]

time = 0.86, size = 190, normalized size = 1.94

method	result
default	$\frac{\sqrt{d} \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right) \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \left(10 \left(2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)\right)^{\frac{3}{2}} \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \operatorname{EllipticF}\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right)}{6d \left(-2 \left(\sin^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d\right)^{\frac{3}{2}} \cos\left(\frac{bx}{2} + \frac{a}{2}\right) \sqrt{d} \left(2 \left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)\right) - 1\right)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)^2/(d*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{6} * (d * (2 * \cos(1/2 * b * x + 1/2 * a) ^ 2 - 1) * \sin(1/2 * b * x + 1/2 * a) ^ 2) ^ (1/2) / d / (-2 * \sin(1/2 * b * x + 1/2 * a) ^ 4 * d + \sin(1/2 * b * x + 1/2 * a) ^ 2 * d) ^ (3/2) / \cos(1/2 * b * x + 1/2 * a) * (10 * (2 * \sin(1/2 * b * x + 1/2 * a) ^ 2 - 1) ^ (3/2) * (\sin(1/2 * b * x + 1/2 * a) ^ 2) ^ (1/2) * \text{EllipticF}(\cos(1/2 * b * x + 1/2 * a), 2 ^ (1/2)) * \cos(1/2 * b * x + 1/2 * a) - 20 * \sin(1/2 * b * x + 1/2 * a) ^ 4 + 20 * \sin(1/2 * b * x + 1/2 * a) ^ 2 - 3) * \sin(1/2 * b * x + 1/2 * a) / (d * (2 * \cos(1/2 * b * x + 1/2 * a) ^ 2 - 1)) ^ (1/2) / b$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")`

[Out] `integrate(csc(b*x + a)^2/(d*cos(b*x + a))^(5/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 129, normalized size = 1.32

$$\frac{-5i\sqrt{2}\sqrt{d}\cos(bx+a)^2\sin(bx+a)\text{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a))+5i\sqrt{2}\sqrt{d}\cos(bx+a)^2\sin(bx+a)\text{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a))-2\sqrt{d\cos(bx+a)}(5\cos(bx+a)^2-2)}{6bd^3\cos(bx+a)^2\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")`

[Out] $\frac{1}{6} * (-5 * I * \sqrt{2} * \sqrt{d} * \cos(b * x + a) ^ 2 * \sin(b * x + a) * \text{weierstrassPInverse}(-4, 0, \cos(b * x + a) + I * \sin(b * x + a)) + 5 * I * \sqrt{2} * \sqrt{d} * \cos(b * x + a) ^ 2 * \sin(b * x + a) * \text{weierstrassPInverse}(-4, 0, \cos(b * x + a) - I * \sin(b * x + a)) - 2 * \sqrt{d * \cos(b * x + a)} * (5 * \cos(b * x + a) ^ 2 - 2)) / (b * d ^ 3 * \cos(b * x + a) ^ 2 * \sin(b * x + a))$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**2/(d*cos(b*x+a))**(5/2),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)^2/(d*cos(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^2 (d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(5/2)),x)

[Out] int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(5/2)), x)

$$3.241 \quad \int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=126

$$\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}} - \frac{21 \sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^4 \sqrt{\cos(a+bx)}} + \frac{7 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} + \frac{21 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}}$$

[Out] $-\csc(b*x+a)/b/d/(d*\cos(b*x+a))^{(5/2)}+7/5*\sin(b*x+a)/b/d/(d*\cos(b*x+a))^{(5/2)}+21/5*\sin(b*x+a)/b/d^3/(d*\cos(b*x+a))^{(1/2)}-21/5*(\cos(1/2*a+1/2*b*x)^2)^{(1/2)}/\cos(1/2*a+1/2*b*x)*\text{EllipticE}(\sin(1/2*a+1/2*b*x),2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}/b/d^4/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2650, 2716, 2721, 2719}

$$-\frac{21E\left(\frac{1}{2}(a+bx) \mid 2\right) \sqrt{d \cos(a+bx)}}{5bd^4 \sqrt{\cos(a+bx)}} + \frac{21 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{7 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} - \frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^2/(d*Cos[a + b*x])^(7/2),x]

[Out] $-(\text{Csc}[a + b*x]/(b*d*(d*\text{Cos}[a + b*x])^{(5/2)})) - (21*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[(a + b*x)/2, 2])/(5*b*d^4*\text{Sqrt}[\text{Cos}[a + b*x]]) + (7*\text{Sin}[a + b*x])/(5*b*d*(d*\text{Cos}[a + b*x])^{(5/2)}) + (21*\text{Sin}[a + b*x])/(5*b*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2650

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2716

Int[((b_.)*sin[(c_.) + (d_.)*(x_.)])^(n_.), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1))), x] + Dist[(n + 2)/(b^2*(n + 1)), Int[(b*Sin[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 2721

Int[((b_)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Dist[(b*Sin[c + d*x])^n/Sin[c + d*x]^n, Int[Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && LtQ[-1, n, 1] && IntegerQ[2*n]

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^2(a+bx)}{(d \cos(a+bx))^{7/2}} dx &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}} + \frac{7}{2} \int \frac{1}{(d \cos(a+bx))^{7/2}} dx \\
 &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}} + \frac{7 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} + \frac{21 \int \frac{1}{(d \cos(a+bx))^{3/2}} dx}{10d^2} \\
 &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}} + \frac{7 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} + \frac{21 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} - \frac{21 \int \sqrt{d \cos(a+bx)}}{5bd^3 \sqrt{d \cos(a+bx)}} \\
 &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}} + \frac{7 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}} + \frac{21 \sin(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}} - \frac{(21 \int \sqrt{d \cos(a+bx)}}{5bd^3 \sqrt{d \cos(a+bx)}}) \\
 &= -\frac{\csc(a+bx)}{bd(d \cos(a+bx))^{5/2}} - \frac{21 \sqrt{d \cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right)}{5bd^4 \sqrt{\cos(a+bx)}} + \frac{7 \sin(a+bx)}{5bd(d \cos(a+bx))^{5/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.13, size = 82, normalized size = 0.65

$$\frac{-5 \cos(a+bx) \cot(a+bx) - 21 \sqrt{\cos(a+bx)} E\left(\frac{1}{2}(a+bx) \mid 2\right) + 16 \sin(a+bx) + 2 \sec(a+bx) \tan(a+bx)}{5bd^3 \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^2/(d*Cos[a + b*x])^(7/2), x]

[Out] (-5*Cos[a + b*x]*Cot[a + b*x] - 21*Sqrt[Cos[a + b*x]]*EllipticE[(a + b*x)/2, 2] + 16*Sin[a + b*x] + 2*Sec[a + b*x]*Tan[a + b*x])/(5*b*d^3*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 407 vs. 2(136) = 272.

time = 0.98, size = 408, normalized size = 3.24

method	result
default	$\frac{\sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right) \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}{\left(168 \operatorname{EllipticE} \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right), \sqrt{2} \right) \sqrt{\frac{1}{2} - \frac{\cos(bx+a)}{2}} \right) \sqrt{2} \left(\sin \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(b*x+a)^2/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/10*(d*(2*cos(1/2*b*x+1/2*a)^2-1)*sin(1/2*b*x+1/2*a)^2)^(1/2)/d^5/(2*sin(1/2*b*x+1/2*a)^2-1)/sin(1/2*b*x+1/2*a)^5/cos(1/2*b*x+1/2*a)/(8*sin(1/2*b*x+1/2*a)^6-12*sin(1/2*b*x+1/2*a)^4+6*sin(1/2*b*x+1/2*a)^2-1)*(168*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^4+336*sin(1/2*b*x+1/2*a)^8-168*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))*(sin(1/2*b*x+1/2*a)^2)^(1/2)*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*cos(1/2*b*x+1/2*a)*sin(1/2*b*x+1/2*a)^2-672*sin(1/2*b*x+1/2*a)^6+42*(2*sin(1/2*b*x+1/2*a)^2-1)^(1/2)*(sin(1/2*b*x+1/2*a)^2)^(1/2)*EllipticE(cos(1/2*b*x+1/2*a),2^(1/2))*cos(1/2*b*x+1/2*a)+448*sin(1/2*b*x+1/2*a)^4-112*sin(1/2*b*x+1/2*a)^2+5)*(-2*sin(1/2*b*x+1/2*a)^4*d+sin(1/2*b*x+1/2*a)^2*d)^(3/2)/(d*(2*cos(1/2*b*x+1/2*a)^2-1))^(1/2)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(b*x + a)^2/(d*cos(b*x + a))^(7/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 145, normalized size = 1.15

$$\frac{-21i\sqrt{2}\sqrt{d}\cos(bx+a)^3\sin(bx+a)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)+i\sin(bx+a)))+21i\sqrt{2}\sqrt{d}\cos(bx+a)^3\sin(bx+a)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(bx+a)-i\sin(bx+a)))}{10kd^4\cos(bx+a)^3\sin(bx+a)} - 2(21\cos(bx+a)^4 - 14\cos(bx+a)^2 - 2)\sqrt{d\cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")
```

```
[Out] 1/10*(-21*I*sqrt(2)*sqrt(d)*cos(b*x + a)^3*sin(b*x + a)*weierstrassZeta(-4,0,weierstrassPInverse(-4,0,cos(b*x + a) + I*sin(b*x + a))) + 21*I*sqrt(2)*sqrt(d)*cos(b*x + a)^3*sin(b*x + a)*weierstrassZeta(-4,0,weierstrassPInverse(-4,0,cos(b*x + a) - I*sin(b*x + a))) - 2*(21*cos(b*x + a)^4 - 14*cos(b*x + a)^2 - 2)*sqrt(d*cos(b*x + a)))/(b*d^4*cos(b*x + a)^3*sin(b*x + a))
```

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**2/(d*cos(b*x+a))**(7/2), x)`

[Out] Timed out

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^2/(d*cos(b*x+a))^(7/2), x, algorithm="giac")`

[Out] `integrate(csc(b*x + a)^2/(d*cos(b*x + a))^(7/2), x)`

Mupad [F]
time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^2 (d \cos(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(7/2)), x)`

[Out] `int(1/(sin(a + b*x)^2*(d*cos(a + b*x))^(7/2)), x)`

3.242 $\int (d \cos(a + bx))^{11/2} \csc^3(a + bx) dx$

Optimal. Leaf size=135

$$\frac{9d^{11/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{9d^{11/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{9d^5 \sqrt{d \cos(a + bx)}}{2b} - \frac{9d^3 (d \cos(a + bx))^{5/2}}{10b} - \frac{d \csc^2(a + bx) (d \cos(a + bx))^{9/2}}{2b}$$

[Out] $9/4*d^{(11/2)*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+9/4*d^{(11/2)*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-9/10*d^3*(d*\cos(b*x+a))^{(5/2)}/b-1/2*d*(d*\cos(b*x+a))^{(9/2)*\csc(b*x+a)^2/b-9/2*d^5*(d*\cos(b*x+a))^{(1/2)}/b}$

Rubi [A]

time = 0.06, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$,

Rules used = {2645, 294, 327, 335, 218, 212, 209}

$$\frac{9d^{11/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{9d^{11/2} \operatorname{tanh}^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{9d^5 \sqrt{d \cos(a + bx)}}{2b} - \frac{9d^3 (d \cos(a + bx))^{5/2}}{10b} - \frac{d \csc^2(a + bx) (d \cos(a + bx))^{9/2}}{2b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d*\operatorname{Cos}[a + b*x])^{(11/2)}*\operatorname{Csc}[a + b*x]^3, x]$

[Out] $(9*d^{(11/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/(4*b) + (9*d^{(11/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/(4*b) - (9*d^5*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/(2*b) - (9*d^3*(d*\operatorname{Cos}[a + b*x])^{(5/2)})/(10*b) - (d*(d*\operatorname{Cos}[a + b*x])^{(9/2)*\operatorname{Csc}[a + b*x]^2)/(2*b)$

Rule 209

$\operatorname{Int}[(a_0 + (b_0)*(x_0)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_0 + (b_0)*(x_0)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_0 + (b_0)*(x_0)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}[\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 327

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{11/2} \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{11/2}}{(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a + bx)\right)}{bd} \\
&= -\frac{d(d \cos(a + bx))^{9/2} \csc^2(a + bx)}{2b} + \frac{(9d)\text{Subst}\left(\int \frac{x^{7/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{9d^3(d \cos(a + bx))^{5/2}}{10b} - \frac{d(d \cos(a + bx))^{9/2} \csc^2(a + bx)}{2b} + \frac{(9d^3)\text{Subst}\left(\int \frac{x^{3/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{9d^5 \sqrt{d \cos(a + bx)}}{2b} - \frac{9d^3(d \cos(a + bx))^{5/2}}{10b} - \frac{d(d \cos(a + bx))^{9/2}}{2b} \\
&= -\frac{9d^5 \sqrt{d \cos(a + bx)}}{2b} - \frac{9d^3(d \cos(a + bx))^{5/2}}{10b} - \frac{d(d \cos(a + bx))^{9/2}}{2b} \\
&= -\frac{9d^5 \sqrt{d \cos(a + bx)}}{2b} - \frac{9d^3(d \cos(a + bx))^{5/2}}{10b} - \frac{d(d \cos(a + bx))^{9/2}}{2b} \\
&= \frac{9d^{11/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{9d^{11/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b}
\end{aligned}$$

Mathematica [A]

time = 1.45, size = 137, normalized size = 1.01

$$\frac{d(d \cos(a + bx))^{9/2} \left(45 \tan^{-1}\left(\sqrt{\cos(a + bx)}\right) + 24 \tanh^{-1}\left(\sqrt{\cos(a + bx)}\right) - 2\sqrt{\cos(a + bx)}(2 \cos(2(a + bx)) + 5 \csc^2(a + bx)) - \frac{21}{2}(8\sqrt{\cos(a + bx)} + \log(1 - \sqrt{\cos(a + bx)}) - \log(1 + \sqrt{\cos(a + bx)}))\right)}{20b \cos^3(a + bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(11/2)*Csc[a + b*x]^3, x]`

```
[Out] (d*(d*Cos[a + b*x])^(9/2)*(45*ArcTan[Sqrt[Cos[a + b*x]]] + 24*ArcTanh[Sqrt[Cos[a + b*x]]] - 2*Sqrt[Cos[a + b*x]]*(2*Cos[2*(a + b*x)] + 5*Csc[a + b*x]^2) - (21*(8*Sqrt[Cos[a + b*x]] + Log[1 - Sqrt[Cos[a + b*x]]] - Log[1 + Sqrt[Cos[a + b*x]]]))/2))/(20*b*Cos[a + b*x]^(9/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 406 vs. 2(107) = 214.

time = 0.82, size = 407, normalized size = 3.01

method	result
--------	--------

default	$-\frac{8d^5 \left(\cos^4\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \sqrt{2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d - d}}{5} + \frac{8d^5 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \sqrt{2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d - d}}{5} + \frac{8d^5 \sqrt{2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d - d}}{5}$
---------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(11/2)*csc(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $(-8/5*d^5*\cos(1/2*b*x+1/2*a)^4*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}+8/5*d^5*\cos(1/2*b*x+1/2*a)^2*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}+8/5*d^5*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}-6*d^5*(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}-1/16*d^5/(\cos(1/2*b*x+1/2*a)+1)*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}+9/8*d^{(11/2)}*\ln((-4*d*\cos(1/2*b*x+1/2*a)+2*d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d)/(\cos(1/2*b*x+1/2*a)+1))+9/8*d^{(11/2)}*\ln((4*d*\cos(1/2*b*x+1/2*a)+2*d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d)/(\cos(1/2*b*x+1/2*a)-1))-1/8*d^5/\cos(1/2*b*x+1/2*a)^2*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}-9/4*d^6/(-d)^{(1/2)}*\ln((-2*d+2*(-d)^{(1/2)}*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)})/\cos(1/2*b*x+1/2*a))+1/16*d^5/(\cos(1/2*b*x+1/2*a)-1)*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}/b$

Maxima [A]

time = 0.51, size = 133, normalized size = 0.99

$$\frac{20 \sqrt{d \cos(bx+a)} d^8}{d^2 \cos(bx+a)^2 - d^2} + 90 d^{13/2} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 45 d^{13/2} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) - 16 (d \cos(bx+a))^{5/2} d^4 - 160 \sqrt{d \cos(bx+a)} d^6$$

40 bd

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(11/2)*csc(b*x+a)^3,x, algorithm="maxima")`

[Out] $1/40*(20*\sqrt{d*\cos(b*x+a)}*d^8/(d^2*\cos(b*x+a)^2-d^2)+90*d^{(13/2)}*\arctan(\sqrt{d*\cos(b*x+a)}/\sqrt{d})-45*d^{(13/2)}*\log((\sqrt{d*\cos(b*x+a)}-\sqrt{d})/(\sqrt{d*\cos(b*x+a)}+\sqrt{d}))-16*(d*\cos(b*x+a))^{(5/2)}*d^4-160*\sqrt{d*\cos(b*x+a)}*d^6)/(b*d)$

Fricas [A]

time = 0.49, size = 419, normalized size = 3.10

$$\frac{90 \sqrt{d \cos(bx+a)} d^8 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 45 \sqrt{d \cos(bx+a)} d^{13/2} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) - 16 (d \cos(bx+a))^{5/2} d^4 - 160 \sqrt{d \cos(bx+a)} d^6}{40 b d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(11/2)*csc(b*x+a)^3,x, algorithm="fricas")`

[Out] $[-1/80*(90*(d^5*\cos(b*x+a)^2-d^5)*\sqrt{-d}*\arctan(2*\sqrt{d*\cos(b*x+a)}*\sqrt{-d}/(d*\cos(b*x+a)+d))-45*(d^5*\cos(b*x+a)^2-d^5)*\sqrt{-d}]*1$

```

og(-(d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1)
- 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*(4*d^5*c
os(b*x + a)^4 + 36*d^5*cos(b*x + a)^2 - 45*d^5)*sqrt(d*cos(b*x + a)))/(b*co
s(b*x + a)^2 - b), -1/80*(90*(d^5*cos(b*x + a)^2 - d^5)*sqrt(d)*arctan(2*sq
rt(d*cos(b*x + a))*sqrt(d)/(d*cos(b*x + a) - d)) - 45*(d^5*cos(b*x + a)^2 -
d^5)*sqrt(d)*log(-(d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(
b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)
) + 8*(4*d^5*cos(b*x + a)^4 + 36*d^5*cos(b*x + a)^2 - 45*d^5)*sqrt(d*cos(b*
x + a)))/(b*cos(b*x + a)^2 - b)]

```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(11/2)*csc(b*x+a)**3,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(11/2)*csc(b*x+a)^3,x, algorithm="giac")
```

```
[Out] integrate((d*cos(b*x + a))^(11/2)*csc(b*x + a)^3, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{11/2}}{\sin(a + b x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(11/2)/sin(a + b*x)^3,x)
```

```
[Out] int((d*cos(a + b*x))^(11/2)/sin(a + b*x)^3, x)
```

3.243 $\int (d \cos(a + bx))^{9/2} \csc^3(a + bx) dx$

Optimal. Leaf size=113

$$-\frac{7d^{9/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{7d^{9/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{7d^3 (d \cos(a + bx))^{3/2}}{6b} - \frac{d(d \cos(a + bx))^{7/2}}{2b}$$

[Out] $-7/4*d^{(9/2)}*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+7/4*d^{(9/2)}*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-7/6*d^3*(d*\cos(b*x+a))^{(3/2)}/b-1/2*d*(d*\cos(b*x+a))^{(7/2)}*\csc(b*x+a)^2/b$

Rubi [A]

time = 0.06, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2645, 294, 327, 335, 304, 209, 212}

$$-\frac{7d^{9/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{7d^{9/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{7d^3 (d \cos(a + bx))^{3/2}}{6b} - \frac{d \csc^2(a + bx) (d \cos(a + bx))^{7/2}}{2b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d*\operatorname{Cos}[a + b*x])^{(9/2)}*\operatorname{Csc}[a + b*x]^3, x]$

[Out] $(-7*d^{(9/2)}*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b) + (7*d^{(9/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b) - (7*d^3*(d*\operatorname{Cos}[a + b*x])^{(3/2)})/(6*b) - (d*(d*\operatorname{Cos}[a + b*x])^{(7/2)}*\operatorname{Csc}[a + b*x]^2)/(2*b)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[p, -1] \ \&\& \operatorname{GtQ}[m+1, n] \ \&\& \operatorname{!} \operatorname{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{9/2} \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{9/2}}{\left(1-\frac{x^2}{d^2}\right)^2} dx, x, d \cos(a + bx)\right)}{bd} \\
&= -\frac{d(d \cos(a + bx))^{7/2} \csc^2(a + bx)}{2b} + \frac{(7d)\text{Subst}\left(\int \frac{x^{5/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{7d^3(d \cos(a + bx))^{3/2}}{6b} - \frac{d(d \cos(a + bx))^{7/2} \csc^2(a + bx)}{2b} + \frac{(7d^3)\text{Subst}\left(\int \frac{x^{3/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{7d^3(d \cos(a + bx))^{3/2}}{6b} - \frac{d(d \cos(a + bx))^{7/2} \csc^2(a + bx)}{2b} + \frac{(7d^3)\text{Subst}\left(\int \frac{x^{1/2}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{7d^3(d \cos(a + bx))^{3/2}}{6b} - \frac{d(d \cos(a + bx))^{7/2} \csc^2(a + bx)}{2b} + \frac{(7d^5)\text{Subst}\left(\int \frac{1}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{7d^{9/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{7d^{9/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.46, size = 78, normalized size = 0.69

$$\frac{d^5 \left((-5 + 2 \cos(2(a + bx))) \cot^2(a + bx) + 21 \sqrt{-\cot^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \csc^2(a + bx)\right) \right)}{6b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(9/2)*Csc[a + b*x]^3,x]

[Out] (d^5*((-5 + 2*cos[2*(a + b*x)])*Cot[a + b*x]^2 + 21*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, Csc[a + b*x]^2]))/(6*b*Sqrt[d*cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 370 vs. 2(89) = 178.

time = 0.88, size = 371, normalized size = 3.28

method	result
--------	--------

default	$-\frac{4d^4 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \sqrt{2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d - d}}{3} - \frac{4d^4 \sqrt{2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d - d}}{3} + 2d^4 \sqrt{d \left(2 \left(\cos^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)\right)}$
---------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(9/2)*csc(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $(-4/3*d^4*\cos(1/2*b*x+1/2*a)^2*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}-4/3*d^4*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}+2*d^4*(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}-1/16*d^4/(\cos(1/2*b*x+1/2*a)+1)*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}+7/8*d^{(9/2)}*\ln((-4*d*\cos(1/2*b*x+1/2*a)+2*d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d)/(\cos(1/2*b*x+1/2*a)+1))+7/8*d^{(9/2)}*\ln((4*d*\cos(1/2*b*x+1/2*a)+2*d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d)/(\cos(1/2*b*x+1/2*a)-1))+1/8*d^4/\cos(1/2*b*x+1/2*a)^2*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}+7/4*d^5/(-d)^{(1/2)}*\ln((-2*d+2*(-d)^{(1/2)}*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)})/\cos(1/2*b*x+1/2*a))+1/16*d^4/(\cos(1/2*b*x+1/2*a)-1)*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)})/b$

Maxima [A]

time = 0.50, size = 118, normalized size = 1.04

$$\frac{12(d \cos(bx+a))^{\frac{3}{2}} d^6}{d^2 \cos(bx+a)^2 - d^2} - 42 d^{\frac{11}{2}} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 21 d^{\frac{11}{2}} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) - 16(d \cos(bx+a))^{\frac{3}{2}} d^4}{24bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a)^3,x, algorithm="maxima")`

[Out] $1/24*(12*(d*\cos(b*x + a))^{(3/2)}*d^6/(d^2*\cos(b*x + a)^2 - d^2) - 42*d^{(11/2)}*\arctan(\sqrt{d*\cos(b*x + a)}/\sqrt{d}) - 21*d^{(11/2)}*\log((\sqrt{d*\cos(b*x + a)} - \sqrt{d})/(\sqrt{d*\cos(b*x + a)} + \sqrt{d}))) - 16*(d*\cos(b*x + a))^{(3/2)}*d^4)/(b*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 199 vs. 2(89) = 178.

time = 0.47, size = 405, normalized size = 3.58

$$\frac{42(d^4 \cos(bx+a)^2 - d^4) \sqrt{-d} \arctan\left(\frac{2\sqrt{d \cos(bx+a)}}{\sqrt{-d}}\right) - 21(d^4 \cos(bx+a)^2 - d^4) \sqrt{-d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{-d}}{\sqrt{d \cos(bx+a)} + \sqrt{-d}}\right) - 16(d^4 \cos(bx+a)^2 - d^4) \sqrt{-d}}{48(b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a)^3,x, algorithm="fricas")`

[Out] $[-1/48*(42*(d^4*\cos(b*x + a)^2 - d^4)*\sqrt{-d}*\arctan(2*\sqrt{d*\cos(b*x + a)})*\sqrt{-d}/(d*\cos(b*x + a) + d) - 21*(d^4*\cos(b*x + a)^2 - d^4)*\sqrt{-d}*1$


```

og(-(d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1)
- 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*(4*d^4*c
os(b*x + a)^3 - 7*d^4*cos(b*x + a))*sqrt(d*cos(b*x + a)))/(b*cos(b*x + a)^2
- b), 1/48*(42*(d^4*cos(b*x + a)^2 - d^4)*sqrt(d)*arctan(2*sqrt(d*cos(b*x
+ a))*sqrt(d)/(d*cos(b*x + a) - d)) + 21*(d^4*cos(b*x + a)^2 - d^4)*sqrt(d)
*log(-(d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1)
+ 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) - 8*(4*d^4*
cos(b*x + a)^3 - 7*d^4*cos(b*x + a))*sqrt(d*cos(b*x + a)))/(b*cos(b*x + a)^
2 - b)]

```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(9/2)*csc(b*x+a)**3,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(9/2)*csc(b*x+a)^3,x, algorithm="giac")
```

```
[Out] integrate((d*cos(b*x + a))^(9/2)*csc(b*x + a)^3, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{9/2}}{\sin(a + b x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(9/2)/sin(a + b*x)^3,x)
```

```
[Out] int((d*cos(a + b*x))^(9/2)/sin(a + b*x)^3, x)
```

3.244 $\int (d \cos(a + bx))^{7/2} \csc^3(a + bx) dx$

Optimal. Leaf size=113

$$\frac{5d^{7/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{5d^{7/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{5d^3 \sqrt{d \cos(a + bx)}}{2b} - \frac{d(d \cos(a + bx))^{5/2}}{2b}$$

[Out] $5/4*d^{(7/2)*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+5/4*d^{(7/2)*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-1/2*d*(d*\cos(b*x+a))^{(5/2)*\csc(b*x+a)^2/b-5/2*d^{(3)*(d*\cos(b*x+a))^{(1/2)}/b}$

Rubi [A]

time = 0.06, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2645, 294, 327, 335, 218, 212, 209}

$$\frac{5d^{7/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{5d^{7/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{5d^3 \sqrt{d \cos(a + bx)}}{2b} - \frac{d \csc^2(a + bx) (d \cos(a + bx))^{5/2}}{2b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d \cos[a + b*x])^{(7/2)} * \csc[a + b*x]^3, x]$

[Out] $(5*d^{(7/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\cos[a + b*x]]/\operatorname{Sqrt}[d]]/(4*b) + (5*d^{(7/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\cos[a + b*x]]/\operatorname{Sqrt}[d]]/(4*b) - (5*d^3*\operatorname{Sqrt}[d*\cos[a + b*x]]/(2*b) - (d*(d*\cos[a + b*x])^{(5/2)*\csc[a + b*x]^2)/(2*b)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{PosQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ \operatorname{NegQ}[a/b] \ \&\& \ (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b, x\} \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 327

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{7/2} \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{7/2}}{(1-x^2)^2} dx, x, d \cos(a + bx)\right)}{bd} \\
&= -\frac{d(d \cos(a + bx))^{5/2} \csc^2(a + bx)}{2b} + \frac{(5d)\text{Subst}\left(\int \frac{x^{3/2}}{1-x^2} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{5d^3 \sqrt{d \cos(a + bx)}}{2b} - \frac{d(d \cos(a + bx))^{5/2} \csc^2(a + bx)}{2b} + \frac{(5d^3)\text{Subst}\left(\int \frac{x^{3/2}}{1-x^2} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{5d^3 \sqrt{d \cos(a + bx)}}{2b} - \frac{d(d \cos(a + bx))^{5/2} \csc^2(a + bx)}{2b} + \frac{(5d^3)\text{Subst}\left(\int \frac{x^{3/2}}{1-x^2} dx, x, d \cos(a + bx)\right)}{4b} \\
&= -\frac{5d^3 \sqrt{d \cos(a + bx)}}{2b} - \frac{d(d \cos(a + bx))^{5/2} \csc^2(a + bx)}{2b} + \frac{(5d^4)\text{Subst}\left(\int \frac{x^{3/2}}{1-x^2} dx, x, d \cos(a + bx)\right)}{4b} \\
&= \frac{5d^{7/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{5d^{7/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b}
\end{aligned}$$

Mathematica [A]

time = 0.84, size = 118, normalized size = 1.04

$$\frac{(d \cos(a + bx))^{7/2} \left(5 \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right) + 3 \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right) - 8 \sqrt{\cos(a + bx)} - 2 \sqrt{\cos(a + bx)} \csc^2(a + bx) - \log\left(1 - \sqrt{\cos(a + bx)}\right) + \log\left(1 + \sqrt{\cos(a + bx)}\right)\right)}{4b \cos^2(a + bx)}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^(7/2)*Csc[a + b*x]^3,x]`

```
[Out] ((d*Cos[a + b*x])^(7/2)*(5*ArcTan[Sqrt[Cos[a + b*x]]] + 3*ArcTanh[Sqrt[Cos[a + b*x]]] - 8*Sqrt[Cos[a + b*x]] - 2*Sqrt[Cos[a + b*x]]*Csc[a + b*x]^2 - Log[1 - Sqrt[Cos[a + b*x]]] + Log[1 + Sqrt[Cos[a + b*x]]]))/(4*b*Cos[a + b*x]^(7/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 309 vs. 2(89) = 178.

time = 0.87, size = 310, normalized size = 2.74

method	result
--------	--------

default	$-2d^3 \sqrt{d \left(2 \left(\cos^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) - 1 \right)} - \frac{d^3 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d}}{16 \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1 \right)} + \frac{5d^{\frac{7}{2}} \ln \left(\frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right)}}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right)} \right)}{8}$
---------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(7/2)*csc(b*x+a)^3,x,method=_RETURNVERBOSE)`

[Out] $(-2*d^3*(d*(2*\cos(1/2*b*x+1/2*a)^2-1))^{(1/2)}-1/16*d^3/(\cos(1/2*b*x+1/2*a)+1))*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}+5/8*d^{(7/2)}*\ln((-4*d*\cos(1/2*b*x+1/2*a)+2*d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d)/(\cos(1/2*b*x+1/2*a)+1))+5/8*d^{(7/2)}*\ln((4*d*\cos(1/2*b*x+1/2*a)+2*d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}-2*d)/(\cos(1/2*b*x+1/2*a)-1))-1/8*d^3/\cos(1/2*b*x+1/2*a)^2*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)}-5/4*d^4/(-d)^{(1/2)}*\ln((-2*d+2*(-d)^{(1/2)}*(2*\cos(1/2*b*x+1/2*a)^2*d-d)^{(1/2)})/\cos(1/2*b*x+1/2*a))+1/16*d^3/(\cos(1/2*b*x+1/2*a)-1)*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{(1/2)}/b$

Maxima [A]

time = 0.50, size = 118, normalized size = 1.04

$$\frac{4 \sqrt{d \cos(bx+a)} d^6}{d^2 \cos(bx+a)^2 - d^2} + 10 d^{\frac{9}{2}} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 5 d^{\frac{9}{2}} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) - 16 \sqrt{d \cos(bx+a)} d^4$$

8bd

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a)^3,x, algorithm="maxima")`

[Out] $1/8*(4*\sqrt{d*\cos(b*x+a)}*d^6/(d^2*\cos(b*x+a)^2-d^2)+10*d^{(9/2)}*\arctan(\sqrt{d*\cos(b*x+a)}/\sqrt{d})-5*d^{(9/2)}*\log((\sqrt{d*\cos(b*x+a)}-s\sqrt{d})/(\sqrt{d*\cos(b*x+a)}+s\sqrt{d}))-16*\sqrt{d*\cos(b*x+a)}*d^4)/(b*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 193 vs. 2(89) = 178.

time = 0.51, size = 393, normalized size = 3.48

$$\frac{10 \left(d^6 \cos(bx+a)^2 - d^6 \right) \sqrt{-d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 5 \left(d^6 \cos(bx+a)^2 - d^6 \right) \sqrt{-d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) + 8 \left(d^6 \cos(bx+a)^2 - 5d^6 \right) \sqrt{d \cos(bx+a)}}{16 \left(\cos(bx+a)^2 - 5 \right)} - \frac{10 \left(d^6 \cos(bx+a)^2 - d^6 \right) \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 5 \left(d^6 \cos(bx+a)^2 - d^6 \right) \sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) + 8 \left(d^6 \cos(bx+a)^2 - 5d^6 \right) \sqrt{d \cos(bx+a)}}{16 \left(\cos(bx+a)^2 - 5 \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a)^3,x, algorithm="fricas")`

[Out] $[-1/16*(10*(d^3*\cos(b*x+a)^2-d^3)*\sqrt{-d}*\arctan(2*\sqrt{d*\cos(b*x+a)})*\sqrt{-d}/(d*\cos(b*x+a)+d)-5*(d^3*\cos(b*x+a)^2-d^3)*\sqrt{-d}*\log(-(d*\cos(b*x+a)^2+4*\sqrt{d*\cos(b*x+a)}*\sqrt{-d}*(\cos(b*x+a)-1)-$

```

6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*(4*d^3*cos(b*x + a)^2 - 5*d^3)*sqrt(d*cos(b*x + a)))/(b*cos(b*x + a)^2 - b), -1/16*(10*(d^3*cos(b*x + a)^2 - d^3)*sqrt(d)*arctan(2*sqrt(d*cos(b*x + a))*sqrt(d)/(d*cos(b*x + a) - d)) - 5*(d^3*cos(b*x + a)^2 - d^3)*sqrt(d)*log(-(d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) + 8*(4*d^3*cos(b*x + a)^2 - 5*d^3)*sqrt(d*cos(b*x + a)))/(b*cos(b*x + a)^2 - b)]

```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(7/2)*csc(b*x+a)**3,x)
```

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(7/2)*csc(b*x+a)^3,x, algorithm="giac")
```

[Out] integrate((d*cos(b*x + a))^(7/2)*csc(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{7/2}}{\sin(a + b x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(7/2)/sin(a + b*x)^3,x)
```

[Out] int((d*cos(a + b*x))^(7/2)/sin(a + b*x)^3, x)

3.245 $\int (d \cos(a + bx))^{5/2} \csc^3(a + bx) dx$

Optimal. Leaf size=91

$$-\frac{3d^{5/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{3d^{5/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{d(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2b}$$

[Out] $-3/4*d^{(5/2)*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+3/4*d^{(5/2)*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-1/2*d*(d*\cos(b*x+a))^{(3/2)*\csc(b*x+a)^2/b}$

Rubi [A]

time = 0.05, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2645, 294, 335, 304, 209, 212}

$$-\frac{3d^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{3d^{5/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{d \csc^2(a + bx)(d \cos(a + bx))^{3/2}}{2b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d*\operatorname{Cos}[a + b*x])^{(5/2)*\operatorname{Csc}[a + b*x]^3, x]$

[Out] $(-3*d^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/(4*b) + (3*d^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]]/(4*b) - (d*(d*\operatorname{Cos}[a + b*x])^{(3/2)*\operatorname{Csc}[a + b*x]^2)/(2*b)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] :> \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] :> \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_)*(x_)^{(m_)*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] :> \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^n*((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[p, -1] \ \&\& \operatorname{GtQ}[m+1, n] \ \&\& \operatorname{!} \operatorname{LtQ}[(m+n*(p+1)+1)/n, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 304

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 335

```
Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)
))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*sin[(e_) + (f_)*(x_)]^(n_), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
 \int (d \cos(a + bx))^{5/2} \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^{5/2}}{(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a + bx)\right)}{bd} \\
 &= -\frac{d(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2b} + \frac{(3d)\text{Subst}\left(\int \frac{\sqrt{x}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4b} \\
 &= -\frac{d(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2b} + \frac{(3d)\text{Subst}\left(\int \frac{x^2}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{2b} \\
 &= -\frac{d(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2b} + \frac{(3d^3)\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{4b} \\
 &= -\frac{3d^{5/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{3d^{5/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.21, size = 65, normalized size = 0.71

$$\frac{d^3 \left(\cot^2(a + bx) - 3 \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \csc^2(a + bx)\right) \right)}{2b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(5/2)*Csc[a + b*x]^3,x]

[Out] -1/2*(d^3*(Cot[a + b*x]^2 - 3*(-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, Csc[a + b*x]^2]))/(b*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(71) = 142.

time = 0.69, size = 286, normalized size = 3.14

method	result
default	$-\frac{d^2 \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d}}{16 \left(\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1 \right)} + \frac{3d^{\frac{5}{2}} \ln \left(\frac{2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} - 4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right)}{8} + \frac{3d^{\frac{5}{2}} \ln \left(\frac{2\sqrt{d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} + 4d \cos \left(\frac{bx}{2} + \frac{a}{2} \right) - 2d}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right) + 1} \right)}{8}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(5/2)*csc(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] (-1/16*d^2/(cos(1/2*b*x+1/2*a)+1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+3/8*d^(5/2)*ln((-4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)+1))+3/8*d^(5/2)*ln((4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)-1))+1/8*d^2/cos(1/2*b*x+1/2*a)^2*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2)+3/4*d^3/(-d)^(1/2)*ln((-2*d+2*(-d)^(1/2)*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2))/cos(1/2*b*x+1/2*a))+1/16*d^2/(cos(1/2*b*x+1/2*a)-1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2))/b

Maxima [A]

time = 0.50, size = 103, normalized size = 1.13

$$\frac{4(d \cos(bx+a))^{\frac{3}{2}} d^4}{d^2 \cos(bx+a)^2 - d^2} - 6d^{\frac{7}{2}} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 3d^{\frac{7}{2}} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)$$

8bd

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^3,x, algorithm="maxima")

[Out] $\frac{1}{8} \cdot (4 \cdot (d \cdot \cos(b \cdot x + a))^{3/2} \cdot d^4 / (d^2 \cdot \cos(b \cdot x + a)^2 - d^2) - 6 \cdot d^{7/2} \cdot \arctan(\sqrt{d \cdot \cos(b \cdot x + a)} / \sqrt{d}) - 3 \cdot d^{7/2} \cdot \log((\sqrt{d \cdot \cos(b \cdot x + a)} - \sqrt{d}) / (\sqrt{d \cdot \cos(b \cdot x + a)} + \sqrt{d}))) / (b \cdot d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 184 vs. 2(71) = 142.

time = 0.40, size = 380, normalized size = 4.18

$$\frac{8 \sqrt{d \cos(bx+a)} d^4 \cos(bx+a) - 6 (d^2 \cos(bx+a) - d^2) \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) + 3 (d^2 \cos(bx+a) - d^2) \sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{8 \sqrt{d \cos(bx+a)} d^4 \cos(bx+a) - 6 (d^2 \cos(bx+a) - d^2) \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) + 3 (d^2 \cos(bx+a) - d^2) \sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{8 (b \cos(bx+a) - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^3,x, algorithm="fricas")

[Out] $\left[\frac{1}{16} \cdot (8 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot d^2 \cdot \cos(b \cdot x + a) - 6 \cdot (d^2 \cdot \cos(b \cdot x + a)^2 - d^2) \cdot \sqrt{-d} \cdot \arctan(1/2 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) + 1) / (d \cdot \cos(b \cdot x + a))) + 3 \cdot (d^2 \cdot \cos(b \cdot x + a)^2 - d^2) \cdot \sqrt{-d} \cdot \log((d \cdot \cos(b \cdot x + a)^2 - 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) - 1) - 6 \cdot d \cdot \cos(b \cdot x + a) + d) / (\cos(b \cdot x + a)^2 + 2 \cdot \cos(b \cdot x + a) + 1))) / (b \cdot \cos(b \cdot x + a)^2 - b), \frac{1}{16} \cdot (8 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot d^2 \cdot \cos(b \cdot x + a) - 6 \cdot (d^2 \cdot \cos(b \cdot x + a)^2 - d^2) \cdot \sqrt{d} \cdot \arctan(1/2 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot (\cos(b \cdot x + a) - 1) / (\sqrt{d} \cdot \cos(b \cdot x + a))) + 3 \cdot (d^2 \cdot \cos(b \cdot x + a)^2 - d^2) \cdot \sqrt{d} \cdot \log((d \cdot \cos(b \cdot x + a)^2 + 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \sqrt{d} \cdot (\cos(b \cdot x + a) + 1) + 6 \cdot d \cdot \cos(b \cdot x + a) + d) / (\cos(b \cdot x + a)^2 - 2 \cdot \cos(b \cdot x + a) + 1))) / (b \cdot \cos(b \cdot x + a)^2 - b) \right]$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^3,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*csc(b*x+a)^3,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(5/2)*csc(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{5/2}}{\sin(a + b x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(5/2)/sin(a + b*x)^3,x)
```

```
[Out] int((d*cos(a + b*x))^(5/2)/sin(a + b*x)^3, x)
```

3.246 $\int (d \cos(a + bx))^{3/2} \csc^3(a + bx) dx$

Optimal. Leaf size=91

$$\frac{d^{3/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{d^{3/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{d \sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2b}$$

[Out] $1/4*d^{(3/2)}*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b+1/4*d^{(3/2)}*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b-1/2*d*\csc(b*x+a)^2*(d*\cos(b*x+a))^{(1/2)}/b$

Rubi [A]

time = 0.05, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2645, 294, 335, 218, 212, 209}

$$\frac{d^{3/2} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{d^{3/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{d \csc^2(a + bx) \sqrt{d \cos(a + bx)}}{2b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d*\cos[a + b*x])^{(3/2)}*\csc[a + b*x]^3,x]$

[Out] $(d^{(3/2)}*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\cos[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b) + (d^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\cos[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b) - (d*\operatorname{Sqrt}[d*\cos[a + b*x]]*\csc[a + b*x]^2)/(2*b)$

Rule 209

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\int (d \cos(a + bx))^{3/2} \csc^3(a + bx) dx = -\frac{\text{Subst}\left(\int \frac{x^{3/2}}{(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a + bx)\right)}{bd}$$

$$= -\frac{d\sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2b} + \frac{d \text{Subst}\left(\int \frac{1}{\sqrt{x} (1-\frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{4b}$$

$$= -\frac{d\sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2b} + \frac{d \text{Subst}\left(\int \frac{1}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{2b}$$

$$= -\frac{d\sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2b} + \frac{d^2 \text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{4b}$$

$$= \frac{d^{3/2} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} + \frac{d^{3/2} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.14, size = 76, normalized size = 0.84

$$\frac{(d \cos(a + bx))^{3/2} (-\cot^2(a + bx))^{3/4} \left(3\sqrt[4]{-\cot^2(a + bx)} + {}_2F_1\left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \csc^2(a + bx)\right) \right) \sec^3(a + bx)}{6b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(3/2)*Csc[a + b*x]^3,x]

[Out] ((d*cos[a + b*x])^(3/2)*(-Cot[a + b*x]^2)^(3/4)*(3*(-Cot[a + b*x]^2)^(1/4) + Hypergeometric2F1[3/4, 3/4, 7/4, Csc[a + b*x]^2]))*Sec[a + b*x]^3)/(6*b)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 279 vs. 2(71) = 142.

time = 0.62, size = 280, normalized size = 3.08

method	result
default	$-\frac{d\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d+d}}{16\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)} + \frac{d^{\frac{3}{2}} \ln\left(\frac{2\sqrt{d}\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d+d} - 4d\cos\left(\frac{bx}{2} + \frac{a}{2}\right) - 2d}}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + 1}\right)}{8} + \frac{d^{\frac{3}{2}} \ln\left(\frac{2\sqrt{d}}{\dots}\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(3/2)*csc(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] (-1/16*d/(cos(1/2*b*x+1/2*a)+1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)+1/8*d^(3/2)*ln((-4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)+1))+1/8*d^(3/2)*ln((4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)-1))-1/8*d/cos(1/2*b*x+1/2*a)^2*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2)-1/4*d^2/(-d)^(1/2)*ln((-2*d+2*(-d)^(1/2)*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2))/cos(1/2*b*x+1/2*a))+1/16*d/(cos(1/2*b*x+1/2*a)-1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2))/b

Maxima [A]

time = 0.52, size = 103, normalized size = 1.13

$$\frac{4\sqrt{d\cos(bx+a)}d^4}{d^2\cos(bx+a)^2-d^2} + 2d^{\frac{5}{2}}\arctan\left(\frac{\sqrt{d\cos(bx+a)}}{\sqrt{d}}\right) - d^{\frac{5}{2}}\log\left(\frac{\sqrt{d\cos(bx+a)}-\sqrt{d}}{\sqrt{d\cos(bx+a)}+\sqrt{d}}\right)}{8bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^3,x, algorithm="maxima")

[Out] 1/8*(4*sqrt(d*cos(b*x + a))*d^4/(d^2*cos(b*x + a)^2 - d^2) + 2*d^(5/2)*arctan(sqrt(d*cos(b*x + a))/sqrt(d)) - d^(5/2)*log((sqrt(d*cos(b*x + a)) - sqrt(d))/(sqrt(d*cos(b*x + a)) + sqrt(d))))/(b*d)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 167 vs. 2(71) = 142.

time = 0.44, size = 347, normalized size = 3.81

$$\frac{2(d \cos(bx+a)^2 - d)\sqrt{-d} \arctan\left(\frac{\sqrt{d \cos(bx+a)} \sqrt{-d} \cos(bx+a)}{2d \cos(bx+a)}\right) - (d \cos(bx+a)^2 - d)\sqrt{-d} \log\left(\frac{d \cos(bx+a) \sqrt{d \cos(bx+a)} \sqrt{-d} \cos(bx+a) - 4d \cos(bx+a)^2}{\cos(bx+a) \sqrt{-d} \cos(bx+a)}\right) - 8 \sqrt{d \cos(bx+a)} d^2 (d \cos(bx+a)^2 - d) \sqrt{-d} \arctan\left(\frac{\sqrt{d \cos(bx+a)} \cos(bx+a)}{2 \sqrt{d \cos(bx+a)}}\right) + (d \cos(bx+a)^2 - d) \sqrt{-d} \log\left(\frac{d \cos(bx+a) \sqrt{d \cos(bx+a)} \sqrt{-d} \cos(bx+a) + 4d \cos(bx+a)^2}{\cos(bx+a) \sqrt{-d} \cos(bx+a)}\right) + 8 \sqrt{d \cos(bx+a)} d}{16(b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^3,x, algorithm="fricas")

[Out] [-1/16*(2*(d*cos(b*x + a)^2 - d)*sqrt(-d)*arctan(1/2*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) + 1)/(d*cos(b*x + a))) - (d*cos(b*x + a)^2 - d)*sqrt(-d)*log((d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1) - 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) - 8*sqrt(d*cos(b*x + a))*d/(b*cos(b*x + a)^2 - b), 1/16*(2*(d*cos(b*x + a)^2 - d)*sqrt(d)*arctan(1/2*sqrt(d*cos(b*x + a))*(cos(b*x + a) - 1)/(sqrt(d)*cos(b*x + a))) + (d*cos(b*x + a)^2 - d)*sqrt(d)*log((d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*d/(b*cos(b*x + a)^2 - b)]

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)**3,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 8009 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*csc(b*x+a)^3,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*csc(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{3/2}}{\sin(a + b x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(3/2)/sin(a + b*x)^3,x)

[Out] int((d*cos(a + b*x))^(3/2)/sin(a + b*x)^3, x)

3.247 $\int \sqrt{d \cos(a + bx)} \csc^3(a + bx) dx$

Optimal. Leaf size=93

$$\frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{\sqrt{d} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2bd}$$

[Out] $-1/2*(d*\cos(b*x+a))^{(3/2)}*\csc(b*x+a)^2/b/d+1/4*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})*d^{(1/2)}/b-1/4*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})*d^{(1/2)}/b$

Rubi [A]

time = 0.04, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2645, 296, 335, 304, 209, 212}

$$\frac{\sqrt{d} \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{\csc^2(a + bx)(d \cos(a + bx))^{3/2}}{2bd} - \frac{\sqrt{d} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^3,x]`

[Out] $(\operatorname{Sqrt}[d]*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b) - (\operatorname{Sqrt}[d]*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b) - ((d*\operatorname{Cos}[a + b*x])^{(3/2)}*\operatorname{Csc}[a + b*x]^2)/(2*b*d)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 296

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^n)^(p + 1)/(a*c*n*(p + 1))), x] + Dist[(m + n*(p + 1) + 1)/(a*n*(p + 1)), Int[(c*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)
))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{\sqrt{x}}{(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2bd} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{4bd} \\ &= -\frac{(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2bd} - \frac{\text{Subst}\left(\int \frac{x^2}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{2bd} \\ &= -\frac{(d \cos(a + bx))^{3/2} \csc^2(a + bx)}{2bd} - \frac{d \text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{4b} \\ &= \frac{\sqrt{d} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} - \frac{\sqrt{d} \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.19, size = 62, normalized size = 0.67

$$\frac{d \left(\cot^2(a + bx) + \sqrt[4]{-\cot^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \csc^2(a + bx)\right) \right)}{2b\sqrt{d\cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Csc[a + b*x]^3,x]

[Out] -1/2*(d*(Cot[a + b*x]^2 + (-Cot[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, Csc[a + b*x]^2]))/(b*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 274 vs. 2(73) = 146.

time = 0.72, size = 275, normalized size = 2.96

method	result
default	$-\frac{\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d+d}}{16\left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right)+1\right)} - \frac{\sqrt{d} \ln\left(\frac{2\sqrt{d} \sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d+d} - 4d\cos\left(\frac{bx}{2} + \frac{a}{2}\right) - 2d}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right)+1}\right)}{8} + \frac{\sqrt{2\left(\cos^2\right)}}{8}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*csc(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] (-1/16/(cos(1/2*b*x+1/2*a)+1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-1/8*d^(1/2)*ln((-4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)+1))+1/8/cos(1/2*b*x+1/2*a)^2*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2)-1/4*d/(-d)^(1/2)*ln((-2*d+2*(-d)^(1/2)*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2))/cos(1/2*b*x+1/2*a))+1/16/(cos(1/2*b*x+1/2*a)-1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-1/8*d^(1/2)*ln((4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)-1)))/b

Maxima [A]

time = 0.53, size = 102, normalized size = 1.10

$$\frac{\frac{4(d\cos(bx+a))^{\frac{3}{2}}d^2}{d^2\cos(bx+a)^2-d^2} + 2d^{\frac{3}{2}}\arctan\left(\frac{\sqrt{d\cos(bx+a)}}{\sqrt{d}}\right) + d^{\frac{3}{2}}\log\left(\frac{\sqrt{d\cos(bx+a)}-\sqrt{d}}{\sqrt{d\cos(bx+a)}+\sqrt{d}}\right)}{8bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^3,x, algorithm="maxima")

[Out] $\frac{1}{8} \cdot (4 \cdot (d \cdot \cos(b \cdot x + a))^{3/2} \cdot d^2 / (d^2 \cdot \cos(b \cdot x + a)^2 - d^2) + 2 \cdot d^{3/2} \cdot \arctan(\sqrt{d \cdot \cos(b \cdot x + a)} / \sqrt{d}) + d^{3/2} \cdot \log((\sqrt{d \cdot \cos(b \cdot x + a)} - \sqrt{d}) / (\sqrt{d \cdot \cos(b \cdot x + a)} + \sqrt{d}))) / (b \cdot d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 164 vs. 2(73) = 146.

time = 0.43, size = 340, normalized size = 3.66

$$\frac{2 \left[\cos(bx+a)^2 - 1 \right] \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)}}{d \cos(bx+a)}\right) + \left[\cos(bx+a)^2 - 1 \right] \sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)} - d \cos(bx+a)}{\sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)} + d \cos(bx+a)}\right) + 8 \sqrt{d \cos(bx+a)} \cos(bx+a) \cdot 2 \left[\cos(bx+a)^2 - 1 \right] \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)}}{2 \sqrt{d \cos(bx+a)}}\right) + \left[\cos(bx+a)^2 - 1 \right] \sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)} - d \cos(bx+a)}{\sqrt{d \cos(bx+a)} \sqrt{d \cos(bx+a)} + d \cos(bx+a)}\right) + 8 \sqrt{d \cos(bx+a)} \cos(bx+a)}{16 (b \cos(bx+a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^3,x, algorithm="fricas")`

[Out] $\left[\frac{1}{16} \cdot (2 \cdot (\cos(b \cdot x + a))^2 - 1) \cdot \sqrt{-d} \cdot \arctan\left(\frac{1}{2} \cdot \sqrt{d \cdot \cos(b \cdot x + a)}\right) \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) + 1) / (d \cdot \cos(b \cdot x + a)) + (\cos(b \cdot x + a))^2 - 1 \cdot \sqrt{-d} \cdot \log\left(\frac{(d \cdot \cos(b \cdot x + a))^2 + 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) - 1) - 6 \cdot d \cdot \cos(b \cdot x + a) + d}{(\cos(b \cdot x + a))^2 + 2 \cdot \cos(b \cdot x + a) + 1}\right) + 8 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \cos(b \cdot x + a) / (b \cdot \cos(b \cdot x + a)^2 - b), \frac{1}{16} \cdot (2 \cdot (\cos(b \cdot x + a))^2 - 1) \cdot \sqrt{d} \cdot \arctan\left(\frac{1}{2} \cdot \sqrt{d \cdot \cos(b \cdot x + a)}\right) \cdot (\cos(b \cdot x + a) - 1) / (\sqrt{d} \cdot \cos(b \cdot x + a)) + (\cos(b \cdot x + a))^2 - 1 \cdot \sqrt{d} \cdot \log\left(\frac{(d \cdot \cos(b \cdot x + a))^2 - 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \sqrt{d} \cdot (\cos(b \cdot x + a) + 1) + 6 \cdot d \cdot \cos(b \cdot x + a) + d}{(\cos(b \cdot x + a))^2 - 2 \cdot \cos(b \cdot x + a) + 1}\right) + 8 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot \cos(b \cdot x + a) / (b \cdot \cos(b \cdot x + a)^2 - b) \right]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(a + bx)} \csc^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**(1/2)*csc(b*x+a)**3,x)`

[Out] `Integral(sqrt(d*cos(a + b*x))*csc(a + b*x)**3, x)`

Giac [A]

time = 5.22, size = 95, normalized size = 1.02

$$\frac{d^3 \left(\frac{2 \sqrt{d \cos(bx+a)} \cos(bx+a)}{(d^2 \cos(bx+a)^2 - d^2) d} + \frac{\arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{-d}}\right)}{\sqrt{-d} d^2} + \frac{\arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{d^{5/2}} \right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*csc(b*x+a)^3,x, algorithm="giac")`

[Out] $\frac{1}{4}d^3(2\sqrt{d\cos(bx+a)}\cos(bx+a)/((d^2\cos(bx+a)^2 - d^2)d) + \arctan(\sqrt{d\cos(bx+a)}/\sqrt{-d})/(\sqrt{-d}d^2) + \arctan(\sqrt{d\cos(bx+a)}/\sqrt{d})/d^{5/2})/b$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{d \cos(a + bx)}}{\sin(a + bx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(a + b*x))^(1/2)/sin(a + b*x)^3,x)`

[Out] `int((d*cos(a + b*x))^(1/2)/sin(a + b*x)^3, x)`

$$3.248 \quad \int \frac{\csc^3(a+bx)}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=93

$$\frac{3 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4b\sqrt{d}} - \frac{3 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4b\sqrt{d}} - \frac{\sqrt{d \cos(a+bx)} \csc^2(a+bx)}{2bd}$$

[Out] $-3/4*\arctan((d*\cos(b*x+a))^(1/2)/d^(1/2))/b/d^(1/2)-3/4*\operatorname{arctanh}((d*\cos(b*x+a))^(1/2)/d^(1/2))/b/d^(1/2)-1/2*\csc(b*x+a)^2*(d*\cos(b*x+a))^(1/2)/b/d$

Rubi [A]

time = 0.05, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2645, 296, 335, 218, 212, 209}

$$\frac{3 \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4b\sqrt{d}} - \frac{\csc^2(a+bx) \sqrt{d \cos(a+bx)}}{2bd} - \frac{3 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4b\sqrt{d}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[a + b*x]^3/\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]], x]$

[Out] $(-3*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b*\operatorname{Sqrt}[d]) - (3*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b*\operatorname{Sqrt}[d]) - (\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]*\operatorname{Csc}[a + b*x]^2)/(2*b*d)$

Rule 209

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& !\operatorname{GtQ}[a/b, 0]$

Rule 296

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^n)^(p + 1)/(a*c*n*(p + 1))), x] + Dist[(m + n*(p + 1) + 1)/(a*n*(p + 1)), Int[(c*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^(p, x), x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned} \int \frac{\csc^3(a + bx)}{\sqrt{d \cos(a + bx)}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{\sqrt{x} (1 - \frac{x^2}{d^2})^2} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{\sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2bd} - \frac{3 \text{Subst}\left(\int \frac{1}{\sqrt{x} (1 - \frac{x^2}{d^2})} dx, x, d \cos(a + bx)\right)}{4bd} \\ &= -\frac{\sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2bd} - \frac{3 \text{Subst}\left(\int \frac{1}{1 - \frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a + bx)}\right)}{2bd} \\ &= -\frac{\sqrt{d \cos(a + bx)} \csc^2(a + bx)}{2bd} - \frac{3 \text{Subst}\left(\int \frac{1}{d - x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{4b} - \frac{3 \text{Subst}\left(\int \frac{1}{1 - x^2} dx, x, \sqrt{d \cos(a + bx)}\right)}{4b} \\ &= -\frac{3 \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b\sqrt{d}} - \frac{3 \tanh^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{d}}\right)}{4b\sqrt{d}} - \frac{\sqrt{d \cos(a + bx)}}{2bd} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.17, size = 69, normalized size = 0.74

$$\frac{d(-\cot^2(a+bx))^{3/4} \left(\sqrt[4]{-\cot^2(a+bx)} - {}_2F_1\left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \csc^2(a+bx)\right) \right)}{2b(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^3/Sqrt[d*Cos[a + b*x]], x]

[Out] (d*(-Cot[a + b*x]^2)^(3/4)*((-Cot[a + b*x]^2)^(1/4) - Hypergeometric2F1[3/4, 3/4, 7/4, Csc[a + b*x]^2]))/(2*b*(d*Cos[a + b*x])^(3/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 282 vs. 2(73) = 146.

time = 0.64, size = 283, normalized size = 3.04

method	result
default	$-\frac{\sqrt{-2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + d}}{16d \left(\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + 1\right)} \frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + d}^{-4d \cos\left(\frac{bx}{2} + \frac{a}{2}\right) - 2d}}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + 1} \frac{{}_2\sqrt{d} \sqrt{-2 \left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right) d + d}^{-4d \cos\left(\frac{bx}{2} + \frac{a}{2}\right) - 2d}}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right) + 1}}{8\sqrt{d}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(b*x+a)^3/(d*cos(b*x+a))^(1/2), x, method=_RETURNVERBOSE)

[Out] (-1/16/d/(cos(1/2*b*x+1/2*a)+1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-3/8/d^(1/2)*ln((-4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)+1))-3/8/d^(1/2)*ln((4*d*cos(1/2*b*x+1/2*a)+2*d^(1/2)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2)-2*d)/(cos(1/2*b*x+1/2*a)-1))-1/8/d/cos(1/2*b*x+1/2*a)^2*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2)+3/4/(-d)^(1/2)*ln((-2*d+2*(-d)^(1/2)*(2*cos(1/2*b*x+1/2*a)^2*d-d)^(1/2))/cos(1/2*b*x+1/2*a))+1/16/d/(cos(1/2*b*x+1/2*a)-1)*(-2*sin(1/2*b*x+1/2*a)^2*d+d)^(1/2))/b

Maxima [A]

time = 0.53, size = 103, normalized size = 1.11

$$\frac{4 \sqrt{d \cos(bx+a)} d^2}{d^2 \cos(bx+a)^2 - d^2} - 6 \sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) + 3 \sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)$$

8bd

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(1/2), x, algorithm="maxima")

[Out] 1/8*(4*sqrt(d*cos(b*x + a))*d^2/(d^2*cos(b*x + a)^2 - d^2) - 6*sqrt(d)*arctan(sqrt(d*cos(b*x + a))/sqrt(d)) + 3*sqrt(d)*log((sqrt(d*cos(b*x + a)) - sqrt(d))/(sqrt(d*cos(b*x + a)) + sqrt(d))))/(b*d)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 161 vs. 2(73) = 146.
time = 0.43, size = 334, normalized size = 3.59

$$\frac{6(\cos(bx+a)^2-1)\sqrt{-d}\arctan\left(\frac{\sqrt{d\cos(bx+a)}\sqrt{-d}\cos(bx+a)}{2d\cos(bx+a)}\right)-3(\cos(bx+a)^2-1)\sqrt{-d}\log\left(\frac{d\cos(bx+a)+\sqrt{d\cos(bx+a)}\sqrt{-d}\cos(bx+a)-d\cos(bx+a)}{d\cos(bx+a)+\sqrt{d\cos(bx+a)}\sqrt{-d}\cos(bx+a)}\right)+8\sqrt{d\cos(bx+a)}\dots}{16(bd\cos(bx+a)^2-bd)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] [1/16*(6*(cos(b*x + a)^2 - 1)*sqrt(-d)*arctan(1/2*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) + 1)/(d*cos(b*x + a))) - 3*(cos(b*x + a)^2 - 1)*sqrt(-d)*log((d*cos(b*x + a)^2 + 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1) - 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a)))/(b*d*cos(b*x + a)^2 - b*d), -1/16*(6*(cos(b*x + a)^2 - 1)*sqrt(d)*arctan(1/2*sqrt(d*cos(b*x + a))*(cos(b*x + a) - 1)/(sqrt(d)*cos(b*x + a))) - 3*(cos(b*x + a)^2 - 1)*sqrt(d)*log((d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) - 8*sqrt(d*cos(b*x + a)))/(b*d*cos(b*x + a)^2 - b*d)]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(a + bx)}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)**3/(d*cos(b*x+a))**(1/2),x)

[Out] Integral(csc(a + b*x)**3/sqrt(d*cos(a + b*x)), x)

Giac [A]

time = 5.10, size = 91, normalized size = 0.98

$$\frac{d^3 \left(\frac{2\sqrt{d\cos(bx+a)}}{(d^2\cos(bx+a)^2-d^2)d^2} + \frac{3\arctan\left(\frac{\sqrt{d\cos(bx+a)}}{\sqrt{-d}}\right)}{\sqrt{-d}d^3} - \frac{3\arctan\left(\frac{\sqrt{d\cos(bx+a)}}{\sqrt{d}}\right)}{d^{\frac{7}{2}}} \right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] 1/4*d^3*(2*sqrt(d*cos(b*x + a))/((d^2*cos(b*x + a)^2 - d^2)*d^2) + 3*arctan(sqrt(d*cos(b*x + a))/sqrt(-d))/(sqrt(-d)*d^3) - 3*arctan(sqrt(d*cos(b*x + a))/sqrt(d))/d^(7/2))/b

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^3 \sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(1/2)), x)

[Out] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(1/2)), x)

$$3.249 \quad \int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=115

$$\frac{5 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{3/2}} - \frac{5 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{3/2}} + \frac{5}{2bd\sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd\sqrt{d \cos(a+bx)}}$$

[Out] 5/4*arctan((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(3/2)-5/4*arctanh((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(3/2)+5/2/b/d/(d*cos(b*x+a))^(1/2)-1/2*csc(b*x+a)^2/b/d/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2645, 296, 331, 335, 304, 209, 212}

$$\frac{5 \text{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{3/2}} - \frac{5 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{3/2}} + \frac{5}{2bd\sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^3/(d*Cos[a + b*x])^(3/2), x]

[Out] (5*ArcTan[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(4*b*d^(3/2)) - (5*ArcTanh[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(4*b*d^(3/2)) + 5/(2*b*d*Sqrt[d*Cos[a + b*x]]) - Csc[a + b*x]^2/(2*b*d*Sqrt[d*Cos[a + b*x]]))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 296

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m+1))*((a+b*x^n)^(p+1)/(a*c*n*(p+1))), x] + Dist[(m+n*(p+1)+1)/(a*n*(p+1)), Int[(c*x)^m*(a+b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p,

x]

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x
)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1)
+ 1)/(a*c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a,
  b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p,
  x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p), x], (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_
Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x
, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] &&
!(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{3/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{3/2}(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a+bx)\right)}{bd} \\
&= -\frac{\csc^2(a+bx)}{2bd\sqrt{d \cos(a+bx)}} - \frac{5\text{Subst}\left(\int \frac{1}{x^{3/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd} \\
&= \frac{5}{2bd\sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd\sqrt{d \cos(a+bx)}} - \frac{5\text{Subst}\left(\int \frac{\sqrt{x}}{1-\frac{x^2}{d^2}} dx, x, d \cos(a+bx)\right)}{4bd^3} \\
&= \frac{5}{2bd\sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd\sqrt{d \cos(a+bx)}} - \frac{5\text{Subst}\left(\int \frac{x^2}{1-\frac{x^2}{d^2}} dx, x, \sqrt{d \cos(a+bx)}\right)}{2bd^3} \\
&= \frac{5}{2bd\sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd\sqrt{d \cos(a+bx)}} - \frac{5\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a+bx)}\right)}{4bd} \\
&= \frac{5 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{3/2}} - \frac{5 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{3/2}} + \frac{5}{2bd\sqrt{d \cos(a+bx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.19, size = 91, normalized size = 0.79

$$\frac{-(-\cot^2(a+bx))^{3/4}(-4+\cot^2(a+bx))+5\cot^2(a+bx) {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \csc^2(a+bx)\right)}{2bd\sqrt{d \cos(a+bx)}(-\cot^2(a+bx))^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^3/(d*Cos[a + b*x])^(3/2), x]

[Out] (-((-Cot[a + b*x]^2)^(3/4)*(-4 + Cot[a + b*x]^2)) + 5*Cot[a + b*x]^2*Hypergeometric2F1[1/4, 1/4, 5/4, Csc[a + b*x]^2])/(2*b*d*Sqrt[d*Cos[a + b*x]]*(-Cot[a + b*x]^2)^(3/4))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 704 vs. 2(91) = 182.

time = 1.04, size = 705, normalized size = 6.13

method	result
--------	--------

default	$-\sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + d} d^{\frac{3}{2}} \sqrt{-d} - \left(\sin^6\left(\frac{bx}{2} + \frac{a}{2}\right)\right) \left(20d^{\frac{5}{2}} \ln\left(\frac{2\sqrt{-d} \sqrt{-2\left(\sin^2\left(\frac{bx}{2} + \frac{a}{2}\right)\right)d + d}}{\cos\left(\frac{bx}{2} + \frac{a}{2}\right)}\right)\right)$
---------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)^3/(d*cos(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{8}d^{7/2}/(-d)^{1/2}/\sin(1/2*b*x+1/2*a)^2/(2*\sin(1/2*b*x+1/2*a)^4-3*\sin(1/2*b*x+1/2*a)^2+1)*(-(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}*d^{3/2}*(-d)^{1/2}-\sin(1/2*b*x+1/2*a)^6*(20*d^{5/2}*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d))+10*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^2+10*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^2+5*(6*d^{5/2}*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d)-4*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}*d^{3/2}*(-d)^{1/2}+3*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^2+3*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^2)*\sin(1/2*b*x+1/2*a)^4-5*(2*d^{5/2}*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d)-4*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}*d^{3/2}*(-d)^{1/2}+(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^2+(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^2)*\sin(1/2*b*x+1/2*a)^2)/b$

Maxima [A]

time = 0.52, size = 117, normalized size = 1.02

$$\frac{10 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{\sqrt{d}} + \frac{5 \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{\sqrt{d}} + \frac{4(5d^2 \cos(bx+a)^2 - 4d^2)}{(d \cos(bx+a))^{\frac{5}{2}} - \sqrt{d \cos(bx+a)} d^2}$$

$8bd$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")`

[Out] $\frac{1}{8}*(10*\arctan(\sqrt{d*\cos(b*x+a)}/\sqrt{d})/\sqrt{d} + 5*\log((\sqrt{d*\cos(b*x+a)} - \sqrt{d})/(\sqrt{d*\cos(b*x+a)} + \sqrt{d}))/\sqrt{d} + 4*(5*d^2*\cos(b*x+a)^2 - 4*d^2)/((d*\cos(b*x+a))^{5/2} - \sqrt{d*\cos(b*x+a)}*d^2))/(b*d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 197 vs. 2(91) = 182.

time = 0.46, size = 406, normalized size = 3.53

$$\frac{10(\cos(bx+a)^2 - \cos(bx+a))\sqrt{d}\arctan\left(\frac{\sqrt{d}\cos(bx+a)\sqrt{d\cos(bx+a)}}{d\cos(bx+a)}\right) - 5(\cos(bx+a)^2 - \cos(bx+a))\sqrt{d}\log\left(\frac{d\cos(bx+a)\sqrt{d\cos(bx+a)}}{d\cos(bx+a)}\right) + 8\sqrt{d}\cos(bx+a)(\cos(bx+a)^2 - 4) - 10(\cos(bx+a)^2 - \cos(bx+a))\sqrt{d}\arctan\left(\frac{\sqrt{d}\cos(bx+a)\sqrt{d\cos(bx+a)}}{d\cos(bx+a)}\right) + 5(\cos(bx+a)^2 - \cos(bx+a))\sqrt{d}\log\left(\frac{d\cos(bx+a)\sqrt{d\cos(bx+a)}}{d\cos(bx+a)}\right) + 8\sqrt{d}\cos(bx+a)(\cos(bx+a)^2 - 4)}{16(d^2\cos(bx+a)^2 - d^2\cos(bx+a))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] [1/16*(10*(cos(b*x + a)^3 - cos(b*x + a))*sqrt(-d)*arctan(1/2*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) + 1)/(d*cos(b*x + a))) - 5*(cos(b*x + a)^3 - cos(b*x + a))*sqrt(-d)*log((d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(-d)*(cos(b*x + a) - 1) - 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 + 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*(5*cos(b*x + a)^2 - 4))/(b*d^2*cos(b*x + a)^3 - b*d^2*cos(b*x + a)), 1/16*(10*(cos(b*x + a)^3 - cos(b*x + a))*sqrt(d)*arctan(1/2*sqrt(d*cos(b*x + a))*(cos(b*x + a) - 1)/(sqrt(d)*cos(b*x + a))) + 5*(cos(b*x + a)^3 - cos(b*x + a))*sqrt(d)*log((d*cos(b*x + a)^2 - 4*sqrt(d*cos(b*x + a))*sqrt(d)*(cos(b*x + a) + 1) + 6*d*cos(b*x + a) + d)/(cos(b*x + a)^2 - 2*cos(b*x + a) + 1)) + 8*sqrt(d*cos(b*x + a))*(5*cos(b*x + a)^2 - 4))/(b*d^2*cos(b*x + a)^3 - b*d^2*cos(b*x + a))]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(a + bx)}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)**3/(d*cos(b*x+a))**(3/2),x)

[Out] Integral(csc(a + b*x)**3/(d*cos(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)^3/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^3 (d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(3/2)),x)
```

```
[Out] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(3/2)), x)
```

$$3.250 \quad \int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=115

$$-\frac{7 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{5/2}} - \frac{7 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{5/2}} + \frac{7}{6bd(d \cos(a+bx))^{3/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{3/2}}$$

[Out] $-7/4*\arctan((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b/d^{(5/2)}-7/4*\operatorname{arctanh}((d*\cos(b*x+a))^{(1/2)}/d^{(1/2)})/b/d^{(5/2)}+7/6/b/d/(d*\cos(b*x+a))^{(3/2)}-1/2*\csc(b*x+a)^2/b/d/(d*\cos(b*x+a))^{(3/2)}$

Rubi [A]

time = 0.05, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2645, 296, 331, 335, 218, 212, 209}

$$-\frac{7 \operatorname{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{5/2}} - \frac{7 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{5/2}} + \frac{7}{6bd(d \cos(a+bx))^{3/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[a + b*x]^3/(d*Cos[a + b*x])^(5/2), x]`

[Out] $(-7*\operatorname{ArcTan}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b*d^{(5/2)}) - (7*\operatorname{ArcTanh}[\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]]/\operatorname{Sqrt}[d]])/(4*b*d^{(5/2)}) + 7/(6*b*d*(d*\operatorname{Cos}[a + b*x])^{(3/2)}) - \operatorname{Csc}[a + b*x]^2/(2*b*d*(d*\operatorname{Cos}[a + b*x])^{(3/2)})$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 296

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^n)^(p + 1)/(a*c*n*(p + 1))), x] + Dist[(m + n*(p + 1) + 1)/(a*n*(p + 1)), Int[(c*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 331

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2645

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{5/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{5/2}(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a+bx)\right)}{bd} \\
&= -\frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{3/2}} - \frac{7\text{Subst}\left(\int \frac{1}{x^{5/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd} \\
&= \frac{7}{6bd(d \cos(a+bx))^{3/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{3/2}} - \frac{7\text{Subst}\left(\int \frac{1}{\sqrt{x}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd^3} \\
&= \frac{7}{6bd(d \cos(a+bx))^{3/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{3/2}} - \frac{7\text{Subst}\left(\int \frac{1}{1-\frac{x^4}{d^2}} dx, x, \sqrt{d \cos(a+bx)}\right)}{2bd^3} \\
&= \frac{7}{6bd(d \cos(a+bx))^{3/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{3/2}} - \frac{7\text{Subst}\left(\int \frac{1}{d-x^2} dx, x, \sqrt{d \cos(a+bx)}\right)}{4bd^2} \\
&= -\frac{7 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{5/2}} - \frac{7 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{5/2}} + \frac{7}{6bd(d \cos(a+bx))^{3/2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.27, size = 92, normalized size = 0.80

$$\frac{\sqrt[4]{-\cot^2(a+bx)} (4 - 3 \cot^2(a+bx)) + 7 \cot^2(a+bx) {}_2F_1\left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \csc^2(a+bx)\right)}{6bd(d \cos(a+bx))^{3/2} \sqrt[4]{-\cot^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^3/(d*Cos[a + b*x])^(5/2), x]

[Out] ((-Cot[a + b*x]^2)^(1/4)*(4 - 3*Cot[a + b*x]^2) + 7*Cot[a + b*x]^2*Hypergeometric2F1[3/4, 3/4, 7/4, Csc[a + b*x]^2])/(6*b*d*(d*Cos[a + b*x])^(3/2)*(-Cot[a + b*x]^2)^(1/4))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 904 vs. 2(91) = 182.

time = 1.19, size = 905, normalized size = 7.87

method	result
--------	--------

default	$3\sqrt{-d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} d^{\frac{5}{2}} - 84 \left(-2 \ln \left(\frac{2\sqrt{-d} \sqrt{-2 \left(\sin^2 \left(\frac{bx}{2} + \frac{a}{2} \right) \right) d + d} - 2d}{\cos \left(\frac{bx}{2} + \frac{a}{2} \right)} \right) \right) d^{\frac{7}{2}} + \sqrt{-d}$
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)^3/(d*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{24}d^{11/2}/(-d)^{1/2}/\sin(1/2*b*x+1/2*a)^2/(4*\sin(1/2*b*x+1/2*a)^6-8*\sin(1/2*b*x+1/2*a)^4+5*\sin(1/2*b*x+1/2*a)^2-1)*(3*(-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}*d^{5/2}-84*(-2*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d))*d^{7/2}+(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^3+(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^3)*\sin(1/2*b*x+1/2*a)^8+168*(-2*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d))*d^{7/2}+(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^3+(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^3)*\sin(1/2*b*x+1/2*a)^6+7*(-6*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d))*d^{7/2}-4*(-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}*d^{5/2}+3*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^3+3*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^3)*\sin(1/2*b*x+1/2*a)^2-7*(-30*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-d))*d^{7/2}-4*(-d)^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}*d^{5/2}+15*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)+1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^3+15*(-d)^{1/2}*\ln(2/(\cos(1/2*b*x+1/2*a)-1))*(d^{1/2}*(-2*\sin(1/2*b*x+1/2*a)^2*d+d)^{1/2}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^3)*\sin(1/2*b*x+1/2*a)^4)/b$

Maxima [A]

time = 0.50, size = 117, normalized size = 1.02

$$\frac{\frac{4(7d^2 \cos^2(bx+a) - 4d^2)}{(d \cos(bx+a))^{\frac{7}{2}} - (d \cos(bx+a))^{\frac{3}{2}} d^2} - \frac{42 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{d^{\frac{3}{2}}} + \frac{21 \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{d^{\frac{3}{2}}}}{24bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")`

[Out] $\frac{1}{24} \cdot (4 \cdot (7 \cdot d^2 \cdot \cos(b \cdot x + a))^2 - 4 \cdot d^2) / ((d \cdot \cos(b \cdot x + a))^{7/2}) - (d \cdot \cos(b \cdot x + a))^{3/2} \cdot d^2 - 42 \cdot \arctan(\sqrt{d \cdot \cos(b \cdot x + a)} / \sqrt{d}) / d^{3/2} + 21 \cdot \log((\sqrt{d \cdot \cos(b \cdot x + a)} - \sqrt{d}) / (\sqrt{d \cdot \cos(b \cdot x + a)} + \sqrt{d})) / d^{3/2} / (b \cdot d)$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 203 vs. 2(91) = 182.
time = 0.43, size = 418, normalized size = 3.63

$$\frac{42(\cos(bx+a)^2 - \cos(bx+a)^2)\sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 21(\cos(bx+a)^2 - \cos(bx+a)^2)\sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) + 8\sqrt{d \cos(bx+a)}(7 \cos(bx+a)^2 - 4)}{48(b^2 \cos(bx+a)^2 - b^2 \cos(bx+a)^2)} \cdot \frac{42(\cos(bx+a)^2 - \cos(bx+a)^2)\sqrt{d} \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right) - 21(\cos(bx+a)^2 - \cos(bx+a)^2)\sqrt{d} \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right) - 8\sqrt{d \cos(bx+a)}(7 \cos(bx+a)^2 - 4)}{48(b^2 \cos(bx+a)^2 - b^2 \cos(bx+a)^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")`

[Out] $\frac{1}{148} \cdot (42 \cdot (\cos(b \cdot x + a))^4 - \cos(b \cdot x + a)^2) \cdot \sqrt{-d} \cdot \arctan(1/2 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) + 1) / (d \cdot \cos(b \cdot x + a)) - 21 \cdot (\cos(b \cdot x + a))^4 - \cos(b \cdot x + a)^2) \cdot \sqrt{-d} \cdot \log((d \cdot \cos(b \cdot x + a))^2 + 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) - 1) - 6 \cdot d \cdot \cos(b \cdot x + a) + d) / (\cos(b \cdot x + a)^2 + 2 \cdot \cos(b \cdot x + a) + 1) + 8 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot (7 \cdot \cos(b \cdot x + a)^2 - 4) / (b \cdot d^3 \cdot \cos(b \cdot x + a)^4 - b \cdot d^3 \cdot \cos(b \cdot x + a)^2), -1/48 \cdot (42 \cdot (\cos(b \cdot x + a))^4 - \cos(b \cdot x + a)^2) \cdot \sqrt{d} \cdot \arctan(1/2 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot (\cos(b \cdot x + a) - 1) / (\sqrt{d} \cdot \cos(b \cdot x + a)) - 21 \cdot (\cos(b \cdot x + a))^4 - \cos(b \cdot x + a)^2) \cdot \sqrt{d} \cdot \log((d \cdot \cos(b \cdot x + a))^2 - 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot \sqrt{d} \cdot (\cos(b \cdot x + a) + 1) + 6 \cdot d \cdot \cos(b \cdot x + a) + d) / (\cos(b \cdot x + a)^2 - 2 \cdot \cos(b \cdot x + a) + 1) - 8 \cdot \sqrt{d \cdot \cos(b \cdot x + a)} \cdot (7 \cdot \cos(b \cdot x + a)^2 - 4) / (b \cdot d^3 \cdot \cos(b \cdot x + a)^4 - b \cdot d^3 \cdot \cos(b \cdot x + a)^2)]$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)**3/(d*cos(b*x+a))**(5/2),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(5/2),x, algorithm="giac")`

[Out] `integrate(csc(b*x + a)^3/(d*cos(b*x + a))^(5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^3 (d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(5/2)), x)

[Out] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(5/2)), x)

$$3.251 \quad \int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=137

$$\frac{9 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{7/2}} - \frac{9 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{7/2}} + \frac{9}{10bd(d \cos(a+bx))^{5/2}} + \frac{9}{2bd^3 \sqrt{d \cos(a+bx)}}$$

[Out] 9/4*arctan((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(7/2)-9/4*arctanh((d*cos(b*x+a))^(1/2)/d^(1/2))/b/d^(7/2)+9/10/b/d/(d*cos(b*x+a))^(5/2)-1/2*csc(b*x+a)^2/b/d/(d*cos(b*x+a))^(5/2)+9/2/b/d^3/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2645, 296, 331, 335, 304, 209, 212}

$$\frac{9 \text{ArcTan}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{7/2}} - \frac{9 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{7/2}} + \frac{9}{2bd^3 \sqrt{d \cos(a+bx)}} + \frac{9}{10bd(d \cos(a+bx))^{5/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[a + b*x]^3/(d*Cos[a + b*x])^(7/2), x]

[Out] (9*ArcTan[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(4*b*d^(7/2))) - (9*ArcTanh[Sqrt[d*Cos[a + b*x]]/Sqrt[d]]/(4*b*d^(7/2))) + 9/(10*b*d*(d*Cos[a + b*x])^(5/2)) + 9/(2*b*d^3*Sqrt[d*Cos[a + b*x]]) - Csc[a + b*x]^2/(2*b*d*(d*Cos[a + b*x])^(5/2))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 296

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m+1))*((a + b*x^n)^(p+1)/(a*c*n*(p+1))), x] + Dist[(m+n*(p+1)+1)/(a*n*(p+1)), Int[(c*x)^m*(a + b*x^n)^(p+1), x], x] /; FreeQ[{a,

b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 331

Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 335

Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2645

Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^(m_.)*sin[(e_.) + (f_.)*(x_)]^(n_.), x_Symbol] := Dist[-(a*f)^(-1), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Cos[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(a+bx)}{(d \cos(a+bx))^{7/2}} dx &= -\frac{\text{Subst}\left(\int \frac{1}{x^{7/2}(1-\frac{x^2}{d^2})^2} dx, x, d \cos(a+bx)\right)}{bd} \\
&= -\frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{5/2}} - \frac{9\text{Subst}\left(\int \frac{1}{x^{7/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd} \\
&= \frac{9}{10bd(d \cos(a+bx))^{5/2}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{5/2}} - \frac{9\text{Subst}\left(\int \frac{1}{x^{3/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd^3} \\
&= \frac{9}{10bd(d \cos(a+bx))^{5/2}} + \frac{9}{2bd^3 \sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{5/2}} - \frac{9\text{Subst}\left(\int \frac{1}{x^{1/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd^3} \\
&= \frac{9}{10bd(d \cos(a+bx))^{5/2}} + \frac{9}{2bd^3 \sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{5/2}} - \frac{9\text{Subst}\left(\int \frac{1}{x^{1/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd^3} \\
&= \frac{9}{10bd(d \cos(a+bx))^{5/2}} + \frac{9}{2bd^3 \sqrt{d \cos(a+bx)}} - \frac{\csc^2(a+bx)}{2bd(d \cos(a+bx))^{5/2}} - \frac{9\text{Subst}\left(\int \frac{1}{x^{1/2}(1-\frac{x^2}{d^2})} dx, x, d \cos(a+bx)\right)}{4bd^3} \\
&= \frac{9 \tan^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{7/2}} - \frac{9 \tanh^{-1}\left(\frac{\sqrt{d \cos(a+bx)}}{\sqrt{d}}\right)}{4bd^{7/2}} + \frac{9}{10bd(d \cos(a+bx))^{5/2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.34, size = 102, normalized size = 0.74

$$\frac{45 \cot^2(a+bx) {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \csc^2(a+bx)\right) + (-\cot^2(a+bx))^{3/4} (40 - 5 \cot^2(a+bx) + 4 \sec^2(a+bx))}{10bd^3 \sqrt{d \cos(a+bx)} (-\cot^2(a+bx))^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[a + b*x]^3/(d*Cos[a + b*x])^(7/2), x]

[Out] (45*Cot[a + b*x]^2*Hypergeometric2F1[1/4, 1/4, 5/4, Csc[a + b*x]^2] + (-Cot[a + b*x]^2)^(3/4)*(40 - 5*Cot[a + b*x]^2 + 4*Sec[a + b*x]^2))/(10*b*d^3*Sqrt[d*Cos[a + b*x]]*(-Cot[a + b*x]^2)^(3/4))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1164 vs. 2(109) = 218.

time = 1.34, size = 1165, normalized size = 8.50

method	result	size
default	Expression too large to display	1165

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(b*x+a)^3/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{40}(-d)^{(1/2)}/d^{(15/2)}/\sin(1/2*b*x+1/2*a)^2/(8*\sin(1/2*b*x+1/2*a)^8-20*\sin(1/2*b*x+1/2*a)^6+18*\sin(1/2*b*x+1/2*a)^4-7*\sin(1/2*b*x+1/2*a)^2+1)*(-5*(-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}*d^{(7/2)}-360*(2*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-d)*d^{(9/2)}+(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^4+(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^4)*\sin(1/2*b*x+1/2*a)^{10}-180*(-10*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-d)*d^{(9/2)}+4*(-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}*d^{(7/2)}-5*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^4-5*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^4)*\sin(1/2*b*x+1/2*a)^8+90*(-18*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-d)*d^{(9/2)}+16*(-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}*d^{(7/2)}-9*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^4-9*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^4)*\sin(1/2*b*x+1/2*a)^6-9*(-70*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-d)*d^{(9/2)}+104*(-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}*d^{(7/2)}-35*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^4-35*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^4)*\sin(1/2*b*x+1/2*a)^4+9*(-10*\ln(2/\cos(1/2*b*x+1/2*a))*((-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-d)*d^{(9/2)}+24*(-d)^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}*d^{(7/2)}-5*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)-1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}+2*d*\cos(1/2*b*x+1/2*a)-d))*d^4-5*(-d)^{(1/2)}*\ln(2/(\cos(1/2*b*x+1/2*a)+1)*(d^{(1/2)}*(-2*\sin(1/2*b*x+1/2*a)^{2*d+d})^{(1/2)}-2*d*\cos(1/2*b*x+1/2*a)-d))*d^4)*\sin(1/2*b*x+1/2*a)^2)/b$

Maxima [A]

time = 0.49, size = 134, normalized size = 0.98

$$\frac{4 \left(45 d^4 \cos^4(bx+a) - 36 d^4 \cos^2(bx+a) - 4 d^4 \right)}{(d \cos(bx+a))^{\frac{9}{2}} d^2 - (d \cos(bx+a))^{\frac{5}{2}} d^4} + \frac{90 \arctan\left(\frac{\sqrt{d \cos(bx+a)}}{\sqrt{d}}\right)}{d^{\frac{5}{2}}} + \frac{45 \log\left(\frac{\sqrt{d \cos(bx+a)} - \sqrt{d}}{\sqrt{d \cos(bx+a)} + \sqrt{d}}\right)}{d^{\frac{5}{2}}}$$

$40 b d$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")

[Out] $\frac{1}{40} \cdot (4 \cdot (45 \cdot d^4 \cdot \cos(b \cdot x + a)^4 - 36 \cdot d^4 \cdot \cos(b \cdot x + a)^2 - 4 \cdot d^4)) / ((d \cdot \cos(b \cdot x + a))^{9/2} \cdot d^2 - (d \cdot \cos(b \cdot x + a))^{5/2} \cdot d^4) + 90 \cdot \arctan(\sqrt{d \cdot \cos(b \cdot x + a)} / \sqrt{d}) / d^{5/2} + 45 \cdot \log((\sqrt{d \cdot \cos(b \cdot x + a)} - \sqrt{d}) / (\sqrt{d \cdot \cos(b \cdot x + a)} + \sqrt{d})) / d^{5/2} / (b \cdot d)$

Fricas [A]

time = 0.45, size = 438, normalized size = 3.20

$\frac{90(\cos(bx+a)^5 - \cos(bx+a)^3)\sqrt{-d}\arctan\left(\frac{1}{2}\sqrt{d\cos(bx+a)}\right) - 45(\cos(bx+a)^5 - \cos(bx+a)^3)\sqrt{-d}\log\left(\frac{\cos(bx+a) + 1}{\cos(bx+a) - 1}\right) + 8(45\cos(bx+a)^4 - 36\cos(bx+a)^2 - 4)\sqrt{d\cos(bx+a)}}{40(d\cos(bx+a))^{9/2}d^2 - (d\cos(bx+a))^{5/2}d^4} + \frac{45(\cos(bx+a)^5 - \cos(bx+a)^3)\sqrt{d}\arctan\left(\frac{1}{2}\sqrt{d\cos(bx+a)}\right) + 45(\cos(bx+a)^5 - \cos(bx+a)^3)\sqrt{d}\log\left(\frac{\cos(bx+a) + 1}{\cos(bx+a) - 1}\right) + 8(45\cos(bx+a)^4 - 36\cos(bx+a)^2 - 4)\sqrt{d\cos(bx+a)}}{40(d\cos(bx+a))^{9/2}d^2 - (d\cos(bx+a))^{5/2}d^4} / (b \cdot d)$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")

[Out] $\left[\frac{1}{80} \cdot (90 \cdot (\cos(b \cdot x + a)^5 - \cos(b \cdot x + a)^3) \cdot \sqrt{-d} \cdot \arctan(1/2 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) + 1) / (d \cdot \cos(b \cdot x + a))) - 45 \cdot (\cos(b \cdot x + a)^5 - \cos(b \cdot x + a)^3) \cdot \sqrt{-d} \cdot \log((d \cdot \cos(b \cdot x + a)^2 - 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot \sqrt{-d} \cdot (\cos(b \cdot x + a) - 1) - 6 \cdot d \cdot \cos(b \cdot x + a) + d) / (\cos(b \cdot x + a)^2 + 2 \cdot \cos(b \cdot x + a) + 1)) + 8 \cdot (45 \cdot \cos(b \cdot x + a)^4 - 36 \cdot \cos(b \cdot x + a)^2 - 4) \cdot \sqrt{d \cdot \cos(b \cdot x + a)}}{(b \cdot d^4 \cdot \cos(b \cdot x + a)^5 - b \cdot d^4 \cdot \cos(b \cdot x + a)^3)}, \frac{1}{80} \cdot (90 \cdot (\cos(b \cdot x + a)^5 - \cos(b \cdot x + a)^3) \cdot \sqrt{d} \cdot \arctan(1/2 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot (\cos(b \cdot x + a) - 1) / (\sqrt{d} \cdot \cos(b \cdot x + a))) + 45 \cdot (\cos(b \cdot x + a)^5 - \cos(b \cdot x + a)^3) \cdot \sqrt{d} \cdot \log((d \cdot \cos(b \cdot x + a)^2 - 4 \cdot \sqrt{d \cdot \cos(b \cdot x + a)}) \cdot \sqrt{d} \cdot (\cos(b \cdot x + a) + 1) + 6 \cdot d \cdot \cos(b \cdot x + a) + d) / (\cos(b \cdot x + a)^2 - 2 \cdot \cos(b \cdot x + a) + 1)) + 8 \cdot (45 \cdot \cos(b \cdot x + a)^4 - 36 \cdot \cos(b \cdot x + a)^2 - 4) \cdot \sqrt{d \cdot \cos(b \cdot x + a)}}{(b \cdot d^4 \cdot \cos(b \cdot x + a)^5 - b \cdot d^4 \cdot \cos(b \cdot x + a)^3)} \right]$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)**3/(d*cos(b*x+a))**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(b*x+a)^3/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate(csc(b*x + a)^3/(d*cos(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(a + bx)^3 (d \cos(a + bx))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(7/2)), x)

[Out] int(1/(sin(a + b*x)^3*(d*cos(a + b*x))^(7/2)), x)

3.252 $\int \sqrt[5]{d \cos(a + bx)} \sin(a + bx) dx$

Optimal. Leaf size=22

$$-\frac{5(d \cos(a + bx))^{6/5}}{6bd}$$

[Out] $-5/6*(d*\cos(b*x+a))^{(6/5)}/b/d$

Rubi [A]

time = 0.02, antiderivative size = 22, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 30}

$$-\frac{5(d \cos(a + bx))^{6/5}}{6bd}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(1/5)}*\text{Sin}[a + b*x], x]$

[Out] $(-5*(d*\text{Cos}[a + b*x])^{(6/5)})/(6*b*d)$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] := \text{Simp}[x^{(m + 1)}/(m + 1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2645

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.))^{(m_.)}*\sin[(e_.) + (f_.)*(x_)]^{(n_.)}, x_Symbol] := \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{((n - 1)/2)}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n - 1)/2] \ \&\& \ !(\text{IntegerQ}[(m - 1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int \sqrt[5]{d \cos(a + bx)} \sin(a + bx) dx &= -\frac{\text{Subst}(\int \sqrt[5]{x} dx, x, d \cos(a + bx))}{bd} \\ &= -\frac{5(d \cos(a + bx))^{6/5}}{6bd} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 22, normalized size = 1.00

$$-\frac{5(d \cos(a + bx))^{6/5}}{6bd}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(1/5)*sin[a + b*x],x]

[Out] $(-5*(d*\cos[a + b*x])^{6/5})/(6*b*d)$

Maple [A]

time = 0.03, size = 19, normalized size = 0.86

method	result	size
derivativedivides	$-\frac{5(d\cos(bx+a))^{6/5}}{6bd}$	19
default	$-\frac{5(d\cos(bx+a))^{6/5}}{6bd}$	19

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/5)*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] $-5/6*(d*\cos(b*x+a))^{6/5}/b/d$

Maxima [A]

time = 0.27, size = 18, normalized size = 0.82

$$-\frac{5(d\cos(bx+a))^{6/5}}{6bd}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/5)*sin(b*x+a),x, algorithm="maxima")

[Out] $-5/6*(d*\cos(b*x + a))^{6/5}/(b*d)$

Fricas [A]

time = 0.35, size = 21, normalized size = 0.95

$$-\frac{5(d\cos(bx+a))^{1/5}\cos(bx+a)}{6b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/5)*sin(b*x+a),x, algorithm="fricas")

[Out] $-5/6*(d*\cos(b*x + a))^{1/5}*\cos(b*x + a)/b$

Sympy [A]

time = 2.20, size = 37, normalized size = 1.68

$$\begin{cases} -\frac{5\sqrt[5]{d\cos(a+bx)}\cos(a+bx)}{6b} & \text{for } b \neq 0 \\ x\sqrt[5]{d\cos(a)}\sin(a) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/5)*sin(b*x+a),x)

[Out] Piecewise((-5*(d*cos(a + b*x))**(1/5)*cos(a + b*x)/(6*b), Ne(b, 0)), (x*(d*cos(a))**(1/5)*sin(a), True))

Giac [A]

time = 3.97, size = 21, normalized size = 0.95

$$\frac{5 (d \cos (bx + a))^{\frac{1}{5}} \cos (bx + a)}{6 b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/5)*sin(b*x+a),x, algorithm="giac")

[Out] -5/6*(d*cos(b*x + a))^(1/5)*cos(b*x + a)/b

Mupad [B]

time = 0.10, size = 18, normalized size = 0.82

$$\frac{5 (d \cos (a + bx))^{6/5}}{6 b d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d*cos(a + b*x))^(1/5),x)

[Out] -(5*(d*cos(a + b*x))^(6/5))/(6*b*d)

3.253 $\int \cos^3(x) \sqrt{\sin(x)} dx$

Optimal. Leaf size=21

$$\frac{2}{3} \sin^{\frac{3}{2}}(x) - \frac{2}{7} \sin^{\frac{7}{2}}(x)$$

[Out] 2/3*sin(x)^(3/2)-2/7*sin(x)^(7/2)

Rubi [A]

time = 0.02, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2644, 14}

$$\frac{2}{3} \sin^{\frac{3}{2}}(x) - \frac{2}{7} \sin^{\frac{7}{2}}(x)$$

Antiderivative was successfully verified.

[In] Int[Cos[x]^3*Sqrt[Sin[x]],x]

[Out] (2*Sin[x]^(3/2))/3 - (2*Sin[x]^(7/2))/7

Rule 14

Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(x) \sqrt{\sin(x)} dx &= \text{Subst} \left(\int \sqrt{x} (1 - x^2) dx, x, \sin(x) \right) \\ &= \text{Subst} \left(\int (\sqrt{x} - x^{5/2}) dx, x, \sin(x) \right) \\ &= \frac{2}{3} \sin^{\frac{3}{2}}(x) - \frac{2}{7} \sin^{\frac{7}{2}}(x) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 18, normalized size = 0.86

$$\frac{1}{21}(11 + 3 \cos(2x)) \sin^{\frac{3}{2}}(x)$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[x]^3*Sqrt[Sin[x]],x]``[Out] ((11 + 3*Cos[2*x])*Sin[x]^(3/2))/21`**Maple [A]**

time = 0.13, size = 14, normalized size = 0.67

method	result	size
default	$\frac{2(\sin^{\frac{3}{2}}(x))}{3} - \frac{2(\sin^{\frac{7}{2}}(x))}{7}$	14

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(x)^3*sin(x)^(1/2),x,method=_RETURNVERBOSE)``[Out] 2/3*sin(x)^(3/2)-2/7*sin(x)^(7/2)`**Maxima [A]**

time = 0.32, size = 13, normalized size = 0.62

$$-\frac{2}{7} \sin(x)^{\frac{7}{2}} + \frac{2}{3} \sin(x)^{\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3*sin(x)^(1/2),x, algorithm="maxima")``[Out] -2/7*sin(x)^(7/2) + 2/3*sin(x)^(3/2)`**Fricas [A]**

time = 0.36, size = 14, normalized size = 0.67

$$\frac{2}{21} (3 \cos(x)^2 + 4) \sin(x)^{\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3*sin(x)^(1/2),x, algorithm="fricas")``[Out] 2/21*(3*cos(x)^2 + 4)*sin(x)^(3/2)`

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 170 vs. 2(19) = 38.

time = 4.70, size = 170, normalized size = 8.10

$$\frac{28\sqrt{2} \sqrt{\frac{\tan(\frac{x}{2})}{\tan^2(\frac{x}{2}) + 1}} \tan^5(\frac{x}{2})}{21 \tan^6(\frac{x}{2}) + 63 \tan^4(\frac{x}{2}) + 63 \tan^2(\frac{x}{2}) + 21} + \frac{8\sqrt{2} \sqrt{\frac{\tan(\frac{x}{2})}{\tan^2(\frac{x}{2}) + 1}} \tan^3(\frac{x}{2})}{21 \tan^6(\frac{x}{2}) + 63 \tan^4(\frac{x}{2}) + 63 \tan^2(\frac{x}{2}) + 21} + \frac{28\sqrt{2} \sqrt{\frac{\tan(\frac{x}{2})}{\tan^2(\frac{x}{2}) + 1}} \tan(\frac{x}{2})}{21 \tan^6(\frac{x}{2}) + 63 \tan^4(\frac{x}{2}) + 63 \tan^2(\frac{x}{2}) + 21}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)**3*sin(x)**(1/2),x)

[Out] 28*sqrt(2)*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**5/(21*tan(x/2)**6 + 63*tan(x/2)**4 + 63*tan(x/2)**2 + 21) + 8*sqrt(2)*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**3/(21*tan(x/2)**6 + 63*tan(x/2)**4 + 63*tan(x/2)**2 + 21) + 28*sqrt(2)*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)/(21*tan(x/2)**6 + 63*tan(x/2)**4 + 63*tan(x/2)**2 + 21)

Giac [A]

time = 5.88, size = 13, normalized size = 0.62

$$-\frac{2}{7} \sin(x)^{\frac{7}{2}} + \frac{2}{3} \sin(x)^{\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3*sin(x)^(1/2),x, algorithm="giac")

[Out] -2/7*sin(x)^(7/2) + 2/3*sin(x)^(3/2)

Mupad [B]

time = 0.47, size = 25, normalized size = 1.19

$$-\frac{\cos(x)^4 \sin(x)^{3/2} {}_2F_1\left(\frac{1}{4}, 2; 3; \cos(x)^2\right)}{4 (\sin(x)^2)^{3/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^3*sin(x)^(1/2),x)

[Out] -(cos(x)^4*sin(x)^(3/2)*hypergeom([1/4, 2], 3, cos(x)^2))/(4*(sin(x)^2)^(3/4))

3.254 $\int \cos^3(x) \sin^{\frac{3}{2}}(x) dx$

Optimal. Leaf size=21

$$\frac{2}{5} \sin^{\frac{5}{2}}(x) - \frac{2}{9} \sin^{\frac{9}{2}}(x)$$

[Out] 2/5*sin(x)^(5/2)-2/9*sin(x)^(9/2)

Rubi [A]

time = 0.02, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2644, 14}

$$\frac{2}{5} \sin^{\frac{5}{2}}(x) - \frac{2}{9} \sin^{\frac{9}{2}}(x)$$

Antiderivative was successfully verified.

[In] Int[Cos[x]^3*Sin[x]^(3/2),x]

[Out] (2*Sin[x]^(5/2))/5 - (2*Sin[x]^(9/2))/9

Rule 14

```
Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2644

```
Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \cos^3(x) \sin^{\frac{3}{2}}(x) dx &= \text{Subst} \left(\int x^{3/2} (1 - x^2) dx, x, \sin(x) \right) \\ &= \text{Subst} \left(\int (x^{3/2} - x^{7/2}) dx, x, \sin(x) \right) \\ &= \frac{2}{5} \sin^{\frac{5}{2}}(x) - \frac{2}{9} \sin^{\frac{9}{2}}(x) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 18, normalized size = 0.86

$$\frac{1}{45}(13 + 5 \cos(2x)) \sin^{\frac{5}{2}}(x)$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[x]^3*Sin[x]^(3/2),x]``[Out] ((13 + 5*Cos[2*x])*Sin[x]^(5/2))/45`**Maple [A]**

time = 0.14, size = 14, normalized size = 0.67

method	result	size
default	$\frac{2(\sin^{\frac{5}{2}}(x))}{5} - \frac{2(\sin^{\frac{9}{2}}(x))}{9}$	14

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(x)^3*sin(x)^(3/2),x,method=_RETURNVERBOSE)``[Out] 2/5*sin(x)^(5/2)-2/9*sin(x)^(9/2)`**Maxima [A]**

time = 0.28, size = 13, normalized size = 0.62

$$-\frac{2}{9} \sin(x)^{\frac{9}{2}} + \frac{2}{5} \sin(x)^{\frac{5}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3*sin(x)^(3/2),x, algorithm="maxima")``[Out] -2/9*sin(x)^(9/2) + 2/5*sin(x)^(5/2)`**Fricas [A]**

time = 0.38, size = 20, normalized size = 0.95

$$-\frac{2}{45} (5 \cos(x)^4 - \cos(x)^2 - 4) \sqrt{\sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3*sin(x)^(3/2),x, algorithm="fricas")``[Out] -2/45*(5*cos(x)^4 - cos(x)^2 - 4)*sqrt(sin(x))`**Sympy [A]**

time = 6.43, size = 24, normalized size = 1.14

$$\frac{8 \sin^{\frac{9}{2}}(x)}{45} + \frac{2 \sin^{\frac{5}{2}}(x) \cos^2(x)}{5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)**3*sin(x)**(3/2),x)`

[Out] `8*sin(x)**(9/2)/45 + 2*sin(x)**(5/2)*cos(x)**2/5`

Giac [A]

time = 6.22, size = 13, normalized size = 0.62

$$-\frac{2}{9} \sin(x)^{\frac{9}{2}} + \frac{2}{5} \sin(x)^{\frac{5}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)^3*sin(x)^(3/2),x, algorithm="giac")`

[Out] `-2/9*sin(x)^(9/2) + 2/5*sin(x)^(5/2)`

Mupad [B]

time = 0.47, size = 25, normalized size = 1.19

$$-\frac{\cos(x)^4 \sin(x)^{5/2} {}_2F_1\left(-\frac{1}{4}, 2; 3; \cos(x)^2\right)}{4 (\sin(x)^2)^{5/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x)^3*sin(x)^(3/2),x)`

[Out] `-(cos(x)^4*sin(x)^(5/2)*hypergeom([-1/4, 2], 3, cos(x)^2))/(4*(sin(x)^2)^(5/4))`

3.255 $\int \cos^3(x) \sin^{\frac{5}{2}}(x) dx$

Optimal. Leaf size=21

$$\frac{2}{7} \sin^{\frac{7}{2}}(x) - \frac{2}{11} \sin^{\frac{11}{2}}(x)$$

[Out] 2/7*sin(x)^(7/2)-2/11*sin(x)^(11/2)

Rubi [A]

time = 0.02, antiderivative size = 21, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2644, 14}

$$\frac{2}{7} \sin^{\frac{7}{2}}(x) - \frac{2}{11} \sin^{\frac{11}{2}}(x)$$

Antiderivative was successfully verified.

[In] Int[Cos[x]^3*Sin[x]^(5/2),x]

[Out] (2*Sin[x]^(7/2))/7 - (2*Sin[x]^(11/2))/11

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] :> Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2644

Int[cos[(e_)+(f_)*(x_)]^(n_)*((a_)*sin[(e_)+(f_)*(x_)])^(m_), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1-x^2/a^2)^((n-1)/2), x], x, a*Sin[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(x) \sin^{\frac{5}{2}}(x) dx &= \text{Subst} \left(\int x^{5/2} (1-x^2) dx, x, \sin(x) \right) \\ &= \text{Subst} \left(\int (x^{5/2} - x^{9/2}) dx, x, \sin(x) \right) \\ &= \frac{2}{7} \sin^{\frac{7}{2}}(x) - \frac{2}{11} \sin^{\frac{11}{2}}(x) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 18, normalized size = 0.86

$$\frac{1}{77}(15 + 7 \cos(2x)) \sin^{\frac{7}{2}}(x)$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[x]^3*Sin[x]^(5/2),x]``[Out] ((15 + 7*Cos[2*x])*Sin[x]^(7/2))/77`**Maple [A]**

time = 0.13, size = 14, normalized size = 0.67

method	result	size
default	$\frac{2(\sin^{\frac{7}{2}}(x))}{7} - \frac{2(\sin^{\frac{11}{2}}(x))}{11}$	14

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(x)^3*sin(x)^(5/2),x,method=_RETURNVERBOSE)``[Out] 2/7*sin(x)^(7/2)-2/11*sin(x)^(11/2)`**Maxima [A]**

time = 0.28, size = 13, normalized size = 0.62

$$-\frac{2}{11} \sin(x)^{\frac{11}{2}} + \frac{2}{7} \sin(x)^{\frac{7}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3*sin(x)^(5/2),x, algorithm="maxima")``[Out] -2/11*sin(x)^(11/2) + 2/7*sin(x)^(7/2)`**Fricas [A]**

time = 0.35, size = 20, normalized size = 0.95

$$-\frac{2}{77} (7 \cos(x)^4 - 3 \cos(x)^2 - 4) \sin(x)^{\frac{3}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(x)^3*sin(x)^(5/2),x, algorithm="fricas")``[Out] -2/77*(7*cos(x)^4 - 3*cos(x)^2 - 4)*sin(x)^(3/2)`**Sympy [A]**

time = 44.49, size = 24, normalized size = 1.14

$$\frac{8 \sin^{\frac{11}{2}}(x)}{77} + \frac{2 \sin^{\frac{7}{2}}(x) \cos^2(x)}{7}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)**3*sin(x)**(5/2),x)`

[Out] `8*sin(x)**(11/2)/77 + 2*sin(x)**(7/2)*cos(x)**2/7`

Giac [A]

time = 6.48, size = 13, normalized size = 0.62

$$-\frac{2}{11} \sin(x)^{\frac{11}{2}} + \frac{2}{7} \sin(x)^{\frac{7}{2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)^3*sin(x)^(5/2),x, algorithm="giac")`

[Out] `-2/11*sin(x)^(11/2) + 2/7*sin(x)^(7/2)`

Mupad [B]

time = 0.45, size = 25, normalized size = 1.19

$$-\frac{\cos(x)^4 \sin(x)^{7/2} {}_2F_1\left(-\frac{3}{4}, 2; 3; \cos(x)^2\right)}{4 (\sin(x)^2)^{7/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x)^3*sin(x)^(5/2),x)`

[Out] `-(cos(x)^4*sin(x)^(7/2)*hypergeom([-3/4, 2], 3, cos(x)^2))/(4*(sin(x)^2)^(7/4))`

$$3.256 \quad \int \frac{\cos^3(x)}{\sqrt{\sin(x)}} dx$$

Optimal. Leaf size=19

$$2\sqrt{\sin(x)} - \frac{2}{5}\sin^{\frac{5}{2}}(x)$$

[Out] -2/5*sin(x)^(5/2)+2*sin(x)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {2644, 14}

$$2\sqrt{\sin(x)} - \frac{2}{5}\sin^{\frac{5}{2}}(x)$$

Antiderivative was successfully verified.

[In] Int[Cos[x]^3/Sqrt[Sin[x]],x]

[Out] 2*Sqrt[Sin[x]] - (2*Sin[x]^(5/2))/5

Rule 14

```
Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]
```

Rule 2644

```
Int[cos[(e_)+(f_)*(x_)]^(n_)*((a_)*sin[(e_)+(f_)*(x_)])^(m_), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1-x^2/a^2)^((n-1)/2), x], x, a*Sin[e+f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \frac{\cos^3(x)}{\sqrt{\sin(x)}} dx &= \text{Subst} \left(\int \frac{1-x^2}{\sqrt{x}} dx, x, \sin(x) \right) \\ &= \text{Subst} \left(\int \left(\frac{1}{\sqrt{x}} - x^{3/2} \right) dx, x, \sin(x) \right) \\ &= 2\sqrt{\sin(x)} - \frac{2}{5}\sin^{\frac{5}{2}}(x) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 0.84

$$\frac{1}{5}(9 + \cos(2x))\sqrt{\sin(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[x]^3/Sqrt[Sin[x]],x]**[Out]** ((9 + Cos[2*x])*Sqrt[Sin[x]])/5**Maple [A]**

time = 0.15, size = 14, normalized size = 0.74

method	result	size
default	$-\frac{2(\sin^{\frac{5}{2}}(x))}{5} + 2(\sqrt{\sin(x)})$	14

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^3/sin(x)^(1/2),x,method=_RETURNVERBOSE)**[Out]** -2/5*sin(x)^(5/2)+2*sin(x)^(1/2)**Maxima [A]**

time = 0.28, size = 13, normalized size = 0.68

$$-\frac{2}{5} \sin(x)^{\frac{5}{2}} + 2 \sqrt{\sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3/sin(x)^(1/2),x, algorithm="maxima")**[Out]** -2/5*sin(x)^(5/2) + 2*sqrt(sin(x))**Fricas [A]**

time = 0.38, size = 12, normalized size = 0.63

$$\frac{2}{5} (\cos(x)^2 + 4) \sqrt{\sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3/sin(x)^(1/2),x, algorithm="fricas")**[Out]** 2/5*(cos(x)^2 + 4)*sqrt(sin(x))**Sympy [B]** Leaf count of result is larger than twice the leaf count of optimal. 323 vs. 2(17) = 34.

time = 5.31, size = 323, normalized size = 17.00

$$\frac{10\sqrt{2}\tan^3(\frac{x}{5})}{5\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}\sqrt{\tan^2(\frac{x}{5})+15}\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}\sqrt{\tan^2(\frac{x}{5})+15}\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}} + \frac{12\sqrt{2}\tan^3(\frac{x}{5})}{5\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}\sqrt{\tan^2(\frac{x}{5})+15}\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}\sqrt{\tan^2(\frac{x}{5})+15}\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}} + \frac{10\sqrt{2}\tan(\frac{x}{5})}{5\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}\sqrt{\tan^2(\frac{x}{5})+15}\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}\sqrt{\tan^2(\frac{x}{5})+15}\sqrt{\frac{\tan(\frac{x}{5})}{\tan^2(\frac{x}{5})+1}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)**3/sin(x)**(1/2),x)

[Out] 10*sqrt(2)*tan(x/2)**5/(5*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**6 + 15*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**4 + 15*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**2 + 5*sqrt(tan(x/2)/(tan(x/2)**2 + 1)))) + 12*sqrt(2)*tan(x/2)**3/(5*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**6 + 15*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**4 + 15*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**2 + 5*sqrt(tan(x/2)/(tan(x/2)**2 + 1)))) + 10*sqrt(2)*tan(x/2)/(5*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**6 + 15*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**4 + 15*sqrt(tan(x/2)/(tan(x/2)**2 + 1))*tan(x/2)**2 + 5*sqrt(tan(x/2)/(tan(x/2)**2 + 1))))

Giac [A]

time = 5.75, size = 13, normalized size = 0.68

$$-\frac{2}{5} \sin(x)^{\frac{5}{2}} + 2 \sqrt{\sin(x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(x)^3/sin(x)^(1/2),x, algorithm="giac")

[Out] -2/5*sin(x)^(5/2) + 2*sqrt(sin(x))

Mupad [B]

time = 0.44, size = 25, normalized size = 1.32

$$\frac{\cos(x)^4 \sqrt{\sin(x)} {}_2F_1\left(\frac{3}{4}, 2; 3; \cos(x)^2\right)}{4 (\sin(x)^2)^{1/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(x)^3/sin(x)^(1/2),x)

[Out] -(cos(x)^4*sin(x)^(1/2)*hypergeom([3/4, 2], 3, cos(x)^2))/(4*(sin(x)^2)^(1/4))

3.257 $\int (d \cos(a + bx))^{9/2} \sqrt{c \sin(a + bx)} dx$

Optimal. Leaf size=132

$$\frac{7d^3(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{30bc} + \frac{d(d \cos(a + bx))^{7/2}(c \sin(a + bx))^{3/2}}{5bc} + \frac{7d^4 \sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4}, \frac{2}{\sin(2a + 2bx)}\right)}{20b \sqrt{\sin(2a + 2bx)}}$$

[Out] $7/30*d^3*(d*\cos(b*x+a))^{(3/2)}*(c*\sin(b*x+a))^{(3/2)}/b/c+1/5*d*(d*\cos(b*x+a))^{(7/2)}*(c*\sin(b*x+a))^{(3/2)}/b/c-7/20*d^4*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2649, 2652, 2719}

$$\frac{7d^4 E\left(a + bx - \frac{\pi}{4}, 2\right) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{20b \sqrt{\sin(2a + 2bx)}} + \frac{7d^3 (c \sin(a + bx))^{3/2} (d \cos(a + bx))^{3/2}}{30bc} + \frac{d (c \sin(a + bx))^{3/2} (d \cos(a + bx))^{7/2}}{5bc}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(9/2)}*\text{Sqrt}[c*\text{Sin}[a + b*x]], x]$

[Out] $(7*d^3*(d*\text{Cos}[a + b*x])^{(3/2)}*(c*\text{Sin}[a + b*x])^{(3/2)})/(30*b*c) + (d*(d*\text{Cos}[a + b*x])^{(7/2)}*(c*\text{Sin}[a + b*x])^{(3/2)})/(5*b*c) + (7*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(20*b*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2649

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[a*(b*\text{Sin}[e + f*x])^{(n + 1)}*((a*\text{Cos}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Sin}[e + f*x])^n*(a*\text{Cos}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x\} \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2652

$\text{Int}[\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[a*\text{Sin}[e + f*x]]*(\text{Sqrt}[b*\text{Cos}[e + f*x]]/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]), \text{Int}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x\}$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x\}$

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{9/2} \sqrt{c \sin(a + bx)} \, dx &= \frac{d(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bc} + \frac{1}{10} (7d^2) \int (d \cos(a + bx))^5 \\
&= \frac{7d^3 (d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{30bc} + \frac{d(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bc} \\
&= \frac{7d^3 (d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{30bc} + \frac{d(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bc} \\
&= \frac{7d^3 (d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{30bc} + \frac{d(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bc}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.07, size = 70, normalized size = 0.53

$$\frac{2d^4 \sqrt{d \cos(a + bx)} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(-\frac{7}{4}, \frac{3}{4}; \frac{7}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)} \tan(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(9/2)*Sqrt[c*Sin[a + b*x]],x]

[Out] (2*d^4*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-7/4, 3/4, 7/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 531 vs. 2(137) = 274.

time = 1.60, size = 532, normalized size = 4.03

method	result
default	$-\frac{\sqrt{c \sin(bx + a)} (d \cos(bx + a))^{\frac{9}{2}} \left(12(\cos^6(bx + a)) \sqrt{2} + 2(\cos^4(bx + a)) \sqrt{2} + 42 \cos(bx + a) \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \right)}{3b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/120/b*(c*sin(b*x+a))^(1/2)*(d*cos(b*x+a))^(9/2)*(12*cos(b*x+a)^6*2^(1/2)+2*cos(b*x+a)^4*2^(1/2)+42*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-21*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos

$$\begin{aligned} & (b*x+a)/\sin(b*x+a))^{(1/2)}*EllipticF(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a)) \\ & ^{(1/2)}, 1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}+42*((-1+\cos \\ & s(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*E \\ & llipticE(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})*((1-\cos(\\ & b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}-21*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x \\ & +a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*EllipticF(((1-\cos(b*x+a)+\sin(\\ & b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a \\ &))^{(1/2)}+7*\cos(b*x+a)^2*2^{(1/2)}-21*\cos(b*x+a)*2^{(1/2)})/\sin(b*x+a)/\cos(b*x+a \\ &)^5*2^{(1/2)} \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^(9/2)*sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*d^4*cos(b*x + a)^4, x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(9/2)*(c*sin(b*x+a))**(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(9/2)*sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + b x))^{9/2} \sqrt{c \sin(a + b x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(9/2)*(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^(9/2)*(c*sin(a + b*x))^(1/2), x)

3.258 $\int (d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)} dx$

Optimal. Leaf size=95

$$\frac{d(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{3bc} + \frac{d^2 \sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a + bx)}}{2b \sqrt{\sin(2a + 2bx)}}$$

[Out] 1/3*d*(d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2)/b/c-1/2*d^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))*(d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2)/b/sin(2*b*x+2*a)^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2649, 2652, 2719}

$$\frac{d^2 E\left(a + bx - \frac{\pi}{4} \mid 2\right) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{2b \sqrt{\sin(2a + 2bx)}} + \frac{d(c \sin(a + bx))^{3/2}(d \cos(a + bx))^{3/2}}{3bc}$$

Antiderivative was successfully verified.

[In] Int[(d*Cos[a + b*x])^(5/2)*Sqrt[c*Sin[a + b*x]],x]

[Out] (d*(d*Cos[a + b*x])^(3/2)*(c*Sin[a + b*x])^(3/2))/(3*b*c) + (d^2*Sqrt[d*Cos[a + b*x]]*EllipticE[a - Pi/4 + b*x, 2]*Sqrt[c*Sin[a + b*x]])/(2*b*Sqrt[Sin[2*a + 2*b*x]])

Rule 2649

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[a*(b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]], x_Symbol] :> Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)} dx &= \frac{d(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{3bc} + \frac{1}{2} d^2 \int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx \\ &= \frac{d(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{3bc} + \frac{\left(d^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} \right)}{2\sqrt{\sin(2a + 2bx)}} \\ &= \frac{d(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{3bc} + \frac{d^2 \sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4}, \sin(2a + 2bx)\right)}{2b\sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.06, size = 70, normalized size = 0.74

$$\frac{2d^2 \sqrt{d \cos(a + bx)} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(-\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)} \tan(a + bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(5/2)*Sqrt[c*Sin[a + b*x]],x]

[Out] (2*d^2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-3/4, 3/4, 7/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 517 vs. 2(106) = 212.

time = 0.15, size = 518, normalized size = 5.45

method	result
default	$-\frac{\sqrt{c \sin(bx + a)} (d \cos(bx + a))^{\frac{5}{2}} \left(2(\cos^4(bx + a)) \sqrt{2} + 6 \cos(bx + a) \sqrt{\frac{-1 + \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}} \sqrt{\frac{-1 + \cos(bx + a)}{\sin(bx + a)}} \right)}{12b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/12/b*(c*sin(b*x+a))^(1/2)*(d*cos(b*x+a))^(5/2)*(2*cos(b*x+a)^4*2^(1/2)+6*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-3*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+6*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2)))

$$\begin{aligned} & b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a) \\ &))^{(1/2)}-3*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)} \\ & *EllipticF(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}, 1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}+\cos(b*x+a)^2*2^{(1/2)} \\ & -3*\cos(b*x+a)*2^{(1/2)})/\sin(b*x+a)/\cos(b*x+a)^3*2^{(1/2)} \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^(5/2)*sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*d^2*cos(b*x + a)^2, x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(5/2)*(c*sin(b*x+a))**(1/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 8570 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(5/2)*sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + b x))^{5/2} \sqrt{c \sin(a + b x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(5/2)*(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^(5/2)*(c*sin(a + b*x))^(1/2), x)

$$3.259 \quad \int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx$$

Optimal. Leaf size=53

$$\frac{\sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a + bx)}}{b \sqrt{\sin(2a + 2bx)}}$$

[Out] $-(\sin(a+1/4*\pi+b*x)^2)^{(1/2)}/\sin(a+1/4*\pi+b*x)*\text{EllipticE}(\cos(a+1/4*\pi+b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2652, 2719}

$$\frac{E\left(a + bx - \frac{\pi}{4} \mid 2\right) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{b \sqrt{\sin(2a + 2bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]], x]

[Out] (Sqrt[d*Cos[a + b*x]]*EllipticE[a - Pi/4 + b*x, 2]*Sqrt[c*Sin[a + b*x]])/(b*Sqrt[Sin[2*a + 2*b*x]])

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]] , x_Symbol] :> Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]] , x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx &= \frac{\left(\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}\right) \int \sqrt{\sin(2a + 2bx)} dx}{\sqrt{\sin(2a + 2bx)}} \\ &= \frac{\sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a + bx)}}{b \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.04, size = 67, normalized size = 1.26

$$\frac{2\sqrt{d\cos(a+bx)}\sqrt[4]{\cos^2(a+bx)}{}_2F_1\left(\frac{1}{4},\frac{3}{4};\frac{7}{4};\sin^2(a+bx)\right)\sqrt{c\sin(a+bx)}\tan(a+bx)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]],x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 3/4, 7/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 504 vs. 2(73) = 146.

time = 0.15, size = 505, normalized size = 9.53

method	result
default	$-\frac{\left(2\cos(bx+a)\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}\operatorname{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}},\frac{\sqrt{2}}{2}\right)\sqrt{\frac{1-\cos(bx+a)}{\sin(bx+a)}}\right)}{3b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/2/b*(2*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+2*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+cos(b*x+a)^2*2^(1/2)-cos(b*x+a)*2^(1/2))*(d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2)/sin(b*x+a)/cos(b*x+a)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)*(c*sin(b*x+a))**(1/2),x)

[Out] Integral(sqrt(c*sin(a + b*x))*sqrt(d*cos(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(1/2), x)

$$3.260 \quad \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{3/2}} dx$$

Optimal. Leaf size=93

$$\frac{2(c \sin(a + bx))^{3/2}}{bcd \sqrt{d \cos(a + bx)}} - \frac{2 \sqrt{d \cos(a + bx)} E(a - \frac{\pi}{4} + bx | 2) \sqrt{c \sin(a + bx)}}{bd^2 \sqrt{\sin(2a + 2bx)}}$$

[Out] $2*(c*\sin(b*x+a))^{(3/2)}/b/c/d/(d*\cos(b*x+a))^{(1/2)}+2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x),2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b/d^2/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2651, 2652, 2719}

$$\frac{2(c \sin(a + bx))^{3/2}}{bcd \sqrt{d \cos(a + bx)}} - \frac{2E(a + bx - \frac{\pi}{4} | 2) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{bd^2 \sqrt{\sin(2a + 2bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(3/2),x]

[Out] $(2*(c*\sin[a + b*x])^{(3/2)})/(b*c*d*\text{Sqrt}[d*\cos[a + b*x]]) - (2*\text{Sqrt}[d*\cos[a + b*x]]*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\sin[a + b*x]])/(b*d^2*\text{Sqrt}[\sin[2*a + 2*b*x]])$

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (b*Sin[e + f*x])^(n + 1))*((a*Cos[e + f*x])^(m + 1))/(a*b*f*(m + 1)), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c \sin(a+bx)}}{(d \cos(a+bx))^{3/2}} dx &= \frac{2(c \sin(a+bx))^{3/2}}{bcd \sqrt{d \cos(a+bx)}} - \frac{2 \int \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)} dx}{d^2} \\ &= \frac{2(c \sin(a+bx))^{3/2}}{bcd \sqrt{d \cos(a+bx)}} - \frac{\left(2 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}\right) \int \sqrt{\sin(2a+2bx)} dx}{d^2 \sqrt{\sin(2a+2bx)}} \\ &= \frac{2(c \sin(a+bx))^{3/2}}{bcd \sqrt{d \cos(a+bx)}} - \frac{2 \sqrt{d \cos(a+bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a+bx)}}{bd^2 \sqrt{\sin(2a+2bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.08, size = 70, normalized size = 0.75

$$\frac{2 \sqrt{d \cos(a+bx)} \sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{3}{4}, \frac{5}{4}; \frac{7}{4}; \sin^2(a+bx)\right) \sqrt{c \sin(a+bx)} \tan(a+bx)}{3bd^2}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(3/2), x]

[Out] (2*sqrt[d*cos[a + b*x]]*(cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[3/4, 5/4, 7/4, Sin[a + b*x]^2]*sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b*d^2)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 492 vs. 2(108) = 216.

time = 1.23, size = 493, normalized size = 5.30

method	result
default	$-\frac{\left(-2 \cos(bx+a) \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)}{3bd^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(3/2), x, method=_RETURNVERBOSE)

[Out] -1/b*(-2*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-2*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)

```
in(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)
)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin
(b*x+a))^(1/2)+((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a
))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)
,1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+cos(b*x+a)*2^(1/
2)-2^(1/2))*cos(b*x+a)*(c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(3/2)/sin(b*x+a)
*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(3/2), x)
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly
1 arguments (2 given)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))**(1/2)/(d*cos(b*x+a))**(3/2),x)
```

```
[Out] Integral(sqrt(c*sin(a + b*x))/(d*cos(a + b*x))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c \sin(a + b x)}}{(d \cos(a + b x))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(3/2),x)

[Out] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(3/2), x)

$$3.261 \quad \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{7/2}} dx$$

Optimal. Leaf size=134

$$\frac{2(c \sin(a + bx))^{3/2}}{5bcd(d \cos(a + bx))^{5/2}} + \frac{4(c \sin(a + bx))^{3/2}}{5bcd^3 \sqrt{d \cos(a + bx)}} - \frac{4\sqrt{d \cos(a + bx)} E(a - \frac{\pi}{4} + bx | 2) \sqrt{c \sin(a + bx)}}{5bd^4 \sqrt{\sin(2a + 2bx)}}$$

[Out] 2/5*(c*sin(b*x+a))^(3/2)/b/c/d/(d*cos(b*x+a))^(5/2)+4/5*(c*sin(b*x+a))^(3/2)/b/c/d^3/(d*cos(b*x+a))^(1/2)+4/5*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))*(d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2)/b/d^4/sin(2*b*x+2*a)^(1/2)

Rubi [A]

time = 0.12, antiderivative size = 134, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2651, 2652, 2719}

$$-\frac{4E(a + bx - \frac{\pi}{4} | 2) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{5bd^4 \sqrt{\sin(2a + 2bx)}} + \frac{4(c \sin(a + bx))^{3/2}}{5bcd^3 \sqrt{d \cos(a + bx)}} + \frac{2(c \sin(a + bx))^{3/2}}{5bcd(d \cos(a + bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(7/2),x]

[Out] (2*(c*Sin[a + b*x])^(3/2))/(5*b*c*d*(d*Cos[a + b*x])^(5/2)) + (4*(c*Sin[a + b*x])^(3/2))/(5*b*c*d^3*Sqrt[d*Cos[a + b*x]]) - (4*Sqrt[d*Cos[a + b*x]]*EllipticE[a - Pi/4 + b*x, 2]*Sqrt[c*Sin[a + b*x]])/(5*b*d^4*Sqrt[Sin[2*a + 2*b*x]])

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-(b*Sin[e + f*x])^(n + 1))*((a*Cos[e + f*x])^(m + 1))/(a*b*f*(m + 1)), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{c \sin(a+bx)}}{(d \cos(a+bx))^{7/2}} dx &= \frac{2(c \sin(a+bx))^{3/2}}{5bcd(d \cos(a+bx))^{5/2}} + \frac{2 \int \frac{\sqrt{c \sin(a+bx)}}{(d \cos(a+bx))^{3/2}} dx}{5d^2} \\ &= \frac{2(c \sin(a+bx))^{3/2}}{5bcd(d \cos(a+bx))^{5/2}} + \frac{4(c \sin(a+bx))^{3/2}}{5bcd^3 \sqrt{d \cos(a+bx)}} - \frac{4 \int \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}{5d^4} \\ &= \frac{2(c \sin(a+bx))^{3/2}}{5bcd(d \cos(a+bx))^{5/2}} + \frac{4(c \sin(a+bx))^{3/2}}{5bcd^3 \sqrt{d \cos(a+bx)}} - \frac{(4 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)})}{5d^4 \sqrt{\sin(2a+2bx)}} \\ &= \frac{2(c \sin(a+bx))^{3/2}}{5bcd(d \cos(a+bx))^{5/2}} + \frac{4(c \sin(a+bx))^{3/2}}{5bcd^3 \sqrt{d \cos(a+bx)}} - \frac{4 \sqrt{d \cos(a+bx)} E(a - \frac{\pi}{4} + bx)}{5bd^4 \sqrt{\sin(2a+2bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.11, size = 70, normalized size = 0.52

$$\frac{2 \sqrt{d \cos(a+bx)} \sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{3}{4}, \frac{9}{4}; \frac{7}{4}; \sin^2(a+bx)\right) \sqrt{c \sin(a+bx)} \tan(a+bx)}{3bd^4}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(7/2), x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[3/4, 9/4, 7/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b*d^4)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 527 vs. 2(139) = 278.

time = 0.16, size = 528, normalized size = 3.94

method	result
default	$\frac{(4(\cos^3(bx+a)) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \text{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right))}{3bd^4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(7/2), x, method=_RETURNVERBOSE)

```
[Out] 1/5/b*(4*cos(b*x+a)^3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-2*cos(b*x+a)^3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+4*cos(b*x+a)^2*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-2*cos(b*x+a)^2*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-2*cos(b*x+a)^3*2^(1/2)+cos(b*x+a)^2*2^(1/2)+2^(1/2))*cos(b*x+a)*(c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(7/2)/sin(b*x+a)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(7/2), x)
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))**(1/2)/(d*cos(b*x+a))**(7/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{c \sin(a + b x)}}{(d \cos(a + b x))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(7/2),x)

[Out] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(7/2), x)

3.262 $\int (d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)} dx$

Optimal. Leaf size=320

$$-\frac{\sqrt{c} d^{3/2} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{4\sqrt{2} b} + \frac{\sqrt{c} d^{3/2} \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{4\sqrt{2} b} + \frac{\sqrt{c} d^{3/2}}{2bc}$$

[Out] $-1/8*d^{(3/2)}*\arctan(1-2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/c^{(1/2)}/(d*\cos(b*x+a))^{(1/2)})*c^{(1/2)}/b*2^{(1/2)}+1/8*d^{(3/2)}*\arctan(1+2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/c^{(1/2)}/(d*\cos(b*x+a))^{(1/2)})*c^{(1/2)}/b*2^{(1/2)}+1/16*d^{(3/2)}*\ln(c^{(1/2)}-2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(1/2)}+c^{(1/2)}*\tan(b*x+a))*c^{(1/2)}/b*2^{(1/2)}-1/16*d^{(3/2)}*\ln(c^{(1/2)}+2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(1/2)}+c^{(1/2)}*\tan(b*x+a))*c^{(1/2)}/b*2^{(1/2)}+1/2*d*(c*\sin(b*x+a))^{(3/2)}*(d*\cos(b*x+a))^{(1/2)}/b/c$

Rubi [A]

time = 0.18, antiderivative size = 320, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2649, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{c} d^{3/2} \text{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{4\sqrt{2} b} + \frac{\sqrt{c} d^{3/2} \text{ArcTan}\left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} + 1\right)}{4\sqrt{2} b} + \frac{\sqrt{c} d^{3/2} \log\left(\frac{-\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx) + \sqrt{c}\right)}{8\sqrt{2} b} - \frac{\sqrt{c} d^{3/2} \log\left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx) + \sqrt{c}\right)}{8\sqrt{2} b} + \frac{d(c \sin(a + bx))^{3/2} \sqrt{d \cos(a + bx)}}{2bc}$$

Antiderivative was successfully verified.

[In] Int[(d*Cos[a + b*x])^(3/2)*Sqrt[c*Sin[a + b*x]],x]

[Out] $-1/4*(\text{Sqrt}[c]*d^{(3/2)}*\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/(\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])]/(\text{Sqrt}[2]*b) + (\text{Sqrt}[c]*d^{(3/2)}*\text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/(\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])]/(4*\text{Sqrt}[2]*b) + (\text{Sqrt}[c]*d^{(3/2)}*\text{Log}[\text{Sqrt}[c] - (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/\text{Sqrt}[d*\text{Cos}[a + b*x]] + \text{Sqrt}[c]*\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b) - (\text{Sqrt}[c]*d^{(3/2)}*\text{Log}[\text{Sqrt}[c] + (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/\text{Sqrt}[d*\text{Cos}[a + b*x]] + \text{Sqrt}[c]*\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b) + (d*\text{Sqrt}[d*\text{Cos}[a + b*x]]*(c*\text{Sin}[a + b*x])^{(3/2)})/(2*b*c)$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a,

b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2649

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n
_), x_Symbol] := Simp[a*(b*Sine[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m - 1)/
(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Sine[e + f*x])^n*(a*
Cos[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] &&
NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sine[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int (d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)} dx &= \frac{d \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bc} + \frac{1}{4} d^2 \int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx \\
&= \frac{d \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bc} + \frac{(cd^3) \text{Subst}\left(\int \frac{x^2}{c^2 + d^2 x^4} dx, x, \frac{c \sin(a + bx)}{\sqrt{d \cos(a + bx)}}\right)}{2b} \\
&= \frac{d \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bc} - \frac{(cd^2) \text{Subst}\left(\int \frac{c - dx^2}{c^2 + d^2 x^4} dx, x, \frac{c \sin(a + bx)}{\sqrt{d \cos(a + bx)}}\right)}{4b} \\
&= \frac{d \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bc} + \frac{(cd) \text{Subst}\left(\int \frac{1}{\frac{c}{d} - \sqrt{2} \sqrt{c} x + x^2} dx, x, \frac{c \sin(a + bx)}{\sqrt{d \cos(a + bx)}}\right)}{8b} \\
&= \frac{\sqrt{c} d^{3/2} \log\left(\sqrt{c} - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx)\right)}{8\sqrt{2} b} \\
&= -\frac{\sqrt{c} d^{3/2} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{4\sqrt{2} b} + \frac{\sqrt{c} d^{3/2} \tan^{-1}\left(\frac{\sqrt{c} \sin(a + bx)}{\sqrt{d \cos(a + bx)}}\right)}{4\sqrt{2} b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.08, size = 70, normalized size = 0.22

$$\frac{2d^2 \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, \frac{3}{4}, \frac{7}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)} \tan(a + bx)}{3b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(3/2)*Sqrt[c*Sin[a + b*x]],x]

[Out] (2*d^2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, 3/4, 7/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b*Sqrt[d*Cos[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.10, size = 514, normalized size = 1.61

method	result
--------	--------

default	$\left(-i \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticPi} \left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \right) \right)$
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{8}b \cdot (-I \cdot ((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} \cdot \operatorname{EllipticPi}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2-1/2I, 1/2 \cdot 2^{1/2})) + I \cdot ((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} \cdot \operatorname{EllipticPi}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2+1/2I, 1/2 \cdot 2^{1/2})) + ((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} \cdot \operatorname{EllipticPi}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2-1/2I, 1/2 \cdot 2^{1/2})) + ((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2} \cdot ((-1+\cos(bx+a))/\sin(bx+a))^{1/2} \cdot \operatorname{EllipticPi}(((1-\cos(bx+a)+\sin(bx+a))/\sin(bx+a))^{1/2}, 1/2+1/2I, 1/2 \cdot 2^{1/2})) + 2 \cdot \cos(bx+a) \cdot 2^{1/2} - 2 \cdot \cos(bx+a) \cdot 2^{1/2} \cdot (d \cdot \cos(bx+a))^{3/2} \cdot (c \cdot \sin(bx+a))^{1/2} \cdot \sin(bx+a) / (-1+\cos(bx+a)) / \cos(bx+a) \cdot 2^{1/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate((d*cos(b*x + a))^(3/2)*sqrt(c*sin(b*x + a)), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2205 vs. 2(238) = 476.

time = 28.54, size = 2205, normalized size = 6.89

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(1/2),x, algorithm="fricas")`

[Out] $\frac{1}{64} \cdot (32 \cdot \sqrt{d \cdot \cos(bx+a)}) \cdot \sqrt{c \cdot \sin(bx+a)} \cdot d \cdot \sin(bx+a) + 4 \cdot \sqrt{(2 \cdot (c^2 \cdot d^6 / b^4))^{1/4} \cdot b \cdot \arctan(((\sqrt{2} \cdot (c^2 \cdot d^6 / b^4))^{1/4} \cdot b \cdot c^3 \cdot d^8 \cdot \cos(bx+a) + \sqrt{2} \cdot (c^2 \cdot d^6 / b^4))^{3/4} \cdot b^3 \cdot c^2 \cdot d^5 \cdot \sin(bx+a)) \cdot \sqrt{d \cdot \cos(bx+a)} \cdot \sqrt{c \cdot \sin(bx+a)} + \sqrt{4 \cdot \sqrt{c^2 \cdot d^6 / b^4}} \cdot b^2 \cdot c^3 \cdot d^7 \cdot \cos$

$$\begin{aligned}
& (b*x + a)*\sin(b*x + a) + c^4*d^{10} - 2*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))}*(2*c^2*d^5*\cos(b*x + a)*\sin(b*x + a) \\
& + \sqrt{c^2*d^6/b^4}*b^2*c*d^2 + (\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c*d^3*\sin(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*\cos(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} \\
&)/(2*c^4*d^{10}*\cos(b*x + a)^2 - c^4*d^{10}) + 4*\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*\arctan(((\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} - \sqrt{4*\sqrt{c^2*d^6/b^4}*b^2*c^3*d^7*\cos(b*x + a)*\sin(b*x + a) + c^4*d^{10} + 2*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))}*(2*c^2*d^5*\cos(b*x + a)*\sin(b*x + a) + \sqrt{c^2*d^6/b^4}*b^2*c*d^2 - (\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c*d^3*\sin(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*\cos(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))}))/((2*c^4*d^{10}*\cos(b*x + a)^2 - c^4*d^{10}) + 4*\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*\arctan(-1/2*(2*c^4*d^{10}*\cos(b*x + a)*\sin(b*x + a) - \sqrt{4*\sqrt{c^2*d^6/b^4}*b^2*c^3*d^7*\cos(b*x + a)*\sin(b*x + a) + c^4*d^{10} + 2*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))}*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c*d^3*\sin(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*\cos(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} + (\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\sin(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\cos(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} - 4*(b^2*c^3*d^7*\cos(b*x + a)^4 - b^2*c^3*d^7*\cos(b*x + a)^2)*\sqrt{c^2*d^6/b^4}))/((2*c^4*d^{10}*\cos(b*x + a)^3 - c^4*d^{10}*\cos(b*x + a))*\sin(b*x + a)) + 4*\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*\arctan(1/2*(2*c^4*d^{10}*\cos(b*x + a)*\sin(b*x + a) + \sqrt{4*\sqrt{c^2*d^6/b^4}*b^2*c^3*d^7*\cos(b*x + a)*\sin(b*x + a) + c^4*d^{10} - 2*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))}*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c*d^3*\sin(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*\cos(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} - (\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\sin(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\cos(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} - 4*(b^2*c^3*d^7*\cos(b*x + a)^4 - b^2*c^3*d^7*\cos(b*x + a)^2)*\sqrt{c^2*d^6/b^4}))/((2*c^4*d^{10}*\cos(b*x + a)^3 - c^4*d^{10}*\cos(b*x + a))*\sin(b*x + a)) - \sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*\log(4*\sqrt{c^2*d^6/b^4}*b^2*c^3*d^7*\cos(b*x + a)*\sin(b*x + a) + c^4*d^{10} + 2*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} + \sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*\log(4*\sqrt{c^2*d^6/b^4}*b^2*c^3*d^7*\cos(b*x + a)*\sin(b*x + a) + c^4*d^{10} - 2*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))} - \sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*\log(1/4*\sqrt{c^2*d^6/b^4}*b^2*c^3*d^7*\cos(b*x + a)*\sin(b*x + a) + 1/16*c^4*d^{10} + 1/8*(\sqrt{2}*(c^2*d^6/b^4)^{(1/4)}*b*c^3*d^8*\cos(b*x + a) + \sqrt{2}*(c^2*d^6/b^4)^{(3/4)}*b^3*c^2*d^5*\sin(b*x + a))*\sqrt{(d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a))}
\end{aligned}$$

$(d \cos(bx + a)) \sqrt{c \sin(bx + a)} + \sqrt{2} (c^2 d^6 / b^4)^{1/4} b \log(1/4 \sqrt{c^2 d^6 / b^4} b^2 c^3 d^7 \cos(bx + a) \sin(bx + a) + 1/16 c^4 d^{10} - 1/8 (\sqrt{2} (c^2 d^6 / b^4)^{1/4} b c^3 d^8 \cos(bx + a) + \sqrt{2} (c^2 d^6 / b^4)^{3/4} b^3 c^2 d^5 \sin(bx + a)) \sqrt{d \cos(bx + a)} \sqrt{c \sin(bx + a)}) / b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sin(a + bx)} (d \cos(a + bx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(3/2)*(c*sin(b*x+a))**(1/2), x)

[Out] Integral(sqrt(c*sin(a + b*x))*(d*cos(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(1/2), x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^(1/2), x)

[Out] int((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^(1/2), x)

$$3.263 \quad \int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx$$

Optimal. Leaf size=280

$$\frac{\sqrt{c} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} \right)}{\sqrt{2} b \sqrt{d}} + \frac{\sqrt{c} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} \right)}{\sqrt{2} b \sqrt{d}} + \sqrt{c} \log \left(\sqrt{c} - \right)$$

[Out] $-1/2*\arctan(1-2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/c^{(1/2)}/(d*\cos(b*x+a))^{(1/2)})*c^{(1/2)}/b*2^{(1/2)}/d^{(1/2)}+1/2*\arctan(1+2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/c^{(1/2)}/(d*\cos(b*x+a))^{(1/2)})*c^{(1/2)}/b*2^{(1/2)}/d^{(1/2)}+1/4*\ln(c^{(1/2)}-2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(1/2)}+c^{(1/2)}*\tan(b*x+a))*c^{(1/2)}/b*2^{(1/2)}/d^{(1/2)}-1/4*\ln(c^{(1/2)}+2^{(1/2)}*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(1/2)}+c^{(1/2)}*\tan(b*x+a))*c^{(1/2)}/b*2^{(1/2)}/d^{(1/2)}$

Rubi [A]

time = 0.13, antiderivative size = 280, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{c} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} \right)}{\sqrt{2} b \sqrt{d}} + \frac{\sqrt{c} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} + 1 \right)}{\sqrt{2} b \sqrt{d}} + \frac{\sqrt{c} \log \left(-\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx) + \sqrt{c} \right)}{2\sqrt{2} b \sqrt{d}} - \frac{\sqrt{c} \log \left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx) + \sqrt{c} \right)}{2\sqrt{2} b \sqrt{d}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sin[a + b*x]]/Sqrt[d*Cos[a + b*x]],x]

[Out] $-\left(\frac{\sqrt{c} \operatorname{ArcTan} \left[1 - \left(\sqrt{2} \sqrt{d} \sqrt{c \sin[a + b*x]} \right) / \left(\sqrt{c} \sqrt{d \cos[a + b*x]} \right) \right]}{\sqrt{2} b \sqrt{d}} \right) + \left(\frac{\sqrt{c} \operatorname{ArcTan} \left[1 + \left(\sqrt{2} \sqrt{d} \sqrt{c \sin[a + b*x]} \right) / \left(\sqrt{c} \sqrt{d \cos[a + b*x]} \right) \right]}{\sqrt{2} b \sqrt{d}} \right) + \left(\frac{\sqrt{c} \operatorname{Log} \left[\sqrt{c} - \left(\sqrt{2} \sqrt{d} \sqrt{c \sin[a + b*x]} \right) / \sqrt{d \cos[a + b*x]} \right]}{2 \sqrt{2} b \sqrt{d}} \right) - \left(\frac{\sqrt{c} \operatorname{Log} \left[\sqrt{c} + \left(\sqrt{2} \sqrt{d} \sqrt{c \sin[a + b*x]} \right) / \sqrt{d \cos[a + b*x]} \right]}{2 \sqrt{2} b \sqrt{d}} \right)$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a,

b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx &= \frac{(2cd) \operatorname{Subst}\left(\int \frac{x^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{b} \\
&= -\frac{c \operatorname{Subst}\left(\int \frac{c - dx^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{b} + \frac{c \operatorname{Subst}\left(\int \frac{c + dx^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{b} \\
&= \frac{c \operatorname{Subst}\left(\int \frac{1}{\frac{c}{d} - \frac{\sqrt{2} \sqrt{c}}{\sqrt{d}} x + x^2} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{2bd} + \frac{c \operatorname{Subst}\left(\int \frac{1}{\frac{c}{d} + \frac{\sqrt{2} \sqrt{c}}{\sqrt{d}} x + x^2} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{2bd} \\
&= \frac{\sqrt{c} \log\left(\sqrt{c} - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx)\right)}{2\sqrt{2} b \sqrt{d}} - \frac{\sqrt{c} \log\left(\sqrt{c} + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx)\right)}{2\sqrt{2} b \sqrt{d}} \\
&= -\frac{\sqrt{c} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{\sqrt{2} b \sqrt{d}} + \frac{\sqrt{c} \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{\sqrt{2} b \sqrt{d}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 67, normalized size = 0.24

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)} \tan(a + bx)}{3b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c*Sin[a + b*x]]/Sqrt[d*Cos[a + b*x]], x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, 3/4, 7/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x])/(3*b*Sqrt[d*Cos[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.10, size = 271, normalized size = 0.97

method	result
default	$ -\frac{\sqrt{c \sin(bx + a)} \left(i \operatorname{EllipticPi}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{1}{2} - \frac{i}{2}, \frac{\sqrt{2}}{2}\right) - i \operatorname{EllipticPi}\left(\sqrt{\frac{1 - \cos(bx + a) + \sin(bx + a)}{\sin(bx + a)}}, \frac{1}{2} + \frac{i}{2}, \frac{\sqrt{2}}{2}\right) \right)}{3b \sqrt{d \cos(bx + a)}} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2/b*(c*sin(b*x+a))^(1/2)*(I*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-I*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))-EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2)))*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)/(-1+cos(b*x+a))/(d*cos(b*x+a))^(1/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(c*sin(b*x + a))/sqrt(d*cos(b*x + a)), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2003 vs. 2(209) = 418.

time = 29.97, size = 2003, normalized size = 7.15

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/4*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*arctan(((sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) + sqrt(4*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*cos(b*x + a)*sin(b*x + a) + c^4 - 2*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)))/(2*c^4*cos(b*x + a)^2 - c^4)) + 1/4*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*arctan(((sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) - sqrt(4*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*cos(b*x + a)*sin
```

```
(b*x + a) + c^4 + 2*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a) +
sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt
t(c*sin(b*x + a)))*(b^2*c*d*sqrt(c^2/(b^4*d^2)) + 2*c^2*cos(b*x + a)*sin(b*
x + a) - (sqrt(2)*b^3*d*(c^2/(b^4*d^2))^(3/4)*cos(b*x + a) + sqrt(2)*b*c*(c
^2/(b^4*d^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)
))/(2*c^4*cos(b*x + a)^2 - c^4) + 1/4*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*arctan
(-1/2*(2*c^4*cos(b*x + a)*sin(b*x + a) - sqrt(4*b^2*c^3*d*sqrt(c^2/(b^4*d^
2))*cos(b*x + a)*sin(b*x + a) + c^4 + 2*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(
3/4)*sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(
d*cos(b*x + a))*sqrt(c*sin(b*x + a)))*(sqrt(2)*b^3*d*(c^2/(b^4*d^2))^(3/4)*
cos(b*x + a) + sqrt(2)*b*c*(c^2/(b^4*d^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b
*x + a))*sqrt(c*sin(b*x + a)) + (sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*co
s(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b
*x + a))*sqrt(c*sin(b*x + a)) - 4*(b^2*c^3*d*cos(b*x + a)^4 - b^2*c^3*d*cos
(b*x + a)^2)*sqrt(c^2/(b^4*d^2)))/((2*c^4*cos(b*x + a)^3 - c^4*cos(b*x + a)
)*sin(b*x + a)) + 1/4*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*arctan(1/2*(2*c^4*cos(
b*x + a)*sin(b*x + a) + sqrt(4*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*cos(b*x + a)*s
in(b*x + a) + c^4 - 2*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a)
+ sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*s
qrt(c*sin(b*x + a)))*(sqrt(2)*b^3*d*(c^2/(b^4*d^2))^(3/4)*cos(b*x + a) + sq
rt(2)*b*c*(c^2/(b^4*d^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*s
in(b*x + a)) - (sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*cos(b*x + a) + sqrt
(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*s
in(b*x + a)) - 4*(b^2*c^3*d*cos(b*x + a)^4 - b^2*c^3*d*cos(b*x + a)^2)*sqrt
(c^2/(b^4*d^2)))/((2*c^4*cos(b*x + a)^3 - c^4*cos(b*x + a))*sin(b*x + a))
- 1/16*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*log(4*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*co
s(b*x + a)*sin(b*x + a) + c^4 + 2*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*
sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos
(b*x + a))*sqrt(c*sin(b*x + a))) + 1/16*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*log(4
*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*cos(b*x + a)*sin(b*x + a) + c^4 - 2*(sqrt(2)
*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^
2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))) - 1/16*sq
rt(2)*(c^2/(b^4*d^2))^(1/4)*log(1/4*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*cos(b*x +
a)*sin(b*x + a) + 1/16*c^4 + 1/8*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*
sin(b*x + a) + sqrt(2)*b*c^3*(c^2/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos
(b*x + a))*sqrt(c*sin(b*x + a))) + 1/16*sqrt(2)*(c^2/(b^4*d^2))^(1/4)*log(1
/4*b^2*c^3*d*sqrt(c^2/(b^4*d^2))*cos(b*x + a)*sin(b*x + a) + 1/16*c^4 - 1/8
*(sqrt(2)*b^3*c^2*d*(c^2/(b^4*d^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^3*(c^2
/(b^4*d^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))**(1/2)/(d*cos(b*x+a))**(1/2),x)`

[Out] `Integral(sqrt(c*sin(a + b*x))/sqrt(d*cos(a + b*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(c*sin(b*x + a))/sqrt(d*cos(b*x + a)), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{c \sin(a + b x)}}{\sqrt{d \cos(a + b x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(1/2),x)`

[Out] `int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(1/2), x)`

$$3.264 \quad \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{5/2}} dx$$

Optimal. Leaf size=37

$$\frac{2(c \sin(a + bx))^{3/2}}{3bcd(d \cos(a + bx))^{3/2}}$$

[Out] $2/3*(c*\sin(b*x+a))^{(3/2)}/b/c/d/(d*\cos(b*x+a))^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2643}

$$\frac{2(c \sin(a + bx))^{3/2}}{3bcd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(5/2), x]`

[Out] $(2*(c*\sin[a + b*x])^{(3/2)})/(3*b*c*d*(d*\cos[a + b*x])^{(3/2)})$

Rule 2643

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] & & NeQ[m, -1]`

Rubi steps

$$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{5/2}} dx = \frac{2(c \sin(a + bx))^{3/2}}{3bcd(d \cos(a + bx))^{3/2}}$$

Mathematica [A]

time = 0.06, size = 37, normalized size = 1.00

$$\frac{2(c \sin(a + bx))^{3/2}}{3bcd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Integrate[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(5/2), x]`

[Out] $(2*(c*\sin[a + b*x])^{(3/2)})/(3*b*c*d*(d*\cos[a + b*x])^{(3/2)})$

Maple [A]

time = 0.12, size = 38, normalized size = 1.03

method	result	size
default	$\frac{2 \sin(bx+a) \cos(bx+a) \sqrt{c \sin(bx+a)}}{3b(d \cos(bx+a))^{\frac{5}{2}}}$	38

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)`

[Out] $2/3/b*\sin(b*x+a)*\cos(b*x+a)*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(5/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(5/2), x)`

Fricas [A]

time = 0.36, size = 42, normalized size = 1.14

$$\frac{2 \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)} \sin(bx+a)}{3bd^3 \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")`

[Out] $2/3*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}*\sin(b*x + a)/(b*d^3*\cos(b*x + a)^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))**(1/2)/(d*cos(b*x+a))**(5/2),x)`

[Out] Integral(sqrt(c*sin(a + b*x))/(d*cos(a + b*x))**(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(5/2), x)

Mupad [B]

time = 0.99, size = 50, normalized size = 1.35

$$\frac{2 \sin(2a + 2bx) \sqrt{c \sin(a + bx)}}{3bd^2 (\cos(2a + 2bx) + 1) \sqrt{d \cos(a + bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(5/2),x)

[Out] (2*sin(2*a + 2*b*x)*(c*sin(a + b*x))^(1/2))/(3*b*d^2*(cos(2*a + 2*b*x) + 1)
*(d*cos(a + b*x))^(1/2))

$$3.265 \quad \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{9/2}} dx$$

Optimal. Leaf size=75

$$\frac{2(c \sin(a + bx))^{3/2}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{8(c \sin(a + bx))^{3/2}}{21bcd^3(d \cos(a + bx))^{3/2}}$$

[Out] $2/7*(c*\sin(b*x+a))^(3/2)/b/c/d/(d*\cos(b*x+a))^(7/2)+8/21*(c*\sin(b*x+a))^(3/2)/b/c/d^3/(d*\cos(b*x+a))^(3/2)$

Rubi [A]

time = 0.07, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2651, 2643}

$$\frac{8(c \sin(a + bx))^{3/2}}{21bcd^3(d \cos(a + bx))^{3/2}} + \frac{2(c \sin(a + bx))^{3/2}}{7bcd(d \cos(a + bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(9/2), x]

[Out] $(2*(c*\sin[a + b*x])^(3/2))/(7*b*c*d*(d*\cos[a + b*x])^(7/2)) + (8*(c*\sin[a + b*x])^(3/2))/(21*b*c*d^3*(d*\cos[a + b*x])^(3/2))$

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(-b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^(n)*((a*Cos[e + f*x])^(m + 2), x), x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{9/2}} dx = \frac{2(c \sin(a + bx))^{3/2}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{4 \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{5/2}} dx}{7d^2}$$

$$= \frac{2(c \sin(a + bx))^{3/2}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{8(c \sin(a + bx))^{3/2}}{21bcd^3(d \cos(a + bx))^{3/2}}$$

Mathematica [A]

time = 0.17, size = 57, normalized size = 0.76

$$\frac{2\sqrt{d \cos(a + bx)} (5 + 2 \cos(2(a + bx))) \sec^4(a + bx) (c \sin(a + bx))^{3/2}}{21bcd^5}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(9/2), x]
```

```
[Out] (2*Sqrt[d*Cos[a + b*x]]*(5 + 2*Cos[2*(a + b*x)])*Sec[a + b*x]^4*(c*Sin[a + b*x])^(3/2))/(21*b*c*d^5)
```

Maple [A]

time = 0.13, size = 50, normalized size = 0.67

method	result	size
default	$\frac{2(4(\cos^2(bx+a))+3) \cos(bx+a) \sqrt{c \sin(bx+a)} \sin(bx+a)}{21b(d \cos(bx+a))^{\frac{9}{2}}}$	50

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(9/2), x, method=_RETURNVERBOSE)
```

```
[Out] 2/21/b*(4*cos(b*x+a)^2+3)*cos(b*x+a)*(c*sin(b*x+a))^(1/2)*sin(b*x+a)/(d*cos(b*x+a))^(9/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(9/2), x, algorithm="maxima")
```

```
[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(9/2), x)
```

Fricas [A]

time = 0.39, size = 54, normalized size = 0.72

$$\frac{2 \sqrt{d \cos(bx + a)} (4 \cos(bx + a)^2 + 3) \sqrt{c \sin(bx + a)} \sin(bx + a)}{21 b d^5 \cos(bx + a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")

[Out] 2/21*sqrt(d*cos(b*x + a))*(4*cos(b*x + a)^2 + 3)*sqrt(c*sin(b*x + a))*sin(b*x + a)/(b*d^5*cos(b*x + a)^4)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(1/2)/(d*cos(b*x+a))**(9/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(9/2), x)

Mupad [B]

time = 2.06, size = 95, normalized size = 1.27

$$\frac{8 \sqrt{c \sin(a + bx)} (11 \sin(2a + 2bx) + 7 \sin(4a + 4bx) + \sin(6a + 6bx))}{21 b d^4 \sqrt{d \cos(a + bx)} (15 \cos(2a + 2bx) + 6 \cos(4a + 4bx) + \cos(6a + 6bx) + 10)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(9/2),x)

[Out] (8*(c*sin(a + b*x))^(1/2)*(11*sin(2*a + 2*b*x) + 7*sin(4*a + 4*b*x) + sin(6*a + 6*b*x)))/(21*b*d^4*(d*cos(a + b*x))^(1/2)*(15*cos(2*a + 2*b*x) + 6*cos(4*a + 4*b*x) + cos(6*a + 6*b*x) + 10))

$$3.266 \quad \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{13/2}} dx$$

Optimal. Leaf size=112

$$\frac{2(c \sin(a + bx))^{3/2}}{11bcd(d \cos(a + bx))^{11/2}} + \frac{16(c \sin(a + bx))^{3/2}}{77bcd^3(d \cos(a + bx))^{7/2}} + \frac{64(c \sin(a + bx))^{3/2}}{231bcd^5(d \cos(a + bx))^{3/2}}$$

[Out] 2/11*(c*sin(b*x+a))^(3/2)/b/c/d/(d*cos(b*x+a))^(11/2)+16/77*(c*sin(b*x+a))^(3/2)/b/c/d^3/(d*cos(b*x+a))^(7/2)+64/231*(c*sin(b*x+a))^(3/2)/b/c/d^5/(d*cos(b*x+a))^(3/2)

Rubi [A]

time = 0.11, antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2651, 2643}

$$\frac{64(c \sin(a + bx))^{3/2}}{231bcd^5(d \cos(a + bx))^{3/2}} + \frac{16(c \sin(a + bx))^{3/2}}{77bcd^3(d \cos(a + bx))^{7/2}} + \frac{2(c \sin(a + bx))^{3/2}}{11bcd(d \cos(a + bx))^{11/2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(13/2), x]

[Out] (2*(c*Sin[a + b*x])^(3/2))/(11*b*c*d*(d*Cos[a + b*x])^(11/2)) + (16*(c*Sin[a + b*x])^(3/2))/(77*b*c*d^3*(d*Cos[a + b*x])^(7/2)) + (64*(c*Sin[a + b*x])^(3/2))/(231*b*c*d^5*(d*Cos[a + b*x])^(3/2))

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{c \sin(a+bx)}}{(d \cos(a+bx))^{13/2}} dx &= \frac{2(c \sin(a+bx))^{3/2}}{11bcd(d \cos(a+bx))^{11/2}} + \frac{8 \int \frac{\sqrt{c \sin(a+bx)}}{(d \cos(a+bx))^{9/2}} dx}{11d^2} \\
&= \frac{2(c \sin(a+bx))^{3/2}}{11bcd(d \cos(a+bx))^{11/2}} + \frac{16(c \sin(a+bx))^{3/2}}{77bcd^3(d \cos(a+bx))^{7/2}} + \frac{32 \int \frac{\sqrt{c \sin(a+bx)}}{(d \cos(a+bx))^{5/2}} dx}{77d^4} \\
&= \frac{2(c \sin(a+bx))^{3/2}}{11bcd(d \cos(a+bx))^{11/2}} + \frac{16(c \sin(a+bx))^{3/2}}{77bcd^3(d \cos(a+bx))^{7/2}} + \frac{64(c \sin(a+bx))^{3/2}}{231bcd^5(d \cos(a+bx))^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.16, size = 67, normalized size = 0.60

$$\frac{2\sqrt{d \cos(a+bx)} (45 + 28 \cos(2(a+bx)) + 4 \cos(4(a+bx))) \sec^6(a+bx) (c \sin(a+bx))^{3/2}}{231bcd^7}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[c*Sin[a + b*x]]/(d*Cos[a + b*x])^(13/2), x]``[Out] (2*sqrt[d*cos[a + b*x]]*(45 + 28*cos[2*(a + b*x)] + 4*cos[4*(a + b*x)])*Sec[a + b*x]^6*(c*Sin[a + b*x])^(3/2))/(231*b*c*d^7)`**Maple [A]**

time = 0.13, size = 60, normalized size = 0.54

method	result	size
default	$\frac{2(32(\cos^4(bx+a))+24(\cos^2(bx+a))+21) \cos(bx+a) \sqrt{c \sin(bx+a)} \sin(bx+a)}{231b(d \cos(bx+a))^{\frac{13}{2}}}$	60

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(13/2), x, method=_RETURNVERBOSE)``[Out] 2/231/b*(32*cos(b*x+a)^4+24*cos(b*x+a)^2+21)*cos(b*x+a)*(c*sin(b*x+a))^(1/2)*sin(b*x+a)/(d*cos(b*x+a))^(13/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(13/2), x, algorithm="maxima")`

[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(13/2), x)

Fricas [A]

time = 0.46, size = 64, normalized size = 0.57

$$\frac{2 (32 \cos (bx + a)^4 + 24 \cos (bx + a)^2 + 21) \sqrt{d \cos (bx + a)} \sqrt{c \sin (bx + a)} \sin (bx + a)}{231 b d^7 \cos (bx + a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(13/2),x, algorithm="fricas")

[Out] 2/231*(32*cos(b*x + a)^4 + 24*cos(b*x + a)^2 + 21)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*sin(b*x + a)/(b*d^7*cos(b*x + a)^6)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(1/2)/(d*cos(b*x+a))**(13/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(13/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a))/(d*cos(b*x + a))^(13/2), x)

Mupad [B]

time = 6.21, size = 216, normalized size = 1.93

$$\frac{\sqrt{c \sin (a + b x)} \left(2 \sin \left(\frac{a}{2} + \frac{b x}{2} \right)^2 - 1 \right) \left(2 \sin \left(\frac{5 a}{2} + \frac{5 b x}{2} \right)^2 + \sin (5 a + 5 b x) \right) \left(\frac{1984 \sin (a + b x) \left(-2 \sin \left(\frac{5 a}{2} + \frac{5 b x}{2} \right)^2 + \sin (5 a + 5 b x) \right) \right)}{231 b d^6} + \frac{256 \sin (3 a + 3 b x) \left(-2 \sin \left(\frac{5 a}{2} + \frac{5 b x}{2} \right)^2 + \sin (5 a + 5 b x) \right) \right)}{77 b d^6} + \frac{128 \sin (5 a + 5 b x) \left(-2 \sin \left(\frac{5 a}{2} + \frac{5 b x}{2} \right)^2 + \sin (5 a + 5 b x) \right) \right)}{231 b d^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(1/2)/(d*cos(a + b*x))^(13/2),x)

[Out] -((c*sin(a + b*x))^(1/2)*(2*sin(a/2 + (b*x)/2)^2 - 1)*(sin(5*a + 5*b*x)*1i + 2*sin((5*a)/2 + (5*b*x)/2)^2 - 1)*((1984*sin(a + b*x)*(sin(5*a + 5*b*x)*1i - 2*sin((5*a)/2 + (5*b*x)/2)^2 + 1))/(231*b*d^6) + (256*sin(3*a + 3*b*x)*(sin(5*a + 5*b*x)*1i - 2*sin((5*a)/2 + (5*b*x)/2)^2 + 1))/(77*b*d^6) + (128*sin(5*a + 5*b*x)*(sin(5*a + 5*b*x)*1i - 2*sin((5*a)/2 + (5*b*x)/2)^2 + 1))/(231*b*d^6))/((32*(sin(a + b*x)^2 - 1)^3*(-d*(2*sin(a/2 + (b*x)/2)^2 - 1))^(1/2))

3.267 $\int (d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2} dx$

Optimal. Leaf size=131

$$\frac{cd\sqrt{d\cos(a+bx)}\sqrt{c\sin(a+bx)}}{6b} - \frac{c(d\cos(a+bx))^{5/2}\sqrt{c\sin(a+bx)}}{3bd} + \frac{c^2d^2F(a-\frac{\pi}{4}+bx|2)\sqrt{\sin(2a+2bx)}}{12b\sqrt{d\cos(a+bx)}\sqrt{c\sin(a+bx)}}$$

[Out] $-1/3*c*(d*\cos(b*x+a))^{(5/2)}*(c*\sin(b*x+a))^{(1/2)}/b/d+1/6*c*d*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b-1/12*c^2*d^2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x),2^{(1/2)})*\sin(2*b*x+2*a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.12, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2648, 2649, 2653, 2720}

$$\frac{c^2d^2\sqrt{\sin(2a+2bx)}F(a+bx-\frac{\pi}{4}|2)}{12b\sqrt{c\sin(a+bx)}\sqrt{d\cos(a+bx)}} - \frac{c\sqrt{c\sin(a+bx)}(d\cos(a+bx))^{5/2}}{3bd} + \frac{cd\sqrt{c\sin(a+bx)}\sqrt{d\cos(a+bx)}}{6b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^{(3/2)}*(c*\text{Sin}[a + b*x])^{(3/2)}, x]$

[Out] $(c*d*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(6*b) - (c*(d*\text{Cos}[a + b*x])^{(5/2)}*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(3*b*d) + (c^2*d^2*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(12*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.))^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{(n)}*(a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2649

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.))^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_)}], x_Symbol] \rightarrow \text{Simp}[a*(b*\text{Sin}[e + f*x])^{(n + 1)}*((a*\text{Cos}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Sin}[e + f*x])^{(n)}*(a*\text{Cos}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2653

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_)]]), x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b$

*Cos[e + f*x]], Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2} dx &= -\frac{c(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bd} + \frac{1}{6}c^2 \int \frac{(d \cos(a + bx))^{3/2}}{\sqrt{c \sin(a + bx)}} dx \\ &= \frac{cd \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{6b} - \frac{c(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bd} \\ &= \frac{cd \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{6b} - \frac{c(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bd} \\ &= \frac{cd \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{6b} - \frac{c(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bd} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.09, size = 71, normalized size = 0.54

$$\frac{2cd \sqrt{d \cos(a + bx)} \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, \frac{5}{4}; \frac{9}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)} \tan^2(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(3/2)*(c*Sin[a + b*x])^(3/2), x]

[Out] (2*c*d*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, 5/4, 9/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]]*Tan[a + b*x]^2)/(5*b)

Maple [A]

time = 0.14, size = 216, normalized size = 1.65

method	result
default	$-\frac{\left(2(\cos^4(bx+a))\sqrt{2} + \sin(bx+a)\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}\right) \text{EllipticF}\left(\frac{1}{2}\left(\frac{c-\sin(bx+a)}{d}\right), \frac{\pi}{2}\right)}{12b \sin(bx+a)(-1)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/12/b*(2*cos(b*x+a)^4*2^(1/2)+sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-2*cos(b*x+a)^3*2^(1/2)-cos(b*x+a)^2*2^(1/2)+cos(b*x+a)*2^(1/2))
*(d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2)/sin(b*x+a)/(-1+cos(b*x+a))/cos(b*x+a)^2*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((d*cos(b*x + a))^(3/2)*(c*sin(b*x + a))^(3/2), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*c*d*cos(b*x + a)*sin(b*x + a), x)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))**(3/2)*(c*sin(b*x+a))**(3/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 4370 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2),x, algorithm="giac")
```

```
[Out] integrate((d*cos(b*x + a))^(3/2)*(c*sin(b*x + a))^(3/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + b x))^{3/2} (c \sin(a + b x))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^(3/2),x)
```

```
[Out] int((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^(3/2), x)
```

$$3.268 \quad \int \frac{(c \sin(a+bx))^{3/2}}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=93

$$-\frac{c\sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}{bd} + \frac{c^2 F(a - \frac{\pi}{4} + bx | 2) \sqrt{\sin(2a + 2bx)}}{2b\sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] $-c*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b/d-1/2*c^2*(\sin(a+1/4*\text{Pi}+b*x))^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticF}(\cos(a+1/4*\text{Pi}+b*x), 2^{(1/2)})*\sin(2*b*x+2*a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2648, 2653, 2720}

$$\frac{c^2 \sqrt{\sin(2a + 2bx)} F(a + bx - \frac{\pi}{4} | 2)}{2b\sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}} - \frac{c\sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{bd}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(3/2)}/\text{Sqrt}[d*\text{Cos}[a + b*x]], x]$

[Out] $-((c*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(b*d)) + (c^2*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(2*b*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2648

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(b*\text{Cos}[e + f*x])^{(n + 1)}*((a*\text{Sin}[e + f*x])^{(m - 1)})/(b*f*(m + n)), x] + \text{Dist}[a^2*((m - 1)/(m + n)), \text{Int}[(b*\text{Cos}[e + f*x])^{n*} (a*\text{Sin}[e + f*x])^{(m - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2653

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{3/2}}{\sqrt{d \cos(a + bx)}} dx &= -\frac{c \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{bd} + \frac{1}{2} c^2 \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx \\ &= -\frac{c \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{bd} + \frac{\left(c^2 \sqrt{\sin(2a + 2bx)} \right) \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx}{2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \\ &= -\frac{c \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{bd} + \frac{c^2 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{2b \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.06, size = 67, normalized size = 0.72

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{5}{4}; \frac{9}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{3/2} \tan(a + bx)}{5b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(3/2)/Sqrt[d*Cos[a + b*x]], x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, 5/4, 9/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(3/2)*Tan[a + b*x])/(5*b*Sqrt[d*Cos[a + b*x]])

Maple [A]

time = 0.13, size = 182, normalized size = 1.96

method	result
default	$-\frac{\left(\sin(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)}{2b(-1+\cos(bx+a)) \sqrt{d \cos(bx+a)} \sin(bx+a)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(1/2), x, method=_RETURNVERBOSE)

[Out] -1/2/b*(sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))+cos(b*x+a)^2*2^(1/2)-cos(b*x+a)*2^(1/2))*(c*sin(b*x+a))^(3/2)/(-1+cos(b*x+a))/(d*cos(b*x+a))^(1/2)/sin(b*x+a)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(3/2)/sqrt(d*cos(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*c*sin(b*x + a)/(d*cos(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(a + bx))^{\frac{3}{2}}}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(3/2)/(d*cos(b*x+a))**(1/2),x)

[Out] Integral((c*sin(a + b*x))**(3/2)/sqrt(d*cos(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)/sqrt(d*cos(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^{3/2}}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(1/2),x)

[Out] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(1/2), x)

$$3.269 \quad \int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=98

$$\frac{2c\sqrt{c \sin(a+bx)}}{3bd(d \cos(a+bx))^{3/2}} - \frac{c^2 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a+2bx)}}{3bd^2 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] 2/3*c*(c*sin(b*x+a))^(1/2)/b/d/(d*cos(b*x+a))^(3/2)+1/3*c^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*sin(2*b*x+2*a)^(1/2)/b/d^2/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2)

Rubi [A]

time = 0.08, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2646, 2653, 2720}

$$\frac{2c\sqrt{c \sin(a+bx)}}{3bd(d \cos(a+bx))^{3/2}} - \frac{c^2 \sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \mid 2\right)}{3bd^2 \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(5/2),x]

[Out] (2*c*Sqrt[c*Sin[a + b*x]])/(3*b*d*(d*Cos[a + b*x])^(3/2)) - (c^2*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[Sin[2*a + 2*b*x]])/(3*b*d^2*Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]])

Rule 2646

Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_)), x_Symbol] := Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{5/2}} dx &= \frac{2c \sqrt{c \sin(a + bx)}}{3bd(d \cos(a + bx))^{3/2}} - \frac{c^2 \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx}{3d^2} \\
&= \frac{2c \sqrt{c \sin(a + bx)}}{3bd(d \cos(a + bx))^{3/2}} - \frac{\left(c^2 \sqrt{\sin(2a + 2bx)}\right) \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx}{3d^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \\
&= \frac{2c \sqrt{c \sin(a + bx)}}{3bd(d \cos(a + bx))^{3/2}} - \frac{c^2 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{3bd^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.11, size = 67, normalized size = 0.68

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{5}{4}, \frac{7}{4}; \frac{9}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{5/2}}{5bcd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(5/2), x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[5/4, 7/4, 9/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(5/2))/(5*b*c*d*(d*Cos[a + b*x])^(3/2))

Maple [A]

time = 0.12, size = 186, normalized size = 1.90

method	result
default	$ \frac{\left(\sin(bx+a) \cos(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)}{\sin(bx+a)}}\right)\right)}{3b(-1+\cos(bx+a))(d \cos(bx+a))^{\frac{5}{2}} \sin(bx+a)} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(5/2), x, method=_RETURNVERBOSE)

[Out] 1/3/b*(sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))+cos(b*x+a)*2^(1/2)-2^(1/2))*cos(b*x+a)*(c*sin(b*x+a))^(3/2)/(-1+cos(b*x+a))/(d*cos(b*x+a))^(5/2)/sin(b*x+a)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(5/2), x)

Fricas [C] Result contains complex when optimal does not.

time = 0.10, size = 106, normalized size = 1.08

$$\frac{\sqrt{icd} c \cos(bx+a)^2 \operatorname{ellipticF}(\cos(bx+a) + i \sin(bx+a), -1) + \sqrt{-icd} c \cos(bx+a)^2 \operatorname{ellipticF}(\cos(bx+a) - i \sin(bx+a), -1) + 2 \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)} c}{3bd^3 \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")

[Out] 1/3*(sqrt(I*c*d)*c*cos(b*x + a)^2*ellipticF(cos(b*x + a) + I*sin(b*x + a), -1) + sqrt(-I*c*d)*c*cos(b*x + a)^2*ellipticF(cos(b*x + a) - I*sin(b*x + a), -1) + 2*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*c)/(b*d^3*cos(b*x + a)^2)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(3/2)/(d*cos(b*x+a))**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3006 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(5/2), x)
```

```
[Out] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(5/2), x)
```

$$3.270 \quad \int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=133

$$\frac{2c\sqrt{c \sin(a+bx)}}{7bd(d \cos(a+bx))^{7/2}} - \frac{2c\sqrt{c \sin(a+bx)}}{21bd^3(d \cos(a+bx))^{3/2}} - \frac{2c^2 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a+2bx)}}{21bd^4 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] $2/7*c*(c*\sin(b*x+a))^{1/2}/b/d/(d*\cos(b*x+a))^{7/2}-2/21*c*(c*\sin(b*x+a))^{1/2}/b/d^3/(d*\cos(b*x+a))^{3/2}+2/21*c^2*(\sin(a+1/4*\pi+b*x)^2)^{1/2}/\sin(a+1/4*\pi+b*x)*\text{EllipticF}(\cos(a+1/4*\pi+b*x),2^{1/2})*\sin(2*b*x+2*a)^{1/2}/b/d^4/(d*\cos(b*x+a))^{1/2}/(c*\sin(b*x+a))^{1/2}$

Rubi [A]

time = 0.11, antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2646, 2651, 2653, 2720}

$$-\frac{2c^2 \sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \mid 2\right)}{21bd^4 \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}} - \frac{2c\sqrt{c \sin(a+bx)}}{21bd^3(d \cos(a+bx))^{3/2}} + \frac{2c\sqrt{c \sin(a+bx)}}{7bd(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] `Int[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(9/2),x]`

[Out] $(2*c*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(7*b*d*(d*\text{Cos}[a + b*x])^{7/2}) - (2*c*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(21*b*d^3*(d*\text{Cos}[a + b*x])^{3/2}) - (2*c^2*\text{EllipticF}[a - \pi/4 + b*x, 2]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(21*b*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2646

`Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])`

Rule 2651

`Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(-b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rule 2653

```
Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{9/2}} dx &= \frac{2c \sqrt{c \sin(a + bx)}}{7bd(d \cos(a + bx))^{7/2}} - \frac{c^2 \int \frac{1}{(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}} dx}{7d^2} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{7bd(d \cos(a + bx))^{7/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{21bd^3(d \cos(a + bx))^{3/2}} - \frac{(2c^2) \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx}{21d^4} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{7bd(d \cos(a + bx))^{7/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{21bd^3(d \cos(a + bx))^{3/2}} - \frac{(2c^2 \sqrt{\sin(2a + 2bx)}) \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx}{21d^4 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{7bd(d \cos(a + bx))^{7/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{21bd^3(d \cos(a + bx))^{3/2}} - \frac{2c^2 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{21bd^4 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.11, size = 70, normalized size = 0.53

$$\frac{2 \cos^2(a + bx)^{7/4} \cot(a + bx) {}_2F_1\left(\frac{5}{4}, \frac{11}{4}; \frac{9}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{7/2}}{5bc^2(d \cos(a + bx))^{9/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(9/2), x]
```

```
[Out] (2*(Cos[a + b*x]^2)^(7/4)*Cot[a + b*x]*Hypergeometric2F1[5/4, 11/4, 9/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(7/2))/(5*b*c^2*(d*Cos[a + b*x])^(9/2))
```

Maple [A]

time = 0.13, size = 215, normalized size = 1.62

method	result
--------	--------

default	$-\frac{\left(-2 \sin(bx+a) \cos^3(bx+a)\right) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)}{21b(-1+\cos(bx+a))}$
---------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(9/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/21/b*(-2*\sin(b*x+a)*\cos(b*x+a)^3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\operatorname{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{(1/2))+\cos(b*x+a)^3*2^{(1/2)}-\cos(b*x+a)^2*2^{(1/2)}-3*\cos(b*x+a)*2^{(1/2)}+3*2^{(1/2)})*\cos(b*x+a)*(c*\sin(b*x+a))^{3/2}/(-1+\cos(b*x+a))/(d*\cos(b*x+a))^{9/2}/\sin(b*x+a)*2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(9/2),x, algorithm="maxima")`

[Out] `integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(9/2), x)`

Fricas [C] Result contains complex when optimal does not.

time = 0.11, size = 119, normalized size = 0.89

$$\frac{2\left(\sqrt{icd}c\cos(bx+a)^4\operatorname{ellipticF}(\cos(bx+a)+i\sin(bx+a),-1)+\sqrt{-icd}c\cos(bx+a)^4\operatorname{ellipticF}(\cos(bx+a)-i\sin(bx+a),-1)-(c\cos(bx+a)^2-3c)\sqrt{d\cos(bx+a)}\sqrt{c\sin(bx+a)}\right)}{21bd^5\cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")`

[Out]
$$\frac{2/21*(\sqrt{I*c*d}*c*\cos(b*x + a)^4*\operatorname{ellipticF}(\cos(b*x + a) + I*\sin(b*x + a), -1) + \sqrt{-I*c*d}*c*\cos(b*x + a)^4*\operatorname{ellipticF}(\cos(b*x + a) - I*\sin(b*x + a), -1) - (c*\cos(b*x + a)^2 - 3*c)*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)})}{(b*d^5*\cos(b*x + a)^4)}$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))**(3/2)/(d*cos(b*x+a))**(9/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(9/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(9/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + b x))^{3/2}}{(d \cos(a + b x))^{9/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(9/2),x)

[Out] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(9/2), x)

3.271 $\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2} dx$

Optimal. Leaf size=320

$$\frac{c^{3/2} \sqrt{d} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)}{4\sqrt{2} b} - \frac{c^{3/2} \sqrt{d} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)}{4\sqrt{2} b} - \frac{c^{3/2} \sqrt{d}}{2bd}$$

[Out] $-1/8*c^{(3/2)*\arctan(-1+2^{(1/2)*c^{(1/2)*(d*\cos(b*x+a))^{(1/2)/d^{(1/2)/(c*\sin(b*x+a))^{(1/2)}}*d^{(1/2)/b*2^{(1/2)}-1/8*c^{(3/2)*\arctan(1+2^{(1/2)*c^{(1/2)*(d*\cos(b*x+a))^{(1/2)/d^{(1/2)/(c*\sin(b*x+a))^{(1/2)}}*d^{(1/2)/b*2^{(1/2)}-1/16*c^{(3/2)*\ln(d^{(1/2)+\cot(b*x+a)*d^{(1/2)-2^{(1/2)*c^{(1/2)*(d*\cos(b*x+a))^{(1/2)/(c*\sin(b*x+a))^{(1/2)}*d^{(1/2)/b*2^{(1/2)+1/16*c^{(3/2)*\ln(d^{(1/2)+\cot(b*x+a)*d^{(1/2)+2^{(1/2)*c^{(1/2)*(d*\cos(b*x+a))^{(1/2)/(c*\sin(b*x+a))^{(1/2)}*d^{(1/2)/b*2^{(1/2)-1/2*c*(d*\cos(b*x+a))^{(3/2)*(c*\sin(b*x+a))^{(1/2)/b/d}}$

Rubi [A]

time = 0.18, antiderivative size = 320, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2648, 2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{c^{3/2} \sqrt{d} \operatorname{ArcTan} \left(\frac{1 - \sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)}{4\sqrt{2} b} - \frac{c^{3/2} \sqrt{d} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} + 1 \right)}{4\sqrt{2} b} - \frac{c^{3/2} \sqrt{d} \log \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} + \sqrt{d} \cot(a + bx) + \sqrt{d} \right)}{8\sqrt{2} b} + \frac{c^{3/2} \sqrt{d} \log \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} + \sqrt{d} \cot(a + bx) + \sqrt{d} \right)}{8\sqrt{2} b} - \frac{c \sqrt{c \sin(a + bx)} (d \cos(a + bx))^{3/2}}{2bd}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^(3/2),x]

[Out] $(c^{(3/2)*\text{Sqrt}[d]*\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])]/(\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])})/(4*\text{Sqrt}[2]*b) - (c^{(3/2)*\text{Sqrt}[d]*\text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])]/(\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])})/(4*\text{Sqrt}[2]*b) - (c^{(3/2)*\text{Sqrt}[d]*\text{Log}[\text{Sqrt}[d] + \text{Sqrt}[d]*\text{Cot}[a + b*x] - (\text{Sqrt}[2]*\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])/\text{Sqrt}[c*\text{Sin}[a + b*x]])})/(8*\text{Sqrt}[2]*b) + (c^{(3/2)*\text{Sqrt}[d]*\text{Log}[\text{Sqrt}[d] + \text{Sqrt}[d]*\text{Cot}[a + b*x] + (\text{Sqrt}[2]*\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])/\text{Sqrt}[c*\text{Sin}[a + b*x]])})/(8*\text{Sqrt}[2]*b) - (c*(d*\text{Cos}[a + b*x])^{(3/2)*\text{Sqrt}[c*\text{Sin}[a + b*x]])}/(2*b*d)$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a,

b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2648

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*SIn[e + f*x])^(m -
1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*
(a*SIn[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1]
&& NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2655

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n
_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^
(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*SIn[e
+ f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0
] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2} dx &= -\frac{c(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}}{2bd} + \frac{1}{4}c^2 \int \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} dx \\
&= -\frac{c(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}}{2bd} - \frac{(c^3 d) \operatorname{Subst}\left(\int \frac{x^2}{d^2 + c^2 x^4} dx, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{2b} \\
&= -\frac{c(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}}{2bd} + \frac{(c^2 d) \operatorname{Subst}\left(\int \frac{d - cx^2}{d^2 + c^2 x^4} dx, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{4b} \\
&= -\frac{c(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}}{2bd} - \frac{(c^{3/2} \sqrt{d}) \operatorname{Subst}\left(\int \frac{\sqrt{d \cos(a + bx)}}{-\frac{d}{c} - x^2} dx, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{4b} \\
&= -\frac{c^{3/2} \sqrt{d} \log\left(\sqrt{d} + \sqrt{d} \cot(a + bx) - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{8\sqrt{2} b} \\
&= \frac{c^{3/2} \sqrt{d} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}}\right)}{4\sqrt{2} b} - \frac{c^{3/2} \sqrt{d} \tan^{-1}\left(\frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{4\sqrt{2} b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.05, size = 67, normalized size = 0.21

$$\frac{2\sqrt{d \cos(a + bx)} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{5}{4}; \frac{9}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{3/2} \tan(a + bx)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^(3/2), x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 5/4, 9/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(3/2)*Tan[a + b*x])/(5*b)

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.10, size = 654, normalized size = 2.04

method	result
--------	--------

default	$\frac{(c \sin(bx+a))^{\frac{3}{2}} \sqrt{d \cos(bx+a)} \left(-i \sin(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \right)}{\dots}$
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/8/b*(c*\sin(b*x+a))^{3/2}*(d*\cos(b*x+a))^{1/2}*(-I*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})+I*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})+\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})+\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-2*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticF(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})+2*\cos(b*x+a)^3*2^{1/2}-2*\cos(b*x+a)^2*2^{1/2})/\sin(b*x+a)/\cos(b*x+a)/(-1+\cos(b*x+a))*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(3/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^(3/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1868 vs. $2(237) = 474$.

time = 17.01, size = 1868, normalized size = 5.84

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(3/2),x, algorithm="fricas")`

```

[Out] -1/32*(16*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*c*cos(b*x + a) + 2*sqrt
(2)*(c^6*d^2/b^4)^(1/4)*b*arctan(1/2*(2*c^10*d^4*cos(b*x + a)*sin(b*x + a)
+ sqrt(4*sqrt(c^6*d^2/b^4)*b^2*c^7*d^3*cos(b*x + a)*sin(b*x + a) + c^10*d^4
+ 2*(sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*sin(b*x + a) + sqrt(2)*(c^6*d^2
/b^4)^(3/4)*b^3*c^5*d^2*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x +
a)))*(sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^3*d*cos(b*x + a) + sqrt(2)*(c^6*d^2/
b^4)^(3/4)*b^3*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) + (s
qrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*cos(b*x + a) + sqrt(2)*(c^6*d^2/b^4)^(
3/4)*b^3*c^5*d^2*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) -
4*(b^2*c^7*d^3*cos(b*x + a)^4 - b^2*c^7*d^3*cos(b*x + a)^2)*sqrt(c^6*d^2/b^
4))/((2*c^10*d^4*cos(b*x + a)^3 - c^10*d^4*cos(b*x + a))*sin(b*x + a))) + 2
*sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*arctan(-1/2*(2*c^10*d^4*cos(b*x + a)*sin(b*x
+ a) - sqrt(4*sqrt(c^6*d^2/b^4)*b^2*c^7*d^3*cos(b*x + a)*sin(b*x + a) + c^
10*d^4 - 2*(sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*sin(b*x + a) + sqrt(2)*(c
^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin
(b*x + a)))*(sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^3*d*cos(b*x + a) + sqrt(2)*(c^
6*d^2/b^4)^(3/4)*b^3*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)
) - (sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*cos(b*x + a) + sqrt(2)*(c^6*d^2/
b^4)^(3/4)*b^3*c^5*d^2*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x +
a)) - 4*(b^2*c^7*d^3*cos(b*x + a)^4 - b^2*c^7*d^3*cos(b*x + a)^2)*sqrt(c^6*
d^2/b^4))/((2*c^10*d^4*cos(b*x + a)^3 - c^10*d^4*cos(b*x + a))*sin(b*x + a)
)) + 2*sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*arctan(1/2*((sqrt(2)*(c^6*d^2/b^4)^(1/
4)*b*c^8*d^3*cos(b*x + a) - sqrt(2)*(c^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*sin(b*x
+ a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) + sqrt(4*sqrt(c^6*d^2/b^4)
*b^2*c^7*d^3*cos(b*x + a)*sin(b*x + a) + c^10*d^4 - 2*(sqrt(2)*(c^6*d^2/b^4)
)^(1/4)*b*c^8*d^3*sin(b*x + a) + sqrt(2)*(c^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*co
s(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)))*(2*c^5*d^2*cos(b*x +
a)*sin(b*x + a) + (sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^3*d*cos(b*x + a) + sqrt
(2)*(c^6*d^2/b^4)^(3/4)*b^3*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b
*x + a))))/(c^10*d^4*cos(b*x + a)*sin(b*x + a))) + 2*sqrt(2)*(c^6*d^2/b^4)^(
1/4)*b*arctan(1/2*((sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*cos(b*x + a) - s
qrt(2)*(c^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*sin(b*x + a))*sqrt(d*cos(b*x + a))*s
qrt(c*sin(b*x + a)) - sqrt(4*sqrt(c^6*d^2/b^4)*b^2*c^7*d^3*cos(b*x + a)*sin
(b*x + a) + c^10*d^4 + 2*(sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*sin(b*x + a)
) + sqrt(2)*(c^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*cos(b*x + a))*sqrt(d*cos(b*x +
a))*sqrt(c*sin(b*x + a)))*(2*c^5*d^2*cos(b*x + a)*sin(b*x + a) - (sqrt(2)*(
c^6*d^2/b^4)^(1/4)*b*c^3*d*cos(b*x + a) + sqrt(2)*(c^6*d^2/b^4)^(3/4)*b^3*s
in(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))))/(c^10*d^4*cos(b*x
+ a)*sin(b*x + a))) - sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*log(4*sqrt(c^6*d^2/b^4)
*b^2*c^7*d^3*cos(b*x + a)*sin(b*x + a) + c^10*d^4 + 2*(sqrt(2)*(c^6*d^2/b^4)
)^(1/4)*b*c^8*d^3*sin(b*x + a) + sqrt(2)*(c^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*co
s(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))) + sqrt(2)*(c^6*d^2/b
^4)^(1/4)*b*log(4*sqrt(c^6*d^2/b^4)*b^2*c^7*d^3*cos(b*x + a)*sin(b*x + a) +
c^10*d^4 - 2*(sqrt(2)*(c^6*d^2/b^4)^(1/4)*b*c^8*d^3*sin(b*x + a) + sqrt(2)
*(c^6*d^2/b^4)^(3/4)*b^3*c^5*d^2*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c

```

$\sin(bx + a)))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^{\frac{3}{2}} \sqrt{d \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)*(c*sin(b*x+a))**(3/2),x)

[Out] Integral((c*sin(a + b*x))**(3/2)*sqrt(d*cos(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(3/2),x)

[Out] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(3/2), x)

$$3.272 \quad \int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=313

$$\frac{c^{3/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{d} \sqrt{c \sin(a+bx)}} \right)}{\sqrt{2} b d^{3/2}} + \frac{c^{3/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{d} \sqrt{c \sin(a+bx)}} \right)}{\sqrt{2} b d^{3/2}} + \frac{c^{3/2} \log \left(\sqrt{d} + \right)}{\sqrt{2} b d^{3/2}}$$

[Out] $\frac{1}{2} c^{3/2} \arctan\left(-1 + 2^{1/2} c^{1/2} (d \cos(bx+a))^{1/2} / d^{1/2} / (c \sin(bx+a))^{1/2}\right) / b d^{3/2} + \frac{1}{2} c^{3/2} \arctan\left(1 + 2^{1/2} c^{1/2} (d \cos(bx+a))^{1/2} / d^{1/2} / (c \sin(bx+a))^{1/2}\right) / b d^{3/2} + \frac{1}{4} c^{3/2} \ln\left(d^{1/2} + \cot(bx+a) d^{1/2} - 2^{1/2} c^{1/2} (d \cos(bx+a))^{1/2} / (c \sin(bx+a))^{1/2}\right) / b d^{3/2} - \frac{1}{4} c^{3/2} \ln\left(d^{1/2} + \cot(bx+a) d^{1/2} + 2^{1/2} c^{1/2} (d \cos(bx+a))^{1/2} / (c \sin(bx+a))^{1/2}\right) / b d^{3/2} + 2 c^{3/2} (c \sin(bx+a))^{1/2} / b d / (d \cos(bx+a))^{1/2}$

Rubi [A]

time = 0.18, antiderivative size = 313, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2646, 2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{c^{3/2} \text{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{d} \sqrt{c \sin(a+bx)}}\right)}{\sqrt{2} b d^{3/2}} + \frac{c^{3/2} \text{ArcTan}\left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{d} \sqrt{c \sin(a+bx)}} + 1\right)}{\sqrt{2} b d^{3/2}} + \frac{c^{3/2} \log\left(-\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} + \sqrt{d} \cot(a+bx) + \sqrt{d}\right)}{2 \sqrt{2} b d^{3/2}} - \frac{c^{3/2} \log\left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} + \sqrt{d} \cot(a+bx) + \sqrt{d}\right)}{2 \sqrt{2} b d^{3/2}} + \frac{2 c \sqrt{c \sin(a+bx)}}{b d \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(3/2), x]

[Out] $-\left(\frac{c^{3/2} \text{ArcTan}\left[1 - \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos[a + b*x]}}{\sqrt{d} \sqrt{c \sin[a + b*x]}}\right)\right]}{\sqrt{2} b d^{3/2}}\right) + \left(\frac{c^{3/2} \text{ArcTan}\left[1 + \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos[a + b*x]}}{\sqrt{d} \sqrt{c \sin[a + b*x]}}\right)\right]}{\sqrt{2} b d^{3/2}}\right) + \frac{c^{3/2} \log\left[\sqrt{d} + \sqrt{d} \cot[a + b*x] - \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos[a + b*x]}}{\sqrt{c \sin[a + b*x]}}\right)\right]}{2 \sqrt{2} b d^{3/2}} - \frac{c^{3/2} \log\left[\sqrt{d} + \sqrt{d} \cot[a + b*x] + \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos[a + b*x]}}{\sqrt{c \sin[a + b*x]}}\right)\right]}{2 \sqrt{2} b d^{3/2}} + \frac{2 c \sqrt{c \sin[a + b*x]}}{b d \sqrt{d \cos[a + b*x]}}$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4

, x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2646

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*Sine[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1)), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sine[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2655

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sine[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{3/2}} dx &= \frac{2c \sqrt{c \sin(a + bx)}}{bd \sqrt{d \cos(a + bx)}} - \frac{c^2 \int \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} dx}{d^2} \\
&= \frac{2c \sqrt{c \sin(a + bx)}}{bd \sqrt{d \cos(a + bx)}} + \frac{(2c^3) \text{Subst}\left(\int \frac{x^2}{d^2 + c^2 x^4} dx, x, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{bd} \\
&= \frac{2c \sqrt{c \sin(a + bx)}}{bd \sqrt{d \cos(a + bx)}} - \frac{c^2 \text{Subst}\left(\int \frac{d - cx^2}{d^2 + c^2 x^4} dx, x, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{bd} + \frac{c^2 \text{Subst}\left(\int \frac{d}{d^2 + c^2 x^4} dx, x, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{bd} \\
&= \frac{2c \sqrt{c \sin(a + bx)}}{bd \sqrt{d \cos(a + bx)}} + \frac{c^{3/2} \text{Subst}\left(\int \frac{\frac{\sqrt{2} \sqrt{d}}{\sqrt{c}} + 2x}{-\frac{d}{c} - \frac{\sqrt{2} \sqrt{d}}{\sqrt{c}} x - x^2} dx, x, \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{2\sqrt{2} bd^{3/2}} + \frac{c^{3/2} \log\left(\sqrt{d} + \sqrt{d} \cot(a + bx) - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{2\sqrt{2} bd^{3/2}} - \frac{c^{3/2} \log\left(\sqrt{d} + \sqrt{d} \cot(a + bx) + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}}\right)}{2\sqrt{2} bd^{3/2}} \\
&= -\frac{c^{3/2} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}}\right)}{\sqrt{2} bd^{3/2}} + \frac{c^{3/2} \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}}\right)}{\sqrt{2} bd^{3/2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.11, size = 67, normalized size = 0.21

$$\frac{2^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{4}, \frac{5}{4}; \frac{9}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{5/2}}{5bcd \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(3/2), x]

[Out] (2*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[5/4, 5/4, 9/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(5/2))/(5*b*c*d*Sqrt[d*Cos[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.12, size = 642, normalized size = 2.05

method	result
default	$\left(i \sin(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticPi} \left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{2} b \left(I \sin(bx+a) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} - \frac{1}{2} I, \frac{1}{2} 2^{1/2} \right) - I \sin(bx+a) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} + \frac{1}{2} I, \frac{1}{2} 2^{1/2} \right) + \sin(bx+a) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} - \frac{1}{2} I, \frac{1}{2} 2^{1/2} \right) + \sin(bx+a) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticPi} \left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} + \frac{1}{2} I, \frac{1}{2} 2^{1/2} \right) - 2 \sin(bx+a) \left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2} \operatorname{EllipticF} \left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)} \right)^{1/2}, \frac{1}{2} 2^{1/2} \right) + 2 \cos(bx+a) 2^{1/2} - 2 2^{1/2} \right) \cos(bx+a) \left(c \sin(bx+a) \right)^{3/2} / (-1+\cos(bx+a)) / (d \cos(bx+a))^{3/2} / \sin(bx+a) 2^{1/2} \right)$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")`

[Out] `integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(3/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1865 vs. $2(237) = 474$.

time = 16.75, size = 1865, normalized size = 5.96

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")`

$d^6)^{1/4} \sin(bx + a) \sqrt{d \cos(bx + a)} \sqrt{c \sin(bx + a)} + 16 \sqrt{d \cos(bx + a)} \sqrt{c \sin(bx + a)} c / (b d^2 \cos(bx + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(3/2)/(d*cos(b*x+a))**(3/2),x)

[Out] Integral((c*sin(a + b*x))**(3/2)/(d*cos(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(3/2),x)

[Out] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(3/2), x)

$$3.273 \quad \int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=37

$$\frac{2(c \sin(a+bx))^{5/2}}{5bcd(d \cos(a+bx))^{5/2}}$$

[Out] $2/5*(c*\sin(b*x+a))^{(5/2)}/b/c/d/(d*\cos(b*x+a))^{(5/2)}$

Rubi [A]

time = 0.04, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2643}

$$\frac{2(c \sin(a+bx))^{5/2}}{5bcd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(3/2)}/(d*\text{Cos}[a + b*x])^{(7/2)}, x]$

[Out] $(2*(c*\text{Sin}[a + b*x])^{(5/2)})/(5*b*c*d*(d*\text{Cos}[a + b*x])^{(5/2)})$

Rule 2643

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e + f*x])^{(m + 1)}*((b*\text{Cos}[e + f*x])^{(n + 1)})/(a*b*f*(m + 1)), x] /;$ FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] & & NeQ[m, -1]

Rubi steps

$$\int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{7/2}} dx = \frac{2(c \sin(a+bx))^{5/2}}{5bcd(d \cos(a+bx))^{5/2}}$$

Mathematica [A]

time = 0.08, size = 40, normalized size = 1.08

$$\frac{2 \cot(a+bx)(c \sin(a+bx))^{7/2}}{5bc^2(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(c*\text{Sin}[a + b*x])^{(3/2)}/(d*\text{Cos}[a + b*x])^{(7/2)}, x]$

[Out] $(2*\text{Cot}[a + b*x]*(c*\text{Sin}[a + b*x])^{(7/2)})/(5*b*c^2*(d*\text{Cos}[a + b*x])^{(7/2)})$

Maple [A]

time = 0.10, size = 38, normalized size = 1.03

method	result	size
default	$\frac{2 \sin(bx+a) \cos(bx+a) (c \sin(bx+a))^{\frac{3}{2}}}{5b(d \cos(bx+a))^{\frac{7}{2}}}$	38

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

[Out] $2/5/b*\sin(b*x+a)*\cos(b*x+a)*(c*\sin(b*x+a))^{(3/2)}/(d*\cos(b*x+a))^{(7/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")`

[Out] `integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(7/2), x)`

Fricas [A]

time = 0.40, size = 50, normalized size = 1.35

$$-\frac{2(c \cos(bx+a)^2 - c) \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)}}{5bd^4 \cos(bx+a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")`

[Out] $-2/5*(c*\cos(b*x + a)^2 - c)*\text{sqrt}(d*\cos(b*x + a))*\text{sqrt}(c*\sin(b*x + a))/(b*d^4*\cos(b*x + a)^3)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))**(3/2)/(d*cos(b*x+a))**(7/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(7/2), x)

Mupad [B]

time = 1.51, size = 64, normalized size = 1.73

$$\frac{2c(\cos(4a + 4bx) - 1)\sqrt{c\sin(a + bx)}}{5bd^3\sqrt{d\cos(a + bx)}(4\cos(2a + 2bx) + \cos(4a + 4bx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(7/2),x)

[Out] -(2*c*(cos(4*a + 4*b*x) - 1)*(c*sin(a + b*x))^(1/2))/(5*b*d^3*(d*cos(a + b*x))^(1/2)*(4*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 3))

$$3.274 \quad \int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{11/2}} dx$$

Optimal. Leaf size=106

$$\frac{2c\sqrt{c \sin(a+bx)}}{9bd(d \cos(a+bx))^{9/2}} - \frac{2c\sqrt{c \sin(a+bx)}}{45bd^3(d \cos(a+bx))^{5/2}} - \frac{8c\sqrt{c \sin(a+bx)}}{45bd^5\sqrt{d \cos(a+bx)}}$$

[Out] $2/9*c*(c*\sin(b*x+a))^{(1/2)}/b/d/(d*\cos(b*x+a))^{(9/2)}-2/45*c*(c*\sin(b*x+a))^{(1/2)}/b/d^3/(d*\cos(b*x+a))^{(5/2)}-8/45*c*(c*\sin(b*x+a))^{(1/2)}/b/d^5/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2646, 2651, 2643}

$$-\frac{8c\sqrt{c \sin(a+bx)}}{45bd^5\sqrt{d \cos(a+bx)}} - \frac{2c\sqrt{c \sin(a+bx)}}{45bd^3(d \cos(a+bx))^{5/2}} + \frac{2c\sqrt{c \sin(a+bx)}}{9bd(d \cos(a+bx))^{9/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(11/2), x]

[Out] $(2*c*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(9*b*d*(d*\text{Cos}[a + b*x])^{(9/2)}) - (2*c*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(45*b*d^3*(d*\text{Cos}[a + b*x])^{(5/2)}) - (8*c*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(45*b*d^5*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2646

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m + 1))

```
/(a*b*f*(m + 1)), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x]
)^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m,
-1] && IntegersQ[2*m, 2*n]
```

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{11/2}} dx &= \frac{2c \sqrt{c \sin(a + bx)}}{9bd(d \cos(a + bx))^{9/2}} - \frac{c^2 \int \frac{1}{(d \cos(a + bx))^{7/2} \sqrt{c \sin(a + bx)}} dx}{9d^2} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{9bd(d \cos(a + bx))^{9/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{45bd^3(d \cos(a + bx))^{5/2}} - \frac{(4c^2) \int \frac{1}{(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}}}{45d^4} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{9bd(d \cos(a + bx))^{9/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{45bd^3(d \cos(a + bx))^{5/2}} - \frac{8c \sqrt{c \sin(a + bx)}}{45bd^5 \sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [A]

time = 0.21, size = 57, normalized size = 0.54

$$\frac{2 \sqrt{d \cos(a + bx)} (7 + 2 \cos(2(a + bx))) \sec^5(a + bx) (c \sin(a + bx))^{5/2}}{45bcd^6}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(11/2), x]
```

```
[Out] (2*Sqrt[d*Cos[a + b*x]]*(7 + 2*Cos[2*(a + b*x)])*Sec[a + b*x]^5*(c*Sin[a +
b*x])^(5/2))/(45*b*c*d^6)
```

Maple [A]

time = 0.10, size = 50, normalized size = 0.47

method	result	size
default	$\frac{2(4(\cos^2(bx+a))+5) \cos(bx+a)(c \sin(bx+a))^{\frac{3}{2}} \sin(bx+a)}{45b(d \cos(bx+a))^{\frac{11}{2}}}$	50

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(11/2), x, method=_RETURNVERBOSE)
```

```
[Out] 2/45/b*(4*cos(b*x+a)^2+5)*cos(b*x+a)*(c*sin(b*x+a))^(3/2)*sin(b*x+a)/(d*cos
(b*x+a))^(11/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(11/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(11/2), x)

Fricas [A]

time = 0.45, size = 61, normalized size = 0.58

$$\frac{2(4c \cos(bx+a)^4 + c \cos(bx+a)^2 - 5c) \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)}}{45bd^6 \cos(bx+a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(11/2),x, algorithm="fricas")

[Out] -2/45*(4*c*cos(b*x + a)^4 + c*cos(b*x + a)^2 - 5*c)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))/(b*d^6*cos(b*x + a)^5)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(11/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(11/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(11/2), x)

Mupad [B]

time = 6.16, size = 207, normalized size = 1.95

$$\frac{\sqrt{c \sin(a+bx)} (2 \sin(2a+2bx)^2 + \sin(4a+4bx) - 1) \left(\frac{32c(-2 \sin(2a+2bx)^2 + \sin(4a+4bx) - 1)}{15bd^6} + \frac{16c(2 \sin(2a+2bx)^2 - 1)(-2 \sin(2a+2bx)^2 + \sin(4a+4bx) - 1)}{45bd^6} + \frac{16c(2 \sin(a+bx)^2 - 1)(-2 \sin(2a+2bx)^2 + \sin(4a+4bx) - 1)}{9bd^6} \right)}{16(\sin(a+bx)^2 - 1)^2 \sqrt{-d \left(2 \sin\left(\frac{a}{2} + \frac{bx}{2}\right)^2 - 1 \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(3/2)/(d*cos(a + b*x))^(11/2),x)

```
[Out] -((c*sin(a + b*x))^(1/2)*(sin(4*a + 4*b*x)*1i + 2*sin(2*a + 2*b*x)^2 - 1)*
(32*c*(sin(4*a + 4*b*x)*1i - 2*sin(2*a + 2*b*x)^2 + 1))/(15*b*d^5) + (16*c*
(2*sin(2*a + 2*b*x)^2 - 1)*(sin(4*a + 4*b*x)*1i - 2*sin(2*a + 2*b*x)^2 + 1)
)/(45*b*d^5) + (16*c*(2*sin(a + b*x)^2 - 1)*(sin(4*a + 4*b*x)*1i - 2*sin(2*
a + 2*b*x)^2 + 1))/(9*b*d^5)))/(16*(sin(a + b*x)^2 - 1)^2*(-d*(2*sin(a/2 +
(b*x)/2)^2 - 1))^(1/2))
```

$$3.275 \quad \int \frac{(c \sin(a+bx))^{3/2}}{(d \cos(a+bx))^{15/2}} dx$$

Optimal. Leaf size=141

$$\frac{2c\sqrt{c \sin(a+bx)}}{13bd(d \cos(a+bx))^{13/2}} - \frac{2c\sqrt{c \sin(a+bx)}}{117bd^3(d \cos(a+bx))^{9/2}} - \frac{16c\sqrt{c \sin(a+bx)}}{585bd^5(d \cos(a+bx))^{5/2}} - \frac{64c\sqrt{c \sin(a+bx)}}{585bd^7\sqrt{d \cos(a+bx)}}$$

[Out] 2/13*c*(c*sin(b*x+a))^(1/2)/b/d/(d*cos(b*x+a))^(13/2)-2/117*c*(c*sin(b*x+a))^(1/2)/b/d^3/(d*cos(b*x+a))^(9/2)-16/585*c*(c*sin(b*x+a))^(1/2)/b/d^5/(d*cos(b*x+a))^(5/2)-64/585*c*(c*sin(b*x+a))^(1/2)/b/d^7/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.16, antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2646, 2651, 2643}

$$\frac{64c\sqrt{c \sin(a+bx)}}{585bd^7\sqrt{d \cos(a+bx)}} - \frac{16c\sqrt{c \sin(a+bx)}}{585bd^5(d \cos(a+bx))^{5/2}} - \frac{2c\sqrt{c \sin(a+bx)}}{117bd^3(d \cos(a+bx))^{9/2}} + \frac{2c\sqrt{c \sin(a+bx)}}{13bd(d \cos(a+bx))^{13/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(15/2), x]

[Out] (2*c*Sqrt[c*Sin[a + b*x]])/(13*b*d*(d*Cos[a + b*x])^(13/2)) - (2*c*Sqrt[c*Sin[a + b*x]])/(117*b*d^3*(d*Cos[a + b*x])^(9/2)) - (16*c*Sqrt[c*Sin[a + b*x]])/(585*b*d^5*(d*Cos[a + b*x])^(5/2)) - (64*c*Sqrt[c*Sin[a + b*x]])/(585*b*d^7*Sqrt[d*Cos[a + b*x]])

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2646

Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[(-a)*(a*Sin[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sin[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(-b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m + 1))

`/(a*b*f*(m + 1)), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{3/2}}{(d \cos(a + bx))^{15/2}} dx &= \frac{2c \sqrt{c \sin(a + bx)}}{13bd(d \cos(a + bx))^{13/2}} - \frac{c^2 \int \frac{1}{(d \cos(a + bx))^{11/2} \sqrt{c \sin(a + bx)}} dx}{13d^2} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{13bd(d \cos(a + bx))^{13/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{117bd^3(d \cos(a + bx))^{9/2}} - \frac{(8c^2) \int \frac{1}{(d \cos(a + bx))^{7/2} \sqrt{c \sin(a + bx)}} dx}{117d^4} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{13bd(d \cos(a + bx))^{13/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{117bd^3(d \cos(a + bx))^{9/2}} - \frac{16c \sqrt{c \sin(a + bx)}}{585bd^5(d \cos(a + bx))^{5/2}} \\ &= \frac{2c \sqrt{c \sin(a + bx)}}{13bd(d \cos(a + bx))^{13/2}} - \frac{2c \sqrt{c \sin(a + bx)}}{117bd^3(d \cos(a + bx))^{9/2}} - \frac{16c \sqrt{c \sin(a + bx)}}{585bd^5(d \cos(a + bx))^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.21, size = 67, normalized size = 0.48

$$\frac{2\sqrt{d \cos(a + bx)} (77 + 36 \cos(2(a + bx)) + 4 \cos(4(a + bx))) \sec^7(a + bx) (c \sin(a + bx))^{5/2}}{585bcd^8}$$

Antiderivative was successfully verified.

`[In] Integrate[(c*Sin[a + b*x])^(3/2)/(d*Cos[a + b*x])^(15/2), x]`

`[Out] (2*Sqrt[d*Cos[a + b*x]]*(77 + 36*Cos[2*(a + b*x)] + 4*Cos[4*(a + b*x)])*Sec[a + b*x]^7*(c*Sin[a + b*x])^(5/2))/(585*b*c*d^8)`

Maple [A]

time = 0.12, size = 60, normalized size = 0.43

method	result	size
default	$\frac{2(32(\cos^4(bx+a))+40(\cos^2(bx+a))+45) \cos(bx+a)(c \sin(bx+a))^{\frac{3}{2}} \sin(bx+a)}{585b(d \cos(bx+a))^{\frac{15}{2}}}$	60

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(15/2), x, method=_RETURNVERBOSE)`

`[Out] 2/585/b*(32*cos(b*x+a)^4+40*cos(b*x+a)^2+45)*cos(b*x+a)*(c*sin(b*x+a))^(3/2)*sin(b*x+a)/(d*cos(b*x+a))^(15/2)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(15/2),x, algorithm="maxima")``[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(15/2), x)`**Fricas [A]**

time = 0.49, size = 73, normalized size = 0.52

$$\frac{2(32c\cos(bx+a)^6 + 8c\cos(bx+a)^4 + 5c\cos(bx+a)^2 - 45c)\sqrt{d\cos(bx+a)}\sqrt{c\sin(bx+a)}}{585bd^8\cos(bx+a)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(15/2),x, algorithm="fricas")``[Out] -2/585*(32*c*cos(b*x + a)^6 + 8*c*cos(b*x + a)^4 + 5*c*cos(b*x + a)^2 - 45*c)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))/(b*d^8*cos(b*x + a)^7)`**Sympy [F(-1)] Timed out**

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))**(3/2)/(d*cos(b*x+a))**(15/2),x)``[Out] Timed out`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((c*sin(b*x+a))^(3/2)/(d*cos(b*x+a))^(15/2),x, algorithm="giac")``[Out] integrate((c*sin(b*x + a))^(3/2)/(d*cos(b*x + a))^(15/2), x)`**Mupad [B]**

time = 6.67, size = 193, normalized size = 1.37

$$\frac{e^{-a6i-bx6i}\sqrt{c\left(\frac{e^{-a1i-bx1i}}{2} - \frac{e^{a1i+bx1i}}{2}\right)}\left(-\frac{3776ce^{a6i+bx6i}}{585bd^7} + \frac{2752ce^{a6i+bx6i}\cos(2a+2bx)}{585bd^7} + \frac{896ce^{a6i+bx6i}\cos(4a+4bx)}{585bd^7} + \frac{128ce^{a6i+bx6i}\cos(6a+6bx)}{585bd^7}\right)}{64\cos(a+bx)^6\sqrt{d\left(\frac{e^{-a1i-bx1i}}{2} + \frac{e^{a1i+bx1i}}{2}\right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((c*\sin(a + b*x))^{3/2}/(d*\cos(a + b*x))^{15/2},x)$

[Out] $-(\exp(-a*6i - b*x*6i)*(c*((\exp(-a*1i - b*x*1i)*1i)/2 - (\exp(a*1i + b*x*1i)*1i)/2))^{1/2}*((2752*c*\exp(a*6i + b*x*6i)*\cos(2*a + 2*b*x))/(585*b*d^7) - (3776*c*\exp(a*6i + b*x*6i))/(585*b*d^7) + (896*c*\exp(a*6i + b*x*6i)*\cos(4*a + 4*b*x))/(585*b*d^7) + (128*c*\exp(a*6i + b*x*6i)*\cos(6*a + 6*b*x))/(585*b*d^7)))/(64*\cos(a + b*x)^6*(d*(\exp(-a*1i - b*x*1i)/2 + \exp(a*1i + b*x*1i)/2))^{1/2})$

3.276 $\int (d \cos(a + bx))^{9/2} (c \sin(a + bx))^{5/2} dx$

Optimal. Leaf size=166

$$\frac{cd^3(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{20b} + \frac{3cd(d \cos(a + bx))^{7/2}(c \sin(a + bx))^{3/2}}{70b} - \frac{c(d \cos(a + bx))^{11/2}(c \sin(a + bx))^{3/2}}{7bd}$$

[Out] $\frac{1}{20}cd^3(d \cos(bx+a))^{3/2}(c \sin(bx+a))^{3/2}/b + \frac{3}{70}cd(d \cos(bx+a))^{7/2}(c \sin(bx+a))^{3/2}/b - \frac{1}{7}c(d \cos(bx+a))^{11/2}(c \sin(bx+a))^{3/2}/b - \frac{3}{40}c^2d^4(\sin(a+1/4\pi+bx))^2 \sqrt{\sin(a+1/4\pi+bx)} \operatorname{EllipticE}(\cos(a+1/4\pi+bx), 2^{1/2})(d \cos(bx+a))^{1/2}(c \sin(bx+a))^{1/2}/\sin(2bx+2a)^{1/2}$

Rubi [A]

time = 0.15, antiderivative size = 166, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2648, 2649, 2652, 2719}

$$\frac{3c^2d^4E(a+bx-\frac{\pi}{4}|2)\sqrt{c \sin(a+bx)}\sqrt{d \cos(a+bx)}}{40b\sqrt{\sin(2a+2bx)}} + \frac{cd^3(c \sin(a+bx))^{3/2}(d \cos(a+bx))^{3/2}}{20b} - \frac{c(c \sin(a+bx))^{3/2}(d \cos(a+bx))^{11/2}}{7bd} + \frac{3cd(c \sin(a+bx))^{3/2}(d \cos(a+bx))^{7/2}}{70b}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d \operatorname{Cos}[a + bx])^{9/2}(c \operatorname{Sin}[a + bx])^{5/2}, x]$

[Out] $(c*d^3*(d \operatorname{Cos}[a + bx])^{3/2}(c \operatorname{Sin}[a + bx])^{3/2})/(20*b) + (3*c*d*(d \operatorname{Cos}[a + bx])^{7/2}(c \operatorname{Sin}[a + bx])^{3/2})/(70*b) - (c*(d \operatorname{Cos}[a + bx])^{11/2}(c \operatorname{Sin}[a + bx])^{3/2})/(7*b*d) + (3*c^2*d^4*\operatorname{Sqrt}[d \operatorname{Cos}[a + bx]]*\operatorname{EllipticE}[a - \operatorname{Pi}/4 + bx, 2]*\operatorname{Sqrt}[c \operatorname{Sin}[a + bx]])/(40*b*\operatorname{Sqrt}[\operatorname{Sin}[2*a + 2*b*x]])$

Rule 2648

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)]))^{(m_)}, x_Symbol] \rightarrow \operatorname{Simp}[(-a)*(b \operatorname{Cos}[e + f*x])^{(n+1)}*((a \operatorname{Sin}[e + f*x])^{(m-1)})/(b*f*(m+n)), x] + \operatorname{Dist}[a^2*((m-1)/(m+n)), \operatorname{Int}[(b \operatorname{Cos}[e + f*x])^n*(a \operatorname{Sin}[e + f*x])^{(m-2)}, x], x] /; \operatorname{FreeQ}\{a, b, e, f, n\}, x] \&\& \operatorname{GtQ}[m, 1] \&\& \operatorname{NeQ}[m+n, 0] \&\& \operatorname{IntegersQ}[2*m, 2*n]$

Rule 2649

$\operatorname{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[a*(b \operatorname{Sin}[e + f*x])^{(n+1)}*((a \operatorname{Cos}[e + f*x])^{(m-1)})/(b*f*(m+n)), x] + \operatorname{Dist}[a^2*((m-1)/(m+n)), \operatorname{Int}[(b \operatorname{Sin}[e + f*x])^n*(a \operatorname{Cos}[e + f*x])^{(m-2)}, x], x] /; \operatorname{FreeQ}\{a, b, e, f, n\}, x] \&\& \operatorname{GtQ}[m, 1] \&\& \operatorname{NeQ}[m+n, 0] \&\& \operatorname{IntegersQ}[2*m, 2*n]$

Rule 2652

```
Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{9/2} (c \sin(a + bx))^{5/2} dx &= -\frac{c(d \cos(a + bx))^{11/2} (c \sin(a + bx))^{3/2}}{7bd} + \frac{1}{14} (3c^2) \int (d \cos(a + bx))^{9/2} (c \sin(a + bx))^{3/2} dx \\ &= \frac{3cd(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{70b} - \frac{c(d \cos(a + bx))^{11/2} (c \sin(a + bx))^{3/2}}{7bd} \\ &= \frac{cd^3(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{20b} + \frac{3cd(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{70b} \\ &= \frac{cd^3(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{20b} + \frac{3cd(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{70b} \\ &= \frac{cd^3(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{20b} + \frac{3cd(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{70b} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.12, size = 72, normalized size = 0.43

$$\frac{2(d \cos(a + bx))^{9/2} \sqrt{\cos^2(a + bx)} {}_2F_1\left(-\frac{7}{4}, \frac{7}{4}; \frac{11}{4}; \sin^2(a + bx)\right) \sec^5(a + bx) (c \sin(a + bx))^{7/2}}{7bc}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(9/2)*(c*Sin[a + b*x])^(5/2), x]
```

```
[Out] (2*(d*Cos[a + b*x])^(9/2)*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-7/4, 7/4, 11/4, Sin[a + b*x]^2]*Sec[a + b*x]^5*(c*Sin[a + b*x])^(7/2))/(7*b*c)
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 544 vs. 2(165) = 330.

time = 0.15, size = 545, normalized size = 3.28

method	result
--------	--------

default	$\left(40\sqrt{2} (\cos^8(bx+a)) - 52(\cos^6(bx+a))\sqrt{2} + 21 \cos(bx+a) \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right) \right)$
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
[Out] 1/560/b*(40*2^(1/2)*cos(b*x+a)^8-52*cos(b*x+a)^6*2^(1/2)+21*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-42*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-2*cos(b*x+a)^4*2^(1/2)+21*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-42*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-7*cos(b*x+a)^2*2^(1/2)+21*cos(b*x+a)*2^(1/2))*(d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(5/2)/sin(b*x+a)^3/cos(b*x+a)^5*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(5/2),x, algorithm="maxima")
[Out] integrate((d*cos(b*x + a))^(9/2)*(c*sin(b*x + a))^(5/2), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(5/2),x, algorithm="fricas")
[Out] integral(-(c^2*d^4*cos(b*x + a)^6 - c^2*d^4*cos(b*x + a)^4)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)), x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(9/2)*(c*sin(b*x+a))**(5/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(9/2)*(c*sin(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(9/2)*(c*sin(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + bx))^{9/2} (c \sin(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(9/2)*(c*sin(a + b*x))^(5/2),x)

[Out] int((d*cos(a + b*x))^(9/2)*(c*sin(a + b*x))^(5/2), x)

3.277 $\int (d \cos(a + bx))^{5/2} (c \sin(a + bx))^{5/2} dx$

Optimal. Leaf size=131

$$\frac{cd(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{10b} - \frac{c(d \cos(a + bx))^{7/2}(c \sin(a + bx))^{3/2}}{5bd} + \frac{3c^2d^2\sqrt{d \cos(a + bx)} E(a - \frac{\pi}{4} + bx)}{20b\sqrt{\sin(2a + 2bx)}}$$

[Out] 1/10*c*d*(d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^(3/2)/b-1/5*c*(d*cos(b*x+a))^(7/2)*(c*sin(b*x+a))^(3/2)/b/d-3/20*c^2*d^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticE(cos(a+1/4*Pi+b*x),2^(1/2))*(d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2)/b/sin(2*b*x+2*a)^(1/2)

Rubi [A]

time = 0.11, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2648, 2649, 2652, 2719}

$$\frac{3c^2d^2E(a + bx - \frac{\pi}{4} | 2) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{20b\sqrt{\sin(2a + 2bx)}} - \frac{c(c \sin(a + bx))^{3/2}(d \cos(a + bx))^{7/2}}{5bd} + \frac{cd(c \sin(a + bx))^{3/2}(d \cos(a + bx))^{3/2}}{10b}$$

Antiderivative was successfully verified.

[In] Int[(d*Cos[a + b*x])^(5/2)*(c*Sin[a + b*x])^(5/2),x]

[Out] (c*d*(d*Cos[a + b*x])^(3/2)*(c*Sin[a + b*x])^(3/2))/(10*b) - (c*(d*Cos[a + b*x])^(7/2)*(c*Sin[a + b*x])^(3/2))/(5*b*d) + (3*c^2*d^2*Sqrt[d*Cos[a + b*x]])*EllipticE[a - Pi/4 + b*x, 2]*Sqrt[c*Sin[a + b*x]]/(20*b*Sqrt[Sin[2*a + 2*b*x]])

Rule 2648

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2649

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[a*(b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2652

```
Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^{5/2} (c \sin(a + bx))^{5/2} dx &= -\frac{c(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bd} + \frac{1}{10} (3c^2) \int (d \cos(a + bx))^{5/2} (c \sin(a + bx))^{3/2} dx \\ &= \frac{cd(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{10b} - \frac{c(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bd} \\ &= \frac{cd(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{10b} - \frac{c(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bd} \\ &= \frac{cd(d \cos(a + bx))^{3/2} (c \sin(a + bx))^{3/2}}{10b} - \frac{c(d \cos(a + bx))^{7/2} (c \sin(a + bx))^{3/2}}{5bd} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.13, size = 70, normalized size = 0.53

$$\frac{2d^2 \sqrt{d \cos(a + bx)} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(-\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{5/2} \tan(a + bx)}{7b}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Cos[a + b*x])^(5/2)*(c*Sin[a + b*x])^(5/2), x]
```

```
[Out] (2*d^2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-3/4, 7/4, 11/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(5/2)*Tan[a + b*x])/(7*b)
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 531 vs. 2(136) = 272.

time = 0.13, size = 532, normalized size = 4.06

method	result
--------	--------

default	$\left(4(\cos^6(bx+a))\sqrt{2} - 6(\cos^4(bx+a))\sqrt{2} + 3\cos(bx+a)\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}\right) \text{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)$
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
[Out] 1/40/b*(4*cos(b*x+a)^6*2^(1/2)-6*cos(b*x+a)^4*2^(1/2)+3*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-6*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+3*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-6*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-cos(b*x+a)^2*2^(1/2)+3*cos(b*x+a)*2^(1/2))*(d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(5/2)/cos(b*x+a)^3/sin(b*x+a)^3*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(5/2),x, algorithm="maxima")
[Out] integrate((d*cos(b*x + a))^(5/2)*(c*sin(b*x + a))^(5/2), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(5/2),x, algorithm="fricas")
[Out] integral(-(c^2*d^2*cos(b*x + a)^4 - c^2*d^2*cos(b*x + a)^2)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)), x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(5/2)*(c*sin(b*x+a))**(5/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(5/2)*(c*sin(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + bx))^{5/2} (c \sin(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(5/2)*(c*sin(a + b*x))^(5/2),x)

[Out] int((d*cos(a + b*x))^(5/2)*(c*sin(a + b*x))^(5/2), x)

3.278 $\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{5/2} dx$

Optimal. Leaf size=95

$$\frac{c(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{3bd} + \frac{c^2 \sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a + bx)}}{2b \sqrt{\sin(2a + 2bx)}}$$

[Out] $-1/3*c*(d*\cos(b*x+a))^{(3/2)}*(c*\sin(b*x+a))^{(3/2)}/b/d-1/2*c^2*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2648, 2652, 2719}

$$\frac{c^2 E\left(a + bx - \frac{\pi}{4} \mid 2\right) \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{2b \sqrt{\sin(2a + 2bx)}} - \frac{c(c \sin(a + bx))^{3/2}(d \cos(a + bx))^{3/2}}{3bd}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^(5/2), x]`

[Out] $-1/3*(c*(d*\text{Cos}[a + b*x])^{(3/2)}*(c*\text{Sin}[a + b*x])^{(3/2)})/(b*d) + (c^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(2*b*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2648

`Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*Sin[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*Sin[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

Rule 2652

`Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] :> Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]] , x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned} \int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{5/2} dx &= -\frac{c(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{3bd} + \frac{1}{2}c^2 \int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx \\ &= -\frac{c(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{3bd} + \frac{(c^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)})}{2\sqrt{d \cos(a + bx)}} \\ &= -\frac{c(d \cos(a + bx))^{3/2}(c \sin(a + bx))^{3/2}}{3bd} + \frac{c^2 \sqrt{d \cos(a + bx)} E(a - bx)}{2b\sqrt{\sin(a + bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.07, size = 67, normalized size = 0.71

$$\frac{2\sqrt{d \cos(a + bx)} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{7}{4}; \frac{11}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{5/2} \tan(a + bx)}{7b}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^(5/2), x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 7/4, 11/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(5/2)*Tan[a + b*x])/(7*b)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 518 vs. 2(106) = 212.

time = 0.14, size = 519, normalized size = 5.46

method	result
default	$\frac{\left(2(\cos^4(bx+a))\sqrt{2} - 6\cos(bx+a)\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}\right) \text{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)}{12b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(5/2), x, method=_RETURNVERBOSE)

[Out] 1/12/b*(2*cos(b*x+a)^4*2^(1/2)-6*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+3*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-6*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-

$$\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}+3*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}*((-1+\cos(b*x+a))/\sin(b*x+a))^{(1/2)}*EllipticF(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)},1/2*2^{(1/2)})*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)}-5*\cos(b*x+a)^2*2^{(1/2)}+3*\cos(b*x+a)*2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(5/2)}/\sin(b*x+a)^3/\cos(b*x+a)*2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^(5/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(-(c^2*cos(b*x + a)^2 - c^2)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)*(c*sin(b*x+a))**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 8570 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(5/2),x)

[Out] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(5/2), x)

$$3.279 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=94

$$\frac{2c(c \sin(a+bx))^{3/2}}{bd\sqrt{d \cos(a+bx)}} - \frac{3c^2 \sqrt{d \cos(a+bx)} E(a - \frac{\pi}{4} + bx | 2) \sqrt{c \sin(a+bx)}}{bd^2 \sqrt{\sin(2a+2bx)}}$$

[Out] $2*c*(c*\sin(b*x+a))^{(3/2)}/b/d/(d*\cos(b*x+a))^{(1/2)}+3*c^2*(\sin(a+1/4*Pi+b*x))^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})*(d*\cos(b*x+a))^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/b/d^2/\sin(2*b*x+2*a)^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2646, 2652, 2719}

$$\frac{2c(c \sin(a+bx))^{3/2}}{bd\sqrt{d \cos(a+bx)}} - \frac{3c^2 E(a+bx - \frac{\pi}{4} | 2) \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}}{bd^2 \sqrt{\sin(2a+2bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^{(5/2)}/(d*\text{Cos}[a + b*x])^{(3/2)}, x]$

[Out] $(2*c*(c*\text{Sin}[a + b*x])^{(3/2)})/(b*d*\text{Sqrt}[d*\text{Cos}[a + b*x]]) - (3*c^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(b*d^2*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.))^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}, x_Symbol] :> \text{Simp}[(-a)*(a*\text{Sin}[e + f*x])^{(m-1)}*((b*\text{Cos}[e + f*x])^{(n+1)})/(b*f*(n+1)), x] + \text{Dist}[a^2*((m-1)/(b^2*(n+1))), \text{Int}[(a*\text{Sin}[e + f*x])^{(m-2)}*(b*\text{Cos}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] || \text{EqQ}[m+n, 0])$

Rule 2652

$\text{Int}[\text{Sqrt}[\cos[(e_.) + (f_.)*(x_)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_)]], x_Symbol] :> \text{Dist}[\text{Sqrt}[a*\text{Sin}[e + f*x]]*(\text{Sqrt}[b*\text{Cos}[e + f*x]]/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]), \text{Int}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned}
\int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{3/2}} dx &= \frac{2c(c \sin(a + bx))^{3/2}}{bd \sqrt{d \cos(a + bx)}} - \frac{(3c^2) \int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)} dx}{d^2} \\
&= \frac{2c(c \sin(a + bx))^{3/2}}{bd \sqrt{d \cos(a + bx)}} - \frac{\left(3c^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}\right) \int \sqrt{\sin(2a + 2bx)}}{d^2 \sqrt{\sin(2a + 2bx)}} \\
&= \frac{2c(c \sin(a + bx))^{3/2}}{bd \sqrt{d \cos(a + bx)}} - \frac{3c^2 \sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a + bx)}}{bd^2 \sqrt{\sin(2a + 2bx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.09, size = 67, normalized size = 0.71

$$\frac{2 \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{4}, \frac{7}{4}, \frac{11}{4}, \sin^2(a + bx)\right) (c \sin(a + bx))^{7/2}}{7bcd \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(5/2)/(d*Cos[a + b*x])^(3/2), x]

[Out] (2*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[5/4, 7/4, 11/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(7/2))/(7*b*c*d*Sqrt[d*Cos[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 507 vs. 2(109) = 218.

time = 0.11, size = 508, normalized size = 5.40

method	result
default	$ \frac{\left(6 \cos(bx+a) \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{1-\cos(bx+a)}{\sin(bx+a)}}\right)}{7bcd} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/2/b*(6*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)-3*cos(b*x+a)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2))*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)+6*((-1+cos(b*x+a)+sin(b*x+a))

$$\frac{1}{\sin(b*x+a)^{1/2}} * \left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)} \right)^{1/2} * \text{EllipticE}\left(\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{1/2}, 1/2*2^{1/2}\right) * \left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)} \right)^{1/2} - 3 * \left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)} \right)^{1/2} * \left(\frac{-1+\cos(b*x+a)}{\sin(b*x+a)} \right)^{1/2} * \text{EllipticF}\left(\left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)}\right)^{1/2}, 1/2*2^{1/2}\right) * \left(\frac{-1+\cos(b*x+a)+\sin(b*x+a)}{\sin(b*x+a)} \right)^{1/2} + \cos(b*x+a)^2 * 2^{1/2} - 3 * \cos(b*x+a) * 2^{1/2} + 2 * 2^{1/2} * \cos(b*x+a) * (c * \sin(b*x+a))^{5/2} / \sin(b*x+a)^3 / (d * \cos(b*x+a))^{3/2} * 2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(-(c^2*cos(b*x + a)^2 - c^2)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))/(d^2*cos(b*x + a)^2), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(3/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 6191 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(3/2), x)

[Out] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(3/2), x)

$$3.280 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{7/2}} dx$$

Optimal. Leaf size=133

$$\frac{2c(c \sin(a+bx))^{3/2}}{5bd(d \cos(a+bx))^{5/2}} - \frac{6c(c \sin(a+bx))^{3/2}}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{6c^2 \sqrt{d \cos(a+bx)} E\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{c \sin(a+bx)}}{5bd^4 \sqrt{\sin(2a+2bx)}}$$

[Out] $2/5*c*(c*\sin(b*x+a))^(3/2)/b/d/(d*\cos(b*x+a))^(5/2)-6/5*c*(c*\sin(b*x+a))^(3/2)/b/d^3/(d*\cos(b*x+a))^(1/2)-6/5*c^2*(\sin(a+1/4*\pi+b*x)^2)^(1/2)/\sin(a+1/4*\pi+b*x)*\text{EllipticE}(\cos(a+1/4*\pi+b*x),2^(1/2))*(d*\cos(b*x+a))^(1/2)*(c*\sin(b*x+a))^(1/2)/b/d^4/\sin(2*b*x+2*a)^(1/2)$

Rubi [A]

time = 0.11, antiderivative size = 133, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2646, 2651, 2652, 2719}

$$\frac{6c^2 E\left(a+bx - \frac{\pi}{4} \mid 2\right) \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}}{5bd^4 \sqrt{\sin(2a+2bx)}} - \frac{6c(c \sin(a+bx))^{3/2}}{5bd^3 \sqrt{d \cos(a+bx)}} + \frac{2c(c \sin(a+bx))^{3/2}}{5bd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^(5/2)/(d*\text{Cos}[a + b*x])^(7/2), x]$

[Out] $(2*c*(c*\text{Sin}[a + b*x])^(3/2))/(5*b*d*(d*\text{Cos}[a + b*x])^(5/2)) - (6*c*(c*\text{Sin}[a + b*x])^(3/2))/(5*b*d^3*\text{Sqrt}[d*\text{Cos}[a + b*x]]) + (6*c^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(5*b*d^4*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*\sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> \text{Simp}[(-a)*(a*\text{Sin}[e + f*x])^(m - 1)*(b*\text{Cos}[e + f*x])^(n + 1)/(b*f*(n + 1)), x] + \text{Dist}[a^2*((m - 1)/(b^2*(n + 1))), \text{Int}[(a*\text{Sin}[e + f*x])^(m - 2)*(b*\text{Cos}[e + f*x])^(n + 2), x], x] /; \text{FreeQ}\{a, b, e, f\}, x \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] \parallel \text{EqQ}[m + n, 0])$

Rule 2651

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*\sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] :> \text{Simp}[(-b*\text{Sin}[e + f*x])^(n + 1)*(a*\text{Cos}[e + f*x])^(m + 1)/(a*b*f*(m + 1)), x] + \text{Dist}[(m + n + 2)/(a^2*(m + 1)), \text{Int}[(b*\text{Sin}[e + f*x])^n*(a*\text{Cos}[e + f*x])^(m + 2), x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2652

```
Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{7/2}} dx &= \frac{2c(c \sin(a + bx))^{3/2}}{5bd(d \cos(a + bx))^{5/2}} - \frac{(3c^2) \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{3/2}} dx}{5d^2} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{5bd(d \cos(a + bx))^{5/2}} - \frac{6c(c \sin(a + bx))^{3/2}}{5bd^3 \sqrt{d \cos(a + bx)}} + \frac{(6c^2) \int \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{5d^4} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{5bd(d \cos(a + bx))^{5/2}} - \frac{6c(c \sin(a + bx))^{3/2}}{5bd^3 \sqrt{d \cos(a + bx)}} + \frac{(6c^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)})}{5d^4 \sqrt{\sin(2a + 2bx)}} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{5bd(d \cos(a + bx))^{5/2}} - \frac{6c(c \sin(a + bx))^{3/2}}{5bd^3 \sqrt{d \cos(a + bx)}} + \frac{6c^2 \sqrt{d \cos(a + bx)} E\left(a - \frac{\pi}{4} + bx\right)}{5bd^4 \sqrt{\sin(2a + 2bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.12, size = 70, normalized size = 0.53

$$\frac{2 \cos^2(a + bx)^{5/4} \cot(a + bx) {}_2F_1\left(\frac{7}{4}, \frac{9}{4}; \frac{11}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{9/2}}{7bc^2(d \cos(a + bx))^{7/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c*Sin[a + b*x])^(5/2)/(d*Cos[a + b*x])^(7/2), x]
```

```
[Out] (2*(Cos[a + b*x]^2)^(5/4)*Cot[a + b*x]*Hypergeometric2F1[7/4, 9/4, 11/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(9/2))/(7*b*c^2*(d*Cos[a + b*x])^(7/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 528 vs. 2(138) = 276.

time = 0.13, size = 529, normalized size = 3.98

method	result
--------	--------

default	$\left(3(\cos^3(bx+a)) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right) \right)$
---------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(7/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{5}b(3\cos(bx+a)^3\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2}\operatorname{EllipticF}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2},\frac{1}{2}2^{1/2}\right)-6\cos(bx+a)^3\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2}\operatorname{EllipticE}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2},\frac{1}{2}2^{1/2}\right)+3\cos(bx+a)^2\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2}\operatorname{EllipticF}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2},\frac{1}{2}2^{1/2}\right)-6\cos(bx+a)^2\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2}\left(\frac{-1+\cos(bx+a)}{\sin(bx+a)}\right)^{1/2}\operatorname{EllipticE}\left(\left(\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}\right)^{1/2},\frac{1}{2}2^{1/2}\right)+3\cos(bx+a)^32^{1/2}-4\cos(bx+a)^22^{1/2}+2^{1/2})\cos(bx+a)(c\sin(bx+a))^{5/2}/\sin(bx+a)^3/(d\cos(bx+a))^{7/2}2^{1/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(7/2),x, algorithm="maxima")`

[Out] `integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(7/2), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(7/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Symbolic function `elliptic_ec` takes exactly 1 arguments (2 given)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(7/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + b x))^{5/2}}{(d \cos(a + b x))^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(7/2),x)

[Out] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(7/2), x)

$$3.281 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{11/2}} dx$$

Optimal. Leaf size=168

$$\frac{2c(c \sin(a+bx))^{3/2}}{9bd(d \cos(a+bx))^{9/2}} - \frac{2c(c \sin(a+bx))^{3/2}}{15bd^3(d \cos(a+bx))^{5/2}} - \frac{4c(c \sin(a+bx))^{3/2}}{15bd^5 \sqrt{d \cos(a+bx)}} + \frac{4c^2 \sqrt{d \cos(a+bx)} E\left(a - \frac{\pi}{4} + bx\right)}{15bd^6 \sqrt{\sin(2a+2bx)}}$$

[Out] $2/9*c*(c*\sin(b*x+a))^(3/2)/b/d/(d*\cos(b*x+a))^(9/2)-2/15*c*(c*\sin(b*x+a))^(3/2)/b/d^3/(d*\cos(b*x+a))^(5/2)-4/15*c*(c*\sin(b*x+a))^(3/2)/b/d^5/(d*\cos(b*x+a))^(1/2)-4/15*c^2*(\sin(a+1/4*Pi+b*x)^2)^(1/2)/\sin(a+1/4*Pi+b*x)*\text{EllipticE}(\cos(a+1/4*Pi+b*x),2^(1/2))*(d*\cos(b*x+a))^(1/2)*(c*\sin(b*x+a))^(1/2)/b/d^6/\sin(2*b*x+2*a)^(1/2)$

Rubi [A]

time = 0.15, antiderivative size = 168, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2646, 2651, 2652, 2719}

$$\frac{4c^2 E\left(a+bx-\frac{\pi}{4}\right) \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}}{15bd^6 \sqrt{\sin(2a+2bx)}} - \frac{4c(c \sin(a+bx))^{3/2}}{15bd^5 \sqrt{d \cos(a+bx)}} - \frac{2c(c \sin(a+bx))^{3/2}}{15bd^3(d \cos(a+bx))^{5/2}} + \frac{2c(c \sin(a+bx))^{3/2}}{9bd(d \cos(a+bx))^{9/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a+b*x])^(5/2)/(d*\text{Cos}[a+b*x])^(11/2),x]$

[Out] $(2*c*(c*\text{Sin}[a+b*x])^(3/2))/(9*b*d*(d*\text{Cos}[a+b*x])^(9/2)) - (2*c*(c*\text{Sin}[a+b*x])^(3/2))/(15*b*d^3*(d*\text{Cos}[a+b*x])^(5/2)) - (4*c*(c*\text{Sin}[a+b*x])^(3/2))/(15*b*d^5*\text{Sqrt}[d*\text{Cos}[a+b*x]]) + (4*c^2*\text{Sqrt}[d*\text{Cos}[a+b*x]]*\text{EllipticE}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[c*\text{Sin}[a+b*x]])/(15*b*d^6*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])$

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> \text{Simp}[(-a)*(a*\text{Sin}[e+f*x])^(m-1)*((b*\text{Cos}[e+f*x])^(n+1)/(b*f*(n+1))), x] + \text{Dist}[a^2*((m-1)/(b^2*(n+1))), \text{Int}[(a*\text{Sin}[e+f*x])^(m-2)*(b*\text{Cos}[e+f*x])^(n+2), x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& (\text{IntegersQ}[2*m, 2*n] || \text{EqQ}[m+n, 0])$

Rule 2651

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> \text{Simp}[(-b*\text{Sin}[e+f*x])^(n+1)*((a*\text{Cos}[e+f*x])^(m+1)/(a*b*f*(m+1))), x] + \text{Dist}[(m+n+2)/(a^2*(m+1)), \text{Int}[(b*\text{Sin}[e+f*x])^(n)*((a*\text{Cos}[e+f*x])^(m+2), x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2652

```
Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]
, x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e
+ 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]
```

Rule 2719

```
Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*
(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{11/2}} dx &= \frac{2c(c \sin(a + bx))^{3/2}}{9bd(d \cos(a + bx))^{9/2}} - \frac{c^2 \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{7/2}} dx}{3d^2} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{9bd(d \cos(a + bx))^{9/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{15bd^3(d \cos(a + bx))^{5/2}} - \frac{(2c^2) \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{3/2}} dx}{15d^4} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{9bd(d \cos(a + bx))^{9/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{15bd^3(d \cos(a + bx))^{5/2}} - \frac{4c(c \sin(a + bx))^{3/2}}{15bd^5 \sqrt{d \cos(a + bx)}} + \frac{(4c^2)}{\sqrt{d \cos(a + bx)}} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{9bd(d \cos(a + bx))^{9/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{15bd^3(d \cos(a + bx))^{5/2}} - \frac{4c(c \sin(a + bx))^{3/2}}{15bd^5 \sqrt{d \cos(a + bx)}} + \frac{(4c^2)}{\sqrt{d \cos(a + bx)}} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{9bd(d \cos(a + bx))^{9/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{15bd^3(d \cos(a + bx))^{5/2}} - \frac{4c(c \sin(a + bx))^{3/2}}{15bd^5 \sqrt{d \cos(a + bx)}} + \frac{4c^2}{\sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.13, size = 72, normalized size = 0.43

$$\frac{2 \cos^5(a + bx) \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{7}{4}, \frac{13}{4}; \frac{11}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{7/2}}{7bc(d \cos(a + bx))^{11/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(c*Sin[a + b*x])^(5/2)/(d*Cos[a + b*x])^(11/2), x]
```

```
[Out] (2*Cos[a + b*x]^5*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[7/4, 13/4, 11/4,
Sin[a + b*x]^2]*(c*Sin[a + b*x])^(7/2))/(7*b*c*(d*Cos[a + b*x])^(11/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 543 vs. 2(167) = 334.

time = 0.12, size = 544, normalized size = 3.24

method	result
default	$\frac{-12(\cos^5(bx+a)) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticE}\left(\sqrt{\frac{1-\cos(bx+a)}{\sin(bx+a)}}\right)}{\sin(bx+a)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(11/2),x,method=_RETURNVERBOSE)
[Out] 1/45/b*(-12*cos(b*x+a)^5*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+
cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)
*EllipticE(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+6*cos(
b*x+a)^5*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b
*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-c
os(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-12*cos(b*x+a)^4*((1-co
s(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+
a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticE(((1-cos(b*x+a)+sin(b
*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+6*cos(b*x+a)^4*((1-cos(b*x+a)+sin(b*x
+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+c
os(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a
))^(1/2),1/2*2^(1/2))+6*2^(1/2)*cos(b*x+a)^5-3*cos(b*x+a)^4*2^(1/2)-8*cos(b
*x+a)^2*2^(1/2)+5*2^(1/2))*cos(b*x+a)*(c*sin(b*x+a))^(5/2)/sin(b*x+a)^3/(d*
cos(b*x+a))^(11/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(11/2),x, algorithm="maxima")
[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(11/2), x)
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(11/2),x, algorithm="fricas")
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly
1 arguments (2 given)
```

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(11/2), x)

[Out] Timed out

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(11/2), x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(11/2), x)

Mupad [F]
time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{11/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(11/2), x)

[Out] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(11/2), x)

$$3.282 \quad \int \frac{(c \sin(a+bx))^{5/2}}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=320

$$\frac{3c^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} \right)}{4\sqrt{2} b\sqrt{d}} + \frac{3c^{5/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} \right)}{4\sqrt{2} b\sqrt{d}} + \frac{3c^{5/2} \log \left(\sqrt{c} \right)}{2bd}$$

[Out] $-3/8*c^{(5/2)*\arctan(1-2^{(1/2)*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/c^{(1/2)/(d*\cos(b*x+a))^{(1/2)})/b*2^{(1/2)/d^{(1/2)}+3/8*c^{(5/2)*\arctan(1+2^{(1/2)*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/c^{(1/2)/(d*\cos(b*x+a))^{(1/2)})/b*2^{(1/2)/d^{(1/2)}+3/16*c^{(5/2)*\ln(c^{(1/2)-2^{(1/2)*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(1/2)+c^{(1/2)*\tan(b*x+a))/b*2^{(1/2)/d^{(1/2)}-3/16*c^{(5/2)*\ln(c^{(1/2)+2^{(1/2)*d^{(1/2)}*(c*\sin(b*x+a))^{(1/2)}/(d*\cos(b*x+a))^{(1/2)+c^{(1/2)*\tan(b*x+a))/b*2^{(1/2)/d^{(1/2)}-1/2*c*(c*\sin(b*x+a))^{(3/2)}*(d*\cos(b*x+a))^{(1/2)/b/d}}$

Rubi [A]

time = 0.17, antiderivative size = 320, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2648, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{3c^{5/2} \text{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} \right)}{4\sqrt{2} b\sqrt{d}} + \frac{3c^{5/2} \text{ArcTan} \left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} + 1 \right)}{4\sqrt{2} b\sqrt{d}} + \frac{3c^{5/2} \log \left(\frac{-\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{d \cos(a+bx)}} + \sqrt{c} \tan(a+bx) + \sqrt{c} \right)}{8\sqrt{2} b\sqrt{d}} - \frac{3c^{5/2} \log \left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{d \cos(a+bx)}} + \sqrt{c} \tan(a+bx) + \sqrt{c} \right)}{8\sqrt{2} b\sqrt{d}} - \frac{c(c \sin(a+bx))^{3/2} \sqrt{d \cos(a+bx)}}{2bd}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^(5/2)/Sqrt[d*Cos[a + b*x]], x]

[Out] $(-3*c^{(5/2)*\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/(\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])})/(4*\text{Sqrt}[2]*b*\text{Sqrt}[d]) + (3*c^{(5/2)*\text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/(\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])})/(4*\text{Sqrt}[2]*b*\text{Sqrt}[d]) + (3*c^{(5/2)*\text{Log}[\text{Sqrt}[c] - (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/\text{Sqrt}[d*\text{Cos}[a + b*x]] + \text{Sqrt}[c]*\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b*\text{Sqrt}[d]) - (3*c^{(5/2)*\text{Log}[\text{Sqrt}[c] + (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])]/\text{Sqrt}[d*\text{Cos}[a + b*x]] + \text{Sqrt}[c]*\text{Tan}[a + b*x]])/(8*\text{Sqrt}[2]*b*\text{Sqrt}[d]) - (c*\text{Sqrt}[d*\text{Cos}[a + b*x]]*(c*\text{Sin}[a + b*x])^{(3/2)})/(2*b*d)$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4)

, x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2648

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*SIN[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*SIN[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*SIN[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{(c \sin(a + bx))^{5/2}}{\sqrt{d \cos(a + bx)}} dx &= -\frac{c \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bd} + \frac{1}{4} (3c^2) \int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx \\
&= -\frac{c \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bd} + \frac{(3c^3 d) \operatorname{Subst} \left(\int \frac{x^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} \right)}{2b} \\
&= -\frac{c \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bd} - \frac{(3c^3) \operatorname{Subst} \left(\int \frac{c - dx^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} \right)}{4b} \\
&= -\frac{c \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}{2bd} + \frac{(3c^3) \operatorname{Subst} \left(\int \frac{1}{\frac{c}{d} - \frac{\sqrt{2} \sqrt{c} x}{\sqrt{d}} + x^2} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} \right)}{8bd} \\
&= \frac{3c^{5/2} \log \left(\sqrt{c} - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx) \right)}{8\sqrt{2} b \sqrt{d}} - \frac{3c^{5/2} \log \left(\sqrt{c} + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx) \right)}{8\sqrt{2} b \sqrt{d}} \\
&= -\frac{3c^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} \right)}{4\sqrt{2} b \sqrt{d}} + \frac{3c^{5/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}} \right)}{4\sqrt{2} b \sqrt{d}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.08, size = 67, normalized size = 0.21

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{7}{4}, \frac{11}{4}, \sin^2(a + bx)\right) (c \sin(a + bx))^{5/2} \tan(a + bx)}{7b \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(5/2)/Sqrt[d*Cos[a + b*x]],x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, 7/4, 11/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(5/2)*Tan[a + b*x])/(7*b*Sqrt[d*Cos[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.11, size = 510, normalized size = 1.59

method	result
--------	--------

default	$-\left(3i \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)$
---------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/8/b*(3*I*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*(-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a)^{1/2}*(-1+\cos(b*x+a))/\sin(b*x+a)^{1/2}*\operatorname{EllipticPi}((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-3*I*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*(-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a)^{1/2}*(-1+\cos(b*x+a))/\sin(b*x+a)^{1/2}*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})-3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*(-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a)^{1/2}*(-1+\cos(b*x+a))/\sin(b*x+a)^{1/2}*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-3*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*(-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a)^{1/2}*(-1+\cos(b*x+a))/\sin(b*x+a)^{1/2}*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})+2*\cos(b*x+a)^2*2^{1/2}-2*\cos(b*x+a)*2^{1/2})*(c*\sin(b*x+a))^{5/2}/(-1+\cos(b*x+a))/(d*\cos(b*x+a))^{1/2}/\sin(b*x+a)*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate((c*sin(b*x + a))^(5/2)/sqrt(d*cos(b*x + a)), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2074 vs. 2(238) = 476.

time = 29.81, size = 2074, normalized size = 6.48

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")`

[Out]
$$-1/64*(32*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}*c^2*\sin(b*x + a) - 12*\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4}*b*d*\arctan(((\sqrt{2})*(c^{10}/(b^4*d^2))^{1/4}*b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4}*b^3*c^8*d*\sin(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} + \sqrt{4*\sqrt{c^{10}/(b^4*d^2)}}*b^4$$

$$\begin{aligned}
& 2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + c^{16} - 2*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} \\
& *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4}*b^3*c^8*d*\sin(b*x + \\
& a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}*(2*c^8*\cos(b*x + a)*\sin(b*x \\
& + a) + \sqrt{c^{10}/(b^4*d^2)}*b^2*c^3*d + (\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4}*b* \\
& c^5*\sin(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4}*b^3*d*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} \\
&)/(2*c^{16}*\cos(b*x + a)^2 - c^{16}) - 12 \\
& *\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4}*b*d*\arctan(((\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} \\
& *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4}*b^3*c^8*d*\sin(b*x + a) \\
&))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} - \sqrt{4*\sqrt{c^{10}/(b^4*d^2)}* \\
& b^2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + c^{16} + 2*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} \\
& *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4}*b^3*c^8*d*\sin(b*x \\
& + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}*(2*c^8*\cos(b*x + a)*\sin(b \\
& *x + a) + \sqrt{c^{10}/(b^4*d^2)}*b^2*c^3*d - (\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4}* \\
& b*c^5*\sin(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4}*b^3*d*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} \\
&))/(2*c^{16}*\cos(b*x + a)^2 - c^{16}) - \\
& 12*\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4}*b*d*\arctan(-1/2*(2*c^{16}*\cos(b*x + a)*\sin(b*x + a) - \sqrt{4*\sqrt{c^{10}/(b^4*d^2)}*b^2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + c^{16} + 2*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\sin(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}))*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^5*\sin(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*d*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} + (\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\sin(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} - 4*(b^2*c^{11}*d*\cos(b*x + a)^4 - b^2*c^{11}*d*\cos(b*x + a)^2)*\sqrt{c^{10}/(b^4*d^2)})))/((2*c^{16}*\cos(b*x + a)^3 - c^{16}*\cos(b*x + a))*\sin(b*x + a)) - 12*\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*d*\arctan(1/2*(2*c^{16}*\cos(b*x + a)*\sin(b*x + a) + \sqrt{4*\sqrt{c^{10}/(b^4*d^2)}*b^2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + c^{16} - 2*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\sin(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}))*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^5*\sin(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*d*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} - (\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\sin(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\cos(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)} - 4*(b^2*c^{11}*d*\cos(b*x + a)^4 - b^2*c^{11}*d*\cos(b*x + a)^2)*\sqrt{c^{10}/(b^4*d^2)})))/((2*c^{16}*\cos(b*x + a)^3 - c^{16}*\cos(b*x + a))*\sin(b*x + a)) + 3*\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*d*\log(2916*\sqrt{c^{10}/(b^4*d^2)}*b^2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + 729*c^{16} + 1458*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\sin(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)})) - 3*\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*d*\log(2916*\sqrt{c^{10}/(b^4*d^2)}*b^2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + 729*c^{16} - 1458*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\sin(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)})) + 3*\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*d*\log(729/4*\sqrt{c^{10}/(b^4*d^2)}*b^2*c^{11}*d*\cos(b*x + a)*\sin(b*x + a) + 729/16*c^{16} + 729/8*(\sqrt{2}*(c^{10}/(b^4*d^2))^{1/4} *b*c^{13}*\cos(b*x + a) + \sqrt{2}*(c^{10}/(b^4*d^2))^{3/4} *b^3*c^8*d*\sin(b*x + a))*\sqrt{d*\cos(b*x + a)}*\sqrt{c*\sin(b*x + a)}))
\end{aligned}$$

(2)*(c^10/(b^4*d^2))^(3/4)*b^3*c^8*d*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))) - 3*sqrt(2)*(c^10/(b^4*d^2))^(1/4)*b*d*log(729/4*sqrt(c^10/(b^4*d^2))*b^2*c^11*d*cos(b*x + a)*sin(b*x + a) + 729/16*c^16 - 729/8*(sqrt(2)*(c^10/(b^4*d^2))^(1/4)*b*c^13*cos(b*x + a) + sqrt(2)*(c^10/(b^4*d^2))^(3/4)*b^3*c^8*d*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))))/(b*d)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(1/2), x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 6191 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(1/2), x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2)/sqrt(d*cos(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(c \sin(a + bx))^{5/2}}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(1/2), x)

[Out] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(1/2), x)

$$3.283 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=315

$$\frac{c^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} \right)}{\sqrt{2} b d^{5/2}} - \frac{c^{5/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} \right)}{\sqrt{2} b d^{5/2}} - c^{5/2} \log \left(\sqrt{c} - \sqrt{d \cos(a+bx)} \right)$$

[Out] $2/3*c*(c*\sin(b*x+a))^(3/2)/b/d/(d*\cos(b*x+a))^(3/2)+1/2*c^(5/2)*\arctan(1-2^(1/2)*d^(1/2)*(c*\sin(b*x+a))^(1/2)/c^(1/2)/(d*\cos(b*x+a))^(1/2))/b/d^(5/2)*2^(1/2)-1/2*c^(5/2)*\arctan(1+2^(1/2)*d^(1/2)*(c*\sin(b*x+a))^(1/2)/c^(1/2)/(d*\cos(b*x+a))^(1/2))/b/d^(5/2)*2^(1/2)-1/4*c^(5/2)*\ln(c^(1/2)-2^(1/2)*d^(1/2)*(c*\sin(b*x+a))^(1/2)/(d*\cos(b*x+a))^(1/2)+c^(1/2)*\tan(b*x+a))/b/d^(5/2)*2^(1/2)+1/4*c^(5/2)*\ln(c^(1/2)+2^(1/2)*d^(1/2)*(c*\sin(b*x+a))^(1/2)/(d*\cos(b*x+a))^(1/2)+c^(1/2)*\tan(b*x+a))/b/d^(5/2)*2^(1/2)$

Rubi [A]

time = 0.18, antiderivative size = 315, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2646, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{c^{5/2} \text{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} \right)}{\sqrt{2} b d^{5/2}} - \frac{c^{5/2} \text{ArcTan} \left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{c} \sqrt{d \cos(a+bx)}} + 1 \right)}{\sqrt{2} b d^{5/2}} - \frac{c^{5/2} \log \left(\frac{-\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{d \cos(a+bx)}} + \sqrt{c} \tan(a+bx) + \sqrt{c} \right)}{2\sqrt{2} b d^{5/2}} + \frac{c^{5/2} \log \left(\frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a+bx)}}{\sqrt{d \cos(a+bx)}} + \sqrt{c} \tan(a+bx) + \sqrt{c} \right)}{2\sqrt{2} b d^{5/2}} + \frac{2c(c \sin(a+bx))^{3/2}}{3bd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*SIn[a + b*x])^(5/2)/(d*Cos[a + b*x])^(5/2), x]

[Out] $(c^(5/2)*\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])]/(\text{Sqrt}[2]*b*d^(5/2)) - (c^(5/2)*\text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(\text{Sqrt}[c]*\text{Sqrt}[d*\text{Cos}[a + b*x]])]/(\text{Sqrt}[2]*b*d^(5/2)) - (c^(5/2)*\text{Log}[\text{Sqrt}[c] - (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/\text{Sqrt}[d*\text{Cos}[a + b*x]] + \text{Sqrt}[c]*\text{Tan}[a + b*x]])/(2*\text{Sqrt}[2]*b*d^(5/2)) + (c^(5/2)*\text{Log}[\text{Sqrt}[c] + (\text{Sqrt}[2]*\text{Sqrt}[d]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/\text{Sqrt}[d*\text{Cos}[a + b*x]] + \text{Sqrt}[c]*\text{Tan}[a + b*x]])/(2*\text{Sqrt}[2]*b*d^(5/2)) + (2*c*(c*\text{Sin}[a + b*x])^(3/2))/(3*b*d*(d*\text{Cos}[a + b*x])^(3/2))$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4)

), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2646

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*Sine[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sine[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sine[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
 \int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{5/2}} dx &= \frac{2c(c \sin(a + bx))^{3/2}}{3bd(d \cos(a + bx))^{3/2}} - \frac{c^2 \int \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} dx}{d^2} \\
 &= \frac{2c(c \sin(a + bx))^{3/2}}{3bd(d \cos(a + bx))^{3/2}} - \frac{(2c^3) \text{Subst}\left(\int \frac{x^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{bd} \\
 &= \frac{2c(c \sin(a + bx))^{3/2}}{3bd(d \cos(a + bx))^{3/2}} + \frac{c^3 \text{Subst}\left(\int \frac{c - dx^2}{c^2 + d^2 x^4} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{bd^2} - \frac{c^3 \text{Subst}\left(\int \frac{1}{\frac{c}{d} - \frac{\sqrt{2} \sqrt{c} x}{\sqrt{d}} + x^2} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{2bd^3} \\
 &= \frac{2c(c \sin(a + bx))^{3/2}}{3bd(d \cos(a + bx))^{3/2}} - \frac{c^3 \text{Subst}\left(\int \frac{1}{\frac{c}{d} - \frac{\sqrt{2} \sqrt{c} x}{\sqrt{d}} + x^2} dx, x, \frac{\sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}}\right)}{2bd^3} \\
 &= -\frac{c^{5/2} \log\left(\sqrt{c} - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx)\right)}{2\sqrt{2} bd^{5/2}} + \frac{c^{5/2} \log\left(\sqrt{c} + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{d \cos(a + bx)}} + \sqrt{c} \tan(a + bx)\right)}{2\sqrt{2} bd^{5/2}} \\
 &= \frac{c^{5/2} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{\sqrt{2} bd^{5/2}} - \frac{c^{5/2} \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt{d} \sqrt{c \sin(a + bx)}}{\sqrt{c} \sqrt{d \cos(a + bx)}}\right)}{\sqrt{2} bd^{5/2}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.09, size = 67, normalized size = 0.21

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{7}{4}, \frac{11}{4}; \sin^2(a + bx)\right) (c \sin(a + bx))^{7/2}}{7bcd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(5/2)/(d*Cos[a + b*x])^(5/2), x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[7/4, 7/4, 11/4, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(7/2))/(7*b*c*d*(d*Cos[a + b*x])^(3/2))

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.11, size = 532, normalized size = 1.69

method	result
default	$\left(-3i \cos(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticPi} \left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(5/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/6/b*(-3*I*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))+3*I*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-3*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))-3*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))+2*cos(b*x+a)*2^(1/2)-2*2^(1/2))*cos(b*x+a)*(c*sin(b*x+a))^(5/2)/(-1+cos(b*x+a))/(d*cos(b*x+a))^(5/2)/sin(b*x+a)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(5/2), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 2268 vs. 2(238) = 476.

time = 30.68, size = 2268, normalized size = 7.20

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(5/2),x, algorithm="fricas")
```

```
[Out] 1/48*(12*sqrt(2)*b*d^3*(c^10/(b^4*d^10))^(1/4)*arctan(-((sqrt(2)*b^3*c^8*d^7*(c^10/(b^4*d^10))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^13*d^2*(c^10/(b^4*d^10
```

$$\begin{aligned}
&))^{1/4} \cos(b*x + a) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)} - \sqrt{4*b^2*c^{11}*d^5*\sqrt{c^{10}/(b^4*d^{10})}} \cos(b*x + a) \sin(b*x + a) + c^{16} - 2*(\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \sin(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)}) * (b^2*c^3*d^5*\sqrt{c^{10}/(b^4*d^{10})} + 2*c^8*\cos(b*x + a) \sin(b*x + a) + (\sqrt{2}*b^3*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \cos(b*x + a) + \sqrt{2}*b*c^5*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \sin(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)})) / (2*c^{16}*\cos(b*x + a)^2 - c^{16}) * \cos(b*x + a)^2 + 12*\sqrt{2}*b*d^3*(c^{10}/(b^4*d^{10}))^{1/4} * \arctan(-(\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \sin(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)} + \sqrt{4*b^2*c^{11}*d^5*\sqrt{c^{10}/(b^4*d^{10})}} \cos(b*x + a) \sin(b*x + a) + c^{16} + 2*(\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \sin(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)}) * (b^2*c^3*d^5*\sqrt{c^{10}/(b^4*d^{10})} + 2*c^8*\cos(b*x + a) \sin(b*x + a) - (\sqrt{2}*b^3*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \cos(b*x + a) + \sqrt{2}*b*c^5*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \sin(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)})) / (2*c^{16}*\cos(b*x + a)^2 - c^{16}) * \cos(b*x + a)^2 - 12*\sqrt{2}*b*d^3*(c^{10}/(b^4*d^{10}))^{1/4} * \arctan(-1/2*(2*c^{16}*\cos(b*x + a) \sin(b*x + a) - \sqrt{4*b^2*c^{11}*d^5*\sqrt{c^{10}/(b^4*d^{10})}} \cos(b*x + a) \sin(b*x + a) + c^{16} + 2*(\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \sin(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)})) * (\sqrt{2}*b^3*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \cos(b*x + a) + \sqrt{2}*b*c^5*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \sin(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)} + (\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \cos(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \sin(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)} - 4*(b^2*c^{11}*d^5*\cos(b*x + a)^4 - b^2*c^{11}*d^5*\cos(b*x + a)^2) \sqrt{c^{10}/(b^4*d^{10})}) / ((2*c^{16}*\cos(b*x + a)^3 - c^{16}*\cos(b*x + a) \sin(b*x + a)) * \cos(b*x + a)^2 - 12*\sqrt{2}*b*d^3*(c^{10}/(b^4*d^{10}))^{1/4} * \arctan(1/2*(2*c^{16}*\cos(b*x + a) \sin(b*x + a) + \sqrt{4*b^2*c^{11}*d^5*\sqrt{c^{10}/(b^4*d^{10})}} \cos(b*x + a) \sin(b*x + a) + c^{16} - 2*(\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \sin(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)})) * (\sqrt{2}*b^3*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \cos(b*x + a) + \sqrt{2}*b*c^5*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \sin(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)} - (\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \cos(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \sin(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)} - 4*(b^2*c^{11}*d^5*\cos(b*x + a)^4 - b^2*c^{11}*d^5*\cos(b*x + a)^2) \sqrt{c^{10}/(b^4*d^{10})}) / ((2*c^{16}*\cos(b*x + a)^3 - c^{16}*\cos(b*x + a) \sin(b*x + a)) * \cos(b*x + a)^2 + 3*\sqrt{2}*b*d^3*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)^2 * \log(4*b^2*c^{11}*d^5*\sqrt{c^{10}/(b^4*d^{10})}} \cos(b*x + a) \sin(b*x + a) + c^{16} + 2*(\sqrt{2}*b^3*c^8*d^7*(c^{10}/(b^4*d^{10}))^{3/4} \sin(b*x + a) + \sqrt{2}*b*c^{13}*d^2*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)) \sqrt{d \cos(b*x + a)} \sqrt{c \sin(b*x + a)})) - 3*\sqrt{2}*b*d^3*(c^{10}/(b^4*d^{10}))^{1/4} \cos(b*x + a)^2 * \log(4*b^2*c^{11}*d^5*\sqrt{c^{10}/(b^4*d^{10})}} \cos(b*x + a) \sin(b*x + a) + c^{16} - 2*(\sqrt{2}*b^3*c^8*d^7*(c^{10}
\end{aligned}$$

```
/(b^4*d^10))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^13*d^2*(c^10/(b^4*d^10))^(1/4)
)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))) + 3*sqrt(2)*b*d^
3*(c^10/(b^4*d^10))^(1/4)*cos(b*x + a)^2*log(1/4*b^2*c^11*d^5*sqrt(c^10/(b^
4*d^10))*cos(b*x + a)*sin(b*x + a) + 1/16*c^16 + 1/8*(sqrt(2)*b^3*c^8*d^7*(
c^10/(b^4*d^10))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^13*d^2*(c^10/(b^4*d^10))^(
1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))) - 3*sqrt(2)*
b*d^3*(c^10/(b^4*d^10))^(1/4)*cos(b*x + a)^2*log(1/4*b^2*c^11*d^5*sqrt(c^10
/(b^4*d^10))*cos(b*x + a)*sin(b*x + a) + 1/16*c^16 - 1/8*(sqrt(2)*b^3*c^8*d
^7*(c^10/(b^4*d^10))^(3/4)*sin(b*x + a) + sqrt(2)*b*c^13*d^2*(c^10/(b^4*d^1
0))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))) + 32*sq
rt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*c^2*sin(b*x + a)/(b*d^3*cos(b*x + a
)^2)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(5/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 6191 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(5/2),x, algorithm="giac")
```

```
[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(5/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(5/2),x)
```

```
[Out] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(5/2), x)
```

$$3.284 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{9/2}} dx$$

Optimal. Leaf size=37

$$\frac{2(c \sin(a+bx))^{7/2}}{7bcd(d \cos(a+bx))^{7/2}}$$

[Out] $2/7*(c*\sin(b*x+a))^(7/2)/b/c/d/(d*\cos(b*x+a))^(7/2)$

Rubi [A]

time = 0.04, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2643}

$$\frac{2(c \sin(a+bx))^{7/2}}{7bcd(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^(5/2)/(d*\text{Cos}[a + b*x])^(9/2), x]$

[Out] $(2*(c*\text{Sin}[a + b*x])^(7/2))/(7*b*c*d*(d*\text{Cos}[a + b*x])^(7/2))$

Rule 2643

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e + f*x])^(m + 1)*((b*\text{Cos}[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m + n + 2, 0] \& \& \text{NeQ}[m, -1]$

Rubi steps

$$\int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{9/2}} dx = \frac{2(c \sin(a+bx))^{7/2}}{7bcd(d \cos(a+bx))^{7/2}}$$

Mathematica [A]

time = 0.10, size = 40, normalized size = 1.08

$$\frac{2 \cot(a+bx)(c \sin(a+bx))^{9/2}}{7bc^2(d \cos(a+bx))^{9/2}}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[(c*\text{Sin}[a + b*x])^(5/2)/(d*\text{Cos}[a + b*x])^(9/2), x]$

[Out] $(2*\text{Cot}[a + b*x]*(c*\text{Sin}[a + b*x])^{(9/2)})/(7*b*c^2*(d*\text{Cos}[a + b*x])^{(9/2)})$

Maple [A]

time = 0.10, size = 38, normalized size = 1.03

method	result	size
default	$\frac{2 \sin(bx+a) \cos(bx+a) (c \sin(bx+a))^{\frac{5}{2}}}{7b(d \cos(bx+a))^{\frac{9}{2}}}$	38

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(9/2),x,method=_RETURNVERBOSE)`

[Out] $2/7/b*\sin(b*x+a)*\cos(b*x+a)*(c*\sin(b*x+a))^{(5/2)}/(d*\cos(b*x+a))^{(9/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(9/2),x, algorithm="maxima")`

[Out] `integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(9/2), x)`

Fricas [A]

time = 0.42, size = 60, normalized size = 1.62

$$\frac{2(c^2 \cos(bx+a)^2 - c^2) \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)} \sin(bx+a)}{7bd^5 \cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(9/2),x, algorithm="fricas")`

[Out] $-2/7*(c^2*\cos(b*x + a)^2 - c^2)*\text{sqrt}(d*\cos(b*x + a))*\text{sqrt}(c*\sin(b*x + a))*\sin(b*x + a)/(b*d^5*\cos(b*x + a)^4)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(9/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(9/2),x, algorithm="giac")
```

```
[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(9/2), x)
```

Mupad [B]

time = 1.78, size = 89, normalized size = 2.41

$$\frac{2c^2 \sqrt{c \sin(a + bx)} (3 \sin(2a + 2bx) - \sin(6a + 6bx))}{7bd^4 \sqrt{d \cos(a + bx)} (15 \cos(2a + 2bx) + 6 \cos(4a + 4bx) + \cos(6a + 6bx) + 10)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(9/2),x)
```

```
[Out] (2*c^2*(c*sin(a + b*x))^(1/2)*(3*sin(2*a + 2*b*x) - sin(6*a + 6*b*x)))/(7*b
*d^4*(d*cos(a + b*x))^(1/2)*(15*cos(2*a + 2*b*x) + 6*cos(4*a + 4*b*x) + cos
(6*a + 6*b*x) + 10))
```

$$3.285 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{13/2}} dx$$

Optimal. Leaf size=106

$$\frac{2c(c \sin(a+bx))^{3/2}}{11bd(d \cos(a+bx))^{11/2}} - \frac{6c(c \sin(a+bx))^{3/2}}{77bd^3(d \cos(a+bx))^{7/2}} - \frac{8c(c \sin(a+bx))^{3/2}}{77bd^5(d \cos(a+bx))^{3/2}}$$

[Out] $2/11*c*(c*\sin(b*x+a))^(3/2)/b/d/(d*\cos(b*x+a))^(11/2)-6/77*c*(c*\sin(b*x+a))^(3/2)/b/d^3/(d*\cos(b*x+a))^(7/2)-8/77*c*(c*\sin(b*x+a))^(3/2)/b/d^5/(d*\cos(b*x+a))^(3/2)$

Rubi [A]

time = 0.11, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2646, 2651, 2643}

$$-\frac{8c(c \sin(a+bx))^{3/2}}{77bd^5(d \cos(a+bx))^{3/2}} - \frac{6c(c \sin(a+bx))^{3/2}}{77bd^3(d \cos(a+bx))^{7/2}} + \frac{2c(c \sin(a+bx))^{3/2}}{11bd(d \cos(a+bx))^{11/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^(5/2)/(d*\text{Cos}[a + b*x])^(13/2), x]$

[Out] $(2*c*(c*\text{Sin}[a + b*x])^(3/2))/(11*b*d*(d*\text{Cos}[a + b*x])^(11/2)) - (6*c*(c*\text{Sin}[a + b*x])^(3/2))/(77*b*d^3*(d*\text{Cos}[a + b*x])^(7/2)) - (8*c*(c*\text{Sin}[a + b*x])^(3/2))/(77*b*d^5*(d*\text{Cos}[a + b*x])^(3/2))$

Rule 2643

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e + f*x])^(m + 1)*((b*\text{Cos}[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /;$ FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\text{Sin}[e + f*x])^(m - 1)*((b*\text{Cos}[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + \text{Dist}[a^2*((m - 1)/(b^2*(n + 1))), \text{Int}[(a*\text{Sin}[e + f*x])^(m - 2)*(b*\text{Cos}[e + f*x])^(n + 2), x], x] /;$ FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2651

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] \rightarrow \text{Simp}[(-b*\text{Sin}[e + f*x])^(n + 1)*((a*\text{Cos}[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + \text{Dist}[(m + n + 2)/(a^2*(m + 1)), \text{Int}[(b*\text{Sin}[e + f*x]$

)^n*(a*cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{13/2}} dx &= \frac{2c(c \sin(a + bx))^{3/2}}{11bd(d \cos(a + bx))^{11/2}} - \frac{(3c^2) \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{9/2}} dx}{11d^2} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{11bd(d \cos(a + bx))^{11/2}} - \frac{6c(c \sin(a + bx))^{3/2}}{77bd^3(d \cos(a + bx))^{7/2}} - \frac{(12c^2) \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{5/2}} dx}{77d^4} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{11bd(d \cos(a + bx))^{11/2}} - \frac{6c(c \sin(a + bx))^{3/2}}{77bd^3(d \cos(a + bx))^{7/2}} - \frac{8c(c \sin(a + bx))^{3/2}}{77bd^5(d \cos(a + bx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.22, size = 57, normalized size = 0.54

$$\frac{2c^4(9 + 2 \cos(2(a + bx))) \tan^5(a + bx)}{77bd^6 \sqrt{d \cos(a + bx)} (c \sin(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^(5/2)/(d*Cos[a + b*x])^(13/2), x]

[Out] (2*c^4*(9 + 2*Cos[2*(a + b*x)])*Tan[a + b*x]^5)/(77*b*d^6*Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^(3/2))

Maple [A]

time = 0.10, size = 50, normalized size = 0.47

method	result	size
default	$\frac{2(4(\cos^2(bx+a))+7) \cos(bx+a)(c \sin(bx+a))^{\frac{5}{2}} \sin(bx+a)}{77b(d \cos(bx+a))^{\frac{13}{2}}}$	50

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(13/2), x, method=_RETURNVERBOSE)

[Out] 2/77/b*(4*cos(b*x+a)^2+7)*cos(b*x+a)*(c*sin(b*x+a))^(5/2)*sin(b*x+a)/(d*cos(b*x+a))^(13/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(13/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(13/2), x)

Fricas [A]

time = 0.48, size = 74, normalized size = 0.70

$$\frac{2(4c^2 \cos(bx+a)^4 + 3c^2 \cos(bx+a)^2 - 7c^2) \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)} \sin(bx+a)}{77bd^7 \cos(bx+a)^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(13/2),x, algorithm="fricas")

[Out] -2/77*(4*c^2*cos(b*x + a)^4 + 3*c^2*cos(b*x + a)^2 - 7*c^2)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*sin(b*x + a)/(b*d^7*cos(b*x + a)^6)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(13/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(13/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(13/2), x)

Mupad [B]

time = 6.34, size = 176, normalized size = 1.66

$$\frac{e^{-a5i-bx5i} \sqrt{c \left(\frac{e^{-a1i-bx1i} 1i}{2} - \frac{e^{a1i+bx1i} 1i}{2} \right)} \left(\frac{96c^2 e^{a5i+bx5i} \sin(3a+3bx)}{77bd^6} + \frac{16c^2 e^{a5i+bx5i} \sin(5a+5bx)}{77bd^6} - \frac{368c^2 e^{a5i+bx5i} \sin(a+bx)}{77bd^6} \right)}{32 \cos(a+bx)^5 \sqrt{d \left(\frac{e^{-a1i-bx1i}}{2} + \frac{e^{a1i+bx1i}}{2} \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(a + b*x))^(5/2)/(d*cos(a + b*x))^(13/2),x)`

[Out] $-(\exp(-a*5i - b*x*5i)*(c*((\exp(-a*1i - b*x*1i)*1i)/2 - (\exp(a*1i + b*x*1i)*1i)/2))^{(1/2)*((96*c^2*\exp(a*5i + b*x*5i)*\sin(3*a + 3*b*x))/(77*b*d^6) + (16*c^2*\exp(a*5i + b*x*5i)*\sin(5*a + 5*b*x))/(77*b*d^6) - (368*c^2*\exp(a*5i + b*x*5i)*\sin(a + b*x))/(77*b*d^6)))/(32*\cos(a + b*x)^5*(d*(\exp(-a*1i - b*x*1i)/2 + \exp(a*1i + b*x*1i)/2))^{(1/2)})$

$$3.286 \quad \int \frac{(c \sin(a+bx))^{5/2}}{(d \cos(a+bx))^{17/2}} dx$$

Optimal. Leaf size=141

$$\frac{2c(c \sin(a+bx))^{3/2}}{15bd(d \cos(a+bx))^{15/2}} - \frac{2c(c \sin(a+bx))^{3/2}}{55bd^3(d \cos(a+bx))^{11/2}} - \frac{16c(c \sin(a+bx))^{3/2}}{385bd^5(d \cos(a+bx))^{7/2}} - \frac{64c(c \sin(a+bx))^{3/2}}{1155bd^7(d \cos(a+bx))^{3/2}}$$

[Out] $2/15*c*(c*\sin(b*x+a))^(3/2)/b/d/(d*\cos(b*x+a))^(15/2)-2/55*c*(c*\sin(b*x+a))^(3/2)/b/d^3/(d*\cos(b*x+a))^(11/2)-16/385*c*(c*\sin(b*x+a))^(3/2)/b/d^5/(d*\cos(b*x+a))^(7/2)-64/1155*c*(c*\sin(b*x+a))^(3/2)/b/d^7/(d*\cos(b*x+a))^(3/2)$

Rubi [A]

time = 0.15, antiderivative size = 141, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2646, 2651, 2643}

$$-\frac{64c(c \sin(a+bx))^{3/2}}{1155bd^7(d \cos(a+bx))^{3/2}} - \frac{16c(c \sin(a+bx))^{3/2}}{385bd^5(d \cos(a+bx))^{7/2}} - \frac{2c(c \sin(a+bx))^{3/2}}{55bd^3(d \cos(a+bx))^{11/2}} + \frac{2c(c \sin(a+bx))^{3/2}}{15bd(d \cos(a+bx))^{15/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^(5/2)/(d*\text{Cos}[a + b*x])^(17/2), x]$

[Out] $(2*c*(c*\text{Sin}[a + b*x])^(3/2))/(15*b*d*(d*\text{Cos}[a + b*x])^(15/2)) - (2*c*(c*\text{Sin}[a + b*x])^(3/2))/(55*b*d^3*(d*\text{Cos}[a + b*x])^(11/2)) - (16*c*(c*\text{Sin}[a + b*x])^(3/2))/(385*b*d^5*(d*\text{Cos}[a + b*x])^(7/2)) - (64*c*(c*\text{Sin}[a + b*x])^(3/2))/(1155*b*d^7*(d*\text{Cos}[a + b*x])^(3/2))$

Rule 2643

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e + f*x])^(m + 1)*((b*\text{Cos}[e + f*x])^(n + 1))/(a*b*f*(m + 1)), x] /;$ FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2646

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\text{Sin}[e + f*x])^(m - 1)*((b*\text{Cos}[e + f*x])^(n + 1))/(b*f*(n + 1)), x] + \text{Dist}[a^2*((m - 1)/(b^2*(n + 1))), \text{Int}[(a*\text{Sin}[e + f*x])^(m - 2)*(b*\text{Cos}[e + f*x])^(n + 2), x], x] /;$ FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2651

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] \rightarrow \text{Simp}[(-b*\text{Sin}[e + f*x])^(n + 1)*((a*\text{Cos}[e + f*x])^(m + 1))$

$/(a*b*f*(m + 1)), x] + \text{Dist}[(m + n + 2)/(a^2*(m + 1)), \text{Int}[(b*\text{Sin}[e + f*x])^n*(a*\text{Cos}[e + f*x])^{m + 2}, x], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a + bx))^{5/2}}{(d \cos(a + bx))^{17/2}} dx &= \frac{2c(c \sin(a + bx))^{3/2}}{15bd(d \cos(a + bx))^{15/2}} - \frac{c^2 \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{13/2}} dx}{5d^2} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{15bd(d \cos(a + bx))^{15/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{55bd^3(d \cos(a + bx))^{11/2}} - \frac{(8c^2) \int \frac{\sqrt{c \sin(a + bx)}}{(d \cos(a + bx))^{9/2}} dx}{55d^4} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{15bd(d \cos(a + bx))^{15/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{55bd^3(d \cos(a + bx))^{11/2}} - \frac{16c(c \sin(a + bx))^{3/2}}{385bd^5(d \cos(a + bx))^{7/2}} \\ &= \frac{2c(c \sin(a + bx))^{3/2}}{15bd(d \cos(a + bx))^{15/2}} - \frac{2c(c \sin(a + bx))^{3/2}}{55bd^3(d \cos(a + bx))^{11/2}} - \frac{16c(c \sin(a + bx))^{3/2}}{385bd^5(d \cos(a + bx))^{7/2}} \end{aligned}$$

Mathematica [A]

time = 0.32, size = 67, normalized size = 0.48

$$\frac{2\sqrt{d \cos(a + bx)} (117 + 44 \cos(2(a + bx)) + 4 \cos(4(a + bx))) \sec^8(a + bx) (c \sin(a + bx))^{7/2}}{1155bcd^9}$$

Antiderivative was successfully verified.

[In] Integrate[(c*SIN[a + b*x])^(5/2)/(d*Cos[a + b*x])^(17/2), x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(117 + 44*Cos[2*(a + b*x)] + 4*Cos[4*(a + b*x)])*Sec[a + b*x]^8*(c*SIN[a + b*x])^(7/2))/(1155*b*c*d^9)

Maple [A]

time = 0.27, size = 60, normalized size = 0.43

method	result	size
default	$\frac{2(32(\cos^4(bx+a)) + 56(\cos^2(bx+a)) + 77) \cos(bx+a) (c \sin(bx+a))^{\frac{5}{2}} \sin(bx+a)}{1155b(d \cos(bx+a))^{\frac{17}{2}}}$	60

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(17/2), x, method=_RETURNVERBOSE)

[Out] 2/1155/b*(32*cos(b*x+a)^4+56*cos(b*x+a)^2+77)*cos(b*x+a)*(c*sin(b*x+a))^(5/2)*sin(b*x+a)/(d*cos(b*x+a))^(17/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(17/2),x, algorithm="maxima")
```

```
[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(17/2), x)
```

Fricas [A]

time = 0.62, size = 87, normalized size = 0.62

$$\frac{2(32c^2 \cos(bx+a)^6 + 24c^2 \cos(bx+a)^4 + 21c^2 \cos(bx+a)^2 - 77c^2) \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)} \sin(bx+a)}{1155bd^9 \cos(bx+a)^8}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(17/2),x, algorithm="fricas")
```

```
[Out] -2/1155*(32*c^2*cos(b*x + a)^6 + 24*c^2*cos(b*x + a)^4 + 21*c^2*cos(b*x + a)^2 - 77*c^2)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*sin(b*x + a)/(b*d^9*cos(b*x + a)^8)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))**(5/2)/(d*cos(b*x+a))**(17/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((c*sin(b*x+a))^(5/2)/(d*cos(b*x+a))^(17/2),x, algorithm="giac")
```

```
[Out] integrate((c*sin(b*x + a))^(5/2)/(d*cos(b*x + a))^(17/2), x)
```

Mupad [B]

time = 6.43, size = 207, normalized size = 1.47

$$\frac{e^{-a7i-bx7i} \sqrt{c \left(\frac{e^{-a1i-bx1i} 1i}{2} - \frac{e^{a1i+bx1i} 1i}{2} \right)} \left(\frac{1216c^2 e^{a7i+bx7i} \sin(3a+3bx)}{385bd^8} + \frac{1024c^2 e^{a7i+bx7i} \sin(5a+5bx)}{1155bd^8} + \frac{128c^2 e^{a7i+bx7i} \sin(7a+7bx)}{1155bd^8} - \frac{3392c^2 e^{a7i+bx7i} \sin(a+bx)}{231bd^8} \right)}{128 \cos(a+bx)^7 \sqrt{d \left(\frac{e^{-a1i-bx1i}}{2} + \frac{e^{a1i+bx1i}}{2} \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((c*\sin(a + b*x))^{5/2}/(d*\cos(a + b*x))^{17/2},x)$

[Out] $-(\exp(-a*7i - b*x*7i)*(c*((\exp(-a*1i - b*x*1i)*1i)/2 - (\exp(a*1i + b*x*1i)*1i)/2))^{1/2}*((1216*c^2*\exp(a*7i + b*x*7i)*\sin(3*a + 3*b*x))/(385*b*d^8) + (1024*c^2*\exp(a*7i + b*x*7i)*\sin(5*a + 5*b*x))/(1155*b*d^8) + (128*c^2*\exp(a*7i + b*x*7i)*\sin(7*a + 7*b*x))/(1155*b*d^8) - (3392*c^2*\exp(a*7i + b*x*7i)*\sin(a + b*x))/(231*b*d^8))/((128*\cos(a + b*x)^7*(d*(\exp(-a*1i - b*x*1i)/2 + \exp(a*1i + b*x*1i)/2))^{1/2}))$

$$3.287 \quad \int \frac{\sin^{\frac{7}{2}}(a+bx)}{\cos^{\frac{7}{2}}(a+bx)} dx$$

Optimal. Leaf size=226

$$\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\log\left(1 + \cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b}$$

[Out] $2/5*\sin(b*x+a)^{(5/2)}/b/\cos(b*x+a)^{(5/2)}-1/2*\arctan(-1+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}-1/2*\arctan(1+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}-1/4*\ln(1+\cot(b*x+a)-2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}+1/4*\ln(1+\cot(b*x+a)+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}-2*\sin(b*x+a)^{(1/2)}/b/\cos(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.10, antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2646, 2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\text{ArcTan}\left(\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{\sqrt{2}b} + \frac{2\sin^{\frac{5}{2}}(a+bx)}{5b\cos^{\frac{5}{2}}(a+bx)} - \frac{2\sqrt{\sin(a+bx)}}{b\sqrt{\cos(a+bx)}} - \frac{\log\left(\cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{2\sqrt{2}b} + \frac{\log\left(\cot(a+bx) + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(7/2)/Cos[a + b*x]^(7/2), x]

[Out] ArcTan[1 - (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(Sqrt[2]*b) - ArcTan[1 + (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(Sqrt[2]*b) - Log[1 + Cot[a + b*x] - (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(2*Sqrt[2]*b) + Log[1 + Cot[a + b*x] + (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(2*Sqrt[2]*b) - (2*Sqrt[Sin[a + b*x]])/(b*Sqrt[Cos[a + b*x]]) + (2*Sin[a + b*x]^(5/2))/(5*b*Cos[a + b*x]^(5/2))

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2646

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*Sine[e + f*x])^(m - 1)*(b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1)), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sine[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])
```

Rule 2655

```
Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sine[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^{\frac{7}{2}}(a+bx)}{\cos^{\frac{7}{2}}(a+bx)} dx &= \frac{2 \sin^{\frac{5}{2}}(a+bx)}{5b \cos^{\frac{5}{2}}(a+bx)} - \int \frac{\sin^{\frac{3}{2}}(a+bx)}{\cos^{\frac{3}{2}}(a+bx)} dx \\
&= -\frac{2\sqrt{\sin(a+bx)}}{b\sqrt{\cos(a+bx)}} + \frac{2 \sin^{\frac{5}{2}}(a+bx)}{5b \cos^{\frac{5}{2}}(a+bx)} + \int \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} dx \\
&= -\frac{2\sqrt{\sin(a+bx)}}{b\sqrt{\cos(a+bx)}} + \frac{2 \sin^{\frac{5}{2}}(a+bx)}{5b \cos^{\frac{5}{2}}(a+bx)} - \frac{2 \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} \\
&= -\frac{2\sqrt{\sin(a+bx)}}{b\sqrt{\cos(a+bx)}} + \frac{2 \sin^{\frac{5}{2}}(a+bx)}{5b \cos^{\frac{5}{2}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} - \text{Subst} \\
&= -\frac{2\sqrt{\sin(a+bx)}}{b\sqrt{\cos(a+bx)}} + \frac{2 \sin^{\frac{5}{2}}(a+bx)}{5b \cos^{\frac{5}{2}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2b} \\
&= -\frac{\log\left(1 + \cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} + \frac{\log\left(1 + \cot(a+bx) + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} \\
&= \frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\log\left(1 + \cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} + \frac{\log\left(1 + \cot(a+bx) + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 57, normalized size = 0.25

$$\frac{2^4 \sqrt{\cos^2(a+bx)} {}_2F_1\left(\frac{9}{4}, \frac{9}{4}; \frac{13}{4}; \sin^2(a+bx)\right) \sin^{\frac{9}{2}}(a+bx)}{9b \sqrt{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^(7/2)/Cos[a + b*x]^(7/2),x]

[Out] (2*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[9/4, 9/4, 13/4, Sin[a + b*x]^2]*Sin[a + b*x]^(9/2))/(9*b*Sqrt[Cos[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.84, size = 692, normalized size = 3.06

method	result
--------	--------

default	$-\frac{5i(\cos^2(bx+a)) \sin(bx+a) \operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{1}{2}-\frac{i}{2}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}}{\dots}$
---------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^(7/2)/cos(b*x+a)^(7/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/10/b*(5*I*\cos(b*x+a)^2*\sin(b*x+a)*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2-1/2*I, 1/2*2^{1/2}))*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}-5*I*\cos(b*x+a)^2*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2+1/2*I, 1/2*2^{1/2}))+5*\cos(b*x+a)^2*\sin(b*x+a)*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2-1/2*I, 1/2*2^{1/2}))*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}+5*\cos(b*x+a)^2*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\operatorname{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2+1/2*I, 1/2*2^{1/2}))-10*\cos(b*x+a)^2*\sin(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*\operatorname{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}, 1/2*2^{1/2}))+12*\cos(b*x+a)^3*2^{1/2}-12*\cos(b*x+a)^2*2^{1/2}-2*\cos(b*x+a)*2^{1/2}+2*2^{1/2})*\sin(b*x+a)^{1/2}/(-1+\cos(b*x+a))/\cos(b*x+a)^{5/2}*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(7/2)/cos(b*x+a)^(7/2),x, algorithm="maxima")`

[Out] `integrate(sin(b*x + a)^(7/2)/cos(b*x + a)^(7/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1282 vs. 2(180) = 360.

time = 14.97, size = 1282, normalized size = 5.67

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(7/2)/cos(b*x+a)^(7/2),x, algorithm="fricas")`

```
[Out] -1/40*(10*sqrt(2)*b*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*
sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(4*b^2*sqrt(b^(-4)
))*cos(b*x + a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) +
sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a
)) + 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + (sqrt(2)*b^3*(b^(-4))^(3/4)
*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*s
qrt(sin(b*x + a)) + 2*cos(b*x + a)*sin(b*x + a) - 4*(b^2*cos(b*x + a)^4 - b
^2*cos(b*x + a)^2)*sqrt(b^(-4)))/((2*cos(b*x + a)^3 - cos(b*x + a))*sin(b*x
+ a))*cos(b*x + a)^3 + 10*sqrt(2)*b*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b
^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sq
rt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3
/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a)
))*sqrt(sin(b*x + a)) + 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + (sqrt(2)*
b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sq
rt(cos(b*x + a))*sqrt(sin(b*x + a)) - 2*cos(b*x + a)*sin(b*x + a) + 4*(b^2*
cos(b*x + a)^4 - b^2*cos(b*x + a)^2)*sqrt(b^(-4)))/((2*cos(b*x + a)^3 - cos
(b*x + a))*sin(b*x + a))*cos(b*x + a)^3 + 10*sqrt(2)*b*(b^(-4))^(1/4)*arct
an(-1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) - sqrt(2)*b*(b^(-4))^(1/4)
)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) - sqrt(4*b^2*sqrt(b^(-
4))*cos(b*x + a)*sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a)
+ sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x +
a)) + 1)*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1
/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 2*cos(b*x + a)*si
n(b*x + a)))/(cos(b*x + a)*sin(b*x + a))*cos(b*x + a)^3 + 10*sqrt(2)*b*(b^
(-4))^(1/4)*arctan(-1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) - sqrt(2)
)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) - sqr
t(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3
/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a)
))*sqrt(sin(b*x + a)) + 1)*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(
2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) - 2
*cos(b*x + a)*sin(b*x + a)))/(cos(b*x + a)*sin(b*x + a))*cos(b*x + a)^3 -
5*sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a)^3*log(4*b^2*sqrt(b^(-4))*cos(b*x +
a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b
^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1) + 5*sq
rt(2)*b*(b^(-4))^(1/4)*cos(b*x + a)^3*log(4*b^2*sqrt(b^(-4))*cos(b*x + a)*
sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-
4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1) + 16*(6*
cos(b*x + a)^2 - 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)))/(b*cos(b*x + a)^
3)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**(7/2)/cos(b*x+a)**(7/2),x)

[Out] Timed out

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^(7/2)/cos(b*x+a)^(7/2),x, algorithm="giac")

[Out] Timed out

Mupad [B]

time = 2.05, size = 44, normalized size = 0.19

$$\frac{2 \sin(a + bx)^{9/2} {}_2F_1\left(-\frac{5}{4}, -\frac{5}{4}; -\frac{1}{4}; \cos(a + bx)^2\right)}{5b \cos(a + bx)^{5/2} (\sin(a + bx)^2)^{9/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^(7/2)/cos(a + b*x)^(7/2),x)

[Out] (2*sin(a + b*x)^(9/2)*hypergeom([-5/4, -5/4], -1/4, cos(a + b*x)^2))/(5*b*cos(a + b*x)^(5/2)*(sin(a + b*x)^2)^(9/4))

$$3.288 \quad \int \frac{\sin^{\frac{3}{2}}(x)}{\cos^{\frac{7}{2}}(x)} dx$$

Optimal. Leaf size=16

$$\frac{2 \sin^{\frac{5}{2}}(x)}{5 \cos^{\frac{5}{2}}(x)}$$

[Out] $2/5*\sin(x)^{(5/2)}/\cos(x)^{(5/2)}$

Rubi [A]

time = 0.02, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2643}

$$\frac{2 \sin^{\frac{5}{2}}(x)}{5 \cos^{\frac{5}{2}}(x)}$$

Antiderivative was successfully verified.

[In] `Int[Sin[x]^(3/2)/Cos[x]^(7/2),x]`

[Out] `(2*Sin[x]^(5/2))/(5*Cos[x]^(5/2))`

Rule 2643

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] & & NeQ[m, -1]`

Rubi steps

$$\int \frac{\sin^{\frac{3}{2}}(x)}{\cos^{\frac{7}{2}}(x)} dx = \frac{2 \sin^{\frac{5}{2}}(x)}{5 \cos^{\frac{5}{2}}(x)}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$\frac{2 \sin^{\frac{5}{2}}(x)}{5 \cos^{\frac{5}{2}}(x)}$$

Antiderivative was successfully verified.

[In] `Integrate[Sin[x]^(3/2)/Cos[x]^(7/2),x]`

[Out] $(2*\sin[x]^{(5/2)})/(5*\cos[x]^{(5/2)})$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 32 vs. $2(10) = 20$.

time = 0.13, size = 33, normalized size = 2.06

method	result	size
default	$-\frac{(\sin^2(x) - \cos^2(x) + 2\cos(x) - 1) \sin^{\frac{5}{2}}(x)}{5(-1 + \cos(x)) \cos(x)^{\frac{7}{2}}}$	33

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(x)^(3/2)/cos(x)^(7/2),x,method=_RETURNVERBOSE)`

[Out] $-1/5*(\sin(x)^2 - \cos(x)^2 + 2*\cos(x) - 1)*\sin(x)^{(5/2)/(-1 + \cos(x))}/\cos(x)^{(7/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^(3/2)/cos(x)^(7/2),x, algorithm="maxima")`

[Out] `integrate(sin(x)^(3/2)/cos(x)^(7/2), x)`

Fricas [A]

time = 0.35, size = 16, normalized size = 1.00

$$\frac{2(\cos(x)^2 - 1)\sqrt{\sin(x)}}{5\cos(x)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^(3/2)/cos(x)^(7/2),x, algorithm="fricas")`

[Out] $-2/5*(\cos(x)^2 - 1)*\sqrt{\sin(x)}/\cos(x)^{(5/2)}$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)**(3/2)/cos(x)**(7/2),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(x)^(3/2)/cos(x)^(7/2),x, algorithm="giac")``[Out] integrate(sin(x)^(3/2)/cos(x)^(7/2), x)`**Mupad [B]**

time = 0.86, size = 53, normalized size = 3.31

$$\frac{8\sqrt{2}\tan\left(\frac{x}{2}\right)^{5/2}\sqrt{1-\tan\left(\frac{x}{2}\right)^2}}{\tan\left(\frac{x}{2}\right)^2\left(\tan\left(\frac{x}{2}\right)^2\left(5\tan\left(\frac{x}{2}\right)^2-15\right)+15\right)-5}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(x)^(3/2)/cos(x)^(7/2),x)``[Out] -(8*2^(1/2)*tan(x/2)^(5/2)*(1 - tan(x/2)^2)^(1/2))/(tan(x/2)^2*(tan(x/2)^2*(5*tan(x/2)^2 - 15) + 15) - 5)`

$$3.289 \quad \int \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} dx$$

Optimal. Leaf size=122

$$-\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}}\right)}{\sqrt{2}} + \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}}\right)}{\sqrt{2}} + \frac{\log\left(1 - \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x)\right)}{2\sqrt{2}} - \frac{\log\left(1 + \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x)\right)}{2\sqrt{2}}$$

[Out] $-1/2*\arctan(1-2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)})*2^{(1/2)}+1/2*\arctan(1+2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)})*2^{(1/2)}+1/4*\ln(1-2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)}+\tan(x))*2^{(1/2)}-1/4*\ln(1+2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)}+\tan(x))*2^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.538$, Rules used = {2654, 303, 1176, 631, 210, 1179, 642}

$$-\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}}\right)}{\sqrt{2}} + \frac{\text{ArcTan}\left(\frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}} + 1\right)}{\sqrt{2}} + \frac{\log\left(\tan(x) - \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}} + 1\right)}{2\sqrt{2}} - \frac{\log\left(\tan(x) + \frac{\sqrt{2}\sqrt{\sin(x)}}{\sqrt{\cos(x)}} + 1\right)}{2\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sin[x]]/Sqrt[Cos[x]], x]

[Out] $-(\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]]]/\text{Sqrt}[2]) + \text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]]]/\text{Sqrt}[2] + \text{Log}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]] + \text{Tan}[x]]/(2*\text{Sqrt}[2]) - \text{Log}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]] + \text{Tan}[x]]/(2*\text{Sqrt}[2])$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} dx &= 2 \operatorname{Subst} \left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= -\operatorname{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) + \operatorname{Subst} \left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= \frac{1}{2} \operatorname{Subst} \left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) + \frac{1}{2} \operatorname{Subst} \left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= \frac{\log \left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x) \right)}{2\sqrt{2}} - \frac{\log \left(1 + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x) \right)}{2\sqrt{2}} + \operatorname{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= -\frac{\tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right)}{\sqrt{2}} + \frac{\tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right)}{\sqrt{2}} + \frac{\log \left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right)}{2\sqrt{2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 38, normalized size = 0.31

$$\frac{2 \cos^2(x)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{3}{4}, \frac{7}{4}; \sin^2(x)\right) \sin^{3/2}(x)}{3 \cos^{3/2}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sin[x]]/Sqrt[Cos[x]],x]

[Out] (2*(Cos[x]^2)^(3/4)*Hypergeometric2F1[3/4, 3/4, 7/4, Sin[x]^2]*Sin[x]^(3/2))/(3*Cos[x]^(3/2))

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.11, size = 166, normalized size = 1.36

method	result
default	$ -\frac{\left(\sin^{3/2}(x)\right) \left(i \operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(x)+\sin(x)}{\sin(x)}}, \frac{1}{2}-\frac{i}{2}, \frac{\sqrt{2}}{2}\right) - i \operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(x)+\sin(x)}{\sin(x)}}, \frac{1}{2}+\frac{i}{2}, \frac{\sqrt{2}}{2}\right) - \operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(x)+\sin(x)}{\sin(x)}}, \frac{1}{2}, \frac{\sqrt{2}}{2}\right)\right)}{3 \cos^{3/2}(x)} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^(1/2)/cos(x)^(1/2),x,method=_RETURNVERBOSE)

```
[Out] -1/2*sin(x)^(3/2)*(I*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,
1/2*2^(1/2))-I*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(
1/2))-EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2))-E
llipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2)))*((-1+cos
(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((1-cos(x)+sin(x))/sin
(x))^(1/2)/(-1+cos(x))/cos(x)^(1/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(x)^(1/2)/cos(x)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(sin(x))/sqrt(cos(x)), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 447 vs. 2(87) = 174.

time = 0.67, size = 447, normalized size = 3.66

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(x)^(1/2)/cos(x)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/4*sqrt(2)*arctan(1/2*(2*cos(x)^3 - 2*cos(x)^2*sin(x) + sqrt(2)*sqrt(2*(sq
rt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x)
+ 1)*sqrt(cos(x))*sqrt(sin(x)) - sqrt(2)*sqrt(cos(x))*sqrt(sin(x)) - 2*cos(
x))/(cos(x)^3 + cos(x)^2*sin(x) - cos(x))) + 1/4*sqrt(2)*arctan(-1/2*(2*cos
(x)^3 - 2*cos(x)^2*sin(x) - sqrt(2)*sqrt(-2*(sqrt(2)*cos(x) + sqrt(2)*sin(
x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)*sqrt(cos(x))*sqrt(sin(
x)) + sqrt(2)*sqrt(cos(x))*sqrt(sin(x)) - 2*cos(x))/(cos(x)^3 + cos(x)^2*sin
(x) - cos(x))) - 1/4*sqrt(2)*arctan(-(sqrt(-2*(sqrt(2)*cos(x) + sqrt(2)*sin
(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)*(sqrt(2)*sqrt(cos(x))
*sqrt(sin(x)) + cos(x) + sin(x)) + sqrt(2)*sqrt(cos(x))*sqrt(sin(x)))/(cos(
x) - sin(x))) - 1/4*sqrt(2)*arctan(-(sqrt(2*(sqrt(2)*cos(x) + sqrt(2)*sin(x)
))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)*(sqrt(2)*sqrt(cos(x))*s
qrt(sin(x)) - cos(x) - sin(x)) + sqrt(2)*sqrt(cos(x))*sqrt(sin(x)))/(cos(x)
- sin(x))) - 1/8*sqrt(2)*log(2*(sqrt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(
x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1) + 1/8*sqrt(2)*log(-2*(sqrt(2)*cos(x)
) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)**(1/2)/cos(x)**(1/2),x)`

[Out] `Integral(sqrt(sin(x))/sqrt(cos(x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^(1/2)/cos(x)^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(sin(x))/sqrt(cos(x)), x)`

Mupad [B]

time = 0.72, size = 25, normalized size = 0.20

$$-\frac{2\sqrt{\cos(x)}\sin(x)^{3/2}{}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \cos(x)^2\right)}{(\sin(x)^2)^{3/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(x)^(1/2)/cos(x)^(1/2),x)`

[Out] `-(2*cos(x)^(1/2)*sin(x)^(3/2)*hypergeom([1/4, 1/4], 5/4, cos(x)^2))/(sin(x)^2)^(3/4)`

$$3.290 \quad \int \frac{\sin^5(x)}{\sqrt{\cos(x)}} dx$$

Optimal. Leaf size=143

$$-\frac{3 \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}}\right)}{4\sqrt{2}} + \frac{3 \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}}\right)}{4\sqrt{2}} + \frac{3 \log\left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x)\right)}{8\sqrt{2}} - \frac{3 \log\left(1 + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x)\right)}{8\sqrt{2}}$$

[Out] $-3/8*\arctan(1-2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)})*2^{(1/2)}+3/8*\arctan(1+2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)})*2^{(1/2)}+3/16*\ln(1-2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)}+\tan(x))*2^{(1/2)}-3/16*\ln(1+2^{(1/2)}*\sin(x)^{(1/2)}/\cos(x)^{(1/2)}+\tan(x))*2^{(1/2)}-1/2*\sin(x)^{(3/2)}*\cos(x)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 143, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.615$, Rules used = {2648, 2654, 303, 1176, 631, 210, 1179, 642}

$$-\frac{3\text{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}}\right)}{4\sqrt{2}} + \frac{3\text{ArcTan}\left(\frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + 1\right)}{4\sqrt{2}} - \frac{1}{2} \sin^{\frac{3}{2}}(x) \sqrt{\cos(x)} + \frac{3 \log\left(\tan(x) - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + 1\right)}{8\sqrt{2}} - \frac{3 \log\left(\tan(x) + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + 1\right)}{8\sqrt{2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^(5/2)/Sqrt[Cos[x]],x]

[Out] $(-3*\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]])/(4*\text{Sqrt}[2]) + (3*\text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]])/(4*\text{Sqrt}[2]) + (3*\text{Log}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]] + \text{Tan}[x])/(8*\text{Sqrt}[2]) - (3*\text{Log}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[x]])/\text{Sqrt}[\text{Cos}[x]] + \text{Tan}[x])/(8*\text{Sqrt}[2]) - (\text{Sqrt}[\text{Cos}[x]]*\text{Sin}[x]^{(3/2)})/2)$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2648

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(b*Cos[e + f*x])^(n + 1)*((a*SIN[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Cos[e + f*x])^n*(a*SIN[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*SIN[e + f*x])^(1/k)/(b*COS[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^{\frac{5}{2}}(x)}{\sqrt{\cos(x)}} dx &= -\frac{1}{2} \sqrt{\cos(x)} \sin^{\frac{3}{2}}(x) + \frac{3}{4} \int \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} dx \\
&= -\frac{1}{2} \sqrt{\cos(x)} \sin^{\frac{3}{2}}(x) + \frac{3}{2} \text{Subst} \left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= -\frac{1}{2} \sqrt{\cos(x)} \sin^{\frac{3}{2}}(x) - \frac{3}{4} \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) + \frac{3}{4} \text{Subst} \left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= -\frac{1}{2} \sqrt{\cos(x)} \sin^{\frac{3}{2}}(x) + \frac{3}{8} \text{Subst} \left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) + \frac{3}{8} \text{Subst} \left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right) \\
&= \frac{3 \log \left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x) \right)}{8\sqrt{2}} - \frac{3 \log \left(1 + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} + \tan(x) \right)}{8\sqrt{2}} - \frac{1}{2} \sqrt{\cos(x)} \sin^{\frac{3}{2}}(x) \\
&= -\frac{3 \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right)}{4\sqrt{2}} + \frac{3 \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right)}{4\sqrt{2}} + \frac{3 \log \left(1 - \frac{\sqrt{2} \sqrt{\sin(x)}}{\sqrt{\cos(x)}} \right)}{8\sqrt{2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 38, normalized size = 0.27

$$\frac{2 \cos^2(x)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; \sin^2(x)\right) \sin^{7/2}(x)}{7 \cos^{3/2}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^(5/2)/Sqrt[Cos[x]],x]

[Out] (2*(Cos[x]^2)^(3/4)*Hypergeometric2F1[3/4, 7/4, 11/4, Sin[x]^2]*Sin[x]^(7/2))/(7*Cos[x]^(3/2))

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.12, size = 2595, normalized size = 18.15

method	result	size
default	Expression too large to display	2595

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^(5/2)/cos(x)^(1/2),x,method=_RETURNVERBOSE)


```

[Out] 1/32*sin(x)^(3/2)*(3*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/s
in(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x)
))^(1/2),1/2-1/2*I,1/2*2^(1/2))+3*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos
(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+
sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))+6*((1-cos(x)+sin(x))/sin(x))^(
1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticP
i(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2))*sin(x)^2*cos(x)^2
+6*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+
cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I
,1/2*2^(1/2))*sin(x)^2*cos(x)^2-12*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+co
s(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)
+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2))*sin(x)^2*cos(x)-12*((1-cos(x)
+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x
))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2)
)*sin(x)^2*cos(x)-3*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/s
in(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x)
))^(1/2),1/2-1/2*I,1/2*2^(1/2))*sin(x)^4+3*I*((1-cos(x)+sin(x))/sin(x))^(1/
2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(
((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(x)^4-6*I*((1-co
s(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/s
in(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(
1/2))*sin(x)^2-3*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(
x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(
1/2),1/2-1/2*I,1/2*2^(1/2))*cos(x)^4+6*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*
((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1
-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(x)^2+3*I*((1-cos(x)
)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(
x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2)
)*cos(x)^4+12*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x)
)^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1
/2),1/2-1/2*I,1/2*2^(1/2))*cos(x)^3-12*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*
((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-
cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*cos(x)^3-18*I*((1-cos(x)
)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(
x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2)
)*cos(x)^2+18*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x)
)^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1
/2),1/2+1/2*I,1/2*2^(1/2))*cos(x)^2+12*I*((1-cos(x)+sin(x))/sin(x))^(1/2)*
((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-
cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2))*cos(x)-12*I*((1-cos(x)+
sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x)
)^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*
cos(x)+4*2^(1/2)*sin(x)^2*cos(x)^2-8*2^(1/2)*sin(x)^2*cos(x)+4*2^(1/2)*sin(
x)^2-4*2^(1/2)*cos(x)^4+16*2^(1/2)*cos(x)^3-24*2^(1/2)*cos(x)^2+16*2^(1/2)*
cos(x)+3*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)

```

```

*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2
-1/2*I,1/2*2^(1/2))*sin(x)^4+3*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)
+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin
(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(x)^4+6*((1-cos(x)+sin(x))/sin
(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*Ell
ipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2))*sin(x)^2+3*
((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos
(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/
2*2^(1/2))*cos(x)^4+6*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/
sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(
x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*sin(x)^2+3*((1-cos(x)+sin(x))/sin(x))^(1/2
)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi((
(1-cos(x)+sin(x))/sin(x))^(1/2),1/2+1/2*I,1/2*2^(1/2))*cos(x)^4-12*((1-cos(
x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin
(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2
))*cos(x)^3-12*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))
^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/
2),1/2+1/2*I,1/2*2^(1/2))*cos(x)^3+18*((1-cos(x)+sin(x))/sin(x))^(1/2)*((-1
+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(1/2)*EllipticPi(((1-cos
(x)+sin(x))/sin(x))^(1/2),1/2-1/2*I,1/2*2^(1/2))*cos(x)^2+18*((1-cos(x)+sin
(x))/sin(x))^(1/2)*((-1+cos(x)+sin(x))/sin(x))^(1/2)*((-1+cos(x))/sin(x))^(
1/2)*EllipticPi(((1-cos(x)+sin(x))/sin(x))^(1/2)...

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^(5/2)/cos(x)^(1/2),x, algorithm="maxima")

[Out] integrate(sin(x)^(5/2)/sqrt(cos(x)), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 457 vs. 2(97) = 194.

time = 0.66, size = 457, normalized size = 3.20

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^(5/2)/cos(x)^(1/2),x, algorithm="fricas")

[Out]
$$-1/2*\sqrt{\cos(x)}*\sin(x)^{3/2} + 3/16*\sqrt{2}*\arctan(1/2*(2*\cos(x)^3 - 2*\cos(x)^2*\sin(x) + \sqrt{2}*\sqrt{2*(\sqrt{2}*\cos(x) + \sqrt{2}*\sin(x))*\sqrt{\cos(x)}}*\sqrt{\sin(x)} + 4*\cos(x)*\sin(x) + 1)*\sqrt{\cos(x)}*\sqrt{\sin(x)} - \sqrt{2}*\sqrt{\cos(x)}*\sqrt{\sin(x)} - 2*\cos(x))/(\cos(x)^3 + \cos(x)^2*\sin(x) - \cos(x))$$

) + 3/16*sqrt(2)*arctan(-1/2*(2*cos(x)^3 - 2*cos(x)^2*sin(x) - sqrt(2)*sqrt(-2*(sqrt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)*sqrt(cos(x))*sqrt(sin(x)) + sqrt(2)*sqrt(cos(x))*sqrt(sin(x)) - 2*cos(x))/(cos(x)^3 + cos(x)^2*sin(x) - cos(x))) - 3/16*sqrt(2)*arctan(-(sqrt(-2*(sqrt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)*(sqrt(2)*sqrt(cos(x))*sqrt(sin(x)) + cos(x) + sin(x)) + sqrt(2)*sqrt(cos(x))*sqrt(sin(x)))/(cos(x) - sin(x))) - 3/16*sqrt(2)*arctan(-(sqrt(2*(sqrt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)*(sqrt(2)*sqrt(cos(x))*sqrt(sin(x)) - cos(x) - sin(x)) + sqrt(2)*sqrt(cos(x))*sqrt(sin(x)))/(cos(x) - sin(x))) - 3/32*sqrt(2)*log(2*(sqrt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1) + 3/32*sqrt(2)*log(-2*(sqrt(2)*cos(x) + sqrt(2)*sin(x))*sqrt(cos(x))*sqrt(sin(x)) + 4*cos(x)*sin(x) + 1)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)**(5/2)/cos(x)**(1/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^(5/2)/cos(x)^(1/2),x, algorithm="giac")

[Out] integrate(sin(x)^(5/2)/sqrt(cos(x)), x)

Mupad [B]

time = 0.81, size = 25, normalized size = 0.17

$$-\frac{2\sqrt{\cos(x)}\sin(x)^{7/2}{}_2F_1\left(-\frac{3}{4}, \frac{1}{4}; \frac{5}{4}; \cos(x)^2\right)}{(\sin(x)^2)^{7/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^(5/2)/cos(x)^(1/2),x)

[Out] -(2*cos(x)^(1/2)*sin(x)^(7/2)*hypergeom([-3/4, 1/4], 5/4, cos(x)^2))/(sin(x)^2)^(7/4)

$$3.291 \quad \int \frac{(d \cos(a+bx))^{7/2}}{\sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=132

$$\frac{5d^3 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}{6bc} + \frac{d(d \cos(a+bx))^{5/2} \sqrt{c \sin(a+bx)}}{3bc} + \frac{5d^4 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a+2bx)}}{12b \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] 1/3*d*(d*cos(b*x+a))^(5/2)*(c*sin(b*x+a))^(1/2)/b/c+5/6*d^3*(d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2)/b/c-5/12*d^4*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*sin(2*b*x+2*a)^(1/2)/b/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2)

Rubi [A]

time = 0.11, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2649, 2653, 2720}

$$\frac{5d^4 \sqrt{\sin(2a+2bx)} F\left(a+bx-\frac{\pi}{4} \mid 2\right)}{12b \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}} + \frac{5d^3 \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}}{6bc} + \frac{d \sqrt{c \sin(a+bx)} (d \cos(a+bx))^{5/2}}{3bc}$$

Antiderivative was successfully verified.

[In] Int[(d*cos[a + b*x])^(7/2)/Sqrt[c*Sin[a + b*x]],x]

[Out] (5*d^3*Sqrt[d*cos[a + b*x]]*Sqrt[c*Sin[a + b*x]])/(6*b*c) + (d*(d*cos[a + b*x])^(5/2)*Sqrt[c*Sin[a + b*x]])/(3*b*c) + (5*d^4*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[Sin[2*a + 2*b*x]])/(12*b*Sqrt[d*cos[a + b*x]]*Sqrt[c*Sin[a + b*x]])

Rule 2649

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a*(b*Sin[e + f*x])^(n + 1)*((a*cos[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*Sin[e + f*x])^n*(a*cos[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{(d \cos(a + bx))^{7/2}}{\sqrt{c \sin(a + bx)}} dx &= \frac{d(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bc} + \frac{1}{6} (5d^2) \int \frac{(d \cos(a + bx))^{3/2}}{\sqrt{c \sin(a + bx)}} dx \\ &= \frac{5d^3 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{6bc} + \frac{d(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bc} + \frac{1}{12} \\ &= \frac{5d^3 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{6bc} + \frac{d(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bc} + \frac{(5}{12} \\ &= \frac{5d^3 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{6bc} + \frac{d(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}}{3bc} + \frac{5d}{12} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.08, size = 70, normalized size = 0.53

$$\frac{2(d \cos(a + bx))^{7/2} \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{5}{4}, \frac{1}{4}; \frac{5}{4}; \sin^2(a + bx)\right) \sec^5(a + bx) \sqrt{c \sin(a + bx)}}{bc}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(7/2)/Sqrt[c*Sin[a + b*x]],x]

[Out] (2*(d*Cos[a + b*x])^(7/2)*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-5/4, 1/4, 5/4, Sin[a + b*x]^2]*Sec[a + b*x]^5*Sqrt[c*Sin[a + b*x]])/(b*c)

Maple [A]

time = 0.20, size = 216, normalized size = 1.64

method	result
default	$\frac{\left(2(\cos^4(bx+a))\sqrt{2} - 5\sin(bx+a)\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}}\right) \text{EllipticF}\left(\frac{1}{2}, \frac{c-d\cos(bx+a)}{c}, 2\right)}{12b(-1+\cos(bx+a))\cos(bx+a)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/12/b*(2*cos(b*x+a)^4*2^(1/2)-5*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)

$$\frac{1}{\sin(b*x+a)^{1/2}} * \text{EllipticF}\left(\frac{(1-\cos(b*x+a)+\sin(b*x+a))}{\sin(b*x+a)}\right)^{1/2}, \\ \frac{1}{2} * 2^{1/2} - 2 * \cos(b*x+a)^3 * 2^{1/2} + 5 * \cos(b*x+a)^2 * 2^{1/2} - 5 * \cos(b*x+a) * 2^{1/2} \\ * (d * \cos(b*x+a))^{7/2} * \sin(b*x+a) / (-1 + \cos(b*x+a)) / \cos(b*x+a)^4 / (c * \sin(b \\ *x+a))^{1/2} * 2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^(7/2)/sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*d^3*cos(b*x + a)^3/(c*sin(b*x + a)), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(7/2)/(c*sin(b*x+a))**(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(7/2)/sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^{7/2}}{\sqrt{c \sin(a + b x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(7/2)/(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^(7/2)/(c*sin(a + b*x))^(1/2), x)

$$3.292 \quad \int \frac{(d \cos(a+bx))^{3/2}}{\sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=92

$$\frac{d \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}{bc} + \frac{d^2 F(a - \frac{\pi}{4} + bx | 2) \sqrt{\sin(2a + 2bx)}}{2b \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] d*(d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^(1/2)/b/c-1/2*d^2*(sin(a+1/4*Pi+b*x)^2)^(1/2)/sin(a+1/4*Pi+b*x)*EllipticF(cos(a+1/4*Pi+b*x),2^(1/2))*sin(2*b*x+2*a)^(1/2)/b/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2649, 2653, 2720}

$$\frac{d^2 \sqrt{\sin(2a + 2bx)} F(a + bx - \frac{\pi}{4} | 2)}{2b \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}} + \frac{d \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}{bc}$$

Antiderivative was successfully verified.

[In] Int[(d*cos[a + b*x])^(3/2)/Sqrt[c*sin[a + b*x]],x]

[Out] (d*Sqrt[d*cos[a + b*x]]*Sqrt[c*sin[a + b*x]])/(b*c) + (d^2*EllipticF[a - Pi/4 + b*x, 2]*Sqrt[Sin[2*a + 2*b*x]])/(2*b*Sqrt[d*cos[a + b*x]]*Sqrt[c*sin[a + b*x]])

Rule 2649

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[a*(b*sin[e + f*x])^(n + 1)*((a*cos[e + f*x])^(m - 1)/(b*f*(m + n))), x] + Dist[a^2*((m - 1)/(m + n)), Int[(b*sin[e + f*x])^n*(a*cos[e + f*x])^(m - 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*sin[e + f*x]]*Sqrt[b*cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
\int \frac{(d \cos(a + bx))^{3/2}}{\sqrt{c \sin(a + bx)}} dx &= \frac{d \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{bc} + \frac{1}{2} d^2 \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx \\
&= \frac{d \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{bc} + \frac{\left(d^2 \sqrt{\sin(2a + 2bx)} \right) \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx}{2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \\
&= \frac{d \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}{bc} + \frac{d^2 F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{2b \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.08, size = 68, normalized size = 0.74

$$\frac{2d^2 \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \sin^2(a + bx)\right) \tan(a + bx)}{b \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^(3/2)/Sqrt[c*Sin[a + b*x]],x]

[Out] (2*d^2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, 1/4, 5/4, Sin[a + b*x]^2]*Tan[a + b*x])/(b*Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]])

Maple [A]

time = 0.14, size = 189, normalized size = 2.05

method	result
default	$ \frac{\left(-\sin(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \right) \right)}{2b(-1+\cos(bx+a)) \cos(bx+a)^2 \sqrt{c \sin(bx+a)}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(3/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/2/b*(-sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+cos(b*x+a)^2*2^(1/2)-cos(b*x+a)*2^(1/2))*sin(b*x+a)*(d*cos(b*x+a))^(3/2)/(-1+cos(b*x+a))/cos(b*x+a)^2/(c*sin(b*x+a))^(1/2)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^(3/2)/sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))*d*cos(b*x + a)/(c*sin(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \cos(a + bx))^{\frac{3}{2}}}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(3/2)/(c*sin(b*x+a))**(1/2),x)

[Out] Integral((d*cos(a + b*x))**(3/2)/sqrt(c*sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)/sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + bx))^{3/2}}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(3/2)/(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^(3/2)/(c*sin(a + b*x))^(1/2), x)

$$3.293 \quad \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx$$

Optimal. Leaf size=53

$$\frac{F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{b \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}$$

[Out] $-(\sin(a+1/4*\text{Pi}+b*x)^2)^{(1/2)}/\sin(a+1/4*\text{Pi}+b*x)*\text{EllipticF}(\cos(a+1/4*\text{Pi}+b*x), 2^{(1/2)})*\sin(2*b*x+2*a)^{(1/2)}/b/(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2653, 2720}

$$\frac{\sqrt{\sin(2a + 2bx)} F\left(a + bx - \frac{\pi}{4} \mid 2\right)}{b \sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]]),x]

[Out] (EllipticF[a - Pi/4 + b*x, 2]*Sqrt[Sin[2*a + 2*b*x]])/(b*Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]])

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx &= \frac{\sqrt{\sin(2a + 2bx)} \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \\ &= \frac{F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{b \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.05, size = 65, normalized size = 1.23

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{3}{4}; \frac{5}{4}; \sin^2(a + bx)\right) \tan(a + bx)}{b \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]]),x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/4, 3/4, 5/4, Sin[a + b*x]^2]*Tan[a + b*x])/(b*Sqrt[d*Cos[a + b*x]]*Sqrt[c*Sin[a + b*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 150 vs. 2(73) = 146.

time = 0.10, size = 151, normalized size = 2.85

method	result
default	$\frac{\text{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}}{b \sqrt{c \sin(bx+a)} \sqrt{d \cos(bx+a)} (-1+\cos(bx+a))}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/b*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*sin(b*x+a)^2/(c*sin(b*x+a))^(1/2)/(d*cos(b*x+a))^(1/2)/(-1+cos(b*x+a))*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))), x)

Fricas [C] Result contains complex when optimal does not.

time = 0.09, size = 60, normalized size = 1.13

$$\frac{\sqrt{icd} \text{ellipticF}(\cos(bx+a) + i \sin(bx+a), -1) + \sqrt{-icd} \text{ellipticF}(\cos(bx+a) - i \sin(bx+a), -1)}{bcd}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")
```

```
[Out] -(sqrt(I*c*d)*ellipticF(cos(b*x + a) + I*sin(b*x + a), -1) + sqrt(-I*c*d)*ellipticF(cos(b*x + a) - I*sin(b*x + a), -1))/(b*c*d)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sin(a + bx)} \sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))**(1/2)/(c*sin(b*x+a))**(1/2),x)
```

```
[Out] Integral(1/(sqrt(c*sin(a + b*x))*sqrt(d*cos(a + b*x))), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(1/(sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(1/2)),x)
```

```
[Out] int(1/((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^(1/2)), x)
```

$$3.294 \quad \int \frac{1}{(d \cos(a+bx))^{5/2} \sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=97

$$\frac{2\sqrt{c \sin(a+bx)}}{3bcd(d \cos(a+bx))^{3/2}} + \frac{2F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a+2bx)}}{3bd^2 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] $2/3*(c*\sin(b*x+a))^{(1/2)}/b/c/d/(d*\cos(b*x+a))^{(3/2)}-2/3*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x),2^{(1/2)})*\sin(2*b*x+2*a)^{(1/2)}/b/d^2/(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2651, 2653, 2720}

$$\frac{2\sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \mid 2\right)}{3bd^2 \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}} + \frac{2\sqrt{c \sin(a+bx)}}{3bcd(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Int[1/((d*Cos[a + b*x])^(5/2)*Sqrt[c*Sin[a + b*x]]),x]`

[Out] $(2*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(3*b*c*d*(d*\text{Cos}[a + b*x])^{(3/2)}) + (2*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(3*b*d^2*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2651

`Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[(- (b*Sin[e + f*x])^(n + 1))*((a*Cos[e + f*x])^(m + 1))/(a*b*f*(m + 1)), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rule 2653

`Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]]), x_Symbol] := Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{1}{(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}} dx &= \frac{2\sqrt{c \sin(a + bx)}}{3bcd(d \cos(a + bx))^{3/2}} + \frac{2 \int \frac{1}{\sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} dx}{3d^2} \\
&= \frac{2\sqrt{c \sin(a + bx)}}{3bcd(d \cos(a + bx))^{3/2}} + \frac{\left(2\sqrt{\sin(2a + 2bx)}\right) \int \frac{1}{\sqrt{\sin(2a + 2bx)}} dx}{3d^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}} \\
&= \frac{2\sqrt{c \sin(a + bx)}}{3bcd(d \cos(a + bx))^{3/2}} + \frac{2F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a + 2bx)}}{3bd^2 \sqrt{d \cos(a + bx)} \sqrt{c \sin(a + bx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.07, size = 65, normalized size = 0.67

$$\frac{2 \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{7}{4}; \frac{5}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)}}{bcd(d \cos(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Cos[a + b*x])^(5/2)*Sqrt[c*Sin[a + b*x]]),x]

[Out] (2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/4, 7/4, 5/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]])/(b*c*d*(d*Cos[a + b*x])^(3/2))

Maple [A]

time = 0.10, size = 185, normalized size = 1.91

method	result
default	$ \frac{\left(-2 \sin(bx+a) \cos(bx+a) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)}{3b(-1+\cos(bx+a))(d \cos(bx+a))^{\frac{5}{2}} \sqrt{c \sin(bx+a)}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*cos(b*x+a))^(5/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/3/b*(-2*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2))*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+cos(b*x+a)*2^(1/2)-2^(1/2))*cos(b*x+a)*sin(b*x+a)/(-1+cos(b*x+a))/(d*cos(b*x+a))^(5/2)/(c*sin(b*x+a))^(1/2)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(5/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/((d*cos(b*x + a))^(5/2)*sqrt(c*sin(b*x + a))), x)
```

Fricas [C] Result contains complex when optimal does not.

time = 0.10, size = 106, normalized size = 1.09

$$\frac{2 \left(\sqrt{icd} \cos(bx+a)^2 \operatorname{ellipticF}(\cos(bx+a) + i \sin(bx+a), -1) + \sqrt{-icd} \cos(bx+a)^2 \operatorname{ellipticF}(\cos(bx+a) - i \sin(bx+a), -1) - \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)} \right)}{3bcd^3 \cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(5/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")
```

```
[Out] -2/3*(sqrt(I*c*d)*cos(b*x + a)^2*ellipticF(cos(b*x + a) + I*sin(b*x + a), -1) + sqrt(-I*c*d)*cos(b*x + a)^2*ellipticF(cos(b*x + a) - I*sin(b*x + a), -1) - sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)))/(b*c*d^3*cos(b*x + a)^2)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))**(5/2)/(c*sin(b*x+a))**(1/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3007 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(5/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(1/((d*cos(b*x + a))^(5/2)*sqrt(c*sin(b*x + a))), x)
```


Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(d \cos(a + bx))^{\frac{5}{2}} \sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d*cos(a + b*x))^(5/2)*(c*sin(a + b*x))^(1/2)),x)

[Out] int(1/((d*cos(a + b*x))^(5/2)*(c*sin(a + b*x))^(1/2)), x)

$$3.295 \quad \int \frac{1}{(d \cos(a+bx))^{9/2} \sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=134

$$\frac{2\sqrt{c \sin(a+bx)}}{7bcd(d \cos(a+bx))^{7/2}} + \frac{4\sqrt{c \sin(a+bx)}}{7bcd^3(d \cos(a+bx))^{3/2}} + \frac{4F\left(a - \frac{\pi}{4} + bx \mid 2\right) \sqrt{\sin(2a+2bx)}}{7bd^4 \sqrt{d \cos(a+bx)} \sqrt{c \sin(a+bx)}}$$

[Out] $2/7*(c*\sin(b*x+a))^{(1/2)}/b/c/d/(d*\cos(b*x+a))^{(7/2)}+4/7*(c*\sin(b*x+a))^{(1/2)}/b/c/d^3/(d*\cos(b*x+a))^{(3/2)}-4/7*(\sin(a+1/4*Pi+b*x)^2)^{(1/2)}/\sin(a+1/4*Pi+b*x)*\text{EllipticF}(\cos(a+1/4*Pi+b*x), 2^{(1/2)})*\sin(2*b*x+2*a)^{(1/2)}/b/d^4/(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 134, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2651, 2653, 2720}

$$\frac{4\sqrt{\sin(2a+2bx)} F\left(a+bx - \frac{\pi}{4} \mid 2\right)}{7bd^4 \sqrt{c \sin(a+bx)} \sqrt{d \cos(a+bx)}} + \frac{4\sqrt{c \sin(a+bx)}}{7bcd^3(d \cos(a+bx))^{3/2}} + \frac{2\sqrt{c \sin(a+bx)}}{7bcd(d \cos(a+bx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Cos[a + b*x])^(9/2)*Sqrt[c*Sin[a + b*x]]), x]

[Out] $(2*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(7*b*c*d*(d*\text{Cos}[a + b*x])^{(7/2)}) + (4*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(7*b*c*d^3*(d*\text{Cos}[a + b*x])^{(3/2)}) + (4*\text{EllipticF}[a - \text{Pi}/4 + b*x, 2]*\text{Sqrt}[\text{Sin}[2*a + 2*b*x]])/(7*b*d^4*\text{Sqrt}[d*\text{Cos}[a + b*x]]*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-(b*Sin[e + f*x])^(n + 1))*((a*Cos[e + f*x])^(m + 1))/(a*b*f*(m + 1)), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^n*(a*Cos[e + f*x])^(m + 2), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(d \cos(a + bx))^{9/2} \sqrt{c \sin(a + bx)}} dx &= \frac{2\sqrt{c \sin(a + bx)}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{6 \int \frac{1}{(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}} dx}{7d^2} \\ &= \frac{2\sqrt{c \sin(a + bx)}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{4\sqrt{c \sin(a + bx)}}{7bcd^3(d \cos(a + bx))^{3/2}} + \frac{4 \int \sqrt{d \cos(a + bx)}}{\sqrt{d \cos(a + bx)}} dx}{7d^4} \\ &= \frac{2\sqrt{c \sin(a + bx)}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{4\sqrt{c \sin(a + bx)}}{7bcd^3(d \cos(a + bx))^{3/2}} + \frac{(4\sqrt{\sin(2a + 2bx)})}{7d^4 \sqrt{d \cos(a + bx)}} \\ &= \frac{2\sqrt{c \sin(a + bx)}}{7bcd(d \cos(a + bx))^{7/2}} + \frac{4\sqrt{c \sin(a + bx)}}{7bcd^3(d \cos(a + bx))^{3/2}} + \frac{4F(a - \frac{\pi}{4} + bx)}{7bd^4 \sqrt{d \cos(a + bx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.09, size = 70, normalized size = 0.52

$$\frac{2 \cos^3(a + bx) \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{11}{4}; \frac{5}{4}; \sin^2(a + bx)\right) \sqrt{c \sin(a + bx)}}{bc(d \cos(a + bx))^{9/2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*cos[a + b*x])^(9/2)*Sqrt[c*Sin[a + b*x]]),x]

[Out] (2*cos[a + b*x]^3*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[1/4, 11/4, 5/4, Sin[a + b*x]^2]*Sqrt[c*Sin[a + b*x]])/(b*c*(d*cos[a + b*x])^(9/2))

Maple [A]

time = 0.13, size = 213, normalized size = 1.59

method	result
default	$\frac{\left(-4 \sin(bx+a) (\cos^3(bx+a)) \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)}{7b(-1+\cos(bx+a))(d \cos(bx+a))^{9/2}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*cos(b*x+a))^(9/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

```
[Out] 1/7/b*(-4*sin(b*x+a)*cos(b*x+a)^3*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2), 1/2*2^(1/2)))+2*cos(b*x+a)^3*2^(1/2)-2*cos(b*x+a)^2*2^(1/2)+cos(b*x+a)*2^(1/2)-2^(1/2))*cos(b*x+a)*sin(b*x+a)/(-1+cos(b*x+a))/(d*cos(b*x+a))^(9/2)/(c*sin(b*x+a))^(1/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(9/2)/(c*sin(b*x+a))^(1/2), x, algorithm="maxima")
```

```
[Out] integrate(1/((d*cos(b*x + a))^(9/2)*sqrt(c*sin(b*x + a))), x)
```

Fricas [C] Result contains complex when optimal does not.

time = 0.13, size = 120, normalized size = 0.90

$$\frac{2(2\sqrt{icd}\cos(bx+a)^4\text{ellipticF}(\cos(bx+a)+i\sin(bx+a), -1)+2\sqrt{-icd}\cos(bx+a)^4\text{ellipticF}(\cos(bx+a)-i\sin(bx+a), -1)-\sqrt{d\cos(bx+a)}(2\cos(bx+a)^2+1)\sqrt{c\sin(bx+a)})}{7bcd^5\cos(bx+a)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(9/2)/(c*sin(b*x+a))^(1/2), x, algorithm="fricas")
```

```
[Out] -2/7*(2*sqrt(I*c*d)*cos(b*x + a)^4*ellipticF(cos(b*x + a) + I*sin(b*x + a), -1) + 2*sqrt(-I*c*d)*cos(b*x + a)^4*ellipticF(cos(b*x + a) - I*sin(b*x + a), -1) - sqrt(d*cos(b*x + a))*(2*cos(b*x + a)^2 + 1)*sqrt(c*sin(b*x + a)))/(b*c*d^5*cos(b*x + a)^4)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))**(9/2)/(c*sin(b*x+a))**(1/2), x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(9/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*cos(b*x + a))^(9/2)*sqrt(c*sin(b*x + a))), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(d \cos(a + bx))^{9/2} \sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d*cos(a + b*x))^(9/2)*(c*sin(a + b*x))^(1/2)),x)

[Out] int(1/((d*cos(a + b*x))^(9/2)*(c*sin(a + b*x))^(1/2)), x)

$$3.296 \quad \int \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} dx$$

Optimal. Leaf size=280

$$\frac{\sqrt{d} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)}{\sqrt{2} b \sqrt{c}} - \frac{\sqrt{d} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)}{\sqrt{2} b \sqrt{c}} - \sqrt{d} \log \left(\sqrt{d} + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)$$

[Out] $-1/2*\arctan(-1+2^{(1/2)}*c^{(1/2)}*(d*\cos(b*x+a))^{(1/2)}/d^{(1/2)}/(c*\sin(b*x+a))^{(1/2)})*d^{(1/2)}/b*2^{(1/2)}/c^{(1/2)}-1/2*\arctan(1+2^{(1/2)}*c^{(1/2)}*(d*\cos(b*x+a))^{(1/2)}/d^{(1/2)}/(c*\sin(b*x+a))^{(1/2)})*d^{(1/2)}/b*2^{(1/2)}/c^{(1/2)}-1/4*\ln(d^{(1/2)}+\cot(b*x+a)*d^{(1/2)}-2^{(1/2)}*c^{(1/2)}*(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)})*d^{(1/2)}/b*2^{(1/2)}/c^{(1/2)}+1/4*\ln(d^{(1/2)}+\cot(b*x+a)*d^{(1/2)}+2^{(1/2)}*c^{(1/2)}*(d*\cos(b*x+a))^{(1/2)}/(c*\sin(b*x+a))^{(1/2)})*d^{(1/2)}/b*2^{(1/2)}/c^{(1/2)}$

Rubi [A]

time = 0.12, antiderivative size = 280, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.280$, Rules used = {2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{d} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} \right)}{\sqrt{2} b \sqrt{c}} - \frac{\sqrt{d} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} + 1 \right)}{\sqrt{2} b \sqrt{c}} - \frac{\sqrt{d} \log \left(-\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} + \sqrt{d} \cot(a + bx) + \sqrt{d} \right)}{2\sqrt{2} b \sqrt{c}} + \frac{\sqrt{d} \log \left(\frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a + bx)}}{\sqrt{d} \sqrt{c \sin(a + bx)}} + \sqrt{d} \cot(a + bx) + \sqrt{d} \right)}{2\sqrt{2} b \sqrt{c}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Cos[a + b*x]]/Sqrt[c*Sin[a + b*x]],x]

[Out] $(\operatorname{Sqrt}[d]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]])]/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c*\operatorname{Sin}[a + b*x]])]/(\operatorname{Sqrt}[2]*b*\operatorname{Sqrt}[c]) - (\operatorname{Sqrt}[d]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]])]/(\operatorname{Sqrt}[d]*\operatorname{Sqrt}[c*\operatorname{Sin}[a + b*x]])]/(\operatorname{Sqrt}[2]*b*\operatorname{Sqrt}[c]) - (\operatorname{Sqrt}[d]*\operatorname{Log}[\operatorname{Sqrt}[d] + \operatorname{Sqrt}[d]*\operatorname{Cot}[a + b*x] - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]])]/\operatorname{Sqrt}[c*\operatorname{Sin}[a + b*x]])]/(2*\operatorname{Sqrt}[2]*b*\operatorname{Sqrt}[c]) + (\operatorname{Sqrt}[d]*\operatorname{Log}[\operatorname{Sqrt}[d] + \operatorname{Sqrt}[d]*\operatorname{Cot}[a + b*x] + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*\operatorname{Sqrt}[d*\operatorname{Cos}[a + b*x]])]/\operatorname{Sqrt}[c*\operatorname{Sin}[a + b*x]])]/(2*\operatorname{Sqrt}[2]*b*\operatorname{Sqrt}[c])$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4

), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2655

Int[(cos[(e_) + (f_)*(x_)])*(a_)^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sin[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} dx &= \frac{(2cd) \operatorname{Subst} \left(\int \frac{x^2}{d^2+c^2x^4} dx, x, \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{b} \\
&= \frac{d \operatorname{Subst} \left(\int \frac{d-cx^2}{d^2+c^2x^4} dx, x, \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{b} - \frac{d \operatorname{Subst} \left(\int \frac{d+cx^2}{d^2+c^2x^4} dx, x, \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{b} \\
&= \frac{\sqrt{d} \operatorname{Subst} \left(\int \frac{\frac{\sqrt{2} \sqrt{d}}{\sqrt{c}} + 2x}{-\frac{d}{c} - \frac{\sqrt{2} \sqrt{d}}{\sqrt{c}} x - x^2} dx, x, \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{2\sqrt{2} b \sqrt{c}} - \frac{\sqrt{d} \operatorname{Subst} \left(\int \frac{\frac{\sqrt{2} \sqrt{d}}{\sqrt{c}}}{-\frac{d}{c} + \frac{\sqrt{2} \sqrt{d}}{\sqrt{c}} x - x^2} dx, x, \frac{\sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{2\sqrt{2} b \sqrt{c}} \\
&= \frac{\sqrt{d} \log \left(\sqrt{d} + \sqrt{d} \cot(a+bx) - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{2\sqrt{2} b \sqrt{c}} + \frac{\sqrt{d} \log \left(\sqrt{d} - \sqrt{d} \cot(a+bx) - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{c \sin(a+bx)}} \right)}{2\sqrt{2} b \sqrt{c}} \\
&= \frac{\sqrt{d} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{d} \sqrt{c \sin(a+bx)}} \right)}{\sqrt{2} b \sqrt{c}} - \frac{\sqrt{d} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{c} \sqrt{d \cos(a+bx)}}{\sqrt{d} \sqrt{c \sin(a+bx)}} \right)}{\sqrt{2} b \sqrt{c}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 65, normalized size = 0.23

$$\frac{2\sqrt{d \cos(a+bx)} \sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \sin^2(a+bx)\right) \tan(a+bx)}{b\sqrt{c \sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]/Sqrt[c*Sin[a + b*x]], x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, Sin[a + b*x]^2]*Tan[a + b*x])/(b*Sqrt[c*Sin[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.12, size = 312, normalized size = 1.11

method	result
default	$ -\frac{\sqrt{d \cos(bx+a)} \sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \left(i \operatorname{EllipticPi} \left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \right) \right)}{b\sqrt{c \sin(a+bx)}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/b*(d*\cos(b*x+a))^{1/2}*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*(I*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-I*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})-2*\text{EllipticF}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2*2^{1/2})+\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})+\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2}))*\sin(b*x+a)^2/(c*\sin(b*x+a))^{1/2}/\cos(b*x+a)/(-1+\cos(b*x+a))*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(d*cos(b*x + a))/sqrt(c*sin(b*x + a)), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1697 vs. 2(208) = 416.

time = 15.44, size = 1697, normalized size = 6.06

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")`

[Out]
$$-1/4*\sqrt{2}*(d^2/(b^4*c^2))^{1/4}*\arctan(1/2*(2*d^4*\cos(b*x + a)*\sin(b*x + a) + \sqrt{4*b^2*c*d^3*\sqrt{d^2/(b^4*c^2)}*\cos(b*x + a)*\sin(b*x + a) + d^4 + 2*(\sqrt{2}*b^3*c*d^2*(d^2/(b^4*c^2))^{3/4}*\cos(b*x + a) + \sqrt{2}*b*d^3*(d^2/(b^4*c^2))^{1/4}*\sin(b*x + a))*\sqrt{d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a)}}*(\sqrt{2}*b^3*c*(d^2/(b^4*c^2))^{3/4}*\sin(b*x + a) + \sqrt{2}*b*d*(d^2/(b^4*c^2))^{1/4}*\cos(b*x + a))*\sqrt{d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a)} + (\sqrt{2}*b^3*c*d^2*(d^2/(b^4*c^2))^{3/4}*\sin(b*x + a) + \sqrt{2}*b*d^3*(d^2/(b^4*c^2))^{1/4}*\cos(b*x + a))*\sqrt{d*\cos(b*x + a))*\sqrt{c*\sin(b*x + a)} - 4*(b^2*c*d^3*\cos(b*x + a)^4 - b^2*c*d^3*\cos(b*x + a)^2)*\sqrt{d^2/(b^4*c^2)})/(2*d^4*\cos(b*x + a)^3 - d^4*\cos(b*x + a)*\sin(b*x + a)) - 1/4*\sqrt{2}*(d^2/(b^4*c^2))^{1/4}*\arctan(-1/2*(2*d^4*\cos(b*x + a)*\sin(b*x + a) - \sqrt{4*b^2$$

```

*c*d^3*sqrt(d^2/(b^4*c^2))*cos(b*x + a)*sin(b*x + a) + d^4 - 2*(sqrt(2)*b^3
*c*d^2*(d^2/(b^4*c^2))^(3/4)*cos(b*x + a) + sqrt(2)*b*d^3*(d^2/(b^4*c^2))^(
1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)))*(sqrt(2)*b^3*
c*(d^2/(b^4*c^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*d*(d^2/(b^4*c^2))^(1/4)*co
s(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) - (sqrt(2)*b^3*c*d^2*
(d^2/(b^4*c^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*d^3*(d^2/(b^4*c^2))^(1/4)*co
s(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) - 4*(b^2*c*d^3*cos(b*
x + a)^4 - b^2*c*d^3*cos(b*x + a)^2)*sqrt(d^2/(b^4*c^2)))/((2*d^4*cos(b*x +
a)^3 - d^4*cos(b*x + a))*sin(b*x + a)) - 1/4*sqrt(2)*(d^2/(b^4*c^2))^(1/4
)*arctan(-1/2*((sqrt(2)*b^3*c*d^2*(d^2/(b^4*c^2))^(3/4)*sin(b*x + a) - sqrt
(2)*b*d^3*(d^2/(b^4*c^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*s
in(b*x + a)) - sqrt(4*b^2*c*d^3*sqrt(d^2/(b^4*c^2))*cos(b*x + a)*sin(b*x +
a) + d^4 - 2*(sqrt(2)*b^3*c*d^2*(d^2/(b^4*c^2))^(3/4)*cos(b*x + a) + sqrt(2
)*b*d^3*(d^2/(b^4*c^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin
(b*x + a)))*(2*d^2*cos(b*x + a)*sin(b*x + a) + (sqrt(2)*b^3*c*(d^2/(b^4*c^2
))^(3/4)*sin(b*x + a) + sqrt(2)*b*d*(d^2/(b^4*c^2))^(1/4)*cos(b*x + a))*sqr
t(d*cos(b*x + a))*sqrt(c*sin(b*x + a))))/(d^4*cos(b*x + a)*sin(b*x + a)) -
1/4*sqrt(2)*(d^2/(b^4*c^2))^(1/4)*arctan(-1/2*((sqrt(2)*b^3*c*d^2*(d^2/(b^
4*c^2))^(3/4)*sin(b*x + a) - sqrt(2)*b*d^3*(d^2/(b^4*c^2))^(1/4)*cos(b*x +
a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) + sqrt(4*b^2*c*d^3*sqrt(d^2/(
b^4*c^2))*cos(b*x + a)*sin(b*x + a) + d^4 + 2*(sqrt(2)*b^3*c*d^2*(d^2/(b^4*
c^2))^(3/4)*cos(b*x + a) + sqrt(2)*b*d^3*(d^2/(b^4*c^2))^(1/4)*sin(b*x + a
))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)))*(2*d^2*cos(b*x + a)*sin(b*x +
a) - (sqrt(2)*b^3*c*(d^2/(b^4*c^2))^(3/4)*sin(b*x + a) + sqrt(2)*b*d*(d^2/(
b^4*c^2))^(1/4)*cos(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))))/(
d^4*cos(b*x + a)*sin(b*x + a)) + 1/8*sqrt(2)*(d^2/(b^4*c^2))^(1/4)*log(4*b
^2*c*d^3*sqrt(d^2/(b^4*c^2))*cos(b*x + a)*sin(b*x + a) + d^4 + 2*(sqrt(2)*b
^3*c*d^2*(d^2/(b^4*c^2))^(3/4)*cos(b*x + a) + sqrt(2)*b*d^3*(d^2/(b^4*c^2)
)^(1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a)) - 1/8*sqrt(
2)*(d^2/(b^4*c^2))^(1/4)*log(4*b^2*c*d^3*sqrt(d^2/(b^4*c^2))*cos(b*x + a)*s
in(b*x + a) + d^4 - 2*(sqrt(2)*b^3*c*d^2*(d^2/(b^4*c^2))^(3/4)*cos(b*x + a)
+ sqrt(2)*b*d^3*(d^2/(b^4*c^2))^(1/4)*sin(b*x + a))*sqrt(d*cos(b*x + a))*s
qrt(c*sin(b*x + a)))

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)/(c*sin(b*x+a))**(1/2), x)

[Out] Integral(sqrt(d*cos(a + b*x))/sqrt(c*sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*cos(b*x + a))/sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{d \cos(a + bx)}}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(1/2)/(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^(1/2)/(c*sin(a + b*x))^(1/2), x)

$$3.297 \quad \int \frac{1}{(d \cos(a+bx))^{3/2} \sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=35

$$\frac{2\sqrt{c \sin(a+bx)}}{bcd\sqrt{d \cos(a+bx)}}$$

[Out] $2*(c*\sin(b*x+a))^(1/2)/b/c/d/(d*\cos(b*x+a))^(1/2)$

Rubi [A]

time = 0.04, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2643}

$$\frac{2\sqrt{c \sin(a+bx)}}{bcd\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] `Int[1/((d*Cos[a + b*x])^(3/2)*Sqrt[c*Sin[a + b*x]]),x]`

[Out] `(2*Sqrt[c*Sin[a + b*x]])/(b*c*d*Sqrt[d*Cos[a + b*x]])`

Rule 2643

`Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] & & NeQ[m, -1]`

Rubi steps

$$\int \frac{1}{(d \cos(a+bx))^{3/2} \sqrt{c \sin(a+bx)}} dx = \frac{2\sqrt{c \sin(a+bx)}}{bcd\sqrt{d \cos(a+bx)}}$$

Mathematica [A]

time = 0.04, size = 36, normalized size = 1.03

$$\frac{\sin(2(a+bx))}{b(d \cos(a+bx))^{3/2} \sqrt{c \sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] `Integrate[1/((d*Cos[a + b*x])^(3/2)*Sqrt[c*Sin[a + b*x]]),x]`

[Out] $\text{Sin}[2*(a + b*x)]/(b*(d*\text{Cos}[a + b*x])^{(3/2)}*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Maple [A]

time = 0.09, size = 38, normalized size = 1.09

method	result	size
default	$\frac{2 \cos(bx+a) \sin(bx+a)}{b(d \cos(bx+a))^{\frac{3}{2}} \sqrt{c \sin(bx+a)}}$	38

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}(1/(d*\text{cos}(b*x+a))^{(3/2)}/(c*\text{sin}(b*x+a))^{(1/2)},x,\text{method}=_RETURNVERBOSE)$

[Out] $2/b*\text{cos}(b*x+a)*\text{sin}(b*x+a)/(d*\text{cos}(b*x+a))^{(3/2)}/(c*\text{sin}(b*x+a))^{(1/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/(d*\text{cos}(b*x+a))^{(3/2)}/(c*\text{sin}(b*x+a))^{(1/2)},x,\text{algorithm}=\text{"maxima"})$

[Out] $\text{integrate}(1/((d*\text{cos}(b*x + a))^{(3/2)}*\text{sqrt}(c*\text{sin}(b*x + a))), x)$

Fricas [A]

time = 0.44, size = 39, normalized size = 1.11

$$\frac{2 \sqrt{d \cos(bx+a)} \sqrt{c \sin(bx+a)}}{bcd^2 \cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/(d*\text{cos}(b*x+a))^{(3/2)}/(c*\text{sin}(b*x+a))^{(1/2)},x,\text{algorithm}=\text{"fricas"})$

[Out] $2*\text{sqrt}(d*\text{cos}(b*x + a))*\text{sqrt}(c*\text{sin}(b*x + a))/(b*c*d^2*\text{cos}(b*x + a))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{c \sin(a + bx)} (d \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/(d*\text{cos}(b*x+a))^{(3/2)}/(c*\text{sin}(b*x+a))^{(1/2)},x)$

[Out] Integral(1/(sqrt(c*sin(a + b*x))*(d*cos(a + b*x))**(3/2)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(3/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*cos(b*x + a))^(3/2)*sqrt(c*sin(b*x + a))), x)

Mupad [B]

time = 0.81, size = 31, normalized size = 0.89

$$\frac{2 \sqrt{c \sin(a + b x)}}{b c d \sqrt{d \cos(a + b x)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^(1/2)),x)

[Out] (2*(c*sin(a + b*x))^(1/2))/(b*c*d*(d*cos(a + b*x))^(1/2))

$$3.298 \quad \int \frac{1}{(d \cos(a+bx))^{7/2} \sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=75

$$\frac{2\sqrt{c \sin(a+bx)}}{5bcd(d \cos(a+bx))^{5/2}} + \frac{8\sqrt{c \sin(a+bx)}}{5bcd^3 \sqrt{d \cos(a+bx)}}$$

[Out] $2/5*(c*\sin(b*x+a))^{(1/2)}/b/c/d/(d*\cos(b*x+a))^{(5/2)}+8/5*(c*\sin(b*x+a))^{(1/2)}/b/c/d^3/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2651, 2643}

$$\frac{8\sqrt{c \sin(a+bx)}}{5bcd^3 \sqrt{d \cos(a+bx)}} + \frac{2\sqrt{c \sin(a+bx)}}{5bcd(d \cos(a+bx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[1/((d*Cos[a + b*x])^(7/2)*Sqrt[c*Sin[a + b*x]]),x]

[Out] (2*Sqrt[c*Sin[a + b*x]])/(5*b*c*d*(d*Cos[a + b*x])^(5/2)) + (8*Sqrt[c*Sin[a + b*x]])/(5*b*c*d^3*Sqrt[d*Cos[a + b*x]])

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] && NeQ[m, -1]

Rule 2651

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] :> Simp[(-b*Sin[e + f*x])^(n + 1)*((a*Cos[e + f*x])^(m + 1)/(a*b*f*(m + 1))), x] + Dist[(m + n + 2)/(a^2*(m + 1)), Int[(b*Sin[e + f*x])^(n)*((a*Cos[e + f*x])^(m + 2)), x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\int \frac{1}{(d \cos(a + bx))^{7/2} \sqrt{c \sin(a + bx)}} dx = \frac{2\sqrt{c \sin(a + bx)}}{5bcd(d \cos(a + bx))^{5/2}} + \frac{4 \int \frac{1}{(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}} dx}{5d^2}$$

$$= \frac{2\sqrt{c \sin(a + bx)}}{5bcd(d \cos(a + bx))^{5/2}} + \frac{8\sqrt{c \sin(a + bx)}}{5bcd^3 \sqrt{d \cos(a + bx)}}$$

Mathematica [A]

time = 0.11, size = 52, normalized size = 0.69

$$\frac{2(3 + 2 \cos(2(a + bx))) \tan(a + bx)}{5bd^2(d \cos(a + bx))^{3/2} \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/((d*Cos[a + b*x])^(7/2)*Sqrt[c*Sin[a + b*x]]),x]
```

```
[Out] (2*(3 + 2*Cos[2*(a + b*x)])*Tan[a + b*x])/(5*b*d^2*(d*Cos[a + b*x])^(3/2)*Sqrt[c*Sin[a + b*x]])
```

Maple [A]

time = 0.10, size = 50, normalized size = 0.67

method	result	size
default	$\frac{2(4(\cos^2(bx+a))+1) \cos(bx+a) \sin(bx+a)}{5b(d \cos(bx+a))^{7/2} \sqrt{c \sin(bx+a)}}$	50

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 2/5/b*(4*cos(b*x+a)^2+1)*cos(b*x+a)*sin(b*x+a)/(d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/((d*cos(b*x + a))^(7/2)*sqrt(c*sin(b*x + a))), x)
```


Fricas [A]

time = 0.40, size = 51, normalized size = 0.68

$$\frac{2 \sqrt{d \cos(bx + a)} (4 \cos(bx + a)^2 + 1) \sqrt{c \sin(bx + a)}}{5 bcd^4 \cos(bx + a)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] 2/5*sqrt(d*cos(b*x + a))*(4*cos(b*x + a)^2 + 1)*sqrt(c*sin(b*x + a))/(b*c*d^4*cos(b*x + a)^3)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))**(7/2)/(c*sin(b*x+a))**(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(7/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*cos(b*x + a))^(7/2)*sqrt(c*sin(b*x + a))), x)

Mupad [B]

time = 1.49, size = 77, normalized size = 1.03

$$\frac{8 \sqrt{c \sin(a + bx)} (5 \cos(2a + 2bx) + \cos(4a + 4bx) + 4)}{5 bcd^3 \sqrt{d \cos(a + bx)} (4 \cos(2a + 2bx) + \cos(4a + 4bx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d*cos(a + b*x))^(7/2)*(c*sin(a + b*x))^(1/2)),x)

[Out] (8*(c*sin(a + b*x))^(1/2)*(5*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 4))/(5*b*c*d^3*(d*cos(a + b*x))^(1/2)*(4*cos(2*a + 2*b*x) + cos(4*a + 4*b*x) + 3))

$$3.299 \quad \int \frac{1}{(d \cos(a+bx))^{11/2} \sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=112

$$\frac{2\sqrt{c \sin(a+bx)}}{9bcd(d \cos(a+bx))^{9/2}} + \frac{16\sqrt{c \sin(a+bx)}}{45bcd^3(d \cos(a+bx))^{5/2}} + \frac{64\sqrt{c \sin(a+bx)}}{45bcd^5 \sqrt{d \cos(a+bx)}}$$

[Out] $2/9*(c*\sin(b*x+a))^{(1/2)}/b/c/d/(d*\cos(b*x+a))^{(9/2)}+16/45*(c*\sin(b*x+a))^{(1/2)}/b/c/d^3/(d*\cos(b*x+a))^{(5/2)}+64/45*(c*\sin(b*x+a))^{(1/2)}/b/c/d^5/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 112, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2651, 2643}

$$\frac{64\sqrt{c \sin(a+bx)}}{45bcd^5 \sqrt{d \cos(a+bx)}} + \frac{16\sqrt{c \sin(a+bx)}}{45bcd^3(d \cos(a+bx))^{5/2}} + \frac{2\sqrt{c \sin(a+bx)}}{9bcd(d \cos(a+bx))^{9/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/((d*\text{Cos}[a + b*x])^{(11/2)}*\text{Sqrt}[c*\text{Sin}[a + b*x]]), x]$

[Out] $(2*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(9*b*c*d*(d*\text{Cos}[a + b*x])^{(9/2)}) + (16*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(45*b*c*d^3*(d*\text{Cos}[a + b*x])^{(5/2)}) + (64*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(45*b*c*d^5*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2643

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e + f*x])^{(m + 1)}*((b*\text{Cos}[e + f*x])^{(n + 1)})/(a*b*f*(m + 1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m + n + 2, 0] \&\& \text{NeQ}[m, -1]$

Rule 2651

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b*\text{Sin}[e + f*x])^{(n + 1)}*((a*\text{Cos}[e + f*x])^{(m + 1)})/(a*b*f*(m + 1)), x] + \text{Dist}[(m + n + 2)/(a^2*(m + 1)), \text{Int}[(b*\text{Sin}[e + f*x])^{(n)}*(a*\text{Cos}[e + f*x])^{(m + 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rubi steps

$$\begin{aligned}
\int \frac{1}{(d \cos(a + bx))^{11/2} \sqrt{c \sin(a + bx)}} dx &= \frac{2\sqrt{c \sin(a + bx)}}{9bcd(d \cos(a + bx))^{9/2}} + \frac{8 \int \frac{1}{(d \cos(a + bx))^{7/2} \sqrt{c \sin(a + bx)}} dx}{9d^2} \\
&= \frac{2\sqrt{c \sin(a + bx)}}{9bcd(d \cos(a + bx))^{9/2}} + \frac{16\sqrt{c \sin(a + bx)}}{45bcd^3(d \cos(a + bx))^{5/2}} + \frac{32 \int \frac{1}{(d \cos(a + bx))^{5/2} \sqrt{c \sin(a + bx)}} dx}{45bcd^5} \\
&= \frac{2\sqrt{c \sin(a + bx)}}{9bcd(d \cos(a + bx))^{9/2}} + \frac{16\sqrt{c \sin(a + bx)}}{45bcd^3(d \cos(a + bx))^{5/2}} + \frac{64\sqrt{c \sin(a + bx)}}{45bcd^5}
\end{aligned}$$

Mathematica [A]

time = 0.16, size = 67, normalized size = 0.60

$$\frac{2\sqrt{d \cos(a + bx)} (21 + 20 \cos(2(a + bx)) + 4 \cos(4(a + bx))) \sec^5(a + bx) \sqrt{c \sin(a + bx)}}{45bcd^6}$$

Antiderivative was successfully verified.

[In] Integrate[1/((d*Cos[a + b*x])^(11/2)*Sqrt[c*Sin[a + b*x]]),x]

[Out] (2*Sqrt[d*Cos[a + b*x]]*(21 + 20*Cos[2*(a + b*x)] + 4*Cos[4*(a + b*x)])*Sec[a + b*x]^5*Sqrt[c*Sin[a + b*x]])/(45*b*c*d^6)

Maple [A]

time = 0.12, size = 60, normalized size = 0.54

method	result	size
default	$\frac{2(32(\cos^4(bx+a))+8(\cos^2(bx+a))+5)\cos(bx+a)\sin(bx+a)}{45b(d\cos(bx+a))^{\frac{11}{2}}\sqrt{c\sin(bx+a)}}$	60

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*cos(b*x+a))^(11/2)/(c*sin(b*x+a))^(1/2),x,method=_RETURNVERBOSE)

[Out] 2/45/b*(32*cos(b*x+a)^4+8*cos(b*x+a)^2+5)*cos(b*x+a)*sin(b*x+a)/(d*cos(b*x+a))^(11/2)/(c*sin(b*x+a))^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(11/2)/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(1/((d*cos(b*x + a))^(11/2)*sqrt(c*sin(b*x + a))), x)

Fricas [A]

time = 0.41, size = 61, normalized size = 0.54

$$\frac{2 (32 \cos (bx + a)^4 + 8 \cos (bx + a)^2 + 5) \sqrt{d \cos (bx + a)} \sqrt{c \sin (bx + a)}}{45 bcd^6 \cos (bx + a)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(11/2)/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] 2/45*(32*cos(b*x + a)^4 + 8*cos(b*x + a)^2 + 5)*sqrt(d*cos(b*x + a))*sqrt(c*sin(b*x + a))/(b*c*d^6*cos(b*x + a)^5)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))**(11/2)/(c*sin(b*x+a))**(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*cos(b*x+a))^(11/2)/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(1/((d*cos(b*x + a))^(11/2)*sqrt(c*sin(b*x + a))), x)

Mupad [B]

time = 3.83, size = 123, normalized size = 1.10

$$\frac{32 \sqrt{c \sin (a + bx)} (162 \cos (2a + 2bx) + 73 \cos (4a + 4bx) + 18 \cos (6a + 6bx) + 2 \cos (8a + 8bx) + 105)}{45 bcd^5 \sqrt{d \cos (a + bx)} (56 \cos (2a + 2bx) + 28 \cos (4a + 4bx) + 8 \cos (6a + 6bx) + \cos (8a + 8bx) + 35)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((d*cos(a + b*x))^(11/2)*(c*sin(a + b*x))^(1/2)),x)

[Out] (32*(c*sin(a + b*x))^(1/2)*(162*cos(2*a + 2*b*x) + 73*cos(4*a + 4*b*x) + 18*cos(6*a + 6*b*x) + 2*cos(8*a + 8*b*x) + 105))/(45*b*c*d^5*(d*cos(a + b*x))^(1/2)*(56*cos(2*a + 2*b*x) + 28*cos(4*a + 4*b*x) + 8*cos(6*a + 6*b*x) + cos(8*a + 8*b*x) + 35))

$$3.300 \quad \int \frac{\sqrt{\cos(a + bx)}}{\sqrt{\sin(a + bx)}} dx$$

Optimal. Leaf size=174

$$\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\log\left(1 + \cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b}$$

[Out] $-1/2*\arctan(-1+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}-1/2*\arctan(1+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}-1/4*\ln(1+\cot(b*x+a))-2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)}/b*2^{(1/2)}+1/4*\ln(1+\cot(b*x+a))+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)}/b*2^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 174, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\text{ArcTan}\left(\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{\sqrt{2}b} - \frac{\log\left(\cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{2\sqrt{2}b} + \frac{\log\left(\cot(a+bx) + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Cos[a + b*x]]/Sqrt[Sin[a + b*x]], x]

[Out] ArcTan[1 - (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(Sqrt[2]*b) - ArcTan[1 + (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(Sqrt[2]*b) - Log[1 + Cot[a + b*x] - (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(2*Sqrt[2]*b) + Log[1 + Cot[a + b*x] + (Sqrt[2]*Sqrt[Cos[a + b*x]])/Sqrt[Sin[a + b*x]]]/(2*Sqrt[2]*b)

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2655

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sin[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} dx &= -\frac{2\text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} - \frac{\text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} \\
&= -\frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2b} - \frac{\text{Subst}\left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2b} \\
&= -\frac{\log\left(1+\cot(a+bx)-\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} + \frac{\log\left(1+\cot(a+bx)+\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} \\
&= \frac{\tan^{-1}\left(1-\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\tan^{-1}\left(1+\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} - \frac{\log\left(1+\cot(a+bx)+\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 55, normalized size = 0.32

$$\frac{2^4 \sqrt{\cos^2(a+bx)} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \sin^2(a+bx)\right) \sqrt{\sin(a+bx)}}{b \sqrt{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Cos[a + b*x]]/Sqrt[Sin[a + b*x]], x]

[Out] (2*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, Sin[a + b*x]^2]*Sqrt[Sin[a + b*x]])/(b*Sqrt[Cos[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.11, size = 292, normalized size = 1.68

method	result
default	$-\frac{\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}} \sqrt{\frac{-1+\cos(bx+a)}{\sin(bx+a)}} \left(i \text{EllipticPi}\left(\sqrt{\frac{1-\cos(bx+a)+\sin(bx+a)}{\sin(bx+a)}}\right)\right)}{b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(1/2)/sin(b*x+a)^(1/2), x, method=_RETURNVERBOSE)

```
[Out] -1/2/b/cos(b*x+a)^(1/2)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*(I*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-I*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))-2*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))+EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2)))*sin(b*x+a)^(3/2)/(-1+cos(b*x+a))*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)^(1/2)/sin(b*x+a)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(cos(b*x + a))/sqrt(sin(b*x + a)), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1185 vs. 2(138) = 276.

time = 14.64, size = 1185, normalized size = 6.81

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)^(1/2)/sin(b*x+a)^(1/2),x, algorithm="fricas")
```

```
[Out] -1/4*sqrt(2)*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + (sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 2*cos(b*x + a)*sin(b*x + a) - 4*(b^2*cos(b*x + a)^4 - b^2*cos(b*x + a)^2)*sqrt(b^(-4)))/((2*cos(b*x + a)^3 - cos(b*x + a))*sin(b*x + a)) - 1/4*sqrt(2)*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + (sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) - 2*cos(b*x + a)*sin(b*x + a) + 4*(b^2*cos(b*x + a)^4 - b^2*cos(b*x + a)^2)*sqrt(b^(-4)))/((2*cos(b*x + a)^3 - cos(b*x + a))*sin(b*x + a)) - 1/4*sqrt(2)*(b^(-4))^(1/4)*arctan(-1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) - sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt
```


$(\sin(bx + a)) - \sqrt{4b^2\sqrt{b^{-4}}}\cos(bx + a)\sin(bx + a) - 2*(\sqrt{2})b^3(b^{-4})^{3/4}\cos(bx + a) + \sqrt{2}b(b^{-4})^{1/4}\sin(bx + a)$
 $)*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)} + 1*((\sqrt{2})b^3(b^{-4})^{3/4}\sin(bx + a) + \sqrt{2}b(b^{-4})^{1/4}\cos(bx + a))*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)}$
 $+ 2*\cos(bx + a)*\sin(bx + a))/(\cos(bx + a)*\sin(bx + a)) - 1/4*\sqrt{2}*(b^{-4})^{1/4}*\arctan(-1/2*((\sqrt{2})b^3(b^{-4})^{3/4}\sin(bx + a) - \sqrt{2}b(b^{-4})^{1/4}\cos(bx + a))*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)})$
 $- \sqrt{4b^2\sqrt{b^{-4}}}\cos(bx + a)\sin(bx + a) + 2*(\sqrt{2})b^3(b^{-4})^{3/4}\cos(bx + a) + \sqrt{2}b(b^{-4})^{1/4}\sin(bx + a)$
 $)*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)} + 1*((\sqrt{2})b^3(b^{-4})^{3/4}\sin(bx + a) + \sqrt{2}b(b^{-4})^{1/4}\cos(bx + a))*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)}$
 $- 2*\cos(bx + a)*\sin(bx + a))/(\cos(bx + a)*\sin(bx + a)) + 1/8*\sqrt{2}*(b^{-4})^{1/4}*\log(4b^2\sqrt{b^{-4}})\cos(bx + a)\sin(bx + a) + 2*(\sqrt{2})b^3(b^{-4})^{3/4}\cos(bx + a) + \sqrt{2}b(b^{-4})^{1/4}\sin(bx + a))*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)}$
 $+ 1 - 1/8*\sqrt{2}*(b^{-4})^{1/4}*\log(4b^2\sqrt{b^{-4}})\cos(bx + a)\sin(bx + a) - 2*(\sqrt{2})b^3(b^{-4})^{3/4}\cos(bx + a) + \sqrt{2}b(b^{-4})^{1/4}\sin(bx + a))*\sqrt{\cos(bx + a)}*\sqrt{\sin(bx + a)} + 1$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\cos(a + bx)}}{\sqrt{\sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(1/2)/sin(b*x+a)**(1/2), x)

[Out] Integral(sqrt(cos(a + b*x))/sqrt(sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(1/2)/sin(b*x+a)^(1/2), x, algorithm="giac")

[Out] integrate(sqrt(cos(b*x + a))/sqrt(sin(b*x + a)), x)

Mupad [B]

time = 1.60, size = 44, normalized size = 0.25

$$-\frac{2 \cos(a + bx)^{3/2} \sqrt{\sin(a + bx)} {}_2F_1\left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \cos(a + bx)^2\right)}{3b (\sin(a + bx)^2)^{1/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^(1/2)/sin(a + b*x)^(1/2),x)
```

```
[Out] -(2*cos(a + b*x)^(3/2)*sin(a + b*x)^(1/2)*hypergeom([3/4, 3/4], 7/4, cos(a + b*x)^2))/(3*b*(sin(a + b*x)^2)^(1/4))
```

$$3.301 \quad \int \frac{\cos^{\frac{3}{2}}(a+bx)}{\sin^{\frac{3}{2}}(a+bx)} dx$$

Optimal. Leaf size=199

$$\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} - \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} - \frac{\log\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right) + \tan(a+bx)}{2\sqrt{2}b}$$

[Out] $\frac{1}{2}\arctan\left(\frac{1-2^{1/2}\sin(bx+a)^{1/2}/\cos(bx+a)^{1/2}}{b2^{1/2}}\right) - \frac{1}{2}\arctan\left(\frac{1+2^{1/2}\sin(bx+a)^{1/2}/\cos(bx+a)^{1/2}}{b2^{1/2}}\right) - \frac{1}{4}\ln\left(\frac{1-2^{1/2}\sin(bx+a)^{1/2}/\cos(bx+a)^{1/2} + \tan(bx+a)}{b2^{1/2}}\right) + \frac{1}{4}\ln\left(\frac{1+2^{1/2}\sin(bx+a)^{1/2}/\cos(bx+a)^{1/2} + \tan(bx+a)}{b2^{1/2}}\right) - \frac{2\cos(bx+a)^{1/2}/b}{\sin(bx+a)^{1/2}}$

Rubi [A]

time = 0.08, antiderivative size = 199, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2647, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} - \frac{\text{ArcTan}\left(\frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + 1\right)}{\sqrt{2}b} - \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} - \frac{\log\left(\tan(a+bx) - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + 1\right)}{2\sqrt{2}b} + \frac{\log\left(\tan(a+bx) + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(3/2)/Sin[a + b*x]^(3/2), x]

[Out] $\text{ArcTan}\left[1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right]/\sqrt{2}b - \text{ArcTan}\left[1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right]/\sqrt{2}b - \text{Log}\left[\frac{1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)}{2\sqrt{2}b}\right] + \text{Log}\left[\frac{1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)}{2\sqrt{2}b}\right] - \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}}$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2647

```
Int[(cos[(e_.) + (f_.)*(x_)])*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_)])^(n_), x_Symbol] := Simp[a*(a*cos[e + f*x])^(m - 1)*((b*sin[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*cos[e + f*x])^(m - 2)*(b*sin[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])
```

Rule 2654

```
Int[(cos[(e_.) + (f_.)*(x_)])*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_)])^m, x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*sin[e + f*x])^(1/k)/(b*cos[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^{\frac{3}{2}}(a+bx)}{\sin^{\frac{3}{2}}(a+bx)} dx &= -\frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} - \int \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} dx \\
&= -\frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} - \frac{2\text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{b} \\
&= -\frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} + \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{b} - \frac{\text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{b} \\
&= -\frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} - \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{2b} - \frac{\text{Subst}\left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{2b} \\
&= -\frac{\log\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)\right)}{2\sqrt{2}b} + \frac{\log\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)\right)}{2\sqrt{2}b} \\
&= \frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} - \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} - \frac{\log\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)\right)}{2\sqrt{2}b} + \frac{\log\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)\right)}{2\sqrt{2}b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 55, normalized size = 0.28

$$-\frac{2\cos^2(a+bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, -\frac{1}{4}; \frac{3}{4}; \sin^2(a+bx)\right)}{b\cos^{\frac{3}{2}}(a+bx)\sqrt{\sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(3/2)/Sin[a + b*x]^(3/2), x]

[Out] (-2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, -1/4, 3/4, Sin[a + b*x]^2])/(b*Cos[a + b*x]^(3/2)*Sqrt[Sin[a + b*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.29, size = 937, normalized size = 4.71

method	result	size
default	Expression too large to display	937

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^(3/2)/sin(b*x+a)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/b*(I*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-I*\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})-\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-\cos(b*x+a)*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})+I*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-I*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})-((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2-1/2*I,1/2*2^{1/2})-((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2}*((-1+\cos(b*x+a))/\sin(b*x+a))^{1/2}*EllipticPi(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{1/2},1/2+1/2*I,1/2*2^{1/2})+2*\cos(b*x+a)*2^{1/2}/\sin(b*x+a)^{1/2}/\cos(b*x+a)^{1/2}*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(3/2)/sin(b*x+a)^(3/2),x, algorithm="maxima")`

[Out] `integrate(cos(b*x + a)^(3/2)/sin(b*x + a)^(3/2), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1545 vs. 2(160) = 320.

time = 27.63, size = 1545, normalized size = 7.76

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(3/2)/sin(b*x+a)^(3/2),x, algorithm="fricas")`

+ a)^(3/2))/(b*cos(b*x + a)^2 - b)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{3}{2}}(a + bx)}{\sin^{\frac{3}{2}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(3/2)/sin(b*x+a)**(3/2),x)

[Out] Integral(cos(a + b*x)**(3/2)/sin(a + b*x)**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(3/2)/sin(b*x+a)^(3/2),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(3/2)/sin(b*x + a)^(3/2), x)

Mupad [B]

time = 1.62, size = 44, normalized size = 0.22

$$-\frac{2 \cos(a + bx)^{5/2} (\sin(a + bx)^2)^{1/4} {}_2F_1\left(\frac{5}{4}, \frac{5}{4}; \frac{9}{4}; \cos(a + bx)^2\right)}{5 b \sqrt{\sin(a + bx)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(3/2)/sin(a + b*x)^(3/2),x)

[Out] -(2*cos(a + b*x)^(5/2)*(sin(a + b*x)^2)^(1/4)*hypergeom([5/4, 5/4], 9/4, cos(a + b*x)^2))/(5*b*sin(a + b*x)^(1/2))

$$3.302 \quad \int \frac{\cos^{\frac{5}{2}}(a+bx)}{\sin^{\frac{5}{2}}(a+bx)} dx$$

Optimal. Leaf size=201

$$\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} + \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} + \frac{\log\left(1 + \cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b}$$

[Out] $-2/3*\cos(b*x+a)^{(3/2)}/b/\sin(b*x+a)^{(3/2)}+1/2*\arctan(-1+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}+1/2*\arctan(1+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}+1/4*\ln(1+\cot(b*x+a)-2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}-1/4*\ln(1+\cot(b*x+a)+2^{(1/2)}*\cos(b*x+a)^{(1/2)}/\sin(b*x+a)^{(1/2)})/b*2^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 201, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2647, 2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} + \frac{\text{ArcTan}\left(\frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{\sqrt{2}b} - \frac{2\cos^{\frac{3}{2}}(a+bx)}{3b\sin^{\frac{3}{2}}(a+bx)} + \frac{\log\left(\cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{2\sqrt{2}b} - \frac{\log\left(\cot(a+bx) + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(5/2)/Sin[a + b*x]^(5/2), x]

[Out] $-(\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Cos}[a + b*x]])/\text{Sqrt}[\text{Sin}[a + b*x]]]/(\text{Sqrt}[2]*b)) + \text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Cos}[a + b*x]])/\text{Sqrt}[\text{Sin}[a + b*x]]]/(\text{Sqrt}[2]*b) + \text{Log}[1 + \text{Cot}[a + b*x] - (\text{Sqrt}[2]*\text{Sqrt}[\text{Cos}[a + b*x]])/\text{Sqrt}[\text{Sin}[a + b*x]]]/(2*\text{Sqrt}[2]*b) - \text{Log}[1 + \text{Cot}[a + b*x] + (\text{Sqrt}[2]*\text{Sqrt}[\text{Cos}[a + b*x]])/\text{Sqrt}[\text{Sin}[a + b*x]]]/(2*\text{Sqrt}[2]*b) - (2*\text{Cos}[a + b*x]^{(3/2)})/(3*b*\text{Sin}[a + b*x]^{(3/2)})$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2647

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[a*(a*cos[e + f*x])^(m - 1)*((b*sin[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*cos[e + f*x])^(m - 2)*(b*sin[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])
```

Rule 2655

```
Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*cos[e + f*x])^(1/k)/(b*sin[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^{\frac{5}{2}}(a+bx)}{\sin^{\frac{5}{2}}(a+bx)} dx &= -\frac{2\cos^{\frac{3}{2}}(a+bx)}{3b\sin^{\frac{3}{2}}(a+bx)} - \int \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}} dx \\
&= -\frac{2\cos^{\frac{3}{2}}(a+bx)}{3b\sin^{\frac{3}{2}}(a+bx)} + \frac{2\text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} \\
&= -\frac{2\cos^{\frac{3}{2}}(a+bx)}{3b\sin^{\frac{3}{2}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} + \frac{\text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{b} \\
&= -\frac{2\cos^{\frac{3}{2}}(a+bx)}{3b\sin^{\frac{3}{2}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2b} + \frac{\text{Subst}\left(\int \frac{1}{1+\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2b} \\
&= \frac{\log\left(1+\cot(a+bx) - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} - \frac{\log\left(1+\cot(a+bx) + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{2\sqrt{2}b} \\
&= -\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} + \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\cos(a+bx)}}{\sqrt{\sin(a+bx)}}\right)}{\sqrt{2}b} + \frac{\log\left(1 + \cot(a+bx)\right)}{b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 57, normalized size = 0.28

$$-\frac{2\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(-\frac{3}{4}, -\frac{3}{4}; \frac{1}{4}; \sin^2(a+bx)\right)}{3b\sqrt{\cos(a+bx)}\sin^{\frac{3}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(5/2)/Sin[a + b*x]^(5/2), x]

[Out] (-2*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[-3/4, -3/4, 1/4, Sin[a + b*x]^2])/(3*b*Sqrt[Cos[a + b*x]]*Sin[a + b*x]^(3/2))

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.10, size = 1261, normalized size = 6.27

method	result	size
default	Expression too large to display	1261

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(b*x+a)^(5/2)/sin(b*x+a)^(5/2),x,method=_RETURNVERBOSE)
[Out] -4/3/b*cos(b*x+a)^(5/2)*(-1+cos(b*x+a))^3*(-3*I*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))+3*I*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))+3*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))+3*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-6*sin(b*x+a)*cos(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))-3*I*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))+3*I*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))+3*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2+1/2*I,1/2*2^(1/2))+3*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticPi(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2-1/2*I,1/2*2^(1/2))-6*sin(b*x+a)*((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2)*((-1+cos(b*x+a))/sin(b*x+a))^(1/2)*EllipticF(((1-cos(b*x+a)+sin(b*x+a))/sin(b*x+a))^(1/2),1/2*2^(1/2))+2*cos(b*x+a)^2*2^(1/2)/sin(b*x+a)^(3/2)/(-1+cos(b*x+a)+sin(b*x+a))^3/(-1+cos(b*x+a)-sin(b*x+a))^3*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)^(5/2)/sin(b*x+a)^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(cos(b*x + a)^(5/2)/sin(b*x + a)^(5/2), x)
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1321 vs. 2(159) = 318.

time = 15.08, size = 1321, normalized size = 6.57

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)^(5/2)/sin(b*x+a)^(5/2),x, algorithm="fricas")
```

```
[Out] 1/24*(6*(sqrt(2)*b*cos(b*x + a)^2 - sqrt(2)*b)*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + (sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 2*cos(b*x + a)*sin(b*x + a) - 4*(b^2*cos(b*x + a)^4 - b^2*cos(b*x + a)^2)*sqrt(b^(-4)))/(2*cos(b*x + a)^3 - cos(b*x + a)*sin(b*x + a)) + 6*(sqrt(2)*b*cos(b*x + a)^2 - sqrt(2)*b)*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1)*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + (sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) - 2*cos(b*x + a)*sin(b*x + a) + 4*(b^2*cos(b*x + a)^4 - b^2*cos(b*x + a)^2)*sqrt(b^(-4)))/(2*cos(b*x + a)^3 - cos(b*x + a)*sin(b*x + a)) - 6*(sqrt(2)*b*cos(b*x + a)^2 - sqrt(2)*b)*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) - sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + sqrt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1)*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 2*cos(b*x + a)*sin(b*x + a)))/(cos(b*x + a)*sin(b*x + a)) - 6*(sqrt(2)*b*cos(b*x + a)^2 - sqrt(2)*b)*(b^(-4))^(1/4)*arctan(1/2*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) - sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + sqrt(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1)*((sqrt(2)*b^3*(b^(-4))^(3/4)*sin(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*cos(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) - 2*cos(b*x + a)*sin(b*x + a)))/(cos(b*x + a)*sin(b*x + a)) - 3*(sqrt(2)*b*cos(b*x + a)^2 - sqrt(2)*b)*(b^(-4))^(1/4)*log(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) + 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))*sqrt(cos(b*x + a))*sqrt(sin(b*x + a)) + 1) + 3*(sqrt(2)*b*cos(b*x + a)^2 - sqrt(2)*b)*(b^(-4))^(1/4)*log(4*b^2*sqrt(b^(-4))*cos(b*x + a)*sin(b*x + a) - 2*(sqrt(2)*b^3*(b^(-4))^(3/4)*cos(b*x + a) + sqrt(2)*b*(b^(-4))^(1/4)*sin(b*x + a))
```

$(-4)^{1/4} \sin(bx + a) \sqrt{\cos(bx + a)} \sqrt{\sin(bx + a) + 1} + 16 \cos(bx + a)^{3/2} \sqrt{\sin(bx + a)} / (b \cos(bx + a)^2 - b)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(5/2)/sin(b*x+a)**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/2)/sin(b*x+a)^(5/2),x, algorithm="giac")

[Out] Timed out

Mupad [B]

time = 1.90, size = 44, normalized size = 0.22

$$\frac{2 \cos(a + bx)^{7/2} (\sin(a + bx)^2)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{7}{4}; \frac{11}{4}; \cos(a + bx)^2\right)}{7 b \sin(a + bx)^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(5/2)/sin(a + b*x)^(5/2),x)

[Out] $-(2 \cos(a + bx)^{7/2} (\sin(a + bx)^2)^{3/4} \text{hypergeom}([7/4, 7/4], 11/4, \cos(a + bx)^2)) / (7 b \sin(a + bx)^{3/2})$

$$3.303 \quad \int \frac{\cos^{\frac{7}{2}}(a+bx)}{\sin^{\frac{7}{2}}(a+bx)} dx$$

Optimal. Leaf size=226

$$\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} + \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} + \frac{\log\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right) + \tan^{-1}\left(\frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{2\sqrt{2}b}$$

[Out] $-2/5*\cos(b*x+a)^{(5/2)}/b/\sin(b*x+a)^{(5/2)}-1/2*\arctan(1-2^{(1/2)}*\sin(b*x+a)^{(1/2)}/\cos(b*x+a)^{(1/2)})/b*2^{(1/2)}+1/2*\arctan(1+2^{(1/2)}*\sin(b*x+a)^{(1/2)}/\cos(b*x+a)^{(1/2)})/b*2^{(1/2)}+1/4*\ln(1-2^{(1/2)}*\sin(b*x+a)^{(1/2)}/\cos(b*x+a)^{(1/2)}+\tan(b*x+a))/b*2^{(1/2)}-1/4*\ln(1+2^{(1/2)}*\sin(b*x+a)^{(1/2)}/\cos(b*x+a)^{(1/2)}+\tan(b*x+a))/b*2^{(1/2)}+2*\cos(b*x+a)^{(1/2)}/b/\sin(b*x+a)^{(1/2)}$

Rubi [A]

time = 0.10, antiderivative size = 226, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2647, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} + \frac{\text{ArcTan}\left(\frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + 1\right)}{\sqrt{2}b} - \frac{2\cos^{\frac{5}{2}}(a+bx)}{5b\sin^{\frac{5}{2}}(a+bx)} + \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} + \frac{\log\left(\tan(a+bx) - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + 1\right)}{2\sqrt{2}b} - \frac{\log\left(\tan(a+bx) + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(7/2)/Sin[a + b*x]^(7/2), x]

[Out] $-(\text{ArcTan}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[a + b*x]])/\text{Sqrt}[\text{Cos}[a + b*x]]]/(\text{Sqrt}[2]*b)) + \text{ArcTan}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[a + b*x]])/\text{Sqrt}[\text{Cos}[a + b*x]]]/(\text{Sqrt}[2]*b) + \text{Log}[1 - (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[a + b*x]])/\text{Sqrt}[\text{Cos}[a + b*x]] + \text{Tan}[a + b*x]]/(2*\text{Sqrt}[2]*b) - \text{Log}[1 + (\text{Sqrt}[2]*\text{Sqrt}[\text{Sin}[a + b*x]])/\text{Sqrt}[\text{Cos}[a + b*x]] + \text{Tan}[a + b*x]]/(2*\text{Sqrt}[2]*b) - (2*\text{Cos}[a + b*x]^{(5/2)})/(5*b*\text{Sin}[a + b*x]^{(5/2)}) + (2*\text{Sqrt}[\text{Cos}[a + b*x]])/(b*\text{Sqrt}[\text{Sin}[a + b*x]])$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2647

```
Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(a*cos[e + f*x])^(m - 1)*((b*sin[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*cos[e + f*x])^(m - 2)*(b*sin[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*sin[e + f*x])^(1/k)/(b*cos[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^{\frac{7}{2}}(a+bx)}{\sin^{\frac{7}{2}}(a+bx)} dx &= -\frac{2\cos^{\frac{5}{2}}(a+bx)}{5b\sin^{\frac{5}{2}}(a+bx)} - \int \frac{\cos^{\frac{3}{2}}(a+bx)}{\sin^{\frac{3}{2}}(a+bx)} dx \\
&= -\frac{2\cos^{\frac{5}{2}}(a+bx)}{5b\sin^{\frac{5}{2}}(a+bx)} + \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} + \int \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} dx \\
&= -\frac{2\cos^{\frac{5}{2}}(a+bx)}{5b\sin^{\frac{5}{2}}(a+bx)} + \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} + \frac{2\text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{b} \\
&= -\frac{2\cos^{\frac{5}{2}}(a+bx)}{5b\sin^{\frac{5}{2}}(a+bx)} + \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} - \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{b} + \dots \\
&= -\frac{2\cos^{\frac{5}{2}}(a+bx)}{5b\sin^{\frac{5}{2}}(a+bx)} + \frac{2\sqrt{\cos(a+bx)}}{b\sqrt{\sin(a+bx)}} + \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{2b} \\
&= \frac{\log\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)\right)}{2\sqrt{2}b} - \frac{\log\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}} + \tan(a+bx)\right)}{2\sqrt{2}b} \\
&= -\frac{\tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} + \frac{\tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{\sin(a+bx)}}{\sqrt{\cos(a+bx)}}\right)}{\sqrt{2}b} + \log\left(1 - \dots\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 57, normalized size = 0.25

$$-\frac{2\cos^2(a+bx)^{3/4} {}_2F_1\left(-\frac{5}{4}, -\frac{5}{4}; -\frac{1}{4}; \sin^2(a+bx)\right)}{5b\cos^{\frac{3}{2}}(a+bx)\sin^{\frac{5}{2}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(7/2)/Sin[a + b*x]^(7/2), x]

[Out] (-2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-5/4, -5/4, -1/4, Sin[a + b*x]^2])/(5*b*Cos[a + b*x]^(3/2)*Sin[a + b*x]^(5/2))

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.12, size = 1934, normalized size = 8.56

method	result	size
--------	--------	------

$$\begin{aligned} &))^{(1/2)}, 1/2+1/2*I, 1/2*2^{(1/2)}-5*I*\cos(b*x+a)^3*((1-\cos(b*x+a)+\sin(b*x+a)) \\ &/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b \\ &*x+a))/\sin(b*x+a))^{(1/2)*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(\\ &(1/2), 1/2-1/2*I, 1/2*2^{(1/2)}-12*\cos(b*x+a)^3*2^{(1/2)}-5*((1-\cos(b*x+a)+\sin(b \\ &*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1 \\ &+\cos(b*x+a))/\sin(b*x+a))^{(1/2)*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b* \\ &x+a))^{(1/2), 1/2+1/2*I, 1/2*2^{(1/2)}-5*((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a)) \\ &^{(1/2)*((-1+\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2)*((-1+\cos(b*x+a))/\sin(b \\ &*x+a))^{(1/2)*\text{EllipticPi}(((1-\cos(b*x+a)+\sin(b*x+a))/\sin(b*x+a))^{(1/2), 1/2-1/ \\ &2*I, 1/2*2^{(1/2)}+10*\cos(b*x+a)*2^{(1/2)}/\sin(b*x+a)^{(5/2)/(-1+\cos(b*x+a)+\sin \\ &(b*x+a))^4/(-1+\cos(b*x+a)-\sin(b*x+a))^4*2^{(1/2)} \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(7/2)/sin(b*x+a)^(7/2),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(7/2)/sin(b*x + a)^(7/2), x)

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 1670 vs. 2(181) = 362.

time = 27.37, size = 1670, normalized size = 7.39

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(7/2)/sin(b*x+a)^(7/2),x, algorithm="fricas")

[Out]
$$\begin{aligned} &-1/80*(32*(6*\cos(b*x + a)^2 - 5)*\text{sqrt}(\cos(b*x + a))*\sin(b*x + a)^{(3/2)} - 20 \\ &*(\text{sqrt}(2)*b*\cos(b*x + a)^4 - 2*\text{sqrt}(2)*b*\cos(b*x + a)^2 + \text{sqrt}(2)*b)*(b^{(-4)} \\ &)^{(1/4)*\arctan(((\text{sqrt}(2)*b^3*(b^{(-4)})^{(3/4)}*\sin(b*x + a) + \text{sqrt}(2)*b*(b^{(-4)} \\ &)^{(1/4)}*\cos(b*x + a))*\text{sqrt}(\cos(b*x + a))*\text{sqrt}(\sin(b*x + a)) + \text{sqrt}(4*b^2* \\ &\text{sqrt}(b^{(-4)})*\cos(b*x + a)*\sin(b*x + a) - 2*(\text{sqrt}(2)*b^3*(b^{(-4)})^{(3/4)}*\sin(\\ &b*x + a) + \text{sqrt}(2)*b*(b^{(-4)})^{(1/4)}*\cos(b*x + a))*\text{sqrt}(\cos(b*x + a))*\text{sqrt}(s \\ &\text{in}(b*x + a)) + 1)*(b^2*\text{sqrt}(b^{(-4)}) + (\text{sqrt}(2)*b^3*(b^{(-4)})^{(3/4)}*\cos(b*x + \\ &a) + \text{sqrt}(2)*b*(b^{(-4)})^{(1/4)}*\sin(b*x + a))*\text{sqrt}(\cos(b*x + a))*\text{sqrt}(\sin(b* \\ &x + a)) + 2*\cos(b*x + a)*\sin(b*x + a)))/(2*\cos(b*x + a)^2 - 1)) - 20*(\text{sqrt}(\\ &2)*b*\cos(b*x + a)^4 - 2*\text{sqrt}(2)*b*\cos(b*x + a)^2 + \text{sqrt}(2)*b)*(b^{(-4)})^{(1/4)} \\ &)*\arctan(((\text{sqrt}(2)*b^3*(b^{(-4)})^{(3/4)}*\sin(b*x + a) + \text{sqrt}(2)*b*(b^{(-4)})^{(1/4)} \\ &)*\cos(b*x + a))*\text{sqrt}(\cos(b*x + a))*\text{sqrt}(\sin(b*x + a)) - \text{sqrt}(4*b^2*\text{sqrt}(b^{(-4)} \\ &)*\cos(b*x + a)*\sin(b*x + a) + 2*(\text{sqrt}(2)*b^3*(b^{(-4)})^{(3/4)}*\sin(b*x + a) \\ &)+ \text{sqrt}(2)*b*(b^{(-4)})^{(1/4)}*\cos(b*x + a))*\text{sqrt}(\cos(b*x + a))*\text{sqrt}(\sin(b*x \\ &+ a)) + 1)*(b^2*\text{sqrt}(b^{(-4)}) - (\text{sqrt}(2)*b^3*(b^{(-4)})^{(3/4)}*\cos(b*x + a) + s \end{aligned}$$

$$\begin{aligned} & \sqrt{2} * b * (b^{-4})^{1/4} * \sin(b*x + a) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} \\ & + 2 * \cos(b*x + a) * \sin(b*x + a) / (2 * \cos(b*x + a)^2 - 1) - 20 * (\sqrt{2} * b * \cos(b*x + a)^4 - 2 * \sqrt{2} * b * \cos(b*x + a)^2 + \sqrt{2} * b * (b^{-4})^{1/4} * \arctan(1/2 * ((\sqrt{2} * b^3 * (b^{-4})^{3/4} * \cos(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \sin(b*x + a)) * \sqrt{4 * b^2 * \sqrt{b^{-4}} * \cos(b*x + a) * \sin(b*x + a)} + 2 * (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \sin(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \cos(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 1) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)}) - (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \cos(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \sin(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} - 2 * \cos(b*x + a) * \sin(b*x + a) + 4 * (b^2 * \cos(b*x + a)^4 - b^2 * \cos(b*x + a)^2) * \sqrt{b^{-4}}) / ((2 * \cos(b*x + a)^3 - \cos(b*x + a)) * \sin(b*x + a)) - 20 * (\sqrt{2} * b * \cos(b*x + a)^4 - 2 * \sqrt{2} * b * \cos(b*x + a)^2 + \sqrt{2} * b * (b^{-4})^{1/4} * \arctan(1/2 * ((\sqrt{2} * b^3 * (b^{-4})^{3/4} * \cos(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \sin(b*x + a)) * \sqrt{4 * b^2 * \sqrt{b^{-4}} * \cos(b*x + a) * \sin(b*x + a)} - 2 * (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \sin(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \cos(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 1) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)}) - (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \cos(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \sin(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 2 * \cos(b*x + a) * \sin(b*x + a) - 4 * (b^2 * \cos(b*x + a)^4 - b^2 * \cos(b*x + a)^2) * \sqrt{b^{-4}}) / ((2 * \cos(b*x + a)^3 - \cos(b*x + a)) * \sin(b*x + a)) + 5 * (\sqrt{2} * b * \cos(b*x + a)^4 - 2 * \sqrt{2} * b * \cos(b*x + a)^2 + \sqrt{2} * b * (b^{-4})^{1/4} * \log(4 * b^2 * \sqrt{b^{-4}} * \cos(b*x + a) * \sin(b*x + a) + 2 * (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \sin(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \cos(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 1) - 5 * (\sqrt{2} * b * \cos(b*x + a)^4 - 2 * \sqrt{2} * b * \cos(b*x + a)^2 + \sqrt{2} * b * (b^{-4})^{1/4} * \log(4 * b^2 * \sqrt{b^{-4}} * \cos(b*x + a) * \sin(b*x + a) - 2 * (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \sin(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \cos(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 1) + 5 * (\sqrt{2} * b * \cos(b*x + a)^4 - 2 * \sqrt{2} * b * \cos(b*x + a)^2 + \sqrt{2} * b * (b^{-4})^{1/4} * \log(1/4 * b^2 * \sqrt{b^{-4}} * \cos(b*x + a) * \sin(b*x + a) + 1/8 * (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \sin(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \cos(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 1/16) - 5 * (\sqrt{2} * b * \cos(b*x + a)^4 - 2 * \sqrt{2} * b * \cos(b*x + a)^2 + \sqrt{2} * b * (b^{-4})^{1/4} * \log(1/4 * b^2 * \sqrt{b^{-4}} * \cos(b*x + a) * \sin(b*x + a) - 1/8 * (\sqrt{2} * b^3 * (b^{-4})^{3/4} * \sin(b*x + a) + \sqrt{2} * b * (b^{-4})^{1/4} * \cos(b*x + a)) * \sqrt{\cos(b*x + a)} * \sqrt{\sin(b*x + a)} + 1/16)) / (b * \cos(b*x + a)^4 - 2 * b * \cos(b*x + a)^2 + b) \end{aligned}$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(7/2)/sin(b*x+a)**(7/2),x)

[Out] Timed out

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(7/2)/sin(b*x+a)^(7/2),x, algorithm="giac")`

[Out] Timed out

Mupad [B]

time = 1.88, size = 44, normalized size = 0.19

$$\frac{2 \cos(a + bx)^{9/2} (\sin(a + bx)^2)^{5/4} {}_2F_1\left(\frac{9}{4}, \frac{9}{4}; \frac{13}{4}; \cos(a + bx)^2\right)}{9 b \sin(a + bx)^{5/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^(7/2)/sin(a + b*x)^(7/2),x)`

[Out] `-(2*cos(a + b*x)^(9/2)*(sin(a + b*x)^2)^(5/4)*hypergeom([9/4, 9/4], 13/4, cos(a + b*x)^2))/(9*b*sin(a + b*x)^(5/2))`

3.304 $\int \cos^4(e + fx) \sqrt[3]{b \sin(e + fx)} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(-\frac{3}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{4/3}}{4bf \sqrt{\cos^2(e + fx)}}$$

[Out] 3/4*cos(f*x+e)*hypergeom([-3/2, 2/3], [5/3], sin(f*x+e)^2)*(b*sin(f*x+e))^(4/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3 \cos(e + fx) (b \sin(e + fx))^{4/3} {}_2F_1\left(-\frac{3}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right)}{4bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^4*(b*Sin[e + f*x])^(1/3), x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[-3/2, 2/3, 5/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(4/3))/(4*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2])*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \cos^4(e + fx) \sqrt[3]{b \sin(e + fx)} dx = \frac{3 \cos(e + fx) {}_2F_1\left(-\frac{3}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{4/3}}{4bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(-\frac{3}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) \sqrt[3]{b \sin(e + fx)} \tan(e + fx)}{4f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[e + f*x]^4*(b*SIN[e + f*x])^(1/3),x]
```

```
[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-3/2, 2/3, 5/3, Sin[e + f*x]^2]*(b*SIN[e + f*x])^(1/3)*Tan[e + f*x])/(4*f)
```

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (\cos^4(fx + e)) (b \sin(fx + e))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(f*x+e)^4*(b*sin(f*x+e))^(1/3),x)
```

```
[Out] int(cos(f*x+e)^4*(b*sin(f*x+e))^(1/3),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)^4*(b*sin(f*x+e))^(1/3),x, algorithm="maxima")
```

```
[Out] integrate((b*sin(f*x + e))^(1/3)*cos(f*x + e)^4, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)^4*(b*sin(f*x+e))^(1/3),x, algorithm="fricas")
```

```
[Out] integral((b*sin(f*x + e))^(1/3)*cos(f*x + e)^4, x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sin(e + fx)} \cos^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)**4*(b*sin(f*x+e))**(1/3),x)
```

```
[Out] Integral((b*sin(e + f*x))**(1/3)*cos(e + f*x)**4, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4*(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(1/3)*cos(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(e + f x)^4 (b \sin(e + f x))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^4*(b*sin(e + f*x))^(1/3),x)

[Out] int(cos(e + f*x)^4*(b*sin(e + f*x))^(1/3), x)

3.305 $\int \cos^2(e + fx) \sqrt[3]{b \sin(e + fx)} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(-\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{4/3}}{4bf \sqrt{\cos^2(e + fx)}}$$

[Out] 3/4*cos(f*x+e)*hypergeom([-1/2, 2/3], [5/3], sin(f*x+e)^2)*(b*sin(f*x+e))^(4/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3 \cos(e + fx) (b \sin(e + fx))^{4/3} {}_2F_1\left(-\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right)}{4bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^2*(b*Sin[e + f*x])^(1/3), x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[-1/2, 2/3, 5/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(4/3))/(4*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \cos^2(e + fx) \sqrt[3]{b \sin(e + fx)} dx = \frac{3 \cos(e + fx) {}_2F_1\left(-\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{4/3}}{4bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(-\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) \sqrt[3]{b \sin(e + fx)} \tan(e + fx)}{4f}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^2*(b*SIN[e + f*x])^(1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/2, 2/3, 5/3, Sin[e + f*x]^2]*(b*SIN[e + f*x])^(1/3)*Tan[e + f*x])/(4*f)

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (\cos^2(fx + e)) (b \sin(fx + e))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^2*(b*sin(f*x+e))^(1/3),x)

[Out] int(cos(f*x+e)^2*(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2*(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(1/3)*cos(f*x + e)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2*(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(1/3)*cos(f*x + e)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sin(e + fx)} \cos^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**2*(b*sin(f*x+e))**(1/3),x)

[Out] Integral((b*sin(e + f*x))**(1/3)*cos(e + f*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(f*x+e)^2*(b*sin(f*x+e))^(1/3),x, algorithm="giac")``[Out] integrate((b*sin(f*x + e))^(1/3)*cos(f*x + e)^2, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(e + f x)^2 (b \sin(e + f x))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(e + f*x)^2*(b*sin(e + f*x))^(1/3),x)``[Out] int(cos(e + f*x)^2*(b*sin(e + f*x))^(1/3), x)`

3.306 $\int \sqrt[3]{b \sin(e + fx)} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{4/3}}{4bf \sqrt{\cos^2(e + fx)}}$$

[Out] $3/4 * \cos(f*x+e) * \text{hypergeom}([1/2, 2/3], [5/3], \sin(f*x+e)^2) * (b * \sin(f*x+e))^{4/3} / b / f / (\cos(f*x+e)^2)^{1/2}$

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(e + fx) (b \sin(e + fx))^{4/3} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right)}{4bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sin[e + f*x])^(1/3), x]

[Out] $(3 * \text{Cos}[e + f*x] * \text{Hypergeometric2F1}[1/2, 2/3, 5/3, \text{Sin}[e + f*x]^2] * (b * \text{Sin}[e + f*x])^{4/3}) / (4 * b * f * \text{Sqrt}[\text{Cos}[e + f*x]^2])$

Rule 2722

Int[((b_.) * sin[(c_.) + (d_.) * (x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x] * ((b * Sin[c + d*x])^(n + 1) / (b * d * (n + 1) * Sqrt[Cos[c + d*x]^2])) * Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \sqrt[3]{b \sin(e + fx)} dx = \frac{3 \cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{4/3}}{4bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{1}{2}, \frac{2}{3}; \frac{5}{3}; \sin^2(e + fx)\right) \sqrt[3]{b \sin(e + fx)} \tan(e + fx)}{4f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sin[e + f*x])^(1/3),x]

[Out] (3*sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/2, 2/3, 5/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(1/3)*Tan[e + f*x])/(4*f)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sin(fx + e))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(f*x+e))^(1/3),x)

[Out] int((b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(1/3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sin(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))**(1/3),x)

[Out] Integral((b*sin(e + f*x))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int (b \sin(e + f x))^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(e + f*x))^(1/3),x)

[Out] int((b*sin(e + f*x))^(1/3), x)

3.307 $\int \sec^2(e + fx) \sqrt[3]{b \sin(e + fx)} dx$

Optimal. Leaf size=58

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{2}{3}, \frac{3}{2}; \frac{5}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{4/3}}{4bf}$$

[Out] 3/4*hypergeom([2/3, 3/2], [5/3], sin(f*x+e)^2)*sec(f*x+e)*(b*sin(f*x+e))^(4/3)*(cos(f*x+e)^2)^(1/2)/b/f

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \sin(e + fx))^{4/3} {}_2F_1\left(\frac{2}{3}, \frac{3}{2}; \frac{5}{3}; \sin^2(e + fx)\right)}{4bf}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^2*(b*Sin[e + f*x])^(1/3), x]

[Out] (3*sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[2/3, 3/2, 5/3, Sin[e + f*x]^2]*Sec[e + f*x]*(b*Sin[e + f*x])^(4/3))/(4*b*f)

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \sec^2(e + fx) \sqrt[3]{b \sin(e + fx)} dx = \frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{2}{3}, \frac{3}{2}; \frac{5}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{4/3}}{4bf}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{2}{3}, \frac{3}{2}; \frac{5}{3}; \sin^2(e + fx)\right) \sqrt[3]{b \sin(e + fx)} \tan(e + fx)}{4f}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[e + f*x]^2*(b*SIN[e + f*x])^(1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[2/3, 3/2, 5/3, Sin[e + f*x]^2]*(b*SIN[e + f*x])^(1/3)*Tan[e + f*x])/(4*f)

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int (\sec^2(fx + e)) (b \sin(fx + e))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^2*(b*sin(f*x+e))^(1/3),x)

[Out] int(sec(f*x+e)^2*(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2*(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(1/3)*sec(f*x + e)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2*(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(1/3)*sec(f*x + e)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{b \sin(e + fx)} \sec^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)**2*(b*sin(f*x+e))**(1/3),x)

[Out] Integral((b*sin(e + f*x))**(1/3)*sec(e + f*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2*(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(1/3)*sec(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \sin(e + f x))^{1/3}}{\cos(e + f x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(e + f*x))^(1/3)/cos(e + f*x)^2,x)

[Out] int((b*sin(e + f*x))^(1/3)/cos(e + f*x)^2, x)

3.308 $\int \sec^4(e + fx) \sqrt[3]{b \sin(e + fx)} dx$

Optimal. Leaf size=58

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{2}{3}, \frac{5}{2}; \frac{5}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{4/3}}{4bf}$$

[Out] $3/4*\text{hypergeom}([2/3, 5/2], [5/3], \sin(f*x+e)^2)*\sec(f*x+e)*(b*\sin(f*x+e))^{4/3}*(\cos(f*x+e)^2)^{(1/2)}/b/f$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \sin(e + fx))^{4/3} {}_2F_1\left(\frac{2}{3}, \frac{5}{2}; \frac{5}{3}; \sin^2(e + fx)\right)}{4bf}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[e + f*x]^4*(b*\text{Sin}[e + f*x])^{(1/3)}, x]$

[Out] $(3*\text{Sqrt}[\text{Cos}[e + f*x]^2]*\text{Hypergeometric2F1}[2/3, 5/2, 5/3, \text{Sin}[e + f*x]^2]*\text{Sec}[e + f*x]*(b*\text{Sin}[e + f*x])^{(4/3)})/(4*b*f)$

Rule 2657

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Cos}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}*((a*\text{Sin}[e + f*x])^{(m + 1)})/(a*f*(m + 1)*(\text{Cos}[e + f*x]^2)^{\text{FracPart}[(n - 1)/2]})*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Sin}[e + f*x]^2], x] /;$ $\text{FreeQ}\{a, b, e, f, m, n\}, x]$

Rubi steps

$$\int \sec^4(e + fx) \sqrt[3]{b \sin(e + fx)} dx = \frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{2}{3}, \frac{5}{2}; \frac{5}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{4/3}}{4bf}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{2}{3}, \frac{5}{2}; \frac{5}{3}; \sin^2(e + fx)\right) \sqrt[3]{b \sin(e + fx)} \tan(e + fx)}{4f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[e + f*x]^4*(b*SIN[e + f*x])^(1/3),x]
```

```
[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[2/3, 5/2, 5/3, Sin[e + f*x]^2]*(b
*SIN[e + f*x])^(1/3)*Tan[e + f*x])/(4*f)
```

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int (\sec^4(fx + e)) (b \sin(fx + e))^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(f*x+e)^4*(b*sin(f*x+e))^(1/3),x)
```

```
[Out] int(sec(f*x+e)^4*(b*sin(f*x+e))^(1/3),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(f*x+e)^4*(b*sin(f*x+e))^(1/3),x, algorithm="maxima")
```

```
[Out] integrate((b*sin(f*x + e))^(1/3)*sec(f*x + e)^4, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(f*x+e)^4*(b*sin(f*x+e))^(1/3),x, algorithm="fricas")
```

```
[Out] integral((b*sin(f*x + e))^(1/3)*sec(f*x + e)^4, x)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(f*x+e)**4*(b*sin(f*x+e))**(1/3),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4*(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(1/3)*sec(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \sin(e + f x))^{1/3}}{\cos(e + f x)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(e + f*x))^(1/3)/cos(e + f*x)^4,x)

[Out] int((b*sin(e + f*x))^(1/3)/cos(e + f*x)^4, x)

$$3.309 \quad \int \cos^4(e + fx)(b \sin(e + fx))^{5/3} dx$$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(-\frac{3}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{8/3}}{8bf \sqrt{\cos^2(e + fx)}}$$

[Out] 3/8*cos(f*x+e)*hypergeom([-3/2, 4/3], [7/3], sin(f*x+e)^2)*(b*sin(f*x+e))^(8/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3 \cos(e + fx)(b \sin(e + fx))^{8/3} {}_2F_1\left(-\frac{3}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right)}{8bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^4*(b*Sin[e + f*x])^(5/3), x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[-3/2, 4/3, 7/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(8/3))/(8*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \cos^4(e + fx)(b \sin(e + fx))^{5/3} dx = \frac{3 \cos(e + fx) {}_2F_1\left(-\frac{3}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{8/3}}{8bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(-\frac{3}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{5/3} \tan(e + fx)}{8f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[e + f*x]^4*(b*Sin[e + f*x])^(5/3),x]
```

```
[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-3/2, 4/3, 7/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(5/3)*Tan[e + f*x])/(8*f)
```

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (\cos^4(fx + e)) (b \sin(fx + e))^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(f*x+e)^4*(b*sin(f*x+e))^(5/3),x)
```

```
[Out] int(cos(f*x+e)^4*(b*sin(f*x+e))^(5/3),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)^4*(b*sin(f*x+e))^(5/3),x, algorithm="maxima")
```

```
[Out] integrate((b*sin(f*x + e))^(5/3)*cos(f*x + e)^4, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)^4*(b*sin(f*x+e))^(5/3),x, algorithm="fricas")
```

```
[Out] integral((b*sin(f*x + e))^(2/3)*b*cos(f*x + e)^4*sin(f*x + e), x)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)**4*(b*sin(f*x+e))**(5/3),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 4846 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(f*x+e)^4*(b*sin(f*x+e))^(5/3),x, algorithm="giac")``[Out] integrate((b*sin(f*x + e))^(5/3)*cos(f*x + e)^4, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(e + f x)^4 (b \sin(e + f x))^{5/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(e + f*x)^4*(b*sin(e + f*x))^(5/3),x)``[Out] int(cos(e + f*x)^4*(b*sin(e + f*x))^(5/3), x)`

3.310 $\int \cos^2(e + fx)(b \sin(e + fx))^{5/3} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(-\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{8/3}}{8bf \sqrt{\cos^2(e + fx)}}$$

[Out] 3/8*cos(f*x+e)*hypergeom([-1/2, 4/3], [7/3], sin(f*x+e)^2)*(b*sin(f*x+e))^(8/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3 \cos(e + fx)(b \sin(e + fx))^{8/3} {}_2F_1\left(-\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right)}{8bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^2*(b*Sin[e + f*x])^(5/3),x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[-1/2, 4/3, 7/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(8/3))/(8*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \cos^2(e + fx)(b \sin(e + fx))^{5/3} dx = \frac{3 \cos(e + fx) {}_2F_1\left(-\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{8/3}}{8bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(-\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{5/3} \tan(e + fx)}{8f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Cos[e + f*x]^2*(b*SIN[e + f*x])^(5/3),x]
```

```
[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/2, 4/3, 7/3, Sin[e + f*x]^2]*(b*SIN[e + f*x])^(5/3)*Tan[e + f*x])/(8*f)
```

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (\cos^2(fx + e)) (b \sin(fx + e))^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(f*x+e)^2*(b*sin(f*x+e))^(5/3),x)
```

```
[Out] int(cos(f*x+e)^2*(b*sin(f*x+e))^(5/3),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)^2*(b*sin(f*x+e))^(5/3),x, algorithm="maxima")
```

```
[Out] integrate((b*sin(f*x + e))^(5/3)*cos(f*x + e)^2, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)^2*(b*sin(f*x+e))^(5/3),x, algorithm="fricas")
```

```
[Out] integral((b*sin(f*x + e))^(2/3)*b*cos(f*x + e)^2*sin(f*x + e), x)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(f*x+e)**2*(b*sin(f*x+e))**(5/3),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3061 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2*(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(5/3)*cos(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \cos(e + f x)^2 (b \sin(e + f x))^{5/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^2*(b*sin(e + f*x))^(5/3),x)

[Out] int(cos(e + f*x)^2*(b*sin(e + f*x))^(5/3), x)

3.311 $\int (b \sin(e + fx))^{5/3} dx$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{8/3}}{8bf \sqrt{\cos^2(e + fx)}}$$

[Out] 3/8*cos(f*x+e)*hypergeom([1/2, 4/3], [7/3], sin(f*x+e)^2)*(b*sin(f*x+e))^(8/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(e + fx) (b \sin(e + fx))^{8/3} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right)}{8bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sin[e + f*x])^(5/3), x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[1/2, 4/3, 7/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(8/3))/(8*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (b \sin(e + fx))^{5/3} dx = \frac{3 \cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{8/3}}{8bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{1}{2}, \frac{4}{3}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{5/3} \tan(e + fx)}{8f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sin[e + f*x])^(5/3),x]

[Out] (3*sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/2, 4/3, 7/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(5/3)*Tan[e + f*x])/(8*f)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sin(fx + e))^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(f*x+e))^(5/3),x)

[Out] int((b*sin(f*x+e))^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(5/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)*b*sin(f*x + e), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sin(e + fx))^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))**(5/3),x)

[Out] Integral((b*sin(e + f*x))**(5/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(5/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int (b \sin(e + f x))^{5/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(e + f*x))^(5/3),x)

[Out] int((b*sin(e + f*x))^(5/3), x)

3.312 $\int \sec^2(e + fx)(b \sin(e + fx))^{5/3} dx$

Optimal. Leaf size=58

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{4}{3}, \frac{3}{2}; \frac{7}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{8/3}}{8bf}$$

[Out] 3/8*hypergeom([4/3, 3/2], [7/3], sin(f*x+e)^2)*sec(f*x+e)*(b*sin(f*x+e))^(8/3)*(cos(f*x+e)^2)^(1/2)/b/f

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \sin(e + fx))^{8/3} {}_2F_1\left(\frac{4}{3}, \frac{3}{2}; \frac{7}{3}; \sin^2(e + fx)\right)}{8bf}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^2*(b*Sin[e + f*x])^(5/3), x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[4/3, 3/2, 7/3, Sin[e + f*x]^2]*Sec[e + f*x]*(b*Sin[e + f*x])^(8/3))/(8*b*f)

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \sec^2(e + fx)(b \sin(e + fx))^{5/3} dx = \frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{4}{3}, \frac{3}{2}; \frac{7}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{8/3}}{8bf}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{4}{3}, \frac{3}{2}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{5/3} \tan(e + fx)}{8f}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[e + f*x]^2*(b*SIN[e + f*x])^(5/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[4/3, 3/2, 7/3, Sin[e + f*x]^2]*(b*SIN[e + f*x])^(5/3)*Tan[e + f*x])/(8*f)

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int (\sec^2(fx + e)) (b \sin(fx + e))^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^2*(b*sin(f*x+e))^(5/3),x)

[Out] int(sec(f*x+e)^2*(b*sin(f*x+e))^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2*(b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(5/3)*sec(f*x + e)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2*(b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)*b*sec(f*x + e)^2*sin(f*x + e), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)**2*(b*sin(f*x+e))**(5/3),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2*(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(5/3)*sec(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \sin(e + f x))^{5/3}}{\cos(e + f x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(e + f*x))^(5/3)/cos(e + f*x)^2,x)

[Out] int((b*sin(e + f*x))^(5/3)/cos(e + f*x)^2, x)

3.313 $\int \sec^4(e + fx)(b \sin(e + fx))^{5/3} dx$

Optimal. Leaf size=58

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{4}{3}, \frac{5}{2}; \frac{7}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{8/3}}{8bf}$$

[Out] 3/8*hypergeom([4/3, 5/2], [7/3], sin(f*x+e)^2)*sec(f*x+e)*(b*sin(f*x+e))^(8/3)*(cos(f*x+e)^2)^(1/2)/b/f

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3\sqrt{\cos^2(e + fx)} \sec(e + fx)(b \sin(e + fx))^{8/3} {}_2F_1\left(\frac{4}{3}, \frac{5}{2}; \frac{7}{3}; \sin^2(e + fx)\right)}{8bf}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^4*(b*Sin[e + f*x])^(5/3), x]

[Out] (3*sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[4/3, 5/2, 7/3, Sin[e + f*x]^2]*Sec[e + f*x]*(b*Sin[e + f*x])^(8/3))/(8*b*f)

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \sec^4(e + fx)(b \sin(e + fx))^{5/3} dx = \frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{4}{3}, \frac{5}{2}; \frac{7}{3}; \sin^2(e + fx)\right) \sec(e + fx)(b \sin(e + fx))^{8/3}}{8bf}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{4}{3}, \frac{5}{2}; \frac{7}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{5/3} \tan(e + fx)}{8f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sec[e + f*x]^4*(b*SIN[e + f*x])^(5/3),x]
```

```
[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[4/3, 5/2, 7/3, Sin[e + f*x]^2]*(b
*SIN[e + f*x])^(5/3)*Tan[e + f*x])/(8*f)
```

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (\sec^4(fx + e)) (b \sin(fx + e))^{\frac{5}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sec(f*x+e)^4*(b*sin(f*x+e))^(5/3),x)
```

```
[Out] int(sec(f*x+e)^4*(b*sin(f*x+e))^(5/3),x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(f*x+e)^4*(b*sin(f*x+e))^(5/3),x, algorithm="maxima")
```

```
[Out] integrate((b*sin(f*x + e))^(5/3)*sec(f*x + e)^4, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(f*x+e)^4*(b*sin(f*x+e))^(5/3),x, algorithm="fricas")
```

```
[Out] integral((b*sin(f*x + e))^(2/3)*b*sec(f*x + e)^4*sin(f*x + e), x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sec(f*x+e)**4*(b*sin(f*x+e))**(5/3),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4*(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(5/3)*sec(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(b \sin(e + f x))^{5/3}}{\cos(e + f x)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sin(e + f*x))^(5/3)/cos(e + f*x)^4,x)

[Out] int((b*sin(e + f*x))^(5/3)/cos(e + f*x)^4, x)

$$3.314 \quad \int \frac{\cos^4(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{3}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right) (b \sin(e+fx))^{2/3}}{2bf \sqrt{\cos^2(e+fx)}}$$

[Out] 3/2*cos(f*x+e)*hypergeom([-3/2, 1/3], [4/3], sin(f*x+e)^2)*(b*sin(f*x+e))^(2/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3 \cos(e+fx) (b \sin(e+fx))^{2/3} {}_2F_1\left(-\frac{3}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^4/(b*Sin[e + f*x])^(1/3), x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[-3/2, 1/3, 4/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(2/3))/(2*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{\cos^4(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx = \frac{3 \cos(e+fx) {}_2F_1\left(-\frac{3}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right) (b \sin(e+fx))^{2/3}}{2bf \sqrt{\cos^2(e+fx)}}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{3}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f \sqrt[3]{b \sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^4/(b*Sin[e + f*x])^(1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-3/2, 1/3, 4/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(1/3))

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int \frac{\cos^4(fx + e)}{(b \sin(fx + e))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^4/(b*sin(f*x+e))^(1/3),x)

[Out] int(cos(f*x+e)^4/(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4/(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(f*x + e)^4/(b*sin(f*x + e))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4/(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)*cos(f*x + e)^4/(b*sin(f*x + e)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^4(e + fx)}{\sqrt[3]{b \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**4/(b*sin(f*x+e))**(1/3),x)

[Out] Integral(cos(e + f*x)**4/(b*sin(e + f*x))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4/(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate(cos(f*x + e)^4/(b*sin(f*x + e))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(e + f x)^4}{(b \sin(e + f x))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^4/(b*sin(e + f*x))^(1/3),x)

[Out] int(cos(e + f*x)^4/(b*sin(e + f*x))^(1/3), x)

$$3.315 \quad \int \frac{\cos^2(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right) (b \sin(e+fx))^{2/3}}{2bf \sqrt{\cos^2(e+fx)}}$$

[Out] 3/2*cos(f*x+e)*hypergeom([-1/2, 1/3],[4/3],sin(f*x+e)^2)*(b*sin(f*x+e))^(2/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3 \cos(e+fx) (b \sin(e+fx))^{2/3} {}_2F_1\left(-\frac{1}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^2/(b*Sin[e + f*x])^(1/3),x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[-1/2, 1/3, 4/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(2/3))/(2*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2657

Int[(cos[(e_) + (f_)*(x_)])*(b_)]^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{\cos^2(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx = \frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right) (b \sin(e+fx))^{2/3}}{2bf \sqrt{\cos^2(e+fx)}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{2}, \frac{1}{3}; \frac{4}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f \sqrt[3]{b \sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^2/(b*Sin[e + f*x])^(1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/2, 1/3, 4/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(1/3))

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(fx + e)}{(b \sin(fx + e))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^2/(b*sin(f*x+e))^(1/3),x)

[Out] int(cos(f*x+e)^2/(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2/(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate(cos(f*x + e)^2/(b*sin(f*x + e))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2/(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)*cos(f*x + e)^2/(b*sin(f*x + e)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(e + fx)}{\sqrt[3]{b \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**2/(b*sin(f*x+e))**(1/3),x)

[Out] Integral(cos(e + f*x)**2/(b*sin(e + f*x))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2/(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate(cos(f*x + e)^2/(b*sin(f*x + e))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(e + f x)^2}{(b \sin(e + f x))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^2/(b*sin(e + f*x))^(1/3),x)

[Out] int(cos(e + f*x)^2/(b*sin(e + f*x))^(1/3), x)

$$3.316 \quad \int \frac{1}{\sqrt[3]{b \sin(e + fx)}} dx$$

Optimal. Leaf size=58

$$\frac{3 \cos(e + fx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{2/3}}{2bf \sqrt{\cos^2(e + fx)}}$$

[Out] 3/2*cos(f*x+e)*hypergeom([1/3, 1/2],[4/3],sin(f*x+e)^2)*(b*sin(f*x+e))^(2/3)/b/f/(cos(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$\frac{3 \cos(e + fx) (b \sin(e + fx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(e + fx)\right)}{2bf \sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sin[e + f*x])^(-1/3),x]

[Out] (3*Cos[e + f*x]*Hypergeometric2F1[1/3, 1/2, 4/3, Sin[e + f*x]^2]*(b*Sin[e + f*x])^(2/3))/(2*b*f*Sqrt[Cos[e + f*x]^2])

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{\sqrt[3]{b \sin(e + fx)}} dx = \frac{3 \cos(e + fx) {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(e + fx)\right) (b \sin(e + fx))^{2/3}}{2bf \sqrt{\cos^2(e + fx)}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3 \sqrt{\cos^2(e + fx)} {}_2F_1\left(\frac{1}{3}, \frac{1}{2}; \frac{4}{3}; \sin^2(e + fx)\right) \tan(e + fx)}{2f \sqrt[3]{b \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sin[e + f*x])^(-1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/3, 1/2, 4/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(1/3))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sin(fx + e))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sin(f*x+e))^(1/3),x)

[Out] int(1/(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(-1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)/(b*sin(f*x + e)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt[3]{b \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))**(1/3),x)

[Out] Integral((b*sin(e + f*x))**(-1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(b \sin(e + f x))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sin(e + f*x))^(1/3),x)

[Out] int(1/(b*sin(e + f*x))^(1/3), x)

$$3.317 \quad \int \frac{\sec^2(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$$

Optimal. Leaf size=58

$$\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(\frac{1}{3}, \frac{3}{2}; \frac{4}{3}; \sin^2(e+fx)\right) \sec(e+fx)(b \sin(e+fx))^{2/3}}{2bf}$$

[Out] 3/2*hypergeom([1/3, 3/2],[4/3],sin(f*x+e)^2)*sec(f*x+e)*(b*sin(f*x+e))^(2/3)*(cos(f*x+e)^2)^(1/2)/b/f

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3\sqrt{\cos^2(e+fx)} \sec(e+fx)(b \sin(e+fx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{3}{2}; \frac{4}{3}; \sin^2(e+fx)\right)}{2bf}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^2/(b*Sin[e + f*x])^(1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/3, 3/2, 4/3, Sin[e + f*x]^2]*Sec[e + f*x]*(b*Sin[e + f*x])^(2/3))/(2*b*f)

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{\sec^2(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx = \frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(\frac{1}{3}, \frac{3}{2}; \frac{4}{3}; \sin^2(e+fx)\right) \sec(e+fx)(b \sin(e+fx))^{2/3}}{2bf}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(\frac{1}{3}, \frac{3}{2}; \frac{4}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f\sqrt[3]{b \sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[e + f*x]^2/(b*Sin[e + f*x])^(1/3),x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/3, 3/2, 4/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(1/3))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(fx + e)}{(b \sin(fx + e))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^2/(b*sin(f*x+e))^(1/3),x)

[Out] int(sec(f*x+e)^2/(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2/(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(f*x + e)^2/(b*sin(f*x + e))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2/(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)*sec(f*x + e)^2/(b*sin(f*x + e)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(e + fx)}{\sqrt[3]{b \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)**2/(b*sin(f*x+e))**(1/3),x)

[Out] Integral(sec(e + f*x)**2/(b*sin(e + f*x))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2/(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate(sec(f*x + e)^2/(b*sin(f*x + e))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(e + f x)^2 (b \sin(e + f x))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(e + f*x)^2*(b*sin(e + f*x))^(1/3)),x)

[Out] int(1/(cos(e + f*x)^2*(b*sin(e + f*x))^(1/3)), x)

$$3.318 \quad \int \frac{\sec^4(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx$$

Optimal. Leaf size=58

$$\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(\frac{1}{3}, \frac{5}{2}; \frac{4}{3}; \sin^2(e+fx)\right) \sec(e+fx)(b \sin(e+fx))^{2/3}}{2bf}$$

[Out] 3/2*hypergeom([1/3, 5/2], [4/3], sin(f*x+e)^2)*sec(f*x+e)*(b*sin(f*x+e))^(2/3)*(cos(f*x+e)^2)^(1/2)/b/f

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$\frac{3\sqrt{\cos^2(e+fx)} \sec(e+fx)(b \sin(e+fx))^{2/3} {}_2F_1\left(\frac{1}{3}, \frac{5}{2}; \frac{4}{3}; \sin^2(e+fx)\right)}{2bf}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^4/(b*Sin[e + f*x])^(1/3), x]

[Out] (3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/3, 5/2, 4/3, Sin[e + f*x]^2]*Sec[e + f*x]*(b*Sin[e + f*x])^(2/3))/(2*b*f)

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{\sec^4(e+fx)}{\sqrt[3]{b \sin(e+fx)}} dx = \frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(\frac{1}{3}, \frac{5}{2}; \frac{4}{3}; \sin^2(e+fx)\right) \sec(e+fx)(b \sin(e+fx))^{2/3}}{2bf}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(\frac{1}{3}, \frac{5}{2}; \frac{4}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f \sqrt[3]{b \sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[e + f*x]^4/(b*Sin[e + f*x])^(1/3),x]

[Out] (3*sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[1/3, 5/2, 4/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(1/3))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(fx + e)}{(b \sin(fx + e))^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^4/(b*sin(f*x+e))^(1/3),x)

[Out] int(sec(f*x+e)^4/(b*sin(f*x+e))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4/(b*sin(f*x+e))^(1/3),x, algorithm="maxima")

[Out] integrate(sec(f*x + e)^4/(b*sin(f*x + e))^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4/(b*sin(f*x+e))^(1/3),x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^(2/3)*sec(f*x + e)^4/(b*sin(f*x + e)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(e + fx)}{\sqrt[3]{b \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)**4/(b*sin(f*x+e))**(1/3),x)

[Out] Integral(sec(e + f*x)**4/(b*sin(e + f*x))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4/(b*sin(f*x+e))^(1/3),x, algorithm="giac")

[Out] integrate(sec(f*x + e)^4/(b*sin(f*x + e))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(e + f x)^4 (b \sin(e + f x))^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(e + f*x)^4*(b*sin(e + f*x))^(1/3)),x)

[Out] int(1/(cos(e + f*x)^4*(b*sin(e + f*x))^(1/3)), x)

$$3.319 \quad \int \frac{\cos^4(e+fx)}{(b \sin(e+fx))^{5/3}} dx$$

Optimal. Leaf size=58

$$-\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{3}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

[Out] $-3/2*\cos(f*x+e)*\text{hypergeom}([-3/2, -1/3], [2/3], \sin(f*x+e)^2)/b/f/(b*\sin(f*x+e))^{\wedge}(2/3)/(\cos(f*x+e)^2)^{\wedge}(1/2)$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$-\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{3}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[e + f*x]^4/(b*\text{Sin}[e + f*x])^{\wedge}(5/3), x]$

[Out] $(-3*\text{Cos}[e + f*x]*\text{Hypergeometric2F1}[-3/2, -1/3, 2/3, \text{Sin}[e + f*x]^2])/(2*b*f*\text{Sqrt}[\text{Cos}[e + f*x]^2]*(b*\text{Sin}[e + f*x])^{\wedge}(2/3))$

Rule 2657

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{\wedge}(n_))*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{\wedge}(m_), x_Symbol] :> \text{Simp}[b^{\wedge}(2*\text{IntPart}[(n - 1)/2] + 1)*(b*\text{Cos}[e + f*x])^{\wedge}(2*\text{FracPart}[(n - 1)/2])*((a*\text{Sin}[e + f*x])^{\wedge}(m + 1)/(a*f*(m + 1)*(\text{Cos}[e + f*x]^2)^{\wedge}\text{FracPart}[(n - 1)/2]))*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Sin}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x]$

Rubi steps

$$\int \frac{\cos^4(e+fx)}{(b \sin(e+fx))^{5/3}} dx = -\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{3}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{3}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f(b \sin(e+fx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^4/(b*Sin[e + f*x])^(5/3),x]

[Out] (-3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-3/2, -1/3, 2/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(5/3))

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int \frac{\cos^4(fx + e)}{(b \sin(fx + e))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^4/(b*sin(f*x+e))^(5/3),x)

[Out] int(cos(f*x+e)^4/(b*sin(f*x+e))^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4/(b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] integrate(cos(f*x + e)^4/(b*sin(f*x + e))^(5/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4/(b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral(-(b*sin(f*x + e))^(1/3)*cos(f*x + e)^4/(b^2*cos(f*x + e)^2 - b^2), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**4/(b*sin(f*x+e))**(5/3),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^4/(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate(cos(f*x + e)^4/(b*sin(f*x + e))^(5/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(e + f x)^4}{(b \sin(e + f x))^{5/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^4/(b*sin(e + f*x))^(5/3),x)

[Out] int(cos(e + f*x)^4/(b*sin(e + f*x))^(5/3), x)

$$3.320 \quad \int \frac{\cos^2(e+fx)}{(b \sin(e+fx))^{5/3}} dx$$

Optimal. Leaf size=58

$$-\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

[Out] -3/2*cos(f*x+e)*hypergeom([-1/2, -1/3], [2/3], sin(f*x+e)^2)/b/f/(b*sin(f*x+e))^2/3/(cos(f*x+e)^2)^1/2

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$-\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[Cos[e + f*x]^2/(b*Sin[e + f*x])^(5/3), x]

[Out] (-3*Cos[e + f*x]*Hypergeometric2F1[-1/2, -1/3, 2/3, Sin[e + f*x]^2])/(2*b*f*sqrt[Cos[e + f*x]^2]*(b*Sin[e + f*x])^(2/3))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{\cos^2(e+fx)}{(b \sin(e+fx))^{5/3}} dx = -\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$-\frac{3 \sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{2}, -\frac{1}{3}; \frac{2}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f(b \sin(e+fx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^2/(b*Sin[e + f*x])^(5/3),x]

[Out] (-3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/2, -1/3, 2/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(5/3))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(fx + e)}{(b \sin(fx + e))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^2/(b*sin(f*x+e))^(5/3),x)

[Out] int(cos(f*x+e)^2/(b*sin(f*x+e))^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2/(b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] integrate(cos(f*x + e)^2/(b*sin(f*x + e))^(5/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2/(b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral(-(b*sin(f*x + e))^(1/3)*cos(f*x + e)^2/(b^2*cos(f*x + e)^2 - b^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^2(e + fx)}{(b \sin(e + fx))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**2/(b*sin(f*x+e))**(5/3),x)

[Out] Integral(cos(e + f*x)**2/(b*sin(e + f*x))**(5/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^2/(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate(cos(f*x + e)^2/(b*sin(f*x + e))^(5/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\cos(e + f x)^2}{(b \sin(e + f x))^{5/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^2/(b*sin(e + f*x))^(5/3),x)

[Out] int(cos(e + f*x)^2/(b*sin(e + f*x))^(5/3), x)

$$3.321 \quad \int \frac{1}{(b \sin(e+fx))^{5/3}} dx$$

Optimal. Leaf size=58

$$-\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

[Out] $-3/2*\cos(f*x+e)*\text{hypergeom}([-1/3, 1/2], [2/3], \sin(f*x+e)^2)/b/f/(b*\sin(f*x+e))^{(2/3)}/(\cos(f*x+e)^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {2722}

$$-\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sin[e + f*x])^(-5/3),x]

[Out] $(-3*\text{Cos}[e + f*x]*\text{Hypergeometric2F1}[-1/3, 1/2, 2/3, \text{Sin}[e + f*x]^2])/(2*b*f*\text{Sqrt}[\text{Cos}[e + f*x]^2]*(b*\text{Sin}[e + f*x])^{(2/3)})$

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int \frac{1}{(b \sin(e+fx))^{5/3}} dx = -\frac{3 \cos(e+fx) {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf \sqrt{\cos^2(e+fx)} (b \sin(e+fx))^{2/3}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$-\frac{3 \sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{1}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f(b \sin(e+fx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sin[e + f*x])^(-5/3),x]

[Out] (-3*sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/3, 1/2, 2/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(5/3))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sin(fx + e))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sin(f*x+e))^(5/3),x)

[Out] int(1/(b*sin(f*x+e))^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^(5/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral(-(b*sin(f*x + e))^(1/3)/(b^2*cos(f*x + e)^2 - b^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sin(e + fx))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))**(5/3),x)

[Out] Integral((b*sin(e + f*x))**(-5/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^(-5/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{(b \sin(e + f x))^{5/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sin(e + f*x))^(5/3),x)

[Out] int(1/(b*sin(e + f*x))^(5/3), x)

$$3.322 \quad \int \frac{\sec^2(e+fx)}{(b \sin(e+fx))^{5/3}} dx$$

Optimal. Leaf size=58

$$-\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{3}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \sec(e+fx)}{2bf(b \sin(e+fx))^{2/3}}$$

[Out] $-3/2*\text{hypergeom}([-1/3, 3/2], [2/3], \sin(f*x+e)^2)*\sec(f*x+e)*(\cos(f*x+e)^2)^{(1/2)}/b/f/(b*\sin(f*x+e))^{(2/3)}$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$-\frac{3\sqrt{\cos^2(e+fx)} \sec(e+fx) {}_2F_1\left(-\frac{1}{3}, \frac{3}{2}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf(b \sin(e+fx))^{2/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[e + f*x]^2/(b*\text{Sin}[e + f*x])^{(5/3)}, x]$

[Out] $(-3*\text{Sqrt}[\text{Cos}[e + f*x]^2]*\text{Hypergeometric2F1}[-1/3, 3/2, 2/3, \text{Sin}[e + f*x]^2]*\text{Sec}[e + f*x])/(2*b*f*(b*\text{Sin}[e + f*x])^{(2/3)})$

Rule 2657

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(b_.))^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Cos}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}*((a*\text{Sin}[e + f*x])^{(m + 1)})/(a*f*(m + 1)*(\text{Cos}[e + f*x]^2)^{\text{FracPart}[(n - 1)/2]})*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Sin}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x]$

Rubi steps

$$\int \frac{\sec^2(e+fx)}{(b \sin(e+fx))^{5/3}} dx = -\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{3}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \sec(e+fx)}{2bf(b \sin(e+fx))^{2/3}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{3}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f(b \sin(e+fx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[e + f*x]^2/(b*Sin[e + f*x])^(5/3),x]

[Out] (-3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/3, 3/2, 2/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(5/3))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(fx + e)}{(b \sin(fx + e))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^2/(b*sin(f*x+e))^(5/3),x)

[Out] int(sec(f*x+e)^2/(b*sin(f*x+e))^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2/(b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] integrate(sec(f*x + e)^2/(b*sin(f*x + e))^(5/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2/(b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral(-(b*sin(f*x + e))^(1/3)*sec(f*x + e)^2/(b^2*cos(f*x + e)^2 - b^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^2(e + fx)}{(b \sin(e + fx))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)**2/(b*sin(f*x+e))**(5/3),x)

[Out] Integral(sec(e + f*x)**2/(b*sin(e + f*x))**(5/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^2/(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate(sec(f*x + e)^2/(b*sin(f*x + e))^(5/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(e + f x)^2 (b \sin(e + f x))^{5/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(e + f*x)^2*(b*sin(e + f*x))^(5/3)),x)

[Out] int(1/(cos(e + f*x)^2*(b*sin(e + f*x))^(5/3)), x)

$$3.323 \quad \int \frac{\sec^4(e+fx)}{(b \sin(e+fx))^{5/3}} dx$$

Optimal. Leaf size=58

$$-\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{5}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \sec(e+fx)}{2bf(b \sin(e+fx))^{2/3}}$$

[Out] $-3/2*\text{hypergeom}([-1/3, 5/2], [2/3], \sin(f*x+e)^2)*\sec(f*x+e)*(\cos(f*x+e)^2)^{(1/2)}/b/f/(b*\sin(f*x+e))^{(2/3)}$

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2657}

$$-\frac{3\sqrt{\cos^2(e+fx)} \sec(e+fx) {}_2F_1\left(-\frac{1}{3}, \frac{5}{2}; \frac{2}{3}; \sin^2(e+fx)\right)}{2bf(b \sin(e+fx))^{2/3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sec}[e + f*x]^4/(b*\text{Sin}[e + f*x])^{(5/3)}, x]$

[Out] $(-3*\text{Sqrt}[\text{Cos}[e + f*x]^2]*\text{Hypergeometric2F1}[-1/3, 5/2, 2/3, \text{Sin}[e + f*x]^2]*\text{Sec}[e + f*x])/(2*b*f*(b*\text{Sin}[e + f*x])^{(2/3)})$

Rule 2657

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] :> \text{Simp}[b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Cos}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}*((a*\text{Sin}[e + f*x])^{(m + 1)})/(a*f*(m + 1)*(\text{Cos}[e + f*x]^2)^{\text{FracPart}[(n - 1)/2]})*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Sin}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x]$

Rubi steps

$$\int \frac{\sec^4(e+fx)}{(b \sin(e+fx))^{5/3}} dx = -\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{5}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \sec(e+fx)}{2bf(b \sin(e+fx))^{2/3}}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.95

$$-\frac{3\sqrt{\cos^2(e+fx)} {}_2F_1\left(-\frac{1}{3}, \frac{5}{2}; \frac{2}{3}; \sin^2(e+fx)\right) \tan(e+fx)}{2f(b \sin(e+fx))^{5/3}}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[e + f*x]^4/(b*Sin[e + f*x])^(5/3),x]

[Out] (-3*Sqrt[Cos[e + f*x]^2]*Hypergeometric2F1[-1/3, 5/2, 2/3, Sin[e + f*x]^2]*Tan[e + f*x])/(2*f*(b*Sin[e + f*x])^(5/3))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(fx + e)}{(b \sin(fx + e))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^4/(b*sin(f*x+e))^(5/3),x)

[Out] int(sec(f*x+e)^4/(b*sin(f*x+e))^(5/3),x)

Maxima [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4/(b*sin(f*x+e))^(5/3),x, algorithm="maxima")

[Out] Timed out

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4/(b*sin(f*x+e))^(5/3),x, algorithm="fricas")

[Out] integral(-(b*sin(f*x + e))^(1/3)*sec(f*x + e)^4/(b^2*cos(f*x + e)^2 - b^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sec^4(e + fx)}{(b \sin(e + fx))^{\frac{5}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)**4/(b*sin(f*x+e))**(5/3),x)

[Out] Integral(sec(e + f*x)**4/(b*sin(e + f*x))**(5/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^4/(b*sin(f*x+e))^(5/3),x, algorithm="giac")

[Out] integrate(sec(f*x + e)^4/(b*sin(f*x + e))^(5/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\cos(e + f x)^4 (b \sin(e + f x))^{5/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(cos(e + f*x)^4*(b*sin(e + f*x))^(5/3)),x)

[Out] int(1/(cos(e + f*x)^4*(b*sin(e + f*x))^(5/3)), x)

$$3.324 \quad \int \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} dx$$

Optimal. Leaf size=128

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - 2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b}$$

[Out] $-1/2*\ln(1+\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)})/b+1/4*\ln(1-\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)}+\sin(b*x+a)^{(4/3)}/\cos(b*x+a)^{(4/3)})/b-1/2*\arctan(1/3*(1-2*\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)})*3^{(1/2)})*3^{(1/2)}/b$

Rubi [A]

time = 0.10, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2654, 281, 298, 31, 648, 632, 210, 642}

$$\frac{\sqrt{3} \text{ArcTan}\left(\frac{1 - 2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(\frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + 1\right)}{2b} + \frac{\log\left(\frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)} - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(1/3)/Cos[a + b*x]^(1/3),x]

[Out] $-1/2*(\text{Sqrt}[3]*\text{ArcTan}[(1 - (2*\text{Sin}[a + b*x]^{(2/3)})/\text{Cos}[a + b*x]^{(2/3)})/\text{Sqrt}[3]])/b - \text{Log}[1 + \text{Sin}[a + b*x]^{(2/3)}/\text{Cos}[a + b*x]^{(2/3)}]/(2*b) + \text{Log}[1 - \text{Sin}[a + b*x]^{(2/3)}/\text{Cos}[a + b*x]^{(2/3)} + \text{Sin}[a + b*x]^{(4/3)}/\text{Cos}[a + b*x]^{(4/3)}]/(4*b)$

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n+1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

Rule 281

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m
+ 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x
^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 298

```
Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-
1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), I
nt[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x
^2), x], x] /; FreeQ[{a, b}, x]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} dx &= \frac{3 \text{Subst}\left(\int \frac{x^3}{1+x^6} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
&= \frac{3 \text{Subst}\left(\int \frac{x}{1+x^3} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\text{Subst}\left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} + \frac{3 \text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} \\
&= -\frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b} - \frac{3 \text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - \frac{2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} - \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 57, normalized size = 0.45

$$\frac{3 \cos^2(a+bx)^{2/3} {}_2F_1\left(\frac{2}{3}, \frac{2}{3}; \frac{5}{3}; \sin^2(a+bx)\right) \sin^{4/3}(a+bx)}{4b \cos^{4/3}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^(1/3)/Cos[a + b*x]^(1/3), x]

[Out] (3*(Cos[a + b*x]^2)^(2/3)*Hypergeometric2F1[2/3, 2/3, 5/3, Sin[a + b*x]^2]*Sin[a + b*x]^(4/3))/(4*b*Cos[a + b*x]^(4/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\sin^{1/3}(bx+a)}{\cos(bx+a)^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3),x)`

[Out] `int(sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3),x, algorithm="maxima")`

[Out] `integrate(sin(b*x + a)^(1/3)/cos(b*x + a)^(1/3), x)`

Fricas [A]

time = 0.42, size = 144, normalized size = 1.12

$$\frac{2\sqrt{3} \arctan\left(-\frac{\sqrt{3} \cos(bx+a) - 2\sqrt{3} \cos(bx+a)^{\frac{1}{3}} \sin(bx+a)^{\frac{2}{3}}}{3 \cos(bx+a)}\right) - 2 \log\left(\frac{\cos(bx+a)^{\frac{1}{3}} \sin(bx+a)^{\frac{2}{3}} + \cos(bx+a)}{\cos(bx+a)}\right) + \log\left(\frac{\cos(bx+a)^2 - \cos(bx+a)^{\frac{4}{3}} \sin(bx+a)^{\frac{2}{3}} + \cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{4}{3}}}{\cos(bx+a)^2}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3),x, algorithm="fricas")`

[Out] `1/4*(2*sqrt(3)*arctan(-1/3*(sqrt(3)*cos(b*x + a) - 2*sqrt(3)*cos(b*x + a)^(1/3)*sin(b*x + a)^(2/3))/cos(b*x + a)) - 2*log((cos(b*x + a)^(1/3)*sin(b*x + a)^(2/3) + cos(b*x + a))/cos(b*x + a)) + log((cos(b*x + a)^2 - cos(b*x + a)^(4/3)*sin(b*x + a)^(2/3) + cos(b*x + a)^(2/3)*sin(b*x + a)^(4/3))/cos(b*x + a)^2))/b`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(1/3)/cos(b*x+a)**(1/3),x)`

[Out] `Integral(sin(a + b*x)**(1/3)/cos(a + b*x)**(1/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3),x, algorithm="giac")`

[Out] integrate(sin(b*x + a)^(1/3)/cos(b*x + a)^(1/3), x)

Mupad [B]

time = 1.24, size = 44, normalized size = 0.34

$$-\frac{3 \cos(a + bx)^{2/3} \sin(a + bx)^{4/3} {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; \cos(a + bx)^2\right)}{2b (\sin(a + bx)^2)^{2/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^(1/3)/cos(a + b*x)^(1/3),x)

[Out] -(3*cos(a + b*x)^(2/3)*sin(a + b*x)^(4/3)*hypergeom([1/3, 1/3], 4/3, cos(a + b*x)^2))/(2*b*(sin(a + b*x)^2)^(2/3))

$$3.325 \quad \int \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} dx$$

Optimal. Leaf size=224

$$\frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} + \frac{\sqrt{3} \log}{b}$$

[Out] arctan(sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3))/b+1/2*arctan(2*sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3)-3^(1/2))/b+1/2*arctan(2*sin(b*x+a)^(1/3)/cos(b*x+a)^(1/3)+3^(1/2))/b+1/4*ln(1+sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3)-sin(b*x+a)^(1/3)*3^(1/2)/cos(b*x+a)^(1/3))*3^(1/2)/b-1/4*ln(1+sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3)+sin(b*x+a)^(1/3)*3^(1/2)/cos(b*x+a)^(1/3))*3^(1/2)/b

Rubi [A]

time = 0.22, antiderivative size = 224, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2654, 301, 648, 632, 210, 642, 209}

$$\frac{\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} + \frac{\text{ArcTan}\left(\frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + \sqrt{3}\right)}{2b} + \frac{\text{ArcTan}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} + \frac{\sqrt{3} \log\left(\frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} - \frac{\sqrt{3}\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + 1\right)}{4b} - \frac{\sqrt{3} \log\left(\frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sqrt{3}\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(2/3)/Cos[a + b*x]^(2/3), x]

[Out] -1/2*ArcTan[Sqrt[3] - (2*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3)]/b + ArcTan[Sqrt[3] + (2*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3)]/(2*b) + ArcTan[Sin[a + b*x]^(1/3)/Cos[a + b*x]^(1/3)]/b + (Sqrt[3]*Log[1 - (Sqrt[3]*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3) + Sin[a + b*x]^(2/3)/Cos[a + b*x]^(2/3)])/(4*b) - (Sqrt[3]*Log[1 + (Sqrt[3]*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3) + Sin[a + b*x]^(2/3)/Cos[a + b*x]^(2/3)])/(4*b)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 301

```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator
[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k
- 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k -
1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k
- 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]
; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m)*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^
(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x]] /; FreeQ[{a, b}, x]
&& IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*SIN[e + f*x])^(1/k)/(b*COS[e +
f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} dx &= \frac{3 \text{Subst}\left(\int \frac{x^4}{1+x^6} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
&= \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} + \frac{\text{Subst}\left(\int \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}x}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
&= \frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} + \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{4b} + \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{4b} \\
&= \frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} + \frac{\sqrt{3} \log\left(1 - \frac{\sqrt{3} \sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\sqrt{3} \log\left(1 - \frac{\sqrt{3} \sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} \\
&= -\frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 57, normalized size = 0.25

$$\frac{3 \cos^2(a+bx)^{5/6} {}_2F_1\left(\frac{5}{6}, \frac{5}{6}; \frac{11}{6}; \sin^2(a+bx)\right) \sin^{5/3}(a+bx)}{5b \cos^{5/3}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^(2/3)/Cos[a + b*x]^(2/3), x]

[Out] (3*(Cos[a + b*x]^2)^(5/6)*Hypergeometric2F1[5/6, 5/6, 11/6, Sin[a + b*x]^2]*Sin[a + b*x]^(5/3))/(5*b*Cos[a + b*x]^(5/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{2}{3}}(bx+a)}{\cos(bx+a)^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3), x)

[Out] int(sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(sin(b*x + a)^(2/3)/cos(b*x + a)^(2/3), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{2}{3}}(a + bx)}{\cos^{\frac{2}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)**(2/3)/cos(b*x+a)**(2/3),x)

[Out] Integral(sin(a + b*x)**(2/3)/cos(a + b*x)**(2/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^(2/3)/cos(b*x+a)^(2/3),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^(2/3)/cos(b*x + a)^(2/3), x)

Mupad [B]

time = 1.05, size = 44, normalized size = 0.20

$$\frac{3 \cos(a + bx)^{1/3} \sin(a + bx)^{5/3} {}_2F_1\left(\frac{1}{6}, \frac{1}{6}; \frac{7}{6}; \cos(a + bx)^2\right)}{b (\sin(a + bx)^2)^{5/6}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^(2/3)/cos(a + b*x)^(2/3),x)

[Out] -(3*cos(a + b*x)^(1/3)*sin(a + b*x)^(5/3)*hypergeom([1/6, 1/6], 7/6, cos(a + b*x)^2))/(b*(sin(a + b*x)^2)^(5/6))

$$3.326 \quad \int \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)} dx$$

Optimal. Leaf size=249

$$\frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} + \sqrt{3} \log$$

[Out] arctan(cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3))/b+1/2*arctan(2*cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3)-3^(1/2))/b+1/2*arctan(2*cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3)+3^(1/2))/b+3*sin(b*x+a)^(1/3)/b/cos(b*x+a)^(1/3)+1/4*ln(1+cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3)-cos(b*x+a)^(1/3)*3^(1/2)/sin(b*x+a)^(1/3))*3^(1/2)/b-1/4*ln(1+cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3)+cos(b*x+a)^(1/3)*3^(1/2)/sin(b*x+a)^(1/3))*3^(1/2)/b

Rubi [A]

time = 0.24, antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2646, 2655, 301, 648, 632, 210, 642, 209}

$$\frac{\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} + \frac{\text{ArcTan}\left(\frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} + \sqrt{3}\right)}{2b} + \frac{\text{ArcTan}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} + \frac{3\sqrt[3]{\sin(a+bx)}}{b\sqrt[3]{\cos(a+bx)}} + \frac{\sqrt{3} \log\left(\frac{\cos^{\frac{2}{3}}(a+bx) - \sqrt{3}\sqrt[3]{\cos(a+bx)}}{\sin^{\frac{2}{3}}(a+bx)} + 1\right)}{4b} - \frac{\sqrt{3} \log\left(\frac{\cos^{\frac{2}{3}}(a+bx) + \sqrt{3}\sqrt[3]{\cos(a+bx)}}{\sin^{\frac{2}{3}}(a+bx)} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(4/3)/Cos[a + b*x]^(4/3), x]

[Out] -1/2*ArcTan[Sqrt[3] - (2*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)]/b + ArcTan[Sqrt[3] + (2*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)]/(2*b) + ArcTan[Cos[a + b*x]^(1/3)/Sin[a + b*x]^(1/3)]/b + (Sqrt[3]*Log[1 + Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3) - (Sqrt[3]*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)])/(4*b) - (Sqrt[3]*Log[1 + Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3) + (Sqrt[3]*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)])/(4*b) + (3*Sin[a + b*x]^(1/3))/(b*Cos[a + b*x]^(1/3))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2]))^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 301

```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator
[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k
- 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k -
1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k
- 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]
; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^
(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x] /; FreeQ[{a, b}, x]
&& IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 2646

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*SIN[e + f*x])^(m - 1)*((b*COS[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*SIN[e + f*x])^(m - 2)*(b*COS[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])
```

Rule 2655

```
Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*COS[e + f*x])^(1/k)/(b*SIN[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)} dx &= \frac{3\sqrt[3]{\sin(a+bx)}}{b\sqrt[3]{\cos(a+bx)}} - \int \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} dx \\
 &= \frac{3\sqrt[3]{\sin(a+bx)}}{b\sqrt[3]{\cos(a+bx)}} + \frac{3\text{Subst}\left(\int \frac{x^4}{1+x^6} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
 &= \frac{3\sqrt[3]{\sin(a+bx)}}{b\sqrt[3]{\cos(a+bx)}} + \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} + \frac{\text{Subst}\left(\int \frac{-\frac{1}{2} + \frac{\sqrt{3}x}{2}}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
 &= \frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} + \frac{3\sqrt[3]{\sin(a+bx)}}{b\sqrt[3]{\cos(a+bx)}} + \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} \\
 &= \frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} + \frac{\sqrt{3} \log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} - \frac{\sqrt{3} \sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} - \frac{\sqrt{3} \log\left(1 - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + \frac{\sqrt{3} \sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} \\
 &= -\frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} + \frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 57, normalized size = 0.23

$$\frac{3\sqrt[6]{\cos^2(a+bx)} {}_2F_1\left(\frac{7}{6}, \frac{7}{6}, \frac{13}{6}; \sin^2(a+bx)\right) \sin^{\frac{7}{3}}(a+bx)}{7b\sqrt[3]{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^(4/3)/Cos[a + b*x]^(4/3), x]

[Out] (3*(Cos[a + b*x]^2)^(1/6)*Hypergeometric2F1[7/6, 7/6, 13/6, Sin[a + b*x]^2]*Sin[a + b*x]^(7/3))/(7*b*Cos[a + b*x]^(1/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{4}{3}}(bx+a)}{\cos(bx+a)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^(4/3)/cos(b*x+a)^(4/3),x)`

[Out] `int(sin(b*x+a)^(4/3)/cos(b*x+a)^(4/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(4/3)/cos(b*x+a)^(4/3),x, algorithm="maxima")`

[Out] `integrate(sin(b*x + a)^(4/3)/cos(b*x + a)^(4/3), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(4/3)/cos(b*x+a)^(4/3),x, algorithm="fricas")`

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{4}{3}}(a + bx)}{\cos^{\frac{4}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(4/3)/cos(b*x+a)**(4/3),x)`

[Out] `Integral(sin(a + b*x)**(4/3)/cos(a + b*x)**(4/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(4/3)/cos(b*x+a)^(4/3),x, algorithm="giac")`

[Out] `integrate(sin(b*x + a)^(4/3)/cos(b*x + a)^(4/3), x)`

Mupad [B]

time = 1.64, size = 44, normalized size = 0.18

$$\frac{3 \sin(a + bx)^{7/3} {}_2F_1\left(-\frac{1}{6}, -\frac{1}{6}; \frac{5}{6}; \cos(a + bx)^2\right)}{b \cos(a + bx)^{1/3} (\sin(a + bx)^2)^{7/6}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^(4/3)/cos(a + b*x)^(4/3), x)``[Out] (3*sin(a + b*x)^(7/3)*hypergeom([-1/6, -1/6], 5/6, cos(a + b*x)^2))/(b*cos(a + b*x)^(1/3)*(sin(a + b*x)^2)^(7/6))`

$$3.327 \quad \int \frac{\sin^{\frac{5}{3}}(a+bx)}{\cos^{\frac{5}{3}}(a+bx)} dx$$

Optimal. Leaf size=155

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - 2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)}$$

[Out] 1/4*ln(1+cos(b*x+a)^(4/3)/sin(b*x+a)^(4/3)-cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3))/b-1/2*ln(1+cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3))/b+3/2*sin(b*x+a)^(2/3)/b/cos(b*x+a)^(2/3)-1/2*arctan(1/3*(1-2*cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3))*3^(1/2))*3^(1/2)/b

Rubi [A]

time = 0.12, antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2646, 2655, 281, 298, 31, 648, 632, 210, 642}

$$\frac{\sqrt{3} \text{ArcTan}\left(\frac{1 - 2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} + \frac{\log\left(\frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + 1\right)}{4b} - \frac{\log\left(\frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + 1\right)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(5/3)/Cos[a + b*x]^(5/3),x]

[Out] -1/2*(Sqrt[3]*ArcTan[(1 - (2*Cos[a + b*x]^(2/3))/Sin[a + b*x]^(2/3))/Sqrt[3]])/b + Log[1 + Cos[a + b*x]^(4/3)/Sin[a + b*x]^(4/3) - Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3)]/(4*b) - Log[1 + Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3)]/(2*b) + (3*Sin[a + b*x]^(2/3))/(2*b*Cos[a + b*x]^(2/3))

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n_+1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 298

Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2646

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*SIN[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*SIN[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2655

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*SIN[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^{\frac{5}{3}}(a+bx)}{\cos^{\frac{5}{3}}(a+bx)} dx &= \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} - \int \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} dx \\
&= \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} + \frac{3 \text{Subst}\left(\int \frac{x^3}{1+x^6} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
&= \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} + \frac{3 \text{Subst}\left(\int \frac{x}{1+x^3} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} + \frac{3 \text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{3 \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)} - \frac{3 \text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - \frac{2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} + \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 57, normalized size = 0.37

$$\frac{3 \sqrt[3]{\cos^2(a+bx)} {}_2F_1\left(\frac{4}{3}, \frac{4}{3}; \frac{7}{3}; \sin^2(a+bx)\right) \sin^{\frac{8}{3}}(a+bx)}{8b \cos^{\frac{2}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^(5/3)/Cos[a + b*x]^(5/3), x]

[Out] (3*(Cos[a + b*x]^2)^(1/3)*Hypergeometric2F1[4/3, 4/3, 7/3, Sin[a + b*x]^2]*Sin[a + b*x]^(8/3))/(8*b*Cos[a + b*x]^(2/3))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{5}{3}}(bx+a)}{\cos^{\frac{5}{3}}(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^(5/3)/cos(b*x+a)^(5/3),x)`

[Out] `int(sin(b*x+a)^(5/3)/cos(b*x+a)^(5/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(5/3)/cos(b*x+a)^(5/3),x, algorithm="maxima")`

[Out] `integrate(sin(b*x + a)^(5/3)/cos(b*x + a)^(5/3), x)`

Fricas [A]

time = 0.53, size = 197, normalized size = 1.27

$$\frac{2\sqrt{3}\arctan\left(\frac{2\sqrt{3}\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{1}{3}}-\sqrt{3}\sin(bx+a)}{3\sin(bx+a)}\right)\cos(bx+a)+\cos(bx+a)\log\left(\frac{4(\cos(bx+a)^2-\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{2}{3}}+\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{2}{3}}-1)}{\cos(bx+a)^2-1}\right)-2\cos(bx+a)\log\left(\frac{-2(\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{1}{3}}+\sin(bx+a))}{\sin(bx+a)}\right)+6\cos(bx+a)^{\frac{1}{3}}\sin(bx+a)^{\frac{2}{3}}}{4b\cos(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(5/3)/cos(b*x+a)^(5/3),x, algorithm="fricas")`

[Out] `1/4*(2*sqrt(3)*arctan(1/3*(2*sqrt(3)*cos(b*x + a)^(2/3)*sin(b*x + a)^(1/3) - sqrt(3)*sin(b*x + a))/sin(b*x + a))*cos(b*x + a) + cos(b*x + a)*log(4*(cos(b*x + a)^2 - cos(b*x + a)^(4/3)*sin(b*x + a)^(2/3) + cos(b*x + a)^(2/3)*sin(b*x + a)^(4/3) - 1)/(cos(b*x + a)^2 - 1)) - 2*cos(b*x + a)*log(-2*(cos(b*x + a)^(2/3)*sin(b*x + a)^(1/3) + sin(b*x + a))/sin(b*x + a)) + 6*cos(b*x + a)^(1/3)*sin(b*x + a)^(2/3)/(b*cos(b*x + a))`

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(5/3)/cos(b*x+a)**(5/3),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(b*x+a)^(5/3)/cos(b*x+a)^(5/3),x, algorithm="giac")

[Out] integrate(sin(b*x + a)^(5/3)/cos(b*x + a)^(5/3), x)

Mupad [B]

time = 1.14, size = 44, normalized size = 0.28

$$\frac{3 \sin(a + bx)^{8/3} {}_2F_1\left(-\frac{1}{3}, -\frac{1}{3}; \frac{2}{3}; \cos(a + bx)^2\right)}{2b \cos(a + bx)^{2/3} (\sin(a + bx)^2)^{4/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^(5/3)/cos(a + b*x)^(5/3),x)

[Out] (3*sin(a + b*x)^(8/3)*hypergeom([-1/3, -1/3], 2/3, cos(a + b*x)^2))/(2*b*cos(a + b*x)^(2/3)*(sin(a + b*x)^2)^(4/3))

$$3.328 \quad \int \frac{\sin^{\frac{7}{3}}(a+bx)}{\cos^{\frac{7}{3}}(a+bx)} dx$$

Optimal. Leaf size=155

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - \frac{2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} + \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b} + \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)}$$

[Out] $\frac{1}{2} \ln(1 + \sin(b*x+a)^{(2/3)} / \cos(b*x+a)^{(2/3)}) / b - \frac{1}{4} \ln(1 - \sin(b*x+a)^{(2/3)} / \cos(b*x+a)^{(2/3)} + \sin(b*x+a)^{(4/3)} / \cos(b*x+a)^{(4/3)}) / b + \frac{3}{4} \sin(b*x+a)^{(4/3)} / b \cos(b*x+a)^{(4/3)} + \frac{1}{2} \arctan(1/3 * (1 - 2 * \sin(b*x+a)^{(2/3)} / \cos(b*x+a)^{(2/3)}) * 3^{(1/2)}) * 3^{(1/2)} / b$

Rubi [A]

time = 0.08, antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2646, 2654, 281, 298, 31, 648, 632, 210, 642}

$$\frac{\sqrt{3} \text{ArcTan}\left(\frac{1 - \frac{2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} + \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} + \frac{\log\left(\frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + 1\right)}{2b} - \frac{\log\left(\frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)} - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Sin[a + b*x]^(7/3)/Cos[a + b*x]^(7/3), x]

[Out] $(\text{Sqrt}[3] * \text{ArcTan}[(1 - (2 * \text{Sin}[a + b*x]^{(2/3)}) / \text{Cos}[a + b*x]^{(2/3)}) / \text{Sqrt}[3]]) / (2 * b) + \text{Log}[1 + \text{Sin}[a + b*x]^{(2/3)} / \text{Cos}[a + b*x]^{(2/3)}] / (2 * b) - \text{Log}[1 - \text{Sin}[a + b*x]^{(2/3)} / \text{Cos}[a + b*x]^{(2/3)} + \text{Sin}[a + b*x]^{(4/3)} / \text{Cos}[a + b*x]^{(4/3)}] / (4 * b) + (3 * \text{Sin}[a + b*x]^{(4/3)}) / (4 * b * \text{Cos}[a + b*x]^{(4/3)})$

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n_ - 1) * ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 298

Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2646

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Simp[(-a)*(a*Sine[e + f*x])^(m - 1)*((b*Cos[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Sine[e + f*x])^(m - 2)*(b*Cos[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegerQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sine[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^{\frac{7}{3}}(a+bx)}{\cos^{\frac{7}{3}}(a+bx)} dx &= \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} - \int \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} dx \\
&= \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} - \frac{3 \text{Subst}\left(\int \frac{x^3}{1+x^6} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
&= \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} - \frac{3 \text{Subst}\left(\int \frac{x}{1+x^3} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{3 \text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} \\
&= \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b} + \frac{3 \sin^{\frac{4}{3}}(a+bx)}{4b \cos^{\frac{4}{3}}(a+bx)} + \frac{3 \text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} \\
&= \frac{\sqrt{3} \tan^{-1}\left(\frac{1 - \frac{2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} + \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 57, normalized size = 0.37

$$\frac{3 \cos^2(a+bx)^{2/3} {}_2F_1\left(\frac{5}{3}, \frac{5}{3}; \frac{8}{3}; \sin^2(a+bx)\right) \sin^{\frac{10}{3}}(a+bx)}{10b \cos^{\frac{4}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[a + b*x]^(7/3)/Cos[a + b*x]^(7/3), x]

[Out] (3*(Cos[a + b*x]^2)^(2/3)*Hypergeometric2F1[5/3, 5/3, 8/3, Sin[a + b*x]^2]*Sin[a + b*x]^(10/3))/(10*b*Cos[a + b*x]^(4/3))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{7}{3}}(bx+a)}{\cos^{\frac{7}{3}}(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(b*x+a)^(7/3)/cos(b*x+a)^(7/3),x)`

[Out] `int(sin(b*x+a)^(7/3)/cos(b*x+a)^(7/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(7/3)/cos(b*x+a)^(7/3),x, algorithm="maxima")`

[Out] `integrate(sin(b*x + a)^(7/3)/cos(b*x + a)^(7/3), x)`

Fricas [A]

time = 0.45, size = 195, normalized size = 1.26

$$\frac{2\sqrt{3}\arctan\left(-\frac{\sqrt{3}\cos(bx+a)-2\sqrt{3}\cos(bx+a)\frac{1}{3}\sin(bx+a)^{\frac{2}{3}}}{3\cos(bx+a)}\right)\cos(bx+a)^2-2\cos(bx+a)^2\log\left(\frac{\cos(bx+a)\frac{1}{3}\sin(bx+a)^{\frac{2}{3}}+\cos(bx+a)}{\cos(bx+a)}\right)+\cos(bx+a)^2\log\left(\frac{\cos(bx+a)^2-\cos(bx+a)\frac{1}{3}\sin(bx+a)^{\frac{2}{3}}+\cos(bx+a)\frac{1}{3}\sin(bx+a)^{\frac{2}{3}}}{\cos(bx+a)^2}\right)-3\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{2}{3}}}{4b\cos(bx+a)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)^(7/3)/cos(b*x+a)^(7/3),x, algorithm="fricas")`

[Out] `-1/4*(2*sqrt(3)*arctan(-1/3*(sqrt(3)*cos(b*x + a) - 2*sqrt(3)*cos(b*x + a)^(1/3)*sin(b*x + a)^(2/3))/cos(b*x + a))*cos(b*x + a)^2 - 2*cos(b*x + a)^2*log((cos(b*x + a)^(1/3)*sin(b*x + a)^(2/3) + cos(b*x + a))/cos(b*x + a)) + cos(b*x + a)^2*log((cos(b*x + a)^2 - cos(b*x + a)^(4/3)*sin(b*x + a)^(2/3) + cos(b*x + a)^(2/3)*sin(b*x + a)^(4/3))/cos(b*x + a)^2) - 3*cos(b*x + a)^(2/3)*sin(b*x + a)^(4/3)/(b*cos(b*x + a)^2)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(b*x+a)**(7/3)/cos(b*x+a)**(7/3),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(b*x+a)^(7/3)/cos(b*x+a)^(7/3),x, algorithm="giac")``[Out] integrate(sin(b*x + a)^(7/3)/cos(b*x + a)^(7/3), x)`**Mupad [B]**

time = 1.64, size = 44, normalized size = 0.28

$$\frac{3 \sin(a + bx)^{10/3} {}_2F_1\left(-\frac{2}{3}, -\frac{2}{3}; \frac{1}{3}; \cos(a + bx)^2\right)}{4 b \cos(a + bx)^{4/3} (\sin(a + bx)^2)^{5/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(a + b*x)^(7/3)/cos(a + b*x)^(7/3),x)``[Out] (3*sin(a + b*x)^(10/3)*hypergeom([-2/3, -2/3], 1/3, cos(a + b*x)^2))/(4*b*cos(a + b*x)^(4/3)*(sin(a + b*x)^2)^(5/3))`

$$3.329 \quad \int \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} dx$$

Optimal. Leaf size=128

$$\frac{\sqrt{3} \tan^{-1} \left(\frac{1 - \frac{2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}}{\sqrt{3}} \right)}{2b} - \frac{\log \left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} \right)}{4b} + \frac{\log \left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} \right)}{2b}$$

[Out] $-1/4*\ln(1+\cos(b*x+a)^{(4/3)}/\sin(b*x+a)^{(4/3)}-\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)})/b+1/2*\ln(1+\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)})/b+1/2*\arctan(1/3*(1-2*\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3}))*3^{(1/2)})*3^{(1/2)}/b$

Rubi [A]

time = 0.05, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2655, 281, 298, 31, 648, 632, 210, 642}

$$\frac{\sqrt{3} \text{ArcTan} \left(\frac{1 - \frac{2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}}{\sqrt{3}} \right)}{2b} - \frac{\log \left(\frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + 1 \right)}{4b} + \frac{\log \left(\frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + 1 \right)}{2b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(1/3)/Sin[a + b*x]^(1/3), x]

[Out] $(\text{Sqrt}[3]*\text{ArcTan}[(1 - (2*\text{Cos}[a + b*x]^{(2/3)})/\text{Sin}[a + b*x]^{(2/3)})/\text{Sqrt}[3]])/(2*b) - \text{Log}[1 + \text{Cos}[a + b*x]^{(4/3)}/\text{Sin}[a + b*x]^{(4/3)} - \text{Cos}[a + b*x]^{(2/3)}/\text{Sin}[a + b*x]^{(2/3)}]/(4*b) + \text{Log}[1 + \text{Cos}[a + b*x]^{(2/3)}/\text{Sin}[a + b*x]^{(2/3)}]/(2*b)$

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n_+1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] & & (LtQ[a, 0] || LtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 298

Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2655

Int[(cos[(e_) + (f_)*(x_)])*(a_)^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sin[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} dx &= -\frac{3\text{Subst}\left(\int \frac{x^3}{1+x^6} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
&= -\frac{3\text{Subst}\left(\int \frac{x}{1+x^3} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\text{Subst}\left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{3\text{Subst}\left(\int \frac{1}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} \\
&= -\frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} + \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{3\text{Subst}\left(\int \frac{1}{-3-x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\sqrt{3} \tan^{-1}\left(\frac{1 - 2\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} + \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 57, normalized size = 0.45

$$\frac{3\sqrt[3]{\cos^2(a+bx)} {}_2F_1\left(\frac{1}{3}, \frac{1}{3}; \frac{4}{3}; \sin^2(a+bx)\right) \sin^{\frac{2}{3}}(a+bx)}{2b \cos^{\frac{2}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(1/3)/Sin[a + b*x]^(1/3), x]

[Out] (3*(Cos[a + b*x]^2)^(1/3)*Hypergeometric2F1[1/3, 1/3, 4/3, Sin[a + b*x]^2]*Sin[a + b*x]^(2/3))/(2*b*Cos[a + b*x]^(2/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{1}{3}}(bx+a)}{\sin(bx+a)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3),x)`

[Out] `int(cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3),x, algorithm="maxima")`

[Out] `integrate(cos(b*x + a)^(1/3)/sin(b*x + a)^(1/3), x)`

Fricas [A]

time = 0.53, size = 152, normalized size = 1.19

$$\frac{2\sqrt{3} \arctan\left(\frac{2\sqrt{3} \cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{1}{3}} - \sqrt{3} \sin(bx+a)}{3 \sin(bx+a)}\right) + \log\left(\frac{4(\cos(bx+a)^2 - \cos(bx+a)^{\frac{4}{3}} \sin(bx+a)^{\frac{2}{3}} + \cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{4}{3}} - 1)}{\cos(bx+a)^2 - 1}\right) - 2 \log\left(\frac{2(\cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{1}{3}} + \sin(bx+a))}{\sin(bx+a)}\right)}{4b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3),x, algorithm="fricas")`

[Out] `-1/4*(2*sqrt(3)*arctan(1/3*(2*sqrt(3)*cos(b*x + a)^(2/3)*sin(b*x + a)^(1/3) - sqrt(3)*sin(b*x + a))/sin(b*x + a)) + log(4*(cos(b*x + a)^2 - cos(b*x + a)^(4/3)*sin(b*x + a)^(2/3) + cos(b*x + a)^(2/3)*sin(b*x + a)^(4/3) - 1)/(cos(b*x + a)^2 - 1)) - 2*log(-2*(cos(b*x + a)^(2/3)*sin(b*x + a)^(1/3) + sin(b*x + a))/sin(b*x + a))/b`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[3]{\cos(a + bx)}}{\sqrt[3]{\sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**(1/3)/sin(b*x+a)**(1/3),x)`

[Out] `Integral(cos(a + b*x)**(1/3)/sin(a + b*x)**(1/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(1/3)/sin(b*x+a)^(1/3),x, algorithm="giac")`

[Out] integrate(cos(b*x + a)^(1/3)/sin(b*x + a)^(1/3), x)

Mupad [B]

time = 1.48, size = 44, normalized size = 0.34

$$-\frac{3 \cos(a + bx)^{4/3} \sin(a + bx)^{2/3} {}_2F_1\left(\frac{2}{3}, \frac{2}{3}; \frac{5}{3}; \cos(a + bx)^2\right)}{4b (\sin(a + bx)^2)^{1/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(1/3)/sin(a + b*x)^(1/3),x)

[Out] -(3*cos(a + b*x)^(4/3)*sin(a + b*x)^(2/3)*hypergeom([2/3, 2/3], 5/3, cos(a + b*x)^2))/(4*b*(sin(a + b*x)^2)^(1/3))

$$3.330 \quad \int \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} dx$$

Optimal. Leaf size=225

$$\frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} - \sqrt{3} \log\left(\dots\right)$$

[Out] $-\arctan(\cos(b*x+a)^{(1/3)}/\sin(b*x+a)^{(1/3)})/b-1/2*\arctan(2*\cos(b*x+a)^{(1/3)}/\sin(b*x+a)^{(1/3)}-3^{(1/2)})/b-1/2*\arctan(2*\cos(b*x+a)^{(1/3)}/\sin(b*x+a)^{(1/3)}+3^{(1/2)})/b-1/4*\ln(1+\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)}-\cos(b*x+a)^{(1/3)}*3^{(1/2)}/\sin(b*x+a)^{(1/3)})*3^{(1/2)}/b+1/4*\ln(1+\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)}+\cos(b*x+a)^{(1/3)}*3^{(1/2)}/\sin(b*x+a)^{(1/3)})*3^{(1/2)}/b$

Rubi [A]

time = 0.20, antiderivative size = 225, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2655, 301, 648, 632, 210, 642, 209}

$$\frac{\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} - \frac{\text{ArcTan}\left(\frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} + \sqrt{3}\right)}{2b} - \frac{\text{ArcTan}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} - \frac{\sqrt{3} \log\left(\frac{\cos^3(a+bx)}{\sin^3(a+bx)} - \frac{\sqrt{3}\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} + 1\right)}{4b} + \frac{\sqrt{3} \log\left(\frac{\cos^3(a+bx)}{\sin^3(a+bx)} + \frac{\sqrt{3}\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3), x]

[Out] ArcTan[Sqrt[3] - (2*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)]/(2*b) - ArcTan[Sqrt[3] + (2*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)]/(2*b) - ArcTan[Cos[a + b*x]^(1/3)/Sin[a + b*x]^(1/3)]/b - (Sqrt[3]*Log[1 + Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3) - (Sqrt[3]*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)])/(4*b) + (Sqrt[3]*Log[1 + Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3) + (Sqrt[3]*Cos[a + b*x]^(1/3))/Sin[a + b*x]^(1/3)])/(4*b)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 301

```

Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator
[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k
- 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k -
1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k
- 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]
; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^
(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x] /; FreeQ[{a, b}, x]
&& IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]

```

Rule 632

```

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]

```

Rule 642

```

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]

```

Rule 648

```

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

```

Rule 2655

```

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n
_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^
(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sin[e
+ f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0
] && LtQ[m, 1]

```

Rubi steps

$$\begin{aligned}
\int \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} dx &= -\frac{3 \operatorname{Subst}\left(\int \frac{x^4}{1+x^6} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
&= -\frac{\operatorname{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} - \frac{\operatorname{Subst}\left(\int \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}x}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
&= -\frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} - \frac{\operatorname{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} - \frac{\operatorname{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} \\
&= -\frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} - \frac{\sqrt{3} \log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} - \frac{\sqrt{3} \sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} + \frac{\sqrt{3} \log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} - \frac{\sqrt{3} \sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{4b} \\
&= \frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 55, normalized size = 0.24

$$\frac{3 \sqrt[6]{\cos^2(a+bx)} {}_2F_1\left(\frac{1}{6}, \frac{1}{6}; \frac{7}{6}; \sin^2(a+bx)\right) \sqrt[3]{\sin(a+bx)}}{b \sqrt[3]{\cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(2/3)/Sin[a + b*x]^(2/3), x]

[Out] (3*(Cos[a + b*x]^2)^(1/6)*Hypergeometric2F1[1/6, 1/6, 7/6, Sin[a + b*x]^2]*Sin[a + b*x]^(1/3))/(b*Cos[a + b*x]^(1/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{2}{3}}(bx+a)}{\sin^{\frac{2}{3}}(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3), x)

[Out] int(cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(2/3)/sin(b*x + a)^(2/3), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{2}{3}}(a + bx)}{\sin^{\frac{2}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(2/3)/sin(b*x+a)**(2/3),x)

[Out] Integral(cos(a + b*x)**(2/3)/sin(a + b*x)**(2/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(2/3)/sin(b*x+a)^(2/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(2/3)/sin(b*x + a)^(2/3), x)

Mupad [B]

time = 1.12, size = 44, normalized size = 0.20

$$\frac{3 \cos(a + bx)^{5/3} \sin(a + bx)^{1/3} {}_2F_1\left(\frac{5}{6}, \frac{5}{6}; \frac{11}{6}; \cos(a + bx)^2\right)}{5 b (\sin(a + bx)^2)^{1/6}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(2/3)/sin(a + b*x)^(2/3),x)

[Out] -(3*cos(a + b*x)^(5/3)*sin(a + b*x)^(1/3)*hypergeom([5/6, 5/6], 11/6, cos(a + b*x)^2))/(5*b*(sin(a + b*x)^2)^(1/6))

$$3.331 \quad \int \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} dx$$

Optimal. Leaf size=250

$$\frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} - \sqrt{3} \log\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)$$

[Out] $-\arctan(\sin(b*x+a)^{(1/3)}/\cos(b*x+a)^{(1/3)})/b-1/2*\arctan(2*\sin(b*x+a)^{(1/3)}/\cos(b*x+a)^{(1/3)}+3^{(1/2)})/b-3*\cos(b*x+a)^{(1/3)}/b/\sin(b*x+a)^{(1/3)}-1/4*\ln(1+\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)}-\sin(b*x+a)^{(1/3)}*3^{(1/2)}/\cos(b*x+a)^{(1/3)})*3^{(1/2)}/b+1/4*\ln(1+\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)}+\sin(b*x+a)^{(1/3)}*3^{(1/2)}/\cos(b*x+a)^{(1/3)})*3^{(1/2)}/b$

Rubi [A]

time = 0.23, antiderivative size = 250, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.381$, Rules used = {2647, 2654, 301, 648, 632, 210, 642, 209}

$$\frac{\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} - \frac{\text{ArcTan}\left(\frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + \sqrt{3}\right)}{2b} - \frac{\text{ArcTan}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} - \frac{3\sqrt[3]{\cos(a+bx)}}{b\sqrt[3]{\sin(a+bx)}} - \frac{\sqrt{3} \log\left(\frac{\sin^2(a+bx)}{\cos^2(a+bx)} - \frac{\sqrt{3}\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + 1\right)}{4b} + \frac{\sqrt{3} \log\left(\frac{\sin^2(a+bx)}{\cos^2(a+bx)} + \frac{\sqrt{3}\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(4/3)/Sin[a + b*x]^(4/3), x]

[Out] ArcTan[Sqrt[3] - (2*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3)]/(2*b) - ArcTan[Sqrt[3] + (2*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3)]/(2*b) - ArcTan[Sin[a + b*x]^(1/3)/Cos[a + b*x]^(1/3)]/b - (Sqrt[3]*Log[1 - (Sqrt[3]*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3) + Sin[a + b*x]^(2/3)/Cos[a + b*x]^(2/3)])/(4*b) + (Sqrt[3]*Log[1 + (Sqrt[3]*Sin[a + b*x]^(1/3))/Cos[a + b*x]^(1/3) + Sin[a + b*x]^(2/3)/Cos[a + b*x]^(2/3)])/(4*b) - (3*Cos[a + b*x]^(1/3))/(b*Sin[a + b*x]^(1/3))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2]))^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 301

```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator
[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k
- 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k -
1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k
- 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]
; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^
(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x] /; FreeQ[{a, b}, x]
&& IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 2647

```
Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(a*Cos[e + f*x])^(m - 1)*((b*Sin[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Cos[e + f*x])^(m - 2)*(b*Sin[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rubi steps

$$\begin{aligned}
 \int \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} dx &= -\frac{3\sqrt[3]{\cos(a+bx)}}{b\sqrt[3]{\sin(a+bx)}} - \int \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} dx \\
 &= -\frac{3\sqrt[3]{\cos(a+bx)}}{b\sqrt[3]{\sin(a+bx)}} - \frac{3\text{Subst}\left(\int \frac{x^4}{1+x^6} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
 &= -\frac{3\sqrt[3]{\cos(a+bx)}}{b\sqrt[3]{\sin(a+bx)}} - \frac{\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} - \frac{\text{Subst}\left(\int \frac{-\frac{1}{2} + \frac{\sqrt{3}x}{2}}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
 &= -\frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} - \frac{3\sqrt[3]{\cos(a+bx)}}{b\sqrt[3]{\sin(a+bx)}} - \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{3}x+x^2} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{4b} \\
 &= -\frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} - \frac{\sqrt{3} \log\left(1 - \frac{\sqrt{3}\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} + \frac{\sqrt{3}}{b} \\
 &= \frac{\tan^{-1}\left(\sqrt{3} - \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\sqrt{3} + \frac{2\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{2b} - \frac{\tan^{-1}\left(\frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 55, normalized size = 0.22

$$-\frac{3\cos^2(a+bx)^{5/6} {}_2F_1\left(-\frac{1}{6}, -\frac{1}{6}; \frac{5}{6}; \sin^2(a+bx)\right)}{b\cos^{\frac{5}{3}}(a+bx)\sqrt[3]{\sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(4/3)/Sin[a + b*x]^(4/3), x]

[Out] (-3*(Cos[a + b*x]^2)^(5/6)*Hypergeometric2F1[-1/6, -1/6, 5/6, Sin[a + b*x]^2])/(b*Cos[a + b*x]^(5/3)*Sin[a + b*x]^(1/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{4}{3}}(bx+a)}{\sin^{\frac{4}{3}}(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^(4/3)/sin(b*x+a)^(4/3),x)`

[Out] `int(cos(b*x+a)^(4/3)/sin(b*x+a)^(4/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(4/3)/sin(b*x+a)^(4/3),x, algorithm="maxima")`

[Out] `integrate(cos(b*x + a)^(4/3)/sin(b*x + a)^(4/3), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(4/3)/sin(b*x+a)^(4/3),x, algorithm="fricas")`

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{4}{3}}(a + bx)}{\sin^{\frac{4}{3}}(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**(4/3)/sin(b*x+a)**(4/3),x)`

[Out] `Integral(cos(a + b*x)**(4/3)/sin(a + b*x)**(4/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(4/3)/sin(b*x+a)^(4/3),x, algorithm="giac")`

[Out] `integrate(cos(b*x + a)^(4/3)/sin(b*x + a)^(4/3), x)`

Mupad [B]

time = 1.70, size = 44, normalized size = 0.18

$$\frac{3 \cos(a + bx)^{7/3} (\sin(a + bx)^2)^{1/6} {}_2F_1\left(\frac{7}{6}, \frac{7}{6}; \frac{13}{6}; \cos(a + bx)^2\right)}{7b \sin(a + bx)^{1/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(4/3)/sin(a + b*x)^(4/3), x)

[Out] -(3*cos(a + b*x)^(7/3)*(sin(a + b*x)^2)^(1/6)*hypergeom([7/6, 7/6], 13/6, cos(a + b*x)^2))/(7*b*sin(a + b*x)^(1/3))

$$3.332 \quad \int \frac{\cos^5(a+bx)}{\sin^5(a+bx)} dx$$

Optimal. Leaf size=155

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - 2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b} - \frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)}$$

[Out] $1/2*\ln(1+\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)})/b-1/4*\ln(1-\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)}+\sin(b*x+a)^{(4/3)}/\cos(b*x+a)^{(4/3)})/b-3/2*\cos(b*x+a)^{(2/3)}/b/\sin(b*x+a)^{(2/3)}+1/2*\arctan(1/3*(1-2*\sin(b*x+a)^{(2/3)}/\cos(b*x+a)^{(2/3)})*3^{(1/2)})*3^{(1/2)}/b$

Rubi [A]

time = 0.08, antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2647, 2654, 281, 298, 31, 648, 632, 210, 642}

$$\frac{\sqrt{3} \text{ArcTan}\left(\frac{1 - 2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} + \frac{\log\left(\frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + 1\right)}{2b} - \frac{\log\left(\frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)} - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + 1\right)}{4b}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^(5/3)/Sin[a + b*x]^(5/3),x]

[Out] $(\text{Sqrt}[3]*\text{ArcTan}[(1 - (2*\text{Sin}[a + b*x]^{(2/3)})/\text{Cos}[a + b*x]^{(2/3)})/\text{Sqrt}[3]])/(2*b) + \text{Log}[1 + \text{Sin}[a + b*x]^{(2/3)}/\text{Cos}[a + b*x]^{(2/3)}]/(2*b) - \text{Log}[1 - \text{Sin}[a + b*x]^{(2/3)}/\text{Cos}[a + b*x]^{(2/3)} + \text{Sin}[a + b*x]^{(4/3)}/\text{Cos}[a + b*x]^{(4/3)}]/(4*b) - (3*\text{Cos}[a + b*x]^{(2/3)})/(2*b*\text{Sin}[a + b*x]^{(2/3)})$

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n_+1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 298

Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2647

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(a*cos[e + f*x])^(m - 1)*((b*sin[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*cos[e + f*x])^(m - 2)*(b*sin[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*sin[e + f*x])^(1/k)/(b*cos[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^{\frac{5}{3}}(a+bx)}{\sin^{\frac{5}{3}}(a+bx)} dx &= -\frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} - \int \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}} dx \\
&= -\frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} - \frac{3 \text{Subst}\left(\int \frac{x^3}{1+x^6} dx, x, \frac{\sqrt[3]{\sin(a+bx)}}{\sqrt[3]{\cos(a+bx)}}\right)}{b} \\
&= -\frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} - \frac{3 \text{Subst}\left(\int \frac{x}{1+x^3} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{3 \text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b} - \frac{3 \cos^{\frac{2}{3}}(a+bx)}{2b \sin^{\frac{2}{3}}(a+bx)} + \frac{3 \text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\sqrt{3} \tan^{-1}\left(\frac{1 - \frac{2 \sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} + \frac{\log\left(1 + \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{\log\left(1 - \frac{\sin^{\frac{2}{3}}(a+bx)}{\cos^{\frac{2}{3}}(a+bx)} + \frac{\sin^{\frac{4}{3}}(a+bx)}{\cos^{\frac{4}{3}}(a+bx)}\right)}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 57, normalized size = 0.37

$$-\frac{3 \cos^2(a+bx)^{2/3} {}_2F_1\left(-\frac{1}{3}, -\frac{1}{3}; \frac{2}{3}; \sin^2(a+bx)\right)}{2b \cos^{\frac{4}{3}}(a+bx) \sin^{\frac{2}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(5/3)/Sin[a + b*x]^(5/3), x]

[Out] (-3*(Cos[a + b*x]^2)^(2/3)*Hypergeometric2F1[-1/3, -1/3, 2/3, Sin[a + b*x]^2])/(2*b*Cos[a + b*x]^(4/3)*Sin[a + b*x]^(2/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{5}{3}}(bx+a)}{\sin^{\frac{5}{3}}(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^(5/3)/sin(b*x+a)^(5/3),x)

[Out] int(cos(b*x+a)^(5/3)/sin(b*x+a)^(5/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/3)/sin(b*x+a)^(5/3),x, algorithm="maxima")

[Out] integrate(cos(b*x + a)^(5/3)/sin(b*x + a)^(5/3), x)

Fricas [A]

time = 0.73, size = 189, normalized size = 1.22

$$\frac{2\sqrt{3}\arctan\left(\frac{-\sqrt{3}\cos(bx+a)-2\sqrt{3}\cos(bx+a)^{\frac{1}{3}}\sin(bx+a)^{\frac{2}{3}}}{3\cos(bx+a)}\right)\sin(bx+a)-2\log\left(\frac{\cos(bx+a)^{\frac{1}{3}}\sin(bx+a)^{\frac{2}{3}}+\cos(bx+a)}{\cos(bx+a)}\right)\sin(bx+a)+\log\left(\frac{\cos(bx+a)^2-\cos(bx+a)^{\frac{4}{3}}\sin(bx+a)^{\frac{2}{3}}+\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{4}{3}}}{\cos(bx+a)^2}\right)\sin(bx+a)+6\cos(bx+a)^{\frac{2}{3}}\sin(bx+a)^{\frac{1}{3}}}{4b\sin(bx+a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/3)/sin(b*x+a)^(5/3),x, algorithm="fricas")

[Out] $-1/4*(2*\sqrt{3}*\arctan(-1/3*(\sqrt{3}*\cos(b*x + a) - 2*\sqrt{3}*\cos(b*x + a)^{\frac{1}{3}}*\sin(b*x + a)^{\frac{2}{3}})/\cos(b*x + a))*\sin(b*x + a) - 2*\log((\cos(b*x + a)^{\frac{1}{3}}*\sin(b*x + a)^{\frac{2}{3}} + \cos(b*x + a))/\cos(b*x + a))*\sin(b*x + a) + \log((\cos(b*x + a)^2 - \cos(b*x + a)^{\frac{4}{3}}*\sin(b*x + a)^{\frac{2}{3}} + \cos(b*x + a)^{\frac{2}{3}}*\sin(b*x + a)^{\frac{4}{3}})/\cos(b*x + a)^2)*\sin(b*x + a) + 6*\cos(b*x + a)^{\frac{2}{3}}*\sin(b*x + a)^{\frac{1}{3}}/(b*\sin(b*x + a))$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**(5/3)/sin(b*x+a)**(5/3),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4369 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^(5/3)/sin(b*x+a)^(5/3),x, algorithm="giac")

[Out] integrate(cos(b*x + a)^(5/3)/sin(b*x + a)^(5/3), x)

Mupad [B]

time = 1.20, size = 44, normalized size = 0.28

$$\frac{3 \cos(a + bx)^{8/3} (\sin(a + bx)^2)^{1/3} {}_2F_1\left(\frac{4}{3}, \frac{4}{3}; \frac{7}{3}; \cos(a + bx)^2\right)}{8 b \sin(a + bx)^{2/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^(5/3)/sin(a + b*x)^(5/3),x)

[Out] -(3*cos(a + b*x)^(8/3)*(sin(a + b*x)^2)^(1/3)*hypergeom([4/3, 4/3], 7/3, cos(a + b*x)^2))/(8*b*sin(a + b*x)^(2/3))

$$3.333 \quad \int \frac{\cos^{\frac{7}{3}}(a+bx)}{\sin^{\frac{7}{3}}(a+bx)} dx$$

Optimal. Leaf size=155

$$\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - \frac{2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} + \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{3 \cos^{\frac{4}{3}}(a+bx)}{4b \sin^{\frac{4}{3}}(a+bx)}$$

[Out] $1/4*\ln(1+\cos(b*x+a)^{(4/3)}/\sin(b*x+a)^{(4/3)}-\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)})/b-1/2*\ln(1+\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)})/b-3/4*\cos(b*x+a)^{(4/3)}/b/\sin(b*x+a)^{(4/3)}-1/2*\arctan(1/3*(1-2*\cos(b*x+a)^{(2/3)}/\sin(b*x+a)^{(2/3)})*3^{(1/2)})*3^{(1/2)}/b$

Rubi [A]

time = 0.08, antiderivative size = 155, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 9, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {2647, 2655, 281, 298, 31, 648, 632, 210, 642}

$$\frac{\sqrt{3} \text{ArcTan}\left(\frac{1 - \frac{2 \cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}}{\sqrt{3}}\right)}{2b} - \frac{3 \cos^{\frac{4}{3}}(a+bx)}{4b \sin^{\frac{4}{3}}(a+bx)} + \frac{\log\left(\frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + 1\right)}{4b} - \frac{\log\left(\frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)} + 1\right)}{2b}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^(7/3)/Sin[a + b*x]^(7/3), x]`

[Out] $-1/2*(\text{Sqrt}[3]*\text{ArcTan}[(1 - (2*\text{Cos}[a + b*x]^{(2/3)})/\text{Sin}[a + b*x]^{(2/3)})/\text{Sqrt}[3]])/b + \text{Log}[1 + \text{Cos}[a + b*x]^{(4/3)}/\text{Sin}[a + b*x]^{(4/3)} - \text{Cos}[a + b*x]^{(2/3)}/\text{Sin}[a + b*x]^{(2/3)}]/(4*b) - \text{Log}[1 + \text{Cos}[a + b*x]^{(2/3)}/\text{Sin}[a + b*x]^{(2/3)}]/(2*b) - (3*\text{Cos}[a + b*x]^{(4/3)})/(4*b*\text{Sin}[a + b*x]^{(4/3)})$

Rule 31

`Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]`

Rule 210

`Int[((a_) + (b_.)*(x_)^2)^(n_), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(n_+1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

Rule 281

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]

Rule 298

Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] := Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 2647

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[a*(a*Cos[e + f*x])^(m - 1)*((b*SIN[e + f*x])^(n + 1)/(b*f*(n + 1))), x] + Dist[a^2*((m - 1)/(b^2*(n + 1))), Int[(a*Cos[e + f*x])^(m - 2)*(b*SIN[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && (IntegersQ[2*m, 2*n] || EqQ[m + n, 0])

Rule 2655

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*SIN[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rubi steps

$$\begin{aligned}
\int \frac{\cos^{\frac{7}{3}}(a+bx)}{\sin^{\frac{7}{3}}(a+bx)} dx &= -\frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} - \int \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}} dx \\
&= -\frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} + \frac{3\text{Subst}\left(\int \frac{x^3}{1+x^6} dx, x, \frac{\sqrt[3]{\cos(a+bx)}}{\sqrt[3]{\sin(a+bx)}}\right)}{b} \\
&= -\frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} + \frac{3\text{Subst}\left(\int \frac{x}{1+x^3} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} - \frac{\text{Subst}\left(\int \frac{1}{1+x} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} + \frac{\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} + \frac{\text{Subst}\left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} + \frac{3\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} - \frac{3\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} \\
&= -\frac{\sqrt{3} \tan^{-1}\left(\frac{1 - 2\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)\sqrt{3}}\right)}{2b} + \frac{\log\left(1 + \frac{\cos^{\frac{4}{3}}(a+bx)}{\sin^{\frac{4}{3}}(a+bx)} - \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{4b} - \frac{\log\left(1 + \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b} - \frac{3\cos^{\frac{4}{3}}(a+bx)}{4b\sin^{\frac{4}{3}}(a+bx)} - \frac{3\text{Subst}\left(\int \frac{1+x}{1-x+x^2} dx, x, \frac{\cos^{\frac{2}{3}}(a+bx)}{\sin^{\frac{2}{3}}(a+bx)}\right)}{2b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 57, normalized size = 0.37

$$-\frac{3\sqrt[3]{\cos^2(a+bx)} {}_2F_1\left(-\frac{2}{3}, -\frac{2}{3}; \frac{1}{3}; \sin^2(a+bx)\right)}{4b\cos^{\frac{2}{3}}(a+bx)\sin^{\frac{4}{3}}(a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^(7/3)/Sin[a + b*x]^(7/3), x]

[Out] (-3*(Cos[a + b*x]^2)^(1/3)*Hypergeometric2F1[-2/3, -2/3, 1/3, Sin[a + b*x]^2])/(4*b*Cos[a + b*x]^(2/3)*Sin[a + b*x]^(4/3))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{7}{3}}(bx + a)}{\sin(bx + a)^{\frac{7}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(b*x+a)^(7/3)/sin(b*x+a)^(7/3),x)`

[Out] `int(cos(b*x+a)^(7/3)/sin(b*x+a)^(7/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(7/3)/sin(b*x+a)^(7/3),x, algorithm="maxima")`

[Out] `integrate(cos(b*x + a)^(7/3)/sin(b*x + a)^(7/3), x)`

Fricas [A]

time = 0.61, size = 219, normalized size = 1.41

$$\frac{2(\sqrt{3} \cos(bx + a)^2 - \sqrt{3}) \arctan\left(\frac{2\sqrt{3} \cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{1}{3}} - \sqrt{3} \sin(bx+a)}{3 \sin(bx+a)}\right) + (\cos(bx + a)^2 - 1) \log\left(\frac{4(\cos(bx+a)^2 - \cos(bx+a)^{\frac{4}{3}} \sin(bx+a)^{\frac{2}{3}} + \cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{4}{3}} - 1)}{\cos(bx+a)^2 - 1}\right) - 2(\cos(bx + a)^2 - 1) \log\left(\frac{2(\cos(bx+a)^{\frac{2}{3}} \sin(bx+a)^{\frac{1}{3}} + \sin(bx+a))}{\sin(bx+a)}\right) + 3 \cos(bx + a)^{\frac{2}{3}} \sin(bx + a)^{\frac{1}{3}}}{4(b \cos(bx + a)^2 - b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(7/3)/sin(b*x+a)^(7/3),x, algorithm="fricas")`

[Out] `1/4*(2*(sqrt(3)*cos(b*x + a)^2 - sqrt(3))*arctan(1/3*(2*sqrt(3)*cos(b*x + a)^(2/3)*sin(b*x + a)^(1/3) - sqrt(3)*sin(b*x + a))/sin(b*x + a)) + (cos(b*x + a)^2 - 1)*log(4*(cos(b*x + a)^2 - cos(b*x + a)^(4/3)*sin(b*x + a)^(2/3) + cos(b*x + a)^(2/3)*sin(b*x + a)^(4/3) - 1)/(cos(b*x + a)^2 - 1)) - 2*(cos(b*x + a)^2 - 1)*log(-2*(cos(b*x + a)^(2/3)*sin(b*x + a)^(1/3) + sin(b*x + a))/sin(b*x + a)) + 3*cos(b*x + a)^(4/3)*sin(b*x + a)^(2/3)/(b*cos(b*x + a)^2 - b)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**(7/3)/sin(b*x+a)**(7/3),x)`

[Out] Timed out

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)^(7/3)/sin(b*x+a)^(7/3),x, algorithm="giac")`

[Out] Timed out

Mupad [B]

time = 1.91, size = 44, normalized size = 0.28

$$\frac{3 \cos(a + bx)^{10/3} (\sin(a + bx)^2)^{2/3} {}_2F_1\left(\frac{5}{3}, \frac{5}{3}; \frac{8}{3}; \cos(a + bx)^2\right)}{10 b \sin(a + bx)^{4/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(a + b*x)^(7/3)/sin(a + b*x)^(7/3),x)`

[Out] `-(3*cos(a + b*x)^(10/3)*(sin(a + b*x)^2)^(2/3)*hypergeom([5/3, 5/3], 8/3, cos(a + b*x)^2))/(10*b*sin(a + b*x)^(4/3))`

$$3.334 \quad \int \frac{\cos^{\frac{2}{3}}(x)}{\sin^{\frac{8}{3}}(x)} dx$$

Optimal. Leaf size=16

$$-\frac{3 \cos^{\frac{5}{3}}(x)}{5 \sin^{\frac{5}{3}}(x)}$$

[Out] $-3/5*\cos(x)^{(5/3)}/\sin(x)^{(5/3)}$

Rubi [A]

time = 0.02, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2643}

$$-\frac{3 \cos^{\frac{5}{3}}(x)}{5 \sin^{\frac{5}{3}}(x)}$$

Antiderivative was successfully verified.

[In] Int[Cos[x]^(2/3)/Sin[x]^(8/3),x]

[Out] $(-3*\cos[x]^{(5/3)})/(5*\sin[x]^{(5/3)})$

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] & & NeQ[m, -1]

Rubi steps

$$\int \frac{\cos^{\frac{2}{3}}(x)}{\sin^{\frac{8}{3}}(x)} dx = -\frac{3 \cos^{\frac{5}{3}}(x)}{5 \sin^{\frac{5}{3}}(x)}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$-\frac{3 \cos^{\frac{5}{3}}(x)}{5 \sin^{\frac{5}{3}}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[x]^(2/3)/Sin[x]^(8/3),x]

[Out] $(-3*\text{Cos}[x]^{(5/3)})/(5*\text{Sin}[x]^{(5/3)})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\cos^{\frac{2}{3}}(x)}{\sin(x)^{\frac{8}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(cos(x)^(2/3)/sin(x)^(8/3),x)`

[Out] `int(cos(x)^(2/3)/sin(x)^(8/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)^(2/3)/sin(x)^(8/3),x, algorithm="maxima")`

[Out] `integrate(cos(x)^(2/3)/sin(x)^(8/3), x)`

Fricas [A]

time = 0.61, size = 18, normalized size = 1.12

$$\frac{3 \cos(x)^{\frac{5}{3}} \sin(x)^{\frac{1}{3}}}{5 (\cos(x)^2 - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)^(2/3)/sin(x)^(8/3),x, algorithm="fricas")`

[Out] `3/5*cos(x)^(5/3)*sin(x)^(1/3)/(cos(x)^2 - 1)`

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(x)**(2/3)/sin(x)**(8/3),x)`

[Out] `Exception raised: SystemError >> excessive stack use: stack is 6189 deep`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(x)^(2/3)/sin(x)^(8/3),x, algorithm="giac")
```

```
[Out] integrate(cos(x)^(2/3)/sin(x)^(8/3), x)
```

Mupad [B]

time = 0.83, size = 10, normalized size = 0.62

$$-\frac{3 \cos(x)^{5/3}}{5 \sin(x)^{5/3}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(x)^(2/3)/sin(x)^(8/3),x)
```

```
[Out] -(3*cos(x)^(5/3))/(5*sin(x)^(5/3))
```

$$3.335 \quad \int \frac{\sin^{\frac{2}{3}}(x)}{\cos^{\frac{8}{3}}(x)} dx$$

Optimal. Leaf size=16

$$\frac{3 \sin^{\frac{5}{3}}(x)}{5 \cos^{\frac{5}{3}}(x)}$$

[Out] $3/5*\sin(x)^{(5/3)}/\cos(x)^{(5/3)}$

Rubi [A]

time = 0.01, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {2643}

$$\frac{3 \sin^{\frac{5}{3}}(x)}{5 \cos^{\frac{5}{3}}(x)}$$

Antiderivative was successfully verified.

[In] Int[Sin[x]^(2/3)/Cos[x]^(8/3),x]

[Out] (3*Sin[x]^(5/3))/(5*Cos[x]^(5/3))

Rule 2643

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Cos[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m + n + 2, 0] & & NeQ[m, -1]

Rubi steps

$$\int \frac{\sin^{\frac{2}{3}}(x)}{\cos^{\frac{8}{3}}(x)} dx = \frac{3 \sin^{\frac{5}{3}}(x)}{5 \cos^{\frac{5}{3}}(x)}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$\frac{3 \sin^{\frac{5}{3}}(x)}{5 \cos^{\frac{5}{3}}(x)}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[x]^(2/3)/Cos[x]^(8/3),x]

[Out] $(3*\sin(x)^{(5/3)})/(5*\cos(x)^{(5/3)})$

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{2}{3}}(x)}{\cos(x)^{\frac{8}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(x)^(2/3)/cos(x)^(8/3),x)`

[Out] `int(sin(x)^(2/3)/cos(x)^(8/3),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^(2/3)/cos(x)^(8/3),x, algorithm="maxima")`

[Out] `integrate(sin(x)^(2/3)/cos(x)^(8/3), x)`

Fricas [A]

time = 0.39, size = 10, normalized size = 0.62

$$\frac{3 \sin(x)^{\frac{5}{3}}}{5 \cos(x)^{\frac{5}{3}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)^(2/3)/cos(x)^(8/3),x, algorithm="fricas")`

[Out] `3/5*sin(x)^(5/3)/cos(x)^(5/3)`

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(x)**(2/3)/cos(x)**(8/3),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 6189 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(x)^(2/3)/cos(x)^(8/3),x, algorithm="giac")

[Out] integrate(sin(x)^(2/3)/cos(x)^(8/3), x)

Mupad [B]

time = 0.82, size = 94, normalized size = 5.88

$$\frac{6 \cdot 2^{2/3} \tan\left(\frac{x}{2}\right)^{5/3} \left(1 - \tan\left(\frac{x}{2}\right)^2\right)^{1/3} + 6 \cdot 2^{2/3} \tan\left(\frac{x}{2}\right)^{11/3} \left(1 - \tan\left(\frac{x}{2}\right)^2\right)^{1/3}}{5 \tan\left(\frac{x}{2}\right)^2 - \tan\left(\frac{x}{2}\right)^2 \left(10 \tan\left(\frac{x}{2}\right)^2 - 5 \tan\left(\frac{x}{2}\right)^2 \left(\tan\left(\frac{x}{2}\right)^2 + 1\right) + 10\right) + 5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(x)^(2/3)/cos(x)^(8/3),x)

[Out] (6*2^(2/3)*tan(x/2)^(5/3)*(1 - tan(x/2)^2)^(1/3) + 6*2^(2/3)*tan(x/2)^(11/3)
)*(1 - tan(x/2)^2)^(1/3)/(5*tan(x/2)^2 - tan(x/2)^2*(10*tan(x/2)^2 - 5*tan
(x/2)^2*(tan(x/2)^2 + 1) + 10) + 5)

3.336 $\int \cos^n(e + fx) \sin^m(e + fx) dx$

Optimal. Leaf size=80

$$-\frac{\cos^{1+n}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1+n)}$$

[Out] $-\cos(f*x+e)^{(1+n)}*\text{hypergeom}([1/2+1/2*n, 1/2-1/2*m], [3/2+1/2*n], \cos(f*x+e)^2)*\sin(f*x+e)^{(-1+m)}*(\sin(f*x+e)^2)^{(1/2-1/2*m)}/f/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.059$, Rules used = {2656}

$$-\frac{\sin^{m-1}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}} \cos^{n+1}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(e + fx)\right)}{f(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[e + f*x]^n*\text{Sin}[e + f*x]^m, x]$

[Out] $-\left(\left(\text{Cos}[e + f*x]^{(1+n)}*\text{Hypergeometric2F1}\left[\frac{(1-m)}{2}, \frac{(1+n)}{2}, \frac{(3+n)}{2}, \text{Cos}[e + f*x]^2\right]*\text{Sin}[e + f*x]^{(-1+m)}*(\text{Sin}[e + f*x]^2)^{\frac{(1-m)}{2}}\right)\right)/\left(f*(1+n)\right)$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^m)*((b_.)*\sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] \rightarrow \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*(a*\text{Cos}[e + f*x])^{(m+1)}/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]}))*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{SimplerQ}[n, m]$

Rubi steps

$$\int \cos^n(e + fx) \sin^m(e + fx) dx = -\frac{\cos^{1+n}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1+n)}$$

Mathematica [A]

time = 0.07, size = 79, normalized size = 0.99

$$\frac{\cos^{-1+n}(e + fx) \cos^2(e + fx)^{\frac{1-n}{2}} {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3+m}{2}; \sin^2(e + fx)\right) \sin^{1+m}(e + fx)}{f(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^n*Sin[e + f*x]^m,x]

[Out] (Cos[e + f*x]^(-1 + n)*(Cos[e + f*x]^2)^((1 - n)/2)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2]*Sin[e + f*x]^(1 + m))/(f*(1 + m))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int (\cos^n(fx + e)) (\sin^m(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^n*sin(f*x+e)^m,x)

[Out] int(cos(f*x+e)^n*sin(f*x+e)^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^n*sin(f*x+e)^m,x, algorithm="maxima")

[Out] integrate(cos(f*x + e)^n*sin(f*x + e)^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^n*sin(f*x+e)^m,x, algorithm="fricas")

[Out] integral(cos(f*x + e)^n*sin(f*x + e)^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^m(e + fx) \cos^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**n*sin(f*x+e)**m,x)

[Out] Integral(sin(e + f*x)**m*cos(e + f*x)**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^n*sin(f*x+e)^m,x, algorithm="giac")

[Out] integrate(cos(f*x + e)^n*sin(f*x + e)^m, x)

Mupad [B]

time = 2.35, size = 71, normalized size = 0.89

$$\frac{\cos(e + f x)^{n+1} \sin(e + f x)^{m+1} {}_2F_1\left(\frac{1}{2} - \frac{m}{2}, \frac{n}{2} + \frac{1}{2}; \frac{n}{2} + \frac{3}{2}; \cos(e + f x)^2\right)}{f (n + 1) (\sin(e + f x)^2)^{\frac{m}{2} + \frac{1}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^n*sin(e + f*x)^m,x)

[Out] -(cos(e + f*x)^(n + 1)*sin(e + f*x)^(m + 1)*hypergeom([1/2 - m/2, n/2 + 1/2], n/2 + 3/2, cos(e + f*x)^2))/(f*(n + 1)*(sin(e + f*x)^2)^(m/2 + 1/2))

3.337 $\int (d \cos(e + fx))^n \sin^m(e + fx) dx$

Optimal. Leaf size=85

$$\frac{(d \cos(e + fx))^{1+n} {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{df(1+n)}$$

[Out] $-(d*\cos(f*x+e))^{(1+n)}*\text{hypergeom}([1/2+1/2*n, 1/2-1/2*m], [3/2+1/2*n], \cos(f*x+e)^2)*\sin(f*x+e)^{-1+m}*(\sin(f*x+e)^2)^{(1/2-1/2*m)}/d/f/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2656}

$$\frac{\sin^{m-1}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}} (d \cos(e + fx))^{n+1} {}_2F_1\left(\frac{1-m}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(e + fx)\right)}{df(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[e + f*x])^n*\text{Sin}[e + f*x]^m, x]$

[Out] $-\left(\left(\left(d*\text{Cos}[e + f*x]\right)^{(1+n)}*\text{Hypergeometric2F1}\left[\frac{(1-m)}{2}, \frac{(1+n)}{2}, \frac{(3+n)}{2}, \text{Cos}[e + f*x]^2\right]*\text{Sin}[e + f*x]^{-1+m}*(\text{Sin}[e + f*x]^2)^{\left(\frac{1-m}{2}\right)}\right)\right)/\left(d*f*(1+n)\right)$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_)}], x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*((a*\text{Cos}[e + f*x])^{(m+1)}/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]})))*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(e + fx))^n \sin^m(e + fx) dx = -\frac{(d \cos(e + fx))^{1+n} {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{df(1+n)}$$

Mathematica [A]

time = 0.09, size = 82, normalized size = 0.96

$$\frac{d(d \cos(e + fx))^{-1+n} \cos^2(e + fx)^{\frac{1-n}{2}} {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3+m}{2}; \sin^2(e + fx)\right) \sin^{1+m}(e + fx)}{f(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[e + f*x])^n*sin[e + f*x]^m,x]

[Out] (d*(d*cos[e + f*x])^(-1 + n)*(Cos[e + f*x]^2)^((1 - n)/2)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2]*Sin[e + f*x]^(1 + m))/(f*(1 + m))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int (d \cos(fx + e))^n (\sin^m(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(f*x+e))^n*sin(f*x+e)^m,x)

[Out] int((d*cos(f*x+e))^n*sin(f*x+e)^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*sin(f*x+e)^m,x, algorithm="maxima")

[Out] integrate((d*cos(f*x + e))^n*sin(f*x + e)^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*sin(f*x+e)^m,x, algorithm="fricas")

[Out] integral((d*cos(f*x + e))^n*sin(f*x + e)^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(e + fx))^n \sin^m(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*sin(f*x+e)^m,x)

[Out] Integral((d*cos(e + f*x)**n*sin(e + f*x)**m, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*sin(f*x+e)^m,x, algorithm="giac")

[Out] integrate((d*cos(f*x + e))^n*sin(f*x + e)^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^m (d \cos(e + f x))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^m*(d*cos(e + f*x))^n,x)

[Out] int(sin(e + f*x)^m*(d*cos(e + f*x))^n, x)

3.338 $\int \cos^n(e + fx)(b \sin(e + fx))^m dx$

Optimal. Leaf size=83

$$-\frac{b \cos^{1+n}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) (b \sin(e + fx))^{-1+m} \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1+n)}$$

[Out] $-b \cos(f*x+e)^{(1+n)} \text{hypergeom}([1/2+1/2*n, 1/2-1/2*m], [3/2+1/2*n], \cos(f*x+e)^2) * (b \sin(f*x+e))^{(-1+m)} * (\sin(f*x+e)^2)^{(1/2-1/2*m)} / f / (1+n)$

Rubi [A]

time = 0.03, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2656}

$$-\frac{b \sin^2(e + fx)^{\frac{1-m}{2}} \cos^{n+1}(e + fx) (b \sin(e + fx))^{m-1} {}_2F_1\left(\frac{1-m}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(e + fx)\right)}{f(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Cos}[e + f*x]^n * (b * \text{Sin}[e + f*x])^m, x]$

[Out] $-((b * \text{Cos}[e + f*x]^{(1+n)} * \text{Hypergeometric2F1}[(1-m)/2, (1+n)/2, (3+n)/2, \text{Cos}[e + f*x]^2] * (b * \text{Sin}[e + f*x])^{(-1+m)} * (\text{Sin}[e + f*x]^2)^{((1-m)/2)}) / (f * (1+n))$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m)} * ((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n)}, x_Symbol] \rightarrow \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}) * (b * \text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])} * ((a * \text{Cos}[e + f*x])^{(m+1)} / (a * f * (m+1) * (\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]})) * \text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int \cos^n(e + fx)(b \sin(e + fx))^m dx = -\frac{b \cos^{1+n}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) (b \sin(e + fx))^{-1+m}}{f(1+n)}$$

Mathematica [A]

time = 0.05, size = 85, normalized size = 1.02

$$\frac{\cos^{-1+n}(e + fx) \cos^2(e + fx)^{\frac{1-n}{2}} {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3+m}{2}; \sin^2(e + fx)\right) \sin(e + fx) (b \sin(e + fx))^m}{f(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[e + f*x]^n*(b*Sin[e + f*x])^m,x]

[Out] (Cos[e + f*x]^(-1 + n)*(Cos[e + f*x]^2)^((1 - n)/2)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2]*Sin[e + f*x]*(b*Sin[e + f*x])^m)/(f*(1 + m))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int (\cos^n(fx + e)) (b \sin(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(f*x+e)^n*(b*sin(f*x+e))^m,x)

[Out] int(cos(f*x+e)^n*(b*sin(f*x+e))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^n*(b*sin(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*sin(f*x + e))^m*cos(f*x + e)^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^n*(b*sin(f*x+e))^m,x, algorithm="fricas")

[Out] integral((b*sin(f*x + e))^m*cos(f*x + e)^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sin(e + fx))^m \cos^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)**n*(b*sin(f*x+e))**m,x)

[Out] Integral((b*sin(e + f*x))**m*cos(e + f*x)**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(f*x+e)^n*(b*sin(f*x+e))^m,x, algorithm="giac")

[Out] integrate((b*sin(f*x + e))^m*cos(f*x + e)^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(e + f x)^n (b \sin(e + f x))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(e + f*x)^n*(b*sin(e + f*x))^m,x)

[Out] int(cos(e + f*x)^n*(b*sin(e + f*x))^m, x)

3.339 $\int (d \cos(e + fx))^n (b \sin(e + fx))^m dx$

Optimal. Leaf size=88

$$\frac{b(d \cos(e + fx))^{1+n} {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) (b \sin(e + fx))^{-1+m} \sin^2(e + fx)^{\frac{1-m}{2}}}{df(1+n)}$$

[Out] $-b*(d*\cos(f*x+e))^{(1+n)}*\text{hypergeom}([1/2+1/2*n, 1/2-1/2*m], [3/2+1/2*n], \cos(f*x+e)^2)*(b*\sin(f*x+e))^{(-1+m)}*(\sin(f*x+e)^2)^{(1/2-1/2*m)}/d/f/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {2656}

$$\frac{b \sin^2(e + fx)^{\frac{1-m}{2}} (b \sin(e + fx))^{m-1} (d \cos(e + fx))^{n+1} {}_2F_1\left(\frac{1-m}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(e + fx)\right)}{df(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[e + f*x])^n*(b*\text{Sin}[e + f*x])^m, x]$

[Out] $-((b*(d*\text{Cos}[e + f*x])^{(1+n)}*\text{Hypergeometric2F1}[(1-m)/2, (1+n)/2, (3+n)/2, \text{Cos}[e + f*x]^2]*(b*\text{Sin}[e + f*x])^{(-1+m)}*(\text{Sin}[e + f*x]^2)^{((1-m)/2)})/(d*f*(1+n))$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*((a*\text{Cos}[e + f*x])^{(m+1)}/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]}))*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(e + fx))^n (b \sin(e + fx))^m dx = -\frac{b(d \cos(e + fx))^{1+n} {}_2F_1\left(\frac{1-m}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(e + fx)\right) (b \sin(e + fx))^{-1+m} \sin^2(e + fx)^{\frac{1-m}{2}}}{df(1+n)}$$

Mathematica [A]

time = 0.07, size = 85, normalized size = 0.97

$$\frac{(d \cos(e + fx))^n \cos^2(e + fx)^{\frac{1-n}{2}} {}_2F_1\left(\frac{1+m}{2}, \frac{1-n}{2}; \frac{3+m}{2}; \sin^2(e + fx)\right) (b \sin(e + fx))^m \tan(e + fx)}{f(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[e + f*x])^n*(b*sin[e + f*x])^m,x]

[Out] ((d*cos[e + f*x])^n*(Cos[e + f*x]^2)^((1 - n)/2)*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2]*(b*sin[e + f*x])^m*Tan[e + f*x])/(f*(1 + m))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int (d \cos (fx + e))^n (b \sin (fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(f*x+e))^n*(b*sin(f*x+e))^m,x)

[Out] int((d*cos(f*x+e))^n*(b*sin(f*x+e))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*(b*sin(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((d*cos(f*x + e))^n*(b*sin(f*x + e))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*(b*sin(f*x+e))^m,x, algorithm="fricas")

[Out] integral((d*cos(f*x + e))^n*(b*sin(f*x + e))^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sin (e + fx))^m (d \cos (e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))**n*(b*sin(f*x+e))**m,x)

[Out] Integral((b*sin(e + f*x))**m*(d*cos(e + f*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(f*x+e))^n*(b*sin(f*x+e))^m,x, algorithm="giac")

[Out] integrate((d*cos(f*x + e))^n*(b*sin(f*x + e))^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(e + f x))^n (b \sin(e + f x))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(e + f*x))^n*(b*sin(e + f*x))^m,x)

[Out] int((d*cos(e + f*x))^n*(b*sin(e + f*x))^m, x)

3.340 $\int \cos^5(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=74

$$\frac{(c \sin(a + bx))^{1+m}}{bc(1+m)} - \frac{2(c \sin(a + bx))^{3+m}}{bc^3(3+m)} + \frac{(c \sin(a + bx))^{5+m}}{bc^5(5+m)}$$

[Out] (c*sin(b*x+a))^(1+m)/b/c/(1+m)-2*(c*sin(b*x+a))^(3+m)/b/c^3/(3+m)+(c*sin(b*x+a))^(5+m)/b/c^5/(5+m)

Rubi [A]

time = 0.05, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2644, 276}

$$\frac{(c \sin(a + bx))^{m+5}}{bc^5(m+5)} - \frac{2(c \sin(a + bx))^{m+3}}{bc^3(m+3)} + \frac{(c \sin(a + bx))^{m+1}}{bc(m+1)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^5*(c*Sin[a + b*x])^m,x]

[Out] (c*Sin[a + b*x])^(1 + m)/(b*c*(1 + m)) - (2*(c*Sin[a + b*x])^(3 + m))/(b*c^3*(3 + m)) + (c*Sin[a + b*x])^(5 + m)/(b*c^5*(5 + m))

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_.) + (b_.)*(x_)^(n_.))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^5(a + bx)(c \sin(a + bx))^m dx &= \frac{\text{Subst}\left(\int x^m \left(1 - \frac{x^2}{c^2}\right)^2 dx, x, c \sin(a + bx)\right)}{bc} \\ &= \frac{\text{Subst}\left(\int \left(x^m - \frac{2x^{2+m}}{c^2} + \frac{x^{4+m}}{c^4}\right) dx, x, c \sin(a + bx)\right)}{bc} \\ &= \frac{(c \sin(a + bx))^{1+m}}{bc(1+m)} - \frac{2(c \sin(a + bx))^{3+m}}{bc^3(3+m)} + \frac{(c \sin(a + bx))^{5+m}}{bc^5(5+m)} \end{aligned}$$

Mathematica [A]

time = 0.23, size = 55, normalized size = 0.74

$$\frac{\sin(a + bx)(c \sin(a + bx))^m \left(\frac{1}{1+m} - \frac{2 \sin^2(a+bx)}{3+m} + \frac{\sin^4(a+bx)}{5+m} \right)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^5*(c*Sin[a + b*x])^m,x]`

```
[Out] (Sin[a + b*x]*(c*Sin[a + b*x])^m*((1 + m)^(-1) - (2*Sin[a + b*x]^2)/(3 + m)
+ Sin[a + b*x]^4/(5 + m)))/b
```

Maple [F]

time = 0.09, size = 0, normalized size = 0.00

$$\int (\cos^5(bx + a)) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^5*(c*sin(b*x+a))^m,x)``[Out] int(cos(b*x+a)^5*(c*sin(b*x+a))^m,x)`**Maxima [A]**

time = 0.29, size = 77, normalized size = 1.04

$$\frac{\frac{c^m \sin(bx+a)^m \sin(bx+a)^5}{m+5} - \frac{2 c^m \sin(bx+a)^m \sin(bx+a)^3}{m+3} + \frac{(c \sin(bx+a))^{m+1}}{c(m+1)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5*(c*sin(b*x+a))^m,x, algorithm="maxima")`

```
[Out] (c^m*sin(b*x + a)^m*sin(b*x + a)^5/(m + 5) - 2*c^m*sin(b*x + a)^m*sin(b*x +
a)^3/(m + 3) + (c*sin(b*x + a))^(m + 1)/(c*(m + 1)))/b
```

Fricas [A]

time = 0.41, size = 70, normalized size = 0.95

$$\frac{((m^2 + 4m + 3) \cos(bx + a)^4 + 4(m + 1) \cos(bx + a)^2 + 8)(c \sin(bx + a))^m \sin(bx + a)}{bm^3 + 9bm^2 + 23bm + 15b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^5*(c*sin(b*x+a))^m,x, algorithm="fricas")`

```
[Out] ((m^2 + 4*m + 3)*cos(b*x + a)^4 + 4*(m + 1)*cos(b*x + a)^2 + 8)*(c*sin(b*x
+ a))^m*sin(b*x + a)/(b*m^3 + 9*b*m^2 + 23*b*m + 15*b)
```

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 2040 vs. $2(60) = 120$.

time = 5.69, size = 2040, normalized size = 27.57

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(cos(b*x+a)**5*(c*sin(b*x+a))**m,x)`

[Out] `Piecewise((x*(c*sin(a))**m*cos(a)**5, Eq(b, 0)), ((log(sin(a + b*x))/b + cos(a + b*x)**2/(2*b*sin(a + b*x)**2) - cos(a + b*x)**4/(4*b*sin(a + b*x)**4))/c**5, Eq(m, -5)), ((16*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**6/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 32*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 16*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 16*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 32*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 16*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - tan(a/2 + b*x/2)**8/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) + 18*tan(a/2 + b*x/2)**4/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2) - 1/(8*b*tan(a/2 + b*x/2)**6 + 16*b*tan(a/2 + b*x/2)**4 + 8*b*tan(a/2 + b*x/2)**2))/c**3, Eq(m, -3)), ((-log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 6*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2)**2 + 1)/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 4*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 6*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 4*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)`

```

)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*tan(a/2 +
  b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 +
  b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*tan(a/2 + b*x/2)**4/(b*tan(a/
  2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan
  (a/2 + b*x/2)**2 + b) - 4*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*
  tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b
  ))/c, Eq(m, -1)), (m**2*(c*sin(a + b*x))**m*sin(a + b*x)*cos(a + b*x)**4/(b
  *m**3 + 9*b*m**2 + 23*b*m + 15*b) + 4*m*(c*sin(a + b*x))**m*sin(a + b*x)**3
  *cos(a + b*x)**2/(b*m**3 + 9*b*m**2 + 23*b*m + 15*b) + 8*m*(c*sin(a + b*x))
  **m*sin(a + b*x)*cos(a + b*x)**4/(b*m**3 + 9*b*m**2 + 23*b*m + 15*b) + 8*(c
  *sin(a + b*x))**m*sin(a + b*x)**5/(b*m**3 + 9*b*m**2 + 23*b*m + 15*b) + 20*
  (c*sin(a + b*x))**m*sin(a + b*x)**3*cos(a + b*x)**2/(b*m**3 + 9*b*m**2 + 23
  *b*m + 15*b) + 15*(c*sin(a + b*x))**m*sin(a + b*x)*cos(a + b*x)**4/(b*m**3
  + 9*b*m**2 + 23*b*m + 15*b), True))

```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 248 vs. 2(74) = 148.

time = 5.65, size = 248, normalized size = 3.35

$(c \sin(bx+a))^m c^{m^2} \sin(bx+a)^5 + 4(c \sin(bx+a))^m c^m \sin(bx+a)^5 - 2(c \sin(bx+a))^m c^{m^2} \sin(bx+a)^5 + 3(c \sin(bx+a))^m c^m \sin(bx+a)^5 - 12(c \sin(bx+a))^m c^m \sin(bx+a)^5 + (c \sin(bx+a))^m c^{m^2} \sin(bx+a) - 10(c \sin(bx+a))^m c^m \sin(bx+a)^5 + 8(c \sin(bx+a))^m c^m \sin(bx+a) + 15(c \sin(bx+a))^m c^m \sin(bx+a) - (c^{m^3} + 9c^{m^2} + 23c^m + 15c^4)bc$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^5*(c*sin(b*x+a))^m,x, algorithm="giac")

```

[Out] ((c*sin(b*x + a))^m*c^5*m^2*sin(b*x + a)^5 + 4*(c*sin(b*x + a))^m*c^5*m*sin
(b*x + a)^5 - 2*(c*sin(b*x + a))^m*c^5*m^2*sin(b*x + a)^3 + 3*(c*sin(b*x +
a))^m*c^5*sin(b*x + a)^5 - 12*(c*sin(b*x + a))^m*c^5*m*sin(b*x + a)^3 + (c*
sin(b*x + a))^m*c^5*m^2*sin(b*x + a) - 10*(c*sin(b*x + a))^m*c^5*sin(b*x +
a)^3 + 8*(c*sin(b*x + a))^m*c^5*m*sin(b*x + a) + 15*(c*sin(b*x + a))^m*c^5*
sin(b*x + a))/((c^4*m^3 + 9*c^4*m^2 + 23*c^4*m + 15*c^4)*b*c)

```

Mupad [B]

time = 1.61, size = 132, normalized size = 1.78

$(c \sin(a + bx))^m (150 \sin(a + bx) + 25 \sin(3a + 3bx) + 3 \sin(5a + 5bx) + 24m \sin(a + bx) + 28m \sin(3a + 3bx) + 4m \sin(5a + 5bx) + 2m^2 \sin(a + bx) + 3m^2 \sin(3a + 3bx) + m^2 \sin(5a + 5bx)) / (16b(m^3 + 9m^2 + 23m + 15))$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^5*(c*sin(a + b*x))^m,x)

```

[Out] ((c*sin(a + b*x))^m*(150*sin(a + b*x) + 25*sin(3*a + 3*b*x) + 3*sin(5*a + 5
*b*x) + 24*m*sin(a + b*x) + 28*m*sin(3*a + 3*b*x) + 4*m*sin(5*a + 5*b*x) +
2*m^2*sin(a + b*x) + 3*m^2*sin(3*a + 3*b*x) + m^2*sin(5*a + 5*b*x)))/(16*b*
(23*m + 9*m^2 + m^3 + 15))

```

3.341 $\int \cos^3(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=50

$$\frac{(c \sin(a + bx))^{1+m}}{bc(1+m)} - \frac{(c \sin(a + bx))^{3+m}}{bc^3(3+m)}$$

[Out] (c*sin(b*x+a))^(1+m)/b/c/(1+m)-(c*sin(b*x+a))^(3+m)/b/c^3/(3+m)

Rubi [A]

time = 0.03, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2644, 14}

$$\frac{(c \sin(a + bx))^{m+1}}{bc(m+1)} - \frac{(c \sin(a + bx))^{m+3}}{bc^3(m+3)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^3*(c*Sin[a + b*x])^m,x]

[Out] (c*Sin[a + b*x])^(1 + m)/(b*c*(1 + m)) - (c*Sin[a + b*x])^(3 + m)/(b*c^3*(3 + m))

Rule 14

Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos^3(a + bx)(c \sin(a + bx))^m dx &= \frac{\text{Subst}\left(\int x^m \left(1 - \frac{x^2}{c^2}\right) dx, x, c \sin(a + bx)\right)}{bc} \\ &= \frac{\text{Subst}\left(\int \left(x^m - \frac{x^{2+m}}{c^2}\right) dx, x, c \sin(a + bx)\right)}{bc} \\ &= \frac{(c \sin(a + bx))^{1+m}}{bc(1+m)} - \frac{(c \sin(a + bx))^{3+m}}{bc^3(3+m)} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 48, normalized size = 0.96

$$\frac{(5 + m + (1 + m) \cos(2(a + bx))) \sin(a + bx) (c \sin(a + bx))^m}{2b(1 + m)(3 + m)}$$

Antiderivative was successfully verified.

`[In] Integrate[Cos[a + b*x]^3*(c*Sin[a + b*x])^m,x]``[Out] ((5 + m + (1 + m)*Cos[2*(a + b*x)])*Sin[a + b*x]*(c*Sin[a + b*x])^m)/(2*b*(1 + m)*(3 + m))`**Maple [C]** Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 1.27, size = 1318, normalized size = 26.36

method	result	size
risch	Expression too large to display	1318

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(b*x+a)^3*(c*sin(b*x+a))^m,x,method=_RETURNVERBOSE)`

```
[Out] 1/8*I/(b*m+3*b)*c^m*(1/2)^m*(exp(2*I*(b*x+a))-1)^m*exp(I*(b*x+a))^(m)*exp(-1/2*I*(-Pi*m*csgn(c*sin(b*x+a))^3+Pi*m*csgn(sin(b*x+a))*csgn(c*sin(b*x+a))^2-Pi*m*csgn(I*c)*csgn(c*sin(b*x+a))^2+Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))^2+Pi*m*csgn(sin(b*x+a))*csgn(I*c)*csgn(c*sin(b*x+a))+Pi*m*csgn(I*c*sin(b*x+a))^3-Pi*m*csgn(sin(b*x+a))^3-Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(sin(b*x+a))^2-Pi*m*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))^2-Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))-Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))-Pi*m*csgn(I*c*sin(b*x+a))^2+Pi*m+6*b*x+6*a)+1/8*I*exp(I*(b*x+a))^(m)*(exp(2*I*(b*x+a))-1)^m*(1/2)^m*c^m/(3+m)/(1+m)/b*(m+9)*exp(-1/2*I*(-Pi*m*csgn(c*sin(b*x+a))^3+Pi*m*csgn(sin(b*x+a))*csgn(c*sin(b*x+a))^2-Pi*m*csgn(I*c)*csgn(c*sin(b*x+a))^2+Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))^2+Pi*m*csgn(sin(b*x+a))*csgn(I*c)*csgn(c*sin(b*x+a))+Pi*m*csgn(I*c*sin(b*x+a))^3-Pi*m*csgn(sin(b*x+a))^3-Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(sin(b*x+a))^2-Pi*m*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))^2-Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))-Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))-Pi*m*csgn(I*c*sin(b*x+a))^2+Pi*m+2*b*x+2*a)-1/8*I*exp(I*(b*x+a))^(m)*(exp(2*I*(b*x+a))-1)^m*(1/2)^m*c^m/(b*m+3*b)*exp(1/2*I*(Pi*m*csgn(c*sin(b*x+a))^3-Pi*m*csgn(sin(b*x+a))*csgn(c*sin(b*x+a))^2+Pi*m*csgn(I*c)*csgn(c*sin(b*x+a))^2-Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))^2-Pi*m*csgn(sin(b*x+a))*csgn(I*c)*csgn(c*sin(b*x+a))-Pi*m*csgn(I*c*sin(b*x+a))^3+Pi*m*csgn(sin(b*x+a))^3+Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(sin(b*x+a))^2+Pi*m*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))^2+Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))+Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))+Pi*m*csgn(I*c*sin(b*x+a))
```

a))^{-2-Pi*m+6*b*x+6*a})-1/8*I*exp(I*(b*x+a))^(-m)*(exp(2*I*(b*x+a))-1)^m*(1/2)^m*c^m/(3+m)/(1+m)/b*(m+9)*exp(1/2*I*(Pi*m*csgn(c*sin(b*x+a))³-Pi*m*csgn(sin(b*x+a))*csgn(c*sin(b*x+a))²+Pi*m*csgn(I*c)*csgn(c*sin(b*x+a))²-Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))²-Pi*m*csgn(sin(b*x+a))*csgn(I*c)*csgn(c*sin(b*x+a))-Pi*m*csgn(I*c*sin(b*x+a))³+Pi*m*csgn(sin(b*x+a))³+Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(sin(b*x+a))²+Pi*m*csgn(I*exp(-I*(b*x+a)))*csgn(sin(b*x+a))²+Pi*m*csgn(I*(exp(2*I*(b*x+a))-1))*csgn(I*exp(-I*(b*x+a))))*csgn(sin(b*x+a))+Pi*m*csgn(c*sin(b*x+a))*csgn(I*c*sin(b*x+a))+Pi*m*csgn(I*c*sin(b*x+a))²-Pi*m+2*b*x+2*a)

Maxima [A]

time = 0.35, size = 53, normalized size = 1.06

$$\frac{\frac{c^m \sin(bx+a)^m \sin(bx+a)^3}{m+3} - \frac{(c \sin(bx+a))^{m+1}}{c(m+1)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] -(c^m*sin(b*x + a)^m*sin(b*x + a)³/(m + 3) - (c*sin(b*x + a))^(m + 1)/(c*(m + 1)))/b

Fricas [A]

time = 0.42, size = 46, normalized size = 0.92

$$\frac{((m + 1) \cos(bx + a)^2 + 2)(c \sin(bx + a))^m \sin(bx + a)}{bm^2 + 4bm + 3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^3*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] ((m + 1)*cos(b*x + a)² + 2)*(c*sin(b*x + a))^m*sin(b*x + a)/(b*m² + 4*b*m + 3*b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 525 vs. 2(37) = 74.

time = 3.18, size = 525, normalized size = 10.50

$$\begin{cases} x(c \sin(a))^m \cos^3(a) & \text{for } b = 0 \\ -\frac{\log(\sin(a+bx)) - \cos^2(a+bx)}{b} & \text{for } m = -3 \\ \frac{\log(\tan^2(\frac{g+\frac{bx}{c}}{c})+1) \tan^4(\frac{g+\frac{bx}{c}}{c}) - 2 \log(\tan^2(\frac{g+\frac{bx}{c}}{c})+1) \tan^2(\frac{g+\frac{bx}{c}}{c}) - \log(\tan^2(\frac{g+\frac{bx}{c}}{c})+1) + \log(\tan(\frac{g+\frac{bx}{c}}{c}) \tan^4(\frac{g+\frac{bx}{c}}{c}) + 2 \log(\tan(\frac{g+\frac{bx}{c}}{c}) \tan^2(\frac{g+\frac{bx}{c}}{c}))}{b \tan^4(\frac{g+\frac{bx}{c}}{c}) + 2b \tan^2(\frac{g+\frac{bx}{c}}{c}) + b} + \frac{\log(\tan(\frac{g+\frac{bx}{c}}{c}))}{b \tan^4(\frac{g+\frac{bx}{c}}{c}) + 2b \tan^2(\frac{g+\frac{bx}{c}}{c}) + b} - \frac{2 \tan^2(\frac{g+\frac{bx}{c}}{c})}{b \tan^4(\frac{g+\frac{bx}{c}}{c}) + 2b \tan^2(\frac{g+\frac{bx}{c}}{c}) + b}}{c} & \text{for } m = -1 \\ \frac{m(c \sin(a+bx))^m \sin(a+bx) \cos^2(a+bx)}{bm^2+4bm+3b} + \frac{2(c \sin(a+bx))^m \sin^2(a+bx)}{bm^2+4bm+3b} + \frac{3(c \sin(a+bx))^m \sin(a+bx) \cos^2(a+bx)}{bm^2+4bm+3b} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**3*(c*sin(b*x+a))**m,x)

```
[Out] Piecewise((x*(c*sin(a))*m*cos(a)**3, Eq(b, 0)), ((-log(sin(a + b*x))/b - c
os(a + b*x)**2/(2*b*sin(a + b*x)**2))/c**3, Eq(m, -3)), ((-log(tan(a/2 + b*
x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x
/2)**2 + b) - 2*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2
 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2)**2 + 1)/
(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2
))*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b
) + 2*log(tan(a/2 + b*x/2))*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*
b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2))/(b*tan(a/2 + b*x/2)**4 +
 2*b*tan(a/2 + b*x/2)**2 + b) - 2*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**
4 + 2*b*tan(a/2 + b*x/2)**2 + b))/c, Eq(m, -1)), (m*(c*sin(a + b*x))*m*sin
(a + b*x)*cos(a + b*x)**2/(b*m**2 + 4*b*m + 3*b) + 2*(c*sin(a + b*x))*m*si
n(a + b*x)**3/(b*m**2 + 4*b*m + 3*b) + 3*(c*sin(a + b*x))*m*sin(a + b*x)*c
os(a + b*x)**2/(b*m**2 + 4*b*m + 3*b), True))
```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 118 vs. 2(50) = 100.

time = 3.18, size = 118, normalized size = 2.36

$$\frac{(c \sin(bx + a))^m c^3 m \sin(bx + a)^3 + (c \sin(bx + a))^m c^3 \sin(bx + a)^3 - (c \sin(bx + a))^m c^3 m \sin(bx + a) - 3(c \sin(bx + a))^m c^3 \sin(bx + a)}{(c^2 m^2 + 4c^2 m + 3c^2)bc}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(cos(b*x+a)^3*(c*sin(b*x+a))^m,x, algorithm="giac")
```

```
[Out] -((c*sin(b*x + a))^m*c^3*m*sin(b*x + a)^3 + (c*sin(b*x + a))^m*c^3*sin(b*x
+ a)^3 - (c*sin(b*x + a))^m*c^3*m*sin(b*x + a) - 3*(c*sin(b*x + a))^m*c^3*s
in(b*x + a))/((c^2*m^2 + 4*c^2*m + 3*c^2)*b*c)
```

Mupad [B]

time = 0.88, size = 62, normalized size = 1.24

$$\frac{(c \sin(a + bx))^m (9 \sin(a + bx) + \sin(3a + 3bx) + m \sin(a + bx) + m \sin(3a + 3bx))}{4b(m^2 + 4m + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(cos(a + b*x)^3*(c*sin(a + b*x))^m,x)
```

```
[Out] ((c*sin(a + b*x))^m*(9*sin(a + b*x) + sin(3*a + 3*b*x) + m*sin(a + b*x) + m
*sin(3*a + 3*b*x)))/(4*b*(4*m + m^2 + 3))
```

3.342 $\int \cos(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=24

$$\frac{(c \sin(a + bx))^{1+m}}{bc(1+m)}$$

[Out] (c*sin(b*x+a))^(1+m)/b/c/(1+m)

Rubi [A]

time = 0.02, antiderivative size = 24, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 30}

$$\frac{(c \sin(a + bx))^{m+1}}{bc(m+1)}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]*(c*Sin[a + b*x])^m,x]

[Out] (c*Sin[a + b*x])^(1 + m)/(b*c*(1 + m))

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \cos(a + bx)(c \sin(a + bx))^m dx &= \frac{\text{Subst}(\int x^m dx, x, c \sin(a + bx))}{bc} \\ &= \frac{(c \sin(a + bx))^{1+m}}{bc(1+m)} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 25, normalized size = 1.04

$$\frac{\sin(a + bx)(c \sin(a + bx))^m}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]*(c*Sin[a + b*x])^m,x]

[Out] (Sin[a + b*x]*(c*Sin[a + b*x])^m)/(b*(1 + m))

Maple [A]

time = 1.11, size = 25, normalized size = 1.04

method	result	size
derivatividivides	$\frac{(c \sin(bx+a))^{1+m}}{bc(1+m)}$	25
default	$\frac{(c \sin(bx+a))^{1+m}}{bc(1+m)}$	25
norman	$\frac{2 \tan\left(\frac{bx+a}{2}\right) e^{m \ln\left(\frac{2c \tan\left(\frac{bx+a}{2}\right)}{1+\tan^2\left(\frac{bx+a}{2}\right)}\right)}}{b(1+m)\left(1+\tan^2\left(\frac{bx+a}{2}\right)\right)}$	66
risch	Expression too large to display	1016

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)*(c*sin(b*x+a))^m,x,method=_RETURNVERBOSE)

[Out] (c*sin(b*x+a))^(1+m)/b/c/(1+m)

Maxima [A]

time = 0.32, size = 24, normalized size = 1.00

$$\frac{(c \sin (bx + a))^{m+1}}{bc(m + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] (c*sin(b*x + a))^(m + 1)/(b*c*(m + 1))

Fricas [A]

time = 0.41, size = 24, normalized size = 1.00

$$\frac{(c \sin (bx + a))^m \sin (bx + a)}{bm + b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] (c*sin(b*x + a))^m*sin(b*x + a)/(b*m + b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 56 vs. $2(17) = 34$.

time = 0.46, size = 56, normalized size = 2.33

$$\begin{cases} \frac{x \cos(a)}{c \sin(a)} & \text{for } b = 0 \wedge m = -1 \\ x(c \sin(a))^m \cos(a) & \text{for } b = 0 \\ \frac{\log(\sin(a+bx))}{bc} & \text{for } m = -1 \\ \frac{(c \sin(a+bx))^m \sin(a+bx)}{bm+b} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*(c*sin(b*x+a))**m,x)

[Out] Piecewise((x*cos(a)/(c*sin(a)), Eq(b, 0) & Eq(m, -1)), (x*(c*sin(a))**m*cos(a), Eq(b, 0)), (log(sin(a + b*x))/(b*c), Eq(m, -1)), ((c*sin(a + b*x))**m*sin(a + b*x)/(b*m + b), True))

Giac [A]

time = 3.24, size = 24, normalized size = 1.00

$$\frac{(c \sin(bx + a))^{m+1}}{bc(m + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] (c*sin(b*x + a))^(m + 1)/(b*c*(m + 1))

Mupad [B]

time = 0.61, size = 25, normalized size = 1.04

$$\frac{\sin(a + bx) (c \sin(a + bx))^m}{b (m + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)*(c*sin(a + b*x))^m,x)

[Out] (sin(a + b*x)*(c*sin(a + b*x))^m)/(b*(m + 1))

3.343 $\int \sec(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=48

$$\frac{{}_2F_1\left(1, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)}$$

[Out] hypergeom([1, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)

Rubi [A]

time = 0.03, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2644, 371}

$$\frac{(c \sin(a + bx))^{m+1} {}_2F_1\left(1, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]*(c*Sin[a + b*x])^m,x]

[Out] (Hypergeometric2F1[1, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n - 1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n - 1)/2] && !(IntegerQ[(m - 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \sec(a + bx)(c \sin(a + bx))^m dx &= \frac{\text{Subst}\left(\int \frac{x^m}{1-x^2} dx, x, c \sin(a + bx)\right)}{bc} \\ &= \frac{{}_2F_1\left(1, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 51, normalized size = 1.06

$$\frac{{}_2F_1\left(1, \frac{1+m}{2}; 1 + \frac{1+m}{2}; \sin^2(a+bx)\right) \sin(a+bx) (c \sin(a+bx))^m}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]*(c*Sin[a + b*x])^m,x]

[Out] (Hypergeometric2F1[1, (1 + m)/2, 1 + (1 + m)/2, Sin[a + b*x]^2]*Sin[a + b*x]*(c*Sin[a + b*x])^m)/(b*(1 + m))

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int \sec(bx + a) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)*(c*sin(b*x+a))^m,x)

[Out] int(sec(b*x+a)*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m*sec(b*x + a), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m \sec(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)*(c*sin(b*x+a))**m,x)`

[Out] `Integral((c*sin(a + b*x))**m*sec(a + b*x), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(b*x+a)*(c*sin(b*x+a))^m,x, algorithm="giac")`

[Out] `integrate((c*sin(b*x + a))^m*sec(b*x + a), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(c \sin(a + bx))^m}{\cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(a + b*x))^m/cos(a + b*x),x)`

[Out] `int((c*sin(a + b*x))^m/cos(a + b*x), x)`

3.344 $\int \sec^3(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=48

$$\frac{{}_2F_1\left(2, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)}$$

[Out] hypergeom([2, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)

Rubi [A]

time = 0.03, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2644, 371}

$$\frac{(c \sin(a + bx))^{m+1} {}_2F_1\left(2, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^3*(c*Sin[a + b*x])^m,x]

[Out] (Hypergeometric2F1[2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m+1)/(c*(m+1)))*Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 2644

Int[cos[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Dist[1/(a*f), Subst[Int[x^m*(1 - x^2/a^2)^((n-1)/2), x], x, a*Sin[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \sec^3(a + bx)(c \sin(a + bx))^m dx &= \frac{\text{Subst}\left(\int \frac{x^m}{(1-\frac{x^2}{c^2})^2} dx, x, c \sin(a + bx)\right)}{bc} \\ &= \frac{{}_2F_1\left(2, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 51, normalized size = 1.06

$$\frac{{}_2F_1\left(2, \frac{1+m}{2}; 1 + \frac{1+m}{2}; \sin^2(a+bx)\right) \sin(a+bx) (c \sin(a+bx))^m}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^3*(c*Sin[a + b*x])^m,x]

[Out] (Hypergeometric2F1[2, (1 + m)/2, 1 + (1 + m)/2, Sin[a + b*x]^2]*Sin[a + b*x]^m)/(b*(1 + m))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int (\sec^3(bx + a)) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^3*(c*sin(b*x+a))^m,x)

[Out] int(sec(b*x+a)^3*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a)^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m*sec(b*x + a)^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m \sec^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**3*(c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m*sec(a + b*x)**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^3*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(c \sin(a + b x))^m}{\cos(a + b x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/cos(a + b*x)^3,x)

[Out] int((c*sin(a + b*x))^m/cos(a + b*x)^3, x)

3.345 $\int \cos^4(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=68

$$\frac{\cos(a + bx) {}_2F_1\left(-\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)\sqrt{\cos^2(a + bx)}}$$

[Out] `cos(b*x+a)*hypergeom([-3/2, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)/(cos(b*x+a)^2)^(1/2)`

Rubi [A]

time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2657}

$$\frac{\cos(a + bx)(c \sin(a + bx))^{m+1} {}_2F_1\left(-\frac{3}{2}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)\sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] `Int[Cos[a + b*x]^4*(c*Sin[a + b*x])^m,x]`

[Out] `(Cos[a + b*x]*Hypergeometric2F1[-3/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m)*Sqrt[Cos[a + b*x]^2])`

Rule 2657

`Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2])]*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]`

Rubi steps

$$\int \cos^4(a + bx)(c \sin(a + bx))^m dx = \frac{\cos(a + bx) {}_2F_1\left(-\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)\sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.04, size = 63, normalized size = 0.93

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(-\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^4*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[-3/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m))

Maple [F]

time = 0.09, size = 0, normalized size = 0.00

$$\int (\cos^4(bx + a)) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^4*(c*sin(b*x+a))^m,x)

[Out] int(cos(b*x+a)^4*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m*cos(b*x + a)^4, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^4*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m*cos(b*x + a)^4, x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**4*(c*sin(b*x+a))**m,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(cos(b*x+a)^4*(c*sin(b*x+a))^m,x, algorithm="giac")``[Out] integrate((c*sin(b*x + a))^m*cos(b*x + a)^4, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(a + bx)^4 (c \sin(a + bx))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(cos(a + b*x)^4*(c*sin(a + b*x))^m,x)``[Out] int(cos(a + b*x)^4*(c*sin(a + b*x))^m, x)`

3.346 $\int \cos^2(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=68

$$\frac{\cos(a + bx) {}_2F_1\left(-\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)\sqrt{\cos^2(a + bx)}}$$

[Out] cos(b*x+a)*hypergeom([-1/2, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)/(cos(b*x+a)^2)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2657}

$$\frac{\cos(a + bx)(c \sin(a + bx))^{m+1} {}_2F_1\left(-\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)\sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Cos[a + b*x]^2*(c*Sin[a + b*x])^m,x]

[Out] (Cos[a + b*x]*Hypergeometric2F1[-1/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m)*Sqrt[Cos[a + b*x]^2])

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \cos^2(a + bx)(c \sin(a + bx))^m dx = \frac{\cos(a + bx) {}_2F_1\left(-\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)\sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.04, size = 63, normalized size = 0.93

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(-\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Cos[a + b*x]^2*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[-1/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m))

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (\cos^2(bx + a)) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(b*x+a)^2*(c*sin(b*x+a))^m,x)

[Out] int(cos(b*x+a)^2*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m*cos(b*x + a)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m*cos(b*x + a)^2, x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)**2*(c*sin(b*x+a))**m,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(cos(b*x+a)^2*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m*cos(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \cos(a + bx)^2 (c \sin(a + bx))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(cos(a + b*x)^2*(c*sin(a + b*x))^m,x)

[Out] int(cos(a + b*x)^2*(c*sin(a + b*x))^m, x)

3.347 $\int (c \sin(a + bx))^m dx$

Optimal. Leaf size=68

$$\frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m) \sqrt{\cos^2(a + bx)}}$$

[Out] `cos(b*x+a)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)/(cos(b*x+a)^2)^(1/2)`

Rubi [A]

time = 0.01, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$\frac{\cos(a + bx)(c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1) \sqrt{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] `Int[(c*Sin[a + b*x])^m,x]`

[Out] `(Cos[a + b*x]*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m)*Sqrt[Cos[a + b*x]^2])`

Rule 2722

`Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

Rubi steps

$$\int (c \sin(a + bx))^m dx = \frac{\cos(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m) \sqrt{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.03, size = 63, normalized size = 0.93

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[1/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^m,x)

[Out] int((c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m,x)

[Out] int((c*sin(a + b*x))^m, x)

3.348 $\int \sec^2(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=68

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sec(a + bx)(c \sin(a + bx))^{1+m}}{bc(1 + m)}$$

[Out] hypergeom([3/2, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*sec(b*x+a)*(c*sin(b*x+a))^(1+m)*(cos(b*x+a)^2)^(1/2)/b/c/(1+m)

Rubi [A]

time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2657}

$$\frac{\sqrt{\cos^2(a + bx)} \sec(a + bx)(c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{3}{2}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m + 1)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^2*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[3/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*Sec[a + b*x]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n_*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \sec^2(a + bx)(c \sin(a + bx))^m dx = \frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sec(a + bx)(c \sin(a + bx))^{1+m}}{bc(1 + m)}$$

Mathematica [A]

time = 0.03, size = 63, normalized size = 0.93

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1 + m)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^2*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[3/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int (\sec^2(bx + a)) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^2*(c*sin(b*x+a))^m,x)

[Out] int(sec(b*x+a)^2*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m*sec(b*x + a)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m \sec^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**2*(c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m*sec(a + b*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^2*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^m}{\cos(a + bx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/cos(a + b*x)^2,x)

[Out] int((c*sin(a + b*x))^m/cos(a + b*x)^2, x)

3.349 $\int \sec^4(a + bx)(c \sin(a + bx))^m dx$

Optimal. Leaf size=68

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sec(a + bx)(c \sin(a + bx))^{1+m}}{bc(1 + m)}$$

[Out] hypergeom([5/2, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*sec(b*x+a)*(c*sin(b*x+a))^(1+m)*(cos(b*x+a)^2)^(1/2)/b/c/(1+m)

Rubi [A]

time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2657}

$$\frac{\sqrt{\cos^2(a + bx)} \sec(a + bx)(c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{5}{2}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m + 1)}$$

Antiderivative was successfully verified.

[In] Int[Sec[a + b*x]^4*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[5/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*Sec[a + b*x]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m))

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \sec^4(a + bx)(c \sin(a + bx))^m dx = \frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sec(a + bx)(c \sin(a + bx))^{1+m}}{bc(1 + m)}$$

Mathematica [A]

time = 0.03, size = 63, normalized size = 0.93

$$\frac{\sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{2}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1 + m)}$$

Antiderivative was successfully verified.

[In] Integrate[Sec[a + b*x]^4*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[Cos[a + b*x]^2]*Hypergeometric2F1[5/2, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m))

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int (\sec^4(bx + a)) (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(b*x+a)^4*(c*sin(b*x+a))^m,x)

[Out] int(sec(b*x+a)^4*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a)^4, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral((c*sin(b*x + a))^m*sec(b*x + a)^4, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m \sec^4(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)**4*(c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m*sec(a + b*x)**4, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(b*x+a)^4*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m*sec(b*x + a)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^m}{\cos(a + bx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/cos(a + b*x)^4,x)

[Out] int((c*sin(a + b*x))^m/cos(a + b*x)^4, x)

3.350 $\int (d \cos(a + bx))^{3/2} (c \sin(a + bx))^m dx$

Optimal. Leaf size=75

$$\frac{d\sqrt{d \cos(a + bx)} {}_2F_1\left(-\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)\sqrt[4]{\cos^2(a + bx)}}$$

[Out] d*hypergeom([-1/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)*(d*cos(b*x+a))^(1/2)/b/c/(1+m)/(cos(b*x+a)^2)^(1/4)

Rubi [A]

time = 0.04, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2657}

$$\frac{d\sqrt{d \cos(a + bx)} (c \sin(a + bx))^{m+1} {}_2F_1\left(-\frac{1}{4}, \frac{m+1}{2}, \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)\sqrt[4]{\cos^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Cos[a + b*x])^(3/2)*(c*Sin[a + b*x])^m,x]

[Out] (d*Sqrt[d*Cos[a + b*x]]*Hypergeometric2F1[-1/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m)*(Cos[a + b*x]^2)^(1/4))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int (d \cos(a + bx))^{3/2} (c \sin(a + bx))^m dx = \frac{d\sqrt{d \cos(a + bx)} {}_2F_1\left(-\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m)\sqrt[4]{\cos^2(a + bx)}}$$

Mathematica [A]

time = 0.10, size = 78, normalized size = 1.04

$$\frac{d^2 \cos^2(a + bx)^{3/4} {}_2F_1\left(-\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1+m)\sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^(3/2)*(c*sin[a + b*x])^m,x]

[Out] (d^2*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[-1/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m)*Sqrt[d*cos[a + b*x]])

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int (d \cos (bx + a))^{\frac{3}{2}} (c \sin (bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x)

[Out] int((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^(3/2)*(c*sin(b*x + a))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m*d*cos(b*x + a), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin (a + bx))^m (d \cos (a + bx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(3/2)*(c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m*(d*cos(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^(3/2)*(c*sin(b*x + a))^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + b x))^{3/2} (c \sin(a + b x))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^m,x)

[Out] int((d*cos(a + b*x))^(3/2)*(c*sin(a + b*x))^m, x)

3.351 $\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^m dx$

Optimal. Leaf size=75

$$\frac{d^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m) \sqrt{d \cos(a + bx)}}$$

[Out] d*(cos(b*x+a)^2)^(1/4)*hypergeom([1/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)
*(c*sin(b*x+a))^(1+m)/b/c/(1+m)/(d*cos(b*x+a))^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2657}

$$\frac{d^4 \sqrt{\cos^2(a + bx)} (c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1) \sqrt{d \cos(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^m,x]

[Out] (d*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m)*Sqrt[d*Cos[a + b*x]])

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^m dx = \frac{d^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^{1+m}}{bc(1+m) \sqrt{d \cos(a + bx)}}$$

Mathematica [A]

time = 0.05, size = 75, normalized size = 1.00

$$\frac{\sqrt{d \cos(a + bx)} \sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (c \sin(a + bx))^m \tan(a + bx)}{b(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Cos[a + b*x]]*(c*Sin[a + b*x])^m,x]

[Out] (Sqrt[d*Cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \sqrt{d \cos(bx + a)} (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x)

[Out] int((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m \sqrt{d \cos(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**(1/2)*(c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m*sqrt(d*cos(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x, algorithm="giac")``[Out] integrate(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^m,x)``[Out] int((d*cos(a + b*x))^(1/2)*(c*sin(a + b*x))^m, x)`

$$3.352 \quad \int \frac{(c \sin(a+bx))^m}{\sqrt{d \cos(a+bx)}} dx$$

Optimal. Leaf size=75

$$\frac{d \cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bc(1+m)(d \cos(a+bx))^{3/2}}$$

[Out] d*(cos(b*x+a)^2)^(3/4)*hypergeom([3/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)/(d*cos(b*x+a))^(3/2)

Rubi [A]

time = 0.03, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2657}

$$\frac{d \cos^2(a+bx)^{3/4} (c \sin(a+bx))^{m+1} {}_2F_1\left(\frac{3}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a+bx)\right)}{bc(m+1)(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^m/Sqrt[d*Cos[a + b*x]],x]

[Out] (d*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m)*(d*Cos[a + b*x])^(3/2))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n_*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{(c \sin(a+bx))^m}{\sqrt{d \cos(a+bx)}} dx = \frac{d \cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bc(1+m)(d \cos(a+bx))^{3/2}}$$

Mathematica [A]

time = 0.05, size = 75, normalized size = 1.00

$$\frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^m \tan(a+bx)}{b(1+m)\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^m/Sqrt[d*Cos[a + b*x]],x]

[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*(1 + m)*Sqrt[d*Cos[a + b*x]])

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(bx + a))^m}{\sqrt{d \cos(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^m/(d*cos(b*x+a))^(1/2),x)

[Out] int((c*sin(b*x+a))^m/(d*cos(b*x+a))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m/sqrt(d*cos(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m/(d*cos(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(a + bx))^m}{\sqrt{d \cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**m/(d*cos(b*x+a))**(1/2),x)

[Out] Integral((c*sin(a + b*x))**m/sqrt(d*cos(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m/sqrt(d*cos(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + b x))^m}{\sqrt{d \cos(a + b x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/(d*cos(a + b*x))^(1/2),x)

[Out] int((c*sin(a + b*x))^m/(d*cos(a + b*x))^(1/2), x)

$$3.353 \quad \int \frac{(c \sin(a+bx))^m}{(d \cos(a+bx))^{3/2}} dx$$

Optimal. Leaf size=77

$$\frac{\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{5}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bcd(1+m) \sqrt{d \cos(a+bx)}}$$

[Out] $(\cos(b*x+a)^2)^{(1/4)} * \text{hypergeom}([5/4, 1/2+1/2*m], [3/2+1/2*m], \sin(b*x+a)^2) * (c*\sin(b*x+a))^{(1+m)}/b/c/d/(1+m)/(d*\cos(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2657}

$$\frac{\sqrt[4]{\cos^2(a+bx)} (c \sin(a+bx))^{m+1} {}_2F_1\left(\frac{5}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a+bx)\right)}{bcd(m+1) \sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c*\text{Sin}[a + b*x])^m/(d*\text{Cos}[a + b*x])^{(3/2)}, x]$

[Out] $((\text{Cos}[a + b*x]^2)^{(1/4)} * \text{Hypergeometric2F1}[5/4, (1 + m)/2, (3 + m)/2, \text{Sin}[a + b*x]^2] * (c*\text{Sin}[a + b*x])^{(1 + m)}) / (b*c*d*(1 + m)*\text{Sqrt}[d*\text{Cos}[a + b*x]])$

Rule 2657

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(b_.)^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_)])^{(m_)}], x_Symbol] :> \text{Simp}[b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Cos}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}*((a*\text{Sin}[e + f*x])^{(m + 1)})/(a*f*(m + 1)*(\text{Cos}[e + f*x]^2)^{\text{FracPart}[(n - 1)/2]}) * \text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Sin}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x]$

Rubi steps

$$\int \frac{(c \sin(a+bx))^m}{(d \cos(a+bx))^{3/2}} dx = \frac{\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{5}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bcd(1+m) \sqrt{d \cos(a+bx)}}$$

Mathematica [A]

time = 0.07, size = 78, normalized size = 1.01

$$\frac{\sqrt{d \cos(a+bx)} \sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{5}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^m \tan(a+bx)}{bd^2(1+m)}$$

Antiderivative was successfully verified.

[In] Integrate[(c*SIN[a + b*x])^m/(d*cos[a + b*x])^(3/2),x]

[Out] (Sqrt[d*cos[a + b*x]]*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[5/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*SIN[a + b*x])^m*Tan[a + b*x])/(b*d^2*(1 + m))

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(bx + a))^m}{(d \cos(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^m/(d*cos(b*x+a))^(3/2),x)

[Out] int((c*sin(b*x+a))^m/(d*cos(b*x+a))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m/(d*cos(b*x + a))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m/(d^2*cos(b*x + a)^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(a + bx))^m}{(d \cos(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a)**m/(d*cos(b*x+a))**(3/2),x)

[Out] Integral((c*sin(a + b*x))**m/(d*cos(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m/(d*cos(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + b x))^m}{(d \cos(a + b x))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/(d*cos(a + b*x))^(3/2),x)

[Out] int((c*sin(a + b*x))^m/(d*cos(a + b*x))^(3/2), x)

$$3.354 \quad \int \frac{(c \sin(a+bx))^m}{(d \cos(a+bx))^{5/2}} dx$$

Optimal. Leaf size=77

$$\frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bcd(1+m)(d \cos(a+bx))^{3/2}}$$

[Out] (cos(b*x+a)^2)^(3/4)*hypergeom([7/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/d/(1+m)/(d*cos(b*x+a))^(3/2)

Rubi [A]

time = 0.04, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2657}

$$\frac{\cos^2(a+bx)^{3/4} (c \sin(a+bx))^{m+1} {}_2F_1\left(\frac{7}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a+bx)\right)}{bcd(m+1)(d \cos(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^m/(d*Cos[a + b*x])^(5/2),x]

[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[7/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*d*(1 + m)*(d*Cos[a + b*x])^(3/2))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\int \frac{(c \sin(a+bx))^m}{(d \cos(a+bx))^{5/2}} dx = \frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bcd(1+m)(d \cos(a+bx))^{3/2}}$$

Mathematica [A]

time = 0.06, size = 78, normalized size = 1.01

$$\frac{\cos^2(a+bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^m \tan(a+bx)}{bd^2(1+m)\sqrt{d \cos(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^m/(d*Cos[a + b*x])^(5/2), x]

[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[7/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^m*Tan[a + b*x])/(b*d^2*(1 + m)*Sqrt[d*Cos[a + b*x]])

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(bx + a))^m}{(d \cos(bx + a))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^m/(d*cos(b*x+a))^(5/2), x)

[Out] int((c*sin(b*x+a))^m/(d*cos(b*x+a))^(5/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(5/2), x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m/(d*cos(b*x + a))^(5/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(5/2), x, algorithm="fricas")

[Out] integral(sqrt(d*cos(b*x + a))*(c*sin(b*x + a))^m/(d^3*cos(b*x + a)^3), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(5/2), x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*cos(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m/(d*cos(b*x + a))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + b x))^m}{(d \cos(a + b x))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/(d*cos(a + b*x))^(5/2),x)

[Out] int((c*sin(a + b*x))^m/(d*cos(a + b*x))^(5/2), x)

3.355 $\int (d \cos(a + bx))^n \sin^5(a + bx) dx$

Optimal. Leaf size=76

$$-\frac{(d \cos(a + bx))^{1+n}}{bd(1+n)} + \frac{2(d \cos(a + bx))^{3+n}}{bd^3(3+n)} - \frac{(d \cos(a + bx))^{5+n}}{bd^5(5+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}/b/d/(1+n)+2*(d*\cos(b*x+a))^{(3+n)}/b/d^3/(3+n)-(d*\cos(b*x+a))^{(5+n)}/b/d^5/(5+n)$

Rubi [A]

time = 0.05, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 276}

$$-\frac{(d \cos(a + bx))^{n+5}}{bd^5(n+5)} + \frac{2(d \cos(a + bx))^{n+3}}{bd^3(n+3)} - \frac{(d \cos(a + bx))^{n+1}}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Sin}[a + b*x]^5, x]$

[Out] $-\left(\frac{d*\text{Cos}[a + b*x]^{(1+n)}}{b*d*(1+n)}\right) + \frac{2*(d*\text{Cos}[a + b*x]^{(3+n)})}{b*d^3*(3+n)} - \frac{d*\text{Cos}[a + b*x]^{(5+n)}}{b*d^5*(5+n)}$

Rule 276

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2645

$\text{Int}[(\cos[(e_*) + (f_*)(x_*)])*(a_*)^{(m_*)}*\sin[(e_*) + (f_*)(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{(n-1)/2}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m, x\} \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^n \sin^5(a + bx) dx &= -\frac{\text{Subst}\left(\int x^n \left(1 - \frac{x^2}{a^2}\right)^2 dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(x^n - \frac{2x^{2+n}}{a^2} + \frac{x^{4+n}}{a^4}\right) dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{1+n}}{bd(1+n)} + \frac{2(d \cos(a + bx))^{3+n}}{bd^3(3+n)} - \frac{(d \cos(a + bx))^{5+n}}{bd^5(5+n)} \end{aligned}$$

Mathematica [A]

time = 0.22, size = 83, normalized size = 1.09

$$\frac{\cos(a+bx)(d\cos(a+bx))^n(89+28n+3n^2-4(7+8n+n^2)\cos(2(a+bx))+(3+4n+n^2)\cos(4(a+bx)))}{8b(1+n)(3+n)(5+n)}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^n*Sin[a + b*x]^5,x]`

```
[Out] -1/8*(Cos[a + b*x]*(d*Cos[a + b*x])^n*(89 + 28*n + 3*n^2 - 4*(7 + 8*n + n^2)
)*Cos[2*(a + b*x)] + (3 + 4*n + n^2)*Cos[4*(a + b*x)]))/(b*(1 + n)*(3 + n)*
(5 + n))
```

Maple [F]

time = 0.14, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (\sin^5(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*cos(b*x+a))^n*sin(b*x+a)^5,x)``[Out] int((d*cos(b*x+a))^n*sin(b*x+a)^5,x)`**Maxima [A]**

time = 0.29, size = 78, normalized size = 1.03

$$-\frac{\frac{d^n \cos(bx+a)^n \cos(bx+a)^5}{n+5} - \frac{2 d^n \cos(bx+a)^n \cos(bx+a)^3}{n+3} + \frac{(d \cos(bx+a))^{n+1}}{d(n+1)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^5,x, algorithm="maxima")`

```
[Out] -(d^n*cos(b*x + a)^n*cos(b*x + a)^5/(n + 5) - 2*d^n*cos(b*x + a)^n*cos(b*x
+ a)^3/(n + 3) + (d*cos(b*x + a))^(n + 1)/(d*(n + 1)))/b
```

Fricas [A]

time = 0.43, size = 84, normalized size = 1.11

$$\frac{((n^2 + 4n + 3)\cos(bx + a)^5 - 2(n^2 + 6n + 5)\cos(bx + a)^3 + (n^2 + 8n + 15)\cos(bx + a))(d\cos(bx + a))^n}{bn^3 + 9bn^2 + 23bn + 15b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^5,x, algorithm="fricas")`

```
[Out] -((n^2 + 4*n + 3)*cos(b*x + a)^5 - 2*(n^2 + 6*n + 5)*cos(b*x + a)^3 + (n^2
+ 8*n + 15)*cos(b*x + a))*(d*cos(b*x + a))^n/(b*n^3 + 9*b*n^2 + 23*b*n + 15
*b)
```

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 2451 vs. $2(60) = 120$.

time = 5.87, size = 2451, normalized size = 32.25

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*sin(b*x+a)**5,x)

[Out] Piecewise((x*(d*cos(a))**n*sin(a)**5, Eq(b, 0)), ((-log(cos(a + b*x))/b + sin(a + b*x)**4/(4*b*cos(a + b*x)**4) - sin(a + b*x)**2/(2*b*cos(a + b*x)**2))/d**5, Eq(n, -5)), ((2*log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) - 4*log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) + 2*log(tan(a/2 + b*x/2) - 1)/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) + 2*log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) - 4*log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) + 2*log(tan(a/2 + b*x/2) + 1)/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) - 2*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) + 4*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) - 2*log(tan(a/2 + b*x/2)**2 + 1)/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) + 4*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b) + 4*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 - 2*b*tan(a/2 + b*x/2)**4 + b))/d**3, Eq(n, -3)), ((-log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 6*log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2) - 1)/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 6*log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 4*log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2) + 1)/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2

```

+ b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2)**2 + 1)*
tan(a/2 + b*x/2)**8/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*
tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 4*log(tan(a/2 + b*x/2)
**2 + 1)*tan(a/2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)*
*6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 6*log(tan(a/2
+ b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2
+ b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) + 4*lo
g(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4*b
*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 +
b) + log(tan(a/2 + b*x/2)**2 + 1)/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*
x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 2*tan(a/
2 + b*x/2)**6/(b*tan(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/
2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2 + b) - 8*tan(a/2 + b*x/2)**4/(b*tan
(a/2 + b*x/2)**8 + 4*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*
tan(a/2 + b*x/2)**2 + b) - 2*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**8 + 4
*b*tan(a/2 + b*x/2)**6 + 6*b*tan(a/2 + b*x/2)**4 + 4*b*tan(a/2 + b*x/2)**2
+ b))/d, Eq(n, -1)), (-n**2*(d*cos(a + b*x))**n*sin(a + b*x)**4*cos(a + b*x
)/(b*n**3 + 9*b*n**2 + 23*b*n + 15*b) - 8*n*(d*cos(a + b*x))**n*sin(a + b*x
)**4*cos(a + b*x)/(b*n**3 + 9*b*n**2 + 23*b*n + 15*b) - 4*n*(d*cos(a + b*x)
)**n*sin(a + b*x)**2*cos(a + b*x)**3/(b*n**3 + 9*b*n**2 + 23*b*n + 15*b) -
15*(d*cos(a + b*x))**n*sin(a + b*x)**4*cos(a + b*x)/(b*n**3 + 9*b*n**2 + 23
*b*n + 15*b) - 20*(d*cos(a + b*x))**n*sin(a + b*x)**2*cos(a + b*x)**3/(b*n*
*3 + 9*b*n**2 + 23*b*n + 15*b) - 8*(d*cos(a + b*x))**n*cos(a + b*x)**5/(b*n
**3 + 9*b*n**2 + 23*b*n + 15*b), True))

```

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 249 vs. 2(76) = 152.

time = 3.04, size = 249, normalized size = 3.28

$\frac{(d \cos(bx+a))^n d^n \cos^2(bx+a)^2 + 4(d \cos(bx+a))^n d^n \cos(bx+a) - 2(d \cos(bx+a))^n d^n \cos^2(bx+a)^2 + 3(d \cos(bx+a))^n d^n \cos(bx+a) - 12(d \cos(bx+a))^n d^n \cos(bx+a)^2 + (d \cos(bx+a))^n d^n \cos^2(bx+a) - 10(d \cos(bx+a))^n d^n \cos(bx+a) + 8(d \cos(bx+a))^n d^n \cos(bx+a) + 15(d \cos(bx+a))^n d^n \cos(bx+a)}{(d^n + 9d^{n-1} + 23d^{n-2} + 15d^{n-3})d}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^5,x, algorithm="giac")

[Out] $-(d \cos(bx+a))^n d^{5n} \cos^2(bx+a)^5 + 4(d \cos(bx+a))^n d^{5n} \cos(bx+a)^5 - 2(d \cos(bx+a))^n d^{5n} \cos^2(bx+a)^3 + 3(d \cos(bx+a))^n d^{5n} \cos(bx+a)^5 - 12(d \cos(bx+a))^n d^{5n} \cos(bx+a)^3 + (d \cos(bx+a))^n d^{5n} \cos^2(bx+a) - 10(d \cos(bx+a))^n d^{5n} \cos(bx+a)^3 + 8(d \cos(bx+a))^n d^{5n} \cos(bx+a) + 15(d \cos(bx+a))^n d^{5n} \cos(bx+a) / ((d^4 n^3 + 9d^4 n^2 + 23d^4 n + 15d^4) b d)$

Mupad [B]

time = 1.54, size = 132, normalized size = 1.74

$\frac{(d \cos(a+bx))^n (150 \cos(a+bx) - 25 \cos(3a+3bx) + 3 \cos(5a+5bx) + 24n \cos(a+bx) - 28n \cos(3a+3bx) + 4n \cos(5a+5bx) + 2n^2 \cos(a+bx) - 3n^2 \cos(3a+3bx) + n^2 \cos(5a+5bx))}{16b(n^3 + 9n^2 + 23n + 15)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(a + b*x)^5*(d*cos(a + b*x))^n,x)`

[Out] $-\left((d \cos(a + b*x))^n (150 \cos(a + b*x) - 25 \cos(3*a + 3*b*x) + 3 \cos(5*a + 5*b*x) + 24*n \cos(a + b*x) - 28*n \cos(3*a + 3*b*x) + 4*n \cos(5*a + 5*b*x) + 2*n^2 \cos(a + b*x) - 3*n^2 \cos(3*a + 3*b*x) + n^2 \cos(5*a + 5*b*x))\right) / (16*b * (23*n + 9*n^2 + n^3 + 15))$

3.356 $\int (d \cos(a + bx))^n \sin^3(a + bx) dx$

Optimal. Leaf size=50

$$-\frac{(d \cos(a + bx))^{1+n}}{bd(1+n)} + \frac{(d \cos(a + bx))^{3+n}}{bd^3(3+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}/b/d/(1+n)+(d*\cos(b*x+a))^{(3+n)}/b/d^3/(3+n)$

Rubi [A]

time = 0.04, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 14}

$$\frac{(d \cos(a + bx))^{n+3}}{bd^3(n+3)} - \frac{(d \cos(a + bx))^{n+1}}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Sin}[a + b*x]^3,x]$

[Out] $-\left(\frac{d*\text{Cos}[a + b*x]^{(1+n)}}{b*d*(1+n)}\right) + \frac{d*\text{Cos}[a + b*x]^{(3+n)}}{b*d^3*(3+n)}$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^{m*u}, x], x] /; \text{FreeQ}[\{c, m\}, x] \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_))] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{InverseFunctionQ}[v]$

Rule 2645

$\text{Int}[(\cos[(e_)] + (f_)*(x_)]*(a_))^{(m_)}*\sin[(e_)] + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{((n-1)/2)}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^n \sin^3(a + bx) dx &= -\frac{\text{Subst}\left(\int x^n \left(1 - \frac{x^2}{d^2}\right) dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{\text{Subst}\left(\int \left(x^n - \frac{x^{2+n}}{d^2}\right) dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{1+n}}{bd(1+n)} + \frac{(d \cos(a + bx))^{3+n}}{bd^3(3+n)} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 50, normalized size = 1.00

$$\frac{\cos(a + bx)(d \cos(a + bx))^n(-5 - n + (1 + n) \cos(2(a + bx)))}{2b(1 + n)(3 + n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^n*Sin[a + b*x]^3,x]**[Out]** (Cos[a + b*x]*(d*Cos[a + b*x])^n*(-5 - n + (1 + n)*Cos[2*(a + b*x)]))/(2*b*(1 + n)*(3 + n))**Maple [C]** Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 1.17, size = 1076, normalized size = 21.52

method	result	size
risch	Expression too large to display	1076

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*sin(b*x+a)^3,x,method=_RETURNVERBOSE)

[Out] $-1/8*(\exp(2*I*(b*x+a))+1)^n*(1/2)^n*d^n*\exp(I*(b*x+a))^{(-n)/(3+n)/(1+n)}/b*(n+9)*\exp(-1/2*I*(\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))- \text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*d*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*\exp(-I*(b*x+a))))-\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))^2-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^2*\text{csgn}(I*\exp(-I*(b*x+a)))+\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^3-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*d*\cos(b*x+a))^3+2*b*x+2*a)-1/8*(\exp(2*I*(b*x+a))+1)^n*(1/2)^n*d^n*\exp(I*(b*x+a))^{(-n)/(3+n)/(1+n)}/b*(n+9)*\exp(1/2*I*(-\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))+\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*d*\cos(b*x+a))^2-\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*\exp(-I*(b*x+a)))+\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^2*\text{csgn}(I*\exp(-I*(b*x+a)))-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^3+\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))^2-\text{Pi}*n*\text{csgn}(I*d*\cos(b*x+a))^3+2*b*x+2*a))+1/8/(b*n+3*b)*\exp(I*(b*x+a))^{(-n)*d^n*(1/2)^n*(\exp(2*I*(b*x+a))+1)^n*\exp(-1/2*I*(\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))- \text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*d*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*\exp(-I*(b*x+a)))-\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))^2-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^2*\text{csgn}(I*\exp(-I*(b*x+a)))+\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^3-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*d*\cos(b*x+a))^3+6*b*x+6*a))+1/8*(\exp(2*I*(b*x+a))+1)^n*(1/2)^n*d^n*\exp(I*(b*x+a))^{(-n)/(b*n+3*b)*\exp(1/2*I*(-\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))+\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*d*\cos(b*x+a))^2-\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*\exp(-I*(b*x+a)))+\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^2*\text{csgn}(I*\exp(-I*(b*x+a)))-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^3-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*d*\cos(b*x+a))^3+6*b*x+6*a))+1/8*(\exp(2*I*(b*x+a))+1)^n*(1/2)^n*d^n*\exp(I*(b*x+a))^{(-n)/(b*n+3*b)*\exp(1/2*I*(-\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))+\text{Pi}*n*\text{csgn}(I*d)*\text{csgn}(I*d*\cos(b*x+a))^2-\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*\exp(-I*(b*x+a)))+\text{Pi}*n*\text{csgn}(I*(\exp(2*I*(b*x+a))+1))*\text{csgn}(I*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^2*\text{csgn}(I*\exp(-I*(b*x+a)))-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))^3-\text{Pi}*n*\text{csgn}(I*\cos(b*x+a))*\text{csgn}(I*d*\cos(b*x+a))^2+\text{Pi}*n*\text{csgn}(I*d*\cos(b*x+a))^3+6*b*x+6*a))$

$(b*x+a)^{3+Pi*n}*csgn(I*cos(b*x+a))*csgn(I*d*cos(b*x+a))^{2-Pi*n}*csgn(I*d*cos(b*x+a))^{3+6*b*x+6*a)}$

Maxima [A]

time = 0.29, size = 52, normalized size = 1.04

$$\frac{\frac{d^n \cos(bx+a)^n \cos(bx+a)^3}{n+3} - \frac{(d \cos(bx+a))^{n+1}}{d(n+1)}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^3,x, algorithm="maxima")

[Out] (d^n*cos(b*x + a)^n*cos(b*x + a)^3/(n + 3) - (d*cos(b*x + a))^(n + 1)/(d*(n + 1)))/b

Fricas [A]

time = 0.38, size = 50, normalized size = 1.00

$$\frac{((n + 1) \cos(bx + a)^3 - (n + 3) \cos(bx + a))(d \cos(bx + a))^n}{bn^2 + 4bn + 3b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^3,x, algorithm="fricas")

[Out] ((n + 1)*cos(b*x + a)^3 - (n + 3)*cos(b*x + a))*(d*cos(b*x + a))^n/(b*n^2 + 4*b*n + 3*b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 688 vs. 2(37) = 74.

time = 2.67, size = 688, normalized size = 13.76

$$\frac{x(d \cos(a))^n \sin^3(a)}{\dots}$$

for b = 0
for n = -3
for n = -1
otherwise

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)**3,x)

[Out] Piecewise((x*(d*cos(a))^n*sin(a)**3, Eq(b, 0)), ((log(cos(a + b*x))/b + sin(a + b*x)**2/(2*b*cos(a + b*x)**2))/d**3, Eq(n, -3)), ((-log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - 2*log(tan(a/2 + b*x/2) - 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - 2*log(tan(a/2 + b*x/2) + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - log(tan(a/2 + b*x/2) + 1)/(b*tan(a/2 + b*x/2)**4

+ 2*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**4/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) + 2*log(tan(a/2 + b*x/2)**2 + 1)*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) + log(tan(a/2 + b*x/2)**2 + 1)/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b) - 2*tan(a/2 + b*x/2)**2/(b*tan(a/2 + b*x/2)**4 + 2*b*tan(a/2 + b*x/2)**2 + b))/d, Eq(n, -1)), (-n*(d*cos(a + b*x))**n*sin(a + b*x)**2*cos(a + b*x)/(b*n**2 + 4*b*n + 3*b) - 3*(d*cos(a + b*x))**n*sin(a + b*x)**2*cos(a + b*x)/(b*n**2 + 4*b*n + 3*b) - 2*(d*cos(a + b*x))**n*cos(a + b*x)**3/(b*n**2 + 4*b*n + 3*b), True))

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 117 vs. 2(50) = 100.

time = 2.69, size = 117, normalized size = 2.34

$$\frac{(d \cos(bx + a))^n d^3 n \cos(bx + a)^3 + (d \cos(bx + a))^n d^3 \cos(bx + a)^3 - (d \cos(bx + a))^n d^3 n \cos(bx + a) - 3(d \cos(bx + a))^n d^3 \cos(bx + a)}{(d^2 n^2 + 4 d^2 n + 3 d^2) b d}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^3,x, algorithm="giac")

[Out] ((d*cos(b*x + a))^n*d^3*n*cos(b*x + a)^3 + (d*cos(b*x + a))^n*d^3*cos(b*x + a)^3 - (d*cos(b*x + a))^n*d^3*n*cos(b*x + a) - 3*(d*cos(b*x + a))^n*d^3*cos(b*x + a))/((d^2*n^2 + 4*d^2*n + 3*d^2)*b*d)

Mupad [B]

time = 0.91, size = 65, normalized size = 1.30

$$\frac{(d \cos(a + bx))^n (9 \cos(a + bx) - \cos(3a + 3bx) + n \cos(a + bx) - n \cos(3a + 3bx))}{4 b (n^2 + 4 n + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^3*(d*cos(a + b*x))^n,x)

[Out] -((d*cos(a + b*x))^n*(9*cos(a + b*x) - cos(3*a + 3*b*x) + n*cos(a + b*x) - n*cos(3*a + 3*b*x)))/(4*b*(4*n + n^2 + 3))

3.357 $\int (d \cos(a + bx))^n \sin(a + bx) dx$

Optimal. Leaf size=25

$$\frac{(d \cos(a + bx))^{1+n}}{bd(1+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}/b/d/(1+n)$

Rubi [A]

time = 0.02, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 30}

$$\frac{(d \cos(a + bx))^{n+1}}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Sin}[a + b*x], x]$

[Out] $-\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)} / (b*d*(1+n))\right)$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \text{ :> Simp}[x^{(m+1)}/(m+1), x] \text{ /; FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2645

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*\sin[(e_.) + (f_.)*(x_.)]^{(n_.)}, x_Symbol] \text{ :> Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{((n-1)/2)}, x], x, a*\text{Cos}[e + f*x]], x] \text{ /; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^n \sin(a + bx) dx &= -\frac{\text{Subst}\left(\int x^n dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{1+n}}{bd(1+n)} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 26, normalized size = 1.04

$$-\frac{\cos(a + bx)(d \cos(a + bx))^n}{b(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*sin[a + b*x],x]

[Out] -((Cos[a + b*x]*(d*cos[a + b*x])^n)/(b*(1 + n)))

Maple [A]

time = 0.93, size = 26, normalized size = 1.04

method	result	size
derivativedivides	$-\frac{(d \cos(bx+a))^{1+n}}{bd(1+n)}$	26
default	$-\frac{(d \cos(bx+a))^{1+n}}{bd(1+n)}$	26
norman	$\frac{(\tan^2(\frac{bx}{2} + \frac{a}{2}))^n e^{n \ln\left(\frac{d(1 - \tan^2(\frac{bx}{2} + \frac{a}{2}))}{1 + \tan^2(\frac{bx}{2} + \frac{a}{2})}\right)}}{b(1+n)} - \frac{e^{n \ln\left(\frac{d(1 - \tan^2(\frac{bx}{2} + \frac{a}{2}))}{1 + \tan^2(\frac{bx}{2} + \frac{a}{2})}\right)}}{b(1+n)}$	120
risch	Expression too large to display	785

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*sin(b*x+a),x,method=_RETURNVERBOSE)

[Out] -(d*cos(b*x+a))^(1+n)/b/d/(1+n)

Maxima [A]

time = 0.30, size = 25, normalized size = 1.00

$$-\frac{(d \cos(bx + a))^{n+1}}{bd(n + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a),x, algorithm="maxima")

[Out] -(d*cos(b*x + a))^(n + 1)/(b*d*(n + 1))

Fricas [A]

time = 0.44, size = 25, normalized size = 1.00

$$-\frac{(d \cos(bx + a))^n \cos(bx + a)}{bn + b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a),x, algorithm="fricas")

[Out] -(d*cos(b*x + a))^n*cos(b*x + a)/(b*n + b)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 60 vs. $2(19) = 38$.

time = 0.83, size = 60, normalized size = 2.40

$$\begin{cases} \frac{x \sin(a)}{d \cos(a)} & \text{for } b = 0 \wedge n = -1 \\ x(d \cos(a))^n \sin(a) & \text{for } b = 0 \\ -\frac{\log(\cos(a+bx))}{bd} & \text{for } n = -1 \\ -\frac{(d \cos(a+bx))^n \cos(a+bx)}{bn+b} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*sin(b*x+a),x)

[Out] Piecewise((x*sin(a)/(d*cos(a)), Eq(b, 0) & Eq(n, -1)), (x*(d*cos(a))**n*sin(a), Eq(b, 0)), (-log(cos(a + b*x))/(b*d), Eq(n, -1)), (-(d*cos(a + b*x))**n*cos(a + b*x)/(b*n + b), True))

Giac [A]

time = 3.77, size = 25, normalized size = 1.00

$$-\frac{(d \cos(bx + a))^{n+1}}{bd(n + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a),x, algorithm="giac")

[Out] -(d*cos(b*x + a))^(n + 1)/(b*d*(n + 1))

Mupad [B]

time = 0.18, size = 26, normalized size = 1.04

$$-\frac{\cos(a + bx) (d \cos(a + bx))^n}{b (n + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)*(d*cos(a + b*x))^n,x)

[Out] -(cos(a + b*x)*(d*cos(a + b*x))^n)/(b*(n + 1))

3.358 $\int (d \cos(a + bx))^n \csc(a + bx) dx$

Optimal. Leaf size=49

$$-\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)}{bd(1+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([1, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)/b/d/(1+n)$

Rubi [A]

time = 0.04, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2645, 371}

$$-\frac{(d \cos(a + bx))^{n+1} {}_2F_1\left(1, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Csc}[a + b*x], x]$

[Out] $-\left(\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)}*\text{Hypergeometric2F1}\left[1, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2\right]\right)/(b*d*(1+n))\right)$

Rule 371

$\text{Int}[(c_*)*(x_*)^{(m_*)}*((a_*) + (b_*)*(x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n + 1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt Q[p, 0] || GtQ[a, 0])

Rule 2645

$\text{Int}[(\cos[(e_*) + (f_*)*(x_*)]*(a_*)^{(m_*)}*\sin[(e_*) + (f_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1 - x^2/a^2)^{(n-1)/2}, x], x, a*\text{Cos}[e + f*x]], x] /;$ FreeQ[{a, e, f, m}, x] && IntegerQ[(n-1)/2] && !(IntegerQ[(m-1)/2] && GtQ[m, 0] && LeQ[m, n])

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^n \csc(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^n}{1-\frac{x^2}{d^2}} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)}{bd(1+n)} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 52, normalized size = 1.06

$$\frac{\cos(a + bx)(d \cos(a + bx))^n {}_2F_1\left(1, \frac{1+n}{2}; 1 + \frac{1+n}{2}; \cos^2(a + bx)\right)}{b(1+n)}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Cos[a + b*x])^n*Csc[a + b*x],x]``[Out] -((Cos[a + b*x]*(d*Cos[a + b*x])^n*Hypergeometric2F1[1, (1 + n)/2, 1 + (1 + n)/2, Cos[a + b*x]^2])/(b*(1 + n)))`**Maple [F]**

time = 0.07, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n \csc(bx + a) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*cos(b*x+a))^n*csc(b*x+a),x)``[Out] int((d*cos(b*x+a))^n*csc(b*x+a),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^n*csc(b*x+a),x, algorithm="maxima")``[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((d*cos(b*x+a))^n*csc(b*x+a),x, algorithm="fricas")``[Out] integral((d*cos(b*x + a))^n*csc(b*x + a), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(a + bx))^n \csc(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))**n*csc(b*x+a),x)`

[Out] `Integral((d*cos(a + b*x))**n*csc(a + b*x), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^n*csc(b*x+a),x, algorithm="giac")`

[Out] `integrate((d*cos(b*x + a))^n*csc(b*x + a), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(d \cos(a + bx))^n}{\sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(a + b*x))^n/sin(a + b*x),x)`

[Out] `int((d*cos(a + b*x))^n/sin(a + b*x), x)`

3.359 $\int (d \cos(a + bx))^n \csc^3(a + bx) dx$

Optimal. Leaf size=49

$$-\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(2, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)}{bd(1+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)/b/d/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 371}

$$-\frac{(d \cos(a + bx))^{n+1} {}_2F_1\left(2, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Csc}[a + b*x]^3, x]$

[Out] $-\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)}*\text{Hypergeometric2F1}\left[2, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2\right]\right)/(b*d*(1+n))$

Rule 371

$\text{Int}[(c*(x_))^{(m_)}*((a_) + (b_)*(x_)^{(n_}))^{(p_)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1)))*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 2645

$\text{Int}[(\cos[(e_) + (f_)*(x_)]*(a_))^{(m_)}*\sin[(e_) + (f_)*(x_)]^{(n_)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1-x^2/a^2)^{((n-1)/2)}, x], x, a*\text{Cos}[e+f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^n \csc^3(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^n}{(1-x^2)^2} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(2, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)}{bd(1+n)} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 154 vs. 2(49) = 98.

time = 1.85, size = 154, normalized size = 3.14

$$\frac{2^{-3-n} \cos(a+bx) (d \cos(a+bx))^n \left({}_2F_1(1, 1+n; 2+n; \cos(a+bx)) + 2^{1+n} {}_2F_1(2, 1+n; 2+n; \cos(a+bx)) + ({}_2F_1(n, 1+n; 2+n; \frac{1}{2} \cos(a+bx) \sec^2(\frac{1}{2}(a+bx))) + {}_2F_1(1+n, 1+n; 2+n; \frac{1}{2} \cos(a+bx) \sec^2(\frac{1}{2}(a+bx))) \sec^2(\frac{1}{2}(a+bx))^{1+n} \right)}{b(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*csc[a + b*x]^3,x]

[Out] -((2^(-3 - n)*Cos[a + b*x]*(d*cos[a + b*x])^n*(2^(1 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, Cos[a + b*x]] + 2^(1 + n)*Hypergeometric2F1[2, 1 + n, 2 + n, Cos[a + b*x]] + (Hypergeometric2F1[n, 1 + n, 2 + n, (Cos[a + b*x]*Sec[(a + b*x)/2]^2)/2] + Hypergeometric2F1[1 + n, 1 + n, 2 + n, (Cos[a + b*x]*Sec[(a + b*x)/2]^2)/2])*(Sec[(a + b*x)/2]^2)^(1 + n)))/(b*(1 + n))

Maple [F]

time = 0.07, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (\csc^3(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*csc(b*x+a)^3,x)

[Out] int((d*cos(b*x+a))^n*csc(b*x+a)^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^3,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^3,x, algorithm="fricas")

[Out] integral((d*cos(b*x + a))^n*csc(b*x + a)^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(a + bx))^n \csc^3(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*csc(b*x+a)**3,x)

[Out] Integral((d*cos(a + b*x))**n*csc(a + b*x)**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^3,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(d \cos(a + bx))^n}{\sin(a + bx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n/sin(a + b*x)^3,x)

[Out] int((d*cos(a + b*x))^n/sin(a + b*x)^3, x)

3.360 $\int (d \cos(a + bx))^n \csc^5(a + bx) dx$

Optimal. Leaf size=49

$$\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(3, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)}{bd(1+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([3, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)/b/d/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2645, 371}

$$\frac{(d \cos(a + bx))^{n+1} {}_2F_1\left(3, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Csc}[a + b*x]^5, x]$

[Out] $-\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)}*\text{Hypergeometric2F1}\left[3, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2\right]\right)/(b*d*(1+n))$

Rule 371

$\text{Int}[\left((c_*)*(x_*)\right)^{(m_*)}*\left((a_*) + (b_*)*(x_*)^{(n_*)}\right)^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[a^p * \left(\frac{c*x}{c*(m+1)}\right)^{(m+1)}*\text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; \text{FreeQ}\{a, b, c, m, n, p\}, x \ \&\& \ !\text{IGtQ}[p, 0] \ \&\& \ (\text{ILtQ}[p, 0] \ || \ \text{GtQ}[a, 0])$

Rule 2645

$\text{Int}[(\cos[(e_*) + (f_*)*(x_*)]*(a_*)\right)^{(m_*)}*\sin[(e_*) + (f_*)*(x_*)]^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[-(a*f)^{-1}, \text{Subst}[\text{Int}[x^m*(1-x^2/a^2)^{(n-1)/2}, x], x, a*\text{Cos}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n-1)/2] \ \&\& \ !(\text{IntegerQ}[(m-1)/2] \ \&\& \ \text{GtQ}[m, 0] \ \&\& \ \text{LeQ}[m, n])$

Rubi steps

$$\begin{aligned} \int (d \cos(a + bx))^n \csc^5(a + bx) dx &= -\frac{\text{Subst}\left(\int \frac{x^n}{(1-\frac{x^2}{d^2})^3} dx, x, d \cos(a + bx)\right)}{bd} \\ &= -\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(3, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)}{bd(1+n)} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 244 vs. 2(49) = 98.

time = 2.85, size = 244, normalized size = 4.98

$$\frac{x^{-n} \cos(a+bx) (\cos(a+bx))^{n-1} \left(3 \cdot 2^{2n} {}_2F_1(1, 1+n; 2+n; \cos(a+bx)) + 3 \cdot 2^{2n} {}_2F_1(2, 1+n; 2+n; \cos(a+bx)) + 2^{2n} {}_2F_1(3, 1+n; 2+n; \cos(a+bx)) + 2 {}_2F_1(-1+n, 1+n; 2+n; \cos(a+bx)) \sec^2\left(\frac{a+bx}{2}\right) \sec^{2n}\left(\frac{a+bx}{2}\right) + 3 {}_2F_1(n, 1+n; 2+n; \cos(a+bx)) \sec^2\left(\frac{a+bx}{2}\right) \sec^{2n}\left(\frac{a+bx}{2}\right) + 3 {}_2F_1(1+n, 1+n; 2+n; \cos(a+bx)) \sec^2\left(\frac{a+bx}{2}\right) \sec^{2n}\left(\frac{a+bx}{2}\right) \right)}{(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Cos[a + b*x])^n*Csc[a + b*x]^5,x]

[Out] -((2^(-5 - n)*Cos[a + b*x]*(d*Cos[a + b*x])^n*(3*2^(1 + n)*Hypergeometric2F1[1, 1 + n, 2 + n, Cos[a + b*x]] + 3*2^(1 + n)*Hypergeometric2F1[2, 1 + n, 2 + n, Cos[a + b*x]] + 2^(2 + n)*Hypergeometric2F1[3, 1 + n, 2 + n, Cos[a + b*x]] + 2*Hypergeometric2F1[-1 + n, 1 + n, 2 + n, (Cos[a + b*x]*Sec[(a + b*x)/2]^2)/2]*(Sec[(a + b*x)/2]^2)^(1 + n) + 3*Hypergeometric2F1[n, 1 + n, 2 + n, (Cos[a + b*x]*Sec[(a + b*x)/2]^2)/2]*(Sec[(a + b*x)/2]^2)^(1 + n) + 3*Hypergeometric2F1[1 + n, 1 + n, 2 + n, (Cos[a + b*x]*Sec[(a + b*x)/2]^2)/2]*(Sec[(a + b*x)/2]^2)^(1 + n)))/(b*(1 + n))

Maple [F]

time = 0.07, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (\csc^5(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*csc(b*x+a)^5,x)

[Out] int((d*cos(b*x+a))^n*csc(b*x+a)^5,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^5,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^5, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^5,x, algorithm="fricas")

[Out] `integral((d*cos(b*x + a))^n*csc(b*x + a)^5, x)`

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^n*csc(b*x+a)**5,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*cos(b*x+a))^n*csc(b*x+a)^5,x, algorithm="giac")`

[Out] `integrate((d*cos(b*x + a))^n*csc(b*x + a)^5, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(d \cos(a + bx))^n}{\sin(a + bx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d*cos(a + b*x))^n/sin(a + b*x)^5,x)`

[Out] `int((d*cos(a + b*x))^n/sin(a + b*x)^5, x)`

3.361 $\int (d \cos(a + bx))^n \sin^4(a + bx) dx$

Optimal. Leaf size=69

$$\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{3}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bd(1+n) \sqrt{\sin^2(a + bx)}}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}\left(\left[-\frac{3}{2}, \frac{1}{2}+\frac{1}{2}*n\right], \left[\frac{3}{2}+\frac{1}{2}*n\right], \cos(b*x+a)^2\right)*\sin(b*x+a)/b/d/(1+n)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2656}

$$\frac{\sin(a + bx)(d \cos(a + bx))^{n+1} {}_2F_1\left(-\frac{3}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1) \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Sin}[a + b*x]^4, x]$

[Out] $-\left(\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)}*\text{Hypergeometric2F1}\left[-\frac{3}{2}, \frac{(1+n)}{2}, \frac{(3+n)}{2}, \text{Cos}[a + b*x]^2\right]*\text{Sin}[a + b*x]\right)\right)/(b*d*(1+n)*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.))^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_)}], x_Symbol] \rightarrow \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*((a*\text{Cos}[e + f*x])^{(m+1)})/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]})*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(a + bx))^n \sin^4(a + bx) dx = -\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{3}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bd(1+n) \sqrt{\sin^2(a + bx)}}$$

Mathematica [A]

time = 0.09, size = 68, normalized size = 0.99

$$\frac{(d \cos(a + bx))^n {}_2F_1\left(-\frac{3}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(2(a + bx))}{2b(1+n) \sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*sin[a + b*x]^4,x]

[Out] $-1/2*((d*\cos[a + b*x])^n*\text{Hypergeometric2F1}[-3/2, (1 + n)/2, (3 + n)/2, \cos[a + b*x]^2]*\sin[2*(a + b*x)])/(b*(1 + n)*\sqrt{\sin[a + b*x]^2})$

Maple [F]

time = 0.29, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (\sin^4(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*sin(b*x+a)^4,x)

[Out] int((d*cos(b*x+a))^n*sin(b*x+a)^4,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^4,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n*sin(b*x + a)^4, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^4,x, algorithm="fricas")

[Out] $\text{integral}((\cos(b*x + a))^4 - 2*\cos(b*x + a)^2 + 1)*(d*\cos(b*x + a))^n, x)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)**4,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^4,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n*sin(b*x + a)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^4 (d \cos(a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^4*(d*cos(a + b*x))^n,x)

[Out] int(sin(a + b*x)^4*(d*cos(a + b*x))^n, x)

3.362 $\int (d \cos(a + bx))^n \sin^2(a + bx) dx$

Optimal. Leaf size=69

$$\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bd(1+n)\sqrt{\sin^2(a + bx)}}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([-1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*\sin(b*x+a)/b/d/(1+n)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2656}

$$\frac{\sin(a + bx)(d \cos(a + bx))^{n+1} {}_2F_1\left(-\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Sin}[a + b*x]^2, x]$

[Out] $-\left(\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)}*\text{Hypergeometric2F1}\left[-1/2, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2*\text{Sin}[a + b*x]\right)\right)/(b*d*(1+n)*\text{Sqrt}[\text{Sin}[a + b*x]^2])$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^n), x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*((a*\text{Cos}[e + f*x])^{(m+1)}/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]})*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x) /; \text{FreeQ}\{a, b, e, f, m, n\}, x\} \&\& \text{SimplerQ}\{n, m\}$

Rubi steps

$$\int (d \cos(a + bx))^n \sin^2(a + bx) dx = -\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bd(1+n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A]

time = 0.07, size = 68, normalized size = 0.99

$$\frac{(d \cos(a + bx))^n {}_2F_1\left(-\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(2(a + bx))}{2b(1+n)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*sin[a + b*x]^2,x]

[Out] $-1/2*((d*\cos[a + b*x])^n*\text{Hypergeometric2F1}[-1/2, (1 + n)/2, (3 + n)/2, \cos[a + b*x]^2]*\sin[2*(a + b*x)])/(b*(1 + n)*\sqrt{\sin[a + b*x]^2})$

Maple [F]

time = 0.37, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (\sin^2(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*sin(b*x+a)^2,x)

[Out] int((d*cos(b*x+a))^n*sin(b*x+a)^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^2,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n*sin(b*x + a)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^2,x, algorithm="fricas")

[Out] integral(-(cos(b*x + a)^2 - 1)*(d*cos(b*x + a))^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(a + bx))^n \sin^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*sin(b*x+a)**2,x)

[Out] Integral((d*cos(a + b*x))**n*sin(a + b*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*sin(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n*sin(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(a + bx)^2 (d \cos(a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(a + b*x)^2*(d*cos(a + b*x))^n,x)

[Out] int(sin(a + b*x)^2*(d*cos(a + b*x))^n, x)

3.363 $\int (d \cos(a + bx))^n dx$

Optimal. Leaf size=69

$$-\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bd(1+n)\sqrt{\sin^2(a + bx)}}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([1/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*\sin(b*x+a)/b/d/(1+n)/(\sin(b*x+a)^2)^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.100$, Rules used = {2722}

$$-\frac{\sin(a + bx)(d \cos(a + bx))^{n+1} {}_2F_1\left(\frac{1}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)\sqrt{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*cos[a + b*x])^n,x]

[Out] $-\left(\left(d*\cos[a + b*x]\right)^{(1 + n)}*\text{Hypergeometric2F1}\left[1/2, (1 + n)/2, (3 + n)/2, \cos[a + b*x]^2\right]*\sin[a + b*x]\right)/(b*d*(1 + n)*\text{Sqrt}[\sin[a + b*x]^2])$

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2, x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rubi steps

$$\int (d \cos(a + bx))^n dx = -\frac{(d \cos(a + bx))^{1+n} {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx)}{bd(1+n)\sqrt{\sin^2(a + bx)}}$$

Mathematica [A]

time = 0.04, size = 64, normalized size = 0.93

$$-\frac{(d \cos(a + bx))^n \cot(a + bx) {}_2F_1\left(\frac{1}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{b(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n,x]

[Out] -((((d*cos[a + b*x])^n*cot[a + b*x]*Hypergeometric2F1[1/2, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]*Sqrt[Sin[a + b*x]^2]))/(b*(1 + n)))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n,x)

[Out] int((d*cos(b*x+a))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n,x, algorithm="fricas")

[Out] integral((d*cos(b*x + a))^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n,x)

[Out] Integral((d*cos(a + b*x))^n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + b x))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n,x)

[Out] int((d*cos(a + b*x))^n, x)

3.364 $\int (d \cos(a + bx))^n \csc^2(a + bx) dx$

Optimal. Leaf size=69

$$\frac{(d \cos(a + bx))^{1+n} \csc(a + bx) {}_2F_1\left(\frac{3}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{bd(1+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\csc(b*x+a)*\text{hypergeom}([3/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*(\sin(b*x+a)^2)^{(1/2)}/b/d/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2656}

$$\frac{\sqrt{\sin^2(a + bx)} \csc(a + bx) (d \cos(a + bx))^{n+1} {}_2F_1\left(\frac{3}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Csc}[a + b*x]^2,x]$

[Out] $-(((d*\text{Cos}[a + b*x])^{(1+n)}*\text{Csc}[a + b*x]*\text{Hypergeometric2F1}[3/2, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2]*\text{Sqrt}[\text{Sin}[a + b*x]^2])/(b*d*(1+n)))$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*((a*\text{Cos}[e + f*x])^{(m+1)}/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]})))*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(a + bx))^n \csc^2(a + bx) dx = -\frac{(d \cos(a + bx))^{1+n} \csc(a + bx) {}_2F_1\left(\frac{3}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{bd(1+n)}$$

Mathematica [A]

time = 0.16, size = 80, normalized size = 1.16

$$\frac{d(d \cos(a + bx))^{-1+n} (-\cot^2(a + bx))^{\frac{1-n}{2}} \csc(a + bx) {}_2F_1\left(\frac{1-n}{2}, 1 - \frac{n}{2}; 2 - \frac{n}{2}; \csc^2(a + bx)\right)}{b(-2+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*csc[a + b*x]^2,x]

[Out] (d*(d*cos[a + b*x])^(-1 + n)*(-Cot[a + b*x]^2)^((1 - n)/2)*Csc[a + b*x]*Hypergeometric2F1[(1 - n)/2, 1 - n/2, 2 - n/2, Csc[a + b*x]^2])/(b*(-2 + n))

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (\csc^2(bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*csc(b*x+a)^2,x)

[Out] int((d*cos(b*x+a))^n*csc(b*x+a)^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^2,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^2,x, algorithm="fricas")

[Out] integral((d*cos(b*x + a))^n*csc(b*x + a)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos(a + bx))^n \csc^2(a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*csc(b*x+a)**2,x)

[Out] Integral((d*cos(a + b*x))**n*csc(a + b*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^2,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + bx))^n}{\sin(a + bx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n/sin(a + b*x)^2,x)

[Out] int((d*cos(a + b*x))^n/sin(a + b*x)^2, x)

3.365 $\int (d \cos(a + bx))^n \csc^4(a + bx) dx$

Optimal. Leaf size=69

$$\frac{(d \cos(a + bx))^{1+n} \csc(a + bx) {}_2F_1\left(\frac{5}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{bd(1+n)}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*csc(b*x+a)*hypergeom([5/2, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*(sin(b*x+a)^2)^{(1/2)}/b/d/(1+n)$

Rubi [A]

time = 0.03, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {2656}

$$\frac{\sqrt{\sin^2(a + bx)} \csc(a + bx) (d \cos(a + bx))^{n+1} {}_2F_1\left(\frac{5}{2}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Csc}[a + b*x]^4,x]$

[Out] $-\left(\left(d*\text{Cos}[a + b*x]\right)^{(1+n)}*\text{Csc}[a + b*x]*\text{Hypergeometric2F1}\left[5/2, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2\right]*\text{Sqrt}[\text{Sin}[a + b*x]^2]\right)/(b*d*(1+n))$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\sin[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*(a*\cos[e + f*x])^{(m+1)}/(a*f*(m+1)*(sin[e + f*x]^2)^{\text{FracPart}[(n-1)/2]})]*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \cos[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(a + bx))^n \csc^4(a + bx) dx = -\frac{(d \cos(a + bx))^{1+n} \csc(a + bx) {}_2F_1\left(\frac{5}{2}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{\sin^2(a + bx)}}{bd(1+n)}$$

Mathematica [A]

time = 0.16, size = 82, normalized size = 1.19

$$\frac{d(d \cos(a + bx))^{-1+n} (-\cot^2(a + bx))^{\frac{1-n}{2}} \csc^3(a + bx) {}_2F_1\left(\frac{1-n}{2}, 2 - \frac{n}{2}; 3 - \frac{n}{2}; \cos^2(a + bx)\right)}{b(-4+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*csc[a + b*x]^4,x]

[Out] (d*(d*cos[a + b*x])^(-1 + n)*(-Cot[a + b*x]^2)^((1 - n)/2)*Csc[a + b*x]^3*Hypergeometric2F1[(1 - n)/2, 2 - n/2, 3 - n/2, Csc[a + b*x]^2])/(b*(-4 + n))

Maple [F]

time = 0.07, size = 0, normalized size = 0.00

$$\int (d \cos (bx + a))^n (\csc^4 (bx + a)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*csc(b*x+a)^4,x)

[Out] int((d*cos(b*x+a))^n*csc(b*x+a)^4,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^4,x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^4, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^4,x, algorithm="fricas")

[Out] integral((d*cos(b*x + a))^n*csc(b*x + a)^4, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \cos (a + bx))^n \csc^4 (a + bx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*csc(b*x+a)**4,x)

[Out] Integral((d*cos(a + b*x))**n*csc(a + b*x)**4, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*csc(b*x+a)^4,x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n*csc(b*x + a)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + bx))^n}{\sin(a + bx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n/sin(a + b*x)^4,x)

[Out] int((d*cos(a + b*x))^n/sin(a + b*x)^4, x)

3.366 $\int (d \cos(a + bx))^n (c \sin(a + bx))^{5/2} dx$

Optimal. Leaf size=76

$$\frac{c(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{3}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) (c \sin(a + bx))^{3/2}}{bd(1+n) \sin^2(a + bx)^{3/4}}$$

[Out] $-c*(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}\left(\left[-\frac{3}{4}, \frac{1}{2}+\frac{1}{2}*n\right], \left[\frac{3}{2}+\frac{1}{2}*n\right], \cos(b*x+a)\right)^{-2}*(c*\sin(b*x+a))^{(3/2)}/b/d/(1+n)/(\sin(b*x+a)^2)^{(3/4)}$

Rubi [A]

time = 0.04, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2656}

$$\frac{c(c \sin(a + bx))^{3/2} (d \cos(a + bx))^{n+1} {}_2F_1\left(-\frac{3}{4}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1) \sin^2(a + bx)^{3/4}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*(c*\text{Sin}[a + b*x])^{(5/2)}, x]$

[Out] $-((c*(d*\text{Cos}[a + b*x])^{(1 + n)}*\text{Hypergeometric2F1}\left[-\frac{3}{4}, \frac{(1 + n)}{2}, \frac{(3 + n)}{2}, \text{Cos}[a + b*x]^2\right]*(c*\text{Sin}[a + b*x])^{(3/2)}))/(b*d*(1 + n)*(\text{Sin}[a + b*x]^2)^{(3/4}))$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}), x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}*((a*\text{Cos}[e + f*x])^{(m + 1)}/(a*f*(m + 1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n - 1)/2]})))*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(a + bx))^n (c \sin(a + bx))^{5/2} dx = -\frac{c(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{3}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) (c \sin(a + bx))^{3/2}}{bd(1+n) \sin^2(a + bx)^{3/4}}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 158 vs. 2(76) = 152.

time = 0.30, size = 158, normalized size = 2.08

$$\frac{(d \cos(a + bx))^n \cot(a + bx) \left(-((3 + n) {}_2F_1\left(-\frac{3}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right)) - (3 + n) {}_2F_1\left(\frac{1}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) + (1 + n) \cos^2(a + bx) {}_2F_1\left(\frac{1}{4}, \frac{3+n}{2}; \frac{5+n}{2}; \cos^2(a + bx)\right) \right) (c \sin(a + bx))^{5/2}}{2b(1+n)(3+n) \sin^2(a + bx)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*(c*sin[a + b*x])^(5/2),x]

[Out] ((d*cos[a + b*x])^n*cot[a + b*x]*(-(3 + n)*Hypergeometric2F1[-3/4, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]) - (3 + n)*Hypergeometric2F1[1/4, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2] + (1 + n)*Cos[a + b*x]^2*Hypergeometric2F1[1/4, (3 + n)/2, (5 + n)/2, Cos[a + b*x]^2])*(c*sin[a + b*x])^(5/2))/(2*b*(1 + n)*(3 + n)*(Sin[a + b*x]^2)^(3/4))

Maple [F]

time = 0.09, size = 0, normalized size = 0.00

$$\int (d \cos(bx + a))^n (c \sin(bx + a))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*(c*sin(b*x+a))^(5/2),x)

[Out] int((d*cos(b*x+a))^n*(c*sin(b*x+a))^(5/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(5/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(5/2)*(d*cos(b*x + a))^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(5/2),x, algorithm="fricas")

[Out] integral(-(c^2*cos(b*x + a)^2 - c^2)*sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n, x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 8570 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(5/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(5/2)*(d*cos(b*x + a))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + bx))^n (c \sin(a + bx))^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n*(c*sin(a + b*x))^(5/2),x)

[Out] int((d*cos(a + b*x))^n*(c*sin(a + b*x))^(5/2), x)

3.367 $\int (d \cos(a + bx))^n (c \sin(a + bx))^{3/2} dx$

Optimal. Leaf size=76

$$-\frac{c(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{1}{4}, \frac{1+n}{2}, \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{c \sin(a + bx)}}{bd(1+n) \sqrt[4]{\sin^2(a + bx)}}$$

[Out] $-c*(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}\left[\left[-\frac{1}{4}, \frac{1}{2}+\frac{1}{2}*n\right], \left[\frac{3}{2}+\frac{1}{2}*n\right], \cos(b*x+a)^2\right]*(c*\sin(b*x+a))^{(1/2)}/b/d/(1+n)/(\sin(b*x+a)^2)^{(1/4)}$

Rubi [A]

time = 0.04, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2656}

$$-\frac{c\sqrt{c \sin(a + bx)} (d \cos(a + bx))^{n+1} {}_2F_1\left(-\frac{1}{4}, \frac{n+1}{2}, \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1) \sqrt[4]{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*(c*\text{Sin}[a + b*x])^{(3/2)}, x]$

[Out] $-((c*(d*\text{Cos}[a + b*x])^{(1 + n)}*\text{Hypergeometric2F1}[-1/4, (1 + n)/2, (3 + n)/2, \text{Cos}[a + b*x]^2]*\text{Sqrt}[c*\text{Sin}[a + b*x]])/(b*d*(1 + n)*(\text{Sin}[a + b*x]^2)^{(1/4)})$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_)]))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2]}*((a*\text{Cos}[e + f*x])^{(m + 1)})/(a*f*(m + 1)*(\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2]}))*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(a + bx))^n (c \sin(a + bx))^{3/2} dx = -\frac{c(d \cos(a + bx))^{1+n} {}_2F_1\left(-\frac{1}{4}, \frac{1+n}{2}, \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt{c \sin(a + bx)}}{bd(1+n) \sqrt[4]{\sin^2(a + bx)}}$$

Mathematica [A]

time = 0.11, size = 76, normalized size = 1.00

$$-\frac{(d \cos(a + bx))^n \cot(a + bx) {}_2F_1\left(-\frac{1}{4}, \frac{1+n}{2}, \frac{3+n}{2}; \cos^2(a + bx)\right) (c \sin(a + bx))^{3/2}}{b(1+n) \sqrt[4]{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*(c*sin[a + b*x])^(3/2),x]

[Out] -((((d*cos[a + b*x])^n*cot[a + b*x]*Hypergeometric2F1[-1/4, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]*(c*sin[a + b*x])^(3/2))/(b*(1 + n)*(Sin[a + b*x]^2)^(1/4))))

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (d \cos (bx + a))^n (c \sin (bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*(c*sin(b*x+a))^(3/2),x)

[Out] int((d*cos(b*x+a))^n*(c*sin(b*x+a))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^(3/2)*(d*cos(b*x + a))^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n*c*sin(b*x + a), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin (a + bx))^{\frac{3}{2}} (d \cos (a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(3/2),x)

[Out] Integral((c*sin(a + b*x))**(3/2)*(d*cos(a + b*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^(3/2)*(d*cos(b*x + a))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + bx))^n (c \sin(a + bx))^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n*(c*sin(a + b*x))^(3/2),x)

[Out] int((d*cos(a + b*x))^n*(c*sin(a + b*x))^(3/2), x)

3.368 $\int (d \cos(a + bx))^n \sqrt{c \sin(a + bx)} dx$

Optimal. Leaf size=76

$$-\frac{c(d \cos(a + bx))^{1+n} {}_2F_1\left(\frac{1}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt[4]{\sin^2(a + bx)}}{bd(1+n) \sqrt{c \sin(a + bx)}}$$

[Out] $-c*(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([1/4, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*(\sin(b*x+a)^2)^{(1/4)}/b/d/(1+n)/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2656}

$$-\frac{c^4 \sqrt{\sin^2(a + bx)} (d \cos(a + bx))^{n+1} {}_2F_1\left(\frac{1}{4}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a + bx)\right)}{bd(n+1) \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n*\text{Sqrt}[c*\text{Sin}[a + b*x]], x]$

[Out] $-((c*(d*\text{Cos}[a + b*x])^{(1 + n)}*\text{Hypergeometric2F1}[1/4, (1 + n)/2, (3 + n)/2, \text{Cos}[a + b*x]^2]*(\text{Sin}[a + b*x]^2)^{(1/4)})/(b*d*(1 + n)*\text{Sqrt}[c*\text{Sin}[a + b*x]])$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2]})*(a*\text{Cos}[e + f*x])^{(m + 1)}/(a*f*(m + 1)*(\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2]}))*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int (d \cos(a + bx))^n \sqrt{c \sin(a + bx)} dx = -\frac{c(d \cos(a + bx))^{1+n} {}_2F_1\left(\frac{1}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sqrt[4]{\sin^2(a + bx)}}{bd(1+n) \sqrt{c \sin(a + bx)}}$$

Mathematica [A]

time = 0.08, size = 82, normalized size = 1.08

$$-\frac{\cos(a + bx)(d \cos(a + bx))^n {}_2F_1\left(\frac{1}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a + bx)\right) \sin(a + bx) \sqrt{c \sin(a + bx)}}{b(1+n) \sin^2(a + bx)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n*Sqrt[c*sin[a + b*x]],x]

[Out] -((Cos[a + b*x]*(d*cos[a + b*x])^n*Hypergeometric2F1[1/4, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]*Sin[a + b*x]*Sqrt[c*sin[a + b*x]])/(b*(1 + n)*(Sin[a + b*x]^2)^(3/4)))

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (d \cos (bx + a))^n \sqrt{c \sin (bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n*(c*sin(b*x+a))^(1/2),x)

[Out] int((d*cos(b*x+a))^n*(c*sin(b*x+a))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c \sin (a + bx)} (d \cos (a + bx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n*(c*sin(b*x+a))**(1/2),x)

[Out] Integral(sqrt(c*sin(a + b*x))*(d*cos(a + b*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n*(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (d \cos(a + bx))^n \sqrt{c \sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n*(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^n*(c*sin(a + b*x))^(1/2), x)

$$3.369 \quad \int \frac{(d \cos(a+bx))^n}{\sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=76

$$-\frac{c(d \cos(a+bx))^{1+n} {}_2F_1\left(\frac{3}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a+bx)\right) \sin^2(a+bx)^{3/4}}{bd(1+n)(c \sin(a+bx))^{3/2}}$$

[Out] $-c*(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([3/4, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*(\sin(b*x+a)^2)^{(3/4)}/b/d/(1+n)/(c*\sin(b*x+a))^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2656}

$$-\frac{c \sin^2(a+bx)^{3/4} (d \cos(a+bx))^{n+1} {}_2F_1\left(\frac{3}{4}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a+bx)\right)}{bd(n+1)(c \sin(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n/\text{Sqrt}[c*\text{Sin}[a + b*x]], x]$

[Out] $-\left(\left(c*(d*\text{Cos}[a + b*x])^{(1+n)}*\text{Hypergeometric2F1}[3/4, (1+n)/2, (3+n)/2, \text{Cos}[a + b*x]^2]*(\text{Sin}[a + b*x]^2)^{(3/4)}\right)/(b*d*(1+n)*(c*\text{Sin}[a + b*x])^{(3/2)})\right)$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sin[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b^{(2*\text{IntPart}[(n-1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n-1)/2])}*(a*\text{Cos}[e + f*x])^{(m+1)}/(a*f*(m+1)*(\text{Sin}[e + f*x]^2)^{\text{FracPart}[(n-1)/2]}))*\text{Hypergeometric2F1}[(1+m)/2, (1-n)/2, (3+m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{SimplerQ}[n, m]$

Rubi steps

$$\int \frac{(d \cos(a+bx))^n}{\sqrt{c \sin(a+bx)}} dx = -\frac{c(d \cos(a+bx))^{1+n} {}_2F_1\left(\frac{3}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a+bx)\right) \sin^2(a+bx)^{3/4}}{bd(1+n)(c \sin(a+bx))^{3/2}}$$

Mathematica [A]

time = 0.09, size = 82, normalized size = 1.08

$$-\frac{\cos(a+bx)(d \cos(a+bx))^n {}_2F_1\left(\frac{3}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a+bx)\right) \sin(a+bx)}{b(1+n)\sqrt{c \sin(a+bx)} \sqrt[4]{\sin^2(a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n/Sqrt[c*Sin[a + b*x]],x]

[Out] -((Cos[a + b*x]*(d*cos[a + b*x])^n*Hypergeometric2F1[3/4, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2*Sin[a + b*x])/(b*(1 + n)*Sqrt[c*Sin[a + b*x]]*(Sin[a + b*x]^2)^(1/4)))

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int \frac{(d \cos (bx + a))^n}{\sqrt{c \sin (bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n/(c*sin(b*x+a))^(1/2),x)

[Out] int((d*cos(b*x+a))^n/(c*sin(b*x+a))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n/sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n/(c*sin(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \cos (a + bx))^n}{\sqrt{c \sin (a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(1/2),x)

[Out] Integral((d*cos(a + b*x))**n/sqrt(c*sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n/sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + bx))^n}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n/(c*sin(a + b*x))^(1/2),x)

[Out] int((d*cos(a + b*x))^n/(c*sin(a + b*x))^(1/2), x)

$$3.370 \quad \int \frac{(d \cos(a+bx))^n}{(c \sin(a+bx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{(d \cos(a+bx))^{1+n} {}_2F_1\left(\frac{5}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a+bx)\right) \sqrt[4]{\sin^2(a+bx)}}{bcd(1+n) \sqrt{c \sin(a+bx)}}$$

[Out] $-(d*\cos(b*x+a))^{(1+n)}*\text{hypergeom}([5/4, 1/2+1/2*n], [3/2+1/2*n], \cos(b*x+a)^2)*(\sin(b*x+a)^2)^{(1/4)}/b/c/d/(1+n)/(c*\sin(b*x+a))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2656}

$$\frac{\sqrt[4]{\sin^2(a+bx)} (d \cos(a+bx))^{n+1} {}_2F_1\left(\frac{5}{4}, \frac{n+1}{2}; \frac{n+3}{2}; \cos^2(a+bx)\right)}{bcd(n+1) \sqrt{c \sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Cos}[a + b*x])^n/(c*\text{Sin}[a + b*x])^{(3/2)}, x]$

[Out] $-(((d*\text{Cos}[a + b*x])^{(1 + n)}*\text{Hypergeometric2F1}[5/4, (1 + n)/2, (3 + n)/2, \text{Cos}[a + b*x]^2]*(\text{Sin}[a + b*x]^2)^{(1/4)})/(b*c*d*(1 + n)*\text{Sqrt}[c*\text{Sin}[a + b*x]]))$

Rule 2656

$\text{Int}[(\cos[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sin[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[(-b^{(2*\text{IntPart}[(n - 1)/2] + 1)}*(b*\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}*((a*\text{Cos}[e + f*x])^{(m + 1)}/(a*f*(m + 1)*(\text{Sin}[e + f*x])^{(2*\text{FracPart}[(n - 1)/2])}))*\text{Hypergeometric2F1}[(1 + m)/2, (1 - n)/2, (3 + m)/2, \text{Cos}[e + f*x]^2], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{SimplerQ}[n, m]$

Rubi steps

$$\int \frac{(d \cos(a+bx))^n}{(c \sin(a+bx))^{3/2}} dx = -\frac{(d \cos(a+bx))^{1+n} {}_2F_1\left(\frac{5}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a+bx)\right) \sqrt[4]{\sin^2(a+bx)}}{bcd(1+n) \sqrt{c \sin(a+bx)}}$$

Mathematica [A]

time = 0.10, size = 79, normalized size = 1.01

$$\frac{(d \cos(a+bx))^n \cot(a+bx) {}_2F_1\left(\frac{5}{4}, \frac{1+n}{2}; \frac{3+n}{2}; \cos^2(a+bx)\right) \sqrt{c \sin(a+bx)} \sqrt[4]{\sin^2(a+bx)}}{bc^2(1+n)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*cos[a + b*x])^n/(c*sin[a + b*x])^(3/2),x]

[Out] -(((d*cos[a + b*x])^n*Cot[a + b*x]*Hypergeometric2F1[5/4, (1 + n)/2, (3 + n)/2, Cos[a + b*x]^2]*Sqrt[c*sin[a + b*x]]*(Sin[a + b*x]^2)^(1/4))/(b*c^2*(1 + n)))

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int \frac{(d \cos(bx + a))^n}{(c \sin(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(b*x+a))^n/(c*sin(b*x+a))^(3/2),x)

[Out] int((d*cos(b*x+a))^n/(c*sin(b*x+a))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((d*cos(b*x + a))^n/(c*sin(b*x + a))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(-sqrt(c*sin(b*x + a))*(d*cos(b*x + a))^n/(c^2*cos(b*x + a)^2 - c^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(d \cos(a + bx))^n}{(c \sin(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))**n/(c*sin(b*x+a))**(3/2),x)

[Out] Integral((d*cos(a + b*x))**n/(c*sin(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*cos(b*x+a))^n/(c*sin(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((d*cos(b*x + a))^n/(c*sin(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(d \cos(a + b x))^n}{(c \sin(a + b x))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*cos(a + b*x))^n/(c*sin(a + b*x))^(3/2),x)

[Out] int((d*cos(a + b*x))^n/(c*sin(a + b*x))^(3/2), x)

3.371 $\int \sqrt{b \sec(e + fx)} \sin^7(e + fx) dx$

Optimal. Leaf size=85

$$\frac{2b^7}{13f(b \sec(e + fx))^{13/2}} - \frac{2b^5}{3f(b \sec(e + fx))^{9/2}} + \frac{6b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}}$$

[Out] $2/13*b^7/f/(b*\sec(f*x+e))^(13/2)-2/3*b^5/f/(b*\sec(f*x+e))^(9/2)+6/5*b^3/f/(b*\sec(f*x+e))^(5/2)-2*b/f/(b*\sec(f*x+e))^(1/2)$

Rubi [A]

time = 0.04, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$\frac{2b^7}{13f(b \sec(e + fx))^{13/2}} - \frac{2b^5}{3f(b \sec(e + fx))^{9/2}} + \frac{6b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^7,x]`

[Out] $(2*b^7)/(13*f*(b*\text{Sec}[e + f*x])^(13/2)) - (2*b^5)/(3*f*(b*\text{Sec}[e + f*x])^(9/2)) + (6*b^3)/(5*f*(b*\text{Sec}[e + f*x])^(5/2)) - (2*b)/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\int \sqrt{b \sec(e + fx)} \sin^7(e + fx) dx = \frac{b^7 \text{Subst} \left(\int \frac{(-1 + \frac{x^2}{b^2})^3}{x^{15/2}} dx, x, b \sec(e + fx) \right)}{f}$$

$$= \frac{b^7 \text{Subst} \left(\int \left(-\frac{1}{x^{15/2}} + \frac{3}{b^2 x^{11/2}} - \frac{3}{b^4 x^{7/2}} + \frac{1}{b^6 x^{3/2}} \right) dx, x, b \sec(e + fx) \right)}{f}$$

$$= \frac{2b^7}{13f(b \sec(e + fx))^{13/2}} - \frac{2b^5}{3f(b \sec(e + fx))^{9/2}} + \frac{6b^3}{5f(b \sec(e + fx))^{5/2}}$$

Mathematica [A]

time = 0.24, size = 58, normalized size = 0.68

$$\frac{(-8939 \cos(e + fx) + 887 \cos(3(e + fx)) - 155 \cos(5(e + fx)) + 15 \cos(7(e + fx))) \sqrt{b \sec(e + fx)}}{6240f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^7,x]**[Out]** ((-8939*Cos[e + f*x] + 887*Cos[3*(e + f*x)] - 155*Cos[5*(e + f*x)] + 15*Cos[7*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/(6240*f)**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 516 vs. 2(71) = 142.

time = 2.74, size = 517, normalized size = 6.08

method	result
default	$-\frac{(-1 + \cos(fx + e))^2 \left(-60(\cos^7(fx + e)) + 260(\cos^5(fx + e)) + 195 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \ln \left(-\frac{2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}}}{(\cos(fx + e) + 1)} \right) \right)}{6240f}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^7*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-1/390/f*(-1+\cos(f*x+e))^2*(-60*\cos(f*x+e)^7+260*\cos(f*x+e)^5+195*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)-1)/\sin(f*x+e)^2*\cos(f*x+e)-195*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)-1)/\sin(f*x+e)^2*\cos(f*x+e)-468*\cos(f*x+e)^3+195*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^(1/2)-c$$

$$\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-1}/\sin(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-195*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-1}/\sin(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)+780*\cos(f*x+e))*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(1/2)}/\sin(f*x+e)^4}$$

Maxima [A]

time = 0.28, size = 67, normalized size = 0.79

$$\frac{2 \left(15 b^6 - \frac{65 b^6}{\cos(fx+e)^2} + \frac{117 b^6}{\cos(fx+e)^4} - \frac{195 b^6}{\cos(fx+e)^6} \right) b}{195 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{13}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] 2/195*(15*b^6 - 65*b^6/cos(f*x + e)^2 + 117*b^6/cos(f*x + e)^4 - 195*b^6/cos(f*x + e)^6)*b/(f*(b/cos(f*x + e))^(13/2))

Fricas [A]

time = 0.40, size = 61, normalized size = 0.72

$$\frac{2 \left(15 \cos(fx+e)^7 - 65 \cos(fx+e)^5 + 117 \cos(fx+e)^3 - 195 \cos(fx+e) \right) \sqrt{\frac{b}{\cos(fx+e)}}}{195 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 2/195*(15*cos(f*x + e)^7 - 65*cos(f*x + e)^5 + 117*cos(f*x + e)^3 - 195*cos(f*x + e))*sqrt(b/cos(f*x + e))/f

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**7*(b*sec(f*x+e))**(1/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 5006 deep

Giac [A]

time = 2.66, size = 108, normalized size = 1.27

$$\frac{2 \left(15 \sqrt{b \cos(fx+e)} b^6 \cos(fx+e)^6 - 65 \sqrt{b \cos(fx+e)} b^6 \cos(fx+e)^4 + 117 \sqrt{b \cos(fx+e)} b^6 \cos(fx+e)^2 - 195 \sqrt{b \cos(fx+e)} b^6 \right) \operatorname{sgn}(\cos(fx+e))}{195 b^6 f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^7*(b*sec(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] 2/195*(15*sqrt(b*cos(f*x + e))*b^6*cos(f*x + e)^6 - 65*sqrt(b*cos(f*x + e))
*b^6*cos(f*x + e)^4 + 117*sqrt(b*cos(f*x + e))*b^6*cos(f*x + e)^2 - 195*sqrt
(b*cos(f*x + e))*b^6)*sgn(cos(f*x + e))/(b^6*f)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^7 \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^7*(b/cos(e + f*x))^(1/2),x)
```

```
[Out] int(sin(e + f*x)^7*(b/cos(e + f*x))^(1/2), x)
```

3.372 $\int \sqrt{b \sec(e + fx)} \sin^5(e + fx) dx$

Optimal. Leaf size=63

$$-\frac{2b^5}{9f(b \sec(e + fx))^{9/2}} + \frac{4b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}}$$

[Out] $-2/9*b^5/f/(b*\sec(f*x+e))^{(9/2)}+4/5*b^3/f/(b*\sec(f*x+e))^{(5/2)}-2*b/f/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$-\frac{2b^5}{9f(b \sec(e + fx))^{9/2}} + \frac{4b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^5,x]`

[Out] $(-2*b^5)/(9*f*(b*\text{Sec}[e + f*x])^{(9/2)}) + (4*b^3)/(5*f*(b*\text{Sec}[e + f*x])^{(5/2)}) - (2*b)/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned} \int \sqrt{b \sec(e+fx)} \sin^5(e+fx) dx &= \frac{b^5 \text{Subst}\left(\int \frac{(-1+\frac{x^2}{b^2})^2}{x^{11/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{b^5 \text{Subst}\left(\int \left(\frac{1}{x^{11/2}} - \frac{2}{b^2 x^{7/2}} + \frac{1}{b^4 x^{3/2}}\right) dx, x, b \sec(e+fx)\right)}{f} \\ &= -\frac{2b^5}{9f(b \sec(e+fx))^{9/2}} + \frac{4b^3}{5f(b \sec(e+fx))^{5/2}} - \frac{2b}{f \sqrt{b \sec(e+fx)}} \end{aligned}$$

Mathematica [A]

time = 0.15, size = 48, normalized size = 0.76

$$\frac{(554 \cos(e+fx) - 47 \cos(3(e+fx)) + 5 \cos(5(e+fx))) \sqrt{b \sec(e+fx)}}{360f}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^5,x]``[Out] -1/360*((554*Cos[e + f*x] - 47*Cos[3*(e + f*x)] + 5*Cos[5*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/f`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 506 vs. 2(53) = 106.

time = 0.26, size = 507, normalized size = 8.05

method	result
default	$\frac{(-1+\cos(fx+e))^2 \left(20(\cos^5(fx+e))-72(\cos^3(fx+e))+45\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} \ln\left(-\frac{2(\cos^2(fx+e))\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}{(\cos(fx+e)+1)^2}\right) \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^5*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/90/f*(-1+cos(f*x+e))^2*(20*cos(f*x+e)^5-72*cos(f*x+e)^3+45*(-cos(f*x+e)/
(cos(f*x+e)+1)^2)^(1/2)*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(
1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/s
in(f*x+e)^2)*cos(f*x+e)-45*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*ln(-2*(2*co
s(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2
*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*cos(f*x+e)+45*ln(-2
*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e
```

$$)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-1}/\sin(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-45*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-1}/\sin(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)+180*\cos(f*x+e))*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(1/2)}/\sin(f*x+e)^4}$$

Maxima [A]

time = 0.27, size = 53, normalized size = 0.84

$$\frac{2 \left(5b^4 - \frac{18b^4}{\cos(fx+e)^2} + \frac{45b^4}{\cos(fx+e)^4} \right) b}{45 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] $-2/45*(5*b^4 - 18*b^4/\cos(f*x + e)^2 + 45*b^4/\cos(f*x + e)^4)*b/(f*(b/\cos(f*x + e))^{(9/2)})$

Fricas [A]

time = 0.39, size = 50, normalized size = 0.79

$$\frac{2 \left(5 \cos(fx + e)^5 - 18 \cos(fx + e)^3 + 45 \cos(fx + e) \right) \sqrt{\frac{b}{\cos(fx + e)}}}{45 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $-2/45*(5*\cos(f*x + e)^5 - 18*\cos(f*x + e)^3 + 45*\cos(f*x + e))*\sqrt{b/\cos(f*x + e))/f$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**5*(b*sec(f*x+e))**(1/2),x)

[Out] Timed out

Giac [A]

time = 3.40, size = 83, normalized size = 1.32

$$\frac{2 \left(5 \sqrt{b \cos(fx + e)} b^4 \cos(fx + e)^4 - 18 \sqrt{b \cos(fx + e)} b^4 \cos(fx + e)^2 + 45 \sqrt{b \cos(fx + e)} b^4 \right) \operatorname{sgn}(\cos(fx + e))}{45 b^4 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^5*(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out]
$$-2/45*(5*\sqrt{b*\cos(f*x + e)}*b^4*\cos(f*x + e)^4 - 18*\sqrt{b*\cos(f*x + e)}*b^4*\cos(f*x + e)^2 + 45*\sqrt{b*\cos(f*x + e)}*b^4)*\operatorname{sgn}(\cos(f*x + e))/(b^4*f)$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + f x)^5 \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^5*(b/cos(e + f*x))^(1/2),x)`

[Out] `int(sin(e + f*x)^5*(b/cos(e + f*x))^(1/2), x)`

3.373 $\int \sqrt{b \sec(e + fx)} \sin^3(e + fx) dx$

Optimal. Leaf size=41

$$\frac{2b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}}$$

[Out] $2/5*b^3/f/(b*\sec(f*x+e))^(5/2)-2*b/f/(b*\sec(f*x+e))^(1/2)$

Rubi [A]

time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 14}

$$\frac{2b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^3,x]

[Out] $(2*b^3)/(5*f*(b*Sec[e + f*x])^(5/2)) - (2*b)/(f*Sqrt[b*Sec[e + f*x]])$

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2702

Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \sqrt{b \sec(e + fx)} \sin^3(e + fx) dx &= \frac{b^3 \text{Subst}\left(\int \frac{-1 + \frac{x^2}{b^2}}{x^{7/2}} dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{b^3 \text{Subst}\left(\int \left(-\frac{1}{x^{7/2}} + \frac{1}{b^2 x^{3/2}}\right) dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{2b^3}{5f(b \sec(e + fx))^{5/2}} - \frac{2b}{f\sqrt{b \sec(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.12, size = 36, normalized size = 0.88

$$\frac{(-17 \cos(e + fx) + \cos(3(e + fx))) \sqrt{b \sec(e + fx)}}{10f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^3,x]**[Out]** ((-17*Cos[e + f*x] + Cos[3*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/(10*f)**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 496 vs. $2(35) = 70$.

time = 0.29, size = 497, normalized size = 12.12

method	result
default	$\frac{(-1 + \cos(fx + e))^2 \left(4(\cos^3(fx + e)) - 5 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \ln \left(-\frac{2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} - (\cos^2(fx + e) + 2 \cos(fx + e))}{\sin(fx + e)^2} \right) \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^3*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] 1/10/f*(-1+cos(f*x+e))^2*(4*cos(f*x+e)^3-5*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*cos(f*x+e)+5*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*cos(f*x+e)-5*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+5*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-20*cos(f*x+e))*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^4

Maxima [A]

time = 0.29, size = 37, normalized size = 0.90

$$\frac{2 \left(b^2 - \frac{5b^2}{\cos(fx+e)^2} \right) b}{5f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] 2/5*(b^2 - 5*b^2/cos(f*x + e)^2)*b/(f*(b/cos(f*x + e))^(5/2))

Fricas [A]

time = 0.36, size = 37, normalized size = 0.90

$$\frac{2(\cos(fx + e)^3 - 5 \cos(fx + e)) \sqrt{\frac{b}{\cos(fx + e)}}}{5f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 2/5*(cos(f*x + e)^3 - 5*cos(f*x + e))*sqrt(b/cos(f*x + e))/f

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**3*(b*sec(f*x+e))**(1/2),x)

[Out] Timed out

Giac [A]

time = 4.77, size = 57, normalized size = 1.39

$$\frac{2\left(\sqrt{b \cos(fx + e)} b^2 \cos(fx + e)^2 - 5 \sqrt{b \cos(fx + e)} b^2\right) \operatorname{sgn}(\cos(fx + e))}{5 b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] 2/5*(sqrt(b*cos(f*x + e))*b^2*cos(f*x + e)^2 - 5*sqrt(b*cos(f*x + e))*b^2)*sgn(cos(f*x + e))/(b^2*f)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + fx)^3 \sqrt{\frac{b}{\cos(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^3*(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^3*(b/cos(e + f*x))^(1/2), x)

3.374 $\int \sqrt{b \sec(e + fx)} \sin(e + fx) dx$

Optimal. Leaf size=18

$$-\frac{2b}{f\sqrt{b\sec(e+fx)}}$$

[Out] $-2*b/f/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 30}

$$-\frac{2b}{f\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x], x]$

[Out] $(-2*b)/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[x^{(m+1)}/(m+1), x] /; \text{FreeQ}[m, x] \ \&\& \ \text{NeQ}[m, -1]$

Rule 2702

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^{(n_.)}*((a_.)*\text{sec}[(e_.) + (f_.)*(x_)]^{(m_.)}, x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1+x^2/a^2)^{((n+1)/2)}, x], x, a*\text{Sec}[e+f*x]], x] /; \text{FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n+1)/2] \ \&\& \ !(\text{IntegerQ}[(m+1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int \sqrt{b \sec(e + fx)} \sin(e + fx) dx &= \frac{b \text{Subst}\left(\int \frac{1}{x^{3/2}} dx, x, b \sec(e + fx)\right)}{f} \\ &= -\frac{2b}{f\sqrt{b\sec(e+fx)}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 18, normalized size = 1.00

$$-\frac{2b}{f\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x],x]
```

```
[Out] (-2*b)/(f*Sqrt[b*Sec[e + f*x]])
```

Maple [A]

time = 0.11, size = 17, normalized size = 0.94

method	result	size
derivativedivides	$-\frac{2b}{f\sqrt{b\sec(fx+e)}}$	17
default	$-\frac{2b}{f\sqrt{b\sec(fx+e)}}$	17
risch	$-\frac{2\sqrt{2}\sqrt{\frac{be^{i(fx+e)}}{e^{2i(fx+e)}+1}}\cos(fx+e)}{f}$	41

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(f*x+e)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -2*b/f/(b*sec(f*x+e))^(1/2)
```

Maxima [A]

time = 0.29, size = 25, normalized size = 1.39

$$-\frac{2\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] -2*sqrt(b/cos(f*x + e))*cos(f*x + e)/f
```

Fricas [A]

time = 0.40, size = 25, normalized size = 1.39

$$-\frac{2\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] -2*sqrt(b/cos(f*x + e))*cos(f*x + e)/f
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \sin(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)*(b*sec(f*x+e))**(1/2),x)**[Out]** Integral(sqrt(b*sec(e + f*x))*sin(e + f*x), x)**Giac [A]**

time = 5.88, size = 24, normalized size = 1.33

$$-\frac{2 \sqrt{b \cos(fx + e)} \operatorname{sgn}(\cos(fx + e))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")**[Out]** -2*sqrt(b*cos(f*x + e))*sgn(cos(f*x + e))/f**Mupad [B]**

time = 0.20, size = 23, normalized size = 1.28

$$-\frac{2 \cos(e + fx) \sqrt{\frac{b}{\cos(e + fx)}}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)*(b/cos(e + f*x))^(1/2),x)**[Out]** -(2*cos(e + f*x)*(b/cos(e + f*x))^(1/2))/f

3.375 $\int \csc(e + fx) \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=58

$$\frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f}$$

[Out] arctan((b*sec(f*x+e))^(1/2)/b^(1/2))*b^(1/2)/f-arctanh((b*sec(f*x+e))^(1/2)/b^(1/2))*b^(1/2)/f

Rubi [A]

time = 0.03, antiderivative size = 58, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2702, 335, 304, 209, 212}

$$\frac{\sqrt{b} \text{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]*Sqrt[b*Sec[e + f*x]],x]

[Out] (Sqrt[b]*ArcTan[Sqrt[b*Sec[e + f*x]]/Sqrt[b]])/f - (Sqrt[b]*ArcTanh[Sqrt[b*Sec[e + f*x]]/Sqrt[b]])/f

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] :> With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
  ))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1
)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc(e + fx) \sqrt{b \sec(e + fx)} dx &= \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{bf} \\ &= \frac{2 \text{Subst}\left(\int \frac{x^2}{-1 + \frac{x^2}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{bf} \\ &= -\frac{b \text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} + \frac{b \text{Subst}\left(\int \frac{1}{b + x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} \\ &= \frac{\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} \end{aligned}$$

Mathematica [A]

time = 0.26, size = 73, normalized size = 1.26

$$\frac{\left(2 \tan^{-1}\left(\sqrt{\sec(e + fx)}\right) + \log\left(1 - \sqrt{\sec(e + fx)}\right) - \log\left(1 + \sqrt{\sec(e + fx)}\right)\right) \sqrt{b \sec(e + fx)}}{2f \sqrt{\sec(e + fx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]*Sqrt[b*Sec[e + f*x]], x]
```

```
[Out] ((2*ArcTan[Sqrt[Sec[e + f*x]]) + Log[1 - Sqrt[Sec[e + f*x]]] - Log[1 + Sqrt
[Sec[e + f*x]])*Sqrt[b*Sec[e + f*x]])/(2*f*Sqrt[Sec[e + f*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 168 vs. 2(46) = 92.

time = 0.20, size = 169, normalized size = 2.91

method	result
default	$\frac{\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)(-1+\cos(fx+e)) \left(\arctan\left(\frac{1}{2\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}\right) - \ln\left(-\frac{2\left(2(\cos^2(fx+e))\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}\right)}{2f \sin(fx+e)^2 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}\right)}{2f \sin(fx+e)^2 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}\right)}{2f \sin(fx+e)^2 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/f*(b/\cos(f*x+e))^{(1/2)}*\cos(f*x+e)*(-1+\cos(f*x+e))*(\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})-\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)/\sin(f*x+e)^2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}$$

Maxima [A]

time = 0.49, size = 75, normalized size = 1.29

$$b \left(\frac{2 \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right)}{\sqrt{b}} + \frac{\log\left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}}\right)}{\sqrt{b}} \right) / 2f$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out]
$$1/2*b*(2*\arctan(\sqrt{b}/\cos(f*x+e))/\sqrt{b})/\sqrt{b} + \log(-(\sqrt{b} - \sqrt{b/\cos(f*x+e)})/(\sqrt{b} + \sqrt{b/\cos(f*x+e)}))/\sqrt{b}/f$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 126 vs. 2(48) = 96.

time = 0.44, size = 265, normalized size = 4.57

$$\left[\frac{2\sqrt{-b} \arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}(\cos(fx+e)+1)}{2b}\right) + \sqrt{-b} \log\left(\frac{b\cos(fx+e)^2 - 4(\cos(fx+e)^2 - \cos(fx+e))\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}} - 6b\cos(fx+e) + b}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right)}{4f}, \dots, \frac{2\sqrt{b} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}(\cos(fx+e)-1)}{2\sqrt{b}}\right) - \sqrt{b} \log\left(\frac{b\cos(fx+e)^2 - 4(\cos(fx+e)^2 + \cos(fx+e))\sqrt{b}\sqrt{\frac{b}{\cos(fx+e)}} + 6b\cos(fx+e) + b}{\cos(fx+e)^2 - 2\cos(fx+e) + 1}\right)}{4f} \right]$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b) + sqrt(-b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)))/f, -1/4*(2*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b)) - sqrt(b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)))/f]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))*csc(e + f*x), x)

Giac [A]

time = 4.29, size = 64, normalized size = 1.10

$$\frac{b^2 \left(\frac{\arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b} - \frac{\arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{b^{3/2}} \right) \operatorname{sgn}(\cos(fx + e))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] b^2*(arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/(sqrt(-b)*b) - arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(3/2))*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\frac{b}{\cos(e + fx)}}}{\sin(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/sin(e + f*x),x)

[Out] int((b/cos(e + f*x))^(1/2)/sin(e + f*x), x)

3.376 $\int \csc^3(e + fx) \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=93

$$\frac{3\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{3\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2bf}$$

[Out] $-1/2*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(3/2)}/b/f+3/4*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})*b^{(1/2)}/f-3/4*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})*b^{(1/2)}/f$

Rubi [A]

time = 0.05, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2702, 294, 335, 304, 209, 212}

$$\frac{3\sqrt{b} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2bf} - \frac{3\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^3*Sqrt[b*Sec[e + f*x]],x]`

[Out] $(3*\operatorname{Sqrt}[b]*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]])/(4*f) - (3*\operatorname{Sqrt}[b]*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]])/(4*f) - (\operatorname{Cot}[e + f*x]^2*(b*\operatorname{Sec}[e + f*x])^{(3/2)})/(2*b*f)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 294

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1
)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned} \int \csc^3(e + fx) \sqrt{b \sec(e + fx)} dx &= \frac{\text{Subst}\left(\int \frac{x^{5/2}}{(-1 + \frac{x^2}{b^2})^2} dx, x, b \sec(e + fx)\right)}{b^3 f} \\ &= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2bf} + \frac{3 \text{Subst}\left(\int \frac{\sqrt{x}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{4bf} \\ &= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2bf} + \frac{3 \text{Subst}\left(\int \frac{x^2}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{2bf} \\ &= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2bf} - \frac{(3b) \text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4f} \\ &= \frac{3\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{3\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} \end{aligned}$$

Mathematica [A]

time = 0.45, size = 95, normalized size = 1.02

$$\frac{\left(-6 \tan^{-1}\left(\sqrt{\sec(e+fx)}\right) - 3 \log\left(1 - \sqrt{\sec(e+fx)}\right) + 3 \log\left(1 + \sqrt{\sec(e+fx)}\right) + \frac{4 \csc^2(e+fx)}{\sqrt{\sec(e+fx)}}\right) \sqrt{b \sec(e+fx)}}{8f \sqrt{\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3*Sqrt[b*Sec[e + f*x]],x]

[Out] -1/8*((-6*ArcTan[Sqrt[Sec[e + f*x]]] - 3*Log[1 - Sqrt[Sec[e + f*x]]] + 3*Log[1 + Sqrt[Sec[e + f*x]]] + (4*Csc[e + f*x]^2)/Sqrt[Sec[e + f*x]])*Sqrt[b*Sec[e + f*x]]/(f*Sqrt[Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 602 vs. 2(73) = 146.

time = 0.27, size = 603, normalized size = 6.48

method	result
default	$\frac{(-1+\cos(fx+e)) \left(8(\cos^2(fx+e)) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 16 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} - 3(\cos^2(fx+e)) \arctan \left(\frac{\cos(fx+e)}{2 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}} \right) \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/8/f*(-1+cos(f*x+e))*(8*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+16*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)-3*cos(f*x+e)^2*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))-cos(f*x+e)^2*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+4*cos(f*x+e)^2*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+8*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+4*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-4*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+3*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))+ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-4*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2))*cos(f*x+e)*(b/cos(f*x+e))^(1/2)/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)/sin(f*x+e)^4

Maxima [A]

time = 0.48, size = 111, normalized size = 1.19

$$b \left(\frac{4 \left(\frac{b}{\cos(fx+e)} \right)^{3/2}}{b^2 - \frac{b^2}{\cos(fx+e)^2}} + \frac{6 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right)}{\sqrt{b}} + \frac{3 \log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right)}{\sqrt{b}} \right) / 8f$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] 1/8*b*(4*(b/cos(f*x + e))^(3/2)/(b^2 - b^2/cos(f*x + e)^2) + 6*arctan(sqrt(b/cos(f*x + e))/sqrt(b))/sqrt(b) + 3*log(-(sqrt(b) - sqrt(b/cos(f*x + e)))/(sqrt(b) + sqrt(b/cos(f*x + e))))/sqrt(b))/f

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 184 vs. 2(77) = 154.

time = 0.48, size = 382, normalized size = 4.11

$$\frac{6(\cos(fx+e)^2-1)\sqrt{-b}\arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + 3(\cos(fx+e)^2-1)\sqrt{-b}\log\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}} - \sqrt{b}}{\sqrt{\frac{b}{\cos(fx+e)}} + \sqrt{b}}\right) + 8\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\cos(fx+e) - 8(\cos(fx+e)^2-1)\sqrt{-b}\arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) - 3(\cos(fx+e)^2-1)\sqrt{-b}\log\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}} - \sqrt{b}}{\sqrt{\frac{b}{\cos(fx+e)}} + \sqrt{b}}\right) - 8\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)}{8f(\cos(fx+e)^2-f)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] [1/16*(6*(cos(f*x + e)^2 - 1)*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e)))*(cos(f*x + e) + 1)/b) + 3*(cos(f*x + e)^2 - 1)*sqrt(-b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 8*sqrt(b/cos(f*x + e))*cos(f*x + e)/(f*cos(f*x + e)^2 - f), -1/16*(6*(cos(f*x + e)^2 - 1)*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e)))*(cos(f*x + e) - 1)/sqrt(b)) - 3*(cos(f*x + e)^2 - 1)*sqrt(b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)) - 8*sqrt(b/cos(f*x + e))*cos(f*x + e)/(f*cos(f*x + e)^2 - f)]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \csc^3(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3*(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))*csc(e + f*x)**3, x)

Giac [A]

time = 4.65, size = 103, normalized size = 1.11

$$b^4 \left(\frac{2\sqrt{b\cos(fx+e)}}{(b^2\cos(fx+e)^2-b^2)b^2} + \frac{3\arctan\left(\frac{\sqrt{b\cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b}b^3} - \frac{3\arctan\left(\frac{\sqrt{b\cos(fx+e)}}{\sqrt{b}}\right)}{b^{\frac{7}{2}}} \right) \operatorname{sgn}(\cos(fx+e))$$

$4f$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] 1/4*b^4*(2*sqrt(b*cos(f*x + e))/((b^2*cos(f*x + e)^2 - b^2)*b^2) + 3*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b)*b^3 - 3*arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(7/2))*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(e+fx)}}}{\sin(e+fx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^3,x)

[Out] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^3, x)

3.377 $\int \csc^5(e + fx) \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=123

$$\frac{21\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{21\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{7 \cot^2(e + fx)(b \sec(e + fx))^{3/2}}{16bf}$$

[Out] $-7/16*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(3/2)}/b/f-1/4*\cot(f*x+e)^4*(b*\sec(f*x+e))^{(7/2)}/b^3/f+21/32*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})*b^{(1/2)}/f-21/32*\arctan(\tanh((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)}))*b^{(1/2)}/f$

Rubi [A]

time = 0.06, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2702, 294, 335, 304, 209, 212}

$$\frac{21\sqrt{b} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{7/2}}{4b^3f} - \frac{7 \cot^2(e + fx)(b \sec(e + fx))^{3/2}}{16bf} - \frac{21\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^5*Sqrt[b*Sec[e + f*x]],x]`

[Out] $(21*\sqrt{b}*\operatorname{ArcTan}[\sqrt{b*\operatorname{Sec}[e + f*x]}/\sqrt{b}])/(32*f) - (21*\sqrt{b}*\operatorname{ArcTanh}[\sqrt{b*\operatorname{Sec}[e + f*x]}/\sqrt{b}])/(32*f) - (7*\operatorname{Cot}[e + f*x]^2*(b*\operatorname{Sec}[e + f*x])^{(3/2)})/(16*b*f) - (\operatorname{Cot}[e + f*x]^4*(b*\operatorname{Sec}[e + f*x])^{(7/2)})/(4*b^3*f)$

Rule 209

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 294

`Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]`

Rule 304

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 335

```
Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)]^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1
)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \csc^5(e + fx) \sqrt{b \sec(e + fx)} \, dx &= \frac{\text{Subst}\left(\int \frac{x^{9/2}}{(-1 + \frac{x^2}{b^2})^3} \, dx, x, b \sec(e + fx)\right)}{b^5 f} \\
&= -\frac{\cot^4(e + fx)(b \sec(e + fx))^{7/2}}{4b^3 f} + \frac{7 \text{Subst}\left(\int \frac{x^{5/2}}{(-1 + \frac{x^2}{b^2})^2} \, dx, x, b \sec(e + fx)\right)}{8b^3 f} \\
&= -\frac{7 \cot^2(e + fx)(b \sec(e + fx))^{3/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{7/2}}{4b^3 f} + \\
&= -\frac{7 \cot^2(e + fx)(b \sec(e + fx))^{3/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{7/2}}{4b^3 f} + \\
&= -\frac{7 \cot^2(e + fx)(b \sec(e + fx))^{3/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{7/2}}{4b^3 f} - \\
&= \frac{21\sqrt{b} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{21\sqrt{b} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f}
\end{aligned}$$

Mathematica [A]

time = 0.63, size = 107, normalized size = 0.87

$$\frac{b(-28 \csc^2(e+fx) - 16 \csc^4(e+fx) + 42 \tan^{-1}(\sqrt{\sec(e+fx)}) \sqrt{\sec(e+fx)} + 21(\log(1 - \sqrt{\sec(e+fx)}) - \log(1 + \sqrt{\sec(e+fx)})) \sqrt{\sec(e+fx)})}{64f\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^5*Sqrt[b*Sec[e + f*x]],x]

```
[Out] (b*(-28*Csc[e + f*x]^2 - 16*Csc[e + f*x]^4 + 42*ArcTan[Sqrt[Sec[e + f*x]]]*
Sqrt[Sec[e + f*x]] + 21*(Log[1 - Sqrt[Sec[e + f*x]]] - Log[1 + Sqrt[Sec[e +
f*x]]])*Sqrt[Sec[e + f*x]])/(64*f*Sqrt[b*Sec[e + f*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1088 vs. 2(99) = 198.

time = 0.27, size = 1089, normalized size = 8.85

method	result	size
default	Expression too large to display	1089

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^5*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

```
[Out] -1/64/f*(72*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*cos(f*x+e)^3+56*cos(f*x+e)
^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)-104*cos(f*x+e)*(-cos(f*x+e)/(cos(f*
x+e)+1)^2)^(3/2)-11*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)
)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f
*x+e)^2)*cos(f*x+e)^3-21*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))*c
os(f*x+e)^3+32*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-c
os(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+
e)^2)*cos(f*x+e)^3-88*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+44*cos(f*x+e)^2*
(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+11*cos(f*x+e)^2*ln(-(2*cos(f*x+e)^2*(-
cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)
/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+21*cos(f*x+e)^2*arctan(1/2/(-cos(
f*x+e)/(cos(f*x+e)+1)^2)^(1/2))-32*cos(f*x+e)^2*ln(-2*(2*cos(f*x+e)^2*(-cos
(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(c
os(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-88*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+
e)+1)^2)^(1/2)+11*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-
cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x
+e)^2)*cos(f*x+e)+21*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))*cos(f
*x+e)-32*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x
+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*
cos(f*x+e)+44*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-11*ln(-(2*cos(f*x+e)^2*(-
cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)
)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-21*arctan(1/2/(-cos(f*x+e)/(cos(
```

$f*x+e)+1)^2)^{(1/2))+32*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)-1)/\sin(f*x+e)^2})*\cos(f*x+e)*(b/\cos(f*x+e))^{(1/2)/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)/\sin(f*x+e)^4}$

Maxima [A]

time = 0.54, size = 145, normalized size = 1.18

$$b \left(\frac{42 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right)}{\sqrt{b}} + \frac{21 \log \left(-\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right)}{\sqrt{b}} + \frac{4 \left(7b^2 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} - 11 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{7}{2}} \right)}{b^4 - \frac{2b^4}{\cos(fx+e)^2} + \frac{b^4}{\cos(fx+e)^4}} \right) / 64f$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] $\frac{1}{64} * b * \left(\frac{42 * \arctan(\sqrt{b/\cos(f*x+e)})/\sqrt{b}}{\sqrt{b}} + 21 * \log\left(-\frac{\sqrt{b} - \sqrt{b/\cos(f*x+e)}}{\sqrt{b} + \sqrt{b/\cos(f*x+e)}}\right) / \sqrt{b} + \frac{4 * (7 * b^2 * (b/\cos(f*x+e))^{3/2} - 11 * (b/\cos(f*x+e))^{7/2})}{b^4 - 2 * b^4 / \cos(f*x+e)^2 + b^4 / \cos(f*x+e)^4} \right) / f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 230 vs. 2(105) = 210.

time = 0.50, size = 474, normalized size = 3.85

$$\frac{\frac{42 \arctan\left(\frac{\sqrt{b/\cos(fx+e)}}{\sqrt{b}}\right)}{\sqrt{b}} + 21 \log\left(-\frac{\sqrt{b} - \sqrt{b/\cos(fx+e)}}{\sqrt{b} + \sqrt{b/\cos(fx+e)}}\right) / \sqrt{b} + \frac{4 \left(7b^2 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} - 11 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{7}{2}} \right)}{b^4 - \frac{2b^4}{\cos(fx+e)^2} + \frac{b^4}{\cos(fx+e)^4}}}{64f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $\frac{1}{128} * (42 * (\cos(f*x+e)^4 - 2 * \cos(f*x+e)^2 + 1) * \sqrt{-b} * \arctan(1/2 * \sqrt{-b} * \sqrt{b/\cos(f*x+e)}) * (\cos(f*x+e) + 1) / b + 21 * (\cos(f*x+e)^4 - 2 * \cos(f*x+e)^2 + 1) * \sqrt{-b} * \log((b * \cos(f*x+e)^2 - 4 * (\cos(f*x+e)^2 - \cos(f*x+e)) * \sqrt{-b} * \sqrt{b/\cos(f*x+e)} - 6 * b * \cos(f*x+e) + b) / (\cos(f*x+e)^2 + 2 * \cos(f*x+e) + 1)) + 8 * (7 * \cos(f*x+e)^3 - 11 * \cos(f*x+e)) * \sqrt{b/\cos(f*x+e)}) / (f * \cos(f*x+e)^4 - 2 * f * \cos(f*x+e)^2 + f), -1/128 * (42 * (\cos(f*x+e)^4 - 2 * \cos(f*x+e)^2 + 1) * \sqrt{b} * \arctan(1/2 * \sqrt{b/\cos(f*x+e)}) * (\cos(f*x+e) - 1) / \sqrt{b}) - 21 * (\cos(f*x+e)^4 - 2 * \cos(f*x+e)^2 + 1) * \sqrt{b} * \log((b * \cos(f*x+e)^2 - 4 * (\cos(f*x+e)^2 - \cos(f*x+e)) * \sqrt{b} * \sqrt{b/\cos(f*x+e)} - 6 * b * \cos(f*x+e) + b) / (\cos(f*x+e)^2 + 2 * \cos(f*x+e) + 1)) + 8 * (7 * \cos(f*x+e)^3 - 11 * \cos(f*x+e)) * \sqrt{b/\cos(f*x+e)}) / (f * \cos(f*x+e)^4 - 2 * f * \cos(f*x+e)^2 + f)$

$\sqrt{b} \log((b \cos(fx + e))^2 - 4(\cos(fx + e))^2 + \cos(fx + e)) \sqrt{b} \sqrt{b/\cos(fx + e)} + 6b \cos(fx + e) + b)/(\cos(fx + e)^2 - 2\cos(fx + e) + 1) - 8(7\cos(fx + e)^3 - 11\cos(fx + e)) \sqrt{b/\cos(fx + e)})/(f \cos(fx + e)^4 - 2f \cos(fx + e)^2 + f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \csc^5(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**5*(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))*csc(e + f*x)**5, x)

Giac [A]

time = 4.77, size = 134, normalized size = 1.09

$$b^6 \left(\frac{21 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^5} - \frac{21 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{b}}\right)}{b^{11/2}} + \frac{2 \left(7 \sqrt{b \cos(fx+e)} b^2 \cos(fx+e)^2 - 11 \sqrt{b \cos(fx+e)} b^2\right)}{(b^2 \cos(fx+e)^2 - b^2)^2 b^4} \right) \operatorname{sgn}(\cos(fx+e))$$

32 f

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] 1/32*b^6*(21*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b)*b^5) - 21*arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(11/2) + 2*(7*sqrt(b*cos(f*x + e))*b^2*cos(f*x + e)^2 - 11*sqrt(b*cos(f*x + e))*b^2)/((b^2*cos(f*x + e)^2 - b^2)^2*b^4)*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(e + fx)}}}{\sin(e + fx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^5,x)

[Out] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^5, x)

3.378 $\int \sqrt{b \sec(e + fx)} \sin^6(e + fx) dx$

Optimal. Leaf size=123

$$\frac{80\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx)|2\right) \sqrt{b \sec(e+fx)}}{77f} - \frac{40b \sin(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{20b \sin^3(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^5(e+fx)}{11f \sqrt{b \sec(e+fx)}}$$

[Out] $-40/77*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}-20/77*b*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(1/2)}-2/11*b*\sin(f*x+e)^5/f/(b*\sec(f*x+e))^{(1/2)}+80/77*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.10, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2707, 3856, 2720}

$$-\frac{2b \sin^5(e+fx)}{11f \sqrt{b \sec(e+fx)}} - \frac{20b \sin^3(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{40b \sin(e+fx)}{77f \sqrt{b \sec(e+fx)}} + \frac{80\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx)|2\right) \sqrt{b \sec(e+fx)}}{77f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x]^6,x]$

[Out] $(80*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(77*f) - (40*b*\text{Sin}[e + f*x])/(77*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (20*b*\text{Sin}[e + f*x]^3)/(77*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (2*b*\text{Sin}[e + f*x]^5)/(11*f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2707

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m+1)}*((b*\text{Sec}[e + f*x])^{(n-1)})/(a*f^{(m+n)})], x] + \text{Dist}[(m+1)/(a^2*(m+n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m+2)}*(b*\text{Sec}[e + f*x])^n, x], x] /;$ FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /;$ FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sqrt{b \sec(e+fx)} \sin^6(e+fx) dx &= -\frac{2b \sin^5(e+fx)}{11f \sqrt{b \sec(e+fx)}} + \frac{10}{11} \int \sqrt{b \sec(e+fx)} \sin^4(e+fx) dx \\
&= -\frac{20b \sin^3(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^5(e+fx)}{11f \sqrt{b \sec(e+fx)}} + \frac{60}{77} \int \sqrt{b \sec(e+fx)} \sin^2(e+fx) dx \\
&= -\frac{40b \sin(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{20b \sin^3(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^5(e+fx)}{11f \sqrt{b \sec(e+fx)}} \\
&= -\frac{40b \sin(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{20b \sin^3(e+fx)}{77f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^5(e+fx)}{11f \sqrt{b \sec(e+fx)}} \\
&= \frac{80 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{77f} - \frac{40b \sin(e+fx)}{77f \sqrt{b \sec(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 73, normalized size = 0.59

$$\frac{\sqrt{b \sec(e+fx)} \left(1280 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) - 435 \sin(2(e+fx)) + 68 \sin(4(e+fx)) - 7 \sin(6(e+fx)) \right)}{1232f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^6,x]**[Out]** (Sqrt[b*Sec[e + f*x]]*(1280*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] - 435*Sin[2*(e + f*x)] + 68*Sin[4*(e + f*x)] - 7*Sin[6*(e + f*x)]))/(1232*f)**Maple [C]** Result contains complex when optimal does not.

time = 0.44, size = 165, normalized size = 1.34

method	result
default	$ -\frac{2(-1+\cos(fx+e)) \left(7(\cos^6(fx+e))+40i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) - 7(\cos^5(fx+e)) \right)}{77f \sin(fx+e)^3} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^6*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
[Out] -2/77/f*(-1+cos(f*x+e))*(7*cos(f*x+e)^6+40*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)-7*cos(f*x+e)^5-24*cos(f*x+e)^4+24*cos(f*x+e)^3+37*cos(f*x+e)^2-37*cos(f*x+e))*cos(f*x+e)+1)^2*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^3

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^6*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))*sin(f*x + e)^6, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 115, normalized size = 0.93

$$\frac{2 \left((7 \cos(fx+e)^5 - 24 \cos(fx+e)^3 + 37 \cos(fx+e)) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) + 20i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e)) - 20i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)) \right)}{7f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^6*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] -2/77*((7*cos(f*x + e)^5 - 24*cos(f*x + e)^3 + 37*cos(f*x + e))*sqrt(b/cos(f*x + e))*sin(f*x + e) + 20*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) - 20*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)))/f
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)**6*(b*sec(f*x+e))**(1/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^6*(b*sec(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))*sin(f*x + e)^6, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^6 \sqrt{\frac{b}{\cos(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^6*(b/cos(e + f*x))^(1/2),x)
```

```
[Out] int(sin(e + f*x)^6*(b/cos(e + f*x))^(1/2), x)
```

3.379 $\int \sqrt{b \sec(e + fx)} \sin^4(e + fx) dx$

Optimal. Leaf size=95

$$\frac{8\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{7f} - \frac{4b \sin(e+fx)}{7f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^3(e+fx)}{7f \sqrt{b \sec(e+fx)}}$$

[Out] $-4/7*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}-2/7*b*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(1/2)}+8/7*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2707, 3856, 2720}

$$-\frac{2b \sin^3(e+fx)}{7f \sqrt{b \sec(e+fx)}} - \frac{4b \sin(e+fx)}{7f \sqrt{b \sec(e+fx)}} + \frac{8\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{7f}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^4,x]

[Out] $(8*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(7*f) - (4*b*\text{Sin}[e + f*x])/(7*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (2*b*\text{Sin}[e + f*x]^3)/(7*f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2707

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sqrt{b \sec(e+fx)} \sin^4(e+fx) dx &= -\frac{2b \sin^3(e+fx)}{7f \sqrt{b \sec(e+fx)}} + \frac{6}{7} \int \sqrt{b \sec(e+fx)} \sin^2(e+fx) dx \\
&= -\frac{4b \sin(e+fx)}{7f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^3(e+fx)}{7f \sqrt{b \sec(e+fx)}} + \frac{4}{7} \int \sqrt{b \sec(e+fx)} dx \\
&= -\frac{4b \sin(e+fx)}{7f \sqrt{b \sec(e+fx)}} - \frac{2b \sin^3(e+fx)}{7f \sqrt{b \sec(e+fx)}} + \frac{1}{7} \left(4 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)} \right) \\
&= \frac{8 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{7f} - \frac{4b \sin(e+fx)}{7f \sqrt{b \sec(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 61, normalized size = 0.64

$$\frac{\sqrt{b \sec(e+fx)} \left(32 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) - 10 \sin(2(e+fx)) + \sin(4(e+fx)) \right)}{28f}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^4,x]`

```
[Out] (Sqrt[b*Sec[e + f*x]]*(32*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] - 10
*Sin[2*(e + f*x)] + Sin[4*(e + f*x)]))/(28*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.28, size = 143, normalized size = 1.51

method	result
default	$ -\frac{2(-1+\cos(fx+e)) \left(4i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) - (\cos^4(fx+e) + \cos^3(fx+e)) \right)}{7f \sin(fx+e)^3} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^4*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] -2/7/f*(-1+cos(f*x+e))*(4*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)
)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)-cos(f*x+e)
^4+cos(f*x+e)^3+3*cos(f*x+e)^2-3*cos(f*x+e))*(cos(f*x+e)+1)^2*(b/cos(f*x+e)
)^(1/2)/sin(f*x+e)^3
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))*sin(f*x + e)^4, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.13, size = 102, normalized size = 1.07

$$\frac{2 \left((\cos(fx + e)^3 - 3 \cos(fx + e)) \sqrt{\frac{b}{\cos(fx + e)}} \sin(fx + e) - 2i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx + e) + i \sin(fx + e)) + 2i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx + e) - i \sin(fx + e)) \right)}{7f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 2/7*((cos(f*x + e)^3 - 3*cos(f*x + e))*sqrt(b/cos(f*x + e))*sin(f*x + e) - 2*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) + 2*I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)))/f

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \sin^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**4*(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))*sin(e + f*x)**4, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*sin(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^4 \sqrt{\frac{b}{\cos(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4*(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^4*(b/cos(e + f*x))^(1/2), x)

3.380 $\int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx$

Optimal. Leaf size=67

$$\frac{4\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} - \frac{2b \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}}$$

[Out] $-2/3*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}+4/3*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2707, 3856, 2720}

$$\frac{4\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} - \frac{2b \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x]^2, x]$

[Out] $(4*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(3*f) - (2*b*\text{Sin}[e + f*x])/(3*f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2707

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] := \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(a*f*(m + n))), x] + \text{Dist}[(m + 1)/(a^2*(m + n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.)^{(n_.)}, x_Symbol] := \text{Dist}[(b*\text{Csc}[c + d*x])^{(n)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx &= -\frac{2b \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{2}{3} \int \sqrt{b \sec(e + fx)} dx \\
&= -\frac{2b \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{1}{3} \left(2 \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{1}{\sqrt{\cos}} \\
&= \frac{4 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} - \frac{2b \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 51, normalized size = 0.76

$$\frac{\sqrt{b \sec(e + fx)} \left(-4 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) + \sin(2(e + fx)) \right)}{3f}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^2,x]``[Out] -1/3*(Sqrt[b*Sec[e + f*x]]*(-4*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] + Sin[2*(e + f*x)]))/f`**Maple [C]** Result contains complex when optimal does not.

time = 0.24, size = 123, normalized size = 1.84

method	result
default	$ -\frac{2(-1+\cos(fx+e)) \left(2i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) + \cos^2(fx+e) - \cos(fx+e) \right)}{3f \sin(fx+e)^3} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^2*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)``[Out] -2/3/f*(-1+cos(f*x+e))*(2*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)+cos(f*x+e)^2-cos(f*x+e))*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^3`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(sin(f*x+e)^2*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] integrate(sqrt(b*sec(f*x + e))*sin(f*x + e)^2, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.12, size = 90, normalized size = 1.34

$$\frac{2 \left(\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e) \sin(fx+e) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e)) - i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)) \right)}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] -2/3*(sqrt(b/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) - I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)))/f

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**2*(b*sec(f*x+e))^(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))*sin(e + f*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*sin(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^2 \sqrt{\frac{b}{\cos(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2*(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^2*(b/cos(e + f*x))^(1/2), x)

3.381 $\int \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=38

$$\frac{2\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{f}$$

[Out] $2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3856, 2720}

$$\frac{2\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{f}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[e + f*x]],x]

[Out] $(2*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/f$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{b \sec(e + fx)} dx &= \left(\sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{1}{\sqrt{\cos(e + fx)}} dx \\ &= \frac{2\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{f} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 38, normalized size = 1.00

$$\frac{2\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]],x]

[Out] (2*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*Sqrt[b*Sec[e + f*x]])/f

Maple [C] Result contains complex when optimal does not.

time = 0.28, size = 98, normalized size = 2.58

method	result	size
default	$-\frac{2i\sqrt{\frac{b}{\cos(fx+e)}}(-1+\cos(fx+e))\sqrt{\frac{1}{\cos(fx+e)+1}}\sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}}\text{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)},i\right)(\cos(fx+e)+1)^2}{f\sin(fx+e)^2}$	98

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] $-2*I/f*(b/\cos(f*x+e))^{1/2}*(-1+\cos(f*x+e))*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*(\cos(f*x+e)+1)^2/\sin(f*x+e)^2$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 61, normalized size = 1.61

$$\frac{-i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{2}\sqrt{b}\text{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $(-I*\text{sqrt}(2)*\text{sqrt}(b)*\text{weierstrassPInverse}(-4,0,\cos(f*x+e)+I*\sin(f*x+e)) + I*\text{sqrt}(2)*\text{sqrt}(b)*\text{weierstrassPInverse}(-4,0,\cos(f*x+e)-I*\sin(f*x+e)))/f$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e)), x)

Mupad [B]

time = 0.56, size = 35, normalized size = 0.92

$$\frac{2 \sqrt{\cos(e + f x)} \sqrt{\frac{b}{\cos(e + f x)}} F\left(\frac{e}{2} + \frac{f x}{2} \middle| 2\right)}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2),x)

[Out] (2*cos(e + f*x)^(1/2)*(b/cos(e + f*x))^(1/2)*ellipticF(e/2 + (f*x)/2, 2))/f

3.382 $\int \csc^2(e + fx) \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=62

$$-\frac{b \csc(e + fx)}{f \sqrt{b \sec(e + fx)}} + \frac{\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{f}$$

[Out] $-b \csc(fx+e)/f/(b \sec(fx+e))^{(1/2)}+(\cos(1/2*fx+1/2*e)^2)^{(1/2)}/\cos(1/2*fx+1/2*e)*\text{EllipticF}(\sin(1/2*fx+1/2*e),2^{(1/2)})*\cos(fx+e)^{(1/2)}*(b \sec(fx+e))^{(1/2)}/f$

Rubi [A]

time = 0.04, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2705, 3856, 2720}

$$\frac{\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{f} - \frac{b \csc(e + fx)}{f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^2*Sqrt[b*Sec[e + f*x]],x]`

[Out] $-\left(\frac{b \csc[e + f*x]}{f \sqrt{b \sec[e + f*x]}}\right) + \left(\frac{\sqrt{\cos[e + f*x]} \text{EllipticF}\left[\frac{e + f*x}{2}, 2\right] \sqrt{b \sec[e + f*x]}}{f}\right)$

Rule 2705

`Int[(csc[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sec[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

Rule 2720

`Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3856

`Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned}
\int \csc^2(e + fx) \sqrt{b \sec(e + fx)} \, dx &= -\frac{b \csc(e + fx)}{f \sqrt{b \sec(e + fx)}} + \frac{1}{2} \int \sqrt{b \sec(e + fx)} \, dx \\
&= -\frac{b \csc(e + fx)}{f \sqrt{b \sec(e + fx)}} + \frac{1}{2} \left(\sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{1}{\sqrt{\cos(e + fx)}} \, dx \\
&= -\frac{b \csc(e + fx)}{f \sqrt{b \sec(e + fx)}} + \frac{\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{f}
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 47, normalized size = 0.76

$$\frac{\left(-\cot(e + fx) + \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right)\right) \sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2*Sqrt[b*Sec[e + f*x]],x]``[Out] ((-Cot[e + f*x] + Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2])*Sqrt[b*Sec[e + f*x]])/f`**Maple [C]** Result contains complex when optimal does not.

time = 0.24, size = 184, normalized size = 2.97

method	result
default	$\frac{(-1 + \cos(fx + e))^2 \left(i \sqrt{\frac{1}{\cos(fx + e) + 1}} \sqrt{\frac{\cos(fx + e)}{\cos(fx + e) + 1}} \operatorname{EllipticF}\left(\frac{i(-1 + \cos(fx + e))}{\sin(fx + e)}, i\right) \sin(fx + e) \cos(fx + e) + i \sqrt{\frac{1}{\cos(fx + e) + 1}} \right)}{f \sin(fx + e)^5}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/f*(-1+cos(f*x+e))^2*(I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-cos(f*x+e))*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^5
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*sec(f*x + e))*csc(f*x + e)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.11, size = 107, normalized size = 1.73

$$\frac{-i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))-2\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)}{2f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] `1/2*(-I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) + I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)) - 2*sqrt(b/cos(f*x + e))*cos(f*x + e))/(f*sin(f*x + e))`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b\sec(e+fx)} \csc^2(e+fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**2*(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(sqrt(b*sec(e + f*x))*csc(e + f*x)**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(b*sec(f*x + e))*csc(f*x + e)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\frac{b}{\cos(e+fx)}}}{\sin(e+fx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^2,x)`

[Out] `int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^2, x)`

3.383 $\int \csc^4(e + fx) \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=95

$$-\frac{5b \csc(e + fx)}{6f \sqrt{b \sec(e + fx)}} - \frac{b \csc^3(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{5 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{6f}$$

[Out] $-5/6*b*\csc(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}-1/3*b*\csc(f*x+e)^3/f/(b*\sec(f*x+e))^{(1/2)}+5/6*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2705, 3856, 2720}

$$-\frac{b \csc^3(e + fx)}{3f \sqrt{b \sec(e + fx)}} - \frac{5b \csc(e + fx)}{6f \sqrt{b \sec(e + fx)}} + \frac{5 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{6f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^4*Sqrt[b*Sec[e + f*x]],x]`

[Out] $(-5*b*Csc[e + f*x])/(6*f*Sqrt[b*Sec[e + f*x]]) - (b*Csc[e + f*x]^3)/(3*f*Sqrt[b*Sec[e + f*x]]) + (5*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*Sqrt[b*Sec[e + f*x]])/(6*f)$

Rule 2705

`Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3856

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned}
\int \csc^4(e+fx) \sqrt{b \sec(e+fx)} dx &= -\frac{b \csc^3(e+fx)}{3f \sqrt{b \sec(e+fx)}} + \frac{5}{6} \int \csc^2(e+fx) \sqrt{b \sec(e+fx)} dx \\
&= -\frac{5b \csc(e+fx)}{6f \sqrt{b \sec(e+fx)}} - \frac{b \csc^3(e+fx)}{3f \sqrt{b \sec(e+fx)}} + \frac{5}{12} \int \sqrt{b \sec(e+fx)} dx \\
&= -\frac{5b \csc(e+fx)}{6f \sqrt{b \sec(e+fx)}} - \frac{b \csc^3(e+fx)}{3f \sqrt{b \sec(e+fx)}} + \frac{1}{12} \left(5 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)} \right. \\
&= -\frac{5b \csc(e+fx)}{6f \sqrt{b \sec(e+fx)}} - \frac{b \csc^3(e+fx)}{3f \sqrt{b \sec(e+fx)}} + \frac{5 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right)}{12}
\end{aligned}$$

Mathematica [A]

time = 0.17, size = 63, normalized size = 0.66

$$\frac{\left(-\cot(e+fx)(5+2\csc^2(e+fx))+5\sqrt{\cos(e+fx)}F\left(\frac{1}{2}(e+fx)\mid 2\right)\right)\sqrt{b\sec(e+fx)}}{6f}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^4*Sqrt[b*Sec[e + f*x]],x]`

```
[Out] ((-(Cot[e + f*x]*(5 + 2*Csc[e + f*x]^2)) + 5*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2])*Sqrt[b*Sec[e + f*x]])/(6*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.41, size = 335, normalized size = 3.53

method	result
default	$ -\frac{(-1+\cos(fx+e))^2 \left(5i \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^3(fx+e) \sin(fx+e) + 5i \operatorname{EllipticF}(\dots)) \right)}{6f} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^4*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/6/f*(-1+cos(f*x+e))^2*(5*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^3*sin(f*x+e)+5*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2*sin(f*x+e)-5*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-5*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e))
```

)-5*cos(f*x+e)^3+7*cos(f*x+e))*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^7

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))*csc(f*x + e)^4, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.14, size = 161, normalized size = 1.69

$$\frac{5\sqrt{2}(i\cos(fx+e)^2-i)\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+5\sqrt{2}(-i\cos(fx+e)^2+i)\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))+2(5\cos(fx+e)^3-7\cos(fx+e))\sqrt{\frac{b}{\cos(fx+e)}}}{12(f\cos(fx+e)^2-f)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] -1/12*(5*sqrt(2)*(I*cos(f*x + e)^2 - I)*sqrt(b)*sin(f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) + 5*sqrt(2)*(-I*cos(f*x + e)^2 + I)*sqrt(b)*sin(f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)) + 2*(5*cos(f*x + e)^3 - 7*cos(f*x + e))*sqrt(b/cos(f*x + e)))/((f*cos(f*x + e)^2 - f)*sin(f*x + e))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{b \sec(e + fx)} \csc^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**4*(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))*csc(e + f*x)**4, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*csc(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(e + f x)}}}{\sin(e + f x)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^4,x)

[Out] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^4, x)

3.384 $\int \csc^6(e + fx) \sqrt{b \sec(e + fx)} dx$

Optimal. Leaf size=123

$$\frac{3b \csc(e + fx)}{4f \sqrt{b \sec(e + fx)}} - \frac{3b \csc^3(e + fx)}{10f \sqrt{b \sec(e + fx)}} - \frac{b \csc^5(e + fx)}{5f \sqrt{b \sec(e + fx)}} + \frac{3 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{4f}$$

[Out] $-3/4*b*\csc(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}-3/10*b*\csc(f*x+e)^3/f/(b*\sec(f*x+e))^{(1/2)}-1/5*b*\csc(f*x+e)^5/f/(b*\sec(f*x+e))^{(1/2)}+3/4*(\cos(1/2*f*x+1/2*e))^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.10, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2705, 3856, 2720}

$$\frac{b \csc^5(e + fx)}{5f \sqrt{b \sec(e + fx)}} - \frac{3b \csc^3(e + fx)}{10f \sqrt{b \sec(e + fx)}} - \frac{3b \csc(e + fx)}{4f \sqrt{b \sec(e + fx)}} + \frac{3 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{4f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^6*\text{Sqrt}[b*\text{Sec}[e + f*x]],x]$

[Out] $(-3*b*\text{Csc}[e + f*x])/(4*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (3*b*\text{Csc}[e + f*x]^3)/(10*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (b*\text{Csc}[e + f*x]^5)/(5*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (3*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(4*f)$

Rule 2705

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[a^2*((m + n - 2)/(m - 1)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{IntegersQ}[2*m, 2*n] \&\& !\text{GtQ}[n, m]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int \csc^6(e+fx) \sqrt{b \sec(e+fx)} dx &= -\frac{b \csc^5(e+fx)}{5f \sqrt{b \sec(e+fx)}} + \frac{9}{10} \int \csc^4(e+fx) \sqrt{b \sec(e+fx)} dx \\
&= -\frac{3b \csc^3(e+fx)}{10f \sqrt{b \sec(e+fx)}} - \frac{b \csc^5(e+fx)}{5f \sqrt{b \sec(e+fx)}} + \frac{3}{4} \int \csc^2(e+fx) \sqrt{b \sec(e+fx)} dx \\
&= -\frac{3b \csc(e+fx)}{4f \sqrt{b \sec(e+fx)}} - \frac{3b \csc^3(e+fx)}{10f \sqrt{b \sec(e+fx)}} - \frac{b \csc^5(e+fx)}{5f \sqrt{b \sec(e+fx)}} + \dots \\
&= -\frac{3b \csc(e+fx)}{4f \sqrt{b \sec(e+fx)}} - \frac{3b \csc^3(e+fx)}{10f \sqrt{b \sec(e+fx)}} - \frac{b \csc^5(e+fx)}{5f \sqrt{b \sec(e+fx)}} + \dots \\
&= -\frac{3b \csc(e+fx)}{4f \sqrt{b \sec(e+fx)}} - \frac{3b \csc^3(e+fx)}{10f \sqrt{b \sec(e+fx)}} - \frac{b \csc^5(e+fx)}{5f \sqrt{b \sec(e+fx)}} + \dots
\end{aligned}$$

Mathematica [A]

time = 0.32, size = 73, normalized size = 0.59

$$\frac{\left(-\cot(e+fx)(15+6\csc^2(e+fx)+4\csc^4(e+fx))+15\sqrt{\cos(e+fx)}F\left(\frac{1}{2}(e+fx)|2\right)\right)\sqrt{b\sec(e+fx)}}{20f}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^6*Sqrt[b*Sec[e + f*x]],x]`

```
[Out] ((-(Cot[e + f*x]*(15 + 6*Csc[e + f*x]^2 + 4*Csc[e + f*x]^4)) + 15*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2])*Sqrt[b*Sec[e + f*x]])/(20*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.29, size = 485, normalized size = 3.94

method	result
default	$\frac{(-1+\cos(fx+e))^2 \left(15i \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^5(fx+e) \sin(fx+e) + 15i \operatorname{EllipticF}\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^6*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/20/f*(-1+cos(f*x+e))^2*(15*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^5*sin(f*x+e)+15*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^4*sin(f*x+e)-30*I*EllipticF
```

$$(I*(-1+\cos(f*x+e))/\sin(f*x+e), I)*(1/(\cos(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*\cos(f*x+e)^3*\sin(f*x+e)-30*I*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e), I)*(1/(\cos(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*\cos(f*x+e)^2*\sin(f*x+e)+15*I*(1/(\cos(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e), I)*\sin(f*x+e)*\cos(f*x+e)+15*I*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e), I)*(1/(\cos(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*\sin(f*x+e)-15*\cos(f*x+e)^5+36*\cos(f*x+e)^3-25*\cos(f*x+e))*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(1/2)}/\sin(f*x+e)^9$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^6*(b*sec(f*x+e))^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))*csc(f*x + e)^6, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 204, normalized size = 1.66

$$\frac{15\sqrt{2}(i\cos(fx+e)^4-2i\cos(fx+e)^2+i)\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+15\sqrt{2}(-i\cos(fx+e)^4+2i\cos(fx+e)^2-i)\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))+2(15\cos(fx+e)^5-36\cos(fx+e)^3+25\cos(fx+e))\sqrt{\frac{b}{\cos(fx+e)}}}{40(f\cos(fx+e)^5-2f\cos(fx+e)^3+f)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^6*(b*sec(f*x+e))^(1/2), x, algorithm="fricas")

[Out] $-1/40*(15*\sqrt{2}*(I*\cos(f*x + e)^4 - 2*I*\cos(f*x + e)^2 + I)*\sqrt{b}*\sin(f*x + e)*\text{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) + 15*\sqrt{2}*(-I*\cos(f*x + e)^4 + 2*I*\cos(f*x + e)^2 - I)*\sqrt{b}*\sin(f*x + e)*\text{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)) + 2*(15*\cos(f*x + e)^5 - 36*\cos(f*x + e)^3 + 25*\cos(f*x + e))*\sqrt{b/\cos(f*x + e)})/((f*\cos(f*x + e)^4 - 2*f*\cos(f*x + e)^2 + f)*\sin(f*x + e))$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**6*(b*sec(f*x+e))**(1/2), x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^6*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*csc(f*x + e)^6, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(e + f x)}}}{\sin(e + f x)^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^6,x)

[Out] int((b/cos(e + f*x))^(1/2)/sin(e + f*x)^6, x)

3.385 $\int (b \sec(e + fx))^{3/2} \sin^7(e + fx) dx$

Optimal. Leaf size=83

$$\frac{2b^7}{11f(b \sec(e + fx))^{11/2}} - \frac{6b^5}{7f(b \sec(e + fx))^{7/2}} + \frac{2b^3}{f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

[Out] $2/11*b^7/f/(b*\sec(f*x+e))^(11/2)-6/7*b^5/f/(b*\sec(f*x+e))^(7/2)+2*b^3/f/(b*\sec(f*x+e))^(3/2)+2*b*(b*\sec(f*x+e))^(1/2)/f$

Rubi [A]

time = 0.04, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$\frac{2b^7}{11f(b \sec(e + fx))^{11/2}} - \frac{6b^5}{7f(b \sec(e + fx))^{7/2}} + \frac{2b^3}{f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^(3/2)*\text{Sin}[e + f*x]^7, x]$

[Out] $(2*b^7)/(11*f*(b*\text{Sec}[e + f*x])^(11/2)) - (6*b^5)/(7*f*(b*\text{Sec}[e + f*x])^(7/2)) + (2*b^3)/(f*(b*\text{Sec}[e + f*x])^(3/2)) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]])/f$

Rule 276

$\text{Int}[(c_*)*(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2702

$\text{Int}[\text{csc}[(e_*) + (f_*)*(x_)]^(n_)*((a_)*\text{sec}[(e_*) + (f_*)*(x_)]^(m_)), x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2)], x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n + 1)/2] \ \&\& \ !(\text{IntegerQ}[(m + 1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\int (b \sec(e + fx))^{3/2} \sin^7(e + fx) dx = \frac{b^7 \text{Subst} \left(\int \frac{(-1 + \frac{x^2}{b^2})^3}{x^{13/2}} dx, x, b \sec(e + fx) \right)}{f}$$

$$= \frac{b^7 \text{Subst} \left(\int \left(-\frac{1}{x^{13/2}} + \frac{3}{b^2 x^{9/2}} - \frac{3}{b^4 x^{5/2}} + \frac{1}{b^6 \sqrt{x}} \right) dx, x, b \sec(e + fx) \right)}{f}$$

$$= \frac{2b^7}{11f(b \sec(e + fx))^{11/2}} - \frac{6b^5}{7f(b \sec(e + fx))^{7/2}} + \frac{2b^3}{f(b \sec(e + fx))^{3/2}}$$

Mathematica [A]

time = 0.10, size = 52, normalized size = 0.63

$$\frac{b(3370 + 809 \cos(2(e + fx)) - 90 \cos(4(e + fx)) + 7 \cos(6(e + fx))) \sqrt{b \sec(e + fx)}}{1232f}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^7,x]``[Out] (b*(3370 + 809*Cos[2*(e + f*x)] - 90*Cos[4*(e + f*x)] + 7*Cos[6*(e + f*x)]) *Sqrt[b*Sec[e + f*x]])/(1232*f)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 968 vs. 2(71) = 142.

time = 1.63, size = 969, normalized size = 11.67

method	result	size
default	Expression too large to display	969

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e)^7,x,method=_RETURNVERBOSE)`

```
[Out] 1/154/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(77*cos(f*x+e)^3*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-77*cos(f*x+e)^3*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+28*cos(f*x+e)^7+231*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-231*cos(f*x+e)^2*(-cos(f*x+e)/(cos
```

$$\begin{aligned} & (f*x+e)+1)^2)^{(3/2)}*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} \\ &)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f \\ & *x+e)^2)+231*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\ln(-2*(2*\cos(f \\ & *x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(- \\ & \cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-231*\cos(f*x+e)*(-\cos(f* \\ & x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1 \\ &)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} \\ & -1)/\sin(f*x+e)^2)-132*\cos(f*x+e)^5+77*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos \\ & (f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+ \\ & 1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}-77*\ln(-2 \\ & *cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e \\ &)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*(-\cos(f*x+e)/(\cos \\ & (f*x+e)+1)^2)^{(3/2)}+308*\cos(f*x+e)^3+308*\cos(f*x+e))*(b/\cos(f*x+e))^{(3/2)}/s \\ & \text{in}(f*x+e)^4 \end{aligned}$$

Maxima [A]

time = 0.31, size = 76, normalized size = 0.92

$$\frac{2b \left(\frac{7b^6}{\left(\frac{b}{\cos(fx+e)}\right)^{\frac{11}{2}}} - \frac{33b^4}{\left(\frac{b}{\cos(fx+e)}\right)^{\frac{7}{2}}} + \frac{77b^2}{\left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}} + 77\sqrt{\frac{b}{\cos(fx+e)}} \right)}{77f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^7,x, algorithm="maxima")

[Out] 2/77*b*(7*b^6/(b/cos(f*x + e))^(11/2) - 33*b^4/(b/cos(f*x + e))^(7/2) + 77*b^2/(b/cos(f*x + e))^(3/2) + 77*sqrt(b/cos(f*x + e)))/f

Fricas [A]

time = 0.38, size = 58, normalized size = 0.70

$$\frac{2(7b \cos(fx+e)^6 - 33b \cos(fx+e)^4 + 77b \cos(fx+e)^2 + 77b) \sqrt{\frac{b}{\cos(fx+e)}}}{77f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^7,x, algorithm="fricas")

[Out] 2/77*(7*b*cos(f*x + e)^6 - 33*b*cos(f*x + e)^4 + 77*b*cos(f*x + e)^2 + 77*b)*sqrt(b/cos(f*x + e))/f

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e)**7,x)`

[Out] Timed out

Giac [A]

time = 4.00, size = 106, normalized size = 1.28

$$\frac{2 \left(7 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e)^5 - 33 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e)^3 + 77 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e) + \frac{77 b^6}{\sqrt{b \cos(fx + e)}} \right) \operatorname{sgn}(\cos(fx + e))}{77 b^4 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^7,x, algorithm="giac")`

[Out] `2/77*(7*sqrt(b*cos(f*x + e))*b^5*cos(f*x + e)^5 - 33*sqrt(b*cos(f*x + e))*b^5*cos(f*x + e)^3 + 77*sqrt(b*cos(f*x + e))*b^5*cos(f*x + e) + 77*b^6/sqrt(b*cos(f*x + e)))*sgn(cos(f*x + e))/(b^4*f)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^7 \left(\frac{b}{\cos(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^7*(b/cos(e + f*x))^(3/2),x)`

[Out] `int(sin(e + f*x)^7*(b/cos(e + f*x))^(3/2), x)`

3.386 $\int (b \sec(e + fx))^{3/2} \sin^5(e + fx) dx$

Optimal. Leaf size=63

$$-\frac{2b^5}{7f(b \sec(e + fx))^{7/2}} + \frac{4b^3}{3f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

[Out] $-2/7*b^5/f/(b*\sec(f*x+e))^{(7/2)}+4/3*b^3/f/(b*\sec(f*x+e))^{(3/2)}+2*b*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$-\frac{2b^5}{7f(b \sec(e + fx))^{7/2}} + \frac{4b^3}{3f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^5, x]$

[Out] $(-2*b^5)/(7*f*(b*\text{Sec}[e + f*x])^{(7/2)}) + (4*b^3)/(3*f*(b*\text{Sec}[e + f*x])^{(3/2)}) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]])/f$

Rule 276

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_*) + (b_*)*(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0]$

Rule 2702

$\text{Int}[\text{csc}[(e_*) + (f_*)*(x_)]^{(n_*)}*((a_*)*\text{sec}[(e_*) + (f_*)*(x_)]^{(m_*)}, x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1 + x^2/a^2)^{(n+1)/2}], x], x, a*\text{Sec}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n+1)/2] \&\& \text{!(IntegerQ}[(m+1)/2] \&\& \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{3/2} \sin^5(e + fx) dx &= \frac{b^5 \text{Subst} \left(\int \frac{(-1 + \frac{x^2}{b^2})^2}{x^{9/2}} dx, x, b \sec(e + fx) \right)}{f} \\
&= \frac{b^5 \text{Subst} \left(\int \left(\frac{1}{x^{9/2}} - \frac{2}{b^2 x^{5/2}} + \frac{1}{b^4 \sqrt{x}} \right) dx, x, b \sec(e + fx) \right)}{f} \\
&= -\frac{2b^5}{7f(b \sec(e + fx))^{7/2}} + \frac{4b^3}{3f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 42, normalized size = 0.67

$$\frac{b(215 + 44 \cos(2(e + fx)) - 3 \cos(4(e + fx))) \sqrt{b \sec(e + fx)}}{84f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^5,x]

[Out] (b*(215 + 44*Cos[2*(e + f*x)] - 3*Cos[4*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/(84*f)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 958 vs. 2(53) = 106.

time = 0.25, size = 959, normalized size = 15.22

method	result	size
default	Expression too large to display	959

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e)^5,x,method=_RETURNVERBOSE)

[Out]
$$\begin{aligned}
& -1/42/f*(\cos(f*x+e)+1)^2*(-1+\cos(f*x+e))^2*(-21*\cos(f*x+e)^3*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+21*\cos(f*x+e)^3*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-63*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+63*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}
\end{aligned}$$

$$2) \cdot \ln(-2 \cos(fx+e)^2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - 1) / \sin(fx+e)^2 + 12 \cos(fx+e)^5 - 63 \cos(fx+e) \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(3/2)} \cdot \ln(-2 \cdot (2 \cos(fx+e)^2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - 1) / \sin(fx+e)^2) + 63 \cos(fx+e) \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(3/2)} \cdot \ln(-2 \cdot (2 \cos(fx+e)^2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - 1) / \sin(fx+e)^2) - 21 \cdot \ln(-2 \cdot (2 \cos(fx+e)^2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - 1) / \sin(fx+e)^2) \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(3/2)} + 21 \cdot \ln(-2 \cdot (2 \cos(fx+e)^2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(1/2)} - 1) / \sin(fx+e)^2) \cdot (-\cos(fx+e)/(\cos(fx+e)+1)^2)^{(3/2)} - 56 \cos(fx+e)^3 - 84 \cos(fx+e) \cdot (b/\cos(fx+e))^{(3/2)} / \sin(fx+e)^4$$

Maxima [A]

time = 0.28, size = 58, normalized size = 0.92

$$-\frac{2b \left(\frac{3b^4}{\left(\frac{b}{\cos(fx+e)}\right)^{\frac{7}{2}}} - \frac{14b^2}{\left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}} - 21 \sqrt{\frac{b}{\cos(fx+e)}} \right)}{21f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^5,x, algorithm="maxima")

[Out] -2/21*b*(3*b^4/(b/cos(f*x + e))^(7/2) - 14*b^2/(b/cos(f*x + e))^(3/2) - 21*sqrt(b/cos(f*x + e)))/f

Fricas [A]

time = 0.38, size = 46, normalized size = 0.73

$$-\frac{2(3b \cos(fx+e)^4 - 14b \cos(fx+e)^2 - 21b) \sqrt{\frac{b}{\cos(fx+e)}}}{21f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^5,x, algorithm="fricas")

[Out] -2/21*(3*b*cos(f*x + e)^4 - 14*b*cos(f*x + e)^2 - 21*b)*sqrt(b/cos(f*x + e))/f

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e)**5,x)

[Out] Timed out

Giac [A]

time = 4.01, size = 81, normalized size = 1.29

$$\frac{2 \left(3 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e)^3 - 14 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e) - \frac{21 b^4}{\sqrt{b \cos(fx + e)}} \right) \operatorname{sgn}(\cos(fx + e))}{21 b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^5,x, algorithm="giac")

[Out] -2/21*(3*sqrt(b*cos(f*x + e))*b^3*cos(f*x + e)^3 - 14*sqrt(b*cos(f*x + e))*b^3*cos(f*x + e) - 21*b^4/sqrt(b*cos(f*x + e)))*sgn(cos(f*x + e))/(b^2*f)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + fx)^5 \left(\frac{b}{\cos(e + fx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^5*(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^5*(b/cos(e + f*x))^(3/2), x)

3.387 $\int (b \sec(e + fx))^{3/2} \sin^3(e + fx) dx$

Optimal. Leaf size=41

$$\frac{2b^3}{3f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

[Out] $2/3*b^3/f/(b*\sec(f*x+e))^(3/2)+2*b*(b*\sec(f*x+e))^(1/2)/f$

Rubi [A]

time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 14}

$$\frac{2b^3}{3f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^(3/2)*\text{Sin}[e + f*x]^3, x]$

[Out] $(2*b^3)/(3*f*(b*\text{Sec}[e + f*x])^(3/2)) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]])/f$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^(m_), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x] \&\& \text{SumQ}[u] \&\& !\text{LinearQ}[u, x] \&\& !\text{MatchQ}[u, (a_ + (b_)*(v_)] /; \text{FreeQ}\{a, b\}, x] \&\& \text{InverseFunctionQ}[v]$

Rule 2702

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^(n_)*((a_.)*\text{sec}[(e_.) + (f_.)*(x_)]^(m_), x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^(m+n-1)/(-1+x^2/a^2)^(n+1)/2], x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n+1)/2] \&\& !(\text{IntegerQ}[(m+1)/2] \&\& \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^{3/2} \sin^3(e + fx) dx &= \frac{b^3 \text{Subst} \left(\int \frac{-1 + \frac{x^2}{b^2}}{x^{5/2}} dx, x, b \sec(e + fx) \right)}{f} \\ &= \frac{b^3 \text{Subst} \left(\int \left(-\frac{1}{x^{5/2}} + \frac{1}{b^2 \sqrt{x}} \right) dx, x, b \sec(e + fx) \right)}{f} \\ &= \frac{2b^3}{3f(b \sec(e + fx))^{3/2}} + \frac{2b\sqrt{b \sec(e + fx)}}{f} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 30, normalized size = 0.73

$$\frac{b(7 + \cos(2(e + fx)))\sqrt{b\sec(e + fx)}}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^3,x]

[Out] (b*(7 + Cos[2*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/(3*f)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 948 vs. 2(35) = 70.

time = 0.24, size = 949, normalized size = 23.15

method	result	size
default	Expression too large to display	949

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e)^3,x,method=_RETURNVERBOSE)

[Out] 1/6/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(3*cos(f*x+e)^3*(-cos(f*x+e))/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e))/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-3*cos(f*x+e)^3*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+9*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-9*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+9*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-9*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+3*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)-3*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+4*cos(f*x+e)^3+12*cos(f*x+e))* (b/cos(f*x+e))^(3/2)/sin(f*x+e)^4

Maxima [A]

time = 0.29, size = 39, normalized size = 0.95

$$\frac{2b \left(\frac{b^2}{\left(\frac{b}{\cos(fx+e)}\right)^{3/2}} + 3 \sqrt{\frac{b}{\cos(fx+e)}} \right)}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^3,x, algorithm="maxima")
```

```
[Out] 2/3*b*(b^2/(b/cos(f*x + e))^(3/2) + 3*sqrt(b/cos(f*x + e)))/f
```

Fricas [A]

time = 0.40, size = 33, normalized size = 0.80

$$\frac{2(b \cos(fx + e)^2 + 3b) \sqrt{\frac{b}{\cos(fx + e)}}}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^3,x, algorithm="fricas")
```

```
[Out] 2/3*(b*cos(f*x + e)^2 + 3*b)*sqrt(b/cos(f*x + e))/f
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e)**3,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 5006 deep
```

Giac [A]

time = 5.94, size = 50, normalized size = 1.22

$$\frac{2 \left(\sqrt{b \cos(fx + e)} b \cos(fx + e) + \frac{3b^2}{\sqrt{b \cos(fx + e)}} \right) \operatorname{sgn}(\cos(fx + e))}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^3,x, algorithm="giac")
```

[Out] $\frac{2}{3}(\sqrt{b\cos(fx + e)} * b\cos(fx + e) + 3*b^2/\sqrt{b\cos(fx + e)}) * \text{sgn}(\cos(fx + e))/f$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + fx)^3 \left(\frac{b}{\cos(e + fx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^3*(b/cos(e + f*x))^(3/2), x)`

[Out] `int(sin(e + f*x)^3*(b/cos(e + f*x))^(3/2), x)`

3.388 $\int (b \sec(e + fx))^{3/2} \sin(e + fx) dx$

Optimal. Leaf size=18

$$\frac{2b\sqrt{b \sec(e + fx)}}{f}$$

[Out] 2*b*(b*sec(f*x+e))^(1/2)/f

Rubi [A]

time = 0.02, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 30}

$$\frac{2b\sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x],x]

[Out] (2*b*Sqrt[b*Sec[e + f*x]])/f

Rule 30

Int[(x_)^(m_.), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^{3/2} \sin(e + fx) dx &= \frac{b \text{Subst}\left(\int \frac{1}{\sqrt{x}} dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{2b\sqrt{b \sec(e + fx)}}{f} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 18, normalized size = 1.00

$$\frac{2b\sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x],x]

[Out] (2*b*Sqrt[b*Sec[e + f*x]])/f

Maple [A]

time = 0.05, size = 17, normalized size = 0.94

method	result	size
derivativedivides	$\frac{2b\sqrt{b\sec(fx+e)}}{f}$	17
default	$\frac{2b\sqrt{b\sec(fx+e)}}{f}$	17

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e),x,method=_RETURNVERBOSE)

[Out] 2*b*(b*sec(f*x+e))^(1/2)/f

Maxima [A]

time = 0.28, size = 25, normalized size = 1.39

$$\frac{2\left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}\cos(fx+e)}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e),x, algorithm="maxima")

[Out] 2*(b/cos(f*x + e))^(3/2)*cos(f*x + e)/f

Fricas [A]

time = 0.38, size = 19, normalized size = 1.06

$$\frac{2b\sqrt{\frac{b}{\cos(fx+e)}}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e),x, algorithm="fricas")

[Out] 2*b*sqrt(b/cos(f*x + e))/f

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b\sec(e+fx))^{\frac{3}{2}}\sin(e+fx)dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e),x)

[Out] Integral((b*sec(e + f*x))**(3/2)*sin(e + f*x), x)

Giac [A]

time = 4.59, size = 27, normalized size = 1.50

$$\frac{2b^2 \operatorname{sgn}(\cos(fx + e))}{\sqrt{b \cos(fx + e)} f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e),x, algorithm="giac")

[Out] 2*b^2*sgn(cos(f*x + e))/(sqrt(b*cos(f*x + e))*f)

Mupad [B]

time = 0.48, size = 18, normalized size = 1.00

$$\frac{2b \sqrt{\frac{b}{\cos(e + fx)}}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)*(b/cos(e + f*x))^(3/2),x)

[Out] (2*b*(b/cos(e + f*x))^(1/2))/f

3.389 $\int \csc(e + fx)(b \sec(e + fx))^{3/2} dx$

Optimal. Leaf size=77

$$-\frac{b^{3/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

[Out] $-b^{(3/2)}*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f-b^{(3/2)}*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f+2*b*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.04, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2702, 327, 335, 218, 212, 209}

$$-\frac{b^{3/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{2b\sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]*(b*Sec[e + f*x])^(3/2),x]`

[Out] $-(b^{(3/2)}*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/f) - (b^{(3/2)}*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/f) + (2*b*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/f$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \csc(e + fx)(b \sec(e + fx))^{3/2} dx &= \frac{\text{Subst}\left(\int \frac{x^{3/2}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{bf} \\
&= \frac{2b\sqrt{b \sec(e + fx)}}{f} + \frac{b \text{Subst}\left(\int \frac{1}{\sqrt{x}(-1 + \frac{x^2}{b^2})} dx, x, b \sec(e + fx)\right)}{f} \\
&= \frac{2b\sqrt{b \sec(e + fx)}}{f} + \frac{(2b) \text{Subst}\left(\int \frac{1}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} \\
&= \frac{2b\sqrt{b \sec(e + fx)}}{f} - \frac{b^2 \text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} - \frac{b^2 \text{Subst}\left(\int \frac{1}{b + x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} \\
&= -\frac{b^{3/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{b^{3/2} \log\left(\frac{1 - \sqrt{b \sec(e + fx)}}{1 + \sqrt{b \sec(e + fx)}}\right)}{f}
\end{aligned}$$

Mathematica [A]

time = 0.66, size = 85, normalized size = 1.10

$$\frac{\left(-2 \tan^{-1}\left(\sqrt{\sec(e + fx)}\right) + \log\left(1 - \sqrt{\sec(e + fx)}\right) - \log\left(1 + \sqrt{\sec(e + fx)}\right) + 4\sqrt{\sec(e + fx)}\right)(b \sec(e + fx))^{3/2}}{2f \sec^{3/2}(e + fx)}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]*(b*Sec[e + f*x])^(3/2),x]

[Out] $((-2*\text{ArcTan}[\text{Sqrt}[\text{Sec}[e + f*x]]] + \text{Log}[1 - \text{Sqrt}[\text{Sec}[e + f*x]]] - \text{Log}[1 + \text{Sqrt}[\text{Sec}[e + f*x]]] + 4*\text{Sqrt}[\text{Sec}[e + f*x]])*(b*\text{Sec}[e + f*x])^{(3/2)}/(2*f*\text{Sec}[e + f*x]^{(3/2)})$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(63) = 126.

time = 0.25, size = 235, normalized size = 3.05

method	result
default	$\frac{(-1 + \cos(fx+e))^3 \left(4 \cos(fx+e) \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} + \arctan\left(\frac{1}{2\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}\right) \cos(fx+e) + \ln\left(-\frac{2\left(2\cos^2(fx+e)\right)}{2f \sin(fx+e)}\right) \right)}{2f \sin(fx+e)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)*(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] $1/2/f*(-1+\cos(f*x+e))^3*(4*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} + \arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})*\cos(f*x+e) + \ln(-2*(2*\cos(f*x+e))^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - \cos(f*x+e)^2 + 2*\cos(f*x+e) - 2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - 1)/\sin(f*x+e)^2)*\cos(f*x+e) + 4*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}*(b/\cos(f*x+e))^{(3/2)}*\cos(f*x+e)^2/\sin(f*x+e)^6/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}$

Maxima [A]

time = 0.51, size = 91, normalized size = 1.18

$$\frac{\left(2\sqrt{b} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) - \sqrt{b} \log\left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}}\right) - 4\sqrt{\frac{b}{\cos(fx+e)}} \right) b}{2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] $-1/2*(2*\text{sqrt}(b)*\arctan(\text{sqrt}(b/\cos(f*x + e))/\text{sqrt}(b)) - \text{sqrt}(b)*\log(-(\text{sqrt}(b) - \text{sqrt}(b/\cos(f*x + e)))/(\text{sqrt}(b) + \text{sqrt}(b/\cos(f*x + e)))) - 4*\text{sqrt}(b/\cos(f*x + e))*b/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 141 vs. 2(66) = 132.

time = 0.42, size = 298, normalized size = 3.87

$$\frac{2\sqrt{-b}\arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\cos(fx+e)+1}\right) + \sqrt{-b}b\log\left(\frac{b\cos(fx+e)^2 + (\cos(fx+e)^2 - \cos(fx+e))\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}} - 4b\cos(fx+e) + b}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right) + 8b\sqrt{\frac{b}{\cos(fx+e)}} + 2b^2\arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{2\sqrt{b}}\right) + b^2\log\left(\frac{b\cos(fx+e)^2 - 4(\cos(fx+e)^2 + \cos(fx+e))\sqrt{b}\sqrt{\frac{b}{\cos(fx+e)}} + 4b\cos(fx+e) + b}{\cos(fx+e)^2 - 2\cos(fx+e) + 1}\right) + 8b\sqrt{\frac{b}{\cos(fx+e)}}}{4f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(-b)*b*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b) + sqrt(-b)*b*log((b*cos(f*x + e)^2 + 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 8*b*sqrt(b/cos(f*x + e)))/f, 1/4*(2*b^(3/2)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b)) + b^(3/2)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)) + 8*b*sqrt(b/cos(f*x + e)))/f]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^{\frac{3}{2}} \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(3/2),x)

[Out] Integral((b*sec(e + f*x))^(3/2)*csc(e + f*x), x)

Giac [A]

time = 5.19, size = 79, normalized size = 1.03

$$\frac{b^4 \left(\frac{\arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^2} + \frac{\arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}} + \frac{2}{\sqrt{b \cos(fx+e)} b^2} \right) \operatorname{sgn}(\cos(fx+e))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] b^4*(arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/(sqrt(-b)*b^2) + arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(5/2) + 2/(sqrt(b*cos(f*x + e))*b^2))*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{3/2}}{\sin(e+fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(3/2)/sin(e + f*x),x)

[Out] int((b/cos(e + f*x))^(3/2)/sin(e + f*x), x)

3.390 $\int \csc^3(e + fx)(b \sec(e + fx))^{3/2} dx$

Optimal. Leaf size=113

$$\frac{5b^{3/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{5b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} + \frac{5b\sqrt{b \sec(e + fx)}}{2f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{5/2}}{2bf}$$

[Out] $-5/4*b^{(3/2)}*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f-5/4*b^{(3/2)}*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f-1/2*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(5/2)}/b/f+5/2*b*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.05, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2702, 294, 327, 335, 218, 212, 209}

$$\frac{5b^{3/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{5b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} + \frac{5b\sqrt{b \sec(e + fx)}}{2f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{5/2}}{2bf}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[e + f*x]^3*(b*\operatorname{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $(-5*b^{(3/2)}*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]])/(4*f) - (5*b^{(3/2)}*\operatorname{ArcTan}[\operatorname{h}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]])/(4*f) + (5*b*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(2*f) - (\operatorname{Cot}[e + f*x]^2*(b*\operatorname{Sec}[e + f*x])^{(5/2)})/(2*b*f)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n *((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 327

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \csc^3(e + fx)(b \sec(e + fx))^{3/2} dx &= \frac{\text{Subst}\left(\int \frac{x^{7/2}}{(-1 + \frac{x^2}{b^2})^2} dx, x, b \sec(e + fx)\right)}{b^3 f} \\
&= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{5/2}}{2bf} + \frac{5 \text{Subst}\left(\int \frac{x^{3/2}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{4bf} \\
&= \frac{5b \sqrt{b \sec(e + fx)}}{2f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{5/2}}{2bf} + \frac{(5b) \text{Subst}\left(\int \frac{x^{3/2}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{4bf} \\
&= \frac{5b \sqrt{b \sec(e + fx)}}{2f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{5/2}}{2bf} + \frac{(5b) \text{Subst}\left(\int \frac{x^{3/2}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{4bf} \\
&= \frac{5b \sqrt{b \sec(e + fx)}}{2f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{5/2}}{2bf} - \frac{(5b^2) \text{Subst}\left(\int \frac{x^{3/2}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{4bf} \\
&= -\frac{5b^{3/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{5b^{3/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f}
\end{aligned}$$

Mathematica [A]

time = 1.63, size = 97, normalized size = 0.86

$$-\frac{(10 \tan^{-1}(\sqrt{\sec(e + fx)}) - 5 \log(1 - \sqrt{\sec(e + fx)}) + 5 \log(1 + \sqrt{\sec(e + fx)}) + 4(-5 + \csc^2(e + fx)) \sqrt{\sec(e + fx)}) (b \sec(e + fx))^{3/2}}{8f \sec^{3/2}(e + fx)}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^3*(b*Sec[e + f*x])^(3/2), x]`

```
[Out] -1/8*((10*ArcTan[Sqrt[Sec[e + f*x]]] - 5*Log[1 - Sqrt[Sec[e + f*x]]] + 5*Log[1 + Sqrt[Sec[e + f*x]]] + 4*(-5 + Csc[e + f*x]^2)*Sqrt[Sec[e + f*x]])*(b*Sec[e + f*x])^(3/2)/(f*Sec[e + f*x]^(3/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 643 vs. 2(89) = 178.

time = 0.23, size = 644, normalized size = 5.70

method	result
--------	--------

default	$(\cos(fx+e)+1)(-1+\cos(fx+e))^3 \left(4 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} (\cos^3(fx+e)+8(\cos^2(fx+e))) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 4 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} \right)$
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^3*(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{8} \frac{1}{f} (\cos(fx+e)+1) (-1+\cos(fx+e))^3 \left(4 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} (\cos^3(fx+e)+8(\cos^2(fx+e))) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 4 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} \right)$$

Maxima [A]

time = 0.69, size = 129, normalized size = 1.14

$$\frac{\left(\frac{4b^2 \sqrt{\frac{b}{\cos(fx+e)}}}{b^2 - \frac{b^2}{\cos(fx+e)^2}} - 10\sqrt{b} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + 5\sqrt{b} \log\left(-\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}}\right) + 16\sqrt{\frac{b}{\cos(fx+e)}} \right) b}{8f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out]
$$\frac{1}{8} (4b^2 \sqrt{b/\cos(fx+e)}) / (b^2 - b^2/\cos(fx+e)^2) - 10 \sqrt{b} \arctan(\sqrt{b/\cos(fx+e)}/\sqrt{b}) + 5 \sqrt{b} \log(-(\sqrt{b} - \sqrt{b/\cos(fx+e)})/(\sqrt{b} + \sqrt{b/\cos(fx+e)})) + 16 \sqrt{b/\cos(fx+e)} * b/f$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 201 vs. 2(94) = 188.

time = 0.48, size = 416, normalized size = 3.68

$$\frac{10 \left((b \cos(fx+e) - b)^2 \sqrt{\frac{b}{\cos(fx+e)}} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + 5 \left((b \cos(fx+e) - b) \sqrt{\frac{b}{\cos(fx+e)}} \log\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}} - \sqrt{b}}{\sqrt{\frac{b}{\cos(fx+e)}} + \sqrt{b}}\right) + 8 \left((b \cos(fx+e) - 4b) \sqrt{\frac{b}{\cos(fx+e)}} \right) \right) + 16 \left((b \cos(fx+e) - b) \sqrt{\frac{b}{\cos(fx+e)}} \right) \right)}{8 \left((b \cos(fx+e) - b) \sqrt{\frac{b}{\cos(fx+e)}} \right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] [1/16*(10*(b*cos(f*x + e)^2 - b)*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e)))*(cos(f*x + e) + 1)/b) + 5*(b*cos(f*x + e)^2 - b)*sqrt(-b)*log((b*cos(f*x + e)^2 + 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 8*(5*b*cos(f*x + e)^2 - 4*b)*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e)^2 - f), 1/16*(10*(b*cos(f*x + e)^2 - b)*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e)))*(cos(f*x + e) - 1)/sqrt(b)) + 5*(b*cos(f*x + e)^2 - b)*sqrt(b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)) + 8*(5*b*cos(f*x + e)^2 - 4*b)*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e)^2 - f)]

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3*(b*sec(f*x+e))**(3/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 5006 deep

Giac [A]

time = 3.77, size = 134, normalized size = 1.19

$$b^6 \left(\frac{5 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^4} + \frac{5 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}} + \frac{2(5b^2 \cos(fx+e)^2 - 4b^2)}{(\sqrt{b \cos(fx+e)} b^2 \cos(fx+e)^2 - \sqrt{b \cos(fx+e)} b^2)^{b^4}} \right) \operatorname{sgn}(\cos(fx+e))$$

$4f$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] 1/4*b^6*(5*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/(sqrt(-b)*b^4) + 5*arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(9/2) + 2*(5*b^2*cos(f*x + e)^2 - 4*b^2)/((sqrt(b*cos(f*x + e))*b^2*cos(f*x + e)^2 - sqrt(b*cos(f*x + e))*b^2)*b^4))*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{3/2}}{\sin(e+fx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(3/2)/sin(e + f*x)^3,x)

[Out] int((b/cos(e + f*x))^(3/2)/sin(e + f*x)^3, x)

3.391 $\int (b \sec(e + fx))^{3/2} \sin^6(e + fx) dx$

Optimal. Leaf size=128

$$-\frac{16b^2 E\left(\frac{1}{2}(e+fx)|2\right)}{3f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}} + \frac{8b^3 \sin(e+fx)}{3f(b\sec(e+fx))^{3/2}} + \frac{20b^3 \sin^3(e+fx)}{9f(b\sec(e+fx))^{3/2}} + \frac{2b\sqrt{b\sec(e+fx)} \sin^5(e+fx)}{f}$$

[Out] $8/3*b^3*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}+20/9*b^3*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(3/2)}-16/3*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}+2*b*\sin(f*x+e)^5*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.11, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2704, 2707, 3856, 2719}

$$\frac{20b^3 \sin^3(e+fx)}{9f(b\sec(e+fx))^{3/2}} + \frac{8b^3 \sin(e+fx)}{3f(b\sec(e+fx))^{3/2}} - \frac{16b^2 E\left(\frac{1}{2}(e+fx)|2\right)}{3f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}} + \frac{2b \sin^5(e+fx) \sqrt{b\sec(e+fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^6, x]$

[Out] $(-16*b^2*\text{EllipticE}[(e + f*x)/2, 2])/((3*f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (8*b^3*\text{Sin}[e + f*x])/((3*f*(b*\text{Sec}[e + f*x])^{(3/2)})) + (20*b^3*\text{Sin}[e + f*x]^3)/((9*f*(b*\text{Sec}[e + f*x])^{(3/2)})) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x]^5)/f$

Rule 2704

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*a*(n - 1)), x] + \text{Dist}[b^2*((m + 1)/(a^2*(n - 1))), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2707

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(a*f*(m + n)), x] + \text{Dist}[(m + 1)/(a^2*(m + n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int (b \sec(e + fx))^{3/2} \sin^6(e + fx) dx &= \frac{2b \sqrt{b \sec(e + fx)} \sin^5(e + fx)}{f} - (10b^2) \int \frac{\sin^4(e + fx)}{\sqrt{b \sec(e + fx)}} dx \\
 &= \frac{20b^3 \sin^3(e + fx)}{9f(b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^5(e + fx)}{f} - \frac{1}{3}(20b^2) \int \frac{\sin^2(e + fx)}{\sqrt{b \sec(e + fx)}} dx \\
 &= \frac{8b^3 \sin(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{20b^3 \sin^3(e + fx)}{9f(b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^5(e + fx)}{f} \\
 &= \frac{8b^3 \sin(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{20b^3 \sin^3(e + fx)}{9f(b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^5(e + fx)}{f} \\
 &= -\frac{16b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{3f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{8b^3 \sin(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{20b^3 \sin^3(e + fx)}{9f(b \sec(e + fx))^{3/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.11, size = 70, normalized size = 0.55

$$\frac{b \sqrt{b \sec(e + fx)} \left(384 \sqrt{\cos(e + fx)} E\left(\frac{1}{2}(e + fx) \mid 2\right) - 158 \sin(e + fx) - 13 \sin(3(e + fx)) + \sin(5(e + fx)) \right)}{72f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^6,x]

[Out] -1/72*(b*Sqrt[b*Sec[e + f*x]]*(384*Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2] - 158*Sin[e + f*x] - 13*Sin[3*(e + f*x)] + Sin[5*(e + f*x)]))/f

Maple [C] Result contains complex when optimal does not.

time = 0.30, size = 330, normalized size = 2.58

method	result
--------	--------

default	$2 \left(24i \operatorname{EllipticE} \left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i \right) \sin(fx+e) \cos(fx+e) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} - 24i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \right)$
---------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e)^6,x,method=_RETURNVERBOSE)
[Out] 2/9/f*(24*I*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)
*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)-24*I*(1/(cos(f*
x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e)
)/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+cos(f*x+e)^6+24*I*sin(f*x+e)*Elliptic
E(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos
(f*x+e)+1))^(1/2)-24*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*
x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-5*cos(f*x+e)^4+
19*cos(f*x+e)^2-24*cos(f*x+e)+9)*(b/cos(f*x+e))^(3/2)*cos(f*x+e)/sin(f*x+e)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^6,x, algorithm="maxima")
[Out] integrate((b*sec(f*x + e))^(3/2)*sin(f*x + e)^6, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 116, normalized size = 0.91

$$\frac{2 \left(12i \sqrt{2} b^{\frac{3}{2}} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e))) - 12i \sqrt{2} b^{\frac{3}{2}} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e))) + (b \cos(fx+e)^4 - 4b \cos(fx+e)^2 - 9b) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) \right)}{9f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^6,x, algorithm="fricas")
[Out] -2/9*(12*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0
, cos(f*x + e) + I*sin(f*x + e))) - 12*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4
, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) + (b*cos(f*
x + e)^4 - 4*b*cos(f*x + e)^2 - 9*b)*sqrt(b/cos(f*x + e))*sin(f*x + e))/f
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e)**6,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^6,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(3/2)*sin(f*x + e)^6, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^6 \left(\frac{b}{\cos(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^6*(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^6*(b/cos(e + f*x))^(3/2), x)

3.392 $\int (b \sec(e + fx))^{3/2} \sin^4(e + fx) dx$

Optimal. Leaf size=98

$$-\frac{24b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{5f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{12b^3 \sin(e + fx)}{5f (b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^3(e + fx)}{f}$$

[Out] $12/5*b^3*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}-24/5*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}+2*b*\sin(f*x+e)^3*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.08, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$,

Rules used = {2704, 2707, 3856, 2719}

$$\frac{12b^3 \sin(e + fx)}{5f (b \sec(e + fx))^{3/2}} - \frac{24b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{5f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{2b \sin^3(e + fx) \sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^4, x]$

[Out] $(-24*b^2*\text{EllipticE}[(e + f*x)/2, 2])/ (5*f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (12*b^3*\text{Sin}[e + f*x])/ (5*f*(b*\text{Sec}[e + f*x])^{(3/2)}) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x]^3)/f$

Rule 2704

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*a^{(n - 1)})], x] + \text{Dist}[b^2*((m + 1)/(a^2*(n - 1))], \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}[a, b, e, f], x] \&\& \text{GtQ}[n, 1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2707

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_)]))^{(n_)}, x_Symbol] :> \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(a*f^{(m + n)})], x] + \text{Dist}[(m + 1)/(a^2*(m + n)], \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}[a, b, e, f, n], x] \&\& \text{LtQ}[m, -1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[c, d], x]$

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int (b \sec(e + fx))^{3/2} \sin^4(e + fx) dx &= \frac{2b \sqrt{b \sec(e + fx)} \sin^3(e + fx)}{f} - (6b^2) \int \frac{\sin^2(e + fx)}{\sqrt{b \sec(e + fx)}} dx \\
 &= \frac{12b^3 \sin(e + fx)}{5f(b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^3(e + fx)}{f} - \frac{1}{5} (12b^2) \int \frac{\sin^2(e + fx)}{\sqrt{b \sec(e + fx)}} dx \\
 &= \frac{12b^3 \sin(e + fx)}{5f(b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^3(e + fx)}{f} - \frac{(12b^2) f}{5 \sqrt{\cos(e + fx)}} \\
 &= -\frac{24b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{5f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{12b^3 \sin(e + fx)}{5f(b \sec(e + fx))^{3/2}} + \frac{2b \sqrt{b \sec(e + fx)} \sin^3(e + fx)}{f}
 \end{aligned}$$

Mathematica [A]

time = 0.09, size = 60, normalized size = 0.61

$$\frac{b \sqrt{b \sec(e + fx)} \left(-48 \sqrt{\cos(e + fx)} E\left(\frac{1}{2}(e + fx) \mid 2\right) + 21 \sin(e + fx) + \sin(3(e + fx)) \right)}{10f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^4,x]

[Out] (b*Sqrt[b*Sec[e + f*x]]*(-48*Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2] + 21*Sin[e + f*x] + Sin[3*(e + f*x)]))/(10*f)

Maple [C] Result contains complex when optimal does not.

time = 0.25, size = 320, normalized size = 3.27

method	result
default	$ \frac{2 \left(12i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - 12i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \right)}{10f} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e)^4,x,method=_RETURNVERBOSE)


```
[Out] -2/5/f*(12*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-12*I*cos(f*x+e)*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)+12*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-12*I*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)+cos(f*x+e)^4-8*cos(f*x+e)^2+12*cos(f*x+e)-5)*cos(f*x+e)*(b/cos(f*x+e))^(3/2)/sin(f*x+e)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^4,x, algorithm="maxima")
```

```
[Out] integrate((b*sec(f*x + e))^(3/2)*sin(f*x + e)^4, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 105, normalized size = 1.07

$$\frac{2 \left(6i \sqrt{2} b^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(fx + e) + i \sin(fx + e))) - 6i \sqrt{2} b^3 \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(fx + e) - i \sin(fx + e))) - (b \cos(fx + e)^2 + 5b) \sqrt{\frac{b}{\cos(fx + e)}} \sin(fx + e) \right)}{5f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^4,x, algorithm="fricas")
```

```
[Out] -2/5*(6*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) - 6*I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) - (b*cos(f*x + e)^2 + 5*b)*sqrt(b/cos(f*x + e))*sin(f*x + e))/f
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e)**4,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 8009 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^4,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(3/2)*sin(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^4 \left(\frac{b}{\cos(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4*(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^4*(b/cos(e + f*x))^(3/2), x)

3.393 $\int (b \sec(e + fx))^{3/2} \sin^2(e + fx) dx$

Optimal. Leaf size=66

$$-\frac{4b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f}$$

[Out] $-4*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}+2*b*\sin(f*x+e)*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.05, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2704, 3856, 2719}

$$\frac{2b \sin(e + fx) \sqrt{b \sec(e + fx)}}{f} - \frac{4b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^2, x]$

[Out] $(-4*b^2*\text{EllipticE}[(e + f*x)/2, 2])/(f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x])/f$

Rule 2704

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_)]*(a_.)^{(m_)}*((b_.)*\sec[(e_.) + (f_.)*(x_)])^{(n_)}], x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*a^{(n - 1)})], x] + \text{Dist}[b^2*((m + 1)/(a^2*(n - 1))], \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}[a, b, e, f], x] \&\& \text{GtQ}[n, 1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[c, d], x]$

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_)}], x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[b, c, d], x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{3/2} \sin^2(e + fx) dx &= \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} - (2b^2) \int \frac{1}{\sqrt{b \sec(e + fx)}} dx \\
&= \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} - \frac{(2b^2) \int \sqrt{\cos(e + fx)} dx}{\sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
&= -\frac{4b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 48, normalized size = 0.73

$$\frac{2b \sqrt{b \sec(e + fx)} \left(-2 \sqrt{\cos(e + fx)} E\left(\frac{1}{2}(e + fx) \mid 2\right) + \sin(e + fx) \right)}{f}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^2,x]``[Out] (2*b*Sqrt[b*Sec[e + f*x]]*(-2*Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2] + Sin[e + f*x]))/f`**Maple [C]** Result contains complex when optimal does not.

time = 0.22, size = 310, normalized size = 4.70

method	result
default	$2 \left(-2i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) + 2i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \right) \sqrt{b \sec(fx+e)}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(3/2)*sin(f*x+e)^2,x,method=_RETURNVERBOSE)`

```
[Out] 2/f*(-2*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+2*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)-2*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)+2*I*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)+cos(f*x+e)^2-2*cos(f*x+e)+1)*cos(f*x+e)*(b/cos(f*x+e))^(3/2)/sin(f*x+e)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^2,x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^(3/2)*sin(f*x + e)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 91, normalized size = 1.38

$$\frac{2 \left(i \sqrt{2} b^{\frac{3}{2}} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(fx + e) + i \sin(fx + e))) - i \sqrt{2} b^{\frac{3}{2}} \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(fx + e) - i \sin(fx + e))) - b \sqrt{\frac{b}{\cos(fx + e)}} \sin(fx + e) \right)}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^2,x, algorithm="fricas")`

[Out] `-2*(I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) - I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) - b*sqrt(b/cos(f*x + e))*sin(f*x + e))/f`

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**(3/2)*sin(f*x+e)**2,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(3/2)*sin(f*x+e)^2,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^(3/2)*sin(f*x + e)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + fx)^2 \left(\frac{b}{\cos(e + fx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^2*(b/cos(e + f*x))^(3/2),x)`

[Out] `int(sin(e + f*x)^2*(b/cos(e + f*x))^(3/2), x)`

3.394 $\int (b \sec(e + fx))^{3/2} dx$

Optimal. Leaf size=66

$$-\frac{2b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f}$$

[Out] $-2*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}+2*b*\sin(f*x+e)*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.03, antiderivative size = 66, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3853, 3856, 2719}

$$\frac{2b \sin(e + fx) \sqrt{b \sec(e + fx)}}{f} - \frac{2b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $(-2*b^2*\text{EllipticE}[(e + f*x)/2, 2])/(f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x])/f$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3853

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{n-1}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{3/2} dx &= \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} - b^2 \int \frac{1}{\sqrt{b \sec(e + fx)}} dx \\
&= \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} - \frac{b^2 \int \sqrt{\cos(e + fx)} dx}{\sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
&= -\frac{2b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{2b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 48, normalized size = 0.73

$$\frac{2b \sqrt{b \sec(e + fx)} \left(-\sqrt{\cos(e + fx)} E\left(\frac{1}{2}(e + fx) \mid 2\right) + \sin(e + fx) \right)}{f}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^(3/2),x]``[Out] (2*b*Sqrt[b*Sec[e + f*x]]*(-(Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2]) + Sin[e + f*x]))/f`**Maple [C]** Result contains complex when optimal does not.

time = 0.26, size = 320, normalized size = 4.85

method	result
default	$-\frac{2(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) \right)}{f}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

```
[Out] -2/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-I*cos(f*x+e)*sin(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)+I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-I*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)+cos(f*x+e)-1)*cos(f*x+e)*(b/cos(f*x+e))^(3/2)/sin(f*x+e)^5
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^(3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 90, normalized size = 1.36

$$\frac{-i\sqrt{2}b^{\frac{3}{2}}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e)))+i\sqrt{2}b^{\frac{3}{2}}\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e)))+2b\sqrt{\frac{b}{\cos(fx+e)}}\sin(fx+e)}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] (-I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + I*sqrt(2)*b^(3/2)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) + 2*b*sqrt(b/cos(f*x + e))*sin(f*x + e))/f

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(3/2),x)

[Out] Integral((b*sec(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \left(\frac{b}{\cos(e + fx)} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(3/2),x)

[Out] int((b/cos(e + f*x))^(3/2), x)

3.395 $\int \csc^2(e + fx)(b \sec(e + fx))^{3/2} dx$

Optimal. Leaf size=90

$$-\frac{3b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{b \csc(e + fx) \sqrt{b \sec(e + fx)}}{f} + \frac{3b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f}$$

[Out] $-3*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}-b*\csc(f*x+e)*(b*\sec(f*x+e))^{(1/2)}/f+3*b*\sin(f*x+e)*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.06, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2705, 3853, 3856, 2719}

$$-\frac{3b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{b \csc(e + fx) \sqrt{b \sec(e + fx)}}{f} + \frac{3b \sin(e + fx) \sqrt{b \sec(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^2*(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $(-3*b^2*\text{EllipticE}[(e + f*x)/2, 2])/(f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (b*\text{Csc}[e + f*x]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/f + (3*b*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x])/f$

Rule 2705

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.)^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] :> \text{Simp}[(-a)*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[a^2*((m + n - 2)/(m - 1)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{IntegersQ}[2*m, 2*n] \&\& !\text{GtQ}[n, m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3853

$\text{Int}[(\csc[(c_.) + (d_.)*(x_.)]*(b_.)^{(n_.)}, x_Symbol] :> \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n - 1)})/(d*(n - 1)), x] + \text{Dist}[b^2*((n - 2)/(n - 1)), \text{Int}[(b*\text{Csc}[c + d*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \csc^2(e + fx)(b \sec(e + fx))^{3/2} dx &= -\frac{b \csc(e + fx) \sqrt{b \sec(e + fx)}}{f} + \frac{3}{2} \int (b \sec(e + fx))^{3/2} dx \\
&= -\frac{b \csc(e + fx) \sqrt{b \sec(e + fx)}}{f} + \frac{3b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} - \frac{1}{2} \int \frac{3b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} dx \\
&= -\frac{b \csc(e + fx) \sqrt{b \sec(e + fx)}}{f} + \frac{3b \sqrt{b \sec(e + fx)} \sin(e + fx)}{f} - \frac{3b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{b \csc(e + fx) \sqrt{b \sec(e + fx)}}{f}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 57, normalized size = 0.63

$$\frac{b \sqrt{b \sec(e + fx)} \left(-\csc(e + fx) - 3 \sqrt{\cos(e + fx)} E\left(\frac{1}{2}(e + fx) \mid 2\right) + 3 \sin(e + fx) \right)}{f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]^2*(b*Sec[e + f*x])^(3/2),x]
```

```
[Out] (b*Sqrt[b*Sec[e + f*x]]*(-Csc[e + f*x] - 3*Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2] + 3*Sin[e + f*x]))/f
```

Maple [C] Result contains complex when optimal does not.

time = 0.23, size = 322, normalized size = 3.58

method	result
default	$ -\frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(3i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - \dots \right)}{f} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)^2*(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)+3*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)+3*cos(f*x+e)-2)*cos(f*x+e)*(b/cos(f*x+e))^(3/2)/sin(f*x+e)^5
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*sec(f*x + e))^(3/2)*csc(f*x + e)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 122, normalized size = 1.36

$$\frac{-3i\sqrt{2}b^{\frac{3}{2}}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e)))+3i\sqrt{2}b^{\frac{3}{2}}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e)))-2(3b\cos(fx+e)^2-2b)\sqrt{\frac{b}{\cos(fx+e)}}}{2f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] 1/2*(-3*I*sqrt(2)*b^(3/2)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*I*sqrt(2)*b^(3/2)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) - 2*(3*b*cos(f*x + e)^2 - 2*b)*sqrt(b/cos(f*x + e))/(f*sin(f*x + e))
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**2*(b*sec(f*x+e))**(3/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(3/2)*csc(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{3/2}}{\sin(e+fx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(3/2)/sin(e + f*x)^2,x)

[Out] int((b/cos(e + f*x))^(3/2)/sin(e + f*x)^2, x)

3.396 $\int \csc^4(e + fx)(b \sec(e + fx))^{3/2} dx$

Optimal. Leaf size=124

$$\frac{7b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{2f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{7b \csc(e + fx) \sqrt{b \sec(e + fx)}}{6f} - \frac{b \csc^3(e + fx) \sqrt{b \sec(e + fx)}}{3f} + \frac{7b \sin(e + fx) \sqrt{b \sec(e + fx)}}{2f}$$

[Out] $-7/2*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}-7/6*b*\csc(f*x+e)*(b*\sec(f*x+e))^{(1/2)}/f-1/3*b*\csc(f*x+e)^3*(b*\sec(f*x+e))^{(1/2)}/f+7/2*b*\sin(f*x+e)*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.09, antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2705, 3853, 3856, 2719}

$$\frac{7b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{2f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{b \csc^3(e + fx) \sqrt{b \sec(e + fx)}}{3f} - \frac{7b \csc(e + fx) \sqrt{b \sec(e + fx)}}{6f} + \frac{7b \sin(e + fx) \sqrt{b \sec(e + fx)}}{2f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^4*(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $(-7*b^2*\text{EllipticE}[(e + f*x)/2, 2])/(2*f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (7*b*\text{Csc}[e + f*x]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(6*f) - (b*\text{Csc}[e + f*x]^3*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(3*f) + (7*b*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sin}[e + f*x])/(2*f)$

Rule 2705

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[a^2*((m + n - 2)/(m - 1)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /;$ FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 3853

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\text{Csc}[c + d*x])^{(n - 1)})/(d*(n - 1)), x] + \text{Dist}[b^2*((n - 2)/(n - 1)), \text{Int}[(b*\text{Csc}[c + d*x])^{(n - 2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \csc^4(e + fx)(b \sec(e + fx))^{3/2} dx &= -\frac{b \csc^3(e + fx) \sqrt{b \sec(e + fx)}}{3f} + \frac{7}{6} \int \csc^2(e + fx)(b \sec(e + fx))^{3/2} dx \\
&= -\frac{7b \csc(e + fx) \sqrt{b \sec(e + fx)}}{6f} - \frac{b \csc^3(e + fx) \sqrt{b \sec(e + fx)}}{3f} + \\
&= -\frac{7b \csc(e + fx) \sqrt{b \sec(e + fx)}}{6f} - \frac{b \csc^3(e + fx) \sqrt{b \sec(e + fx)}}{3f} + \\
&= -\frac{7b \csc(e + fx) \sqrt{b \sec(e + fx)}}{6f} - \frac{b \csc^3(e + fx) \sqrt{b \sec(e + fx)}}{3f} + \\
&= -\frac{7b^2 E\left(\frac{1}{2}(e + fx) \mid 2\right)}{2f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{7b \csc(e + fx) \sqrt{b \sec(e + fx)}}{6f}
\end{aligned}$$

Mathematica [A]

time = 0.15, size = 77, normalized size = 0.62

$$\frac{b(-21 + 7 \csc^2(e + fx) + 2 \csc^4(e + fx) + 21 \sqrt{\cos(e + fx)} \csc(e + fx) E\left(\frac{1}{2}(e + fx) \mid 2\right)) \sqrt{b \sec(e + fx)} \sin(e + fx)}{6f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]^4*(b*Sec[e + f*x])^(3/2), x]
```

```
[Out] -1/6*(b*(-21 + 7*Csc[e + f*x]^2 + 2*Csc[e + f*x]^4 + 21*Sqrt[Cos[e + f*x]]*
Csc[e + f*x]*EllipticE[(e + f*x)/2, 2])*Sqrt[b*Sec[e + f*x]]*Sin[e + f*x])/
f
```

Maple [C] Result contains complex when optimal does not.

time = 0.24, size = 622, normalized size = 5.02

method	result
default	$ \frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(21i \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^3(fx+e) \sin(fx+e)) \right)}{6f} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)^4*(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
[Out] 1/6/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(21*I*sin(f*x+e)*cos(f*x+e)^3*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)-21*I*sin(f*x+e)*cos(f*x+e)^3*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)+21*I*sin(f*x+e)*cos(f*x+e)^2*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)-21*I*sin(f*x+e)*cos(f*x+e)^2*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)-21*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+21*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)-21*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)+21*I*sin(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)+21*cos(f*x+e)^3-14*cos(f*x+e)^2-21*cos(f*x+e)+12)*cos(f*x+e)*(b/cos(f*x+e))^(3/2)/sin(f*x+e)^7
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((b*sec(f*x + e))^(3/2)*csc(f*x + e)^4, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 180, normalized size = 1.45

$$\frac{21\sqrt{2}(ib\cos(fx+e)^2-ib)\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e)))+21\sqrt{2}(-ib\cos(fx+e)^2+ib)\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e)))+2(21b\cos(fx+e)^4-35b\cos(fx+e)^2+12b)\sqrt{\frac{b}{\cos(fx+e)}}}{12(f\cos(fx+e)^2-f)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] -1/12*(21*sqrt(2)*(I*b*cos(f*x + e)^2 - I*b)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + 21*sqrt(2)*(-I*b*cos(f*x + e)^2 + I*b)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) + 2*(21*b*cos(f*x + e)^4 - 35*b*cos(f*x + e)^2 + 12*b)*sqrt(b/cos(f*x + e))/((f*cos(f*x + e)^2 - f)*sin(f*x + e))
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**4*(b*sec(f*x+e))**(3/2),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 8009 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(3/2),x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^(3/2)*csc(f*x + e)^4, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{3/2}}{\sin(e+fx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(e + f*x))^(3/2)/sin(e + f*x)^4,x)`

[Out] `int((b/cos(e + f*x))^(3/2)/sin(e + f*x)^4, x)`

3.397 $\int (b \sec(e + fx))^{5/2} \sin^7(e + fx) dx$

Optimal. Leaf size=85

$$\frac{2b^7}{9f(b \sec(e + fx))^{9/2}} - \frac{6b^5}{5f(b \sec(e + fx))^{5/2}} + \frac{6b^3}{f\sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

[Out] $2/9*b^7/f/(b*\sec(f*x+e))^(9/2)-6/5*b^5/f/(b*\sec(f*x+e))^(5/2)+2/3*b*(b*\sec(f*x+e))^(3/2)/f+6*b^3/f/(b*\sec(f*x+e))^(1/2)$

Rubi [A]

time = 0.04, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$\frac{2b^7}{9f(b \sec(e + fx))^{9/2}} - \frac{6b^5}{5f(b \sec(e + fx))^{5/2}} + \frac{6b^3}{f\sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^(5/2)*\text{Sin}[e + f*x]^7, x]$

[Out] $(2*b^7)/(9*f*(b*\text{Sec}[e + f*x])^(9/2)) - (6*b^5)/(5*f*(b*\text{Sec}[e + f*x])^(5/2)) + (6*b^3)/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*(b*\text{Sec}[e + f*x])^(3/2))/(3*f)$

Rule 276

$\text{Int}[(c_*)*(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x \ \&\& \ \text{IGtQ}[p, 0]$

Rule 2702

$\text{Int}[\text{csc}[(e_*) + (f_*)*(x_)]^(n_)*((a_)*\text{sec}[(e_*) + (f_*)*(x_)]^(m_)), x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2)], x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n + 1)/2] \ \&\& \ !(\text{IntegerQ}[(m + 1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{5/2} \sin^7(e + fx) dx &= \frac{b^7 \text{Subst} \left(\int \frac{(-1 + \frac{x^2}{b^2})^3}{x^{11/2}} dx, x, b \sec(e + fx) \right)}{f} \\
&= \frac{b^7 \text{Subst} \left(\int \left(-\frac{1}{x^{11/2}} + \frac{3}{b^2 x^{7/2}} - \frac{3}{b^4 x^{3/2}} + \frac{\sqrt{x}}{b^6} \right) dx, x, b \sec(e + fx) \right)}{f} \\
&= \frac{2b^7}{9f(b \sec(e + fx))^{9/2}} - \frac{6b^5}{5f(b \sec(e + fx))^{5/2}} + \frac{6b^3}{f \sqrt{b \sec(e + fx)}} + \frac{2}{f}
\end{aligned}$$

Mathematica [A]

time = 0.31, size = 52, normalized size = 0.61

$$\frac{b(2366 + 1803 \cos(2(e + fx)) - 78 \cos(4(e + fx)) + 5 \cos(6(e + fx)))(b \sec(e + fx))^{3/2}}{720f}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x]^7,x]`

```
[Out] (b*(2366 + 1803*Cos[2*(e + f*x)] - 78*Cos[4*(e + f*x)] + 5*Cos[6*(e + f*x)])
*(b*Sec[e + f*x])^(3/2))/(720*f)
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 531 vs. 2(71) = 142.

time = 0.23, size = 532, normalized size = 6.26

method	result
default	$\frac{(-1 + \cos(fx + e))^2 \left(20(\cos^6(fx + e)) - 108(\cos^4(fx + e)) - 135 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \right) \ln \left(-\frac{2 \left(2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \right)}{\dots} \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(5/2)*sin(f*x+e)^7,x,method=_RETURNVERBOSE)`

```
[Out] 1/90/f*(-1+cos(f*x+e))^2*(20*cos(f*x+e)^6-108*cos(f*x+e)^4-135*(-cos(f*x+e)
/(cos(f*x+e)+1)^(1/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^(1/2)
)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^(1/2)-1
)/sin(f*x+e)^2)*cos(f*x+e)^2+135*(-cos(f*x+e)/(cos(f*x+e)+1)^(1/2)*ln(-
(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^(1/2)-cos(f*x+e)^2+2*cos(f*x+
e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^(1/2)-1)/sin(f*x+e)^2)*cos(f*x+e)^2-135
```

$$\begin{aligned} & *(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)+135*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)+540*\cos(f*x+e)^2+60)*\cos(f*x+e)*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(5/2)}/\sin(f*x+e)^4 \end{aligned}$$

Maxima [A]

time = 0.29, size = 70, normalized size = 0.82

$$\frac{2 \left(15 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} + \frac{5b^6 - \frac{27b^6}{\cos(fx+e)^2} + \frac{135b^6}{\cos(fx+e)^4}}{\left(\frac{b}{\cos(fx+e)} \right)^{\frac{9}{2}}} \right) b}{45 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^7,x, algorithm="maxima")

[Out] $2/45*(15*(b/\cos(f*x + e))^{(3/2)} + (5*b^6 - 27*b^6/\cos(f*x + e)^2 + 135*b^6/\cos(f*x + e)^4)/(b/\cos(f*x + e))^{(9/2)})*b/f$

Fricas [A]

time = 0.40, size = 75, normalized size = 0.88

$$\frac{2(5b^2 \cos(fx+e)^6 - 27b^2 \cos(fx+e)^4 + 135b^2 \cos(fx+e)^2 + 15b^2) \sqrt{\frac{b}{\cos(fx+e)}}}{45 f \cos(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^7,x, algorithm="fricas")

[Out] $2/45*(5*b^2*\cos(f*x + e)^6 - 27*b^2*\cos(f*x + e)^4 + 135*b^2*\cos(f*x + e)^2 + 15*b^2)*\sqrt{b/\cos(f*x + e)}/(f*\cos(f*x + e))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e)**7,x)

[Out] Timed out

Giac [A]

time = 5.24, size = 108, normalized size = 1.27

$$\frac{2 \left(5 \sqrt{b \cos(fx + e)} b^4 \cos(fx + e)^4 - 27 \sqrt{b \cos(fx + e)} b^4 \cos(fx + e)^2 + 135 \sqrt{b \cos(fx + e)} b^4 + \frac{15 b^5}{\sqrt{b \cos(fx + e)} \cos(fx + e)} \right) \operatorname{sgn}(\cos(fx + e))}{45 b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^7,x, algorithm="giac")

[Out] 2/45*(5*sqrt(b*cos(f*x + e))*b^4*cos(f*x + e)^4 - 27*sqrt(b*cos(f*x + e))*b^4*cos(f*x + e)^2 + 135*sqrt(b*cos(f*x + e))*b^4 + 15*b^5/(sqrt(b*cos(f*x + e))*cos(f*x + e)))*sgn(cos(f*x + e))/(b^2*f)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^7 \left(\frac{b}{\cos(e + f x)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^7*(b/cos(e + f*x))^(5/2),x)**[Out]** int(sin(e + f*x)^7*(b/cos(e + f*x))^(5/2), x)

3.398 $\int (b \sec(e + fx))^{5/2} \sin^5(e + fx) dx$

Optimal. Leaf size=63

$$-\frac{2b^5}{5f(b \sec(e + fx))^{5/2}} + \frac{4b^3}{f\sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

[Out] $-2/5*b^5/f/(b*\sec(f*x+e))^{(5/2)}+2/3*b*(b*\sec(f*x+e))^{(3/2)}/f+4*b^3/f/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$-\frac{2b^5}{5f(b \sec(e + fx))^{5/2}} + \frac{4b^3}{f\sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(5/2)}*\text{Sin}[e + f*x]^5, x]$

[Out] $(-2*b^5)/(5*f*(b*\text{Sec}[e + f*x])^{(5/2)}) + (4*b^3)/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*(b*\text{Sec}[e + f*x])^{(3/2)})/(3*f)$

Rule 276

$\text{Int}[(c_*)*(x_)^{(m_*)}*((a_) + (b_*)*(x_)^{(n_)})^{(p_*)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n\}, x] \&\& \text{IGtQ}[p, 0]$

Rule 2702

$\text{Int}[\text{csc}[(e_*) + (f_*)*(x_)]^{(n_*)}*((a_*)*\text{sec}[(e_*) + (f_*)*(x_)]^{(m_*)}, x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1 + x^2/a^2)^{((n+1)/2)}, x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n+1)/2] \&\& !(\text{IntegerQ}[(m+1)/2] \&\& \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{5/2} \sin^5(e + fx) dx &= \frac{b^5 \text{Subst} \left(\int \frac{\left(-1 + \frac{x^2}{b^2}\right)^2}{x^{7/2}} dx, x, b \sec(e + fx) \right)}{f} \\
&= \frac{b^5 \text{Subst} \left(\int \left(\frac{1}{x^{7/2}} - \frac{2}{b^2 x^{3/2}} + \frac{\sqrt{x}}{b^4} \right) dx, x, b \sec(e + fx) \right)}{f} \\
&= -\frac{2b^5}{5f(b \sec(e + fx))^{5/2}} + \frac{4b^3}{f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.25, size = 42, normalized size = 0.67

$$\frac{b(151 + 108 \cos(2(e + fx)) - 3 \cos(4(e + fx)))(b \sec(e + fx))^{3/2}}{60f}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x]^5,x]``[Out] (b*(151 + 108*Cos[2*(e + f*x)] - 3*Cos[4*(e + f*x)])*(b*Sec[e + f*x])^(3/2))/(60*f)`**Maple [B]** Leaf count of result is larger than twice the leaf count of optimal. 521 vs. 2(53) = 106.

time = 0.26, size = 522, normalized size = 8.29

method	result
default	$ \frac{(-1 + \cos(fx + e))^2 \left(6(\cos^4(fx + e)) + 15 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \ln \left(-\frac{2 \left(2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} - (\cos^2(fx + e)) + 2 \cos(fx + e) \right)}{\sin(fx + e)^2} \right) \right)}{15f} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(5/2)*sin(f*x+e)^5,x,method=_RETURNVERBOSE)`

```
[Out] -1/15/f*(-1+cos(f*x+e))^2*(6*cos(f*x+e)^4+15*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e))^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2*cos(f*x+e)^2-15*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2*cos(f*x+e)^2+15*(-cos(f*x+e)/(cos(f
```

$*x+e)+1)^2)^{(1/2)}*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)-15*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)-60*\cos(f*x+e)^2-10)*\cos(f*x+e)*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(5/2)}/\sin(f*x+e)^4$

Maxima [A]

time = 0.30, size = 55, normalized size = 0.87

$$\frac{2 \left(5 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} - \frac{3 \left(b^4 - \frac{10b^4}{\cos(fx+e)^2} \right)}{\left(\frac{b}{\cos(fx+e)} \right)^{\frac{5}{2}}} \right) b}{15 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^5,x, algorithm="maxima")

[Out] 2/15*(5*(b/cos(f*x + e))^(3/2) - 3*(b^4 - 10*b^4/cos(f*x + e)^2)/(b/cos(f*x + e))^(5/2))*b/f

Fricas [A]

time = 0.42, size = 61, normalized size = 0.97

$$\frac{2 \left(3b^2 \cos(fx + e)^4 - 30b^2 \cos(fx + e)^2 - 5b^2 \right) \sqrt{\frac{b}{\cos(fx + e)}}}{15 f \cos(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^5,x, algorithm="fricas")

[Out] -2/15*(3*b^2*cos(f*x + e)^4 - 30*b^2*cos(f*x + e)^2 - 5*b^2)*sqrt(b/cos(f*x + e))/(f*cos(f*x + e))

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e)**5,x)

[Out] Timed out

Giac [A]

time = 3.55, size = 80, normalized size = 1.27

$$\frac{2 \left(3 \sqrt{b \cos(fx + e)} b^2 \cos(fx + e)^2 - 30 \sqrt{b \cos(fx + e)} b^2 - \frac{5b^3}{\sqrt{b \cos(fx + e)} \cos(fx + e)} \right) \operatorname{sgn}(\cos(fx + e))}{15 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^5,x, algorithm="giac")

[Out] -2/15*(3*sqrt(b*cos(f*x + e))*b^2*cos(f*x + e)^2 - 30*sqrt(b*cos(f*x + e))*
b^2 - 5*b^3/(sqrt(b*cos(f*x + e))*cos(f*x + e)))*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + f x)^5 \left(\frac{b}{\cos(e + f x)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^5*(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^5*(b/cos(e + f*x))^(5/2), x)

3.399 $\int (b \sec(e + fx))^{5/2} \sin^3(e + fx) dx$

Optimal. Leaf size=41

$$\frac{2b^3}{f\sqrt{b\sec(e+fx)}} + \frac{2b(b\sec(e+fx))^{3/2}}{3f}$$

[Out] $2/3*b*(b*\sec(f*x+e))^(3/2)/f+2*b^3/f/(b*\sec(f*x+e))^(1/2)$

Rubi [A]

time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 14}

$$\frac{2b^3}{f\sqrt{b\sec(e+fx)}} + \frac{2b(b\sec(e+fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^(5/2)*\text{Sin}[e + f*x]^3, x]$

[Out] $(2*b^3)/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*(b*\text{Sec}[e + f*x])^(3/2))/(3*f)$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^(m_), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m, x\} \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{InverseFunctionQ}[v]]$

Rule 2702

$\text{Int}[\text{csc}[e_] + (f_)*(x_)]^(n_)*((a_)*\sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^(m+n-1)/(-1+x^2/a^2)^(n+1)/2], x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}\{a, e, f, m, x\} \ \&\& \ \text{IntegerQ}[(n+1)/2] \ \&\& \ !(\text{IntegerQ}[(m+1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^{5/2} \sin^3(e + fx) dx &= \frac{b^3 \text{Subst}\left(\int \frac{-1 + \frac{x^2}{b^2}}{x^{3/2}} dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{b^3 \text{Subst}\left(\int \left(-\frac{1}{x^{3/2}} + \frac{\sqrt{x}}{b^2}\right) dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{2b^3}{f\sqrt{b\sec(e+fx)}} + \frac{2b(b\sec(e+fx))^{3/2}}{3f} \end{aligned}$$

Mathematica [A]

time = 0.15, size = 32, normalized size = 0.78

$$\frac{b(5 + 3 \cos(2(e + fx)))(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x]^3,x]

[Out] (b*(5 + 3*Cos[2*(e + f*x)])*(b*Sec[e + f*x])^(3/2))/(3*f)

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 356 vs. 2(35) = 70.

time = 0.22, size = 357, normalized size = 8.71

method	result
default	$(-1 + \cos(fx + e)) \left(12(\cos^3(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} - 3(\cos^2(fx + e)) \ln \left(-\frac{2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} - (\cos^2(fx + e))}{\sin(fx + e)} \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(5/2)*sin(f*x+e)^3,x,method=_RETURNVERBOSE)

[Out]
$$-1/6/f*(-1+\cos(f*x+e))*(12*\cos(f*x+e)^3*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - 3*\cos(f*x+e)^2*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - \cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+3*\cos(f*x+e)^2*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - \cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+12*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+4*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+4*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2))*\cos(f*x+e)*(b/\cos(f*x+e))^{(5/2)}/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}/\sin(f*x+e)^2$$

Maxima [A]

time = 0.28, size = 38, normalized size = 0.93

$$\frac{2 \left(\frac{3b^2}{\sqrt{\frac{b}{\cos(fx + e)}}} + \left(\frac{b}{\cos(fx + e)} \right)^{\frac{3}{2}} \right) b}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^3,x, algorithm="maxima")

[Out] $2/3*(3*b^2/\sqrt{b/\cos(f*x + e)} + (b/\cos(f*x + e))^{(3/2)})*b/f$

Fricas [A]

time = 0.38, size = 45, normalized size = 1.10

$$\frac{2(3b^2 \cos(fx + e)^2 + b^2) \sqrt{\frac{b}{\cos(fx + e)}}}{3f \cos(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^3,x, algorithm="fricas")

[Out] $2/3*(3*b^2*\cos(f*x + e)^2 + b^2)*\sqrt{b/\cos(f*x + e)}/(f*\cos(f*x + e))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e)**3,x)

[Out] Timed out

Giac [A]

time = 4.45, size = 53, normalized size = 1.29

$$\frac{2 \left(3 \sqrt{b \cos(fx + e)} b + \frac{b^2}{\sqrt{b \cos(fx + e)} \cos(fx + e)} \right) b \operatorname{sgn}(\cos(fx + e))}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^3,x, algorithm="giac")

[Out] $2/3*(3*\sqrt{b*\cos(f*x + e)}*b + b^2/(\sqrt{b*\cos(f*x + e)}*\cos(f*x + e)))*b*\operatorname{sgn}(\cos(f*x + e))/f$

Mupad [B]

time = 0.91, size = 50, normalized size = 1.22

$$\frac{b^2 \sqrt{\frac{b}{\cos(e + fx)}} \left(\frac{13 \cos(e + fx)}{3} + \cos(3e + 3fx) \right)}{f (\cos(2e + 2fx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^3*(b/cos(e + f*x))^(5/2),x)

[Out] $(b^2*(b/\cos(e + f*x))^{(1/2)}*((13*\cos(e + f*x))/3 + \cos(3*e + 3*f*x)))/(f*(\cos(2*e + 2*f*x) + 1))$

3.400 $\int (b \sec(e + fx))^{5/2} \sin(e + fx) dx$

Optimal. Leaf size=20

$$\frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

[Out] $2/3*b*(b*\sec(f*x+e))^{(3/2)}/f$

Rubi [A]

time = 0.02, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 30}

$$\frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(5/2)}*\text{Sin}[e + f*x],x]$

[Out] $(2*b*(b*\text{Sec}[e + f*x])^{(3/2)})/(3*f)$

Rule 30

$\text{Int}[(x_)^{(m_.)}, x_Symbol] \text{ :> Simp}[x^{(m + 1)}/(m + 1), x] \text{ /; FreeQ}[m, x] \ \&\& \ \text{N eQ}[m, -1]$

Rule 2702

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^{(n_.)}*((a_.)*\sec[(e_.) + (f_.)*(x_)]^{(m_.)}, x_Symbol] \text{ :> Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m + n - 1)}/(-1 + x^2/a^2)^{((n + 1)/2)}, x], x, a*\text{Sec}[e + f*x]], x] \text{ /; FreeQ}[\{a, e, f, m\}, x] \ \&\& \ \text{IntegerQ}[(n + 1)/2] \ \&\& \ !(\text{IntegerQ}[(m + 1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^{5/2} \sin(e + fx) dx &= \frac{b \text{Subst}(\int \sqrt{x} dx, x, b \sec(e + fx))}{f} \\ &= \frac{2b(b \sec(e + fx))^{3/2}}{3f} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 20, normalized size = 1.00

$$\frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x],x]

[Out] (2*b*(b*Sec[e + f*x])^(3/2))/(3*f)

Maple [A]

time = 0.04, size = 17, normalized size = 0.85

method	result	size
derivativedivides	$\frac{2b(b \sec(fx+e))^{\frac{3}{2}}}{3f}$	17
default	$\frac{2b(b \sec(fx+e))^{\frac{3}{2}}}{3f}$	17

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(5/2)*sin(f*x+e),x,method=_RETURNVERBOSE)

[Out] 2/3*b*(b*sec(f*x+e))^(3/2)/f

Maxima [A]

time = 0.28, size = 25, normalized size = 1.25

$$\frac{2 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{5}{2}} \cos(fx+e)}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e),x, algorithm="maxima")

[Out] 2/3*(b/cos(f*x + e))^(5/2)*cos(f*x + e)/f

Fricas [A]

time = 0.41, size = 30, normalized size = 1.50

$$\frac{2b^2 \sqrt{\frac{b}{\cos(fx+e)}}}{3f \cos(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e),x, algorithm="fricas")

[Out] 2/3*b^2*sqrt(b/cos(f*x + e))/(f*cos(f*x + e))

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e),x)`

[Out] Timed out

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(17) = 34.
time = 4.58, size = 36, normalized size = 1.80

$$\frac{2b^3 \operatorname{sgn}(\cos(fx + e))}{3 \sqrt{b \cos(fx + e)} f \cos(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e),x, algorithm="giac")`

[Out] `2/3*b^3*sgn(cos(f*x + e))/(sqrt(b*cos(f*x + e))*f*cos(f*x + e))`

Mupad [B]

time = 0.62, size = 39, normalized size = 1.95

$$\frac{4b^2 \cos(e + fx) \sqrt{\frac{b}{\cos(e + fx)}}}{3f (\cos(2e + 2fx) + 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)*(b/cos(e + f*x))^(5/2),x)`

[Out] `(4*b^2*cos(e + f*x)*(b/cos(e + f*x))^(1/2))/(3*f*(cos(2*e + 2*f*x) + 1))`

3.401 $\int \csc(e + fx)(b \sec(e + fx))^{5/2} dx$

Optimal. Leaf size=78

$$\frac{b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

[Out] $b^{(5/2)*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f - b^{(5/2)*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f + 2/3*b*(b*\sec(f*x+e))^{(3/2)}/f}$

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2702, 327, 335, 304, 209, 212}

$$\frac{b^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{2b(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[e + f*x]*(b*\operatorname{Sec}[e + f*x])^{(5/2)}, x]$

[Out] $(b^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/f - (b^{(5/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/f + (2*b*(b*\operatorname{Sec}[e + f*x])^{(3/2)})/(3*f))$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 304

$\operatorname{Int}[(x_)^2/((a_ + (b_)*(x_)^4), x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r + s*x^2), x], x] - \operatorname{Dist}[s/(2*b), \operatorname{Int}[1/(r - s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a/b, 0]$

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \csc(e + fx)(b \sec(e + fx))^{5/2} dx &= \frac{\text{Subst}\left(\int \frac{x^{5/2}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{bf} \\
&= \frac{2b(b \sec(e + fx))^{3/2}}{3f} + \frac{b \text{Subst}\left(\int \frac{\sqrt{x}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{f} \\
&= \frac{2b(b \sec(e + fx))^{3/2}}{3f} + \frac{(2b) \text{Subst}\left(\int \frac{x^2}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} \\
&= \frac{2b(b \sec(e + fx))^{3/2}}{3f} - \frac{b^3 \text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} + \frac{b^3 \text{Subst}\left(\int \frac{1}{b + x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} \\
&= \frac{b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} - \frac{b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f} + \frac{2b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{f}
\end{aligned}$$

Mathematica [A]

time = 0.14, size = 87, normalized size = 1.12

$$\frac{(b \sec(e + fx))^{5/2} \left(6 \tan^{-1}\left(\sqrt{\sec(e + fx)}\right) + 3 \log\left(1 - \sqrt{\sec(e + fx)}\right) - 3 \log\left(1 + \sqrt{\sec(e + fx)}\right) + 4 \sec^{3/2}(e + fx)\right)}{6f \sec^{5/2}(e + fx)}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]*(b*Sec[e + f*x])^(5/2),x]

[Out] ((b*Sec[e + f*x])^(5/2)*(6*ArcTan[Sqrt[Sec[e + f*x]]] + 3*Log[1 - Sqrt[Sec[e + f*x]]] - 3*Log[1 + Sqrt[Sec[e + f*x]]] + 4*Sec[e + f*x]^(3/2)))/(6*f*Sec[e + f*x]^(5/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 236 vs. 2(62) = 124.

time = 0.22, size = 237, normalized size = 3.04

method	result
default	$\frac{(-1 + \cos(fx + e)) \left(3(\cos^2(fx + e)) \ln \left(-\frac{2 \left(2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} - (\cos^2(fx + e)) + 2\cos(fx + e) - 2 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \right)}{\sin(fx + e)^2} \right)}{6f \sqrt{-\dots}} \right)}{6f \sqrt{-\dots}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)*(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)

[Out] 1/6/f*(-1+cos(f*x+e))*(3*cos(f*x+e)^2*ln(-2*(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)-3*cos(f*x+e)^2*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))-4*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-4*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))*cos(f*x+e)*(b/cos(f*x+e))^(5/2)/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)/sin(f*x+e)^2

Maxima [A]

time = 0.51, size = 91, normalized size = 1.17

$$\frac{\left(6 b^{\frac{3}{2}} \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx + e)}}}{\sqrt{b}} \right) + 3 b^{\frac{3}{2}} \log \left(-\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx + e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx + e)}}} \right) + 4 \left(\frac{b}{\cos(fx + e)} \right)^{\frac{3}{2}} b \right)}{6 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] 1/6*(6*b^(3/2)*arctan(sqrt(b/cos(f*x + e))/sqrt(b)) + 3*b^(3/2)*log(-(sqrt(b) - sqrt(b/cos(f*x + e)))/(sqrt(b) + sqrt(b/cos(f*x + e)))) + 4*(b/cos(f*x + e))^(3/2))*b/f

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 167 vs. 2(65) = 130.

time = 0.46, size = 354, normalized size = 4.54

$$\frac{6\sqrt{-b}^b \arctan\left(\frac{\sqrt{-b}}{\cos(fx+e)}\right) \cos(fx+e) + 3\sqrt{-b}^b \cos(fx+e) \log\left(\frac{b \cos(fx+e)^2 - 4(\cos(fx+e)^2 - \cos(fx+e))\sqrt{-b} \sqrt{\frac{b}{\cos(fx+e)}} - 4b \cos(fx+e)}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right) + 8b^b \sqrt{\frac{b}{\cos(fx+e)}}}{12f \cos(fx+e)} - \frac{6b^b \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) \cos(fx+e) - 3b^b \cos(fx+e) \log\left(\frac{b \cos(fx+e)^2 - 4(\cos(fx+e)^2 - \cos(fx+e))\sqrt{b} \sqrt{\frac{b}{\cos(fx+e)}} - 4b \cos(fx+e)}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right) - 8b^b \sqrt{\frac{b}{\cos(fx+e)}}}{12f \cos(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] [1/12*(6*sqrt(-b)*b^2*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b)*cos(f*x + e) + 3*sqrt(-b)*b^2*cos(f*x + e)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) + 8*b^2*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e)), -1/12*(6*b^(5/2)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b))*cos(f*x + e) - 3*b^(5/2)*cos(f*x + e)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)) - 8*b^2*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e))]

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(5/2),x)

[Out] Timed out

Giac [A]

time = 4.31, size = 91, normalized size = 1.17

$$\frac{b^6 \left(\frac{3 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^3} - \frac{3 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{b}}\right)}{b^{\frac{7}{2}}} + \frac{2}{\sqrt{b \cos(fx+e)} b^3 \cos(fx+e)} \right) \operatorname{sgn}(\cos(fx+e))}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] 1/3*b^6*(3*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b)*b^3 - 3*arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(7/2) + 2/(sqrt(b*cos(f*x + e))*b^3*cos(f*x + e)))*sgn(cos(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{5/2}}{\sin(e+fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(5/2)/sin(e + f*x),x)

[Out] int((b/cos(e + f*x))^(5/2)/sin(e + f*x), x)

3.402 $\int \csc^3(e + fx)(b \sec(e + fx))^{5/2} dx$

Optimal. Leaf size=113

$$\frac{7b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{7b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} + \frac{7b(b \sec(e + fx))^{3/2}}{6f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{7/2}}{2bf}$$

[Out] $7/4*b^{(5/2)*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f-7/4*b^{(5/2)*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f+7/6*b*(b*\sec(f*x+e))^{(3/2)}/f-1/2*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(7/2)}/b/f}$

Rubi [A]

time = 0.06, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2702, 294, 327, 335, 304, 209, 212}

$$\frac{7b^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} - \frac{7b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4f} + \frac{7b(b \sec(e + fx))^{3/2}}{6f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{7/2}}{2bf}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[e + f*x]^3*(b*\operatorname{Sec}[e + f*x])^{(5/2)}, x]$

[Out] $(7*b^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(4*f) - (7*b^{(5/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(4*f) + (7*b*(b*\operatorname{Sec}[e + f*x])^{(3/2)})/(6*f) - (\operatorname{Cot}[e + f*x]^2*(b*\operatorname{Sec}[e + f*x])^{(7/2)})/(2*b*f)$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[c^{(n - 1)}*(c*x)^{(m - n + 1)}*((a + b*x^n)^{(p + 1)}/(b*n*(p + 1))), x] - \operatorname{Dist}[c^{(n - 1)}*((m - n + 1)/(b*n*(p + 1))), \operatorname{Int}[(c*x)^{(m - n)}*(a + b*x^n)^{(p + 1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[p, -1] \ \&\& \operatorname{GtQ}[m + 1, n] \ \&\& \operatorname{!} \operatorname{LtQ}[m + n*(p + 1) + 1, n, 0] \ \&\& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 327

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n
- 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \csc^3(e+fx)(b \sec(e+fx))^{5/2} dx &= \frac{\text{Subst}\left(\int \frac{x^{9/2}}{\left(-1+\frac{x^2}{b^2}\right)^2} dx, x, b \sec(e+fx)\right)}{b^3 f} \\
&= -\frac{\cot^2(e+fx)(b \sec(e+fx))^{7/2}}{2bf} + \frac{7 \text{Subst}\left(\int \frac{x^{5/2}}{-1+\frac{x^2}{b^2}} dx, x, b \sec(e+fx)\right)}{4bf} \\
&= \frac{7b(b \sec(e+fx))^{3/2}}{6f} - \frac{\cot^2(e+fx)(b \sec(e+fx))^{7/2}}{2bf} + \frac{(7b) \text{Subst}\left(\int \frac{x^{3/2}}{-1+\frac{x^2}{b^2}} dx, x, b \sec(e+fx)\right)}{4bf} \\
&= \frac{7b(b \sec(e+fx))^{3/2}}{6f} - \frac{\cot^2(e+fx)(b \sec(e+fx))^{7/2}}{2bf} + \frac{(7b) \text{Subst}\left(\int \frac{x^{1/2}}{-1+\frac{x^2}{b^2}} dx, x, b \sec(e+fx)\right)}{4bf} \\
&= \frac{7b(b \sec(e+fx))^{3/2}}{6f} - \frac{\cot^2(e+fx)(b \sec(e+fx))^{7/2}}{2bf} - \frac{(7b^3) \text{Subst}\left(\int \frac{1}{-1+\frac{x^2}{b^2}} dx, x, b \sec(e+fx)\right)}{4bf} \\
&= \frac{7b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4f} - \frac{7b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4f}
\end{aligned}$$

Mathematica [A]

time = 1.36, size = 109, normalized size = 0.96

$$\frac{b^3(-12 \csc^2(e+fx) + 42 \tan^{-1}(\sqrt{\sec(e+fx)}) \sqrt{\sec(e+fx)} + 21(\log(1 - \sqrt{\sec(e+fx)}) - \log(1 + \sqrt{\sec(e+fx)})) \sqrt{\sec(e+fx)} + 16 \sec^2(e+fx))}{24f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3*(b*Sec[e + f*x])^(5/2), x]

[Out] (b^3*(-12*Csc[e + f*x]^2 + 42*ArcTan[Sqrt[Sec[e + f*x]]]*Sqrt[Sec[e + f*x]] + 21*(Log[1 - Sqrt[Sec[e + f*x]]] - Log[1 + Sqrt[Sec[e + f*x]]])*Sqrt[Sec[e + f*x]] + 16*Sec[e + f*x]^2))/(24*f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 698 vs. 2(89) = 178.

time = 0.32, size = 699, normalized size = 6.19

method	result
--------	--------

default	$\frac{(-1+\cos(fx+e)) \left(24 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} (\cos^4(fx+e)) + 48 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} (\cos^3(fx+e)) + 24 (\cos^2(fx+e)) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right) \right)}{\dots}$
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^3*(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/24/f*(-1+\cos(f*x+e))*(24*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\cos(f*x+e) \\ & ^4+48*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\cos(f*x+e)^3+24*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)} \\ & +24*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^4 \\ & -21*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})*\cos(f*x+e)^4-3*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^4 \\ & -4*\cos(f*x+e)^3*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-28*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-24*\cos(f*x+e)^2*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+21*\cos(f*x+e)^2*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}))+3*\cos(f*x+e)^2*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+16*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+16*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})*\cos(f*x+e)*(b/\cos(f*x+e))^{(5/2)}/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}/\sin(f*x+e)^4 \end{aligned}$$

Maxima [A]

time = 0.49, size = 129, normalized size = 1.14

$$\frac{\left(\frac{12b^2 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}}}{b^2 - \frac{b^2}{\cos(fx+e)^2}} + 42b^{\frac{3}{2}} \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right) + 21b^{\frac{3}{2}} \log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right) + 16 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} \right) b}{24f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out]
$$\begin{aligned} & 1/24*(12*b^2*(b/\cos(f*x + e))^{(3/2)}/(b^2 - b^2/\cos(f*x + e)^2) + 42*b^{(3/2)} \\ & *arctan(sqrt(b/\cos(f*x + e))/sqrt(b)) + 21*b^{(3/2)}*log(-sqrt(b) - sqrt(b/\cos(f*x + e)))/sqrt(b) + sqrt(b/\cos(f*x + e))) + 16*(b/\cos(f*x + e))^{(3/2)} \\ &)*b/f \end{aligned}$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 234 vs. 2(94) = 188.

time = 0.51, size = 482, normalized size = 4.27

$$\frac{\frac{42 \sqrt{b \cos(fx + e)} \sqrt{-b} \arctan\left(\frac{\sqrt{-b} \sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right) + 21 \sqrt{b \cos(fx + e)} \sqrt{-b} \log\left(\frac{\sqrt{b \cos(fx + e)} \sqrt{-b}}{\sqrt{-b}}\right) + 8 \sqrt{b \cos(fx + e)} \sqrt{-b} \log\left(\frac{\sqrt{b \cos(fx + e)} \sqrt{-b}}{\sqrt{-b}}\right) + 8 \sqrt{b \cos(fx + e)} \sqrt{-b} \log\left(\frac{\sqrt{b \cos(fx + e)} \sqrt{-b}}{\sqrt{-b}}\right)}{48 \sqrt{b \cos(fx + e)} \sqrt{-b}} \dots$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] [1/48*(42*(b^2*cos(f*x + e))^3 - b^2*cos(f*x + e))*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b) + 21*(b^2*cos(f*x + e))^3 - b^2*cos(f*x + e))*sqrt(-b)*log((b*cos(f*x + e))^2 - 4*(cos(f*x + e))^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1) + 8*(7*b^2*cos(f*x + e)^2 - 4*b^2)*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e)^3 - f*cos(f*x + e)), -1/48*(42*(b^2*cos(f*x + e))^3 - b^2*cos(f*x + e))*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b)) - 21*(b^2*cos(f*x + e))^3 - b^2*cos(f*x + e))*sqrt(b)*log((b*cos(f*x + e))^2 - 4*(cos(f*x + e))^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1) - 8*(7*b^2*cos(f*x + e)^2 - 4*b^2)*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e)^3 - f*cos(f*x + e))]

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3*(b*sec(f*x+e))**(5/2),x)

[Out] Timed out

Giac [A]

time = 4.33, size = 128, normalized size = 1.13

$$\frac{b^8 \left(\frac{6 \sqrt{b \cos(fx + e)}}{(b^2 \cos(fx + e)^2 - b^2) b^4} + \frac{21 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^5} - \frac{21 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{b^{\frac{11}{2}} \sqrt{b}} + \frac{8}{\sqrt{b \cos(fx + e)} b^5 \cos(fx + e)} \right) \operatorname{sgn}(\cos(fx + e))}{12 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] 1/12*b^8*(6*sqrt(b*cos(f*x + e))/((b^2*cos(f*x + e))^2 - b^2)*b^4) + 21*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b)*b^5) - 21*arctan(sqrt(b*cos(f*x

+ e))/sqrt(b))/b^(11/2) + 8/(sqrt(b*cos(f*x + e))*b^5*cos(f*x + e))*sgn(c
os(f*x + e))/f

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{5/2}}{\sin(e+fx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^3,x)

[Out] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^3, x)

3.403 $\int \csc^5(e + fx)(b \sec(e + fx))^{5/2} dx$

Optimal. Leaf size=143

$$\frac{77b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{77b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} + \frac{77b(b \sec(e + fx))^{3/2}}{48f} - \frac{11 \cot^2(e + fx)}{16bf}$$

[Out] $77/32*b^{(5/2)*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f-77/32*b^{(5/2)*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f+77/48*b*(b*\sec(f*x+e))^{(3/2)}/f-11/16*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(7/2)}/b/f-1/4*\cot(f*x+e)^4*(b*\sec(f*x+e))^{(11/2)}/b^3/f}$

Rubi [A]

time = 0.07, antiderivative size = 143, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2702, 294, 327, 335, 304, 209, 212}

$$\frac{77b^{5/2} \operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{77b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{11/2}}{4b^3 f} + \frac{77b(b \sec(e + fx))^{3/2}}{48f} - \frac{11 \cot^2(e + fx)(b \sec(e + fx))^{7/2}}{16bf}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[e + f*x]^5*(b*\operatorname{Sec}[e + f*x])^{(5/2)}, x]$

[Out] $(77*b^{(5/2)*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(32*f) - (77*b^{(5/2)*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(32*f) + (77*b*(b*\operatorname{Sec}[e + f*x])^{(3/2)})/(48*f) - (11*\operatorname{Cot}[e + f*x]^2*(b*\operatorname{Sec}[e + f*x])^{(7/2)})/(16*b*f) - (\operatorname{Cot}[e + f*x]^4*(b*\operatorname{Sec}[e + f*x])^{(11/2)})/(4*b^3*f)}$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{PosQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 294

$\operatorname{Int}[(c_)*(x_)^{(m_)}*((a_ + (b_)*(x_)^{(n_)})^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[c^{(n-1)}*(c*x)^{(m-n+1)}*((a + b*x^n)^{(p+1)}/(b*n*(p+1))), x] - \operatorname{Dist}[c^{(n-1)}*((m-n+1)/(b*n*(p+1))), \operatorname{Int}[(c*x)^{(m-n)}*(a + b*x^n)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, b, c\}, x] \ \&\& \operatorname{IGtQ}[n, 0] \ \&\& \operatorname{LtQ}[p, -1] \ \&\& \operatorname{GtQ}[m+1, n] \ \&\& \operatorname{!}I$

LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 327

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 335

Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned}
\int \csc^5(e + fx)(b \sec(e + fx))^{5/2} dx &= \frac{\text{Subst}\left(\int \frac{x^{13/2}}{\left(-1 + \frac{x^2}{b^2}\right)^3} dx, x, b \sec(e + fx)\right)}{b^5 f} \\
&= -\frac{\cot^4(e + fx)(b \sec(e + fx))^{11/2}}{4b^3 f} + \frac{11 \text{Subst}\left(\int \frac{x^{9/2}}{\left(-1 + \frac{x^2}{b^2}\right)^2} dx, x, b \sec(e + fx)\right)}{8b^3 f} \\
&= -\frac{11 \cot^2(e + fx)(b \sec(e + fx))^{7/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{11/2}}{4b^3 f} \\
&= \frac{77b(b \sec(e + fx))^{3/2}}{48f} - \frac{11 \cot^2(e + fx)(b \sec(e + fx))^{7/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{11/2}}{4b^3 f} \\
&= \frac{77b(b \sec(e + fx))^{3/2}}{48f} - \frac{11 \cot^2(e + fx)(b \sec(e + fx))^{7/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{11/2}}{4b^3 f} \\
&= \frac{77b(b \sec(e + fx))^{3/2}}{48f} - \frac{11 \cot^2(e + fx)(b \sec(e + fx))^{7/2}}{16bf} - \frac{\cot^4(e + fx)(b \sec(e + fx))^{11/2}}{4b^3 f} \\
&= \frac{77b^{5/2} \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f} - \frac{77b^{5/2} \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{32f}
\end{aligned}$$

Mathematica [A]

time = 0.93, size = 119, normalized size = 0.83

$$\frac{b^3(-180 \csc^2(e + fx) - 48 \csc^4(e + fx) + 462 \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right) \sqrt{\sec(e + fx)} + 231(\log(1 - \sqrt{\sec(e + fx)}) - \log(1 + \sqrt{\sec(e + fx)})) \sqrt{\sec(e + fx)} + 128 \sec^2(e + fx))}{192f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^5*(b*Sec[e + f*x])^(5/2), x]

[Out] (b^3*(-180*Csc[e + f*x]^2 - 48*Csc[e + f*x]^4 + 462*ArcTan[Sqrt[Sec[e + f*x]]]*Sqrt[Sec[e + f*x]] + 231*(Log[1 - Sqrt[Sec[e + f*x]]] - Log[1 + Sqrt[Sec[e + f*x]]])*Sqrt[Sec[e + f*x]] + 128*Sec[e + f*x]^2))/(192*f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1160 vs. 2(115) = 230.

time = 0.20, size = 1161, normalized size = 8.12

method	result	size
default	Expression too large to display	1161

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^5*(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/192/f*(408*\cos(f*x+e)^5*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}+360*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\cos(f*x+e)^4+288*\cos(f*x+e)^5*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-231*\cos(f*x+e)^5*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})-57*\cos(f*x+e)^5*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-504*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\cos(f*x+e)^3+100*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}*\cos(f*x+e)^4-288*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^4+231*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})*\cos(f*x+e)^4+57*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^4-456*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}-456*\cos(f*x+e)^3*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-288*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^3+231*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})*\cos(f*x+e)^3+57*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^3+484*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+288*\cos(f*x+e)^2*\ln(-2*(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-231*\cos(f*x+e)^2*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})-57*\cos(f*x+e)^2*\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-128*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}*\cos(f*x+e)*(b/\cos(f*x+e))^{(5/2)}/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}/\sin(f*x+e)^4 \end{aligned}$$

Maxima [A]

time = 0.50, size = 163, normalized size = 1.14

$$\frac{\left(462 b^{\frac{3}{2}} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + 231 b^{\frac{3}{2}} \log\left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}}\right) + 128 \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}} + \frac{12 \left(15 b^4 \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}} - 19 b^2 \left(\frac{b}{\cos(fx+e)}\right)^{\frac{7}{2}}\right)}{b^4 - \frac{2 b^4}{\cos(fx+e)^2} + \frac{b^4}{\cos(fx+e)^4}} \right) b}{192 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5*(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] $\frac{1}{192} \cdot (462 \cdot b^{3/2} \cdot \arctan(\sqrt{b/\cos(fx+e)})/\sqrt{b}) + 231 \cdot b^{3/2} \cdot \log(-(\sqrt{b} - \sqrt{b/\cos(fx+e)})/(\sqrt{b} + \sqrt{b/\cos(fx+e)})) + 128 \cdot (b/\cos(fx+e))^{3/2} + 12 \cdot (15 \cdot b^4 \cdot (b/\cos(fx+e))^{3/2} - 19 \cdot b^2 \cdot (b/\cos(fx+e))^{7/2}) / (b^4 - 2 \cdot b^4/\cos(fx+e)^2 + b^4/\cos(fx+e)^4) \cdot b/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(122) = 244.

time = 0.51, size = 584, normalized size = 4.08

$$\frac{\left(\frac{1}{192} \cdot (462 \cdot b^{3/2} \cdot \arctan(\sqrt{b/\cos(fx+e)})/\sqrt{b}) + 231 \cdot b^{3/2} \cdot \log(-(\sqrt{b} - \sqrt{b/\cos(fx+e)})/(\sqrt{b} + \sqrt{b/\cos(fx+e)})) + 128 \cdot (b/\cos(fx+e))^{3/2} + 12 \cdot (15 \cdot b^4 \cdot (b/\cos(fx+e))^{3/2} - 19 \cdot b^2 \cdot (b/\cos(fx+e))^{7/2}) / (b^4 - 2 \cdot b^4/\cos(fx+e)^2 + b^4/\cos(fx+e)^4) \cdot b/f \right)}{b^4 - 2 \cdot b^4/\cos(fx+e)^2 + b^4/\cos(fx+e)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5*(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] $\left[\frac{1}{384} \cdot (462 \cdot (b^2 \cdot \cos(fx+e))^5 - 2 \cdot b^2 \cdot \cos(fx+e)^3 + b^2 \cdot \cos(fx+e)) \cdot \sqrt{-b} \cdot \arctan(1/2 \cdot \sqrt{-b} \cdot \sqrt{b/\cos(fx+e)}) \cdot (\cos(fx+e) + 1)/b + 2 \cdot 31 \cdot (b^2 \cdot \cos(fx+e))^5 - 2 \cdot b^2 \cdot \cos(fx+e)^3 + b^2 \cdot \cos(fx+e)) \cdot \sqrt{-b} \cdot \log((b \cdot \cos(fx+e))^2 - 4 \cdot (\cos(fx+e))^2 - \cos(fx+e)) \cdot \sqrt{-b} \cdot \sqrt{b/\cos(fx+e)} - 6 \cdot b \cdot \cos(fx+e) + b) / (\cos(fx+e)^2 + 2 \cdot \cos(fx+e) + 1) + 8 \cdot (77 \cdot b^2 \cdot \cos(fx+e)^4 - 121 \cdot b^2 \cdot \cos(fx+e)^2 + 32 \cdot b^2) \cdot \sqrt{b/\cos(fx+e)} / (f \cdot \cos(fx+e)^5 - 2 \cdot f \cdot \cos(fx+e)^3 + f \cdot \cos(fx+e)), -1/384 \cdot (462 \cdot (b^2 \cdot \cos(fx+e))^5 - 2 \cdot b^2 \cdot \cos(fx+e)^3 + b^2 \cdot \cos(fx+e)) \cdot \sqrt{b} \cdot \arctan(1/2 \cdot \sqrt{b/\cos(fx+e)}) \cdot (\cos(fx+e) - 1)/\sqrt{b} - 231 \cdot (b^2 \cdot \cos(fx+e))^5 - 2 \cdot b^2 \cdot \cos(fx+e)^3 + b^2 \cdot \cos(fx+e)) \cdot \sqrt{b} \cdot \log((b \cdot \cos(fx+e))^2 - 4 \cdot (\cos(fx+e))^2 + \cos(fx+e)) \cdot \sqrt{b} \cdot \sqrt{b/\cos(fx+e)} + 6 \cdot b \cdot \cos(fx+e) + b) / (\cos(fx+e)^2 - 2 \cdot \cos(fx+e) + 1) - 8 \cdot (77 \cdot b^2 \cdot \cos(fx+e)^4 - 121 \cdot b^2 \cdot \cos(fx+e)^2 + 32 \cdot b^2) \cdot \sqrt{b/\cos(fx+e)} / (f \cdot \cos(fx+e)^5 - 2 \cdot f \cdot \cos(fx+e)^3 + f \cdot \cos(fx+e)) \right]$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**5*(b*sec(f*x+e))**(5/2),x)

[Out] Timed out

Giac [A]

time = 3.53, size = 159, normalized size = 1.11

$$b^{10} \left(\frac{231 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b} \cdot b^7} - \frac{231 \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{b}}\right)}{b^{1/2}} + \frac{6 \left(15 \sqrt{b \cos(fx+e)} \cdot b^2 \cos^2(fx+e) - 19 \sqrt{b \cos(fx+e)} \cdot b^2 \right)}{(b^2 \cos^2(fx+e) - b^2)^2 \cdot b^6} + \frac{64}{\sqrt{b \cos(fx+e)} \cdot b^7 \cos(fx+e)} \right) \operatorname{sgn}(\cos(fx+e))$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5*(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] $\frac{1}{96}b^{10} \left(\frac{231 \arctan(\sqrt{b \cos(fx + e)})/\sqrt{-b}}{\sqrt{-b}b^7} - 231 \arctan(\sqrt{b \cos(fx + e)})/\sqrt{b} \right) / b^{15/2} + 6 \left(\frac{15 \sqrt{b \cos(fx + e)} b^2 \cos(fx + e)^2 - 19 \sqrt{b \cos(fx + e)} b^2}{(b^2 \cos(fx + e)^2 - b^2)^2 b^6} + 64 / (\sqrt{b \cos(fx + e)} b^7 \cos(fx + e)) \right) \operatorname{sgn}(\cos(fx + e)) / f$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)} \right)^{5/2}}{\sin(e+fx)^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^5,x)

[Out] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^5, x)

3.404 $\int (b \sec(e + fx))^{5/2} \sin^6(e + fx) dx$

Optimal. Leaf size=130

$$-\frac{80b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{21f} + \frac{40b^3 \sin(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{20b^3 \sin^3(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^5(e + fx)}{3f}$$

[Out] $2/3*b*(b*\sec(f*x+e))^{(3/2)}*\sin(f*x+e)^5/f+40/21*b^3*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}+20/21*b^3*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(1/2)}-80/21*b^2*(\cos(1/2*f*x+1/2*e))^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.10, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2704, 2707, 3856, 2720}

$$\frac{20b^3 \sin^3(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{40b^3 \sin(e + fx)}{21f \sqrt{b \sec(e + fx)}} - \frac{80b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{21f} + \frac{2b \sin^5(e + fx)(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(5/2)}*\text{Sin}[e + f*x]^6,x]$

[Out] $(-80*b^2*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(21*f) + (40*b^3*\text{Sin}[e + f*x])/(21*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (20*b^3*\text{Sin}[e + f*x]^3)/(21*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^5)/(3*f)$

Rule 2704

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(f*a*(n - 1))), x] + \text{Dist}[b^2*((m + 1)/(a^2*(n - 1))), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{IntegersQ}[2*m, 2*n]$

Rule 2707

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(a*f*(m + n))), x] + \text{Dist}[(m + 1)/(a^2*(m + n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x \ \&\& \ \text{LtQ}[m, -1] \ \&\& \ \text{NeQ}[m + n, 0] \ \&\& \ \text{IntegersQ}[2*m, 2*n]$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int (b \sec(e + fx))^{5/2} \sin^6(e + fx) dx &= \frac{2b(b \sec(e + fx))^{3/2} \sin^5(e + fx)}{3f} - \frac{1}{3}(10b^2) \int \sqrt{b \sec(e + fx)} \sin^4(e + fx) dx \\
 &= \frac{20b^3 \sin^3(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^5(e + fx)}{3f} - \frac{1}{7}(20b^2) \int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx \\
 &= \frac{40b^3 \sin(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{20b^3 \sin^3(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^5(e + fx)}{3f} - \frac{1}{7}(20b^2) \int \sqrt{b \sec(e + fx)} dx \\
 &= \frac{40b^3 \sin(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{20b^3 \sin^3(e + fx)}{21f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^5(e + fx)}{3f} - \frac{80b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{21f} + \frac{40b^3 \sin^3(e + fx)}{21f \sqrt{b \sec(e + fx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.17, size = 74, normalized size = 0.57

$$\frac{b^2 \sqrt{b \sec(e + fx)} \left(320 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) - 58 \sin(2(e + fx)) + 3 \sin(4(e + fx)) - 56 \tan(e + fx) \right)}{84f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x]^6,x]

[Out] -1/84*(b^2*Sqrt[b*Sec[e + f*x]]*(320*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] - 58*Sin[2*(e + f*x)] + 3*Sin[4*(e + f*x)] - 56*Tan[e + f*x]))/f

Maple [C] Result contains complex when optimal does not.

time = 0.31, size = 168, normalized size = 1.29

method	result
--------	--------

default	$\frac{2(-1+\cos(fx+e)) \left(40i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - 3(\cos^5(fx+e)) + 3 \right)}{21f \sin(fx+e)^3}$
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(f*x+e))^(5/2)*sin(f*x+e)^6,x,method=_RETURNVERBOSE)`

[Out] $2/21/f*(-1+\cos(f*x+e))*(40*I*(1/(\cos(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*\operatorname{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)-3*\cos(f*x+e)^5+3*\cos(f*x+e)^4+16*\cos(f*x+e)^3-16*\cos(f*x+e)^2+7*\cos(f*x+e)-7)*\cos(f*x+e)*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(5/2)}/\sin(f*x+e)^3$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^6,x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^(5/2)*sin(f*x + e)^6, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.18, size = 140, normalized size = 1.08

$$\frac{2 \left(-20i \sqrt{2} b^{\frac{5}{2}} \cos(fx+e) \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e)) + 20i \sqrt{2} b^{\frac{5}{2}} \cos(fx+e) \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)) + (3b^2 \cos(fx+e)^4 - 16b^2 \cos(fx+e)^2 - 7b^2) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) \right)}{21f \cos(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^6,x, algorithm="fricas")`

[Out] $-2/21*(-20*I*\sqrt{2}*b^{(5/2)}*\cos(f*x + e)*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) + 20*I*\sqrt{2}*b^{(5/2)}*\cos(f*x + e)*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)) + (3*b^2*\cos(f*x + e)^4 - 16*b^2*\cos(f*x + e)^2 - 7*b^2)*\sqrt{b/\cos(f*x + e)}*\sin(f*x + e))/(f*\cos(f*x + e))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e)**6,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^6,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(5/2)*sin(f*x + e)^6, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^6 \left(\frac{b}{\cos(e + f x)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^6*(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^6*(b/cos(e + f*x))^(5/2), x)

3.405 $\int (b \sec(e + fx))^{5/2} \sin^4(e + fx) dx$

Optimal. Leaf size=100

$$-\frac{8b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{4b^3 \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^3(e + fx)}{3f}$$

[Out] $2/3*b*(b*\sec(f*x+e))^{(3/2)}*\sin(f*x+e)^3/f+4/3*b^3*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(1/2)}-8/3*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.07, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2704, 2707, 3856, 2720}

$$\frac{4b^3 \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} - \frac{8b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b \sin^3(e + fx)(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(5/2)}*\text{Sin}[e + f*x]^4, x]$

[Out] $(-8*b^2*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(3*f) + (4*b^3*\text{Sin}[e + f*x])/(3*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (2*b*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^3)/(3*f)$

Rule 2704

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(f*a*(n - 1))), x] + \text{Dist}[b^2*((m + 1)/(a^2*(n - 1))), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2707

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(a*f*(m + n))), x] + \text{Dist}[(m + 1)/(a^2*(m + n)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{NeQ}[m + n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^{5/2} \sin^4(e + fx) dx &= \frac{2b(b \sec(e + fx))^{3/2} \sin^3(e + fx)}{3f} - (2b^2) \int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx \\ &= \frac{4b^3 \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^3(e + fx)}{3f} - \frac{1}{3} (4b^2) \int \sqrt{b \sec(e + fx)} \sin^2(e + fx) dx \\ &= \frac{4b^3 \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{2b(b \sec(e + fx))^{3/2} \sin^3(e + fx)}{3f} - \frac{1}{3} (4b^2 \sqrt{b \sec(e + fx)}) \int \sin^2(e + fx) dx \\ &= -\frac{8b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{4b^3 \sin(e + fx)}{3f \sqrt{b \sec(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 64, normalized size = 0.64

$$\frac{b^2 \sqrt{b \sec(e + fx)} \left(8 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) - \sin(2(e + fx)) - 2 \tan(e + fx) \right)}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x]^4,x]

[Out] -1/3*(b^2*Sqrt[b*Sec[e + f*x]]*(8*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] - Sin[2*(e + f*x)] - 2*Tan[e + f*x]))/f

Maple [C] Result contains complex when optimal does not.

time = 0.26, size = 144, normalized size = 1.44

method	result
default	$\frac{2(-1+\cos(fx+e)) \left(4i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) + \cos^3(fx+e) - (\cos(fx+e))^3 \right)}{3f \sin(fx+e)^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(5/2)*sin(f*x+e)^4,x,method=_RETURNVERBOSE)

[Out] $\frac{2}{3}f(-1+\cos(f*x+e))*(4*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)+\cos(f*x+e)^3-\cos(f*x+e)^2+\cos(f*x+e)-1)*\cos(f*x+e)*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{5/2}/\sin(f*x+e)^3$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^4,x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^(5/2)*sin(f*x + e)^4, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 124, normalized size = 1.24

$$\frac{2 \left(-2i \sqrt{2} b^{\frac{1}{2}} \cos(fx+e) \text{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e)) + 2i \sqrt{2} b^{\frac{1}{2}} \cos(fx+e) \text{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)) - (b^2 \cos(fx+e)^2 + b^2) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) \right)}{3f \cos(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^4,x, algorithm="fricas")`

[Out] $-2/3*(-2*I*\sqrt{2}*b^{(5/2)}*\cos(f*x + e)*\text{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) + 2*I*\sqrt{2}*b^{(5/2)}*\cos(f*x + e)*\text{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)) - (b^2*\cos(f*x + e)^2 + b^2)*\sqrt{(b/\cos(f*x + e))*\sin(f*x + e))/(f*\cos(f*x + e))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e)**4,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^4,x, algorithm="giac")`

[Out] integrate((b*sec(f*x + e))^(5/2)*sin(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^4 \left(\frac{b}{\cos(e + f x)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4*(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^4*(b/cos(e + f*x))^(5/2), x)

3.406 $\int (b \sec(e + fx))^{5/2} \sin^2(e + fx) dx$

Optimal. Leaf size=70

$$-\frac{4b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f}$$

[Out] $2/3*b*(b*\sec(f*x+e))^{3/2}*\sin(f*x+e)/f-4/3*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.05, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2704, 3856, 2720}

$$\frac{2b \sin(e + fx)(b \sec(e + fx))^{3/2}}{3f} - \frac{4b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{5/2}*\text{Sin}[e + f*x]^2,x]$

[Out] $(-4*b^2*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(3*f) + (2*b*(b*\text{Sec}[e + f*x])^{3/2}*\text{Sin}[e + f*x])/(3*f)$

Rule 2704

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Csc}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(f*a*(n - 1))), x] + \text{Dist}[b^2*((m + 1)/(a^2*(n - 1))), \text{Int}[(a*\text{Csc}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, x\} \&\& \text{GtQ}[n, 1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{5/2} \sin^2(e + fx) dx &= \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f} - \frac{1}{3}(2b^2) \int \sqrt{b \sec(e + fx)} dx \\
&= \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f} - \frac{1}{3} \left(2b^2 \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \\
&= -\frac{4b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.10, size = 52, normalized size = 0.74

$$\frac{2b^2 \sqrt{b \sec(e + fx)} \left(-2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) + \tan(e + fx) \right)}{3f}$$

Antiderivative was successfully verified.

```
[In] Integrate[(b*Sec[e + f*x])^(5/2)*Sin[e + f*x]^2,x]
```

```
[Out] (2*b^2*Sqrt[b*Sec[e + f*x]]*(-2*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] + Tan[e + f*x]))/(3*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.22, size = 126, normalized size = 1.80

method	result
default	$\frac{2(-1+\cos(fx+e)) \left(2i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) + \cos(fx+e) - 1 \right) \cos(fx+e)}{3f \sin(fx+e)^3}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*sec(f*x+e))^(5/2)*sin(f*x+e)^2,x,method=_RETURNVERBOSE)
```

```
[Out] 2/3/f*(-1+cos(f*x+e))*(2*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+cos(f*x+e)-1)*cos(f*x+e)*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(5/2)/sin(f*x+e)^3
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^2,x, algorithm="maxima")
```

[Out] integrate((b*sec(f*x + e))^(5/2)*sin(f*x + e)^2, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.11, size = 110, normalized size = 1.57

$$\frac{2 \left(-i \sqrt{2} b^{\frac{5}{2}} \cos(fx + e) \operatorname{weierstrassPInverse}(-4, 0, \cos(fx + e) + i \sin(fx + e)) + i \sqrt{2} b^{\frac{5}{2}} \cos(fx + e) \operatorname{weierstrassPInverse}(-4, 0, \cos(fx + e) - i \sin(fx + e)) - b^2 \sqrt{\frac{b}{\cos(fx + e)}} \sin(fx + e) \right)}{3 f \cos(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^2,x, algorithm="fricas")

[Out] $-2/3 * (-I * \sqrt{2} * b^{(5/2)} * \cos(f*x + e) * \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I * \sin(f*x + e)) + I * \sqrt{2} * b^{(5/2)} * \cos(f*x + e) * \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I * \sin(f*x + e)) - b^2 * \sqrt{b / \cos(f*x + e)} * \sin(f*x + e)) / (f * \cos(f*x + e))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(5/2)*sin(f*x+e)**2,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2)*sin(f*x+e)^2,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(5/2)*sin(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^2 \left(\frac{b}{\cos(e + f x)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2*(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^2*(b/cos(e + f*x))^(5/2), x)

3.407 $\int (b \sec(e + fx))^{5/2} dx$

Optimal. Leaf size=70

$$\frac{2b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f}$$

[Out] $2/3*b*(b*\sec(f*x+e))^{(3/2)*\sin(f*x+e)/f+2/3*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.02, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3853, 3856, 2720}

$$\frac{2b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b \sin(e + fx)(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^{(5/2)},x]$

[Out] $(2*b^2*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(3*f) + (2*b*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x])/(3*f)$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3853

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n-1)}/(d*(n-1))), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\text{Csc}[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \&\& \text{GtQ}[n, 1] \&\& \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n-1)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int (b \sec(e + fx))^{5/2} dx &= \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f} + \frac{1}{3} b^2 \int \sqrt{b \sec(e + fx)} dx \\
&= \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f} + \frac{1}{3} \left(b^2 \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{1}{\sqrt{\cos(e + fx)}} dx \\
&= \frac{2b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b(b \sec(e + fx))^{3/2} \sin(e + fx)}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 51, normalized size = 0.73

$$\frac{2b^2 \sqrt{b \sec(e + fx)} \left(\sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) + \tan(e + fx) \right)}{3f}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^(5/2),x]`

```
[Out] (2*b^2*Sqrt[b*Sec[e + f*x]]*(Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] + Tan[e + f*x]))/(3*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.20, size = 128, normalized size = 1.83

method	result
default	$ -\frac{2(-1+\cos(fx+e)) \left(i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - \cos(fx+e)+1 \right) \cos(fx+e)}{3f \sin(fx+e)^3} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

```
[Out] -2/3/f*(-1+cos(f*x+e))*(I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-cos(f*x+e)+1)*cos(f*x+e)*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(5/2)/sin(f*x+e)^3
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out] integrate((b*sec(f*x + e))^(5/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.12, size = 110, normalized size = 1.57

$$\frac{-i\sqrt{2}b^{\frac{5}{2}}\cos(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{2}b^{\frac{5}{2}}\cos(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))+2b^2\sqrt{\frac{b}{\cos(fx+e)}}\sin(fx+e)}{3f\cos(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] $\frac{1}{3}*(-I*\sqrt{2}*b^{(5/2)}*\cos(f*x + e)*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) + I*\sqrt{2}*b^{(5/2)}*\cos(f*x + e)*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)) + 2*b^2*\sqrt{b/\cos(f*x + e)}*\sin(f*x + e))/(f*\cos(f*x + e))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(5/2),x)

[Out] Integral((b*sec(e + f*x))**(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(e + fx)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(5/2),x)

[Out] int((b/cos(e + f*x))^(5/2), x)

3.408 $\int \csc^2(e + fx)(b \sec(e + fx))^{5/2} dx$

Optimal. Leaf size=98

$$-\frac{5b^3 \csc(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{5b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b \csc(e + fx)(b \sec(e + fx))^{3/2}}{3f}$$

[Out] $2/3*b*csc(f*x+e)*(b*sec(f*x+e))^{(3/2)}/f-5/3*b^3*csc(f*x+e)/f/(b*sec(f*x+e))^{(1/2)}+5/3*b^2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*sec(f*x+e))^{(1/2)}/f$

Rubi [A]

time = 0.07, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2706, 2705, 3856, 2720}

$$-\frac{5b^3 \csc(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{5b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} + \frac{2b \csc(e + fx)(b \sec(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^2*(b*\text{Sec}[e + f*x])^{(5/2)}, x]$

[Out] $(-5*b^3*\text{Csc}[e + f*x])/(3*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (5*b^2*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(3*f) + (2*b*\text{Csc}[e + f*x]*(b*\text{Sec}[e + f*x])^{(3/2)})/(3*f)$

Rule 2705

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(m - 1)), x] + \text{Dist}[a^2*((m + n - 2)/(m - 1)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m - 2)}*(b*\text{Sec}[e + f*x])^{(n)}, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{IntegersQ}[2*m, 2*n] \&\& !\text{GtQ}[n, m]$

Rule 2706

$\text{Int}[(\text{csc}[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\text{sec}[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[a*b*(a*\text{Csc}[e + f*x])^{(m - 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(f*(n - 1)), x] + \text{Dist}[b^2*((m + n - 2)/(n - 1)), \text{Int}[(a*\text{Csc}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n - 2)}, x], x] /; \text{FreeQ}\{a, b, e, f, m\}, x] \&\& \text{GtQ}[n, 1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\text{csc}[c + d*x] + (d_*)*(x_*))*(b_*)^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n)}*\text{Sin}[c + d*x]^{-n}, \text{Int}[1/\text{Sin}[c + d*x]^{-n}, x], x] /;$ FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \csc^2(e + fx)(b \sec(e + fx))^{5/2} dx &= \frac{2b \csc(e + fx)(b \sec(e + fx))^{3/2}}{3f} + \frac{1}{3}(5b^2) \int \csc^2(e + fx) \sqrt{b \sec(e + fx)} dx \\ &= -\frac{5b^3 \csc(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{2b \csc(e + fx)(b \sec(e + fx))^{3/2}}{3f} + \frac{1}{6}(5b^2) \int \csc^2(e + fx) \sqrt{b \sec(e + fx)} dx \\ &= -\frac{5b^3 \csc(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{2b \csc(e + fx)(b \sec(e + fx))^{3/2}}{3f} + \frac{1}{6}(5b^2) \int \csc^2(e + fx) \sqrt{b \sec(e + fx)} dx \\ &= -\frac{5b^3 \csc(e + fx)}{3f \sqrt{b \sec(e + fx)}} + \frac{5b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3f} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 67, normalized size = 0.68

$$\frac{b \left(2 - 3 \cot^2(e + fx) + 5 \cos^{\frac{3}{2}}(e + fx) \csc(e + fx) F\left(\frac{1}{2}(e + fx) \mid 2\right) \right) (b \sec(e + fx))^{3/2} \sin(e + fx)}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^2*(b*Sec[e + f*x])^(5/2), x]

[Out] (b*(2 - 3*Cot[e + f*x]^2 + 5*Cos[e + f*x]^(3/2)*Csc[e + f*x]*EllipticF[(e + f*x)/2, 2])*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x])/(3*f)

Maple [C] Result contains complex when optimal does not.

time = 0.34, size = 202, normalized size = 2.06

method	result
default	$\frac{(-1 + \cos(fx + e))^2 \left(5i \text{EllipticF}\left(\frac{i(-1 + \cos(fx + e))}{\sin(fx + e)}, i\right) \sqrt{\frac{1}{\cos(fx + e) + 1}} \sqrt{\frac{\cos(fx + e)}{\cos(fx + e) + 1}} (\cos^2(fx + e) \sin(fx + e) + 5i \sqrt{\frac{1}{\cos(fx + e) + 1}}) \right)}{3f}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^2*(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)

[Out] 1/3/f*(-1+cos(f*x+e))^2*(5*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2*sin(f*x

$+e)+5*I*(1/(\cos(f*x+e)+1))^{(1/2)}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{(1/2)}*Elliptic$
 $F(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)-5*\cos(f*x+e)^2+2)*$
 $\cos(f*x+e)*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{(5/2)}/\sin(f*x+e)^5$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^(5/2)*csc(f*x + e)^2, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 143, normalized size = 1.46

$$\frac{-5i\sqrt{2}b^{\frac{5}{2}}\cos(fx+e)\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+5i\sqrt{2}b^{\frac{5}{2}}\cos(fx+e)\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))-2(5b^2\cos(fx+e)^2-2b^2)\sqrt{\frac{b}{\cos(fx+e)}}}{6f\cos(fx+e)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

[Out] `1/6*(-5*I*sqrt(2)*b^(5/2)*cos(f*x + e)*sin(f*x + e)*weierstrassPInverse(-4,`
`0, cos(f*x + e) + I*sin(f*x + e)) + 5*I*sqrt(2)*b^(5/2)*cos(f*x + e)*sin(f`
`*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)) - 2*(5*b^`
`2*cos(f*x + e)^2 - 2*b^2)*sqrt(b/cos(f*x + e)))/(f*cos(f*x + e)*sin(f*x + e`
`))`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**2*(b*sec(f*x+e))**(5/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^2*(b*sec(f*x+e))^(5/2),x, algorithm="giac")`

[Out] integrate((b*sec(f*x + e))^(5/2)*csc(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{5/2}}{\sin(e+fx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^2,x)

[Out] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^2, x)

3.409 $\int \csc^4(e + fx)(b \sec(e + fx))^{5/2} dx$

Optimal. Leaf size=123

$$-\frac{5b^3 \csc(e + fx)}{2f \sqrt{b \sec(e + fx)}} + \frac{5b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{2f} + \frac{b \csc(e + fx)(b \sec(e + fx))^{3/2}}{f}$$

[Out] b*csc(f*x+e)*(b*sec(f*x+e))^(3/2)/f-1/3*b*csc(f*x+e)^3*(b*sec(f*x+e))^(3/2)/f-5/2*b^3*csc(f*x+e)/f/(b*sec(f*x+e))^(1/2)+5/2*b^2*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2))*cos(f*x+e)^(1/2)*(b*sec(f*x+e))^(1/2)/f

Rubi [A]

time = 0.11, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2705, 2706, 3856, 2720}

$$-\frac{5b^3 \csc(e + fx)}{2f \sqrt{b \sec(e + fx)}} + \frac{5b^2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{2f} - \frac{b \csc^3(e + fx)(b \sec(e + fx))^{3/2}}{3f} + \frac{b \csc(e + fx)(b \sec(e + fx))^{3/2}}{f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^4*(b*Sec[e + f*x])^(5/2),x]

[Out] (-5*b^3*Csc[e + f*x])/(2*f*Sqrt[b*Sec[e + f*x]]) + (5*b^2*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*Sqrt[b*Sec[e + f*x]])/(2*f) + (b*Csc[e + f*x]*(b*Sec[e + f*x])^(3/2))/f - (b*Csc[e + f*x]^3*(b*Sec[e + f*x])^(3/2))/(3*f)

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2706

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[a*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(n - 1))), x] + Dist[b^2*((m + n - 2)/(n - 1)), Int[(a*Csc[e + f*x])^m*(b*Sec[e + f*x])^(n - 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && GtQ[n, 1] && IntegersQ[2*m, 2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)(x_.)]*(b_.))^{\wedge}(n_.), x_Symbol] \text{ :> Dist}[(b*\text{Csc}[c + d*x])^{\wedge}n*\text{Sin}[c + d*x]^{\wedge}n, \text{Int}[1/\text{Sin}[c + d*x]^{\wedge}n, x], x] \text{ /; FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned} \int \csc^4(e+fx)(b \sec(e+fx))^{5/2} dx &= -\frac{b \csc^3(e+fx)(b \sec(e+fx))^{3/2}}{3f} + \frac{3}{2} \int \csc^2(e+fx)(b \sec(e+fx))^{3/2} dx \\ &= \frac{b \csc(e+fx)(b \sec(e+fx))^{3/2}}{f} - \frac{b \csc^3(e+fx)(b \sec(e+fx))^{3/2}}{3f} + \frac{3}{2} \int \csc^2(e+fx)(b \sec(e+fx))^{3/2} dx \\ &= -\frac{5b^3 \csc(e+fx)}{2f \sqrt{b \sec(e+fx)}} + \frac{b \csc(e+fx)(b \sec(e+fx))^{3/2}}{f} - \frac{b \csc^3(e+fx)(b \sec(e+fx))^{3/2}}{3f} \\ &= -\frac{5b^3 \csc(e+fx)}{2f \sqrt{b \sec(e+fx)}} + \frac{b \csc(e+fx)(b \sec(e+fx))^{3/2}}{f} - \frac{b \csc^3(e+fx)(b \sec(e+fx))^{3/2}}{3f} \\ &= -\frac{5b^3 \csc(e+fx)}{2f \sqrt{b \sec(e+fx)}} + \frac{5b^2 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{2f} \end{aligned}$$

Mathematica [A]

time = 0.30, size = 79, normalized size = 0.64

$$\frac{b \left(4 - \cot^2(e+fx) (11 + 2 \csc^2(e+fx)) + 15 \cos^{3/2}(e+fx) \csc(e+fx) F\left(\frac{1}{2}(e+fx) \mid 2\right) \right) (b \sec(e+fx))^{3/2} \sin(e+fx)}{6f}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^4*(b*Sec[e + f*x])^(5/2), x]

[Out] (b*(4 - Cot[e + f*x]^2*(11 + 2*Csc[e + f*x]^2) + 15*Cos[e + f*x]^(3/2)*Csc[e + f*x]*EllipticF[(e + f*x)/2, 2])*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x])/(6*f)

Maple [C] Result contains complex when optimal does not.

time = 0.28, size = 352, normalized size = 2.86

method	result
default	$-\frac{(-1+\cos(fx+e))^2 \left(15i \text{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^4(fx+e) \sin(fx+e) + 15i \text{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \right) \right)}{6f}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^4*(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/6/f*(-1+\cos(f*x+e))^2*(15*I*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\cos(f*x+e)^4*\sin(f*x+e)+15*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\cos(f*x+e)^3*\sin(f*x+e)*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)-15*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\cos(f*x+e)^2*\sin(f*x+e)*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)-15*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)-15*\cos(f*x+e)^4+21*\cos(f*x+e)^2-4)*\cos(f*x+e)*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{5/2}/\sin(f*x+e)^7$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^(5/2)*csc(f*x + e)^4, x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 209, normalized size = 1.70

$$\frac{15\sqrt{2}(i^b\cos(fx+e)^3 - i^b\cos(fx+e))\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e) + i\sin(fx+e)) + 15\sqrt{2}(-i^b\cos(fx+e)^3 + i^b\cos(fx+e))\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e) - i\sin(fx+e)) + 2(15i^b\cos(fx+e)^4 - 21i^b\cos(fx+e)^2 + 4i^b)\sqrt{\frac{b}{\cos(fx+e)}}}{12(f\cos(fx+e)^3 - f\cos(fx+e))\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

[Out]
$$-1/12*(15*\sqrt{2}*(I*b^2*\cos(f*x + e)^3 - I*b^2*\cos(f*x + e))*\sqrt{b}*\sin(f*x + e)*\text{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) + 15*\sqrt{2}*(-I*b^2*\cos(f*x + e)^3 + I*b^2*\cos(f*x + e))*\sqrt{b}*\sin(f*x + e)*\text{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)) + 2*(15*b^2*\cos(f*x + e)^4 - 21*b^2*\cos(f*x + e)^2 + 4*b^2)*\sqrt{b/\cos(f*x + e)})/((f*\cos(f*x + e))^3 - f*\cos(f*x + e))*\sin(f*x + e)$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**4*(b*sec(f*x+e))**(5/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4*(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(5/2)*csc(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^{5/2}}{\sin(e+fx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^4,x)

[Out] int((b/cos(e + f*x))^(5/2)/sin(e + f*x)^4, x)

$$3.410 \quad \int \frac{\sin^7(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=87

$$\frac{2b^7}{15f(b \sec(e+fx))^{15/2}} - \frac{6b^5}{11f(b \sec(e+fx))^{11/2}} + \frac{6b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

[Out] $2/15*b^7/f/(b*\sec(f*x+e))^{(15/2)}-6/11*b^5/f/(b*\sec(f*x+e))^{(11/2)}+6/7*b^3/f/(b*\sec(f*x+e))^{(7/2)}-2/3*b/f/(b*\sec(f*x+e))^{(3/2)}$

Rubi [A]

time = 0.04, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$\frac{2b^7}{15f(b \sec(e+fx))^{15/2}} - \frac{6b^5}{11f(b \sec(e+fx))^{11/2}} + \frac{6b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^7/Sqrt[b*Sec[e + f*x]],x]

[Out] $(2*b^7)/(15*f*(b*\text{Sec}[e + f*x])^{(15/2)}) - (6*b^5)/(11*f*(b*\text{Sec}[e + f*x])^{(11/2)}) + (6*b^3)/(7*f*(b*\text{Sec}[e + f*x])^{(7/2)}) - (2*b)/(3*f*(b*\text{Sec}[e + f*x])^{(3/2)})$

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_.), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\int \frac{\sin^7(e+fx)}{\sqrt{b \sec(e+fx)}} dx = \frac{b^7 \text{Subst} \left(\int \frac{(-1+\frac{x^2}{b^2})^3}{x^{17/2}} dx, x, b \sec(e+fx) \right)}{f}$$

$$= \frac{b^7 \text{Subst} \left(\int \left(-\frac{1}{x^{17/2}} + \frac{3}{b^2 x^{13/2}} - \frac{3}{b^4 x^{9/2}} + \frac{1}{b^6 x^{5/2}} \right) dx, x, b \sec(e+fx) \right)}{f}$$

$$= \frac{2b^7}{15f(b \sec(e+fx))^{15/2}} - \frac{6b^5}{11f(b \sec(e+fx))^{11/2}} + \frac{6b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{3f(b \sec(e+fx))^{3/2}}{f}$$

Mathematica [A]

time = 0.16, size = 52, normalized size = 0.60

$$\frac{b(-7410 + 4035 \cos(2(e+fx)) - 798 \cos(4(e+fx)) + 77 \cos(6(e+fx)))}{18480 f (b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^7/Sqrt[b*Sec[e + f*x]], x]``[Out] (b*(-7410 + 4035*Cos[2*(e + f*x)] - 798*Cos[4*(e + f*x)] + 77*Cos[6*(e + f*x)]))/(18480*f*(b*Sec[e + f*x])^(3/2))`**Maple [A]**

time = 0.29, size = 56, normalized size = 0.64

method	result	size
default	$\frac{2(77(\cos^6(fx+e)) - 315(\cos^4(fx+e)) + 495(\cos^2(fx+e)) - 385)\cos(fx+e)}{1155f \sqrt{\frac{b}{\cos(fx+e)}}}$	56

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^7/(b*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)``[Out] 2/1155/f*(77*cos(f*x+e)^6-315*cos(f*x+e)^4+495*cos(f*x+e)^2-385)*cos(f*x+e)/(b/cos(f*x+e))^(1/2)`**Maxima [A]**

time = 0.28, size = 67, normalized size = 0.77

$$\frac{2 \left(77 b^6 - \frac{315 b^6}{\cos(fx+e)^2} + \frac{495 b^6}{\cos(fx+e)^4} - \frac{385 b^6}{\cos(fx+e)^6} \right) b}{1155 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{15}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] 2/1155*(77*b^6 - 315*b^6/cos(f*x + e)^2 + 495*b^6/cos(f*x + e)^4 - 385*b^6/cos(f*x + e)^6)*b/(f*(b/cos(f*x + e))^(15/2))

Fricas [A]

time = 0.39, size = 66, normalized size = 0.76

$$\frac{2(77 \cos(fx + e)^8 - 315 \cos(fx + e)^6 + 495 \cos(fx + e)^4 - 385 \cos(fx + e)^2) \sqrt{\frac{b}{\cos(fx + e)}}}{1155 bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 2/1155*(77*cos(f*x + e)^8 - 315*cos(f*x + e)^6 + 495*cos(f*x + e)^4 - 385*cos(f*x + e)^2)*sqrt(b/cos(f*x + e))/(b*f)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**7/(b*sec(f*x+e))**(1/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [A]

time = 4.16, size = 117, normalized size = 1.34

$$\frac{2(77 \sqrt{b \cos(fx + e)} b^7 \cos(fx + e)^7 - 315 \sqrt{b \cos(fx + e)} b^7 \cos(fx + e)^5 + 495 \sqrt{b \cos(fx + e)} b^7 \cos(fx + e)^3 - 385 \sqrt{b \cos(fx + e)} b^7 \cos(fx + e))}{1155 b^8 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] 2/1155*(77*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e)^7 - 315*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e)^5 + 495*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e)^3 - 385*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e))/(b^8*f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + fx)^7}{\sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^7/(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^7/(b/cos(e + f*x))^(1/2), x)

$$3.411 \quad \int \frac{\sin^5(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=65

$$-\frac{2b^5}{11f(b \sec(e+fx))^{11/2}} + \frac{4b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

[Out] $-2/11*b^5/f/(b*\sec(f*x+e))^{(11/2)}+4/7*b^3/f/(b*\sec(f*x+e))^{(7/2)}-2/3*b/f/(b*\sec(f*x+e))^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$-\frac{2b^5}{11f(b \sec(e+fx))^{11/2}} + \frac{4b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^5/Sqrt[b*Sec[e + f*x]],x]

[Out] $(-2*b^5)/(11*f*(b*Sec[e + f*x])^{(11/2)}) + (4*b^3)/(7*f*(b*Sec[e + f*x])^{(7/2)}) - (2*b)/(3*f*(b*Sec[e + f*x])^{(3/2)})$

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2)], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\int \frac{\sin^5(e+fx)}{\sqrt{b \sec(e+fx)}} dx = \frac{b^5 \text{Subst} \left(\int \frac{(-1+\frac{x^2}{b^2})^2}{x^{13/2}} dx, x, b \sec(e+fx) \right)}{f}$$

$$= \frac{b^5 \text{Subst} \left(\int \left(\frac{1}{x^{13/2}} - \frac{2}{b^2 x^{9/2}} + \frac{1}{b^4 x^{5/2}} \right) dx, x, b \sec(e+fx) \right)}{f}$$

$$= -\frac{2b^5}{11f(b \sec(e+fx))^{11/2}} + \frac{4b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

Mathematica [A]

time = 0.12, size = 42, normalized size = 0.65

$$\frac{b(-415 + 180 \cos(2(e+fx)) - 21 \cos(4(e+fx)))}{924f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^5/Sqrt[b*Sec[e + f*x]],x]``[Out] (b*(-415 + 180*Cos[2*(e + f*x)] - 21*Cos[4*(e + f*x)]))/(924*f*(b*Sec[e + f*x])^(3/2))`**Maple [A]**

time = 0.21, size = 46, normalized size = 0.71

method	result	size
default	$-\frac{2(21(\cos^4(fx+e)) - 66(\cos^2(fx+e)) + 77)\cos(fx+e)}{231f\sqrt{\frac{b}{\cos(fx+e)}}}$	46

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^5/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)``[Out] -2/231/f*(21*cos(f*x+e)^4-66*cos(f*x+e)^2+77)*cos(f*x+e)/(b/cos(f*x+e))^(1/2)`**Maxima [A]**

time = 0.27, size = 53, normalized size = 0.82

$$-\frac{2 \left(21 b^4 - \frac{66 b^4}{\cos(fx+e)^2} + \frac{77 b^4}{\cos(fx+e)^4} \right) b}{231 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{11}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] $-2/231*(21*b^4 - 66*b^4/\cos(f*x + e)^2 + 77*b^4/\cos(f*x + e)^4)*b/(f*(b/\cos(f*x + e))^{(11/2)})$

Fricas [A]

time = 0.39, size = 55, normalized size = 0.85

$$\frac{2(21 \cos(fx + e)^6 - 66 \cos(fx + e)^4 + 77 \cos(fx + e)^2) \sqrt{\frac{b}{\cos(fx + e)}}}{231bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $-2/231*(21*\cos(f*x + e)^6 - 66*\cos(f*x + e)^4 + 77*\cos(f*x + e)^2)*\text{sqrt}(b/\cos(f*x + e))/(b*f)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**5/(b*sec(f*x+e))**(1/2),x)

[Out] Timed out

Giac [A]

time = 3.66, size = 92, normalized size = 1.42

$$\frac{2(21 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e)^5 - 66 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e)^3 + 77 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e))}{231 b^6 f \text{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] $-2/231*(21*\text{sqrt}(b*\cos(f*x + e))*b^5*\cos(f*x + e)^5 - 66*\text{sqrt}(b*\cos(f*x + e))*b^5*\cos(f*x + e)^3 + 77*\text{sqrt}(b*\cos(f*x + e))*b^5*\cos(f*x + e))/(b^6*f*\text{sgn}(\cos(f*x + e)))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(e + fx)^5}{\sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^5/(b/cos(e + f*x))^(1/2),x)
```

```
[Out] int(sin(e + f*x)^5/(b/cos(e + f*x))^(1/2), x)
```

$$3.412 \quad \int \frac{\sin^3(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=43

$$\frac{2b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

[Out] $2/7*b^3/f/(b*\sec(f*x+e))^{(7/2)}-2/3*b/f/(b*\sec(f*x+e))^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 14}

$$\frac{2b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^3/Sqrt[b*Sec[e + f*x]],x]

[Out] $(2*b^3)/(7*f*(b*Sec[e + f*x])^{(7/2)}) - (2*b)/(3*f*(b*Sec[e + f*x])^{(3/2)})$

Rule 14

Int[(u_)*((c_)*(x_))^(m_), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2702

Int[csc[(e_.) + (f_)*(x_)]^(n_)*((a_)*sec[(e_.) + (f_)*(x_)]^(m_)), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m+n-1)/(-1+x^2/a^2)^((n+1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n+1)/2] && !(IntegerQ[(m+1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= \frac{b^3 \text{Subst}\left(\int \frac{-1+\frac{x^2}{b^2}}{x^{9/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{b^3 \text{Subst}\left(\int \left(-\frac{1}{x^{9/2}} + \frac{1}{b^2 x^{5/2}}\right) dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{2b^3}{7f(b \sec(e+fx))^{7/2}} - \frac{2b}{3f(b \sec(e+fx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 32, normalized size = 0.74

$$\frac{b(-11 + 3 \cos(2(e + fx)))}{21f(b \sec(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^3/Sqrt[b*Sec[e + f*x]],x]

[Out] (b*(-11 + 3*Cos[2*(e + f*x)]))/(21*f*(b*Sec[e + f*x])^(3/2))

Maple [A]

time = 0.19, size = 36, normalized size = 0.84

method	result	size
default	$\frac{2(3(\cos^2(fx+e))-7)\cos(fx+e)}{21f\sqrt{\frac{b}{\cos(fx+e)}}}$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^3/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] 2/21/f*(3*cos(f*x+e)^2-7)*cos(f*x+e)/(b/cos(f*x+e))^(1/2)

Maxima [A]

time = 0.31, size = 39, normalized size = 0.91

$$\frac{2\left(3b^2 - \frac{7b^2}{\cos(fx+e)^2}\right)b}{21f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{7}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] 2/21*(3*b^2 - 7*b^2/cos(f*x + e)^2)*b/(f*(b/cos(f*x + e))^(7/2))

Fricas [A]

time = 0.38, size = 44, normalized size = 1.02

$$\frac{2(3 \cos(fx + e)^4 - 7 \cos(fx + e)^2) \sqrt{\frac{b}{\cos(fx + e)}}}{21bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $2/21*(3*\cos(f*x + e)^4 - 7*\cos(f*x + e)^2)*\sqrt{b/\cos(f*x + e)}/(b*f)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**3/(b*sec(f*x+e))**(1/2),x)`

[Out] Timed out

Giac [A]

time = 4.05, size = 67, normalized size = 1.56

$$\frac{2 \left(3 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e)^3 - 7 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e) \right)}{21 b^4 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] $2/21*(3*\sqrt{b*\cos(f*x + e)}*b^3*\cos(f*x + e)^3 - 7*\sqrt{b*\cos(f*x + e)}*b^3*\cos(f*x + e))/b^4*f*\operatorname{sgn}(\cos(f*x + e))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(e + f x)^3}{\sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^3/(b/cos(e + f*x))^(1/2),x)`

[Out] `int(sin(e + f*x)^3/(b/cos(e + f*x))^(1/2), x)`

$$3.413 \quad \int \frac{\sin(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=20

$$-\frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

[Out] $-2/3*b/f/(b*\sec(f*x+e))^{(3/2)}$

Rubi [A]

time = 0.02, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 30}

$$-\frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]/Sqrt[b*Sec[e + f*x]],x]

[Out] $(-2*b)/(3*f*(b*Sec[e + f*x])^{(3/2)})$

Rule 30

Int[(x_)^(m_), x_Symbol] :> Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2702

Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= \frac{b \text{Subst}\left(\int \frac{1}{x^{5/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= -\frac{2b}{3f(b \sec(e+fx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 20, normalized size = 1.00

$$-\frac{2b}{3f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]/Sqrt[b*Sec[e + f*x]],x]

[Out] $(-2*b)/(3*f*(b*\text{Sec}[e + f*x])^{(3/2)})$

Maple [A]

time = 0.05, size = 17, normalized size = 0.85

method	result	size
derivativedivides	$-\frac{2b}{3f(b\sec(fx+e))^{\frac{3}{2}}}$	17
default	$-\frac{2b}{3f(b\sec(fx+e))^{\frac{3}{2}}}$	17

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] $-2/3*b/f/(b*\text{sec}(f*x+e))^{(3/2)}$

Maxima [A]

time = 0.34, size = 25, normalized size = 1.25

$$-\frac{2 \cos(fx + e)}{3f \sqrt{\frac{b}{\cos(fx + e)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] $-2/3*\cos(f*x + e)/(f*\text{sqrt}(b/\cos(f*x + e)))$

Fricas [A]

time = 0.39, size = 30, normalized size = 1.50

$$-\frac{2 \sqrt{\frac{b}{\cos(fx + e)}} \cos(fx + e)^2}{3bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $-2/3*\text{sqrt}(b/\cos(f*x + e))*\cos(f*x + e)^2/(b*f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sin(e + f*x)/sqrt(b*sec(e + f*x)), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 36 vs. 2(17) = 34.
time = 3.97, size = 36, normalized size = 1.80

$$-\frac{2 \sqrt{b \cos(fx + e)} \cos(fx + e)}{3 b f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] -2/3*sqrt(b*cos(f*x + e))*cos(f*x + e)/(b*f*sgn(cos(f*x + e)))

Mupad [B]

time = 0.54, size = 28, normalized size = 1.40

$$-\frac{2 \cos(e + fx)^2 \sqrt{\frac{b}{\cos(e + fx)}}}{3 b f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)/(b/cos(e + f*x))^(1/2),x)

[Out] -(2*cos(e + f*x)^2*(b/cos(e + f*x))^(1/2))/(3*b*f)

$$3.414 \quad \int \frac{\csc(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=59

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b} f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b} f}$$

[Out] $-\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f/b^{(1/2)}-\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f/b^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.263$, Rules used = {2702, 335, 218, 212, 209}

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b} f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{\sqrt{b} f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]/Sqrt[b*Sec[e + f*x]],x]`

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(\operatorname{Sqrt}[b]*f)) - \operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(\operatorname{Sqrt}[b]*f)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 335

```
Int[((c_.)*(x_)^(m_))*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\int \frac{\csc(e + fx)}{\sqrt{b \sec(e + fx)}} dx = \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(-1 + \frac{x^2}{b^2})} dx, x, b \sec(e + fx)\right)}{bf}$$

$$= \frac{2 \text{Subst}\left(\int \frac{1}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{bf}$$

$$= -\frac{\text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f} - \frac{\text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{f}$$

$$= -\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{\sqrt{b} f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{\sqrt{b} f}$$

Mathematica [A]

time = 0.08, size = 73, normalized size = 1.24

$$\frac{\left(2 \tan^{-1}\left(\sqrt{\sec(e + fx)}\right) - \log\left(1 - \sqrt{\sec(e + fx)}\right) + \log\left(1 + \sqrt{\sec(e + fx)}\right)\right) \sqrt{\sec(e + fx)}}{2f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]/Sqrt[b*Sec[e + f*x]],x]
```

```
[Out] -1/2*((2*ArcTan[Sqrt[Sec[e + f*x]]] - Log[1 - Sqrt[Sec[e + f*x]]] + Log[1 +
  Sqrt[Sec[e + f*x]]])*Sqrt[Sec[e + f*x]])/(f*Sqrt[b*Sec[e + f*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(47) = 94$.
time = 0.21, size = 161, normalized size = 2.73

method	result
default	$\frac{(-1 + \cos(fx + e)) \left(\arctan \left(\frac{1}{2 \sqrt{\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}}} \right) + \ln \left(-\frac{2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} - (\cos^2(fx + e) + 2 \cos(fx + e))}{\sin(fx + e)^2} \right)}{2f \sin(fx + e)^2 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} \sqrt{\frac{b}{\cos(fx + e)}} \right)}{}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-1/2/f*(-1+\cos(f*x+e))*(\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}))+\ln(-2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)/\sin(f*x+e)^2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}/(b/\cos(f*x+e))^{(1/2)}$

Maxima [A]

time = 0.50, size = 76, normalized size = 1.29

$$\frac{b \left(\frac{2 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx + e)}}}{\sqrt{b}} \right)}{b^{3/2}} - \frac{\log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx + e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx + e)}}} \right)}{b^{3/2}} \right)}{2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] $-1/2*b*(2*\arctan(\sqrt{b/\cos(f*x + e)})/\sqrt{b})/b^{(3/2)} - \log(-(\sqrt{b} - \sqrt{b/\cos(f*x + e)})/(\sqrt{b} + \sqrt{b/\cos(f*x + e)}))/b^{(3/2)}/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 128 vs. $2(49) = 98$.

time = 0.44, size = 271, normalized size = 4.59

$$\frac{2\sqrt{-b} \arctan \left(\frac{\sqrt{-b} \sqrt{\frac{b}{\cos(fx + e)}} (\cos(fx + e) + 1)}{2b} \right) - \sqrt{-b} \log \left(\frac{b \cos(fx + e)^2 - 4(\cos(fx + e)^2 - \cos(fx + e)) \sqrt{-b} \sqrt{\frac{b}{\cos(fx + e)}} - 4b \cos(fx + e) + b}{\cos(fx + e)^2 + 2 \cos(fx + e) + 1} \right)}{4bf} - \frac{2\sqrt{b} \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx + e)}} (\cos(fx + e) - 1)}{2\sqrt{b}} \right) + \sqrt{b} \log \left(\frac{b \cos(fx + e)^2 - 4(\cos(fx + e)^2 + \cos(fx + e)) \sqrt{b} \sqrt{\frac{b}{\cos(fx + e)}} + 4b \cos(fx + e) + b}{\cos(fx + e)^2 - 2 \cos(fx + e) + 1} \right)}{4bf}}{}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] [1/4*(2*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b) - sqrt(-b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)))/(b*f), 1/4*(2*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b)) + sqrt(b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)))/(b*f)]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(1/2),x)

[Out] Integral(csc(e + f*x)/sqrt(b*sec(e + f*x)), x)

Giac [A]

time = 4.13, size = 59, normalized size = 1.00

$$\frac{\frac{\arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b}} + \frac{\arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{\sqrt{b}}}{f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] (arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b) + arctan(sqrt(b*cos(f*x + e))/sqrt(b))/sqrt(b))/(f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(e + fx) \sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)*(b/cos(e + f*x))^(1/2)),x)

[Out] int(1/(sin(e + f*x)*(b/cos(e + f*x))^(1/2)), x)

$$3.415 \quad \int \frac{\csc^3(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=93

$$\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4\sqrt{b}f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4\sqrt{b}f} - \frac{\cot^2(e+fx)\sqrt{b \sec(e+fx)}}{2bf}$$

[Out] $-1/4*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f/b^{(1/2)}-1/4*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f/b^{(1/2)}-1/2*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(1/2)}/b/f$

Rubi [A]

time = 0.05, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2702, 294, 335, 218, 212, 209}

$$\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4\sqrt{b}f} - \frac{\cot^2(e+fx)\sqrt{b \sec(e+fx)}}{2bf} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4\sqrt{b}f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^3/Sqrt[b*Sec[e + f*x]],x]`

[Out] $-1/4*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(\operatorname{Sqrt}[b]*f) - \operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(4*\operatorname{Sqrt}[b]*f) - (\operatorname{Cot}[e + f*x]^2*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(2*b*f)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b`

, 0]

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(e + fx)}{\sqrt{b \sec(e + fx)}} dx &= \frac{\text{Subst}\left(\int \frac{x^{3/2}}{(-1 + \frac{x^2}{b^2})^2} dx, x, b \sec(e + fx)\right)}{b^3 f} \\
&= -\frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2bf} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x} (-1 + \frac{x^2}{b^2})} dx, x, b \sec(e + fx)\right)}{4bf} \\
&= -\frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2bf} + \frac{\text{Subst}\left(\int \frac{1}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{2bf} \\
&= -\frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2bf} - \frac{\text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4f} - \frac{\text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4f} \\
&= -\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4\sqrt{b} f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4\sqrt{b} f} - \frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2bf}
\end{aligned}$$

Mathematica [A]

time = 0.86, size = 93, normalized size = 1.00

$$\frac{\left(-2 \tan^{-1}\left(\sqrt{\sec(e+fx)}\right) + \log\left(1 - \sqrt{\sec(e+fx)}\right) - \log\left(1 + \sqrt{\sec(e+fx)}\right) - \frac{4 \csc^2(e+fx)}{\sec^{\frac{3}{2}}(e+fx)}\right) \sqrt{\sec(e+fx)}}{8f\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3/Sqrt[b*Sec[e + f*x]], x]

[Out] ((-2*ArcTan[Sqrt[Sec[e + f*x]]] + Log[1 - Sqrt[Sec[e + f*x]]] - Log[1 + Sqrt[Sec[e + f*x]]] - (4*Csc[e + f*x]^2)/Sec[e + f*x]^(3/2))*Sqrt[Sec[e + f*x]])/(8*f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 424 vs. 2(73) = 146.

time = 0.25, size = 425, normalized size = 4.57

method	result
default	$\frac{(-1+\cos(fx+e)) \left(8(\cos^2(fx+e)) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 16 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 8 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} - 4(\cos^2(fx+e)) \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3/(b*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)

[Out] -1/8/f*(-1+cos(f*x+e))*(8*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+16*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+8*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)-4*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))-cos(f*x+e)^2*ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+4*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))+ln(-2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2))/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)/sin(f*x+e)^4/(b/cos(f*x+e))^(1/2)

Maxima [A]

time = 0.52, size = 110, normalized size = 1.18

$$b \left(\frac{4 \sqrt{\frac{b}{\cos(fx+e)}}}{b^2 - \frac{b^2}{\cos(fx+e)^2}} - \frac{2 \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right)}{b^{\frac{3}{2}}} + \frac{\log\left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}}\right)}{b^{\frac{3}{2}}} \right) \frac{1}{8f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] 1/8*b*(4*sqrt(b/cos(f*x + e))/(b^2 - b^2/cos(f*x + e)^2) - 2*arctan(sqrt(b/cos(f*x + e))/sqrt(b))/b^(3/2) + log(-(sqrt(b) - sqrt(b/cos(f*x + e)))/(sqrt(b) + sqrt(b/cos(f*x + e))))/b^(3/2))/f

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(77) = 154.

time = 0.46, size = 389, normalized size = 4.18

$$\frac{2(\cos(fx+e)^2-1)\sqrt{-b}\arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + 8\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)^2 - (\cos(fx+e)^2-1)\sqrt{-b}\log\left(\frac{b\cos(fx+e)^2-4(\cos(fx+e)^2-\cos(fx+e))\sqrt{-b}}{(\cos(fx+e)^2-2\cos(fx+e)+1)\sqrt{b}}\right)}{16(b\cos(fx+e)^2-bf)} + \frac{2(\cos(fx+e)^2-1)\sqrt{-b}\arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + 8\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)^2 + (\cos(fx+e)^2-1)\sqrt{-b}\log\left(\frac{b\cos(fx+e)^2-4(\cos(fx+e)^2+\cos(fx+e))\sqrt{-b}}{(\cos(fx+e)^2-2\cos(fx+e)+1)\sqrt{b}}\right)}{16(b\cos(fx+e)^2-bf)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] [1/16*(2*(cos(f*x + e)^2 - 1)*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b) + 8*sqrt(b/cos(f*x + e))*cos(f*x + e)^2 - (cos(f*x + e)^2 - 1)*sqrt(-b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)))/(b*f*cos(f*x + e)^2 - b*f), 1/16*(2*(cos(f*x + e)^2 - 1)*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b)) + 8*sqrt(b/cos(f*x + e))*cos(f*x + e)^2 + (cos(f*x + e)^2 - 1)*sqrt(b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)))/(b*f*cos(f*x + e)^2 - b*f)]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**3/(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(csc(e + f*x)**3/sqrt(b*sec(e + f*x)), x)`

Giac [A]

time = 4.31, size = 110, normalized size = 1.18

$$\frac{b^2 \left(\frac{2 \sqrt{b \cos(fx + e)} \cos(fx + e)}{(b^2 \cos(fx + e)^2 - b^2) b} + \frac{\arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^2} + \frac{\arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}} \right)}{4 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `1/4*b^2*(2*sqrt(b*cos(f*x + e))*cos(f*x + e)/((b^2*cos(f*x + e)^2 - b^2)*b) + arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/(sqrt(-b)*b^2) + arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(5/2))/(f*sgn(cos(f*x + e)))`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^3 \sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^3*(b/cos(e + f*x))^(1/2)),x)`

[Out] `int(1/(sin(e + f*x)^3*(b/cos(e + f*x))^(1/2)), x)`

$$3.416 \quad \int \frac{\csc^5(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=123

$$\frac{5 \tan^{-1} \left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}} \right)}{32\sqrt{b} f} - \frac{5 \tanh^{-1} \left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}} \right)}{32\sqrt{b} f} - \frac{5 \cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16bf} - \frac{\cot^4(e+fx)}{b/f}$$

[Out] $-1/4*\cot(f*x+e)^4*(b*\sec(f*x+e))^{(5/2)}/b^3/f-5/32*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f/b^{(1/2)}-5/32*\operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/f/b^{(1/2)}-5/16*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(1/2)}/b/f$

Rubi [A]

time = 0.06, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2702, 294, 335, 218, 212, 209}

$$\frac{5 \operatorname{ArcTan} \left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}} \right)}{32\sqrt{b} f} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{5/2}}{4b^3 f} - \frac{5 \cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16bf} - \frac{5 \tanh^{-1} \left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}} \right)}{32\sqrt{b} f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^5/Sqrt[b*Sec[e + f*x]],x]`

[Out] $(-5*\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]])/(32*\operatorname{Sqrt}[b]*f) - (5*\operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]])/(32*\operatorname{Sqrt}[b]*f) - (5*\operatorname{Cot}[e + f*x]^2*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(16*b*f) - (\operatorname{Cot}[e + f*x]^4*(b*\operatorname{Sec}[e + f*x])^{(5/2)})/(4*b^3*f)$

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b`

, 0]

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p, x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^5(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= \frac{\text{Subst}\left(\int \frac{x^{7/2}}{(-1+\frac{x^2}{b^2})^3} dx, x, b \sec(e+fx)\right)}{b^5 f} \\
&= -\frac{\cot^4(e+fx)(b \sec(e+fx))^{5/2}}{4b^3 f} + \frac{5 \text{Subst}\left(\int \frac{x^{3/2}}{(-1+\frac{x^2}{b^2})^2} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{5 \cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16bf} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{5/2}}{4b^3 f} + \frac{5 \text{Subst}\left(\int \frac{1}{\sqrt{b \sec(e+fx)}} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{5 \cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16bf} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{5/2}}{4b^3 f} + \frac{5 \text{Subst}\left(\int \frac{1}{\sqrt{b \sec(e+fx)}} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{5 \cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16bf} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{5/2}}{4b^3 f} - \frac{5 \text{Subst}\left(\int \frac{1}{\sqrt{b \sec(e+fx)}} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{5 \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32\sqrt{b} f} - \frac{5 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32\sqrt{b} f} - \frac{5 \cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16bf}
\end{aligned}$$

Mathematica [A]

time = 1.62, size = 107, normalized size = 0.87

$$\frac{(10 \tan^{-1}(\sqrt{\sec(e+fx)}) - 5 \log(1 - \sqrt{\sec(e+fx)}) + 5 \log(1 + \sqrt{\sec(e+fx)}) + 4(-5 + \csc^2(e+fx) + 4 \csc^4(e+fx)) \sqrt{\sec(e+fx)}) \sqrt{\sec(e+fx)}}{64f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^5/Sqrt[b*Sec[e + f*x]],x]

[Out] -1/64*((10*ArcTan[Sqrt[Sec[e + f*x]]] - 5*Log[1 - Sqrt[Sec[e + f*x]]] + 5*Log[1 + Sqrt[Sec[e + f*x]]] + 4*(-5 + Csc[e + f*x]^2 + 4*Csc[e + f*x]^4)*Sqrt[Sec[e + f*x]]*Sqrt[Sec[e + f*x]])/(f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 728 vs. 2(99) = 198.

time = 0.25, size = 729, normalized size = 5.93

method	result
--------	--------

default	$40\left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}\right)^{\frac{3}{2}}(\cos^3(fx+e))+24(\cos^2(fx+e))\left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}\right)^{\frac{3}{2}}-20(\cos^3(fx+e))\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}-5\ln\left(-\right)$
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^5/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/64/f*(40*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\cos(f*x+e)^3+24*\cos(f*x+e) \\ & ^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}-20*\cos(f*x+e)^3*(-\cos(f*x+e)/(\cos(f \\ & *x+e)+1)^2)^{(1/2)}-5*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2} \\ &)-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f \\ & *x+e)^2)*\cos(f*x+e)^3-5*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)})*\co \\ & s(f*x+e)^3-72*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}+40*\cos(f*x+e) \\ & ^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+5*\cos(f*x+e)^2*\ln(-(2*\cos(f*x+e)^2* \\ & (-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+ \\ & e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)+5*\cos(f*x+e)^2*\arctan(1/2/(-\cos \\ & (f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}))-56*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}-20 \\ & *\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+5*\ln(-(2*\cos(f*x+e)^2*(-\co \\ & s(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/ \\ & \cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)+5*\arctan(1/2/(-\cos(f*x+e) \\ &)/(\cos(f*x+e)+1)^2)^{(1/2)})*\cos(f*x+e)-5*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(c \\ & os(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+ \\ & 1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-5*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1 \\ & /2)}))/\sin(f*x+e)^4/(b/\cos(f*x+e))^(1/2)/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2} \\ &) \end{aligned}$$

Maxima [A]

time = 0.55, size = 145, normalized size = 1.18

$$b \left(\frac{4 \left(5b^2 \sqrt{\frac{b}{\cos(fx+e)}} - 9 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{5}{2}} \right)}{b^4 - \frac{2b^4}{\cos(fx+e)^2} + \frac{b^4}{\cos(fx+e)^4}} - \frac{10 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right)}{b^{\frac{3}{2}}} + \frac{5 \log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right)}{b^{\frac{3}{2}}} \right)$$

64 f

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] $\frac{1}{64}b*(4*(5*b^2*\sqrt{b/\cos(f*x + e)} - 9*(b/\cos(f*x + e))^{(5/2)})/(b^4 - 2*b^4/\cos(f*x + e)^2 + b^4/\cos(f*x + e)^4) - 10*\arctan(\sqrt{b/\cos(f*x + e)})/\sqrt{b})/b^{(3/2)} + 5*\log(-(\sqrt{b} - \sqrt{b/\cos(f*x + e)}))/(\sqrt{b} + \sqrt{b/\cos(f*x + e)})/b^{(3/2)}/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 236 vs. $2(105) = 210$.
time = 0.46, size = 486, normalized size = 3.95

$$\frac{\frac{10 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \arctan\left(\frac{\sqrt{b}}{\sqrt{b \cos(fx + e)}}\right) - 5 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{128 \cos(fx + e)^2 - 25 \cos(fx + e) + 1} \frac{\frac{10 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \arctan\left(\frac{\sqrt{b}}{\sqrt{b \cos(fx + e)}}\right) + 8 \cos(fx + e)^2 - 9 \cos(fx + e)}{\cos(fx + e)} \frac{10 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \arctan\left(\frac{\sqrt{b}}{\sqrt{b \cos(fx + e)}}\right) + 5 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{128 \cos(fx + e)^2 - 25 \cos(fx + e) + 1} \frac{10 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \arctan\left(\frac{\sqrt{b}}{\sqrt{b \cos(fx + e)}}\right) + 8 \cos(fx + e)^2 - 9 \cos(fx + e)}{\cos(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] $\left[\frac{1}{128} * (10 * (\cos(f*x + e))^4 - 2 * \cos(f*x + e)^2 + 1) * \sqrt{-b} * \arctan\left(\frac{1}{2} * \sqrt{-b} * \sqrt{\frac{b}{\cos(f*x + e)}} * (\cos(f*x + e) + 1) / b\right) - 5 * (\cos(f*x + e))^4 - 2 * \cos(f*x + e)^2 + 1) * \sqrt{-b} * \log\left(\frac{(b * \cos(f*x + e))^2 - 4 * (\cos(f*x + e))^2 - \cos(f*x + e)}{\sqrt{-b} * \sqrt{\frac{b}{\cos(f*x + e)}}} - 6 * b * \cos(f*x + e) + b\right) / (\cos(f*x + e))^2 + 2 * \cos(f*x + e) + 1) + 8 * (5 * \cos(f*x + e)^4 - 9 * \cos(f*x + e)^2) * \sqrt{b / \cos(f*x + e)} \right] / (b * f * \cos(f*x + e)^4 - 2 * b * f * \cos(f*x + e)^2 + b * f), \frac{1}{128} * (10 * (\cos(f*x + e))^4 - 2 * \cos(f*x + e)^2 + 1) * \sqrt{b} * \arctan\left(\frac{1}{2} * \sqrt{b} * \sqrt{\frac{b}{\cos(f*x + e)}} * (\cos(f*x + e) - 1) / \sqrt{b}\right) + 5 * (\cos(f*x + e))^4 - 2 * \cos(f*x + e)^2 + 1) * \sqrt{b} * \log\left(\frac{(b * \cos(f*x + e))^2 - 4 * (\cos(f*x + e))^2 + \cos(f*x + e)}{\sqrt{b} * \sqrt{\frac{b}{\cos(f*x + e)}}} + 6 * b * \cos(f*x + e) + b\right) / (\cos(f*x + e))^2 - 2 * \cos(f*x + e) + 1) + 8 * (5 * \cos(f*x + e)^4 - 9 * \cos(f*x + e)^2) * \sqrt{b / \cos(f*x + e)} \right] / (b * f * \cos(f*x + e)^4 - 2 * b * f * \cos(f*x + e)^2 + b * f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^5(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**5/(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(csc(e + f*x)**5/sqrt(b*sec(e + f*x)), x)`

Giac [A]

time = 4.46, size = 143, normalized size = 1.16

$$\frac{b^4 \left(\frac{5 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^4} + \frac{5 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{b^{\frac{3}{2}}} + \frac{2 \left(5 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e)^3 - 9 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e) \right)}{(b^2 \cos(fx + e)^2 - b^2)^{\frac{3}{2}} b^4} \right)}{32 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] $\frac{1}{32}b^4(5\arctan(\sqrt{b\cos(fx+e)})/\sqrt{-b})/(\sqrt{-b}b^4) + 5\arctan(\sqrt{b\cos(fx+e)})/\sqrt{b}/b^{9/2} + 2(5\sqrt{b\cos(fx+e)}b^3\cos(fx+e)^3 - 9\sqrt{b\cos(fx+e)}b^3\cos(fx+e))/((b^2\cos(fx+e)^2 - b^2)^2b^4)/(f\operatorname{sgn}(\cos(fx+e)))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e+fx)^5 \sqrt{\frac{b}{\cos(e+fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^5*(b/cos(e + f*x))^(1/2)),x)

[Out] int(1/(sin(e + f*x)^5*(b/cos(e + f*x))^(1/2)), x)

$$3.417 \quad \int \frac{\sin^6(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=123

$$\frac{16E\left(\frac{1}{2}(e+fx) \mid 2\right)}{39f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}} - \frac{8b\sin(e+fx)}{39f(b\sec(e+fx))^{3/2}} - \frac{20b\sin^3(e+fx)}{117f(b\sec(e+fx))^{3/2}} - \frac{2b\sin^5(e+fx)}{13f(b\sec(e+fx))^{3/2}}$$

[Out] $-8/39*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}-20/117*b*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(3/2)}-2/13*b*\sin(f*x+e)^5/f/(b*\sec(f*x+e))^{(3/2)}+16/39*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.10, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2707, 3856, 2719}

$$-\frac{2b\sin^5(e+fx)}{13f(b\sec(e+fx))^{3/2}} - \frac{20b\sin^3(e+fx)}{117f(b\sec(e+fx))^{3/2}} - \frac{8b\sin(e+fx)}{39f(b\sec(e+fx))^{3/2}} + \frac{16E\left(\frac{1}{2}(e+fx) \mid 2\right)}{39f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^6/Sqrt[b*Sec[e + f*x]],x]

[Out] $(16*\text{EllipticE}[(e+f*x)/2, 2])/(39*f*\text{Sqrt}[\text{Cos}[e+f*x]]*\text{Sqrt}[b*\text{Sec}[e+f*x]]) - (8*b*\text{Sin}[e+f*x])/(39*f*(b*\text{Sec}[e+f*x])^{(3/2)}) - (20*b*\text{Sin}[e+f*x]^3)/(117*f*(b*\text{Sec}[e+f*x])^{(3/2)}) - (2*b*\text{Sin}[e+f*x]^5)/(13*f*(b*\text{Sec}[e+f*x])^{(3/2)})$

Rule 2707

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^6(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{2b \sin^5(e+fx)}{13f(b \sec(e+fx))^{3/2}} + \frac{10}{13} \int \frac{\sin^4(e+fx)}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{20b \sin^3(e+fx)}{117f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^5(e+fx)}{13f(b \sec(e+fx))^{3/2}} + \frac{20}{39} \int \frac{\sin^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{8b \sin(e+fx)}{39f(b \sec(e+fx))^{3/2}} - \frac{20b \sin^3(e+fx)}{117f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^5(e+fx)}{13f(b \sec(e+fx))^{3/2}} + \frac{8}{39} \int \frac{1}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{8b \sin(e+fx)}{39f(b \sec(e+fx))^{3/2}} - \frac{20b \sin^3(e+fx)}{117f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^5(e+fx)}{13f(b \sec(e+fx))^{3/2}} + \frac{8}{39} \sqrt{\frac{b \sec(e+fx)}{\cos(e+fx)}} \\
&= \frac{16E\left(\frac{1}{2}(e+fx) \mid 2\right)}{39f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{8b \sin(e+fx)}{39f(b \sec(e+fx))^{3/2}} - \frac{20b \sin^3(e+fx)}{117f(b \sec(e+fx))^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.33, size = 73, normalized size = 0.59

$$\frac{768E\left(\frac{1}{2}(e+fx) \mid 2\right) - 317 \sin(2(e+fx)) + 76 \sin(4(e+fx)) - 9 \sin(6(e+fx))}{1872f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^6/Sqrt[b*Sec[e + f*x]],x]`

```
[Out] ((768*EllipticE[(e + f*x)/2, 2])/Sqrt[Cos[e + f*x]] - 317*Sin[2*(e + f*x)]
+ 76*Sin[4*(e + f*x)] - 9*Sin[6*(e + f*x)])/(1872*f*Sqrt[b*Sec[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.51, size = 338, normalized size = 2.75

method	result
default	$ -\frac{2 \left(-9(\cos^8(fx+e)) + 24i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} - 24i \sqrt{\frac{1}{\cos(fx+e)+1}} \right)}{1872f \sqrt{b \sec(e+fx)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^6/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] -2/117/f*(-9*cos(f*x+e)^8+24*I*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*si
n(f*x+e)*cos(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1
```

/2)-24*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+37*cos(f*x+e)^6+24*I*sin(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)-24*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-59*cos(f*x+e)^4+55*cos(f*x+e)^2-24*cos(f*x+e))*(b/cos(f*x+e))^(1/2)/sin(f*x+e)/b

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^6/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sin(f*x + e)^6/sqrt(b*sec(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 126, normalized size = 1.02

$$\frac{2 \left((9 \cos(fx+e)^6 - 28 \cos(fx+e)^4 + 31 \cos(fx+e)^2) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) - 12i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e))) + 12i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e))) \right)}{117bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^6/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] -2/117*((9*cos(f*x + e)^6 - 28*cos(f*x + e)^4 + 31*cos(f*x + e)^2)*sqrt(b/cos(f*x + e))*sin(f*x + e) - 12*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + 12*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))))/(b*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^6(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**6/(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sin(e + f*x)**6/sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^6/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sin(f*x + e)^6/sqrt(b*sec(f*x + e)), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^6}{\sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^6/(b/cos(e + f*x))^(1/2),x)`

[Out] `int(sin(e + f*x)^6/(b/cos(e + f*x))^(1/2), x)`

$$3.418 \quad \int \frac{\sin^4(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=95

$$\frac{8E\left(\frac{1}{2}(e+fx) \mid 2\right)}{15f\sqrt{\cos(e+fx)}\sqrt{b \sec(e+fx)}} - \frac{4b \sin(e+fx)}{15f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^3(e+fx)}{9f(b \sec(e+fx))^{3/2}}$$

[Out] $-4/15*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}-2/9*b*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(3/2)}+8/15*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2707, 3856, 2719}

$$-\frac{2b \sin^3(e+fx)}{9f(b \sec(e+fx))^{3/2}} - \frac{4b \sin(e+fx)}{15f(b \sec(e+fx))^{3/2}} + \frac{8E\left(\frac{1}{2}(e+fx) \mid 2\right)}{15f\sqrt{\cos(e+fx)}\sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^4/Sqrt[b*Sec[e + f*x]], x]

[Out] $(8*\text{EllipticE}[(e+f*x)/2, 2])/((15*f*\text{Sqrt}[\text{Cos}[e+f*x]]*\text{Sqrt}[b*\text{Sec}[e+f*x]]) - (4*b*\text{Sin}[e+f*x])/((15*f*(b*\text{Sec}[e+f*x])^{(3/2)})) - (2*b*\text{Sin}[e+f*x]^3)/(9*f*(b*\text{Sec}[e+f*x])^{(3/2)}))$

Rule 2707

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^4(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{2b \sin^3(e+fx)}{9f(b \sec(e+fx))^{3/2}} + \frac{2}{3} \int \frac{\sin^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{4b \sin(e+fx)}{15f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^3(e+fx)}{9f(b \sec(e+fx))^{3/2}} + \frac{4}{15} \int \frac{1}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{4b \sin(e+fx)}{15f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^3(e+fx)}{9f(b \sec(e+fx))^{3/2}} + \frac{4 \int \sqrt{\cos(e+fx)} dx}{15 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
&= \frac{8E\left(\frac{1}{2}(e+fx) \mid 2\right)}{15f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{4b \sin(e+fx)}{15f(b \sec(e+fx))^{3/2}} - \frac{2b \sin^3(e+fx)}{9f(b \sec(e+fx))^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.24, size = 63, normalized size = 0.66

$$\frac{192E\left(\frac{1}{2}(e+fx) \mid 2\right)}{\sqrt{\cos(e+fx)}} - 68 \sin(2(e+fx)) + 10 \sin(4(e+fx))}{360f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^4/Sqrt[b*Sec[e + f*x]],x]`

```
[Out] ((192*EllipticE[(e + f*x)/2, 2])/Sqrt[Cos[e + f*x]] - 68*Sin[2*(e + f*x)] + 10*Sin[4*(e + f*x)])/(360*f*Sqrt[b*Sec[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.23, size = 328, normalized size = 3.45

method	result
default	$2 \left(12i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - 12i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^4/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 2/45/f*(12*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-12*I*cos(f*x+e)*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)-5*cos(f*x+e)^6+12*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-12*I*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(
```

$\cos(f*x+e)+1)^{(1/2)}*EllipticE(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)+16*\cos(f*x+e)^4-23*\cos(f*x+e)^2+12*\cos(f*x+e)*(b/\cos(f*x+e))^{(1/2)}/\sin(f*x+e)/b$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sin(f*x + e)^4/sqrt(b*sec(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 115, normalized size = 1.21

$$2 \left(\frac{(5 \cos(fx+e)^4 - 11 \cos(fx+e)^2) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) + 6i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e))) - 6i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)))}{45bf} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $\frac{2}{45} * ((5 * \cos(f*x + e)^4 - 11 * \cos(f*x + e)^2) * \sqrt{b / \cos(f*x + e)} * \sin(f*x + e) + 6 * I * \sqrt{2} * \sqrt{b} * \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I * \sin(f*x + e))) - 6 * I * \sqrt{2} * \sqrt{b} * \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I * \sin(f*x + e)))) / (b * f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^4(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**4/(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sin(e + f*x)**4/sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^4/sqrt(b*sec(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^4}{\sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4/(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^4/(b/cos(e + f*x))^(1/2), x)

$$3.419 \quad \int \frac{\sin^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=67

$$\frac{4E\left(\frac{1}{2}(e+fx) \mid 2\right)}{5f\sqrt{\cos(e+fx)}\sqrt{b \sec(e+fx)}} - \frac{2b \sin(e+fx)}{5f(b \sec(e+fx))^{3/2}}$$

[Out] $-2/5*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}+4/5*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2707, 3856, 2719}

$$\frac{4E\left(\frac{1}{2}(e+fx) \mid 2\right)}{5f\sqrt{\cos(e+fx)}\sqrt{b \sec(e+fx)}} - \frac{2b \sin(e+fx)}{5f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^2/Sqrt[b*Sec[e + f*x]],x]

[Out] $(4*\text{EllipticE}[(e+f*x)/2, 2])/(5*f*\text{Sqrt}[\text{Cos}[e+f*x]]*\text{Sqrt}[b*\text{Sec}[e+f*x]]) - (2*b*\text{Sin}[e+f*x])/(5*f*(b*\text{Sec}[e+f*x])^{(3/2)})$

Rule 2707

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{2b \sin(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{2}{5} \int \frac{1}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{2b \sin(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{2 \int \sqrt{\cos(e+fx)} dx}{5 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
&= \frac{4E\left(\frac{1}{2}(e+fx) \mid 2\right)}{5f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{2b \sin(e+fx)}{5f(b \sec(e+fx))^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 60, normalized size = 0.90

$$\frac{\sqrt{b \sec(e+fx)} \left(-8 \sqrt{\cos(e+fx)} E\left(\frac{1}{2}(e+fx) \mid 2\right) + \sin(e+fx) + \sin(3(e+fx)) \right)}{10bf}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^2/Sqrt[b*Sec[e + f*x]],x]`

```
[Out] -1/10*(Sqrt[b*Sec[e + f*x]]*(-8*Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2] + Sin[e + f*x] + Sin[3*(e + f*x)]))/(b*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.27, size = 318, normalized size = 4.75

method	result
default	$ -\frac{2 \left(2i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} - 2i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \right)}{10bf} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^2/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] -2/5/f*(2*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)-2*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)+2*I*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)-2*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-cos(f*x+e)^4+3*cos(f*x+e)^2-2*cos(f*x+e))*(b/cos(f*x+e))^(1/2)/sin(f*x+e)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sin(f*x + e)^2/sqrt(b*sec(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.12, size = 101, normalized size = 1.51

$$\frac{2 \left(\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)^2 \sin(fx+e) - i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e))) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e))) \right)}{5bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $-2/5 * (\sqrt{b/\cos(f*x+e)} * \cos(f*x+e)^2 * \sin(f*x+e) - I * \sqrt{2} * \sqrt{b} * \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x+e) + I * \sin(f*x+e))) + I * \sqrt{2} * \sqrt{b} * \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x+e) - I * \sin(f*x+e)))) / (b*f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^2(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**2/(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sin(e + f*x)**2/sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^2/sqrt(b*sec(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + fx)^2}{\sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2/(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^2/(b/cos(e + f*x))^(1/2), x)

$$3.420 \quad \int \frac{1}{\sqrt{b \sec(e + fx)}} dx$$

Optimal. Leaf size=38

$$\frac{2E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}}$$

[Out] $2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3856, 2719}

$$\frac{2E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[b*Sec[e + f*x]],x]

[Out] $(2*\text{EllipticE}[(e + f*x)/2, 2])/(f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \sec(e + fx)}} dx &= \frac{\int \sqrt{\cos(e + fx)} dx}{\sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} \\ &= \frac{2E\left(\frac{1}{2}(e + fx) \mid 2\right)}{f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 38, normalized size = 1.00

$$\frac{2E\left(\frac{1}{2}(e+fx)\middle|2\right)}{f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/Sqrt[b*Sec[e + f*x]],x]``[Out] (2*EllipticE[(e + f*x)/2, 2])/(f*Sqrt[Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])`**Maple [C]** Result contains complex when optimal does not.

time = 0.44, size = 306, normalized size = 8.05

method	result
risch	$-\frac{i\sqrt{2}}{f\sqrt{\frac{be^{i(fx+e)}}{e^{2i(fx+e)}+1}}}-i\left(-\frac{2(b e^{2i(fx+e)}+b)}{b\sqrt{e^{i(fx+e)}}(b e^{2i(fx+e)}+b)}+\frac{i\sqrt{-i(e^{i(fx+e)}+i)}\sqrt{2}\sqrt{i(e^{i(fx+e)}-i)}\sqrt{ie^{i(fx+e)}}}{b\sqrt{e^{i(fx+e)}}(b e^{2i(fx+e)}+b)}\right)$
default	$2\left(i\sqrt{\frac{1}{\cos(fx+e)+1}}\sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}}\text{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)},i\right)\sin(fx+e)\cos(fx+e)-i\text{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)},i\right)\sin(fx+e)\right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 2/f*(I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF
(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-I*cos(f*x+e)*sin(f*x
+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos
(f*x+e)/(cos(f*x+e)+1))^(1/2)+I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(
1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-I*sin(
f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE
(I*(-1+cos(f*x+e))/sin(f*x+e),I)-cos(f*x+e)^2+cos(f*x+e))*(b/cos(f*x+e))^(
1/2)/sin(f*x+e)/b
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")``[Out] integrate(1/sqrt(b*sec(f*x + e)), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.11, size = 70, normalized size = 1.84

$$\frac{i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))) - i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e) - i\sin(fx+e)))}{bf}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] `(I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) - I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))))/(b*f)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(1/sqrt(b*sec(e + f*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/sqrt(b*sec(f*x + e)), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{1}{\sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b/cos(e + f*x))^(1/2),x)`

[Out] `int(1/(b/cos(e + f*x))^(1/2), x)`

$$3.421 \quad \int \frac{\csc^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=63

$$-\frac{b \csc(e+fx)}{f(b \sec(e+fx))^{3/2}} - \frac{E\left(\frac{1}{2}(e+fx) \mid 2\right)}{f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

[Out] $-b \csc(f*x+e)/f/(b*\sec(f*x+e))^{3/2} - (\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2705, 3856, 2719}

$$-\frac{b \csc(e+fx)}{f(b \sec(e+fx))^{3/2}} - \frac{E\left(\frac{1}{2}(e+fx) \mid 2\right)}{f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^2/Sqrt[b*Sec[e + f*x]],x]`

[Out] $-\left(\frac{b \csc[e + f*x]}{f(b \sec[e + f*x])^{3/2}}\right) - \text{EllipticE}\left[\frac{e + f*x}{2}, 2\right] / (f \sqrt{\cos[e + f*x]} \sqrt{b \sec[e + f*x]})$

Rule 2705

`Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3856

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned}
\int \frac{\csc^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{b \csc(e+fx)}{f(b \sec(e+fx))^{3/2}} - \frac{1}{2} \int \frac{1}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{b \csc(e+fx)}{f(b \sec(e+fx))^{3/2}} - \frac{\int \sqrt{\cos(e+fx)} dx}{2\sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
&= -\frac{b \csc(e+fx)}{f(b \sec(e+fx))^{3/2}} - \frac{E\left(\frac{1}{2}(e+fx) \mid 2\right)}{f\sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 48, normalized size = 0.76

$$\frac{-\cot(e+fx) - \frac{E\left(\frac{1}{2}(e+fx) \mid 2\right)}{\sqrt{\cos(e+fx)}}}{f\sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2/Sqrt[b*Sec[e + f*x]], x]``[Out] (-Cot[e + f*x] - EllipticE[(e + f*x)/2, 2]/Sqrt[Cos[e + f*x]])/(f*Sqrt[b*Sec[e + f*x]])`**Maple [C]** Result contains complex when optimal does not.

time = 0.23, size = 316, normalized size = 5.02

method	result
default	$ -\frac{(-1+\cos(fx+e))^2 \left(i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \right)}{f \sqrt{b \sec(e+fx)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2/(b*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)`

```

[Out] -1/f*(-1+cos(f*x+e))^2*(I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*sin(f*x+e)*cos(f*x+e)-I*cos(f*x+e)*sin(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e), I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)+I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-I*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e), I)+cos(f*x+e))*(cos(f*x+e)+1)^2*(b/cos(f*x+e))^(1/2)/b/sin(f*x+e)^5

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^2/sqrt(b*sec(f*x + e)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 118, normalized size = 1.87

$$\frac{-i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e)))+i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e)))-2\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)^2}{2bf\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/2*(-I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) - 2*sqrt(b/cos(f*x + e))*cos(f*x + e)^2/(b*f*sin(f*x + e))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**2/(b*sec(f*x+e))**(1/2),x)
```

```
[Out] Integral(csc(e + f*x)**2/sqrt(b*sec(e + f*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(csc(f*x + e)^2/sqrt(b*sec(f*x + e)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(e + f x)^2 \sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^2*(b/cos(e + f*x))^(1/2)),x)

[Out] int(1/(sin(e + f*x)^2*(b/cos(e + f*x))^(1/2)), x)

$$3.422 \quad \int \frac{\csc^4(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=95

$$-\frac{b \csc(e+fx)}{2f(b \sec(e+fx))^{3/2}} - \frac{b \csc^3(e+fx)}{3f(b \sec(e+fx))^{3/2}} - \frac{E\left(\frac{1}{2}(e+fx) \mid 2\right)}{2f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

[Out] $-1/2*b*\csc(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}-1/3*b*\csc(f*x+e)^3/f/(b*\sec(f*x+e))^{(3/2)}-1/2*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2705, 3856, 2719}

$$-\frac{b \csc^3(e+fx)}{3f(b \sec(e+fx))^{3/2}} - \frac{b \csc(e+fx)}{2f(b \sec(e+fx))^{3/2}} - \frac{E\left(\frac{1}{2}(e+fx) \mid 2\right)}{2f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^4/Sqrt[b*Sec[e + f*x]],x]

[Out] $-1/2*(b*\text{Csc}[e + f*x])/(f*(b*\text{Sec}[e + f*x])^{(3/2)}) - (b*\text{Csc}[e + f*x]^3)/(3*f*(b*\text{Sec}[e + f*x])^{(3/2)}) - \text{EllipticE}[(e + f*x)/2, 2]/(2*f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

$$e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticE}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)^2-3*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)+3*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticE}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)-3*I*\text{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\sin(f*x+e)+3*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\text{EllipticE}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)+3*\cos(f*x+e)^3-2*\cos(f*x+e)^2-3*\cos(f*x+e))*(\cos(f*x+e)+1)^2*(b/\cos(f*x+e))^{1/2}/b/\sin(f*x+e)^7$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(csc(f*x + e)^4/sqrt(b*sec(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 171, normalized size = 1.80

$$\frac{3\sqrt{2}(i\cos(fx+e)^2-i)\sqrt{b}\sin(fx+e)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e)))+3\sqrt{2}(-i\cos(fx+e)^2+i)\sqrt{b}\sin(fx+e)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e)))+2(3\cos(fx+e)^4-5\cos(fx+e)^2)\sqrt{\frac{b}{\cos(fx+e)}}}{12(bf\cos(fx+e)^2-bf)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $-1/12*(3*\text{sqrt}(2)*(I*\cos(f*x + e)^2 - I)*\text{sqrt}(b)*\sin(f*x + e)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e))) + 3*\text{sqrt}(2)*(-I*\cos(f*x + e)^2 + I)*\text{sqrt}(b)*\sin(f*x + e)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e))) + 2*(3*\cos(f*x + e)^4 - 5*\cos(f*x + e)^2)*\text{sqrt}(b/\cos(f*x + e)))/((b*f*\cos(f*x + e)^2 - b*f)*\sin(f*x + e))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^4(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**4/(b*sec(f*x+e))**(1/2),x)

[Out] Integral(csc(e + f*x)**4/sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(1/2),x, algorithm="giac")``[Out] integrate(csc(f*x + e)^4/sqrt(b*sec(f*x + e)), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^4 \sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(sin(e + f*x)^4*(b/cos(e + f*x))^(1/2)),x)``[Out] int(1/(sin(e + f*x)^4*(b/cos(e + f*x))^(1/2)), x)`

$$3.423 \quad \int \frac{\csc^6(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=123

$$-\frac{7b \csc(e+fx)}{20f(b \sec(e+fx))^{3/2}} - \frac{7b \csc^3(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} - \frac{7E\left(\frac{1}{2}(e+fx) \mid 2\right)}{20f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

[Out] $-7/20*b*\csc(f*x+e)/f/(b*\sec(f*x+e))^{(3/2)}-7/30*b*\csc(f*x+e)^3/f/(b*\sec(f*x+e))^{(3/2)}-1/5*b*\csc(f*x+e)^5/f/(b*\sec(f*x+e))^{(3/2)}-7/20*(\cos(1/2*f*x+1/2*e))^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.10, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2705, 3856, 2719}

$$-\frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} - \frac{7b \csc^3(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{7b \csc(e+fx)}{20f(b \sec(e+fx))^{3/2}} - \frac{7E\left(\frac{1}{2}(e+fx) \mid 2\right)}{20f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^6/Sqrt[b*Sec[e + f*x]],x]

[Out] $(-7*b*Csc[e + f*x])/(20*f*(b*Sec[e + f*x])^{(3/2)}) - (7*b*Csc[e + f*x]^3)/(30*f*(b*Sec[e + f*x])^{(3/2)}) - (b*Csc[e + f*x]^5)/(5*f*(b*Sec[e + f*x])^{(3/2)}) - (7*EllipticE[(e + f*x)/2, 2])/(20*f*Sqrt[Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])$

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\csc^6(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7}{10} \int \frac{\csc^4(e+fx)}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{7b \csc^3(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7}{20} \int \frac{\csc^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{7b \csc(e+fx)}{20f(b \sec(e+fx))^{3/2}} - \frac{7b \csc^3(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} - \frac{7}{40} \int \frac{1}{\sqrt{b \sec(e+fx)}} dx \\
&= -\frac{7b \csc(e+fx)}{20f(b \sec(e+fx))^{3/2}} - \frac{7b \csc^3(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} - \frac{7}{40\sqrt{cb}} \int \frac{1}{\sqrt{\sec(e+fx)}} dx \\
&= -\frac{7b \csc(e+fx)}{20f(b \sec(e+fx))^{3/2}} - \frac{7b \csc^3(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \csc^5(e+fx)}{5f(b \sec(e+fx))^{3/2}} - \frac{7}{20f\sqrt{cb}} \int \frac{1}{\sqrt{\sec(e+fx)}} dx
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 86, normalized size = 0.70

$$\frac{\left(-21 + 7 \csc^2(e+fx) + 2 \csc^4(e+fx) + 12 \csc^6(e+fx) + 21 \sqrt{\cos(e+fx)} \csc(e+fx) E\left(\frac{1}{2}(e+fx) \middle| 2\right)\right) \tan(e+fx)}{60f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^6/Sqrt[b*Sec[e + f*x]], x]`

```
[Out] -1/60*((-21 + 7*Csc[e + f*x]^2 + 2*Csc[e + f*x]^4 + 12*Csc[e + f*x]^6 + 21*
Sqrt[Cos[e + f*x]]*Csc[e + f*x]*EllipticE[(e + f*x)/2, 2])*Tan[e + f*x])/(f
*Sqrt[b*Sec[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.28, size = 918, normalized size = 7.46

method	result	size
default	Expression too large to display	918

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^6/(b*sec(f*x+e))^(1/2), x, method=_RETURNVERBOSE)`

```
[Out] -1/60/f*(-1+cos(f*x+e))^2*(-42*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*
(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2*sin
(f*x+e)+42*I*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e), I)*(1/(cos(f*x+e)+1))^(1/2)
```

$$\begin{aligned} & 1/2) * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \cos(f*x+e)^2 * \sin(f*x+e) - 42 * I * \text{Elliptic} \\ & \text{cF}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * (1 / (\cos(f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos \\ & (f*x+e)+1))^{1/2} * \cos(f*x+e)^3 * \sin(f*x+e) + 21 * I * (1 / (\cos(f*x+e)+1))^{1/2} * (\cos \\ & (f*x+e) / (\cos(f*x+e)+1))^{1/2} * \text{EllipticF}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * \sin \\ & (f*x+e) * \cos(f*x+e)^5 + 21 * I * \text{EllipticF}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * (1 / (\cos \\ & (f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \sin(f*x+e) - 21 * I * (1 / (\cos \\ & (f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \text{EllipticE}(I * (-1 + \cos(f \\ & *x+e)) / \sin(f*x+e), I) * \sin(f*x+e) * \cos(f*x+e)^4 + 21 * I * (1 / (\cos(f*x+e)+1))^{1/2} * \\ & (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \text{EllipticF}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) \\ & * \sin(f*x+e) * \cos(f*x+e)^4 - 21 * I * \text{EllipticE}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * \cos \\ & (f*x+e) * (1 / (\cos(f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \sin(f*x+ \\ & e) + 21 * I * (1 / (\cos(f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \text{Elliptic} \\ & \text{F}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * \sin(f*x+e) * \cos(f*x+e) - 21 * I * (1 / (\cos(f*x+e) \\ & +1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \text{EllipticE}(I * (-1 + \cos(f*x+e)) / \sin \\ & (f*x+e), I) * \sin(f*x+e) + 42 * I * \text{EllipticE}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * (1 / (\cos \\ & (f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \cos(f*x+e)^3 * \sin(f*x+ \\ & e) - 21 * I * (1 / (\cos(f*x+e)+1))^{1/2} * (\cos(f*x+e) / (\cos(f*x+e)+1))^{1/2} * \text{Elliptic} \\ & \text{E}(I * (-1 + \cos(f*x+e)) / \sin(f*x+e), I) * \sin(f*x+e) * \cos(f*x+e)^5 + 21 * \cos(f*x+e)^5 - 1 \\ & 4 * \cos(f*x+e)^4 - 42 * \cos(f*x+e)^3 + 26 * \cos(f*x+e)^2 + 21 * \cos(f*x+e) * (\cos(f*x+e)+ \\ & 1)^2 * (b / \cos(f*x+e))^{1/2} / b / \sin(f*x+e)^9 \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(1/2), x, algorithm="maxima")

[Out] integrate(csc(f*x + e)^6/sqrt(b*sec(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 216, normalized size = 1.76

$$\frac{21\sqrt{2}(1\cos(fx+e)^2 - 2\cos(fx+e)^2 + 1)\sqrt{\sin(fx+e)}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(fx+e) + i\sin(fx+e))) + 21\sqrt{2}(-1\cos(fx+e)^2 + 2\cos(fx+e)^2 - 1)\sqrt{\sin(fx+e)}\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(fx+e) - i\sin(fx+e))) + 2(21\cos(fx+e)^6 - 56\cos(fx+e)^4 + 47\cos(fx+e)^2)\sqrt{\frac{b}{\cos(fx+e)}}}{120(b\cos(fx+e)^2 - 2b\cos(fx+e) + bf)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(1/2), x, algorithm="fricas")

[Out]
$$\begin{aligned} & -1/120 * (21 * \text{sqrt}(2) * (I * \cos(f*x + e)^4 - 2 * I * \cos(f*x + e)^2 + I) * \text{sqrt}(b) * \sin(\\ & f*x + e) * \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I \\ & * \sin(f*x + e))) + 21 * \text{sqrt}(2) * (-I * \cos(f*x + e)^4 + 2 * I * \cos(f*x + e)^2 - I) * \text{sqrt} \\ & (b) * \sin(f*x + e) * \text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(f \\ & *x + e) - I * \sin(f*x + e))) + 2 * (21 * \cos(f*x + e)^6 - 56 * \cos(f*x + e)^4 + 47 * \\ & \cos(f*x + e)^2) * \text{sqrt}(b / \cos(f*x + e))) / ((b * f * \cos(f*x + e)^4 - 2 * b * f * \cos(f*x \\ & + e)^2 + b * f) * \sin(f*x + e)) \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^6(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csc(f*x+e)**6/(b*sec(f*x+e))**(1/2),x)``[Out] Integral(csc(e + f*x)**6/sqrt(b*sec(e + f*x)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(1/2),x, algorithm="giac")``[Out] integrate(csc(f*x + e)^6/sqrt(b*sec(f*x + e)), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + fx)^6 \sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(sin(e + f*x)^6*(b/cos(e + f*x))^(1/2)),x)``[Out] int(1/(sin(e + f*x)^6*(b/cos(e + f*x))^(1/2)), x)`

$$3.424 \quad \int \frac{\sin^7(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=87

$$\frac{2b^7}{17f(b \sec(e+fx))^{17/2}} - \frac{6b^5}{13f(b \sec(e+fx))^{13/2}} + \frac{2b^3}{3f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

[Out] $2/17*b^7/f/(b*\sec(f*x+e))^{(17/2)}-6/13*b^5/f/(b*\sec(f*x+e))^{(13/2)}+2/3*b^3/f/(b*\sec(f*x+e))^{(9/2)}-2/5*b/f/(b*\sec(f*x+e))^{(5/2)}$

Rubi [A]

time = 0.04, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$\frac{2b^7}{17f(b \sec(e+fx))^{17/2}} - \frac{6b^5}{13f(b \sec(e+fx))^{13/2}} + \frac{2b^3}{3f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[e + f*x]^7/(b*Sec[e + f*x])^(3/2),x]`

[Out] $(2*b^7)/(17*f*(b*Sec[e + f*x])^{(17/2)}) - (6*b^5)/(13*f*(b*Sec[e + f*x])^{(13/2)}) + (2*b^3)/(3*f*(b*Sec[e + f*x])^{(9/2)}) - (2*b)/(5*f*(b*Sec[e + f*x])^{(5/2)})$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\int \frac{\sin^7(e+fx)}{(b \sec(e+fx))^{3/2}} dx = \frac{b^7 \text{Subst} \left(\int \frac{(-1+\frac{x^2}{b^2})^3}{x^{19/2}} dx, x, b \sec(e+fx) \right)}{f}$$

$$= \frac{b^7 \text{Subst} \left(\int \left(-\frac{1}{x^{19/2}} + \frac{3}{b^2 x^{15/2}} - \frac{3}{b^4 x^{11/2}} + \frac{1}{b^6 x^{7/2}} \right) dx, x, b \sec(e+fx) \right)}{f}$$

$$= \frac{2b^7}{17f(b \sec(e+fx))^{17/2}} - \frac{6b^5}{13f(b \sec(e+fx))^{13/2}} + \frac{2b^3}{3f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

Mathematica [A]

time = 0.30, size = 52, normalized size = 0.60

$$\frac{b(-10766 + 8365 \cos(2(e+fx)) - 1890 \cos(4(e+fx)) + 195 \cos(6(e+fx)))}{53040 f (b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^7/(b*Sec[e + f*x])^(3/2), x]``[Out] (b*(-10766 + 8365*Cos[2*(e + f*x)] - 1890*Cos[4*(e + f*x)] + 195*Cos[6*(e + f*x)]))/(53040*f*(b*Sec[e + f*x])^(5/2))`**Maple [A]**

time = 0.26, size = 56, normalized size = 0.64

method	result	size
default	$\frac{2(195(\cos^6(fx+e)) - 765(\cos^4(fx+e)) + 1105(\cos^2(fx+e)) - 663)\cos(fx+e)}{3315f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}}$	56

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^7/(b*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)``[Out] 2/3315/f*(195*cos(f*x+e)^6-765*cos(f*x+e)^4+1105*cos(f*x+e)^2-663)*cos(f*x+e)/(b/cos(f*x+e))^(3/2)`**Maxima [A]**

time = 0.29, size = 67, normalized size = 0.77

$$\frac{2 \left(195 b^6 - \frac{765 b^6}{\cos(fx+e)^2} + \frac{1105 b^6}{\cos(fx+e)^4} - \frac{663 b^6}{\cos(fx+e)^6} \right) b}{3315 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{17}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] $2/3315*(195*b^6 - 765*b^6/\cos(f*x + e)^2 + 1105*b^6/\cos(f*x + e)^4 - 663*b^6/\cos(f*x + e)^6)*b/(f*(b/\cos(f*x + e))^(17/2))$

Fricas [A]

time = 0.43, size = 66, normalized size = 0.76

$$\frac{2 \left(195 \cos(fx + e)^9 - 765 \cos(fx + e)^7 + 1105 \cos(fx + e)^5 - 663 \cos(fx + e)^3 \right) \sqrt{\frac{b}{\cos(fx + e)}}}{3315 b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] $2/3315*(195*\cos(f*x + e)^9 - 765*\cos(f*x + e)^7 + 1105*\cos(f*x + e)^5 - 663*\cos(f*x + e)^3)*\text{sqrt}(b/\cos(f*x + e))/(b^2*f)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**7/(b*sec(f*x+e))**(3/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [A]

time = 4.75, size = 119, normalized size = 1.37

$$\frac{2 \left(195 \sqrt{b \cos(fx + e)} b^8 \cos(fx + e)^8 - 765 \sqrt{b \cos(fx + e)} b^8 \cos(fx + e)^6 + 1105 \sqrt{b \cos(fx + e)} b^8 \cos(fx + e)^4 - 663 \sqrt{b \cos(fx + e)} b^8 \cos(fx + e)^2 \right)}{3315 b^{10} f \text{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] $2/3315*(195*\text{sqrt}(b*\cos(f*x + e))*b^8*\cos(f*x + e)^8 - 765*\text{sqrt}(b*\cos(f*x + e))*b^8*\cos(f*x + e)^6 + 1105*\text{sqrt}(b*\cos(f*x + e))*b^8*\cos(f*x + e)^4 - 663*\text{sqrt}(b*\cos(f*x + e))*b^8*\cos(f*x + e)^2)/(b^{10}*f*\text{sgn}(\cos(f*x + e)))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^7}{\left(\frac{b}{\cos(e + f x)} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^7/(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^7/(b/cos(e + f*x))^(3/2), x)

$$3.425 \quad \int \frac{\sin^5(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=65

$$-\frac{2b^5}{13f(b \sec(e+fx))^{13/2}} + \frac{4b^3}{9f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

[Out] $-2/13*b^5/f/(b*\sec(f*x+e))^{(13/2)}+4/9*b^3/f/(b*\sec(f*x+e))^{(9/2)}-2/5*b/f/(b*\sec(f*x+e))^{(5/2)}$

Rubi [A]

time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$-\frac{2b^5}{13f(b \sec(e+fx))^{13/2}} + \frac{4b^3}{9f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[e + f*x]^5/(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $(-2*b^5)/(13*f*(b*\text{Sec}[e + f*x])^{(13/2)}) + (4*b^3)/(9*f*(b*\text{Sec}[e + f*x])^{(9/2)}) - (2*b)/(5*f*(b*\text{Sec}[e + f*x])^{(5/2)})$

Rule 276

$\text{Int}[\text{((c_.)*(x_))}^{(m_.)}*\text{((a_.) + (b_.)*(x_)^{(n_)})}^{(p_.)}, x_Symbol] \text{ :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[\{a, b, c, m, n\}, x] \&\& IGtQ[p, 0]$

Rule 2702

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^{(n_.)}*\text{((a_.)*\text{sec}[(e_.) + (f_.)*(x_)])}^{(m_.)}, x_Symbol] \text{ :> Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m + n - 1)}/(-1 + x^2/a^2)^{((n + 1)/2)}, x], x, a*\text{Sec}[e + f*x]], x] /; FreeQ[\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n + 1)/2] \&\& \text{!(IntegerQ}[(m + 1)/2] \&\& \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned}
\int \frac{\sin^5(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= \frac{b^5 \text{Subst} \left(\int \frac{(-1+\frac{x^2}{b^2})^2}{x^{15/2}} dx, x, b \sec(e+fx) \right)}{f} \\
&= \frac{b^5 \text{Subst} \left(\int \left(\frac{1}{x^{15/2}} - \frac{2}{b^2 x^{11/2}} + \frac{1}{b^4 x^{7/2}} \right) dx, x, b \sec(e+fx) \right)}{f} \\
&= -\frac{2b^5}{13f(b \sec(e+fx))^{13/2}} + \frac{4b^3}{9f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}
\end{aligned}$$

Mathematica [A]

time = 0.18, size = 42, normalized size = 0.65

$$\frac{b(-551 + 340 \cos(2(e+fx)) - 45 \cos(4(e+fx)))}{2340f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^5/(b*Sec[e + f*x])^(3/2), x]``[Out] (b*(-551 + 340*Cos[2*(e + f*x)] - 45*Cos[4*(e + f*x)]))/(2340*f*(b*Sec[e + f*x])^(5/2))`**Maple [A]**

time = 0.20, size = 46, normalized size = 0.71

method	result	size
default	$-\frac{2(45 \cos^4(fx+e) - 130 \cos^2(fx+e) + 117) \cos(fx+e)}{585 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}}}$	46

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^5/(b*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)``[Out] -2/585/f*(45*cos(f*x+e)^4-130*cos(f*x+e)^2+117)*cos(f*x+e)/(b/cos(f*x+e))^(3/2)`**Maxima [A]**

time = 0.29, size = 53, normalized size = 0.82

$$-\frac{2 \left(45 b^4 - \frac{130 b^4}{\cos(fx+e)^2} + \frac{117 b^4}{\cos(fx+e)^4} \right) b}{585 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{13}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] -2/585*(45*b^4 - 130*b^4/cos(f*x + e)^2 + 117*b^4/cos(f*x + e)^4)*b/(f*(b/cos(f*x + e))^(13/2))

Fricas [A]

time = 0.41, size = 55, normalized size = 0.85

$$\frac{2(45 \cos(fx + e)^7 - 130 \cos(fx + e)^5 + 117 \cos(fx + e)^3) \sqrt{\frac{b}{\cos(fx + e)}}}{585 b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] -2/585*(45*cos(f*x + e)^7 - 130*cos(f*x + e)^5 + 117*cos(f*x + e)^3)*sqrt(b/cos(f*x + e))/(b^2*f)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**5/(b*sec(f*x+e))**(3/2),x)

[Out] Timed out

Giac [A]

time = 3.89, size = 94, normalized size = 1.45

$$\frac{2(45 \sqrt{b \cos(fx + e)} b^6 \cos(fx + e)^6 - 130 \sqrt{b \cos(fx + e)} b^6 \cos(fx + e)^4 + 117 \sqrt{b \cos(fx + e)} b^6 \cos(fx + e)^2)}{585 b^8 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] -2/585*(45*sqrt(b*cos(f*x + e))*b^6*cos(f*x + e)^6 - 130*sqrt(b*cos(f*x + e))*b^6*cos(f*x + e)^4 + 117*sqrt(b*cos(f*x + e))*b^6*cos(f*x + e)^2)/(b^8*f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(e + f x)^5}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^5/(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^5/(b/cos(e + f*x))^(3/2), x)

$$3.426 \quad \int \frac{\sin^3(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=43

$$\frac{2b^3}{9f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

[Out] $2/9*b^3/f/(b*\sec(f*x+e))^(9/2)-2/5*b/f/(b*\sec(f*x+e))^(5/2)$

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 14}

$$\frac{2b^3}{9f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^3/(b*Sec[e + f*x])^(3/2),x]

[Out] $(2*b^3)/(9*f*(b*Sec[e + f*x])^(9/2)) - (2*b)/(5*f*(b*Sec[e + f*x])^(5/2))$

Rule 14

Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m+n-1)/(-1+x^2/a^2)^((n+1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n+1)/2] && !(IntegerQ[(m+1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= \frac{b^3 \text{Subst}\left(\int \frac{-1+\frac{x^2}{b^2}}{x^{11/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{b^3 \text{Subst}\left(\int \left(-\frac{1}{x^{11/2}} + \frac{1}{b^2 x^{7/2}}\right) dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{2b^3}{9f(b \sec(e+fx))^{9/2}} - \frac{2b}{5f(b \sec(e+fx))^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.13, size = 32, normalized size = 0.74

$$\frac{b(-13 + 5 \cos(2(e + fx)))}{45f(b \sec(e + fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^3/(b*Sec[e + f*x])^(3/2),x]

[Out] (b*(-13 + 5*Cos[2*(e + f*x)]))/(45*f*(b*Sec[e + f*x])^(5/2))

Maple [A]

time = 0.17, size = 36, normalized size = 0.84

method	result	size
default	$\frac{2(5(\cos^2(fx+e))-9)\cos(fx+e)}{45f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}}$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^3/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] 2/45/f*(5*cos(f*x+e)^2-9)*cos(f*x+e)/(b/cos(f*x+e))^(3/2)

Maxima [A]

time = 0.29, size = 39, normalized size = 0.91

$$\frac{2\left(5b^2 - \frac{9b^2}{\cos(fx+e)^2}\right)b}{45f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{9}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] 2/45*(5*b^2 - 9*b^2/cos(f*x + e)^2)*b/(f*(b/cos(f*x + e))^(9/2))

Fricas [A]

time = 0.41, size = 44, normalized size = 1.02

$$\frac{2(5 \cos(fx + e)^5 - 9 \cos(fx + e)^3) \sqrt{\frac{b}{\cos(fx + e)}}}{45b^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] $2/45*(5*\cos(f*x + e)^5 - 9*\cos(f*x + e)^3)*\sqrt{b/\cos(f*x + e)}/(b^2*f)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**3/(b*sec(f*x+e))**(3/2),x)`

[Out] Timed out

Giac [A]

time = 2.81, size = 69, normalized size = 1.60

$$\frac{2 \left(5 \sqrt{b \cos(fx + e)} b^4 \cos(fx + e)^4 - 9 \sqrt{b \cos(fx + e)} b^4 \cos(fx + e)^2 \right)}{45 b^6 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(3/2),x, algorithm="giac")`

[Out] $2/45*(5*\sqrt{b*\cos(f*x + e)}*b^4*\cos(f*x + e)^4 - 9*\sqrt{b*\cos(f*x + e)}*b^4*\cos(f*x + e)^2)/(b^6*f*\operatorname{sgn}(\cos(f*x + e)))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(e + f x)^3}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^3/(b/cos(e + f*x))^(3/2),x)`

[Out] `int(sin(e + f*x)^3/(b/cos(e + f*x))^(3/2), x)`

$$3.427 \quad \int \frac{\sin(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=20

$$-\frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

[Out] $-2/5*b/f/(b*\sec(f*x+e))^{(5/2)}$

Rubi [A]

time = 0.02, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 30}

$$-\frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]/(b*Sec[e + f*x])^(3/2), x]

[Out] $(-2*b)/(5*f*(b*Sec[e + f*x])^{(5/2)})$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2702

Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= \frac{b \text{Subst}\left(\int \frac{1}{x^{7/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= -\frac{2b}{5f(b \sec(e+fx))^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 20, normalized size = 1.00

$$-\frac{2b}{5f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]/(b*Sec[e + f*x])^(3/2),x]

[Out] (-2*b)/(5*f*(b*Sec[e + f*x])^(5/2))

Maple [A]

time = 0.05, size = 17, normalized size = 0.85

method	result	size
derivativedivides	$-\frac{2b}{5f(b\sec(fx+e))^{\frac{5}{2}}}$	17
default	$-\frac{2b}{5f(b\sec(fx+e))^{\frac{5}{2}}}$	17

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] -2/5*b/f/(b*sec(f*x+e))^(5/2)

Maxima [A]

time = 0.28, size = 25, normalized size = 1.25

$$-\frac{2 \cos(fx + e)}{5f \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] -2/5*cos(f*x + e)/(f*(b/cos(f*x + e))^(3/2))

Fricas [A]

time = 0.39, size = 30, normalized size = 1.50

$$-\frac{2 \sqrt{\frac{b}{\cos(fx + e)}} \cos(fx + e)^3}{5b^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] -2/5*sqrt(b/cos(f*x + e))*cos(f*x + e)^3/(b^2*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))**(3/2),x)

[Out] Integral(sin(e + f*x)/(b*sec(e + f*x))**(3/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(17) = 34.
time = 3.70, size = 38, normalized size = 1.90

$$-\frac{2 \sqrt{b \cos (f x+e)} \cos (f x+e)^2}{5 b^2 f \operatorname{sgn}(\cos (f x+e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] -2/5*sqrt(b*cos(f*x + e))*cos(f*x + e)^2/(b^2*f*sgn(cos(f*x + e)))

Mupad [B]

time = 0.57, size = 28, normalized size = 1.40

$$-\frac{2 \cos (e+f x)^3 \sqrt{\frac{b}{\cos (e+f x)}}}{5 b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)/(b/cos(e + f*x))^(3/2),x)

[Out] -(2*cos(e + f*x)^3*(b/cos(e + f*x))^(1/2))/(5*b^2*f)

$$3.428 \quad \int \frac{\csc(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f} + \frac{2}{bf \sqrt{b \sec(e+fx)}}$$

[Out] arctan((b*sec(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f-arctanh((b*sec(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f+2/b/f/(b*sec(f*x+e))^(1/2)

Rubi [A]

time = 0.04, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2702, 331, 335, 304, 209, 212}

$$\frac{\text{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{3/2}f} + \frac{2}{bf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]/(b*Sec[e + f*x])^(3/2),x]

[Out] ArcTan[Sqrt[b*Sec[e + f*x]]/Sqrt[b]]/(b^(3/2)*f) - ArcTanh[Sqrt[b*Sec[e + f*x]]/Sqrt[b]]/(b^(3/2)*f) + 2/(b*f*Sqrt[b*Sec[e + f*x]])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))), Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc(e + fx)}{(b \sec(e + fx))^{3/2}} dx &= \frac{\text{Subst}\left(\int \frac{1}{x^{3/2}(-1 + \frac{x^2}{b^2})} dx, x, b \sec(e + fx)\right)}{bf} \\
&= \frac{2}{bf \sqrt{b \sec(e + fx)}} + \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{b^3 f} \\
&= \frac{2}{bf \sqrt{b \sec(e + fx)}} + \frac{2 \text{Subst}\left(\int \frac{x^2}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{b^3 f} \\
&= \frac{2}{bf \sqrt{b \sec(e + fx)}} - \frac{\text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{bf} + \frac{\text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{bf} \\
&= \frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{b^{3/2} f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{b^{3/2} f} + \frac{2}{bf \sqrt{b \sec(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 1.38, size = 89, normalized size = 1.14

$$\frac{4 + 2 \tan^{-1}(\sqrt{\sec(e + fx)}) \sqrt{\sec(e + fx)} + \left(\log(1 - \sqrt{\sec(e + fx)}) - \log(1 + \sqrt{\sec(e + fx)})\right) \sqrt{\sec(e + fx)}}{2bf \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]/(b*Sec[e + f*x])^(3/2),x]

[Out] (4 + 2*ArcTan[Sqrt[Sec[e + f*x]]]*Sqrt[Sec[e + f*x]] + (Log[1 - Sqrt[Sec[e + f*x]]] - Log[1 + Sqrt[Sec[e + f*x]]])*Sqrt[Sec[e + f*x]])/(2*b*f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 220 vs. 2(64) = 128.

time = 0.21, size = 221, normalized size = 2.83

method	result
default	$\frac{(-1 + \cos(fx + e)) \left(4 \cos(fx + e) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}} + \arctan\left(\frac{1}{2 \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}}}\right) - \ln\left(-\frac{2(\cos^2(fx + e)) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}}}{\cos(fx + e)}\right) \right)}{2f \sin(fx + e)^2 \left(\frac{b}{\cos(fx + e)}\right)^{\frac{3}{2}} \cos(fx + e) \sqrt{-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2}}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] -1/2/f*(-1+cos(f*x+e))*(4*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))-ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+4*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))/sin(f*x+e)^2/(b/cos(f*x+e))^(3/2)/cos(f*x+e)/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)

Maxima [A]

time = 0.51, size = 93, normalized size = 1.19

$$b \left(\frac{2 \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx + e)}}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}} + \frac{\log\left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx + e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx + e)}}}\right)}{b^{\frac{5}{2}}} + \frac{4}{b^2 \sqrt{\frac{b}{\cos(fx + e)}}} \right) \frac{1}{2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] $\frac{1}{2} \sqrt{b} (2 \arctan(\sqrt{b/\cos(fx+e)})/\sqrt{b})/b^{5/2} + \log(-(\sqrt{b} - \sqrt{b/\cos(fx+e)}))/(\sqrt{b} + \sqrt{b/\cos(fx+e)})/b^{5/2} + 4/(b^2 \sqrt{b/\cos(fx+e)})/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 164 vs. 2(67) = 134.

time = 0.50, size = 338, normalized size = 4.33

$$\frac{2\sqrt{-b} \arctan\left(\frac{z\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\cos(fx+e)}\right) + 8\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e) - \sqrt{-b} \log\left(\frac{\cos(fx+e)^2 + (\cos(fx+e)^2 - \cos(fx+e))\sqrt{-b}}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right)}{4b^2 f} + \frac{2\sqrt{b} \arctan\left(\frac{z\sqrt{b}\sqrt{\frac{b}{\cos(fx+e)}}}{\cos(fx+e)}\right) + 8\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e) + \sqrt{b} \log\left(\frac{\cos(fx+e)^2 + (\cos(fx+e)^2 - \cos(fx+e))\sqrt{b}}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right)}{4b^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{4} (2\sqrt{-b} \arctan(2\sqrt{-b} \sqrt{b/\cos(fx+e)}) \cos(fx+e) / (b \cos(fx+e) + b) + 8\sqrt{b/\cos(fx+e)} \cos(fx+e) - \sqrt{-b} \log(-b \cos(fx+e)^2 + 4(\cos(fx+e)^2 - \cos(fx+e)) \sqrt{-b} \sqrt{b/\cos(fx+e)} - 6b \cos(fx+e) + b) / (\cos(fx+e)^2 + 2\cos(fx+e) + 1)) / (b^2 f) + \frac{1}{4} (2\sqrt{b} \arctan(2\sqrt{b} \sqrt{b/\cos(fx+e)}) \cos(fx+e) / (b \cos(fx+e) - b) + 8\sqrt{b/\cos(fx+e)} \cos(fx+e) + \sqrt{b} \log(-b \cos(fx+e)^2 - 4(\cos(fx+e)^2 + \cos(fx+e)) \sqrt{b} \sqrt{b/\cos(fx+e)} + 6b \cos(fx+e) + b) / (\cos(fx+e)^2 - 2\cos(fx+e) + 1)) / (b^2 f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))**(3/2),x)

[Out] Integral(csc(e + f*x)/(b*sec(e + f*x))**(3/2), x)

Giac [A]

time = 3.49, size = 77, normalized size = 0.99

$$\frac{b \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b}} - \sqrt{b} \arctan\left(\frac{\sqrt{b \cos(fx+e)}}{\sqrt{b}}\right) + 2 \sqrt{b \cos(fx+e)} \frac{1}{b^2 f \operatorname{sgn}(\cos(fx+e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] (b*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b) - sqrt(b)*arctan(sqrt(b*cos(f*x + e))/sqrt(b)) + 2*sqrt(b*cos(f*x + e)))/(b^2*f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x) \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)*(b/cos(e + f*x))^(3/2)), x)

$$3.429 \quad \int \frac{\csc^3(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=93

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{3/2}f} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{3/2}f} - \frac{\cot^2(e+fx)(b \sec(e+fx))^{3/2}}{2b^3f}$$

[Out] $-1/4*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/b^{(3/2)}/f+1/4*\arctanh((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/b^{(3/2)}/f-1/2*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(3/2)}/b^3/f$

Rubi [A]

time = 0.05, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2702, 296, 335, 304, 209, 212}

$$-\frac{\text{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{3/2}f} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{3/2}f} - \frac{\cot^2(e+fx)(b \sec(e+fx))^{3/2}}{2b^3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^3/(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $-1/4*\text{ArcTan}[\text{Sqrt}[b*\text{Sec}[e + f*x]]/\text{Sqrt}[b]]/(b^{(3/2)*f}) + \text{ArcTanh}[\text{Sqrt}[b*\text{Sec}[e + f*x]]/\text{Sqrt}[b]]/(4*b^{(3/2)*f}) - (\text{Cot}[e + f*x]^2*(b*\text{Sec}[e + f*x])^{(3/2)})/(2*b^3*f)$

Rule 209

$\text{Int}[(a_+ + (b_-)*(x_-)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{GtQ}[b, 0])$

Rule 212

$\text{Int}[(a_+ + (b_-)*(x_-)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[-b, 2]))*\text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{NegQ}[a/b] \ \&\& \ (\text{GtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 296

$\text{Int}[(c_+*(x_+))^{(m_+)}*((a_+ + (b_-)*(x_-)^n)^{(p_+)}), x_Symbol] \rightarrow \text{Simp}[(-(c*x)^{(m+1))*((a + b*x^n)^{(p+1)})/(a*c*n*(p+1)), x] + \text{Dist}[(m + n*(p + 1) + 1)/(a*n*(p + 1)), \text{Int}[(c*x)^m*(a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}\{a, b, c, m\}, x \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{IntBinomialQ}[a, b, c, n, m, p]$

x]

Rule 304

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 335

```
Int[((c_)*(x_)^(m_))*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
  Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))^(p), x], x, (c*x)^(1/k)], x]] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1
)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(e + fx)}{(b \sec(e + fx))^{3/2}} dx &= \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{(-1 + \frac{x^2}{b^2})^2} dx, x, b \sec(e + fx)\right)}{b^3 f} \\
&= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2b^3 f} - \frac{\text{Subst}\left(\int \frac{\sqrt{x}}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{4b^3 f} \\
&= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2b^3 f} - \frac{\text{Subst}\left(\int \frac{x^2}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{2b^3 f} \\
&= -\frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2b^3 f} + \frac{\text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4bf} - \frac{\text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4bf} \\
&= -\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4b^{3/2} f} + \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4b^{3/2} f} - \frac{\cot^2(e + fx)(b \sec(e + fx))^{3/2}}{2b^3 f}
\end{aligned}$$

Mathematica [A]

time = 0.34, size = 98, normalized size = 1.05

$$\frac{-4 \csc^2(e + fx) - 2 \tan^{-1}\left(\sqrt{\sec(e + fx)}\right) \sqrt{\sec(e + fx)} + \left(-\log\left(1 - \sqrt{\sec(e + fx)}\right) + \log\left(1 + \sqrt{\sec(e + fx)}\right)\right) \sqrt{\sec(e + fx)}}{8bf\sqrt{b\sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3/(b*Sec[e + f*x])^(3/2), x]

[Out] $(-4*\text{Csc}[e + f*x]^2 - 2*\text{ArcTan}[\text{Sqrt}[\text{Sec}[e + f*x]]]*\text{Sqrt}[\text{Sec}[e + f*x]] + (-\text{Log}[1 - \text{Sqrt}[\text{Sec}[e + f*x]]] + \text{Log}[1 + \text{Sqrt}[\text{Sec}[e + f*x]]])* \text{Sqrt}[\text{Sec}[e + f*x]])/(8*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 425 vs. 2(73) = 146.

time = 0.22, size = 426, normalized size = 4.58

method	result
default	$- \frac{(-1 + \cos(fx + e)) \left(8(\cos^2(fx + e)) \left(-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2} \right)^{\frac{3}{2}} + 16 \cos(fx + e) \left(-\frac{\cos(fx + e)}{(\cos(fx + e) + 1)^2} \right)^{\frac{3}{2}} + (\cos^2(fx + e)) \arctan \left(\frac{\cos(fx + e)}{2\sqrt{-\cos(fx + e)}} \right) \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3/(b*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)

[Out] $-1/8/f*(-1+\cos(f*x+e))*(8*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)} + 16*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)} + \cos(f*x+e)^2*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}) - \cos(f*x+e)^2*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - \cos(f*x+e)^2 + 2*\cos(f*x+e) - 2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - 1)/\sin(f*x+e)^2) + 8*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)} + 4*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - 4*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - \arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}) + \ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - \cos(f*x+e)^2 + 2*\cos(f*x+e) - 2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} - 1)/\sin(f*x+e)^2))/\cos(f*x+e)/\sin(f*x+e)^4/(b/\cos(f*x+e))^{(3/2)}/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}$

Maxima [A]

time = 0.49, size = 111, normalized size = 1.19

$$b \left(\frac{4 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}}}{b^4 - \frac{b^4}{\cos(fx+e)^2}} - \frac{2 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right)}{b^{\frac{5}{2}}} - \frac{\log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right)}{b^{\frac{5}{2}}} \right) \frac{1}{8f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] 1/8*b*(4*(b/cos(f*x + e))^(3/2)/(b^4 - b^4/cos(f*x + e)^2) - 2*arctan(sqrt(b/cos(f*x + e))/sqrt(b))/b^(5/2) - log(-(sqrt(b) - sqrt(b/cos(f*x + e)))/(sqrt(b) + sqrt(b/cos(f*x + e))))/b^(5/2))/f

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 189 vs. 2(77) = 154.

time = 0.53, size = 392, normalized size = 4.22

$$\frac{2(\cos(fx+e)^2-1)\sqrt{-b} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + (\cos(fx+e)^2-1)\sqrt{-b} \log\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}} - \sqrt{b}}{\sqrt{\frac{b}{\cos(fx+e)}} + \sqrt{b}}\right) - 8\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e) + 2(\cos(fx+e)^2-1)\sqrt{-b} \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) + (\cos(fx+e)^2-1)\sqrt{-b} \log\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}} - \sqrt{b}}{\sqrt{\frac{b}{\cos(fx+e)}} + \sqrt{b}}\right) + 8\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)}{16(b^2 \cos(fx+e) - b^2 f)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] [-1/16*(2*(cos(f*x + e)^2 - 1)*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e)))*(cos(f*x + e) + 1)/b + (cos(f*x + e)^2 - 1)*sqrt(-b)*log((b*cos(f*x + e)^2 - 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)) - 8*sqrt(b/cos(f*x + e))*cos(f*x + e))/(b^2*f*cos(f*x + e)^2 - b^2*f), 1/16*(2*(cos(f*x + e)^2 - 1)*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e)))*(cos(f*x + e) - 1)/sqrt(b)) + (cos(f*x + e)^2 - 1)*sqrt(b)*log((b*cos(f*x + e)^2 + 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)) + 8*sqrt(b/cos(f*x + e))*cos(f*x + e))/(b^2*f*cos(f*x + e)^2 - b^2*f)]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3/(b*sec(f*x+e))**(3/2),x)

[Out] Integral(csc(e + f*x)**3/(b*sec(e + f*x))**(3/2), x)

Giac [A]

time = 3.50, size = 98, normalized size = 1.05

$$\frac{\frac{2\sqrt{b\cos(fx+e)}}{b^2\cos(fx+e)^2-b^2} - \frac{\arctan\left(\frac{\sqrt{b\cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b}b} + \frac{\arctan\left(\frac{\sqrt{b\cos(fx+e)}}{\sqrt{b}}\right)}{b^{\frac{3}{2}}}}{4f\operatorname{sgn}(\cos(fx+e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] 1/4*(2*sqrt(b*cos(f*x + e))/(b^2*cos(f*x + e)^2 - b^2) - arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b)*b + arctan(sqrt(b*cos(f*x + e))/sqrt(b))/b^(3/2))/(f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + fx)^3 \left(\frac{b}{\cos(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^3*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^3*(b/cos(e + f*x))^(3/2)), x)

$$3.430 \quad \int \frac{\csc^5(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=123

$$-\frac{3 \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{3/2}f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{3/2}f} - \frac{3 \cot^2(e+fx)(b \sec(e+fx))^{3/2}}{16b^3f} - \frac{\cot^4(e+fx)}{16b^3f}$$

[Out] -3/32*arctan((b*sec(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f+3/32*arctanh((b*sec(f*x+e))^(1/2)/b^(1/2))/b^(3/2)/f-3/16*cot(f*x+e)^2*(b*sec(f*x+e))^(3/2)/b^3/f-1/4*cot(f*x+e)^4*(b*sec(f*x+e))^(3/2)/b^3/f

Rubi [A]

time = 0.06, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2702, 294, 296, 335, 304, 209, 212}

$$-\frac{3 \text{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{3/2}f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{3/2}f} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{3/2}}{4b^3f} - \frac{3 \cot^2(e+fx)(b \sec(e+fx))^{3/2}}{16b^3f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^5/(b*Sec[e + f*x])^(3/2),x]

[Out] (-3*ArcTan[Sqrt[b*Sec[e + f*x]]/Sqrt[b]]/(32*b^(3/2)*f) + (3*ArcTanh[Sqrt[b*Sec[e + f*x]]/Sqrt[b]]/(32*b^(3/2)*f) - (3*Cot[e + f*x]^2*(b*Sec[e + f*x])^(3/2))/(16*b^3*f) - (Cot[e + f*x]^4*(b*Sec[e + f*x])^(3/2))/(4*b^3*f)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 294

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(n-1)*(c*x)^(m-n+1)*((a+b*x^n)^(p+1)/(b*n*(p+1))), x] - Dist[c^n*((m-n+1)/(b*n*(p+1))), Int[(c*x)^(m-n)*(a+b*x^n)^(p+1), x], x] /; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m+1, n] && !I

$\text{LtQ}[(m + n*(p + 1) + 1)/n, 0] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 296

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[(- (c*x)^{(m+1)}) * ((a + b*x^n)^{(p+1}) / (a*c*n*(p+1))), x] + \text{Dist}[(m + n*(p + 1) + 1) / (a*n*(p + 1)), \text{Int}[(c*x)^m * (a + b*x^n)^{(p+1)}, x], x] /; \text{FreeQ}\{a, b, c, m\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[p, -1] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 304

$\text{Int}[(x_)^2 / ((a_*) + (b_*)(x_)^4), x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Dist}[s/(2*b), \text{Int}[1/(r + s*x^2), x], x] - \text{Dist}[s/(2*b), \text{Int}[1/(r - s*x^2), x], x] /; \text{FreeQ}\{a, b\}, x] \&\& !\text{GtQ}[a/b, 0]$

Rule 335

$\text{Int}[(c_*)(x_)^{(m_*)}((a_*) + (b_*)(x_)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{With}\{k = \text{Denominator}[m]\}, \text{Dist}[k/c, \text{Subst}[\text{Int}[x^{(k*(m+1) - 1)} * (a + b*(x^{(k*n)})/c^n)^p, x], x, (c*x)^{(1/k)}], x] /; \text{FreeQ}\{a, b, c, p\}, x] \&\& \text{IGtQ}[n, 0] \&\& \text{FractionQ}[m] \&\& \text{IntBinomialQ}[a, b, c, n, m, p, x]$

Rule 2702

$\text{Int}[\text{csc}[(e_*) + (f_*)(x_)]^{(n_*)} * ((a_*) * \text{sec}[(e_*) + (f_*)(x_)])^{(m_*)}, x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)} / (-1 + x^2/a^2)^{(n+1)/2}], x], x, a*\text{Sec}[e + f*x], x] /; \text{FreeQ}\{a, e, f, m\}, x] \&\& \text{IntegerQ}[(n+1)/2] \&\& !(IntegerQ[(m+1)/2] \&\& \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned}
\int \frac{\csc^5(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= \frac{\text{Subst}\left(\int \frac{x^{5/2}}{(-1+\frac{x^2}{b^2})^3} dx, x, b \sec(e+fx)\right)}{b^5 f} \\
&= -\frac{\cot^4(e+fx)(b \sec(e+fx))^{3/2}}{4b^3 f} + \frac{3\text{Subst}\left(\int \frac{\sqrt{x}}{(-1+\frac{x^2}{b^2})^2} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{3 \cot^2(e+fx)(b \sec(e+fx))^{3/2}}{16b^3 f} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{3/2}}{4b^3 f} - \frac{3\text{Subst}\left(\int \frac{\sqrt{x}}{(-1+\frac{x^2}{b^2})} dx, x, b \sec(e+fx)\right)}{16b^3 f} \\
&= -\frac{3 \cot^2(e+fx)(b \sec(e+fx))^{3/2}}{16b^3 f} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{3/2}}{4b^3 f} - \frac{3\text{Subst}\left(\int \frac{\sqrt{x}}{(-1+\frac{x^2}{b^2})} dx, x, b \sec(e+fx)\right)}{16b^3 f} \\
&= -\frac{3 \cot^2(e+fx)(b \sec(e+fx))^{3/2}}{16b^3 f} - \frac{\cot^4(e+fx)(b \sec(e+fx))^{3/2}}{4b^3 f} + \frac{3\text{Subst}\left(\int \frac{\sqrt{x}}{(-1+\frac{x^2}{b^2})} dx, x, b \sec(e+fx)\right)}{16b^3 f} \\
&= -\frac{3 \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{3/2} f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{3/2} f} - \frac{3 \cot^2(e+fx)(b \sec(e+fx))^{3/2}}{16b^3 f}
\end{aligned}$$

Mathematica [A]

time = 0.47, size = 109, normalized size = 0.89

$$\frac{4 \csc^2(e+fx) - 16 \csc^4(e+fx) - 6 \tan^{-1}\left(\frac{\sqrt{\sec(e+fx)}}{\sqrt{b}}\right) \sqrt{\sec(e+fx)} + 3\left(-\log\left(1 - \sqrt{\sec(e+fx)}\right) + \log\left(1 + \sqrt{\sec(e+fx)}\right)\right) \sqrt{\sec(e+fx)}}{64bf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^5/(b*Sec[e + f*x])^(3/2), x]`

```
[Out] (4*Csc[e + f*x]^2 - 16*Csc[e + f*x]^4 - 6*ArcTan[Sqrt[Sec[e + f*x]])*Sqrt[Sec[e + f*x]] + 3*(-Log[1 - Sqrt[Sec[e + f*x]]] + Log[1 + Sqrt[Sec[e + f*x]])*Sqrt[Sec[e + f*x]])/(64*b*f*Sqrt[b*Sec[e + f*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 728 vs. 2(99) = 198.

time = 0.24, size = 729, normalized size = 5.93

method	result
--------	--------

default	$8\left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}\right)^{\frac{3}{2}}(\cos^3(fx+e))-8(\cos^2(fx+e))\left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}\right)^{\frac{3}{2}}-3\ln\left(-\frac{2(\cos^2(fx+e))\sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}-(\cos^2(fx+e))}{\sin(fx+e)^2}\right)$
---------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^5/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -1/64/f*(8*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}*\cos(f*x+e)^3-8*\cos(f*x+e)^2 \\ & *(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(3/2)}-3*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(c \\ & \cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+ \\ & 1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)^3+3*\arctan(1/2/(-\cos(f*x+e)/(\cos(f* \\ & x+e)+1)^2)^{(1/2}))*\cos(f*x+e)^3-40*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2) \\ & ^{(3/2)}+12*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+3*\cos(f*x+e)^2* \\ & \ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos \\ & (f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)-3*\cos(f*x+e \\ &)^2*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2}))-24*(-\cos(f*x+e)/(\cos(f \\ & *x+e)+1)^2)^{(3/2)}-24*\cos(f*x+e)*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}+3*\ln(- \\ & (2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x \\ & +e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/\sin(f*x+e)^2)*\cos(f*x+e)-3*\ar \\ & ctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2}))*\cos(f*x+e)+12*(-\cos(f*x+e)/(\\ & \cos(f*x+e)+1)^2)^{(1/2)}-3*\ln(-(2*\cos(f*x+e)^2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2) \\ & ^{(1/2)}-\cos(f*x+e)^2+2*\cos(f*x+e)-2*(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)}-1)/ \\ & \sin(f*x+e)^2)+3*\arctan(1/2/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2}))) / \sin(f*x+e \\ &)^4/(b/\cos(f*x+e))^(3/2)/\cos(f*x+e)/(-\cos(f*x+e)/(\cos(f*x+e)+1)^2)^{(1/2)} \end{aligned}$$

Maxima [A]

time = 0.50, size = 144, normalized size = 1.17

$$b \left(\frac{4 \left(b^2 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} + 3 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{7}{2}} \right)}{b^6 - \frac{2b^6}{\cos(fx+e)^2} + \frac{b^6}{\cos(fx+e)^4}} + \frac{6 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right)}{b^{\frac{5}{2}}} + \frac{3 \log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right)}{b^{\frac{5}{2}}} \right) / 64 f$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] $-1/64*b*(4*(b^2*(b/\cos(f*x + e))^{(3/2)} + 3*(b/\cos(f*x + e))^{(7/2)))/(b^6 - 2*b^6/\cos(f*x + e)^2 + b^6/\cos(f*x + e)^4) + 6*\arctan(\sqrt{b/\cos(f*x + e)})/\sqrt{b})/b^{(5/2)} + 3*\log(-(\sqrt{b} - \sqrt{b/\cos(f*x + e)}))/(\sqrt{b} + \sqrt{b/\cos(f*x + e)})/b^{(5/2)}/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(105) = 210$.
time = 0.47, size = 490, normalized size = 3.98

$$\frac{\frac{6 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right) + 3 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{128 \sqrt{b \cos(fx + e)} \sqrt{b^2 \cos(fx + e)^4 - 2 b^2 f \cos(fx + e)^2 + b^2 f}} + \frac{6 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \log\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right) + \frac{6 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{\sqrt{b}} \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right) + 3 \cos(fx + e)^2 - 2 \cos(fx + e) + 1}{128 \sqrt{b \cos(fx + e)} \sqrt{b^2 \cos(fx + e)^4 - 2 b^2 f \cos(fx + e)^2 + b^2 f}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")`

[Out] $[-1/128*(6*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{-b}*\arctan(1/2*\sqrt{-b}*\sqrt{b/\cos(f*x + e)}*(\cos(f*x + e) + 1)/b) + 3*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{-b}*\log((b*\cos(f*x + e)^2 - 4*(\cos(f*x + e)^2 - \cos(f*x + e))*\sqrt{-b}*\sqrt{b/\cos(f*x + e)} - 6*b*\cos(f*x + e) + b)/(\cos(f*x + e)^2 + 2*\cos(f*x + e) + 1)) + 8*(\cos(f*x + e)^3 + 3*\cos(f*x + e))*\sqrt{b/\cos(f*x + e)})/(b^2*f*\cos(f*x + e)^4 - 2*b^2*f*\cos(f*x + e)^2 + b^2*f), 1/128*(6*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{b}*\arctan(1/2*\sqrt{b/\cos(f*x + e)}*(\cos(f*x + e) - 1)/\sqrt{b}) + 3*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{b}*\log((b*\cos(f*x + e)^2 + 4*(\cos(f*x + e)^2 + \cos(f*x + e))*\sqrt{b}*\sqrt{b/\cos(f*x + e)} + 6*b*\cos(f*x + e) + b)/(\cos(f*x + e)^2 - 2*\cos(f*x + e) + 1)) - 8*(\cos(f*x + e)^3 + 3*\cos(f*x + e))*\sqrt{b/\cos(f*x + e)})/(b^2*f*\cos(f*x + e)^4 - 2*b^2*f*\cos(f*x + e)^2 + b^2*f)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^5(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**5/(b*sec(f*x+e))**(3/2),x)`

[Out] `Integral(csc(e + f*x)**5/(b*sec(e + f*x))**(3/2), x)`

Giac [A]

time = 3.62, size = 135, normalized size = 1.10

$$\frac{b^2 \left(\frac{3 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^3} - \frac{3 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{b^{\frac{3}{2}}} + \frac{2 \left(\sqrt{b \cos(fx + e)} b^2 \cos(fx + e)^2 + 3 \sqrt{b \cos(fx + e)} b^2 \right)}{(b^2 \cos(fx + e)^2 - b^2)^2 b^2} \right)}{32 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out]
$$-1/32*b^2*(3*\arctan(\sqrt{b*\cos(f*x + e)})/\sqrt{-b})/(\sqrt{-b}*b^3) - 3*\arctan(\sqrt{b*\cos(f*x + e)})/\sqrt{b})/b^{7/2} + 2*(\sqrt{b*\cos(f*x + e)}*b^2*\cos(f*x + e)^2 + 3*\sqrt{b*\cos(f*x + e)}*b^2)/((b^2*\cos(f*x + e)^2 - b^2)^{2*b^2})/(f*\text{sgn}(\cos(f*x + e)))$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^5 \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^5*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^5*(b/cos(e + f*x))^(3/2)), x)

$$3.431 \quad \int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=126

$$\frac{8\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{77b^2 f} - \frac{12b \sin(e+fx)}{77f(b \sec(e+fx))^{5/2}} + \frac{8 \sin(e+fx)}{77bf \sqrt{b \sec(e+fx)}} - \frac{2b \sin^3(e+fx)}{11f(b \sec(e+fx))^{5/2}}$$

[Out] -12/77*b*sin(f*x+e)/f/(b*sec(f*x+e))^(5/2)-2/11*b*sin(f*x+e)^3/f/(b*sec(f*x+e))^(5/2)+8/77*sin(f*x+e)/b/f/(b*sec(f*x+e))^(1/2)+8/77*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2))*cos(f*x+e)^(1/2)*(b*sec(f*x+e))^(1/2)/b^2/f

Rubi [A]

time = 0.09, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2707, 3854, 3856, 2720}

$$\frac{8\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{77b^2 f} - \frac{2b \sin^3(e+fx)}{11f(b \sec(e+fx))^{5/2}} + \frac{8 \sin(e+fx)}{77bf \sqrt{b \sec(e+fx)}} - \frac{12b \sin(e+fx)}{77f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^4/(b*Sec[e + f*x])^(3/2),x]

[Out] (8*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*Sqrt[b*Sec[e + f*x]])/(77*b^2*f) - (12*b*Sin[e + f*x])/(77*f*(b*Sec[e + f*x])^(5/2)) + (8*Sin[e + f*x])/(77*b*f*Sqrt[b*Sec[e + f*x]]) - (2*b*Sin[e + f*x]^3)/(11*f*(b*Sec[e + f*x])^(5/2))

Rule 2707

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_.)])^n, x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= -\frac{2b \sin^3(e+fx)}{11f(b \sec(e+fx))^{5/2}} + \frac{6}{11} \int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx \\
 &= -\frac{12b \sin(e+fx)}{77f(b \sec(e+fx))^{5/2}} - \frac{2b \sin^3(e+fx)}{11f(b \sec(e+fx))^{5/2}} + \frac{12}{77} \int \frac{1}{(b \sec(e+fx))^{3/2}} dx \\
 &= -\frac{12b \sin(e+fx)}{77f(b \sec(e+fx))^{5/2}} + \frac{8 \sin(e+fx)}{77bf \sqrt{b \sec(e+fx)}} - \frac{2b \sin^3(e+fx)}{11f(b \sec(e+fx))^{5/2}} + \frac{4}{77} \int \frac{1}{\sqrt{b \sec(e+fx)}} dx \\
 &= -\frac{12b \sin(e+fx)}{77f(b \sec(e+fx))^{5/2}} + \frac{8 \sin(e+fx)}{77bf \sqrt{b \sec(e+fx)}} - \frac{2b \sin^3(e+fx)}{11f(b \sec(e+fx))^{5/2}} + \frac{4 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right)}{77b^2 f} \\
 &= \frac{8 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{77b^2 f} - \frac{12b \sin(e+fx)}{77f(b \sec(e+fx))^{5/2}} + \frac{4 \sqrt{\cos(e+fx)}}{77b}
 \end{aligned}$$

Mathematica [A]

time = 0.12, size = 81, normalized size = 0.64

$$\frac{\sec^2(e+fx) \left(128 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) - 5 \sin(2(e+fx)) - 24 \sin(4(e+fx)) + 7 \sin(6(e+fx)) \right)}{1232f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^4/(b*Sec[e + f*x])^(3/2), x]

[Out] (Sec[e + f*x]^2*(128*sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] - 5*Sin[2*(e + f*x)] - 24*Sin[4*(e + f*x)] + 7*Sin[6*(e + f*x)])/(1232*f*(b*Sec[e + f*x])^(3/2))

Maple [C] Result contains complex when optimal does not.

time = 0.28, size = 173, normalized size = 1.37

method	result
--------	--------

default	$\frac{2(\cos(fx+e)+1)^2(-1+\cos(fx+e))\left(7(\cos^6(fx+e))-7(\cos^5(fx+e))-4i\sqrt{\frac{1}{\cos(fx+e)+1}}\sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}}\operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}\right)\right)}{77f\cos(fx+e)^2\sin(fx+e)^3\left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}}$
---------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(f*x+e)^4/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

[Out] $2/77/f*(\cos(f*x+e)+1)^2*(-1+\cos(f*x+e))*(7*\cos(f*x+e)^6-7*\cos(f*x+e)^5-4*I*\operatorname{EllipticF}(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\sin(f*x+e)-13*\cos(f*x+e)^4+13*\cos(f*x+e)^3+4*\cos(f*x+e)^2-4*\cos(f*x+e))/\cos(f*x+e)^2/\sin(f*x+e)^3/(b/\cos(f*x+e))^{3/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate(sin(f*x + e)^4/(b*sec(f*x + e))^(3/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.12, size = 118, normalized size = 0.94

$$\frac{2\left((7\cos(fx+e)^5-13\cos(fx+e)^3+4\cos(fx+e))\sqrt{\frac{b}{\cos(fx+e)}}\sin(fx+e)-2i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+2i\sqrt{2}\sqrt{b}\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))\right)}{77b^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")`

[Out] $2/77*((7*\cos(f*x + e)^5 - 13*\cos(f*x + e)^3 + 4*\cos(f*x + e))*\operatorname{sqrt}(b/\cos(f*x + e))*\sin(f*x + e) - 2*I*\operatorname{sqrt}(2)*\operatorname{sqrt}(b)*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) + 2*I*\operatorname{sqrt}(2)*\operatorname{sqrt}(b)*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)))/(b^2*f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^4(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**4/(b*sec(f*x+e))**(3/2),x)`

[Out] Integral(sin(e + f*x)**4/(b*sec(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^4/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^4}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4/(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^4/(b/cos(e + f*x))^(3/2), x)

$$3.432 \quad \int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=98

$$\frac{4\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{21b^2 f} - \frac{2b \sin(e+fx)}{7f(b \sec(e+fx))^{5/2}} + \frac{4 \sin(e+fx)}{21bf \sqrt{b \sec(e+fx)}}$$

[Out] $-2/7*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(5/2)}+4/21*\sin(f*x+e)/b/f/(b*\sec(f*x+e))^{(1/2)}+4/21*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e), 2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{2/f}$

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2707, 3854, 3856, 2720}

$$\frac{4\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{21b^2 f} + \frac{4 \sin(e+fx)}{21bf \sqrt{b \sec(e+fx)}} - \frac{2b \sin(e+fx)}{7f(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[e + f*x]^2/(b*Sec[e + f*x])^(3/2), x]`

[Out] $(4*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(21*b^{2*f}) - (2*b*\text{Sin}[e + f*x])/(7*f*(b*\text{Sec}[e + f*x])^{(5/2)}) + (4*\text{Sin}[e + f*x])/(21*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2707

`Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3854

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{n*} \text{Sin}[c + d*x]^{n-1}, \text{Int}[1/\text{Sin}[c + d*x]^{n-1}, x], x] /;$ FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= -\frac{2b \sin(e+fx)}{7f(b \sec(e+fx))^{5/2}} + \frac{2}{7} \int \frac{1}{(b \sec(e+fx))^{3/2}} dx \\ &= -\frac{2b \sin(e+fx)}{7f(b \sec(e+fx))^{5/2}} + \frac{4 \sin(e+fx)}{21bf \sqrt{b \sec(e+fx)}} + \frac{2 \int \sqrt{b \sec(e+fx)} dx}{21b^2} \\ &= -\frac{2b \sin(e+fx)}{7f(b \sec(e+fx))^{5/2}} + \frac{4 \sin(e+fx)}{21bf \sqrt{b \sec(e+fx)}} + \frac{(2 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)})}{21b^2} \\ &= \frac{4 \sqrt{\cos(e+fx)} F(\frac{1}{2}(e+fx)|2) \sqrt{b \sec(e+fx)}}{21b^2 f} - \frac{2b \sin(e+fx)}{7f(b \sec(e+fx))^{5/2}} + \frac{2 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}{21b^2} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 71, normalized size = 0.72

$$\frac{\sec^2(e+fx) \left(16 \sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx)|2\right) + 2 \sin(2(e+fx)) - 3 \sin(4(e+fx)) \right)}{84f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^2/(b*Sec[e + f*x])^(3/2), x]

[Out] (Sec[e + f*x]^2*(16*sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] + 2*Sin[2*(e + f*x)] - 3*Sin[4*(e + f*x)])/(84*f*(b*Sec[e + f*x])^(3/2))

Maple [C] Result contains complex when optimal does not.

time = 0.23, size = 153, normalized size = 1.56

method	result
default	$-\frac{2(\cos(fx+e)+1)^2(-1+\cos(fx+e)) \left(2i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \text{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) + 3(\cos^4(fx+e)) \right)}{21f \cos(fx+e)^2 \sin(fx+e)^3 \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^2/(b*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)

[Out]
$$-2/21/f*(\cos(f*x+e)+1)^2*(-1+\cos(f*x+e))*(2*I*EllipticF(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\sin(f*x+e)+3*\cos(f*x+e)^4-3*\cos(f*x+e)^3-2*\cos(f*x+e)^2+2*\cos(f*x+e))/\cos(f*x+e)^2/\sin(f*x+e)^3/(b/\cos(f*x+e))^{3/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate(sin(f*x + e)^2/(b*sec(f*x + e))^(3/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.15, size = 107, normalized size = 1.09

$$\frac{2 \left((3 \cos(fx+e)^3 - 2 \cos(fx+e)) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e)) - i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)) \right)}{21b^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")`

[Out]
$$-2/21*((3*\cos(f*x + e)^3 - 2*\cos(f*x + e))*\sqrt{b/\cos(f*x + e)}*\sin(f*x + e) + I*\sqrt{2}*\sqrt{b}*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e)) - I*\sqrt{2}*\sqrt{b}*\operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e)))/(b^2*f)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^2(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**2/(b*sec(f*x+e))**(3/2),x)`

[Out] `Integral(sin(e + f*x)**2/(b*sec(e + f*x))**(3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^2/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^2}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2/(b/cos(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^2/(b/cos(e + f*x))^(3/2), x)

$$3.433 \quad \int \frac{1}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=72

$$\frac{2\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{3b^2 f} + \frac{2 \sin(e+fx)}{3bf \sqrt{b \sec(e+fx)}}$$

[Out] 2/3*sin(f*x+e)/b/f/(b*sec(f*x+e))^(1/2)+2/3*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2))*cos(f*x+e)^(1/2)*(b*sec(f*x+e))^(1/2)/b^2/f

Rubi [A]

time = 0.03, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3854, 3856, 2720}

$$\frac{2\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{3b^2 f} + \frac{2 \sin(e+fx)}{3bf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^(-3/2),x]

[Out] (2*sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*sqrt[b*Sec[e + f*x]])/(3*b^2*f) + (2*Sin[e + f*x])/(3*b*f*sqrt[b*Sec[e + f*x]])

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(b \sec(e + fx))^{3/2}} dx &= \frac{2 \sin(e + fx)}{3bf \sqrt{b \sec(e + fx)}} + \frac{\int \sqrt{b \sec(e + fx)} dx}{3b^2} \\
&= \frac{2 \sin(e + fx)}{3bf \sqrt{b \sec(e + fx)}} + \frac{\left(\sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{1}{\sqrt{\cos(e + fx)}} dx}{3b^2} \\
&= \frac{2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) \sqrt{b \sec(e + fx)}}{3b^2 f} + \frac{2 \sin(e + fx)}{3bf \sqrt{b \sec(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 59, normalized size = 0.82

$$\frac{\sec^2(e + fx) \left(2 \sqrt{\cos(e + fx)} F\left(\frac{1}{2}(e + fx) \mid 2\right) + \sin(2(e + fx)) \right)}{3f(b \sec(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(-3/2),x]**[Out]** (Sec[e + f*x]^2*(2*Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2] + Sin[2*(e + f*x)])/(3*f*(b*Sec[e + f*x])^(3/2))**Maple [C]** Result contains complex when optimal does not.

time = 0.19, size = 131, normalized size = 1.82

method	result
default	$ -\frac{2(\cos(fx+e)+1)^2(-1+\cos(fx+e)) \left(i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) - (\cos^2(fx+e) - 1) \right)}{3f \sin(fx+e)^3 \cos(fx+e)^2 \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}}} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)**[Out]** -2/3/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))*(I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-cos(f*x+e)^2+cos(f*x+e))/sin(f*x+e)^3/cos(f*x+e)^2/(b/cos(f*x+e))^(3/2)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^(-3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.11, size = 94, normalized size = 1.31

$$2 \sqrt{\frac{b}{\cos(fx+e)}} \frac{\cos(fx+e) \sin(fx+e) - i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e)) + i \sqrt{2} \sqrt{b} \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e))}{3b^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] 1/3*(2*sqrt(b/cos(f*x + e))*cos(f*x + e)*sin(f*x + e) - I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) + I*sqrt(2)*sqrt(b)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)))/(b^2*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))**(3/2),x)

[Out] Integral((b*sec(e + f*x))**(-3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^(-3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{b}{\cos(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b/cos(e + f*x))^(3/2),x)

[Out] int(1/(b/cos(e + f*x))^(3/2), x)

$$3.434 \quad \int \frac{\csc^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=68

$$-\frac{\csc(e+fx)}{bf\sqrt{b\sec(e+fx)}} - \frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b\sec(e+fx)}}{b^2f}$$

[Out] $-\csc(f*x+e)/b/f/(b*\sec(f*x+e))^{(1/2)}-(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticF}(\sin(1/2*f*x+1/2*e),2^{(1/2)})*\cos(f*x+e)^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^2/f$

Rubi [A]

time = 0.05, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2703, 3856, 2720}

$$-\frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b\sec(e+fx)}}{b^2f} - \frac{\csc(e+fx)}{bf\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^2/(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $-(\text{Csc}[e + f*x]/(b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]])) - (\text{Sqrt}[\text{Cos}[e + f*x]]*\text{EllipticF}[(e + f*x)/2, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])/(b^2*f)$

Rule 2703

$\text{Int}[(\csc[(e_) + (f_)*(x_)]*(a_))^{(m_)}*((b_)*\sec[(e_) + (f_)*(x_)])^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\text{Csc}[e + f*x])^{(m-1)}*((b*\text{Sec}[e + f*x])^{(n+1)})/(f*b*(m-1)), x] + \text{Dist}[a^2*((n+1)/(b^2*(m-1))), \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegersQ}[2*m, 2*n]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\csc[(c_) + (d_)*(x_)]*(b_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int \frac{\csc^2(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= -\frac{\csc(e+fx)}{bf \sqrt{b \sec(e+fx)}} - \frac{\int \sqrt{b \sec(e+fx)} dx}{2b^2} \\
&= -\frac{\csc(e+fx)}{bf \sqrt{b \sec(e+fx)}} - \frac{\left(\sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}\right) \int \frac{1}{\sqrt{\cos(e+fx)}} dx}{2b^2} \\
&= -\frac{\csc(e+fx)}{bf \sqrt{b \sec(e+fx)}} - \frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b \sec(e+fx)}}{b^2 f}
\end{aligned}$$

Mathematica [A]

time = 0.08, size = 58, normalized size = 0.85

$$-\frac{\sqrt{\cos(e+fx)} \csc(e+fx) - F\left(\frac{1}{2}(e+fx) \mid 2\right)}{f \cos^{\frac{3}{2}}(e+fx) (b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2/(b*Sec[e + f*x])^(3/2), x]``[Out] (-Sqrt[Cos[e + f*x]]*Csc[e + f*x]) - EllipticF[(e + f*x)/2, 2])/(f*Cos[e + f*x]^(3/2)*(b*Sec[e + f*x])^(3/2))`**Maple [C]** Result contains complex when optimal does not.

time = 0.20, size = 191, normalized size = 2.81

method	result
default	$ -\frac{(-1+\cos(fx+e))^2 \left(i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) + i \sqrt{\frac{1}{\cos(fx+e)+1}} \right)}{f \sin(fx+e)^5 \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}} \cos(fx+e)^5} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2/(b*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] -1/f*(-1+cos(f*x+e))^2*(I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*sin(f*x+e)*cos(f*x+e)+I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)+cos(f*x+e))*(cos(f*x+e)+1)^2/sin(f*x+e)^5/(b/cos(f*x+e))^(3/2)/cos(f*x+e)^2
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate(csc(f*x + e)^2/(b*sec(f*x + e))^(3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.14, size = 110, normalized size = 1.62

$$\frac{i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))-i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))-2\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)}{2b^2f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] 1/2*(I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) - I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)) - 2*sqrt(b/cos(f*x + e))*cos(f*x + e))/(b^2*f*sin(f*x + e))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**2/(b*sec(f*x+e))**(3/2),x)

[Out] Integral(csc(e + f*x)**2/(b*sec(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^2/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + fx)^2 \left(\frac{b}{\cos(e + fx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^2*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^2*(b/cos(e + f*x))^(3/2)), x)

$$3.435 \quad \int \frac{\csc^4(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=102

$$\frac{\csc(e+fx)}{6bf\sqrt{b\sec(e+fx)}} - \frac{\csc^3(e+fx)}{3bf\sqrt{b\sec(e+fx)}} - \frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b\sec(e+fx)}}{6b^2f}$$

[Out] 1/6*csc(f*x+e)/b/f/(b*sec(f*x+e))^(1/2)-1/3*csc(f*x+e)^3/b/f/(b*sec(f*x+e))^(1/2)-1/6*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2))*cos(f*x+e)^(1/2)*(b*sec(f*x+e))^(1/2)/b^2/f

Rubi [A]

time = 0.08, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2703, 2705, 3856, 2720}

$$-\frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b\sec(e+fx)}}{6b^2f} - \frac{\csc^3(e+fx)}{3bf\sqrt{b\sec(e+fx)}} + \frac{\csc(e+fx)}{6bf\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^4/(b*Sec[e + f*x])^(3/2),x]

[Out] Csc[e + f*x]/(6*b*f*Sqrt[b*Sec[e + f*x]]) - Csc[e + f*x]^3/(3*b*f*Sqrt[b*Sec[e + f*x]]) - (Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*Sqrt[b*Sec[e + f*x]])/(6*b^2*f)

Rule 2703

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Dist[a^2*((n + 1)/(b^2*(m - 1))), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]`

Rubi steps

$$\begin{aligned} \int \frac{\csc^4(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= -\frac{\csc^3(e+fx)}{3bf \sqrt{b \sec(e+fx)}} - \frac{\int \csc^2(e+fx) \sqrt{b \sec(e+fx)} dx}{6b^2} \\ &= \frac{\csc(e+fx)}{6bf \sqrt{b \sec(e+fx)}} - \frac{\csc^3(e+fx)}{3bf \sqrt{b \sec(e+fx)}} - \frac{\int \sqrt{b \sec(e+fx)} dx}{12b^2} \\ &= \frac{\csc(e+fx)}{6bf \sqrt{b \sec(e+fx)}} - \frac{\csc^3(e+fx)}{3bf \sqrt{b \sec(e+fx)}} - \frac{\left(\sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}\right)}{12b^2} \\ &= \frac{\csc(e+fx)}{6bf \sqrt{b \sec(e+fx)}} - \frac{\csc^3(e+fx)}{3bf \sqrt{b \sec(e+fx)}} - \frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right)}{6b^2 f} \end{aligned}$$

Mathematica [A]

time = 0.17, size = 62, normalized size = 0.61

$$\frac{\csc(e+fx) - 2 \csc^3(e+fx) - \frac{F\left(\frac{1}{2}(e+fx) \mid 2\right)}{\sqrt{\cos(e+fx)}}}{6bf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^4/(b*Sec[e + f*x])^(3/2), x]`

`[Out] (Csc[e + f*x] - 2*Csc[e + f*x]^3 - EllipticF[(e + f*x)/2, 2]/Sqrt[Cos[e + f*x]])/(6*b*f*Sqrt[b*Sec[e + f*x]])`

Maple [C] Result contains complex when optimal does not.

time = 0.22, size = 343, normalized size = 3.36

method	result
default	$\frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(i \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^3(fx+e) \sin(fx+e)) \right)}{6bf \sqrt{b \sec(e+fx)}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^4/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{6}f(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2(I\cos(fx+e)^3\sin(fx+e)*(1/(\cos(fx+e)+1))^{1/2}+(\cos(fx+e)/(\cos(fx+e)+1))^{1/2})\text{EllipticF}(I(-1+\cos(fx+e))/\sin(fx+e),I)+I\cos(fx+e)^2\sin(fx+e)*(1/(\cos(fx+e)+1))^{1/2}+(\cos(fx+e)/(\cos(fx+e)+1))^{1/2})\text{EllipticF}(I(-1+\cos(fx+e))/\sin(fx+e),I)-I(1/(\cos(fx+e)+1))^{1/2}+(\cos(fx+e)/(\cos(fx+e)+1))^{1/2})\text{EllipticF}(I(-1+\cos(fx+e))/\sin(fx+e),I)*\sin(fx+e)\cos(fx+e)-I\text{EllipticF}(I(-1+\cos(fx+e))/\sin(fx+e),I)*(1/(\cos(fx+e)+1))^{1/2}+(\cos(fx+e)/(\cos(fx+e)+1))^{1/2})\sin(fx+e)-\cos(fx+e)^3-\cos(fx+e))/\cos(fx+e)^2/\sin(fx+e)^7/(b/\cos(fx+e))^{3/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate(csc(f*x + e)^4/(b*sec(f*x + e))^(3/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 161, normalized size = 1.58

$$\frac{\sqrt{2}(i\cos(fx+e)^2-i)\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+\sqrt{2}(-i\cos(fx+e)^2+i)\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))+2(\cos(fx+e)^3+\cos(fx+e))\sqrt{\frac{b}{\cos(fx+e)}}}{12(b^2f\cos(fx+e)^2-b^2f)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")`

[Out] $\frac{1}{12}(\sqrt{2}(I\cos(fx+e)^2-I)\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e)+I\sin(fx+e))+\sqrt{2}(-I\cos(fx+e)^2+I)\sqrt{b}\sin(fx+e)\text{weierstrassPInverse}(-4,0,\cos(fx+e)-I\sin(fx+e))+2(\cos(fx+e)^3+\cos(fx+e))\sqrt{b/\cos(fx+e)})/(b^2f\cos(fx+e)^2-b^2f)\sin(fx+e)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^4(e+fx)}{(b\sec(e+fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**4/(b*sec(f*x+e))**(3/2),x)`

[Out] `Integral(csc(e + f*x)**4/(b*sec(e + f*x))**(3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(3/2),x, algorithm="giac")``[Out] integrate(csc(f*x + e)^4/(b*sec(f*x + e))^(3/2), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^4 \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(sin(e + f*x)^4*(b/cos(e + f*x))^(3/2)),x)``[Out] int(1/(sin(e + f*x)^4*(b/cos(e + f*x))^(3/2)), x)`

$$3.436 \quad \int \frac{\csc^6(e+fx)}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=132

$$\frac{\csc(e+fx)}{12bf\sqrt{b\sec(e+fx)}} + \frac{\csc^3(e+fx)}{30bf\sqrt{b\sec(e+fx)}} - \frac{\csc^5(e+fx)}{5bf\sqrt{b\sec(e+fx)}} - \frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b\sec(e+fx)}}{12b^2f}$$

[Out] 1/12*csc(f*x+e)/b/f/(b*sec(f*x+e))^(1/2)+1/30*csc(f*x+e)^3/b/f/(b*sec(f*x+e))^(1/2)-1/5*csc(f*x+e)^5/b/f/(b*sec(f*x+e))^(1/2)-1/12*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticF(sin(1/2*f*x+1/2*e),2^(1/2))*cos(f*x+e)^(1/2)*(b*sec(f*x+e))^(1/2)/b^2/f

Rubi [A]

time = 0.10, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2703, 2705, 3856, 2720}

$$-\frac{\sqrt{\cos(e+fx)} F\left(\frac{1}{2}(e+fx) \mid 2\right) \sqrt{b\sec(e+fx)}}{12b^2f} - \frac{\csc^5(e+fx)}{5bf\sqrt{b\sec(e+fx)}} + \frac{\csc^3(e+fx)}{30bf\sqrt{b\sec(e+fx)}} + \frac{\csc(e+fx)}{12bf\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^6/(b*Sec[e + f*x])^(3/2),x]

[Out] Csc[e + f*x]/(12*b*f*Sqrt[b*Sec[e + f*x]]) + Csc[e + f*x]^3/(30*b*f*Sqrt[b*Sec[e + f*x]]) - Csc[e + f*x]^5/(5*b*f*Sqrt[b*Sec[e + f*x]]) - (Sqrt[Cos[e + f*x]]*EllipticF[(e + f*x)/2, 2]*Sqrt[b*Sec[e + f*x]])/(12*b^2*f)

Rule 2703

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Dist[a^2*((n + 1)/(b^2*(m - 1))), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^6(e+fx)}{(b \sec(e+fx))^{3/2}} dx &= -\frac{\csc^5(e+fx)}{5bf \sqrt{b \sec(e+fx)}} - \frac{\int \csc^4(e+fx) \sqrt{b \sec(e+fx)} dx}{10b^2} \\
 &= \frac{\csc^3(e+fx)}{30bf \sqrt{b \sec(e+fx)}} - \frac{\csc^5(e+fx)}{5bf \sqrt{b \sec(e+fx)}} - \frac{\int \csc^2(e+fx) \sqrt{b \sec(e+fx)} dx}{12b^2} \\
 &= \frac{\csc(e+fx)}{12bf \sqrt{b \sec(e+fx)}} + \frac{\csc^3(e+fx)}{30bf \sqrt{b \sec(e+fx)}} - \frac{\csc^5(e+fx)}{5bf \sqrt{b \sec(e+fx)}} - \frac{\int \sqrt{b \sec(e+fx)} dx}{12b^2} \\
 &= \frac{\csc(e+fx)}{12bf \sqrt{b \sec(e+fx)}} + \frac{\csc^3(e+fx)}{30bf \sqrt{b \sec(e+fx)}} - \frac{\csc^5(e+fx)}{5bf \sqrt{b \sec(e+fx)}} - \frac{\int \sqrt{\cos(e+fx)} dx}{12b^2} \\
 &= \frac{\csc(e+fx)}{12bf \sqrt{b \sec(e+fx)}} + \frac{\csc^3(e+fx)}{30bf \sqrt{b \sec(e+fx)}} - \frac{\csc^5(e+fx)}{5bf \sqrt{b \sec(e+fx)}} - \frac{\int \sqrt{\cos(e+fx)} dx}{12b^2}
 \end{aligned}$$

Mathematica [A]

time = 0.24, size = 74, normalized size = 0.56

$$\frac{5 \csc(e+fx) + 2 \csc^3(e+fx) - 12 \csc^5(e+fx) - \frac{5F(\frac{1}{2}(e+fx)|2)}{\sqrt{\cos(e+fx)}}}{60bf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^6/(b*Sec[e + f*x])^(3/2), x]

[Out] (5*Csc[e + f*x] + 2*Csc[e + f*x]^3 - 12*Csc[e + f*x]^5 - (5*EllipticF[(e + f*x)/2, 2])/Sqrt[Cos[e + f*x]])/(60*b*f*Sqrt[b*Sec[e + f*x]])

Maple [C] Result contains complex when optimal does not.

time = 0.26, size = 493, normalized size = 3.73

method	result
default	$-\frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(5i \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^5(fx+e)) \sin(fx+e) \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)^6/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/60/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(5*I*cos(f*x+e)^5*sin(f*x+e)*(1/
(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos
(f*x+e))/sin(f*x+e),I)+5*I*cos(f*x+e)^4*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)
*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I
)-10*I*cos(f*x+e)^3*sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*
x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)-10*I*cos(f*x+e)^2*
sin(f*x+e)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*Ellip
ticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)+5*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+
e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+
e)*cos(f*x+e)+5*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+
1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-5*cos(f*x+e)^5+12*co
s(f*x+e)^3+5*cos(f*x+e))/cos(f*x+e)^2/sin(f*x+e)^9/(b/cos(f*x+e))^(3/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^6/(b*sec(f*x + e))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 214, normalized size = 1.62

$$\frac{5\sqrt{2}(-i\cos(fx+e)^5+2i\cos(fx+e)^3-i)\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))+5\sqrt{2}(i\cos(fx+e)^5-2i\cos(fx+e)^3+i)\sqrt{b}\sin(fx+e)\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))-2(5\cos(fx+e)^9-12\cos(fx+e)^7-5\cos(fx+e)^5)\sqrt{\frac{b}{\cos(fx+e)}}}{120(b^2f\cos(fx+e)^4-2b^2f\cos(fx+e)^2+b^2f)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] -1/120*(5*sqrt(2)*(-I*cos(f*x + e)^4 + 2*I*cos(f*x + e)^2 - I)*sqrt(b)*sin(
f*x + e)*weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e)) + 5*sqrt
(2)*(I*cos(f*x + e)^4 - 2*I*cos(f*x + e)^2 + I)*sqrt(b)*sin(f*x + e)*weiers
trassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)) - 2*(5*cos(f*x + e)^5 -
```

$12*\cos(f*x + e)^3 - 5*\cos(f*x + e))*\sqrt{b/\cos(f*x + e))}/((b^2*f*\cos(f*x + e)^4 - 2*b^2*f*\cos(f*x + e)^2 + b^2*f)*\sin(f*x + e))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^6(e + fx)}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**6/(b*sec(f*x+e))**(3/2),x)

[Out] Integral(csc(e + f*x)**6/(b*sec(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^6/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + fx)^6 \left(\frac{b}{\cos(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^6*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^6*(b/cos(e + f*x))^(3/2)), x)

$$3.437 \quad \int \frac{\sin^7(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=87

$$\frac{2b^7}{19f(b \sec(e+fx))^{19/2}} - \frac{2b^5}{5f(b \sec(e+fx))^{15/2}} + \frac{6b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

[Out] $2/19*b^7/f/(b*\sec(f*x+e))^{(19/2)}-2/5*b^5/f/(b*\sec(f*x+e))^{(15/2)}+6/11*b^3/f/(b*\sec(f*x+e))^{(11/2)}-2/7*b/f/(b*\sec(f*x+e))^{(7/2)}$

Rubi [A]

time = 0.04, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$\frac{2b^7}{19f(b \sec(e+fx))^{19/2}} - \frac{2b^5}{5f(b \sec(e+fx))^{15/2}} + \frac{6b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^7/(b*Sec[e + f*x])^(5/2),x]

[Out] $(2*b^7)/(19*f*(b*Sec[e + f*x])^{(19/2)}) - (2*b^5)/(5*f*(b*Sec[e + f*x])^{(15/2)}) + (6*b^3)/(11*f*(b*Sec[e + f*x])^{(11/2)}) - (2*b)/(7*f*(b*Sec[e + f*x])^{(7/2)})$

Rule 276

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] :> Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)]^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin^7(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= \frac{b^7 \text{Subst} \left(\int \frac{\left(-1 + \frac{x^2}{b^2}\right)^3}{x^{21/2}} dx, x, b \sec(e+fx) \right)}{f} \\ &= \frac{b^7 \text{Subst} \left(\int \left(-\frac{1}{x^{21/2}} + \frac{3}{b^2 x^{17/2}} - \frac{3}{b^4 x^{13/2}} + \frac{1}{b^6 x^{9/2}}\right) dx, x, b \sec(e+fx) \right)}{f} \\ &= \frac{2b^7}{19f(b \sec(e+fx))^{19/2}} - \frac{2b^5}{5f(b \sec(e+fx))^{15/2}} + \frac{6b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}} \end{aligned}$$

Mathematica [A]

time = 0.26, size = 62, normalized size = 0.71

$$\frac{\cos^4(e+fx)(-15226 + 14287 \cos(2(e+fx)) - 3542 \cos(4(e+fx)) + 385 \cos(6(e+fx))) \sqrt{b \sec(e+fx)}}{117040b^3 f}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^7/(b*Sec[e + f*x])^(5/2), x]`
`[Out] (Cos[e + f*x]^4*(-15226 + 14287*Cos[2*(e + f*x)] - 3542*Cos[4*(e + f*x)] + 385*Cos[6*(e + f*x)])*Sqrt[b*Sec[e + f*x]]/(117040*b^3*f)`
Maple [A]

time = 0.27, size = 56, normalized size = 0.64

method	result	size
default	$\frac{2(385(\cos^6(fx+e)) - 1463(\cos^4(fx+e)) + 1995(\cos^2(fx+e)) - 1045)\cos(fx+e)}{7315f \left(\frac{b}{\cos(fx+e)}\right)^{\frac{5}{2}}}$	56

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^7/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)`
`[Out] 2/7315/f*(385*cos(f*x+e)^6-1463*cos(f*x+e)^4+1995*cos(f*x+e)^2-1045)*cos(f*x+e)/(b/cos(f*x+e))^(5/2)`
Maxima [A]

time = 0.29, size = 67, normalized size = 0.77

$$\frac{2 \left(385 b^6 - \frac{1463 b^6}{\cos(fx+e)^2} + \frac{1995 b^6}{\cos(fx+e)^4} - \frac{1045 b^6}{\cos(fx+e)^6} \right) b}{7315 f \left(\frac{b}{\cos(fx+e)} \right)^{\frac{19}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] $2/7315*(385*b^6 - 1463*b^6/\cos(f*x + e)^2 + 1995*b^6/\cos(f*x + e)^4 - 1045*b^6/\cos(f*x + e)^6)*b/(f*(b/\cos(f*x + e))^{(19/2)})$

Fricas [A]

time = 0.41, size = 66, normalized size = 0.76

$$\frac{2(385 \cos(fx + e)^{10} - 1463 \cos(fx + e)^8 + 1995 \cos(fx + e)^6 - 1045 \cos(fx + e)^4) \sqrt{\frac{b}{\cos(fx + e)}}}{7315 b^3 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] $2/7315*(385*\cos(f*x + e)^{10} - 1463*\cos(f*x + e)^8 + 1995*\cos(f*x + e)^6 - 1045*\cos(f*x + e)^4)*\sqrt{b/\cos(f*x + e)}/(b^3*f)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**7/(b*sec(f*x+e))**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 5007 deep

Giac [A]

time = 5.32, size = 119, normalized size = 1.37

$$\frac{2(385 \sqrt{b \cos(fx + e)} b^9 \cos(fx + e)^9 - 1463 \sqrt{b \cos(fx + e)} b^9 \cos(fx + e)^7 + 1995 \sqrt{b \cos(fx + e)} b^9 \cos(fx + e)^5 - 1045 \sqrt{b \cos(fx + e)} b^9 \cos(fx + e)^3)}{7315 b^{12} f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^7/(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] $2/7315*(385*\sqrt{b*\cos(f*x + e)}*b^9*\cos(f*x + e)^9 - 1463*\sqrt{b*\cos(f*x + e)}*b^9*\cos(f*x + e)^7 + 1995*\sqrt{b*\cos(f*x + e)}*b^9*\cos(f*x + e)^5 - 1045*\sqrt{b*\cos(f*x + e)}*b^9*\cos(f*x + e)^3)/(b^{12}*f*\operatorname{sgn}(\cos(f*x + e)))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^7}{\left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^7/(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^7/(b/cos(e + f*x))^(5/2), x)

$$3.438 \quad \int \frac{\sin^5(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=65

$$-\frac{2b^5}{15f(b \sec(e+fx))^{15/2}} + \frac{4b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

[Out] $-2/15*b^5/f/(b*\sec(f*x+e))^{(15/2)}+4/11*b^3/f/(b*\sec(f*x+e))^{(11/2)}-2/7*b/f/(b*\sec(f*x+e))^{(7/2)}$

Rubi [A]

time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 276}

$$-\frac{2b^5}{15f(b \sec(e+fx))^{15/2}} + \frac{4b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[e + f*x]^5/(b*Sec[e + f*x])^(5/2),x]`

[Out] $(-2*b^5)/(15*f*(b*Sec[e + f*x])^{(15/2)}) + (4*b^3)/(11*f*(b*Sec[e + f*x])^{(11/2)}) - (2*b)/(7*f*(b*Sec[e + f*x])^{(7/2)})$

Rule 276

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_.), x_Symbol] := Int[Exp andIntegrand[(c*x)^m*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned}
\int \frac{\sin^5(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= \frac{b^5 \text{Subst} \left(\int \frac{(-1+\frac{x^2}{b^2})^2}{x^{17/2}} dx, x, b \sec(e+fx) \right)}{f} \\
&= \frac{b^5 \text{Subst} \left(\int \left(\frac{1}{x^{17/2}} - \frac{2}{b^2 x^{13/2}} + \frac{1}{b^4 x^{9/2}} \right) dx, x, b \sec(e+fx) \right)}{f} \\
&= -\frac{2b^5}{15f(b \sec(e+fx))^{15/2}} + \frac{4b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}
\end{aligned}$$

Mathematica [A]

time = 0.16, size = 52, normalized size = 0.80

$$\frac{\cos^4(e+fx)(-711+532\cos(2(e+fx))-77\cos(4(e+fx)))\sqrt{b \sec(e+fx)}}{4620b^3f}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^5/(b*Sec[e + f*x])^(5/2), x]``[Out] (Cos[e + f*x]^4*(-711 + 532*Cos[2*(e + f*x)] - 77*Cos[4*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/(4620*b^3*f)`**Maple [A]**

time = 0.20, size = 46, normalized size = 0.71

method	result	size
default	$-\frac{2(77\cos^4(fx+e)-210(\cos^2(fx+e))+165)\cos(fx+e)}{1155f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{5}{2}}}$	46

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^5/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)``[Out] -2/1155/f*(77*cos(f*x+e)^4-210*cos(f*x+e)^2+165)*cos(f*x+e)/(b/cos(f*x+e))^(5/2)`**Maxima [A]**

time = 0.30, size = 53, normalized size = 0.82

$$-\frac{2\left(77b^4 - \frac{210b^4}{\cos(fx+e)^2} + \frac{165b^4}{\cos(fx+e)^4}\right)b}{1155f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{15}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] -2/1155*(77*b^4 - 210*b^4/cos(f*x + e)^2 + 165*b^4/cos(f*x + e)^4)*b/(f*(b/cos(f*x + e))^(15/2))

Fricas [A]

time = 0.37, size = 55, normalized size = 0.85

$$\frac{2 \left(77 \cos (fx + e)^8 - 210 \cos (fx + e)^6 + 165 \cos (fx + e)^4 \right) \sqrt{\frac{b}{\cos (fx + e)}}}{1155 b^3 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] -2/1155*(77*cos(f*x + e)^8 - 210*cos(f*x + e)^6 + 165*cos(f*x + e)^4)*sqrt(b/cos(f*x + e))/(b^3*f)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**5/(b*sec(f*x+e))**(5/2),x)

[Out] Timed out

Giac [A]

time = 4.33, size = 94, normalized size = 1.45

$$\frac{2 \left(77 \sqrt{b \cos (fx + e)} b^7 \cos (fx + e)^7 - 210 \sqrt{b \cos (fx + e)} b^7 \cos (fx + e)^5 + 165 \sqrt{b \cos (fx + e)} b^7 \cos (fx + e)^3 \right)}{1155 b^{10} f \operatorname{sgn}(\cos (fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^5/(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] -2/1155*(77*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e)^7 - 210*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e)^5 + 165*sqrt(b*cos(f*x + e))*b^7*cos(f*x + e)^3)/(b^10*f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin (e + f x)^5}{\left(\frac{b}{\cos (e + f x)} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^5/(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^5/(b/cos(e + f*x))^(5/2), x)

$$3.439 \quad \int \frac{\sin^3(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=43

$$\frac{2b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

[Out] $2/11*b^3/f/(b*\sec(f*x+e))^{(11/2)}-2/7*b/f/(b*\sec(f*x+e))^{(7/2)}$

Rubi [A]

time = 0.03, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2702, 14}

$$\frac{2b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[e + f*x]^3/(b*Sec[e + f*x])^(5/2),x]`

[Out] $(2*b^3)/(11*f*(b*Sec[e + f*x])^{(11/2)}) - (2*b)/(7*f*(b*Sec[e + f*x])^{(7/2)})$

Rule 14

`Int[(u_)*((c_.)*(x_))^(m_.), x_Symbol] := Int[ExpandIntegrand[(c*x)^m*u, x], x] /; FreeQ[{c, m}, x] && SumQ[u] && !LinearQ[u, x] && !MatchQ[u, (a_ + (b_.)*(v_)) /; FreeQ[{a, b}, x] && InverseFunctionQ[v]]`

Rule 2702

`Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m+n-1)/(-1+x^2/a^2)^((n+1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n+1)/2] && !(IntegerQ[(m+1)/2] && LtQ[0, m, n])`

Rubi steps

$$\begin{aligned} \int \frac{\sin^3(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= \frac{b^3 \text{Subst}\left(\int \frac{-1+\frac{x^2}{b^2}}{x^{13/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{b^3 \text{Subst}\left(\int \left(-\frac{1}{x^{13/2}} + \frac{1}{b^2 x^{9/2}}\right) dx, x, b \sec(e+fx)\right)}{f} \\ &= \frac{2b^3}{11f(b \sec(e+fx))^{11/2}} - \frac{2b}{7f(b \sec(e+fx))^{7/2}} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 42, normalized size = 0.98

$$\frac{\cos^4(e + fx)(-15 + 7 \cos(2(e + fx))) \sqrt{b \sec(e + fx)}}{77b^3 f}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^3/(b*Sec[e + f*x])^(5/2), x]

[Out] (Cos[e + f*x]^4*(-15 + 7*Cos[2*(e + f*x)])*Sqrt[b*Sec[e + f*x]])/(77*b^3*f)

Maple [A]

time = 0.18, size = 36, normalized size = 0.84

method	result	size
default	$\frac{2(7(\cos^2(fx+e))-11)\cos(fx+e)}{77f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{5}{2}}}$	36

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^3/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)

[Out] 2/77/f*(7*cos(f*x+e)^2-11)*cos(f*x+e)/(b/cos(f*x+e))^(5/2)

Maxima [A]

time = 0.29, size = 39, normalized size = 0.91

$$\frac{2\left(7b^2 - \frac{11b^2}{\cos(fx+e)^2}\right)b}{77f\left(\frac{b}{\cos(fx+e)}\right)^{\frac{11}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(5/2), x, algorithm="maxima")

[Out] 2/77*(7*b^2 - 11*b^2/cos(f*x + e)^2)*b/(f*(b/cos(f*x + e))^(11/2))

Fricas [A]

time = 0.40, size = 44, normalized size = 1.02

$$\frac{2(7 \cos(fx + e)^6 - 11 \cos(fx + e)^4) \sqrt{\frac{b}{\cos(fx + e)}}}{77b^3 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(5/2), x, algorithm="fricas")

[Out] $2/77*(7*\cos(f*x + e)^6 - 11*\cos(f*x + e)^4)*\sqrt{b/\cos(f*x + e)}/(b^3*f)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**3/(b*sec(f*x+e))**(5/2),x)`

[Out] Timed out

Giac [A]

time = 4.73, size = 69, normalized size = 1.60

$$\frac{2 \left(7 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e)^5 - 11 \sqrt{b \cos(fx + e)} b^5 \cos(fx + e)^3 \right)}{77 b^8 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^3/(b*sec(f*x+e))^(5/2),x, algorithm="giac")`

[Out] $2/77*(7*\sqrt{b*\cos(f*x + e)}*b^5*\cos(f*x + e)^5 - 11*\sqrt{b*\cos(f*x + e)}*b^5*\cos(f*x + e)^3)/(b^8*f*\operatorname{sgn}(\cos(f*x + e)))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sin(e + f x)^3}{\left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^3/(b/cos(e + f*x))^(5/2),x)`

[Out] `int(sin(e + f*x)^3/(b/cos(e + f*x))^(5/2), x)`

$$3.440 \quad \int \frac{\sin(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=20

$$-\frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

[Out] $-2/7*b/f/(b*\sec(f*x+e))^{(7/2)}$

Rubi [A]

time = 0.03, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 30}

$$-\frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]/(b*Sec[e + f*x])^(5/2), x]

[Out] $(-2*b)/(7*f*(b*Sec[e + f*x])^{(7/2)})$

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2702

Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \frac{\sin(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= \frac{b \text{Subst}\left(\int \frac{1}{x^{9/2}} dx, x, b \sec(e+fx)\right)}{f} \\ &= -\frac{2b}{7f(b \sec(e+fx))^{7/2}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 20, normalized size = 1.00

$$-\frac{2b}{7f(b \sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]/(b*Sec[e + f*x])^(5/2),x]

[Out] $(-2*b)/(7*f*(b*\text{Sec}[e + f*x])^{(7/2)})$

Maple [A]

time = 0.05, size = 17, normalized size = 0.85

method	result	size
derivativedivides	$-\frac{2b}{7f(b\sec(fx+e))^{\frac{7}{2}}}$	17
default	$-\frac{2b}{7f(b\sec(fx+e))^{\frac{7}{2}}}$	17

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)/(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)

[Out] $-2/7*b/f/(b*\text{sec}(f*x+e))^{(7/2)}$

Maxima [A]

time = 0.28, size = 25, normalized size = 1.25

$$-\frac{2 \cos(fx + e)}{7f \left(\frac{b}{\cos(fx+e)}\right)^{\frac{5}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] $-2/7*\cos(f*x + e)/(f*(b/\cos(f*x + e))^{(5/2)})$

Fricas [A]

time = 0.41, size = 30, normalized size = 1.50

$$-\frac{2 \sqrt{\frac{b}{\cos(fx + e)}} \cos(fx + e)^4}{7b^3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] $-2/7*\text{sqrt}(b/\cos(f*x + e))*\cos(f*x + e)^4/(b^3*f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin(e + fx)}{(b \sec(e + fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))**(5/2),x)

[Out] Integral(sin(e + f*x)/(b*sec(e + f*x))**(5/2), x)

Giac [B] Leaf count of result is larger than twice the leaf count of optimal. 38 vs. 2(17) = 34.
time = 4.26, size = 38, normalized size = 1.90

$$-\frac{2 \sqrt{b \cos(fx + e)} \cos(fx + e)^3}{7 b^3 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] -2/7*sqrt(b*cos(f*x + e))*cos(f*x + e)^3/(b^3*f*sgn(cos(f*x + e)))

Mupad [B]

time = 0.57, size = 28, normalized size = 1.40

$$-\frac{2 \cos(e + fx)^4 \sqrt{\frac{b}{\cos(e + fx)}}}{7 b^3 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)/(b/cos(e + f*x))^(5/2),x)

[Out] -(2*cos(e + f*x)^4*(b/cos(e + f*x))^(1/2))/(7*b^3*f)

$$3.441 \quad \int \frac{\csc(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=81

$$-\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{5/2}f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{5/2}f} + \frac{2}{3bf(b \sec(e+fx))^{3/2}}$$

[Out] $-\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/b^{(5/2)}/f - \operatorname{arctanh}((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/b^{(5/2)}/f + 2/3/b/f/(b*\sec(f*x+e))^{(3/2)}$

Rubi [A]

time = 0.04, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.316$, Rules used = {2702, 331, 335, 218, 212, 209}

$$-\frac{\operatorname{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{5/2}f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{b^{5/2}f} + \frac{2}{3bf(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Csc}[e + f*x]/(b*\operatorname{Sec}[e + f*x])^{(5/2)}, x]$

[Out] $-(\operatorname{ArcTan}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(b^{(5/2)*f})) - \operatorname{ArcTanh}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]/\operatorname{Sqrt}[b]]/(b^{(5/2)*f}) + 2/(3*b*f*(b*\operatorname{Sec}[e + f*x])^{(3/2)})$

Rule 209

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{PosQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{GtQ}[b, 0])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b] \&\& (\operatorname{GtQ}[a, 0] \parallel \operatorname{LtQ}[b, 0])$

Rule 218

$\operatorname{Int}[(a_ + (b_)*(x_)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& !\operatorname{GtQ}[a/b, 0]$

Rule 331

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*c^n*(m + 1))], Int[(c*x)^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && LtQ[m, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n)]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2)], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc(e + fx)}{(b \sec(e + fx))^{5/2}} dx &= \frac{\text{Subst}\left(\int \frac{1}{x^{5/2}(-1 + \frac{x^2}{b^2})} dx, x, b \sec(e + fx)\right)}{bf} \\
&= \frac{2}{3bf(b \sec(e + fx))^{3/2}} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x}(-1 + \frac{x^2}{b^2})} dx, x, b \sec(e + fx)\right)}{b^3 f} \\
&= \frac{2}{3bf(b \sec(e + fx))^{3/2}} + \frac{2 \text{Subst}\left(\int \frac{1}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{b^3 f} \\
&= \frac{2}{3bf(b \sec(e + fx))^{3/2}} - \frac{\text{Subst}\left(\int \frac{1}{b-x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{b^2 f} - \frac{\text{Subst}\left(\int \frac{1}{b+x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{b^2 f} \\
&= -\frac{\tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{b^{5/2} f} - \frac{\tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{b^{5/2} f} + \frac{2}{3bf(b \sec(e + fx))^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.16, size = 90, normalized size = 1.11

$$\frac{\left(-6 \tan^{-1}\left(\sqrt{\sec(e+fx)}\right) + 3 \log\left(1 - \sqrt{\sec(e+fx)}\right) - 3 \log\left(1 + \sqrt{\sec(e+fx)}\right) + \frac{4}{\sec^{\frac{3}{2}}(e+fx)}\right) \sqrt{\sec(e+fx)}}{6b^2 f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]/(b*Sec[e + f*x])^(5/2), x]

[Out] ((-6*ArcTan[Sqrt[Sec[e + f*x]]] + 3*Log[1 - Sqrt[Sec[e + f*x]]] - 3*Log[1 + Sqrt[Sec[e + f*x]]] + 4/Sec[e + f*x]^(3/2))*Sqrt[Sec[e + f*x]])/(6*b^2*f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 376 vs. $2(65) = 130$.

time = 0.21, size = 377, normalized size = 4.65

method	result
default	$\frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(3 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} \arctan\left(\frac{1}{2 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}}}\right) \cos(fx+e) + 3 \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)

[Out] -1/6/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(3*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))*cos(f*x+e)+3*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*cos(f*x+e)+3*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+3*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-4*cos(f*x+e)^2/sin(f*x+e)^4/(b/cos(f*x+e))^(5/2)/cos(f*x+e)^3

Maxima [A]

time = 0.51, size = 94, normalized size = 1.16

$$\frac{b \left(6 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right) - 3 \log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right) - \frac{4}{b^2 \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}}} \right)}{6f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] $-1/6*b*(6*\arctan(\sqrt{b/\cos(f*x+e)})/\sqrt{b})/b^{7/2} - 3*\log(-(\sqrt{b} - \sqrt{b/\cos(f*x+e)})/(\sqrt{b} + \sqrt{b/\cos(f*x+e)}))/b^{7/2} - 4/(b^2*(b/\cos(f*x+e))^{3/2})/f$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 167 vs. 2(68) = 136.

time = 0.49, size = 343, normalized size = 4.23

$$\frac{8\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)^2 + 6\sqrt{-b} \arctan\left(\frac{\pm\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\cos(fx+e)}\right) - 3\sqrt{-b} \log\left(\frac{\cos(fx+e)^2 - 4(\cos(fx+e)^2 - \cos(fx+e))\sqrt{-b}}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right)}{12b^3f} - \frac{8\sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)^2 - 6\sqrt{b} \arctan\left(\frac{\pm\sqrt{b}\sqrt{\frac{b}{\cos(fx+e)}}}{\cos(fx+e)}\right) + 3\sqrt{b} \log\left(\frac{\cos(fx+e)^2 - 4(\cos(fx+e)^2 + \cos(fx+e))\sqrt{b}}{\cos(fx+e)^2 - 2\cos(fx+e) + 1}\right)}{12b^3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] $[1/12*(8*\sqrt{b/\cos(f*x+e)}*\cos(f*x+e)^2 + 6*\sqrt{-b}*\arctan(2*\sqrt{-b}*\sqrt{b/\cos(f*x+e)}*\cos(f*x+e)/(b*\cos(f*x+e) + b)) - 3*\sqrt{-b}*\log(-(b*\cos(f*x+e)^2 - 4*(\cos(f*x+e)^2 - \cos(f*x+e))*\sqrt{-b}*\sqrt{b/\cos(f*x+e)} - 6*b*\cos(f*x+e) + b)/(\cos(f*x+e)^2 + 2*\cos(f*x+e) + 1)))/(b^3*f), 1/12*(8*\sqrt{b/\cos(f*x+e)}*\cos(f*x+e)^2 - 6*\sqrt{b}*\arctan(2*\sqrt{b}*\sqrt{b/\cos(f*x+e)}*\cos(f*x+e)/(b*\cos(f*x+e) - b)) + 3*\sqrt{b}*\log(-(b*\cos(f*x+e)^2 - 4*(\cos(f*x+e)^2 + \cos(f*x+e))*\sqrt{b}*\sqrt{b/\cos(f*x+e)} + 6*b*\cos(f*x+e) + b)/(\cos(f*x+e)^2 - 2*\cos(f*x+e) + 1)))/(b^3*f)]$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(e+fx)}{(b \sec(e+fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))**(5/2),x)

[Out] Integral(csc(e + f*x)/(b*sec(e + f*x))**(5/2), x)

Giac [A]

time = 4.64, size = 86, normalized size = 1.06

$$\frac{3b \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b}} + 3\sqrt{b} \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right) + 2\sqrt{b \cos(fx + e)} \cos(fx + e)}{3b^3 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] 1/3*(3*b*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b) + 3*sqrt(b)*arctan(sqrt(b*cos(f*x + e))/sqrt(b)) + 2*sqrt(b*cos(f*x + e))*cos(f*x + e))/(b^3*f*sgn(cos(f*x + e)))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + fx) \left(\frac{b}{\cos(e+fx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)*(b/cos(e + f*x))^(5/2)),x)

[Out] int(1/(sin(e + f*x)*(b/cos(e + f*x))^(5/2)), x)

$$3.442 \quad \int \frac{\csc^3(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=93

$$\frac{3 \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{5/2}f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{5/2}f} - \frac{\cot^2(e+fx)\sqrt{b \sec(e+fx)}}{2b^3f}$$

[Out] $3/4*\arctan((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/b^{(5/2)}/f+3/4*\arctanh((b*\sec(f*x+e))^{(1/2)}/b^{(1/2)})/b^{(5/2)}/f-1/2*\cot(f*x+e)^2*(b*\sec(f*x+e))^{(1/2)}/b^3/f$

Rubi [A]

time = 0.05, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {2702, 296, 335, 218, 212, 209}

$$\frac{3 \text{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{5/2}f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{4b^{5/2}f} - \frac{\cot^2(e+fx)\sqrt{b \sec(e+fx)}}{2b^3f}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^3/(b*Sec[e + f*x])^(5/2),x]`

[Out] `(3*ArcTan[Sqrt[b*Sec[e + f*x]]/Sqrt[b]])/(4*b^(5/2)*f) + (3*ArcTanh[Sqrt[b*Sec[e + f*x]]/Sqrt[b]])/(4*b^(5/2)*f) - (Cot[e + f*x]^2*Sqrt[b*Sec[e + f*x]])/(2*b^3*f)`

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]`

Rule 296

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(c*x)^(m + 1))*((a + b*x^n)^(p + 1)/(a*c*n*(p + 1))), x] + Dist[(m + n*(p + 1) + 1)/(a*n*(p + 1)), Int[(c*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a, b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k = Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n))^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && FractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\int \frac{\csc^3(e + fx)}{(b \sec(e + fx))^{5/2}} dx = \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x} \left(-1 + \frac{x^2}{b^2}\right)^2} dx, x, b \sec(e + fx)\right)}{b^3 f}$$

$$= -\frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2b^3 f} - \frac{3 \text{Subst}\left(\int \frac{1}{\sqrt{x} \left(-1 + \frac{x^2}{b^2}\right)} dx, x, b \sec(e + fx)\right)}{4b^3 f}$$

$$= -\frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2b^3 f} - \frac{3 \text{Subst}\left(\int \frac{1}{-1 + \frac{x^4}{b^2}} dx, x, \sqrt{b \sec(e + fx)}\right)}{2b^3 f}$$

$$= -\frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2b^3 f} + \frac{3 \text{Subst}\left(\int \frac{1}{b - x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4b^2 f} + \frac{3 \text{Subst}\left(\int \frac{1}{b + x^2} dx, x, \sqrt{b \sec(e + fx)}\right)}{4b^2 f}$$

$$= \frac{3 \tan^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4b^{5/2} f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e + fx)}}{\sqrt{b}}\right)}{4b^{5/2} f} - \frac{\cot^2(e + fx) \sqrt{b \sec(e + fx)}}{2b^3 f}$$

Mathematica [A]

time = 1.61, size = 98, normalized size = 1.05

$$\frac{\left(6 \tan^{-1}\left(\sqrt{\sec(e+fx)}\right) - 3 \log\left(1 - \sqrt{\sec(e+fx)}\right) + 3 \log\left(1 + \sqrt{\sec(e+fx)}\right) - \frac{4 \csc^2(e+fx)}{\sec^{\frac{3}{2}}(e+fx)}\right) \sqrt{\sec(e+fx)}}{8b^2 f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3/(b*Sec[e + f*x])^(5/2),x]

[Out] ((6*ArcTan[Sqrt[Sec[e + f*x]]] - 3*Log[1 - Sqrt[Sec[e + f*x]]] + 3*Log[1 + Sqrt[Sec[e + f*x]]] - (4*Csc[e + f*x]^2)/Sec[e + f*x]^(3/2))*Sqrt[Sec[e + f*x]])/(8*b^2*f*Sqrt[b*Sec[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 436 vs. $2(73) = 146$.

time = 0.22, size = 437, normalized size = 4.70

method	result
default	$\frac{(-1+\cos(fx+e)) \left(8(\cos^2(fx+e)) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 16 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} - 4(\cos^2(fx+e)) \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} \right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3/(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)

[Out] -1/8/f*(-1+cos(f*x+e))*(8*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+16*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)-4*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)+3*cos(f*x+e)^2*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))+3*cos(f*x+e)^2*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2)+8*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(3/2)+4*cos(f*x+e)*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-3*arctan(1/2/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2))-3*ln(-(2*cos(f*x+e)^2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-cos(f*x+e)^2+2*cos(f*x+e)-2*(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)-1)/sin(f*x+e)^2))/cos(f*x+e)^2/sin(f*x+e)^4/(b/cos(f*x+e))^(5/2)/(-cos(f*x+e)/(cos(f*x+e)+1)^2)^(1/2)

Maxima [A]

time = 0.48, size = 111, normalized size = 1.19

$$b \left(\frac{4 \sqrt{\frac{b}{\cos(fx+e)}}}{b^4 - \frac{b^4}{\cos(fx+e)^2}} + \frac{6 \arctan\left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right)}{b^{\frac{7}{2}}} - \frac{3 \log\left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}}\right)}{b^{\frac{7}{2}}} \right) \frac{1}{8f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")

[Out] 1/8*b*(4*sqrt(b/cos(f*x + e))/(b^4 - b^4/cos(f*x + e)^2) + 6*arctan(sqrt(b/cos(f*x + e))/sqrt(b))/b^(7/2) - 3*log(-(sqrt(b) - sqrt(b/cos(f*x + e)))/(sqrt(b) + sqrt(b/cos(f*x + e))))/b^(7/2))/f

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 192 vs. 2(77) = 154.

time = 0.47, size = 398, normalized size = 4.28

$$\frac{4(\cos(fx+e)^2-1)\sqrt{-b}\arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) - 8\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)^2 + 3(\cos(fx+e)^2-1)\sqrt{-b}\log\left(\frac{b\cos(fx+e)^2 + 4(\cos(fx+e)^2 - \cos(fx+e))\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}} - 6b\cos(fx+e) + b}{\cos(fx+e)^2 + 2\cos(fx+e) + 1}\right)}{8(b^3f\cos(fx+e)^2 - b^3f)} - \frac{4(\cos(fx+e)^2-1)\sqrt{-b}\arctan\left(\frac{\sqrt{-b}\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}}\right) - 8\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)^2 - 3(\cos(fx+e)^2-1)\sqrt{-b}\log\left(\frac{b\cos(fx+e)^2 + 4(\cos(fx+e)^2 + \cos(fx+e))\sqrt{b}\sqrt{\frac{b}{\cos(fx+e)}} + 6b\cos(fx+e) + b}{\cos(fx+e)^2 - 2\cos(fx+e) + 1}\right)}{8(b^3f\cos(fx+e)^2 - b^3f)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")

[Out] [-1/16*(6*(cos(f*x + e)^2 - 1)*sqrt(-b)*arctan(1/2*sqrt(-b)*sqrt(b/cos(f*x + e))*(cos(f*x + e) + 1)/b) - 8*sqrt(b/cos(f*x + e))*cos(f*x + e)^2 + 3*(cos(f*x + e)^2 - 1)*sqrt(-b)*log((b*cos(f*x + e)^2 + 4*(cos(f*x + e)^2 - cos(f*x + e))*sqrt(-b)*sqrt(b/cos(f*x + e)) - 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 + 2*cos(f*x + e) + 1)))/(b^3*f*cos(f*x + e)^2 - b^3*f), -1/16*(6*(cos(f*x + e)^2 - 1)*sqrt(b)*arctan(1/2*sqrt(b/cos(f*x + e))*(cos(f*x + e) - 1)/sqrt(b)) - 8*sqrt(b/cos(f*x + e))*cos(f*x + e)^2 - 3*(cos(f*x + e)^2 - 1)*sqrt(b)*log((b*cos(f*x + e)^2 + 4*(cos(f*x + e)^2 + cos(f*x + e))*sqrt(b)*sqrt(b/cos(f*x + e)) + 6*b*cos(f*x + e) + b)/(cos(f*x + e)^2 - 2*cos(f*x + e) + 1)))/(b^3*f*cos(f*x + e)^2 - b^3*f)]

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(e + fx)}{(b \sec(e + fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**3/(b*sec(f*x+e))**(5/2),x)`

[Out] `Integral(csc(e + f*x)**3/(b*sec(e + f*x))**(5/2), x)`

Giac [A]

time = 4.38, size = 107, normalized size = 1.15

$$\frac{\frac{2\sqrt{b\cos(fx+e)}b\cos(fx+e)}{b^2\cos(fx+e)^2-b^2} - \frac{3\arctan\left(\frac{\sqrt{b\cos(fx+e)}}{\sqrt{-b}}\right)}{\sqrt{-b}} - \frac{3\arctan\left(\frac{\sqrt{b\cos(fx+e)}}{\sqrt{b}}\right)}{\sqrt{b}}}{4b^2f\operatorname{sgn}(\cos(fx+e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^3/(b*sec(f*x+e))^(5/2),x, algorithm="giac")`

[Out] `1/4*(2*sqrt(b*cos(f*x + e))*b*cos(f*x + e)/(b^2*cos(f*x + e)^2 - b^2) - 3*arctan(sqrt(b*cos(f*x + e))/sqrt(-b))/sqrt(-b) - 3*arctan(sqrt(b*cos(f*x + e))/sqrt(b))/sqrt(b))/(b^2*f*sgn(cos(f*x + e)))`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e+fx)^3 \left(\frac{b}{\cos(e+fx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^3*(b/cos(e + f*x))^(5/2)),x)`

[Out] `int(1/(sin(e + f*x)^3*(b/cos(e + f*x))^(5/2)), x)`

$$3.443 \quad \int \frac{\csc^5(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=123

$$\frac{3 \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{5/2}f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{5/2}f} - \frac{\cot^2(e+fx)\sqrt{b \sec(e+fx)}}{16b^3f} - \frac{\cot^4(e+fx)\sqrt{b \sec(e+fx)}}{4b^3f}$$

[Out] 3/32*arctan((b*sec(f*x+e))^(1/2)/b^(1/2))/b^(5/2)/f+3/32*arctanh((b*sec(f*x+e))^(1/2)/b^(1/2))/b^(5/2)/f-1/16*cot(f*x+e)^2*(b*sec(f*x+e))^(1/2)/b^3/f-1/4*cot(f*x+e)^4*(b*sec(f*x+e))^(1/2)/b^3/f

Rubi [A]

time = 0.06, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {2702, 294, 296, 335, 218, 212, 209}

$$\frac{3 \text{ArcTan}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{5/2}f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{5/2}f} - \frac{\cot^4(e+fx)\sqrt{b \sec(e+fx)}}{4b^3f} - \frac{\cot^2(e+fx)\sqrt{b \sec(e+fx)}}{16b^3f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^5/(b*Sec[e + f*x])^(5/2),x]

[Out] (3*ArcTan[Sqrt[b*Sec[e + f*x]]/Sqrt[b]]/(32*b^(5/2)*f) + (3*ArcTanh[Sqrt[b*Sec[e + f*x]]/Sqrt[b]]/(32*b^(5/2)*f) - (Cot[e + f*x]^2*Sqrt[b*Sec[e + f*x]]/(16*b^3*f) - (Cot[e + f*x]^4*Sqrt[b*Sec[e + f*x]]/(4*b^3*f)

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !I
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 296

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(-(
c*x)^(m + 1))*((a + b*x^n)^(p + 1)/(a*c*n*(p + 1))), x] + Dist[(m + n*(p +
1) + 1)/(a*n*(p + 1)), Int[(c*x)^m*(a + b*x^n)^(p + 1), x], x] /; FreeQ[{a,
b, c, m}, x] && IGtQ[n, 0] && LtQ[p, -1] && IntBinomialQ[a, b, c, n, m, p,
x]
```

Rule 335

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := With[{k =
Denominator[m]}, Dist[k/c, Subst[Int[x^(k*(m + 1) - 1)*(a + b*(x^(k*n)/c^n
))]^p, x], x, (c*x)^(1/k)], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && F
ractionQ[m] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2702

```
Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_S
ymbol] := Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2
], x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)
/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])
```

Rubi steps

$$\begin{aligned}
\int \frac{\csc^5(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= \frac{\text{Subst}\left(\int \frac{x^{3/2}}{(-1+\frac{x^2}{b^2})^3} dx, x, b \sec(e+fx)\right)}{b^5 f} \\
&= -\frac{\cot^4(e+fx) \sqrt{b \sec(e+fx)}}{4b^3 f} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{x} (-1+\frac{x^2}{b^2})^2} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{\cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16b^3 f} - \frac{\cot^4(e+fx) \sqrt{b \sec(e+fx)}}{4b^3 f} - \frac{3 \text{Subst}\left(\int \frac{1}{\sqrt{x}} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{\cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16b^3 f} - \frac{\cot^4(e+fx) \sqrt{b \sec(e+fx)}}{4b^3 f} - \frac{3 \text{Subst}\left(\int \frac{1}{-1+x} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= -\frac{\cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16b^3 f} - \frac{\cot^4(e+fx) \sqrt{b \sec(e+fx)}}{4b^3 f} + \frac{3 \text{Subst}\left(\int \frac{1}{b-x^2} dx, x, b \sec(e+fx)\right)}{8b^3 f} \\
&= \frac{3 \tan^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{5/2} f} + \frac{3 \tanh^{-1}\left(\frac{\sqrt{b \sec(e+fx)}}{\sqrt{b}}\right)}{32b^{5/2} f} - \frac{\cot^2(e+fx) \sqrt{b \sec(e+fx)}}{16b^3 f}
\end{aligned}$$

Mathematica [A]

time = 1.89, size = 110, normalized size = 0.89

$$\frac{\left(6 \tan^{-1}\left(\sqrt{\sec(e+fx)}\right) - 3 \log\left(1 - \sqrt{\sec(e+fx)}\right) + 3 \log\left(1 + \sqrt{\sec(e+fx)}\right) - \frac{2(5+3 \cos(2(e+fx))) \csc^4(e+fx)}{\sec^{\frac{3}{2}}(e+fx)}\right) \sqrt{\sec(e+fx)}}{64b^2 f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^5/(b*Sec[e + f*x])^(5/2), x]`

```
[Out] ((6*ArcTan[Sqrt[Sec[e + f*x]]] - 3*Log[1 - Sqrt[Sec[e + f*x]]] + 3*Log[1 + Sqrt[Sec[e + f*x]]] - (2*(5 + 3*Cos[2*(e + f*x)])*Csc[e + f*x]^4)/Sec[e + f*x]^(3/2))*Sqrt[Sec[e + f*x]])/(64*b^2*f*Sqrt[b*Sec[e + f*x]])
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 736 vs. 2(99) = 198.

time = 0.24, size = 737, normalized size = 5.99

method	result
--------	--------

default	$24 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} (\cos^3(fx+e)+40(\cos^2(fx+e)) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} -12(\cos^3(fx+e)) \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} -3 \ln \left(-\frac{2(\cos(fx+e)+1)^2}{\cos(fx+e)+1} \right)$
---------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^5/(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{64f} \left(24 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} \cos^3(fx+e) + 40 \cos^2(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} - 12 \cos^3(fx+e) \sqrt{-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2}} - 3 \ln \left(-\frac{2(\cos(fx+e)+1)^2}{\cos(fx+e)+1} \right) \right. \\ \left. - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - 1 \right) / \sin(fx+e)^2 * \cos(fx+e)^3 - 3 \arctan \left(\frac{1/2}{-\cos(fx+e)/(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} * \cos(fx+e)^3 + 8 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} + 24 \cos(fx+e)^2 * \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} + 3 \cos(fx+e)^2 \ln \left(-\frac{2 \cos(fx+e)^2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - 1 \right) / \sin(fx+e)^2 + 3 \cos(fx+e)^2 \arctan \left(\frac{1/2}{-\cos(fx+e)/(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - 8 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{3}{2}} - 12 \cos(fx+e) \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} + 3 \ln \left(-\frac{2 \cos(fx+e)^2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - 1 \right) / \sin(fx+e)^2 * \cos(fx+e) + 3 \arctan \left(\frac{1/2}{-\cos(fx+e)/(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} * \cos(fx+e) - 3 \ln \left(-\frac{2 \cos(fx+e)^2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - \cos(fx+e)^2 + 2 \cos(fx+e) - 2 \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} - 1 \right) / \sin(fx+e)^2 - 3 \arctan \left(\frac{1/2}{-\cos(fx+e)/(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}} \right) / \sin(fx+e)^4 / (b/\cos(fx+e))^{5/2} / \cos(fx+e)^2 / \left(-\frac{\cos(fx+e)}{(\cos(fx+e)+1)^2} \right)^{\frac{1}{2}}$$

Maxima [A]

time = 0.54, size = 143, normalized size = 1.16

$$\frac{b \left(\frac{4 \left(3b^2 \sqrt{\frac{b}{\cos(fx+e)}} + \left(\frac{b}{\cos(fx+e)} \right)^{\frac{3}{2}} \right)}{b^6 - \frac{2b^6}{\cos(fx+e)^2} + \frac{b^6}{\cos(fx+e)^4}} - \frac{6 \arctan \left(\frac{\sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b}} \right)}{b^{\frac{7}{2}}} + \frac{3 \log \left(\frac{\sqrt{b} - \sqrt{\frac{b}{\cos(fx+e)}}}{\sqrt{b} + \sqrt{\frac{b}{\cos(fx+e)}}} \right)}{b^{\frac{7}{2}}} \right)}{64f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out]
$$-1/64*b*(4*(3*b^2*\sqrt{b/\cos(f*x + e)} + (b/\cos(f*x + e))^{5/2})/(b^6 - 2*b^6/\cos(f*x + e)^2 + b^6/\cos(f*x + e)^4) - 6*\arctan(\sqrt{b/\cos(f*x + e)})/\sqrt{t(b)}/b^{7/2} + 3*\log(-(\sqrt{b} - \sqrt{b/\cos(f*x + e)}))/(\sqrt{b} + \sqrt{b/\cos(f*x + e)}))/b^{7/2})/f$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 240 vs. $2(105) = 210$.

time = 0.44, size = 494, normalized size = 4.02

$$\frac{\frac{6(\cos(x+e)^2-2\cos(x+e)+1)\sqrt{-b}\arctan\left(\frac{\sqrt{b}\cos(x+e)}{\sqrt{-b}}\right)+3(\cos(x+e)^2-2\cos(x+e)+1)\sqrt{-b}\log\left(\frac{b\cos(x+e)^2+4(\cos(x+e)^2-\cos(x+e))\sqrt{-b}}{(\cos(x+e)^2+2\cos(x+e)+1)}\right)+8(3\cos(x+e)^4+\cos(x+e)^2)\sqrt{b/\cos(x+e)}}{128(b^2\cos(x+e)^2-2b^2\cos(x+e)+b^2)}}{\frac{6(\cos(x+e)^2-2\cos(x+e)+1)\sqrt{-b}\arctan\left(\frac{\sqrt{b}\cos(x+e)}{\sqrt{-b}}\right)+3(\cos(x+e)^2-2\cos(x+e)+1)\sqrt{-b}\log\left(\frac{b\cos(x+e)^2+4(\cos(x+e)^2-\cos(x+e))\sqrt{-b}}{(\cos(x+e)^2+2\cos(x+e)+1)}\right)+8(3\cos(x+e)^4+\cos(x+e)^2)\sqrt{b/\cos(x+e)}}{128(b^2\cos(x+e)^2-2b^2\cos(x+e)+b^2)}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & [-1/128*(6*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{-b}*\arctan(1/2*\sqrt{b/\cos(f*x + e)}*(\cos(f*x + e) + 1)/b) + 3*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{-b}*\log((b*\cos(f*x + e)^2 + 4*(\cos(f*x + e)^2 - \cos(f*x + e))*\sqrt{-b}*\sqrt{b/\cos(f*x + e)} - 6*b*\cos(f*x + e) + b)/(\cos(f*x + e)^2 + 2*\cos(f*x + e) + 1)) + 8*(3*\cos(f*x + e)^4 + \cos(f*x + e)^2)*\sqrt{b/\cos(f*x + e)})/(b^3*f*\cos(f*x + e)^4 - 2*b^3*f*\cos(f*x + e)^2 + b^3*f), \\ & -1/128*(6*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{b}*\arctan(1/2*\sqrt{b/\cos(f*x + e)}*(\cos(f*x + e) - 1)/\sqrt{b}) - 3*(\cos(f*x + e)^4 - 2*\cos(f*x + e)^2 + 1)*\sqrt{b}*\log((b*\cos(f*x + e)^2 + 4*(\cos(f*x + e)^2 + \cos(f*x + e))*\sqrt{b}*\sqrt{b/\cos(f*x + e)} + 6*b*\cos(f*x + e) + b)/(\cos(f*x + e)^2 - 2*\cos(f*x + e) + 1)) + 8*(3*\cos(f*x + e)^4 + \cos(f*x + e)^2)*\sqrt{b/\cos(f*x + e)})/(b^3*f*\cos(f*x + e)^4 - 2*b^3*f*\cos(f*x + e)^2 + b^3*f)] \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^5(e + fx)}{(b \sec(e + fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**5/(b*sec(f*x+e))**(5/2),x)`

[Out] `Integral(csc(e + f*x)**5/(b*sec(e + f*x))**(5/2), x)`

Giac [A]

time = 3.78, size = 139, normalized size = 1.13

$$\frac{\frac{3 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{-b}}\right)}{\sqrt{-b} b^2} + \frac{3 \arctan\left(\frac{\sqrt{b \cos(fx + e)}}{\sqrt{b}}\right)}{b^{\frac{5}{2}}} + \frac{2 \left(3 \sqrt{b \cos(fx + e)} b^3 \cos(fx + e)^3 + \sqrt{b \cos(fx + e)} b^3 \cos(fx + e)\right)}{(b^2 \cos(fx + e)^2 - b^2)^2 b^2}}{32 f \operatorname{sgn}(\cos(fx + e))}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5/(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out]
$$-1/32*(3*\arctan(\sqrt{b*\cos(f*x + e)})/\sqrt{-b})/(\sqrt{-b}*b^2) + 3*\arctan(\sqrt{b*\cos(f*x + e)})/\sqrt{b})/b^{5/2} + 2*(3*\sqrt{b*\cos(f*x + e)}*b^3*\cos(f*x + e)^3 + \sqrt{b*\cos(f*x + e)}*b^3*\cos(f*x + e))/((b^2*\cos(f*x + e)^2 - b^2)^{2*b^2})/(f*\operatorname{sgn}(\cos(f*x + e)))$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^5 \left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^5*(b/cos(e + f*x))^(5/2)),x)

[Out] int(1/(sin(e + f*x)^5*(b/cos(e + f*x))^(5/2)), x)

$$3.444 \quad \int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=126

$$\frac{8E\left(\frac{1}{2}(e+fx) \mid 2\right)}{65b^2f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}} - \frac{4b\sin(e+fx)}{39f(b\sec(e+fx))^{7/2}} + \frac{8\sin(e+fx)}{195bf(b\sec(e+fx))^{3/2}} - \frac{2b\sin^3(e+fx)}{13f(b\sec(e+fx))^{7/2}}$$

[Out] $-4/39*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(7/2)}+8/195*\sin(f*x+e)/b/f/(b*\sec(f*x+e))^{(3/2)}-2/13*b*\sin(f*x+e)^3/f/(b*\sec(f*x+e))^{(7/2)}+8/65*(\cos(1/2*f*x+1/2*e))^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/b^2/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.09, antiderivative size = 126, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2707, 3854, 3856, 2719}

$$\frac{8E\left(\frac{1}{2}(e+fx) \mid 2\right)}{65b^2f\sqrt{\cos(e+fx)}\sqrt{b\sec(e+fx)}} - \frac{2b\sin^3(e+fx)}{13f(b\sec(e+fx))^{7/2}} + \frac{8\sin(e+fx)}{195bf(b\sec(e+fx))^{3/2}} - \frac{4b\sin(e+fx)}{39f(b\sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^4/(b*Sec[e + f*x])^(5/2),x]

[Out] $(8*\text{EllipticE}[(e+f*x)/2,2])/(65*b^2*f*\text{Sqrt}[\text{Cos}[e+f*x]]*\text{Sqrt}[b*\text{Sec}[e+f*x]]) - (4*b*\text{Sin}[e+f*x])/(39*f*(b*\text{Sec}[e+f*x])^{(7/2)}) + (8*\text{Sin}[e+f*x])/(195*b*f*(b*\text{Sec}[e+f*x])^{(3/2)}) - (2*b*\text{Sin}[e+f*x]^3)/(13*f*(b*\text{Sec}[e+f*x])^{(7/2)})$

Rule 2707

Int[(csc[(e.) + (f.)*(x_)]*(a.))^(m_)*((b.)*sec[(e.) + (f.)*(x_)]^(n_)), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]

Rule 2719

Int[Sqrt[sin[(c.) + (d.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c.) + (d.)*(x_)]*(b.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= -\frac{2b \sin^3(e+fx)}{13f(b \sec(e+fx))^{7/2}} + \frac{6}{13} \int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx \\
 &= -\frac{4b \sin(e+fx)}{39f(b \sec(e+fx))^{7/2}} - \frac{2b \sin^3(e+fx)}{13f(b \sec(e+fx))^{7/2}} + \frac{4}{39} \int \frac{1}{(b \sec(e+fx))^{5/2}} dx \\
 &= -\frac{4b \sin(e+fx)}{39f(b \sec(e+fx))^{7/2}} + \frac{8 \sin(e+fx)}{195bf(b \sec(e+fx))^{3/2}} - \frac{2b \sin^3(e+fx)}{13f(b \sec(e+fx))^{7/2}} + \frac{4}{65} \int \frac{1}{(b \sec(e+fx))^{3/2}} dx \\
 &= -\frac{4b \sin(e+fx)}{39f(b \sec(e+fx))^{7/2}} + \frac{8 \sin(e+fx)}{195bf(b \sec(e+fx))^{3/2}} - \frac{2b \sin^3(e+fx)}{13f(b \sec(e+fx))^{7/2}} + \frac{8 \sin(e+fx)}{195bf(b \sec(e+fx))^{3/2}} \\
 &= \frac{8E\left(\frac{1}{2}(e+fx) \mid 2\right)}{65b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{4b \sin(e+fx)}{39f(b \sec(e+fx))^{7/2}} + \frac{8 \sin(e+fx)}{195bf(b \sec(e+fx))^{3/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.36, size = 83, normalized size = 0.66

$$\frac{192E\left(\frac{1}{2}(e+fx) \mid 2\right) + \cos^{\frac{3}{2}}(e+fx)(-6 \sin(e+fx) - 55 \sin(3(e+fx)) + 15 \sin(5(e+fx)))}{1560f \cos^{\frac{5}{2}}(e+fx)(b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^4/(b*Sec[e + f*x])^(5/2), x]

[Out] (192*EllipticE[(e + f*x)/2, 2] + Cos[e + f*x]^(3/2)*(-6*Sin[e + f*x] - 55*Sin[3*(e + f*x)] + 15*Sin[5*(e + f*x)])/(1560*f*Cos[e + f*x]^(5/2)*(b*Sec[e + f*x])^(5/2))

Maple [C] Result contains complex when optimal does not.

time = 0.24, size = 343, normalized size = 2.72

method	result
--------	--------

default	$-\frac{2\left(15(\cos^8(fx+e))-40(\cos^6(fx+e))+12i\operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)},i\right)\sin(fx+e)\cos(fx+e)\sqrt{\frac{1}{\cos(fx+e)+1}}\sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}}\right)}{b^5}$
---------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(f*x+e)^4/(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$-\frac{2}{195} \frac{15 \cos^8(fx+e) - 40 \cos^6(fx+e) + 12 I \cos(fx+e) \sin(fx+e) \operatorname{EllipticE}\left(\frac{I(-1+\cos(fx+e))}{\sin(fx+e)}, I\right) \left(\frac{1}{\cos(fx+e)+1}\right)^{1/2} \left(\frac{\cos(fx+e)}{\cos(fx+e)+1}\right)^{1/2} - 12 I \left(\frac{1}{\cos(fx+e)+1}\right)^{1/2} \left(\frac{\cos(fx+e)}{\cos(fx+e)+1}\right)^{1/2} \operatorname{EllipticF}\left(\frac{I(-1+\cos(fx+e))}{\sin(fx+e)}, I\right) \sin(fx+e) \cos(fx+e) + 12 I \sin(fx+e) \operatorname{EllipticE}\left(\frac{I(-1+\cos(fx+e))}{\sin(fx+e)}, I\right) \left(\frac{1}{\cos(fx+e)+1}\right)^{1/2} \left(\frac{\cos(fx+e)}{\cos(fx+e)+1}\right)^{1/2} - 12 I \operatorname{EllipticF}\left(\frac{I(-1+\cos(fx+e))}{\sin(fx+e)}, I\right) \left(\frac{1}{\cos(fx+e)+1}\right)^{1/2} \left(\frac{\cos(fx+e)}{\cos(fx+e)+1}\right)^{1/2} \sin(fx+e) + 29 \cos^4(fx+e) + 8 \cos^2(fx+e) - 12 \cos(fx+e)}{b^5 \cos^3(fx+e)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out] `integrate(sin(f*x + e)^4/(b*sec(f*x + e))^(5/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 126, normalized size = 1.00

$$\frac{2\left(\left(15\cos^5(fx+e)-25\cos^3(fx+e)+4\cos(fx+e)\right)\sqrt{\frac{b}{\cos(fx+e)}}\sin(fx+e)+6i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4,0,\cos(fx+e)+i\sin(fx+e))-6i\sqrt{2}\sqrt{b}\operatorname{weierstrassZeta}(-4,0,\cos(fx+e)-i\sin(fx+e))\right)}{195b^5f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

[Out]
$$\frac{2}{195} \left((15 \cos^5(fx+e) - 25 \cos^3(fx+e) + 4 \cos(fx+e))^2 \sqrt{b/\cos(fx+e)} \sin(fx+e) + 6 I \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + I \sin(fx+e))) - 6 I \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - I \sin(fx+e))) \right) / (b^3 f)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**4/(b*sec(f*x+e))**(5/2),x)`

[Out] `Integral(sin(e + f*x)**4/(b*sec(e + f*x))**(5/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^4/(b*sec(f*x+e))^(5/2),x, algorithm="giac")`

[Out] `integrate(sin(f*x + e)^4/(b*sec(f*x + e))^(5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^4}{\left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^4/(b/cos(e + f*x))^(5/2),x)`

[Out] `int(sin(e + f*x)^4/(b/cos(e + f*x))^(5/2), x)`

$$3.445 \quad \int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=98

$$\frac{4E\left(\frac{1}{2}(e+fx) \mid 2\right)}{15b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{2b \sin(e+fx)}{9f(b \sec(e+fx))^{7/2}} + \frac{4 \sin(e+fx)}{45bf(b \sec(e+fx))^{3/2}}$$

[Out] $-2/9*b*\sin(f*x+e)/f/(b*\sec(f*x+e))^{(7/2)}+4/45*\sin(f*x+e)/b/f/(b*\sec(f*x+e))^{(3/2)}+4/15*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/b^2/f/\cos(f*x+e)^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2707, 3854, 3856, 2719}

$$\frac{4E\left(\frac{1}{2}(e+fx) \mid 2\right)}{15b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} + \frac{4 \sin(e+fx)}{45bf(b \sec(e+fx))^{3/2}} - \frac{2b \sin(e+fx)}{9f(b \sec(e+fx))^{7/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sin[e + f*x]^2/(b*Sec[e + f*x])^(5/2),x]`

[Out] $(4*\text{EllipticE}[(e+f*x)/2, 2])/(15*b^2*f*\text{Sqrt}[\text{Cos}[e+f*x]]*\text{Sqrt}[b*\text{Sec}[e+f*x]]) - (2*b*\text{Sin}[e+f*x])/(9*f*(b*\text{Sec}[e+f*x])^{(7/2)}) + (4*\text{Sin}[e+f*x])/(45*b*f*(b*\text{Sec}[e+f*x])^{(3/2)})$

Rule 2707

`Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] := Simp[b*(a*Csc[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + n))), x] + Dist[(m + 1)/(a^2*(m + n)), Int[(a*Csc[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && NeQ[m + n, 0] && IntegersQ[2*m, 2*n]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3854

`Int[(csc[(c_.) + (d_.)*(x_.)]*(b_.))^(n_.), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= -\frac{2b \sin(e+fx)}{9f(b \sec(e+fx))^{7/2}} + \frac{2}{9} \int \frac{1}{(b \sec(e+fx))^{5/2}} dx \\
 &= -\frac{2b \sin(e+fx)}{9f(b \sec(e+fx))^{7/2}} + \frac{4 \sin(e+fx)}{45bf(b \sec(e+fx))^{3/2}} + \frac{2 \int \frac{1}{\sqrt{b \sec(e+fx)}} dx}{15b^2} \\
 &= -\frac{2b \sin(e+fx)}{9f(b \sec(e+fx))^{7/2}} + \frac{4 \sin(e+fx)}{45bf(b \sec(e+fx))^{3/2}} + \frac{2 \int \sqrt{\cos(e+fx)} dx}{15b^2 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
 &= \frac{4E\left(\frac{1}{2}(e+fx)|2\right)}{15b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{2b \sin(e+fx)}{9f(b \sec(e+fx))^{7/2}} + \frac{4 \sin(e+fx)}{45bf(b \sec(e+fx))^{3/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.29, size = 66, normalized size = 0.67

$$\frac{\frac{96E\left(\frac{1}{2}(e+fx)|2\right)}{\sqrt{\cos(e+fx)}} - 4 \sin(2(e+fx)) - 10 \sin(4(e+fx))}{360b^2 f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^2/(b*Sec[e + f*x])^(5/2), x]

[Out] ((96*EllipticE[(e + f*x)/2, 2])/Sqrt[Cos[e + f*x]] - 4*Sin[2*(e + f*x)] - 10*Sin[4*(e + f*x)])/(360*b^2*f*Sqrt[b*Sec[e + f*x]])

Maple [C] Result contains complex when optimal does not.

time = 0.21, size = 333, normalized size = 3.40

method	result
default	$ \frac{2(\cos^6(fx+e))}{9} + \frac{4i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e)}{15} - \frac{4i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}\right)}{15} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^2/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)

[Out] $2/45/f*(5*\cos(f*x+e)^6+6*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*EllipticF(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*\sin(f*x+e)*\cos(f*x+e)-6*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\cos(f*x+e)*\sin(f*x+e)*EllipticE(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)+6*I*EllipticF(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\sin(f*x+e)-6*I*(1/(\cos(f*x+e)+1))^{1/2}*(\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2}*\sin(f*x+e)*EllipticE(I*(-1+\cos(f*x+e))/\sin(f*x+e),I)-7*\cos(f*x+e)^4-4*\cos(f*x+e)^2+6*\cos(f*x+e))/\cos(f*x+e)^3/\sin(f*x+e)/(b/\cos(f*x+e))^{5/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out] `integrate(sin(f*x + e)^2/(b*sec(f*x + e))^(5/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 115, normalized size = 1.17

$$\frac{2 \left((5 \cos(fx+e)^4 - 2 \cos(fx+e)^2) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e) - 3i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e))) + 3i \sqrt{2} \sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e))) \right)}{45 b^3 f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

[Out] $-2/45*((5*\cos(f*x + e)^4 - 2*\cos(f*x + e)^2)*\sqrt{b/\cos(f*x + e)}*\sin(f*x + e) - 3*I*\sqrt{2}*\sqrt{b}*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e))) + 3*I*\sqrt{2}*\sqrt{b}*\operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e))))/(b^3*f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^2(e + fx)}{(b \sec(e + fx))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**2/(b*sec(f*x+e))**(5/2),x)`

[Out] `Integral(sin(e + f*x)**2/(b*sec(e + f*x))**(5/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(b*sec(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^2/(b*sec(f*x + e))^(5/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^2}{\left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2/(b/cos(e + f*x))^(5/2),x)

[Out] int(sin(e + f*x)^2/(b/cos(e + f*x))^(5/2), x)

$$3.446 \quad \int \frac{1}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{6E\left(\frac{1}{2}(e+fx) \mid 2\right)}{5b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} + \frac{2 \sin(e+fx)}{5bf(b \sec(e+fx))^{3/2}}$$

[Out] 2/5*sin(f*x+e)/b/f/(b*sec(f*x+e))^(3/2)+6/5*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticE(sin(1/2*f*x+1/2*e),2^(1/2))/b^2/f/cos(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3854, 3856, 2719}

$$\frac{6E\left(\frac{1}{2}(e+fx) \mid 2\right)}{5b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} + \frac{2 \sin(e+fx)}{5bf(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^(-5/2),x]

[Out] (6*EllipticE[(e + f*x)/2, 2])/(5*b^2*f*Sqrt[Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) + (2*Sin[e + f*x])/(5*b*f*(b*Sec[e + f*x])^(3/2))

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(b \sec(e + fx))^{5/2}} dx &= \frac{2 \sin(e + fx)}{5bf(b \sec(e + fx))^{3/2}} + \frac{3 \int \frac{1}{\sqrt{b \sec(e + fx)}} dx}{5b^2} \\
&= \frac{2 \sin(e + fx)}{5bf(b \sec(e + fx))^{3/2}} + \frac{3 \int \sqrt{\cos(e + fx)} dx}{5b^2 \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
&= \frac{6E\left(\frac{1}{2}(e + fx) \mid 2\right)}{5b^2 f \sqrt{\cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{2 \sin(e + fx)}{5bf(b \sec(e + fx))^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 60, normalized size = 0.83

$$\frac{\sqrt{b \sec(e + fx)} \left(12 \sqrt{\cos(e + fx)} E\left(\frac{1}{2}(e + fx) \mid 2\right) + \sin(e + fx) + \sin(3(e + fx)) \right)}{10b^3 f}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^(-5/2),x]**[Out]** (Sqrt[b*Sec[e + f*x]]*(12*Sqrt[Cos[e + f*x]]*EllipticE[(e + f*x)/2, 2] + Sin[e + f*x] + Sin[3*(e + f*x)]))/(10*b^3*f)**Maple [C]** Result contains complex when optimal does not.

time = 0.25, size = 321, normalized size = 4.46

method	result
default	$ -\frac{2 \left(3i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} - 3i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \right)}{\dots} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)

[Out]
$$-\frac{2}{5} \frac{1}{f} \left(3I \left(\frac{1}{(\cos(f*x+e)+1)} \right)^{1/2} \left(\frac{\cos(f*x+e)}{(\cos(f*x+e)+1)} \right)^{1/2} \operatorname{EllipticE}\left(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I\right) \sin(f*x+e) \cos(f*x+e) - 3I \left(\frac{1}{(\cos(f*x+e)+1)} \right)^{1/2} \left(\frac{\cos(f*x+e)}{(\cos(f*x+e)+1)} \right)^{1/2} \operatorname{EllipticF}\left(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I\right) \sin(f*x+e) \cos(f*x+e) + 3I \left(\frac{1}{(\cos(f*x+e)+1)} \right)^{1/2} \left(\frac{\cos(f*x+e)}{(\cos(f*x+e)+1)} \right)^{1/2} \operatorname{EllipticE}\left(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I\right) \sin(f*x+e) \cos(f*x+e) - 3I \operatorname{EllipticF}\left(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I\right) \left(\frac{1}{(\cos(f*x+e)+1)} \right)^{1/2} \left(\frac{\cos(f*x+e)}{(\cos(f*x+e)+1)} \right)^{1/2} \sin(f*x+e) + \cos(f*x+e)^4 + 2 \cos(f*x+e)^2 - 3 \cos(f*x+e) \right) / (b/\cos(f*x+e))^{5/2} / \cos(f*x+e)^3 / \sin(f*x+e)$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")``[Out] integrate((b*sec(f*x + e))^(-5/2), x)`**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.15, size = 102, normalized size = 1.42

$$2 \sqrt{\frac{b}{\cos(fx+e)} \cos(fx+e)^2 \sin(fx+e) + 3i\sqrt{2}\sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) + i \sin(fx+e))) - 3i\sqrt{2}\sqrt{b} \operatorname{weierstrassZeta}(-4, 0, \operatorname{weierstrassPInverse}(-4, 0, \cos(fx+e) - i \sin(fx+e)))}$$

$$5b^3f$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

```
[Out] 1/5*(2*sqrt(b/cos(f*x + e))*cos(f*x + e)^2*sin(f*x + e) + 3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) - 3*I*sqrt(2)*sqrt(b)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e)))/(b^3*f)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(b \sec(e + fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(b*sec(f*x+e))**(5/2),x)``[Out] Integral((b*sec(e + f*x))**(-5/2), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(b*sec(f*x+e))^(5/2),x, algorithm="giac")``[Out] integrate((b*sec(f*x + e))^(-5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{b}{\cos(e+fx)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b/cos(e + f*x))^(5/2),x)

[Out] int(1/(b/cos(e + f*x))^(5/2), x)

$$3.447 \quad \int \frac{\csc^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=68

$$-\frac{\csc(e+fx)}{bf(b \sec(e+fx))^{3/2}} - \frac{3E\left(\frac{1}{2}(e+fx) \mid 2\right)}{b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

[Out] $-\csc(f*x+e)/b/f/(b*\sec(f*x+e))^{(3/2)}-3*(\cos(1/2*f*x+1/2*e)^2)^{(1/2)}/\cos(1/2*f*x+1/2*e)*\text{EllipticE}(\sin(1/2*f*x+1/2*e),2^{(1/2)})/b^2/f/\cos(f*x+e)^{(1/2)/(b*\sec(f*x+e))^{(1/2)}}$

Rubi [A]

time = 0.05, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {2703, 3856, 2719}

$$-\frac{3E\left(\frac{1}{2}(e+fx) \mid 2\right)}{b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\csc(e+fx)}{bf(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^2/(b*\text{Sec}[e + f*x])^{(5/2)}, x]$

[Out] $-(\text{Csc}[e + f*x]/(b*f*(b*\text{Sec}[e + f*x])^{(3/2)})) - (3*\text{EllipticE}[(e + f*x)/2, 2])/(b^2*f*\text{Sqrt}[\text{Cos}[e + f*x]]*\text{Sqrt}[b*\text{Sec}[e + f*x]])$

Rule 2703

$\text{Int}[(\csc[(e_.) + (f_.)*(x_.)]*(a_.))^{(m_.)}*((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-a)*(a*\text{Csc}[e + f*x])^{(m-1)}*((b*\text{Sec}[e + f*x])^{(n+1)/(f*b*(m-1)}), x] + \text{Dist}[a^2*((n+1)/(b^2*(m-1))], \text{Int}[(a*\text{Csc}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{LtQ}[n, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_.)*(x_.)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^{(n)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned}
\int \frac{\csc^2(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= -\frac{\csc(e+fx)}{bf(b \sec(e+fx))^{3/2}} - \frac{3 \int \frac{1}{\sqrt{b \sec(e+fx)}} dx}{2b^2} \\
&= -\frac{\csc(e+fx)}{bf(b \sec(e+fx))^{3/2}} - \frac{3 \int \sqrt{\cos(e+fx)} dx}{2b^2 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
&= -\frac{\csc(e+fx)}{bf(b \sec(e+fx))^{3/2}} - \frac{3E\left(\frac{1}{2}(e+fx) \mid 2\right)}{b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.13, size = 51, normalized size = 0.75

$$\frac{-\cot(e+fx) - \frac{3E\left(\frac{1}{2}(e+fx) \mid 2\right)}{\sqrt{\cos(e+fx)}}}{b^2 f \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2/(b*Sec[e + f*x])^(5/2), x]``[Out] (-Cot[e + f*x] - (3*EllipticE[(e + f*x)/2, 2])/Sqrt[Cos[e + f*x]])/(b^2*f*Sqrt[b*Sec[e + f*x]])`**Maple [C]** Result contains complex when optimal does not.

time = 0.23, size = 313, normalized size = 4.60

method	result
default	$ -\frac{3i \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} \operatorname{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e) \cos(fx+e) - 3i \operatorname{EllipticE}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sin(fx+e)}{b^2 f \sqrt{b \sec(fx+e)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)`

```
[Out] -1/f*(3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*sin(f*x+e)*cos(f*x+e)-3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e), I)*sin(f*x+e)+3*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e), I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)-3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*sin(f*x+e)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e), I)-2*cos(f*x+e)^2+3*cos(f*x+e))/sin(f*x+e)/(b/cos(f*x+e))^(5/2)/cos(f*x+e)^3
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^2/(b*sec(f*x + e))^(5/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 118, normalized size = 1.74

$$\frac{-3i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))) + 3i\sqrt{2}\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))) - 2\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)^2}{2b^2f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")
```

```
[Out] 1/2*(-3*I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*I*sqrt(2)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) - 2*sqrt(b/cos(f*x + e))*cos(f*x + e)^2/(b^3*f*sin(f*x + e))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(e + fx)}{(b \sec(e + fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**2/(b*sec(f*x+e))**(5/2),x)
```

```
[Out] Integral(csc(e + f*x)**2/(b*sec(e + f*x))**(5/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(b*sec(f*x+e))^(5/2),x, algorithm="giac")
```

```
[Out] integrate(csc(f*x + e)^2/(b*sec(f*x + e))^(5/2), x)
```


Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^2 \left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^2*(b/cos(e + f*x))^(5/2)), x)

[Out] int(1/(sin(e + f*x)^2*(b/cos(e + f*x))^(5/2)), x)

$$3.448 \quad \int \frac{\csc^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=102

$$\frac{\csc(e+fx)}{2bf(b \sec(e+fx))^{3/2}} - \frac{\csc^3(e+fx)}{3bf(b \sec(e+fx))^{3/2}} + \frac{E(\frac{1}{2}(e+fx)|2)}{2b^2f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

[Out] 1/2*csc(f*x+e)/b/f/(b*sec(f*x+e))^(3/2)-1/3*csc(f*x+e)^3/b/f/(b*sec(f*x+e))^(3/2)+1/2*(cos(1/2*f*x+1/2*e)^2)^(1/2)/cos(1/2*f*x+1/2*e)*EllipticE(sin(1/2*f*x+1/2*e),2^(1/2))/b^2/f/cos(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2)

Rubi [A]

time = 0.08, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2703, 2705, 3856, 2719}

$$\frac{E(\frac{1}{2}(e+fx)|2)}{2b^2f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\csc^3(e+fx)}{3bf(b \sec(e+fx))^{3/2}} + \frac{\csc(e+fx)}{2bf(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^4/(b*Sec[e + f*x])^(5/2),x]

[Out] Csc[e + f*x]/(2*b*f*(b*Sec[e + f*x])^(3/2)) - Csc[e + f*x]^3/(3*b*f*(b*Sec[e + f*x])^(3/2)) + EllipticE[(e + f*x)/2, 2]/(2*b^2*f*Sqrt[Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])

Rule 2703

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Dist[a^2*((n + 1)/(b^2*(m - 1))), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_.)]*(a_.))^(m_)*((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

$\text{Int}[(\text{csc}[(c_.) + (d_.)*(x_)]*(b_.)^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n * \text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /;$ $\text{FreeQ}\{b, c, d\}, x\} \&\& \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned} \int \frac{\csc^4(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= -\frac{\csc^3(e+fx)}{3bf(b \sec(e+fx))^{3/2}} - \frac{\int \frac{\csc^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx}{2b^2} \\ &= \frac{\csc(e+fx)}{2bf(b \sec(e+fx))^{3/2}} - \frac{\csc^3(e+fx)}{3bf(b \sec(e+fx))^{3/2}} + \frac{\int \frac{1}{\sqrt{b \sec(e+fx)}} dx}{4b^2} \\ &= \frac{\csc(e+fx)}{2bf(b \sec(e+fx))^{3/2}} - \frac{\csc^3(e+fx)}{3bf(b \sec(e+fx))^{3/2}} + \frac{\int \sqrt{\cos(e+fx)} dx}{4b^2 \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \\ &= \frac{\csc(e+fx)}{2bf(b \sec(e+fx))^{3/2}} - \frac{\csc^3(e+fx)}{3bf(b \sec(e+fx))^{3/2}} + \frac{E(\frac{1}{2}(e+fx)|2)}{2b^2 f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} \end{aligned}$$

Mathematica [A]

time = 0.19, size = 79, normalized size = 0.77

$$\frac{(-3 + 5 \csc^2(e+fx) - 2 \csc^4(e+fx) + 3 \sqrt{\cos(e+fx)} \csc(e+fx) E(\frac{1}{2}(e+fx)|2)) \sqrt{b \sec(e+fx)} \sin(e+fx)}{6b^3 f}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^4/(b*Sec[e + f*x])^(5/2), x]

[Out] ((-3 + 5*Csc[e + f*x]^2 - 2*Csc[e + f*x]^4 + 3*Sqrt[Cos[e + f*x]]*Csc[e + f*x]*EllipticE[(e + f*x)/2, 2])*Sqrt[b*Sec[e + f*x]]*Sin[e + f*x])/(6*b^3*f)

Maple [C] Result contains complex when optimal does not.

time = 0.25, size = 623, normalized size = 6.11

method	result
default	$-\frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(3i \text{EllipticF}\left(\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}, i\right) \sqrt{\frac{1}{\cos(fx+e)+1}} \sqrt{\frac{\cos(fx+e)}{\cos(fx+e)+1}} (\cos^3(fx+e) \sin(fx+e)) \right)}{6b^3 f}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^4/(b*sec(f*x+e))^(5/2), x, method=_RETURNVERBOSE)

```
[Out] -1/6/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(3*I*EllipticF(I*(-1+cos(f*x+e)))/
sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*co
s(f*x+e)^3*sin(f*x+e)-3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+
1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)^3
+3*I*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(co
s(f*x+e)/(cos(f*x+e)+1))^(1/2)*cos(f*x+e)^2*sin(f*x+e)-3*I*(1/(cos(f*x+e)+1
))^^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(
f*x+e),I)*sin(f*x+e)*cos(f*x+e)^2-3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/
(cos(f*x+e)+1))^(1/2)*EllipticF(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*
cos(f*x+e)+3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(f*x+e)+1))^(1/2)*E
llipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)*cos(f*x+e)-3*I*Elliptic
F(I*(-1+cos(f*x+e))/sin(f*x+e),I)*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos
(f*x+e)+1))^(1/2)*sin(f*x+e)+3*I*(1/(cos(f*x+e)+1))^(1/2)*(cos(f*x+e)/(cos(
f*x+e)+1))^(1/2)*EllipticE(I*(-1+cos(f*x+e))/sin(f*x+e),I)*sin(f*x+e)+3*cos
(f*x+e)^3+2*cos(f*x+e)^2-3*cos(f*x+e))/cos(f*x+e)^3/sin(f*x+e)^7/(b/cos(f*x
+e))^(5/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^4/(b*sec(f*x + e))^(5/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 175, normalized size = 1.72

$$\frac{3\sqrt{2}(-i\cos(fx+e)^2+i)\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e))) + 3\sqrt{2}(i\cos(fx+e)^2-i)\sqrt{b}\sin(fx+e)\operatorname{weierstrassZeta}(-4,0,\operatorname{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e))) - 2(3\cos(fx+e)^4 - \cos(fx+e)^2)\sqrt{\frac{b}{\cos(fx+e)}}}{12(b^2f\cos(fx+e)^2 - b^2f)\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")
```

```
[Out] -1/12*(3*sqrt(2)*(-I*cos(f*x + e)^2 + I)*sqrt(b)*sin(f*x + e)*weierstrassZe
ta(-4, 0, weierstrassPInverse(-4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*sq
rt(2)*(I*cos(f*x + e)^2 - I)*sqrt(b)*sin(f*x + e)*weierstrassZeta(-4, 0, we
ierstrassPInverse(-4, 0, cos(f*x + e) - I*sin(f*x + e))) - 2*(3*cos(f*x + e
)^4 - cos(f*x + e)^2)*sqrt(b/cos(f*x + e)))/((b^3*f*cos(f*x + e)^2 - b^3*f
*sin(f*x + e))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^4(e + fx)}{(b \sec(e + fx))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**4/(b*sec(f*x+e))**(5/2),x)`

[Out] `Integral(csc(e + f*x)**4/(b*sec(e + f*x))**(5/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4/(b*sec(f*x+e))^(5/2),x, algorithm="giac")`

[Out] `integrate(csc(f*x + e)^4/(b*sec(f*x + e))^(5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^4 \left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^4*(b/cos(e + f*x))^(5/2)),x)`

[Out] `int(1/(sin(e + f*x)^4*(b/cos(e + f*x))^(5/2)), x)`

$$3.449 \quad \int \frac{\csc^6(e+fx)}{(b \sec(e+fx))^{5/2}} dx$$

Optimal. Leaf size=132

$$\frac{3 \csc(e+fx)}{20bf(b \sec(e+fx))^{3/2}} + \frac{\csc^3(e+fx)}{10bf(b \sec(e+fx))^{3/2}} - \frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} + \frac{3E\left(\frac{1}{2}(e+fx) \mid 2\right)}{20b^2f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

[Out] 3/20*csc(f*x+e)/b/f/(b*sec(f*x+e))^(3/2)+1/10*csc(f*x+e)^3/b/f/(b*sec(f*x+e))^(3/2)-1/5*csc(f*x+e)^5/b/f/(b*sec(f*x+e))^(3/2)+3/20*(cos(1/2*f*x+1/2*e))^2^(1/2)/cos(1/2*f*x+1/2*e)*EllipticE(sin(1/2*f*x+1/2*e),2^(1/2))/b^2/f/cos(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2)

Rubi [A]

time = 0.11, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {2703, 2705, 3856, 2719}

$$\frac{3E\left(\frac{1}{2}(e+fx) \mid 2\right)}{20b^2f \sqrt{\cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} + \frac{\csc^3(e+fx)}{10bf(b \sec(e+fx))^{3/2}} + \frac{3 \csc(e+fx)}{20bf(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^6/(b*Sec[e + f*x])^(5/2),x]

[Out] (3*Csc[e + f*x])/(20*b*f*(b*Sec[e + f*x])^(3/2)) + Csc[e + f*x]^3/(10*b*f*(b*Sec[e + f*x])^(3/2)) - Csc[e + f*x]^5/(5*b*f*(b*Sec[e + f*x])^(3/2)) + (3*EllipticE[(e + f*x)/2, 2])/(20*b^2*f*Sqrt[Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])

Rule 2703

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Simp[(-a)*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n + 1)/(f*b*(m - 1))), x] + Dist[a^2*((n + 1)/(b^2*(m - 1))), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && GtQ[m, 1] && LtQ[n, -1] && IntegersQ[2*m, 2*n]

Rule 2705

Int[(csc[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sec[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Simp[(-a)*b*(a*Csc[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - 1))), x] + Dist[a^2*((m + n - 2)/(m - 1)), Int[(a*Csc[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && IntegersQ[2*m, 2*n] && !GtQ[n, m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
 \int \frac{\csc^6(e+fx)}{(b \sec(e+fx))^{5/2}} dx &= -\frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} - \frac{3 \int \frac{\csc^4(e+fx)}{\sqrt{b \sec(e+fx)}} dx}{10b^2} \\
 &= \frac{\csc^3(e+fx)}{10bf(b \sec(e+fx))^{3/2}} - \frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} - \frac{3 \int \frac{\csc^2(e+fx)}{\sqrt{b \sec(e+fx)}} dx}{20b^2} \\
 &= \frac{3 \csc(e+fx)}{20bf(b \sec(e+fx))^{3/2}} + \frac{\csc^3(e+fx)}{10bf(b \sec(e+fx))^{3/2}} - \frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} + \frac{3 \int -}{40b^2} \\
 &= \frac{3 \csc(e+fx)}{20bf(b \sec(e+fx))^{3/2}} + \frac{\csc^3(e+fx)}{10bf(b \sec(e+fx))^{3/2}} - \frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} + \frac{3 \int -}{40b^2} \\
 &= \frac{3 \csc(e+fx)}{20bf(b \sec(e+fx))^{3/2}} + \frac{\csc^3(e+fx)}{10bf(b \sec(e+fx))^{3/2}} - \frac{\csc^5(e+fx)}{5bf(b \sec(e+fx))^{3/2}} + \frac{3 \int -}{20b^2}
 \end{aligned}$$

Mathematica [A]

time = 0.14, size = 87, normalized size = 0.66

$$\frac{(-3 + \csc^2(e+fx) + 6 \csc^4(e+fx) - 4 \csc^6(e+fx) + 3 \sqrt{\cos(e+fx)} \csc(e+fx) E(\frac{1}{2}(e+fx)|2)) \sqrt{b \sec(e+fx)} \sin(e+fx)}{20b^3 f}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^6/(b*Sec[e + f*x])^(5/2), x]

[Out] ((-3 + Csc[e + f*x]^2 + 6*Csc[e + f*x]^4 - 4*Csc[e + f*x]^6 + 3*Sqrt[Cos[e + f*x]]*Csc[e + f*x]*EllipticE[(e + f*x)/2, 2])*Sqrt[b*Sec[e + f*x]]*Sin[e + f*x])/(20*b^3*f)

Maple [C] Result contains complex when optimal does not.

time = 0.28, size = 923, normalized size = 6.99

method	result	size
default	Expression too large to display	923

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(csc(f*x+e)^6/(b*sec(f*x+e))^(5/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{20} f \frac{(\cos(f*x+e)+1)^2 (-1+\cos(f*x+e))^2 (-3I \frac{1}{(\cos(f*x+e)+1)})^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticE}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \cos(f*x+e)^4 \sin(f*x+e) + 3I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticF}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \cos(f*x+e)^4 \sin(f*x+e) + 3I \text{EllipticF}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \sin(f*x+e) + 6I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticE}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \cos(f*x+e)^3 \sin(f*x+e) - 3I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \cos(f*x+e) \text{EllipticE}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \sin(f*x+e) + 3I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticF}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \sin(f*x+e) \cos(f*x+e) + 6I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticE}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \cos(f*x+e)^2 \sin(f*x+e) - 6I \cos(f*x+e)^2 \sin(f*x+e) \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticF}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) - 3I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \sin(f*x+e) \text{EllipticE}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) - 3I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticE}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \cos(f*x+e)^5 \sin(f*x+e) + 3I \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticF}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) \cos(f*x+e)^5 \sin(f*x+e) - 6I \cos(f*x+e)^3 \sin(f*x+e) \frac{1}{(\cos(f*x+e)+1)}^{1/2} (\cos(f*x+e)/(\cos(f*x+e)+1))^{1/2} \text{EllipticF}(I \frac{-1+\cos(f*x+e)}{\sin(f*x+e)}, I) + 3 \cos(f*x+e)^5 - 2 \cos(f*x+e)^4 - 6 \cos(f*x+e)^3 - 2 \cos(f*x+e)^2 + 3 \cos(f*x+e) / \cos(f*x+e)^3 \sin(f*x+e)^9 / (b/\cos(f*x+e))^{5/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(5/2),x, algorithm="maxima")`

[Out] `integrate(csc(f*x + e)^6/(b*sec(f*x + e))^(5/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 220, normalized size = 1.67

$3\sqrt{2}(-1+\cos(fx+e)+2i\cos(fx+e)^2-i)\sqrt{b}\sin(fx+e)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(fx+e)+i\sin(fx+e)))+3\sqrt{2}(i\cos(fx+e)^3-2i\cos(fx+e)^2+i)\sqrt{b}\sin(fx+e)\text{weierstrassZeta}(-4,0,\text{weierstrassPInverse}(-4,0,\cos(fx+e)-i\sin(fx+e)))-2(3\cos(fx+e)^9-8\cos(fx+e)^8+\cos(fx+e)^7)\sqrt{\frac{b}{\cos(fx+e)}}$
 $40(b^3\cos(fx+e)^3-2b^2\cos(fx+e)^2+bf)\sin(fx+e)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(5/2),x, algorithm="fricas")`

[Out]
$$-1/40*(3*\sqrt{2})*(-I*\cos(f*x + e)^4 + 2*I*\cos(f*x + e)^2 - I)*\sqrt{b}*\sin(f*x + e)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(f*x + e) + I*\sin(f*x + e))) + 3*\sqrt{2}*(I*\cos(f*x + e)^4 - 2*I*\cos(f*x + e)^2 + I)*\sqrt{b}*\sin(f*x + e)*\text{weierstrassZeta}(-4, 0, \text{weierstrassPInverse}(-4, 0, \cos(f*x + e) - I*\sin(f*x + e))) - 2*(3*\cos(f*x + e)^6 - 8*\cos(f*x + e)^4 + \cos(f*x + e)^2)*\sqrt{b/\cos(f*x + e)})/((b^3*f*\cos(f*x + e)^4 - 2*b^3*f*\cos(f*x + e)^2 + b^3*f)*\sin(f*x + e))$$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**6/(b*sec(f*x+e))**(5/2),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^6/(b*sec(f*x+e))^(5/2),x, algorithm="giac")`

[Out] `integrate(csc(f*x + e)^6/(b*sec(f*x + e))^(5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^6 \left(\frac{b}{\cos(e + f x)}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^6*(b/cos(e + f*x))^(5/2)),x)`

[Out] `int(1/(sin(e + f*x)^6*(b/cos(e + f*x))^(5/2)), x)`

3.450 $\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx$

Optimal. Leaf size=449

$$\frac{21a^{9/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{32\sqrt{2} \sqrt{b} f} + \frac{21a^{9/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{32\sqrt{2} \sqrt{b} f}$$

[Out] $-7/16*a^3*b*(a*\sin(f*x+e))^{3/2}/f/(b*\sec(f*x+e))^{1/2}-1/4*a*b*(a*\sin(f*x+e))^{7/2}/f/(b*\sec(f*x+e))^{1/2}-21/64*a^{9/2}*arctan(1-2^{1/2}*b^{1/2}*(a*\sin(f*x+e))^{1/2}/a^{1/2}/(b*\cos(f*x+e))^{1/2})*(b*\cos(f*x+e))^{1/2}*(b*\sec(f*x+e))^{1/2}/f*2^{1/2}/b^{1/2}+21/64*a^{9/2}*arctan(1+2^{1/2}*b^{1/2}*(a*\sin(f*x+e))^{1/2}/a^{1/2}/(b*\cos(f*x+e))^{1/2})*(b*\cos(f*x+e))^{1/2}*(b*\sec(f*x+e))^{1/2}/f*2^{1/2}/b^{1/2}+21/128*a^{9/2}*ln(a^{1/2}-2^{1/2}*b^{1/2}*(a*\sin(f*x+e))^{1/2}/(b*\cos(f*x+e))^{1/2}+a^{1/2}*tan(f*x+e))*(b*\cos(f*x+e))^{1/2}*(b*\sec(f*x+e))^{1/2}/f*2^{1/2}/b^{1/2}-21/128*a^{9/2}*ln(a^{1/2}+2^{1/2}*b^{1/2}*(a*\sin(f*x+e))^{1/2}/(b*\cos(f*x+e))^{1/2}+a^{1/2}*tan(f*x+e))*(b*\cos(f*x+e))^{1/2}*(b*\sec(f*x+e))^{1/2}/f*2^{1/2}/b^{1/2}$

Rubi [A]

time = 0.31, antiderivative size = 449, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {2663, 2665, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{21a^{9/2} \sqrt{b \sec(e + fx)} \sqrt{b \cos(e + fx)} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{32\sqrt{2} \sqrt{b} f} + \frac{21a^{9/2} \sqrt{b \sec(e + fx)} \sqrt{b \cos(e + fx)} \operatorname{ArcTan} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{32\sqrt{2} \sqrt{b} f} + \frac{21a^{9/2} \sqrt{b \sec(e + fx)} \sqrt{b \cos(e + fx)} \ln \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{2} \tan(e + fx) + \sqrt{2} \right)}{64\sqrt{2} \sqrt{b} f} - \frac{21a^{9/2} \sqrt{b \sec(e + fx)} \sqrt{b \cos(e + fx)} \ln \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} - \sqrt{2} \tan(e + fx) + \sqrt{2} \right)}{64\sqrt{2} \sqrt{b} f} + \frac{21a^{9/2} \sqrt{b \sec(e + fx)} \sqrt{b \cos(e + fx)} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{16\sqrt{2} \sqrt{b} f} - \frac{21a^{9/2} \sqrt{b \sec(e + fx)} \sqrt{b \cos(e + fx)} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{16\sqrt{2} \sqrt{b} f}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(9/2),x]`

[Out] $(-21*a^{9/2}*ArcTan[1 - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*\cos[e + f*x]])]*Sqrt[b*\cos[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(32*Sqrt[2]*Sqrt[b]*f) + (21*a^{9/2}*ArcTan[1 + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*\cos[e + f*x]])]*Sqrt[b*\cos[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(32*Sqrt[2]*Sqrt[b]*f) + (21*a^{9/2}*Sqrt[b*\cos[e + f*x]]*Log[Sqrt[a] - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*\cos[e + f*x]] + Sqrt[a]*Tan[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(64*Sqrt[2]*Sqrt[b]*f) - (21*a^{9/2}*Sqrt[b*\cos[e + f*x]]*Log[Sqrt[a] + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*\cos[e + f*x]] + Sqrt[a]*Tan[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(64*Sqrt[2]*Sqrt[b]*f) - (7*a^3*b*(a*Sin[e + f*x])^{3/2})/(16*f*Sqrt[b*\sec[e + f*x]]) - (a*b*(a*Sin[e + f*x])^{7/2})/(4*f*Sqrt[b*\sec[e + f*x]])$

Rule 210

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &`

& (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] & & AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] & & EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2654

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sine[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] & & LtQ[m, 1]

Rule 2663

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Simp[(-a)*b*(a*Sin[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n
- 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*Sin[e + f*x])^(m
- 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1]
&& NeQ[m - n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +
f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte
gerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx &= -\frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} + \frac{1}{8} (7a^2) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2} dx \\
&= -\frac{7a^3 b (a \sin(e + fx))^{3/2}}{16f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} + \frac{1}{32} (21a^4) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2} dx \\
&= -\frac{7a^3 b (a \sin(e + fx))^{3/2}}{16f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} + \frac{1}{32} (21a^4 \sqrt{b \cos(e + fx)}) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2} dx \\
&= -\frac{7a^3 b (a \sin(e + fx))^{3/2}}{16f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} + \frac{(21a^5 b \sqrt{b \cos(e + fx)})}{32} \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{1/2} dx \\
&= -\frac{7a^3 b (a \sin(e + fx))^{3/2}}{16f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} - \frac{(21a^5 \sqrt{b \cos(e + fx)})}{32} \int \sqrt{b \sec(e + fx)} dx \\
&= -\frac{7a^3 b (a \sin(e + fx))^{3/2}}{16f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} + \frac{(21a^5 \sqrt{b \cos(e + fx)})}{32} \log \left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} \right) + \sqrt{b \cos(e + fx)} \\
&= -\frac{7a^3 b (a \sin(e + fx))^{3/2}}{16f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{7/2}}{4f \sqrt{b \sec(e + fx)}} + \frac{21a^{9/2} \sqrt{b \cos(e + fx)} \log \left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} \right) + \sqrt{b \cos(e + fx)}}{64\sqrt{2} \sqrt{b} f} \\
&= -\frac{21a^{9/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)}}{32\sqrt{2} \sqrt{b} f}
\end{aligned}$$

Mathematica [A]

time = 1.39, size = 169, normalized size = 0.38

$$\frac{a^4 \cot(e + fx) \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} \left(4(-9 + 2 \cos(2(e + fx))) \sin^2(e + fx) + 21\sqrt{2} \tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} - 21\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{2} \sqrt{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} \right)}{64f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(9/2),x]

[Out] (a^4*Cot[e + f*x]*Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]]*(4*(-9 + 2*Cos[2*(e + f*x)])*Sin[e + f*x]^2 + 21*Sqrt[2]*ArcTan[(-1 + Sqrt[Tan[e + f*x]^2])/(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))]*(Tan[e + f*x]^2)^(1/4) - 21*Sqrt[2]*ArcTanh[(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))/(1 + Sqrt[Tan[e + f*x]^2])]*(Tan[e + f*x]^2)^(1/4)))/(64*f)

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 1.40, size = 538, normalized size = 1.20

method	result
default	$-\frac{\left(21i \sqrt{\frac{1 - \cos(fx+e) + \sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e) - 1 + \cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1 + \cos(fx+e)}{\sin(fx+e)}} \text{EllipticPi} \left(\sqrt{\frac{1 - \cos(fx+e) + \sin(fx+e)}{\sin(fx+e)}} \right) \right)}{64f}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^(9/2)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/64/f*(21*I*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-21*I*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))-8*cos(f*x+e)^4*2^(1/2)-21*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-21*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))+8*cos(f*x+e)^3*2^(1/2)+22*cos(f*x+e)^2*2^(1/2)-22*cos(f*x+e)*2^(1/2))*(a*sin(f*x+e))^(9/2)*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^3/(-1+cos(f*x+e))*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(9/2)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(9/2), x)
```

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(9/2)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] Timed out
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))**(9/2)*(b*sec(f*x+e))**(1/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(9/2)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(9/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a \sin(e + f x))^{9/2} \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*sin(e + f*x))^(9/2)*(b/cos(e + f*x))^(1/2),x)
```

```
[Out] int((a*sin(e + f*x))^(9/2)*(b/cos(e + f*x))^(1/2), x)
```

3.451 $\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2} dx$

Optimal. Leaf size=414

$$\frac{3a^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} + 3a^{5/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b}}{\sqrt{a}} \right)}{4\sqrt{2} \sqrt{b} f}$$

[Out] $-1/2*a*b*(a*\sin(f*x+e))^{(3/2)}/f/(b*\sec(f*x+e))^{(1/2)}-3/8*a^{(5/2)*\arctan(1-2^{(1/2)*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/a^{(1/2)}/(b*\cos(f*x+e))^{(1/2)})*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f*2^{(1/2)}/b^{(1/2)}+3/8*a^{(5/2)*\arctan(1+2^{(1/2)*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/a^{(1/2)}/(b*\cos(f*x+e))^{(1/2)})*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f*2^{(1/2)}/b^{(1/2)}+3/16*a^{(5/2)*\ln(a^{(1/2)}-2^{(1/2)*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}+a^{(1/2)*\tan(f*x+e)})*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f*2^{(1/2)}/b^{(1/2)}-3/16*a^{(5/2)*\ln(a^{(1/2)}+2^{(1/2)*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}+a^{(1/2)*\tan(f*x+e)})*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/f*2^{(1/2)}/b^{(1/2)}$

Rubi [A]

time = 0.24, antiderivative size = 414, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {2663, 2665, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{3a^{5/2} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) + 3a^{5/2} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + 1 \right) + 3a^{5/2} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \log \left(\frac{-\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) + \sqrt{a} \right) + 3a^{5/2} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \log \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) + \sqrt{a} \right) - \frac{ab(a \sin(e + fx))^{5/2}}{2f \sqrt{b \cos(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]]*(a*\operatorname{Sin}[e + f*x])^{(5/2)}, x]$

[Out] $(-3*a^{(5/2)*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\operatorname{Sin}[e + f*x]])]/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]])]*\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]]*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(4*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*f) + (3*a^{(5/2)*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\operatorname{Sin}[e + f*x]])]/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]])]*\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]]*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(4*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*f) + (3*a^{(5/2)*\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]]*\operatorname{Log}[\operatorname{Sqrt}[a] - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\operatorname{Sin}[e + f*x]])]/\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]] + \operatorname{Sqrt}[a]*\operatorname{Tan}[e + f*x]]*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(8*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*f) - (3*a^{(5/2)*\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]]*\operatorname{Log}[\operatorname{Sqrt}[a] + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\operatorname{Sin}[e + f*x]])]/\operatorname{Sqrt}[b*\operatorname{Cos}[e + f*x]] + \operatorname{Sqrt}[a]*\operatorname{Tan}[e + f*x]]*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])/(8*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*f) - (a*b*(a*\operatorname{Sin}[e + f*x])^{(3/2)})/(2*f*\operatorname{Sqrt}[b*\operatorname{Sec}[e + f*x]])$

Rule 210

$\operatorname{Int}[(a + (b \cdot x)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-\operatorname{Rt}[-a, 2]*\operatorname{Rt}[-b, 2])^{(-1)}* \operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[-a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ \operatorname{PosQ}[a/b] \ \& \ \& \ (\operatorname{LtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 303

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rule 2663

```
Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := Simp[(-a)*b*(a*Sin[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n
```


- 1)/(f*(m - n)), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*Sin[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rubi steps

$$\begin{aligned}
 \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2} dx &= -\frac{ab(a \sin(e + fx))^{3/2}}{2f \sqrt{b \sec(e + fx)}} + \frac{1}{4}(3a^2) \int \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} dx \\
 &= -\frac{ab(a \sin(e + fx))^{3/2}}{2f \sqrt{b \sec(e + fx)}} + \frac{1}{4} \left(3a^2 \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \\
 &= -\frac{ab(a \sin(e + fx))^{3/2}}{2f \sqrt{b \sec(e + fx)}} + \frac{(3a^3 b \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{4} \\
 &= -\frac{ab(a \sin(e + fx))^{3/2}}{2f \sqrt{b \sec(e + fx)}} - \frac{(3a^3 \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{4} \\
 &= -\frac{ab(a \sin(e + fx))^{3/2}}{2f \sqrt{b \sec(e + fx)}} + \frac{(3a^3 \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{4} \\
 &= \frac{3a^{5/2} \sqrt{b \cos(e + fx)} \log \left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} \right) + \sqrt{b \cos(e + fx)} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{8\sqrt{2} \sqrt{b} f} \\
 &= -\frac{3a^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)}}{4\sqrt{2} \sqrt{b} f}
 \end{aligned}$$

Mathematica [A]

time = 0.76, size = 157, normalized size = 0.38

$$\frac{a^2 \cot(e + fx) \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} \left(4 \sin^2(e + fx) - 3\sqrt{2} \tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} + 3\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{2} \sqrt{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} \right)}{8f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(5/2),x]

[Out] $-\frac{1}{8}(a^2 \cot[e + f x] \sqrt{b \sec[e + f x]} \sqrt{a \sin[e + f x]} (4 \sin[e + f x]^2 - 3 \sqrt{2} \operatorname{ArcTan}[-1 + \sqrt{\tan[e + f x]^2}]) / (\sqrt{2} (\tan[e + f x]^2)^{1/4})) + 3 \sqrt{2} \operatorname{ArcTanh}[(\sqrt{2} (\tan[e + f x]^2)^{1/4}) / (1 + \sqrt{\tan[e + f x]^2})] (\tan[e + f x]^2)^{1/4}) / f$

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.27, size = 512, normalized size = 1.24

method	result
default	$-\left(3i \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^(5/2)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] $-\frac{1}{8}f(3I \operatorname{EllipticPi}(((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2}), 1/2-1/2I, 1/2*2^{1/2}) * ((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2} * ((\sin(fx+e)-1+\cos(fx+e))/\sin(fx+e))^{1/2} * ((-1+\cos(fx+e))/\sin(fx+e))^{1/2} - 3I * ((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2} * ((\sin(fx+e)-1+\cos(fx+e))/\sin(fx+e))^{1/2} * ((-1+\cos(fx+e))/\sin(fx+e))^{1/2} * \operatorname{EllipticPi}(((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2}), 1/2+1/2I, 1/2*2^{1/2}) - 3 * ((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2} * ((\sin(fx+e)-1+\cos(fx+e))/\sin(fx+e))^{1/2} * ((-1+\cos(fx+e))/\sin(fx+e))^{1/2} * \operatorname{EllipticPi}(((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2}), 1/2-1/2I, 1/2*2^{1/2}) - 3 * ((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2} * ((\sin(fx+e)-1+\cos(fx+e))/\sin(fx+e))^{1/2} * ((-1+\cos(fx+e))/\sin(fx+e))^{1/2} * \operatorname{EllipticPi}(((1-\cos(fx+e)+\sin(fx+e))/\sin(fx+e))^{1/2}), 1/2+1/2I, 1/2*2^{1/2}) + 2 * \cos(fx+e)^2 * 2^{1/2} - 2 * \cos(fx+e) * 2^{1/2}) * (a * \sin(fx+e))^{5/2} * (b / \cos(fx+e))^{1/2} / (-1 + \cos(fx+e)) / \sin(fx+e) * 2^{1/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(5/2)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(5/2), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sin(f*x+e))^(5/2)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sin(f*x+e))**(5/2)*(b*sec(f*x+e))**(1/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sin(f*x+e))^(5/2)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a \sin(e + f x))^{5/2} \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sin(e + f*x))^(5/2)*(b/cos(e + f*x))^(1/2),x)`

[Out] `int((a*sin(e + f*x))^(5/2)*(b/cos(e + f*x))^(1/2), x)`

3.452 $\int \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} dx$

Optimal. Leaf size=376

$$\frac{\sqrt{a} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} + \sqrt{a} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{\sqrt{2} \sqrt{b} f}$$

[Out] $-1/2*\arctan(1-2^{(1/2)*b^{(1/2)*(a*\sin(f*x+e))^{(1/2)/a^{(1/2)/(b*\cos(f*x+e))^{(1/2)}}*a^{(1/2)*(b*\cos(f*x+e))^{(1/2)*(b*\sec(f*x+e))^{(1/2)/f*2^{(1/2)/b^{(1/2)+1/2}}*\arctan(1+2^{(1/2)*b^{(1/2)*(a*\sin(f*x+e))^{(1/2)/a^{(1/2)/(b*\cos(f*x+e))^{(1/2)}}*a^{(1/2)*(b*\cos(f*x+e))^{(1/2)*(b*\sec(f*x+e))^{(1/2)/f*2^{(1/2)/b^{(1/2)+1/4}}*\ln(a^{(1/2)-2^{(1/2)*b^{(1/2)*(a*\sin(f*x+e))^{(1/2)/(b*\cos(f*x+e))^{(1/2)+a^{(1/2)*\tan(f*x+e))})*a^{(1/2)*(b*\cos(f*x+e))^{(1/2)*(b*\sec(f*x+e))^{(1/2)/f*2^{(1/2)/b^{(1/2)-1/4}}*\ln(a^{(1/2)+2^{(1/2)*b^{(1/2)*(a*\sin(f*x+e))^{(1/2)/(b*\cos(f*x+e))^{(1/2)+a^{(1/2)*\tan(f*x+e))})*a^{(1/2)*(b*\cos(f*x+e))^{(1/2)*(b*\sec(f*x+e))^{(1/2)/f*2^{(1/2)/b^{(1/2)}}}$

Rubi [A]

time = 0.17, antiderivative size = 376, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.320$, Rules used = {2665, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) + \sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + 1 \right) + \sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \log \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) + \sqrt{a} \right) + \sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \log \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) + \sqrt{a} \right)}{2\sqrt{2} \sqrt{b} f}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]],x]

[Out] $-((\operatorname{Sqrt}[a] \operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2] \operatorname{Sqrt}[b] \operatorname{Sqrt}[a \operatorname{Sin}[e + f*x]])]/(\operatorname{Sqrt}[a] \operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]])]) \operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]] \operatorname{Sqrt}[b \operatorname{Sec}[e + f*x]])/(\operatorname{Sqrt}[2] \operatorname{Sqrt}[b] * f) + (\operatorname{Sqrt}[a] \operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2] \operatorname{Sqrt}[b] \operatorname{Sqrt}[a \operatorname{Sin}[e + f*x]])]/(\operatorname{Sqrt}[a] \operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]])]) \operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]] \operatorname{Sqrt}[b \operatorname{Sec}[e + f*x]])/(\operatorname{Sqrt}[2] \operatorname{Sqrt}[b] * f) + (\operatorname{Sqrt}[a] \operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]] \operatorname{Log}[\operatorname{Sqrt}[a] - (\operatorname{Sqrt}[2] \operatorname{Sqrt}[b] \operatorname{Sqrt}[a \operatorname{Sin}[e + f*x]])/\operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]] + \operatorname{Sqrt}[a] \operatorname{Tan}[e + f*x]] \operatorname{Sqrt}[b \operatorname{Sec}[e + f*x]])/(2 \operatorname{Sqrt}[2] \operatorname{Sqrt}[b] * f) - (\operatorname{Sqrt}[a] \operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]] \operatorname{Log}[\operatorname{Sqrt}[a] + (\operatorname{Sqrt}[2] \operatorname{Sqrt}[b] \operatorname{Sqrt}[a \operatorname{Sin}[e + f*x]])/\operatorname{Sqrt}[b \operatorname{Cos}[e + f*x]] + \operatorname{Sqrt}[a] \operatorname{Tan}[e + f*x]] \operatorname{Sqrt}[b \operatorname{Sec}[e + f*x]])/(2 \operatorname{Sqrt}[2] \operatorname{Sqrt}[b] * f)$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] & AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2654

```
Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^m_, x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^m_, x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte
```

gerQ[m - 1/2] && IntegerQ[n - 1/2]

Rubi steps

$$\begin{aligned}
 \int \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} dx &= \left(\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{\sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} dx \\
 &= \frac{\left(2ab \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \text{Subst} \left(\int \frac{x^2}{a^2 + b^2 x^4} dx, x, \frac{\sqrt{a}}{\sqrt{b}} \right)}{f} \\
 &= - \frac{\left(a \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \text{Subst} \left(\int \frac{a - bx^2}{a^2 + b^2 x^4} dx, x, \frac{\sqrt{a}}{\sqrt{b}} \right)}{f} \\
 &= \frac{\left(a \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \text{Subst} \left(\int \frac{1}{\frac{a}{b} - \frac{\sqrt{2} \sqrt{a}}{\sqrt{b}} x + x^2} dx, \right)}{2bf} \\
 &= \frac{\sqrt{a} \sqrt{b \cos(e + fx)} \log \left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} + \sqrt{a} \right)}{2\sqrt{2} \sqrt{b} f} \\
 &= - \frac{\sqrt{a} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{\sqrt{2} \sqrt{b} f}
 \end{aligned}$$

Mathematica [A]

time = 0.44, size = 122, normalized size = 0.32

$$\frac{\left(\tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt[4]{\tan^2(e + fx)}} \right) - \tanh^{-1} \left(\frac{\sqrt{2} \sqrt[4]{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \right) \cot(e + fx) \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} \sqrt[4]{\tan^2(e + fx)}}{\sqrt{2} f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]],x]

[Out] ((ArcTan[(-1 + Sqrt[Tan[e + f*x]^2])]/(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))]) - ArcTanh[(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))/(1 + Sqrt[Tan[e + f*x]^2])])*Cot[e + f*x]*Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]]*(Tan[e + f*x]^2)^(1/4)/(Sqrt[2]*f)

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.28, size = 273, normalized size = 0.73

method	result
default	$-\frac{\sqrt{a \sin(fx + e)} \sqrt{\frac{b}{\cos(fx + e)}} \sqrt{\frac{1 - \cos(fx + e) + \sin(fx + e)}{\sin(fx + e)}} \sqrt{\frac{\sin(fx + e) - 1 + \cos(fx + e)}{\sin(fx + e)}} \sqrt{\frac{-1 + \cos(fx + e)}{\sin(fx + e)}}}{\left(i \text{ Ell} \right)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sin(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/f*(a*\sin(f*x+e))^{(1/2)}*(b/\cos(f*x+e))^{(1/2)}*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*(I*\text{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})-I*\text{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})-\text{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})-\text{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)}))*\sin(f*x+e)/(-1+\cos(f*x+e))*2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sin(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e)), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sin(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{a \sin(e + fx)} \sqrt{b \sec(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))**(1/2)*(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(a*sin(e + f*x))*sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{a \sin(e + f x)} \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(1/2)*(b/cos(e + f*x))^(1/2),x)

[Out] int((a*sin(e + f*x))^(1/2)*(b/cos(e + f*x))^(1/2), x)

$$3.453 \quad \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{3/2}} dx$$

Optimal. Leaf size=33

$$-\frac{2b}{af \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

[Out] $-2*b/a/f/(b*\sec(f*x+e))^{(1/2)}/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 33, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2658}

$$-\frac{2b}{af \sqrt{a \sin(e + fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[e + f*x]]/(a*\text{Sin}[e + f*x])^{(3/2)}, x]$

[Out] $(-2*b)/(a*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2658

$\text{Int}[(b_*)*\sec[(e_*) + (f_*)*(x_*)]^{(n_*)}*((a_*)*\sin[(e_*) + (f_*)*(x_*)])^{(m_*)}, x_Symbol] :> \text{Simp}[b*(a*\text{Sin}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(a*f*(m + 1)), x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \text{EqQ}[m - n + 2, 0] \& \& \text{NeQ}[m, -1]$

Rubi steps

$$\int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{3/2}} dx = -\frac{2b}{af \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

Mathematica [A]

time = 0.06, size = 37, normalized size = 1.12

$$-\frac{\sqrt{b \sec(e + fx)} \sin(2(e + fx))}{f(a \sin(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(3/2),x]

[Out] -((Sqrt[b*Sec[e + f*x]]*Sin[2*(e + f*x)])/(f*(a*Sin[e + f*x])^(3/2)))

Maple [A]

time = 0.23, size = 40, normalized size = 1.21

method	result	size
default	$-\frac{2 \sin(fx+e) \cos(fx+e) \sqrt{\frac{b}{\cos(fx+e)}}}{f(a \sin(fx+e))^{\frac{3}{2}}}$	40

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] -2/f*sin(f*x+e)*cos(f*x+e)*(b/cos(f*x+e))^(1/2)/(a*sin(f*x+e))^(3/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(3/2), x)

Fricas [A]

time = 0.43, size = 48, normalized size = 1.45

$$-\frac{2 \sqrt{a \sin(fx+e)} \sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)}{a^2 f \sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(3/2),x, algorithm="fricas")

[Out] -2*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e))*cos(f*x + e)/(a^2*f*sin(f*x + e))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(1/2)/(a*sin(f*x+e))**(3/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))/(a*sin(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(3/2), x)

Mupad [B]

time = 0.87, size = 36, normalized size = 1.09

$$-\frac{2 \cos(e + f x) \sqrt{\frac{b}{\cos(e + f x)}}}{a f \sqrt{a \sin(e + f x)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(3/2),x)

[Out] -(2*cos(e + f*x)*(b/cos(e + f*x))^(1/2))/(a*f*(a*sin(e + f*x))^(1/2))

$$3.454 \quad \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{7/2}} dx$$

Optimal. Leaf size=71

$$-\frac{2b}{5af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2}} - \frac{8b}{5a^3 f \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

[Out] $-2/5*b/a/f/(a*\sin(f*x+e))^{(5/2)}/(b*\sec(f*x+e))^{(1/2)}-8/5*b/a^3/f/(b*\sec(f*x+e))^{(1/2)}/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2664, 2658}

$$-\frac{8b}{5a^3 f \sqrt{a \sin(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{2b}{5af (a \sin(e + fx))^{5/2} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(7/2),x]`

[Out] $(-2*b)/(5*a*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(5/2)}) - (8*b)/(5*a^3*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2658

`Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m - n + 2, 0] & & NeQ[m, -1]`

Rule 2664

`Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] := Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rubi steps

$$\int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{7/2}} dx = -\frac{2b}{5af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2}} + \frac{4 \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{3/2}} dx}{5a^2}$$

$$= -\frac{2b}{5af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2}} - \frac{8b}{5a^3 f \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

Mathematica [A]

time = 0.14, size = 52, normalized size = 0.73

$$\frac{2(-3 + 2 \cos(2(e + fx))) \cot(e + fx) \sqrt{b \sec(e + fx)}}{5a^2 f (a \sin(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(7/2),x]``[Out] (2*(-3 + 2*Cos[2*(e + f*x)])*Cot[e + f*x]*Sqrt[b*Sec[e + f*x]])/(5*a^2*f*(a*Sin[e + f*x])^(3/2))`**Maple [A]**

time = 0.24, size = 52, normalized size = 0.73

method	result	size
default	$\frac{2(4(\cos^2(fx+e))-5) \cos(fx+e) \sqrt{\frac{b}{\cos(fx+e)}} \sin(fx+e)}{5f(a \sin(fx+e))^{\frac{7}{2}}}$	52

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(7/2),x,method=_RETURNVERBOSE)``[Out] 2/5/f*(4*cos(f*x+e)^2-5)*cos(f*x+e)*(b/cos(f*x+e))^(1/2)*sin(f*x+e)/(a*sin(f*x+e))^(7/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(7/2),x, algorithm="maxima")``[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(7/2), x)`

Fricas [A]

time = 0.51, size = 79, normalized size = 1.11

$$\frac{2(4 \cos(fx + e)^3 - 5 \cos(fx + e)) \sqrt{a \sin(fx + e)} \sqrt{\frac{b}{\cos(fx + e)}}}{5(a^4 f \cos(fx + e)^2 - a^4 f) \sin(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(7/2),x, algorithm="fricas")
```

```
[Out] -2/5*(4*cos(f*x + e)^3 - 5*cos(f*x + e))*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e))/((a^4*f*cos(f*x + e)^2 - a^4*f)*sin(f*x + e))
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))**(1/2)/(a*sin(f*x+e))**(7/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(7/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(7/2), x)
```

Mupad [B]

time = 1.81, size = 83, normalized size = 1.17

$$\frac{4 \sqrt{\frac{b}{\cos(e + fx)}} (3 \cos(e + fx) - 4 \cos(3e + 3fx) + \cos(5e + 5fx))}{5a^3 f \sqrt{a \sin(e + fx)} (\cos(4e + 4fx) - 4 \cos(2e + 2fx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(7/2),x)
```

```
[Out] -(4*(b/cos(e + f*x))^(1/2)*(3*cos(e + f*x) - 4*cos(3*e + 3*f*x) + cos(5*e + 5*f*x)))/(5*a^3*f*(a*sin(e + f*x))^(1/2)*(cos(4*e + 4*f*x) - 4*cos(2*e + 2*f*x) + 3))
```

$$3.455 \quad \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{11/2}} dx$$

Optimal. Leaf size=106

$$\frac{2b}{9af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2}} - \frac{16b}{45a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2}} - \frac{64b}{45a^5 f \sqrt{b \sec(e + fx)}}$$

[Out] $-2/9*b/a/f/(a*\sin(f*x+e))^(9/2)/(b*\sec(f*x+e))^(1/2)-16/45*b/a^3/f/(a*\sin(f*x+e))^(5/2)/(b*\sec(f*x+e))^(1/2)-64/45*b/a^5/f/(b*\sec(f*x+e))^(1/2)/(a*\sin(f*x+e))^(1/2)$

Rubi [A]

time = 0.11, antiderivative size = 106, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.080$, Rules used = {2664, 2658}

$$\frac{64b}{45a^5 f \sqrt{a \sin(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{16b}{45a^3 f (a \sin(e + fx))^{5/2} \sqrt{b \sec(e + fx)}} - \frac{2b}{9af (a \sin(e + fx))^{9/2} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(11/2), x]

[Out] $(-2*b)/(9*a*f*Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(9/2)) - (16*b)/(45*a^3*f*Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(5/2)) - (64*b)/(45*a^5*f*Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]])$

Rule 2658

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m - n + 2, 0] && NeQ[m, -1]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{11/2}} dx = -\frac{2b}{9af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2}} + \frac{8 \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{7/2}} dx}{9a^2}$$

$$= -\frac{2b}{9af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2}} - \frac{16b}{45a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2}}$$

$$= -\frac{2b}{9af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2}} - \frac{16b}{45a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2}}$$

Mathematica [A]

time = 0.15, size = 65, normalized size = 0.61

$$\frac{2b(-21 + 20 \cos(2(e + fx)) - 4 \cos(4(e + fx))) \csc^5(e + fx) \sqrt{a \sin(e + fx)}}{45a^6 f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(11/2), x]``[Out] (2*b*(-21 + 20*Cos[2*(e + f*x)] - 4*Cos[4*(e + f*x)])*Csc[e + f*x]^5*Sqrt[a*Sin[e + f*x]])/(45*a^6*f*Sqrt[b*Sec[e + f*x]])`**Maple [A]**

time = 0.26, size = 62, normalized size = 0.58

method	result	size
default	$-\frac{2(32(\cos^4(fx+e))-72(\cos^2(fx+e))+45)\cos(fx+e)\sqrt{\frac{b}{\cos(fx+e)}}\sin(fx+e)}{45f(a\sin(fx+e))^{\frac{11}{2}}}$	62

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(11/2), x, method=_RETURNVERBOSE)``[Out] -2/45/f*(32*cos(f*x+e)^4-72*cos(f*x+e)^2+45)*cos(f*x+e)*(b/cos(f*x+e))^(1/2)*sin(f*x+e)/(a*sin(f*x+e))^(11/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(11/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(11/2), x)

Fricas [A]

time = 0.50, size = 104, normalized size = 0.98

$$\frac{2 \left(32 \cos (f x + e)^5 - 72 \cos (f x + e)^3 + 45 \cos (f x + e) \right) \sqrt{a \sin (f x + e)} \sqrt{\frac{b}{\cos (f x + e)}}}{45 \left(a^6 f \cos (f x + e)^4 - 2 a^6 f \cos (f x + e)^2 + a^6 f \right) \sin (f x + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(11/2),x, algorithm="fricas")

[Out] -2/45*(32*cos(f*x + e)^5 - 72*cos(f*x + e)^3 + 45*cos(f*x + e))*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e))/((a^6*f*cos(f*x + e)^4 - 2*a^6*f*cos(f*x + e)^2 + a^6*f)*sin(f*x + e))

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(11/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(11/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(11/2), x)

Mupad [B]

time = 5.69, size = 169, normalized size = 1.59

$$\frac{e^{-e 5i - f x 5i} \sqrt{\frac{b}{\frac{e^{-e 1i - f x 1i}}{2} + \frac{e^{e 1i + f x 1i}}{2}}} \left(\frac{352 \cos(e + f x) e^{e 5i + f x 5i}}{45 a^5 f} - \frac{256 e^{e 5i + f x 5i} \cos(3e + 3f x)}{45 a^5 f} + \frac{64 e^{e 5i + f x 5i} \cos(5e + 5f x)}{45 a^5 f} \right)}{16 \sin(e + f x)^4 \sqrt{a \left(\frac{e^{-e 1i - f x 1i} 1i}{2} - \frac{e^{e 1i + f x 1i} 1i}{2} \right)}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((b/\cos(e + f*x))^{1/2}/(a*\sin(e + f*x))^{11/2},x)$

[Out] $-(\exp(-e*5i - f*x*5i)*(b/(\exp(-e*1i - f*x*1i)/2 + \exp(e*1i + f*x*1i)/2))^{1/2}*((352*\cos(e + f*x)*\exp(e*5i + f*x*5i))/(45*a^5*f) - (256*\exp(e*5i + f*x*5i)*\cos(3*e + 3*f*x))/(45*a^5*f) + (64*\exp(e*5i + f*x*5i)*\cos(5*e + 5*f*x))/(45*a^5*f)))/(16*\sin(e + f*x)^4*(a*((\exp(-e*1i - f*x*1i)*1i)/2 - (\exp(e*1i + f*x*1i)*1i)/2))^{1/2})$

3.456 $\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2} dx$

Optimal. Leaf size=128

$$-\frac{5a^3b\sqrt{a\sin(e+fx)}}{6f\sqrt{b\sec(e+fx)}} - \frac{ab(a\sin(e+fx))^{5/2}}{3f\sqrt{b\sec(e+fx)}} + \frac{5a^4F(e-\frac{\pi}{4}+fx|2)\sqrt{b\sec(e+fx)}\sqrt{\sin(2e+2fx)}}{12f\sqrt{a\sin(e+fx)}}$$

[Out] $-1/3*a*b*(a*\sin(f*x+e))^{(5/2)}/f/(b*\sec(f*x+e))^{(1/2)}-5/6*a^3*b*(a*\sin(f*x+e))^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}-5/12*a^4*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticF}(\cos(e+1/4*Pi+f*x),2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2663, 2665, 2653, 2720}

$$\frac{5a^4\sqrt{\sin(2e+2fx)}F(e+fx-\frac{\pi}{4}|2)\sqrt{b\sec(e+fx)}}{12f\sqrt{a\sin(e+fx)}} - \frac{5a^3b\sqrt{a\sin(e+fx)}}{6f\sqrt{b\sec(e+fx)}} - \frac{ab(a\sin(e+fx))^{5/2}}{3f\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(7/2)}, x]$

[Out] $(-5*a^3*b*\text{Sqrt}[a*\text{Sin}[e + f*x]])/(6*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (a*b*(a*\text{Sin}[e + f*x])^{(5/2)})/(3*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (5*a^4*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(12*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2663

$\text{Int}[(b_.)*\sec[(e_.) + (f_.)*(x_.)]^{(n)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m)}, x_Symbol] \rightarrow \text{Simp}[(-a)*b*(a*\text{Sin}[e + f*x])^{(m-1)}*(b*\text{Sec}[e + f*x])^{(n-1)}/(f*(m-n)), x] + \text{Dist}[a^2*((m-1)/(m-n)), \text{Int}[(a*\text{Sin}[e + f*x])^{(m-2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{GtQ}[m, 1] \&\& \text{NeQ}[m-n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2665

$\text{Int}[(b_.)*\sec[(e_.) + (f_.)*(x_.)]^{(n)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Cos}[e + f*x])^n*(b*\text{Sec}[e + f*x])^n, \text{Int}[(a*\text{Sin}[e +$

$f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2} dx &= -\frac{ab(a \sin(e + fx))^{5/2}}{3f \sqrt{b \sec(e + fx)}} + \frac{1}{6}(5a^2) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2} dx \\ &= -\frac{5a^3 b \sqrt{a \sin(e + fx)}}{6f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{5/2}}{3f \sqrt{b \sec(e + fx)}} + \frac{1}{12}(5a^4) \int \frac{\sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}}{\sqrt{a \sin(e + fx)}} dx \\ &= -\frac{5a^3 b \sqrt{a \sin(e + fx)}}{6f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{5/2}}{3f \sqrt{b \sec(e + fx)}} + \frac{1}{12} \left(5a^4 \sqrt{b \cos(e + fx)} \int \frac{1}{\sqrt{a \sin(e + fx)}} dx \right) \\ &= -\frac{5a^3 b \sqrt{a \sin(e + fx)}}{6f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{5/2}}{3f \sqrt{b \sec(e + fx)}} + \frac{5a^4 \sqrt{b \sec(e + fx)} F\left(e - \frac{\pi}{4}, \frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \sec^2(e + fx)\right)}{12} \\ &= -\frac{5a^3 b \sqrt{a \sin(e + fx)}}{6f \sqrt{b \sec(e + fx)}} - \frac{ab(a \sin(e + fx))^{5/2}}{3f \sqrt{b \sec(e + fx)}} + \frac{5a^4 F\left(e - \frac{\pi}{4}, \frac{1}{2}, \frac{3}{4}, \frac{3}{2}, \sec^2(e + fx)\right)}{12} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 15.70, size = 90, normalized size = 0.70

$$\frac{a^3 b \sqrt{a \sin(e + fx)} \left(2(-6 + \cos(2(e + fx))) + 5 \csc^2(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4} \right)}{12f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(7/2),x]

[Out] (a^3*b*Sqrt[a*Sin[e + f*x]]*(2*(-6 + Cos[2*(e + f*x)]) + 5*Csc[e + f*x]^2*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4)))/(12*f*Sqrt[b*Sec[e + f*x]])

Maple [A]

time = 0.36, size = 212, normalized size = 1.66

method	result
default	$-\frac{\left(5\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{12f\sin(fx+e)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*sin(f*x+e))^(7/2)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/12/f*(5*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)-2*cos(f*x+e)^4*2^(1/2)+2*cos(f*x+e)^3*2^(1/2)+7*cos(f*x+e)^2*2^(1/2)-7*cos(f*x+e)*2^(1/2))*(a*sin(f*x+e))^(7/2)*(b/cos(f*x+e))^(1/2)/sin(f*x+e)^3/(-1+cos(f*x+e))*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(7/2)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(7/2), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(7/2)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] integral(-(a^3*cos(f*x + e)^2 - a^3)*sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e))*sin(f*x + e), x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))**(7/2)*(b*sec(f*x+e))**(1/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(7/2)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(7/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \sin(e + f x))^{7/2} \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(7/2)*(b/cos(e + f*x))^(1/2),x)

[Out] int((a*sin(e + f*x))^(7/2)*(b/cos(e + f*x))^(1/2), x)

3.457 $\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2} dx$

Optimal. Leaf size=91

$$-\frac{ab\sqrt{a\sin(e+fx)}}{f\sqrt{b\sec(e+fx)}} + \frac{a^2 F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b\sec(e+fx)} \sqrt{\sin(2e+2fx)}}{2f\sqrt{a\sin(e+fx)}}$$

[Out] $-a*b*(a*\sin(f*x+e))^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}-1/2*a^2*(\sin(e+1/4*\text{Pi}+f*x)^2)^{(1/2)}/\sin(e+1/4*\text{Pi}+f*x)*\text{EllipticF}(\cos(e+1/4*\text{Pi}+f*x),2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.10, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2663, 2665, 2653, 2720}

$$\frac{a^2 \sqrt{\sin(2e + 2fx)} F\left(e + fx - \frac{\pi}{4} \mid 2\right) \sqrt{b \sec(e + fx)}}{2f \sqrt{a \sin(e + fx)}} - \frac{ab \sqrt{a \sin(e + fx)}}{f \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(3/2),x]`

[Out] $-((a*b*\text{Sqrt}[a*\text{Sin}[e + f*x]])/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]])) + (a^2*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(2*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

`Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)])], x_Symbol] := Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

Rule 2663

`Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[(-a)*b*(a*Sin[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*Sin[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]`

Rule 2665

`Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte`

gerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2} dx &= -\frac{ab \sqrt{a \sin(e + fx)}}{f \sqrt{b \sec(e + fx)}} + \frac{1}{2} a^2 \int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx \\
 &= -\frac{ab \sqrt{a \sin(e + fx)}}{f \sqrt{b \sec(e + fx)}} + \frac{1}{2} \left(a^2 \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \right) \int \frac{1}{\sqrt{a \sin(e + fx)}} dx \\
 &= -\frac{ab \sqrt{a \sin(e + fx)}}{f \sqrt{b \sec(e + fx)}} + \frac{\left(a^2 \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)} \right) \int \frac{1}{\sqrt{a \sin(e + fx)}} dx}{2 \sqrt{a \sin(e + fx)}} \\
 &= -\frac{ab \sqrt{a \sin(e + fx)}}{f \sqrt{b \sec(e + fx)}} + \frac{a^2 F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}{2f \sqrt{a \sin(e + fx)}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.92, size = 66, normalized size = 0.73

$$\frac{{}_2F_1\left(-\frac{1}{2}, -\frac{1}{4}; \frac{1}{2}; \sec^2(e + fx)\right) (b \sec(e + fx))^{3/2} (a \sin(e + fx))^{5/2}}{abf (-\tan^2(e + fx))^{5/4}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(3/2), x]

[Out] (Hypergeometric2F1[-1/2, -1/4, 1/2, Sec[e + f*x]^2]*(b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(5/2))/(a*b*f*(-Tan[e + f*x]^2)^(5/4))

Maple [A]

time = 0.29, size = 184, normalized size = 2.02

method	result
default	$ -\frac{\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \text{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}, \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{2f(-1+\cos(fx+e))\sin(fx+e)} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*sin(f*x+e))^(3/2)*(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2/f*(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)+cos(f*x+e)^2*2^(1/2)-cos(f*x+e)*2^(1/2))*(a*sin(f*x+e))^(3/2)*(b/cos(f*x+e))^(1/2)/(-1+cos(f*x+e))/sin(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(3/2)*(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(3/2), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(3/2)*(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e))*a*sin(f*x + e), x)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))**(3/2)*(b*sec(f*x+e))**(1/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 6437 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(3/2)*(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))*(a*sin(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \sin(e + f x))^{3/2} \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(3/2)*(b/cos(e + f*x))^(1/2),x)

[Out] int((a*sin(e + f*x))^(3/2)*(b/cos(e + f*x))^(1/2), x)

$$3.458 \quad \int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx$$

Optimal. Leaf size=53

$$\frac{F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}{f \sqrt{a \sin(e + fx)}}$$

[Out] $-(\sin(e+1/4*\text{Pi}+f*x)^2)^{(1/2)}/\sin(e+1/4*\text{Pi}+f*x)*\text{EllipticF}(\cos(e+1/4*\text{Pi}+f*x), 2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {2665, 2653, 2720}

$$\frac{\sqrt{\sin(2e + 2fx)} F\left(e + fx - \frac{\pi}{4} \mid 2\right) \sqrt{b \sec(e + fx)}}{f \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[b*Sec[e + f*x]]/Sqrt[a*Sin[e + f*x]],x]`

[Out] `(EllipticF[e - Pi/4 + f*x, 2]*Sqrt[b*Sec[e + f*x]]*Sqrt[Sin[2*e + 2*f*x]])/(f*Sqrt[a*Sin[e + f*x]])`

Rule 2653

`Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)])], x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

Rule 2665

`Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rubi steps

$$\begin{aligned}
\int \frac{\sqrt{b \sec(e+fx)}}{\sqrt{a \sin(e+fx)}} dx &= \left(\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \right) \int \frac{1}{\sqrt{b \cos(e+fx)} \sqrt{a \sin(e+fx)}} dx \\
&= \frac{\left(\sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)} \right) \int \frac{1}{\sqrt{\sin(2e+2fx)}} dx}{\sqrt{a \sin(e+fx)}} \\
&= \frac{F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}{f \sqrt{a \sin(e+fx)}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.28, size = 66, normalized size = 1.25

$$\frac{\cot(e+fx) {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e+fx)\right) \sqrt{b \sec(e+fx)} (-\tan^2(e+fx))^{3/4}}{f \sqrt{a \sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]/Sqrt[a*Sin[e + f*x]],x]

[Out] (Cot[e + f*x]*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*Sqrt[b*Sec[e + f*x]]*(-Tan[e + f*x]^2)^(3/4))/(f*Sqrt[a*Sin[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(73) = 146.

time = 0.25, size = 153, normalized size = 2.89

method	result
default	$ -\frac{\sqrt{\frac{b}{\cos(fx+e)}} (\sin^2(fx+e)) \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \text{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)}{f \sqrt{a \sin(fx+e)} (-1+\cos(fx+e))} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/f*(b/cos(f*x+e))^(1/2)*sin(f*x+e)^2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))/(a*sin(f*x+e))^(1/2)/(-1+cos(f*x+e))*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(b*sec(f*x + e))/sqrt(a*sin(f*x + e)), x)

Fricas [C] Result contains complex when optimal does not.

time = 0.16, size = 61, normalized size = 1.15

$$\frac{\sqrt{iab} \operatorname{ellipticF}(\cos(fx + e) + i \sin(fx + e), -1) + \sqrt{-iab} \operatorname{ellipticF}(\cos(fx + e) - i \sin(fx + e), -1)}{af}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(1/2),x, algorithm="fricas")

[Out] -(sqrt(I*a*b)*ellipticF(cos(f*x + e) + I*sin(f*x + e), -1) + sqrt(-I*a*b)*ellipticF(cos(f*x + e) - I*sin(f*x + e), -1))/(a*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**(1/2)/(a*sin(f*x+e))**(1/2),x)

[Out] Integral(sqrt(b*sec(e + f*x))/sqrt(a*sin(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(b*sec(f*x + e))/sqrt(a*sin(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\frac{b}{\cos(e + fx)}}}{\sqrt{a \sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(1/2),x)

[Out] int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(1/2), x)

$$3.459 \quad \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{5/2}} dx$$

Optimal. Leaf size=95

$$\frac{2b}{3af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} + \frac{2F(e - \frac{\pi}{4} + fx | 2) \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}{3a^2 f \sqrt{a \sin(e + fx)}}$$

[Out] $-2/3*b/a/f/(a*\sin(f*x+e))^{(3/2)}/(b*\sec(f*x+e))^{(1/2)}-2/3*(\sin(e+1/4*\text{Pi}+f*x)^2)^{(1/2)}/\sin(e+1/4*\text{Pi}+f*x)*\text{EllipticF}(\cos(e+1/4*\text{Pi}+f*x),2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)*\sin(2*f*x+2*e)^{(1/2)}/a^2/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2664, 2665, 2653, 2720}

$$\frac{2\sqrt{\sin(2e + 2fx)} F(e + fx - \frac{\pi}{4} | 2) \sqrt{b \sec(e + fx)}}{3a^2 f \sqrt{a \sin(e + fx)}} - \frac{2b}{3af (a \sin(e + fx))^{3/2} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[b*\text{Sec}[e + f*x]]/(a*\text{Sin}[e + f*x])^{(5/2)}, x]$

[Out] $(-2*b)/(3*a*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(3/2)}) + (2*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(3*a^2*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)]])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f, x\}$

Rule 2664

$\text{Int}(((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_)}), x_Symbol] \rightarrow \text{Simp}[b*(a*\text{Sin}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)}/(a*f*(m + 1))), x] + \text{Dist}[(m - n + 2)/(a^2*(m + 1)), \text{Int}[(a*\text{Sin}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, n\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2665

$\text{Int}(((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_)}), x_Symbol] \rightarrow \text{Dist}[(b*\text{Cos}[e + f*x])^n*(b*\text{Sec}[e + f*x])^n, \text{Int}[(a*\text{Sin}[e +$

$f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x\} \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] :> \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{5/2}} dx &= -\frac{2b}{3af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} + \frac{2 \int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx}{3a^2} \\ &= -\frac{2b}{3af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} + \frac{(2 \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{3a^2} \\ &= -\frac{2b}{3af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} + \frac{(2 \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)})}{3a^2 \sqrt{a \sin(e + fx)}} \\ &= -\frac{2b}{3af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} + \frac{2F(e - \frac{\pi}{4} + fx | 2) \sqrt{b \sec(e + fx)}}{3a^2 f \sqrt{a \sin(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.33, size = 75, normalized size = 0.79

$$\frac{2 \cot(e + fx) \sqrt{b \sec(e + fx)} \left(-1 + {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4} \right)}{3a^2 f \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(5/2),x]

[Out] (2*Cot[e + f*x]*Sqrt[b*Sec[e + f*x]]*(-1 + Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4)))/(3*a^2*f*Sqrt[a*Sin[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 277 vs. 2(106) = 212.

time = 0.24, size = 278, normalized size = 2.93

method	result
default	$-\frac{\left(-2 \cos(fx+e) \sin(fx+e) \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/3/f*(-2*cos(f*x+e)*sin(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)+cos(f*x+e)*2^(1/2))*sin(f*x+e)*(b/cos(f*x+e))^(1/2)/(a*sin(f*x+e))^(5/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(5/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(5/2), x)
```

Fricas [C] Result contains complex when optimal does not.

time = 0.12, size = 133, normalized size = 1.40

$$\frac{2 \left(\sqrt{iab} (\cos(fx+e)^2 - 1) \operatorname{ellipticF}(\cos(fx+e) + i \sin(fx+e), -1) + \sqrt{-iab} (\cos(fx+e)^2 - 1) \operatorname{ellipticF}(\cos(fx+e) - i \sin(fx+e), -1) - \sqrt{a \sin(fx+e)} \sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e) \right)}{3 (a^3 f \cos(fx+e)^2 - a^3 f)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(5/2),x, algorithm="fricas")
```

```
[Out] -2/3*(sqrt(I*a*b)*(cos(f*x + e)^2 - 1)*ellipticF(cos(f*x + e) + I*sin(f*x + e), -1) + sqrt(-I*a*b)*(cos(f*x + e)^2 - 1)*ellipticF(cos(f*x + e) - I*sin(f*x + e), -1) - sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e))*cos(f*x + e))/(a^3*f*cos(f*x + e)^2 - a^3*f)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**(1/2)/(a*sin(f*x+e))**(5/2),x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 6438 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(5/2),x, algorithm="giac")`

[Out] `integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(5/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(e + f x)}}}{(a \sin(e + f x))^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(5/2),x)`

[Out] `int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(5/2), x)`

$$3.460 \quad \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{9/2}} dx$$

Optimal. Leaf size=130

$$\frac{2b}{7af\sqrt{b\sec(e+fx)}(a\sin(e+fx))^{7/2}} - \frac{4b}{7a^3f\sqrt{b\sec(e+fx)}(a\sin(e+fx))^{3/2}} + \frac{4F(e - \frac{\pi}{4} + fx|2)\sqrt{b\sec(e+fx)}}{7a^4f\sqrt{a\sin(e+fx)}}$$

[Out] $-2/7*b/a/f/(a*\sin(f*x+e))^{(7/2)}/(b*\sec(f*x+e))^{(1/2)}-4/7*b/a^3/f/(a*\sin(f*x+e))^{(3/2)}/(b*\sec(f*x+e))^{(1/2)}-4/7*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticF}(\cos(e+1/4*Pi+f*x),2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)*\sin(2*f*x+2*e)^{(1/2)}/a^4/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2664, 2665, 2653, 2720}

$$\frac{4\sqrt{\sin(2e+2fx)}F(e+fx-\frac{\pi}{4}|2)\sqrt{b\sec(e+fx)}}{7a^4f\sqrt{a\sin(e+fx)}} - \frac{4b}{7a^3f(a\sin(e+fx))^{3/2}\sqrt{b\sec(e+fx)}} - \frac{2b}{7af(a\sin(e+fx))^{7/2}\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(9/2),x]

[Out] $(-2*b)/(7*a*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(7/2)}) - (4*b)/(7*a^3*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(3/2)}) + (4*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(7*a^4*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] := Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{9/2}} dx &= -\frac{2b}{7af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} + \frac{6 \int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{5/2}} dx}{7a^2} \\ &= -\frac{2b}{7af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} - \frac{4b}{7a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^3} \\ &= -\frac{2b}{7af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} - \frac{4b}{7a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^3} \\ &= -\frac{2b}{7af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} - \frac{4b}{7a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^3} \\ &= -\frac{2b}{7af \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} - \frac{4b}{7a^3 f \sqrt{b \sec(e + fx)} (a \sin(e + fx))^3} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.76, size = 111, normalized size = 0.85

$$\frac{2 \cos(2(e + fx))(b \sec(e + fx))^{3/2} \left((-2 + \cos(2(e + fx))) \csc^2(e + fx) + {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4} \right)}{7a^3 b f (-2 + \sec^2(e + fx)) (a \sin(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[b*Sec[e + f*x]]/(a*Sin[e + f*x])^(9/2),x]
```

```
[Out] (-2*Cos[2*(e + f*x)]*(b*Sec[e + f*x])^(3/2)*((-2 + Cos[2*(e + f*x)])*Csc[e + f*x]^2 + 2*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4))/(7*a^3*b*f*(-2 + Sec[e + f*x]^2)*(a*Sin[e + f*x])^(3/2))
```

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 531 vs. $2(135) = 270$.

time = 0.26, size = 532, normalized size = 4.09

method	result
default	$\frac{\left(-4\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}, \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(9/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{7} \frac{1}{f} \left(-4 \left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \operatorname{EllipticF}\left(\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}, \frac{1}{2} \sqrt{2} \right) \sin(fx+e) \cos(fx+e)^3 - 4 \left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \operatorname{EllipticF}\left(\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}, \frac{1}{2} \sqrt{2} \right) \sin(fx+e) \cos(fx+e)^2 + 4 \cos(fx+e) \sin(fx+e) \left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \operatorname{EllipticF}\left(\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}, \frac{1}{2} \sqrt{2} \right) + 4 \left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \left(\frac{-1+\cos(fx+e)}{\sin(fx+e)} \right)^{1/2} \operatorname{EllipticF}\left(\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}, \frac{1}{2} \sqrt{2} \right) \sin(fx+e) + 2 \cos(fx+e)^3 \sqrt{2} - 3 \cos(fx+e)^2 \sqrt{2} \right) \left(\frac{b}{\cos(fx+e)} \right)^{1/2} \sin(fx+e) / \left(a \sin(fx+e) \right)^{9/2} \sqrt{2} \right)$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(9/2),x,algorithm="maxima")`

[Out] `integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(9/2), x)`

Fricas [C] Result contains complex when optimal does not.

time = 0.12, size = 185, normalized size = 1.42

$$\frac{2 \left(2 (\cos(fx+e)^4 - 2 \cos(fx+e)^2 + 1) \sqrt{1-ab} \operatorname{ellipticF}(\cos(fx+e) + i \sin(fx+e), -1) + 2 (\cos(fx+e)^4 - 2 \cos(fx+e)^2 + 1) \sqrt{-1-ab} \operatorname{ellipticF}(\cos(fx+e) - i \sin(fx+e), -1) - (2 \cos(fx+e)^3 - 3 \cos(fx+e)) \sqrt{a \sin(fx+e)} \sqrt{\frac{b}{\cos(fx+e)}} \right)}{7 (a^2 f \cos(fx+e)^3 - 2 a^2 f \cos(fx+e)^2 + a^2 f)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(9/2),x,algorithm="fricas")`

[Out]
$$-2/7 * (2 * (\cos(fx + e))^4 - 2 * \cos(fx + e)^2 + 1) * \sqrt{I * a * b} * \operatorname{ellipticF}(\cos(fx + e) + I * \sin(fx + e), -1) + 2 * (\cos(fx + e))^4 - 2 * \cos(fx + e)^2 + 1) * s$$

```

qrt(-I*a*b)*ellipticF(cos(f*x + e) - I*sin(f*x + e), -1) - (2*cos(f*x + e)^
3 - 3*cos(f*x + e))*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e)))/(a^5*f*cos(f
*x + e)^4 - 2*a^5*f*cos(f*x + e)^2 + a^5*f)

```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))**(1/2)/(a*sin(f*x+e))**(9/2),x)
```

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(9/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(b*sec(f*x + e))/(a*sin(f*x + e))^(9/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{b}{\cos(e + f x)}}}{(a \sin(e + f x))^{9/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(9/2),x)
```

```
[Out] int((b/cos(e + f*x))^(1/2)/(a*sin(e + f*x))^(9/2), x)
```

$$3.461 \quad \int \frac{\sin^{\frac{9}{2}}(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=115

$$-\frac{7b \sin^{\frac{3}{2}}(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7E(e - \frac{\pi}{4} + fx|2) \sqrt{\sin(e+fx)}}{20f \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}$$

[Out] $-7/30*b*\sin(f*x+e)^{(3/2)}/f/(b*\sec(f*x+e))^{(3/2)}-1/5*b*\sin(f*x+e)^{(7/2)}/f/(b*\sec(f*x+e))^{(3/2)}-7/20*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticE}(\cos(e+1/4*Pi+f*x),2^{(1/2)})*\sin(f*x+e)^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}/\sin(2*f*x+2*e)^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {2663, 2665, 2652, 2719}

$$-\frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} - \frac{7b \sin^{\frac{3}{2}}(e+fx)}{30f(b \sec(e+fx))^{3/2}} + \frac{7\sqrt{\sin(e+fx)} E(e+fx - \frac{\pi}{4}|2)}{20f \sqrt{\sin(2e+2fx)} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^(9/2)/Sqrt[b*Sec[e + f*x]],x]

[Out] $(-7*b*\sin[e + f*x]^{(3/2)})/(30*f*(b*\sec[e + f*x])^{(3/2)}) - (b*\sin[e + f*x]^{(7/2)})/(5*f*(b*\sec[e + f*x])^{(3/2)}) + (7*\text{EllipticE}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(20*f*\text{Sqrt}[b*\sec[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])$

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]] , x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2663

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[(-a)*b*(a*Sin[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*Sin[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +

$f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \ \&\& \ \text{IntegerQ}[m - 1/2] \ \&\& \ \text{IntegerQ}[n - 1/2]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x_Symbol] \ :> \ \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\sin^{\frac{9}{2}}(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7}{10} \int \frac{\sin^{\frac{5}{2}}(e+fx)}{\sqrt{b \sec(e+fx)}} dx \\ &= -\frac{7b \sin^{\frac{3}{2}}(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7}{20} \int \frac{\sqrt{\sin(e+fx)}}{\sqrt{b \sec(e+fx)}} dx \\ &= -\frac{7b \sin^{\frac{3}{2}}(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7 \int \sqrt{b \cos(e+fx)} \sqrt{\sin(e+fx)}}{20 \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} dx \\ &= -\frac{7b \sin^{\frac{3}{2}}(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{(7 \sqrt{\sin(e+fx)}) \int \sqrt{\sin(2e+2fx)}}{20 \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}} dx \\ &= -\frac{7b \sin^{\frac{3}{2}}(e+fx)}{30f(b \sec(e+fx))^{3/2}} - \frac{b \sin^{\frac{7}{2}}(e+fx)}{5f(b \sec(e+fx))^{3/2}} + \frac{7E(e - \frac{\pi}{4} + fx|2) \sqrt{\sin(e+fx)}}{20f \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.39, size = 86, normalized size = 0.75

$$\frac{b \left(23 - 26 \cos(2(e+fx)) + 3 \cos(4(e+fx)) + 42 {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \sec^2(e+fx)\right) \sqrt[4]{-\tan^2(e+fx)} \right)}{120f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^(9/2)/Sqrt[b*Sec[e + f*x]],x]

[Out] -1/120*(b*(23 - 26*Cos[2*(e + f*x)] + 3*Cos[4*(e + f*x)] + 42*Hypergeometric2F1[-1/2, 1/4, 1/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(1/4)))/(f*(b*Sec[e + f*x])^(3/2)*Sqrt[Sin[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 523 vs. 2(120) = 240.

time = 0.37, size = 524, normalized size = 4.56

method	result
default	$-\frac{\left(12\sqrt{2}(\cos^6(fx+e))+42\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\right)}{\text{EllipticE}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/120/f*(12*2^(1/2)*cos(f*x+e)^6+42*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)-21*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)-38*cos(f*x+e)^4*2^(1/2)+42*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-21*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+47*cos(f*x+e)^2*2^(1/2)-21*cos(f*x+e)^2*(1/2))/sin(f*x+e)^(1/2)/(b/cos(f*x+e))^(1/2)/cos(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(f*x + e)^(9/2)/sqrt(b*sec(f*x + e)), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] integral((cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(b*sec(f*x + e))*sqrt(sin(f*x + e))/(b*sec(f*x + e)), x)
```


Sympy [F(-1)] Timed out
 time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**(9/2)/(b*sec(f*x+e))**(1/2), x)

[Out] Timed out

Giac [F]
 time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2), x, algorithm="giac")

[Out] integrate(sin(f*x + e)^(9/2)/sqrt(b*sec(f*x + e)), x)

Mupad [F]
 time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^{9/2}}{\sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^(9/2)/(b/cos(e + f*x))^(1/2), x)

[Out] int(sin(e + f*x)^(9/2)/(b/cos(e + f*x))^(1/2), x)

$$3.462 \quad \int \frac{\sin^{\frac{5}{2}}(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=85

$$-\frac{b \sin^{\frac{3}{2}}(e+fx)}{3f(b \sec(e+fx))^{3/2}} + \frac{E(e - \frac{\pi}{4} + fx|2) \sqrt{\sin(e+fx)}}{2f \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}$$

[Out] $-1/3*b*\sin(f*x+e)^{(3/2)}/f/(b*\sec(f*x+e))^{(3/2)}-1/2*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticE}(\cos(e+1/4*Pi+f*x),2^{(1/2)})*\sin(f*x+e)^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}/\sin(2*f*x+2*e)^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 85, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {2663, 2665, 2652, 2719}

$$\frac{\sqrt{\sin(e+fx)} E(e+fx - \frac{\pi}{4}|2)}{2f \sqrt{\sin(2e+2fx)} \sqrt{b \sec(e+fx)}} - \frac{b \sin^{\frac{3}{2}}(e+fx)}{3f(b \sec(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^(5/2)/Sqrt[b*Sec[e + f*x]],x]

[Out] $-1/3*(b*\sin[e + f*x]^{(3/2)})/(f*(b*\sec[e + f*x])^{(3/2)}) + (\text{EllipticE}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[\sin[e + f*x]])/(2*f*\text{Sqrt}[b*\sec[e + f*x]]*\text{Sqrt}[\sin[2*e + 2*f*x]])$

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_)]] , x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2663

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Simp[(-a)*b*(a*Sin[e + f*x])^(m-1)*((b*Sec[e + f*x])^(n-1)/(f*(m-n))), x] + Dist[a^2*((m-1)/(m-n)), Int[(a*Sin[e + f*x])^(m-2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m-n, 0] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +

$f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{\sin^{\frac{5}{2}}(e + fx)}{\sqrt{b \sec(e + fx)}} dx &= -\frac{b \sin^{\frac{3}{2}}(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{1}{2} \int \frac{\sqrt{\sin(e + fx)}}{\sqrt{b \sec(e + fx)}} dx \\ &= -\frac{b \sin^{\frac{3}{2}}(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{\int \sqrt{b \cos(e + fx)} \sqrt{\sin(e + fx)} dx}{2\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} \\ &= -\frac{b \sin^{\frac{3}{2}}(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{\sqrt{\sin(e + fx)} \int \sqrt{\sin(2e + 2fx)} dx}{2\sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}} \\ &= -\frac{b \sin^{\frac{3}{2}}(e + fx)}{3f(b \sec(e + fx))^{3/2}} + \frac{E(e - \frac{\pi}{4} + fx | 2) \sqrt{\sin(e + fx)}}{2f \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.22, size = 74, normalized size = 0.87

$$\frac{b \left(-1 + \cos(2(e + fx)) - 3 {}_2F_1 \left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \sec^2(e + fx) \right) \sqrt[4]{-\tan^2(e + fx)} \right)}{6f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sin[e + f*x]^(5/2)/Sqrt[b*Sec[e + f*x]],x]

[Out] (b*(-1 + Cos[2*(e + f*x)] - 3*Hypergeometric2F1[-1/2, 1/4, 1/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(1/4)))/(6*f*(b*Sec[e + f*x])^(3/2)*Sqrt[Sin[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 510 vs. 2(96) = 192.

time = 0.29, size = 511, normalized size = 6.01

method	result
--------	--------

default	$-\left(6\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\text{EllipticE}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}, \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\right)\right)$
---------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/12/f*(6*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticE}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})*\cos(f*x+e)-3*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})*\cos(f*x+e)-2*\cos(f*x+e)^4*2^{1/2}+6*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticE}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})-3*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))+5*\cos(f*x+e)^2*2^{1/2}-3*\cos(f*x+e)*2^{1/2}))/\sin(f*x+e)^{1/2}/(b/\cos(f*x+e))^{1/2}/\cos(f*x+e)*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(sin(f*x + e)^(5/2)/sqrt(b*sec(f*x + e)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] `integral(-(cos(f*x + e))^2 - 1)*sqrt(b*sec(f*x + e))*sqrt(sin(f*x + e))/(b*sec(f*x + e)), x)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**(5/2)/(b*sec(f*x+e))**(1/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sin(f*x + e)^(5/2)/sqrt(b*sec(f*x + e)), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^{5/2}}{\sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^(5/2)/(b/cos(e + f*x))^(1/2),x)`

[Out] `int(sin(e + f*x)^(5/2)/(b/cos(e + f*x))^(1/2), x)`

$$3.463 \quad \int \frac{\sqrt{\sin(e + fx)}}{\sqrt{b \sec(e + fx)}} dx$$

Optimal. Leaf size=51

$$\frac{E\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{\sin(e + fx)}}{f \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}$$

[Out] $-(\sin(e+1/4\pi+fx)^2)^{(1/2)}/\sin(e+1/4\pi+fx)*\text{EllipticE}(\cos(e+1/4\pi+fx), 2^{(1/2)})*\sin(f*x+e)^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}/\sin(2*f*x+2*e)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {2665, 2652, 2719}

$$\frac{\sqrt{\sin(e + fx)} E\left(e + fx - \frac{\pi}{4} \mid 2\right)}{f \sqrt{\sin(2e + 2fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[Sin[e + f*x]]/Sqrt[b*Sec[e + f*x]],x]

[Out] (EllipticE[e - Pi/4 + f*x, 2]*Sqrt[Sin[e + f*x]])/(f*Sqrt[b*Sec[e + f*x]]*Sqrt[Sin[2*e + 2*f*x]])

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]] , x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_.)]] , x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\int \frac{\sqrt{\sin(e+fx)}}{\sqrt{b \sec(e+fx)}} dx = \frac{\int \frac{\sqrt{b \cos(e+fx)} \sqrt{\sin(e+fx)}}{\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} dx}{\sqrt{\sin(e+fx)} \int \frac{\sqrt{\sin(2e+2fx)}}{\sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}} dx}$$

$$= \frac{E(e - \frac{\pi}{4} + fx | 2) \sqrt{\sin(e+fx)}}{f \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.81, size = 60, normalized size = 1.18

$$\frac{{}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \sec^2(e+fx)\right) \sqrt[4]{-\tan^2(e+fx)}}{f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[Sin[e + f*x]]/Sqrt[b*Sec[e + f*x]],x]

[Out] -((b*Hypergeometric2F1[-1/2, 1/4, 1/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(1/4))/(f*(b*Sec[e + f*x])^(3/2)*Sqrt[Sin[e + f*x]]))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 496 vs. 2(71) = 142.

time = 0.29, size = 497, normalized size = 9.75

method	result
default	$-\frac{\left(2 \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \text{EllipticE}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{f \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] -1/2/f*(2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)-((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)+2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))

$$\int \frac{\sqrt{\sin(fx+e)}}{\sqrt{b \sec(fx+e)}} dx$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(sin(f*x + e))/sqrt(b*sec(f*x + e)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(b*sec(f*x + e))*sqrt(sin(f*x + e))/(b*sec(f*x + e)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\sin(e + fx)}}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**(1/2)/(b*sec(f*x+e))**(1/2),x)

[Out] Integral(sqrt(sin(e + f*x))/sqrt(b*sec(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(sin(f*x + e))/sqrt(b*sec(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{\sin(e + f x)}}{\sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^(1/2)/(b/cos(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^(1/2)/(b/cos(e + f*x))^(1/2), x)

$$3.464 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{3}{2}}(e + fx)} dx$$

Optimal. Leaf size=81

$$-\frac{2b}{f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}} - \frac{2E(e - \frac{\pi}{4} + fx | 2) \sqrt{\sin(e + fx)}}{f \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}$$

[Out] $-2*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(1/2)}+2*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticE}(\cos(e+1/4*Pi+f*x),2^{(1/2)})*\sin(f*x+e)^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}/\sin(2*f*x+2*e)^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {2664, 2665, 2652, 2719}

$$-\frac{2b}{f \sqrt{\sin(e + fx)} (b \sec(e + fx))^{3/2}} - \frac{2 \sqrt{\sin(e + fx)} E(e + fx - \frac{\pi}{4} | 2)}{f \sqrt{\sin(2e + 2fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(3/2)),x]

[Out] $(-2*b)/(f*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sqrt}[\text{Sin}[e + f*x]]) - (2*\text{EllipticE}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])$

Rule 2652

Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte

gerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{3}{2}}(e+fx)} dx &= -\frac{2b}{f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}} - 2 \int \frac{\sqrt{\sin(e+fx)}}{\sqrt{b \sec(e+fx)}} dx \\ &= -\frac{2b}{f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}} - \frac{2 \int \sqrt{b \cos(e+fx)} \sqrt{\sin(e+fx)}}{\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} dx \\ &= -\frac{2b}{f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}} - \frac{(2 \sqrt{\sin(e+fx)}) \int \sqrt{\sin(2e+2fx)}}{\sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}} dx \\ &= -\frac{2b}{f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}} - \frac{2E(e - \frac{\pi}{4} + fx | 2) \sqrt{\sin(e+fx)}}{f \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.19, size = 63, normalized size = 0.78

$$\frac{2b \left(-1 + {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \sec^2(e+fx)\right) \sqrt[4]{-\tan^2(e+fx)} \right)}{f(b \sec(e+fx))^{3/2} \sqrt{\sin(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(3/2)),x]

[Out] (2*b*(-1 + Hypergeometric2F1[-1/2, 1/4, 1/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(1/4)))/(f*(b*Sec[e + f*x])^(3/2)*Sqrt[Sin[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 481 vs. 2(96) = 192.

time = 0.21, size = 482, normalized size = 5.95

method	result
--------	--------

default	$-\frac{\left(-2\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\right)\text{EllipticE}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)}{\dots}$
---------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/f*(-2*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticE}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})*\cos(f*x+e)+((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})*\cos(f*x+e)-2*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticE}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))+((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))+\cos(f*x+e)*2^{1/2})/\cos(f*x+e)/\sin(f*x+e)^{1/2}/(b/\cos(f*x+e))^{1/2}*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(3/2)), x)`

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] Exception raised: TypeError >> Symbolic function `elliptic_ec` takes exactly 1 arguments (2 given)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{3}{2}}(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)**(3/2)/(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(1/(sqrt(b*sec(e + f*x))*sin(e + f*x)**(3/2)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(3/2)), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^{3/2} \sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^(3/2)*(b/cos(e + f*x))^(1/2)),x)`

[Out] `int(1/(sin(e + f*x)^(3/2)*(b/cos(e + f*x))^(1/2)), x)`

$$3.465 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{7}{2}}(e + fx)} dx$$

Optimal. Leaf size=115

$$\frac{2b}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)} - \frac{4b}{5f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}} - \frac{4E(e - \frac{\pi}{4} + fx | 2) \sqrt{\sin(e + fx)}}{5f \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}$$

[Out] $-2/5*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(5/2)}-4/5*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(1/2)}+4/5*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticE}(\cos(e+1/4*Pi+f*x), 2^{(1/2)})*\sin(f*x+e)^{(1/2)}/f/(b*\sec(f*x+e))^{(1/2)}/\sin(2*f*x+2*e)^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.174$, Rules used = {2664, 2665, 2652, 2719}

$$\frac{2b}{5f \sin^{\frac{5}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{4b}{5f \sqrt{\sin(e + fx)} (b \sec(e + fx))^{3/2}} - \frac{4 \sqrt{\sin(e + fx)} E(e + fx - \frac{\pi}{4} | 2)}{5f \sqrt{\sin(2e + 2fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(7/2)),x]`

[Out] $(-2*b)/(5*f*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^{(5/2)}) - (4*b)/(5*f*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sqrt}[\text{Sin}[e + f*x]]) - (4*\text{EllipticE}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(5*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])$

Rule 2652

`Int[Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]], x_Symbol] := Dist[Sqrt[a*Sin[e + f*x]]*(Sqrt[b*Cos[e + f*x]]/Sqrt[Sin[2*e + 2*f*x]]), Int[Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]`

Rule 2664

`Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/(a*f*(m + 1))), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]`

Rule 2665

`Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +`

$f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_.) + (d_.)*(x_.)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{7}{2}}(e + fx)} dx &= -\frac{2b}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)} + \frac{2}{5} \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{3}{2}}(e + fx)} dx \\ &= -\frac{2b}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)} - \frac{4b}{5f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}} \\ &= -\frac{2b}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)} - \frac{4b}{5f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}} \\ &= -\frac{2b}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)} - \frac{4b}{5f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}} \\ &= -\frac{2b}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)} - \frac{4b}{5f(b \sec(e + fx))^{3/2} \sqrt{\sin(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.35, size = 82, normalized size = 0.71

$$\frac{2b \left(-2 + \cos(2(e + fx)) + 2 {}_2F_1\left(-\frac{1}{2}, \frac{1}{4}; \frac{1}{2}; \sec^2(e + fx)\right) \sin^2(e + fx) \sqrt[4]{-\tan^2(e + fx)} \right)}{5f(b \sec(e + fx))^{3/2} \sin^{\frac{5}{2}}(e + fx)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(7/2)),x]

[Out] (2*b*(-2 + Cos[2*(e + f*x)] + 2*Hypergeometric2F1[-1/2, 1/4, 1/2, Sec[e + f*x]^2]*Sin[e + f*x]^2*(-Tan[e + f*x]^2)^(1/4)))/(5*f*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^(5/2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 1029 vs. 2(120) = 240.

time = 0.23, size = 1030, normalized size = 8.96

method	result	size
default	Expression too large to display	1030

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/sin(f*x+e)^(7/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 16/5/f*(-1+cos(f*x+e))^4*(4*cos(f*x+e)^3*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-2*cos(f*x+e)^3*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+4*cos(f*x+e)^2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-2*cos(f*x+e)^2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-4*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)+2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)-2*cos(f*x+e)^3*2^(1/2)-4*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+cos(f*x+e)^2*2^(1/2)+2*cos(f*x+e)*2^(1/2)/sin(f*x+e)^(5/2)/(b/cos(f*x+e))^(1/2)/(sin(f*x+e)-1+cos(f*x+e))/(-1+cos(f*x+e)-sin(f*x+e))/(sin(f*x+e)^2+cos(f*x+e)^2-2*cos(f*x+e)+1)^3*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(f*x+e)^(7/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(7/2)), x)
```

Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(f*x+e)^(7/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] Exception raised: TypeError >> Symbolic function elliptic_ec takes exactly 1 arguments (2 given)
```

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(f*x+e)**(7/2)/(b*sec(f*x+e))**(1/2),x)
```

```
[Out] Timed out
```

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/sin(f*x+e)^(7/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(7/2)), x)
```

Mupad [F]
time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^{7/2} \sqrt{\frac{b}{\cos(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(sin(e + f*x)^(7/2)*(b/cos(e + f*x))^(1/2)),x)
```

```
[Out] int(1/(sin(e + f*x)^(7/2)*(b/cos(e + f*x))^(1/2)), x)
```

$$3.466 \quad \int \frac{\sin^3(e+fx)}{\sqrt{b \sec(e+fx)}} dx$$

Optimal. Leaf size=363

$$\frac{\sqrt{b} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e+fx)}}{\sqrt{b} \sqrt{\sin(e+fx)}} \right)}{4\sqrt{2} f \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\sqrt{b} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b \cos(e+fx)}}{\sqrt{b} \sqrt{\sin(e+fx)}} \right)}{4\sqrt{2} f \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\sqrt{b} \log \left(\sqrt{b} + \sqrt{b} \cot \right)}{8\sqrt{2} f \sqrt{b \cos(e+fx)}}$$

[Out] 1/8*arctan(1-2^(1/2)*(b*cos(f*x+e))^(1/2)/b^(1/2)/sin(f*x+e)^(1/2))*b^(1/2)/f*2^(1/2)/(b*cos(f*x+e))^(1/2)/(b*sec(f*x+e))^(1/2)-1/8*arctan(1+2^(1/2)*(b*cos(f*x+e))^(1/2)/b^(1/2)/sin(f*x+e)^(1/2))*b^(1/2)/f*2^(1/2)/(b*cos(f*x+e))^(1/2)/(b*sec(f*x+e))^(1/2)-1/16*ln(b^(1/2)+cot(f*x+e)*b^(1/2)-2^(1/2)*(b*cos(f*x+e))^(1/2)/sin(f*x+e)^(1/2))*b^(1/2)/f*2^(1/2)/(b*cos(f*x+e))^(1/2)/(b*sec(f*x+e))^(1/2)+1/16*ln(b^(1/2)+cot(f*x+e)*b^(1/2)+2^(1/2)*(b*cos(f*x+e))^(1/2)/sin(f*x+e)^(1/2))*b^(1/2)/f*2^(1/2)/(b*cos(f*x+e))^(1/2)/(b*sec(f*x+e))^(1/2)-1/2*b*sin(f*x+e)^(1/2)/f/(b*sec(f*x+e))^(3/2)

Rubi [A]

time = 0.18, antiderivative size = 363, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.391$, Rules used = {2663, 2665, 2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{b} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e+fx)}}{\sqrt{b} \sqrt{\sin(e+fx)}} \right)}{4\sqrt{2} f \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\sqrt{b} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b \cos(e+fx)}}{\sqrt{b} \sqrt{\sin(e+fx)}} + 1 \right)}{4\sqrt{2} f \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{b \sqrt{\sin(e+fx)}}{2f(b \sec(e+fx))^{3/2}} - \frac{\sqrt{b} \log \left(\sqrt{b} \cot(e+fx) - \frac{\sqrt{2} \sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}} + \sqrt{b} \right)}{8\sqrt{2} f \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} + \frac{\sqrt{b} \log \left(\sqrt{b} \cot(e+fx) + \frac{\sqrt{2} \sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}} + \sqrt{b} \right)}{8\sqrt{2} f \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^(3/2)/Sqrt[b*Sec[e + f*x]],x]

[Out] (Sqrt[b]*ArcTan[1 - (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/(Sqrt[b]*Sqrt[Sin[e + f*x]])]/(4*Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) - (Sqrt[b]*ArcTan[1 + (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/(Sqrt[b]*Sqrt[Sin[e + f*x]])]/(4*Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) - (Sqrt[b]*Log[Sqrt[b] + Sqrt[b]*Cot[e + f*x] - (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[e + f*x]])/(8*Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) + (Sqrt[b]*Log[Sqrt[b] + Sqrt[b]*Cot[e + f*x] + (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[e + f*x]])/(8*Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) - (b*Sqrt[Sin[e + f*x]])/(2*f*(b*Sec[e + f*x])^(3/2))

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] & AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2655

```
Int[(cos[(e_) + (f_)*(x_)]*(a_.))^m_)*((b_)*sin[(e_) + (f_)*(x_)])^n_, x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sin[e + f*x])^(1/k)], x] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]
```

Rule 2663

```
Int[((b_)*sec[(e_) + (f_)*(x_)])^n_)*((a_)*sin[(e_) + (f_)*(x_)])^m_, x_Symbol] := Simp[(-a)*b*(a*SIN[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*SIN[e + f*x])^m,
```

- 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rubi steps

$$\begin{aligned}
 \int \frac{\sin^{\frac{3}{2}}(e+fx)}{\sqrt{b \sec(e+fx)}} dx &= -\frac{b\sqrt{\sin(e+fx)}}{2f(b \sec(e+fx))^{3/2}} + \frac{1}{4} \int \frac{1}{\sqrt{b \sec(e+fx)} \sqrt{\sin(e+fx)}} dx \\
 &= -\frac{b\sqrt{\sin(e+fx)}}{2f(b \sec(e+fx))^{3/2}} + \frac{\int \frac{\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}} dx}{4\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
 &= -\frac{b\sqrt{\sin(e+fx)}}{2f(b \sec(e+fx))^{3/2}} - \frac{b \operatorname{Subst}\left(\int \frac{x^2}{b^2+x^4} dx, x, \frac{\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}}\right)}{2f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
 &= -\frac{b\sqrt{\sin(e+fx)}}{2f(b \sec(e+fx))^{3/2}} + \frac{b \operatorname{Subst}\left(\int \frac{b-x^2}{b^2+x^4} dx, x, \frac{\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}}\right)}{4f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{b \operatorname{Subst}\left(\int \frac{b+x^2}{b^2+x^4} dx, x, \frac{\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}}\right)}{4f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
 &= -\frac{b\sqrt{\sin(e+fx)}}{2f(b \sec(e+fx))^{3/2}} - \frac{\sqrt{b} \operatorname{Subst}\left(\int \frac{\sqrt{2}\sqrt{b}+2x}{-b-\sqrt{2}\sqrt{b}x-x^2} dx, x, \frac{\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}}\right)}{8\sqrt{2}f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
 &= -\frac{\sqrt{b} \log\left(\sqrt{b} + \sqrt{b} \cot(e+fx) - \frac{\sqrt{2}\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}}\right)}{8\sqrt{2}f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} + \frac{\sqrt{b} \log\left(\sqrt{b} + \sqrt{b} \tan(e+fx) + \frac{\sqrt{2}\sqrt{b \cos(e+fx)}}{\sqrt{\sin(e+fx)}}\right)}{8\sqrt{2}f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} \\
 &= \frac{\sqrt{b} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt{b \cos(e+fx)}}{\sqrt{b}\sqrt{\sin(e+fx)}}\right)}{4\sqrt{2}f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}} - \frac{\sqrt{b} \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt{b \cos(e+fx)}}{\sqrt{b}\sqrt{\sin(e+fx)}}\right)}{4\sqrt{2}f\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.99, size = 145, normalized size = 0.40

$$\frac{b\left(-4\sin^2(e+fx) + \sqrt{2} \tan^{-1}\left(\frac{-1+\sqrt{\tan^2(e+fx)}}{\sqrt{2}\sqrt[4]{\tan^2(e+fx)}}\right) \tan^2(e+fx)^{3/4} + \sqrt{2} \tanh^{-1}\left(\frac{\sqrt{2}\sqrt[4]{\tan^2(e+fx)}}{1+\sqrt{\tan^2(e+fx)}}\right) \tan^2(e+fx)^{3/4}\right)}{8f(b \sec(e+fx))^{3/2} \sin^{\frac{3}{2}}(e+fx)}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[e + f*x]^(3/2)/Sqrt[b*Sec[e + f*x]],x]
```

```
[Out] (b*(-4*Sin[e + f*x]^2 + Sqrt[2]*ArcTan[(-1 + Sqrt[Tan[e + f*x]^2])]/(Sqrt[2]
*(Tan[e + f*x]^2)^(1/4)))*(Tan[e + f*x]^2)^(3/4) + Sqrt[2]*ArcTanh[(Sqrt[2]
*(Tan[e + f*x]^2)^(1/4))/(1 + Sqrt[Tan[e + f*x]^2])]*(Tan[e + f*x]^2)^(3/4)
))/(8*f*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^(3/2))
```

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.20, size = 648, normalized size = 1.79

method	result
default	$\left(i \sin(fx+e) \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \text{EllipticPi}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/8/f*(I*sin(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+
e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*Ellip
ticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))-I
*sin(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos
(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((
1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-sin(f*x+e
)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/s
in(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+
e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))-sin(f*x+e)*((1-cos(
f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))
^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x
+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))+2*((1-cos(f*x+e)+sin(f*x+e))/
sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x
+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/
2),1/2*2^(1/2))*sin(f*x+e)-2*cos(f*x+e)^3*2^(1/2)+2*cos(f*x+e)^2*2^(1/2))*s
in(f*x+e)^(1/2)/(-1+cos(f*x+e))/(b/cos(f*x+e))^(1/2)/cos(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(f*x + e)^(3/2)/sqrt(b*sec(f*x + e)), x)
```

Fricas [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] Timed out

Sympy [F]
time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^{\frac{3}{2}}(e + fx)}{\sqrt{b \sec(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)**(3/2)/(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(sin(e + f*x)**(3/2)/sqrt(b*sec(e + f*x)), x)`

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sin(f*x+e)^(3/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sin(f*x + e)^(3/2)/sqrt(b*sec(f*x + e)), x)`

Mupad [F]
time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sin(e + fx)^{3/2}}{\sqrt{\frac{b}{\cos(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^(3/2)/(b/cos(e + f*x))^(1/2),x)`

[Out] `int(sin(e + f*x)^(3/2)/(b/cos(e + f*x))^(1/2), x)`

$$3.467 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sqrt{\sin(e + fx)}} dx$$

Optimal. Leaf size=328

$$\frac{\sqrt{b} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right) - \sqrt{b} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right) - \sqrt{b} \log \left(\sqrt{b} + \sqrt{b} \cot \right)}{\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{\sqrt{b} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right) - \sqrt{b} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right) - \sqrt{b} \log \left(\sqrt{b} + \sqrt{b} \cot \right)}{2\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}$$

[Out] $1/2*\arctan(1-2^{(1/2)}*(b*\cos(f*x+e))^{(1/2)}/b^{(1/2)}/\sin(f*x+e)^{(1/2)})*b^{(1/2)}/f*2^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}-1/2*\arctan(1+2^{(1/2)}*(b*\cos(f*x+e))^{(1/2)}/b^{(1/2)}/\sin(f*x+e)^{(1/2)})*b^{(1/2)}/f*2^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}-1/4*\ln(b^{(1/2)}+\cot(f*x+e)*b^{(1/2)}-2^{(1/2)}*(b*\cos(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)})*b^{(1/2)}/f*2^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}+1/4*\ln(b^{(1/2)}+\cot(f*x+e)*b^{(1/2)}+2^{(1/2)}*(b*\cos(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)})*b^{(1/2)}/f*2^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}/(b*\sec(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 328, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.348$, Rules used = {2665, 2655, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{b} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right) - \sqrt{b} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} + 1 \right) - \sqrt{b} \log \left(\sqrt{b} \cot(e + fx) - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} + \sqrt{b} \right) + \sqrt{b} \log \left(\sqrt{b} \cot(e + fx) + \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} + \sqrt{b} \right)}{\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{\sqrt{b} \operatorname{ArcTan} \left(\frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} + 1 \right) - \sqrt{b} \operatorname{ArcTan} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right) - \sqrt{b} \log \left(\sqrt{b} \cot(e + fx) - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} + \sqrt{b} \right) + \sqrt{b} \log \left(\sqrt{b} \cot(e + fx) + \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} + \sqrt{b} \right)}{2\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[b*Sec[e + f*x]]*Sqrt[Sin[e + f*x]]),x]

[Out] (Sqrt[b]*ArcTan[1 - (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/(Sqrt[b]*Sqrt[Sin[e + f*x]])]/(Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) - (Sqrt[b]*ArcTan[1 + (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/(Sqrt[b]*Sqrt[Sin[e + f*x]])]/(Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) - (Sqrt[b]*Log[Sqrt[b] + Sqrt[b]*Cot[e + f*x] - (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[e + f*x]])]/(2*Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]]) + (Sqrt[b]*Log[Sqrt[b] + Sqrt[b]*Cot[e + f*x] + (Sqrt[2]*Sqrt[b*Cos[e + f*x]])/Sqrt[Sin[e + f*x]])]/(2*Sqrt[2]*f*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2655

```
Int[(cos[(e_) + (f_)*(x_)])*(a_)^(m)*((b_)*sin[(e_) + (f_)*(x_)])^(n
_), x_Symbol] := With[{k = Denominator[m]}, Dist[(-k)*a*(b/f), Subst[Int[x^
(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Cos[e + f*x])^(1/k)/(b*Sin[e
+ f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0
] && LtQ[m, 1]
```

Rule 2665

```
Int[((b_)*sec[(e_) + (f_)*(x_)])^(n)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +
f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte
```


gerQ[m - 1/2] && IntegerQ[n - 1/2]

Rubi steps

$$\begin{aligned}
 \int \frac{1}{\sqrt{b \sec(e + fx)} \sqrt{\sin(e + fx)}} dx &= \frac{\int \frac{\sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} dx}{\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
 &= \frac{(2b) \text{Subst} \left(\int \frac{x^2}{b^2 + x^4} dx, x, \frac{\sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
 &= \frac{b \text{Subst} \left(\int \frac{b - x^2}{b^2 + x^4} dx, x, \frac{\sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{b \text{Subst} \left(\int \frac{b + x^2}{b^2 + x^4} dx, x, \frac{\sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
 &= \frac{\sqrt{b} \text{Subst} \left(\int \frac{\sqrt{2} \sqrt{b} + 2x}{-b - \sqrt{2} \sqrt{b} x - x^2} dx, x, \frac{\sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{2\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{\sqrt{b} \text{Subst} \left(\int \frac{\sqrt{2} \sqrt{b} - 2x}{-b - \sqrt{2} \sqrt{b} x - x^2} dx, x, \frac{\sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{2\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
 &= \frac{\sqrt{b} \log \left(\sqrt{b} + \sqrt{b} \cot(e + fx) - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{2\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} + \frac{\sqrt{b} \log \left(\sqrt{b} - \sqrt{b} \cot(e + fx) - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{\sin(e + fx)}} \right)}{2\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} \\
 &= \frac{\sqrt{b} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right)}{\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}} - \frac{\sqrt{b} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b \cos(e + fx)}}{\sqrt{b} \sqrt{\sin(e + fx)}} \right)}{\sqrt{2} f \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}
 \end{aligned}$$

Mathematica [A]

time = 0.52, size = 113, normalized size = 0.34

$$\frac{b \left(\tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt[4]{\tan^2(e + fx)}} \right) + \tanh^{-1} \left(\frac{\sqrt{2} \sqrt[4]{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \right) \tan^2(e + fx)^{3/4}}{\sqrt{2} f (b \sec(e + fx))^{3/2} \sin^{3/2}(e + fx)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sqrt[Sin[e + f*x]]),x]

[Out] (b*(ArcTan[(-1 + Sqrt[Tan[e + f*x]^2)]/(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))]) + ArcTanh[(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))/(1 + Sqrt[Tan[e + f*x]^2])])*(Tan[e + f*x]^2)^(3/4))/(Sqrt[2]*f*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^(3/2))

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.23, size = 304, normalized size = 0.93

method	result
default	$-\frac{\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \left(i \operatorname{EllipticPi} \left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \right) \right)}{1}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/2/f*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*(I*\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2-1/2*I,1/2*2^{1/2})-I*\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2+1/2*I,1/2*2^{1/2})-2*\operatorname{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})+\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2-1/2*I,1/2*2^{1/2})+\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2+1/2*I,1/2*2^{1/2}))*\sin(f*x+e)^{3/2}/(b/\cos(f*x+e))^{1/2}/\cos(f*x+e)/(-1+\cos(f*x+e))*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(1/(sqrt(b*sec(f*x + e))*sqrt(sin(f*x + e))), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{b \sec(e + fx)} \sqrt{\sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)**(1/2)/(b*sec(f*x+e))**(1/2),x)`

[Out] `Integral(1/(sqrt(b*sec(e + f*x))*sqrt(sin(e + f*x))), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(1/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/(sqrt(b*sec(f*x + e))*sqrt(sin(f*x + e))), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\sqrt{\sin(e + f x)} \sqrt{\frac{b}{\cos(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^(1/2)*(b/cos(e + f*x))^(1/2)),x)`

[Out] `int(1/(sin(e + f*x)^(1/2)*(b/cos(e + f*x))^(1/2)), x)`

$$3.468 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{5}{2}}(e + fx)} dx$$

Optimal. Leaf size=30

$$-\frac{2b}{3f(b \sec(e + fx))^{3/2} \sin^{\frac{3}{2}}(e + fx)}$$

[Out] $-2/3*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(3/2)}$

Rubi [A]

time = 0.02, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {2658}

$$-\frac{2b}{3f \sin^{\frac{3}{2}}(e + fx)(b \sec(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Int[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(5/2)),x]`

[Out] $(-2*b)/(3*f*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^{(3/2)})$

Rule 2658

`Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sine[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m - n + 2, 0] & & NeQ[m, -1]`

Rubi steps

$$\int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{5}{2}}(e + fx)} dx = -\frac{2b}{3f(b \sec(e + fx))^{3/2} \sin^{\frac{3}{2}}(e + fx)}$$

Mathematica [A]

time = 0.07, size = 30, normalized size = 1.00

$$-\frac{2b}{3f(b \sec(e + fx))^{3/2} \sin^{\frac{3}{2}}(e + fx)}$$

Antiderivative was successfully verified.

[In] `Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(5/2)),x]`

[Out] $(-2*b)/(3*f*(b*\text{Sec}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^{(3/2)})$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 69 vs. $2(24) = 48$.

time = 0.18, size = 70, normalized size = 2.33

method	result	size
default	$-\frac{8 \cos(fx+e)(-1+\cos(fx+e))^2}{3f \sin(fx+e)^{\frac{3}{2}} (\sin^2(fx+e)+\cos^2(fx+e)-2 \cos(fx+e)+1)^2 \sqrt{\frac{b}{\cos(fx+e)}}}$	70

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-8/3/f*\cos(f*x+e)*(-1+\cos(f*x+e))^2/\sin(f*x+e)^{(3/2)}/(\sin(f*x+e)^2+\cos(f*x+e)^2-2*\cos(f*x+e)+1)^2/(b/\cos(f*x+e))^{(1/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] `integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(5/2)), x)`

Fricas [A]

time = 0.41, size = 52, normalized size = 1.73

$$\frac{2 \sqrt{\frac{b}{\cos(fx+e)}} \cos(fx+e)^2 \sqrt{\sin(fx+e)}}{3 (bf \cos(fx+e)^2 - bf)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")`

[Out] $2/3*\text{sqrt}(b/\cos(f*x + e))*\cos(f*x + e)^2*\text{sqrt}(\sin(f*x + e))/(b*f*\cos(f*x + e)^2 - b*f)$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)**(5/2)/(b*sec(f*x+e))**(1/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 6438 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(5/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(5/2)), x)

Mupad [B]

time = 1.25, size = 57, normalized size = 1.90

$$\frac{\sqrt{\frac{b}{\cos(e + f x)}} (\sin(e + f x) + \sin(3e + 3f x))}{3 b f \sqrt{\sin(e + f x)} (\cos(2e + 2f x) - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^(5/2)*(b/cos(e + f*x))^(1/2)),x)

[Out] ((b/cos(e + f*x))^(1/2)*(sin(e + f*x) + sin(3*e + 3*f*x)))/(3*b*f*sin(e + f*x)^(1/2)*(cos(2*e + 2*f*x) - 1))

$$3.469 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{9}{2}}(e + fx)} dx$$

Optimal. Leaf size=61

$$-\frac{2b}{7f(b \sec(e + fx))^{3/2} \sin^{\frac{7}{2}}(e + fx)} - \frac{8b}{21f(b \sec(e + fx))^{3/2} \sin^{\frac{3}{2}}(e + fx)}$$

[Out] $-2/7*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(7/2)}-8/21*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(3/2)}$

Rubi [A]

time = 0.05, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2664, 2658}

$$-\frac{8b}{21f \sin^{\frac{3}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{2b}{7f \sin^{\frac{7}{2}}(e + fx)(b \sec(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(9/2)),x]

[Out] $(-2*b)/(7*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(7/2)}) - (8*b)/(21*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(3/2)})$

Rule 2658

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sine[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m - n + 2, 0] && NeQ[m, -1]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sine[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sine[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{9}{2}}(e + fx)} dx &= -\frac{2b}{7f(b \sec(e + fx))^{3/2} \sin^{\frac{7}{2}}(e + fx)} + \frac{4}{7} \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{5}{2}}(e + fx)} dx \\ &= -\frac{2b}{7f(b \sec(e + fx))^{3/2} \sin^{\frac{7}{2}}(e + fx)} - \frac{8b}{21f(b \sec(e + fx))^{3/2} \sin^{\frac{3}{2}}(e + fx)} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 42, normalized size = 0.69

$$\frac{2b(-5 + 2 \cos(2(e + fx)))}{21f(b \sec(e + fx))^{3/2} \sin^{7/2}(e + fx)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(9/2)),x]

[Out] (2*b*(-5 + 2*Cos[2*(e + f*x)]))/(21*f*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^(7/2))

Maple [A]

time = 0.19, size = 82, normalized size = 1.34

method	result	size
default	$\frac{32 \cos(fx+e)(4(\cos^2(fx+e)-7)(-1+\cos(fx+e))^4}{21f \sin(fx+e)^{7/2}(\sin^2(fx+e)+\cos^2(fx+e)-2\cos(fx+e)+1)^4 \sqrt{\frac{b}{\cos(fx+e)}}$	82

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] 32/21/f*cos(f*x+e)*(4*cos(f*x+e)^2-7)*(-1+cos(f*x+e))^4/sin(f*x+e)^(7/2)/(sin(f*x+e)^2+cos(f*x+e)^2-2*cos(f*x+e)+1)^4/(b/cos(f*x+e))^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(9/2)), x)

Fricas [A]

time = 0.42, size = 78, normalized size = 1.28

$$\frac{2(4 \cos(fx + e)^4 - 7 \cos(fx + e)^2) \sqrt{\frac{b}{\cos(fx + e)}} \sqrt{\sin(fx + e)}}{21(bf \cos(fx + e)^4 - 2bf \cos(fx + e)^2 + bf)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $2/21*(4*\cos(f*x + e)^4 - 7*\cos(f*x + e)^2)*\sqrt{b/\cos(f*x + e)}*\sqrt{\sin(f*x + e)}/(b*f*\cos(f*x + e)^4 - 2*b*f*\cos(f*x + e)^2 + b*f)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)**(9/2)/(b*sec(f*x+e))**(1/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/sin(f*x+e)^(9/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(9/2)), x)`

Mupad [B]

time = 2.66, size = 103, normalized size = 1.69

$$4 \sqrt{\frac{b}{\cos(e + f x)}} \frac{(11 \sin(e + f x) + 4 \sin(3e + 3f x) - 6 \sin(5e + 5f x) + \sin(7e + 7f x))}{21 b f \sqrt{\sin(e + f x)} (15 \cos(2e + 2f x) - 6 \cos(4e + 4f x) + \cos(6e + 6f x) - 10)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(sin(e + f*x)^(9/2)*(b/cos(e + f*x))^(1/2)),x)`

[Out] $(4*(b/\cos(e + f*x))^(1/2)*(11*\sin(e + f*x) + 4*\sin(3*e + 3*f*x) - 6*\sin(5*e + 5*f*x) + \sin(7*e + 7*f*x)))/(21*b*f*\sin(e + f*x)^(1/2)*(15*\cos(2*e + 2*f*x) - 6*\cos(4*e + 4*f*x) + \cos(6*e + 6*f*x) - 10))$

$$3.470 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{13}{2}}(e + fx)} dx$$

Optimal. Leaf size=91

$$\frac{2b}{11f(b \sec(e + fx))^{3/2} \sin^{\frac{11}{2}}(e + fx)} - \frac{16b}{77f(b \sec(e + fx))^{3/2} \sin^{\frac{7}{2}}(e + fx)} - \frac{64b}{231f(b \sec(e + fx))^{3/2} \sin^{\frac{3}{2}}(e + fx)}$$

[Out] $-2/11*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(11/2)}-16/77*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(7/2)}-64/231*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(3/2)}$

Rubi [A]

time = 0.08, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$,

Rules used = {2664, 2658}

$$\frac{64b}{231f \sin^{\frac{3}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{16b}{77f \sin^{\frac{7}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{2b}{11f \sin^{\frac{11}{2}}(e + fx)(b \sec(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(13/2)),x]

[Out] $(-2*b)/(11*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(11/2)}) - (16*b)/(77*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(7/2)}) - (64*b)/(231*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(3/2)})$

Rule 2658

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m - n + 2, 0] & & NeQ[m, -1]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{13}{2}}(e+fx)} dx &= -\frac{2b}{11f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)} + \frac{8}{11} \int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{9}{2}}(e+fx)} dx \\ &= -\frac{2b}{11f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)} - \frac{16b}{77f(b \sec(e+fx))^{3/2} \sin^{\frac{7}{2}}(e+fx)} \\ &= -\frac{2b}{11f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)} - \frac{16b}{77f(b \sec(e+fx))^{3/2} \sin^{\frac{7}{2}}(e+fx)} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 52, normalized size = 0.57

$$\frac{2b(-45 + 28 \cos(2(e+fx)) - 4 \cos(4(e+fx)))}{231f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(13/2)),x]``[Out] (2*b*(-45 + 28*Cos[2*(e + f*x)] - 4*Cos[4*(e + f*x)]))/(231*f*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^(11/2))`**Maple [A]**

time = 0.22, size = 92, normalized size = 1.01

method	result	size
default	$-\frac{128 \cos(fx+e)(32(\cos^4(fx+e))-88(\cos^2(fx+e))+77)(-1+\cos(fx+e))^6}{231f \sin(fx+e)^{\frac{11}{2}}(\sin^2(fx+e)+\cos^2(fx+e)-2\cos(fx+e)+1)^6 \sqrt{\frac{b}{\cos(fx+e)}}}$	92

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/sin(f*x+e)^(13/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)``[Out] -128/231/f*cos(f*x+e)*(32*cos(f*x+e)^4-88*cos(f*x+e)^2+77)*(-1+cos(f*x+e))^6/sin(f*x+e)^(11/2)/(sin(f*x+e)^2+cos(f*x+e)^2-2*cos(f*x+e)+1)^6/(b/cos(f*x+e))^(1/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/sin(f*x+e)^(13/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")`

[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(13/2)), x)

Fricas [A]

time = 0.45, size = 103, normalized size = 1.13

$$\frac{2(32 \cos(fx + e)^6 - 88 \cos(fx + e)^4 + 77 \cos(fx + e)^2) \sqrt{\frac{b}{\cos(fx + e)}} \sqrt{\sin(fx + e)}}{231(bf \cos(fx + e)^6 - 3bf \cos(fx + e)^4 + 3bf \cos(fx + e)^2 - bf)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(13/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 2/231*(32*cos(f*x + e)^6 - 88*cos(f*x + e)^4 + 77*cos(f*x + e)^2)*sqrt(b/cos(f*x + e))*sqrt(sin(f*x + e))/(b*f*cos(f*x + e)^6 - 3*b*f*cos(f*x + e)^4 + 3*b*f*cos(f*x + e)^2 - b*f)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)**(13/2)/(b*sec(f*x+e))^(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(13/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(13/2)), x)

Mupad [B]

time = 6.24, size = 163, normalized size = 1.79

$$\frac{e^{-e6i - fx6i} \sqrt{\frac{b}{\frac{e^{-e11 - fx11}}{2} + \frac{e^{e11 + fx11}}{2}}} \left(\frac{e^{e6i + fx6i} 992i}{231bf} + \frac{e^{e6i + fx6i} \cos(2e + 2fx) 608i}{231bf} - \frac{e^{e6i + fx6i} \cos(4e + 4fx) 320i}{231bf} + \frac{e^{e6i + fx6i} \cos(6e + 6fx) 64i}{231bf} \right) i}{32 \sin(e + fx)^{11/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^(13/2)*(b/cos(e + f*x))^(1/2)),x)

[Out] (exp(-e*6i - f*x*6i)*(b/(exp(-e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2)))^(1/2)*((exp(e*6i + f*x*6i)*992i)/(231*b*f) + (exp(e*6i + f*x*6i)*cos(2*e + 2*f*x)*608i)/(231*b*f) - (exp(e*6i + f*x*6i)*cos(4*e + 4*f*x)*320i)/(231*b*f) + (exp(e*6i + f*x*6i)*cos(6*e + 6*f*x)*64i)/(231*b*f))*1i/(32*sin(e + f*x)^(11/2))

$$3.471 \quad \int \frac{1}{\sqrt{b \sec(e + fx)} \sin^{\frac{17}{2}}(e + fx)} dx$$

Optimal. Leaf size=121

$$\frac{2b}{15f(b \sec(e + fx))^{3/2} \sin^{\frac{15}{2}}(e + fx)} - \frac{8b}{55f(b \sec(e + fx))^{3/2} \sin^{\frac{11}{2}}(e + fx)} - \frac{64b}{385f(b \sec(e + fx))^{3/2} \sin^{\frac{7}{2}}(e + fx)}$$

[Out] $-2/15*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(15/2)}-8/55*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(11/2)}-64/385*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(7/2)}-256/1155*b/f/(b*\sec(f*x+e))^{(3/2)}/\sin(f*x+e)^{(3/2)}$

Rubi [A]

time = 0.11, antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2664, 2658}

$$\frac{256b}{1155f \sin^{\frac{3}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{64b}{385f \sin^{\frac{7}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{8b}{55f \sin^{\frac{11}{2}}(e + fx)(b \sec(e + fx))^{3/2}} - \frac{2b}{15f \sin^{\frac{15}{2}}(e + fx)(b \sec(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(17/2)),x]

[Out] $(-2*b)/(15*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(15/2)}) - (8*b)/(55*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(11/2)}) - (64*b)/(385*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(7/2)}) - (256*b)/(1155*f*(b*Sec[e + f*x])^{(3/2)}*Sin[e + f*x]^{(3/2)})$

Rule 2658

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sine[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] /; FreeQ[{a, b, e, f, m, n}, x] && EqQ[m - n + 2, 0] && NeQ[m, -1]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_.), x_Symbol] :> Simp[b*(a*Sine[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1))/(a*f*(m + 1)), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sine[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rubi steps

$$\begin{aligned}
\int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{17}{2}}(e+fx)} dx &= -\frac{2b}{15f(b \sec(e+fx))^{3/2} \sin^{\frac{15}{2}}(e+fx)} + \frac{4}{5} \int \frac{1}{\sqrt{b \sec(e+fx)} \sin^{\frac{13}{2}}(e+fx)} dx \\
&= -\frac{2b}{15f(b \sec(e+fx))^{3/2} \sin^{\frac{15}{2}}(e+fx)} - \frac{8b}{55f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)} \\
&= -\frac{2b}{15f(b \sec(e+fx))^{3/2} \sin^{\frac{15}{2}}(e+fx)} - \frac{8b}{55f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)} \\
&= -\frac{2b}{15f(b \sec(e+fx))^{3/2} \sin^{\frac{15}{2}}(e+fx)} - \frac{8b}{55f(b \sec(e+fx))^{3/2} \sin^{\frac{11}{2}}(e+fx)}
\end{aligned}$$

Mathematica [A]

time = 0.19, size = 62, normalized size = 0.51

$$\frac{2b(-195 + 150 \cos(2(e+fx)) - 36 \cos(4(e+fx)) + 4 \cos(6(e+fx)))}{1155f(b \sec(e+fx))^{3/2} \sin^{\frac{15}{2}}(e+fx)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(Sqrt[b*Sec[e + f*x]]*Sin[e + f*x]^(17/2)),x]

[Out] (2*b*(-195 + 150*Cos[2*(e + f*x)] - 36*Cos[4*(e + f*x)] + 4*Cos[6*(e + f*x)]))/(1155*f*(b*Sec[e + f*x])^(3/2)*Sin[e + f*x]^(15/2))

Maple [A]

time = 0.27, size = 102, normalized size = 0.84

method	result	size
default	$\frac{512 \cos(fx+e)(128(\cos^6(fx+e)) - 480(\cos^4(fx+e)) + 660(\cos^2(fx+e)) - 385)(-1 + \cos(fx+e))^8}{1155f \sin(fx+e)^{\frac{15}{2}} (\sin^2(fx+e) + \cos^2(fx+e) - 2 \cos(fx+e) + 1)^8 \sqrt{\frac{b}{\cos(fx+e)}}}$	102

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/sin(f*x+e)^(17/2)/(b*sec(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out] 512/1155/f*cos(f*x+e)*(128*cos(f*x+e)^6-480*cos(f*x+e)^4+660*cos(f*x+e)^2-385)*(-1+cos(f*x+e))^8/sin(f*x+e)^(15/2)/(sin(f*x+e)^2+cos(f*x+e)^2-2*cos(f*x+e)+1)^8/(b/cos(f*x+e))^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(17/2)/(b*sec(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(17/2)), x)

Fricas [A]

time = 0.43, size = 126, normalized size = 1.04

$$\frac{2(128 \cos(fx + e)^8 - 480 \cos(fx + e)^6 + 660 \cos(fx + e)^4 - 385 \cos(fx + e)^2) \sqrt{\frac{b}{\cos(fx + e)}} \sqrt{\sin(fx + e)}}{1155 (bf \cos(fx + e)^8 - 4bf \cos(fx + e)^6 + 6bf \cos(fx + e)^4 - 4bf \cos(fx + e)^2 + bf)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(17/2)/(b*sec(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 2/1155*(128*cos(f*x + e)^8 - 480*cos(f*x + e)^6 + 660*cos(f*x + e)^4 - 385*cos(f*x + e)^2)*sqrt(b/cos(f*x + e))*sqrt(sin(f*x + e))/(b*f*cos(f*x + e)^8 - 4*b*f*cos(f*x + e)^6 + 6*b*f*cos(f*x + e)^4 - 4*b*f*cos(f*x + e)^2 + b*f)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)**(17/2)/(b*sec(f*x+e))**(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/sin(f*x+e)^(17/2)/(b*sec(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(1/(sqrt(b*sec(f*x + e))*sin(f*x + e)^(17/2)), x)

Mupad [B]

time = 6.41, size = 192, normalized size = 1.59

$$\frac{e^{-e 8i - f x 8i} \sqrt{\frac{b}{\frac{e^{-e 11 - f x 11}}{2} + \frac{e^{e 11 + f x 11}}{2}}} \left(\frac{e^{e 8i + f x 8i} 1024i}{77 b f} + \frac{e^{e 8i + f x 8i} \cos(2e + 2fx) 384i}{55 b f} - \frac{e^{e 8i + f x 8i} \cos(4e + 4fx) 5248i}{1155 b f} + \frac{e^{e 8i + f x 8i} \cos(6e + 6fx) 256i}{165 b f} - \frac{e^{e 8i + f x 8i} \cos(8e + 8fx) 256i}{1155 b f} \right) \operatorname{li}}{128 \sin(e + fx)^{15/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^(17/2)*(b/cos(e + f*x))^(1/2)),x)

```
[Out] (exp(- e*8i - f*x*8i)*(b/(exp(- e*1i - f*x*1i)/2 + exp(e*1i + f*x*1i)/2))^(
1/2)*((exp(e*8i + f*x*8i)*1024i)/(77*b*f) + (exp(e*8i + f*x*8i)*cos(2*e + 2
*f*x)*384i)/(55*b*f) - (exp(e*8i + f*x*8i)*cos(4*e + 4*f*x)*5248i)/(1155*b*
f) + (exp(e*8i + f*x*8i)*cos(6*e + 6*f*x)*256i)/(165*b*f) - (exp(e*8i + f*x
*8i)*cos(8*e + 8*f*x)*256i)/(1155*b*f))*1i)/(128*sin(e + f*x)^(15/2))
```


$$3.472 \quad \int \frac{(a \sin(e+fx))^{9/2}}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=490

$$\frac{7a^{9/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} - 7a^{9/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}}{128 \sqrt{2} b^{5/2} f}$$

[Out] $-7/192*a^3*(a*\sin(f*x+e))^{(3/2)}/b/f/(b*\sec(f*x+e))^{(1/2)}-1/48*a*(a*\sin(f*x+e))^{(7/2)}/b/f/(b*\sec(f*x+e))^{(1/2)}+1/6*(a*\sin(f*x+e))^{(11/2)}/a/b/f/(b*\sec(f*x+e))^{(1/2)}-7/256*a^{(9/2)}*\arctan(1-2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/a^{(1/2)/(b*\cos(f*x+e))^{(1/2)}}*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}+7/256*a^{(9/2)}*\arctan(1+2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/a^{(1/2)/(b*\cos(f*x+e))^{(1/2)}}*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}+7/512*a^{(9/2)}*\ln(a^{(1/2)}-2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}+a^{(1/2)}*\tan(f*x+e))*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}-7/512*a^{(9/2)}*\ln(a^{(1/2)}+2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}+a^{(1/2)}*\tan(f*x+e))*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}$

Rubi [A]

time = 0.37, antiderivative size = 490, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2662, 2663, 2665, 2654, 303, 1176, 631, 210, 1179, 642}

$\frac{1}{128 \sqrt{2} b^{5/2} f} \left(7a^{9/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} - 7a^{9/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \right)$

Antiderivative was successfully verified.

[In] Int[(a*Sin[e + f*x])^(9/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] $(-7*a^{(9/2)}*ArcTan[1 - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*\cos[e + f*x]])]*Sqrt[b*\cos[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(128*Sqrt[2]*b^{(5/2)}*f) + (7*a^{(9/2)}*ArcTan[1 + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*\cos[e + f*x]])]*Sqrt[b*\cos[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(128*Sqrt[2]*b^{(5/2)}*f) + (7*a^{(9/2)}*Sqrt[b*\cos[e + f*x]]*Log[Sqrt[a] - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*\cos[e + f*x]] + Sqrt[a]*Tan[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(256*Sqrt[2]*b^{(5/2)}*f) - (7*a^{(9/2)}*Sqrt[b*\cos[e + f*x]]*Log[Sqrt[a] + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*\cos[e + f*x]] + Sqrt[a]*Tan[e + f*x]]*Sqrt[b*\sec[e + f*x]])/(256*Sqrt[2]*b^{(5/2)}*f) - (7*a^3*(a*\sin[e + f*x])^{(3/2)})/(192*b*f*Sqrt[b*\sec[e + f*x]]) - (a*(a*\sin[e + f*x])^{(7/2)})/(48*b*f*Sqrt[b*\sec[e + f*x]]) + (a*\sin[e + f*x])^{(11/2)}/(6*a*b*f*Sqrt[b*\sec[e + f*x]])$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 2654

Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_)])^m, x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e + f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] && LtQ[m, 1]

Rule 2662

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a*b*f*(m - n))), x] - Dist[(n + 1)/(b^2*(m - n)), Int[(a*Sin[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2663

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Simp[(-a)*b*(a*Sin[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n - 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*Sin[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a \sin(e + fx))^{9/2}}{(b \sec(e + fx))^{3/2}} dx &= \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{12b^2} \\
&= -\frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^2) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^4) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^4) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^5) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^5) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^5) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^5) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= -\frac{7a^3(a \sin(e + fx))^{3/2}}{192bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{7/2}}{48bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{11/2}}{6abf \sqrt{b \sec(e + fx)}} + \frac{(7a^5) \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{9/2} dx}{96b^2} \\
&= \frac{7a^{9/2} \sqrt{b \cos(e + fx)} \log \left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) \right)}{256 \sqrt{2} b^{5/2} f} \\
&= -\frac{7a^{9/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{128 \sqrt{2} b^{5/2} f} +
\end{aligned}$$

Mathematica [A]

time = 0.98, size = 176, normalized size = 0.36

$$\frac{a^5 \left(4(-3 + 14 \cos(2(e + fx)) - 4 \cos(4(e + fx))) \sin^2(e + fx) - 21\sqrt{2} \tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt[3]{\tan^2(e + fx)}} \right) \sqrt[3]{\tan^2(e + fx)} + 21\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{2} \sqrt[3]{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \sqrt[3]{\tan^2(e + fx)} \right)}{768bf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sin[e + f*x])^(9/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] $-1/768*(a^5*(4*(-3 + 14*\cos[2*(e + f*x)] - 4*\cos[4*(e + f*x)])*\sin[e + f*x]^2 - 21*\sqrt{2}*\arctan[(-1 + \sqrt{\tan[e + f*x]^2})/(\sqrt{2}*(\tan[e + f*x]^2)^{(1/4)})])*(\tan[e + f*x]^2)^{(1/4)} + 21*\sqrt{2}*\operatorname{arctanh}[(\sqrt{2}*(\tan[e + f*x]^2)^{(1/4)})/(1 + \sqrt{\tan[e + f*x]^2})]*(\tan[e + f*x]^2)^{(1/4)})/(b*f*\sqrt{b*\sec[e + f*x]}*\sqrt{a*\sin[e + f*x]})$

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.29, size = 572, normalized size = 1.17

method	result
default	$-\frac{\left(-64\sqrt{2}(\cos^6(fx+e))+64\sqrt{2}(\cos^5(fx+e))+21i\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sin(f*x+e))^(9/2)/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`

[Out] $-1/768/f*(-64*2^{(1/2)}*\cos(f*x+e)^6+64*2^{(1/2)}*\cos(f*x+e)^5+21*I*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})-21*I*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})+120*\cos(f*x+e)^4*2^{(1/2)}-120*\cos(f*x+e)^3*2^{(1/2)}-21*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2-1/2*I,1/2*2^{(1/2)})-21*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*\operatorname{EllipticPi}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})-42*\cos(f*x+e)^2*2^{(1/2)}+42*\cos(f*x+e)*2^{(1/2)}*(a*\sin(f*x+e))^{(9/2)/(-1+\cos(f*x+e))/\sin(f*x+e)^3/\cos(f*x+e)^2/(b/\cos(f*x+e))^{(3/2)}*2^{(1/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a*sin(f*x+e))^(9/2)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate((a*sin(f*x + e))^(9/2)/(b*sec(f*x + e))^(3/2), x)`

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(9/2)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))**(9/2)/(b*sec(f*x+e))**(3/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(9/2)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((a*sin(f*x + e))^(9/2)/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a \sin(e + f x))^{9/2}}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(9/2)/(b/cos(e + f*x))^(3/2),x)

[Out] int((a*sin(e + f*x))^(9/2)/(b/cos(e + f*x))^(3/2), x)

$$3.473 \quad \int \frac{(a \sin(e+fx))^{5/2}}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=453

$$\frac{3a^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} + 3a^{5/2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)}}{32\sqrt{2} b^{5/2} f}$$

[Out] $-1/16*a*(a*\sin(f*x+e))^{(3/2)}/b/f/(b*\sec(f*x+e))^{(1/2)}+1/4*(a*\sin(f*x+e))^{(7/2)}/a/b/f/(b*\sec(f*x+e))^{(1/2)}-3/64*a^{(5/2)}*\arctan(1-2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/a^{(1/2)}/(b*\cos(f*x+e))^{(1/2)})*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}+3/64*a^{(5/2)}*\arctan(1+2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/a^{(1/2)}/(b*\cos(f*x+e))^{(1/2)})*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}+3/128*a^{(5/2)}*\ln(a^{(1/2)}-2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}+a^{(1/2)}*\tan(f*x+e))*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}-3/128*a^{(5/2)}*\ln(a^{(1/2)}+2^{(1/2)}*b^{(1/2)}*(a*\sin(f*x+e))^{(1/2)}/(b*\cos(f*x+e))^{(1/2)}+a^{(1/2)}*\tan(f*x+e))*(b*\cos(f*x+e))^{(1/2)}*(b*\sec(f*x+e))^{(1/2)}/b^{(5/2)}/f*2^{(1/2)}$

Rubi [A]

time = 0.30, antiderivative size = 453, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.400$, Rules used = {2662, 2663, 2665, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{3a^{5/2} \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}}\right) + 3a^{5/2} \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}}\right) + 3a^{5/2} \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \log\left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} + \sqrt{a} \tan(e+fx) + \sqrt{a}\right) - 3a^{5/2} \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \log\left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} - \sqrt{a} \tan(e+fx) + \sqrt{a}\right)}{32\sqrt{2} b^{5/2} f}$$

Antiderivative was successfully verified.

[In] Int[(a*Sin[e + f*x])^(5/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] $(-3*a^{(5/2)}*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\sin[e + f*x]])]/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b*\cos[e + f*x]])]*\operatorname{Sqrt}[b*\cos[e + f*x]]*\operatorname{Sqrt}[b*\sec[e + f*x]]/(32*\operatorname{Sqrt}[2]*b^{(5/2)}*f) + (3*a^{(5/2)}*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\sin[e + f*x]])]/(\operatorname{Sqrt}[a]*\operatorname{Sqrt}[b*\cos[e + f*x]])]*\operatorname{Sqrt}[b*\cos[e + f*x]]*\operatorname{Sqrt}[b*\sec[e + f*x]]/(32*\operatorname{Sqrt}[2]*b^{(5/2)}*f) + (3*a^{(5/2)}*\operatorname{Sqrt}[b*\cos[e + f*x]]*\operatorname{Log}[\operatorname{Sqrt}[a] - (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\sin[e + f*x]])/\operatorname{Sqrt}[b*\cos[e + f*x]] + \operatorname{Sqrt}[a]*\tan[e + f*x]]*\operatorname{Sqrt}[b*\sec[e + f*x]])/(64*\operatorname{Sqrt}[2]*b^{(5/2)}*f) - (3*a^{(5/2)}*\operatorname{Sqrt}[b*\cos[e + f*x]]*\operatorname{Log}[\operatorname{Sqrt}[a] + (\operatorname{Sqrt}[2]*\operatorname{Sqrt}[b]*\operatorname{Sqrt}[a*\sin[e + f*x]])/\operatorname{Sqrt}[b*\cos[e + f*x]] + \operatorname{Sqrt}[a]*\tan[e + f*x]]*\operatorname{Sqrt}[b*\sec[e + f*x]])/(64*\operatorname{Sqrt}[2]*b^{(5/2)}*f) - (a*(a*\sin[e + f*x])^{(3/2)})/(16*b*f*\operatorname{Sqrt}[b*\sec[e + f*x]]) + (a*\sin[e + f*x])^{(7/2)}/(4*a*b*f*\operatorname{Sqrt}[b*\sec[e + f*x]])$

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &

& (LtQ[a, 0] || LtQ[b, 0])

Rule 303

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[
e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)])*(b_)^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sine[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rule 2662


```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Simp[(a*SIN[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a
*b*f*(m - n))), x] - Dist[(n + 1)/(b^2*(m - n)), Int[(a*SIN[e + f*x])^m*(b*
Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] &&
NeQ[m - n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2663

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Simp[(-a)*b*(a*SIN[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n
- 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*SIN[e + f*x])^(m
- 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1]
&& NeQ[m - n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] :> Dist[(b*cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*SIN[e +
f*x])^m/(b*cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte
gerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a \sin(e + fx))^{5/2}}{(b \sec(e + fx))^{3/2}} dx &= \frac{(a \sin(e + fx))^{7/2}}{4abf \sqrt{b \sec(e + fx)}} + \frac{\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2} dx}{8b^2} \\
&= -\frac{a(a \sin(e + fx))^{3/2}}{16bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{7/2}}{4abf \sqrt{b \sec(e + fx)}} + \frac{(3a^2) \int \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} dx}{32b^2} \\
&= -\frac{a(a \sin(e + fx))^{3/2}}{16bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{7/2}}{4abf \sqrt{b \sec(e + fx)}} + \frac{(3a^2 \sqrt{b \cos(e + fx)}) \sqrt{b \sec(e + fx)}}{32b^2} \\
&= -\frac{a(a \sin(e + fx))^{3/2}}{16bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{7/2}}{4abf \sqrt{b \sec(e + fx)}} + \frac{(3a^3 \sqrt{b \cos(e + fx)}) \sqrt{b \sec(e + fx)}}{32b^2} \\
&= -\frac{a(a \sin(e + fx))^{3/2}}{16bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{7/2}}{4abf \sqrt{b \sec(e + fx)}} - \frac{(3a^3 \sqrt{b \cos(e + fx)}) \sqrt{b \sec(e + fx)}}{32b^2} \\
&= -\frac{a(a \sin(e + fx))^{3/2}}{16bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{7/2}}{4abf \sqrt{b \sec(e + fx)}} + \frac{(3a^3 \sqrt{b \cos(e + fx)}) \sqrt{b \sec(e + fx)}}{32b^2} \\
&= \frac{3a^{5/2} \sqrt{b \cos(e + fx)} \log \left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) \right)}{64\sqrt{2} b^{5/2} f} \\
&= -\frac{3a^{5/2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{32\sqrt{2} b^{5/2} f} + \dots
\end{aligned}$$

Mathematica [A]

time = 0.82, size = 165, normalized size = 0.36

$$\frac{a^3 \left(4 - 6 \cos(2(e + fx)) + 2 \cos(4(e + fx)) + 3\sqrt{2} \tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} - 3\sqrt{2} \tanh^{-1} \left(\frac{\sqrt{2} \sqrt{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} \right)}{64bf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sin[e + f*x])^(5/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] (a^3*(4 - 6*Cos[2*(e + f*x)] + 2*Cos[4*(e + f*x)] + 3*Sqrt[2]*ArcTan[(-1 + Sqrt[Tan[e + f*x]^2)]/(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))])*(Tan[e + f*x]^2)^(1/2)

/4) - 3*sqrt[2]*ArcTanh[(sqrt[2]*(tan[e + f*x]^2)^(1/4))/(1 + sqrt[tan[e + f*x]^2])*(tan[e + f*x]^2)^(1/4)]/(64*b*f*sqrt[b*sec[e + f*x]]*sqrt[a*sin[e + f*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.

time = 0.25, size = 546, normalized size = 1.21

method	result
default	$-\frac{\left(3i\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\operatorname{EllipticPi}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{64bf\sqrt{b\sec[e+fx]}\sqrt{a\sin[e+fx]}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^(5/2)/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out]
$$-\frac{1}{64}f(3I\operatorname{EllipticPi}\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2},1/2-1/2I,1/2*2^{1/2})\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}-3I\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\operatorname{EllipticPi}\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2},1/2+1/2I,1/2*2^{1/2})+8\cos(fx+e)^4*2^{1/2}-3\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\operatorname{EllipticPi}\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2},1/2-1/2I,1/2*2^{1/2})-3\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\left(\frac{-1+\cos(fx+e)}{\sin(fx+e)}\right)^{1/2}\operatorname{EllipticPi}\left(\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}\right)^{1/2},1/2+1/2I,1/2*2^{1/2})-8\cos(fx+e)^3*2^{1/2}-6\cos(fx+e)^2*2^{1/2}+6\cos(fx+e)*2^{1/2})(a\sin(fx+e))^{5/2}/(-1+\cos(fx+e))/\cos(fx+e)^2/\sin(fx+e)/(b/\cos(fx+e))^{3/2}*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(5/2)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((a*sin(f*x + e))^(5/2)/(b*sec(f*x + e))^(3/2), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(5/2)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))**(5/2)/(b*sec(f*x+e))**(3/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(5/2)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((a*sin(f*x + e))^(5/2)/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a \sin(e + f x))^{5/2}}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(5/2)/(b/cos(e + f*x))^(3/2),x)

[Out] int((a*sin(e + f*x))^(5/2)/(b/cos(e + f*x))^(3/2), x)

$$3.474 \quad \int \frac{\sqrt{a \sin(e + fx)}}{(b \sec(e + fx))^{3/2}} dx$$

Optimal. Leaf size=418

$$\frac{\sqrt{a} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{4\sqrt{2} b^{5/2} f} + \frac{\sqrt{a} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{4\sqrt{2} b^{5/2} f}$$

```
[Out] 1/2*(a*sin(f*x+e))^(3/2)/a/b/f/(b*sec(f*x+e))^(1/2)-1/8*arctan(1-2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/a^(1/2)/(b*cos(f*x+e))^(1/2))*a^(1/2)*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/b^(5/2)/f*2^(1/2)+1/8*arctan(1+2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/a^(1/2)/(b*cos(f*x+e))^(1/2))*a^(1/2)*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/b^(5/2)/f*2^(1/2)+1/16*ln(a^(1/2)-2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/(b*cos(f*x+e))^(1/2)+a^(1/2)*tan(f*x+e))*a^(1/2)*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/b^(5/2)/f*2^(1/2)-1/16*ln(a^(1/2)+2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/(b*cos(f*x+e))^(1/2)+a^(1/2)*tan(f*x+e))*a^(1/2)*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/b^(5/2)/f*2^(1/2)
```

Rubi [A]

time = 0.23, antiderivative size = 418, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {2662, 2665, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \operatorname{Arctan} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{4\sqrt{2} b^{5/2} f} + \frac{\sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \operatorname{Arctan} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} \right)}{4\sqrt{2} b^{5/2} f} + \frac{\sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \log \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) + \sqrt{a} \right)}{8\sqrt{2} b^{5/2} f} - \frac{\sqrt{a} \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)} \log \left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx) + \sqrt{a} \right)}{8\sqrt{2} b^{5/2} f} + \frac{(a \sin(e + fx))^{3/2}}{2ab \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[a*Sin[e + f*x]]/(b*Sec[e + f*x])^(3/2),x]

```
[Out] -1/4*(Sqrt[a]*ArcTan[1 - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*Cos[e + f*x]])]*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])/(Sqrt[2]*b^(5/2)*f) + (Sqrt[a]*ArcTan[1 + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*Cos[e + f*x]])]*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])/(4*Sqrt[2]*b^(5/2)*f) + (Sqrt[a]*Sqrt[b*Cos[e + f*x]]*Log[Sqrt[a] - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*Cos[e + f*x]]] + Sqrt[a]*Tan[e + f*x])*Sqrt[b*Sec[e + f*x]])/(8*Sqrt[2]*b^(5/2)*f) - (Sqrt[a]*Sqrt[b*Cos[e + f*x]]*Log[Sqrt[a] + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*Cos[e + f*x]]] + Sqrt[a]*Tan[e + f*x])*Sqrt[b*Sec[e + f*x]])/(8*Sqrt[2]*b^(5/2)*f) + (a*Sin[e + f*x])^(3/2)/(2*a*b*f*Sqrt[b*Sec[e + f*x]])
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2654

```
Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rule 2662

```
Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m
_), x_Symbol] := Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a
```

*b*f*(m - n)), x] - Dist[(n + 1)/(b^2*(m - n)), Int[(a*Sin[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m - n, 0] && IntegerQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rubi steps

$$\begin{aligned}
 \int \frac{\sqrt{a \sin(e + fx)}}{(b \sec(e + fx))^{3/2}} dx &= \frac{(a \sin(e + fx))^{3/2}}{2abf \sqrt{b \sec(e + fx)}} + \frac{\int \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)} dx}{4b^2} \\
 &= \frac{(a \sin(e + fx))^{3/2}}{2abf \sqrt{b \sec(e + fx)}} + \frac{\left(\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}\right) \int \frac{\sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}}}{4b^2} \\
 &= \frac{(a \sin(e + fx))^{3/2}}{2abf \sqrt{b \sec(e + fx)}} + \frac{\left(a \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}\right) \text{Subst}\left(\int \frac{x^2}{a^2 + b^2 x^4} dx\right)}{2bf} \\
 &= \frac{(a \sin(e + fx))^{3/2}}{2abf \sqrt{b \sec(e + fx)}} - \frac{\left(a \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}\right) \text{Subst}\left(\int \frac{a - bx^2}{a^2 + b^2 x^4} dx\right)}{4b^2 f} \\
 &= \frac{(a \sin(e + fx))^{3/2}}{2abf \sqrt{b \sec(e + fx)}} + \frac{\left(a \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}\right) \text{Subst}\left(\int \frac{1}{\frac{a}{b} - \sqrt{2} \frac{1}{\sqrt{b}}}\right)}{8b^3 f} \\
 &= \frac{\sqrt{a} \sqrt{b \cos(e + fx)} \log\left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} + \sqrt{a} \tan(e + fx)\right)}{8\sqrt{2} b^{5/2} f} \\
 &= -\frac{\sqrt{a} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}}\right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{4\sqrt{2} b^{5/2} f}
 \end{aligned}$$

Mathematica [A]

time = 0.63, size = 151, normalized size = 0.36

$$\frac{a \left(4 \sin^2(e + fx) + \sqrt{2} \tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} - \sqrt{2} \tanh^{-1} \left(\frac{\sqrt{2} \sqrt{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \sqrt{\tan^2(e + fx)} \right)}{8bf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[a*Sin[e + f*x]]/(b*Sec[e + f*x])^(3/2), x]

[Out] (a*(4*Sin[e + f*x]^2 + Sqrt[2]*ArcTan[(-1 + Sqrt[Tan[e + f*x]^2])]/(Sqrt[2]*(Tan[e + f*x]^2)^(1/4)))*(Tan[e + f*x]^2)^(1/4) - Sqrt[2]*ArcTanh[(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))/(1 + Sqrt[Tan[e + f*x]^2])]*(Tan[e + f*x]^2)^(1/4)))/(8*b*f*Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]])

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.22, size = 516, normalized size = 1.23

method	result
default	$\left(i \sqrt{\frac{1 - \cos(fx+e) + \sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e) - 1 + \cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1 + \cos(fx+e)}{\sin(fx+e)}} \text{EllipticPi} \left(\sqrt{\frac{1 - \cos(fx+e) + \sin(fx+e)}{\sin(fx+e)}}, \frac{1}{2} + \right. \right.$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^(1/2)/(b*sec(f*x+e))^(3/2), x, method=_RETURNVERBOSE)

[Out] 1/8/f*(I*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2+1/2*I, 1/2*2^(1/2))-I*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2-1/2*I, 1/2*2^(1/2))+((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2+1/2*I, 1/2*2^(1/2))+((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2-1/2*I, 1/2*2^(1/2))+2*cos(f*x+e)^2*2^(1/2)-2*cos(f*x+e)*2^(1/2))*(a*sin(f*x+e))^(1/2)*sin(f*x+e)/(-1+cos(f*x+e))/cos(f*x+e)^2/(b/cos(f*x+e))^(3/2)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(1/2)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate(sqrt(a*sin(f*x + e))/(b*sec(f*x + e))^(3/2), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(1/2)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a \sin(e + fx)}}{(b \sec(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))**(1/2)/(b*sec(f*x+e))**(3/2),x)

[Out] Integral(sqrt(a*sin(e + f*x))/(b*sec(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(1/2)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(sqrt(a*sin(f*x + e))/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{a \sin(e + fx)}}{\left(\frac{b}{\cos(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(1/2)/(b/cos(e + f*x))^(3/2),x)

[Out] int((a*sin(e + f*x))^(1/2)/(b/cos(e + f*x))^(3/2), x)

$$3.475 \quad \int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{3/2}} dx$$

Optimal. Leaf size=411

$$\frac{\tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right) \sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} \right)}{\sqrt{2} a^{3/2} b^{5/2} f}$$

[Out] 1/2*arctan(1-2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/a^(1/2)/(b*cos(f*x+e))^(1/2))*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/a^(3/2)/b^(5/2)/f*2^(1/2)-1/2*arctan(1+2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/a^(1/2)/(b*cos(f*x+e))^(1/2))*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/a^(3/2)/b^(5/2)/f*2^(1/2)-1/4*ln(a^(1/2)-2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/(b*cos(f*x+e))^(1/2)+a^(1/2)*tan(f*x+e))*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/a^(3/2)/b^(5/2)/f*2^(1/2)+1/4*ln(a^(1/2)+2^(1/2)*b^(1/2)*(a*sin(f*x+e))^(1/2)/(b*cos(f*x+e))^(1/2)+a^(1/2)*tan(f*x+e))*(b*cos(f*x+e))^(1/2)*(b*sec(f*x+e))^(1/2)/a^(3/2)/b^(5/2)/f*2^(1/2)-2/a/b/f/(b*sec(f*x+e))^(1/2)/(a*sin(f*x+e))^(1/2)

Rubi [A]

time = 0.24, antiderivative size = 411, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.360$, Rules used = {2661, 2665, 2654, 303, 1176, 631, 210, 1179, 642}

$$\frac{\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}}\right)}{\sqrt{2} a^{3/2} b^{5/2} f} - \frac{\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \operatorname{ArcTan}\left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} + 1\right)}{\sqrt{2} a^{3/2} b^{5/2} f} - \frac{\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \log\left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} + \sqrt{a} \tan(e+fx) + \sqrt{a}\right)}{\sqrt{2} a^{3/2} b^{5/2} f} + \frac{\sqrt{b \cos(e+fx)} \sqrt{b \sec(e+fx)} \log\left(\frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e+fx)}}{\sqrt{a} \sqrt{b \cos(e+fx)}} + \sqrt{a} \tan(e+fx) + \sqrt{a}\right)}{\sqrt{2} a^{3/2} b^{5/2} f} - \frac{2}{ab \sqrt{a \sin(e+fx)} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(3/2)),x]

[Out] (ArcTan[1 - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*Cos[e + f*x]])]*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])/(Sqrt[2]*a^(3/2)*b^(5/2)*f) - (ArcTan[1 + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/(Sqrt[a]*Sqrt[b*Cos[e + f*x]])]*Sqrt[b*Cos[e + f*x]]*Sqrt[b*Sec[e + f*x]])/(Sqrt[2]*a^(3/2)*b^(5/2)*f) - (Sqrt[b*Cos[e + f*x]]*Log[Sqrt[a] - (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*Cos[e + f*x]] + Sqrt[a]*Tan[e + f*x]]*Sqrt[b*Sec[e + f*x]])/(2*Sqrt[2]*a^(3/2)*b^(5/2)*f) + (Sqrt[b*Cos[e + f*x]]*Log[Sqrt[a] + (Sqrt[2]*Sqrt[b]*Sqrt[a*Sin[e + f*x]])/Sqrt[b*Cos[e + f*x]] + Sqrt[a]*Tan[e + f*x]]*Sqrt[b*Sec[e + f*x]])/(2*Sqrt[2]*a^(3/2)*b^(5/2)*f) - 2/(a*b*f*Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 2654

```
Int[(cos[(e_.) + (f_.)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := With[{k = Denominator[m]}, Dist[k*a*(b/f), Subst[Int[x^(k*
(m + 1) - 1)/(a^2 + b^2*x^(2*k)), x], x, (a*Sin[e + f*x])^(1/k)/(b*Cos[e +
f*x])^(1/k)], x]] /; FreeQ[{a, b, e, f}, x] && EqQ[m + n, 0] && GtQ[m, 0] &
& LtQ[m, 1]
```

Rule 2661

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a
```

$*b*f*(m + 1))), x] - \text{Dist}[(n + 1)/(a^2*b^2*(m + 1)), \text{Int}[(a*\text{Sin}[e + f*x])^{(m + 2)}*(b*\text{Sec}[e + f*x])^{(n + 2)}, x], x] /; \text{FreeQ}[\{a, b, e, f\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2665

$\text{Int}[(b*sec[e + f*x])^n*(a*sin[e + f*x])^m, x_Symbol] := \text{Dist}[(b*\text{Cos}[e + f*x])^n*(b*\text{Sec}[e + f*x])^n, \text{Int}[(a*\text{Sin}[e + f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \&\& \text{IntegerQ}[m - 1/2] \&\& \text{IntegerQ}[n - 1/2]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \sec(e + fx))^{3/2} (a \sin(e + fx))^{3/2}} dx &= -\frac{2}{abf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}} - \frac{\int \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}{a^2 b^2} \\ &= -\frac{2}{abf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}} - \frac{(\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{a^2 b^2} \\ &= -\frac{2}{abf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}} - \frac{(2\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{a^2 b^2} \\ &= -\frac{2}{abf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}} + \frac{(\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{a^2 b^2} \\ &= -\frac{2}{abf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}} - \frac{(\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{a^2 b^2} \\ &= -\frac{\sqrt{b \cos(e + fx)} \log\left(\sqrt{a} - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{b \cos(e + fx)}} + \sqrt{a}\right)}{2\sqrt{2} a^{3/2} b^{5/2} f} \\ &= \frac{\tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt{b} \sqrt{a \sin(e + fx)}}{\sqrt{a} \sqrt{b \cos(e + fx)}}\right) \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}}{\sqrt{2} a^{3/2} b^{5/2} f} \end{aligned}$$

Mathematica [A]

time = 0.58, size = 144, normalized size = 0.35

$$\frac{4 + \sqrt{2} \tan^{-1} \left(\frac{-1 + \sqrt{\tan^2(e + fx)}}{\sqrt{2} \sqrt[4]{\tan^2(e + fx)}} \right) \sqrt[4]{\tan^2(e + fx)} - \sqrt{2} \tanh^{-1} \left(\frac{\sqrt{2} \sqrt[4]{\tan^2(e + fx)}}{1 + \sqrt{\tan^2(e + fx)}} \right) \sqrt[4]{\tan^2(e + fx)}}{2abf \sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(3/2)),x]
```

```
[Out] -1/2*(4 + Sqrt[2]*ArcTan[(-1 + Sqrt[Tan[e + f*x]^2])/(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))]*(Tan[e + f*x]^2)^(1/4) - Sqrt[2]*ArcTanh[(Sqrt[2]*(Tan[e + f*x]^2)^(1/4))/(1 + Sqrt[Tan[e + f*x]^2])]*(Tan[e + f*x]^2)^(1/4)]/(a*b*f*Sqrt[b*Sec[e + f*x]]*Sqrt[a*Sin[e + f*x]])
```

Maple [C] Result contains higher order function than in optimal. Order 4 vs. order 3.
time = 0.22, size = 957, normalized size = 2.33

method	result	size
default	Expression too large to display	957

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2/f*(I*cos(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-I*cos(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))-cos(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-cos(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))+I*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-I*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))-((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))-((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2-1/2*I,1/2*2^(1/2))-((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticPi(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2+1/2*I,1/2*2^(1/2))
```

$$2)*((-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*EllipticPi(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2+1/2*I,1/2*2^{(1/2)})+2*\cos(f*x+e)*2^{(1/2)}*\sin(f*x+e)/\cos(f*x+e)^2/(b/\cos(f*x+e))^{(3/2)}/(a*\sin(f*x+e))^{(3/2)}*2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(3/2)), x)

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(3/2),x, algorithm="fricas")

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))**(3/2)/(a*sin(f*x+e))**(3/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(3/2)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{(a \sin(e + f x))^{3/2} \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*sin(e + f*x))^(3/2)*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/((a*sin(e + f*x))^(3/2)*(b/cos(e + f*x))^(3/2)), x)

$$3.476 \quad \int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{7/2}} dx$$

Optimal. Leaf size=35

$$-\frac{2b}{5af(b \sec(e+fx))^{5/2} (a \sin(e+fx))^{5/2}}$$

[Out] $-2/5*b/a/f/(b*\sec(f*x+e))^{(5/2)}/(a*\sin(f*x+e))^{(5/2)}$

Rubi [A]

time = 0.04, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.040$, Rules used = {2658}

$$-\frac{2b}{5af(a \sin(e+fx))^{5/2} (b \sec(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/((b*\text{Sec}[e + f*x])^{(3/2)}*(a*\text{Sin}[e + f*x])^{(7/2)}),x]$

[Out] $(-2*b)/(5*a*f*(b*\text{Sec}[e + f*x])^{(5/2)}*(a*\text{Sin}[e + f*x])^{(5/2)})$

Rule 2658

$\text{Int}(((b_.)*\sec[(e_.) + (f_.)*(x_.)])^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol) \rightarrow \text{Simp}[b*(a*\text{Sin}[e + f*x])^{(m + 1)}*((b*\text{Sec}[e + f*x])^{(n - 1)})/(a*f*(m + 1)), x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x \ \&\& \ \text{EqQ}[m - n + 2, 0] \ \& \ \text{NeQ}[m, -1]$

Rubi steps

$$\int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{7/2}} dx = -\frac{2b}{5af(b \sec(e+fx))^{5/2} (a \sin(e+fx))^{5/2}}$$

Mathematica [A]

time = 0.09, size = 45, normalized size = 1.29

$$-\frac{2 \cot^3(e+fx) \sqrt{b \sec(e+fx)} \sqrt{a \sin(e+fx)}}{5a^4 b^2 f}$$

Antiderivative was successfully verified.

[In] $\text{Integrate}[1/((b*\text{Sec}[e + f*x])^{(3/2)}*(a*\text{Sin}[e + f*x])^{(7/2)}),x]$

[Out] $(-2*\text{Cot}[e + f*x]^3*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[a*\text{Sin}[e + f*x]])/(5*a^4*b^2*f)$

Maple [A]

time = 0.18, size = 40, normalized size = 1.14

method	result	size
default	$-\frac{2 \cos(fx+e) \sin(fx+e)}{5f \left(\frac{b}{\cos(fx+e)}\right)^{\frac{3}{2}} (a \sin(fx+e))^{\frac{7}{2}}}$	40

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(7/2),x,method=_RETURNVERBOSE)`

[Out] $-2/5/f*\cos(f*x+e)*\sin(f*x+e)/(b/\cos(f*x+e))^{3/2}/(a*\sin(f*x+e))^{7/2}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(7/2),x, algorithm="maxima")`

[Out] `integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(7/2)), x)`

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 73 vs. 2(31) = 62.

time = 0.42, size = 73, normalized size = 2.09

$$\frac{2 \sqrt{a \sin(fx + e)} \sqrt{\frac{b}{\cos(fx + e)}} \cos(fx + e)^3}{5 (a^4 b^2 f \cos(fx + e)^2 - a^4 b^2 f) \sin(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(7/2),x, algorithm="fricas")`

[Out] $2/5*\text{sqrt}(a*\sin(f*x + e))*\text{sqrt}(b/\cos(f*x + e))*\cos(f*x + e)^3/((a^4*b^2*f*\cos(f*x + e)^2 - a^4*b^2*f)*\sin(f*x + e))$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))**(3/2)/(a*sin(f*x+e))**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(7/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(7/2)), x)

Mupad [B]

time = 1.86, size = 84, normalized size = 2.40

$$\frac{\sqrt{\frac{b}{\cos(e + fx)}} (\cos(3e + 3fx) - 2 \cos(e + fx) + \cos(5e + 5fx))}{5a^3 b^2 f \sqrt{a \sin(e + fx)} (\cos(4e + 4fx) - 4 \cos(2e + 2fx) + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*sin(e + f*x))^(7/2)*(b/cos(e + f*x))^(3/2)),x)

[Out] ((b/cos(e + f*x))^(1/2)*(cos(3*e + 3*f*x) - 2*cos(e + f*x) + cos(5*e + 5*f*x)))/(5*a^3*b^2*f*(a*sin(e + f*x))^(1/2)*(cos(4*e + 4*f*x) - 4*cos(2*e + 2*f*x) + 3))

$$3.477 \quad \int \frac{(a \sin(e+fx))^{7/2}}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=172

$$-\frac{a^3 \sqrt{a \sin(e+fx)}}{12bf \sqrt{b \sec(e+fx)}} - \frac{a(a \sin(e+fx))^{5/2}}{30bf \sqrt{b \sec(e+fx)}} + \frac{(a \sin(e+fx))^{9/2}}{5abf \sqrt{b \sec(e+fx)}} + \frac{a^4 F(e - \frac{\pi}{4} + fx | 2) \sqrt{b \sec(e+fx)}}{24b^2 f \sqrt{a \sin(e+fx)}}$$

[Out] $-1/30*a*(a*\sin(f*x+e))^{(5/2)}/b/f/(b*\sec(f*x+e))^{(1/2)}+1/5*(a*\sin(f*x+e))^{(9/2)}/a/b/f/(b*\sec(f*x+e))^{(1/2)}-1/12*a^3*(a*\sin(f*x+e))^{(1/2)}/b/f/(b*\sec(f*x+e))^{(1/2)}-1/24*a^4*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticF}(\cos(e+1/4*Pi+f*x), 2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/b^2/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.19, antiderivative size = 172, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2662, 2663, 2665, 2653, 2720}

$$\frac{a^4 \sqrt{\sin(2e+2fx)} F(e+fx - \frac{\pi}{4} | 2) \sqrt{b \sec(e+fx)}}{24b^2 f \sqrt{a \sin(e+fx)}} - \frac{a^3 \sqrt{a \sin(e+fx)}}{12bf \sqrt{b \sec(e+fx)}} + \frac{(a \sin(e+fx))^{9/2}}{5abf \sqrt{b \sec(e+fx)}} - \frac{a(a \sin(e+fx))^{5/2}}{30bf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a*\text{Sin}[e + f*x])^{(7/2)}/(b*\text{Sec}[e + f*x])^{(3/2)}, x]$

[Out] $-1/12*(a^3*\text{Sqrt}[a*\text{Sin}[e + f*x]])/(b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) - (a*(a*\text{Sin}[e + f*x])^{(5/2)})/(30*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (a*\text{Sin}[e + f*x])^{(9/2)}/(5*a*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]) + (a^4*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]])*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(24*b^2*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e + f*x]]*\text{Sqrt}[b*\text{Cos}[e + f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2662

$\text{Int}[(b_.)*\sec[(e_.) + (f_.)*(x_.)]^{(n_.)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e + f*x])^{(m+1)}*((b*\text{Sec}[e + f*x])^{(n+1)})/(a*b*f*(m-n)), x] - \text{Dist}[(n+1)/(b^2*(m-n)), \text{Int}[(a*\text{Sin}[e + f*x])^{(m)}*(b*\text{Sec}[e + f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f, m\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{NeQ}[m-n, 0] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2663

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Simp[(-a)*b*(a*SIn[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n
- 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*SIn[e + f*x])^(m
- 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1]
&& NeQ[m - n, 0] && IntegersQ[2*m, 2*n]
```

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*SIn[e +
f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte
gerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a \sin(e + fx))^{7/2}}{(b \sec(e + fx))^{3/2}} dx &= \frac{(a \sin(e + fx))^{9/2}}{5abf \sqrt{b \sec(e + fx)}} + \frac{\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2} dx}{10b^2} \\
&= -\frac{a(a \sin(e + fx))^{5/2}}{30bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{9/2}}{5abf \sqrt{b \sec(e + fx)}} + \frac{a^2 \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{5/2} dx}{12b^2} \\
&= -\frac{a^3 \sqrt{a \sin(e + fx)}}{12bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{5/2}}{30bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{9/2}}{5abf \sqrt{b \sec(e + fx)}} + \frac{a^4 \int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2} dx}{10b^2} \\
&= -\frac{a^3 \sqrt{a \sin(e + fx)}}{12bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{5/2}}{30bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{9/2}}{5abf \sqrt{b \sec(e + fx)}} + \frac{(a^4 \sqrt{a \sin(e + fx)}) \int \sqrt{b \sec(e + fx)} dx}{10b^2} \\
&= -\frac{a^3 \sqrt{a \sin(e + fx)}}{12bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{5/2}}{30bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{9/2}}{5abf \sqrt{b \sec(e + fx)}} + \frac{(a^4 \sqrt{a \sin(e + fx)}) \int \sqrt{b \sec(e + fx)} dx}{10b^2} \\
&= -\frac{a^3 \sqrt{a \sin(e + fx)}}{12bf \sqrt{b \sec(e + fx)}} - \frac{a(a \sin(e + fx))^{5/2}}{30bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{9/2}}{5abf \sqrt{b \sec(e + fx)}} + \frac{a^4 F(\arcsin(\frac{a \sin(e + fx)}{\sqrt{a \sin(e + fx)}}))}{10b^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.63, size = 103, normalized size = 0.60

$$\frac{a^5 \left(-4 + 17 \cos(2(e + fx)) - 16 \cos(4(e + fx)) + 3 \cos(6(e + fx)) - 20 {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4} \right)}{480bf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sin[e + f*x])^(7/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] -1/480*(a^5*(-4 + 17*Cos[2*(e + f*x)] - 16*Cos[4*(e + f*x)] + 3*Cos[6*(e + f*x)] - 20*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4)))/(b*f*Sqrt[b*Sec[e + f*x]]*(a*Sin[e + f*x])^(3/2))

Maple [A]

time = 0.25, size = 246, normalized size = 1.43

method	result
default	$\frac{\left(12\sqrt{2} (\cos^6(fx+e)) - 12\sqrt{2} (\cos^5(fx+e)) - 22(\cos^4(fx+e))\sqrt{2} - 5\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \right)}{120f(-$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^(7/2)/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] 1/120/f*(12*2^(1/2)*cos(f*x+e)^6-12*2^(1/2)*cos(f*x+e)^5-22*cos(f*x+e)^4*2^(1/2)-5*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)+22*cos(f*x+e)^3*2^(1/2)+5*cos(f*x+e)^2*2^(1/2)-5*cos(f*x+e)*2^(1/2))*(a*sin(f*x+e))^(7/2)/(-1+cos(f*x+e))/(b/cos(f*x+e))^(3/2)/cos(f*x+e)^2/sin(f*x+e)^3*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(7/2)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((a*sin(f*x + e))^(7/2)/(b*sec(f*x + e))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(7/2)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")
[Out] integral(-(a^3*cos(f*x + e)^2 - a^3)*sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e))*sin(f*x + e)/(b^2*sec(f*x + e)^2), x)
Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00
```

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))**(7/2)/(b*sec(f*x+e))**(3/2),x)
[Out] Timed out
Giac [F]
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a*sin(f*x+e))^(7/2)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")
[Out] integrate((a*sin(f*x + e))^(7/2)/(b*sec(f*x + e))^(3/2), x)
Mupad [F]
time = 0.00, size = -1, normalized size = -0.01
```

$$\int \frac{(a \sin(e + f x))^{7/2}}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*sin(e + f*x))^(7/2)/(b/cos(e + f*x))^(3/2),x)
[Out] int((a*sin(e + f*x))^(7/2)/(b/cos(e + f*x))^(3/2), x)
```

$$3.478 \quad \int \frac{(a \sin(e+fx))^{3/2}}{(b \sec(e+fx))^{3/2}} dx$$

Optimal. Leaf size=135

$$-\frac{a\sqrt{a\sin(e+fx)}}{6bf\sqrt{b\sec(e+fx)}} + \frac{(a\sin(e+fx))^{5/2}}{3abf\sqrt{b\sec(e+fx)}} + \frac{a^2 F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b\sec(e+fx)} \sqrt{\sin(2e+2fx)}}{12b^2 f \sqrt{a\sin(e+fx)}}$$

[Out] 1/3*(a*sin(f*x+e))^(5/2)/a/b/f/(b*sec(f*x+e))^(1/2)-1/6*a*(a*sin(f*x+e))^(1/2)/b/f/(b*sec(f*x+e))^(1/2)-1/12*a^2*(sin(e+1/4*Pi+f*x)^2)^(1/2)/sin(e+1/4*Pi+f*x)*EllipticF(cos(e+1/4*Pi+f*x),2^(1/2))*(b*sec(f*x+e))^(1/2)*sin(2*f*x+2*e)^(1/2)/b^2/f/(a*sin(f*x+e))^(1/2)

Rubi [A]

time = 0.15, antiderivative size = 135, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2662, 2663, 2665, 2653, 2720}

$$\frac{a^2 \sqrt{\sin(2e+2fx)} F\left(e+fx-\frac{\pi}{4} \mid 2\right) \sqrt{b\sec(e+fx)}}{12b^2 f \sqrt{a\sin(e+fx)}} + \frac{(a\sin(e+fx))^{5/2}}{3abf\sqrt{b\sec(e+fx)}} - \frac{a\sqrt{a\sin(e+fx)}}{6bf\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[(a*Sin[e + f*x])^(3/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] -1/6*(a*Sqrt[a*Sin[e + f*x]])/(b*f*Sqrt[b*Sec[e + f*x]]) + (a*Sin[e + f*x])^(5/2)/(3*a*b*f*Sqrt[b*Sec[e + f*x]]) + (a^2*EllipticF[e - Pi/4 + f*x, 2]*Sqrt[b*Sec[e + f*x]]*Sqrt[Sin[2*e + 2*f*x]])/(12*b^2*f*Sqrt[a*Sin[e + f*x]])

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2662

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a*b*f*(m - n))), x] - Dist[(n + 1)/(b^2*(m - n)), Int[(a*Sin[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]

Rule 2663

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(-a)*b*(a*Sin[e + f*x])^(m - 1)*((b*Sec[e + f*x])^(n

- 1)/(f*(m - n))), x] + Dist[a^2*((m - 1)/(m - n)), Int[(a*Sin[e + f*x])^(m - 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && GtQ[m, 1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_)*sec[(e_.) + (f_)*(x_)])^(n_)*((a_)*sin[(e_.) + (f_)*(x_)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{(a \sin(e + fx))^{3/2}}{(b \sec(e + fx))^{3/2}} dx &= \frac{(a \sin(e + fx))^{5/2}}{3abf \sqrt{b \sec(e + fx)}} + \frac{\int \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2} dx}{6b^2} \\
 &= -\frac{a \sqrt{a \sin(e + fx)}}{6bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{5/2}}{3abf \sqrt{b \sec(e + fx)}} + \frac{a^2 \int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx}{12b^2} \\
 &= -\frac{a \sqrt{a \sin(e + fx)}}{6bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{5/2}}{3abf \sqrt{b \sec(e + fx)}} + \frac{(a^2 \sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)})}{12b^2} \\
 &= -\frac{a \sqrt{a \sin(e + fx)}}{6bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{5/2}}{3abf \sqrt{b \sec(e + fx)}} + \frac{(a^2 \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)})}{12b^2 \sqrt{a \sin(e + fx)}} \\
 &= -\frac{a \sqrt{a \sin(e + fx)}}{6bf \sqrt{b \sec(e + fx)}} + \frac{(a \sin(e + fx))^{5/2}}{3abf \sqrt{b \sec(e + fx)}} + \frac{a^2 F(e - \frac{\pi}{4} + fx | 2) \sqrt{b \sec(e + fx)}}{12b^2 f \sqrt{a \sin(e + fx)}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 0.31, size = 87, normalized size = 0.64

$$\frac{a \sqrt{a \sin(e + fx)} \left(-2 \cos(2(e + fx)) + \csc^2(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4} \right)}{12bf \sqrt{b \sec(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(a*Sin[e + f*x])^(3/2)/(b*Sec[e + f*x])^(3/2),x]

[Out] (a*Sqrt[a*Sin[e + f*x]]*(-2*Cos[2*(e + f*x)] + Csc[e + f*x]^2*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2*(-Tan[e + f*x]^2)^(3/4)))/(12*b*f*Sqrt[b*Sec[e + f*x]])

Maple [A]

time = 0.26, size = 218, normalized size = 1.61

method	result
default	$-\frac{\left(2(\cos^4(fx+e))\sqrt{2} + \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\right) \text{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)}{\sin(fx+e)}}\right)}{12f(-1+\cos(fx+e))\sin(fx+e)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(f*x+e))^(3/2)/(b*sec(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] -1/12/f*(2*cos(f*x+e)^4*2^(1/2)+((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2))*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)-2*cos(f*x+e)^3*2^(1/2)-cos(f*x+e)^2*2^(1/2)+cos(f*x+e)*2^(1/2))*(a*sin(f*x+e))^(3/2)/(-1+cos(f*x+e))/sin(f*x+e)/(b/cos(f*x+e))^(3/2)/cos(f*x+e)^2*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(3/2)/(b*sec(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((a*sin(f*x + e))^(3/2)/(b*sec(f*x + e))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(3/2)/(b*sec(f*x+e))^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e))*a*sin(f*x + e)/(b^2*sec(f*x + e)^2), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))**(3/2)/(b*sec(f*x+e))**(3/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3435 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a*sin(f*x+e))^(3/2)/(b*sec(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((a*sin(f*x + e))^(3/2)/(b*sec(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(a \sin(e + f x))^{3/2}}{\left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^(3/2)/(b/cos(e + f*x))^(3/2),x)

[Out] int((a*sin(e + f*x))^(3/2)/(b/cos(e + f*x))^(3/2), x)

$$3.479 \quad \int \frac{1}{(b \sec(e+fx))^{3/2} \sqrt{a \sin(e+fx)}} dx$$

Optimal. Leaf size=94

$$\frac{\sqrt{a \sin(e+fx)}}{abf \sqrt{b \sec(e+fx)}} + \frac{F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}{2b^2 f \sqrt{a \sin(e+fx)}}$$

[Out] (a*sin(f*x+e))^(1/2)/a/b/f/(b*sec(f*x+e))^(1/2)-1/2*(sin(e+1/4*Pi+f*x)^2)^(1/2)/sin(e+1/4*Pi+f*x)*EllipticF(cos(e+1/4*Pi+f*x),2^(1/2))*(b*sec(f*x+e))^(1/2)*sin(2*f*x+2*e)^(1/2)/b^2/f/(a*sin(f*x+e))^(1/2)

Rubi [A]

time = 0.11, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2662, 2665, 2653, 2720}

$$\frac{\sqrt{\sin(2e+2fx)} F\left(e+fx-\frac{\pi}{4} \mid 2\right) \sqrt{b \sec(e+fx)}}{2b^2 f \sqrt{a \sin(e+fx)}} + \frac{\sqrt{a \sin(e+fx)}}{abf \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((b*Sec[e + f*x])^(3/2)*Sqrt[a*Sin[e + f*x]]),x]

[Out] Sqrt[a*Sin[e + f*x]]/(a*b*f*Sqrt[b*Sec[e + f*x]]) + (EllipticF[e - Pi/4 + f*x, 2]*Sqrt[b*Sec[e + f*x]]*Sqrt[Sin[2*e + 2*f*x]])/(2*b^2*f*Sqrt[a*Sin[e + f*x]])

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2662

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1))/(a*b*f*(m - n)), x] - Dist[(n + 1)/(b^2*(m - n)), Int[(a*Sin[e + f*x])^m*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f, m}, x] && LtQ[n, -1] && NeQ[m - n, 0] && IntegersQ[2*m, 2*n]

Rule 2665

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +

$f*x])^m/(b*\text{Cos}[e + f*x])^n, x], x] /; \text{FreeQ}[\{a, b, e, f, m, n\}, x] \ \&\& \ \text{IntegerQ}[m - 1/2] \ \&\& \ \text{IntegerQ}[n - 1/2]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_.)]], x_Symbol] := \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \sec(e + fx))^{3/2} \sqrt{a \sin(e + fx)}} dx &= \frac{\sqrt{a \sin(e + fx)}}{abf \sqrt{b \sec(e + fx)}} + \frac{\int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx}{2b^2} \\ &= \frac{\sqrt{a \sin(e + fx)}}{abf \sqrt{b \sec(e + fx)}} + \frac{\left(\sqrt{b \cos(e + fx)} \sqrt{b \sec(e + fx)}\right) \int \frac{1}{\sqrt{b \cos(e + fx)}} dx}{2b^2} \\ &= \frac{\sqrt{a \sin(e + fx)}}{abf \sqrt{b \sec(e + fx)}} + \frac{\left(\sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}\right) \int \frac{1}{\sqrt{\sin(2e + 2fx)}} dx}{2b^2 \sqrt{a \sin(e + fx)}} \\ &= \frac{\sqrt{a \sin(e + fx)}}{abf \sqrt{b \sec(e + fx)}} + \frac{F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b \sec(e + fx)} \sqrt{\sin(2e + 2fx)}}{2b^2 f \sqrt{a \sin(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.41, size = 84, normalized size = 0.89

$$\frac{\cot(e + fx) \sqrt{b \sec(e + fx)} \left(-1 + \cos(2(e + fx)) - {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4}\right)}{2b^2 f \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((b*Sec[e + f*x])^(3/2)*Sqrt[a*Sin[e + f*x]]),x]

[Out] -1/2*(Cot[e + f*x]*Sqrt[b*Sec[e + f*x]]*(-1 + Cos[2*(e + f*x)] - Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4)))/(b^2*f*Sqrt[a*Sin[e + f*x]])

Maple [A]

time = 0.27, size = 190, normalized size = 2.02

method	result
--------	--------

default	$-\frac{\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\operatorname{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}},\frac{b}{\cos(fx+e)}\right)\right)^{\frac{3}{2}}}{2f(-1+\cos(fx+e))\cos(fx+e)^2\sqrt{a\sin(fx+e)}}$
---------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
[Out] -1/2/f*(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)-cos(f*x+e)^2*2^(1/2)+cos(f*x+e)*2^(1/2))*sin(f*x+e)/(-1+cos(f*x+e))/cos(f*x+e)^2/(b/cos(f*x+e))^(3/2)/(a*sin(f*x+e))^(1/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(1/2),x, algorithm="maxima")
[Out] integrate(1/((b*sec(f*x + e))^(3/2)*sqrt(a*sin(f*x + e))), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(1/2),x, algorithm="fricas")
[Out] integral(sqrt(b*sec(f*x + e))*sqrt(a*sin(f*x + e))/(a*b^2*sec(f*x + e)^2*sin(f*x + e)), x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))**(3/2)/(a*sin(f*x+e))**(1/2),x)
```

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*sqrt(a*sin(f*x + e))), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sqrt{a \sin(e + f x)} \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*sin(e + f*x))^(1/2)*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/((a*sin(e + f*x))^(1/2)*(b/cos(e + f*x))^(3/2)), x)

$$3.480 \quad \int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{5/2}} dx$$

Optimal. Leaf size=100

$$-\frac{2}{3abf \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{3/2}} \frac{F\left(e - \frac{\pi}{4} + fx \mid 2\right) \sqrt{b \sec(e+fx)} \sqrt{\sin(2e+2fx)}}{3a^2b^2f \sqrt{a \sin(e+fx)}}$$

[Out] $-2/3/a/b/f/(a*\sin(f*x+e))^{(3/2)}/(b*\sec(f*x+e))^{(1/2)}+1/3*(\sin(e+1/4*Pi+f*x))^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticF}(\cos(e+1/4*Pi+f*x), 2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/a^2/b^2/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {2661, 2665, 2653, 2720}

$$-\frac{\sqrt{\sin(2e+2fx)} F\left(e+fx-\frac{\pi}{4} \mid 2\right) \sqrt{b \sec(e+fx)}}{3a^2b^2f \sqrt{a \sin(e+fx)}} - \frac{2}{3abf (a \sin(e+fx))^{3/2} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/((b*\text{Sec}[e+f*x])^{(3/2)}*(a*\text{Sin}[e+f*x])^{(5/2)}),x]$

[Out] $-2/(3*a*b*f*\text{Sqrt}[b*\text{Sec}[e+f*x]]*(a*\text{Sin}[e+f*x])^{(3/2)}) - (\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e+f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(3*a^2*b^2*f*\text{Sqrt}[a*\text{Sin}[e+f*x]])$

Rule 2653

$\text{Int}[1/(\text{Sqrt}[\cos[(e_.) + (f_.)*(x_.)]*(b_.)]*\text{Sqrt}[(a_.)*\sin[(e_.) + (f_.)*(x_.)])], x_Symbol] \rightarrow \text{Dist}[\text{Sqrt}[\text{Sin}[2*e + 2*f*x]]/(\text{Sqrt}[a*\text{Sin}[e+f*x]]*\text{Sqrt}[b*\text{Cos}[e+f*x]]), \text{Int}[1/\text{Sqrt}[\text{Sin}[2*e + 2*f*x]], x], x] /; \text{FreeQ}\{a, b, e, f\}, x]$

Rule 2661

$\text{Int}[(b_.)*\sec[(e_.) + (f_.)*(x_.)]^{(n)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m)}], x_Symbol] \rightarrow \text{Simp}[(a*\text{Sin}[e+f*x])^{(m+1)}*((b*\text{Sec}[e+f*x])^{(n+1)})/(a*b*f*(m+1)), x] - \text{Dist}[(n+1)/(a^2*b^2*(m+1)), \text{Int}[(a*\text{Sin}[e+f*x])^{(m+2)}*(b*\text{Sec}[e+f*x])^{(n+2)}, x], x] /; \text{FreeQ}\{a, b, e, f\}, x] \&\& \text{LtQ}[n, -1] \&\& \text{LtQ}[m, -1] \&\& \text{IntegersQ}[2*m, 2*n]$

Rule 2665

$\text{Int}[(b_.)*\sec[(e_.) + (f_.)*(x_.)]^{(n)}*((a_.)*\sin[(e_.) + (f_.)*(x_.)])^{(m)}], x_Symbol] \rightarrow \text{Dist}[(b*\text{Cos}[e+f*x])^{(n)}*(b*\text{Sec}[e+f*x])^{(n)}, \text{Int}[(a*\text{Sin}[e+f*x])^{(m)}/(b*\text{Cos}[e+f*x])^{(n)}, x], x] /; \text{FreeQ}\{a, b, e, f, m, n\}, x] \&\& \text{Inte}$

gerQ[m - 1/2] && IntegerQ[n - 1/2]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \sec(e + fx))^{3/2} (a \sin(e + fx))^{5/2}} dx &= -\frac{2}{3abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} - \frac{\int \frac{\sqrt{b \sec(e + fx)}}{\sqrt{a \sin(e + fx)}} dx}{3a^2b^2} \\ &= -\frac{2}{3abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} - \frac{\left(\sqrt{b \cos(e + fx)} \sqrt{a \sin(e + fx)}\right)}{3a^2b^2} \\ &= -\frac{2}{3abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} - \frac{\left(\sqrt{b \sec(e + fx)} \sqrt{a \sin(e + fx)}\right)}{3a^2b^2} \\ &= -\frac{2}{3abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{3/2}} - \frac{F\left(e - \frac{\pi}{4} + fx \mid 2\right)}{3a^2b^2} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.38, size = 78, normalized size = 0.78

$$\frac{\cot(e + fx) \sqrt{b \sec(e + fx)} \left(2 + {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4}\right)}{3a^2b^2 f \sqrt{a \sin(e + fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(5/2)),x]

[Out] -1/3*(Cot[e + f*x]*Sqrt[b*Sec[e + f*x]]*(2 + Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4)))/(a^2*b^2*f*Sqrt[a*Sin[e + f*x]])

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 283 vs. 2(111) = 222.

time = 0.20, size = 284, normalized size = 2.84

method	result
default	$-\frac{\cos(fx+e)\sin(fx+e)\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}}\text{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(5/2),x,method=_RETURNVERBOSE)
[Out] -1/3/f*(cos(f*x+e)*sin(f*x+e)*((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*
((sin(f*x+e)-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(
1/2)*EllipticF(((1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+((
1-cos(f*x+e)+sin(f*x+e))/sin(f*x+e))^(1/2)*((sin(f*x+e)-1+cos(f*x+e))/sin(f
*x+e))^(1/2)*((-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((1-cos(f*x+e)+si
n(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)+cos(f*x+e)*2^(1/2))*sin
(f*x+e)/(b/cos(f*x+e))^(3/2)/(a*sin(f*x+e))^(5/2)/cos(f*x+e)^2*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(5/2),x, algorithm="maxima"
)
```

```
[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(5/2)), x)
```

Fricas [C] Result contains complex when optimal does not.

time = 0.12, size = 139, normalized size = 1.39

$$\frac{\sqrt{iab}(\cos(fx+e)^2-1)\text{ellipticF}(\cos(fx+e)+i\sin(fx+e),-1)+\sqrt{-iab}(\cos(fx+e)^2-1)\text{ellipticF}(\cos(fx+e)-i\sin(fx+e),-1)+2\sqrt{a\sin(fx+e)}\sqrt{\frac{b}{\cos(fx+e)}}\cos(fx+e)}{3(a^3b^2f\cos(fx+e)^2-a^3b^2f)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(5/2),x, algorithm="fricas"
)
```

```
[Out] 1/3*(sqrt(I*a*b)*(cos(f*x + e)^2 - 1)*ellipticF(cos(f*x + e) + I*sin(f*x +
e), -1) + sqrt(-I*a*b)*(cos(f*x + e)^2 - 1)*ellipticF(cos(f*x + e) - I*sin(
f*x + e), -1) + 2*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e))*cos(f*x + e))/(
a^3*b^2*f*cos(f*x + e)^2 - a^3*b^2*f)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))**(3/2)/(a*sin(f*x+e))**(5/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(5/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(5/2)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a \sin(e + f x))^{5/2} \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*sin(e + f*x))^(5/2)*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/((a*sin(e + f*x))^(5/2)*(b/cos(e + f*x))^(3/2)), x)

$$3.481 \quad \int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{9/2}} dx$$

Optimal. Leaf size=137

$$-\frac{2}{7abf \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{7/2}} + \frac{2}{21a^3bf \sqrt{b \sec(e+fx)} (a \sin(e+fx))^{3/2}} - \frac{2F\left(e - \frac{\pi}{4} + fx \mid 2\right)}{21a^4b^2}$$

[Out] $-2/7/a/b/f/(a*\sin(f*x+e))^{(7/2)}/(b*\sec(f*x+e))^{(1/2)}+2/21/a^3/b/f/(a*\sin(f*x+e))^{(3/2)}/(b*\sec(f*x+e))^{(1/2)}+2/21*(\sin(e+1/4*\text{Pi}+f*x)^2)^{(1/2)}/\sin(e+1/4*\text{Pi}+f*x)*\text{EllipticF}(\cos(e+1/4*\text{Pi}+f*x),2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/a^4/b^2/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2661, 2664, 2665, 2653, 2720}

$$-\frac{2\sqrt{\sin(2e+2fx)} F\left(e+fx-\frac{\pi}{4} \mid 2\right) \sqrt{b \sec(e+fx)}}{21a^4b^2f \sqrt{a \sin(e+fx)}} + \frac{2}{21a^3bf (a \sin(e+fx))^{3/2} \sqrt{b \sec(e+fx)}} - \frac{2}{7abf (a \sin(e+fx))^{7/2} \sqrt{b \sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(9/2)),x]

[Out] $-2/(7*a*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(7/2)}) + 2/(21*a^3*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(3/2)}) - (2*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(21*a^4*b^2*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2661

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] - Dist[(n + 1)/(a^2*b^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && LtQ[n, -1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2664

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/

```
(a*f*(m + 1)), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n]
```

Rule 2665

```
Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && IntegerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \sec(e + fx))^{3/2} (a \sin(e + fx))^{9/2}} dx &= -\frac{2}{7abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} - \frac{\int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{5/2}} dx}{7a^2b^2} \\ &= -\frac{2}{7abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} + \frac{2}{21a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{7abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} + \frac{2}{21a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{7abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} + \frac{2}{21a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{7abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{7/2}} + \frac{2}{21a^3bf \sqrt{b \sec(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.64, size = 119, normalized size = 0.87

$$\frac{\cos(2(e + fx)) \csc^4(e + fx) \sqrt{a \sin(e + fx)} \left((5 + \cos(2(e + fx))) \sec^2(e + fx) - {}_2F_1\left(\frac{1}{2}, \frac{3}{4}; \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{7/4} \right)}{21a^5bf \sqrt{b \sec(e + fx)} (-2 + \sec^2(e + fx))}$$

Antiderivative was successfully verified.

[In] Integrate[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(9/2)),x]

[Out] (Cos[2*(e + f*x)]*Csc[e + f*x]^4*Sqrt[a*Sin[e + f*x]]*((5 + Cos[2*(e + f*x)])*Sec[e + f*x]^2 - 2*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(7/4)))/(21*a^5*b*f*Sqrt[b*Sec[e + f*x]]*(-2 + Sec[e + f*x]^2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 538 vs. $2(142) = 284$.

time = 0.23, size = 539, normalized size = 3.93

method	result
default	$-2 \sqrt{\frac{1 - \cos(fx+e) + \sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e) - 1 + \cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1 + \cos(fx+e)}{\sin(fx+e)}} \text{EllipticF}\left(\sqrt{\frac{1 - \cos(fx+e) + \sin(fx+e)}{\sin(fx+e)}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(9/2),x,method=_RETURNVERBOSE)

[Out]
$$-1/21/f*(-2*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))*\sin(f*x+e)*\cos(f*x+e)^3-2*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))*\sin(f*x+e)*\cos(f*x+e)^2+2*\cos(f*x+e)*\sin(f*x+e)*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))+2*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))*\sin(f*x+e)+\cos(f*x+e)^3*2^{1/2}+2*\cos(f*x+e)*2^{1/2})*\sin(f*x+e)/\cos(f*x+e)^2/(b/\cos(f*x+e))^{3/2}/(a*\sin(f*x+e))^{9/2}*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(9/2),x, algorithm="maxima")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(9/2)), x)

Fricas [C] Result contains complex when optimal does not.

time = 0.12, size = 190, normalized size = 1.39

$$\frac{2 \left((\cos(fx+e)^4 - 2\cos(fx+e)^2 + 1) \sqrt{ab} \operatorname{ellipticF}(\cos(fx+e) + i \sin(fx+e), -1) + (\cos(fx+e)^4 - 2\cos(fx+e)^2 + 1) \sqrt{-iab} \operatorname{ellipticF}(\cos(fx+e) - i \sin(fx+e), -1) - (\cos(fx+e)^3 + 2\cos(fx+e)) \sqrt{a \sin(fx+e)} \sqrt{\frac{b}{\cos(fx+e)}} \right)}{21 (a^2 b^2 f \cos(fx+e)^4 - 2 a^2 b^2 f \cos(fx+e)^2 + a^2 b^2 f)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(9/2),x, algorithm="fricas")

[Out] 2/21*((cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(I*a*b)*ellipticF(cos(f*x + e) + I*sin(f*x + e), -1) + (cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*sqrt(-I*a*b)*ellipticF(cos(f*x + e) - I*sin(f*x + e), -1) - (cos(f*x + e)^3 + 2*cos(f*x + e))*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e)))/(a^5*b^2*f*cos(f*x + e)^4 - 2*a^5*b^2*f*cos(f*x + e)^2 + a^5*b^2*f)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))**(3/2)/(a*sin(f*x+e))**(9/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(9/2),x, algorithm="giac")

[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(9/2)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a \sin(e + f x))^{9/2} \left(\frac{b}{\cos(e + f x)} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*sin(e + f*x))^(9/2)*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/((a*sin(e + f*x))^(9/2)*(b/cos(e + f*x))^(3/2)), x)

$$3.482 \quad \int \frac{1}{(b \sec(e+fx))^{3/2} (a \sin(e+fx))^{13/2}} dx$$

Optimal. Leaf size=174

$$-\frac{2}{11abf\sqrt{b\sec(e+fx)}(a\sin(e+fx))^{11/2}} + \frac{2}{77a^3bf\sqrt{b\sec(e+fx)}(a\sin(e+fx))^{7/2}} + \frac{2}{77a^5bf\sqrt{b\sec(e+fx)}(a\sin(e+fx))^{3/2}}$$

[Out] $-2/11/a/b/f/(a*\sin(f*x+e))^{(11/2)}/(b*\sec(f*x+e))^{(1/2)}+2/77/a^3/b/f/(a*\sin(f*x+e))^{(7/2)}/(b*\sec(f*x+e))^{(1/2)}+4/77/a^5/b/f/(a*\sin(f*x+e))^{(3/2)}/(b*\sec(f*x+e))^{(1/2)}+4/77*(\sin(e+1/4*Pi+f*x)^2)^{(1/2)}/\sin(e+1/4*Pi+f*x)*\text{EllipticF}(\cos(e+1/4*Pi+f*x), 2^{(1/2)})*(b*\sec(f*x+e))^{(1/2)}*\sin(2*f*x+2*e)^{(1/2)}/a^6/b^2/f/(a*\sin(f*x+e))^{(1/2)}$

Rubi [A]

time = 0.20, antiderivative size = 174, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {2661, 2664, 2665, 2653, 2720}

$$-\frac{4\sqrt{\sin(2e+2fx)}F(e+fx-\frac{\pi}{4}|2)\sqrt{b\sec(e+fx)}}{77a^6b^2f\sqrt{a\sin(e+fx)}} + \frac{4}{77a^5bf(a\sin(e+fx))^{3/2}\sqrt{b\sec(e+fx)}} + \frac{2}{77a^3bf(a\sin(e+fx))^{7/2}\sqrt{b\sec(e+fx)}} - \frac{2}{11abf(a\sin(e+fx))^{11/2}\sqrt{b\sec(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(13/2)),x]

[Out] $-2/(11*a*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(11/2)}) + 2/(77*a^3*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(7/2)}) + 4/(77*a^5*b*f*\text{Sqrt}[b*\text{Sec}[e + f*x]]*(a*\text{Sin}[e + f*x])^{(3/2)}) - (4*\text{EllipticF}[e - \text{Pi}/4 + f*x, 2]*\text{Sqrt}[b*\text{Sec}[e + f*x]]*\text{Sqrt}[\text{Sin}[2*e + 2*f*x]])/(77*a^6*b^2*f*\text{Sqrt}[a*\text{Sin}[e + f*x]])$

Rule 2653

Int[1/(Sqrt[cos[(e_.) + (f_.)*(x_.)]*(b_.)]*Sqrt[(a_.)*sin[(e_.) + (f_.)*(x_.)]]), x_Symbol] :> Dist[Sqrt[Sin[2*e + 2*f*x]]/(Sqrt[a*Sin[e + f*x]]*Sqrt[b*Cos[e + f*x]]), Int[1/Sqrt[Sin[2*e + 2*f*x]], x], x] /; FreeQ[{a, b, e, f}, x]

Rule 2661

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] :> Simp[(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n + 1)/(a*b*f*(m + 1))), x] - Dist[(n + 1)/(a^2*b^2*(m + 1)), Int[(a*Sin[e + f*x])^(m + 2)*(b*Sec[e + f*x])^(n + 2), x], x] /; FreeQ[{a, b, e, f}, x] && LtQ[n, -1] && LtQ[m, -1] && IntegersQ[2*m, 2*n]

Rule 2664

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Simp[b*(a*Sin[e + f*x])^(m + 1)*((b*Sec[e + f*x])^(n - 1)/
(a*f*(m + 1))), x] + Dist[(m - n + 2)/(a^2*(m + 1)), Int[(a*Sin[e + f*x])^(
m + 2)*(b*Sec[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, n}, x] && LtQ[m, -1
] && IntegersQ[2*m, 2*n]
```

Rule 2665

```
Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_)])^(m
_), x_Symbol] := Dist[(b*Cos[e + f*x])^n*(b*Sec[e + f*x])^n, Int[(a*Sin[e +
f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && Inte
gerQ[m - 1/2] && IntegerQ[n - 1/2]
```

Rule 2720

```
Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2
)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{1}{(b \sec(e + fx))^{3/2} (a \sin(e + fx))^{13/2}} dx &= -\frac{2}{11abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{11/2}} - \frac{\int \frac{\sqrt{b \sec(e + fx)}}{(a \sin(e + fx))^{9/2}}}{11a^2b^2} \\ &= -\frac{2}{11abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{11/2}} + \frac{2}{77a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{11abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{11/2}} + \frac{2}{77a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{11abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{11/2}} + \frac{2}{77a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{11abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{11/2}} + \frac{2}{77a^3bf \sqrt{b \sec(e + fx)}} \\ &= -\frac{2}{11abf \sqrt{b \sec(e + fx)} (a \sin(e + fx))^{11/2}} + \frac{2}{77a^3bf \sqrt{b \sec(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 4 in optimal.

time = 10.88, size = 131, normalized size = 0.75

$$\frac{2 \cot(2(e + fx)) \csc(2(e + fx)) \sqrt{a \sin(e + fx)} \left((23 + 6 \cos(2(e + fx)) - \cos(4(e + fx))) \csc^4(e + fx) + 8 {}_2F_1\left(\frac{1}{2}, \frac{3}{4}, \frac{3}{2}; \sec^2(e + fx)\right) (-\tan^2(e + fx))^{3/4} \right)}{77a^7bf\sqrt{b\sec(e + fx)}(-2 + \sec^2(e + fx))}$$

Antiderivative was successfully verified.

[In] Integrate[1/((b*Sec[e + f*x])^(3/2)*(a*Sin[e + f*x])^(13/2)),x]

[Out] (2*Cot[2*(e + f*x)]*Csc[2*(e + f*x)]*Sqrt[a*Sin[e + f*x]]*((23 + 6*Cos[2*(e + f*x)] - Cos[4*(e + f*x)])*Csc[e + f*x]^4 + 8*Hypergeometric2F1[1/2, 3/4, 3/2, Sec[e + f*x]^2]*(-Tan[e + f*x]^2)^(3/4)))/(77*a^7*b*f*Sqrt[b*Sec[e + f*x]]*(-2 + Sec[e + f*x]^2))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 792 vs. $2(173) = 346$.

time = 0.26, size = 793, normalized size = 4.56

method	result
default	$-\frac{\left(4(\cos^5(fx+e)) \sin(fx+e) \sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{\sin(fx+e)-1+\cos(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-1+\cos(fx+e)}{\sin(fx+e)}} \text{EllipticF}\left(\sqrt{\frac{1-\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(13/2),x,method=_RETURNVERBOSE)

[Out]
$$-1/77/f*(4*\cos(f*x+e)^5*\sin(f*x+e)*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2})))+4*\cos(f*x+e)^4*\sin(f*x+e)*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))-8*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))*\sin(f*x+e)*\cos(f*x+e)^3-8*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))*\sin(f*x+e)*\cos(f*x+e)^2-2*2^{1/2}*\cos(f*x+e)^5+4*\cos(f*x+e)*\sin(f*x+e)*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))+4*((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2}*((\sin(f*x+e)-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*\text{EllipticF}(((1-\cos(f*x+e)+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2}))*\sin(f*x+e)+5*\cos(f*x+e)^3*2^{1/2}+4*\cos(f*x+e)*2^{1/2}*\sin(f*x+e)/\cos(f*x+e)^2/(b/\cos(f*x+e))^{3/2}/(a*\sin(f*x+e))^{13/2}*2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(13/2),x, algorithm="maxima")
```

```
[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(13/2)), x)
```

Fricas [C] Result contains complex when optimal does not.

time = 0.12, size = 246, normalized size = 1.41

$$\frac{2 \left(2 (\cos(fx+e)^6 - 3 \cos(fx+e)^4 + 3 \cos(fx+e)^2 - 1) \sqrt{ab} \operatorname{ellipticF}(\cos(fx+e) + i \sin(fx+e), -1) + 2 (\cos(fx+e)^6 - 3 \cos(fx+e)^4 + 3 \cos(fx+e)^2 - 1) \sqrt{-1ab} \operatorname{ellipticF}(\cos(fx+e) - i \sin(fx+e), -1) - (2 \cos(fx+e)^5 - 5 \cos(fx+e)^3 - 4 \cos(fx+e)) \sqrt{a \sin(fx+e)} \sqrt{\frac{b}{\cos(fx+e)}} \right)}{77 (a^{10} b^2 \cos(fx+e)^6 - 3 a^{10} b^2 \cos(fx+e)^4 + 3 a^{10} b^2 \cos(fx+e)^2 - a^{10} b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(13/2),x, algorithm="fricas")
```

```
[Out] 2/77*(2*(cos(f*x + e)^6 - 3*cos(f*x + e)^4 + 3*cos(f*x + e)^2 - 1)*sqrt(I*a*b)*ellipticF(cos(f*x + e) + I*sin(f*x + e), -1) + 2*(cos(f*x + e)^6 - 3*cos(f*x + e)^4 + 3*cos(f*x + e)^2 - 1)*sqrt(-I*a*b)*ellipticF(cos(f*x + e) - I*sin(f*x + e), -1) - (2*cos(f*x + e)^5 - 5*cos(f*x + e)^3 - 4*cos(f*x + e))*sqrt(a*sin(f*x + e))*sqrt(b/cos(f*x + e)))/(a^7*b^2*f*cos(f*x + e)^6 - 3*a^7*b^2*f*cos(f*x + e)^4 + 3*a^7*b^2*f*cos(f*x + e)^2 - a^7*b^2*f)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(13/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(b*sec(f*x+e))^(3/2)/(a*sin(f*x+e))^(13/2),x, algorithm="giac")
```

```
[Out] integrate(1/((b*sec(f*x + e))^(3/2)*(a*sin(f*x + e))^(13/2)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{(a \sin(e + f x))^{13/2} \left(\frac{b}{\cos(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*sin(e + f*x))^(13/2)*(b/cos(e + f*x))^(3/2)),x)

[Out] int(1/((a*sin(e + f*x))^(13/2)*(b/cos(e + f*x))^(3/2)), x)

3.483 $\int (d \sec(a + bx))^{5/2} (c \sin(a + bx))^m dx$

Optimal. Leaf size=75

$$\frac{d \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (d \sec(a + bx))^{3/2} (c \sin(a + bx))^{1+m}}{bc(1+m)}$$

[Out] d*(cos(b*x+a)^2)^(3/4)*hypergeom([7/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2) *(d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^(1+m)/b/c/(1+m)

Rubi [A]

time = 0.08, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2657}

$$\frac{d \cos^2(a + bx)^{3/4} (d \sec(a + bx))^{3/2} (c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{7}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(d*Sec[a + b*x])^(5/2)*(c*Sin[a + b*x])^m,x]

[Out] (d*(Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[7/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(d*Sec[a + b*x])^(3/2)*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m], x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m], x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (d \sec(a + bx))^{5/2} (c \sin(a + bx))^m dx &= (d^2 (d \cos(a + bx))^{3/2} (d \sec(a + bx))^{3/2}) \int \frac{(c \sin(a + bx))^m}{(d \cos(a + bx))^{5/2}} dx \\ &= \frac{d \cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{7}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (d \sec(a + bx))^{3/2}}{bc(1+m)} \end{aligned}$$

Mathematica [A]

time = 9.75, size = 96, normalized size = 1.28

$$\frac{2 \cot(a + bx) {}_2F_1\left(\frac{1}{4}(5 - 2m), \frac{1-m}{2}; \frac{1}{4}(9 - 2m); \sec^2(a + bx)\right) (d \sec(a + bx))^{5/2} (c \sin(a + bx))^m (-\tan^2(a + bx))^{\frac{1-m}{2}}}{b(-5 + 2m)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(5/2)*(c*Sin[a + b*x])^m,x]

[Out] (-2*Cot[a + b*x]*Hypergeometric2F1[(5 - 2*m)/4, (1 - m)/2, (9 - 2*m)/4, Sec[a + b*x]^2]*(d*Sec[a + b*x])^(5/2)*(c*Sin[a + b*x])^m*(-Tan[a + b*x]^2)^((1 - m)/2))/(b*(-5 + 2*m))

Maple [F]

time = 0.14, size = 0, normalized size = 0.00

$$\int (d \sec(bx + a))^{\frac{5}{2}} (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(5/2)*(c*sin(b*x+a))^m,x)

[Out] int((d*sec(b*x+a))^(5/2)*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((d*sec(b*x + a))^(5/2)*(c*sin(b*x + a))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m*d^2*sec(b*x + a)^2, x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))**(5/2)*(c*sin(b*x+a))**m,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(5/2)*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate((d*sec(b*x + a))^(5/2)*(c*sin(b*x + a))^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + b x))^m \left(\frac{d}{\cos(a + b x)} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m*(d/cos(a + b*x))^(5/2),x)

[Out] int((c*sin(a + b*x))^m*(d/cos(a + b*x))^(5/2), x)

3.484 $\int (d \sec(a + bx))^{3/2} (c \sin(a + bx))^m dx$

Optimal. Leaf size=75

$$\frac{d^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sqrt{d \sec(a + bx)} (c \sin(a + bx))^{1+m}}{bc(1+m)}$$

[Out] d*(cos(b*x+a)^2)^(1/4)*hypergeom([5/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)
*(c*sin(b*x+a))^(1+m)*(d*sec(b*x+a))^(1/2)/b/c/(1+m)

Rubi [A]

time = 0.07, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2657}

$$\frac{d^4 \sqrt{\cos^2(a + bx)} \sqrt{d \sec(a + bx)} (c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{5}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bc(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(d*Sec[a + b*x])^(3/2)*(c*Sin[a + b*x])^m,x]

[Out] (d*(Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[5/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*Sqrt[d*Sec[a + b*x]]*(c*Sin[a + b*x])^(1 + m))/(b*c*(1 + m))

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_.))^(n_)*((a_.)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2667

Int[((b_.)*sec[(e_) + (f_)*(x_)])^(n_)*((a_.)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (d \sec(a + bx))^{3/2} (c \sin(a + bx))^m dx &= \left(d^2 \sqrt{d \cos(a + bx)} \sqrt{d \sec(a + bx)} \right) \int \frac{(c \sin(a + bx))^m}{(d \cos(a + bx))^{3/2}} dx \\ &= \frac{d^4 \sqrt{\cos^2(a + bx)} {}_2F_1\left(\frac{5}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sqrt{d \sec(a + bx)}}{bc(1+m)} \end{aligned}$$

Mathematica [A]

time = 5.86, size = 96, normalized size = 1.28

$$\frac{2 \cot(a + bx) {}_2F_1\left(\frac{1}{4}(3 - 2m), \frac{1-m}{2}; \frac{1}{4}(7 - 2m); \sec^2(a + bx)\right) (d \sec(a + bx))^{3/2} (c \sin(a + bx))^m (-\tan^2(a + bx))^{\frac{1-m}{2}}}{b(-3 + 2m)}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Sec[a + b*x])^(3/2)*(c*Sin[a + b*x])^m,x]

[Out] (-2*Cot[a + b*x]*Hypergeometric2F1[(3 - 2*m)/4, (1 - m)/2, (7 - 2*m)/4, Sec[a + b*x]^2]*(d*Sec[a + b*x])^(3/2)*(c*Sin[a + b*x])^m*(-Tan[a + b*x]^2)^((1 - m)/2))/(b*(-3 + 2*m))

Maple [F]

time = 0.12, size = 0, normalized size = 0.00

$$\int (d \sec(bx + a))^{\frac{3}{2}} (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x)

[Out] int((d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate((d*sec(b*x + a))^(3/2)*(c*sin(b*x + a))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m*d*sec(b*x + a), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*sec(b*x+a))**(3/2)*(c*sin(b*x+a))**m,x)`

[Out] Exception raised: SystemError >> excessive stack use: stack is 6437 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^m,x, algorithm="giac")`

[Out] `integrate((d*sec(b*x + a))^(3/2)*(c*sin(b*x + a))^m, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^m \left(\frac{d}{\cos(a + bx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(a + b*x))^m*(d/cos(a + b*x))^(3/2),x)`

[Out] `int((c*sin(a + b*x))^m*(d/cos(a + b*x))^(3/2), x)`

3.485 $\int \sqrt{d \sec(a + bx)} (c \sin(a + bx))^m dx$

Optimal. Leaf size=77

$$\frac{\cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (d \sec(a + bx))^{3/2} (c \sin(a + bx))^{1+m}}{bcd(1 + m)}$$

[Out] (cos(b*x+a)^2)^(3/4)*hypergeom([3/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(d*sec(b*x+a))^(3/2)*(c*sin(b*x+a))^(1+m)/b/c/d/(1+m)

Rubi [A]

time = 0.06, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$,

Rules used = {2666, 2657}

$$\frac{\cos^2(a + bx)^{3/4} (d \sec(a + bx))^{3/2} (c \sin(a + bx))^{m+1} {}_2F_1\left(\frac{3}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a + bx)\right)}{bcd(m + 1)}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Sec[a + b*x]]*(c*Sin[a + b*x])^m,x]

[Out] ((Cos[a + b*x]^2)^(3/4)*Hypergeometric2F1[3/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(d*Sec[a + b*x])^(3/2)*(c*Sin[a + b*x])^(1 + m))/(b*c*d*(1 + m))

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^n]*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2666

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^n]*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] := Dist[(1/b^2)*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \sqrt{d \sec(a + bx)} (c \sin(a + bx))^m dx &= \frac{((d \cos(a + bx))^{3/2} (d \sec(a + bx))^{3/2}) \int \frac{(c \sin(a + bx))^m}{\sqrt{d \cos(a + bx)}} dx}{d^2} \\ &= \frac{\cos^2(a + bx)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) (d \sec(a + bx))^{3/2} (c \sin(a + bx))^{1+m}}{bcd(1 + m)} \end{aligned}$$

Mathematica [A]

time = 5.69, size = 106, normalized size = 1.38

$$\frac{\csc^2(a + bx) {}_2F_1\left(\frac{1}{4}(1 - 2m), \frac{1-m}{2}; \frac{1}{4}(5 - 2m); \sec^2(a + bx)\right) \sqrt{d \sec(a + bx)} (c \sin(a + bx))^m \sin(2(a + bx)) (-\tan^2(a + bx))^{\frac{1-m}{2}}}{b(-1 + 2m)}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Sec[a + b*x]]*(c*Sin[a + b*x])^m,x]

[Out] -((Csc[a + b*x]^2*Hypergeometric2F1[(1 - 2*m)/4, (1 - m)/2, (5 - 2*m)/4, Sec[a + b*x]^2]*Sqrt[d*Sec[a + b*x]]*(c*Sin[a + b*x])^m*Sin[2*(a + b*x)]*(-Tan[a + b*x]^2)^(1 - m)/2))/(b*(-1 + 2*m))

Maple [F]

time = 0.12, size = 0, normalized size = 0.00

$$\int \sqrt{d \sec(bx + a)} (c \sin(bx + a))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*sec(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x)

[Out] int((d*sec(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x, algorithm="maxima")

[Out] integrate(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x, algorithm="fricas")

[Out] integral(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c \sin(a + bx))^m \sqrt{d \sec(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))**(1/2)*(c*sin(b*x+a))**m,x)

[Out] Integral((c*sin(a + b*x))**m*sqrt(d*sec(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*sec(b*x+a))^(1/2)*(c*sin(b*x+a))^m,x, algorithm="giac")

[Out] integrate(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (c \sin(a + bx))^m \sqrt{\frac{d}{\cos(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m*(d/cos(a + b*x))^(1/2),x)

[Out] int((c*sin(a + b*x))^m*(d/cos(a + b*x))^(1/2), x)

$$3.486 \quad \int \frac{(c \sin(a+bx))^m}{\sqrt{d \sec(a+bx)}} dx$$

Optimal. Leaf size=77

$$\frac{\sqrt[4]{\cos^2(a+bx)} {}_2F_1\left(\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) \sqrt{d \sec(a+bx)} (c \sin(a+bx))^{1+m}}{bcd(1+m)}$$

[Out] (cos(b*x+a)^2)^(1/4)*hypergeom([1/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)*(d*sec(b*x+a))^(1/2)/b/c/d/(1+m)

Rubi [A]

time = 0.07, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2666, 2657}

$$\frac{\sqrt[4]{\cos^2(a+bx)} \sqrt{d \sec(a+bx)} (c \sin(a+bx))^{m+1} {}_2F_1\left(\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a+bx)\right)}{bcd(m+1)}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^m/Sqrt[d*Sec[a + b*x]], x]

[Out] ((Cos[a + b*x]^2)^(1/4)*Hypergeometric2F1[1/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*Sqrt[d*Sec[a + b*x]]*(c*Sin[a + b*x])^(1 + m))/(b*c*d*(1 + m))

Rule 2657

Int[(cos[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2666

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(1/b^2)*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

Rubi steps

$$\int \frac{(c \sin(a + bx))^m}{\sqrt{d \sec(a + bx)}} dx = \frac{\left(\sqrt{d \cos(a + bx)} \sqrt{d \sec(a + bx)}\right) \int \sqrt{d \cos(a + bx)} (c \sin(a + bx))^m dx}{d^2}$$

$$= \frac{\sqrt[4]{\cos^2(a + bx)} {}_2F_1\left(\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a + bx)\right) \sqrt{d \sec(a + bx)} (c \sin(a + bx))^{1+m}}{bcd(1+m)}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 18.67, size = 289, normalized size = 3.75

$$\frac{\text{Sc}(3+m)F_1\left(\frac{1+m}{2}; -\frac{1}{2}, \frac{3}{2}+m; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(a+bx)\right), -\tan^2\left(\frac{1}{2}(a+bx)\right)\right) \cos^4\left(\frac{1}{2}(a+bx)\right) \sin^2\left(\frac{1}{2}(a+bx)\right) (c \sin(a+bx))^{1+m}}{b(1+m) \left((3+2m)F_1\left(\frac{1+m}{2}; -\frac{1}{2}, \frac{3}{2}+m; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(a+bx)\right), -\tan^2\left(\frac{1}{2}(a+bx)\right)\right) + F_1\left(\frac{1+m}{2}; \frac{1}{2}, \frac{3}{2}+m; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(a+bx)\right), -\tan^2\left(\frac{1}{2}(a+bx)\right)\right) (-1 + \cos(a+bx)) + (3+m)F_1\left(\frac{1+m}{2}; -\frac{1}{2}, \frac{3}{2}+m; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(a+bx)\right), -\tan^2\left(\frac{1}{2}(a+bx)\right)\right) (1 + \cos(a+bx)) \right) \sqrt{d \sec(a+bx)}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(c*Sin[a + b*x])^m/Sqrt[d*Sec[a + b*x]],x]

[Out] (8*c*(3 + m)*AppellF1[(1 + m)/2, -1/2, 3/2 + m, (3 + m)/2, Tan[(a + b*x)/2]^2, -Tan[(a + b*x)/2]^2]*Cos[(a + b*x)/2]^4*Sin[(a + b*x)/2]^2*(c*Sin[a + b*x])^(-1 + m))/(b*(1 + m)*(((3 + 2*m)*AppellF1[(3 + m)/2, -1/2, 5/2 + m, (5 + m)/2, Tan[(a + b*x)/2]^2, -Tan[(a + b*x)/2]^2] + AppellF1[(3 + m)/2, 1/2, 3/2 + m, (5 + m)/2, Tan[(a + b*x)/2]^2, -Tan[(a + b*x)/2]^2])*(-1 + Cos[a + b*x]) + (3 + m)*AppellF1[(1 + m)/2, -1/2, 3/2 + m, (3 + m)/2, Tan[(a + b*x)/2]^2, -Tan[(a + b*x)/2]^2]*(1 + Cos[a + b*x]))*Sqrt[d*Sec[a + b*x]])

Maple [F]

time = 0.12, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(bx + a))^m}{\sqrt{d \sec(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^m/(d*sec(b*x+a))^(1/2),x)

[Out] int((c*sin(b*x+a))^m/(d*sec(b*x+a))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m/sqrt(d*sec(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m/(d*sec(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(a + bx))^m}{\sqrt{d \sec(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(1/2),x)

[Out] Integral((c*sin(a + b*x))^m/sqrt(d*sec(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m/sqrt(d*sec(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + bx))^m}{\sqrt{\frac{d}{\cos(a + bx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/(d/cos(a + b*x))^(1/2),x)

[Out] int((c*sin(a + b*x))^m/(d/cos(a + b*x))^(1/2), x)

$$3.487 \quad \int \frac{(c \sin(a+bx))^m}{(d \sec(a+bx))^{3/2}} dx$$

Optimal. Leaf size=77

$$\frac{{}_2F_1\left(-\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bcd(1+m) \sqrt[4]{\cos^2(a+bx)} \sqrt{d \sec(a+bx)}}$$

[Out] hypergeom([-1/4, 1/2+1/2*m], [3/2+1/2*m], sin(b*x+a)^2)*(c*sin(b*x+a))^(1+m)/b/c/d/(1+m)/(cos(b*x+a)^2)^(1/4)/(d*sec(b*x+a))^(1/2)

Rubi [A]

time = 0.08, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2666, 2657}

$$\frac{(c \sin(a+bx))^{m+1} {}_2F_1\left(-\frac{1}{4}, \frac{m+1}{2}; \frac{m+3}{2}; \sin^2(a+bx)\right)}{bcd(m+1) \sqrt[4]{\cos^2(a+bx)} \sqrt{d \sec(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(c*Sin[a + b*x])^m/(d*Sec[a + b*x])^(3/2),x]

[Out] (Hypergeometric2F1[-1/4, (1 + m)/2, (3 + m)/2, Sin[a + b*x]^2]*(c*Sin[a + b*x])^(1 + m))/(b*c*d*(1 + m)*(Cos[a + b*x]^2)^(1/4)*Sqrt[d*Sec[a + b*x]])

Rule 2657

Int[(cos[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Simp[b^(2*IntPart[(n - 1)/2] + 1)*(b*Cos[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Sin[e + f*x])^(m + 1)/(a*f*(m + 1)*(Cos[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Sin[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rule 2666

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] := Dist[(1/b^2)*(b*Cos[e + f*x])^(n + 1)*(b*Sec[e + f*x])^(n + 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n] && LtQ[n, 1]

Rubi steps

$$\begin{aligned} \int \frac{(c \sin(a+bx))^m}{(d \sec(a+bx))^{3/2}} dx &= \frac{\int (d \cos(a+bx))^{3/2} (c \sin(a+bx))^m dx}{d^2 \sqrt{d \cos(a+bx)} \sqrt{d \sec(a+bx)}} \\ &= \frac{{}_2F_1\left(-\frac{1}{4}, \frac{1+m}{2}; \frac{3+m}{2}; \sin^2(a+bx)\right) (c \sin(a+bx))^{1+m}}{bcd(1+m) \sqrt[4]{\cos^2(a+bx)} \sqrt{d \sec(a+bx)}} \end{aligned}$$

Mathematica [A]

time = 17.30, size = 116, normalized size = 1.51

$$\frac{2c \cos(2(a + bx)) {}_2F_1\left(\frac{1}{4}(-3 - 2m), \frac{1-m}{2}; \frac{1}{4}(1 - 2m); \sec^2(a + bx)\right) (c \sin(a + bx))^{-1+m} (-\tan^2(a + bx))^{\frac{1-m}{2}}}{bd(3 + 2m) \sqrt{d \sec(a + bx)} (-2 + \sec^2(a + bx))}$$

Antiderivative was successfully verified.

[In] Integrate[(c*Sin[a + b*x])^m/(d*Sec[a + b*x])^(3/2), x]

[Out] (2*c*Cos[2*(a + b*x)]*Hypergeometric2F1[(-3 - 2*m)/4, (1 - m)/2, (1 - 2*m)/4, Sec[a + b*x]^2]*(c*Sin[a + b*x])^(-1 + m)*(-Tan[a + b*x]^2)^((1 - m)/2)) / (b*d*(3 + 2*m)*Sqrt[d*Sec[a + b*x]]*(-2 + Sec[a + b*x]^2))

Maple [F]

time = 0.11, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(bx + a))^m}{(d \sec(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(b*x+a))^m/(d*sec(b*x+a))^(3/2), x)

[Out] int((c*sin(b*x+a))^m/(d*sec(b*x+a))^(3/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(3/2), x, algorithm="maxima")

[Out] integrate((c*sin(b*x + a))^m/(d*sec(b*x + a))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(3/2), x, algorithm="fricas")

[Out] integral(sqrt(d*sec(b*x + a))*(c*sin(b*x + a))^m/(d^2*sec(b*x + a)^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(c \sin(a + bx))^m}{(d \sec(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))**m/(d*sec(b*x+a))**(3/2),x)

[Out] Integral((c*sin(a + b*x))**m/(d*sec(a + b*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((c*sin(b*x+a))^m/(d*sec(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((c*sin(b*x + a))^m/(d*sec(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(c \sin(a + b x))^m}{\left(\frac{d}{\cos(a + b x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((c*sin(a + b*x))^m/(d/cos(a + b*x))^(3/2),x)

[Out] int((c*sin(a + b*x))^m/(d/cos(a + b*x))^(3/2), x)

3.488 $\int \sec^n(e + fx) \sin^m(e + fx) dx$

Optimal. Leaf size=86

$$-\frac{{}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) \sec^{-1+n}(e + fx) \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1-n)}$$

[Out] -hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], cos(f*x+e)^2)*sec(f*x+e)^(-1+n)*sin(f*x+e)^(-1+m)*(sin(f*x+e)^2)^(1/2-1/2*m)/f/(1-n)

Rubi [A]

time = 0.06, antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2667, 2656}

$$\frac{\sin^{m-1}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}} \sec^{n-1}(e + fx) {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^n*Sin[e + f*x]^m,x]

[Out] -((Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*Sec[e + f*x]^(-1 + n)*Sin[e + f*x]^(-1 + m)*(Sin[e + f*x]^2)^((1 - m)/2))/(f*(1 - n)))

Rule 2656

Int[(cos[(e_) + (f_)*(x_)])*(a_)^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*(a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2])*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^n(e + fx) \sin^m(e + fx) dx &= (\cos^n(e + fx) \sec^n(e + fx)) \int \cos^{-n}(e + fx) \sin^m(e + fx) dx \\ &= -\frac{{}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) \sec^{-1+n}(e + fx) \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1-n)} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 1.01, size = 285, normalized size = 3.31

$$\frac{4(3+m)F_1\left(\frac{1+m}{2}, n, 1+m-n; \frac{3+m}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \cos^2\left(\frac{1}{2}(e+fx)\right) \sec^2(e+fx) \sin\left(\frac{1}{2}(e+fx)\right) \sin^m(e+fx)}{f(1+m)\left((3+m)F_1\left(\frac{1+m}{2}, n, 1+m-n; \frac{3+m}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) (1+\cos(e+fx)) - 4\left((1+m-n)F_1\left(\frac{1+m}{2}, n, 2+m-n; \frac{3+m}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - nF_1\left(\frac{1+m}{2}, 1+n, 1+m-n; \frac{3+m}{2}, \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right)\right) \sin^2\left(\frac{1}{2}(e+fx)\right)\right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Sec[e + f*x]^n*Sin[e + f*x]^m,x]

[Out] (4*(3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*Sec[e + f*x]^n*Sin[(e + f*x)/2]*Sin[e + f*x]^m)/(f*(1 + m)*((3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(1 + Cos[e + f*x]) - 4*((1 + m - n)*AppellF1[(3 + m)/2, n, 2 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - n*AppellF1[(3 + m)/2, 1 + n, 1 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sin[(e + f*x)/2]^2))

Maple [F]

time = 0.10, size = 0, normalized size = 0.00

$$\int (\sec^n (fx + e)) (\sin^m (fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^n*sin(f*x+e)^m,x)

[Out] int(sec(f*x+e)^n*sin(f*x+e)^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^n*sin(f*x+e)^m,x, algorithm="maxima")

[Out] integrate(sec(f*x + e)^n*sin(f*x + e)^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^n*sin(f*x+e)^m,x, algorithm="fricas")

[Out] `integral(sec(f*x + e)^n*sin(f*x + e)^m, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sin^m(e + fx) \sec^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(f*x+e)**n*sin(f*x+e)**m,x)`

[Out] `Integral(sin(e + f*x)**m*sec(e + f*x)**n, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(f*x+e)^n*sin(f*x+e)^m,x, algorithm="giac")`

[Out] `integrate(sec(f*x + e)^n*sin(f*x + e)^m, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^m \left(\frac{1}{\cos(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^m*(1/cos(e + f*x))^n,x)`

[Out] `int(sin(e + f*x)^m*(1/cos(e + f*x))^n, x)`

3.489 $\int \sec^n(e + fx)(a \sin(e + fx))^m dx$

Optimal. Leaf size=89

$$-\frac{a {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) \sec^{-1+n}(e + fx)(a \sin(e + fx))^{-1+m} \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1-n)}$$

[Out] -a*hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], cos(f*x+e)^2)*sec(f*x+e)^(-1+n)*(a*sin(f*x+e))^(1+m)*(sin(f*x+e)^2)^(1/2-1/2*m)/f/(1-n)

Rubi [A]

time = 0.06, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2667, 2656}

$$-\frac{a \sin^2(e + fx)^{\frac{1-m}{2}} \sec^{n-1}(e + fx)(a \sin(e + fx))^{m-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Sec[e + f*x]^n*(a*Sin[e + f*x])^m,x]

[Out] -((a*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*Sec[e + f*x]^(-1 + n)*(a*Sin[e + f*x])^(-1 + m)*(Sin[e + f*x]^2)^((1 - m)/2))/(f*(1 - n)))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_.)])^n, x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^n*((a_.)*sin[(e_.) + (f_.)*(x_.)])^m, x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \sec^n(e + fx)(a \sin(e + fx))^m dx &= (\cos^n(e + fx) \sec^n(e + fx)) \int \cos^{-n}(e + fx)(a \sin(e + fx))^m dx \\ &= -\frac{a {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) \sec^{-1+n}(e + fx)(a \sin(e + fx))^{-1}}{f(1-n)} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.20, size = 287, normalized size = 3.22

$$\frac{4(3+m)F_1\left(\frac{4+m}{2}; n, 1+m-n, \frac{4+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \cos^2\left(\frac{1}{2}(e+fx)\right) \sec^2(e+fx) \sin\left(\frac{1}{2}(e+fx)\right) (a \sin(e+fx))^m}{f(1+m) \left((3+m)F_1\left(\frac{4+m}{2}; n, 1+m-n, \frac{4+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) (1+\cos(e+fx)) - 4(1+m-n)F_1\left(\frac{4+m}{2}; n, 2+m-n, \frac{4+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - nF_1\left(\frac{4+m}{2}; 1+n, 1+m-n, \frac{4+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \sin^2\left(\frac{1}{2}(e+fx)\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Sec[e + f*x]^n*(a*Sin[e + f*x])^m,x]

[Out] (4*(3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*Sec[e + f*x]^n*Sin[(e + f*x)/2]*(a*Sin[e + f*x])^m)/(f*(1 + m)*((3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(1 + Cos[e + f*x]) - 4*((1 + m - n)*AppellF1[(3 + m)/2, n, 2 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - n*AppellF1[(3 + m)/2, 1 + n, 1 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2))*Sin[(e + f*x)/2]^2)

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (\sec^n(fx + e)) (a \sin(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sec(f*x+e)^n*(a*sin(f*x+e))^m,x)

[Out] int(sec(f*x+e)^n*(a*sin(f*x+e))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^n*(a*sin(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((a*sin(f*x + e))^m*sec(f*x + e)^n, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sec(f*x+e)^n*(a*sin(f*x+e))^m,x, algorithm="fricas")

[Out] `integral((a*sin(f*x + e))^m*sec(f*x + e)^n, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sin(e + fx))^m \sec^n(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(f*x+e)**n*(a*sin(f*x+e))**m,x)`

[Out] `Integral((a*sin(e + f*x))**m*sec(e + f*x)**n, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(sec(f*x+e)^n*(a*sin(f*x+e))^m,x, algorithm="giac")`

[Out] `integrate((a*sin(f*x + e))^m*sec(f*x + e)^n, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \sin(e + fx))^m \left(\frac{1}{\cos(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sin(e + f*x))^m*(1/cos(e + f*x))^n,x)`

[Out] `int((a*sin(e + f*x))^m*(1/cos(e + f*x))^n, x)`

3.490 $\int (b \sec(e + fx))^n \sin^m(e + fx) dx$

Optimal. Leaf size=89

$$\frac{b {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin^{-1+m}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1-n)}$$

[Out] -b*hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))⁻⁽¹⁺ⁿ⁾*sin(f*x+e)^(-1+m)*(sin(f*x+e)^2)^(1/2-1/2*m)/f/(1-n)

Rubi [A]

time = 0.06, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2667, 2656}

$$\frac{b \sin^{m-1}(e + fx) \sin^2(e + fx)^{\frac{1-m}{2}} (b \sec(e + fx))^{n-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])ⁿ*Sin[e + f*x]^m, x]

[Out] -((b*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^(-1 + n)*Sin[e + f*x]^(-1 + m)*(Sin[e + f*x]^2)^{((1 - m)/2)})/(f*(1 - n)))

Rule 2656

Int[(cos[(e_) + (f_)*(x_)])*(a_)^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^{(2*IntPart[(n - 1)/2] + 1)}*(b*Sin[e + f*x])^{(2*FracPart[(n - 1)/2])}*(a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^{FracPart[(n - 1)/2]}))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[b²*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])ⁿ, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin^m(e + fx) dx &= (b^2 (b \cos(e + fx))^{-1+n} (b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} \sin^m(e + fx) dx \\ &= -\frac{b {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin^{-1+m}(e + fx)}{f(1-n)} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.19, size = 287, normalized size = 3.22

$$\frac{4(3+m)F_1\left(\frac{1+m}{2}, 1+m-n; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \cos^2\left(\frac{1}{2}(e+fx)\right) (b \sec(e+fx))^n \sin\left(\frac{1}{2}(e+fx)\right) \sin^m(e+fx)}{f(1+m) \left((3+m)F_1\left(\frac{1+m}{2}, 1+m-n; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) (1+\cos(e+fx)) - 4 \left((1+m-n)F_1\left(\frac{1+m}{2}, n, 2+m-n; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - nF_1\left(\frac{3+m}{2}, 1+n, 1+m-n; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \right) \sin^2\left(\frac{1}{2}(e+fx)\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x]^m,x]

[Out] (4*(3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*(b*Sec[e + f*x])^n*Sin[(e + f*x)/2]*Sin[e + f*x]^m)/(f*(1 + m)*((3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(1 + Cos[e + f*x]) - 4*((1 + m - n)*AppellF1[(3 + m)/2, n, 2 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - n*AppellF1[(3 + m)/2, 1 + n, 1 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sin[(e + f*x)/2]^2)

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n (\sin^m(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^n*sin(f*x+e)^m,x)

[Out] int((b*sec(f*x+e))^n*sin(f*x+e)^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^m,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e)^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^m,x, algorithm="fricas")

[Out] `integral((b*sec(f*x + e))^n*sin(f*x + e)^m, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \sin^m(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*sin(f*x+e)^m,x)`

[Out] `Integral((b*sec(e + f*x))^n*sin(e + f*x)^m, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*sin(f*x+e)^m,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^n*sin(f*x + e)^m, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^m \left(\frac{b}{\cos(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^m*(b/cos(e + f*x))^n,x)`

[Out] `int(sin(e + f*x)^m*(b/cos(e + f*x))^n, x)`

3.491 $\int (b \sec(e + fx))^n (a \sin(e + fx))^m dx$

Optimal. Leaf size=92

$$\frac{ab {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} (a \sin(e + fx))^{-1+m} \sin^2(e + fx)^{\frac{1-m}{2}}}{f(1-n)}$$

[Out] -a*b*hypergeom([1/2-1/2*n, 1/2-1/2*m], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^{(-1+n)*(a*sin(f*x+e))^{(-1+m)*(sin(f*x+e)^2)^{(1/2-1/2*m)/f/(1-n)}}

Rubi [A]

time = 0.07, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2667, 2656}

$$\frac{ab \sin^2(e + fx)^{\frac{1-m}{2}} (a \sin(e + fx))^{m-1} (b \sec(e + fx))^{n-1} {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m,x]

[Out] -((a*b*Hypergeometric2F1[(1 - m)/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^{(-1 + n)*(a*Sin[e + f*x])^{(-1 + m)*(Sin[e + f*x]^2)^{(1 - m)/2}})/(f*(1 - n)))

Rule 2656

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] := Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_)*sec[(e_) + (f_)*(x_)])^(n_)*((a_)*sin[(e_) + (f_)*(x_)])^(m_), x_Symbol] := Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n (a \sin(e + fx))^m dx &= (b^2 (b \cos(e + fx))^{-1+n} (b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} (a \sin(e + fx))^m dx \\ &= -\frac{ab {}_2F_1\left(\frac{1-m}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} (a \sin(e + fx))^m}{f(1-n)} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.19, size = 289, normalized size = 3.14

$$\frac{4(3+m)F_1\left(\frac{1+m}{2}; n, 1+m-n; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\cos^2\left(\frac{1}{2}(e+fx)\right)\sec(e+fx)\right)^n \sin\left(\frac{1}{2}(e+fx)\right) (a \sin(e+fx))^m}{f(1+m) \left((3+m)F_1\left(\frac{1+m}{2}; n, 1+m-n; \frac{3+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) (1+\cos(e+fx)) - 4(1+m-n)F_1\left(\frac{1+m}{2}; n, 2+m-n; \frac{5+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) - nF_1\left(\frac{1+m}{2}; 1+n, 1+m-n; \frac{5+m}{2}; \tan^2\left(\frac{1}{2}(e+fx)\right), -\tan^2\left(\frac{1}{2}(e+fx)\right)\right) \sin^2\left(\frac{1}{2}(e+fx)\right) \right)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Sec[e + f*x])^n*(a*Sin[e + f*x])^m,x]

[Out] (4*(3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Cos[(e + f*x)/2]^3*(b*Sec[e + f*x])^n*Sin[(e + f*x)/2]*(a*Sin[e + f*x])^m)/(f*(1 + m)*((3 + m)*AppellF1[(1 + m)/2, n, 1 + m - n, (3 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(1 + Cos[e + f*x]) - 4*((1 + m - n)*AppellF1[(3 + m)/2, n, 2 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] - n*AppellF1[(3 + m)/2, 1 + n, 1 + m - n, (5 + m)/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2))*Sin[(e + f*x)/2]^2)

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n (a \sin(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^n*(a*sin(f*x+e))^m,x)

[Out] int((b*sec(f*x+e))^n*(a*sin(f*x+e))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*(a*sin(f*x + e))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="fricas")

[Out] `integral((b*sec(f*x + e))^n*(a*sin(f*x + e))^m, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sin(e + fx))^m (b \sec(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*(a*sin(f*x+e))^m,x)`

[Out] `Integral((a*sin(e + f*x))^m*(b*sec(e + f*x))^n, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^n*(a*sin(f*x + e))^m, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \sin(e + fx))^m \left(\frac{b}{\cos(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*sin(e + f*x))^m*(b/cos(e + f*x))^n,x)`

[Out] `int((a*sin(e + f*x))^m*(b/cos(e + f*x))^n, x)`

3.492 $\int (b \sec(e + fx))^n \sin^5(e + fx) dx$

Optimal. Leaf size=80

$$-\frac{b^5(b \sec(e + fx))^{-5+n}}{f(5-n)} + \frac{2b^3(b \sec(e + fx))^{-3+n}}{f(3-n)} - \frac{b(b \sec(e + fx))^{-1+n}}{f(1-n)}$$

[Out] $-b^5*(b*\sec(f*x+e))^{(-5+n)}/f/(5-n)+2*b^3*(b*\sec(f*x+e))^{(-3+n)}/f/(3-n)-b*(b*\sec(f*x+e))^{(-1+n)}/f/(1-n)$

Rubi [A]

time = 0.05, antiderivative size = 80, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 276}

$$-\frac{b^5(b \sec(e + fx))^{n-5}}{f(5-n)} + \frac{2b^3(b \sec(e + fx))^{n-3}}{f(3-n)} - \frac{b(b \sec(e + fx))^{n-1}}{f(1-n)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^n*\text{Sin}[e + f*x]^5, x]$

[Out] $-((b^5*(b*\text{Sec}[e + f*x])^{(-5 + n)})/(f*(5 - n))) + (2*b^3*(b*\text{Sec}[e + f*x])^{(-3 + n)})/(f*(3 - n)) - (b*(b*\text{Sec}[e + f*x])^{(-1 + n)})/(f*(1 - n))$

Rule 276

$\text{Int}[(c_*)(x_*)^{(m_*)}*((a_*) + (b_*)(x_*)^{(n_*)})^{(p_*)}, x_Symbol] := \text{Int}[\text{Exp andIntegrand}[(c*x)^m*(a + b*x^n)^p, x], x] /; \text{FreeQ}\{a, b, c, m, n, x\} \&\& \text{IGtQ}[p, 0]$

Rule 2702

$\text{Int}[\text{csc}[(e_*) + (f_*)(x_*)]^{(n_*)}*((a_*)*\sec[(e_*) + (f_*)(x_*)])^{(m_*)}, x_Symbol] := \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1 + x^2/a^2)^{((n+1)/2)}, x], x, a*\text{Sec}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m, x\} \&\& \text{IntegerQ}[(n+1)/2] \&\& !(\text{IntegerQ}[(m+1)/2] \&\& \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin^5(e + fx) dx &= \frac{b^5 \text{Subst}\left(\int x^{-6+n} \left(-1 + \frac{x^2}{b^2}\right)^2 dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{b^5 \text{Subst}\left(\int \left(x^{-6+n} - \frac{2x^{-4+n}}{b^2} + \frac{x^{-2+n}}{b^4}\right) dx, x, b \sec(e + fx)\right)}{f} \\ &= -\frac{b^5(b \sec(e + fx))^{-5+n}}{f(5-n)} + \frac{2b^3(b \sec(e + fx))^{-3+n}}{f(3-n)} - \frac{b(b \sec(e + fx))^{-1+n}}{f(1-n)} \end{aligned}$$

Mathematica [A]

time = 0.27, size = 80, normalized size = 1.00

$$\frac{b(89 - 28n + 3n^2 - 4(7 - 8n + n^2) \cos(2(e + fx)) + (3 - 4n + n^2) \cos(4(e + fx))) (b \sec(e + fx))^{-1+n}}{8f(-5 + n)(-3 + n)(-1 + n)}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x]^5,x]`

```
[Out] (b*(89 - 28*n + 3*n^2 - 4*(7 - 8*n + n^2)*Cos[2*(e + f*x)] + (3 - 4*n + n^2)*Cos[4*(e + f*x)])*(b*Sec[e + f*x])^(-1 + n)/(8*f*(-5 + n)*(-3 + n)*(-1 + n))
```

Maple [F]

time = 0.23, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n (\sin^5(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^n*sin(f*x+e)^5,x)``[Out] int((b*sec(f*x+e))^n*sin(f*x+e)^5,x)`**Maxima [A]**

time = 0.32, size = 91, normalized size = 1.14

$$\frac{\frac{b^n \cos(fx+e)^{-n} \cos(fx+e)^5}{n-5} - \frac{2b^n \cos(fx+e)^{-n} \cos(fx+e)^3}{n-3} + \frac{b^n \cos(fx+e)^{-n} \cos(fx+e)}{n-1}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^5,x, algorithm="maxima")`

```
[Out] (b^n*cos(f*x + e)^(-n)*cos(f*x + e)^5/(n - 5) - 2*b^n*cos(f*x + e)^(-n)*cos(f*x + e)^3/(n - 3) + b^n*cos(f*x + e)^(-n)*cos(f*x + e)/(n - 1))/f
```

Fricas [A]

time = 0.38, size = 89, normalized size = 1.11

$$\frac{((n^2 - 4n + 3) \cos(fx + e)^5 - 2(n^2 - 6n + 5) \cos(fx + e)^3 + (n^2 - 8n + 15) \cos(fx + e)) \left(\frac{b}{\cos(fx+e)}\right)^n}{fn^3 - 9fn^2 + 23fn - 15f}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^5,x, algorithm="fricas")`

[Out] $((n^2 - 4n + 3)\cos(fx + e)^5 - 2(n^2 - 6n + 5)\cos(fx + e)^3 + (n^2 - 8n + 15)\cos(fx + e))\cdot(b/\cos(fx + e))^n/(fn^3 - 9fn^2 + 23fn - 15f)$

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**n*sin(f*x+e)**5,x)`

[Out] Timed out

Giac [F]
time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*sin(f*x+e)^5,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^n*sin(f*x + e)^5, x)`

Mupad [B]
time = 1.60, size = 134, normalized size = 1.68

$$\left(\frac{b}{\cos(e+fx)}\right)^n \frac{(150 \cos(e+fx) - 25 \cos(3e+3fx) + 3 \cos(5e+5fx) - 24n \cos(e+fx) + 28n \cos(3e+3fx) - 4n \cos(5e+5fx) + 2n^2 \cos(e+fx) - 3n^2 \cos(3e+3fx) + n^2 \cos(5e+5fx))}{16f(n^3 - 9n^2 + 23n - 15)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^5*(b/cos(e + f*x))^n,x)`

[Out] $((b/\cos(e + fx))^n \cdot (150 \cos(e + fx) - 25 \cos(3e + 3fx) + 3 \cos(5e + 5fx) - 24n \cos(e + fx) + 28n \cos(3e + 3fx) - 4n \cos(5e + 5fx) + 2n^2 \cos(e + fx) - 3n^2 \cos(3e + 3fx) + n^2 \cos(5e + 5fx))) / (16f \cdot (23n - 9n^2 + n^3 - 15))$

3.493 $\int (b \sec(e + fx))^n \sin^3(e + fx) dx$

Optimal. Leaf size=52

$$\frac{b^3(b \sec(e + fx))^{-3+n}}{f(3-n)} - \frac{b(b \sec(e + fx))^{-1+n}}{f(1-n)}$$

[Out] $b^3*(b*\sec(f*x+e))^{(-3+n)}/f/(3-n)-b*(b*\sec(f*x+e))^{(-1+n)}/f/(1-n)$

Rubi [A]

time = 0.04, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 14}

$$\frac{b^3(b \sec(e + fx))^{n-3}}{f(3-n)} - \frac{b(b \sec(e + fx))^{n-1}}{f(1-n)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(b*\text{Sec}[e + f*x])^n*\text{Sin}[e + f*x]^3, x]$

[Out] $(b^3*(b*\text{Sec}[e + f*x])^{(-3 + n)})/(f*(3 - n)) - (b*(b*\text{Sec}[e + f*x])^{(-1 + n)})/(f*(1 - n))$

Rule 14

$\text{Int}[(u_)*((c_)*(x_))^{(m_)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(c*x)^m*u, x], x] /; \text{FreeQ}\{c, m\}, x \ \&\& \ \text{SumQ}[u] \ \&\& \ !\text{LinearQ}[u, x] \ \&\& \ !\text{MatchQ}[u, (a_ + (b_)*(v_)) /; \text{FreeQ}\{a, b\}, x] \ \&\& \ \text{InverseFunctionQ}[v]$

Rule 2702

$\text{Int}[\text{csc}[(e_.) + (f_.)*(x_)]^{(n_.)}*((a_.)*\text{sec}[(e_.) + (f_.)*(x_)]^{(m_)}], x_Symbol] \rightarrow \text{Dist}[1/(f*a^n), \text{Subst}[\text{Int}[x^{(m+n-1)}/(-1+x^2/a^2)^{((n+1)/2)}, x], x, a*\text{Sec}[e + f*x]], x] /; \text{FreeQ}\{a, e, f, m\}, x \ \&\& \ \text{IntegerQ}[(n+1)/2] \ \&\& \ !(\text{IntegerQ}[(m+1)/2] \ \&\& \ \text{LtQ}[0, m, n])$

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin^3(e + fx) dx &= \frac{b^3 \text{Subst}\left(\int x^{-4+n} \left(-1 + \frac{x^2}{b^2}\right) dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{b^3 \text{Subst}\left(\int \left(-x^{-4+n} + \frac{x^{-2+n}}{b^2}\right) dx, x, b \sec(e + fx)\right)}{f} \\ &= \frac{b^3(b \sec(e + fx))^{-3+n}}{f(3-n)} - \frac{b(b \sec(e + fx))^{-1+n}}{f(1-n)} \end{aligned}$$

Mathematica [A]

time = 0.11, size = 47, normalized size = 0.90

$$\frac{b(5 - n + (-1 + n) \cos(2(e + fx)))(b \sec(e + fx))^{-1+n}}{2f(-3 + n)(-1 + n)}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x]^3,x]``[Out] -1/2*(b*(5 - n + (-1 + n)*Cos[2*(e + f*x)])*(b*Sec[e + f*x])^(-1 + n))/(f*(-3 + n)*(-1 + n))`**Maple [C]** Result contains higher order function than in optimal. Order 9 vs. order 3.

time = 1.22, size = 1732, normalized size = 33.31

method	result	size
risch	Expression too large to display	1732

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^n*sin(f*x+e)^3,x,method=_RETURNVERBOSE)`

```
[Out] -1/8/(f*n-3*f)*2^n*(exp(2*I*(f*x+e))+1)^(-n)*b^n*exp(I*(f*x+e))^n*exp(-1/2*I*(Pi*n*csgn(I/(exp(2*I*(f*x+e))+1))*csgn(I*exp(I*(f*x+e))))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))-Pi*n*csgn(I/(exp(2*I*(f*x+e))+1))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2-Pi*n*csgn(I*exp(I*(f*x+e)))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2+Pi*n*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^3-Pi*n*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2+Pi*n*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2+Pi*n*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^3-Pi*n*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2*csgn(I*b)+6*f*x+6*e))-1/8*exp(I*(f*x+e))^n*b^n*(exp(2*I*(f*x+e))+1)^(-n)*2^n/(f*n-3*f)*exp(1/2*I*(-Pi*n*csgn(I/(exp(2*I*(f*x+e))+1))*csgn(I*exp(I*(f*x+e))))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))+Pi*n*csgn(I/(exp(2*I*(f*x+e))+1))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2+Pi*n*csgn(I*exp(I*(f*x+e)))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2-Pi*n*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^3+Pi*n*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2-Pi*n*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2-Pi*n*csgn(I*b)-Pi*n*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^3+Pi*n*csgn(I*b*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2*csgn(I*b)+6*f*x+6*e))+1/8*exp(I*(f*x+e))^n*b^n*(exp(2*I*(f*x+e))+1)^(-n)*2^n/(-3+n)/(-1+n)/f*(n-9)*exp(-1/2*I*(Pi*n*csgn(I/(exp(2*I*(f*x+e))+1))*csgn(I*exp(I*(f*x+e))))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))-Pi*n*csgn(I/(exp(2*I*(f*x+e))+1))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2-Pi*n*csgn(I*exp(I*(f*x+e)))*csgn(I*exp(I*(f*x+e)))/(exp(2*I*(f*x+e))+1))^2+Pi*n*csgn(I*exp(I*(f*x+e)))/(e
```

$$\begin{aligned} & \exp(2I*(f*x+e))+1)^{3-Pi*n}*csgn(I*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))*csgn \\ & (I*b*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))^{2+Pi*n}*csgn(I*\exp(I*(f*x+e)))/(\exp \\ & (2I*(f*x+e))+1))*csgn(I*b*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))*csgn(I*b)+P \\ & i*n*csgn(I*b*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))^{3-Pi*n}*csgn(I*b*\exp(I*(f* \\ & x+e)))/(\exp(2I*(f*x+e))+1))^{2}*csgn(I*b)+2*f*x+2*e))+1/8*\exp(I*(f*x+e))^n*b^ \\ & n*(\exp(2I*(f*x+e))+1)^{-n}*2^n/(-3+n)/(-1+n)/f*(n-9)*\exp(1/2*I*(-Pi*n*csgn \\ & (I/(\exp(2I*(f*x+e))+1))*csgn(I*\exp(I*(f*x+e))))*csgn(I*\exp(I*(f*x+e)))/(\exp \\ & (2I*(f*x+e))+1))+Pi*n*csgn(I/(\exp(2I*(f*x+e))+1))*csgn(I*\exp(I*(f*x+e)))/(e \\ & xp(2I*(f*x+e))+1))^{2+Pi*n}*csgn(I*\exp(I*(f*x+e)))*csgn(I*\exp(I*(f*x+e)))/(e \\ & xp(2I*(f*x+e))+1))^{2-Pi*n}*csgn(I*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))^{3+Pi* \\ & n*csgn(I*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))*csgn(I*b*\exp(I*(f*x+e)))/(\exp \\ & (2I*(f*x+e))+1))^{2-Pi*n}*csgn(I*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))*csgn(I* \\ & b*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1))*csgn(I*b)-Pi*n*csgn(I*b*\exp(I*(f*x+e \\ &)))/(\exp(2I*(f*x+e))+1))^{3+Pi*n}*csgn(I*b*\exp(I*(f*x+e)))/(\exp(2I*(f*x+e))+1 \\ &))^{2}*csgn(I*b)+2*f*x+2*e)) \end{aligned}$$

Maxima [A]

time = 0.28, size = 63, normalized size = 1.21

$$\frac{\frac{b^n \cos(fx+e)^{-n} \cos(fx+e)^3}{n-3} - \frac{b^n \cos(fx+e)^{-n} \cos(fx+e)}{n-1}}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^3,x, algorithm="maxima")

[Out] $-(b^n*\cos(f*x + e)^{-n}*\cos(f*x + e)^3/(n - 3) - b^n*\cos(f*x + e)^{-n}*\cos(f*x + e)/(n - 1))/f$

Fricas [A]

time = 0.35, size = 56, normalized size = 1.08

$$\frac{((n-1)\cos(fx+e)^3 - (n-3)\cos(fx+e))\left(\frac{b}{\cos(fx+e)}\right)^n}{fn^2 - 4fn + 3f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^3,x, algorithm="fricas")

[Out] $-((n-1)*\cos(f*x + e)^3 - (n-3)*\cos(f*x + e))*(b/\cos(f*x + e))^n/(f*n^2 - 4*f*n + 3*f)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)**3,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^3,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e)^3, x)

Mupad [B]

time = 0.94, size = 67, normalized size = 1.29

$$\frac{\left(\frac{b}{\cos(e+fx)}\right)^n (9 \cos(e+fx) - \cos(3e+3fx) - n \cos(e+fx) + n \cos(3e+3fx))}{4f(n^2 - 4n + 3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^3*(b/cos(e + f*x))^n,x)

[Out] -((b/cos(e + f*x))^n*(9*cos(e + f*x) - cos(3*e + 3*f*x) - n*cos(e + f*x) + n*cos(3*e + 3*f*x)))/(4*f*(n^2 - 4*n + 3))

3.494 $\int (b \sec(e + fx))^n \sin(e + fx) dx$

Optimal. Leaf size=25

$$-\frac{b(b \sec(e + fx))^{-1+n}}{f(1-n)}$$

[Out] -b*(b*sec(f*x+e))⁽⁻¹⁺ⁿ⁾/f/(1-n)

Rubi [A]

time = 0.02, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {2702, 30}

$$-\frac{b(b \sec(e + fx))^{n-1}}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])ⁿ*Sin[e + f*x],x]

[Out] -((b*(b*Sec[e + f*x])^(-1 + n))/(f*(1 - n)))

Rule 30

Int[(x_)^(m_), x_Symbol] := Simp[x^(m + 1)/(m + 1), x] /; FreeQ[m, x] && NeQ[m, -1]

Rule 2702

Int[csc[(e_) + (f_)*(x_)]^(n_)*((a_)*sec[(e_) + (f_)*(x_)]^(m_), x_Symbol] := Dist[1/(f*aⁿ), Subst[Int[x^(m + n - 1)/(-1 + x²/a²)^{((n + 1)/2)}, x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin(e + fx) dx &= \frac{b \text{Subst}(\int x^{-2+n} dx, x, b \sec(e + fx))}{f} \\ &= -\frac{b(b \sec(e + fx))^{-1+n}}{f(1-n)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 22, normalized size = 0.88

$$\frac{b(b \sec(e + fx))^{-1+n}}{f(-1 + n)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x],x]

[Out] (b*(b*Sec[e + f*x])^(-1 + n))/(f*(-1 + n))

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 119 vs. $2(25) = 50$.

time = 0.78, size = 120, normalized size = 4.80

method	result	size
norman	$\frac{e^{n \ln\left(\frac{b(1+\tan^2(\frac{fx}{2}+\frac{e}{2}))}{1-\tan^2(\frac{fx}{2}+\frac{e}{2})}\right)} f^{(-1+n)} - \frac{(\tan^2(\frac{fx}{2}+\frac{e}{2}))^n e^{n \ln\left(\frac{b(1+\tan^2(\frac{fx}{2}+\frac{e}{2}))}{1-\tan^2(\frac{fx}{2}+\frac{e}{2})}\right)} f^{(-1+n)}}{1+\tan^2\left(\frac{fx}{2}+\frac{e}{2}\right)}$	120
risch	Expression too large to display	930

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^n*sin(f*x+e),x,method=_RETURNVERBOSE)

[Out] (1/f/(-1+n)*exp(n*ln(b*(1+tan(1/2*f*x+1/2*e)^2)/(1-tan(1/2*f*x+1/2*e)^2)))-1/f/(-1+n)*tan(1/2*f*x+1/2*e)^2*exp(n*ln(b*(1+tan(1/2*f*x+1/2*e)^2)/(1-tan(1/2*f*x+1/2*e)^2))))/(1+tan(1/2*f*x+1/2*e)^2)

Maxima [A]

time = 0.31, size = 30, normalized size = 1.20

$$\frac{b^n \cos(fx + e)^{-n} \cos(fx + e)}{f(n - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e),x, algorithm="maxima")

[Out] b^n*cos(f*x + e)^(-n)*cos(f*x + e)/(f*(n - 1))

Fricas [A]

time = 0.37, size = 30, normalized size = 1.20

$$\frac{\left(\frac{b}{\cos(fx+e)}\right)^n \cos(fx + e)}{fn - f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e),x, algorithm="fricas")

[Out] (b/cos(f*x + e))^n*cos(f*x + e)/(f*n - f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \sin(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**n*sin(f*x+e),x)

[Out] Integral((b*sec(e + f*x))**n*sin(e + f*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e),x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e), x)

Mupad [B]

time = 0.19, size = 27, normalized size = 1.08

$$\frac{\cos(e + fx) \left(\frac{b}{\cos(e+fx)}\right)^n}{f(n-1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)*(b/cos(e + f*x))^n,x)

[Out] (cos(e + f*x)*(b/cos(e + f*x))^n)/(f*(n - 1))

3.495 $\int \csc(e + fx)(b \sec(e + fx))^n dx$

Optimal. Leaf size=49

$$-\frac{{}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \sec^2(e + fx)\right) (b \sec(e + fx))^{1+n}}{bf(1+n)}$$

[Out] -hypergeom([1, 1/2+1/2*n], [3/2+1/2*n], sec(f*x+e)^2)*(b*sec(f*x+e))^(1+n)/b/f/(1+n)

Rubi [A]

time = 0.03, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$,

Rules used = {2702, 371}

$$-\frac{(b \sec(e + fx))^{n+1} {}_2F_1\left(1, \frac{n+1}{2}; \frac{n+3}{2}; \sec^2(e + fx)\right)}{bf(n+1)}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]*(b*Sec[e + f*x])^n,x]

[Out] -((Hypergeometric2F1[1, (1 + n)/2, (3 + n)/2, Sec[e + f*x]^2]*(b*Sec[e + f*x])^(1 + n))/(b*f*(1 + n)))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p *((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILt Q[p, 0] || GtQ[a, 0])

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^((n + 1)/2), x], x, a*Sec[e + f*x]], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \csc(e + fx)(b \sec(e + fx))^n dx &= \frac{\text{Subst}\left(\int \frac{x^n}{-1 + \frac{x^2}{b^2}} dx, x, b \sec(e + fx)\right)}{bf} \\ &= -\frac{{}_2F_1\left(1, \frac{1+n}{2}; \frac{3+n}{2}; \sec^2(e + fx)\right) (b \sec(e + fx))^{1+n}}{bf(1+n)} \end{aligned}$$

Mathematica [A]

time = 0.24, size = 92, normalized size = 1.88

$$\frac{\left({}_2F_1(1, -n; 1 - n; \cos(e + fx)) - 2^n {}_2F_1(-n, -n; 1 - n; \frac{1}{2} \cos(e + fx) \sec^2(\frac{1}{2}(e + fx))) \sec^2(\frac{1}{2}(e + fx))^{-n}\right) (b \sec(e + fx))^n}{2fn}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]*(b*Sec[e + f*x])^n,x]

```
[Out] ((Hypergeometric2F1[1, -n, 1 - n, Cos[e + f*x]] - (2^n*Hypergeometric2F1[-n, -n, 1 - n, (Cos[e + f*x]*Sec[(e + f*x)/2]^2)/2]))/(Sec[(e + f*x)/2]^2)^n*(b*Sec[e + f*x])^n)/(2*f*n)
```

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int \csc(fx + e) (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)*(b*sec(f*x+e))^n,x)

[Out] int(csc(f*x+e)*(b*sec(f*x+e))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*csc(f*x + e), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(b*sec(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*sec(f*x + e))^n*csc(f*x + e), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(b*sec(f*x+e))**n,x)`

[Out] `Integral((b*sec(e + f*x))**n*csc(e + f*x), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)*(b*sec(f*x+e))^n,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^n*csc(f*x + e), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^n}{\sin(e+fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(e + f*x))^n/sin(e + f*x),x)`

[Out] `int((b/cos(e + f*x))^n/sin(e + f*x), x)`

3.496 $\int \csc^3(e + fx)(b \sec(e + fx))^n dx$

Optimal. Leaf size=48

$$\frac{{}_2F_1\left(2, \frac{3+n}{2}; \frac{5+n}{2}; \sec^2(e + fx)\right) (b \sec(e + fx))^{3+n}}{b^3 f(3+n)}$$

[Out] hypergeom([2, 3/2+1/2*n], [5/2+1/2*n], sec(f*x+e)^2)*(b*sec(f*x+e))^(3+n)/b^3/f/(3+n)

Rubi [A]

time = 0.04, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2702, 371}

$$\frac{(b \sec(e + fx))^{n+3} {}_2F_1\left(2, \frac{n+3}{2}; \frac{n+5}{2}; \sec^2(e + fx)\right)}{b^3 f(n+3)}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^3*(b*Sec[e + f*x])^n,x]

[Out] (Hypergeometric2F1[2, (3 + n)/2, (5 + n)/2, Sec[e + f*x]^2]*(b*Sec[e + f*x])^(3 + n))/(b^3*f*(3 + n))

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p*((c*x)^(m + 1)/(c*(m + 1)))*Hypergeometric2F1[-p, (m + 1)/n, (m + 1)/n + 1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 2702

Int[csc[(e_.) + (f_.)*(x_)]^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_)])^(m_), x_Symbol] :> Dist[1/(f*a^n), Subst[Int[x^(m + n - 1)/(-1 + x^2/a^2)^(n + 1)/2], x], x, a*Sec[e + f*x], x] /; FreeQ[{a, e, f, m}, x] && IntegerQ[(n + 1)/2] && !(IntegerQ[(m + 1)/2] && LtQ[0, m, n])

Rubi steps

$$\begin{aligned} \int \csc^3(e + fx)(b \sec(e + fx))^n dx &= \frac{\text{Subst}\left(\int \frac{x^{2+n}}{(-1+\frac{x^2}{b^2})^2} dx, x, b \sec(e + fx)\right)}{b^3 f} \\ &= \frac{{}_2F_1\left(2, \frac{3+n}{2}; \frac{5+n}{2}; \sec^2(e + fx)\right) (b \sec(e + fx))^{3+n}}{b^3 f(3+n)} \end{aligned}$$

Mathematica [B] Leaf count is larger than twice the leaf count of optimal. 201 vs. 2(48) = 96.

time = 3.09, size = 201, normalized size = 4.19

$$\frac{b(b \sec(e + fx))^{-1+n} \left({}_2F_1(1, 1-n; 2-n; \cos(e+fx)) + 2 {}_2F_1(2, 1-n; 2-n; \cos(e+fx)) + 2^2 {}_2F_1(1-n, -n; 2-n; \frac{1}{2} \cos(e+fx) \sec^2(\frac{1}{2}(e+fx))) \sec^2(\frac{1}{2}(e+fx))^{-1+n} + 2^n {}_2F_1(1-n, 1-n; 2-n; \frac{1}{2} \cos(e+fx) \sec^2(\frac{1}{2}(e+fx))) \sec^{-n}(e+fx) (\cos^2(\frac{1}{2}(e+fx)) \sec(e+fx))^{-1+n} \right)}{8f^{(-1+n)}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[Csc[e + f*x]^3*(b*Sec[e + f*x])^n,x]

[Out] (b*(b*Sec[e + f*x])^(-1 + n)*(2*Hypergeometric2F1[1, 1 - n, 2 - n, Cos[e + f*x]] + 2*Hypergeometric2F1[2, 1 - n, 2 - n, Cos[e + f*x]] + 2^n*Hypergeometric2F1[1 - n, -n, 2 - n, (Cos[e + f*x]*Sec[(e + f*x)/2]^2)/2]*(Sec[(e + f*x)/2]^2)^(1 - n) + 2^n*Hypergeometric2F1[1 - n, 1 - n, 2 - n, (Cos[e + f*x]*Sec[(e + f*x)/2]^2)/2]*Sec[e + f*x]^(1 - n)*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^(-1 + n)))/(8*f*(-1 + n))

Maple [F]

time = 0.05, size = 0, normalized size = 0.00

$$\int (\csc^3(fx + e)) (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3*(b*sec(f*x+e))^n,x)

[Out] int(csc(f*x+e)^3*(b*sec(f*x+e))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*csc(f*x + e)^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*sec(f*x + e))^n*csc(f*x + e)^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \csc^3(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3*(b*sec(f*x+e))**n,x)

[Out] Integral((b*sec(e + f*x))**n*csc(e + f*x)**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(b*sec(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^n*csc(f*x + e)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^n}{\sin(e+fx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^n/sin(e + f*x)^3,x)

[Out] int((b/cos(e + f*x))^n/sin(e + f*x)^3, x)

3.497 $\int (b \sec(e + fx))^n \sin^6(e + fx) dx$

Optimal. Leaf size=73

$$-\frac{b {}_2F_1\left(-\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n)\sqrt{\sin^2(e + fx)}}$$

[Out] -b*hypergeom([-5/2, 1/2-1/2*n], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^(1-n)*sin(f*x+e)/f/(1-n)/(sin(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2712, 2656}

$$-\frac{b \sin(e + fx)(b \sec(e + fx))^{n-1} {}_2F_1\left(-\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)\sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^n*Sin[e + f*x]^6,x]

[Out] -((b*Hypergeometric2F1[-5/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^(1 - n)*Sin[e + f*x])/(f*(1 - n)*Sqrt[Sin[e + f*x]^2]))

Rule 2656

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2712

Int[(csc[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(a^2/b^2)*(a*Sec[e + f*x])^(m - 1)*(b*Csc[e + f*x])^(n + 1)*(a*Cos[e + f*x])^(m - 1)*(b*Sin[e + f*x])^(n + 1), Int[1/((a*Cos[e + f*x])^m*(b*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin^6(e + fx) dx &= (b^2 (b \cos(e + fx))^{-1+n} (b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} \sin^6(e + fx) dx \\ &= -\frac{b {}_2F_1\left(-\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 25.27, size = 8327, normalized size = 114.07

Result too large to show

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x]^6,x]

[Out] Result too large to show

Maple [F]

time = 0.26, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n (\sin^6(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^n*sin(f*x+e)^6,x)

[Out] int((b*sec(f*x+e))^n*sin(f*x+e)^6,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^6,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e)^6, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^6,x, algorithm="fricas")

[Out] integral(-(cos(f*x + e))^6 - 3*cos(f*x + e)^4 + 3*cos(f*x + e)^2 - 1)*(b*sec(f*x + e))^n, x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))**n*sin(f*x+e)**6,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep
```

Giac [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^6,x, algorithm="giac")
```

```
[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e)^6, x)
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.01
```

$$\int \sin(e + f x)^6 \left(\frac{b}{\cos(e + f x)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^6*(b/cos(e + f*x))^n,x)
```

```
[Out] int(sin(e + f*x)^6*(b/cos(e + f*x))^n, x)
```

3.498 $\int (b \sec(e + fx))^n \sin^4(e + fx) dx$

Optimal. Leaf size=73

$$\frac{b {}_2F_1\left(-\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n) \sqrt{\sin^2(e + fx)}}$$

[Out] -b*hypergeom([-3/2, 1/2-1/2*n], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^(1-n)*sin(f*x+e)/f/(1-n)/(sin(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2712, 2656}

$$\frac{b \sin(e + fx) (b \sec(e + fx))^{n-1} {}_2F_1\left(-\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n) \sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^n*Sin[e + f*x]^4,x]

[Out] -((b*Hypergeometric2F1[-3/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^(1 - n)*Sin[e + f*x])/((f*(1 - n)*Sqrt[Sin[e + f*x]^2]))

Rule 2656

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^(FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2712

Int[(csc[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(a^2/b^2)*(a*Sec[e + f*x])^(m - 1)*(b*Csc[e + f*x])^(n + 1)*(a*Cos[e + f*x])^(m - 1)*(b*Sin[e + f*x])^(n + 1), Int[1/((a*Cos[e + f*x])^m*(b*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin^4(e + fx) dx &= (b^2 (b \cos(e + fx))^{-1+n} (b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} \sin^4(e + fx) dx \\ &= -\frac{b {}_2F_1\left(-\frac{3}{2}, \frac{1-n}{2}, \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n) \sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 22.01, size = 6192, normalized size = 84.82

Result too large to show

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x]^4,x]

[Out] Result too large to show

Maple [F]

time = 0.24, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n (\sin^4(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^n*sin(f*x+e)^4,x)

[Out] int((b*sec(f*x+e))^n*sin(f*x+e)^4,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^4,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e)^4, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^4,x, algorithm="fricas")

[Out] integral((cos(f*x + e)^4 - 2*cos(f*x + e)^2 + 1)*(b*sec(f*x + e))^n, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \sin^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**n*sin(f*x+e)**4,x)

[Out] Integral((b*sec(e + f*x))**n*sin(e + f*x)**4, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n*sin(f*x+e)^4,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^n*sin(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^4 \left(\frac{b}{\cos(e + f x)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4*(b/cos(e + f*x))^n,x)

[Out] int(sin(e + f*x)^4*(b/cos(e + f*x))^n, x)

3.499 $\int (b \sec(e + fx))^n \sin^2(e + fx) dx$

Optimal. Leaf size=73

$$-\frac{b {}_2F_1\left(-\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n)\sqrt{\sin^2(e + fx)}}$$

[Out] -b*hypergeom([-1/2, 1/2-1/2*n], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^(1-n)*sin(f*x+e)/f/(1-n)/(sin(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2712, 2656}

$$-\frac{b \sin(e + fx) (b \sec(e + fx))^{n-1} {}_2F_1\left(-\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)\sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^n*Sin[e + f*x]^2,x]

[Out] -((b*Hypergeometric2F1[-1/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^(1 - n)*Sin[e + f*x])/(f*(1 - n)*Sqrt[Sin[e + f*x]^2]))

Rule 2656

Int[(cos[(e_) + (f_)*(x_)]*(a_))^(m_)*((b_)*sin[(e_) + (f_)*(x_)])^(n_), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2712

Int[(csc[(e_) + (f_)*(x_)]*(b_))^(n_)*((a_)*sec[(e_) + (f_)*(x_)])^(m_), x_Symbol] :> Dist[(a^2/b^2)*(a*Sec[e + f*x])^(m - 1)*(b*Csc[e + f*x])^(n + 1)*(a*Cos[e + f*x])^(m - 1)*(b*Sin[e + f*x])^(n + 1), Int[1/((a*Cos[e + f*x])^m*(b*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n \sin^2(e + fx) dx &= (b^2 (b \cos(e + fx))^{-1+n} (b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} \sin^2(e + fx) dx \\ &= -\frac{b {}_2F_1\left(-\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 18.01, size = 4143, normalized size = 56.75

Result too large to show

Warning: Unable to verify antiderivative.

[In] Integrate[(b*Sec[e + f*x])^n*Sin[e + f*x]^2,x]

[Out] (24*(Sec[(e + f*x)/2]^2)^(-3 + n)*(b*Sec[e + f*x])^n*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*Sin[e + f*x]^2*Tan[(e + f*x)/2]*((AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2)/(3*AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-2 + n)*AppellF1[3/2, n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 + n, 2 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2) - AppellF1[1/2, n, 3 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]/(3*AppellF1[1/2, n, 3 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-3 + n)*AppellF1[3/2, n, 4 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 + n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)))/(f*(12*(Sec[(e + f*x)/2]^2)^(-2 + n)*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*((AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2)/(3*AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-2 + n)*AppellF1[3/2, n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 + n, 2 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2) - AppellF1[1/2, n, 3 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]/(3*AppellF1[1/2, n, 3 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-3 + n)*AppellF1[3/2, n, 4 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 + n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)) + 24*(-3 + n)*(Sec[(e + f*x)/2]^2)^(-3 + n)*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*Tan[(e + f*x)/2]^2*((AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2)/(3*AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-2 + n)*AppellF1[3/2, n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 + n, 2 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2) - AppellF1[1/2, n, 3 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]/(3*AppellF1[1/2, n, 3 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-3 + n)*AppellF1[3/2, n, 4 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 + n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)) + 24*(Sec[(e + f*x)/2]^2)^(-3 + n)*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*Tan[(e + f*x)/2]^2*((AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/((3*AppellF1[1/2, n, 2 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*((-2 + n)*AppellF1[3/2, n, 3 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + n*AppellF1[3/2, 1 +

$n, 2 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2) * \tan[(e + fx)/2]^2 + (\sec[(e + fx)/2]^2 * (-1/3 * ((2 - n) * \text{AppellF1}[3/2, n, 3 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) + (n * \text{AppellF1}[3/2, 1 + n, 2 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2])) / (3 * \text{AppellF1}[1/2, n, 2 - n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + 2 * ((-2 + n) * \text{AppellF1}[3/2, n, 3 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + n * \text{AppellF1}[3/2, 1 + n, 2 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \tan[(e + fx)/2]^2) - (-1/3 * ((3 - n) * \text{AppellF1}[3/2, n, 4 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) + (n * \text{AppellF1}[3/2, 1 + n, 3 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2])) / (3 * \text{AppellF1}[1/2, n, 3 - n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + 2 * ((-3 + n) * \text{AppellF1}[3/2, n, 4 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + n * \text{AppellF1}[3/2, 1 + n, 3 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \tan[(e + fx)/2]^2) - (\text{AppellF1}[1/2, n, 2 - n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * (2 * ((-2 + n) * \text{AppellF1}[3/2, n, 3 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + n * \text{AppellF1}[3/2, 1 + n, 2 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2] + 3 * (-1/3 * ((2 - n) * \text{AppellF1}[3/2, n, 3 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) + (n * \text{AppellF1}[3/2, 1 + n, 2 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2])) / 3) + 2 * \tan[(e + fx)/2]^2 * ((-2 + n) * ((-3 * (3 - n) * \text{AppellF1}[5/2, n, 4 - n, 7/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 5 + (3 * n * \text{AppellF1}[5/2, 1 + n, 3 - n, 7/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 5) + n * ((-3 * (2 - n) * \text{AppellF1}[5/2, 1 + n, 3 - n, 7/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 5 + (3 * (1 + n) * \text{AppellF1}[5/2, 2 + n, 2 - n, 7/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 5)))) / (3 * \text{AppellF1}[1/2, n, 2 - n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + 2 * ((-2 + n) * \text{AppellF1}[3/2, n, 3 - n, 5/2, \tan[(e + fx)...$

Maple [F]

time = 0.17, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n (\sin^2(fx + e)) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(f*x+e))^n*sin(f*x+e)^2,x)

[Out] int((b*sec(f*x+e))^n*sin(f*x+e)^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*sin(f*x+e)^2,x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^n*sin(f*x + e)^2, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*sin(f*x+e)^2,x, algorithm="fricas")`

[Out] `integral(-(cos(f*x + e)^2 - 1)*(b*sec(f*x + e))^n, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \sin^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))**n*sin(f*x+e)**2,x)`

[Out] `Integral((b*sec(e + f*x))**n*sin(e + f*x)**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(f*x+e))^n*sin(f*x+e)^2,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^n*sin(f*x + e)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^2 \left(\frac{b}{\cos(e + fx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(sin(e + f*x)^2*(b/cos(e + f*x))^n,x)`

[Out] `int(sin(e + f*x)^2*(b/cos(e + f*x))^n, x)`

3.500 $\int (b \sec(e + fx))^n dx$

Optimal. Leaf size=73

$$-\frac{b {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sin(e + fx)}{f(1-n)\sqrt{\sin^2(e + fx)}}$$

[Out] -b*hypergeom([1/2, 1/2-1/2*n], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^(1-n)*sin(f*x+e)/f/(1-n)/(sin(f*x+e)^2)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {3857, 2722}

$$-\frac{b \sin(e + fx)(b \sec(e + fx))^{n-1} {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)\sqrt{\sin^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[e + f*x])^n,x]

[Out] -((b*Hypergeometric2F1[1/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^(1 - n)*Sin[e + f*x])/(f*(1 - n)*Sqrt[Sin[e + f*x]^2]))

Rule 2722

Int[((b_.)*sin[(c_.) + (d_.)*(x_)])^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]

Rule 3857

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(b*Csc[c + d*x])^(n - 1)*((Sin[c + d*x]/b)^(n - 1)*Int[1/(Sin[c + d*x]/b)^n, x]), x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int (b \sec(e + fx))^n dx &= \left(\frac{\cos(e + fx)}{b}\right)^n (b \sec(e + fx))^n \int \left(\frac{\cos(e + fx)}{b}\right)^{-n} dx \\ &= -\frac{\cos(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^n \sin(e + fx)}{f(1-n)\sqrt{\sin^2(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 61, normalized size = 0.84

$$\frac{\cot(e + fx) {}_2F_1\left(\frac{1}{2}, \frac{n}{2}; \frac{2+n}{2}; \sec^2(e + fx)\right) (b \sec(e + fx))^n \sqrt{-\tan^2(e + fx)}}{fn}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[e + f*x])^n,x]``[Out] (Cot[e + f*x]*Hypergeometric2F1[1/2, n/2, (2 + n)/2, Sec[e + f*x]^2]*(b*Sec[e + f*x])^n*Sqrt[-Tan[e + f*x]^2])/(f*n)`**Maple [F]**

time = 0.01, size = 0, normalized size = 0.00

$$\int (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(f*x+e))^n,x)``[Out] int((b*sec(f*x+e))^n,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(f*x+e))^n,x, algorithm="maxima")``[Out] integrate((b*sec(f*x + e))^n, x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(f*x+e))^n,x, algorithm="fricas")``[Out] integral((b*sec(f*x + e))^n, x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))**n,x)

[Out] Integral((b*sec(e + f*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^n, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{b}{\cos(e + f x)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^n,x)

[Out] int((b/cos(e + f*x))^n, x)

3.501 $\int \csc^2(e + fx)(b \sec(e + fx))^n dx$

Optimal. Leaf size=73

$$-\frac{b \csc(e + fx) {}_2F_1\left(\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sqrt{\sin^2(e + fx)}}{f(1-n)}$$

[Out] -b*csc(f*x+e)*hypergeom([3/2, 1/2-1/2*n], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^{(-1+n)*(sin(f*x+e)^2)^{(1/2)}/f/(1-n)}

Rubi [A]

time = 0.06, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2712, 2656}

$$-\frac{b \sqrt{\sin^2(e + fx)} \csc(e + fx) (b \sec(e + fx))^{n-1} {}_2F_1\left(\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^2*(b*Sec[e + f*x])^n,x]

[Out] -((b*Csc[e + f*x]*Hypergeometric2F1[3/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^{(-1 + n)*Sqrt[Sin[e + f*x]^2]})/(f*(1 - n)))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2712

Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[(a^2/b^2)*(a*Sec[e + f*x])^(m - 1)*(b*Csc[e + f*x])^(n + 1)*(a*Cos[e + f*x])^(m - 1)*(b*Sin[e + f*x])^(n + 1), Int[1/((a*Cos[e + f*x])^m*(b*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\begin{aligned} \int \csc^2(e + fx)(b \sec(e + fx))^n dx &= (b^2(b \cos(e + fx))^{-1+n}(b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} \csc^2(e + fx) dx \\ &= -\frac{b \csc(e + fx) {}_2F_1\left(\frac{3}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sqrt{\sin^2(e + fx)}}{f(1-n)} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 13.74, size = 2638, normalized size = 36.14

Result too large to show

Warning: Unable to verify antiderivative.

[In] Integrate[Csc[e + f*x]^2*(b*Sec[e + f*x])^n,x]

[Out] $(\text{Cot}[(e + f*x)/2] * \text{Csc}[e + f*x]^2 * (b * \text{Sec}[e + f*x])^n * (\text{Cos}[(e + f*x)/2]^2 * \text{Sec}[e + f*x])^n * (-\text{AppellF1}[-1/2, n, -n, 1/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * (\text{Cos}[e + f*x] * \text{Sec}[(e + f*x)/2]^2)^n) + (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * (\text{Sec}[(e + f*x)/2]^2)^n * \text{Tan}[(e + f*x)/2]^2) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]) * \text{Tan}[(e + f*x)/2]^2)) / (2 * f * (-1/4 * (\text{Csc}[(e + f*x)/2]^2 * (\text{Cos}[(e + f*x)/2]^2 * \text{Sec}[e + f*x])^n * (-\text{AppellF1}[-1/2, n, -n, 1/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * (\text{Cos}[e + f*x] * \text{Sec}[(e + f*x)/2]^2)^n) + (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * (\text{Sec}[(e + f*x)/2]^2)^n * \text{Tan}[(e + f*x)/2]^2) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]) * \text{Tan}[(e + f*x)/2]^2)) + (\text{Cot}[(e + f*x)/2] * (\text{Cos}[(e + f*x)/2]^2 * \text{Sec}[e + f*x])^n * (-((\text{Cos}[e + f*x] * \text{Sec}[(e + f*x)/2]^2)^n * (-n * \text{AppellF1}[1/2, n, 1 - n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * \text{Sec}[(e + f*x)/2]^2 * \text{Tan}[(e + f*x)/2]) - n * \text{AppellF1}[1/2, 1 + n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * \text{Sec}[(e + f*x)/2]^2 * \text{Tan}[(e + f*x)/2])) - n * \text{AppellF1}[-1/2, n, -n, 1/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * (\text{Cos}[e + f*x] * \text{Sec}[(e + f*x)/2]^2)^{-1 + n} * (-\text{Sec}[(e + f*x)/2]^2 * \text{Sin}[e + f*x]) + \text{Cos}[e + f*x] * \text{Sec}[(e + f*x)/2]^2 * \text{Tan}[(e + f*x)/2]) + (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * (\text{Sec}[(e + f*x)/2]^2)^{1 + n} * \text{Tan}[(e + f*x)/2]) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]) * \text{Tan}[(e + f*x)/2]^2) + (3 * n * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * \text{Sec}[(e + f*x)/2]^2)^n * \text{Tan}[(e + f*x)/2]^3) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2]) * \text{Tan}[(e + f*x)/2]^2) + (3 * (\text{Sec}[(e + f*x)/2]^2)^n * \text{Tan}[(e + f*x)/2]^2 * ((n * \text{AppellF1}[3/2, n, 1 - n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * \text{Sec}[(e + f*x)/2]^2 * \text{Tan}[(e + f*x)/2]) / 3 + (n * \text{AppellF1}[3/2, 1 + n, -n, 5/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] * \text{Sec}[(e + f*x)/2]^2 * \text{Tan}[(e + f*x)/2]) / 3)) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \text{Tan}[(e + f*x)/2]^2, -\text{Tan}[(e + f*x)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1$

- n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2) - (3*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^2*(2*n*(AppellF1[3/2, n, 1 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2] + 3*((n*AppellF1[3/2, n, 1 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/3 + (n*AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/3) + 2*n*Tan[(e + f*x)/2]^2*((-3*(1 - n)*AppellF1[5/2, n, 2 - n, 7/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/5 + (6*n*AppellF1[5/2, 1 + n, 1 - n, 7/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/5 + (3*(1 + n)*AppellF1[5/2, 2 + n, -n, 7/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/5)))/(3*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*n*(AppellF1[3/2, n, 1 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))/2 + (n*Cot[(e + f*x)/2]*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^(-1 + n)*(-(AppellF1[-1/2, n, -n, 1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n) + (3*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*(Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^2)/(3*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*n*(AppellF1[3/2, n, 1 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2))*(-(Cos[(e + f*x)/2]*Sec[e + f*x]*Sin[(e + f*x)/2]) + Cos[(e + f*x)/2]^2*Sec[e + f*x]*Tan[e + f*x]))/2))

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (\csc^2(fx + e)) (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^2*(b*sec(f*x+e))^n,x)

[Out] int(csc(f*x+e)^2*(b*sec(f*x+e))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(b*sec(f*x+e))^n,x, algorithm="maxima")

[Out] integrate((b*sec(f*x + e))^n*csc(f*x + e)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(b*sec(f*x+e))^n,x, algorithm="fricas")

[Out] integral((b*sec(f*x + e))^n*csc(f*x + e)^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \csc^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**2*(b*sec(f*x+e))**n,x)

[Out] Integral((b*sec(e + f*x))**n*csc(e + f*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(b*sec(f*x+e))^n,x, algorithm="giac")

[Out] integrate((b*sec(f*x + e))^n*csc(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^n}{\sin(e+fx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(e + f*x))^n/sin(e + f*x)^2,x)

[Out] int((b/cos(e + f*x))^n/sin(e + f*x)^2, x)

3.502 $\int \csc^4(e + fx)(b \sec(e + fx))^n dx$

Optimal. Leaf size=73

$$-\frac{b \csc(e + fx) {}_2F_1\left(\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sqrt{\sin^2(e + fx)}}{f(1-n)}$$

[Out] -b*csc(f*x+e)*hypergeom([5/2, 1/2-1/2*n], [3/2-1/2*n], cos(f*x+e)^2)*(b*sec(f*x+e))^{(-1+n)*(sin(f*x+e)^2)^{(1/2)}/f/(1-n)}

Rubi [A]

time = 0.06, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {2712, 2656}

$$-\frac{b \sqrt{\sin^2(e + fx)} \csc(e + fx) (b \sec(e + fx))^{n-1} {}_2F_1\left(\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right)}{f(1-n)}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^4*(b*Sec[e + f*x])^n,x]

[Out] -((b*Csc[e + f*x]*Hypergeometric2F1[5/2, (1 - n)/2, (3 - n)/2, Cos[e + f*x]^2]*(b*Sec[e + f*x])^{(-1 + n)*Sqrt[Sin[e + f*x]^2]})/(f*(1 - n)))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2712

Int[(csc[(e_.) + (f_.)*(x_.)]*(b_.))^(n_.)*((a_.)*sec[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[(a^2/b^2)*(a*Sec[e + f*x])^(m - 1)*(b*Csc[e + f*x])^(n + 1)*(a*Cos[e + f*x])^(m - 1)*(b*Sin[e + f*x])^(n + 1), Int[1/((a*Cos[e + f*x])^m*(b*Sin[e + f*x])^n), x], x] /; FreeQ[{a, b, e, f, m, n}, x]

Rubi steps

$$\begin{aligned} \int \csc^4(e + fx)(b \sec(e + fx))^n dx &= (b^2(b \cos(e + fx))^{-1+n}(b \sec(e + fx))^{-1+n}) \int (b \cos(e + fx))^{-n} \csc^4(e + fx) dx \\ &= -\frac{b \csc(e + fx) {}_2F_1\left(\frac{5}{2}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(e + fx)\right) (b \sec(e + fx))^{-1+n} \sqrt{\sin^2(e + fx)}}{f(1-n)} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 15.36, size = 3833, normalized size = 52.51

Result too large to show

Warning: Unable to verify antiderivative.

[In] Integrate[Csc[e + f*x]^4*(b*Sec[e + f*x])^n,x]

[Out] (Cot[(e + f*x)/2]^3*Csc[e + f*x]^4*(b*Sec[e + f*x])^n*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*(-(AppellF1[-3/2, n, -n, -1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n - 9*AppellF1[-1/2, n, -n, 1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^2 + AppellF1[3/2, n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^6 + (27*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^4)/(3*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*n*(AppellF1[3/2, n, 1 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)))/(24*f*(-1/16*(Cot[(e + f*x)/2]^2*Csc[(e + f*x)/2]^2*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*(-(AppellF1[-3/2, n, -n, -1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n - 9*AppellF1[-1/2, n, -n, 1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^2 + AppellF1[3/2, n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^6 + (27*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*(Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^4)/(3*AppellF1[1/2, n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + 2*n*(AppellF1[3/2, n, 1 - n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2] + AppellF1[3/2, 1 + n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2])*Tan[(e + f*x)/2]^2)) + (Cot[(e + f*x)/2]^3*(Cos[(e + f*x)/2]^2*Sec[e + f*x])^n*(-9*AppellF1[-1/2, n, -n, 1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*Sec[(e + f*x)/2]^2*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2] + 3*AppellF1[3/2, n, -n, 5/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*Sec[(e + f*x)/2]^2*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^5 - (Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*(3*n*AppellF1[-1/2, n, 1 - n, 1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2] + 3*n*AppellF1[-1/2, 1 + n, -n, 1/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2]) - 9*(Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^2*(-(n*AppellF1[1/2, n, 1 - n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2]) - n*AppellF1[1/2, 1 + n, -n, 3/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2]*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2]) + (Cos[e + f*x]*Sec[(e + f*x)/2]^2)^n*Tan[(e + f*x)/2]^6*((3*n*AppellF1[5/2, n, 1 - n, 7/2, Tan[(e + f*x)/2]^2, -Tan[(e + f*x)/2]^2)*Sec[(e + f*x)/2]^2*Tan[(e + f*x)/2])/5 + (3*n*AppellF1[5

$$\begin{aligned} & /2, 1 + n, -n, 7/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2 * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 5 - n * \text{AppellF1}[-3/2, n, -n, -1/2, \tan[(e + fx)/2]^2, \\ & -\tan[(e + fx)/2]^2 * (\cos[e + fx] * \sec[(e + fx)/2]^2)^{-1 + n} * (-\sec[(e + fx)/2]^2 * \sin[e + fx]) + \cos[e + fx] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) \\ & - 9 * n * \text{AppellF1}[-1/2, n, -n, 1/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2 * (\cos[e + fx] * \sec[(e + fx)/2]^2)^{-1 + n} * \tan[(e + fx)/2]^2 * (-\sec[(e + fx)/2]^2 * \sin[e + fx]) + \cos[e + fx] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) \\ & + n * \text{AppellF1}[3/2, n, -n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2 * (\cos[e + fx] * \sec[(e + fx)/2]^2)^{-1 + n} * \tan[(e + fx)/2]^6 * (-\sec[(e + fx)/2]^2 * \sin[e + fx]) + \cos[e + fx] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) \\ & + (54 * \text{AppellF1}[1/2, n, -n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2 * (\sec[(e + fx)/2]^2)^{1 + n} * \tan[(e + fx)/2]^3) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \tan[(e + fx)/2]^2) + (27 * n * \text{AppellF1}[1/2, n, -n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2 * (\sec[(e + fx)/2]^2)^n * \tan[(e + fx)/2]^5) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \tan[(e + fx)/2]^2) + (27 * (\sec[(e + fx)/2]^2)^n * \tan[(e + fx)/2]^4 * ((n * \text{AppellF1}[3/2, n, 1 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 3 + (n * \text{AppellF1}[3/2, 1 + n, -n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2]) / 3)) / (3 * \text{AppellF1}[1/2, n, -n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + 2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \tan[(e + fx)/2]^2) - (27 * \text{AppellF1}[1/2, n, -n, 3/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2 * (\sec[(e + fx)/2]^2)^n * \tan[(e + fx)/2]^4 * (2 * n * (\text{AppellF1}[3/2, n, 1 - n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2] + \text{AppellF1}[3/2, 1 + n, -n, 5/2, \tan[(e + fx)/2]^2, -\tan[(e + fx)/2]^2]) * \sec[(e + fx)/2]^2 * \tan[(e + fx)/2] + 3 * (...
\end{aligned}$$

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int (\csc^4(fx + e)) (b \sec(fx + e))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^4*(b*sec(f*x+e))^n,x)

[Out] int(csc(f*x+e)^4*(b*sec(f*x+e))^n,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4*(b*sec(f*x+e))^n,x, algorithm="maxima")`

[Out] `integrate((b*sec(f*x + e))^n*csc(f*x + e)^4, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4*(b*sec(f*x+e))^n,x, algorithm="fricas")`

[Out] `integral((b*sec(f*x + e))^n*csc(f*x + e)^4, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(e + fx))^n \csc^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)**4*(b*sec(f*x+e))**n,x)`

[Out] `Integral((b*sec(e + f*x))**n*csc(e + f*x)**4, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(csc(f*x+e)^4*(b*sec(f*x+e))^n,x, algorithm="giac")`

[Out] `integrate((b*sec(f*x + e))^n*csc(f*x + e)^4, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(e+fx)}\right)^n}{\sin(e+fx)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((b/cos(e + f*x))^n/sin(e + f*x)^4,x)`

[Out] `int((b/cos(e + f*x))^n/sin(e + f*x)^4, x)`

3.503 $\int (b \sec(a + bx))^n (c \sin(a + bx))^{3/2} dx$

Optimal. Leaf size=76

$$\frac{c {}_2F_1\left(-\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) (b \sec(a + bx))^{-1+n} \sqrt{c \sin(a + bx)}}{(1-n) \sqrt[4]{\sin^2(a + bx)}}$$

[Out] -c*hypergeom([-1/4, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*(b*sec(b*x+a))^(-1+n)*(c*sin(b*x+a))^(1/2)/(1-n)/(sin(b*x+a)^2)^(1/4)

Rubi [A]

time = 0.08, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2656}

$$\frac{c \sqrt{c \sin(a + bx)} (b \sec(a + bx))^{n-1} {}_2F_1\left(-\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right)}{(1-n) \sqrt[4]{\sin^2(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[a + b*x])^n*(c*Sin[a + b*x])^(3/2),x]

[Out] -((c*Hypergeometric2F1[-1/4, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*(b*Sec[a + b*x])^(-1 + n)*Sqrt[c*Sin[a + b*x]])/((1 - n)*(Sin[a + b*x]^2)^(1/4)))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_), x_Symbol] := Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^(FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_), x_Symbol] := Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\int (b \sec(a + bx))^n (c \sin(a + bx))^{3/2} dx = (b^2 (b \cos(a + bx))^{-1+n} (b \sec(a + bx))^{-1+n}) \int (b \cos(a + bx))^{-n} (c \sin(a + bx))^{3/2} dx$$

$$= -\frac{c {}_2F_1\left(-\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) (b \sec(a + bx))^{-1+n} \sqrt{c \sin(a + bx)}}{(1-n)^4 \sqrt{\sin^2(a + bx)}}$$

Mathematica [A]

time = 42.08, size = 104, normalized size = 1.37

$$\frac{2 \cos^2(a + bx)^{\frac{1}{2}(-1+n)} (b \sec(a + bx))^{-1+n} (c \sin(a + bx))^{5/2} \left(9 {}_2F_1\left(\frac{5}{4}, \frac{1}{2}(-1+n); \frac{9}{4}; \sin^2(a + bx)\right) + 5 {}_2F_1\left(\frac{9}{4}, \frac{1+n}{2}; \frac{13}{4}; \sin^2(a + bx)\right) \sin^2(a + bx)\right)}{45c}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[a + b*x])^n*(c*Sin[a + b*x])^(3/2),x]

[Out] (2*(Cos[a + b*x]^2)^((-1 + n)/2)*(b*Sec[a + b*x])^(-1 + n)*(c*Sin[a + b*x])^(5/2)*(9*Hypergeometric2F1[5/4, (-1 + n)/2, 9/4, Sin[a + b*x]^2] + 5*Hypergeometric2F1[9/4, (1 + n)/2, 13/4, Sin[a + b*x]^2]*Sin[a + b*x]^2))/(45*c)

Maple [F]

time = 0.08, size = 0, normalized size = 0.00

$$\int (b \sec(bx + a))^n (c \sin(bx + a))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(b*x+a))^n*(c*sin(b*x+a))^(3/2),x)**[Out]** int((b*sec(b*x+a))^n*(c*sin(b*x+a))^(3/2),x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(3/2),x, algorithm="maxima")**[Out]** integrate((c*sin(b*x + a))^(3/2)*(b*sec(b*x + a))^n, x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(3/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(c*sin(b*x + a))*(b*sec(b*x + a))^n*c*sin(b*x + a), x)
```

Sympy [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(b*x+a))**n*(c*sin(b*x+a))**(3/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 6437 deep
```

Giac [F]

```
time = 0.00, size = 0, normalized size = 0.00
```

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(3/2),x, algorithm="giac")
```

```
[Out] integrate((c*sin(b*x + a))^(3/2)*(b*sec(b*x + a))^n, x)
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.01
```

$$\int (c \sin(a + bx))^{3/2} \left(\frac{b}{\cos(a + bx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((c*sin(a + b*x))^(3/2)*(b/cos(a + b*x))^n,x)
```

```
[Out] int((c*sin(a + b*x))^(3/2)*(b/cos(a + b*x))^n, x)
```

3.504 $\int (b \sec(a + bx))^n \sqrt{c \sin(a + bx)} dx$

Optimal. Leaf size=76

$$\frac{{}_2F_1\left(\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) (b \sec(a + bx))^{-1+n} \sqrt[4]{\sin^2(a + bx)}}{(1-n) \sqrt{c \sin(a + bx)}}$$

[Out] -c*hypergeom([1/4, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*(b*sec(b*x+a))^(-1+n)*(sin(b*x+a)^2)^(1/4)/(1-n)/(c*sin(b*x+a))^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$,

Rules used = {2667, 2656}

$$\frac{c \sqrt[4]{\sin^2(a + bx)} (b \sec(a + bx))^{n-1} {}_2F_1\left(\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right)}{(1-n) \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[a + b*x])^n*Sqrt[c*Sin[a + b*x]],x]

[Out] -((c*Hypergeometric2F1[1/4, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*(b*Sec[a + b*x])^(-1 + n)*(Sin[a + b*x]^2)^(1/4))/((1 - n)*Sqrt[c*Sin[a + b*x]]))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\int (b \sec(a + bx))^n \sqrt{c \sin(a + bx)} dx = (b^2 (b \cos(a + bx))^{-1+n} (b \sec(a + bx))^{-1+n}) \int (b \cos(a + bx))^{-n} \sqrt{c \sin(a + bx)} dx$$

$$= -\frac{c {}_2F_1\left(\frac{1}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) (b \sec(a + bx))^{-1+n} \sqrt{\sin^2(a + bx)}}{(1-n) \sqrt{c \sin(a + bx)}}$$

Mathematica [A]

time = 10.13, size = 75, normalized size = 0.99

$$\frac{\cos^2(a + bx)^{\frac{1}{2}(-1+n)} {}_2F_1\left(\frac{3}{4}, \frac{1+n}{2}; \frac{7}{4}; \sin^2(a + bx)\right) (b \sec(a + bx))^n \sqrt{c \sin(a + bx)} \sin(2(a + bx))}{3b}$$

Antiderivative was successfully verified.

`[In] Integrate[(b*Sec[a + b*x])^n*Sqrt[c*Sin[a + b*x]],x]``[Out] ((Cos[a + b*x]^2)^((-1 + n)/2)*Hypergeometric2F1[3/4, (1 + n)/2, 7/4, Sin[a + b*x]^2]*(b*Sec[a + b*x])^n*Sqrt[c*Sin[a + b*x]]*Sin[2*(a + b*x)])/(3*b)`**Maple [F]**

time = 0.08, size = 0, normalized size = 0.00

$$\int (b \sec(bx + a))^n \sqrt{c \sin(bx + a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((b*sec(b*x+a))^n*(c*sin(b*x+a))^(1/2),x)``[Out] int((b*sec(b*x+a))^n*(c*sin(b*x+a))^(1/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(1/2),x, algorithm="maxima")``[Out] integrate(sqrt(c*sin(b*x + a))*(b*sec(b*x + a))^n, x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(1/2),x, algorithm="fricas")`

[Out] `integral(sqrt(c*sin(b*x + a))*(b*sec(b*x + a))^n, x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (b \sec(a + bx))^n \sqrt{c \sin(a + bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(1/2),x)`

[Out] `Integral((b*sec(a + b*x))^n*sqrt(c*sin(a + b*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((b*sec(b*x+a))^n*(c*sin(b*x+a))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(c*sin(b*x + a))*(b*sec(b*x + a))^n, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sqrt{c \sin(a + bx)} \left(\frac{b}{\cos(a + bx)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((c*sin(a + b*x))^(1/2)*(b/cos(a + b*x))^n,x)`

[Out] `int((c*sin(a + b*x))^(1/2)*(b/cos(a + b*x))^n, x)`

$$3.505 \quad \int \frac{(b \sec(a+bx))^n}{\sqrt{c \sin(a+bx)}} dx$$

Optimal. Leaf size=76

$$-\frac{{}_2F_1\left(\frac{3}{4}, \frac{1-n}{2}, \frac{3-n}{2}; \cos^2(a+bx)\right) (b \sec(a+bx))^{-1+n} \sin^2(a+bx)^{3/4}}{(1-n)(c \sin(a+bx))^{3/2}}$$

[Out] -c*hypergeom([3/4, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*(b*sec(b*x+a))^(-1+n)*(sin(b*x+a)^2)^(3/4)/(1-n)/(c*sin(b*x+a))^(3/2)

Rubi [A]

time = 0.07, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {2667, 2656}

$$-\frac{c \sin^2(a+bx)^{3/4} (b \sec(a+bx))^{n-1} {}_2F_1\left(\frac{3}{4}, \frac{1-n}{2}, \frac{3-n}{2}; \cos^2(a+bx)\right)}{(1-n)(c \sin(a+bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[a + b*x])^n/Sqrt[c*Sin[a + b*x]],x]

[Out] -((c*Hypergeometric2F1[3/4, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*(b*Sec[a + b*x])^(-1 + n)*(Sin[a + b*x]^2)^(3/4))/((1 - n)*(c*Sin[a + b*x])^(3/2)))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_)]*(a_.))^m*((b_.)*sin[(e_.) + (f_.)*(x_)])^n, x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^(FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_)])^n*((a_.)*sin[(e_.) + (f_.)*(x_)])^m, x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\begin{aligned} \int \frac{(b \sec(a+bx))^n}{\sqrt{c \sin(a+bx)}} dx &= (b^2 (b \cos(a+bx))^{-1+n} (b \sec(a+bx))^{-1+n}) \int \frac{(b \cos(a+bx))^{-n}}{\sqrt{c \sin(a+bx)}} dx \\ &= -\frac{{}_2F_1\left(\frac{3}{4}, \frac{1-n}{2}, \frac{3-n}{2}; \cos^2(a+bx)\right) (b \sec(a+bx))^{-1+n} \sin^2(a+bx)^{3/4}}{(1-n)(c \sin(a+bx))^{3/2}} \end{aligned}$$

Mathematica [A]

time = 10.11, size = 72, normalized size = 0.95

$$\frac{\cos^2(a + bx)^{\frac{1}{2}(-1+n)} {}_2F_1\left(\frac{1}{4}, \frac{1+n}{2}; \frac{5}{4}; \sin^2(a + bx)\right) (b \sec(a + bx))^n \sin(2(a + bx))}{b \sqrt{c \sin(a + bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[a + b*x])^n/Sqrt[c*Sin[a + b*x]],x]

[Out] ((Cos[a + b*x]^2)^((-1 + n)/2)*Hypergeometric2F1[1/4, (1 + n)/2, 5/4, Sin[a + b*x]^2]*(b*Sec[a + b*x])^n*Sin[2*(a + b*x)]/(b*Sqrt[c*Sin[a + b*x]])

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(bx + a))^n}{\sqrt{c \sin(bx + a)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(b*x+a))^n/(c*sin(b*x+a))^(1/2),x)

[Out] int((b*sec(b*x+a))^n/(c*sin(b*x+a))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(1/2),x, algorithm="maxima")

[Out] integrate((b*sec(b*x + a))^n/sqrt(c*sin(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(c*sin(b*x + a))*(b*sec(b*x + a))^n/(c*sin(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(a + bx))^n}{\sqrt{c \sin(a + bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a)**n/(c*sin(b*x+a))**(1/2),x)

[Out] Integral((b*sec(a + b*x)**n/sqrt(c*sin(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(1/2),x, algorithm="giac")

[Out] integrate((b*sec(b*x + a))^n/sqrt(c*sin(b*x + a)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(a+bx)}\right)^n}{\sqrt{c \sin(a+bx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(a + b*x))^n/(c*sin(a + b*x))^(1/2),x)

[Out] int((b/cos(a + b*x))^n/(c*sin(a + b*x))^(1/2), x)

$$3.506 \quad \int \frac{(b \sec(a+bx))^n}{(c \sin(a+bx))^{3/2}} dx$$

Optimal. Leaf size=78

$$\frac{{}_2F_1\left(\frac{5}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a+bx)\right) (b \sec(a+bx))^{-1+n} \sqrt[4]{\sin^2(a+bx)}}{c(1-n) \sqrt{c \sin(a+bx)}}$$

[Out] -hypergeom([5/4, 1/2-1/2*n], [3/2-1/2*n], cos(b*x+a)^2)*(b*sec(b*x+a))^(1-n)
*(sin(b*x+a)^2)^(1/4)/c/(1-n)/(c*sin(b*x+a))^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 78, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$,
Rules used = {2667, 2656}

$$\frac{\sqrt[4]{\sin^2(a+bx)} (b \sec(a+bx))^{n-1} {}_2F_1\left(\frac{5}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a+bx)\right)}{c(1-n) \sqrt{c \sin(a+bx)}}$$

Antiderivative was successfully verified.

[In] Int[(b*Sec[a + b*x])^n/(c*Sin[a + b*x])^(3/2), x]

[Out] -((Hypergeometric2F1[5/4, (1 - n)/2, (3 - n)/2, Cos[a + b*x]^2]*(b*Sec[a + b*x])^(1 - n)*(Sin[a + b*x]^2)^(1/4))/(c*(1 - n)*Sqrt[c*Sin[a + b*x]]))

Rule 2656

Int[(cos[(e_.) + (f_.)*(x_.)]*(a_.))^(m_.)*((b_.)*sin[(e_.) + (f_.)*(x_.)])^(n_.), x_Symbol] :> Simp[(-b^(2*IntPart[(n - 1)/2] + 1))*(b*Sin[e + f*x])^(2*FracPart[(n - 1)/2])*((a*Cos[e + f*x])^(m + 1)/(a*f*(m + 1)*(Sin[e + f*x]^2)^(FracPart[(n - 1)/2]))*Hypergeometric2F1[(1 + m)/2, (1 - n)/2, (3 + m)/2, Cos[e + f*x]^2], x] /; FreeQ[{a, b, e, f, m, n}, x] && SimplerQ[n, m]

Rule 2667

Int[((b_.)*sec[(e_.) + (f_.)*(x_.)])^(n_.)*((a_.)*sin[(e_.) + (f_.)*(x_.)])^(m_.), x_Symbol] :> Dist[b^2*(b*Cos[e + f*x])^(n - 1)*(b*Sec[e + f*x])^(n - 1), Int[(a*Sin[e + f*x])^m/(b*Cos[e + f*x])^n, x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]

Rubi steps

$$\int \frac{(b \sec(a + bx))^n}{(c \sin(a + bx))^{3/2}} dx = (b^2 (b \cos(a + bx))^{-1+n} (b \sec(a + bx))^{-1+n}) \int \frac{(b \cos(a + bx))^{-n}}{(c \sin(a + bx))^{3/2}} dx$$

$$= - \frac{{}_2F_1\left(\frac{5}{4}, \frac{1-n}{2}; \frac{3-n}{2}; \cos^2(a + bx)\right) (b \sec(a + bx))^{-1+n} \sqrt[4]{\sin^2(a + bx)}}{c(1-n) \sqrt{c \sin(a + bx)}}$$

Mathematica [A]

time = 10.11, size = 73, normalized size = 0.94

$$- \frac{\cos^2(a + bx)^{\frac{1}{2}(-1+n)} {}_2F_1\left(-\frac{1}{4}, \frac{1+n}{2}; \frac{3}{4}; \sin^2(a + bx)\right) (b \sec(a + bx))^n \sin(2(a + bx))}{b(c \sin(a + bx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Sec[a + b*x])^n/(c*Sin[a + b*x])^(3/2),x]

[Out] -((((Cos[a + b*x]^2)^((-1 + n)/2)*Hypergeometric2F1[-1/4, (1 + n)/2, 3/4, Sin[a + b*x]^2]*(b*Sec[a + b*x])^n*Sin[2*(a + b*x)])/(b*(c*Sin[a + b*x])^(3/2))))

Maple [F]

time = 0.06, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(bx + a))^n}{(c \sin(bx + a))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*sec(b*x+a))^n/(c*sin(b*x+a))^(3/2),x)

[Out] int((b*sec(b*x+a))^n/(c*sin(b*x+a))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(3/2),x, algorithm="maxima")

[Out] integrate((b*sec(b*x + a))^n/(c*sin(b*x + a))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(3/2),x, algorithm="fricas")

[Out] integral(-sqrt(c*sin(b*x + a))*(b*sec(b*x + a))^n/(c^2*cos(b*x + a)^2 - c^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(b \sec(a + bx))^n}{(c \sin(a + bx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(3/2),x)

[Out] Integral((b*sec(a + b*x))^n/(c*sin(a + b*x))^(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*sec(b*x+a))^n/(c*sin(b*x+a))^(3/2),x, algorithm="giac")

[Out] integrate((b*sec(b*x + a))^n/(c*sin(b*x + a))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{b}{\cos(a+bx)}\right)^n}{(c \sin(a + bx))^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b/cos(a + b*x))^n/(c*sin(a + b*x))^(3/2),x)

[Out] int((b/cos(a + b*x))^n/(c*sin(a + b*x))^(3/2), x)

3.507 $\int \sqrt{d \csc(e + fx)} \sin^4(e + fx) dx$

Optimal. Leaf size=100

$$-\frac{2d^3 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{10 \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{21f}$$

[Out] $-2/7*d^3*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(5/2)}-10/21*d*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(1/2)}-10/21*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.05, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2720}

$$-\frac{2d^3 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{10 \sqrt{\sin(e + fx)} F\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right) \sqrt{d \csc(e + fx)}}{21f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sin}[e + f*x]^4, x]$

[Out] $(-2*d^3*\text{Cos}[e + f*x])/(7*f*(d*\text{Csc}[e + f*x])^{(5/2)}) - (10*d*\text{Cos}[e + f*x])/(21*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (10*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(21*f)$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3854

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_*))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\csc[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\csc[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856


```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \sqrt{d \csc(e + fx)} \sin^4(e + fx) dx &= d^4 \int \frac{1}{(d \csc(e + fx))^{7/2}} dx \\ &= -\frac{2d^3 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} + \frac{1}{7}(5d^2) \int \frac{1}{(d \csc(e + fx))^{3/2}} dx \\ &= -\frac{2d^3 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{5}{21} \int \sqrt{d \csc(e + fx)} dx \\ &= -\frac{2d^3 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{1}{21} \left(5 \sqrt{d \csc(e + fx)} \right) \\ &= -\frac{2d^3 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{10 \sqrt{d \csc(e + fx)}}{21} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 67, normalized size = 0.67

$$\frac{\sqrt{d \csc(e + fx)} \left(40F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} + 26 \sin(2(e + fx)) - 3 \sin(4(e + fx)) \right)}{84f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x]^4,x]
```

```
[Out] -1/84*(Sqrt[d*Csc[e + f*x]]*(40*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Si
n[e + f*x]] + 26*Sin[2*(e + f*x)] - 3*Sin[4*(e + f*x)]))/f
```

Maple [C] Result contains complex when optimal does not.

time = 1.50, size = 214, normalized size = 2.14

method	result
default	$\frac{\sin(fx+e) \sqrt{\frac{d}{\sin(fx+e)}} \left(-5i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e) - i \sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e) - i \sin(fx+e)}{\sin(fx+e)}}\right) \right)}{2}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(f*x+e)^4*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/21/f*sin(f*x+e)*(d/sin(f*x+e))^(1/2)*(-5*I*(-I*(-1+cos(f*x+e)))/sin(f*x+e)
)^(1/2)*sin(f*x+e)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e)^(1/2)*Elliptic
F(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*((I*cos(f*x+e)
)-I+sin(f*x+e))/sin(f*x+e)^(1/2)+3*cos(f*x+e)^4*2^(1/2)-3*cos(f*x+e)^3*2^(
1/2)-8*cos(f*x+e)^2*2^(1/2)+8*cos(f*x+e)*2^(1/2))/(-1+cos(f*x+e))*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^4*(d*csc(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e)^4, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 103, normalized size = 1.03

$$\frac{2(3 \cos(fx + e)^3 - 8 \cos(fx + e)) \sqrt{\frac{d}{\sin(fx + e)}} \sin(fx + e) - 5i \sqrt{2i d} \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e)) + 5i \sqrt{-2i d} \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e))}{21 f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^4*(d*csc(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/21*(2*(3*cos(f*x + e)^3 - 8*cos(f*x + e))*sqrt(d/sin(f*x + e))*sin(f*x +
e) - 5*I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e
)) + 5*I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x +
e)))/f
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} \sin^4(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)**4*(d*csc(f*x+e))**(1/2),x)
```

```
[Out] Integral(sqrt(d*csc(e + f*x))*sin(e + f*x)**4, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^4*(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^4 \sqrt{\frac{d}{\sin(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4*(d/sin(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^4*(d/sin(e + f*x))^(1/2), x)

3.508 $\int \sqrt{d \csc(e + fx)} \sin^3(e + fx) dx$

Optimal. Leaf size=75

$$-\frac{2d^2 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{6dE\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{5f\sqrt{d \csc(e + fx)}\sqrt{\sin(e + fx)}}$$

[Out] $-2/5*d^2*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(3/2)}-6/5*d*(\sin(1/2*e+1/4*Pi+1/2*f*x))^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2719}

$$\frac{6dE\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right)}{5f\sqrt{\sin(e + fx)}\sqrt{d \csc(e + fx)}} - \frac{2d^2 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x]^3,x]`

[Out] $(-2*d^2*\cos[e + f*x])/(5*f*(d*\csc[e + f*x])^{(3/2)}) + (6*d*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(5*f*\text{Sqrt}[d*\csc[e + f*x]]*\text{Sqrt}[\sin[e + f*x]])$

Rule 16

`Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2719

`Int[Sqrt[sin[(c_) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3854

`Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3856

`Int[(csc[(c_) + (d_)*(x_)]*(b_))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&`

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sqrt{d \csc(e + fx)} \sin^3(e + fx) dx &= d^3 \int \frac{1}{(d \csc(e + fx))^{5/2}} dx \\
&= -\frac{2d^2 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{1}{5}(3d) \int \frac{1}{\sqrt{d \csc(e + fx)}} dx \\
&= -\frac{2d^2 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{(3d) \int \sqrt{\sin(e + fx)} dx}{5\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \\
&= -\frac{2d^2 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{6dE\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{5f\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 62, normalized size = 0.83

$$\frac{2\sqrt{d \csc(e + fx)} \left(3E\left(\frac{1}{4}(-2e + \pi - 2fx) \middle| 2\right) \sqrt{\sin(e + fx)} + \cos(e + fx) \sin^2(e + fx)\right)}{5f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x]^3,x]**[Out]** (-2*Sqrt[d*Csc[e + f*x]]*(3*EllipticE[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]] + Cos[e + f*x]*Sin[e + f*x]^2))/(5*f)**Maple [C]** Result contains complex when optimal does not.

time = 0.14, size = 538, normalized size = 7.17

method	result
default	$\sqrt{\frac{d}{\sin(fx+e)}} \left(-6 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \right) \text{EllipticE}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^3*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
[Out] 1/5/f*(d/sin(f*x+e))^(1/2)*(-6*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+3*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*

$$\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*EllipticF(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2*2^{(1/2)})+\cos(f*x+e)^3*2^{(1/2)}-6*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*EllipticE(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2*2^{(1/2)})+3*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*EllipticF(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2*2^{(1/2)})-4*\cos(f*x+e)*2^{(1/2)}+3*2^{(1/2)})*2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3*(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e)^3, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 100, normalized size = 1.33

$$\frac{2(\cos(fx+e)^3 - \cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}} + 3\sqrt{2i d} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e))) + 3\sqrt{-2i d} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e)))}{5f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3*(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 1/5*(2*(cos(f*x + e)^3 - cos(f*x + e))*sqrt(d/sin(f*x + e)) + 3*sqrt(2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e))))/f

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**3*(d*csc(f*x+e))**(1/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^3*(d*csc(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e)^3, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^3 \sqrt{\frac{d}{\sin(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^3*(d/sin(e + f*x))^(1/2),x)
```

```
[Out] int(sin(e + f*x)^3*(d/sin(e + f*x))^(1/2), x)
```

3.509 $\int \sqrt{d \csc(e + fx)} \sin^2(e + fx) dx$

Optimal. Leaf size=72

$$-\frac{2d \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{2\sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e + fx)}}{3f}$$

[Out] $-2/3*d*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(1/2)}-2/3*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2720}

$$\frac{2\sqrt{\sin(e + fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e + fx)}}{3f} - \frac{2d \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x]^2,x]`

[Out] $(-2*d*\cos[e + f*x])/(3*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*f)$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2720

`Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3854

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]`

Rule 3856

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&`

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sqrt{d \csc(e + fx)} \sin^2(e + fx) dx &= d^2 \int \frac{1}{(d \csc(e + fx))^{3/2}} dx \\
&= -\frac{2d \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{1}{3} \int \sqrt{d \csc(e + fx)} dx \\
&= -\frac{2d \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{1}{3} \left(\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)} \right) \int \frac{1}{\sqrt{\sin(e + fx)}} dx \\
&= -\frac{2d \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{2 \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 55, normalized size = 0.76

$$\frac{\sqrt{d \csc(e + fx)} \left(2F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} + \sin(2(e + fx)) \right)}{3f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x]^2,x]**[Out]** -1/3*(Sqrt[d*Csc[e + f*x]]*(2*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]] + Sin[2*(e + f*x)]))/f**Maple [C]** Result contains complex when optimal does not.

time = 0.21, size = 187, normalized size = 2.60

method	result
default	$ -\frac{\sin(fx+e) \sqrt{\frac{d}{\sin(fx+e)}} \left(i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}}\right) \right)}{3f(-1+\cos(fx+e))} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(f*x+e)^2*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
[Out] -1/3/f*sin(f*x+e)*(d/sin(f*x+e))^(1/2)*(I*sin(f*x+e)*(-I*(-1+cos(f*x+e)))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+cos(f*x+e)^2*2^(1/2)-cos(f*x+e)*2^(1/2))/(-1+cos(f*x+e))*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^2*(d*csc(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 89, normalized size = 1.24

$$\frac{2\sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) \sin(fx+e) + i\sqrt{2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e)) - i\sqrt{-2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e))}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^2*(d*csc(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] -1/3*(2*sqrt(d/sin(f*x + e))*cos(f*x + e)*sin(f*x + e) + I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) - I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/f
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} \sin^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)**2*(d*csc(f*x+e))**(1/2),x)
```

```
[Out] Integral(sqrt(d*csc(e + f*x))*sin(e + f*x)**2, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^2*(d*csc(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e)^2, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^2 \sqrt{\frac{d}{\sin(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^2*(d/sin(e + f*x))^(1/2),x)
```

```
[Out] int(sin(e + f*x)^2*(d/sin(e + f*x))^(1/2), x)
```

3.510 $\int \sqrt{d \csc(e + fx)} \sin(e + fx) dx$

Optimal. Leaf size=44

$$\frac{2dE\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{f\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] -2*d*(sin(1/2*e+1/4*Pi+1/2*f*x)^2)^(1/2)/sin(1/2*e+1/4*Pi+1/2*f*x)*EllipticE(cos(1/2*e+1/4*Pi+1/2*f*x),2^(1/2))/f/(d*csc(f*x+e))^(1/2)/sin(f*x+e)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3856, 2719}

$$\frac{2dE\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right)}{f\sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x],x]

[Out] (2*d*EllipticE[(e - Pi/2 + f*x)/2, 2])/(f*Sqrt[d*Csc[e + f*x]]*Sqrt[Sin[e + f*x]])

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \sqrt{d \csc(e + fx)} \sin(e + fx) dx &= d \int \frac{1}{\sqrt{d \csc(e + fx)}} dx \\
&= \frac{d \int \sqrt{\sin(e + fx)} dx}{\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \\
&= \frac{2dE\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 43, normalized size = 0.98

$$-\frac{2dE\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sqrt[d*Csc[e + f*x]]*Sin[e + f*x],x]``[Out] (-2*d*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/(f*Sqrt[d*Csc[e + f*x]]*Sqrt[Sin[e + f*x]])`**Maple [C]** Result contains complex when optimal does not.

time = 0.20, size = 513, normalized size = 11.66

method	result
risch	$ -\frac{(e^{2i(fx+e)}-1)\sqrt{2}}{f} \sqrt{\frac{ide^{i(fx+e)}}{e^{2i(fx+e)}-1}} e^{-i(fx+e)} + \left(-\frac{2i(id e^{2i(fx+e)} - id)}{d \sqrt{e^{i(fx+e)}(id e^{2i(fx+e)} - id)}} \frac{\sqrt{e^{i(fx+e)} + 1} \sqrt{-2e^{i(fx+e)}}}{\sqrt{e^{i(fx+e)}(id e^{2i(fx+e)} - id)}} \right) $
default	$ -\frac{\left(2 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-i \cos(fx+e)+\sin(fx+e)+i}{\sin(fx+e)}} \right) \text{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\right)}{\dots} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] -1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2))*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2))*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2))*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))
```

$$\frac{f*x+e)}{\sin(f*x+e)}^{1/2} * ((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2} * ((-I*\cos(f*x+e)+\sin(f*x+e)+I)/\sin(f*x+e))^{1/2} * \text{EllipticE}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2}, 1/2*2^{1/2}) - (-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2} * ((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2} * ((-I*\cos(f*x+e)+\sin(f*x+e)+I)/\sin(f*x+e))^{1/2} * \text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2}, 1/2*2^{1/2}) + \cos(f*x+e)*2^{1/2} - 2^{1/2}) * (d/\sin(f*x+e))^{1/2} * 2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)*(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 63, normalized size = 1.43

$$\frac{\sqrt{2i d} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e))) + \sqrt{-2i d} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e)))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)*(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] (sqrt(2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e))))/f

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} \sin(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)*(d*csc(f*x+e))^(1/2),x)

[Out] Integral(sqrt(d*csc(e + f*x))*sin(e + f*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)*(d*csc(f*x+e))^(1/2),x, algorithm="giac")
```

```
[Out] integrate(sqrt(d*csc(f*x + e))*sin(f*x + e), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + f x) \sqrt{\frac{d}{\sin(e + f x)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)*(d/sin(e + f*x))^(1/2),x)
```

```
[Out] int(sin(e + f*x)*(d/sin(e + f*x))^(1/2), x)
```

3.511 $\int \sqrt{d \csc(e + fx)} dx$

Optimal. Leaf size=43

$$\frac{2\sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e + fx)}}{f}$$

[Out] $-2*(\sin(1/2*e+1/4*\pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*\pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*\pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$,

Rules used = {3856, 2720}

$$\frac{2\sqrt{\sin(e + fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[d*Csc[e + f*x]],x]

[Out] $(2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \pi/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/f$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \sqrt{d \csc(e + fx)} dx &= \left(\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)} \right) \int \frac{1}{\sqrt{\sin(e + fx)}} dx \\ &= \frac{2\sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e + fx)}}{f} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 42, normalized size = 0.98

$$\frac{2\sqrt{d\csc(e+fx)} F\left(\frac{1}{4}(-2e+\pi-2fx)|2\right)\sqrt{\sin(e+fx)}}{f}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[d*Csc[e + f*x]],x]**[Out]** (-2*Sqrt[d*Csc[e + f*x]]*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]])/f**Maple [C]** Result contains complex when optimal does not.

time = 0.12, size = 162, normalized size = 3.77

method	result
default	$-\frac{i\sqrt{2}\sqrt{\frac{d}{\sin(fx+e)}}(-1+\cos(fx+e))\sqrt{\frac{i\cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\sqrt{\frac{-i\cos(fx+e)+\sin(fx+e)+i}{\sin(fx+e)}}\sqrt{\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}}}{f\sin(fx+e)^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)

[Out]
$$-I/f*2^{(1/2)}*(d/\sin(f*x+e))^{(1/2)}*(-1+\cos(f*x+e))*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*((-I*\cos(f*x+e)+\sin(f*x+e)+I)/\sin(f*x+e))^{(1/2)}*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}*EllipticF(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2*2^{(1/2)})/\sin(f*x+e)^2*(\cos(f*x+e)+1)^2$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(1/2),x, algorithm="maxima")**[Out]** integrate(sqrt(d*csc(f*x + e)), x)**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 59, normalized size = 1.37

$$\frac{-i\sqrt{2id}\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $(-I\sqrt{2I*d}\text{weierstrassPInverse}(4, 0, \cos(f*x + e) + I\sin(f*x + e)) + I\sqrt{-2I*d}\text{weierstrassPInverse}(4, 0, \cos(f*x + e) - I\sin(f*x + e)))/f$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(f*x+e))**(1/2),x)`

[Out] `Integral(sqrt(d*csc(e + f*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((d*csc(f*x+e))^(1/2),x, algorithm="giac")`

[Out] `integrate(sqrt(d*csc(f*x + e)), x)`

Mupad [B]

time = 0.62, size = 63, normalized size = 1.47

$$\frac{2 \sqrt{\sin(e + fx)} F\left(\operatorname{asin}\left(\frac{\sqrt{2} \sqrt{1 - \sin(e + fx)}}{2}\right) \middle| 2\right) \sqrt{\frac{d}{\sin(e + fx)}} \sqrt{\cos(e + fx)^2}}{f \cos(e + fx)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((d/sin(e + f*x))^(1/2),x)`

[Out] `-(2*sin(e + f*x)^(1/2)*ellipticF(asin((2^(1/2)*(1 - sin(e + f*x))^(1/2))/2), 2)*(d/sin(e + f*x))^(1/2)*(cos(e + f*x)^2)^(1/2))/(f*cos(e + f*x))`

3.512 $\int \csc(e + fx) \sqrt{d \csc(e + fx)} dx$

Optimal. Leaf size=68

$$-\frac{2 \cos(e + fx) \sqrt{d \csc(e + fx)}}{f} - \frac{2dE\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2*\cos(f*x+e)*(d*\csc(f*x+e))^{(1/2)}/f+2*d*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 68, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3853, 3856, 2719}

$$-\frac{2 \cos(e + fx) \sqrt{d \csc(e + fx)}}{f} - \frac{2dE\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right)}{f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]*Sqrt[d*Csc[e + f*x]],x]`

[Out] $(-2*\text{Cos}[e + f*x]*\text{Sqrt}[d*\text{Csc}[e + f*x]])/f - (2*d*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m + n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3853

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n - 1)/(d*(n - 1)), x] + Dist[b^2*((n - 2)/(n - 1)), Int[(b*Csc[c + d*x])^(n - 2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3856

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&`

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \csc(e+fx) \sqrt{d \csc(e+fx)} dx &= \frac{\int (d \csc(e+fx))^{3/2} dx}{d} \\
&= -\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{f} - d \int \frac{1}{\sqrt{d \csc(e+fx)}} dx \\
&= -\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{f} - \frac{d \int \sqrt{\sin(e+fx)} dx}{\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}} \\
&= -\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{f} - \frac{2dE\left(\frac{1}{2}(e-\frac{\pi}{2}+fx) \mid 2\right)}{f \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.10, size = 57, normalized size = 0.84

$$\frac{(d \csc(e+fx))^{3/2} \left(2E\left(\frac{1}{4}(-2e+\pi-2fx) \mid 2\right) \sin^{\frac{3}{2}}(e+fx) - \sin(2(e+fx)) \right)}{df}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]*Sqrt[d*Csc[e + f*x]],x]`

```
[Out] ((d*Csc[e + f*x])^(3/2)*(2*EllipticE[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]
^(3/2) - Sin[2*(e + f*x)]))/(d*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.15, size = 514, normalized size = 7.56

method	result
default	$\frac{\left(2 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}} \right) \text{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)}{\sin(fx+e)}}\right)}{d}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)
*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/s
```

$$\sin(fx+e)^{1/2} * (-(\cos(fx+e) - I - \sin(fx+e)) / \sin(fx+e))^{1/2} * \text{EllipticF}(\frac{(\cos(fx+e) - I + \sin(fx+e)) / \sin(fx+e)}{1/2 * 2^{1/2}}, 1/2 * 2^{1/2}) + 2 * (-I * (-1 + \cos(fx+e)) / \sin(fx+e))^{1/2} * ((\cos(fx+e) - I + \sin(fx+e)) / \sin(fx+e))^{1/2} * (-(\cos(fx+e) - I - \sin(fx+e)) / \sin(fx+e))^{1/2} * \text{EllipticE}(\frac{(\cos(fx+e) - I + \sin(fx+e)) / \sin(fx+e)}{1/2 * 2^{1/2}}, 1/2 * 2^{1/2}) - (-I * (-1 + \cos(fx+e)) / \sin(fx+e))^{1/2} * ((\cos(fx+e) - I + \sin(fx+e)) / \sin(fx+e))^{1/2} * (-(\cos(fx+e) - I - \sin(fx+e)) / \sin(fx+e))^{1/2} * \text{EllipticF}(\frac{(\cos(fx+e) - I + \sin(fx+e)) / \sin(fx+e)}{1/2}, 1/2 * 2^{1/2}) - 2^{1/2}) * (d / \sin(fx+e))^{1/2} * 2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(f*x + e))*csc(f*x + e), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 86, normalized size = 1.26

$$\frac{2 \sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) + \sqrt{2id} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e))) + \sqrt{-2id} \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e)))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] $-(2 * \sqrt{d / \sin(fx + e)}) * \cos(fx + e) + \sqrt{2 * I * d} * \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(fx + e) + I * \sin(fx + e))) + \sqrt{-2 * I * d} * \text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(fx + e) - I * \sin(fx + e))) / f$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))**(1/2),x)

[Out] Integral(sqrt(d*csc(e + f*x))*csc(e + f*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(f*x + e))*csc(f*x + e), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{d}{\sin(e + f x)}}}{\sin(e + f x)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(e + f*x))^(1/2)/sin(e + f*x),x)

[Out] int((d/sin(e + f*x))^(1/2)/sin(e + f*x), x)

3.513 $\int \csc^2(e + fx) \sqrt{d \csc(e + fx)} dx$

Optimal. Leaf size=74

$$-\frac{2 \cos(e + fx)(d \csc(e + fx))^{3/2}}{3df} + \frac{2\sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e + fx)}}{3f}$$

[Out] $-2/3*\cos(f*x+e)*(d*\csc(f*x+e))^{(3/2)}/d/f-2/3*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.03, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2720}

$$\frac{2\sqrt{\sin(e + fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e + fx)}}{3f} - \frac{2 \cos(e + fx)(d \csc(e + fx))^{3/2}}{3df}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^2*\text{Sqrt}[d*\text{Csc}[e + f*x]], x]$

[Out] $(-2*\text{Cos}[e + f*x]*(d*\text{Csc}[e + f*x])^{(3/2)})/(3*d*f) + (2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*f)$

Rule 16

$\text{Int}[(u_.)*(v_)^{(m_.)}*((b_.)*(v_))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3853

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*((b*\csc[c + d*x])^{(n-1)})/(d*(n-1)), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\csc[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\&$

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \csc^2(e + fx) \sqrt{d \csc(e + fx)} dx &= \frac{\int (d \csc(e + fx))^{5/2} dx}{d^2} \\
&= -\frac{2 \cos(e + fx) (d \csc(e + fx))^{3/2}}{3df} + \frac{1}{3} \int \sqrt{d \csc(e + fx)} dx \\
&= -\frac{2 \cos(e + fx) (d \csc(e + fx))^{3/2}}{3df} + \frac{1}{3} \left(\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)} \right) \\
&= -\frac{2 \cos(e + fx) (d \csc(e + fx))^{3/2}}{3df} + \frac{2 \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right)\right)}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 55, normalized size = 0.74

$$\frac{2(d \csc(e + fx))^{3/2} \left(\cos(e + fx) + F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sin^{\frac{3}{2}}(e + fx) \right)}{3df}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2*Sqrt[d*Csc[e + f*x]],x]`

```
[Out] (-2*(d*Csc[e + f*x])^(3/2)*(Cos[e + f*x] + EllipticF[(-2*e + Pi - 2*f*x)/4,
2]*Sin[e + f*x]^(3/2)))/(3*d*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 319, normalized size = 4.31

method	result
default	$ \sqrt{\frac{d}{\sin(fx+e)}} (\cos(fx+e)+1)^2 (-1+\cos(fx+e))^2 \left(i \sqrt{\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}} \right) $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/3/f*(d/sin(f*x+e))^(1/2)*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(I*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*sin(f*x+e)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)+I*sin(f*x+e)
```


$$e) * (-I * (-1 + \cos(f*x+e)) / \sin(f*x+e))^{(1/2)} * ((I * \cos(f*x+e) - I + \sin(f*x+e)) / \sin(f*x+e))^{(1/2)} * (-I * \cos(f*x+e) - I - \sin(f*x+e)) / \sin(f*x+e)^{(1/2)} * \text{EllipticF}(((I * \cos(f*x+e) - I + \sin(f*x+e)) / \sin(f*x+e))^{(1/2)}, 1/2 * 2^{(1/2)}) - \cos(f*x+e) * 2^{(1/2)} / \sin(f*x+e)^5 * 2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(d*csc(f*x + e))*csc(f*x + e)^2, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 105, normalized size = 1.42

$$\frac{-i \sqrt{2id} \sin(fx + e) \text{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e)) + i \sqrt{-2id} \sin(fx + e) \text{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e)) - 2 \sqrt{\frac{d}{\sin(fx + e)}} \cos(fx + e)}{3 f \sin(fx + e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 1/3*(-I*sqrt(2*I*d)*sin(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) + I *sin(f*x + e)) + I*sqrt(-2*I*d)*sin(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)) - 2*sqrt(d/sin(f*x + e))*cos(f*x + e))/(f*sin(f*x + e))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} \csc^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**2*(d*csc(f*x+e))**(1/2),x)

[Out] Integral(sqrt(d*csc(e + f*x))*csc(e + f*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(f*x + e))*csc(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{d}{\sin(e + f x)}}}{\sin(e + f x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(e + f*x))^(1/2)/sin(e + f*x)^2,x)

[Out] int((d/sin(e + f*x))^(1/2)/sin(e + f*x)^2, x)

3.514 $\int \csc^3(e + fx) \sqrt{d \csc(e + fx)} dx$

Optimal. Leaf size=100

$$\frac{6 \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5d^2 f} - \frac{6dE\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{5f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2/5*\cos(f*x+e)*(d*\csc(f*x+e))^{(5/2)}/d^2/f-6/5*\cos(f*x+e)*(d*\csc(f*x+e))^{(1/2)}/f+6/5*d*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 100, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2719}

$$\frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5d^2 f} - \frac{6 \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{6dE\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right)}{5f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^3*\text{Sqrt}[d*\text{Csc}[e + f*x]], x]$

[Out] $(-6*\text{Cos}[e + f*x]*\text{Sqrt}[d*\text{Csc}[e + f*x]])/(5*f) - (2*\text{Cos}[e + f*x]*(d*\text{Csc}[e + f*x])^{(5/2)})/(5*d^2*f) - (6*d*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(5*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3853

$\text{Int}[(\csc[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\csc[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[b^2*(n-2)/(n-1), \text{Int}[(b*\csc[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \& \ \text{IntegerQ}[2*n]$

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
 \int \csc^3(e + fx) \sqrt{d \csc(e + fx)} dx &= \frac{\int (d \csc(e + fx))^{7/2} dx}{d^3} \\
 &= -\frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5d^2 f} + \frac{3 \int (d \csc(e + fx))^{3/2} dx}{5d} \\
 &= -\frac{6 \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5d^2 f} \\
 &= -\frac{6 \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5d^2 f} \\
 &= -\frac{6 \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5d^2 f}
 \end{aligned}$$

Mathematica [A]

time = 0.11, size = 68, normalized size = 0.68

$$\frac{2\sqrt{d \csc(e + fx)} \left(3 \cos(e + fx) + \cot(e + fx) \csc(e + fx) - 3E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} \right)}{5f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]^3*Sqrt[d*Csc[e + f*x]],x]
```

```
[Out] (-2*Sqrt[d*Csc[e + f*x]]*(3*Cos[e + f*x] + Cot[e + f*x]*Csc[e + f*x] - 3*EllipticE[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]]))/(5*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.14, size = 1055, normalized size = 10.55

method	result	size
default	Expression too large to display	1055

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)^3*(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/5/f*(3*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x
```

```

+e))^(1/2), 1/2*2^(1/2))*cos(f*x+e)^3*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)-
6*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+
e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1
/2), 1/2*2^(1/2))*cos(f*x+e)^3*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)+3*((I*c
os(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin
(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2
*2^(1/2))*cos(f*x+e)^2*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)-6*((I*cos(f*x+
e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e)
)^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2*2^(1/2
))*cos(f*x+e)^2*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)-3*cos(f*x+e)*(-I*(-1+
cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2
))*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-
I+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2*2^(1/2))+6*cos(f*x+e)*(-I*(-1+cos(f*x+e
))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*co
s(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x
+e))/sin(f*x+e))^(1/2), 1/2*2^(1/2))-3*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)
*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e
))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/
2), 1/2*2^(1/2))+6*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+si
n(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)
*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2*2^(1/2))+3*co
s(f*x+e)^2*2^(1/2)-cos(f*x+e)*2^(1/2)-3*2^(1/2))*(d/sin(f*x+e))^(1/2)/sin(f
*x+e)^2*2^(1/2)

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^3*(d*csc(f*x+e))^(1/2), x, algorithm="maxima")
```

```
[Out] integrate(sqrt(d*csc(f*x + e))*csc(f*x + e)^3, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 138, normalized size = 1.38

$$\frac{3(\cos(fx+e)^2-1)\sqrt{2i d} \operatorname{weierstrassZeta}(4,0, \operatorname{weierstrassPInverse}(4,0, \cos(fx+e) + i \sin(fx+e))) + 3(\cos(fx+e)^2-1)\sqrt{-2i d} \operatorname{weierstrassZeta}(4,0, \operatorname{weierstrassPInverse}(4,0, \cos(fx+e) - i \sin(fx+e))) + 2(3 \cos(fx+e)^3 - 4 \cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}}}{5(f \cos(fx+e)^2 - f)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^3*(d*csc(f*x+e))^(1/2), x, algorithm="fricas")
```

```
[Out] -1/5*(3*(cos(f*x + e)^2 - 1)*sqrt(2*I*d)*weierstrassZeta(4, 0, weierstrassP
Inverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*(cos(f*x + e)^2 - 1)*sqrt
(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*s
```

$\ln(f*x + e))) + 2*(3*\cos(f*x + e)^3 - 4*\cos(f*x + e))*\sqrt{d/\sin(f*x + e)})$
 $/(f*\cos(f*x + e)^2 - f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{d \csc(e + fx)} \csc^3(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3*(d*csc(f*x+e))**(1/2),x)

[Out] Integral(sqrt(d*csc(e + f*x))*csc(e + f*x)**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3*(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sqrt(d*csc(f*x + e))*csc(f*x + e)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{d}{\sin(e + fx)}}}{\sin(e + fx)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(e + f*x))^(1/2)/sin(e + f*x)^3,x)

[Out] int((d/sin(e + f*x))^(1/2)/sin(e + f*x)^3, x)

3.515 $\int (d \csc(e + fx))^{3/2} \sin^5(e + fx) dx$

Optimal. Leaf size=103

$$-\frac{2d^4 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d^2 \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{10d \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{21f}$$

[Out] $-2/7*d^4*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(5/2)}-10/21*d^2*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(1/2)}-10/21*d*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.05, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2720}

$$-\frac{2d^4 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d^2 \cos(e + fx)}{21f \sqrt{d \csc(e + fx)}} + \frac{10d \sqrt{\sin(e + fx)} F\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right) \sqrt{d \csc(e + fx)}}{21f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^5, x]$

[Out] $(-2*d^4*\text{Cos}[e + f*x])/(7*f*(d*\text{Csc}[e + f*x])^{(5/2)}) - (10*d^2*\text{Cos}[e + f*x])/(21*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (10*d*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(21*f)$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3854

$\text{Int}[(\csc[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\csc[c + d*x])^{(n+1)}/(b*d^n)), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\csc[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int (d \csc(e + fx))^{3/2} \sin^5(e + fx) dx &= d^5 \int \frac{1}{(d \csc(e + fx))^{7/2}} dx \\ &= -\frac{2d^4 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} + \frac{1}{7}(5d^3) \int \frac{1}{(d \csc(e + fx))^{3/2}} dx \\ &= -\frac{2d^4 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d^2 \cos(e + fx)}{21f\sqrt{d \csc(e + fx)}} + \frac{1}{21}(5d) \int \sqrt{d \csc(e + fx)} dx \\ &= -\frac{2d^4 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d^2 \cos(e + fx)}{21f\sqrt{d \csc(e + fx)}} + \frac{1}{21} \left(5d\sqrt{d \csc(e + fx)} \right. \\ &= -\frac{2d^4 \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10d^2 \cos(e + fx)}{21f\sqrt{d \csc(e + fx)}} + \frac{10d\sqrt{d \csc(e + fx)}}{21} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 68, normalized size = 0.66

$$\frac{d\sqrt{d \csc(e + fx)} \left(40F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} + 26 \sin(2(e + fx)) - 3 \sin(4(e + fx)) \right)}{84f}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Csc[e + f*x])^(3/2)*Sin[e + f*x]^5,x]
```

```
[Out] -1/84*(d*Sqrt[d*Csc[e + f*x]]*(40*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[
Sin[e + f*x]] + 26*Sin[2*(e + f*x)] - 3*Sin[4*(e + f*x)]))/f
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 216, normalized size = 2.10

method	result
default	$\left(-5i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{i}{\sin(fx+e)}} \right) / f$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*csc(f*x+e))^(3/2)*sin(f*x+e)^5,x,method=_RETURNVERBOSE)
```



```
[Out] 1/21/f*(-5*I*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*sin(f*x+e)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)+3*cos(f*x+e)^4*2^(1/2)-3*cos(f*x+e)^3*2^(1/2)-8*cos(f*x+e)^2*2^(1/2)+8*cos(f*x+e)*2^(1/2))*(d/sin(f*x+e))^(3/2)*sin(f*x+e)^2/(-1+cos(f*x+e))*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^5,x, algorithm="maxima")
```

```
[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^5, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.13, size = 107, normalized size = 1.04

$$\frac{2(3d\cos(fx+e)^3 - 8d\cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}} \sin(fx+e) - 5i\sqrt{2id} \operatorname{dweierstrassPInverse}(4, 0, \cos(fx+e) + i\sin(fx+e)) + 5i\sqrt{-2id} \operatorname{dweierstrassPInverse}(4, 0, \cos(fx+e) - i\sin(fx+e))}{21f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^5,x, algorithm="fricas")
```

```
[Out] 1/21*(2*(3*d*cos(f*x + e)^3 - 8*d*cos(f*x + e))*sqrt(d/sin(f*x + e))*sin(f*x + e) - 5*I*sqrt(2*I*d)*d*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) + 5*I*sqrt(-2*I*d)*d*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/f
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^5,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^5,x, algorithm="giac")
```

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^5, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^5 \left(\frac{d}{\sin(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^5*(d/sin(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^5*(d/sin(e + f*x))^(3/2), x)

3.516 $\int (d \csc(e + fx))^{3/2} \sin^4(e + fx) dx$

Optimal. Leaf size=77

$$-\frac{2d^3 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{6d^2 E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \mid 2\right)}{5f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2/5*d^3*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(3/2)}-6/5*d^2*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2719}

$$\frac{6d^2 E\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \mid 2\right)}{5f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}} - \frac{2d^3 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^4,x]$

[Out] $(-2*d^3*\text{Cos}[e + f*x])/(5*f*(d*\text{Csc}[e + f*x])^{(3/2)}) + (6*d^2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(5*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3854

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_*)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\csc[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\csc[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_*)^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x] \ \&\&$

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int (d \csc(e + fx))^{3/2} \sin^4(e + fx) dx &= d^4 \int \frac{1}{(d \csc(e + fx))^{5/2}} dx \\
&= -\frac{2d^3 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{1}{5}(3d^2) \int \frac{1}{\sqrt{d \csc(e + fx)}} dx \\
&= -\frac{2d^3 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{(3d^2) \int \sqrt{\sin(e + fx)} dx}{5\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \\
&= -\frac{2d^3 \cos(e + fx)}{5f(d \csc(e + fx))^{3/2}} + \frac{6d^2 E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{5f\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.13, size = 62, normalized size = 0.81

$$\frac{2(d \csc(e + fx))^{3/2} \left(3E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sin^{\frac{3}{2}}(e + fx) + \cos(e + fx) \sin^3(e + fx) \right)}{5f}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[e + f*x])^(3/2)*Sin[e + f*x]^4,x]**[Out]** (-2*(d*Csc[e + f*x])^(3/2)*(3*EllipticE[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]^(3/2) + Cos[e + f*x]*Sin[e + f*x]^3))/(5*f)**Maple [C]** Result contains complex when optimal does not.

time = 0.12, size = 544, normalized size = 7.06

method	result
default	$ \frac{\left(-6 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)+i-\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{5f} $

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d*csc(f*x+e))^(3/2)*sin(f*x+e)^4,x,method=_RETURNVERBOSE)
[Out] 1/5/f*(-6*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+3*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2))

$$\begin{aligned} & e)) / \sin(f*x+e))^{(1/2)} * (- (I*\cos(f*x+e) - I - \sin(f*x+e)) / \sin(f*x+e))^{(1/2)} * \text{EllipticF} \\ & ((I*\cos(f*x+e) - I + \sin(f*x+e)) / \sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)} + \cos(f*x+e) \\ & ^3*2^{(1/2)} - 6*(-I*(-1+\cos(f*x+e)) / \sin(f*x+e))^{(1/2)} * ((I*\cos(f*x+e) - I + \sin(f*x \\ & +e)) / \sin(f*x+e))^{(1/2)} * (- (I*\cos(f*x+e) - I - \sin(f*x+e)) / \sin(f*x+e))^{(1/2)} * \text{EllipticE} \\ & (((I*\cos(f*x+e) - I + \sin(f*x+e)) / \sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)} + 3*(-I*(-1 \\ & +\cos(f*x+e)) / \sin(f*x+e))^{(1/2)} * ((I*\cos(f*x+e) - I + \sin(f*x+e)) / \sin(f*x+e))^{(1/2)} \\ &) * (- (I*\cos(f*x+e) - I - \sin(f*x+e)) / \sin(f*x+e))^{(1/2)} * \text{EllipticF}(((I*\cos(f*x+e) \\ & - I + \sin(f*x+e)) / \sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)} - 4*\cos(f*x+e)*2^{(1/2)} + 3*2^{(1/2)} \\ &)) * (d/\sin(f*x+e))^{(3/2)} * \sin(f*x+e)*2^{(1/2)} \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^4,x, algorithm="maxima")

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^4, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 105, normalized size = 1.36

$$\frac{3\sqrt{2i}d \operatorname{dweierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e))) + 3\sqrt{-2i}d \operatorname{dweierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e))) + 2(d \cos(fx+e)^3 - d \cos(fx+e)) \sqrt{\frac{d}{\sin(fx+e)}}}{5f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^4,x, algorithm="fricas")

[Out] $\frac{1}{5} * (3 * \sqrt{2 * I * d} * d * \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(f * x + e) + I * \sin(f * x + e))) + 3 * \sqrt{-2 * I * d} * d * \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(f * x + e) - I * \sin(f * x + e))) + 2 * (d * \cos(f * x + e)^3 - d * \cos(f * x + e)) * \sqrt{d / \sin(f * x + e)}) / f$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))**(3/2)*sin(f*x+e)**4,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 8008 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^4,x, algorithm="giac")

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + f x)^4 \left(\frac{d}{\sin(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^4*(d/sin(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^4*(d/sin(e + f*x))^(3/2), x)

3.517 $\int (d \csc(e + fx))^{3/2} \sin^3(e + fx) dx$

Optimal. Leaf size=75

$$-\frac{2d^2 \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{2d \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{3f}$$

[Out] $-2/3*d^2*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(1/2)}-2/3*d*(\sin(1/2*e+1/4*Pi+1/2*f*x))^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.04, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2720}

$$\frac{2d \sqrt{\sin(e + fx)} F\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right) \sqrt{d \csc(e + fx)}}{3f} - \frac{2d^2 \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^3, x]$

[Out] $(-2*d^2*\text{Cos}[e + f*x])/(3*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (2*d*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*f)$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3854

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\text{Csc}[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\text{Csc}[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\&$

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int (d \csc(e + fx))^{3/2} \sin^3(e + fx) dx &= d^3 \int \frac{1}{(d \csc(e + fx))^{3/2}} dx \\
&= -\frac{2d^2 \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{1}{3} d \int \sqrt{d \csc(e + fx)} dx \\
&= -\frac{2d^2 \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{1}{3} \left(d \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)} \right) \int \frac{1}{\sqrt{\sin(e + fx)}} dx \\
&= -\frac{2d^2 \cos(e + fx)}{3f \sqrt{d \csc(e + fx)}} + \frac{2d \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 56, normalized size = 0.75

$$-\frac{d \sqrt{d \csc(e + fx)} \left(2F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} + \sin(2(e + fx)) \right)}{3f}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Csc[e + f*x])^(3/2)*Sin[e + f*x]^3,x]``[Out] -1/3*(d*Sqrt[d*Csc[e + f*x]]*(2*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]] + Sin[2*(e + f*x)]))/f`**Maple [C]** Result contains complex when optimal does not.

time = 0.11, size = 189, normalized size = 2.52

method	result
default	$-\frac{\left(i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{i}{-1+\cos(fx+e)}} \right)}{3f(-1+\cos(fx+e))}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*csc(f*x+e))^(3/2)*sin(f*x+e)^3,x,method=_RETURNVERBOSE)`

```
[Out] -1/3/f*(I*sin(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+cos(f*x+e)^2*2^(1/2)-cos(f*x+e)*2^(1/2))*(d/sin(f*x+e))^(3/2)*sin(f*x+e)^2/(-1+cos(f*x+e))*2^(1/2)
```


Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^3,x, algorithm="maxima")
```

```
[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^3, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 92, normalized size = 1.23

$$\frac{2d\sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) \sin(fx+e) + i\sqrt{2id} \operatorname{dweierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e)) - i\sqrt{-2id} \operatorname{dweierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e))}{3f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^3,x, algorithm="fricas")
```

```
[Out] -1/3*(2*d*sqrt(d/sin(f*x + e))*cos(f*x + e)*sin(f*x + e) + I*sqrt(2*I*d)*d*
weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) - I*sqrt(-2*I*d)*d*
*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/f
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))**(3/2)*sin(f*x+e)**3,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 5005 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^3,x, algorithm="giac")
```

```
[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^3, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \sin(e + fx)^3 \left(\frac{d}{\sin(e + fx)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(e + f*x)^3*(d/sin(e + f*x))^(3/2),x)
```

```
[Out] int(sin(e + f*x)^3*(d/sin(e + f*x))^(3/2), x)
```

3.518 $\int (d \csc(e + fx))^{3/2} \sin^2(e + fx) dx$

Optimal. Leaf size=46

$$\frac{2d^2 E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2*d^2*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {16, 3856, 2719}

$$\frac{2d^2 E\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right)}{f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x]^2, x]$

[Out] $(2*d^2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n\}, x \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\text{sin}[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3856

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_*))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

Rubi steps

$$\begin{aligned} \int (d \csc(e + fx))^{3/2} \sin^2(e + fx) dx &= d^2 \int \frac{1}{\sqrt{d \csc(e + fx)}} dx \\ &= \frac{d^2 \int \sqrt{\sin(e + fx)} dx}{\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \\ &= \frac{2d^2 E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 45, normalized size = 0.98

$$\frac{2d^2 E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[(d*Csc[e + f*x])^(3/2)*Sin[e + f*x]^2,x]``[Out] (-2*d^2*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/(f*Sqrt[d*Csc[e + f*x]]*Sqrt[Sin[e + f*x]])`**Maple [C]** Result contains complex when optimal does not.

time = 0.13, size = 531, normalized size = 11.54

method	result
risch	$-\frac{(e^{2i(fx+e)}-1)\sqrt{2}}{f} d \sqrt{\frac{ide^{i(fx+e)}}{e^{2i(fx+e)}-1}} e^{-i(fx+e)} + \left(-\frac{2i(id e^{2i(fx+e)}-id)}{d \sqrt{e^{i(fx+e)}(id e^{2i(fx+e)}-id)}} - \frac{\sqrt{e^{i(fx+e)}+1} \sqrt{-2e}}{\dots} \right)$
default	$-\left(2 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \text{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((d*csc(f*x+e))^(3/2)*sin(f*x+e)^2,x,method=_RETURNVERBOSE)`

```
[Out] -1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2)))/f
```

$$\frac{f*x+e)}{\sin(f*x+e))^{1/2}*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{1/2}*EllipticE(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2))-(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{1/2}*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{1/2}*EllipticF(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{1/2},1/2*2^{1/2))+\cos(f*x+e)*2^{1/2}-2^{1/2})*(d/\sin(f*x+e))^{3/2}*sin(f*x+e)*2^{1/2}}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^2,x, algorithm="maxima")

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^2, x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 65, normalized size = 1.41

$$\frac{\sqrt{2i d} \operatorname{dweierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e))) + \sqrt{-2i d} \operatorname{dweierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e)))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^2,x, algorithm="fricas")

[Out] (sqrt(2*I*d)*d*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + sqrt(-2*I*d)*d*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e))))/f

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))**(3/2)*sin(f*x+e)**2,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3003 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e)^2,x, algorithm="giac")

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + f x)^2 \left(\frac{d}{\sin(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2*(d/sin(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^2*(d/sin(e + f*x))^(3/2), x)

3.519 $\int (d \csc(e + fx))^{3/2} \sin(e + fx) dx$

Optimal. Leaf size=44

$$\frac{2d\sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e + fx)}}{f}$$

[Out] $-2*d*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/f$

Rubi [A]

time = 0.02, antiderivative size = 44, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3856, 2720}

$$\frac{2d\sqrt{\sin(e + fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[e + f*x])^{(3/2)}*\text{Sin}[e + f*x], x]$

[Out] $(2*d*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/f$

Rule 16

$\text{Int}[(u_*)*(v_)^{(m_*)}*((b_*)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 3856

$\text{Int}[(\csc[(c_*) + (d_*)*(x_)]*(b_*))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\text{Csc}[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /;$ FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int (d \csc(e + fx))^{3/2} \sin(e + fx) dx &= d \int \sqrt{d \csc(e + fx)} dx \\ &= \left(d \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)} \right) \int \frac{1}{\sqrt{\sin(e + fx)}} dx \\ &= \frac{2d \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{f} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 43, normalized size = 0.98

$$\frac{2d \sqrt{d \csc(e + fx)} F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)}}{f}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d*Csc[e + f*x])^(3/2)*Sin[e + f*x],x]
```

```
[Out] (-2*d*Sqrt[d*Csc[e + f*x]]*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]])/f
```

Maple [C] Result contains complex when optimal does not.

time = 0.13, size = 162, normalized size = 3.68

method	result
default	$-\frac{i\sqrt{2} (\cos(fx+e)+1)^2 \operatorname{EllipticF}\left(\sqrt{\frac{i\cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{-i\cos(fx+e)+\sin(fx+e)}{\sin(fx+e)}}}{f \sin(fx+e)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((d*csc(f*x+e))^(3/2)*sin(f*x+e),x,method=_RETURNVERBOSE)
```

```
[Out] -I/f*2^(1/2)*(cos(f*x+e)+1)^2*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-1+cos(f*x+e))*(d/sin(f*x+e))^(3/2)/sin(f*x+e)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e),x, algorithm="maxima")

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 61, normalized size = 1.39

$$\frac{-i\sqrt{2id}\operatorname{dweierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}\operatorname{dweierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e),x, algorithm="fricas")

[Out] (-I*sqrt(2*I*d)*d*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) + I*sqrt(-2*I*d)*d*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/f

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(e + fx))^{\frac{3}{2}} \sin(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))**(3/2)*sin(f*x+e),x)

[Out] Integral((d*csc(e + f*x))**(3/2)*sin(e + f*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2)*sin(f*x+e),x, algorithm="giac")

[Out] integrate((d*csc(f*x + e))^(3/2)*sin(f*x + e), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \sin(e + fx) \left(\frac{d}{\sin(e + fx)} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)*(d/sin(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)*(d/sin(e + f*x))^(3/2), x)

3.520 $\int (d \csc(e + fx))^{3/2} dx$

Optimal. Leaf size=71

$$-\frac{2d \cos(e + fx) \sqrt{d \csc(e + fx)}}{f} - \frac{2d^2 E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2*d*\cos(f*x+e)*(d*\csc(f*x+e))^{(1/2)}/f+2*d^2*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3853, 3856, 2719}

$$-\frac{2d^2 E\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right)}{f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}} - \frac{2d \cos(e + fx) \sqrt{d \csc(e + fx)}}{f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d*\text{Csc}[e + f*x])^{(3/2)}, x]$

[Out] $(-2*d*\text{Cos}[e + f*x]*\text{Sqrt}[d*\text{Csc}[e + f*x]])/f - (2*d^2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3853

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\csc[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\csc[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{GtQ}[n, 1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.))^{(n_.)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^{(n-1)}*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{EqQ}[n^2, 1/4]$

Rubi steps

$+e)/\sin(f*x+e))^{(1/2)}*EllipticF(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)},1/2*2^{(1/2)}-2^{(1/2)})*\sin(f*x+e)*(d/\sin(f*x+e))^{(3/2)}*2^{(1/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(f*x + e))^(3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 89, normalized size = 1.25

$$\frac{2d\sqrt{\frac{d}{\sin(fx+e)}}\cos(fx+e)+\sqrt{2id}d\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e)))+\sqrt{-2id}d\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e)))}{f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] $-(2*d*\sqrt{d/\sin(f*x + e)}*\cos(f*x + e) + \sqrt{2*I*d}*d*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(f*x + e) + I*\sin(f*x + e))) + \sqrt{-2*I*d}*d*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(f*x + e) - I*\sin(f*x + e))))/f$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(e + fx))^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))**(3/2),x)

[Out] Integral((d*csc(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{d}{\sin(e + f x)} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(e + f*x))^(3/2),x)

[Out] int((d/sin(e + f*x))^(3/2), x)

3.521 $\int \csc(e + fx)(d \csc(e + fx))^{3/2} dx$

Optimal. Leaf size=72

$$-\frac{2 \cos(e + fx)(d \csc(e + fx))^{3/2}}{3f} + \frac{2d \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e + fx)}}{3f}$$

[Out] $-2/3 \cos(fx+e) \cdot (d \csc(fx+e))^{3/2} / f - 2/3 d \cdot (\sin(1/2 \cdot e + 1/4 \cdot \pi + 1/2 \cdot fx))^{1/2} \cdot (1/2) / \sin(1/2 \cdot e + 1/4 \cdot \pi + 1/2 \cdot fx) \cdot \text{EllipticF}(\cos(1/2 \cdot e + 1/4 \cdot \pi + 1/2 \cdot fx), 2^{1/2}) \cdot (d \csc(fx+e))^{1/2} \cdot \sin(fx+e)^{1/2} / f$

Rubi [A]

time = 0.03, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3853, 3856, 2720}

$$\frac{2d \sqrt{\sin(e + fx)} F\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right) \sqrt{d \csc(e + fx)}}{3f} - \frac{2 \cos(e + fx)(d \csc(e + fx))^{3/2}}{3f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + fx] \cdot (d \cdot \text{Csc}[e + fx])^{3/2}, x]$

[Out] $(-2 \cdot \text{Cos}[e + fx] \cdot (d \cdot \text{Csc}[e + fx])^{3/2}) / (3 \cdot f) + (2 \cdot d \cdot \text{Sqrt}[d \cdot \text{Csc}[e + fx]] \cdot \text{EllipticF}[(e - \pi/2 + fx)/2, 2] \cdot \text{Sqrt}[\text{Sin}[e + fx]]) / (3 \cdot f)$

Rule 16

$\text{Int}[(u \cdot v)^m \cdot (b \cdot v)^n, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u \cdot (b \cdot v)^{m+n}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[c] + (d \cdot x)], x_Symbol] \rightarrow \text{Simp}[(2/d) \cdot \text{EllipticF}[(1/2) \cdot (c - \pi/2 + d \cdot x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 3853

$\text{Int}[(\csc[c] + (d \cdot x)) \cdot (b \cdot x)^n, x_Symbol] \rightarrow \text{Simp}[(-b) \cdot \text{Cos}[c + d \cdot x] \cdot ((b \cdot \csc[c + d \cdot x])^{n-1} / (d \cdot (n-1))), x] + \text{Dist}[b^2 \cdot ((n-2)/(n-1)), \text{Int}[(b \cdot \csc[c + d \cdot x])^{n-2}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2 \cdot n]

Rule 3856

$\text{Int}[(\csc[c] + (d \cdot x)) \cdot (b \cdot x)^n, x_Symbol] \rightarrow \text{Dist}[(b \cdot \csc[c + d \cdot x])^n \cdot \text{Sin}[c + d \cdot x]^n, \text{Int}[1/\text{Sin}[c + d \cdot x]^n, x], x] /;$ FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \csc(e+fx)(d \csc(e+fx))^{3/2} dx &= \frac{\int (d \csc(e+fx))^{5/2} dx}{d} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3f} + \frac{1}{3} d \int \sqrt{d \csc(e+fx)} dx \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3f} + \frac{1}{3} \left(d \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)} \right. \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3f} + \frac{2d \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right)\right)}{3f}
\end{aligned}$$

Mathematica [A]

time = 0.13, size = 58, normalized size = 0.81

$$\frac{(d \csc(e+fx))^{5/2} \left(2F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sin^{5/2}(e+fx) + \sin(2(e+fx)) \right)}{3df}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]*(d*Csc[e + f*x])^(3/2),x]``[Out] -1/3*((d*Csc[e + f*x])^(5/2)*(2*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]^(5/2) + Sin[2*(e + f*x)]))/(d*f)`**Maple** [C] Result contains complex when optimal does not.

time = 0.12, size = 319, normalized size = 4.43

method	result
default	$ \frac{(-1+\cos(fx+e))^2 \left(i \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \sin(fx+e) \operatorname{EllipticF}\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right)\right) \right)}{3df} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)*(d*csc(f*x+e))^(3/2),x,method=_RETURNVERBOSE)`
`[Out] 1/3/f*(-1+cos(f*x+e))^2*(I*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*sin(f*x+e)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)+I*sin(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)`

2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)*2^(1/2))*(cos(f*x+e)+1)^2*(d/sin(f*x+e))^(3/2)/sin(f*x+e)^4*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate((d*csc(f*x + e))^(3/2)*csc(f*x + e), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 108, normalized size = 1.50

$$\frac{-i\sqrt{2id}d\sin(fx+e)\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}d\sin(fx+e)\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))-2d\sqrt{\frac{d}{\sin(fx+e)}}\cos(fx+e)}{3f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] 1/3*(-I*sqrt(2*I*d)*d*sin(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) + I*sqrt(-2*I*d)*d*sin(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)) - 2*d*sqrt(d/sin(f*x + e))*cos(f*x + e))/(f*sin(f*x + e))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(e + fx))^{\frac{3}{2}} \csc(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(3/2),x)

[Out] Integral((d*csc(e + f*x))^(3/2)*csc(e + f*x), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)*(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(f*x + e))^(3/2)*csc(f*x + e), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(e+fx)}\right)^{3/2}}{\sin(e+fx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(e + f*x))^(3/2)/sin(e + f*x),x)

[Out] int((d/sin(e + f*x))^(3/2)/sin(e + f*x), x)

3.522 $\int \csc^2(e + fx)(d \csc(e + fx))^{3/2} dx$

Optimal. Leaf size=103

$$-\frac{6d \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5df} - \frac{6d^2 E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{5f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2/5*\cos(f*x+e)*(d*\csc(f*x+e))^{(5/2)}/d/f-6/5*d*\cos(f*x+e)*(d*\csc(f*x+e))^{(1/2)}/f+6/5*d^2*(\sin(1/2*e+1/4*\text{Pi}+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*\text{Pi}+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*\text{Pi}+1/2*f*x),2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2719}

$$-\frac{6d^2 E\left(\frac{1}{2}(e + fx - \frac{\pi}{2}) \mid 2\right)}{5f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}} - \frac{2 \cos(e + fx) (d \csc(e + fx))^{5/2}}{5df} - \frac{6d \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^2*(d*\text{Csc}[e + f*x])^{(3/2)}, x]$

[Out] $(-6*d*\text{Cos}[e + f*x]*\text{Sqrt}[d*\text{Csc}[e + f*x]])/(5*f) - (2*\text{Cos}[e + f*x]*(d*\text{Csc}[e + f*x])^{(5/2)})/(5*d*f) - (6*d^2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(5*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

$\text{Int}[(u_.)*(v_)^{(m_.)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}[\{b, n\}, x] \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}[\{c, d\}, x]$

Rule 3853

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\csc[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\csc[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}[\{b, c, d\}, x] \ \&\& \ \text{GtQ}[n, 1] \ \& \ \text{IntegerQ}[2*n]$

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n], x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \csc^2(e + fx)(d \csc(e + fx))^{3/2} dx &= \frac{\int (d \csc(e + fx))^{7/2} dx}{d^2} \\ &= -\frac{2 \cos(e + fx)(d \csc(e + fx))^{5/2}}{5df} + \frac{3}{5} \int (d \csc(e + fx))^{3/2} dx \\ &= -\frac{6d \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx)(d \csc(e + fx))^{5/2}}{5df} \\ &= -\frac{6d \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx)(d \csc(e + fx))^{5/2}}{5df} \\ &= -\frac{6d \cos(e + fx) \sqrt{d \csc(e + fx)}}{5f} - \frac{2 \cos(e + fx)(d \csc(e + fx))^{5/2}}{5df} \end{aligned}$$

Mathematica [A]

time = 0.18, size = 68, normalized size = 0.66

$$\frac{(d \csc(e + fx))^{5/2} \left(-7 \cos(e + fx) + 3 \cos(3(e + fx)) + 12E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sin^{5/2}(e + fx) \right)}{10df}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]^2*(d*Csc[e + f*x])^(3/2),x]
```

```
[Out] ((d*Csc[e + f*x])^(5/2)*(-7*Cos[e + f*x] + 3*Cos[3*(e + f*x)] + 12*Elliptic
E[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]^(5/2)))/(10*d*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.15, size = 1055, normalized size = 10.24

method	result	size
default	Expression too large to display	1055

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)^2*(d*csc(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/5/f*(d/sin(f*x+e))^(3/2)*(3*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)
)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-
```

```

I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)^3*(-I*(-1+cos(f*x+e
))/sin(f*x+e))^(1/2)-6*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*
cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f
*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)^3*(-I*(-1+cos(f*x+e))/sin(
f*x+e))^(1/2)+3*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x
+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/
sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)^2*(-I*(-1+cos(f*x+e))/sin(f*x+e))
^(1/2)-6*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-s
in(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x
+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)^2*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)-
3*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x
+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*Elli
pticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+6*cos(f*x
+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(
f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I
*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-3*(-I*(-1+cos(f*x+
e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*c
os(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*
x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+6*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2
)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+
e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1
/2),1/2*2^(1/2))+3*cos(f*x+e)^2*2^(1/2)-cos(f*x+e)*2^(1/2)-3*2^(1/2))/sin(f
*x+e)*2^(1/2)

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2*(d*csc(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((d*csc(f*x + e))^(3/2)*csc(f*x + e)^2, x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 148, normalized size = 1.44

$$\frac{3(d \cos(fx+e)^2 - d)\sqrt{2d} \operatorname{weierstrassZeta}(4,0, \operatorname{weierstrassPInverse}(4,0, \cos(fx+e) + i \sin(fx+e))) + 3(d \cos(fx+e)^2 - d)\sqrt{-2d} \operatorname{weierstrassZeta}(4,0, \operatorname{weierstrassPInverse}(4,0, \cos(fx+e) - i \sin(fx+e))) + 2(3d \cos(fx+e)^2 - 4d \cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}}}{5(f \cos(fx+e)^2 - f)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2*(d*csc(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] -1/5*(3*(d*cos(f*x + e)^2 - d)*sqrt(2*I*d)*weierstrassZeta(4, 0, weierstras
sPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*(d*cos(f*x + e)^2 - d)*
sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) -
```

$I*\sin(f*x + e))) + 2*(3*d*\cos(f*x + e)^3 - 4*d*\cos(f*x + e))*\sqrt{d/\sin(f*x + e)))/(f*\cos(f*x + e)^2 - f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (d \csc(e + fx))^{\frac{3}{2}} \csc^2(e + fx) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**2*(d*csc(f*x+e))**(3/2),x)

[Out] Integral((d*csc(e + f*x))**(3/2)*csc(e + f*x)**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2*(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate((d*csc(f*x + e))^(3/2)*csc(f*x + e)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{d}{\sin(e+fx)}\right)^{3/2}}{\sin(e+fx)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((d/sin(e + f*x))^(3/2)/sin(e + f*x)^2,x)

[Out] int((d/sin(e + f*x))^(3/2)/sin(e + f*x)^2, x)

$$3.523 \quad \int \frac{\sin^3(e+fx)}{\sqrt{d \csc(e+fx)}} dx$$

Optimal. Leaf size=102

$$-\frac{2d^2 \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} - \frac{10 \cos(e+fx)}{21f \sqrt{d \csc(e+fx)}} + \frac{10 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right) \sqrt{\sin(e+fx)}}{21df}$$

[Out] $-2/7*d^2*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(5/2)}-10/21*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(1/2)}-10/21*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/d/f$

Rubi [A]

time = 0.05, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2720}

$$-\frac{2d^2 \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} - \frac{10 \cos(e+fx)}{21f \sqrt{d \csc(e+fx)}} + \frac{10 \sqrt{\sin(e+fx)} F\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right) \sqrt{d \csc(e+fx)}}{21df}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^3/Sqrt[d*Csc[e + f*x]],x]

[Out] $(-2*d^2*\text{Cos}[e + f*x])/(7*f*(d*\text{Csc}[e + f*x])^{(5/2)}) - (10*\text{Cos}[e + f*x])/(21*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (10*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(21*d*f)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d^n)), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n], x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^3(e+fx)}{\sqrt{d \csc(e+fx)}} dx &= d^3 \int \frac{1}{(d \csc(e+fx))^{7/2}} dx \\
&= -\frac{2d^2 \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} + \frac{1}{7}(5d) \int \frac{1}{(d \csc(e+fx))^{3/2}} dx \\
&= -\frac{2d^2 \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} - \frac{10 \cos(e+fx)}{21f \sqrt{d \csc(e+fx)}} + \frac{5 \int \sqrt{d \csc(e+fx)} dx}{21d} \\
&= -\frac{2d^2 \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} - \frac{10 \cos(e+fx)}{21f \sqrt{d \csc(e+fx)}} + \frac{(5 \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)})}{21d} \\
&= -\frac{2d^2 \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} - \frac{10 \cos(e+fx)}{21f \sqrt{d \csc(e+fx)}} + \frac{10 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + \right.\right.} \\
&\quad \left.\left. \right)}{21df}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 70, normalized size = 0.69

$$\frac{\sqrt{d \csc(e+fx)} \left(40F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e+fx)} + 26 \sin(2(e+fx)) - 3 \sin(4(e+fx)) \right)}{84df}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[e + f*x]^3/Sqrt[d*Csc[e + f*x]],x]
```

```
[Out] -1/84*(Sqrt[d*Csc[e + f*x]]*(40*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Si
n[e + f*x]] + 26*Sin[2*(e + f*x)] - 3*Sin[4*(e + f*x)]))/(d*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 208, normalized size = 2.04

method	result
default	$ -\frac{\left(5i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\right)}{21f(-1+\cos(fx+e))} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(sin(f*x+e)^3/(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
[Out] -1/21/f*(5*I*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)
-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin
(f*x+e))^(1/2),1/2*2^(1/2))*sin(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2
)-3*cos(f*x+e)^4*2^(1/2)+3*cos(f*x+e)^3*2^(1/2)+8*cos(f*x+e)^2*2^(1/2)-8*co
s(f*x+e)*2^(1/2))/(-1+cos(f*x+e))/(d/sin(f*x+e))^(1/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^3/(d*csc(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(f*x + e)^3/sqrt(d*csc(f*x + e)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 106, normalized size = 1.04

$$\frac{2(3 \cos(fx + e)^3 - 8 \cos(fx + e)) \sqrt{\frac{d}{\sin(fx + e)}} \sin(fx + e) - 5i \sqrt{2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e)) + 5i \sqrt{-2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e))}{21df}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^3/(d*csc(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] 1/21*(2*(3*cos(f*x + e)^3 - 8*cos(f*x + e))*sqrt(d/sin(f*x + e))*sin(f*x +
e) - 5*I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e
)) + 5*I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x +
e)))/(d*f)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)**3/(d*csc(f*x+e))**(1/2),x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^3/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^3/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^3}{\sqrt{\frac{d}{\sin(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^3/(d/sin(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^3/(d/sin(e + f*x))^(1/2), x)

$$3.524 \quad \int \frac{\sin^2(e+fx)}{\sqrt{d \csc(e+fx)}} dx$$

Optimal. Leaf size=72

$$-\frac{2d \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{6E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{5f \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}$$

[Out] $-2/5*d*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(3/2)}-6/5*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2719}

$$\frac{6E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right)}{5f \sqrt{\sin(e+fx)} \sqrt{d \csc(e+fx)}} - \frac{2d \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^2/Sqrt[d*Csc[e + f*x]],x]

[Out] $(-2*d*\text{Cos}[e + f*x])/(5*f*(d*\text{Csc}[e + f*x])^{(3/2)}) + (6*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(5*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d^n)), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sin^2(e+fx)}{\sqrt{d \csc(e+fx)}} dx &= d^2 \int \frac{1}{(d \csc(e+fx))^{5/2}} dx \\
&= -\frac{2d \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{3}{5} \int \frac{1}{\sqrt{d \csc(e+fx)}} dx \\
&= -\frac{2d \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{3 \int \sqrt{\sin(e+fx)} dx}{5 \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}} \\
&= -\frac{2d \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{6E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{5f \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 57, normalized size = 0.79

$$\frac{-\frac{12E\left(\frac{1}{4}(-2e+\pi-2fx) \mid 2\right)}{\sqrt{\sin(e+fx)}} - 2 \sin(2(e+fx))}{10f \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]^2/Sqrt[d*Csc[e + f*x]],x]`

```
[Out] ((-12*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/Sqrt[Sin[e + f*x]] - 2*Sin[2*(e + f*x)])/(10*f*Sqrt[d*Csc[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.14, size = 546, normalized size = 7.58

method	result
default	$ \frac{\left(-6 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)+i+\sin(fx+e)}{\sin(fx+e)}}\right)\right)}{10f \sqrt{d \csc(e+fx)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)^2/(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/5/f*(-6*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+3
```

```
*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+
e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*Ellip
ticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+cos(f*x+e)
^3*2^(1/2)-6*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x
+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*Elli
pticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+3*(-I*(-1
+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/
2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)
-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-4*cos(f*x+e)*2^(1/2)+3*2^(1/2
)))/(d/sin(f*x+e))^(1/2)/sin(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sin(f*x + e)^2/sqrt(d*csc(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 103, normalized size = 1.43

$$\frac{2(\cos(fx+e)^3 - \cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}} + 3\sqrt{2d}\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))) + 3\sqrt{-2d}\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e)))}{5df}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] 1/5*(2*(cos(f*x + e)^3 - cos(f*x + e))*sqrt(d/sin(f*x + e)) + 3*sqrt(2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + 3*sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e))))/(d*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^2(e + fx)}{\sqrt{d \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)**2/(d*csc(f*x+e))**(1/2),x)

[Out] Integral(sin(e + f*x)**2/sqrt(d*csc(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^2/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^2}{\sqrt{\frac{d}{\sin(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2/(d/sin(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)^2/(d/sin(e + f*x))^(1/2), x)

$$3.525 \quad \int \frac{\sin(e+fx)}{\sqrt{d \csc(e+fx)}} dx$$

Optimal. Leaf size=74

$$-\frac{2 \cos(e+fx)}{3f \sqrt{d \csc(e+fx)}} + \frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{3df}$$

[Out] $-2/3*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(1/2)}-2/3*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)*\sin(f*x+e)^{(1/2)}/d/f$

Rubi [A]

time = 0.03, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3854, 3856, 2720}

$$\frac{2 \sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e+fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{3df} - \frac{2 \cos(e+fx)}{3f \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]/Sqrt[d*Csc[e + f*x]],x]

[Out] $(-2*\text{Cos}[e + f*x])/(3*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*d*f)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d^n)), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sin(e+fx)}{\sqrt{d \csc(e+fx)}} dx &= d \int \frac{1}{(d \csc(e+fx))^{3/2}} dx \\
&= -\frac{2 \cos(e+fx)}{3f \sqrt{d \csc(e+fx)}} + \frac{\int \sqrt{d \csc(e+fx)} dx}{3d} \\
&= -\frac{2 \cos(e+fx)}{3f \sqrt{d \csc(e+fx)}} + \frac{\left(\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}\right) \int \frac{1}{\sqrt{\sin(e+fx)}} dx}{3d} \\
&= -\frac{2 \cos(e+fx)}{3f \sqrt{d \csc(e+fx)}} + \frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \mid 2\right) \sqrt{\sin(e+fx)}}{3df}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 64, normalized size = 0.86

$$-\frac{d \csc^2(e+fx) \left(2F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e+fx)} + \sin(2(e+fx))\right)}{3f(d \csc(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]/Sqrt[d*Csc[e + f*x]], x]``[Out] -1/3*(d*Csc[e + f*x]^2*(2*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]] + Sin[2*(e + f*x)])/(f*(d*Csc[e + f*x])^(3/2))`**Maple** [C] Result contains complex when optimal does not.

time = 0.14, size = 181, normalized size = 2.45

method	result
default	$-\frac{\left(i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{i}\right)}{3f(-1+\cos(fx+e)) \sqrt{\frac{d}{\sin(fx+e)}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)/(d*csc(f*x+e))^(1/2), x, method=_RETURNVERBOSE)``[Out] -1/3/f*(I*sin(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1`

$$\frac{1}{2} * \text{EllipticF}\left(\frac{(I * \cos(f*x+e) - I + \sin(f*x+e)) / \sin(f*x+e)}{\sin(f*x+e)}\right)^{1/2}, 1/2 * 2^{1/2} + \cos(f*x+e)^2 * 2^{1/2} - \cos(f*x+e) * 2^{1/2} / (-1 + \cos(f*x+e)) / (d / \sin(f*x+e))^{1/2} * 2^{1/2}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(sin(f*x + e)/sqrt(d*csc(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.18, size = 92, normalized size = 1.24

$$\frac{2 \sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) \sin(fx+e) + i \sqrt{2id} \text{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e)) - i \sqrt{-2id} \text{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e))}{3df}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out]
$$-1/3 * (2 * \sqrt{d / \sin(f*x + e)} * \cos(f*x + e) * \sin(f*x + e) + I * \sqrt{2 * I * d} * \text{weierstrassPInverse}(4, 0, \cos(f*x + e) + I * \sin(f*x + e)) - I * \sqrt{-2 * I * d} * \text{weierstrassPInverse}(4, 0, \cos(f*x + e) - I * \sin(f*x + e))) / (d * f)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin(e + fx)}{\sqrt{d \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(1/2),x)

[Out] Integral(sin(e + f*x)/sqrt(d*csc(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)}{\sqrt{\frac{d}{\sin(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)/(d/sin(e + f*x))^(1/2),x)

[Out] int(sin(e + f*x)/(d/sin(e + f*x))^(1/2), x)

$$3.526 \quad \int \frac{1}{\sqrt{d \csc(e + fx)}} dx$$

Optimal. Leaf size=43

$$\frac{2E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

[Out] $-2*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {3856, 2719}

$$\frac{2E\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right)}{f \sqrt{\sin(e + fx)} \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

[In] Int[1/Sqrt[d*Csc[e + f*x]],x]

[Out] $(2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{1}{\sqrt{d \csc(e + fx)}} dx &= \frac{\int \sqrt{\sin(e + fx)} dx}{\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \\ &= \frac{2E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 42, normalized size = 0.98

$$\frac{2E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right)}{f \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/Sqrt[d*Csc[e + f*x]],x]
```

```
[Out] (-2*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/(f*Sqrt[d*Csc[e + f*x]]*Sqrt[Sin[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 533, normalized size = 12.40

method	result
risch	$-\frac{i\sqrt{2}}{f \sqrt{\frac{id e^{i(fx+e)}}{e^{2i(fx+e)} - 1}}} + i \left(-\frac{2i(id e^{2i(fx+e)} - id)}{d \sqrt{e^{i(fx+e)}} (id e^{2i(fx+e)} - id)} - \frac{\sqrt{e^{i(fx+e)} + 1} \sqrt{-2e^{i(fx+e)} + 2} \sqrt{-e^{i(fx+e)}}}{\sqrt{id}} \right)$
default	$-\frac{\left(2 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \right) \text{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\right)}{f \sqrt{\frac{id e^{i(fx+e)}}{e^{2i(fx+e)} - 1}}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)
```

```
[Out] -1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+cos(f*x+e)*2^(1/2)-2^(1/2))/(d/sin(f*x+e))^(1/2)/sin(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt(d*csc(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.13, size = 66, normalized size = 1.53

$$\frac{\sqrt{2i d} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e))) + \sqrt{-2i d} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e)))}{df}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] (sqrt(2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e))))/(d*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{d \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(f*x+e))^(1/2),x)

[Out] Integral(1/sqrt(d*csc(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(1/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sqrt{\frac{d}{\sin(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d/sin(e + f*x))^(1/2),x)

[Out] int(1/(d/sin(e + f*x))^(1/2), x)

$$3.527 \quad \int \frac{\csc(e+fx)}{\sqrt{d \csc(e+fx)}} dx$$

Optimal. Leaf size=46

$$\frac{2\sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{df}$$

[Out] $-2*(\sin(1/2*e+1/4*\text{Pi}+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*\text{Pi}+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*\text{Pi}+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/d/f$

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3856, 2720}

$$\frac{2\sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{df}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]/Sqrt[d*Csc[e + f*x]],x]

[Out] $(2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(d*f)$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_)^(n_.), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_.), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\int \frac{\csc(e+fx)}{\sqrt{d \csc(e+fx)}} dx = \frac{\int \sqrt{d \csc(e+fx)} dx}{d}$$

$$= \frac{\left(\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)} \right) \int \frac{1}{\sqrt{\sin(e+fx)}} dx}{d}$$

$$= \frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{df}$$

Mathematica [A]

time = 0.01, size = 45, normalized size = 0.98

$$\frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{4}(-2e + \pi - 2fx) \middle| 2\right) \sqrt{\sin(e+fx)}}{df}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]/Sqrt[d*Csc[e + f*x]],x]``[Out] (-2*Sqrt[d*Csc[e + f*x]]*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]])/(d*f)`**Maple [C]** Result contains complex when optimal does not.

time = 0.19, size = 165, normalized size = 3.59

method	result
default	$-\frac{i(-1+\cos(fx+e)) \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\right)}{f \sqrt{\frac{d}{\sin(fx+e)}} \sin(fx+e)^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)/(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`
`[Out] -I/f*(-1+cos(f*x+e))*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*((cos(f*x+e)+1)^2*2^(1/2)/(d/sin(f*x+e))^(1/2)/sin(f*x+e))^3`
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(1/2),x, algorithm="maxima")

[Out] integrate(csc(f*x + e)/sqrt(d*csc(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.16, size = 62, normalized size = 1.35

$$\frac{-i\sqrt{2id}\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))}{df}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(1/2),x, algorithm="fricas")

[Out] (-I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) + I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/(d*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(e + fx)}{\sqrt{d \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(1/2),x)

[Out] Integral(csc(e + f*x)/sqrt(d*csc(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(e + fx) \sqrt{\frac{d}{\sin(e + fx)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)*(d/sin(e + f*x))^(1/2)),x)

[Out] int(1/(sin(e + f*x)*(d/sin(e + f*x))^(1/2)), x)

$$3.528 \quad \int \frac{\csc^2(e+fx)}{\sqrt{d \csc(e+fx)}} dx$$

Optimal. Leaf size=70

$$-\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{df} - \frac{2E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{f \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}$$

[Out] $-2*\cos(f*x+e)*(d*\csc(f*x+e))^{(1/2)}/d/f+2*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 70, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2719}

$$-\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{df} - \frac{2E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right)}{f \sqrt{\sin(e+fx)} \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^2/Sqrt[d*Csc[e + f*x]],x]

[Out] $(-2*\text{Cos}[e + f*x]*\text{Sqrt}[d*\text{Csc}[e + f*x]])/(d*f) - (2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] :> Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[(-b)*Cos[c + d*x]*((b*Csc[c + d*x])^(n-1)/(d*(n-1))), x] + Dist[b^2*((n-2)/(n-1)), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\csc^2(e+fx)}{\sqrt{d \csc(e+fx)}} dx &= \frac{\int (d \csc(e+fx))^{3/2} dx}{d^2} \\
&= -\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{df} - \int \frac{1}{\sqrt{d \csc(e+fx)}} dx \\
&= -\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{df} - \frac{\int \sqrt{\sin(e+fx)} dx}{\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}} \\
&= -\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{df} - \frac{2E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{f \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 52, normalized size = 0.74

$$\frac{-2 \cot(e+fx) + \frac{2E\left(\frac{1}{4}(-2e+\pi-2fx) \middle| 2\right)}{\sqrt{\sin(e+fx)}}}{f \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2/Sqrt[d*Csc[e + f*x]],x]`

```
[Out] (-2*Cot[e + f*x] + (2*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/Sqrt[Sin[e + f*x]])/(f*Sqrt[d*Csc[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.13, size = 522, normalized size = 7.46

method	result
default	$ \frac{\left(2 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)}{\sin(fx+e)}}\right)\right)}{f \sqrt{d \csc(e+fx)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2/(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)`

```
[Out] 1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)
```



```
*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(
f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/s
in(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(
((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*(-I*(-1+cos(f
*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-
I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin
(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/
2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x
+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(
1/2),1/2*2^(1/2))-2^(1/2))/(d/sin(f*x+e))^(1/2)/sin(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(d*csc(f*x+e))^(1/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^2/sqrt(d*csc(f*x + e)), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 89, normalized size = 1.27

$$\frac{2\sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) + \sqrt{2id} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e))) + \sqrt{-2id} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e)))}{df}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^2/(d*csc(f*x+e))^(1/2),x, algorithm="fricas")
```

```
[Out] -(2*sqrt(d/sin(f*x + e))*cos(f*x + e) + sqrt(2*I*d)*weierstrassZeta(4, 0, w
eierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + sqrt(-2*I*d)*wei
erstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)
)))/(d*f)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(e + fx)}{\sqrt{d \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**2/(d*csc(f*x+e))**(1/2),x)
```

```
[Out] Integral(csc(e + f*x)**2/sqrt(d*csc(e + f*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^2/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^2 \sqrt{\frac{d}{\sin(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^2*(d/sin(e + f*x))^(1/2)),x)

[Out] int(1/(sin(e + f*x)^2*(d/sin(e + f*x))^(1/2)), x)

$$3.529 \quad \int \frac{\csc^3(e+fx)}{\sqrt{d \csc(e+fx)}} dx$$

Optimal. Leaf size=77

$$-\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^2 f} + \frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{3df}$$

[Out] $-2/3*\cos(f*x+e)*(d*\csc(f*x+e))^{(3/2)}/d^2/f-2/3*(\sin(1/2*e+1/4*Pi+1/2*f*x))^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)*\sin(f*x+e)^{(1/2)}/d/f$

Rubi [A]

time = 0.03, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2720}

$$\frac{2 \sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{3df} - \frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^2 f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^3/\text{Sqrt}[d*\text{Csc}[e + f*x]], x]$

[Out] $(-2*\text{Cos}[e + f*x]*(d*\text{Csc}[e + f*x])^{(3/2)})/(3*d^2*f) + (2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*d*f)$

Rule 16

$\text{Int}[(u_*)*(v_*)^{(m_*)}*((b_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /;$ FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_*) + (d_*)*(x_*)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /;$ FreeQ[{c, d}, x]

Rule 3853

$\text{Int}[(\csc[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\csc[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\csc[c + d*x])^{(n-2)}, x], x] /;$ FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3856

$\text{Int}[(\csc[(c_*) + (d_*)*(x_*)]*(b_*))^{(n_*)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /;$ FreeQ[{b, c, d}, x] &&

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(e+fx)}{\sqrt{d \csc(e+fx)}} dx &= \frac{\int (d \csc(e+fx))^{5/2} dx}{d^3} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^2 f} + \frac{\int \sqrt{d \csc(e+fx)} dx}{3d} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^2 f} + \frac{\left(\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}\right) \int \frac{1}{\sqrt{\sin(e+fx)}} dx}{3d} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^2 f} + \frac{2\sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{3df}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 60, normalized size = 0.78

$$-\frac{2 \csc^2(e+fx) \left(\cos(e+fx) + F\left(\frac{1}{4}(-2e + \pi - 2fx) \middle| 2\right) \sin^{\frac{3}{2}}(e+fx) \right)}{3f \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Integrate[Csc[e + f*x]^3/Sqrt[d*Csc[e + f*x]],x]**[Out]** (-2*Csc[e + f*x]^2*(Cos[e + f*x] + EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]^(3/2)))/(3*f*Sqrt[d*Csc[e + f*x]])**Maple [C]** Result contains complex when optimal does not.

time = 0.12, size = 319, normalized size = 4.14

method	result
default	$\frac{(-1+\cos(fx+e))^2 \left(i \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \sin(fx+e) \operatorname{EllipticF}\left[\frac{1}{4}(-2e+\pi-2fx) \middle 2\right] \right)}{3f \sqrt{d \csc(e+fx)}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(csc(f*x+e)^3/(d*csc(f*x+e))^(1/2),x,method=_RETURNVERBOSE)**[Out]** 1/3/f*(-1+cos(f*x+e))^2*(I*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2))*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)

$$e)^{(1/2)} * \sin(f*x+e) * \text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)}) * \cos(f*x+e) + I * \sin(f*x+e) * (-I * (-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)} * ((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)} * (-I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)} * \text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)}) - \cos(f*x+e) * 2^{(1/2)} * (\cos(f*x+e)+1)^2 / \sin(f*x+e)^6 / (d/\sin(f*x+e))^{(1/2)} * 2^{(1/2)}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(d*csc(f*x+e))^(1/2), x, algorithm="maxima")

[Out] integrate(csc(f*x + e)^3/sqrt(d*csc(f*x + e)), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 108, normalized size = 1.40

$$\frac{-i\sqrt{2id}\sin(fx+e)\text{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}\sin(fx+e)\text{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))-2\sqrt{\frac{d}{\sin(fx+e)}}\cos(fx+e)}{3d^f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(d*csc(f*x+e))^(1/2), x, algorithm="fricas")

[Out] $\frac{1}{3} * (-I * \sqrt{2 * I * d} * \sin(f*x + e) * \text{weierstrassPInverse}(4, 0, \cos(f*x + e) + I * \sin(f*x + e)) + I * \sqrt{-2 * I * d} * \sin(f*x + e) * \text{weierstrassPInverse}(4, 0, \cos(f*x + e) - I * \sin(f*x + e)) - 2 * \sqrt{d / \sin(f*x + e)} * \cos(f*x + e)) / (d * f * \sin(f*x + e))$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(e + fx)}{\sqrt{d \csc(e + fx)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**3/(d*csc(f*x+e))**(1/2), x)

[Out] Integral(csc(e + f*x)**3/sqrt(d*csc(e + f*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(d*csc(f*x+e))^(1/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^3/sqrt(d*csc(f*x + e)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^3 \sqrt{\frac{d}{\sin(e + f x)}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^3*(d/sin(e + f*x))^(1/2)),x)

[Out] int(1/(sin(e + f*x)^3*(d/sin(e + f*x))^(1/2)), x)

$$3.530 \quad \int \frac{\sin^2(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=103

$$-\frac{2d \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}} - \frac{10 \cos(e+fx)}{21df \sqrt{d \csc(e+fx)}} + \frac{10 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{21d^2 f}$$

[Out] $-2/7*d*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(5/2)}-10/21*\cos(f*x+e)/d/f/(d*\csc(f*x+e))^{(1/2)}-10/21*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)}*\sin(f*x+e)^{(1/2)}/d^2/f$

Rubi [A]

time = 0.06, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3854, 3856, 2720}

$$\frac{10 \sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e+fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{21d^2 f} - \frac{10 \cos(e+fx)}{21df \sqrt{d \csc(e+fx)}} - \frac{2d \cos(e+fx)}{7f(d \csc(e+fx))^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[Sin[e + f*x]^2/(d*Csc[e + f*x])^(3/2),x]

[Out] $(-2*d*\text{Cos}[e + f*x])/(7*f*(d*\text{Csc}[e + f*x])^{(5/2)}) - (10*\text{Cos}[e + f*x])/(21*d*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (10*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(21*d^2*f)$

Rule 16

Int[(u_)*(v_)^(m_)*((b_)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_)+(d_)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_)+(d_)*(x_)]*(b_))^(n_), x_Symbol] := Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n+1)/(b*d^n)), x] + Dist[(n+1)/(b^2*n), Int[(b*Csc[c + d*x])^(n+2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n, x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned}
\int \frac{\sin^2(e + fx)}{(d \csc(e + fx))^{3/2}} dx &= d^2 \int \frac{1}{(d \csc(e + fx))^{7/2}} dx \\
&= -\frac{2d \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} + \frac{5}{7} \int \frac{1}{(d \csc(e + fx))^{3/2}} dx \\
&= -\frac{2d \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10 \cos(e + fx)}{21df \sqrt{d \csc(e + fx)}} + \frac{5 \int \sqrt{d \csc(e + fx)} dx}{21d^2} \\
&= -\frac{2d \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10 \cos(e + fx)}{21df \sqrt{d \csc(e + fx)}} + \frac{\left(5 \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}\right)}{21d^2} \\
&= -\frac{2d \cos(e + fx)}{7f(d \csc(e + fx))^{5/2}} - \frac{10 \cos(e + fx)}{21df \sqrt{d \csc(e + fx)}} + \frac{10 \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2}\right)\right)}{21d^2 f}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 70, normalized size = 0.68

$$\frac{\sqrt{d \csc(e + fx)} \left(40F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} + 26 \sin(2(e + fx)) - 3 \sin(4(e + fx))\right)}{84d^2 f}$$

Antiderivative was successfully verified.

```
[In] Integrate[Sin[e + f*x]^2/(d*Csc[e + f*x])^(3/2),x]
```

```
[Out] -1/84*(Sqrt[d*Csc[e + f*x]]*(40*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]] + 26*Sin[2*(e + f*x)] - 3*Sin[4*(e + f*x)]))/(d^2*f)
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 216, normalized size = 2.10

method	result
default	$ \frac{\left(-5i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{\frac{i \cos(fx+e)-i \sin(fx+e)}{\sin(fx+e)}}\right)}{21f(-1+\cos(fx+e))\left(\frac{d}{\sin(fx+e)}\right)} $

Verification of antiderivative is not currently implemented for this CAS.


```
[In] int(sin(f*x+e)^2/(d*csc(f*x+e))^(3/2),x,method=_RETURNVERBOSE)
[Out] 1/21/f*(-5*I*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*sin(f*x+e)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)+3*cos(f*x+e)^4*2^(1/2)-3*cos(f*x+e)^3*2^(1/2)-8*cos(f*x+e)^2*2^(1/2)+8*cos(f*x+e)*2^(1/2))/(-1+cos(f*x+e))/(d/sin(f*x+e))^(3/2)/sin(f*x+e)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^2/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(f*x + e)^2/(d*csc(f*x + e))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 106, normalized size = 1.03

$$\frac{2(3 \cos(fx + e)^3 - 8 \cos(fx + e)) \sqrt{\frac{d}{\sin(fx + e)}} \sin(fx + e) - 5i \sqrt{2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e)) + 5i \sqrt{-2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e))}{21 d^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)^2/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] 1/21*(2*(3*cos(f*x + e)^3 - 8*cos(f*x + e))*sqrt(d/sin(f*x + e))*sin(f*x + e) - 5*I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) + 5*I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/(d^2*f)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin^2(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)**2/(d*csc(f*x+e))**(3/2),x)
```

```
[Out] Integral(sin(e + f*x)**2/(d*csc(e + f*x))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)^2/(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)^2/(d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)^2}{\left(\frac{d}{\sin(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)^2/(d/sin(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)^2/(d/sin(e + f*x))^(3/2), x)

$$3.531 \quad \int \frac{\sin(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=74

$$-\frac{2 \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{6E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{5df \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}$$

[Out] $-2/5*\cos(f*x+e)/f/(d*\csc(f*x+e))^{(3/2)}-6/5*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/d/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 74, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.210$, Rules used = {16, 3854, 3856, 2719}

$$\frac{6E\left(\frac{1}{2}\left(e+fx - \frac{\pi}{2}\right) \middle| 2\right)}{5df \sqrt{\sin(e+fx)} \sqrt{d \csc(e+fx)}} - \frac{2 \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sin}[e + f*x]/(d*\text{Csc}[e + f*x])^{(3/2)}, x]$

[Out] $(-2*\text{Cos}[e + f*x])/(5*f*(d*\text{Csc}[e + f*x])^{(3/2)}) + (6*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(5*d*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

$\text{Int}[(u_)*(v_)^{(m_)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2719

$\text{Int}[\text{Sqrt}[\sin[(c_.) + (d_)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticE}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d\}, x]$

Rule 3854

$\text{Int}[(\csc[(c_.) + (d_)*(x_)]*(b_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[\text{Cos}[c + d*x]*((b*\csc[c + d*x])^{(n+1)})/(b*d^n), x] + \text{Dist}[(n+1)/(b^2*n), \text{Int}[(b*\csc[c + d*x])^{(n+2)}, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\& \ \text{LtQ}[n, -1] \ \&\& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_)*(x_)]*(b_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d\}, x \ \&\&$

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\sin(e+fx)}{(d \csc(e+fx))^{3/2}} dx &= d \int \frac{1}{(d \csc(e+fx))^{5/2}} dx \\
&= -\frac{2 \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{3 \int \frac{1}{\sqrt{d \csc(e+fx)}} dx}{5d} \\
&= -\frac{2 \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{3 \int \sqrt{\sin(e+fx)} dx}{5d \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}} \\
&= -\frac{2 \cos(e+fx)}{5f(d \csc(e+fx))^{3/2}} + \frac{6E\left(\frac{1}{2}(e-\frac{\pi}{2}+fx) \mid 2\right)}{5df \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 60, normalized size = 0.81

$$\frac{-\frac{12E\left(\frac{1}{4}(-2e+\pi-2fx) \mid 2\right)}{\sqrt{\sin(e+fx)}} - 2 \sin(2(e+fx))}{10df \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Sin[e + f*x]/(d*Csc[e + f*x])^(3/2), x]`

```
[Out] ((-12*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/Sqrt[Sin[e + f*x]] - 2*Sin[2*(e + f*x)])/(10*d*f*Sqrt[d*Csc[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.13, size = 547, normalized size = 7.39

method	result
default	$ -\frac{\left(6 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \right) \text{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\right)}{10df \sqrt{d \csc(e+fx)}} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(sin(f*x+e)/(d*csc(f*x+e))^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] -1/5/f*(6*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)
```

```

/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-3
*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+
e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*Ellip
ticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+6*(-I*(-1+
cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2
)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-
I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-3*(-I*(-1+cos(f*x+e))/sin(f*x+
e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-
sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*
x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)^3*2^(1/2)+4*cos(f*x+e)*2^(1/2)-3*2^(1/2
)))/(d/sin(f*x+e))^(3/2)/sin(f*x+e)^2*2^(1/2)

```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(sin(f*x + e)/(d*csc(f*x + e))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 103, normalized size = 1.39

$$\frac{2(\cos(fx+e)^3 - \cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}} + 3\sqrt{2d}\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))) + 3\sqrt{-2id}\operatorname{weierstrassZeta}(4,0,\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e)))}{5d^2f}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] 1/5*(2*(cos(f*x + e)^3 - cos(f*x + e))*sqrt(d/sin(f*x + e)) + 3*sqrt(2*I*d)
*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x +
e))) + 3*sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(
f*x + e) - I*sin(f*x + e))))/(d^2*f)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sin(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(sin(f*x+e)/(d*csc(f*x+e))**(3/2),x)
```

```
[Out] Integral(sin(e + f*x)/(d*csc(e + f*x))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(sin(f*x+e)/(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(sin(f*x + e)/(d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sin(e + f x)}{\left(\frac{d}{\sin(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(sin(e + f*x)/(d/sin(e + f*x))^(3/2),x)

[Out] int(sin(e + f*x)/(d/sin(e + f*x))^(3/2), x)

$$3.532 \quad \int \frac{1}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=77

$$-\frac{2 \cos(e+fx)}{3df \sqrt{d \csc(e+fx)}} + \frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{3d^2 f}$$

[Out] $-2/3*\cos(f*x+e)/d/f/(d*\csc(f*x+e))^{(1/2)}-2/3*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)*\sin(f*x+e)^{(1/2)}/d^2/f$

Rubi [A]

time = 0.03, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {3854, 3856, 2720}

$$\frac{2 \sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e+fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{3d^2 f} - \frac{2 \cos(e+fx)}{3df \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[(d*Csc[e + f*x])^(-3/2),x]

[Out] $(-2*\text{Cos}[e + f*x])/(3*d*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]) + (2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*d^2*f)$

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] :> Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3854

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Simp[Cos[c + d*x]*((b*Csc[c + d*x])^(n + 1)/(b*d^n)), x] + Dist[(n + 1)/(b^2*n), Int[(b*Csc[c + d*x])^(n + 2), x], x] /; FreeQ[{b, c, d}, x] && LtQ[n, -1] && IntegerQ[2*n]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] :> Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{1}{(d \csc(e + fx))^{3/2}} dx &= -\frac{2 \cos(e + fx)}{3df \sqrt{d \csc(e + fx)}} + \frac{\int \sqrt{d \csc(e + fx)} dx}{3d^2} \\
&= -\frac{2 \cos(e + fx)}{3df \sqrt{d \csc(e + fx)}} + \frac{\left(\sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}\right) \int \frac{1}{\sqrt{\sin(e + fx)}} dx}{3d^2} \\
&= -\frac{2 \cos(e + fx)}{3df \sqrt{d \csc(e + fx)}} + \frac{2 \sqrt{d \csc(e + fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \mid 2\right) \sqrt{\sin(e + fx)}}{3d^2 f}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 63, normalized size = 0.82

$$-\frac{\csc^2(e + fx) \left(2F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sqrt{\sin(e + fx)} + \sin(2(e + fx))\right)}{3f(d \csc(e + fx))^{3/2}}$$

Antiderivative was successfully verified.

[In] Integrate[(d*Csc[e + f*x])^(-3/2),x]**[Out]** -1/3*(Csc[e + f*x]^2*(2*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sqrt[Sin[e + f*x]] + Sin[2*(e + f*x)])/(f*(d*Csc[e + f*x])^(3/2))**Maple [C]** Result contains complex when optimal does not.

time = 0.10, size = 189, normalized size = 2.45

method	result
default	$-\frac{\left(i \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sin(fx+e) \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{i}\right)}{3f(-1+\cos(fx+e))\left(\frac{d}{\sin(fx+e)}\right)^{\frac{3}{2}} \sin(fx+e)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(d*csc(f*x+e))^(3/2),x,method=_RETURNVERBOSE)

[Out] -1/3/f*(I*sin(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+cos(f*x+e)^2*2^(1/2)-cos(f*x+e)*2^(1/2))/(-1+cos(f*x+e))/(d/sin(f*x+e))^(3/2)/sin(f*x+e)*2^(1/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")`

[Out] `integrate((d*csc(f*x + e))^(-3/2), x)`

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.10, size = 92, normalized size = 1.19

$$\frac{2\sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) \sin(fx+e) + i\sqrt{2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e)) - i\sqrt{-2id} \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e))}{3d^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")`

[Out] `-1/3*(2*sqrt(d/sin(f*x + e))*cos(f*x + e)*sin(f*x + e) + I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) - I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/(d^2*f)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(f*x+e))**(3/2),x)`

[Out] `Integral((d*csc(e + f*x))**(-3/2), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(d*csc(f*x+e))^(3/2),x, algorithm="giac")`

[Out] `integrate((d*csc(f*x + e))^(-3/2), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\left(\frac{d}{\sin(e+fx)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(d/sin(e + f*x))^(3/2),x)`

[Out] `int(1/(d/sin(e + f*x))^(3/2), x)`

$$3.533 \quad \int \frac{\csc(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=46

$$\frac{2E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{df \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}$$

[Out] -2*(sin(1/2*e+1/4*Pi+1/2*f*x)^2)^(1/2)/sin(1/2*e+1/4*Pi+1/2*f*x)*EllipticE(cos(1/2*e+1/4*Pi+1/2*f*x),2^(1/2))/d/f/(d*csc(f*x+e))^(1/2)/sin(f*x+e)^(1/2)

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {16, 3856, 2719}

$$\frac{2E\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right)}{df \sqrt{\sin(e+fx)} \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]/(d*Csc[e + f*x])^(3/2),x]

[Out] (2*EllipticE[(e - Pi/2 + f*x)/2, 2])/(d*f*Sqrt[d*Csc[e + f*x]]*Sqrt[Sin[e + f*x]])

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\int \frac{\csc(e + fx)}{(d \csc(e + fx))^{3/2}} dx = \frac{\int \frac{1}{\sqrt{d \csc(e + fx)}} dx}{d}$$

$$= \frac{\int \sqrt{\sin(e + fx)} dx}{d \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

$$= \frac{2E\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right)}{df \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

Mathematica [A]

time = 0.02, size = 45, normalized size = 0.98

$$-\frac{2E\left(\frac{1}{4}(-2e + \pi - 2fx) \middle| 2\right)}{df \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]/(d*Csc[e + f*x])^(3/2), x]``[Out] (-2*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/(d*f*Sqrt[d*Csc[e + f*x]]*Sqrt[Sin[e + f*x]])`**Maple [C]** Result contains complex when optimal does not.

time = 0.12, size = 533, normalized size = 11.59

method	result
risch	$-\frac{i\sqrt{2}}{fd\sqrt{\frac{ide^{i(fx+e)}}{e^{2i(fx+e)}-1}}} + i\left(-\frac{2i(id e^{2i(fx+e)} - id)}{d\sqrt{e^{i(fx+e)}}(id e^{2i(fx+e)} - id)} - \frac{\sqrt{e^{i(fx+e)} + 1} \sqrt{-2e^{i(fx+e)} + 2} \sqrt{-e^{i(fx+e)}}}{fd}\right)$
default	$-\left(2\cos(fx+e)\sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}}\sqrt{\frac{i\cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\sqrt{-\frac{i\cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}}\right)\text{EllipticE}\left(\sqrt{\frac{i\cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)/(d*csc(f*x+e))^(3/2), x, method=_RETURNVERBOSE)`
`[Out] -1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2*2^(1/2))-cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)`

```
sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF
(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*(-I*(-1+cos(
f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-
(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+si
n(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1
/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*
x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(
1/2),1/2*2^(1/2))+cos(f*x+e)*2^(1/2)-2^(1/2))/(d/sin(f*x+e))^(3/2)/sin(f*x
+e)^2*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)/(d*csc(f*x + e))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 66, normalized size = 1.43

$$\frac{\sqrt{2i d} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) + i \sin(fx + e))) + \sqrt{-2i d} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx + e) - i \sin(fx + e)))}{d^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] (sqrt(2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) +
I*sin(f*x + e))) + sqrt(-2*I*d)*weierstrassZeta(4, 0, weierstrassPInverse(
4, 0, cos(f*x + e) - I*sin(f*x + e))))/(d^2*f)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)/(d*csc(f*x+e))**(3/2),x)
```

```
[Out] Integral(csc(e + f*x)/(d*csc(e + f*x))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)/(d*csc(f*x+e))^(3/2),x, algorithm="giac")
```

```
[Out] integrate(csc(f*x + e)/(d*csc(f*x + e))^(3/2), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(e + f x) \left(\frac{d}{\sin(e + f x)} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(sin(e + f*x)*(d/sin(e + f*x))^(3/2)),x)
```

```
[Out] int(1/(sin(e + f*x)*(d/sin(e + f*x))^(3/2)), x)
```

$$3.534 \quad \int \frac{\csc^2(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=46

$$\frac{2\sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{d^2 f}$$

[Out] -2*(sin(1/2*e+1/4*Pi+1/2*f*x)^2)^(1/2)/sin(1/2*e+1/4*Pi+1/2*f*x)*EllipticF(cos(1/2*e+1/4*Pi+1/2*f*x),2^(1/2))*(d*csc(f*x+e))^(1/2)*sin(f*x+e)^(1/2)/d^2/f

Rubi [A]

time = 0.02, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$,

Rules used = {16, 3856, 2720}

$$\frac{2\sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e + fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{d^2 f}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^2/(d*Csc[e + f*x])^(3/2),x]

[Out] (2*sqrt[d*Csc[e + f*x]]*EllipticF[(e - Pi/2 + f*x)/2, 2]*sqrt[Sin[e + f*x]])/(d^2*f)

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2720

Int[1/Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticF[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3856

Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] && EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned} \int \frac{\csc^2(e+fx)}{(d \csc(e+fx))^{3/2}} dx &= \frac{\int \sqrt{d \csc(e+fx)} dx}{d^2} \\ &= \frac{\left(\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)} \right) \int \frac{1}{\sqrt{\sin(e+fx)}} dx}{d^2} \\ &= \frac{2\sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{d^2 f} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 45, normalized size = 0.98

$$-\frac{2\sqrt{d \csc(e+fx)} F\left(\frac{1}{4}(-2e + \pi - 2fx) \middle| 2\right) \sqrt{\sin(e+fx)}}{d^2 f}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^2/(d*Csc[e + f*x])^(3/2), x]``[Out] (-2*sqrt[d*Csc[e + f*x]]*EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*sqrt[Sin[e + f*x]])/(d^2*f)`**Maple [C]** Result contains complex when optimal does not.

time = 0.12, size = 165, normalized size = 3.59

method	result
default	$-\frac{i\sqrt{2} (\cos(fx+e)+1)^2 \operatorname{EllipticF}\left(\sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}}, \frac{\sqrt{2}}{2}\right) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}}}{f \left(\frac{d}{\sin(fx+e)}\right)^{\frac{3}{2}} \sin(fx+e)^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^2/(d*csc(f*x+e))^(3/2), x, method=_RETURNVERBOSE)``[Out] -I/f*2^(1/2)*(cos(f*x+e)+1)^2*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2), 1/2*2^(1/2))*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*(-I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e)^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-1+cos(f*x+e))/(d/sin(f*x+e))^(3/2)/sin(f*x+e)^4`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")

[Out] integrate(csc(f*x + e)^2/(d*csc(f*x + e))^(3/2), x)

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 0.14, size = 62, normalized size = 1.35

$$\frac{-i\sqrt{2id}\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))}{d^2f}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] (-I*sqrt(2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e)) + I*sqrt(-2*I*d)*weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)))/(d^2*f)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^2(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**2/(d*csc(f*x+e))**(3/2),x)

[Out] Integral(csc(e + f*x)**2/(d*csc(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^2/(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^2/(d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1}{\sin(e + fx)^2 \left(\frac{d}{\sin(e+fx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^2*(d/sin(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^2*(d/sin(e + f*x))^(3/2)), x)

$$3.535 \quad \int \frac{\csc^3(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=73

$$-\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{d^2 f} - \frac{2E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{df \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}$$

[Out] $-2*\cos(f*x+e)*(d*\csc(f*x+e))^{(1/2)}/d^2/f+2*(\sin(1/2*e+1/4*Pi+1/2*f*x)^2)^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticE}(\cos(1/2*e+1/4*Pi+1/2*f*x),2^{(1/2)})/d/f/(d*\csc(f*x+e))^{(1/2)}/\sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2719}

$$-\frac{2 \cos(e+fx) \sqrt{d \csc(e+fx)}}{d^2 f} - \frac{2E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right)}{df \sqrt{\sin(e+fx)} \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] `Int[Csc[e + f*x]^3/(d*Csc[e + f*x])^(3/2),x]`

[Out] $(-2*\text{Cos}[e + f*x]*\text{Sqrt}[d*\text{Csc}[e + f*x]])/(d^2*f) - (2*\text{EllipticE}[(e - \text{Pi}/2 + f*x)/2, 2])/(d*f*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{Sqrt}[\text{Sin}[e + f*x]])$

Rule 16

`Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]`

Rule 2719

`Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]`

Rule 3853

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Dist[b^2*((n-2)/(n-1)), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]`

Rule 3856

`Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^(n_), x_Symbol] := Dist[(b*Csc[c + d*x])^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&`

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\csc^3(e + fx)}{(d \csc(e + fx))^{3/2}} dx &= \frac{\int (d \csc(e + fx))^{3/2} dx}{d^3} \\
&= -\frac{2 \cos(e + fx) \sqrt{d \csc(e + fx)}}{d^2 f} - \frac{\int \frac{1}{\sqrt{d \csc(e + fx)}} dx}{d} \\
&= -\frac{2 \cos(e + fx) \sqrt{d \csc(e + fx)}}{d^2 f} - \frac{\int \sqrt{\sin(e + fx)} dx}{d \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}} \\
&= -\frac{2 \cos(e + fx) \sqrt{d \csc(e + fx)}}{d^2 f} - \frac{2E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{df \sqrt{d \csc(e + fx)} \sqrt{\sin(e + fx)}}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 55, normalized size = 0.75

$$\frac{-2 \cot(e + fx) + \frac{2E\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right)}{\sqrt{\sin(e + fx)}}}{df \sqrt{d \csc(e + fx)}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^3/(d*Csc[e + f*x])^(3/2), x]`

```
[Out] (-2*Cot[e + f*x] + (2*EllipticE[(-2*e + Pi - 2*f*x)/4, 2])/Sqrt[Sin[e + f*x]])/(d*f*Sqrt[d*Csc[e + f*x]])
```

Maple [C] Result contains complex when optimal does not.

time = 0.11, size = 510, normalized size = 6.99

method	result
default	$ \frac{\left(2 \cos(fx+e) \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{\frac{-i \cos(fx+e)+\sin(fx+e)+i}{\sin(fx+e)}} \operatorname{EllipticE}\left(\sqrt{\frac{i \cos(fx+e)}{\sin(fx+e)}}\right)\right)}{d^2 f} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^3/(d*csc(f*x+e))^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/f*(2*cos(f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2))
```

```
*EllipticE(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(
f*x+e)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/s
in(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2)*EllipticF(
((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))+2*(-I*(-1+cos(f
*x+e))/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*((-
I*cos(f*x+e)+sin(f*x+e)+I)/sin(f*x+e))^(1/2)*EllipticE(((I*cos(f*x+e)-I+sin
(f*x+e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/
2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*((-I*cos(f*x+e)+sin(f*x+e
)+I)/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(
1/2),1/2*2^(1/2))-2^(1/2))/(d/sin(f*x+e))^(3/2)/sin(f*x+e)^2*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^3/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^3/(d*csc(f*x + e))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 89, normalized size = 1.22

$$\frac{2\sqrt{\frac{d}{\sin(fx+e)}} \cos(fx+e) + \sqrt{2id} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) + i \sin(fx+e))) + \sqrt{-2id} \operatorname{weierstrassZeta}(4, 0, \operatorname{weierstrassPInverse}(4, 0, \cos(fx+e) - i \sin(fx+e)))}{d^2 f}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^3/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] -(2*sqrt(d/sin(f*x + e))*cos(f*x + e) + sqrt(2*I*d)*weierstrassZeta(4, 0, w
eierstrassPInverse(4, 0, cos(f*x + e) + I*sin(f*x + e))) + sqrt(-2*I*d)*wei
eierstrassZeta(4, 0, weierstrassPInverse(4, 0, cos(f*x + e) - I*sin(f*x + e)
)))/(d^2*f)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^3(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**3/(d*csc(f*x+e))**(3/2),x)
```

```
[Out] Integral(csc(e + f*x)**3/(d*csc(e + f*x))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^3/(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^3/(d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^3 \left(\frac{d}{\sin(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^3*(d/sin(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^3*(d/sin(e + f*x))^(3/2)), x)

$$3.536 \quad \int \frac{\csc^4(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=77

$$-\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^3 f} + \frac{2 \sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \middle| 2\right) \sqrt{\sin(e+fx)}}{3d^2 f}$$

[Out] $-2/3*\cos(f*x+e)*(d*\csc(f*x+e))^{(3/2)}/d^3/f-2/3*(\sin(1/2*e+1/4*Pi+1/2*f*x))^{(1/2)}/\sin(1/2*e+1/4*Pi+1/2*f*x)*\text{EllipticF}(\cos(1/2*e+1/4*Pi+1/2*f*x), 2^{(1/2)})*(d*\csc(f*x+e))^{(1/2)*\sin(f*x+e)^{(1/2)}/d^2/f$

Rubi [A]

time = 0.03, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2720}

$$\frac{2 \sqrt{\sin(e+fx)} F\left(\frac{1}{2}\left(e+fx - \frac{\pi}{2}\right) \middle| 2\right) \sqrt{d \csc(e+fx)}}{3d^2 f} - \frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^3 f}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Csc}[e + f*x]^4/(d*\text{Csc}[e + f*x])^{(3/2)}, x]$

[Out] $(-2*\text{Cos}[e + f*x]*(d*\text{Csc}[e + f*x])^{(3/2)})/(3*d^3*f) + (2*\text{Sqrt}[d*\text{Csc}[e + f*x]]*\text{EllipticF}[(e - \text{Pi}/2 + f*x)/2, 2]*\text{Sqrt}[\text{Sin}[e + f*x]])/(3*d^2*f)$

Rule 16

$\text{Int}[(u_.)*(v_)^{(m_.)}*((b_)*(v_))^{(n_)}, x_Symbol] \rightarrow \text{Dist}[1/b^m, \text{Int}[u*(b*v)^{(m+n)}, x], x] /; \text{FreeQ}\{b, n, x\} \ \&\& \ \text{IntegerQ}[m]$

Rule 2720

$\text{Int}[1/\text{Sqrt}[\sin[(c_.) + (d_.)*(x_)]], x_Symbol] \rightarrow \text{Simp}[(2/d)*\text{EllipticF}[(1/2)*(c - \text{Pi}/2 + d*x), 2], x] /; \text{FreeQ}\{c, d, x\}$

Rule 3853

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.)^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(-b)*\text{Cos}[c + d*x]*(b*\csc[c + d*x])^{(n-1)}/(d*(n-1)), x] + \text{Dist}[b^2*((n-2)/(n-1)), \text{Int}[(b*\csc[c + d*x])^{(n-2)}, x], x] /; \text{FreeQ}\{b, c, d, x\} \ \&\& \ \text{GtQ}[n, 1] \ \& \ \text{IntegerQ}[2*n]$

Rule 3856

$\text{Int}[(\csc[(c_.) + (d_.)*(x_)]*(b_.)^{(n_)}, x_Symbol] \rightarrow \text{Dist}[(b*\csc[c + d*x])^n*\text{Sin}[c + d*x]^n, \text{Int}[1/\text{Sin}[c + d*x]^n, x], x] /; \text{FreeQ}\{b, c, d, x\} \ \&\&$

EqQ[n^2, 1/4]

Rubi steps

$$\begin{aligned}
\int \frac{\csc^4(e+fx)}{(d \csc(e+fx))^{3/2}} dx &= \frac{\int (d \csc(e+fx))^{5/2} dx}{d^4} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^3 f} + \frac{\int \sqrt{d \csc(e+fx)} dx}{3d^2} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^3 f} + \frac{\left(\sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}\right) \int \frac{1}{\sqrt{\sin(e+fx)}} dx}{3d^2} \\
&= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{3/2}}{3d^3 f} + \frac{2\sqrt{d \csc(e+fx)} F\left(\frac{1}{2}\left(e - \frac{\pi}{2} + fx\right) \mid 2\right) \sqrt{\sin(e+fx)}}{3d^2 f}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 60, normalized size = 0.78

$$\frac{2 \csc^3(e+fx) \left(\cos(e+fx) + F\left(\frac{1}{4}(-2e + \pi - 2fx) \mid 2\right) \sin^{\frac{3}{2}}(e+fx) \right)}{3f(d \csc(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[Csc[e + f*x]^4/(d*Csc[e + f*x])^(3/2), x]`

```
[Out] (-2*Csc[e + f*x]^3*(Cos[e + f*x] + EllipticF[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]^(3/2)))/(3*f*(d*Csc[e + f*x])^(3/2))
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 319, normalized size = 4.14

method	result
default	$ \frac{(\cos(fx+e)+1)^2(-1+\cos(fx+e))^2 \left(i \sqrt{\frac{i \cos(fx+e)-i+\sin(fx+e)}{\sin(fx+e)}} \sqrt{-\frac{i(-1+\cos(fx+e))}{\sin(fx+e)}} \sqrt{-\frac{i \cos(fx+e)-i-\sin(fx+e)}{\sin(fx+e)}} \right)}{\sin(fx+e)} $

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(csc(f*x+e)^4/(d*csc(f*x+e))^(3/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/3/f*(cos(f*x+e)+1)^2*(-1+cos(f*x+e))^2*(I*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-I*(-1+cos(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*sin(f*x+e)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+e))
```

```
)/sin(f*x+e))^(1/2),1/2*2^(1/2))*cos(f*x+e)+I*sin(f*x+e)*(-I*(-1+cos(f*x+e)
)/sin(f*x+e))^(1/2)*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos
(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticF(((I*cos(f*x+e)-I+sin(f*x+
e))/sin(f*x+e))^(1/2),1/2*2^(1/2))-cos(f*x+e)*2^(1/2))/sin(f*x+e)^7/(d/sin(
f*x+e))^(3/2)*2^(1/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^4/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(csc(f*x + e)^4/(d*csc(f*x + e))^(3/2), x)
```

Fricas [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.10, size = 108, normalized size = 1.40

$$\frac{-i\sqrt{2id}\sin(fx+e)\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e))+i\sqrt{-2id}\sin(fx+e)\operatorname{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e))-2\sqrt{\frac{d}{\sin(fx+e)}}\cos(fx+e)}{3d^2f\sin(fx+e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)^4/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")
```

```
[Out] 1/3*(-I*sqrt(2*I*d)*sin(f*x + e)*weierstrassPInverse(4, 0, cos(f*x + e) + I
*sin(f*x + e)) + I*sqrt(-2*I*d)*sin(f*x + e)*weierstrassPInverse(4, 0, cos(
f*x + e) - I*sin(f*x + e)) - 2*sqrt(d/sin(f*x + e))*cos(f*x + e))/(d^2*f*si
n(f*x + e))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^4(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(csc(f*x+e)**4/(d*csc(f*x+e))**(3/2),x)
```

```
[Out] Integral(csc(e + f*x)**4/(d*csc(e + f*x))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^4/(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^4/(d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + f x)^4 \left(\frac{d}{\sin(e + f x)}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^4*(d/sin(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^4*(d/sin(e + f*x))^(3/2)), x)

$$3.537 \quad \int \frac{\csc^5(e+fx)}{(d \csc(e+fx))^{3/2}} dx$$

Optimal. Leaf size=105

$$\frac{6 \cos(e+fx) \sqrt{d \csc(e+fx)}}{5d^2 f} - \frac{2 \cos(e+fx) (d \csc(e+fx))^{5/2}}{5d^4 f} - \frac{6E\left(\frac{1}{2}(e - \frac{\pi}{2} + fx) \mid 2\right)}{5df \sqrt{d \csc(e+fx)} \sqrt{\sin(e+fx)}}$$

[Out] $-2/5 \cos(f*x+e) * (d * \csc(f*x+e))^{(5/2)} / d^4 / f - 6/5 \cos(f*x+e) * (d * \csc(f*x+e))^{(1/2)} / d^2 / f + 6/5 * (\sin(1/2 * e + 1/4 * \text{Pi} + 1/2 * f * x))^2)^{(1/2)} / \sin(1/2 * e + 1/4 * \text{Pi} + 1/2 * f * x) * \text{EllipticE}(\cos(1/2 * e + 1/4 * \text{Pi} + 1/2 * f * x), 2^{(1/2)}) / d / f / (d * \csc(f*x+e))^{(1/2)} / \sin(f*x+e)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 105, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.190$, Rules used = {16, 3853, 3856, 2719}

$$\frac{2 \cos(e+fx) (d \csc(e+fx))^{5/2}}{5d^4 f} - \frac{6 \cos(e+fx) \sqrt{d \csc(e+fx)}}{5d^2 f} - \frac{6E\left(\frac{1}{2}(e+fx - \frac{\pi}{2}) \mid 2\right)}{5df \sqrt{\sin(e+fx)} \sqrt{d \csc(e+fx)}}$$

Antiderivative was successfully verified.

[In] Int[Csc[e + f*x]^5/(d*Csc[e + f*x])^(3/2),x]

[Out] $(-6 * \text{Cos}[e + f * x] * \text{Sqrt}[d * \text{Csc}[e + f * x]]) / (5 * d^2 * f) - (2 * \text{Cos}[e + f * x] * (d * \text{Csc}[e + f * x])^{(5/2)}) / (5 * d^4 * f) - (6 * \text{EllipticE}[(e - \text{Pi}/2 + f * x) / 2, 2]) / (5 * d * f * \text{Sqrt}[d * \text{Csc}[e + f * x]] * \text{Sqrt}[\text{Sin}[e + f * x]])$

Rule 16

Int[(u_.)*(v_)^(m_.)*((b_.)*(v_))^(n_), x_Symbol] := Dist[1/b^m, Int[u*(b*v)^(m+n), x], x] /; FreeQ[{b, n}, x] && IntegerQ[m]

Rule 2719

Int[Sqrt[sin[(c_.) + (d_.)*(x_)]], x_Symbol] := Simp[(2/d)*EllipticE[(1/2)*(c - Pi/2 + d*x), 2], x] /; FreeQ[{c, d}, x]

Rule 3853

Int[(csc[(c_.) + (d_.)*(x_)])*(b_.)^(n_), x_Symbol] := Simp[(-b)*Cos[c + d*x]*(b*Csc[c + d*x])^(n-1)/(d*(n-1)), x] + Dist[b^2*((n-2)/(n-1)), Int[(b*Csc[c + d*x])^(n-2), x], x] /; FreeQ[{b, c, d}, x] && GtQ[n, 1] && IntegerQ[2*n]

Rule 3856

```
Int[(csc[(c_.) + (d_.)*(x_)]*(b_.))^n_], x_Symbol] := Dist[(b*Csc[c + d*x]
)^n*Sin[c + d*x]^n, Int[1/Sin[c + d*x]^n, x], x] /; FreeQ[{b, c, d}, x] &&
EqQ[n^2, 1/4]
```

Rubi steps

$$\begin{aligned} \int \frac{\csc^5(e+fx)}{(d \csc(e+fx))^{3/2}} dx &= \frac{\int (d \csc(e+fx))^{7/2} dx}{d^5} \\ &= -\frac{2 \cos(e+fx)(d \csc(e+fx))^{5/2}}{5d^4 f} + \frac{3 \int (d \csc(e+fx))^{3/2} dx}{5d^3} \\ &= -\frac{6 \cos(e+fx) \sqrt{d \csc(e+fx)}}{5d^2 f} - \frac{2 \cos(e+fx)(d \csc(e+fx))^{5/2}}{5d^4 f} - \frac{3 \int \sqrt{d \csc(e+fx)} dx}{5d^3} \\ &= -\frac{6 \cos(e+fx) \sqrt{d \csc(e+fx)}}{5d^2 f} - \frac{2 \cos(e+fx)(d \csc(e+fx))^{5/2}}{5d^4 f} - \frac{3 \int \sqrt{d \csc(e+fx)} dx}{5d \sqrt{d \csc(e+fx)}} \\ &= -\frac{6 \cos(e+fx) \sqrt{d \csc(e+fx)}}{5d^2 f} - \frac{2 \cos(e+fx)(d \csc(e+fx))^{5/2}}{5d^4 f} - \frac{6E\left(\frac{1}{2}\right)}{5df \sqrt{d \csc(e+fx)}} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 73, normalized size = 0.70

$$\frac{\csc^4(e+fx) \left(-7 \cos(e+fx) + 3 \cos(3(e+fx)) + 12E\left(\frac{1}{4}(-2e+\pi-2fx) \mid 2\right) \sin^{\frac{5}{2}}(e+fx) \right)}{10f(d \csc(e+fx))^{3/2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[Csc[e + f*x]^5/(d*Csc[e + f*x])^(3/2), x]
```

```
[Out] (Csc[e + f*x]^4*(-7*Cos[e + f*x] + 3*Cos[3*(e + f*x)] + 12*EllipticE[(-2*e + Pi - 2*f*x)/4, 2]*Sin[e + f*x]^(5/2)))/(10*f*(d*Csc[e + f*x])^(3/2))
```

Maple [C] Result contains complex when optimal does not.

time = 0.12, size = 1054, normalized size = 10.04

method	result	size
default	Expression too large to display	1054

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(csc(f*x+e)^5/(d*csc(f*x+e))^(3/2), x, method=_RETURNVERBOSE)
```

```
[Out] -1/5/f*(6*((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))^(1/2)*(-(I*cos(f*x+e)-I-sin(f*x+e))/sin(f*x+e))^(1/2)*EllipticE((I*cos(f*x+e)-I+sin(f*x+e))/sin(f*x+e))
```

$$\begin{aligned} & x+e))^{(1/2)}, 1/2*2^{(1/2)})*\cos(f*x+e)^3*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & -3*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x \\ & +e))/\sin(f*x+e))^{(1/2)}*\text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)}) \\ & *\cos(f*x+e)^3*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}+6*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *(-I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*\text{EllipticE}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/ \\ & 2*2^{(1/2)})*\cos(f*x+e)^2*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}-3*((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *(-I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*\text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)}) \\ & *\cos(f*x+e)^2*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)}-6*\cos(f*x+e)*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *\text{EllipticE}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)})+3*\cos(f*x+e)*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *\text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)})-6*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *\text{EllipticE}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)})+3*(-I*(-1+\cos(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}*(-(I*\cos(f*x+e)-I-\sin(f*x+e))/\sin(f*x+e))^{(1/2)} \\ & *\text{EllipticF}(((I*\cos(f*x+e)-I+\sin(f*x+e))/\sin(f*x+e))^{(1/2)}, 1/2*2^{(1/2)})-3*\cos(f*x+e)^2*2^{(1/2)}+\cos(f*x+e)*2^{(1/2)}+3*2^{(1/2)}/\sin(f*x+e)^{4/(d/\sin(f*x+e))^{(3/2)}*2^{(1/2)}} \end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5/(d*csc(f*x+e))^(3/2),x, algorithm="maxima")**[Out]** integrate(csc(f*x + e)^5/(d*csc(f*x + e))^(3/2), x)**Fricas [C]** Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.09, size = 144, normalized size = 1.37

$$\frac{3(\cos(fx+e)^2-1)\sqrt{2d}\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(fx+e)+i\sin(fx+e)))+3(\cos(fx+e)^2-1)\sqrt{-2id}\text{weierstrassZeta}(4,0,\text{weierstrassPInverse}(4,0,\cos(fx+e)-i\sin(fx+e)))+2(3\cos(fx+e)^3-4\cos(fx+e))\sqrt{\frac{d}{\sin(fx+e)}}}{5(d^2f\cos(fx+e)^2-d^2f)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5/(d*csc(f*x+e))^(3/2),x, algorithm="fricas")

[Out] $-1/5*(3*(\cos(f*x + e)^2 - 1)*\text{sqrt}(2*I*d)*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(f*x + e) + I*\sin(f*x + e))) + 3*(\cos(f*x + e)^2 - 1)*\text{sqrt}(-2*I*d)*\text{weierstrassZeta}(4, 0, \text{weierstrassPInverse}(4, 0, \cos(f*x + e) - I*s$

$\ln(fx + e)) + 2*(3*\cos(fx + e)^3 - 4*\cos(fx + e))*\sqrt{d/\sin(fx + e))}$
 $/(d^2*f*\cos(fx + e)^2 - d^2*f)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\csc^5(e + fx)}{(d \csc(e + fx))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)**5/(d*csc(f*x+e))**(3/2),x)

[Out] Integral(csc(e + f*x)**5/(d*csc(e + f*x))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(csc(f*x+e)^5/(d*csc(f*x+e))^(3/2),x, algorithm="giac")

[Out] integrate(csc(f*x + e)^5/(d*csc(f*x + e))^(3/2), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{\sin(e + fx)^5 \left(\frac{d}{\sin(e+fx)}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(sin(e + f*x)^5*(d/sin(e + f*x))^(3/2)),x)

[Out] int(1/(sin(e + f*x)^5*(d/sin(e + f*x))^(3/2)), x)

3.538 $\int (b \csc(e + fx))^n (a \sin(e + fx))^m dx$

Optimal. Leaf size=87

$$\frac{\cos(e + fx)(b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m - n); \frac{1}{2}(3 + m - n); \sin^2(e + fx)\right) (a \sin(e + fx))^{1+m}}{af(1 + m - n)\sqrt{\cos^2(e + fx)}}$$

[Out] `cos(f*x+e)*(b*csc(f*x+e))^n*hypergeom([1/2, 1/2+1/2*m-1/2*n], [3/2+1/2*m-1/2*n], sin(f*x+e)^2)*(a*sin(f*x+e))^(1+m)/a/f/(1+m-n)/(cos(f*x+e)^2)^(1/2)`

Rubi [A]

time = 0.05, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {2668, 2722}

$$\frac{\cos(e + fx)(a \sin(e + fx))^{m+1}(b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(m - n + 1); \frac{1}{2}(m - n + 3); \sin^2(e + fx)\right)}{af(m - n + 1)\sqrt{\cos^2(e + fx)}}$$

Antiderivative was successfully verified.

[In] `Int[(b*Csc[e + f*x])^n*(a*Sin[e + f*x])^m,x]`

[Out] `(Cos[e + f*x]*(b*Csc[e + f*x])^n*Hypergeometric2F1[1/2, (1 + m - n)/2, (3 + m - n)/2, Sin[e + f*x]^2]*(a*Sin[e + f*x])^(1 + m))/(a*f*(1 + m - n)*Sqrt[Cos[e + f*x]^2])`

Rule 2668

`Int[(csc[(e_) + (f_)*(x_)]*(b_.))^n*((a_)*sin[(e_) + (f_)*(x_)])^m, x_Symbol] :> Dist[(a*b)^IntPart[n]*(a*Sin[e + f*x])^FracPart[n]*(b*Csc[e + f*x])^FracPart[n], Int[(a*Sin[e + f*x])^(m - n), x], x] /; FreeQ[{a, b, e, f, m, n}, x] && !IntegerQ[m] && !IntegerQ[n]`

Rule 2722

`Int[((b_)*sin[(c_) + (d_)*(x_)])^n, x_Symbol] :> Simp[Cos[c + d*x]*((b*Sin[c + d*x])^(n + 1)/(b*d*(n + 1)*Sqrt[Cos[c + d*x]^2]))*Hypergeometric2F1[1/2, (n + 1)/2, (n + 3)/2, Sin[c + d*x]^2], x] /; FreeQ[{b, c, d, n}, x] && !IntegerQ[2*n]`

Rubi steps

$$\begin{aligned} \int (b \csc(e + fx))^n (a \sin(e + fx))^m dx &= ((b \csc(e + fx))^n (a \sin(e + fx))^n) \int (a \sin(e + fx))^{m-n} dx \\ &= \frac{\cos(e + fx)(b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2}, \frac{1}{2}(1 + m - n); \frac{1}{2}(3 + m - n); \sin^2(e + fx)\right) (a \sin(e + fx))^{1+m}}{af(1 + m - n)\sqrt{\cos^2(e + fx)}} \end{aligned}$$

Mathematica [A]

time = 7.15, size = 102, normalized size = 1.17

$$\frac{2(b \csc(e + fx))^n {}_2F_1\left(\frac{1}{2}(1 + m - n), 1 + m - n; \frac{1}{2}(3 + m - n); -\tan^2\left(\frac{1}{2}(e + fx)\right)\right) \sec^2\left(\frac{1}{2}(e + fx)\right)^{m-n} (a \sin(e + fx))^m \tan\left(\frac{1}{2}(e + fx)\right)}{f(1 + m - n)}$$

Antiderivative was successfully verified.

[In] Integrate[(b*Csc[e + f*x])^n*(a*Sin[e + f*x])^m,x]

[Out] (2*(b*Csc[e + f*x])^n*Hypergeometric2F1[(1 + m - n)/2, 1 + m - n, (3 + m - n)/2, -Tan[(e + f*x)/2]^2]*(Sec[(e + f*x)/2]^2)^(m - n)*(a*Sin[e + f*x])^m*Tan[(e + f*x)/2])/(f*(1 + m - n))

Maple [F]

time = 0.07, size = 0, normalized size = 0.00

$$\int (b \csc(fx + e))^n (a \sin(fx + e))^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((b*csc(f*x+e))^n*(a*sin(f*x+e))^m,x)

[Out] int((b*csc(f*x+e))^n*(a*sin(f*x+e))^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="maxima")

[Out] integrate((b*csc(f*x + e))^n*(a*sin(f*x + e))^m, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="fricas")

[Out] integral((b*csc(f*x + e))^n*(a*sin(f*x + e))^m, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a \sin(e + fx))^m (b \csc(e + fx))^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))**n*(a*sin(f*x+e))**m,x)

[Out] Integral((a*sin(e + f*x))**m*(b*csc(e + f*x))**n, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((b*csc(f*x+e))^n*(a*sin(f*x+e))^m,x, algorithm="giac")

[Out] integrate((b*csc(f*x + e))^n*(a*sin(f*x + e))^m, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a \sin(e + f x))^m \left(\frac{b}{\sin(e + f x)} \right)^n dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*sin(e + f*x))^m*(b/sin(e + f*x))^n,x)

[Out] int((a*sin(e + f*x))^m*(b/sin(e + f*x))^n, x)

Chapter 4

Appendix

Local contents

4.1	Download section	2114
4.2	Listing of Grading functions	2114

4.1 Download section

The following zip files contain the raw integrals used in this test.

Mathematica format Mathematica_syntax.zip

Maple and Mupad format Maple_syntax.zip

Sympy format SYMPY_syntax.zip

Sage math format SAGE_syntax.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*           is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*           antiderivative*)
(* "A" if result can be considered optimal*)
```

```

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A","none"}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","none"}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)

```

(*9 = unknown function*)

```

ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
            If[ElementaryFunctionQ[Head[expn]],
              Max[3, ExpnType[expn[[1]]]],
              If[SpecialFunctionQ[Head[expn]],
                Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
                If[HypergeometricFunctionQ[Head[expn]],
                  Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
                  If[AppellFunctionQ[Head[expn]],
                    Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
                    If[Head[expn]===RootSum,
                      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
                      If[Head[expn]===Integrate || Head[expn]===Int,
                        Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
                        9]]]]]]]]]]]]

```

```

ElementaryFunctionQ[func_] :=
  MemberQ[{
    Exp, Log,
    Sin, Cos, Tan, Cot, Sec, Csc,
    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
    Sinh, Cosh, Tanh, Coth, Sech, Cschn,
    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCschn
  }, func]

```

```

SpecialFunctionQ[func_] :=
  MemberQ[{
    Erf, Erfc, Erfi,
    FresnelS, FresnelC,

```

```

ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]

```

4.2.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

  leaf_count_result:=leafcount(result);
  #do NOT call ExpnType() if leaf size is too large. Recursion problem
  if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
  fi;

  leaf_count_optimal := leafcount(optimal);
  ExpnType_result := ExpnType(result);
  ExpnType_optimal := ExpnType(optimal);

```

```

    if debug then
        print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
    fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#   is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
#   antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A","";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of r
                    convert(leaf_count_result,string)," vs. $2 (" ,
                    convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_co
            end if
        else #result contains complex but optimal is not
            if debug then
                print("result contains complex but optimal is not");
            fi;
            return "C","Result contains complex when optimal does not.";
        fi;
    else # result do not contain complex

```

```

    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A","";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string),")=",convert(2*leaf_cou

    fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                  convert(ExpnType_result,string)," vs. order ",
                  convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function

```

```

# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`) or type(expn,'*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,

```



```

    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

4.2.4 SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or instan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```



```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```