

Computer algebra independent integration tests

Summer 2022 edition

5-Inverse-trig-functions/5.3-Inverse-tangent/149-5.3.3-d+e-x-^m-
a+b-arctan-c-x^n-^p

Nasser M. Abbasi

September 27, 2022 Compiled on September 27, 2022 at 4:52am

Contents

1	Introduction	3
2	detailed summary tables of results	19
3	Listing of integrals	33
4	Appendix	211

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	list of integrals that has no closed form antiderivative	11
1.5	List of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	13
1.7	Timing	13
1.8	Verification	14
1.9	Important notes about some of the results	14
1.10	Design of the test system	17

This report gives the result of running the computer algebra independent integration test. The download section in the appendix contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [31]. This is test number [149].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.1 (June 29, 2022) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.0.1 on windows 10.
3. Maple 2022.1 (June 1, 2022) on windows 10.
4. Maxima 5.46 (April 13, 2022) using Lisp SBCL 2.1.11.debian on Linux via sagemath 9.6.
5. Fricas 1.3.8 (June 21, 2022) based on sbcl 2.1.11.debian on Linux via sagemath 9.6.
6. Giac/Xcas 1.9.0-13 (July 3, 2022) on Linux via sagemath 9.6.
7. Sympy 1.10.1 (March 20, 2022) Using Python 3.10.4 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly from Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or Hypergeometric2F1 functions. RootSum and RootOf are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (31)	0.00 (0)
Maple	96.77 (30)	3.23 (1)
Mathematica	83.87 (26)	16.13 (5)
Mupad	45.16 (14)	54.84 (17)
Maxima	45.16 (14)	54.84 (17)
Fricas	35.48 (11)	64.52 (20)
Sympy	35.48 (11)	64.52 (20)
Giac	19.35 (6)	80.65 (25)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	77.42	3.23	3.23	16.13
Maple	48.39	19.35	29.03	3.23
Maxima	45.16	0.00	0.00	54.84
Fricas	22.58	12.90	0.00	64.52
Giac	19.35	0.00	0.00	80.65
Sympy	16.13	6.45	12.90	64.52
Mupad	N/A	38.71	0.00	54.84

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sageMath and the CAS (applicable only to FriCAS, Maxima and

Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	5	80.00 %	20.00 %	0.00 %
Maple	1	100.00 %	0.00 %	0.00 %
Fricas	20	80.00 %	15.00 %	5.00 %
Giac	25	92.00 %	8.00 %	0.00 %
Maxima	17	82.35 %	11.76 %	5.88 %
Sympy	20	60.00 %	40.00 %	0.00 %
Mupad	17	100.00 %	0.00 %	0.00 %

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.


Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.35	351.45	0.94	270.00	1.00
Mathematica	10.39	487.81	1.17	276.00	1.11
Maple	2.49	2341.33	4.45	351.50	1.37
Maxima	0.45	217.07	1.02	220.00	1.02
Fricas	1.14	625.00	2.96	190.00	1.32
Sympy	6.66	1411.64	7.88	345.00	1.82
Giac	2.10	176.67	0.68	217.00	0.92
Mupad	1.34	460.57	1.62	238.00	1.43

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used from the above table.



1.4 list of integrals that has no closed form antiderivative

{26, 27}

1.5 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {25}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each `integrate` call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime'
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the `integrate` command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for Rubi and Mathematica.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
```

```
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError.`

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount = 1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.


The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand, the_variable)
```

Which gives $\sin(x)^{2/2}$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified.

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,...}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	20
2.2	Detailed conclusion table per each integral for all CAS systems	23
2.3	Detailed conclusion table specific for Rubi results	30

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	21
2.1.2	Mathematica	21
2.1.3	Maple	21
2.1.4	Maxima	21
2.1.5	FriCAS	21
2.1.6	Sympy	22
2.1.7	Giac	22
2.1.8	Mupad	22

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 13, 14, 15, 16, 17, 21, 22, 24, 26, 27, 28, 29, 31 }

B grade: { 25 }

C grade: { 23 }

F grade: { 12, 18, 19, 20, 30 }

2.1.3 Maple

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 21, 22, 24, 26, 27, 29, 31 }

B grade: { 9, 10, 11, 13, 14, 28 }

C grade: { 12, 15, 16, 17, 18, 19, 20, 23, 30 }

F grade: { 25 }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 6, 7, 8, 21, 22, 24, 26, 28, 29, 31 }

B grade: { }

C grade: { }

F grade: { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 25, 27, 30 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 6, 26, 27 }

B grade: { 7, 8, 21, 22 }

C grade: { }

F grade: { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 24, 25, 28, 29, 30, 31 }

2.1.6 Sympy

A grade: { 3, 4, 21, 28, 29 }

B grade: { 1, 2 }

C grade: { 6, 7, 8, 22 }

F grade: { 5, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 24, 25, 26, 27, 30, 31 }

2.1.7 Giac

A grade: { 21, 22, 26, 27, 28, 29 }

B grade: { }

C grade: { }

F grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 24, 25, 30, 31 }

2.1.8 Mupad

A grade: { 26, 27 }

B grade: { 1, 2, 3, 4, 6, 7, 21, 22, 24, 28, 29, 31 }

C grade: { }

F grade: { 5, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 23, 25, 30 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column N.S. in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, **Mathematica** was abbreviated to MMA.

	Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
MMA	grade	A	A	A	A	A	A	B	F	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	184	184	255	241	248	257	345	0	273
	N.S.	1	1.00	1.39	1.31	1.35	1.40	1.88	0.00	1.48
	time (sec)	N/A	0.100	0.323	0.251	0.467	1.383	0.441	0.000	1.049

	Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
MMA	grade	A	A	A	A	A	A	B	F	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	144	144	218	179	184	190	262	0	197
	N.S.	1	1.00	1.51	1.24	1.28	1.32	1.82	0.00	1.37
	time (sec)	N/A	0.085	0.300	0.116	0.486	1.494	0.348	0.000	0.802

	Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
MMA	grade	A	A	A	A	A	A	A	F	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	103	103	163	123	126	132	160	0	127
	N.S.	1	1.00	1.58	1.19	1.22	1.28	1.55	0.00	1.23
	time (sec)	N/A	0.063	0.225	0.112	0.474	1.540	0.269	0.000	0.353

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	77	78	73	73	87	0	67
N.S.	1	1.00	1.01	1.03	0.96	0.96	1.14	0.00	0.88
time (sec)	N/A	0.044	0.007	0.045	0.478	1.187	0.162	0.000	0.410

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	138	138	245	178	0	0	0	0	-1
N.S.	1	1.00	1.78	1.29	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.143	0.056	0.000	0.000	0.000	0.000	0.000

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	C	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	111	130	106	114	682	0	112
N.S.	1	1.00	1.13	1.33	1.08	1.16	6.96	0.00	1.14
time (sec)	N/A	0.037	0.133	0.102	0.471	1.136	1.485	0.000	3.460

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	C	F	B	
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	146	146	192	188	206	299	2866	0	591
N.S.	1	1.00	1.32	1.29	1.41	2.05	19.63	0.00	4.05
time (sec)	N/A	0.086	0.220	0.143	0.480	1.358	3.419	0.000	4.428

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	C	F	F	
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	206	206	254	286	354	620	9202	0	-1
N.S.	1	1.00	1.23	1.39	1.72	3.01	44.67	0.00	-0.00
time (sec)	N/A	0.130	0.427	0.196	0.491	1.991	7.258	0.000	0.000

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	376	376	472	880	0	0	0	0	-1
N.S.	1	1.00	1.26	2.34	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.394	0.524	0.356	0.000	0.000	0.000	0.000	0.000

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	270	270	312	700	0	0	0	0	-1
N.S.	1	1.00	1.16	2.59	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.264	0.352	0.232	0.000	0.000	0.000	0.000	0.000

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	172	340	0	0	0	0	-1
N.S.	1	1.00	1.01	1.99	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.204	0.137	0.124	0.000	0.000	0.000	0.000	0.000

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	C	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	223	223	0	1324	0	0	0	0	-1
N.S.	1	1.00	0.00	5.94	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.033	78.300	2.899	0.000	0.000	0.000	0.000	0.000

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	341	341	300	740	0	0	0	0	-1
N.S.	1	1.00	0.88	2.17	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.258	1.956	2.340	0.000	0.000	0.000	0.000	0.000

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-1)	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	496	496	479	965	0	0	0	0	-1
N.S.	1	1.00	0.97	1.95	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.393	4.051	5.139	0.000	0.000	0.000	0.000	0.000

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	652	652	855	3456	0	0	0	0	-1
N.S.	1	1.00	1.31	5.30	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.822	1.211	22.015	0.000	0.000	0.000	0.000	0.000

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	411	411	621	2921	0	0	0	0	-1
N.S.	1	1.00	1.51	7.11	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.524	0.718	7.533	0.000	0.000	0.000	0.000	0.000

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	264	264	342	7195	0	0	0	0	-1
N.S.	1	1.00	1.30	27.25	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.384	0.321	1.451	0.000	0.000	0.000	0.000	0.000

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	C	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	320	320	0	2664	0	0	0	0	-1
N.S.	1	1.00	0.00	8.32	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.039	137.065	2.777	0.000	0.000	0.000	0.000	0.000

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	C	F	F	F	F(-1)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	499	499	0	3050	0	0	0	0	-1
N.S.	1	1.00	0.00	6.11	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.364	85.555	2.879	0.000	0.000	0.000	0.000	0.000

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	C	F(-1)	F	F(-1)	F(-1)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	936	936	0	41275	0	0	0	0	-1
N.S.	1	1.00	0.00	44.10	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.770	45.789	14.874	0.000	0.000	0.000	0.000	0.000

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	250	250	252	363	323	4652	403	308	419
N.S.	1	1.00	1.01	1.45	1.29	18.61	1.61	1.23	1.68
time (sec)	N/A	0.197	2.008	0.437	0.479	1.435	11.146	1.149	3.366

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	C	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	192	192	153	168	170	538	1266	184	203
N.S.	1	1.00	0.80	0.88	0.89	2.80	6.59	0.96	1.06
time (sec)	N/A	0.129	0.070	0.092	0.464	1.051	7.522	0.594	2.534

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	C	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	501	501	326	138	0	0	0	0	-1
N.S.	1	1.00	0.65	0.28	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.649	22.379	0.119	0.000	0.000	0.000	0.000	0.000

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	F(-1)	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	328	328	321	433	286	0	0	0	883
N.S.	1	1.00	0.98	1.32	0.87	0.00	0.00	0.00	2.69
time (sec)	N/A	0.349	0.473	0.182	0.483	0.000	0.000	0.000	0.661

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	B	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	1325	1325	5420	0	0	0	0	0	-1
N.S.	1	1.00	4.09	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	1.624	30.640	0.095	0.000	0.000	0.000	0.000	0.000

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	A	A	F(-1)	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	0	0	0	0	0	0	0	-1
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.04
time (sec)	N/A	0.019	47.024	0.027	0.000	0.000	0.000	0.000	0.000

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	N/A	A	A	A	F(-2)	A	F(-1)	A	A
verified	N/A	N/A	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	23	0	0	0	0	0	0	0	-1
N.S.	1	0.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.04
time (sec)	N/A	0.019	44.473	0.221	0.000	0.000	0.000	0.000	0.000

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	F(-1)	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	315	315	297	518	280	0	151	318	988
N.S.	1	1.00	0.94	1.64	0.89	0.00	0.48	1.01	3.14
time (sec)	N/A	0.496	102.002	9.230	0.567	0.000	28.537	9.908	0.517

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	F(-2)	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	285	285	310	315	234	0	104	250	485
N.S.	1	1.00	1.09	1.11	0.82	0.00	0.36	0.88	1.70
time (sec)	N/A	0.404	0.064	0.246	0.474	0.000	12.693	0.952	0.436

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F(-1)	C	F	F	F(-1)	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	739	739	0	172	0	0	0	0	-1
N.S.	1	1.00	0.00	0.23	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.930	180.003	0.108	0.000	0.000	0.000	0.000	0.000

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	F(-1)	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	906	906	536	1220	449	0	0	0	2105
N.S.	1	1.00	0.59	1.35	0.50	0.00	0.00	0.00	2.32
time (sec)	N/A	0.936	10.004	0.398	0.490	0.000	0.000	0.000	0.776

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [9] had the largest ratio of [18]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	5	1.00	16	0.312
2	A	6	5	1.00	16	0.312
3	A	6	5	1.00	16	0.312
4	A	6	5	1.00	14	0.357
5	A	4	4	1.00	16	0.250
6	A	6	6	1.00	16	0.375
7	A	7	6	1.00	16	0.375
8	A	7	6	1.00	16	0.375
9	A	19	14	1.00	18	0.778
10	A	15	12	1.00	18	0.667
11	A	12	9	1.00	16	0.562
12	A	1	1	1.00	18	0.056
13	A	13	9	1.00	18	0.500
14	A	19	15	1.00	18	0.833
15	A	29	15	1.00	18	0.833
16	A	20	13	1.00	18	0.722
17	A	14	10	1.00	16	0.625
18	A	1	1	1.00	18	0.056
19	A	10	8	1.00	18	0.444
20	A	23	12	1.00	18	0.667
21	A	17	13	1.00	18	0.722
22	A	16	12	1.00	16	0.750
23	A	19	8	1.00	18	0.444
24	A	18	13	1.00	18	0.722
25	A	77	26	1.00	18	1.444

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	<u>number of rules</u> <u>integrand leaf size</u>
26	A	0	0	0.00	0	0.000
27	A	0	0	0.00	0	0.000
28	A	24	15	1.00	18	0.833
29	A	22	12	1.00	16	0.750
30	A	25	8	1.00	18	0.444
31	A	34	15	1.00	18	0.833

Chapter 3

Listing of integrals

Local contents

3.1	$\int (d + ex)^4(a + b\text{ArcTan}(cx)) dx$	34
3.2	$\int (d + ex)^3(a + b\text{ArcTan}(cx)) dx$	38
3.3	$\int (d + ex)^2(a + b\text{ArcTan}(cx)) dx$	42
3.4	$\int (d + ex)(a + b\text{ArcTan}(cx)) dx$	46
3.5	$\int \frac{a+b\text{ArcTan}(cx)}{d+ex} dx$	50
3.6	$\int \frac{a+b\text{ArcTan}(cx)}{(d+ex)^2} dx$	54
3.7	$\int \frac{a+b\text{ArcTan}(cx)}{(d+ex)^3} dx$	59
3.8	$\int \frac{a+b\text{ArcTan}(cx)}{(d+ex)^4} dx$	65
3.9	$\int (d + ex)^3(a + b\text{ArcTan}(cx))^2 dx$	71
3.10	$\int (d + ex)^2(a + b\text{ArcTan}(cx))^2 dx$	78
3.11	$\int (d + ex)(a + b\text{ArcTan}(cx))^2 dx$	84
3.12	$\int \frac{(a+b\text{ArcTan}(cx))^2}{d+ex} dx$	89
3.13	$\int \frac{(a+b\text{ArcTan}(cx))^2}{(d+ex)^2} dx$	92
3.14	$\int \frac{(a+b\text{ArcTan}(cx))^2}{(d+ex)^3} dx$	97
3.15	$\int (d + ex)^3(a + b\text{ArcTan}(cx))^3 dx$	103
3.16	$\int (d + ex)^2(a + b\text{ArcTan}(cx))^3 dx$	112
3.17	$\int (d + ex)(a + b\text{ArcTan}(cx))^3 dx$	120
3.18	$\int \frac{(a+b\text{ArcTan}(cx))^3}{d+ex} dx$	125
3.19	$\int \frac{(a+b\text{ArcTan}(cx))^3}{(d+ex)^2} dx$	129
3.20	$\int \frac{(a+b\text{ArcTan}(cx))^3}{(d+ex)^3} dx$	135
3.21	$\int (d + ex)^2(a + b\text{ArcTan}(cx^2)) dx$	141
3.22	$\int (d + ex)(a + b\text{ArcTan}(cx^2)) dx$	149
3.23	$\int \frac{a+b\text{ArcTan}(cx^2)}{d+ex} dx$	155

3.24	$\int \frac{a+b\text{ArcTan}(cx^2)}{(d+ex)^2} dx$	160
3.25	$\int (d+ex)(a+b\text{ArcTan}(cx^2))^2 dx$	167
3.26	$\int \frac{(a+b\text{ArcTan}(cx^2))^2}{d+ex} dx$	176
3.27	$\int \frac{(a+b\text{ArcTan}(cx^2))^2}{(d+ex)^2} dx$	179
3.28	$\int (d+ex)^2(a+b\text{ArcTan}(cx^3)) dx$	183
3.29	$\int (d+ex)(a+b\text{ArcTan}(cx^3)) dx$	190
3.30	$\int \frac{a+b\text{ArcTan}(cx^3)}{d+ex} dx$	197
3.31	$\int \frac{a+b\text{ArcTan}(cx^3)}{(d+ex)^2} dx$	202

$$3.1 \quad \int (d + ex)^4 (a + b \text{ArcTan}(cx)) dx$$

Optimal. Leaf size=184

$$\frac{-bde(2c^2d^2 - e^2)x}{c^3} - \frac{be^2(10c^2d^2 - e^2)x^2}{10c^3} - \frac{bde^3x^3}{3c} - \frac{be^4x^4}{20c} - \frac{bd(c^4d^4 - 10c^2d^2e^2 + 5e^4)\text{ArcTan}(cx)}{5c^4e} + \frac{(d + ex)^5(a + b\text{ArcTan}(cx))}{5e}$$

$$[\text{Out}] \quad -b*d*e*(2*c^2*d^2 - e^2)*x/c^3 - 1/10*b*e^2*(10*c^2*d^2 - e^2)*x^2/c^3 - 1/3*b*d*e^3*x^3/c^3 - 1/20*b*e^4*x^4/c^3 - 1/5*b*d*(c^4*d^4 - 10*c^2*d^2 - e^2 + 5*e^4)*\arctan(cx)/c^4/e + 1/5*(e*x + d)^5*(a + b*\arctan(cx))/e - 1/10*b*(5*c^4*d^4 - 10*c^2*d^2 - e^2 + 5*e^4)*\ln(c^2*x^2 + 1)/c^5$$

Rubi [A]

time = 0.10, antiderivative size = 184, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.312, Rules used = {4972, 716, 649, 209, 266}

$$\frac{(d + ex)^5(a + b\text{ArcTan}(cx))}{5e} - \frac{bd\text{ArcTan}(cx)(c^4d^4 - 10c^2d^2e^2 + 5e^4)}{5c^4e} - \frac{be^2x^2(10c^2d^2 - e^2)}{10c^3} - \frac{bdex(2c^2d^2 - e^2)}{c^3} - \frac{b(5c^4d^4 - 10c^2d^2e^2 + e^4)\log(c^2x^2 + 1)}{10c^5} - \frac{bde^3x^3}{3c} - \frac{be^4x^4}{20c}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d + e*x)^4*(a + b*\text{ArcTan}[c*x]), x]$

$$[\text{Out}] \quad -((b*d*e*(2*c^2*d^2 - e^2)*x)/c^3) - (b*e^2*(10*c^2*d^2 - e^2)*x^2)/(10*c^3) - (b*d*e^3*x^3)/(3*c) - (b*e^4*x^4)/(20*c) - (b*d*(c^4*d^4 - 10*c^2*d^2 - e^2 + 5*e^4)*\text{ArcTan}[c*x])/(5*c^4*e) + ((d + e*x)^5*(a + b*\text{ArcTan}[c*x]))/(5*e) - (b*(5*c^4*d^4 - 10*c^2*d^2 - e^2 + 5*e^4)*\text{Log}[1 + c^2*x^2])/(10*c^5)$$

Rule 209

$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*a \text{rcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \text{||} \text{GtQ}[b, 0])$

Rule 266

$\text{Int}[(x_)^{(m_)} / ((a_) + (b_)*(x_)^{(n_)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}[\{a, b, m, n\}, x] \&& \text{EqQ}[m, n - 1]$

Rule 649

$\text{Int}[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{Dist}[d, \text{Int}[1/(a + c*x^2), x], x] + \text{Dist}[e, \text{Int}[x/(a + c*x^2), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{!NiceSqrtQ}[-a]*c]$

Rule 716

$\text{Int}[((d_) + (e_)*(x_))^{(m_)} / ((a_) + (c_)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{Int}[\text{PolynomialDivide}[(d + e*x)^m, a + c*x^2, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{NeQ}[$

```
c*d^2 + a*e^2, 0] && IGtQ[m, 1] && (NeQ[d, 0] || GtQ[m, 2])
```

Rule 4972

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))*((d_) + (e_)*(x_))^(q_), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned} \int (d+ex)^4 (a+b\tan^{-1}(cx)) \, dx &= \frac{(d+ex)^5 (a+b\tan^{-1}(cx))}{5e} - \frac{(bc) \int \frac{(d+ex)^5}{1+c^2x^2} \, dx}{5e} \\ &= \frac{(d+ex)^5 (a+b\tan^{-1}(cx))}{5e} - \frac{(bc) \int \left(\frac{5de^2(2c^2d^2-e^2)}{c^4} + \frac{e^3(10c^2d^2-e^2)x}{c^4} + \frac{5de^5}{c^5} \right)}{5e} \\ &= -\frac{bde(2c^2d^2-e^2)x}{c^3} - \frac{be^2(10c^2d^2-e^2)x^2}{10c^3} - \frac{bde^3x^3}{3c} - \frac{be^4x^4}{20c} + \frac{(d+ex)}{5e} \\ &= -\frac{bde(2c^2d^2-e^2)x}{c^3} - \frac{be^2(10c^2d^2-e^2)x^2}{10c^3} - \frac{bde^3x^3}{3c} - \frac{be^4x^4}{20c} + \frac{(d+ex)}{5e} \\ &= -\frac{bde(2c^2d^2-e^2)x}{c^3} - \frac{be^2(10c^2d^2-e^2)x^2}{10c^3} - \frac{bde^3x^3}{3c} - \frac{be^4x^4}{20c} - \frac{bd(c^4d^4-e^4)}{20c} \end{aligned}$$

Mathematica [A]

time = 0.32, size = 255, normalized size = 1.39

$$\frac{(d+ex)^5(a+b\tan^{-1}(cx)) - \frac{b\left(c^2e^2x(-6e^2(10d+cx)+c^2(120d^3+60d^2cx+20d^2x^2+3c^3x^3))+6(-10c^2d^2e^2(\sqrt{-c^2}d+e)+e^4(5\sqrt{-c^2}d+e)+e^4d^4(\sqrt{-c^2}d+5e))\log(1-\sqrt{-c^2}x)-6(c^4d^4(\sqrt{-c^2}d-5e)-10c^2d^2(\sqrt{-c^2}d-e))^2+(5\sqrt{-c^2}d-e)e^4\log(1+\sqrt{-c^2}x)\right)}{12e^3}}{5e}$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)^4*(a + b*ArcTan[c*x]), x]`

[Out] $((d + e*x)^5*(a + b*ArcTan[c*x])) - (b*(c^2*e^2*x*(-6*e^2*(10*d + e*x) + c^2*(120*d^3 + 60*d^2*e*x + 20*d^2*e^2*x^2 + 3*e^3*x^3)) + 6*(-10*c^2*d^2*e^2*(Sqrt[-c^2]*d + e) + e^4*(5*Sqrt[-c^2]*d + e) + c^4*d^4*(Sqrt[-c^2]*d + 5*e))*Log[1 - Sqrt[-c^2]*x] - 6*(c^4*d^4*(Sqrt[-c^2]*d - 5*e) - 10*c^2*d^2*(Sqrt[-c^2]*d - e)*e^2 + (5*Sqrt[-c^2]*d - e)*e^4)*Log[1 + Sqrt[-c^2]*x]))/(12*c^5))/(5*e)$

Maple [A]

time = 0.25, size = 241, normalized size = 1.31

method	result
derivativedivides	$\frac{(ce^{}x^{}+cd^{})^5a}{5c^4e}+b\arctan(cx)d^4cx+2bce\arctan(cx)d^3x^2+2bc e^2\arctan(cx)d^2x^3+bc e^3\arctan(cx)dx^4+\frac{bc e^4\arctan(cx)x^5}{5}-2$
default	$\frac{(ce^{}x^{}+cd^{})^5a}{5c^4e}+b\arctan(cx)d^4cx+2bce\arctan(cx)d^3x^2+2bc e^2\arctan(cx)d^2x^3+bc e^3\arctan(cx)dx^4+\frac{bc e^4\arctan(cx)x^5}{5}-2$
risch	$\frac{ie^4b x^5 \ln(-icx+1)}{10} + \frac{ib d^5 \ln(c^2 x^2+1)}{20e} + ieb d^3 x^2 \ln(-icx+1) + \frac{ib d^4 x \ln(-icx+1)}{2} + \frac{x^5 e^4 a}{5} - \frac{i(ex+cd^5)}{5}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((e*x+d)^4*(a+b*arctan(c*x)),x,method=_RETURNVERBOSE)
[Out] 1/c*(1/5*(c*e*x+c*d)^5*a/c^4/e+b*b*arctan(c*x)*d^4*c*x+2*b*c*e*arctan(c*x)*d^3*x^2+2*b*c*e^2*arctan(c*x)*d^2*x^3+b*c*e^3*arctan(c*x)*d*x^4+1/5*b*c*e^4*a*rctan(c*x)*x^5-2*b*e*d^3*x-b*e^2*d^2*x^2-1/3*b*e^3*d*x^3-1/20*b*e^4*x^4+b/c^2*e^3*d*x+1/10*b/c^2*e^4*x^2-1/2*b*ln(c^2*x^2+1)*d^4+b/c^2*e^2*ln(c^2*x^2+1)*d^2-1/10*b/c^4*e^4*ln(c^2*x^2+1)+2*b/c*e*arctan(c*x)*d^3-b/c^3*e^3*arcta n(c*x)*d)
```

Maxima [A]

time = 0.47, size = 248, normalized size = 1.35

$$\frac{1}{5} a x^5 e^4 + a d x^4 e^3 + 2 a d^2 x^3 e^2 + 2 a d^3 x^2 e + a d^4 x + 2 \left(x^5 \arctan(cx) - c \left(\frac{x}{c^2} - \frac{\arctan(cx)}{c^3} \right) \right) b d^5 e + \left(2 c x \arctan(cx) - \log(x^2 + 1) \right) b d^4 e + \left(2 x^3 \arctan(cx) - c \left(\frac{x^2}{c^2} - \frac{\log(x^2 + 1)}{c^3} \right) \right) b d^3 e^2 + \frac{1}{3} \left(3 x^4 \arctan(cx) - c \left(\frac{c^2 x^3 - 3 x}{c^4} + \frac{3 \arctan(cx)}{c^5} \right) \right) b d^2 e^3 + \frac{1}{20} \left(4 x^5 \arctan(cx) - c \left(\frac{c^2 x^4 - 2 x^2}{c^5} + \frac{2 \log(x^2 + 1)}{c^6} \right) \right) b e^4$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^4*(a+b*arctan(c*x)),x, algorithm="maxima")
[Out] 1/5*a*x^5*e^4 + a*d*x^4*e^3 + 2*a*d^2*x^3*e^2 + 2*a*d^3*x^2*e + a*d^4*x + 2*(x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*b*d^3*e + 1/2*(2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*b*d^4/c + (2*x^3*arctan(c*x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c^4))*b*d^2*e^2 + 1/3*(3*x^4*arctan(c*x) - c*((c^2*x^3 - 3*x)/c^4 + 3*arctan(c*x)/c^5))*b*d*e^3 + 1/20*(4*x^5*arctan(c*x) - c*((c^2*x^4 - 2*x^2)/c^4 + 2*log(c^2*x^2 + 1)/c^6))*b*e^4
```

Fricas [A]

time = 1.38, size = 257, normalized size = 1.40

$$\frac{60 a^2 d^4 x + 12 (b c^2 x^2 e^4 + 10 b c^2 d^2 x^3 e^2 + 5 b c^2 d^4 x + 5 (b c^2 d x^4 - b c d^3 e) e^3 + 10 (b c^2 d^2 x^2 + b c^3 d^2 e) \arctan(cx) + 3 (4 a c^2 x^5 - b c^3 x^4 + 2 b c^2 x^2) e^4 + 20 (3 a c^2 d x^4 - b c^4 d x^3 + 3 b c^2 d x^2) e^3 + 60 (2 a c^2 d^2 x^3 - b c^4 d^2 x^2) e^2 + 120 (a c^2 d^3 x^2 - b c^4 d^3 x) e - 6 (5 b c^4 d^4 - 10 b c^2 d^2 e^2 + b e^4) \log(x^2 + 1)}{60 c^5}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^4*(a+b*arctan(c*x)),x, algorithm="fricas")
[Out] 1/60*(60*a*c^5*d^4*x + 12*(b*c^5*x^5*e^4 + 10*b*c^5*d^2*x^3*e^2 + 5*b*c^5*d*x^4 + 5*(b*c^5*d*x^4 - b*c^4*d*x^3)*e^3 + 10*(b*c^5*d^3*x^2 + b*c^4*d*x^3)*e)*arctan(c*x) + 3*(4*a*c^5*x^5 - b*c^6*x^4 + 2*b*c^5*x^2)*e^4 + 20*(3*a*c^5*d*x^4 - b*c^6*d*x^3 + 3*b*c^5*d*x^2)*e^3 + 60*(2*a*c^5*d^2*x^3 - b*c^6*d^2*x^2)*e^2 + 120*(a*c^5*d^3*x^2 - b*c^6*d^3*x)*e + 20*(5*b*c^6*d^4*x^4 - 10*b*c^4*d^2*x^2 + b*e^4)*log(x^2 + 1))
```

$$\begin{aligned}
& - b*c^4*d*x^3 + 3*b*c^2*d*x)*e^3 + 60*(2*a*c^5*d^2*x^3 - b*c^4*d^2*x^2)*e^2 \\
& + 120*(a*c^5*d^3*x^2 - b*c^4*d^3*x)*e - 6*(5*b*c^4*d^4 - 10*b*c^2*d^2*e^2 \\
& + b*e^4)*log(c^2*x^2 + 1))/c^5
\end{aligned}$$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 345 vs. $\frac{2(170)}{2} = 340$.

time = 0.44, size = 345, normalized size = 1.88

$$\left\{ \begin{array}{l} ad^4x + 2ad^3ex^2 + 2ad^2e^2x^3 + ade^3x^4 + \frac{ae^5x^5}{5} + bd^4x \operatorname{atan}(cx) + 2bd^3ex^2 \operatorname{atan}(cx) + 2bd^2e^2x^3 \operatorname{atan}(cx) + bde^3x^4 \operatorname{atan}(cx) + \frac{be^4x^5 \operatorname{atan}(cx)}{5} - \frac{bd^4 \log(x^2 + \frac{1}{c^2})}{2c} - \frac{2bd^3ex}{c} - \frac{bd^2e^2x^2}{c} - \frac{bd^3x^3}{3c} - \frac{bd^4x^4}{20c} + \frac{2bd^2e^3 \operatorname{atan}(cx)}{c^2} + \frac{bd^2x^2 \log(x^2 + \frac{1}{c^2})}{c^2} + \frac{bd^3x}{c^2} + \frac{bd^2x^2}{10c^2} - \frac{bd^3 \operatorname{atan}(cx)}{c^2} - \frac{bd^4 \log(x^2 + \frac{1}{c^2})}{10c^2} \\ \text{for } c \neq 0 \\ \text{otherwise} \end{array} \right.$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)**4*(a+b*atan(c*x)),x)`

[Out] `Piecewise((a*d**4*x + 2*a*d**3*e*x**2 + 2*a*d**2*e**2*x**3 + a*d*e**3*x**4 + a*e**4*x**5/5 + b*d**4*x*atan(c*x) + 2*b*d**3*e*x**2*atan(c*x) + 2*b*d**2*e**2*x**3*atan(c*x) + b*d**3*x**4*atan(c*x) + b*e**4*x**5*atan(c*x)/5 - b*d**4*log(x**2 + c*(-2))/(2*c) - 2*b*d**3*e*x/c - b*d**2*e**2*x**2/c - b*d**3*x**3/(3*c) - b*e**4*x**4/(20*c) + 2*b*d**3*e*atan(c*x)/c**2 + b*d**2*e**2*log(x**2 + c*(-2))/c**3 + b*d**3*x/c**3 + b*e**4*x**2/(10*c**3) - b*d**3*x*atan(c*x)/c**4 - b*e**4*log(x**2 + c*(-2))/(10*c**5), Ne(c, 0)), (a*(d**4*x + 2*d**3*e*x**2 + 2*d**2*e**2*x**3 + d*e**3*x**4 + e**4*x**5/5), True))`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^4*(a+b*arctan(c*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B]

time = 1.05, size = 273, normalized size = 1.48

$$\frac{a e^4 x^5}{5} + a d^4 x - \frac{b d^4 \ln(c^2 x^2 + 1)}{2 c} - \frac{b e^4 \ln(c^2 x^2 + 1)}{10 c^5} + 2 a d^2 e^2 x^3 - \frac{b e^4 x^4}{10 c^3} + b d^4 x \operatorname{atan}(cx) + 2 a d^3 e x^2 + a d e^3 x^4 + \frac{b e^4 x^5 \operatorname{atan}(cx)}{5} - \frac{2 b d^3 e x}{c} + \frac{b d^2 e^2 x}{c^2} + \frac{2 b d^3 e \operatorname{atan}(cx)}{c^2} - \frac{b d^2 e^3 \operatorname{atan}(cx)}{c^4} + 2 b d^2 e^2 x^2 \operatorname{atan}(cx) + b d^2 e^2 x^4 \operatorname{atan}(cx) - \frac{b d^2 e^2 x^3}{3 c} + 2 b d^2 e^2 x^5 \operatorname{atan}(cx) + \frac{b d^2 e^2 \ln(c^2 x^2 + 1)}{c^3} - \frac{b d^2 e^2 x^2}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))*(d + e*x)^4,x)`

[Out] `(a*e^4*x^5)/5 + a*d^4*x - (b*d^4*log(c^2*x^2 + 1))/(2*c) - (b*e^4*log(c^2*x^2 + 1))/(10*c^5) + 2*a*d^2*e^2*x^3 - (b*e^4*x^4)/(20*c) + (b*e^4*x^2)/(10*c^3) + b*d^4*x*atan(c*x) + 2*a*d^3*x^2 + a*d^2*e^3*x^4 + (b*e^4*x^5*atan(c*x))/5 - (2*b*d^3*x^2)/c + (b*d^2*e^3*x^3)/c^3 + (2*b*d^3*x^4*atan(c*x))/c^2 - (b*d^2*e^3*atan(c*x))/c^4 + 2*b*d^2*x^2*atan(c*x) + b*d^2*x^4*atan(c*x) - (b*d^2*x^3*atan(c*x))/3*c + 2*b*d^2*x^2*atan(c*x) + (b*d^2*x^4*atan(c*x))/c^3 - (b*d^2*x^2*atan(c*x))/c`

3.2 $\int (d + ex)^3(a + b \text{ArcTan}(cx)) dx$

Optimal. Leaf size=144

$$\frac{be(6c^2d^2 - e^2)x}{4c^3} - \frac{bde^2x^2}{2c} - \frac{be^3x^3}{12c} - \frac{b(c^4d^4 - 6c^2d^2e^2 + e^4)\text{ArcTan}(cx)}{4c^4e} + \frac{(d + ex)^4(a + b\text{ArcTan}(cx))}{4e} - \frac{bd(cx^2 + 1)^3}{4c^3}$$

[Out] $-1/4*b*e*(6*c^2*d^2 - e^2)*x/c^3 - 1/2*b*d*e^2*x^2/c - 1/12*b*e^3*x^3/c - 1/4*b*(c^4*d^4 - 6*c^2*d^2*e^2 + e^4)*\text{arctan}(c*x)/c^4/e + 1/4*(e*x + d)^4*(a + b*\text{arctan}(c*x))/e - 1/2*b*d*(c*d - e)*(c*d + e)*\ln(c^2*x^2 + 1)/c^3$

Rubi [A]

time = 0.09, antiderivative size = 144, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.312, Rules used = {4972, 716, 649, 209, 266}

$$\frac{(d + ex)^4(a + b\text{ArcTan}(cx))}{4e} - \frac{b\text{ArcTan}(cx)(c^4d^4 - 6c^2d^2e^2 + e^4)}{4c^4e} - \frac{bex(6c^2d^2 - e^2)}{4c^3} - \frac{bd(cd - e)(cd + e)\log(c^2x^2 + 1)}{2c^3} - \frac{bde^2x^2}{2c} - \frac{be^3x^3}{12c}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d + e*x)^3*(a + b*\text{ArcTan}[c*x]), x]$

[Out] $-1/4*(b*e*(6*c^2*d^2 - e^2)*x)/c^3 - (b*d*e^2*x^2)/(2*c) - (b*e^3*x^3)/(12*c) - (b*(c^4*d^4 - 6*c^2*d^2*e^2 + e^4)*\text{ArcTan}[c*x])/(4*c^4*e) + ((d + e*x)^4*(a + b*\text{ArcTan}[c*x]))/(4*e) - (b*d*(c*d - e)*(c*d + e)*\text{Log}[1 + c^2*x^2])/(2*c^3)$

Rule 209

$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_{\text{Symbol}}] := \text{Simp}[(1/(Rt[a, 2]*Rt[b, 2]))*a \text{rcTan}[Rt[b, 2]*(x/Rt[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \text{ || } \text{GtQ}[b, 0])$

Rule 266

$\text{Int}[(x_)^{(m_)} / ((a_) + (b_)*(x_)^{(n_)}), x_{\text{Symbol}}] := \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}[\{a, b, m, n\}, x] \&& \text{EqQ}[m, n - 1]$

Rule 649

$\text{Int}[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_{\text{Symbol}}] := \text{Dist}[d, \text{Int}[1/(a + c*x^2), x], x] + \text{Dist}[e, \text{Int}[x/(a + c*x^2), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{!NiceSqrtQ}[-a]*c]$

Rule 716

$\text{Int}[((d_) + (e_)*(x_))^{(m_)} / ((a_) + (c_)*(x_)^2), x_{\text{Symbol}}] := \text{Int}[\text{PolynomialDivide}[(d + e*x)^m, a + c*x^2, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{NeQ}[$

```
c*d^2 + a*e^2, 0] && IGtQ[m, 1] && (NeQ[d, 0] || GtQ[m, 2])
```

Rule 4972

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))*((d_) + (e_)*(x_))^(q_), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned} \int (d+ex)^3 (a+b\tan^{-1}(cx)) \, dx &= \frac{(d+ex)^4 (a+b\tan^{-1}(cx))}{4e} - \frac{(bc) \int \frac{(d+ex)^4}{1+c^2x^2} \, dx}{4e} \\ &= \frac{(d+ex)^4 (a+b\tan^{-1}(cx))}{4e} - \frac{(bc) \int \left(\frac{e^2(6c^2d^2-e^2)}{c^4} + \frac{4de^3x}{c^2} + \frac{e^4x^2}{c^2} + \frac{c^4d^4-6e^4}{c^8} \right) \, dx}{4e} \\ &= -\frac{be(6c^2d^2-e^2)x}{4c^3} - \frac{bde^2x^2}{2c} - \frac{be^3x^3}{12c} + \frac{(d+ex)^4 (a+b\tan^{-1}(cx))}{4e} - \frac{b}{4c^4} \\ &= -\frac{be(6c^2d^2-e^2)x}{4c^3} - \frac{bde^2x^2}{2c} - \frac{be^3x^3}{12c} + \frac{(d+ex)^4 (a+b\tan^{-1}(cx))}{4e} - \frac{(b^2-4c^2)d^4}{4c^4} \\ &= -\frac{be(6c^2d^2-e^2)x}{4c^3} - \frac{bde^2x^2}{2c} - \frac{be^3x^3}{12c} - \frac{b(c^4d^4-6c^2d^2e^2+e^4)\tan^{-1}(cx)}{4c^4e} \end{aligned}$$

Mathematica [A]

time = 0.30, size = 218, normalized size = 1.51

$$\frac{(d+ex)^4(a+b\text{ArcTan}(cx)) - \frac{bc(2\sqrt{-c^2}e^2x(-3e^2+c^2(18d^2+6dex+e^2x^2))-3(c^4d^4+e^3(4\sqrt{-c^2}d+e)-2c^2d^2e(2\sqrt{-c^2}d+3e))\log(1-\sqrt{-c^2}x)+3(c^4d^4+2c^2d^2(2\sqrt{-c^2}d-3e)e+e^3(-4\sqrt{-c^2}d+e))\log(1+\sqrt{-c^2}x))}{6(-c^2)^{5/2}}}{4e}$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)^3*(a + b*ArcTan[c*x]), x]`

[Out] $((d + e*x)^4*(a + b*\text{ArcTan}[c*x]) - (b*c*(2*\text{Sqrt}[-c^2]*e^2*x*(-3*e^2 + c^2*(18*d^2 + 6*d*e*x + e^2*x^2)) - 3*(c^4*d^4 + e^3*(4*\text{Sqrt}[-c^2]*d + e) - 2*c^2*d^2*e*(2*\text{Sqrt}[-c^2]*d + 3*e))*\text{Log}[1 - \text{Sqrt}[-c^2]*x] + 3*(c^4*d^4 + 2*c^2*d^2*(2*\text{Sqrt}[-c^2]*d - 3*e)*e + e^3*(-4*\text{Sqrt}[-c^2]*d + e))*\text{Log}[1 + \text{Sqrt}[-c^2]*x]))/(6*(-c^2)^(5/2)))/(4*e)$

Maple [A]

time = 0.12, size = 179, normalized size = 1.24

method	result
--------	--------

derivative divides	$\frac{(ce+cd)^4 a}{4c^3 e} + b \arctan(cx) d^3 cx + \frac{3bce \arctan(cx) d^2 x^2}{2} + bc e^2 \arctan(cx) d x^3 + \frac{bc e^3 \arctan(cx) x^4}{4} - \frac{3be d^2 x}{2} - \frac{b e^2 d x^2}{2} - \frac{b e^3 x^3}{12} + \frac{c}{c}$
default	$\frac{(ce+cd)^4 a}{4c^3 e} + b \arctan(cx) d^3 cx + \frac{3bce \arctan(cx) d^2 x^2}{2} + bc e^2 \arctan(cx) d x^3 + \frac{bc e^3 \arctan(cx) x^4}{4} - \frac{3be d^2 x}{2} - \frac{b e^2 d x^2}{2} - \frac{b e^3 x^3}{12} + \frac{c}{c}$
risch	$\frac{ib d^4 \ln(c^2 x^2 + 1)}{16e} + \frac{3eb d^2 \arctan(cx)}{2c^2} + \frac{ie^2 bd x^3 \ln(-icx + 1)}{2} - \frac{i(ex+d)^4 b \ln(icx + 1)}{8e} + \frac{ib d^3 x \ln(-icx + 1)}{2} + \frac{ie^3}{c}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((e*x+d)^3*(a+b*arctan(c*x)),x,method=_RETURNVERBOSE)
[Out] 1/c*(1/4*(c*e*x+c*d)^4*a/c^3/e+b*arctan(c*x)*d^3*c*x+3/2*b*c*e*arctan(c*x)*d^2*x^2+2*b*c*e^2*arctan(c*x)*d*x^3+1/4*b*c*e^3*arctan(c*x)*x^4-3/2*b*e*d^2*x-1/2*b*e^2*d*x^2-1/12*b*e^3*x^3+1/4*b/c^2*e^3*x-1/2*b*ln(c^2*x^2+1)*d^3+1/2*b/c^2*e^2*ln(c^2*x^2+1)*d+3/2*b/c*e*arctan(c*x)*d^2-1/4*b/c^3*e^3*arctan(c*x))
```

Maxima [A]

time = 0.49, size = 184, normalized size = 1.28

$$\frac{1}{4}ax^4e^3 + adx^3e^2 + \frac{3}{2}ad^2x^2e + ad^3x + \frac{3}{2}\left(x^2\arctan(cx) - c\left(\frac{x}{c^2} - \frac{\arctan(cx)}{c^3}\right)\right)bd^2e + \frac{(2cx\arctan(cx) - \log(c^2x^2 + 1))bd^3}{2c} + \frac{1}{2}\left(2x^3\arctan(cx) - c\left(\frac{x^2}{c^2} - \frac{\log(c^2x^2 + 1)}{c^3}\right)\right)bde^2 + \frac{1}{12}\left(3x^4\arctan(cx) - c\left(\frac{c^2x^3 - 3x}{c^4} + \frac{3\arctan(cx)}{c^5}\right)\right)be^3$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^3*(a+b*arctan(c*x)),x, algorithm="maxima")
[Out] 1/4*a*x^4*e^3 + a*d*x^3*e^2 + 3/2*a*d^2*x^2*e + a*d^3*x + 3/2*(x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*b*d^2*e + 1/2*(2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*b*d^3/c + 1/2*(2*x^3*arctan(c*x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c^4))*b*d*e^2 + 1/12*(3*x^4*arctan(c*x) - c*((c^2*x^3 - 3*x)/c^4 + 3*arctan(c*x)/c^5))*b*e^3
```

Fricas [A]

time = 1.49, size = 190, normalized size = 1.32

$$\frac{12ac^4d^3x + 3(4bc^4dx^3e^2 + 4bc^4d^3x + (bc^4x^4 - b)e^3 + 6(bc^4d^2x^2 + bc^2d^2)e)\arctan(cx) + (3ac^4x^4 - bc^3x^3 + 3bcx)e^3 + 6(2ac^4dx^3 - bc^3dx^2)e^2 + 18(ac^4d^2x^2 - bc^3d^2x)e - 6(bc^3d^3 - bcde^2)\log(c^2x^2 + 1)}{12c^4}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^3*(a+b*arctan(c*x)),x, algorithm="fricas")
[Out] 1/12*(12*a*c^4*d^3*x + 3*(4*b*c^4*d*x^3*e^2 + 4*b*c^4*d^3*x + (b*c^4*x^4 - b)*e^3 + 6*(b*c^4*d^2*x^2 + b*c^2*d^2)*e)*arctan(c*x) + (3*a*c^4*x^4 - b*c^3*x^3 + 3*b*c*x)*e^3 + 6*(2*a*c^4*d*x^3 - b*c^3*d*x^2)*e^2 + 18*(a*c^4*d^2*x^2 - b*c^3*d^2*x)*e - 6*(b*c^3*d^3 - b*c*d*e^2)*log(c^2*x^2 + 1))/c^4
```

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 262 vs. 2(129) = 258.

time = 0.35, size = 262, normalized size = 1.82

$$\begin{cases} ad^3x + \frac{3ad^2ex^2}{2} + ade^2x^3 + \frac{ae^3x^4}{4} + bd^3x \operatorname{atan}(cx) + \frac{3bd^2ex^2 \operatorname{atan}(cx)}{2} + bde^2x^3 \operatorname{atan}(cx) + \frac{be^3x^4 \operatorname{atan}(cx)}{4} - \frac{bd^3 \log(x^2 + \frac{1}{c^2})}{2c} - \frac{3bd^2ex}{2c} - \frac{bd^2e^2x^2}{2c^2} - \frac{be^3x^3}{12c} + \frac{3bd^2e \operatorname{atan}(cx)}{2c^3} + \frac{bde^2 \log(x^2 + \frac{1}{c^2})}{2c^3} + \frac{be^3x}{4c^3} - \frac{be^3 \operatorname{atan}(cx)}{4c^4} & \text{for } c \neq 0 \\ a\left(d^3x + \frac{3d^2ex^2}{2} + de^2x^3 + \frac{e^3x^4}{4}\right) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)**3*(a+b*atan(c*x)),x)`

[Out] Piecewise((a*d**3*x + 3*a*d**2*e*x**2/2 + a*d**2*x**3 + a*e**3*x**4/4 + b*d**3*x*atan(c*x) + 3*b*d**2*e*x**2*atan(c*x)/2 + b*d**2*x**3*atan(c*x) + b*e**3*x**4*atan(c*x)/4 - b*d**3*log(x**2 + c*(-2))/(2*c) - 3*b*d**2*e*x/(2*c) - b*d**2*x**2/(2*c) - b*e**3*x**3/(12*c) + 3*b*d**2*e*atan(c*x)/(2*c**2) + b*d**2*log(x**2 + c*(-2))/(2*c**3) + b*e**3*x/(4*c**3) - b*e**3*atan(c*x)/(4*c**4), Ne(c, 0)), (a*(d**3*x + 3*d**2*e*x**2/2 + d*e**2*x**3 + e**3*x**4/4), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^3*(a+b*arctan(c*x)),x, algorithm="giac")`

[Out] sage0*x

Mupad [B]

time = 0.80, size = 197, normalized size = 1.37

$$\frac{a e^3 x^4 + a d^3 x - \frac{b d^3 \ln(c^2 x^2 + 1)}{2c} - \frac{b e^3 x^3}{12c} + b d^3 x \operatorname{atan}(cx) + \frac{3 a d^2 e x^2}{2} + a d e^2 x^3 + \frac{b e^3 x}{4c^3} - \frac{b e^3 \operatorname{atan}(cx)}{4c^4} + \frac{b e^3 x^4 \operatorname{atan}(cx)}{4} - \frac{3 b d^2 e x}{2c} + \frac{3 b d^2 e^2 \operatorname{atan}(cx)}{2c^2} + \frac{3 b d^2 e x^2 \operatorname{atan}(cx)}{2} + b d e^2 x^3 \operatorname{atan}(cx) + \frac{b d e^2 \ln(c^2 x^2 + 1)}{2c^3} - \frac{b d e^2 x^2}{2c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))*(d + e*x)^3,x)`

[Out] $(a \cdot e^3 \cdot x^4)/4 + a \cdot d^3 \cdot x - (b \cdot d^3 \cdot \log(c^2 \cdot x^2 + 1))/(2 \cdot c) - (b \cdot e^3 \cdot x^3)/(12 \cdot c) + b \cdot d^3 \cdot x \cdot \operatorname{atan}(c \cdot x) + (3 \cdot a \cdot d^2 \cdot e \cdot x^2)/2 + a \cdot d \cdot e^2 \cdot x^3 + (b \cdot e^3 \cdot x)/(4 \cdot c^3) - (b \cdot e^3 \cdot \operatorname{atan}(c \cdot x))/(4 \cdot c^4) + (b \cdot e^3 \cdot x^4 \cdot \operatorname{atan}(c \cdot x))/4 - (3 \cdot b \cdot d^2 \cdot e \cdot x)/(2 \cdot c) + (3 \cdot b \cdot d^2 \cdot e \cdot \operatorname{atan}(c \cdot x))/(2 \cdot c^2) + (3 \cdot b \cdot d^2 \cdot e \cdot x^2 \cdot \operatorname{atan}(c \cdot x))/2 + b \cdot d \cdot e^2 \cdot x^3 \cdot \operatorname{atan}(c \cdot x) + (b \cdot d \cdot e^2 \cdot \log(c^2 \cdot x^2 + 1))/(2 \cdot c^3) - (b \cdot d \cdot e^2 \cdot x^2)/(2 \cdot c)$

$$3.3 \quad \int (d + ex)^2 (a + b \text{ArcTan}(cx)) dx$$

Optimal. Leaf size=103

$$-\frac{bdex}{c} - \frac{be^2x^2}{6c} - \frac{bd\left(d^2 - \frac{3e^2}{c^2}\right) \text{ArcTan}(cx)}{3e} + \frac{(d + ex)^3(a + b \text{ArcTan}(cx))}{3e} - \frac{b(3c^2d^2 - e^2) \log(1 + c^2x^2)}{6c^3}$$

[Out] $-\frac{b*d*e*x/c - 1/6*b*e^2*x^2/c - 1/3*b*d*(d^2 - 3e^2/c^2)*\text{arctan}(c*x)/e + 1/3*(e*x + d)^3*(a + b*\text{arctan}(c*x))/e - 1/6*b*(3*c^2*d^2 - e^2)*\ln(c^2*x^2 + 1)/c^3}{3e}$

Rubi [A]

time = 0.06, antiderivative size = 103, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.312, Rules used = {4972, 716, 649, 209, 266}

$$\frac{(d + ex)^3(a + b \text{ArcTan}(cx))}{3e} - \frac{bd\text{ArcTan}(cx)\left(d^2 - \frac{3e^2}{c^2}\right)}{3e} - \frac{b(3c^2d^2 - e^2) \log(c^2x^2 + 1)}{6c^3} - \frac{bdex}{c} - \frac{be^2x^2}{6c}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(d + e*x)^2 * (a + b*\text{ArcTan}[c*x]), x]$

[Out] $-\frac{((b*d*e*x)/c) - (b*e^2*x^2)/(6*c) - (b*d*(d^2 - (3*e^2)/c^2)*\text{ArcTan}[c*x])/(3*e) + ((d + e*x)^3*(a + b*\text{ArcTan}[c*x]))/(3*e) - (b*(3*c^2*d^2 - e^2)*\text{Log}[1 + c^2*x^2])/(6*c^3)}{3e}$

Rule 209

$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \Rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*a \text{rcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \text{ || } \text{GtQ}[b, 0])$

Rule 266

$\text{Int}[(x_)^{m_}/((a_) + (b_)*(x_)^{n_}), x_Symbol] \Rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}[\{a, b, m, n\}, x] \&& \text{EqQ}[m, n - 1]$

Rule 649

$\text{Int}[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] \Rightarrow \text{Dist}[d, \text{Int}[1/(a + c*x^2), x], x] + \text{Dist}[e, \text{Int}[x/(a + c*x^2), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{!NiceSqrtQ}[(-a)*c]$

Rule 716

$\text{Int}[((d_) + (e_)*(x_))^{m_}/((a_) + (c_)*(x_)^2), x_Symbol] \Rightarrow \text{Int}[\text{PolynomialDivide}[(d + e*x)^m, a + c*x^2, x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{NeQ}[$

$c*d^2 + a*e^2, 0] \&& \text{IGtQ}[m, 1] \&& (\text{NeQ}[d, 0] \text{ || } \text{GtQ}[m, 2])$

Rule 4972

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))*((d_) + (e_)*(x_))^(q_), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/((e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned} \int (d+ex)^2 (a+b\tan^{-1}(cx)) \, dx &= \frac{(d+ex)^3 (a+b\tan^{-1}(cx))}{3e} - \frac{(bc) \int \frac{(d+ex)^3}{1+c^2x^2} \, dx}{3e} \\ &= \frac{(d+ex)^3 (a+b\tan^{-1}(cx))}{3e} - \frac{(bc) \int \left(\frac{3de^2}{c^2} + \frac{e^3x}{c^2} + \frac{c^2d^3-3de^2+e(3c^2d^2-e^2)x}{c^2(1+c^2x^2)} \right)}{3e} \\ &= -\frac{bdex}{c} - \frac{be^2x^2}{6c} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx))}{3e} - \frac{b \int \frac{c^2d^3-3de^2+e(3c^2d^2-e^2)x}{1+c^2x^2}}{3ce} \\ &= -\frac{bdex}{c} - \frac{be^2x^2}{6c} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx))}{3e} - \frac{1}{3} \left(bd \left(\frac{cd^2}{e} - \frac{3e}{c} \right) \right) \int \\ &= -\frac{bdex}{c} - \frac{be^2x^2}{6c} - \frac{bd \left(d^2 - \frac{3e^2}{c^2} \right) \tan^{-1}(cx)}{3e} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx))}{3e} \end{aligned}$$

Mathematica [A]

time = 0.23, size = 163, normalized size = 1.58

$$\frac{(d+ex)^3(a+b\text{ArcTan}(cx)) - \frac{b \left(c^2 e^2 x (6 d+e x)+\left(-e^2 \left(3 \sqrt{-c^2} d+e\right)+c^2 d^2 \left(\sqrt{-c^2} d+3 e\right)\right) \log \left(1-\sqrt{-c^2} x\right)-\left(c^2 d^2 \left(\sqrt{-c^2} d-3 e\right)+e^2 \left(-3 \sqrt{-c^2} d+e\right)\right) \log \left(1+\sqrt{-c^2} x\right)\right)}{2 c^3}}{3e}$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)^2*(a + b*ArcTan[c*x]), x]`

[Out] $((d + e*x)^3*(a + b*ArcTan[c*x]) - (b*(c^2*e^2*x*(6*d + e*x) + (-e^2*(3*Sqrt[-c^2]*d + e)) + c^2*d^2*(Sqrt[-c^2]*d + 3*e)*Log[1 - Sqrt[-c^2]*x] - (c^2*d^2*(Sqrt[-c^2]*d - 3*e) + e^2*(-3*Sqrt[-c^2]*d + e))*Log[1 + Sqrt[-c^2]*x]))/(2*c^3))/(3*e)$

Maple [A]

time = 0.11, size = 123, normalized size = 1.19

method	result
--------	--------

derivative divides	$\frac{(ce^2+cd)^3 a}{3c^2 e} + b \arctan(cx) d^2 cx + bce \arctan(cx) dx^2 + \frac{bc e^2 \arctan(cx) x^3}{3} - bedx - \frac{be^2 x^2}{6} - \frac{b \ln(c^2 x^2 + 1) d^2}{2} + \frac{b e^2 \ln(c^2 x^2 + 1)}{6c^2} +$
default	$\frac{(ce^2+cd)^3 a}{3c^2 e} + b \arctan(cx) d^2 cx + bce \arctan(cx) dx^2 + \frac{bc e^2 \arctan(cx) x^3}{3} - bedx - \frac{be^2 x^2}{6} - \frac{b \ln(c^2 x^2 + 1) d^2}{2} + \frac{b e^2 \ln(c^2 x^2 + 1)}{6c^2} +$
risch	$-\frac{i(ex+d)^3 b \ln(icx+1)}{6e} + \frac{iebd x^2 \ln(-icx+1)}{2} + \frac{ib d^2 x \ln(-icx+1)}{2} - \frac{bd^2 \ln(c^2 x^2 + 1)}{2c} + \frac{ebd \arctan(cx)}{c^2} + \frac{ib d^3}{c}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)^2*(a+b*arctan(c*x)),x,method=_RETURNVERBOSE)`

[Out] $1/c*(1/3*(c*e*x+c*d)^3*a/c^2/e+b*arctan(c*x)*d^2*c*x+b*c*e*arctan(c*x)*d*x^2+1/3*b*c*e^2*arctan(c*x)*x^3-b*e*d*x-1/6*b*e^2*x^2-1/2*b*ln(c^2*x^2+1)*d^2+1/6*b/c^2*e^2*ln(c^2*x^2+1)+b/c*e*arctan(c*x)*d)$

Maxima [A]

time = 0.47, size = 126, normalized size = 1.22

$$\frac{1}{3} ax^3 e^2 + adx^2 e + ad^2 x + \left(x^2 \arctan(cx) - c \left(\frac{x}{c^2} - \frac{\arctan(cx)}{c^3} \right) \right) bde + \frac{(2cx \arctan(cx) - \log(c^2 x^2 + 1)) bd^2}{2c} + \frac{1}{6} \left(2x^3 \arctan(cx) - c \left(\frac{x^2}{c^2} - \frac{\log(c^2 x^2 + 1)}{c^4} \right) \right) be^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x)),x, algorithm="maxima")`

[Out] $1/3*a*x^3*e^2 + a*d*x^2*e + a*d^2*x + (x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*b*d*e + 1/2*(2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*b*d^2/c + 1/6*(2*x^3*arctan(c*x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c^4))*b*e^2$

Fricas [A]

time = 1.54, size = 132, normalized size = 1.28

$$\frac{6 ac^3 d^2 x + 2(bc^3 x^3 e^2 + 3bc^3 d^2 x + 3(bc^3 dx^2 + bcd)e) \arctan(cx) + (2ac^3 x^3 - bc^2 x^2)e^2 + 6(ac^3 dx^2 - bc^2 dx)e - (3bc^2 d^2 - be^2) \log(c^2 x^2 + 1)}{6c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x)),x, algorithm="fricas")`

[Out] $1/6*(6*a*c^3*d^2*x + 2*(b*c^3*x^3*e^2 + 3*b*c^3*d^2*x + 3*(b*c^3*d*x^2 + b*c*d)*e)*arctan(c*x) + (2*a*c^3*x^3 - b*c^2*x^2)*e^2 + 6*(a*c^3*d*x^2 - b*c^2*d*x)*e - (3*b*c^2*d^2 - b*e^2)*log(c^2*x^2 + 1))/c^3$

Sympy [A]

time = 0.27, size = 160, normalized size = 1.55

$$\begin{cases} ad^2 x + adex^2 + \frac{ae^2 x^3}{3} + bd^2 x \operatorname{atan}(cx) + bdex^2 \operatorname{atan}(cx) + \frac{be^2 x^3 \operatorname{atan}(cx)}{3} - \frac{bd^2 \log\left(\frac{x^2 + 1}{c^2}\right)}{2c} - \frac{bdex}{c} - \frac{be^2 x^2}{6c} + \frac{bde \operatorname{atan}(cx)}{c^2} + \frac{be^2 \log\left(\frac{x^2 + 1}{c^2}\right)}{6c^3} & \text{for } c \neq 0 \\ a(d^2 x + dex^2 + \frac{e^2 x^3}{3}) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)**2*(a+b*atan(c*x)),x)`

[Out] `Piecewise((a*d**2*x + a*d*e*x**2 + a*e**2*x**3/3 + b*d**2*x*atan(c*x) + b*d*e*x**2*atan(c*x) + b*e**2*x**3*atan(c*x)/3 - b*d**2*log(x**2 + c*(-2))/(2*c) - b*d*e*x/c - b*e**2*x**2/(6*c) + b*d*e*atan(c*x)/c**2 + b*e**2*log(x**2 + c*(-2))/(6*c**3), Ne(c, 0)), (a*(d**2*x + d*e*x**2 + e**2*x**3/3), True))`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B]

time = 0.35, size = 127, normalized size = 1.23

$$\frac{ae^2 x^3}{3} + ad^2 x - \frac{bd^2 \ln(c^2 x^2 + 1)}{2c} + \frac{be^2 \ln(c^2 x^2 + 1)}{6c^3} - \frac{be^2 x^2}{6c} + ade x^2 + bd^2 x \operatorname{atan}(cx) + \frac{be^2 x^3 \operatorname{atan}(cx)}{3} - \frac{bde x}{c} + \frac{bde \operatorname{atan}(cx)}{c^2} + bde x^2 \operatorname{atan}(cx)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))*(d + e*x)^2,x)`

[Out] `(a*e^2*x^3)/3 + a*d^2*x - (b*d^2*log(c^2*x^2 + 1))/(2*c) + (b*e^2*log(c^2*x^2 + 1))/(6*c^3) - (b*e^2*x^2)/(6*c) + a*d*e*x^2 + b*d^2*x*atan(c*x) + (b*e^2*x^3*atan(c*x))/3 - (b*d*e*x)/c + (b*d*e*atan(c*x))/c^2 + b*d*e*x^2*atan(c*x)`

3.4 $\int (d + ex)(a + b \operatorname{ArcTan}(cx)) dx$

Optimal. Leaf size=76

$$-\frac{bex}{2c} - \frac{b\left(d^2 - \frac{e^2}{c^2}\right) \operatorname{ArcTan}(cx)}{2e} + \frac{(d+ex)^2(a+b\operatorname{ArcTan}(cx))}{2e} - \frac{bd \log(1+c^2x^2)}{2c}$$

[Out] $-1/2*b*e*x/c - 1/2*b*(d^2 - e^2/c^2)*\operatorname{arctan}(c*x)/e + 1/2*(e*x+d)^2*(a+b*\operatorname{arctan}(c*x))/e - 1/2*b*d*\ln(c^2*x^2+1)/c$

Rubi [A]

time = 0.04, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {4972, 716, 649, 209, 266}

$$\frac{(d+ex)^2(a+b\operatorname{ArcTan}(cx))}{2e} - \frac{b\operatorname{ArcTan}(cx)\left(d^2 - \frac{e^2}{c^2}\right)}{2e} - \frac{bd \log(c^2x^2 + 1)}{2c} - \frac{bex}{2c}$$

Antiderivative was successfully verified.

[In] Int[(d + e*x)*(a + b*ArcTan[c*x]), x]

[Out] $-1/2*(b*e*x)/c - (b*(d^2 - e^2/c^2)*\operatorname{ArcTan}(c*x))/(2*e) + ((d + e*x)^2*(a + b*\operatorname{ArcTan}(c*x)))/(2*e) - (b*d*\log(1 + c^2*x^2))/(2*c)$

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 266

Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> Simp[Log[RemoveContent[t[a + b*x^n, x]]/(b*n)], x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]

Rule 649

Int[((d_) + (e_.)*(x_))/((a_) + (c_.)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && !NiceSqrtQ[(-a)*c]

Rule 716

Int[((d_) + (e_.)*(x_))^(m_)/((a_) + (c_.)*(x_)^2), x_Symbol] :> Int[PolynomialDivide[(d + e*x)^m, a + c*x^2, x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && IGtQ[m, 1] && (NeQ[d, 0] || GtQ[m, 2])

Rule 4972

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.*))*((d_) + (e_.*)(x_))^q, x_Symbol]
  :> Simp[(d + e*x)^q*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^q/(1 + c^2*x^2), x], x] /; FreeQ[{a, b,
  c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned}
 \int (d + ex) (a + b \tan^{-1}(cx)) \, dx &= \frac{(d + ex)^2 (a + b \tan^{-1}(cx))}{2e} - \frac{(bc) \int \frac{(d+ex)^2}{1+c^2x^2} \, dx}{2e} \\
 &= \frac{(d + ex)^2 (a + b \tan^{-1}(cx))}{2e} - \frac{(bc) \int \left(\frac{e^2}{c^2} + \frac{c^2d^2-e^2+2c^2dex}{c^2(1+c^2x^2)} \right) \, dx}{2e} \\
 &= -\frac{bex}{2c} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))}{2e} - \frac{b \int \frac{c^2d^2-e^2+2c^2dex}{1+c^2x^2} \, dx}{2ce} \\
 &= -\frac{bex}{2c} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))}{2e} - (bcd) \int \frac{x}{1+c^2x^2} \, dx - \frac{(b(cd - e))(}{2c} \\
 &= -\frac{bex}{2c} - \frac{b \left(d^2 - \frac{e^2}{c^2} \right) \tan^{-1}(cx)}{2e} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))}{2e} - \frac{bd \log(1 + c^2x^2)}{2c}
 \end{aligned}$$

Mathematica [A]

time = 0.01, size = 77, normalized size = 1.01

$$adx - \frac{bex}{2c} + \frac{1}{2}aex^2 + \frac{be \operatorname{ArcTan}(cx)}{2c^2} + bdx \operatorname{ArcTan}(cx) + \frac{1}{2}bex^2 \operatorname{ArcTan}(cx) - \frac{bd \log(1 + c^2x^2)}{2c}$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)*(a + b*ArcTan[c*x]), x]`[Out] $a*d*x - (b*e*x)/(2*c) + (a*e*x^2)/2 + (b*e*ArcTan[c*x])/(2*c^2) + b*d*x*ArcTan[c*x] + (b*e*x^2*ArcTan[c*x])/2 - (b*d*Log[1 + c^2*x^2])/(2*c)$ Maple [A]

time = 0.04, size = 78, normalized size = 1.03

method	result
derivativedivides	$\frac{a(d c^2 x + \frac{1}{2} e c^2 x^2)}{c} + b \arctan(cx) dx + \frac{b c \arctan(cx) e x^2}{2} - \frac{b e x}{2} - \frac{b d \ln(c^2 x^2 + 1)}{2} + \frac{b e \arctan(cx)}{2c}$
default	$\frac{a(d c^2 x + \frac{1}{2} e c^2 x^2)}{c} + b \arctan(cx) dx + \frac{b c \arctan(cx) e x^2}{2} - \frac{b e x}{2} - \frac{b d \ln(c^2 x^2 + 1)}{2} + \frac{b e \arctan(cx)}{2c}$

risch	$-\frac{ib(ex^2+2dx)\ln(icx+1)}{4} + \frac{ibe x^2 \ln(-icx+1)}{4} + \frac{ibdx \ln(-icx+1)}{2} + \frac{ae x^2}{2} + adx - \frac{bd \ln(c^2 x^2 + 1)}{2c} - \frac{bex}{2c} + \dots$
-------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)*(a+b*arctan(c*x)),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{c} \left(\frac{a}{c} \left(\frac{d c^2 x^2 + 1/2 e c^2 x^2 + 1}{2} \right) + b \arctan(cx) \left(d c x + 1/2 b c \arctan(cx) * e x^2 - 1/2 b e x - 1/2 b d \ln(c^2 x^2 + 1) + 1/2 b/c e \arctan(cx) \right) \right)$

Maxima [A]

time = 0.48, size = 73, normalized size = 0.96

$$\frac{1}{2} a x^2 e + a d x + \frac{1}{2} \left(x^2 \arctan(cx) - c \left(\frac{x}{c^2} - \frac{\arctan(cx)}{c^3} \right) \right) b e + \frac{(2 c x \arctan(cx) - \log(c^2 x^2 + 1)) b d}{2 c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x)),x, algorithm="maxima")`

[Out] $\frac{1}{2} \left(2 a x^2 e + a d x + \frac{1}{2} \left(x^2 \arctan(cx) - c \left(\frac{x}{c^2} - \frac{\arctan(cx)}{c^3} \right) \right) b e + \frac{1}{2} \left(2 c x \arctan(cx) - \log(c^2 x^2 + 1) \right) b d \right) / c$

Fricas [A]

time = 1.19, size = 73, normalized size = 0.96

$$\frac{2 a c^2 d x - b c d \log(c^2 x^2 + 1) + (2 b c^2 d x + (b c^2 x^2 + b) e) \arctan(cx) + (a c^2 x^2 - b c x) e}{2 c^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x)),x, algorithm="fricas")`

[Out] $\frac{1}{2} \left(2 a c^2 d x - b c d \log(c^2 x^2 + 1) + (2 b c^2 d x + (b c^2 x^2 + b) e) \arctan(cx) + (a c^2 x^2 - b c x) e \right) / c^2$

Sympy [A]

time = 0.16, size = 87, normalized size = 1.14

$$\begin{cases} adx + \frac{aex^2}{2} + bdx \operatorname{atan}(cx) + \frac{bex^2 \operatorname{atan}(cx)}{2} - \frac{bd \log\left(x^2 + \frac{1}{c^2}\right)}{2c} - \frac{bex}{2c} + \frac{be \operatorname{atan}(cx)}{2c^2} & \text{for } c \neq 0 \\ a \left(dx + \frac{ex^2}{2} \right) & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*atan(c*x)),x)`

[Out] `Piecewise((a*d*x + a*e*x**2/2 + b*d*x*atan(c*x) + b*e*x**2*atan(c*x)/2 - b*d*log(x**2 + c*(-2))/(2*c) - b*e*x/(2*c) + b*e*atan(c*x)/(2*c**2), Ne(c, 0)), (a*(d*x + e*x**2/2), True))`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B]

time = 0.41, size = 67, normalized size = 0.88

$$adx + \frac{ae x^2}{2} + bdx \operatorname{atan}(cx) - \frac{be x}{2c} + \frac{be \operatorname{atan}(cx)}{2c^2} + \frac{be x^2 \operatorname{atan}(cx)}{2} - \frac{bd \ln(c^2 x^2 + 1)}{2c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))*(d + e*x),x)`

[Out] `a*d*x + (a*e*x^2)/2 + b*d*x*atan(cx) - (b*e*x)/(2*c) + (b*e*atan(cx))/(2*c^2) + (b*e*x^2*atan(cx))/2 - (b*d*log(c^2*x^2 + 1))/(2*c)`

$$3.5 \quad \int \frac{a+b\text{ArcTan}(cx)}{d+ex} dx$$

Optimal. Leaf size=138

$$-\frac{(a + b\text{ArcTan}(cx)) \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a + b\text{ArcTan}(cx)) \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{ib\text{PolyLog}(2, 1 - \frac{2}{1-icx})}{2e} - \frac{ib\text{PolyLog}(2, 1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)})}{2e}$$

[Out] $-(a+b*\arctan(c*x))*\ln(2/(1-I*c*x))/e+(a+b*\arctan(c*x))*\ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e+1/2*I*b*polylog(2,1-2/(1-I*c*x))/e-1/2*I*b*polylog(2,1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e$

Rubi [A]

time = 0.07, antiderivative size = 138, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.250, Rules used = {4966, 2449, 2352, 2497}

$$\frac{(a + b\text{ArcTan}(cx)) \log\left(\frac{2c(d+ex)}{(1-icx)(cd+ie)}\right)}{e} - \frac{\log\left(\frac{2}{1-icx}\right) (a + b\text{ArcTan}(cx))}{e} - \frac{ib\text{Li}_2\left(1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{2e} + \frac{ib\text{Li}_2\left(1 - \frac{2}{1-icx}\right)}{2e}$$

Antiderivative was successfully verified.

[In] Int[(a + b*ArcTan[c*x])/(d + e*x), x]

[Out] $-(((a + b*\arctan(c*x))*\text{Log}[2/(1 - I*c*x)])/e) + ((a + b*\arctan(c*x))*\text{Log}[(2*c*(d + e*x))/(c*d + I*e)*(1 - I*c*x)])/e + ((I/2)*b*\text{PolyLog}[2, 1 - 2/(1 - I*c*x)]/e - ((I/2)*b*\text{PolyLog}[2, 1 - (2*c*(d + e*x))/((c*d + I*e)*(1 - I*c*x))])/e$

Rule 2352

Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]

Rule 2449

Int[Log[(c_)/((d_) + (e_)*(x_))] / ((f_) + (g_)*(x_)^2), x_Symbol] :> Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]

Rule 2497

Int[Log[u_]*(Pq_)^(m_), x_Symbol] :> With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]]

Rule 4966

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_))/((d_) + (e_.)*(x_)), x_Symbol] :> Si
mp[(-(a + b*ArcTan[c*x]))*(Log[2/(1 - I*c*x])/e), x] + (Dist[b*(c/e), Int[Log[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e
*x)/((c*d + I*e)*(1 - I*c*x)))]/(1 + c^2*x^2), x], x] + Simp[(a + b*ArcTan[
c*x])*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x]) /; FreeQ[{a,
b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rubi steps

$$\begin{aligned} \int \frac{a + b \tan^{-1}(cx)}{d + ex} dx &= -\frac{(a + b \tan^{-1}(cx)) \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a + b \tan^{-1}(cx)) \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{(bc) \int \frac{\log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{1+c^2x^2} dx}{e} \\ &= -\frac{(a + b \tan^{-1}(cx)) \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a + b \tan^{-1}(cx)) \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} - \frac{ib \text{Li}_2(1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)})}{e} \\ &= -\frac{(a + b \tan^{-1}(cx)) \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a + b \tan^{-1}(cx)) \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{ib \text{Li}_2(1 - \frac{2c(d+ex)}{(cd+ie)(1-icx)})}{2e} \end{aligned}$$

Mathematica [A]

time = 0.14, size = 245, normalized size = 1.78

$$\frac{a \log(d+ex) + b \text{ArcTan}(cx) \left(\frac{1}{2} \log(1+c^2 x^2) + \log(\sin(\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex))) + \frac{1}{2} (-i (r - 2 \text{ArcTan}(cx))^2 - i (\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex))^2 + (r - 2 \text{ArcTan}(ex)) \log(1 + e^{-i \text{ArcTan}(ex)}) + 2 (\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex)) \log\left(1 - e^{i (\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex))}\right) - (r - 2 \text{ArcTan}(ex)) \log\left(\frac{e^{i (\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex))}}{\sqrt{1+c^2 x^2}}\right) - 2 (\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex)) \log(\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex)) - i \text{PolyLog}(2, e^{-i \text{ArcTan}(ex)}) - i \text{PolyLog}\left(2, e^{i (\text{ArcTan}(\frac{d}{2}) + \text{ArcTan}(ex))}\right)\right)}{e}$$

Antiderivative was successfully verified.

[In] `Integrate[(a + b*ArcTan[c*x])/(d + e*x), x]`

[Out] `(a*Log[d + e*x] + b*ArcTan[c*x]*(Log[1 + c^2*x^2]/2 + Log[Sin[ArcTan[(c*d)/e] + ArcTan[c*x]]]) + (b*((-1/4*I)*(Pi - 2*ArcTan[c*x])^2 - I*(ArcTan[(c*d)/e] + ArcTan[c*x])^2 + (Pi - 2*ArcTan[c*x])*Log[1 + E^((-2*I)*ArcTan[c*x])] + 2*(ArcTan[(c*d)/e] + ArcTan[c*x])*Log[1 - E^((-2*I)*(ArcTan[(c*d)/e] + ArcTan[c*x]))] - (Pi - 2*ArcTan[c*x])*Log[2/Sqrt[1 + c^2*x^2]] - 2*(ArcTan[(c*d)/e] + ArcTan[c*x])*Log[2*Sin[ArcTan[(c*d)/e] + ArcTan[c*x]]] - I*PolyLog[2, -E^((-2*I)*ArcTan[c*x])] - I*PolyLog[2, E^((-2*I)*(ArcTan[(c*d)/e] + ArcTan[c*x]))])))/2)/e`

Maple [A]

time = 0.06, size = 178, normalized size = 1.29

method	result
derivative divides	$\frac{ac \ln(cx+d)}{e} + \frac{bc \ln(cx+d) \arctan(cx)}{e} + \frac{ibc \ln(cx+d) \ln\left(\frac{-cex+ie}{cd+ie}\right)}{2e} - \frac{ibc \ln(cx+d) \ln\left(\frac{cex+ie}{-cd+ie}\right)}{2e} + \frac{ibc \operatorname{dilog}\left(\frac{-cex+ie}{cd+ie}\right)}{2e} - \frac{ibc \operatorname{dilog}\left(\frac{cex+ie}{-cd+ie}\right)}{2e}$

default	$\frac{ac \ln(cx+d)}{e} + \frac{bc \ln(cx+d) \arctan(cx)}{e} + \frac{ibc \ln(cx+d) \ln\left(\frac{-cx+ie}{cd+ie}\right)}{2e} - \frac{ibc \ln(cx+d) \ln\left(\frac{ce+ie}{cd+ie}\right)}{2e} + \frac{ibc \operatorname{dilog}\left(\frac{-ce+ie}{cd+ie}\right)}{2e} - \frac{ibc \operatorname{dilog}\left(\frac{ce+ie}{cd+ie}\right)}{2e}$
risch	$\frac{ib \operatorname{dilog}\left(\frac{-icd+(-icx+1)e-e}{-icd-e}\right)}{2e} + \frac{ib \ln(-icx+1) \ln\left(\frac{-icd+(-icx+1)e-e}{-icd-e}\right)}{2e} + \frac{a \ln(icd-(-icx+1)e+e)}{e} - \frac{ib \operatorname{dilog}\left(\frac{icd+(-icx+1)e-e}{-icd-e}\right)}{2e}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a+b*arctan(c*x))/(e*x+d),x,method=_RETURNVERBOSE)
[Out] 1/c*(a*c*ln(c*e*x+c*d)/e+b*c*ln(c*e*x+c*d)/e*arctan(c*x)+1/2*I*b*c*ln(c*e*x+c*d)/e*ln((I*e-c*e*x)/(c*d+I*e))-1/2*I*b*c*ln(c*e*x+c*d)/e*ln((I*e+c*e*x)/(I*e-c*d))+1/2*I*b*c/e*dilog((I*e-c*e*x)/(c*d+I*e))-1/2*I*b*c/e*dilog((I*e+c*e*x)/(I*e-c*d)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))/(e*x+d),x, algorithm="maxima")
[Out] a*e^(-1)*log(x*e + d) + 2*b*integrate(1/2*arctan(c*x)/(x*e + d), x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))/(e*x+d),x, algorithm="fricas")
[Out] integral((b*arctan(c*x) + a)/(x*e + d), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{a + b \operatorname{atan}(cx)}{d + ex} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*atan(c*x))/(e*x+d),x)
[Out] Integral((a + b*atan(c*x))/(d + e*x), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(cx))/(e*x+d),x, algorithm="giac")`
[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{a + b \operatorname{atan}(cx)}{d + e x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(cx))/(d + e*x),x)`
[Out] `int((a + b*atan(cx))/(d + e*x), x)`

3.6 $\int \frac{a+b\text{ArcTan}(cx)}{(d+ex)^2} dx$

Optimal. Leaf size=98

$$\frac{bc^2 d \text{ArcTan}(cx)}{e(c^2 d^2 + e^2)} - \frac{a + b \text{ArcTan}(cx)}{e(d + ex)} + \frac{bc \log(d + ex)}{c^2 d^2 + e^2} - \frac{bc \log(1 + c^2 x^2)}{2(c^2 d^2 + e^2)}$$

[Out] $b*c^2*d*\arctan(c*x)/e/(c^2*d^2+e^2)+(-a-b*\arctan(c*x))/e/(e*x+d)+b*c*\ln(e*x+d)/(c^2*d^2+e^2)-1/2*b*c*\ln(c^2*x^2+1)/(c^2*d^2+e^2)$

Rubi [A]

time = 0.04, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4972, 720, 31, 649, 209, 266}

$$-\frac{a + b \text{ArcTan}(cx)}{e(d + ex)} + \frac{bc^2 d \text{ArcTan}(cx)}{e(c^2 d^2 + e^2)} - \frac{bc \log(c^2 x^2 + 1)}{2(c^2 d^2 + e^2)} + \frac{bc \log(d + ex)}{c^2 d^2 + e^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x])/(d + e*x)^2, x]$

[Out] $(b*c^2*d*\text{ArcTan}[c*x])/(\text{e}*(c^2*d^2 + e^2)) - (a + b*\text{ArcTan}[c*x])/(\text{e}*(d + e*x)) + (b*c*\text{Log}[d + e*x])/(\text{c}^2*\text{d}^2 + e^2) - (b*c*\text{Log}[1 + c^2*x^2])/(2*(c^2*d^2 + e^2))$

Rule 31

$\text{Int}[((a_) + (b_)*(x_))^{(-1)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}[\{a, b\}, x]$

Rule 209

$\text{Int}[((a_) + (b_)*(x_))^{(-1)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*a \text{rcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \text{ || } \text{GtQ}[b, 0])$

Rule 266

$\text{Int}[(x_)^{(m_)}/((a_) + (b_)*(x_))^{(n_)}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}[\{a, b, m, n\}, x] \&& \text{EqQ}[m, n - 1]$

Rule 649

$\text{Int}[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{Dist}[d, \text{Int}[1/(a + c*x^2), x], x] + \text{Dist}[e, \text{Int}[x/(a + c*x^2), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{!NiceSqrtQ}[(-a)*c]$

Rule 720

```
Int[1/(((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)), x_Symbol] :> Dist[e^2/(c*d^2 + a*e^2), Int[1/(d + e*x), x], x] + Dist[1/(c*d^2 + a*e^2), Int[(c*d - c*e*x)/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0]
```

Rule 4972

```
Int[((a_) + ArcTan[(c_)*(x_)]*(b_))*((d_) + (e_)*(x_))^(q_), x_Symbol]
:> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned} \int \frac{a + b \tan^{-1}(cx)}{(d + ex)^2} dx &= -\frac{a + b \tan^{-1}(cx)}{e(d + ex)} + \frac{(bc) \int \frac{1}{(d+ex)(1+c^2x^2)} dx}{e} \\ &= -\frac{a + b \tan^{-1}(cx)}{e(d + ex)} + \frac{(bc) \int \frac{c^2d - c^2ex}{1+c^2x^2} dx}{e(c^2d^2 + e^2)} + \frac{(bce) \int \frac{1}{d+ex} dx}{c^2d^2 + e^2} \\ &= -\frac{a + b \tan^{-1}(cx)}{e(d + ex)} + \frac{bc \log(d + ex)}{c^2d^2 + e^2} - \frac{(bc^3) \int \frac{x}{1+c^2x^2} dx}{c^2d^2 + e^2} + \frac{(bc^3d) \int \frac{1}{1+c^2x^2} dx}{e(c^2d^2 + e^2)} \\ &= \frac{bc^2d \tan^{-1}(cx)}{e(c^2d^2 + e^2)} - \frac{a + b \tan^{-1}(cx)}{e(d + ex)} + \frac{bc \log(d + ex)}{c^2d^2 + e^2} - \frac{bc \log(1 + c^2x^2)}{2(c^2d^2 + e^2)} \end{aligned}$$

Mathematica [A]

time = 0.13, size = 111, normalized size = 1.13

$$\frac{-\frac{a+b \text{ArcTan}(cx)}{d+ex}+\frac{bc \left(\left(\sqrt{-c^2} d-e\right) \log \left(1-\sqrt{-c^2} x\right)-\left(\sqrt{-c^2} d+e\right) \log \left(1+\sqrt{-c^2} x\right)+2 e \log (d+ex)\right)}{2 (c^2 d^2+e^2)}}{e}$$

Antiderivative was successfully verified.

[In] `Integrate[(a + b*ArcTan[c*x])/(d + e*x)^2, x]`

[Out] $\frac{\left(-((a+b \text{ArcTan}[c x])/(d+e x))+\frac{(b+c ((\text{Sqrt}[-c^2] * d-e) * \text{Log}[1-\text{Sqrt}[-c^2] * x]-(\text{Sqrt}[-c^2] * d+e) * \text{Log}[1+\text{Sqrt}[-c^2] * x]+2 e * \text{Log}[d+e x]))/(2 * (c^2 * d^2+e^2))\right)/e$

Maple [A]

time = 0.10, size = 130, normalized size = 1.33

method	result
derivativedivides	$\frac{-\frac{a c^2}{(cex+cd)e} - \frac{b c^2 \arctan(cx)}{(cex+cd)e} - \frac{b c^2 \ln(c^2 x^2 + 1)}{2(c^2 d^2 + e^2)} + \frac{b c^3 d \arctan(cx)}{e(c^2 d^2 + e^2)} + \frac{b c^2 \ln(cex+cd)}{c^2 d^2 + e^2}}{c}$
default	$\frac{-\frac{a c^2}{(cex+cd)e} - \frac{b c^2 \arctan(cx)}{(cex+cd)e} - \frac{b c^2 \ln(c^2 x^2 + 1)}{2(c^2 d^2 + e^2)} + \frac{b c^3 d \arctan(cx)}{e(c^2 d^2 + e^2)} + \frac{b c^2 \ln(cex+cd)}{c^2 d^2 + e^2}}{c}$
risch	$\frac{ib \ln(icx+1)}{2e(ex+d)} + \frac{-ib c^2 d^2 \ln(-icx+1) - ib e^2 \ln(-icx+1) + 2 \ln(-ex-d) bc e^2 x + 2 \ln(-ex-d) bcde - 2a c^2 d^2 - 2a e^2 - \ln((-ex-d)^2 bcde)}{2e(ex+d)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x))/(e*x+d)^2,x,method=_RETURNVERBOSE)`

[Out]
$$\frac{1}{c} * (-a*c^2/(c*e*x+c*d)/e - b*c^2/(c*e*x+c*d)/e*\arctan(c*x) - 1/2*b*c^2/(c^2*d^2+e^2)*\ln(c^2*x^2+1) + b*c^3/e/(c^2*d^2+e^2)*d*\arctan(c*x) + b*c^2/(c^2*d^2+e^2)*\ln(c*e*x+c*d))$$

Maxima [A]

time = 0.47, size = 106, normalized size = 1.08

$$\frac{1}{2} \left(\left(\frac{2 cd \arctan(cx)}{c^2 d^2 e + e^3} - \frac{\log(c^2 x^2 + 1)}{c^2 d^2 + e^2} + \frac{2 \log(xe + d)}{c^2 d^2 + e^2} \right) c - \frac{2 \arctan(cx)}{xe^2 + de} \right) b - \frac{a}{xe^2 + de}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))/(e*x+d)^2,x, algorithm="maxima")`

[Out]
$$\frac{1}{2} * ((2*c*d*\arctan(c*x)/(c^2*d^2*e + e^3) - \log(c^2*x^2 + 1)/(c^2*d^2 + e^2) + 2*\log(xe + d)/(c^2*d^2 + e^2))*c - 2*\arctan(c*x)/(x*e^2 + d*e))*b - a/(x*e^2 + d*e)$$

Fricas [A]

time = 1.14, size = 114, normalized size = 1.16

$$\frac{2 ac^2 d^2 - 2 (bc^2 d x e - be^2) \arctan(cx) + 2 ae^2 + (bc x e^2 + bcde) \log(c^2 x^2 + 1) - 2 (bc x e^2 + bcde) \log(xe + d)}{2 (c^2 d^2 x e^2 + c^2 d^3 e + x e^4 + de^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))/(e*x+d)^2,x, algorithm="fricas")`

[Out]
$$\frac{-1/2*(2*a*c^2*d^2 - 2*(b*c^2*d*x*e - b*e^2)*\arctan(c*x) + 2*a*e^2 + (b*c*x*e^2 + b*c*d*e)*\log(c^2*x^2 + 1) - 2*(b*c*x*e^2 + b*c*d*e)*\log(xe + d))/(c^2*d^2*x*e^2 + c^2*d^3*e + x*e^4 + d*e^3)}$$

Sympy [C] Result contains complex when optimal does not.

time = 1.49, size = 682, normalized size = 6.96

$$\begin{cases} -\frac{a}{de^2x^2} & \text{for } c = 0 \\ -\frac{2ad}{2d^2e+2de^2x} + \frac{i b d \operatorname{atanh}\left(\frac{cx}{d}\right)}{2d^2e+2de^2x} + \frac{i b e x \operatorname{atanh}\left(\frac{cx}{d}\right)}{2d^2e+2de^2x} - \frac{i b e x \operatorname{atanh}\left(\frac{dx}{e}\right)}{2d^2e+2de^2x} \\ -\frac{2ad}{2d^2e+2de^2x} - \frac{i b d \operatorname{atanh}\left(\frac{dx}{e}\right)}{2d^2e+2de^2x} - \frac{i b e x \operatorname{atanh}\left(\frac{dx}{e}\right)}{2d^2e+2de^2x} + \frac{i b e x \operatorname{atanh}\left(\frac{cx}{d}\right)}{2d^2e+2de^2x} \\ \infty \left(ax + bx \operatorname{atan}(cx) - \frac{b \log\left(x^2 + \frac{1}{c^2}\right)}{2c} \right) & \text{for } c = -\frac{ie}{d} \\ \frac{ax + bx \operatorname{atan}(cx) - \frac{b \log\left(x^2 + \frac{1}{c^2}\right)}{2c}}{d^2} & \text{for } c = \frac{ie}{d} \\ -\frac{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} - \frac{2ae^2}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} + \frac{2be^2 \operatorname{dex} \operatorname{atan}(cx)}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} - \frac{bde \log\left(x^2 + \frac{1}{c^2}\right)}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} + \frac{2bde \log\left(\frac{d}{e} + x\right)}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} - \frac{bcx^2 \log\left(x^2 + \frac{1}{c^2}\right)}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} + \frac{2bcx^2 \log\left(\frac{d}{e} + x\right)}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} - \frac{2be^2 \operatorname{atan}(cx)}{2c^2d^3e+2c^2d^2e^2x+2de^3+2e^4x} & \text{for } d = -ex \\ \text{otherwise} & \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x))/(e*x+d)**2,x)`

[Out] Piecewise((-a/(d*e + e**2*x), Eq(c, 0)), (-2*a*d/(2*d**2*e + 2*d*e**2*x) + I*b*d*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x) + I*b*d/(2*d**2*e + 2*d*e**2*x) - I*b*e*x*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x), Eq(c, -I*e/d)), (-2*a*d/(2*d**2*e + 2*d*e**2*x) - I*b*d*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x) - I*b*d/(2*d**2*e + 2*d*e**2*x) + I*b*e*x*atanh(e*x/d)/(2*d**2*e + 2*d*e**2*x), Eq(c, I*e/d)), (zoo*(a*x + b*x*atan(c*x) - b*log(x**2 + c*(-2))/(2*c)), Eq(d, -e*x)), ((a*x + b*x*atan(c*x) - b*log(x**2 + c*(-2))/(2*c))/d**2, Eq(e, 0)), (-2*a*c**2*d**2/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) - 2*a*e**2/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) + 2*b*c**2*d*x*atan(c*x)/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) - b*c*d*e*log(x**2 + c*(-2))/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) + 2*b*c*d*e*log(d/e + x)/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) - b*c*x**2*log(x**2 + c*(-2))/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) + 2*b*c*x**2*log(d/e + x)/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x) - 2*b*x**2*atan(c*x)/(2*c**2*d**3*e + 2*c**2*d**2*e**2*x + 2*d*e**3 + 2*e**4*x), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))/(e*x+d)^2,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B]

time = 3.46, size = 112, normalized size = 1.14

$$\frac{d^2 \left(b c \ln(d + e x) - \frac{b c \ln(c^2 x^2 + 1)}{2} + a c^2 x + b c^2 x \operatorname{atan}(c x) \right) - d e \left(b \operatorname{atan}(c x) - b c x \ln(d + e x) + \frac{b c x \ln(c^2 x^2 + 1)}{2} \right) + a e^2 x}{d (c^2 d^2 + e^2) (d + e x)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))/(d + e*x)^2,x)`

```
[Out] (d^2*(b*c*log(d + e*x) - (b*c*log(c^2*x^2 + 1))/2 + a*c^2*x + b*c^2*x*atan(c*x)) - d*e*(b*atan(c*x) - b*c*x*log(d + e*x) + (b*c*x*log(c^2*x^2 + 1))/2) + a*e^2*x)/(d*(e^2 + c^2*d^2)*(d + e*x))
```

3.7 $\int \frac{a+b\text{ArcTan}(cx)}{(d+ex)^3} dx$

Optimal. Leaf size=146

$$-\frac{bc}{2(c^2d^2 + e^2)(d+ex)} + \frac{bc^2(cd-e)(cd+e)\text{ArcTan}(cx)}{2e(c^2d^2 + e^2)^2} - \frac{a+b\text{ArcTan}(cx)}{2e(d+ex)^2} + \frac{bc^3d\log(d+ex)}{(c^2d^2 + e^2)^2} - \frac{bc^3d\log(1+e^2)}{2(c^2d^2 + e^2)}$$

[Out] $-1/2*b*c/(c^2*d^2 + e^2)/(e*x + d) + 1/2*b*c^2*(c*d - e)*(c*d + e)*\arctan(c*x)/e/(c^2*d^2 + e^2)^2 + 1/2*(-a - b*\arctan(c*x))/e/(e*x + d)^2 + b*c^3*d*\ln(e*x + d)/(c^2*d^2 + e^2)^2 - 1/2*b*c^3*d*\ln(c^2*x^2 + 1)/(c^2*d^2 + e^2)^2$

Rubi [A]

time = 0.09, antiderivative size = 146, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.375, Rules used = {4972, 724, 815, 649, 209, 266}

$$-\frac{a+b\text{ArcTan}(cx)}{2e(d+ex)^2} + \frac{bc^2\text{ArcTan}(cx)(cd-e)(cd+e)}{2e(c^2d^2 + e^2)^2} - \frac{bc}{2(c^2d^2 + e^2)(d+ex)} - \frac{bc^3d\log(c^2x^2 + 1)}{2(c^2d^2 + e^2)^2} + \frac{bc^3d\log(d+ex)}{(c^2d^2 + e^2)^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x])/(d + e*x)^3, x]$

[Out] $-1/2*(b*c)/((c^2*d^2 + e^2)*(d + e*x)) + (b*c^2*(c*d - e)*(c*d + e))*\text{ArcTan}[c*x]/(2*e*(c^2*d^2 + e^2)^2) - (a + b*\text{ArcTan}[c*x])/((2*e*(d + e*x)^2) + (b*c^3*d*\log[d + e*x])/(c^2*d^2 + e^2)^2 - (b*c^3*d*\log[1 + c^2*x^2])/(2*(c^2*d^2 + e^2)^2)$

Rule 209

$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \text{ || } \text{GtQ}[b, 0])$

Rule 266

$\text{Int}[(x_)^{(m_)} / ((a_) + (b_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}[\{a, b, m, n\}, x] \&& \text{EqQ}[m, n - 1]$

Rule 649

$\text{Int}[(d_) + (e_)*(x_)/((a_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Dist}[d, \text{Int}[1/(a + c*x^2), x], x] + \text{Dist}[e, \text{Int}[x/(a + c*x^2), x], x] /; \text{FreeQ}[\{a, c, d, e\}, x] \&& \text{!NiceSqrtQ}[(-a)*c]$

Rule 724

```
Int[((d_) + (e_)*(x_))^(m_)/((a_) + (c_)*(x_)^2), x_Symbol] :> Simp[e*((d + e*x)^(m + 1)/((m + 1)*(c*d^2 + a*e^2))), x] + Dist[c/(c*d^2 + a*e^2), Int[(d + e*x)^(m + 1)*((d - e*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[m, -1]
```

Rule 815

```
Int[((d_) + (e_)*(x_))^(m_)*((f_.) + (g_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]
```

Rule 4972

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))*((d_) + (e_)*(x_))^(q_), x_Symbol] :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{a + b \tan^{-1}(cx)}{(d + ex)^3} dx &= -\frac{a + b \tan^{-1}(cx)}{2e(d + ex)^2} + \frac{(bc) \int \frac{1}{(d+ex)^2(1+c^2x^2)} dx}{2e} \\
&= -\frac{bc}{2(c^2d^2 + e^2)(d + ex)} - \frac{a + b \tan^{-1}(cx)}{2e(d + ex)^2} + \frac{(bc^3) \int \frac{d-ex}{(d+ex)(1+c^2x^2)} dx}{2e(c^2d^2 + e^2)} \\
&= -\frac{bc}{2(c^2d^2 + e^2)(d + ex)} - \frac{a + b \tan^{-1}(cx)}{2e(d + ex)^2} + \frac{(bc^3) \int \left(\frac{2de^2}{(c^2d^2 + e^2)(d+ex)} + \frac{c^2d^2 - e^2 - 2c^2d}{(c^2d^2 + e^2)(1+c^2x^2)} \right) dx}{2e(c^2d^2 + e^2)} \\
&= -\frac{bc}{2(c^2d^2 + e^2)(d + ex)} - \frac{a + b \tan^{-1}(cx)}{2e(d + ex)^2} + \frac{bc^3 d \log(d + ex)}{(c^2d^2 + e^2)^2} + \frac{(bc^3) \int \frac{c^2d^2 - e^2 - 2c^2d}{1+c^2x^2} dx}{2e(c^2d^2 + e^2)} \\
&= -\frac{bc}{2(c^2d^2 + e^2)(d + ex)} - \frac{a + b \tan^{-1}(cx)}{2e(d + ex)^2} + \frac{bc^3 d \log(d + ex)}{(c^2d^2 + e^2)^2} - \frac{(bc^5 d) \int \frac{x}{1+c^2x^2} dx}{(c^2d^2 + e^2)^2} \\
&= -\frac{bc}{2(c^2d^2 + e^2)(d + ex)} + \frac{bc^2(cd - e)(cd + e) \tan^{-1}(cx)}{2e(c^2d^2 + e^2)^2} - \frac{a + b \tan^{-1}(cx)}{2e(d + ex)^2} + \frac{bc^3 d}{(c^2d^2 + e^2)^2}
\end{aligned}$$

Mathematica [A]

time = 0.22, size = 192, normalized size = 1.32

$$-\frac{2(a + b \text{ArcTan}(cx)) + \frac{bc(d+ex) \left(2e(c^2d^2 + e^2) - \left(c^2d \left(\sqrt{-c^2} d - 2e\right) - \sqrt{-c^2} e^2\right) (d+ex) \log \left(1 - \sqrt{-c^2} x\right) - \left(\sqrt{-c^2} e^2 - c^2 d \left(\sqrt{-c^2} d + 2e\right)\right) (d+ex) \log \left(1 + \sqrt{-c^2} x\right) - 4c^2 d e (d+ex) \log (d+ex)\right)}{(c^2d^2 + e^2)^2}}{4e(d + ex)^2}$$

Antiderivative was successfully verified.

[In] `Integrate[(a + b*ArcTan[c*x])/(d + e*x)^3, x]`

[Out]
$$\begin{aligned} & -\frac{1}{4} \cdot \frac{2(a + b \operatorname{ArcTan}[c x]) + (b c (d + e x) \cdot (2 e (c^2 d^2 + e^2) - (c^2 d^2 + e^2) \operatorname{Log}[1 - \sqrt{-c^2} x] - (\sqrt{-c^2} e^2 - c^2 d (\sqrt{-c^2} d + 2 e)) \cdot (d + e x) \operatorname{Log}[1 + \sqrt{-c^2} x] - 4 c^2 d e (d + e x) \operatorname{Log}[d + e x]))}{(c^2 d^2 + e^2)^2} \\ & \cdot (e (d + e x)^2) \end{aligned}$$

Maple [A]

time = 0.14, size = 188, normalized size = 1.29

method	result
derivativedivides	$\begin{aligned} & -\frac{\frac{a c^3}{2(cex+cd)^2e} - \frac{b c^3 \arctan(cx)}{2(cex+cd)^2e} + \frac{b c^5 \arctan(cx)d^2}{2e(c^2d^2+e^2)^2} - \frac{b c^4 d \ln(c^2x^2+1)}{2(c^2d^2+e^2)^2} - \frac{b c^3 e \arctan(cx)}{2(c^2d^2+e^2)^2} - \frac{b c^3}{2(c^2d^2+e^2)(cex+cd)} + \frac{b c^4 d \ln(cex+cd)}{(c^2d^2+e^2)^2}}{c} \\ & -\frac{\frac{a c^3}{2(cex+cd)^2e} - \frac{b c^3 \arctan(cx)}{2(cex+cd)^2e} + \frac{b c^5 \arctan(cx)d^2}{2e(c^2d^2+e^2)^2} - \frac{b c^4 d \ln(c^2x^2+1)}{2(c^2d^2+e^2)^2} - \frac{b c^3 e \arctan(cx)}{2(c^2d^2+e^2)^2} - \frac{b c^3}{2(c^2d^2+e^2)(cex+cd)} + \frac{b c^4 d \ln(cex+cd)}{(c^2d^2+e^2)^2}}{c} \end{aligned}$
default	
risch	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x))/(e*x+d)^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 1/c * (-1/2*a*c^3/(c*e*x+c*d)^2/e - 1/2*b*c^3/(c*e*x+c*d)^2/e*\arctan(cx) + 1/2*b*c^5/e/(c^2*d^2+e^2)^2*\arctan(cx)*d^2 - 1/2*b*c^4/(c^2*d^2+e^2)^2*d*\ln(c^2*x^2+1) - 1/2*b*c^3/e/(c^2*d^2+e^2)^2*\arctan(cx) - 1/2*b*c^3/(c^2*d^2+e^2)/(c*e*x+c*d) + b*c^4*d/(c^2*d^2+e^2)^2*\ln(c*e*x+c*d)) \end{aligned}$$

Maxima [A]

time = 0.48, size = 206, normalized size = 1.41

$$-\frac{1}{2} \left(\left(\frac{c^2 d \log(c^2 x^2 + 1)}{c^4 d^4 + 2 c^2 d^2 e^2 + e^4} - \frac{2 c^2 d \log(xe + d)}{c^4 d^4 + 2 c^2 d^2 e^2 + e^4} - \frac{(c^4 d^2 - c^2 e^2) \arctan(cx)}{(c^4 d^4 e + 2 c^2 d^2 e^3 + e^5)c} + \frac{1}{c^2 d^3 + (c^2 d^2 e + e^3)x + de^2} \right) c + \frac{\arctan(cx)}{x^2 e^3 + 2 dxe^2 + d^2 e} \right) b - \frac{a}{2(x^2 e^3 + 2 dxe^2 + d^2 e)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))/(e*x+d)^3,x, algorithm="maxima")`

[Out]
$$\begin{aligned} & -1/2*((c^2*d*log(c^2*x^2 + 1)/(c^4*d^4 + 2*c^2*d^2*e^2 + e^4) - 2*c^2*d*log(xe + d)/(c^4*d^4 + 2*c^2*d^2*e^2 + e^4) - (c^4*d^2 - c^2*e^2)*\arctan(cx))/((c^4*d^4*e + 2*c^2*d^2*e^3 + e^5)*c) + 1/(c^2*d^3 + (c^2*d^2*e + e^3)*x + d*e^2)*c + \arctan(cx)/(x^2*e^3 + 2*d*x*e^2 + d^2*e))*b - 1/2*a/(x^2*e^3 + 2*d*x*e^2 + d^2*e) \end{aligned}$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 299 vs. 2(137) = 274.

time = 1.36, size = 299, normalized size = 2.05

$$\frac{a c^4 d^4 + b c^3 d^3 e + b c d e^3 - (2 b c^4 d^2 x e - 2 b c^3 d x e^2 - (b c^2 x^2 + b) e^4 + (b c^4 d^2 x^2 - 3 b c^2 d^2) e^2) \arctan(cx) + (b c x + a) e^4 + (b c^3 d^2 x + 2 a c^2 d^2) e^2 + (b c^3 d x^2 + 2 b c^2 d^2 x e^3 + 2 b c^2 d^2 x e^2 + b c^2 d^2 e) \log(c^2 x^2 + 1) - 2(b c^3 d x^2 e^3 + 2 b c^3 d^2 x e^2 + b c^3 d^2 e) \log(xe + d)}{2(2 c^4 d^3 x e^2 + c^4 d^4 e + 4 c^2 d^3 x e^4 + x^2 e^4 + 2 d x e^6 + (2 c^2 d^2 x^2 + d^2) e^5 + (c^4 d^4 x^2 + 2 c^2 d^4) e^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))/(e*x+d)^3,x, algorithm="fricas")
[Out] -1/2*(a*c^4*d^4 + b*c^3*d^3*e + b*c*d*e^3 - (2*b*c^4*d^3*x*e - 2*b*c^2*d*x*
e^3 - (b*c^2*x^2 + b)*e^4 + (b*c^4*d^2*x^2 - 3*b*c^2*d^2)*e^2)*arctan(c*x)
+ (b*c*x + a)*e^4 + (b*c^3*d^2*x + 2*a*c^2*d^2)*e^2 + (b*c^3*d*x^2*e^3 + 2*
b*c^3*d^2*x*x^2 + b*c^3*d^3*e)*log(c^2*x^2 + 1) - 2*(b*c^3*d*x^2*e^3 + 2*b*
c^3*d^2*x*x^2 + b*c^3*d^3*e)*log(x*x + d))/(2*c^4*d^5*x*x^2 + c^4*d^6*e + 4*
c^2*d^3*x*x^4 + x^2*e^7 + 2*d*x*x^6 + (2*c^2*d^2*x^2 + d^2)*e^5 + (c^4*d^4*
x^2 + 2*c^2*d^4)*e^3)
```

Sympy [C] Result contains complex when optimal does not.

time = 3.42, size = 2866, normalized size = 19.63

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*atan(c*x))/(e*x+d)**3,x)
```

```
[Out] Piecewise((a*x/d**3, Eq(c, 0) & Eq(e, 0)), (-4*a*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + 3*I*b*d**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + 2*I*b*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - 2*I*b*d*x*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - I*b*e**2*x**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2), Eq(c, -I*e/d)), (-4*a*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - 3*I*b*d**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - 2*I*b*d**2/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + 2*I*b*d*x*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) - I*b*d*x/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2) + I*b*e**2*x**2*atanh(e*x/d)/(8*d**4*e + 16*d**3*e**2*x + 8*d**2*e**3*x**2), Eq(c, I*e/d)), ((a*x + b*x*atan(c*x) - b*log(x**2 + c*(-2))/(2*c))/d**3, Eq(e, 0)), (-a/(2*d**2*e + 4*d*e**2*x + 2*e**3*x**2), Eq(c, 0)), (-a*c**4*d**4/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*x**2 + 2*c**4*d**5*x**2 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) - 2*a*c**2*d**2*x**2/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*x**2 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*x**5 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) - a*e**4/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*x**2 + 4*c**2*d**4*x**3 + 8*c**2*d**3*x**4*x + 4*c**2*d**2*x**5*x**2 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) + 4*c**2*d**3*x**2 + 4*c**2*d**4*x**3 + 8*c**2*d**3*x**4*x + 4*c**2*d**2*x**5*x**2 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) + 2*b*c**4*d**3*x*x*atan(c*x)/(2*c**4*d**6*e + 4*c**4*d**5*x**2 + 2*c**4*d**4*x**3 + 4*c**2*d**4*x**4*x + 8*c**2*d**3*x**5*x**2 + 2*c**2*d**2*x**6*x + 2*c**4*d**5*x**2 + 4*c**2*d**4*x**3 + 8*c**2*d**3*x**4*x + 4*c**2*d**2*x**5*x**2 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) + b*c**4*d**2*x*x**2*atan(c*x)/(2*c**4*d**6*e + 4*c**4*d**5*x**2 + 2*c**4*d**4*x**3 + 4*c**2*d**4*x**4*x + 8*c**2*d**3*x**5*x**2 + 2*c**2*d**2*x**6*x + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) - b*c**3*d**3*x*log(x**2 + c*(-2))/(2*c**4*d**6*e + 4*c**4*d**5*x**2 + 2*c**4*d**4*x**3 + 4*c**2*d**4*x**4*x + 8*c**2*d**3*x**5*x**2 + 4*c**2*d**2*x**6*x + 4*c**2*d**5*x**2 + 4*c**2*d**4*x**3 + 8*c**2*d**3*x**4*x + 4*c**2*d**2*x**5*x**2 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2) - b*c**3*d**2*x**2 + 4*c**2*d**4*x**3 + 8*c**2*d**3*x**4*x + 4*c**2*d**2*x**5*x**2 + 2*d**2*x**5 + 4*d*x**6*x + 2*e**7*x**2)
```

$$\begin{aligned}
& 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) + 2*b*c**3*d**3*e*log(d/e + x)/(2*c \\
& **4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 \\
& + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2 \\
& *e**7*x**2) - b*c**3*d**3*e/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d \\
& **4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x** \\
& 2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) - 2*b*c**3*d**2*e**2*x*log(x**2 \\
& + c*(-2))/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4 \\
& *c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 \\
& + 4*d*e**6*x + 2*e**7*x**2) + 4*b*c**3*d**2*e**2*x*log(d/e + x)/(2*c**4*d** \\
& 6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c** \\
& 2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x \\
& **2) - b*c**3*d**2*e**2*x/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4 \\
& *e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 \\
& + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) - b*c**3*d**3*x**2*log(x**2 + c \\
&*(-2))/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c** \\
& 2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4* \\
&d*e**6*x + 2*e**7*x**2) + 2*b*c**3*d**3*x**2*log(d/e + x)/(2*c**4*d**6*e \\
& + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d* \\
& *3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) \\
& - 3*b*c**2*d**2*e**2*atan(c*x)/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c** \\
& 4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5 \\
& *x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) - 2*b*c**2*d**3*x*atan(c* \\
&x)/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d** \\
& 4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e** \\
& 6*x + 2*e**7*x**2) - b*c**2*e**4*x**2*atan(c*x)/(2*c**4*d**6*e + 4*c**4*d** \\
& 5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + \\
& 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) - b*c*d*e** \\
& 3/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4 \\
& *e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6 \\
& *x + 2*e**7*x**2) - b*c*e**4*x/(2*c**4*d**6*e + 4*c**4*d**5*e**2*x + 2*c**4 \\
& *d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c**2*d**3*e**4*x + 4*c**2*d**2*e**5*x \\
& **2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7*x**2) - b*e**4*atan(c*x)/(2*c**4*d \\
& **6*e + 4*c**4*d**5*e**2*x + 2*c**4*d**4*e**3*x**2 + 4*c**2*d**4*e**3 + 8*c \\
& **2*d**3*e**4*x + 4*c**2*d**2*e**5*x**2 + 2*d**2*e**5 + 4*d*e**6*x + 2*e**7 \\
& *x**2), \text{True})
\end{aligned}$$
Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*arctan(c*x))/(e*x+d)^3,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 4.43, size = 591, normalized size = 4.05

$$\frac{c \left(e^{i \pi} d^2 + \frac{b d^2 e^{i \pi}}{c} + e^{i \pi} c^2 \right)}{d \left(c^2 e^{i \pi} + d^2 e^{i \pi} \right)} - \frac{\ln \left(c^2 e^{i \pi} \right)}{2 c} + \frac{e^2 \left(b c^2 d^2 e^{i \pi} + b d^2 e^{i \pi} + b^2 e^{i \pi} \right)}{c^2 d^2 e^{i \pi} + 2 c^2 d e^{i \pi} + c^2 e^{i \pi} x^2 + d^2 + 2 d e^{i \pi} x + e^{i \pi} x^2} + \frac{e^{i \pi} \left(b c^2 d^2 e^{i \pi} + b d^2 e^{i \pi} + b^2 e^{i \pi} \right)}{c^2 d^2 e^{i \pi} + 2 c^2 d e^{i \pi} + c^2} - \frac{b c^2 d \ln(d + e x)}{c^2 d^2 + 2 c^2 d^2 e^{i \pi} + e^2} - \frac{b c^2 d \ln(c^2 x^2 + 1)}{2 \left(c^4 d^4 + 2 c^2 d^2 e^{i \pi} + e^4 \right)} + \frac{\operatorname{atan}\left(\frac{c^2 d}{\sqrt{c^2 e^{i \pi}}}\right) \left(c^2\right)^{5/2} \left(e^4 d^4 + 8 e^2 d^2 e^2 + 2 e^4\right) \left(3 c^4 d^4 + 26 c^2 d^2 e^2 + 4 e^2 e^4\right) \left(27 b c^{10} d^{10} + 23 b c^8 d^8 e^2 - 34 b c^6 d^6 e^4 - 26 b c^4 d^4 e^6 + 7 b c^2 d^2 e^8 + 3 b e^{10}\right)}{2 c \left(81 c^{20} d^{20} e + 1662 c^{18} d^{18} e^2 + 11515 c^{16} d^{16} e^4 + 32306 c^{14} d^{14} e^6 + 43705 c^{12} d^{12} e^8 + 28142 c^{10} d^{10} e^{10} + 4857 c^{14} d^8 e^{12} - 3650 c^{12} d^6 e^{14} - 2054 c^{10} d^4 e^{16} - 380 c^8 d^2 e^{18} - 24 e^2 e^{20}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))/(d + e*x)^3,x)`

[Out] $\frac{((x*(a*e^2 + a*c^2*d^2 + (b*c*d*e)/2))/(d*(e^2 + c^2*d^2)) - (b*atan(c*x))/(2*e) + (x^2*((a*e^3)/2 + (b*c*d*e^2)/2 + (a*c^2*d^2*e)/2))/(d^2*(e^2 + c^2*d^2)) + (x^4*((a*c^2*e^3)/2 + (a*c^4*d^2*e)/2 + (b*c^3*d*e^2)/2))/(d^2*(e^2 + c^2*d^2)) + (x^3*((a*c^4*d^2 + a*c^2*e^2 + (b*c^3*d*e)/2))/(d*(e^2 + c^2*d^2)) - (b*c^2*x^2*atan(c*x))/(2*e))/(d^2 + e^2*x^2 + 2*d*e*x + c^2*d^2*x^2 + c^2*e^2*x^4 + 2*c^2*d*e*x^3) + (b*c^3*d*log(d + e*x))/(e^4 + c^4*d^4 + 2*c^2*d^2*e^2) - (b*c^3*d*log(c^2*x^2 + 1))/(2*(e^4 + c^4*d^4 + 2*c^2*d^2*e^2)) + (\operatorname{atan}(c^2*x)/(c^2)^{(1/2)})*(c^2)^{(7/2)}*(2*e^4 + c^4*d^4 + 8*c^2*d^2*e^2)*(3*c^6*d^4 + 4*c^2*e^4 + 26*c^4*d^2*e^2)*(3*b*c^10 + 27*b*c^10*d^10 + 7*b*c^2*d^2*e^8 - 26*b*c^4*d^4*e^6 - 34*b*c^6*d^6*e^4 + 23*b*c^8*d^8*e^2))/(2*c*(81*c^26*d^20*e - 24*c^6*e^21 - 380*c^8*d^2*e^19 - 2054*c^10*d^4*e^17 - 3650*c^12*d^6*e^15 + 4857*c^14*d^8*e^13 + 28142*c^16*d^10*e^11 + 43705*c^18*d^12*e^9 + 32306*c^20*d^14*e^7 + 11515*c^22*d^16*e^5 + 1662*c^24*d^18*e^3))$

3.8 $\int \frac{a+b\text{ArcTan}(cx)}{(d+ex)^4} dx$

Optimal. Leaf size=206

$$-\frac{bc}{6(c^2d^2 + e^2)(d+ex)^2} - \frac{2bc^3d}{3(c^2d^2 + e^2)^2(d+ex)} + \frac{bc^4d(c^2d^2 - 3e^2)\text{ArcTan}(cx)}{3e(c^2d^2 + e^2)^3} - \frac{a+b\text{ArcTan}(cx)}{3e(d+ex)^3} + \frac{bc^3(3c^2d^2 - 3e^2)\log(d+ex)}{3(c^2d^2 + e^2)^3}$$

[Out] $-1/6*b*c/(c^2*d^2 + e^2)/(e*x + d)^2 - 2/3*b*c^3*d/(c^2*d^2 + e^2)^2/(e*x + d) + 1/3*b*c^4*d*(c^2*d^2 - 3e^2)*\text{arctan}(c*x)/e/(c^2*d^2 + e^2)^3 + 1/3*(-a - b*\text{arctan}(c*x))/e/(e*x + d)^3 + 1/3*b*c^3*(3*c^2*d^2 - e^2)*\ln(e*x + d)/(c^2*d^2 + e^2)^3 - 1/6*b*c^3*(3*c^2*d^2 - e^2)*\ln(c^2*x^2 + 1)/(c^2*d^2 + e^2)^3$

Rubi [A]

time = 0.13, antiderivative size = 206, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {4972, 724, 815, 649, 209, 266}

$$-\frac{a+b\text{ArcTan}(cx)}{3e(d+ex)^3} + \frac{bc^4d\text{ArcTan}(cx)(c^2d^2 - 3e^2)}{3e(c^2d^2 + e^2)^3} - \frac{bc}{6(c^2d^2 + e^2)(d+ex)^2} - \frac{bc^3(3c^2d^2 - e^2)\log(c^2x^2 + 1)}{6(c^2d^2 + e^2)^3} - \frac{2bc^3d}{3(c^2d^2 + e^2)^2(d+ex)} + \frac{bc^3(3c^2d^2 - e^2)\log(d+ex)}{3(c^2d^2 + e^2)^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x])/(d + e*x)^4, x]$

[Out] $-1/6*(b*c)/((c^2*d^2 + e^2)*(d + e*x)^2) - (2*b*c^3*d)/(3*(c^2*d^2 + e^2)^2*(d + e*x)) + (b*c^4*d*(c^2*d^2 - 3e^2)*\text{ArcTan}[c*x])/((3*e*(c^2*d^2 + e^2)^3) - (a + b*\text{ArcTan}[c*x])/((3*e*(d + e*x)^3) + (b*c^3*(3*c^2*d^2 - e^2)*\text{Log}[d + e*x])/((3*(c^2*d^2 + e^2)^3) - (b*c^3*(3*c^2*d^2 - e^2)*\text{Log}[1 + c^2*x^2]))/(6*(c^2*d^2 + e^2)^3)$

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 649

```
Int[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 724

```
Int[((d_) + (e_)*(x_))^(m_)/((a_) + (c_)*(x_)^2), x_Symbol] :> Simp[e*((d + e*x)^(m + 1)/((m + 1)*(c*d^2 + a*e^2))), x] + Dist[c/(c*d^2 + a*e^2), In t[(d + e*x)^(m + 1)*((d - e*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, m}, x] && NeQ[c*d^2 + a*e^2, 0] && LtQ[m, -1]
```

Rule 815

```
Int[((((d_) + (e_)*(x_))^(m_)*(f_*) + (g_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(d + e*x)^m*((f + g*x)/(a + c*x^2)), x], x] /; FreeQ[{a, c, d, e, f, g}, x] && NeQ[c*d^2 + a*e^2, 0] && IntegerQ[m]
```

Rule 4972

```
Int[((a_*) + ArcTan[(c_)*(x_)]*(b_))*((d_ + (e_)*(x_))^(q_), x_Symbol] :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/(e*(q + 1))), x] - Dist[b*(c/(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e, q}, x] && NeQ[q, -1]
```

Rubi steps

$$\begin{aligned}
\int \frac{a + b \tan^{-1}(cx)}{(d + ex)^4} dx &= -\frac{a + b \tan^{-1}(cx)}{3e(d + ex)^3} + \frac{(bc) \int \frac{1}{(d+ex)^3(1+c^2x^2)} dx}{3e} \\
&= -\frac{bc}{6(c^2d^2 + e^2)(d + ex)^2} - \frac{a + b \tan^{-1}(cx)}{3e(d + ex)^3} + \frac{(bc^3) \int \frac{d-ex}{(d+ex)^2(1+c^2x^2)} dx}{3e(c^2d^2 + e^2)} \\
&= -\frac{bc}{6(c^2d^2 + e^2)(d + ex)^2} - \frac{a + b \tan^{-1}(cx)}{3e(d + ex)^3} + \frac{(bc^3) \int \left(\frac{2de^2}{(c^2d^2 + e^2)(d+ex)^2} - \frac{e^2(-3c^2d^2 + 2c^2d^2 + e^2)}{(c^2d^2 + e^2)^2(d+ex)} \right) dx}{3e(c^2d^2 + e^2)} \\
&= -\frac{bc}{6(c^2d^2 + e^2)(d + ex)^2} - \frac{2bc^3d}{3(c^2d^2 + e^2)^2(d + ex)} - \frac{a + b \tan^{-1}(cx)}{3e(d + ex)^3} + \frac{bc^3(3c^2d^2 - 2c^2d^2 - e^2)}{3(c^2d^2 + e^2)} \\
&= -\frac{bc}{6(c^2d^2 + e^2)(d + ex)^2} - \frac{2bc^3d}{3(c^2d^2 + e^2)^2(d + ex)} - \frac{a + b \tan^{-1}(cx)}{3e(d + ex)^3} + \frac{bc^3(3c^2d^2 - 2c^2d^2 - e^2)}{3(c^2d^2 + e^2)} \\
&= -\frac{bc}{6(c^2d^2 + e^2)(d + ex)^2} - \frac{2bc^3d}{3(c^2d^2 + e^2)^2(d + ex)} + \frac{bc^4d(c^2d^2 - 3e^2)\tan^{-1}(cx)}{3e(c^2d^2 + e^2)^3}
\end{aligned}$$

Mathematica [A]

time = 0.43, size = 254, normalized size = 1.23

$$\frac{2(a + b \text{ArcTan}(cx)) + \frac{bc(d+ex) \left(e(c^2d^2 + e^2)^2 + 4c^2de(c^2d^2 + e^2)(d+ex) - c^2 \left(c^2d^2 \left(\sqrt{-c^2} d - 3e\right) + c^2 \left(-3\sqrt{-c^2} d + e\right)\right) (d+ex)^2 \log \left(1 - \sqrt{-c^2} x\right) - c^2 \left(e^2 \left(3\sqrt{-c^2} d + e\right) - c^2 d^2 \left(\sqrt{-c^2} d + 3e\right)\right) (d+ex)^2 \log \left(1 + \sqrt{-c^2} x\right) - 2c^2 e (3c^2d^2 - c^2) (d+ex)^2 \log (d+ex)}{(c^2d^2 + e^2)^3}}{6e(d + ex)^3}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + b*ArcTan[c*x])/(d + e*x)^4, x]
[Out] -1/6*(2*(a + b*ArcTan[c*x]) + (b*c*(d + e*x)*(e*(c^2*d^2 + e^2)^2 + 4*c^2*d^2)*e*(c^2*d^2 + e^2)*(d + e*x) - c^2*(c^2*d^2*(Sqrt[-c^2]*d - 3*e) + e^2*(-3*Sqrt[-c^2]*d + e)))*(d + e*x)^2*Log[1 - Sqrt[-c^2]*x] - c^2*(e^2*(3*Sqrt[-c^2]*d + e) - c^2*d^2*(Sqrt[-c^2]*d + 3*e))*(d + e*x)^2*Log[1 + Sqrt[-c^2]*x] - 2*c^2*e*(3*c^2*d^2 - e^2)*(d + e*x)^2*Log[d + e*x]))/(c^2*d^2 + e^2)^3)/(e*(d + e*x)^3)
```

Maple [A]

time = 0.20, size = 286, normalized size = 1.39

method	result
derivativedivides	$\frac{-\frac{a c^4}{3(cex+cd)^3 e} - \frac{b c^4 \arctan(cx)}{3(cex+cd)^3 e} - \frac{b c^6 \ln(c^2 x^2 + 1) d^2}{2(c^2 d^2 + e^2)^3} + \frac{b c^4 e^2 \ln(c^2 x^2 + 1)}{6(c^2 d^2 + e^2)^3} + \frac{b c^7 \arctan(cx) d^3}{3e(c^2 d^2 + e^2)^3} - \frac{b c^5 e \arctan(cx) d}{(c^2 d^2 + e^2)^3} - \frac{b c^4}{6(c^2 d^2 + e^2)(cex+cd)^3}}{c}$
default	$\frac{-\frac{a c^4}{3(cex+cd)^3 e} - \frac{b c^4 \arctan(cx)}{3(cex+cd)^3 e} - \frac{b c^6 \ln(c^2 x^2 + 1) d^2}{2(c^2 d^2 + e^2)^3} + \frac{b c^4 e^2 \ln(c^2 x^2 + 1)}{6(c^2 d^2 + e^2)^3} + \frac{b c^7 \arctan(cx) d^3}{3e(c^2 d^2 + e^2)^3} - \frac{b c^5 e \arctan(cx) d}{(c^2 d^2 + e^2)^3} - \frac{b c^4}{6(c^2 d^2 + e^2)(cex+cd)^3}}{c}$
risch	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a+b*arctan(c*x))/(e*x+d)^4,x,method=_RETURNVERBOSE)
[Out] 1/c*(-1/3*a*c^4/(c*e*x+c*d)^3/e-1/3*b*c^4/(c*e*x+c*d)^3/e*arctan(c*x)-1/2*b*c^6/(c^2*d^2+e^2)^3*ln(c^2*x^2+1)*d^2+1/6*b*c^4*e^2/(c^2*d^2+e^2)^3*ln(c^2*x^2+1)+1/3*b*c^7/e/(c^2*d^2+e^2)^3*arctan(c*x)*d^3-b*c^5*e/(c^2*d^2+e^2)^3*arctan(c*x)*d-1/6*b*c^4/(c^2*d^2+e^2)/(c*e*x+c*d)^2+b*c^6/(c^2*d^2+e^2)^3*ln(c*e*x+c*d)*d^2-1/3*b*c^4*e^2/(c^2*d^2+e^2)^3*ln(c*e*x+c*d)-2/3*b*c^5*d/(c^2*d^2+e^2)^2/(c*e*x+c*d))
```

Maxima [A]

time = 0.49, size = 354, normalized size = 1.72

$$-\frac{1}{6} \left(c \left(\frac{(3 c^4 d^2 - c^2 e^2) \log(c^2 x^2 + 1)}{c^6 d^6 + 3 c^4 d^4 e^2 + 3 c^2 d^2 e^4 + e^6} - \frac{2 (3 c^4 d^2 - c^2 e^2) \log(xe + d)}{c^6 d^6 + 3 c^4 d^4 e^2 + 3 c^2 d^2 e^4 + e^6} + \frac{4 c^2 dxe + 5 c^2 d^2 + e^2}{c^6 d^6 + 2 c^2 d^4 e^2 + (c^4 d^2 e^2 + 2 c^2 d^2 e^4 + e^6)x^2 + d^2 e^4 + 2 (c^4 d^2 e + 2 c^2 d^2 e^3 + de^5)x} - \frac{2 (c^6 d^3 - 3 c^4 d e^2) \arctan(cx)}{(c^6 d^6 e + 3 c^4 d^4 e^3 + 3 c^2 d^2 e^5 + e^7)c} \right) + \frac{2 \arctan(cx)}{x^2 e^4 + 3 dx^2 e^3 + 3 d^2 x e^2 + d^3 e} \right) b - \frac{a}{3 (x^2 e^4 + 3 dx^2 e^3 + 3 d^2 x e^2 + d^3 e)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))/(e*x+d)^4,x, algorithm="maxima")
[Out] -1/6*(c*((3*c^4*d^2 - c^2*e^2)*log(c^2*x^2 + 1)/(c^6*d^6 + 3*c^4*d^4*e^2 + 3*c^2*d^2*e^4 + e^6) - 2*(3*c^4*d^2 - c^2*e^2)*log(x*e + d)/(c^6*d^6 + 3*c^4*d^4*e^2 + 3*c^2*d^2*e^4 + e^6) + (4*c^2*d*x*e + 5*c^2*d^2 + e^2)/(c^4*d^6 + 2*c^2*d^4*e^2 + (c^4*d^2 + 2*c^2*d^2*e^4 + e^6)*x^2 + d^2*e^4 + 2*c^2*d^2*e^3 + de^5)*x - 2*(c^6*d^3 - 3*c^4*d*e^2)*arctan(c*x))/((c^6*d^6*e + 3*c^4*d^4*e^3 + 3*c^2*d^2*e^5 + e^7)*c) + 2*arctan(c*x)/(x
```

$$\begin{aligned} & -3e^4 + 3d*x^2*e^3 + 3d^2*x*e^2 + d^3*e) * b - 1/3*a/(x^3*e^4 + 3d*x^2*e^3 + 3d^2*x*e^2 + d^3*e) \end{aligned}$$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 620 vs. $2(188) = 376$.

time = 1.99, size = 620, normalized size = 3.01

$$\begin{aligned} & 2(a^2d^2 - 5ab^2d^2 - 2c^2b^2d^2 - 3b^2cd^2 + b^4d^2 - 3b^2d^2a^2 + b^2cd^2a^2 - 9b^2d^2c^2) + 3(c^2b^2d^2 - 2b^2cd^2a^2) \text{atan}(ac) + (bc + 2ab^2d^2 + 4b^2cd^2a^2 - 10b^2d^2c^2) + 2(b^2d^2a^2 - 3b^2cd^2a^2 - 2(b^2d^2a^2 + 2cd^2a^2b^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) + b^2d^2a^2 - 2b^2cd^2a^2) + 2(b^2d^2a^2 + 2cd^2a^2b^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) + (9b^2d^2a^2 - bc^2d^2a^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) + 2(b^2d^2a^2 - bc^2d^2a^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) \\ & + 6(3b^2d^2a^2 + 2cd^2a^2b^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) + 3(b^2d^2a^2 + 2cd^2a^2b^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) + 3(b^2d^2a^2 + 2cd^2a^2b^2) \ln(b^2d^2a^2 + 2cd^2a^2b^2) \end{aligned}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))/(e*x+d)^4,x, algorithm="fricas")
```

```
[Out] -1/6*(2*a*c^6*d^6 + 5*b*c^5*d^5*e - 2*(3*b*c^6*d^5*x*e - 3*b*c^4*d*x^3*e^5 - b*e^6 - 3*(3*b*c^4*d^2*x^2 + b*c^2*d^2)*e^4 + (b*c^6*d^3*x^3 - 9*b*c^4*d^3*x)*e^3 + 3*(b*c^6*d^4*x^2 - 2*b*c^4*d^4)*e^2)*arctan(c*x) + (b*c*x + 2*a)*e^6 + (4*b*c^3*d*x^2 + b*c*d)*e^5 + 2*(5*b*c^3*d^2*x + 3*a*c^2*d^2)*e^4 + 2*(2*b*c^5*d^3*x^2 + 3*b*c^3*d^3)*e^3 + 3*(3*b*c^5*d^4*x + 2*a*c^4*d^4)*e^2 + (9*b*c^5*d^4*x*e^2 + 3*b*c^5*d^5*e - b*c^3*x^3*e^6 - 3*b*c^3*d*x^2*e^5 + 3*(b*c^5*d^2*x^3 - b*c^3*d^2*x)*e^4 + (9*b*c^5*d^3*x^2 - b*c^3*d^3)*e^3)*log(c^2*x^2 + 1) - 2*(9*b*c^5*d^4*x*e^2 + 3*b*c^5*d^5*e - b*c^3*x^3*e^6 - 3*b*c^3*d*x^2*e^5 + 3*(b*c^5*d^2*x^3 - b*c^3*d^2*x)*e^4 + (9*b*c^5*d^3*x^2 - b*c^3*d^3)*e^3)*log(x*e + d))/(3*c^6*d^8*x*e^2 + c^6*d^9*e + x^3*e^10 + 3*d*x^2*e^9 + 3*(c^2*d^2*x^3 + d^2*x)*e^8 + (9*c^2*d^3*x^2 + d^3)*e^7 + 3*(c^4*d^4*x^3 + 3*c^2*d^4*x)*e^6 + 3*(3*c^4*d^5*x^2 + c^2*d^5)*e^5 + (c^6*d^6*x^3 + 9*c^4*d^6*x)*e^4 + 3*(c^6*d^7*x^2 + c^4*d^7)*e^3)
```

Sympy [C] Result contains complex when optimal does not.

time = 7.26, size = 9202, normalized size = 44.67

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x))/(e*x+d)**4,x)`

```
[Out] Piecewise((a*x/d**4, Eq(c, 0) & Eq(e, 0)), (-a/(3*d**3*e + 9*d**2*e**2*x + 9*d**3*x**2 + 3*e**4*x**3), Eq(c, 0)), ((a*x + b*x*atan(c*x) - b*log(x**2 + c**(-2))/(2*c))/d**4, Eq(e, 0)), (-24*a*d**3/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 21*I*b*d**3*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 10*I*b*d**3/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 9*I*b*d**2*e*x*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 9*I*b*d**2*e*x/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 9*I*b*d**2*x**2*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 3*I*b*d**2*x**2/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 3*I*b*e**3*x**3*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3))
```

$$\begin{aligned}
& 6*d**4*e**3*x**2 + 72*d**3*e**4*x**3), \text{Eq}(c, -I*e/d)), (-24*a*d**3/(72*d**6 \\
& *e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 21*I*b*d** \\
& 3*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3* \\
& e**4*x**3) - 10*I*b*d**3/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 \\
& + 72*d**3*e**4*x**3) + 9*I*b*d**2*e*x*atanh(e*x/d)/(72*d**6*e + 216*d**5*e* \\
& *2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) - 9*I*b*d**2*e*x/(72*d**6*e \\
& + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3) + 9*I*b*d*e**2* \\
& x**2*atanh(e*x/d)/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d* \\
& *3*e**4*x**3) - 3*I*b*d*e**2*x**2/(72*d**6*e + 216*d**5*e**2*x + 216*d**4*e \\
& **3*x**2 + 72*d**3*e**4*x**3) + 3*I*b*e**3*x**3*atanh(e*x/d)/(72*d**6*e + 2 \\
& 16*d**5*e**2*x + 216*d**4*e**3*x**2 + 72*d**3*e**4*x**3), \text{Eq}(c, I*e/d)), (- \\
& 2*a*c**6*d**6/(6*c**6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 \\
& + 6*c**6*d**6*e**4*x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c** \\
& 4*d**5*e**5*x**2 + 18*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d** \\
& 4*e**6*x + 54*c**2*d**3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + \\
& 18*d**2*e**8*x + 18*d**9*x**2 + 6*e**10*x**3) - 6*a*c**4*d**4*e**2/(6*c** \\
& 6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c**6*d**6*e**4* \\
& x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5*e**5*x**2 + 1 \\
& 8*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6*x + 54*c**2*d \\
& **3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d**2*e**8*x + 18* \\
& d**9*x**2 + 6*e**10*x**3) - 6*a*c**2*d**2*e**4/(6*c**6*d**9*e + 18*c**6*d \\
& **8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c**6*d**6*e**4*x**3 + 18*c**4*d**7* \\
& e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5*e**5*x**2 + 18*c**4*d**4*e**6*x** \\
& 3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6*x + 54*c**2*d**3*e**7*x**2 + 18*c \\
& **2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d**2*e**8*x + 18*d**9*x**2 + 6*e**1 \\
& 0*x**3) - 2*a*e**6/(6*c**6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3 \\
& *x**2 + 6*c**6*d**6*e**4*x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 5 \\
& 4*c**4*d**5*e**5*x**2 + 18*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c** \\
& 2*d**4*e**6*x + 54*c**2*d**3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e* \\
& *7 + 18*d**2*e**8*x + 18*d**9*x**2 + 6*e**10*x**3) + 6*b*c**6*d**5*e*x*at \\
& an(c*x)/(6*c**6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c \\
& **6*d**6*e**4*x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5* \\
& e**5*x**2 + 18*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6* \\
& x + 54*c**2*d**3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d** \\
& 2*e**8*x + 18*d**9*x**2 + 6*e**10*x**3) + 6*b*c**6*d**4*e**2*x**2*atan(c* \\
& x)/(6*c**6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c**6*d \\
& **6*e**4*x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5*e**5* \\
& x**2 + 18*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6*x + \\
& 54*c**2*d**3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d**2*e** \\
& 8*x + 18*d**9*x**2 + 6*e**10*x**3) + 2*b*c**6*d**3*e**3*x**3*atan(c*x)/(6 \\
& *c**6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c**6*d**6*e \\
& **4*x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5*e**5*x**2 \\
& + 18*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6*x + 54*c \\
& *2*d**3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d**2*e**8*x + \\
& 18*d**9*x**2 + 6*e**10*x**3) - 3*b*c**5*d**5*e*log(x**2 + c*(-2))/(6*c*
\end{aligned}$$

$$\begin{aligned}
& *6*d**9*e + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c**6*d**6*e**4 \\
& *x**3 + 18*c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5*e**5*x**2 + \\
& 18*c**4*d**4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6*x + 54*c**2* \\
& d**3*e**7*x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d**2*e**8*x + 18 \\
& *d**9*x**2 + 6*e**10*x**3) + 6*b*c**5*d**5*e*log(d/e + x)/(6*c**6*d**9*e \\
& + 18*c**6*d**8*e**2*x + 18*c**6*d**7*e**3*x**2 + 6*c**6*d**6*e**4*x**3 + 18 \\
& *c**4*d**7*e**3 + 54*c**4*d**6*e**4*x + 54*c**4*d**5*e**5*x**2 + 18*c**4*d* \\
& *4*e**6*x**3 + 18*c**2*d**5*e**5 + 54*c**2*d**4*e**6*x + 54*c**2*d**3*e**7* \\
& x**2 + 18*c**2*d**2*e**8*x**3 + 6*d**3*e**7 + 18*d**2*e**8*x + 18*d**9*x* \\
& *2 + 6*e**10*x**3) - 5*b*c**5*d**5*e/(6*c**6*d*...
\end{aligned}$$
Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))/(e*x+d)^4,x, algorithm="giac")`[Out] `sage0*x`Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{a + b \operatorname{atan}(cx)}{(d + ex)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))/(d + e*x)^4,x)`[Out] `int((a + b*atan(c*x))/(d + e*x)^4, x)`

$$3.9 \quad \int (d + ex)^3 (a + b \operatorname{ArcTan}(cx))^2 dx$$

Optimal. Leaf size=376

$$\frac{b^2 de^2 x}{c^2} - \frac{abe(6c^2 d^2 - e^2) x}{2c^3} + \frac{b^2 e^3 x^2}{12c^2} - \frac{b^2 de^2 \operatorname{ArcTan}(cx)}{c^3} - \frac{b^2 e(6c^2 d^2 - e^2) x \operatorname{ArcTan}(cx)}{2c^3} - \frac{bde^2 x^2 (a + b \operatorname{ArcTan}(cx))}{c}$$

[Out] $b^{2-d} e^{2-x} / c^{2-1/2} * a * b * e * (6*c^{2-d} 2-e^2) * x / c^{3+1/12} * b^{2-e^3} x^2 / c^{2-b^2-d} * e^{2-arctan(cx)}/c^{3-1/2} * b^{2-e^*(6*c^{2-d} 2-e^2) * x * arctan(cx)}/c^{3-b*d} e^{2*x^2} * (a+b*arctan(cx))/c-1/6*b^3 e^{3*x^3} * (a+b*arctan(cx))/c+I*d*(c*d-e)*(c*d+e)*(a+b*arctan(cx))^2/c^{3-1/4} * (c^4 d^4-6*c^2 d^2 e^2+e^4) * (a+b*arctan(cx))^2/c^{4/e+1/4} * (e*x+d)^4 * (a+b*arctan(cx))^2/e+2*b*d*(c*d-e)*(c*d+e)*(a+b*arctan(cx))*ln(2/(1+I*c*x))/c^{3-1/12} * b^{2-e^3} * ln(c^{2*x^2+1})/c^{4+1/4} * b^{2-e^*(6*c^2-d^2-e^2)*ln(c^{2*x^2+1})/c^4+I*b^{2-d*(c*d-e)*(c*d+e)}*polylog(2,1-2/(1+I*c*x))/c^3$

Rubi [A]

time = 0.39, antiderivative size = 376, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 14, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.778$, Rules used = {4974, 4930, 266, 4946, 327, 209, 272, 45, 5104, 5004, 5040, 4964, 2449, 2352}

$$\frac{id(cd - e)(cd + e)(a + b \operatorname{ArcTan}(cx))^2}{c^2} - \frac{2id(cd - e)(cd + e) \log\left(\frac{c}{\sqrt{cd - e^2}}\right)(a + b \operatorname{ArcTan}(cx))^2}{c^2} - \frac{(cd^2 - 6c^2 d^2 e^2 + e^4)(a + b \operatorname{ArcTan}(cx))^2}{c^2} - \frac{id e^2 x^2 (a + b \operatorname{ArcTan}(cx))^2}{c} + \frac{(d + e x)^2 (a + b \operatorname{ArcTan}(cx))^2}{4c} - \frac{ib e^2 x^2 (a + b \operatorname{ArcTan}(cx))}{c^2} - \frac{ib \operatorname{ArcTan}(cx)}{c^2} - \frac{b e^2 x \operatorname{ArcTan}(cx)}{c^2} (6c^2 d^2 - e^2) + \frac{ib^2 d (cd - e)(cd + e) L_1(1 - \frac{e^2}{cd}))}{c^2} + \frac{ib^2 d e^2 x}{c^2} + \frac{ib^2 e^2 x^3}{12c^2} + \frac{ib^2 (6c^2 d^2 - e^2) \log(c^2 x^2 + 1)}{c^2} - \frac{ib^2 e^2 \log(c^2 x^2 + 1)}{12c^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)^3 * (a + b*\operatorname{ArcTan}[c*x])^2, x]$

[Out] $(b^{2-d} e^{2-x} / c^{2-1/2} * a * b * e * (6*c^{2-d} 2-e^2) * x) / (2*c^{3-}) + (b^{2-e^3} x^2) / (12*c^{3-}) - (b^{2-d} e^{2-ArcTan[c*x]}) / c^{3-} - (b^{2-e^*(6*c^{2-d} 2-e^2) * x * ArcTan[c*x]}) / (2*c^{3-}) - (b*d^2 e^{2*x^2} * (a + b*\operatorname{ArcTan}[c*x])) / c - (b^3 e^{3*x^3} * (a + b*\operatorname{ArcTan}[c*x])) / (6*c) + (I*d*(c*d - e)*(c*d + e)*(a + b*\operatorname{ArcTan}[c*x]))^2 / c^{3-} - ((c^4 * d^4 - 6*c^2 * d^2 * e^2 + e^4) * (a + b*\operatorname{ArcTan}[c*x]))^2 / (4*c^{4-e}) + ((d + e*x)^4 * (a + b*\operatorname{ArcTan}[c*x]))^2 / (4*c^{4-e}) + (2*b*d*(c*d - e)*(c*d + e)*(a + b*\operatorname{ArcTan}[c*x])) * Log[2/(1 + I*c*x)] / c^{3-} - (b^{2-e^3} * Log[1 + c^{2*x^2}]) / (12*c^{4-}) + (b^{2-e^*(6*c^{2-d} 2-e^2) * Log[1 + c^{2*x^2}]}) / (4*c^{4-}) + (I*b^{2-d*(c*d - e)*(c*d + e)} * PolyLog[2, 1 - 2/(1 + I*c*x)]) / c^{3-}$

Rule 45

```
Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LessQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
```

```
, 0] || GtQ[b, 0])
```

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_ .)*(x_)^(n_.)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_ .)*(x_)^(n_.))^p_, x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 327

```
Int[((c_ .)*(x_))^(m_)*((a_) + (b_ .)*(x_)^(n_.))^p_, x_Symbol] :> Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2352

```
Int[Log[(c_ .)*(x_)]/((d_) + (e_ .)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_ .)/((d_) + (e_ .)*(x_))]/((f_) + (g_ .)*(x_)^2), x_Symbol] :> Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 4930

```
Int[((a_ .) + ArcTan[(c_ .)*(x_)^(n_.)]*(b_ .))^p_, x_Symbol] :> Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```

Rule 4946

```
Int[((a_ .) + ArcTan[(c_ .)*(x_)^(n_.)]*(b_ .))^p_*(x_)^m_, x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Dist[b*c*n*(p/(m + 1)), Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

Rule 4964

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)), x_Symbol]
  :> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e),
    Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)),
    x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_)*((d_) + (e_)*(x_))^(q_), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Dist[b*c*(p/(e*(q + 1))),
    Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2),
    x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b,
    c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_)*(x_)^2),
  x_Symbol) :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Dist[1/(c*d),
  Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
  d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*(f_) + (g_)*(x_))^(m_.))/((d_) + (e_)*(x_)^2),
  x_Symbol) :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p/(d + e*x^2),
  (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int (d+ex)^3 (a+b\tan^{-1}(cx))^2 dx &= \frac{(d+ex)^4 (a+b\tan^{-1}(cx))^2}{4e} - \frac{(bc) \int \left(\frac{e^2(6c^2d^2-e^2)(a+b\tan^{-1}(cx))}{c^4} + \frac{4de^3x}{c^2} \right)}{4e} \\
&= \frac{(d+ex)^4 (a+b\tan^{-1}(cx))^2}{4e} - \frac{b \int \frac{(c^4d^4-6c^2d^2e^2+e^4+4c^2d(cd-e)e(cd+e)x)(a+b\tan^{-1}(cx))}{1+c^2x^2}}{2c^3e} \\
&= -\frac{abe(6c^2d^2-e^2)x}{2c^3} - \frac{bde^2x^2(a+b\tan^{-1}(cx))}{c} - \frac{be^3x^3(a+b\tan^{-1}(cx))}{6c} \\
&= \frac{b^2de^2x}{c^2} - \frac{abe(6c^2d^2-e^2)x}{2c^3} - \frac{b^2e(6c^2d^2-e^2)x\tan^{-1}(cx)}{2c^3} - \frac{bde^2x^2(a+b\tan^{-1}(cx))}{c} \\
&= \frac{b^2de^2x}{c^2} - \frac{abe(6c^2d^2-e^2)x}{2c^3} - \frac{b^2de^2\tan^{-1}(cx)}{c^3} - \frac{b^2e(6c^2d^2-e^2)x\tan^{-1}(cx)}{2c^3} \\
&= \frac{b^2de^2x}{c^2} - \frac{abe(6c^2d^2-e^2)x}{2c^3} + \frac{b^2e^3x^2}{12c^2} - \frac{b^2de^2\tan^{-1}(cx)}{c^3} - \frac{b^2e(6c^2d^2-e^2)x\tan^{-1}(cx)}{2c^3} \\
&= \frac{b^2de^2x}{c^2} - \frac{abe(6c^2d^2-e^2)x}{2c^3} + \frac{b^2e^3x^2}{12c^2} - \frac{b^2de^2\tan^{-1}(cx)}{c^3} - \frac{b^2e(6c^2d^2-e^2)x\tan^{-1}(cx)}{2c^3} \\
&= \frac{b^2de^2x}{c^2} - \frac{abe(6c^2d^2-e^2)x}{2c^3} + \frac{b^2e^3x^2}{12c^2} - \frac{b^2de^2\tan^{-1}(cx)}{c^3} - \frac{b^2e(6c^2d^2-e^2)x\tan^{-1}(cx)}{2c^3}
\end{aligned}$$

Mathematica [A]

time = 0.52, size = 472, normalized size = 1.26

Antiderivative was successfully verified.

[In] Integrate[(d + e*x)^3*(a + b*ArcTan[c*x])^2, x]

```
[Out] (b^2*e^3 + 12*a^2*c^4*d^3*x - 36*a*b*c^3*d^2*e*x + 12*b^2*c^2*d*e^2*x + 6*a*b*c*e^3*x + 18*a^2*c^4*d^2*e*x^2 - 12*a*b*c^3*d*e^2*x^2 + b^2*c^2*e^3*x^2 + 12*a^2*c^4*d*e^2*x^3 - 2*a*b*c^3*e^3*x^3 + 3*a^2*c^4*e^3*x^4 + 3*b^2*((-4*I)*c^3*d^3 + 6*c^2*d^2*e + (4*I)*c*d*e^2 - e^3 + c^4*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3))*ArcTan[c*x]^2 + 2*b*ArcTan[c*x]*(-(b*c*e*(18*c^2*d^2*x + e^2*x*(-3 + c^2*x^2) + 6*d*(e + c^2*e*x^2))) + 3*a*(6*c^2*d^2*e - e^3 + c^4*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3)) + 12*b*c*d*(c^2*d^2 - e^2)*Log[1 + E^((2*I)*ArcTan[c*x]))] - 12*a*b*c^3*d^3*Log[1 + c^2*x^2] + 18*b^2*c^2*d^2*e*Log[1 + c^2*x^2] + 12*a*b*c*d*e^2*Log[1 + c^2*x^2] - 4*b^2*e^3*Log[1 + c^2*x^2] - (12*I)*b^2*c*d*(c^2*d^2 - e^2)*PolyLog[2, -E^((2*I)*ArcTan[c*x]))])/(12*c^4)
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 879 vs. $2(356) = 712$.

time = 0.36, size = 880, normalized size = 2.34 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((e*x+d)^3*(a+b*arctan(c*x))^2,x,method=_RETURNVERBOSE)
[Out] 1/c*(3*a*b/c*e*arctan(c*x)*d^2+3*a*b*c*e*arctan(c*x)*d^2*x^2+2*a*b*c*e^2*arctan(c*x)*d*x^3+1/2*I*b^2/c^2*x^2*d*ln(c^2*x^2+1)*ln(c*x-I)-1/2*I*b^2/c^2*e^2*d*ln(-1/2*I*(c*x+I))*ln(c*x-I)-1/2*I*b^2/c^2*x^2*d*ln(c^2*x^2+1)*ln(c*x+I)+1/2*I*b^2/c^2*x^2*d*ln(c*x+I)*ln(1/2*I*(c*x-I))-3*a*b*e*d^2*x-a*b*e^2*d*x^2+1/2*a*b*c*e^3*arctan(c*x)*x^4-3*b^2*e*arctan(c*x)*d^2*x-b^2*e^2*arctan(c*x)*d*x^2+3/2*b^2*c*e*arctan(c*x)^2*d^2*x^2+b^2*c*e^2*arctan(c*x)^2*d*x^3+1/2*I*b^2/c^2*x^2*d*dilog(1/2*I*(c*x-I))-1/4*I*b^2/c^2*x^2*d*ln(c*x-I)^2-1/6*a*b*e^3*x^3+1/2*a*b/c^2*x^2*a*b*arctan(c*x)*d^3*c*x+b^2/c*d*e^2*x+1/4*b^2*c*e^3*arctan(c*x)^2*x^4-1/6*b^2*x^3*arctan(c*x)*x^3+1/2*b^2/c^2*x^3*arctan(c*x)*x+a*b/c^2*x^2*ln(c^2*x^2+1)*d+b^2/c^2*x^2*arctan(c*x)*ln(c^2*x^2+1)*d-1/2*I*b^2/c^2*x^2*d*dilog(-1/2*I*(c*x+I))+1/4*I*b^2/c^2*x^2*d*ln(c*x+I)^2-1/2*a*b/c^3*x^3*arctan(c*x)-1/2*I*b^2*d^3*ln(c^2*x^2+1)*ln(c*x-I)+1/2*I*b^2*d^3*ln(-1/2*I*(c*x+I))*ln(c*x-I)+1/2*I*b^2*d^3*ln(c^2*x^2+1)*ln(c*x+I)-1/2*I*b^2*d^3*ln(c*x+I)*ln(1/2*I*(c*x-I))+3/2*b^2/c*e*arctan(c*x)^2*d^2*x+3/2*b^2/c*e*ln(c^2*x^2+1)*d^2-b^2/c^2*x^2*d*arctan(c*x)+1/12*b^2/c*x^2+b^2*arctan(c*x)^2*d^3*c*x+1/4*(c*e*x+c*d)^4*a^2/c^3/e-a*b*ln(c^2*x^2+1)*d^3-b^2*arctan(c*x)*ln(c^2*x^2+1)*d^3-1/4*b^2/c^3*x^3*arctan(c*x)^2-1/3*b^2/c^3*x^3*ln(c^2*x^2+1)-1/4*I*b^2*d^3*ln(c*x+I)^2-1/2*I*b^2*d^3*dilog(1/2*I*(c*x-I))+1/4*I*b^2*d^3*ln(c*x-I)^2+1/2*I*b^2*d^3*dilog(-1/2*I*(c*x+I)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^3*(a+b*arctan(c*x))^2,x, algorithm="maxima")
[Out] 36*b^2*c^2*d^2*e*integrate(1/16*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^2*c^2*d^2*e*integrate(1/16*x^3*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 12*b^2*c^2*d^3*integrate(1/16*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 6*b^2*c^2*d^2*e*integrate(1/16*x^3*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + b^2*c^2*d^3*integrate(1/16*x^2*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 4*b^2*c^2*d^3*integrate(1/16*x^2*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + 1/4*b^2*d^3*arctan(c*x)^3/c + 1/4*a^2*x^4*e^3 + a^2*d*x^3*e^2 + 3/2*a^2*d^2*x^2*e + 36*b^2*c^2*d*e^2*integrate(1/16*x^4*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^2*c^2*d^2*e^2*integrate(1/16*x^4*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 4*b^2*c^2*d^2*e^2*integrate(1/16*x^4*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) - 12*b^2*c*d^2*e*integrate(1/16*x^2*arctan(c*x)/(c^2*x^2 + 1), x) - 8*b^2*c*d^3*integrate(1/16*x*arctan(c*x)^2/(c^2*x^2 + 1), x)
```

$$\begin{aligned}
& \frac{(c*x)/(c^2*x^2 + 1), x) + a^2*d^3*x + 3*(x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*a*b*d^2*e + 12*b^2*c^2*e^3*integrate(1/16*x^5*arctan(c*x)^2/(c^2*x^2 + 1), x) + b^2*c^2*e^3*integrate(1/16*x^5*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + b^2*c^2*e^3*integrate(1/16*x^5*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) - 8*b^2*c*d*e^2*integrate(1/16*x^3*arctan(c*x)/(c^2*x^2 + 1), x) + 36*b^2*d^2*e*integrate(1/16*x*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^2*d^2*e*integrate(1/16*x*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + b^2*d^3*integrate(1/16*x*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + (2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*a*b*d^3/c + (2*x^3*arctan(c*x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c^4))*a*b*d*e^2 - 2*b^2*c*e^3*integrate(1/16*x^4*arctan(c*x)/(c^2*x^2 + 1), x) + 36*b^2*d^2*e^2*integrate(1/16*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^2*d^2*e^2*integrate(1/16*x^2*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 1/6*(3*x^4*arctan(c*x) - c*((c^2*x^3 - 3*x)/c^4 + 3*arctan(c*x)/c^5))*a*b*e^3 + 12*b^2*e^3*integrate(1/16*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + b^2*e^3*integrate(1/16*x^3*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 1/16*(b^2*x^4*e^3 + 4*b^2*d*x^3*e^2 + 6*b^2*d^2*x^2*e + 4*b^2*d^3*x)*arctan(c*x)^2 - 1/64*(b^2*x^4*e^3 + 4*b^2*d*x^3*e^2 + 6*b^2*d^2*x^2*e + 4*b^2*d^3*x)*log(c^2*x^2 + 1)^2
\end{aligned}$$
Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^3*(a+b*arctan(c*x))^2,x, algorithm="fricas")`
[Out] `integral(a^2*x^3*e^3 + 3*a^2*d*x^2*e^2 + 3*a^2*d^2*x*e + a^2*d^3 + (b^2*x^3 *e^3 + 3*b^2*d*x^2*e^2 + 3*b^2*d^2*x*e + b^2*d^3)*arctan(c*x)^2 + 2*(a*b*x^3 *e^3 + 3*a*b*d*x^2*e^2 + 3*a*b*d^2*x*e + a*b*d^3)*arctan(c*x), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx))^2 (d + ex)^3 \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)**3*(a+b*atan(c*x))**2,x)`
[Out] `Integral((a + b*atan(c*x))**2*(d + e*x)**3, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^3*(a+b*arctan(c*x))^2,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a + b \operatorname{atan}(c x))^2 (d + e x)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^2*(d + e*x)^3,x)`

[Out] `int((a + b*atan(c*x))^2*(d + e*x)^3, x)`

3.10 $\int (d + ex)^2(a + b \operatorname{ArcTan}(cx))^2 dx$

Optimal. Leaf size=270

$$-\frac{2abdex}{c} + \frac{b^2e^2x}{3c^2} - \frac{b^2e^2\operatorname{ArcTan}(cx)}{3c^3} - \frac{2b^2dex\operatorname{ArcTan}(cx)}{c} - \frac{be^2x^2(a + b\operatorname{ArcTan}(cx))}{3c} + \frac{i(3c^2d^2 - e^2)(a + b\operatorname{ArcTan}(cx))}{3c^3}$$

[Out] $-2*a*b*d*e*x/c+1/3*b^2*e^2*x/c^2-1/3*b^2*e^2*arctan(c*x)/c^3-2*b^2*d*e*x*arctan(c*x)/c-1/3*b*e^2*x^2*(a+b*arctan(c*x))/c+1/3*I*(3*c^2*d^2-e^2)*(a+b*arctan(c*x))^2/c^3-1/3*d*(d^2-3*e^2/c^2)*(a+b*arctan(c*x))^2/e+1/3*(e*x+d)^3*(a+b*arctan(c*x))^2/e+2/3*b*(3*c^2*d^2-e^2)*(a+b*arctan(c*x))*ln(2/(1+I*c*x))/c^3+b^2*d*e*ln(c^2*x^2+1)/c^2+1/3*I*b^2*(3*c^2*d^2-e^2)*polylog(2,1-2/(1+I*c*x))/c^3$

Rubi [A]

time = 0.26, antiderivative size = 270, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {4974, 4930, 266, 4946, 327, 209, 5104, 5004, 5040, 4964, 2449, 2352}

$$-\frac{d\left(d^2 - \frac{b^2 e^2}{c^2}\right)(a + b\operatorname{ArcTan}(cx))^2}{3e} + \frac{i(3c^2d^2 - e^2)(a + b\operatorname{ArcTan}(cx))^2}{3c^3} + \frac{2b(3c^2d^2 - e^2)\log(\frac{c}{1+I*c*x})}{3c^3}(a + b\operatorname{ArcTan}(cx)) + \frac{(d + ex)^3(a + b\operatorname{ArcTan}(cx))^2}{3e} - \frac{be^2x^2(a + b\operatorname{ArcTan}(cx))}{3c} - \frac{2abdex}{c} - \frac{b^2e^2\operatorname{ArcTan}(cx)}{3c^3} - \frac{2b^2dex\operatorname{ArcTan}(cx)}{c} + \frac{b^2de\log(e^2x^2 + 1)}{c^2} + \frac{b^2e^2x}{3c^2} + \frac{ib^2(3c^2d^2 - e^2)\text{Li}_2(1 - \frac{ex}{1+I*c*x})}{3c^3}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)^2*(a + b*\operatorname{ArcTan}[c*x])^2, x]$

[Out] $(-2*a*b*d*e*x)/c + (b^2*e^2*x)/(3*c^2) - (b^2*x^2*\operatorname{ArcTan}[c*x])/(3*c^3) - (2*b^2*d*e*x*\operatorname{ArcTan}[c*x])/c - (b*x^2*x^2*(a + b*\operatorname{ArcTan}[c*x]))/(3*c) + ((I/3)*(3*c^2*d^2 - e^2)*(a + b*\operatorname{ArcTan}[c*x])^2)/c^3 - (d*(d^2 - (3*e^2)/c^2)*(a + b*\operatorname{ArcTan}[c*x])^2)/(3*e) + ((d + e*x)^3*(a + b*\operatorname{ArcTan}[c*x])^2)/(3*e) + (2*b*(3*c^2*d^2 - e^2)*(a + b*\operatorname{ArcTan}[c*x])*Log[2/(1 + I*c*x)])/(3*c^3) + (b^2*d*e*Log[1 + c^2*x^2])/c^2 + ((I/3)*b^2*(3*c^2*d^2 - e^2)*PolyLog[2, 1 - 2/(1 + I*c*x)])/c^3$

Rule 209

$\operatorname{Int}[(a_.) + (b_.)*(x_.)^2]^{(-1)}, x_{\text{Symbol}}] :> \operatorname{Simp}[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&& \operatorname{PosQ}[a/b] \&& (\operatorname{GtQ}[a, 0] \|\operatorname{GtQ}[b, 0])$

Rule 266

$\operatorname{Int}[(x_.)^{(m_.)}/((a_.) + (b_.)*(x_.)^{(n_.)}), x_{\text{Symbol}}] :> \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \operatorname{FreeQ}[\{a, b, m, n\}, x] \&& \operatorname{EqQ}[m, n - 1]$

Rule 327

$\operatorname{Int}[((c_.)*(x_.)^{(m_.)})*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, x_{\text{Symbol}}] :> \operatorname{Simp}[c^{(n - 1)}*(c*x)^{(m - n + 1)}*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - \operatorname{Dist}[\dots]$

```
a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x],  
x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p  
+ 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2352

```
Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLo  
g[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_)/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] :> Dist  
[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{  
c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 4930

```
Int[((a_*) + ArcTan[(c_)*(x_)^(n_.)]*(b_.))^p, x_Symbol] :> Simp[x*(a  
+ b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^p  
- 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] &&  
(EqQ[n, 1] || EqQ[p, 1])
```

Rule 4946

```
Int[((a_*) + ArcTan[(c_)*(x_)^(n_.)]*(b_.))^p*(x_)^m, x_Symbol] :>  
Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Dist[b*c*n*(p/(m +  
1)), Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x]  
/; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] &&  
IntegerQ[m])) && NeQ[m, -1]
```

Rule 4964

```
Int[((a_*) + ArcTan[(c_)*(x_)]*(b_.))^p/((d_) + (e_)*(x_)), x_Symbol]  
:> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(  
p/e), Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)),  
x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_*) + ArcTan[(c_)*(x_)]*(b_.))^p*((d_) + (e_)*(x_))^q, x_Symbol]  
:> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - D  
ist[b*c*(p/(e*(q + 1))), Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1),  
(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] &&  
IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^^(p_.)/((d_) + (e_)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^^(p_.)*(x_))/((d_) + (e_)*(x_)^2), x_Symbol] :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Dist[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^^(p_.)*(f_) + (g_)*(x_))^(m_.))/((d_) + (e_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p / (d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned} \int (d + ex)^2 (a + b \tan^{-1}(cx))^2 dx &= \frac{(d + ex)^3 (a + b \tan^{-1}(cx))^2}{3e} - \frac{(2bc) \int \left(\frac{3de^2(a + b \tan^{-1}(cx))}{c^2} + \frac{e^3 x (a + b \tan^{-1}(cx))}{c^2} \right)}{3e} \\ &= \frac{(d + ex)^3 (a + b \tan^{-1}(cx))^2}{3e} - \frac{(2b) \int \frac{(c^2 d^3 - 3de^2 + e(3c^2 d^2 - e^2)x)(a + b \tan^{-1}(cx))}{1 + c^2 x^2}}{3ce} \\ &= -\frac{2abdex}{c} - \frac{be^2 x^2 (a + b \tan^{-1}(cx))}{3c} + \frac{(d + ex)^3 (a + b \tan^{-1}(cx))^2}{3e} - \frac{2abdex}{c} + \frac{b^2 e^2 x}{3c^2} - \frac{2b^2 dex \tan^{-1}(cx)}{c} - \frac{be^2 x^2 (a + b \tan^{-1}(cx))}{3c} \\ &= -\frac{2abdex}{c} + \frac{b^2 e^2 x}{3c^2} - \frac{b^2 e^2 \tan^{-1}(cx)}{3c^3} - \frac{2b^2 dex \tan^{-1}(cx)}{c} - \frac{be^2 x^2 (a + b \tan^{-1}(cx))}{3c} \\ &= -\frac{2abdex}{c} + \frac{b^2 e^2 x}{3c^2} - \frac{b^2 e^2 \tan^{-1}(cx)}{3c^3} - \frac{2b^2 dex \tan^{-1}(cx)}{c} - \frac{be^2 x^2 (a + b \tan^{-1}(cx))}{3c} \\ &= -\frac{2abdex}{c} + \frac{b^2 e^2 x}{3c^2} - \frac{b^2 e^2 \tan^{-1}(cx)}{3c^3} - \frac{2b^2 dex \tan^{-1}(cx)}{c} - \frac{be^2 x^2 (a + b \tan^{-1}(cx))}{3c} \end{aligned}$$

Mathematica [A]

time = 0.35, size = 312, normalized size = 1.16

$$\frac{3a^3c^3d^3x - 6abc^2dex + b^2cx^2 + 3a^2c^2dex^2 - abc^2e^2x^2 + a^2c^2e^2x^3 + b^2(-3ac^2d^2 + 3cde + ie^2 + e^2x(3d^2 + 3dex + e^2x^2)) \operatorname{ArcTan}(cx)^2 + b\operatorname{ArcTan}(cx)^3 + 2ac^2x(3d^2 + 3dex + e^2x^2) + 2b(3c^2d^2 - e^2) \log(1 + e^{2\operatorname{ArcTan}(cx)}) - 3abc^2d^2 \log(1 + e^2x^2) + 3b^2cde \log(1 + e^2x^2) + abc^2 \log(1 + e^2x^2) - b^2(3c^2d^2 - e^2) \operatorname{PolyLog}(2, -e^{2\operatorname{ArcTan}(cx)})}{3c^2}$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)^2*(a + b*ArcTan[c*x])^2, x]`

[Out] $(3a^2c^3d^2x^2 - 6abc^2e^2x^2 + b^2c^2e^2x^3 + 3a^2c^3d^2e^2x^2 - a*b*c^2e^2x^2 + a^2c^3e^2x^3 + b^2((-3*I)*c^2d^2 + 3*c*d*e + I*e^2 + c^3*x*(3*d^2 + 3*dex + e^2*x^2))*\operatorname{ArcTan}(c*x)^2 + b*\operatorname{ArcTan}(c*x)*(6*a*c*d*e - b*e*(e + 6*c^2*d*x + c^2*e*x^2) + 2*a*c^3*x*(3*d^2 + 3*dex + e^2*x^2) + 2*b*(3*c^2*d^2 - e^2)*\operatorname{Log}[1 + e^{2\operatorname{ArcTan}(c*x)}]) - 3*a*b*c^2*d^2*\operatorname{Log}[1 + e^2*x^2] + 3*b^2*c*d*e*\operatorname{Log}[1 + e^2*x^2] + a*b*e^2*\operatorname{Log}[1 + e^2*x^2] - I*b^2*(3*c^2*d^2 - e^2)*\operatorname{PolyLog}[2, -e^{((2*I)*\operatorname{ArcTan}(c*x))}])/(3*c^3)$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 699 vs. 2(250) = 500.

time = 0.23, size = 700, normalized size = 2.59 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)^2*(a+b*arctan(c*x))^2, x, method=_RETURNVERBOSE)`

[Out] $\frac{1}{c}*(2*a*b*c*e*\operatorname{arctan}(c*x)*d*x^2 + b^2*c*e*\operatorname{arctan}(c*x)^2*d*x^2 - 2*b^2*e*\operatorname{arctan}(c*x)*d*x^2 - 2*a*b*d*e*x^2 + 3*a*b*c*e^2*\operatorname{arctan}(c*x)*x^3 + 1/6*I*b^2/c^2*\ln(c*x - I)*\ln(c^2*x^2 + 1)*e^2 - 1/6*I*b^2/c^2*\ln(-1/2*I*(c*x + I))*\ln(c*x - I)*e^2 - 1/6*I*b^2/c^2*\ln(c*x + I)*\ln(c^2*x^2 + 1)*e^2 + 1/6*I*b^2/c^2*\ln(1/2*I*(c*x - I))*\ln(c*x + I)*e^2 + 2*a*b/c*e*\operatorname{arctan}(c*x)*d - 1/3*b^2*e^2*\operatorname{arctan}(c*x)*x^2 + 1/3*b^2*c*e^2*\operatorname{arctan}(c*x)^2*x^3 - 1/3*a*b*e^2*x^2 + 2*a*b*\operatorname{arctan}(c*x)*d^2 - 2*c*x + 1/3*(c*e*x + c*d)^3*a^2/2/c^2/e - 1/3*b^2/c^2*\ln(c^2*x^2 + 1)*d^2 + 1/2*I*b^2*\operatorname{dilog}(-1/2*I*(c*x + I))*d^2 - 1/4*I*b^2*\ln(c*x + I)*d^2 - 1/2*I*b^2*\operatorname{dilog}(1/2*I*(c*x - I))*d^2 - a*b*\ln(c^2*x^2 + 1)*d^2 + 1/3*b^2/c^2*\ln(c^2*x^2 + 1)*d^2 + 1/3*b^2/c^2*\operatorname{arctan}(c*x)*\ln(c^2*x^2 + 1) + b^2/c*e*\operatorname{arctan}(c*x)^2*d - 1/2*I*b^2*\ln(1/2*I*(c*x - I))*\ln(c*x + I)*d^2 - 1/2*I*b^2*\ln(c*x - I)*\ln(c^2*x^2 + 1)*d^2 + 1/2*I*b^2*\ln(-1/2*I*(c*x + I))*\ln(c*x - I)*d^2 + 1/2*I*b^2*\ln(c*x + I)*\ln(c^2*x^2 + 1)*d^2 - 1/12*I*b^2/c^2*\ln(c*x - I)^2*e^2 - 1/6*I*b^2/c^2*\operatorname{dilog}(-1/2*I*(c*x + I))*e^2 + 1/12*I*b^2/c^2*\ln(c*x + I)^2*e^2 + 1/6*I*b^2/c^2*\operatorname{dilog}(1/2*I*(c*x - I))*e^2 + 1/3*a*b/c^2*e^2*\ln(c^2*x^2 + 1))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^2*(a+b*arctan(c*x))^2,x, algorithm="maxima")
[Out] 72*b^2*c^2*d*e*integrate(1/48*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + 6*b^2*c
^2*d*e*integrate(1/48*x^3*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 36*b^2*c^2
*d^2*integrate(1/48*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 12*b^2*c^2*d*e*in
tegrate(1/48*x^3*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + 3*b^2*c^2*d^2*integra
te(1/48*x^2*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 12*b^2*c^2*d^2*integrate
(1/48*x^2*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + 1/4*b^2*d^2*arctan(c*x)^3/c
+ 1/3*a^2*x^3*e^2 + a^2*d*x^2*e + 36*b^2*c^2*e^2*integrate(1/48*x^4*arctan(
c*x)^2/(c^2*x^2 + 1), x) + 3*b^2*c^2*e^2*integrate(1/48*x^4*log(c^2*x^2 + 1)
)^2/(c^2*x^2 + 1), x) + 4*b^2*c^2*e^2*integrate(1/48*x^4*log(c^2*x^2 + 1)/(
c^2*x^2 + 1), x) - 24*b^2*c*d*e*integrate(1/48*x^2*arctan(c*x)/(c^2*x^2 + 1
), x) - 24*b^2*c*d^2*integrate(1/48*x*arctan(c*x)/(c^2*x^2 + 1), x) + a^2*d
^2*x + 2*(x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*a*b*d*e - 8*b^2*c*
e^2*integrate(1/48*x^3*arctan(c*x)/(c^2*x^2 + 1), x) + 72*b^2*d*e*integrate
(1/48*x*arctan(c*x)^2/(c^2*x^2 + 1), x) + 6*b^2*d*e*integrate(1/48*x*log(c^
2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 3*b^2*d^2*integrate(1/48*log(c^2*x^2 + 1)^
2/(c^2*x^2 + 1), x) + (2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*a*b*d^2/c + 1/
3*(2*x^3*arctan(c*x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c^4))*a*b*e^2 + 36*b^2
*e^2*integrate(1/48*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^2*e^2*integra
te(1/48*x^2*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 1/12*(b^2*x^3*e^2 + 3*b^
2*d*x^2*e + 3*b^2*d^2*x)*arctan(c*x)^2 - 1/48*(b^2*x^3*e^2 + 3*b^2*d*x^2*e
+ 3*b^2*d^2*x)*log(c^2*x^2 + 1)^2
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^2*(a+b*arctan(c*x))^2,x, algorithm="fricas")
```

```
[Out] integral(a^2*x^2*e^2 + 2*a^2*d*x*e + a^2*d^2 + (b^2*x^2*e^2 + 2*b^2*d*x*e +
b^2*d^2)*arctan(c*x)^2 + 2*(a*b*x^2*e^2 + 2*a*b*d*x*e + a*b*d^2)*arctan(c*
x), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx))^2 (d + ex)^2 \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)**2*(a+b*atan(c*x))**2,x)
```

```
[Out] Integral((a + b*atan(c*x))**2*(d + e*x)**2, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x))^2,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a + b \operatorname{atan}(c x))^2 (d + e x)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^2*(d + e*x)^2,x)`

[Out] `int((a + b*atan(c*x))^2*(d + e*x)^2, x)`

3.11 $\int (d + ex)(a + b \operatorname{ArcTan}(cx))^2 dx$

Optimal. Leaf size=171

$$-\frac{abex}{c} - \frac{b^2 ex \operatorname{ArcTan}(cx)}{c} + \frac{id(a + b \operatorname{ArcTan}(cx))^2}{c} - \frac{\left(d^2 - \frac{e^2}{c^2}\right) (a + b \operatorname{ArcTan}(cx))^2}{2e} + \frac{(d + ex)^2 (a + b \operatorname{ArcTan}(cx))^2}{2e}$$

[Out] $-a*b*e*x/c - b^2 e*x*\arctan(c*x)/c + I*d*(a+b*\arctan(c*x))^2/c - 1/2*(d^2 - e^2/c^2)*(a+b*\arctan(c*x))^2/c + id(a+b*\arctan(c*x))^2/c - (d+ex)^2(a+b*\arctan(c*x))^2/(2e) + (d+ex)^2(a+b*\arctan(c*x))^2/(2e)$
 $-(a+b*\arctan(c*x))^2/e + 1/2*(e*x+d)^2*(a+b*\arctan(c*x))^2/e + 2*b*d*(a+b*\arctan(c*x))*\ln(2/(1+I*c*x))/c + 1/2*b^2 e^2*\ln(c^2*x^2+1)/c^2 + I*b^2 d^2*polylog(2, 1 - 2/(1+I*c*x))/c$

Rubi [A]

time = 0.20, antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 9, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.562,
 Rules used = {4974, 4930, 266, 5104, 5004, 5040, 4964, 2449, 2352}

$$-\frac{\left(d^2 - \frac{e^2}{c^2}\right) (a + b \operatorname{ArcTan}(cx))^2}{2e} + \frac{(d + ex)^2 (a + b \operatorname{ArcTan}(cx))^2}{2e} + \frac{id(a + b \operatorname{ArcTan}(cx))^2}{c} + \frac{2bd \log\left(\frac{2}{1+icx}\right) (a + b \operatorname{ArcTan}(cx))}{c} - \frac{abex}{c} - \frac{b^2 ex \operatorname{ArcTan}(cx)}{c} + \frac{b^2 e \log(c^2 x^2 + 1)}{2c^2} + \frac{ib^2 d \text{Li}_2\left(1 - \frac{2}{icx+1}\right)}{c}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)*(a + b*\operatorname{ArcTan}[c*x])^2, x]$

[Out] $-\left((a*b*e*x)/c\right) - \left(b^2 e*x*\operatorname{ArcTan}[c*x]/c\right) + \left(I*d*(a + b*\operatorname{ArcTan}[c*x])^2/c\right) - \left((d^2 - e^2/c^2)*(a + b*\operatorname{ArcTan}[c*x])^2/(2*e)\right) + \left((d + e*x)^2*(a + b*\operatorname{ArcTan}[c*x])^2/(2*e)\right) + \left(2*b*d*(a + b*\operatorname{ArcTan}[c*x])*Log[2/(1 + I*c*x)]/c\right) + \left(b^2 e*\operatorname{Log}[1 + c^2*x^2]/(2*c^2)\right) + \left(I*b^2 d^2*\operatorname{PolyLog}[2, 1 - 2/(1 + I*c*x)]/c\right)$

Rule 266

$\operatorname{Int}[(x_.)^m*((a_.) + (b_.)*(x_.)^n), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \operatorname{FreeQ}[\{a, b, m, n\}, x] \&& \operatorname{EqQ}[m, n - 1]$

Rule 2352

$\operatorname{Int}[\operatorname{Log}[(c_.)*(x_.)]/((d_.) + (e_.)*(x_.)), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-e^{(-1)})*\operatorname{PolyLog}[2, 1 - c*x], x] /; \operatorname{FreeQ}[\{c, d, e\}, x] \&& \operatorname{EqQ}[e + c*d, 0]$

Rule 2449

$\operatorname{Int}[\operatorname{Log}[(c_.)/((d_.) + (e_.)*(x_.))]/((f_.) + (g_.)*(x_.)^2), x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[-e/g, \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \operatorname{FreeQ}[\{c, d, e, f, g\}, x] \&& \operatorname{EqQ}[c, 2*d] \&& \operatorname{EqQ}[e^2*f + d^2*g, 0]$

Rule 4930

$\operatorname{Int}[((a_.) + \operatorname{ArcTan}[(c_.)*(x_.)^n]*(b_.))^p, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[x*(a + b*\operatorname{ArcTan}[c*x^n])^p, x] - \operatorname{Dist}[b*c*n*p, \operatorname{Int}[x^n*((a + b*\operatorname{ArcTan}[c*x^n])^p, x)] /; \operatorname{FreeQ}[\{a, b, c, d, e, f, g\}, x] \&& \operatorname{EqQ}[c, 2*d] \&& \operatorname{EqQ}[e^2*f + d^2*g, 0]$

```
- 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] &&
(EqQ[n, 1] || EqQ[p, 1])
```

Rule 4964

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)), x_Symbol]
  :> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e),
  Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)),
  x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_)*((d_) + (e_)*(x_))^(q_), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Dist[b*c*(p/(e*(q + 1))),
  Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_)*(x_)^2),
x_Symbol) :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Dist[1/(c*d),
Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*(f_) + (g_)*(x_))^(m_.))/(((d_) + (e_)*(x_)^2),
x_Symbol) :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int (d + ex) (a + b \tan^{-1}(cx))^2 dx &= \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^2}{2e} - \frac{(bc) \int \left(\frac{e^2 (a + b \tan^{-1}(cx))}{c^2} + \frac{(c^2 d^2 - e^2 + 2c^2 dex)}{c^2 (1+c^2 x^2)} \right) dx}{e} \\
&= \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^2}{2e} - \frac{b \int \frac{(c^2 d^2 - e^2 + 2c^2 dex)(a + b \tan^{-1}(cx))}{1 + c^2 x^2} dx}{ce} - \frac{(be) \int \frac{c^2 d^2 (1 - \frac{e^2}{c^2 d^2}) (a + b \tan^{-1}(cx))}{1 + c^2 x^2} dx}{ce} \\
&= -\frac{abex}{c} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^2}{2e} - \frac{b \int \frac{c^2 d^2 (1 - \frac{e^2}{c^2 d^2}) (a + b \tan^{-1}(cx))}{1 + c^2 x^2} dx}{ce} \\
&= -\frac{abex}{c} - \frac{b^2 ex \tan^{-1}(cx)}{c} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^2}{2e} - (2bcd) \int \frac{x}{1 + c^2 x^2} dx \\
&= -\frac{abex}{c} - \frac{b^2 ex \tan^{-1}(cx)}{c} + \frac{id(a + b \tan^{-1}(cx))^2}{c} - \frac{\left(d^2 - \frac{e^2}{c^2}\right) (a + b \tan^{-1}(cx))^2}{2e} \\
&= -\frac{abex}{c} - \frac{b^2 ex \tan^{-1}(cx)}{c} + \frac{id(a + b \tan^{-1}(cx))^2}{c} - \frac{\left(d^2 - \frac{e^2}{c^2}\right) (a + b \tan^{-1}(cx))^2}{2e} \\
&= -\frac{abex}{c} - \frac{b^2 ex \tan^{-1}(cx)}{c} + \frac{id(a + b \tan^{-1}(cx))^2}{c} - \frac{\left(d^2 - \frac{e^2}{c^2}\right) (a + b \tan^{-1}(cx))^2}{2e} \\
&= -\frac{abex}{c} - \frac{b^2 ex \tan^{-1}(cx)}{c} + \frac{id(a + b \tan^{-1}(cx))^2}{c} - \frac{\left(d^2 - \frac{e^2}{c^2}\right) (a + b \tan^{-1}(cx))^2}{2e}
\end{aligned}$$

Mathematica [A]

time = 0.14, size = 172, normalized size = 1.01

$$\frac{2a^2c^2dx - 2abceix + a^2c^2ex^2 + b^2(-i + cx)(2cd + ie + cex)\text{ArcTan}(cx)^2 + 2b\text{ArcTan}(cx)(-bceix + a(e + 2c^2dx + c^2ex^2) + 2bcd\log(1 + e^{2i\text{ArcTan}(cx)})) - 2abcd\log(1 + c^2x^2) + b^2e\log(1 + c^2x^2) - 2ib^2cd\text{PolyLog}(2, -e^{2i\text{ArcTan}(cx)})}{2c^2}$$

Antiderivative was successfully verified.

[In] Integrate[(d + e*x)*(a + b*ArcTan[c*x])^2, x]

[Out] $(2*a^2*c^2*d*x - 2*a*b*c*e*x + a^2*c^2*e*x^2 + b^2*(-I + c*x)*(2*c*d + I*e + c*e*x)*\text{ArcTan}[c*x]^2 + 2*b*\text{ArcTan}[c*x]*(-(b*c*e*x) + a*(e + 2*c^2*d*x + c^2*e*x^2) + 2*b*c*d*\text{Log}[1 + E^{((2*I)*\text{ArcTan}[c*x])}] - 2*a*b*c*d*\text{Log}[1 + c^2*x^2] + b^2*e*\text{Log}[1 + c^2*x^2] - (2*I)*b^2*c*d*\text{PolyLog}[2, -E^{((2*I)*\text{ArcTan}[c*x])}])/(2*c^2)$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 339 vs. $2(161) = 322$.

time = 0.12, size = 340, normalized size = 1.99

method	result
--------	--------

derivative divides	$\frac{a^2(d c^2 x + \frac{1}{2} e c^2 x^2)}{c} + b^2 \arctan(cx)^2 d x + \frac{b^2 c \arctan(cx)^2 e x^2}{2} - b^2 \ln(c^2 x^2 + 1) \arctan(cx) d + \frac{b^2 \arctan(cx)^2 e}{2c} - b^2 \arctan(cx) e x^2$
default	$\frac{a^2(d c^2 x + \frac{1}{2} e c^2 x^2)}{c} + b^2 \arctan(cx)^2 d x + \frac{b^2 c \arctan(cx)^2 e x^2}{2} - b^2 \ln(c^2 x^2 + 1) \arctan(cx) d + \frac{b^2 \arctan(cx)^2 e}{2c} - b^2 \arctan(cx) e x^2$
risch	$a^2 dx + \frac{7b^2 e \ln(c^2 x^2 + 1)}{16c^2} - \frac{abex}{c} + \frac{ea^2}{2c^2} + \frac{eba \arctan(cx)}{c^2} - \frac{abd \ln(c^2 x^2 + 1)}{c} + \frac{a^2 e x^2}{2} + \frac{ib^2 \ln(\frac{1}{2} - \frac{icx}{2}) \ln(\frac{1}{2} + \frac{icx}{2})}{c}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)*(a+b*arctan(c*x))^2,x,method=_RETURNVERBOSE)`

[Out] $1/c*(a^2/c*(d*c^2*x+1/2*e*c^2*x^2)+b^2*arctan(c*x)^2*d*c*x+1/2*b^2*c*arctan(c*x)^2*e*x^2-b^2*ln(c^2*x^2+1)*arctan(c*x)*d+1/2*b^2*c*arctan(c*x)^2*e-b^2*arctan(c*x)*e*x+1/2*b^2*c*e*ln(c^2*x^2+1)-1/2*I*b^2*d*dilog(1/2*I*(c*x-I))+1/2*I*b^2*d*ln(c*x+I)*ln(c^2*x^2+1)-1/4*I*b^2*d*ln(c*x+I)^2-1/2*I*b^2*d*ln(c*x+I)*ln(1/2*I*(c*x-I))+1/2*I*b^2*d*ln(c*x-I)*ln(-1/2*I*(c*x+I))-1/2*I*b^2*d*ln(c*x-I)*ln(c^2*x^2+1)+1/2*I*b^2*d*dilog(-1/2*I*(c*x+I))+1/4*I*b^2*d*ln(c*x-I)^2+2*a*b*arctan(c*x)*d*c*x+a*b*c*arctan(c*x)*e*x^2-a*b*e*x-a*b*d*ln(c^2*x^2+1)+a*b/c*e*arctan(c*x))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x))^2,x, algorithm="maxima")`

[Out] $12*b^2*c^2*e*integrate(1/16*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + b^2*c^2*e*integrate(1/16*x^3*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 12*b^2*c^2*d*integrate(1/16*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 2*b^2*c^2*e*integrate(1/16*x^3*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + b^2*c^2*d*integrate(1/16*x^2*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 4*b^2*c^2*d*integrate(1/16*x^2*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + 1/4*b^2*d*arctan(c*x)^3/c + 1/2*a^2*x^2*e - 4*b^2*c*e*integrate(1/16*x^2*arctan(c*x)/(c^2*x^2 + 1), x) - 8*b^2*c*d*integrate(1/16*x*arctan(c*x)/(c^2*x^2 + 1), x) + a^2*d*x + (x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*a*b*e + 12*b^2*e*integrate(1/16*x*arctan(c*x)^2/(c^2*x^2 + 1), x) + b^2*e*integrate(1/16*x*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + b^2*d*integrate(1/16*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + (2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*a*b*d/c + 1/8*(b^2*x^2*e + 2*b^2*d*x)*arctan(c*x)^2 - 1/32*(b^2*x^2*e + 2*b^2*d*x)*log(c^2*x^2 + 1)^2$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x))^2,x, algorithm="fricas")`

[Out] `integral(a^2*x*e + a^2*d + (b^2*x*e + b^2*d)*arctan(c*x)^2 + 2*(a*b*x*e + a*b*d)*arctan(c*x), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx))^2 (d + ex) \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*atan(c*x))**2,x)`

[Out] `Integral((a + b*atan(c*x))**2*(d + e*x), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x))^2,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int (a + b \operatorname{atan}(cx))^2 (d + e x) \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^2*(d + e*x),x)`

[Out] `int((a + b*atan(c*x))^2*(d + e*x), x)`

3.12 $\int \frac{(a+b\text{ArcTan}(cx))^2}{d+ex} dx$

Optimal. Leaf size=223

$$-\frac{(a+b\text{ArcTan}(cx))^2 \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a+b\text{ArcTan}(cx))^2 \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{ib(a+b\text{ArcTan}(cx))\text{PolyLog}(2, \frac{2c(d+ex)}{(cd+ie)(1-icx)})}{e}$$

[Out] $-(a+b*\text{arctan}(c*x))^2 \ln(2/(1-I*c*x))/e + (a+b*\text{arctan}(c*x))^2 \ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e + I*b*(a+b*\text{arctan}(c*x))*\text{polylog}(2, 1-2/(1-I*c*x))/e - I*b*(a+b*\text{arctan}(c*x))*\text{polylog}(2, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e - 1/2*b^2*\text{polylog}(3, 1-2/(1-I*c*x))/e + 1/2*b^2*\text{polylog}(3, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e$

Rubi [A]

time = 0.03, antiderivative size = 223, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.056, Rules used = {4968}

$$-\frac{ib(a+b\text{ArcTan}(cx))\text{Li}_2\left(1-\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{(a+b\text{ArcTan}(cx))^2 \log\left(\frac{2c(d+ex)}{(1-icx)(cd+ie)}\right)}{e} + \frac{ib\text{Li}_2\left(1-\frac{2}{1-icx}\right)(a+b\text{ArcTan}(cx))}{e} - \frac{\log\left(\frac{2}{1-icx}\right)(a+b\text{ArcTan}(cx))^2}{e} + \frac{b^2\text{Li}_3\left(1-\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{2e} - \frac{b^2\text{Li}_3\left(1-\frac{2}{1-icx}\right)}{2e}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a+b*\text{ArcTan}[c*x])^2/(d+e*x), x]$

[Out] $-((a+b*\text{ArcTan}[c*x])^2 \text{Log}[2/(1-I*c*x)]/e) + ((a+b*\text{ArcTan}[c*x])^2 \text{Log}[2*c*(d+e*x)/((c*d+I*e)*(1-I*c*x))]/e) + (I*b*(a+b*\text{ArcTan}[c*x])*\text{PolyLog}[2, 1-2/(1-I*c*x)]/e) - (I*b*(a+b*\text{ArcTan}[c*x])*\text{PolyLog}[2, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))]/e) - (b^2*\text{PolyLog}[3, 1-2/(1-I*c*x)]/(2*e)) + (b^2*\text{PolyLog}[3, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))]/(2*e))$

Rule 4968

```
Int[((a_) + ArcTan[(c_)*(x_)]*(b_))^(2/((d_) + (e_)*(x_)), x_Symbol] :>
Simp[(-(a + b*ArcTan[c*x])^2)*Log[2/(1 - I*c*x)]/e, x] + (Simp[(a + b*ArcTan[c*x])^(2*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x] + Simp[I*b*(a + b*ArcTan[c*x])*(PolyLog[2, 1 - 2/(1 - I*c*x)]/e), x] - Simp[I*b*(a + b*ArcTan[c*x])*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x] - Simp[b^(2*(PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x] + Simp[b^(2*(PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x]) /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^(2*d^2 + e^2, 0]
```

Rubi steps

$$\int \frac{(a+b\tan^{-1}(cx))^2}{d+ex} dx = -\frac{(a+b\tan^{-1}(cx))^2 \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a+b\tan^{-1}(cx))^2 \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{ib(a+b\tan^{-1}(cx))\text{PolyLog}(2, \frac{2c(d+ex)}{(cd+ie)(1-icx)})}{e}$$

Mathematica [F]

time = 78.30, size = 0, normalized size = 0.00

$$\int \frac{(a + b \operatorname{ArcTan}(cx))^2}{d + ex} dx$$

Verification is not applicable to the result.

[In] `Integrate[(a + b*ArcTan[c*x])^2/(d + e*x), x]`
[Out] `Integrate[(a + b*ArcTan[c*x])^2/(d + e*x), x]`

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 2.90, size = 1324, normalized size = 5.94

method	result	size
derivativedivides	Expression too large to display	1324
default	Expression too large to display	1324

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x))^2/(e*x+d), x, method=_RETURNVERBOSE)`
[Out]
$$\begin{aligned} & 1/c*(a^2*c*ln(c*e*x+c*d)/e+b^2*c*ln(c*e*x+c*d)/e*arctan(c*x)^2-b^2*c/e*arctan(c*x)^2*ln(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)-1/2*I*b^2*c/e*arctan(c*x)^2*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))^\text{2}*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1))*Pi-I*b^2*c*arctan(c*x)*polylog(2, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)+I*a*b*c*ln(c*e*x+c*d)/e*ln((I*e-c*e*x)/(c*d+I*e))+1/2*I*b^2*c/e*arctan(c*x)^2*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))^\text{3}*Pi-I*a*b*c*ln(c*e*x+c*d)/e*ln((I*e+c*e*x)/(I*e-c*d))-1/2*b^2*c/e*polylog(3, -(1+I*c*x)^2/(c^2*x^2+1))+b^2*c^2/e*d/(-I*e+c*d)*arctan(c*x)^2*ln(1-(I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))+I*a*b*c/e*dilog((I*e-c*e*x)/(c*d+I*e))+1/2*b^2*c^2/e*d/(-I*e+c*d)*polylog(3, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))+b^2*c*arctan(c*x)^2*ln(1-(I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)+I*b^2*c/e*arctan(c*x)*polylog(2, -(1+I*c*x)^2/(c^2*x^2+1))+1/2*b^2*c*polylog(3, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)+2*a*b*c*ln(c*e*x+c*d)/e*arctan(c*x)-1/2*I*b^2*c/e*arctan(c*x)^2*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))^\text{2}*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))*Pi-I*a*b*c/e*dilog((I*e+c*e*x)/(I*e-c*d))-I*b^2*c^2/e*d/(-I*e+c*d)*arctan(c*x)*polylog(2, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))+1/2*I*b^2*c/e*arctan(c*x)^2*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1))*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))*Pi) \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))^2/(e*x+d),x, algorithm="maxima")`[Out] $a^2 e^{-1} \log(x e + d) + \text{integrate}(1/16*(12 b^2 \arctan(c x)^2 + b^2 \log(c x)^2 x^2 + 1)^2 + 32 a b \arctan(c x))/(x e + d), x)$ **Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))^2/(e*x+d),x, algorithm="fricas")`[Out] `integral(b^2 arctan(c x)^2 + 2 a b arctan(c x) + a^2)/(x e + d), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + b \operatorname{atan}(cx))^2}{d + ex} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x))**2/(e*x+d),x)`[Out] `Integral((a + b*atan(c*x))**2/(d + e*x), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))^2/(e*x+d),x, algorithm="giac")`[Out] `sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a + b \operatorname{atan}(cx))^2}{d + ex} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^(2/1)/(d + e*x),x)`[Out] `int((a + b*atan(c*x))^(2/1)/(d + e*x), x)`

$$3.13 \quad \int \frac{(a+b\text{ArcTan}(cx))^2}{(d+ex)^2} dx$$

Optimal. Leaf size=341

$$\frac{ic(a+b\text{ArcTan}(cx))^2}{c^2d^2+e^2} + \frac{c^2d(a+b\text{ArcTan}(cx))^2}{e(c^2d^2+e^2)} - \frac{(a+b\text{ArcTan}(cx))^2}{e(d+ex)} - \frac{2bc(a+b\text{ArcTan}(cx))\log(\frac{2}{1-icx})}{c^2d^2+e^2} +$$

[Out] $I*c*(a+b*\text{arctan}(c*x))^2/(c^2*d^2+e^2)+c^2*d*(a+b*\text{arctan}(c*x))^2/e/(c^2*d^2+e^2)-(a+b*\text{arctan}(c*x))^2/e/(e*x+d)-2*b*c*(a+b*\text{arctan}(c*x))*\ln(2/(1-I*c*x))/$
 $(c^2*d^2+e^2)+2*b*c*(a+b*\text{arctan}(c*x))*\ln(2/(1+I*c*x))/(c^2*d^2+e^2)+2*b*c*(a+b*\text{arctan}(c*x))*\ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)+I*b^2*c*\text{polylog}(2,1-2/(1-I*c*x))/(c^2*d^2+e^2)+I*b^2*c*\text{polylog}(2,1-2/(1+I*c*x))/(c^2*d^2+e^2)-I*b^2*c*\text{polylog}(2,1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)$

Rubi [A]

time = 0.26, antiderivative size = 341, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.500,
 Rules used = {4974, 4966, 2449, 2352, 2497, 5104, 5004, 5040, 4964}

$$\frac{ic(a+b\text{ArcTan}(cx))^2}{c^2d^2+e^2} + \frac{c^2d(a+b\text{ArcTan}(cx))^2}{e(c^2d^2+e^2)} - \frac{2bc\log(\frac{2}{1-icx})(a+b\text{ArcTan}(cx))}{c^2d^2+e^2} + \frac{2bc\log(\frac{2}{1+icx})(a+b\text{ArcTan}(cx))}{c^2d^2+e^2} + \frac{2bc(a+b\text{ArcTan}(cx))\log(\frac{2c(d+ex)}{(1-icx)(cd+ea)})}{c^2d^2+e^2} - \frac{(a+b\text{ArcTan}(cx))^2}{e(d+ex)} - \frac{ib^2c\text{Li}_2(1-\frac{2}{1-icx})}{c^2d^2+e^2} + \frac{ib^2c\text{Li}_2(1-\frac{2}{1+icx})}{c^2d^2+e^2} + \frac{ib^2c\text{Li}_2(1-\frac{2}{icx+1})}{c^2d^2+e^2} - \frac{ib^2c\text{Li}_2(1-\frac{2c(d+ex)}{(cd+ea)(1-icx)})}{c^2d^2+e^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a+b*\text{ArcTan}[c*x])^2/(d+e*x)^2, x]$

[Out] $(I*c*(a+b*\text{ArcTan}[c*x])^2)/(c^2*d^2+e^2) + (c^2*d*(a+b*\text{ArcTan}[c*x])^2)/$
 $(e*(c^2*d^2+e^2)) - (a+b*\text{ArcTan}[c*x])^2/(e*(d+e*x)) - (2*b*c*(a+b*\text{ArcTan}[c*x])*Log[2/(1-I*c*x)])/(c^2*d^2+e^2) + (2*b*c*(a+b*\text{ArcTan}[c*x])*Log[2/(1+I*c*x)])/(c^2*d^2+e^2) + (2*b*c*(a+b*\text{ArcTan}[c*x])*Log[(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))])/(c^2*d^2+e^2) + (I*b^2*c*\text{PolyLog}[2, 1-2/(1-I*c*x)])/(c^2*d^2+e^2) + (I*b^2*c*\text{PolyLog}[2, 1-2/(1+I*c*x)])/(c^2*d^2+e^2) - (I*b^2*c*\text{PolyLog}[2, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))])/(c^2*d^2+e^2)$

Rule 2352

$\text{Int}[\text{Log}[(c_*)(x_*)]/((d_*)+(e_*)(x_*)), x_{\text{Symbol}}] \Rightarrow \text{Simp}[(-e^{(-1)})*\text{PolyLog}[2, 1-c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \&& \text{EqQ}[e+c*d, 0]$

Rule 2449

$\text{Int}[\text{Log}[(c_*)/((d_*)+(e_*)(x_*))]/((f_*)+(g_*)(x_*)^2), x_{\text{Symbol}}] \Rightarrow \text{Dist}[-e/g, \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1-2*d*x), x], x, 1/(d+e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \&& \text{EqQ}[c, 2*d] \&& \text{EqQ}[e^2*f + d^2*g, 0]$

Rule 2497

```
Int[Log[u_]*(Pq_)^(m_), x_Symbol] :> With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simplify[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]]
```

Rule 4964

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_.) + (e_)*(x_)), x_Symbol]
  :> Simplify[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e), Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4966

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))/((d_.) + (e_)*(x_)), x_Symbol] :> Simplify[(-(a + b*ArcTan[c*x]))*(Log[2/(1 - I*c*x)]/e), x] + (Dist[b*(c/e), Int[Log[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e)*x)/(c*d + I*e)*(1 - I*c*x))]/(1 + c^2*x^2), x], x] + Simplify[(a + b*ArcTan[c*x])*(Log[2*c*((d + e)*x)/(c*d + I*e)*(1 - I*c*x))]/e), x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_)*((d_.) + (e_)*(x_))^q, x_Symbol] :> Simplify[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Dist[b*c*(p/(e*(q + 1))), Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_.) + (e_)*(x_)^2), x_Symbol] :> Simplify[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*(x_))/((d_.) + (e_)*(x_)^2), x_Symbol] :> Simplify[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*c*(p + 1))), x] - Dist[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((a_) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*((f_) + (g_)*(x_))^(m_.))/((d_) + (e_)*(x_)^2, x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a + b \tan^{-1}(cx))^2}{(d + ex)^2} dx &= -\frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} + \frac{(2bc) \int \left(\frac{e^2(a+b \tan^{-1}(cx))}{(c^2d^2+e^2)(d+ex)} + \frac{c^2(d-ex)(a+b \tan^{-1}(cx))}{(c^2d^2+e^2)(1+c^2x^2)} \right) dx}{e} \\
&= -\frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} + \frac{(2bc^3) \int \frac{(d-ex)(a+b \tan^{-1}(cx))}{1+c^2x^2} dx}{e(c^2d^2+e^2)} + \frac{(2bce) \int \frac{a+b \tan^{-1}(cx)}{d+ex} dx}{c^2d^2+e^2} \\
&= -\frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} - \frac{2bc(a + b \tan^{-1}(cx)) \log(\frac{2}{1-icx})}{c^2d^2+e^2} + \frac{2bc(a + b \tan^{-1}(cx))}{c^2d^2+e^2} \\
&= -\frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} - \frac{2bc(a + b \tan^{-1}(cx)) \log(\frac{2}{1-icx})}{c^2d^2+e^2} + \frac{2bc(a + b \tan^{-1}(cx))}{c^2d^2+e^2} \\
&= \frac{ic(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^2}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} - \frac{2bc(a + b \tan^{-1}(cx))}{e(d + ex)} \\
&= \frac{ic(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^2}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} - \frac{2bc(a + b \tan^{-1}(cx))}{e(d + ex)} \\
&= \frac{ic(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^2}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} - \frac{2bc(a + b \tan^{-1}(cx))}{e(d + ex)} \\
&= \frac{ic(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^2}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^2}{e(d + ex)} - \frac{2bc(a + b \tan^{-1}(cx))}{e(d + ex)}
\end{aligned}$$

Mathematica [A]

time = 1.96, size = 300, normalized size = 0.88

$$\frac{a^2}{e(d + ex)} + \frac{ab(-2(e - c^2dx) \operatorname{ArcTan}(cx) + c(d + ex)(2 \log(d + ex) - \log(1 + c^2x^2)))}{(c^2d^2 + e^2)(d + ex)} + \frac{b^2 \left(-\frac{e^{i \operatorname{ArcTan}(\frac{a}{d})} (\operatorname{ArcTan}(cx))^2 + \frac{e \operatorname{ArcTan}(cx)}{dx}}{\sqrt{1 + \frac{c^2d^2}{e^2}} x} + \frac{a \left(-i \left(x - i \operatorname{ArcTan}(\frac{a}{d}) \right) \operatorname{ArcTan}(cx) - i \log(1 + x \operatorname{ArcTan}(\frac{a}{d})) - 2 \left(\operatorname{ArcTan}(\frac{a}{d}) + \operatorname{ArcTan}(cx) \right) \log \left(1 - x \left(\operatorname{ArcTan}(\frac{a}{d}) + \operatorname{ArcTan}(cx) \right) \right) - \frac{1}{2} \log(1 + x^2) + i \operatorname{ArcTan}(\frac{a}{d}) \log(x \operatorname{ArcTan}(\frac{a}{d}) + \operatorname{ArcTan}(cx)) + \text{PolyLog} \left(z, x^2 \left(\operatorname{ArcTan}(\frac{a}{d}) + \operatorname{ArcTan}(cx) \right) \right) \right)}{c^2d^2 + e^2} \right)}{d}$$

Antiderivative was successfully verified.

[In] `Integrate[(a + b*ArcTan[c*x])^2/(d + e*x)^2, x]`

[Out] $-(a^2/(e*(d + e*x))) + (a*b*(-2*(e - c^2*d*x)*ArcTan[c*x] + c*(d + e*x)*(2*Log[c*(d + e*x)] - Log[1 + c^2*x^2])))/((c^2*d^2 + e^2)*(d + e*x)) + (b^2*((E^(I*ArcTan[(c*d)/e])*ArcTan[c*x]^2)/(Sqrt[1 + (c^2*d^2)/e^2]*e)) + (x*A$

$$\text{rcTan}[c*x]^2/(d + e*x) - (c*d*((-I)*(Pi - 2*ArcTan[(c*d)/e])*ArcTan[c*x] - Pi*Log[1 + E^((-2*I)*ArcTan[c*x])] - 2*(ArcTan[(c*d)/e] + ArcTan[c*x])*Log[1 - E^((2*I)*(ArcTan[(c*d)/e] + ArcTan[c*x]))]) - (Pi*Log[1 + c^2*x^2])/2 + 2*ArcTan[(c*d)/e]*Log[Sin[ArcTan[(c*d)/e] + ArcTan[c*x]]] + I*PolyLog[2, E^((2*I)*(ArcTan[(c*d)/e] + ArcTan[c*x]))]))/(c^2*d^2 + e^2))/d$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 739 vs. 2(329) = 658.

time = 2.34, size = 740, normalized size = 2.17

method	result
derivativedivides	$-\frac{a^2 c^2}{(cex+cd)e} - \frac{b^2 c^2 \arctan(cx)^2}{(cex+cd)e} - \frac{b^2 c^2 \arctan(cx) \ln(c^2 x^2 + 1)}{c^2 d^2 + e^2} + \frac{b^2 c^3 d \arctan(cx)^2}{e (c^2 d^2 + e^2)} + \frac{2 b^2 c^2 \arctan(cx) \ln(cex+cd)}{c^2 d^2 + e^2} + \frac{i b^2 c^2 \ln(cex+cd)}{c^2 d^2 + e^2}$
default	$-\frac{a^2 c^2}{(cex+cd)e} - \frac{b^2 c^2 \arctan(cx)^2}{(cex+cd)e} - \frac{b^2 c^2 \arctan(cx) \ln(c^2 x^2 + 1)}{c^2 d^2 + e^2} + \frac{b^2 c^3 d \arctan(cx)^2}{e (c^2 d^2 + e^2)} + \frac{2 b^2 c^2 \arctan(cx) \ln(cex+cd)}{c^2 d^2 + e^2} + \frac{i b^2 c^2 \ln(cex+cd)}{c^2 d^2 + e^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a+b*arctan(c*x))^2/(e*x+d)^2,x,method=_RETURNVERBOSE)

[Out]
$$\begin{aligned} & 1/c*(-a^2*c^2/(c*e*x+c*d)/e-b^2*c^2/(c*e*x+c*d)/e*arctan(c*x)^2-b^2*c^2*arc \\ & tan(c*x)/(c^2*d^2+e^2)*ln(c^2*x^2+1)+b^2*c^3/e/(c^2*d^2+e^2)*d*arctan(c*x)^2 \\ & +2*b^2*c^2*arctan(c*x)/(c^2*d^2+e^2)*ln(c*e*x+c*d)+1/2*I*b^2*c^2/(c^2*d^2+e^2) \\ & *ln(c*x+I)*ln(c^2*x^2+1)+1/2*I*b^2*c^2/(c^2*d^2+e^2)*dilog(-1/2*I*(c*x+I)) \\ & -1/2*I*b^2*c^2/(c^2*d^2+e^2)*ln(c*x-I)*ln(c^2*x^2+1)+1/4*I*b^2*c^2/(c^2*d^2+e^2) \\ & *ln(c*x-I)^2-I*b^2*c^2/(c^2*d^2+e^2)*dilog((I*e+c*e*x)/(I*e-c*d))-1/2*I*b^2*c^2/(c^2*d^2+e^2)*dilog(1/2*I*(c*x-I))+1/2*I*b^2*c^2/(c^2*d^2+e^2) \\ & *ln(c*x-I)*ln(-1/2*I*(c*x+I))+I*b^2*c^2/(c^2*d^2+e^2)*dilog((I*e-c*e*x)/(c*d+I*e))-1/2*I*b^2*c^2/(c^2*d^2+e^2)*ln(c*x+I)*ln(1/2*I*(c*x-I))+I*b^2*c^2/(c^2*d^2+e^2)*ln(c*e*x+c*d)*ln((I*e-c*e*x)/(c*d+I*e))-1/4*I*b^2*c^2/(c^2*d^2+e^2)*ln(c*x+I)^2-I*b^2*c^2/(c^2*d^2+e^2)*ln(c*e*x+c*d)*ln((I*e+c*e*x)/(I*e-c*d))-2*a*b*c^2/(c*e*x+c*d)/e*arctan(c*x)-a*b*c^2/(c^2*d^2+e^2)*ln(c^2*x^2+1)+2*a*b*c^3/e/(c^2*d^2+e^2)*d*arctan(c*x)+2*a*b*c^2/(c^2*d^2+e^2)*ln(c*e*x+c*d)) \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a+b*arctan(c*x))^2/(e*x+d)^2,x, algorithm="maxima")

[Out]
$$((2*c*d*arctan(c*x)/(c^2*d^2*2*e + e^3) - log(c^2*x^2 + 1)/(c^2*d^2 + e^2) + 2*log(x*e + d)/(c^2*d^2 + e^2))*c - 2*arctan(c*x)/(x*e^2 + d*e))*a*b - 1/16$$

```
*(4*arctan(c*x)^2 - 16*(x*e^2 + d*e)*integrate(1/16*(12*(c^2*x^2*e + e)*arctan(c*x)^2 + (c^2*x^2*e + e)*log(c^2*x^2 + 1)^2 + 8*(c*x*e + c*d)*arctan(c*x) - 4*(c^2*x^2*e + c^2*d*x)*log(c^2*x^2 + 1))/(c^2*x^4*e^3 + 2*c^2*d*x^3*e^2 + (c^2*d^2*e + e^3)*x^2 + 2*d*x*e^2 + d^2*e), x) - log(c^2*x^2 + 1)^2)*b^2/(x*e^2 + d*e) - a^2/(x*e^2 + d*e)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^2/(e*x+d)^2,x, algorithm="fricas")
[Out] integral((b^2*arctan(c*x)^2 + 2*a*b*arctan(c*x) + a^2)/(x^2*e^2 + 2*d*x*e + d^2), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + b \operatorname{atan}(cx))^2}{(d + ex)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*atan(c*x))**2/(e*x+d)**2,x)
[Out] Integral((a + b*atan(c*x))**2/(d + e*x)**2, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^2/(e*x+d)^2,x, algorithm="giac")
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a + b \operatorname{atan}(cx))^2}{(d + e x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a + b*atan(c*x))^2/(d + e*x)^2,x)
[Out] int((a + b*atan(c*x))^2/(d + e*x)^2, x)
```

$$3.14 \quad \int \frac{(a+b\text{ArcTan}(cx))^2}{(d+ex)^3} dx$$

Optimal. Leaf size=496

$$\frac{b^2 c^3 d \text{ArcTan}(cx)}{(c^2 d^2 + e^2)^2} - \frac{b c (a + b \text{ArcTan}(cx))}{(c^2 d^2 + e^2) (d + ex)} + \frac{i c^3 d (a + b \text{ArcTan}(cx))^2}{(c^2 d^2 + e^2)^2} + \frac{c^2 (cd - e) (cd + e) (a + b \text{ArcTan}(cx))^2}{2 e (c^2 d^2 + e^2)^2}$$

[Out] $b^{2+}c^{3+d}\arctan(c*x)/(c^{2+d^2+e^2})^{2-b*c*(a+b*\arctan(c*x))/(c^{2+d^2+e^2})/(e*x+d)+I*c^{3+d*(a+b*\arctan(c*x))^{2/(c^{2+d^2+e^2})^{2+1/2*c^{2*(c*d-e)*(c*d+e)*}}(a+b*\arctan(c*x))^{2/e}/(c^{2+d^2+e^2})^{2-1/2*(a+b*\arctan(c*x))^{2/e}/(e*x+d)^{2-2*b*c^{3+d*(a+b*\arctan(c*x))*ln(2/(1-I*c*x))/(c^{2+d^2+e^2})^{2+2*b*c^{3+d*(a+b*\arctan(c*x))*ln(2/(1+I*c*x))/(c^{2+d^2+e^2})^{2+b*c^{2*2*e*ln(e*x+d)/(c^{2+d^2+e^2})^{2+2*b*c^{3+d*(a+b*\arctan(c*x))*ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^{2+d^2+e^2})^{2-1/2*b^{2*c^{2*2*e*ln(c^{2*x^2+1})/(c^{2+d^2+e^2})^{2+I*b^{2*c^{3+d*polylog(2,1-2/(1-I*c*x)))/(c^{2+d^2+e^2})^{2+I*b^{2*c^{3+d*polylog(2,1-2/(1+I*c*x)))/(c^{2+d^2+e^2})^{2-I*b^{2*c^{3+d*polylog(2,1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^{2+d^2+e^2})^{2}}$

Rubi [A]

time = 0.39, antiderivative size = 496, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 15, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {4974, 4972, 720, 31, 649, 209, 266, 4966, 2449, 2352, 2497, 5104, 5004, 5040, 4964}

$$\frac{c^2 (cd - e) (cd + e) (a + b \text{ArcTan}(cx))^2}{2 e (c^2 d^2 + e^2)^2} - \frac{b c (a + b \text{ArcTan}(cx))}{(c^2 d^2 + e^2) (d + ex)} + \frac{i c^3 d (a + b \text{ArcTan}(cx))^2}{(c^2 d^2 + e^2)^2} - \frac{2 b^2 d \log \left(\frac{c^2 d}{c^2 d^2 + e^2}\right) (a + b \text{ArcTan}(cx))}{(c^2 d^2 + e^2)^2} + \frac{2 b c^2 d \log \left(\frac{c^2 d}{c^2 d^2 + e^2}\right) (a + b \text{ArcTan}(cx)) \log \left(\frac{2 i (d + e x)}{(1 - e x) (d + e x)}\right)}{(c^2 d^2 + e^2)^2} - \frac{(a + b \text{ArcTan}(cx))^2}{2 e (d + ex)^2} + \frac{i b^2 c^2 d \text{ArcTan}(cx)}{(c^2 d^2 + e^2)^2} - \frac{b^2 c^2 e \log (c^2 d^2 + 1)}{2 (c^2 d^2 + e^2)^2} + \frac{i b^2 c^2 e \log (d + ex)}{(c^2 d^2 + e^2)^2} + \frac{b^2 c^2 d \text{Li}_2 \left(1 - \frac{c^2 d}{d + e x}\right)}{(c^2 d^2 + e^2)^2} + \frac{i b^2 c^2 d \text{Li}_2 \left(1 - \frac{2 i (d + e x)}{(1 - e x) (d + e x)}\right)}{(c^2 d^2 + e^2)^2} - \frac{i b^2 c^2 d \text{Li}_2 \left(1 - \frac{2 i (d + e x)}{(1 + e x) (d + e x)}\right)}{(c^2 d^2 + e^2)^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x])^2/(d + e*x)^3, x]$

[Out] $(b^{2+}c^{3+d}\text{ArcTan}[c*x])/((c^{2+d^2+e^2})^{2-(b*c*(a+b*\text{ArcTan}[c*x]))/((c^{2+d^2+e^2})^2*(d+e*x))+(I*c^{3+d*(a+b*\text{ArcTan}[c*x])^2}/(c^{2+d^2+e^2})^{2+(c^{2*(c*d-e)*(c*d+e)*(a+b*\text{ArcTan}[c*x])^2})/(2*e*(c^{2+d^2+e^2})^{2-(a+b*\text{ArcTan}[c*x])^2/(2*e*(d+e*x)^2)-(2*b*c^{3+d*(a+b*\text{ArcTan}[c*x])*Log[2/(1-I*c*x)])/(c^{2+d^2+e^2})^{2+(2*b*c^{3+d*(a+b*\text{ArcTan}[c*x])*Log[2/(1+I*c*x)])/(c^{2+d^2+e^2})^{2+(b^{2+}c^{2+e*Log[d+e*x]}/(c^{2+d^2+e^2})^{2+(2*b*c^{3+d*(a+b*\text{ArcTan}[c*x])*Log[(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))]/(c^{2+d^2+e^2})^{2-(b^{2+}c^{2+e*Log[1+c^{2+2*x^2}]})/(2*(c^{2+d^2+e^2})^{2+(I*b^{2+}c^{3+d*PolyLog[2,1-2/(1-I*c*x)])/(c^{2+d^2+e^2})^{2+(I*b^{2+}c^{3+d*PolyLog[2,1-2/(1+I*c*x)])/(c^{2+d^2+e^2})^{2-(I*b^{2+}c^{3+d*PolyLog[2,1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))])/(c^{2+d^2+e^2})^{2}}$

Rule 31

$\text{Int}[(a_+ + b_-)*(x_-)^{(-1)}, x_{\text{Symbol}}] := \text{Simp}[\text{Log}[\text{RemoveContent}[a_+ + b_*x, x]/b, x] /; \text{FreeQ}[\{a, b\}, x]$

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 649

```
Int[((d_) + (e_.)*(x_))/((a_) + (c_.)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 720

```
Int[1/(((d_) + (e_.)*(x_))*(a_) + (c_.)*(x_)^2), x_Symbol] :> Dist[e^2/(c*d^2 + a*e^2), Int[1/(d + e*x), x], x] + Dist[1/(c*d^2 + a*e^2), Int[(c*d - c*e*x)/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0]
```

Rule 2352

```
Int[Log[(c_.)*(x_)]/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_.)/((d_) + (e_.)*(x_))]/((f_) + (g_.)*(x_)^2), x_Symbol] :> Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 2497

```
Int[Log[u_]*(Pq_)^(m_), x_Symbol] :> With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u, x][[2]], Expon[Pq, x]]]
```

Rule 4964

```
Int[((a_) + ArcTan[(c_.)*(x_)*(b_.))^(p_.)/((d_) + (e_.)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e), Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)),
```

```
x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4966

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))/((d_) + (e_.)*(x_)), x_Symbol] :> Si
mp[(-(a + b*ArcTan[c*x]))*(Log[2/(1 - I*c*x)]/e), x] + (Dist[b*(c/e), Int[L
og[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e
*x)/((c*d + I*e)*(1 - I*c*x)))]/(1 + c^2*x^2), x], x] + Simp[(a + b*ArcTan[
c*x])*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x]) /; FreeQ[{a,
b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rule 4972

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))*((d_) + (e_.*(x_))^(q_.), x_Symbol]
:> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])/((e*(q + 1))), x] - Dist[b*(c/
(e*(q + 1))), Int[(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b,
c, d, e, q}, x] && NeQ[q, -1]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_)*((d_) + (e_.*(x_))^(q_.), x_Sy
mbol] :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - D
ist[b*c*(p/(e*(q + 1))), Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1),
(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] &&
IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.)/((d_) + (e_.*(x_))^2), x_Symbo
l] :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b,
c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.)*((d_) + (e_.*(x_))^2),
x_Symbol] :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Di
st[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.)*((f_) + (g_.*(x_))^(m_.)))/((d_
) + (e_.*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p
/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGt
Q[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a + b \tan^{-1}(cx))^2}{(d + ex)^3} dx &= -\frac{(a + b \tan^{-1}(cx))^2}{2e(d + ex)^2} + \frac{(bc) \int \left(\frac{e^2(a+b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d+ex)^2} + \frac{2c^2 de^2(a+b \tan^{-1}(cx))}{(c^2 d^2 + e^2)^2(d+ex)} + \frac{(c^4 d^2 - c^2 e^2 - 2c^2 d e^2)}{(c^2 d^2 + e^2)^2} \right) dx}{e} \\
&= -\frac{(a + b \tan^{-1}(cx))^2}{2e(d + ex)^2} + \frac{(bc) \int \frac{(c^4 d^2 - c^2 e^2 - 2c^2 d e^2)(a+b \tan^{-1}(cx))}{1+c^2 x^2} dx}{e (c^2 d^2 + e^2)^2} + \frac{(2bc^3 de) \int \frac{a+b \tan^{-1}(cx)}{c^2 d^2 + e^2} dx}{(c^2 d^2 + e^2)^2} \\
&= -\frac{bc(a + b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d + ex)} - \frac{(a + b \tan^{-1}(cx))^2}{2e(d + ex)^2} - \frac{2bc^3 d(a + b \tan^{-1}(cx)) \log(\frac{2}{1-icx})}{(c^2 d^2 + e^2)^2} \\
&= -\frac{bc(a + b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d + ex)} - \frac{(a + b \tan^{-1}(cx))^2}{2e(d + ex)^2} - \frac{2bc^3 d(a + b \tan^{-1}(cx)) \log(\frac{2}{1-icx})}{(c^2 d^2 + e^2)^2} \\
&= -\frac{bc(a + b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d + ex)} + \frac{ic^3 d(a + b \tan^{-1}(cx))^2}{(c^2 d^2 + e^2)^2} + \frac{c^2(cd - e)(cd + e)(a + b \tan^{-1}(cx))}{2e(c^2 d^2 + e^2)^2} \\
&= \frac{b^2 c^3 d \tan^{-1}(cx)}{(c^2 d^2 + e^2)^2} - \frac{bc(a + b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d + ex)} + \frac{ic^3 d(a + b \tan^{-1}(cx))^2}{(c^2 d^2 + e^2)^2} + \frac{c^2(cd - e)(cd + e)(a + b \tan^{-1}(cx))}{2e(c^2 d^2 + e^2)^2} \\
&= \frac{b^2 c^3 d \tan^{-1}(cx)}{(c^2 d^2 + e^2)^2} - \frac{bc(a + b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d + ex)} + \frac{ic^3 d(a + b \tan^{-1}(cx))^2}{(c^2 d^2 + e^2)^2} + \frac{c^2(cd - e)(cd + e)(a + b \tan^{-1}(cx))}{2e(c^2 d^2 + e^2)^2} \\
&= \frac{b^2 c^3 d \tan^{-1}(cx)}{(c^2 d^2 + e^2)^2} - \frac{bc(a + b \tan^{-1}(cx))}{(c^2 d^2 + e^2)(d + ex)} + \frac{ic^3 d(a + b \tan^{-1}(cx))^2}{(c^2 d^2 + e^2)^2} + \frac{c^2(cd - e)(cd + e)(a + b \tan^{-1}(cx))}{2e(c^2 d^2 + e^2)^2}
\end{aligned}$$

Mathematica [A]

time = 4.05, size = 479, normalized size = 0.97

$$\frac{x^3}{2(d + ex)^2} + \frac{ib(-c^2 + e^2)(d + ex) - e^2(c^2d^2 + 2de + e^2)(a) \operatorname{ArcTan}(ax) + (cd + ex)(-c^2d^2 - e^2 + 2c^2(d + ex)\log(d + ex) - c^2(d + ex)\log(1 + e^2x^2))}{2(d + ex)^2(d + ex)^2} + \frac{\frac{ib(-c^2 + e^2)(d + ex)(a) \operatorname{ArcTan}(ax) + (cd + ex)(-c^2d^2 - e^2 + 2c^2(d + ex)\log(d + ex) - c^2(d + ex)\log(1 + e^2x^2))}{2(d + ex)^2(d + ex)^2} + \frac{ib(-c^2 + e^2)(d + ex)(a) \operatorname{ArcTan}(ax) + (cd + ex)(-c^2d^2 - e^2 + 2c^2(d + ex)\log(d + ex) - c^2(d + ex)\log(1 + e^2x^2))}{2(d + ex)^2(d + ex)^2}}{2(d + ex)^2(d + ex)^2}$$

Antiderivative was successfully verified.

[In] `Integrate[(a + b*ArcTan[c*x])^2/(d + e*x)^3, x]`

[Out]
$$\begin{aligned}
&-1/2*a^2/(e*(d + e*x)^2) + (a*b*((-e^3 + c^4*d^2*x^2*(2*d + e*x) - c^2*e*(3*d^2 + 2*d*e*x + e^2*x^2))*\operatorname{ArcTan}[c*x] + c*(d + e*x)*(-(c^2*d^2) - e^2 + 2*c^2*d*(d + e*x)*\operatorname{Log}[c*(d + e*x)] - c^2*d*(d + e*x)*\operatorname{Log}[1 + c^2*x^2])))/((c^2*d^2 + e^2)^2*(d + e*x)^2) + (b^2*c^2*((-2*E^(I*\operatorname{ArcTan}[(c*d)/e])*c*\operatorname{ArcTan}[c*x]^2)/(Sqrt[1 + (c^2*d^2)/e^2]*e) - (e*(1 + c^2*x^2)*\operatorname{ArcTan}[c*x]^2)/(c^2*(d + e*x)^2) + (2*x*\operatorname{ArcTan}[c*x]*(e + c*d*\operatorname{ArcTan}[c*x]))/(c*d*(d + e*x)) + (-2*e^2*\operatorname{ArcTan}[c*x] + 2*c*d*e*\operatorname{Log}[(c*(d + e*x))/Sqrt[1 + c^2*x^2]]))/(c^3*d^3 + c*
\end{aligned}$$

$$d e^2) - (2 c d ((-I) (\Pi - 2 \operatorname{ArcTan}[(c d)/e]) \operatorname{ArcTan}[c x] - \Pi \log[1 + e^{(-2 I) \operatorname{ArcTan}[c x]}]) - 2 (\operatorname{ArcTan}[(c d)/e] + \operatorname{ArcTan}[c x]) \log[1 - e^{((2 I) (\operatorname{ArcTan}[(c d)/e] + \operatorname{ArcTan}[c x])}] - (\Pi \log[1 + c^2 x^2])/2 + 2 \operatorname{ArcTan}[(c d)/e] \log[\sin[\operatorname{ArcTan}[(c d)/e] + \operatorname{ArcTan}[c x]]] + I \operatorname{PolyLog}[2, e^{((2 I) (\operatorname{ArcTan}[(c d)/e] + \operatorname{ArcTan}[c x]))}])/(c^2 d^2 + e^2))/(2 (c^2 d^2 + e^2))$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 964 vs. 2(478) = 956.

time = 5.14, size = 965, normalized size = 1.95 Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a+b*arctan(c*x))^2/(e*x+d)^3,x,method=_RETURNVERBOSE)
[Out] 1/c*(-a*b*c^3/(c*e*x+c*d)^2/e*arctan(c*x)+I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*e*x+c*d)*ln((I*e-c*e*x)/(c*d+I*e))-1/2*I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*x+I)*ln(1/2*I*(c*x-I))+1/2*I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*x+I)*ln(c^2*x^2+1)-I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*e*x+c*d)*ln((I*e+c*e*x)/(I*e-c*d))-1/2*I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*x-I)*ln(c^2*x^2+1)+1/2*I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*x-I)*ln(-1/2*I*(c*x+I))+a*b*c^5/e/(c^2*d^2+e^2)^2*arctan(c*x)*d^2-a*b*c^4/(c^2*d^2+e^2)^2*d*ln(c^2*x^2+1)-a*b*c^3*e/(c^2*d^2+e^2)^2*arctan(c*x)+2*a*b*c^4*d/(c^2*d^2+e^2)^2*ln(c*e*x+c*d)+1/2*b^2*c^5/e/(c^2*d^2+e^2)^2*arctan(c*x)^2*d^2-b^2*c^4*arctan(c*x)/(c^2*d^2+e^2)^2*d*ln(c^2*x^2+1)+2*b^2*c^4*arctan(c*x)*d/(c^2*d^2+e^2)^2*ln(c*e*x+c*d)+I*b^2*c^4/(c^2*d^2+e^2)^2*d*dilog((I*e-c*e*x)/(c*d+I*e))-1/4*I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*x+I)^2-1/2*I*b^2*c^4/(c^2*d^2+e^2)^2*d*dilog(1/2*I*(c*x-I))-I*b^2*c^4/(c^2*d^2+e^2)^2*d*dilog((I*e+c*e*x)/(I*e-c*d))+1/4*I*b^2*c^4/(c^2*d^2+e^2)^2*d*ln(c*x-I)^2+1/2*I*b^2*c^4/(c^2*d^2+e^2)^2*d*dilog(-1/2*I*(c*x+I))-a*b*c^3/(c^2*d^2+e^2)/(c*e*x+c*d)-1/2*b^2*c^3/(c*e*x+c*d)^2/e*arctan(c*x)^2-1/2*b^2*c^3*e/(c^2*d^2+e^2)^2*ln(c^2*x^2+1)+b^2*c^4/(c^2*d^2+e^2)^2*d*arctan(c*x)+b^2*c^3*e/(c^2*d^2+e^2)^2*ln(c*e*x+c*d)-1/2*a^2*c^3/(c*e*x+c*d)^2/e)
```

Maxima [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^2/(e*x+d)^3,x, algorithm="maxima")
```

```
[Out] Timed out
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^2/(e*x+d)^3,x, algorithm="fricas")
[Out] integral((b^2*arctan(c*x)^2 + 2*a*b*arctan(c*x) + a^2)/(x^3*e^3 + 3*d*x^2*e^2 + 3*d^2*x*e + d^3), x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*atan(c*x))**2/(e*x+d)**3,x)
```

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^2/(e*x+d)^3,x, algorithm="giac")
```

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a + b \operatorname{atan}(c x))^2}{(d + e x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a + b*atan(c*x))^2/(d + e*x)^3,x)
```

[Out] int((a + b*atan(c*x))^2/(d + e*x)^3, x)

$$3.15 \quad \int (d + ex)^3 (a + b \operatorname{ArcTan}(cx))^3 dx$$

Optimal. Leaf size=652

$$\frac{3ab^2de^2x}{c^2} - \frac{b^3e^3x}{4c^3} + \frac{b^3e^3 \operatorname{ArcTan}(cx)}{4c^4} + \frac{3b^3de^2x \operatorname{ArcTan}(cx)}{c^2} + \frac{b^2e^3x^2(a + b \operatorname{ArcTan}(cx))}{4c^2} - \frac{3bde^2(a + b \operatorname{ArcTan}(cx))}{2c^3}$$

[Out]
$$\begin{aligned} & 3*a*b^2*d*e^2*x/c^2 - 1/4*b^3*e^3*x/c^3 + 1/4*b^3*e^3*\arctan(cx)/c^4 + 3*b^3*d*e^2*x*\arctan(cx)/c^2 + 3*b^3*d*e^2*x*\arctan(cx)/c^3 + 3*I*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))*\text{polylog}(2,1) \\ & - 2/(1+I*c*x))/c^3 - 3/4*I*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^4 - 3/4*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^3 - 3/4*(c^4*d^4 - 6*c^2*d^2*e^2)*(\arctan(cx))^3/c^4 + e^{1/4}*(e*x+d)^4*(a+b*\arctan(cx))^3/e^{1/2} + 2*b^2*d^2*x^2*(a+b*\arctan(cx))^2/c^3 - 3/4*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^2 + I*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^1 - 3/4*(c^4*d^4 - 6*c^2*d^2*e^2)*(\arctan(cx))^3/c^2 + 2*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^1 - 3/4*I*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^0 + 2*b^2*d*(c*d-e)*(c*d+e)*\ln(2/(1+I*c*x))/c^4 - 3/2*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2*\ln(2/(1+I*c*x))/c^3 - 3/2*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2*\ln(2/(1+I*c*x))/c^2 - 3/4*I*b^2*d*(c*d-e)*(c*d+e)*\text{polylog}(2,1) - 2/(1+I*c*x))/c^4 + 1/4*I*b^2*d*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^2/c^3 - 3/2*b^2*d*(c*d-e)*(c*d+e)*\text{polylog}(3,1) - 2/(1+I*c*x))/c^3 \end{aligned}$$

Rubi [A]

time = 0.82, antiderivative size = 652, normalized size of antiderivative = 1.00, number of steps used = 29, number of rules used = 15, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {4974, 4930, 5040, 4964, 2449, 2352, 4946, 5036, 266, 5004, 327, 209, 5104, 5114, 6745}

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)^3 * (a + b*\operatorname{ArcTan}[c*x])^3, x]$

[Out]
$$\begin{aligned} & (3*a*b^2*d*e^2*x)/c^2 - (b^3*e^3*x)/(4*c^3) + (b^3*e^3*\operatorname{ArcTan}[c*x])/(4*c^4) \\ & + (3*b^3*d*e^2*x*\operatorname{ArcTan}[c*x])/c^2 + (b^2*e^3*x^2*(a + b*\operatorname{ArcTan}[c*x]))/(4*c^2) \\ & - (3*b^2*d*e^2*(a + b*\operatorname{ArcTan}[c*x])^2)/(2*c^3) + ((I/4)*b^2*e^3*(a + b*\operatorname{ArcTan}[c*x])^2)/c^4 \\ & - (((3*I)/4)*b^2*e^3*(6*c^2*d^2 - e^2)*(a + b*\operatorname{ArcTan}[c*x])^2)/c^4 \\ & - (3*b^2*e^3*(6*c^2*d^2 - e^2)*(a + b*\operatorname{ArcTan}[c*x])^2)/(4*c^3) - (3*b^2*d^2*x^2*(a + b*\operatorname{ArcTan}[c*x])^2)/(2*c) \\ & - (b^2*e^3*x^3*(a + b*\operatorname{ArcTan}[c*x])^2)/(4*c) + (I*d*(c*d - e)*(c*d + e)*(a + b*\operatorname{ArcTan}[c*x])^3)/c^3 - ((c^4*d^4 - 6*c^2*d^2 \\ & - 2*e^2 + e^4)*(a + b*\operatorname{ArcTan}[c*x])^3)/(4*c^4) + ((d + e*x)^4*(a + b*\operatorname{ArcTan}[c*x])^3)/(4*c^4) \\ & + (b^2*e^3*(a + b*\operatorname{ArcTan}[c*x])*Log[2/(1 + I*c*x)])/(2*c^4) - (3*b^2*e^3*(6*c^2*d^2 - e^2)*(a + b*\operatorname{ArcTan}[c*x])*Log[2/(1 + I*c*x)])/(2*c^4) \\ & + (3*b^2*d*(c*d - e)*(c*d + e)*(a + b*\operatorname{ArcTan}[c*x])^2*Log[2/(1 + I*c*x)])/(c^3) - (3*b^3*d^2*e^2*Log[1 + c^2*x^2])/(2*c^3) + ((I/4)*b^3*e^3*PolyLog[2, 1 - 2/(1 + I*c*x)])/(c^4) - (((3*I)/4)*b^3*e^3*(6*c^2*d^2 - e^2)*PolyLog[2, 1 - 2/(1 + I*c*x)])/(c^4) \end{aligned}$$

$$\frac{1 + I c x)]/c^4 + ((3 I) b^2 d (c d - e) (c d + e) (a + b \operatorname{ArcTan}[c x]) \operatorname{PolyLog}[2, 1 - 2/(1 + I c x)]/c^3 + (3 b^3 d (c d - e) (c d + e)) \operatorname{PolyLog}[3, 1 - 2/(1 + I c x)]/(2 c^3)$$
Rule 209

$$\operatorname{Int}[(a_.) + (b_.) (x_)^2)^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(Rt[a, 2] * Rt[b, 2])) * A \operatorname{rcTan}[Rt[b, 2] * (x/Rt[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{PosQ}[a/b] \& (\operatorname{GtQ}[a, 0] \mid\mid \operatorname{GtQ}[b, 0])$$
Rule 266

$$\operatorname{Int}[(x_)^{m_.}/((a_.) + (b_.) (x_)^{n_.}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a + b x^n, x]]/(b n), x] /; \operatorname{FreeQ}[\{a, b, m, n\}, x] \& \operatorname{EqQ}[m, n - 1]$$
Rule 327

$$\operatorname{Int}[(c_.) (x_)^{m_.} ((a_.) + (b_.) (x_)^{n_.})^{p_.}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[c^{(n - 1)} (c x)^{(m - n + 1)} ((a + b x^n)^{p + 1})/(b (m + n p + 1)), x] - \operatorname{Dist}[a c^{n_*} ((m - n + 1)/(b (m + n p + 1))), \operatorname{Int}[(c x)^{(m - n)} (a + b x^n)^p, x], x] /; \operatorname{FreeQ}[\{a, b, c, p\}, x] \& \operatorname{IGtQ}[n, 0] \& \operatorname{GtQ}[m, n - 1] \& \operatorname{NeQ}[m + n p + 1, 0] \& \operatorname{IntBinomialQ}[a, b, c, n, m, p, x]$$
Rule 2352

$$\operatorname{Int}[\operatorname{Log}[(c_.) (x_)]/((d_.) + (e_.) (x_)), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(-e^{-1}) \operatorname{PolyLog}[2, 1 - c x], x] /; \operatorname{FreeQ}[\{c, d, e\}, x] \& \operatorname{EqQ}[e + c d, 0]$$
Rule 2449

$$\operatorname{Int}[\operatorname{Log}[(c_.)/((d_.) + (e_.) (x_))] / ((f_.) + (g_.) (x_)^2), x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[-e/g, \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[2 d x]/(1 - 2 d x), x], x, 1/(d + e x)], x] /; \operatorname{FreeQ}[\{c, d, e, f, g\}, x] \& \operatorname{EqQ}[c, 2 d] \& \operatorname{EqQ}[e^{2 f} + d^{2 g}, 0]$$
Rule 4930

$$\operatorname{Int}[(a_.) + \operatorname{ArcTan}[(c_.) (x_)^{n_.}] (b_.)^{p_.}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[x (a + b \operatorname{ArcTan}[c x^n])^p, x] - \operatorname{Dist}[b c^{n_*} p, \operatorname{Int}[x^{n_*} ((a + b \operatorname{ArcTan}[c x^n])^{p - 1})/(1 + c^{2 n}), x], x] /; \operatorname{FreeQ}[\{a, b, c, n\}, x] \& \operatorname{IGtQ}[p, 0] \& (\operatorname{EqQ}[n, 1] \mid\mid \operatorname{EqQ}[p, 1])$$
Rule 4946

$$\operatorname{Int}[(a_.) + \operatorname{ArcTan}[(c_.) (x_)^{n_.}] (b_.)^{p_.}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[x^{(m + 1)} ((a + b \operatorname{ArcTan}[c x^n])^p)/(m + 1), x] - \operatorname{Dist}[b c^{n_*} p/(m + 1), \operatorname{Int}[x^{(m + n)} ((a + b \operatorname{ArcTan}[c x^n])^{p - 1})/(1 + c^{2 n}), x], x] /; \operatorname{FreeQ}[\{a, b, c, m, n\}, x] \& \operatorname{IGtQ}[p, 0] \& (\operatorname{EqQ}[p, 1] \mid\mid \operatorname{EqQ}[n, 1])$$

```
IntegerQ[m])) && NeQ[m, -1]
```

Rule 4964

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)), x_Symbol]
  :> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e),
  Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)),
  x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_)*((d_) + (e_)*(x_))^(q_.), x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Dist[b*c*(p/(e*(q + 1))),
  Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5036

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*((f_)*(x_))^(m_.))/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Dist[f^2/e, Int[(f*x)^(m - 2)*(a + b*ArcTan[c*x])^p, x], x] - Dist[d*(f^2/e), Int[(f*x)^(m - 2)*((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Dist[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*((f_) + (g_)*(x_))^(m_.))/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rule 5114

```

Int[(Log[u_]*((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.))/((d_) + (e_.)*(x_)^2),
  x_Symbol] :> Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)),
  x] + Dist[b*p*(I/2), Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]

```

Rule 6745

```

Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]

```

Rubi steps

$$\begin{aligned}
\int (d + ex)^3 (a + b \tan^{-1}(cx))^3 dx &= \frac{(d + ex)^4 (a + b \tan^{-1}(cx))^3}{4e} - \frac{(3bc) \int \left(\frac{e^2 (6c^2 d^2 - e^2) (a + b \tan^{-1}(cx))^2}{c^4} + \frac{4de}{c^2} \right)}{4e} \\
&= \frac{(d + ex)^4 (a + b \tan^{-1}(cx))^3}{4e} - \frac{(3b) \int \frac{(c^4 d^4 - 6c^2 d^2 e^2 + e^4 + 4c^2 d(c d - e)e(c d + e)x)}{1 + c^2 x^2}}{4c^3 e} \\
&= -\frac{3be(6c^2 d^2 - e^2) x (a + b \tan^{-1}(cx))^2}{4c^3} - \frac{3bde^2 x^2 (a + b \tan^{-1}(cx))^2}{2c} - \\
&= -\frac{3ibe(6c^2 d^2 - e^2) (a + b \tan^{-1}(cx))^2}{4c^4} - \frac{3be(6c^2 d^2 - e^2) x (a + b \tan^{-1}(cx))^2}{4c^3} \\
&= \frac{3ab^2 de^2 x}{c^2} + \frac{b^2 e^3 x^2 (a + b \tan^{-1}(cx))}{4c^2} - \frac{3bde^2 (a + b \tan^{-1}(cx))^2}{2c^3} + \frac{ibe^5 x^3}{c^5} \\
&= \frac{3ab^2 de^2 x}{c^2} - \frac{b^3 e^3 x}{4c^3} + \frac{3b^3 de^2 x \tan^{-1}(cx)}{c^2} + \frac{b^2 e^3 x^2 (a + b \tan^{-1}(cx))}{4c^2} - \\
&= \frac{3ab^2 de^2 x}{c^2} - \frac{b^3 e^3 x}{4c^3} + \frac{b^3 e^3 \tan^{-1}(cx)}{4c^4} + \frac{3b^3 de^2 x \tan^{-1}(cx)}{c^2} + \frac{b^2 e^3 x^2 (a + b \tan^{-1}(cx))}{4c^2} \\
&= \frac{3ab^2 de^2 x}{c^2} - \frac{b^3 e^3 x}{4c^3} + \frac{b^3 e^3 \tan^{-1}(cx)}{4c^4} + \frac{3b^3 de^2 x \tan^{-1}(cx)}{c^2} + \frac{b^2 e^3 x^2 (a + b \tan^{-1}(cx))}{4c^2}
\end{aligned}$$

Mathematica [A]

time = 1.21, size = 855, normalized size = 1.31

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)^3*(a + b*ArcTan[c*x])^3, x]`

[Out]
$$\begin{aligned} & (a^2*c*(4*a*c^3*d^3 + 3*b*e*(-6*c^2*d^2 + e^2))*x + 6*a^2*c^3*d*e*(a*c*d - b*e)*x^2 + a^2*c^3*e^2*(4*a*c*d - b*e)*x^3 + a^3*c^4*e^3*x^4 + 3*a^2*b*(6*c^2*d^2*e - e^3)*ArcTan[c*x] + 3*a^2*b*c^4*x*(4*d^3 + 6*d^2*e*x + 4*d*e^2*x^2 + e^3*x^3)*ArcTan[c*x] + a*b^2*e^3*(1 + c^2*x^2 + (6*c*x - 2*c^3*x^3)*ArcTan[c*x] + 3*(-1 + c^4*x^4)*ArcTan[c*x]^2 - 4*\text{Log}[1 + c^2*x^2]) - 6*a^2*b*c*d*(c^2*d^2 - e^2)*\text{Log}[1 + c^2*x^2] + 18*a*b^2*c^2*d^2*e*(-2*c*x*ArcTan[c*x] + (1 + c^2*x^2)*ArcTan[c*x]^2 + \text{Log}[1 + c^2*x^2]) + 12*a*b^2*c^3*d^3*(ArcTan[c*x]*((-I + c*x)*ArcTan[c*x] + 2*\text{Log}[1 + E^((2*I)*ArcTan[c*x])]) - I*\text{PolyLog}[2, -E^((2*I)*ArcTan[c*x])] + 12*a*b^2*c*d^2*e^2*(c*x + (I + c^3*x^3)*ArcTan[c*x]^2 - ArcTan[c*x]*(1 + c^2*x^2 + 2*\text{Log}[1 + E^((2*I)*ArcTan[c*x])]) + I*\text{PolyLog}[2, -E^((2*I)*ArcTan[c*x])] + 6*b^3*c^2*d^2*e*(ArcTan[c*x]*((3*I - 3*c*x)*ArcTan[c*x] + (1 + c^2*x^2)*ArcTan[c*x]^2 - 6*\text{Log}[1 + E^((2*I)*ArcTan[c*x])] + (3*I)*\text{PolyLog}[2, -E^((2*I)*ArcTan[c*x])] + b^3*e^3*(-(c*x) - (4*I - 3*c*x + c^3*x^3)*ArcTan[c*x]^2 + (-1 + c^4*x^4)*ArcTan[c*x]^3 + ArcTan[c*x]*(1 + c^2*x^2 + 8*\text{Log}[1 + E^((2*I)*ArcTan[c*x])]) - (4*I)*\text{PolyLog}[2, -E^((2*I)*ArcTan[c*x])] + 2*b^3*c*d^2*e^2*(6*c*x*ArcTan[c*x] - 3*ArcTan[c*x]^2 - 3*c^2*x^2*ArcTan[c*x]^2 + (2*I)*ArcTan[c*x]^3 + 2*c^3*x^3*ArcTan[c*x]^3 - 6*ArcTan[c*x]^2*\text{Log}[1 + E^((2*I)*ArcTan[c*x])] - 3*\text{Log}[1 + c^2*x^2] + (6*I)*ArcTan[c*x]*\text{PolyLog}[2, -E^((2*I)*ArcTan[c*x])] - 3*\text{PolyLog}[3, -E^((2*I)*ArcTan[c*x])] + 2*b^3*c^3*d^3*(2*ArcTan[c*x]^2*((-I + c*x)*ArcTan[c*x] + 3*\text{Log}[1 + E^((2*I)*ArcTan[c*x])]) - (6*I)*ArcTan[c*x]*\text{PolyLog}[2, -E^((2*I)*ArcTan[c*x])] + 3*\text{PolyLog}[3, -E^((2*I)*ArcTan[c*x])))/(4*c^4) \end{aligned}$$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 22.02, size = 3456, normalized size = 5.30

method	result	size
derivativedivides	Expression too large to display	3456
default	Expression too large to display	3456

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)^3*(a+b*arctan(c*x))^3,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & 1/c*(-3/4*I*b^3/c^2*e^2*Pi*d*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2+3/2*I*b^3/c^2*e^2*Pi*d*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1))*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2+3/4*I*b^3/c^2*e^2*Pi*d*csgn(I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})^2*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))*arctan(c*x)^2-3/2*I*b^3/c^2*e^2*Pi*d*csgn(I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})^2*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))*arctan(c*x)^2-3/2*I*b^3/c^2*e^2*Pi*d*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))*arctan(c*x)^2+1/4*a*b^2/c*e^3*x^2-3/4*I*b^3/c^2*e^2*Pi*d*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2-3/4*I*b^3/c^2*e^2*Pi*d*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))+csgn(I*((1+I*c*x)^2/(c^2*x^2+1))^2)*arctan(c*x)^2+3/4*I*b^3Pi \end{aligned}$$

$$\begin{aligned}
& *d^3*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/(\\
& (1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2-3/4*a*b^2/c^3*e^3*arctan(c*x) \\
& ^{-2-a*b^2/c^3*e^3*ln(c^2*x^2+1)-3*a*b^2*arctan(c*x)*ln(c^2*x^2+1)*d^3+3/2*I* \\
& a*b^2*d^3*dilog(-1/2*I*(c*x+I))+3/4*I*a*b^2*d^3*ln(c*x-I)^2-3/2*I*a*b^2*d^3 \\
& *dilog(1/2*I*(c*x-I))-3/4*I*a*b^2*d^3*ln(c*x+I)^2-3/4*a^2*b/c^3*arctan(c*x) \\
& *e^3-1/4*b^3/c^2*e^3*x+b^3*arctan(c*x)^3*d^3*c*x-3/2*b^3/c^2*e^2*d*arctan(c \\
& *x)^2+3/2*b^3/c*e*arctan(c*x)^3*d^2+3*b^3/c^2*e^2*d*ln((1+I*c*x)^2/(c^2*x^2 \\
& +1)+1)-3/2*b^3/c^2*e^2*d*polylog(3,-(1+I*c*x)^2/(c^2*x^2+1))+2*b^3/c^3*e^3* \\
& arctan(c*x)*ln(1+I*(1+I*c*x)/(c^2*x^2+1)^(1/2))+2*b^3/c^3*e^3*arctan(c*x)*l \\
& n(1-I*(1+I*c*x)/(c^2*x^2+1)^(1/2))-3*I*b^3*d^3*arctan(c*x)*polylog(2,-(1+I* \\
& c*x)^2/(c^2*x^2+1))-I*b^3/c^3*e^3*arctan(c*x)^2-2*I*b^3/c^3*e^3*dilog(1-I*(\\
& 1+I*c*x)/(c^2*x^2+1)^(1/2))-2*I*b^3/c^3*e^3*dilog(1+I*(1+I*c*x)/(c^2*x^2+1) \\
& ^{(1/2)})-3/2*I*a*b^2/c^2*e^2*d*ln(c^2*x^2+1)*ln(c*x+I)-3/2*I*a*b^2/c^2*e^2*d \\
& *ln(-1/2*I*(c*x+I))*ln(c*x-I)+3/2*I*a*b^2/c^2*e^2*d*ln(c^2*x^2+1)*ln(c*x-I) \\
& +3/2*I*a*b^2/c^2*e^2*d*ln(c*x+I)*ln(1/2*I*(c*x-I))-3/4*I*b^3/c^2*e^2*Pi*d*c \\
& sgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^3*arctan(c*x)^2+3/4*I*b^3/c^2*e^2*Pi*d \\
& *csgn(I*((1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^3*arctan(c*x) \\
&)^2+3/4*I*b^3/c^2*e^2*Pi*d*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))^3*arctan(c*x)^2- \\
& 3/4*I*b^3*Pi*d^3*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*(1+I*c*x)^2/(\\
& c^2*x^2+1))*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*a \\
& rctan(c*x)^2-9*a*b^2*e*arctan(c*x)*d^2*x-3*a*b^2*e^2*arctan(c*x)*d*x^2+9/2* \\
& a*b^2*c*e*arctan(c*x)^2*d^2*x^2+3*a*b^2*c*e^2*arctan(c*x)^2*d*x^3+9/2*a^2*b \\
& *c*e*arctan(c*x)*d^2*x^2+1/4*(c*e*x+c*d)^4*a^3/c^3/e-3/2*a^2*b*ln(c^2*x^2+1) \\
&)*d^3+3*b^3*d^3*ln((1+I*c*x)/(c^2*x^2+1)^(1/2))*arctan(c*x)^2-3/2*b^3*arcta \\
& n(c*x)^2*ln(c^2*x^2+1)*d^3-1/4*I*b^3/c^3*e^3-I*b^3*d^3*arctan(c*x)^3-1/4*b^ \\
& 3/c^3*e^3*arctan(c*x)^3+1/4*b^3/c^3*arctan(c*x)*e^3+3*b^3*ln(2)*d^3*arctan(\\
& c*x)^2+3/4*I*b^3*Pi*d^3*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1))^2*csgn(I*((1+I* \\
& c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2+3/4*I*b^3*Pi*d^3*csgn(I*(1+I*c*x)^2/(\\
& c^2*x^2+1))*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2) \\
& ^2*arctan(c*x)^2+3/2*I*b^3*Pi*d^3*csgn(I*(1+I*c*x)/(c^2*x^2+1)^(1/2))*csgn(I \\
& *(1+I*c*x)^2/(c^2*x^2+1))^2*arctan(c*x)^2-3/2*I*b^3*Pi*d^3*csgn(I*((1+I*c*x) \\
&)^2/(c^2*x^2+1)+1)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2-9 \\
& /2*b^3*e*arctan(c*x)^2*d^2*x-3/2*b^3*e^2*arctan(c*x)^2*d*x^2+3/2*b^3*c*e*ar \\
& ctan(c*x)^3*d^2*x^2+b^3*c*e^2*arctan(c*x)^3*d*x^3+3/2*I*a*b^2/c^2*e^2*d*arctan(\\
& c*x)*x+3*a*b^2/c^2*e^2*arctan(c*x)*ln(c^2*x^2+1)*d+3/2*I*a*b^2/c^2*e^2*d*d \\
& ilog(1/2*I*(c*x-I))-3/2*I*a*b^2/c^2*e^2*d*dilog(-1/2*I*(c*x+I))-3/4*I*a*b^2/c^2* \\
& e^2*d*ln(c*x-I)^2+3/4*I*a*b^2/c^2*e^2*d*ln(c*x+I)^2-9/2*a^2*b*d^2*e*x-3/2*a \\
& ^2*b*d*e^2*x^2+3/4*a^2*b*c*e^3*arctan(c*x)*x^4+3/4*a*b^2*c*e^3*arctan(c*x) \\
& ^2*x^4-1/2*a*b^2*x^3*arctan(c*x)*x^3+3/2*a*b^2/c^2*e^3*arctan(c*x)*x+3*a*b^2 \\
& /c*d*e^2*x-1/4*b^3*c^3*arctan(c*x)^2*x^3+3/4*b^3/c^2*e^3*arctan(c*x)^2*x+1/ \\
& 4*b^3/c*e^3*arctan(c*x)*x^2+9*I*b^3/c*e*d^2*dilog(1-I*(1+I*c*x)/(c^2*x^2+1) \\
& ^{(1/2)})+9*I*b^3/c*e*d^2*dilog(1+I*(1+I*c*x)/(c^2*x^2+1)^(1/2))-3*I*b^3/c^2* \\
& e^2*d*arctan(c*x)+9/2*I*b^3/c*e*d^2*arctan(c*x)^2-1/4*a^2*b*e^3*x^3+3/4*a^2 \\
& *b/c^2*e^3*x+3*a^2*b*arctan(c*x)*d^3*c*x+3/2*a^2*b/c^2*e^2*ln(c^2*x^2+1)*d+ \\
& 9/2*a^2*b/c*e*arctan(c*x)*d^2+9/2*a*b^2/c*e*ln(c^2*x^2+1)*d^2+9/2*a*b^2/c*e
\end{aligned}$$

$$\begin{aligned}
& * \arctan(c*x)^2 * d^2 - 3*a*b^2/c^2 * e^2 * d * \arctan(c*x) + 3/2 * I * a * b^2 * d^3 * \ln(c^2 * x^2 + 1) * \ln(c*x + I) + 3/2 * I * a * b^2 * d^3 * \ln(-1/2 * I * (c*x + I)) * \ln(c*x - I) - 3/2 * I * a * b^2 * d^3 * \ln(c^2 * x^2 + 1) * \ln(c*x - I) - 3/2 * I * a * b^2 * d^3 * \ln(c*x + I) * \ln(1/2 * I * (c*x - I)) + 1/4 * b^3 * c * e^3 * \arctan(c*x)^3 * x^4 + 3 * I * b^3 / c^2 * e^2 * d * \arctan(c*x) * \text{polylog}(2, -(1 + I * c * x)^2 / (c^2 * x^2 + 1)) + 3/2 * b^3 / c^2 * e^2 * \arctan(c*x)^2 * \ln(c^2 * x^2 + 1) * d + I * b^3 / c^2 * e^2 * d * \arctan(c*x)^3 - 3 * b^3 / c^2 * e^2 * \ln(2) * d * \arctan(c*x)^2 - 9 * b^3 / c * e * d^2 * \arctan(c*x) * \ln(1 + I * (1 + I * c * x)) / (c^2 * x^2 + 1)^{(1/2)} - 9 * b^3 / c * e * d^2 * \arctan(c*x) * \ln(1 - I * (1 + I * c * x)) / (c^2 * x^2 + 1)^{(1/2)} - 3 * b^3 / c^2 * e^2 * d * \ln((1 + I * c * x) / (c^2 * x^2 + 1)^{(1/2)}) * \arctan(c*x)^2 - 3/4 * I * b^3 * \text{Pi} * d^3 * \text{csgn}(I * (1 + I * c * x)^2 / (c^2 * x^2 + 1)) ^ 3 * \arctan(c*x)^2 - 3/4 * I * b^3 * \text{Pi} * d^3 * \text{csgn}(I * (1 + I * c * x)^2 / (c^2 * x^2 + 1) + 1) ^ 2) ^ 3 * \arctan(c*x)^2 + 3/4 * I * b^3 * \text{Pi} * d^3 * \text{csgn}(I * ((1 + I * c * x)^2 / (c^2 * x^2 + 1) + 1) ^ 2) ^ 3 * \arctan(c*x)^2 + 3 * a * b^2 * \arctan(c*x)^2 * d^3 * c...
\end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^3*(a+b*arctan(c*x))^3,x, algorithm="maxima")
[Out] 7/32*b^3*d^3*arctan(c*x)^4/c + 336*b^3*c^2*d^2*e*integrate(1/128*x^3*arctan(c*x)^3/(c^2*x^2 + 1), x) + 36*b^3*c^2*d^2*e*integrate(1/128*x^3*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 1152*a*b^2*c^2*d^2*e*integrate(1/128*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + 112*b^3*c^2*d^3*integrate(1/128*x^2*arctan(c*x)^3/(c^2*x^2 + 1), x) + 72*b^3*c^2*d^2*e*integrate(1/128*x^3*arctan(c*x)*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + 12*b^3*c^2*d^3*integrate(1/128*x^2*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 384*a*b^2*c^2*d^3*integrate(1/128*x^2*arctan(c*x)*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + 48*b^3*c^2*d^3*integrate(1/128*x^2*arctan(c*x)*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + a*b^2*d^3*a*arctan(c*x)^3/c + 1/4*a^3*x^4*e^3 + a^3*d*x^3*e^2 + 3/2*a^3*d^2*x^2*e + 336*b^3*c^2*d*e^2*integrate(1/128*x^4*arctan(c*x)^3/(c^2*x^2 + 1), x) + 36*b^3*c^2*d^2*e^2*integrate(1/128*x^4*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 1152*a*b^2*c^2*d^2*e^2*integrate(1/128*x^4*arctan(c*x)^2/(c^2*x^2 + 1), x) + 48*b^3*c^2*d^2*e^2*integrate(1/128*x^4*arctan(c*x)*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) - 72*b^3*c*d^2*e*integrate(1/128*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 18*b^3*c*d^2*e*integrate(1/128*x^2*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) - 48*b^3*c*d^3*integrate(1/128*x*arctan(c*x)^2/(c^2*x^2 + 1), x) + 12*b^3*c*d^3*integrate(1/128*x*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + a^3*d^3*x + 9/2*(x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*a^2*b*d^2*e + 112*b^3*c^2*e^3*integrate(1/128*x^5*arctan(c*x)^3/(c^2*x^2 + 1), x) + 12*b^3*c^2*e^3*integrate(1/128*x^5*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 384*a*b^2*c^2*e^3*integrate(1/128*x^5*arctan(c*x)^2/(c^2*x^2 + 1), x) + 12*b^3*c^2*e^3*integrate(1/128*x^5*arctan(c*x)*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) - 48*b^3*c*d^2*e^2*integrate(1/128*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x)
```

```

) + 12*b^3*c*d*e^2*integrate(1/128*x^3*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x)
+ 336*b^3*d^2*e*integrate(1/128*x*arctan(c*x)^3/(c^2*x^2 + 1), x) + 36*b^3
*d^2*e*integrate(1/128*x*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) +
1152*a*b^2*d^2*e*integrate(1/128*x*arctan(c*x)^2/(c^2*x^2 + 1), x) + 12*b^
3*d^3*integrate(1/128*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 3/
2*(2*c*x*arctan(c*x) - log(c^2*x^2 + 1))*a^2*b*d^3/c + 3/2*(2*x^3*arctan(c*
x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c^4))*a^2*b*d*e^2 - 12*b^3*c*e^3*integra
te(1/128*x^4*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^3*c*e^3*integrate(1/128*
x^4*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 336*b^3*d*e^2*integrate(1/128*x^
2*arctan(c*x)^3/(c^2*x^2 + 1), x) + 36*b^3*d*e^2*integrate(1/128*x^2*arctan(
c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 1152*a*b^2*d^2*e^2*integrate(1/1
28*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x) + 1/4*(3*x^4*arctan(c*x) - c*((c^2*x
^3 - 3*x)/c^4 + 3*arctan(c*x)/c^5))*a^2*b*e^3 + 112*b^3*e^3*integrate(1/128*
x^3*arctan(c*x)^3/(c^2*x^2 + 1), x) + 12*b^3*e^3*integrate(1/128*x^3*arcta
n(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 384*a*b^2*e^3*integrate(1/128*
x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + 1/32*(b^3*x^4*e^3 + 4*b^3*d*x^3*e^2
+ 6*b^3*d^2*x^2*e + 4*b^3*d^3*x)*arctan(c*x)^3 - 3/128*(b^3*x^4*e^3 + 4*b^3
*d*x^3*e^2 + 6*b^3*d^2*x^2*e + 4*b^3*d^3*x)*arctan(c*x)*log(c^2*x^2 + 1)^2

```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```

[In] integrate((e*x+d)^3*(a+b*arctan(c*x))^3,x, algorithm="fricas")
[Out] integral(a^3*x^3*e^3 + 3*a^3*d*x^2*e^2 + 3*a^3*d^2*x*e + a^3*d^3 + (b^3*x^3
*e^3 + 3*b^3*d*x^2*e^2 + 3*b^3*d^2*x*e + b^3*d^3)*arctan(c*x)^3 + 3*(a*b^2*
x^3*e^3 + 3*a*b^2*d*x^2*e^2 + 3*a*b^2*d^2*x*e + a*b^2*d^3)*arctan(c*x)^2 +
3*(a^2*b*x^3*e^3 + 3*a^2*b*d*x^2*e^2 + 3*a^2*b*d^2*x*e + a^2*b*d^3)*arctan(
c*x), x)

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx))^3 (d + ex)^3 \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

```

[In] integrate((e*x+d)**3*(a+b*atan(c*x))**3,x)
[Out] Integral((a + b*atan(c*x))**3*(d + e*x)**3, x)

```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^3*(a+b*arctan(c*x))^3,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a + b \operatorname{atan}(c x))^3 (d + e x)^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^3*(d + e*x)^3,x)`

[Out] `int((a + b*atan(c*x))^3*(d + e*x)^3, x)`

$$\text{3.16} \quad \int (d + ex)^2 (a + b \text{ArcTan}(cx))^3 dx$$

Optimal. Leaf size=411

$$\frac{ab^2 e^2 x}{c^2} + \frac{b^3 e^2 x \text{ArcTan}(cx)}{c^2} - \frac{3ibde(a + b \text{ArcTan}(cx))^2}{c^2} - \frac{be^2(a + b \text{ArcTan}(cx))^2}{2c^3} - \frac{3bdex(a + b \text{ArcTan}(cx))^2}{c}$$

[Out] $a*b^2e^2x/c^2+b^3e^2x\text{ArcTan}(cx)/c^2-3*I*b*d*e*(a+b*\text{arctan}(c*x))^2/c^2-1/2*b*e^2*(a+b*\text{arctan}(c*x))^2/c^3-3*b*d*e*x*(a+b*\text{arctan}(c*x))^2/c-1/2*b*e^2*x^2*(a+b*\text{arctan}(c*x))^2/c+1/3*I*(3*c^2*d^2-e^2)*(a+b*\text{arctan}(c*x))^3/c^3-1/3*d*(d^2-3*e^2/c^2)*(a+b*\text{arctan}(c*x))^3/e+1/3*(e*x+d)^3*(a+b*\text{arctan}(c*x))^3/e-6*b^2*d*e*(a+b*\text{arctan}(c*x))*\ln(2/(1+I*c*x))/c^2+b*(3*c^2*d^2-e^2)*(a+b*\text{arctan}(c*x))^2*\ln(2/(1+I*c*x))/c^3-1/2*b^3*e^2*\ln(c^2*x^2+1)/c^3-3*I*b^3*d*e*\text{polylog}(2,1-2/(1+I*c*x))/c^2+I*b^2*(3*c^2*d^2-e^2)*(a+b*\text{arctan}(c*x))*\text{polylog}(2,1-2/(1+I*c*x))/c^3+1/2*b^3*(3*c^2*d^2-e^2)*\text{polylog}(3,1-2/(1+I*c*x))/c^3$

Rubi [A]

time = 0.52, antiderivative size = 411, normalized size of antiderivative = 1.00, number of steps used = 20, number of rules used = 13, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.722, Rules used = {4974, 4930, 5040, 4964, 2449, 2352, 4946, 5036, 266, 5004, 5104, 5114, 6745}

$$\frac{ab^2 d e \log \left(\frac{1}{c^2} \frac{1}{x}\right)}{c^2} (a + b \text{ArcTan}(cx)) + \frac{ab^2 (b^2 e^2 d^2 - e^4) \ln(1 - \frac{1}{c^2 x})}{c^2} (a + b \text{ArcTan}(cx)) - \frac{be^2 (a + b \text{ArcTan}(cx))^2}{2c^3} + \frac{d (d^2 - \frac{3e^2}{c^2}) (a + b \text{ArcTan}(cx))^3}{3c} - \frac{3 ib d (a + b \text{ArcTan}(cx))^2}{c^2} - \frac{i (3 b^2 d^2 - e^2) (a + b \text{ArcTan}(cx))^2}{2c^2} + \frac{i (3 b^2 d^2 - e^2) \log \left(\frac{1}{c^2} \frac{1}{x}\right)}{c^2} (a + b \text{ArcTan}(cx))^2 - \frac{3 ib d (a + b \text{ArcTan}(cx))^2}{c} - \frac{(d + e x)^2 (a + b \text{ArcTan}(cx))^3}{2c} + \frac{\ln(e x^2) (a + b \text{ArcTan}(cx))^2}{2c} - \frac{ab^2 e^2 x}{c^2} + \frac{b^3 e^2 x \text{ArcTan}(cx)}{c^2} - \frac{3ib^2 d k_3 (1 - \frac{1}{c^2 x})}{c^2} + \frac{b^2 (b^2 e^2 d^2 - e^4) \ln(1 - \frac{1}{c^2 x})}{2c^2} - \frac{b^2 e^2 \log (e^2 x^2 + 1)}{2c^2}$$

Antiderivative was successfully verified.

$$[\text{In}] \quad \text{Int}[(d + e*x)^2*(a + b*\text{ArcTan}[c*x])^3, x]$$

[Out] $(a*b^2e^2x/c^2 + (b^3e^2x\text{ArcTan}[c*x])/c^2 - ((3*I)*b*d*e*(a + b*\text{ArcTan}[c*x]))^2/c^2 - (b*e^2*(a + b*\text{ArcTan}[c*x]))^2/(2*c^3) - (3*b*d*e*x*(a + b*\text{ArcTan}[c*x]))^2/c - (b*e^2*x^2*(a + b*\text{ArcTan}[c*x]))^2/(2*c) + ((I/3)*(3*c^2*d^2 - e^2)*(a + b*\text{ArcTan}[c*x]))^3/c^3 - (d*(d^2 - (3*e^2)/c^2)*(a + b*\text{ArcTan}[c*x]))^3/(3*e) + ((d + e*x)^3*(a + b*\text{ArcTan}[c*x]))^3/(3*e) - (6*b^2*d*e*(a + b*\text{ArcTan}[c*x])*Log[2/(1 + I*c*x)])/c^2 + (b*(3*c^2*d^2 - e^2)*(a + b*\text{ArcTan}[c*x]))^2*Log[2/(1 + I*c*x)]/c^3 - (b^3*e^2*Log[1 + c^2*x^2])/(2*c^3) - ((3*I)*b^3*d*e*\text{PolyLog}[2, 1 - 2/(1 + I*c*x)])/c^2 + (I*b^2*(3*c^2*d^2 - e^2)*(a + b*\text{ArcTan}[c*x])*PolyLog[2, 1 - 2/(1 + I*c*x)])/c^3 + (b^3*(3*c^2*d^2 - e^2)*PolyLog[3, 1 - 2/(1 + I*c*x)])/(2*c^3)$

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> Simp[Log[RemoveContent[t[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 2352

```
Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_)/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] :> Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 4930

```
Int[((a_*) + ArcTan[(c_)*(x_)^(n_)]*(b_.))^p, x_Symbol] :> Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```

Rule 4946

```
Int[((a_*) + ArcTan[(c_)*(x_)^(n_)]*(b_.))^p*(x_)^m, x_Symbol] :> Simp[x^(m + 1)*((a + b*ArcTan[c*x^n])^p/(m + 1)), x] - Dist[b*c*n*(p/(m + 1)), Int[x^(m + n)*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 0] && (EqQ[p, 1] || (EqQ[n, 1] && IntegerQ[m])) && NeQ[m, -1]
```

Rule 4964

```
Int[((a_*) + ArcTan[(c_)*(x_)]*(b_.))^p/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e), Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_*) + ArcTan[(c_)*(x_)]*(b_.))^p*((d_) + (e_)*(x_)^q), x_Symbol] :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Dist[b*c*(p/(e*(q + 1))), Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2)], x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_*) + ArcTan[(c_)*(x_)]*(b_.))^p/((d_) + (e_)*(x_)^2), x_Symbol] :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5036

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*((f_)*(x_))^(m_))/((d_) + (e_)*(x_)^2), x_Symbol] :> Dist[f^2/e, Int[(f*x)^(m - 2)*(a + b*ArcTan[c*x])^p, x], x] - Dist[d*(f^2/e), Int[(f*x)^(m - 2)*((a + b*ArcTan[c*x])^p/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e, f}, x] && GtQ[p, 0] && GtQ[m, 1]
```

Rule 5040

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*((x_)))/((d_) + (e_)*(x_)^2), x_Symbol] :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Dist[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.)*((f_) + (g_)*(x_))^(m_.))/((d_) + (e_)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rule 5114

```
Int[(Log[u_]*((a_.) + ArcTan[(c_)*(x_)]*(b_.))^(p_.))/((d_) + (e_)*(x_)^2), x_Symbol] :> Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Dist[b*p*(I/2), Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]
```

Rule 6745

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simp[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Rubi steps

$$\begin{aligned}
\int (d+ex)^2 (a+b\tan^{-1}(cx))^3 dx &= \frac{(d+ex)^3 (a+b\tan^{-1}(cx))^3}{3e} - \frac{(bc) \int \left(\frac{3de^2(a+b\tan^{-1}(cx))^2}{c^2} + \frac{e^3x(a+b\tan^{-1}(cx))^2}{c^2} \right) dx}{3e} \\
&= \frac{(d+ex)^3 (a+b\tan^{-1}(cx))^3}{3e} - \frac{b \int \frac{(c^2d^3-3de^2+e(3c^2d^2-e^2)x)(a+b\tan^{-1}(cx))^2}{1+c^2x^2} dx}{ce} \\
&= -\frac{3bdex(a+b\tan^{-1}(cx))^2}{c} - \frac{be^2x^2(a+b\tan^{-1}(cx))^2}{2c} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx))^3}{3e} \\
&= -\frac{3ibde(a+b\tan^{-1}(cx))^2}{c^2} - \frac{3bdex(a+b\tan^{-1}(cx))^2}{c} - \frac{be^2x^2(a+b\tan^{-1}(cx))^2}{2c} \\
&= \frac{ab^2e^2x}{c^2} - \frac{3ibde(a+b\tan^{-1}(cx))^2}{c^2} - \frac{be^2(a+b\tan^{-1}(cx))^2}{2c^3} - \frac{3bdex(a+b\tan^{-1}(cx))^3}{c^3} \\
&= \frac{ab^2e^2x}{c^2} + \frac{b^3e^2x\tan^{-1}(cx)}{c^2} - \frac{3ibde(a+b\tan^{-1}(cx))^2}{c^2} - \frac{be^2(a+b\tan^{-1}(cx))^2}{2c^3} \\
&= \frac{ab^2e^2x}{c^2} + \frac{b^3e^2x\tan^{-1}(cx)}{c^2} - \frac{3ibde(a+b\tan^{-1}(cx))^2}{c^2} - \frac{be^2(a+b\tan^{-1}(cx))^2}{2c^3} \\
&= \frac{ab^2e^2x}{c^2} + \frac{b^3e^2x\tan^{-1}(cx)}{c^2} - \frac{3ibde(a+b\tan^{-1}(cx))^2}{c^2} - \frac{be^2(a+b\tan^{-1}(cx))^2}{2c^3}
\end{aligned}$$

Mathematica [A]

time = 0.72, size = 621, normalized size = 1.51

Antiderivative was successfully verified.

[In] Integrate[(d + e*x)^2*(a + b*ArcTan[c*x])^3, x]

[Out]

$$\begin{aligned}
&(6*a^2*c^2*d*(a*c*d - 3*b*e)*x + 3*a^2*c^2*e*(2*a*c*d - b*e)*x^2 + 2*a^3*c^3*e^2*x^3 + 18*a^2*b*c*d*e*ArcTan[c*x] + 6*a^2*b*c^3*x*(3*d^2 + 3*d*e*x + e^2*x^2)*ArcTan[c*x] - 3*a^2*b*(3*c^2*d^2 - e^2)*Log[1 + c^2*x^2] + 18*a*b^2*c*d*e*(-2*c*x*ArcTan[c*x] + (1 + c^2*x^2)*ArcTan[c*x]^2 + Log[1 + c^2*x^2]) + 18*a*b^2*c^2*d^2*(ArcTan[c*x]*((-I + c*x)*ArcTan[c*x] + 2*Log[1 + E^((2*I)*ArcTan[c*x])]) - I*PolyLog[2, -E^((2*I)*ArcTan[c*x])]) + 6*a*b^2*e^2*(c*x + (I + c^3*x^3)*ArcTan[c*x]^2 - ArcTan[c*x]*(1 + c^2*x^2) + 2*Log[1 + E^((2*I)*ArcTan[c*x])]) + I*PolyLog[2, -E^((2*I)*ArcTan[c*x])] + 6*b^3*c*d*e*(ArcTan[c*x]*((3*I - 3*c*x)*ArcTan[c*x] + (1 + c^2*x^2)*ArcTan[c*x]^2 - 6*Log[1 + E^((2*I)*ArcTan[c*x])]) + (3*I)*PolyLog[2, -E^((2*I)*ArcTan[c*x])]) + b^3*e^2*(6*c*x*ArcTan[c*x] - 3*ArcTan[c*x]^2 - 3*c^2*x^2*ArcTan[c*x]^2 + 6*c*x*ArcTan[c*x]^3 - 3*c*x*Log[1 + E^((2*I)*ArcTan[c*x])]) + 3*c*x^2*ArcTan[c*x]^3 - 3*c*x^2*Log[1 + E^((2*I)*ArcTan[c*x])])
\end{aligned}$$

$$(2*I)*ArcTan[c*x]^3 + 2*c^3*x^3*ArcTan[c*x]^3 - 6*ArcTan[c*x]^2*Log[1 + E^(-(2*I)*ArcTan[c*x])] - 3*Log[1 + c^2*x^2] + (6*I)*ArcTan[c*x]*PolyLog[2, -E^(-(2*I)*ArcTan[c*x])] - 3*PolyLog[3, -E^(-(2*I)*ArcTan[c*x])] + 3*b^3*c^2*d^2*(2*ArcTan[c*x]^2*((-I + c*x)*ArcTan[c*x] + 3*Log[1 + E^((2*I)*ArcTan[c*x])]) - (6*I)*ArcTan[c*x]*PolyLog[2, -E^((2*I)*ArcTan[c*x])] + 3*PolyLog[3, -E^((2*I)*ArcTan[c*x])))/(6*c^3)$$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 7.53, size = 2921, normalized size = 7.11

method	result	size
derivativedivides	Expression too large to display	2921
default	Expression too large to display	2921

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((e*x+d)^2*(a+b*arctan(c*x))^3,x,method=_RETURNVERBOSE)
[Out] 1/c*(I*b^3/c^2*e^2*arctan(c*x)*polylog(2,-(1+I*c*x)^2/(c^2*x^2+1))+1/4*I*b^3/c^2*e^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))^3*arctan(c*x)^2+1/4*I*b^3/c^2*e^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^3*arctan(c*x)^2-3*b^3*e*arctan(c*x)^2*d*x-a*b^2*e^2*arctan(c*x)*x^2+a*b^2*c*e^2*a*rctan(c*x)^2*x^3-3*a^2*b*d*e*x+a^2*b*c*e^2*arctan(c*x)*x^3-3/2*a^2*b*ln(c^2*x^2+1)*d^2-1/2*b^3/c^2*e^2*arctan(c*x)^2+b^3/c^2*e^2*ln((1+I*c*x)^2/(c^2*x^2+1)+1)-1/2*b^3/c^2*e^2*polylog(3,-(1+I*c*x)^2/(c^2*x^2+1))+3*b^3*d^2*ln(2)*arctan(c*x)^2+3*b^3*ln((1+I*c*x)/(c^2*x^2+1)^(1/2))*d^2*arctan(c*x)^2-3/2*b^3*arctan(c*x)^2*ln(c^2*x^2+1)*d^2-I*b^3*d^2*arctan(c*x)^3-1/2*I*b^3/c^2*e^2*Pi*csgn(I*(1+I*c*x)/(c^2*x^2+1)^(1/2))*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))^2*arctan(c*x)^2+1/4*I*b^3/c^2*e^2*Pi*csgn(I*(1+I*c*x)/(c^2*x^2+1)^(1/2))^2*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))*arctan(c*x)^2-1/4*I*b^3/c^2*e^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)+1)^2*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2-1/4*I*b^3/c^2*e^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2-1/4*I*b^3/c^2*e^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2-3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2-1/4*I*b^3/c^2*e^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2+3*a*b^2*c*e*arctan(c*x)^2*d*x^2-6*a*b^2*e*arctan(c*x)*d*x+3*a^2*b*c*e*arctan(c*x)*d*x^2+b^3*c*e*arctan(c*x)^3*d*x^2+1/2*I*a*b^2/c^2*e^2*ln(1/2*I*(c*x-I))*ln(c*x+I)-1/2*I*a*b^2/c^2*e^2*ln(-1/2*I*(c*x+I))*ln(c*x-I)+1/2*I*a*b^2/c^2*e^2*ln(c*x-I)*ln(c^2*x^2+1)-1/2*I*a*b^2/c^2*e^2*ln(c*x+I)*ln(c^2*x^2+1)+1/4*I*b^3/c^2*e^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*arctan(c*x)^2+1/3*(c*e*x+c*d)^3*a^3/c^2/e+3/2*b^3*polylog(3,-(1+I*c*x)^2/(c^2*x^2+1))*d^2+3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))*d^2+3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)))
```

$$\begin{aligned}
& x^2/(c^2*x^2+1)+1)^2)*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2+3/4*I*b^3*d^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))* \\
& csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2-3/4*I*b^3*d^2*Pi*csgn(I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})^2*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)+1)^2*arctan(c*x)^2-3/2*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2+3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1))^{2*arctan(c*x)^2+3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2-1/4*I*b^3/c^2*e^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2*arctan(c*x)^2-6*b^3/c*e*d*arctan(c*x)*ln(1+I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})-6*b^3/c*e*d*arctan(c*x)*ln(1-I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})+6*I*b^3/c*e*d*dilog(1-I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})+6*I*b^3/c*e*d*dilog(1+I*(1+I*c*x)/(c^2*x^2+1)^{(1/2)})+3*I*b^3/c*e*d*arctan(c*x)^2-3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1))^3*arctan(c*x)^2-3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^3*arctan(c*x)^2+3/4*I*b^3*d^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^3*arctan(c*x)^2+a*b^2/c^2*e^2*arctan(c*x)*ln(c^2*x^2+1)+3*a*b^2/c*e*ln(c^2*x^2+1)*d+3*a*b^2/c*e*arctan(c*x)^2*d-3/2*I*a*b^2*ln(c*x-I)*ln(c^2*x^2+1)*d^2-3/2*I*a*b^2*ln(1/2*I*(c*x-I))*ln(c*x+I)*d^2+3/2*I*a*b^2*ln(-1/2*I*(c*x+I))*ln(c*x-I)*d^2+3/2*I*a*b^2*ln(c*x+I)*ln(c^2*x^2+1)*d^2+1/2*I*a*b^2/c^2*e^2*dilog(1/2*I*(c*x-I))-1/2*I*a*b^2/c^2*e^2*dilog(-1/2*I*(c*x+I))+1/4*I*a*b^2/c^2*e^2*ln(c*x+I)^2-1/4*I*a*b^2/c^2*e^2*ln(c*x-I)^2+3*a^2*b/c*e*arctan(c*x)*d+1/3*b^3*c*e^2*arctan(c*x)^3*x^3-1/2*b^3*e^2*arctan(c*x)^2*x^2+b^3/c^2*e^2*arctan(c*x)*x+a*b^2/c*e^2*x+3*a*b^2*arctan(c*x)^2*d^2*2*c*x-1/2*a^2*b*e^2*x^2+3*a^2*b*arctan(c*x)*d^2*2*c*x+b^3*arctan(c*x)^3*d^2*2*c*x-b^3/c^2*e^2*ln(2)*arctan(c*x)^2+1/2*b^3/c^2*e^2*arctan(c*x)^2*ln(c^2*x^2+1)-b^3/c^2*e^2*ln((1+I*c*x)/(c^2*x^2+1)^{(1/2})*arctan(c*x)^2+b^3/c*e*arctan(c*x)^3*d-3*I*b^3*arctan(c*x)*polylog(2,-(1+I*c*x)^2/(c^2*x^2+1))*d^2+1/3*I*b^3/c^2*e^2*arctan(c*x)^3-I*b^3/c^2*e^2*arctan(c*x)-3*a*b^2*arctan(c*x)*ln(c^2*x^2+1)*d^2-a*b^2/c^2*e^2*arctan(c*x)-3/2*I*a*b^2*dilog(1/2*I*(c*x-I))*d^2+3/2*I*a*b^2*dilog(-1/2*I*(c*x+I))*d^2+3/4*I*a*b^2*ln(c*x-I)^2*d^2-3/4*I*a*b^2*ln(c*x+I)^2*d^2+1/2*a^2*b/c^2*e^2*ln(c^2*x^2+1)
\end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^2*(a+b*arctan(c*x))^3,x, algorithm="maxima")
[Out] 7/32*b^3*d^2*arctan(c*x)^4/c + 56*b^3*c^2*d*e*integrate(1/32*x^3*arctan(c*x)^3/(c^2*x^2 + 1), x) + 6*b^3*c^2*d*e*integrate(1/32*x^3*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 192*a*b^2*c^2*d*e*integrate(1/32*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + 28*b^3*c^2*d^2*integrate(1/32*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x)
```

```

3/(c^2*x^2 + 1), x) + 12*b^3*c^2*d*e*integrate(1/32*x^3*arctan(c*x)*log(c^2
*x^2 + 1)/(c^2*x^2 + 1), x) + 3*b^3*c^2*d^2*integrate(1/32*x^2*arctan(c*x)*
log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 96*a*b^2*c^2*d^2*integrate(1/32*x^2*
arctan(c*x)^2/(c^2*x^2 + 1), x) + 12*b^3*c^2*d^2*integrate(1/32*x^2*arctan(
c*x)*log(c^2*x^2 + 1)/(c^2*x^2 + 1), x) + a*b^2*d^2*arctan(c*x)^3/c + 1/3*a
^3*x^3*e^2 + a^3*d*x^2*e + 28*b^3*c^2*e^2*integrate(1/32*x^4*arctan(c*x)^3/
(c^2*x^2 + 1), x) + 3*b^3*c^2*e^2*integrate(1/32*x^4*arctan(c*x)*log(c^2*x^
2 + 1)^2/(c^2*x^2 + 1), x) + 96*a*b^2*c^2*e^2*integrate(1/32*x^4*arctan(c*x)^
2/(c^2*x^2 + 1), x) + 4*b^3*c^2*e^2*integrate(1/32*x^4*arctan(c*x)*log(c^
2*x^2 + 1)/(c^2*x^2 + 1), x) - 12*b^3*c*d*e*integrate(1/32*x^2*arctan(c*x)^
2/(c^2*x^2 + 1), x) + 3*b^3*c*d*e*integrate(1/32*x^2*log(c^2*x^2 + 1)^2/(c^
2*x^2 + 1), x) - 12*b^3*c*d^2*integrate(1/32*x*arctan(c*x)^2/(c^2*x^2 + 1),
x) + 3*b^3*c*d^2*integrate(1/32*x*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + a
^3*d^2*x + 3*(x^2*arctan(c*x) - c*(x/c^2 - arctan(c*x)/c^3))*a^2*b*d*e - 4*
b^3*c*e^2*integrate(1/32*x^3*arctan(c*x)^2/(c^2*x^2 + 1), x) + b^3*c*e^2*in
tegrate(1/32*x^3*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 56*b^3*d*e*integrat
e(1/32*x*arctan(c*x)^3/(c^2*x^2 + 1), x) + 6*b^3*d*e*integrate(1/32*x*arcta
n(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 192*a*b^2*d*e*integrate(1/32*
x*arctan(c*x)^2/(c^2*x^2 + 1), x) + 3*b^3*d^2*integrate(1/32*arctan(c*x)*lo
g(c^2*x^2 + 1)^2/(c^2*x^2 + 1), x) + 3/2*(2*c*x*arctan(c*x) - log(c^2*x^2 +
1))*a^2*b*d^2/c + 1/2*(2*x^3*arctan(c*x) - c*(x^2/c^2 - log(c^2*x^2 + 1)/c
^4))*a^2*b*e^2 + 28*b^3*e^2*integrate(1/32*x^2*arctan(c*x)^3/(c^2*x^2 + 1),
x) + 3*b^3*e^2*integrate(1/32*x^2*arctan(c*x)*log(c^2*x^2 + 1)^2/(c^2*x^2
+ 1), x) + 96*a*b^2*e^2*integrate(1/32*x^2*arctan(c*x)^2/(c^2*x^2 + 1), x)
+ 1/24*(b^3*x^3*e^2 + 3*b^3*d*x^2*e + 3*b^3*d^2*x)*arctan(c*x)^3 - 1/32*(b^
3*x^3*e^2 + 3*b^3*d*x^2*e + 3*b^3*d^2*x)*arctan(c*x)*log(c^2*x^2 + 1)^2

```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```

[In] integrate((e*x+d)^2*(a+b*arctan(c*x))^3,x, algorithm="fricas")
[Out] integral(a^3*x^2*e^2 + 2*a^3*d*x*e + a^3*d^2 + (b^3*x^2*e^2 + 2*b^3*d*x*e +
b^3*d^2)*arctan(c*x)^3 + 3*(a*b^2*x^2*e^2 + 2*a*b^2*d*x*e + a*b^2*d^2)*arc
tan(c*x)^2 + 3*(a^2*b*x^2*e^2 + 2*a^2*b*d*x*e + a^2*b*d^2)*arctan(c*x), x)

```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx))^3 (d + ex)^2 \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)**2*(a+b*atan(c*x))**3,x)`
[Out] `Integral((a + b*atan(c*x))**3*(d + e*x)**2, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x))^3,x, algorithm="giac")`
[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a + b \operatorname{atan}(c x))^3 (d + e x)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^3*(d + e*x)^2,x)`
[Out] `int((a + b*atan(c*x))^3*(d + e*x)^2, x)`

$$3.17 \quad \int (d + ex)(a + b \operatorname{ArcTan}(cx))^3 dx$$

Optimal. Leaf size=264

$$\frac{-\frac{3ibe(a + b \operatorname{ArcTan}(cx))^2}{c^2} - \frac{3bex(a + b \operatorname{ArcTan}(cx))^2}{2c} + \frac{id(a + b \operatorname{ArcTan}(cx))^3}{c}}{2e} - \frac{\left(d^2 - \frac{e^2}{c^2}\right)(a + b \operatorname{ArcTan}(cx))}{2e}$$

[Out] $-3/2*I*b*e*(a+b*\arctan(c*x))^2/c^2-3/2*b*e*x*(a+b*\arctan(c*x))^2/c+I*d*(a+b*\arctan(c*x))^3/c-1/2*(d^2-e^2/c^2)*(a+b*\arctan(c*x))^3/e+1/2*(e*x+d)^2*(a+b*\arctan(c*x))^3/e-3*b^2*e*(a+b*\arctan(c*x))*\ln(2/(1+I*c*x))/c^2+3*b*d*(a+b*\arctan(c*x))^2*\ln(2/(1+I*c*x))/c-3/2*I*b^3*e*\text{polylog}(2,1-2/(1+I*c*x))/c^2+3*I*b^2*d*(a+b*\arctan(c*x))*\text{polylog}(2,1-2/(1+I*c*x))/c+3/2*b^3*d*\text{polylog}(3,1-2/(1+I*c*x))/c$

Rubi [A]

time = 0.38, antiderivative size = 264, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 10, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.625,
Rules used = {4974, 4930, 5040, 4964, 2449, 2352, 5104, 5004, 5114, 6745}

$$-\frac{3b^2e \log \left(\frac{d}{1+I c x}\right) (a+b \operatorname{ArcTan}(cx))}{c^2}+\frac{3b^2 d L_2\left(1-\frac{d}{1+I c x}\right) (a+b \operatorname{ArcTan}(cx))}{c}-\frac{\left(d^2-\frac{e^2}{c^2}\right) (a+b \operatorname{ArcTan}(cx))^3}{2 e}-\frac{3 i b e (a+b \operatorname{ArcTan}(cx))^2}{2 c^2}+\frac{(d+e x)^2 (a+b \operatorname{ArcTan}(cx))^3}{2 e}+\frac{i d (a+b \operatorname{ArcTan}(cx))^3}{c}+\frac{3 b d \log \left(\frac{d}{1+I c x}\right) (a+b \operatorname{ArcTan}(cx))^2}{c}-\frac{3 b e x (a+b \operatorname{ArcTan}(cx))^2}{2 c}-\frac{3 i b^2 e L_2\left(1-\frac{d}{1+I c x}\right)}{2 c^2}+\frac{3 b^2 d L_3\left(1-\frac{d}{1+I c x}\right)}{2 c}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)*(a + b*\operatorname{ArcTan}[c*x])^3, x]$

[Out] $(((-3*I)/2)*b*e*(a + b*\operatorname{ArcTan}[c*x])^2)/c^2 - (3*b*e*x*(a + b*\operatorname{ArcTan}[c*x])^2)/(2*c) + (I*d*(a + b*\operatorname{ArcTan}[c*x])^3)/c - ((d^2 - e^2/c^2)*(a + b*\operatorname{ArcTan}[c*x])^3)/(2*e) + ((d + e*x)^2*(a + b*\operatorname{ArcTan}[c*x])^3)/(2*e) - (3*b^2*e*(a + b*\operatorname{ArcTan}[c*x])*Log[2/(1 + I*c*x)])/c^2 + (3*b*d*(a + b*\operatorname{ArcTan}[c*x])^2*Log[2/(1 + I*c*x)])/c - (((3*I)/2)*b^3*e*PolyLog[2, 1 - 2/(1 + I*c*x)])/c^2 + ((3*I)*b^2*d*(a + b*\operatorname{ArcTan}[c*x])*PolyLog[2, 1 - 2/(1 + I*c*x)])/c + (3*b^3*d*PolyLog[3, 1 - 2/(1 + I*c*x)])/(2*c)$

Rule 2352

$\operatorname{Int}[\operatorname{Log}[(c_*)(x_*)]/((d_) + (e_*)(x_)), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[-e^{(-1)} * \operatorname{PolyLog}[2, 1 - c*x], x] /; \operatorname{FreeQ}[\{c, d, e\}, x] \& \operatorname{EqQ}[e + c*d, 0]$

Rule 2449

$\operatorname{Int}[\operatorname{Log}[(c_*)/((d_) + (e_*)(x_))]/((f_) + (g_*)(x_)^2), x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[-e/g, \operatorname{Subst}[\operatorname{Int}[\operatorname{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \operatorname{FreeQ}[\{c, d, e, f, g\}, x] \& \operatorname{EqQ}[c, 2*d] \& \operatorname{EqQ}[e^2*f + d^2*g, 0]$

Rule 4930

$\operatorname{Int}[((a_*) + \operatorname{ArcTan}[(c_*)(x_*)^{(n_*)}](b_*))^p, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[x*(a + b*\operatorname{ArcTan}[c*x^n])^p, x] - \operatorname{Dist}[b*c*n*p, \operatorname{Int}[x^n*((a + b*\operatorname{ArcTan}[c*x^n])^p$

```
- 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] &&
(EqQ[n, 1] || EqQ[p, 1])
```

Rule 4964

```
Int[((a_.) + ArcTan[(c_)*(x_)*(b_.)])^(p_.)/((d_) + (e_)*(x_)), x_Symbol]
  :> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e),
  Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)),
  x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_)*(x_)*(b_.)])^(p_)*((d_) + (e_)*(x_))^q, x_Symbol]
  :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - Dist[b*c*(p/(e*(q + 1))),
  Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (d + e*x)^(q + 1)/(1 + c^2*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] &&
  IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_)*(x_)*(b_.)])^(p_.)/((d_) + (e_)*(x_)^2), x_Symbol]
  :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_)*(x_)*(b_.)])^(p_.)*(x_))/((d_) + (e_)*(x_)^2),
x_Symbol) :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Dist[1/(c*d),
Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_)*(x_)*(b_.)])^(p_.)*(x_))/((f_) + (g_)*(x_)^m))/((d_) + (e_)*(x_)^2,
x_Symbol) :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rule 5114

```
Int[(Log[u_]*((a_.) + ArcTan[(c_)*(x_)*(b_.)])^(p_.))/((d_) + (e_)*(x_)^2),
x_Symbol) :> Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)), x] + Dist[b*p*(I/2),
Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]
```

Rule 6745

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simplify[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Rubi steps

$$\begin{aligned}
\int (d + ex) (a + b \tan^{-1}(cx))^3 dx &= \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^3}{2e} - \frac{(3bc) \int \left(\frac{e^2 (a + b \tan^{-1}(cx))^2}{c^2} + \frac{(c^2 d^2 - e^2 + 2c^2 de)}{c^2 (1 + c^2 x^2)} \right) dx}{2e} \\
&= \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^3}{2e} - \frac{(3b) \int \frac{(c^2 d^2 - e^2 + 2c^2 dex)(a + b \tan^{-1}(cx))^2}{1 + c^2 x^2} dx}{2ce} \\
&= -\frac{3bex(a + b \tan^{-1}(cx))^2}{2c} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^3}{2e} - \frac{(3b) \int \left(\frac{c^2 d^2 - e^2 + 2c^2 dex}{1 + c^2 x^2} \right) dx}{2ce} \\
&= -\frac{3ibe(a + b \tan^{-1}(cx))^2}{2c^2} - \frac{3bex(a + b \tan^{-1}(cx))^2}{2c} + \frac{(d + ex)^2 (a + b \tan^{-1}(cx))^3}{2e} \\
&= -\frac{3ibe(a + b \tan^{-1}(cx))^2}{2c^2} - \frac{3bex(a + b \tan^{-1}(cx))^2}{2c} + \frac{id(a + b \tan^{-1}(cx))^3}{c} \\
&= -\frac{3ibe(a + b \tan^{-1}(cx))^2}{2c^2} - \frac{3bex(a + b \tan^{-1}(cx))^2}{2c} + \frac{id(a + b \tan^{-1}(cx))^3}{c} \\
&= -\frac{3ibe(a + b \tan^{-1}(cx))^2}{2c^2} - \frac{3bex(a + b \tan^{-1}(cx))^2}{2c} + \frac{id(a + b \tan^{-1}(cx))^3}{c}
\end{aligned}$$

Mathematica [A]

time = 0.32, size = 342, normalized size = 1.30

$$a^2(-2a^2 + 3a)x + a^2x^2 + 3a^2x\text{ArcTan}(ax) + 3a^2x^3(2d + cx)\text{ArcTan}(ax) - 3a^2x^4\log(1 + c^2x^2) + 3a^2x^5(-2c\text{ArcTan}(ax) + (1 + c^2x^2)\text{ArcTan}(ax) + \log(1 + c^2x^2)) + 6a^2x^6\text{ArcTan}(ax)(-1 + c^2x)\text{ArcTan}(ax) + 2\log(1 + c^2x^2)(-2a^2\text{ArcTan}(ax) - 2d\log(2, -c^2x^2\text{ArcTan}(ax))) + 3b^2x^7\log(2, -c^2x^2\text{ArcTan}(ax)) + b^2x^8(2a\text{ArcTan}(ax)^2 + (-4 + c^2)x\text{ArcTan}(ax) + 4\log(1 + c^2x^2)\text{ArcTan}(ax)) + 3b^2x^9\log(2, -c^2x^2\text{ArcTan}(ax)) + b^2x^{10}(4a\text{ArcTan}(ax)\log(2, -c^2x^2\text{ArcTan}(ax)) + (-4a + 2c^2)x\text{ArcTan}(ax)\log(2, -c^2x^2\text{ArcTan}(ax)) + 4\log(1 + c^2x^2)\text{ArcTan}(ax))$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)*(a + b*ArcTan[c*x])^3,x]`

[Out]
$$(a^2*c*(2*a*c*d - 3*b*e)*x + a^3*c^2*e*x^2 + 3*a^2*b*e*\text{ArcTan}[c*x] + 3*a^2*b*c^2*x*(2*d + e*x)*\text{ArcTan}[c*x] - 3*a^2*b*c*d*\text{Log}[1 + c^2*x^2] + 3*a*b^2*e*(-2*c*x*\text{ArcTan}[c*x] + (1 + c^2*x^2)*\text{ArcTan}[c*x]^2 + \text{Log}[1 + c^2*x^2]) + 6*a*b^2*c*d*(\text{ArcTan}[c*x]*((-I + c*x)*\text{ArcTan}[c*x] + 2*\text{Log}[1 + E^{((2*I)*\text{ArcTan}[c*x])}]))$$

$$*x])) - I*PolyLog[2, -E^((2*I)*ArcTan[c*x])] + b^3 e*(ArcTan[c*x]*((3*I - 3*c*x)*ArcTan[c*x] + (1 + c^2*x^2)*ArcTan[c*x]^2 - 6*Log[1 + E^((2*I)*ArcTan[c*x])]) + (3*I)*PolyLog[2, -E^((2*I)*ArcTan[c*x])]) + b^3 c*d*(2*ArcTan[c*x]^2*((-I + c*x)*ArcTan[c*x] + 3*Log[1 + E^((2*I)*ArcTan[c*x])]) - (6*I)*ArcTan[c*x]*PolyLog[2, -E^((2*I)*ArcTan[c*x])] + 3*PolyLog[3, -E^((2*I)*ArcTan[c*x])]))/(2*c^2)$$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 1.45, size = 7195, normalized size = 27.25

method	result	size
derivativedivides	Expression too large to display	7195
default	Expression too large to display	7195

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)*(a+b*arctan(c*x))^3,x,method=_RETURNVERBOSE)`
[Out] result too large to display

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x))^3,x, algorithm="maxima")`
[Out]
$$\begin{aligned} & 7/32 b^3 d \arctan(c x)^4/c + 56 b^3 c^2 e \int \frac{1}{64 x^3 \arctan(c x)^3} \\ & (c^2 x^2 + 1), x + 6 b^3 c^2 e \int \frac{1}{64 x^3 \arctan(c x)} \log(c^2 x^2 \\ & + 1)^2/(c^2 x^2 + 1), x + 192 a b^2 c^2 e \int \frac{1}{64 x^3 \arctan(c x)^2} \\ & /(c^2 x^2 + 1), x + 56 b^3 c^2 d \int \frac{1}{64 x^2 \arctan(c x)^3/(c^2 x^2 \\ & + 1)}, x + 12 b^3 c^2 e \int \frac{1}{64 x^3 \arctan(c x)} \log(c^2 x^2 + 1)/(c^2 x^2 + 1), x \\ & + 6 b^3 c^2 d \int \frac{1}{64 x^2 \arctan(c x)} \log(c^2 x^2 + 1)^2/(c^2 x^2 + 1), x + 192 a b^2 c^2 d \int \frac{1}{64 x^2 \arctan(c x)^2/(c^2 x^2 + 1)}, x \\ & + 24 b^3 c^2 d \int \frac{1}{64 x^2 \arctan(c x)} \log(c^2 x^2 + 1)/(c^2 x^2 + 1), x + a b^2 d \arctan(c x)^3/c + 1/2 a^3 x^2 e - 12 b^3 c \\ & *e \int \frac{1}{64 x^2 \arctan(c x)^2/(c^2 x^2 + 1)}, x + 3 b^3 c e \int \frac{1}{64 x^2 \log(c^2 x^2 + 1)^2/(c^2 x^2 + 1)}, x - 24 b^3 c d \int \frac{1}{64 x \arctan(c x)^2/(c^2 x^2 + 1)}, x + 6 b^3 c d \int \frac{1}{64 x \log(c^2 x^2 + 1)^2/(c^2 x^2 + 1)}, x + a^3 d x + 3/2 (x^2 \arctan(c x) - c (x/c^2 - \arctan(c x)/c^3)) a^2 b e + 56 b^3 e \int \frac{1}{64 x \arctan(c x)^3/(c^2 x^2 + 1)}, x + 6 b^3 e \int \frac{1}{64 x \arctan(c x)} \log(c^2 x^2 + 1)^2/(c^2 x^2 + 1), x + 192 a b^2 e \int \frac{1}{64 x \arctan(c x)^2/(c^2 x^2 + 1)}, x + 6 b^3 d \int \frac{1}{64 \arctan(c x) \log(c^2 x^2 + 1)^2/(c^2 x^2 + 1)}, x + 3/2 (2 c x \arctan(c x) - \log(c^2 x^2 + 1)) a^2 b d/c + 1/16 (b^3 x^2 e + 2 b^3\end{aligned}$$

```
*d*x)*arctan(c*x)^3 - 3/64*(b^3*x^2*e + 2*b^3*d*x)*arctan(c*x)*log(c^2*x^2
+ 1)^2
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x))^3,x, algorithm="fricas")`

[Out] `integral(a^3*x*e + a^3*d + (b^3*x*e + b^3*d)*arctan(c*x)^3 + 3*(a*b^2*x*e + a*b^2*d)*arctan(c*x)^2 + 3*(a^2*b*x*e + a^2*b*d)*arctan(c*x), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx))^3 (d + ex) \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*atan(c*x))**3,x)`

[Out] `Integral((a + b*atan(c*x))**3*(d + e*x), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x))^3,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a + b \operatorname{atan}(cx))^3 (d + e x) \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^3*(d + e*x),x)`

[Out] `int((a + b*atan(c*x))^3*(d + e*x), x)`

3.18 $\int \frac{(a+b\text{ArcTan}(cx))^3}{d+ex} dx$

Optimal. Leaf size=320

$$-\frac{(a+b\text{ArcTan}(cx))^3 \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a+b\text{ArcTan}(cx))^3 \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{3ib(a+b\text{ArcTan}(cx))^2 \text{PolyLog}(1, \frac{2c(d+ex)}{(cd+ie)(1-icx)})}{2e}$$

[Out] $-(a+b*\arctan(cx))^3 \ln(2/(1-I*c*x))/e + (a+b*\arctan(cx))^3 \ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e + 3/2*I*b*(a+b*\arctan(cx))^2 \text{polylog}(2, 1-2/(1-I*c*x))/e - 3/2*I*b*(a+b*\arctan(cx))^2 \text{polylog}(2, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e - 3/2*b^2*(a+b*\arctan(cx))*\text{polylog}(3, 1-2/(1-I*c*x))/e + 3/2*b^2*(a+b*\arctan(cx))*\text{polylog}(3, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e - 3/4*I*b^3*\text{polylog}(4, 1-2/(1-I*c*x))/e + 3/4*I*b^3*\text{polylog}(4, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/e$

Rubi [A]

time = 0.04, antiderivative size = 320, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.056, Rules used = {4970}

$$\frac{3b^2(a+b\text{ArcTan}(cx))\text{Li}_1\left(1-\frac{2(c(d+ex))}{(cd+ie)(1-icx)}\right)}{2e} - \frac{3b^2\text{Li}_1\left(1-\frac{2}{1-icx}\right)(a+b\text{ArcTan}(cx))}{2e} - \frac{3ib(a+b\text{ArcTan}(cx))^2\text{Li}_2\left(1-\frac{2(c(d+ex))}{(cd+ie)(1-icx)}\right)}{2e} + \frac{(a+b\text{ArcTan}(cx))^3 \log\left(\frac{2c(d+ex)}{(1-icx)(cd+ie)}\right)}{e} + \frac{3i\text{Li}_2\left(1-\frac{2}{1-icx}\right)(a+b\text{ArcTan}(cx))^2}{2e} - \frac{\log\left(\frac{2}{1-icx}\right)(a+b\text{ArcTan}(cx))^3}{e} + \frac{3ib^2\text{Li}_3\left(1-\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{4e} - \frac{3ib^2\text{Li}_4\left(1-\frac{2}{1-icx}\right)}{4e}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a+b*\text{ArcTan}[c*x])^3/(d+e*x), x]$

[Out] $-(((a+b*\text{ArcTan}[c*x])^3 \text{Log}[2/(1-I*c*x)])/e) + ((a+b*\text{ArcTan}[c*x])^3 \text{Log}[(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))]/e) + (((3*I)/2)*b*(a+b*\text{ArcTan}[c*x])^2 \text{PolyLog}[2, 1-2/(1-I*c*x)]/e - (((3*I)/2)*b*(a+b*\text{ArcTan}[c*x])^2 \text{PolyLog}[2, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))]/e - (3*b^2*(a+b*\text{ArcTan}[c*x])* \text{PolyLog}[3, 1-2/(1-I*c*x)]/(2*e) + (3*b^2*(a+b*\text{ArcTan}[c*x])* \text{PolyLog}[3, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))]/(2*e) - (((3*I)/4)*b^3*\text{PolyLog}[4, 1-2/(1-I*c*x)]/e + (((3*I)/4)*b^3*\text{PolyLog}[4, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))])/e$

Rule 4970

```
Int[((a_.) + ArcTan[(c_.)*(x_.)*(b_.)])^3/((d_) + (e_)*(x_)), x_Symbol] :>
Simp[(-(a + b*ArcTan[c*x])^3)*(Log[2/(1 - I*c*x)]/e), x] + (Simp[(a + b*ArcTan[c*x])^3*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x] + Simp[3*I*b*(a + b*ArcTan[c*x])^2*(PolyLog[2, 1 - 2/(1 - I*c*x)]/(2*e)), x] - Simp[3*I*b*(a + b*ArcTan[c*x])^2*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x] - Simp[3*b^2*(a + b*ArcTan[c*x])* (PolyLog[3, 1 - 2/(1 - I*c*x)]/(2*e)), x] + Simp[3*b^2*(a + b*ArcTan[c*x])* (PolyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x] - Simp[3*I*b^3*(PolyLog[4, 1 - 2/(1 - I*c*x)]/(4*e)), x] + Simp[3*I*b^3*(PolyLog[4, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(4*e)), x]) /; FreeQ[{a, b, c, d, e}, x] &
```

& $\text{NeQ}[c^2 d^2 + e^2, 0]$

Rubi steps

$$\int \frac{(a + b \tan^{-1}(cx))^3}{d + ex} dx = -\frac{(a + b \tan^{-1}(cx))^3 \log\left(\frac{2}{1-icx}\right)}{e} + \frac{(a + b \tan^{-1}(cx))^3 \log\left(\frac{2c(d+ex)}{(cd+ie)(1-icx)}\right)}{e} + \frac{3ib(a + b \tan^{-1}(cx))^3}{e}$$

Mathematica [F]

time = 137.07, size = 0, normalized size = 0.00

$$\int \frac{(a + b \text{ArcTan}(cx))^3}{d + ex} dx$$

Verification is not applicable to the result.

[In] `Integrate[(a + b*ArcTan[c*x])^3/(d + e*x), x]`

[Out] `Integrate[(a + b*ArcTan[c*x])^3/(d + e*x), x]`

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
time = 2.78, size = 2664, normalized size = 8.32

method	result	size
derivativedivides	Expression too large to display	2664
default	Expression too large to display	2664

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x))^3/(e*x+d), x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & \frac{1}{c} \left(3a^2 b c \ln(c e x + c d) / e \arctan(c x) + a^3 c \ln(c e x + c d) / e + 1/2 I b^3 c \right. \\ & \left. / e \arctan(c x) \right)^3 c \operatorname{sgn}(I(-I e (1+I c x)^2 / (c^2 x^2 + 1) + c d (1+I c x)^2 / (c^2 x^2 + 1) + I e + c d) / ((1+I c x)^2 / (c^2 x^2 + 1) + 1))^{3/2} \operatorname{Pi} + 3/2 I a^2 b c \ln(c e x + c d) / e \ln((I e - c e x) / (c d + I e)) - 3/2 I a^2 b^2 c \ln(c e x + c d) / e \ln((I e + c e x) / (I e - c d)) + 3/2 a b^2 c^2 / e d (-I e + c d) \operatorname{polylog}(3, (I e - c d) / (c d + I e) * (1+I c x)^2 / (c^2 x^2 + 1)) + 3 I a b^2 c / e \arctan(c x) \operatorname{polylog}(2, -(1+I c x)^2 / (c^2 x^2 + 1)) - 3 I a b^2 c \arctan(c x) \operatorname{polylog}(2, (I e - c d) / (c d + I e) * (1+I c x)^2 / (c^2 x^2 + 1)) / (e + I c d) - 3/2 a b^2 c / e \operatorname{polylog}(3, -(1+I c x)^2 / (c^2 x^2 + 1)) + 3/2 a b^2 c \operatorname{polylog}(3, (I e - c d) / (c d + I e) * (1+I c x)^2 / (c^2 x^2 + 1)) / (e + I c d) + b^3 c \ln(c e x + c d) / e \arctan(c x) \right)^3 - b^3 c \operatorname{arctan}(c x) \operatorname{ln}(-I e (1+I c x)^2 / (c^2 x^2 + 1) + c d (1+I c x)^2 / (c^2 x^2 + 1) + I e + c d) - 3/2 b^3 c / e \arctan(c x) \operatorname{polylog}(3, -(1+I c x)^2 / (c^2 x^2 + 1)) + b^3 c \arctan(c x) \operatorname{ln}(1 - (I e - c d) / (c d + I e) * (1+I c x)^2 / (c^2 x^2 + 1)) / (e + I c d) + 3/2 b^3 c \operatorname{arctan}(c x) \operatorname{polylog}(3, (I e - c d) / (c d + I e) * (1+I c x)^2 / (c^2 x^2 + 1)) / (e + I c d) \end{aligned}$$

$$\begin{aligned}
& \text{I*}e\text{-c*d})/(c*d+I*c*x)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)+3/4*I*b^3*c*\text{polylog}(4 \\
& , (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)-3/4*I*b^3*c/e*\text{poly} \\
& \text{og}(4, -(1+I*c*x)^2/(c^2*x^2+1))+3/2*I*a^2*b*c/e*\text{dilog}((I*e-c*x)/(c*d+I*e))) \\
& -3/2*I*a^2*b*c/e*\text{dilog}((I*e+c*x)/(I*e-c*d))+3*a*b^2*c*\ln(c*x+c*d)/e*\text{arc} \\
& \tan(c*x)^2-3*a*b^2*c/e*\arctan(c*x)^2*\ln(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1 \\
& +I*c*x)^2/(c^2*x^2+1)+I*e+c*d)+3*a*b^2*c*\arctan(c*x)^2*\ln(1-(I*e-c*d)/(c*d+ \\
& I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)-3/2*I*b^3*c*\arctan(c*x)^2*\text{polylog}(2 \\
& , (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(e+I*c*d)+3/2*I*b^3*c/e*\arcta \\
& n(c*x)^2*\text{polylog}(2, -(1+I*c*x)^2/(c^2*x^2+1))+3/2*I*a*b^2*c/e*\arctan(c*x)^2* \\
& \text{csgn}(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/(\\
& (1+I*c*x)^2/(c^2*x^2+1)+1))*\text{csgn}(I/((1+I*c*x)^2/(c^2*x^2+1)+1))*\text{csgn}(I*(-I* \\
& e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))*\text{Pi}-1/2*I*b^ \\
& 3*c/e*\arctan(c*x)^3*\text{csgn}(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c \\
& ^2*x^2+1)+I*e+c*d)/(1+I*c*x)^2/(c^2*x^2+1)+1))^2*\text{csgn}(I*(-I*e*(1+I*c*x)^2/ \\
& (c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))*\text{Pi}-3/2*I*b^3*c^2/e*d/(-I* \\
& e+c*d)*\arctan(c*x)^2*\text{polylog}(2, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1)) \\
& -1/2*I*b^3*c/e*\arctan(c*x)^3*\text{csgn}(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I* \\
& c*x)^2/(c^2*x^2+1)+I*e+c*d)/(1+I*c*x)^2/(c^2*x^2+1)+1))^2*\text{csgn}(I/((1+I*c*x) \\
&)^2/(c^2*x^2+1)+1))*\text{Pi}+3*a*b^2*c^2/e*d/(-I*e+c*d)*\arctan(c*x)^2*\ln(1-(I*e-c \\
& *d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))+3/2*I*a*b^2*c/e*\arctan(c*x)^2*\text{csgn}(I \\
& *(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/(1+I*c*x) \\
&)^2/(c^2*x^2+1)+1))^3*\text{Pi}+1/2*I*b^3*c/e*\arctan(c*x)^3*\text{csgn}(I*(-I*e*(1+I*c*x) \\
&)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/(1+I*c*x)^2/(c^2*x^2+1) \\
& +1))*\text{csgn}(I/((1+I*c*x)^2/(c^2*x^2+1)+1))*\text{csgn}(I*(-I*e*(1+I*c*x)^2/(c^2*x \\
& ^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))*\text{Pi}-3/2*I*a*b^2*c/e*\arctan(c*x)^ \\
& 2*\text{csgn}(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d) \\
& /(1+I*c*x)^2/(c^2*x^2+1)+1))^2*\text{csgn}(I/((1+I*c*x)^2/(c^2*x^2+1)+1))*\text{Pi}-3/2* \\
& I*a*b^2*c/e*\arctan(c*x)^2*\text{csgn}(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x) \\
&)^2/(c^2*x^2+1)+I*e+c*d)/(1+I*c*x)^2/(c^2*x^2+1)+1))^2*\text{csgn}(I*(-I*e*(1+I*c*x) \\
&)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))*\text{Pi}-3*I*a*b^2*c^2/e* \\
& d/(-I*e+c*d)*\arctan(c*x)*\text{polylog}(2, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2 \\
& +1))+b^3*c^2/e*d/(-I*e+c*d)*\arctan(c*x)^3*\ln(1-(I*e-c*d)/(c*d+I*e)*(1+I*c*x) \\
&)^2/(c^2*x^2+1))+3/2*b^3*c^2/e*d/(-I*e+c*d)*\arctan(c*x)*\text{polylog}(3, (I*e-c*d) \\
& /(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))+3/4*I*b^3*c^2/e*d/(-I*e+c*d)*\text{polylog}(4, \\
& (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1)))
\end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^3/(e*x+d),x, algorithm="maxima")
[Out] a^3*e^(-1)*log(x*e + d) + integrate(1/32*(28*b^3*arctan(c*x)^3 + 3*b^3*arctan(c*x)^5)/(x*d + e), x)
```

$\text{an}(c*x)*\log(c^2*x^2 + 1)^2 + 96*a*b^2*\arctan(c*x)^2 + 96*a^2*b*\arctan(c*x)) / (x*e + d), x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b*\arctan(c*x))^3/(e*x+d), x, \text{algorithm}=\text{"fricas"})$

[Out] $\text{integral}((b^3*\arctan(c*x)^3 + 3*a*b^2*\arctan(c*x)^2 + 3*a^2*b*\arctan(c*x) + a^3)/(x*e + d), x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + b \tan(cx))^3}{d + ex} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b*\atan(c*x))^3/(e*x+d), x)$

[Out] $\text{Integral}((a + b*\atan(c*x))^3/(d + e*x), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b*\arctan(c*x))^3/(e*x+d), x, \text{algorithm}=\text{"giac"})$

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a + b \tan(cx))^3}{d + ex} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((a + b*\atan(c*x))^3/(d + e*x), x)$

[Out] $\text{int}((a + b*\atan(c*x))^3/(d + e*x), x)$

$$3.19 \quad \int \frac{(a+b\text{ArcTan}(cx))^3}{(d+ex)^2} dx$$

Optimal. Leaf size=499

$$\frac{ic(a+b\text{ArcTan}(cx))^3}{c^2d^2+e^2} + \frac{c^2d(a+b\text{ArcTan}(cx))^3}{e(c^2d^2+e^2)} - \frac{(a+b\text{ArcTan}(cx))^3}{e(d+ex)} - \frac{3bc(a+b\text{ArcTan}(cx))^2 \log\left(\frac{2}{1-icx}\right)}{c^2d^2+e^2} + \dots$$

[Out] $I*c*(a+b*\arctan(c*x))^3/(c^2*d^2+e^2)+c^2*d*(a+b*\arctan(c*x))^3/e/(c^2*d^2+e^2)-(a+b*\arctan(c*x))^3/e/(e*x+d)-3*b*c*(a+b*\arctan(c*x))^2*\ln(2/(1-I*c*x))/(c^2*d^2+e^2)+3*b*c*(a+b*\arctan(c*x))^2*\ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)+3*I*b^2*c*(a+b*\arctan(c*x))*\text{polylog}(2,1-2/(1-I*c*x))/(c^2*d^2+e^2)+3*I*b^2*c*(a+b*\arctan(c*x))*\text{polylog}(2,1-2/(1+I*c*x))/(c^2*d^2+e^2)-3*I*b^2*c*(a+b*\arctan(c*x))*\text{polylog}(2,1-2/(1-I*c*x))/(c^2*d^2+e^2)-3/2*b^3*c*\text{polylog}(3,1-2/(1-I*c*x))/(c^2*d^2+e^2)+3/2*b^3*c*\text{polylog}(3,1-2/(1+I*c*x))/(c^2*d^2+e^2)+3/2*b^3*c*\text{polylog}(3,1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)+3/2*b^3*c*\text{polylog}(3,1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)$

Rubi [A]

time = 0.36, antiderivative size = 499, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {4974, 4968, 5104, 5004, 5040, 4964, 5114, 6745}

$$\frac{3b^2\text{Li}_2\left(1-\frac{e^{2ix}}{c^2d^2+e^2}\right)(a+b\text{ArcTan}(cx))}{c^2d^2+e^2} + \frac{3b^2\text{Li}_2\left(1-\frac{e^{-2ix}}{c^2d^2+e^2}\right)(a+b\text{ArcTan}(cx))}{c^2d^2+e^2} - \frac{3b^2c(a+b\text{ArcTan}(cx))\text{Li}_2\left(1-\frac{e^{2ix}}{c^2d^2+e^2}\right)}{c^2d^2+e^2} + \frac{ic(a+b\text{ArcTan}(cx))^2}{c^2d^2+e^2} - \frac{3ie\log\left(\frac{e^{2ix}}{c^2d^2+e^2}\right)(a+b\text{ArcTan}(cx))^2}{c^2d^2+e^2} + \frac{3ie\log\left(\frac{e^{-2ix}}{c^2d^2+e^2}\right)(a+b\text{ArcTan}(cx))^2}{c^2d^2+e^2} + \frac{3bc(a+b\text{ArcTan}(cx))^2 \log\left(\frac{2e^{2ix}}{1-e^{2icx}(1-icx)}\right)}{c^2d^2+e^2} - \frac{(a+b\text{ArcTan}(cx))^2}{c^2d^2+e^2} - \frac{3b^2\text{Li}_2\left(1-\frac{e^{2ix}}{c^2d^2+e^2}\right)}{2(c^2d^2+e^2)} + \frac{3b^2\text{Li}_2\left(1-\frac{e^{-2ix}}{c^2d^2+e^2}\right)}{2(c^2d^2+e^2)} + \frac{3b^2\text{Li}_2\left(1-\frac{e^{2ix}}{c^2d^2+e^2}\right)}{2(c^2d^2+e^2)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a+b*\text{ArcTan}[c*x])^3/(d+e*x)^2, x]$

[Out] $(I*c*(a+b*\text{ArcTan}[c*x])^3/(c^2*d^2+e^2) + (c^2*d*(a+b*\text{ArcTan}[c*x])^3)/(e*(c^2*d^2+e^2)) - (a+b*\text{ArcTan}[c*x])^3/(e*(d+e*x)) - (3*b*c*(a+b*\text{ArcTan}[c*x])^2*\text{Log}[2/(1-I*c*x)])/(c^2*d^2+e^2) + (3*b*c*(a+b*\text{ArcTan}[c*x])^2*\text{Log}[2/(1+I*c*x)])/(c^2*d^2+e^2) + (3*b*c*(a+b*\text{ArcTan}[c*x])^2*\text{PolyLog}[2, 1-2/(1-I*c*x)])/(c^2*d^2+e^2) + ((3*I)*b^2*c*(a+b*\text{ArcTan}[c*x])* \text{PolyLog}[2, 1-2/(1+I*c*x)])/(c^2*d^2+e^2) - ((3*I)*b^2*c*(a+b*\text{ArcTan}[c*x])* \text{PolyLog}[2, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))])/(c^2*d^2+e^2) - (3*b^3*c*\text{PolyLog}[3, 1-2/(1-I*c*x)])/(2*(c^2*d^2+e^2)) + (3*b^3*c*\text{PolyLog}[3, 1-2/(1+I*c*x)])/(2*(c^2*d^2+e^2)) + (3*b^3*c*\text{PolyLog}[3, 1-(2*c*(d+e*x))/((c*d+I*e)*(1-I*c*x))])/(2*(c^2*d^2+e^2))$

Rule 4964

```
Int[((a_) + ArcTan[(c_)*(x_)]*(b_))^(p_)/((d_) + (e_)*(x_)), x_Symbol]
:> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(
```

```
p/e), Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4968

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.)]^2/((d_) + (e_.)*(x_)), x_Symbol] :>
Simp[(-(a + b*ArcTan[c*x])^2)*(Log[2/(1 - I*c*x)]/e), x] + (Simp[(a + b*Arc
Tan[c*x])^2*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x] + Simp[I
*b*(a + b*ArcTan[c*x])*(PolyLog[2, 1 - 2/(1 - I*c*x)]/e), x] - Simp[I*b*(a
+ b*ArcTan[c*x])*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e),
x] - Simp[b^2*(PolyLog[3, 1 - 2/(1 - I*c*x)]/(2*e)), x] + Simp[b^2*(Po
lyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x]) /; Free
Q[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.)]^(p_)*((d_) + (e_.)*(x_))^(q_.), x_Sy
mbol] :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - D
ist[b*c*(p/(e*(q + 1))), Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1), (
d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] &&
IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.)]^(p_.)/((d_) + (e_.)*(x_)^2), x_Symbo
l] :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b,
c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.)]^(p_.)*((x_))/((d_) + (e_.)*(x_)^2),
x_Symbol] :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Di
st[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.)]^(p_.)*((f_) + (g_.)*(x_))^(m_.)/((
d_) + (e_.)*(x_)^2), x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p
/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGt
Q[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rule 5114

```
Int[(Log[u_]*((a_.) + ArcTan[(c_.)*(x_)]*(b_.)]^(p_.))/((d_) + (e_.)*(x_)^2),
x_Symbol] :> Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d))
```

```
, x] + Dist[b*p*(I/2), Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]
```

Rule 6745

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simpl[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a + b \tan^{-1}(cx))^3}{(d + ex)^2} dx &= -\frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} + \frac{(3bc) \int \left(\frac{e^2(a+b \tan^{-1}(cx))^2}{(c^2d^2+e^2)(d+ex)} + \frac{c^2(d-ex)(a+b \tan^{-1}(cx))^2}{(c^2d^2+e^2)(1+c^2x^2)} \right) dx}{e} \\
&= -\frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} + \frac{(3bc^3) \int \frac{(d-ex)(a+b \tan^{-1}(cx))^2}{1+c^2x^2} dx}{e(c^2d^2+e^2)} + \frac{(3bce) \int \frac{(a+b \tan^{-1}(cx))^2}{d+ex} dx}{c^2d^2+e^2} \\
&= -\frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} - \frac{3bc(a + b \tan^{-1}(cx))^2 \log(\frac{2}{1-icx})}{c^2d^2+e^2} + \frac{3bc(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} \\
&= -\frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} - \frac{3bc(a + b \tan^{-1}(cx))^2 \log(\frac{2}{1-icx})}{c^2d^2+e^2} + \frac{3bc(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} \\
&= \frac{ic(a + b \tan^{-1}(cx))^3}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^3}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} - \frac{3bc(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} \\
&= \frac{ic(a + b \tan^{-1}(cx))^3}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^3}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} - \frac{3bc(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2} \\
&= \frac{ic(a + b \tan^{-1}(cx))^3}{c^2d^2+e^2} + \frac{c^2d(a + b \tan^{-1}(cx))^3}{e(c^2d^2+e^2)} - \frac{(a + b \tan^{-1}(cx))^3}{e(d + ex)} - \frac{3bc(a + b \tan^{-1}(cx))^2}{c^2d^2+e^2}
\end{aligned}$$

Mathematica [F]

time = 85.55, size = 0, normalized size = 0.00

$$\int \frac{(a + b \text{ArcTan}(cx))^3}{(d + ex)^2} dx$$

Verification is not applicable to the result.

[In] $\int (a + b \arctan(cx))^3 / (d + ex)^2 dx$
 [Out] $\int (a + b \arctan(cx))^3 / (d + ex)^2, x$

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.
 time = 2.88, size = 3050, normalized size = 6.11

method	result	size
derivativedivides	Expression too large to display	3050
default	Expression too large to display	3050

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((a+b \arctan(cx))^3 / (e*x+d)^2, x, \text{method}=_RETURNVERBOSE)$
 [Out]
$$\begin{aligned} & 1/c*(b^3*c^3/e*\arctan(cx)^3/(c^2*d^2+e^2)*d-a^3*c^2/(c*e*x+c*d)/e+3/2*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(cx)^2*Pi*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1))*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))-3/4*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(cx)^2*Pi*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)-3/2*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*x+I)*ln(1/2*I*(c*x-I))+3/2*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*x-I)*ln(-1/2*I*(c*x+I))+3*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*e*x+c*d)*ln((I*e-c*e*x)/(c*d+I*e))+3/2*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*x+I)*ln(c^2*x^2+1)-3*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*e*x+c*d)*ln((I*e+c*e*x)/(I*e-c*d))+3*a^2*b*c^3/e/(c^2*d^2+e^2)*d*\arctan(cx)+3/2*b^3*c^2*e*polylog(3, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(c^2*d^2+e^2)/(e+I*c*d)+3/2*b^3*c^3/(c^2*d^2+e^2)*d/(-I*e+c*d)*polylog(3, (I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))-3*a*b^2*c^2/(c^2*d^2+e^2)*d*\arctan(cx)-3*a*b^2*c^2*arctan(cx)/(c^2*d^2+e^2)*ln(c^2*x^2+1)+6*a*b^2*c^2*arctan(cx)/(c^2*d^2+e^2)*ln(c*e*x+c*d)-3*I*a*b^2*c^2/(c^2*d^2+e^2)*dilog((I*e+c*e*x)/(I*e-c*d))-3/4*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*x+I)^2-3/2*I*a*b^2*c^2/(c^2*d^2+e^2)*dilog(1/2*I*(c*x-I))+3/4*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*x-I)^2+3/2*I*a*b^2*c^2/(c^2*d^2+e^2)*dilog(-1/2*I*(c*x+I))+3*I*a*b^2*c^2/(c^2*d^2+e^2)*dilog((I*e-c*e*x)/(c*d+I*e))-3*a^2*b*c^2/(c*e*x+c*d)/e*\arctan(cx)+3*a*b^2*c^3/e/(c^2*d^2+e^2)*d*\arctan(cx)^2-3/2*I*a*b^2*c^2/(c^2*d^2+e^2)*ln(c*x-I)*ln(c^2*x^2+1)+3*b^3*c^3/(c^2*d^2+e^2)*d/(-I*e+c*d)*arctan(cx)^2*ln(1-(I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))-3/4*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(cx)^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)^2-3+3/2*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(cx)^2*Pi*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))^-3-3/4*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(cx)^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))^2-3+3*b^3*c^2*e*\arctan(cx)^2*ln(1-(I*e-c*d)/(c*d+I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(c^2*d^2+e^2)/(e+I*c*d)-3*I*b^3*c^3/(c^2*d^2+e^2)*d/(-I*e+c*d)*arctan(cx)*polylog(2, (I*e-c*d)/(c*d+I*e)* \end{aligned}$$

$$\begin{aligned}
& (1+I*c*x)^2/(c^2*x^2+1)) - 3*I*b^3*c^2*e*\arctan(c*x)*\text{polylog}(2, (I*e-c*d)/(c*d + I*e)*(1+I*c*x)^2/(c^2*x^2+1))/(c^2*d^2+e^2)/(e+I*c*d) + 3/4*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1))^2*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2) - 3/2*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))^2*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1)) - 3/2*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1))*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+1)^2) + 3/2*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I*((1+I*c*x)/(c^2*x^2+1))^(1/2))*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))^2 - 3/2*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+1)^2+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)/((1+I*c*x)^2/(c^2*x^2+1)+1))^2*csgn(I*(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d))/((1+I*c*x)^2/(c^2*x^2+1)+1)^2*csgn(I*((1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d)) + 3/4*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I/((1+I*c*x)^2/(c^2*x^2+1)+1)^2)*csgn(I*(1+I*c*x)^2/(c^2*x^2+1)/((1+I*c*x)^2/(c^2*x^2+1)+1)^2) + 3/4*I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*Pi*csgn(I*(1+I*c*x)^2/(c^2*x^2+1))/((1+I*c*x)^2/(c^2*x^2+1)+1)^2) - b^3*c^2/(c*e*x+c*d)/e*\arctan(c*x)^3 - 3/2*b^3*c^2*a*\arctan(c*x)^2/(c^2*d^2+e^2)*ln(c^2*x^2+1) + 3*b^3*c^2*\arctan(c*x)^2/(c^2*d^2+e^2)*ln(c*e*x+c*d) + 3*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*ln((1+I*c*x)/(c^2*x^2+1))^(1/2) - 3*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*ln(-I*e*(1+I*c*x)^2/(c^2*x^2+1)+c*d*(1+I*c*x)^2/(c^2*x^2+1)+I*e+c*d) + 3*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^2*ln(2) - I*b^3*c^2/(c^2*d^2+e^2)*\arctan(c*x)^3 - 3/2*a^2*b*c^2/(c^2*d^2+e^2)*ln(c^2*x^2+1) + 3*a^2*b*c^2/(c^2*d^2+e^2)*ln(c*e*x+c*d)
\end{aligned}$$
Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^3/(e*x+d)^2,x, algorithm="maxima")
[Out] 3/2*((2*c*d*arctan(c*x)/(c^2*d^2+e^2) - log(c^2*x^2+1)/(c^2*d^2+e^2) + 2*log(x*e+d)/(c^2*d^2+e^2))*c - 2*arctan(c*x)/(x*e^2+d*e))*a^2*b - a^3/(x*e^2+d*e) - 1/32*(4*b^3*arctan(c*x)^3 - 3*b^3*arctan(c*x)*log(c^2*x^2+1)^2 - 32*(x*e^2+d*e)*integrate(1/32*(28*(b^3*c^2*x^2+e+b^3*c)*a*\arctan(c*x)^3 + 12*(8*a*b^2*c^2*x^2+e+b^3*c*x+e+b^3*c*d+8*a*b^2*e)*\arctan(c*x)^2 - 12*(b^3*c^2*x^2+e+b^3*c^2*d*x)*\arctan(c*x)*log(c^2*x^2+1) - 3*(b^3*c*x+e+b^3*c*d - (b^3*c^2*x^2+e+b^3*c)*\arctan(c*x))*log(c^2*x^2+1)^2)/(c^2*x^4+e^3+2*c^2*d*x^3+e^2+(c^2*d^2+e^2)*x^2+2*d*x+e^2+d^2), x))/(x*e^2+d*e)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^3/(e*x+d)^2,x, algorithm="fricas")
[Out] integral((b^3*arctan(c*x)^3 + 3*a*b^2*arctan(c*x)^2 + 3*a^2*b*arctan(c*x) +
a^3)/(x^2*e^2 + 2*d*x*e + d^2), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a + b \operatorname{atan}(cx))^3}{(d + ex)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*atan(c*x))**3/(e*x+d)**2,x)
[Out] Integral((a + b*atan(c*x))**3/(d + e*x)**2, x)
```

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x))^3/(e*x+d)^2,x, algorithm="giac")
[Out] Timed out
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a + b \operatorname{atan}(cx))^3}{(d + ex)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a + b*atan(c*x))^3/(d + e*x)^2,x)
[Out] int((a + b*atan(c*x))^3/(d + e*x)^2, x)
```

$$3.20 \quad \int \frac{(a+b\text{ArcTan}(cx))^3}{(d+ex)^3} dx$$

Optimal. Leaf size=936

$$\frac{3bc^3d(a+b\text{ArcTan}(cx))^2}{2(c^2d^2+e^2)^2} + \frac{3ibc^2e(a+b\text{ArcTan}(cx))^2}{2(c^2d^2+e^2)^2} - \frac{3bc(a+b\text{ArcTan}(cx))^2}{2(c^2d^2+e^2)(d+ex)} + \frac{ic^3d(a+b\text{ArcTan}(cx))^3}{(c^2d^2+e^2)^2} + \frac{ci^2b^2d^2(a+b\text{ArcTan}(cx))^4}{(c^2d^2+e^2)^3}$$

[Out] $\frac{3}{2}b*c^3*d*(a+b*\arctan(cx))^2/(c^2*d^2+e^2)^2 + \frac{3ibc^2e(a+b*\arctan(cx))^2}{2(c^2d^2+e^2)^2} - \frac{3bc(a+b*\arctan(cx))^2}{2(c^2d^2+e^2)(d+ex)} + \frac{ic^3d(a+b*\arctan(cx))^3}{(c^2d^2+e^2)^2} + \frac{ci^2b^2d^2(a+b*\arctan(cx))^4}{(c^2d^2+e^2)^3}$

$\frac{3}{2}b*c^3*d*(a+b*\arctan(cx))^2/(c^2*d^2+e^2)^2 - 3/2*b*c*(a+b*\arctan(cx))^2/(c^2*d^2+e^2)/(e*x+d) - 3*I*b^2*c^3*d*(a+b*\arctan(cx))*\text{polylog}(2, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)^2 + 1/2*c^2*(c*d-e)*(c*d+e)*(a+b*\arctan(cx))^3/e/(c^2*d^2+e^2)^2 - 1/2*(a+b*\arctan(cx))^3/e/(e*x+d)^2 - 3*b^2*c^2*e*(a+b*\arctan(cx))*\ln(2/(1-I*c*x))/(c^2*d^2+e^2)^2 - 3*b*c^3*d*(a+b*\arctan(cx))^2*\ln(2/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3*b^2*c^2*e*(a+b*\arctan(cx))*\ln(2/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3*b*c^3*d*(a+b*\arctan(cx))^2*\ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3*b*c^3*d*(a+b*\arctan(cx))^2*\ln(2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)^2 - 3/2*I*b^2*c^2*e*\text{polylog}(2, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3*I*b^2*c^3*d*(a+b*\arctan(cx))*\text{polylog}(2, 1-2/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3/2*I*b^2*c^2*e*\text{polylog}(2, 1-2/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3/2*I*b^2*c^3*d*(a+b*\arctan(cx))^2/(c^2*d^2+e^2)^2 + I*c^3*d*(a+b*\arctan(cx))^3/(c^2*d^2+e^2)^2 + 3*I*b^2*c^3*d*(a+b*\arctan(cx))*\text{polylog}(2, 1-2/(1-I*c*x))/(c^2*d^2+e^2)^2 - 3/2*b^2*c^3*d*\text{polylog}(3, 1-2/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3/2*b^2*c^3*d*\text{polylog}(3, 1-2/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3/2*b^2*c^3*d*\text{polylog}(3, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)^2 + 3/2*b^2*c^3*d*\text{polylog}(3, 1-2*c*(e*x+d)/(c*d+I*e)/(1-I*c*x))/(c^2*d^2+e^2)^2$

Rubi [A]

time = 0.77, antiderivative size = 936, normalized size of antiderivative = 1.00, number of steps used = 23, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.667, Rules used = {4974, 4966, 2449, 2352, 2497, 5104, 5004, 5040, 4964, 4968, 5114, 6745}



Antiderivative was successfully verified.

[In] $\text{Int}[(a+b*\text{ArcTan}[c*x])^3/(d+e*x)^3, x]$

[Out] $(3*b*c^3*d*(a+b*\text{ArcTan}[c*x])^2)/(2*(c^2*d^2+e^2)^2) + (((3*I)/2)*b*c^2*e*(a+b*\text{ArcTan}[c*x])^2)/(c^2*d^2+e^2)^2 - (3*b*c*(a+b*\text{ArcTan}[c*x])^2)/(2*(c^2*d^2+e^2)*(d+e*x)) + (I*c^3*d*(a+b*\text{ArcTan}[c*x])^3)/(c^2*d^2+e^2)^2 + (c^2*(c*d-e)*(c*d+e)*(a+b*\text{ArcTan}[c*x])^3)/(2*e*(c^2*d^2+e^2)^2) - (a+b*\text{ArcTan}[c*x])^3/(2*e*(d+e*x)^2) - (3*b^2*c^2*e*(a+b*\text{ArcTan}[c*x])*Log[2/(1-I*c*x)])/(c^2*d^2+e^2)^2 - (3*b*c^3*d*(a+b*\text{ArcTan}[c*x])^2*Log[2/(1-I*c*x)])/(c^2*d^2+e^2)^2 + (3*b^2*c^2*e*(a+b*\text{ArcTan}[c*x])*Log[2/(1+I*c*x)])/(c^2*d^2+e^2)^2 + (3*b*c^3*d*(a+b*\text{ArcTan}[c*x])^2)$

$$\begin{aligned}
& 2*\text{Log}[2/(1 + I*c*x)]/(c^2*d^2 + e^2)^2 + (3*b^2*c^2*e*(a + b*\text{ArcTan}[c*x])* \\
& \text{Log}[(2*c*(d + e*x))/((c*d + I*e)*(1 - I*c*x))]/(c^2*d^2 + e^2)^2 + (3*b*c^3*d*(a + b*\text{ArcTan}[c*x])^2*\text{Log}[(2*c*(d + e*x))/((c*d + I*e)*(1 - I*c*x))])/ \\
& (c^2*d^2 + e^2)^2 + (((3*I)/2)*b^3*c^2*e*\text{PolyLog}[2, 1 - 2/(1 - I*c*x)])/(c^2*d^2 + e^2)^2 + ((3*I)*b^2*c^3*d*(a + b*\text{ArcTan}[c*x])* \\
& \text{PolyLog}[2, 1 - 2/(1 - I*c*x)])/(c^2*d^2 + e^2)^2 + (((3*I)/2)*b^3*c^2*e*\text{PolyLog}[2, 1 - 2/(1 + I*c*x)])/(c^2*d^2 + e^2)^2 + ((3*I)*b^2*c^3*d*(a + b*\text{ArcTan}[c*x])* \\
& \text{PolyLog}[2, 1 - 2/(1 + I*c*x)])/(c^2*d^2 + e^2)^2 - (((3*I)/2)*b^3*c^2*e*\text{PolyLog}[2, 1 - \\
& (2*c*(d + e*x))/((c*d + I*e)*(1 - I*c*x))]/(c^2*d^2 + e^2)^2 - ((3*I)*b^2*c^3*d*(a + b*\text{ArcTan}[c*x])* \\
& \text{PolyLog}[2, 1 - (2*c*(d + e*x))/((c*d + I*e)*(1 - I*c*x))]/(c^2*d^2 + e^2)^2 - (3*b^3*c^3*d*\text{PolyLog}[3, 1 - 2/(1 - I*c*x)])/(\\
& 2*(c^2*d^2 + e^2)^2) + (3*b^3*c^3*d*\text{PolyLog}[3, 1 - 2/(1 + I*c*x)])/(2*(c^2*d^2 + e^2)^2) + (3*b^3*c^3*d*\text{PolyLog}[3, 1 - \\
& (2*c*(d + e*x))/((c*d + I*e)*(1 - I*c*x))]/(2*(c^2*d^2 + e^2)^2)
\end{aligned}$$
Rule 2352

$$\text{Int}[\text{Log}[(c_*)*(x_*)]/((d_) + (e_*)*(x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[-e^{(-1)}*\text{PolyLog}[2, 1 - c*x], x] /; \text{FreeQ}[\{c, d, e\}, x] \&& \text{EqQ}[e + c*d, 0]$$
Rule 2449

$$\text{Int}[\text{Log}[(c_*)/((d_) + (e_*)*(x_))]/((f_) + (g_*)*(x_)^2), x_{\text{Symbol}}] \rightarrow \text{Dist}[-e/g, \text{Subst}[\text{Int}[\text{Log}[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; \text{FreeQ}[\{c, d, e, f, g\}, x] \&& \text{EqQ}[c, 2*d] \&& \text{EqQ}[e^2*f + d^2*g, 0]$$
Rule 2497

$$\text{Int}[\text{Log}[u_*]*(Pq_*)^{(m_)}, x_{\text{Symbol}}] \rightarrow \text{With}[\{C = \text{FullSimplify}[Pq^m*((1 - u)/D[u, x])]\}, \text{Simp}[C*\text{PolyLog}[2, 1 - u], x] /; \text{FreeQ}[C, x] /; \text{IntegerQ}[m] \&& \text{PolyQ}[Pq, x] \&& \text{RationalFunctionQ}[u, x] \&& \text{LeQ}[\text{RationalFunctionExponents}[u, x][[2]], \text{Expon}[Pq, x]]]$$
Rule 4964

$$\begin{aligned}
& \text{Int}[((a_*) + \text{ArcTan}[(c_*)*(x_*)]*(b_*)^{(p_*)}/((d_) + (e_*)*(x_)), x_{\text{Symbol}}] \\
& \rightarrow \text{Simp}[(-(a + b*\text{ArcTan}[c*x])^p)*(\text{Log}[2/(1 + e*(x/d))]/e), x] + \text{Dist}[b*c*(p/e), \text{Int}[(a + b*\text{ArcTan}[c*x])^{(p - 1)}*(\text{Log}[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x] /; \text{FreeQ}[\{a, b, c, d, e\}, x] \&& \text{IGtQ}[p, 0] \&& \text{EqQ}[c^2*d^2 + e^2, 0]
\end{aligned}$$
Rule 4966

$$\begin{aligned}
& \text{Int}[((a_*) + \text{ArcTan}[(c_*)*(x_*)]*(b_*)/((d_) + (e_*)*(x_)), x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(a + b*\text{ArcTan}[c*x]))*(\text{Log}[2/(1 - I*c*x)]/e), x] + (\text{Dist}[b*(c/e), \text{Int}[\text{Log}[2/(1 - I*c*x)]/(1 + c^2*x^2), x], x] - \text{Dist}[b*(c/e), \text{Int}[\text{Log}[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(1 + c^2*x^2), x], x] + \text{Simp}[(a + b*\text{ArcTan}[c*x])*(\text{Log}[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x]) /; \text{FreeQ}[\{a,
\end{aligned}$$

```
b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rule 4968

```
Int[((a_.) + ArcTan[(c_.)*(x_)]*(b_.))^(2)/((d_) + (e_.)*(x_)), x_Symbol] :>
Simp[(-(a + b*ArcTan[c*x])^2)*(Log[2/(1 - I*c*x)]/e), x] + (Simp[(a + b*Arc
Tan[c*x])^2*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x] + Simp[I
*b*(a + b*ArcTan[c*x])*(PolyLog[2, 1 - 2/(1 - I*c*x)]/e), x] - Simp[I*b*(a
+ b*ArcTan[c*x])*(PolyLog[2, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e),
x] - Simp[b^2*(PolyLog[3, 1 - 2/(1 - I*c*x)]/(2*e)), x] + Simp[b^2*(Po
lyLog[3, 1 - 2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(2*e)), x]) /; Free
Q[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rule 4974

```
Int[((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_)*((d_) + (e_.*(x_))^(q_.), x_Sy
mbol] :> Simp[(d + e*x)^(q + 1)*((a + b*ArcTan[c*x])^p/(e*(q + 1))), x] - D
ist[b*c*(p/(e*(q + 1))), Int[ExpandIntegrand[(a + b*ArcTan[c*x])^(p - 1),
(d + e*x)^(q + 1)/(1 + c^2*x^2), x], x], x] /; FreeQ[{a, b, c, d, e}, x] &&
IGtQ[p, 1] && IntegerQ[q] && NeQ[q, -1]
```

Rule 5004

```
Int[((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.)/((d_) + (e_.*(x_))^2), x_Symb
ol] :> Simp[(a + b*ArcTan[c*x])^(p + 1)/(b*c*d*(p + 1)), x] /; FreeQ[{a, b,
c, d, e, p}, x] && EqQ[e, c^2*d] && NeQ[p, -1]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_.*(x_))^2),
x_Symbol] :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Di
st[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5104

```
Int[((((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.)*((f_) + (g_.*(x_))^(m_.)))/((
d_) + (e_.*(x_))^2), x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x])^p
/(d + e*x^2), (f + g*x)^m, x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && IGt
Q[p, 0] && EqQ[e, c^2*d] && IGtQ[m, 0]
```

Rule 5114

```
Int[(Log[u_]*((a_.) + ArcTan[(c_.*(x_)]*(b_.))^(p_.))/((d_) + (e_.*(x_))^2
), x_Symbol] :> Simp[(-I)*(a + b*ArcTan[c*x])^p*(PolyLog[2, 1 - u]/(2*c*d)),
x] + Dist[b*p*(I/2), Int[(a + b*ArcTan[c*x])^(p - 1)*(PolyLog[2, 1 - u]/(2*c*d)),
x]] /; FreeQ[{a, b, c, d, e, p}, x] && EqQ[u, 1 - e^2/d^2]
```

```
d + e*x^2)), x], x] /; FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[e, c^2*d] && EqQ[(1 - u)^2 - (1 - 2*(I/(I - c*x)))^2, 0]
```

Rule 6745

```
Int[(u_)*PolyLog[n_, v_], x_Symbol] :> With[{w = DerivativeDivides[v, u*v, x]}, Simpl[w*PolyLog[n + 1, v], x] /; !FalseQ[w]] /; FreeQ[n, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{(a + b \tan^{-1}(cx))^3}{(d + ex)^3} dx &= -\frac{(a + b \tan^{-1}(cx))^3}{2e(d + ex)^2} + \frac{(3bc) \int \left(\frac{e^2(a+b \tan^{-1}(cx))^2}{(c^2d^2+e^2)(d+ex)^2} + \frac{2c^2de^2(a+b \tan^{-1}(cx))^2}{(c^2d^2+e^2)^2(d+ex)} + \frac{(c^4d^2-c^2e^2)(a+b \tan^{-1}(cx))^2}{(c^2d^2+e^2)^3(d+ex)} \right) dx}{2e} \\
&= -\frac{(a + b \tan^{-1}(cx))^3}{2e(d + ex)^2} + \frac{(3bc) \int \frac{(c^4d^2-c^2e^2-2c^4dex)(a+b \tan^{-1}(cx))^2}{1+c^2x^2} dx}{2e(c^2d^2+e^2)^2} + \frac{(3bc^3de) \int \frac{(a+b \tan^{-1}(cx))^3}{(c^2d^2+e^2)^2} dx}{(c^2d^2+e^2)^3} \\
&= -\frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} - \frac{(a + b \tan^{-1}(cx))^3}{2e(d + ex)^2} - \frac{3bc^3d(a + b \tan^{-1}(cx))^2 \log(\frac{1-cx}{1+cx})}{(c^2d^2+e^2)^2} \\
&= -\frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} - \frac{(a + b \tan^{-1}(cx))^3}{2e(d + ex)^2} - \frac{3bc^3d(a + b \tan^{-1}(cx))^2 \log(\frac{1-cx}{1+cx})}{(c^2d^2+e^2)^2} \\
&= -\frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} + \frac{ic^3d(a + b \tan^{-1}(cx))^3}{(c^2d^2+e^2)^2} + \frac{c^2(cd-e)(cd+e)(a+b \tan^{-1}(cx))^2}{2e(c^2d^2+e^2)^2} \\
&= -\frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} + \frac{ic^3d(a + b \tan^{-1}(cx))^3}{(c^2d^2+e^2)^2} + \frac{c^2(cd-e)(cd+e)(a+b \tan^{-1}(cx))^2}{2e(c^2d^2+e^2)^2} \\
&= \frac{3bc^3d(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} + \frac{3ibc^2e(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} - \frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} + \\
&= \frac{3bc^3d(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} + \frac{3ibc^2e(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} - \frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} + \\
&= \frac{3bc^3d(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} + \frac{3ibc^2e(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} - \frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} + \\
&= \frac{3bc^3d(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} + \frac{3ibc^2e(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)^2} - \frac{3bc(a + b \tan^{-1}(cx))^2}{2(c^2d^2+e^2)(d + ex)} +
\end{aligned}$$

Mathematica [F]

time = 45.79, size = 0, normalized size = 0.00

$$\int \frac{(a + b \operatorname{ArcTan}(cx))^3}{(d + ex)^3} dx$$

Verification is not applicable to the result.

[In] `Integrate[(a + b*ArcTan[c*x])^3/(d + e*x)^3, x]`

[Out] `Integrate[(a + b*ArcTan[c*x])^3/(d + e*x)^3, x]`

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 14.87, size = 41275, normalized size = 44.10

method	result	size
derivativedivides	Expression too large to display	41275
default	Expression too large to display	41275

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x))^3/(e*x+d)^3, x, method=_RETURNVERBOSE)`

[Out] result too large to display

Maxima [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))^3/(e*x+d)^3, x, algorithm="maxima")`

[Out] Timed out

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))^3/(e*x+d)^3, x, algorithm="fricas")`

[Out] `integral((b^3*arctan(c*x)^3 + 3*a*b^2*arctan(c*x)^2 + 3*a^2*b*arctan(c*x) + a^3)/(x^3*e^3 + 3*d*x^2*e^2 + 3*d^2*x*e + d^3), x)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x))**3/(e*x+d)**3,x)`

[Out] Timed out

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x))^3/(e*x+d)^3,x, algorithm="giac")`

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(a + b \operatorname{atan}(c x))^3}{(d + e x)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x))^3/(d + e*x)^3,x)`

[Out] `int((a + b*atan(c*x))^3/(d + e*x)^3, x)`

$$\mathbf{3.21} \quad \int (d + ex)^2 (a + b \operatorname{ArcTan}(cx^2)) dx$$

Optimal. Leaf size=250

$$-\frac{2be^2x}{3c} - \frac{bd^3 \operatorname{ArcTan}(cx^2)}{3e} + \frac{(d+ex)^3 (a+b \operatorname{ArcTan}(cx^2))}{3e} + \frac{b(3cd^2-e^2) \operatorname{ArcTan}\left(1-\sqrt{2}\sqrt{c}x\right)}{3\sqrt{2}c^{3/2}} - \frac{b(3cd^2-}$$

[Out] $-2/3*b*e^2*x/c-1/3*b*d^3*\arctan(c*x^2)/e+1/3*(e*x+d)^3*(a+b*\arctan(c*x^2))/e-1/2*b*d^2*e*\ln(c^2*x^4+1)/c-1/6*b*(3*c*d^2-e^2)*\arctan(-1+x*2^(1/2)*c^(1/2))/c^(3/2)*2^(1/2)-1/6*b*(3*c*d^2-e^2)*\arctan(1+x*2^(1/2)*c^(1/2))/c^(3/2)*2^(1/2)-1/12*b*(3*c*d^2+e^2)*\ln(1+c*x^2-x*2^(1/2)*c^(1/2))/c^(3/2)*2^(1/2)+1/12*b*(3*c*d^2+e^2)*\ln(1+c*x^2+x*2^(1/2)*c^(1/2))/c^(3/2)*2^(1/2)$

Rubi [A]

time = 0.20, antiderivative size = 250, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 13, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.722$, Rules used = {4980, 1845, 1262, 649, 209, 266, 1294, 1182, 1176, 631, 210, 1179, 642}

$$\frac{(d+ex)^3 (a+b \operatorname{ArcTan}(cx^2))}{3e} + \frac{b \operatorname{ArcTan}\left(1-\sqrt{2}\sqrt{c}x\right) (3cd^2-e^2)}{3\sqrt{2}c^{3/2}} - \frac{b \operatorname{ArcTan}\left(\sqrt{2}\sqrt{c}x+1\right) (3cd^2-e^2)}{3\sqrt{2}c^{3/2}} - \frac{bd^3 \operatorname{ArcTan}(cx^2)}{3e} - \frac{b(3cd^2+e^2) \log(cx^2-\sqrt{2}\sqrt{c}x+1)}{6\sqrt{2}c^{3/2}} + \frac{b(3cd^2+e^2) \log(cx^2+\sqrt{2}\sqrt{c}x+1)}{6\sqrt{2}c^{3/2}} - \frac{bde \log(c^2x^4+1)}{2c} - \frac{2be^2x}{3c}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d+e*x)^2*(a+b*\operatorname{ArcTan}[c*x^2]), x]$

[Out] $(-2*b*e^2*x)/(3*c) - (b*d^3*\operatorname{ArcTan}[c*x^2])/(3*e) + ((d+e*x)^3*(a+b*\operatorname{ArcTan}[c*x^2]))/(3*e) + (b*(3*c*d^2-e^2)*\operatorname{ArcTan}[1-\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x])/(3*\operatorname{Sqrt}[2]*c^(3/2)) - (b*(3*c*d^2-e^2)*\operatorname{ArcTan}[1+\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x])/(3*\operatorname{Sqrt}[2]*c^(3/2)) - (b*(3*c*d^2+e^2)*\operatorname{Log}[1-\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x+c*x^2])/(6*\operatorname{Sqrt}[2]*c^(3/2)) + (b*(3*c*d^2+e^2)*\operatorname{Log}[1+\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x+c*x^2])/(6*\operatorname{Sqrt}[2]*c^(3/2)) - (b*d*e*\operatorname{Log}[1+c^2*x^4])/(2*c)$

Rule 209

$\operatorname{Int}[((a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \Rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{PosQ}[a/b] \& \operatorname{GtQ}[a, 0] \mid\mid \operatorname{GtQ}[b, 0])$

Rule 210

$\operatorname{Int}[((a_)+(b_)*(x_)^2)^{-1}, x_Symbol] \Rightarrow \operatorname{Simp}[(-(Rt[-a, 2]*Rt[-b, 2])^{(-1)})*\operatorname{ArcTan}[\operatorname{Rt}[-b, 2]*(x/Rt[-a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \& \operatorname{PosQ}[a/b] \& (\operatorname{LtQ}[a, 0] \mid\mid \operatorname{LtQ}[b, 0])$

Rule 266

$\operatorname{Int}[(x_)^{(m_)}/((a_)+(b_)*(x_)^{(n_)}), x_Symbol] \Rightarrow \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a+b*x^n, x]]/(b*n), x] /; \operatorname{FreeQ}[\{a, b, m, n\}, x] \& \operatorname{EqQ}[m, n-1]$

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> Simplify[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 649

```
Int[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 1182

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[a*c, 2]}, Dist[(d*q + a*e)/(2*a*c), Int[(q + c*x^2)/(a + c*x^4), x], x] + Dist[(d*q - a*e)/(2*a*c), Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]
```

Rule 1262

```
Int[(x_)*((d_) + (e_)*(x_)^2)^(q_)*((a_) + (c_)*(x_)^4)^(p_), x_Symbol] :> Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]
```

Rule 1294

```
Int[((f_)*(x_))^(m_)*(d_) + (e_)*(x_)^2)*((a_) + (c_)*(x_)^4)^(p_), x_
Symbol] :> Simp[e*f*(f*x)^(m - 1)*((a + c*x^4)^(p + 1)/(c*(m + 4*p + 3))), 
x] - Dist[f^2/(c*(m + 4*p + 3)), Int[(f*x)^(m - 2)*(a + c*x^4)^p*(a*e*(m - 
1) - c*d*(m + 4*p + 3)*x^2), x], x] /; FreeQ[{a, c, d, e, f, p}, x] && GtQ[ 
m, 1] && NeQ[m + 4*p + 3, 0] && IntegerQ[2*p] && (IntegerQ[p] || IntegerQ[m])]
```

Rule 1845

```
Int[((Pq_)*((c_)*(x_))^(m_.))/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> With[ 
{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]*x^(n/2) 
)/(c^ii*(a + b*x^n))), {ii, 0, n/2 - 1}]}, Int[v, x] /; SumQ[v]] /; FreeQ[{ 
a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n
```

Rule 4980

```
Int[((a_) + ArcTan[(c_)*(x_)^(n_)]*(b_.))*((d_) + (e_)*(x_)^(m_.)), x_Sy 
mbol] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcTan[c*x^n])/((e*(m + 1))), x] - D 
ist[b*c*(n/(e*(m + 1))), Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 + c^2*x^(2*n))) 
], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\int (d+ex)^2 (a+b\tan^{-1}(cx^2)) \, dx &= \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{b \int \frac{2cx(d+ex)^3}{1+c^2x^4} \, dx}{3e} \\
&= \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{(2bc) \int \frac{x(d+ex)^3}{1+c^2x^4} \, dx}{3e} \\
&= \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{(2bc) \int \left(\frac{x(d^3+3de^2x^2)}{1+c^2x^4} + \frac{x^2(3d^2e+e^3x^2)}{1+c^2x^4} \right) \, dx}{3e} \\
&= \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{(2bc) \int \frac{x(d^3+3de^2x^2)}{1+c^2x^4} \, dx}{3e} - \frac{(2bc) \int \frac{x^2(3d^2e+e^3x^2)}{1+c^2x^4} \, dx}{3e} \\
&= -\frac{2be^2x}{3c} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} + \frac{(2b) \int \frac{e^3-3c^2d^2ex^2}{1+c^2x^4} \, dx}{3ce} - \frac{(bc) \int \frac{e^3-3c^2d^2ex^2}{1+c^2x^4} \, dx}{3e} \\
&= -\frac{2be^2x}{3c} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{(bcd^3) \text{Subst}(\int \frac{1}{1+c^2x^2} \, dx, x, \sqrt{c^2+x^2})}{3e} \\
&= -\frac{2be^2x}{3c} - \frac{bd^3 \tan^{-1}(cx^2)}{3e} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{bde \log(1+\sqrt{c^2+x^2})}{2} \\
&= -\frac{2be^2x}{3c} - \frac{bd^3 \tan^{-1}(cx^2)}{3e} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} - \frac{b(3cd^2+3de^2)x}{3e} \\
&= -\frac{2be^2x}{3c} - \frac{bd^3 \tan^{-1}(cx^2)}{3e} + \frac{(d+ex)^3 (a+b\tan^{-1}(cx^2))}{3e} + \frac{b(3cd^2+3de^2)}{3e}
\end{aligned}$$

Mathematica [A]

time = 2.01, size = 252, normalized size = 1.01

$$\frac{1}{12} \left(12ad^2x - \frac{8be^2x}{c} + 12ade^2x^2 + 4ae^2x^3 + 4bx(3d^2+3dex+e^2x^2) \text{ArcTan}(cx^2) + \frac{2\sqrt{2}b(3cd^2-e^2) \text{ArcTan}(1-\sqrt{2}\sqrt{c}x)}{c^{3/2}} - \frac{2\sqrt{2}b(3cd^2-e^2) \text{ArcTan}(1+\sqrt{2}\sqrt{c}x)}{c^{3/2}} - \frac{\sqrt{2}b(3cd^2+e^2) \log(1-\sqrt{2}\sqrt{c}x+cx^2)}{c^{3/2}} + \frac{\sqrt{2}b(3cd^2+e^2) \log(1+\sqrt{2}\sqrt{c}x+cx^2)}{c^{3/2}} - \frac{6bde \log(1+c^2x^4)}{c} \right)$$

Antiderivative was successfully verified.

[In] Integrate[(d + e*x)^2*(a + b*ArcTan[c*x^2]), x]

[Out] $(12*a*d^2*x - (8*b*e^2*x)/c + 12*a*d*e*x^2 + 4*a*e^2*x^3 + 4*b*x*(3*d^2 + 3*d*e*x + e^2*x^2)*\text{ArcTan}[c*x^2] + (2*\text{Sqrt}[2]*b*(3*c*d^2 - e^2)*\text{ArcTan}[1 - Sqrt[2]*Sqrt[c]*x])/c^{(3/2)} - (2*\text{Sqrt}[2]*b*(3*c*d^2 - e^2)*\text{ArcTan}[1 + Sqrt[2]*Sqrt[c]*x])/c^{(3/2)} - (\text{Sqrt}[2]*b*(3*c*d^2 + e^2)*\text{Log}[1 - Sqrt[2]*Sqrt[c]*x + c*x^2])/c^{(3/2)} + (\text{Sqrt}[2]*b*(3*c*d^2 + e^2)*\text{Log}[1 + Sqrt[2]*Sqrt[c]*x + c*x^2])/c^{(3/2)} - (6*b*d*e*\text{Log}[1 + c^2*x^4])/c)/12$

Maple [A]

time = 0.44, size = 363, normalized size = 1.45

method	result
default	$\frac{(ex+d)^3 a}{3e} + \frac{b e^2 \arctan(cx^2) x^3}{3} + b e \arctan(cx^2) x^2 d + b \arctan(cx^2) x d^2 + \frac{b d^3 \arctan(cx^2)}{3e} - \frac{2 b e^2 x}{3c} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)^2*(a+b*arctan(c*x^2)),x,method=_RETURNVERBOSE)`

[Out] $1/3*(e*x+d)^3*a/e+1/3*b*e^2*arctan(c*x^2)*x^3+b*e*arctan(c*x^2)*x^2*d+b*arc\tan(c*x^2)*x*d^2+1/3*b*d^3*arctan(c*x^2)/e-2/3*b*e^2*x/c+1/12*b*e^2/c*(1/c^2)^{(1/4)*2^(1/2)*ln((x^2+(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2)))/(x^2-(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2)))+1/6*b*e^2/c*(1/c^2)^(1/4)*2^(1/2)*arctan(2^(1/2)/(1/c^2)^(1/4)*x+1)+1/6*b*e^2/c*(1/c^2)^(1/4)*2^(1/2)*arctan(2^(1/2)/(1/c^2)^(1/4)*x-1)-1/3*b/e*c*d^3/(c^2)^(1/2)*arctan(x^2*(c^2)^(1/2))-1/4*b/c*d^2/(1/c^2)^(1/4)*2^(1/2)*ln((x^2-(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2))/(x^2+(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2)))-1/2*b/c*d^2/(1/c^2)^(1/4)*2^(1/2)*arctan(2^(1/2)/(1/c^2)^(1/4)*x+1)-1/2*b/c*d^2/(1/c^2)^(1/4)*2^(1/2)*arctan(2^(1/2)/(1/c^2)^(1/4)*x-1)-1/2*b*d*e*ln(c^2*x^4+1)/c$

Maxima [A]

time = 0.48, size = 323, normalized size = 1.29

$$\frac{1}{3} ax^4 + adx^2 - \frac{1}{2} \left(\frac{2 \sqrt{c} \arctan\left(\frac{\sqrt{c}(1+x\sqrt{c})}{\sqrt{c}}\right)}{c^2} + \frac{2 \sqrt{c} \arctan\left(\frac{\sqrt{c}(1-x\sqrt{c})}{\sqrt{c}}\right)}{c^2} - \frac{\sqrt{c} \log\left(cx^2 + \sqrt{c}\sqrt{c}x + 1\right)}{c^2} + \frac{\sqrt{c} \log\left(cx^2 - \sqrt{c}\sqrt{c}x + 1\right)}{c^2} - 4x \arctan(cx^2) \right) bd^2 + ad^2 + \frac{1}{12} \left(\frac{8x}{c^2} - \frac{i\sqrt{c} \operatorname{atan}\left(\frac{\sqrt{c}(1+x\sqrt{c})}{\sqrt{c}}\right)}{c^2} + \frac{i\sqrt{c} \operatorname{atan}\left(\frac{\sqrt{c}(1-x\sqrt{c})}{\sqrt{c}}\right)}{c^2} + \frac{\sqrt{c} \ln\left(cx^2 + \sqrt{c}\sqrt{c}x + 1\right)}{c^2} - \frac{\sqrt{c} \ln\left(cx^2 - \sqrt{c}\sqrt{c}x + 1\right)}{c^2} \right) bc^2 + \frac{(2cx^3 \operatorname{atan}(cx^2) - \log(cx^2 + 1))bd}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x^2)),x, algorithm="maxima")`

[Out] $1/3*a*x^3*e^2 + a*d*x^2*e - 1/4*(c*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*c*x + s\qrt(2)*sqrt(c))/sqrt(c))/c^(3/2) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*c*x - sqrt(2)*sqrt(c))/sqrt(c))/c^(3/2) - sqrt(2)*log(c*x^2 + sqrt(2)*sqrt(c)*x + 1)/c^(3/2) + sqrt(2)*log(c*x^2 - sqrt(2)*sqrt(c)*x + 1)/c^(3/2))) - 4*x*arctan(c*x^2)*b*d^2 + a*d^2*x + 1/12*(4*x^3*arctan(c*x^2) - c*(8*x/c^2 - (2*sqrt(2)*arctan(1/2*sqrt(2)*(2*c*x + sqrt(2)*sqrt(c))/sqrt(c))/sqrt(c) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*c*x - sqrt(2)*sqrt(c))/sqrt(c))/sqrt(c) + sqrt(2)*log(c*x^2 + sqrt(2)*sqrt(c)*x + 1)/sqrt(c) - sqrt(2)*log(c*x^2 - sqrt(2)*sqrt(c)*x + 1)/sqrt(c))/c^2)*b*e^2 + 1/2*(2*c*x^2*arctan(c*x^2) - log(c^2*x^4 + 1))*b*d*e/c$

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 4652 vs. 2(196) = 392.

time = 1.43, size = 4652, normalized size = 18.61

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^2*(a+b*arctan(c*x^2)),x, algorithm="fricas")
[Out] 1/12*(972*a*b^4*c^5*d^9*x^2*e + 972*a*b^4*c^5*d^10*x + 216*a*b^4*c^3*d^5*x^2*e^5 + 216*a*b^4*c^3*d^6*x*e^4 - 4*sqrt(2)*c^7*sqrt((81*b^2*c^4*d^8 + 18*b^2*c^2*d^4*e^4 - 6*c^4*d^2*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*e^2 + b^2*e^8)/(81*b^2*c^4*d^8 - 18*b^2*c^2*d^4*e^4 + b^2*e^8))*(81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)^(3/4)*sqrt((81*b^4*c^4*d^8 - 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*arctan(sqrt(2)*sqrt(6561*b^6*c^8*d^16*x^2 - 162*b^6*c^4*d^8*x^2*e^8 + b^6*x^2*e^16 + sqrt(2)*(729*b^5*c^7*d^12*x*e^2 - 81*b^5*c^5*d^8*x*e^6 - 9*b^5*c^3*d^4*x*e^10 + b^5*c*x*e^14 + 3*(81*b^3*c^9*d^10*x - 18*b^3*c^7*d^6*x*e^4 + b^3*c^5*d^2*x*e^8)*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6))*sqrt((81*b^2*c^4*d^8 + 18*b^2*c^2*d^4*e^4 - 6*c^4*d^2*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*e^2 + b^2*e^8)/(81*b^2*c^4*d^8 - 18*b^2*c^2*d^4*e^4 + b^2*e^8)))*((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)^(1/4) + (729*b^4*c^8*d^12 - 81*b^4*c^6*d^8*e^4 - 9*b^4*c^4*d^4*e^8 + b^4*c^2*e^12)*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6))*c^11*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*sqrt((81*b^4*c^4*d^8 - 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*e^2 + 3*(9*b^2*c^11*d^6 + b^2*c^9*d^2*e^4)*sqrt((81*b^4*c^4*d^8 - 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6))*sqrt((81*b^2*c^4*d^8 + 18*b^2*c^2*d^4*e^4 - 6*c^4*d^2*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*e^2 + b^2*e^8)/(81*b^2*c^4*d^8 - 18*b^2*c^2*d^4*e^4 + b^2*e^8)))*((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)^(3/4) + sqrt(2)*((81*b^3*c^15*d^8*x^2 - b^3*c^11*x^10)*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6) + 3*(729*b^5*c^15*d^14*x + 81*b^5*c^13*d^10*x*e^4 - 9*b^5*c^11*d^6*x*e^8 - b^5*c^9*d^2*x*e^12)*sqrt((81*b^4*c^4*d^8 - 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6))*sqrt((81*b^2*c^4*d^8 + 18*b^2*c^2*d^4*e^4 - 6*c^4*d^2*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*e^2 + b^2*e^8)/(81*b^2*c^4*d^8 - 18*b^2*c^2*d^4*e^4 + b^2*e^8))*((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)^(3/4) + (6561*b^6*c^14*d^16 + 1458*b^6*c^12*d^12*e^4 - 18*b^6*c^8*d^4*e^12 - b^6*c^6*e^16)*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*sqrt((81*b^4*c^4*d^8 - 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6))/(531441*b^10*c^12*d^24 + 118098*b^10*c^10*d^20*e^4 - 6561*b^10*c^8*d^16*e^8 - 2916*b^10*c^6*d^12*e^12 - 81*b^10*c^4*d^8*e^16 + 18*b^10*c^2*d^4*e^20 + b^10*e^24)) - 4*sqrt(2)*c^7*sqrt((81*b^2*c^4*d^8 + 18*b^2*c^2*d^4*e^4 - 6*c^4*d^2*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*e^2 + b^2*e^8)/(81*b^2*c^4*d^8 - 18*b^2*c^2*d^4*e^4 + b^2*e^8))*((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)*arctan(sqrt(2)*sqrt(6561*b^6*c^8*d^16*x^2 - 162*b^6*c^4*d^8*x^2 - 81*b^5*c^5*d^8*x^6 - 9*b^5*c^3*d^4*x^10 + b^5*c*x^14 + 3*(81*b^3*c^9*d^10*x - 18*b^3*c^7*d^6*x^4 + b^3*c^5*d^2*x^8)*sqrt((81*b^4*c^4*d^8 + 18*b^4*c^2*d^4*e^4 + b^4*e^8)/c^6)))
```

$$\begin{aligned}
& + b^{4}e^{8})/c^6)) * \sqrt{((81*b^{2}c^{4}d^{8} + 18*b^{2}c^{2}d^{4}e^{4} - 6*c^{4}d^{2}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * e^{2} + b^{2}e^{8})/(81*b^{2}c^{4}d^{8} - 18*b^{2}c^{2}d^{4}e^{4} + b^{2}e^{8})) * ((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)^{(1/4)} + (729*b^{4}c^{8}d^{12} - 81*b^{4}c^{6}d^{8}e^{4} - 9*b^{4}c^{4}d^{4}e^{8} + b^{4}c^{2}e^{12}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * (c^{11} * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * \sqrt{((81*b^{4}c^{4}d^{8} - 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * e^{2} + 3 * (9*b^{2} * c^{11}d^{6} + b^{2}c^{9}d^{2}e^{4}) * \sqrt{((81*b^{4}c^{4}d^{8} - 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * \sqrt{((81*b^{2}c^{4}d^{8} + 18*b^{2}c^{2}d^{4}e^{4} - 6*c^{4}d^{2}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * e^{2} + b^{2}e^{8})/(81*b^{2}c^{4}d^{8} - 18*b^{2}c^{2}d^{4}e^{4} + b^{2}e^{8})) * ((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)^{(3/4)} + \sqrt{2} * ((81*b^{3}c^{15}d^{8}x^{2} - b^{3}c^{11}x^{10}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * \sqrt{((81*b^{4}c^{4}d^{8} - 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * \sqrt{((81*b^{5}c^{15}d^{14}x + 81*b^{5}c^{13}d^{10}x^{4} - 9*b^{5}c^{11}d^{6}x^{8} - b^{5}c^{9}d^{2}x^{12}) * \sqrt{((81*b^{4}c^{4}d^{8} - 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * \sqrt{((81*b^{2}c^{4}d^{8} + 18*b^{2}c^{2}d^{4}e^{4} - 6*c^{4}d^{2}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * e^{2} + b^{2}e^{8})/(81*b^{2}c^{4}d^{8} - 18*b^{2}c^{2}d^{4}e^{4} + b^{2}e^{8})) * ((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)^{(3/4)} - (6561*b^{6}c^{14}d^{16} + 1458*b^{6}c^{12}d^{12}e^{4} - 18*b^{6}c^{8}d^{4}e^{12} - b^{6}c^{6}e^{16}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} - 6*c^{4}d^{2}) * \sqrt{((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * \sqrt{((81*b^{4}c^{4}d^{8} - 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)} * e^{2} + b^{2}e^{8})/(81*b^{2}c^{4}d^{8} - 18*b^{2}c^{2}d^{4}e^{4} + b^{2}e^{8})) * ((81*b^{4}c^{4}d^{8} + 18*b^{4}c^{2}d^{4}e^{4} + b^{4}e^{8})/c^6)^{(3/4)} - (531441*b^{10}c^{12}d^{24} + 118098*b^{10}c^{1}0*d^{20}e^{4} - 6561*b^{10}c^{8}d^{16}e^{8} - 2916*b^{10}c^{6}d^{12}e^{12} - 81*b^{10}c^{4}*d^{8}e^{16} + 18*b^{10}c^{2}d^{4}e^{20} + b^{10}e^{24}) + 12*a*b^{4}c*d*x^{2}e^{9} + 12*a*b^{4}c*d^{2}x^{8} + 4*(81*b^{5}c^{5}d^{8}x^{3}e^{2} + 243*b^{5}c^{5}d^{9}x^{2}e^{6} + 243*b^{5}c^{5}d^{10}x + 18*b^{5}c^{3}d^{4}x^{3}e^{6} + 54*b...} \\
\end{aligned}$$

Sympy [A]

time = 11.15, size = 403, normalized size = 1.61

$$\begin{cases}
ad^2x + adex^2 + \frac{ae^2x^3}{3} + bd^2x \operatorname{atan}(cx^2) + bde^2x \operatorname{atan}(cx^2) + \frac{be^2x^3 \operatorname{atan}(cx^2)}{3} - \frac{bd^2 \log\left(x - \sqrt{\frac{-b}{2c}}\right)}{c\sqrt{-\frac{b}{2c}}} + \frac{bd^2 \log\left(x^2 + \sqrt{\frac{-b}{2c}}\right)}{c\sqrt{-\frac{b}{2c}}} - \frac{bd^2 \operatorname{atan}\left(\sqrt{\frac{-b}{2c}}\right)}{c\sqrt{-\frac{b}{2c}}} - \frac{bde \log\left(x^2 + \sqrt{\frac{-b}{2c}}\right)}{c} - \frac{2be^2x}{3c} - \frac{bd^2 \operatorname{atan}(cx^2)}{c^2(-\frac{b}{2c})^{\frac{3}{2}}} + \frac{bd^2 \operatorname{atan}(cx^2)}{c^2\sqrt{-\frac{b}{2c}}} - \frac{be^2 \operatorname{atan}(cx^2)}{3c^2\sqrt{-\frac{b}{2c}}} + \frac{be^2 \log\left(x - \sqrt{\frac{-b}{2c}}\right)}{6c^2(-\frac{b}{2c})^{\frac{3}{2}}} - \frac{be^2 \log\left(x^2 + \sqrt{\frac{-b}{2c}}\right)}{3c^2(-\frac{b}{2c})^{\frac{3}{2}}} & \text{for } c \neq 0 \\
a(d^2x + dex^2 + \frac{e^2x^3}{3}) & \text{otherwise}
\end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)**2*(a+b*atan(c*x**2)),x)
[Out] Piecewise((a*d**2*x + a*d*e*x**2 + a*e**2*x**3/3 + b*d**2*x*atan(c*x**2) +
b*d*e*x**2*atan(c*x**2) + b*e**2*x**3*atan(c*x**2)/3 - b*d**2*log(x - (-1/c)**2)**(1/4)/(c*(-1/c**2)**(1/4)) + b*d**2*log(x**2 + sqrt(-1/c**2))/(2*c*(-1/c**2)**(1/4)) - b*d**2*atan(x/(-1/c**2)**(1/4))/(c*(-1/c**2)**(1/4)) - b*d*e*log(x**2 + sqrt(-1/c**2))/c - 2*b*e**2*x/(3*c) - b*d**2*atan(c*x**2)/(c**2*(-1/c**2)**(3/4)) + b*d*e*atan(c*x**2)/(c**2*sqrt(-1/c**2)) - b*e**2*a*atan(c*x**2)/(3*c**2*(-1/c**2)**(1/4)) + b*e**2*log(x - (-1/c**2)**(1/4))/(3*c**3*(-1/c**2)**(3/4)) - b*e**2*log(x**2 + sqrt(-1/c**2))/(6*c**3*(-1/c**2)**(3/4)) - b*e**2*atan(x/(-1/c**2)**(1/4))/(3*c**3*(-1/c**2)**(3/4)), Ne(c, 0)), (a*(d**2*x + d*e*x**2 + e**2*x**3/3), True))
```

Giac [A]

time = 1.15, size = 308, normalized size = 1.23

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)^2*(a+b*a*arctan(c*x^2)),x, algorithm="giac")
[Out] -1/2*b*d*e*log(c^2*x^4 + 1)/c + 1/3*(b*c*e^2*x^3*arctan(c*x^2) + a*c*e^2*x^3 + 3*b*c*d*e*x^2*arctan(c*x^2) + 3*a*c*d*e*x^2 + 3*b*c*d^2*x*arctan(c*x^2) + 3*a*c*d^2*x - 2*b*e^2*x)/c - 1/12*sqrt(2)*(3*b*c^2*d^2 + b*e^2*abs(c))*log(x^2 - sqrt(2)*x/sqrt(abs(c)) + 1/abs(c))/(c*abs(c)^(3/2)) - 1/6*sqrt(2)*(3*b*c^2*d^2*sqrt(abs(c)) - b*e^2*abs(c)^(3/2))*arctan(1/2*sqrt(2)*(2*x + sqrt(2)/sqrt(abs(c)))*sqrt(abs(c)))/c^3 - 1/6*sqrt(2)*(3*b*c^2*d^2*sqrt(abs(c)) - b*e^2*abs(c)^(3/2))*arctan(1/2*sqrt(2)*(2*x - sqrt(2)/sqrt(abs(c)))*sqrt(abs(c)))/c^3 + 1/12*sqrt(2)*(3*b*c^2*d^2*sqrt(abs(c)) + b*e^2*abs(c)^(3/2))*log(x^2 + sqrt(2)*x/sqrt(abs(c)) + 1/abs(c))/c^3
```

Mupad [B]

time = 3.37, size = 419, normalized size = 1.68

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int ((a + b \cdot \arctan(c \cdot x^2)) \cdot (d + e \cdot x)^2) dx$

```
[Out] (a*e^2*x^3)/3 + a*d^2*x + (b*e^2*x^3*atan(c*x^2))/3 + a*d*e*x^2 - (3*b*d^2*log(3*c*x*(1i/(9*c)))^(1/2) - 1)*(1i/(9*c))^(1/2))/2 + (3*b*d^2*log(3*c*x*(1i/(9*c)))^(1/2) + 1)*(1i/(9*c))^(1/2))/2 - (b*d^2*log(3*c*x*(1i/(9*c)))^(1/2) + 1i)*(1i/(9*c))^(1/2)*3i)/2 + (b*d^2*log(c*x*(1i/(9*c)))^(1/2)*3i + 1)*(1i/(9*c))^(1/2)*3i)/2 - (2*b*e^2*x)/(3*c) + b*d^2*x*atan(c*x^2) + (b*e^2*log(3*c*x*(1i/(9*c)))^(1/2) - 1)*(1i/(9*c))^(1/2)*1i)/(2*c) - (b*e^2*log(3*c*x*(1i/(9*c)))^(1/2) + 1)*(1i/(9*c))^(1/2)*1i)/(2*c) + (b*e^2*log(3*c*x*(1i/(9*c)))^(1/2) + 1i)*(1i/(9*c))^(1/2))/(2*c) - (b*e^2*log(c*x*(1i/(9*c)))^(1/2)*3i + 1)*(1i/(9*c))^(1/2))/(2*c) + b*d*e*x^2*atan(c*x^2) - (b*d*e*log(3*c*x*(1i/(9*c)))^(1/2) - 1))/(2*c) - (b*d*e*log(3*c*x*(1i/(9*c)))^(1/2) + 1))/(2*c) - (b*d*e*log(3*c*x*(1i/(9*c)))^(1/2) + 1i))/(2*c) - (b*d*e*log(c*x*(1i/(9*c)))^(1/2)*3i + 1))/(2*c)
```

$$\mathbf{3.22} \quad \int (d + ex) (a + b \operatorname{ArcTan}(cx^2)) \, dx$$

Optimal. Leaf size=192

$$-\frac{bd^2 \operatorname{ArcTan}(cx^2)}{2e} + \frac{(d + ex)^2 (a + b \operatorname{ArcTan}(cx^2))}{2e} + \frac{bd \operatorname{ArcTan}\left(1 - \sqrt{2} \sqrt{c} x\right)}{\sqrt{2} \sqrt{c}} - \frac{bd \operatorname{ArcTan}\left(1 + \sqrt{2} \sqrt{c} x\right)}{\sqrt{2} \sqrt{c}}$$

[Out] $-1/2*b*d^2*arctan(c*x^2)/e + 1/2*(e*x+d)^2*(a+b*arctan(c*x^2))/e - 1/4*b*e*ln(c^2*x^4+1)/c - 1/2*b*d*arctan(-1+x*2^(1/2)*c^(1/2))*2^(1/2)/c^(1/2) - 1/2*b*d*arctan(1+x*2^(1/2)*c^(1/2))*2^(1/2)/c^(1/2) - 1/4*b*d*ln(1+c*x^2-x*2^(1/2)*c^(1/2))*2^(1/2)/c^(1/2) + 1/4*b*d*ln(1+c*x^2+x*2^(1/2)*c^(1/2))*2^(1/2)/c^(1/2)$

Rubi [A]

time = 0.13, antiderivative size = 192, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {4980, 1845, 303, 1176, 631, 210, 1179, 642, 1262, 649, 209, 266}

$$\frac{(d + ex)^2 (a + b \operatorname{ArcTan}(cx^2))}{2e} - \frac{bd^2 \operatorname{ArcTan}(cx^2)}{2e} + \frac{bd \operatorname{ArcTan}\left(1 - \sqrt{2} \sqrt{c} x\right)}{\sqrt{2} \sqrt{c}} - \frac{bd \operatorname{ArcTan}\left(\sqrt{2} \sqrt{c} x + 1\right)}{\sqrt{2} \sqrt{c}} - \frac{be \log(c^2 x^4 + 1)}{4c} - \frac{bd \log(cx^2 - \sqrt{2} \sqrt{c} x + 1)}{2\sqrt{2} \sqrt{c}} + \frac{bd \log(cx^2 + \sqrt{2} \sqrt{c} x + 1)}{2\sqrt{2} \sqrt{c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)*(a + b*\operatorname{ArcTan}[c*x^2]), x]$

[Out] $-1/2*(b*d^2*\operatorname{ArcTan}[c*x^2])/e + ((d + e*x)^2*(a + b*\operatorname{ArcTan}[c*x^2]))/(2*e) + (b*d*\operatorname{ArcTan}[1 - \operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x])/(\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]) - (b*d*\operatorname{ArcTan}[1 + \operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x])/(\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]) - (b*d*\operatorname{Log}[1 - \operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x + c*x^2])/(2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]) + (b*d*\operatorname{Log}[1 + \operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]*x + c*x^2])/(2*\operatorname{Sqrt}[2]*\operatorname{Sqrt}[c]) - (b*e*\operatorname{Log}[1 + c^2*x^4])/(4*c)$

Rule 209

$\operatorname{Int}[(a_1 + b_1)*(x_1^2)^{-1}, x_1] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a_1, 2]*\operatorname{Rt}[b_1, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b_1, 2]*(x/\operatorname{Rt}[a_1, 2])], x_1] /; \operatorname{FreeQ}[\{a_1, b_1\}, x_1] \& \operatorname{PosQ}[a/b] \&& (\operatorname{GtQ}[a_1, 0] \|\| \operatorname{GtQ}[b_1, 0])$

Rule 210

$\operatorname{Int}[(a_1 + b_1)*(x_1^2)^{-1}, x_1] \rightarrow \operatorname{Simp}[(-(Rt[-a_1, 2]*Rt[-b_1, 2])^{(-1)})*\operatorname{ArcTan}[\operatorname{Rt}[-b_1, 2]*(x/Rt[-a_1, 2])], x_1] /; \operatorname{FreeQ}[\{a_1, b_1\}, x_1] \& \operatorname{PosQ}[a/b] \& (\operatorname{LtQ}[a_1, 0] \|\| \operatorname{LtQ}[b_1, 0])$

Rule 266

$\operatorname{Int}[(x_1^{m_1})/((a_1 + b_1)*(x_1^{n_1})), x_1] \rightarrow \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a_1 + b_1*x_1^{n_1}, x_1]]/(b*n_1), x_1] /; \operatorname{FreeQ}[\{a_1, b_1, m_1, n_1\}, x_1] \& \operatorname{EqQ}[m_1, n_1 - 1]$

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] :> With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> SImp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 649

```
Int[((d_) + (e_.)*(x_))/((a_) + (c_.)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 1262

```
Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_)*((a_) + (c_.)*(x_)^4)^(p_), x_Symbol] :> Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ[{a, c, d, e, p, q}, x]
```

Rule 1845

```
Int[((Pq_)*((c_ .)*(x_ ))^(m_ .))/((a_ ) + (b_ .)*(x_ )^(n_ )), x_Symbol] :> With[
{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]*x^(n/2))
/(c^ii*(a + b*x^n))), {ii, 0, n/2 - 1}]}, Int[v, x] /; SumQ[v] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n
```

Rule 4980

```
Int[((a_.) + ArcTan[(c_ .)*(x_ )^(n_ )]*(b_ .))*((d_ ) + (e_ .)*(x_ ))^(m_ .), x_Symbol] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcTan[c*x^n])/((e*(m + 1))), x] - Dist[b*c*(n/(e*(m + 1))), Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\int (d + ex) (a + b \tan^{-1}(cx^2)) \, dx &= \int (a(d + ex) + b(d + ex) \tan^{-1}(cx^2)) \, dx \\
&= \frac{a(d + ex)^2}{2e} + b \int (d + ex) \tan^{-1}(cx^2) \, dx \\
&= \frac{a(d + ex)^2}{2e} + b \int (d \tan^{-1}(cx^2) + ex \tan^{-1}(cx^2)) \, dx \\
&= \frac{a(d + ex)^2}{2e} + (bd) \int \tan^{-1}(cx^2) \, dx + (be) \int x \tan^{-1}(cx^2) \, dx \\
&= \frac{a(d + ex)^2}{2e} + bdx \tan^{-1}(cx^2) + \frac{1}{2}bex^2 \tan^{-1}(cx^2) - (2bcd) \int \frac{x^2}{1 + c^2x^4} \, dx \\
&= \frac{a(d + ex)^2}{2e} + bdx \tan^{-1}(cx^2) + \frac{1}{2}bex^2 \tan^{-1}(cx^2) - \frac{be \log(1 + c^2x^4)}{4c} + \\
&\quad \frac{a(d + ex)^2}{2e} + bdx \tan^{-1}(cx^2) + \frac{1}{2}bex^2 \tan^{-1}(cx^2) - \frac{be \log(1 + c^2x^4)}{4c} - \\
&= \frac{a(d + ex)^2}{2e} + bdx \tan^{-1}(cx^2) + \frac{1}{2}bex^2 \tan^{-1}(cx^2) - \frac{bd \log(1 - \sqrt{2} \sqrt{c} x)}{2\sqrt{2} \sqrt{c}} + \\
&= \frac{a(d + ex)^2}{2e} + bdx \tan^{-1}(cx^2) + \frac{1}{2}bex^2 \tan^{-1}(cx^2) + \frac{bd \tan^{-1}(1 - \sqrt{2} \sqrt{c} x)}{\sqrt{2} \sqrt{c}}
\end{aligned}$$

Mathematica [A]

time = 0.07, size = 153, normalized size = 0.80

$$adx + \frac{1}{2}aex^2 + bdx \text{ArcTan}(cx^2) + \frac{1}{2}bex^2 \text{ArcTan}(cx^2) - \frac{bd(-2 \text{ArcTan}(1 - \sqrt{2} \sqrt{c} x) + 2 \text{ArcTan}(1 + \sqrt{2} \sqrt{c} x) + \log(1 - \sqrt{2} \sqrt{c} x + cx^2) - \log(1 + \sqrt{2} \sqrt{c} x + cx^2))}{2\sqrt{2} \sqrt{c}} - \frac{be \log(1 + c^2x^4)}{4c}$$

Antiderivative was successfully verified.

[In] `Integrate[(d + e*x)*(a + b*ArcTan[c*x^2]), x]`

[Out] $a*d*x + (a*e*x^2)/2 + b*d*x*\text{ArcTan}[c*x^2] + (b*e*x^2*\text{ArcTan}[c*x^2])/2 - (b*d*(-2*\text{ArcTan}[1 - \sqrt{2}*\sqrt{c}*x] + 2*\text{ArcTan}[1 + \sqrt{2}*\sqrt{c}*x] + \log[1 - \sqrt{2}*\sqrt{c}*x + c*x^2] - \log[1 + \sqrt{2}*\sqrt{c}*x + c*x^2]))/(2*\sqrt{2}*\sqrt{c}) - (b*e*\log[1 + c^2*x^4])/(4*c)$

Maple [A]

time = 0.09, size = 168, normalized size = 0.88

method	result
default	$a\left(\frac{1}{2}e x^2 + dx\right) + \frac{b \arctan(cx^2)x^2 e}{2} + b \arctan(cx^2) dx - \frac{bd\sqrt{2} \ln\left(\frac{x^2 - \left(\frac{1}{c^2}\right)^{\frac{1}{4}}x\sqrt{2} + \sqrt{\frac{1}{c^2}}}{x^2 + \left(\frac{1}{c^2}\right)^{\frac{1}{4}}x\sqrt{2} + \sqrt{\frac{1}{c^2}}}\right)}{4c\left(\frac{1}{c^2}\right)^{\frac{1}{4}}} - \frac{bd\sqrt{2} \arctan\left(\frac{x\sqrt{2}}{\sqrt{\frac{1}{c^2}}}\right)}{2}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((e*x+d)*(a+b*arctan(c*x^2)),x,method=_RETURNVERBOSE)
```

[Out] $a*(1/2*e*x^2+d*x)+1/2*b*\arctan(c*x^2)*x^2+e+b*\arctan(c*x^2)*d*x-1/4*b*d/c/(1/c^2)^{(1/4)}*2^{(1/2)}*\ln((x^2-(1/c^2)^{(1/4)}*x*2^{(1/2)}+(1/c^2)^{(1/2)})/(x^2+(1/c^2)^{(1/4)}*x*2^{(1/2)}+(1/c^2)^{(1/2)}))-1/2*b*d/c/(1/c^2)^{(1/4)}*2^{(1/2)}*\arctan(2^{(1/2)}/(1/c^2)^{(1/4)}*x+1)-1/2*b*d/c/(1/c^2)^{(1/4)}*2^{(1/2)}*\arctan(2^{(1/2)}/(1/c^2)^{(1/4)}*x-1)-1/4*b*e*\ln(c^2*x^4+1)/c$

Maxima [A]

time = 0.46, size = 170, normalized size = 0.89

$$\frac{1}{2}ax^2e - \frac{1}{4}\left(c\left(\frac{2\sqrt{2}\arctan\left(\frac{\sqrt{2}(2cx+\sqrt{2}\sqrt{c})}{2\sqrt{c}}\right)}{c^{\frac{3}{2}}} + \frac{2\sqrt{2}\arctan\left(\frac{\sqrt{2}(2cx-\sqrt{2}\sqrt{c})}{2\sqrt{c}}\right)}{c^{\frac{3}{2}}} - \frac{\sqrt{2}\log(cx^2+\sqrt{2}\sqrt{c}x+1)}{c^{\frac{3}{2}}} + \frac{\sqrt{2}\log(cx^2-\sqrt{2}\sqrt{c}x+1)}{c^{\frac{3}{2}}}\right) - 4x\arctan(cx^2)\right)bd + adx + \frac{(2cx^2\arctan(cx^2) - \log(c^2x^4+1))be}{4c}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)*(a+b*arctan(c*x^2)),x, algorithm="maxima")
```

```
[Out] 1/2*a*x^2*e - 1/4*(c*(2*sqrt(2)*arctan(1/2*sqrt(2)*(2*c*x + sqrt(2)*sqrt(c)))/sqrt(c))/c^(3/2) + 2*sqrt(2)*arctan(1/2*sqrt(2)*(2*c*x - sqrt(2)*sqrt(c))/sqrt(c))/c^(3/2) - sqrt(2)*log(c*x^2 + sqrt(2)*sqrt(c)*x + 1)/c^(3/2) + sqrt(2)*log(c*x^2 - sqrt(2)*sqrt(c)*x + 1)/c^(3/2)) - 4*x*arctan(c*x^2))*b*d + a*d*x + 1/4*(2*c*x^2*arctan(c*x^2) - log(c^2*x^4 + 1))*b*e/c
```

Fricas [B] Leaf count of result is larger than twice the leaf count of optimal. 538 vs. $2(150) = 300$.

time = 1.05, size = 538, normalized size = 2.80

$$2ab^2\partial_x^2v^2+4ab^2\partial_x^2v+4\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2b^2v^2\arctan\left(\frac{\sqrt{-2}\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2v^2-\sqrt{2}\left(\frac{\partial_x v^2+v^2}{v^2}\right)^2b^2v^2+\sqrt{\frac{b^2v^2}{v^2}}\left(\frac{\partial_x v}{v^2}\right)^2}{v^2}\right)+4\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2b^2v^2\arctan\left(\frac{\sqrt{-2}\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2v^2-\sqrt{2}\left(\frac{\partial_x v^2+v^2}{v^2}\right)^2b^2v^2+\sqrt{\frac{b^2v^2}{v^2}}\left(\frac{\partial_x v}{v^2}\right)^2}{v^2}\right)+2\left(b^2\partial_x^2v+2b^2\partial_x^2v\right)\arctan\left(\frac{\partial_x v}{v^2}\right)-\left(b^2\partial_x^2v+\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2b^2v^2\right)\log\left(\left(b^2\partial_x^2v+\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2b^2v^2\right)\sqrt{b^2\partial_x^2v^2+\sqrt{\frac{b^2v^2}{v^2}}b^2v^2}\right)-\left(b^2\partial_x^2v+\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2b^2v^2\right)\log\left(\left(b^2\partial_x^2v+\sqrt{2}\left(\frac{\partial_x v}{v^2}\right)^2b^2v^2\right)\sqrt{b^2\partial_x^2v^2+\sqrt{\frac{b^2v^2}{v^2}}b^2v^2}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)*(a+b*arctan(c*x^2)),x, algorithm="fricas")
[Out] 1/4*(2*a*b^4*c*d^4*x^2*e + 4*a*b^4*c*d^5*x + 4*sqrt(2)*(b^4*d^4/c^2)^(1/4)*
b^4*c*d^4*arctan(-(b^8*d^8 + sqrt(2)*(b^4*d^4/c^2)^(5/4)*b^3*c^3*d^3*x - sq
rt(2)*sqrt(b^6*d^6*x^2 + sqrt(2)*(b^4*d^4/c^2)^(3/4)*b^3*c*d^3*x + sqrt(b^4
*d^4/c^2)*b^4*d^4)*(b^4*d^4/c^2)^(5/4)*c^3)/(b^8*d^8)) + 4*sqrt(2)*(b^4*d^4
/c^2)^(1/4)*b^4*c*d^4*arctan((b^8*d^8 - sqrt(2)*(b^4*d^4/c^2)^(5/4)*b^3*c^3
*d^3*x + sqrt(2)*sqrt(b^6*d^6*x^2 - sqrt(2)*(b^4*d^4/c^2)^(3/4)*b^3*c*d^3*x
+ sqrt(b^4*d^4/c^2)*b^4*d^4)*(b^4*d^4/c^2)^(5/4)*c^3)/(b^8*d^8)) + 2*(b^5*
c*d^4*x^2*e + 2*b^5*c*d^5*x)*arctan(c*x^2) - (b^5*d^4*e - sqrt(2)*(b^4*d^4/c^2)
^(1/4)*b^4*c*d^4)*log(b^6*d^6*x^2 + sqrt(2)*(b^4*d^4/c^2)^(3/4)*b^3*c*d^3*x
+ sqrt(b^4*d^4/c^2)*b^4*d^4) - (b^5*d^4*e + sqrt(2)*(b^4*d^4/c^2)^(1/4)
)*b^4*c*d^4)*log(b^6*d^6*x^2 - sqrt(2)*(b^4*d^4/c^2)^(3/4)*b^3*c*d^3*x + sq
rt(b^4*d^4/c^2)*b^4*d^4))/(b^4*c*d^4)
```

Sympy [C] Result contains complex when optimal does not.

time = 7.52, size = 1266, normalized size = 6.59

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((e*x+d)*(a+b*atan(c*x**2)),x)
```

```
[Out] Piecewise(((a - oo*I*b)*(d*x + e*x**2/2), Eq(c, -I/x**2)), ((a + oo*I*b)*(d
*x + e*x**2/2), Eq(c, I/x**2)), (a*(d*x + e*x**2/2), Eq(c, 0)), (2*a*c**5*d
*x**5*(-1/c**2)**(11/4)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4))
+ a*c**5*e*x**6*(-1/c**2)**(11/4)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ 2*a*c**3*d*x*(-1/c**2)**(11/4)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ a*c**3*e*x**2*(-1/c**2)**(11/4)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ 2*c**3*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4) + a*c**3*e*x**2*(-1/c**2)**(11/4)
/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4) + 2*b*c**5*d*x
**5*(-1/c**2)**(11/4)*atan(c*x**2)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ b*c**5*e*x**6*(-1/c**2)**(11/4)*atan(c*x**2)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ 2*c**4*(-1/c**2)**(11/4)*log(x - (-1/c**2)**(1/4))/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ b*c**4*d*x**4*(-1/c**2)**(5/2)*log(x**2 + sqrt(-1/c**2))/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ 2*b*c**3*d*x*(-1/c**2)**(11/4)*atan(c*x**2)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
+ 2*c**3*(-1/c**2)**(11/4) + b*c**3*e*x**2*(-1/c**2)**(11/4)*atan(c*x**2)/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4)
- 2*b*c**2*d*(-1/c**2)**(5/2)*log(x - (-1/c**2)**(1/4))/(2*c**5*x**4*(-1/c**2)**(11/4) +
```

```

2*c**3*(-1/c**2)**(11/4)) + b*c**2*d*(-1/c**2)**(5/2)*log(x**2 + sqrt(-1/c
**2))/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4) - 2*b*c**2
*d*(-1/c**2)**(5/2)*atan(x/(-1/c**2)**(1/4))/(2*c**5*x**4*(-1/c**2)**(11/4)
+ 2*c**3*(-1/c**2)**(11/4) - b*c**2*e*(-1/c**2)**(11/4)*log(x**2 + sqrt(-
1/c**2))/(2*c**5*x**4*(-1/c**2)**(11/4) + 2*c**3*(-1/c**2)**(11/4) - 2*b*d
*x**4*atan(c*x**2)/(2*c**6*x**4*(-1/c**2)**(11/4) + 2*c**4*(-1/c**2)**(11/4
)) - 2*b*d*atan(c*x**2)/(2*c**8*x**4*(-1/c**2)**(11/4) + 2*c**6*(-1/c**2)**(1
1/4)) + b*e*x**4*(-1/c**2)**(1/4)*atan(c*x**2)/(2*c**6*x**4*(-1/c**2)**(1
1/4) + 2*c**4*(-1/c**2)**(11/4)) + b*e*(-1/c**2)**(1/4)*atan(c*x**2)/(2*c**
8*x**4*(-1/c**2)**(11/4) + 2*c**6*(-1/c**2)**(11/4)), True))

```

Giac [A]

time = 0.59, size = 184, normalized size = 0.96

$$\frac{\frac{1}{2}bex^2 \arctan(cx^2) + \frac{1}{2}aex^2 + bdx \arctan(cx^2) + adx - \frac{\sqrt{2}bcd \arctan\left(\frac{1}{2}\sqrt{2}\left(2x + \frac{\sqrt{c}}{\sqrt{|c|}}\right)\sqrt{|c|}\right)}{2|c|^{\frac{3}{2}}} - \frac{\sqrt{2}bcd \arctan\left(\frac{1}{2}\sqrt{2}\left(2x - \frac{\sqrt{c}}{\sqrt{|c|}}\right)\sqrt{|c|}\right)}{2|c|^{\frac{3}{2}}} + \frac{(\sqrt{2}bcd\sqrt{|c|} - bce)\log\left(x^2 + \frac{\sqrt{c}}{\sqrt{|c|}} + \frac{1}{|c|}\right)}{4c^2} - \frac{(\sqrt{2}bcd\sqrt{|c|} + bce)\log\left(x^2 - \frac{\sqrt{c}}{\sqrt{|c|}} + \frac{1}{|c|}\right)}{4c^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^2)),x, algorithm="giac")`

[Out] $\frac{1}{2}b^2e^2x^2\arctan(cx^2) + \frac{1}{2}a^2e^2x^2 + b^2d^2x^2\arctan(cx^2) + a^2d^2x - \frac{1}{2}\sqrt{2}b^2c^2d^2\arctan\left(\frac{1}{2}\sqrt{2}(2x + \sqrt{\frac{c}{|c|}})\right)\sqrt{|c|} + \frac{1}{2}\sqrt{2}b^2c^2d^2\arctan\left(\frac{1}{2}\sqrt{2}(2x - \sqrt{\frac{c}{|c|}})\right)\sqrt{|c|} + \frac{1}{4}b^2c^2e^2\log(x^2 + \sqrt{\frac{c}{|c|}} + \frac{1}{|c|}) - \frac{1}{4}b^2c^2e^2\log(x^2 - \sqrt{\frac{c}{|c|}} + \frac{1}{|c|}) + \frac{1}{4}b^2c^2d^2\sqrt{|c|} + \frac{1}{4}b^2c^2d^2\sqrt{|c|}$

Mupad [B]

time = 2.53, size = 203, normalized size = 1.06

$$adx + \frac{ae x^2}{2} + bdx \arctan(cx^2) - \frac{be \ln(x\sqrt{-c\text{II}} - 1)}{4c} - \frac{be \ln(x\sqrt{-c\text{II}} + 1)}{4c} - \frac{be \ln(x\sqrt{c\text{II}} - 1)}{4c} - \frac{be \ln(x\sqrt{c\text{II}} + 1)}{4c} + \frac{be x^2 \arctan(cx^2)}{2} + \frac{bd \ln(x\sqrt{-c\text{II}} - 1)\sqrt{-c\text{II}}}{2c} + \frac{bd \ln(x\sqrt{-c\text{II}} + 1)\sqrt{-c\text{II}}}{2c} - \frac{bd \ln(x\sqrt{c\text{II}} - 1)\sqrt{c\text{II}}}{2c} + \frac{bd \ln(x\sqrt{c\text{II}} + 1)\sqrt{c\text{II}}}{2c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^2))*(d + e*x),x)`

[Out] $a^2d^2x + (a^2e^2x^2)/2 + b^2d^2x^2\arctan(cx^2) - (b^2e^2\log(x*(-c\text{II})^{(1/2)} - 1))/(4*c) - (b^2e^2\log(x*(-c\text{II})^{(1/2)} + 1))/(4*c) - (b^2e^2\log(x*(c\text{II})^{(1/2)} - 1))/(4*c) - (b^2e^2\log(x*(c\text{II})^{(1/2)} + 1))/(4*c) + (b^2e^2x^2\arctan(cx^2))/2 - (b^2d^2\log(x*(-c\text{II})^{(1/2)} - 1)*(-c\text{II})^{(1/2)})/(2*c) + (b^2d^2\log(x*(-c\text{II})^{(1/2)} + 1)*(-c\text{II})^{(1/2)})/(2*c) + (b^2d^2\log(x*(c\text{II})^{(1/2)} - 1)*(c\text{II})^{(1/2)})/(2*c) + (b^2d^2\log(x*(c\text{II})^{(1/2)} + 1)*(c\text{II})^{(1/2)})/(2*c)$

$$3.23 \quad \int \frac{a+b\text{ArcTan}(cx^2)}{d+ex} dx$$

Optimal. Leaf size=501

$$\frac{(a + b\text{ArcTan}(cx^2)) \log(d + ex)}{e} + \frac{bc \log\left(\frac{e(1 - \sqrt[4]{-c^2}x)}{\sqrt[4]{-c^2}^{d+e}}\right) \log(d + ex)}{2\sqrt{-c^2}e} + \frac{bc \log\left(-\frac{e(1 + \sqrt[4]{-c^2}x)}{\sqrt[4]{-c^2}^{d-e}}\right) \log(d + ex)}{2\sqrt{-c^2}e}$$

[Out] $(a+b*\arctan(c*x^2))*\ln(e*x+d)/e+1/2*b*c*\ln(e*(1-(-c^2)^(1/4)*x)/((-c^2)^(1/4)*d+e))*\ln(e*x+d)/e/(-c^2)^(1/2)+1/2*b*c*\ln(-e*(1+(-c^2)^(1/4)*x)/((-c^2)^(1/4)*d-e))*\ln(e*x+d)/e/(-c^2)^(1/2)-1/2*b*c*\ln(e*x+d)*\ln(e*(1-x*(-(-c^2)^(1/2))/(e+d*(-(-c^2)^(1/2))))/e/(-c^2)^(1/2)-1/2*b*c*\ln(e*x+d)*\ln(-e*(1+x*(-(-c^2)^(1/2))/(e+d*(-(-c^2)^(1/2))))/e/(-c^2)^(1/2)+1/2*b*c*polylog(2,(-c^2)^(1/4)*(e*x+d)/((-c^2)^(1/4)*d-e))/e/(-c^2)^(1/2)+1/2*b*c*polylog(2,(-c^2)^(1/4)*(e*x+d)/((-c^2)^(1/4)*d+e))/e/(-c^2)^(1/2)-1/2*b*c*polylog(2,(e*x+d)*(-(-c^2)^(1/2))^(1/2)/(-e+d*(-(-c^2)^(1/2))^(1/2))/e/(-c^2)^(1/2)-1/2*b*c*polylog(2,(e*x+d)*(-(-c^2)^(1/2))^(1/2)/(e+d*(-(-c^2)^(1/2))^(1/2))))/e/(-c^2)^(1/2)$

Rubi [A]

time = 0.65, antiderivative size = 501, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 8, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {4976, 281, 209, 2463, 266, 2441, 2440, 2438}

$$\frac{\log(d+ex)(a+b\text{ArcTan}(cx^2))}{e} + \frac{bc\text{Li}_2\left(\frac{\sqrt{-c^2}x(d+ex)}{\sqrt{-c^2}^{d+e}}\right)}{2\sqrt{-c^2}e} - \frac{bc\text{Li}_2\left(\frac{\sqrt{-c^2}x(d+ex)}{\sqrt{-c^2}^{d-e}}\right)}{2\sqrt{-c^2}e} + \frac{bc\text{Li}_2\left(\frac{\sqrt{-c^2}x(d+ex)}{\sqrt{-c^2}^{d+e}}\right)}{2\sqrt{-c^2}e} - \frac{bc\log(d+ex)\log\left(\frac{e(1-\sqrt{-c^2}x)}{\sqrt{-c^2}^{d+e}}\right)}{2\sqrt{-c^2}e} + \frac{bc\log(d+ex)\log\left(\frac{e(1+\sqrt{-c^2}x)}{\sqrt{-c^2}^{d-e}}\right)}{2\sqrt{-c^2}e} - \frac{bc\log(d+ex)\log\left(\frac{e(1-\sqrt{-c^2}x+1)}{\sqrt{-c^2}^{d+e}}\right)}{2\sqrt{-c^2}e} - \frac{bc\log(d+ex)\log\left(\frac{e(1+\sqrt{-c^2}x+1)}{\sqrt{-c^2}^{d-e}}\right)}{2\sqrt{-c^2}e}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x^2])/(d + e*x), x]$

[Out] $((a + b\text{ArcTan}[c*x^2])*Log[d + e*x])/e + (b*c*Log[(e*(1 - (-c^2)^(1/4)*x))/((-c^2)^(1/4)*d + e)]*Log[d + e*x])/(2*Sqrt[-c^2]*e) + (b*c*Log[-((e*(1 + (-c^2)^(1/4)*x))/((-c^2)^(1/4)*d - e))]*Log[d + e*x])/(2*Sqrt[-c^2]*e) - (b*c*Log[(e*(1 - Sqrt[-Sqrt[-c^2]]*x))/(-Sqrt[-Sqrt[-c^2]]*d + e)]*Log[d + e*x])/(2*Sqrt[-c^2]*e) - (b*c*Log[-((e*(1 + Sqrt[-Sqrt[-c^2]]*x))/(-Sqrt[-Sqrt[-c^2]]*d - e))]/(Sqrt[-Sqrt[-c^2]]*d - e))*Log[d + e*x]/(2*Sqrt[-c^2]*e) + (b*c*PolyLog[2, ((-c^2)^(1/4)*(d + e*x))/((-c^2)^(1/4)*d - e)]/(2*Sqrt[-c^2]*e) - (b*c*PolyLog[2, (Sqrt[-Sqrt[-c^2]]*(d + e*x))/(-Sqrt[-Sqrt[-c^2]]*d - e)]/(2*Sqrt[-c^2]*e) + (b*c*PolyLog[2, ((-c^2)^(1/4)*(d + e*x))/((-c^2)^(1/4)*d + e)]/(2*Sqrt[-c^2]*e) - (b*c*PolyLog[2, (Sqrt[-Sqrt[-c^2]]*(d + e*x))/(-Sqrt[-Sqrt[-c^2]]*d + e)]/(2*Sqrt[-c^2]*e))$

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 281

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 2438

```
Int[Log[(c_)*(d_) + (e_)*(x_)^(n_.)]/(x_), x_Symbol] :> Simp[-PolyLog[2, (-c)*e*x^n]/n, x] /; FreeQ[{c, d, e, n}, x] && EqQ[c*d, 1]
```

Rule 2440

```
Int[((a_) + Log[(c_)*(d_) + (e_)*(x_)])*(b_.))/((f_.) + (g_)*(x_), x_Symbol) :> Dist[1/g, Subst[Int[(a + b*Log[1 + c*e*(x/g)])/x, x], x, f + g*x], x] /; FreeQ[{a, b, c, d, e, f, g}, x] && NeQ[e*f - d*g, 0] && EqQ[g + c*(e*f - d*g), 0]
```

Rule 2441

```
Int[((a_) + Log[(c_)*(d_) + (e_)*(x_)])^(n_.)*(b_.))/((f_.) + (g_)*(x_), x_Symbol) :> Simp[Log[e*((f + g*x)/(e*f - d*g))]*((a + b*Log[c*(d + e*x)^n])/g), x] - Dist[b*e*(n/g), Int[Log[(e*(f + g*x))/(e*f - d*g)]/(d + e*x), x], x] /; FreeQ[{a, b, c, d, e, f, g, n}, x] && NeQ[e*f - d*g, 0]
```

Rule 2463

```
Int[((a_) + Log[(c_)*(d_) + (e_)*(x_)])^(n_.)*(b_.))^(p_)*((h_.)*(x_)^(m_)*((f_) + (g_)*(x_)^(r_.))^(q_), x_Symbol) :> Int[ExpandIntegrand[(a + b*Log[c*(d + e*x)^n])^p, (h*x)^m*(f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q, r}, x] && IntegerQ[m] && IntegerQ[q]
```

Rule 4976

```
Int[((a_) + ArcTan[(c_)*(x_)^(n_.)]*(b_.))/((d_) + (e_)*(x_), x_Symbol) :> Simp[Log[d + e*x]*((a + b*ArcTan[c*x^n])/e), x] - Dist[b*c*(n/e), Int[x^(n - 1)*(Log[d + e*x]/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, n}
```

```
}, x] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \int \frac{a + b \tan^{-1}(cx^2)}{d + ex} dx &= \int \left(\frac{a}{d + ex} + \frac{b \tan^{-1}(cx^2)}{d + ex} \right) dx \\ &= \frac{a \log(d + ex)}{e} + b \int \frac{\tan^{-1}(cx^2)}{d + ex} dx \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 22.38, size = 326, normalized size = 0.65

$$\frac{a \log(d + ex)}{e} + \frac{b \left(2 \operatorname{ArcTan}(cx^2) \log(d + ex) + i \left(\log(d + ex) \log \left(1 - \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e + \sqrt{-1} x} \right) + \log(d + ex) \log \left(1 - \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e - (-1)^{1/4} x} \right) - \log(d + ex) \log \left(1 - \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e - (-1)^{3/4} x} \right) - \log(d + ex) \log \left(1 - \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e - (-1)^{5/4} x} \right) + \operatorname{PolyLog} \left(2, \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e + \sqrt{-1} x} \right) + \operatorname{PolyLog} \left(2, \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e - (-1)^{1/4} x} \right) - \operatorname{PolyLog} \left(2, \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e - (-1)^{3/4} x} \right) - \operatorname{PolyLog} \left(2, \frac{\sqrt{c} \operatorname{atan}(cx)}{\sqrt{c} e - (-1)^{5/4} x} \right) \right) \right)}{2e}$$

Antiderivative was successfully verified.

```
[In] Integrate[(a + b*ArcTan[c*x^2])/(d + e*x), x]
```

```
[Out] (a*Log[d + e*x])/e + (b*(2*ArcTan[c*x^2]*Log[d + e*x] + I*(Log[d + e*x]*Log[1 - (Sqrt[c]*(d + e*x))/(Sqrt[c]*d - (-1)^(1/4)*e)] + Log[d + e*x]*Log[1 - (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + (-1)^(1/4)*e)] - Log[d + e*x]*Log[1 - (Sqrt[c]*(d + e*x))/(Sqrt[c]*d - (-1)^(3/4)*e)] - Log[d + e*x]*Log[1 - (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + (-1)^(3/4)*e)] + PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d - (-1)^(1/4)*e)] + PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + (-1)^(1/4)*e)] - PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d - (-1)^(3/4)*e)] - PolyLog[2, (Sqrt[c]*(d + e*x))/(Sqrt[c]*d + (-1)^(3/4)*e)]))/)/(2*e)
```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.12, size = 138, normalized size = 0.28

method	result
default	$\frac{a \ln(ex+d)}{e} + \frac{b \ln(ex+d) \arctan(cx^2)}{e} - \frac{be \left(\sum_{\substack{-R1=\text{RootOf}(c^2-Z^4-4c^2d-Z^3+6c^2d^2-Z^2-4c^2d^3-Z+c^2d^4+e^4)}} \frac{\ln(ex+d) \ln \left(\frac{-ex+d}{R1} \right)}{2c} \right)}{R1}$
risch	$\frac{ib \ln(ex+d) \ln(-icx^2+1)}{2e} - \frac{ib \ln(ex+d) \ln \left(\frac{e\sqrt{-ic}-(ex+d)c+cd}{e\sqrt{-ic}+cd} \right)}{2e} - \frac{ib \ln(ex+d) \ln \left(\frac{e\sqrt{-ic}+(ex+d)c-cd}{e\sqrt{-ic}-cd} \right)}{2e} - ib \operatorname{dilog} \left(\frac{e\sqrt{-ic}}{ex+d} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a+b*arctan(c*x^2))/(e*x+d), x, method=_RETURNVERBOSE)
```

[Out] $a \ln(e*x+d)/e + b \ln(e*x+d)/e * \arctan(c*x^2) - 1/2 * b*e/c * \text{sum}(1/(_R1^2 - 2*_R1*d + d^2) * (\ln(e*x+d) * \ln((-e*x+_R1-d)/_R1) + \text{dilog}((-e*x+_R1-d)/_R1)), _R1=\text{RootOf}(_Z^4 * c^2 - 4*_Z^3*c^2*d + 6*_Z^2*c^2*d^2 - 4*_Z*c^2*d^3 + c^2*d^4 + e^4))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^2))/(e*x+d),x, algorithm="maxima")`

[Out] $a*e^{(-1)}*\log(x*e + d) + 2*b*\text{integrate}(1/2*\arctan(c*x^2)/(x*e + d), x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^2))/(e*x+d),x, algorithm="fricas")`

[Out] `integral((b*arctan(c*x^2) + a)/(x*e + d), x)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x**2))/(e*x+d),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^2))/(e*x+d),x, algorithm="giac")`

[Out] `integrate((b*arctan(c*x^2) + a)/(e*x + d), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{a + b \operatorname{atan}(c x^2)}{d + e x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int((a + b\arctan(cx^2))/(d + ex), x)$

[Out] $\int((a + b\arctan(cx^2))/(d + ex), x)$

$$3.24 \quad \int \frac{a+b\text{ArcTan}(cx^2)}{(d+ex)^2} dx$$

Optimal. Leaf size=328

$$\frac{bc^2d^3\text{ArcTan}(cx^2)}{e(c^2d^4+e^4)} - \frac{a+b\text{ArcTan}(cx^2)}{e(d+ex)} + \frac{b\sqrt{c}(cd^2-e^2)\text{ArcTan}\left(1-\sqrt{2}\sqrt{c}x\right)}{\sqrt{2}(c^2d^4+e^4)} - \frac{b\sqrt{c}(cd^2-e^2)\text{ArcTan}\left(1+\sqrt{2}\sqrt{c}x\right)}{\sqrt{2}(c^2d^4+e^4)}$$

[Out] $b*c^2*d^3*\arctan(c*x^2)/e/(c^2*d^4+e^4)+(-a-b*\arctan(c*x^2))/e/(e*x+d)-2*b*c*d*e*\ln(e*x+d)/(c^2*d^4+e^4)+1/2*b*c*d*e*\ln(c^2*x^4+1)/(c^2*d^4+e^4)-1/2*b*(c*d^2-e^2)*\arctan(-1+x*2^(1/2)*c^(1/2))*c^(1/2)/(c^2*d^4+e^4)*2^(1/2)-1/2*b*(c*d^2-e^2)*\arctan(1+x*2^(1/2)*c^(1/2))*c^(1/2)/(c^2*d^4+e^4)*2^(1/2)-1/4*b*(c*d^2+e^2)*\ln(1+c*x^2-x*2^(1/2)*c^(1/2))*c^(1/2)/(c^2*d^4+e^4)*2^(1/2)+1/4*b*(c*d^2+e^2)*\ln(1+c*x^2+x*2^(1/2)*c^(1/2))*c^(1/2)/(c^2*d^4+e^4)*2^(1/2)$

Rubi [A]

time = 0.35, antiderivative size = 328, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 13, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.722, Rules used = {4980, 6857, 1890, 1182, 1176, 631, 210, 1179, 642, 1262, 649, 209, 266}

$$-\frac{a+b\text{ArcTan}(cx^2)}{e(d+ex)} + \frac{bc^2d^3\text{ArcTan}(cx^2)}{e(c^2d^4+e^4)} + \frac{b\sqrt{c}\text{ArcTan}\left(1-\sqrt{2}\sqrt{c}x\right)(cd^2-e^2)}{\sqrt{2}(c^2d^4+e^4)} - \frac{b\sqrt{c}\text{ArcTan}\left(\sqrt{2}\sqrt{c}x+1\right)(cd^2-e^2)}{\sqrt{2}(c^2d^4+e^4)} + \frac{bcde\log(c^2x^4+1)}{2(c^2d^4+e^4)} - \frac{2bcde\log(d+ex)}{c^2d^4+e^4} - \frac{b\sqrt{c}(cd^2+e^2)\log(cx^2-\sqrt{2}\sqrt{c}x+1)}{2\sqrt{2}(c^2d^4+e^4)} + \frac{b\sqrt{c}(cd^2+e^2)\log(cx^2+\sqrt{2}\sqrt{c}x+1)}{2\sqrt{2}(c^2d^4+e^4)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x^2])/(d + e*x)^2, x]$

[Out] $(b*c^2*d^3*\text{ArcTan}[c*x^2])/(\text{e}*(\text{c}^2*\text{d}^4 + \text{e}^4)) - (a + b*\text{ArcTan}[c*x^2])/(\text{e}*(\text{d} + \text{e}*x)) + (\text{b}*\text{Sqrt}[\text{c}]*(\text{c}*\text{d}^2 - \text{e}^2)*\text{ArcTan}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{c}]*x])/(\text{Sqrt}[2]*(\text{c}^2*\text{d}^4 + \text{e}^4)) - (\text{b}*\text{Sqrt}[\text{c}]*(\text{c}*\text{d}^2 - \text{e}^2)*\text{ArcTan}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{c}]*x])/(\text{Sqrt}[2]*(\text{c}^2*\text{d}^4 + \text{e}^4)) - (2*\text{b}*\text{c}*\text{d}*\text{e}*\text{Log}[\text{d} + \text{e}*x])/(\text{c}^2*\text{d}^4 + \text{e}^4) - (\text{b}*\text{Sqrt}[\text{c}]*(\text{c}*\text{d}^2 + \text{e}^2)*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[\text{c}]*x + \text{c}*\text{x}^2])/(\text{2}*\text{Sqrt}[2]*(\text{c}^2*\text{d}^4 + \text{e}^4)) + (\text{b}*\text{Sqrt}[\text{c}]*(\text{c}*\text{d}^2 + \text{e}^2)*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[\text{c}]*x + \text{c}*\text{x}^2])/(\text{2}*\text{Sqrt}[2]*(\text{c}^2*\text{d}^4 + \text{e}^4)) + (\text{b}*\text{c}*\text{d}*\text{e}*\text{Log}[1 + \text{c}^2*\text{x}^4])/(\text{2}*(\text{c}^2*\text{d}^4 + \text{e}^4))$

Rule 209

$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \Rightarrow \text{Simp}[(1/(\text{Rt}[\text{a}, 2]*\text{Rt}[\text{b}, 2]))*\text{ArcTan}[\text{Rt}[\text{b}, 2]*(\text{x}/\text{Rt}[\text{a}, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{PosQ}[\text{a}/\text{b}] \& \text{GtQ}[\text{a}, 0] \text{ || } \text{GtQ}[\text{b}, 0])$

Rule 210

$\text{Int}[((a_) + (b_)*(x_)^2)^{-1}, \text{x_Symbol}] \Rightarrow \text{Simp}[(-(Rt[-a, 2]*Rt[-b, 2])^{(-1)})*\text{ArcTan}[\text{Rt}[-b, 2]*(\text{x}/\text{Rt}[-a, 2])], \text{x}] /; \text{FreeQ}[\{\text{a}, \text{b}\}, \text{x}] \& \text{PosQ}[\text{a}/\text{b}] \& (\text{LtQ}[\text{a}, 0] \text{ || } \text{LtQ}[\text{b}, 0])$

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_ .)*(x_)^(n_.)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 631

```
Int[((a_) + (b_ .)*(x_) + (c_ .)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_ .)*(x_))/((a_) + (b_ .)*(x_) + (c_ .)*(x_)^2), x_Symbol] :> Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 649

```
Int[((d_) + (e_ .)*(x_))/((a_) + (c_ .)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 1176

```
Int[((d_) + (e_ .)*(x_)^2)/((a_) + (c_ .)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x]], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_ .)*(x_)^2)/((a_) + (c_ .)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 1182

```
Int[((d_) + (e_ .)*(x_)^2)/((a_) + (c_ .)*(x_)^4), x_Symbol] :> With[{q = Rt[a*c, 2]}, Dist[(d*q + a*e)/(2*a*c), Int[(q + c*x^2)/(a + c*x^4), x], x] + Dist[(d*q - a*e)/(2*a*c), Int[(q - c*x^2)/(a + c*x^4), x], x] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 + a*e^2, 0] && NeQ[c*d^2 - a*e^2, 0] && NegQ[(-a)*c]
```

Rule 1262

```
Int[(x_)*((d_) + (e_.)*(x_)^2)^(q_.)*((a_) + (c_.)*(x_)^4)^(p_.), x_Symbol]
:> Dist[1/2, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^2], x] /; FreeQ
[{a, c, d, e, p, q}, x]
```

Rule 1890

```
Int[(Pq_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = Sum[x^ii*((Coeff
[Pq, x, ii] + Coeff[Pq, x, n/2 + ii])*x^(n/2))/(a + b*x^n)), {ii, 0, n/2 - 1
}], Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[n/2,
0] && Expon[Pq, x] < n
```

Rule 4980

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_)]*(b_.))*((d_) + (e_.)*(x_))^(m_.), x_Sy
mbol] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcTan[c*x^n])/(e*(m + 1))), x] - D
ist[b*c*(n/(e*(m + 1))), Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 + c^2*x^(2*n))
), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rule 6857

```
Int[(u_)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionE
xpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ
[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{a + b \tan^{-1}(cx^2)}{(d + ex)^2} dx &= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} + \frac{b \int \frac{2cx}{(d+ex)(1+c^2x^4)} dx}{e} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} + \frac{(2bc) \int \frac{x}{(d+ex)(1+c^2x^4)} dx}{e} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} + \frac{(2bc) \int \left(-\frac{de^3}{(c^2d^4+e^4)(d+ex)} + \frac{e^3+c^2d^3x-c^2d^2ex^2+c^2de^2x^3}{(c^2d^4+e^4)(1+c^2x^4)} \right) dx}{e} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} - \frac{2bcde \log(d + ex)}{c^2d^4 + e^4} + \frac{(2bc) \int \frac{e^3+c^2d^3x-c^2d^2ex^2+c^2de^2x^3}{1+c^2x^4} dx}{e(c^2d^4 + e^4)} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} - \frac{2bcde \log(d + ex)}{c^2d^4 + e^4} + \frac{(2bc) \int \left(\frac{e^3-c^2d^2ex^2}{1+c^2x^4} + \frac{x(c^2d^3+c^2de^2x^2)}{1+c^2x^4} \right) dx}{e(c^2d^4 + e^4)} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} - \frac{2bcde \log(d + ex)}{c^2d^4 + e^4} + \frac{(2bc) \int \frac{e^3-c^2d^2ex^2}{1+c^2x^4} dx}{e(c^2d^4 + e^4)} + \frac{(2bc) \int \frac{x(c^2d^3+c^2de^2x^2)}{1+c^2x^4} dx}{e(c^2d^4 + e^4)} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} - \frac{2bcde \log(d + ex)}{c^2d^4 + e^4} + \frac{(bc) \text{Subst} \left(\int \frac{c^2d^3+c^2de^2x}{1+c^2x^2} dx, x, x^2 \right)}{e(c^2d^4 + e^4)} - \frac{(b) \text{Subst} \left(\int \frac{c^2d^3+c^2de^2x^2}{1+c^2x^4} dx, x, x^2 \right)}{e(c^2d^4 + e^4)} \\
&= -\frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} - \frac{2bcde \log(d + ex)}{c^2d^4 + e^4} + \frac{(bc^3d^3) \text{Subst} \left(\int \frac{1}{1+c^2x^2} dx, x, x^2 \right)}{e(c^2d^4 + e^4)} + \frac{(bc^3d^3) \text{Subst} \left(\int \frac{1}{1+c^2x^4} dx, x, x^2 \right)}{e(c^2d^4 + e^4)} \\
&= \frac{bc^2d^3 \tan^{-1}(cx^2)}{e(c^2d^4 + e^4)} - \frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} - \frac{2bcde \log(d + ex)}{c^2d^4 + e^4} - \frac{b\sqrt{c}(cd^2 + e^2) \log(1 - \sqrt{2}\sqrt{c}x)}{2\sqrt{2}(c^2d^4 + e^4)} \\
&= \frac{bc^2d^3 \tan^{-1}(cx^2)}{e(c^2d^4 + e^4)} - \frac{a + b \tan^{-1}(cx^2)}{e(d + ex)} + \frac{b\sqrt{c}(cd^2 - e^2) \tan^{-1}(1 - \sqrt{2}\sqrt{c}x)}{\sqrt{2}(c^2d^4 + e^4)} - \frac{b\sqrt{c}(cd^2 + e^2) \log(1 + \sqrt{2}\sqrt{c}x)}{\sqrt{2}(c^2d^4 + e^4)} - \frac{b\sqrt{c}(cd^2 - e^2) \log(1 - \sqrt{2}\sqrt{c}x)}{\sqrt{2}(c^2d^4 + e^4)}
\end{aligned}$$

Mathematica [A]

time = 0.47, size = 321, normalized size = 0.98

$$\frac{-4a(c^2d^4 + e^4) + 4b(c^2d^4 + e^4) \text{ArcTan}(cx^2) + 2b\sqrt{c} \left(2c^{1/2}d^4 - \sqrt{2}cd^2e + \sqrt{2}e^4 \right) (d + ex) \text{ArcTan} \left(1 - \sqrt{2}\sqrt{c}x \right) + 2b\sqrt{c} \left(2c^{1/2}d^4 + \sqrt{2}cd^2e - \sqrt{2}e^4 \right) (d + ex) \text{ArcTan} \left(1 + \sqrt{2}\sqrt{c}x \right) + 8bcde^2(d + ex) \log(d + ex) + \sqrt{2}b\sqrt{c}e(cd^2 + e^2)(d + ex) \log(1 - \sqrt{2}\sqrt{c}x + cx^2) - \sqrt{2}b\sqrt{c}e(cd^2 + e^2)(d + ex) \log(1 + \sqrt{2}\sqrt{c}x + cx^2) - 2bcde^2(d + ex) \log(1 + c^2x^4)}{4c(c^2d^4 + e^4)(d + ex)}$$

Antiderivative was successfully verified.

[In] `Integrate[(a + b*ArcTan[c*x^2])/(d + e*x)^2, x]`

[Out] `-1/4*(4*a*(c^2*d^4 + e^4) + 4*b*(c^2*d^4 + e^4)*ArcTan[c*x^2] + 2*b*Sqrt[c]*(2*c^(3/2)*d^3 - Sqrt[2]*c*d^2*e + Sqrt[2]*e^3)*(d + e*x)*ArcTan[1 - Sqrt[2]*Sqrt[c]*x] + 2*b*Sqrt[c]*(2*c^(3/2)*d^3 + Sqrt[2]*c*d^2*e - Sqrt[2]*e^3)`

$$\begin{aligned} & * (d + e*x) * \text{ArcTan}[1 + \text{Sqrt}[2] * \text{Sqrt}[c] * x] + 8*b*c*d*e^2*(d + e*x)*\text{Log}[d + e*x] \\ & + \text{Sqrt}[2]*b*\text{Sqrt}[c]*e*(c*d^2 + e^2)*(d + e*x)*\text{Log}[1 - \text{Sqrt}[2]*\text{Sqrt}[c]*x \\ & + c*x^2] - \text{Sqrt}[2]*b*\text{Sqrt}[c]*e*(c*d^2 + e^2)*(d + e*x)*\text{Log}[1 + \text{Sqrt}[2]*\text{Sqrt}[c]*x \\ & + c*x^2] - 2*b*c*d*e^2*(d + e*x)*\text{Log}[1 + c^2*x^4])/(e*(c^2*d^4 + e^4) \\ & *(d + e*x)) \end{aligned}$$

Maple [A]

time = 0.18, size = 433, normalized size = 1.32

method	result
default	$-\frac{a}{(ex+d)e} - \frac{b \arctan(cx^2)}{(ex+d)e} - \frac{2bcde \ln(ex+d)}{c^2 d^4 + e^4} + \frac{b e^2 c \left(\frac{1}{c^2}\right)^{\frac{1}{4}} \sqrt{2} \arctan\left(\frac{\sqrt{2} \frac{x}{\left(\frac{1}{c^2}\right)^{\frac{1}{4}}} + 1}{\left(\frac{1}{c^2}\right)^{\frac{1}{4}}}\right)}{2c^2 d^4 + 2e^4} + \frac{b e^2 c \left(\frac{1}{c^2}\right)^{\frac{1}{4}} \sqrt{2} \arctan\left(\frac{\sqrt{2} \frac{x}{\left(\frac{1}{c^2}\right)^{\frac{1}{4}}} - 1}{\left(\frac{1}{c^2}\right)^{\frac{1}{4}}}\right)}{2c^2 d^4 + 2e^4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x^2))/(e*x+d)^2,x,method=_RETURNVERBOSE)`

[Out]

$$\begin{aligned} & -a/(e*x+d)/e - b/(e*x+d)/e*\arctan(c*x^2) - 2*b*c*d*e*\ln(e*x+d)/(c^2*d^4+e^4)+1/2*b*e^2*c/(c^2*d^4+e^4)*(1/c^2)^(1/4)*2^(1/2)*\arctan(2^(1/2)/(1/c^2)^(1/4)*x+1)+1/2*b*e^2*c/(c^2*d^4+e^4)*(1/c^2)^(1/4)*2^(1/2)*\arctan(2^(1/2)/(1/c^2)^(1/4)*x-1)+1/4*b*e^2*c/(c^2*d^4+e^4)*(1/c^2)^(1/4)*2^(1/2)*\ln((x^2+(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2))/((x^2-(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2)))+b/e*c^3/(c^2*d^4+e^4)*d^3/(c^2)^(1/2)*\arctan(x^2*(c^2)^(1/2))-1/4*b*c/(c^2*d^4+e^4)*d^2/(1/c^2)^(1/4)*2^(1/2)*\ln((x^2-(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2))/(x^2+(1/c^2)^(1/4)*x*2^(1/2)+(1/c^2)^(1/2)))-1/2*b*c/(c^2*d^4+e^4)*d^2/(1/c^2)^(1/4)*2^(1/2)*\arctan(2^(1/2)/(1/c^2)^(1/4)*x+1)-1/2*b*c/(c^2*d^4+e^4)*d^2/(1/c^2)^(1/4)*2^(1/2)*\arctan(2^(1/2)/(1/c^2)^(1/4)*x-1)+1/2*b*c*d*e*\ln(c^2*x^4+1)/(c^2*d^4+e^4) \end{aligned}$$
Maxima [A]

time = 0.48, size = 286, normalized size = 0.87

$$\begin{aligned} & \frac{1}{4} \left(\frac{8 de \log(xe + d)}{c^2 d^4 + e^4} - \frac{\sqrt{2} (\alpha^2 e + \sqrt{2} \sqrt{c} d e^2 + e^3) \log(\alpha^2 + \sqrt{2} \sqrt{c} x + 1)}{\sqrt{c}} - \frac{\sqrt{2} (\alpha^2 e - \sqrt{2} \sqrt{c} d e^2 + e^3) \log(\alpha^2 - \sqrt{2} \sqrt{c} x + 1)}{\sqrt{c}} - \frac{2 (2 \alpha^2 d^3 + \sqrt{2} \alpha^2 d^2 e - \sqrt{2} \sqrt{c} e^3) \arctan\left(\frac{\sqrt{2} (x e + \sqrt{2} \sqrt{c})}{z \sqrt{c}}\right)}{c} + \frac{2 (2 \alpha^2 d^3 - \sqrt{2} \alpha^2 d^2 e + \sqrt{2} \sqrt{c} e^3) \arctan\left(\frac{\sqrt{2} (x e - \sqrt{2} \sqrt{c})}{z \sqrt{c}}\right)}{c} \right) c + \frac{4 \arctan(cx^2)}{xe^2 + de} b - \frac{a}{xe^2 + de} \end{aligned}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^2))/(e*x+d)^2,x, algorithm="maxima")`

[Out]

$$\begin{aligned} & -1/4*((8*d*e*log(xe + d)/(c^2*d^4 + e^4) - (\sqrt{2)*(c*d^2*e + \sqrt{2)*sqrt(c)*sqrt(d)*sqrt(e^3)*log(c*x^2 + sqrt(2)*sqrt(c)*x + 1)/sqrt(c) - sqrt(2)*(c*d^2*e - sqrt(2)*sqrt(c)*d*e^2 + e^3)*log(c*x^2 - sqrt(2)*sqrt(c)*x + 1)/sqrt(c) - 2*(2*c^2*d^3 + sqrt(2)*c^(3/2)*d^2*e - sqrt(2)*sqrt(c)*e^3)*arctan(1/2*sqrt(2)*(2*c*x + sqrt(2)*sqrt(c))/sqrt(c))/c + 2*(2*c^2*d^3 - sqrt(2)*c^3) \end{aligned}$$

$\frac{3}{2}d^2e + \sqrt{2}\sqrt{c}e^3 \operatorname{arctan}\left(\frac{1}{2}\sqrt{2}(2cx - \sqrt{2}\sqrt{c})\right) / (c^2d^4e + e^5) * c + 4\operatorname{arctan}(cx^2) / (x^2e^2 + d^2e) * b - a / (x^2e^2 + d^2e)$

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(cx^2))/(e*x+d)^2,x, algorithm="fricas")`

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(cx**2))/(e*x+d)**2,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(cx^2))/(e*x+d)^2,x, algorithm="giac")`

[Out] undef

Mupad [B]

time = 0.66, size = 883, normalized size = 2.69

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(cx^2))/(d + e*x)^2,x)`

[Out] `symsum(log((320*root(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)^4*c^8*e^9*x - 128*root(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)^4*c^10*d^5*e^4 + 16*b^4*c^10*e*x - 8*root(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)^4*c^8*e^9*x)), z)`

$$\begin{aligned}
& 4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)*b^3*c^9*e^3 + \\
& 384*\text{root}(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2* \\
& 2*e^2*z^2 + b^4*c^2, z, k)^4*c^8*d*e^8 + 8*\text{root}(16*c^2*d^4*e^4*z^4 + 16*e^8 \\
& *z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)*b^3*c^11*d \\
& ^3*x - 320*\text{root}(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2* \\
& c^2*d^2*e^2*z^2 + b^4*c^2, z, k)^3*b*c^9*d^2*e^5 - 192*\text{root}(16*c^2*d^4*e^4* \\
& z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k) \\
&)^4*c^10*d^4*e^5*x + 32*\text{root}(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5 \\
& *z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)^3*b*c^11*d^5*e^2*x + 64*\text{root}(\\
& 16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 \\
& + b^4*c^2, z, k)^2*b^2*c^10*d^2*e^3*x - 416*\text{root}(16*c^2*d^4*e^4*z^4 + 16*e^ \\
& 8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k)^3*b*c^9*d \\
& *e^6*x)/e^2)*\text{root}(16*c^2*d^4*e^4*z^4 + 16*e^8*z^4 - 32*b*c*d*e^5*z^3 + 8*b^ \\
& 2*c^2*d^2*e^2*z^2 + b^4*c^2, z, k), k, 1, 4) - a/(d*e + e^2*x) - (b*atan(c* \\
& x^2))/(d*e + e^2*x) - (2*b*c*d*e*log(d + e*x))/(e^4 + c^2*d^4)
\end{aligned}$$

$$3.25 \quad \int (d + ex) (a + b \operatorname{ArcTan}(cx^2))^2 dx$$

Optimal. Leaf size=1325

$$a^2 dx - \frac{2(-1)^{3/4} ab d \operatorname{ArcTan}((-1)^{3/4} \sqrt{c} x)}{\sqrt{c}} + \frac{(-1)^{3/4} b^2 d \operatorname{ArcTan}((-1)^{3/4} \sqrt{c} x)^2}{\sqrt{c}} + \frac{i e (a + b \operatorname{ArcTan}(c x^2))^2}{2 c} + \frac{1}{2} e$$

Rubi [A]

time = 1.62, antiderivative size = 1325, normalized size of antiderivative = 1.00, number of steps used = 77, number of rules used = 26, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 1.444, Rules used = {4982, 4932, 2498, 327, 209, 2500, 2526, 2520, 12, 5040, 4964, 2449, 2352, 212,

2636, 211, 5048, 4966, 2497, 214, 6139, 6057, 6131, 6055, 4948, 4930}

Antiderivative was successfully verified.

```
[In] Int[(d + e*x)*(a + b*ArcTan[c*x^2])^2, x]
[Out] a^2*d*x - (2*(-1)^(3/4)*a*b*d*ArcTan[(-1)^(3/4)*Sqrt[c]*x])/Sqrt[c] + ((-1)^(3/4)*b^2*d*ArcTan[(-1)^(3/4)*Sqrt[c]*x]^2)/Sqrt[c] + ((I/2)*e*(a + b*ArcTan[c*x^2])^2)/c + (e*x^2*(a + b*ArcTan[c*x^2])^2)/2 + (2*(-1)^(3/4)*a*b*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])/Sqrt[c] - ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x]^2)/Sqrt[c] + (2*(-1)^(1/4)*b^2*d*ArcTan[(-1)^(3/4)*Sqrt[c]*x])*Log[2/(1 - (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] - (2*(-1)^(1/4)*b^2*d*ArcTan[(-1)^(3/4)*Sqrt[c]*x])*Log[2/(1 + (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] + ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[(Sqrt[2]*((-1)^(1/4) + Sqrt[c]*x))/(1 + (-1)^(1/4)*Sqrt[c]*x))/Sqrt[c] + (2*(-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[2/(1 - (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] - (2*(-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[2/(1 + (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] + ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[-((Sqrt[2]*((-1)^(3/4) + Sqrt[c]*x))/(1 + (-1)^(3/4)*Sqrt[c]*x))]/Sqrt[c] + ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[((1 + I)*(1 + (-1)^(1/4)*Sqrt[c]*x))/(1 + (-1)^(3/4)*Sqrt[c]*x)]/Sqrt[c] + ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[((1 - I)*(1 + (-1)^(3/4)*Sqrt[c]*x))/(1 + (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] + I*a*b*d*x*Log[1 - I*c*x^2] + ((-1)^(1/4)*b^2*d*ArcTan[(-1)^(3/4)*Sqrt[c]*x])*Log[1 - I*c*x^2]/Sqrt[c] - ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[1 - I*c*x^2]/Sqrt[c] - (b^2*d*x*Log[1 - I*c*x^2]^2)/4 + (b*e*(a + b*ArcTan[c*x^2]))*Log[2/(1 + I*c*x^2)]/c - I*a*b*d*x*Log[1 + I*c*x^2] - ((-1)^(1/4)*b^2*d*ArcTan[(-1)^(3/4)*Sqrt[c]*x])*Log[1 + I*c*x^2]/Sqrt[c] + ((-1)^(1/4)*b^2*d*ArcTanh[(-1)^(3/4)*Sqrt[c]*x])*Log[1 + I*c*x^2]/Sqrt[c] + (b^2*d*x*Log[1 - I*c*x^2])*Log[1 + I*c*x^2]/2 - (b^2*d*x*Log[1 + I*c*x^2]^2)/4 + ((-1)^(3/4)*b^2*d*PolyLog[2, 1 - 2/(1 - (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] + ((-1)^(3/4)*b^2*d*PolyLog[2, 1 - 2/(1 + (-1)^(1/4)*Sqrt[c]*x)]/Sqrt[c] - ((-1)^(3/4)*b^2*d*PolyLog[2, 1 - (Sqrt[2]*((-1)^(1/4) + Sqrt[c]*x))/(1 + (-1)^(1/4)*Sqrt[c]*x)])/(2*Sqrt[c]) + ((-1)^(1/4)*b^2*d*PolyLog[2, 1 - 2/(1 - (-1)^(3/4)*Sqrt[c]*x)]/Sqrt[c] + ((-1)^(1/4)*b^2*d*PolyLog[2, 1 - 2/(1 + (-1)^(3/4)*Sqrt[c]*x)]/Sqrt[c] - ((-1)^(1/4)*b^2*d*PolyLog[2, 1 + (Sqrt[2]*((-1)^(3/4) + Sqrt[c]*x))/(1 + (-1)^(3/4)*Sqrt[c]*x)])/(2*Sqrt[c]) - ((-1)^(1/4)*b^2*d*PolyLog[2, 1 - ((1 + I)*(1 + (-1)^(1/4)*Sqrt[c]*x))/Sqrt[c]]/(1 + (-1)^(3/4)*Sqrt[c]*x)]/(2*Sqrt[c]) - ((-1)^(3/4)*b^2*d*PolyLog[2, 1 - ((1 - I)*(1 + (-1)^(3/4)*Sqrt[c]*x))/(1 + (-1)^(1/4)*Sqrt[c]*x)]/(2*Sqrt[c]) + ((I/2)*b^2*e*PolyLog[2, 1 - 2/(1 + I*c*x^2)])/c
```

Rule 12

```
Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 211

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 327

```
Int[((c_)*(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> Simp[c^(n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*(m + n*p + 1))), x] - Dist[a*c^n*((m - n + 1)/(b*(m + n*p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, c, p}, x] && IGtQ[n, 0] && GtQ[m, n - 1] && NeQ[m + n*p + 1, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 2352

```
Int[Log[(c_)*(x_)]/((d_) + (e_)*(x_)), x_Symbol] :> Simp[(-e^(-1))*PolyLog[2, 1 - c*x], x] /; FreeQ[{c, d, e}, x] && EqQ[e + c*d, 0]
```

Rule 2449

```
Int[Log[(c_)/((d_) + (e_)*(x_))]/((f_) + (g_)*(x_)^2), x_Symbol] :> Dist[-e/g, Subst[Int[Log[2*d*x]/(1 - 2*d*x), x], x, 1/(d + e*x)], x] /; FreeQ[{c, d, e, f, g}, x] && EqQ[c, 2*d] && EqQ[e^2*f + d^2*g, 0]
```

Rule 2497

```
Int[Log[u_]*(Pq_)^(m_), x_Symbol] :> With[{C = FullSimplify[Pq^m*((1 - u)/D[u, x])]}, Simp[C*PolyLog[2, 1 - u], x] /; FreeQ[C, x]] /; IntegerQ[m] && PolyQ[Pq, x] && RationalFunctionQ[u, x] && LeQ[RationalFunctionExponents[u,
```

$x] [[2]], \text{Expon}[Pq, x]]$

Rule 2498

```
Int[Log[(c_.)*((d_) + (e_.)*(x_)^(n_))^(p_.)], x_Symbol] :> Simp[x*Log[c*(d + e*x^n)^p], x] - Dist[e*n*p, Int[x^n/(d + e*x^n), x], x] /; FreeQ[{c, d, e, n, p}, x]
```

Rule 2500

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.*(x_)^(n_))^(p_.))^(b_.))]^(q_), x_Symbol] :> Simp[x*(a + b*Log[c*(d + e*x^n)^p])^q, x] - Dist[b*e*n*p*q, Int[x^n*((a + b*Log[c*(d + e*x^n)^p])^(q - 1)/(d + e*x^n)), x], x] /; FreeQ[{a, b, c, d, e, n, p}, x] && IGtQ[q, 0] && (EqQ[q, 1] || IntegerQ[n])
```

Rule 2520

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.*(x_)^(n_))^(p_.))^(b_.))]/((f_) + (g_.)*x_)^2, x_Symbol] :> With[{u = IntHide[1/(f + g*x^2), x]}, Simp[u*(a + b*Log[c*(d + e*x^n)^p]), x] - Dist[b*e*n*p, Int[u*(x^(n - 1)/(d + e*x^n)), x], x] /; FreeQ[{a, b, c, d, e, f, g, n, p}, x] && IntegerQ[n]
```

Rule 2526

```
Int[((a_.) + Log[(c_.)*((d_) + (e_.*(x_)^(n_))^(p_.))^(b_.))]^(q_.*(x_)^(m_.*((f_) + (g_.)*x_)^(s_))^(r_.)), x_Symbol] :> Int[ExpandIntegrand[(a + b*Log[c*(d + e*x^n)^p])^q, x^m*(f + g*x^s)^r, x], x] /; FreeQ[{a, b, c, d, e, f, g, m, n, p, q, r, s}, x] && IGtQ[q, 0] && IntegerQ[m] && IntegerQ[r] && IntegerQ[s]
```

Rule 2636

```
Int[Log[v_]*Log[w_], x_Symbol] :> Simp[x*Log[v]*Log[w], x] + (-Int[Simplify[Integrand[x*Log[w]*(D[v, x]/v), x], x] - Int[Simplify[Integrand[x*Log[v]*(D[w, x]/w), x], x]] /; InverseFunctionFreeQ[v, x] && InverseFunctionFreeQ[w, x]
```

Rule 4930

```
Int[((a_.) + ArcTan[(c_.*(x_)^(n_.))^(b_.))]^(p_.), x_Symbol] :> Simp[x*(a + b*ArcTan[c*x^n])^p, x] - Dist[b*c*n*p, Int[x^n*((a + b*ArcTan[c*x^n])^(p - 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, n}, x] && IGtQ[p, 0] && (EqQ[n, 1] || EqQ[p, 1])
```

Rule 4932

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_)]*(b_.))^(p_), x_Symbol] :> Int[ExpandIn-
tegrand[(a + (I*b*Log[1 - I*c*x^n])/2 - (I*b*Log[1 + I*c*x^n])/2)^p, x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[p, 1] && IGtQ[n, 0]
```

Rule 4948

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_)]*(b_.))^(p_)*(x_)^(m_.), x_Symbol] :>
Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*ArcTan[c*x])^p, x],
x, x^n], x] /; FreeQ[{a, b, c, m, n}, x] && IGtQ[p, 1] && IntegerQ[Simplify
[(m + 1)/n]]
```

Rule 4964

```
Int[((a_.) + ArcTan[(c_.)*(x_)*(b_.)])^(p_.)/((d_) + (e_.)*(x_)), x_Symbol]
:> Simp[(-(a + b*ArcTan[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e),
Int[(a + b*ArcTan[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 + c^2*x^2)), x], x] /;
FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 + e^2, 0]
```

Rule 4966

```
Int[((a_.) + ArcTan[(c_.)*(x_)*(b_.)])/((d_) + (e_.)*(x_)), x_Symbol] :> Si-
mp[(-(a + b*ArcTan[c*x]))*(Log[2/(1 - I*c*x)]/e), x] + (Dist[b*(c/e), Int[Log[2/(1 -
I*c*x)]/(1 + c^2*x^2), x], x] - Dist[b*(c/e), Int[Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/(1 + c^2*x^2), x], x] + Simp[(a + b*ArcTan[c*x])*(Log[2*c*((d + e*x)/((c*d + I*e)*(1 - I*c*x)))]/e), x]) /;
FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 + e^2, 0]
```

Rule 4982

```
Int[((a_.) + ArcTan[(c_.)*(x_)^(n_)]*(b_.))^(p_)*((d_) + (e_.)*(x_))^(m_.),
x_Symbol] :> Int[ExpandIntegrand[(a + b*ArcTan[c*x^n])^p, (d + e*x)^m, x], x]
/; FreeQ[{a, b, c, d, e, n}, x] && IGtQ[p, 1] && IGtQ[m, 0]
```

Rule 5040

```
Int[((((a_.) + ArcTan[(c_.)*(x_)*(b_.)])^(p_)*(x_))/((d_) + (e_.)*(x_)^2),
x_Symbol) :> Simp[(-I)*((a + b*ArcTan[c*x])^(p + 1)/(b*e*(p + 1))), x] - Di-
st[1/(c*d), Int[(a + b*ArcTan[c*x])^p/(I - c*x), x], x] /; FreeQ[{a, b, c,
d, e}, x] && EqQ[e, c^2*d] && IGtQ[p, 0]
```

Rule 5048

```
Int[((((a_.) + ArcTan[(c_.)*(x_)*(b_.)])*(x_)^(m_.))/((d_) + (e_.)*(x_)^2),
x_Symbol) :> Int[ExpandIntegrand[a + b*ArcTan[c*x], x^m/(d + e*x^2), x], x]
/; FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0])
```

Rule 6055

```
Int[((a_.) + ArcTanh[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)), x_Symbol]
  :> Simp[(-(a + b*ArcTanh[c*x])^p)*(Log[2/(1 + e*(x/d))]/e), x] + Dist[b*c*(p/e),
  Int[(a + b*ArcTanh[c*x])^(p - 1)*(Log[2/(1 + e*(x/d))]/(1 - c^2*x^2)), x], x] /;
  FreeQ[{a, b, c, d, e}, x] && IGtQ[p, 0] && EqQ[c^2*d^2 - e^2, 0]
```

Rule 6057

```
Int[((a_.) + ArcTanh[(c_)*(x_)]*(b_.))^(p_.)/((d_) + (e_)*(x_)), x_Symbol]
  :> Simp[(-(a + b*ArcTanh[c*x]))*(Log[2/(1 + c*x)]/e), x] + (Dist[b*(c/e),
  Int[Log[2/(1 + c*x)]/(1 - c^2*x^2), x], x] - Dist[b*(c/e),
  Int[Log[2*c*((d + e*x)/(c*d + e)*(1 + c*x))]/(1 - c^2*x^2), x], x] + Simp[(a + b*ArcTanh[c*x])
  *(Log[2*c*((d + e*x)/(c*d + e)*(1 + c*x))]/e), x]) /;
  FreeQ[{a, b, c, d, e}, x] && NeQ[c^2*d^2 - e^2, 0]
```

Rule 6131

```
Int[((((a_.) + ArcTanh[(c_)*(x_)]*(b_.))^(p_.)*(x_))/((d_) + (e_)*(x_)^2),
  x_Symbol) :> Simp[(a + b*ArcTanh[c*x])^(p + 1)/(b*e*(p + 1)), x] + Dist[1/(
  c*d), Int[(a + b*ArcTanh[c*x])^p/(1 - c*x), x], x] /;
  FreeQ[{a, b, c, d, e}, x] && EqQ[c^2*d + e, 0] && IGtQ[p, 0]
```

Rule 6139

```
Int[((((a_.) + ArcTanh[(c_)*(x_)]*(b_.))*(x_)^(m_.))/((d_) + (e_)*(x_)^2),
  x_Symbol) :> Int[ExpandIntegrand[a + b*ArcTanh[c*x], x^m/(d + e*x^2), x],
  x] /;
  FreeQ[{a, b, c, d, e}, x] && IntegerQ[m] && !(EqQ[m, 1] && NeQ[a, 0])
```

Rubi steps

$$\begin{aligned}
\int (d+ex) \left(a + b \tan^{-1}(cx^2) \right)^2 dx &= \int \left(a^2(d+ex) + 2ab(d+ex) \tan^{-1}(cx^2) + b^2(d+ex) \tan^{-1}(cx^2)^2 \right) dx \\
&= \frac{a^2(d+ex)^2}{2e} + (2ab) \int (d+ex) \tan^{-1}(cx^2) dx + b^2 \int (d+ex) \tan^{-1}(cx^2)^2 dx \\
&= \frac{a^2(d+ex)^2}{2e} + (2ab) \int (d \tan^{-1}(cx^2) + ex \tan^{-1}(cx^2)) dx + b^2 \int (d \tan^{-1}(cx^2)^2 + ex \tan^{-1}(cx^2)^2) dx \\
&= \frac{a^2(d+ex)^2}{2e} + (2abd) \int \tan^{-1}(cx^2) dx + (b^2d) \int \tan^{-1}(cx^2)^2 dx + (b^2e) \int ex \tan^{-1}(cx^2)^2 dx \\
&= \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) + (b^2d) \int \left(-\frac{1}{4} \frac{1}{1+c^2x^2} \right) dx \\
&= \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) - \frac{abe \log(1+c^2x^2)}{2c} \\
&\quad + \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) - \frac{1}{4} b^2 dx \log^2(1+c^2x^2) \\
&= \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) - \frac{1}{4} b^2 dx \log^2(1+c^2x^2) \\
&= \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) + \frac{\sqrt{2} abd \tan^{-1}(cx^2)}{c} \\
&= -\frac{1}{2} b^2 ex^2 + \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) + \frac{\sqrt{2} abd \tan^{-1}(cx^2)}{c} \\
&= -4b^2 dx - \frac{1}{2} b^2 ex^2 + \frac{a^2(d+ex)^2}{2e} + 2abdx \tan^{-1}(cx^2) + abex^2 \tan^{-1}(cx^2) \\
&= -\frac{1}{2} b^2 ex^2 + \frac{a^2(d+ex)^2}{2e} - \frac{2\sqrt{-1} b^2 d \tan^{-1}((-1)^{3/4} \sqrt{c} x)}{\sqrt{c}} + \frac{(-1)^{3/4} b^2 d \tan^{-1}((-1)^{3/4} \sqrt{c} x)}{\sqrt{c}} \\
&= \frac{a^2(d+ex)^2}{2e} + \frac{(-1)^{3/4} b^2 d \tan^{-1}((-1)^{3/4} \sqrt{c} x)^2}{\sqrt{c}} + 2abdx \tan^{-1}(cx^2) \\
&= \frac{a^2(d+ex)^2}{2e} + \frac{(-1)^{3/4} b^2 d \tan^{-1}((-1)^{3/4} \sqrt{c} x)^2}{\sqrt{c}} + 2abdx \tan^{-1}(cx^2) \\
&= \frac{a^2(d+ex)^2}{2e} + \frac{(-1)^{3/4} b^2 d \tan^{-1}((-1)^{3/4} \sqrt{c} x)^2}{\sqrt{c}} + 2abdx \tan^{-1}(cx^2)
\end{aligned}$$

Mathematica [B] Both result and optimal contain complex but leaf count is larger than twice the leaf count of optimal. 5420 vs. 2(1325) = 2650.
 time = 30.64, size = 5420, normalized size = 4.09

Result too large to show

Warning: Unable to verify antiderivative.

[In] `Integrate[(d + e*x)*(a + b*ArcTan[c*x^2])^2, x]`
 [Out] Result too large to show

Maple [F]

time = 0.10, size = 0, normalized size = 0.00

$$\int (ex + d) (a + b \arctan(cx^2))^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)*(a+b*arctan(c*x^2))^2, x)`
 [Out] `int((e*x+d)*(a+b*arctan(c*x^2))^2, x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^2))^2, x, algorithm="maxima")`
 [Out] `12*b^2*c^2*e*integrate(1/16*x^5*arctan(c*x^2)^2/(c^2*x^4 + 1), x) + b^2*c^2 *e*integrate(1/16*x^5*log(c^2*x^4 + 1)^2/(c^2*x^4 + 1), x) + 12*b^2*c^2*d*i ntegrate(1/16*x^4*arctan(c*x^2)^2/(c^2*x^4 + 1), x) + 4*b^2*c^2*e*integrate (1/16*x^5*log(c^2*x^4 + 1)/(c^2*x^4 + 1), x) + b^2*c^2*d*integrate(1/16*x^4 *log(c^2*x^4 + 1)^2/(c^2*x^4 + 1), x) + 8*b^2*c^2*d*integrate(1/16*x^4*log(c^2*x^4 + 1)/(c^2*x^4 + 1), x) + 1/2*a^2*x^2*e + 1/8*b^2*arctan(c*x^2)^3*e/c - 8*b^2*c*e*integrate(1/16*x^3*arctan(c*x^2)/(c^2*x^4 + 1), x) - 16*b^2*c *d*integrate(1/16*x^2*arctan(c*x^2)/(c^2*x^4 + 1), x) - 1/2*(c*(2*sqrt(2)*a rctan(1/2*sqrt(2)*(2*c*x + sqrt(2)*sqrt(c))/sqrt(c))/c^(3/2) + 2*sqrt(2)*ar ctan(1/2*sqrt(2)*(2*c*x - sqrt(2)*sqrt(c))/sqrt(c))/c^(3/2) - sqrt(2)*log(c *x^2 + sqrt(2)*sqrt(c)*x + 1)/c^(3/2) + sqrt(2)*log(c*x^2 - sqrt(2)*sqrt(c) *x + 1)/c^(3/2)) - 4*x*arctan(c*x^2))*a*b*d + a^2*d*x + b^2*e*integrate(1/1 6*x*log(c^2*x^4 + 1)^2/(c^2*x^4 + 1), x) + 12*b^2*d*integrate(1/16*arctan(c *x^2)^2/(c^2*x^4 + 1), x) + b^2*d*integrate(1/16*log(c^2*x^4 + 1)^2/(c^2*x^4 + 1), x) + 1/8*(b^2*x^2*e + 2*b^2*d*x)*arctan(c*x^2)^2 + 1/2*(2*c*x^2*arc`

$\tan(c*x^2) - \log(c^2*x^4 + 1)*a*b*e/c - 1/32*(b^2*x^2*e + 2*b^2*d*x)*\log(c^2*x^4 + 1)^2$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^2))^2,x, algorithm="fricas")`

[Out] `integral(a^2*x*e + a^2*d + (b^2*x*e + b^2*d)*arctan(c*x^2)^2 + 2*(a*b*x*e + a*b*d)*arctan(c*x^2), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (a + b \operatorname{atan}(cx^2))^2 (d + ex) \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*atan(c*x**2))**2,x)`

[Out] `Integral((a + b*atan(c*x**2))**2*(d + e*x), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^2))^2,x, algorithm="giac")`

[Out] `integrate((e*x + d)*(b*arctan(c*x^2) + a)^2, x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int (a + b \operatorname{atan}(cx^2))^2 (d + ex) \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^2))^2*(d + e*x),x)`

[Out] `int((a + b*atan(c*x^2))^2*(d + e*x), x)`

3.26 $\int \frac{(a+b\text{ArcTan}(cx^2))^2}{d+ex} dx$

Optimal. Leaf size=23

$$\text{Int}\left(\frac{(a+b\text{ArcTan}(cx^2))^2}{d+ex}, x\right)$$

[Out] Unintegrable((a+b*arctan(c*x^2))^2/(e*x+d),x)

Rubi [A]

time = 0.02, antiderivative size = 0, normalized size of antiderivative = 0.00, number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{(a+b\text{ArcTan}(cx^2))^2}{d+ex} dx$$

Verification is not applicable to the result.

[In] Int[(a + b*ArcTan[c*x^2])^2/(d + e*x),x]

[Out] Defer[Int][(a + b*ArcTan[c*x^2])^2/(d + e*x), x]

Rubi steps

$$\begin{aligned} \int \frac{(a+b\tan^{-1}(cx^2))^2}{d+ex} dx &= \int \left(\frac{a^2}{d+ex} + \frac{2ab\tan^{-1}(cx^2)}{d+ex} + \frac{b^2\tan^{-1}(cx^2)^2}{d+ex} \right) dx \\ &= \frac{a^2 \log(d+ex)}{e} + (2ab) \int \frac{\tan^{-1}(cx^2)}{d+ex} dx + b^2 \int \frac{\tan^{-1}(cx^2)^2}{d+ex} dx \end{aligned}$$

Mathematica [A]

time = 47.02, size = 0, normalized size = 0.00

$$\int \frac{(a+b\text{ArcTan}(cx^2))^2}{d+ex} dx$$

Verification is not applicable to the result.

[In] Integrate[(a + b*ArcTan[c*x^2])^2/(d + e*x),x]

[Out] Integrate[(a + b*ArcTan[c*x^2])^2/(d + e*x), x]

Maple [A]

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{(a+b\arctan(cx^2))^2}{ex+d} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int ((a+b\arctan(cx^2))^2/(ex+d), x)$
 [Out] $\int ((a+b\arctan(cx^2))^2/(ex+d), x)$

Maxima [A]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b\arctan(cx^2))^2/(ex+d), x, \text{algorithm}=\text{"maxima"})$
 [Out] $a^2e^{-1}\log(ex + d) + \text{integrate}((b^2\arctan(cx^2)^2 + 2ab\arctan(cx^2))/x, x)$

Fricas [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b\arctan(cx^2))^2/(ex+d), x, \text{algorithm}=\text{"fricas"})$
 [Out] $\text{integral}((b^2\arctan(cx^2)^2 + 2ab\arctan(cx^2) + a^2)/(x^2e + d), x)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b\arctan(cx^2))^2/(ex+d), x)$
 [Out] Timed out

Giac [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b\arctan(cx^2))^2/(ex+d), x, \text{algorithm}=\text{"giac"})$
 [Out] $\text{integrate}((b\arctan(cx^2) + a)^2/(ex + d), x)$

Mupad [A]

time = 0.00, size = -1, normalized size = -0.04

$$\int \frac{(a + b \operatorname{atan}(c x^2))^2}{d + e x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^2))^2/(d + e*x),x)`[Out] `int((a + b*atan(c*x^2))^2/(d + e*x), x)`

$$3.27 \quad \int \frac{(a+b\text{ArcTan}(cx^2))^2}{(d+ex)^2} dx$$

Optimal. Leaf size=23

$$\text{Int}\left(\frac{(a+b\text{ArcTan}(cx^2))^2}{(d+ex)^2}, x\right)$$

[Out] Unintegrable((a+b*arctan(c*x^2))^2/(e*x+d)^2, x)

Rubi [A]

time = 0.02, antiderivative size = 0, normalized size of antiderivative = 0.00, number of steps used = 0, number of rules used = 0, integrand size = 0, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.000, Rules used = {}

$$\int \frac{(a+b\text{ArcTan}(cx^2))^2}{(d+ex)^2} dx$$

Verification is not applicable to the result.

[In] Int[(a + b*ArcTan[c*x^2])^2/(d + e*x)^2, x]

[Out] Defer[Int][(a + b*ArcTan[c*x^2])^2/(d + e*x)^2, x]

Rubi steps

$$\begin{aligned}
\int \frac{(a + b \tan^{-1}(cx^2))^2}{(d + ex)^2} dx &= \int \left(\frac{a^2}{(d + ex)^2} + \frac{2ab \tan^{-1}(cx^2)}{(d + ex)^2} + \frac{b^2 \tan^{-1}(cx^2)^2}{(d + ex)^2} \right) dx \\
&= -\frac{a^2}{e(d + ex)} + (2ab) \int \frac{\tan^{-1}(cx^2)}{(d + ex)^2} dx + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \frac{(2ab) \int \frac{2cx}{(d+ex)(1+c^2x^4)} dx}{e} \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \frac{(4abc) \int \frac{x}{(d+ex)(1+c^2x^4)} dx}{e} \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \frac{(4abc) \int \left(-\frac{de^3}{(c^2d^4+e^4)(d+ex)} \right) dx}{e} \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} + b^2 \int \frac{\tan^{-1}(cx^2)^2}{(d + ex)^2} dx + \\
&= -\frac{a^2}{e(d + ex)} + \frac{2abc^2d^3 \tan^{-1}(cx^2)}{e(c^2d^4 + e^4)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} - \frac{4abcde \log(d + ex)}{c^2d^4 + e^4} - \\
&= -\frac{a^2}{e(d + ex)} + \frac{2abc^2d^3 \tan^{-1}(cx^2)}{e(c^2d^4 + e^4)} - \frac{2ab \tan^{-1}(cx^2)}{e(d + ex)} + \frac{\sqrt{2}ab\sqrt{c}(cd^2 - e^2)\tan^{-1}(cx^2)}{c^2d^4} -
\end{aligned}$$

Mathematica [A]

time = 44.47, size = 0, normalized size = 0.00

$$\int \frac{(a + b \operatorname{ArcTan}(cx^2))^2}{(d + ex)^2} dx$$

Verification is not applicable to the result.

[In] $\text{Integrate}[(a + b \arctan(c x^2))^2 / (d + e x)^2, x]$
[Out] $\text{Integrate}[(a + b \arctan(c x^2))^2 / (d + e x)^2, x]$

Maple [A]

time = 0.22, size = 0, normalized size = 0.00

$$\int \frac{(a + b \arctan(cx^2))^2}{(ex + d)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((a+b \arctan(c x^2))^2 / (e x + d)^2, x)$
[Out] $\text{int}((a+b \arctan(c x^2))^2 / (e x + d)^2, x)$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b \arctan(c x^2))^2 / (e x + d)^2, x, \text{algorithm}=\text{"maxima"})$
[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b \arctan(c x^2))^2 / (e x + d)^2, x, \text{algorithm}=\text{"fricas"})$
[Out] $\text{integral}((b^2 \arctan(c x^2)^2 + 2 a b \arctan(c x^2) + a^2) / (x^2 e^2 + 2 d x e + d^2), x)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}((a+b \arctan(c x^2))^2 / (e x + d)^2, x)$
[Out] Timed out

Giac [A]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^2))^2/(e*x+d)^2,x, algorithm="giac")`

[Out] `integrate((b*arctan(c*x^2) + a)^2/(e*x + d)^2, x)`

Mupad [A]

time = 0.00, size = -1, normalized size = -0.04

$$\int \frac{(a + b \operatorname{atan}(c x^2))^2}{(d + e x)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^2))^2/(d + e*x)^2,x)`

[Out] `int((a + b*atan(c*x^2))^2/(d + e*x)^2, x)`

$$3.28 \quad \int (d + ex)^2 (a + b \operatorname{ArcTan}(cx^3)) dx$$

Optimal. Leaf size=315

$$-\frac{bde \operatorname{ArcTan}(\sqrt[3]{c} x)}{c^{2/3}} - \frac{bd^3 \operatorname{ArcTan}(cx^3)}{3e} + \frac{(d + ex)^3 (a + b \operatorname{ArcTan}(cx^3))}{3e} + \frac{bde \operatorname{ArcTan}(\sqrt{3} - 2\sqrt[3]{c} x)}{2c^{2/3}} - \frac{bde \operatorname{ArcTan}(\sqrt{3} + 2\sqrt[3]{c} x)}{2c^{2/3}}$$

[Out]
$$\begin{aligned} & -b*d*e*\arctan(c^{(1/3)*x})/c^{(2/3)} - 1/3*b*d^3*\arctan(c*x^3)/e + 1/3*(e*x+d)^3*(a \\ & + b*\arctan(c*x^3))/e - 1/2*b*d^2*e*\arctan(2*c^{(1/3)*x} - 3^{(1/2)})/c^{(2/3)} - 1/2*b*d^2*e \\ & * \arctan(2*c^{(1/3)*x} + 3^{(1/2)})/c^{(2/3)} + 1/2*b*d^2*ln(1 + c^{(2/3)*x^2})/c^{(1/3)} - 1/ \\ & 4*b*d^2*2*ln(1 - c^{(2/3)*x^2} + c^{(4/3)*x^4})/c^{(1/3)} - 1/6*b^2*e^2*2*ln(c^2*x^6 + 1)/c^{(1/2)} \\ & *b*d^2*2*\arctan(1/3*(1 - 2*c^{(2/3)*x^2})*3^{(1/2)})*3^{(1/2)}/c^{(1/3)} - 1/4*b*d^2*e*ln(1 \\ & + c^{(2/3)*x^2} - c^{(1/3)*x*3^{(1/2)}})*3^{(1/2)}/c^{(2/3)} + 1/4*b*d^2*e*ln(1 + c^{(2/3)*x^2} + \\ & c^{(1/3)*x*3^{(1/2)}})*3^{(1/2)}/c^{(2/3)} \end{aligned}$$

Rubi [A]

time = 0.50, antiderivative size = 315, normalized size of antiderivative = 1.00, number of steps used = 24, number of rules used = 15, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {4980, 1845, 281, 298, 31, 648, 631, 210, 642, 301, 632, 209, 1483, 649, 266}

$$\frac{(d + ex)^3 (a + b \operatorname{ArcTan}(cx^3))}{3e} + \frac{\sqrt{3} bd^2 \operatorname{ArcTan}\left(\frac{c^{2/3} x^2}{\sqrt{3}}\right)}{2\sqrt{c}} - \frac{bde \operatorname{ArcTan}(\sqrt[3]{c} x)}{c^{2/3}} + \frac{bde \operatorname{ArcTan}\left(\sqrt{3} - 2\sqrt[3]{c} x\right)}{2c^{2/3}} - \frac{bde \operatorname{ArcTan}\left(2\sqrt[3]{c} x + \sqrt{3}\right)}{2c^{2/3}} - \frac{bd^2 \operatorname{ArcTan}(cx^3)}{3e} + \frac{bd^2 \log(c^{2/3} x^2 + 1)}{2\sqrt{c}} - \frac{bd^2 \log(c^{2/3} x^4 - c^{2/3} x^2 + 1)}{4\sqrt{c}} - \frac{\sqrt{3} bde \log(c^{2/3} x^2 - \sqrt{3} \sqrt[3]{c} x + 1)}{4c^{2/3}} + \frac{\sqrt{3} bde \log(c^{2/3} x^2 + \sqrt{3} \sqrt[3]{c} x + 1)}{4c^{2/3}} - \frac{be^2 \log(c^2 x^6 + 1)}{6c}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)^2*(a + b*\operatorname{ArcTan}[c*x^3]), x]$

[Out]
$$\begin{aligned} & -((b*d^2*e*\operatorname{ArcTan}[c^{(1/3)*x}])/c^{(2/3)}) - (b*d^3*\operatorname{ArcTan}[c*x^3])/(3*e) + ((d + \\ & e*x)^3*(a + b*\operatorname{ArcTan}[c*x^3]))/(3*e) + (b*d^2*e*\operatorname{ArcTan}[\operatorname{Sqrt}[3] - 2*c^{(1/3)*x}]) \\ & /(2*c^{(2/3)}) - (b*d^2*e*\operatorname{ArcTan}[\operatorname{Sqrt}[3] + 2*c^{(1/3)*x}])/(2*c^{(2/3)}) + (\operatorname{Sqrt}[3] \\ & *b*d^2*\operatorname{ArcTan}[(1 - 2*c^{(2/3)*x^2})/\operatorname{Sqrt}[3]])/(2*c^{(1/3)}) + (b*d^2*\operatorname{Log}[1 + c^{(2/3)*x^2}])/(2*c^{(1/3)}) - (\operatorname{Sqrt}[3]*b*d^2*e*\operatorname{Log}[1 - \operatorname{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}])/(4*c^{(2/3)}) + (\operatorname{Sqrt}[3]*b*d^2*e*\operatorname{Log}[1 + \operatorname{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}])/(4*c^{(2/3)}) - (b*d^2*\operatorname{Log}[1 - c^{(2/3)*x^2} + c^{(4/3)*x^4}])/(4*c^{(1/3)}) - \\ & (b^2 e^2 \operatorname{Log}[1 + c^2 x^6])/(6*c) \end{aligned}$$

Rule 31

$\operatorname{Int}[(a_ + b_)*x_{}^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[\operatorname{Log}[\operatorname{RemoveContent}[a + b*x, x]/b, x] /; \operatorname{FreeQ}[\{a, b\}, x]$

Rule 209

$\operatorname{Int}[(a_ + b_)*x_{}^{-2}^{-1}, x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[b, 2]))*\operatorname{ArcTan}[\operatorname{Rt}[b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&& \operatorname{PosQ}[a/b] \&& (\operatorname{GtQ}[a, 0] \mid\mid \operatorname{GtQ}[b, 0])$

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 266

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> Simp[Log[RemoveContent[a + b*x^n, x]]/(b*n), x] /; FreeQ[{a, b, m, n}, x] && EqQ[m, n - 1]
```

Rule 281

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 298

```
Int[(x_)/((a_) + (b_.)*(x_)^3), x_Symbol] :> Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]
```

Rule 301

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] :> Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k - 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 632

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
```

$x] \&& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 649

```
Int[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(
a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e
}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 1483

```
Int[(x_)^(m_)*(a_ + (c_)*(x_)^(n2_))^(p_)*(d_ + (e_)*(x_)^(n_))^(q
_), x_Symbol] :> Dist[1/n, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^n
], x] /; FreeQ[{a, c, d, e, m, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[Simplify
[m - n + 1], 0]
```

Rule 1845

```
Int[((Pq_)*((c_)*(x_)^(m_))/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> With[
{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]*x^(n/2)
)/(c^ii*(a + b*x^n))), {ii, 0, n/2 - 1}]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a,
b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n
```

Rule 4980

```
Int[((a_) + ArcTan[(c_)*(x_)^(n_)]*(b_))*(d_ + (e_)*(x_)^(m_)), x_Sy
mbol] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcTan[c*x^n])/(e*(m + 1))), x] - D
ist[b*c*(n/(e*(m + 1))), Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 + c^2*x^(2*n))
), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\int (d+ex)^2 (a+b\tan^{-1}(cx^3)) \, dx &= \int (a(d+ex)^2 + b(d+ex)^2 \tan^{-1}(cx^3)) \, dx \\
&= \frac{a(d+ex)^3}{3e} + b \int (d+ex)^2 \tan^{-1}(cx^3) \, dx \\
&= \frac{a(d+ex)^3}{3e} + b \int (d^2 \tan^{-1}(cx^3) + 2dex \tan^{-1}(cx^3) + e^2 x^2 \tan^{-1}(cx^3)) \, dx \\
&= \frac{a(d+ex)^3}{3e} + (bd^2) \int \tan^{-1}(cx^3) \, dx + (2bde) \int x \tan^{-1}(cx^3) \, dx + \\
&= \frac{a(d+ex)^3}{3e} + bd^2 x \tan^{-1}(cx^3) + bdex^2 \tan^{-1}(cx^3) + \frac{1}{3}be^2 x^3 \tan^{-1}(cx^3) \\
&= \frac{a(d+ex)^3}{3e} + bd^2 x \tan^{-1}(cx^3) + bdex^2 \tan^{-1}(cx^3) + \frac{1}{3}be^2 x^3 \tan^{-1}(cx^3) \\
&= \frac{a(d+ex)^3}{3e} - \frac{bde \tan^{-1}(\sqrt[3]{c}x)}{c^{2/3}} + bd^2 x \tan^{-1}(cx^3) + bdex^2 \tan^{-1}(cx^3) \\
&= \frac{a(d+ex)^3}{3e} - \frac{bde \tan^{-1}(\sqrt[3]{c}x)}{c^{2/3}} + bd^2 x \tan^{-1}(cx^3) + bdex^2 \tan^{-1}(cx^3) \\
&= \frac{a(d+ex)^3}{3e} - \frac{bde \tan^{-1}(\sqrt[3]{c}x)}{c^{2/3}} + bd^2 x \tan^{-1}(cx^3) + bdex^2 \tan^{-1}(cx^3)
\end{aligned}$$

Mathematica [A]

time = 102.00, size = 297, normalized size = 0.94

$$\frac{12acdx^2 + 12acd^2x^2 + 4acx^2d^2 - 12b\sqrt{c}dc\text{ArcTan}(\sqrt[3]{c}x) + 4bcx(3d^2 + 3dex + c^2x^2)\text{ArcTan}(cx^3) + 6b\sqrt{c}d(\sqrt{3}\sqrt[3]{c}x + e)\text{ArcTan}(\sqrt{3} - 2\sqrt[3]{c}x) + 6b\sqrt{c}d(\sqrt{3}\sqrt[3]{c}d - e)\text{ArcTan}(\sqrt{3} + 2\sqrt[3]{c}x) + 6b\sqrt{c}d^2\log(1 + c^{2/3}x^2) - 3b\sqrt{c}d(\sqrt[3]{c}d + \sqrt{3}e)\log(1 - \sqrt{3}\sqrt[3]{c}x + c^{2/3}x^2) - 3b\sqrt{c}d(\sqrt[3]{c}d - \sqrt{3}e)\log(1 + \sqrt{3}\sqrt[3]{c}x + c^{2/3}x^2) - 2b^2\log(1 + c^2x^2)}{12c}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d + e*x)^2*(a + b*ArcTan[c*x^3]), x]
[Out] (12*a*c*d^2*x + 12*a*c*d*e*x^2 + 4*a*c*e^2*x^3 - 12*b*c^(1/3)*d*e*ArcTan[c^(1/3)*x] + 4*b*c*x*(3*d^2 + 3*d*e*x + e^2*x^2)*ArcTan[c*x^3] + 6*b*c^(1/3)*d*(Sqrt[3]*c^(1/3)*d + e)*ArcTan[Sqrt[3] - 2*c^(1/3)*x] + 6*b*c^(1/3)*d*(Sqrt[3]*c^(1/3)*d - e)*ArcTan[Sqrt[3] + 2*c^(1/3)*x] + 6*b*c^(2/3)*d^2*Log[1 + c^(2/3)*x^2] - 3*b*c^(1/3)*d*(c^(1/3)*d + Sqrt[3]*e)*Log[1 - Sqrt[3]*c^(1/3)*x] - 3*b*c^(1/3)*d*(c^(1/3)*d + Sqrt[3]*e)*Log[1 - Sqrt[3]*c^(1/3)*x] - 2b^2*Log(1 + c^2*x^2)
```

$$\frac{1}{3}x + c^{(2/3)}x^2] - 3b*c^{(1/3)}*d*(c^{(1/3)}*d - \text{Sqrt}[3]*e)*\text{Log}[1 + \text{Sqrt}[3]]*c^{(1/3)}*x + c^{(2/3)}x^2] - 2b*e^2*\text{Log}[1 + c^2*x^6])/(12*c)$$

Maple [B] Leaf count of result is larger than twice the leaf count of optimal. 517 vs. $\frac{2(244)}{2} = 488$.

time = 9.23, size = 518, normalized size = 1.64

method	result
default	$\frac{(ex+d)^3 a}{3e} + \frac{b e^2 \arctan(cx^3) x^3}{3} + b e \arctan(cx^3) x^2 d + b \arctan(cx^3) x d^2 + \frac{b d^3 \arctan(cx^3)}{3e} - \frac{b e c \ln(x^2 - \sqrt{c^2 x^4 + 1})}{3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)^2*(a+b*arctan(c*x^3)),x,method=_RETURNVERBOSE)`

[Out] $1/3*(e*x+d)^3*a/e+1/3*b*e^2*\arctan(c*x^3)*x^3+b*e*\arctan(c*x^3)*x^2*d+b*arc\tan(c*x^3)*x*d^2+1/3*b*d^3*\arctan(c*x^3)/e-1/4*b*e*c*\ln(x^2-3^(1/2)*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*3^(1/2)*(1/c^2)^{(5/6)}*d-1/4*b*c*\ln(x^2-3^(1/2)*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*(1/c^2)^{(2/3)}*d^2-1/6*b*e^2*c*\ln(x^2-3^(1/2)*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})-1/2*b*e/c/(1/c^2)^{(1/6})*\arctan(2*x/(1/c^2)^{(1/6)}-3^(1/2)*d-1/2*b*c*(1/c^2)^{(2/3})*\arctan(2*x/(1/c^2)^{(1/6)}-3^(1/2))*3^(1/2)*(1/c^2)^{(5/6})*d-1/4*b*c*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6})*x+(1/c^2)^{(1/3})*3^(1/2)*(1/c^2)^{(5/6})*d-1/4*b*c*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6})*x+(1/c^2)^{(1/3})*(1/c^2)^{(2/3})*d^2-1/6*b*e^2*c*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6})*x+(1/c^2)^{(1/3})*3^(1/2)*(1/c^2)^{(5/6})*d-1/4*b*c*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6})*x+(1/c^2)^{(1/3})*(1/c^2)^{(2/3})*d^2-1/3*b*e*c*(1/c^2)^{(1/2})*\arctan(2*x/(1/c^2)^{(1/6)}+3^(1/2)*d-1/3*b*e*c*(1/c^2)^{(1/2})*\arctan(2*x/(1/c^2)^{(1/6)}+3^(1/2)*d^2+1/2*b*c*(1/c^2)^{(2/3})*\ln(x^2+(1/c^2)^{(1/3})*(1/c^2)^{(2/3})*d^2-1/6*b*e^2*c*\ln(x^2+(1/c^2)^{(1/3}))+1/3*b*e*c*(1/c^2)^{(1/2})*\arctan(x/(1/c^2)^{(1/6})*d-3*b*e/c/(1/c^2)^{(1/6})*\arctan(x/(1/c^2)^{(1/6})*d$

Maxima [A]

time = 0.57, size = 280, normalized size = 0.89

$$\frac{1}{3}ax^2 + adx^2e - \frac{1}{4}\left(\frac{2\sqrt{3}\arctan\left(\frac{\sqrt{3}(x^2-a^2)}{x^2}\right)}{c^{\frac{3}{2}}} + \frac{\log\left(c^{\frac{3}{2}}x^4 - c^{\frac{3}{2}}x^2 + 1\right)}{c^{\frac{3}{2}}} - \frac{2\log\left(\frac{dx+1}{dx-1}\right)}{c^{\frac{3}{2}}}\right) - 4x\arctan(cx^3) + \frac{1}{4}\left(4x^2\arctan(cx^3) + \left(\frac{\sqrt{3}\log\left(c^{\frac{3}{2}}x^2 + \sqrt{3}c^{\frac{1}{2}}x + 1\right)}{c^{\frac{3}{2}}} - \frac{\sqrt{3}\log\left(c^{\frac{3}{2}}x^2 - \sqrt{3}c^{\frac{1}{2}}x + 1\right)}{c^{\frac{3}{2}}} - \frac{4\arctan\left(cx^2\right)}{c^{\frac{3}{2}}} - \frac{2\arctan\left(\frac{ia^2x+\sqrt{3}a^2}{x^2}\right)}{c^{\frac{3}{2}}} - \frac{2\arctan\left(\frac{ia^2x-\sqrt{3}a^2}{x^2}\right)}{c^{\frac{3}{2}}}\right)\right) + \frac{(2cx^2\arctan(cx^3) - \log(c^2x^4 + 1))de^2}{6c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x^3)),x, algorithm="maxima")`

[Out] $1/3*a*x^3*e^2 + a*d*x^2*e - 1/4*(c*(2*\text{sqrt}(3)*\arctan(1/3*\text{sqrt}(3)*(2*c^{(4/3)}*x^2 - c^{(2/3)})/c^{(2/3)})/c^{(4/3)} + \log(c^{(4/3)}*x^4 - c^{(2/3)}*x^2 + 1)/c^{(4/3)} - 2*\log((c^{(2/3)}*x^2 + 1)/c^{(2/3)})/c^{(4/3)}) - 4*x*\arctan(c*x^3)*b*d^2 + a*d^2*x + 1/4*(4*x^2*\arctan(c*x^3) + c*(\text{sqrt}(3)*\log(c^{(2/3)}*x^2 + \text{sqrt}(3)*c^{(1/3)}*x + 1)/c^{(5/3)} - \text{sqrt}(3)*\log(c^{(2/3)}*x^2 - \text{sqrt}(3)*c^{(1/3)}*x + 1)/c^{(5/3)})$

$$\begin{aligned} & \sim (5/3) - 4 \operatorname{arctan}(c^{(1/3)}x)/c^{(5/3)} - 2 \operatorname{arctan}((2*c^{(2/3)}x + \sqrt{3})c^{(1/3)})/c^{(5/3)} - 2 \operatorname{arctan}((2*c^{(2/3)}x - \sqrt{3})c^{(1/3)})/c^{(5/3)}) * b*d*e + 1/6 * (2*c*x^3 * \operatorname{arctan}(c*x^3) - \log(c^2*x^6 + 1)) * b*e^2/c \end{aligned}$$

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x^3)),x, algorithm="fricas")`

[Out] Timed out

Sympy [A]

time = 28.54, size = 151, normalized size = 0.48

$$ad^2x + ade^2x^3 + \frac{ae^2x^3}{3} - 3bcd^2 \operatorname{RootSum}(216t^3c^4 + 1, (t \mapsto t \log(36t^2c^2 + x^2))) - 3bcde \operatorname{RootSum}(46656t^8c^{10} + 1, (t \mapsto t \log(7776t^5c^8 + x))) + bd^2x \operatorname{atan}(cx^3) + bdex^2 \operatorname{atan}(cx^3) + be^2 \left(\begin{cases} 0 & \text{for } c = 0 \\ \frac{x^2 \operatorname{atan}(cx^3)}{3} - \frac{\log(c^2x^6 + 1)}{6c} & \text{otherwise} \end{cases} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)**2*(a+b*atan(c*x**3)),x)`

[Out] $a*d^{**2}*x + a*d*e*x**2 + a*e**2*x**3/3 - 3*b*c*d**2*\operatorname{RootSum}(216*_t^{**3}*c^{**4} + 1, \operatorname{Lambda}(_t, _t*\log(36*_t^{**2}*c^{**2} + x^{**2}))) - 3*b*c*d*e*\operatorname{RootSum}(46656*_t^{**6}*c^{**10} + 1, \operatorname{Lambda}(_t, _t*\log(7776*_t^{**5}*c^{**8} + x))) + b*d^{**2}*x*\operatorname{atan}(c*x^{**3}) + b*d*e*x**2*\operatorname{atan}(c*x^{**3}) + b*e**2* \operatorname{Piecewise}((0, \operatorname{Eq}(c, 0)), (x^{**3}*\operatorname{atan}(c*x^{**3})/3 - \log(c^{**2}*x^{**6} + 1)/(6*c), \operatorname{True}))$

Giac [A]

time = 9.91, size = 318, normalized size = 1.01

$$\begin{aligned} & \frac{1}{3}bc^2x^2\operatorname{atan}(cx^2) + \frac{1}{3}ae^2x^2 + bde^2x^2\operatorname{atan}(cx^2) + ade^2x^2 + bdf^2x^2\operatorname{atan}(cx^2) + ad^2x - \frac{bde\operatorname{atan}(x)^3}{|c|^3} + \frac{(\sqrt{3}bdc^2|x|^3 - bde|x|^3)\operatorname{atan}\left(\left(2x + \frac{\sqrt{3}c}{|c|}\right)|x|^3\right)}{2c^2} - \frac{(\sqrt{3}bdc^2|x|^3 + bde|x|^3)\operatorname{atan}\left(\left(2x - \frac{\sqrt{3}c}{|c|}\right)|x|^3\right)}{2c^2} + \frac{(3\sqrt{3}bdc^2|x|^3 - 3bdc^2|x|^3 + 2bc^2)\log\left(x^2 - \frac{\sqrt{3}c^2}{|c|^2} + \frac{1}{4c^2}\right)}{12c^2} - \frac{(3\sqrt{3}bdc^2|x|^3 + 3bdc^2|x|^3 + 2bc^2)\log\left(x^2 - \frac{\sqrt{3}c^2}{|c|^2} + \frac{1}{4c^2}\right)}{12c^2} + \frac{(3bdc^2|x|^3 - bce^2)\log\left(x^2 + \frac{1}{4c^2}\right)}{6c^2} \end{aligned}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)^2*(a+b*arctan(c*x^3)),x, algorithm="giac")`

[Out] $1/3*b^2e^2*x^3* \operatorname{arctan}(c*x^3) + 1/3*a*e^2*x^3 + b*d^2e*x^2* \operatorname{arctan}(c*x^3) + a*d^2*x^2 + b*d^2*x*\operatorname{arctan}(c*x^3) + a*d^2*x - b*c*d*e*\operatorname{arctan}(x*abs(c)^{(1/3)})/a*bs(c)^{(5/3)} + 1/2*(\sqrt{3})*b*c*d^2*abs(c)^{(2/3)} - b*c*d*e*abs(c)^{(1/3})*\operatorname{atan}((2*x + \sqrt{3}/abs(c)^{(1/3}))*abs(c)^{(1/3)})/c^2 - 1/2*(\sqrt{3})*b*c*d^2*a*bs(c)^{(2/3)} + b*c*d*e*abs(c)^{(1/3})*\operatorname{arctan}((2*x - \sqrt{3}/abs(c)^{(1/3}))*abs(c)^{(1/3)})/c^2 + 1/12*(3*\sqrt{3})*b*c*d^2*abs(c)^{(1/3)} - 3*b*c*d^2*abs(c)^{(2/3)} - 2*b*c*e^2)*\log(x^2 + \sqrt{3}*\operatorname{abs}(c)^{(1/3)} + 1/\operatorname{abs}(c)^{(2/3)})/c^2 - 1/12*(3*\sqrt{3})*b*c*d^2*abs(c)^{(1/3)} + 3*b*c*d^2*abs(c)^{(2/3)} + 2*b*c*e^2)*\log(x^2 - \sqrt{3}*\operatorname{abs}(c)^{(1/3)} + 1/\operatorname{abs}(c)^{(2/3)})/c^2 + 1/6*(3*b*c*d^2*abs(c)^{(2/3)} - b*c*e^2)*\log(x^2 + 1/\operatorname{abs}(c)^{(2/3)})/c^2$

Mupad [B]

time = 0.52, size = 988, normalized size = 3.14

Verification of antiderivative is not currently implemented for this CAS.

[In] $\int ((a + b \cdot \text{atan}(c \cdot x^3)) \cdot (d + e \cdot x)^2, x)$

[Out]
$$\begin{aligned} & \text{atan}(c \cdot x^3) \cdot ((b \cdot e^2 \cdot x^3)/3 + b \cdot d^2 \cdot x + b \cdot d \cdot e \cdot x^2) + \text{symsum}(\log(x \cdot (6 \cdot b^5 \cdot c^7 \\ & \cdot d^2 \cdot e^8 - 162 \cdot b^5 \cdot c^9 \cdot d^8 \cdot e^2)) + \text{root}(46656 \cdot a^6 \cdot c^6 + 46656 \cdot a^5 \cdot b \cdot c^5 \cdot e^2 \\ & + 19440 \cdot a^4 \cdot b^2 \cdot c^4 \cdot e^4 + 4320 \cdot a^3 \cdot b^3 \cdot c^3 \cdot e^6 - 11664 \cdot a^3 \cdot b^3 \cdot c^5 \cdot d^6 + 20 \\ & 412 \cdot a^2 \cdot b^4 \cdot c^4 \cdot d^6 \cdot e^2 + 540 \cdot a^2 \cdot b^4 \cdot c^2 \cdot e^8 - 972 \cdot a \cdot b^5 \cdot c^3 \cdot d^6 \cdot e^4 + 36 \cdot \\ & a \cdot b^5 \cdot c \cdot e^{10} - 54 \cdot b^6 \cdot c^2 \cdot d^6 \cdot e^6 + 729 \cdot b^6 \cdot c^4 \cdot d^{12} + b^6 \cdot e^{12}, a, k) \cdot (x \cdot (\\ & 486 \cdot b^4 \cdot c^{10} \cdot d^8 + 90 \cdot b^4 \cdot c^8 \cdot d^2 \cdot e^6) - \text{root}(46656 \cdot a^6 \cdot c^6 + 46656 \cdot a^5 \cdot b \cdot c \\ & ^5 \cdot e^2 + 19440 \cdot a^4 \cdot b^2 \cdot c^4 \cdot e^4 + 4320 \cdot a^3 \cdot b^3 \cdot c^3 \cdot e^6 - 11664 \cdot a^3 \cdot b^3 \cdot c^5 \cdot d \\ & ^6 + 20412 \cdot a^2 \cdot b^4 \cdot c^4 \cdot d^6 \cdot e^2 + 540 \cdot a^2 \cdot b^4 \cdot c^2 \cdot e^8 - 972 \cdot a \cdot b^5 \cdot c^3 \cdot d^6 \cdot e^4 \\ & + 36 \cdot a \cdot b^5 \cdot c \cdot e^{10} - 54 \cdot b^6 \cdot c^2 \cdot d^6 \cdot e^6 + 729 \cdot b^6 \cdot c^4 \cdot d^{12} + b^6 \cdot e^{12}, a, k) \cdot (\text{root}(46656 \cdot a^6 \cdot c^6 + 46656 \cdot a^5 \cdot b \cdot c^5 \cdot e^2 + 19440 \cdot a^4 \cdot b^2 \cdot c^4 \cdot e^4 + 4320 \\ & \cdot a^3 \cdot b^3 \cdot c^3 \cdot e^6 - 11664 \cdot a^3 \cdot b^3 \cdot c^5 \cdot d^6 + 20412 \cdot a^2 \cdot b^4 \cdot c^4 \cdot d^6 \cdot e^2 + 540 \cdot \\ & a^2 \cdot b^4 \cdot c^2 \cdot e^8 - 972 \cdot a \cdot b^5 \cdot c^3 \cdot d^6 \cdot e^4 + 36 \cdot a \cdot b^5 \cdot c \cdot e^{10} - 54 \cdot b^6 \cdot c^2 \cdot d^6 \cdot \\ & e^6 + 729 \cdot b^6 \cdot c^4 \cdot d^{12} + b^6 \cdot e^{12}, a, k) \cdot (3888 \cdot b^2 \cdot c^{10} \cdot d^3 \cdot e + 3888 \cdot \text{root}(4 \\ & 6656 \cdot a^6 \cdot c^6 + 46656 \cdot a^5 \cdot b \cdot c^5 \cdot e^2 + 19440 \cdot a^4 \cdot b^2 \cdot c^4 \cdot e^4 + 4320 \cdot a^3 \cdot b^3 \cdot c \\ & ^3 \cdot e^6 - 11664 \cdot a^3 \cdot b^3 \cdot c^5 \cdot d^6 + 20412 \cdot a^2 \cdot b^4 \cdot c^4 \cdot d^6 \cdot e^2 + 540 \cdot a^2 \cdot b^4 \cdot c^2 \cdot e^8 \\ & - 972 \cdot a \cdot b^5 \cdot c^3 \cdot d^6 \cdot e^4 + 36 \cdot a \cdot b^5 \cdot c \cdot e^{10} - 54 \cdot b^6 \cdot c^2 \cdot d^6 \cdot e^6 + 729 \cdot \\ & b^6 \cdot c^4 \cdot d^{12} + b^6 \cdot e^{12}, a, k) \cdot b \cdot c^{11} \cdot d^2 \cdot x + 648 \cdot b^2 \cdot c^{10} \cdot d^2 \cdot e^2 \cdot x) + 972 \\ & \cdot b^3 \cdot c^9 \cdot d^3 \cdot e^3 - 324 \cdot b^3 \cdot c^9 \cdot d^2 \cdot e^4 \cdot x) - 243 \cdot b^5 \cdot c^9 \cdot d^9 \cdot e + 9 \cdot b^5 \cdot c^7 \cdot \\ & d^3 \cdot e^7) \cdot \text{root}(46656 \cdot a^6 \cdot c^6 + 46656 \cdot a^5 \cdot b \cdot c^5 \cdot e^2 + 19440 \cdot a^4 \cdot b^2 \cdot c^4 \cdot e^4 + \\ & 4320 \cdot a^3 \cdot b^3 \cdot c^3 \cdot e^6 - 11664 \cdot a^3 \cdot b^3 \cdot c^5 \cdot d^6 + 20412 \cdot a^2 \cdot b^4 \cdot c^4 \cdot d^6 \cdot e^2 + \\ & 540 \cdot a^2 \cdot b^4 \cdot c^2 \cdot e^8 - 972 \cdot a \cdot b^5 \cdot c^3 \cdot d^6 \cdot e^4 + 36 \cdot a \cdot b^5 \cdot c \cdot e^{10} - 54 \cdot b^6 \cdot c^2 \\ & \cdot d^6 \cdot e^6 + 729 \cdot b^6 \cdot c^4 \cdot d^{12} + b^6 \cdot e^{12}, a, k), k, 1, 6) + (a \cdot e^{2 \cdot x^3})/3 + a \\ & \cdot d^2 \cdot x + a \cdot d \cdot e \cdot x^2 \end{aligned}$$

$$3.29 \quad \int (d + ex) (a + b \operatorname{ArcTan}(cx^3)) \, dx$$

Optimal. Leaf size=285

$$-\frac{be \operatorname{ArcTan}(\sqrt[3]{c} x)}{2 c^{2/3}} - \frac{bd^2 \operatorname{ArcTan}(cx^3)}{2e} + \frac{(d + ex)^2 (a + b \operatorname{ArcTan}(cx^3))}{2e} + \frac{be \operatorname{ArcTan}(\sqrt{3} - 2\sqrt[3]{c} x)}{4 c^{2/3}} - \frac{be \operatorname{ArcTan}(\sqrt{3} + 2\sqrt[3]{c} x)}{4 c^{2/3}}$$

[Out] $-1/2*b*e*\arctan(c^{(1/3)*x})/c^{(2/3)} - 1/2*b*d^2*\arctan(c*x^3)/e + 1/2*(e*x+d)^2*(a+b*\arctan(c*x^3))/e - 1/4*b*e*\arctan(2*c^{(1/3)*x}-3^{(1/2)})/c^{(2/3)} - 1/4*b*e*a*\arctan(2*c^{(1/3)*x}+3^{(1/2)})/c^{(2/3)} + 1/2*b*d*\ln(1+c^{(2/3)*x^2})/c^{(1/3)} - 1/4*b*d*\ln(1-c^{(2/3)*x^2}+c^{(4/3)*x^4})/c^{(1/3)} + 1/2*b*d*\arctan(1/3*(1-2*c^{(2/3)*x^2})*3^{(1/2)})*3^{(1/2)}/c^{(1/3)} - 1/8*b*e*\ln(1+c^{(2/3)*x^2}-c^{(1/3)*x*3^{(1/2)}})*3^{(1/2)}/c^{(2/3)} + 1/8*b*e*\ln(1+c^{(2/3)*x^2}+c^{(1/3)*x*3^{(1/2)}})*3^{(1/2)}/c^{(2/3)}$

Rubi [A]

time = 0.40, antiderivative size = 285, normalized size of antiderivative = 1.00, number of steps used = 22, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {4980, 1845, 281, 209, 298, 31, 648, 631, 210, 642, 301, 632}

$$\frac{(d + ex)^2 (a + b \operatorname{ArcTan}(cx^3))}{2e} + \frac{\sqrt{3} bd \operatorname{ArcTan}\left(\frac{1+\sqrt[3]{c} x^2}{\sqrt{3}}\right)}{2\sqrt[3]{c}} - \frac{be \operatorname{ArcTan}(\sqrt[3]{c} x)}{2 c^{2/3}} + \frac{be \operatorname{ArcTan}(\sqrt{3} - 2\sqrt[3]{c} x)}{4 c^{2/3}} - \frac{be \operatorname{ArcTan}(\sqrt{3} + 2\sqrt[3]{c} x)}{4 c^{2/3}} - \frac{bd^2 \operatorname{ArcTan}(cx^3)}{2e} + \frac{bd \log(c^{2/3} x^2 + 1)}{2\sqrt[3]{c}} - \frac{bd \log(c^{4/3} x^4 - c^{2/3} x^2 + 1)}{4\sqrt[3]{c}} - \frac{\sqrt{3} be \log(c^{2/3} x^2 - \sqrt{3} \sqrt[3]{c} x + 1)}{8 c^{2/3}} + \frac{\sqrt{3} be \log(c^{2/3} x^2 + \sqrt{3} \sqrt[3]{c} x + 1)}{8 c^{2/3}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(d + e*x)*(a + b*\operatorname{ArcTan}[c*x^3]), x]$

[Out] $-1/2*(b*e*\operatorname{ArcTan}[c^{(1/3)*x}])/c^{(2/3)} - (b*d^2*\operatorname{ArcTan}[c*x^3])/(2*e) + ((d + e*x)^2*(a + b*\operatorname{ArcTan}[c*x^3]))/(2*e) + (b*e*\operatorname{ArcTan}[\operatorname{Sqrt}[3] - 2*c^{(1/3)*x}])/(4*c^{(2/3)}) - (b*e*\operatorname{ArcTan}[\operatorname{Sqrt}[3] + 2*c^{(1/3)*x}])/(4*c^{(2/3)}) + (\operatorname{Sqrt}[3]*b*d*\operatorname{ArcTan}[(1 - 2*c^{(2/3)*x^2})/\operatorname{Sqrt}[3]])/(2*c^{(1/3)}) + (b*d*\operatorname{Log}[1 + c^{(2/3)*x^2}]/(2*c^{(1/3)}) - (\operatorname{Sqrt}[3]*b*e*\operatorname{Log}[1 - \operatorname{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}])/(8*c^{(2/3)}) + (\operatorname{Sqrt}[3]*b*e*\operatorname{Log}[1 + \operatorname{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}])/(8*c^{(2/3)}) - (b*d*\operatorname{Log}[1 - c^{(2/3)*x^2} + c^{(4/3)*x^4}])/(4*c^{(1/3)})$

Rule 31

```
Int[((a_) + (b_)*(x_))^(-1), x_Symbol] :> Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]
```

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 210

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] :> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 281

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] :> With[{k = GCD[m + 1, n]}, Dist[1/k, Subst[Int[x^((m + 1)/k - 1)*(a + b*x^(n/k))^p, x], x, x^k], x] /; k != 1] /; FreeQ[{a, b, p}, x] && IGtQ[n, 0] && IntegerQ[m]
```

Rule 298

```
Int[(x_)/((a_) + (b_)*(x_)^3), x_Symbol] :> Dist[-(3*Rt[a, 3]*Rt[b, 3])^(-1), Int[1/(Rt[a, 3] + Rt[b, 3]*x), x], x] + Dist[1/(3*Rt[a, 3]*Rt[b, 3]), Int[(Rt[a, 3] + Rt[b, 3]*x)/(Rt[a, 3]^2 - Rt[a, 3]*Rt[b, 3]*x + Rt[b, 3]^2*x^2), x], x] /; FreeQ[{a, b}, x]
```

Rule 301

```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k - 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] :> Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> Simplify[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] :> D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1845

```
Int[((Pq_)*((c_.*(x_))^(m_.))/((a_) + (b_.*(x_)^(n_))), x_Symbol] :> With[
{v = Sum[(c*x)^(m + ii)*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]]*x^(n/2)
)/(c^ii*(a + b*x^n))), {ii, 0, n/2 - 1}]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b, c, m}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n
```

Rule 4980

```
Int[((a_.) + ArcTan[(c_.*(x_)^(n_))*(b_.)]*((d_) + (e_.*(x_))^(m_.)), x_Sy
mbol] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcTan[c*x^n])/(e*(m + 1))), x] - D
ist[b*c*(n/(e*(m + 1))), Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 + c^2*x^(2*n)))
], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rubi steps

$$\begin{aligned}
\int (d+ex) (a+b \tan^{-1}(cx^3)) dx &= \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} - \frac{b \int \frac{3cx^2(d+ex)^2}{1+c^2x^6} dx}{2e} \\
&= \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} - \frac{(3bc) \int \frac{x^2(d+ex)^2}{1+c^2x^6} dx}{2e} \\
&= \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} - \frac{(3bc) \int \left(\frac{d^2x^2}{1+c^2x^6} + \frac{2dex^3}{1+c^2x^6} + \frac{e^2x^4}{1+c^2x^6} \right) dx}{2e} \\
&= \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} - (3bcd) \int \frac{x^3}{1+c^2x^6} dx - \frac{(3bcd^2) \int \frac{x^2}{1+c^2x^6}}{2e} \\
&= \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} - \frac{1}{2} (3bcd) \text{Subst} \left(\int \frac{x}{1+c^2x^3} dx, x, x^2 \right) - \\
&= -\frac{be \tan^{-1}(\sqrt[3]{c} x)}{2c^{2/3}} - \frac{bd^2 \tan^{-1}(cx^3)}{2e} + \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} + \\
&= -\frac{be \tan^{-1}(\sqrt[3]{c} x)}{2c^{2/3}} - \frac{bd^2 \tan^{-1}(cx^3)}{2e} + \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} + \\
&= -\frac{be \tan^{-1}(\sqrt[3]{c} x)}{2c^{2/3}} - \frac{bd^2 \tan^{-1}(cx^3)}{2e} + \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} + \\
&= -\frac{be \tan^{-1}(\sqrt[3]{c} x)}{2c^{2/3}} - \frac{bd^2 \tan^{-1}(cx^3)}{2e} + \frac{(d+ex)^2 (a+b \tan^{-1}(cx^3))}{2e} +
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 310, normalized size = 1.09

$$adx + \frac{1}{2} uax^2 - \frac{be \text{ArcTan}(\sqrt[3]{c} x)}{2c^{2/3}} + bdx \text{ArcTan}(cx^3) + \frac{1}{2} bex^2 \text{ArcTan}(cx^3) + \frac{be \text{ArcTan}(\sqrt[3]{c} - 2\sqrt[3]{c} x)}{4c^{2/3}} - \frac{be \text{ArcTan}(\sqrt[3]{c} + 2\sqrt[3]{c} x)}{4c^{2/3}} - \frac{\sqrt[3]{b} \ln \log(1 + \sqrt[3]{c} \sqrt[3]{c} x + c^{2/3} x^2)}{8c^{2/3}} + \frac{\sqrt[3]{b} \ln \log(1 + \sqrt[3]{c} \sqrt[3]{c} x + c^{2/3} x^2)}{8c^{2/3}} - \frac{bd(-2\sqrt[3]{b} \text{ArcTan}(\sqrt[3]{c} - 2\sqrt[3]{c} x) - 2\sqrt[3]{b} \text{ArcTan}(\sqrt[3]{c} + 2\sqrt[3]{c} x) - 2 \log(1 + c^{2/3} x^2) + \log(1 + \sqrt[3]{c} \sqrt[3]{c} x + c^{2/3} x^2) + \log(1 + \sqrt[3]{c} \sqrt[3]{c} x + c^{2/3} x^2))}{4\sqrt[3]{c}}$$

Antiderivative was successfully verified.

```
[In] Integrate[(d + e*x)*(a + b*ArcTan[c*x^3]), x]
[Out] a*d*x + (a*e*x^2)/2 - (b*e*ArcTan[c^(1/3)*x])/(2*c^(2/3)) + b*d*x*ArcTan[c*x^3] + (b*e*x^2*ArcTan[c*x^3])/2 + (b*e*ArcTan[Sqrt[3] - 2*c^(1/3)*x])/(4*c^(2/3)) - (b*e*ArcTan[Sqrt[3] + 2*c^(1/3)*x])/(4*c^(2/3)) - (Sqrt[3]*b*e*Log[1 - Sqrt[3]*c^(1/3)*x + c^(2/3)*x^2])/(8*c^(2/3)) + (Sqrt[3]*b*e*Log[1 + Sqrt[3]*c^(1/3)*x + c^(2/3)*x^2])/(8*c^(2/3)) - (b*d*(-2*Sqrt[3]*ArcTan[Sqr[t[3] - 2*c^(1/3)*x] - 2*Sqrt[3]*ArcTan[Sqr[t[3] + 2*c^(1/3)*x]] - 2*Log[1 + c
```

$$\begin{aligned} & \sim (2/3)*x^2] + \text{Log}[1 - \text{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}] + \text{Log}[1 + \text{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}])/(4*c^{(1/3)}) \end{aligned}$$

Maple [A]

time = 0.25, size = 315, normalized size = 1.11

method	result
default	$a\left(\frac{1}{2}e x^2 + dx\right) + \frac{b \arctan(cx^3)x^2 e}{2} + b \arctan(cx^3) dx - \frac{b c^3 \ln\left(x^2 - \sqrt{3}\left(\frac{1}{c^2}\right)^{\frac{1}{6}}x + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)\left(\frac{1}{c^2}\right)^{\frac{5}{3}}d}{4} - \frac{b c \ln\left(x^2 + \sqrt{3}\left(\frac{1}{c^2}\right)^{\frac{1}{6}}x + \left(\frac{1}{c^2}\right)^{\frac{1}{3}}\right)\left(\frac{1}{c^2}\right)^{\frac{5}{3}}e}{4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((e*x+d)*(a+b*arctan(c*x^3)),x,method=_RETURNVERBOSE)`

[Out]

$$\begin{aligned} & a*(1/2*e*x^2+d*x)+1/2*b*arctan(c*x^3)*x^2+e+b*arctan(c*x^3)*d*x-1/4*b*c^3*x^2 \\ & -3^{(1/2)}*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*(1/c^2)^{(5/3)}*d-1/8*b*c*\ln(x^2 \\ & -3^{(1/2)}*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*3^{(1/2)}*(1/c^2)^{(5/6)}*e-1/2*b*c^3*(\\ & 1/c^2)^{(5/3)}*arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*3^{(1/2)}*d-1/4*b/c/(1/c^2)^{(1/6)}* \\ & arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*e+1/8*b*c*\ln(x^2+3^{(1/2)}*(1/c^2)^{(1/6)} \\ &)*x+(1/c^2)^{(1/3)})*3^{(1/2)}*(1/c^2)^{(5/6)}*e-1/4*b*c*\ln(x^2+3^{(1/2)}*(1/c^2)^{(1/6)}* \\ & x+(1/c^2)^{(1/3)})*(1/c^2)^{(2/3)}*d-1/4*b/c/(1/c^2)^{(1/6)}*arctan(2*x/(1/c^2)^{(1/6)}+3^{(1/2)} \\ &)*e+1/2*b*c*(1/c^2)^{(2/3)}*arctan(2*x/(1/c^2)^{(1/6)}+3^{(1/2)})*3^{(1/2)}*d+\ln(x^2+(1/c^2)^{(1/3)})-1/2*b/c*e/(1/c^2)^{(1/6)}* \\ & arctan(x/(1/c^2)^{(1/6)}) \end{aligned}$$
Maxima [A]

time = 0.47, size = 234, normalized size = 0.82

$$\frac{1}{2}ax^2e - \frac{1}{4}\left(c\left(\frac{2\sqrt{3}\arctan\left(\frac{\sqrt{3}(x^{\frac{2}{3}}x^{\frac{4}{3}}-c^{\frac{2}{3}})}{c^{\frac{3}{2}}}\right)}{c^{\frac{3}{2}}} + \frac{\log\left(c^{\frac{1}{2}}x^2 - c^{\frac{3}{2}}x^2 + 1\right)}{c^{\frac{3}{2}}} - \frac{2\log\left(\frac{x^{\frac{2}{3}}+1}{x^{\frac{1}{3}}}\right)}{c^{\frac{3}{2}}}\right) - 4x\arctan(cx^3)\right)bd + adx + \frac{1}{8}\left(4x^2\arctan(cx^3) + c\left(\frac{\sqrt{3}\log\left(c^{\frac{1}{2}}x^2 + \sqrt{3}c^{\frac{1}{2}}x + 1\right)}{c^{\frac{3}{2}}} - \frac{\sqrt{3}\log\left(c^{\frac{1}{2}}x^2 - \sqrt{3}c^{\frac{1}{2}}x + 1\right)}{c^{\frac{3}{2}}} - \frac{4\arctan\left(c^{\frac{1}{2}}x\right)}{c^{\frac{3}{2}}} - \frac{2\arctan\left(\frac{2c^{\frac{1}{2}}x + \sqrt{3}c^{\frac{1}{2}}}{c^{\frac{3}{2}}}\right)}{c^{\frac{3}{2}}} - \frac{2\arctan\left(\frac{2c^{\frac{1}{2}}x - \sqrt{3}c^{\frac{1}{2}}}{c^{\frac{3}{2}}}\right)}{c^{\frac{3}{2}}}\right)\right)be$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^3)),x, algorithm="maxima")`

[Out]

$$\begin{aligned} & 1/2*a*x^2*e - 1/4*(c*(2*sqrt(3)*arctan(1/3*sqrt(3)*(2*c^(4/3)*x^2 - c^(2/3)))/c^(2/3))/c^(4/3) + \log(c^(4/3)*x^4 - c^(2/3)*x^2 + 1)/c^(4/3) - 2*\log((c^(2/3)*x^2 + 1)/c^(2/3))/c^(4/3)) - 4*x*arctan(c*x^3)*b*d + a*d*x + 1/8*(4*x^2*arctan(c*x^3) + c*(sqrt(3)*log(c^(2/3)*x^2 + sqrt(3)*c^(1/3)*x + 1)/c^(5/3) - sqrt(3)*log(c^(2/3)*x^2 - sqrt(3)*c^(1/3)*x + 1)/c^(5/3) - 4*arctan(c^(1/3)*x)/c^(5/3) - 2*arctan((2*c^(2/3)*x + sqrt(3)*c^(1/3))/c^(1/3))/c^(5/3) - 2*arctan((2*c^(2/3)*x - sqrt(3)*c^(1/3))/c^(1/3))/c^(5/3)))*b*e \end{aligned}$$
Fricas [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^3)),x, algorithm="fricas")`

[Out] Exception raised: RuntimeError >> no explicit roots found

Sympy [A]

time = 12.69, size = 104, normalized size = 0.36

$$adx + \frac{aex^2}{2} - 3bcd \operatorname{RootSum}(216t^3c^4 + 1, (t \mapsto t \log(36t^2c^2 + x^2))) - \frac{3bce \operatorname{RootSum}(46656t^6c^{10} + 1, (t \mapsto t \log(7776t^5c^8 + x)))}{2} + bdx \operatorname{atan}(cx^3) + \frac{bex^2 \operatorname{atan}(cx^3)}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*atan(c*x**3)),x)`

[Out] $a*d*x + a*e*x**2/2 - 3*b*c*d*\operatorname{RootSum}(216*_t**3*c**4 + 1, \operatorname{Lambda}(_t, _t*\log(36*_t**2*c**2 + x**2))) - 3*b*c*e*\operatorname{RootSum}(46656*_t**6*c**10 + 1, \operatorname{Lambda}(_t, _t*\log(7776*_t**5*c**8 + x))/2 + b*d*x*\operatorname{atan}(c*x**3) + b*e*x**2*\operatorname{atan}(c*x**3)/2$

Giac [A]

time = 0.95, size = 250, normalized size = 0.88

$$\frac{1}{2}bcx^2\arctan(cx^3) + \frac{1}{2}acx^2 + bdx\arctan(cx^3) + adx + \frac{bd|c|^{\frac{3}{2}}\log\left(x^2 + \frac{1}{|c|^{\frac{1}{2}}}\right)}{2c} - \frac{bcr\arctan\left(x|c|^{\frac{1}{2}}\right)}{2|c|^{\frac{1}{2}}} + \frac{(2\sqrt{3}\operatorname{ber}|c|^{\frac{3}{2}} - bcr|c|^{\frac{1}{2}})\arctan\left(\left(2x + \frac{\sqrt{3}}{|c|^{\frac{1}{2}}}\right)|c|^{\frac{1}{2}}\right)}{4c^2} - \frac{(2\sqrt{3}\operatorname{ber}|c|^{\frac{3}{2}} + bcr|c|^{\frac{1}{2}})\arctan\left(\left(2x - \frac{\sqrt{3}}{|c|^{\frac{1}{2}}}\right)|c|^{\frac{1}{2}}\right)}{4c^2} + \frac{(\sqrt{3}\operatorname{ber}|c|^{\frac{3}{2}} - 2\operatorname{ber}|c|^{\frac{1}{2}})\log\left(x^2 + \frac{\sqrt{3}x}{|c|^{\frac{1}{2}}} + \frac{1}{|c|^{\frac{1}{2}}}\right)}{8cx} - \frac{(\sqrt{3}\operatorname{ber}|c|^{\frac{3}{2}} + 2\operatorname{ber}|c|^{\frac{1}{2}})\log\left(x^2 - \frac{\sqrt{3}x}{|c|^{\frac{1}{2}}} + \frac{1}{|c|^{\frac{1}{2}}}\right)}{8cx}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((e*x+d)*(a+b*arctan(c*x^3)),x, algorithm="giac")`

[Out] $1/2*b*e*x^2*\arctan(c*x^3) + 1/2*a*e*x^2 + b*d*x*\arctan(c*x^3) + a*d*x + 1/2*b*d*abs(c)^{(2/3)}*\log(x^2 + 1/abs(c)^{(2/3)})/c - 1/2*b*c*e*\arctan(x*abs(c)^{(1/3)}/abs(c)^{(5/3)} + 1/4*(2*sqrt(3)*b*c*d*abs(c)^{(2/3)} - b*c*e*abs(c)^{(1/3)})*\arctan((2*x + sqrt(3)/abs(c)^{(1/3})*abs(c)^{(1/3)})/c^2 - 1/4*(2*sqrt(3)*b*c*d*abs(c)^{(2/3)} + b*c*e*abs(c)^{(1/3)})*\arctan((2*x - sqrt(3)/abs(c)^{(1/3})*abs(c)^{(1/3)})/c^2 + 1/8*(sqrt(3)*b*c*e*abs(c)^{(1/3)} - 2*b*c*d*abs(c)^{(2/3)})*\log(x^2 + sqrt(3)*x/abs(c)^{(1/3)} + 1/abs(c)^{(2/3)})/c^2 - 1/8*(sqrt(3)*b*c*e*abs(c)^{(1/3)} + 2*b*c*d*abs(c)^{(2/3)})*\log(x^2 - sqrt(3)*x/abs(c)^{(1/3)} + 1/abs(c)^{(2/3)})/c^2$

Mupad [B]

time = 0.44, size = 485, normalized size = 1.70

$$\operatorname{atan}(c*x^3) \left(\frac{b^2 c^2 d^2 e^2}{x^2} + \left(\sum_{n=1}^{\infty} \left(\frac{(-1)^n (2096 a^6 c^4 - 1024 a^3 b^3 c^3 + 576 a^2 b^2 c^2 + 48 a b^5 c^4 + 64 b^6 c^2) (-1)^n (2096 a^6 c^4 - 1024 a^3 b^3 c^3 + 576 a^2 b^2 c^2 - 48 a b^5 c^4 + 64 b^6 c^2) n!}{(2096 a^6 c^4 - 1024 a^3 b^3 c^3 + 576 a^2 b^2 c^2 - 48 a b^5 c^4 + 64 b^6 c^2) n!} \right) \frac{2096 a^6 c^4 - 1024 a^3 b^3 c^3 + 576 a^2 b^2 c^2 - 48 a b^5 c^4 + 64 b^6 c^2}{x^{2n+2}} \right) + \frac{2096 a^6 c^4 - 1024 a^3 b^3 c^3 + 576 a^2 b^2 c^2 - 48 a b^5 c^4 + 64 b^6 c^2}{x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^3))*(d + e*x),x)`

[Out] $\operatorname{atan}(c*x^3)*(b*d*x + (b*e*x^2)/2) + \operatorname{symsum}(\log(-\operatorname{root}(4096*a^6*c^4 - 1024*a^3*b^3*c^3 + 576*a^2*b^2*c^2 + 48*a*b^5*c^4 + 64*b^6*c^2)*x^n, n)) + \frac{2096 a^6 c^4 - 1024 a^3 b^3 c^3 + 576 a^2 b^2 c^2 - 48 a b^5 c^4 + 64 b^6 c^2}{x^2}$

$$\begin{aligned} & 6 + b^6 \cdot e^6, a, k) * (\text{root}(4096*a^6*c^4 - 1024*a^3*b^3*c^3*d^3 + 576*a^2*b^4*c^2*d^2 - 48*a*b^5*c*d*e^4 + 64*b^6*c^2*d^6 + b^6 \cdot e^6, a, k) * (\text{root}(4096*a^6*c^4 - 1024*a^3*b^3*c^3*d^3 + 576*a^2*b^4*c^2*d^2 \cdot e^2 - 48*a*b^5*c*d*e^4 + 64*b^6*c^2*d^6 + b^6 \cdot e^6, a, k) * (1944*b^2*c^10*d*e - 486*b^2*c^10 \cdot e^2*x + 3888*\text{root}(4096*a^6*c^4 - 1024*a^3*b^3*c^3*d^3 + 576*a^2*b^4*c^2*d^2 \cdot e^2 - 48*a*b^5*c*d*e^4 + 64*b^6*c^2*d^6 + b^6 \cdot e^6, a, k) * b*c^11*d*x) - (243*b^3*c^9 \cdot e^3)/2 - 486*b^4*c^10*d^4*x) - (243*b^5*c^9*d^4 \cdot e)/2 - (243*b^5*c^9*d^3 \cdot e^2*x)/4) * \text{root}(4096*a^6*c^4 - 1024*a^3*b^3*c^3*d^3 + 576*a^2*b^4*c^2*d^2 \cdot e^2 - 48*a*b^5*c*d*e^4 + 64*b^6*c^2*d^6 + b^6 \cdot e^6, a, k), k, 1, 6) + a*d*x + (a \cdot e \cdot x^2)/2 \end{aligned}$$

3.30 $\int \frac{a+b\text{ArcTan}(cx^3)}{d+ex} dx$

Optimal. Leaf size=739

$$\frac{(a + b\text{ArcTan}(cx^3)) \log(d + ex)}{e} + \frac{bc \log\left(\frac{e\left(1-\sqrt[6]{-c^2}x\right)}{\sqrt[6]{-c^2}d+e}\right) \log(d + ex)}{\frac{2\sqrt{-c^2}}{e}} - \frac{bc \log\left(-\frac{e\left(1+\sqrt[6]{-c^2}x\right)}{\sqrt[6]{-c^2}d-e}\right) \log(d + ex)}{\frac{2\sqrt{-c^2}}{e}}$$

[Out] $(a+b*\arctan(c*x^3))*\ln(e*x+d)/e+1/2*b*c*\ln(e*(1-(-c^2)^(1/6)*x)/((-c^2)^(1/6)*d+e))*\ln(e*x+d)/e/(-c^2)^(1/2)-1/2*b*c*\ln(-e*(1+(-c^2)^(1/6)*x)/((-c^2)^(1/6)*d-e))*\ln(e*x+d)/e/(-c^2)^(1/2)+1/2*b*c*\ln(-e*((-1)^(1/3)+(-c^2)^(1/6)*x)/((-c^2)^(1/6)*d-(-1)^(1/3)*e))*\ln(e*x+d)/e/(-c^2)^(1/2)-1/2*b*c*\ln(-e*(-1)^(2/3)+(-c^2)^(1/6)*x)/((-c^2)^(1/6)*d-(-1)^(2/3)*e))*\ln(e*x+d)/e/(-c^2)^(1/2)+1/2*b*c*\ln((-1)^(2/3)*e*(1+(-1)^(1/3)*(-c^2)^(1/6)*x)/((-c^2)^(1/6)*d+(-1)^(1/3)*e))*\ln(e*x+d)/e/(-c^2)^(1/2)-1/2*b*c*\ln((-1)^(1/3)*e*(1+(-1)^(2/3)*(-c^2)^(1/6)*x)/((-c^2)^(1/6)*d+(-1)^(1/3)*e))*\ln(e*x+d)/e/(-c^2)^(1/2)-1/2*b*c*polylog(2, (-c^2)^(1/6)*(e*x+d)/((-c^2)^(1/6)*d-e))/e/(-c^2)^(1/2)+1/2*b*c*polylog(2, (-c^2)^(1/6)*(e*x+d)/((-c^2)^(1/6)*d+e))/e/(-c^2)^(1/2)+1/2*b*c*polylog(2, (-c^2)^(1/6)*(e*x+d)/((-c^2)^(1/6)*d-(-1)^(1/3)*e))/e/(-c^2)^(1/2)-1/2*b*c*polylog(2, (-c^2)^(1/6)*(e*x+d)/((-c^2)^(1/6)*d+(-1)^(1/3)*e))/e/(-c^2)^(1/2)-1/2*b*c*polylog(2, (-c^2)^(1/6)*(e*x+d)/((-c^2)^(1/6)*d-(-1)^(2/3)*e))/e/(-c^2)^(1/2)+1/2*b*c*polylog(2, (-c^2)^(1/6)*(e*x+d)/((-c^2)^(1/6)*d+(-1)^(2/3)*e))/e/(-c^2)^(1/2)$

Rubi [A]

time = 0.93, antiderivative size = 739, normalized size of antiderivative = 1.00, number of steps used = 25, number of rules used = 8, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.444$, Rules used = {4976, 281, 209, 2463, 266, 2441, 2440, 2438}

$$\frac{\log(d+ex)(a+b\text{ArcTan}(cx^3))}{e} + \frac{\text{bcln}_1\left(\frac{\sqrt[6]{-c^2}x}{\sqrt[6]{-c^2}d+e}\right)}{2\sqrt{-c^2}e} + \frac{\text{bcln}_1\left(\frac{\sqrt[6]{-c^2}x}{\sqrt[6]{-c^2}d-e}\right)}{2\sqrt{-c^2}e} - \frac{\text{bcln}_1\left(\frac{\sqrt[6]{-c^2}x}{\sqrt[6]{-c^2}d+e}\right)}{2\sqrt{-c^2}e} + \frac{\text{bcln}_1\left(\frac{\sqrt[6]{-c^2}x}{\sqrt[6]{-c^2}d-e}\right)}{2\sqrt{-c^2}e} + \frac{\text{bcln}(d+ex)\log\left(\frac{\sqrt[6]{-c^2}x+1}{\sqrt[6]{-c^2}x-1}\right)}{2\sqrt{-c^2}e} + \frac{\text{bcln}(d+ex)\log\left(\frac{\sqrt[6]{-c^2}x+\sqrt{-1}}{\sqrt[6]{-c^2}x-\sqrt{-1}}\right)}{2\sqrt{-c^2}e} - \frac{\text{bcln}(d+ex)\log\left(\frac{\sqrt[6]{-c^2}x+\sqrt{-1}}{\sqrt[6]{-c^2}x-\sqrt{-1}}\right)}{2\sqrt{-c^2}e} + \frac{\text{bcln}(d+ex)\log\left(\frac{(-1)^{1/6}\sqrt[6]{-c^2}x+\sqrt{-1}}{\sqrt[6]{-c^2}x-\sqrt{-1}}\right)}{2\sqrt{-c^2}e} - \frac{\text{bcln}(d+ex)\log\left(\frac{(-1)^{1/6}\sqrt[6]{-c^2}x+\sqrt{-1}}{\sqrt[6]{-c^2}x-\sqrt{-1}}\right)}{2\sqrt{-c^2}e}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x^3])/(d + e*x), x]$

[Out] $((a + b*\text{ArcTan}[c*x^3])*Log[d + e*x])/e + (b*c*Log[(e*(1 - (-c^2)^(1/6)*x))/((-c^2)^(1/6)*d + e)]*Log[d + e*x])/(2*.Sqrt[-c^2]*e) - (b*c*Log[-((e*(1 + (-c^2)^(1/6)*x))/((-c^2)^(1/6)*d - e))]*Log[d + e*x])/(2*.Sqrt[-c^2]*e) + (b*c*Log[-((e*((-1)^(1/3) + (-c^2)^(1/6)*x))/((-c^2)^(1/6)*d - (-1)^(1/3)*e))]*Log[d + e*x])/(2*.Sqrt[-c^2]*e) - (b*c*Log[-((e*((-1)^(2/3) + (-c^2)^(1/6)*x))/((-c^2)^(1/6)*d - (-1)^(2/3)*e))]*Log[d + e*x])/(2*.Sqrt[-c^2]*e) + (b*c*Log[((-1)^(2/3)*e*(1 + (-1)^(1/3)*(-c^2)^(1/6)*x))/((-c^2)^(1/6)*d + (-1)^(2/3)*e)]*Log[d + e*x])/(2*.Sqrt[-c^2]*e) - (b*c*Log[((-1)^(1/3)*e*(1 + (-1)^(2/3)*(-c^2)^(1/6)*x))/((-c^2)^(1/6)*d + (-1)^(1/3)*e)]*Log[d + e*x])/(2*.Sqrt[-c^2]*e) - (b*c*PolyLog[2, ((-c^2)^(1/6)*(d + e*x))/((-c^2)^(1/6)*d - e)]$

$$\frac{b c \text{PolyLog}[2, ((-c^2)^{(1/6)} (d + e x)) / ((-c^2)^{(1/6)} d + e)]}{2 \sqrt{-c^2} e} + \frac{b c \text{PolyLog}[2, ((-c^2)^{(1/6)} (d + e x)) / ((-c^2)^{(1/6)} d - (-1)^{(1/3)} e)]}{2 \sqrt{-c^2} e} - \frac{b c \text{PolyLog}[2, ((-c^2)^{(1/6)} (d + e x)) / ((-c^2)^{(1/6)} d + (-1)^{(1/3)} e)]}{2 \sqrt{-c^2} e} - \frac{b c \text{PolyLog}[2, ((-c^2)^{(1/6)} (d + e x)) / ((-c^2)^{(1/6)} d - (-1)^{(2/3)} e)]}{2 \sqrt{-c^2} e} + \frac{b c \text{PolyLog}[2, ((-c^2)^{(1/6)} (d + e x)) / ((-c^2)^{(1/6)} d + (-1)^{(2/3)} e)]}{2 \sqrt{-c^2} e}$$
Rule 209

$$\text{Int}[(a_.) + (b_.) (x_)^2, x] \rightarrow \text{Simp}[(1/(Rt[a, 2] * Rt[b, 2])) * \text{ArcTan}[Rt[b, 2] * (x/Rt[a, 2])], x] \text{ /; FreeQ}\{a, b\}, x \text{ \&& PosQ}\{a/b\} \text{ \&& (GtQ}\{a, 0\} \text{ || GtQ}\{b, 0\})$$
Rule 266

$$\text{Int}[(x_)^{(m_.)} / ((a_.) + (b_.) (x_)^{(n_.)}), x] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b x^n, x]] / (b n), x] \text{ /; FreeQ}\{a, b, m, n\}, x \text{ \&& EqQ}\{m, n - 1\}$$
Rule 281

$$\text{Int}[(x_)^{(m_.)} ((a_.) + (b_.) (x_)^{(n_.)})^{(p_.)}, x] \rightarrow \text{With}\{k = \text{GCD}[m + 1, n]\}, \text{Dist}[1/k, \text{Subst}[\text{Int}[x^{((m + 1)/k - 1)} (a + b x^{(n/k)})^p, x], x, x^k], x] \text{ /; k != 1} \text{ /; FreeQ}\{a, b, p\}, x \text{ \&& IGtQ}\{n, 0\} \text{ \&& IntegerQ}\{m\}]$$
Rule 2438

$$\text{Int}[\text{Log}[(c_.) ((d_.) + (e_.) (x_)^{(n_.)})], x] \rightarrow \text{Simp}[-\text{PolyLog}[2, (-c) * e * x^n] / n, x] \text{ /; FreeQ}\{c, d, e, n\}, x \text{ \&& EqQ}\{c * d, 1\}$$
Rule 2440

$$\text{Int}[(a_.) + \text{Log}[(c_.) ((d_.) + (e_.) (x_))] * (b_.) / ((f_.) + (g_.) (x_)), x] \rightarrow \text{Dist}[1/g, \text{Subst}[\text{Int}[(a + b \text{Log}[1 + c e (x/g)]) / x, x], x, f + g x], x] \text{ /; FreeQ}\{a, b, c, d, e, f, g\}, x \text{ \&& NeQ}\{e * f - d * g, 0\} \text{ \&& EqQ}\{g + c (e * f - d * g), 0\}]$$
Rule 2441

$$\text{Int}[(a_.) + \text{Log}[(c_.) ((d_.) + (e_.) (x_))^{(n_.)} * (b_.) / ((f_.) + (g_.) (x_)), x] \rightarrow \text{Simp}[\text{Log}[e * ((f + g x) / (e * f - d * g))] * ((a + b \text{Log}[c (d + e x)^n]) / g), x] - \text{Dist}[b * e * (n/g), \text{Int}[\text{Log}[(e * (f + g x)) / (e * f - d * g)] / (d + e x), x], x] \text{ /; FreeQ}\{a, b, c, d, e, f, g, n\}, x \text{ \&& NeQ}\{e * f - d * g, 0\}]$$
Rule 2463

$$\text{Int}[(a_.) + \text{Log}[(c_.) ((d_.) + (e_.) (x_))^{(n_.)} * (b_.)^{(p_.)} * ((h_.) * (x_))^{(m_.)} * ((f_.) + (g_.) * (x_)^{(r_.)})^{(q_.)}, x] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a$$

```
+ b*Log[c*(d + e*x)^n]^p, (h*x)^m*(f + g*x^r)^q, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n, p, q, r}, x] && IntegerQ[m] && IntegerQ[q]
```

Rule 4976

```
Int[((a_.) + ArcTan[(c_)*(x_)^(n_)]*(b_.))/((d_) + (e_)*(x_)), x_Symbol]
:› Simp[Log[d + e*x]*((a + b*ArcTan[c*x^n])/e), x] - Dist[b*c*(n/e), Int[x^(n - 1)*(Log[d + e*x]/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, n}, x] && IntegerQ[n]
```

Rubi steps

$$\begin{aligned} \int \frac{a + b \tan^{-1}(cx^3)}{d + ex} dx &= \int \left(\frac{a}{d + ex} + \frac{b \tan^{-1}(cx^3)}{d + ex} \right) dx \\ &= \frac{a \log(d + ex)}{e} + b \int \frac{\tan^{-1}(cx^3)}{d + ex} dx \end{aligned}$$

Mathematica [F]

time = 180.00, size = 0, normalized size = 0.00

\$Aborted

Verification is not applicable to the result.

[In] `Integrate[(a + b*ArcTan[c*x^3])/(d + e*x), x]`

[Out] \$Aborted

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 4.

time = 0.11, size = 172, normalized size = 0.23

method	result
default	$\frac{a \ln(ex+d)}{e} + \frac{b \ln(ex+d) \arctan(cx^3)}{e} - \frac{b e^2 \sum_{R1=\text{RootOf}(-Z^6 c^2 - 6 c^2 d - Z^5 + 15 c^2 d^2 - Z^4 - 20 c^2 d^3 - Z^3 + 15 c^2 d^4 - Z^2 - 6 c^2 d^5 - Z + c^2 d^6)}{2c}$
risch	$\frac{i b \ln(ex+d) \ln(-i c x^3 + 1)}{2e} - \frac{i b \sum_{R1=\text{RootOf}(c - Z^3 - 3 - Z^{2cd+3} - Z^{cd^2-cd^3+e^3} \text{RootOf}(-Z^{2+1, index=1}))} \left(\ln(ex+d) \ln\left(\frac{-ex+R1}{R1}\right)\right)}{2e}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x^3))/(e*x+d), x, method=_RETURNVERBOSE)`

[Out] $a \ln(e*x+d)/e + b \ln(e*x+d)/e * \arctan(c*x^3) - 1/2 * b * e^2 / c * \text{sum}(1/(_R1^3 - 3*_R1^2 * d + 3*_R1*d^2 - d^3) * (\ln(e*x+d) * \ln((-e*x+_R1-d)/_R1) + \text{dilog}((-e*x+_R1-d)/_R1)), _R1 = \text{RootOf}(_Z^6*c^2 - 6*_Z^5*c^2*d + 15*_Z^4*c^2*d^2 - 20*_Z^3*c^2*d^3 + 15*_Z^2*c^2*d^4 - 6*_Z*c^2*d^5 + c^2*d^6 + e^6))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^3))/(e*x+d),x, algorithm="maxima")`

[Out] $a*e^{(-1)}*\log(x*e + d) + 2*b*\text{integrate}(1/2*\arctan(c*x^3)/(x*e + d), x)$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^3))/(e*x+d),x, algorithm="fricas")`

[Out] $\text{integral}((b*\arctan(c*x^3) + a)/(x*e + d), x)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x**3))/(e*x+d),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^3))/(e*x+d),x, algorithm="giac")`

[Out] $\text{integrate}((b*\arctan(c*x^3) + a)/(e*x + d), x)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{a + b \operatorname{atan}(c x^3)}{d + e x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^3))/(d + e*x),x)`[Out] `int((a + b*atan(c*x^3))/(d + e*x), x)`

3.31 $\int \frac{a+b\text{ArcTan}(cx^3)}{(d+ex)^2} dx$

Optimal. Leaf size=906

$$-\frac{bc^{2/3}de^3\text{ArcTan}\left(\sqrt[3]{c}x\right)}{c^2d^6+e^6}+\frac{bc^2d^5\text{ArcTan}(cx^3)}{e(c^2d^6+e^6)}-\frac{a+b\text{ArcTan}(cx^3)}{e(d+ex)}+\frac{bc^{2/3}d\left(\sqrt{3}cd^3+e^3\right)\text{ArcTan}\left(\sqrt{3}-2\sqrt[3]{c}x\right)}{2(c^2d^6+e^6)}$$

[Out] $-b*c^{(2/3)*d*e^3*\arctan(c^{(1/3)*x})/(c^{2*d^6+e^6})+b*c^{2*d^5*\arctan(c*x^3)/e/(c^{2*d^6+e^6})+(-a-b*\arctan(c*x^3))/e/(e*x+d)+3*b*c*d^2*e^2*\ln(e*x+d)/(c^{2*d^6+e^6})+1/2*b*c^{(5/3)*d^4*\ln(1+c^{(2/3)*x^2})/(c^{2*d^6+e^6})-1/2*b*c*d^2*e^2*(1/2)*(c^{2*x^6+1})/(c^{2*d^6+e^6})+1/2*b*c^{(2/3)*d*\arctan(2*c^{(1/3)*x+3^(1/2)})*(-e^3+c*d^3*3^(1/2))}/(c^{2*d^6+e^6})-1/2*b*c^{(2/3)*d*\arctan(2*c^{(1/3)*x-3^(1/2)})*(e^3+c*d^3*3^(1/2))}/(c^{2*d^6+e^6})-1/4*b*c^{(2/3)*d*\ln(1+c^{(2/3)*x^2-c^{(1/3)*x^3}}/(c^{2*d^6+e^6})-1/4*b*c^{(2/3)*d*\ln(1+c^{(2/3)*x^2+c^{(1/3)*x^3}}/(c^{2*d^6+e^6})-1/2*b*c^{(5/3)*e*\ln((-c^2)^{(1/6)}+c^{(2/3)*x})*(-e^3+d^3*(-c^2)^{(1/2)})}/(-c^2)^{(2/3)}/(c^{2*d^6+e^6})+1/4*b*c^{(5/3)*e*\ln((-c^2)^{(1/3)}-c^{(2/3)*(-c^2)^{(1/6)}}*x+c^{(4/3)*x^2})*(-e^3+d^3*(-c^2)^{(1/2)})}/(-c^2)^{(2/3)}/(c^{2*d^6+e^6})-1/2*b*c^{(5/3)*e*\arctan(1/3*(c^{(4/3)+2*(-c^2)^{(5/6)*x}}/c^{(4/3)*3^(1/2)})*3^(1/2)*(-e^3+d^3*(-c^2)^{(1/2)})}/(-c^2)^{(2/3)}/(c^{2*d^6+e^6})+1/2*b*c^{(5/3)*e*\ln((-c^2)^{(1/6)}-c^{(2/3)*x})*(e^3+d^3*(-c^2)^{(1/2)})}/(-c^2)^{(2/3)}/(c^{2*d^6+e^6})-1/4*b*c^{(5/3)*e*\ln((-c^2)^{(1/3)}+c^{(2/3)*(-c^2)^{(1/6)}*x+c^{(4/3)*x^2}}*(e^3+d^3*(-c^2)^{(1/2)})}/(-c^2)^{(2/3)}/(c^{2*d^6+e^6})+1/2*b*c^{(5/3)*e*\arctan(1/3*(1+2*c^{(2/3)*x}/(-c^2)^{(1/6)})*3^(1/2))*3^(1/2)*(e^3+d^3*(-c^2)^{(1/2)})}/(-c^2)^{(2/3)}/(c^{2*d^6+e^6})$

Rubi [A]

time = 0.94, antiderivative size = 906, normalized size of antiderivative = 1.00, number of steps used = 34, number of rules used = 15, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}}$ = 0.833, Rules used = {4980, 6857, 1890, 1430, 649, 209, 266, 648, 631, 210, 642, 1525, 298, 31, 1483}

$\frac{a c^{2/3} d e^3 \text{ArcTan}\left(\sqrt[3]{c} x\right)}{c^2 d^6+e^6}+\frac{b c^2 d^5 \text{ArcTan}(c x^3)}{e \left(c^2 d^6+e^6\right)}-\frac{a+b \text{ArcTan}(c x^3)}{e (d+e x)}+\frac{b c^{2/3} d \left(\sqrt{3} c d^3+e^3\right) \text{ArcTan}\left(\sqrt{3}-2 \sqrt[3]{c} x\right)}{2 \left(c^2 d^6+e^6\right)}$

Antiderivative was successfully verified.

[In] $\text{Int}[(a + b*\text{ArcTan}[c*x^3])/(d + e*x)^2, x]$

[Out] $-\left((b*c^{(2/3)*d*e^3*\arctan(c^{(1/3)*x})/(c^{2*d^6+e^6})+(b*c^{2*d^5*\arctan(c*x^3)/e/(c^{2*d^6+e^6})-(a+b*\arctan(c*x^3))/e/(e*(d+e*x))+(b*c^{(2/3)*d*(\text{Sqrt}[3]*c*d^3+e^3)*\arctan(\text{Sqrt}[3]-2*c^{(1/3)*x})/(2*(c^{2*d^6+e^6})+(b*c^{(2/3)*d*(\text{Sqrt}[3]*c*d^3-e^3)*\arctan(\text{Sqrt}[3]+2*c^{(1/3)*x})/(2*(c^{2*d^6+e^6})+(\text{Sqrt}[3]*b*c^{(5/3)*e*(\text{Sqrt}[-c^2]*d^3+e^3)*\arctan[(1+(2*c^{(2/3)*x}/(-c^2)^{(1/6)})/\text{Sqrt}[3]])/(2*(-c^2)^{(2/3)*(c^{2*d^6+e^6})}-(S$

$$\begin{aligned}
& \text{qrt[3]*b*c}^{(5/3)}*e*(\text{Sqrt}[-c^2]*d^3 - e^3)*\text{ArcTan}[(c^{(4/3)} + 2*(-c^2)^{(5/6)}*x)/(\text{Sqrt}[3]*c^{(4/3)})]/(2*(-c^2)^{(2/3)}*(c^2*d^6 + e^6)) + (b*c}^{(5/3)}*e*(\text{Sqr}\\
& \text{t}[-c^2]*d^3 + e^3)*\text{Log}[(-c^2)^{(1/6)} - c^{(2/3)*x}]/(2*(-c^2)^{(2/3)}*(c^2*d^6\\
& + e^6)) - (b*c}^{(5/3)}*e*(\text{Sqrt}[-c^2]*d^3 - e^3)*\text{Log}[(-c^2)^{(1/6)} + c^{(2/3)*x}]/(2*(-c^2)^{(2/3)}*(c^2*d^6 + e^6)) + (3*b*c*d^2*e^2*\text{Log}[d + e*x])/(c^2*d^6\\
& + e^6) + (b*c}^{(5/3)}*d^4*\text{Log}[1 + c^{(2/3)*x^2}]/(2*(c^2*d^6 + e^6)) - (b*c}^{(2/3)}*d*(c*d^3 - \text{Sqrt}[3]*e^3)*\text{Log}[1 - \text{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}]/(4*(c^2*d^6 + e^6)) - (b*c}^{(2/3)}*d*(c*d^3 + \text{Sqrt}[3]*e^3)*\text{Log}[1 + \text{Sqrt}[3]*c^{(1/3)*x} + c^{(2/3)*x^2}]/(4*(c^2*d^6 + e^6)) + (b*c}^{(5/3)}*e*(\text{Sqrt}[-c^2]*d^3 - e^3)*\text{Log}[(-c^2)^{(1/3)} - c^{(2/3)*(-c^2)^{(1/6)*x} + c^{(4/3)*x^2}}]/(4*(-c^2)^{(2/3)}*(c^2*d^6 + e^6)) - (b*c}^{(5/3)}*e*(\text{Sqrt}[-c^2]*d^3 + e^3)*\text{Log}[(-c^2)^{(1/3)} + c^{(2/3)*(-c^2)^{(1/6)*x} + c^{(4/3)*x^2}}]/(4*(-c^2)^{(2/3)}*(c^2*d^6 + e^6)) - (b*c*d^2*e^2*\text{Log}[1 + c^2*x^6])/(2*(c^2*d^6 + e^6))
\end{aligned}$$
Rule 31

$$\text{Int}[(a_ + b_)*(x_)^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}[\{a, b\}, x]$$
Rule 209

$$\text{Int}[(a_ + b_)*(x_)^2^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(1/\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{GtQ}[a, 0] \text{ || } \text{GtQ}[b, 0])$$
Rule 210

$$\text{Int}[(a_ + b_)*(x_)^2^{-1}, x_{\text{Symbol}}] \rightarrow \text{Simp}[(-(Rt[-a, 2]*Rt[-b, 2])^{(-1)})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/Rt[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&& \text{PosQ}[a/b] \&& (\text{LtQ}[a, 0] \text{ || } \text{LtQ}[b, 0])$$
Rule 266

$$\text{Int}[(x_)^{(m_)}/((a_ + b_)*(x_)^{(n_)}), x_{\text{Symbol}}] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x^n, x]]/(b*n), x] /; \text{FreeQ}[\{a, b, m, n\}, x] \&& \text{EqQ}[m, n - 1]$$
Rule 298

$$\text{Int}[(x_)/((a_ + b_)*(x_)^3), x_{\text{Symbol}}] \rightarrow \text{Dist}[-(3*\text{Rt}[a, 3]*\text{Rt}[b, 3])^{(-1)}, \text{Int}[1/(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x), x], x] + \text{Dist}[1/(3*\text{Rt}[a, 3]*\text{Rt}[b, 3]), \text{Int}[(\text{Rt}[a, 3] + \text{Rt}[b, 3]*x)/(\text{Rt}[a, 3]^2 - \text{Rt}[a, 3]*\text{Rt}[b, 3]*x + \text{Rt}[b, 3]^2*x^2), x], x] /; \text{FreeQ}[\{a, b\}, x]$$
Rule 631

$$\text{Int}[(a_ + b_)*(x_) + (c_)*(x_)^2^{-1}, x_{\text{Symbol}}] \rightarrow \text{With}[\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Dist}[-2/b, \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)$$

```
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] :> D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 649

```
Int[((d_) + (e_)*(x_))/((a_) + (c_)*(x_)^2), x_Symbol] :> Dist[d, Int[1/(
a + c*x^2), x], x] + Dist[e, Int[x/(a + c*x^2), x], x] /; FreeQ[{a, c, d, e
}, x] && !NiceSqrtQ[(-a)*c]
```

Rule 1430

```
Int[((d_) + (e_)*(x_)^3)/((a_) + (c_)*(x_)^6), x_Symbol] :> With[{q = Rt[
c/a, 6]}, Dist[1/(3*a*q^2), Int[(q^2*d - e*x)/(1 + q^2*x^2), x], x] + (Dist
[1/(6*a*q^2), Int[(2*q^2*d - (Sqrt[3]*q^3*d - e)*x)/(1 - Sqrt[3]*q*x + q^2*
x^2), x], x] + Dist[1/(6*a*q^2), Int[(2*q^2*d + (Sqrt[3]*q^3*d + e)*x)/(1 +
Sqrt[3]*q*x + q^2*x^2), x], x])] /; FreeQ[{a, c, d, e}, x] && NeQ[c*d^2 +
a*e^2, 0] && PosQ[c/a]
```

Rule 1483

```
Int[(x_)^(m_)*(a_ + (c_)*(x_)^(n2_))^(p_)*(d_ + (e_)*(x_)^(n_))^(q
_), x_Symbol] :> Dist[1/n, Subst[Int[(d + e*x)^q*(a + c*x^2)^p, x], x, x^n
], x] /; FreeQ[{a, c, d, e, m, n, p, q}, x] && EqQ[n2, 2*n] && EqQ[Simplify
[m - n + 1], 0]
```

Rule 1525

```
Int[((f_)*(x_)^(m_)*(d_ + (e_)*(x_)^(n_)))/((a_) + (c_)*(x_)^(n2_))
, x_Symbol] :> With[{q = Rt[(-a)*c, 2]}, Dist[-(e/2 + c*(d/(2*q))), Int[(f*
x)^m/(q - c*x^n), x], x] + Dist[e/2 - c*(d/(2*q)), Int[(f*x)^m/(q + c*x^n),
x], x]] /; FreeQ[{a, c, d, e, f, m}, x] && EqQ[n2, 2*n] && IGtQ[n, 0]
```

Rule 1890

```
Int[(Pq_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> With[{v = Sum[x^ii*((Coeff[Pq, x, ii] + Coeff[Pq, x, n/2 + ii]*x^(n/2))/(a + b*x^n)), {ii, 0, n/2 - 1}]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && PolyQ[Pq, x] && IGtQ[n/2, 0] && Expon[Pq, x] < n
```

Rule 4980

```
Int[((a_.) + ArcTan[(c_)*(x_)^(n_)]*(b_.))*(d_) + (e_)*(x_)^(m_), x_Symbol] :> Simp[(d + e*x)^(m + 1)*((a + b*ArcTan[c*x^n])/(e*(m + 1))), x] - Dist[b*c*(n/(e*(m + 1))), Int[x^(n - 1)*((d + e*x)^(m + 1)/(1 + c^2*x^(2*n))), x], x] /; FreeQ[{a, b, c, d, e, m, n}, x] && NeQ[m, -1]
```

Rule 6857

```
Int[(u_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] :> With[{v = RationalFunctionExpand[u/(a + b*x^n), x]}, Int[v, x] /; SumQ[v]] /; FreeQ[{a, b}, x] && IGtQ[n, 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{a + b \tan^{-1}(cx^3)}{(d+ex)^2} dx &= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{b \int \frac{3cx^2}{(d+ex)(1+c^2x^6)} dx}{e} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{(3bc) \int \frac{x^2}{(d+ex)(1+c^2x^6)} dx}{e} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{(3bc) \int \left(\frac{d^2e^4}{(c^2d^6+e^6)(d+ex)} + \frac{(d-ex)(-e^4+c^2d^4x^2+c^2d^2e^2x^4)}{(c^2d^6+e^6)(1+c^2x^6)} \right) dx}{e} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{3bcd^2e^2 \log(d+ex)}{c^2d^6+e^6} + \frac{(3bc) \int \frac{(d-ex)(-e^4+c^2d^4x^2+c^2d^2e^2x^4)}{1+c^2x^6} dx}{e(c^2d^6+e^6)} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{3bcd^2e^2 \log(d+ex)}{c^2d^6+e^6} + \frac{(3bc) \int \left(\frac{-de^4-c^2d^4ex^3}{1+c^2x^6} + \frac{x(e^5+c^2d^3e^2x^3)}{1+c^2x^6} \right) dx}{e(c^2d^6+e^6)} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{3bcd^2e^2 \log(d+ex)}{c^2d^6+e^6} + \frac{(3bc) \int \frac{-de^4-c^2d^4ex^3}{1+c^2x^6} dx}{e(c^2d^6+e^6)} + \frac{(3bc) \int \frac{x(e^5+c^2d^3e^2x^3)}{1+c^2x^6} dx}{e(c^2d^6+e^6)} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{3bcd^2e^2 \log(d+ex)}{c^2d^6+e^6} + \frac{(b\sqrt[3]{c}) \int \frac{-2c^{2/3}de^4 - (c^2d^4e - \sqrt{3}cd^2e^4)x}{1-\sqrt{3}\sqrt[3]{c}x+c^{2/3}x^2} dx}{2e(c^2d^6+e^6)} \\
&= -\frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{3bcd^2e^2 \log(d+ex)}{c^2d^6+e^6} + \frac{(bc^{7/3}d^4) \int \frac{x}{1+c^{2/3}x^2} dx}{c^2d^6+e^6} + \frac{(bc^3d^5) \text{Sub}}{e(d+ex)} \\
&= -\frac{bc^{2/3}de^3 \tan^{-1}(\sqrt[3]{c}x)}{c^2d^6+e^6} + \frac{bc^2d^5 \tan^{-1}(cx^3)}{e(c^2d^6+e^6)} - \frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{bc^{5/3}e(\sqrt{-c^2}x)}{2} \\
&= -\frac{bc^{2/3}de^3 \tan^{-1}(\sqrt[3]{c}x)}{c^2d^6+e^6} + \frac{bc^2d^5 \tan^{-1}(cx^3)}{e(c^2d^6+e^6)} - \frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{bc^{2/3}d(\sqrt{3}cx)}{c^2d^6+e^6} \\
&= -\frac{bc^{2/3}de^3 \tan^{-1}(\sqrt[3]{c}x)}{c^2d^6+e^6} + \frac{bc^2d^5 \tan^{-1}(cx^3)}{e(c^2d^6+e^6)} - \frac{a + b \tan^{-1}(cx^3)}{e(d+ex)} + \frac{bc^{2/3}d(\sqrt{3}cx)}{c^2d^6+e^6}
\end{aligned}$$

Mathematica [A]

time = 10.00, size = 536, normalized size = 0.59

Antiderivative was successfully verified

[In] `Integrate[(a + b*ArcTan[c*x^3])/(d + e*x)^2, x]`

[Out]
$$\begin{aligned} & (-4*a*c^{(1/3)}*(c^2*d^6 + e^6) - 4*b*c*d*(c^{(4/3)}*d^4 - c^{(2/3)}*d^2*e^2 + e^4)*(d + e*x)*\text{ArcTan}[c^{(1/3)}*x] - 4*b*c^{(1/3)}*(c^2*d^6 + e^6)*\text{ArcTan}[c*x^3] \\ & - 2*b*c^{(2/3)}*(2*c^{(5/3)}*d^5 - \text{Sqrt}[3]*c^{(4/3)}*d^4*e + c*d^3*e^2 - c^{(1/3)}*d*e^4 + \text{Sqrt}[3]*e^5)*(d + e*x)*\text{ArcTan}[\text{Sqrt}[3] - 2*c^{(1/3)}*x] + 2*b*c^{(2/3)}*(2*c^{(5/3)}*d^5 + \text{Sqrt}[3]*c^{(4/3)}*d^4*e + c*d^3*e^2 - c^{(1/3)}*d*e^4 - \text{Sqrt}[3]*e^5)*(d + e*x)*\text{ArcTan}[\text{Sqrt}[3] + 2*c^{(1/3)}*x] + 12*b*c^{(4/3)}*d^2*e^3*(d + e*x)*\text{Log}[d + e*x] + 2*b*e*(c^2*d^4 + c^{(2/3)}*e^4)*(d + e*x)*\text{Log}[1 + c^{(2/3)}*x^2] - b*c^{(2/3)}*e*(c^{(4/3)}*d^4 - \text{Sqrt}[3]*c*d^3*e - \text{Sqrt}[3]*c^{(1/3)}*d*e^3 + e^4)*(d + e*x)*\text{Log}[1 - \text{Sqrt}[3]*c^{(1/3)}*x + c^{(2/3)}*x^2] - b*c^{(2/3)}*e*(c^{(4/3)}*d^4 + \text{Sqrt}[3]*c*d^3*e + \text{Sqrt}[3]*c^{(1/3)}*d*e^3 + e^4)*(d + e*x)*\text{Log}[1 + \text{Sqrt}[3]*c^{(1/3)}*x + c^{(2/3)}*x^2] - 2*b*c^{(4/3)}*d^2*e^3*(d + e*x)*\text{Log}[1 + c^2*x^6])/((4*c^{(1/3)}*e*(c^2*d^6 + e^6)*(d + e*x))) \end{aligned}$$

Maple [A]

time = 0.40, size = 1220, normalized size = 1.35

method	result	size
default	Expression too large to display	1220

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a+b*arctan(c*x^3))/(e*x+d)^2, x, method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -a/(e*x+d)/e+3*b*c*d^2*e^2*\ln(e*x+d)/(c^2*d^6+e^6)+b*e^4*c^3/(c^2*d^6+e^6)*(1/c^2)^{(4/3)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*3^{(1/2)}-1/2*b*e^3*c^3/(c^2*d^6+e^6)*(1/c^2)^{(7/6)}*\arctan(2*x/(1/c^2)^{(1/6)}+3^{(1/2)})*d-b*e^3*c^3/(c^2*d^6+e^6)*(1/c^2)^{(7/6)}*\arctan(x/(1/c^2)^{(1/6})*d+1/2*b*e*c/(c^2*d^6+e^6)/(1/c^2)^{(1/6)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*d^3-1/2*b*e^4*c/(c^2*d^6+e^6)*(1/c^2)^{(1/3)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*3^{(1/2)}+b/e*c^3/(c^2*d^6+e^6)*(1/c^2)^{(1/2)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*d^5+1/2*b*e*c/(c^2*d^6+e^6)/(1/c^2)^{(1/6)}*\arctan(2*x/(1/c^2)^{(1/6)}+3^{(1/2)})*d^3-1/2*b*e^4*c/(c^2*d^6+e^6)*(1/c^2)^{(1/3)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*3^{(1/2)}+b/e*c^3/(c^2*d^6+e^6)*(1/c^2)^{(1/2)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*d^5-b/e*c^3/(c^2*d^6+e^6)/(1/c^2)^{(1/6)}*\arctan(2*x/(1/c^2)^{(1/6)}+3^{(1/2)})*d^3-1/2*b*c^3/(c^2*d^6+e^6)*(1/c^2)^{(2/3)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*3^{(1/2)}*d^4+1/2*b*c^3/(c^2*d^6+e^6)*(1/c^2)^{(2/3)}*\arctan(2*x/(1/c^2)^{(1/6)}+3^{(1/2)})*3^{(1/2)}*d^4-1/2*b*e^3*c^3/(c^2*d^6+e^6)*(1/c^2)^{(2/3)}*\arctan(2*x/(1/c^2)^{(1/6)}-3^{(1/2)})*3^{(1/2)}*d^5+1/4*b*e^3*c^3/(c^2*d^6+e^6)*\ln(x^2-3^{(1/2)}*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*3^{(1/2)}*(1/c^2)^{(7/6)}*d-1/4*b*e^3*c^3/(c^2*d^6+e^6)*\ln(x^2+3^{(1/2)}*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*3^{(1/2)}*(1/c^2)^{(7/6)}*d-1/4*b*e*c^3/(c^2*d^6+e^6)*\ln(x^2-3^{(1/2)}*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*3^{(1/2)}*(1/c^2)^{(5/6)}*d^3+1/2*b*e^4*c/(c^2*d^6+e^6)*\ln(x^2+(1/c^2)^{(1/3)})*(1/c^2)^{(1/3)}-1/2*b*e^2*c/(c^2*d^6+e^6)*\ln(x^2+(1/c^2)^{(1/3)})*d^2-1/4*b*c^3/(c^2*d^6+e^6)*\ln(x^2-3^{(1/2)}*(1/c^2)^{(1/6)}*x+(1/c^2)^{(1/3)})*3^{(1/2)}*(1/c^2)^{(5/6)}*d^3 \end{aligned}$$

$$\begin{aligned}
& /2)*(1/c^2)^{(1/6)*x+(1/c^2)^(1/3)}*(1/c^2)^(2/3)*d^4-1/4*b*c^3/(c^2*d^6+e^6) \\
&)*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6)*x+(1/c^2)^(1/3)}*(1/c^2)^(2/3)*d^4+1/2*b*c^3 \\
& /(c^2*d^6+e^6)*\ln(x^2+(1/c^2)^(1/3))*(1/c^2)^(2/3)*d^4-1/4*b*e^4*c/(c^2*d^6 \\
& +e^6)*\ln(x^2-3^(1/2)*(1/c^2)^{(1/6)*x+(1/c^2)^(1/3)}*(1/c^2)^(1/3)-1/2*b*e^2 \\
& *c/(c^2*d^6+e^6)*\ln(x^2-3^(1/2)*(1/c^2)^{(1/6)*x+(1/c^2)^(1/3)}*d^2-1/4*b*e^ \\
& 4*c/(c^2*d^6+e^6)*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6)*x+(1/c^2)^(1/3)}*(1/c^2)^(1/ \\
& 3)-1/2*b*e^2*c/(c^2*d^6+e^6)*\ln(x^2+3^(1/2)*(1/c^2)^{(1/6)*x+(1/c^2)^(1/3)}* \\
& d^2-b/(e*x+d)/e*\arctan(c*x^3)
\end{aligned}$$

Maxima [A]

time = 0.49, size = 449, normalized size = 0.50

$$\frac{1}{4} \left(\left(\frac{12 d^2 e^2 \log(xe + d)}{c^2 d^6 + e^6} - \frac{e (\sqrt{3} d^{1/2} e^{1/2} \rho^{1/2} \sigma^{1/2} \text{atan}(\sqrt{3} \rho))}{c^2} - \frac{2 (\sqrt{3} d^{1/2} e^{1/2} \rho^{1/2} \sigma^{1/2} \text{atan}(\sqrt{3} \rho)) \text{atan}\left(\frac{\sqrt{3} d^{1/2} e^{1/2}}{c}\right)}{c^2} + \frac{2 (\sqrt{3} d^{1/2} e^{1/2} \rho^{1/2} \sigma^{1/2} \text{atan}(\sqrt{3} \rho)) \text{atan}\left(\frac{\sqrt{3} d^{1/2} e^{1/2}}{c}\right)}{c^2} + \frac{(\sqrt{3} d^{1/2} e^{1/2} \rho^{1/2} \sigma^{1/2} \text{atan}(\sqrt{3} \rho)) \text{atan}\left(\frac{\sqrt{3} d^{1/2} e^{1/2}}{c}\right)}{c^2} + \frac{(\sqrt{3} d^{1/2} e^{1/2} \rho^{1/2} \sigma^{1/2} \text{atan}(\sqrt{3} \rho)) \text{atan}\left(\frac{\sqrt{3} d^{1/2} e^{1/2}}{c}\right)}{c^2} - \frac{2 (\sqrt{3} d^{1/2} e^{1/2} \rho^{1/2} \sigma^{1/2} \text{atan}(\sqrt{3} \rho)) \text{atan}\left(\frac{\sqrt{3} d^{1/2} e^{1/2}}{c}\right)}{c^2} \right) c - \frac{4 \arctan(cx^3)}{xe^2 + dc} \right) b - \frac{a}{xe^2 + dc}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x^3))/(e*x+d)^2,x, algorithm="maxima")
[Out] 1/4*((12*d^2*e^2*log(xe + d)/(c^2*d^6 + e^6) - (4*(c^(8/3)*d^5 - c^2*d^3*e^2 + c^(4/3)*d*e^4)*arctan(c^(1/3)*x)/c^(5/3) - 2*(sqrt(3)*c^(8/3)*d^4*e + 2*c^3*d^5 + c^(7/3)*d^3*e^2 - c^(5/3)*d*e^4 - sqrt(3)*c^(4/3)*e^5)*arctan((2*c^(2/3)*x + sqrt(3)*c^(1/3))/c^(1/3))/c^2 + 2*(sqrt(3)*c^(8/3)*d^4*e - 2*c^3*d^5 - c^(7/3)*d^3*e^2 + c^(5/3)*d*e^4 - sqrt(3)*c^(4/3)*e^5)*arctan((2*c^(2/3)*x - sqrt(3)*c^(1/3))/c^(1/3))/c^2 + (c^(8/3)*d^4*e + sqrt(3)*c^(7/3))*d^3*e^2 + 2*c^2*d^2*e^3 + sqrt(3)*c^(5/3)*d*e^4 + c^(4/3)*e^5)*log(c^(2/3)*x^2 + sqrt(3)*c^(1/3)*x + 1)/c^2 + (c^(8/3)*d^4*e - sqrt(3)*c^(7/3)*d^3*e^2 + 2*c^2*d^2*e^3 - sqrt(3)*c^(5/3)*d*e^4 + c^(4/3)*e^5)*log(c^(2/3)*x^2 - sqrt(3)*c^(1/3)*x + 1)/c^2 - 2*(c^(8/3)*d^4*e - c^2*d^2*e^3 + c^(4/3)*e^5)*log(c^(2/3)*x^2 + 1)/c^2)/(c^2*d^6*e + e^7))*c - 4*arctan(c*x^3)/(x*e^2 + d*e))*b - a/(x*e^2 + d*e)
```

Fricas [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a+b*arctan(c*x^3))/(e*x+d)^2,x, algorithm="fricas")
```

[Out] Timed out

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*atan(c*x**3))/(e*x+d)**2,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((a+b*arctan(c*x^3))/(e*x+d)^2,x, algorithm="giac")`

[Out] sage0*x

Mupad [B]

time = 0.78, size = 2105, normalized size = 2.32

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a + b*atan(c*x^3))/(d + e*x)^2,x)`

[Out] `symsum(log((729*b^6*c^14*d*e^2 + 54432*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^6*c^12*d^15*x + 729*b^6*c^14*e^3*x - 31104*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^6*c^14*d^7*e^8 - 243*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)*b^5*c^15*d^5 + 62208*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^6*c^12*d^14 + 5832*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^2*b^4*c^14*d^3*e^4 - 1944*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^2*z^2 + b^6*c^2, z, k)^3*b^3*c^15*d^7*e^2 + 15552*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^4*b^2*c^14*d^5*e^6 - 10692*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^3*b^3*c^13*d^8 + 101088*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^5*b*c^13*d^10 - 3888*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^5*b*c^15*d^9*e^4 - 12636*root(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 +`

$$\begin{aligned}
& 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c \\
& \sim 2*d^2*e^2*z^2 + b^6*c^2, z, k)^3 * b^3*c^13*e^9*x - 38880*\text{root}(64*c^2*d^6*e^6*z^6 \\
& + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3 \\
& *c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^6 * c^14*d^6*e^9*x + 116 \\
& 640*\text{root}(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2 \\
& *d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^5 * b*c^13*d^2*e^11*x \\
& - 11664*\text{root}(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c \\
& *d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2 \\
& *z^2 + b^6*c^2, z, k)^5 * b*c^15*d^8*e^5*x + 11664*\text{root}(64*c^2*d^6*e^6*z^6 \\
& + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6 \\
& *z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^2 * b^4*c^14*d^2*e^5*x - 3888* \\
& \text{root}(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4 \\
& *e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k)^3 * b^3 \\
& *c^15*d^6*e^3*x + 38880*\text{root}(64*c^2*d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2 \\
& *e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 + 12*b^4*c^2*d^2*e^2 \\
& *z^2 + b^6*c^2, z, k)^4 * b^2*c^14*d^4*e^7*x - 243*\text{root}(64*c^2*d^6*e^6*z^6 + 6 \\
& 4*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3*c*e^6*z^3 \\
& + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k) * b^5*c^15*d^4*e*x) / e^4) * \text{root}(64*c^2 \\
& *d^6*e^6*z^6 + 64*e^12*z^6 + 192*b*c*d^2*e^8*z^5 + 48*b^2*c^2*d^4*e^4*z^4 - 16*b^3 \\
& *c*e^6*z^3 + 12*b^4*c^2*d^2*e^2*z^2 + b^6*c^2, z, k), k, 1, 6) - \\
& a/(d*e + e^2*x) - (b*atan(c*x^3))/(d*e + e^2*x) + (3*b*c*d^2*e^2*log(d + e*x)) / (e^6 + c^2*d^6)
\end{aligned}$$

Chapter 4

Appendix

Local contents

4.1 Download section	212
4.2 Listing of Grading functions	212

4.1 Download section

The following zip files contain the raw integrals used in this test.

Mathematica format Mathematica_syntax.zip

Maple and Mupad format Maple_syntax.zip

Sympy format SYMPY_syntax.zip

Sage math format SAGE_syntax.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(* Small rewrite of logic in main function to make it*)
(* match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*      GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*      is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*      antiderivative*)
(* "A" if result can be considered optimal*)
```

```

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
expnResult = ExpnType[result];
expnOptimal = ExpnType[optimal];
leafCountResult = LeafCount[result];
leafCountOptimal = LeafCount[optimal];

(*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
If[expnResult<=expnOptimal,
  If[Not[FreeQ[result,Complex]], (*result contains complex*)
    If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","none"}
        ,(*ELSE*)
        finalresult={"B","Both result and optimal contain complex but leaf count
      ]
      ,(*ELSE*)
      finalresult={"C","Result contains complex when optimal does not."}
    ]
    ,(*ELSE*)(*result does not contains complex*)
    If[leafCountResult<=2*leafCountOptimal,
      finalresult={"A","none"}
      ,(*ELSE*)
      finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $"
    ]
    ]
  ,(*ELSE*)(*expnResult>expnOptimal*)
  If[FreeQ[result,Integrate] && FreeQ[result,Int],
    finalresult={"C","Result contains higher order function than in optimal. Order "<
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];
finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hypergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)

```

```
(*9 = unknown function*)
```

```
ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
  If[ListQ[expn],
    Max[Map[ExpnType, expn]],
  If[Head[expn] === Power,
    If[IntegerQ[expn[[2]]],
      ExpnType[expn[[1]]],
    If[Head[expn[[2]]] === Rational,
      If[IntegerQ[expn[[1]]] || Head[expn[[1]]] === Rational,
        1,
      Max[ExpnType[expn[[1]]], 2]],
      Max[ExpnType[expn[[1]]], ExpnType[expn[[2]], 3]]],
    If[Head[expn] === Plus || Head[expn] === Times,
      Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
    If[ElementaryFunctionQ[Head[expn]],
      Max[3, ExpnType[expn[[1]]]],
    If[SpecialFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
    If[HypergeometricFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
    If[AppellFunctionQ[Head[expn]],
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
    If[Head[expn] === RootSum,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
    If[Head[expn] === Integrate || Head[expn] === Int,
      Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
    9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
  MemberQ[{  

    Exp, Log,  

    Sin, Cos, Tan, Cot, Sec, Csc,  

    ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,  

    Sinh, Cosh, Tanh, Coth, Sech, CsCh,  

    ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsCh  

  }, func]
```

```
SpecialFunctionQ[func_] :=
  MemberQ[{  

    Erf, Erfc, Erfi,  

    FresnelS, FresnelC,
```

```

ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
MemberQ[{Hypergeometric1F1, Hypergeometric2F1, HypergeometricPFQ}, func]

AppellFunctionQ[func_] :=
MemberQ[{AppellF1}, func]

```

4.2.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#           if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#           see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

      leaf_count_result:=leafcount(result);
      #do NOT call ExpnType() if leaf size is too large. Recursion problem
      if leaf_count_result > 500000 then
          return "B","result has leaf size over 500,000. Avoiding possible recursion issues
      fi;

      leaf_count_optimal := leafcount(optimal);
      ExpnType_result := ExpnType(result);
      ExpnType_optimal := ExpnType(optimal);

```

```

if debug then
    print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
# is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
# antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A","");
            else
                return "B",cat("Both result and optimal contain complex but leaf count of r
                                convert(leaf_count_result,string)," vs. $2 (
                                convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_c
                end if
            else #result contains complex but optimal is not
                if debug then
                    print("result contains complex but optimal is not");
                fi;
                return "C","Result contains complex when optimal does not.";
            fi;
        else # result do not contain complex
    fi;
fi;

```

```

        # this assumes optimal do not as well. No check is needed here.
        if debug then
            print("result do not contain complex, this assumes optimal do not as well")
        fi;
        if leaf_count_result<=2*leaf_count_optimal then
            if debug then
                print("leaf_count_result<=2*leaf_count_optimal");
            fi;
            return "A","");
        else
            if debug then
                print("leaf_count_result>2*leaf_count_optimal");
            fi;
            return "B",cat("Leaf count of result is larger than twice the leaf count of optimal",
                           convert(leaf_count_result,string)," vs. $2(",
                           convert(leaf_count_optimal,string),")=",convert(2*leaf_count_optimal,string));
        fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                   convert(ExpnType_result,string)," vs. order ",
                   convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hypergeometric function
# 6 = appell function
# 7 = rootsum function

```

```

# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'`^`) then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'`+`) or type(expn,'`*`) then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
  9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,

```

```

arcsin,arccos,arctan,arccot,arcsec,arccsc,
sinh,cosh,tanh,coth,sech,csch,
arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func,[
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func,[Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func,[AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```
#Dec 24, 2019. Nasser M. Abbasi:
#          Port of original Maple grading function by
#          Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#          added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
                   asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
                   asinh,acosh,atanh,acoth,asech,acsch
                   ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
                    fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
                    gamma,loggamma,digamma,zeta,polylog,LambertW,
                    elliptic_f,elliptic_e,elliptic_pi,exp_polar
                    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False
    except:
        return False
```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn))  #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow):  #type(expn,'`^`)
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'`+') or type(expn,'`*`)
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

#print ("Enter grade_antiderivative for sageMath")
#print("Enter grade_antiderivative, result=",result, " optimal=",optimal)

leaf_count_result = leaf_count(result)
leaf_count_optimal = leaf_count(optimal)

#print("leaf_count_result=",leaf_count_result)
#print("leaf_count_optimal=",leaf_count_optimal)

expnType_result = expnType(result)
expnType_optimal = expnType(optimal)

if str(result).find("Integral") != -1:
    grade = "F"
    grade_annotation = ""
else:
    if expnType_result <= expnType_optimal:
        if result.has(I):
            if optimal.has(I): #both result and optimal complex
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation ="Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal."
            else: #result contains complex but optimal is not
                grade = "C"
                grade_annotation ="Result contains complex when optimal does not."
        else: # result do not contain complex, this assumes optimal do not as well
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = ""
            else:
                grade = "B"
                grade_annotation ="Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count(result)-2*leaf_count(optimal))
    else:
        grade = "C"
        grade_annotation ="Result contains higher order function than in optimal. Order "+str(ExpnType_result-ExpnType_optimal)

```

```
#print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation
```

4.2.4 SageMath grading function

```
#Dec 24, 2019. Nasser: Ported original Maple grading function by
#           Albert Rich to use with Sagemath. This is used to
#           grade Fricas, Giac and Maxima results.
#Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#           'arctan2','floor','abs','log_integral'
#June 4, 2022 Made default grade_annotation "none" instead of "" due
#           issue later when reading the file.
#July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    """
    Return the tree size of this expression.
    """
    #print("Enter tree_size, expr is ",expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow:  #isinstance(expr,Pow):
        if expr.operands()[1]==1/2: #expr.args[1] == Rational(1,2):
            if debug: print ("expr is sqrt")
            return True
        else:
            return False
    else:
        return False
```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arccoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
    if m:
        print ("func ", func , " is special_function")
    else:
        print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric']  #[appellf1] can't find this in sage

```

```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

#thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-
try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__)
    return False


def expnType(expn):

    if debug:
        print (">>>>Enter expnType, expn=", expn)
        print (">>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

        return 1
    else:
        return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    else:
        return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or isinstan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

        leaf_count_result = tree_size(result) #leaf_count(result)
        leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

        #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger than optimal"
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)
    else:
        grade = "C"
        grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_result)

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```