

Computer algebra independent integration tests

Summer 2022 edition

5-Inverse-trig-functions/5.3-Inverse-tangent/152-5.3.6-Exponentials-
of-inverse-tangent

Nasser M. Abbasi

September 27, 2022

Compiled on September 27, 2022 at 6:14am

Contents

1	Introduction	3
2	detailed summary tables of results	19
3	Listing of integrals	117
4	Appendix	1719

Chapter 1

Introduction

Local contents

1.1	Listing of CAS systems tested	4
1.2	Results	5
1.3	Time and leaf size Performance	9
1.4	list of integrals that has no closed form antiderivative	11
1.5	List of integrals solved by CAS but has no known antiderivative	12
1.6	list of integrals solved by CAS but failed verification	13
1.7	Timing	13
1.8	Verification	14
1.9	Important notes about some of the results	14
1.10	Design of the test system	17

This report gives the result of running the computer algebra independent integration test. The download section in the appendix contains links to download the problems in plain text format used for all CAS systems.

The number of integrals in this report is [385]. This is test number [152].

1.1 Listing of CAS systems tested

The following are the CAS systems tested:

1. Mathematica 13.1 (June 29, 2022) on windows 10.
2. Rubi 4.16.1 (Dec 19, 2018) on Mathematica 13.0.1 on windows 10.
3. Maple 2022.1 (June 1, 2022) on windows 10.
4. Maxima 5.46 (April 13, 2022) using Lisp SBCL 2.1.11.debian on Linux via sagemath 9.6.
5. Fricas 1.3.8 (June 21, 2022) based on sbcl 2.1.11.debian on Linux via sagemath 9.6.
6. Giac/Xcas 1.9.0-13 (July 3, 2022) on Linux via sagemath 9.6.
7. Sympy 1.10.1 (March 20, 2022) Using Python 3.10.4 on Linux.
8. Mupad using Matlab 2021a with Symbolic Math Toolbox Version 8.7 on windows 10.

Maxima and Fricas and Giac are called using Sagemath. This was done using Sagemath `integrate` command by changing the name of the algorithm to use the different CAS systems.

Sympy was called directly from Python.

1.2 Results

Important note: A number of problems in this test suite have no antiderivative in closed form. This means the antiderivative of these integrals can not be expressed in terms of elementary, special functions or `Hypergeometric2F1` functions. `RootSum` and `RootOf` are not allowed.

If a CAS returns the above integral unevaluated within the time limit, then the result is counted as passed and assigned an A grade.

However, if CAS times out, then it is assigned an F grade even if the integral is not integrable, as this implies CAS could not determine that the integral is not integrable in the time limit.

If a CAS returns an antiderivative to such an integral, it is assigned an A grade automatically and this special result is listed in the introduction section of each individual test report to make it easy to identify as this can be important result to investigate.

The results given in in the table below reflects the above.

System	% solved	% Failed
Rubi	100.00 (385)	0.00 (0)
Mathematica	95.58 (368)	4.42 (17)
Fricas	74.29 (286)	25.71 (99)
Maple	52.73 (203)	47.27 (182)
Mupad	38.18 (147)	61.82 (238)
Maxima	34.81 (134)	65.19 (251)
Giac	30.39 (117)	69.61 (268)
Sympy	22.34 (86)	77.66 (299)

Table 1.1: Percentage solved for each CAS

The table below gives additional break down of the grading of quality of the antiderivatives generated by each CAS. The grading is given using the letters A,B,C and F with A being the best quality. The grading is accomplished by comparing the antiderivative generated with the optimal antiderivatives included in the test suite. The following table describes the meaning of these grades.

grade	description
A	Integral was solved and antiderivative is optimal in quality and leaf size.
B	Integral was solved and antiderivative is optimal in quality but leaf size is larger than twice the optimal antiderivatives leaf size.
C	Integral was solved and antiderivative is non-optimal in quality. This can be due to one or more of the following reasons <ol style="list-style-type: none"> 1. antiderivative contains a hypergeometric function and the optimal antiderivative does not. 2. antiderivative contains a special function and the optimal antiderivative does not. 3. antiderivative contains the imaginary unit and the optimal antiderivative does not.
F	Integral was not solved. Either the integral was returned unevaluated within the time limit, or it timed out, or CAS hanged or crashed or an exception was raised.


Table 1.2: Description of grading applied to integration result

Grading is implemented for all CAS systems. Based on the above, the following table summarizes the grading for this test suite.


System	% A grade	% B grade	% C grade	% F grade
Rubi	100.00	0.00	0.00	0.00
Mathematica	65.71	0.00	29.87	4.42
Fricas	59.74	14.55	0.00	25.71
Maple	29.87	21.30	1.56	47.27
Maxima	22.34	12.47	0.00	65.19
Giac	21.82	8.57	0.00	69.61
Sympy	16.10	6.23	0.00	77.66
Mupad	N/A	38.18	0.00	61.82

Table 1.3: Antiderivative Grade distribution of each CAS

The following is a Bar chart illustration of the data in the above table.



The figure below compares the CAS systems for each grade level.



The following table shows the distribution of the different types of failure for each CAS. There are 3 types of reasons why it can fail. The first is when CAS returns back the input within the time limit, which means it could not solve it. This is the typical normal failure **F**.

The second is due to time out. CAS could not solve the integral within the 3 minutes time limit which is assigned **F(-1)**.

The third is due to an exception generated. Assigned **F(-2)**. This most likely indicates an interface problem between sagemath and the CAS (applicable only to FriCAS, Maxima and

Giac) or it could be an indication of an internal error in CAS. This type of error requires more investigations to determine the cause.

System	Number failed	Percentage normal failure	Percentage time-out failure	Percentage exception failure
Rubi	0	0.00 %	0.00 %	0.00 %
Mathematica	17	100.00 %	0.00 %	0.00 %
Maple	182	100.00 %	0.00 %	0.00 %
Fricas	99	100.00 %	0.00 %	0.00 %
Giac	268	56.72 %	0.75 %	42.54 %
Maxima	251	95.62 %	0.00 %	4.38 %
Sympy	299	83.28 %	13.04 %	3.68 %
Mupad	238	99.58 %	0.42 %	0.00 %

Table 1.4: Failure statistics for each CAS

1.3 Time and leaf size Performance

The table below summarizes the performance of each CAS system in terms of time used and leaf size of results.

Mean size is the average leaf size produced by the CAS (before any normalization). The Normalized mean is relative to the mean size of the optimal anti-derivative given in the input files.

For example, if CAS has **Normalized mean** of 3, then the mean size of its leaf size is 3 times as large as the mean size of the optimal leaf size.

Median size is value of leaf size where half the values are larger than this and half are smaller (before any normalization). i.e. The Middle value.



Similarly the **Normalized median** is relative to the median leaf size of the optimal.

For example, if a CAS has Normalized median of 1.2, then its median is 1.2 as large as the median leaf size of the optimal.

System	Mean time (sec)	Mean size	Normalized mean	Median size	Normalized median
Rubi	0.10	142.49	1.01	95.00	1.00
Mathematica	0.06	85.19	0.80	76.50	0.82
Maple	0.11	300.62	2.57	87.00	1.37
Maxima	0.37	221.49	1.80	64.50	1.25
Fricas	2.98	176.45	1.31	118.50	1.00
Sympy	3.37	92.74	1.64	50.00	1.14
Giac	0.44	114.00	1.40	70.00	1.14
Mupad	0.54	68.97	1.09	52.00	0.95

Table 1.5: Time and leaf size performance for each CAS

The following are bar charts for the normalized leafsize and time used from the above table.



1.4 list of integrals that has no closed form antiderivative

{

1.5 List of integrals solved by CAS but has no known antiderivative

Rubi {}

Mathematica {}

Maple {}

Maxima {}

Fricas {}

Sympy {}

Giac {}

Mupad {}

1.6 list of integrals solved by CAS but failed verification

The following are integrals solved by CAS but the verification phase failed to verify the anti-derivative produced is correct. This does not mean necessarily that the anti-derivative is wrong, as additional methods of verification might be needed, or more time is needed (3 minutes time limit was used). These integrals are listed here to make it easier to do further investigation to determine why it was not possible to verify the result produced.

Rubi {}

Mathematica {6, 7, 8, 9, 10, 23, 24, 25, 26, 39, 40, 41, 42, 43, 56, 57, 58, 59, 60, 140, 141, 142, 143, 164, 165, 168, 169, 170, 184, 195, 196, 197, 207, 208, 209, 210, 211, 212, 302, 304, 307, 309, 311, 313, 316, 318, 329, 331, 334, 336, 365}

Maple Verification phase not implemented yet.

Maxima Verification phase not implemented yet.

Fricas Verification phase not implemented yet.

Sympy Verification phase not implemented yet.

Giac Verification phase not implemented yet.

Mupad Verification phase not implemented yet.

1.7 Timing

The command `AbsoluteTiming[]` was used in Mathematica to obtain the elapsed time for each integrate call. In Maple, the command `Usage` was used as in the following example

```
cpu_time := Usage(assign ('result_of_int',int(expr,x)),output='realtime')
```

For all other CAS systems, the elapsed time to complete each integral was found by taking the difference between the time after the call completed from the time before the call was made. This was done using Python's `time.time()` call.

All elapsed times shown are in seconds. A time limit of 3 CPU minutes was used for each integral. If the integrate command did not complete within this time limit, the integral was aborted and considered to have failed and assigned an F grade. The time used by failed integrals due to time out was not counted in the final statistics.

1.8 Verification

A verification phase was applied on the result of integration for `Rubi` and `Mathematica`.

Future version of this report will implement verification for the other CAS systems. For the integrals whose result was not run through a verification phase, it is assumed that the antiderivative was correct.

Verification phase also had 3 minutes time out. An integral whose result was not verified could still be correct, but further investigation is needed on those integrals. These integrals were marked in the summary table below and also in each integral separate section so they are easy to identify and locate.

1.9 Important notes about some of the results

1.9.1 Important note about Maxima results

Since tests were run in a batch mode, and using an automated script, then any integral where Maxima needed an interactive response from the user to answer a question during the evaluation of the integral will fail.

The exception raised is `ValueError`. Therefore Maxima results is lower than what would result if Maxima was run directly and each question was answered correctly.

The percentage of such failures were not counted for each test file, but for an example, for the `Timofeev` test file, there were about 14 such integrals out of total 705, or about 2 percent. This percentage can be higher or lower depending on the specific input test file.

Such integrals can be identified by looking at the output of the integration in each section for Maxima. The exception message will indicate the cause of error.

Maxima `integrate` was run using SageMath with the following settings set by default

```
'besselexpand : true'
'display2d : false'
'domain : complex'
'keepfloat : true'
'load(to_poly_solve)'
'load(simplify_sum)'
'load(abs_integrate)' 'load(diag)'
```

SageMath automatic loading of Maxima `abs_integrate` was found to cause some problems. So the following code was added to disable this effect.

```
from sage.interfaces.maxima_lib import maxima_lib
maxima_lib.set('extra_definite_integration_methods', '[]')
```

```
maxima_lib.set('extra_integration_methods', '[]')
```

See <https://ask.sagemath.org/question/43088/integrate-results-that-are-different-from-using-maxima/> for reference.

1.9.2 Important note about FriCAS result

There were few integrals which failed due to SageMath interface and not because FriCAS system could not do the integration.

These will fail With error `Exception raised: NotImplementedError`.

The number of such cases seems to be very small. About 1 or 2 percent of all integrals. These can be identified by looking at the exception message given in the result.

1.9.3 Important note about finding leaf size of antiderivative

For Mathematica, Rubi, and Maple, the builtin system function `LeafSize` was used to find the leaf size of each antiderivative.

The other CAS systems (SageMath and Sympy) do not have special builtin function for this purpose at this time. Therefore the leaf size for Fricas and Sympy antiderivative was determined using the following function, thanks to user `slelievre` at https://ask.sagemath.org/question/57123/could-we-have-a-leaf_count-function-in-base-sagemath/

```
def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)
```

For Sympy, which was called directly from Python, the following code was used to obtain the leafsize of its result

```
try:
    # 1.7 is a fudge factor since it is low side from actual leaf count
    leafCount = round(1.7*count_ops(anti))

except Exception as ee:
    leafCount =1
```

1.9.4 Important note about Mupad results

Matlab's symbolic toolbox does not have a leaf count function to measure the size of the antiderivative. Maple was used to determine the leaf size of Mupad output by post processing Mupad result.

Currently no grading of the antiderivative for Mupad is implemented. If it can integrate the problem, it was assigned a B grade automatically as a placeholder. In the future, when grading function is implemented for Mupad, the tests will be rerun again.

The following is an example of using Matlab's symbolic toolbox (Mupad) to solve an integral

```
integrand = evalin(symengine, 'cos(x)*sin(x)')
the_variable = evalin(symengine, 'x')
anti = int(integrand,the_variable)
```

Which gives $\sin(x)^2/2$

1.10 Design of the test system

The following diagram gives a high level view of the current test build system.



High level overview of the CAS independent integration test build system

One record (line) per one integral result. The line is CSV comma separated. This is description of each record

1. integer, the problem number.
2. integer. 0 for failed, 1 for passed, -1 for timeout, -2 for CAS specific exception. (this is not the grade field)
3. integer. Leaf size of result.
4. integer. Leaf size of the optimal antiderivative.
5. number. CPU time used to solve this integral. 0 if failed.
6. string. The integral in Latex format
7. string. The input used in CAS own syntax.
8. string. The result (antiderivative) produced by CAS in Latex format
9. string. The optimal antiderivative in Latex format.
10. integer. 0 or 1. Indicates if problem has known antiderivative or not
11. String. The result (antiderivative) in CAS own syntax.
12. String. The grade of the antiderivative. Can be "A", "B", "C", or "F"
13. String. Small string description of why the grade was given.
14. integer. 1 if result was verified or 0 if not verified.

The following fields are present only in Rubi Table file

15. integer. Number of steps used.
16. integer. Number of rules used.
17. integer. Integrand leaf size.
18. real number. Ratio. Field 16 over field 17
19. String of form "{n,n,..}" which is list of the rules used by Rubi
20. String. The optimal antiderivative in Mathematica syntax

Chapter 2

detailed summary tables of results

Local contents

2.1	List of integrals sorted by grade for each CAS	20
2.2	Detailed conclusion table per each integral for all CAS systems	26
2.3	Detailed conclusion table specific for Rubi results	104

2.1 List of integrals sorted by grade for each CAS

Local contents

2.1.1	Rubi	21
2.1.2	Mathematica	21
2.1.3	Maple	22
2.1.4	Maxima	22
2.1.5	FriCAS	23
2.1.6	Sympy	23
2.1.7	Giac	24
2.1.8	Mupad	24

2.1.1 Rubi

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385 }

B grade: { }

C grade: { }

F grade: { }

2.1.2 Mathematica

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 70, 88, 97, 134, 135, 136, 137, 138, 139, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 295, 296, 297, 298, 299, 300, 301, 303, 304, 305, 306, 307, 308, 309, 310, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 344, 345, 346, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 365, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385 }

B grade: { }

C grade: { 61, 62, 63, 64, 65, 66, 67, 68, 69, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 89, 90, 91, 92, 93, 94, 95, 96, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 140, 141, 142, 143, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, }

232, 233, 234, 235, 248, 249, 250, 251, 252, 263, 264, 265, 266, 277, 278, 279, 280, 291, 292, 293, 294, 302, 311, 343, 347 }

F grade: { 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 236, 364, 366, 367, 368, 369, 370 }

2.1.3 Maple

A grade: { 1, 2, 3, 5, 7, 8, 9, 10, 11, 12, 13, 14, 16, 17, 18, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 44, 45, 46, 47, 48, 49, 50, 51, 140, 141, 167, 168, 169, 198, 199, 200, 201, 202, 203, 204, 205, 206, 248, 249, 250, 251, 252, 256, 257, 258, 263, 264, 265, 266, 270, 271, 272, 277, 278, 279, 280, 284, 285, 286, 291, 292, 293, 294, 298, 299, 300, 301, 303, 305, 306, 308, 310, 312, 314, 315, 316, 317, 319, 321, 323, 324, 325, 326, 327, 328, 330, 332, 333, 335, 343, 347, 372, 373, 374, 375, 376, 377, 378, 380, 381, 382, 383, 384, 385 }

B grade: { 4, 6, 15, 19, 20, 21, 22, 23, 35, 36, 37, 38, 39, 40, 41, 42, 43, 52, 53, 54, 55, 56, 57, 58, 59, 60, 162, 163, 164, 165, 166, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 207, 208, 209, 210, 211, 212, 213, 214, 215, 302, 304, 307, 309, 311, 313, 318, 320, 322, 329, 331, 334, 336, 379 }

C grade: { 134, 135, 136, 137, 138, 139 }

F grade: { 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 253, 254, 255, 259, 260, 261, 262, 267, 268, 269, 273, 274, 275, 276, 281, 282, 283, 287, 288, 289, 290, 295, 296, 297, 337, 338, 339, 340, 341, 342, 344, 345, 346, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371 }

2.1.4 Maxima

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 44, 45, 46, 47, 48, 49, 50, 51, 52, 54, 55, 166, 191, 192, 193, 198, 199, 200, 201, 202, 203, 211, 248, 263, 277, 291, 301, 303, 304, 306, 307, 308, 315, 316, 317, 318, 322, 323, 325, 326, 333, 334, 335, 343, 372, 383, 384 }

B grade: { 53, 162, 163, 164, 165, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 204, 205, 206, 207, 208, 209, 210, 302, 305, 309, 319, 320, 321, 327, 336, 375, 376, 377, 385 }

C grade: { }

F grade: { 40, 41, 42, 43, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 194, 195, 196, 197,

212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 293, 294, 295, 296, 297, 298, 299, 300, 310, 311, 312, 313, 314, 324, 328, 329, 330, 331, 332, 337, 338, 339, 340, 341, 342, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 373, 374, 378, 379, 380, 381, 382 }

2.1.5 FriCAS

A grade: { 1, 2, 3, 4, 5, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 58, 59, 60, 61, 62, 63, 64, 66, 67, 68, 69, 70, 71, 72, 73, 74, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 94, 95, 96, 97, 98, 99, 100, 101, 103, 104, 105, 106, 107, 108, 109, 110, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 162, 163, 164, 165, 166, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 189, 190, 191, 192, 193, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 216, 217, 218, 219, 221, 222, 223, 226, 227, 228, 229, 231, 232, 233, 248, 249, 250, 251, 252, 256, 257, 258, 263, 264, 265, 266, 270, 271, 272, 277, 278, 279, 280, 284, 285, 286, 291, 292, 293, 294, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 319, 320, 321, 322, 325, 326, 327, 328, 329, 330, 331, 334, 335, 336, 343, 347, 372, 373, 374, 377, 378 }

B grade: { 6, 7, 23, 24, 39, 40, 56, 57, 65, 75, 93, 102, 111, 112, 167, 168, 169, 170, 185, 186, 187, 188, 194, 195, 196, 197, 212, 213, 214, 215, 220, 224, 225, 230, 234, 235, 310, 311, 312, 313, 314, 315, 316, 317, 318, 323, 324, 332, 333, 375, 376, 379, 380, 381, 384, 385 }

C grade: { }

F grade: { 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 253, 254, 255, 259, 260, 261, 262, 267, 268, 269, 273, 274, 275, 276, 281, 282, 283, 287, 288, 289, 290, 295, 296, 297, 337, 338, 339, 340, 341, 342, 344, 345, 346, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 382, 383 }

2.1.6 SymPy

A grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 27, 28, 29, 30, 31, 32, 33, 34, 44, 45, 46, 47, 48, 49, 50, 51, 141, 166, 171, 172, 173, 174, 175, 198, 199, 200, 201, 202, 248, 263, 277, 291, 301, 303, 305, 306, 308, 319, 321, 323, 324, 326, 377, 378 }

B grade: { 136, 137, 176, 177, 178, 179, 203, 204, 205, 206, 249, 250, 251, 252, 264, 265, 266, 278, 279, 292, 343, 375, 376, 379 }

C grade: { }

F grade: { 19, 20, 21, 22, 23, 24, 25, 26, 35, 36, 37, 38, 39, 40, 41, 42, 43, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 138, 139, 140, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155,

156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 167, 168, 169, 170, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 293, 294, 295, 296, 297, 298, 299, 300, 302, 304, 307, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 320, 322, 325, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 380, 381, 382, 383, 384, 385 }

2.1.7 Giac

A grade: { 2, 4, 5, 11, 12, 13, 14, 15, 16, 17, 18, 27, 28, 29, 30, 31, 32, 33, 34, 37, 38, 44, 45, 49, 50, 51, 162, 163, 164, 165, 166, 167, 168, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 189, 190, 191, 192, 193, 194, 195, 202, 207, 208, 209, 210, 248, 263, 277, 291, 301, 302, 303, 305, 306, 308, 309, 311, 313, 316, 318, 319, 321, 322, 323, 324, 325, 326, 331, 334, 343, 377 }

B grade: { 6, 7, 8, 9, 10, 39, 46, 47, 48, 169, 170, 184, 185, 196, 197, 198, 199, 200, 201, 203, 204, 205, 206, 211, 212, 320, 327, 329, 336, 375, 376, 378, 379 }

C grade: { }

F grade: { 1, 3, 19, 20, 21, 22, 23, 24, 25, 26, 35, 36, 40, 41, 42, 43, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 186, 187, 188, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 292, 293, 294, 295, 296, 297, 298, 299, 300, 304, 307, 310, 312, 314, 315, 317, 328, 330, 332, 333, 335, 337, 338, 339, 340, 341, 342, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 380, 381, 382, 383, 384, 385 }

2.1.8 Mupad

A grade: { }

B grade: { 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 166, 167, 168, 171, 172, 173, 174, 175, 176, 177, 178, 179, 198, 199, 200, 201, 202, 203, 204, 205, 206, 248, 249, 250, 251, 252, 256, 257, 258, 263, 264, 265, 266, 270, 271, 272, 277, 278, 279, 280, 284, 285, 286, 291, 292, 293, 294, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 334, 335, 336, 343, 347, 372, 373, 376, 377, 378, 379, 380, 381, 384, 385 }

C grade: { }

F grade: { 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 169, 170, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 253, 254, 255, 259, 260, 261, 262, 267, 268, 269, 273, 274, 275, 276, 281, 282, 283, 287, 288, 289, 290, 295, 296, 297, 310, 311, 312, 313, 314, 315, 316, 317, 318, 332, 333, 337, 338, 339, 340, 341, 342, 344, 345, 346, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 374, 375, 382, 383 }

2.2 Detailed conclusion table per each integral for all CAS systems

Detailed conclusion table per each integral is given by table below. The elapsed time is in seconds. For failed result it is given as F(-1) if the failure was due to timeout. It is given as F(-2) if the failure was due to an exception being raised, which could indicate a bug in the system. If the failure was due to integral not being evaluated within the time limit, then it is given just an F.

In this table, the column N.S. in the table below, which stands for **normalized size** is defined as $\frac{\text{antiderivative leaf size}}{\text{optimal antiderivative leaf size}}$. To help make the table fit, Mathematica was abbreviated to MMA.

	Problem 1	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
viated to MMA.	grade	A	A	A	A	A	A	A	F(-2)	B
	verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
	size	113	113	64	142	100	67	138	0	98
	N.S.	1	1.00	0.57	1.26	0.88	0.59	1.22	0.00	0.87
	time (sec)	N/A	0.065	0.038	0.101	0.266	1.370	3.534	0.000	0.098

Problem 2	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	56	117	81	59	119	70	85
N.S.	1	1.00	0.62	1.30	0.90	0.66	1.32	0.78	0.94
time (sec)	N/A	0.049	0.029	0.073	0.267	1.244	3.550	0.421	0.425

Problem 3	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	46	92	62	51	75	0	71
N.S.	1	1.00	0.61	1.23	0.83	0.68	1.00	0.00	0.95
time (sec)	N/A	0.036	0.025	0.063	0.267	1.943	2.142	0.000	0.411

Problem 4	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	38	72	42	43	51	53	51
N.S.	1	1.00	0.90	1.71	1.00	1.02	1.21	1.26	1.21
time (sec)	N/A	0.014	0.020	0.056	0.264	1.341	2.063	0.416	0.041

Problem 5	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	26	48	25	37	68	41	32
N.S.	1	1.00	0.90	1.66	0.86	1.28	2.34	1.41	1.10
time (sec)	N/A	0.008	0.013	0.075	0.251	1.610	0.742	0.415	0.037

Problem 6	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	A	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	29	48	18	58	53	68	32
N.S.	1	1.00	1.16	1.92	0.72	2.32	2.12	2.72	1.28
time (sec)	N/A	0.027	0.013	0.068	0.254	1.354	2.087	0.428	0.037

Problem 7	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	47	34	29	66	26	75	33
N.S.	1	1.00	1.24	0.89	0.76	1.74	0.68	1.97	0.87
time (sec)	N/A	0.029	0.022	0.079	0.264	1.200	1.459	0.420	0.037

Problem 8	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	57	53	48	83	48	153	52
N.S.	1	1.00	0.90	0.84	0.76	1.32	0.76	2.43	0.83
time (sec)	N/A	0.038	0.030	0.077	0.275	5.372	1.975	0.422	0.040

Problem 9	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	70	75	67	92	75	161	74
N.S.	1	1.00	0.78	0.83	0.74	1.02	0.83	1.79	0.82
time (sec)	N/A	0.051	0.037	0.087	0.251	4.203	2.103	0.406	0.036

Problem 10	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	76	97	86	101	122	237	95
N.S.	1	1.00	0.67	0.86	0.76	0.89	1.08	2.10	0.84
time (sec)	N/A	0.063	0.042	0.079	0.252	4.921	3.501	0.432	0.031

Problem 11	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	48	63	56	46	41	46	43
N.S.	1	1.00	1.00	1.31	1.17	0.96	0.85	0.96	0.90
time (sec)	N/A	0.028	0.017	0.100	0.460	3.282	0.058	0.416	0.418

Problem 12	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	39	55	47	37	31	37	36
N.S.	1	1.00	1.00	1.41	1.21	0.95	0.79	0.95	0.92
time (sec)	N/A	0.023	0.011	0.076	0.463	4.105	0.062	0.394	0.421

Problem 13	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	29	46	38	29	22	29	27
N.S.	1	1.00	1.00	1.59	1.31	1.00	0.76	1.00	0.93
time (sec)	N/A	0.016	0.010	0.095	0.478	2.995	0.050	0.406	0.062

Problem 14	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	30	30	28	21	12	15	19
N.S.	1	1.00	1.58	1.58	1.47	1.11	0.63	0.79	1.00
time (sec)	N/A	0.007	0.010	0.060	0.485	1.801	0.050	0.406	0.043

Problem 15	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	13	33	21	15	17	12	14
N.S.	1	1.00	1.00	2.54	1.62	1.15	1.31	0.92	1.08
time (sec)	N/A	0.015	0.006	0.071	0.468	1.871	0.086	0.412	0.439

Problem 16	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	26	26	26	43	31	26	32	21	17
N.S.	1	1.00	1.00	1.65	1.19	1.00	1.23	0.81	0.65
time (sec)	N/A	0.017	0.008	0.107	0.534	2.729	0.083	0.416	0.062

Problem 17	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	36	52	42	39	42	31	27
N.S.	1	1.00	1.00	1.44	1.17	1.08	1.17	0.86	0.75
time (sec)	N/A	0.020	0.009	0.109	0.474	1.862	0.120	0.399	0.079

Problem 18	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	48	48	48	60	51	47	54	39	34
N.S.	1	1.00	1.00	1.25	1.06	0.98	1.12	0.81	0.71
time (sec)	N/A	0.023	0.011	0.089	0.488	2.621	0.115	0.390	0.072

Problem 19	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	137	80	286	114	88	0	0	137
N.S.	1	1.00	0.58	2.09	0.83	0.64	0.00	0.00	1.00
time (sec)	N/A	0.438	0.046	0.132	0.289	2.381	0.000	0.000	0.456

Problem 20	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	63	236	95	80	0	0	114
N.S.	1	1.00	0.62	2.31	0.93	0.78	0.00	0.00	1.12
time (sec)	N/A	0.399	0.039	0.119	0.270	2.644	0.000	0.000	0.063

Problem 21	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	54	193	76	72	0	0	104
N.S.	1	1.00	0.59	2.10	0.83	0.78	0.00	0.00	1.13
time (sec)	N/A	0.227	0.043	0.105	0.277	2.340	0.000	0.000	0.427

Problem 22	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	42	128	57	60	0	0	72
N.S.	1	1.00	0.70	2.13	0.95	1.00	0.00	0.00	1.20
time (sec)	N/A	0.034	0.028	0.101	0.351	1.872	0.000	0.000	0.422

Problem 23	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	55	101	46	100	0	0	73
N.S.	1	1.00	1.08	1.98	0.90	1.96	0.00	0.00	1.43
time (sec)	N/A	0.547	0.029	0.076	0.303	2.169	0.000	0.000	0.428

Problem 24	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	61	80	60	109	0	0	75
N.S.	1	1.00	0.97	1.27	0.95	1.73	0.00	0.00	1.19
time (sec)	N/A	0.464	0.038	0.111	0.286	2.248	0.000	0.000	0.059

Problem 25	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	79	105	81	130	0	0	99
N.S.	1	1.00	0.86	1.14	0.88	1.41	0.00	0.00	1.08
time (sec)	N/A	0.508	0.056	0.131	0.270	3.097	0.000	0.000	0.426

Problem 26	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	117	117	89	141	100	139	0	0	116
N.S.	1	1.00	0.76	1.21	0.85	1.19	0.00	0.00	0.99
time (sec)	N/A	0.513	0.054	0.130	0.267	3.341	0.000	0.000	0.415

Problem 27	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	65	67	77	70	56	60	60
N.S.	1	1.00	1.00	1.03	1.18	1.08	0.86	0.92	0.92
time (sec)	N/A	0.035	0.031	0.099	0.480	2.197	0.113	0.408	0.431

Problem 28	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	58	67	62	44	51	51
N.S.	1	1.00	1.00	1.09	1.26	1.17	0.83	0.96	0.96
time (sec)	N/A	0.028	0.024	0.090	0.497	2.262	0.109	0.397	0.059

Problem 29	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	45	45	45	50	60	53	36	43	43
N.S.	1	1.00	1.00	1.11	1.33	1.18	0.80	0.96	0.96
time (sec)	N/A	0.020	0.019	0.073	0.481	3.619	0.092	0.402	0.064

Problem 30	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	31	31	42	33	44	43	22	25	32
N.S.	1	1.00	1.35	1.06	1.42	1.39	0.71	0.81	1.03
time (sec)	N/A	0.010	0.018	0.085	0.482	1.651	0.093	0.403	0.427

Problem 31	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	22	18	10	13	14
N.S.	1	1.00	1.00	0.94	1.38	1.12	0.62	0.81	0.88
time (sec)	N/A	0.017	0.008	0.091	0.478	2.431	0.112	0.406	0.077

Problem 32	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	38	44	53	60	44	35	37
N.S.	1	1.00	1.00	1.16	1.39	1.58	1.16	0.92	0.97
time (sec)	N/A	0.021	0.018	0.088	0.483	1.651	0.156	0.417	0.450

Problem 33	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	52	53	69	77	58	46	43
N.S.	1	1.00	1.00	1.02	1.33	1.48	1.12	0.88	0.83
time (sec)	N/A	0.025	0.029	0.118	0.491	2.051	0.181	0.417	0.468

Problem 34	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	62	61	77	86	70	54	55
N.S.	1	1.00	1.00	0.98	1.24	1.39	1.13	0.87	0.89
time (sec)	N/A	0.029	0.028	0.105	0.482	3.122	0.198	0.411	0.130

Problem 35	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	56	241	76	59	0	0	85
N.S.	1	1.00	0.62	2.68	0.84	0.66	0.00	0.00	0.94
time (sec)	N/A	0.049	0.031	0.105	0.503	2.389	0.000	0.000	0.063

Problem 36	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	75	75	46	167	59	51	0	0	71
N.S.	1	1.00	0.61	2.23	0.79	0.68	0.00	0.00	0.95
time (sec)	N/A	0.034	0.024	0.078	0.482	2.050	0.000	0.000	0.416

Problem 37	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	38	150	42	43	0	53	51
N.S.	1	1.00	0.90	3.57	1.00	1.02	0.00	1.26	1.21
time (sec)	N/A	0.014	0.021	0.075	0.483	2.302	0.000	0.423	0.404

Problem 38	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	26	100	25	37	0	41	32
N.S.	1	1.00	0.90	3.45	0.86	1.28	0.00	1.41	1.10
time (sec)	N/A	0.007	0.014	0.074	0.469	2.264	0.000	0.407	0.405

Problem 39	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	25	25	29	121	26	58	0	68	32
N.S.	1	1.00	1.16	4.84	1.04	2.32	0.00	2.72	1.28
time (sec)	N/A	0.025	0.012	0.074	0.478	1.084	0.000	0.412	0.038

Problem 40	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F(-2)	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	47	195	0	66	0	0	33
N.S.	1	1.00	1.24	5.13	0.00	1.74	0.00	0.00	0.87
time (sec)	N/A	0.025	0.020	0.074	0.000	1.924	0.000	0.000	0.036

Problem 41	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	57	217	0	83	0	0	52
N.S.	1	1.00	0.90	3.44	0.00	1.32	0.00	0.00	0.83
time (sec)	N/A	0.036	0.030	0.070	0.000	1.571	0.000	0.000	0.039

Problem 42	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F(-2)	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	70	273	0	92	0	0	74
N.S.	1	1.00	0.78	3.03	0.00	1.02	0.00	0.00	0.82
time (sec)	N/A	0.049	0.035	0.072	0.000	1.506	0.000	0.000	0.035

Problem 43	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	113	113	76	290	0	101	0	0	95
N.S.	1	1.00	0.67	2.57	0.00	0.89	0.00	0.00	0.84
time (sec)	N/A	0.070	0.042	0.076	0.000	2.485	0.000	0.000	0.033

Problem 44	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	49	48	44	46	41	68	43
N.S.	1	1.00	1.00	0.98	0.90	0.94	0.84	1.39	0.88
time (sec)	N/A	0.027	0.015	0.081	0.268	2.641	0.063	0.413	0.056

Problem 45	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	40	35	37	31	58	36
N.S.	1	1.00	1.00	1.00	0.88	0.92	0.78	1.45	0.90
time (sec)	N/A	0.022	0.011	0.072	0.256	1.883	0.063	0.412	0.415

Problem 46	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	31	28	29	22	52	27
N.S.	1	1.00	1.00	1.03	0.93	0.97	0.73	1.73	0.90
time (sec)	N/A	0.016	0.010	0.069	0.273	1.671	0.052	0.414	0.420

Problem 47	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	30	19	16	21	14	65	19
N.S.	1	1.00	1.50	0.95	0.80	1.05	0.70	3.25	0.95
time (sec)	N/A	0.008	0.010	0.073	0.280	1.557	0.052	0.396	0.409

Problem 48	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	14	14	14	14	12	15	17	44	14
N.S.	1	1.00	1.00	1.00	0.86	1.07	1.21	3.14	1.00
time (sec)	N/A	0.015	0.006	0.072	0.263	1.806	0.075	0.414	0.451

Problem 49	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	27	27	27	25	34	26	32	34	17
N.S.	1	1.00	1.00	0.93	1.26	0.96	1.19	1.26	0.63
time (sec)	N/A	0.017	0.008	0.104	0.256	2.186	0.079	0.410	0.420

Problem 50	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	37	34	50	39	42	54	26
N.S.	1	1.00	1.00	0.92	1.35	1.05	1.14	1.46	0.70
time (sec)	N/A	0.019	0.009	0.085	0.260	2.044	0.101	0.411	0.068

Problem 51	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	49	44	57	47	54	67	33
N.S.	1	1.00	1.00	0.90	1.16	0.96	1.10	1.37	0.67
time (sec)	N/A	0.024	0.012	0.104	0.256	1.739	0.116	0.407	0.430

Problem 52	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	137	137	80	682	216	88	0	0	138
N.S.	1	1.00	0.58	4.98	1.58	0.64	0.00	0.00	1.01
time (sec)	N/A	0.427	0.041	0.125	0.483	1.839	0.000	0.000	0.466

Problem 53	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	63	619	181	80	0	0	115
N.S.	1	1.00	0.62	6.07	1.77	0.78	0.00	0.00	1.13
time (sec)	N/A	0.400	0.038	0.119	0.489	2.060	0.000	0.000	0.071

Problem 54	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	60	463	112	72	0	0	105
N.S.	1	1.00	0.65	5.03	1.22	0.78	0.00	0.00	1.14
time (sec)	N/A	0.232	0.031	0.101	0.468	2.560	0.000	0.000	0.421

Problem 55	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	42	257	65	60	0	0	73
N.S.	1	1.00	0.70	4.28	1.08	1.00	0.00	0.00	1.22
time (sec)	N/A	0.031	0.027	0.110	0.467	2.634	0.000	0.000	0.416

Problem 56	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	55	649	0	100	0	0	74
N.S.	1	1.00	1.06	12.48	0.00	1.92	0.00	0.00	1.42
time (sec)	N/A	0.467	0.032	0.090	0.000	5.002	0.000	0.000	0.433

Problem 57	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	64	64	61	580	0	109	0	0	76
N.S.	1	1.00	0.95	9.06	0.00	1.70	0.00	0.00	1.19
time (sec)	N/A	0.452	0.038	0.101	0.000	4.649	0.000	0.000	0.422

Problem 58	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	79	551	0	130	0	0	100
N.S.	1	1.00	0.85	5.92	0.00	1.40	0.00	0.00	1.08
time (sec)	N/A	0.483	0.059	0.118	0.000	5.204	0.000	0.000	0.432

Problem 59	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	118	118	89	829	0	139	0	0	117
N.S.	1	1.00	0.75	7.03	0.00	1.18	0.00	0.00	0.99
time (sec)	N/A	0.497	0.054	0.124	0.000	3.926	0.000	0.000	0.066

Problem 60	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	139	139	95	937	0	146	0	0	139
N.S.	1	1.00	0.68	6.74	0.00	1.05	0.00	0.00	1.00
time (sec)	N/A	0.534	0.060	0.124	0.000	2.237	0.000	0.000	0.434

Problem 61	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	339	339	82	0	0	244	0	0	-1
N.S.	1	1.00	0.24	0.00	0.00	0.72	0.00	0.00	-0.00
time (sec)	N/A	0.170	0.027	0.015	0.000	2.047	0.000	0.000	0.000

Problem 62	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	295	295	63	0	0	236	0	0	-1
N.S.	1	1.00	0.21	0.00	0.00	0.80	0.00	0.00	-0.00
time (sec)	N/A	0.132	0.014	0.006	0.000	2.826	0.000	0.000	0.000

Problem 63	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	268	268	41	0	0	209	0	0	-1
N.S.	1	1.00	0.15	0.00	0.00	0.78	0.00	0.00	-0.00
time (sec)	N/A	0.109	0.022	0.006	0.000	2.152	0.000	0.000	0.000

Problem 64	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	267	267	97	0	0	243	0	0	-1
N.S.	1	1.00	0.36	0.00	0.00	0.91	0.00	0.00	-0.00
time (sec)	N/A	0.126	0.027	0.005	0.000	2.681	0.000	0.000	0.000

Problem 65	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	71	0	0	151	0	0	-1
N.S.	1	1.00	0.77	0.00	0.00	1.64	0.00	0.00	-0.01
time (sec)	N/A	0.023	0.012	0.006	0.000	3.084	0.000	0.000	0.000

Problem 66	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	81	0	0	175	0	0	-1
N.S.	1	1.00	0.61	0.00	0.00	1.33	0.00	0.00	-0.01
time (sec)	N/A	0.030	0.015	0.006	0.000	3.727	0.000	0.000	0.000

Problem 67	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	170	93	0	0	184	0	0	-1
N.S.	1	1.00	0.55	0.00	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.045	0.018	0.006	0.000	2.698	0.000	0.000	0.000

Problem 68	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	99	0	0	192	0	0	-1
N.S.	1	1.00	0.49	0.00	0.00	0.95	0.00	0.00	-0.00
time (sec)	N/A	0.057	0.020	0.006	0.000	2.739	0.000	0.000	0.000

Problem 69	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	240	240	111	0	0	200	0	0	-1
N.S.	1	1.00	0.46	0.00	0.00	0.83	0.00	0.00	-0.00
time (sec)	N/A	0.072	0.026	0.006	0.000	1.388	0.000	0.000	0.000

Problem 70	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	337	228	0	0	254	0	0	-1
N.S.	1	1.00	0.68	0.00	0.00	0.75	0.00	0.00	-0.00
time (sec)	N/A	0.158	0.240	0.009	0.000	2.586	0.000	0.000	0.000

Problem 71	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	339	339	82	0	0	247	0	0	-1
N.S.	1	1.00	0.24	0.00	0.00	0.73	0.00	0.00	-0.00
time (sec)	N/A	0.154	0.033	0.009	0.000	2.086	0.000	0.000	0.000

Problem 72	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	295	295	61	0	0	239	0	0	-1
N.S.	1	1.00	0.21	0.00	0.00	0.81	0.00	0.00	-0.00
time (sec)	N/A	0.126	0.014	0.010	0.000	3.296	0.000	0.000	0.000

Problem 73	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	268	268	41	0	0	215	0	0	-1
N.S.	1	1.00	0.15	0.00	0.00	0.80	0.00	0.00	-0.00
time (sec)	N/A	0.105	0.025	0.009	0.000	2.298	0.000	0.000	0.000

Problem 74	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	267	267	96	0	0	243	0	0	-1
N.S.	1	1.00	0.36	0.00	0.00	0.91	0.00	0.00	-0.00
time (sec)	N/A	0.120	0.023	0.008	0.000	3.516	0.000	0.000	0.000

Problem 75	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	68	0	0	157	0	0	-1
N.S.	1	1.00	0.74	0.00	0.00	1.71	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.011	0.007	0.000	1.527	0.000	0.000	0.000

Problem 76	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	81	0	0	179	0	0	-1
N.S.	1	1.00	0.61	0.00	0.00	1.36	0.00	0.00	-0.01
time (sec)	N/A	0.031	0.013	0.007	0.000	1.885	0.000	0.000	0.000

Problem 77	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	170	93	0	0	187	0	0	-1
N.S.	1	1.00	0.55	0.00	0.00	1.10	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.017	0.010	0.000	2.727	0.000	0.000	0.000

Problem 78	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	99	0	0	195	0	0	-1
N.S.	1	1.00	0.49	0.00	0.00	0.97	0.00	0.00	-0.00
time (sec)	N/A	0.059	0.021	0.009	0.000	3.068	0.000	0.000	0.000

Problem 79	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-2)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	373	373	96	0	0	251	0	0	-1
N.S.	1	1.00	0.26	0.00	0.00	0.67	0.00	0.00	-0.00
time (sec)	N/A	0.180	0.032	0.013	0.000	1.880	0.000	0.000	0.000

Problem 80	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-2)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	371	371	86	0	0	244	0	0	-1
N.S.	1	1.00	0.23	0.00	0.00	0.66	0.00	0.00	-0.00
time (sec)	N/A	0.177	0.024	0.010	0.000	1.948	0.000	0.000	0.000

Problem 81	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	324	324	72	0	0	236	0	0	-1
N.S.	1	1.00	0.22	0.00	0.00	0.73	0.00	0.00	-0.00
time (sec)	N/A	0.150	0.026	0.010	0.000	1.963	0.000	0.000	0.000

Problem 82	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	299	299	41	0	0	209	0	0	-1
N.S.	1	1.00	0.14	0.00	0.00	0.70	0.00	0.00	-0.00
time (sec)	N/A	0.125	0.032	0.008	0.000	2.557	0.000	0.000	0.000

Problem 83	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	293	293	112	0	0	267	0	0	-1
N.S.	1	1.00	0.38	0.00	0.00	0.91	0.00	0.00	-0.00
time (sec)	N/A	0.156	0.028	0.008	0.000	1.248	0.000	0.000	0.000

Problem 84	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	87	0	0	152	0	0	-1
N.S.	1	1.00	0.72	0.00	0.00	1.26	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.015	0.009	0.000	1.945	0.000	0.000	0.000

Problem 85	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	99	0	0	176	0	0	-1
N.S.	1	1.00	0.61	0.00	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.037	0.018	0.009	0.000	2.019	0.000	0.000	0.000

Problem 86	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	203	106	0	0	184	0	0	-1
N.S.	1	1.00	0.52	0.00	0.00	0.91	0.00	0.00	-0.00
time (sec)	N/A	0.058	0.024	0.009	0.000	1.935	0.000	0.000	0.000

Problem 87	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	233	233	118	0	0	192	0	0	-1
N.S.	1	1.00	0.51	0.00	0.00	0.82	0.00	0.00	-0.00
time (sec)	N/A	0.072	0.027	0.010	0.000	1.929	0.000	0.000	0.000

Problem 88	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	337	272	0	0	255	0	0	-1
N.S.	1	1.00	0.81	0.00	0.00	0.76	0.00	0.00	-0.00
time (sec)	N/A	0.157	0.285	0.008	0.000	1.312	0.000	0.000	0.000

Problem 89	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	339	339	73	0	0	247	0	0	-1
N.S.	1	1.00	0.22	0.00	0.00	0.73	0.00	0.00	-0.00
time (sec)	N/A	0.159	0.025	0.010	0.000	1.982	0.000	0.000	0.000

Problem 90	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	295	295	63	0	0	238	0	0	-1
N.S.	1	1.00	0.21	0.00	0.00	0.81	0.00	0.00	-0.00
time (sec)	N/A	0.137	0.011	0.009	0.000	1.760	0.000	0.000	0.000

Problem 91	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	268	268	41	0	0	213	0	0	-1
N.S.	1	1.00	0.15	0.00	0.00	0.79	0.00	0.00	-0.00
time (sec)	N/A	0.113	0.024	0.007	0.000	2.144	0.000	0.000	0.000

Problem 92	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	267	267	96	0	0	243	0	0	-1
N.S.	1	1.00	0.36	0.00	0.00	0.91	0.00	0.00	-0.00
time (sec)	N/A	0.125	0.020	0.006	0.000	1.274	0.000	0.000	0.000

Problem 93	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	69	0	0	156	0	0	-1
N.S.	1	1.00	0.75	0.00	0.00	1.70	0.00	0.00	-0.01
time (sec)	N/A	0.025	0.010	0.009	0.000	4.966	0.000	0.000	0.000

Problem 94	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	81	0	0	178	0	0	-1
N.S.	1	1.00	0.61	0.00	0.00	1.35	0.00	0.00	-0.01
time (sec)	N/A	0.031	0.013	0.009	0.000	3.470	0.000	0.000	0.000

Problem 95	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	170	92	0	0	187	0	0	-1
N.S.	1	1.00	0.54	0.00	0.00	1.10	0.00	0.00	-0.01
time (sec)	N/A	0.046	0.017	0.013	0.000	4.573	0.000	0.000	0.000

Problem 96	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	99	0	0	195	0	0	-1
N.S.	1	1.00	0.49	0.00	0.00	0.97	0.00	0.00	-0.00
time (sec)	N/A	0.059	0.023	0.010	0.000	2.386	0.000	0.000	0.000

Problem 97	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	337	337	272	0	0	251	0	0	-1
N.S.	1	1.00	0.81	0.00	0.00	0.74	0.00	0.00	-0.00
time (sec)	N/A	0.158	0.281	0.013	0.000	2.133	0.000	0.000	0.000

Problem 98	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	339	339	73	0	0	243	0	0	-1
N.S.	1	1.00	0.22	0.00	0.00	0.72	0.00	0.00	-0.00
time (sec)	N/A	0.159	0.024	0.010	0.000	2.499	0.000	0.000	0.000

Problem 99	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	295	295	63	0	0	236	0	0	-1
N.S.	1	1.00	0.21	0.00	0.00	0.80	0.00	0.00	-0.00
time (sec)	N/A	0.135	0.013	0.011	0.000	4.536	0.000	0.000	0.000

Problem 100	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	268	268	39	0	0	209	0	0	-1
N.S.	1	1.00	0.15	0.00	0.00	0.78	0.00	0.00	-0.00
time (sec)	N/A	0.114	0.038	0.010	0.000	2.820	0.000	0.000	0.000

Problem 101	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	267	267	97	0	0	243	0	0	-1
N.S.	1	1.00	0.36	0.00	0.00	0.91	0.00	0.00	-0.00
time (sec)	N/A	0.131	0.022	0.011	0.000	3.226	0.000	0.000	0.000

Problem 102	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	69	0	0	152	0	0	-1
N.S.	1	1.00	0.75	0.00	0.00	1.65	0.00	0.00	-0.01
time (sec)	N/A	0.024	0.012	0.014	0.000	1.980	0.000	0.000	0.000

Problem 103	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	132	132	81	0	0	176	0	0	-1
N.S.	1	1.00	0.61	0.00	0.00	1.33	0.00	0.00	-0.01
time (sec)	N/A	0.032	0.014	0.012	0.000	3.475	0.000	0.000	0.000

Problem 104	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	170	170	93	0	0	184	0	0	-1
N.S.	1	1.00	0.55	0.00	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.047	0.018	0.012	0.000	2.814	0.000	0.000	0.000

Problem 105	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	99	0	0	192	0	0	-1
N.S.	1	1.00	0.49	0.00	0.00	0.95	0.00	0.00	-0.00
time (sec)	N/A	0.060	0.023	0.012	0.000	4.654	0.000	0.000	0.000

Problem 106	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	373	373	100	0	0	304	0	0	-1
N.S.	1	1.00	0.27	0.00	0.00	0.82	0.00	0.00	-0.00
time (sec)	N/A	0.182	0.030	0.015	0.000	3.128	0.000	0.000	0.000

Problem 107	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	371	371	91	0	0	296	0	0	-1
N.S.	1	1.00	0.25	0.00	0.00	0.80	0.00	0.00	-0.00
time (sec)	N/A	0.177	0.026	0.009	0.000	2.466	0.000	0.000	0.000

Problem 108	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	324	324	63	0	0	289	0	0	-1
N.S.	1	1.00	0.19	0.00	0.00	0.89	0.00	0.00	-0.00
time (sec)	N/A	0.153	0.019	0.007	0.000	5.488	0.000	0.000	0.000

Problem 109	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	299	299	39	0	0	261	0	0	-1
N.S.	1	1.00	0.13	0.00	0.00	0.87	0.00	0.00	-0.00
time (sec)	N/A	0.126	0.037	0.006	0.000	4.226	0.000	0.000	0.000

Problem 110	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	293	293	106	0	0	329	0	0	-1
N.S.	1	1.00	0.36	0.00	0.00	1.12	0.00	0.00	-0.00
time (sec)	N/A	0.158	0.042	0.007	0.000	2.353	0.000	0.000	0.000

Problem 111	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	69	0	0	212	0	0	-1
N.S.	1	1.00	0.57	0.00	0.00	1.75	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.014	0.007	0.000	2.952	0.000	0.000	0.000

Problem 112	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	81	0	0	238	0	0	-1
N.S.	1	1.00	0.50	0.00	0.00	1.46	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.015	0.007	0.000	2.258	0.000	0.000	0.000

Problem 113	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	203	203	93	0	0	246	0	0	-1
N.S.	1	1.00	0.46	0.00	0.00	1.21	0.00	0.00	-0.00
time (sec)	N/A	0.059	0.020	0.010	0.000	4.811	0.000	0.000	0.000

Problem 114	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-2)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	233	233	99	0	0	254	0	0	-1
N.S.	1	1.00	0.42	0.00	0.00	1.09	0.00	0.00	-0.00
time (sec)	N/A	0.075	0.024	0.013	0.000	2.438	0.000	0.000	0.000

Problem 115	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	319	319	73	0	0	208	0	0	-1
N.S.	1	1.00	0.23	0.00	0.00	0.65	0.00	0.00	-0.00
time (sec)	N/A	0.292	0.027	0.005	0.000	3.188	0.000	0.000	0.000

Problem 116	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	278	278	57	0	0	195	0	0	-1
N.S.	1	1.00	0.21	0.00	0.00	0.70	0.00	0.00	-0.00
time (sec)	N/A	0.260	0.016	0.007	0.000	1.696	0.000	0.000	0.000

Problem 117	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	262	262	34	0	0	195	0	0	-1
N.S.	1	1.00	0.13	0.00	0.00	0.74	0.00	0.00	-0.00
time (sec)	N/A	0.245	0.016	0.006	0.000	2.568	0.000	0.000	0.000

Problem 118	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	430	430	90	0	0	339	0	0	-1
N.S.	1	1.00	0.21	0.00	0.00	0.79	0.00	0.00	-0.00
time (sec)	N/A	0.363	0.020	0.004	0.000	1.755	0.000	0.000	0.000

Problem 119	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	253	253	64	0	0	211	0	0	-1
N.S.	1	1.00	0.25	0.00	0.00	0.83	0.00	0.00	-0.00
time (sec)	N/A	0.121	0.010	0.005	0.000	1.193	0.000	0.000	0.000

Problem 120	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	280	280	72	0	0	234	0	0	-1
N.S.	1	1.00	0.26	0.00	0.00	0.84	0.00	0.00	-0.00
time (sec)	N/A	0.136	0.012	0.005	0.000	2.289	0.000	0.000	0.000

Problem 121	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	319	319	81	0	0	243	0	0	-1
N.S.	1	1.00	0.25	0.00	0.00	0.76	0.00	0.00	-0.00
time (sec)	N/A	0.153	0.014	0.004	0.000	1.584	0.000	0.000	0.000

Problem 122	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	177	177	73	0	0	117	0	0	-1
N.S.	1	1.00	0.41	0.00	0.00	0.66	0.00	0.00	-0.01
time (sec)	N/A	0.040	0.025	0.005	0.000	1.404	0.000	0.000	0.000

Problem 123	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	54	0	0	116	0	0	-1
N.S.	1	1.00	0.39	0.00	0.00	0.83	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.015	0.005	0.000	1.429	0.000	0.000	0.000

Problem 124	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	34	0	0	104	0	0	-1
N.S.	1	1.00	0.29	0.00	0.00	0.90	0.00	0.00	-0.01
time (sec)	N/A	0.014	0.018	0.006	0.000	2.131	0.000	0.000	0.000

Problem 125	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	90	0	0	145	0	0	-1
N.S.	1	1.00	0.55	0.00	0.00	0.89	0.00	0.00	-0.01
time (sec)	N/A	0.026	0.018	0.005	0.000	2.358	0.000	0.000	0.000

Problem 126	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	111	111	59	0	0	120	0	0	-1
N.S.	1	1.00	0.53	0.00	0.00	1.08	0.00	0.00	-0.01
time (sec)	N/A	0.021	0.009	0.004	0.000	2.517	0.000	0.000	0.000

Problem 127	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	69	0	0	138	0	0	-1
N.S.	1	1.00	0.49	0.00	0.00	0.97	0.00	0.00	-0.01
time (sec)	N/A	0.028	0.011	0.004	0.000	1.856	0.000	0.000	0.000

Problem 128	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	741	741	83	0	0	435	0	0	-1
N.S.	1	1.00	0.11	0.00	0.00	0.59	0.00	0.00	-0.00
time (sec)	N/A	0.553	0.029	0.007	0.000	2.372	0.000	0.000	0.000

Problem 129	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	689	689	63	0	0	428	0	0	-1
N.S.	1	1.00	0.09	0.00	0.00	0.62	0.00	0.00	-0.00
time (sec)	N/A	0.378	0.014	0.007	0.000	2.775	0.000	0.000	0.000

Problem 130	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	674	674	41	0	0	383	0	0	-1
N.S.	1	1.00	0.06	0.00	0.00	0.57	0.00	0.00	-0.00
time (sec)	N/A	0.330	0.021	0.003	0.000	2.139	0.000	0.000	0.000

Problem 131	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	859	859	97	0	0	509	0	0	-1
N.S.	1	1.00	0.11	0.00	0.00	0.59	0.00	0.00	-0.00
time (sec)	N/A	0.408	0.026	0.005	0.000	2.287	0.000	0.000	0.000

Problem 132	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	328	328	71	0	0	345	0	0	-1
N.S.	1	1.00	0.22	0.00	0.00	1.05	0.00	0.00	-0.00
time (sec)	N/A	0.094	0.012	0.003	0.000	2.210	0.000	0.000	0.000

Problem 133	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	364	364	84	0	0	381	0	0	-1
N.S.	1	1.00	0.23	0.00	0.00	1.05	0.00	0.00	-0.00
time (sec)	N/A	0.119	0.015	0.006	0.000	2.801	0.000	0.000	0.000

Problem 134	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	114	94	748	0	0	0	0	-1
N.S.	1	1.00	0.82	6.56	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.032	0.164	0.000	0.000	0.000	0.000	0.000

Problem 135	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	58	417	0	0	0	0	-1
N.S.	1	1.00	1.16	8.34	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.031	0.018	0.095	0.000	0.000	0.000	0.000	0.000

Problem 136	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	175	0	0	129	0	-1
N.S.	1	1.00	0.74	4.49	0.00	0.00	3.31	0.00	-0.03
time (sec)	N/A	0.016	0.007	0.099	0.000	0.000	2.050	0.000	0.000

Problem 137	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	B	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	39	39	29	158	0	0	136	0	-1
N.S.	1	1.00	0.74	4.05	0.00	0.00	3.49	0.00	-0.03
time (sec)	N/A	0.017	0.007	0.085	0.000	0.000	2.630	0.000	0.000

Problem 138	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	58	428	0	0	0	0	-1
N.S.	1	1.00	1.16	8.56	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.017	0.105	0.000	0.000	0.000	0.000	0.000

Problem 139	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	C	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	94	1196	0	0	0	0	-1
N.S.	1	1.00	0.82	10.40	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.061	0.030	0.173	0.000	0.000	0.000	0.000	0.000

Problem 140	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	F	F(-2)	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	159	159	113	146	0	0	0	0	-1
N.S.	1	1.00	0.71	0.92	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.661	0.063	0.077	0.000	0.000	0.000	0.000	0.000

Problem 141	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	F	A	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	85	71	0	0	95	0	-1
N.S.	1	1.00	1.08	0.90	0.00	0.00	1.20	0.00	-0.01
time (sec)	N/A	0.030	0.033	0.064	0.000	0.000	1.789	0.000	0.000

Problem 142	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	85	0	0	0	0	0	-1
N.S.	1	1.00	1.08	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.029	0.033	0.017	0.000	0.000	0.000	0.000	0.000

Problem 143	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	F	F	F(-2)	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	159	159	113	0	0	0	0	0	-1
N.S.	1	1.00	0.71	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.590	0.054	0.017	0.000	0.000	0.000	0.000	0.000

Problem 144	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-1)	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.024	0.216	0.005	0.000	0.000	0.000	0.000	0.000

Problem 145	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-2)	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.018	0.175	0.003	0.000	0.000	0.000	0.000	0.000

Problem 146	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.019	0.170	0.003	0.000	0.000	0.000	0.000	0.000

Problem 147	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.019	0.193	0.004	0.000	0.000	0.000	0.000	0.000

Problem 148	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.019	0.190	0.004	0.000	0.000	0.000	0.000	0.000

Problem 149	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-2)	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.019	0.224	0.004	0.000	0.000	0.000	0.000	0.000

Problem 150	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.018	0.148	0.005	0.000	0.000	0.000	0.000	0.000

Problem 151	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.018	0.150	0.006	0.000	0.000	0.000	0.000	0.000

Problem 152	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-1)	F(-2)	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	36	36	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.03
time (sec)	N/A	0.020	0.168	0.003	0.000	0.000	0.000	0.000	0.000

Problem 153	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.023	0.175	0.020	0.000	0.000	0.000	0.000	0.000

Problem 154	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	158	0	0	0	0	0	-1
N.S.	1	1.00	0.92	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.084	0.611	0.010	0.000	0.000	0.000	0.000	0.000

Problem 155	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	159	159	116	0	0	0	0	0	-1
N.S.	1	1.00	0.73	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.061	0.046	0.007	0.000	0.000	0.000	0.000	0.000

Problem 156	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	107	107	105	0	0	0	0	0	-1
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.032	0.022	0.007	0.000	0.000	0.000	0.000	0.000

Problem 157	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	71	71	53	0	0	0	0	0	-1
N.S.	1	1.00	0.75	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.009	0.026	0.005	0.000	0.000	0.000	0.000	0.000

Problem 158	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	125	125	106	0	0	0	0	0	-1
N.S.	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.038	0.020	0.004	0.000	0.000	0.000	0.000	0.000

Problem 159	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	82	0	0	0	0	0	-1
N.S.	1	1.00	1.04	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.022	0.013	0.004	0.000	0.000	0.000	0.000	0.000

Problem 160	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	114	0	0	0	0	0	-1
N.S.	1	1.00	0.95	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.035	0.026	0.004	0.000	0.000	0.000	0.000	0.000

Problem 161	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	119	0	0	0	0	0	-1
N.S.	1	1.00	0.70	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.051	0.044	0.004	0.000	0.000	0.000	0.000	0.000

Problem 162	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	276	276	157	1260	749	177	0	205	-1
N.S.	1	1.00	0.57	4.57	2.71	0.64	0.00	0.74	-0.00
time (sec)	N/A	0.147	0.178	0.112	0.268	2.375	0.000	0.468	0.000

Problem 163	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	201	121	749	529	139	0	155	-1
N.S.	1	1.00	0.60	3.73	2.63	0.69	0.00	0.77	-0.00
time (sec)	N/A	0.138	0.089	0.111	0.280	2.361	0.000	0.451	0.000

Problem 164	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	135	436	351	106	0	113	-1
N.S.	1	1.00	0.79	2.55	2.05	0.62	0.00	0.66	-0.01
time (sec)	N/A	0.091	0.115	0.089	0.264	1.367	0.000	0.436	0.000

Problem 165	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	108	238	209	79	0	75	-1
N.S.	1	1.00	0.98	2.16	1.90	0.72	0.00	0.68	-0.01
time (sec)	N/A	0.054	0.086	0.107	0.258	2.603	0.000	0.464	0.000

Problem 166	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	28	164	62	60	36	51	97
N.S.	1	1.00	0.54	3.15	1.19	1.15	0.69	0.98	1.87
time (sec)	N/A	0.026	0.017	0.084	0.289	1.537	1.761	0.440	1.092

Problem 167	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	142	107	233	144	0	112	118
N.S.	1	1.00	1.60	1.20	2.62	1.62	0.00	1.26	1.33
time (sec)	N/A	0.065	0.052	0.099	0.273	2.503	0.000	0.492	1.149

Problem 168	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	A	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	120	152	239	224	0	145	218
N.S.	1	1.00	0.92	1.17	1.84	1.72	0.00	1.12	1.68
time (sec)	N/A	0.052	0.058	0.108	0.289	1.529	0.000	0.488	1.676

Problem 169	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	201	201	154	287	424	452	0	471	-1
N.S.	1	1.00	0.77	1.43	2.11	2.25	0.00	2.34	-0.00
time (sec)	N/A	0.114	0.096	0.105	0.278	2.089	0.000	0.529	0.000

Problem 170	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	283	283	247	529	644	690	0	884	-1
N.S.	1	1.00	0.87	1.87	2.28	2.44	0.00	3.12	-0.00
time (sec)	N/A	0.161	0.220	0.128	0.265	1.193	0.000	0.487	0.000

Problem 171	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	92	92	92	228	149	105	110	123	201
N.S.	1	1.00	1.00	2.48	1.62	1.14	1.20	1.34	2.18
time (sec)	N/A	0.065	0.052	0.158	0.470	2.573	0.253	0.427	0.585

Problem 172	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	72	170	116	77	75	83	153
N.S.	1	1.00	1.00	2.36	1.61	1.07	1.04	1.15	2.12
time (sec)	N/A	0.044	0.049	0.132	0.473	2.328	0.188	0.450	0.532

Problem 173	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	54	135	87	53	46	53	107
N.S.	1	1.00	1.00	2.50	1.61	0.98	0.85	0.98	1.98
time (sec)	N/A	0.035	0.028	0.109	0.464	2.344	0.147	0.446	0.513

Problem 174	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	37	99	64	35	29	35	60
N.S.	1	1.00	1.00	2.68	1.73	0.95	0.78	0.95	1.62
time (sec)	N/A	0.023	0.018	0.098	0.465	1.592	0.103	0.424	0.129

Problem 175	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	20	20	32	51	46	22	14	16	21
N.S.	1	1.00	1.60	2.55	2.30	1.10	0.70	0.80	1.05
time (sec)	N/A	0.009	0.011	0.097	0.480	1.774	0.066	0.425	0.464

Problem 176	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	31	110	78	27	100	33	32
N.S.	1	1.00	0.82	2.89	2.05	0.71	2.63	0.87	0.84
time (sec)	N/A	0.025	0.015	0.109	0.472	2.660	0.448	0.457	0.704

Problem 177	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	55	55	39	161	126	40	156	61	98
N.S.	1	1.00	0.71	2.93	2.29	0.73	2.84	1.11	1.78
time (sec)	N/A	0.032	0.020	0.126	0.481	2.124	0.335	0.446	0.637

Problem 178	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	63	211	188	69	228	89	154
N.S.	1	1.00	0.83	2.78	2.47	0.91	3.00	1.17	2.03
time (sec)	N/A	0.040	0.027	0.145	0.483	2.098	0.460	0.424	0.693

Problem 179	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	88	267	263	94	286	126	199
N.S.	1	1.00	0.95	2.87	2.83	1.01	3.08	1.35	2.14
time (sec)	N/A	0.045	0.038	0.158	0.486	2.255	0.621	0.421	0.718

Problem 180	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	324	324	249	4893	3081	264	0	334	-1
N.S.	1	1.00	0.77	15.10	9.51	0.81	0.00	1.03	-0.00
time (sec)	N/A	0.197	0.325	0.215	0.296	1.719	0.000	0.456	0.000

Problem 181	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	249	249	201	2972	2295	216	0	285	-1
N.S.	1	1.00	0.81	11.94	9.22	0.87	0.00	1.14	-0.00
time (sec)	N/A	0.175	0.186	0.180	0.290	2.825	0.000	0.461	0.000

Problem 182	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	227	227	160	1785	1608	174	0	243	-1
N.S.	1	1.00	0.70	7.86	7.08	0.77	0.00	1.07	-0.00
time (sec)	N/A	0.121	0.170	0.161	0.293	2.112	0.000	0.466	0.000

Problem 183	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	132	1052	1108	136	0	209	-1
N.S.	1	1.00	0.81	6.45	6.80	0.83	0.00	1.28	-0.01
time (sec)	N/A	0.085	0.118	0.148	0.277	3.037	0.000	0.454	0.000

Problem 184	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	45	617	736	99	0	180	-1
N.S.	1	1.00	0.48	6.56	7.83	1.05	0.00	1.91	-0.01
time (sec)	N/A	0.031	0.036	0.135	0.292	2.485	0.000	0.464	0.000

Problem 185	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	B	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	134	134	196	485	733	356	0	252	-1
N.S.	1	1.00	1.46	3.62	5.47	2.66	0.00	1.88	-0.01
time (sec)	N/A	0.095	0.530	0.115	0.275	1.916	0.000	0.557	0.000

Problem 186	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	176	176	145	631	992	389	0	0	-1
N.S.	1	1.00	0.82	3.59	5.64	2.21	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.104	0.203	0.273	2.385	0.000	0.000	0.000

Problem 187	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	264	264	194	946	1536	574	0	0	-1
N.S.	1	1.00	0.73	3.58	5.82	2.17	0.00	0.00	-0.00
time (sec)	N/A	0.127	0.160	0.207	0.272	6.055	0.000	0.000	0.000

Problem 188	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	338	338	282	1649	2313	839	0	0	-1
N.S.	1	1.00	0.83	4.88	6.84	2.48	0.00	0.00	-0.00
time (sec)	N/A	0.241	0.283	0.260	0.284	4.745	0.000	0.000	0.000

Problem 189	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	276	276	158	1152	456	177	0	205	-1
N.S.	1	1.00	0.57	4.17	1.65	0.64	0.00	0.74	-0.00
time (sec)	N/A	0.162	0.177	0.168	0.507	4.002	0.000	0.452	0.000

Problem 190	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	201	201	118	681	308	139	0	155	-1
N.S.	1	1.00	0.59	3.39	1.53	0.69	0.00	0.77	-0.00
time (sec)	N/A	0.139	0.110	0.120	0.492	4.483	0.000	0.437	0.000

Problem 191	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	171	171	162	485	161	106	0	113	-1
N.S.	1	1.00	0.95	2.84	0.94	0.62	0.00	0.66	-0.01
time (sec)	N/A	0.098	0.209	0.099	0.472	4.603	0.000	0.471	0.000

Problem 192	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	110	110	131	237	97	79	0	75	-1
N.S.	1	1.00	1.19	2.15	0.88	0.72	0.00	0.68	-0.01
time (sec)	N/A	0.056	0.093	0.113	0.483	4.729	0.000	0.446	0.000

Problem 193	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	52	52	28	125	35	60	0	51	-1
N.S.	1	1.00	0.54	2.40	0.67	1.15	0.00	0.98	-0.02
time (sec)	N/A	0.024	0.016	0.093	0.469	2.963	0.000	0.435	0.000

Problem 194	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-2)	B	F	A	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	142	260	0	144	0	76	-1
N.S.	1	1.00	1.60	2.92	0.00	1.62	0.00	0.85	-0.01
time (sec)	N/A	0.053	0.049	0.092	0.000	2.527	0.000	0.464	0.000

Problem 195	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	130	130	119	546	0	224	0	145	-1
N.S.	1	1.00	0.92	4.20	0.00	1.72	0.00	1.12	-0.01
time (sec)	N/A	0.054	0.053	0.102	0.000	3.837	0.000	0.476	0.000

Problem 196	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	201	201	154	1008	0	452	0	471	-1
N.S.	1	1.00	0.77	5.01	0.00	2.25	0.00	2.34	-0.00
time (sec)	N/A	0.111	0.093	0.122	0.000	4.305	0.000	0.484	0.000

Problem 197	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	283	282	247	1511	0	690	0	884	-1
N.S.	1	1.00	0.87	5.34	0.00	2.44	0.00	3.12	-0.00
time (sec)	N/A	0.179	0.233	0.135	0.000	3.912	0.000	0.494	0.000

Problem 198	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	99	99	95	125	105	105	114	215	165
N.S.	1	1.00	0.96	1.26	1.06	1.06	1.15	2.17	1.67
time (sec)	N/A	0.068	0.054	0.142	0.264	3.206	0.254	0.450	0.166

Problem 199	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	77	85	73	77	76	158	129
N.S.	1	1.00	1.00	1.10	0.95	1.00	0.99	2.05	1.68
time (sec)	N/A	0.046	0.047	0.128	0.253	2.903	0.200	0.435	0.538

Problem 200	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	59	59	55	59	53	53	49	109	90
N.S.	1	1.00	0.93	1.00	0.90	0.90	0.83	1.85	1.53
time (sec)	N/A	0.036	0.025	0.091	0.294	3.317	0.157	0.436	0.537

Problem 201	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	40	40	40	39	36	35	29	72	51
N.S.	1	1.00	1.00	0.98	0.90	0.88	0.72	1.80	1.28
time (sec)	N/A	0.031	0.017	0.084	0.260	4.044	0.122	0.419	0.503

Problem 202	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	23	23	32	22	19	22	15	37	21
N.S.	1	1.00	1.39	0.96	0.83	0.96	0.65	1.61	0.91
time (sec)	N/A	0.010	0.011	0.072	0.265	2.546	0.071	0.447	0.060

Problem 203	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	34	42	47	27	99	70	34
N.S.	1	1.00	0.83	1.02	1.15	0.66	2.41	1.71	0.83
time (sec)	N/A	0.028	0.016	0.097	0.262	3.912	0.468	0.450	0.715

Problem 204	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	62	62	42	69	110	40	158	95	100
N.S.	1	1.00	0.68	1.11	1.77	0.65	2.55	1.53	1.61
time (sec)	N/A	0.032	0.022	0.112	0.273	4.084	0.347	0.442	0.654

Problem 205	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	81	66	109	160	69	226	142	156
N.S.	1	0.98	0.80	1.31	1.93	0.83	2.72	1.71	1.88
time (sec)	N/A	0.039	0.031	0.132	0.277	2.298	0.511	0.445	0.698

Problem 206	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	104	102	91	150	218	94	286	183	199
N.S.	1	0.98	0.88	1.44	2.10	0.90	2.75	1.76	1.91
time (sec)	N/A	0.046	0.038	0.151	0.282	2.684	0.673	0.436	0.772

Problem 207	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F(-1)	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	324	324	299	1362	1368	264	0	336	-1
N.S.	1	1.00	0.92	4.20	4.22	0.81	0.00	1.04	-0.00
time (sec)	N/A	0.202	0.373	0.224	0.511	2.303	0.000	0.463	0.000

Problem 208	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F(-1)	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	249	249	244	983	979	216	0	285	-1
N.S.	1	1.00	0.98	3.95	3.93	0.87	0.00	1.14	-0.00
time (sec)	N/A	0.172	0.285	0.193	0.485	3.916	0.000	0.454	0.000

Problem 209	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	229	229	198	799	624	174	0	243	-1
N.S.	1	1.00	0.86	3.49	2.72	0.76	0.00	1.06	-0.00
time (sec)	N/A	0.117	0.237	0.168	0.481	2.994	0.000	0.464	0.000

Problem 210	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	163	163	157	595	293	136	0	210	-1
N.S.	1	1.00	0.96	3.65	1.80	0.83	0.00	1.29	-0.01
time (sec)	N/A	0.082	0.231	0.158	0.474	1.897	0.000	0.456	0.000

Problem 211	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	94	94	45	327	103	99	0	182	-1
N.S.	1	1.00	0.48	3.48	1.10	1.05	0.00	1.94	-0.01
time (sec)	N/A	0.030	0.034	0.139	0.463	2.163	0.000	0.489	0.000

Problem 212	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	B	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	134	134	189	1067	0	356	0	253	-1
N.S.	1	1.00	1.41	7.96	0.00	2.66	0.00	1.89	-0.01
time (sec)	N/A	0.083	0.456	0.123	0.000	1.904	0.000	0.526	0.000

Problem 213	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	178	178	145	1543	0	389	0	0	-1
N.S.	1	1.00	0.81	8.67	0.00	2.19	0.00	0.00	-0.01
time (sec)	N/A	0.076	0.125	0.189	0.000	4.205	0.000	0.000	0.000

Problem 214	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	264	264	194	2328	0	574	0	0	-1
N.S.	1	1.00	0.73	8.82	0.00	2.17	0.00	0.00	-0.00
time (sec)	N/A	0.124	0.177	0.220	0.000	1.959	0.000	0.000	0.000

Problem 215	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-2)	B	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	339	339	275	3637	0	839	0	0	-1
N.S.	1	1.00	0.81	10.73	0.00	2.47	0.00	0.00	-0.00
time (sec)	N/A	0.249	0.293	0.302	0.000	2.533	0.000	0.000	0.000

Problem 216	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	494	494	121	0	0	554	0	0	-1
N.S.	1	1.00	0.24	0.00	0.00	1.12	0.00	0.00	-0.00
time (sec)	N/A	0.305	0.061	0.019	0.000	4.162	0.000	0.000	0.000

Problem 217	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	410	410	81	0	0	415	0	0	-1
N.S.	1	1.00	0.20	0.00	0.00	1.01	0.00	0.00	-0.00
time (sec)	N/A	0.219	0.031	0.009	0.000	4.341	0.000	0.000	0.000

Problem 218	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	338	338	45	0	0	255	0	0	-1
N.S.	1	1.00	0.13	0.00	0.00	0.75	0.00	0.00	-0.00
time (sec)	N/A	0.144	0.014	0.005	0.000	3.150	0.000	0.000	0.000

Problem 219	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	395	395	124	0	0	414	0	0	-1
N.S.	1	1.00	0.31	0.00	0.00	1.05	0.00	0.00	-0.00
time (sec)	N/A	0.210	0.065	0.006	0.000	2.809	0.000	0.000	0.000

Problem 220	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	205	205	110	0	0	598	0	0	-1
N.S.	1	1.00	0.54	0.00	0.00	2.92	0.00	0.00	-0.00
time (sec)	N/A	0.106	0.021	0.006	0.000	3.255	0.000	0.000	0.000

Problem 221	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-2)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	494	494	121	0	0	561	0	0	-1
N.S.	1	1.00	0.24	0.00	0.00	1.14	0.00	0.00	-0.00
time (sec)	N/A	0.288	0.062	0.010	0.000	3.019	0.000	0.000	0.000

Problem 222	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	410	410	79	0	0	431	0	0	-1
N.S.	1	1.00	0.19	0.00	0.00	1.05	0.00	0.00	-0.00
time (sec)	N/A	0.206	0.031	0.012	0.000	2.947	0.000	0.000	0.000

Problem 223	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	338	338	45	0	0	268	0	0	-1
N.S.	1	1.00	0.13	0.00	0.00	0.79	0.00	0.00	-0.00
time (sec)	N/A	0.146	0.015	0.013	0.000	3.578	0.000	0.000	0.000

Problem 224	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	427	427	122	0	0	690	0	0	-1
N.S.	1	1.00	0.29	0.00	0.00	1.62	0.00	0.00	-0.00
time (sec)	N/A	0.210	0.061	0.010	0.000	2.650	0.000	0.000	0.000

Problem 225	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	211	211	106	0	0	694	0	0	-1
N.S.	1	1.00	0.50	0.00	0.00	3.29	0.00	0.00	-0.00
time (sec)	N/A	0.116	0.018	0.008	0.000	3.971	0.000	0.000	0.000

Problem 226	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	494	494	99	0	0	561	0	0	-1
N.S.	1	1.00	0.20	0.00	0.00	1.14	0.00	0.00	-0.00
time (sec)	N/A	0.289	0.064	0.015	0.000	2.553	0.000	0.000	0.000

Problem 227	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	410	410	84	0	0	421	0	0	-1
N.S.	1	1.00	0.20	0.00	0.00	1.03	0.00	0.00	-0.00
time (sec)	N/A	0.220	0.029	0.009	0.000	2.768	0.000	0.000	0.000

Problem 228	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	338	338	45	0	0	266	0	0	-1
N.S.	1	1.00	0.13	0.00	0.00	0.79	0.00	0.00	-0.00
time (sec)	N/A	0.145	0.016	0.009	0.000	3.798	0.000	0.000	0.000

Problem 229	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	395	395	126	0	0	470	0	0	-1
N.S.	1	1.00	0.32	0.00	0.00	1.19	0.00	0.00	-0.00
time (sec)	N/A	0.162	0.022	0.009	0.000	5.508	0.000	0.000	0.000

Problem 230	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	210	210	107	0	0	707	0	0	-1
N.S.	1	1.00	0.51	0.00	0.00	3.37	0.00	0.00	-0.00
time (sec)	N/A	0.095	0.020	0.008	0.000	8.002	0.000	0.000	0.000

Problem 231	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	494	494	98	0	0	555	0	0	-1
N.S.	1	1.00	0.20	0.00	0.00	1.12	0.00	0.00	-0.00
time (sec)	N/A	0.293	0.065	0.019	0.000	5.347	0.000	0.000	0.000

Problem 232	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	410	410	84	0	0	433	0	0	-1
N.S.	1	1.00	0.20	0.00	0.00	1.06	0.00	0.00	-0.00
time (sec)	N/A	0.213	0.027	0.010	0.000	6.181	0.000	0.000	0.000

Problem 233	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	A	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	338	338	43	0	0	255	0	0	-1
N.S.	1	1.00	0.13	0.00	0.00	0.75	0.00	0.00	-0.00
time (sec)	N/A	0.146	0.016	0.010	0.000	4.288	0.000	0.000	0.000

Problem 234	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	427	427	128	0	0	629	0	0	-1
N.S.	1	1.00	0.30	0.00	0.00	1.47	0.00	0.00	-0.00
time (sec)	N/A	0.204	0.023	0.011	0.000	7.016	0.000	0.000	0.000

Problem 235	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	F	F	B	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	211	211	107	0	0	613	0	0	-1
N.S.	1	1.00	0.51	0.00	0.00	2.91	0.00	0.00	-0.00
time (sec)	N/A	0.109	0.024	0.013	0.000	5.544	0.000	0.000	0.000

Problem 236	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	140	140	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.680	0.020	0.000	0.000	0.000	0.000	0.000

Problem 237	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	260	260	243	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.140	1.141	0.008	0.000	0.000	0.000	0.000	0.000

Problem 238	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	220	220	160	0	0	0	0	0	-1
N.S.	1	1.00	0.73	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.108	0.115	0.023	0.000	0.000	0.000	0.000	0.000

Problem 239	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	147	147	128	0	0	0	0	0	-1
N.S.	1	1.00	0.87	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.044	0.096	0.006	0.000	0.000	0.000	0.000	0.000

Problem 240	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	91	91	60	0	0	0	0	0	-1
N.S.	1	1.00	0.66	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.011	0.026	0.005	0.000	0.000	0.000	0.000	0.000

Problem 241	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	191	191	170	0	0	0	0	0	-1
N.S.	1	1.00	0.89	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.069	0.028	0.005	0.000	0.000	0.000	0.000	0.000

Problem 242	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	128	128	125	0	0	0	0	0	-1
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.034	0.018	0.005	0.000	0.000	0.000	0.000	0.000

Problem 243	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-1)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	207	207	173	0	0	0	0	0	-1
N.S.	1	1.00	0.84	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.088	0.053	0.005	0.000	0.000	0.000	0.000	0.000

Problem 244	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	102	102	102	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.022	0.021	0.000	0.000	0.000	0.000	0.000

Problem 245	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.013	0.013	0.000	0.000	0.000	0.000	0.000

Problem 246	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	61	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.022	0.011	0.006	0.000	0.000	0.000	0.000	0.000

Problem 247	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	45	0	0	0	0	0	-1
N.S.	1	1.00	0.75	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.016	0.006	0.000	0.000	0.000	0.000	0.000

Problem 248	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	13	13	35	13	12	12	12	12	12
N.S.	1	1.00	2.69	1.00	0.92	0.92	0.92	0.92	0.92
time (sec)	N/A	0.018	0.007	0.066	0.494	2.995	0.511	0.431	0.534

Problem 249	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	60	39	0	39	95	0	44
N.S.	1	1.00	1.20	0.78	0.00	0.78	1.90	0.00	0.88
time (sec)	N/A	0.037	0.017	0.092	0.000	3.353	1.432	0.000	0.556

Problem 250	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	83	83	114	55	0	66	223	0	74
N.S.	1	1.00	1.37	0.66	0.00	0.80	2.69	0.00	0.89
time (sec)	N/A	0.057	0.205	0.083	0.000	3.722	4.116	0.000	0.624

Problem 251	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	116	116	123	71	0	93	398	0	104
N.S.	1	1.00	1.06	0.61	0.00	0.80	3.43	0.00	0.90
time (sec)	N/A	0.083	0.203	0.073	0.000	2.606	10.797	0.000	0.687

Problem 252	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	149	149	153	87	0	120	620	0	134
N.S.	1	1.00	1.03	0.58	0.00	0.81	4.16	0.00	0.90
time (sec)	N/A	0.109	0.195	0.089	0.000	2.534	25.658	0.000	0.738

Problem 253	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	98	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.060	0.018	0.013	0.000	0.000	0.000	0.000	0.000

Problem 254	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	97	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.014	0.010	0.000	0.000	0.000	0.000	0.000

Problem 255	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	93	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.053	0.016	0.013	0.000	0.000	0.000	0.000	0.000

Problem 256	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	35	37	0	42	0	0	33
N.S.	1	1.00	1.00	1.06	0.00	1.20	0.00	0.00	0.94
time (sec)	N/A	0.026	0.015	0.077	0.000	3.108	0.000	0.000	0.602

Problem 257	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	72	72	60	54	0	70	0	0	78
N.S.	1	1.00	0.83	0.75	0.00	0.97	0.00	0.00	1.08
time (sec)	N/A	0.055	0.027	0.078	0.000	3.337	0.000	0.000	0.646

Problem 258	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	108	108	79	70	0	97	0	0	120
N.S.	1	1.00	0.73	0.65	0.00	0.90	0.00	0.00	1.11
time (sec)	N/A	0.084	0.031	0.071	0.000	3.072	0.000	0.000	0.666

Problem 259	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	90	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.055	0.021	0.020	0.000	0.000	0.000	0.000	0.000

Problem 260	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.012	0.013	0.000	0.000	0.000	0.000	0.000

Problem 261	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	51	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.019	0.011	0.006	0.000	0.000	0.000	0.000	0.000

Problem 262	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	0	0	0	0	0	-1
N.S.	1	1.00	0.80	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.020	0.004	0.000	0.000	0.000	0.000	0.000

Problem 263	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	34	16	15	15	15	15	15
N.S.	1	1.00	1.89	0.89	0.83	0.83	0.83	0.83	0.83
time (sec)	N/A	0.021	0.006	0.065	0.479	2.409	0.546	0.405	0.532

Problem 264	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	55	40	0	40	99	0	46
N.S.	1	1.00	1.04	0.75	0.00	0.75	1.87	0.00	0.87
time (sec)	N/A	0.042	0.014	0.079	0.000	2.763	1.491	0.000	0.577

Problem 265	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	86	57	0	68	231	0	79
N.S.	1	1.00	0.98	0.65	0.00	0.77	2.62	0.00	0.90
time (sec)	N/A	0.062	0.098	0.084	0.000	2.512	4.111	0.000	0.623

Problem 266	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	122	73	0	95	410	0	111
N.S.	1	1.00	0.99	0.59	0.00	0.77	3.33	0.00	0.90
time (sec)	N/A	0.087	0.186	0.076	0.000	2.736	10.726	0.000	0.666

Problem 267	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.057	0.018	0.010	0.000	0.000	0.000	0.000	0.000

Problem 268	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.017	0.013	0.000	0.000	0.000	0.000	0.000

Problem 269	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.014	0.013	0.000	0.000	0.000	0.000	0.000

Problem 270	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	37	37	37	39	0	44	0	0	35
N.S.	1	1.00	1.00	1.05	0.00	1.19	0.00	0.00	0.95
time (sec)	N/A	0.028	0.012	0.059	0.000	1.763	0.000	0.000	0.608

Problem 271	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	76	76	62	56	0	72	0	0	80
N.S.	1	1.00	0.82	0.74	0.00	0.95	0.00	0.00	1.05
time (sec)	N/A	0.057	0.025	0.100	0.000	3.269	0.000	0.000	0.219

Problem 272	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	114	114	81	72	0	99	0	0	122
N.S.	1	1.00	0.71	0.63	0.00	0.87	0.00	0.00	1.07
time (sec)	N/A	0.092	0.026	0.074	0.000	2.099	0.000	0.000	0.694

Problem 273	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	101	101	102	0	0	0	0	0	-1
N.S.	1	1.00	1.01	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.021	0.020	0.000	0.000	0.000	0.000	0.000

Problem 274	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	63	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.029	0.012	0.013	0.000	0.000	0.000	0.000	0.000

Problem 275	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	61	61	61	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.020	0.011	0.009	0.000	0.000	0.000	0.000	0.000

Problem 276	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	45	0	0	0	0	0	-1
N.S.	1	1.00	0.75	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.020	0.005	0.000	0.000	0.000	0.000	0.000

Problem 277	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	36	16	23	15	15	15	15
N.S.	1	1.00	2.25	1.00	1.44	0.94	0.94	0.94	0.94
time (sec)	N/A	0.020	0.008	0.064	0.276	1.465	4.054	0.425	0.526

Problem 278	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	60	41	0	41	116	0	47
N.S.	1	1.00	1.11	0.76	0.00	0.76	2.15	0.00	0.87
time (sec)	N/A	0.041	0.020	0.100	0.000	2.261	22.630	0.000	0.556

Problem 279	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	91	57	0	68	291	0	80
N.S.	1	1.00	1.02	0.64	0.00	0.76	3.27	0.00	0.90
time (sec)	N/A	0.062	0.116	0.085	0.000	1.478	103.566	0.000	0.617

Problem 280	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	127	73	0	95	0	0	112
N.S.	1	1.00	1.02	0.59	0.00	0.77	0.00	0.00	0.90
time (sec)	N/A	0.089	0.256	0.082	0.000	2.414	0.000	0.000	0.678

Problem 281	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-2)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	98	98	98	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.017	0.013	0.000	0.000	0.000	0.000	0.000

Problem 282	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	97	97	97	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.054	0.015	0.013	0.000	0.000	0.000	0.000	0.000

Problem 283	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	93	93	93	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.018	0.011	0.000	0.000	0.000	0.000	0.000

Problem 284	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	37	39	0	44	0	0	35
N.S.	1	1.00	0.97	1.03	0.00	1.16	0.00	0.00	0.92
time (sec)	N/A	0.028	0.017	0.090	0.000	2.657	0.000	0.000	0.626

Problem 285	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	62	56	0	72	0	0	81
N.S.	1	1.00	0.81	0.73	0.00	0.94	0.00	0.00	1.05
time (sec)	N/A	0.057	0.031	0.088	0.000	2.783	0.000	0.000	0.650

Problem 286	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	81	72	0	99	0	0	123
N.S.	1	1.00	0.70	0.63	0.00	0.86	0.00	0.00	1.07
time (sec)	N/A	0.090	0.035	0.070	0.000	2.125	0.000	0.000	0.667

Problem 287	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	90	90	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.052	0.018	0.020	0.000	0.000	0.000	0.000	0.000

Problem 288	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	53	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.028	0.012	0.013	0.000	0.000	0.000	0.000	0.000

Problem 289	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	51	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.019	0.011	0.007	0.000	0.000	0.000	0.000	0.000

Problem 290	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	46	46	37	0	0	0	0	0	-1
N.S.	1	1.00	0.80	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.009	0.020	0.003	0.000	0.000	0.000	0.000	0.000

Problem 291	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	34	18	23	15	19	15	15
N.S.	1	1.00	1.89	1.00	1.28	0.83	1.06	0.83	0.83
time (sec)	N/A	0.020	0.009	0.064	0.275	3.825	8.604	0.402	0.532

Problem 292	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	B	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	55	42	0	40	124	0	47
N.S.	1	1.00	1.02	0.78	0.00	0.74	2.30	0.00	0.87
time (sec)	N/A	0.040	0.018	0.089	0.000	2.693	44.951	0.000	0.566

Problem 293	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	85	59	0	68	0	0	79
N.S.	1	1.00	0.96	0.66	0.00	0.76	0.00	0.00	0.89
time (sec)	N/A	0.066	0.121	0.079	0.000	4.046	0.000	0.000	0.621

Problem 294	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	124	124	121	75	0	95	0	0	111
N.S.	1	1.00	0.98	0.60	0.00	0.77	0.00	0.00	0.90
time (sec)	N/A	0.094	0.247	0.079	0.000	6.903	0.000	0.000	0.672

Problem 295	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	88	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.067	0.018	0.012	0.000	0.000	0.000	0.000	0.000

Problem 296	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.016	0.012	0.000	0.000	0.000	0.000	0.000

Problem 297	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	87	87	87	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.019	0.013	0.000	0.000	0.000	0.000	0.000

Problem 298	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	37	41	0	44	0	0	35
N.S.	1	1.00	0.97	1.08	0.00	1.16	0.00	0.00	0.92
time (sec)	N/A	0.029	0.016	0.076	0.000	4.465	0.000	0.000	0.178

Problem 299	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-2)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	77	77	62	58	0	72	0	0	81
N.S.	1	1.00	0.81	0.75	0.00	0.94	0.00	0.00	1.05
time (sec)	N/A	0.058	0.032	0.073	0.000	4.919	0.000	0.000	0.633

Problem 300	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	81	74	0	99	0	0	123
N.S.	1	1.00	0.70	0.64	0.00	0.86	0.00	0.00	1.07
time (sec)	N/A	0.096	0.035	0.072	0.000	4.170	0.000	0.000	0.661

Problem 301	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	50	50	42	34	63	53	36	30	49
N.S.	1	1.00	0.84	0.68	1.26	1.06	0.72	0.60	0.98
time (sec)	N/A	0.037	0.018	0.083	0.477	3.255	0.158	0.412	0.135

Problem 302	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	B	A	F	A	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	48	224	112	86	0	24	92
N.S.	1	1.00	0.66	3.07	1.53	1.18	0.00	0.33	1.26
time (sec)	N/A	0.031	0.010	0.088	0.266	2.730	0.000	0.459	0.530

Problem 303	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	30	30	30	28	43	31	19	24	28
N.S.	1	1.00	1.00	0.93	1.43	1.03	0.63	0.80	0.93
time (sec)	N/A	0.028	0.013	0.079	0.457	3.586	0.080	0.405	0.491

Problem 304	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	52	87	40	54	0	0	55
N.S.	1	1.00	1.27	2.12	0.98	1.32	0.00	0.00	1.34
time (sec)	N/A	0.025	0.021	0.070	0.248	4.769	0.000	0.000	0.489

Problem 305	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	15	15	15	26	24	15	8	12	15
N.S.	1	1.00	1.00	1.73	1.60	1.00	0.53	0.80	1.00
time (sec)	N/A	0.021	0.006	0.061	0.459	3.079	0.014	0.449	0.474

Problem 306	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	16	16	16	15	12	15	10	12	15
N.S.	1	1.00	1.00	0.94	0.75	0.94	0.62	0.75	0.94
time (sec)	N/A	0.022	0.005	0.074	0.252	3.219	0.015	0.424	0.475

Problem 307	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	F	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	41	41	56	149	33	54	0	0	56
N.S.	1	1.00	1.37	3.63	0.80	1.32	0.00	0.00	1.37
time (sec)	N/A	0.026	0.042	0.082	0.466	3.295	0.000	0.000	0.486

Problem 308	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	32	32	32	30	41	31	19	24	29
N.S.	1	1.00	1.00	0.94	1.28	0.97	0.59	0.75	0.91
time (sec)	N/A	0.030	0.012	0.065	0.253	2.999	0.083	0.435	0.493

Problem 309	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	A	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	73	73	82	305	107	86	0	24	93
N.S.	1	1.00	1.12	4.18	1.47	1.18	0.00	0.33	1.27
time (sec)	N/A	0.029	0.055	0.087	0.472	5.120	0.000	0.463	0.102

Problem 310	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	131	69	84	0	364	0	0	-1
N.S.	1	1.00	0.53	0.64	0.00	2.78	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.028	0.095	0.000	4.245	0.000	0.000	0.000

Problem 311	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	B	F	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	71	526	0	186	0	132	-1
N.S.	1	1.00	0.74	5.48	0.00	1.94	0.00	1.38	-0.01
time (sec)	N/A	0.064	0.018	0.142	0.000	2.809	0.000	0.509	0.000

Problem 312	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	55	61	0	357	0	0	-1
N.S.	1	1.00	0.65	0.73	0.00	4.25	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.024	0.091	0.000	2.814	0.000	0.000	0.000

Problem 313	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	91	204	0	152	0	70	-1
N.S.	1	1.00	1.44	3.24	0.00	2.41	0.00	1.11	-0.02
time (sec)	N/A	0.044	0.025	0.079	0.000	2.738	0.000	0.455	0.000

Problem 314	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	42	42	42	53	0	253	0	0	-1
N.S.	1	1.00	1.00	1.26	0.00	6.02	0.00	0.00	-0.02
time (sec)	N/A	0.051	0.012	0.080	0.000	3.617	0.000	0.000	0.000

Problem 315	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	43	43	43	42	15	253	0	0	-1
N.S.	1	1.00	1.00	0.98	0.35	5.88	0.00	0.00	-0.02
time (sec)	N/A	0.048	0.012	0.088	0.276	2.803	0.000	0.000	0.000

Problem 316	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	63	63	117	87	40	152	0	70	-1
N.S.	1	1.00	1.86	1.38	0.63	2.41	0.00	1.11	-0.02
time (sec)	N/A	0.045	0.042	0.077	0.465	3.013	0.000	0.444	0.000

Problem 317	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	60	66	35	357	0	0	-1
N.S.	1	1.00	0.70	0.77	0.41	4.15	0.00	0.00	-0.01
time (sec)	N/A	0.058	0.024	0.099	0.262	4.536	0.000	0.000	0.000

Problem 318	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	B	F	A	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	96	96	132	188	76	186	0	132	-1
N.S.	1	1.00	1.38	1.96	0.79	1.94	0.00	1.38	-0.01
time (sec)	N/A	0.060	0.059	0.087	0.459	2.267	0.000	0.474	0.000

Problem 319	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	35	35	24	20	59	35	39	18	21
N.S.	1	1.00	0.69	0.57	1.69	1.00	1.11	0.51	0.60
time (sec)	N/A	0.032	0.014	0.088	0.477	5.413	0.143	0.399	0.097

Problem 320	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	47	269	95	75	0	111	41
N.S.	1	1.00	0.70	4.01	1.42	1.12	0.00	1.66	0.61
time (sec)	N/A	0.031	0.013	0.091	0.253	3.280	0.000	0.423	0.515

Problem 321	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	18	15	35	21	20	12	24
N.S.	1	1.00	0.95	0.79	1.84	1.11	1.05	0.63	1.26
time (sec)	N/A	0.027	0.016	0.079	0.470	1.660	0.089	0.401	0.500

Problem 322	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	48	104	45	51	0	67	33
N.S.	1	1.00	0.72	1.55	0.67	0.76	0.00	1.00	0.49
time (sec)	N/A	0.030	0.010	0.082	0.262	2.557	0.000	0.443	0.496

Problem 323	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	28	28	21	38	28	49	32	36	23
N.S.	1	1.00	0.75	1.36	1.00	1.75	1.14	1.29	0.82
time (sec)	N/A	0.033	0.013	0.083	0.455	2.495	0.108	0.415	0.482

Problem 324	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	29	29	21	43	0	49	34	36	25
N.S.	1	1.00	0.72	1.48	0.00	1.69	1.17	1.24	0.86
time (sec)	N/A	0.032	0.015	0.081	0.000	3.161	0.107	0.413	0.482

Problem 325	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	48	93	58	51	0	67	31
N.S.	1	1.00	0.72	1.39	0.87	0.76	0.00	1.00	0.46
time (sec)	N/A	0.030	0.011	0.065	0.474	2.658	0.000	0.431	0.060

Problem 326	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	19	19	18	16	13	21	22	13	24
N.S.	1	1.00	0.95	0.84	0.68	1.11	1.16	0.68	1.26
time (sec)	N/A	0.024	0.017	0.067	0.254	2.687	0.088	0.419	0.051

Problem 327	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	F	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	67	67	47	92	99	75	0	111	40
N.S.	1	1.00	0.70	1.37	1.48	1.12	0.00	1.66	0.60
time (sec)	N/A	0.029	0.012	0.083	0.259	7.369	0.000	0.445	0.516

Problem 328	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	95	95	56	48	0	101	0	0	48
N.S.	1	1.00	0.59	0.51	0.00	1.06	0.00	0.00	0.51
time (sec)	N/A	0.062	0.022	0.087	0.000	2.979	0.000	0.000	1.620

Problem 329	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	77	940	0	66	0	134	46
N.S.	1	1.00	1.12	13.62	0.00	0.96	0.00	1.94	0.67
time (sec)	N/A	0.049	0.026	0.090	0.000	3.429	0.000	0.444	0.979

Problem 330	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	57	42	0	71	0	0	41
N.S.	1	1.00	1.16	0.86	0.00	1.45	0.00	0.00	0.84
time (sec)	N/A	0.054	0.029	0.079	0.000	7.347	0.000	0.000	1.245

Problem 331	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F	A	F	A	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	78	398	0	47	0	76	32
N.S.	1	1.00	1.44	7.37	0.00	0.87	0.00	1.41	0.59
time (sec)	N/A	0.040	0.021	0.105	0.000	3.634	0.000	0.428	0.652

Problem 332	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	88	88	51	58	0	317	0	0	-1
N.S.	1	1.00	0.58	0.66	0.00	3.60	0.00	0.00	-0.01
time (sec)	N/A	0.059	0.025	0.082	0.000	5.398	0.000	0.000	0.000

Problem 333	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	89	89	60	86	52	317	0	0	-1
N.S.	1	1.00	0.67	0.97	0.58	3.56	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.028	0.082	0.269	3.309	0.000	0.000	0.000

Problem 334	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	A	A	F	A	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	54	54	78	137	59	47	0	76	32
N.S.	1	1.00	1.44	2.54	1.09	0.87	0.00	1.41	0.59
time (sec)	N/A	0.041	0.025	0.074	0.272	1.789	0.000	0.481	0.652

Problem 335	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	57	43	29	71	0	0	49
N.S.	1	1.00	1.16	0.88	0.59	1.45	0.00	0.00	1.00
time (sec)	N/A	0.052	0.033	0.076	0.269	2.541	0.000	0.000	1.065

Problem 336	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	B	A	F	B	B
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	69	69	77	307	119	66	0	134	45
N.S.	1	1.00	1.12	4.45	1.72	0.96	0.00	1.94	0.65
time (sec)	N/A	0.050	0.032	0.072	0.271	2.909	0.000	0.470	0.996

Problem 337	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	86	86	90	0	0	0	0	0	-1
N.S.	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.043	0.017	0.014	0.000	0.000	0.000	0.000	0.000

Problem 338	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	84	84	88	0	0	0	0	0	-1
N.S.	1	1.00	1.05	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.031	0.014	0.006	0.000	0.000	0.000	0.000	0.000

Problem 339	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	81	81	56	0	0	0	0	0	-1
N.S.	1	1.00	0.69	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.014	0.019	0.004	0.000	0.000	0.000	0.000	0.000

Problem 340	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	131	206	141	0	0	0	0	0	-1
N.S.	1	1.57	1.08	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.170	0.089	0.014	0.000	0.000	0.000	0.000	0.000

Problem 341	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	164	164	77	0	0	0	0	0	-1
N.S.	1	1.00	0.47	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.102	0.098	0.013	0.000	0.000	0.000	0.000	0.000

Problem 342	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	122	122	104	0	0	0	0	0	-1
N.S.	1	1.00	0.85	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.056	0.100	0.010	0.000	0.000	0.000	0.000	0.000

Problem 343	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	A	A	B	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	18	18	42	18	17	17	26	17	17
N.S.	1	1.00	2.33	1.00	0.94	0.94	1.44	0.94	0.94
time (sec)	N/A	0.021	0.007	0.066	0.490	1.834	0.598	0.408	0.634

Problem 344	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	122	120	0	0	0	0	0	-1
N.S.	1	1.88	1.85	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.069	0.032	0.011	0.000	0.000	0.000	0.000	0.000

Problem 345	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	90	166	142	0	0	0	0	0	-1
N.S.	1	1.84	1.58	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.093	0.040	0.011	0.000	0.000	0.000	0.000	0.000

Problem 346	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	126	233	174	0	0	0	0	0	-1
N.S.	1	1.85	1.38	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.129	0.060	0.010	0.000	0.000	0.000	0.000	0.000

Problem 347	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	C	A	F	A	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	181	181	165	166	0	298	0	0	281
N.S.	1	1.00	0.91	0.92	0.00	1.65	0.00	0.00	1.55
time (sec)	N/A	0.133	0.356	0.089	0.000	2.794	0.000	0.000	0.869

Problem 348	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	118	0	0	0	0	0	-1
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.079	0.084	0.013	0.000	0.000	0.000	0.000	0.000

Problem 349	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	117	0	0	0	0	0	-1
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.040	0.010	0.000	0.000	0.000	0.000	0.000

Problem 350	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	117	0	0	0	0	0	-1
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.070	0.027	0.013	0.000	0.000	0.000	0.000	0.000

Problem 351	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F(-1)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	283	283	217	0	0	0	0	0	-1
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.220	0.187	0.013	0.000	0.000	0.000	0.000	0.000

Problem 352	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	280	280	214	0	0	0	0	0	-1
N.S.	1	1.00	0.76	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.200	0.150	0.013	0.000	0.000	0.000	0.000	0.000

Problem 353	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	322	322	248	0	0	0	0	0	-1
N.S.	1	1.00	0.77	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.255	0.222	0.013	0.000	0.000	0.000	0.000	0.000

Problem 354	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	291	291	206	0	0	0	0	0	-1
N.S.	1	1.00	0.71	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.245	0.136	0.014	0.000	0.000	0.000	0.000	0.000

Problem 355	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	202	202	187	0	0	0	0	0	-1
N.S.	1	1.00	0.93	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.133	0.104	0.013	0.000	0.000	0.000	0.000	0.000

Problem 356	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	117	0	0	0	0	0	-1
N.S.	1	1.00	0.98	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.062	0.021	0.000	0.000	0.000	0.000	0.000	0.000

Problem 357	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	121	121	120	0	0	0	0	0	-1
N.S.	1	1.00	0.99	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.139	0.032	0.011	0.000	0.000	0.000	0.000	0.000

Problem 358	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	196	196	142	0	0	0	0	0	-1
N.S.	1	1.00	0.72	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.162	0.046	0.013	0.000	0.000	0.000	0.000	0.000

Problem 359	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	281	281	159	0	0	0	0	0	-1
N.S.	1	1.00	0.57	0.00	0.00	0.00	0.00	0.00	-0.00
time (sec)	N/A	0.180	0.057	0.013	0.000	0.000	0.000	0.000	0.000

Problem 360	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F(-2)	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	120	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.079	0.049	0.013	0.000	0.000	0.000	0.000	0.000

Problem 361	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	120	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.078	0.034	0.011	0.000	0.000	0.000	0.000	0.000

Problem 362	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	120	120	120	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.080	0.033	0.013	0.000	0.000	0.000	0.000	0.000

Problem 363	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	123	123	123	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.087	0.038	0.011	0.000	0.000	0.000	0.000	0.000

Problem 364	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	49	49	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.051	0.351	0.020	0.000	0.000	0.000	0.000	0.000

Problem 365	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	NO	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	96	0	0	0	0	0	-1
N.S.	1	1.00	1.88	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.065	0.116	0.011	0.000	0.000	0.000	0.000	0.000

Problem 366	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.064	0.348	0.031	0.000	0.000	0.000	0.000	0.000

Problem 367	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	51	51	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.02
time (sec)	N/A	0.065	0.552	0.038	0.000	0.000	0.000	0.000	0.000

Problem 368	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	79	79	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.141	0.249	0.026	0.000	0.000	0.000	0.000	0.000

Problem 369	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.154	0.337	0.024	0.000	0.000	0.000	0.000	0.000

Problem 370	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	F	F	F	F	F(-2)	F	F
verified	N/A	Yes	N/A	TBD	TBD	TBD	TBD	TBD	TBD
size	82	82	0	0	0	0	0	0	-1
N.S.	1	1.00	0.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.152	0.442	0.026	0.000	0.000	0.000	0.000	0.000

Problem 371	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	F	F	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	115	115	115	0	0	0	0	0	-1
N.S.	1	1.00	1.00	0.00	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.064	0.026	0.019	0.000	0.000	0.000	0.000	0.000

Problem 372	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	39	41	76	42	0	0	54
N.S.	1	1.00	0.74	0.77	1.43	0.79	0.00	0.00	1.02
time (sec)	N/A	0.048	0.022	0.142	0.277	3.337	0.000	0.000	0.658

Problem 373	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	53	53	39	41	0	44	0	0	54
N.S.	1	1.00	0.74	0.77	0.00	0.83	0.00	0.00	1.02
time (sec)	N/A	0.044	0.019	0.099	0.000	14.583	0.000	0.000	0.591

Problem 374	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F	A	F(-2)	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	60	60	55	62	0	78	0	0	-1
N.S.	1	1.00	0.92	1.03	0.00	1.30	0.00	0.00	-0.02
time (sec)	N/A	0.080	0.018	0.107	0.000	7.026	0.000	0.000	0.000

Problem 375	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	B	B	F(-1)
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	34	292	379	439	299	-1
N.S.	1	1.00	0.95	0.89	7.68	9.97	11.55	7.87	-0.03
time (sec)	N/A	0.054	0.779	0.548	0.619	9.674	3.693	0.430	0.000

Problem 376	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	35	155	169	194	139	160
N.S.	1	1.00	0.95	0.92	4.08	4.45	5.11	3.66	4.21
time (sec)	N/A	0.057	0.156	0.302	0.494	3.905	1.023	0.412	3.581

Problem 377	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	A	A	A	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	34	62	49	54	45	47
N.S.	1	1.00	0.95	0.89	1.63	1.29	1.42	1.18	1.24
time (sec)	N/A	0.054	0.027	0.110	0.460	3.103	0.250	0.419	0.640

Problem 378	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	A	A	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	62	0	49	53	80	65
N.S.	1	1.00	0.95	1.63	0.00	1.29	1.39	2.11	1.71
time (sec)	N/A	0.055	0.026	0.112	0.000	2.640	0.227	0.429	0.588

Problem 379	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	B	F(-2)	B	B	B	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	38	38	36	218	0	169	192	139	159
N.S.	1	1.00	0.95	5.74	0.00	4.45	5.05	3.66	4.18
time (sec)	N/A	0.056	0.153	0.316	0.000	2.668	0.851	0.399	3.400

Problem 380	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	63	57	0	496	0	0	46
N.S.	1	1.00	0.97	0.88	0.00	7.63	0.00	0.00	0.71
time (sec)	N/A	0.144	0.362	0.257	0.000	4.952	0.000	0.000	2.210

Problem 381	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	B	F	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	65	57	0	192	0	0	48
N.S.	1	1.00	1.00	0.88	0.00	2.95	0.00	0.00	0.74
time (sec)	N/A	0.148	0.075	0.128	0.000	3.067	0.000	0.000	1.598

Problem 382	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	F(-2)	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	142	142	74	87	0	0	0	0	-1
N.S.	1	1.00	0.52	0.61	0.00	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.164	0.040	0.104	0.000	0.000	0.000	0.000	0.000

Problem 383	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	F	F	F	F
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	143	143	75	86	54	0	0	0	-1
N.S.	1	1.00	0.52	0.60	0.38	0.00	0.00	0.00	-0.01
time (sec)	N/A	0.159	0.045	0.108	0.274	0.000	0.000	0.000	0.000

Problem 384	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	A	B	F(-1)	F	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	65	57	93	192	0	0	57
N.S.	1	1.00	1.00	0.88	1.43	2.95	0.00	0.00	0.88
time (sec)	N/A	0.146	0.080	0.132	0.273	3.609	0.000	0.000	1.589

Problem 385	Optimal	Rubi	MMA	Maple	Maxima	Fricas	Sympy	Giac	Mupad
grade	A	A	A	A	B	B	F(-1)	F(-2)	B
verified	N/A	Yes	Yes	TBD	TBD	TBD	TBD	TBD	TBD
size	65	65	63	57	274	496	0	0	47
N.S.	1	1.00	0.97	0.88	4.22	7.63	0.00	0.00	0.72
time (sec)	N/A	0.152	0.392	0.203	0.341	3.072	0.000	0.000	3.089

2.3 Detailed conclusion table specific for Rubi results

The following table is specific to Rubi. It gives additional statistics for each integral. the column **steps** is the number of steps used by Rubi to obtain the antiderivative. The **rules** column is the number of unique rules used. The **integrand size** column is the leaf size of the integrand. Finally the ratio $\frac{\text{number of rules}}{\text{integrand size}}$ is given. The larger this ratio is, the harder the integral was to solve. In this test, problem number [374] had the largest ratio of [35]

Table 2.1: Rubi specific breakdown of results for each integral

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
1	A	6	4	1.00	14	0.286
2	A	5	4	1.00	14	0.286
3	A	7	5	1.00	14	0.357
4	A	3	3	1.00	12	0.250
5	A	3	3	1.00	10	0.300
6	A	6	6	1.00	14	0.429
7	A	5	5	1.00	14	0.357
8	A	6	6	1.00	14	0.429
9	A	7	6	1.00	14	0.429
10	A	8	6	1.00	14	0.429
11	A	3	2	1.00	14	0.143
12	A	3	2	1.00	14	0.143
13	A	3	2	1.00	12	0.167
14	A	3	2	1.00	10	0.200
15	A	3	2	1.00	14	0.143
16	A	3	2	1.00	14	0.143
17	A	3	2	1.00	14	0.143
18	A	3	2	1.00	14	0.143
19	A	14	11	1.00	14	0.786
20	A	10	9	1.00	14	0.643
21	A	9	7	1.00	12	0.583
22	A	5	5	1.00	10	0.500
23	A	8	7	1.00	14	0.500
24	A	8	7	1.00	14	0.500
25	A	12	8	1.00	14	0.571

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
26	A	14	9	1.00	14	0.643
27	A	3	2	1.00	14	0.143
28	A	3	2	1.00	14	0.143
29	A	3	2	1.00	12	0.167
30	A	3	2	1.00	10	0.200
31	A	3	2	1.00	14	0.143
32	A	3	2	1.00	14	0.143
33	A	3	2	1.00	14	0.143
34	A	3	2	1.00	14	0.143
35	A	5	4	1.00	14	0.286
36	A	7	5	1.00	14	0.357
37	A	3	3	1.00	12	0.250
38	A	3	3	1.00	10	0.300
39	A	6	6	1.00	14	0.429
40	A	5	5	1.00	14	0.357
41	A	6	6	1.00	14	0.429
42	A	7	6	1.00	14	0.429
43	A	8	6	1.00	14	0.429
44	A	3	2	1.00	14	0.143
45	A	3	2	1.00	14	0.143
46	A	3	2	1.00	12	0.167
47	A	3	2	1.00	10	0.200
48	A	3	2	1.00	14	0.143
49	A	3	2	1.00	14	0.143
50	A	3	2	1.00	14	0.143
51	A	3	2	1.00	14	0.143
52	A	14	11	1.00	14	0.786
53	A	10	9	1.00	14	0.643
54	A	9	7	1.00	12	0.583
55	A	5	5	1.00	10	0.500
56	A	8	7	1.00	14	0.500
57	A	8	7	1.00	14	0.500
58	A	12	8	1.00	14	0.571
59	A	14	9	1.00	14	0.643
60	A	19	9	1.00	14	0.643

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
61	A	15	12	1.00	16	0.750
62	A	14	11	1.00	14	0.786
63	A	13	10	1.00	12	0.833
64	A	17	14	1.00	16	0.875
65	A	6	6	1.00	16	0.375
66	A	7	7	1.00	16	0.438
67	A	9	8	1.00	16	0.500
68	A	10	8	1.00	16	0.500
69	A	11	8	1.00	16	0.500
70	A	15	12	1.00	16	0.750
71	A	15	12	1.00	16	0.750
72	A	14	11	1.00	14	0.786
73	A	13	10	1.00	12	0.833
74	A	17	14	1.00	16	0.875
75	A	6	6	1.00	16	0.375
76	A	7	7	1.00	16	0.438
77	A	9	8	1.00	16	0.500
78	A	10	8	1.00	16	0.500
79	A	16	13	1.00	16	0.812
80	A	16	12	1.00	16	0.750
81	A	15	11	1.00	14	0.786
82	A	14	11	1.00	12	0.917
83	A	19	16	1.00	16	1.000
84	A	7	6	1.00	16	0.375
85	A	8	7	1.00	16	0.438
86	A	10	9	1.00	16	0.562
87	A	11	9	1.00	16	0.562
88	A	15	12	1.00	16	0.750
89	A	15	12	1.00	16	0.750
90	A	14	11	1.00	14	0.786
91	A	13	10	1.00	12	0.833
92	A	17	14	1.00	16	0.875
93	A	6	6	1.00	16	0.375
94	A	7	7	1.00	16	0.438
95	A	9	8	1.00	16	0.500

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
96	A	10	8	1.00	16	0.500
97	A	15	12	1.00	16	0.750
98	A	15	12	1.00	16	0.750
99	A	14	11	1.00	14	0.786
100	A	13	10	1.00	12	0.833
101	A	17	14	1.00	16	0.875
102	A	6	6	1.00	16	0.375
103	A	7	7	1.00	16	0.438
104	A	9	8	1.00	16	0.500
105	A	10	8	1.00	16	0.500
106	A	16	13	1.00	16	0.812
107	A	16	12	1.00	16	0.750
108	A	15	11	1.00	14	0.786
109	A	14	11	1.00	12	0.917
110	A	19	16	1.00	16	1.000
111	A	7	6	1.00	16	0.375
112	A	8	7	1.00	16	0.438
113	A	10	9	1.00	16	0.562
114	A	11	9	1.00	16	0.562
115	A	16	12	1.00	14	0.857
116	A	15	11	1.00	12	0.917
117	A	14	10	1.00	10	1.000
118	A	25	13	1.00	14	0.929
119	A	13	9	1.00	14	0.643
120	A	14	10	1.00	14	0.714
121	A	16	11	1.00	14	0.786
122	A	5	5	1.00	14	0.357
123	A	4	4	1.00	12	0.333
124	A	3	3	1.00	10	0.300
125	A	4	4	1.00	14	0.286
126	A	3	3	1.00	14	0.214
127	A	4	4	1.00	14	0.286
128	A	27	13	1.00	16	0.812
129	A	26	12	1.00	14	0.857
130	A	25	11	1.00	12	0.917

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
131	A	39	20	1.00	16	1.250
132	A	16	13	1.00	16	0.812
133	A	17	14	1.00	16	0.875
134	A	4	4	1.00	14	0.286
135	A	4	4	1.00	14	0.286
136	A	3	3	1.00	14	0.214
137	A	3	3	1.00	14	0.214
138	A	4	4	1.00	14	0.286
139	A	4	4	1.00	14	0.286
140	A	9	5	1.00	14	0.357
141	A	4	3	1.00	14	0.214
142	A	4	3	1.00	14	0.214
143	A	9	5	1.00	14	0.357
144	A	2	2	1.00	16	0.125
145	A	2	2	1.00	16	0.125
146	A	2	2	1.00	16	0.125
147	A	2	2	1.00	16	0.125
148	A	2	2	1.00	16	0.125
149	A	2	2	1.00	16	0.125
150	A	2	2	1.00	12	0.167
151	A	2	2	1.00	12	0.167
152	A	2	2	1.00	16	0.125
153	A	2	2	1.00	15	0.133
154	A	4	4	1.00	15	0.267
155	A	4	4	1.00	15	0.267
156	A	3	3	1.00	13	0.231
157	A	2	2	1.00	11	0.182
158	A	4	4	1.00	15	0.267
159	A	2	2	1.00	15	0.133
160	A	3	3	1.00	15	0.200
161	A	5	5	1.00	15	0.333
162	A	8	8	1.00	16	0.500
163	A	7	7	1.00	16	0.438
164	A	7	7	1.00	16	0.438
165	A	6	6	1.00	14	0.429

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
166	A	5	5	1.00	12	0.417
167	A	8	8	1.00	16	0.500
168	A	4	4	1.00	16	0.250
169	A	5	5	1.00	16	0.312
170	A	7	6	1.00	16	0.375
171	A	3	2	1.00	16	0.125
172	A	3	2	1.00	16	0.125
173	A	3	2	1.00	16	0.125
174	A	3	2	1.00	14	0.143
175	A	3	2	1.00	12	0.167
176	A	3	2	1.00	16	0.125
177	A	3	2	1.00	16	0.125
178	A	3	2	1.00	16	0.125
179	A	3	2	1.00	16	0.125
180	A	9	8	1.00	16	0.500
181	A	8	8	1.00	16	0.500
182	A	8	7	1.00	16	0.438
183	A	7	6	1.00	14	0.429
184	A	6	6	1.00	12	0.500
185	A	8	8	1.00	16	0.500
186	A	5	4	1.00	16	0.250
187	A	6	5	1.00	16	0.312
188	A	8	7	1.00	16	0.438
189	A	8	8	1.00	16	0.500
190	A	7	7	1.00	16	0.438
191	A	7	7	1.00	16	0.438
192	A	6	6	1.00	14	0.429
193	A	5	5	1.00	12	0.417
194	A	8	8	1.00	16	0.500
195	A	4	4	1.00	16	0.250
196	A	5	5	1.00	16	0.312
197	A	7	6	1.00	16	0.375
198	A	3	2	1.00	16	0.125
199	A	3	2	1.00	16	0.125
200	A	3	2	1.00	16	0.125

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
201	A	3	2	1.00	14	0.143
202	A	3	2	1.00	12	0.167
203	A	3	2	1.00	16	0.125
204	A	3	2	1.00	16	0.125
205	A	3	2	0.98	16	0.125
206	A	3	2	0.98	16	0.125
207	A	9	8	1.00	16	0.500
208	A	8	8	1.00	16	0.500
209	A	8	7	1.00	16	0.438
210	A	7	6	1.00	14	0.429
211	A	6	6	1.00	12	0.500
212	A	8	8	1.00	16	0.500
213	A	5	4	1.00	16	0.250
214	A	6	5	1.00	16	0.312
215	A	8	7	1.00	16	0.438
216	A	15	12	1.00	18	0.667
217	A	14	11	1.00	16	0.688
218	A	13	10	1.00	14	0.714
219	A	15	12	1.00	18	0.667
220	A	6	6	1.00	18	0.333
221	A	15	12	1.00	18	0.667
222	A	14	11	1.00	16	0.688
223	A	13	10	1.00	14	0.714
224	A	18	15	1.00	18	0.833
225	A	6	6	1.00	18	0.333
226	A	15	12	1.00	18	0.667
227	A	14	11	1.00	16	0.688
228	A	13	10	1.00	14	0.714
229	A	14	11	1.00	18	0.611
230	A	5	5	1.00	18	0.278
231	A	15	12	1.00	18	0.667
232	A	14	11	1.00	16	0.688
233	A	13	10	1.00	14	0.714
234	A	18	15	1.00	18	0.833
235	A	6	6	1.00	18	0.333

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
236	A	4	3	1.00	14	0.214
237	A	4	4	1.00	14	0.286
238	A	4	4	1.00	14	0.286
239	A	3	3	1.00	12	0.250
240	A	2	2	1.00	10	0.200
241	A	5	5	1.00	14	0.357
242	A	2	2	1.00	14	0.143
243	A	3	3	1.00	14	0.214
244	A	3	3	1.00	19	0.158
245	A	2	2	1.00	19	0.105
246	A	2	2	1.00	17	0.118
247	A	2	2	1.00	6	0.333
248	A	1	1	1.00	19	0.053
249	A	2	2	1.00	19	0.105
250	A	3	2	1.00	19	0.105
251	A	4	2	1.00	19	0.105
252	A	5	2	1.00	19	0.105
253	A	3	3	1.00	21	0.143
254	A	3	3	1.00	21	0.143
255	A	3	3	1.00	21	0.143
256	A	1	1	1.00	21	0.048
257	A	2	2	1.00	21	0.095
258	A	3	2	1.00	21	0.095
259	A	3	3	1.00	21	0.143
260	A	2	2	1.00	21	0.095
261	A	2	2	1.00	19	0.105
262	A	2	2	1.00	8	0.250
263	A	1	1	1.00	21	0.048
264	A	2	2	1.00	21	0.095
265	A	3	2	1.00	21	0.095
266	A	4	2	1.00	21	0.095
267	A	3	3	1.00	23	0.130
268	A	3	3	1.00	23	0.130
269	A	3	3	1.00	23	0.130
270	A	1	1	1.00	23	0.043

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
271	A	2	2	1.00	23	0.087
272	A	3	2	1.00	23	0.087
273	A	3	3	1.00	21	0.143
274	A	2	2	1.00	21	0.095
275	A	2	2	1.00	19	0.105
276	A	2	2	1.00	8	0.250
277	A	1	1	1.00	21	0.048
278	A	2	2	1.00	21	0.095
279	A	3	2	1.00	21	0.095
280	A	4	2	1.00	21	0.095
281	A	3	3	1.00	23	0.130
282	A	3	3	1.00	23	0.130
283	A	3	3	1.00	23	0.130
284	A	1	1	1.00	23	0.043
285	A	2	2	1.00	23	0.087
286	A	3	2	1.00	23	0.087
287	A	3	3	1.00	21	0.143
288	A	2	2	1.00	21	0.095
289	A	2	2	1.00	19	0.105
290	A	2	2	1.00	8	0.250
291	A	1	1	1.00	21	0.048
292	A	2	2	1.00	21	0.095
293	A	3	2	1.00	21	0.095
294	A	4	2	1.00	21	0.095
295	A	3	3	1.00	23	0.130
296	A	3	3	1.00	23	0.130
297	A	3	3	1.00	23	0.130
298	A	1	1	1.00	23	0.043
299	A	2	2	1.00	23	0.087
300	A	3	2	1.00	23	0.087
301	A	3	2	1.00	24	0.083
302	A	5	4	1.00	24	0.167
303	A	3	2	1.00	24	0.083
304	A	4	4	1.00	24	0.167
305	A	2	2	1.00	24	0.083

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
306	A	2	2	1.00	24	0.083
307	A	4	4	1.00	24	0.167
308	A	3	2	1.00	24	0.083
309	A	5	4	1.00	24	0.167
310	A	4	3	1.00	25	0.120
311	A	5	5	1.00	25	0.200
312	A	4	3	1.00	25	0.120
313	A	4	4	1.00	25	0.160
314	A	3	3	1.00	25	0.120
315	A	3	3	1.00	25	0.120
316	A	4	4	1.00	25	0.160
317	A	4	3	1.00	25	0.120
318	A	5	5	1.00	25	0.200
319	A	3	2	1.00	24	0.083
320	A	3	3	1.00	24	0.125
321	A	2	2	1.00	24	0.083
322	A	3	3	1.00	24	0.125
323	A	4	3	1.00	24	0.125
324	A	4	3	1.00	24	0.125
325	A	3	3	1.00	24	0.125
326	A	2	2	1.00	24	0.083
327	A	3	3	1.00	24	0.125
328	A	4	3	1.00	25	0.120
329	A	3	3	1.00	25	0.120
330	A	3	3	1.00	25	0.120
331	A	3	3	1.00	25	0.120
332	A	5	4	1.00	25	0.160
333	A	5	4	1.00	25	0.160
334	A	3	3	1.00	25	0.120
335	A	3	3	1.00	25	0.120
336	A	3	3	1.00	25	0.120
337	A	2	2	1.00	21	0.095
338	A	2	2	1.00	19	0.105
339	A	2	2	1.00	8	0.250
340	A	4	4	1.57	24	0.167

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
341	A	4	4	1.00	24	0.167
342	A	3	3	1.00	22	0.136
343	A	1	1	1.00	21	0.048
344	A	3	3	1.88	24	0.125
345	A	5	5	1.84	24	0.208
346	A	6	6	1.85	24	0.250
347	A	4	2	1.00	21	0.095
348	A	3	3	1.00	23	0.130
349	A	3	3	1.00	23	0.130
350	A	3	3	1.00	23	0.130
351	A	5	5	1.00	26	0.192
352	A	5	5	1.00	26	0.192
353	A	5	5	1.00	26	0.192
354	A	5	5	1.00	26	0.192
355	A	4	4	1.00	24	0.167
356	A	3	3	1.00	23	0.130
357	A	3	3	1.00	26	0.115
358	A	4	4	1.00	26	0.154
359	A	6	6	1.00	26	0.231
360	A	3	3	1.00	23	0.130
361	A	3	3	1.00	23	0.130
362	A	3	3	1.00	23	0.130
363	A	3	3	1.00	23	0.130
364	A	2	2	1.00	22	0.091
365	A	2	2	1.00	24	0.083
366	A	2	2	1.00	24	0.083
367	A	2	2	1.00	24	0.083
368	A	3	3	1.00	26	0.115
369	A	3	3	1.00	26	0.115
370	A	3	3	1.00	26	0.115
371	A	3	3	1.00	21	0.143
372	A	3	3	1.00	24	0.125
373	A	3	3	1.00	24	0.125
374	A	1	1	1.00	35	0.029
375	A	2	2	1.00	26	0.077

Continued on next page

Table 2.1 – continued from previous page

#	grade	number of steps used	number of unique rules	normalized antiderivative leaf size	integrand leaf size	$\frac{\text{number of rules}}{\text{integrand leaf size}}$
376	A	2	2	1.00	26	0.077
377	A	2	2	1.00	26	0.077
378	A	2	2	1.00	26	0.077
379	A	2	2	1.00	26	0.077
380	A	3	3	1.00	28	0.107
381	A	3	3	1.00	28	0.107
382	A	4	3	1.00	28	0.107
383	A	4	3	1.00	28	0.107
384	A	3	3	1.00	28	0.107
385	A	3	3	1.00	28	0.107

Chapter 3

Listing of integrals

Local contents

3.1	$\int e^{i\text{ArcTan}(ax)} x^4 dx$	118
3.2	$\int e^{i\text{ArcTan}(ax)} x^3 dx$	122
3.3	$\int e^{i\text{ArcTan}(ax)} x^2 dx$	126
3.4	$\int e^{i\text{ArcTan}(ax)} x dx$	130
3.5	$\int e^{i\text{ArcTan}(ax)} dx$	134
3.6	$\int \frac{e^{i\text{ArcTan}(ax)}}{x} dx$	137
3.7	$\int \frac{e^{i\text{ArcTan}(ax)}}{x^2} dx$	141
3.8	$\int \frac{e^{i\text{ArcTan}(ax)}}{x^3} dx$	145
3.9	$\int \frac{e^{i\text{ArcTan}(ax)}}{x^4} dx$	149
3.10	$\int \frac{e^{i\text{ArcTan}(ax)}}{x^5} dx$	154
3.11	$\int e^{2i\text{ArcTan}(ax)} x^3 dx$	159
3.12	$\int e^{2i\text{ArcTan}(ax)} x^2 dx$	162
3.13	$\int e^{2i\text{ArcTan}(ax)} x dx$	165
3.14	$\int e^{2i\text{ArcTan}(ax)} dx$	168
3.15	$\int \frac{e^{2i\text{ArcTan}(ax)}}{x} dx$	171
3.16	$\int \frac{e^{2i\text{ArcTan}(ax)}}{x^2} dx$	174
3.17	$\int \frac{e^{2i\text{ArcTan}(ax)}}{x^3} dx$	177
3.18	$\int \frac{e^{2i\text{ArcTan}(ax)}}{x^4} dx$	180
3.19	$\int e^{3i\text{ArcTan}(ax)} x^3 dx$	183
3.20	$\int e^{3i\text{ArcTan}(ax)} x^2 dx$	190
3.21	$\int e^{3i\text{ArcTan}(ax)} x dx$	195
3.22	$\int e^{3i\text{ArcTan}(ax)} dx$	200
3.23	$\int \frac{e^{3i\text{ArcTan}(ax)}}{x} dx$	204
3.24	$\int \frac{e^{3i\text{ArcTan}(ax)}}{x^2} dx$	208
3.25	$\int \frac{e^{3i\text{ArcTan}(ax)}}{x^3} dx$	212

3.26	$\int \frac{e^{3i\text{ArcTan}(ax)}}{x^4} dx$	217
3.27	$\int e^{4i\text{ArcTan}(ax)} x^3 dx$	222
3.28	$\int e^{4i\text{ArcTan}(ax)} x^2 dx$	225
3.29	$\int e^{4i\text{ArcTan}(ax)} x dx$	228
3.30	$\int e^{4i\text{ArcTan}(ax)} dx$	231
3.31	$\int \frac{e^{4i\text{ArcTan}(ax)}}{x} dx$	234
3.32	$\int \frac{e^{4i\text{ArcTan}(ax)}}{x^2} dx$	237
3.33	$\int \frac{e^{4i\text{ArcTan}(ax)}}{x^3} dx$	240
3.34	$\int \frac{e^{4i\text{ArcTan}(ax)}}{x^4} dx$	243
3.35	$\int e^{-i\text{ArcTan}(ax)} x^3 dx$	246
3.36	$\int e^{-i\text{ArcTan}(ax)} x^2 dx$	250
3.37	$\int e^{-i\text{ArcTan}(ax)} x dx$	254
3.38	$\int e^{-i\text{ArcTan}(ax)} dx$	258
3.39	$\int \frac{e^{-i\text{ArcTan}(ax)}}{x} dx$	261
3.40	$\int \frac{e^{-i\text{ArcTan}(ax)}}{x^2} dx$	265
3.41	$\int \frac{e^{-i\text{ArcTan}(ax)}}{x^3} dx$	269
3.42	$\int \frac{e^{-i\text{ArcTan}(ax)}}{x^4} dx$	273
3.43	$\int \frac{e^{-i\text{ArcTan}(ax)}}{x^5} dx$	277
3.44	$\int e^{-2i\text{ArcTan}(ax)} x^3 dx$	281
3.45	$\int e^{-2i\text{ArcTan}(ax)} x^2 dx$	284
3.46	$\int e^{-2i\text{ArcTan}(ax)} x dx$	287
3.47	$\int e^{-2i\text{ArcTan}(ax)} dx$	290
3.48	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{x} dx$	293
3.49	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{x^2} dx$	296
3.50	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{x^3} dx$	299
3.51	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{x^4} dx$	302
3.52	$\int e^{-3i\text{ArcTan}(ax)} x^3 dx$	305
3.53	$\int e^{-3i\text{ArcTan}(ax)} x^2 dx$	312
3.54	$\int e^{-3i\text{ArcTan}(ax)} x dx$	317
3.55	$\int e^{-3i\text{ArcTan}(ax)} dx$	322
3.56	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{x} dx$	326
3.57	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{x^2} dx$	330
3.58	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{x^3} dx$	334
3.59	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{x^4} dx$	339
3.60	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{x^5} dx$	344
3.61	$\int e^{\frac{1}{2}i\text{ArcTan}(ax)} x^2 dx$	349
3.62	$\int e^{\frac{1}{2}i\text{ArcTan}(ax)} x dx$	355
3.63	$\int e^{\frac{1}{2}i\text{ArcTan}(ax)} dx$	361
3.64	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x} dx$	366

3.65	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^2} dx$	372
3.66	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^3} dx$	376
3.67	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^4} dx$	381
3.68	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^5} dx$	386
3.69	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^6} dx$	391
3.70	$\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x^3 dx$	396
3.71	$\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x^2 dx$	402
3.72	$\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x dx$	408
3.73	$\int e^{\frac{3}{2}i\text{ArcTan}(ax)} dx$	414
3.74	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(ax)}}{x} dx$	419
3.75	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(ax)}}{x^2} dx$	425
3.76	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(ax)}}{x^3} dx$	429
3.77	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(ax)}}{x^4} dx$	434
3.78	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(ax)}}{x^5} dx$	439
3.79	$\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x^3 dx$	444
3.80	$\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x^2 dx$	450
3.81	$\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x dx$	456
3.82	$\int e^{\frac{5}{2}i\text{ArcTan}(ax)} dx$	462
3.83	$\int \frac{e^{\frac{5}{2}i\text{ArcTan}(ax)}}{x} dx$	468
3.84	$\int \frac{e^{\frac{5}{2}i\text{ArcTan}(ax)}}{x^2} dx$	474
3.85	$\int \frac{e^{\frac{5}{2}i\text{ArcTan}(ax)}}{x^3} dx$	478
3.86	$\int \frac{e^{\frac{5}{2}i\text{ArcTan}(ax)}}{x^4} dx$	483
3.87	$\int \frac{e^{\frac{5}{2}i\text{ArcTan}(ax)}}{x^5} dx$	488
3.88	$\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^3 dx$	493
3.89	$\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^2 dx$	499
3.90	$\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x dx$	505
3.91	$\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} dx$	511
3.92	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x} dx$	516
3.93	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^2} dx$	522
3.94	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^3} dx$	526
3.95	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^4} dx$	531
3.96	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^5} dx$	536
3.97	$\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x^3 dx$	541
3.98	$\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x^2 dx$	547
3.99	$\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x dx$	553
3.100	$\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} dx$	559

3.101	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)} x}{x} dx$	564
3.102	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)} x}{x^2} dx$	570
3.103	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)} x}{x^3} dx$	574
3.104	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)} x}{x^4} dx$	579
3.105	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)} x}{x^5} dx$	584
3.106	$\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^3 dx$	589
3.107	$\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^2 dx$	595
3.108	$\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x dx$	601
3.109	$\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} dx$	607
3.110	$\int \frac{e^{-\frac{5}{2}i\text{ArcTan}(ax)} x}{x} dx$	613
3.111	$\int \frac{e^{-\frac{5}{2}i\text{ArcTan}(ax)} x}{x^2} dx$	619
3.112	$\int \frac{e^{-\frac{5}{2}i\text{ArcTan}(ax)} x}{x^3} dx$	623
3.113	$\int \frac{e^{-\frac{5}{2}i\text{ArcTan}(ax)} x}{x^4} dx$	628
3.114	$\int \frac{e^{-\frac{5}{2}i\text{ArcTan}(ax)} x}{x^5} dx$	633
3.115	$\int e^{\frac{1}{3}i\text{ArcTan}(x)} x^2 dx$	638
3.116	$\int e^{\frac{1}{3}i\text{ArcTan}(x)} x dx$	644
3.117	$\int e^{\frac{1}{3}i\text{ArcTan}(x)} dx$	650
3.118	$\int \frac{e^{\frac{1}{3}i\text{ArcTan}(x)} x}{x} dx$	655
3.119	$\int \frac{e^{\frac{1}{3}i\text{ArcTan}(x)} x}{x^2} dx$	661
3.120	$\int \frac{e^{\frac{1}{3}i\text{ArcTan}(x)} x}{x^3} dx$	666
3.121	$\int \frac{e^{\frac{1}{3}i\text{ArcTan}(x)} x}{x^4} dx$	671
3.122	$\int e^{\frac{2}{3}i\text{ArcTan}(x)} x^2 dx$	677
3.123	$\int e^{\frac{2}{3}i\text{ArcTan}(x)} x dx$	681
3.124	$\int e^{\frac{2}{3}i\text{ArcTan}(x)} dx$	685
3.125	$\int \frac{e^{\frac{2}{3}i\text{ArcTan}(x)} x}{x} dx$	688
3.126	$\int \frac{e^{\frac{2}{3}i\text{ArcTan}(x)} x}{x^2} dx$	692
3.127	$\int \frac{e^{\frac{2}{3}i\text{ArcTan}(x)} x}{x^3} dx$	695
3.128	$\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x^2 dx$	699
3.129	$\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x dx$	706
3.130	$\int e^{\frac{1}{4}i\text{ArcTan}(ax)} dx$	712
3.131	$\int \frac{e^{\frac{1}{4}i\text{ArcTan}(ax)} x}{x} dx$	718
3.132	$\int \frac{e^{\frac{1}{4}i\text{ArcTan}(ax)} x}{x^2} dx$	726
3.133	$\int \frac{e^{\frac{1}{4}i\text{ArcTan}(ax)} x}{x^3} dx$	732
3.134	$\int e^{6i\text{ArcTan}(ax)} x^m dx$	738
3.135	$\int e^{4i\text{ArcTan}(ax)} x^m dx$	742
3.136	$\int e^{2i\text{ArcTan}(ax)} x^m dx$	746

3.137	$\int e^{-2i\text{ArcTan}(ax)} x^m dx$	749
3.138	$\int e^{-4i\text{ArcTan}(ax)} x^m dx$	752
3.139	$\int e^{-6i\text{ArcTan}(ax)} x^m dx$	756
3.140	$\int e^{3i\text{ArcTan}(ax)} x^m dx$	760
3.141	$\int e^{i\text{ArcTan}(ax)} x^m dx$	764
3.142	$\int e^{-i\text{ArcTan}(ax)} x^m dx$	767
3.143	$\int e^{-3i\text{ArcTan}(ax)} x^m dx$	770
3.144	$\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x^m dx$	774
3.145	$\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x^m dx$	777
3.146	$\int e^{\frac{1}{2}i\text{ArcTan}(ax)} x^m dx$	780
3.147	$\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^m dx$	783
3.148	$\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x^m dx$	786
3.149	$\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^m dx$	789
3.150	$\int e^{\frac{2\text{ArcTan}(x)}{3}} x^m dx$	792
3.151	$\int e^{\frac{\text{ArcTan}(x)}{3}} x^m dx$	795
3.152	$\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x^m dx$	798
3.153	$\int e^{in\text{ArcTan}(ax)} x^m dx$	801
3.154	$\int e^{in\text{ArcTan}(ax)} x^3 dx$	804
3.155	$\int e^{in\text{ArcTan}(ax)} x^2 dx$	808
3.156	$\int e^{in\text{ArcTan}(ax)} x dx$	812
3.157	$\int e^{in\text{ArcTan}(ax)} dx$	815
3.158	$\int \frac{e^{in\text{ArcTan}(ax)}}{x} dx$	818
3.159	$\int \frac{e^{in\text{ArcTan}(ax)}}{x^2} dx$	821
3.160	$\int \frac{e^{in\text{ArcTan}(ax)}}{x^3} dx$	824
3.161	$\int \frac{e^{in\text{ArcTan}(ax)}}{x^4} dx$	827
3.162	$\int e^{i\text{ArcTan}(a+bx)} x^4 dx$	831
3.163	$\int e^{i\text{ArcTan}(a+bx)} x^3 dx$	839
3.164	$\int e^{i\text{ArcTan}(a+bx)} x^2 dx$	845
3.165	$\int e^{i\text{ArcTan}(a+bx)} x dx$	851
3.166	$\int e^{i\text{ArcTan}(a+bx)} dx$	855
3.167	$\int \frac{e^{i\text{ArcTan}(a+bx)}}{x} dx$	859
3.168	$\int \frac{e^{i\text{ArcTan}(a+bx)}}{x^2} dx$	864
3.169	$\int \frac{e^{i\text{ArcTan}(a+bx)}}{x^3} dx$	868
3.170	$\int \frac{e^{i\text{ArcTan}(a+bx)}}{x^4} dx$	873
3.171	$\int e^{2i\text{ArcTan}(a+bx)} x^4 dx$	880
3.172	$\int e^{2i\text{ArcTan}(a+bx)} x^3 dx$	883
3.173	$\int e^{2i\text{ArcTan}(a+bx)} x^2 dx$	886
3.174	$\int e^{2i\text{ArcTan}(a+bx)} x dx$	889
3.175	$\int e^{2i\text{ArcTan}(a+bx)} dx$	892
3.176	$\int \frac{e^{2i\text{ArcTan}(a+bx)}}{x} dx$	895
3.177	$\int \frac{e^{2i\text{ArcTan}(a+bx)}}{x^2} dx$	898

3.178	$\int \frac{e^{2i\text{ArcTan}(a+bx)}}{x^3} dx$	901
3.179	$\int \frac{e^{2i\text{ArcTan}(a+bx)}}{x^4} dx$	905
3.180	$\int e^{3i\text{ArcTan}(a+bx)} x^4 dx$	909
3.181	$\int e^{3i\text{ArcTan}(a+bx)} x^3 dx$	918
3.182	$\int e^{3i\text{ArcTan}(a+bx)} x^2 dx$	926
3.183	$\int e^{3i\text{ArcTan}(a+bx)} x dx$	933
3.184	$\int e^{3i\text{ArcTan}(a+bx)} dx$	940
3.185	$\int \frac{e^{3i\text{ArcTan}(a+bx)}}{x} dx$	946
3.186	$\int \frac{e^{3i\text{ArcTan}(a+bx)}}{x^2} dx$	952
3.187	$\int \frac{e^{3i\text{ArcTan}(a+bx)}}{x^3} dx$	958
3.188	$\int \frac{e^{3i\text{ArcTan}(a+bx)}}{x^4} dx$	965
3.189	$\int e^{-i\text{ArcTan}(a+bx)} x^4 dx$	973
3.190	$\int e^{-i\text{ArcTan}(a+bx)} x^3 dx$	979
3.191	$\int e^{-i\text{ArcTan}(a+bx)} x^2 dx$	985
3.192	$\int e^{-i\text{ArcTan}(a+bx)} x dx$	990
3.193	$\int e^{-i\text{ArcTan}(a+bx)} dx$	994
3.194	$\int \frac{e^{-i\text{ArcTan}(a+bx)}}{x} dx$	998
3.195	$\int \frac{e^{-i\text{ArcTan}(a+bx)}}{x^2} dx$	1003
3.196	$\int \frac{e^{-i\text{ArcTan}(a+bx)}}{x^3} dx$	1008
3.197	$\int \frac{e^{-i\text{ArcTan}(a+bx)}}{x^4} dx$	1013
3.198	$\int e^{-2i\text{ArcTan}(a+bx)} x^4 dx$	1019
3.199	$\int e^{-2i\text{ArcTan}(a+bx)} x^3 dx$	1022
3.200	$\int e^{-2i\text{ArcTan}(a+bx)} x^2 dx$	1025
3.201	$\int e^{-2i\text{ArcTan}(a+bx)} x dx$	1028
3.202	$\int e^{-2i\text{ArcTan}(a+bx)} dx$	1031
3.203	$\int \frac{e^{-2i\text{ArcTan}(a+bx)}}{x} dx$	1034
3.204	$\int \frac{e^{-2i\text{ArcTan}(a+bx)}}{x^2} dx$	1037
3.205	$\int \frac{e^{-2i\text{ArcTan}(a+bx)}}{x^3} dx$	1040
3.206	$\int \frac{e^{-2i\text{ArcTan}(a+bx)}}{x^4} dx$	1044
3.207	$\int e^{-3i\text{ArcTan}(a+bx)} x^4 dx$	1048
3.208	$\int e^{-3i\text{ArcTan}(a+bx)} x^3 dx$	1055
3.209	$\int e^{-3i\text{ArcTan}(a+bx)} x^2 dx$	1061
3.210	$\int e^{-3i\text{ArcTan}(a+bx)} x dx$	1067
3.211	$\int e^{-3i\text{ArcTan}(a+bx)} dx$	1073
3.212	$\int \frac{e^{-3i\text{ArcTan}(a+bx)}}{x} dx$	1078
3.213	$\int \frac{e^{-3i\text{ArcTan}(a+bx)}}{x^2} dx$	1084
3.214	$\int \frac{e^{-3i\text{ArcTan}(a+bx)}}{x^3} dx$	1089
3.215	$\int \frac{e^{-3i\text{ArcTan}(a+bx)}}{x^4} dx$	1095
3.216	$\int e^{\frac{1}{2}i\text{ArcTan}(a+bx)} x^2 dx$	1102
3.217	$\int e^{\frac{1}{2}i\text{ArcTan}(a+bx)} x dx$	1108

3.218	$\int e^{\frac{1}{2}i\text{ArcTan}(a+bx)} dx$	1114
3.219	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(a+bx)}}{x} dx$	1119
3.220	$\int \frac{e^{\frac{1}{2}i\text{ArcTan}(a+bx)}}{x^2} dx$	1125
3.221	$\int e^{\frac{3}{2}i\text{ArcTan}(a+bx)} x^2 dx$	1130
3.222	$\int e^{\frac{3}{2}i\text{ArcTan}(a+bx)} x dx$	1136
3.223	$\int e^{\frac{3}{2}i\text{ArcTan}(a+bx)} dx$	1142
3.224	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(a+bx)}}{x} dx$	1147
3.225	$\int \frac{e^{\frac{3}{2}i\text{ArcTan}(a+bx)}}{x^2} dx$	1153
3.226	$\int e^{-\frac{1}{2}i\text{ArcTan}(a+bx)} x^2 dx$	1158
3.227	$\int e^{-\frac{1}{2}i\text{ArcTan}(a+bx)} x dx$	1164
3.228	$\int e^{-\frac{1}{2}i\text{ArcTan}(a+bx)} dx$	1170
3.229	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(a+bx)}}{x} dx$	1175
3.230	$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(a+bx)}}{x^2} dx$	1181
3.231	$\int e^{-\frac{3}{2}i\text{ArcTan}(a+bx)} x^2 dx$	1186
3.232	$\int e^{-\frac{3}{2}i\text{ArcTan}(a+bx)} x dx$	1192
3.233	$\int e^{-\frac{3}{2}i\text{ArcTan}(a+bx)} dx$	1198
3.234	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(a+bx)}}{x} dx$	1203
3.235	$\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(a+bx)}}{x^2} dx$	1209
3.236	$\int e^{n\text{ArcTan}(a+bx)} x^m dx$	1214
3.237	$\int e^{n\text{ArcTan}(a+bx)} x^3 dx$	1217
3.238	$\int e^{n\text{ArcTan}(a+bx)} x^2 dx$	1221
3.239	$\int e^{n\text{ArcTan}(a+bx)} x dx$	1225
3.240	$\int e^{n\text{ArcTan}(a+bx)} dx$	1228
3.241	$\int \frac{e^{n\text{ArcTan}(a+bx)}}{x} dx$	1231
3.242	$\int \frac{e^{n\text{ArcTan}(a+bx)}}{x^2} dx$	1235
3.243	$\int \frac{e^{n\text{ArcTan}(a+bx)}}{x^3} dx$	1238
3.244	$\int e^{\text{ArcTan}(ax)} (c + a^2 cx^2)^p dx$	1241
3.245	$\int e^{\text{ArcTan}(ax)} (c + a^2 cx^2)^2 dx$	1244
3.246	$\int e^{\text{ArcTan}(ax)} (c + a^2 cx^2) dx$	1247
3.247	$\int e^{\text{ArcTan}(ax)} dx$	1250
3.248	$\int \frac{e^{\text{ArcTan}(ax)}}{c + a^2 cx^2} dx$	1253
3.249	$\int \frac{e^{\text{ArcTan}(ax)}}{(c + a^2 cx^2)^2} dx$	1256
3.250	$\int \frac{e^{\text{ArcTan}(ax)}}{(c + a^2 cx^2)^3} dx$	1259
3.251	$\int \frac{e^{\text{ArcTan}(ax)}}{(c + a^2 cx^2)^4} dx$	1262
3.252	$\int \frac{e^{\text{ArcTan}(ax)}}{(c + a^2 cx^2)^5} dx$	1266
3.253	$\int e^{\text{ArcTan}(ax)} (c + a^2 cx^2)^{3/2} dx$	1270
3.254	$\int e^{\text{ArcTan}(ax)} \sqrt{c + a^2 cx^2} dx$	1273

3.255	$\int \frac{e^{\text{ArcTan}(ax)}}{\sqrt{c+a^2cx^2}} dx$	1276
3.256	$\int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1279
3.257	$\int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$	1282
3.258	$\int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$	1285
3.259	$\int e^{2\text{ArcTan}(ax)}(c+a^2cx^2)^p dx$	1288
3.260	$\int e^{2\text{ArcTan}(ax)}(c+a^2cx^2)^2 dx$	1291
3.261	$\int e^{2\text{ArcTan}(ax)}(c+a^2cx^2) dx$	1294
3.262	$\int e^{2\text{ArcTan}(ax)} dx$	1297
3.263	$\int \frac{e^{2\text{ArcTan}(ax)}}{c+a^2cx^2} dx$	1300
3.264	$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$	1303
3.265	$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$	1306
3.266	$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$	1309
3.267	$\int e^{2\text{ArcTan}(ax)}(c+a^2cx^2)^{3/2} dx$	1313
3.268	$\int e^{2\text{ArcTan}(ax)}\sqrt{c+a^2cx^2} dx$	1316
3.269	$\int \frac{e^{2\text{ArcTan}(ax)}}{\sqrt{c+a^2cx^2}} dx$	1319
3.270	$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1322
3.271	$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$	1325
3.272	$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$	1328
3.273	$\int e^{-\text{ArcTan}(ax)}(c+a^2cx^2)^p dx$	1331
3.274	$\int e^{-\text{ArcTan}(ax)}(c+a^2cx^2)^2 dx$	1334
3.275	$\int e^{-\text{ArcTan}(ax)}(c+a^2cx^2) dx$	1337
3.276	$\int e^{-\text{ArcTan}(ax)} dx$	1340
3.277	$\int \frac{e^{-\text{ArcTan}(ax)}}{c+a^2cx^2} dx$	1343
3.278	$\int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$	1346
3.279	$\int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$	1349
3.280	$\int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$	1353
3.281	$\int e^{-\text{ArcTan}(ax)}(c+a^2cx^2)^{3/2} dx$	1356
3.282	$\int e^{-\text{ArcTan}(ax)}\sqrt{c+a^2cx^2} dx$	1359
3.283	$\int \frac{e^{-\text{ArcTan}(ax)}}{\sqrt{c+a^2cx^2}} dx$	1362
3.284	$\int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1365
3.285	$\int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$	1368
3.286	$\int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$	1371
3.287	$\int e^{-2\text{ArcTan}(ax)}(c+a^2cx^2)^p dx$	1374
3.288	$\int e^{-2\text{ArcTan}(ax)}(c+a^2cx^2)^2 dx$	1377

3.289	$\int e^{-2\text{ArcTan}(ax)}(c + a^2cx^2) dx$	1380
3.290	$\int e^{-2\text{ArcTan}(ax)} dx$	1383
3.291	$\int \frac{e^{-2\text{ArcTan}(ax)}}{c+a^2cx^2} dx$	1386
3.292	$\int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$	1389
3.293	$\int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$	1392
3.294	$\int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$	1395
3.295	$\int e^{-2\text{ArcTan}(ax)}(c + a^2cx^2)^{3/2} dx$	1398
3.296	$\int e^{-2\text{ArcTan}(ax)}\sqrt{c + a^2cx^2} dx$	1401
3.297	$\int \frac{e^{-2\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1404
3.298	$\int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1407
3.299	$\int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$	1410
3.300	$\int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$	1413
3.301	$\int \frac{e^{5i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1416
3.302	$\int \frac{e^{4i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1419
3.303	$\int \frac{e^{3i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1423
3.304	$\int \frac{e^{2i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1426
3.305	$\int \frac{e^{i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1430
3.306	$\int \frac{e^{-i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1433
3.307	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1436
3.308	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1440
3.309	$\int \frac{e^{-4i\text{ArcTan}(ax)}}{\sqrt{1 + a^2x^2}} dx$	1443
3.310	$\int \frac{e^{5i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1447
3.311	$\int \frac{e^{4i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1451
3.312	$\int \frac{e^{3i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1455
3.313	$\int \frac{e^{2i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1459
3.314	$\int \frac{e^{i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1463
3.315	$\int \frac{e^{-i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1467
3.316	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1471
3.317	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$	1475

3.318	$\int \frac{e^{-4i\text{ArcTan}(ax)}}{\sqrt{c+a^2cx^2}} dx$	1479
3.319	$\int \frac{e^{5i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1483
3.320	$\int \frac{e^{4i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1486
3.321	$\int \frac{e^{3i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1490
3.322	$\int \frac{e^{2i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1493
3.323	$\int \frac{e^{i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1497
3.324	$\int \frac{e^{-i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1500
3.325	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1503
3.326	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1507
3.327	$\int \frac{e^{-4i\text{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$	1510
3.328	$\int \frac{e^{5i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1514
3.329	$\int \frac{e^{4i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1518
3.330	$\int \frac{e^{3i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1522
3.331	$\int \frac{e^{2i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1526
3.332	$\int \frac{e^{i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1530
3.333	$\int \frac{e^{-i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1534
3.334	$\int \frac{e^{-2i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1538
3.335	$\int \frac{e^{-3i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1542
3.336	$\int \frac{e^{-4i\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$	1546
3.337	$\int e^{n\text{ArcTan}(ax)}(c+a^2cx^2)^2 dx$	1550
3.338	$\int e^{n\text{ArcTan}(ax)}(c+a^2cx^2) dx$	1553
3.339	$\int e^{n\text{ArcTan}(ax)} dx$	1556
3.340	$\int \frac{e^{n\text{ArcTan}(ax)}x^3}{c+a^2cx^2} dx$	1559
3.341	$\int \frac{e^{n\text{ArcTan}(ax)}x^2}{c+a^2cx^2} dx$	1563
3.342	$\int \frac{e^{n\text{ArcTan}(ax)}x}{c+a^2cx^2} dx$	1567
3.343	$\int \frac{e^{n\text{ArcTan}(ax)}}{c+a^2cx^2} dx$	1570
3.344	$\int \frac{e^{n\text{ArcTan}(ax)}}{x(c+a^2cx^2)} dx$	1573
3.345	$\int \frac{e^{n\text{ArcTan}(ax)}}{x^2(c+a^2cx^2)} dx$	1576
3.346	$\int \frac{e^{n\text{ArcTan}(ax)}}{x^3(c+a^2cx^2)} dx$	1580
3.347	$\int \frac{e^{n\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$	1584
3.348	$\int e^{n\text{ArcTan}(ax)}(c+a^2cx^2)^{3/2} dx$	1588

3.349	$\int e^{n\text{ArcTan}(ax)} \sqrt{c + a^2 cx^2} dx$	1591
3.350	$\int \frac{e^{n\text{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$	1594
3.351	$\int e^{n\text{ArcTan}(ax)} x^2 (c + a^2 cx^2)^{3/2} dx$	1597
3.352	$\int e^{n\text{ArcTan}(ax)} x^2 \sqrt{c + a^2 cx^2} dx$	1601
3.353	$\int \frac{e^{n\text{ArcTan}(ax)} x^3}{\sqrt{c + a^2 cx^2}} dx$	1605
3.354	$\int \frac{e^{n\text{ArcTan}(ax)} x^2}{\sqrt{c + a^2 cx^2}} dx$	1609
3.355	$\int \frac{e^{n\text{ArcTan}(ax)} x}{\sqrt{c + a^2 cx^2}} dx$	1613
3.356	$\int \frac{e^{n\text{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$	1617
3.357	$\int \frac{e^{n\text{ArcTan}(ax)}}{x \sqrt{c + a^2 cx^2}} dx$	1620
3.358	$\int \frac{e^{n\text{ArcTan}(ax)}}{x^2 \sqrt{c + a^2 cx^2}} dx$	1623
3.359	$\int \frac{e^{n\text{ArcTan}(ax)}}{x^3 \sqrt{c + a^2 cx^2}} dx$	1627
3.360	$\int e^{n\text{ArcTan}(ax)} \sqrt[3]{c + a^2 cx^2} dx$	1631
3.361	$\int \frac{e^{n\text{ArcTan}(ax)}}{\sqrt[3]{c + a^2 cx^2}} dx$	1634
3.362	$\int \frac{e^{n\text{ArcTan}(ax)}}{(c + a^2 cx^2)^{2/3}} dx$	1637
3.363	$\int \frac{e^{n\text{ArcTan}(ax)}}{(c + a^2 cx^2)^{4/3}} dx$	1640
3.364	$\int e^{n\text{ArcTan}(ax)} x^m (c + a^2 cx^2) dx$	1643
3.365	$\int \frac{e^{n\text{ArcTan}(ax)} x^m}{c + a^2 cx^2} dx$	1646
3.366	$\int \frac{e^{n\text{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^2} dx$	1649
3.367	$\int \frac{e^{n\text{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^3} dx$	1652
3.368	$\int \frac{e^{n\text{ArcTan}(ax)} x^m}{\sqrt{c + a^2 cx^2}} dx$	1655
3.369	$\int \frac{e^{n\text{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^{3/2}} dx$	1658
3.370	$\int \frac{e^{n\text{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^{5/2}} dx$	1661
3.371	$\int e^{n\text{ArcTan}(ax)} (c + a^2 cx^2)^p dx$	1664
3.372	$\int e^{-2ip\text{ArcTan}(ax)} (c + a^2 cx^2)^p dx$	1667
3.373	$\int e^{2ip\text{ArcTan}(ax)} (c + a^2 cx^2)^p dx$	1671
3.374	$\int e^{in\text{ArcTan}(ax)} x^2 (c + a^2 cx^2)^{-1 - \frac{n^2}{2}} dx$	1674
3.375	$\int \frac{e^{6i\text{ArcTan}(ax)} x^2}{(c + a^2 cx^2)^{19}} dx$	1677
3.376	$\int \frac{e^{4i\text{ArcTan}(ax)} x^2}{(c + a^2 cx^2)^9} dx$	1681
3.377	$\int \frac{e^{2i\text{ArcTan}(ax)} x^2}{(c + a^2 cx^2)^3} dx$	1685
3.378	$\int \frac{e^{-2i\text{ArcTan}(ax)} x^2}{(c + a^2 cx^2)^3} dx$	1688
3.379	$\int \frac{e^{-4i\text{ArcTan}(ax)} x^2}{(c + a^2 cx^2)^9} dx$	1691

3.380	$\int \frac{e^{5i\text{ArcTan}(ax)x^2}}{(c+a^2cx^2)^{27/2}} dx$	1695
3.381	$\int \frac{e^{3i\text{ArcTan}(ax)x^2}}{(c+a^2cx^2)^{11/2}} dx$	1699
3.382	$\int \frac{e^{i\text{ArcTan}(ax)x^2}}{(c+a^2cx^2)^{3/2}} dx$	1703
3.383	$\int \frac{e^{-i\text{ArcTan}(ax)x^2}}{(c+a^2cx^2)^{3/2}} dx$	1707
3.384	$\int \frac{e^{-3i\text{ArcTan}(ax)x^2}}{(c+a^2cx^2)^{11/2}} dx$	1711
3.385	$\int \frac{e^{-5i\text{ArcTan}(ax)x^2}}{(c+a^2cx^2)^{27/2}} dx$	1715

3.1 $\int e^{i\text{ArcTan}(ax)} x^4 dx$

Optimal. Leaf size=113

$$-\frac{4ix^2\sqrt{1+a^2x^2}}{15a^3} + \frac{x^3\sqrt{1+a^2x^2}}{4a^2} + \frac{ix^4\sqrt{1+a^2x^2}}{5a} + \frac{(64i-45ax)\sqrt{1+a^2x^2}}{120a^5} + \frac{3\sinh^{-1}(ax)}{8a^5}$$

[Out] $3/8*\text{arcsinh}(a*x)/a^5-4/15*I*x^2*(a^2*x^2+1)^{(1/2)}/a^3+1/4*x^3*(a^2*x^2+1)^{(1/2)}/a^2+1/5*I*x^4*(a^2*x^2+1)^{(1/2)}/a+1/120*(64*I-45*a*x)*(a^2*x^2+1)^{(1/2)}/a^5$

Rubi [A]

time = 0.07, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5168, 847, 794, 221}

$$\frac{3\sinh^{-1}(ax)}{8a^5} + \frac{ix^4\sqrt{a^2x^2+1}}{5a} + \frac{x^3\sqrt{a^2x^2+1}}{4a^2} + \frac{(-45ax+64i)\sqrt{a^2x^2+1}}{120a^5} - \frac{4ix^2\sqrt{a^2x^2+1}}{15a^3}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])*x^4,x]

[Out] $(((-4*I)/15)*x^2*\text{Sqrt}[1+a^2*x^2])/a^3 + (x^3*\text{Sqrt}[1+a^2*x^2])/(4*a^2) + ((I/5)*x^4*\text{Sqrt}[1+a^2*x^2])/a + ((64*I-45*a*x)*\text{Sqrt}[1+a^2*x^2])/(120*a^5) + (3*\text{ArcSinh}[a*x])/(8*a^5)$

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 794

Int[((d_.) + (e_.)*(x_))*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x)*((a + c*x^2)^(p + 1)/(2*c*(p + 1)*(2*p + 3))), x] - Dist[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && !LeQ[p, -1]

Rule 847

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[g*(d + e*x)^m*((a + c*x^2)^(p + 1)/(c*(m + 2*p + 2))), x] + Dist[1/(c*(m + 2*p + 2)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^p*Simp[c*d*f*(m + 2*p + 2) - a*e*g*m + c*(e*f*(m + 2*p + 2) + d*g*m)*x, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 + a*e^2, 0] && GtQ[m, 0] && NeQ[m + 2*p + 2, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p])

) && !(IGtQ[m, 0] && EqQ[f, 0])

Rule 5168

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*n + 1)/2)/((1 + I*a*x)^(I*n - 1)/2)*Sqrt[1 + a^2*x^2]], x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int e^{i \tan^{-1}(ax)} x^4 dx &= \int \frac{x^4(1 + iax)}{\sqrt{1 + a^2x^2}} dx \\
 &= \frac{ix^4\sqrt{1 + a^2x^2}}{5a} + \frac{\int \frac{x^3(-4ia + 5a^2x)}{\sqrt{1 + a^2x^2}} dx}{5a^2} \\
 &= \frac{x^3\sqrt{1 + a^2x^2}}{4a^2} + \frac{ix^4\sqrt{1 + a^2x^2}}{5a} + \frac{\int \frac{x^2(-15a^2 - 16ia^3x)}{\sqrt{1 + a^2x^2}} dx}{20a^4} \\
 &= -\frac{4ix^2\sqrt{1 + a^2x^2}}{15a^3} + \frac{x^3\sqrt{1 + a^2x^2}}{4a^2} + \frac{ix^4\sqrt{1 + a^2x^2}}{5a} + \frac{\int \frac{x(32ia^3 - 45a^4x)}{\sqrt{1 + a^2x^2}} dx}{60a^6} \\
 &= -\frac{4ix^2\sqrt{1 + a^2x^2}}{15a^3} + \frac{x^3\sqrt{1 + a^2x^2}}{4a^2} + \frac{ix^4\sqrt{1 + a^2x^2}}{5a} + \frac{(64i - 45ax)\sqrt{1 + a^2x^2}}{120a^5} + \frac{3 \int}{120a^5} \\
 &= -\frac{4ix^2\sqrt{1 + a^2x^2}}{15a^3} + \frac{x^3\sqrt{1 + a^2x^2}}{4a^2} + \frac{ix^4\sqrt{1 + a^2x^2}}{5a} + \frac{(64i - 45ax)\sqrt{1 + a^2x^2}}{120a^5} + \frac{3 \operatorname{si}}{120a^5}
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 64, normalized size = 0.57

$$\frac{\sqrt{1 + a^2x^2} (64i - 45ax - 32ia^2x^2 + 30a^3x^3 + 24ia^4x^4) + 45 \sinh^{-1}(ax)}{120a^5}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a*x])*x^4,x]

[Out] (Sqrt[1 + a^2*x^2]*(64*I - 45*a*x - (32*I)*a^2*x^2 + 30*a^3*x^3 + (24*I)*a^4*x^4) + 45*ArcSinh[a*x])/(120*a^5)

Maple [A]

time = 0.10, size = 142, normalized size = 1.26

method	result
--------	--------

risch	$\frac{i(24a^4x^4 - 30ia^3x^3 - 32a^2x^2 + 45iax + 64)\sqrt{a^2x^2 + 1}}{120a^5} + \frac{3 \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + 1}\right)}{8a^4\sqrt{a^2}}$
meijerg	$\frac{-\sqrt{\pi} x (a^2)^{\frac{5}{2}} (-10a^2x^2 + 15)\sqrt{a^2x^2 + 1}}{20a^4} + \frac{3\sqrt{\pi} (a^2)^{\frac{5}{2}} \operatorname{arcsinh}(ax)}{4a^5} + \frac{i\left(-\frac{16\sqrt{\pi}}{15} + \frac{\sqrt{\pi} (6a^4x^4 - 8a^2x^2 + 16)\sqrt{a^2x^2 + 1}}{15}\right)}{2a^5\sqrt{\pi}}$
default	$ia \left(\frac{x^4\sqrt{a^2x^2 + 1}}{5a^2} - \frac{4\left(\frac{x^2\sqrt{a^2x^2 + 1}}{3a^2} - \frac{2\sqrt{a^2x^2 + 1}}{3a^4}\right)}{5a^2} \right) + \frac{x^3\sqrt{a^2x^2 + 1}}{4a^2} - \frac{3\left(\frac{x\sqrt{a^2x^2 + 1}}{2a^2} - \frac{\ln\left(\frac{a}{\sqrt{a^2x^2 + 1}}\right)}{4a^2}\right)}{4a^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^4,x,method=_RETURNVERBOSE)`

[Out] $I*a*(1/5*x^4/a^2*(a^2*x^2+1)^{(1/2)} - 4/5/a^2*(1/3*x^2*(a^2*x^2+1)^{(1/2)}/a^2 - 2/3*(a^2*x^2+1)^{(1/2)}/a^4) + 1/4*x^3*(a^2*x^2+1)^{(1/2)}/a^2 - 3/4/a^2*(1/2*x*(a^2*x^2+1)^{(1/2)}/a^2 - 1/2/a^2*\ln(a^2*x/(a^2)^{(1/2)} + (a^2*x^2+1)^{(1/2)})/(a^2)^{(1/2)})$

Maxima [A]

time = 0.27, size = 100, normalized size = 0.88

$$\frac{i\sqrt{a^2x^2 + 1}x^4}{5a} + \frac{\sqrt{a^2x^2 + 1}x^3}{4a^2} - \frac{4i\sqrt{a^2x^2 + 1}x^2}{15a^3} - \frac{3\sqrt{a^2x^2 + 1}x}{8a^4} + \frac{3\operatorname{arsinh}(ax)}{8a^5} + \frac{8i\sqrt{a^2x^2 + 1}}{15a^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^4,x, algorithm="maxima")`

[Out] $1/5*I*\sqrt{a^2*x^2 + 1}*x^4/a + 1/4*\sqrt{a^2*x^2 + 1}*x^3/a^2 - 4/15*I*\sqrt{a^2*x^2 + 1}*x^2/a^3 - 3/8*\sqrt{a^2*x^2 + 1}*x/a^4 + 3/8*\operatorname{arcsinh}(a*x)/a^5 + 8/15*I*\sqrt{a^2*x^2 + 1}/a^5$

Fricas [A]

time = 1.37, size = 67, normalized size = 0.59

$$\frac{(24ia^4x^4 + 30a^3x^3 - 32ia^2x^2 - 45ax + 64i)\sqrt{a^2x^2 + 1} - 45 \log(-ax + \sqrt{a^2x^2 + 1})}{120a^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^4,x, algorithm="fricas")`

[Out] $1/120*((24*I*a^4*x^4 + 30*a^3*x^3 - 32*I*a^2*x^2 - 45*a*x + 64*I)*\sqrt{a^2*x^2 + 1} - 45*\log(-a*x + \sqrt{a^2*x^2 + 1}))/a^5$

Sympy [A]

time = 3.53, size = 138, normalized size = 1.22

$$ia \left(\begin{cases} \frac{x^4 \sqrt{a^2 x^2 + 1}}{5a^2} - \frac{4x^2 \sqrt{a^2 x^2 + 1}}{15a^4} + \frac{8 \sqrt{a^2 x^2 + 1}}{15a^6} & \text{for } a \neq 0 \\ \frac{x^6}{6} & \text{otherwise} \end{cases} \right) + \frac{x^5}{4\sqrt{a^2 x^2 + 1}} - \frac{x^3}{8a^2 \sqrt{a^2 x^2 + 1}} - \frac{3x}{8a^4 \sqrt{a^2 x^2 + 1}} + \frac{3 \operatorname{asinh}(ax)}{8a^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)*x**4,x)

[Out] I*a*Piecewise((x**4*sqrt(a**2*x**2 + 1)/(5*a**2) - 4*x**2*sqrt(a**2*x**2 + 1)/(15*a**4) + 8*sqrt(a**2*x**2 + 1)/(15*a**6), Ne(a, 0)), (x**6/6, True)) + x**5/(4*sqrt(a**2*x**2 + 1)) - x**3/(8*a**2*sqrt(a**2*x**2 + 1)) - 3*x/(8*a**4*sqrt(a**2*x**2 + 1)) + 3*asinh(a*x)/(8*a**5)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^4,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.10, size = 98, normalized size = 0.87

$$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{x^3 (a^2)^{3/2}}{4a^4} - \frac{3x \sqrt{a^2}}{8a^4} + \frac{a 8i}{15 (a^2)^{5/2}} - \frac{a^3 x^2 4i}{15 (a^2)^{5/2}} + \frac{a^5 x^4 1i}{5 (a^2)^{5/2}} \right)}{\sqrt{a^2}} + \frac{3 \operatorname{asinh}(x \sqrt{a^2})}{8 a^4 \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*(a*x*1i + 1))/(a^2*x^2 + 1)^(1/2),x)

[Out] ((a^2*x^2 + 1)^(1/2)*((a*8i)/(15*(a^2)^(5/2)) - (a^3*x^2*4i)/(15*(a^2)^(5/2))) + (x^3*(a^2)^(3/2))/(4*a^4) + (a^5*x^4*1i)/(5*(a^2)^(5/2)) - (3*x*(a^2)^(1/2))/(8*a^4))/(a^2)^(1/2) + (3*asinh(x*(a^2)^(1/2)))/(8*a^4*(a^2)^(1/2))

3.2 $\int e^{i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=90

$$\frac{x^2 \sqrt{1+a^2x^2}}{3a^2} + \frac{ix^3 \sqrt{1+a^2x^2}}{4a} - \frac{(16+9iax)\sqrt{1+a^2x^2}}{24a^4} + \frac{3i \sinh^{-1}(ax)}{8a^4}$$

[Out] $3/8*I*\text{arcsinh}(a*x)/a^4+1/3*x^2*(a^2*x^2+1)^{(1/2)}/a^2+1/4*I*x^3*(a^2*x^2+1)^{(1/2)}/a-1/24*(16+9*I*a*x)*(a^2*x^2+1)^{(1/2)}/a^4$

Rubi [A]

time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5168, 847, 794, 221}

$$\frac{3i \sinh^{-1}(ax)}{8a^4} + \frac{x^2 \sqrt{a^2x^2+1}}{3a^2} + \frac{ix^3 \sqrt{a^2x^2+1}}{4a} - \frac{(16+9iax)\sqrt{a^2x^2+1}}{24a^4}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])*x^3,x]

[Out] $(x^2*\text{Sqrt}[1+a^2*x^2])/(3*a^2) + ((I/4)*x^3*\text{Sqrt}[1+a^2*x^2])/a - ((16+(9*I)*a*x)*\text{Sqrt}[1+a^2*x^2])/(24*a^4) + (((3*I)/8)*\text{ArcSinh}[a*x])/a^4$

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 794

Int[((d_.) + (e_.)*(x_))*((f_.) + (g_.)*(x_))*((a_.) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x)*((a + c*x^2)^(p + 1)/(2*c*(p + 1)*(2*p + 3))), x] - Dist[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && !LeQ[p, -1]

Rule 847

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[g*(d + e*x)^m*((a + c*x^2)^(p + 1)/(c*(m + 2*p + 2))), x] + Dist[1/(c*(m + 2*p + 2)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^p*Simp[c*d*f*(m + 2*p + 2) - a*e*g*m + c*(e*f*(m + 2*p + 2) + d*g*m)*x, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 + a*e^2, 0] && GtQ[m, 0] && NeQ[m + 2*p + 2, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p]) && !(IGtQ[m, 0] && EqQ[f, 0])

Rule 5168

Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(((I*n + 1)/2)/((1 + I*a*x)^(((I*n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int e^{i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1 + iax)}{\sqrt{1 + a^2x^2}} dx \\
 &= \frac{ix^3\sqrt{1 + a^2x^2}}{4a} + \frac{\int \frac{x^2(-3ia + 4a^2x)}{\sqrt{1 + a^2x^2}} dx}{4a^2} \\
 &= \frac{x^2\sqrt{1 + a^2x^2}}{3a^2} + \frac{ix^3\sqrt{1 + a^2x^2}}{4a} + \frac{\int \frac{x(-8a^2 - 9ia^3x)}{\sqrt{1 + a^2x^2}} dx}{12a^4} \\
 &= \frac{x^2\sqrt{1 + a^2x^2}}{3a^2} + \frac{ix^3\sqrt{1 + a^2x^2}}{4a} - \frac{(16 + 9iax)\sqrt{1 + a^2x^2}}{24a^4} + \frac{(3i) \int \frac{1}{\sqrt{1 + a^2x^2}} dx}{8a^3} \\
 &= \frac{x^2\sqrt{1 + a^2x^2}}{3a^2} + \frac{ix^3\sqrt{1 + a^2x^2}}{4a} - \frac{(16 + 9iax)\sqrt{1 + a^2x^2}}{24a^4} + \frac{3i \sinh^{-1}(ax)}{8a^4}
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 56, normalized size = 0.62

$$\frac{\sqrt{1 + a^2x^2}(-16 - 9iax + 8a^2x^2 + 6ia^3x^3) + 9i \sinh^{-1}(ax)}{24a^4}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a*x])*x^3,x]

[Out] (Sqrt[1 + a^2*x^2]*(-16 - (9*I)*a*x + 8*a^2*x^2 + (6*I)*a^3*x^3) + (9*I)*ArcSinh[a*x])/(24*a^4)

Maple [A]

time = 0.07, size = 117, normalized size = 1.30

method	result
risch	$\frac{i(6a^3x^3 - 8a^2x^2 - 9ax + 16i)\sqrt{a^2x^2 + 1}}{24a^4} + \frac{3i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + 1}\right)}{8a^3\sqrt{a^2}}$

meijerg	$\frac{\frac{4\sqrt{\pi}}{3} - \frac{\sqrt{\pi}(-4a^2x^2+8)\sqrt{a^2x^2+1}}{6}}{2a^4\sqrt{\pi}} + i \left(\frac{-\sqrt{\pi} x (a^2)^{\frac{5}{2}} (-10a^2x^2+15)\sqrt{a^2x^2+1}}{20a^4} + \frac{3\sqrt{\pi} (a^2)^{\frac{5}{2}} \operatorname{arcsinh}(ax)}{4a^5} \right)}{2a^3\sqrt{\pi}\sqrt{a^2}}$
default	$ia \left(\frac{x^3\sqrt{a^2x^2+1}}{4a^2} - \frac{\frac{x\sqrt{a^2x^2+1}}{2a^2} - \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a^2\sqrt{a^2}}}{4a^2} \right) + \frac{x^2\sqrt{a^2x^2+1}}{3a^2} - \frac{2\sqrt{a^2x^2+1}}{3a^4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^3,x,method=_RETURNVERBOSE)`

[Out] $I*a*(1/4*x^3*(a^2*x^2+1)^{(1/2)}/a^2-3/4/a^2*(1/2*x*(a^2*x^2+1)^{(1/2)}/a^2-1/2/a^2*\ln(a^2*x/(a^2)^{(1/2)}+(a^2*x^2+1)^{(1/2)})/(a^2)^{(1/2)}))+1/3*x^2*(a^2*x^2+1)^{(1/2)}/a^2-2/3*(a^2*x^2+1)^{(1/2)}/a^4$

Maxima [A]

time = 0.27, size = 81, normalized size = 0.90

$$\frac{i\sqrt{a^2x^2+1}x^3}{4a} + \frac{\sqrt{a^2x^2+1}x^2}{3a^2} - \frac{3i\sqrt{a^2x^2+1}x}{8a^3} + \frac{3i\operatorname{arsinh}(ax)}{8a^4} - \frac{2\sqrt{a^2x^2+1}}{3a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^3,x, algorithm="maxima")`

[Out] $1/4*I*\sqrt{a^2*x^2+1}*x^3/a + 1/3*\sqrt{a^2*x^2+1}*x^2/a^2 - 3/8*I*\sqrt{a^2*x^2+1}*x/a^3 + 3/8*I*\operatorname{arcsinh}(a*x)/a^4 - 2/3*\sqrt{a^2*x^2+1}/a^4$

Fricas [A]

time = 1.24, size = 59, normalized size = 0.66

$$\frac{(6i a^3 x^3 + 8 a^2 x^2 - 9i a x - 16)\sqrt{a^2 x^2 + 1} - 9i \log(-a x + \sqrt{a^2 x^2 + 1})}{24 a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^3,x, algorithm="fricas")`

[Out] $1/24*((6*I*a^3*x^3 + 8*a^2*x^2 - 9*I*a*x - 16)*\sqrt{a^2*x^2+1} - 9*I*\log(-a*x + \sqrt{a^2*x^2+1}))/a^4$

Sympy [A]

time = 3.55, size = 119, normalized size = 1.32

$$\frac{iax^5}{4\sqrt{a^2x^2+1}} + \begin{cases} \frac{x^2\sqrt{a^2x^2+1}}{3a^2} - \frac{2\sqrt{a^2x^2+1}}{3a^4} & \text{for } a \neq 0 \\ \frac{x^4}{4} & \text{otherwise} \end{cases} - \frac{ix^3}{8a\sqrt{a^2x^2+1}} - \frac{3ix}{8a^3\sqrt{a^2x^2+1}} + \frac{3i\operatorname{asinh}(ax)}{8a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)*x**3,x)

[Out] I*a*x**5/(4*sqrt(a**2*x**2 + 1)) + Piecewise((x**2*sqrt(a**2*x**2 + 1)/(3*a**2) - 2*sqrt(a**2*x**2 + 1)/(3*a**4), Ne(a, 0)), (x**4/4, True)) - I*x**3/(8*a*sqrt(a**2*x**2 + 1)) - 3*I*x/(8*a**3*sqrt(a**2*x**2 + 1)) + 3*I*asinh(a*x)/(8*a**4)

Giac [A]

time = 0.42, size = 70, normalized size = 0.78

$$-\frac{1}{24} \sqrt{a^2 x^2 + 1} \left(\left(2x \left(-\frac{3ix}{a} - \frac{4}{a^2} \right) + \frac{9i}{a^3} \right) x + \frac{16}{a^4} \right) - \frac{3i \log \left(-x|a| + \sqrt{a^2 x^2 + 1} \right)}{8a^3|a|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^3,x, algorithm="giac")

[Out] -1/24*sqrt(a^2*x^2 + 1)*((2*x*(-3*I*x/a - 4/a^2) + 9*I/a^3)*x + 16/a^4) - 3/8*I*log(-x*abs(a) + sqrt(a^2*x^2 + 1))/(a^3*abs(a))

Mupad [B]

time = 0.43, size = 85, normalized size = 0.94

$$\frac{\operatorname{asinh}\left(x\sqrt{a^2}\right) 3i}{8a^3\sqrt{a^2}} - \frac{\sqrt{a^2x^2+1} \left(\frac{2}{3(a^2)^{3/2}} - \frac{a^2x^2}{3(a^2)^{3/2}} - \frac{x^3(a^2)^{3/2} 1i}{4a^3} + \frac{x\sqrt{a^2} 3i}{8a^3} \right)}{\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a*x*1i + 1))/(a^2*x^2 + 1)^(1/2),x)

[Out] (asinh(x*(a^2)^(1/2))*3i)/(8*a^3*(a^2)^(1/2)) - ((a^2*x^2 + 1)^(1/2)*(2/(3*(a^2)^(3/2)) - (a^2*x^2)/(3*(a^2)^(3/2)) - (x^3*(a^2)^(3/2)*1i)/(4*a^3) + (x*(a^2)^(1/2)*3i)/(8*a^3)))/(a^2)^(1/2)

3.3 $\int e^{i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=75

$$-\frac{i\sqrt{1+a^2x^2}}{a^3} + \frac{x\sqrt{1+a^2x^2}}{2a^2} + \frac{i(1+a^2x^2)^{3/2}}{3a^3} - \frac{\sinh^{-1}(ax)}{2a^3}$$

[Out] $1/3*I*(a^2*x^2+1)^{(3/2)}/a^3-1/2*\text{arcsinh}(a*x)/a^3-I*(a^2*x^2+1)^{(1/2)}/a^3+1/2*x*(a^2*x^2+1)^{(1/2)}/a^2$

Rubi [A]

time = 0.04, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5168, 811, 655, 201, 221}

$$-\frac{\sinh^{-1}(ax)}{2a^3} + \frac{x\sqrt{a^2x^2+1}}{2a^2} + \frac{i(a^2x^2+1)^{3/2}}{3a^3} - \frac{i\sqrt{a^2x^2+1}}{a^3}$$

Antiderivative was successfully verified.

[In] `Int[E^(I*ArcTan[a*x])*x^2,x]`

[Out] $((-I)*\text{Sqrt}[1+a^2*x^2])/a^3 + (x*\text{Sqrt}[1+a^2*x^2])/(2*a^2) + ((I/3)*(1+a^2*x^2)^{(3/2)})/a^3 - \text{ArcSinh}[a*x]/(2*a^3)$

Rule 201

`Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x*((a + b*x^n)^p/(n*p + 1)), x] + Dist[a*n*(p/(n*p + 1)), Int[(a + b*x^n)^(p - 1), x], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])`

Rule 221

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 655

`Int[((d_) + (e_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*((a + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Dist[d, Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]`

Rule 811

`Int[(x_)^2*((f_) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/c, Int[(f + g*x)*(a + c*x^2)^(p + 1), x], x] - Dist[a/c, Int[(f + g*x)*`

$(a + c*x^2)^p, x]$, $x]$ /; FreeQ[{a, c, f, g, p}, x] && EqQ[a*g^2 + f^2*c, 0]

Rule 5168

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*n + 1)/2)/((1 + I*a*x)^(I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int e^{i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1 + iax)}{\sqrt{1 + a^2x^2}} dx \\
 &= -\frac{\int \frac{1+iax}{\sqrt{1+a^2x^2}} dx}{a^2} + \frac{\int (1+iax)\sqrt{1+a^2x^2} dx}{a^2} \\
 &= -\frac{i\sqrt{1+a^2x^2}}{a^3} + \frac{i(1+a^2x^2)^{3/2}}{3a^3} - \frac{\int \frac{1}{\sqrt{1+a^2x^2}} dx}{a^2} + \frac{\int \sqrt{1+a^2x^2} dx}{a^2} \\
 &= -\frac{i\sqrt{1+a^2x^2}}{a^3} + \frac{x\sqrt{1+a^2x^2}}{2a^2} + \frac{i(1+a^2x^2)^{3/2}}{3a^3} - \frac{\sinh^{-1}(ax)}{a^3} + \frac{\int \frac{1}{\sqrt{1+a^2x^2}} dx}{2a^2} \\
 &= -\frac{i\sqrt{1+a^2x^2}}{a^3} + \frac{x\sqrt{1+a^2x^2}}{2a^2} + \frac{i(1+a^2x^2)^{3/2}}{3a^3} - \frac{\sinh^{-1}(ax)}{2a^3}
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 46, normalized size = 0.61

$$\frac{(-4i + 3ax + 2ia^2x^2)\sqrt{1+a^2x^2} - 3\sinh^{-1}(ax)}{6a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a*x])*x^2,x]

[Out] ((-4*I + 3*a*x + (2*I)*a^2*x^2)*Sqrt[1 + a^2*x^2] - 3*ArcSinh[a*x])/(6*a^3)

Maple [A]

time = 0.06, size = 92, normalized size = 1.23

method	result	size
risch	$\frac{i(2a^2x^2 - 3iax - 4)\sqrt{a^2x^2 + 1}}{6a^3} - \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + 1}\right)}{2a^2\sqrt{a^2}}$	67

default	$ia \left(\frac{x^2 \sqrt{a^2 x^2 + 1}}{3a^2} - \frac{2\sqrt{a^2 x^2 + 1}}{3a^4} \right) + \frac{x\sqrt{a^2 x^2 + 1}}{2a^2} - \frac{\ln \left(\frac{a^2 x}{\sqrt{a^2}} + \sqrt{a^2 x^2 + 1} \right)}{2a^2 \sqrt{a^2}}$	92
meijerg	$\frac{\sqrt{\pi} x (a^2)^{\frac{3}{2}} \sqrt{a^2 x^2 + 1}}{a^2} - \frac{\sqrt{\pi} (a^2)^{\frac{3}{2}} \operatorname{arcsinh}(ax)}{a^3} + i \left(\frac{4\sqrt{\pi}}{3} - \frac{\sqrt{\pi} (-4a^2 x^2 + 8) \sqrt{a^2 x^2 + 1}}{6} \right)$	98

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2,x,method=_RETURNVERBOSE)`

[Out] $I*a*(1/3*x^2*(a^2*x^2+1)^{(1/2)}/a^2-2/3*(a^2*x^2+1)^{(1/2)}/a^4)+1/2*x*(a^2*x^2+1)^{(1/2)}/a^2-1/2/a^2*\ln(a^2*x/(a^2)^{(1/2)}+(a^2*x^2+1)^{(1/2)})/(a^2)^{(1/2)}$

Maxima [A]

time = 0.27, size = 62, normalized size = 0.83

$$\frac{i \sqrt{a^2 x^2 + 1} x^2}{3 a} + \frac{\sqrt{a^2 x^2 + 1} x}{2 a^2} - \frac{\operatorname{arsinh}(a x)}{2 a^3} - \frac{2 i \sqrt{a^2 x^2 + 1}}{3 a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2,x, algorithm="maxima")`

[Out] $1/3*I*\sqrt{a^2*x^2 + 1}*x^2/a + 1/2*\sqrt{a^2*x^2 + 1}*x/a^2 - 1/2*\operatorname{arcsinh}(a*x)/a^3 - 2/3*I*\sqrt{a^2*x^2 + 1}/a^3$

Fricas [A]

time = 1.94, size = 51, normalized size = 0.68

$$\frac{\sqrt{a^2 x^2 + 1} (2i a^2 x^2 + 3 a x - 4i) + 3 \log(-a x + \sqrt{a^2 x^2 + 1})}{6 a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2,x, algorithm="fricas")`

[Out] $1/6*(\sqrt{a^2*x^2 + 1}*(2*I*a^2*x^2 + 3*a*x - 4*I) + 3*\log(-a*x + \sqrt{a^2*x^2 + 1}))/a^3$

Sympy [A]

time = 2.14, size = 75, normalized size = 1.00

$$ia \left(\begin{cases} \frac{x^2 \sqrt{a^2 x^2 + 1}}{3a^2} - \frac{2\sqrt{a^2 x^2 + 1}}{3a^4} & \text{for } a \neq 0 \\ \frac{x^4}{4} & \text{otherwise} \end{cases} \right) + \frac{x\sqrt{a^2 x^2 + 1}}{2a^2} - \frac{\operatorname{asinh}(ax)}{2a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)*x**2,x)

[Out] I*a*Piecewise((x**2*sqrt(a**2*x**2 + 1)/(3*a**2) - 2*sqrt(a**2*x**2 + 1)/(3*a**4), Ne(a, 0)), (x**4/4, True)) + x*sqrt(a**2*x**2 + 1)/(2*a**2) - asinh(a*x)/(2*a**3)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.41, size = 71, normalized size = 0.95

$$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{x \sqrt{a^2}}{2a^2} - \frac{a 2i}{3(a^2)^{3/2}} + \frac{a^3 x^2 1i}{3(a^2)^{3/2}} \right)}{\sqrt{a^2}} - \frac{\operatorname{asinh}\left(x \sqrt{a^2}\right)}{2 a^2 \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1))/(a^2*x^2 + 1)^(1/2),x)

[Out] ((a^2*x^2 + 1)^(1/2)*((a^3*x^2*1i)/(3*(a^2)^(3/2)) - (a*2i)/(3*(a^2)^(3/2)) + (x*(a^2)^(1/2))/(2*a^2)))/(a^2)^(1/2) - asinh(x*(a^2)^(1/2))/(2*a^2*(a^2)^(1/2))

3.4 $\int e^{i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=42

$$\frac{(2 + iax)\sqrt{1 + a^2x^2}}{2a^2} - \frac{i \sinh^{-1}(ax)}{2a^2}$$

[Out] $-1/2*I*\text{arcsinh}(a*x)/a^2+1/2*(2+I*a*x)*(a^2*x^2+1)^{(1/2)}/a^2$

Rubi [A]

time = 0.01, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5168, 794, 221}

$$\frac{(2 + iax)\sqrt{a^2x^2 + 1}}{2a^2} - \frac{i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I*\text{ArcTan}[a*x])*x}, x]$

[Out] $((2 + I*a*x)*\text{Sqrt}[1 + a^2*x^2])/(2*a^2) - ((I/2)*\text{ArcSinh}[a*x])/a^2$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b]$

Rule 794

$\text{Int}[(d_) + (e_)*(x_))*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x*((a + c*x^2)^{(p + 1)/(2*c*(p + 1)*(2*p + 3))}), x] - \text{Dist}[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), \text{Int}[(a + c*x^2)^p, x], x] /; \text{FreeQ}\{a, c, d, e, f, g, p\}, x \ \&\& \ !\text{Le} \ Q[p, -1]$

Rule 5168

$\text{Int}[E^{(\text{ArcTan}[(a_)*(x_)]*(n_))*x)^{(m_)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{((I*n + 1)/2)/((1 + I*a*x)^{((I*n - 1)/2)*\text{Sqrt}[1 + a^2*x^2]})}), x] /; \text{Free} \ Q\{a, m\}, x \ \&\& \ \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
\int e^{i \tan^{-1}(ax)} x dx &= \int \frac{x(1+iax)}{\sqrt{1+a^2x^2}} dx \\
&= \frac{(2+iax)\sqrt{1+a^2x^2}}{2a^2} - \frac{i \int \frac{1}{\sqrt{1+a^2x^2}} dx}{2a} \\
&= \frac{(2+iax)\sqrt{1+a^2x^2}}{2a^2} - \frac{i \sinh^{-1}(ax)}{2a^2}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 38, normalized size = 0.90

$$\frac{(2+iax)\sqrt{1+a^2x^2} - i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*ArcTan[a*x])*x,x]``[Out] ((2 + I*a*x)*Sqrt[1 + a^2*x^2] - I*ArcSinh[a*x])/(2*a^2)`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 71 vs. 2(34) = 68.

time = 0.06, size = 72, normalized size = 1.71

method	result	size
risch	$\frac{i(ax-2i)\sqrt{a^2x^2+1}}{2a^2} - \frac{i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a\sqrt{a^2}}$	59
default	$ia \left(\frac{x\sqrt{a^2x^2+1}}{2a^2} - \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a^2\sqrt{a^2}} \right) + \frac{\sqrt{a^2x^2+1}}{a^2}$	72
meijerg	$\frac{-2\sqrt{\pi} + 2\sqrt{\pi} \sqrt{a^2x^2+1}}{2a^2\sqrt{\pi}} + i \left(\frac{\sqrt{\pi} x (a^2)^{\frac{3}{2}} \sqrt{a^2x^2+1}}{a^2} - \frac{\sqrt{\pi} (a^2)^{\frac{3}{2}} \operatorname{arcsinh}(ax)}{a^3} \right)$	88

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2)*x,x,method=_RETURNVERBOSE)``[Out] I*a*(1/2*x*(a^2*x^2+1)^(1/2)/a^2-1/2/a^2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)/a^2`

Maxima [A]

time = 0.26, size = 42, normalized size = 1.00

$$\frac{i \sqrt{a^2 x^2 + 1} x}{2a} - \frac{i \operatorname{arsinh}(ax)}{2a^2} + \frac{\sqrt{a^2 x^2 + 1}}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x,x, algorithm="maxima")``[Out] 1/2*I*sqrt(a^2*x^2 + 1)*x/a - 1/2*I*arcsinh(a*x)/a^2 + sqrt(a^2*x^2 + 1)/a^2`**Fricas [A]**

time = 1.34, size = 43, normalized size = 1.02

$$\frac{\sqrt{a^2 x^2 + 1} (i a x + 2) + i \log(-a x + \sqrt{a^2 x^2 + 1})}{2 a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x,x, algorithm="fricas")``[Out] 1/2*(sqrt(a^2*x^2 + 1)*(I*a*x + 2) + I*log(-a*x + sqrt(a^2*x^2 + 1)))/a^2`**Sympy [A]**

time = 2.06, size = 51, normalized size = 1.21

$$\begin{cases} \frac{x^2}{2} & \text{for } a^2 = 0 \\ \frac{\sqrt{a^2 x^2 + 1}}{a^2} & \text{otherwise} \end{cases} + \frac{i x \sqrt{a^2 x^2 + 1}}{2a} - \frac{i \operatorname{asinh}(ax)}{2a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)*x,x)``[Out] Piecewise((x**2/2, Eq(a**2, 0)), (sqrt(a**2*x**2 + 1)/a**2, True)) + I*x*sqrt(a**2*x**2 + 1)/(2*a) - I*asinh(a*x)/(2*a**2)`**Giac [A]**

time = 0.42, size = 53, normalized size = 1.26

$$-\frac{1}{2} \sqrt{a^2 x^2 + 1} \left(-\frac{i x}{a} - \frac{2}{a^2} \right) + \frac{i \log(-x|a| + \sqrt{a^2 x^2 + 1})}{2 a |a|}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x,x, algorithm="giac")`

[Out] $-1/2*\sqrt{a^2*x^2 + 1}*(-I*x/a - 2/a^2) + 1/2*I*\log(-x*abs(a) + \sqrt{a^2*x^2 + 1})/(a*abs(a))$

Mupad [B]

time = 0.04, size = 51, normalized size = 1.21

$$\frac{\left(\frac{1}{\sqrt{a^2}} + \frac{x\sqrt{a^2} \operatorname{li}}{2a}\right) \sqrt{a^2 x^2 + 1} - \frac{\operatorname{asinh}\left(x\sqrt{a^2}\right) \operatorname{li}}{2a}}{\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{int}\left(\frac{x*(a*x*\operatorname{li} + 1)}{(a^2*x^2 + 1)^{(1/2)}, x}\right)$

[Out] $\left(\frac{1}{(a^2)^{(1/2)} + (x*(a^2)^{(1/2)*\operatorname{li}})/(2*a)}\right)*(a^2*x^2 + 1)^{(1/2)} - (\operatorname{asinh}(x*(a^2)^{(1/2))*\operatorname{li}})/(2*a))/(a^2)^{(1/2)}$

3.5 $\int e^{i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=29

$$\frac{i\sqrt{1+a^2x^2}}{a} + \frac{\sinh^{-1}(ax)}{a}$$

[Out] arcsinh(a*x)/a+I*(a^2*x^2+1)^(1/2)/a

Rubi [A]

time = 0.01, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5167, 655, 221}

$$\frac{\sinh^{-1}(ax)}{a} + \frac{i\sqrt{a^2x^2+1}}{a}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x]),x]

[Out] (I*Sqrt[1 + a^2*x^2])/a + ArcSinh[a*x]/a

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 655

Int[((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] :> Simp[e*((a + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Dist[d, Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]

Rule 5167

Int[E^(ArcTan[(a_)*(x_)])*(n_), x_Symbol] :> Int[(1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[a, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{i \tan^{-1}(ax)} dx &= \int \frac{1 + iax}{\sqrt{1 + a^2x^2}} dx \\ &= \frac{i\sqrt{1 + a^2x^2}}{a} + \int \frac{1}{\sqrt{1 + a^2x^2}} dx \\ &= \frac{i\sqrt{1 + a^2x^2}}{a} + \frac{\sinh^{-1}(ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 26, normalized size = 0.90

$$\frac{i\sqrt{1+a^2x^2} + \sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*ArcTan[a*x]),x]``[Out] (I*Sqrt[1 + a^2*x^2] + ArcSinh[a*x])/a`**Maple [A]**

time = 0.08, size = 48, normalized size = 1.66

method	result	size
meijerg	$\frac{\operatorname{arcsinh}(ax)}{a} + \frac{i(-2\sqrt{\pi} + 2\sqrt{\pi}\sqrt{a^2x^2+1})}{2a\sqrt{\pi}}$	41
default	$\frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{\sqrt{a^2}} + \frac{i\sqrt{a^2x^2+1}}{a}$	48
risch	$\frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{\sqrt{a^2}} + \frac{i\sqrt{a^2x^2+1}}{a}$	48

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)``[Out] ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2)+I*(a^2*x^2+1)^(1/2)/a`**Maxima [A]**

time = 0.25, size = 25, normalized size = 0.86

$$\frac{\operatorname{arsinh}(ax)}{a} + \frac{i\sqrt{a^2x^2+1}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2),x, algorithm="maxima")``[Out] arcsinh(a*x)/a + I*sqrt(a^2*x^2 + 1)/a`**Fricas [A]**

time = 1.61, size = 37, normalized size = 1.28

$$\frac{i\sqrt{a^2x^2+1} - \log\left(-ax + \sqrt{a^2x^2+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2),x, algorithm="fricas")

[Out] (I*sqrt(a^2*x^2 + 1) - log(-a*x + sqrt(a^2*x^2 + 1)))/a

Sympy [A]

time = 0.74, size = 68, normalized size = 2.34

$$ia \left(\begin{cases} \frac{x^2}{2} & \text{for } a^2 = 0 \\ \frac{\sqrt{a^2 x^2 + 1}}{a^2} & \text{otherwise} \end{cases} \right) + \begin{cases} \sqrt{-\frac{1}{a^2}} \operatorname{asin}(x\sqrt{-a^2}) & \text{for } a^2 < 0 \\ \sqrt{\frac{1}{a^2}} \operatorname{asinh}(x\sqrt{a^2}) & \text{for } a^2 > 0 \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2),x)

[Out] I*a*Piecewise((x**2/2, Eq(a**2, 0)), (sqrt(a**2*x**2 + 1)/a**2, True)) + Piecewise((sqrt(-1/a**2)*asin(x*sqrt(-a**2)), a**2 < 0), (sqrt(a**(-2))*asinh(x*sqrt(a**2)), a**2 > 0))

Giac [A]

time = 0.42, size = 41, normalized size = 1.41

$$-\frac{\log(-x|a| + \sqrt{a^2 x^2 + 1})}{|a|} + \frac{i \sqrt{a^2 x^2 + 1}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] -log(-x*abs(a) + sqrt(a^2*x^2 + 1))/abs(a) + I*sqrt(a^2*x^2 + 1)/a

Mupad [B]

time = 0.04, size = 32, normalized size = 1.10

$$\frac{\operatorname{asinh}(x\sqrt{a^2})}{\sqrt{a^2}} + \frac{\sqrt{a^2 x^2 + 1} \operatorname{li}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2),x)

[Out] ((a^2*x^2 + 1)^(1/2)*1i)/a + asinh(x*(a^2)^(1/2))/(a^2)^(1/2)

3.6 $\int \frac{e^{i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=25

$$i \sinh^{-1}(ax) - \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] I*arcsinh(a*x)-arctanh((a^2*x^2+1)^(1/2))

Rubi [A]

time = 0.03, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 858, 221, 272, 65, 214}

$$- \tanh^{-1}\left(\sqrt{a^2x^2+1}\right) + i \sinh^{-1}(ax)$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])/x,x]

[Out] I*ArcSinh[a*x] - ArcTanh[Sqrt[1 + a^2*x^2]]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 858

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] + D
ist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d,
e, f, g, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{i \tan^{-1}(ax)}}{x} dx &= \int \frac{1 + iax}{x\sqrt{1 + a^2x^2}} dx \\
&= (ia) \int \frac{1}{\sqrt{1 + a^2x^2}} dx + \int \frac{1}{x\sqrt{1 + a^2x^2}} dx \\
&= i \sinh^{-1}(ax) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
&= i \sinh^{-1}(ax) + \frac{\text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + x^2} dx, x, \sqrt{1 + a^2x^2} \right)}{a^2} \\
&= i \sinh^{-1}(ax) - \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 29, normalized size = 1.16

$$i \sinh^{-1}(ax) + \log(x) - \log \left(1 + \sqrt{1 + a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a*x])/x,x]

[Out] I*ArcSinh[a*x] + Log[x] - Log[1 + Sqrt[1 + a^2*x^2]]

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 47 vs. 2(22) = 44.

time = 0.07, size = 48, normalized size = 1.92

method	result	size
--------	--------	------

default	$\frac{ia \ln\left(\frac{a^2 x}{\sqrt{a^2}} + \sqrt{a^2 x^2 + 1}\right)}{\sqrt{a^2}} - \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2 x^2 + 1}}\right)$	48
meijerg	$\frac{-2\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2 x^2 + 1}}{2}\right) + (-2\ln(2) + 2\ln(x) + \ln(a^2))\sqrt{\pi}}{2\sqrt{\pi}} + i \operatorname{arsinh}(ax)$	53

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)/(a^2*x^2+1)^(1/2)/x,x,method=_RETURNVERBOSE)`

[Out] `I*a*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2))`

Maxima [A]

time = 0.25, size = 18, normalized size = 0.72

$$i \operatorname{arsinh}(ax) - \operatorname{arsinh}\left(\frac{1}{a|x|}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x,x, algorithm="maxima")`

[Out] `I*arcsinh(a*x) - arcsinh(1/(a*abs(x)))`

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(21) = 42$.

time = 1.35, size = 58, normalized size = 2.32

$$-\log\left(-ax + \sqrt{a^2 x^2 + 1} + 1\right) - i \log\left(-ax + \sqrt{a^2 x^2 + 1}\right) + \log\left(-ax + \sqrt{a^2 x^2 + 1} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x,x, algorithm="fricas")`

[Out] `-log(-a*x + sqrt(a^2*x^2 + 1) + 1) - I*log(-a*x + sqrt(a^2*x^2 + 1)) + log(-a*x + sqrt(a^2*x^2 + 1) - 1)`

Sympy [A]

time = 2.09, size = 53, normalized size = 2.12

$$ia \left(\begin{cases} \sqrt{-\frac{1}{a^2}} \operatorname{asin}(x\sqrt{-a^2}) & \text{for } a^2 < 0 \\ \sqrt{\frac{1}{a^2}} \operatorname{arsinh}(x\sqrt{a^2}) & \text{for } a^2 > 0 \end{cases} \right) - \operatorname{arsinh}\left(\frac{1}{ax}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/x,x)

[Out] I*a*Piecewise((sqrt(-1/a**2)*asin(x*sqrt(-a**2)), a**2 < 0), (sqrt(a**(-2))*asinh(x*sqrt(a**2)), a**2 > 0)) - asinh(1/(a*x))

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(21) = 42$.
time = 0.43, size = 68, normalized size = 2.72

$$-\frac{ia \log\left(-x|a| + \sqrt{a^2 x^2 + 1}\right)}{|a|} - \log\left(\left|-x|a| + \sqrt{a^2 x^2 + 1} + 1\right|\right) + \log\left(\left|-x|a| + \sqrt{a^2 x^2 + 1} - 1\right|\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x,x, algorithm="giac")

[Out] -I*a*log(-x*abs(a) + sqrt(a^2*x^2 + 1))/abs(a) - log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) + 1)) + log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) - 1))

Mupad [B]

time = 0.04, size = 32, normalized size = 1.28

$$-\operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right) + \frac{a \operatorname{asinh}\left(x \sqrt{a^2}\right) \operatorname{li}}{\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)/(x*(a^2*x^2 + 1)^(1/2)),x)

[Out] (a*asinh(x*(a^2)^(1/2))*1i)/(a^2)^(1/2) - atanh((a^2*x^2 + 1)^(1/2))

3.7 $\int \frac{e^{i \operatorname{ArcTan}(ax)}}{x^2} dx$

Optimal. Leaf size=38

$$-\frac{\sqrt{1+a^2x^2}}{x} - ia \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-I*a*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-(a^2*x^2+1)^{(1/2)}/x$

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5168, 821, 272, 65, 214}

$$-\frac{\sqrt{a^2x^2+1}}{x} - ia \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*\operatorname{ArcTan}[a*x])}/x^2, x]$

[Out] $-(\operatorname{Sqrt}[1+a^2*x^2]/x) - I*a*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 272

$\operatorname{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.)})^{(p_.)}, x_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n] - 1)*(a + b*x)^p}], x, x^n], x] /; \operatorname{FreeQ}[\{a, b, m, n, p\}, x] \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]]$

Rule 821

$\operatorname{Int}[(d_. + (e_.)*(x_.))^{(m_.)}*((f_.) + (g_.)*(x_.))*((a_.) + (c_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(-e*f - d*g)*(d + e*x)^{(m+1)}*((a + c*x^2)^{(p+1)})/(2*(p+1)*(c*d^2 + a*e^2)), x] + \operatorname{Dist}[(c*d*f + a*e*g)/(c*d^2 + a*e^2), \operatorname{Int}[(d + e*x)^{(m+1)}*(a + c*x^2)^p, x], x] /; \operatorname{FreeQ}[\{a, c, d, e, f, g, m,$

p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 5168

Int[E^(ArcTan[a_.]*(x_.))*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*n + 1)/2)/((1 + I*a*x)^(I*n - 1)/2)*Sqrt[1 + a^2*x^2]], x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{1 + iax}{x^2 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} + (ia) \int \frac{1}{x \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} + \frac{1}{2}(ia) \text{Subst} \left(\int \frac{1}{x \sqrt{1 + a^2 x}} dx, x, x^2 \right) \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} + \frac{i \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2 x^2} \right)}{a} \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} - ia \tanh^{-1} \left(\sqrt{1 + a^2 x^2} \right)
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 47, normalized size = 1.24

$$-\frac{\sqrt{1 + a^2 x^2}}{x} + ia \log(x) - ia \log \left(1 + \sqrt{1 + a^2 x^2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a*x])/x^2,x]

[Out] -(Sqrt[1 + a^2*x^2]/x) + I*a*Log[x] - I*a*Log[1 + Sqrt[1 + a^2*x^2]]

Maple [A]

time = 0.08, size = 34, normalized size = 0.89

method	result	size
default	$-ia \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2 x^2 + 1}} \right) - \frac{\sqrt{a^2 x^2 + 1}}{x}$	34

risch	$-ia \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right) - \frac{\sqrt{a^2x^2+1}}{x}$	34
meijerg	$-\frac{\sqrt{a^2x^2+1}}{x} + \frac{ia\left(-2\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2}\right) + (-2\ln(2) + 2\ln(x) + \ln(a^2))\sqrt{\pi}\right)}{2\sqrt{\pi}}$	64

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

[Out] `-I*a*arctanh(1/(a^2*x^2+1)^(1/2))-(a^2*x^2+1)^(1/2)/x`

Maxima [A]

time = 0.26, size = 29, normalized size = 0.76

$$-ia \operatorname{arsinh}\left(\frac{1}{a|x|}\right) - \frac{\sqrt{a^2x^2+1}}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^2,x, algorithm="maxima")`

[Out] `-I*a*arcsinh(1/(a*abs(x))) - sqrt(a^2*x^2 + 1)/x`

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 66 vs. $2(32) = 64$.

time = 1.20, size = 66, normalized size = 1.74

$$\frac{-iax \log\left(-ax + \sqrt{a^2x^2+1} + 1\right) + iax \log\left(-ax + \sqrt{a^2x^2+1} - 1\right) - ax - \sqrt{a^2x^2+1}}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^2,x, algorithm="fricas")`

[Out] `(-I*a*x*log(-a*x + sqrt(a^2*x^2 + 1) + 1) + I*a*x*log(-a*x + sqrt(a^2*x^2 + 1) - 1) - a*x - sqrt(a^2*x^2 + 1))/x`

Sympy [A]

time = 1.46, size = 26, normalized size = 0.68

$$-a\sqrt{1 + \frac{1}{a^2x^2}} - ia \operatorname{asinh}\left(\frac{1}{ax}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/x**2,x)`

[Out] `-a*sqrt(1 + 1/(a**2*x**2)) - I*a*asinh(1/(a*x))`

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 75 vs. $2(32) = 64$.

time = 0.42, size = 75, normalized size = 1.97

$$-i a \log \left(\left| -x|a| + \sqrt{a^2 x^2 + 1} + 1 \right| \right) + i a \log \left(\left| -x|a| + \sqrt{a^2 x^2 + 1} - 1 \right| \right) + \frac{2|a|}{\left(|x|a| - \sqrt{a^2 x^2 + 1} \right)^2 - 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^2,x, algorithm="giac")

[Out] -I*a*log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) + 1)) + I*a*log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) - 1)) + 2*abs(a)/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)

Mupad [B]

time = 0.04, size = 33, normalized size = 0.87

$$-\frac{\sqrt{a^2 x^2 + 1}}{x} - a \operatorname{atanh} \left(\sqrt{a^2 x^2 + 1} \right) \operatorname{li}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)/(x^2*(a^2*x^2 + 1)^(1/2)),x)

[Out] - a*atanh((a^2*x^2 + 1)^(1/2))*1i - (a^2*x^2 + 1)^(1/2)/x

3.8 $\int \frac{e^{i\text{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=63

$$-\frac{\sqrt{1+a^2x^2}}{2x^2} - \frac{ia\sqrt{1+a^2x^2}}{x} + \frac{1}{2}a^2 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] 1/2*a^2*arctanh((a^2*x^2+1)^(1/2))-1/2*(a^2*x^2+1)^(1/2)/x^2-I*a*(a^2*x^2+1)^(1/2)/x

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 849, 821, 272, 65, 214}

$$-\frac{ia\sqrt{a^2x^2+1}}{x} - \frac{\sqrt{a^2x^2+1}}{2x^2} + \frac{1}{2}a^2 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])/x^3,x]

[Out] -1/2*Sqrt[1 + a^2*x^2]/x^2 - (I*a*Sqrt[1 + a^2*x^2])/x + (a^2*ArcTanh[Sqrt[1 + a^2*x^2]])/2

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 821

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] :> Simp[(-(e*f - d*g))*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1
```

```
)/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2),
Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,
p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]
```

Rule 849

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/(
(m + 1)*(c*d^2 + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 + a*e^2)), Int[(d +
e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
+ 2*p + 3)*x, x], x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 +
a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*
p])
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{1 + iax}{x^3 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} - \frac{1}{2} \int \frac{-2ia + a^2 x}{x^2 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} - \frac{ia \sqrt{1 + a^2 x^2}}{x} - \frac{1}{2} a^2 \int \frac{1}{x \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} - \frac{ia \sqrt{1 + a^2 x^2}}{x} - \frac{1}{4} a^2 \text{Subst} \left(\int \frac{1}{x \sqrt{1 + a^2 x}} dx, x, x^2 \right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} - \frac{ia \sqrt{1 + a^2 x^2}}{x} - \frac{1}{2} \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2 x^2} \right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} - \frac{ia \sqrt{1 + a^2 x^2}}{x} + \frac{1}{2} a^2 \tanh^{-1} \left(\sqrt{1 + a^2 x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 57, normalized size = 0.90

$$\frac{1}{2} \left(\frac{(-1 - 2iax)\sqrt{1 + a^2 x^2}}{x^2} - a^2 \log(x) + a^2 \log \left(1 + \sqrt{1 + a^2 x^2} \right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a*x])/x^3,x]

[Out] (((-1 - (2*I)*a*x)*Sqrt[1 + a^2*x^2])/x^2 - a^2*Log[x] + a^2*Log[1 + Sqrt[1 + a^2*x^2]])/2

Maple [A]

time = 0.08, size = 53, normalized size = 0.84

method	result
default	$-\frac{\sqrt{a^2x^2+1}}{2x^2} + \frac{a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{2} - \frac{ia\sqrt{a^2x^2+1}}{x}$
risch	$-\frac{i(2a^3x^3-ia^2x^2+2ax-i)}{2x^2\sqrt{a^2x^2+1}} + \frac{a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{2}$
meijerg	$\frac{a^2 \left(\frac{\sqrt{\pi} (4a^2x^2+8)}{8a^2x^2} - \frac{\sqrt{\pi} \sqrt{a^2x^2+1}}{a^2x^2} + \sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2}\right) - \frac{(1-2\ln(2)+2\ln(x)+\ln(a^2))\sqrt{\pi}}{2} - \frac{\sqrt{\pi}}{x^2a^2} \right)}{2\sqrt{\pi}} - \frac{ia\sqrt{a^2x^2+1}}{x}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^3,x,method=_RETURNVERBOSE)

[Out] -1/2*(a^2*x^2+1)^(1/2)/x^2+1/2*a^2*arctanh(1/(a^2*x^2+1)^(1/2))-I*a*(a^2*x^2+1)^(1/2)/x

Maxima [A]

time = 0.28, size = 48, normalized size = 0.76

$$\frac{1}{2} a^2 \operatorname{arsinh}\left(\frac{1}{a|x|}\right) - \frac{i \sqrt{a^2x^2+1} a}{x} - \frac{\sqrt{a^2x^2+1}}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^3,x, algorithm="maxima")

[Out] 1/2*a^2*arcsinh(1/(a*abs(x))) - I*sqrt(a^2*x^2 + 1)*a/x - 1/2*sqrt(a^2*x^2 + 1)/x^2

Fricas [A]

time = 5.37, size = 83, normalized size = 1.32

$$\frac{a^2x^2 \log\left(-ax + \sqrt{a^2x^2+1} + 1\right) - a^2x^2 \log\left(-ax + \sqrt{a^2x^2+1} - 1\right) - 2ia^2x^2 + \sqrt{a^2x^2+1}(-2iax - 1)}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^3,x, algorithm="fricas")

[Out] 1/2*(a^2*x^2*log(-a*x + sqrt(a^2*x^2 + 1) + 1) - a^2*x^2*log(-a*x + sqrt(a^2*x^2 + 1) - 1) - 2*I*a^2*x^2 + sqrt(a^2*x^2 + 1)*(-2*I*a*x - 1))/x^2

Sympy [A]

time = 1.97, size = 48, normalized size = 0.76

$$-ia^2 \sqrt{1 + \frac{1}{a^2 x^2}} + \frac{a^2 \operatorname{asinh}\left(\frac{1}{ax}\right)}{2} - \frac{a \sqrt{1 + \frac{1}{a^2 x^2}}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/x**3,x)

[Out] -I*a**2*sqrt(1 + 1/(a**2*x**2)) + a**2*asinh(1/(a*x))/2 - a*sqrt(1 + 1/(a**2*x**2))/(2*x)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 153 vs. 2(51) = 102.

time = 0.42, size = 153, normalized size = 2.43

$$\frac{1}{2} a^2 \log\left(\left|-x|a| + \sqrt{a^2 x^2 + 1}\right| + 1\right) - \frac{1}{2} a^2 \log\left(\left|-x|a| + \sqrt{a^2 x^2 + 1}\right| - 1\right) + \frac{\left(x|a| - \sqrt{a^2 x^2 + 1}\right)^3 a^2 + 2i\left(x|a| - \sqrt{a^2 x^2 + 1}\right)^2 a|a| + \left(x|a| - \sqrt{a^2 x^2 + 1}\right) a^2 - 2i a|a|}{\left(\left(x|a| - \sqrt{a^2 x^2 + 1}\right)^2 - 1\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^3,x, algorithm="giac")

[Out] 1/2*a^2*log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) + 1)) - 1/2*a^2*log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) - 1)) + ((x*abs(a) - sqrt(a^2*x^2 + 1))^3*a^2 + 2*I*(x*abs(a) - sqrt(a^2*x^2 + 1))^2*a*abs(a) + (x*abs(a) - sqrt(a^2*x^2 + 1))*a^2 - 2*I*a*abs(a))/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)^2

Mupad [B]

time = 0.04, size = 52, normalized size = 0.83

$$\frac{a^2 \operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right)}{2} - \frac{\sqrt{a^2 x^2 + 1}}{2 x^2} - \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)/(x^3*(a^2*x^2 + 1)^(1/2)),x)

[Out] (a^2*atanh((a^2*x^2 + 1)^(1/2)))/2 - (a^2*x^2 + 1)^(1/2)/(2*x^2) - (a*(a^2*x^2 + 1)^(1/2)*1i)/x

3.9 $\int \frac{e^{i \operatorname{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=90

$$-\frac{\sqrt{1+a^2x^2}}{3x^3} - \frac{ia\sqrt{1+a^2x^2}}{2x^2} + \frac{2a^2\sqrt{1+a^2x^2}}{3x} + \frac{1}{2}ia^3 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $1/2*I*a^3*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/3*(a^2*x^2+1)^{(1/2)}/x^3-1/2*I*a*(a^2*x^2+1)^{(1/2)}/x^2+2/3*a^2*(a^2*x^2+1)^{(1/2)}/x$

Rubi [A]

time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 849, 821, 272, 65, 214}

$$\frac{2a^2\sqrt{a^2x^2+1}}{3x} - \frac{ia\sqrt{a^2x^2+1}}{2x^2} - \frac{\sqrt{a^2x^2+1}}{3x^3} + \frac{1}{2}ia^3 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*\operatorname{ArcTan}[a*x])}/x^4, x]$

[Out] $-1/3*\operatorname{Sqrt}[1+a^2*x^2]/x^3 - ((I/2)*a*\operatorname{Sqrt}[1+a^2*x^2])/x^2 + (2*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(3*x) + (I/2)*a^3*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 272

$\operatorname{Int}[(x_.)^{(m_.)}*((a_.) + (b_.)*(x_.)^{(n_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n] - 1)*(a + b*x)^p}], x, x^n], x] /; \operatorname{FreeQ}\{a, b, m, n, p\}, x] \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]]$

Rule 821

$\operatorname{Int}[(d_. + (e_.)*(x_.))^{(m_.)}*((f_.) + (g_.)*(x_.))*((a_.) + (c_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(-e*f - d*g)*(d + e*x)^{(m+1)}*((a + c*x^2)^{(p+1)}$

)/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2),
 Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,
 p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 849

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_.) + (c_.)*(x_)^2)^(p
 _.), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/(
 (m + 1)*(c*d^2 + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 + a*e^2)), Int[(d +
 e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
 + 2*p + 3)*x, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 +
 a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*
 p])

Rule 5168

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
 x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
 Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{1 + iax}{x^4 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{1}{3} \int \frac{-3ia + 2a^2 x}{x^3 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{1}{6} \int \frac{-4a^2 - 3ia^3 x}{x^2 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} - \frac{1}{2} (ia^3) \int \frac{1}{x \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} - \frac{1}{4} (ia^3) \text{Subst} \left(\int \frac{1}{x \sqrt{1 + a^2 x}} dx, x, \sqrt{1 + a^2 x^2} \right) \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} - \frac{1}{2} (ia) \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2 x^2} \right) \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} + \frac{1}{2} ia^3 \tanh^{-1} \left(\sqrt{1 + a^2 x^2} \right)
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 70, normalized size = 0.78

$$\frac{1}{6} \left(\frac{\sqrt{1+a^2x^2}(-2-3iax+4a^2x^2)}{x^3} - 3ia^3 \log(x) + 3ia^3 \log\left(1 + \sqrt{1+a^2x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a*x])/x^4,x]

[Out] ((Sqrt[1 + a^2*x^2]*(-2 - (3*I)*a*x + 4*a^2*x^2))/x^3 - (3*I)*a^3*Log[x] + (3*I)*a^3*Log[1 + Sqrt[1 + a^2*x^2]])/6

Maple [A]

time = 0.09, size = 75, normalized size = 0.83

method	result
risch	$\frac{4a^4x^4 - 3ia^3x^3 + 2a^2x^2 - 3iax - 2}{6x^3\sqrt{a^2x^2 + 1}} + \frac{ia^3 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2 + 1}}\right)}{2}$
default	$-\frac{\sqrt{a^2x^2 + 1}}{3x^3} + \frac{2a^2\sqrt{a^2x^2 + 1}}{3x} + ia \left(-\frac{\sqrt{a^2x^2 + 1}}{2x^2} + \frac{a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2 + 1}}\right)}{2} \right)$
meijerg	$-\frac{(-2a^2x^2+1)\sqrt{a^2x^2+1}}{3x^3} + \frac{ia^3 \left(\frac{\sqrt{\pi} (4a^2x^2+8)}{8a^2x^2} - \frac{\sqrt{\pi} \sqrt{a^2x^2+1}}{a^2x^2} + \sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2}\right) \right)}{2\sqrt{\pi}} - \frac{(1-2\ln(2)+2\ln)}{2\sqrt{\pi}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^4,x,method=_RETURNVERBOSE)

[Out] -1/3*(a^2*x^2+1)^(1/2)/x^3+2/3*a^2*(a^2*x^2+1)^(1/2)/x+I*a*(-1/2*(a^2*x^2+1)^(1/2)/x^2+1/2*a^2*arctanh(1/(a^2*x^2+1)^(1/2)))

Maxima [A]

time = 0.25, size = 67, normalized size = 0.74

$$\frac{1}{2}ia^3 \operatorname{arsinh}\left(\frac{1}{a|x|}\right) + \frac{2\sqrt{a^2x^2+1}a^2}{3x} - \frac{i\sqrt{a^2x^2+1}a}{2x^2} - \frac{\sqrt{a^2x^2+1}}{3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^4,x, algorithm="maxima")

[Out] 1/2*I*a^3*arcsinh(1/(a*abs(x))) + 2/3*sqrt(a^2*x^2 + 1)*a^2/x - 1/2*I*sqrt(a^2*x^2 + 1)*a/x^2 - 1/3*sqrt(a^2*x^2 + 1)/x^3

Fricas [A]

time = 4.20, size = 92, normalized size = 1.02

$$\frac{3i a^3 x^3 \log(-ax + \sqrt{a^2 x^2 + 1}) + 1 - 3i a^3 x^3 \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + 4 a^3 x^3 + (4 a^2 x^2 - 3i a x - 2) \sqrt{a^2 x^2 + 1}}{6 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^4,x, algorithm="fricas")

[Out] 1/6*(3*I*a^3*x^3*log(-a*x + sqrt(a^2*x^2 + 1) + 1) - 3*I*a^3*x^3*log(-a*x + sqrt(a^2*x^2 + 1) - 1) + 4*a^3*x^3 + (4*a^2*x^2 - 3*I*a*x - 2)*sqrt(a^2*x^2 + 1))/x^3

Sympy [A]

time = 2.10, size = 75, normalized size = 0.83

$$\frac{2a^3 \sqrt{1 + \frac{1}{a^2 x^2}}}{3} + \frac{ia^3 \operatorname{asinh}\left(\frac{1}{ax}\right)}{2} - \frac{ia^2 \sqrt{1 + \frac{1}{a^2 x^2}}}{2x} - \frac{a \sqrt{1 + \frac{1}{a^2 x^2}}}{3x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/x**4,x)

[Out] 2*a**3*sqrt(1 + 1/(a**2*x**2))/3 + I*a**3*asinh(1/(a*x))/2 - I*a**2*sqrt(1 + 1/(a**2*x**2))/(2*x) - a*sqrt(1 + 1/(a**2*x**2))/(3*x**2)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 161 vs. 2(70) = 140.

time = 0.41, size = 161, normalized size = 1.79

$$\frac{\frac{1}{2} i a^3 \log(|-x|a + \sqrt{a^2 x^2 + 1}) + 1 - \frac{1}{2} i a^3 \log(|-x|a + \sqrt{a^2 x^2 + 1} - 1) - \frac{-3i(x|a - \sqrt{a^2 x^2 + 1})^5 a^3 - 12(x|a - \sqrt{a^2 x^2 + 1})^2 a^2 |a| + 3(i|x|a - i\sqrt{a^2 x^2 + 1}) a^3 + 4 a^2 |a|}{3((x|a - \sqrt{a^2 x^2 + 1})^2 - 1)^3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^4,x, algorithm="giac")

[Out] 1/2*I*a^3*log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) + 1)) - 1/2*I*a^3*log(abs(-x*abs(a) + sqrt(a^2*x^2 + 1) - 1)) - 1/3*(-3*I*(x*abs(a) - sqrt(a^2*x^2 + 1))^5*a^3 - 12*(x*abs(a) - sqrt(a^2*x^2 + 1))^2*a^2*abs(a) + 3*(I*x*abs(a) - I*sqrt(a^2*x^2 + 1))*a^3 + 4*a^2*abs(a))/((x*abs(a) - sqrt(a^2*x^2 + 1))^2 - 1)^3

Mupad [B]

time = 0.04, size = 74, normalized size = 0.82

$$\frac{a^3 \operatorname{atan}\left(\sqrt{a^2 x^2 + 1} \operatorname{li}\right)}{2} - \frac{\sqrt{a^2 x^2 + 1}}{3 x^3} + \frac{2 a^2 \sqrt{a^2 x^2 + 1}}{3 x} - \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{2 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*x*1i + 1)/(x^4*(a^2*x^2 + 1)^(1/2)),x)
```

```
[Out] (a^3*atan((a^2*x^2 + 1)^(1/2)*1i))/2 - (a^2*x^2 + 1)^(1/2)/(3*x^3) - (a*(a^2*x^2 + 1)^(1/2)*1i)/(2*x^2) + (2*a^2*(a^2*x^2 + 1)^(1/2))/(3*x)
```

3.10 $\int \frac{e^{i \operatorname{ArcTan}(ax)}}{x^5} dx$

Optimal. Leaf size=113

$$-\frac{\sqrt{1+a^2x^2}}{4x^4} - \frac{ia\sqrt{1+a^2x^2}}{3x^3} + \frac{3a^2\sqrt{1+a^2x^2}}{8x^2} + \frac{2ia^3\sqrt{1+a^2x^2}}{3x} - \frac{3}{8}a^4 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-3/8*a^4*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/4*(a^2*x^2+1)^{(1/2)}/x^4-1/3*I*a*(a^2*x^2+1)^{(1/2)}/x^3+3/8*a^2*(a^2*x^2+1)^{(1/2)}/x^2+2/3*I*a^3*(a^2*x^2+1)^{(1/2)}/x$

Rubi [A]

time = 0.06, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 849, 821, 272, 65, 214}

$$\frac{3a^2\sqrt{a^2x^2+1}}{8x^2} - \frac{\sqrt{a^2x^2+1}}{4x^4} - \frac{ia\sqrt{a^2x^2+1}}{3x^3} - \frac{3}{8}a^4 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right) + \frac{2ia^3\sqrt{a^2x^2+1}}{3x}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*\operatorname{ArcTan}[a*x])}/x^5, x]$

[Out] $-1/4*\operatorname{Sqrt}[1+a^2*x^2]/x^4 - ((I/3)*a*\operatorname{Sqrt}[1+a^2*x^2])/x^3 + (3*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(8*x^2) + (((2*I)/3)*a^3*\operatorname{Sqrt}[1+a^2*x^2])/x - (3*a^4*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]])/8$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^{n, x}], x, (a+b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 272

$\operatorname{Int}[(x_.)^{(m_.)*((a_.) + (b_.)*(x_.)^{(n_.))^{(p_.)}, x_Symbol] := \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n]-1)*(a+b*x)^p}, x], x, x^n], x] /; \operatorname{FreeQ}[\{a, b, m, n, p\}, x] \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]]$

Rule 821

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(-e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)
)/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2),
Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,
p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]
```

Rule 849

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/(
(m + 1)*(c*d^2 + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 + a*e^2)), Int[(d +
e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
+ 2*p + 3)*x, x], x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 +
a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*
p])
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{1 + iax}{x^5 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{1}{4} \int \frac{-4ia + 3a^2 x}{x^4 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{1}{12} \int \frac{-9a^2 - 8ia^3 x}{x^3 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} - \frac{1}{24} \int \frac{16ia^3 - 9a^4 x}{x^2 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} + \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} + \frac{1}{8}(3a^4) \int \frac{1}{x\sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} + \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} + \frac{1}{16}(3a^4) \text{Subst}\left(\int \frac{1}{x\sqrt{1 + a^2 x^2}} dx\right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} + \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} + \frac{1}{8}(3a^2) \text{Subst}\left(\int \frac{1}{x\sqrt{1 + a^2 x^2}} dx\right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} + \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} - \frac{3}{8}a^4 \tanh^{-1}\left(\sqrt{1 + a^2 x^2}\right)
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 76, normalized size = 0.67

$$\frac{1}{24} \left(\frac{\sqrt{1+a^2x^2}(-6-8iax+9a^2x^2+16ia^3x^3)}{x^4} + 9a^4 \log(x) - 9a^4 \log\left(1 + \sqrt{1+a^2x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a*x])/x^5,x]

[Out] ((Sqrt[1 + a^2*x^2]*(-6 - (8*I)*a*x + 9*a^2*x^2 + (16*I)*a^3*x^3))/x^4 + 9*a^4*Log[x] - 9*a^4*Log[1 + Sqrt[1 + a^2*x^2]])/24

Maple [A]

time = 0.08, size = 97, normalized size = 0.86

method	result
risch	$\frac{i(16a^5x^5-9ia^4x^4+8a^3x^3-3ia^2x^2-8ax+6i)}{24x^4\sqrt{a^2x^2+1}} - \frac{3a^4 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{8}$
default	$ia \left(-\frac{\sqrt{a^2x^2+1}}{3x^3} + \frac{2a^2\sqrt{a^2x^2+1}}{3x} \right) - \frac{\sqrt{a^2x^2+1}}{4x^4} - \frac{3a^2 \left(-\frac{\sqrt{a^2x^2+1}}{2x^2} + \frac{a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{2} \right)}{4}$
meijerg	$a^4 \left(\frac{\sqrt{\pi}(-7a^4x^4-8a^2x^2+8)}{16a^4x^4} - \frac{\sqrt{\pi}(-12a^2x^2+8)\sqrt{a^2x^2+1}}{16a^4x^4} - \frac{3\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2}\right)}{4} \right) + \frac{3\left(\frac{7}{8}-2\ln(2)+2\ln(x)+\ln(a^2)\right)}{8}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^5,x,method=_RETURNVERBOSE)

[Out] I*a*(-1/3*(a^2*x^2+1)^(1/2)/x^3+2/3*a^2*(a^2*x^2+1)^(1/2)/x)-1/4*(a^2*x^2+1)^(1/2)/x^4-3/4*a^2*(-1/2*(a^2*x^2+1)^(1/2)/x^2+1/2*a^2*arctanh(1/(a^2*x^2+1)^(1/2)))

Maxima [A]

time = 0.25, size = 86, normalized size = 0.76

$$-\frac{3}{8} a^4 \operatorname{arsinh}\left(\frac{1}{a|x|}\right) + \frac{2i\sqrt{a^2x^2+1}a^3}{3x} + \frac{3\sqrt{a^2x^2+1}a^2}{8x^2} - \frac{i\sqrt{a^2x^2+1}a}{3x^3} - \frac{\sqrt{a^2x^2+1}}{4x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^5,x, algorithm="maxima")

[Out] $-3/8*a^4*\operatorname{arcsinh}(1/(a*\operatorname{abs}(x))) + 2/3*I*\sqrt{a^2*x^2 + 1}*a^3/x + 3/8*\sqrt{a^2*x^2 + 1}*a^2/x^2 - 1/3*I*\sqrt{a^2*x^2 + 1}*a/x^3 - 1/4*\sqrt{a^2*x^2 + 1}/x^4$

Fricas [A]

time = 4.92, size = 101, normalized size = 0.89

$$\frac{9a^4x^4 \log(-ax + \sqrt{a^2x^2 + 1} + 1) - 9a^4x^4 \log(-ax + \sqrt{a^2x^2 + 1} - 1) - 16ia^4x^4 - (16ia^3x^3 + 9a^2x^2 - 8iax - 6)\sqrt{a^2x^2 + 1}}{24x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^5,x, algorithm="fricas")`

[Out] $-1/24*(9*a^4*x^4*\log(-a*x + \sqrt{a^2*x^2 + 1}) + 1) - 9*a^4*x^4*\log(-a*x + \sqrt{a^2*x^2 + 1} - 1) - 16*I*a^4*x^4 - (16*I*a^3*x^3 + 9*a^2*x^2 - 8*I*a*x - 6)*\sqrt{a^2*x^2 + 1})/x^4$

Sympy [A]

time = 3.50, size = 122, normalized size = 1.08

$$\frac{2ia^4\sqrt{1+\frac{1}{a^2x^2}}}{3} - \frac{3a^4\operatorname{asinh}\left(\frac{1}{ax}\right)}{8} + \frac{3a^3}{8x\sqrt{1+\frac{1}{a^2x^2}}} - \frac{ia^2\sqrt{1+\frac{1}{a^2x^2}}}{3x^2} + \frac{a}{8x^3\sqrt{1+\frac{1}{a^2x^2}}} - \frac{1}{4ax^5\sqrt{1+\frac{1}{a^2x^2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/x**5,x)`

[Out] $2*I*a**4*\sqrt{1 + 1/(a**2*x**2)}/3 - 3*a**4*\operatorname{asinh}(1/(a*x))/8 + 3*a**3/(8*x*\sqrt{1 + 1/(a**2*x**2)}) - I*a**2*\sqrt{1 + 1/(a**2*x**2)}/(3*x**2) + a/(8*x**3*\sqrt{1 + 1/(a**2*x**2)}) - 1/(4*a*x**5*\sqrt{1 + 1/(a**2*x**2)})$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(89) = 178.

time = 0.43, size = 237, normalized size = 2.10

$$\frac{-\frac{3}{8}a^4\log(-|x|a + \sqrt{a^2x^2 + 1} + 1) + \frac{3}{8}a^4\log(-|x|a + \sqrt{a^2x^2 + 1} - 1) - \frac{9(|x|a - \sqrt{a^2x^2 + 1})^7a^4 - 33(|x|a - \sqrt{a^2x^2 + 1})^5a^4 - 48i(|x|a - \sqrt{a^2x^2 + 1})^4a^3 - 33(|x|a - \sqrt{a^2x^2 + 1})^3a^4 + 64i(|x|a - \sqrt{a^2x^2 + 1})^2a^3 + 9(|x|a - \sqrt{a^2x^2 + 1})a^4 - 16ia^3}{12((|x|a - \sqrt{a^2x^2 + 1})^2 - 1)^4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/x^5,x, algorithm="giac")`

[Out] $-3/8*a^4*\log(\operatorname{abs}(-x*\operatorname{abs}(a) + \sqrt{a^2*x^2 + 1}) + 1) + 3/8*a^4*\log(\operatorname{abs}(-x*\operatorname{abs}(a) + \sqrt{a^2*x^2 + 1} - 1)) - 1/12*(9*(x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))^7*a^4 - 33*(x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))^5*a^4 - 48*I*(x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))^4*a^3*\operatorname{abs}(a) - 33*(x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))^3*a^4 + 64*I*(x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))^2*a^3*\operatorname{abs}(a) + 9*(x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))*a^4 - 16*I*a^3*\operatorname{abs}(a))/((x*\operatorname{abs}(a) - \sqrt{a^2*x^2 + 1}))^2 - 1)^4$

Mupad [B]

time = 0.03, size = 95, normalized size = 0.84

$$\frac{a^4 \operatorname{atan}\left(\sqrt{a^2 x^2 + 1} \operatorname{li}\right) 3i}{8} - \frac{\sqrt{a^2 x^2 + 1}}{4 x^4} - \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{3 x^3} + \frac{3 a^2 \sqrt{a^2 x^2 + 1}}{8 x^2} + \frac{a^3 \sqrt{a^2 x^2 + 1} 2i}{3 x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*x*1i + 1)/(x^5*(a^2*x^2 + 1)^(1/2)),x)`

[Out] `(a^4*atan((a^2*x^2 + 1)^(1/2)*1i)*3i)/8 - (a^2*x^2 + 1)^(1/2)/(4*x^4) - (a*(a^2*x^2 + 1)^(1/2)*1i)/(3*x^3) + (3*a^2*(a^2*x^2 + 1)^(1/2))/(8*x^2) + (a^3*(a^2*x^2 + 1)^(1/2)*2i)/(3*x)`

3.11 $\int e^{2i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=48

$$-\frac{2ix}{a^3} + \frac{x^2}{a^2} + \frac{2ix^3}{3a} - \frac{x^4}{4} - \frac{2\log(i+ax)}{a^4}$$

[Out] $-2*I*x/a^3+x^2/a^2+2/3*I*x^3/a-1/4*x^4-2*\ln(I+a*x)/a^4$

Rubi [A]

time = 0.03, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-\frac{2\log(ax+i)}{a^4} - \frac{2ix}{a^3} + \frac{x^2}{a^2} + \frac{2ix^3}{3a} - \frac{x^4}{4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}*x^3, x]$

[Out] $((-2*I)*x)/a^3 + x^2/a^2 + (((2*I)/3)*x^3)/a - x^4/4 - (2*\text{Log}[I + a*x])/a^4$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2)})/(1 + I*a*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1+iax)}{1-iax} dx \\ &= \int \left(-\frac{2i}{a^3} + \frac{2x}{a^2} + \frac{2ix^2}{a} - x^3 - \frac{2}{a^3(i+ax)} \right) dx \\ &= -\frac{2ix}{a^3} + \frac{x^2}{a^2} + \frac{2ix^3}{3a} - \frac{x^4}{4} - \frac{2\log(i+ax)}{a^4} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 48, normalized size = 1.00

$$-\frac{2ix}{a^3} + \frac{x^2}{a^2} + \frac{2ix^3}{3a} - \frac{x^4}{4} - \frac{2\log(i+ax)}{a^4}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a*x])*x^3,x]``[Out] ((-2*I)*x)/a^3 + x^2/a^2 + (((2*I)/3)*x^3)/a - x^4/4 - (2*Log[I + a*x])/a^4`**Maple [A]**

time = 0.10, size = 63, normalized size = 1.31

method	result	size
risch	$-\frac{x^4}{4} + \frac{2ix^3}{3a} + \frac{x^2}{a^2} - \frac{2ix}{a^3} - \frac{\ln(a^2x^2+1)}{a^4} + \frac{2i \arctan(ax)}{a^4}$	55
default	$-\frac{\frac{1}{4}a^3x^4 + \frac{2}{3}ia^2x^3 + ax^2 - 2ix}{a^3} + \frac{-\frac{\ln(a^2x^2+1)}{a} + \frac{2i \arctan(ax)}{a}}{a^3}$	63
meijerg	$\frac{a^2x^2 - \ln(a^2x^2+1)}{2a^4} + \frac{i \left(-\frac{2x(a^2)^{\frac{5}{2}}(-5a^2x^2+15)}{15a^4} + \frac{2(a^2)^{\frac{5}{2}} \arctan(ax)}{a^5} \right)}{a^3 \sqrt{a^2}} - \frac{-\frac{x^2a^2(-3a^2x^2+6)}{6} + \ln(a^2x^2+1)}{2a^4}$	108

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^2/(a^2*x^2+1)*x^3,x,method=_RETURNVERBOSE)``[Out] 1/a^3*(-1/4*a^3*x^4+2/3*I*a^2*x^3+a*x^2-2*I*x)+1/a^3*(-1/a*ln(a^2*x^2+1)+2*I*arctan(a*x)/a)`**Maxima [A]**

time = 0.46, size = 56, normalized size = 1.17

$$-\frac{3a^3x^4 - 8ia^2x^3 - 12ax^2 + 24ix}{12a^3} + \frac{2i \arctan(ax)}{a^4} - \frac{\log(a^2x^2 + 1)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^3,x, algorithm="maxima")``[Out] -1/12*(3*a^3*x^4 - 8*I*a^2*x^3 - 12*a*x^2 + 24*I*x)/a^3 + 2*I*arctan(a*x)/a^4 - log(a^2*x^2 + 1)/a^4`**Fricas [A]**

time = 3.28, size = 46, normalized size = 0.96

$$-\frac{3a^4x^4 - 8ia^3x^3 - 12a^2x^2 + 24iax + 24 \log\left(\frac{ax+i}{a}\right)}{12a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^3,x, algorithm="fricas")

[Out] -1/12*(3*a^4*x^4 - 8*I*a^3*x^3 - 12*a^2*x^2 + 24*I*a*x + 24*log((a*x + I)/a))/a^4

Sympy [A]

time = 0.06, size = 41, normalized size = 0.85

$$-\frac{x^4}{4} + \frac{2ix^3}{3a} + \frac{x^2}{a^2} - \frac{2ix}{a^3} - \frac{2 \log(ax + i)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)*x**3,x)

[Out] -x**4/4 + 2*I*x**3/(3*a) + x**2/a**2 - 2*I*x/a**3 - 2*log(a*x + I)/a**4

Giac [A]

time = 0.42, size = 46, normalized size = 0.96

$$-\frac{3a^4x^4 - 8ia^3x^3 - 12a^2x^2 + 24iax}{12a^4} - \frac{2 \log(ax + i)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^3,x, algorithm="giac")

[Out] -1/12*(3*a^4*x^4 - 8*I*a^3*x^3 - 12*a^2*x^2 + 24*I*a*x)/a^4 - 2*log(a*x + I)/a^4

Mupad [B]

time = 0.42, size = 43, normalized size = 0.90

$$\frac{x^2}{a^2} - \frac{x^4}{4} - \frac{2 \ln\left(x + \frac{1i}{a}\right)}{a^4} - \frac{x 2i}{a^3} + \frac{x^3 2i}{3a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a*x*1i + 1)^2)/(a^2*x^2 + 1),x)

[Out] (x^3*2i)/(3*a) - (x*2i)/a^3 - x^4/4 - (2*log(x + 1i/a))/a^4 + x^2/a^2

3.12 $\int e^{2i \operatorname{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=39

$$\frac{2x}{a^2} + \frac{ix^2}{a} - \frac{x^3}{3} - \frac{2i \log(i + ax)}{a^3}$$

[Out] $2*x/a^2+I*x^2/a-1/3*x^3-2*I*\ln(I+ax)/a^3$

Rubi [A]

time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-\frac{2i \log(ax + i)}{a^3} + \frac{2x}{a^2} + \frac{ix^2}{a} - \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}*x^2,x]$

[Out] $(2*x)/a^2 + (I*x^2)/a - x^3/3 - ((2*I)*\text{Log}[I + a*x])/a^3$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_))*((c_. + (d_.)*(x_))^{(n_.)*((e_. + (f_.)*(x_))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_)]*(n_.))*(x_)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, m, n\}, x\} \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1 + iax)}{1 - iax} dx \\ &= \int \left(\frac{2}{a^2} + \frac{2ix}{a} - x^2 - \frac{2i}{a^2(i + ax)} \right) dx \\ &= \frac{2x}{a^2} + \frac{ix^2}{a} - \frac{x^3}{3} - \frac{2i \log(i + ax)}{a^3} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 39, normalized size = 1.00

$$\frac{2x}{a^2} + \frac{ix^2}{a} - \frac{x^3}{3} - \frac{2i \log(i + ax)}{a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a*x])*x^2,x]

[Out] (2*x)/a^2 + (I*x^2)/a - x^3/3 - ((2*I)*Log[I + a*x])/a^3

Maple [A]

time = 0.08, size = 55, normalized size = 1.41

method	result	size
risch	$\frac{2x}{a^2} - \frac{x^3}{3} + \frac{ix^2}{a} - \frac{i \ln(a^2x^2+1)}{a^3} - \frac{2 \arctan(ax)}{a^3}$	47
default	$\frac{2x - \frac{1}{3}a^2x^3 + ia x^2}{a^2} + \frac{-\frac{i \ln(a^2x^2+1)}{a} - \frac{2 \arctan(ax)}{a}}{a^2}$	55
meijerg	$\frac{2x(a^2)^{\frac{3}{2}}}{a^2} - \frac{2(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3} + \frac{i(a^2x^2 - \ln(a^2x^2+1))}{a^3} - \frac{2x(a^2)^{\frac{5}{2}}(-5a^2x^2+15)}{15a^4} + \frac{2(a^2)^{\frac{5}{2}} \arctan(ax)}{a^5}$	110

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^2/(a^2*x^2+1)*x^2,x,method=_RETURNVERBOSE)

[Out] 1/a^2*(2*x-1/3*a^2*x^3+I*a*x^2)+1/a^2*(-I/a*ln(a^2*x^2+1)-2*arctan(a*x)/a)

Maxima [A]

time = 0.46, size = 47, normalized size = 1.21

$$-\frac{a^2x^3 - 3i ax^2 - 6x}{3a^2} - \frac{2 \arctan(ax)}{a^3} - \frac{i \log(a^2x^2 + 1)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^2,x, algorithm="maxima")

[Out] -1/3*(a^2*x^3 - 3*I*a*x^2 - 6*x)/a^2 - 2*arctan(a*x)/a^3 - I*log(a^2*x^2 + 1)/a^3

Fricas [A]

time = 4.11, size = 37, normalized size = 0.95

$$-\frac{a^3x^3 - 3i a^2x^2 - 6ax + 6i \log\left(\frac{ax+i}{a}\right)}{3a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^2,x, algorithm="fricas")

[Out] -1/3*(a^3*x^3 - 3*I*a^2*x^2 - 6*a*x + 6*I*log((a*x + I)/a))/a^3

Sympy [A]

time = 0.06, size = 31, normalized size = 0.79

$$-\frac{x^3}{3} + \frac{ix^2}{a} + \frac{2x}{a^2} - \frac{2i \log(ax + i)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)*x**2,x)

[Out] -x**3/3 + I*x**2/a + 2*x/a**2 - 2*I*log(a*x + I)/a**3

Giac [A]

time = 0.39, size = 37, normalized size = 0.95

$$-\frac{a^3x^3 - 3ia^2x^2 - 6ax}{3a^3} - \frac{2i \log(ax + i)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^2,x, algorithm="giac")

[Out] -1/3*(a^3*x^3 - 3*I*a^2*x^2 - 6*a*x)/a^3 - 2*I*log(a*x + I)/a^3

Mupad [B]

time = 0.42, size = 36, normalized size = 0.92

$$\frac{2x}{a^2} - \frac{\ln\left(x + \frac{1i}{a}\right) 2i}{a^3} - \frac{x^3}{3} + \frac{x^2 1i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1)^2)/(a^2*x^2 + 1),x)

[Out] (2*x)/a^2 - (log(x + 1i/a)*2i)/a^3 - x^3/3 + (x^2*1i)/a

3.13 $\int e^{2i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=29

$$\frac{2ix}{a} - \frac{x^2}{2} + \frac{2\log(i+ax)}{a^2}$$

[Out] $2*I*x/a - 1/2*x^2 + 2*\ln(I+a*x)/a^2$

Rubi [A]

time = 0.02, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5170, 78}

$$\frac{2\log(ax+i)}{a^2} + \frac{2ix}{a} - \frac{x^2}{2}$$

Antiderivative was successfully verified.

[In] `Int[E^((2*I)*ArcTan[a*x])*x,x]`

[Out] `((2*I)*x)/a - x^2/2 + (2*Log[I + a*x])/a^2`

Rule 78

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0]) || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(ax)} x dx &= \int \frac{x(1+iax)}{1-iax} dx \\ &= \int \left(\frac{2i}{a} - x + \frac{2}{a(i+ax)} \right) dx \\ &= \frac{2ix}{a} - \frac{x^2}{2} + \frac{2\log(i+ax)}{a^2} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 29, normalized size = 1.00

$$\frac{2ix}{a} - \frac{x^2}{2} + \frac{2 \log(i + ax)}{a^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a*x])*x,x]**[Out]** ((2*I)*x)/a - x^2/2 + (2*Log[I + a*x])/a^2**Maple [A]**

time = 0.10, size = 46, normalized size = 1.59

method	result	size
risch	$-\frac{x^2}{2} + \frac{2ix}{a} + \frac{\ln(a^2x^2+1)}{a^2} - \frac{2i \arctan(ax)}{a^2}$	38
default	$-\frac{1}{2}ax^2 + \frac{2ix}{a} + \frac{\ln(a^2x^2+1)}{a} - \frac{2i \arctan(ax)}{a}$	46
meijerg	$\frac{\ln(a^2x^2+1)}{2a^2} + \frac{i \left(\frac{2x(a^2)^{\frac{3}{2}}}{a^2} - \frac{2(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3} \right)}{a\sqrt{a^2}} - \frac{a^2x^2 - \ln(a^2x^2+1)}{2a^2}$	79

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^2/(a^2*x^2+1)*x,x,method=_RETURNVERBOSE)**[Out]** 1/a*(-1/2*a*x^2+2*I*x)+1/a*(1/a*ln(a^2*x^2+1)-2*I*arctan(a*x)/a)**Maxima [A]**

time = 0.48, size = 38, normalized size = 1.31

$$-\frac{ax^2 - 4ix}{2a} - \frac{2i \arctan(ax)}{a^2} + \frac{\log(a^2x^2 + 1)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x,x, algorithm="maxima")**[Out]** -1/2*(a*x^2 - 4*I*x)/a - 2*I*arctan(a*x)/a^2 + log(a^2*x^2 + 1)/a^2**Fricas [A]**

time = 3.00, size = 29, normalized size = 1.00

$$-\frac{a^2x^2 - 4i ax - 4 \log\left(\frac{ax+i}{a}\right)}{2a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x,x, algorithm="fricas")

[Out] -1/2*(a^2*x^2 - 4*I*a*x - 4*log((a*x + I)/a))/a^2

Sympy [A]

time = 0.05, size = 22, normalized size = 0.76

$$-\frac{x^2}{2} + \frac{2ix}{a} + \frac{2 \log(ax + i)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)*x,x)

[Out] -x**2/2 + 2*I*x/a + 2*log(a*x + I)/a**2

Giac [A]

time = 0.41, size = 29, normalized size = 1.00

$$-\frac{a^2x^2 - 4i ax}{2a^2} + \frac{2 \log(ax + i)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x,x, algorithm="giac")

[Out] -1/2*(a^2*x^2 - 4*I*a*x)/a^2 + 2*log(a*x + I)/a^2

Mupad [B]

time = 0.06, size = 27, normalized size = 0.93

$$\frac{2 \ln\left(x + \frac{i}{a}\right)}{a^2} - \frac{x^2}{2} + \frac{x 2i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*(a*x*1i + 1)^2)/(a^2*x^2 + 1),x)

[Out] (2*log(x + 1i/a))/a^2 + (x*2i)/a - x^2/2

3.14 $\int e^{2i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=19

$$-x + \frac{2i \log(i + ax)}{a}$$

[Out] $-x+2*I*\ln(I+a*x)/a$

Rubi [A]

time = 0.01, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5169, 45}

$$-x + \frac{2i \log(ax + i)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}, x]$

[Out] $-x + ((2*I)*\text{Log}[I + a*x])/a$

Rule 45

$\text{Int}[(a_. + (b_.)(x_.))^{(m_.)}((c_.) + (d_.)(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5169

$\text{Int}[E^{(\text{ArcTan}[(a_.)(x_.)]*(n_.))}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}, x] /;$ FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(ax)} dx &= \int \frac{1 + iax}{1 - iax} dx \\ &= \int \left(-1 + \frac{2i}{i + ax} \right) dx \\ &= -x + \frac{2i \log(i + ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 30, normalized size = 1.58

$$-x + \frac{2\text{ArcTan}(ax)}{a} + \frac{i \log(1 + a^2 x^2)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a*x]),x]

[Out] $-x + (2*\text{ArcTan}[a*x])/a + (I*\text{Log}[1 + a^2*x^2])/a$

Maple [A]

time = 0.06, size = 30, normalized size = 1.58

method	result	size
default	$-x + \frac{i \ln(a^2x^2+1)}{a} + \frac{2 \arctan(ax)}{a}$	30
risch	$-x + \frac{i \ln(a^2x^2+1)}{a} + \frac{2 \arctan(ax)}{a}$	30
meijerg	$\frac{\arctan(ax)}{a} + \frac{i \ln(a^2x^2+1)}{a} - \frac{\frac{2x(a^2)^{\frac{3}{2}}}{a^2} - \frac{2(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3}}{2\sqrt{a^2}}$	59

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^2/(a^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] $-x + I/a * \ln(a^2*x^2+1) + 2*\arctan(a*x)/a$

Maxima [A]

time = 0.48, size = 28, normalized size = 1.47

$$-x + \frac{2 \arctan(ax)}{a} + \frac{i \log(a^2x^2 + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1),x, algorithm="maxima")

[Out] $-x + 2*\arctan(a*x)/a + I*\log(a^2*x^2 + 1)/a$

Fricas [A]

time = 1.80, size = 21, normalized size = 1.11

$$\frac{ax - 2i \log\left(\frac{ax+i}{a}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1),x, algorithm="fricas")

[Out] $-(a*x - 2*I*\log((a*x + I)/a))/a$

Sympy [A]

time = 0.05, size = 12, normalized size = 0.63

$$-x + \frac{2i \log(ax + i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1),x)

[Out] $-x + 2*I*\log(ax + I)/a$

Giac [A]

time = 0.41, size = 15, normalized size = 0.79

$$-x + \frac{2i \log(ax + i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1),x, algorithm="giac")

[Out] $-x + 2*I*\log(ax + I)/a$

Mupad [B]

time = 0.04, size = 19, normalized size = 1.00

$$-x + \frac{\ln\left(x + \frac{1i}{a}\right) 2i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/(a^2*x^2 + 1),x)

[Out] $(\log(x + 1i/a)*2i)/a - x$

3.15 $\int \frac{e^{2i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=13

$$\log(x) - 2 \log(i + ax)$$

[Out] `ln(x)-2*ln(I+a*x)`

Rubi [A]

time = 0.01, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$\log(x) - 2 \log(ax + i)$$

Antiderivative was successfully verified.

[In] `Int[E^((2*I)*ArcTan[a*x])/x,x]`

[Out] `Log[x] - 2*Log[I + a*x]`

Rule 78

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)}}{x} dx &= \int \frac{1 + iax}{x(1 - iax)} dx \\ &= \int \left(\frac{1}{x} - \frac{2a}{i + ax} \right) dx \\ &= \log(x) - 2 \log(i + ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 13, normalized size = 1.00

$$\log(x) - 2\log(i + ax)$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a*x])/x,x]

[Out] Log[x] - 2*Log[I + a*x]

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 32 vs. 2(12) = 24.

time = 0.07, size = 33, normalized size = 2.54

method	result	size
risch	$-\ln(a^2x^2 + 1) + 2i \arctan(ax) + \ln(-x)$	25
meijerg	$-\ln(a^2x^2 + 1) + \ln(x) + \frac{\ln(a^2)}{2} + 2i \arctan(ax)$	29
default	$2a \left(-\frac{\ln(a^2x^2+1)}{2a} + \frac{i \arctan(ax)}{a} \right) + \ln(x)$	33

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^2/(a^2*x^2+1)/x,x,method=_RETURNVERBOSE)

[Out] 2*a*(-1/2/a*ln(a^2*x^2+1)+I*arctan(a*x)/a)+ln(x)

Maxima [A]

time = 0.47, size = 21, normalized size = 1.62

$$2i \arctan(ax) - \log(a^2x^2 + 1) + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x,x, algorithm="maxima")

[Out] 2*I*arctan(a*x) - log(a^2*x^2 + 1) + log(x)

Fricas [A]

time = 1.87, size = 15, normalized size = 1.15

$$\log(x) - 2 \log\left(\frac{ax + i}{a}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x,x, algorithm="fricas")

[Out] log(x) - 2*log((a*x + I)/a)

Sympy [A]

time = 0.09, size = 17, normalized size = 1.31

$$\log(3ax) - 2\log(3ax + 3i)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)/x,x)

[Out] log(3*a*x) - 2*log(3*a*x + 3*I)

Giac [A]

time = 0.41, size = 12, normalized size = 0.92

$$-2 \log(ax + i) + \log(|x|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x,x, algorithm="giac")

[Out] -2*log(a*x + I) + log(abs(x))

Mupad [B]

time = 0.44, size = 14, normalized size = 1.08

$$\ln(x) - 2 \ln\left(x + \frac{1i}{a}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/(x*(a^2*x^2 + 1)),x)

[Out] log(x) - 2*log(x + 1i/a)

3.16 $\int \frac{e^{2i \operatorname{ArcTan}(ax)}}{x^2} dx$

Optimal. Leaf size=26

$$-\frac{1}{x} + 2ia \log(x) - 2ia \log(i + ax)$$

[Out] $-1/x+2*I*a*\ln(x)-2*I*a*\ln(I+a*x)$

Rubi [A]

time = 0.02, antiderivative size = 26, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$2ia \log(x) - 2ia \log(ax + i) - \frac{1}{x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}/x^2, x]$

[Out] $-x^{(-1)} + (2*I)*a*\text{Log}[x] - (2*I)*a*\text{Log}[I + a*x]$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_))*((c_. + (d_.)*(x_))^{(n_.)*((e_. + (f_.)*(x_))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_)]*(n_.))*(x_)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, m, n\}, x\} \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{1 + iax}{x^2(1 - iax)} dx \\ &= \int \left(\frac{1}{x^2} + \frac{2ia}{x} - \frac{2ia^2}{i + ax} \right) dx \\ &= -\frac{1}{x} + 2ia \log(x) - 2ia \log(i + ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 26, normalized size = 1.00

$$-\frac{1}{x} + 2ia \log(x) - 2ia \log(i + ax)$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a*x])/x^2,x]``[Out] -x^(-1) + (2*I)*a*Log[x] - (2*I)*a*Log[I + a*x]`**Maple [A]**

time = 0.11, size = 43, normalized size = 1.65

method	result	size
risch	$-\frac{1}{x} - 2a \arctan(ax) - ia \ln(a^2x^2 + 1) + 2ia \ln(x)$	34
default	$-2a^2 \left(\frac{i \ln(a^2x^2+1)}{2a} + \frac{\arctan(ax)}{a} \right) - \frac{1}{x} + 2ia \ln(x)$	43
meijerg	$\frac{a^2 \left(-\frac{2}{x\sqrt{a^2}} - \frac{2a \arctan(ax)}{\sqrt{a^2}} \right)}{2\sqrt{a^2}} + ia(-\ln(a^2x^2 + 1) + 2 \ln(x) + \ln(a^2)) - a \arctan(ax)$	67

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^2/(a^2*x^2+1)/x^2,x,method=_RETURNVERBOSE)``[Out] -2*a^2*(1/2*I/a*ln(a^2*x^2+1)+arctan(a*x)/a)-1/x+2*I*a*ln(x)`**Maxima [A]**

time = 0.53, size = 31, normalized size = 1.19

$$-2a \arctan(ax) - ia \log(a^2x^2 + 1) + 2ia \log(x) - \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^2,x, algorithm="maxima")``[Out] -2*a*arctan(a*x) - I*a*log(a^2*x^2 + 1) + 2*I*a*log(x) - 1/x`**Fricas [A]**

time = 2.73, size = 26, normalized size = 1.00

$$\frac{2i ax \log(x) - 2i ax \log\left(\frac{ax+i}{a}\right) - 1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^2,x, algorithm="fricas")`

[Out] $(2*I*a*x*\log(x) - 2*I*a*x*\log((a*x + I)/a) - 1)/x$

Sympy [A]

time = 0.08, size = 32, normalized size = 1.23

$$-2a(-i \log(4a^2x) + i \log(4a^2x + 4ia)) - \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**2/(a**2*x**2+1)/x**2,x)`

[Out] $-2*a*(-I*\log(4*a**2*x) + I*\log(4*a**2*x + 4*I*a)) - 1/x$

Giac [A]

time = 0.42, size = 21, normalized size = 0.81

$$-2i a \log(ax + i) + 2i a \log(|x|) - \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^2/(a^2*x^2+1)/x^2,x, algorithm="giac")`

[Out] $-2*I*a*\log(a*x + I) + 2*I*a*\log(\text{abs}(x)) - 1/x$

Mupad [B]

time = 0.06, size = 17, normalized size = 0.65

$$-4a \operatorname{atan}(2ax + 1i) - \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*x*1i + 1)^2/(x^2*(a^2*x^2 + 1)),x)`

[Out] $-4*a*\operatorname{atan}(2*a*x + 1i) - 1/x$

3.17 $\int \frac{e^{2i \operatorname{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=36

$$-\frac{1}{2x^2} - \frac{2ia}{x} - 2a^2 \log(x) + 2a^2 \log(i + ax)$$

[Out] $-1/2/x^2 - 2*I*a/x - 2*a^2*\ln(x) + 2*a^2*\ln(I+a*x)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-2a^2 \log(x) + 2a^2 \log(ax + i) - \frac{2ia}{x} - \frac{1}{2x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}/x^3, x]$

[Out] $-1/2*1/x^2 - ((2*I)*a)/x - 2*a^2*\text{Log}[x] + 2*a^2*\text{Log}[I + a*x]$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol)] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, m, n\}, x\} \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{1 + iax}{x^3(1 - iax)} dx \\ &= \int \left(\frac{1}{x^3} + \frac{2ia}{x^2} - \frac{2a^2}{x} + \frac{2a^3}{i + ax} \right) dx \\ &= -\frac{1}{2x^2} - \frac{2ia}{x} - 2a^2 \log(x) + 2a^2 \log(i + ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 36, normalized size = 1.00

$$-\frac{1}{2x^2} - \frac{2ia}{x} - 2a^2 \log(x) + 2a^2 \log(i + ax)$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a*x])/x^3,x]``[Out] -1/2*1/x^2 - ((2*I)*a)/x - 2*a^2*Log[x] + 2*a^2*Log[I + a*x]`**Maple [A]**

time = 0.11, size = 52, normalized size = 1.44

method	result	size
risch	$-\frac{2iax - \frac{1}{2}}{x^2} - 2ia^2 \arctan(ax) + a^2 \ln(a^2x^2 + 1) - 2a^2 \ln(x)$	44
default	$-2a^3 \left(-\frac{\ln(a^2x^2 + 1)}{2a} + \frac{i \arctan(ax)}{a} \right) - \frac{1}{2x^2} - \frac{2ia}{x} - 2a^2 \ln(x)$	52
meijerg	$\frac{a^2 \left(\ln(a^2x^2 + 1) - 2 \ln(x) - \ln(a^2) - \frac{1}{a^2x^2} \right)}{2} + \frac{ia^3 \left(-\frac{2}{x\sqrt{a^2}} - \frac{2a \arctan(ax)}{\sqrt{a^2}} \right)}{\sqrt{a^2}} - \frac{a^2 \left(-\ln(a^2x^2 + 1) + 2 \ln(x) + \ln(a^2) \right)}{2}$	96

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^2/(a^2*x^2+1)/x^3,x,method=_RETURNVERBOSE)``[Out] -2*a^3*(-1/2/a*ln(a^2*x^2+1)+I*arctan(a*x)/a)-1/2/x^2-2*I*a/x-2*a^2*ln(x)`**Maxima [A]**

time = 0.47, size = 42, normalized size = 1.17

$$-2i a^2 \arctan(ax) + a^2 \log(a^2x^2 + 1) - 2a^2 \log(x) - \frac{4i ax + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^3,x, algorithm="maxima")``[Out] -2*I*a^2*arctan(a*x) + a^2*log(a^2*x^2 + 1) - 2*a^2*log(x) - 1/2*(4*I*a*x + 1)/x^2`**Fricas [A]**

time = 1.86, size = 39, normalized size = 1.08

$$-\frac{4a^2x^2 \log(x) - 4a^2x^2 \log\left(\frac{ax+i}{a}\right) + 4i ax + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^3,x, algorithm="fricas")

[Out] -1/2*(4*a^2*x^2*log(x) - 4*a^2*x^2*log((a*x + I)/a) + 4*I*a*x + 1)/x^2

Sympy [A]

time = 0.12, size = 42, normalized size = 1.17

$$-2a^2(\log(4a^3x) - \log(4a^3x + 4ia^2)) - \frac{4iax + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)/x**3,x)

[Out] -2*a**2*(log(4*a**3*x) - log(4*a**3*x + 4*I*a**2)) - (4*I*a*x + 1)/(2*x**2)

Giac [A]

time = 0.40, size = 31, normalized size = 0.86

$$2a^2 \log(ax + i) - 2a^2 \log(|x|) - \frac{4iax + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^3,x, algorithm="giac")

[Out] 2*a^2*log(a*x + I) - 2*a^2*log(abs(x)) - 1/2*(4*I*a*x + 1)/x^2

Mupad [B]

time = 0.08, size = 27, normalized size = 0.75

$$-a^2 \operatorname{atan}(2ax + 1i) 4i - \frac{\frac{1}{2} + ax 2i}{x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/(x^3*(a^2*x^2 + 1)),x)

[Out] - a^2*atan(2*a*x + 1i)*4i - (a*x*2i + 1/2)/x^2

3.18 $\int \frac{e^{2i \operatorname{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=48

$$-\frac{1}{3x^3} - \frac{ia}{x^2} + \frac{2a^2}{x} - 2ia^3 \log(x) + 2ia^3 \log(i + ax)$$

[Out] $-1/3/x^3 - I*a/x^2 + 2*a^2/x - 2*I*a^3*\ln(x) + 2*I*a^3*\ln(I+a*x)$

Rubi [A]

time = 0.02, antiderivative size = 48, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-2ia^3 \log(x) + 2ia^3 \log(ax + i) + \frac{2a^2}{x} - \frac{ia}{x^2} - \frac{1}{3x^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])/x^4}, x]$

[Out] $-1/3*1/x^3 - (I*a)/x^2 + (2*a^2)/x - (2*I)*a^3*\text{Log}[x] + (2*I)*a^3*\text{Log}[I + a*x]$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}), x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0]) \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{1 + iax}{x^4(1 - iax)} dx \\ &= \int \left(\frac{1}{x^4} + \frac{2ia}{x^3} - \frac{2a^2}{x^2} - \frac{2ia^3}{x} + \frac{2ia^4}{i + ax} \right) dx \\ &= -\frac{1}{3x^3} - \frac{ia}{x^2} + \frac{2a^2}{x} - 2ia^3 \log(x) + 2ia^3 \log(i + ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 48, normalized size = 1.00

$$-\frac{1}{3x^3} - \frac{ia}{x^2} + \frac{2a^2}{x} - 2ia^3 \log(x) + 2ia^3 \log(i + ax)$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a*x])/x^4,x]``[Out] -1/3*1/x^3 - (I*a)/x^2 + (2*a^2)/x - (2*I)*a^3*Log[x] + (2*I)*a^3*Log[I + a*x]`**Maple [A]**

time = 0.09, size = 60, normalized size = 1.25

method	result
risch	$\frac{2a^2x^2 - iax - \frac{1}{3}}{x^3} - 2ia^3 \ln(-x) + 2a^3 \arctan(ax) + ia^3 \ln(a^2x^2 + 1)$
default	$2a^4 \left(\frac{i \ln(a^2x^2 + 1)}{2a} + \frac{\arctan(ax)}{a} \right) - \frac{1}{3x^3} - 2ia^3 \ln(x) - \frac{ia}{x^2} + \frac{2a^2}{x}$
meijerg	$\frac{a^4 \left(\frac{2a^2}{x(a^2)^{\frac{3}{2}}} - \frac{2}{3x^3(a^2)^{\frac{3}{2}}} + \frac{2a^3 \arctan(ax)}{(a^2)^{\frac{3}{2}}} \right)}{2\sqrt{a^2}} + ia^3 (\ln(a^2x^2 + 1) - 2 \ln(x) - \ln(a^2) - \frac{1}{a^2x^2}) - \frac{a^4 \left(-\frac{2}{x\sqrt{a^2}} - \frac{2a \arctan(ax)}{a^2} \right)}{2\sqrt{a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^2/(a^2*x^2+1)/x^4,x,method=_RETURNVERBOSE)``[Out] 2*a^4*(1/2*I/a*ln(a^2*x^2+1)+arctan(a*x)/a)-1/3/x^3-2*I*a^3*ln(x)-I*a/x^2+2*a^2/x`**Maxima [A]**

time = 0.49, size = 51, normalized size = 1.06

$$2a^3 \arctan(ax) + ia^3 \log(a^2x^2 + 1) - 2ia^3 \log(x) + \frac{6a^2x^2 - 3iax - 1}{3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^4,x, algorithm="maxima")``[Out] 2*a^3*arctan(a*x) + I*a^3*log(a^2*x^2 + 1) - 2*I*a^3*log(x) + 1/3*(6*a^2*x^2 - 3*I*a*x - 1)/x^3`**Fricas [A]**

time = 2.62, size = 47, normalized size = 0.98

$$\frac{-6ia^3x^3 \log(x) + 6ia^3x^3 \log\left(\frac{ax+i}{a}\right) + 6a^2x^2 - 3iax - 1}{3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^4,x, algorithm="fricas")

[Out] $\frac{1}{3}*(-6*I*a^3*x^3*\log(x) + 6*I*a^3*x^3*\log((a*x + I)/a) + 6*a^2*x^2 - 3*I*a*x - 1)/x^3$

Sympy [A]

time = 0.12, size = 54, normalized size = 1.12

$$-2a^3(i \log(4a^4x) - i \log(4a^4x + 4ia^3)) - \frac{-6a^2x^2 + 3iax + 1}{3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)/x**4,x)

[Out] $-2*a**3*(I*\log(4*a**4*x) - I*\log(4*a**4*x + 4*I*a**3)) - (-6*a**2*x**2 + 3*I*a*x + 1)/(3*x**3)$

Giac [A]

time = 0.39, size = 39, normalized size = 0.81

$$2i a^3 \log(ax + i) - 2i a^3 \log(|x|) + \frac{6 a^2 x^2 - 3i a x - 1}{3 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/x^4,x, algorithm="giac")

[Out] $2*I*a^3*\log(a*x + I) - 2*I*a^3*\log(\text{abs}(x)) + 1/3*(6*a^2*x^2 - 3*I*a*x - 1)/x^3$

Mupad [B]

time = 0.07, size = 34, normalized size = 0.71

$$4 a^3 \operatorname{atan}(2 a x + i) - \frac{-2 a^2 x^2 + a x i + \frac{1}{3}}{x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/(x^4*(a^2*x^2 + 1)),x)

[Out] $4*a^3*\operatorname{atan}(2*a*x + 1i) - (a*x*1i - 2*a^2*x^2 + 1/3)/x^3$

3.19 $\int e^{3i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=137

$$\frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{27\sqrt{1+a^2x^2}}{4a^4} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i-3ax)\sqrt{1+a^2x^2}}{8a^4} - \frac{51i\sinh^{-1}(ax)}{8a^4}$$

[Out] $-51/8*I*\text{arcsinh}(a*x)/a^4+(1+I*a*x)^3/a^4/(a^2*x^2+1)^{(1/2)}+27/4*(a^2*x^2+1)^{(1/2)}/a^4-x^2*(a^2*x^2+1)^{(1/2)}/a^2-1/4*I*x^3*(a^2*x^2+1)^{(1/2)}/a-9/8*I*(2*I-3*a*x)*(a^2*x^2+1)^{(1/2)}/a^4$

Rubi [A]

time = 0.44, antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5168, 1647, 1607, 12, 866, 1649, 1829, 27, 757, 655, 221}

$$-\frac{51i\sinh^{-1}(ax)}{8a^4} - \frac{x^2\sqrt{a^2x^2+1}}{a^2} - \frac{ix^3\sqrt{a^2x^2+1}}{4a} - \frac{9i(-3ax+2i)\sqrt{a^2x^2+1}}{8a^4} + \frac{27\sqrt{a^2x^2+1}}{4a^4} + \frac{(1+iax)^3}{a^4\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)*\text{ArcTan}[a*x])}*x^3, x]$

[Out] $(1 + I*a*x)^3/(a^4*\text{Sqrt}[1 + a^2*x^2]) + (27*\text{Sqrt}[1 + a^2*x^2])/(4*a^4) - (x^2*\text{Sqrt}[1 + a^2*x^2])/a^2 - ((I/4)*x^3*\text{Sqrt}[1 + a^2*x^2])/a - (((9*I)/8)*(2*I - 3*a*x)*\text{Sqrt}[1 + a^2*x^2])/a^4 - (((51*I)/8)*\text{ArcSinh}[a*x])/a^4$

Rule 12

$\text{Int}[(a_)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_)*(v_)] /; \text{FreeQ}[b, x]$

Rule 27

$\text{Int}[(u_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[u*\text{Cancel}[(b/2 + c*x)^{(2*p)}/c^p], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{EqQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{IntegerQ}[p]$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b]$

Rule 655

$\text{Int}[(d_)+(e_)*(x_))*((a_)+(c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*((a+c*x^2)^{(p+1)}/(2*c*(p+1))), x] + \text{Dist}[d, \text{Int}[(a+c*x^2)^p, x], x] /$

; FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]

Rule 757

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^(m - 1)*((a + c*x^2)^(p + 1)/(c*(m + 2*p + 1))), x] + Dist[1/(c*(m + 2*p + 1)), Int[(d + e*x)^(m - 2)*Simp[c*d^2*(m + 2*p + 1) - a*e^2*(m - 1) + 2*c*d*e*(m + p)*x, x]*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && If[RationalQ[m], GtQ[m, 1], SumSimplerQ[m, -2]] && NeQ[m + 2*p + 1, 0] && IntQuadraticQ[a, 0, c, d, e, m, p, x]

Rule 866

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))^(n_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[d^(2*m)/a^m, Int[(f + g*x)^n*((a + c*x^2)^(m + p))/(d - e*x)^m], x], x] /; FreeQ[{a, c, d, e, f, g, n, p}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[f, 0] && ILtQ[m, -1] && !(IGtQ[n, 0] && ILtQ[m + n, 0] && !GtQ[p, 1])

Rule 1607

Int[(u_)*((a_)*(x_)^(p_) + (b_)*(x_)^(q_))^(n_), x_Symbol] := Int[u*x^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] && PosQ[q - p]

Rule 1647

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[d*e, Int[(d + e*x)^(m - 1)*PolynomialQuotient[Pq, a*e + c*d*x, x]*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && PolyQ[Pq, x] && EqQ[c*d^2 + a*e^2, 0] && EqQ[PolynomialRemainder[Pq, a*e + c*d*x, x], 0]

Rule 1649

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a*e + c*d*x, x], f = PolynomialRemainder[Pq, a*e + c*d*x, x]}, Simp[(-d)*f*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*a*e*(p + 1))), x] + Dist[d/(2*a*(p + 1)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^(p + 1)*ExpandToSum[2*a*e*(p + 1)*Q + f*(m + 2*p + 2), x], x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && EqQ[c*d^2 + a*e^2, 0] && ILtQ[p + 1/2, 0] && GtQ[m, 0]

Rule 1829

Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], e = Coeff[Pq, x, Expon[Pq, x]]}, Simp[e*x^(q - 1)*((a + b*x^2)^(p + 1))/(b*(

```

q + 2*p + 1))), x] + Dist[1/(b*(q + 2*p + 1)), Int[(a + b*x^2)^p*ExpandToSum
m[b*(q + 2*p + 1)*Pq - a*e*(q - 1)*x^(q - 2) - b*e*(q + 2*p + 1)*x^q, x], x
], x]] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x] && !LeQ[p, -1]

```

Rule 5168

```

Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{3i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1+iax)^2}{(1-iax)\sqrt{1+a^2x^2}} dx \\
&= - \left((ia) \int \frac{\sqrt{1+a^2x^2} \left(\frac{ix^3}{a} - x^4 \right)}{(1-iax)^2} dx \right) \\
&= - \left((ia) \int \frac{\left(\frac{i}{a} - x \right) x^3 \sqrt{1+a^2x^2}}{(1-iax)^2} dx \right) \\
&= a^2 \int \frac{x^3(1+a^2x^2)^{3/2}}{a^2(1-iax)^3} dx \\
&= \int \frac{x^3(1+a^2x^2)^{3/2}}{(1-iax)^3} dx \\
&= \int \frac{x^3(1+iax)^3}{(1+a^2x^2)^{3/2}} dx \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} - \int \frac{(1+iax)^2 \left(\frac{3i}{a^3} - \frac{x}{a^2} - \frac{ix^2}{a} \right)}{\sqrt{1+a^2x^2}} dx \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{\int \frac{\frac{12i}{a} - 28x - 27iax^2 + 12a^2x^3}{\sqrt{1+a^2x^2}} dx}{4a^2} \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{\int \frac{36ia - 108a^2x - 81ia^3x^2}{\sqrt{1+a^2x^2}} dx}{12a^4} \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{\int -\frac{9ia(-2i+3ax)^2}{\sqrt{1+a^2x^2}} dx}{12a^4} \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} + \frac{(3i) \int \frac{(-2i+3ax)^2}{\sqrt{1+a^2x^2}} dx}{4a^3} \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i-3ax)\sqrt{1+a^2x^2}}{8a^4} + \frac{(3i) \int}{8a^4} \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{27\sqrt{1+a^2x^2}}{4a^4} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i-3ax)\sqrt{1+a^2x^2}}{8a^4} \\
&= \frac{(1+iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{27\sqrt{1+a^2x^2}}{4a^4} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i-3ax)\sqrt{1+a^2x^2}}{8a^4}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 80, normalized size = 0.58

$$\sqrt{1+a^2x^2} \left(\frac{6}{a^4} + \frac{19ix}{8a^3} - \frac{x^2}{a^2} - \frac{ix^3}{4a} + \frac{4i}{a^4(i+ax)} \right) - \frac{51i \sinh^{-1}(ax)}{8a^4}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x])*x^3,x]

[Out] Sqrt[1 + a^2*x^2]*(6/a^4 + (((19*I)/8)*x)/a^3 - x^2/a^2 - ((I/4)*x^3)/a + (4*I)/(a^4*(I + a*x))) - (((51*I)/8)*ArcSinh[a*x])/a^4

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 285 vs. 2(114) = 228.

time = 0.13, size = 286, normalized size = 2.09

method	result
risch	$-\frac{i(2a^3x^3-8ia^2x^2-19ax+48i)\sqrt{a^2x^2+1}}{8a^4} - \frac{51i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{8a^3\sqrt{a^2}} + \frac{4i\sqrt{\left(x+\frac{i}{a}\right)^2a^2-2ia\left(x+\frac{i}{a}\right)}}{a^5\left(x+\frac{i}{a}\right)}$
meijerg	$\frac{-2\sqrt{\pi} + \frac{\sqrt{\pi}(4a^2x^2+8)}{4\sqrt{a^2x^2+1}}}{a^4\sqrt{\pi}} + \frac{3i\left(\frac{\sqrt{\pi}x(a^2)^{\frac{5}{2}}(5a^2x^2+15)}{10a^4\sqrt{a^2x^2+1}} - \frac{3\sqrt{\pi}(a^2)^{\frac{5}{2}}\operatorname{arcsinh}(ax)}{2a^5}\right)}{a^3\sqrt{\pi}\sqrt{a^2}} - \frac{3\left(\frac{8\sqrt{\pi}}{3} - \frac{\sqrt{\pi}(-2a^4x^4+8a^2x^2+1)}{6\sqrt{a^2x^2+1}}\right)}{a^4\sqrt{\pi}}$
default	$-ia^3 \left(\frac{x^5}{4a^2\sqrt{a^2x^2+1}} - \frac{\left(\frac{x^3}{2a^2\sqrt{a^2x^2+1}} - \frac{\left(\frac{x}{a^2\sqrt{a^2x^2+1}} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{a^2\sqrt{a^2}} \right)}{2a^2} \right)}{4a^2} \right) - 3a^2 \left(\dots \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^3,x,method=_RETURNVERBOSE)

[Out] -I*a^3*(1/4*x^5/a^2/(a^2*x^2+1)^(1/2)-5/4/a^2*(1/2*x^3/a^2/(a^2*x^2+1)^(1/2)-3/2/a^2*(-x/a^2/(a^2*x^2+1)^(1/2)+1/a^2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2)))-3*a^2*(1/3*x^4/a^2/(a^2*x^2+1)^(1/2)-4/3/a^2*(x^2/a^2/(a^2*x^2+1)^(1/2)+2/a^4/(a^2*x^2+1)^(1/2)))+3*I*a*(1/2*x^3/a^2/(a^2*x^2+1)

$$\frac{(1/2)-3/2/a^2*(-x/a^2/(a^2*x^2+1)^{(1/2)}+1/a^2*\ln(a^2*x/(a^2)^{(1/2)}+(a^2*x^2+1)^{(1/2)))/(a^2)^{(1/2)))+x^2/a^2/(a^2*x^2+1)^{(1/2)}+2/a^4/(a^2*x^2+1)^{(1/2)}$$

Maxima [A]

time = 0.29, size = 114, normalized size = 0.83

$$-\frac{iax^5}{4\sqrt{a^2x^2+1}} - \frac{x^4}{\sqrt{a^2x^2+1}} + \frac{17ix^3}{8\sqrt{a^2x^2+1}a} + \frac{5x^2}{\sqrt{a^2x^2+1}a^2} + \frac{51ix}{8\sqrt{a^2x^2+1}a^3} - \frac{51i \operatorname{arsinh}(ax)}{8a^4} + \frac{10}{\sqrt{a^2x^2+1}a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^3,x, algorithm="maxima")

[Out] -1/4*I*a*x^5/sqrt(a^2*x^2 + 1) - x^4/sqrt(a^2*x^2 + 1) + 17/8*I*x^3/(sqrt(a^2*x^2 + 1)*a) + 5*x^2/(sqrt(a^2*x^2 + 1)*a^2) + 51/8*I*x/(sqrt(a^2*x^2 + 1)*a^3) - 51/8*I*arcsinh(a*x)/a^4 + 10/(sqrt(a^2*x^2 + 1)*a^4)

Fricas [A]

time = 2.38, size = 88, normalized size = 0.64

$$\frac{32iax - 51(-iax + 1)\log(-ax + \sqrt{a^2x^2 + 1}) + (-2ia^4x^4 - 6a^3x^3 + 11ia^2x^2 + 29ax + 80i)\sqrt{a^2x^2 + 1} - 32}{8(a^5x + ia^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^3,x, algorithm="fricas")

[Out] 1/8*(32*I*a*x - 51*(-I*a*x + 1)*log(-a*x + sqrt(a^2*x^2 + 1)) + (-2*I*a^4*x^4 - 6*a^3*x^3 + 11*I*a^2*x^2 + 29*a*x + 80*I)*sqrt(a^2*x^2 + 1) - 32)/(a^5*x + I*a^4)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i\left(\int \frac{ix^3}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ax^4}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}}\right) dx + \int \frac{a^3x^6}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ia^2x^5}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}}\right) dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)*x**3,x)

[Out] -I*(Integral(I*x**3/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) + Integral(-3*a*x**4/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) + Integral(a**3*x**6/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) + Integral(-3*I*a**2*x**5/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x))

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^3,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in
 dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.46, size = 137, normalized size = 1.00

$$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{4}{(a^2)^{3/2}} + \frac{2\sqrt{a^2}}{a^4} - \frac{x^2 \sqrt{a^2}}{a^2} - \frac{x^3 (a^2)^{3/2} 1i}{4a^3} + \frac{x \sqrt{a^2} 19i}{8a^3} \right)}{\sqrt{a^2}} - \frac{\operatorname{asinh}\left(x \sqrt{a^2}\right) 51i}{8a^3 \sqrt{a^2}} + \frac{\sqrt{a^2 x^2 + 1} 4i}{a^3 \left(x \sqrt{a^2} + \frac{\sqrt{a^2} 1i}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a*x*1i + 1)^3)/(a^2*x^2 + 1)^(3/2),x)

[Out] ((a^2*x^2 + 1)^(1/2)*(4/(a^2)^(3/2) + (2*(a^2)^(1/2))/a^4 - (x^2*(a^2)^(1/2))/a^2 - (x^3*(a^2)^(3/2)*1i)/(4*a^3) + (x*(a^2)^(1/2)*19i)/(8*a^3)))/(a^2)^(1/2) - (asinh(x*(a^2)^(1/2))*51i)/(8*a^3*(a^2)^(1/2)) + ((a^2*x^2 + 1)^(1/2)*4i)/(a^3*(((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))

3.20 $\int e^{3i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=102

$$\frac{i(1+iax)^3}{a^3\sqrt{1+a^2x^2}} + \frac{(28i-3ax)\sqrt{1+a^2x^2}}{6a^3} + \frac{i(3+iax)^2\sqrt{1+a^2x^2}}{3a^3} + \frac{11\sinh^{-1}(ax)}{2a^3}$$

[Out] 11/2*arcsinh(a*x)/a^3+I*(1+I*a*x)^3/a^3/(a^2*x^2+1)^(1/2)+1/6*(28*I-3*a*x)*(a^2*x^2+1)^(1/2)/a^3+1/3*I*(3+I*a*x)^2*(a^2*x^2+1)^(1/2)/a^3

Rubi [A]

time = 0.40, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {5168, 1647, 1607, 12, 866, 1649, 1668, 794, 221}

$$\frac{11\sinh^{-1}(ax)}{2a^3} + \frac{i(1+iax)^3}{a^3\sqrt{a^2x^2+1}} + \frac{i(3+iax)^2\sqrt{a^2x^2+1}}{3a^3} + \frac{(-3ax+28i)\sqrt{a^2x^2+1}}{6a^3}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a*x])*x^2,x]

[Out] (I*(1+I*a*x)^3)/(a^3*Sqrt[1+a^2*x^2]) + ((28*I-3*a*x)*Sqrt[1+a^2*x^2])/(6*a^3) + ((I/3)*(3+I*a*x)^2*Sqrt[1+a^2*x^2])/a^3 + (11*ArcSinh[a*x])/(2*a^3)

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 794

Int[((d_.) + (e_.)*(x_))*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[((e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x)*((a + c*x^2)^(p + 1)/(2*c*(p + 1)*(2*p + 3))), x] - Dist[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && !LeQ[p, -1]

Rule 866

Int[((d_) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))^(n_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[d^(2*m)/a^m, Int[(f + g*x)^n*((a + c*x^2)^(m + p))

```
/(d - e*x)^m), x], x] /; FreeQ[{a, c, d, e, f, g, n, p}, x] && NeQ[e*f - d*
g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[f, 0] && ILtQ[m, -1]
&& !(IGtQ[n, 0] && ILtQ[m + n, 0] && !GtQ[p, 1])
```

Rule 1607

```
Int[(u_.)*((a_.)*(x_)^(p_.) + (b_.)*(x_)^(q_.))^(n_.), x_Symbol] := Int[u*x
^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] &&
PosQ[q - p]
```

Rule 1647

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[d*e, Int[(d + e*x)^(m - 1)*PolynomialQuotient[Pq, a*e + c*d*x, x]*(
a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && PolyQ[Pq, x]
&& EqQ[c*d^2 + a*e^2, 0] && EqQ[PolynomialRemainder[Pq, a*e + c*d*x, x], 0]
```

Rule 1649

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :
> With[{Q = PolynomialQuotient[Pq, a*e + c*d*x, x], f = PolynomialRemainder
[Pq, a*e + c*d*x, x]}, Simp[(-d)*f*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*a*e*
(p + 1))), x] + Dist[d/(2*a*(p + 1)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^(p
+ 1)*ExpandToSum[2*a*e*(p + 1)*Q + f*(m + 2*p + 2), x], x], x] /; FreeQ[{a
, c, d, e}, x] && PolyQ[Pq, x] && EqQ[c*d^2 + a*e^2, 0] && ILtQ[p + 1/2, 0]
&& GtQ[m, 0]
```

Rule 1668

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :
> With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)
^(m + q - 1)*((a + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1))), x] + Di
st[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + c*x^2)^p*ExpandToSum[c
*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)
^(q - 2)*(a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - 2*c*d*e*(m + q + p)
*x), x], x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, c, d,
e, m, p}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && !(EqQ[d, 0] && T
rue) && !(IGtQ[m, 0] && RationalQ[a, c, d, e] && (IntegerQ[p] || ILtQ[p +
1/2, 0]))
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)^(n_)])*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{3i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1+iax)^2}{(1-iax)\sqrt{1+a^2x^2}} dx \\
&= -\left((ia) \int \frac{\sqrt{1+a^2x^2} \left(\frac{ix^2}{a} - x^3\right)}{(1-iax)^2} dx \right) \\
&= -\left((ia) \int \frac{\left(\frac{i}{a} - x\right) x^2 \sqrt{1+a^2x^2}}{(1-iax)^2} dx \right) \\
&= a^2 \int \frac{x^2(1+a^2x^2)^{3/2}}{a^2(1-iax)^3} dx \\
&= \int \frac{x^2(1+a^2x^2)^{3/2}}{(1-iax)^3} dx \\
&= \int \frac{x^2(1+iax)^3}{(1+a^2x^2)^{3/2}} dx \\
&= \frac{i(1+iax)^3}{a^3\sqrt{1+a^2x^2}} - \int \frac{\left(-\frac{3}{a^2} - \frac{ix}{a}\right)(1+iax)^2}{\sqrt{1+a^2x^2}} dx \\
&= \frac{i(1+iax)^3}{a^3\sqrt{1+a^2x^2}} + \frac{i(3+iax)^2\sqrt{1+a^2x^2}}{3a^3} + \frac{1}{3} \int \frac{\left(-\frac{3}{a^2} - \frac{ix}{a}\right)(-5-3iax)}{\sqrt{1+a^2x^2}} dx \\
&= \frac{i(1+iax)^3}{a^3\sqrt{1+a^2x^2}} + \frac{(28i-3ax)\sqrt{1+a^2x^2}}{6a^3} + \frac{i(3+iax)^2\sqrt{1+a^2x^2}}{3a^3} + \frac{11}{2a^2} \int \frac{1}{\sqrt{1+a^2x^2}} dx \\
&= \frac{i(1+iax)^3}{a^3\sqrt{1+a^2x^2}} + \frac{(28i-3ax)\sqrt{1+a^2x^2}}{6a^3} + \frac{i(3+iax)^2\sqrt{1+a^2x^2}}{3a^3} + \frac{11 \sinh^{-1}(ax)}{2a^3}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 63, normalized size = 0.62

$$\frac{\sqrt{1+a^2x^2} \frac{(-52+19iax-7a^2x^2-2ia^3x^3)}{i+ax} + 33 \sinh^{-1}(ax)}{6a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x])*x^2,x]**[Out]** ((Sqrt[1+a^2*x^2]*(-52+(19*I)*a*x-7*a^2*x^2-(2*I)*a^3*x^3))/(I+a*x)+33*ArcSinh[a*x])/(6*a^3)**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 235 vs. 2(85) = 170.

time = 0.12, size = 236, normalized size = 2.31

method	result
risch	$-\frac{i(2a^2x^2-9iax-28)\sqrt{a^2x^2+1}}{6a^3} + \frac{11\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a^2\sqrt{a^2}} - \frac{4\sqrt{\left(x+\frac{i}{a}\right)^2a^2-2ia\left(x+\frac{i}{a}\right)}}{a^4\left(x+\frac{i}{a}\right)}$
meijerg	$-\frac{\sqrt{\pi}x(a^2)^{\frac{3}{2}}}{a^2\sqrt{a^2x^2+1}} + \frac{\sqrt{\pi}(a^2)^{\frac{3}{2}}\operatorname{arcsinh}(ax)}{a^3} + \frac{3i\left(-2\sqrt{\pi} + \frac{\sqrt{\pi}(4a^2x^2+8)}{4\sqrt{a^2x^2+1}}\right)}{a^3\sqrt{\pi}} - \frac{3\left(\frac{\sqrt{\pi}x(a^2)^{\frac{5}{2}}(5a^2x^2+15)}{10a^4\sqrt{a^2x^2+1}} - \frac{3\sqrt{\pi}(a^2)^{\frac{3}{2}}}{2a^3}\right)}{a^2\sqrt{\pi}\sqrt{a^2}}$
default	$-ia^3\left(\frac{x^4}{3a^2\sqrt{a^2x^2+1}} - \frac{4\left(\frac{x^2}{a^2\sqrt{a^2x^2+1}} + \frac{2}{a^4\sqrt{a^2x^2+1}}\right)}{3a^2}\right) - 3a^2\left(\frac{x^3}{2a^2\sqrt{a^2x^2+1}} - \frac{3\left(-\frac{x}{a^2\sqrt{a^2x^2+1}}\right)}{a^2\sqrt{\pi}\sqrt{a^2}}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2,x,method=_RETURNVERBOSE)`

[Out]
$$-Ia^3\left(\frac{1}{3}x^4/a^2/(a^2x^2+1)^{(1/2)} - \frac{4}{3}x^3/a^2/(a^2x^2+1)^{(1/2)} + \frac{2}{a^4}(a^2x^2+1)^{(1/2)}\right) - 3a^2\left(\frac{1}{2}x^3/a^2/(a^2x^2+1)^{(1/2)} - \frac{3}{2}x^2/a^2/(a^2x^2+1)^{(1/2)} + \frac{1}{a^2}\ln(a^2x/(a^2)^{(1/2)} + (a^2x^2+1)^{(1/2)})/(a^2)^{(1/2)}\right) + 3Ia\left(x^2/a^2/(a^2x^2+1)^{(1/2)} + \frac{2}{a^4}(a^2x^2+1)^{(1/2)} - x/a^2/(a^2x^2+1)^{(1/2)} + \frac{1}{a^2}\ln(a^2x/(a^2)^{(1/2)} + (a^2x^2+1)^{(1/2)})/(a^2)^{(1/2)}\right)$$

Maxima [A]

time = 0.27, size = 95, normalized size = 0.93

$$-\frac{iax^4}{3\sqrt{a^2x^2+1}} - \frac{3x^3}{2\sqrt{a^2x^2+1}} + \frac{13ix^2}{3\sqrt{a^2x^2+1}a} - \frac{11x}{2\sqrt{a^2x^2+1}a^2} + \frac{11\operatorname{arsinh}(ax)}{2a^3} + \frac{26i}{3\sqrt{a^2x^2+1}a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2,x, algorithm="maxima")`

[Out]
$$-1/3Iax^4/\sqrt{a^2x^2+1} - 3/2x^3/\sqrt{a^2x^2+1} + 13/3Ix^2/(\sqrt{a^2x^2+1}a) - 11/2x/(\sqrt{a^2x^2+1}a^2) + 11/2\operatorname{arcsinh}(ax)/a^3 + 26/3I/(\sqrt{a^2x^2+1}a^3)$$

Fricas [A]

time = 2.64, size = 80, normalized size = 0.78

$$\frac{24ax + 33(ax+i)\log\left(-ax + \sqrt{a^2x^2+1}\right) - (-2ia^3x^3 - 7a^2x^2 + 19iax - 52)\sqrt{a^2x^2+1} + 24i}{6(a^4x + ia^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2,x, algorithm="fricas")

[Out] $-1/6*(24*a*x + 33*(a*x + I)*\log(-a*x + \sqrt{a^2*x^2 + 1}) - (-2*I*a^3*x^3 - 7*a^2*x^2 + 19*I*a*x - 52)*\sqrt{a^2*x^2 + 1} + 24*I)/(a^4*x + I*a^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \left(\int \frac{ix^2}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ax^3}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx + \int \frac{a^3x^5}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ia^2x^4}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)*x**2,x)

[Out] $-I*(\text{Integral}(I*x**2/(a**2*x**2*\sqrt{a**2*x**2 + 1} + \sqrt{a**2*x**2 + 1}), x) + \text{Integral}(-3*a*x**3/(a**2*x**2*\sqrt{a**2*x**2 + 1} + \sqrt{a**2*x**2 + 1}), x) + \text{Integral}(a**3*x**5/(a**2*x**2*\sqrt{a**2*x**2 + 1} + \sqrt{a**2*x**2 + 1}), x) + \text{Integral}(-3*I*a**2*x**4/(a**2*x**2*\sqrt{a**2*x**2 + 1} + \sqrt{a**2*x**2 + 1}), x))$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.06, size = 114, normalized size = 1.12

$$\frac{11 \operatorname{asinh}\left(x \sqrt{a^2}\right)}{2 a^2 \sqrt{a^2}} - \frac{\sqrt{a^2 x^2 + 1} \left(\frac{3 x \sqrt{a^2}}{2 a^2} - \frac{a 14 i}{3 (a^2)^{3/2}} + \frac{a^3 x^2 1 i}{3 (a^2)^{3/2}} \right)}{\sqrt{a^2}} - \frac{4 \sqrt{a^2 x^2 + 1}}{a^2 \left(x \sqrt{a^2} + \frac{\sqrt{a^2} 1 i}{a} \right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1)^3)/(a^2*x^2 + 1)^(3/2),x)

[Out] $(11*\operatorname{asinh}(x*(a^2)^{(1/2)}))/(2*a^2*(a^2)^{(1/2)}) - ((a^2*x^2 + 1)^{(1/2)}*((a^3*x^2*1i)/(3*(a^2)^{(3/2)}) - (a*14i)/(3*(a^2)^{(3/2)}) + (3*x*(a^2)^{(1/2)}))/(2*a^2)))/(a^2)^{(1/2)} - (4*(a^2*x^2 + 1)^{(1/2)})/(a^2*(((a^2)^{(1/2)}*1i)/a + x*(a^2)^{(1/2)}))*(a^2)^{(1/2)}$

3.21 $\int e^{3i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=92

$$-\frac{9\sqrt{1+a^2x^2}}{2a^2} - \frac{3(1+a^2x^2)^{3/2}}{2a^2(1-iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1-iax)^3} + \frac{9i \sinh^{-1}(ax)}{2a^2}$$

[Out] $-3/2*(a^2*x^2+1)^{(3/2)}/a^2/(1-I*a*x)-(a^2*x^2+1)^{(5/2)}/a^2/(1-I*a*x)^3+9/2*I*\text{arcsinh}(a*x)/a^2-9/2*(a^2*x^2+1)^{(1/2)}/a^2$

Rubi [A]

time = 0.23, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {5168, 1647, 1607, 12, 807, 679, 221}

$$-\frac{(a^2x^2+1)^{5/2}}{a^2(1-iax)^3} - \frac{3(a^2x^2+1)^{3/2}}{2a^2(1-iax)} - \frac{9\sqrt{a^2x^2+1}}{2a^2} + \frac{9i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a*x])*x,x]

[Out] $(-9*\text{Sqrt}[1+a^2*x^2])/(2*a^2) - (3*(1+a^2*x^2)^{(3/2)})/(2*a^2*(1-I*a*x)) - (1+a^2*x^2)^{(5/2)}/(a^2*(1-I*a*x)^3) + ((9*I)/2)*\text{ArcSinh}[a*x]/a^2$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 679

Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[2*c*d*(p/(e^(2*(m + 2*p + 1)))), Int[(d + e*x)^(m + 1)*(a + c*x^2)^(p - 1), x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 + a*e^2, 0] && GtQ[p, 0] && (LeQ[-2, m, 0] || EqQ[m + p + 1, 0]) && NeQ[m + 2*p + 1, 0] && IntegerQ[2*p]

Rule 807

Int[((d_) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d*g - e*f)*(d + e*x)^m*((a + c*x^2)^(p + 1))/(2*c*d*(m

```

+ p + 1))), x] + Dist[(m*(g*c*d + c*e*f) + 2*e*c*f*(p + 1))/(e*(2*c*d)*(m
+ p + 1)), Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e
, f, g, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && ((LtQ[m, -1] && !IGtQ[m + p
+ 1, 0]) || (LtQ[m, 0] && LtQ[p, -1]) || EqQ[m + 2*p + 2, 0]) && NeQ[m + p
+ 1, 0]

```

Rule 1607

```

Int[(u_.)*((a_.)*(x_)^(p_.) + (b_.)*(x_)^(q_.))^(n_.), x_Symbol] := Int[u*x
^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] &&
PosQ[q - p]

```

Rule 1647

```

Int[(Pq_)*((d_) + (e_.)*(x_)^(m_.))*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[d*e, Int[(d + e*x)^(m - 1)*PolynomialQuotient[Pq, a*e + c*d*x, x]*(
a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && PolyQ[Pq, x]
&& EqQ[c*d^2 + a*e^2, 0] && EqQ[PolynomialRemainder[Pq, a*e + c*d*x, x], 0]

```

Rule 5168

```

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{3i \tan^{-1}(ax)} x dx &= \int \frac{x(1+iax)^2}{(1-iax)\sqrt{1+a^2x^2}} dx \\
&= -\left((ia) \int \frac{\left(\frac{ix}{a} - x^2\right) \sqrt{1+a^2x^2}}{(1-iax)^2} dx \right) \\
&= -\left((ia) \int \frac{\left(\frac{i}{a} - x\right) x \sqrt{1+a^2x^2}}{(1-iax)^2} dx \right) \\
&= a^2 \int \frac{x(1+a^2x^2)^{3/2}}{a^2(1-iax)^3} dx \\
&= \int \frac{x(1+a^2x^2)^{3/2}}{(1-iax)^3} dx \\
&= -\frac{(1+a^2x^2)^{5/2}}{a^2(1-iax)^3} + \frac{(3i) \int \frac{(1+a^2x^2)^{3/2}}{(1-iax)^2} dx}{a} \\
&= -\frac{3(1+a^2x^2)^{3/2}}{2a^2(1-iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1-iax)^3} + \frac{(9i) \int \frac{\sqrt{1+a^2x^2}}{1-iax} dx}{2a} \\
&= -\frac{9\sqrt{1+a^2x^2}}{2a^2} - \frac{3(1+a^2x^2)^{3/2}}{2a^2(1-iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1-iax)^3} + \frac{(9i) \int \frac{1}{\sqrt{1+a^2x^2}} dx}{2a} \\
&= -\frac{9\sqrt{1+a^2x^2}}{2a^2} - \frac{3(1+a^2x^2)^{3/2}}{2a^2(1-iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1-iax)^3} + \frac{9i \sinh^{-1}(ax)}{2a^2}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 54, normalized size = 0.59

$$-\frac{i \left(\frac{\sqrt{1+a^2x^2} (14-5iax+a^2x^2)}{i+ax} - 9 \sinh^{-1}(ax) \right)}{2a^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x])*x,x]

[Out] ((-1/2*I)*((Sqrt[1+a^2*x^2]*(14-(5*I)*a*x+a^2*x^2))/(I+a*x)-9*ArcSinh[a*x]))/a^2

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 192 vs. $2(77) = 154$.

time = 0.10, size = 193, normalized size = 2.10

method	result
risch	$-\frac{i(ax-6i)\sqrt{a^2x^2+1}}{2a^2} + \frac{9i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a\sqrt{a^2}} - \frac{4i\sqrt{\left(x+\frac{i}{a}\right)^2 a^2 - 2ia\left(x+\frac{i}{a}\right)}}{a^3\left(x+\frac{i}{a}\right)}$
meijerg	$\frac{\sqrt{\pi} - \frac{\sqrt{\pi}}{\sqrt{a^2x^2+1}}}{a^2\sqrt{\pi}} + \frac{3i\left(-\frac{\sqrt{\pi}x(a^2)^{\frac{3}{2}}}{a^2\sqrt{a^2x^2+1}} + \frac{\sqrt{\pi}(a^2)^{\frac{3}{2}}\operatorname{arcsinh}(ax)}{a^3}\right)}{a\sqrt{\pi}\sqrt{a^2}} - \frac{3\left(-2\sqrt{\pi} + \frac{\sqrt{\pi}(4a^2x^2+8)}{4\sqrt{a^2x^2+1}}\right)}{a^2\sqrt{\pi}} - \frac{i\left(\frac{\sqrt{\pi}}{10}\right)}{10}$
default	$-ia^3 \left(\frac{x^3}{2a^2\sqrt{a^2x^2+1}} - \frac{3\left(\frac{x}{a^2\sqrt{a^2x^2+1}} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{a^2\sqrt{a^2}}\right)}{2a^2} \right) - 3a^2 \left(\frac{x^2}{a^2\sqrt{a^2x^2+1}} + \frac{1}{a^4} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x,x,method=_RETURNVERBOSE)`

[Out]
$$-I*a^3*(1/2*x^3/a^2/(a^2*x^2+1)^(1/2)-3/2/a^2*(-x/a^2/(a^2*x^2+1)^(1/2)+1/a^2*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2))-3*a^2*(x^2/a^2/(a^2*x^2+1)^(1/2)+2/a^4/(a^2*x^2+1)^(1/2))+3*I*a*(-x/a^2/(a^2*x^2+1)^(1/2)+1/a^2*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2))-1/a^2/(a^2*x^2+1)^(1/2)$$

Maxima [A]

time = 0.28, size = 76, normalized size = 0.83

$$-\frac{iax^3}{2\sqrt{a^2x^2+1}} - \frac{3x^2}{\sqrt{a^2x^2+1}} - \frac{9ix}{2\sqrt{a^2x^2+1}a} + \frac{9i \operatorname{arsinh}(ax)}{2a^2} - \frac{7}{\sqrt{a^2x^2+1}a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x,x, algorithm="maxima")`

[Out]
$$-1/2*I*a*x^3/\operatorname{sqrt}(a^2*x^2+1) - 3*x^2/\operatorname{sqrt}(a^2*x^2+1) - 9/2*I*x/(\operatorname{sqrt}(a^2*x^2+1)*a) + 9/2*I*\operatorname{arcsinh}(a*x)/a^2 - 7/(\operatorname{sqrt}(a^2*x^2+1)*a^2)$$

Fricas [A]

time = 2.34, size = 72, normalized size = 0.78

$$\frac{-8iax - 9(iax - 1)\log\left(-ax + \sqrt{a^2x^2+1}\right) + \sqrt{a^2x^2+1}(-ia^2x^2 - 5ax - 14i) + 8}{2(a^3x + ia^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x,x, algorithm="fricas")

[Out] 1/2*(-8*I*a*x - 9*(I*a*x - 1)*log(-a*x + sqrt(a^2*x^2 + 1)) + sqrt(a^2*x^2 + 1)*(-I*a^2*x^2 - 5*a*x - 14*I) + 8)/(a^3*x + I*a^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i\left(\int \frac{ix}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ax^2}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}}\right) dx + \int \frac{a^3x^4}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ia^2x^3}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}}\right) dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)*x,x)

[Out] -I*(Integral(I*x/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) + Integral(-3*a*x**2/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) + Integral(a**3*x**4/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) + Integral(-3*I*a**2*x**3/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.43, size = 104, normalized size = 1.13

$$-\frac{\sqrt{a^2x^2+1} \left(\frac{3\sqrt{a^2}}{a^2} + \frac{x\sqrt{a^2}}{2a} \right)}{\sqrt{a^2}} + \frac{\operatorname{asinh}\left(x\sqrt{a^2}\right) 9i}{2a\sqrt{a^2}} - \frac{\sqrt{a^2x^2+1} 4i}{a \left(x\sqrt{a^2} + \frac{\sqrt{a^2}}{a} \right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*(a*x*1i + 1)^3)/(a^2*x^2 + 1)^(3/2),x)

[Out] (asinh(x*(a^2)^(1/2))*9i)/(2*a*(a^2)^(1/2)) - ((a^2*x^2 + 1)^(1/2)*((3*(a^2)^(1/2))/a^2 + (x*(a^2)^(1/2)*1i)/(2*a)))/(a^2)^(1/2) - ((a^2*x^2 + 1)^(1/2)*4i)/(a*((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))

3.22 $\int e^{3i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=60

$$-\frac{2i(1+iax)^2}{a\sqrt{1+a^2x^2}} - \frac{3i\sqrt{1+a^2x^2}}{a} - \frac{3\sinh^{-1}(ax)}{a}$$

[Out] $-3*\text{arcsinh}(a*x)/a-2*I*(1+I*a*x)^2/a/(a^2*x^2+1)^{(1/2)}-3*I*(a^2*x^2+1)^{(1/2)}/a$

Rubi [A]

time = 0.03, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5167, 867, 683, 655, 221}

$$-\frac{2i(1+iax)^2}{a\sqrt{a^2x^2+1}} - \frac{3i\sqrt{a^2x^2+1}}{a} - \frac{3\sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)*\text{ArcTan}[a*x])}, x]$

[Out] $((-2*I)*(1+I*a*x)^2)/(a*\text{Sqrt}[1+a^2*x^2]) - ((3*I)*\text{Sqrt}[1+a^2*x^2])/a - (3*\text{ArcSinh}[a*x])/a$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /;$ $\text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b]$

Rule 655

$\text{Int}[(d_) + (e_)*(x_)]*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*((a + c*x^2)^{(p+1)})/(2*c*(p+1)), x] + \text{Dist}[d, \text{Int}[(a + c*x^2)^p, x], x] /;$ $\text{FreeQ}\{a, c, d, e, p\}, x\} \ \&\& \ \text{NeQ}[p, -1]$

Rule 683

$\text{Int}[(d_) + (e_)*(x_)]^{(m_)}*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*(d + e*x)^{(m-1)}*((a + c*x^2)^{(p+1)})/(c*(p+1)), x] - \text{Dist}[e^2*((m+p)/(c*(p+1))), \text{Int}[(d + e*x)^{(m-2)}*(a + c*x^2)^{(p+1)}, x], x] /;$ $\text{FreeQ}\{a, c, d, e\}, x\} \ \&\& \ \text{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ \text{GtQ}[m, 1] \ \&\& \ \text{IntegerQ}[2*p]$

Rule 867

$\text{Int}[(d_) + (e_)*(x_)]^{(m_)}*((f_) + (g_)*(x_)]^{(n_)}*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[d^{(2*m)}/a^m, \text{Int}[(f + g*x)^n*(a + c*x^2)^{(m+p)}]$

$(d - e*x)^m, x], x] /;$ FreeQ[{a, c, d, e, f, g, n, p}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && ILtQ[m, 0] && IntegerQ[n]

Rule 5167

Int[E^(ArcTan[(a_.)*(x_)]*(n_)), x_Symbol] := Int[(1 - I*a*x)^((I*n + 1)/2) / ((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[a, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{3i \tan^{-1}(ax)} dx &= \int \frac{(1 + iax)^2}{(1 - iax)\sqrt{1 + a^2x^2}} dx \\ &= \int \frac{(1 + iax)^3}{(1 + a^2x^2)^{3/2}} dx \\ &= -\frac{2i(1 + iax)^2}{a\sqrt{1 + a^2x^2}} - 3 \int \frac{1 + iax}{\sqrt{1 + a^2x^2}} dx \\ &= -\frac{2i(1 + iax)^2}{a\sqrt{1 + a^2x^2}} - \frac{3i\sqrt{1 + a^2x^2}}{a} - 3 \int \frac{1}{\sqrt{1 + a^2x^2}} dx \\ &= -\frac{2i(1 + iax)^2}{a\sqrt{1 + a^2x^2}} - \frac{3i\sqrt{1 + a^2x^2}}{a} - \frac{3 \sinh^{-1}(ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 42, normalized size = 0.70

$$\frac{\sqrt{1 + a^2x^2} \left(-i + \frac{4}{i+ax}\right)}{a} - \frac{3 \sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x]), x]

[Out] (Sqrt[1 + a^2*x^2]*(-I + 4/(I + a*x)))/a - (3*ArcSinh[a*x])/a

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 127 vs. 2(53) = 106.

time = 0.10, size = 128, normalized size = 2.13

method	result
risch	$-\frac{i\sqrt{a^2x^2 + 1}}{a} - \frac{3 \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + 1}\right)}{\sqrt{a^2}} + \frac{4\sqrt{\left(x + \frac{i}{a}\right)^2 a^2 - 2ia\left(x + \frac{i}{a}\right)}}{a^2\left(x + \frac{i}{a}\right)}$

default	$\frac{x}{\sqrt{a^2x^2+1}} - ia^3 \left(\frac{x^2}{a^2\sqrt{a^2x^2+1}} + \frac{2}{a^4\sqrt{a^2x^2+1}} \right) - 3a^2 \left(-\frac{x}{a^2\sqrt{a^2x^2+1}} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2}\right)}{a^2\sqrt{a^2}} \right)$
meijerg	$\frac{x}{\sqrt{a^2x^2+1}} + \frac{3i \left(\sqrt{\pi} - \frac{\sqrt{\pi}}{\sqrt{a^2x^2+1}} \right)}{a\sqrt{\pi}} - \frac{3 \left(-\frac{\sqrt{\pi} x (a^2)^{\frac{3}{2}}}{a^2\sqrt{a^2x^2+1}} + \frac{\sqrt{\pi} (a^2)^{\frac{3}{2}} \operatorname{arcsinh}(ax)}{a^3} \right)}{\sqrt{\pi} \sqrt{a^2}} - \frac{i \left(-2\sqrt{\pi} + \frac{\sqrt{\pi}}{4\sqrt{a^2}} \right)}{a\sqrt{\pi}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^3/(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $x/(a^2x^2+1)^{(1/2)} - I*a^3*(x^2/a^2/(a^2x^2+1)^{(1/2)} + 2/a^4/(a^2x^2+1)^{(1/2)}) - 3*a^2*(-x/a^2/(a^2x^2+1)^{(1/2)} + 1/a^2*\ln(a^2*x/(a^2)^{(1/2)} + (a^2x^2+1)^{(1/2)})/(a^2)^{(1/2)}) - 3*I/a/(a^2x^2+1)^{(1/2)}$

Maxima [A]

time = 0.35, size = 57, normalized size = 0.95

$$-\frac{iax^2}{\sqrt{a^2x^2+1}} + \frac{4x}{\sqrt{a^2x^2+1}} - \frac{3 \operatorname{arsinh}(ax)}{a} - \frac{5i}{\sqrt{a^2x^2+1}a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out] $-I*a*x^2/\sqrt{a^2x^2+1} + 4*x/\sqrt{a^2x^2+1} - 3*\operatorname{arcsinh}(a*x)/a - 5*I/(\sqrt{a^2x^2+1}*a)$

Fricas [A]

time = 1.87, size = 60, normalized size = 1.00

$$\frac{4ax + 3(ax + i) \log(-ax + \sqrt{a^2x^2 + 1}) + \sqrt{a^2x^2 + 1}(-iax + 5) + 4i}{a^2x + ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2),x, algorithm="fricas")`

[Out] $(4*a*x + 3*(a*x + I)*\log(-a*x + \sqrt{a^2x^2 + 1}) + \sqrt{a^2x^2 + 1}*(-I*a*x + 5) + 4*I)/(a^2x + I*a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \left(\int \frac{i}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ax}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx + \int \frac{a^3x^3}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ia^2x^2}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2),x)

[Out] -I*(Integral(I/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) +
Integral(-3*a*x/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) +
Integral(a**3*x**3/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)),
x) + Integral(-3*I*a**2*x**2/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**
2 + 1))), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2),x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.42, size = 72, normalized size = 1.20

$$-\frac{\sqrt{a^2 x^2 + 1} \operatorname{li}}{a} - \frac{3 \operatorname{asinh}\left(x \sqrt{a^2}\right)}{\sqrt{a^2}} + \frac{4 \sqrt{a^2 x^2 + 1}}{\left(x \sqrt{a^2} + \frac{\sqrt{a^2} \operatorname{li}}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^3/(a^2*x^2 + 1)^(3/2),x)

[Out] (4*(a^2*x^2 + 1)^(1/2))/((((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))
- (3*asinh(x*(a^2)^(1/2)))/(a^2)^(1/2) - ((a^2*x^2 + 1)^(1/2)*1i)/a

3.23 $\int \frac{e^{3i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=51

$$\frac{4i\sqrt{1+a^2x^2}}{i+ax} - i \sinh^{-1}(ax) - \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-I*\operatorname{arcsinh}(a*x)-\operatorname{arctanh}((a^2*x^2+1)^{(1/2}))+4*I*(a^2*x^2+1)^{(1/2)/(I+a*x)$

Rubi [A]

time = 0.55, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5168, 6874, 221, 272, 65, 214, 665}

$$\frac{4i\sqrt{a^2x^2+1}}{ax+i} - \tanh^{-1}\left(\sqrt{a^2x^2+1}\right) - i \sinh^{-1}(ax)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)*\operatorname{ArcTan}[a*x])/x}, x]$

[Out] $((4*I)*\operatorname{Sqrt}[1+a^2*x^2]/(I+a*x) - I*\operatorname{ArcSinh}[a*x] - \operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]])$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^{n}}, x], x, (a+b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_. + (b_.)*(x_.)^2], x_Symbol] := \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}[\{a, b\}, x] \&\& \operatorname{GtQ}[a, 0] \&\& \operatorname{PosQ}[b]$

Rule 272

$\operatorname{Int}[(x_.)^{(m_.)*((a_.) + (b_.)*(x_.)^{(n_.))^{(p_.)}, x_Symbol] := \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n]-1)*(a+b*x)^p}, x], x, x^n], x] /; \operatorname{FreeQ}[\{a, b, m, n, p\}, x] \&\& \operatorname{IntegerQ}[\operatorname{Simplify}[(m+1)/n]]$

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d,
e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2,
0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1 + iax)^2}{x(1 - iax)\sqrt{1 + a^2x^2}} dx \\
&= \int \left(-\frac{ia}{\sqrt{1 + a^2x^2}} + \frac{1}{x\sqrt{1 + a^2x^2}} - \frac{4a}{(i + ax)\sqrt{1 + a^2x^2}} \right) dx \\
&= -\left((ia) \int \frac{1}{\sqrt{1 + a^2x^2}} dx \right) - (4a) \int \frac{1}{(i + ax)\sqrt{1 + a^2x^2}} dx + \int \frac{1}{x\sqrt{1 + a^2x^2}} dx \\
&= \frac{4i\sqrt{1 + a^2x^2}}{i + ax} - i \sinh^{-1}(ax) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
&= \frac{4i\sqrt{1 + a^2x^2}}{i + ax} - i \sinh^{-1}(ax) + \frac{\text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2x^2} \right)}{a^2} \\
&= \frac{4i\sqrt{1 + a^2x^2}}{i + ax} - i \sinh^{-1}(ax) - \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 1.08

$$\frac{4i\sqrt{1 + a^2x^2}}{i + ax} - i \sinh^{-1}(ax) + \log(x) - \log \left(1 + \sqrt{1 + a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((3*I)*ArcTan[a*x])/x,x]

[Out] ((4*I)*Sqrt[1 + a^2*x^2])/(I + a*x) - I*ArcSinh[a*x] + Log[x] - Log[1 + Sqrt[1 + a^2*x^2]]

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 100 vs. 2(44) = 88.
time = 0.08, size = 101, normalized size = 1.98

method	result
default	$-ia^3 \left(-\frac{x}{a^2 \sqrt{a^2 x^2 + 1}} + \frac{\ln\left(\frac{a^2 x}{\sqrt{a^2}} + \sqrt{a^2 x^2 + 1}\right)}{a^2 \sqrt{a^2}} \right) + \frac{4}{\sqrt{a^2 x^2 + 1}} + \frac{3iax}{\sqrt{a^2 x^2 + 1}} - \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2 x^2 + 1}}\right)$
meijerg	$\frac{-\sqrt{\pi} + \frac{\sqrt{\pi}}{\sqrt{a^2 x^2 + 1}} - \sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2 x^2 + 1}}{2}\right) + \frac{(2 - 2 \ln(2) + 2 \ln(x) + \ln(a^2)) \sqrt{\pi}}{2}}{\sqrt{\pi}} + \frac{3iax}{\sqrt{a^2 x^2 + 1}} - \frac{3\left(\sqrt{\pi} - \frac{1}{\sqrt{a^2 x^2 + 1}}\right)}{2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x,x,method=_RETURNVERBOSE)

[Out] -I*a^3*(-x/a^2/(a^2*x^2+1)^(1/2)+1/a^2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2))+4/(a^2*x^2+1)^(1/2)+3*I*a*x/(a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2))

Maxima [A]

time = 0.30, size = 46, normalized size = 0.90

$$\frac{4i ax}{\sqrt{a^2 x^2 + 1}} + \frac{4}{\sqrt{a^2 x^2 + 1}} - i \operatorname{arsinh}(ax) - \operatorname{arsinh}\left(\frac{1}{a|x|}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x,x, algorithm="maxima")

[Out] 4*I*a*x/sqrt(a^2*x^2 + 1) + 4/sqrt(a^2*x^2 + 1) - I*arcsinh(a*x) - arcsinh(1/(a*abs(x)))

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 100 vs. 2(41) = 82.

time = 2.17, size = 100, normalized size = 1.96

$$\frac{4i ax - (ax + i) \log(-ax + \sqrt{a^2 x^2 + 1} + 1) + (i ax - 1) \log(-ax + \sqrt{a^2 x^2 + 1}) + (ax + i) \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + 4i \sqrt{a^2 x^2 + 1} - 4}{ax + i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x,x, algorithm="fricas")

[Out] $(4*I*a*x - (a*x + I)*\log(-a*x + \sqrt{a^2*x^2 + 1}) + 1) + (I*a*x - 1)*\log(-a*x + \sqrt{a^2*x^2 + 1}) + (a*x + I)*\log(-a*x + \sqrt{a^2*x^2 + 1}) - 1) + 4*I*\sqrt{a^2*x^2 + 1} - 4)/(a*x + I)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \left(\int \frac{i}{a^2 x^3 \sqrt{a^2 x^2 + 1} + x \sqrt{a^2 x^2 + 1}} dx + \int \left(-\frac{3ax}{a^2 x^3 \sqrt{a^2 x^2 + 1} + x \sqrt{a^2 x^2 + 1}} \right) dx + \int \frac{a^3 x^3}{a^2 x^3 \sqrt{a^2 x^2 + 1} + x \sqrt{a^2 x^2 + 1}} dx + \int \left(-\frac{3ia^2 x^2}{a^2 x^3 \sqrt{a^2 x^2 + 1} + x \sqrt{a^2 x^2 + 1}} \right) dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)/x,x)

[Out] $-I*(\text{Integral}(I/(a**2*x**3*\sqrt{a**2*x**2 + 1}) + x*\sqrt{a**2*x**2 + 1}), x) + \text{Integral}(-3*a*x/(a**2*x**3*\sqrt{a**2*x**2 + 1}) + x*\sqrt{a**2*x**2 + 1}), x) + \text{Integral}(a**3*x**3/(a**2*x**3*\sqrt{a**2*x**2 + 1}) + x*\sqrt{a**2*x**2 + 1}), x) + \text{Integral}(-3*I*a**2*x**2/(a**2*x**3*\sqrt{a**2*x**2 + 1}) + x*\sqrt{a**2*x**2 + 1}), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.43, size = 73, normalized size = 1.43

$$-\operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right) - \frac{a \operatorname{asinh}\left(x \sqrt{a^2}\right) \operatorname{li}}{\sqrt{a^2}} + \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{\left(x \sqrt{a^2} + \frac{\sqrt{a^2} \operatorname{li}}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^3/(x*(a^2*x^2 + 1)^(3/2)),x)

[Out] $(a*(a^2*x^2 + 1)^{(1/2)}*4i)/((((a^2)^{(1/2)}*1i)/a + x*(a^2)^{(1/2)})*(a^2)^{(1/2)}) - (a*\operatorname{asinh}(x*(a^2)^{(1/2)})*1i)/(a^2)^{(1/2)} - \operatorname{atanh}((a^2*x^2 + 1)^{(1/2)})$

3.24 $\int \frac{e^{3i \operatorname{ArcTan}(ax)}}{x^2} dx$

Optimal. Leaf size=63

$$-\frac{\sqrt{1+a^2x^2}}{x} - \frac{4a\sqrt{1+a^2x^2}}{i+ax} - 3ia \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-3*I*a*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-(a^2*x^2+1)^{(1/2)}/x-4*a*(a^2*x^2+1)^{(1/2)}/(I+a*x)$

Rubi [A]

time = 0.46, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5168, 6874, 270, 272, 65, 214, 665}

$$-\frac{4a\sqrt{a^2x^2+1}}{ax+i} - \frac{\sqrt{a^2x^2+1}}{x} - 3ia \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)*\operatorname{ArcTan}[a*x])/x^2}, x]$

[Out] $-(\operatorname{Sqrt}[1+a^2*x^2]/x) - (4*a*\operatorname{Sqrt}[1+a^2*x^2])/(I+a*x) - (3*I)*a*\operatorname{ArcTan}h[\operatorname{Sqrt}[1+a^2*x^2]]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b)^n}, x], x, (a+bx)^{(1/p)}], x]] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 270

$\operatorname{Int}[(c_.)*(x_.)^{(m_.)*((a_.) + (b_.)*(x_.)^{(n_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(c*x)^{(m+1)*((a+bx^n)^{(p+1)/(a*c*(m+1))}], x] /; \operatorname{FreeQ}\{a, b, c, m, n, p\}, x] \&\& \operatorname{EqQ}[(m+1)/n + p + 1, 0] \&\& \operatorname{NeQ}[m, -1]$

Rule 272

$\operatorname{Int}[(x_.)^{(m_.)*((a_.) + (b_.)*(x_.)^{(n_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Dist}[1/n, \operatorname{Subst}[\operatorname{Int}[x^{(\operatorname{Simplify}[(m+1)/n - 1)*(a+bx)^p}, x], x, x^n], x] /; \operatorname{FreeQ}\{a, b$

, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 665

Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d, e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2, 0]

Rule 5168

Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n + 1)/2)/((1 + I*a*x)^(I*(n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{3i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1 + iax)^2}{x^2(1 - iax)\sqrt{1 + a^2x^2}} dx \\
 &= \int \left(\frac{1}{x^2\sqrt{1 + a^2x^2}} + \frac{3ia}{x\sqrt{1 + a^2x^2}} - \frac{4ia^2}{(i + ax)\sqrt{1 + a^2x^2}} \right) dx \\
 &= (3ia) \int \frac{1}{x\sqrt{1 + a^2x^2}} dx - (4ia^2) \int \frac{1}{(i + ax)\sqrt{1 + a^2x^2}} dx + \int \frac{1}{x^2\sqrt{1 + a^2x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2x^2}}{x} - \frac{4a\sqrt{1 + a^2x^2}}{i + ax} + \frac{1}{2}(3ia)\text{Subst}\left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2\right) \\
 &= -\frac{\sqrt{1 + a^2x^2}}{x} - \frac{4a\sqrt{1 + a^2x^2}}{i + ax} + \frac{(3i)\text{Subst}\left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2x^2}\right)}{a} \\
 &= -\frac{\sqrt{1 + a^2x^2}}{x} - \frac{4a\sqrt{1 + a^2x^2}}{i + ax} - 3ia \tanh^{-1}\left(\sqrt{1 + a^2x^2}\right)
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 61, normalized size = 0.97

$$\sqrt{1 + a^2x^2} \left(-\frac{1}{x} - \frac{4a}{i + ax} \right) + 3ia \log(x) - 3ia \log\left(1 + \sqrt{1 + a^2x^2}\right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((3*I)*ArcTan[a*x])/x^2,x]

[Out] Sqrt[1 + a^2*x^2]*(-x^(-1) - (4*a)/(I + a*x)) + (3*I)*a*Log[x] - (3*I)*a*Log[1 + Sqrt[1 + a^2*x^2]]

Maple [A]

time = 0.11, size = 80, normalized size = 1.27

method	result
default	$\frac{ia}{\sqrt{a^2x^2+1}} - \frac{5a^2x}{\sqrt{a^2x^2+1}} + 3ia \left(\frac{1}{\sqrt{a^2x^2+1}} - \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2x^2+1}} \right) \right) - \frac{1}{x\sqrt{a^2x^2+1}}$
risch	$-\frac{\sqrt{a^2x^2+1}}{x} + ia \left(-3 \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2x^2+1}} \right) + \frac{4i\sqrt{\left(x+\frac{i}{a}\right)^2 a^2 - 2ia\left(x+\frac{i}{a}\right)}}{a\left(x+\frac{i}{a}\right)} \right)$
meijerg	$-\frac{2a^2x^2+1}{x\sqrt{a^2x^2+1}} + \frac{3ia \left(-\sqrt{\pi} + \frac{\sqrt{\pi}}{\sqrt{a^2x^2+1}} - \sqrt{\pi} \ln \left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2} \right) + \frac{(2-2\ln(2)+2\ln(x)+\ln(a^2))\sqrt{\pi}}{2} \right)}{\sqrt{\pi}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^2,x,method=_RETURNVERBOSE)

[Out] I*a/(a^2*x^2+1)^(1/2)-5*a^2*x/(a^2*x^2+1)^(1/2)+3*I*a*(1/(a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2)))-1/x/(a^2*x^2+1)^(1/2)

Maxima [A]

time = 0.29, size = 60, normalized size = 0.95

$$-\frac{5a^2x}{\sqrt{a^2x^2+1}} - 3ia \operatorname{arsinh} \left(\frac{1}{a|x|} \right) + \frac{4ia}{\sqrt{a^2x^2+1}} - \frac{1}{\sqrt{a^2x^2+1}x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^2,x, algorithm="maxima")

[Out] -5*a^2*x/sqrt(a^2*x^2+1) - 3*I*a*arcsinh(1/(a*abs(x))) + 4*I*a/sqrt(a^2*x^2+1) - 1/(sqrt(a^2*x^2+1)*x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 109 vs. 2(53) = 106.

time = 2.25, size = 109, normalized size = 1.73

$$\frac{5a^2x^2 + 5iax + 3(i a^2x^2 - ax) \log(-ax + \sqrt{a^2x^2+1} + 1) + 3(-i a^2x^2 + ax) \log(-ax + \sqrt{a^2x^2+1} - 1) + \sqrt{a^2x^2+1} (5ax + i)}{ax^2 + ix}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^2,x, algorithm="fricas")

[Out] $-(5*a^2*x^2 + 5*I*a*x + 3*(I*a^2*x^2 - a*x)*\log(-a*x + \sqrt{a^2*x^2 + 1}) + 1) + 3*(-I*a^2*x^2 + a*x)*\log(-a*x + \sqrt{a^2*x^2 + 1} - 1) + \sqrt{a^2*x^2 + 1}*(5*a*x + I)/(a*x^2 + I*x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \left(\int \frac{i}{a^2 x^4 \sqrt{a^2 x^2 + 1} + x^2 \sqrt{a^2 x^2 + 1}} dx + \int \left(-\frac{3ax}{a^2 x^4 \sqrt{a^2 x^2 + 1} + x^2 \sqrt{a^2 x^2 + 1}} \right) dx + \int \frac{a^3 x^3}{a^2 x^4 \sqrt{a^2 x^2 + 1} + x^2 \sqrt{a^2 x^2 + 1}} dx + \int \left(-\frac{3ia^2 x^2}{a^2 x^4 \sqrt{a^2 x^2 + 1} + x^2 \sqrt{a^2 x^2 + 1}} \right) dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)/x**2,x)

[Out] $-I*(\text{Integral}(I/(a**2*x**4*\sqrt{a**2*x**2 + 1}) + x**2*\sqrt{a**2*x**2 + 1}), x) + \text{Integral}(-3*a*x/(a**2*x**4*\sqrt{a**2*x**2 + 1}) + x**2*\sqrt{a**2*x**2 + 1}), x) + \text{Integral}(a**3*x**3/(a**2*x**4*\sqrt{a**2*x**2 + 1}) + x**2*\sqrt{a**2*x**2 + 1}), x) + \text{Integral}(-3*I*a**2*x**2/(a**2*x**4*\sqrt{a**2*x**2 + 1}) + x**2*\sqrt{a**2*x**2 + 1}), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^2,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.06, size = 75, normalized size = 1.19

$$-a \operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right) 3i - \frac{\sqrt{a^2 x^2 + 1}}{x} - \frac{4a^2 \sqrt{a^2 x^2 + 1}}{\left(x \sqrt{a^2} + \frac{\sqrt{a^2} 1i}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^3/(x^2*(a^2*x^2 + 1)^(3/2)),x)

[Out] $-a*\operatorname{atanh}((a^2*x^2 + 1)^(1/2))*3i - (a^2*x^2 + 1)^(1/2)/x - (4*a^2*(a^2*x^2 + 1)^(1/2))/(((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))$

3.25 $\int \frac{e^{3i \operatorname{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=92

$$-\frac{\sqrt{1+a^2x^2}}{2x^2} - \frac{3ia\sqrt{1+a^2x^2}}{x} - \frac{4ia^2\sqrt{1+a^2x^2}}{i+ax} + \frac{9}{2}a^2 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $9/2*a^2*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/2*(a^2*x^2+1)^{(1/2)}/x^2-3*I*a*(a^2*x^2+1)^{(1/2)}/x-4*I*a^2*(a^2*x^2+1)^{(1/2)}/(I+a*x)$

Rubi [A]

time = 0.51, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {5168, 6874, 272, 44, 65, 214, 270, 665}

$$-\frac{4ia^2\sqrt{a^2x^2+1}}{ax+i} - \frac{3ia\sqrt{a^2x^2+1}}{x} - \frac{\sqrt{a^2x^2+1}}{2x^2} + \frac{9}{2}a^2 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)*\operatorname{ArcTan}[a*x])}/x^3, x]$

[Out] $-1/2*\operatorname{Sqrt}[1+a^2*x^2]/x^2 - ((3*I)*a*\operatorname{Sqrt}[1+a^2*x^2])/x - ((4*I)*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(I+a*x) + (9*a^2*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]])/2$

Rule 44

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)} / ((b*c - a*d)*(m + 1))), x] - \operatorname{Dist}[d*((m + n + 2) / ((b*c - a*d)*(m + 1))), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{ILtQ}[m, -1] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{LtQ}[n, 0]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*
x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n,
p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d,
e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2,
0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1+iax)^2}{x^3(1-iax)\sqrt{1+a^2x^2}} dx \\
&= \int \left(\frac{1}{x^3\sqrt{1+a^2x^2}} + \frac{3ia}{x^2\sqrt{1+a^2x^2}} - \frac{4a^2}{x\sqrt{1+a^2x^2}} + \frac{4a^3}{(i+ax)\sqrt{1+a^2x^2}} \right) dx \\
&= (3ia) \int \frac{1}{x^2\sqrt{1+a^2x^2}} dx - (4a^2) \int \frac{1}{x\sqrt{1+a^2x^2}} dx + (4a^3) \int \frac{1}{(i+ax)\sqrt{1+a^2x^2}} dx \\
&= -\frac{3ia\sqrt{1+a^2x^2}}{x} - \frac{4ia^2\sqrt{1+a^2x^2}}{i+ax} + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x^2\sqrt{1+a^2x}} dx, x, x^2 \right) - (2a^2) \text{Subst} \left(\int \frac{1}{x\sqrt{1+a^2x}} dx, x, x^2 \right) \\
&= -\frac{\sqrt{1+a^2x^2}}{2x^2} - \frac{3ia\sqrt{1+a^2x^2}}{x} - \frac{4ia^2\sqrt{1+a^2x^2}}{i+ax} - 4 \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1+a^2x^2} \right) \\
&= -\frac{\sqrt{1+a^2x^2}}{2x^2} - \frac{3ia\sqrt{1+a^2x^2}}{x} - \frac{4ia^2\sqrt{1+a^2x^2}}{i+ax} + 4a^2 \tanh^{-1} \left(\sqrt{1+a^2x^2} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{x^2\sqrt{1+a^2x}} dx, x, x^2 \right) \\
&= -\frac{\sqrt{1+a^2x^2}}{2x^2} - \frac{3ia\sqrt{1+a^2x^2}}{x} - \frac{4ia^2\sqrt{1+a^2x^2}}{i+ax} + \frac{9}{2} a^2 \tanh^{-1} \left(\sqrt{1+a^2x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 79, normalized size = 0.86

$$\sqrt{1+a^2x^2} \left(-\frac{1}{2x^2} - \frac{3ia}{x} - \frac{4ia^2}{i+ax} \right) - \frac{9}{2} a^2 \log(x) + \frac{9}{2} a^2 \log \left(1 + \sqrt{1+a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

`[In] Integrate[E^((3*I)*ArcTan[a*x])/x^3,x]``[Out] Sqrt[1+a^2*x^2]*(-1/2*1/x^2 - ((3*I)*a)/x - ((4*I)*a^2)/(I+a*x)) - (9*a^2*Log[x])/2 + (9*a^2*Log[1+Sqrt[1+a^2*x^2]])/2`**Maple [A]**

time = 0.13, size = 105, normalized size = 1.14

method	result
default	$-\frac{ia^3x}{\sqrt{a^2x^2+1}} - \frac{1}{2x^2\sqrt{a^2x^2+1}} - \frac{9a^2 \left(\frac{1}{\sqrt{a^2x^2+1}} - \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2x^2+1}} \right) \right)}{2} + 3ia \left(-\frac{1}{x\sqrt{a^2x^2+1}} \right)$
risch	$-\frac{i(6a^3x^3-ia^2x^2+6ax-i)}{2x^2\sqrt{a^2x^2+1}} - \frac{a^2 \left(-9 \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2x^2+1}} \right) + \frac{{}_8i \sqrt{(x+\frac{i}{a})^2 a^2 - 2ia(x+\frac{i}{a})}}{a(x+\frac{i}{a})} \right)}{2}$

meijerg	$\frac{a^2 \left(\frac{\sqrt{\pi} (20a^2x^2+8)}{16a^2x^2} - \frac{\sqrt{\pi} (24a^2x^2+8)}{16a^2x^2\sqrt{a^2x^2+1}} + \frac{3\sqrt{\pi} \ln\left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2}\right)}{2} - \frac{3\left(\frac{5}{3} - 2\ln(2) + 2\ln(x) + \ln(a^2)\right)\sqrt{\pi}}{4} - \frac{\sqrt{\pi}}{2x^2a^2} \right)}{\sqrt{\pi}}$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^3,x,method=_RETURNVERBOSE)`

[Out] $-I*a^3*x/(a^2*x^2+1)^{(1/2)} - 1/2/x^2/(a^2*x^2+1)^{(1/2)} - 9/2*a^2*(1/(a^2*x^2+1)^{(1/2)} - \operatorname{arctanh}(1/(a^2*x^2+1)^{(1/2)})) + 3*I*a*(-1/x/(a^2*x^2+1)^{(1/2)} - 2*a^2*x/(a^2*x^2+1)^{(1/2)})$

Maxima [A]

time = 0.27, size = 81, normalized size = 0.88

$$-\frac{7i a^3 x}{\sqrt{a^2 x^2 + 1}} + \frac{9}{2} a^2 \operatorname{arsinh}\left(\frac{1}{a|x|}\right) - \frac{9 a^2}{2 \sqrt{a^2 x^2 + 1}} - \frac{3i a}{\sqrt{a^2 x^2 + 1} x} - \frac{1}{2 \sqrt{a^2 x^2 + 1} x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^3,x, algorithm="maxima")`

[Out] $-7*I*a^3*x/\sqrt{a^2*x^2+1} + 9/2*a^2*\operatorname{arcsinh}(1/(a*\operatorname{abs}(x))) - 9/2*a^2/\sqrt{a^2*x^2+1} - 3*I*a/(\sqrt{a^2*x^2+1}*x) - 1/2/(\sqrt{a^2*x^2+1}*x^2)$

Fricas [A]

time = 3.10, size = 130, normalized size = 1.41

$$\frac{-14i a^3 x^3 + 14 a^2 x^2 + 9(a^3 x^3 + i a^2 x^2) \log(-ax + \sqrt{a^2 x^2 + 1} + 1) - 9(a^3 x^3 + i a^2 x^2) \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + \sqrt{a^2 x^2 + 1} (-14i a^2 x^2 + 5 a x - i)}{2(a x^3 + i x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^3,x, algorithm="fricas")`

[Out] $1/2*(-14*I*a^3*x^3 + 14*a^2*x^2 + 9*(a^3*x^3 + I*a^2*x^2)*\log(-a*x + \sqrt{a^2*x^2+1} + 1) - 9*(a^3*x^3 + I*a^2*x^2)*\log(-a*x + \sqrt{a^2*x^2+1} - 1) + \sqrt{a^2*x^2+1}*(-14*I*a^2*x^2 + 5*a*x - I))/(a*x^3 + I*x^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \left(\int \frac{i}{a^2 x^5 \sqrt{a^2 x^2 + 1} + x^3 \sqrt{a^2 x^2 + 1}} dx + \int \left(-\frac{3ax}{a^2 x^5 \sqrt{a^2 x^2 + 1} + x^3 \sqrt{a^2 x^2 + 1}} \right) dx + \int \frac{a^3 x^3}{a^2 x^5 \sqrt{a^2 x^2 + 1} + x^3 \sqrt{a^2 x^2 + 1}} dx + \int \left(-\frac{3ia^2 x^2}{a^2 x^5 \sqrt{a^2 x^2 + 1} + x^3 \sqrt{a^2 x^2 + 1}} \right) dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)/x**3,x)`

```
[Out] -I*(Integral(I/(a**2*x**5*sqrt(a**2*x**2 + 1) + x**3*sqrt(a**2*x**2 + 1)),
x) + Integral(-3*a*x/(a**2*x**5*sqrt(a**2*x**2 + 1) + x**3*sqrt(a**2*x**2 +
1)), x) + Integral(a**3*x**3/(a**2*x**5*sqrt(a**2*x**2 + 1) + x**3*sqrt(a*
**2*x**2 + 1)), x) + Integral(-3*I*a**2*x**2/(a**2*x**5*sqrt(a**2*x**2 + 1)
+ x**3*sqrt(a**2*x**2 + 1)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^3,x, algorithm="giac")
```

```
[Out] undef
```

Mupad [B]

time = 0.43, size = 99, normalized size = 1.08

$$-\frac{a^2 \operatorname{atan}\left(\sqrt{a^2 x^2 + 1}\right) 9i}{2} - \frac{\sqrt{a^2 x^2 + 1}}{2x^2} - \frac{a \sqrt{a^2 x^2 + 1} 3i}{x} - \frac{a^3 \sqrt{a^2 x^2 + 1} 4i}{\left(x \sqrt{a^2} + \frac{\sqrt{a^2}}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*x*1i + 1)^3/(x^3*(a^2*x^2 + 1)^(3/2)),x)
```

```
[Out] - (a^2*atan((a^2*x^2 + 1)^(1/2)*1i)*9i)/2 - (a^2*x^2 + 1)^(1/2)/(2*x^2) - (
a*(a^2*x^2 + 1)^(1/2)*3i)/x - (a^3*(a^2*x^2 + 1)^(1/2)*4i)/(((a^2)^(1/2)*1
i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))
```

3.26 $\int \frac{e^{3i \operatorname{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=117

$$-\frac{\sqrt{1+a^2x^2}}{3x^3} - \frac{3ia\sqrt{1+a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1+a^2x^2}}{3x} + \frac{4a^3\sqrt{1+a^2x^2}}{i+ax} + \frac{11}{2}ia^3 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $11/2*I*a^3*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/3*(a^2*x^2+1)^{(1/2)}/x^3-3/2*I*a*(a^2*x^2+1)^{(1/2)}/x^2+14/3*a^2*(a^2*x^2+1)^{(1/2)}/x+4*a^3*(a^2*x^2+1)^{(1/2)}/(I+a*x)$

Rubi [A]

time = 0.51, antiderivative size = 117, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {5168, 6874, 277, 270, 272, 44, 65, 214, 665}

$$\frac{14a^2\sqrt{a^2x^2+1}}{3x} - \frac{3ia\sqrt{a^2x^2+1}}{2x^2} - \frac{\sqrt{a^2x^2+1}}{3x^3} + \frac{4a^3\sqrt{a^2x^2+1}}{ax+i} + \frac{11}{2}ia^3 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)*\operatorname{ArcTan}[a*x])/x^4}, x]$

[Out] $-1/3*\operatorname{Sqrt}[1+a^2*x^2]/x^3 - (((3*I)/2)*a*\operatorname{Sqrt}[1+a^2*x^2])/x^2 + (14*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(3*x) + (4*a^3*\operatorname{Sqrt}[1+a^2*x^2])/(I+a*x) + ((11*I)/2)*a^3*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]]$

Rule 44

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^{(n+1)}/((b*c - a*d)*(m+1))), x] - \operatorname{Dist}[d*((m+n+2)/((b*c - a*d)*(m+1))), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{(-1)}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b]

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 277

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^n)^(p + 1)/(a*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*(m + 1))), Int[x^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n + p + 1], 0] && NeQ[m, -1]
```

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d, e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2, 0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1+iax)^2}{x^4(1-iax)\sqrt{1+a^2x^2}} dx \\
&= \int \left(\frac{1}{x^4\sqrt{1+a^2x^2}} + \frac{3ia}{x^3\sqrt{1+a^2x^2}} - \frac{4a^2}{x^2\sqrt{1+a^2x^2}} - \frac{4ia^3}{x\sqrt{1+a^2x^2}} + \frac{4ia^4}{(i+ax)\sqrt{1+a^2x^2}} \right) dx \\
&= (3ia) \int \frac{1}{x^3\sqrt{1+a^2x^2}} dx - (4a^2) \int \frac{1}{x^2\sqrt{1+a^2x^2}} dx - (4ia^3) \int \frac{1}{x\sqrt{1+a^2x^2}} dx + (4ia^4) \int \frac{1}{(i+ax)\sqrt{1+a^2x^2}} dx \\
&= -\frac{\sqrt{1+a^2x^2}}{3x^3} + \frac{4a^2\sqrt{1+a^2x^2}}{x} + \frac{4a^3\sqrt{1+a^2x^2}}{i+ax} + \frac{1}{2}(3ia)\text{Subst}\left(\int \frac{1}{x^2\sqrt{1+a^2x}} dx, x\right) \\
&= -\frac{\sqrt{1+a^2x^2}}{3x^3} - \frac{3ia\sqrt{1+a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1+a^2x^2}}{3x} + \frac{4a^3\sqrt{1+a^2x^2}}{i+ax} - (4ia)\text{Subst}\left(\int \frac{1}{x\sqrt{1+a^2x}} dx, x\right) \\
&= -\frac{\sqrt{1+a^2x^2}}{3x^3} - \frac{3ia\sqrt{1+a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1+a^2x^2}}{3x} + \frac{4a^3\sqrt{1+a^2x^2}}{i+ax} + 4ia^3 \tanh^{-1}\left(\frac{\sqrt{1+a^2x^2}}{i+ax}\right) \\
&= -\frac{\sqrt{1+a^2x^2}}{3x^3} - \frac{3ia\sqrt{1+a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1+a^2x^2}}{3x} + \frac{4a^3\sqrt{1+a^2x^2}}{i+ax} + \frac{11}{2}ia^3 \tanh^{-1}\left(\frac{\sqrt{1+a^2x^2}}{i+ax}\right)
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 89, normalized size = 0.76

$$\frac{1}{6} \left(\frac{\sqrt{1+a^2x^2}(-2i+7ax+19ia^2x^2+52a^3x^3)}{x^3(i+ax)} - 33ia^3 \log(x) + 33ia^3 \log\left(1+\sqrt{1+a^2x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

`[In] Integrate[E^((3*I)*ArcTan[a*x])/x^4,x]`

```
[Out] ((Sqrt[1 + a^2*x^2]*(-2*I + 7*a*x + (19*I)*a^2*x^2 + 52*a^3*x^3))/(x^3*(I + a*x)) - (33*I)*a^3*Log[x] + (33*I)*a^3*Log[1 + Sqrt[1 + a^2*x^2]])/6
```

Maple [A]

time = 0.13, size = 141, normalized size = 1.21

method	result
risch	$ \frac{28a^4x^4-9ia^3x^3+26a^2x^2-9iax-2}{6x^3\sqrt{a^2x^2+1}} - \frac{ia^3 \left(-11 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right) + \frac{\sqrt{(x+\frac{i}{a})^2 a^2 - 2ia(x+\frac{i}{a})}}{a(x+\frac{i}{a})} \right)}{2} $

default	$-\frac{1}{3x^3\sqrt{a^2x^2+1}} - \frac{13a^2\left(-\frac{1}{x\sqrt{a^2x^2+1}} - \frac{2a^2x}{\sqrt{a^2x^2+1}}\right)}{3} + 3ia\left(-\frac{1}{2x^2\sqrt{a^2x^2+1}} - \frac{3a^2\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{\sqrt{a^2x^2+1}}\right)$
meijerg	$-\frac{-8a^4x^4-4a^2x^2+1}{3x^3\sqrt{a^2x^2+1}} + \frac{3ia^3\left(\frac{\sqrt{\pi}(20a^2x^2+8)}{16a^2x^2} - \frac{\sqrt{\pi}(24a^2x^2+8)}{16a^2x^2\sqrt{a^2x^2+1}} + \frac{3\sqrt{\pi}\ln\left(\frac{1}{2} + \frac{\sqrt{a^2x^2+1}}{2}\right)}{2}\right)}{\sqrt{\pi}} - \frac{3\left(\frac{5}{3} - 2\ln(2) + 2\ln(x)\right)}{4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^4,x,method=_RETURNVERBOSE)`

[Out]
$$-1/3/x^3/(a^2*x^2+1)^{(1/2)} - 13/3*a^2*(-1/x/(a^2*x^2+1)^{(1/2)} - 2*a^2*x/(a^2*x^2+1)^{(1/2)}) + 3*I*a*(-1/2/x^2/(a^2*x^2+1)^{(1/2)} - 3/2*a^2*(1/(a^2*x^2+1)^{(1/2)} - \operatorname{arctanh}(1/(a^2*x^2+1)^{(1/2)}))) - I*a^3*(1/(a^2*x^2+1)^{(1/2)} - \operatorname{arctanh}(1/(a^2*x^2+1)^{(1/2)}))$$

Maxima [A]

time = 0.27, size = 100, normalized size = 0.85

$$\frac{26a^4x}{3\sqrt{a^2x^2+1}} + \frac{11}{2}i a^3 \operatorname{arsinh}\left(\frac{1}{a|x|}\right) - \frac{11i a^3}{2\sqrt{a^2x^2+1}} + \frac{13a^2}{3\sqrt{a^2x^2+1}x} - \frac{3ia}{2\sqrt{a^2x^2+1}x^2} - \frac{1}{3\sqrt{a^2x^2+1}x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^4,x, algorithm="maxima")`

[Out]
$$26/3*a^4*x/\operatorname{sqrt}(a^2*x^2+1) + 11/2*I*a^3*\operatorname{arcsinh}(1/(a*\operatorname{abs}(x))) - 11/2*I*a^3/\operatorname{sqrt}(a^2*x^2+1) + 13/3*a^2/(\operatorname{sqrt}(a^2*x^2+1)*x) - 3/2*I*a/(\operatorname{sqrt}(a^2*x^2+1)*x^2) - 1/3/(\operatorname{sqrt}(a^2*x^2+1)*x^3)$$

Fricas [A]

time = 3.34, size = 139, normalized size = 1.19

$$\frac{52a^4x^4 + 52i a^3x^3 - 33(-i a^4x^4 + a^3x^3) \log(-ax + \sqrt{a^2x^2+1} + 1) - 33(i a^4x^4 - a^3x^3) \log(-ax + \sqrt{a^2x^2+1} - 1) + (52a^3x^3 + 19i a^2x^2 + 7ax - 2i)\sqrt{a^2x^2+1}}{6(ax^4 + ix^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^4,x, algorithm="fricas")`

[Out]
$$1/6*(52*a^4*x^4 + 52*I*a^3*x^3 - 33*(-I*a^4*x^4 + a^3*x^3)*\log(-a*x + \operatorname{sqrt}(a^2*x^2+1) + 1) - 33*(I*a^4*x^4 - a^3*x^3)*\log(-a*x + \operatorname{sqrt}(a^2*x^2+1) - 1) + (52*a^3*x^3 + 19*I*a^2*x^2 + 7*a*x - 2*I)*\operatorname{sqrt}(a^2*x^2+1))/(a*x^4 + I*x^3)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i\left(\int \frac{i}{a^2x^6\sqrt{a^2x^2+1} + x^4\sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ax}{a^2x^6\sqrt{a^2x^2+1} + x^4\sqrt{a^2x^2+1}}\right) dx + \int \frac{a^3x^3}{a^2x^6\sqrt{a^2x^2+1} + x^4\sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ia^2x^2}{a^2x^6\sqrt{a^2x^2+1} + x^4\sqrt{a^2x^2+1}}\right) dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)/x**4,x)

[Out] -I*(Integral(I/(a**2*x**6*sqrt(a**2*x**2 + 1) + x**4*sqrt(a**2*x**2 + 1)), x) + Integral(-3*a*x/(a**2*x**6*sqrt(a**2*x**2 + 1) + x**4*sqrt(a**2*x**2 + 1)), x) + Integral(a**3*x**3/(a**2*x**6*sqrt(a**2*x**2 + 1) + x**4*sqrt(a**2*x**2 + 1)), x) + Integral(-3*I*a**2*x**2/(a**2*x**6*sqrt(a**2*x**2 + 1) + x**4*sqrt(a**2*x**2 + 1)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/x^4,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.41, size = 116, normalized size = 0.99

$$\frac{11a^3 \operatorname{atan}\left(\sqrt{a^2x^2+1} \operatorname{li}\right)}{2} - \frac{\sqrt{a^2x^2+1}}{3x^3} - \frac{a\sqrt{a^2x^2+1} \operatorname{li}}{2x^2} + \frac{14a^2\sqrt{a^2x^2+1}}{3x} + \frac{4a^4\sqrt{a^2x^2+1}}{\left(x\sqrt{a^2} + \frac{\sqrt{a^2} \operatorname{li}}{a}\right)\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^3/(x^4*(a^2*x^2 + 1)^(3/2)),x)

[Out] (11*a^3*atan((a^2*x^2 + 1)^(1/2)*1i))/2 - (a^2*x^2 + 1)^(1/2)/(3*x^3) - (a*(a^2*x^2 + 1)^(1/2)*3i)/(2*x^2) + (14*a^2*(a^2*x^2 + 1)^(1/2))/(3*x) + (4*a^4*(a^2*x^2 + 1)^(1/2))/(((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))

3.27 $\int e^{4i \operatorname{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=65

$$\frac{12ix}{a^3} - \frac{4x^2}{a^2} - \frac{4ix^3}{3a} + \frac{x^4}{4} + \frac{4i}{a^4(i+ax)} + \frac{16 \log(i+ax)}{a^4}$$

[Out] $12*I*x/a^3 - 4*x^2/a^2 - 4/3*I*x^3/a + 1/4*x^4 + 4*I/a^4/(I+a*x) + 16*\ln(I+a*x)/a^4$

Rubi [A]

time = 0.04, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$,

Rules used = {5170, 90}

$$\frac{4i}{a^4(ax+i)} + \frac{16 \log(ax+i)}{a^4} + \frac{12ix}{a^3} - \frac{4x^2}{a^2} - \frac{4ix^3}{3a} + \frac{x^4}{4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(4*I)*\text{ArcTan}[a*x]}*x^3, x]$

[Out] $((12*I)*x)/a^3 - (4*x^2)/a^2 - (((4*I)/3)*x^3)/a + x^4/4 + (4*I)/(a^4*(I + a*x)) + (16*\text{Log}[I + a*x])/a^4$

Rule 90

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))

Rule 5170

$\text{Int}[E^{\text{ArcTan}[(a_.)*(x_.)]*(n_.)*x^{(m_.)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{I*(n/2)})/(1 + I*a*x)^{I*(n/2)}], x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{4i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1+iax)^2}{(1-iax)^2} dx \\ &= \int \left(\frac{12i}{a^3} - \frac{8x}{a^2} - \frac{4ix^2}{a} + x^3 - \frac{4i}{a^3(i+ax)^2} + \frac{16}{a^3(i+ax)} \right) dx \\ &= \frac{12ix}{a^3} - \frac{4x^2}{a^2} - \frac{4ix^3}{3a} + \frac{x^4}{4} + \frac{4i}{a^4(i+ax)} + \frac{16 \log(i+ax)}{a^4} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 65, normalized size = 1.00

$$\frac{12ix}{a^3} - \frac{4x^2}{a^2} - \frac{4ix^3}{3a} + \frac{x^4}{4} + \frac{4i}{a^4(i+ax)} + \frac{16 \log(i+ax)}{a^4}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x])*x^3,x]`

```
[Out] ((12*I)*x)/a^3 - (4*x^2)/a^2 - (((4*I)/3)*x^3)/a + x^4/4 + (4*I)/(a^4*(I + a*x)) + (16*Log[I + a*x])/a^4
```

Maple [A]

time = 0.10, size = 67, normalized size = 1.03

method	result
default	$-\frac{-\frac{1}{4}a^3x^4 + \frac{4}{3}ia^2x^3 + 4ax^2 - 12ix}{a^3} - \frac{4\left(-\frac{i}{a(ax+i)} - \frac{4 \ln(ax+i)}{a}\right)}{a^3}$
risch	$\frac{x^4}{4} - \frac{4ix^3}{3a} - \frac{4x^2}{a^2} + \frac{12ix}{a^3} + \frac{4i}{a^4(ax+i)} + \frac{8 \ln(a^2x^2+1)}{a^4} - \frac{16i \arctan(ax)}{a^4}$
meijerg	$-\frac{\frac{a^2x^2}{a^2x^2+1} + \ln(a^2x^2+1)}{2a^4} + \frac{2i\left(\frac{x(a^2)^{\frac{5}{2}}(10a^2x^2+15)}{5a^4(a^2x^2+1)} - \frac{3(a^2)^{\frac{5}{2}} \arctan(ax)}{a^5}\right)}{a^3\sqrt{a^2}} - \frac{3\left(\frac{x^2a^2(3a^2x^2+6)}{3a^2x^2+3} - 2 \ln(a^2x^2+1)\right)}{a^4} - \frac{2i\left(-\frac{x(a^2)^{\frac{7}{2}}}{2}\right)}{a^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2*x^3,x,method=_RETURNVERBOSE)`

```
[Out] -1/a^3*(-1/4*a^3*x^4+4/3*I*a^2*x^3+4*a*x^2-12*I*x)-4/a^3*(-I/a/(I+a*x)-4/a*ln(I+a*x))
```

Maxima [A]

time = 0.48, size = 77, normalized size = 1.18

$$-\frac{4(-iax-1)}{a^6x^2+a^4} + \frac{3a^3x^4 - 16ia^2x^3 - 48ax^2 + 144ix}{12a^3} - \frac{16i \arctan(ax)}{a^4} + \frac{8 \log(a^2x^2+1)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^3,x, algorithm="maxima")`

```
[Out] -4*(-I*a*x - 1)/(a^6*x^2 + a^4) + 1/12*(3*a^3*x^4 - 16*I*a^2*x^3 - 48*a*x^2 + 144*I*x)/a^3 - 16*I*arctan(a*x)/a^4 + 8*log(a^2*x^2 + 1)/a^4
```

Fricas [A]

time = 2.20, size = 70, normalized size = 1.08

$$\frac{3a^5x^5 - 13ia^4x^4 - 32a^3x^3 + 96ia^2x^2 - 144ax + 192(ax+i) \log\left(\frac{ax+i}{a}\right) + 48i}{12(a^5x + ia^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^3,x, algorithm="fricas")

[Out] 1/12*(3*a^5*x^5 - 13*I*a^4*x^4 - 32*a^3*x^3 + 96*I*a^2*x^2 - 144*a*x + 192*(a*x + I)*log((a*x + I)/a) + 48*I)/(a^5*x + I*a^4)

Sympy [A]

time = 0.11, size = 56, normalized size = 0.86

$$\frac{x^4}{4} + \frac{4i}{a^5x + ia^4} - \frac{4ix^3}{3a} - \frac{4x^2}{a^2} + \frac{12ix}{a^3} + \frac{16 \log(ax + i)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2*x**3,x)

[Out] x**4/4 + 4*I/(a**5*x + I*a**4) - 4*I*x**3/(3*a) - 4*x**2/a**2 + 12*I*x/a**3 + 16*log(a*x + I)/a**4

Giac [A]

time = 0.41, size = 60, normalized size = 0.92

$$\frac{16 \log(ax + i)}{a^4} + \frac{4i}{(ax + i)a^4} + \frac{3a^8x^4 - 16ia^7x^3 - 48a^6x^2 + 144ia^5x}{12a^8}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^3,x, algorithm="giac")

[Out] 16*log(a*x + I)/a^4 + 4*I/((a*x + I)*a^4) + 1/12*(3*a^8*x^4 - 16*I*a^7*x^3 - 48*a^6*x^2 + 144*I*a^5*x)/a^8

Mupad [B]

time = 0.43, size = 60, normalized size = 0.92

$$\frac{16 \ln\left(x + \frac{1i}{a}\right)}{a^4} + \frac{x^4}{4} - \frac{4x^2}{a^2} + \frac{4i}{a^5\left(x + \frac{1i}{a}\right)} + \frac{x 12i}{a^3} - \frac{x^3 4i}{3a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a*x*1i + 1)^4)/(a^2*x^2 + 1)^2,x)

[Out] 4i/(a^5*(x + 1i/a)) + (16*log(x + 1i/a))/a^4 + (x*12i)/a^3 + x^4/4 - (x^3*4i)/(3*a) - (4*x^2)/a^2

3.28 $\int e^{4i \operatorname{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=53

$$-\frac{8x}{a^2} - \frac{2ix^2}{a} + \frac{x^3}{3} - \frac{4}{a^3(i+ax)} + \frac{12i \log(i+ax)}{a^3}$$

[Out] $-8*x/a^2 - 2*I*x^2/a + 1/3*x^3 - 4/a^3/(I+a*x) + 12*I*\ln(I+a*x)/a^3$

Rubi [A]

time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 90}

$$-\frac{4}{a^3(ax+i)} + \frac{12i \log(ax+i)}{a^3} - \frac{8x}{a^2} - \frac{2ix^2}{a} + \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] `Int[E^((4*I)*ArcTan[a*x])*x^2,x]`

[Out] $(-8*x)/a^2 - ((2*I)*x^2)/a + x^3/3 - 4/(a^3*(I + a*x)) + ((12*I)*\text{Log}[I + a*x])/a^3$

Rule 90

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))`

Rule 5170

`Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]`

Rubi steps

$$\begin{aligned} \int e^{4i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1+iax)^2}{(1-iax)^2} dx \\ &= \int \left(-\frac{8}{a^2} - \frac{4ix}{a} + x^2 + \frac{4}{a^2(i+ax)^2} + \frac{12i}{a^2(i+ax)} \right) dx \\ &= -\frac{8x}{a^2} - \frac{2ix^2}{a} + \frac{x^3}{3} - \frac{4}{a^3(i+ax)} + \frac{12i \log(i+ax)}{a^3} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 53, normalized size = 1.00

$$-\frac{8x}{a^2} - \frac{2ix^2}{a} + \frac{x^3}{3} - \frac{4}{a^3(i+ax)} + \frac{12i \log(i+ax)}{a^3}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x])*x^2,x]``[Out] (-8*x)/a^2 - ((2*I)*x^2)/a + x^3/3 - 4/(a^3*(I + a*x)) + ((12*I)*Log[I + a*x])/a^3`**Maple [A]**

time = 0.09, size = 58, normalized size = 1.09

method	result
default	$-\frac{8x - \frac{1}{3}a^2x^3 + 2iax^2}{a^2} + \frac{-\frac{4}{a(ax+i)} + \frac{12i \ln(ax+i)}{a}}{a^2}$
risch	$-\frac{8x}{a^2} + \frac{x^3}{3} - \frac{2ix^2}{a} - \frac{4}{a^3(ax+i)} + \frac{6i \ln(a^2x^2+1)}{a^3} + \frac{12 \arctan(ax)}{a^3}$
meijerg	$\frac{-\frac{x(a^2)^{\frac{3}{2}}}{a^2(a^2x^2+1)} + \frac{(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3}}{2a^2\sqrt{a^2}} + \frac{2i\left(-\frac{a^2x^2}{a^2x^2+1} + \ln(a^2x^2+1)\right)}{a^3} - \frac{3\left(\frac{x(a^2)^{\frac{5}{2}}(10a^2x^2+15)}{5a^4(a^2x^2+1)} - \frac{3(a^2)^{\frac{5}{2}} \arctan(ax)}{a^5}\right)}{a^2\sqrt{a^2}} - \frac{2i\left(\frac{x^2a^2}{3a^2}\right)}{a^2\sqrt{a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2*x^2,x,method=_RETURNVERBOSE)``[Out] -1/a^2*(8*x-1/3*a^2*x^3+2*I*a*x^2)+4/a^2*(-1/a/(I+a*x)+3*I*ln(I+a*x)/a)`**Maxima [A]**

time = 0.50, size = 67, normalized size = 1.26

$$-\frac{4(ax-i)}{a^5x^2+a^3} + \frac{a^2x^3-6iax^2-24x}{3a^2} + \frac{12 \arctan(ax)}{a^3} + \frac{6i \log(a^2x^2+1)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^2,x, algorithm="maxima")``[Out] -4*(a*x - I)/(a^5*x^2 + a^3) + 1/3*(a^2*x^3 - 6*I*a*x^2 - 24*x)/a^2 + 12*arctan(a*x)/a^3 + 6*I*log(a^2*x^2 + 1)/a^3`**Fricas [A]**

time = 2.26, size = 62, normalized size = 1.17

$$\frac{a^4x^4 - 5ia^3x^3 - 18a^2x^2 - 24iax - 36(-iax+1) \log\left(\frac{ax+i}{a}\right) - 12}{3(a^4x+ia^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^2,x, algorithm="fricas")

[Out] 1/3*(a^4*x^4 - 5*I*a^3*x^3 - 18*a^2*x^2 - 24*I*a*x - 36*(-I*a*x + 1)*log((a*x + I)/a) - 12)/(a^4*x + I*a^3)

Sympy [A]

time = 0.11, size = 44, normalized size = 0.83

$$\frac{x^3}{3} - \frac{4}{a^4x + ia^3} - \frac{2ix^2}{a} - \frac{8x}{a^2} + \frac{12i \log(ax + i)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2*x**2,x)

[Out] x**3/3 - 4/(a**4*x + I*a**3) - 2*I*x**2/a - 8*x/a**2 + 12*I*log(a*x + I)/a**3

Giac [A]

time = 0.40, size = 51, normalized size = 0.96

$$\frac{12i \log(ax + i)}{a^3} - \frac{4}{(ax + i)a^3} + \frac{a^6x^3 - 6ia^5x^2 - 24a^4x}{3a^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^2,x, algorithm="giac")

[Out] 12*I*log(a*x + I)/a^3 - 4/((a*x + I)*a^3) + 1/3*(a^6*x^3 - 6*I*a^5*x^2 - 24*a^4*x)/a^6

Mupad [B]

time = 0.06, size = 51, normalized size = 0.96

$$\frac{x^3}{3} + \frac{\ln\left(x + \frac{1i}{a}\right) 12i}{a^3} - \frac{8x}{a^2} - \frac{4}{a^4\left(x + \frac{1i}{a}\right)} - \frac{x^2 2i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1)^4)/(a^2*x^2 + 1)^2,x)

[Out] (log(x + 1i/a)*12i)/a^3 - 4/(a^4*(x + 1i/a)) - (8*x)/a^2 + x^3/3 - (x^2*2i)/a

3.29 $\int e^{4i \operatorname{ArcTan}(ax)} x dx$

Optimal. Leaf size=45

$$-\frac{4ix}{a} + \frac{x^2}{2} - \frac{4i}{a^2(i+ax)} - \frac{8 \log(i+ax)}{a^2}$$

[Out] $-4*I*x/a+1/2*x^2-4*I/a^2/(I+a*x)-8*\ln(I+a*x)/a^2$

Rubi [A]

time = 0.02, antiderivative size = 45, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5170, 78}

$$-\frac{4i}{a^2(ax+i)} - \frac{8 \log(ax+i)}{a^2} - \frac{4ix}{a} + \frac{x^2}{2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((4*I)*\text{ArcTan}[a*x])*x}, x]$

[Out] $((-4*I)*x)/a + x^2/2 - (4*I)/(a^2*(I + a*x)) - (8*\text{Log}[I + a*x])/a^2$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[a_.)*(x_.)]*(n_.)*(x_.)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{4i \tan^{-1}(ax)} x dx &= \int \frac{x(1+iax)^2}{(1-iax)^2} dx \\ &= \int \left(-\frac{4i}{a} + x + \frac{4i}{a(i+ax)^2} - \frac{8}{a(i+ax)} \right) dx \\ &= -\frac{4ix}{a} + \frac{x^2}{2} - \frac{4i}{a^2(i+ax)} - \frac{8 \log(i+ax)}{a^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 45, normalized size = 1.00

$$-\frac{4ix}{a} + \frac{x^2}{2} - \frac{4i}{a^2(i+ax)} - \frac{8\log(i+ax)}{a^2}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x])*x,x]``[Out] ((-4*I)*x)/a + x^2/2 - (4*I)/(a^2*(I + a*x)) - (8*Log[I + a*x])/a^2`**Maple [A]**

time = 0.07, size = 50, normalized size = 1.11

method	result
default	$-\frac{-\frac{1}{2}ax^2+4ix}{a} + \frac{-\frac{4i}{a(ax+i)} - \frac{8\ln(ax+i)}{a}}{a}$
risch	$\frac{x^2}{2} - \frac{4ix}{a} - \frac{4i}{a^2(ax+i)} - \frac{4\ln(a^2x^2+1)}{a^2} + \frac{8i\arctan(ax)}{a^2}$
meijerg	$\frac{x^2}{2a^2x^2+2} + \frac{2i\left(-\frac{x(a^2)^{\frac{3}{2}}}{a^2(a^2x^2+1)} + \frac{(a^2)^{\frac{3}{2}}\arctan(ax)}{a^3}\right)}{a\sqrt{a^2}} - \frac{3\left(-\frac{a^2x^2}{a^2x^2+1} + \ln(a^2x^2+1)\right)}{a^2} - \frac{2i\left(\frac{x(a^2)^{\frac{5}{2}}(10a^2x^2+15)}{5a^4(a^2x^2+1)} - \frac{3(a^2)^{\frac{5}{2}}\arctan(ax)}{a^5}\right)}{a\sqrt{a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2*x,x,method=_RETURNVERBOSE)``[Out] -1/a*(-1/2*a*x^2+4*I*x)+4/a*(-I/a/(I+a*x)-2/a*ln(I+a*x))`**Maxima [A]**

time = 0.48, size = 60, normalized size = 1.33

$$-\frac{4(iax+1)}{a^4x^2+a^2} + \frac{ax^2-8ix}{2a} + \frac{8i\arctan(ax)}{a^2} - \frac{4\log(a^2x^2+1)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x,x, algorithm="maxima")``[Out] -4*(I*a*x + 1)/(a^4*x^2 + a^2) + 1/2*(a*x^2 - 8*I*x)/a + 8*I*arctan(a*x)/a^2 - 4*log(a^2*x^2 + 1)/a^2`**Fricas [A]**

time = 3.62, size = 53, normalized size = 1.18

$$\frac{a^3x^3 - 7ia^2x^2 + 8ax - 16(ax+i)\log\left(\frac{ax+i}{a}\right) - 8i}{2(a^3x + ia^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x,x, algorithm="fricas")

[Out] $\frac{1}{2}(a^3x^3 - 7Ia^2x^2 + 8ax - 16(ax + I)\log((ax + I)/a) - 8I)/(a^3x + Ia^2)$

Sympy [A]

time = 0.09, size = 36, normalized size = 0.80

$$\frac{x^2}{2} - \frac{4i}{a^3x + ia^2} - \frac{4ix}{a} - \frac{8 \log(ax + i)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2*x,x)

[Out] $x^{**2}/2 - 4*I/(a^{**3}*x + I*a^{**2}) - 4*I*x/a - 8*\log(ax + I)/a^{**2}$

Giac [A]

time = 0.40, size = 43, normalized size = 0.96

$$-\frac{8 \log(ax + i)}{a^2} + \frac{a^4x^2 - 8ia^3x}{2a^4} - \frac{4i}{(ax + i)a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x,x, algorithm="giac")

[Out] $-8*\log(ax + I)/a^2 + 1/2*(a^4*x^2 - 8*I*a^3*x)/a^4 - 4*I/((ax + I)*a^2)$

Mupad [B]

time = 0.06, size = 43, normalized size = 0.96

$$\frac{x^2}{2} - \frac{8 \ln\left(x + \frac{1i}{a}\right)}{a^2} - \frac{4i}{a^3\left(x + \frac{1i}{a}\right)} - \frac{x 4i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*(a*x*1i + 1)^4)/(a^2*x^2 + 1)^2,x)

[Out] $x^2/2 - (8*\log(x + 1i/a))/a^2 - (x*4i)/a - 4i/(a^3*(x + 1i/a))$

3.30 $\int e^{4i \operatorname{ArcTan}(ax)} dx$

Optimal. Leaf size=31

$$x + \frac{4}{a(i+ax)} - \frac{4i \log(i+ax)}{a}$$

[Out] x+4/a/(I+a*x)-4*I*ln(I+a*x)/a

Rubi [A]

time = 0.01, antiderivative size = 31, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5169, 45}

$$\frac{4}{a(ax+i)} - \frac{4i \log(ax+i)}{a} + x$$

Antiderivative was successfully verified.

[In] Int[E^((4*I)*ArcTan[a*x]),x]

[Out] x + 4/(a*(I + a*x)) - ((4*I)*Log[I + a*x])/a

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]

Rule 5169

Int[E^(ArcTan[(a_.)*(x_)^(n_.)]), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{4i \tan^{-1}(ax)} dx &= \int \frac{(1+iax)^2}{(1-iax)^2} dx \\ &= \int \left(1 - \frac{4}{(i+ax)^2} - \frac{4i}{i+ax} \right) dx \\ &= x + \frac{4}{a(i+ax)} - \frac{4i \log(i+ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 1.35

$$x + \frac{4}{a(i + ax)} - \frac{4\text{ArcTan}(ax)}{a} - \frac{2i \log(1 + a^2x^2)}{a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x]),x]``[Out] x + 4/(a*(I + a*x)) - (4*ArcTan[a*x])/a - ((2*I)*Log[1 + a^2*x^2])/a`**Maple [A]**

time = 0.08, size = 33, normalized size = 1.06

method	result
default	$x - 4a \left(-\frac{1}{a^2(ax+i)} + \frac{i \ln(ax+i)}{a^2} \right)$
risch	$x + \frac{4}{a(ax+i)} - \frac{2i \ln(a^2x^2+1)}{a} - \frac{4 \arctan(ax)}{a}$
meijerg	$\frac{\frac{2x\sqrt{a^2}}{2a^2x^2+2} + \frac{\sqrt{a^2} \arctan(ax)}{a}}{2\sqrt{a^2}} + \frac{2iax^2}{a^2x^2+1} - \frac{3 \left(-\frac{x(a^2)^{\frac{3}{2}}}{a^2(a^2x^2+1)} + \frac{(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3} \right)}{\sqrt{a^2}} - \frac{2i \left(-\frac{a^2x^2}{a^2x^2+1} + \ln(a^2x^2+1) \right)}{a} + \frac{x(a^2)^{\frac{5}{2}}}{5a^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2,x,method=_RETURNVERBOSE)``[Out] x-4*a*(-1/a^2/(I+a*x)+I/a^2*ln(I+a*x))`**Maxima [A]**

time = 0.48, size = 44, normalized size = 1.42

$$x + \frac{4(ax - i)}{a^3x^2 + a} - \frac{4 \arctan(ax)}{a} - \frac{2i \log(a^2x^2 + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2,x, algorithm="maxima")``[Out] x + 4*(a*x - I)/(a^3*x^2 + a) - 4*arctan(a*x)/a - 2*I*log(a^2*x^2 + 1)/a`**Fricas [A]**

time = 1.65, size = 43, normalized size = 1.39

$$\frac{a^2x^2 + iax - 4(iax - 1) \log\left(\frac{ax+i}{a}\right) + 4}{a^2x + ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2,x, algorithm="fricas")

[Out] (a^2*x^2 + I*a*x - 4*(I*a*x - 1)*log((a*x + I)/a) + 4)/(a^2*x + I*a)

Sympy [A]

time = 0.09, size = 22, normalized size = 0.71

$$x + \frac{4}{a^2x + ia} - \frac{4i \log(ax + i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2,x)

[Out] x + 4/(a**2*x + I*a) - 4*I*log(a*x + I)/a

Giac [A]

time = 0.40, size = 25, normalized size = 0.81

$$x - \frac{4i \log(ax + i)}{a} + \frac{4}{(ax + i)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2,x, algorithm="giac")

[Out] x - 4*I*log(a*x + I)/a + 4/((a*x + I)*a)

Mupad [B]

time = 0.43, size = 32, normalized size = 1.03

$$x + \frac{4}{a^2 \left(x + \frac{1i}{a}\right)} - \frac{\ln\left(x + \frac{1i}{a}\right) 4i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(a^2*x^2 + 1)^2,x)

[Out] x + 4/(a^2*(x + 1i/a)) - (log(x + 1i/a)*4i)/a

3.31 $\int \frac{e^{4i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=16

$$\frac{4i}{i + ax} + \log(x)$$

[Out] 4*I/(I+a*x)+ln(x)

Rubi [A]

time = 0.02, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 90}

$$\log(x) + \frac{4i}{ax + i}$$

Antiderivative was successfully verified.

[In] Int[E^((4*I)*ArcTan[a*x])/x,x]

[Out] (4*I)/(I + a*x) + Log[x]

Rule 90

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1 + iax)^2}{x(1 - iax)^2} dx \\ &= \int \left(\frac{1}{x} - \frac{4ia}{(i + ax)^2} \right) dx \\ &= \frac{4i}{i + ax} + \log(x) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$\frac{4i}{i + ax} + \log(x)$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x])/x,x]``[Out] (4*I)/(I + a*x) + Log[x]`**Maple [A]**

time = 0.09, size = 15, normalized size = 0.94

method	result
default	$\frac{4i}{ax+i} + \ln(x)$
risch	$\frac{4i}{ax+i} + \ln(-x)$
norman	$\frac{-4a^2x^2+4iax}{a^2x^2+1} + \ln(x)$
meijerg	$-\frac{a^2x^2}{2a^2x^2+2} + \frac{1}{2} + \ln(x) + \frac{\ln(a^2)}{2} + \frac{2ia \left(\frac{2x\sqrt{a^2}}{2a^2x^2+2} + \frac{\sqrt{a^2} \arctan(ax)}{a} \right)}{\sqrt{a^2}} - \frac{7a^2x^2}{2(a^2x^2+1)} - \frac{2ia \left(-\frac{x(a^2)^{\frac{3}{2}}}{a^2(a^2x^2+1)} + \frac{(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3} \right)}{\sqrt{a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2/x,x,method=_RETURNVERBOSE)``[Out] 4*I/(I+a*x)+ln(x)`**Maxima [A]**

time = 0.48, size = 22, normalized size = 1.38

$$-\frac{4(-i ax - 1)}{a^2x^2 + 1} + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x,x, algorithm="maxima")``[Out] -4*(-I*a*x - 1)/(a^2*x^2 + 1) + log(x)`**Fricas [A]**

time = 2.43, size = 18, normalized size = 1.12

$$\frac{(ax + i) \log(x) + 4i}{ax + i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x,x, algorithm="fricas")

[Out] ((a*x + I)*log(x) + 4*I)/(a*x + I)

Sympy [A]

time = 0.11, size = 10, normalized size = 0.62

$$\log(x) + \frac{4i}{ax + i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2/x,x)

[Out] log(x) + 4*I/(a*x + I)

Giac [A]

time = 0.41, size = 13, normalized size = 0.81

$$\frac{4i}{ax + i} + \log(|x|)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x,x, algorithm="giac")

[Out] 4*I/(a*x + I) + log(abs(x))

Mupad [B]

time = 0.08, size = 14, normalized size = 0.88

$$\ln(x) + \frac{4i}{ax + 1i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(x*(a^2*x^2 + 1)^2),x)

[Out] log(x) + 4i/(a*x + 1i)

3.32 $\int \frac{e^{4i \operatorname{ArcTan}(ax)}}{x^2} dx$

Optimal. Leaf size=38

$$-\frac{1}{x} - \frac{4a}{i+ax} + 4ia \log(x) - 4ia \log(i+ax)$$

[Out] $-1/x - 4*a/(I+a*x) + 4*I*a*\ln(x) - 4*I*a*\ln(I+a*x)$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 90}

$$-\frac{4a}{ax+i} + 4ia \log(x) - 4ia \log(ax+i) - \frac{1}{x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(4*I)*\text{ArcTan}[a*x]}/x^2, x]$

[Out] $-x^{(-1)} - (4*a)/(I + a*x) + (4*I)*a*\text{Log}[x] - (4*I)*a*\text{Log}[I + a*x]$

Rule 90

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, p\}, x\} \&\& \text{IntegersQ}\{m, n\} \&\& (\text{IntegerQ}\{p\} \|\ (\text{GtQ}\{m, 0\} \&\& \text{GeQ}\{n, -1\}))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*x_.^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, m, n\}, x\} \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1+iax)^2}{x^2(1-iax)^2} dx \\ &= \int \left(\frac{1}{x^2} + \frac{4ia}{x} + \frac{4a^2}{(i+ax)^2} - \frac{4ia^2}{i+ax} \right) dx \\ &= -\frac{1}{x} - \frac{4a}{i+ax} + 4ia \log(x) - 4ia \log(i+ax) \end{aligned}$$

Mathematica [A]

time = 0.02, size = 38, normalized size = 1.00

$$-\frac{1}{x} - \frac{4a}{i+ax} + 4ia \log(x) - 4ia \log(i+ax)$$

Antiderivative was successfully verified.

[In] Integrate[E^((4*I)*ArcTan[a*x])/x^2,x]

[Out] -x^(-1) - (4*a)/(I + a*x) + (4*I)*a*Log[x] - (4*I)*a*Log[I + a*x]

Maple [A]

time = 0.09, size = 44, normalized size = 1.16

method	result
default	$-4a^2 \left(\frac{1}{a(ax+i)} + \frac{i \ln(ax+i)}{a} \right) - \frac{1}{x} + 4ia \ln(x)$
risch	$\frac{-5ax-i}{(ax+i)x} + 4ia \ln(x) - 4a \arctan(ax) - 2ia \ln(a^2x^2 + 1)$
meijerg	$\frac{a^2 \left(-\frac{2(3a^2x^2+2)}{x\sqrt{a^2(2a^2x^2+2)}} - \frac{3a \arctan(ax)}{\sqrt{a^2}} \right)}{2\sqrt{a^2}} + 2ia \left(-\frac{2a^2x^2}{2a^2x^2+2} - \ln(a^2x^2 + 1) + 1 + 2 \ln(x) + \ln(a^2) \right) - \frac{3a^2}{\dots}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^4/(a^2*x^2+1)^2/x^2,x,method=_RETURNVERBOSE)

[Out] -4*a^2*(1/a/(I+a*x)+I*ln(I+a*x)/a)-1/x+4*I*a*ln(x)

Maxima [A]

time = 0.48, size = 53, normalized size = 1.39

$$-4a \arctan(ax) - 2ia \log(a^2x^2 + 1) + 4ia \log(x) - \frac{5a^2x^2 - 4iax + 1}{a^2x^3 + x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^2,x, algorithm="maxima")

[Out] -4*a*arctan(a*x) - 2*I*a*log(a^2*x^2 + 1) + 4*I*a*log(x) - (5*a^2*x^2 - 4*I*a*x + 1)/(a^2*x^3 + x)

Fricas [A]

time = 1.65, size = 60, normalized size = 1.58

$$\frac{5ax + 4(-ia^2x^2 + ax) \log(x) + 4(ia^2x^2 - ax) \log\left(\frac{ax+i}{a}\right) + i}{ax^2 + ix}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^2,x, algorithm="fricas")

[Out] $-(5ax + 4(-Ia^2x^2 + ax)\log(x) + 4(Ia^2x^2 - ax)\log((ax + I)/a + I)/(ax^2 + Ix))$

Sympy [A]

time = 0.16, size = 44, normalized size = 1.16

$$4a(i \log(8a^2x) - i \log(8a^2x + 8ia)) + \frac{-5ax - i}{ax^2 + ix}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2/x**2,x)

[Out] $4a*(I*\log(8a**2*x) - I*\log(8a**2*x + 8*I*a)) + (-5*a*x - I)/(a*x**2 + I*x)$

Giac [A]

time = 0.42, size = 35, normalized size = 0.92

$$-4i a \log(ax + i) + 4i a \log(|x|) - \frac{5ax + i}{ax^2 + ix}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^2,x, algorithm="giac")

[Out] $-4*I*a*\log(ax + I) + 4*I*a*\log(\text{abs}(x)) - (5*a*x + I)/(a*x^2 + I*x)$

Mupad [B]

time = 0.45, size = 37, normalized size = 0.97

$$-8a \operatorname{atan}(2ax + 1i) - \frac{5x + \frac{1i}{a}}{x^2 + \frac{x1i}{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(x^2*(a^2*x^2 + 1)^2),x)

[Out] $-8*a*\operatorname{atan}(2*a*x + 1i) - (5*x + 1i/a)/((x*1i)/a + x^2)$

3.33 $\int \frac{e^{4i \operatorname{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=52

$$-\frac{1}{2x^2} - \frac{4ia}{x} - \frac{4ia^2}{i+ax} - 8a^2 \log(x) + 8a^2 \log(i+ax)$$

[Out] $-1/2/x^2-4*I*a/x-4*I*a^2/(I+a*x)-8*a^2*\ln(x)+8*a^2*\ln(I+a*x)$

Rubi [A]

time = 0.03, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 90}

$$-\frac{4ia^2}{ax+i} - 8a^2 \log(x) + 8a^2 \log(ax+i) - \frac{4ia}{x} - \frac{1}{2x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(4*I)*\text{ArcTan}[a*x]}/x^3, x]$

[Out] $-1/2*1/x^2 - ((4*I)*a)/x - ((4*I)*a^2)/(I + a*x) - 8*a^2*\text{Log}[x] + 8*a^2*\text{Log}[I + a*x]$

Rule 90

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] := \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, p}, x] && IntegersQ[m, n] && (IntegerQ[p] || (GtQ[m, 0] && GeQ[n, -1]))

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] := \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1+iax)^2}{x^3(1-iax)^2} dx \\ &= \int \left(\frac{1}{x^3} + \frac{4ia}{x^2} - \frac{8a^2}{x} + \frac{4ia^3}{(i+ax)^2} + \frac{8a^3}{i+ax} \right) dx \\ &= -\frac{1}{2x^2} - \frac{4ia}{x} - \frac{4ia^2}{i+ax} - 8a^2 \log(x) + 8a^2 \log(i+ax) \end{aligned}$$

Mathematica [A]

time = 0.03, size = 52, normalized size = 1.00

$$-\frac{1}{2x^2} - \frac{4ia}{x} - \frac{4ia^2}{i+ax} - 8a^2 \log(x) + 8a^2 \log(i+ax)$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x])/x^3,x]``[Out] -1/2*1/x^2 - ((4*I)*a)/x - ((4*I)*a^2)/(I + a*x) - 8*a^2*Log[x] + 8*a^2*Log[I + a*x]`**Maple [A]**

time = 0.12, size = 53, normalized size = 1.02

method	result
default	$-4a^3 \left(\frac{i}{a(ax+i)} - \frac{2 \ln(ax+i)}{a} \right) - \frac{1}{2x^2} - \frac{4ia}{x} - 8a^2 \ln(x)$
risch	$-\frac{8ia^2x^2 + \frac{7}{2}ax - \frac{1}{2}i}{(ax+i)x^2} - 8a^2 \ln(x) - 8ia^2 \arctan(ax) + 4a^2 \ln(a^2x^2 + 1)$
meijerg	$\frac{a^2 \left(\frac{3a^2x^2}{3a^2x^2+3} + 2 \ln(a^2x^2+1) - 1 - 4 \ln(x) - 2 \ln(a^2) - \frac{1}{a^2x^2} \right)}{2} + \frac{2ia^3 \left(-\frac{2(3a^2x^2+2)}{x\sqrt{a^2(2a^2x^2+2)}} - \frac{3a \arctan(ax)}{\sqrt{a^2}} \right)}{\sqrt{a^2}} - 3a^2 \left(-\frac{2a^2x^2}{2a^2x^2+2} - \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2/x^3,x,method=_RETURNVERBOSE)``[Out] -4*a^3*(I/a/(I+a*x)-2/a*ln(I+a*x))-1/2/x^2-4*I*a/x-8*a^2*ln(x)`**Maxima [A]**

time = 0.49, size = 69, normalized size = 1.33

$$-8ia^2 \arctan(ax) + 4a^2 \log(a^2x^2 + 1) - 8a^2 \log(x) + \frac{-16ia^3x^3 - 9a^2x^2 - 8iax - 1}{2(a^2x^4 + x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^3,x, algorithm="maxima")``[Out] -8*I*a^2*arctan(a*x) + 4*a^2*log(a^2*x^2 + 1) - 8*a^2*log(x) + 1/2*(-16*I*a^3*x^3 - 9*a^2*x^2 - 8*I*a*x - 1)/(a^2*x^4 + x^2)`**Fricas [A]**

time = 2.05, size = 77, normalized size = 1.48

$$\frac{-16ia^2x^2 + 7ax - 16(a^3x^3 + ia^2x^2) \log(x) + 16(a^3x^3 + ia^2x^2) \log\left(\frac{ax+i}{a}\right) - i}{2(a^3x^3 + ix^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^3,x, algorithm="fricas")

[Out] $\frac{1}{2}*(-16*I*a^2*x^2 + 7*a*x - 16*(a^3*x^3 + I*a^2*x^2)*\log(x) + 16*(a^3*x^3 + I*a^2*x^2)*\log((a*x + I)/a) - I)/(a*x^3 + I*x^2)$

Sympy [A]

time = 0.18, size = 58, normalized size = 1.12

$$8a^2(-\log(16a^3x) + \log(16a^3x + 16ia^2)) + \frac{-16ia^2x^2 + 7ax - i}{2ax^3 + 2ix^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2/x**3,x)

[Out] $8*a**2*(-\log(16*a**3*x) + \log(16*a**3*x + 16*I*a**2)) + (-16*I*a**2*x**2 + 7*a*x - I)/(2*a*x**3 + 2*I*x**2)$

Giac [A]

time = 0.42, size = 46, normalized size = 0.88

$$8a^2 \log(ax + i) - 8a^2 \log(|x|) - \frac{16ia^2x^2 - 7ax + i}{2(ax + i)x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^3,x, algorithm="giac")

[Out] $8*a^2*\log(a*x + I) - 8*a^2*\log(\text{abs}(x)) - 1/2*(16*I*a^2*x^2 - 7*a*x + I)/((a*x + I)*x^2)$

Mupad [B]

time = 0.47, size = 43, normalized size = 0.83

$$-a^2 \operatorname{atan}(2ax + 1i) 16i + \frac{8a^2x^2 + \frac{ax7i}{2} + \frac{1}{2}}{x^2(-1 + ax1i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(x^3*(a^2*x^2 + 1)^2),x)

[Out] $((a*x*7i)/2 + 8*a^2*x^2 + 1/2)/(x^2*(a*x*1i - 1)) - a^2*\operatorname{atan}(2*a*x + 1i)*16i$

3.34 $\int \frac{e^{4i \operatorname{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=62

$$-\frac{1}{3x^3} - \frac{2ia}{x^2} + \frac{8a^2}{x} + \frac{4a^3}{i+ax} - 12ia^3 \log(x) + 12ia^3 \log(i+ax)$$

[Out] $-1/3/x^3 - 2*I*a/x^2 + 8*a^2/x + 4*a^3/(I+a*x) - 12*I*a^3*\ln(x) + 12*I*a^3*\ln(I+a*x)$

Rubi [A]

time = 0.03, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 90}

$$\frac{4a^3}{ax+i} - 12ia^3 \log(x) + 12ia^3 \log(ax+i) + \frac{8a^2}{x} - \frac{2ia}{x^2} - \frac{1}{3x^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((4*I)*\text{ArcTan}[a*x])/x^4}, x]$

[Out] $-1/3*1/x^3 - ((2*I)*a)/x^2 + (8*a^2)/x + (4*a^3)/(I + a*x) - (12*I)*a^3*\text{Log}[x] + (12*I)*a^3*\text{Log}[I + a*x]$

Rule 90

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, p\}, x] \&\& \text{IntegersQ}[m, n] \&\& (\text{IntegerQ}[p] \|\| (\text{GtQ}[m, 0] \&\& \text{GeQ}[n, -1]))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1+iax)^2}{x^4(1-iax)^2} dx \\ &= \int \left(\frac{1}{x^4} + \frac{4ia}{x^3} - \frac{8a^2}{x^2} - \frac{12ia^3}{x} - \frac{4a^4}{(i+ax)^2} + \frac{12ia^4}{i+ax} \right) dx \\ &= -\frac{1}{3x^3} - \frac{2ia}{x^2} + \frac{8a^2}{x} + \frac{4a^3}{i+ax} - 12ia^3 \log(x) + 12ia^3 \log(i+ax) \end{aligned}$$

Mathematica [A]

time = 0.03, size = 62, normalized size = 1.00

$$-\frac{1}{3x^3} - \frac{2ia}{x^2} + \frac{8a^2}{x} + \frac{4a^3}{i+ax} - 12ia^3 \log(x) + 12ia^3 \log(i+ax)$$

Antiderivative was successfully verified.

`[In] Integrate[E^((4*I)*ArcTan[a*x])/x^4,x]`

```
[Out] -1/3*1/x^3 - ((2*I)*a)/x^2 + (8*a^2)/x + (4*a^3)/(I + a*x) - (12*I)*a^3*Log[x] + (12*I)*a^3*Log[I + a*x]
```

Maple [A]

time = 0.10, size = 61, normalized size = 0.98

method	result
default	$4a^4 \left(\frac{1}{a(ax+i)} + \frac{3i \ln(ax+i)}{a} \right) - \frac{1}{3x^3} - 12ia^3 \ln(x) - \frac{2ia}{x^2} + \frac{8a^2}{x}$
risch	$\frac{12a^3x^3+6ia^2x^2+\frac{5}{3}ax-\frac{1}{3}i}{(ax+i)x^3} + 12a^3 \arctan(ax) + 6ia^3 \ln(a^2x^2+1) - 12ia^3 \ln(-x)$
meijerg	$\frac{a^4 \left(-\frac{2(-15a^4x^4-10a^2x^2+2)}{3x^3(a^2)^{\frac{3}{2}}(2a^2x^2+2)} + \frac{5a^3 \arctan(ax)}{(a^2)^{\frac{3}{2}}} \right)}{2\sqrt{a^2}} + 2ia^3 \left(\frac{3a^2x^2}{3a^2x^2+3} + 2 \ln(a^2x^2+1) - 1 - 4 \ln(x) - 2 \ln(a^2) - \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^4/(a^2*x^2+1)^2/x^4,x,method=_RETURNVERBOSE)`

```
[Out] 4*a^4*(1/a/(I+a*x)+3*I*ln(I+a*x)/a)-1/3/x^3-12*I*a^3*ln(x)-2*I*a/x^2+8*a^2/x
```

Maxima [A]

time = 0.48, size = 77, normalized size = 1.24

$$12a^3 \arctan(ax) + 6ia^3 \log(a^2x^2+1) - 12ia^3 \log(x) + \frac{36a^4x^4 - 18ia^3x^3 + 23a^2x^2 - 6iax - 1}{3(a^2x^5 + x^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^4,x, algorithm="maxima")`

```
[Out] 12*a^3*arctan(a*x) + 6*I*a^3*log(a^2*x^2 + 1) - 12*I*a^3*log(x) + 1/3*(36*a^4*x^4 - 18*I*a^3*x^3 + 23*a^2*x^2 - 6*I*a*x - 1)/(a^2*x^5 + x^3)
```

Fricas [A]

time = 3.12, size = 86, normalized size = 1.39

$$\frac{36a^3x^3 + 18ia^2x^2 + 5ax - 36(i a^4x^4 - a^3x^3) \log(x) - 36(-i a^4x^4 + a^3x^3) \log\left(\frac{ax+i}{a}\right) - i}{3(ax^4 + ix^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^4,x, algorithm="fricas")

[Out] 1/3*(36*a^3*x^3 + 18*I*a^2*x^2 + 5*a*x - 36*(I*a^4*x^4 - a^3*x^3)*log(x) - 36*(-I*a^4*x^4 + a^3*x^3)*log((a*x + I)/a) - I)/(a*x^4 + I*x^3)

Sympy [A]

time = 0.20, size = 70, normalized size = 1.13

$$12a^3(-i \log(24a^4x) + i \log(24a^4x + 24ia^3)) + \frac{36a^3x^3 + 18ia^2x^2 + 5ax - i}{3ax^4 + 3ix^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2/x**4,x)

[Out] 12*a**3*(-I*log(24*a**4*x) + I*log(24*a**4*x + 24*I*a**3)) + (36*a**3*x**3 + 18*I*a**2*x**2 + 5*a*x - I)/(3*a*x**4 + 3*I*x**3)

Giac [A]

time = 0.41, size = 54, normalized size = 0.87

$$12i a^3 \log(ax + i) - 12i a^3 \log(|x|) + \frac{36 a^3 x^3 + 18i a^2 x^2 + 5 a x - i}{3(ax + i)x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/x^4,x, algorithm="giac")

[Out] 12*I*a^3*log(a*x + I) - 12*I*a^3*log(abs(x)) + 1/3*(36*a^3*x^3 + 18*I*a^2*x^2 + 5*a*x - I)/((a*x + I)*x^3)

Mupad [B]

time = 0.13, size = 55, normalized size = 0.89

$$24 a^3 \operatorname{atan}(2 a x + 1 i) + \frac{\frac{5 x}{3} + 12 a^2 x^3 + a x^2 6 i - \frac{1 i}{3 a}}{x^4 + \frac{x^3 1 i}{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(x^4*(a^2*x^2 + 1)^2),x)

[Out] 24*a^3*atan(2*a*x + 1i) + ((5*x)/3 + a*x^2*6i - 1i/(3*a) + 12*a^2*x^3)/(x^4 + (x^3*1i)/a)

3.35 $\int e^{-i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=90

$$\frac{x^2\sqrt{1+a^2x^2}}{3a^2} - \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{(16-9iax)\sqrt{1+a^2x^2}}{24a^4} - \frac{3i\sinh^{-1}(ax)}{8a^4}$$

[Out] $-3/8*I*\text{arcsinh}(a*x)/a^4+1/3*x^2*(a^2*x^2+1)^{(1/2)}/a^2-1/4*I*x^3*(a^2*x^2+1)^{(1/2)}/a-1/24*(16-9*I*a*x)*(a^2*x^2+1)^{(1/2)}/a^4$

Rubi [A]

time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5168, 847, 794, 221}

$$-\frac{3i\sinh^{-1}(ax)}{8a^4} + \frac{x^2\sqrt{a^2x^2+1}}{3a^2} - \frac{ix^3\sqrt{a^2x^2+1}}{4a} - \frac{(16-9iax)\sqrt{a^2x^2+1}}{24a^4}$$

Antiderivative was successfully verified.

[In] Int[x^3/E^(I*ArcTan[a*x]),x]

[Out] $(x^2*\text{Sqrt}[1+a^2*x^2])/(3*a^2) - ((I/4)*x^3*\text{Sqrt}[1+a^2*x^2])/a - ((16 - (9*I)*a*x)*\text{Sqrt}[1+a^2*x^2])/(24*a^4) - (((3*I)/8)*\text{ArcSinh}[a*x])/a^4$

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 794

Int[((d_) + (e_)*(x_))*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[((e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x)*((a + c*x^2)^(p + 1)/(2*c*(p + 1)*(2*p + 3))), x] - Dist[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && !LeQ[p, -1]

Rule 847

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[g*(d + e*x)^m*((a + c*x^2)^(p + 1)/(c*(m + 2*p + 2))), x] + Dist[1/(c*(m + 2*p + 2)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^p*Simp[c*d*f*(m + 2*p + 2) - a*e*g*m + c*(e*f*(m + 2*p + 2) + d*g*m)*x, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 + a*e^2, 0] && GtQ[m, 0] && NeQ[m + 2*p + 2, 0] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*p]) && !(IGtQ[m, 0] && EqQ[f, 0])

Rule 5168

`Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(((I*n + 1)/2)/((1 + I*a*x)^(((I*n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]`

Rubi steps

$$\begin{aligned}
 \int e^{-i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1 - iax)}{\sqrt{1 + a^2x^2}} dx \\
 &= -\frac{ix^3\sqrt{1 + a^2x^2}}{4a} + \frac{\int \frac{x^2(3ia + 4a^2x)}{\sqrt{1 + a^2x^2}} dx}{4a^2} \\
 &= \frac{x^2\sqrt{1 + a^2x^2}}{3a^2} - \frac{ix^3\sqrt{1 + a^2x^2}}{4a} + \frac{\int \frac{x(-8a^2 + 9ia^3x)}{\sqrt{1 + a^2x^2}} dx}{12a^4} \\
 &= \frac{x^2\sqrt{1 + a^2x^2}}{3a^2} - \frac{ix^3\sqrt{1 + a^2x^2}}{4a} - \frac{(16 - 9iax)\sqrt{1 + a^2x^2}}{24a^4} - \frac{(3i) \int \frac{1}{\sqrt{1 + a^2x^2}} dx}{8a^3} \\
 &= \frac{x^2\sqrt{1 + a^2x^2}}{3a^2} - \frac{ix^3\sqrt{1 + a^2x^2}}{4a} - \frac{(16 - 9iax)\sqrt{1 + a^2x^2}}{24a^4} - \frac{3i \sinh^{-1}(ax)}{8a^4}
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 56, normalized size = 0.62

$$\frac{\sqrt{1 + a^2x^2}(-16 + 9iax + 8a^2x^2 - 6ia^3x^3) - 9i \sinh^{-1}(ax)}{24a^4}$$

Antiderivative was successfully verified.

[In] `Integrate[x^3/E^(I*ArcTan[a*x]), x]`

[Out] `(Sqrt[1 + a^2*x^2]*(-16 + (9*I)*a*x + 8*a^2*x^2 - (6*I)*a^3*x^3) - (9*I)*ArcSinh[a*x])/(24*a^4)`

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 240 vs. 2(73) = 146.

time = 0.10, size = 241, normalized size = 2.68

method	result
risch	$-\frac{i(6a^3x^3 + 8ia^2x^2 - 9ax - 16i)\sqrt{a^2x^2 + 1}}{24a^4} - \frac{3i \ln\left(\frac{a^2x}{\sqrt{a^2x^2 + 1}} + \sqrt{a^2x^2 + 1}\right)}{8a^3\sqrt{a^2}}$

default	$-i \left(\frac{x(a^2x^2+1)^{\frac{3}{2}}}{4a^2} - \frac{x\sqrt{a^2x^2+1}}{4a^2} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2\sqrt{a^2}} \right) + \frac{(a^2x^2+1)^{\frac{3}{2}}}{3a^4} + i \left(\frac{x\sqrt{a^2x^2+1}}{2} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2\sqrt{a^2}} \right) a^3$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^3/(1+I*a*x)*(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-I/a*(1/4*x*(a^2*x^2+1)^(3/2)/a^2-1/4/a^2*(1/2*x*(a^2*x^2+1)^(1/2)+1/2*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2))+1/3/a^4*(a^2*x^2+1)^(3/2)+I/a^3*(1/2*x*(a^2*x^2+1)^(1/2)+1/2*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2))-1/a^4*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+I*a*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))$$

Maxima [A]

time = 0.50, size = 76, normalized size = 0.84

$$-\frac{i(a^2x^2+1)^{\frac{3}{2}}x}{4a^3} + \frac{5i\sqrt{a^2x^2+1}x}{8a^3} + \frac{(a^2x^2+1)^{\frac{3}{2}}}{3a^4} - \frac{3i \operatorname{arsinh}(ax)}{8a^4} - \frac{\sqrt{a^2x^2+1}}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="maxima")`

[Out]
$$-1/4*I*(a^2*x^2+1)^(3/2)*x/a^3+5/8*I*\sqrt{a^2*x^2+1}*x/a^3+1/3*(a^2*x^2+1)^(3/2)/a^4-3/8*I*\operatorname{arcsinh}(a*x)/a^4-\sqrt{a^2*x^2+1}/a^4$$

Fricas [A]

time = 2.39, size = 59, normalized size = 0.66

$$\frac{(-6i a^3 x^3 + 8 a^2 x^2 + 9i a x - 16) \sqrt{a^2 x^2 + 1} + 9i \log(-a x + \sqrt{a^2 x^2 + 1})}{24 a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="fricas")`

[Out]
$$1/24*((-6*I*a^3*x^3+8*a^2*x^2+9*I*a*x-16)*\sqrt{a^2*x^2+1}+9*I*\log(-a*x+\sqrt{a^2*x^2+1}))/a^4$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^3 \sqrt{a^2 x^2 + 1}}{a x - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3/(1+I*a*x)*(a**2*x**2+1)**(1/2),x)

[Out] -I*Integral(x**3*sqrt(a**2*x**2 + 1)/(a*x - I), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.06, size = 85, normalized size = 0.94

$$\frac{\operatorname{asinh}\left(x\sqrt{a^2}\right) 3i}{8a^3\sqrt{a^2}} - \frac{\sqrt{a^2x^2+1}\left(\frac{2}{3(a^2)^{3/2}} - \frac{a^2x^2}{3(a^2)^{3/2}} + \frac{x^3(a^2)^{3/2}i}{4a^3} - \frac{x\sqrt{a^2}3i}{8a^3}\right)}{\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a^2*x^2 + 1)^(1/2))/(a*x*1i + 1),x)

[Out] - (asinh(x*(a^2)^(1/2))*3i)/(8*a^3*(a^2)^(1/2)) - ((a^2*x^2 + 1)^(1/2)*(2/(
3*(a^2)^(3/2)) - (a^2*x^2)/(3*(a^2)^(3/2)) + (x^3*(a^2)^(3/2)*1i)/(4*a^3) -
(x*(a^2)^(1/2)*3i)/(8*a^3)))/(a^2)^(1/2)

3.36 $\int e^{-i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=75

$$\frac{i\sqrt{1+a^2x^2}}{a^3} + \frac{x\sqrt{1+a^2x^2}}{2a^2} - \frac{i(1+a^2x^2)^{3/2}}{3a^3} - \frac{\sinh^{-1}(ax)}{2a^3}$$

[Out] $-1/3*I*(a^2*x^2+1)^{(3/2)}/a^3-1/2*\text{arcsinh}(a*x)/a^3+I*(a^2*x^2+1)^{(1/2)}/a^3+1/2*x*(a^2*x^2+1)^{(1/2)}/a^2$

Rubi [A]

time = 0.03, antiderivative size = 75, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5168, 811, 655, 201, 221}

$$-\frac{\sinh^{-1}(ax)}{2a^3} + \frac{x\sqrt{a^2x^2+1}}{2a^2} - \frac{i(a^2x^2+1)^{3/2}}{3a^3} + \frac{i\sqrt{a^2x^2+1}}{a^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{(I*\text{ArcTan}[a*x])}, x]$

[Out] $(I*\text{Sqrt}[1+a^2*x^2])/a^3 + (x*\text{Sqrt}[1+a^2*x^2])/(2*a^2) - ((I/3)*(1+a^2*x^2)^{(3/2)})/a^3 - \text{ArcSinh}[a*x]/(2*a^3)$

Rule 201

$\text{Int}[(a_+ + (b_+)*(x_+)^{n_+})^{p_+}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^p/(n*p + 1)), x] + \text{Dist}[a*n*(p/(n*p + 1)), \text{Int}[(a + b*x^n)^{p-1}, x], x] /;$ FreeQ[{a, b}, x] && IGtQ[n, 0] && GtQ[p, 0] && (IntegerQ[2*p] || (EqQ[n, 2] && IntegerQ[4*p]) || (EqQ[n, 2] && IntegerQ[3*p]) || LtQ[Denominator[p + 1/n], Denominator[p]])

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_+ + (b_+)*(x_+)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /;$ FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 655

$\text{Int}[(d_+ + (e_+)*(x_+))*((a_+ + (c_+)*(x_+)^2)^{p_+}), x_Symbol] \rightarrow \text{Simp}[e*((a + c*x^2)^{p+1}/(2*c*(p+1))), x] + \text{Dist}[d, \text{Int}[(a + c*x^2)^p, x], x] /;$ FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]

Rule 811

$\text{Int}[(x_+)^2*((f_+ + (g_+)*(x_+))*((a_+ + (c_+)*(x_+)^2)^{p_+}), x_Symbol] \rightarrow \text{Dist}[1/c, \text{Int}[(f + g*x)*(a + c*x^2)^{p+1}, x], x] - \text{Dist}[a/c, \text{Int}[(f + g*x)*$

$(a + c*x^2)^p, x] /; \text{FreeQ}\{a, c, f, g, p\}, x] \ \&\& \ \text{EqQ}[a*g^2 + f^2*c, 0]$

Rule 5168

$\text{Int}[E^{(\text{ArcTan}[a_*](x_*)^{(n_*)})}(x_*)^{(m_*)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{((I*n + 1)/2)} / ((1 + I*a*x)^{((I*n - 1)/2)} * \text{Sqrt}[1 + a^2*x^2])), x] /; \text{FreeQ}\{a, m\}, x] \ \&\& \ \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{-i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1 - iax)}{\sqrt{1 + a^2x^2}} dx \\ &= -\frac{\int \frac{1 - iax}{\sqrt{1 + a^2x^2}} dx}{a^2} + \frac{\int (1 - iax)\sqrt{1 + a^2x^2} dx}{a^2} \\ &= \frac{i\sqrt{1 + a^2x^2}}{a^3} - \frac{i(1 + a^2x^2)^{3/2}}{3a^3} - \frac{\int \frac{1}{\sqrt{1 + a^2x^2}} dx}{a^2} + \frac{\int \sqrt{1 + a^2x^2} dx}{a^2} \\ &= \frac{i\sqrt{1 + a^2x^2}}{a^3} + \frac{x\sqrt{1 + a^2x^2}}{2a^2} - \frac{i(1 + a^2x^2)^{3/2}}{3a^3} - \frac{\sinh^{-1}(ax)}{a^3} + \frac{\int \frac{1}{\sqrt{1 + a^2x^2}} dx}{2a^2} \\ &= \frac{i\sqrt{1 + a^2x^2}}{a^3} + \frac{x\sqrt{1 + a^2x^2}}{2a^2} - \frac{i(1 + a^2x^2)^{3/2}}{3a^3} - \frac{\sinh^{-1}(ax)}{2a^3} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 46, normalized size = 0.61

$$\frac{(4i + 3ax - 2ia^2x^2)\sqrt{1 + a^2x^2} - 3\sinh^{-1}(ax)}{6a^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^(I*ArcTan[a*x]),x]

[Out] ((4*I + 3*a*x - (2*I)*a^2*x^2)*Sqrt[1 + a^2*x^2] - 3*ArcSinh[a*x])/(6*a^3)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 166 vs. $2(61) = 122$.

time = 0.08, size = 167, normalized size = 2.23

method	result
risch	$-\frac{i(2a^2x^2 + 3iax - 4)\sqrt{a^2x^2 + 1}}{6a^3} - \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2 + 1}\right)}{2a^2\sqrt{a^2}}$

default	$-\frac{i(a^2x^2+1)^{\frac{3}{2}}}{3a^3} + \frac{x\sqrt{a^2x^2+1}}{2} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{a^2} + \frac{i\left(\sqrt{\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia+(x-\frac{i}{a})}{\sqrt{\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)}}\right)}{a^3}\right)}{a^3}$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-1/3*I*(a^2*x^2+1)^{(3/2)}/a^3+1/a^2*(1/2*x*(a^2*x^2+1)^{(1/2)}+1/2*\ln(a^2*x/(a^2)^{(1/2)}+(a^2*x^2+1)^{(1/2)))/(a^2)^{(1/2)}+I/a^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^{(1/2)}+I*a*\ln((I*a+(x-I/a)*a^2)/(a^2)^{(1/2)}+((x-I/a)^2*a^2+2*I*a*(x-I/a))^{(1/2)))/(a^2)^{(1/2)}$$

Maxima [A]

time = 0.48, size = 59, normalized size = 0.79

$$\frac{\sqrt{a^2x^2+1}x}{2a^2} - \frac{i(a^2x^2+1)^{\frac{3}{2}}}{3a^3} - \frac{\operatorname{arsinh}(ax)}{2a^3} + \frac{i\sqrt{a^2x^2+1}}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="maxima")`

[Out]
$$1/2*\sqrt{a^2*x^2+1}*x/a^2 - 1/3*I*(a^2*x^2+1)^{(3/2)}/a^3 - 1/2*\operatorname{arcsinh}(a*x)/a^3 + I*\sqrt{a^2*x^2+1}/a^3$$

Fricas [A]

time = 2.05, size = 51, normalized size = 0.68

$$\frac{\sqrt{a^2x^2+1}(-2ia^2x^2+3ax+4i)+3\log(-ax+\sqrt{a^2x^2+1})}{6a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="fricas")`

[Out]
$$1/6*(\sqrt{a^2*x^2+1}*(-2*I*a^2*x^2+3*a*x+4*I)+3*\log(-a*x+\sqrt{a^2*x^2+1}))/a^3$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^2 \sqrt{a^2x^2+1}}{ax-i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*a*x)*(a**2*x**2+1)**(1/2),x)

[Out] -I*Integral(x**2*sqrt(a**2*x**2 + 1)/(a*x - I), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.42, size = 71, normalized size = 0.95

$$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{x \sqrt{a^2}}{2a^2} + \frac{a 2i}{3(a^2)^{3/2}} - \frac{a^3 x^2 1i}{3(a^2)^{3/2}} \right)}{\sqrt{a^2}} - \frac{\operatorname{asinh}\left(x \sqrt{a^2}\right)}{2 a^2 \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a^2*x^2 + 1)^(1/2))/(a*x*1i + 1),x)

[Out] ((a^2*x^2 + 1)^(1/2)*((a*2i)/(3*(a^2)^(3/2)) - (a^3*x^2*1i)/(3*(a^2)^(3/2))
+ (x*(a^2)^(1/2))/(2*a^2)))/(a^2)^(1/2) - asinh(x*(a^2)^(1/2))/(2*a^2*(a^2
)^(1/2))

3.37 $\int e^{-i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=42

$$\frac{(2 - iax)\sqrt{1 + a^2x^2}}{2a^2} + \frac{i \sinh^{-1}(ax)}{2a^2}$$

[Out] $1/2*I*\text{arcsinh}(a*x)/a^2 + 1/2*(2 - I*a*x)*(a^2*x^2 + 1)^{(1/2)}/a^2$

Rubi [A]

time = 0.01, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5168, 794, 221}

$$\frac{\sqrt{a^2x^2 + 1} (2 - iax)}{2a^2} + \frac{i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

[In] Int[x/E^(I*ArcTan[a*x]),x]

[Out] $((2 - I*a*x)*\text{Sqrt}[1 + a^2*x^2])/(2*a^2) + ((I/2)*\text{ArcSinh}[a*x])/a^2$

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 794

Int[((d_) + (e_)*(x_))*((f_) + (g_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] :> Simp[((e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x)*((a + c*x^2)^(p + 1)/(2*c*(p + 1)*(2*p + 3))), x] - Dist[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && !LeQ[p, -1]

Rule 5168

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] :> Int[x^m*((1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-i \tan^{-1}(ax)} x dx &= \int \frac{x(1 - iax)}{\sqrt{1 + a^2 x^2}} dx \\ &= \frac{(2 - iax)\sqrt{1 + a^2 x^2}}{2a^2} + \frac{i \int \frac{1}{\sqrt{1 + a^2 x^2}} dx}{2a} \\ &= \frac{(2 - iax)\sqrt{1 + a^2 x^2}}{2a^2} + \frac{i \sinh^{-1}(ax)}{2a^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 38, normalized size = 0.90

$$\frac{(2 - iax)\sqrt{1 + a^2 x^2} + i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

`[In] Integrate[x/E^(I*ArcTan[a*x]),x]``[Out] ((2 - I*a*x)*Sqrt[1 + a^2*x^2] + I*ArcSinh[a*x])/(2*a^2)`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 149 vs. $2(34) = 68$.

time = 0.08, size = 150, normalized size = 3.57

method	result
risch	$-\frac{i(ax+2i)\sqrt{a^2x^2+1}}{2a^2} + \frac{i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a\sqrt{a^2}}$
default	$-\frac{i \left(\frac{x\sqrt{a^2x^2+1}}{2} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2\sqrt{a^2}} \right)}{a} + \frac{\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2}{\sqrt{a^2}} + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}\right)}{a^2}}{a^2}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x/(1+I*a*x)*(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`
`[Out] -I/a*(1/2*x*(a^2*x^2+1)^(1/2)+1/2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2))+1/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+I*a*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))`
Maxima [A]

time = 0.48, size = 42, normalized size = 1.00

$$-\frac{i \sqrt{a^2 x^2 + 1} x}{2a} + \frac{i \operatorname{arsinh}(ax)}{2a^2} + \frac{\sqrt{a^2 x^2 + 1}}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="maxima")

[Out] $-1/2*I*\sqrt{a^2*x^2 + 1}*x/a + 1/2*I*\operatorname{arcsinh}(a*x)/a^2 + \sqrt{a^2*x^2 + 1}/a^2$

Fricas [A]

time = 2.30, size = 43, normalized size = 1.02

$$\frac{\sqrt{a^2x^2 + 1}(-iax + 2) - i \log(-ax + \sqrt{a^2x^2 + 1})}{2a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="fricas")

[Out] $1/2*(\sqrt{a^2*x^2 + 1}*(-I*a*x + 2) - I*\log(-a*x + \sqrt{a^2*x^2 + 1}))/a^2$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x\sqrt{a^2x^2 + 1}}{ax - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*a*x)*(a**2*x**2+1)**(1/2),x)

[Out] $-I*\operatorname{Integral}(x*\sqrt{a**2*x**2 + 1}/(a*x - I), x)$

Giac [A]

time = 0.42, size = 53, normalized size = 1.26

$$-\frac{1}{2}\sqrt{a^2x^2 + 1}\left(\frac{ix}{a} - \frac{2}{a^2}\right) - \frac{i \log(-x|a| + \sqrt{a^2x^2 + 1})}{2a|a|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] $-1/2*\sqrt{a^2*x^2 + 1}*(I*x/a - 2/a^2) - 1/2*I*\log(-x*\operatorname{abs}(a) + \sqrt{a^2*x^2 + 1})/(a*\operatorname{abs}(a))$

Mupad [B]

time = 0.40, size = 51, normalized size = 1.21

$$\frac{\left(\frac{1}{\sqrt{a^2}} - \frac{x\sqrt{a^2}}{2a}\right)\sqrt{a^2x^2 + 1} + \frac{\operatorname{asinh}\left(x\sqrt{a^2}\right)}{2a}}{\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x*(a^2*x^2 + 1)^(1/2))/(a*x+1),x)
```

```
[Out] ((1/(a^2)^(1/2) - (x*(a^2)^(1/2)*1)/(2*a))*(a^2*x^2 + 1)^(1/2) + (asinh(x*(a^2)^(1/2))*1)/(2*a))/(a^2)^(1/2)
```

3.38 $\int e^{-i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=29

$$-\frac{i\sqrt{1+a^2x^2}}{a} + \frac{\sinh^{-1}(ax)}{a}$$

[Out] arcsinh(a*x)/a-I*(a^2*x^2+1)^(1/2)/a

Rubi [A]

time = 0.01, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5167, 655, 221}

$$\frac{\sinh^{-1}(ax)}{a} - \frac{i\sqrt{a^2x^2+1}}{a}$$

Antiderivative was successfully verified.

[In] Int[E^((-I)*ArcTan[a*x]),x]

[Out] ((-I)*Sqrt[1 + a^2*x^2])/a + ArcSinh[a*x]/a

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 655

Int[((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] :> Simp[e*((a + c*x^2)^(p + 1)/(2*c*(p + 1))), x] + Dist[d, Int[(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]

Rule 5167

Int[E^(ArcTan[(a_)*(x_)])*(n_), x_Symbol] :> Int[(1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[a, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-i \tan^{-1}(ax)} dx &= \int \frac{1 - iax}{\sqrt{1 + a^2x^2}} dx \\ &= -\frac{i\sqrt{1 + a^2x^2}}{a} + \int \frac{1}{\sqrt{1 + a^2x^2}} dx \\ &= -\frac{i\sqrt{1 + a^2x^2}}{a} + \frac{\sinh^{-1}(ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 26, normalized size = 0.90

$$\frac{-i\sqrt{1+a^2x^2} + \sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((-I)*ArcTan[a*x]),x]``[Out] ((-I)*Sqrt[1 + a^2*x^2] + ArcSinh[a*x])/a`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(26) = 52$.

time = 0.07, size = 100, normalized size = 3.45

method	result	size
risch	$-\frac{i\sqrt{a^2x^2+1}}{a} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{\sqrt{a^2}}$	48
default	$i \left(\frac{\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2}{\sqrt{a^2}} + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}\right)}{\sqrt{a^2}} \right)$	100

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)`
`[Out] -I/a*(((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+I*a*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)))/(a^2)^(1/2))`
Maxima [A]

time = 0.47, size = 25, normalized size = 0.86

$$\frac{\operatorname{arsinh}(ax)}{a} - \frac{i\sqrt{a^2x^2+1}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="maxima")``[Out] arcsinh(a*x)/a - I*sqrt(a^2*x^2 + 1)/a`**Fricas [A]**

time = 2.26, size = 37, normalized size = 1.28

$$\frac{-i\sqrt{a^2x^2+1} - \log\left(-ax + \sqrt{a^2x^2+1}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="fricas")

[Out] (-I*sqrt(a^2*x^2 + 1) - log(-a*x + sqrt(a^2*x^2 + 1)))/a

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 x^2 + 1}}{ax - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2),x)

[Out] -I*Integral(sqrt(a**2*x**2 + 1)/(a*x - I), x)

Giac [A]

time = 0.41, size = 41, normalized size = 1.41

$$\frac{\log\left(-x|a| + \sqrt{a^2 x^2 + 1}\right)}{|a|} - \frac{i \sqrt{a^2 x^2 + 1}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] -log(-x*abs(a) + sqrt(a^2*x^2 + 1))/abs(a) - I*sqrt(a^2*x^2 + 1)/a

Mupad [B]

time = 0.40, size = 32, normalized size = 1.10

$$\frac{\operatorname{asinh}\left(x \sqrt{a^2}\right)}{\sqrt{a^2}} - \frac{\sqrt{a^2 x^2 + 1} \operatorname{li}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(1/2)/(a*x*1i + 1),x)

[Out] asinh(x*(a^2)^(1/2))/(a^2)^(1/2) - ((a^2*x^2 + 1)^(1/2)*1i)/a

$$3.39 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=25

$$-i \sinh^{-1}(ax) - \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] -I*arcsinh(a*x)-arctanh((a^2*x^2+1)^(1/2))

Rubi [A]

time = 0.03, antiderivative size = 25, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 858, 221, 272, 65, 214}

$$-\tanh^{-1}\left(\sqrt{a^2x^2+1}\right) - i \sinh^{-1}(ax)$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a*x])*x),x]

[Out] (-I)*ArcSinh[a*x] - ArcTanh[Sqrt[1 + a^2*x^2]]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 858


```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Dist[g/e, Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] + D
ist[(e*f - d*g)/e, Int[(d + e*x)^m*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d,
e, f, g, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && !IGtQ[m, 0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(ax)}}{x} dx &= \int \frac{1 - iax}{x\sqrt{1 + a^2x^2}} dx \\
&= -\left((ia) \int \frac{1}{\sqrt{1 + a^2x^2}} dx \right) + \int \frac{1}{x\sqrt{1 + a^2x^2}} dx \\
&= -i \sinh^{-1}(ax) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
&= -i \sinh^{-1}(ax) + \frac{\text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2x^2} \right)}{a^2} \\
&= -i \sinh^{-1}(ax) - \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 29, normalized size = 1.16

$$-i \sinh^{-1}(ax) + \log(x) - \log \left(1 + \sqrt{1 + a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[1/(E^(I*ArcTan[a*x])*x), x]
```

```
[Out] (-I)*ArcSinh[a*x] + Log[x] - Log[1 + Sqrt[1 + a^2*x^2]]
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 120 vs. $2(22) = 44$.

time = 0.07, size = 121, normalized size = 4.84

method	result
--------	--------

default	$-\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia\left(x - \frac{i}{a}\right)} - \frac{ia \ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2 + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia\left(x - \frac{i}{a}\right)}}{\sqrt{a^2}}\right)}{\sqrt{a^2}} + \sqrt{a^2 x^2 + 1}$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x,x,method=_RETURNVERBOSE)`

[Out] $-\left(\left(x - \frac{i}{a}\right)^2 a^2 + 2Ia\left(x - \frac{i}{a}\right)\right)^{1/2} - Ia \ln\left(\frac{Ia + \left(x - \frac{i}{a}\right)a^2}{\left(x - \frac{i}{a}\right)^2 a^2 + 2Ia\left(x - \frac{i}{a}\right)}\right) / \left(\left(x - \frac{i}{a}\right)^2 a^2 + 2Ia\left(x - \frac{i}{a}\right)\right)^{1/2} + \left(\left(x - \frac{i}{a}\right)^2 a^2 + 2Ia\left(x - \frac{i}{a}\right)\right)^{1/2} - \operatorname{arctanh}\left(\frac{1}{\left(x - \frac{i}{a}\right)^2 a^2 + 2Ia\left(x - \frac{i}{a}\right)}\right)$

Maxima [A]

time = 0.48, size = 26, normalized size = 1.04

$$-i a \left(\frac{\operatorname{arsinh}(ax)}{a} - \frac{i \operatorname{arsinh}\left(\frac{1}{a|x|}\right)}{a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x,x, algorithm="maxima")`

[Out] $-Ia * (\operatorname{arcsinh}(ax)/a - I * \operatorname{arcsinh}(1/(a*|x|)))/a$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 58 vs. $2(21) = 42$.

time = 1.08, size = 58, normalized size = 2.32

$$-\log\left(-ax + \sqrt{a^2 x^2 + 1} + 1\right) + i \log\left(-ax + \sqrt{a^2 x^2 + 1}\right) + \log\left(-ax + \sqrt{a^2 x^2 + 1} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x,x, algorithm="fricas")`

[Out] $-\log(-ax + \sqrt{a^2 x^2 + 1} + 1) + I * \log(-ax + \sqrt{a^2 x^2 + 1}) + \log(-ax + \sqrt{a^2 x^2 + 1} - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 x^2 + 1}}{ax^2 - ix} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/x,x)`

[Out] $-I \cdot \text{Integral}(\sqrt{a^2 x^2 + 1} / (a x^2 - I x), x)$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 68 vs. $2(21) = 42$.

time = 0.41, size = 68, normalized size = 2.72

$$\frac{i a \log\left(-x|a| + \sqrt{a^2 x^2 + 1}\right)}{|a|} - \log\left(\left|-x|a| + \sqrt{a^2 x^2 + 1} + 1\right|\right) + \log\left(\left|-x|a| + \sqrt{a^2 x^2 + 1} - 1\right|\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x,x, algorithm="giac")`

[Out] $I a \log(-x \operatorname{abs}(a) + \sqrt{a^2 x^2 + 1}) / \operatorname{abs}(a) - \log(\operatorname{abs}(-x \operatorname{abs}(a) + \sqrt{a^2 x^2 + 1}) + 1) + \log(\operatorname{abs}(-x \operatorname{abs}(a) + \sqrt{a^2 x^2 + 1}) - 1)$

Mupad [B]

time = 0.04, size = 32, normalized size = 1.28

$$-\operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right) - \frac{a \operatorname{asinh}\left(x \sqrt{a^2}\right) i}{\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^2*x^2 + 1)^(1/2)/(x*(a*x*1i + 1)),x)`

[Out] $-\operatorname{atanh}((a^2 x^2 + 1)^{1/2}) - (a \operatorname{asinh}(x (a^2)^{1/2}) * 1i) / (a^2)^{1/2}$

$$3.40 \quad \int \frac{e^{-i\text{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=38

$$-\frac{\sqrt{1+a^2x^2}}{x} + ia \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] I*a*arctanh((a^2*x^2+1)^(1/2))-(a^2*x^2+1)^(1/2)/x

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5168, 821, 272, 65, 214}

$$-\frac{\sqrt{a^2x^2+1}}{x} + ia \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a*x])*x^2),x]

[Out] -(Sqrt[1 + a^2*x^2]/x) + I*a*ArcTanh[Sqrt[1 + a^2*x^2]]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 821

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(-(e*f - d*g))*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1))/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2), Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,

p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 5168

Int[E^(ArcTan[a_]*(x_))*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{1 - iax}{x^2 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} - (ia) \int \frac{1}{x \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} - \frac{1}{2}(ia) \text{Subst} \left(\int \frac{1}{x \sqrt{1 + a^2 x}} dx, x, x^2 \right) \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} - \frac{i \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2 x^2} \right)}{a} \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{x} + ia \tanh^{-1} \left(\sqrt{1 + a^2 x^2} \right)
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 47, normalized size = 1.24

$$-\frac{\sqrt{1 + a^2 x^2}}{x} - ia \log(x) + ia \log \left(1 + \sqrt{1 + a^2 x^2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^(I*ArcTan[a*x])*x^2), x]

[Out] -(Sqrt[1 + a^2*x^2]/x) - I*a*Log[x] + I*a*Log[1 + Sqrt[1 + a^2*x^2]]

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 194 vs. 2(33) = 66.

time = 0.07, size = 195, normalized size = 5.13

method	result
risch	$-\frac{\sqrt{a^2 x^2 + 1}}{x} + ia \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2 x^2 + 1}} \right)$

default	$ia \left(\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)} + \frac{ia \ln \left(\frac{ia + \left(x - \frac{i}{a}\right)a^2 + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}}{\sqrt{a^2}} \right)}{\sqrt{a^2}} \right) - ia \left(\sqrt{a^2 x^2 + 1} \right)$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

[Out] $I*a*((x-I/a)^2*a^2+2*I*a*(x-I/a))^{(1/2)}+I*a*\ln((I*a+(x-I/a)*a^2)/(a^2)^{(1/2)}+(x-I/a)^2*a^2+2*I*a*(x-I/a))^{(1/2)})/(a^2)^{(1/2)}-I*a*((a^2*x^2+1)^{(1/2)}-\arctanh(1/(a^2*x^2+1)^{(1/2)}))-1/x*(a^2*x^2+1)^{(3/2)}+2*a^2*(1/2*x*(a^2*x^2+1)^{(1/2)}+1/2*\ln(a^2*x/(a^2)^{(1/2)}+(a^2*x^2+1)^{(1/2)}))/(a^2)^{(1/2)}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^2,x, algorithm="maxima")`

[Out] `integrate(sqrt(a^2*x^2 + 1)/((I*a*x + 1)*x^2), x)`

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 66 vs. 2(32) = 64.

time = 1.92, size = 66, normalized size = 1.74

$$\frac{iax \log(-ax + \sqrt{a^2x^2 + 1} + 1) - iax \log(-ax + \sqrt{a^2x^2 + 1} - 1) - ax - \sqrt{a^2x^2 + 1}}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^2,x, algorithm="fricas")`

[Out] $(I*a*x*\log(-a*x + \sqrt{a^2*x^2 + 1} + 1) - I*a*x*\log(-a*x + \sqrt{a^2*x^2 + 1} - 1) - a*x - \sqrt{a^2*x^2 + 1})/x$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2x^2 + 1}}{ax^3 - ix^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/x**2,x)`

[Out] $-I \cdot \text{Integral}(\sqrt{a^2 x^2 + 1} / (a x^3 - I x^2), x)$

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^2,x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
 dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.04, size = 33, normalized size = 0.87

$$-\frac{\sqrt{a^2 x^2 + 1}}{x} + a \operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right) \operatorname{li}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^2*x^2 + 1)^(1/2)/(x^2*(a*x*1i + 1)),x)`

[Out] $a \operatorname{atanh}((a^2 x^2 + 1)^{1/2}) * 1i - (a^2 x^2 + 1)^{1/2} / x$

3.41 $\int \frac{e^{-i\text{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=63

$$-\frac{\sqrt{1+a^2x^2}}{2x^2} + \frac{ia\sqrt{1+a^2x^2}}{x} + \frac{1}{2}a^2 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $1/2*a^2*\text{arctanh}((a^2*x^2+1)^{(1/2)})-1/2*(a^2*x^2+1)^{(1/2)}/x^2+I*a*(a^2*x^2+1)^{(1/2)}/x$

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 849, 821, 272, 65, 214}

$$\frac{ia\sqrt{a^2x^2+1}}{x} - \frac{\sqrt{a^2x^2+1}}{2x^2} + \frac{1}{2}a^2 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(I*ArcTan[a*x])*x^3),x]`

[Out] $-1/2*\text{Sqrt}[1+a^2*x^2]/x^2+(I*a*\text{Sqrt}[1+a^2*x^2])/x+(a^2*\text{ArcTanh}[\text{Sqrt}[1+a^2*x^2]])/2$

Rule 65

`Int[((a_.)+(b_.)*(x_))^(m_)*((c_.)+(d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^n, x], x, (a+b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c-a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 214

`Int[((a_.)+(b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 272

`Int[(x_)^(m_.)*((a_.)+(b_.)*(x_)^(n_))^(p_), x_Symbol] :> Dist[1/n, Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m+1)/n]]`

Rule 821

`Int[((d_.)+(e_.)*(x_))^(m_)*((f_.)+(g_.)*(x_))*((a_.)+(c_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(-(e*f-d*g))*(d+e*x)^(m+1)*((a+c*x^2)^(p+1`

)/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2),
Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,
p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]

Rule 849

Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/(
(m + 1)*(c*d^2 + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 + a*e^2)), Int[(d +
e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
+ 2*p + 3)*x, x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 +
a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*
p])

Rule 5168

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{1 - iax}{x^3 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} - \frac{1}{2} \int \frac{2ia + a^2 x}{x^2 \sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} + \frac{ia\sqrt{1 + a^2 x^2}}{x} - \frac{1}{2} a^2 \int \frac{1}{x\sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} + \frac{ia\sqrt{1 + a^2 x^2}}{x} - \frac{1}{4} a^2 \text{Subst}\left(\int \frac{1}{x\sqrt{1 + a^2 x}} dx, x, x^2\right) \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} + \frac{ia\sqrt{1 + a^2 x^2}}{x} - \frac{1}{2} \text{Subst}\left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2 x^2}\right) \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{2x^2} + \frac{ia\sqrt{1 + a^2 x^2}}{x} + \frac{1}{2} a^2 \tanh^{-1}\left(\sqrt{1 + a^2 x^2}\right)
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 57, normalized size = 0.90

$$\frac{1}{2} \left(\frac{(-1 + 2iax)\sqrt{1 + a^2 x^2}}{x^2} - a^2 \log(x) + a^2 \log\left(1 + \sqrt{1 + a^2 x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^(I*ArcTan[a*x])*x^3),x]

[Out] (((-1 + (2*I)*a*x)*Sqrt[1 + a^2*x^2])/x^2 - a^2*Log[x] + a^2*Log[1 + Sqrt[1 + a^2*x^2]])/2

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 216 vs. 2(52) = 104.
time = 0.07, size = 217, normalized size = 3.44

method	result
risch	$\frac{i(2a^3x^3+ia^2x^2+2ax+i)}{2x^2\sqrt{a^2x^2+1}} + \frac{a^2 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{2}$
default	$a^2 \left(\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2}{\sqrt{a^2}} + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}\right)}{\sqrt{a^2}} \right) - \frac{(a^2x^2+1)^{\frac{3}{2}}}{2x^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^3,x,method=_RETURNVERBOSE)

[Out] a^2*(((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+I*a*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)))/(a^2)^(1/2))-1/2/x^2*(a^2*x^2+1)^(3/2)-1/2*a^2*((a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2)))-I*a*(-1/x*(a^2*x^2+1)^(3/2)+2*a^2*(1/2*x*(a^2*x^2+1)^(1/2)+1/2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2)))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^3,x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^2 + 1)/((I*a*x + 1)*x^3), x)

Fricas [A]

time = 1.57, size = 83, normalized size = 1.32

$$\frac{a^2x^2 \log\left(-ax + \sqrt{a^2x^2+1} + 1\right) - a^2x^2 \log\left(-ax + \sqrt{a^2x^2+1} - 1\right) + 2ia^2x^2 + \sqrt{a^2x^2+1} (2iax - 1)}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^3,x, algorithm="fricas")

[Out] 1/2*(a^2*x^2*log(-a*x + sqrt(a^2*x^2 + 1) + 1) - a^2*x^2*log(-a*x + sqrt(a^2*x^2 + 1) - 1) + 2*I*a^2*x^2 + sqrt(a^2*x^2 + 1)*(2*I*a*x - 1))/x^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 x^2 + 1}}{a x^4 - i x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/x**3,x)

[Out] -I*Integral(sqrt(a**2*x**2 + 1)/(a*x**4 - I*x**3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^3,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.04, size = 52, normalized size = 0.83

$$\frac{a^2 \operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right)}{2} - \frac{\sqrt{a^2 x^2 + 1}}{2 x^2} + \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(1/2)/(x^3*(a*x*1i + 1)),x)

[Out] (a^2*atanh((a^2*x^2 + 1)^(1/2)))/2 - (a^2*x^2 + 1)^(1/2)/(2*x^2) + (a*(a^2*x^2 + 1)^(1/2)*1i)/x

$$3.42 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)}}{x^4} dx$$

Optimal. Leaf size=90

$$-\frac{\sqrt{1+a^2x^2}}{3x^3} + \frac{ia\sqrt{1+a^2x^2}}{2x^2} + \frac{2a^2\sqrt{1+a^2x^2}}{3x} - \frac{1}{2}ia^3 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-1/2*I*a^3*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/3*(a^2*x^2+1)^{(1/2)}/x^3+1/2*I*a*(a^2*x^2+1)^{(1/2)}/x^2+2/3*a^2*(a^2*x^2+1)^{(1/2)}/x$

Rubi [A]

time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 849, 821, 272, 65, 214}

$$\frac{2a^2\sqrt{a^2x^2+1}}{3x} + \frac{ia\sqrt{a^2x^2+1}}{2x^2} - \frac{\sqrt{a^2x^2+1}}{3x^3} - \frac{1}{2}ia^3 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(I*ArcTan[a*x])*x^4),x]`

[Out] $-1/3*\operatorname{Sqrt}[1+a^2*x^2]/x^3 + ((I/2)*a*\operatorname{Sqrt}[1+a^2*x^2])/x^2 + (2*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(3*x) - (I/2)*a^3*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]]$

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^n, x], x, (a+b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c-a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 272

`Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m+1)/n]]`

Rule 821

`Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol] := Simp[(-(e*f-d*g))*(d+e*x)^(m+1)*((a+c*x^2)^(p+1`

```
)/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2),
Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,
p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]
```

Rule 849

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/((
m + 1)*(c*d^2 + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 + a*e^2)), Int[(d +
e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
+ 2*p + 3)*x, x], x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 +
a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*
p])
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{1 - iax}{x^4 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} - \frac{1}{3} \int \frac{3ia + 2a^2 x}{x^3 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} + \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{1}{6} \int \frac{-4a^2 + 3ia^3 x}{x^2 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} + \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} + \frac{1}{2} (ia^3) \int \frac{1}{x \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} + \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} + \frac{1}{4} (ia^3) \text{Subst} \left(\int \frac{1}{x \sqrt{1 + a^2 x^2}} dx, x, \sqrt{1 + a^2 x^2} \right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} + \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} + \frac{1}{2} (ia) \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2 x^2} \right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{3x^3} + \frac{ia\sqrt{1 + a^2 x^2}}{2x^2} + \frac{2a^2 \sqrt{1 + a^2 x^2}}{3x} - \frac{1}{2} ia^3 \tanh^{-1} \left(\sqrt{1 + a^2 x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 70, normalized size = 0.78

$$\frac{1}{6} \left(\frac{\sqrt{1+a^2x^2}(-2+3iax+4a^2x^2)}{x^3} + 3ia^3 \log(x) - 3ia^3 \log\left(1 + \sqrt{1+a^2x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^(I*ArcTan[a*x])*x^4),x]

[Out] ((Sqrt[1 + a^2*x^2]*(-2 + (3*I)*a*x + 4*a^2*x^2))/x^3 + (3*I)*a^3*Log[x] - (3*I)*a^3*Log[1 + Sqrt[1 + a^2*x^2]])/6

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 272 vs. 2(72) = 144.

time = 0.07, size = 273, normalized size = 3.03

method	result
risch	$\frac{4a^4x^4+3ia^3x^3+2a^2x^2+3iax-2}{6x^3\sqrt{a^2x^2+1}} - \frac{ia^3 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{2}$
default	$-ia^3 \left(\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2}{\sqrt{a^2}} + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}\right)}{\sqrt{a^2}} \right) - \frac{(a^2x^2+1)}{3x^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^4,x,method=_RETURNVERBOSE)

[Out] -I*a^3*(((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+I*a*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)))/(a^2)^(1/2))-1/3/x^3*(a^2*x^2+1)^(3/2)-I*a*(-1/2/x^2*(a^2*x^2+1)^(3/2)+1/2*a^2*((a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2))))+I*a^3*((a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2)))-a^2*(-1/x*(a^2*x^2+1)^(3/2)+2*a^2*(1/2*x*(a^2*x^2+1)^(1/2)+1/2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^4,x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^2 + 1)/((I*a*x + 1)*x^4), x)

Fricas [A]

time = 1.51, size = 92, normalized size = 1.02

$$\frac{-3i a^3 x^3 \log(-ax + \sqrt{a^2 x^2 + 1} + 1) + 3i a^3 x^3 \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + 4 a^3 x^3 + (4 a^2 x^2 + 3i a x - 2) \sqrt{a^2 x^2 + 1}}{6 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^4,x, algorithm="fricas")`

```
[Out] 1/6*(-3*I*a^3*x^3*log(-a*x + sqrt(a^2*x^2 + 1) + 1) + 3*I*a^3*x^3*log(-a*x
+ sqrt(a^2*x^2 + 1) - 1) + 4*a^3*x^3 + (4*a^2*x^2 + 3*I*a*x - 2)*sqrt(a^2*x
^2 + 1))/x^3
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 x^2 + 1}}{a x^5 - i x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/x**4,x)``[Out] -I*Integral(sqrt(a**2*x**2 + 1)/(a*x**5 - I*x**4), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^4,x, algorithm="giac")`

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [B]

time = 0.04, size = 74, normalized size = 0.82

$$\frac{2 a^2 \sqrt{a^2 x^2 + 1}}{3 x} - \frac{\sqrt{a^2 x^2 + 1}}{3 x^3} - \frac{a^3 \operatorname{atan}\left(\sqrt{a^2 x^2 + 1} \operatorname{li}\right)}{2} + \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{2 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a^2*x^2 + 1)^(1/2)/(x^4*(a*x*1i + 1)),x)`

```
[Out] (a*(a^2*x^2 + 1)^(1/2)*1i)/(2*x^2) - (a^2*x^2 + 1)^(1/2)/(3*x^3) - (a^3*ata
n((a^2*x^2 + 1)^(1/2)*1i))/2 + (2*a^2*(a^2*x^2 + 1)^(1/2))/(3*x)
```

3.43 $\int \frac{e^{-i\text{ArcTan}(ax)}}{x^5} dx$

Optimal. Leaf size=113

$$-\frac{\sqrt{1+a^2x^2}}{4x^4} + \frac{ia\sqrt{1+a^2x^2}}{3x^3} + \frac{3a^2\sqrt{1+a^2x^2}}{8x^2} - \frac{2ia^3\sqrt{1+a^2x^2}}{3x} - \frac{3}{8}a^4 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-3/8*a^4*\text{arctanh}((a^2*x^2+1)^{(1/2)})-1/4*(a^2*x^2+1)^{(1/2)}/x^4+1/3*I*a*(a^2*x^2+1)^{(1/2)}/x^3+3/8*a^2*(a^2*x^2+1)^{(1/2)}/x^2-2/3*I*a^3*(a^2*x^2+1)^{(1/2)}/x$

Rubi [A]

time = 0.07, antiderivative size = 113, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5168, 849, 821, 272, 65, 214}

$$\frac{3a^2\sqrt{a^2x^2+1}}{8x^2} - \frac{\sqrt{a^2x^2+1}}{4x^4} + \frac{ia\sqrt{a^2x^2+1}}{3x^3} - \frac{3}{8}a^4 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right) - \frac{2ia^3\sqrt{a^2x^2+1}}{3x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{(I*\text{ArcTan}[a*x])}*x^5), x]$

[Out] $-1/4*\text{Sqrt}[1 + a^2*x^2]/x^4 + ((I/3)*a*\text{Sqrt}[1 + a^2*x^2])/x^3 + (3*a^2*\text{Sqrt}[1 + a^2*x^2])/(8*x^2) - (((2*I)/3)*a^3*\text{Sqrt}[1 + a^2*x^2])/x - (3*a^4*\text{ArcTanh}[\text{Sqrt}[1 + a^2*x^2]])/8$

Rule 65

$\text{Int}[(a_.) + (b_.)*(x_)^m]*((c_.) + (d_.)*(x_)^n), x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^n}, x], x, (a+b*x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\text{Int}[(a_) + (b_.)*(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Rule 272

$\text{Int}[(x_)^m]*((a_) + (b_.)*(x_)^n)^p, x_Symbol] \rightarrow \text{Dist}[1/n, \text{Subst}[\text{Int}[x^{(\text{Simplify}[(m+1)/n]-1)*(a+b*x)^p}, x], x, x^n], x] /; \text{FreeQ}[\{a, b, m, n, p\}, x] \&\& \text{IntegerQ}[\text{Simplify}[(m+1)/n]]$

Rule 821


```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(-e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)
)/(2*(p + 1)*(c*d^2 + a*e^2)), x] + Dist[(c*d*f + a*e*g)/(c*d^2 + a*e^2),
Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, f, g, m,
p}, x] && NeQ[c*d^2 + a*e^2, 0] && EqQ[Simplify[m + 2*p + 3], 0]
```

Rule 849

```
Int[((d_.) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p
_.), x_Symbol] := Simp[(e*f - d*g)*(d + e*x)^(m + 1)*((a + c*x^2)^(p + 1)/(
(m + 1)*(c*d^2 + a*e^2))), x] + Dist[1/((m + 1)*(c*d^2 + a*e^2)), Int[(d +
e*x)^(m + 1)*(a + c*x^2)^p*Simp[(c*d*f + a*e*g)*(m + 1) - c*(e*f - d*g)*(m
+ 2*p + 3)*x, x], x], x] /; FreeQ[{a, c, d, e, f, g, p}, x] && NeQ[c*d^2 +
a*e^2, 0] && LtQ[m, -1] && (IntegerQ[m] || IntegerQ[p] || IntegersQ[2*m, 2*
p])
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{1 - iax}{x^5 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} - \frac{1}{4} \int \frac{4ia + 3a^2 x}{x^4 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{1}{12} \int \frac{-9a^2 + 8ia^3 x}{x^3 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} - \frac{1}{24} \int \frac{-16ia^3 - 9a^4 x}{x^2 \sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} - \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} + \frac{1}{8}(3a^4) \int \frac{1}{x\sqrt{1 + a^2 x^2}} dx \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} - \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} + \frac{1}{16}(3a^4) \text{Subst}\left(\frac{1}{u\sqrt{1 + a^2 u^2}}, u, x\right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} - \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} + \frac{1}{8}(3a^2) \text{Subst}\left(\frac{1}{u\sqrt{1 + a^2 u^2}}, u, x\right) \\
&= -\frac{\sqrt{1 + a^2 x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2 x^2}}{3x^3} + \frac{3a^2\sqrt{1 + a^2 x^2}}{8x^2} - \frac{2ia^3\sqrt{1 + a^2 x^2}}{3x} - \frac{3}{8}a^4 \tanh^{-1}\left(\frac{ax}{\sqrt{1 + a^2 x^2}}\right)
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 76, normalized size = 0.67

$$\frac{1}{24} \left(\frac{\sqrt{1+a^2x^2}(-6+8iax+9a^2x^2-16ia^3x^3)}{x^4} + 9a^4 \log(x) - 9a^4 \log\left(1 + \sqrt{1+a^2x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

`[In] Integrate[1/(E^(I*ArcTan[a*x])*x^5),x]`

`[Out] ((Sqrt[1 + a^2*x^2]*(-6 + (8*I)*a*x + 9*a^2*x^2 - (16*I)*a^3*x^3))/x^4 + 9*a^4*Log[x] - 9*a^4*Log[1 + Sqrt[1 + a^2*x^2]])/24`

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 289 vs. 2(91) = 182.

time = 0.08, size = 290, normalized size = 2.57

method	result
risch	$-\frac{i(16a^5x^5+9ia^4x^4+8a^3x^3+3ia^2x^2-8ax-6i)}{24x^4\sqrt{a^2x^2+1}} - \frac{3a^4 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right)}{8}$
default	$-a^4 \left(\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2 + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}}{\sqrt{a^2}} + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}\right)}{\sqrt{a^2}} \right) + \frac{ia(a^2x^2+1)}{3x^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^5,x,method=_RETURNVERBOSE)`

`[Out] -a^4*(((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+I*a*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)))/(a^2)^(1/2))+1/3*I*a/x^3*(a^2*x^2+1)^(3/2)-5/4*a^2*(-1/2/x^2*(a^2*x^2+1)^(3/2)+1/2*a^2*((a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2))))+a^4*((a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2)))-1/4/x^4*(a^2*x^2+1)^(3/2)+I*a^3*(-1/x*(a^2*x^2+1)^(3/2)+2*a^2*(1/2*x*(a^2*x^2+1)^(1/2)+1/2*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2)))`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^5,x, algorithm="maxima")`

[Out] integrate(sqrt(a^2*x^2 + 1)/((I*a*x + 1)*x^5), x)

Fricas [A]

time = 2.49, size = 101, normalized size = 0.89

$$\frac{9a^4x^4 \log(-ax + \sqrt{a^2x^2 + 1} + 1) - 9a^4x^4 \log(-ax + \sqrt{a^2x^2 + 1} - 1) + 16ia^4x^4 - (-16ia^3x^3 + 9a^2x^2 + 8iax - 6)\sqrt{a^2x^2 + 1}}{24x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^5,x, algorithm="fricas")

[Out] -1/24*(9*a^4*x^4*log(-a*x + sqrt(a^2*x^2 + 1) + 1) - 9*a^4*x^4*log(-a*x + sqrt(a^2*x^2 + 1) - 1) + 16*I*a^4*x^4 - (-16*I*a^3*x^3 + 9*a^2*x^2 + 8*I*a*x - 6)*sqrt(a^2*x^2 + 1))/x^4

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2x^2 + 1}}{ax^6 - ix^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/x**5,x)

[Out] -I*Integral(sqrt(a**2*x**2 + 1)/(a*x**6 - I*x**5), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/x^5,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.03, size = 95, normalized size = 0.84

$$\frac{a^4 \operatorname{atan}\left(\sqrt{a^2x^2 + 1} \operatorname{li}\right) 3i}{8} - \frac{\sqrt{a^2x^2 + 1}}{4x^4} + \frac{a\sqrt{a^2x^2 + 1} \operatorname{li}}{3x^3} + \frac{3a^2\sqrt{a^2x^2 + 1}}{8x^2} - \frac{a^3\sqrt{a^2x^2 + 1} 2i}{3x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(1/2)/(x^5*(a*x*1i + 1)),x)

[Out] (a^4*atan((a^2*x^2 + 1)^(1/2)*1i)*3i)/8 - (a^2*x^2 + 1)^(1/2)/(4*x^4) + (a*(a^2*x^2 + 1)^(1/2)*1i)/(3*x^3) + (3*a^2*(a^2*x^2 + 1)^(1/2))/(8*x^2) - (a^3*(a^2*x^2 + 1)^(1/2)*2i)/(3*x)

3.44 $\int e^{-2i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=49

$$\frac{2ix}{a^3} + \frac{x^2}{a^2} - \frac{2ix^3}{3a} - \frac{x^4}{4} - \frac{2\log(i-ax)}{a^4}$$

[Out] $2*I*x/a^3+x^2/a^2-2/3*I*x^3/a-1/4*x^4-2*\ln(I-a*x)/a^4$

Rubi [A]

time = 0.03, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-\frac{2\log(-ax+i)}{a^4} + \frac{2ix}{a^3} + \frac{x^2}{a^2} - \frac{2ix^3}{3a} - \frac{x^4}{4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{((2*I)*\text{ArcTan}[a*x])}, x]$

[Out] $((2*I)*x)/a^3 + x^2/a^2 - (((2*I)/3)*x^3)/a - x^4/4 - (2*\text{Log}[I - a*x])/a^4$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) || \text{EqQ}[p, 1] || (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] || \text{LeQ}[9*p + 5*(n + 2), 0] || \text{GeQ}[n + p + 1, 0] || (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*x^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}[\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1-iax)}{1+iax} dx \\ &= \int \left(\frac{2i}{a^3} + \frac{2x}{a^2} - \frac{2ix^2}{a} - x^3 - \frac{2}{a^3(-i+ax)} \right) dx \\ &= \frac{2ix}{a^3} + \frac{x^2}{a^2} - \frac{2ix^3}{3a} - \frac{x^4}{4} - \frac{2\log(i-ax)}{a^4} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 49, normalized size = 1.00

$$\frac{2ix}{a^3} + \frac{x^2}{a^2} - \frac{2ix^3}{3a} - \frac{x^4}{4} - \frac{2 \log(i - ax)}{a^4}$$

Antiderivative was successfully verified.

`[In] Integrate[x^3/E^((2*I)*ArcTan[a*x]),x]``[Out] ((2*I)*x)/a^3 + x^2/a^2 - (((2*I)/3)*x^3)/a - x^4/4 - (2*Log[I - a*x])/a^4`**Maple [A]**

time = 0.08, size = 48, normalized size = 0.98

method	result	size
default	$-\frac{\frac{1}{4}a^3x^4 + \frac{2}{3}ia^2x^3 - ax^2 - 2ix}{a^3} - \frac{2 \ln(-ax+i)}{a^4}$	48
risch	$-\frac{x^4}{4} - \frac{2ix^3}{3a} + \frac{x^2}{a^2} + \frac{2ix}{a^3} - \frac{\ln(a^2x^2+1)}{a^4} - \frac{2i \arctan(ax)}{a^4}$	55
meijerg	$-\frac{ixa(-3a^4x^4 - 5ia^3x^3 + 10a^2x^2 + 30iax + 60)}{12(iax+1)a^4} + 5 \ln(iax+1) + \frac{-\frac{iax(2a^2x^2 + 6iax + 12)}{4(iax+1)} + 3 \ln(iax+1)}{a^4}$	108

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^3/(1+I*a*x)^2*(a^2*x^2+1),x,method=_RETURNVERBOSE)``[Out] -1/a^3*(1/4*a^3*x^4+2/3*I*a^2*x^3-a*x^2-2*I*x)-2*ln(I-a*x)/a^4`**Maxima [A]**

time = 0.27, size = 44, normalized size = 0.90

$$-\frac{i(-3ia^3x^4 + 8a^2x^3 + 12iax^2 - 24x)}{12a^3} - \frac{2 \log(iax + 1)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="maxima")``[Out] -1/12*I*(-3*I*a^3*x^4 + 8*a^2*x^3 + 12*I*a*x^2 - 24*x)/a^3 - 2*log(I*a*x + 1)/a^4`**Fricas [A]**

time = 2.64, size = 46, normalized size = 0.94

$$-\frac{3a^4x^4 + 8ia^3x^3 - 12a^2x^2 - 24iax + 24 \log\left(\frac{ax-i}{a}\right)}{12a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="fricas")`

[Out] $-1/12*(3*a^4*x^4 + 8*I*a^3*x^3 - 12*a^2*x^2 - 24*I*a*x + 24*\log((a*x - I)/a))/a^4$

Sympy [A]

time = 0.06, size = 41, normalized size = 0.84

$$-\frac{x^4}{4} - \frac{2ix^3}{3a} + \frac{x^2}{a^2} + \frac{2ix}{a^3} - \frac{2\log(ax - i)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**3/(1+I*a*x)**2*(a**2*x**2+1),x)`

[Out] $-x^{**4}/4 - 2*I*x^{**3}/(3*a) + x^{**2}/a^{**2} + 2*I*x/a^{**3} - 2*\log(a*x - I)/a^{**4}$

Giac [A]

time = 0.41, size = 68, normalized size = 1.39

$$\frac{(i ax + 1)^4 \left(\frac{20}{i ax + 1} - \frac{54}{(i ax + 1)^2} + \frac{84}{(i ax + 1)^3} - 3 \right)}{12 a^4} + \frac{2 \log\left(\frac{1}{\sqrt{a^2 x^2 + 1} |a|}\right)}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="giac")`

[Out] $1/12*(I*a*x + 1)^4*(20/(I*a*x + 1) - 54/(I*a*x + 1)^2 + 84/(I*a*x + 1)^3 - 3)/a^4 + 2*\log(1/(\sqrt{a^2*x^2 + 1}*abs(a)))/a^4$

Mupad [B]

time = 0.06, size = 43, normalized size = 0.88

$$\frac{x^2}{a^2} - \frac{x^4}{4} - \frac{2 \ln\left(x - \frac{1i}{a}\right)}{a^4} + \frac{x 2i}{a^3} - \frac{x^3 2i}{3 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^3*(a^2*x^2 + 1))/(a*x*1i + 1)^2,x)`

[Out] $(x*2i)/a^3 - (2*\log(x - 1i/a))/a^4 - x^4/4 - (x^3*2i)/(3*a) + x^2/a^2$

3.45 $\int e^{-2i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=40

$$\frac{2x}{a^2} - \frac{ix^2}{a} - \frac{x^3}{3} + \frac{2i \log(i - ax)}{a^3}$$

[Out] $2*x/a^2 - I*x^2/a - 1/3*x^3 + 2*I*\ln(I - a*x)/a^3$

Rubi [A]

time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$\frac{2i \log(-ax + i)}{a^3} + \frac{2x}{a^2} - \frac{ix^2}{a} - \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{((2*I)*\text{ArcTan}[a*x])}, x]$

[Out] $(2*x)/a^2 - (I*x^2)/a - x^3/3 + ((2*I)*\text{Log}[I - a*x])/a^3$

Rule 78

$\text{Int}[(a + b*x)^n * (c + d*x)^m * (e + f*x)^p, x] := \text{Int}[\text{ExpandIntegrand}[(a + b*x)^n * (c + d*x)^m * (e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5170

$\text{Int}[E^{\text{ArcTan}[a*x]^n} * (x^m), x] := \text{Int}[x^m * ((1 - I*a*x)^{I*(n/2)} / (1 + I*a*x)^{I*(n/2)}), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1 - iax)}{1 + iax} dx \\ &= \int \left(\frac{2}{a^2} - \frac{2ix}{a} - x^2 + \frac{2i}{a^2(-i + ax)} \right) dx \\ &= \frac{2x}{a^2} - \frac{ix^2}{a} - \frac{x^3}{3} + \frac{2i \log(i - ax)}{a^3} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 40, normalized size = 1.00

$$\frac{2x}{a^2} - \frac{ix^2}{a} - \frac{x^3}{3} + \frac{2i \log(i - ax)}{a^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^((2*I)*ArcTan[a*x]),x]

[Out] (2*x)/a^2 - (I*x^2)/a - x^3/3 + ((2*I)*Log[I - a*x])/a^3

Maple [A]

time = 0.07, size = 40, normalized size = 1.00

method	result	size
default	$-\frac{\frac{1}{3}a^2x^3+iax^2-2x}{a^2} + \frac{2i \ln(-ax+i)}{a^3}$	40
risch	$-\frac{x^3}{3} - \frac{ix^2}{a} + \frac{2x}{a^2} + \frac{i \ln(a^2x^2+1)}{a^3} - \frac{2 \arctan(ax)}{a^3}$	47
meijerg	$-\frac{i \left(\frac{ixa(-5ia^3x^3+10a^2x^2+30iax+60)}{15iax+15} - 4 \ln(iax+1) \right)}{a^3} + \frac{i \left(\frac{iax(3iax+6)}{3iax+3} - 2 \ln(iax+1) \right)}{a^3}$	95

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*a*x)^2*(a^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] -1/a^2*(1/3*a^2*x^3+I*a*x^2-2*x)+2*I*ln(I-a*x)/a^3

Maxima [A]

time = 0.26, size = 35, normalized size = 0.88

$$-\frac{a^2x^3 + 3i ax^2 - 6x}{3a^2} + \frac{2i \log(iax + 1)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="maxima")

[Out] -1/3*(a^2*x^3 + 3*I*a*x^2 - 6*x)/a^2 + 2*I*log(I*a*x + 1)/a^3

Fricas [A]

time = 1.88, size = 37, normalized size = 0.92

$$-\frac{a^3x^3 + 3i a^2x^2 - 6ax - 6i \log\left(\frac{ax-i}{a}\right)}{3a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="fricas")

[Out] $-1/3*(a^3*x^3 + 3*I*a^2*x^2 - 6*a*x - 6*I*\log((a*x - I)/a))/a^3$

Sympy [A]

time = 0.06, size = 31, normalized size = 0.78

$$-\frac{x^3}{3} - \frac{ix^2}{a} + \frac{2x}{a^2} + \frac{2i \log(ax - i)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2/(1+I*a*x)**2*(a**2*x**2+1),x)`

[Out] $-x**3/3 - I*x**2/a + 2*x/a**2 + 2*I*\log(a*x - I)/a**3$

Giac [A]

time = 0.41, size = 58, normalized size = 1.45

$$\frac{i(iax + 1)^3 \left(\frac{6}{iax+1} - \frac{15}{(iax+1)^2} - 1 \right)}{3a^3} - \frac{2i \log\left(\frac{1}{\sqrt{a^2x^2 + 1}|a|}\right)}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="giac")`

[Out] $1/3*I*(I*a*x + 1)^3*(6/(I*a*x + 1) - 15/(I*a*x + 1)^2 - 1)/a^3 - 2*I*\log(1/(\sqrt{a^2*x^2 + 1}*abs(a)))/a^3$

Mupad [B]

time = 0.41, size = 36, normalized size = 0.90

$$\frac{\ln\left(x - \frac{1i}{a}\right) 2i}{a^3} + \frac{2x}{a^2} - \frac{x^3}{3} - \frac{x^2 1i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2*(a^2*x^2 + 1))/(a*x*1i + 1)^2,x)`

[Out] $(\log(x - 1i/a)*2i)/a^3 + (2*x)/a^2 - x^3/3 - (x^2*1i)/a$

3.46 $\int e^{-2i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=30

$$-\frac{2ix}{a} - \frac{x^2}{2} + \frac{2\log(i-ax)}{a^2}$$

[Out] $-2*I*x/a-1/2*x^2+2*\ln(I-a*x)/a^2$

Rubi [A]

time = 0.02, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5170, 78}

$$\frac{2\log(-ax+i)}{a^2} - \frac{2ix}{a} - \frac{x^2}{2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x/E^{((2*I)*\text{ArcTan}[a*x])}, x]$

[Out] $((-2*I)*x)/a - x^2/2 + (2*\text{Log}[I - a*x])/a^2$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_))*((c_. + (d_.)*(x_))^{(n_.)*((e_. + (f_.)*(x_))^{(p_.)}, x_Symbol)] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) || \text{EqQ}[p, 1] || (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] || \text{LeQ}[9*p + 5*(n + 2), 0] || \text{GeQ}[n + p + 1, 0] || (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_)]*(n_.))*(x_)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}[\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(ax)} x dx &= \int \frac{x(1 - iax)}{1 + iax} dx \\ &= \int \left(-\frac{2i}{a} - x + \frac{2}{a(-i + ax)} \right) dx \\ &= -\frac{2ix}{a} - \frac{x^2}{2} + \frac{2\log(i-ax)}{a^2} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 30, normalized size = 1.00

$$-\frac{2ix}{a} - \frac{x^2}{2} + \frac{2\log(i - ax)}{a^2}$$

Antiderivative was successfully verified.

`[In] Integrate[x/E^((2*I)*ArcTan[a*x]),x]``[Out] ((-2*I)*x)/a - x^2/2 + (2*Log[I - a*x])/a^2`**Maple [A]**

time = 0.07, size = 31, normalized size = 1.03

method	result	size
default	$-\frac{\frac{1}{2}ax^2+2ix}{a} + \frac{2\ln(-ax+i)}{a^2}$	31
risch	$-\frac{x^2}{2} - \frac{2ix}{a} + \frac{\ln(a^2x^2+1)}{a^2} + \frac{2i\arctan(ax)}{a^2}$	38
meijerg	$-\frac{iax(2a^2x^2+6iax+12)}{4(iax+1)} + 3\ln(iax+1) - \frac{-\frac{iax}{iax+1} + \ln(iax+1)}{a^2}$	74

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x/(1+I*a*x)^2*(a^2*x^2+1),x,method=_RETURNVERBOSE)``[Out] -1/a*(1/2*a*x^2+2*I*x)+2*ln(I-a*x)/a^2`**Maxima [A]**

time = 0.27, size = 28, normalized size = 0.93

$$\frac{i(iax^2 - 4x)}{2a} + \frac{2\log(iax + 1)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="maxima")``[Out] 1/2*I*(I*a*x^2 - 4*x)/a + 2*log(I*a*x + 1)/a^2`**Fricas [A]**

time = 1.67, size = 29, normalized size = 0.97

$$-\frac{a^2x^2 + 4iax - 4\log\left(\frac{ax-i}{a}\right)}{2a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="fricas")`

[Out] $-1/2*(a^2*x^2 + 4*I*a*x - 4*\log((a*x - I)/a))/a^2$

Sympy [A]

time = 0.05, size = 22, normalized size = 0.73

$$-\frac{x^2}{2} - \frac{2ix}{a} + \frac{2\log(ax - i)}{a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*a*x)**2*(a**2*x**2+1),x)`

[Out] $-x^{**2}/2 - 2*I*x/a + 2*\log(a*x - I)/a^{**2}$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 52 vs. $2(24) = 48$.

time = 0.41, size = 52, normalized size = 1.73

$$\frac{i \left(\frac{(i a x + 1)^2 \left(-\frac{6i}{i a x + 1} + i \right)}{a} - \frac{4i \log \left(\frac{1}{\sqrt{a^2 x^2 + 1}} \right)}{a} \right)}{2 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="giac")`

[Out] $-1/2*I*((I*a*x + 1)^2*(-6*I/(I*a*x + 1) + I)/a - 4*I*\log(1/(\sqrt{a^2*x^2 + 1})*\text{abs}(a)))/a/a$

Mupad [B]

time = 0.42, size = 27, normalized size = 0.90

$$\frac{2 \ln \left(x - \frac{1i}{a} \right)}{a^2} - \frac{x^2}{2} - \frac{x 2i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x*(a^2*x^2 + 1))/(a*x*1i + 1)^2,x)`

[Out] $(2*\log(x - 1i/a))/a^2 - (x*2i)/a - x^2/2$

3.47 $\int e^{-2i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=20

$$-x - \frac{2i \log(i - ax)}{a}$$

[Out] $-x-2*I*\ln(I-a*x)/a$

Rubi [A]

time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5169, 45}

$$-x - \frac{2i \log(-ax + i)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((-2*I)*\text{ArcTan}[a*x])}, x]$

[Out] $-x - ((2*I)*\text{Log}[I - a*x])/a$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5169

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}, x] /;$ FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(ax)} dx &= \int \frac{1 - iax}{1 + iax} dx \\ &= \int \left(-1 - \frac{2i}{-i + ax} \right) dx \\ &= -x - \frac{2i \log(i - ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 30, normalized size = 1.50

$$-x + \frac{2\text{ArcTan}(ax)}{a} - \frac{i \log(1 + a^2x^2)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((-2*I)*ArcTan[a*x]),x]

[Out] $-x + (2*\text{ArcTan}[a*x])/a - (I*\text{Log}[1 + a^2*x^2])/a$

Maple [A]

time = 0.07, size = 19, normalized size = 0.95

method	result	size
default	$-x - \frac{2i \ln(-ax+i)}{a}$	19
risch	$-x - \frac{i \ln(a^2x^2+1)}{a} + \frac{2 \arctan(ax)}{a}$	30
meijerg	$\frac{i \left(\frac{iax(3iax+6)}{3iax+3} - 2 \ln(iax+1) \right)}{a} + \frac{x}{iax+1}$	51

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^2*(a^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] $-x-2*I*\ln(I-a*x)/a$

Maxima [A]

time = 0.28, size = 16, normalized size = 0.80

$$-x - \frac{2i \log(iax + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="maxima")

[Out] $-x - 2*I*\log(I*a*x + 1)/a$

Fricas [A]

time = 1.56, size = 21, normalized size = 1.05

$$\frac{ax + 2i \log\left(\frac{ax-i}{a}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="fricas")

[Out] $-(a*x + 2*I*\log((a*x - I)/a))/a$

Sympy [A]

time = 0.05, size = 14, normalized size = 0.70

$$-x - \frac{2i \log(ax - i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**2*(a**2*x**2+1),x)

[Out] $-x - 2*I*\log(a*x - I)/a$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 65 vs. $2(16) = 32$.

time = 0.40, size = 65, normalized size = 3.25

$$a^2 \left(\frac{i(iax + 1)}{a^3} + \frac{2i \log\left(\frac{1}{\sqrt{a^2x^2 + 1}|a|}\right)}{a^3} - \frac{i}{(iax + 1)a^3} \right) + \frac{i}{(iax + 1)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="giac")

[Out] $a^2*(I*(I*a*x + 1)/a^3 + 2*I*\log(1/(\sqrt{a^2*x^2 + 1}*abs(a)))/a^3 - I/((I*a*x + 1)*a^3)) + I/((I*a*x + 1)*a)$

Mupad [B]

time = 0.41, size = 19, normalized size = 0.95

$$-x - \frac{\ln\left(x - \frac{1i}{a}\right) 2i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)/(a*x*1i + 1)^2,x)

[Out] $-x - (\log(x - 1i/a)*2i)/a$

$$3.48 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=14

$$\log(x) - 2 \log(i - ax)$$

[Out] ln(x)-2*ln(I-a*x)

Rubi [A]

time = 0.02, antiderivative size = 14, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$\log(x) - 2 \log(-ax + i)$$

Antiderivative was successfully verified.

[In] Int[1/(E^((2*I)*ArcTan[a*x])*x),x]

[Out] Log[x] - 2*Log[I - a*x]

Rule 78

```
Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{x} dx &= \int \frac{1 - iax}{x(1 + iax)} dx \\ &= \int \left(\frac{1}{x} - \frac{2a}{-i + ax} \right) dx \\ &= \log(x) - 2 \log(i - ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 14, normalized size = 1.00

$$\log(x) - 2\log(i - ax)$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((2*I)*ArcTan[a*x]))*x],x]

[Out] Log[x] - 2*Log[I - a*x]

Maple [A]

time = 0.07, size = 14, normalized size = 1.00

method	result	size
default	$\ln(x) - 2\ln(-ax + i)$	14
risch	$\ln(x) - \ln(a^2x^2 + 1) - 2i \arctan(ax)$	23
meijerg	$\frac{iax}{iax+1} - 2\ln(iax + 1) - \frac{2iax}{2iax+2} + 1 + \ln(x) + \ln(ia)$	48

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)/x,x,method=_RETURNVERBOSE)

[Out] ln(x)-2*ln(I-a*x)

Maxima [A]

time = 0.26, size = 12, normalized size = 0.86

$$-2\log(iax + 1) + \log(x)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x,x, algorithm="maxima")

[Out] -2*log(I*a*x + 1) + log(x)

Fricas [A]

time = 1.81, size = 15, normalized size = 1.07

$$\log(x) - 2\log\left(\frac{ax - i}{a}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x,x, algorithm="fricas")

[Out] log(x) - 2*log((a*x - I)/a)

Sympy [A]

time = 0.07, size = 17, normalized size = 1.21

$$\log(3ax) - 2\log(3ax - 3i)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**2*(a**2*x**2+1)/x,x)

[Out] log(3*a*x) - 2*log(3*a*x - 3*I)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 44 vs. 2(12) = 24.

time = 0.41, size = 44, normalized size = 3.14

$$i a \left(-\frac{i \log\left(\frac{i}{i a x + 1} - i\right)}{a} - \frac{i \log\left(\frac{1}{\sqrt{a^2 x^2 + 1} |a|}\right)}{a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x,x, algorithm="giac")

[Out] I*a*(-I*log(I/(I*a*x + 1) - I)/a - I*log(1/(sqrt(a^2*x^2 + 1)*abs(a)))/a)

Mupad [B]

time = 0.45, size = 14, normalized size = 1.00

$$\ln(x) - 2 \ln\left(x - \frac{1i}{a}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)/(x*(a*x*1i + 1)^2),x)

[Out] log(x) - 2*log(x - 1i/a)

$$3.49 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=27

$$-\frac{1}{x} - 2ia \log(x) + 2ia \log(i - ax)$$

[Out] $-1/x - 2I*a*\ln(x) + 2I*a*\ln(I - a*x)$

Rubi [A]

time = 0.02, antiderivative size = 27, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-2ia \log(x) + 2ia \log(-ax + i) - \frac{1}{x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((2*I)*\text{ArcTan}[a*x])}*x^2), x]$

[Out] $-x^{(-1)} - (2*I)*a*\text{Log}[x] + (2*I)*a*\text{Log}[I - a*x]$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2)})/(1 + I*a*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{1 - iax}{x^2(1 + iax)} dx \\ &= \int \left(\frac{1}{x^2} - \frac{2ia}{x} + \frac{2ia^2}{-i + ax} \right) dx \\ &= -\frac{1}{x} - 2ia \log(x) + 2ia \log(i - ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 27, normalized size = 1.00

$$-\frac{1}{x} - 2ia \log(x) + 2ia \log(i - ax)$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a*x])*x^2),x]``[Out] -x^(-1) - (2*I)*a*Log[x] + (2*I)*a*Log[I - a*x]`**Maple [A]**

time = 0.10, size = 25, normalized size = 0.93

method	result	size
default	$-\frac{1}{x} - 2ia \ln(x) + 2ia \ln(-ax + i)$	25
risch	$-\frac{1}{x} - 2ia \ln(x) - 2a \arctan(ax) + ia \ln(a^2x^2 + 1)$	34
meijerg	$\frac{a^2x}{iax+1} + ia\left(\frac{3iax}{3iax+3} + 2 \ln(iax + 1) - 1 - 2 \ln(x) - 2 \ln(ia) + \frac{i}{xa}\right)$	66

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)/x^2,x,method=_RETURNVERBOSE)``[Out] -1/x-2*I*a*ln(x)+2*I*a*ln(I-a*x)`**Maxima [A]**

time = 0.26, size = 34, normalized size = 1.26

$$2ia \log(iax + 1) - 2ia \log(x) - \frac{ax - i}{ax^2 - ix}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^2,x, algorithm="maxima")``[Out] 2*I*a*log(I*a*x + 1) - 2*I*a*log(x) - (a*x - I)/(a*x^2 - I*x)`**Fricas [A]**

time = 2.19, size = 26, normalized size = 0.96

$$\frac{-2iax \log(x) + 2iax \log\left(\frac{ax-i}{a}\right) - 1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^2,x, algorithm="fricas")``[Out] (-2*I*a*x*log(x) + 2*I*a*x*log((a*x - I)/a) - 1)/x`

Sympy [A]

time = 0.08, size = 32, normalized size = 1.19

$$-2a(i \log(4a^2x) - i \log(4a^2x - 4ia)) - \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**2*(a**2*x**2+1)/x**2,x)

[Out] -2*a*(I*log(4*a**2*x) - I*log(4*a**2*x - 4*I*a)) - 1/x

Giac [A]

time = 0.41, size = 34, normalized size = 1.26

$$-2i a \log\left(\frac{i}{i a x + 1} - i\right) - \frac{a}{\frac{i}{i a x + 1} - i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^2,x, algorithm="giac")

[Out] -2*I*a*log(I/(I*a*x + 1) - I) - a/(I/(I*a*x + 1) - I)

Mupad [B]

time = 0.42, size = 17, normalized size = 0.63

$$-4 a \operatorname{atan}(2 a x - i) - \frac{1}{x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)/(x^2*(a*x*1i + 1)^2),x)

[Out] - 4*a*atan(2*a*x - 1i) - 1/x

3.50 $\int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=37

$$-\frac{1}{2x^2} + \frac{2ia}{x} - 2a^2 \log(x) + 2a^2 \log(i - ax)$$

[Out] $-1/2/x^2+2*I*a/x-2*a^2*\ln(x)+2*a^2*\ln(I-a*x)$

Rubi [A]

time = 0.02, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$-2a^2 \log(x) + 2a^2 \log(-ax + i) + \frac{2ia}{x} - \frac{1}{2x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((2*I)*\text{ArcTan}[a*x])}*x^3), x]$

[Out] $-1/2*1/x^2 + ((2*I)*a)/x - 2*a^2*\text{Log}[x] + 2*a^2*\text{Log}[I - a*x]$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*x_.^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2)})/(1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, m, n\}, x\} \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{1 - iax}{x^3(1 + iax)} dx \\ &= \int \left(\frac{1}{x^3} - \frac{2ia}{x^2} - \frac{2a^2}{x} + \frac{2a^3}{-i + ax} \right) dx \\ &= -\frac{1}{2x^2} + \frac{2ia}{x} - 2a^2 \log(x) + 2a^2 \log(i - ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 37, normalized size = 1.00

$$-\frac{1}{2x^2} + \frac{2ia}{x} - 2a^2 \log(x) + 2a^2 \log(i - ax)$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a*x]))*x^3, x]``[Out] -1/2*1/x^2 + ((2*I)*a)/x - 2*a^2*Log[x] + 2*a^2*Log[I - a*x]`**Maple [A]**

time = 0.08, size = 34, normalized size = 0.92

method	result
default	$-\frac{1}{2x^2} + \frac{2ia}{x} - 2a^2 \ln(x) + 2a^2 \ln(-ax + i)$
risch	$\frac{2iax - \frac{1}{2}}{x^2} - 2a^2 \ln(-x) + 2ia^2 \arctan(ax) + a^2 \ln(a^2x^2 + 1)$
meijerg	$a^2 \left(-\frac{2iax}{2iax+2} - \ln(iax + 1) + 1 + \ln(x) + \ln(ia) \right) - a^2 \left(-\frac{4iax}{4iax+4} - 3 \ln(iax + 1) + 1 + 3 \ln(x) + \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)/x^3,x,method=_RETURNVERBOSE)``[Out] -1/2/x^2+2*I*a/x-2*a^2*ln(x)+2*a^2*ln(I-a*x)`**Maxima [A]**

time = 0.26, size = 50, normalized size = 1.35

$$2a^2 \log(iax + 1) - 2a^2 \log(x) - \frac{4a^2x^2 - 3iax + 1}{2iax^3 + 2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^3,x, algorithm="maxima")``[Out] 2*a^2*log(I*a*x + 1) - 2*a^2*log(x) - (4*a^2*x^2 - 3*I*a*x + 1)/(2*I*a*x^3 + 2*x^2)`**Fricas [A]**

time = 2.04, size = 39, normalized size = 1.05

$$\frac{4a^2x^2 \log(x) - 4a^2x^2 \log\left(\frac{ax-i}{a}\right) - 4iax + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^3,x, algorithm="fricas")`

[Out] $-1/2*(4*a^2*x^2*\log(x) - 4*a^2*x^2*\log((a*x - I)/a) - 4*I*a*x + 1)/x^2$

Sympy [A]

time = 0.10, size = 42, normalized size = 1.14

$$-2a^2(\log(4a^3x) - \log(4a^3x - 4ia^2)) - \frac{-4iax + 1}{2x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)**2*(a**2*x**2+1)/x**3,x)`

[Out] $-2*a**2*(\log(4*a**3*x) - \log(4*a**3*x - 4*I*a**2)) - (-4*I*a*x + 1)/(2*x**2)$

Giac [A]

time = 0.41, size = 54, normalized size = 1.46

$$-2a^2 \log\left(-\frac{1}{iax+1} + 1\right) + \frac{5a^2 - \frac{6a^2}{iax+1}}{2\left(\frac{i}{iax+1} - i\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^3,x, algorithm="giac")`

[Out] $-2*a^2*\log(-1/(I*a*x + 1) + 1) + 1/2*(5*a^2 - 6*a^2/(I*a*x + 1))/(I/(I*a*x + 1) - I)^2$

Mupad [B]

time = 0.07, size = 26, normalized size = 0.70

$$a^2 \operatorname{atan}(2ax - i) 4i + \frac{-\frac{1}{2} + ax 2i}{x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^2*x^2 + 1)/(x^3*(a*x+1i + 1)^2),x)`

[Out] $a^2*\operatorname{atan}(2*a*x - 1i)*4i + (a*x*2i - 1/2)/x^2$

3.51 $\int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=49

$$-\frac{1}{3x^3} + \frac{ia}{x^2} + \frac{2a^2}{x} + 2ia^3 \log(x) - 2ia^3 \log(i - ax)$$

[Out] $-1/3/x^3 + I*a/x^2 + 2*a^2/x + 2*I*a^3*\ln(x) - 2*I*a^3*\ln(I-a*x)$

Rubi [A]

time = 0.02, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5170, 78}

$$2ia^3 \log(x) - 2ia^3 \log(-ax + i) + \frac{2a^2}{x} + \frac{ia}{x^2} - \frac{1}{3x^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((2*I)*\text{ArcTan}[a*x])})*x^4), x]$

[Out] $-1/3*1/x^3 + (I*a)/x^2 + (2*a^2)/x + (2*I)*a^3*\text{Log}[x] - (2*I)*a^3*\text{Log}[I - a*x]$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}), x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) || \text{EqQ}[p, 1] || (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] || \text{LeQ}[9*p + 5*(n + 2), 0] || \text{GeQ}[n + p + 1, 0] || (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)}, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{1 - iax}{x^4(1 + iax)} dx \\ &= \int \left(\frac{1}{x^4} - \frac{2ia}{x^3} - \frac{2a^2}{x^2} + \frac{2ia^3}{x} - \frac{2ia^4}{-i + ax} \right) dx \\ &= -\frac{1}{3x^3} + \frac{ia}{x^2} + \frac{2a^2}{x} + 2ia^3 \log(x) - 2ia^3 \log(i - ax) \end{aligned}$$

Mathematica [A]

time = 0.01, size = 49, normalized size = 1.00

$$-\frac{1}{3x^3} + \frac{ia}{x^2} + \frac{2a^2}{x} + 2ia^3 \log(x) - 2ia^3 \log(i - ax)$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a*x])*x^4),x]``[Out] -1/3*1/x^3 + (I*a)/x^2 + (2*a^2)/x + (2*I)*a^3*Log[x] - (2*I)*a^3*Log[I - a*x]`**Maple [A]**

time = 0.10, size = 44, normalized size = 0.90

method	result
default	$-\frac{1}{3x^3} + \frac{ia}{x^2} + \frac{2a^2}{x} + 2ia^3 \ln(x) - 2ia^3 \ln(-ax + i)$
risch	$\frac{2a^2x^2+iax-\frac{1}{3}}{x^3} + 2ia^3 \ln(-x) + 2a^3 \arctan(ax) - ia^3 \ln(a^2x^2 + 1)$
meijerg	$ia^3 \left(\frac{3iax}{3iax+3} + 2 \ln(iax + 1) - 1 - 2 \ln(x) - 2 \ln(ia) + \frac{i}{xa} \right) - ia^3 \left(\frac{5iax}{5iax+5} + 4 \ln(iax + 1) - 1 - 4 \ln(x) \right)$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)/x^4,x,method=_RETURNVERBOSE)``[Out] -1/3/x^3+I*a/x^2+2*a^2/x+2*I*a^3*ln(x)-2*I*a^3*ln(I-a*x)`**Maxima [A]**

time = 0.26, size = 57, normalized size = 1.16

$$-2ia^3 \log(iax + 1) + 2ia^3 \log(x) + \frac{6ia^3x^3 + 3a^2x^2 + 2iax - 1}{3iax^4 + 3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^4,x, algorithm="maxima")``[Out] -2*I*a^3*log(I*a*x + 1) + 2*I*a^3*log(x) + (6*I*a^3*x^3 + 3*a^2*x^2 + 2*I*a*x - 1)/(3*I*a*x^4 + 3*x^3)`**Fricas [A]**

time = 1.74, size = 47, normalized size = 0.96

$$\frac{6ia^3x^3 \log(x) - 6ia^3x^3 \log\left(\frac{ax-i}{a}\right) + 6a^2x^2 + 3iax - 1}{3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^4,x, algorithm="fricas")`

[Out] $\frac{1}{3}*(6*I*a^3*x^3*\log(x) - 6*I*a^3*x^3*\log((a*x - I)/a) + 6*a^2*x^2 + 3*I*a*x - 1)/x^3$

Sympy [A]

time = 0.12, size = 54, normalized size = 1.10

$$-2a^3(-i \log(4a^4x) + i \log(4a^4x - 4ia^3)) - \frac{-6a^2x^2 - 3iax + 1}{3x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)**2*(a**2*x**2+1)/x**4,x)`

[Out] $-2*a**3*(-I*\log(4*a**4*x) + I*\log(4*a**4*x - 4*I*a**3)) - (-6*a**2*x**2 - 3*I*a*x + 1)/(3*x**3)$

Giac [A]

time = 0.41, size = 67, normalized size = 1.37

$$2i a^3 \log\left(\frac{i}{i a x + 1} - i\right) - \frac{10 a^3 - \frac{24 a^3}{i a x + 1} + \frac{15 a^3}{(i a x + 1)^2}}{3 \left(\frac{i}{i a x + 1} - i\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/x^4,x, algorithm="giac")`

[Out] $2*I*a^3*\log(I/(I*a*x + 1) - I) - 1/3*(10*a^3 - 24*a^3/(I*a*x + 1) + 15*a^3/(I*a*x + 1)^2)/(I/(I*a*x + 1) - I)^3$

Mupad [B]

time = 0.43, size = 33, normalized size = 0.67

$$4 a^3 \operatorname{atan}(2 a x - i) + \frac{2 a^2 x^2 + a x i - \frac{1}{3}}{x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^2*x^2 + 1)/(x^4*(a*x*1i + 1)^2),x)`

[Out] $4*a^3*\operatorname{atan}(2*a*x - 1i) + (a*x*1i + 2*a^2*x^2 - 1/3)/x^3$

3.52 $\int e^{-3i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=137

$$\frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{27\sqrt{1+a^2x^2}}{4a^4} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i+3ax)\sqrt{1+a^2x^2}}{8a^4} + \frac{51i\sinh^{-1}(ax)}{8a^4}$$

[Out] $51/8*I*\text{arcsinh}(a*x)/a^4+(1-I*a*x)^3/a^4/(a^2*x^2+1)^{(1/2)}+27/4*(a^2*x^2+1)^{(1/2)}/a^4-x^2*(a^2*x^2+1)^{(1/2)}/a^2+1/4*I*x^3*(a^2*x^2+1)^{(1/2)}/a-9/8*I*(2*I+3*a*x)*(a^2*x^2+1)^{(1/2)}/a^4$

Rubi [A]

time = 0.43, antiderivative size = 137, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5168, 1647, 1607, 12, 866, 1649, 1829, 27, 757, 655, 221}

$$\frac{51i\sinh^{-1}(ax)}{8a^4} - \frac{x^2\sqrt{a^2x^2+1}}{a^2} + \frac{ix^3\sqrt{a^2x^2+1}}{4a} - \frac{9i(3ax+2i)\sqrt{a^2x^2+1}}{8a^4} + \frac{27\sqrt{a^2x^2+1}}{4a^4} + \frac{(1-iax)^3}{a^4\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{((3*I)*\text{ArcTan}[a*x])}, x]$

[Out] $(1 - I*a*x)^3/(a^4*\text{Sqrt}[1 + a^2*x^2]) + (27*\text{Sqrt}[1 + a^2*x^2])/(4*a^4) - (x^2*\text{Sqrt}[1 + a^2*x^2])/a^2 + ((I/4)*x^3*\text{Sqrt}[1 + a^2*x^2])/a - (((9*I)/8)*(2*I + 3*a*x)*\text{Sqrt}[1 + a^2*x^2])/a^4 + (((51*I)/8)*\text{ArcSinh}[a*x])/a^4$

Rule 12

$\text{Int}[(a_)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_)*(v_)] /; \text{FreeQ}[b, x]$

Rule 27

$\text{Int}[(u_)*((a_) + (b_)*(x_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Int}[u*\text{Cancel}[(b/2 + c*x)^{(2*p)}/c^p], x] /; \text{FreeQ}[\{a, b, c\}, x] \ \&\& \ \text{EqQ}[b^2 - 4*a*c, 0] \ \&\& \ \text{IntegerQ}[p]$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{GtQ}[a, 0] \ \&\& \ \text{PosQ}[b]$

Rule 655

$\text{Int}[(d_)+(e_)*(x_))*((a_)+(c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*((a+c*x^2)^{(p+1)}/(2*c*(p+1))), x] + \text{Dist}[d, \text{Int}[(a+c*x^2)^p, x], x] /$

; FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]

Rule 757

Int[((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^(m - 1)*((a + c*x^2)^(p + 1)/(c*(m + 2*p + 1))), x] + Dist[1/(c*(m + 2*p + 1)), Int[(d + e*x)^(m - 2)*Simp[c*d^2*(m + 2*p + 1) - a*e^2*(m - 1) + 2*c*d*e*(m + p)*x, x]*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e, m, p}, x] && NeQ[c*d^2 + a*e^2, 0] && If[RationalQ[m], GtQ[m, 1], SumSimplerQ[m, -2]] && NeQ[m + 2*p + 1, 0] && IntQuadraticQ[a, 0, c, d, e, m, p, x]

Rule 866

Int[((d_) + (e_)*(x_))^(m_)*((f_) + (g_)*(x_)^n)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[d^(2*m)/a^m, Int[(f + g*x)^n*((a + c*x^2)^(m + p))/(d - e*x)^m], x], x] /; FreeQ[{a, c, d, e, f, g, n, p}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[f, 0] && ILtQ[m, -1] && !(IGtQ[n, 0] && ILtQ[m + n, 0] && !GtQ[p, 1])

Rule 1607

Int[(u_)*((a_)*(x_)^(p_) + (b_)*(x_)^(q_))^(n_), x_Symbol] := Int[u*x^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] && PosQ[q - p]

Rule 1647

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[d*e, Int[(d + e*x)^(m - 1)*PolynomialQuotient[Pq, a*e + c*d*x, x]*(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && PolyQ[Pq, x] && EqQ[c*d^2 + a*e^2, 0] && EqQ[PolynomialRemainder[Pq, a*e + c*d*x, x], 0]

Rule 1649

Int[(Pq_)*((d_) + (e_)*(x_))^(m_)*((a_) + (c_)*(x_)^2)^(p_), x_Symbol] := With[{Q = PolynomialQuotient[Pq, a*e + c*d*x, x], f = PolynomialRemainder[Pq, a*e + c*d*x, x]}, Simp[(-d)*f*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*a*e*(p + 1))), x] + Dist[d/(2*a*(p + 1)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^(p + 1)*ExpandToSum[2*a*e*(p + 1)*Q + f*(m + 2*p + 2), x], x], x] /; FreeQ[{a, c, d, e}, x] && PolyQ[Pq, x] && EqQ[c*d^2 + a*e^2, 0] && ILtQ[p + 1/2, 0] && GtQ[m, 0]

Rule 1829

Int[(Pq_)*((a_) + (b_)*(x_)^2)^(p_), x_Symbol] := With[{q = Expon[Pq, x], e = Coeff[Pq, x, Expon[Pq, x]]}, Simp[e*x^(q - 1)*((a + b*x^2)^(p + 1))/(b*(

```

q + 2*p + 1))), x] + Dist[1/(b*(q + 2*p + 1)), Int[(a + b*x^2)^p*ExpandToSu
m[b*(q + 2*p + 1)*Pq - a*e*(q - 1)*x^(q - 2) - b*e*(q + 2*p + 1)*x^q, x], x
], x]] /; FreeQ[{a, b, p}, x] && PolyQ[Pq, x] && !LeQ[p, -1]

```

Rule 5168

```

Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1-iax)^2}{(1+iax)\sqrt{1+a^2x^2}} dx \\
&= (ia) \int \frac{\sqrt{1+a^2x^2} \left(-\frac{ix^3}{a} - x^4\right)}{(1+iax)^2} dx \\
&= (ia) \int \frac{\left(-\frac{i}{a} - x\right) x^3 \sqrt{1+a^2x^2}}{(1+iax)^2} dx \\
&= a^2 \int \frac{x^3(1+a^2x^2)^{3/2}}{a^2(1+iax)^3} dx \\
&= \int \frac{x^3(1+a^2x^2)^{3/2}}{(1+iax)^3} dx \\
&= \int \frac{x^3(1-iax)^3}{(1+a^2x^2)^{3/2}} dx \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} - \int \frac{(1-iax)^2 \left(-\frac{3i}{a^3} - \frac{x}{a^2} + \frac{ix^2}{a}\right)}{\sqrt{1+a^2x^2}} dx \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{\int \frac{-\frac{12i}{a} - 28x + 27iax^2 + 12a^2x^3}{\sqrt{1+a^2x^2}} dx}{4a^2} \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{\int \frac{-36ia - 108a^2x + 81ia^3x^2}{\sqrt{1+a^2x^2}} dx}{12a^4} \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{\int \frac{9ia(2i+3ax)^2}{\sqrt{1+a^2x^2}} dx}{12a^4} \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{(3i) \int \frac{(2i+3ax)^2}{\sqrt{1+a^2x^2}} dx}{4a^3} \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i+3ax)\sqrt{1+a^2x^2}}{8a^4} - \frac{(3i)}{8a^4} \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{27\sqrt{1+a^2x^2}}{4a^4} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i+3ax)\sqrt{1+a^2x^2}}{8a^4} \\
&= \frac{(1-iax)^3}{a^4\sqrt{1+a^2x^2}} + \frac{27\sqrt{1+a^2x^2}}{4a^4} - \frac{x^2\sqrt{1+a^2x^2}}{a^2} + \frac{ix^3\sqrt{1+a^2x^2}}{4a} - \frac{9i(2i+3ax)\sqrt{1+a^2x^2}}{8a^4}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 80, normalized size = 0.58

$$\sqrt{1+a^2x^2} \left(\frac{6}{a^4} - \frac{19ix}{8a^3} - \frac{x^2}{a^2} + \frac{ix^3}{4a} - \frac{4i}{a^4(-i+ax)} \right) + \frac{51i \sinh^{-1}(ax)}{8a^4}$$

Antiderivative was successfully verified.

[In] Integrate[x^3/E^((3*I)*ArcTan[a*x]),x]

[Out] Sqrt[1+a^2*x^2]*(6/a^4 - ((19*I)/8)*x)/a^3 - x^2/a^2 + ((I/4)*x^3)/a - (4*I)/(a^4*(-I+a*x)) + (((51*I)/8)*ArcSinh[a*x])/a^4

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 681 vs. 2(114) = 228.

time = 0.12, size = 682, normalized size = 4.98

method	result
risch	$\frac{i(2a^3x^3+8ia^2x^2-19ax-48i)\sqrt{a^2x^2+1}}{8a^4} + \frac{51i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{8a^3\sqrt{a^2}} - \frac{4i\sqrt{\left(x-\frac{i}{a}\right)^2a^2+2ia\left(x-\frac{i}{a}\right)}}{a^5\left(x-\frac{i}{a}\right)}$
default	$i \left(\frac{x(a^2x^2+1)^{\frac{3}{2}}}{4} + \frac{3x\sqrt{a^2x^2+1}}{8} + \frac{3 \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{8\sqrt{a^2}} \right) - \frac{3 \left(\frac{\left(x-\frac{i}{a}\right)^2a^2+2ia\left(x-\frac{i}{a}\right)}{3} \right)^{\frac{3}{2}} + ia \left(\frac{(2\left(x-\frac{i}{a}\right)a^2+2ia)\sqrt{\dots}}{\dots} \right)}{a^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)

[Out] I/a^3*(1/4*x*(a^2*x^2+1)^(3/2)+3/8*x*(a^2*x^2+1)^(1/2)+3/8*ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2))-3/a^4*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2)))-3*I/a^5*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))))+1/a^6*(I/a/(x-I/a)^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)-2*I*a*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))))))

Maxima [A]

time = 0.48, size = 216, normalized size = 1.58

$$\frac{(a^2x^2+1)^{\frac{3}{2}}}{a^6x^2-2ia^3x-a^4} + \frac{3(a^2x^2+1)^{\frac{3}{2}}}{2ia^5x+2a^4} + \frac{6\sqrt{a^2x^2+1}}{ia^5x+a^4} + \frac{i(a^2x^2+1)^{\frac{3}{2}}}{4a^3} + \frac{3i\sqrt{a^2x^2+1}x}{8a^3} - \frac{3i\sqrt{-a^2x^2+4iax+3}x}{2a^3} - \frac{(a^2x^2+1)^{\frac{3}{2}}}{a^4} + \frac{3i \arcsin(ax+2)}{2a^4} + \frac{63i \operatorname{arsinh}(ax)}{8a^4} + \frac{9\sqrt{a^2x^2+1}}{2a^4} - \frac{3\sqrt{-a^2x^2+4iax+3}}{a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="maxima")

[Out] (a^2*x^2 + 1)^(3/2)/(a^6*x^2 - 2*I*a^5*x - a^4) + 3*(a^2*x^2 + 1)^(3/2)/(2*I*a^5*x + 2*a^4) + 6*sqrt(a^2*x^2 + 1)/(I*a^5*x + a^4) + 1/4*I*(a^2*x^2 + 1)^(3/2)*x/a^3 + 3/8*I*sqrt(a^2*x^2 + 1)*x/a^3 - 3/2*I*sqrt(-a^2*x^2 + 4*I*a*x + 3)*x/a^3 - (a^2*x^2 + 1)^(3/2)/a^4 + 3/2*I*arcsin(I*a*x + 2)/a^4 + 63/8*I*arcsinh(a*x)/a^4 + 9/2*sqrt(a^2*x^2 + 1)/a^4 - 3*sqrt(-a^2*x^2 + 4*I*a*x + 3)/a^4

Fricas [A]

time = 1.84, size = 88, normalized size = 0.64

$$\frac{-32i ax - 51 (i ax + 1) \log(-ax + \sqrt{a^2 x^2 + 1}) + (2i a^4 x^4 - 6 a^3 x^3 - 11i a^2 x^2 + 29 ax - 80i) \sqrt{a^2 x^2 + 1} - 32}{8(a^5 x - i a^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="fricas")

[Out] 1/8*(-32*I*a*x - 51*(I*a*x + 1)*log(-a*x + sqrt(a^2*x^2 + 1)) + (2*I*a^4*x^4 - 6*a^3*x^3 - 11*I*a^2*x^2 + 29*a*x - 80*I)*sqrt(a^2*x^2 + 1) - 32)/(a^5*x - I*a^4)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{x^3 \sqrt{a^2 x^2 + 1}}{a^3 x^3 - 3i a^2 x^2 - 3ax + i} dx + \int \frac{a^2 x^5 \sqrt{a^2 x^2 + 1}}{a^3 x^3 - 3i a^2 x^2 - 3ax + i} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3/(1+I*a*x)**3*(a**2*x**2+1)**(3/2),x)

[Out] I*(Integral(x**3*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x) + Integral(a**2*x**5*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x))

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 0.47, size = 138, normalized size = 1.01

$$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{4}{(a^2)^{3/2}} + \frac{2\sqrt{a^2}}{a^4} - \frac{x^2 \sqrt{a^2}}{a^2} + \frac{x^3 (a^2)^{3/2} 1i}{4a^3} - \frac{x \sqrt{a^2} 19i}{8a^3} \right)}{\sqrt{a^2}} + \frac{\operatorname{asinh}(x \sqrt{a^2}) 51i}{8a^3 \sqrt{a^2}} + \frac{\sqrt{a^2 x^2 + 1} 4i}{a^3 \left(-x \sqrt{a^2} + \frac{\sqrt{a^2} 1i}{a} \right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^3*(a^2*x^2 + 1)^(3/2))/(a*x*1i + 1)^3,x)`

[Out] `((a^2*x^2 + 1)^(1/2)*(4/(a^2)^(3/2) + (2*(a^2)^(1/2))/a^4 - (x^2*(a^2)^(1/2))/a^2 + (x^3*(a^2)^(3/2)*1i)/(4*a^3) - (x*(a^2)^(1/2)*19i)/(8*a^3)))/(a^2)^(1/2) + (asinh(x*(a^2)^(1/2))*51i)/(8*a^3*(a^2)^(1/2)) + ((a^2*x^2 + 1)^(1/2)*4i)/(a^3*((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2))`

3.53 $\int e^{-3i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=102

$$-\frac{i(1-iax)^3}{a^3\sqrt{1+a^2x^2}} - \frac{i(3-iax)^2\sqrt{1+a^2x^2}}{3a^3} - \frac{(28i+3ax)\sqrt{1+a^2x^2}}{6a^3} + \frac{11\sinh^{-1}(ax)}{2a^3}$$

[Out] $11/2*\text{arcsinh}(a*x)/a^3 - I*(1-I*a*x)^3/a^3/(a^2*x^2+1)^{(1/2)} - 1/3*I*(3-I*a*x)^2*(a^2*x^2+1)^{(1/2)}/a^3 - 1/6*(28*I+3*a*x)*(a^2*x^2+1)^{(1/2)}/a^3$

Rubi [A]

time = 0.40, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {5168, 1647, 1607, 12, 866, 1649, 1668, 794, 221}

$$\frac{11\sinh^{-1}(ax)}{2a^3} - \frac{i(1-iax)^3}{a^3\sqrt{a^2x^2+1}} - \frac{i(3-iax)^2\sqrt{a^2x^2+1}}{3a^3} - \frac{(3ax+28i)\sqrt{a^2x^2+1}}{6a^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{((3*I)*\text{ArcTan}[a*x])}, x]$

[Out] $((-I)*(1-I*a*x)^3)/(a^3*\text{Sqrt}[1+a^2*x^2]) - ((I/3)*(3-I*a*x)^2*\text{Sqrt}[1+a^2*x^2])/a^3 - ((28*I+3*a*x)*\text{Sqrt}[1+a^2*x^2])/(6*a^3) + (11*\text{ArcSinh}[a*x])/(2*a^3)$

Rule 12

$\text{Int}[(a_*)(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \&\& \text{!MatchQ}[u, (b_*)(v_)] /; \text{FreeQ}[b, x]$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_*) + (b_*)(x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{GtQ}[a, 0] \&\& \text{PosQ}[b]$

Rule 794

$\text{Int}[((d_*) + (e_*)(x_))*((f_*) + (g_*)(x_))*((a_*) + (c_*)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[((e*f + d*g)*(2*p + 3) + 2*e*g*(p + 1)*x)*((a + c*x^2)^{(p + 1})/(2*c*(p + 1)*(2*p + 3))), x] - \text{Dist}[(a*e*g - c*d*f*(2*p + 3))/(c*(2*p + 3)), \text{Int}[(a + c*x^2)^p, x], x] /; \text{FreeQ}[\{a, c, d, e, f, g, p\}, x] \&\& \text{!LeQ}[p, -1]$

Rule 866

$\text{Int}[((d_*) + (e_*)(x_))^{(m_)*}((f_*) + (g_*)(x_))^{(n_)*}((a_*) + (c_*)(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[d^{(2*m)}/a^m, \text{Int}[(f + g*x)^n*((a + c*x^2)^{(m + p)})]$

```
/(d - e*x)^m), x], x] /; FreeQ[{a, c, d, e, f, g, n, p}, x] && NeQ[e*f - d*
g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[f, 0] && ILtQ[m, -1]
&& !(IGtQ[n, 0] && ILtQ[m + n, 0] && !GtQ[p, 1])
```

Rule 1607

```
Int[(u_.)*((a_.)*(x_)^(p_.) + (b_.)*(x_)^(q_.))^(n_.), x_Symbol] := Int[u*x
^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] &&
PosQ[q - p]
```

Rule 1647

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[d*e, Int[(d + e*x)^(m - 1)*PolynomialQuotient[Pq, a*e + c*d*x, x]*(
a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && PolyQ[Pq, x]
&& EqQ[c*d^2 + a*e^2, 0] && EqQ[PolynomialRemainder[Pq, a*e + c*d*x, x], 0]
```

Rule 1649

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :
> With[{Q = PolynomialQuotient[Pq, a*e + c*d*x, x], f = PolynomialRemainder
[Pq, a*e + c*d*x, x]}, Simp[(-d)*f*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*a*e*
(p + 1))), x] + Dist[d/(2*a*(p + 1)), Int[(d + e*x)^(m - 1)*(a + c*x^2)^(p
+ 1)*ExpandToSum[2*a*e*(p + 1)*Q + f*(m + 2*p + 2), x], x], x] /; FreeQ[{a
, c, d, e}, x] && PolyQ[Pq, x] && EqQ[c*d^2 + a*e^2, 0] && ILtQ[p + 1/2, 0]
&& GtQ[m, 0]
```

Rule 1668

```
Int[(Pq_)*((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :
> With[{q = Expon[Pq, x], f = Coeff[Pq, x, Expon[Pq, x]]}, Simp[f*(d + e*x)
^(m + q - 1)*((a + c*x^2)^(p + 1)/(c*e^(q - 1)*(m + q + 2*p + 1))), x] + Di
st[1/(c*e^q*(m + q + 2*p + 1)), Int[(d + e*x)^m*(a + c*x^2)^p*ExpandToSum[c
*e^q*(m + q + 2*p + 1)*Pq - c*f*(m + q + 2*p + 1)*(d + e*x)^q - f*(d + e*x)
^(q - 2)*(a*e^2*(m + q - 1) - c*d^2*(m + q + 2*p + 1) - 2*c*d*e*(m + q + p)
*x), x], x], x] /; GtQ[q, 1] && NeQ[m + q + 2*p + 1, 0] /; FreeQ[{a, c, d,
e, m, p}, x] && PolyQ[Pq, x] && NeQ[c*d^2 + a*e^2, 0] && !(EqQ[d, 0] && T
rue) && !(IGtQ[m, 0] && RationalQ[a, c, d, e] && (IntegerQ[p] || ILtQ[p +
1/2, 0]))
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)^(n_)])*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1-iax)^2}{(1+iax)\sqrt{1+a^2x^2}} dx \\
&= (ia) \int \frac{\sqrt{1+a^2x^2} \left(-\frac{ix^2}{a} - x^3\right)}{(1+iax)^2} dx \\
&= (ia) \int \frac{\left(-\frac{i}{a} - x\right) x^2 \sqrt{1+a^2x^2}}{(1+iax)^2} dx \\
&= a^2 \int \frac{x^2(1+a^2x^2)^{3/2}}{a^2(1+iax)^3} dx \\
&= \int \frac{x^2(1+a^2x^2)^{3/2}}{(1+iax)^3} dx \\
&= \int \frac{x^2(1-iax)^3}{(1+a^2x^2)^{3/2}} dx \\
&= -\frac{i(1-iax)^3}{a^3\sqrt{1+a^2x^2}} - \int \frac{\left(-\frac{3}{a^2} + \frac{ix}{a}\right) (1-iax)^2}{\sqrt{1+a^2x^2}} dx \\
&= -\frac{i(1-iax)^3}{a^3\sqrt{1+a^2x^2}} - \frac{i(3-iax)^2\sqrt{1+a^2x^2}}{3a^3} + \frac{1}{3} \int \frac{\left(-\frac{3}{a^2} + \frac{ix}{a}\right) (-5+3iax)}{\sqrt{1+a^2x^2}} dx \\
&= -\frac{i(1-iax)^3}{a^3\sqrt{1+a^2x^2}} - \frac{i(3-iax)^2\sqrt{1+a^2x^2}}{3a^3} - \frac{(28i+3ax)\sqrt{1+a^2x^2}}{6a^3} + \frac{11}{2a^2} \int \frac{1}{\sqrt{1+a^2x^2}} dx \\
&= -\frac{i(1-iax)^3}{a^3\sqrt{1+a^2x^2}} - \frac{i(3-iax)^2\sqrt{1+a^2x^2}}{3a^3} - \frac{(28i+3ax)\sqrt{1+a^2x^2}}{6a^3} + \frac{11 \sinh^{-1}(ax)}{2a^3}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 63, normalized size = 0.62

$$\frac{\sqrt{1+a^2x^2} \frac{(-52-19iax-7a^2x^2+2ia^3x^3)}{-i+ax} + 33 \sinh^{-1}(ax)}{6a^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^((3*I)*ArcTan[a*x]), x]

[Out] ((Sqrt[1 + a^2*x^2]*(-52 - (19*I)*a*x - 7*a^2*x^2 + (2*I)*a^3*x^3))/(-I + a*x) + 33*ArcSinh[a*x])/(6*a^3)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 618 vs. 2(85) = 170.

time = 0.12, size = 619, normalized size = 6.07

method	result
risch	$\frac{i(2a^2x^2+9iax-28)\sqrt{a^2x^2+1}}{6a^3} + \frac{11\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a^2\sqrt{a^2}} - \frac{4\sqrt{\left(x-\frac{i}{a}\right)^2a^2+2ia\left(x-\frac{i}{a}\right)}}{a^4\left(x-\frac{i}{a}\right)}$
default	$\frac{i\left(\frac{\left(x-\frac{i}{a}\right)^2a^2+2ia\left(x-\frac{i}{a}\right)}{3}\right)^{\frac{3}{2}} + ia\left(\frac{2\left(x-\frac{i}{a}\right)a^2+2ia}{4a^2}\sqrt{\left(x-\frac{i}{a}\right)^2a^2+2ia\left(x-\frac{i}{a}\right)} + \frac{\ln\left(\frac{ia+\left(x-\frac{i}{a}\right)a^2}{\sqrt{a^2}} + \sqrt{\left(x-\frac{i}{a}\right)^2a^2}\right)}{2\sqrt{a^2}}\right)}{a^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{I}{a^3} \left(\frac{1}{3} \left(\left(x - \frac{I}{a} \right)^2 a^2 + 2Ia \left(x - \frac{I}{a} \right) \right)^{\frac{3}{2}} + Ia \left(\frac{2 \left(x - \frac{I}{a} \right) a^2 + 2ia}{4a^2} \sqrt{\left(x - \frac{I}{a} \right)^2 a^2 + 2ia \left(x - \frac{I}{a} \right)} + \frac{\ln \left(\frac{ia + \left(x - \frac{I}{a} \right) a^2}{\sqrt{a^2}} + \sqrt{\left(x - \frac{I}{a} \right)^2 a^2} \right)}{2\sqrt{a^2}} \right) \right)$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 181 vs. $2(80) = 160$.
time = 0.49, size = 181, normalized size = 1.77

$$\frac{i(a^2x^2+1)^{\frac{3}{2}}}{a^5x^2-2ia^4x-a^3} - \frac{i(a^2x^2+1)^{\frac{3}{2}}}{ia^4x+a^3} - \frac{6i\sqrt{a^2x^2+1}}{ia^4x+a^3} - \frac{\sqrt{-a^2x^2+4iax+3}x}{2a^2} + \frac{i(a^2x^2+1)^{\frac{3}{2}}}{3a^3} + \frac{\arcsin(iax+2)}{2a^3} + \frac{6\operatorname{arsinh}(ax)}{a^3} - \frac{3i\sqrt{a^2x^2+1}}{a^3} + \frac{i\sqrt{-a^2x^2+4iax+3}}{a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out]
$$-I(a^2x^2+1)^{\frac{3}{2}}/(a^5x^2-2Ia^4x-a^3) - I(a^2x^2+1)^{\frac{3}{2}}/(Ia^4x+a^3) - 6I\sqrt{a^2x^2+1}/(Ia^4x+a^3) - 1/2\sqrt{-a^2x^2+4Ia^4x+3}x/a^2 + 1/3I(a^2x^2+1)^{\frac{3}{2}}/a^3 + 1/2\arcsin(Ia^4x+2)/a^3 + 6\operatorname{arsinh}(a^4x)/a^3 - 3I\sqrt{a^2x^2+1}/a^3 + I\sqrt{-a^2x^2+4Ia^4x+3}/a^3$$

Fricas [A]

time = 2.06, size = 80, normalized size = 0.78

$$\frac{24ax + 33(ax - i) \log\left(-ax + \sqrt{a^2x^2 + 1}\right) - (2ia^3x^3 - 7a^2x^2 - 19iax - 52)\sqrt{a^2x^2 + 1} - 24i}{6(a^4x - ia^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="fricas")

[Out] $-1/6*(24*a*x + 33*(a*x - I)*\log(-a*x + \sqrt{a^2*x^2 + 1}) - (2*I*a^3*x^3 - 7*a^2*x^2 - 19*I*a*x - 52)*\sqrt{a^2*x^2 + 1} - 24*I)/(a^4*x - I*a^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{x^2 \sqrt{a^2 x^2 + 1}}{a^3 x^3 - 3i a^2 x^2 - 3a x + i} dx + \int \frac{a^2 x^4 \sqrt{a^2 x^2 + 1}}{a^3 x^3 - 3i a^2 x^2 - 3a x + i} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*a*x)**3*(a**2*x**2+1)**(3/2),x)

[Out] $I*(\text{Integral}(x**2*\sqrt{a**2*x**2 + 1}/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x) + \text{Integral}(a**2*x**4*\sqrt{a**2*x**2 + 1}/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x))$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.07, size = 115, normalized size = 1.13

$$\frac{11 \operatorname{asinh}(x \sqrt{a^2})}{2 a^2 \sqrt{a^2}} - \frac{\sqrt{a^2 x^2 + 1} \left(\frac{3x \sqrt{a^2}}{2 a^2} + \frac{a^{14} i}{3 (a^2)^{3/2}} - \frac{a^3 x^2 1 i}{3 (a^2)^{3/2}} \right)}{\sqrt{a^2}} + \frac{4 \sqrt{a^2 x^2 + 1}}{a^2 \left(-x \sqrt{a^2} + \frac{\sqrt{a^2} 1 i}{a} \right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a^2*x^2 + 1)^(3/2))/(a*x*1i + 1)^3,x)

[Out] $(11*\operatorname{asinh}(x*(a^2)^{(1/2)}))/(2*a^2*(a^2)^{(1/2)}) - ((a^2*x^2 + 1)^{(1/2)}*((a^{14} i)/(3*(a^2)^{(3/2)}) - (a^3*x^2*1i)/(3*(a^2)^{(3/2)}) + (3*x*(a^2)^{(1/2)}))/(2*a^2)))/(a^2)^{(1/2)} + (4*(a^2*x^2 + 1)^{(1/2)})/(a^2*((a^2)^{(1/2)}*1i)/a - x*(a^2)^{(1/2)})*(a^2)^{(1/2)}$

3.54 $\int e^{-3i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=92

$$-\frac{9\sqrt{1+a^2x^2}}{2a^2} - \frac{3(1+a^2x^2)^{3/2}}{2a^2(1+iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1+iax)^3} - \frac{9i \sinh^{-1}(ax)}{2a^2}$$

[Out] $-3/2*(a^2*x^2+1)^{(3/2)}/a^2/(1+I*a*x)-(a^2*x^2+1)^{(5/2)}/a^2/(1+I*a*x)^3-9/2*I*\text{arcsinh}(a*x)/a^2-9/2*(a^2*x^2+1)^{(1/2)}/a^2$

Rubi [A]

time = 0.23, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 7, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.583$, Rules used = {5168, 1647, 1607, 12, 807, 679, 221}

$$-\frac{(a^2x^2+1)^{5/2}}{a^2(1+iax)^3} - \frac{3(a^2x^2+1)^{3/2}}{2a^2(1+iax)} - \frac{9\sqrt{a^2x^2+1}}{2a^2} - \frac{9i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

[In] `Int[x/E^((3*I)*ArcTan[a*x]),x]`

[Out] $(-9*\text{Sqrt}[1+a^2*x^2])/(2*a^2) - (3*(1+a^2*x^2)^{(3/2)})/(2*a^2*(1+I*a*x)) - (1+a^2*x^2)^{(5/2)}/(a^2*(1+I*a*x)^3) - (((9*I)/2)*\text{ArcSinh}[a*x])/a^2$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 221

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 679

`Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d + e*x)^(m + 1)*((a + c*x^2)^p/(e*(m + 2*p + 1))), x] - Dist[2*c*d*(p/(e^2*(m + 2*p + 1))), Int[(d + e*x)^(m + 1)*(a + c*x^2)^(p - 1), x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 + a*e^2, 0] && GtQ[p, 0] && (LeQ[-2, m, 0] || EqQ[m + p + 1, 0]) && NeQ[m + 2*p + 1, 0] && IntegerQ[2*p]`

Rule 807

`Int[((d_) + (e_.)*(x_))^(m_)*((f_.) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[(d*g - e*f)*(d + e*x)^m*(a + c*x^2)^(p + 1)/(2*c*d*(m + 1)), x]`


```

+ p + 1))), x] + Dist[(m*(g*c*d + c*e*f) + 2*e*c*f*(p + 1))/(e*(2*c*d)*(m
+ p + 1)), Int[(d + e*x)^(m + 1)*(a + c*x^2)^p, x], x] /; FreeQ[{a, c, d, e
, f, g, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && ((LtQ[m, -1] && !IGtQ[m + p
+ 1, 0]) || (LtQ[m, 0] && LtQ[p, -1]) || EqQ[m + 2*p + 2, 0]) && NeQ[m + p
+ 1, 0]

```

Rule 1607

```

Int[(u_.)*((a_.)*(x_)^(p_.) + (b_.)*(x_)^(q_.))^(n_.), x_Symbol] := Int[u*x
^(n*p)*(a + b*x^(q - p))^n, x] /; FreeQ[{a, b, p, q}, x] && IntegerQ[n] &&
PosQ[q - p]

```

Rule 1647

```

Int[(Pq_)*((d_) + (e_.)*(x_)^(m_.))*((a_) + (c_.)*(x_)^2)^(p_.), x_Symbol]
:= Dist[d*e, Int[(d + e*x)^(m - 1)*PolynomialQuotient[Pq, a*e + c*d*x, x]*(
a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, m, p}, x] && PolyQ[Pq, x]
&& EqQ[c*d^2 + a*e^2, 0] && EqQ[PolynomialRemainder[Pq, a*e + c*d*x, x], 0]

```

Rule 5168

```

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(ax)} x dx &= \int \frac{x(1-iax)^2}{(1+iax)\sqrt{1+a^2x^2}} dx \\
&= (ia) \int \frac{\left(-\frac{ix}{a} - x^2\right) \sqrt{1+a^2x^2}}{(1+iax)^2} dx \\
&= (ia) \int \frac{\left(-\frac{i}{a} - x\right) x \sqrt{1+a^2x^2}}{(1+iax)^2} dx \\
&= a^2 \int \frac{x(1+a^2x^2)^{3/2}}{a^2(1+iax)^3} dx \\
&= \int \frac{x(1+a^2x^2)^{3/2}}{(1+iax)^3} dx \\
&= \frac{(1+a^2x^2)^{5/2}}{a^2(1+iax)^3} - \frac{(3i) \int \frac{(1+a^2x^2)^{3/2}}{(1+iax)^2} dx}{a} \\
&= -\frac{3(1+a^2x^2)^{3/2}}{2a^2(1+iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1+iax)^3} - \frac{(9i) \int \frac{\sqrt{1+a^2x^2}}{1+iax} dx}{2a} \\
&= -\frac{9\sqrt{1+a^2x^2}}{2a^2} - \frac{3(1+a^2x^2)^{3/2}}{2a^2(1+iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1+iax)^3} - \frac{(9i) \int \frac{1}{\sqrt{1+a^2x^2}} dx}{2a} \\
&= -\frac{9\sqrt{1+a^2x^2}}{2a^2} - \frac{3(1+a^2x^2)^{3/2}}{2a^2(1+iax)} - \frac{(1+a^2x^2)^{5/2}}{a^2(1+iax)^3} - \frac{9i \sinh^{-1}(ax)}{2a^2}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 60, normalized size = 0.65

$$\sqrt{1+a^2x^2} \left(-\frac{3}{a^2} + \frac{ix}{2a} + \frac{4i}{a^2(-i+ax)} \right) - \frac{9i \sinh^{-1}(ax)}{2a^2}$$

Antiderivative was successfully verified.

`[In] Integrate[x/E^((3*I)*ArcTan[a*x]),x]`

```
[Out] Sqrt[1 + a^2*x^2]*(-3/a^2 + ((I/2)*x)/a + (4*I)/(a^2*(-I + a*x))) - (((9*I)/2)*ArcSinh[a*x])/a^2
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 462 vs. $2(77) = 154$.

time = 0.10, size = 463, normalized size = 5.03

method	result
--------	--------

risch	$\frac{i(ax+6i)\sqrt{a^2x^2+1}}{2a^2} - \frac{9i \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{2a\sqrt{a^2}} + \frac{4i\sqrt{\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)}}{a^3\left(x-\frac{i}{a}\right)}$
default	$i \left(-\frac{i\left(\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)\right)^{\frac{5}{2}}}{a\left(x-\frac{i}{a}\right)^2} + 3ia \left(\frac{\left(\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)\right)^{\frac{3}{2}}}{3} + ia \left(\frac{\left(2\left(x-\frac{i}{a}\right)a^2 + 2ia\right)\sqrt{\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)}}{4a^2} \right) \right) \right) + \frac{\quad}{a^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & I/a^3 \cdot (-I/a/(x-I/a)^2 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{5/2} + 3Ia \cdot (1/3 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{3/2} + Ia \cdot (1/4 \cdot (2 \cdot (x-I/a) \cdot a^2 + 2Ia) / a^2 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{1/2} + 1/2 \cdot \ln((Ia + (x-I/a) \cdot a^2) / (a^2)^{1/2} + ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{1/2})) / (a^2)^{1/2})) - 1/a^4 \cdot (I/a/(x-I/a)^3 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{5/2} - 2Ia \cdot (-I/a/(x-I/a)^2 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{5/2} + 3Ia \cdot (1/3 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{3/2} + Ia \cdot (1/4 \cdot (2 \cdot (x-I/a) \cdot a^2 + 2Ia) / a^2 \cdot ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{1/2} + 1/2 \cdot \ln((Ia + (x-I/a) \cdot a^2) / (a^2)^{1/2} + ((x-I/a)^2 a^2 + 2Ia(x-I/a))^{1/2})) / (a^2)^{1/2})) \end{aligned}$$

Maxima [A]

time = 0.47, size = 112, normalized size = 1.22

$$-\frac{(a^2x^2+1)^{\frac{3}{2}}}{a^4x^2-2ia^3x-a^2} - \frac{(a^2x^2+1)^{\frac{3}{2}}}{2ia^3x+2a^2} - \frac{6\sqrt{a^2x^2+1}}{ia^3x+a^2} - \frac{9i \operatorname{arsinh}(ax)}{2a^2} - \frac{3\sqrt{a^2x^2+1}}{2a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out]
$$-(a^2x^2+1)^{3/2}/(a^4x^2-2Ia^3x-a^2) - (a^2x^2+1)^{3/2}/(2Ia^3x+2a^2) - 6\sqrt{a^2x^2+1}/(Ia^3x+a^2) - 9/2I \operatorname{arcsinh}(ax)/a^2 - 3/2\sqrt{a^2x^2+1}/a^2$$

Fricas [A]

time = 2.56, size = 72, normalized size = 0.78

$$\frac{8iax - 9(-iax - 1) \log\left(-ax + \sqrt{a^2x^2+1}\right) + \sqrt{a^2x^2+1}(ia^2x^2 - 5ax + 14i) + 8}{2(a^3x - ia^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="fricas")`

[Out] $\frac{1}{2}*(8*I*a*x - 9*(-I*a*x - 1)*\log(-a*x + \sqrt{a^2*x^2 + 1}) + \sqrt{a^2*x^2 + 1}*(I*a^2*x^2 - 5*a*x + 14*I) + 8)/(a^3*x - I*a^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{x\sqrt{a^2x^2+1}}{a^3x^3-3ia^2x^2-3ax+i} dx + \int \frac{a^2x^3\sqrt{a^2x^2+1}}{a^3x^3-3ia^2x^2-3ax+i} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*a*x)**3*(a**2*x**2+1)**(3/2),x)`

[Out] `I*(Integral(x*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x) + Integral(a**2*x**3*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x))`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="giac")`

[Out] `undef`

Mupad [B]

time = 0.42, size = 105, normalized size = 1.14

$$\frac{\sqrt{a^2x^2+1} \left(\frac{3\sqrt{a^2}}{a^2} - \frac{x\sqrt{a^2}}{2a} \right)}{\sqrt{a^2}} - \frac{\operatorname{asinh}\left(x\sqrt{a^2}\right) 9i}{2a\sqrt{a^2}} - \frac{\sqrt{a^2x^2+1} 4i}{a \left(-x\sqrt{a^2} + \frac{\sqrt{a^2}}{a} \right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x*(a^2*x^2 + 1)^(3/2))/(a*x*I + 1)^3,x)`

[Out] `-((a^2*x^2 + 1)^(1/2)*((3*(a^2)^(1/2))/a^2 - (x*(a^2)^(1/2)*I)/(2*a)))/(a^2)^(1/2) - (asinh(x*(a^2)^(1/2))*9i)/(2*a*(a^2)^(1/2)) - ((a^2*x^2 + 1)^(1/2)*4i)/(a*((a^2)^(1/2)*I)/a - x*(a^2)^(1/2))*(a^2)^(1/2))`

3.55 $\int e^{-3i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=60

$$\frac{2i(1-iax)^2}{a\sqrt{1+a^2x^2}} + \frac{3i\sqrt{1+a^2x^2}}{a} - \frac{3\sinh^{-1}(ax)}{a}$$

[Out] $-3*\text{arcsinh}(a*x)/a+2*I*(1-I*a*x)^2/a/(a^2*x^2+1)^{(1/2)}+3*I*(a^2*x^2+1)^{(1/2)}/a$

Rubi [A]

time = 0.03, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5167, 867, 683, 655, 221}

$$\frac{2i(1-iax)^2}{a\sqrt{a^2x^2+1}} + \frac{3i\sqrt{a^2x^2+1}}{a} - \frac{3\sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(-3*I)*\text{ArcTan}[a*x]}, x]$

[Out] $((2*I)*(1-I*a*x)^2)/(a*\text{Sqrt}[1+a^2*x^2]) + ((3*I)*\text{Sqrt}[1+a^2*x^2])/a - (3*\text{ArcSinh}[a*x])/a$

Rule 221

$\text{Int}[1/\text{Sqrt}[(a_) + (b_)*(x_)^2], x_Symbol] \rightarrow \text{Simp}[\text{ArcSinh}[\text{Rt}[b, 2]*(x/\text{Sqrt}[a])]/\text{Rt}[b, 2], x] /;$ FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 655

$\text{Int}[((d_) + (e_)*(x_))*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*((a + c*x^2)^{(p+1)}/(2*c*(p+1))), x] + \text{Dist}[d, \text{Int}[(a + c*x^2)^p, x], x] /;$ FreeQ[{a, c, d, e, p}, x] && NeQ[p, -1]

Rule 683

$\text{Int}[((d_) + (e_)*(x_))^{(m_)}*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*(d + e*x)^{(m-1)}*((a + c*x^2)^{(p+1)}/(c*(p+1))), x] - \text{Dist}[e^2*((m+p)/(c*(p+1))), \text{Int}[(d + e*x)^{(m-2)}*(a + c*x^2)^{(p+1)}, x], x] /;$ FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && GtQ[m, 1] && IntegerQ[2*p]

Rule 867

$\text{Int}[((d_) + (e_)*(x_))^{(m_)}*((f_) + (g_)*(x_))^{(n_)}*((a_) + (c_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[d^{(2*m)}/a^m, \text{Int}[(f + g*x)^n*(a + c*x^2)^{(m+p)}/$

$(d - e*x)^m, x], x] /;$ FreeQ[{a, c, d, e, f, g, n, p}, x] && NeQ[e*f - d*g, 0] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && ILtQ[m, 0] && IntegerQ[n]

Rule 5167

Int[E^(ArcTan[(a_.)*(x_)]*(n_)), x_Symbol] := Int[(1 - I*a*x)^((I*n + 1)/2) / ((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[a, x] && IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-3i \tan^{-1}(ax)} dx &= \int \frac{(1 - iax)^2}{(1 + iax)\sqrt{1 + a^2x^2}} dx \\ &= \int \frac{(1 - iax)^3}{(1 + a^2x^2)^{3/2}} dx \\ &= \frac{2i(1 - iax)^2}{a\sqrt{1 + a^2x^2}} - 3 \int \frac{1 - iax}{\sqrt{1 + a^2x^2}} dx \\ &= \frac{2i(1 - iax)^2}{a\sqrt{1 + a^2x^2}} + \frac{3i\sqrt{1 + a^2x^2}}{a} - 3 \int \frac{1}{\sqrt{1 + a^2x^2}} dx \\ &= \frac{2i(1 - iax)^2}{a\sqrt{1 + a^2x^2}} + \frac{3i\sqrt{1 + a^2x^2}}{a} - \frac{3 \sinh^{-1}(ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 42, normalized size = 0.70

$$\frac{\sqrt{1 + a^2x^2} \left(i + \frac{4}{-i+ax} \right)}{a} - \frac{3 \sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((-3*I)*ArcTan[a*x]), x]

[Out] (Sqrt[1 + a^2*x^2]*(I + 4/(-I + a*x)))/a - (3*ArcSinh[a*x])/a

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 256 vs. 2(53) = 106.

time = 0.11, size = 257, normalized size = 4.28

method	result
risch	$\frac{i\sqrt{a^2x^2+1}}{a} - \frac{3 \ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{\sqrt{a^2}} + \frac{4\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia\left(x - \frac{i}{a}\right)}}{a^2\left(x - \frac{i}{a}\right)}$

default	$i \left(\frac{i \left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{5}{2}}}{a \left(x - \frac{i}{a} \right)^3} - 2ia \left(- \frac{i \left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{5}{2}}}{a \left(x - \frac{i}{a} \right)^2} + 3ia \left(\frac{\left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{3}{2}}}{3} + ia \left(\frac{2 \left(x - \frac{i}{a} \right) a^2 + 2ia}{a^3} \sqrt{\left(x - \frac{i}{a} \right)^2 a^2 + 2ia} \right) \right)$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $I/a^3*(I/a/(x-I/a)^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)-2*I*a*(-I/a/(x-I/a))^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))))$

Maxima [A]

time = 0.47, size = 65, normalized size = 1.08

$$\frac{i(a^2x^2 + 1)^{\frac{3}{2}}}{a^3x^2 - 2ia^2x - a} - \frac{3 \operatorname{arsinh}(ax)}{a} + \frac{6i\sqrt{a^2x^2 + 1}}{ia^2x + a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out] $I*(a^2*x^2 + 1)^(3/2)/(a^3*x^2 - 2*I*a^2*x - a) - 3*\operatorname{arcsinh}(a*x)/a + 6*I*\operatorname{sqrt}(a^2*x^2 + 1)/(I*a^2*x + a)$

Fricas [A]

time = 2.63, size = 60, normalized size = 1.00

$$\frac{4ax + 3(ax - i) \log(-ax + \sqrt{a^2x^2 + 1}) + \sqrt{a^2x^2 + 1}(iax + 5) - 4i}{a^2x - ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="fricas")`

[Out] $(4*a*x + 3*(a*x - I)*\log(-a*x + \operatorname{sqrt}(a^2*x^2 + 1)) + \operatorname{sqrt}(a^2*x^2 + 1)*(I*a*x + 5) - 4*I)/(a^2*x - I*a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2x^2 + 1}}{a^3x^3 - 3ia^2x^2 - 3ax + i} dx + \int \frac{a^2x^2\sqrt{a^2x^2 + 1}}{a^3x^3 - 3ia^2x^2 - 3ax + i} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2),x)

[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.42, size = 73, normalized size = 1.22

$$\frac{\sqrt{a^2 x^2 + 1} \operatorname{li}}{a} - \frac{3 \operatorname{asinh}\left(x \sqrt{a^2}\right)}{\sqrt{a^2}} - \frac{4 \sqrt{a^2 x^2 + 1}}{\left(-x \sqrt{a^2} + \frac{\sqrt{a^2} \operatorname{li}}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(a*x*1i + 1)^3,x)

[Out] ((a^2*x^2 + 1)^(1/2)*1i)/a - (3*asinh(x*(a^2)^(1/2)))/(a^2)^(1/2) - (4*(a^2*x^2 + 1)^(1/2))/((((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2))

$$3.56 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=52

$$\frac{4i\sqrt{1+a^2x^2}}{i-ax} + i \sinh^{-1}(ax) - \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] I*arcsinh(a*x)-arctanh((a^2*x^2+1)^(1/2))+4*I*(a^2*x^2+1)^(1/2)/(I-a*x)

Rubi [A]

time = 0.47, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5168, 6874, 221, 272, 65, 214, 665}

$$\frac{4i\sqrt{a^2x^2+1}}{-ax+i} - \tanh^{-1}\left(\sqrt{a^2x^2+1}\right) + i \sinh^{-1}(ax)$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a*x])*x),x]

[Out] ((4*I)*Sqrt[1+a^2*x^2])/(I-a*x) + I*ArcSinh[a*x] - ArcTanh[Sqrt[1+a^2*x^2]]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^n, x], x, (a+b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c-a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m+1)/n]]

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d,
e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2,
0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2])), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1 - iax)^2}{x(1 + iax)\sqrt{1 + a^2x^2}} dx \\
&= \int \left(\frac{ia}{\sqrt{1 + a^2x^2}} + \frac{1}{x\sqrt{1 + a^2x^2}} - \frac{4a}{(-i + ax)\sqrt{1 + a^2x^2}} \right) dx \\
&= (ia) \int \frac{1}{\sqrt{1 + a^2x^2}} dx - (4a) \int \frac{1}{(-i + ax)\sqrt{1 + a^2x^2}} dx + \int \frac{1}{x\sqrt{1 + a^2x^2}} dx \\
&= \frac{4i\sqrt{1 + a^2x^2}}{i - ax} + i \sinh^{-1}(ax) + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
&= \frac{4i\sqrt{1 + a^2x^2}}{i - ax} + i \sinh^{-1}(ax) + \frac{\text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2x^2} \right)}{a^2} \\
&= \frac{4i\sqrt{1 + a^2x^2}}{i - ax} + i \sinh^{-1}(ax) - \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 1.06

$$-\frac{4i\sqrt{1 + a^2x^2}}{-i + ax} + i \sinh^{-1}(ax) + \log(x) - \log \left(1 + \sqrt{1 + a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x]))*x],x]

[Out] ((-4*I)*Sqrt[1 + a^2*x^2])/(-I + a*x) + I*ArcSinh[a*x] + Log[x] - Log[1 + Sqrt[1 + a^2*x^2]]

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 648 vs. $2(45) = 90$.

time = 0.09, size = 649, normalized size = 12.48

method	result
default	$-\frac{\left(x-\frac{i}{a}\right)^2 a^2+2ia\left(x-\frac{i}{a}\right)^{\frac{3}{2}}}{3}-ia\left(\frac{2\left(x-\frac{i}{a}\right)a^2+2ia\sqrt{\left(x-\frac{i}{a}\right)^2 a^2+2ia\left(x-\frac{i}{a}\right)}}{4a^2}+\frac{\ln\left(\frac{ia+\left(x-\frac{i}{a}\right)a^2+\sqrt{\left(x-\frac{i}{a}\right)^2 a^2+2ia\left(x-\frac{i}{a}\right)}}{\sqrt{a^2}}+\sqrt{\left(x-\frac{i}{a}\right)^2 a^2+2ia\left(x-\frac{i}{a}\right)}\right)}{2}\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x,x,method=_RETURNVERBOSE)

[Out] -1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)-I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))+1/3*(a^2*x^2+1)^(3/2)+(a^2*x^2+1)^(1/2)-arctanh(1/(a^2*x^2+1)^(1/2))+I/a*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))))+1/a^2*(I/a/(x-I/a)^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)-2*I*a*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))))))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x,x, algorithm="maxima")

[Out] integrate((a^2*x^2 + 1)^(3/2)/((I*a*x + 1)^3*x), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(41) = 82$.

time = 5.00, size = 100, normalized size = 1.92

$$\frac{-4i ax - (ax - i) \log(-ax + \sqrt{a^2 x^2 + 1} + 1) + (-i ax - 1) \log(-ax + \sqrt{a^2 x^2 + 1}) + (ax - i) \log(-ax + \sqrt{a^2 x^2 + 1} - 1) - 4i \sqrt{a^2 x^2 + 1} - 4}{ax - i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x,x, algorithm="fricas")

[Out] (-4*I*a*x - (a*x - I)*log(-a*x + sqrt(a^2*x^2 + 1) + 1) + (-I*a*x - 1)*log(-a*x + sqrt(a^2*x^2 + 1)) + (a*x - I)*log(-a*x + sqrt(a^2*x^2 + 1) - 1) - 4*I*sqrt(a^2*x^2 + 1) - 4)/(a*x - I)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2 x^2 + 1}}{a^3 x^4 - 3i a^2 x^3 - 3a x^2 + i x} dx + \int \frac{a^2 x^2 \sqrt{a^2 x^2 + 1}}{a^3 x^4 - 3i a^2 x^3 - 3a x^2 + i x} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/x,x)

[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**3*x**4 - 3*I*a**2*x**3 - 3*a*x**2 + I*x), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**3*x**4 - 3*I*a**2*x**3 - 3*a*x**2 + I*x), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.43, size = 74, normalized size = 1.42

$$-\operatorname{atanh}\left(\sqrt{a^2 x^2 + 1}\right) + \frac{a \operatorname{asinh}\left(x \sqrt{a^2}\right) i}{\sqrt{a^2}} + \frac{a \sqrt{a^2 x^2 + 1} 4i}{\left(-x \sqrt{a^2} + \frac{\sqrt{a^2} i}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(x*(a*x*1i + 1)^3),x)

[Out] (a*asinh(x*(a^2)^(1/2))*1i)/(a^2)^(1/2) - atanh((a^2*x^2 + 1)^(1/2)) + (a*(a^2*x^2 + 1)^(1/2)*4i)/(((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2)

$$3.57 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=64

$$-\frac{\sqrt{1+a^2x^2}}{x} + \frac{4a\sqrt{1+a^2x^2}}{i-ax} + 3ia \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] 3*I*a*arctanh((a^2*x^2+1)^(1/2))-(a^2*x^2+1)^(1/2)/x+4*a*(a^2*x^2+1)^(1/2)/(I-a*x)

Rubi [A]

time = 0.45, antiderivative size = 64, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5168, 6874, 270, 272, 65, 214, 665}

$$\frac{4a\sqrt{a^2x^2+1}}{-ax+i} - \frac{\sqrt{a^2x^2+1}}{x} + 3ia \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a*x])*x^2),x]

[Out] -(Sqrt[1+a^2*x^2]/x) + (4*a*Sqrt[1+a^2*x^2])/(I-a*x) + (3*I)*a*ArcTanh[Sqrt[1+a^2*x^2]]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b)+d*(x^p/b))^n, x], x, (a+b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c-a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 270

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m+1)*((a+b*x^n)^(p+1)/(a*c*(m+1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m+1)/n+p+1, 0] && NeQ[m, -1]

Rule 272

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m+1)/n]-1)*(a+b*x)^p, x], x, x^n], x] /; FreeQ[{a, b

, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]

Rule 665

Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d, e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2, 0]

Rule 5168

Int[E^(ArcTan[(a_.)*(x_)])*(n_)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*n + 1)/2)/((1 + I*a*x)^(I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]

Rule 6874

Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-3i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1 - iax)^2}{x^2(1 + iax)\sqrt{1 + a^2x^2}} dx \\
 &= \int \left(\frac{1}{x^2\sqrt{1 + a^2x^2}} - \frac{3ia}{x\sqrt{1 + a^2x^2}} + \frac{4ia^2}{(-i + ax)\sqrt{1 + a^2x^2}} \right) dx \\
 &= -\left((3ia) \int \frac{1}{x\sqrt{1 + a^2x^2}} dx \right) + (4ia^2) \int \frac{1}{(-i + ax)\sqrt{1 + a^2x^2}} dx + \int \frac{1}{x^2\sqrt{1 + a^2x^2}} \\
 &= -\frac{\sqrt{1 + a^2x^2}}{x} + \frac{4a\sqrt{1 + a^2x^2}}{i - ax} - \frac{1}{2}(3ia) \text{Subst} \left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
 &= -\frac{\sqrt{1 + a^2x^2}}{x} + \frac{4a\sqrt{1 + a^2x^2}}{i - ax} - \frac{(3i) \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2x^2} \right)}{a} \\
 &= -\frac{\sqrt{1 + a^2x^2}}{x} + \frac{4a\sqrt{1 + a^2x^2}}{i - ax} + 3ia \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right)
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 61, normalized size = 0.95

$$\sqrt{1 + a^2x^2} \left(-\frac{1}{x} - \frac{4a}{-i + ax} \right) - 3ia \log(x) + 3ia \log \left(1 + \sqrt{1 + a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x])*x^2),x]

[Out] Sqrt[1 + a^2*x^2]*(-x^(-1) - (4*a)/(-I + a*x)) - (3*I)*a*Log[x] + (3*I)*a*Log[1 + Sqrt[1 + a^2*x^2]]

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 579 vs. 2(56) = 112.

time = 0.10, size = 580, normalized size = 9.06

method	result
risch	$-\frac{\sqrt{a^2x^2+1}}{x} + ia \left(\frac{4i \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}}{a \left(x - \frac{i}{a}\right)} + 3 \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2x^2+1}} \right) \right)$
default	$9ia \left(\frac{\left(\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)\right)^{\frac{3}{2}}}{3} + ia \left(\frac{\left(2\left(x - \frac{i}{a}\right)a^2 + 2ia\right) \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}}{4a^2} + \frac{\ln\left(\frac{ia + \left(x - \frac{i}{a}\right)a^2}{\sqrt{a^2}} + \sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia \left(x - \frac{i}{a}\right)}\right)}{\sqrt{a^2}} \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^2,x,method=_RETURNVERBOSE)

[Out] $9*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))-2*I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)-I/a*(I/a/(x-I/a)^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)-2*I*a*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))/(a^2)^(1/2))))-3*I*a*(1/3*(a^2*x^2+1)^(3/2)+(a^2*x^2+1)^(1/2)-\operatorname{arctanh}(1/(a^2*x^2+1)^(1/2)))-1/x*(a^2*x^2+1)^(5/2)+4*a^2*(1/4*x*(a^2*x^2+1)^(3/2)+3/8*x*(a^2*x^2+1)^(1/2)+3/8*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2))/(a^2)^(1/2))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^2,x, algorithm="maxima")

[Out] integrate((a^2*x^2 + 1)^(3/2)/((I*a*x + 1)^3*x^2), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 109 vs. 2(53) = 106.

time = 4.65, size = 109, normalized size = 1.70

$$\frac{5a^2x^2 - 5iax + 3(-ia^2x^2 - ax)\log(-ax + \sqrt{a^2x^2 + 1}) + 3(ia^2x^2 + ax)\log(-ax + \sqrt{a^2x^2 + 1} - 1) + \sqrt{a^2x^2 + 1}(5ax - i)}{ax^2 - ix}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^2,x, algorithm="fricas")

[Out] -(5*a^2*x^2 - 5*I*a*x + 3*(-I*a^2*x^2 - a*x)*log(-a*x + sqrt(a^2*x^2 + 1) + 1) + 3*(I*a^2*x^2 + a*x)*log(-a*x + sqrt(a^2*x^2 + 1) - 1) + sqrt(a^2*x^2 + 1)*(5*a*x - I))/(a*x^2 - I*x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2x^2 + 1}}{a^3x^5 - 3ia^2x^4 - 3ax^3 + ix^2} dx + \int \frac{a^2x^2\sqrt{a^2x^2 + 1}}{a^3x^5 - 3ia^2x^4 - 3ax^3 + ix^2} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/x**2,x)

[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**3*x**5 - 3I*a**2*x**4 - 3*a*x**3 + I*x**2), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**3*x**5 - 3I*a**2*x**4 - 3*a*x**3 + I*x**2), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^2,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.42, size = 76, normalized size = 1.19

$$a \operatorname{atanh}\left(\sqrt{a^2x^2 + 1}\right) 3i - \frac{\sqrt{a^2x^2 + 1}}{x} + \frac{4a^2\sqrt{a^2x^2 + 1}}{\left(-x\sqrt{a^2} + \frac{\sqrt{a^2}}{a}1i\right)\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(x^2*(a*x+1i + 1)^3),x)

[Out] a*atanh((a^2*x^2 + 1)^(1/2))*3i - (a^2*x^2 + 1)^(1/2)/x + (4*a^2*(a^2*x^2 + 1)^(1/2))/(((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2))

$$3.58 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{x^3} dx$$

Optimal. Leaf size=93

$$-\frac{\sqrt{1+a^2x^2}}{2x^2} + \frac{3ia\sqrt{1+a^2x^2}}{x} - \frac{4ia^2\sqrt{1+a^2x^2}}{i-ax} + \frac{9}{2}a^2 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $9/2*a^2*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/2*(a^2*x^2+1)^{(1/2)}/x^2+3*I*a*(a^2*x^2+1)^{(1/2)}/x-4*I*a^2*(a^2*x^2+1)^{(1/2)}/(I-a*x)$

Rubi [A]

time = 0.48, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 12, number of rules used = 8, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.571$, Rules used = {5168, 6874, 272, 44, 65, 214, 270, 665}

$$-\frac{4ia^2\sqrt{a^2x^2+1}}{-ax+i} + \frac{3ia\sqrt{a^2x^2+1}}{x} - \frac{\sqrt{a^2x^2+1}}{2x^2} + \frac{9}{2}a^2 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(E^{((3*I)*\operatorname{ArcTan}[a*x])})x^3), x]$

[Out] $-1/2*\operatorname{Sqrt}[1+a^2*x^2]/x^2 + ((3*I)*a*\operatorname{Sqrt}[1+a^2*x^2])/x - ((4*I)*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(I-a*x) + (9*a^2*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]])/2$

Rule 44

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^{(n + 1)})/((b*c - a*d)*(m + 1)), x] - \operatorname{Dist}[d*((m + n + 2)/((b*c - a*d)*(m + 1))), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n, x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x] \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& \operatorname{ILtQ}\{m, -1\} \&\& \operatorname{IntegerQ}\{n\} \&\& \operatorname{LtQ}\{n, 0\}$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{With}\{p = \operatorname{Denominator}\{m\}\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& \operatorname{LtQ}\{-1, m, 0\} \&\& \operatorname{LeQ}\{-1, n, 0\} \&\& \operatorname{LeQ}\{\operatorname{Denominator}\{n\}, \operatorname{Denominator}\{m\}\} \&\& \operatorname{IntLinearQ}\{a, b, c, d, m, n, x\}$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}\{a/b\}$

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*
x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n,
p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[
Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b
, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d,
e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2,
0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)^(n_)])*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1 - iax)^2}{x^3(1 + iax)\sqrt{1 + a^2x^2}} dx \\
&= \int \left(\frac{1}{x^3\sqrt{1 + a^2x^2}} - \frac{3ia}{x^2\sqrt{1 + a^2x^2}} - \frac{4a^2}{x\sqrt{1 + a^2x^2}} + \frac{4a^3}{(-i + ax)\sqrt{1 + a^2x^2}} \right) dx \\
&= -\left((3ia) \int \frac{1}{x^2\sqrt{1 + a^2x^2}} dx \right) - (4a^2) \int \frac{1}{x\sqrt{1 + a^2x^2}} dx + (4a^3) \int \frac{1}{(-i + ax)\sqrt{1 + a^2x^2}} dx \\
&= \frac{3ia\sqrt{1 + a^2x^2}}{x} - \frac{4ia^2\sqrt{1 + a^2x^2}}{i - ax} + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x^2\sqrt{1 + a^2x}} dx, x, x^2 \right) - (2a^2) \text{Subst} \left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{2x^2} + \frac{3ia\sqrt{1 + a^2x^2}}{x} - \frac{4ia^2\sqrt{1 + a^2x^2}}{i - ax} - 4 \text{Subst} \left(\int \frac{1}{-\frac{1}{a^2} + \frac{x^2}{a^2}} dx, x, \sqrt{1 + a^2x^2} \right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{2x^2} + \frac{3ia\sqrt{1 + a^2x^2}}{x} - \frac{4ia^2\sqrt{1 + a^2x^2}}{i - ax} + 4a^2 \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{x^2\sqrt{1 + a^2x}} dx, x, x^2 \right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{2x^2} + \frac{3ia\sqrt{1 + a^2x^2}}{x} - \frac{4ia^2\sqrt{1 + a^2x^2}}{i - ax} + \frac{9}{2} a^2 \tanh^{-1} \left(\sqrt{1 + a^2x^2} \right)
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 79, normalized size = 0.85

$$\sqrt{1 + a^2x^2} \left(-\frac{1}{2x^2} + \frac{3ia}{x} + \frac{4ia^2}{-i + ax} \right) - \frac{9}{2} a^2 \log(x) + \frac{9}{2} a^2 \log \left(1 + \sqrt{1 + a^2x^2} \right)$$

Warning: Unable to verify antiderivative.

`[In] Integrate[1/(E^((3*I)*ArcTan[a*x])*x^3), x]``[Out] Sqrt[1 + a^2*x^2]*(-1/2*1/x^2 + ((3*I)*a)/x + ((4*I)*a^2)/(-I + a*x)) - (9*a^2*Log[x])/2 + (9*a^2*Log[1 + Sqrt[1 + a^2*x^2]])/2`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 550 vs. 2(78) = 156.

time = 0.12, size = 551, normalized size = 5.92

method	result
risch	$\frac{i(6a^3x^3 + ia^2x^2 + 6ax + i)}{2x^2\sqrt{a^2x^2 + 1}} - \frac{a^2 \left(-\frac{8i\sqrt{(x - \frac{i}{a})^2 a^2 + 2ia(x - \frac{i}{a})}}{a(x - \frac{i}{a})} - 9 \operatorname{arctanh} \left(\frac{1}{\sqrt{a^2x^2 + 1}} \right) \right)}{2}$

default	$6a^2 \left(\frac{\left((x - \frac{i}{a})^2 a^2 + 2ia(x - \frac{i}{a}) \right)^{\frac{3}{2}}}{3} + ia \left(\frac{(2(x - \frac{i}{a})a^2 + 2ia) \sqrt{(x - \frac{i}{a})^2 a^2 + 2ia(x - \frac{i}{a})}}{4a^2} + \frac{\ln \left(\frac{ia + (x - \frac{i}{a})a^2 + \sqrt{(x - \frac{i}{a})^2 a^2 + 2ia(x - \frac{i}{a})}}{\sqrt{a^2}} + \sqrt{(x - \frac{i}{a})^2 a^2 + 2ia(x - \frac{i}{a})} \right)}{4a^2} \right) \right)$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^3,x,method=_RETURNVERBOSE)`

[Out] $6a^2 \left(\frac{1}{3} \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{3/2} + Ia \left(\frac{1}{4} \left(2(x - I/a) a^2 + 2Ia(x - I/a) \right) / a^2 \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{1/2} + \frac{1}{2} \ln \left(\frac{Ia + (x - I/a) a^2}{a^2} \right)^{1/2} + \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{1/2} \right) / a^2 \right) - Ia \left(-I/a \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{5/2} + 3Ia \left(\frac{1}{3} \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{3/2} + Ia \left(\frac{1}{4} \left(2(x - I/a) a^2 + 2Ia(x - I/a) \right) / a^2 \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{1/2} + \frac{1}{2} \ln \left(\frac{Ia + (x - I/a) a^2}{a^2} \right)^{1/2} + \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{1/2} \right) / a^2 \right) \right) - \frac{1}{2} x^2 \left(a^2 x^2 + 1 \right)^{5/2} - \frac{9}{2} a^2 \left(\frac{1}{3} \left(a^2 x^2 + 1 \right)^{3/2} + \left(a^2 x^2 + 1 \right)^{1/2} - \operatorname{arctanh} \left(\frac{1}{\left(a^2 x^2 + 1 \right)^{1/2}} \right) \right) - I/a \left((x - I/a)^3 \left((x - I/a)^2 a^2 + 2Ia(x - I/a) \right)^{5/2} - 3Ia \left(-1/x \left(a^2 x^2 + 1 \right)^{5/2} + 4a^2 \left(\frac{1}{4} x \left(a^2 x^2 + 1 \right)^{3/2} + \frac{3}{8} x \left(a^2 x^2 + 1 \right)^{1/2} + \frac{3}{8} \ln \left(a^2 x / a^2 \right)^{1/2} + \left(a^2 x^2 + 1 \right)^{1/2} \right) / a^2 \right) \right)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^3,x, algorithm="maxima")`

[Out] `integrate((a^2*x^2 + 1)^(3/2)/((I*a*x + 1)^3*x^3), x)`

Fricas [A]

time = 5.20, size = 130, normalized size = 1.40

$$\frac{14i a^3 x^3 + 14a^2 x^2 + 9(a^3 x^3 - i a^2 x^2) \log(-ax + \sqrt{a^2 x^2 + 1} + 1) - 9(a^3 x^3 - i a^2 x^2) \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + \sqrt{a^2 x^2 + 1} (14i a^2 x^2 + 5ax + i)}{2(a x^3 - i x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^3,x, algorithm="fricas")`

[Out] $\frac{1}{2} \left(14Ia^3 x^3 + 14a^2 x^2 + 9(a^3 x^3 - Ia^2 x^2) \log(-ax + \sqrt{a^2 x^2 + 1} + 1) - 9(a^3 x^3 - Ia^2 x^2) \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + \sqrt{a^2 x^2 + 1} (14Ia^2 x^2 + 5a^2 x + I) \right) / (a^3 x^3 - Ix^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2 x^2 + 1}}{a^3 x^6 - 3ia^2 x^5 - 3ax^4 + ix^3} dx + \int \frac{a^2 x^2 \sqrt{a^2 x^2 + 1}}{a^3 x^6 - 3ia^2 x^5 - 3ax^4 + ix^3} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/x**3,x)

[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**3*x**6 - 3*I*a**2*x**5 - 3*a*x**4 + I*x**3), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**3*x**6 - 3*I*a**2*x**5 - 3*a*x**4 + I*x**3), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^3,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.43, size = 100, normalized size = 1.08

$$-\frac{a^2 \operatorname{atan}\left(\sqrt{a^2 x^2 + 1}\right) 9i}{2} - \frac{\sqrt{a^2 x^2 + 1}}{2 x^2} + \frac{a \sqrt{a^2 x^2 + 1} 3i}{x} - \frac{a^3 \sqrt{a^2 x^2 + 1} 4i}{\left(-x \sqrt{a^2} + \frac{\sqrt{a^2}}{a}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(x^3*(a*x*1i + 1)^3),x)

[Out] (a*(a^2*x^2 + 1)^(1/2)*3i)/x - (a^2*x^2 + 1)^(1/2)/(2*x^2) - (a^2*atan((a^2*x^2 + 1)^(1/2)*1i)*9i)/2 - (a^3*(a^2*x^2 + 1)^(1/2)*4i)/(((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2))

$$3.59 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{x^4} dx$$

Optimal. Leaf size=118

$$-\frac{\sqrt{1+a^2x^2}}{3x^3} + \frac{3ia\sqrt{1+a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1+a^2x^2}}{3x} - \frac{4a^3\sqrt{1+a^2x^2}}{i-ax} - \frac{11}{2}ia^3 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-11/2*I*a^3*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/3*(a^2*x^2+1)^{(1/2)}/x^3+3/2*I*a*(a^2*x^2+1)^{(1/2)}/x^2+14/3*a^2*(a^2*x^2+1)^{(1/2)}/x-4*a^3*(a^2*x^2+1)^{(1/2)}/(I-a*x)$

Rubi [A]

time = 0.50, antiderivative size = 118, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 9, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {5168, 6874, 277, 270, 272, 44, 65, 214, 665}

$$\frac{14a^2\sqrt{a^2x^2+1}}{3x} + \frac{3ia\sqrt{a^2x^2+1}}{2x^2} - \frac{\sqrt{a^2x^2+1}}{3x^3} - \frac{4a^3\sqrt{a^2x^2+1}}{-ax+i} - \frac{11}{2}ia^3 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right)$$

Antiderivative was successfully verified.

[In] `Int[1/(E^((3*I)*ArcTan[a*x])*x^4),x]`

[Out] $-1/3*\operatorname{Sqrt}[1+a^2*x^2]/x^3 + (((3*I)/2)*a*\operatorname{Sqrt}[1+a^2*x^2])/x^2 + (14*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(3*x) - (4*a^3*\operatorname{Sqrt}[1+a^2*x^2])/(I-a*x) - ((11*I)/2)*a^3*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]]$

Rule 44

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*((m + n + 2)/((b*c - a*d)*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, -1] && !IntegerQ[n] && LtQ[n, 0]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 277

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^n)^(p + 1)/(a*(m + 1))), x] - Dist[b*((m + n*(p + 1) + 1)/(a*(m + 1))), Int[x^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n + p + 1], 0] && NeQ[m, -1]
```

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d, e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2, 0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1 - iax)^2}{x^4(1 + iax)\sqrt{1 + a^2x^2}} dx \\
&= \int \left(\frac{1}{x^4\sqrt{1 + a^2x^2}} - \frac{3ia}{x^3\sqrt{1 + a^2x^2}} - \frac{4a^2}{x^2\sqrt{1 + a^2x^2}} + \frac{4ia^3}{x\sqrt{1 + a^2x^2}} - \frac{4ia^4}{(-i + ax)\sqrt{1 + a^2x^2}} \right) dx \\
&= - \left((3ia) \int \frac{1}{x^3\sqrt{1 + a^2x^2}} dx \right) - (4a^2) \int \frac{1}{x^2\sqrt{1 + a^2x^2}} dx + (4ia^3) \int \frac{1}{x\sqrt{1 + a^2x^2}} dx \\
&= -\frac{\sqrt{1 + a^2x^2}}{3x^3} + \frac{4a^2\sqrt{1 + a^2x^2}}{x} - \frac{4a^3\sqrt{1 + a^2x^2}}{i - ax} - \frac{1}{2}(3ia)\text{Subst}\left(\int \frac{1}{x^2\sqrt{1 + a^2x}} dx, \sqrt{1 + a^2x^2}\right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{3x^3} + \frac{3ia\sqrt{1 + a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1 + a^2x^2}}{3x} - \frac{4a^3\sqrt{1 + a^2x^2}}{i - ax} + (4ia)\text{Subst}\left(\int \frac{1}{x\sqrt{1 + a^2x}} dx, \sqrt{1 + a^2x^2}\right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{3x^3} + \frac{3ia\sqrt{1 + a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1 + a^2x^2}}{3x} - \frac{4a^3\sqrt{1 + a^2x^2}}{i - ax} - 4ia^3 \tanh^{-1}\left(\frac{1}{\sqrt{1 + a^2x^2}}\right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{3x^3} + \frac{3ia\sqrt{1 + a^2x^2}}{2x^2} + \frac{14a^2\sqrt{1 + a^2x^2}}{3x} - \frac{4a^3\sqrt{1 + a^2x^2}}{i - ax} - \frac{11}{2}ia^3 \tanh^{-1}\left(\frac{1}{\sqrt{1 + a^2x^2}}\right)
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 89, normalized size = 0.75

$$\frac{1}{6} \left(\frac{\sqrt{1 + a^2x^2} (2i + 7ax - 19ia^2x^2 + 52a^3x^3)}{x^3(-i + ax)} + 33ia^3 \log(x) - 33ia^3 \log\left(1 + \sqrt{1 + a^2x^2}\right) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x])*x^4), x]**[Out]** ((Sqrt[1 + a^2*x^2]*(2*I + 7*a*x - (19*I)*a^2*x^2 + 52*a^3*x^3))/(x^3*(-I + a*x)) + (33*I)*a^3*Log[x] - (33*I)*a^3*Log[1 + Sqrt[1 + a^2*x^2]])/6**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 828 vs. 2(97) = 194.

time = 0.12, size = 829, normalized size = 7.03

method	result
risch	$ \frac{28a^4x^4 + 9ia^3x^3 + 26a^2x^2 + 9iax - 2}{6x^3\sqrt{a^2x^2 + 1}} + \frac{ia^3 \left(-\frac{8i\sqrt{\left(x - \frac{i}{a}\right)^2 a^2 + 2ia\left(x - \frac{i}{a}\right)}}{a\left(x - \frac{i}{a}\right)} - 11 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2 + 1}}\right) \right)}{2} $

default	$-10ia^3 \left(\frac{\left((x - \frac{i}{a})^2 a^2 + 2ia(x - \frac{i}{a}) \right)^{\frac{3}{2}}}{3} + ia \left(\frac{(2(x - \frac{i}{a})a^2 + 2ia) \sqrt{\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right)}}{4a^2} + \frac{\ln \left(\frac{ia + \left(x - \frac{i}{a} \right) a^2}{\sqrt{a^2}} \right)}{1} \right) \right)$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^4,x,method=_RETURNVERBOSE)`

[Out]
$$-10Ia^3 \left(\frac{1}{3} \left(\frac{(x-I/a)^2 a^2 + 2Ia(x-I/a)}{3} \right)^{\frac{3}{2}} + Ia \left(\frac{(2(x-I/a)a^2 + 2Ia) \sqrt{\left(x - \frac{I}{a} \right)^2 a^2 + 2Ia \left(x - \frac{I}{a} \right)}}{4a^2} + \frac{\ln \left(\frac{Ia + \left(x - \frac{I}{a} \right) a^2}{\sqrt{a^2}} \right)}{1} \right) \right)$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^4,x, algorithm="maxima")`

[Out] `integrate((a^2*x^2 + 1)^(3/2)/((I*a*x + 1)^3*x^4), x)`

Fricas [A]

time = 3.93, size = 139, normalized size = 1.18

$$\frac{52a^4x^4 - 52ia^3x^3 - 33(i a^4x^4 + a^3x^3) \log(-ax + \sqrt{a^2x^2 + 1} + 1) - 33(-i a^4x^4 - a^3x^3) \log(-ax + \sqrt{a^2x^2 + 1} - 1) + (52a^3x^3 - 19ia^2x^2 + 7ax + 2i)\sqrt{a^2x^2 + 1}}{6(ax^4 - ix^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^4,x, algorithm="fricas")`

[Out] $\frac{1}{6}(52a^4x^4 - 52Ia^3x^3 - 33(Ia^4x^4 + a^3x^3)\log(-ax + \sqrt{a^2x^2 + 1}) + 1) - 33(-Ia^4x^4 - a^3x^3)\log(-ax + \sqrt{a^2x^2 + 1}) - 1) + (52a^3x^3 - 19Ia^2x^2 + 7ax + 2I)\sqrt{a^2x^2 + 1})/(a^4x^4 - Ix^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2x^2 + 1}}{a^3x^7 - 3ia^2x^6 - 3ax^5 + ix^4} dx + \int \frac{a^2x^2\sqrt{a^2x^2 + 1}}{a^3x^7 - 3ia^2x^6 - 3ax^5 + ix^4} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/x**4,x)

[Out] $I*(\text{Integral}(\sqrt{a^2x^2 + 1}/(a^3x^7 - 3Ia^2x^6 - 3ax^5 + Ix^4), x) + \text{Integral}(a^2x^2\sqrt{a^2x^2 + 1}/(a^3x^7 - 3Ia^2x^6 - 3ax^5 + Ix^4), x))$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^4,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.07, size = 117, normalized size = 0.99

$$\frac{14a^2\sqrt{a^2x^2+1}}{3x} - \frac{\sqrt{a^2x^2+1}}{3x^3} + \frac{a\sqrt{a^2x^2+1}3i}{2x^2} - \frac{11a^3\operatorname{atan}\left(\frac{\sqrt{a^2x^2+1}}{a}i\right)}{2} - \frac{4a^4\sqrt{a^2x^2+1}}{\left(-x\sqrt{a^2} + \frac{\sqrt{a^2}}{a}i\right)\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(x^4*(a*x*I + 1)^3),x)

[Out] $(a*(a^2x^2 + 1)^{(1/2)}*3i)/(2x^2) - (a^2x^2 + 1)^{(1/2)}/(3x^3) - (11a^3*\operatorname{atan}((a^2x^2 + 1)^{(1/2)}*i))/2 + (14a^2*(a^2x^2 + 1)^{(1/2)})/(3x) - (4a^4*(a^2x^2 + 1)^{(1/2)})/(((a^2)^{(1/2)}*i)/a - x*(a^2)^{(1/2)}*(a^2)^{(1/2)})$

$$3.60 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=139

$$-\frac{\sqrt{1+a^2x^2}}{4x^4} + \frac{ia\sqrt{1+a^2x^2}}{x^3} + \frac{19a^2\sqrt{1+a^2x^2}}{8x^2} - \frac{6ia^3\sqrt{1+a^2x^2}}{x} + \frac{4ia^4\sqrt{1+a^2x^2}}{i-ax} - \frac{51}{8}a^4 \tanh^{-1}\left(\sqrt{1+a^2x^2}\right)$$

[Out] $-51/8*a^4*\operatorname{arctanh}((a^2*x^2+1)^{(1/2)})-1/4*(a^2*x^2+1)^{(1/2)}/x^4+I*a*(a^2*x^2+1)^{(1/2)}/x^3+19/8*a^2*(a^2*x^2+1)^{(1/2)}/x^2-6*I*a^3*(a^2*x^2+1)^{(1/2)}/x+4*I*a^4*(a^2*x^2+1)^{(1/2)}/(I-a*x)$

Rubi [A]

time = 0.53, antiderivative size = 139, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 9, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {5168, 6874, 272, 44, 65, 214, 277, 270, 665}

$$\frac{19a^2\sqrt{a^2x^2+1}}{8x^2} - \frac{\sqrt{a^2x^2+1}}{4x^4} + \frac{ia\sqrt{a^2x^2+1}}{x^3} + \frac{4ia^4\sqrt{a^2x^2+1}}{-ax+i} - \frac{51}{8}a^4 \tanh^{-1}\left(\sqrt{a^2x^2+1}\right) - \frac{6ia^3\sqrt{a^2x^2+1}}{x}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(E^{((3*I)*\operatorname{ArcTan}[a*x])}*x^5),x]$

[Out] $-1/4*\operatorname{Sqrt}[1+a^2*x^2]/x^4+(I*a*\operatorname{Sqrt}[1+a^2*x^2])/x^3+(19*a^2*\operatorname{Sqrt}[1+a^2*x^2])/(8*x^2)-((6*I)*a^3*\operatorname{Sqrt}[1+a^2*x^2])/x+((4*I)*a^4*\operatorname{Sqrt}[1+a^2*x^2])/(I-a*x)-(51*a^4*\operatorname{ArcTanh}[\operatorname{Sqrt}[1+a^2*x^2]])/8$

Rule 44

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^{(n+1)} / ((b*c - a*d)*(m+1))), x] - \operatorname{Dist}[d*((m+n+2) / ((b*c - a*d)*(m+1))), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n, x] /; \operatorname{FreeQ}\{a, b, c, d, n\}, x] \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& \operatorname{ILtQ}[m, -1] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{LtQ}[n, 0]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \operatorname{With}\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 214

$\operatorname{Int}[(a_. + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2]/a)*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x] \&\& \operatorname{NegQ}[a/b]$

Rule 270

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[(c*x)^(m + 1)*((a + b*x^n)^(p + 1)/(a*c*(m + 1))), x] /; FreeQ[{a, b, c, m, n, p}, x] && EqQ[(m + 1)/n + p + 1, 0] && NeQ[m, -1]
```

Rule 272

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[1/n, Subst[Int[x^(Simplify[(m + 1)/n] - 1)*(a + b*x)^p, x], x, x^n], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n]]
```

Rule 277

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[x^(m + 1)*((a + b*x^n)^(p + 1)/(a*(m + 1))), x] - Dist[b*(m + n*(p + 1) + 1)/(a*(m + 1)), Int[x^(m + n)*(a + b*x^n)^p, x], x] /; FreeQ[{a, b, m, n, p}, x] && IntegerQ[Simplify[(m + 1)/n + p + 1], 0] && NeQ[m, -1]
```

Rule 665

```
Int[((d_) + (e_.)*(x_))^(m_.)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[e*(d + e*x)^m*((a + c*x^2)^(p + 1)/(2*c*d*(p + 1))), x] /; FreeQ[{a, c, d, e, m, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && EqQ[m + 2*p + 2, 0]
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{(1 - iax)^2}{x^5(1 + iax)\sqrt{1 + a^2x^2}} dx \\
&= \int \left(\frac{1}{x^5\sqrt{1 + a^2x^2}} - \frac{3ia}{x^4\sqrt{1 + a^2x^2}} - \frac{4a^2}{x^3\sqrt{1 + a^2x^2}} + \frac{4ia^3}{x^2\sqrt{1 + a^2x^2}} + \frac{4a^4}{x\sqrt{1 + a^2x^2}} \right) dx \\
&= -\left((3ia) \int \frac{1}{x^4\sqrt{1 + a^2x^2}} dx \right) - (4a^2) \int \frac{1}{x^3\sqrt{1 + a^2x^2}} dx + (4ia^3) \int \frac{1}{x^2\sqrt{1 + a^2x^2}} dx \\
&= \frac{ia\sqrt{1 + a^2x^2}}{x^3} - \frac{4ia^3\sqrt{1 + a^2x^2}}{x} + \frac{4ia^4\sqrt{1 + a^2x^2}}{i - ax} + \frac{1}{2} \text{Subst} \left(\int \frac{1}{x^3\sqrt{1 + a^2x}} dx, x \right) \\
&= -\frac{\sqrt{1 + a^2x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2x^2}}{x^3} + \frac{2a^2\sqrt{1 + a^2x^2}}{x^2} - \frac{6ia^3\sqrt{1 + a^2x^2}}{x} + \frac{4ia^4\sqrt{1 + a^2x^2}}{i - ax} \\
&= -\frac{\sqrt{1 + a^2x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2x^2}}{x^3} + \frac{19a^2\sqrt{1 + a^2x^2}}{8x^2} - \frac{6ia^3\sqrt{1 + a^2x^2}}{x} + \frac{4ia^4\sqrt{1 + a^2x^2}}{i - ax} \\
&= -\frac{\sqrt{1 + a^2x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2x^2}}{x^3} + \frac{19a^2\sqrt{1 + a^2x^2}}{8x^2} - \frac{6ia^3\sqrt{1 + a^2x^2}}{x} + \frac{4ia^4\sqrt{1 + a^2x^2}}{i - ax} \\
&= -\frac{\sqrt{1 + a^2x^2}}{4x^4} + \frac{ia\sqrt{1 + a^2x^2}}{x^3} + \frac{19a^2\sqrt{1 + a^2x^2}}{8x^2} - \frac{6ia^3\sqrt{1 + a^2x^2}}{x} + \frac{4ia^4\sqrt{1 + a^2x^2}}{i - ax}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 95, normalized size = 0.68

$$\frac{1}{8} \left(\frac{\sqrt{1 + a^2x^2} (2i + 6ax - 11ia^2x^2 - 29a^3x^3 - 80ia^4x^4)}{x^4(-i + ax)} + 51a^4 \log(x) - 51a^4 \log(1 + \sqrt{1 + a^2x^2}) \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x]))*x^5), x]

[Out] ((Sqrt[1 + a^2*x^2]*(2*I + 6*a*x - (11*I)*a^2*x^2 - 29*a^3*x^3 - (80*I)*a^4*x^4))/(x^4*(-I + a*x)) + 51*a^4*Log[x] - 51*a^4*Log[1 + Sqrt[1 + a^2*x^2]])/8

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 936 vs. 2(117) = 234.

time = 0.12, size = 937, normalized size = 6.74

method	result
--------	--------

risch	$\frac{i(48a^5x^5+19ia^4x^4+40a^3x^3+17ia^2x^2-8ax-2i)}{8x^4\sqrt{a^2x^2+1}} + \frac{a^4 \left(-\frac{32i\sqrt{(x-\frac{i}{a})^2a^2+2ia(x-\frac{i}{a})}}{a(x-\frac{i}{a})} - 51 \operatorname{arctanh}\left(\frac{1}{\sqrt{a^2x^2+1}}\right) \right)}{8}$
default	$5ia^3 \left(-\frac{i\left((x-\frac{i}{a})^2a^2+2ia(x-\frac{i}{a})\right)^{\frac{5}{2}}}{a\left(x-\frac{i}{a}\right)^2} + 3ia \left(\frac{\left((x-\frac{i}{a})^2a^2+2ia(x-\frac{i}{a})\right)^{\frac{3}{2}}}{3} + ia \left(\frac{(2(x-\frac{i}{a})a^2+2ia)\sqrt{\left(x-\frac{i}{a}\right)^2a^2+2ia}}{4a^2} \right) \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^5,x,method=_RETURNVERBOSE)`

[Out] $5*I*a^3*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)))/(a^2)^(1/2)))-15*a^4*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)))/(a^2)^(1/2))-3*I*a*(-1/3/x^3*(a^2*x^2+1)^(5/2)+2/3*a^2*(-1/x*(a^2*x^2+1)^(5/2)+4*a^2*(1/4*x*(a^2*x^2+1)^(3/2)+3/8*x*(a^2*x^2+1)^(1/2)+3/8*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2)))-23/4*a^2*(-1/2/x^2*(a^2*x^2+1)^(5/2)+3/2*a^2*(1/3*(a^2*x^2+1)^(3/2)+(a^2*x^2+1)^(1/2)-\operatorname{arctanh}(1/(a^2*x^2+1)^(1/2))))+15*a^4*(1/3*(a^2*x^2+1)^(3/2)+(a^2*x^2+1)^(1/2)-\operatorname{arctanh}(1/(a^2*x^2+1)^(1/2)))-1/4/x^4*(a^2*x^2+1)^(5/2)+a^2*(I/a/(x-I/a)^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)-2*I*a*(-I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(5/2)+3*I*a*(1/3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)+I*a*(1/4*(2*(x-I/a)*a^2+2*I*a)/a^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)+1/2*\ln((I*a+(x-I/a)*a^2)/(a^2)^(1/2)+((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))))) + 10*I*a^3*(-1/x*(a^2*x^2+1)^(5/2)+4*a^2*(1/4*x*(a^2*x^2+1)^(3/2)+3/8*x*(a^2*x^2+1)^(1/2)+3/8*\ln(a^2*x/(a^2)^(1/2)+(a^2*x^2+1)^(1/2)))/(a^2)^(1/2))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^5,x, algorithm="maxima")`

[Out] `integrate((a^2*x^2 + 1)^(3/2)/((I*a*x + 1)^3*x^5), x)`

Fricas [A]

time = 2.24, size = 146, normalized size = 1.05

$$\frac{-80i a^5 x^5 - 80 a^4 x^4 - 51(a^5 x^5 - i a^4 x^4) \log(-ax + \sqrt{a^2 x^2 + 1} + 1) + 51(a^5 x^5 - i a^4 x^4) \log(-ax + \sqrt{a^2 x^2 + 1} - 1) + (-80i a^4 x^4 - 29 a^3 x^3 - 11i a^2 x^2 + 6 a x + 2i) \sqrt{a^2 x^2 + 1}}{8(a x^5 - i x^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^5,x, algorithm="fricas")

[Out] 1/8*(-80*I*a^5*x^5 - 80*a^4*x^4 - 51*(a^5*x^5 - I*a^4*x^4)*log(-a*x + sqrt(a^2*x^2 + 1) + 1) + 51*(a^5*x^5 - I*a^4*x^4)*log(-a*x + sqrt(a^2*x^2 + 1) - 1) + (-80*I*a^4*x^4 - 29*a^3*x^3 - 11*I*a^2*x^2 + 6*a*x + 2*I)*sqrt(a^2*x^2 + 1))/(a*x^5 - I*x^4)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2 x^2 + 1}}{a^3 x^8 - 3i a^2 x^7 - 3a x^6 + i x^5} dx + \int \frac{a^2 x^2 \sqrt{a^2 x^2 + 1}}{a^3 x^8 - 3i a^2 x^7 - 3a x^6 + i x^5} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/x**5,x)

[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**3*x**8 - 3*I*a**2*x**7 - 3*a*x**6 + I*x**5), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**3*x**8 - 3*I*a**2*x**7 - 3*a*x**6 + I*x**5), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/x^5,x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.43, size = 139, normalized size = 1.00

$$\frac{a^4 \operatorname{atan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a}\right) 51i}{8} - \frac{\sqrt{a^2 x^2 + 1}}{4x^4} + \frac{a \sqrt{a^2 x^2 + 1} \operatorname{li}}{x^3} + \frac{19a^2 \sqrt{a^2 x^2 + 1}}{8x^2} - \frac{a^3 \sqrt{a^2 x^2 + 1} 6i}{x} + \frac{a^5 \sqrt{a^2 x^2 + 1} 4i}{\left(-x \sqrt{a^2} + \frac{\sqrt{a^2}}{a} \operatorname{li}\right) \sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(x^5*(a*x*1i + 1)^3),x)

[Out] (a^4*atan((a^2*x^2 + 1)^(1/2)*1i)*51i)/8 - (a^2*x^2 + 1)^(1/2)/(4*x^4) + (a*(a^2*x^2 + 1)^(1/2)*1i)/x^3 + (19*a^2*(a^2*x^2 + 1)^(1/2))/(8*x^2) - (a^3*(a^2*x^2 + 1)^(1/2)*6i)/x + (a^5*(a^2*x^2 + 1)^(1/2)*4i)/(((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2))

3.61 $\int e^{\frac{1}{2}i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=339

$$\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} + \frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3}$$

[Out] $-3/8*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^3-1/12*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^3+1/3*x*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^2+3/16*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3-3/16*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3+2^{(1/2)}-3/32*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3+3/32*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3+2^{(1/2)}$

Rubi [A]

time = 0.17, antiderivative size = 339, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5170, 92, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{3i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} - \frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{3i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} + \frac{3i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((I/2)*\text{ArcTan}[a*x])} * x^2, x]$

[Out] $(((-3*I)/8)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/a^3 - ((I/12)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)})/a^3 + (x*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)})/(3*a^2) + (((3*I)/8)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3) - (((3*I)/8)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3) - (((3*I)/16)*\text{Log}[1 + \text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3) + (((3*I)/16)*\text{Log}[1 + \text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 92

```
Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(
d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)
^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b
*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ
[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 210

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{2}i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2 \sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx \\
&= \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} + \frac{\int \frac{(-1-\frac{iax}{2})\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx}{3a^2} \\
&= -\frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} - \frac{3 \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx}{8a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2} \\
&= -\frac{3i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} - \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{12a^3} + \frac{x(1-iax)^{3/4}(1+iax)^{5/4}}{3a^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 82, normalized size = 0.24

$$\frac{(1-iax)^{3/4} \left(\sqrt[4]{1+iax} (-i+5ax+4ia^2x^2) - 6i\sqrt[4]{2} {}_2F_1\left(-\frac{1}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax)\right) \right)}{12a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])*x^2,x]

[Out] ((1 - I*a*x)^(3/4)*((1 + I*a*x)^(1/4)*(-I + 5*a*x + (4*I)*a^2*x^2) - (6*I)*2^(1/4)*Hypergeometric2F1[-1/4, 3/4, 7/4, (1 - I*a*x)/2]))/(12*a^3)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^2,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^2,x, algorithm="maxima")

[Out] integrate(x^2*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)

Fricas [A]

time = 2.05, size = 244, normalized size = 0.72

$$\frac{12a^3\sqrt{\frac{9i}{64a^6}}\log\left(\frac{9i}{64a^6}\sqrt{\frac{9i}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-12a^3\sqrt{\frac{9i}{64a^6}}\log\left(-\frac{9i}{64a^6}\sqrt{\frac{9i}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)+12a^3\sqrt{\frac{9i}{64a^6}}\log\left(\frac{9i}{64a^6}\sqrt{\frac{9i}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-12a^3\sqrt{\frac{9i}{64a^6}}\log\left(-\frac{9i}{64a^6}\sqrt{\frac{9i}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-(8a^3x^3-2i a^2x^2-ax-11i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{24a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^2,x, algorithm="fricas")

[Out] $-1/24*(12*a^3*\sqrt{9/64*I/a^6}*\log(8/3*I*a^3*\sqrt{9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 12*a^3*\sqrt{9/64*I/a^6}*\log(-8/3*I*a^3*\sqrt{9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) + 12*a^3*\sqrt{-9/64*I/a^6}*\log(8/3*I*a^3*\sqrt{-9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 12*a^3*\sqrt{-9/64*I/a^6}*\log(-8/3*I*a^3*\sqrt{-9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - (8*a^3*x^3 - 2*I*a^2*x^2 - a*x - 11*I)*\sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)})/a^3$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)*x**2,x)

[Out] Integral(x**2*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)

[Out] int(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)

3.62 $\int e^{\frac{1}{2}i \text{ArcTan}(ax)} x dx$

Optimal. Leaf size=295

$$\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4} (1+iax)^{5/4}}{2a^2} - \frac{\text{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2} a^2} + \frac{\text{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2} a^2}$$

[Out] $\frac{1}{4}*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^2 + \frac{1}{2}*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^2 - \frac{1}{8}*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2 + \frac{1}{8}*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2 + \frac{1}{16}*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2 + \frac{1}{16}*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2 - \frac{1}{8\sqrt{2}}*\frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{a^2} + \frac{1}{8\sqrt{2}}*\frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{a^2}$

Rubi [A]

time = 0.13, antiderivative size = 295, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$-\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2} a^2} + \frac{\text{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2} a^2} + \frac{(1-iax)^{3/4} (1+iax)^{5/4}}{2a^2} + \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2} a^2} - \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2} a^2}$$

Antiderivative was successfully verified.

[In] Int[E^((I/2)*ArcTan[a*x])*x, x]

[Out] $((1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(1/4)})/(4*a^2) + ((1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(5/4)})/(2*a^2) - \text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(4*\text{Sqrt}[2]*a^2) + \text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(4*\text{Sqrt}[2]*a^2) + \text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(8*\text{Sqrt}[2]*a^2) - \text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(8*\text{Sqrt}[2]*a^2)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*(b*c - a*d)/(b*(m + n + 1)), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +

```
d*(x^p/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_.), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{2}i \tan^{-1}(ax)} x dx &= \int \frac{x \sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx \\
&= \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{i \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx}{4a} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{i \int \frac{1}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx}{8a} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} + \frac{\text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax}\right)}{2a^2} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} + \frac{\text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2a^2} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4a^2} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} + \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8a^2} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} + \frac{\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2} a^2} \\
&= \frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{\tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2} a^2} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 63, normalized size = 0.21

$$\frac{(1-iax)^{3/4} \left(3(1+iax)^{5/4} + 2\sqrt[4]{2} {}_2F_1\left(-\frac{1}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax)\right) \right)}{6a^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])*x,x]

[Out] ((1 - I*a*x)^(3/4)*(3*(1 + I*a*x)^(5/4) + 2*2^(1/4)*Hypergeometric2F1[-1/4, 3/4, 7/4, (1 - I*a*x)/2]))/(6*a^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x,x)``[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x,x, algorithm="maxima")``[Out] integrate(x*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)`**Fricas [A]**

time = 2.83, size = 236, normalized size = 0.80

$$\frac{2a^2\sqrt{\frac{i}{16a^4}}\log\left(4a^2\sqrt{\frac{i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)-2a^2\sqrt{\frac{i}{16a^4}}\log\left(-4a^2\sqrt{\frac{i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)+2a^2\sqrt{\frac{-i}{16a^4}}\log\left(4a^2\sqrt{\frac{-i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)-2a^2\sqrt{\frac{-i}{16a^4}}\log\left(-4a^2\sqrt{\frac{-i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)-(2a^2x^2-iax+3)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{4a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x,x, algorithm="fricas")`

```
[Out] -1/4*(2*a^2*sqrt(1/16*I/a^4)*log(4*a^2*sqrt(1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(1/16*I/a^4)*log(-4*a^2*sqrt(1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*a^2*sqrt(-1/16*I/a^4)*log(4*a^2*sqrt(-1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(-1/16*I/a^4)*log(-4*a^2*sqrt(-1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - (2*a^2*x^2 - I*a*x + 3)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a^2
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)*x,x)`

[Out] Integral(x*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)

[Out] int(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)

3.63 $\int e^{\frac{1}{2}i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=268

$$\frac{i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{2\sqrt{2}a}$$

[Out] $I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a - 1/2*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)} + 1/2*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)} + 1/4*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)} + (1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)} - 1/4*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)} + (1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 268, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5169, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$-\frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} + \frac{i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a} - \frac{i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I/2)*\text{ArcTan}[a*x]}, x]$

[Out] $(I*(1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(1/4)})/a - (I*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) + (I*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) + ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) - ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Den}$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 338

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],

$x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

Rule 5169

$\text{Int}[E^{\text{ArcTan}[(a_.)*(x_)]*(n_.)}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}, x] /; \text{FreeQ}\{a, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
 \int e^{\frac{1}{2}i \tan^{-1}(ax)} dx &= \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} + \frac{1}{2} \int \frac{1}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} + \frac{(2i) \text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax}\right)}{a} \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} + \frac{(2i) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} - \frac{i \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} + \frac{i \text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)}{a} \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} + \frac{i \text{Subst}\left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2a} + \frac{i \text{Subst}\left(\int \frac{1}{1+\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)}{2a} \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} + \frac{i \log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2\sqrt{2} a} - \frac{i \log\left(1 + \frac{\sqrt{1+iax}}{\sqrt{1-iax}} - \frac{\sqrt{2} \sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)}{2\sqrt{2} a} \\
 &= \frac{i(1-iax)^{3/4} \sqrt[4]{1+iax}}{a} - \frac{i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2} a} + \frac{i \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2} a}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 41, normalized size = 0.15

$$\frac{8ie^{\frac{5}{2}i \text{ArcTan}(ax)} {}_2F_1\left(\frac{5}{4}, 2; \frac{9}{4}; -e^{2i \text{ArcTan}(ax)}\right)}{5a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x]),x]

[Out] (((-8*I)/5)*E^(((5*I)/2)*ArcTan[a*x])*Hypergeometric2F1[5/4, 2, 9/4, -E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)

Fricas [A]

time = 2.15, size = 209, normalized size = 0.78

$$\frac{a\sqrt{\frac{i}{a^2}} \log\left(ia\sqrt{\frac{i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - a\sqrt{\frac{i}{a^2}} \log\left(-ia\sqrt{\frac{i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + a\sqrt{\frac{-i}{a^2}} \log\left(ia\sqrt{\frac{-i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - a\sqrt{\frac{-i}{a^2}} \log\left(-ia\sqrt{\frac{-i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + 2(ax+i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="fricas")

[Out] 1/2*(a*sqrt(I/a^2)*log(I*a*sqrt(I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(I/a^2)*log(-I*a*sqrt(I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + a*sqrt(-I/a^2)*log(I*a*sqrt(-I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(-I/a^2)*log(-I*a*sqrt(-I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*(a*x + I)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2),x)
```

```
[Out] Integral(sqrt((I*a*x + 1)/sqrt(a**2*x**2 + 1)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)
```


3.64 $\int \frac{e^{\frac{1}{2}i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=267

$$-2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - 2 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-2 \operatorname{arctan}\left(\frac{(1+I*a*x)^{1/4}}{(1-I*a*x)^{1/4}}\right) - 2 \operatorname{arctanh}\left(\frac{(1+I*a*x)^{1/4}}{(1-I*a*x)^{1/4}}\right) - \frac{1}{2} \ln\left(\frac{1 - (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} + (1-I*a*x)^{1/2}}\right) - \frac{1}{2} \ln\left(\frac{1 + (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} + (1-I*a*x)^{1/2}}\right) + \operatorname{arctan}\left(\frac{1 - (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} * 2^{1/2}}\right) - \operatorname{arctan}\left(\frac{1 + (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} * 2^{1/2}}\right)$

Rubi [A]

time = 0.13, antiderivative size = 267, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5170, 132, 65, 338, 303, 1176, 631, 210, 1179, 642, 95, 218, 212, 209}

$$-2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} + \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - 2 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(I/2) \operatorname{ArcTan}[a*x]} / x, x\right]$

[Out] $-2 \operatorname{ArcTan}\left[\frac{(1+I*a*x)^{1/4}}{(1-I*a*x)^{1/4}}\right] + \operatorname{Sqrt}[2] \operatorname{ArcTan}\left[1 - \frac{\operatorname{Sqrt}[2] * (1-I*a*x)^{1/4}}{(1+I*a*x)^{1/4}}\right] - \operatorname{Sqrt}[2] \operatorname{ArcTan}\left[1 + \frac{\operatorname{Sqrt}[2] * (1-I*a*x)^{1/4}}{(1+I*a*x)^{1/4}}\right] - 2 \operatorname{ArcTanh}\left[\frac{(1+I*a*x)^{1/4}}{(1-I*a*x)^{1/4}}\right] - \operatorname{Log}\left[\frac{1 + \operatorname{Sqrt}[1-I*a*x]}{\operatorname{Sqrt}[1+I*a*x]}\right] - \frac{\operatorname{Sqrt}[2] * (1-I*a*x)^{1/4}}{(1+I*a*x)^{1/4}} / \operatorname{Sqrt}[2] + \operatorname{Log}\left[\frac{1 + \operatorname{Sqrt}[1-I*a*x]}{\operatorname{Sqrt}[1+I*a*x]}\right] + \frac{\operatorname{Sqrt}[2] * (1-I*a*x)^{1/4}}{(1+I*a*x)^{1/4}} / \operatorname{Sqrt}[2]$

Rule 65

$\operatorname{Int}\left[\left((a_{.}) + (b_{.}) * (x_{.})\right)^{m_{.}} * \left((c_{.}) + (d_{.}) * (x_{.})\right)^{n_{.}}, x_{\text{Symbol}}\right] \rightarrow \operatorname{With}\left[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}\left[p/b, \operatorname{Subst}\left[\operatorname{Int}\left[x^{p*(m+1)-1} * (c - a*(d/b) + d*(x^p/b))^{n_{.}}, x\right], x, (a + b*x)^{1/p}\right], x\right] /; \operatorname{FreeQ}\{a, b, c, d\}, x\} \&\& \operatorname{NeQ}\left[b*c - a*d, 0\right] \&\& \operatorname{LtQ}\{-1, m, 0\} \&\& \operatorname{LeQ}\{-1, n, 0\} \&\& \operatorname{LeQ}\left[\operatorname{Denominator}[n], \operatorname{Denominator}[m]\right] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 95

$\operatorname{Int}\left[\left(\left((a_{.}) + (b_{.}) * (x_{.})\right)^{m_{.}} * \left((c_{.}) + (d_{.}) * (x_{.})\right)^{n_{.}}\right) / \left((e_{.}) + (f_{.}) * (x_{.})\right), x_{\text{Symbol}}\right] \rightarrow \operatorname{With}\left[\{q = \operatorname{Denominator}[m]\}, \operatorname{Dist}\left[q, \operatorname{Subst}\left[\operatorname{Int}\left[x^{q*(m+1)-1} / (b*e - a*f - (d*e - c*f) * x^q), x\right], x, (a + b*x)^{1/q} / (c + d*x)^{1/q}\right]\right]$

```
], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n]
&& LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 132

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^(m - 1), x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{2}i \tan^{-1}(ax)}}{x} dx &= \int \frac{\sqrt[4]{1+iax}}{x\sqrt[4]{1-iax}} dx \\
&= (ia) \int \frac{1}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx + \int \frac{1}{x\sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\left(4\text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax}\right)\right) + 4\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) \\
&= -\left(2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)\right) - 2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 4 \\
&= -2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + 2\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) \\
&= -2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\text{Subst}\left(\int \frac{\sqrt{2}+2x}{-1-\sqrt{2}x-x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}} \\
&= -2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}} \\
&= -2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 97, normalized size = 0.36

$$\frac{2(1-iax)^{3/4} \left(\sqrt[4]{2} (1+iax)^{3/4} {}_2F_1\left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax)\right) + 2 {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}\right) \right)}{3(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])/x,x]

[Out] (-2*(1 - I*a*x)^(3/4)*(2^(1/4)*(1 + I*a*x)^(3/4)*Hypergeometric2F1[3/4, 3/4, 7/4, (1 - I*a*x)/2] + 2*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/((3*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x, algorithm="maxima")`

[Out] `integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))/x, x)`

Fricas [A]

time = 2.68, size = 243, normalized size = 0.91

$$\frac{1}{2}\sqrt{4i}\log\left(\frac{1}{2}\sqrt{4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \frac{1}{2}\sqrt{4i}\log\left(-\frac{1}{2}\sqrt{4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \frac{1}{2}\sqrt{-4i}\log\left(\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \frac{1}{2}\sqrt{-4i}\log\left(-\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - i\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + i\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) + \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x, algorithm="fricas")`

[Out] `1/2*sqrt(4*I)*log(1/2*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/2*sqrt(4*I)*log(-1/2*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(-4*I)*log(1/2*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/2*sqrt(-4*I)*log(-1/2*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x,x)`

[Out] `Integral(sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))/x, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x, x)
```

3.65 $\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^2} dx$

Optimal. Leaf size=92

$$-\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{x} - ia\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-I*a*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-I*a*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.02, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5170, 96, 95, 218, 212, 209}

$$-ia\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{x} - ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((I/2)*\text{ArcTan}[a*x])/x^2}, x]$

[Out] $-(((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x) - I*a*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}] - I*a*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]$

Rule 95

$\text{Int}[(((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)})/((e_.) + (f_.)*(x_.)), x_Symbol] \rightarrow \text{With}\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$

Rule 96

$\text{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*(e + f*x)^{(p+1)}/((m+1)*(b*e - a*f)), x] - \text{Dist}[n*((d*e - c*f)/((m+1)*(b*e - a*f))), \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p\}, x \ \&\& \ \text{EqQ}[m + n + p + 2, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimplerQ}[p, 1]) \ \&\& \ \text{NeQ}[m, -1]$

Rule 209

$\text{Int}[((a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))* \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b\}, x \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{GtQ}[a$

, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{\frac{1}{2}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{\sqrt[4]{1+iax}}{x^2 \sqrt[4]{1-iax}} dx \\
 &= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{x} + \frac{1}{2}(ia) \int \frac{1}{x \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
 &= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{x} + (2ia) \text{Subst} \left(\int \frac{1}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{x} - (ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - (ia) \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{x} - ia \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - ia \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 71, normalized size = 0.77

$$-\frac{i(1-iax)^{3/4} \left(-3i + 3ax + 2ax {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax} \right) \right)}{3x(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])/x^2,x]

[Out] ((-1/3*I)*(1 - I*a*x)^(3/4)*(-3*I + 3*a*x + 2*a*x*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(x*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x, algorithm="maxima")

[Out] integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))/x^2, x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 151 vs. 2(64) = 128.

time = 3.08, size = 151, normalized size = 1.64

$$\frac{-i ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) + i ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2(-i ax + 1)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x, algorithm="fricas")

[Out] 1/2*(-I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(-I*a*x + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**2,x)
```

```
[Out] Integral(sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))/x**2, x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^2,x)
```

```
[Out] int(((a*x*i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^2, x)
```

3.66 $\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=132

$$-\frac{ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} + \frac{1}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{1}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-1/4*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-1/2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/x^2+1/4*a^2*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+1/4*a^2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.03, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5170, 98, 96, 95, 218, 212, 209}

$$\frac{1}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{1}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} - \frac{ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((I/2)*\text{ArcTan}[a*x])}/x^3,x]$

[Out] $((-1/4*I)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x - ((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)})/(2*x^2) + (a^2*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/4 + (a^2*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/4$

Rule 95

$\text{Int}[(((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)})/((e_.) + (f_.)*(x_)), x_Symbol] \rightarrow \text{With}\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$

Rule 96

$\text{Int}[(a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*(e + f*x)^{(p+1)}/((m+1)*(b*e - a*f)), x] - \text{Dist}[n*((d*e - c*f)/((m+1)*(b*e - a*f))), \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p\}, x \ \&\& \ \text{EqQ}[m + n + p + 2, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ (\text{SumSimplerQ}[m, 1] \ || \ !\text{SumSimplerQ}[p, 1]) \ \&\& \ \text{NeQ}[m, -1]$

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{2}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{\sqrt[4]{1+iax}}{x^3 \sqrt[4]{1-iax}} dx \\
&= -\frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} + \frac{1}{4}(ia) \int \frac{\sqrt[4]{1+iax}}{x^2 \sqrt[4]{1-iax}} dx \\
&= -\frac{ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} - \frac{1}{8}a^2 \int \frac{1}{x \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} - \frac{1}{2}a^2 \text{Subst} \left(\int \frac{1}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
&= -\frac{ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} + \frac{1}{4}a^2 \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
&= -\frac{ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{3/4}(1+iax)^{5/4}}{2x^2} + \frac{1}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{1}{4}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 81, normalized size = 0.61

$$\frac{(1-iax)^{3/4} (-6 - 15iax + 9a^2x^2 + 2a^2x^2 {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}))}{12x^2(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])/x^3,x]

[Out] (((1 - I*a*x)^(3/4)*(-6 - (15*I)*a*x + 9*a^2*x^2 + 2*a^2*x^2*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)])))/(12*x^2*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x, algorithm="maxima")
```

```
[Out] integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))/x^3, x)
```

Fricas [A]

time = 3.73, size = 175, normalized size = 1.33

$$\frac{a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) + ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) - ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) - a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) - 2(3a^2x^2 + iax + 2)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{8x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x, algorithm="fricas")
```

```
[Out] 1/8*(a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(3*a^2*x^2 + I*a*x + 2)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^2
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**3,x)
```

```
[Out] Integral(sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))/x**3, x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
```

the root of a polynomial with parameters. This might be wrong. The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^3,x)

[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^3, x)

3.67 $\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=170

$$-\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{12x^2} + \frac{11a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{24x} + \frac{3}{8}ia^3\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-1/3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^3-5/12*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^2+11/24*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x+3/8*I*a^3*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+3/8*I*a^3*\arctanh((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.05, antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 218, 212, 209}

$$\frac{3}{8}ia^3\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{3}{8}ia^3 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{11a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{24x} - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{12x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((I/2)*\text{ArcTan}[a*x])}/x^4, x]$

[Out] $-1/3*((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^3 - (((5*I)/12)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^2 + (11*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(24*x) + ((3*I)/8)*a^3*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}] + ((3*I)/8)*a^3*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]$

Rule 95

$\text{Int}[(((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)})/((e_.) + (f_.)*(x_)), x_Symbol] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$

Rule 101

$\text{Int}[((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x] - \text{Dist}[1/((m+1)*(b*e - a*f)), \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x]$


```
(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m
+ n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1]
] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || Integ
ersQ[p, m + n])
```

Rule 156

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{2}i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{\sqrt[4]{1+iax}}{x^4 \sqrt[4]{1-iax}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{1}{3} \int \frac{\frac{5ia}{2} - 2a^2x}{x^3 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} - \frac{1}{6} \int \frac{\frac{11a^2}{4} + \frac{5}{2}ia^3x}{x^2 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} + \frac{1}{6} \int \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{x \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} - \frac{1}{6} \int \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} - \frac{1}{4} \int \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} + \frac{1}{8} \int \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} - \frac{5ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} + \frac{3}{8} \int \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 93, normalized size = 0.55

$$\frac{(1-iax)^{3/4} (-8 - 18iax + 21a^2x^2 + 11ia^3x^3 + 6ia^3x^3 {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}, \frac{i+ax}{i-ax}))}{24x^3(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])/x^4,x]

[Out] ((1 - I*a*x)^(3/4)*(-8 - (18*I)*a*x + 21*a^2*x^2 + (11*I)*a^3*x^3 + (6*I)*a^3*x^3*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(24*x^3*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x, algorithm="maxima")`

[Out] `integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))/x^4, x)`

Fricas [A]

time = 2.70, size = 184, normalized size = 1.08

$$\frac{9i a^3 x^3 \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{ax + i}} + 1\right) - 9 a^3 x^3 \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{ax + i}} + i\right) + 9 a^3 x^3 \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{ax + i}} - i\right) - 9i a^3 x^3 \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{ax + i}} - 1\right) - 2(11i a^3 x^3 - a^2 x^2 + 2i ax + 8) \sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{ax + i}}}{48 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x, algorithm="fricas")`

[Out] `1/48*(9*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 9*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 9*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 9*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(11*I*a^3*x^3 - a^2*x^2 + 2*I*a*x + 8)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^3`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(ax - i)}{\sqrt{a^2 x^2 + 1}}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**4,x)`

[Out] `Integral(sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))/x**4, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.01
```

$$\int \frac{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^4,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^4, x)
```

$$3.68 \quad \int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=202

$$\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{96x^2} + \frac{83ia^3(1-iax)^{3/4}\sqrt[4]{1+iax}}{192x}$$

[Out] $-1/4*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^4-7/24*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^3+29/96*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^2+83/192*I*a^3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-11/64*a^4*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-11/64*a^4*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.06, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 218, 212, 209}

$$-\frac{11}{64}a^4\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{11}{64}a^4\operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{83ia^3(1-iax)^{3/4}\sqrt[4]{1+iax}}{192x} + \frac{29a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{96x^2} - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{24x^3}$$

Antiderivative was successfully verified.

[In] Int[E^((I/2)*ArcTan[a*x])/x^5,x]

[Out] $-1/4*((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^4 - (((7*I)/24)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^3 + (29*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(96*x^2) + (((83*I)/192)*a^3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x - (11*a^4*\operatorname{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64 - (11*a^4*\operatorname{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 101

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m+1)*(c + d*x)^n*(e + f*x)^(p+1)

)/((m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 156

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{2}i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{\sqrt[4]{1+iax}}{x^5 \sqrt[4]{1-iax}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} + \frac{1}{4} \int \frac{\frac{7ia}{2} - 3a^2x}{x^4 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} - \frac{1}{12} \int \frac{\frac{29a^2}{4} + 7ia^3x}{x^3 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x^4} - \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x^3} + \frac{29a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{96x^2} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 99, normalized size = 0.49

$$\frac{(1-iax)^{3/4} (48 + 104iax - 114a^2x^2 - 141ia^3x^3 + 83a^4x^4 + 22a^4x^4 {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}))}{192x^4(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])/x^5,x]

[Out] -1/192*((1 - I*a*x)^(3/4)*(48 + (104*I)*a*x - 114*a^2*x^2 - (141*I)*a^3*x^3 + 83*a^4*x^4 + 22*a^4*x^4*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x]]))/(x^4*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x, algorithm="maxima")`

[Out] `integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))/x^5, x)`

Fricas [A]

time = 2.74, size = 192, normalized size = 0.95

$$\frac{33a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + 33i a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - 33i a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 33a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2(83a^4x^4 + 25i a^3x^3 + 2a^2x^2 - 8i ax - 48)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{384x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x, algorithm="fricas")`

[Out] `-1/384*(33*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 33*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 33*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 33*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(83*a^4*x^4 + 25*I*a^3*x^3 + 2*a^2*x^2 - 8*I*a*x - 48)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^4`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**5,x)`

[Out] `Integral(sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))/x**5, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^5,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^5, x)
```

3.69 $\int \frac{e^{\frac{1}{2}i\text{ArcTan}(ax)}}{x^6} dx$

Optimal. Leaf size=240

$$\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{48x^3} + \frac{269ia^3(1-iax)^{3/4}\sqrt[4]{1+iax}}{960x^2}$$

[Out] $-1/5*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^5-9/40*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^4+11/48*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^3+269/960*I*a^3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^2-611/1920*a^4*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-31/128*I*a^5*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-31/128*I*a^5*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.07, antiderivative size = 240, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 218, 212, 209}

$$-\frac{31}{128}ia^5\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{31}{128}ia^5\operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{611a^4(1-iax)^{3/4}\sqrt[4]{1+iax}}{1920x} + \frac{269ia^3(1-iax)^{3/4}\sqrt[4]{1+iax}}{960x^2} + \frac{11a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{48x^3} - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{40x^4}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((I/2)*\operatorname{ArcTan}[a*x])}/x^6, x]$

[Out] $-1/5*((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^5 - (((9*I)/40)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^4 + (11*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(48*x^3) + (((269*I)/960)*a^3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^2 - (611*a^4*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(1920*x) - ((31*I)/128)*a^5*\operatorname{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}] - ((31*I)/128)*a^5*\operatorname{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[u, (b_*)(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 95

$\operatorname{Int}[(((a_.) + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)})/((e_.) + (f_.)*(x_)), x_Symbol] \rightarrow \operatorname{With}[\{q = \operatorname{Denominator}[m]\}, \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 101

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_], x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1
))/(m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(
m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m
+ n + p + 2)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1
] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || Integ
ersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/(m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 209

```

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

Rule 218

```

Int[((a_.) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{2}i \tan^{-1}(ax)}}{x^6} dx &= \int \frac{\sqrt[4]{1+iax}}{x^6 \sqrt[4]{1-iax}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} + \frac{1}{5} \int \frac{\frac{9ia}{2} - 4a^2x}{x^5 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} - \frac{1}{20} \int \frac{\frac{55a^2}{4} + \frac{27}{2}ia^3x}{x^4 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{5x^5} - \frac{9ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{40x^4} + \frac{11a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{48x^3} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 111, normalized size = 0.46

$$\frac{(1-iax)^{3/4} (-384 - 816iax + 872a^2x^2 + 978ia^3x^3 - 1149a^4x^4 - 611ia^5x^5 - 310ia^5x^5 {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}))}{1920x^5(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a*x])/x^6,x]

[Out] ((1 - I*a*x)^(3/4)*(-384 - (816*I)*a*x + 872*a^2*x^2 + (978*I)*a^3*x^3 - 1149*a^4*x^4 - (611*I)*a^5*x^5 - (310*I)*a^5*x^5*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(1920*x^5*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}}{x^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^6,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^6,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^6,x, algorithm="maxima")

[Out] integrate(sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))/x^6, x)

Fricas [A]

time = 1.39, size = 200, normalized size = 0.83

$$\frac{-465i a^5 x^5 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + 465 a^5 x^5 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - 465 a^5 x^5 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) + 465i a^5 x^5 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2(-611i a^5 x^5 + 73 a^4 x^4 - 98i a^3 x^3 - 8 a^2 x^2 + 48i a x + 384) \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{3840 x^6}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^6,x, algorithm="fricas")

[Out] 1/3840*(-465*I*a^5*x^5*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 465*a^5*x^5*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 465*a^5*x^5*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + 465*I*a^5*x^5*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(-611*I*a^5*x^5 + 73*a^4*x^4 - 98*I*a^3*x^3 - 8*a^2*x^2 + 48*I*a*x + 384)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^5

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}}}{x^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**6,x)

[Out] Integral(sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))/x**6, x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^6,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}}{x^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*I + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^6,x)

[Out] int(((a*x*I + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)/x^6, x)

3.70 $\int e^{\frac{3}{2}i \text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=337

$$-\frac{41\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2\sqrt[4]{1-iax}(1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}(11+4iax)}{32a^4} + \frac{123\text{ArcTan}\left(\frac{1-\sqrt[4]{1-iax}}{1+\sqrt[4]{1-iax}}\right)}{64a^4}$$

```
[Out] -41/64*(1-I*a*x)^(1/4)*(1+I*a*x)^(3/4)/a^4+1/4*x^2*(1-I*a*x)^(1/4)*(1+I*a*x)^(7/4)/a^2-1/32*(1-I*a*x)^(1/4)*(1+I*a*x)^(7/4)*(11+4*I*a*x)/a^4+123/128*arctan(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))/a^4*2^(1/2)-123/128*arctan(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))/a^4*2^(1/2)+123/256*ln(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))/a^4*2^(1/2)-123/256*ln(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))/a^4*2^(1/2)
```

Rubi [A]

time = 0.16, antiderivative size = 337, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5170, 102, 152, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{123\text{ArcTan}\left(\frac{1-\sqrt[4]{1-iax}}{1+\sqrt[4]{1-iax}}\right)}{64\sqrt[4]{a^4}} - \frac{123\text{ArcTan}\left(\frac{1+\sqrt[4]{1-iax}}{1+\sqrt[4]{1+iax}}\right)}{64\sqrt[4]{a^4}} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}(11+4iax)}{32a^4} - \frac{41\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{123\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{128\sqrt[4]{a^4}} - \frac{123\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{128\sqrt[4]{a^4}} + \frac{x^2\sqrt[4]{1-iax}(1+iax)^{7/4}}{4a^2}$$

Antiderivative was successfully verified.

[In] Int[E^(((3*I)/2)*ArcTan[a*x])*x^3,x]

```
[Out] (-41*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(3/4))/(64*a^4) + (x^2*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(7/4))/(4*a^2) - ((1 - I*a*x)^(1/4)*(1 + I*a*x)^(7/4)*(11 + (4*I)*a*x))/(32*a^4) + (123*ArcTan[1 - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/(64*Sqrt[2]*a^4) - (123*ArcTan[1 + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/(64*Sqrt[2]*a^4) + (123*Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/(128*Sqrt[2]*a^4) - (123*Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/(128*Sqrt[2]*a^4)
```

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 102

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_], x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a
+ b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b
*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*
(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p
}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))*(g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n},
x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
```


b}], x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int e^{\frac{3}{2}i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1+iax)^{3/4}}{(1-iax)^{3/4}} dx \\
&= \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} + \frac{\int \frac{x(1+iax)^{3/4}(-2-\frac{3iax}{2})}{(1-iax)^{3/4}} dx}{4a^2} \\
&= \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} + \frac{(41i) \int \frac{(1+iax)^{3/4}}{(1-iax)^{3/4}} dx}{64a^3} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4} \\
&= -\frac{41 \sqrt[4]{1-iax} (1+iax)^{3/4}}{64a^4} + \frac{x^2 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^2} - \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}(11+4iax)}{32a^4}
\end{aligned}$$

Mathematica [A]

time = 0.24, size = 228, normalized size = 0.68

$$\frac{512e^{\frac{3}{2}\text{ArcTan}(ax)}}{(1+e^{2\text{ArcTan}(ax)})^4} - \frac{1152e^{\frac{3}{2}\text{ArcTan}(ax)}}{(1+e^{2\text{ArcTan}(ax)})^3} + \frac{1008e^{\frac{3}{2}\text{ArcTan}(ax)}}{(1+e^{2\text{ArcTan}(ax)})^2} - \frac{532e^{\frac{3}{2}\text{ArcTan}(ax)}}{(1+e^{2\text{ArcTan}(ax)})} - 123(-1)^{3/4} \log(\sqrt{-1} - e^{\frac{1}{2}i\text{ArcTan}(ax)}) - 123\sqrt{-1} \log((-1)^{3/4} - e^{\frac{1}{2}i\text{ArcTan}(ax)}) + 123(-1)^{3/4} \log(\sqrt{-1} + e^{\frac{1}{2}i\text{ArcTan}(ax)}) + 123\sqrt{-1} \log((-1)^{3/4} + e^{\frac{1}{2}i\text{ArcTan}(ax)})$$

128a⁴

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])*x^3,x]

[Out] ((512*E^(((3*I)/2)*ArcTan[a*x]))/(1 + E^((2*I)*ArcTan[a*x]))^4 - (1152*E^(((3*I)/2)*ArcTan[a*x]))/(1 + E^((2*I)*ArcTan[a*x]))^3 + (1008*E^(((3*I)/2)*ArcTan[a*x]))/(1 + E^((2*I)*ArcTan[a*x]))^2 - (532*E^(((3*I)/2)*ArcTan[a*x]))

)/(1 + E^((2*I)*ArcTan[a*x])) - 123*(-1)^(3/4)*Log[(-1)^(1/4) - E^((I/2)*ArcTan[a*x])] - 123*(-1)^(1/4)*Log[(-1)^(3/4) - E^((I/2)*ArcTan[a*x])] + 123*(-1)^(3/4)*Log[(-1)^(1/4) + E^((I/2)*ArcTan[a*x])] + 123*(-1)^(1/4)*Log[(-1)^(3/4) + E^((I/2)*ArcTan[a*x])]/(128*a^4)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} x^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^3,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^3,x, algorithm="maxima")

[Out] integrate(x^3*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [A]

time = 2.59, size = 254, normalized size = 0.75

$$\frac{32a^4\sqrt{\frac{15129}{4096a^2}}\log\left(\frac{64i a^4\sqrt{\frac{15129}{4096a^2}} + \sqrt{i\sqrt{a^2x^2+1}}}{az+i}\right) - 32a^4\sqrt{\frac{15129}{4096a^2}}\log\left(\frac{64i a^4\sqrt{\frac{15129}{4096a^2}} + \sqrt{i\sqrt{a^2x^2+1}}}{az+i}\right) - 32a^4\sqrt{\frac{15129}{4096a^2}}\log\left(\frac{64i a^4\sqrt{\frac{15129}{4096a^2}} + \sqrt{i\sqrt{a^2x^2+1}}}{az+i}\right) + 32a^4\sqrt{\frac{15129}{4096a^2}}\log\left(\frac{64i a^4\sqrt{\frac{15129}{4096a^2}} + \sqrt{i\sqrt{a^2x^2+1}}}{az+i}\right) + (16i a^3x^3 + 24a^2x^2 - 30i a^2x - 63)\sqrt{a^2x^2+1}\sqrt{\frac{i\sqrt{a^2x^2+1}}{az+i}}}{64a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^3,x, algorithm="fricas")

[Out] 1/64*(32*a^4*sqrt(15129/4096*I/a^8)*log(64/123*I*a^4*sqrt(15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 32*a^4*sqrt(15129/4096*I/a^8)*log(-64/123*I*a^4*sqrt(15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 32*a^4*sqrt(-15129/4096*I/a^8)*log(64/123*I*a^4*sqrt(-15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 32*a^4*sqrt(-15129/4096*I/a^8)*log(-64/123*I*a^4*sqrt(-15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + (16*I*a^3*x^3 + 24*a^2*x^2 - 30*I*a*x - 63)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/a^4

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \left(\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)*x**3,x)
```

```
[Out] Integral(x**3*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^3,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^3 \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)
```

3.71 $\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=339

$$-\frac{17i\sqrt[4]{1-iax}(1+iax)^{3/4}}{24a^3} - \frac{i\sqrt[4]{1-iax}(1+iax)^{7/4}}{4a^3} + \frac{x\sqrt[4]{1-iax}(1+iax)^{7/4}}{3a^2} + \frac{17i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3}$$

```
[Out] -17/24*I*(1-I*a*x)^(1/4)*(1+I*a*x)^(3/4)/a^3-1/4*I*(1-I*a*x)^(1/4)*(1+I*a*x)^(7/4)/a^3+1/3*x*(1-I*a*x)^(1/4)*(1+I*a*x)^(7/4)/a^2+17/16*I*arctan(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))/a^3*2^(1/2)-17/16*I*arctan(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))/a^3*2^(1/2)+17/32*I*ln(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))/a^3*2^(1/2)-17/32*I*ln(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))/a^3*2^(1/2)
```

Rubi [A]

time = 0.15, antiderivative size = 339, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5170, 92, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{17i\text{ArcTan}\left(\frac{1 - \sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{17i\text{ArcTan}\left(\frac{1 + \sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{i\sqrt[4]{1-iax}(1+iax)^{7/4}}{4a^3} - \frac{17i\sqrt[4]{1-iax}(1+iax)^{3/4}}{24a^3} + \frac{17i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} - \frac{17i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} + \frac{x\sqrt[4]{1-iax}(1+iax)^{7/4}}{3a^2}$$

Antiderivative was successfully verified.

[In] Int[E^(((3*I)/2)*ArcTan[a*x])*x^2,x]

```
[Out] (((-17*I)/24)*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(3/4))/a^3 - ((I/4)*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(7/4))/a^3 + (x*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(7/4))/(3*a^2) + (((17*I)/8)*ArcTan[1 - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/((Sqrt[2]*a^3) - (((17*I)/8)*ArcTan[1 + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/((Sqrt[2]*a^3) + (((17*I)/16)*Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/(Sqrt[2]*a^3) - (((17*I)/16)*Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)])/(Sqrt[2]*a^3))
```

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 92

```
Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(
d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)
^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b
*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ
[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{3}{2}i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1+iax)^{3/4}}{(1-iax)^{3/4}} dx \\
&= \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} + \frac{\int \frac{(1+iax)^{3/4}(-1-\frac{3iax}{2})}{(1-iax)^{3/4}} dx}{3a^2} \\
&= -\frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17 \int \frac{(1+iax)^{3/4}}{(1-iax)^{3/4}} dx}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2} \\
&= -\frac{17i^4 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24a^3} - \frac{i^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{4a^3} + \frac{x^4 \sqrt[4]{1-iax} (1+iax)^{7/4}}{3a^2} - \frac{17}{24a^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 82, normalized size = 0.24

$$\frac{\sqrt[4]{1-iax} ((1+iax)^{3/4} (-3i + 7ax + 4ia^2x^2) - 34i2^{3/4} {}_2F_1(-\frac{3}{4}, \frac{1}{4}; \frac{5}{4}; \frac{1}{2}(1-iax)))}{12a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])*x^2,x]

[Out] ((1 - I*a*x)^(1/4)*((1 + I*a*x)^(3/4)*(-3*I + 7*a*x + (4*I)*a^2*x^2) - (34*I)*2^(3/4)*Hypergeometric2F1[-3/4, 1/4, 5/4, (1 - I*a*x)/2]))/(12*a^3)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^2,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^2,x, algorithm="maxima")

[Out] integrate(x^2*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [A]

time = 2.09, size = 247, normalized size = 0.73

$$\frac{12a^3 \sqrt{\frac{289i}{64a^6}} \log\left(\frac{8}{17}a^3 \sqrt{\frac{289i}{64a^6}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 12a^3 \sqrt{\frac{289i}{64a^6}} \log\left(-\frac{8}{17}a^3 \sqrt{\frac{289i}{64a^6}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 12a^3 \sqrt{\frac{289i}{64a^6}} \log\left(\frac{8}{17}a^3 \sqrt{\frac{289i}{64a^6}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + 12a^3 \sqrt{\frac{289i}{64a^6}} \log\left(-\frac{8}{17}a^3 \sqrt{\frac{289i}{64a^6}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \sqrt{a^2x^2+1} (8i a^2 x^2 + 14ax - 23i) \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{24a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^2,x, algorithm="fricas")

[Out] $-1/24*(12*a^3*\sqrt{289/64*I/a^6}*\log(8/17*a^3*\sqrt{289/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 12*a^3*\sqrt{289/64*I/a^6}*\log(-8/17*a^3*\sqrt{289/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 12*a^3*\sqrt{-289/64*I/a^6}*\log(8/17*a^3*\sqrt{-289/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) + 12*a^3*\sqrt{-289/64*I/a^6}*\log(-8/17*a^3*\sqrt{-289/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - \sqrt{a^2*x^2 + 1}*(8*I*a^2*x^2 + 14*a*x - 23*I)*\sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)})/a^3$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \left(\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)*x**2,x)

[Out] Integral($x^{**2}*(I*(a*x - I)/\sqrt{a^{**2}*x^{**2} + 1})^{**}(3/2)$, x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)

[Out] int(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)

3.72 $\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=295

$$\frac{3\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2a^2} - \frac{9\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \frac{9\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)}{4\sqrt{2}a^2}$$

[Out] $3/4*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^2+1/2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(7/4)}/a^2-9/8*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2*2^{(1/2)}+9/8*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2*2^{(1/2)}-9/16*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^2*2^{(1/2)}+9/16*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^2*2^{(1/2)}$

Rubi [A]

time = 0.13, antiderivative size = 295, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$-\frac{9\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \frac{9\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2a^2} + \frac{3\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{9\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2} + \frac{9\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)/2)*\text{ArcTan}[a*x]}*x, x]$

[Out] $(3*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/(4*a^2) + ((1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(7/4)})/(2*a^2) - (9*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4})])/(4*\text{Sqrt}[2]*a^2) + (9*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4})])/(4*\text{Sqrt}[2]*a^2) - (9*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4})])/(8*\text{Sqrt}[2]*a^2) + (9*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4})])/(8*\text{Sqrt}[2]*a^2)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) +$

```
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{3}{2}i \tan^{-1}(ax)} x dx &= \int \frac{x(1+iax)^{3/4}}{(1-iax)^{3/4}} dx \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} - \frac{(3i) \int \frac{(1+iax)^{3/4}}{(1-iax)^{3/4}} dx}{4a} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} - \frac{(9i) \int \frac{1}{(1-iax)^{3/4} \sqrt[4]{1+iax}} dx}{8a} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} + \frac{9 \text{Subst} \left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-iax} \right)}{2a^2} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} + \frac{9 \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{2a^2} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} + \frac{9 \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{4a^2} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} + \frac{9 \text{Subst} \left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{8a^2} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} - \frac{9 \log \left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{8\sqrt{2} a^2} \\
&= \frac{3\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{\sqrt[4]{1-iax} (1+iax)^{7/4}}{2a^2} - \frac{9 \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{4\sqrt{2} a^2} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 61, normalized size = 0.21

$$\frac{\sqrt[4]{1-iax} \left((1+iax)^{7/4} + 6 \cdot 2^{3/4} {}_2F_1 \left(-\frac{3}{4}, \frac{1}{4}; \frac{5}{4}; \frac{1}{2}(1-iax) \right) \right)}{2a^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])*x,x]

[Out] ((1 - I*a*x)^(1/4)*((1 + I*a*x)^(7/4) + 6*2^(3/4)*Hypergeometric2F1[-3/4, 1/4, 5/4, (1 - I*a*x)/2]))/(2*a^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x,x, algorithm="maxima")

[Out] integrate(x*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [A]

time = 3.30, size = 239, normalized size = 0.81

$$\frac{2a^2\sqrt{\frac{81i}{16a^4}} \log\left(\frac{\frac{1}{2}i a^2\sqrt{\frac{81i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{\frac{1}{2}i a^2\sqrt{\frac{81i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) - 2a^2\sqrt{\frac{81i}{16a^4}} \log\left(-\frac{1}{2}i a^2\sqrt{\frac{81i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 2a^2\sqrt{-\frac{81i}{16a^4}} \log\left(\frac{\frac{1}{2}i a^2\sqrt{\frac{81i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{\frac{1}{2}i a^2\sqrt{\frac{81i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) + 2a^2\sqrt{\frac{81i}{16a^4}} \log\left(-\frac{1}{2}i a^2\sqrt{\frac{81i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \sqrt{a^2x^2+1} (2i ax + 5) \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{4a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x,x, algorithm="fricas")

[Out] -1/4*(2*a^2*sqrt(81/16*I/a^4)*log(4/9*I*a^2*sqrt(81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(81/16*I/a^4)*log(-4/9*I*a^2*sqrt(81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(-81/16*I/a^4)*log(4/9*I*a^2*sqrt(-81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*a^2*sqrt(-81/16*I/a^4)*log(-4/9*I*a^2*sqrt(-81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - sqrt(a^2*x^2 + 1)*(2*I*a*x + 5)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \left(\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)*x,x)

[Out] Integral(x*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*((a*x*I + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)

[Out] int(x*((a*x*I + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)

3.73 $\int e^{\frac{3}{2}i \operatorname{ArcTan}(ax)} dx$

Optimal. Leaf size=268

$$\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} - \frac{3i \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{3i \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - 3i \log\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)$$

[Out] $I*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a-3/2*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}+3/2*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}-3/4*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}+3/4*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 268, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5169, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$-\frac{3i \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{3i \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} - \frac{3i \log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a} + \frac{3i \log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)/2)*\operatorname{ArcTan}[a*x]}, x]$

[Out] $(I*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/a - ((3*I)*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*a) + ((3*I)*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*a) - (((3*I)/2)*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a*x]/\operatorname{Sqrt}[1 + I*a*x] - (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*a) + (((3*I)/2)*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a*x]/\operatorname{Sqrt}[1 + I*a*x] + (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*a)$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] := \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \operatorname{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \operatorname{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{GtQ}[n, 0] \&\& \operatorname{NeQ}[m + n + 1, 0] \&\& !(\operatorname{IGtQ}[m, 0] \&\& (!\operatorname{IntegerQ}[n] || (\operatorname{GtQ}[m, 0] \&\& \operatorname{LtQ}[m - n, 0]))) \&\& !\operatorname{ILtQ}[m + n + 2, 0] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] := \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^{p/b})^n), x], x, (a + b*x)^{(1/p)}], x]] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{LeQ}[-1, n, 0] \&\& \operatorname{LeQ}[\operatorname{Denominator}[n], \operatorname{Denominator}[m]]$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],

$x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$

Rule 5169

$\text{Int}[E^{\text{ArcTan}[(a_.)*(x_.)]*(n_.)}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*x)^{I*(n/2)} / (1 + I*a*x)^{I*(n/2)}, x] /; \text{FreeQ}[\{a, n\}, x] \ \&\& \ !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
 \int e^{\frac{3}{2}i \tan^{-1}(ax)} dx &= \int \frac{(1 + iax)^{3/4}}{(1 - iax)^{3/4}} dx \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} + \frac{3}{2} \int \frac{1}{(1 - iax)^{3/4} \sqrt[4]{1 + iax}} dx \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} + \frac{(6i) \text{Subst}\left(\int \frac{1}{\sqrt[4]{2 - x^4}} dx, x, \sqrt[4]{1 - iax}\right)}{a} \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} + \frac{(6i) \text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{a} \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} + \frac{(3i) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{a} + \frac{(3i) \text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{a} \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} + \frac{(3i) \text{Subst}\left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{2a} + \frac{(3i) \text{Subst}\left(\int \frac{1}{1+\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{2a} \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} - \frac{3i \log\left(1 + \frac{\sqrt{1 - iax}}{\sqrt{1 + iax}} - \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{2\sqrt{2} a} + \frac{3i \log\left(1 + \frac{\sqrt{1 + iax}}{\sqrt{1 - iax}} - \frac{\sqrt{2} \sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}}\right)}{2\sqrt{2} a} \\
 &= \frac{i\sqrt[4]{1 - iax} (1 + iax)^{3/4}}{a} - \frac{3i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{\sqrt{2} a} + \frac{3i \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{\sqrt{2} a}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 41, normalized size = 0.15

$$\frac{8ie^{\frac{7}{2}i \text{ArcTan}(ax)} {}_2F_1\left(\frac{7}{4}, 2; \frac{11}{4}; -e^{2i \text{ArcTan}(ax)}\right)}{7a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x]),x]

[Out] (((-8*I)/7)*E^(((7*I)/2)*ArcTan[a*x])*Hypergeometric2F1[7/4, 2, 11/4, -E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [A]

time = 2.30, size = 215, normalized size = 0.80

$$\frac{a\sqrt{\frac{9i}{a^2}} \log\left(\frac{1}{3}a\sqrt{\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - a\sqrt{\frac{9i}{a^2}} \log\left(-\frac{1}{3}a\sqrt{\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - a\sqrt{\frac{9i}{a^2}} \log\left(\frac{1}{3}a\sqrt{-\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + a\sqrt{\frac{9i}{a^2}} \log\left(-\frac{1}{3}a\sqrt{-\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + 2i\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="fricas")

[Out] 1/2*(a*sqrt(9*I/a^2)*log(1/3*a*sqrt(9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(9*I/a^2)*log(-1/3*a*sqrt(9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(-9*I/a^2)*log(1/3*a*sqrt(-9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + a*sqrt(-9*I/a^2)*log(-1/3*a*sqrt(-9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*I*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2),x)
```

```
[Out] Integral(((I*a*x + 1)/sqrt(a**2*x**2 + 1))**(3/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)
```

$$3.74 \quad \int \frac{e^{\frac{3}{2}i \operatorname{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=267

$$2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - 2 \tanh^{-1}$$

[Out] 2*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-2*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))+1/2*ln(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))*2^(1/2)-1/2*ln(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))*2^(1/2)+arctan(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))*2^(1/2)-arctan(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))*2^(1/2)

Rubi [A]

time = 0.12, antiderivative size = 267, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5170, 132, 65, 246, 217, 1179, 642, 1176, 631, 210, 95, 304, 209, 212}

$$2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right) - \log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] Int[E^(((3*I)/2)*ArcTan[a*x])/x,x]

[Out] 2*ArcTan[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)] + Sqrt[2]*ArcTan[1 - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)] - Sqrt[2]*ArcTan[1 + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)] - 2*ArcTanh[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)] + Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)]/Sqrt[2] - Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)]/Sqrt[2]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)]

], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 132

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^(m, x), x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 304

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{3}{2}i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1+iax)^{3/4}}{x(1-iax)^{3/4}} dx \\
&= (ia) \int \frac{1}{(1-iax)^{3/4} \sqrt[4]{1+iax}} dx + \int \frac{1}{x(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= -\left(4\text{Subst}\left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-iax}\right)\right) + 4\text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) \\
&= -\left(2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)\right) + 2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{\text{Subst}\left(\int \frac{\sqrt{2}+2x}{-1-\sqrt{2}x-x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}} \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}} \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 96, normalized size = 0.36

$$-22^{3/4} \sqrt[4]{1-iax} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \frac{1}{2}(1-iax)\right) - \frac{4\sqrt[4]{1-iax} {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; -\frac{1-iax}{-1-iax}\right)}{\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])/x,x]

[Out] -2*2^(3/4)*(1 - I*a*x)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, (1 - I*a*x)/2] - (4*(1 - I*a*x)^(1/4)*Hypergeometric2F1[1/4, 1, 5/4, -((1 - I*a*x)/(-1 - I*a*x))])/(1 + I*a*x)^(1/4)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x, algorithm="maxima")`

[Out] `integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)/x, x)`

Fricas [A]

time = 3.52, size = 243, normalized size = 0.91

$$\frac{1}{2}\sqrt{4i}\log\left(\frac{1}{2}\sqrt{4i} + \sqrt{\frac{\sqrt{4i^2+1}}{ax+i}}\right) - \frac{1}{2}\sqrt{4i}\log\left(-\frac{1}{2}\sqrt{4i} + \sqrt{\frac{\sqrt{4i^2+1}}{ax+i}}\right) - \frac{1}{2}\sqrt{-4i}\log\left(\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{\sqrt{4i^2+1}}{ax+i}}\right) + \frac{1}{2}\sqrt{-4i}\log\left(-\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{\sqrt{4i^2+1}}{ax+i}}\right) - \log\left(\sqrt{\frac{\sqrt{4i^2+1}}{ax+i}} + 1\right) + i\log\left(\sqrt{\frac{\sqrt{4i^2+1}}{ax+i}} + i\right) - i\log\left(\sqrt{\frac{\sqrt{4i^2+1}}{ax+i}} - i\right) + \log\left(\sqrt{\frac{\sqrt{4i^2+1}}{ax+i}} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x, algorithm="fricas")`

[Out] `1/2*sqrt(4*I)*log(1/2*I*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/2*sqrt(4*I)*log(-1/2*I*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/2*sqrt(-4*I)*log(1/2*I*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(-4*I)*log(-1/2*I*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x,x)`

[Out] `Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)/x, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{3/2}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x, x)
```

$$3.75 \quad \int \frac{e^{\frac{3}{2}i \operatorname{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=92

$$-\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} + 3ia \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 3ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x+3*I*a*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-3*I*a*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.02, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5170, 96, 95, 304, 209, 212}

$$3ia \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} - 3ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] `Int[E^(((3*I)/2)*ArcTan[a*x])/x^2,x]`

[Out] $-\left(\frac{(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}}{x}\right) + (3*I)*a*\operatorname{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}] - (3*I)*a*\operatorname{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]$

Rule 95

`Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

Rule 96

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

Rule 209

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a`

, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{\frac{3}{2}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1+iax)^{3/4}}{x^2(1-iax)^{3/4}} dx \\
 &= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} + \frac{1}{2}(3ia) \int \frac{1}{x(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
 &= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} + (6ia) \text{Subst} \left(\int \frac{x^2}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} - (3ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + (3ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} + 3ia \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - 3ia \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 68, normalized size = 0.74

$$-\frac{i\sqrt[4]{1-iax}(-i+ax+6ax {}_2F_1(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}))}{x^4\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])/x^2,x]

[Out] ((-I)*(1 - I*a*x)^(1/4)*(-I + a*x + 6*a*x*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(x*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2,x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)/x^2, x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 157 vs. $2(64) = 128$.

time = 1.53, size = 157, normalized size = 1.71

$$\frac{-3i ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - 3ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + 3ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) + 3i ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2,x, algorithm="fricas")

[Out] 1/2*(-3*I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 3*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 3*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + 3*I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**2,x)
```

```
[Out] Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)/x**2, x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{3/2}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x^2,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x^2, x)
```

3.76 $\int \frac{e^{\frac{3}{2}i\text{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=132

$$-\frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} - \frac{9}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{9}{4}a^2\text{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-3/4*I*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x-1/2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(7/4)}/x^2-9/4*a^2*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+9/4*a^2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.03, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5170, 98, 96, 95, 304, 209, 212}

$$-\frac{9}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{9}{4}a^2\text{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x}$$

Antiderivative was successfully verified.

[In] `Int[E^(((3*I)/2)*ArcTan[a*x])/x^3,x]`

[Out] $(((-3*I)/4)*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x - ((1-I*a*x)^{(1/4)}*(1+I*a*x)^{(7/4)}/(2*x^2) - (9*a^2*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/4 + (9*a^2*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/4)$

Rule 95

```
Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 96

```
Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]
```

Rule 98


```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])

```

Rule 209

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

```

Rule 304

```

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{3}{2}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1+iax)^{3/4}}{x^3(1-iax)^{3/4}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} + \frac{1}{4}(3ia) \int \frac{(1+iax)^{3/4}}{x^2(1-iax)^{3/4}} dx \\
&= -\frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} - \frac{1}{8}(9a^2) \int \frac{1}{x(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} - \frac{1}{2}(9a^2) \text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \frac{1+iax}{1-iax}\right) \\
&= -\frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} + \frac{1}{4}(9a^2) \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{1+iax}{1-iax}\right) \\
&= -\frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x} - \frac{\sqrt[4]{1-iax}(1+iax)^{7/4}}{2x^2} - \frac{9}{4}a^2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{9}{4}a^2
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 81, normalized size = 0.61

$$\frac{\sqrt[4]{1-iax}(-2-7iax+5a^2x^2+18a^2x^2{}_2F_1(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}))}{4x^2\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])/x^3,x]

[Out] ((1-I*a*x)^(1/4)*(-2-(7*I)*a*x+5*a^2*x^2+18*a^2*x^2*Hypergeometric2F1[1/4, 1, 5/4, (I+a*x)/(I-a*x)]))/(4*x^2*(1+I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x, algorithm="maxima")``[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)/x^3, x)`**Fricas [A]**

time = 1.88, size = 179, normalized size = 1.36

$$\frac{9a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - 9ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + 9ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 9a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2\sqrt{a^2x^2+1}(5i ax + 2)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{8x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x, algorithm="fricas")`

`[Out] 1/8*(9*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 9*I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 9*I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 9*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*sqrt(a^2*x^2 + 1)*(5*I*a*x + 2)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^2`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**3,x)``[Out] Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)/x**3, x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
 the root of a polynomial with parameters. This might be wrong.The choice wa
 s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+ax}{\sqrt{a^2x^2+1}}\right)^{3/2}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x+1)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x)

[Out] int(((a*x+1)/(a^2*x^2+1)^(1/2))^(3/2)/x^3, x)

$$3.77 \quad \int \frac{e^{\frac{3}{2}i \operatorname{ArcTan}(ax)}}{x^4} dx$$

Optimal. Leaf size=170

$$-\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} - \frac{17}{8}ia^3 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-1/3*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^3-7/12*I*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^2+23/24*a^2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x-17/8*I*a^3*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+17/8*I*a^3*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.05, antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 304, 209, 212}

$$-\frac{17}{8}ia^3 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{17}{8}ia^3 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} - \frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)/2)*\operatorname{ArcTan}[a*x]}/x^4, x]$

[Out] $-1/3*((1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/x^3 - (((7*I)/12)*a*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/x^2 + (23*a^2*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/(24*x) - ((17*I)/8)*a^3*\operatorname{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] + ((17*I)/8)*a^3*\operatorname{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}]$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match}Q[u, (b_)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 95

$\operatorname{Int}[(((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)})/((e_.) + (f_.)*(x_.))], x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Denominator}[m]\}, \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 101

$\operatorname{Int}[(a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x] - \operatorname{Dist}[1/((m+1)*(b*e - a*f)), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x]$

$(m + 1)(c + dx)^{n-1}(e + fx)^p \text{Simp}[d^*e^n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x] /;$ FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 156

$\text{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}((c_.) + (d_.)*(x_.)^{(n_.)}((e_.) + (f_.)*(x_.)^{(p_.)}((g_.) + (h_.)*(x_.)^{(n_.)}))], x_Symbol] := \text{Simp}[(b*g - a*h)*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*(e + f*x)^{(p + 1)} / ((m + 1)*(b*c - a*d)*(b*e - a*f))], x] + \text{Dist}[1 / ((m + 1)*(b*c - a*d)*(b*e - a*f)), \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p \text{Simp}[a*d*f*g - b*(d*e + c*f)*g + b*c*e*h*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /;$ FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

Rule 209

$\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \text{Simp}[(1 / (\text{Rt}[a, 2]*\text{Rt}[b, 2]))* \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

$\text{Int}[(a_.) + (b_.)*(x_.)^2)^{-1}, x_Symbol] := \text{Simp}[(1 / (\text{Rt}[a, 2]*\text{Rt}[-b, 2]))* \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /;$ FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

$\text{Int}[(x_.)^2 / ((a_.) + (b_.)*(x_.)^4), x_Symbol] := \text{With}[\{r = \text{Numerator}[\text{Rt}[-a/b, 2]], s = \text{Denominator}[\text{Rt}[-a/b, 2]]\}, \text{Dist}[s/(2*b), \text{Int}[1/(r + s*x^2), x], x] - \text{Dist}[s/(2*b), \text{Int}[1/(r - s*x^2), x], x]] /;$ FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)] * (n_.) * (x_.)^{(m_.)})}, x_Symbol] := \text{Int}[x^m * ((1 - I*a*x)^{(I*(n/2))} / (1 + I*a*x)^{(I*(n/2))}), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{3}{2}i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1+iax)^{3/4}}{x^4(1-iax)^{3/4}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} + \frac{1}{3} \int \frac{\frac{7ia}{2} - 2a^2x}{x^3(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} - \frac{1}{6} \int \frac{\frac{23a^2}{4} + \frac{7}{2}ia^3x}{x^2(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} + \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} - \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} - \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} + \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} - \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{23a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} -
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 93, normalized size = 0.55

$$\frac{\sqrt[4]{1-iax}(-8-22iax+37a^2x^2+23ia^3x^3+102ia^3x^3{}_2F_1(\frac{1}{4}, 1; \frac{5}{4}, \frac{i+ax}{i-ax}))}{24x^3\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])/x^4,x]

[Out] ((1 - I*a*x)^(1/4)*(-8 - (22*I)*a*x + 37*a^2*x^2 + (23*I)*a^3*x^3 + (102*I)*a^3*x^3*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(24*x^3*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x, algorithm="maxima")`

[Out] `integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)/x^4, x)`

Fricas [A]

time = 2.73, size = 187, normalized size = 1.10

$$\frac{51i a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2 x^2 + 1}}{ax + i}} + 1\right) + 51 a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2 x^2 + 1}}{ax + i}} + i\right) - 51 a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2 x^2 + 1}}{ax + i}} - i\right) - 51i a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2 x^2 + 1}}{ax + i}} - 1\right) + 2(23 a^2 x^2 - 14i ax - 8)\sqrt{a^2 x^2 + 1} \sqrt{\frac{i\sqrt{a^2 x^2 + 1}}{ax + i}}}{48 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x, algorithm="fricas")`

[Out] `1/48*(51*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 51*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 51*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 51*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) + 2*(23*a^2*x^2 - 14*I*a*x - 8)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^3`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(ax-i)}{\sqrt{a^2 x^2 + 1}}\right)^{\frac{3}{2}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**4,x)`

[Out] `Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)/x**4, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.01
```

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{3/2}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x^4,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x^4, x)
```

$$3.78 \quad \int \frac{e^{\frac{3}{2}i \operatorname{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=202

$$-\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{63ia^3\sqrt[4]{1-iax}(1+iax)^{3/4}}{64x}$$

[Out] $-1/4*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^4-3/8*I*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^3+15/32*a^2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^2+63/64*I*a^3*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x+123/64*a^4*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-123/64*a^4*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.06, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 304, 209, 212}

$$\frac{123}{64}a^4 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{123}{64}a^4 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{63ia^3\sqrt[4]{1-iax}(1+iax)^{3/4}}{64x} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} - \frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)/2)*\operatorname{ArcTan}[a*x]}/x^5, x]$

[Out] $-1/4*((1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/x^4 - (((3*I)/8)*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/x^3 + (15*a^2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/(32*x^2) + (((63*I)/64)*a^3*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/x + (123*a^4*\operatorname{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64 - (123*a^4*\operatorname{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match}Q[u, (b_*)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 95

$\operatorname{Int}[(((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)})/((e_.) + (f_.)*(x_)), x_Symbol] \rightarrow \operatorname{With}[q = \operatorname{Denominator}[m], \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{q*(m+1)-1}/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 101

$\operatorname{Int}[((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)}$

)/((m + 1)*(b*e - a*f))), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 156

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{3}{2}i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{(1+iax)^{3/4}}{x^5(1-iax)^{3/4}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} + \frac{1}{4} \int \frac{\frac{9ia}{2} - 3a^2x}{x^4(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} - \frac{1}{12} \int \frac{\frac{45a^2}{4} + 9ia^3x}{x^3(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{1}{2} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{x^2(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{6}{5} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{x(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{6}{5} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{6}{5} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{1-iax} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{6}{5} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{1-iax} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{6}{5} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{1-iax} dx \\
&= -\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} - \frac{3ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{8x^3} + \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{32x^2} + \frac{6}{5} \int \frac{15a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{1-iax} dx
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 99, normalized size = 0.49

$$-\frac{\sqrt[4]{1-iax}(16+40iax-54a^2x^2-93ia^3x^3+63a^4x^4+246a^4x^4{}_2F_1(\frac{1}{4}, 1; \frac{5}{4}, \frac{i+ax}{i-ax}))}{64x^4\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])/x^5,x]

[Out] -1/64*((1-I*a*x)^(1/4)*(16+(40*I)*a*x-54*a^2*x^2-(93*I)*a^3*x^3+63*a^4*x^4+246*a^4*x^4*Hypergeometric2F1[1/4,1,5/4,(I+a*x)/(I-a*x)]))/(x^4*(1+I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x, algorithm="maxima")`

[Out] `integrate((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)/x^5, x)`

Fricas [A]

time = 3.07, size = 195, normalized size = 0.97

$$\frac{123a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) - 123ia^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) + 123ia^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) - 123a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) + 2(-63ia^3x^3 - 30a^2x^2 + 24iax + 16)\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{128x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x, algorithm="fricas")`

[Out] `-1/128*(123*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 123*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 123*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 123*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) + 2*(-63*I*a^3*x^3 - 30*a^2*x^2 + 24*I*a*x + 16)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^4`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**5,x)`

[Out] `Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)/x**5, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+axi}{\sqrt{a^2x^2+1}}\right)^{3/2}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x^5,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)/x^5, x)
```

3.79 $\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=373

$$\frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i-452ax)(1-iax)^{3/4}}{96a^4}$$

[Out] $475/64*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^4-4*I*x^3*(1+I*a*x)^{(5/4)}/a/(1-I*a*x)^{(1/4)}-17/4*x^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^2-1/96*I*(521*I-452*a*x)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^4-475/128*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4)}/a^4*2^{(1/2)}+475/128*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4)}/a^4*2^{(1/2)}+475/256*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)}/a^4*2^{(1/2)}-475/256*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)}/a^4*2^{(1/2)}$

Rubi [A]

time = 0.18, antiderivative size = 373, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 13, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.812$, Rules used = {5170, 99, 158, 152, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{475\text{ArcTan}\left(\frac{1-\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} + \frac{475\text{ArcTan}\left(\frac{1+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} - \frac{i(-452ax+521i)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} + \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} + \frac{475\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{128\sqrt{2}a^4} - \frac{475\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{128\sqrt{2}a^4} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt{1-iax}}$$

Antiderivative was successfully verified.

[In] Int[E^(((5*I)/2)*ArcTan[a*x])*x^3,x]

[Out] $(475*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(64*a^4) - ((4*I)*x^3*(1+I*a*x)^{(5/4)})/(a*(1-I*a*x)^{(1/4)}) - (17*x^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)})/(4*a^2) - ((I/96)*(521*I-452*a*x)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)})/a^4 - (475*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(64*\text{Sqrt}[2]*a^4) + (475*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(64*\text{Sqrt}[2]*a^4) + (475*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(128*\text{Sqrt}[2]*a^4) - (475*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(128*\text{Sqrt}[2]*a^4)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n)/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 99

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_, x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(
m + 1))), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*
(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{
a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ
[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))*(g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 158

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n +
1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p
+ 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n +
p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p
+ 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /
; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p +
2, 0] && IntegerQ[m]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303


```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] & AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{5}{2}i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1+iax)^{5/4}}{(1-iax)^{5/4}} dx \\
&= -\frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} + \frac{(4i) \int \frac{x^2\sqrt[4]{1+iax} (3+\frac{17iax}{4})}{\sqrt[4]{1-iax}} dx}{a} \\
&= -\frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} + \frac{i \int \frac{x\sqrt[4]{1+iax} (-\frac{17ia}{2} + \frac{113a^2x}{8})}{\sqrt[4]{1-iax}} dx}{a^3} \\
&= -\frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4} \\
&= \frac{475(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} - \frac{4ix^3(1+iax)^{5/4}}{a^4\sqrt[4]{1-iax}} - \frac{17x^2(1-iax)^{3/4}(1+iax)^{5/4}}{4a^2} - \frac{i(521i - 452ax)(1-iax)^{3/4}(1+iax)^{5/4}}{96a^4}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 96, normalized size = 0.26

$$\frac{-\sqrt[4]{1+iax} (-i+ax)^2 (59-5iax+6a^2x^2) + 380\sqrt[4]{2} (1-iax) {}_2F_1\left(-\frac{5}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax)\right)}{24a^4\sqrt[4]{1-iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])*x^3,x]

[Out] $(-((1 + I*a*x)^{(1/4)}*(-I + a*x)^2*(59 - (5*I)*a*x + 6*a^2*x^2)) + 380*2^{(1/4)}*(1 - I*a*x)*\text{Hypergeometric2F1}[-5/4, 3/4, 7/4, (1 - I*a*x)/2])/(24*a^4*(1 - I*a*x)^{(1/4)})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}} x^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^3,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^3,x, algorithm="maxima")

[Out] integrate(x^3*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)

Fricas [A]

time = 1.88, size = 251, normalized size = 0.67

$$\frac{96a^4\sqrt{\frac{225625}{4096a^2}} \log\left(\frac{\frac{25}{24}a^4\sqrt{\frac{225625}{4096a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+1}}}{-96a^4\sqrt{\frac{225625}{4096a^2}} \log\left(-\frac{25}{24}a^4\sqrt{\frac{225625}{4096a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+1}}\right) + 96a^4\sqrt{\frac{225625}{4096a^2}} \log\left(\frac{25}{24}a^4\sqrt{\frac{225625}{4096a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+1}}\right) - 96a^4\sqrt{-\frac{225625}{4096a^2}} \log\left(-\frac{25}{24}a^4\sqrt{\frac{225625}{4096a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+1}}\right) + (48a^4x^4 - 136Ia^3x^3 - 226a^2x^2 + 521Ia*x - 2467)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+1}}}{192a^4}\right)}{192a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^3,x, algorithm="fricas")

[Out] $-1/192*(96*a^4*\text{sqrt}(225625/4096*I/a^8)*\log(64/475*a^4*\text{sqrt}(225625/4096*I/a^8) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - 96*a^4*\text{sqrt}(225625/4096*I/a^8)*\log(-64/475*a^4*\text{sqrt}(225625/4096*I/a^8) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) + 96*a^4*\text{sqrt}(-225625/4096*I/a^8)*\log(64/475*a^4*\text{sqrt}(-225625/4096*I/a^8) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - 96*a^4*\text{sqrt}(-225625/4096*I/a^8)*\log(-64/475*a^4*\text{sqrt}(-225625/4096*I/a^8) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) + (48*a^4*x^4 - 136*I*a^3*x^3 - 226*a^2*x^2 + 521*I*a*x - 2467)*\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)))/a^4$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)*x**3,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3877 deep

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^3,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^3 \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)

[Out] int(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)

3.80 $\int e^{\frac{5}{2}i \operatorname{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=371

$$\frac{55i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3\sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} - \frac{55i \operatorname{ArcTan}\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{55i \operatorname{ArcTan}\left(\frac{1+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{3a^3} + \frac{2i(1+iax)^{9/4}}{a^3\sqrt[4]{1-iax}} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{55i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} + \frac{55i \log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{16\sqrt{2}a^3} - \frac{55i \log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{16\sqrt{2}a^3}$$

[Out] $55/8*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^3+11/4*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^3+2*I*(1+I*a*x)^{(9/4)}/a^3/(1-I*a*x)^{(1/4)}+1/3*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(9/4)}/a^3-55/16*I*\operatorname{arctan}(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3*2^{(1/2)}+55/16*I*\operatorname{arctan}(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3*2^{(1/2)}+55/32*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3*2^{(1/2)}-55/32*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3*2^{(1/2)}$

Rubi [A]

time = 0.18, antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5170, 91, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$-\frac{55i \operatorname{ArcTan}\left(\frac{1-\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{55i \operatorname{ArcTan}\left(\frac{1+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{i(1-iax)^{3/4}(1+iax)^{5/4}}{3a^3} + \frac{2i(1+iax)^{9/4}}{a^3\sqrt[4]{1-iax}} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{55i(1-iax)^{3/4}\sqrt[4]{1+iax}}{8a^3} + \frac{55i \log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{16\sqrt{2}a^3} - \frac{55i \log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{16\sqrt{2}a^3}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((5*I)/2)*\operatorname{ArcTan}[a*x]}*x^2, x]$

[Out] $((55*I)/8)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^3 + ((11*I)/4)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^3 + ((2*I)*(1+I*a*x)^{(9/4)})/(a^3*(1-I*a*x)^{(1/4)}) + ((I/3)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(9/4)})/a^3 - ((55*I)/8)*\operatorname{ArcTan}[1-(\operatorname{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]/(\operatorname{Sqrt}[2]*a^3) + ((55*I)/8)*\operatorname{ArcTan}[1+(\operatorname{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]/(\operatorname{Sqrt}[2]*a^3) + ((55*I)/16)*\operatorname{Log}[1+\operatorname{Sqrt}[1-I*a*x]/\operatorname{Sqrt}[1+I*a*x]-(\operatorname{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]/(\operatorname{Sqrt}[2]*a^3) - ((55*I)/16)*\operatorname{Log}[1+\operatorname{Sqrt}[1-I*a*x]/\operatorname{Sqrt}[1+I*a*x]+(\operatorname{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]/(\operatorname{Sqrt}[2]*a^3)$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \operatorname{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \operatorname{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 91

```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] := Simp[(b*c - a*d)^(2*(c + d*x)^(n + 1))*((e + f*x)^(p + 1)
/(d^2*(d*e - c*f)*(n + 1))), x] - Dist[1/(d^2*(d*e - c*f)*(n + 1)), Int[(c
+ d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1)
+ c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n
+ 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] ||
(EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p,
1])))
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &&
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{5}{2}i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1+iax)^{5/4}}{(1-iax)^{5/4}} dx \\
&= \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} - \frac{(2i) \int \frac{(1+iax)^{5/4} \left(\frac{5ia}{2} - \frac{a^2 x}{2}\right)}{\sqrt[4]{1-iax}} dx}{a^3} \\
&= \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} + \frac{11 \int \frac{(1+iax)^{5/4}}{\sqrt[4]{1-iax}} dx}{2a^2} \\
&= \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} + \frac{55 \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx}{8a^2} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3} \\
&= \frac{55i(1-iax)^{3/4} \sqrt[4]{1+iax}}{8a^3} + \frac{11i(1-iax)^{3/4}(1+iax)^{5/4}}{4a^3} + \frac{2i(1+iax)^{9/4}}{a^3 \sqrt[4]{1-iax}} + \frac{i(1-iax)^{3/4}(1+iax)^{9/4}}{3a^3}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 86, normalized size = 0.23

$$\frac{-\sqrt[4]{1+iax} (-i+ax)^2(7i+ax) + 44\sqrt[4]{2} (i+ax) {}_2F_1\left(-\frac{5}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax)\right)}{3a^3 \sqrt[4]{1-iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])*x^2,x]

[Out] $(-(1 + I*a*x)^{1/4}*(-I + a*x)^{2*(7*I + a*x)} + 44*2^{1/4}*(I + a*x)*\text{Hypergeometric2F1}[-5/4, 3/4, 7/4, (1 - I*a*x)/2])/(3*a^3*(1 - I*a*x)^{1/4})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)*x^2,x)

[Out] int((((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)*x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)*x^2,x, algorithm="maxima")

[Out] integrate(x^2*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)

Fricas [A]

time = 1.95, size = 244, normalized size = 0.66

$$\frac{12a^3\sqrt{\frac{3025}{64a^6}} \log\left(\frac{5}{11}ia^3\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1\sqrt{a^2x^2+1}}{ax+i}}\right) - 12a^3\sqrt{\frac{3025}{64a^6}} \log\left(-\frac{5}{11}ia^3\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1\sqrt{a^2x^2+1}}{ax+i}}\right) + 12a^3\sqrt{\frac{3025}{64a^6}} \log\left(\frac{5}{11}ia^3\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1\sqrt{a^2x^2+1}}{ax+i}}\right) - 12a^3\sqrt{\frac{3025}{64a^6}} \log\left(-\frac{5}{11}ia^3\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1\sqrt{a^2x^2+1}}{ax+i}}\right) - (8a^3x^3 - 26i a^2x^2 - 61ax - 287i)\sqrt{\frac{1\sqrt{a^2x^2+1}}{ax+i}}}{24a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)*x^2,x, algorithm="fricas")

[Out] $\frac{1}{24}*(12*a^3*\text{sqrt}(3025/64*I/a^6)*\log(8/55*I*a^3*\text{sqrt}(3025/64*I/a^6)) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - 12*a^3*\text{sqrt}(3025/64*I/a^6)*\log(-8/55*I*a^3*\text{sqrt}(3025/64*I/a^6)) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) + 12*a^3*\text{sqrt}(-3025/64*I/a^6)*\log(8/55*I*a^3*\text{sqrt}(-3025/64*I/a^6)) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - 12*a^3*\text{sqrt}(-3025/64*I/a^6)*\log(-8/55*I*a^3*\text{sqrt}(-3025/64*I/a^6)) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - (8*a^3*x^3 - 26*I*a^2*x^2 - 61*a*x - 287*I)*\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)))/a^3$

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)*x**2,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3061 deep
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^2,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int x^2 \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)
```

```
[Out] int(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

3.81 $\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=324

$$\frac{25(1-iax)^{3/4}\sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2\sqrt[4]{1-iax}} + \frac{25\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2}$$

[Out] $-25/4*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^2-5/2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(5/4)}/a^2-2*(1+I*a*x)^{(9/4)}/a^2/(1-I*a*x)^{(1/4)}+25/8*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2*2^{(1/2)}-25/8*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2*2^{(1/2)}-25/16*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^2*2^{(1/2)}+25/16*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^2*2^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 324, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 79, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{25\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{25\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{2(1+iax)^{9/4}}{a^2\sqrt[4]{1-iax}} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{25(1-iax)^{3/4}\sqrt[4]{1+iax}}{4a^2} - \frac{25\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2} + \frac{25\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((5*I)/2)*\text{ArcTan}[a*x]}*x, x]$

[Out] $(-25*(1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(1/4)})/(4*a^2) - (5*(1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(5/4)})/(2*a^2) - (2*(1 + I*a*x)^{(9/4)})/(a^2*(1 - I*a*x)^{(1/4)}) + (25*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(4*\text{Sqrt}[2]*a^2) - (25*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(4*\text{Sqrt}[2]*a^2) - (25*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(8*\text{Sqrt}[2]*a^2) + (25*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(8*\text{Sqrt}[2]*a^2)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) +$

$d*(x^p/b)^n, x, (a + b*x)^{1/p}, x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 79

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(-(b*e - a*f))*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^(p + 1), x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || (EqQ[e, 0] || (EqQ[c, 0] || LtQ[p, n]))))

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 338

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,

e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] :> Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int e^{\frac{5}{2}i \tan^{-1}(ax)} x dx &= \int \frac{x(1+iax)^{5/4}}{(1-iax)^{5/4}} dx \\
&= -\frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} + \frac{(5i) \int \frac{(1+iax)^{5/4}}{\sqrt[4]{1-iax}} dx}{a} \\
&= -\frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} + \frac{(25i) \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} + \frac{(25i) \int \frac{1}{\sqrt[4]{1-iax}} dx}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} - \frac{25 \text{Subst}\left(\int \frac{1}{\sqrt[4]{1-iax}} dx\right)}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} - \frac{25 \text{Subst}\left(\int \frac{1}{\sqrt[4]{1-iax}} dx\right)}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} + \frac{25 \text{Subst}\left(\int \frac{1}{\sqrt[4]{1-iax}} dx\right)}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} - \frac{25 \text{Subst}\left(\int \frac{1}{\sqrt[4]{1-iax}} dx\right)}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} - \frac{25 \log\left(1 + \sqrt[4]{1-iax}\right)}{4a} \\
&= -\frac{25(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} - \frac{5(1-iax)^{3/4}(1+iax)^{5/4}}{2a^2} - \frac{2(1+iax)^{9/4}}{a^2 \sqrt[4]{1-iax}} + \frac{25 \tan^{-1}\left(1 + \sqrt[4]{1-iax}\right)}{4a}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 72, normalized size = 0.22

$$\frac{2\left(-3(1+iax)^{9/4} + 20i\sqrt[4]{2}(i+ax) {}_2F_1\left(-\frac{5}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax)\right)\right)}{3a^2 \sqrt[4]{1-iax}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])*x,x]

[Out] $(2*(-3*(1 + I*a*x)^{(9/4)} + (20*I)*2^{(1/4)}*(I + a*x)*\text{Hypergeometric2F1}[-5/4, 3/4, 7/4, (1 - I*a*x)/2]))/(3*a^2*(1 - I*a*x)^{(1/4)})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x,x, algorithm="maxima")`

[Out] `integrate(x*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)`

Fricas [A]

time = 1.96, size = 236, normalized size = 0.73

$$\frac{2a^2\sqrt{\frac{625i}{16a^4}}\log\left(\frac{\frac{5}{2}a^2\sqrt{\frac{625i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{-2a^2\sqrt{\frac{625i}{16a^4}}\log\left(-\frac{5}{2}a^2\sqrt{\frac{625i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + 2a^2\sqrt{\frac{625i}{16a^4}}\log\left(\frac{5}{2}a^2\sqrt{\frac{625i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 2a^2\sqrt{\frac{625i}{16a^4}}\log\left(-\frac{5}{2}a^2\sqrt{\frac{625i}{16a^4}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - (2a^2x^2 - 9iax + 43)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)}{4a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x,x, algorithm="fricas")`

[Out] $\frac{1}{4}*(2*a^2*\text{sqrt}(625/16*I/a^4)*\log(4/25*a^2*\text{sqrt}(625/16*I/a^4) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*\text{sqrt}(625/16*I/a^4)*\log(-4/25*a^2*\text{sqrt}(625/16*I/a^4) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) + 2*a^2*\text{sqrt}(-625/16*I/a^4)*\log(4/25*a^2*\text{sqrt}(-625/16*I/a^4) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*\text{sqrt}(-625/16*I/a^4)*\log(-4/25*a^2*\text{sqrt}(-625/16*I/a^4) + \text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I))) - (2*a^2*x^2 - 9*I*a*x + 43)*\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)))/a^2$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)*x,x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
 dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)

[Out] int(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)

3.82 $\int e^{\frac{5}{2}i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=299

$$\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} + \frac{5i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{5i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a}$$

[Out] $-5*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a-4*I*(1+I*a*x)^{(5/4)}/a/(1-I*a*x)^{(1/4)}+5/2*I*\text{arctan}(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}-5/2*I*\text{arctan}(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}-5/4*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}+5/4*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}$

Rubi [A]

time = 0.13, antiderivative size = 299, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {5169, 49, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{5i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{5i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - \frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{5i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a} + \frac{5i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((5*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $((-5*I)*(1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(1/4)})/a - ((4*I)*(1 + I*a*x)^{(5/4)})/(a*(1 - I*a*x)^{(1/4)}) + ((5*I)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) - ((5*I)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) - (((5*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) + (((5*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a)$

Rule 49

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + 1))), x] - \text{Dist}[d*(n/(b*(m + 1))), \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{LtQ}[m, -1] \&\& !(\text{IntegerQ}[n] \&\& !\text{IntegerQ}[m]) \&\& !(\text{ILeQ}[m + n + 2, 0] \&\& (\text{FractionQ}[m] || \text{GeQ}[2*n + m + 1, 0])) \& \& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/($

```
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5169

```
Int[E^(ArcTan[(a_)*(x_)])*(n_.), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1
+ I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{5}{2}i \tan^{-1}(ax)} dx &= \int \frac{(1+iax)^{5/4}}{(1-iax)^{5/4}} dx \\
&= -\frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - 5 \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - \frac{5}{2} \int \frac{1}{\sqrt[4]{1-iax}(1+iax)^{3/4}} dx \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - \frac{(10i)\text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax}\right)}{a} \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - \frac{(10i)\text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} + \frac{(5i)\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - \frac{(5i)\text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2a} \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} - \frac{5i \log\left(1 + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2\sqrt{2}a} \\
&= -\frac{5i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{4i(1+iax)^{5/4}}{a\sqrt[4]{1-iax}} + \frac{5i \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{5i \tan^{-1}\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 41, normalized size = 0.14

$$-\frac{8ie^{\frac{9}{2}i \text{ArcTan}(ax)} {}_2F_1\left(2, \frac{9}{4}, \frac{13}{4}; -e^{2i \text{ArcTan}(ax)}\right)}{9a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] (((-8*I)/9)*E^(((9*I)/2)*ArcTan[a*x])*Hypergeometric2F1[2, 9/4, 13/4, -E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)

Fricas [A]

time = 2.56, size = 209, normalized size = 0.70

$$\frac{a\sqrt{\frac{25i}{a^2}} \log\left(\frac{\frac{1}{2}ia\sqrt{\frac{25i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{a}\right) - a\sqrt{\frac{25i}{a^2}} \log\left(-\frac{1}{2}ia\sqrt{\frac{25i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + a\sqrt{-\frac{25i}{a^2}} \log\left(\frac{\frac{1}{2}ia\sqrt{-\frac{25i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{a}\right) - a\sqrt{-\frac{25i}{a^2}} \log\left(-\frac{1}{2}ia\sqrt{-\frac{25i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + 2(ax+9i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="fricas")

[Out] -1/2*(a*sqrt(25*I/a^2)*log(1/5*I*a*sqrt(25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(25*I/a^2)*log(-1/5*I*a*sqrt(25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + a*sqrt(-25*I/a^2)*log(1/5*I*a*sqrt(-25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(-25*I/a^2)*log(-1/5*I*a*sqrt(-25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*(a*x + 9*I)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/a

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2),x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)

[Out] int(((a*x*i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)

3.83 $\int \frac{e^{\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=293

$$\frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 2\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)$$

[Out] $8*(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}-2*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$
 $-2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+1/2*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)})$
 $/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2))*2^{(1/2)}-1/2*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)})$
 $/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2))*2^{(1/2)}-\operatorname{arctan}(1-(1-I*a*x)^{(1/4)}*2^{(1/2)})$
 $/(1+I*a*x)^{(1/4))*2^{(1/2)}+\operatorname{arctan}(1+(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4))*2^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 293, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 16, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5170, 100, 21, 132, 65, 338, 303, 1176, 631, 210, 1179, 642, 95, 218, 212, 209}

$$-2\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} + \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - 2\operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((5*I)/2)*\operatorname{ArcTan}[a*x]}/x, x]$

[Out] $(8*(1 + I*a*x)^{(1/4)})/(1 - I*a*x)^{(1/4)} - 2*\operatorname{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] - \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}] + \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}] - 2*\operatorname{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] + \operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a*x]/\operatorname{Sqrt}[1 + I*a*x] - (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/\operatorname{Sqrt}[2] - \operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a*x]/\operatorname{Sqrt}[1 + I*a*x] + (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/\operatorname{Sqrt}[2]$

Rule 21

$\operatorname{Int}[(u_*)*((a_*) + (b_*)*(v_*))^{(m_*)}*((c_*) + (d_*)*(v_*))^{(n_*)}, x_Symbol] \rightarrow \operatorname{Dist}[(b/d)^m, \operatorname{Int}[u*(c + d*v)^{(m+n)}, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])

Rule 65

$\operatorname{Int}[(a_*) + (b_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}), x_Symbol] \rightarrow \operatorname{With}[\{p = \operatorname{Denominator}[m]\}, \operatorname{Dist}[p/b, \operatorname{Subst}[\operatorname{Int}[x^{(p*(m+1)-1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ

[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 132

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt

Q[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] & AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 338

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{5}{2}i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1+iax)^{5/4}}{x(1-iax)^{5/4}} dx \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} + \frac{(4i) \int \frac{-\frac{ia}{4} - \frac{a^2x}{4}}{x\sqrt[4]{1-iax}(1+iax)^{3/4}} dx}{a} \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} + \int \frac{(1-iax)^{3/4}}{x(1+iax)^{3/4}} dx \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - (ia) \int \frac{1}{\sqrt[4]{1-iax}(1+iax)^{3/4}} dx + \int \frac{1}{x\sqrt[4]{1-iax}(1+iax)^{3/4}} dx \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} + 4\text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax}\right) + 4\text{Subst}\left(\int \frac{1}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{\text{Subst}\left(\int \frac{\sqrt{2}+2x}{-1-\sqrt{2}x-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)}{\sqrt{2}} \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{\sqrt{2}} \\
&= \frac{8\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2} \tan^{-1}\left(\frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 112, normalized size = 0.38

$$\frac{4\left(3 + 3iax + 3\sqrt[4]{2}(1 + iax)^{3/4} {}_2F_1\left(-\frac{1}{4}, -\frac{1}{4}; \frac{3}{4}; \frac{1}{2}(1 - iax)\right) + (-1 + iax) {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}\right)\right)}{3\sqrt[4]{1 - iax}(1 + iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])/x,x]

[Out] (4*(3 + (3*I)*a*x + 3*2^(1/4)*(1 + I*a*x)^(3/4)*Hypergeometric2F1[-1/4, -1/4, 3/4, (1 - I*a*x)/2] + (-1 + I*a*x)*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)])/(3*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x,x, algorithm="maxima")

[Out] integrate((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)/x, x)

Fricas [A]

time = 1.25, size = 267, normalized size = 0.91

$$-\frac{1}{2}\sqrt{a}\log\left(\frac{1}{2}\sqrt{a} + \sqrt{\frac{i\sqrt{a^2+1}}{az+1}}\right) + \frac{1}{2}\sqrt{a}\log\left(-\frac{1}{2}\sqrt{a} + \sqrt{\frac{i\sqrt{a^2+1}}{az+1}}\right) - \frac{1}{2}\sqrt{-a}\log\left(\frac{1}{2}\sqrt{-a} + \sqrt{\frac{i\sqrt{a^2+1}}{az+1}}\right) + \frac{1}{2}\sqrt{-a}\log\left(-\frac{1}{2}\sqrt{-a} + \sqrt{\frac{i\sqrt{a^2+1}}{az+1}}\right) + 8\sqrt{\frac{i\sqrt{a^2+1}}{az+1}} - \log\left(\sqrt{\frac{i\sqrt{a^2+1}}{az+1}} + 1\right) - i\log\left(\sqrt{\frac{i\sqrt{a^2+1}}{az+1}} + i\right) + i\log\left(\sqrt{\frac{i\sqrt{a^2+1}}{az+1}} - i\right) + \log\left(\sqrt{\frac{i\sqrt{a^2+1}}{az+1}} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x,x, algorithm="fricas")

[Out] -1/2*sqrt(4*I)*log(1/2*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(4*I)*log(-1/2*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/

$2\sqrt{-4I}\log(1/2\sqrt{-4I} + \sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)}) + 1/2\sqrt{-4I}\log(-1/2\sqrt{-4I} + \sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)}) + 8\sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)} - \log(\sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)} + 1) - I\log(\sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)} + I) + I\log(\sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)} - I) + \log(\sqrt{I\sqrt{a^2x^2 + 1}/(ax + I)} - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x,x)

[Out] Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(5/2)/x, x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2x^2+1}}\right)^{5/2}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)/x,x)

[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)/x, x)

$$3.84 \quad \int \frac{e^{\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=121

$$\frac{10ia\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - \frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} - 5ia \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 5ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] 10*I*a*(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)-(1+I*a*x)^(5/4)/x/(1-I*a*x)^(1/4)-5*I*a*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-5*I*a*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))

Rubi [A]

time = 0.03, antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5170, 96, 95, 218, 212, 209}

$$-5ia \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} + \frac{10ia\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - 5ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] Int[E^(((5*I)/2)*ArcTan[a*x])/x^2,x]

[Out] ((10*I)*a*(1+I*a*x)^(1/4))/(1-I*a*x)^(1/4) - (1+I*a*x)^(5/4)/(x*(1-I*a*x)^(1/4)) - (5*I)*a*ArcTan[(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)] - (5*I)*a*ArcTanh[(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m+1)*(c + d*x)^n*(e + f*x)^(p+1)/((m+1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m+1)*(b*e - a*f))], Int[(a + b*x)^(m+1)*(c + d*x)^(n-1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))], x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
 \int \frac{e^{\frac{5}{2}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1+iax)^{5/4}}{x^2(1-iax)^{5/4}} dx \\
 &= -\frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} + \frac{1}{2}(5ia) \int \frac{\sqrt[4]{1+iax}}{x(1-iax)^{5/4}} dx \\
 &= \frac{10ia\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - \frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} + \frac{1}{2}(5ia) \int \frac{1}{x\sqrt[4]{1-iax}(1+iax)^{3/4}} dx \\
 &= \frac{10ia\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - \frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} + (10ia) \text{Subst} \left(\int \frac{1}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= \frac{10ia\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - \frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} - (5ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - (5ia) \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= \frac{10ia\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} - \frac{(1+iax)^{5/4}}{x\sqrt[4]{1-iax}} - 5ia \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - 5ia \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 87, normalized size = 0.72

$$\frac{-3(1 - 8iax + 9a^2x^2) - 10ax(i + ax) {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}\right)}{3x\sqrt[4]{1-iax}(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])/x^2,x]

[Out] (-3*(1 - (8*I)*a*x + 9*a^2*x^2) - 10*a*x*(I + a*x)*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)])/(3*x*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)/x^2, x)

Fricas [A]

time = 1.95, size = 152, normalized size = 1.26

$$\frac{-5iax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + 5ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - 5ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) + 5iax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2(-9iax + 1)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x, algorithm="fricas")

[Out] 1/2*(-5*I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 5*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 5*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + 5*I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(-9*I*a*x + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**2,x)

[Out] Timed out

Giac [F(-2)]
time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]
time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+ax}{\sqrt{a^2x^2+1}}\right)^{5/2}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x+1)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x)

[Out] int(((a*x+1)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x)

$$3.85 \quad \int \frac{e^{\frac{5}{2}i\text{ArcTan}(ax)}}{x^3} dx$$

Optimal. Leaf size=163

$$-\frac{25a^2\sqrt[4]{1+iax}}{2\sqrt[4]{1-iax}} - \frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}} - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} + \frac{25}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{25}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-25/2*a^2*(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}-5/4*I*a*(1+I*a*x)^{(5/4)}/x/(1-I*a*x)^{(1/4)}-1/2*(1+I*a*x)^{(9/4)}/x^2/(1-I*a*x)^{(1/4)}+25/4*a^2*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+25/4*a^2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.04, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5170, 98, 96, 95, 218, 212, 209}

$$\frac{25}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{25a^2\sqrt[4]{1+iax}}{2\sqrt[4]{1-iax}} + \frac{25}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} - \frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((5*I)/2)*\text{ArcTan}[a*x]}/x^3, x]$

[Out] $(-25*a^2*(1 + I*a*x)^{(1/4)})/(2*(1 - I*a*x)^{(1/4)}) - (((5*I)/4)*a*(1 + I*a*x)^{(5/4)})/(x*(1 - I*a*x)^{(1/4)}) - (1 + I*a*x)^{(9/4)}/(2*x^2*(1 - I*a*x)^{(1/4)}) + (25*a^2*\text{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}])/4 + (25*a^2*\text{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}])/4$

Rule 95

$\text{Int}[\frac{(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}}{(e_.) + (f_.)*(x_.)}, x_Symbol] :> \text{With}[\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{EqQ}[m + n + 1, 0] \&\& \text{RationalQ}[n] \&\& \text{LtQ}[-1, m, 0] \&\& \text{SimplerQ}[a + b*x, c + d*x]$

Rule 96

$\text{Int}[\frac{(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}}{x_Symbol] :> \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x] - \text{Dist}[n*((d*e - c*f)/((m+1)*(b*e - a*f))], \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, p\}, x] \&\& \text{EqQ}[m + n + p + 2, 0] \&\& \text{GtQ}[n, 0] \&\& (\text{SumSimplerQ}[m, 1] || !\text{SumSimplerQ}[p, 1]) \&\& \text{NeQ}[m, -1]$

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{5}{2}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1+iax)^{5/4}}{x^3(1-iax)^{5/4}} dx \\
&= -\frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} + \frac{1}{4}(5ia) \int \frac{(1+iax)^{5/4}}{x^2(1-iax)^{5/4}} dx \\
&= -\frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}} - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} - \frac{1}{8}(25a^2) \int \frac{\sqrt[4]{1+iax}}{x(1-iax)^{5/4}} dx \\
&= -\frac{25a^2\sqrt[4]{1+iax}}{2\sqrt[4]{1-iax}} - \frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}} - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} - \frac{1}{8}(25a^2) \int \frac{1}{x\sqrt[4]{1-iax}(1+iax)} dx \\
&= -\frac{25a^2\sqrt[4]{1+iax}}{2\sqrt[4]{1-iax}} - \frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}} - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} - \frac{1}{2}(25a^2) \text{Subst} \left(\int \frac{1}{-1+x^4} dx, x \right) \\
&= -\frac{25a^2\sqrt[4]{1+iax}}{2\sqrt[4]{1-iax}} - \frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}} - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} + \frac{1}{4}(25a^2) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x \right) \\
&= -\frac{25a^2\sqrt[4]{1+iax}}{2\sqrt[4]{1-iax}} - \frac{5ia(1+iax)^{5/4}}{4x\sqrt[4]{1-iax}} - \frac{(1+iax)^{9/4}}{2x^2\sqrt[4]{1-iax}} + \frac{25}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{25}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{25}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 99, normalized size = 0.61

$$\frac{-6 - 33iax - 102a^2x^2 - 129ia^3x^3 + 50a^2x^2(1-iax) {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}\right)}{12x^2\sqrt[4]{1-iax}(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])/x^3,x]

[Out] (-6 - (33*I)*a*x - 102*a^2*x^2 - (129*I)*a^3*x^3 + 50*a^2*x^2*(1 - I*a*x))*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]/(12*x^2*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^3,x)

[Out] $\int \left(\frac{(1+I*a*x)}{(a^2*x^2+1)^{(1/2)}} \right)^{(5/2)} / x^3, x$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}\left(\left(\frac{(1+I*a*x)}{(a^2*x^2+1)^{(1/2)}}\right)^{(5/2)} / x^3, x, \text{algorithm}="maxima"\right)$

[Out] $\text{integrate}\left(\left(\frac{(I*a*x + 1)}{\sqrt{a^2*x^2 + 1}}\right)^{(5/2)} / x^3, x\right)$

Fricas [A]

time = 2.02, size = 176, normalized size = 1.08

$$\frac{25a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + 25ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - 25ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 25a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2(43a^2x^2 + 9iax + 2)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{8x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}\left(\left(\frac{(1+I*a*x)}{(a^2*x^2+1)^{(1/2)}}\right)^{(5/2)} / x^3, x, \text{algorithm}="fricas"\right)$

[Out] $\frac{1}{8} * (25*a^2*x^2 * \log(\sqrt{(I*\sqrt{a^2*x^2 + 1}) / (a*x + I)} + 1) + 25*I*a^2*x^2 * \log(\sqrt{(I*\sqrt{a^2*x^2 + 1}) / (a*x + I)} + I) - 25*I*a^2*x^2 * \log(\sqrt{(I*\sqrt{a^2*x^2 + 1}) / (a*x + I)} - I) - 25*a^2*x^2 * \log(\sqrt{(I*\sqrt{a^2*x^2 + 1}) / (a*x + I)} - 1) - 2*(43*a^2*x^2 + 9*I*a*x + 2) * \sqrt{(I*\sqrt{a^2*x^2 + 1}) / (a*x + I)}) / x^2$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}\left(\left(\frac{(1+I*a*x)}{(a^2*x^2+1)^{(1/2)}}\right)^{(5/2)} / x^3, x\right)$

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}\left(\left(\frac{(1+I*a*x)}{(a^2*x^2+1)^{(1/2)}}\right)^{(5/2)} / x^3, x, \text{algorithm}="giac"\right)$

[Out] Exception raised: TypeError >> An error occurred running a Giac command: INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+axi}{\sqrt{a^2x^2+1}}\right)^{5/2}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)/x^3, x)

[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)/x^3, x)

$$3.86 \quad \int \frac{e^{\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^4} dx$$

Optimal. Leaf size=203

$$-\frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} + \frac{55}{8}ia^3 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{55}{8}ia^3$$

[Out] $-287/24*I*a^3*(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}-1/3*(1+I*a*x)^{(1/4)}/x^3/(1-I*a*x)^{(1/4)}-13/12*I*a*(1+I*a*x)^{(1/4)}/x^2/(1-I*a*x)^{(1/4)}+61/24*a^2*(1+I*a*x)^{(1/4)}/x/(1-I*a*x)^{(1/4)}+55/8*I*a^3*\operatorname{arctan}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+55/8*I*a^3*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.06, antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {5170, 100, 156, 160, 12, 95, 218, 212, 209}

$$\frac{55}{8}ia^3 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} + \frac{55}{8}ia^3 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((5*I)/2)*\operatorname{ArcTan}[a*x]}/x^4, x]$

[Out] $(((-287*I)/24)*a^3*(1 + I*a*x)^{(1/4)})/(1 - I*a*x)^{(1/4)} - (1 + I*a*x)^{(1/4)}/(3*x^3*(1 - I*a*x)^{(1/4)}) - (((13*I)/12)*a*(1 + I*a*x)^{(1/4)})/(x^2*(1 - I*a*x)^{(1/4)}) + (61*a^2*(1 + I*a*x)^{(1/4)})/(24*x*(1 - I*a*x)^{(1/4)}) + ((55*I)/8)*a^3*\operatorname{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] + ((55*I)/8)*a^3*\operatorname{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}]$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match}Q[u, (b_)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 95

$\operatorname{Int}[(((a_.) + (b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.))}/((e_.) + (f_.)*(x_))), x_Symbol] \rightarrow \operatorname{With}[{q = \operatorname{Denominator}[m]}, \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{q*(m+1)-1}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}[{a, b, c, d, e, f}, x] \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 100

$\operatorname{Int}[((a_.) + (b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.)*((e_.) + (f_.)*(x_))^{(p_.)}], x_Symbol] \rightarrow \operatorname{Simp}[(b*c - a*d)*(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}$

```

*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m
+ 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(
n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*
(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a,
b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2
*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n + 1)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 160

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n + 1)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m + n + p + 2,
0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || (!NeQ[n, -1] && SumSimplerQ[n,
1]) && !(NeQ[p, -1] && SumSimplerQ[p, 1]))

```

Rule 209

```

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

Rule 218

```

Int[((a_.) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b

```

, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{5}{2}i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1+iax)^{5/4}}{x^4(1-iax)^{5/4}} dx \\
&= -\frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{1}{3} \int \frac{-\frac{13ia}{2} + 6a^2x}{x^3(1-iax)^{5/4}(1+iax)^{3/4}} dx \\
&= -\frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{1}{6} \int \frac{-\frac{61a^2}{4} - 13ia^3x}{x^2(1-iax)^{5/4}(1+iax)^{3/4}} dx \\
&= -\frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} - \frac{1}{6} \int \frac{\frac{165ia^3}{8} - \frac{61a^4x}{4}}{x(1-iax)^{5/4}(1+iax)^{3/4}} dx \\
&= -\frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} - \frac{i \int \frac{\sqrt[4]{1+iax}}{16x\sqrt[4]{1-iax}} dx}{16} \\
&= -\frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} - \frac{1}{16}(55ia^3) \int \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx \\
&= -\frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} - \frac{1}{4}(55ia^3) \operatorname{Subst}\left(\int \frac{\sqrt[4]{1+u}}{\sqrt[4]{1-u}} du, ax\right) \\
&= -\frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} + \frac{1}{8}(55ia^3) \operatorname{Subst}\left(\int \frac{\sqrt[4]{1+u}}{\sqrt[4]{1-u}} du, ax\right) \\
&= -\frac{287ia^3\sqrt[4]{1+iax}}{24\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{3x^3\sqrt[4]{1-iax}} - \frac{13ia\sqrt[4]{1+iax}}{12x^2\sqrt[4]{1-iax}} + \frac{61a^2\sqrt[4]{1+iax}}{24x\sqrt[4]{1-iax}} + \frac{55}{8}ia^3 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 106, normalized size = 0.52

$$\frac{-8 - 34iax + 87a^2x^2 - 226ia^3x^3 + 287a^4x^4 + 110a^3x^3(i+ax) {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}\right)}{24x^3\sqrt[4]{1-iax}(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])/x^4,x]

[Out] $(-8 - (34*I)*a*x + 87*a^2*x^2 - (226*I)*a^3*x^3 + 287*a^4*x^4 + 110*a^3*x^3*(I + a*x)*\text{Hypergeometric2F1}[3/4, 1, 7/4, (I + a*x)/(I - a*x)])/(24*x^3*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)/x^4, x)

Fricas [A]

time = 1.93, size = 184, normalized size = 0.91

$$\frac{165i a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) - 165 a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) + 165 a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) - 165i a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) - 2(287i a^3 x^3 - 61 a^2 x^2 + 26i a x + 8) \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{48 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x, algorithm="fricas")

[Out] $\frac{1}{48}*(165*I*a^3*x^3*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) + 1) - 165*a^3*x^3*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) + I) + 165*a^3*x^3*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) - I) - 165*I*a^3*x^3*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(287*I*a^3*x^3 - 61*a^2*x^2 + 26*I*a*x + 8)*\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)))/x^3$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**4,x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{\left(\frac{1+axi}{\sqrt{a^2x^2+1}}\right)^{5/2}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)/x^4,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)/x^4, x)
```

$$3.87 \quad \int \frac{e^{\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=233

$$\frac{2467a^4\sqrt[4]{1+iax}}{192\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{4x^4\sqrt[4]{1-iax}} - \frac{17ia\sqrt[4]{1+iax}}{24x^3\sqrt[4]{1-iax}} + \frac{113a^2\sqrt[4]{1+iax}}{96x^2\sqrt[4]{1-iax}} + \frac{521ia^3\sqrt[4]{1+iax}}{192x\sqrt[4]{1-iax}} - \frac{475}{64}a^4 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] 2467/192*a^4*(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)-1/4*(1+I*a*x)^(1/4)/x^4/(1-I*a*x)^(1/4)-17/24*I*a*(1+I*a*x)^(1/4)/x^3/(1-I*a*x)^(1/4)+113/96*a^2*(1+I*a*x)^(1/4)/x^2/(1-I*a*x)^(1/4)+521/192*I*a^3*(1+I*a*x)^(1/4)/x/(1-I*a*x)^(1/4)-475/64*a^4*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-475/64*a^4*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))

Rubi [A]

time = 0.07, antiderivative size = 233, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 9, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {5170, 100, 156, 160, 12, 95, 218, 212, 209}

$$-\frac{475}{64}a^4 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{2467a^4\sqrt[4]{1+iax}}{192\sqrt[4]{1-iax}} - \frac{475}{64}a^4 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{521ia^3\sqrt[4]{1+iax}}{192x\sqrt[4]{1-iax}} + \frac{113a^2\sqrt[4]{1+iax}}{96x^2\sqrt[4]{1-iax}} - \frac{\sqrt[4]{1+iax}}{4x^4\sqrt[4]{1-iax}} - \frac{17ia\sqrt[4]{1+iax}}{24x^3\sqrt[4]{1-iax}}$$

Antiderivative was successfully verified.

[In] Int[E^(((5*I)/2)*ArcTan[a*x])/x^5,x]

[Out] (2467*a^4*(1 + I*a*x)^(1/4))/(192*(1 - I*a*x)^(1/4)) - (1 + I*a*x)^(1/4)/(4*x^4*(1 - I*a*x)^(1/4)) - (((17*I)/24)*a*(1 + I*a*x)^(1/4))/(x^3*(1 - I*a*x)^(1/4)) + (113*a^2*(1 + I*a*x)^(1/4))/(96*x^2*(1 - I*a*x)^(1/4)) + (((521*I)/192)*a^3*(1 + I*a*x)^(1/4))/(x*(1 - I*a*x)^(1/4)) - (475*a^4*ArcTan[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)])/64 - (475*a^4*ArcTanh[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)])/64

Rule 12

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 160

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m + n + p + 2, 0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || (! (NeQ[n, -1] && SumSimplerQ[n, 1]) && !(NeQ[p, -1] && SumSimplerQ[p, 1])))

```

Rule 209

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

```

Rule 218

```

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]

```

+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{\frac{5}{2}i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{(1 + iax)^{5/4}}{x^5(1 - iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{1}{4} \int \frac{-\frac{17ia}{2} + 8a^2x}{x^4(1 - iax)^{5/4}(1 + iax)^{3/4}} dx \\
 &= -\frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{1}{12} \int \frac{-\frac{113a^2}{4} - \frac{51}{2}ia^3x}{x^3(1 - iax)^{5/4}(1 + iax)^{3/4}} dx \\
 &= -\frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} - \frac{1}{24} \int \frac{\frac{521ia^3}{8} - \frac{113a^4x}{2}}{x^2(1 - iax)^{5/4}(1 + iax)^{3/4}} dx \\
 &= -\frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} + \frac{521ia^3\sqrt[4]{1 + iax}}{192x\sqrt[4]{1 - iax}} + \frac{1}{24} \int \frac{1}{x(1 - iax)^{5/4}(1 + iax)^{3/4}} dx \\
 &= \frac{2467a^4\sqrt[4]{1 + iax}}{192\sqrt[4]{1 - iax}} - \frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} + \frac{521ia^3\sqrt[4]{1 + iax}}{192x\sqrt[4]{1 - iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 + iax}}{192\sqrt[4]{1 - iax}} - \frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} + \frac{521ia^3\sqrt[4]{1 + iax}}{192x\sqrt[4]{1 - iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 + iax}}{192\sqrt[4]{1 - iax}} - \frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} + \frac{521ia^3\sqrt[4]{1 + iax}}{192x\sqrt[4]{1 - iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 + iax}}{192\sqrt[4]{1 - iax}} - \frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} + \frac{521ia^3\sqrt[4]{1 + iax}}{192x\sqrt[4]{1 - iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 + iax}}{192\sqrt[4]{1 - iax}} - \frac{\sqrt[4]{1 + iax}}{4x^4\sqrt[4]{1 - iax}} - \frac{17ia\sqrt[4]{1 + iax}}{24x^3\sqrt[4]{1 - iax}} + \frac{113a^2\sqrt[4]{1 + iax}}{96x^2\sqrt[4]{1 - iax}} + \frac{521ia^3\sqrt[4]{1 + iax}}{192x\sqrt[4]{1 - iax}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 118, normalized size = 0.51

$$\frac{-48 - 184iax + 362a^2x^2 + 747ia^3x^3 + 1946a^4x^4 + 2467ia^5x^5 + 950ia^4x^4(i + ax) {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}\right)}{192x^4\sqrt[4]{1 - iax} (1 + iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])/x^5,x]

[Out] (-48 - (184*I)*a*x + 362*a^2*x^2 + (747*I)*a^3*x^3 + 1946*a^4*x^4 + (2467*I)*a^5*x^5 + (950*I)*a^4*x^4*(I + a*x)*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)])/(192*x^4*(1 - I*a*x)^(1/4)*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)/x^5, x)

Fricas [A]

time = 1.93, size = 192, normalized size = 0.82

$$\frac{1425 a^4 x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) + 1425i a^4 x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) - 1425i a^4 x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) - 1425 a^4 x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) - 2(2467 a^4 x^4 + 521i a^3 x^3 + 226 a^2 x^2 - 136i a x - 48)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{384 x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x, algorithm="fricas")

[Out] -1/384*(1425*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 1425*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 1425*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 1425*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(2467*a^4*x^4 + 521*I*a^3*x^3 + 226*a^2*x^2 - 136*I*a*x - 48)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^4

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**5,x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{\left(\frac{1+ax}{\sqrt{a^2x^2+1}}\right)^{5/2}}{x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x+1)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x)
```

```
[Out] int(((a*x+1)/(a^2*x^2+1)^(1/2))^(5/2)/x^5, x)
```

3.88 $\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=337

$$-\frac{11\sqrt{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}(25-4iax)}{96a^4} - \frac{11\text{ArcTan}\left(\frac{1-\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4}$$

[Out] $-11/64*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^4+1/4*x^2*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^2-1/96*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}*(25-4*I*a*x)/a^4-11/128*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^4*2^{(1/2)}+11/128*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^4*2^{(1/2)}-11/256*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^4*2^{(1/2)}+11/256*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^4*2^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 337, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$,

Rules used = {5170, 102, 152, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$-\frac{11\text{ArcTan}\left(\frac{1-\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} + \frac{11\text{ArcTan}\left(\frac{1+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}(25-4iax)}{96a^4} - \frac{11\sqrt{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{11\log\left(\frac{\sqrt{1-iax}-\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}+1\right)}{128\sqrt{2}a^4} + \frac{11\log\left(\frac{\sqrt{1-iax}+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}+1\right)}{128\sqrt{2}a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{((I/2)*\text{ArcTan}[a*x])}, x]$

[Out] $(-11*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/(64*a^4) + (x^2*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)})/(4*a^2) - ((1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}*(25-(4*I)*a*x))/(96*a^4) - (11*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(64*\text{Sqrt}[2]*a^4) + (11*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(64*\text{Sqrt}[2]*a^4) - (11*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(128*\text{Sqrt}[2]*a^4) + (11*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(128*\text{Sqrt}[2]*a^4)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(IGtQ[m, 0] \&\& (!IntegerQ[n] || (GtQ[m, 0] \&\& LtQ[m - n, 0]))) \&\& !ILtQ[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 102

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_, x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a
+ b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b
*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*
(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p
}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_
))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
```

b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)^(n_)])*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int e^{-\frac{1}{2}i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3 \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx \\
&= \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} + \frac{\int \frac{x(-2+\frac{iax}{2})\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx}{4a^2} \\
&= \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}(25-4iax)}{96a^4} - \frac{(11i) \int \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx}{64a^3} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4} \\
&= -\frac{11\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{96a^4}
\end{aligned}$$

Mathematica [A]

time = 0.28, size = 272, normalized size = 0.81

$$\frac{1536 \frac{\sqrt[4]{1-iax} \operatorname{ArcTan}(ax)}{(1+iax)^{3/4}} - 3200 \frac{\sqrt[4]{1-iax} \operatorname{ArcTan}(ax)}{(1+iax)^{3/4}} + 2512 \frac{\sqrt[4]{1-iax} \operatorname{ArcTan}(ax)}{(1+iax)^{3/4}} - \frac{984 \sqrt[4]{1-iax} \operatorname{ArcTan}(ax)}{1+iax} + 33(-1)^{3/4} \log(e^{-2i \operatorname{ArcTan}(ax)}(\sqrt{-1} - e^{i \operatorname{ArcTan}(ax)})) + 33\sqrt{-1} \log(e^{-2i \operatorname{ArcTan}(ax)}((-1)^{3/4} - e^{i \operatorname{ArcTan}(ax)})) - 33(-1)^{3/4} \log(e^{-2i \operatorname{ArcTan}(ax)}(\sqrt{-1} + e^{i \operatorname{ArcTan}(ax)})) - 33\sqrt{-1} \log(e^{-2i \operatorname{ArcTan}(ax)}((-1)^{3/4} + e^{i \operatorname{ArcTan}(ax)})}{384a^4}$$

Antiderivative was successfully verified.

[In] Integrate[x^3/E^((I/2)*ArcTan[a*x]), x]

[Out] ((1536*E^(((15*I)/2)*ArcTan[a*x]))/(1 + E^((2*I)*ArcTan[a*x]))^4 - (3200*E^(((11*I)/2)*ArcTan[a*x]))/(1 + E^((2*I)*ArcTan[a*x]))^3 + (2512*E^(((7*I)/2

) * ArcTan[a*x]) / (1 + E^((2*I)*ArcTan[a*x]))^2 - (980 * E^(((3*I)/2) * ArcTan[a*x])) / (1 + E^((2*I)*ArcTan[a*x])) + 33 * (-1)^(3/4) * Log[(-1)^(1/4) - E^((I/2) * ArcTan[a*x])] / E^((2*I)*ArcTan[a*x]) + 33 * (-1)^(1/4) * Log[(-1)^(3/4) - E^((I/2) * ArcTan[a*x])] / E^((2*I)*ArcTan[a*x]) - 33 * (-1)^(3/4) * Log[(-1)^(1/4) + E^((I/2) * ArcTan[a*x])] / E^((2*I)*ArcTan[a*x]) - 33 * (-1)^(1/4) * Log[(-1)^(3/4) + E^((I/2) * ArcTan[a*x])] / E^((2*I)*ArcTan[a*x]) / (384 * a^4)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x^3}{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)

[Out] int(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="maxima")

[Out] integrate(x^3/sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)

Fricas [A]

time = 1.31, size = 255, normalized size = 0.76

$$\frac{96a^4\sqrt{\frac{121i}{4096a^8}}\log\left(\frac{3i+1}{4096a^8}\sqrt{\frac{121i}{4096a^8}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-96a^4\sqrt{\frac{121i}{4096a^8}}\log\left(\frac{3i+1}{4096a^8}\sqrt{\frac{121i}{4096a^8}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-96a^4\sqrt{\frac{121i}{4096a^8}}\log\left(\frac{3i+1}{4096a^8}\sqrt{\frac{121i}{4096a^8}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)+96a^4\sqrt{\frac{121i}{4096a^8}}\log\left(\frac{3i+1}{4096a^8}\sqrt{\frac{121i}{4096a^8}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-(-48i^2a^3+56a^2x^2+58iax-83)\sqrt{a^2x^2+1}\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{192a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="fricas")

[Out] -1/192*(96*a^4*sqrt(121/4096*I/a^8)*log(64/11*I*a^4*sqrt(121/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 96*a^4*sqrt(121/4096*I/a^8)*log(-64/11*I*a^4*sqrt(121/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 96*a^4*sqrt(-121/4096*I/a^8)*log(64/11*I*a^4*sqrt(-121/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 96*a^4*sqrt(-121/4096*I/a^8)*log(-64/11*I*a^4*sqrt(-121/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - (-48*I*a^3*x^3 + 56*a^2*x^2 + 58*I*a*x - 83)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/a^4

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^3}{\sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x**3/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2),x)``[Out] Integral(x**3/sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="giac")`

`[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
 dex_m & i,const vecteur & l) Error: Bad Argument Value`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3}{\sqrt{\frac{1+ax\text{li}}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^3/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)``[Out] int(x^3/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)`

3.89 $\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=339

$$\frac{3i\sqrt[4]{1-iax}(1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} + \frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3}$$

[Out] $\frac{3}{8}I*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^3 + \frac{1}{12}I*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^3 + \frac{1}{3}x*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^2 + \frac{3}{16}I*\text{arctan}(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3 + \frac{2^{(1/2)}-3}{16}I*\text{arctan}(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3 + \frac{2^{(1/2)}+3}{32}I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}) + \frac{2^{(1/2)}-3}{32}I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}) + \frac{2^{(1/2)}+3}{32}I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}) + \frac{2^{(1/2)}-3}{32}I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3$

Rubi [A]

time = 0.16, antiderivative size = 339, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$,

Rules used = {5170, 92, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{3i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{i(1+iax)^{3/4}(1-iax)^{5/4}}{12a^3} + \frac{3i(1+iax)^{3/4}\sqrt[4]{1-iax}}{8a^3} + \frac{3i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} - \frac{3i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} + \frac{x(1+iax)^{3/4}(1-iax)^{5/4}}{3a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{((I/2)*\text{ArcTan}[a*x])}, x]$

[Out] $\left(\frac{3I}{8}\right)*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^3 + \left(\frac{I}{12}\right)*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^3 + \frac{x*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}}{3*a^2} + \left(\frac{3I}{8}\right)*\text{ArcTan}\left[1 - \frac{\text{Sqrt}[2]*(1-I*a*x)^{(1/4)}}{(1+I*a*x)^{(1/4)}}\right]/(\text{Sqrt}[2]*a^3) - \left(\frac{3I}{8}\right)*\text{ArcTan}\left[1 + \frac{\text{Sqrt}[2]*(1-I*a*x)^{(1/4)}}{(1+I*a*x)^{(1/4)}}\right]/(\text{Sqrt}[2]*a^3) + \left(\frac{3I}{16}\right)*\text{Log}\left[1 + \frac{\text{Sqrt}[1-I*a*x]}{\text{Sqrt}[1+I*a*x]} - \frac{\text{Sqrt}[2]*(1-I*a*x)^{(1/4)}}{(1+I*a*x)^{(1/4)}}\right]/(\text{Sqrt}[2]*a^3) - \left(\frac{3I}{16}\right)*\text{Log}\left[1 + \frac{\text{Sqrt}[1-I*a*x]}{\text{Sqrt}[1+I*a*x]} + \frac{\text{Sqrt}[2]*(1-I*a*x)^{(1/4)}}{(1+I*a*x)^{(1/4)}}\right]/(\text{Sqrt}[2]*a^3)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_. + (d_.)*(x_))^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 92

```
Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(
d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)
^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b
*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ
[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 210

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_.) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_.) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{1}{2}i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2 \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx \\
&= \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} + \frac{\int \frac{(-1+\frac{iax}{2})\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx}{3a^2} \\
&= \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} - \frac{3 \int \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx}{8a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} \\
&= \frac{3i\sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} + \frac{i(1-iax)^{5/4}(1+iax)^{3/4}}{12a^3} + \frac{x(1-iax)^{5/4}(1+iax)^{3/4}}{3a^2} +
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 73, normalized size = 0.22

$$\frac{(1-iax)^{5/4} (5(1+iax)^{3/4}(i+4ax) - 9i2^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{5}{4}; \frac{9}{4}; \frac{1}{2}(1-iax)\right))}{60a^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^((I/2)*ArcTan[a*x]),x]

[Out] ((1 - I*a*x)^(5/4)*(5*(1 + I*a*x)^(3/4)*(I + 4*a*x) - (9*I)*2^(3/4)*Hypergeometric2F1[1/4, 5/4, 9/4, (1 - I*a*x)/2]))/(60*a^3)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)**[Out]** int(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="maxima")**[Out]** integrate(x^2/sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)**Fricas [A]**

time = 1.98, size = 247, normalized size = 0.73

$$\frac{12a^3\sqrt{\frac{9i}{64a^2}}\log\left(\frac{3}{2}a^3\sqrt{\frac{9i}{64a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 12a^3\sqrt{\frac{9i}{64a^2}}\log\left(-\frac{3}{2}a^3\sqrt{\frac{9i}{64a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 12a^3\sqrt{-\frac{9i}{64a^2}}\log\left(\frac{3}{2}a^3\sqrt{-\frac{9i}{64a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + 12a^3\sqrt{-\frac{9i}{64a^2}}\log\left(-\frac{3}{2}a^3\sqrt{-\frac{9i}{64a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \sqrt{a^2x^2+1}(-8i a^2x^2 + 10ax + 11i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{24a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="fricas")

[Out] $-1/24*(12*a^3*\sqrt{9/64*I/a^6}*\log(8/3*a^3*\sqrt{9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 12*a^3*\sqrt{9/64*I/a^6}*\log(-8/3*a^3*\sqrt{9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 12*a^3*\sqrt{-9/64*I/a^6}*\log(8/3*a^3*\sqrt{-9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) + 12*a^3*\sqrt{-9/64*I/a^6}*\log(-8/3*a^3*\sqrt{-9/64*I/a^6} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - \sqrt{a^2*x^2 + 1}*(-8*I*a^2*x^2 + 10*a*x + 11*I)*\sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)})/a^3$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**2/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2), x)
```

```
[Out] Integral(x**2/sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2), x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2}{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)
```

```
[Out] int(x^2/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)
```

3.90 $\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=295

$$\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2}$$

[Out] $1/4*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^2+1/2*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^2+1/8*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2*2^{(1/2)}-1/8*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^2*2^{(1/2)}+1/16*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^2*2^{(1/2)}-1/16*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^2*2^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 295, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \frac{(1+iax)^{3/4}(1-iax)^{5/4}}{2a^2} + \frac{(1+iax)^{3/4}\sqrt[4]{1-iax}}{4a^2} + \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2} - \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2}$$

Antiderivative was successfully verified.

[In] Int[x/E^((I/2)*ArcTan[a*x]), x]

[Out] $((1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/(4*a^2) + ((1 - I*a*x)^{(5/4)}*(1 + I*a*x)^{(3/4)})/(2*a^2) + \text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(4*\text{Sqrt}[2]*a^2) - \text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(4*\text{Sqrt}[2]*a^2) + \text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(8*\text{Sqrt}[2]*a^2) - \text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/(8*\text{Sqrt}[2]*a^2)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*(b*c - a*d)/(b*(m + n + 1)), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +

```
d*(x^p/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{1}{2}i \tan^{-1}(ax)} x dx &= \int \frac{x \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx \\
&= \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{i \int \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx}{4a} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{i \int \frac{1}{(1-iax)^{3/4} \sqrt[4]{1+iax}} dx}{8a} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{\text{Subst}\left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-iax}\right)}{2a^2} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{\text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2a^2} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4a^2} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{\text{Subst}\left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8a^2} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2} a^2} \\
&= \frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4a^2} + \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{\tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2} a^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 63, normalized size = 0.21

$$\frac{(1-iax)^{5/4} (5(1+iax)^{3/4} - 2^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{5}{4}; \frac{9}{4}; \frac{1}{2}(1-iax)\right))}{10a^2}$$

Antiderivative was successfully verified.

[In] Integrate[x/E^((I/2)*ArcTan[a*x]),x]

[Out] (((1 - I*a*x)^(5/4)*(5*(1 + I*a*x)^(3/4) - 2^(3/4)*Hypergeometric2F1[1/4, 5/4, 9/4, (1 - I*a*x)/2]))/(10*a^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x}{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)``[Out] int(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="maxima")``[Out] integrate(x/sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)`**Fricas [A]**

time = 1.76, size = 238, normalized size = 0.81

$$\frac{2a^2\sqrt{\frac{i}{16a^4}}\log\left(4ia^2\sqrt{\frac{i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)-2a^2\sqrt{\frac{i}{16a^4}}\log\left(-4ia^2\sqrt{\frac{i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)-2a^2\sqrt{-\frac{i}{16a^4}}\log\left(4ia^2\sqrt{-\frac{i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)+2a^2\sqrt{-\frac{i}{16a^4}}\log\left(-4ia^2\sqrt{-\frac{i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)+\sqrt{a^2x^2+1}(-2iax+3)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{4a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="fricas")`

```
[Out] 1/4*(2*a^2*sqrt(1/16*I/a^4)*log(4*I*a^2*sqrt(1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(1/16*I/a^4)*log(-4*I*a^2*sqrt(1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(-1/16*I/a^4)*log(4*I*a^2*sqrt(-1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*a^2*sqrt(-1/16*I/a^4)*log(-4*I*a^2*sqrt(-1/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + sqrt(a^2*x^2 + 1)*(-2*I*a*x + 3)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a^2
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x}{\sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2),x)
```

```
[Out] Integral(x/sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x}{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)
```

```
[Out] int(x/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)
```

3.91 $\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=268

$$\frac{i^4\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} - \frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{i\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{2\sqrt{2}a}$$

[Out] $-I*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a-1/2*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}+1/2*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}-1/4*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}+1/4*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 268, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5169, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$-\frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{i^4\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} - \frac{i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a} + \frac{i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(-1/2*I)*\text{ArcTan}[a*x]}, x]$

[Out] $((-I)*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/a - (I*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) + (I*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) - ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) + ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILTQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] :> \text{With}\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Den}$

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x],

$x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

Rule 5169

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)])*(n_.)}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}, x] /; \text{FreeQ}[\{a, n\}, x] \&\& \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
 \int e^{-\frac{1}{2}i \tan^{-1}(ax)} dx &= \int \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} + \frac{1}{2} \int \frac{1}{(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} + \frac{(2i)\text{Subst}\left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-iax}\right)}{a} \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} + \frac{(2i)\text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} + \frac{i\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} + \frac{i\text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a} \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} + \frac{i\text{Subst}\left(\int \frac{1}{1-\sqrt{2}xx^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2a} + \frac{i\text{Subst}\left(\int \frac{1}{1+\sqrt{2}xx^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2a} \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} - \frac{i \log\left(1 + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2\sqrt{2}a} + \frac{i \log\left(1 + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{2\sqrt{2}a} \\
 &= -\frac{i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} - \frac{i \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{i \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 41, normalized size = 0.15

$$\frac{8ie^{\frac{3}{2}i \text{ArcTan}(ax)} {}_2F_1\left(\frac{3}{4}, 2; \frac{7}{4}; -e^{2i \text{ArcTan}(ax)}\right)}{3a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((-1/2*I)*ArcTan[a*x]),x]

[Out] (((-8*I)/3)*E^(((3*I)/2)*ArcTan[a*x])*Hypergeometric2F1[3/4, 2, 7/4, -E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)

Fricas [A]

time = 2.14, size = 213, normalized size = 0.79

$$\frac{a\sqrt{\frac{i}{a^2}} \log\left(a\sqrt{\frac{i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - a\sqrt{\frac{i}{a^2}} \log\left(-a\sqrt{\frac{i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - a\sqrt{\frac{-i}{a^2}} \log\left(a\sqrt{\frac{-i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + a\sqrt{\frac{-i}{a^2}} \log\left(-a\sqrt{\frac{-i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - 2i\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="fricas")

[Out] 1/2*(a*sqrt(I/a^2)*log(a*sqrt(I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(I/a^2)*log(-a*sqrt(I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(-I/a^2)*log(a*sqrt(-I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + a*sqrt(-I/a^2)*log(-a*sqrt(-I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*I*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2),x)
```

```
[Out] Integral(1/sqrt((I*a*x + 1)/sqrt(a**2*x**2 + 1)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)
```

```
[Out] int(1/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)
```

3.92

$$\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=267

$$2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2}\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2}\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - 2\tanh^{-1}$$

[Out] 2*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-2*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-1/2*ln(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))*2^(1/2)+1/2*ln(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))*2^(1/2)-arctan(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))*2^(1/2)+arctan(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))*2^(1/2)

Rubi [A]

time = 0.13, antiderivative size = 267, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5170, 132, 65, 246, 217, 1179, 642, 1176, 631, 210, 95, 304, 209, 212}

$$2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2}\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2}\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} + \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - 2\tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(E^((I/2)*ArcTan[a*x]))*x], x]

[Out] 2*ArcTan[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)] - Sqrt[2]*ArcTan[1 - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)] + Sqrt[2]*ArcTan[1 + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)] - 2*ArcTanh[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)] - Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] - (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)]/Sqrt[2] + Log[1 + Sqrt[1 - I*a*x]/Sqrt[1 + I*a*x] + (Sqrt[2]*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4)]/Sqrt[2]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 95

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)]

], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n]
&& LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 132

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^(m - 1), x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 246

Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 304


```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
  2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{1}{2}i \tan^{-1}(ax)}}{x} dx &= \int \frac{\sqrt[4]{1-iax}}{x\sqrt[4]{1+iax}} dx \\
&= -\left((ia) \int \frac{1}{(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \right) + \int \frac{1}{x(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= 4\text{Subst}\left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-iax}\right) + 4\text{Subst}\left(\int \frac{x^2}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) \\
&= -\left(2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)\right) + 2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + 2\text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\text{Subst}\left(\int \frac{\sqrt{2}+2x}{-1-\sqrt{2}x-x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}} \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\log\left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}} \\
&= 2 \tan^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2} \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 96, normalized size = 0.36

$$2^{3/4} \sqrt[4]{1-iax} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; \frac{1}{2}(1-iax)\right) - \frac{4\sqrt[4]{1-iax} {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; -\frac{1-iax}{-1-iax}\right)}{\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a*x]))*x),x]

[Out] 2*2^(3/4)*(1 - I*a*x)^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, (1 - I*a*x)/2] - (4*(1 - I*a*x)^(1/4)*Hypergeometric2F1[1/4, 1, 5/4, -((1 - I*a*x)/(-1 - I*a*x))])/(1 + I*a*x)^(1/4)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x)`

[Out] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x, algorithm="maxima")`

[Out] `integrate(1/(x*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))), x)`

Fricas [A]

time = 1.27, size = 243, normalized size = 0.91

$$\frac{1}{2}\sqrt{4i}\log\left(\frac{1}{2}\sqrt{4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + \frac{1}{2}\sqrt{4i}\log\left(-\frac{1}{2}\sqrt{4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) + \frac{1}{2}\sqrt{-4i}\log\left(\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \frac{1}{2}\sqrt{-4i}\log\left(-\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right) - \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + i\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - i\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) + \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x, algorithm="fricas")`

[Out] `-1/2*sqrt(4*I)*log(1/2*I*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(4*I)*log(-1/2*I*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(-4*I)*log(1/2*I*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/2*sqrt(-4*I)*log(-1/2*I*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x \sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x,x)`

[Out] `Integral(1/(x*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)),x)
```

```
[Out] int(1/(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)), x)
```

3.93 $\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^2} dx$

Optimal. Leaf size=92

$$-\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} - ia\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x-I*a*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+I*a*\arctanh((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.03, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5170, 96, 95, 304, 209, 212}

$$-ia\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{x} + ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] `Int[1/(E^((I/2)*ArcTan[a*x])*x^2),x]`

[Out] $-(((1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/x) - I*a*\text{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] + I*a*\text{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}]$

Rule 95

```
Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 96

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1)/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
```

, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-\frac{1}{2}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{\sqrt[4]{1-iax}}{x^2 \sqrt[4]{1+iax}} dx \\
 &= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{x} - \frac{1}{2}(ia) \int \frac{1}{x(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
 &= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{x} - (2ia) \text{Subst} \left(\int \frac{x^2}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{x} + (ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - (ia) \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
 &= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{x} - ia \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + ia \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 69, normalized size = 0.75

$$\frac{i \sqrt[4]{1-iax} (i-ax + 2ax {}_2F_1(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}))}{x \sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a*x])*x^2),x]

[Out] (I*(1 - I*a*x)^(1/4)*(I - a*x + 2*a*x*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(x*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}} x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x, algorithm="maxima")

[Out] integrate(1/(x^2*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 156 vs. 2(64) = 128.

time = 4.97, size = 156, normalized size = 1.70

$$\frac{iax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - iax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x, algorithm="fricas")

[Out] 1/2*(I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - I - I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^2 \sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**2,x)
```

```
[Out] Integral(1/(x**2*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^2,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^2 \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)),x)
```

```
[Out] int(1/(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)), x)
```


3.94 $\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=132

$$\frac{ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} - \frac{1}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{1}{4}a^2\text{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] 1/4*I*a*(1-I*a*x)^(1/4)*(1+I*a*x)^(3/4)/x-1/2*(1-I*a*x)^(5/4)*(1+I*a*x)^(3/4)/x^2-1/4*a^2*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))+1/4*a^2*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))

Rubi [A]

time = 0.03, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5170, 98, 96, 95, 304, 209, 212}

$$-\frac{1}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{1}{4}a^2\text{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} + \frac{ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((I/2)*ArcTan[a*x])*x^3),x]

[Out] ((I/4)*a*(1-I*a*x)^(1/4)*(1+I*a*x)^(3/4))/x - ((1-I*a*x)^(5/4)*(1+I*a*x)^(3/4))/(2*x^2) - (a^2*ArcTan[(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)])/4 + (a^2*ArcTanh[(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)])/4

Rule 95

```
Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 96

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]
```

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{1}{2}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{\sqrt[4]{1-iax}}{x^3 \sqrt[4]{1+iax}} dx \\
&= -\frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} - \frac{1}{4}(ia) \int \frac{\sqrt[4]{1-iax}}{x^2 \sqrt[4]{1+iax}} dx \\
&= \frac{ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{4x} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} - \frac{1}{8}a^2 \int \frac{1}{x(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= \frac{ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{4x} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} - \frac{1}{2}a^2 \text{Subst} \left(\int \frac{x^2}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
&= \frac{ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{4x} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} + \frac{1}{4}a^2 \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
&= \frac{ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{4x} - \frac{(1-iax)^{5/4}(1+iax)^{3/4}}{2x^2} - \frac{1}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{1}{4}a^2 \ln \left| \frac{\sqrt[4]{1+iax} + \sqrt[4]{1-iax}}{\sqrt[4]{1+iax} - \sqrt[4]{1-iax}} \right|
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 81, normalized size = 0.61

$$\frac{\sqrt[4]{1-iax} (-2 + iax - 3a^2x^2 + 2a^2x^2 {}_2F_1(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}))}{4x^2 \sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a*x]))*x^3, x]

[Out] (((1 - I*a*x)^(1/4)*(-2 + I*a*x - 3*a^2*x^2 + 2*a^2*x^2*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)])))/(4*x^2*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}}} x^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3, x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x, algorithm="maxima")
```

```
[Out] integrate(1/(x^3*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))), x)
```

Fricas [A]

time = 3.47, size = 178, normalized size = 1.35

$$\frac{a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2\sqrt{a^2x^2+1}(-3iax+2)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{8x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x, algorithm="fricas")
```

```
[Out] 1/8*(a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*sqrt(a^2*x^2 + 1)*(-3*I*a*x + 2)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^2
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^3 \sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**3,x)
```

```
[Out] Integral(1/(x**3*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^3,x, algorithm="giac")
```

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
 the root of a polynomial with parameters. This might be wrong.The choice wa
 s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^3 \sqrt{\frac{1 + a x 1i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)),x)

[Out] int(1/(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)), x)

3.95 $\int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=170

$$-\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} + \frac{5ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2} + \frac{11a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} + \frac{3}{8}ia^3\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-1/3*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^3+5/12*I*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^2+11/24*a^2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x+3/8*I*a^3*\text{arctan}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-3/8*I*a^3*\text{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.05, antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 304, 209, 212}

$$\frac{3}{8}ia^3\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{3}{8}ia^3\text{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{11a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x} - \frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{3x^3} + \frac{5ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{12x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((I/2)*\text{ArcTan}[a*x])}*x^4), x]$

[Out] $-1/3*((1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/x^3 + (((5*I)/12)*a*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/x^2 + (11*a^2*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/(24*x) + ((3*I)/8)*a^3*\text{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] - ((3*I)/8)*a^3*\text{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}]$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]$

Rule 95

$\text{Int}[(((a_*) + (b_*)*(x_))^{(m_)*}((c_*) + (d_*)*(x_))^{(n_)})/((e_*) + (f_*)*(x_)), x_Symbol] \rightarrow \text{With}[q = \text{Denominator}[m], \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \text{FreeQ}[a, b, c, d, e, f], x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$

Rule 101

$\text{Int}[((a_*) + (b_*)*(x_))^{(m_)*}((c_*) + (d_*)*(x_))^{(n_)*}((e_*) + (f_*)*(x_))^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x] - \text{Dist}[1/((m+1)*(b*e - a*f)), \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x]$

```
(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m
+ n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1]
&& GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || Integ
ersQ[p, m + n])
```

Rule 156

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{1}{2}i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{\sqrt[4]{1-iax}}{x^4 \sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{1}{3} \int \frac{-\frac{5ia}{2} - 2a^2x}{x^3(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{5ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{12x^2} - \frac{1}{6} \int \frac{\frac{11a^2}{4} - \frac{5}{2}ia^3x}{x^2(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{5ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{12x^2} + \frac{11a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x} + \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{5ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{12x^2} + \frac{11a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x} + \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{5ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{12x^2} + \frac{11a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x} + \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{5ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{12x^2} + \frac{11a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x} - \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{3x^3} + \frac{5ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{12x^2} + \frac{11a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x} +
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 92, normalized size = 0.54

$$\frac{\sqrt[4]{1-iax} (-8 + 2iax + a^2x^2 + 11ia^3x^3 - 18ia^3x^3 {}_2F_1(\frac{1}{4}, 1; \frac{5}{4}, \frac{i+ax}{i-ax}))}{24x^3 \sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a*x])*x^4),x]

[Out] ((1 - I*a*x)^(1/4)*(-8 + (2*I)*a*x + a^2*x^2 + (11*I)*a^3*x^3 - (18*I)*a^3*x^3*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(24*x^3*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}} x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x)`

[Out] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x, algorithm="maxima")`

[Out] `integrate(1/(x^4*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))), x)`

Fricas [A]

time = 4.57, size = 187, normalized size = 1.10

$$\frac{-9ia^3x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) - 9a^3x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) + 9a^3x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) + 9ia^3x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) + 2(11a^2x^2 + 10iax - 8)\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{48x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x, algorithm="fricas")`

[Out] `1/48*(-9*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 9*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 9*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + 9*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) + 2*(11*a^2*x^2 + 10*I*a*x - 8)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^3`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^4 \sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**4,x)`

[Out] `Integral(1/(x**4*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^4,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^4 \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^4*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)),x)
```

```
[Out] int(1/(x^4*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)), x)
```

$$3.96 \quad \int \frac{e^{-\frac{1}{2}i\text{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=202

$$-\frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} + \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x^3} + \frac{29a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{96x^2} - \frac{83ia^3\sqrt[4]{1-iax}(1+iax)^{3/4}}{192x}$$

[Out] $-1/4*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^4+7/24*I*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^3+29/96*a^2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x^2-83/192*I*a^3*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/x+11/64*a^4*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-11/64*a^4*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.06, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 304, 209, 212}

$$\frac{11}{64}a^4\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{11}{64}a^4\operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{83ia^3\sqrt[4]{1-iax}(1+iax)^{3/4}}{192x} + \frac{29a^2\sqrt[4]{1-iax}(1+iax)^{3/4}}{96x^2} - \frac{\sqrt[4]{1-iax}(1+iax)^{3/4}}{4x^4} + \frac{7ia\sqrt[4]{1-iax}(1+iax)^{3/4}}{24x^3}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((I/2)*ArcTan[a*x])*x^5),x]

[Out] $-1/4*((1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/x^4 + (((7*I)/24)*a*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/x^3 + (29*a^2*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/(96*x^2) - (((83*I)/192)*a^3*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/x + (11*a^4*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64 - (11*a^4*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 101

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m+1)*(c + d*x)^n*(e + f*x)^(p+1)

)/((m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 156

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 304

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)^(n_.)]*(x_)^(m_.)), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{1}{2}i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{\sqrt[4]{1-iax}}{x^5 \sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{1}{4} \int \frac{-\frac{7ia}{2} - 3a^2x}{x^4(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} - \frac{1}{12} \int \frac{\frac{29a^2}{4} - 7ia^3x}{x^3(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots \\
&= -\frac{\sqrt[4]{1-iax} (1+iax)^{3/4}}{4x^4} + \frac{7ia \sqrt[4]{1-iax} (1+iax)^{3/4}}{24x^3} + \frac{29a^2 \sqrt[4]{1-iax} (1+iax)^{3/4}}{96x^2} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 99, normalized size = 0.49

$$\frac{\sqrt[4]{1-iax} (-48 + 8iax + 2a^2x^2 - 25ia^3x^3 + 83a^4x^4 - 66a^4x^4 {}_2F_1(\frac{1}{4}, 1; \frac{5}{4}, \frac{i+ax}{i-ax}))}{192x^4 \sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a*x]))*x^5),x]

[Out] ((1 - I*a*x)^(1/4)*(-48 + (8*I)*a*x + 2*a^2*x^2 - (25*I)*a^3*x^3 + 83*a^4*x^4 - 66*a^4*x^4*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(192*x^4*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{iax+1}{\sqrt{a^2x^2+1}}} x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x)`

[Out] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x, algorithm="maxima")`

[Out] `integrate(1/(x^5*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1))), x)`

Fricas [A]

time = 2.39, size = 195, normalized size = 0.97

$$\frac{33a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - 33ia^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + 33ia^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 33a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) + 2(83ia^3x^3 - 58a^2x^2 - 56iax + 48)\sqrt{a^2x^2+1} \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{384x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x, algorithm="fricas")`

[Out] `-1/384*(33*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 33*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 33*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 33*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) + 2*(83*I*a^3*x^3 - 58*a^2*x^2 - 56*I*a*x + 48)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^4`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^5 \sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)/x**5,x)`

[Out] `Integral(1/(x**5*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1))), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)/x^5,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x^5 \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^5*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)),x)
```

```
[Out] int(1/(x^5*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2)), x)
```

3.97 $\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=337

$$-\frac{41(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4}\sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4}\sqrt[4]{1+iax}(11-4iax)}{32a^4} - \frac{123\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4}$$

[Out] $-41/64*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^4+1/4*x^2*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)}/a^2-1/32*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)}*(11-4*I*a*x)/a^4-123/128*\text{arctan}\left(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}\right)/a^4-123/128*\text{arctan}\left(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}\right)/a^4+123/256*\ln\left(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)}\right)/a^4+123/256*\ln\left(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)}\right)/a^4$

Rubi [A]

time = 0.16, antiderivative size = 337, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$,

Rules used = {5170, 102, 152, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$-\frac{123\text{ArcTan}\left(1-\frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} + \frac{123\text{ArcTan}\left(1+\frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} - \frac{(1-iax)^{7/4}\sqrt[4]{1+iax}(11-4iax)}{32a^4} - \frac{41(1-iax)^{3/4}\sqrt[4]{1+iax}}{64a^4} + \frac{123\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{128\sqrt{2}a^4} - \frac{123\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}} + 1\right)}{128\sqrt{2}a^4} + \frac{x^2(1-iax)^{7/4}\sqrt[4]{1+iax}}{4a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{((3*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $(-41*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(64*a^4) + (x^2*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)})/(4*a^2) - ((1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)}*(11-(4*I)*a*x))/(32*a^4) - (123*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(64*\text{Sqrt}[2]*a^4) + (123*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(64*\text{Sqrt}[2]*a^4) + (123*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(128*\text{Sqrt}[2]*a^4) - (123*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)})]/(128*\text{Sqrt}[2]*a^4)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$
 $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m + n + 1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0]))) \ \&\& \ !\text{ILTQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 102

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_, x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x
)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a
+ b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b
*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*
(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p
}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_
))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
```

$^{(1/n)}$, x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
 $^{(-1)}$] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
 Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
 /(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
 & EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
 eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)^(n_)])*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a
 x)^(I(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
 rQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int e^{-\frac{3}{2}i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1-iax)^{3/4}}{(1+iax)^{3/4}} dx \\
&= \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} + \frac{\int \frac{x(1-iax)^{3/4}(-2+\frac{3iax}{2})}{(1+iax)^{3/4}} dx}{4a^2} \\
&= \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} - \frac{(41i) \int \frac{(1-iax)^{3/4}}{(1+iax)^{3/4}} dx}{64a^3} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4} \\
&= -\frac{41(1-iax)^{3/4} \sqrt[4]{1+iax}}{64a^4} + \frac{x^2(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^2} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax} (11-4iax)}{32a^4}
\end{aligned}$$

Mathematica [A]

time = 0.28, size = 272, normalized size = 0.81

$$\frac{\frac{512 \sqrt[4]{1+iax} \operatorname{ArcTan}(ax)}{(1+iax \operatorname{ArcTan}(ax))^2} - \frac{1152 \sqrt[4]{1+iax} \operatorname{ArcTan}(ax)}{(1+iax \operatorname{ArcTan}(ax))^2} + \frac{1008 \sqrt[4]{1+iax} \operatorname{ArcTan}(ax)}{(1+iax \operatorname{ArcTan}(ax))^2} - \frac{532 \sqrt[4]{1+iax} \operatorname{ArcTan}(ax)}{1+iax \operatorname{ArcTan}(ax)} + 123 \sqrt[4]{-1} \log(e^{-2i \operatorname{ArcTan}(ax)} (\sqrt{-1} - e^{\frac{1}{2}i \operatorname{ArcTan}(ax)})) + 123(-1)^{3/4} \log(e^{-2i \operatorname{ArcTan}(ax)} ((-1)^{3/4} - e^{\frac{1}{2}i \operatorname{ArcTan}(ax)})) - 123 \sqrt[4]{-1} \log(e^{-2i \operatorname{ArcTan}(ax)} (\sqrt{-1} + e^{\frac{1}{2}i \operatorname{ArcTan}(ax)})) - 123(-1)^{3/4} \log(e^{-2i \operatorname{ArcTan}(ax)} ((-1)^{3/4} + e^{\frac{1}{2}i \operatorname{ArcTan}(ax)}))}{128a^4}$$

Antiderivative was successfully verified.

[In] Integrate[x^3/E^(((3*I)/2)*ArcTan[a*x]), x]

[Out] $\frac{((512 * E^{(((13 * I) / 2) * \operatorname{ArcTan}[a * x])}) / (1 + E^{((2 * I) * \operatorname{ArcTan}[a * x])}))^4 - (1152 * E^{((9 * I) / 2) * \operatorname{ArcTan}[a * x]}) / (1 + E^{((2 * I) * \operatorname{ArcTan}[a * x])})^3 + (1008 * E^{((5 * I) / 2) * \operatorname{ArcTan}[a * x]}) / (1 + E^{((2 * I) * \operatorname{ArcTan}[a * x])})^2 - (532 * E^{((I) / 2) * \operatorname{ArcTan}[a * x]}) / (1 + E^{((2 * I) * \operatorname{ArcTan}[a * x])}) + 123 * (-1)^{(1/4)} * \operatorname{Log}[((-1)^{(1/4)} - E^{((I) / 2) * \operatorname{ArcTan}[a * x]})]$

$\frac{\operatorname{an}[a*x])}{E^{((2*I)*\operatorname{ArcTan}[a*x])}} + 123*(-1)^{(3/4)}*\operatorname{Log}[((-1)^{(3/4)} - E^{((I/2)*\operatorname{ArcTan}[a*x])})/E^{((2*I)*\operatorname{ArcTan}[a*x])}] - 123*(-1)^{(1/4)}*\operatorname{Log}[((-1)^{(1/4)} + E^{((I/2)*\operatorname{ArcTan}[a*x])})/E^{((2*I)*\operatorname{ArcTan}[a*x])}] - 123*(-1)^{(3/4)}*\operatorname{Log}[((-1)^{(3/4)} + E^{((I/2)*\operatorname{ArcTan}[a*x])})/E^{((2*I)*\operatorname{ArcTan}[a*x])}]]/(128*a^4)$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x^3}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

[Out] int(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="maxima")

[Out] integrate(x^3/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [A]

time = 2.13, size = 251, normalized size = 0.74

$$\frac{32 a^4 \sqrt{\frac{15129}{4096 a^2}} \log\left(\frac{\frac{64}{123} a^4 \sqrt{\frac{15129}{4096 a^2}} + \sqrt{\frac{15129}{4096 a^2}}}{ax+1}\right) - 32 a^4 \sqrt{\frac{15129}{4096 a^2}} \log\left(-\frac{\frac{64}{123} a^4 \sqrt{\frac{15129}{4096 a^2}} + \sqrt{\frac{15129}{4096 a^2}}}{ax+1}\right) + 32 a^4 \sqrt{\frac{15129}{4096 a^2}} \log\left(\frac{\frac{64}{123} a^4 \sqrt{\frac{15129}{4096 a^2}} + \sqrt{\frac{15129}{4096 a^2}}}{ax+1}\right) - 32 a^4 \sqrt{\frac{15129}{4096 a^2}} \log\left(-\frac{\frac{64}{123} a^4 \sqrt{\frac{15129}{4096 a^2}} + \sqrt{\frac{15129}{4096 a^2}}}{ax+1}\right) + (16 a^4 x^4 + 40 a^3 x^3 - 54 a^2 x^2 - 93 a x + 63) \sqrt{\frac{15129}{4096 a^2}}}{64 a^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="fricas")

[Out] -1/64*(32*a^4*sqrt(15129/4096*I/a^8)*log(64/123*a^4*sqrt(15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 32*a^4*sqrt(15129/4096*I/a^8)*log(-64/123*a^4*sqrt(15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 32*a^4*sqrt(-15129/4096*I/a^8)*log(64/123*a^4*sqrt(-15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 32*a^4*sqrt(-15129/4096*I/a^8)*log(-64/123*a^4*sqrt(-15129/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + (16*a^4*x^4 + 40*I*a^3*x^3 - 54*a^2*x^2 - 93*I*a*x + 63)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/a^4

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^3}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2),x)

[Out] Integral(x**3/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3}{\left(\frac{1+ax \text{ li}}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)

[Out] int(x^3/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)

3.98 $\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=339

$$\frac{17i(1-iax)^{3/4}\sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4}\sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4}\sqrt[4]{1+iax}}{3a^2} + \frac{17i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3}$$

[Out] $17/24*I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^3+1/4*I*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)}/a^3+1/3*x*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)}/a^2+17/16*I*arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3+2^{(1/2)}-17/16*I*arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3+2^{(1/2)}-17/32*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3+2^{(1/2)}+17/32*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3+2^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 339, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5170, 92, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{17i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{17i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{i\sqrt[4]{1+iax}(1-iax)^{7/4}}{4a^3} + \frac{17i\sqrt[4]{1+iax}(1-iax)^{3/4}}{24a^3} - \frac{17i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} + \frac{17i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{16\sqrt{2}a^3} + \frac{x\sqrt[4]{1+iax}(1-iax)^{7/4}}{3a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{((3*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $((17*I)/24)*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a^3 + ((I/4)*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)})/a^3 + (x*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)})/(3*a^2) + (((17*I)/8)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3) - (((17*I)/8)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3) - (((17*I)/16)*\text{Log}[1 + \text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3) + (((17*I)/16)*\text{Log}[1 + \text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a^3)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_. + (d_.)*(x_))^{(n_)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(IGtQ[m, 0] \&\& (!IntegerQ[n] || (GtQ[m, 0] \&\& LtQ[m - n, 0]))) \&\& !ILtQ[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 92

```
Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(
d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)
^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b
*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ
[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 210

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{3}{2}i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1-iax)^{3/4}}{(1+iax)^{3/4}} dx \\
&= \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} + \frac{\int \frac{(1-iax)^{3/4}(-1+\frac{3iax}{2})}{(1+iax)^{3/4}} dx}{3a^2} \\
&= \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{17 \int \frac{(1-iax)^{3/4}}{(1+iax)^{3/4}} dx}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} + \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} - \frac{1}{24a^2} \\
&= \frac{17i(1-iax)^{3/4} \sqrt[4]{1+iax}}{24a^3} + \frac{i(1-iax)^{7/4} \sqrt[4]{1+iax}}{4a^3} + \frac{x(1-iax)^{7/4} \sqrt[4]{1+iax}}{3a^2} + \frac{1}{24a^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 73, normalized size = 0.22

$$\frac{(1-iax)^{7/4} \left(7\sqrt[4]{1+iax} (3i+4ax) - 17i\sqrt[4]{2} {}_2F_1\left(\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; \frac{1}{2}(1-iax)\right) \right)}{84a^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^(((3*I)/2)*ArcTan[a*x]),x]

[Out] ((1 - I*a*x)^(7/4)*(7*(1 + I*a*x)^(1/4)*(3*I + 4*a*x) - (17*I)*2^(1/4)*Hypergeometric2F1[3/4, 7/4, 11/4, (1 - I*a*x)/2]))/(84*a^3)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)``[Out] int(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="maxima")``[Out] integrate(x^2/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)`**Fricas [A]**

time = 2.50, size = 243, normalized size = 0.72

$$\frac{12a^3\sqrt{\frac{289}{64a^6}}\log\left(\frac{\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}\right)-12a^3\sqrt{\frac{289}{64a^6}}\log\left(\frac{-\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}\right)+12a^3\sqrt{\frac{289}{64a^6}}\log\left(\frac{\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}\right)-12a^3\sqrt{\frac{289}{64a^6}}\log\left(\frac{-\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{\frac{289}{64a^6}\sqrt{\frac{289}{64a^6}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}\right)+(8a^3x^3+22Ia^2x^2-37ax-23I)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{24a^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="fricas")`

```
[Out] -1/24*(12*a^3*sqrt(289/64*I/a^6)*log(8/17*I*a^3*sqrt(289/64*I/a^6) + sqrt(I
*sqrt(a^2*x^2 + 1)/(a*x + I))) - 12*a^3*sqrt(289/64*I/a^6)*log(-8/17*I*a^3*
sqrt(289/64*I/a^6) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 12*a^3*sqrt(-28
9/64*I/a^6)*log(8/17*I*a^3*sqrt(-289/64*I/a^6) + sqrt(I*sqrt(a^2*x^2 + 1)/(
a*x + I))) - 12*a^3*sqrt(-289/64*I/a^6)*log(-8/17*I*a^3*sqrt(-289/64*I/a^6)
+ sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + (8*a^3*x^3 + 22*I*a^2*x^2 - 37*a*
x - 23*I)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/a^3
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2),x)

[Out] Integral(x**2/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2}{\left(\frac{1+ax}{\sqrt{a^2x^2+1}}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/((a*x+1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)

[Out] int(x^2/((a*x+1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)

3.99 $\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=295

$$\frac{3(1-iax)^{3/4}\sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4}\sqrt[4]{1+iax}}{2a^2} + \frac{9\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{9\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2}$$

[Out] $\frac{3}{4}(1-Iax)^{3/4}(1+Iax)^{1/4}/a^2 + \frac{1}{2}(1-Iax)^{7/4}(1+Iax)^{1/4}/a^2 + \frac{9}{4\sqrt{2}}\frac{\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a^2} - \frac{9}{4\sqrt{2}}\frac{\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{a^2}$

Rubi [A]

time = 0.13, antiderivative size = 295, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{9\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{9\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \frac{\sqrt[4]{1+iax}(1-iax)^{7/4}}{2a^2} + \frac{3\sqrt[4]{1+iax}(1-iax)^{3/4}}{4a^2} - \frac{9\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2} + \frac{9\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x/E^{((3*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $(3*(1 - Iax)^{3/4}(1 + Iax)^{1/4})/(4a^2) + ((1 - Iax)^{7/4}(1 + Iax)^{1/4})/(2a^2) + (9*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - Iax)^{1/4})/(1 + Iax)^{1/4}])/(4*\text{Sqrt}[2]*a^2) - (9*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - Iax)^{1/4})/(1 + Iax)^{1/4}])/(4*\text{Sqrt}[2]*a^2) - (9*\text{Log}[1 + \text{Sqrt}[1 - Iax]/\text{Sqrt}[1 + Iax] - (\text{Sqrt}[2]*(1 - Iax)^{1/4})/(1 + Iax)^{1/4}])/(8*\text{Sqrt}[2]*a^2) + (9*\text{Log}[1 + \text{Sqrt}[1 - Iax]/\text{Sqrt}[1 + Iax] + (\text{Sqrt}[2]*(1 - Iax)^{1/4})/(1 + Iax)^{1/4}])/(8*\text{Sqrt}[2]*a^2)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_.)*((c_. + (d_.)*(x_))^{(n_.)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^n/(b*(m + n + 1))}, x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))], \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_))^{(m_.)*((c_. + (d_.)*(x_))^{(n_.)}, x_Symbol] :> \text{With}\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) +$

```
d*(x^p/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_.), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{3}{2}i \tan^{-1}(ax)} x dx &= \int \frac{x(1-iax)^{3/4}}{(1+iax)^{3/4}} dx \\
&= \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} + \frac{(3i) \int \frac{(1-iax)^{3/4}}{(1+iax)^{3/4}} dx}{4a} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} + \frac{(9i) \int \frac{1}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx}{8a} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} - \frac{9 \text{Subst} \left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax} \right)}{2a^2} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} - \frac{9 \text{Subst} \left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{2a^2} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} + \frac{9 \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{4a^2} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} - \frac{9 \text{Subst} \left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{8a^2} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} - \frac{9 \log \left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{8\sqrt{2} a^2} \\
&= \frac{3(1-iax)^{3/4} \sqrt[4]{1+iax}}{4a^2} + \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2a^2} + \frac{9 \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{4\sqrt{2} a^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 63, normalized size = 0.21

$$\frac{(1-iax)^{7/4} \left(7\sqrt[4]{1+iax} - 3\sqrt[4]{2} {}_2F_1\left(\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; \frac{1}{2}(1-iax)\right) \right)}{14a^2}$$

Antiderivative was successfully verified.

[In] Integrate[x/E^(((3*I)/2)*ArcTan[a*x]),x]

[Out] (((1 - I*a*x)^(7/4)*(7*(1 + I*a*x)^(1/4) - 3*2^(1/4)*Hypergeometric2F1[3/4, 7/4, 11/4, (1 - I*a*x)/2]))/(14*a^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

[Out] int(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="maxima")

[Out] integrate(x/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [A]

time = 4.54, size = 236, normalized size = 0.80

$$\frac{2a^2\sqrt{\frac{81i}{16a^4}}\log\left(\frac{\frac{1}{2}a^2\sqrt{\frac{81i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{\frac{1}{2}a^2\sqrt{\frac{81i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-2a^2\sqrt{\frac{81i}{16a^4}}\log\left(-\frac{1}{2}a^2\sqrt{\frac{81i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)+2a^2\sqrt{\frac{81i}{16a^4}}\log\left(\frac{\frac{1}{2}a^2\sqrt{-\frac{81i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{\frac{1}{2}a^2\sqrt{-\frac{81i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right)-2a^2\sqrt{\frac{81i}{16a^4}}\log\left(-\frac{1}{2}a^2\sqrt{-\frac{81i}{16a^4}}+\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}\right)-(2a^2x^2+7iax-5)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{4a^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="fricas")

[Out] 1/4*(2*a^2*sqrt(81/16*I/a^4)*log(4/9*a^2*sqrt(81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(81/16*I/a^4)*log(-4/9*a^2*sqrt(81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*a^2*sqrt(-81/16*I/a^4)*log(4/9*a^2*sqrt(-81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*a^2*sqrt(-81/16*I/a^4)*log(-4/9*a^2*sqrt(-81/16*I/a^4) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - (2*a^2*x^2 + 7*I*a*x - 5)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2),x)

[Out] Integral(x/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x}{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)

[Out] int(x/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)

3.100 $\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=268

$$\frac{i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} - \frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{3i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{3i\log\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a}$$

[Out] $-I*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/a-3/2*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}+3/2*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}+3/4*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}-3/4*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}$

Rubi [A]

time = 0.11, antiderivative size = 268, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5169, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$-\frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{3i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{i(1-iax)^{3/4}\sqrt[4]{1+iax}}{a} + \frac{3i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a} - \frac{3i\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((-3*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $((-I)*(1 - I*a*x)^{(3/4)}*(1 + I*a*x)^{(1/4)})/a - ((3*I)*\text{ArcTan}[1 - (\text{Sqrt}[2]* (1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a) + ((3*I)*\text{ArcTan}[1 + (\text{Sqrt}[2]* (1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a) + (((3*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a) - (((3*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]) / (\text{Sqrt}[2]*a)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))], x] + \text{Dist}[n*(b*c - a*d)/(b*(m + n + 1)), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILTQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) + d*(x^p/b))^{(n)}, x], x, (a + b*x)^{(1/p)}], x]] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 303

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 338

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x], x]

$x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}[\{a, c, d, e\}, x] \ \&\& \ \text{EqQ}[c*d^2 - a*e^2, 0] \ \&\& \ \text{NegQ}[d*e]$

Rule 5169

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))}, x_Symbol] \ :> \ \text{Int}[(1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}, x] /; \text{FreeQ}[\{a, n\}, x] \ \&\& \ !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
 \int e^{-\frac{3}{2}i \tan^{-1}(ax)} dx &= \int \frac{(1 - iax)^{3/4}}{(1 + iax)^{3/4}} dx \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} + \frac{3}{2} \int \frac{1}{\sqrt[4]{1 - iax} (1 + iax)^{3/4}} dx \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} + \frac{(6i) \text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1 - iax}\right)}{a} \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} + \frac{(6i) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{a} \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} - \frac{(3i) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{a} + \frac{(3i) \text{Subst}\left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{a} \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} + \frac{(3i) \text{Subst}\left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{2a} + \frac{(3i) \text{Subst}\left(\int \frac{1}{1+\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{2a} \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} + \frac{3i \log\left(1 + \frac{\sqrt{1 - iax}}{\sqrt{1 + iax}} - \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{2\sqrt{2} a} - \frac{3i \log\left(1 + \frac{\sqrt{1 + iax}}{\sqrt{1 - iax}} - \frac{\sqrt{2} \sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}}\right)}{2\sqrt{2} a} \\
 &= -\frac{i(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{a} - \frac{3i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{\sqrt{2} a} + \frac{3i \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}}\right)}{\sqrt{2} a}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 39, normalized size = 0.15

$$\frac{8ie^{\frac{1}{2}i \text{ArcTan}(ax)} {}_2F_1\left(\frac{1}{4}, 2; \frac{5}{4}; -e^{2i \text{ArcTan}(ax)}\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((−3*I)/2)*ArcTan[a*x]),x]

[Out] ((−8*I)*E^((I/2)*ArcTan[a*x])*Hypergeometric2F1[1/4, 2, 5/4, −E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(-3/2), x)

Fricas [A]

time = 2.82, size = 209, normalized size = 0.78

$$\frac{a\sqrt{\frac{9i}{a^2}} \log\left(\frac{\frac{1}{3}ia\sqrt{\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{\frac{1}{3}ia\sqrt{\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) - a\sqrt{\frac{9i}{a^2}} \log\left(\frac{-\frac{1}{3}ia\sqrt{\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{-\frac{1}{3}ia\sqrt{\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) + a\sqrt{-\frac{9i}{a^2}} \log\left(\frac{\frac{1}{3}ia\sqrt{-\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{\frac{1}{3}ia\sqrt{-\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) - a\sqrt{-\frac{9i}{a^2}} \log\left(\frac{-\frac{1}{3}ia\sqrt{-\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{-\frac{1}{3}ia\sqrt{-\frac{9i}{a^2}} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) - 2(ax+i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="fricas")

[Out] 1/2*(a*sqrt(9*I/a^2)*log(1/3*I*a*sqrt(9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(9*I/a^2)*log(-1/3*I*a*sqrt(9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + a*sqrt(-9*I/a^2)*log(1/3*I*a*sqrt(-9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - a*sqrt(-9*I/a^2)*log(-1/3*I*a*sqrt(-9*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 2*(a*x + I)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/a

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2),x)
```

```
[Out] Integral(((I*a*x + 1)/sqrt(a**2*x**2 + 1))**(-3/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(1/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)
```

3.101 $\int \frac{e^{-\frac{3}{2}i \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=267

$$-2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - 2 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-2 \operatorname{arctan}\left(\frac{(1+I*a*x)^{1/4}}{(1-I*a*x)^{1/4}}\right) - 2 \operatorname{arctanh}\left(\frac{(1+I*a*x)^{1/4}}{(1-I*a*x)^{1/4}}\right) + \frac{1}{2} \ln\left(\frac{1 - (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} + (1-I*a*x)^{1/2}}\right) + \frac{1}{2} \ln\left(\frac{1 + (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} + (1-I*a*x)^{1/2}}\right) - \operatorname{arctan}\left(\frac{1 - (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} * 2^{1/2}}\right) + \operatorname{arctan}\left(\frac{1 + (1-I*a*x)^{1/4} * 2^{1/2}}{(1+I*a*x)^{1/4} * 2^{1/2}}\right)$

Rubi [A]

time = 0.13, antiderivative size = 267, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5170, 132, 65, 338, 303, 1176, 631, 210, 1179, 642, 95, 218, 212, 209}

$$-2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - \frac{\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - 2 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] `Int[1/(E^((3*I)/2)*ArcTan[a*x])*x], x]`

[Out] $-2 \operatorname{ArcTan}\left[\frac{1 + I*a*x}{(1 - I*a*x)^{1/4}}\right] - \operatorname{Sqrt}[2] \operatorname{ArcTan}\left[1 - \frac{\operatorname{Sqrt}[2] * (1 - I*a*x)^{1/4}}{1 + I*a*x}\right] + \operatorname{Sqrt}[2] \operatorname{ArcTan}\left[1 + \frac{\operatorname{Sqrt}[2] * (1 - I*a*x)^{1/4}}{1 + I*a*x}\right] - 2 \operatorname{ArcTanh}\left[\frac{1 + I*a*x}{(1 - I*a*x)^{1/4}}\right] + \operatorname{Log}\left[\frac{1 + \operatorname{Sqrt}[1 - I*a*x]}{\operatorname{Sqrt}[1 + I*a*x]}\right] - \frac{\operatorname{Sqrt}[2] * (1 - I*a*x)^{1/4}}{(1 + I*a*x)^{1/4}} / \operatorname{Sqrt}[2] - \operatorname{Log}\left[\frac{1 + \operatorname{Sqrt}[1 - I*a*x]}{\operatorname{Sqrt}[1 + I*a*x]}\right] + \frac{\operatorname{Sqrt}[2] * (1 - I*a*x)^{1/4}}{(1 + I*a*x)^{1/4}} / \operatorname{Sqrt}[2]$

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 95

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)]`

```
], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n]
&& LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 132

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^(m - 1), x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338


```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{3}{2}i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1-iax)^{3/4}}{x(1+iax)^{3/4}} dx \\
&= -\left((ia) \int \frac{1}{\sqrt[4]{1-iax} (1+iax)^{3/4}} dx \right) + \int \frac{1}{x \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= 4\text{Subst} \left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-iax} \right) + 4\text{Subst} \left(\int \frac{1}{-1+x^4} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \\
&= -\left(2\text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) \right) - 2\text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \\
&= -2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - 2\text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right) \\
&= -2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{\text{Subst} \left(\int \frac{\sqrt{2}+2x}{-1-\sqrt{2}x-x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{\sqrt{2}} \\
&= -2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{\log \left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{\sqrt{2}} \\
&= -2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) - \sqrt{2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right) + \sqrt{2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 97, normalized size = 0.36

$$\frac{2(1-iax)^{3/4} \left(\sqrt[4]{2} (1+iax)^{3/4} {}_2F_1 \left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; \frac{1}{2}(1-iax) \right) - 2 {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax} \right) \right)}{3(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a*x]))*x], x]

[Out] (2*(1 - I*a*x)^(3/4)*(2^(1/4)*(1 + I*a*x)^(3/4)*Hypergeometric2F1[3/4, 3/4, 7/4, (1 - I*a*x)/2] - 2*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(3*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}} \right)^{\frac{3}{2}} x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x)`

[Out] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x, algorithm="maxima")`

[Out] `integrate(1/(x*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)), x)`

Fricas [A]

time = 3.23, size = 243, normalized size = 0.91

$$-\frac{1}{2}\sqrt{4i}\log\left(\frac{1}{2}\sqrt{4i} + \sqrt{\frac{\sqrt{4i^2+1}}{4i+1}}\right) + \frac{1}{2}\sqrt{4i}\log\left(-\frac{1}{2}\sqrt{4i} + \sqrt{\frac{\sqrt{4i^2+1}}{4i+1}}\right) - \frac{1}{2}\sqrt{-4i}\log\left(\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{\sqrt{4i^2+1}}{4i+1}}\right) + \frac{1}{2}\sqrt{-4i}\log\left(-\frac{1}{2}\sqrt{-4i} + \sqrt{\frac{\sqrt{4i^2+1}}{4i+1}}\right) - \log\left(\sqrt{\frac{\sqrt{4i^2+1}}{4i+1}} + 1\right) - i\log\left(\sqrt{\frac{\sqrt{4i^2+1}}{4i+1}} + i\right) + i\log\left(\sqrt{\frac{\sqrt{4i^2+1}}{4i+1}} - i\right) + \log\left(\sqrt{\frac{\sqrt{4i^2+1}}{4i+1}} - 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x, algorithm="fricas")`

[Out] `-1/2*sqrt(4*I)*log(1/2*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(4*I)*log(-1/2*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1/2*sqrt(-4*I)*log(1/2*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 1/2*sqrt(-4*I)*log(-1/2*sqrt(-4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 1*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + I*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x \left(\frac{i(ax-i)}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x,x)`

[Out] `Integral(1/(x*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x \left(\frac{1+ax1i}{\sqrt{a^2 x^2 + 1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)),x)
```

```
[Out] int(1/(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)), x)
```

$$3.102 \quad \int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=92

$$-\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{x} + 3ia\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + 3ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x+3*I*a*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+3*I*a*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.02, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5170, 96, 95, 218, 212, 209}

$$3ia\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{x} + 3ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(E^(((3*I)/2)*ArcTan[a*x])*x^2),x]

[Out] $-\left(\frac{(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}}{x} + (3*I)*a*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}] + (3*I)*a*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]\right)$

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1)/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a

, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-\frac{3}{2}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1 - iax)^{3/4}}{x^2(1 + iax)^{3/4}} dx \\
 &= -\frac{(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{x} - \frac{1}{2}(3ia) \int \frac{1}{x \sqrt[4]{1 - iax} (1 + iax)^{3/4}} dx \\
 &= -\frac{(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{x} - (6ia) \text{Subst} \left(\int \frac{1}{-1 + x^4} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) \\
 &= -\frac{(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{x} + (3ia) \text{Subst} \left(\int \frac{1}{1 - x^2} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + (3ia) \text{Subst} \left(\int \frac{1}{1 + x^2} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) \\
 &= -\frac{(1 - iax)^{3/4} \sqrt[4]{1 + iax}}{x} + 3ia \tan^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + 3ia \tanh^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 69, normalized size = 0.75

$$\frac{i(1 - iax)^{3/4} (i - ax + 2ax {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}))}{x(1 + iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a*x]))*x^2, x]

[Out] (I*(1 - I*a*x)^(3/4)*(I - a*x + 2*a*x*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(x*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}} x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2, x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2, x, algorithm="maxima")

[Out] integrate(1/(x^2*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 152 vs. 2(64) = 128.

time = 1.98, size = 152, normalized size = 1.65

$$\frac{3i ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - 3ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + 3ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 3i ax \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) - 2(-iax+1)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2, x, algorithm="fricas")

[Out] 1/2*(3*I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 3*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 3*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 3*I*a*x*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(-I*a*x + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^2 \left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**2,x)

[Out] Integral(1/(x**2*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^2 \left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)),x)

[Out] int(1/(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)), x)

$$3.103 \quad \int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)}}{x^3} dx$$

Optimal. Leaf size=132

$$\frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{7/4}\sqrt[4]{1+iax}}{2x^2} + \frac{9}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{9}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $3/4*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-1/2*(1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)}/x^2+9/4*a^2*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+9/4*a^2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.03, antiderivative size = 132, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5170, 98, 96, 95, 218, 212, 209}

$$\frac{9}{4}a^2\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{9}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{7/4}\sqrt[4]{1+iax}}{2x^2} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(((3*I)/2)*ArcTan[a*x])*x^3),x]

[Out] $((((3*I)/4)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x - ((1-I*a*x)^{(7/4)}*(1+I*a*x)^{(1/4)})/(2*x^2) + (9*a^2*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]))/4 + (9*a^2*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/4$

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{3}{2}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1-iax)^{3/4}}{x^3(1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2x^2} - \frac{1}{4}(3ia) \int \frac{(1-iax)^{3/4}}{x^2(1+iax)^{3/4}} dx \\
&= \frac{3ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2x^2} - \frac{1}{8}(9a^2) \int \frac{1}{x \sqrt[4]{1-iax} (1+iax)} dx \\
&= \frac{3ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2x^2} - \frac{1}{2}(9a^2) \text{Subst} \left(\int \frac{1}{-1+x^4} dx, x \right) \\
&= \frac{3ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2x^2} + \frac{1}{4}(9a^2) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x \right) \\
&= \frac{3ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{4x} - \frac{(1-iax)^{7/4} \sqrt[4]{1+iax}}{2x^2} + \frac{9}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{9}{4}a
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 81, normalized size = 0.61

$$\frac{(1-iax)^{3/4} (-2 + 3iax - 5a^2x^2 + 6a^2x^2 {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}))}{4x^2(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a*x]))*x^3, x]

[Out] ((1 - I*a*x)^(3/4)*(-2 + (3*I)*a*x - 5*a^2*x^2 + 6*a^2*x^2*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(4*x^2*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}} \right)^{\frac{3}{2}} x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3, x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x, algorithm="maxima")``[Out] integrate(1/(x^3*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)), x)`**Fricas [A]**

time = 3.47, size = 176, normalized size = 1.33

$$\frac{9a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) + 9ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) - 9ia^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 9a^2x^2 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right) + 2(5a^2x^2 + 7iax - 2)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{8x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x, algorithm="fricas")`

`[Out] 1/8*(9*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 9*I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 9*I*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 9*a^2*x^2*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) + 2*(5*a^2*x^2 + 7*I*a*x - 2)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^2`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^3 \left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}} \right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**3,x)``[Out] Integral(1/(x**3*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^3,x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
 the root of a polynomial with parameters. This might be wrong.The choice wa
 s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^3 \left(\frac{1+axi}{\sqrt{a^2 x^2 + 1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)),x)

[Out] int(1/(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)), x)

3.104 $\int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)}}{x^4} dx$

Optimal. Leaf size=170

$$-\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{12x^2} + \frac{23a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{24x} - \frac{17}{8}ia^3\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-1/3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^3+7/12*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^2+23/24*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-17/8*I*a^3*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-17/8*I*a^3*\arctanh((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.05, antiderivative size = 170, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 218, 212, 209}

$$-\frac{17}{8}ia^3\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{17}{8}ia^3 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{23a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{24x} - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{12x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((3*I)/2)*\text{ArcTan}[a*x]})*x^4), x]$

[Out] $-1/3*((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^3 + (((7*I)/12)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^2 + (23*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(24*x) - ((17*I)/8)*a^3*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}] - ((17*I)/8)*a^3*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}]$

Rule 12

$\text{Int}[(a_*)*(u_), x_Symbol] \rightarrow \text{Dist}[a, \text{Int}[u, x], x] /; \text{FreeQ}[a, x] \ \&\& \ !\text{MatchQ}[u, (b_*)*(v_)] /; \text{FreeQ}[b, x]$

Rule 95

$\text{Int}[(((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)})/((e_.) + (f_.)*(x_)), x_Symbol] \rightarrow \text{With}[q = \text{Denominator}[m], \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \text{FreeQ}[a, b, c, d, e, f], x] \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$

Rule 101

$\text{Int}[((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x] - \text{Dist}[1/((m+1)*(b*e - a*f)), \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x]$

```
(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m
+ n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1]
] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || Integ
ersQ[p, m + n])
```

Rule 156

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{3}{2}i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1-iax)^{3/4}}{x^4(1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{1}{3} \int \frac{-\frac{7ia}{2} - 2a^2x}{x^3 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} - \frac{1}{6} \int \frac{\frac{23a^2}{4} - \frac{7}{2}ia^3x}{x^2 \sqrt[4]{1-iax} (1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{23a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} + \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{23a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} + \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{23a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} + \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{23a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} - \\
&= -\frac{(1-iax)^{3/4} \sqrt[4]{1+iax}}{3x^3} + \frac{7ia(1-iax)^{3/4} \sqrt[4]{1+iax}}{12x^2} + \frac{23a^2(1-iax)^{3/4} \sqrt[4]{1+iax}}{24x} -
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 93, normalized size = 0.55

$$\frac{(1-iax)^{3/4} (-8 + 6iax + 9a^2x^2 + 23ia^3x^3 - 34ia^3x^3 {}_2F_1(\frac{3}{4}, 1; \frac{7}{4}; \frac{i+ax}{i-ax}))}{24x^3(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a*x]))*x^4], x]

[Out] ((1 - I*a*x)^(3/4)*(-8 + (6*I)*a*x + 9*a^2*x^2 + (23*I)*a^3*x^3 - (34*I)*a^3*x^3*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(24*x^3*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}} x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x)`

[Out] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x, algorithm="maxima")`

[Out] `integrate(1/(x^4*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)), x)`

Fricas [A]

time = 2.81, size = 184, normalized size = 1.08

$$\frac{-51i a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) + 51 a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) - 51 a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) + 51i a^3 x^3 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) - 2(23i a^3 x^3 - 37 a^2 x^2 - 22i a x + 8) \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{48 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x, algorithm="fricas")`

[Out] `1/48*(-51*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 51*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 51*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) + 51*I*a^3*x^3*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) - 2*(23*I*a^3*x^3 - 37*a^2*x^2 - 22*I*a*x + 8)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^3`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^4 \left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}} \right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**4,x)`

[Out] `Integral(1/(x**4*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^4,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in
 dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^4 \left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^4*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)),x)

[Out] int(1/(x^4*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)), x)

$$3.105 \quad \int \frac{e^{-\frac{3}{2}i\text{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=202

$$-\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} - \frac{63ia^3(1-iax)^{3/4}\sqrt[4]{1+iax}}{64x}$$

[Out] $-1/4*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^4+3/8*I*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^3+15/32*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x^2-63/64*I*a^3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)}/x-123/64*a^4*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-123/64*a^4*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.06, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5170, 101, 156, 12, 95, 218, 212, 209}

$$-\frac{123}{64}a^4\text{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{123}{64}a^4\operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{63ia^3(1-iax)^{3/4}\sqrt[4]{1+iax}}{64x} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} - \frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(((3*I)/2)*ArcTan[a*x])*x^5),x]

[Out] $-1/4*((1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^4 + (((3*I)/8)*a*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x^3 + (15*a^2*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/(32*x^2) - (((63*I)/64)*a^3*(1-I*a*x)^{(3/4)}*(1+I*a*x)^{(1/4)})/x - (123*a^4*\text{ArcTan}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64 - (123*a^4*\text{ArcTanh}[(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/64$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_)]/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 101

Int[((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m+1)*(c + d*x)^n*(e + f*x)^(p+1)

)/((m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 156

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{3}{2}i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{(1-iax)^{3/4}}{x^5(1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{1}{4} \int \frac{-\frac{9ia}{2} - 3a^2x}{x^4\sqrt[4]{1-iax}(1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} - \frac{1}{12} \int \frac{\frac{45a^2}{4} - 9ia^3x}{x^3\sqrt[4]{1-iax}(1+iax)^{3/4}} dx \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots \\
&= -\frac{(1-iax)^{3/4}\sqrt[4]{1+iax}}{4x^4} + \frac{3ia(1-iax)^{3/4}\sqrt[4]{1+iax}}{8x^3} + \frac{15a^2(1-iax)^{3/4}\sqrt[4]{1+iax}}{32x^2} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 99, normalized size = 0.49

$$\frac{(1-iax)^{3/4}(-16+8iax+6a^2x^2-33ia^3x^3+63a^4x^4-82a^4x^4{}_2F_1(\frac{3}{4}, 1; \frac{7}{4}, \frac{i+ax}{i-ax}))}{64x^4(1+iax)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a*x]))*x^5), x]

[Out] ((1 - I*a*x)^(3/4)*(-16 + (8*I)*a*x + 6*a^2*x^2 - (33*I)*a^3*x^3 + 63*a^4*x^4 - 82*a^4*x^4*Hypergeometric2F1[3/4, 1, 7/4, (I + a*x)/(I - a*x)]))/(64*x^4*(1 + I*a*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}} x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x)`

[Out] `int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x, algorithm="maxima")`

[Out] `integrate(1/(x^5*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2)), x)`

Fricas [A]

time = 4.65, size = 192, normalized size = 0.95

$$\frac{123a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) + 123ia^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) - 123ia^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) - 123a^4x^4 \log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right) + 2(63a^4x^4 + 93ia^3x^3 - 54a^2x^2 - 40iax + 16)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{128x^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x, algorithm="fricas")`

[Out] `-1/128*(123*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 123*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) - 123*I*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 123*a^4*x^4*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1) + 2*(63*a^4*x^4 + 93*I*a^3*x^3 - 54*a^2*x^2 - 40*I*a*x + 16)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/x^4`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^5 \left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}} \right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)/x**5,x)`

[Out] `Integral(1/(x**5*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2)), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)/x^5,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x^5 \left(\frac{1+axi}{\sqrt{a^2x^2+1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^5*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)),x)
```

```
[Out] int(1/(x^5*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2)), x)
```

3.106 $\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=373

$$\frac{4ix^3(1-iax)^{5/4}}{a^4\sqrt{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{i(1-iax)^{5/4}(1+iax)^{3/4}(521i)}{96a^4}$$

[Out] $4*I*x^3*(1-I*a*x)^(5/4)/a/(1+I*a*x)^(1/4)+475/64*(1-I*a*x)^(1/4)*(1+I*a*x)^(3/4)/a^4-17/4*x^2*(1-I*a*x)^(5/4)*(1+I*a*x)^(3/4)/a^2-1/96*I*(1-I*a*x)^(5/4)*(1+I*a*x)^(3/4)*(521*I+452*a*x)/a^4+475/128*\arctan(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))/a^4*2^(1/2)-475/128*\arctan(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4))/a^4*2^(1/2)+475/256*\ln(1-(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))/a^4*2^(1/2)-475/256*\ln(1+(1-I*a*x)^(1/4)*2^(1/2)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/2)/(1+I*a*x)^(1/2))/a^4*2^(1/2)$

Rubi [A]

time = 0.18, antiderivative size = 373, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 13, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.812$, Rules used = {5170, 99, 158, 152, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{475\text{ArcTan}\left(\frac{1-\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} - \frac{475\text{ArcTan}\left(\frac{1+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}\right)}{64\sqrt{2}a^4} - \frac{i(1-iax)^{5/4}(1+iax)^{3/4}(452ax+521i)}{96a^4} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} + \frac{475\log\left(\frac{\sqrt{1-iax}-\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}+1\right)}{128\sqrt{2}a^4} - \frac{475\log\left(\frac{\sqrt{1-iax}+\sqrt{2}\sqrt{1-iax}}{\sqrt{1+iax}}+1\right)}{128\sqrt{2}a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} + \frac{475i(1-iax)^{5/4}}{96a^4}$$

Antiderivative was successfully verified.

[In] Int[x^3/E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] $((4*I)*x^3*(1-I*a*x)^(5/4))/(a*(1+I*a*x)^(1/4)) + (475*(1-I*a*x)^(1/4)*(1+I*a*x)^(3/4))/(64*a^4) - (17*x^2*(1-I*a*x)^(5/4)*(1+I*a*x)^(3/4))/(4*a^2) - ((I/96)*(1-I*a*x)^(5/4)*(1+I*a*x)^(3/4)*(521*I+452*a*x))/a^4 + (475*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a*x)^(1/4))/(1+I*a*x)^(1/4)])/ (64*\text{Sqrt}[2]*a^4) - (475*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a*x)^(1/4))/(1+I*a*x)^(1/4)])/ (64*\text{Sqrt}[2]*a^4) + (475*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^(1/4))/(1+I*a*x)^(1/4)])/ (128*\text{Sqrt}[2]*a^4) - (475*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^(1/4))/(1+I*a*x)^(1/4)])/ (128*\text{Sqrt}[2]*a^4)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 99

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_, x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p/(b*(
m + 1)), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*
(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{
a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ
[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 158

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n +
1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p
+ 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n +
p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p
+ 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /
; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p +
2, 0] && IntegerQ[m]
```

Rule 210

```
Int[((a_.) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{5}{2}i \tan^{-1}(ax)} x^3 dx &= \int \frac{x^3(1-iax)^{5/4}}{(1+iax)^{5/4}} dx \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} - \frac{(4i) \int \frac{x^2 \sqrt[4]{1-iax} (3-\frac{17iax}{4})}{\sqrt[4]{1+iax}} dx}{a} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{i \int \frac{x \sqrt[4]{1-iax} (\frac{17ia}{2} + \frac{113a^2x}{8})}{\sqrt[4]{1+iax}} dx}{a^3} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - \frac{i(1-iax)^{5/4}(1+iax)^{3/4}(521i + \dots)}{96a^4} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - i(1-iax)^{5/4}(1+iax)^{3/4} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - i(1-iax)^{5/4}(1+iax)^{3/4} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - i(1-iax)^{5/4}(1+iax)^{3/4} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - i(1-iax)^{5/4}(1+iax)^{3/4} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - i(1-iax)^{5/4}(1+iax)^{3/4} \\
&= \frac{4ix^3(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{475\sqrt[4]{1-iax}(1+iax)^{3/4}}{64a^4} - \frac{17x^2(1-iax)^{5/4}(1+iax)^{3/4}}{4a^2} - i(1-iax)^{5/4}(1+iax)^{3/4}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 100, normalized size = 0.27

$$\frac{\sqrt[4]{1-iax} (i+ax)^2 \left(3(59+5iax+6a^2x^2) - 95 \cdot 2^{3/4} \sqrt[4]{1+iax} {}_2F_1\left(\frac{1}{4}, \frac{9}{4}; \frac{13}{4}; \frac{1}{2}(1-iax)\right) \right)}{72a^4 \sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[x^3/E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] -1/72*((1 - I*a*x)^(1/4)*(I + a*x)^2*(3*(59 + (5*I)*a*x + 6*a^2*x^2) - 95*2^(3/4)*(1 + I*a*x)^(1/4)*Hypergeometric2F1[1/4, 9/4, 13/4, (1 - I*a*x)/2]))/(a^4*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{x^3}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

[Out] int(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="maxima")

[Out] integrate(x^3/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)

Fricas [A]

time = 3.13, size = 304, normalized size = 0.82

$$\frac{96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}} \log\left(\frac{\frac{225625}{4096a^8} + \sqrt{\frac{1-\sqrt{a^2x^2+1}}{ax+1}}}{-96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}}}\right) - 96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}} \log\left(-\frac{\frac{225625}{4096a^8} + \sqrt{\frac{1-\sqrt{a^2x^2+1}}{ax+1}}}{-96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}}}\right) - 96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}} \log\left(\frac{\frac{225625}{4096a^8} + \sqrt{\frac{1-\sqrt{a^2x^2+1}}{ax+1}}}{-96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}}}\right) + 96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}} \log\left(\frac{\frac{225625}{4096a^8} + \sqrt{\frac{1-\sqrt{a^2x^2+1}}{ax+1}}}{-96(a^2 - ia^4) \sqrt{\frac{225625}{4096a^8}}}\right) + (48ia^2x^4 - 130a^2x^3 - 226a^2x^2 + 521ax - 2467i) \sqrt{a^2x^2+1} \sqrt{\frac{1-\sqrt{a^2x^2+1}}{ax+1}}}{192(a^2 - ia^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="fricas")

[Out] 1/192*(96*(a^5*x - I*a^4)*sqrt(225625/4096*I/a^8)*log(64/475*I*a^4*sqrt(225625/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 96*(a^5*x - I*a^4)*sqrt(225625/4096*I/a^8)*log(-64/475*I*a^4*sqrt(225625/4096*I/a^8) + sqrt(I

```
*sqrt(a^2*x^2 + 1)/(a*x + I))) - 96*(a^5*x - I*a^4)*sqrt(-225625/4096*I/a^8
)*log(64/475*I*a^4*sqrt(-225625/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x
+ I))) + 96*(a^5*x - I*a^4)*sqrt(-225625/4096*I/a^8)*log(-64/475*I*a^4*sqr
t(-225625/4096*I/a^8) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + (48*I*a^4*x^
4 - 136*a^3*x^3 - 226*I*a^2*x^2 + 521*a*x - 2467*I)*sqrt(a^2*x^2 + 1)*sqrt(
I*sqrt(a^2*x^2 + 1)/(a*x + I)))/(a^5*x - I*a^4)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^3}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**3/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2), x)
```

```
[Out] Integral(x**3/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(5/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^3/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2), x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3}{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2x^2+1}}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^3/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

```
[Out] int(x^3/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

3.107 $\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=371

$$\frac{2i(1-iax)^{9/4}}{a^3\sqrt[4]{1+iax}} - \frac{55i\sqrt[4]{1-iax}(1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} - 55i\text{Ar}$$

[Out] $-2*I*(1-I*a*x)^{(9/4)}/a^3/(1+I*a*x)^{(1/4)}-55/8*I*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^3-11/4*I*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^3-1/3*I*(1-I*a*x)^{(9/4)}*(1+I*a*x)^{(3/4)}/a^3-55/16*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3*2^{(1/2)}+55/16*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a^3*2^{(1/2)}-55/32*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3*2^{(1/2)}+55/32*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a^3*2^{(1/2)}$

Rubi [A]

time = 0.18, antiderivative size = 371, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.750$, Rules used = {5170, 91, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{55i\text{ArcTan}\left(1-\frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} + \frac{55i\text{ArcTan}\left(1+\frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{8\sqrt{2}a^3} - \frac{i(1+iax)^{3/4}(1-iax)^{3/4}}{3a^3} - \frac{2i(1-iax)^{5/4}}{a^2\sqrt[4]{1+iax}} - \frac{11i(1+iax)^{3/4}(1-iax)^{3/4}}{4a^3} - \frac{55i(1+iax)^{3/4}\sqrt[4]{1-iax}}{8a^3} - \frac{55i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}-\frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}+1\right)}{16\sqrt{2}a^3} + \frac{55i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}+\frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}+1\right)}{16\sqrt{2}a^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{((5*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $((-2*I)*(1-I*a*x)^{(9/4)})/(a^3*(1+I*a*x)^{(1/4)}) - (((55*I)/8)*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/a^3 - (((11*I)/4)*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)})/a^3 - ((I/3)*(1-I*a*x)^{(9/4)}*(1+I*a*x)^{(3/4)})/a^3 - (((55*I)/8)*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(a^3*\text{Sqrt}[2]) + (((55*I)/8)*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(a^3*\text{Sqrt}[2]) - (((55*I)/16)*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(a^3*\text{Sqrt}[2]) + (((55*I)/16)*\text{Log}[1+\text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(a^3*\text{Sqrt}[2])$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_. + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 91

```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)
/(d^2*(d*e - c*f)*(n + 1))), x] - Dist[1/(d^2*(d*e - c*f)*(n + 1)), Int[(c
+ d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1)
+ c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n
+ 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] ||
(EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p,
1])))
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{5}{2}i \tan^{-1}(ax)} x^2 dx &= \int \frac{x^2(1-iax)^{5/4}}{(1+iax)^{5/4}} dx \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} + \frac{(2i) \int \frac{(1-iax)^{5/4} \left(-\frac{5ia}{2} - \frac{a^2 x}{2}\right)}{\sqrt[4]{1+iax}} dx}{a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} + \frac{11 \int \frac{(1-iax)^{5/4}}{\sqrt[4]{1+iax}} dx}{2a^2} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} + \frac{55 \int \frac{1}{\sqrt[4]{1+iax}} dx}{2a^2} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3} \\
&= -\frac{2i(1-iax)^{9/4}}{a^3 \sqrt[4]{1+iax}} - \frac{55i \sqrt[4]{1-iax} (1+iax)^{3/4}}{8a^3} - \frac{11i(1-iax)^{5/4}(1+iax)^{3/4}}{4a^3} - \frac{i(1-iax)^{9/4}(1+iax)^{3/4}}{3a^3}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 91, normalized size = 0.25

$$-\frac{\sqrt[4]{1-iax} (i+ax)^2 \left(-21i+3ax+11i2^{3/4}\sqrt[4]{1+iax} {}_2F_1\left(\frac{1}{4}, \frac{9}{4}; \frac{13}{4}; \frac{1}{2}(1-iax)\right)\right)}{9a^3 \sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] $-1/9*((1 - I*a*x)^{(1/4)}*(I + a*x)^2*(-21*I + 3*a*x + (11*I)*2^{(3/4)}*(1 + I*a*x)^{(1/4)}*Hypergeometric2F1[1/4, 9/4, 13/4, (1 - I*a*x)/2]))/(a^3*(1 + I*a*x)^{(1/4)})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

[Out] int(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="maxima")

[Out] integrate(x^2/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)

Fricas [A]

time = 2.47, size = 296, normalized size = 0.80

$$\frac{12(a^2x - ia^2)\sqrt{\frac{3025}{64a^6}} \log\left(\frac{\frac{1}{2}a^2\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1+\sqrt{a^2x^2+1}}{az+i}}}{-12(a^2x - ia^2)\sqrt{\frac{3025}{64a^6}} \log\left(-\frac{1}{2}a^2\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1+\sqrt{a^2x^2+1}}{az+i}}\right) - 12(a^2x - ia^2)\sqrt{\frac{3025}{64a^6}} \log\left(\frac{\frac{1}{2}a^2\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1+\sqrt{a^2x^2+1}}{az+i}}}{24(a^2x - ia^2)}\right) + 12(a^2x - ia^2)\sqrt{\frac{3025}{64a^6}} \log\left(-\frac{1}{2}a^2\sqrt{\frac{3025}{64a^6}} + \sqrt{\frac{1+\sqrt{a^2x^2+1}}{az+i}}\right) + (8a^2x^2 - 26a^2x^2 - 61iaz - 287)\sqrt{a^2x^2+1}\sqrt{\frac{1+\sqrt{a^2x^2+1}}{az+i}}\right)}{24(a^2x - ia^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="fricas")

[Out] $1/24*(12*(a^4*x - I*a^3)*sqrt(3025/64*I/a^6)*log(8/55*a^3*sqrt(3025/64*I/a^6) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 12*(a^4*x - I*a^3)*sqrt(3025/64*I/a^6)*log(-8/55*a^3*sqrt(3025/64*I/a^6) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - 12*(a^4*x - I*a^3)*sqrt(-3025/64*I/a^6)*log(8/55*a^3*sqrt(-3025/64*I/a^6) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 12*(a^4*x - I*a^3)*sqrt(-3025/64*I/a^6)*log(-8/55*a^3*sqrt(-3025/64*I/a^6) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + (8*I*a^3*x^3 - 26*a^2*x^2 - 61*I*a*x - 287)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/(a^4*x - I*a^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**2/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2),x)
```

```
[Out] Integral(x**2/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(5/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2}{\left(\frac{1+ax \text{ li}}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)
```

```
[Out] int(x^2/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

3.108 $\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=324

$$\frac{2(1-iax)^{9/4}}{a^2\sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{25\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \dots$$

[Out] $-2*(1-I*a*x)^{(9/4)}/a^2/(1+I*a*x)^{(1/4)}-25/4*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a^2-5/2*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)}/a^2-25/8*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)/(1+I*a*x)^{(1/4)})}/a^2*2^{(1/2)}+25/8*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)/(1+I*a*x)^{(1/4)})}/a^2*2^{(1/2)}-25/16*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)/(1+I*a*x)^{(1/4)})+(1-I*a*x)^{(1/2)/(1+I*a*x)^{(1/2)})}/a^2*2^{(1/2)}+25/16*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)/(1+I*a*x)^{(1/4)})+(1-I*a*x)^{(1/2)/(1+I*a*x)^{(1/2)})}/a^2*2^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 324, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 79, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{25\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} + \frac{25\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{4\sqrt{2}a^2} - \frac{2(1-iax)^{9/4}}{a^2\sqrt[4]{1+iax}} - \frac{5(1+iax)^{3/4}(1-iax)^{3/4}}{2a^2} - \frac{25(1+iax)^{3/4}\sqrt[4]{1-iax}}{4a^2} - \frac{25\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2} + \frac{25\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{8\sqrt{2}a^2}$$

Antiderivative was successfully verified.

[In] Int[x/E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] $(-2*(1-I*a*x)^{(9/4)})/(a^2*(1+I*a*x)^{(1/4)}) - (25*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)})/(4*a^2) - (5*(1-I*a*x)^{(5/4)}*(1+I*a*x)^{(3/4)})/(2*a^2) - (25*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(4*\text{Sqrt}[2]*a^2) + (25*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(4*\text{Sqrt}[2]*a^2) - (25*\text{Log}[1 + \text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] - (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(8*\text{Sqrt}[2]*a^2) + (25*\text{Log}[1 + \text{Sqrt}[1-I*a*x]/\text{Sqrt}[1+I*a*x] + (\text{Sqrt}[2]*(1-I*a*x)^{(1/4)})/(1+I*a*x)^{(1/4)}])/(8*\text{Sqrt}[2]*a^2)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +

$d*(x^p/b)^n, x, (a + b*x)^{1/p}, x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 79

$\text{Int}[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(f*(p + 1)*(c*f - d*e))), x] - \text{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), \text{Int}[(c + d*x)^n*(e + f*x)^{(p + 1)}, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \ \&\& \ \text{LtQ}[p, -1] \ \&\& \ (\ !\text{LtQ}[n, -1] \ || \ \text{IntegerQ}[p] \ || \ \!(\text{IntegerQ}[n] \ || \ \!(\text{EqQ}[e, 0] \ || \ \!(\text{EqQ}[c, 0] \ || \ \text{LtQ}[p, n])))$

Rule 210

$\text{Int}[(a_.) + (b_.)*(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 217

$\text{Int}[(a_.) + (b_.)*(x_)^4]^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*r), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 246

$\text{Int}[(a_.) + (b_.)*(x_)^n]^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[a^{(p + 1/n)}, \text{Subst}[\text{Int}[1/(1 - b*x^n)^{(p + 1/n + 1)}, x], x, x/(a + b*x^n)^{1/n}], x] /; \text{FreeQ}\{a, b\}, x\} \ \&\& \ \text{IGtQ}[n, 0] \ \&\& \ \text{LtQ}[-1, p, 0] \ \&\& \ \text{NeQ}[p, -2^{(-1)}] \ \&\& \ \text{IntegerQ}[p + 1/n]$

Rule 631

$\text{Int}[(a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2]^{-1}, x_Symbol] \rightarrow \text{With}\{q = 1 - 4*\text{Simplify}[a*(c/b^2)]\}, \text{Dist}[-2/b, \text{Subst}[\text{Int}[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; \text{RationalQ}[q] \ \&\& \ (\text{EqQ}[q^2, 1] \ || \ \!\text{RationalQ}[b^2 - 4*a*c]) /; \text{FreeQ}\{a, b, c\}, x\} \ \&\& \ \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 642

$\text{Int}[(d_.) + (e_.)*(x_)]/((a_.) + (b_.)*(x_.) + (c_.)*(x_.)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d,$

$e\}, x] \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 1176

$\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[2*(d/e), 2]\}, \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e + q*x + x^2, x], x], x] + \text{Dist}[e/(2*c), \text{Int}[1/\text{Simp}[d/e - q*x + x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{PosQ}[d*e]$

Rule 1179

$\text{Int}[\frac{(d_.) + (e_.)x^2}{(a_.) + (c_.)x^4}, x_Symbol] \rightarrow \text{With}[\{q = \text{Rt}[-2*(d/e), 2]\}, \text{Dist}[e/(2*c*q), \text{Int}[(q - 2*x)/\text{Simp}[d/e + q*x - x^2, x], x], x] + \text{Dist}[e/(2*c*q), \text{Int}[(q + 2*x)/\text{Simp}[d/e - q*x - x^2, x], x], x]] /; \text{FreeQ}\{a, c, d, e\}, x] \&\& \text{EqQ}[c*d^2 - a*e^2, 0] \&\& \text{NegQ}[d*e]$

Rule 5170

$\text{Int}[E^{\text{ArcTan}[(a_.)x]}*(n_.)x^{(m_.)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, m, n\}, x] \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
\int e^{-\frac{5}{2}i \tan^{-1}(ax)} x dx &= \int \frac{x(1-iax)^{5/4}}{(1+iax)^{5/4}} dx \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{(5i) \int \frac{(1-iax)^{5/4}}{\sqrt[4]{1+iax}} dx}{a} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{(25i) \int \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx}{4a} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{(25i) \int \frac{1}{(1+iax)^{5/4}} dx}{(1)} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{25 \text{Subst} \left(\int \frac{1}{(1+iax)^{5/4}} dx \right)}{(1)} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{25 \text{Subst} \left(\int \frac{1}{(1+iax)^{5/4}} dx \right)}{(1)} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{25 \text{Subst} \left(\int \frac{1}{(1+iax)^{5/4}} dx \right)}{(1)} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{25 \text{Subst} \left(\int \frac{1}{(1+iax)^{5/4}} dx \right)}{(1)} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} + \frac{25 \log \left(1 + \frac{iax}{\sqrt[4]{1+iax}} \right)}{(1)} \\
&= -\frac{2(1-iax)^{9/4}}{a^2 \sqrt[4]{1+iax}} - \frac{25\sqrt[4]{1-iax}(1+iax)^{3/4}}{4a^2} - \frac{5(1-iax)^{5/4}(1+iax)^{3/4}}{2a^2} - \frac{25 \tan^{-1} \left(\frac{iax}{\sqrt[4]{1+iax}} \right)}{(1)}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 63, normalized size = 0.19

$$\frac{2(1-iax)^{9/4} \left(-\frac{9}{\sqrt[4]{1+iax}} + 5 \cdot 2^{3/4} {}_2F_1 \left(\frac{1}{4}, \frac{9}{4}; \frac{13}{4}; \frac{1}{2}(1-iax) \right) \right)}{9a^2}$$

Antiderivative was successfully verified.

[In] Integrate[x/E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] $(2*(1 - I*a*x)^{(9/4)}*(-9/(1 + I*a*x)^{(1/4)} + 5*2^{(3/4)}*Hypergeometric2F1[1/4, 9/4, 13/4, (1 - I*a*x)/2]))/(9*a^2)$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)`

[Out] `int(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="maxima")`

[Out] `integrate(x/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)`

Fricas [A]

time = 5.49, size = 289, normalized size = 0.89

$$\frac{2(a^2x - ia^2)\sqrt{\frac{625}{16a^4}} \log\left(\frac{\frac{625}{16a^4}\sqrt{\frac{625}{16a^4} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{ax+i}\right) - 2(a^2x - ia^2)\sqrt{\frac{625}{16a^4}} \log\left(\frac{-\frac{625}{16a^4}\sqrt{\frac{625}{16a^4} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{ax+i}\right) - 2(a^2x - ia^2)\sqrt{\frac{625}{16a^4}} \log\left(\frac{\frac{625}{16a^4}\sqrt{-\frac{625}{16a^4} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{ax+i}\right) + 2(a^2x - ia^2)\sqrt{\frac{625}{16a^4}} \log\left(\frac{-\frac{625}{16a^4}\sqrt{-\frac{625}{16a^4} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}}{ax+i}\right) - \sqrt{a^2x^2+1}(2ia^2x - 9ax + 43i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{4(a^2x - ia^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="fricas")`

[Out] $-1/4*(2*(a^3*x - I*a^2)*\sqrt{625/16*I/a^4}*\log(4/25*I*a^2*\sqrt{625/16*I/a^4} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - 2*(a^3*x - I*a^2)*\sqrt{625/16*I/a^4}*\log(-4/25*I*a^2*\sqrt{625/16*I/a^4} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)})) - 2*(a^3*x - I*a^2)*\sqrt{-625/16*I/a^4}*\log(4/25*I*a^2*\sqrt{-625/16*I/a^4} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) + 2*(a^3*x - I*a^2)*\sqrt{-625/16*I/a^4}*\log(-4/25*I*a^2*\sqrt{-625/16*I/a^4} + \sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)}) - \sqrt{a^2*x^2 + 1}*(2*I*a^2*x^2 - 9*a*x + 43*I)*\sqrt{I*\sqrt{a^2*x^2 + 1}/(a*x + I)))/(a^3*x - I*a^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2), x)
```

```
[Out] Integral(x/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(5/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2), x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x}{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

```
[Out] int(x/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

3.109 $\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=299

$$\frac{4i(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{5i\sqrt[4]{1-iax}(1+iax)^{3/4}}{a} + \frac{5i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{5i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a}$$

[Out] $4*I*(1-I*a*x)^{(5/4)}/a/(1+I*a*x)^{(1/4)}+5*I*(1-I*a*x)^{(1/4)}*(1+I*a*x)^{(3/4)}/a+5/2*I*\arctan(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}-5/2*I*\arctan(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)})/a*2^{(1/2)}+5/4*I*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}-5/4*I*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2)})/a*2^{(1/2)}$

Rubi [A]

time = 0.13, antiderivative size = 299, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {5169, 49, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{5i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} - \frac{5i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{\sqrt{2}a} + \frac{4i(1-iax)^{5/4}}{a\sqrt[4]{1+iax}} + \frac{5i(1+iax)^{3/4}\sqrt[4]{1-iax}}{a} + \frac{5i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a} - \frac{5i\log\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{2\sqrt{2}a}$$

Antiderivative was successfully verified.

[In] Int[E^(((−5*I)/2)*ArcTan[a*x]), x]

[Out] $((4*I)*(1 - I*a*x)^{(5/4)})/(a*(1 + I*a*x)^{(1/4)}) + ((5*I)*(1 - I*a*x)^{(1/4)}*(1 + I*a*x)^{(3/4)})/a + ((5*I)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) - ((5*I)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) + (((5*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] - (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a) - (((5*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a*x]/\text{Sqrt}[1 + I*a*x] + (\text{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}])/(\text{Sqrt}[2]*a)$

Rule 49

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*(b*c - a*d)/(

```
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5169

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1
+ I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{5}{2}i \tan^{-1}(ax)} dx &= \int \frac{(1-iax)^{5/4}}{(1+iax)^{5/4}} dx \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} - 5 \int \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} - \frac{5}{2} \int \frac{1}{(1-iax)^{3/4} \sqrt[4]{1+iax}} dx \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} - \frac{(10i) \text{Subst} \left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-iax} \right)}{a} \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} - \frac{(10i) \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{a} \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} - \frac{(5i) \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{a} \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} - \frac{(5i) \text{Subst} \left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{2a} \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} + \frac{5i \log \left(1 + \frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{2\sqrt{2} a} \\
&= \frac{4i(1-iax)^{5/4}}{a^4 \sqrt[4]{1+iax}} + \frac{5i \sqrt[4]{1-iax} (1+iax)^{3/4}}{a} + \frac{5i \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} \right)}{\sqrt{2} a} - \frac{5i \tan^{-1} \left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} \right)}{\sqrt{2} a}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 39, normalized size = 0.13

$$\frac{8ie^{-\frac{1}{2}i \text{ArcTan}(ax)} {}_2F_1\left(-\frac{1}{4}, 2, \frac{3}{4}, -e^{2i \text{ArcTan}(ax)}\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((−5*I)/2)*ArcTan[a*x]),x]

[Out] ((8*I)*Hypergeometric2F1[−1/4, 2, 3/4, −E^((2*I)*ArcTan[a*x])])/(a*E^((I/2)*ArcTan[a*x]))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)``[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="maxima")``[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(-5/2), x)`**Fricas [A]**

time = 4.23, size = 261, normalized size = 0.87

$$\frac{(a^2x - ia)\sqrt{\frac{25i}{a^2}} \log\left(\frac{1}{2}a\sqrt{\frac{25i}{a^2} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) - (a^2x - ia)\sqrt{\frac{25i}{a^2}} \log\left(-\frac{1}{2}a\sqrt{\frac{25i}{a^2} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) - (a^2x - ia)\sqrt{-\frac{25i}{a^2}} \log\left(\frac{1}{2}a\sqrt{-\frac{25i}{a^2} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) + (a^2x - ia)\sqrt{-\frac{25i}{a^2}} \log\left(-\frac{1}{2}a\sqrt{-\frac{25i}{a^2} + \sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}\right) + 2\sqrt{a^2x^2+1}(-iax-9)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}}{2(a^2x - ia)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="fricas")`

```
[Out] -1/2*((a^2*x - I*a)*sqrt(25*I/a^2)*log(1/5*a*sqrt(25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - (a^2*x - I*a)*sqrt(25*I/a^2)*log(-1/5*a*sqrt(25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - (a^2*x - I*a)*sqrt(-25*I/a^2)*log(1/5*a*sqrt(-25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + (a^2*x - I*a)*sqrt(-25*I/a^2)*log(-1/5*a*sqrt(-25*I/a^2) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) + 2*sqrt(a^2*x^2 + 1)*(-I*a*x - 9)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)))/(a^2*x - I*a)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2),x)

[Out] Integral(((I*a*x + 1)/sqrt(a**2*x**2 + 1))**(-5/2), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)

[Out] int(1/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)

$$3.110 \quad \int \frac{e^{-\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=293

$$\frac{8\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 2\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)$$

[Out] $8*(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}+2*\arctan((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})-2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+1/2*\ln(1-(1-I*a*x)^{(1/4)}*2^{(1/2)})/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2))*2^{(1/2)}-1/2*\ln(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/2)}/(1+I*a*x)^{(1/2))*2^{(1/2)}+ \operatorname{arctan}(1-(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4))*2^{(1/2)}-\operatorname{arctan}(1+(1-I*a*x)^{(1/4)}*2^{(1/2)}/(1+I*a*x)^{(1/4))*2^{(1/2)}}$

Rubi [A]

time = 0.16, antiderivative size = 293, normalized size of antiderivative = 1.00, number of steps used = 19, number of rules used = 16, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5170, 100, 21, 132, 65, 246, 217, 1179, 642, 1176, 631, 210, 95, 304, 209, 212}

$$2\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right) + \frac{8\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} - \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - \frac{\log\left(\frac{\sqrt{1-iax}}{\sqrt{1+iax}} + \frac{\sqrt{2}\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 1\right)}{\sqrt{2}} - 2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(((5*I)/2)*ArcTan[a*x])*x), x]`

[Out] $(8*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)} + 2*\operatorname{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] + \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}] - \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}] - 2*\operatorname{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}] + \operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a*x]/\operatorname{Sqrt}[1 + I*a*x] - (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/\operatorname{Sqrt}[2] - \operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a*x]/\operatorname{Sqrt}[1 + I*a*x] + (\operatorname{Sqrt}[2]*(1 - I*a*x)^{(1/4)})/(1 + I*a*x)^{(1/4)}]/\operatorname{Sqrt}[2]$

Rule 21

`Int[(u_.)*((a_.) + (b_.)*(v_))^(m_.)*((c_.) + (d_.)*(v_))^(n_.), x_Symbol] :> Dist[(b/d)^m, Int[u*(c + d*v)^(m + n), x], x] /; FreeQ[{a, b, c, d, n}, x] && EqQ[b*c - a*d, 0] && IntegerQ[m] && (!IntegerQ[n] || SimplerQ[c + d*x, a + b*x])`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ`

$[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 95

$\text{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{With}[\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)} / (b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)} / (c + d*x)^{(1/q)}], x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{EqQ}[m + n + 1, 0] \&\& \text{RationalQ}[n] \&\& \text{LtQ}[-1, m, 0] \&\& \text{SimplerQ}[a + b*x, c + d*x]$

Rule 100

$\text{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)*(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}*((e + f*x)^{(p+1)} / (b*(b*e - a*f)*(m+1))), x] + \text{Dist}[1 / (b*(b*e - a*f)*(m+1)), \text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-2)}*(e + f*x)^p * \text{Simp}[a*d*(d*e*(n-1) + c*f*(p+1)) + b*c*(d*e*(m-n+2) - c*f*(m+p+2)) + d*(a*d*f*(n+p) + b*(d*e*(m+1) - c*f*(m+n+p+1))]*x, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, p\}, x] \&\& \text{LtQ}[m, -1] \&\& \text{GtQ}[n, 1] \&\& (\text{IntegersQ}[2*m, 2*n, 2*p] \|\| \text{IntegersQ}[m, n+p] \|\| \text{IntegersQ}[p, m+n])$

Rule 132

$\text{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[b*d^{(m+n)}*f^p, \text{Int}[(a + b*x)^{(m-1)} / (c + d*x)^m, x], x] + \text{Int}[(a + b*x)^{(m-1)}*((e + f*x)^p / (c + d*x)^m * \text{ExpandToSum}[(a + b*x)*(c + d*x)^{(-p-1)} - (b*d^{(-p-1)}*f^p) / (e + f*x)^p], x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, m, n\}, x] \&\& \text{EqQ}[m + n + p + 1, 0] \&\& \text{ILtQ}[p, 0] \&\& (\text{GtQ}[m, 0] \|\| \text{SumSimplerQ}[m, -1] \|\| !(\text{GtQ}[n, 0] \|\| \text{SumSimplerQ}[n, -1]))$

Rule 209

$\text{Int}[(a + b*(x^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(1 / (\text{Rt}[a, 2]*\text{Rt}[b, 2])) * \text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{GtQ}[a, 0] \|\| \text{GtQ}[b, 0])$

Rule 210

$\text{Int}[(a + b*(x^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{(-1)} * \text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \|\| \text{LtQ}[b, 0])$

Rule 212

$\text{Int}[(a + b*(x^2)^{-1}), x_Symbol] \rightarrow \text{Simp}[(1 / (\text{Rt}[a, 2]*\text{Rt}[-b, 2])) * \text{ArcTanh}[\text{Rt}[-b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{NegQ}[a/b] \&\& (\text{GtQ}[a, 0] \|\| \text{GtQ}[b, 0])$

$Q[a, 0] \parallel LtQ[b, 0]$)

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a
/b, 0]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x])/b], x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Free
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{5}{2}i \tan^{-1}(ax)}}{x} dx &= \int \frac{(1 - iax)^{5/4}}{x(1 + iax)^{5/4}} dx \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - \frac{(4i) \int \frac{\frac{ia - a^2x}{4} \sqrt[4]{1 + iax}}{x(1 - iax)^{3/4} \sqrt[4]{1 + iax}} dx}{a} \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} + \int \frac{(1 + iax)^{3/4}}{x(1 - iax)^{3/4}} dx \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} + (ia) \int \frac{1}{(1 - iax)^{3/4} \sqrt[4]{1 + iax}} dx + \int \frac{1}{x(1 - iax)^{3/4} \sqrt[4]{1 + iax}} dx \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - 4 \text{Subst} \left(\int \frac{1}{\sqrt[4]{2 - x^4}} dx, x, \sqrt[4]{1 - iax} \right) + 4 \text{Subst} \left(\int \frac{x^2}{-1 + x^4} dx, x, \sqrt[4]{1 - iax} \right) \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - 2 \text{Subst} \left(\int \frac{1}{1 - x^2} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + 2 \text{Subst} \left(\int \frac{1}{1 + x^2} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} + 2 \tan^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) - 2 \text{Subst} \left(\int \frac{1 - x^2}{1 + x^4} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} + 2 \tan^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + \frac{\text{Subst} \left(\int \frac{\sqrt{2} + 2}{-1 - \sqrt{2} x} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right)}{\sqrt{2}} \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} + 2 \tan^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + \frac{\log \left(1 + \frac{\sqrt{1 - iax}}{\sqrt{1 + iax}} \right)}{\sqrt{2}} \\
&= \frac{8\sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} + 2 \tan^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + \sqrt{2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} \right) - \sqrt{2} \tan^{-1} \left(1 + \frac{\sqrt{2} \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.04, size = 106, normalized size = 0.36

$$\frac{\sqrt[4]{1-iax} \left(20 - 20 {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}\right) + 2^{3/4}(1-iax)\sqrt[4]{1+iax} {}_2F_1\left(\frac{5}{4}, \frac{5}{4}; \frac{9}{4}; \frac{1}{2}(1-iax)\right) \right)}{5\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((5*I)/2)*ArcTan[a*x]))*x), x]

[Out] ((1 - I*a*x)^(1/4)*(20 - 20*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]) + 2^(3/4)*(1 - I*a*x)*(1 + I*a*x)^(1/4)*Hypergeometric2F1[5/4, 5/4, 9/4, (1 - I*a*x)/2]))/(5*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}} x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)/x,x)

[Out] int(1/(((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)/x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)/x,x, algorithm="maxima")

[Out] integrate(1/(x*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)), x)

Fricas [A]

time = 2.35, size = 329, normalized size = 1.12

$$\frac{\sqrt{a}(ax-1)\log\left(\frac{1}{2}\sqrt{a} + \sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}}\right) - \sqrt{a}(ax-1)\log\left(-\frac{1}{2}\sqrt{a} + \sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}}\right) - \sqrt{a}(ax-1)\log\left(\frac{1}{2}\sqrt{a} + \sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}}\right) + \sqrt{a}(ax-1)\log\left(-\frac{1}{2}\sqrt{a} + \sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}}\right) - 2(ax-1)\log\left(\sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}} + 1\right) - 2(-ax-1)\log\left(\sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}} + 1\right) - 2(ax+1)\log\left(\sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}} - 1\right) + 2(-ax-1)\log\left(\sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}} - 1\right) - 16\sqrt{a^2x^2+1}\sqrt{\frac{\sqrt{a^2x^2+1}}{ax+1}}}{2(ax-1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(((1+I*a*x)/(a^2*x^2+1)^(1/2)))^(5/2)/x,x, algorithm="fricas")

[Out] 1/2*(sqrt(4*I)*(a*x - I)*log(1/2*I*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))) - sqrt(4*I)*(a*x - I)*log(-1/2*I*sqrt(4*I) + sqrt(I*sqrt(a^2*x^2 +

$$\frac{1}{(ax + I)} - \sqrt{-4I}(ax - I) \log\left(\frac{1}{2}I\sqrt{-4I} + \sqrt{I\sqrt{a^2x^2 + 1}}\right) + \sqrt{-4I}(ax - I) \log\left(-\frac{1}{2}I\sqrt{-4I} + \sqrt{I\sqrt{a^2x^2 + 1}}\right) - 2(ax - I) \log\left(\sqrt{I\sqrt{a^2x^2 + 1}}\right) + \sqrt{-4I} - 2(-Iax - 1) \log\left(\sqrt{I\sqrt{a^2x^2 + 1}}\right) + I - 2(Iax + 1) \log\left(\sqrt{I\sqrt{a^2x^2 + 1}}\right) - I + 2(ax - I) \log\left(\sqrt{I\sqrt{a^2x^2 + 1}}\right) - 1 - 16I\sqrt{a^2x^2 + 1} \sqrt{I\sqrt{a^2x^2 + 1}}\right) / (ax - I)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x \left(\frac{i(ax-i)}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x,x)

[Out] Integral(1/(x*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(5/2)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x \left(\frac{1+ax \operatorname{li}}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)),x)

[Out] int(1/(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)), x)

$$3.111 \quad \int \frac{e^{-\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=121

$$-\frac{10ia\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{(1-iax)^{5/4}}{x\sqrt[4]{1+iax}} - 5ia \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + 5ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] -10*I*a*(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-(1-I*a*x)^(5/4)/x/(1+I*a*x)^(1/4)-5*I*a*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))+5*I*a*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))

Rubi [A]

time = 0.03, antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5170, 96, 95, 304, 209, 212}

$$-5ia \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{5/4}}{x\sqrt[4]{1+iax}} - \frac{10ia\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} + 5ia \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] Int[1/(E^(((5*I)/2)*ArcTan[a*x]))*x^2),x]

[Out] ((-10*I)*a*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4) - (1 - I*a*x)^(5/4)/(x*(1 + I*a*x)^(1/4)) - (5*I)*a*ArcTan[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)] + (5*I)*a*ArcTanh[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 209

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 304

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-\frac{5}{2}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1 - iax)^{5/4}}{x^2(1 + iax)^{5/4}} dx \\
 &= -\frac{(1 - iax)^{5/4}}{x^4 \sqrt[4]{1 + iax}} - \frac{1}{2}(5ia) \int \frac{\sqrt[4]{1 - iax}}{x(1 + iax)^{5/4}} dx \\
 &= -\frac{10ia \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - \frac{(1 - iax)^{5/4}}{x \sqrt[4]{1 + iax}} - \frac{1}{2}(5ia) \int \frac{1}{x(1 - iax)^{3/4} \sqrt[4]{1 + iax}} dx \\
 &= -\frac{10ia \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - \frac{(1 - iax)^{5/4}}{x \sqrt[4]{1 + iax}} - (10ia) \text{Subst} \left(\int \frac{x^2}{-1 + x^4} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) \\
 &= -\frac{10ia \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - \frac{(1 - iax)^{5/4}}{x \sqrt[4]{1 + iax}} + (5ia) \text{Subst} \left(\int \frac{1}{1 - x^2} dx, x, \frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) - (5ia) \int \frac{1}{x(1 - iax)^{3/4} \sqrt[4]{1 + iax}} dx \\
 &= -\frac{10ia \sqrt[4]{1 - iax}}{\sqrt[4]{1 + iax}} - \frac{(1 - iax)^{5/4}}{x \sqrt[4]{1 + iax}} - 5ia \tan^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right) + 5ia \tanh^{-1} \left(\frac{\sqrt[4]{1 + iax}}{\sqrt[4]{1 - iax}} \right)
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 69, normalized size = 0.57

$$\frac{i\sqrt[4]{1-iax} \left(i - 9ax + 10ax {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}\right)\right)}{x\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((5*I)/2)*ArcTan[a*x]))*x^2, x]

[Out] (I*(1 - I*a*x)^(1/4)*(I - 9*a*x + 10*a*x*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(x*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}} x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2, x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2, x, algorithm="maxima")

[Out] integrate(1/(x^2*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 212 vs. 2(83) = 166.

time = 2.95, size = 212, normalized size = 1.75

$$\frac{2\sqrt{a^2x^2+1}(9ax-i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 5(-ia^2x^2-ax)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right) - 5(a^2x^2-iax)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right) + 5(a^2x^2-iax)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right) + 5(i a^2x^2+ax)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right)}{2(ax^2-ix)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2, x, algorithm="fricas")

[Out] -1/2*(2*sqrt(a^2*x^2 + 1)*(9*a*x - I)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 5*(-I*a^2*x^2 - a*x)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 5*(a^2*x^2 - I*a*x)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 5*(a^2*x^2 - I

$*a*x)*\log(\sqrt{I*\sqrt{a^2*x^2 + 1}}/(a*x + I)) - I) + 5*(I*a^2*x^2 + a*x)*\log(\sqrt{I*\sqrt{a^2*x^2 + 1}}/(a*x + I)) - 1)/(a*x^2 - I*x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^2 \left(\frac{i(ax-i)}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**2,x)

[Out] Integral(1/(x**2*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(5/2)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^2 \left(\frac{1+ax \ 1i}{\sqrt{a^2x^2 + 1}} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)),x)

[Out] int(1/(x^2*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)), x)

$$3.112 \quad \int \frac{e^{-\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^3} dx$$

Optimal. Leaf size=163

$$-\frac{25a^2\sqrt[4]{1-iax}}{2\sqrt[4]{1+iax}} + \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}} - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} - \frac{25}{4}a^2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{25}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] $-25/2*a^2*(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}+5/4*I*a*(1-I*a*x)^{(5/4)}/x/(1+I*a*x)^{(1/4)}-1/2*(1-I*a*x)^{(9/4)}/x^2/(1+I*a*x)^{(1/4)}-25/4*a^2*\operatorname{arctan}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})+25/4*a^2*\operatorname{arctanh}((1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)})$

Rubi [A]

time = 0.04, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5170, 98, 96, 95, 304, 209, 212}

$$-\frac{25}{4}a^2 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{25a^2\sqrt[4]{1-iax}}{2\sqrt[4]{1+iax}} + \frac{25}{4}a^2 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} + \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(((5*I)/2)*ArcTan[a*x])*x^3),x]`

[Out] $(-25*a^2*(1 - I*a*x)^{(1/4)})/(2*(1 + I*a*x)^{(1/4)}) + (((5*I)/4)*a*(1 - I*a*x)^{(5/4)})/(x*(1 + I*a*x)^{(1/4)}) - (1 - I*a*x)^{(9/4)}/(2*x^2*(1 + I*a*x)^{(1/4)}) - (25*a^2*\operatorname{ArcTan}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}])/4 + (25*a^2*\operatorname{ArcTanh}[(1 + I*a*x)^{(1/4)}/(1 - I*a*x)^{(1/4)}])/4$

Rule 95

`Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

Rule 96

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

Rule 98

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])

```

Rule 209

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

```

Rule 304

```

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{5}{2}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1-iax)^{5/4}}{x^3(1+iax)^{5/4}} dx \\
&= -\frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} - \frac{1}{4}(5ia) \int \frac{(1-iax)^{5/4}}{x^2(1+iax)^{5/4}} dx \\
&= \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}} - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} - \frac{1}{8}(25a^2) \int \frac{\sqrt[4]{1-iax}}{x(1+iax)^{5/4}} dx \\
&= -\frac{25a^2\sqrt[4]{1-iax}}{2\sqrt[4]{1+iax}} + \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}} - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} - \frac{1}{8}(25a^2) \int \frac{1}{x(1-iax)^{3/4}\sqrt[4]{1+iax}} dx \\
&= -\frac{25a^2\sqrt[4]{1-iax}}{2\sqrt[4]{1+iax}} + \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}} - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} - \frac{1}{2}(25a^2) \text{Subst} \left(\int \frac{x^2}{-1+x^4} dx, x \right) \\
&= -\frac{25a^2\sqrt[4]{1-iax}}{2\sqrt[4]{1+iax}} + \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}} - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} + \frac{1}{4}(25a^2) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x \right) \\
&= -\frac{25a^2\sqrt[4]{1-iax}}{2\sqrt[4]{1+iax}} + \frac{5ia(1-iax)^{5/4}}{4x\sqrt[4]{1+iax}} - \frac{(1-iax)^{9/4}}{2x^2\sqrt[4]{1+iax}} - \frac{25}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{25}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right) + \frac{25}{4}a^2 \tan^{-1} \left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 81, normalized size = 0.50

$$\frac{\sqrt[4]{1-iax} (-2 + 9iax - 43a^2x^2 + 50a^2x^2 {}_2F_1(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}))}{4x^2\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((5*I)/2)*ArcTan[a*x])*x^3), x]

[Out] ((1 - I*a*x)^(1/4)*(-2 + (9*I)*a*x - 43*a^2*x^2 + 50*a^2*x^2*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(4*x^2*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}} x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^3, x)

[Out] $\int \frac{1}{((1+I*a*x)/(a^2*x^2+1)^{(1/2)})^{(5/2)}/x^3, x}$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/((1+I*a*x)/(a^2*x^2+1)^{(1/2)})^{(5/2)}/x^3, x, \text{algorithm}="maxima")$

[Out] $\text{integrate}(1/(x^3*((I*a*x + 1)/\text{sqrt}(a^2*x^2 + 1))^{(5/2)}), x)$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 238 vs. $2(111) = 222$.

time = 2.26, size = 238, normalized size = 1.46

$$\frac{2\sqrt{a^2x^2+1}(-43ia^2x^2-9ax-2i)\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-25(a^2x^3-ia^2x^2)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+1\right)+25(i a^2x^3+a^2x^2)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}+i\right)+25(-ia^2x^3-a^2x^2)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-i\right)+25(a^2x^3-ia^2x^2)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}}-1\right)}{8(ax^3-ix^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/((1+I*a*x)/(a^2*x^2+1)^{(1/2)})^{(5/2)}/x^3, x, \text{algorithm}="fricas")$

[Out] $-1/8*(2*\text{sqrt}(a^2*x^2 + 1)*(-43*I*a^2*x^2 - 9*a*x - 2*I)*\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) - 25*(a^3*x^3 - I*a^2*x^2)*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) + 1) + 25*(I*a^3*x^3 + a^2*x^2)*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) + I) + 25*(-I*a^3*x^3 - a^2*x^2)*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) - I) + 25*(a^3*x^3 - I*a^2*x^2)*\log(\text{sqrt}(I*\text{sqrt}(a^2*x^2 + 1)/(a*x + I)) - 1))/(a*x^3 - I*x^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^3 \left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}} \right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{integrate}(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**3, x)$

[Out] $\text{Integral}(1/(x**3*(I*(a*x - I)/\text{sqrt}(a**2*x**2 + 1))**(5/2)), x)$

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^3,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for
 the root of a polynomial with parameters. This might be wrong.The choice wa
 s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1}{x^3 \left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)),x)

[Out] int(1/(x^3*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)), x)

$$3.113 \quad \int \frac{e^{-\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^4} dx$$

Optimal. Leaf size=203

$$\frac{287ia^3\sqrt[4]{1-iax}}{24\sqrt[4]{1+iax}} - \frac{\sqrt[4]{1-iax}}{3x^3\sqrt[4]{1+iax}} + \frac{13ia^4\sqrt[4]{1-iax}}{12x^2\sqrt[4]{1+iax}} + \frac{61a^2\sqrt[4]{1-iax}}{24x\sqrt[4]{1+iax}} + \frac{55}{8}ia^3\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{55}{8}ia^3 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)$$

[Out] 287/24*I*a^3*(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-1/3*(1-I*a*x)^(1/4)/x^3/(1+I*a*x)^(1/4)+13/12*I*a*(1-I*a*x)^(1/4)/x^2/(1+I*a*x)^(1/4)+61/24*a^2*(1-I*a*x)^(1/4)/x/(1+I*a*x)^(1/4)+55/8*I*a^3*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-55/8*I*a^3*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))

Rubi [A]

time = 0.06, antiderivative size = 203, normalized size of antiderivative = 1.00, number of steps used = 10, number of rules used = 9, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {5170, 100, 156, 160, 12, 95, 304, 209, 212}

$$\frac{55}{8}ia^3\operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{287ia^3\sqrt[4]{1-iax}}{24\sqrt[4]{1+iax}} - \frac{55}{8}ia^3 \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{61a^2\sqrt[4]{1-iax}}{24x\sqrt[4]{1+iax}} - \frac{\sqrt[4]{1-iax}}{3x^3\sqrt[4]{1+iax}} + \frac{13ia^4\sqrt[4]{1-iax}}{12x^2\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(((5*I)/2)*ArcTan[a*x]))*x^4], x]

[Out] (((287*I)/24)*a^3*(1 - I*a*x)^(1/4))/(1 + I*a*x)^(1/4) - (1 - I*a*x)^(1/4)/(3*x^3*(1 + I*a*x)^(1/4)) + (((13*I)/12)*a*(1 - I*a*x)^(1/4))/(x^2*(1 + I*a*x)^(1/4)) + (61*a^2*(1 - I*a*x)^(1/4))/(24*x*(1 + I*a*x)^(1/4)) + ((55*I)/8)*a^3*ArcTan[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)] - ((55*I)/8)*a^3*ArcTanh[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)]

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)

```

*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m
+ 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(
n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*
(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a,
b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2
*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 160

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m + n + p + 2,
0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || (! (NeQ[n, -1] && SumSimplerQ[n,
1]) && !(NeQ[p, -1] && SumSimplerQ[p, 1])))

```

Rule 209

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

Rule 304

```

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x
] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a

```


/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-\frac{5}{2}i \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1 - iax)^{5/4}}{x^4(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} - \frac{1}{3} \int \frac{\frac{13ia}{2} + 6a^2x}{x^3(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{1}{6} \int \frac{-\frac{61a^2}{4} + 13ia^3x}{x^2(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{61a^2\sqrt[4]{1 - iax}}{24x\sqrt[4]{1 + iax}} - \frac{1}{6} \int \frac{-\frac{165ia^3}{8} - \frac{61a^4x}{4}}{x(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= \frac{287ia^3\sqrt[4]{1 - iax}}{24\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{61a^2\sqrt[4]{1 - iax}}{24x\sqrt[4]{1 + iax}} + \frac{i \int \frac{1}{16x(1 - iax)^{5/4}} dx}{16} \\
 &= \frac{287ia^3\sqrt[4]{1 - iax}}{24\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{61a^2\sqrt[4]{1 - iax}}{24x\sqrt[4]{1 + iax}} + \frac{1}{16} (55ia^3) S \\
 &= \frac{287ia^3\sqrt[4]{1 - iax}}{24\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{61a^2\sqrt[4]{1 - iax}}{24x\sqrt[4]{1 + iax}} + \frac{1}{4} (55ia^3) S \\
 &= \frac{287ia^3\sqrt[4]{1 - iax}}{24\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{61a^2\sqrt[4]{1 - iax}}{24x\sqrt[4]{1 + iax}} - \frac{1}{8} (55ia^3) S \\
 &= \frac{287ia^3\sqrt[4]{1 - iax}}{24\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{3x^3\sqrt[4]{1 + iax}} + \frac{13ia\sqrt[4]{1 - iax}}{12x^2\sqrt[4]{1 + iax}} + \frac{61a^2\sqrt[4]{1 - iax}}{24x\sqrt[4]{1 + iax}} + \frac{55}{8} ia^3 \tan^{-1}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 93, normalized size = 0.46

$$\frac{\sqrt[4]{1 - iax} \left(-8 + 26iax + 61a^2x^2 + 287ia^3x^3 - 330ia^3x^3 {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}\right) \right)}{24x^3\sqrt[4]{1 + iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((5*I)/2)*ArcTan[a*x])*x^4),x]

[Out] ((1 - I*a*x)^(1/4)*(-8 + (26*I)*a*x + 61*a^2*x^2 + (287*I)*a^3*x^3 - (330*I)*a^3*x^3*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)]))/(24*x^3*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}} x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x, algorithm="maxima")

[Out] integrate(1/(x^4*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)), x)

Fricas [A]

time = 4.81, size = 246, normalized size = 1.21

$$\frac{2(287a^3x^3 - 61ia^2x^2 + 26ax + 8i)\sqrt{a^2x^2+1}\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 165(i a^4x^4 + a^3x^3)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + 1\right) - 165(a^4x^4 - ia^3x^3)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} + i\right) + 165(a^4x^4 - ia^3x^3)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - i\right) - 165(-ia^4x^4 - a^3x^3)\log\left(\sqrt{\frac{i\sqrt{a^2x^2+1}}{ax+i}} - 1\right)}{48(ax^4 - i x^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x, algorithm="fricas")

[Out] 1/48*(2*(287*a^3*x^3 - 61*I*a^2*x^2 + 26*a*x + 8*I)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 165*(I*a^4*x^4 + a^3*x^3)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) - 165*(a^4*x^4 - I*a^3*x^3)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 165*(a^4*x^4 - I*a^3*x^3)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 165*(-I*a^4*x^4 - a^3*x^3)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1))/(a*x^4 - I*x^3)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**4,x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^4,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{1}{x^4 \left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^4*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)),x)
```

```
[Out] int(1/(x^4*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)), x)
```

$$3.114 \quad \int \frac{e^{-\frac{5}{2}i \operatorname{ArcTan}(ax)}}{x^5} dx$$

Optimal. Leaf size=233

$$\frac{2467a^4\sqrt[4]{1-iax}}{192\sqrt[4]{1+iax}} - \frac{\sqrt[4]{1-iax}}{4x^4\sqrt[4]{1+iax}} + \frac{17ia\sqrt[4]{1-iax}}{24x^3\sqrt[4]{1+iax}} + \frac{113a^2\sqrt[4]{1-iax}}{96x^2\sqrt[4]{1+iax}} - \frac{521ia^3\sqrt[4]{1-iax}}{192x\sqrt[4]{1+iax}} + \frac{475}{64}a^4 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)$$

[Out] 2467/192*a^4*(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-1/4*(1-I*a*x)^(1/4)/x^4/(1+I*a*x)^(1/4)+17/24*I*a*(1-I*a*x)^(1/4)/x^3/(1+I*a*x)^(1/4)+113/96*a^2*(1-I*a*x)^(1/4)/x^2/(1+I*a*x)^(1/4)-521/192*I*a^3*(1-I*a*x)^(1/4)/x/(1+I*a*x)^(1/4)+475/64*a^4*arctan((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))-475/64*a^4*arctanh((1+I*a*x)^(1/4)/(1-I*a*x)^(1/4))

Rubi [A]

time = 0.07, antiderivative size = 233, normalized size of antiderivative = 1.00, number of steps used = 11, number of rules used = 9, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.562$, Rules used = {5170, 100, 156, 160, 12, 95, 304, 209, 212}

$$\frac{475}{64}a^4 \operatorname{ArcTan}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) + \frac{2467a^4\sqrt[4]{1-iax}}{192\sqrt[4]{1+iax}} - \frac{475}{64}a^4 \tanh^{-1}\left(\frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right) - \frac{521ia^3\sqrt[4]{1-iax}}{192x\sqrt[4]{1+iax}} + \frac{113a^2\sqrt[4]{1-iax}}{96x^2\sqrt[4]{1+iax}} - \frac{\sqrt[4]{1-iax}}{4x^4\sqrt[4]{1+iax}} + \frac{17ia\sqrt[4]{1-iax}}{24x^3\sqrt[4]{1+iax}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(((5*I)/2)*ArcTan[a*x])*x^5), x]

[Out] (2467*a^4*(1 - I*a*x)^(1/4))/(192*(1 + I*a*x)^(1/4)) - (1 - I*a*x)^(1/4)/(4*x^4*(1 + I*a*x)^(1/4)) + (((17*I)/24)*a*(1 - I*a*x)^(1/4))/(x^3*(1 + I*a*x)^(1/4)) + (113*a^2*(1 - I*a*x)^(1/4))/(96*x^2*(1 + I*a*x)^(1/4)) - (((521*I)/192)*a^3*(1 - I*a*x)^(1/4))/(x*(1 + I*a*x)^(1/4)) + (475*a^4*ArcTan[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)])/64 - (475*a^4*ArcTanh[(1 + I*a*x)^(1/4)/(1 - I*a*x)^(1/4)])/64

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_], x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)
*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m
+ 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(
n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*
(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a,
b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2
*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)
*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 160

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)
*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m + n + p + 2,
0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || (!NeQ[n, -1] && SumSimplerQ[n,
1]) && !(NeQ[p, -1] && SumSimplerQ[p, 1]))

```

Rule 209

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

Rule 304

```

Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b,
2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x

```

] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-\frac{5}{2}i \tan^{-1}(ax)}}{x^5} dx &= \int \frac{(1 - iax)^{5/4}}{x^5(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} - \frac{1}{4} \int \frac{\frac{17ia}{2} + 8a^2x}{x^4(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{1}{12} \int \frac{-\frac{113a^2}{4} + \frac{51}{2}ia^3x}{x^3(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{1}{24} \int \frac{-\frac{521ia^3}{8} - \frac{113a^4x}{2}}{x^2(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= -\frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{521ia^3\sqrt[4]{1 - iax}}{192x\sqrt[4]{1 + iax}} + \frac{1}{24} \int \frac{1}{x(1 - iax)^{3/4}(1 + iax)^{5/4}} dx \\
 &= \frac{2467a^4\sqrt[4]{1 - iax}}{192\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{521ia^3\sqrt[4]{1 - iax}}{192x\sqrt[4]{1 + iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 - iax}}{192\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{521ia^3\sqrt[4]{1 - iax}}{192x\sqrt[4]{1 + iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 - iax}}{192\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{521ia^3\sqrt[4]{1 - iax}}{192x\sqrt[4]{1 + iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 - iax}}{192\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{521ia^3\sqrt[4]{1 - iax}}{192x\sqrt[4]{1 + iax}} \\
 &= \frac{2467a^4\sqrt[4]{1 - iax}}{192\sqrt[4]{1 + iax}} - \frac{\sqrt[4]{1 - iax}}{4x^4\sqrt[4]{1 + iax}} + \frac{17ia\sqrt[4]{1 - iax}}{24x^3\sqrt[4]{1 + iax}} + \frac{113a^2\sqrt[4]{1 - iax}}{96x^2\sqrt[4]{1 + iax}} - \frac{521ia^3\sqrt[4]{1 - iax}}{192x\sqrt[4]{1 + iax}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 99, normalized size = 0.42

$$\frac{\sqrt[4]{1 - iax} \left(-48 + 136iax + 226a^2x^2 - 521ia^3x^3 + 2467a^4x^4 - 2850a^4x^4 {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{i+ax}{i-ax}\right) \right)}{192x^4\sqrt[4]{1 + iax}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((5*I)/2)*ArcTan[a*x]))*x^5,x]

[Out] (((1 - I*a*x)^(1/4)*(-48 + (136*I)*a*x + 226*a^2*x^2 - (521*I)*a^3*x^3 + 2467*a^4*x^4 - 2850*a^4*x^4*Hypergeometric2F1[1/4, 1, 5/4, (I + a*x)/(I - a*x)])))/(192*x^4*(1 + I*a*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}} x^5} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x)

[Out] int(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x, algorithm="maxima")

[Out] integrate(1/(x^5*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2)), x)

Fricas [A]

time = 2.44, size = 254, normalized size = 1.09

$$\frac{2(2467i a^4 x^4 + 521 a^3 x^3 + 226i a^2 x^2 - 136 a x - 48) \sqrt{a^2 x^2 + 1} \sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{a x + i} + 1} + 1425 (a^2 x^2 - i a^4 x^4) \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{a x + i} + 1}\right) + 1425 (-i a^2 x^2 - a^4 x^4) \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{a x + i} + i}\right) + 1425 (i a^2 x^2 + a^4 x^4) \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{a x + i} - i}\right) - 1425 (a^2 x^2 - i a^4 x^4) \log\left(\sqrt{\frac{i \sqrt{a^2 x^2 + 1}}{a x + i} - 1}\right)}{384 (a x^2 - i x^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x, algorithm="fricas")

[Out] -1/384*(2*(2467*I*a^4*x^4 + 521*a^3*x^3 + 226*I*a^2*x^2 - 136*a*x - 48*I)*sqrt(a^2*x^2 + 1)*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1425*(a^5*x^5 - I*a^4*x^4)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + 1) + 1425*(-I*a^5*x^5 - a^4*x^4)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) + I) + 1425*(I*a^5*x^5 + a^4*x^4)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - I) - 1425*(a^5*x^5 - I*a^4*x^4)*log(sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)) - 1))/(a*x^5 - I*x^4)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)/x**5,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3063 deep
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)/x^5,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \frac{1}{x^5 \left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}} \right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^5*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)),x)
```

```
[Out] int(1/(x^5*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2)), x)
```


3.115 $\int e^{\frac{1}{3}i\text{ArcTan}(x)} x^2 dx$

Optimal. Leaf size=319

$$-\frac{19}{54}i(1-ix)^{5/6}\sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6}x + \frac{19}{162}i\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right)$$

[Out] -19/54*I*(1-I*x)^(5/6)*(1+I*x)^(1/6)-1/18*I*(1-I*x)^(5/6)*(1+I*x)^(7/6)+1/3*(1-I*x)^(5/6)*(1+I*x)^(7/6)*x-19/81*I*arctan((1-I*x)^(1/6)/(1+I*x)^(1/6))-19/162*I*arctan(2*(1-I*x)^(1/6)/(1+I*x)^(1/6)-3^(1/2))-19/162*I*arctan(2*(1-I*x)^(1/6)/(1+I*x)^(1/6)+3^(1/2))-19/324*I*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3))-(1-I*x)^(1/6)*3^(1/2)/(1+I*x)^(1/6))*3^(1/2)+19/324*I*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3)+(1-I*x)^(1/6)*3^(1/2)/(1+I*x)^(1/6))*3^(1/2)

Rubi [A]

time = 0.29, antiderivative size = 319, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 12, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {5170, 92, 81, 52, 65, 338, 301, 648, 632, 210, 642, 209}

$$\frac{19}{162}\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) - \frac{19}{162}i\text{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) - \frac{19}{81}i\text{ArcTan}\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{3}(1-ix)^{5/6}x(1+ix)^{7/6} - \frac{1}{18}(1-ix)^{5/6}(1+ix)^{7/6} - \frac{19}{54}(1-ix)^{5/6}\sqrt[6]{1+ix} - \frac{19i\log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} - \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right)}{108\sqrt{3}} + \frac{19i\log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right)}{108\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[E^((I/3)*ArcTan[x])*x^2,x]

[Out] ((-19*I)/54)*(1 - I*x)^(5/6)*(1 + I*x)^(1/6) - (I/18)*(1 - I*x)^(5/6)*(1 + I*x)^(7/6) + ((1 - I*x)^(5/6)*(1 + I*x)^(7/6)*x)/3 + ((19*I)/162)*ArcTan[Sqrt[3] - (2*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)] - ((19*I)/162)*ArcTan[Sqrt[3] + (2*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)] - ((19*I)/81)*ArcTan[(1 - I*x)^(1/6)/(1 + I*x)^(1/6)] - (((19*I)/108)*Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/3) - (Sqrt[3]*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)])/Sqrt[3] + (((19*I)/108)*Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/3) + (Sqrt[3]*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)])/Sqrt[3]

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +

```
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 92

```
Int[((a_.) + (b_.)*(x_))^(2)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(
p_.), x_Symbol] :> Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(
d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)
^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b
*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ
[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 301

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] :> Module[{r = Numerator
[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*cos[(2*k
- 1)*m*(Pi/n)] - s*cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*cos[(2*k
- 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*cos[(2*k - 1)*m*(Pi/n)] + s*cos[(2*k
- 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]
; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^(
m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x]] /; FreeQ[{a, b}, x]
&& IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{3}i \tan^{-1}(x)} x^2 dx &= \int \frac{\sqrt[6]{1+ix} x^2}{\sqrt[6]{1-ix}} dx \\
&= \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x + \frac{1}{3} \int \frac{\left(-1 - \frac{ix}{3}\right) \sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} dx \\
&= -\frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{54} \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} dx \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{162}i \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{27}i \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{27}i \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{81}i \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{81}i \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x - \frac{19}{81}i \\
&= -\frac{19}{54}i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{18}i(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3}(1-ix)^{5/6}(1+ix)^{7/6} x + \frac{19}{162}i
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 73, normalized size = 0.23

$$\frac{1}{90}(1-ix)^{5/6} \left(5\sqrt[6]{1+ix} (-i+7x+6ix^2) - 38i\sqrt[6]{2} {}_2F_1\left(-\frac{1}{6}, \frac{5}{6}; \frac{11}{6}; \frac{1}{2} - \frac{ix}{2}\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x])*x^2,x]

[Out] ((1 - I*x)^(5/6)*(5*(1 + I*x)^(1/6)*(-I + 7*x + (6*I)*x^2) - (38*I)*2^(1/6)*Hypergeometric2F1[-1/6, 5/6, 11/6, 1/2 - (I/2)*x]))/90

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{ix + 1}{\sqrt{x^2 + 1}} \right)^{\frac{1}{3}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x^2,x)**[Out]** int(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x^2,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x^2,x, algorithm="maxima")**[Out]** integrate(x^2*((I*x + 1)/sqrt(x^2 + 1))^(1/3), x)**Fricas [A]**

time = 3.19, size = 208, normalized size = 0.65

$$\frac{19}{384}(-i\sqrt{x+1})\log\left(\frac{1}{2}\sqrt{x+1} + \frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right) - \frac{19}{384}(-i\sqrt{x-1})\log\left(\frac{1}{2}\sqrt{x-1} + \frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right) - \frac{19}{384}(i\sqrt{x+1})\log\left(\frac{1}{2}\sqrt{x+1} + \frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right) - \frac{19}{384}(i\sqrt{x-1})\log\left(\frac{1}{2}\sqrt{x-1} + \frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right) + \frac{1}{54}(18x^3 - 3ix^2 - x - 22i)\left(\frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right) + \frac{19}{162}\log\left(\frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right) + \frac{19}{162}\log\left(\frac{(\sqrt{x^2+1})^{1/3}}{x+1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x^2,x, algorithm="fricas")

[Out] -19/324*(-I*sqrt(3) + 1)*log(1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2*I) - 19/324*(-I*sqrt(3) - 1)*log(1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2*I) - 19/324*(I*sqrt(3) + 1)*log(-1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2*I) - 19/324*(I*sqrt(3) - 1)*log(-1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2*I) + 1/54*(18*x^3 - 3*I*x^2 - x - 22*I) * (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 19/162*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) + I) + 19/162*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) - I)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \sqrt[3]{\frac{i(x-i)}{\sqrt{x^2+1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3)*x**2,x)

[Out] Integral(x**2*(I*(x - I)/sqrt(x**2 + 1))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x^2,x, algorithm="giac")

[Out] integrate(x^2*((I*x + 1)/sqrt(x^2 + 1))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \left(\frac{1 + x i}{\sqrt{x^2 + 1}} \right)^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3),x)

[Out] int(x^2*((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3), x)

3.116 $\int e^{\frac{1}{3}i \operatorname{ArcTan}(x)} x dx$

Optimal. Leaf size=278

$$\frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} - \frac{1}{18}\operatorname{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{18}\operatorname{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right)$$

[Out] 1/6*(1-I*x)^(5/6)*(1+I*x)^(1/6)+1/2*(1-I*x)^(5/6)*(1+I*x)^(7/6)+1/9*arctan((1-I*x)^(1/6)/(1+I*x)^(1/6))+1/18*arctan(2*(1-I*x)^(1/6)/(1+I*x)^(1/6)-3^(1/2))+1/18*arctan(2*(1-I*x)^(1/6)/(1+I*x)^(1/6)+3^(1/2))+1/36*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3)-(1-I*x)^(1/6)*3^(1/2)/(1+I*x)^(1/6))*3^(1/2)-1/36*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3)+(1-I*x)^(1/6)*3^(1/2)/(1+I*x)^(1/6))*3^(1/2)

Rubi [A]

time = 0.26, antiderivative size = 278, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 11, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {5170, 81, 52, 65, 338, 301, 648, 632, 210, 642, 209}

$$-\frac{1}{18}\operatorname{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{18}\operatorname{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{9}\operatorname{ArcTan}\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{\log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} - \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right)}{12\sqrt{3}} - \frac{\log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right)}{12\sqrt{3}}$$

Antiderivative was successfully verified.

[In] Int[E^((I/3)*ArcTan[x])*x,x]

[Out] ((1 - I*x)^(5/6)*(1 + I*x)^(1/6))/6 + ((1 - I*x)^(5/6)*(1 + I*x)^(7/6))/2 - ArcTan[Sqrt[3] - (2*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)]/18 + ArcTan[Sqrt[3] + (2*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)]/18 + ArcTan[(1 - I*x)^(1/6)/(1 + I*x)^(1/6)]/9 + Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/3) - (Sqrt[3]*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)]/(12*Sqrt[3]) - Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/6) + (Sqrt[3]*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)]/(12*Sqrt[3])

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 301

Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := Module[{r = Numerator[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*cos[(2*k - 1)*m*(Pi/n)] - s*cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*cos[(2*k - 1)*m*(Pi/n)] + s*cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]

Rule 338

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]

Rule 632

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{3}i \tan^{-1}(x)} x dx &= \int \frac{\sqrt[6]{1+ix} x}{\sqrt[6]{1-ix}} dx \\
&= \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} - \frac{1}{6}i \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} dx \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} - \frac{1}{18}i \int \frac{1}{\sqrt[6]{1-ix}(1+ix)^{5/6}} dx \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3} \text{Subst} \left(\int \frac{x^4}{(2-x^6)^{5/6}} dx, x, \sqrt[6]{1-ix} \right) \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{3} \text{Subst} \left(\int \frac{x^4}{1+x^6} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{9} \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{9} \tan^{-1} \left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{1}{36} \text{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} + \frac{1}{9} \tan^{-1} \left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{\log \left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}} \right)}{36} \\
&= \frac{1}{6}(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{1}{2}(1-ix)^{5/6}(1+ix)^{7/6} - \frac{1}{18} \tan^{-1} \left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{1}{18} \log \left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 57, normalized size = 0.21

$$\frac{1}{10}(1-ix)^{5/6} \left(5(1+ix)^{7/6} + 2\sqrt[6]{2} {}_2F_1 \left(-\frac{1}{6}, \frac{5}{6}; \frac{11}{6}; \frac{1}{2} - \frac{ix}{2} \right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x])*x,x]

[Out] ((1 - I*x)^(5/6)*(5*(1 + I*x)^(7/6) + 2*2^(1/6)*Hypergeometric2F1[-1/6, 5/6, 11/6, 1/2 - (I/2)*x]))/10

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{ix+1}{\sqrt{x^2+1}} \right)^{\frac{1}{3}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x,x)`

[Out] `int(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x,x, algorithm="maxima")`

[Out] `integrate(x*((I*x + 1)/sqrt(x^2 + 1))^(1/3), x)`

Fricas [A]

time = 1.70, size = 195, normalized size = 0.70

$$\frac{1}{36}(\sqrt{3}+i)\log\left(\frac{1}{2}\sqrt{3}+\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+\frac{1}{36}(\sqrt{3}-i)\log\left(\frac{1}{2}\sqrt{3}+\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}-\frac{1}{36}(\sqrt{3}-i)\log\left(-\frac{1}{2}\sqrt{3}+\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+\frac{1}{36}(\sqrt{3}+i)\log\left(-\frac{1}{2}\sqrt{3}+\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+\frac{1}{6}(3x^2-ix+4)\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}-\frac{1}{18}i\log\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+\frac{1}{18}i\log\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x,x, algorithm="fricas")`

[Out] `-1/36*(sqrt(3) + I)*log(1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2 *I) - 1/36*(sqrt(3) - I)*log(1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2*I) + 1/36*(sqrt(3) - I)*log(-1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2*I) + 1/36*(sqrt(3) + I)*log(-1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2*I) + 1/6*(3*x^2 - I*x + 4)*(I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/18*I*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) + I) + 1/18*I*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) - I)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 \sqrt[3]{\frac{i(x-i)}{\sqrt{x^2+1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3)*x,x)`

[Out] `Integral(x*(I*(x - I)/sqrt(x**2 + 1))**(1/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)*x,x, algorithm="giac")

[Out] integrate(x*((I*x + 1)/sqrt(x^2 + 1))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \left(\frac{1 + x \text{I}}{\sqrt{x^2 + 1}} \right)^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*((x*I + 1)/(x^2 + 1)^(1/2))^(1/3),x)

[Out] int(x*((x*I + 1)/(x^2 + 1)^(1/2))^(1/3), x)

3.117 $\int e^{\frac{1}{3}i\text{ArcTan}(x)} dx$

Optimal. Leaf size=262

$$i(1-ix)^{5/6}\sqrt[6]{1+ix} - \frac{1}{3}i\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{3}i\text{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{2}{3}i\text{ArcTan}\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right)$$

[Out] $I*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)}+2/3*I*\arctan((1-I*x)^{(1/6)}/(1+I*x)^{(1/6)})+1/3*I*\arctan(2*(1-I*x)^{(1/6)}/(1+I*x)^{(1/6)}-3^{(1/2)})+1/3*I*\arctan(2*(1-I*x)^{(1/6)}/(1+I*x)^{(1/6)}+3^{(1/2)})+1/6*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}-(1-I*x)^{(1/6)}*3^{(1/2)}/(1+I*x)^{(1/6)})+1/6*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}+(1-I*x)^{(1/6)}*3^{(1/2)}/(1+I*x)^{(1/6)})+1/6*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}-(1-I*x)^{(1/6)}*3^{(1/2)}/(1+I*x)^{(1/6)})+1/6*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}+(1-I*x)^{(1/6)}*3^{(1/2)}/(1+I*x)^{(1/6)})+1/6*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}-(1-I*x)^{(1/6)}*3^{(1/2)}/(1+I*x)^{(1/6)})+1/6*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}+(1-I*x)^{(1/6)}*3^{(1/2)}/(1+I*x)^{(1/6)})$

Rubi [A]

time = 0.25, antiderivative size = 262, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 10, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 1.000$, Rules used = {5169, 52, 65, 338, 301, 648, 632, 210, 642, 209}

$$-\frac{1}{3}i\text{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{1}{3}i\text{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \frac{2}{3}i\text{ArcTan}\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + i(1-ix)^{5/6}\sqrt[6]{1+ix} + \frac{i\log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} - \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right)}{2\sqrt{3}} - \frac{i\log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right)}{2\sqrt{3}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I/3)*\text{ArcTan}[x]}, x]$

[Out] $I*(1 - I*x)^{(5/6)}*(1 + I*x)^{(1/6)} - (I/3)*\text{ArcTan}[\text{Sqrt}[3] - (2*(1 - I*x)^{(1/6)})/(1 + I*x)^{(1/6)}] + (I/3)*\text{ArcTan}[\text{Sqrt}[3] + (2*(1 - I*x)^{(1/6)})/(1 + I*x)^{(1/6)}] + ((2*I)/3)*\text{ArcTan}[(1 - I*x)^{(1/6)}/(1 + I*x)^{(1/6)}] + ((I/2)*\text{Log}[1 + (1 - I*x)^{(1/3)}/(1 + I*x)^{(1/3)} - (\text{Sqrt}[3]*(1 - I*x)^{(1/6)})/(1 + I*x)^{(1/6)}])/(\text{Sqrt}[3]) - ((I/2)*\text{Log}[1 + (1 - I*x)^{(1/3)}/(1 + I*x)^{(1/3)} + (\text{Sqrt}[3]*(1 - I*x)^{(1/6)})/(1 + I*x)^{(1/6)}])/(\text{Sqrt}[3])$

Rule 52

$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] := \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^n / (b*(m+n+1)), x] + \text{Dist}[n * (b*c - a*d) / (b*(m+n+1)), \text{Int}[(a + b*x)^m * (c + d*x)^{n-1}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m+n+1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m-n, 0]))) \ \&\& \ !\text{ILtQ}[m+n+2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] := \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{p*(m+1)-1} * (c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^{(1/p)}], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}$

$[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 209

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(1/(\text{Rt}[a, 2]*\text{Rt}[b, 2]))*\text{ArcTan}[\text{Rt}[b, 2]*(x/\text{Rt}[a, 2])], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{PosQ}[a/b] \&\& (\text{GtQ}[a, 0] \parallel \text{GtQ}[b, 0])$

Rule 210

$\text{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \parallel \text{LtQ}[b, 0])$

Rule 301

$\text{Int}(x_)^{(m_)} / ((a_ + (b_)*(x_)^n))^{(p_)}, x_Symbol] \rightarrow \text{Module}\{r = \text{Numerator}[\text{Rt}[a/b, n]], s = \text{Denominator}[\text{Rt}[a/b, n]], k, u\}, \text{Simp}[u = \text{Int}[(r*\text{Cos}[(2*k - 1)*m*(\text{Pi}/n)] - s*\text{Cos}[(2*k - 1)*(m + 1)*(\text{Pi}/n)]*x)/(r^2 - 2*r*s*\text{Cos}[(2*k - 1)*(\text{Pi}/n)]*x + s^2*x^2), x] + \text{Int}[(r*\text{Cos}[(2*k - 1)*m*(\text{Pi}/n)] + s*\text{Cos}[(2*k - 1)*(m + 1)*(\text{Pi}/n)]*x)/(r^2 + 2*r*s*\text{Cos}[(2*k - 1)*(\text{Pi}/n)]*x + s^2*x^2), x]; 2*(-1)^{(m/2)}*(r^{(m + 2)} / (a*n*s^m))*\text{Int}[1/(r^2 + s^2*x^2), x] + \text{Dist}[2*(r^{(m + 1)} / (a*n*s^m)), \text{Sum}[u, \{k, 1, (n - 2)/4\}], x], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{IGtQ}[(n - 2)/4, 0] \&\& \text{IGtQ}[m, 0] \&\& \text{LtQ}[m, n - 1] \&\& \text{PosQ}[a/b]$

Rule 338

$\text{Int}(x_)^{(m_)} * ((a_ + (b_)*(x_)^n))^{(p_)}, x_Symbol] \rightarrow \text{Dist}[a^{(p + (m + 1)/n)}, \text{Subst}[\text{Int}[x^m / (1 - b*x^n)^{(p + (m + 1)/n + 1)}, x], x, x/(a + b*x^n)^{(1/n)}], x] /; \text{FreeQ}\{a, b, x\} \&\& \text{IGtQ}[n, 0] \&\& \text{LtQ}[-1, p, 0] \&\& \text{NeQ}[p, -2^{(-1)}] \&\& \text{IntegersQ}[m, p + (m + 1)/n]$

Rule 632

$\text{Int}(((a_.) + (b_)*(x_) + (c_)*(x_)^2)^{-1}), x_Symbol] \rightarrow \text{Dist}[-2, \text{Subst}[\text{Int}[1/\text{Simp}[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; \text{FreeQ}\{a, b, c, x\} \&\& \text{NeQ}[b^2 - 4*a*c, 0]$

Rule 642

$\text{Int}(((d_.) + (e_)*(x_)) / ((a_.) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] \rightarrow \text{Simp}[d*(\text{Log}[\text{RemoveContent}[a + b*x + c*x^2, x]]/b), x] /; \text{FreeQ}\{a, b, c, d, e, x\} \&\& \text{EqQ}[2*c*d - b*e, 0]$

Rule 648

```
Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 5169

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{3}i \tan^{-1}(x)} dx &= \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} dx \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} + \frac{1}{3} \int \frac{1}{\sqrt[6]{1-ix} (1+ix)^{5/6}} dx \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} + 2i \operatorname{Subst} \left(\int \frac{x^4}{(2-x^6)^{5/6}} dx, x, \sqrt[6]{1-ix} \right) \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} + 2i \operatorname{Subst} \left(\int \frac{x^4}{1+x^6} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} + \frac{2}{3} i \operatorname{Subst} \left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{2}{3} i \operatorname{Subst} \left(\int \frac{-\frac{1}{2} + \frac{\sqrt{3}}{2}}{1 - \sqrt{3}x} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} + \frac{2}{3} i \tan^{-1} \left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{1}{6} i \operatorname{Subst} \left(\int \frac{1}{1 - \sqrt{3}x + x^2} dx, x, \frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} + \frac{2}{3} i \tan^{-1} \left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{i \log \left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}} - \frac{\sqrt{3} \sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right)}{2\sqrt{3}} \\
&= i(1-ix)^{5/6} \sqrt[6]{1+ix} - \frac{1}{3} i \tan^{-1} \left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \frac{1}{3} i \tan^{-1} \left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 34, normalized size = 0.13

$$-\frac{12}{7} i e^{\frac{7}{3} i \operatorname{ArcTan}(x)} {}_2F_1 \left(\frac{7}{6}, 2; \frac{13}{6}; -e^{2i \operatorname{ArcTan}(x)} \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x]),x]

[Out] ((-12*I)/7)*E^(((7*I)/3)*ArcTan[x])*Hypergeometric2F1[7/6, 2, 13/6, -E^((2*I)*ArcTan[x])]

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{ix + 1}{\sqrt{x^2 + 1}} \right)^{\frac{1}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(1/3),x)

[Out] int(((1+I*x)/(x^2+1)^(1/2))^(1/3),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3),x, algorithm="maxima")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3), x)

Fricas [A]

time = 2.57, size = 195, normalized size = 0.74

$$\frac{1}{6}(-i\sqrt{3}+1)\log\left(\frac{1}{2}\sqrt{3}+\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}+\frac{1}{2}\right)+\frac{1}{6}(-i\sqrt{3}-1)\log\left(\frac{1}{2}\sqrt{3}+\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}-\frac{1}{2}\right)+\frac{1}{6}(i\sqrt{3}+1)\log\left(-\frac{1}{2}\sqrt{3}+\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}+\frac{1}{2}\right)+\frac{1}{6}(i\sqrt{3}-1)\log\left(-\frac{1}{2}\sqrt{3}+\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}-\frac{1}{2}\right)+(x+i)\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}+\frac{1}{3}\log\left(\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}+i\right)-\frac{1}{3}\log\left(\left(\frac{i\sqrt{3}+1}{x+i}\right)^{\frac{1}{3}}-i\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3),x, algorithm="fricas")

[Out] 1/6*(-I*sqrt(3) + 1)*log(1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2*I) + 1/6*(-I*sqrt(3) - 1)*log(1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2*I) + 1/6*(I*sqrt(3) + 1)*log(-1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2*I) + 1/6*(I*sqrt(3) - 1)*log(-1/2*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2*I) + (x + I)*(I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/3*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) + I) - 1/3*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) - I)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{\frac{ix + 1}{\sqrt{x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3),x)

[Out] Integral(((I*x + 1)/sqrt(x**2 + 1))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3),x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \left(\frac{1 + x \text{li}}{\sqrt{x^2 + 1}} \right)^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3),x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3), x)

$$3.118 \quad \int \frac{e^{\frac{1}{3}i \operatorname{ArcTan}(x)}}{x} dx$$

Optimal. Leaf size=430

$$\operatorname{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) - \operatorname{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \sqrt{3} \operatorname{ArcTan}\left(\frac{1 - \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}}\right) - \sqrt{3} \operatorname{ArcTan}\left(\frac{1 + \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}}\right)$$

[Out] -2*arctan((1-I*x)^(1/6)/(1+I*x)^(1/6))-arctan(2*(1-I*x)^(1/6)/(1+I*x)^(1/6)-3^(1/2))-arctan(2*(1-I*x)^(1/6)/(1+I*x)^(1/6)+3^(1/2))-2*arctanh((1+I*x)^(1/6)/(1-I*x)^(1/6))+1/2*ln(1-(1+I*x)^(1/6)/(1-I*x)^(1/6)+(1+I*x)^(1/3)/(1-I*x)^(1/3))-1/2*ln(1+(1+I*x)^(1/6)/(1-I*x)^(1/6)+(1+I*x)^(1/3)/(1-I*x)^(1/3))+arctan(1/3*(1-2*(1+I*x)^(1/6)/(1-I*x)^(1/6))*3^(1/2))*3^(1/2)-arctan(1/3*(1+2*(1+I*x)^(1/6)/(1-I*x)^(1/6))*3^(1/2))*3^(1/2)-1/2*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3)-(1-I*x)^(1/6)*3^(1/2)/(1+I*x)^(1/6))*3^(1/2)+1/2*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3)+(1-I*x)^(1/6)*3^(1/2)/(1+I*x)^(1/6))*3^(1/2)

Rubi [A]

time = 0.36, antiderivative size = 430, normalized size of antiderivative = 1.00, number of steps used = 25, number of rules used = 13, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.929$, Rules used = {5170, 132, 65, 338, 301, 648, 632, 210, 642, 209, 95, 216, 212}

$$\operatorname{ArcTan}\left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) - \operatorname{ArcTan}\left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) + \sqrt{3} \operatorname{ArcTan}\left(\frac{1 - \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}}\right) - \sqrt{3} \operatorname{ArcTan}\left(\frac{1 + \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}}\right) - 2 \operatorname{ArcTan}\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}}\right) - \frac{1}{2} \sqrt{3} \log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} - \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right) + \frac{1}{2} \sqrt{3} \log\left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + \frac{\sqrt{3}\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} + 1\right) + \frac{1}{2} \log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) - \frac{1}{2} \log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) - 2 \operatorname{tanh}^{-1}\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)$$

Antiderivative was successfully verified.

[In] Int[E^((I/3)*ArcTan[x])/x,x]

[Out] ArcTan[Sqrt[3] - (2*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)] - ArcTan[Sqrt[3] + (2*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)] + Sqrt[3]*ArcTan[(1 - (2*(1 + I*x)^(1/6)))/(1 - I*x)^(1/6)]/Sqrt[3] - Sqrt[3]*ArcTan[(1 + (2*(1 + I*x)^(1/6)))/(1 - I*x)^(1/6)]/Sqrt[3] - 2*ArcTan[(1 - I*x)^(1/6)/(1 + I*x)^(1/6)] - 2*ArcTanh[(1 + I*x)^(1/6)/(1 - I*x)^(1/6)] - (Sqrt[3]*Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/3) - (Sqrt[3]*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)])/2 + (Sqrt[3]*Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/3) + (Sqrt[3]*(1 - I*x)^(1/6))/(1 + I*x)^(1/6)])/2 + Log[1 - (1 + I*x)^(1/6)/(1 - I*x)^(1/6) + (1 + I*x)^(1/3)/(1 - I*x)^(1/3)]/2 - Log[1 + (1 + I*x)^(1/6)/(1 - I*x)^(1/6) + (1 + I*x)^(1/3)/(1 - I*x)^(1/3)]/2

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den

ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 95

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 132

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 216

Int[((a_) + (b_.)*(x_)^(n_))^(n_)*((c_.) + (d_.)*(x_))^(n_)/((e_.) + (f_.)*(x_)), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n], k, u], Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n))*Int[1/(r^2 - s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x], x] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]

Rule 301

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_.)), x_Symbol] := Module[{r = Numerator
[Rt[a/b, n]], s = Denominator[Rt[a/b, n]], k, u}, Simp[u = Int[(r*Cos[(2*k
- 1)*m*(Pi/n)] - s*Cos[(2*k - 1)*(m + 1)*(Pi/n)]*x)/(r^2 - 2*r*s*Cos[(2*k -
1)*(Pi/n)]*x + s^2*x^2), x] + Int[(r*Cos[(2*k - 1)*m*(Pi/n)] + s*Cos[(2*k
- 1)*(m + 1)*(Pi/n)]*x)/(r^2 + 2*r*s*Cos[(2*k - 1)*(Pi/n)]*x + s^2*x^2), x]
; 2*(-1)^(m/2)*(r^(m + 2)/(a*n*s^m))*Int[1/(r^2 + s^2*x^2), x] + Dist[2*(r^
(m + 1)/(a*n*s^m)), Sum[u, {k, 1, (n - 2)/4}], x], x] /; FreeQ[{a, b}, x]
&& IGtQ[(n - 2)/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && PosQ[a/b]
```

Rule 338

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_.))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_.) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{3}i \tan^{-1}(x)}}{x} dx &= \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix} x} dx \\
&= i \int \frac{1}{\sqrt[6]{1-ix} (1+ix)^{5/6}} dx + \int \frac{1}{\sqrt[6]{1-ix} (1+ix)^{5/6} x} dx \\
&= - \left(6 \text{Subst} \left(\int \frac{x^4}{(2-x^6)^{5/6}} dx, x, \sqrt[6]{1-ix} \right) \right) + 6 \text{Subst} \left(\int \frac{1}{-1+x^6} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= - \left(2 \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \right) - 2 \text{Subst} \left(\int \frac{1-\frac{x}{2}}{1-x+x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -2 \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) + \frac{1}{2} \text{Subst} \left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) - \frac{1}{2} \text{Subst} \left(\int \frac{1}{1+x} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -2 \tan^{-1} \left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) - 2 \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) + \frac{1}{2} \log \left(1 - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix}} \right) \\
&= \sqrt{3} \tan^{-1} \left(\frac{1 - \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right) - \sqrt{3} \tan^{-1} \left(\frac{1 + \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right) - 2 \tan^{-1} \left(\frac{\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) \\
&= \tan^{-1} \left(\sqrt{3} - \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) - \tan^{-1} \left(\sqrt{3} + \frac{2\sqrt[6]{1-ix}}{\sqrt[6]{1+ix}} \right) + \sqrt{3} \tan^{-1} \left(\frac{1 - \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 90, normalized size = 0.21

$$\frac{3(1-ix)^{5/6} \left(\sqrt[6]{2} (1+ix)^{5/6} {}_2F_1 \left(\frac{5}{6}, \frac{5}{6}, \frac{11}{6}; \frac{1}{2} - \frac{ix}{2} \right) + 2 {}_2F_1 \left(\frac{5}{6}, 1, \frac{11}{6}; \frac{i+x}{i-x} \right) \right)}{5(1+ix)^{5/6}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x])/x,x]

[Out] (-3*(1 - I*x)^(5/6)*(2^(1/6)*(1 + I*x)^(5/6)*Hypergeometric2F1[5/6, 5/6, 11/6, 1/2 - (I/2)*x] + 2*Hypergeometric2F1[5/6, 1, 11/6, (I + x)/(I - x)])/(5*(1 + I*x)^(5/6))

Maple [F]

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3)/x,x)

[Out] Integral((I*(x - I)/sqrt(x**2 + 1))**(1/3)/x, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x,x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{1/3}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x,x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x, x)

$$3.119 \quad \int \frac{e^{\frac{1}{3}i \operatorname{ArcTan}(x)}}{x^2} dx$$

Optimal. Leaf size=253

$$\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} + \frac{i \operatorname{ArcTan}\left(\frac{1-2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{\sqrt{3}} - \frac{i \operatorname{ArcTan}\left(\frac{1+2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{\sqrt{3}} - \frac{2}{3} i \tanh^{-1}\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right) + \frac{1}{6} i$$

[Out] $-(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)}/x-2/3*I*\operatorname{arctanh}((1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})+1/6*I*\ln(1-(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)})-1/6*I*\ln(1+(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)})+1/3*I*\operatorname{arctan}(1/3*(1-2*(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)))*3^{(1/2)})*3^{(1/2)}-1/3*I*\operatorname{arctan}(1/3*(1+2*(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)))*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.12, antiderivative size = 253, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 9, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.643$, Rules used = {5170, 96, 95, 216, 648, 632, 210, 642, 212}

$$\frac{i \operatorname{ArcTan}\left(\frac{1-2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{\sqrt{3}} - \frac{i \operatorname{ArcTan}\left(\frac{1+2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{\sqrt{3}} - \frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} + \frac{1}{6} i \log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) - \frac{1}{6} i \log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) - \frac{2}{3} i \tanh^{-1}\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((I/3)*\operatorname{ArcTan}[x])/x^2}, x]$

[Out] $-(((1-I*x)^{(5/6)}*(1+I*x)^{(1/6)})/x) + (I*\operatorname{ArcTan}[(1-(2*(1+I*x)^{(1/6)})/(1-I*x)^{(1/6)})/\operatorname{Sqrt}[3]])/\operatorname{Sqrt}[3] - (I*\operatorname{ArcTan}[(1+(2*(1+I*x)^{(1/6)})/(1-I*x)^{(1/6)})/\operatorname{Sqrt}[3]])/\operatorname{Sqrt}[3] - ((2*I)/3)*\operatorname{ArcTanh}[(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}] + (I/6)*\operatorname{Log}[1-(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}] - (I/6)*\operatorname{Log}[1+(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}]$

Rule 95

$\operatorname{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}]/((e_.) + (f_.)*(x_.)), x_Symbol] \rightarrow \operatorname{With}[q = \operatorname{Denominator}[m], \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \operatorname{FreeQ}[a, b, c, d, e, f], x] \&\& \operatorname{EqQ}[m + n + 1, 0] \&\& \operatorname{RationalQ}[n] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 96

$\operatorname{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)}$

)/((m + 1)*(b*e - a*f))), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))),
 Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b,
 c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimpler
 Q[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 210

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
 -1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
 & (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
 ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
 Q[a, 0] || LtQ[b, 0])

Rule 216

Int[((a_) + (b_)*(x_)^(n_))(-1), x_Symbol] := Module[{r = Numerator[Rt[-a
 /b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*
 Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2
 *k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n))*I
 nt[1/(r^2 - s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x],
 x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]

Rule 632

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
 nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
 x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
 ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
 t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
 [2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{3}i \tan^{-1}(x)}}{x^2} dx &= \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix} x^2} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} + \frac{1}{3}i \int \frac{1}{\sqrt[6]{1-ix} (1+ix)^{5/6} x} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} + 2i \operatorname{Subst} \left(\int \frac{1}{-1+x^6} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} - \frac{2}{3}i \operatorname{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) - \frac{2}{3}i \operatorname{Subst} \left(\int \frac{1-\frac{x}{2}}{1-x+x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} - \frac{2}{3}i \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) + \frac{1}{6}i \operatorname{Subst} \left(\int \frac{-1+2x}{1-x+x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} - \frac{2}{3}i \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) + \frac{1}{6}i \log \left(1 - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{x} + \frac{i \tan^{-1} \left(\frac{1 - \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{i \tan^{-1} \left(\frac{1 + \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right)}{\sqrt{3}} - \frac{2}{3}i \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 64, normalized size = 0.25

$$\frac{i(1-ix)^{5/6} (-5i + 5x + 2x {}_2F_1(\frac{5}{6}, 1; \frac{11}{6}; \frac{i+x}{i-x}))}{5(1+ix)^{5/6} x}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x])/x^2,x]

[Out] ((-1/5*I)*(1 - I*x)^(5/6)*(-5*I + 5*x + 2*x*Hypergeometric2F1[5/6, 1, 11/6, (I + x)/(I - x)]))/((1 + I*x)^(5/6)*x)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{ix+1}{\sqrt{x^2+1}}\right)^{\frac{1}{3}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^2,x)**[Out]** int(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^2,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^2,x, algorithm="maxima")**[Out]** integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x^2, x)**Fricas [A]**

time = 1.19, size = 211, normalized size = 0.83

$$\frac{(\sqrt{3}x-iz)\log\left(\frac{1}{2}\sqrt{3} + \left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}} + i\right) + (\sqrt{3}x+iz)\log\left(\frac{1}{2}\sqrt{3} + \left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}} - i\right) - (\sqrt{3}x+iz)\log\left(-\frac{1}{2}\sqrt{3} + \left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}} + i\right) - (\sqrt{3}x-iz)\log\left(-\frac{1}{2}\sqrt{3} + \left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}} - i\right) - 2ix\log\left(\left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}} + 1\right) + 2ix\log\left(\left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}} - 1\right) - 6(-ix+1)\left(\frac{\sqrt{3}x+1}{2}\right)^{\frac{1}{3}}}{6x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^2,x, algorithm="fricas")

[Out] 1/6*((sqrt(3)*x - I*x)*log(1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2) + (sqrt(3)*x + I*x)*log(1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2) - (sqrt(3)*x + I*x)*log(-1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2) - (sqrt(3)*x - I*x)*log(-1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2) - 2*I*x*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1) + 2*I*x*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1) - 6*(-I*x + 1)*(I*sqrt(x^2 + 1)/(x + I))^(1/3))/x

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[3]{\frac{i(x-i)}{\sqrt{x^2+1}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3)/x**2,x)

[Out] Integral((I*(x - I)/sqrt(x**2 + 1))**(1/3)/x**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^2,x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{1/3}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x^2,x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x^2, x)

$$3.120 \quad \int \frac{e^{\frac{1}{3}i \operatorname{ArcTan}(x)}}{x^3} dx$$

Optimal. Leaf size=280

$$\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} - \frac{\operatorname{ArcTan}\left(\frac{1-2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{6\sqrt{3}} + \frac{\operatorname{ArcTan}\left(\frac{1+2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{6\sqrt{3}} + \frac{1}{9} \operatorname{tanh}^{-1}\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)$$

[Out] $-1/2*(1-I*x)^{(5/6)}*(1+I*x)^{(7/6)}/x^2-1/6*I*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)}/x+1/9*\operatorname{arctanh}((1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})-1/36*\ln(1-(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}+1/36*\ln(1+(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)})-1/18*\arctan(1/3*(1-2*(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})*3^{(1/2)})*3^{(1/2)}+1/18*\arctan(1/3*(1+2*(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 280, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 10, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {5170, 98, 96, 95, 216, 648, 632, 210, 642, 212}

$$\frac{\operatorname{ArcTan}\left(\frac{1-2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{6\sqrt{3}} + \frac{\operatorname{ArcTan}\left(\frac{1+2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{6\sqrt{3}} - \frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} - \frac{1}{36} \log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) + \frac{1}{36} \log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) + \frac{1}{9} \operatorname{tanh}^{-1}\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((I/3)*\operatorname{ArcTan}[x])}/x^3, x]$

[Out] $-1/2*((1-I*x)^{(5/6)}*(1+I*x)^{(7/6)})/x^2 - ((I/6)*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)})/x - \operatorname{ArcTan}[(1-(2*(1+I*x)^{(1/6)})/(1-I*x)^{(1/6)})/\operatorname{Sqrt}[3]]/(6*\operatorname{Sqrt}[3]) + \operatorname{ArcTan}[(1+(2*(1+I*x)^{(1/6)})/(1-I*x)^{(1/6)})/\operatorname{Sqrt}[3]]/(6*\operatorname{Sqrt}[3]) + \operatorname{ArcTanh}[(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}]/9 - \operatorname{Log}[1-(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)} + (1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}]/36 + \operatorname{Log}[1+(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)} + (1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}]/36$

Rule 95

$\operatorname{Int}[(((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)})/((e_.) + (f_.)*(x_.)), x_Symbol] := \operatorname{With}[\{q = \operatorname{Denominator}[m]\}, \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \operatorname{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \operatorname{EqQ}[m + n + 1, 0] \&\& \operatorname{RationalQ}[n] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 96

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^ (p_.), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1
))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))),
Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b,
c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimpler
Q[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

```

Rule 98

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^ (p_.), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x
))^ (p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Dist[(a*d*f*(m + 1) + b*
c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*
x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n
, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m
, 1])

```

Rule 210

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*
ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (Gt
Q[a, 0] || LtQ[b, 0])

```

Rule 216

```

Int[((a_) + (b_.)*(x_)^(n_))^(n_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^ (p_.), x_Symbol] := Module[{r = Numerator[Rt[-a
/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*
Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2
*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n))*I
nt[1/(r^2 - s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x],
x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]

```

Rule 632

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[In
t[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
x] && NeQ[b^2 - 4*a*c, 0]

```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := D
ist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), In
t[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{3}i \tan^{-1}(x)}}{x^3} dx &= \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix} x^3} dx \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} + \frac{1}{6}i \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix} x^2} dx \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} - \frac{1}{18} \int \frac{1}{\sqrt[6]{1-ix} (1+ix)^{5/6} x} dx \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} - \frac{1}{3} \text{Subst} \left(\int \frac{1}{-1+x^6} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} + \frac{1}{9} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} + \frac{1}{9} \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) - \frac{1}{36} \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} + \frac{1}{9} \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) - \frac{1}{36} \log \left(1 - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6}(1+ix)^{7/6}}{2x^2} - \frac{i(1-ix)^{5/6}\sqrt[6]{1+ix}}{6x} - \frac{\tan^{-1} \left(\frac{1 - \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right)}{6\sqrt{3}} + \frac{\tan^{-1} \left(\frac{1 + \frac{2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}}{\sqrt{3}} \right)}{6\sqrt{3}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 72, normalized size = 0.26

$$\frac{(1 - ix)^{5/6} (5(-3 - 7ix + 4x^2) + 2x^2 {}_2F_1(\frac{5}{6}, 1; \frac{11}{6}; \frac{i+x}{i-x}))}{30(1 + ix)^{5/6}x^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x])/x^3,x]

[Out] ((1 - I*x)^(5/6)*(5*(-3 - (7*I)*x + 4*x^2) + 2*x^2*Hypergeometric2F1[5/6, 1, 11/6, (I + x)/(I - x)]))/(30*(1 + I*x)^(5/6)*x^2)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{ix+1}{\sqrt{x^2+1}}\right)^{\frac{1}{3}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^3,x)

[Out] int(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^3,x, algorithm="maxima")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x^3, x)

Fricas [A]

time = 2.29, size = 234, normalized size = 0.84

$$\frac{2x^2 \log\left(\left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+1\right) - 2x^2 \log\left(\left(\frac{\sqrt{x^2+1}}{x-i}\right)^{\frac{1}{3}}-1\right) + (i\sqrt{3}x^2+x^2) \log\left(\frac{1}{3}\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+1\right) + (i\sqrt{3}x^2-x^2) \log\left(\frac{1}{3}\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x-i}\right)^{\frac{1}{3}}-1\right) + (-i\sqrt{3}x^2+x^2) \log\left(-\frac{1}{3}\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+1\right) + (-i\sqrt{3}x^2-x^2) \log\left(-\frac{1}{3}\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x-i}\right)^{\frac{1}{3}}-1\right) - 6(4x^2+ix+3)\left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}}{36x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^3,x, algorithm="fricas")

[Out] 1/36*(2*x^2*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1) - 2*x^2*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1) + (I*sqrt(3)*x^2 + x^2)*log(1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2) + (I*sqrt(3)*x^2 - x^2)*log(1/2*I*sqrt(3)

) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2) + (-I*sqrt(3)*x^2 + x^2)*log(-1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2) + (-I*sqrt(3)*x^2 - x^2)*log(-1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2) - 6*(4*x^2 + I*x + 3)*(I*sqrt(x^2 + 1)/(x + I))^(1/3))/x^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[3]{\frac{i(x-i)}{\sqrt{x^2+1}}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3)/x**3,x)

[Out] Integral((I*(x - I)/sqrt(x**2 + 1))**(1/3)/x**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^3,x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{1/3}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x^3,x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x^3, x)

$$3.121 \quad \int \frac{e^{\frac{1}{3}i\text{ArcTan}(x)}}{x^4} dx$$

Optimal. Leaf size=319

$$\frac{(1-ix)^{5/6}\sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6}\sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6}\sqrt[6]{1+ix}}{27x} - \frac{19i\text{ArcTan}\left(\frac{1-2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{54\sqrt{3}} + \frac{19i\text{A...}}{54\sqrt{3}}$$

[Out] $-1/3*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)}/x^3-7/18*I*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)}/x^2+11/27*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)}/x+19/81*I*\text{arctanh}((1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})-19/324*I*\ln(1-(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3}))+19/324*I*\ln(1+(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)})-19/162*I*\arctan(1/3*(1-2*(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})*3^{(1/2)})*3^{(1/2)}+19/162*I*\arctan(1/3*(1+2*(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)})*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 319, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 11, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.786$, Rules used = {5170, 101, 156, 12, 95, 216, 648, 632, 210, 642, 212}

$$-\frac{19i\text{ArcTan}\left(\frac{1-2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{54\sqrt{3}} + \frac{19i\text{ArcTan}\left(\frac{1+2\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)}{54\sqrt{3}} - \frac{(1-ix)^{5/6}\sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6}\sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6}\sqrt[6]{1+ix}}{27x} - \frac{19}{324}i\log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} - \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) + \frac{19}{324}i\log\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} + 1\right) + \frac{19}{81}i\text{tanh}^{-1}\left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}}\right)$$

Antiderivative was successfully verified.

[In] Int[E^((I/3)*ArcTan[x])/x^4,x]

[Out] $-1/3*((1-I*x)^{(5/6)}*(1+I*x)^{(1/6)})/x^3 - (((7*I)/18)*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)})/x^2 + (11*(1-I*x)^{(5/6)}*(1+I*x)^{(1/6)})/(27*x) - (((19*I)/54)*\text{ArcTan}[(1-(2*(1+I*x)^{(1/6)})/(1-I*x)^{(1/6)})/\text{Sqrt}[3]])/\text{Sqrt}[3] + (((19*I)/54)*\text{ArcTan}[(1+(2*(1+I*x)^{(1/6)})/(1-I*x)^{(1/6)})/\text{Sqrt}[3]])/\text{Sqrt}[3] + ((19*I)/81)*\text{ArcTanh}[(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}] - ((19*I)/324)*\text{Log}[1-(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}] + ((19*I)/324)*\text{Log}[1+(1+I*x)^{(1/6)}/(1-I*x)^{(1/6)}+(1+I*x)^{(1/3)}/(1-I*x)^{(1/3)}]$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_)]/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1))

$- 1)/(b*e - a*f - (d*e - c*f)*x^q)$, x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)
], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n]
 && LtQ[-1, m, 0] && SimplrQ[a + b*x, c + d*x]

Rule 101

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1)/((m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 156

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h]*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 216

Int[((a_) + (b_.)*(x_)^(n_))^(n_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Module[{r = Numerator[Rt[-a/b, n]], s = Denominator[Rt[-a/b, n]], k, u}, Simp[u = Int[(r - s*Cos[(2*k*Pi)/n]*x)/(r^2 - 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x] + Int[(r + s*Cos[(2*k*Pi)/n]*x)/(r^2 + 2*r*s*Cos[(2*k*Pi)/n]*x + s^2*x^2), x]; 2*(r^2/(a*n))*Int[1/(r^2 - s^2*x^2), x] + Dist[2*(r/(a*n)), Sum[u, {k, 1, (n - 2)/4}], x], x]] /; FreeQ[{a, b}, x] && IGtQ[(n - 2)/4, 0] && NegQ[a/b]

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_.) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{3}i \tan^{-1}(x)}}{x^4} dx &= \int \frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix} x^4} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} + \frac{1}{3} \int \frac{\frac{7i}{3} - 2x}{\sqrt[6]{1-ix} (1+ix)^{5/6} x^3} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} - \frac{1}{6} \int \frac{\frac{22}{9} + \frac{7ix}{3}}{\sqrt[6]{1-ix} (1+ix)^{5/6} x^2} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} + \frac{1}{6} \int -\frac{1}{27\sqrt[6]{1-ix}} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} - \frac{19}{162} i \int \frac{1}{\sqrt[6]{1-ix}} dx \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} - \frac{19}{27} i \text{Subst} \left(\frac{1}{\sqrt[6]{1-ix}}, x \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} + \frac{19}{81} i \text{Subst} \left(\frac{1}{\sqrt[6]{1-ix}}, x \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} + \frac{19}{81} i \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} + \frac{19}{81} i \tanh^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right) \\
&= -\frac{(1-ix)^{5/6} \sqrt[6]{1+ix}}{3x^3} - \frac{7i(1-ix)^{5/6} \sqrt[6]{1+ix}}{18x^2} + \frac{11(1-ix)^{5/6} \sqrt[6]{1+ix}}{27x} - \frac{19i \tan^{-1} \left(\frac{\sqrt[6]{1+ix}}{\sqrt[6]{1-ix}} \right)}{81}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 81, normalized size = 0.25

$$\frac{(1-ix)^{5/6} (5(-18-39ix+43x^2+22ix^3) + 38ix^3 {}_2F_1\left(\frac{5}{6}, 1; \frac{11}{6}; \frac{i+x}{i-x}\right))}{270(1+ix)^{5/6} x^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/3)*ArcTan[x])/x^4,x]

[Out] ((1 - I*x)^(5/6)*(5*(-18 - (39*I)*x + 43*x^2 + (22*I)*x^3) + (38*I)*x^3*Hypergeometric2F1[5/6, 1, 11/6, (I + x)/(I - x)]))/(270*(1 + I*x)^(5/6)*x^3)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{ix+1}{\sqrt{x^2+1}}\right)^{\frac{1}{3}}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^4,x)``[Out] int(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^4,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^4,x, algorithm="maxima")``[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x^4, x)`**Fricas [A]**

time = 1.58, size = 243, normalized size = 0.76

$$\frac{38i^3 \log\left(\left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+1\right) - 38i^3 \log\left(\left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}-1\right) - 19(\sqrt{3}x^3 - ix^3) \log\left(\frac{1}{2}i\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+i\right) - 19(\sqrt{3}x^3 + ix^3) \log\left(\frac{1}{2}i\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}-i\right) + 19(\sqrt{3}x^3 - ix^3) \log\left(-\frac{1}{2}i\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+i\right) + 19(\sqrt{3}x^3 + ix^3) \log\left(-\frac{1}{2}i\sqrt{3} + \left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}-i\right) - 6(22ix^3 - x^2 + 3ix + 18)\left(\frac{\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}}{324x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^4,x, algorithm="fricas")`

```
[Out] 1/324*(38*I*x^3*log((I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1) - 38*I*x^3*log((I*
sqrt(x^2 + 1)/(x + I))^(1/3) - 1) - 19*(sqrt(3)*x^3 - I*x^3)*log(1/2*I*sqrt
(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2) - 19*(sqrt(3)*x^3 + I*x^3)*log
(1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) - 1/2) + 19*(sqrt(3)*x^3 +
I*x^3)*log(-1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) + 1/2) + 19*(s
qrt(3)*x^3 - I*x^3)*log(-1/2*I*sqrt(3) + (I*sqrt(x^2 + 1)/(x + I))^(1/3) -
1/2) - 6*(22*I*x^3 - x^2 + 3*I*x + 18)*(I*sqrt(x^2 + 1)/(x + I))^(1/3))/x^3
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(1/3)/x**4,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(1/3)/x^4,x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(1/3)/x^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{1/3}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x^4,x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(1/3)/x^4, x)

3.122 $\int e^{\frac{2}{3}i \operatorname{ArcTan}(x)} x^2 dx$

Optimal. Leaf size=177

$$-\frac{11}{27}i(1-ix)^{2/3}\sqrt[3]{1+ix} - \frac{1}{9}i(1-ix)^{2/3}(1+ix)^{4/3} + \frac{1}{3}(1-ix)^{2/3}(1+ix)^{4/3}x + \frac{22i \operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{27\sqrt{3}}$$

[Out] $-11/27*I*(1-I*x)^{(2/3)}*(1+I*x)^{(1/3)} - 1/9*I*(1-I*x)^{(2/3)}*(1+I*x)^{(4/3)} + 1/3*(1-I*x)^{(2/3)}*(1+I*x)^{(4/3)}*x + 11/27*I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}) + 11/81*I*\ln(1+I*x) + 22/81*I*\arctan(1/3*3^{(1/2)} - 2/3*(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 177, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5170, 92, 81, 52, 62}

$$\frac{22i \operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{27\sqrt{3}} + \frac{1}{3}(1-ix)^{2/3}x(1+ix)^{4/3} - \frac{1}{9}i(1-ix)^{2/3}(1+ix)^{4/3} - \frac{11}{27}i(1-ix)^{2/3}\sqrt[3]{1+ix} + \frac{11}{27}i \log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) + \frac{11}{81}i \log(1+ix)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((2*I)/3)*\operatorname{ArcTan}[x]}*x^2, x]$

[Out] $((-11*I)/27)*(1 - I*x)^{(2/3)}*(1 + I*x)^{(1/3)} - (I/9)*(1 - I*x)^{(2/3)}*(1 + I*x)^{(4/3)} + ((1 - I*x)^{(2/3)}*(1 + I*x)^{(4/3)}*x)/3 + (((22*I)/27)*\operatorname{ArcTan}[1/\operatorname{Sqrt}[3] - (2*(1 - I*x)^{(1/3)})/(\operatorname{Sqrt}[3]*(1 + I*x)^{(1/3)})])/\operatorname{Sqrt}[3] + ((11*I)/27)*\operatorname{Log}[1 + (1 - I*x)^{(1/3)}/(1 + I*x)^{(1/3)}] + ((11*I)/81)*\operatorname{Log}[1 + I*x]$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \operatorname{Dist}[n*(b*c - a*d)/(b*(m + n + 1)), \operatorname{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 62

$\operatorname{Int}[1/(((a_.) + (b_.)*(x_.))^{(1/3)}*((c_.) + (d_.)*(x_.))^{(2/3)}), x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Rt}[-d/b, 3]\}, \operatorname{Simp}[\operatorname{Sqrt}[3]*(q/d)*\operatorname{ArcTan}[1/\operatorname{Sqrt}[3] - 2*q*((a + b*x)^{(1/3)}/(\operatorname{Sqrt}[3]*(c + d*x)^{(1/3)}))], x] + (\operatorname{Simp}[3*(q/(2*d))*\operatorname{Log}[q*((a + b*x)^{(1/3)}/(c + d*x)^{(1/3)}) + 1], x] + \operatorname{Simp}[(q/(2*d))*\operatorname{Log}[c + d*x], x]) /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NegQ[d/b]

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92

Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int e^{\frac{2}{3}i \tan^{-1}(x)} x^2 dx &= \int \frac{\sqrt[3]{1+ix} x^2}{\sqrt[3]{1-ix}} dx \\
 &= \frac{1}{3}(1-ix)^{2/3}(1+ix)^{4/3}x + \frac{1}{3} \int \frac{(-1 - \frac{2ix}{3}) \sqrt[3]{1+ix}}{\sqrt[3]{1-ix}} dx \\
 &= -\frac{1}{9}i(1-ix)^{2/3}(1+ix)^{4/3} + \frac{1}{3}(1-ix)^{2/3}(1+ix)^{4/3}x - \frac{11}{27} \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix}} dx \\
 &= -\frac{11}{27}i(1-ix)^{2/3}\sqrt[3]{1+ix} - \frac{1}{9}i(1-ix)^{2/3}(1+ix)^{4/3} + \frac{1}{3}(1-ix)^{2/3}(1+ix)^{4/3}x - \frac{22}{81} \\
 &= -\frac{11}{27}i(1-ix)^{2/3}\sqrt[3]{1+ix} - \frac{1}{9}i(1-ix)^{2/3}(1+ix)^{4/3} + \frac{1}{3}(1-ix)^{2/3}(1+ix)^{4/3}x + \frac{22i}{81}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 73, normalized size = 0.41

$$\frac{1}{18}(1-ix)^{2/3} \left(2\sqrt[3]{1+ix} (-i + 4x + 3ix^2) - 11i\sqrt[3]{2} {}_2F_1\left(-\frac{1}{3}, \frac{2}{3}; \frac{5}{3}; \frac{1}{2} - \frac{ix}{2}\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^(((2*I)/3)*ArcTan[x])*x^2,x]

[Out] ((1 - I*x)^(2/3)*(2*(1 + I*x)^(1/3)*(-I + 4*x + (3*I)*x^2) - (11*I)*2^(1/3)*Hypergeometric2F1[-1/3, 2/3, 5/3, 1/2 - (I/2)*x]))/18

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{ix + 1}{\sqrt{x^2 + 1}} \right)^{\frac{2}{3}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x^2,x)

[Out] int(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x^2,x, algorithm="maxima")

[Out] integrate(x^2*((I*x + 1)/sqrt(x^2 + 1))^(2/3), x)

Fricas [A]

time = 1.40, size = 117, normalized size = 0.66

$$-\frac{11}{81}(\sqrt{3} + i) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + \frac{1}{2}i\sqrt{3} - \frac{1}{2}\right) + \frac{11}{81}(\sqrt{3} - i) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - \frac{1}{2}i\sqrt{3} - \frac{1}{2}\right) + \frac{1}{27}(9x^3 - 3ix^2 - 2x - 14i)\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + \frac{22}{81}i \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x^2,x, algorithm="fricas")

[Out] -11/81*(sqrt(3) + I)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1/2*I*sqrt(3) - 1/2) + 11/81*(sqrt(3) - I)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1/2*I*sqrt(3) - 1/2) + 1/27*(9*x^3 - 3*I*x^2 - 2*x - 14*I)*(I*sqrt(x^2 + 1)/(x + I))^(2/3) + 22/81*I*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(2/3)*x**2,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x^2,x, algorithm="giac")

[Out] integrate(x^2*((I*x + 1)/sqrt(x^2 + 1))^(2/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^2 \left(\frac{1 + x \text{Ii}}{\sqrt{x^2 + 1}} \right)^{2/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3),x)

[Out] int(x^2*((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3), x)

3.123 $\int e^{\frac{2}{3}i \operatorname{ArcTan}(x)} x dx$

Optimal. Leaf size=140

$$\frac{1}{3}(1-ix)^{2/3}\sqrt[3]{1+ix} + \frac{1}{2}(1-ix)^{2/3}(1+ix)^{4/3} - \frac{2\operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{3\sqrt{3}} - \frac{1}{3}\log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) - \frac{1}{9}\log(1+ix)$$

[Out] $\frac{1}{3}(1-I*x)^{(2/3)}*(1+I*x)^{(1/3)} + \frac{1}{2}(1-I*x)^{(2/3)}*(1+I*x)^{(4/3)} - \frac{1}{3}\ln(1+(1-I*x)^{(1/3))/(1+I*x)^{(1/3)}) - \frac{1}{9}\ln(1+I*x) - \frac{2}{9}\operatorname{arctan}(1/3*3^{(1/2)} - 2/3*(1-I*x)^{(1/3))/(1+I*x)^{(1/3)}*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5170, 81, 52, 62}

$$-\frac{2\operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{3\sqrt{3}} + \frac{1}{2}(1-ix)^{2/3}(1+ix)^{4/3} + \frac{1}{3}(1-ix)^{2/3}\sqrt[3]{1+ix} - \frac{1}{3}\log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) - \frac{1}{9}\log(1+ix)$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((2*I)/3)*\operatorname{ArcTan}[x]}*x, x]$

[Out] $((1 - I*x)^{(2/3)}*(1 + I*x)^{(1/3)})/3 + ((1 - I*x)^{(2/3)}*(1 + I*x)^{(4/3)})/2 - (2*\operatorname{ArcTan}[1/\operatorname{Sqrt}[3] - (2*(1 - I*x)^{(1/3)})/(\operatorname{Sqrt}[3]*(1 + I*x)^{(1/3)})])/(3*\operatorname{Sqrt}[3]) - \operatorname{Log}[1 + (1 - I*x)^{(1/3)}/(1 + I*x)^{(1/3)}]/3 - \operatorname{Log}[1 + I*x]/9$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \operatorname{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \operatorname{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{GtQ}[n, 0] \ \&\& \ \operatorname{NeQ}[m + n + 1, 0] \ \&\& \ !(\operatorname{IGtQ}[m, 0] \ \&\& \ (!\operatorname{IntegerQ}[n] \ || \ (\operatorname{GtQ}[m, 0] \ \&\& \ \operatorname{LtQ}[m - n, 0]))) \ \&\& \ !\operatorname{ILtQ}[m + n + 2, 0] \ \&\& \ \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 62

$\operatorname{Int}[1/(((a_.) + (b_.)*(x_.))^{(1/3)}*((c_.) + (d_.)*(x_.))^{(2/3)}), x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Rt}[-d/b, 3]\}, \operatorname{Simp}[\operatorname{Sqrt}[3]*(q/d)*\operatorname{ArcTan}[1/\operatorname{Sqrt}[3] - 2*q*((a + b*x)^{(1/3)}/(\operatorname{Sqrt}[3]*(c + d*x)^{(1/3)}))], x] + (\operatorname{Simp}[3*(q/(2*d))*\operatorname{Log}[q*((a + b*x)^{(1/3)}/(c + d*x)^{(1/3)}) + 1], x] + \operatorname{Simp}[(q/(2*d))*\operatorname{Log}[c + d*x], x]) /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \ \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \ \operatorname{NegQ}[d/b]$

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{2}{3}i \tan^{-1}(x)} x \, dx &= \int \frac{\sqrt[3]{1+ix} x}{\sqrt[3]{1-ix}} \, dx \\
&= \frac{1}{2}(1-ix)^{2/3}(1+ix)^{4/3} - \frac{1}{3}i \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix}} \, dx \\
&= \frac{1}{3}(1-ix)^{2/3}\sqrt[3]{1+ix} + \frac{1}{2}(1-ix)^{2/3}(1+ix)^{4/3} - \frac{2}{9}i \int \frac{1}{\sqrt[3]{1-ix}(1+ix)^{2/3}} \, dx \\
&= \frac{1}{3}(1-ix)^{2/3}\sqrt[3]{1+ix} + \frac{1}{2}(1-ix)^{2/3}(1+ix)^{4/3} - \frac{2 \tan^{-1}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{3\sqrt{3}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 54, normalized size = 0.39

$$\frac{1}{2}(1-ix)^{2/3} \left((1+ix)^{4/3} + \sqrt[3]{2} {}_2F_1\left(-\frac{1}{3}, \frac{2}{3}; \frac{5}{3}; \frac{1}{2} - \frac{ix}{2}\right) \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^(((2*I)/3)*ArcTan[x])*x,x]

[Out] ((1 - I*x)^(2/3)*((1 + I*x)^(4/3) + 2^(1/3)*Hypergeometric2F1[-1/3, 2/3, 5/3, 1/2 - (I/2)*x]))/2

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{ix + 1}{\sqrt{x^2 + 1}} \right)^{\frac{2}{3}} x \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x,x)`

[Out] `int(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x,x, algorithm="maxima")`

[Out] `integrate(x*((I*x + 1)/sqrt(x^2 + 1))^(2/3), x)`

Fricas [A]

time = 1.43, size = 116, normalized size = 0.83

$$-\frac{1}{9}(i\sqrt{3}-1)\log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}}+\frac{1}{2}i\sqrt{3}-\frac{1}{2}\right)-\frac{1}{9}(-i\sqrt{3}-1)\log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}}-\frac{1}{2}i\sqrt{3}-\frac{1}{2}\right)+\frac{1}{6}(3x^2-2ix+5)\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}}-\frac{2}{9}\log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}}+1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x,x, algorithm="fricas")`

[Out] `-1/9*(I*sqrt(3) - 1)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1/2*I*sqrt(3) - 1/2) - 1/9*(-I*sqrt(3) - 1)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1/2*I*sqrt(3) - 1/2) + 1/6*(3*x^2 - 2*I*x + 5)*(I*sqrt(x^2 + 1)/(x + I))^(2/3) - 2/9*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1)`

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x**2+1)**(1/2))**(2/3)*x,x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)*x,x, algorithm="giac")`

[Out] integrate(x*((I*x + 1)/sqrt(x^2 + 1))^(2/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x \left(\frac{1 + x \text{I}}{\sqrt{x^2 + 1}} \right)^{2/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*((x*I + 1)/(x^2 + 1)^(1/2))^(2/3), x)

[Out] int(x*((x*I + 1)/(x^2 + 1)^(1/2))^(2/3), x)

3.124 $\int e^{\frac{2}{3}i\text{ArcTan}(x)} dx$

Optimal. Leaf size=116

$$i(1-ix)^{2/3}\sqrt[3]{1+ix} - \frac{2i\text{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{\sqrt{3}} - i\log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) - \frac{1}{3}i\log(1+ix)$$

[Out] $I*(1-I*x)^{(2/3)}*(1+I*x)^{(1/3)} - I*\ln(1+(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}) - 1/3*I*\ln(1+I*x) - 2/3*I*\arctan(1/3*3^{(1/2)} - 2/3*(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.01, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.300$, Rules used = {5169, 52, 62}

$$-\frac{2i\text{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{\sqrt{3}} + i(1-ix)^{2/3}\sqrt[3]{1+ix} - i\log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) - \frac{1}{3}i\log(1+ix)$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)/3)*\text{ArcTan}[x]}, x]$

[Out] $I*(1 - I*x)^{(2/3)}*(1 + I*x)^{(1/3)} - ((2*I)*\text{ArcTan}[1/\text{Sqrt}[3] - (2*(1 - I*x))^{(1/3)}/(\text{Sqrt}[3]*(1 + I*x)^{(1/3)})]/\text{Sqrt}[3] - I*\text{Log}[1 + (1 - I*x)^{(1/3)}/(1 + I*x)^{(1/3)}] - (I/3)*\text{Log}[1 + I*x])$

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILTQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 62

```
Int[1/(((a_.) + (b_.)*(x_))^(1/3)*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] :=
With[{q = Rt[-d/b, 3]}, Simp[Sqrt[3]*(q/d)*ArcTan[1/Sqrt[3] - 2*q*((a + b*
x)^(1/3)/(Sqrt[3]*(c + d*x)^(1/3))], x] + (Simp[3*(q/(2*d))*Log[q*((a + b*
x)^(1/3)/(c + d*x)^(1/3)) + 1], x] + Simp[(q/(2*d))*Log[c + d*x], x])] /; F
reeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NegQ[d/b]
```

Rule 5169

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.)), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{\frac{2}{3}i \tan^{-1}(x)} dx &= \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix}} dx \\ &= i(1-ix)^{2/3} \sqrt[3]{1+ix} + \frac{2}{3} \int \frac{1}{\sqrt[3]{1-ix} (1+ix)^{2/3}} dx \\ &= i(1-ix)^{2/3} \sqrt[3]{1+ix} - \frac{2i \tan^{-1} \left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3} \sqrt[3]{1+ix}} \right)}{\sqrt{3}} - i \log \left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}} \right) - \frac{1}{3}i \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 34, normalized size = 0.29

$$-\frac{3}{2} i e^{\frac{8}{3} i \text{ArcTan}(x)} {}_2F_1 \left(\frac{4}{3}, 2; \frac{7}{3}; -e^{2i \text{ArcTan}(x)} \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^(((2*I)/3)*ArcTan[x]), x]

[Out] ((-3*I)/2)*E^(((8*I)/3)*ArcTan[x])*Hypergeometric2F1[4/3, 2, 7/3, -E^((2*I)*ArcTan[x])]

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{ix + 1}{\sqrt{x^2 + 1}} \right)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((((1+I*x)/(x^2+1))^(1/2))^(2/3), x)

[Out] int((((1+I*x)/(x^2+1))^(1/2))^(2/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3),x, algorithm="maxima")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3), x)

Fricas [A]

time = 2.13, size = 104, normalized size = 0.90

$$\frac{1}{3}(\sqrt{3} + i) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + \frac{1}{2}i\sqrt{3} - \frac{1}{2}\right) - \frac{1}{3}(\sqrt{3} - i) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - \frac{1}{2}i\sqrt{3} - \frac{1}{2}\right) + (x+i)\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - \frac{2}{3}i \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + 1\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3),x, algorithm="fricas")

[Out] 1/3*(sqrt(3) + I)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1/2*I*sqrt(3) - 1/2) - 1/3*(sqrt(3) - I)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1/2*I*sqrt(3) - 1/2) + (x + I)*(I*sqrt(x^2 + 1)/(x + I))^(2/3) - 2/3*I*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{ix + 1}{\sqrt{x^2 + 1}} \right)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(2/3),x)

[Out] Integral(((I*x + 1)/sqrt(x**2 + 1))**(2/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3),x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \left(\frac{1 + x i}{\sqrt{x^2 + 1}} \right)^{\frac{2}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3),x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3), x)

$$3.125 \quad \int \frac{e^{\frac{2}{3}i \operatorname{ArcTan}(x)}}{x} dx$$

Optimal. Leaf size=163

$$\sqrt{3} \operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right) + \sqrt{3} \operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right) + \frac{3}{2} \log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) + \frac{3}{2} \log\left(1 - \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right)$$

```
[Out] 3/2*ln(1+(1-I*x)^(1/3)/(1+I*x)^(1/3))+3/2*ln((1-I*x)^(1/3)-(1+I*x)^(1/3))+1/2*ln(1+I*x)-1/2*ln(x)+arctan(1/3*3^(1/2)-2/3*(1-I*x)^(1/3)/(1+I*x)^(1/3)*3^(1/2))*3^(1/2)+arctan(1/3*3^(1/2)+2/3*(1-I*x)^(1/3)/(1+I*x)^(1/3)*3^(1/2))*3^(1/2)
```

Rubi [A]

time = 0.03, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5170, 132, 62, 93}

$$\sqrt{3} \operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right) + \sqrt{3} \operatorname{ArcTan}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right) + \frac{3}{2} \log\left(1 + \frac{\sqrt[3]{1-ix}}{\sqrt[3]{1+ix}}\right) + \frac{3}{2} \log\left(\sqrt[3]{1-ix} - \sqrt[3]{1+ix}\right) + \frac{1}{2} \log(1+ix) - \frac{\log(x)}{2}$$

Antiderivative was successfully verified.

```
[In] Int[E^(((2*I)/3)*ArcTan[x])/x,x]
```

```
[Out] Sqrt[3]*ArcTan[1/Sqrt[3] - (2*(1 - I*x)^(1/3))/(Sqrt[3]*(1 + I*x)^(1/3))] + Sqrt[3]*ArcTan[1/Sqrt[3] + (2*(1 - I*x)^(1/3))/(Sqrt[3]*(1 + I*x)^(1/3))] + (3*Log[1 + (1 - I*x)^(1/3)/(1 + I*x)^(1/3)])/2 + (3*Log[(1 - I*x)^(1/3) - (1 + I*x)^(1/3)])/2 + Log[1 + I*x]/2 - Log[x]/2
```

Rule 62

```
Int[1/(((a_.) + (b_.)*(x_))^(1/3)*((c_.) + (d_.)*(x_))^(2/3)), x_Symbol] := With[{q = Rt[-d/b, 3]}, Simp[Sqrt[3]*(q/d)*ArcTan[1/Sqrt[3] - 2*q*((a + b*x)^(1/3)/(Sqrt[3]*(c + d*x)^(1/3))]], x] + (Simp[3*(q/(2*d))*Log[q*((a + b*x)^(1/3)/(c + d*x)^(1/3)) + 1], x] + Simp[(q/(2*d))*Log[c + d*x], x])] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && NegQ[d/b]
```

Rule 93

```
Int[1/(((a_.) + (b_.)*(x_))^(1/3)*((c_.) + (d_.)*(x_))^(2/3)*((e_.) + (f_.)*(x_))), x_Symbol] := With[{q = Rt[(d*e - c*f)/(b*e - a*f), 3]}, Simp[(-Sqrt[3])*q*(ArcTan[1/Sqrt[3] + 2*q*((a + b*x)^(1/3)/(Sqrt[3]*(c + d*x)^(1/3))])/ (d*e - c*f)], x] + (Simp[q*(Log[e + f*x]/(2*(d*e - c*f))), x] - Simp[3*q*(Log[q*(a + b*x)^(1/3) - (c + d*x)^(1/3)]/(2*(d*e - c*f))), x])] /; FreeQ[{a, b, c, d, e, f}, x]
```

Rule 132

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m+n)*f^p, Int[(a+b*x)^(m-1)/(c+d*x)^m, x], x] + Int[(a+b*x)^(m-1)*((e+f*x)^p/(c+d*x)^m)*ExpandToSum[(a+b*x)*(c+d*x)^(-p-1) - (b*d^(-p-1)*f^p)/(e+f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m+n+p+1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1-I*a*x)^(I*(n/2))/(1+I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n-1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\frac{2}{3}i \tan^{-1}(x)}}{x} dx &= \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix} x} dx \\ &= i \int \frac{1}{\sqrt[3]{1-ix} (1+ix)^{2/3}} dx + \int \frac{1}{\sqrt[3]{1-ix} (1+ix)^{2/3} x} dx \\ &= \sqrt{3} \tan^{-1} \left(\frac{1}{\sqrt{3}} - \frac{2\sqrt[3]{1-ix}}{\sqrt{3} \sqrt[3]{1+ix}} \right) + \sqrt{3} \tan^{-1} \left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3} \sqrt[3]{1+ix}} \right) + \frac{3}{2} \log \left(1 + \right. \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 90, normalized size = 0.55

$$\frac{3(1-ix)^{2/3} \left(\sqrt[3]{2} (1+ix)^{2/3} {}_2F_1\left(\frac{2}{3}, \frac{2}{3}; \frac{5}{3}; \frac{1}{2} - \frac{ix}{2}\right) + 2 {}_2F_1\left(\frac{2}{3}, 1; \frac{5}{3}; \frac{i+x}{i-x}\right) \right)}{4(1+ix)^{2/3}}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(((2*I)/3)*ArcTan[x])/x,x]
```

```
[Out] (-3*(1-I*x)^(2/3)*(2^(1/3)*(1+I*x)^(2/3)*Hypergeometric2F1[2/3, 2/3, 5/3, 1/2 - (I/2)*x] + 2*Hypergeometric2F1[2/3, 1, 5/3, (I+x)/(I-x)])/(4*(1+I*x)^(2/3))
```

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{ix+1}{\sqrt{x^2+1}} \right)^{\frac{2}{3}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x,x)`

[Out] `int(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x,x, algorithm="maxima")`

[Out] `integrate((I*x + 1)/sqrt(x^2 + 1))^(2/3)/x, x)`

Fricas [A]

time = 2.36, size = 145, normalized size = 0.89

$$\frac{1}{2}(i\sqrt{3}-1)\log\left(\frac{\sqrt{3}(ix-1)+x+2i\sqrt{x^2+1}\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+i}{2(x+i)}\right)+\frac{1}{2}(-i\sqrt{3}-1)\log\left(\frac{\sqrt{3}(-ix+1)+x+2i\sqrt{x^2+1}\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+i}{2(x+i)}\right)+\log\left(-\frac{x-i\sqrt{x^2+1}\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{1}{3}}+i}{x+i}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x,x, algorithm="fricas")`

[Out] `1/2*(I*sqrt(3) - 1)*log(1/2*(sqrt(3)*(I*x - 1) + x + 2*I*sqrt(x^2 + 1)*(I*sqrt(x^2 + 1)/(x + I)))^(1/3) + I)/(x + I)) + 1/2*(-I*sqrt(3) - 1)*log(1/2*(sqrt(3)*(-I*x + 1) + x + 2*I*sqrt(x^2 + 1)*(I*sqrt(x^2 + 1)/(x + I)))^(1/3) + I)/(x + I)) + log(-(x - I*sqrt(x^2 + 1)*(I*sqrt(x^2 + 1)/(x + I)))^(1/3) + I)/(x + I))`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{i(x-i)}{\sqrt{x^2+1}}\right)^{\frac{2}{3}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*x)/(x**2+1)**(1/2))**(2/3)/x,x)`

[Out] `Integral((I*(x - I)/sqrt(x**2 + 1))**(2/3)/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x,x, algorithm="giac")
```

```
[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3)/x, x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{2/3}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3)/x,x)
```

```
[Out] int((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3)/x, x)
```

$$3.126 \quad \int \frac{e^{\frac{2}{3}i\text{ArcTan}(x)}}{x^2} dx$$

Optimal. Leaf size=111

$$-\frac{(1-ix)^{2/3}\sqrt[3]{1+ix}}{x} + \frac{2i\text{ArcTan}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{\sqrt{3}} + i\log\left(\sqrt[3]{1-ix} - \sqrt[3]{1+ix}\right) - \frac{1}{3}i\log(x)$$

[Out] $-(1-I*x)^{(2/3)}*(1+I*x)^{(1/3)}/x+I*\ln((1-I*x)^{(1/3)}-(1+I*x)^{(1/3)})-1/3*I*\ln(x)+2/3*I*\arctan(1/3*3^{(1/2)}+2/3*(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 111, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5170, 96, 93}

$$\frac{2i\text{ArcTan}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{\sqrt{3}} - \frac{(1-ix)^{2/3}\sqrt[3]{1+ix}}{x} + i\log\left(\sqrt[3]{1-ix} - \sqrt[3]{1+ix}\right) - \frac{1}{3}i\log(x)$$

Antiderivative was successfully verified.

[In] Int[E^(((2*I)/3)*ArcTan[x])/x^2,x]

[Out] $-(((1-I*x)^{(2/3)}*(1+I*x)^{(1/3)})/x) + ((2*I)*\text{ArcTan}[1/\text{Sqrt}[3] + (2*(1-I*x)^{(1/3)})/(\text{Sqrt}[3]*(1+I*x)^{(1/3)})])/\text{Sqrt}[3] + I*\text{Log}[(1-I*x)^{(1/3)} - (1+I*x)^{(1/3)}] - (I/3)*\text{Log}[x]$

Rule 93

Int[1/(((a_.) + (b_.)*(x_))^(1/3)*((c_.) + (d_.)*(x_))^(2/3)*((e_.) + (f_.)*(x_))), x_Symbol] :> With[{q = Rt[(d*e - c*f)/(b*e - a*f), 3]}, Simp[(-Sqrt[3])*q*(ArcTan[1/Sqrt[3] + 2*q*((a + b*x)^(1/3)/(Sqrt[3]*(c + d*x)^(1/3))])/(d*e - c*f), x] + (Simp[q*(Log[e + f*x]/(2*(d*e - c*f))), x] - Simp[3*q*(Log[q*(a + b*x)^(1/3) - (c + d*x)^(1/3)]/(2*(d*e - c*f))), x])] /; FreeQ[{a, b, c, d, e, f}, x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1)/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\frac{2}{3}i \tan^{-1}(x)}}{x^2} dx &= \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix} x^2} dx \\ &= -\frac{(1-ix)^{2/3} \sqrt[3]{1+ix}}{x} + \frac{2}{3}i \int \frac{1}{\sqrt[3]{1-ix} (1+ix)^{2/3} x} dx \\ &= -\frac{(1-ix)^{2/3} \sqrt[3]{1+ix}}{x} + \frac{2i \tan^{-1}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{\sqrt{3}} + i \log\left(\sqrt[3]{1-ix} - \sqrt[3]{1+ix}\right) \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 59, normalized size = 0.53

$$\frac{i(1-ix)^{2/3} \left(-i+x+x {}_2F_1\left(\frac{2}{3}, 1; \frac{5}{3}; \frac{i+x}{i-x}\right)\right)}{(1+ix)^{2/3} x}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((2*I)/3)*ArcTan[x])/x^2,x]

[Out] ((-I)*(1 - I*x)^(2/3)*(-I + x + x*Hypergeometric2F1[2/3, 1, 5/3, (I + x)/(I - x)]))/((1 + I*x)^(2/3)*x)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{ix+1}{\sqrt{x^2+1}}\right)^{\frac{2}{3}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^2,x)

[Out] int(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^2,x, algorithm="maxima")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3)/x^2, x)

Fricas [A]

time = 2.52, size = 120, normalized size = 1.08

$$\frac{(\sqrt{3}x - ix) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right) - (\sqrt{3}x + ix) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right) + 2ix \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - 1\right) - 3(-ix + 1)\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}}}{3x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^2,x, algorithm="fricas")

[Out] 1/3*((sqrt(3)*x - I*x)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1/2*I*sqrt(3) + 1/2) - (sqrt(3)*x + I*x)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1/2*I*sqrt(3) + 1/2) + 2*I*x*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1) - 3*(-I*x + 1)*(I*sqrt(x^2 + 1)/(x + I))^(2/3))/x

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(2/3)/x**2,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^2,x, algorithm="giac")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3)/x^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{2/3}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3)/x^2,x)

[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3)/x^2, x)

$$3.127 \quad \int \frac{e^{\frac{2}{3}i\text{ArcTan}(x)}}{x^3} dx$$

Optimal. Leaf size=142

$$\frac{(1-ix)^{2/3}(1+ix)^{4/3}}{2x^2} - \frac{i(1-ix)^{2/3}\sqrt[3]{1+ix}}{3x} - \frac{2\text{ArcTan}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{3\sqrt{3}} - \frac{1}{3}\log\left(\sqrt[3]{1-ix} - \sqrt[3]{1+ix}\right)$$

[Out] $-1/2*(1-I*x)^{(2/3)}*(1+I*x)^{(4/3)}/x^2-1/3*I*(1-I*x)^{(2/3)}*(1+I*x)^{(1/3)}/x-1/3*\ln((1-I*x)^{(1/3)}-(1+I*x)^{(1/3}))+1/9*\ln(x)-2/9*\arctan(1/3*3^{(1/2)}+2/3*(1-I*x)^{(1/3)}/(1+I*x)^{(1/3)}*3^{(1/2)})*3^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5170, 98, 96, 93}

$$\frac{2\text{ArcTan}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{3\sqrt{3}} - \frac{(1-ix)^{2/3}(1+ix)^{4/3}}{2x^2} - \frac{i(1-ix)^{2/3}\sqrt[3]{1+ix}}{3x} - \frac{1}{3}\log\left(\sqrt[3]{1-ix} - \sqrt[3]{1+ix}\right) + \frac{\log(x)}{9}$$

Antiderivative was successfully verified.

[In] `Int[E^(((2*I)/3)*ArcTan[x])/x^3,x]`

[Out] $-1/2*((1-I*x)^{(2/3)}*(1+I*x)^{(4/3)}/x^2 - ((I/3)*(1-I*x)^{(2/3)}*(1+I*x)^{(1/3)}/x - (2*\text{ArcTan}[1/\text{Sqrt}[3] + (2*(1-I*x)^{(1/3)})/(\text{Sqrt}[3]*(1+I*x)^{(1/3)})])/((3*\text{Sqrt}[3]) - \text{Log}[(1-I*x)^{(1/3)} - (1+I*x)^{(1/3)}]/3 + \text{Log}[x]/9$

Rule 93

`Int[1/(((a_.) + (b_.)*(x_))^(1/3)*((c_.) + (d_.)*(x_))^(2/3)*((e_.) + (f_.)*(x_))), x_Symbol] :> With[{q = Rt[(d*e - c*f)/(b*e - a*f), 3]}, Simp[(-Sqrt[3])*q*(ArcTan[1/Sqrt[3] + 2*q*((a + b*x)^(1/3)/(Sqrt[3]*(c + d*x)^(1/3))]/(d*e - c*f)), x] + (Simp[q*(Log[e + f*x]/(2*(d*e - c*f))), x] - Simp[3*q*(Log[q*(a + b*x)^(1/3) - (c + d*x)^(1/3)]/(2*(d*e - c*f))), x])] /; FreeQ[{a, b, c, d, e, f}, x]`

Rule 96

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{\frac{2}{3}i \tan^{-1}(x)}}{x^3} dx &= \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix} x^3} dx \\ &= -\frac{(1-ix)^{2/3}(1+ix)^{4/3}}{2x^2} + \frac{1}{3}i \int \frac{\sqrt[3]{1+ix}}{\sqrt[3]{1-ix} x^2} dx \\ &= -\frac{(1-ix)^{2/3}(1+ix)^{4/3}}{2x^2} - \frac{i(1-ix)^{2/3}\sqrt[3]{1+ix}}{3x} - \frac{2}{9} \int \frac{1}{\sqrt[3]{1-ix} (1+ix)^{2/3} x} dx \\ &= -\frac{(1-ix)^{2/3}(1+ix)^{4/3}}{2x^2} - \frac{i(1-ix)^{2/3}\sqrt[3]{1+ix}}{3x} - \frac{2 \tan^{-1}\left(\frac{1}{\sqrt{3}} + \frac{2\sqrt[3]{1-ix}}{\sqrt{3}\sqrt[3]{1+ix}}\right)}{3\sqrt{3}} - \frac{1}{3} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 69, normalized size = 0.49

$$\frac{(1-ix)^{2/3}(-3-8ix+5x^2+2x^2 {}_2F_1\left(\frac{2}{3}, 1; \frac{5}{3}; \frac{i+x}{i-x}\right))}{6(1+ix)^{2/3}x^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((2*I)/3)*ArcTan[x])/x^3,x]

[Out] ((1 - I*x)^(2/3)*(-3 - (8*I)*x + 5*x^2 + 2*x^2*Hypergeometric2F1[2/3, 1, 5/3, (I + x)/(I - x)]))/(6*(1 + I*x)^(2/3)*x^2)

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{ix+1}{\sqrt{x^2+1}}\right)^{\frac{2}{3}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^3,x)

[Out] int(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^3,x, algorithm="maxima")

[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3)/x^3, x)

Fricas [A]

time = 1.86, size = 138, normalized size = 0.97

$$\frac{4x^2 \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - 1\right) + 2(-i\sqrt{3}x^2 - x^2) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} + \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right) + 2(i\sqrt{3}x^2 - x^2) \log\left(\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}} - \frac{1}{2}i\sqrt{3} + \frac{1}{2}\right) + 3(5x^2 + 2ix + 3)\left(\frac{i\sqrt{x^2+1}}{x+i}\right)^{\frac{2}{3}}}{18x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^3,x, algorithm="fricas")

[Out] -1/18*(4*x^2*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1) + 2*(-I*sqrt(3)*x^2 - x^2)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) + 1/2*I*sqrt(3) + 1/2) + 2*(I*sqrt(3)*x^2 - x^2)*log((I*sqrt(x^2 + 1)/(x + I))^(2/3) - 1/2*I*sqrt(3) + 1/2) + 3*(5*x^2 + 2*I*x + 3)*(I*sqrt(x^2 + 1)/(x + I))^(2/3))/x^2

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*x)/(x**2+1)**(1/2))**(2/3)/x**3,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*x)/(x^2+1)^(1/2))^(2/3)/x^3,x, algorithm="giac")``[Out] integrate(((I*x + 1)/sqrt(x^2 + 1))^(2/3)/x^3, x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\left(\frac{1+xi}{\sqrt{x^2+1}}\right)^{2/3}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3)/x^3,x)``[Out] int(((x*1i + 1)/(x^2 + 1)^(1/2))^(2/3)/x^3, x)`

3.128 $\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=741

$$\frac{11i(1-iax)^{7/8}\sqrt[8]{1+iax}}{32a^3} - \frac{i(1-iax)^{7/8}(1+iax)^{9/8}}{24a^3} + \frac{x(1-iax)^{7/8}(1+iax)^{9/8}}{3a^2} + \frac{11i\sqrt{2+\sqrt{2}}\text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}(1-iax)^{1/8}}{(1+iax)^{1/8}}\right)}{3a^2}$$

[Out] $-11/32*I*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)}/a^3-1/24*I*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)}/a^3+1/3*x*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)}/a^2+11/128*I*\arctan((-2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}/a^3-11/128*I*\arctan((2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}/a^3-11/256*I*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}-(1-I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)})*(2-2^{(1/2)})^{(1/2)}/a^3+11/256*I*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)})*(2-2^{(1/2)})^{(1/2)}/a^3+11/128*I*\arctan((-2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2+2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}/a^3-11/128*I*\arctan((2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2+2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}/a^3-11/256*I*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}-(1-I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)})*(2+2^{(1/2)})^{(1/2)}/a^3+11/256*I*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)})*(2+2^{(1/2)})^{(1/2)}/a^3$

Rubi [A]

time = 0.55, antiderivative size = 741, normalized size of antiderivative = 1.00, number of steps used = 27, number of rules used = 13, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.812$, Rules used = {5170, 92, 81, 52, 65, 338, 305, 1136, 1183, 648, 632, 210, 642}

$$\frac{11i\sqrt{2+\sqrt{2}}\text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}(1-iax)^{1/8}}{(1+iax)^{1/8}}\right)}{3a^2} - \frac{i(1-iax)^{7/8}(1+iax)^{9/8}}{24a^3} + \frac{x(1-iax)^{7/8}(1+iax)^{9/8}}{3a^2} + \frac{11i(1-iax)^{7/8}\sqrt[8]{1+iax}}{32a^3}$$

Antiderivative was successfully verified.

[In] Int[E^((I/4)*ArcTan[a*x])*x^2,x]

[Out] $(((-11*I)/32)*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)}/a^3 - ((I/24)*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)}/a^3 + (x*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)})/(3*a^2) + (((11*I)/128)*\text{Sqrt}[2+\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2-\text{Sqrt}[2]] - (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2+\text{Sqrt}[2]])]/a^3 + (((11*I)/128)*\text{Sqrt}[2-\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2+\text{Sqrt}[2]] - (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2-\text{Sqrt}[2]])]/a^3 - (((11*I)/128)*\text{Sqrt}[2+\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2-\text{Sqrt}[2]] + (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2+\text{Sqrt}[2]])]/a^3 - (((11*I)/128)*\text{Sqrt}[2-\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2+\text{Sqrt}[2]] + (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2-\text{Sqrt}[2]])]/a^3 - (((11*I)/$

256)*Sqrt[2 - Sqrt[2]]*Log[1 + (1 - I*a*x)^(1/4)/(1 + I*a*x)^(1/4) - (Sqrt[2 - Sqrt[2]]*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))]/a^3 + (((11*I)/256)*Sqrt[2 - Sqrt[2]]*Log[1 + (1 - I*a*x)^(1/4)/(1 + I*a*x)^(1/4) + (Sqrt[2 - Sqrt[2]]*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))]/a^3 - (((11*I)/256)*Sqrt[2 + Sqrt[2]]*Log[1 + (1 - I*a*x)^(1/4)/(1 + I*a*x)^(1/4) - (Sqrt[2 + Sqrt[2]]*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))]/a^3 + (((11*I)/256)*Sqrt[2 + Sqrt[2]]*Log[1 + (1 - I*a*x)^(1/4)/(1 + I*a*x)^(1/4) + (Sqrt[2 + Sqrt[2]]*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))]/a^3

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92

Int[((a_.) + (b_.)*(x_))^2*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &

& (LtQ[a, 0] || LtQ[b, 0])

Rule 305

```
Int[(x_)^(m_)/((a_) + (b_)*(x_)^(n_)), x_Symbol] := With[{r = Numerator[Rt[a/b, 4]], s = Denominator[Rt[a/b, 4]]}, Dist[s^3/(2*Sqrt[2]*b*r), Int[x^(m - n/4)/(r^2 - Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] - Dist[s^3/(2*Sqrt[2]*b*r), Int[x^(m - n/4)/(r^2 + Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] /; FreeQ[{a, b}, x] && IGtQ[n/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && GtQ[a/b, 0]
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1136

```
Int[((d_)*(x_))^(m_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[d^3*(d*x)^(m - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 1))), x] - Dist[d^4/(c*(m + 4*p + 1)), Int[(d*x)^(m - 4)*Simp[a*(m - 3) + b*(m + 2*p - 1)*x^2, x]*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 3] && NeQ[m + 4*p + 1, 0] && IntegerQ[2*p] && (IntegerQ[p] || IntegerQ[m])
```


Rule 1183

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] :
> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int
[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r +
(d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 83, normalized size = 0.11

$$\frac{(1 - iax)^{7/8} \left(7\sqrt[8]{1 + iax} (-i + 9ax + 8ia^2x^2) - 66i\sqrt[8]{2} {}_2F_1\left(-\frac{1}{8}, \frac{7}{8}; \frac{15}{8}; \frac{1}{2}(1 - iax)\right) \right)}{168a^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/4)*ArcTan[a*x])*x^2,x]

[Out] ((1 - I*a*x)^(7/8)*(7*(1 + I*a*x)^(1/8)*(-I + 9*a*x + (8*I)*a^2*x^2) - (66*I)*2^(1/8)*Hypergeometric2F1[-1/8, 7/8, 15/8, (1 - I*a*x)/2]))/(168*a^3)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{1}{4}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^2,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^2,x, algorithm="maxima")

[Out] integrate(x^2*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(1/4), x)

Fricas [A]

time = 2.37, size = 435, normalized size = 0.59

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^2,x, algorithm="fricas")

[Out] 1/96*(96*I*a^3*(14641/268435456*I/a^12)^(1/4)*log(128/11*a^3*(14641/268435456*I/a^12)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) - 96*a^3*(14641/268435456*I/a^12)^(1/4)*log(128/11*I*a^3*(14641/268435456*I/a^12)^(1/4) + (I

```
*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) + 96*a^3*(14641/268435456*I/a^12)^(1/4)
)*log(-128/11*I*a^3*(14641/268435456*I/a^12)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(
a*x + I))^(1/4)) - 96*I*a^3*(14641/268435456*I/a^12)^(1/4)*log(-128/11*a^3*
(14641/268435456*I/a^12)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) + 9
6*I*a^3*(-14641/268435456*I/a^12)^(1/4)*log(128/11*a^3*(-14641/268435456*I/
a^12)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) - 96*a^3*(-14641/26843
5456*I/a^12)^(1/4)*log(128/11*I*a^3*(-14641/268435456*I/a^12)^(1/4) + (I*sq
rt(a^2*x^2 + 1)/(a*x + I))^(1/4)) + 96*a^3*(-14641/268435456*I/a^12)^(1/4)*
log(-128/11*I*a^3*(-14641/268435456*I/a^12)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a
*x + I))^(1/4)) - 96*I*a^3*(-14641/268435456*I/a^12)^(1/4)*log(-128/11*a^3*
(-14641/268435456*I/a^12)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) +
(32*a^3*x^3 - 4*I*a^2*x^2 - a*x - 37*I)*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/
4))/a^3
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \sqrt[4]{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4)*x**2,x)

[Out] Integral(x**2*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(1/4), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \left(\frac{1 + ax \operatorname{li}}{\sqrt{a^2x^2 + 1}} \right)^{1/4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((a*x*li + 1)/(a^2*x^2 + 1)^(1/2))^(1/4),x)

[Out] int(x^2*((a*x*li + 1)/(a^2*x^2 + 1)^(1/2))^(1/4), x)

3.129 $\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x dx$

Optimal. Leaf size=689

$$\frac{(1-iax)^{7/8}\sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} - \frac{\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} - \frac{\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2}$$

[Out] $1/8*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)}/a^2+1/2*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)}/a^2-1/32*\arctan((-2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}/a^2+1/32*\arctan((2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2+2^{(1/2)})^{(1/2)})/(2-2^{(1/2)})^{(1/2)})*(2-2^{(1/2)})^{(1/2)}/a^2+1/64*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}-(1-I*a*x)^{(1/8)}*(2-2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)}*(2-2^{(1/2)})^{(1/2)}/a^2-1/64*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/8)}*(2-2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)}*(2-2^{(1/2)})^{(1/2)}/a^2-1/32*\arctan((-2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}/a^2+1/32*\arctan((2*(1-I*a*x)^{(1/8)}/(1+I*a*x)^{(1/8)}+(2-2^{(1/2)})^{(1/2)})/(2+2^{(1/2)})^{(1/2)})*(2+2^{(1/2)})^{(1/2)}/a^2+1/64*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}-(1-I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/a^2-1/64*\ln(1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)}+(1-I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/(1+I*a*x)^{(1/8)}*(2+2^{(1/2)})^{(1/2)}/a^2$

Rubi [A]

time = 0.38, antiderivative size = 689, normalized size of antiderivative = 1.00, number of steps used = 26, number of rules used = 12, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.857$, Rules used = {5170, 81, 52, 65, 338, 305, 1136, 1183, 648, 632, 210, 642}

$$\frac{\sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} - \frac{\sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} + \frac{\sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} - \frac{\sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} + \frac{(1-\text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)) \sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} - \frac{(1-\text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)) \sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} + \frac{\sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} - \frac{\sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} + \frac{\sqrt{2+\sqrt{2}} \arctan\left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} - \frac{\sqrt{2-\sqrt{2}} \arctan\left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2}$$

Antiderivative was successfully verified.

[In] Int[E^((I/4)*ArcTan[a*x])*x,x]

[Out] $((1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)})/(8*a^2) + ((1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)})/(2*a^2) - (\text{Sqrt}[2+\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2-\text{Sqrt}[2]] - (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2+\text{Sqrt}[2]])]/(32*a^2) - (\text{Sqrt}[2-\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2+\text{Sqrt}[2]] - (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2-\text{Sqrt}[2]])]/(32*a^2) + (\text{Sqrt}[2+\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2-\text{Sqrt}[2]] + (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2+\text{Sqrt}[2]])]/(32*a^2) + (\text{Sqrt}[2-\text{Sqrt}[2]]*\text{ArcTan}[(\text{Sqrt}[2+\text{Sqrt}[2]] + (2*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})/\text{Sqrt}[2-\text{Sqrt}[2]])]/(32*a^2) + (\text{Sqrt}[2-\text{Sqrt}[2]]*\text{Log}[1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)} - (\text{Sqrt}[2-\text{Sqrt}[2]]*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})]/(64*a^2) - (\text{Sqrt}[2-\text{Sqrt}[2]]*\text{Log}[1+(1-I*a*x)^{(1/4)}/(1+I*a*x)^{(1/4)} + (\text{Sqrt}[2-\text{Sqrt}[2]]*(1-I*a*x)^{(1/8)})/(1+I*a*x)^{(1/8)})]/(64*a^2)$

)]/(64*a^2) + (Sqrt[2 + Sqrt[2]]*Log[1 + (1 - I*a*x)^(1/4)/(1 + I*a*x)^(1/4) - (Sqrt[2 + Sqrt[2]]*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8)])/((64*a^2) - (Sqrt[2 + Sqrt[2]]*Log[1 + (1 - I*a*x)^(1/4)/(1 + I*a*x)^(1/4) + (Sqrt[2 + Sqrt[2]]*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8)])/((64*a^2)

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) + d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Denominator[m]] && IntLinearQ[a, b, c, d, m, n, x]

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 305

Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{r = Numerator[Rt[a/b, 4]], s = Denominator[Rt[a/b, 4]]}, Dist[s^3/(2*Sqrt[2]*b*r), Int[x^(m - n/4)/(r^2 - Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] - Dist[s^3/(2*Sqrt[2]*b*r), Int[x^(m - n/4)/(r^2 + Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] /; FreeQ[{a, b}, x] && IGtQ[n/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && GtQ[a/b, 0]

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1136

```
Int[((d_)*(x_))^(m_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[d^3*(d*x)^(m - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 1))), x] - Dist[d^4/(c*(m + 4*p + 1)), Int[(d*x)^(m - 4)*Simp[a*(m - 3) + b*(m + 2*p - 1)*x^2, x]*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 3] && NeQ[m + 4*p + 1, 0] && IntegerQ[2*p] && (IntegerQ[p] || IntegerQ[m])
```

Rule 1183

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{4}i \tan^{-1}(ax)} x dx &= \int \frac{x \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} dx \\
&= \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} - \frac{i \int \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} dx}{8a} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} - \frac{i \int \frac{1}{\sqrt[8]{1-iax} (1+iax)^{7/8}} dx}{32a} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} + \frac{\text{Subst}\left(\int \frac{x^6}{(2-x^8)^{7/8}} dx, x, \sqrt[8]{1-iax}\right)}{4a^2} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} + \frac{\text{Subst}\left(\int \frac{x^6}{1+x^8} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{4a^2} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} + \frac{\text{Subst}\left(\int \frac{x^4}{1-\sqrt{2}x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{8\sqrt{2}a^2} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} - \frac{\text{Subst}\left(\int \frac{1-\sqrt{2}x^2}{1-\sqrt{2}x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{8\sqrt{2}a^2} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} + \frac{\text{Subst}\left(\int \frac{\sqrt{2-\sqrt{2}}-(1-\sqrt{2})x}{1-\sqrt{2-\sqrt{2}}x+x^2} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{16\sqrt{2}(2-\sqrt{2})} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} + \frac{\sqrt{\frac{1}{2}(3-2\sqrt{2})} \text{Subst}\left(\int \frac{1}{1-\sqrt{2}x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{32a^2} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} + \frac{\sqrt{2-\sqrt{2}} \log\left(1 + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}}\right)}{64a^2} \\
&= \frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{8a^2} + \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2a^2} - \frac{\sqrt{2+\sqrt{2}} \tan^{-1}\left(\frac{\sqrt{2-\sqrt{2}}-x}{\sqrt{2+\sqrt{2}}+x}\right)}{32a^2}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 63, normalized size = 0.09

$$\frac{(1 - iax)^{7/8} \left(7(1 + iax)^{9/8} + 2\sqrt[8]{2} {}_2F_1\left(-\frac{1}{8}, \frac{7}{8}; \frac{15}{8}; \frac{1}{2}(1 - iax)\right) \right)}{14a^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/4)*ArcTan[a*x])*x,x]

[Out] ((1 - I*a*x)^(7/8)*(7*(1 + I*a*x)^(9/8) + 2*2^(1/8)*Hypergeometric2F1[-1/8, 7/8, 15/8, (1 - I*a*x)/2]))/(14*a^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{1}{4}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x,x, algorithm="maxima")

[Out] integrate(x*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(1/4), x)

Fricas [A]

time = 2.77, size = 428, normalized size = 0.62

.....

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x,x, algorithm="fricas")

[Out] -1/8*(8*a^2*(1/1048576*I/a^8)^(1/4)*log(32*a^2*(1/1048576*I/a^8)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) + 8*I*a^2*(1/1048576*I/a^8)^(1/4)*log(32*I*a^2*(1/1048576*I/a^8)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) -

$$8*I*a^2*(1/1048576*I/a^8)^{(1/4)}*\log(-32*I*a^2*(1/1048576*I/a^8)^{(1/4)} + (I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)} - 8*a^2*(1/1048576*I/a^8)^{(1/4)}*\log(-32*a^2*(1/1048576*I/a^8)^{(1/4)} + (I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)} + 8*a^2*(-1/1048576*I/a^8)^{(1/4)}*\log(32*a^2*(-1/1048576*I/a^8)^{(1/4)} + (I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)} + 8*I*a^2*(-1/1048576*I/a^8)^{(1/4)}*\log(32*I*a^2*(-1/1048576*I/a^8)^{(1/4)} + (I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)} - 8*I*a^2*(-1/1048576*I/a^8)^{(1/4)}*\log(-32*I*a^2*(-1/1048576*I/a^8)^{(1/4)} + (I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)} - 8*a^2*(-1/1048576*I/a^8)^{(1/4)}*\log(-32*a^2*(-1/1048576*I/a^8)^{(1/4)} + (I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)} - (4*a^2*x^2 - I*a*x + 5)*(I*\sqrt{a^2*x^2 + 1})/(a*x + I))^{(1/4)}/a^2$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^4 \sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4)*x,x)

[Out] Integral(x*(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(1/4), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \left(\frac{1 + ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}} \right)^{1/4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4),x)

[Out] int(x*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4), x)

3.130 $\int e^{\frac{1}{4}i\text{ArcTan}(ax)} dx$

Optimal. Leaf size=674

$$\frac{i(1-iax)^{7/8}\sqrt[8]{1+iax}}{a} - \frac{i\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)}{4a} - \frac{i\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}}{\sqrt[8]{1+iax}}\right)}{4a}$$

[Out] I*(1-I*a*x)^(7/8)*(1+I*a*x)^(1/8)/a-1/4*I*arctan((-2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2+2^(1/2))^(1/2))/(2-2^(1/2))^(1/2))*(2-2^(1/2))^(1/2)/a+1/4*I*arctan((2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2+2^(1/2))^(1/2))/(2-2^(1/2))^(1/2))*(2-2^(1/2))^(1/2)/a+1/8*I*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-(1-I*a*x)^(1/8)*(2-2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*(2-2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*(2-2^(1/2))^(1/2)/a-1/8*I*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/8)*(2-2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*(2-2^(1/2))^(1/2)/a-1/4*I*arctan((-2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2+2^(1/2))^(1/2))/(2+2^(1/2))^(1/2))*(2+2^(1/2))^(1/2)/a+1/4*I*arctan((2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2-2^(1/2))^(1/2))/(2+2^(1/2))^(1/2))*(2+2^(1/2))^(1/2)/a+1/8*I*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-(1-I*a*x)^(1/8)*(2+2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*(2+2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*(2+2^(1/2))^(1/2)/a-1/8*I*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/8)*(2+2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*(2+2^(1/2))^(1/2)/a

Rubi [A]

time = 0.33, antiderivative size = 674, normalized size of antiderivative = 1.00, number of steps used = 25, number of rules used = 11, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.917$, Rules used = {5169, 52, 65, 338, 305, 1136, 1183, 648, 632, 210, 642}

$$\frac{1}{\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)} - \frac{1}{\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)} - \frac{1}{\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)} - \frac{1}{\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)} - \frac{1}{\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)} - \frac{1}{\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)} - \frac{1}{\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)} - \frac{1}{\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)} - \frac{1}{\sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}\right)} - \frac{1}{\sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}} \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}$$

Antiderivative was successfully verified.

[In] Int[E^((I/4)*ArcTan[a*x]),x]

[Out] (I*(1-I*a*x)^(7/8)*(1+I*a*x)^(1/8))/a - ((I/4)*Sqrt[2+Sqrt[2]]*ArcTan[(Sqrt[2-Sqrt[2]] - (2*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8))/Sqrt[2+Sqrt[2]])/a - ((I/4)*Sqrt[2-Sqrt[2]]*ArcTan[(Sqrt[2+Sqrt[2]] - (2*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8))/Sqrt[2-Sqrt[2]])/a + ((I/4)*Sqrt[2+Sqrt[2]]*ArcTan[(Sqrt[2-Sqrt[2]] + (2*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8))/Sqrt[2+Sqrt[2]])/a + ((I/4)*Sqrt[2-Sqrt[2]]*ArcTan[(Sqrt[2+Sqrt[2]] + (2*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8))/Sqrt[2-Sqrt[2]])/a + ((I/8)*Sqrt[2-Sqrt[2]]*Log[1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4) - (Sqrt[2-Sqrt[2]]*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8)])/a - ((I/8)*Sqrt[2-Sqrt[2]]*Log[1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4) + (Sqrt[2-Sqrt[2]]*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8)])/a + ((I/8)*Sqrt[2+Sqrt[2]]*Log[1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4) - (Sqrt[2+Sqrt[2]]*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8)])/a + ((I/8)*Sqrt[2+Sqrt[2]]*Log[1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4) + (Sqrt[2+Sqrt[2]]*(1-I*a*x)^(1/8))/(1+I*a*x)^(1/8)])/a

$$\frac{1}{a} \left(\frac{(1 + I a x)^{1/4}}{(1 + I a x)^{1/8}} - \frac{\sqrt{2 + \sqrt{2}} (1 - I a x)^{1/8}}{(1 + I a x)^{1/4}} \right) + \frac{\sqrt{2 + \sqrt{2}} (1 - I a x)^{1/8}}{(1 + I a x)^{1/4}}$$
Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 305

```
Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{r = Numerator[R
t[a/b, 4]], s = Denominator[Rt[a/b, 4]]}, Dist[s^3/(2*sqrt[2]*b*r), Int[x^(
m - n/4)/(r^2 - sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] - Dist[s^3/(2*sq
rt[2]*b*r), Int[x^(m - n/4)/(r^2 + sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x],
x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && G
tQ[a/b, 0]
```

Rule 338

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 632

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[I
nt[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c},
```

x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 648

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]

Rule 1136

Int[((d_)*(x_))^(m_)*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[d^3*(d*x)^(m - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 1))), x] - Dist[d^4/(c*(m + 4*p + 1)), Int[(d*x)^(m - 4)*Simp[a*(m - 3) + b*(m + 2*p - 1)*x^2, x]*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 3] && NeQ[m + 4*p + 1, 0] && IntegerQ[2*p] && (IntegerQ[p] || IntegerQ[m])

Rule 1183

Int[((d_) + (e_)*(x_)^2)/((a_) + (b_)*(x_)^2 + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r + (d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]

Rule 5169

Int[E^(ArcTan[(a_)*(x_)])*(n_), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{4}i \tan^{-1}(ax)} dx &= \int \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} dx \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{1}{4} \int \frac{1}{\sqrt[8]{1-iax} (1+iax)^{7/8}} dx \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{(2i) \text{Subst} \left(\int \frac{x^6}{(2-x^8)^{7/8}} dx, x, \sqrt[8]{1-iax} \right)}{a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{(2i) \text{Subst} \left(\int \frac{x^6}{1+x^8} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{i \text{Subst} \left(\int \frac{x^4}{1-\sqrt{2} x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{\sqrt{2} a} - \frac{i \text{Subst} \left(\int \frac{x^4}{1+\sqrt{2} x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{\sqrt{2} a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} - \frac{i \text{Subst} \left(\int \frac{1-\sqrt{2} x^2}{1-\sqrt{2} x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{\sqrt{2} a} + \frac{i \text{Subst} \left(\int \frac{1+\sqrt{2} x^2}{1+\sqrt{2} x^2+x^4} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{\sqrt{2} a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{i \text{Subst} \left(\int \frac{\sqrt{2-\sqrt{2}} - (1-\sqrt{2})x}{1-\sqrt{2-\sqrt{2}} x+x^2} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{2\sqrt{2} (2-\sqrt{2}) a} + \frac{i \text{Subst} \left(\int \frac{\sqrt{2+\sqrt{2}} - (1+\sqrt{2})x}{1+\sqrt{2+\sqrt{2}} x+x^2} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{2\sqrt{2} (2+\sqrt{2}) a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{\left(i\sqrt{\frac{1}{2}} (3-2\sqrt{2}) \right) \text{Subst} \left(\int \frac{1}{1-\sqrt{2+\sqrt{2}} x+x^2} dx, x, \frac{\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{4a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} + \frac{i\sqrt{2-\sqrt{2}} \log \left(1 + \frac{\sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} - \frac{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}}{\sqrt[8]{1+iax}} \right)}{8a} \\
&= \frac{i(1-iax)^{7/8} \sqrt[8]{1+iax}}{a} - \frac{i\sqrt{2+\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2-\sqrt{2}} - \sqrt[8]{1-iax}}{\sqrt{2+\sqrt{2}} \sqrt[8]{1+iax}} \right)}{4a} - \frac{i\sqrt{2-\sqrt{2}} \tan^{-1} \left(\frac{\sqrt{2+\sqrt{2}} - \sqrt[8]{1+iax}}{\sqrt{2-\sqrt{2}} \sqrt[8]{1-iax}} \right)}{4a}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 41, normalized size = 0.06

$$\frac{16ie^{\frac{9}{4}i\text{ArcTan}(ax)} {}_2F_1\left(\frac{9}{8}, 2; \frac{17}{8}; -e^{2i\text{ArcTan}(ax)}\right)}{9a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/4)*ArcTan[a*x]), x]

[Out] (((-16*I)/9)*E^(((9*I)/4)*ArcTan[a*x])*Hypergeometric2F1[9/8, 2, 17/8, -E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{1}{4}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4), x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4), x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(1/4), x)

Fricas [A]

time = 2.14, size = 383, normalized size = 0.57

$$\frac{-i\sqrt[4]{a}\log\left(\frac{1+iax}{\sqrt{a^2x^2+1}}\right) + i\sqrt[4]{a}\log\left(\frac{1-iax}{\sqrt{a^2x^2+1}}\right) - i\sqrt[4]{a}\log\left(\frac{1+iax}{\sqrt{a^2x^2+1}}\right) + \left(\frac{\sqrt{a^2x^2+1}}{\sqrt{a^2x^2+1}}\right) + i\sqrt[4]{a}\log\left(\frac{1-iax}{\sqrt{a^2x^2+1}}\right) + \left(\frac{\sqrt{a^2x^2+1}}{\sqrt{a^2x^2+1}}\right) + i\sqrt[4]{a}\log\left(\frac{1+iax}{\sqrt{a^2x^2+1}}\right) - i\sqrt[4]{a}\log\left(\frac{1-iax}{\sqrt{a^2x^2+1}}\right) + \left(\frac{\sqrt{a^2x^2+1}}{\sqrt{a^2x^2+1}}\right) + i\sqrt[4]{a}\log\left(\frac{1-iax}{\sqrt{a^2x^2+1}}\right) + \left(\frac{\sqrt{a^2x^2+1}}{\sqrt{a^2x^2+1}}\right) - i\sqrt[4]{a}\log\left(\frac{1+iax}{\sqrt{a^2x^2+1}}\right) - i\sqrt[4]{a}\log\left(\frac{1-iax}{\sqrt{a^2x^2+1}}\right) + \left(\frac{\sqrt{a^2x^2+1}}{\sqrt{a^2x^2+1}}\right)}{4\sqrt[4]{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4), x, algorithm="fricas")

[Out] (-I*a*(1/256*I/a^4)^(1/4)*log(4*a*(1/256*I/a^4)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) + a*(1/256*I/a^4)^(1/4)*log(4*I*a*(1/256*I/a^4)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) - a*(1/256*I/a^4)^(1/4)*log(-4*I*a*(1/256*I/a^4)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) + I*a*(1/256*I/a^4)^(1/4)*log(-4*a*(1/256*I/a^4)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4)) - I*a*(-1/256*I/a^4)^(1/4)*log(4*a*(-1/256*I/a^4)^(1/4) + (I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4))

$$2x^2 + 1)/(ax + I)^{(1/4)} + a(-1/256I/a^4)^{(1/4)} \log(4Ia(-1/256I/a^4)^{(1/4)} + (I\sqrt{a^2x^2 + 1})/(ax + I))^{(1/4)} - a(-1/256I/a^4)^{(1/4)} \log(-4Ia(-1/256I/a^4)^{(1/4)} + (I\sqrt{a^2x^2 + 1})/(ax + I))^{(1/4)} + Ia(-1/256I/a^4)^{(1/4)} \log(-4a(-1/256I/a^4)^{(1/4)} + (I\sqrt{a^2x^2 + 1})/(ax + I))^{(1/4)} + (ax + I)(I\sqrt{a^2x^2 + 1})/(ax + I)^{(1/4)}/a$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[4]{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4), x)

[Out] Integral(((I*a*x + 1)/sqrt(a**2*x**2 + 1))**(1/4), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4), x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \left(\frac{1 + ax \operatorname{li}}{\sqrt{a^2x^2 + 1}} \right)^{1/4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4), x)

[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4), x)

$$3.131 \quad \int \frac{e^{\frac{1}{4}i\text{ArcTan}(ax)}}{x} dx$$

Optimal. Leaf size=859

$$-2\text{ArcTan}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \sqrt{2+\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2-\sqrt{2}} - \frac{2\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}}{\sqrt{2+\sqrt{2}}}\right) + \sqrt{2-\sqrt{2}} \text{ArcTan}\left(\frac{\sqrt{2+\sqrt{2}}}{\sqrt{2-\sqrt{2}}}\right)$$

```
[Out] -2*arctan((1+I*a*x)^(1/8)/(1-I*a*x)^(1/8))-2*arctanh((1+I*a*x)^(1/8)/(1-I*a*x)^(1/8))+1/2*ln(1+(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)-(1+I*a*x)^(1/8)*2^(1/2)/(1-I*a*x)^(1/8))*2^(1/2)-1/2*ln(1+(1+I*a*x)^(1/4)/(1-I*a*x)^(1/4)+(1+I*a*x)^(1/8)*2^(1/2)/(1-I*a*x)^(1/8))*2^(1/2)+arctan(1-(1+I*a*x)^(1/8)*2^(1/2)/(1-I*a*x)^(1/8))*2^(1/2)-arctan(1+(1+I*a*x)^(1/8)*2^(1/2)/(1-I*a*x)^(1/8))*2^(1/2)+arctan((-2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2+2^(1/2))^(1/2))/(2-2^(1/2))^(1/2))*2^(1/2)-arctan((2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2+2^(1/2))^(1/2))/(2-2^(1/2))^(1/2))*2^(1/2)-1/2*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-(1-I*a*x)^(1/8)*(2-2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*2^(1/2)-1/2*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/8)*(2-2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*2^(1/2)+arctan((-2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2-2^(1/2))^(1/2))/(2+2^(1/2))^(1/2))*2^(1/2)-arctan((2*(1-I*a*x)^(1/8)/(1+I*a*x)^(1/8)+(2-2^(1/2))^(1/2))/(2+2^(1/2))^(1/2))*2^(1/2)-1/2*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)-(1-I*a*x)^(1/8)*(2+2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*2^(1/2)+1/2*ln(1+(1-I*a*x)^(1/4)/(1+I*a*x)^(1/4)+(1-I*a*x)^(1/8)*(2+2^(1/2))^(1/2)/(1+I*a*x)^(1/8))*2^(1/2))^(1/2)
```

Rubi [A]

time = 0.41, antiderivative size = 859, normalized size of antiderivative = 1.00, number of steps used = 39, number of rules used = 20, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 1.250$, Rules used = {5170, 132, 65, 338, 305, 1136, 1183, 648, 632, 210, 642, 95, 220, 218, 212, 209, 217, 1179, 1176, 631}

Antiderivative was successfully verified.

[In] Int[E^((I/4)*ArcTan[a*x])/x,x]

```
[Out] -2*ArcTan[(1 + I*a*x)^(1/8)/(1 - I*a*x)^(1/8)] + Sqrt[2 + Sqrt[2]]*ArcTan[(Sqrt[2 - Sqrt[2]] - (2*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))/Sqrt[2 + Sqrt[2]]] + Sqrt[2 - Sqrt[2]]*ArcTan[(Sqrt[2 + Sqrt[2]] - (2*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))/Sqrt[2 - Sqrt[2]]] - Sqrt[2 + Sqrt[2]]*ArcTan[(Sqrt[2 - Sqrt[2]] + (2*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))/Sqrt[2 + Sqrt[2]]] - Sqrt[2 - Sqrt[2]]*ArcTan[(Sqrt[2 + Sqrt[2]] + (2*(1 - I*a*x)^(1/8))/(1 + I*a*x)^(1/8))/Sqrt[2 - Sqrt[2]]]
```

$$\begin{aligned} & x^{1/8})/\sqrt{2 - \sqrt{2}}] + \sqrt{2} \operatorname{ArcTan}[1 - (\sqrt{2}*(1 + I*a*x)^{1/8}) \\ &)/(1 - I*a*x)^{1/8}] - \sqrt{2} \operatorname{ArcTan}[1 + (\sqrt{2}*(1 + I*a*x)^{1/8})/(1 - \\ & I*a*x)^{1/8}] - 2 \operatorname{ArcTanh}[(1 + I*a*x)^{1/8}/(1 - I*a*x)^{1/8}] - (\sqrt{2 - \sqrt{2}} \\ & \sqrt{2}) * \operatorname{Log}[1 + (1 - I*a*x)^{1/4}/(1 + I*a*x)^{1/4}] - (\sqrt{2 - \sqrt{2}}) * \\ & (1 - I*a*x)^{1/8})/(1 + I*a*x)^{1/8}))/2 + (\sqrt{2 - \sqrt{2}}) * \operatorname{Log}[1 + (1 - \\ & I*a*x)^{1/4}/(1 + I*a*x)^{1/4} + (\sqrt{2 - \sqrt{2}}) * (1 - I*a*x)^{1/8})/(1 + \\ & I*a*x)^{1/8}))/2 - (\sqrt{2 + \sqrt{2}}) * \operatorname{Log}[1 + (1 - I*a*x)^{1/4}/(1 + I*a*x) \\ &)^{1/4} - (\sqrt{2 + \sqrt{2}}) * (1 - I*a*x)^{1/8})/(1 + I*a*x)^{1/8}))/2 + (\sqrt{2 + \sqrt{2}}) * \\ & \operatorname{Log}[1 + (1 - I*a*x)^{1/4}/(1 + I*a*x)^{1/4} + (\sqrt{2 + \sqrt{2}}) * (1 - I*a*x)^{1/8})/(1 + I*a*x) \\ &)^{1/4} - (\sqrt{2 + \sqrt{2}}) * (1 - I*a*x)^{1/8})/(1 + I*a*x)^{1/8}))/2 + \operatorname{Log}[1 - (\sqrt{2}*(1 + I*a*x) \\ &)^{1/8})/(1 - I*a*x)^{1/8} + (1 + I*a*x)^{1/4}/(1 - I*a*x)^{1/4}]/\sqrt{2} \\ & - \operatorname{Log}[1 + (\sqrt{2}*(1 + I*a*x)^{1/8})/(1 - I*a*x)^{1/8} + (1 + I*a*x)^{1/4} \\ & / (1 - I*a*x)^{1/4}]/\sqrt{2} \end{aligned}$$

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 95

```
Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x
_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1)
- 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)
], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n]
&& LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 132

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x
_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x
)^(m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(
a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; Fre
eQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (
GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*A
rcTan[Rt[b, 2]*(x/Rt[a, 2]), x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a
, 0] || GtQ[b, 0])
```

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 212

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

Rule 217

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 218

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 220

Int[((a_) + (b_.)*(x_)^(n_))(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^(n/2)), x], x] + Dist[r/(2*a), Int[1/(r + s*x^(n/2)), x], x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 1] && !GtQ[a/b, 0]

Rule 305

Int[(x_)^(m_.)/((a_) + (b_.)*(x_)^(n_)), x_Symbol] := With[{r = Numerator[Rt[a/b, 4]], s = Denominator[Rt[a/b, 4]]}, Dist[s^3/(2*Sqrt[2]*b*r), Int[x^(m - n/4)/(r^2 - Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x] - Dist[s^3/(2*Sqrt[2]*b*r), Int[x^(m - n/4)/(r^2 + Sqrt[2]*r*s*x^(n/4) + s^2*x^(n/2)), x], x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 0] && IGtQ[m, 0] && LtQ[m, n - 1] && GtQ[a/b, 0]

Rule 338

Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m + 1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2]

$^{-(-1)}$ && IntegersQ[m, p + (m + 1)/n]

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 632

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := Dist[-2, Subst[Int[1/Simp[b^2 - 4*a*c - x^2, x], x], x, b + 2*c*x], x] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 648

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Dist[(2*c*d - b*e)/(2*c), Int[1/(a + b*x + c*x^2), x], x] + Dist[e/(2*c), Int[(b + 2*c*x)/(a + b*x + c*x^2), x], x] /; FreeQ[{a, b, c, d, e}, x] && NeQ[2*c*d - b*e, 0] && NeQ[b^2 - 4*a*c, 0] && !NiceSqrtQ[b^2 - 4*a*c]
```

Rule 1136

```
Int[((d_)*(x_)^(m_))*((a_) + (b_)*(x_)^2 + (c_)*(x_)^4)^(p_), x_Symbol] := Simp[d^3*(d*x)^(m - 3)*((a + b*x^2 + c*x^4)^(p + 1)/(c*(m + 4*p + 1))), x] - Dist[d^4/(c*(m + 4*p + 1)), Int[(d*x)^(m - 4)*Simp[a*(m - 3) + b*(m + 2*p - 1)*x^2, x]*(a + b*x^2 + c*x^4)^p, x], x] /; FreeQ[{a, b, c, d, p}, x] && NeQ[b^2 - 4*a*c, 0] && GtQ[m, 3] && NeQ[m + 4*p + 1, 0] && IntegerQ[2*p] && (IntegerQ[p] || IntegerQ[m])
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 1183

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (b_.)*(x_)^2 + (c_.)*(x_)^4), x_Symbol] :
> With[{q = Rt[a/c, 2]}, With[{r = Rt[2*q - b/c, 2]}, Dist[1/(2*c*q*r), Int
[(d*r - (d - e*q)*x)/(q - r*x + x^2), x], x] + Dist[1/(2*c*q*r), Int[(d*r +
(d - e*q)*x)/(q + r*x + x^2), x], x]]] /; FreeQ[{a, b, c, d, e}, x] && NeQ
[b^2 - 4*a*c, 0] && NeQ[c*d^2 - b*d*e + a*e^2, 0] && NegQ[b^2 - 4*a*c]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{4}i \tan^{-1}(ax)}}{x} dx &= \int \frac{\sqrt[8]{1+iax}}{x\sqrt[8]{1-iax}} dx \\
&= (ia) \int \frac{1}{\sqrt[8]{1-iax} (1+iax)^{7/8}} dx + \int \frac{1}{x\sqrt[8]{1-iax} (1+iax)^{7/8}} dx \\
&= -\left(8\text{Subst}\left(\int \frac{x^6}{(2-x^8)^{7/8}} dx, x, \sqrt[8]{1-iax}\right)\right) + 8\text{Subst}\left(\int \frac{1}{-1+x^8} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) \\
&= -\left(4\text{Subst}\left(\int \frac{1}{1-x^4} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)\right) - 4\text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - 8 \\
&= -\left(2\text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)\right) - 2\text{Subst}\left(\int \frac{1}{1+x^2} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - 2 \\
&= -2 \tan^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \frac{\text{Subst}\left(\int \frac{\sqrt{2}+2x}{-1-\sqrt{2}x-x^2} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}{\sqrt{2}} \\
&= -2 \tan^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - 2 \tanh^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \frac{\log\left(1 - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} + \frac{\sqrt[4]{1+iax}}{\sqrt[4]{1-iax}}\right)}{\sqrt{2}} \\
&= -2 \tan^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - \sqrt{2} \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) \\
&= -2 \tan^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - \sqrt{2} \tan^{-1}\left(1 + \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) \\
&= -2 \tan^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \sqrt{2+\sqrt{2}} \tan^{-1}\left(\frac{\sqrt{2-\sqrt{2}} - \frac{2\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}}{\sqrt{2+\sqrt{2}}}\right) + \sqrt{2-\sqrt{2}} \tan^{-1}\left(\frac{\sqrt{2-\sqrt{2}} - \frac{2\sqrt[8]{1-iax}}{\sqrt[8]{1+iax}}}{\sqrt{2-\sqrt{2}}}\right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 97, normalized size = 0.11

$$\frac{4(1-iax)^{7/8} \left(\sqrt[8]{2} (1+iax)^{7/8} {}_2F_1\left(\frac{7}{8}, \frac{7}{8}; \frac{15}{8}; \frac{1}{2}(1-iax)\right) + 2 {}_2F_1\left(\frac{7}{8}, 1; \frac{15}{8}; \frac{i+ax}{i-ax}\right) \right)}{7(1+iax)^{7/8}}$$

Antiderivative was successfully verified.

$t(a^2x^2 + 1)/(ax + I)^{(1/4)} - (-I)^{(1/4)} \log(-(-I)^{(1/4)} + (I\sqrt{a^2x^2 + 1})/(ax + I)^{(1/4)}) - \log((I\sqrt{a^2x^2 + 1})/(ax + I)^{(1/4)} + 1) - I \log((I\sqrt{a^2x^2 + 1})/(ax + I)^{(1/4)} + I) + I \log((I\sqrt{a^2x^2 + 1})/(ax + I)^{(1/4)} - I) + \log((I\sqrt{a^2x^2 + 1})/(ax + I)^{(1/4)} - 1)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[4]{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4)/x,x)

[Out] Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(1/4)/x, x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2x^2 + 1}}\right)^{1/4}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*I + 1)/(a^2*x^2 + 1)^(1/2))^(1/4)/x,x)

[Out] int(((a*x*I + 1)/(a^2*x^2 + 1)^(1/2))^(1/4)/x, x)

$$3.132 \quad \int \frac{e^{\frac{1}{4}i\text{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=328

$$-\frac{(1-iax)^{7/8}\sqrt[8]{1+iax}}{x} - \frac{1}{2}ia\text{ArcTan}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) + \frac{ia\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}{2\sqrt{2}} - \frac{ia\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}{2\sqrt{2}}$$

[Out] $-(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)}/x-1/2*I*a*\arctan((1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)})-1/2*I*a*\operatorname{arctanh}((1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)})+1/4*I*a*\arctan(1-(1+I*a*x)^{(1/8)*2^{(1/2)}}/(1-I*a*x)^{(1/8))*2^{(1/2)}-1/4*I*a*\arctan(1+(1+I*a*x)^{(1/8)*2^{(1/2)}}/(1-I*a*x)^{(1/8))*2^{(1/2)}+1/8*I*a*\ln(1+(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}-(1+I*a*x)^{(1/8)*2^{(1/2)}}/(1-I*a*x)^{(1/8))*2^{(1/2)}-1/8*I*a*\ln(1+(1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}+(1+I*a*x)^{(1/8)*2^{(1/2)}}/(1-I*a*x)^{(1/8))*2^{(1/2)})$

Rubi [A]

time = 0.09, antiderivative size = 328, normalized size of antiderivative = 1.00, number of steps used = 16, number of rules used = 13, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.812$, Rules used = {5170, 96, 95, 220, 218, 212, 209, 217, 1179, 642, 1176, 631, 210}

$$-\frac{1}{2}ia\text{ArcTan}\left(\frac{\sqrt{1+iax}}{\sqrt{1-iax}}\right) + \frac{ia\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{1+iax}}{\sqrt{1-iax}}\right)}{2\sqrt{2}} - \frac{ia\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt{1+iax}}{\sqrt{1-iax}}\right)}{2\sqrt{2}} - \frac{(1-iax)^{7/8}\sqrt{1+iax}}{x} + \frac{ia\log\left(\frac{\sqrt{1+iax}}{\sqrt{1-iax}} - \frac{\sqrt{2}\sqrt{1+iax}}{\sqrt{1-iax}} + 1\right)}{4\sqrt{2}} - \frac{ia\log\left(\frac{\sqrt{1+iax}}{\sqrt{1-iax}} + \frac{\sqrt{2}\sqrt{1+iax}}{\sqrt{1-iax}} + 1\right)}{4\sqrt{2}} - \frac{1}{2}ia\operatorname{tanh}^{-1}\left(\frac{\sqrt{1+iax}}{\sqrt{1-iax}}\right)$$

Antiderivative was successfully verified.

[In] Int[E^((I/4)*ArcTan[a*x])/x^2,x]

[Out] $-(((1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)})/x) - (I/2)*a*\text{ArcTan}[(1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)}] + ((I/2)*a*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)}])/\text{Sqrt}[2] - ((I/2)*a*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)}])/\text{Sqrt}[2] - (I/2)*a*\text{ArcTanh}[(1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)}] + ((I/4)*a*\text{Log}[1 - (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)} + (1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/\text{Sqrt}[2] - ((I/4)*a*\text{Log}[1 + (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)} + (1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/\text{Sqrt}[2]$

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

```

Rule 209

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

```

Rule 210

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1)*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

```

Rule 212

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])

```

Rule 217

```

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

```

Rule 218

```

Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

```

Rule 220

```

Int[((a_) + (b_.)*(x_)^(n_))^(n_ - 1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^(n/2)), x], x] + Dist[r/(2*a), Int[1/(r + s*x^(n/2)), x], x]] /; FreeQ[{a, b}, x] && IGtQ[n/4, 1] && !GtQ[a/b, 0]

```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5170

```
Int[E^(ArcTan[(a_)*(x_)])*(n_)*(x_)^(m_), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{4}i \tan^{-1}(ax)}}{x^2} dx &= \int \frac{\sqrt[8]{1+iax}}{x^2 \sqrt[8]{1-iax}} dx \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} + \frac{1}{4}(ia) \int \frac{1}{x \sqrt[8]{1-iax} (1+iax)^{7/8}} dx \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} + (2ia) \text{Subst} \left(\int \frac{1}{-1+x^8} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} - (ia) \text{Subst} \left(\int \frac{1}{1-x^4} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) - (ia) \text{Subst} \left(\int \frac{1}{1-x^4} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} - \frac{1}{2}(ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) - \frac{1}{2}(ia) \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} - \frac{1}{2} ia \tan^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) - \frac{1}{2} ia \tanh^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) - \frac{1}{4} ia \tanh^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} - \frac{1}{2} ia \tan^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) - \frac{1}{2} ia \tanh^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) + \frac{1}{4} ia \tanh^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{(1-iax)^{7/8} \sqrt[8]{1+iax}}{x} - \frac{1}{2} ia \tan^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) + \frac{ia \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right)}{2\sqrt{2}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 71, normalized size = 0.22

$$-\frac{i(1-iax)^{7/8} (-7i + 7ax + 2ax {}_2F_1(\frac{7}{8}, 1; \frac{15}{8}; \frac{i+ax}{i-ax}))}{7x(1+iax)^{7/8}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/4)*ArcTan[a*x])/x^2,x]

[Out] ((-1/7*I)*(1 - I*a*x)^(7/8)*(-7*I + 7*a*x + 2*a*x*Hypergeometric2F1[7/8, 1, 15/8, (I + a*x)/(I - a*x)]))/(x*(1 + I*a*x)^(7/8))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}} \right)^{\frac{1}{4}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^2,x)`

[Out] `int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^2,x, algorithm="maxima")`

[Out] `integrate((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(1/4)/x^2, x)`

Fricas [A]

time = 2.21, size = 345, normalized size = 1.05

$$\frac{-i \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) + \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) - \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) + i \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) + \sqrt{a^2 x^2 + 1} \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) - \sqrt{a^2 x^2 + 1} \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) + \sqrt{a^2 x^2 + 1} \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) - \sqrt{a^2 x^2 + 1} \operatorname{arctan}\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right) - 4(-i a x + 1)\left(\frac{\sqrt{a^2 x^2 + 1}}{a x}\right)^{\frac{1}{4}}}{4 x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^2,x, algorithm="fricas")`

[Out] $\frac{1}{4} * (-I * a * x * \log((I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} + 1) + a * x * \log((I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} + I) - a * x * \log((I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} - I) + I * a * x * \log((I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} - 1) + \sqrt{a^2 * x^2 + 1} * x * \log((a * (I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} + I * \sqrt{a^2 * x^2 + 1}) / a) - \sqrt{a^2 * x^2 + 1} * x * \log((a * (I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} - I * \sqrt{a^2 * x^2 + 1}) / a) + \sqrt{a^2 * x^2 + 1} * x * \log((a * (I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} + I * \sqrt{a^2 * x^2 + 1}) / a) - \sqrt{a^2 * x^2 + 1} * x * \log((a * (I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} - I * \sqrt{a^2 * x^2 + 1}) / a) - 4 * (-I * a * x + 1) * (I * \sqrt{a^2 * x^2 + 1}) / (a * x + I))^{\frac{1}{4}} / x$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[4]{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4)/x**2,x)`

[Out] `Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(1/4)/x**2, x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{1/4}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4)/x^2,x)

[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4)/x^2, x)

3.133 $\int \frac{e^{\frac{1}{4}i\text{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=364

$$\frac{ia(1-iax)^{7/8}\sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{16}a^2\text{ArcTan}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - \frac{a^2\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}{16\sqrt{2}}$$

[Out] $-1/8*I*a*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)}/x-1/2*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)}/x^2+1/16*a^2*\arctan((1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)})+1/16*a^2*\operatorname{arctanh}((1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)})-1/32*a^2*\arctan(1-(1+I*a*x)^{(1/8)}*2^{(1/2)})/(1-I*a*x)^{(1/8)}*2^{(1/2)}+1/32*a^2*\arctan(1+(1+I*a*x)^{(1/8)}*2^{(1/2)})/(1-I*a*x)^{(1/8)}*2^{(1/2)}-1/64*a^2*\ln(1+(1+I*a*x)^{(1/4)})/(1-I*a*x)^{(1/4)}-(1+I*a*x)^{(1/8)}*2^{(1/2)}/(1-I*a*x)^{(1/8)}*2^{(1/2)}+1/64*a^2*\ln(1+(1+I*a*x)^{(1/4)})/(1-I*a*x)^{(1/4)}+(1+I*a*x)^{(1/8)}*2^{(1/2)}/(1-I*a*x)^{(1/8)}*2^{(1/2)}$

Rubi [A]

time = 0.12, antiderivative size = 364, normalized size of antiderivative = 1.00, number of steps used = 17, number of rules used = 14, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.875$, Rules used = {5170, 98, 96, 95, 220, 218, 212, 209, 217, 1179, 642, 1176, 631, 210}

$$\frac{1}{16}a^2\text{ArcTan}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - \frac{a^2\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}{16\sqrt{2}} + \frac{a^2\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right)}{16\sqrt{2}} - \frac{a^2\log\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} - \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} + 1\right)}{32\sqrt{2}} + \frac{a^2\log\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} + \frac{\sqrt{2}\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} + 1\right)}{32\sqrt{2}} + \frac{1}{16}a^2\operatorname{tanh}^{-1}\left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}}\right) - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} - \frac{ia(1-iax)^{7/8}\sqrt[8]{1+iax}}{8x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I/4)*\text{ArcTan}[a*x]}/x^3, x]$

[Out] $((-1/8*I)*a*(1-I*a*x)^{(7/8)}*(1+I*a*x)^{(1/8)}/x - ((1-I*a*x)^{(7/8)}*(1+I*a*x)^{(9/8)})/(2*x^2) + (a^2*\text{ArcTan}[(1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)}])/16 - (a^2*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)}])/ (16*\text{Sqrt}[2]) + (a^2*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)}])/ (16*\text{Sqrt}[2]) + (a^2*\text{ArcTanh}[(1+I*a*x)^{(1/8)}/(1-I*a*x)^{(1/8)}])/16 - (a^2*\text{Log}[1 - (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)} + (1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/ (32*\text{Sqrt}[2]) + (a^2*\text{Log}[1 + (\text{Sqrt}[2]*(1+I*a*x)^{(1/8)})/(1-I*a*x)^{(1/8)} + (1+I*a*x)^{(1/4)}/(1-I*a*x)^{(1/4)}])/ (32*\text{Sqrt}[2])$

Rule 95

$\text{Int}[\frac{(a_1 + b_1*x_1)^{m_1}*(c_1 + d_1*x_1)^{n_1}}{(e_1 + f_1*x_1)}, x_Symbol] := \text{With}[\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \text{FreeQ}[\{a, b, c, d, e, f\}, x] \&\& \text{EqQ}[m + n + 1, 0] \&\& \text{RationalQ}[n] \&\& \text{LtQ}[-1, m, 0] \&\& \text{SimplerQ}[a + b*x, c + d*x]$

Rule 96

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]
```

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 209

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])
```

Rule 212

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
```


+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 220

Int[((a_) + (b_.)*(x_)^(n_))^(n_ - 1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^(n/2)), x], x] + Dist[r/(2*a), Int[1/(r + s*x^(n/2)), x], x] /; FreeQ[{a, b}, x] && IGtQ[n/4, 1] && !GtQ[a/b, 0]

Rule 631

Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(n_ - 1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c]) /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)^(n_.)]*(x_)^(m_.)), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{4}i \tan^{-1}(ax)}}{x^3} dx &= \int \frac{\sqrt[8]{1+iax}}{x^3 \sqrt[8]{1-iax}} dx \\
&= -\frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{8}(ia) \int \frac{\sqrt[8]{1+iax}}{x^2 \sqrt[8]{1-iax}} dx \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} - \frac{1}{32}a^2 \int \frac{1}{x \sqrt[8]{1-iax} (1+iax)^{7/8}} dx \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} - \frac{1}{4}a^2 \text{Subst} \left(\int \frac{1}{-1+x^8} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{8}a^2 \text{Subst} \left(\int \frac{1}{1-x^4} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{16}a^2 \text{Subst} \left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{16}a^2 \tan^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) + \frac{1}{16}a^2 \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{16}a^2 \tan^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) + \frac{1}{16}a^2 \\
&= -\frac{ia(1-iax)^{7/8} \sqrt[8]{1+iax}}{8x} - \frac{(1-iax)^{7/8}(1+iax)^{9/8}}{2x^2} + \frac{1}{16}a^2 \tan^{-1} \left(\frac{\sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} \right) - \frac{1}{16}a^2
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 84, normalized size = 0.23

$$\frac{(1-iax)^{7/8} (7(-4-9iax+5a^2x^2) + 2a^2x^2 {}_2F_1(\frac{7}{8}, 1; \frac{15}{8}; \frac{i+ax}{i-ax}))}{56x^2(1+iax)^{7/8}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/4)*ArcTan[a*x])/x^3,x]

[Out] ((1 - I*a*x)^(7/8)*(7*(-4 - (9*I)*a*x + 5*a^2*x^2) + 2*a^2*x^2*Hypergeometric2F1[7/8, 1, 15/8, (I + a*x)/(I - a*x)]))/(56*x^2*(1 + I*a*x)^(7/8))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{1}{4}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^3,x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^3,x, algorithm="maxima")

[Out] integrate(((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(1/4)/x^3, x)

Fricas [A]

time = 2.80, size = 381, normalized size = 1.05

$$\frac{a^2 x^2 \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} + 1\right) + i a^2 x^2 \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} + i\right) - i a^2 x^2 \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} - i\right) - a^2 x^2 \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} - 1\right) + \sqrt{a^2 x^2} \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} + \sqrt{-i a^2}\right) - \sqrt{a^2 x^2} \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} - \sqrt{-i a^2}\right) + \sqrt{-i a^2 x^2} \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} + \sqrt{-i a^2}\right) - \sqrt{-i a^2 x^2} \log\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I} - \sqrt{-i a^2}\right) - 4(2 a^2 x^2 + i a x + 4)\left(\frac{\sqrt{a^2 x^2 + 1}}{a x + I}\right)^{\frac{1}{4}}}{32 x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^3,x, algorithm="fricas")

[Out] 1/32*(a^2*x^2*log((I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) + 1) + I*a^2*x^2*log((I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) + I) - I*a^2*x^2*log((I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) - I) - a^2*x^2*log((I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) - 1) + sqrt(I*a^4)*x^2*log((a^2*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) + sqrt(I*a^4))/a^2) - sqrt(I*a^4)*x^2*log((a^2*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) - sqrt(I*a^4))/a^2) + sqrt(-I*a^4)*x^2*log((a^2*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) + sqrt(-I*a^4))/a^2) - sqrt(-I*a^4)*x^2*log((a^2*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4) - sqrt(-I*a^4))/a^2) - 4*(5*a^2*x^2 + I*a*x + 4)*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4))/x^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt[4]{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4)/x**3,x)
```

```
[Out] Integral((I*(a*x - I)/sqrt(a**2*x**2 + 1))**(1/4)/x**3, x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)/x^3,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{1/4}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4)/x^3,x)
```

```
[Out] int(((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4)/x^3, x)
```

3.134 $\int e^{6i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=114

$$-\frac{x^{1+m}(1+iax)^2}{(1+m)(1-iax)^2} + \frac{4ix^{1+m}(i(1+m)^2 + a(3+3m+m^2)x)}{(1+m)(1-iax)^2} + \frac{2(3+4m+2m^2)x^{1+m} {}_2F_1(1, 1+m; 2+m, 1+m)}{1+m}$$

[Out] $-x^{1+m}*(1+I*a*x)^2/(1+m)/(1-I*a*x)^2+4*I*x^{1+m}*(I*(1+m)^2+a*(m^2+3*m+3)*x)/(1+m)/(1-I*a*x)^2+2*(2*m^2+4*m+3)*x^{1+m}*hypergeom([1, 1+m], [2+m], I*a*x)/(1+m)$

Rubi [A]

time = 0.07, antiderivative size = 114, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5170, 102, 150, 66}

$$\frac{2(2m^2 + 4m + 3)x^{m+1} {}_2F_1(1, m+1; m+2; iax)}{m+1} + \frac{4ix^{m+1}(a(m^2 + 3m + 3)x + i(m+1)^2)}{(m+1)(1-iax)^2} - \frac{(1+iax)^2 x^{m+1}}{(m+1)(1-iax)^2}$$

Antiderivative was successfully verified.

[In] Int[E^((6*I)*ArcTan[a*x])*x^m,x]

[Out] $-((x^{1+m}*(1+I*a*x)^2)/((1+m)*(1-I*a*x)^2)) + ((4*I)*x^{1+m}*(I*(1+m)^2 + a*(3+3*m+m^2)*x))/((1+m)*(1-I*a*x)^2) + (2*(3+4*m+2*m^2)*x^{1+m}*Hypergeometric2F1[1, 1+m, 2+m, I*a*x])/((1+m))$

Rule 66

Int[((b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m+1)/(b*(m+1)))*Hypergeometric2F1[-n, m+1, m+2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))

Rule 102

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a+b*x)^(m-1)*(c+d*x)^(n+1)*((e+f*x)^(p+1)/(d*f*(m+n+p+1))), x] + Dist[1/(d*f*(m+n+p+1)), Int[(a+b*x)^(m-2)*(c+d*x)^n*(e+f*x)^p*Simp[a^2*d*f*(m+n+p+1) - b*(b*c*e*(m-1) + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(2*m+n+p) - b*(d*e*(m+n) + c*f*(m+p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m+n+p+1, 0] && IntegerQ[m]

Rule 150

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[((b^3*c*e*g*(m+2) - a^3*d*f*h*(

```

n + 2) - a^2*b*(c*f*h*m - d*(f*g + e*h)*(m + n + 3)) - a*b^2*(c*(f*g + e*h)
+ d*e*g*(2*m + n + 4)) + b*(a^2*d*f*h*(m - n) - a*b*(2*c*f*h*(m + 1) - d*(
f*g + e*h)*(n + 1)) + b^2*(c*(f*g + e*h)*(m + 1) - d*e*g*(m + n + 2)))*x)/(
b^2*(b*c - a*d)^2*(m + 1)*(m + 2))*(a + b*x)^(m + 1)*(c + d*x)^(n + 1), x]
+ Dist[f*(h/b^2) - (d*(m + n + 3)*(a^2*d*f*h*(m - n) - a*b*(2*c*f*h*(m + 1)
) - d*(f*g + e*h)*(n + 1)) + b^2*(c*(f*g + e*h)*(m + 1) - d*e*g*(m + n + 2)
)))/(b^2*(b*c - a*d)^2*(m + 1)*(m + 2)), Int[(a + b*x)^(m + 2)*(c + d*x)^n,
x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && (LtQ[m, -2] || (EqQ[
m + n + 3, 0] && !LtQ[n, -2]))

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Integ
rQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{6i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 + iax)^3}{(1 - iax)^3} dx \\
&= -\frac{x^{1+m} (1 + iax)^2}{(1 + m)(1 - iax)^2} + \frac{i \int \frac{x^m (1 + iax)(-2ia(1+m) + 2a^2(3+m)x)}{(1 - iax)^3} dx}{a(1 + m)} \\
&= -\frac{x^{1+m} (1 + iax)^2}{(1 + m)(1 - iax)^2} + \frac{4ix^{1+m}(i(1 + m)^2 + a(3 + 3m + m^2)x)}{(1 + m)(1 - iax)^2} + \frac{2(3 + 4m + 2m^2)x}{(1 + m)(1 - iax)^2} \\
&= -\frac{x^{1+m} (1 + iax)^2}{(1 + m)(1 - iax)^2} + \frac{4ix^{1+m}(i(1 + m)^2 + a(3 + 3m + m^2)x)}{(1 + m)(1 - iax)^2} + \frac{2(3 + 4m + 2m^2)x}{(1 + m)(1 - iax)^2}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 94, normalized size = 0.82

$$\frac{x^{1+m}(5 - 10iax - a^2x^2 + 4m(2 - 3iax) + m^2(4 - 4iax) + 2(3 + 4m + 2m^2)(i + ax)^2 {}_2F_1(1, 1 + m; 2 + m; iax))}{(1 + m)(i + ax)^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^((6*I)*ArcTan[a*x])*x^m,x]
```

```
[Out] (x^(1 + m)*(5 - (10*I)*a*x - a^2*x^2 + 4*m*(2 - (3*I)*a*x) + m^2*(4 - (4*I)
*a*x) + 2*(3 + 4*m + 2*m^2)*(I + a*x)^2*Hypergeometric2F1[1, 1 + m, 2 + m,
I*a*x]))/((1 + m)*(I + a*x)^2)

```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 5.

time = 0.16, size = 748, normalized size = 6.56

method	result
meijerg	$(a^2)^{-\frac{1}{2}-\frac{m}{2}} \left(\frac{x^{1+m} (a^2)^{\frac{1}{2}+\frac{m}{2}} (-a^2 m^2 x^2 + 2a^2 m x^2 + 3a^2 x^2 - m^2 + 4m + 5)}{2(1+m)(a^2 x^2 + 1)^2} + \frac{4x^{1+m} (a^2)^{\frac{1}{2}+\frac{m}{2}} \left(\frac{1}{16} m^3 - \frac{3}{16} m^2 - \frac{1}{16} m + \frac{3}{16} \right) \Phi(-a^2 x^2, 1, \frac{1}{2} + \frac{m}{2})}{1+m} \right)$
	4

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^6/(a^2*x^2+1)^3*x^m,x,method=_RETURNVERBOSE)

[Out] $\frac{1}{4} (a^2)^{-1/2-1/2*m} (1/2/(1+m) * x^{(1+m)} * (a^2)^{(1/2+1/2*m)} * (-a^2*m^2*x^2+2*a^2*m*x^2+3*a^2*x^2-m^2+4*m+5) / (a^2*x^2+1)^2 + 4/(1+m) * x^{(1+m)} * (a^2)^{(1/2+1/2*m)} * (1/16*m^3-3/16*m^2-1/16*m+3/16) * \text{LerchPhi}(-a^2*x^2, 1, 1/2+1/2*m)) + 3/2 * I / a * (a^2)^{-1/2*m} * (1/2*x^m * (a^2)^{(1/2*m)} * (a^2*m*x^2+m-2) / (a^2*x^2+1)^2 - 1/4*x^m * (a^2)^{(1/2*m)} * (-2+m) * m * \text{LerchPhi}(-a^2*x^2, 1, 1/2*m)) - 15/4 * (a^2)^{-1/2-1/2*m} * (1/2*x^{(1+m)} * (a^2)^{(3/2+1/2*m)} * (a^2*m*x^2+a^2*x^2+m-1) / (a^2*x^2+1)^2 / a^2 - 1/4*x^{(1+m)} * (a^2)^{(3/2+1/2*m)} * (1+m) * (-1+m) / a^2 * \text{LerchPhi}(-a^2*x^2, 1, 1/2+1/2*m)) - 5 * I * (a^2)^{-1/2*m} / a * (-1/2*x^m * (a^2)^{(1/2*m)} * (a^2*m*x^2+4*a^2*x^2+m+2) / (a^2*x^2+1)^2 + 1/4*x^m * (a^2)^{(1/2*m)} * m * (2+m) * \text{LerchPhi}(-a^2*x^2, 1, 1/2*m)) + 15/4 * (a^2)^{-1/2-1/2*m} * (-1/2*x^{(1+m)} * (a^2)^{(1/2*m+5/2)} * (a^2*m*x^2+5*a^2*x^2+m+3) / a^4 / (a^2*x^2+1)^2 + 1/4*x^{(1+m)} * (a^2)^{(1/2*m+5/2)} * (m^2+4*m+3) / a^4 * \text{LerchPhi}(-a^2*x^2, 1, 1/2+1/2*m)) + 3/2 * I * (a^2)^{-1/2*m} / a * (1/2*x^m * (a^2)^{(1/2*m)} * (8*a^4*x^4+a^2*m^2*x^2+8*a^2*m*x^2+16*a^2*x^2+m^2+6*m+8) / (a^2*x^2+1)^2 / m - 1/4*x^m * (a^2)^{(1/2*m)} * (m^2+6*m+8) * \text{LerchPhi}(-a^2*x^2, 1, 1/2*m)) - 1/4 * (a^2)^{-1/2-1/2*m} * (1/2*x^{(1+m)} * (a^2)^{(7/2+1/2*m)} * (8*a^4*x^4+a^2*m^2*x^2+10*a^2*m*x^2+25*a^2*x^2+m^2+8*m+15) / (a^2*x^2+1)^2 / (1+m) / a^6 - 1/4*x^{(1+m)} * (a^2)^{(7/2+1/2*m)} * (m^2+8*m+15) / a^6 * \text{LerchPhi}(-a^2*x^2, 1, 1/2+1/2*m))$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^6/(a^2*x^2+1)^3*x^m,x, algorithm="maxima")

[Out] integrate((I*a*x + 1)^6*x^m/(a^2*x^2 + 1)^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^6/(a^2*x^2+1)^3*x^m,x, algorithm="fricas")

[Out] integral(-(a³*x³ - 3*I*a²*x² - 3*a*x + I)*x^m/(a³*x³ + 3*I*a²*x² - 3*a*x - I), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \left(\frac{x^m}{a^3 x^3 + 3a^2 x^2 + 3a^2 x^2 + 1} \right) dx - \int \frac{15a^2 x^2 x^m}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} dx - \int \left(\frac{15a^2 x^2 x^m}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} \right) dx - \int \frac{a^6 x^2 x^m}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} dx - \int \left(\frac{6ia x x^m}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} \right) dx - \int \frac{20a^3 x^2 x^m}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} dx - \int \left(\frac{6ia^3 x^2 x^m}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**6/(a**2*x**2+1)**3*x**m,x)

[Out] -Integral(-x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x) - Integral(15*a**2*x**2*x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x) - Integral(-15*a**4*x**4*x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x) - Integral(a**6*x**6*x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x) - Integral(-6*I*a*x*x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x) - Integral(20*I*a**3*x**3*x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x) - Integral(-6*I*a**5*x**5*x**m/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^6/(a^2*x^2+1)^3*x^m,x, algorithm="giac")

[Out] integrate((I*a*x + 1)^6*x^m/(a^2*x^2 + 1)^3, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m (1 + a x i)^6}{(a^2 x^2 + 1)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a*x*1i + 1)^6)/(a^2*x^2 + 1)^3,x)

[Out] int((x^m*(a*x*1i + 1)^6)/(a^2*x^2 + 1)^3, x)

3.135 $\int e^{4i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=50

$$\frac{x^{1+m}}{1+m} + \frac{4x^{1+m}}{1-iax} - 4x^{1+m} {}_2F_1(1, 1+m; 2+m; iax)$$

[Out] $x^{(1+m)}/(1+m)+4*x^{(1+m)}/(1-I*a*x)-4*x^{(1+m)}*\text{hypergeom}([1, 1+m], [2+m], I*a*x)$

Rubi [A]

time = 0.03, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5170, 91, 81, 66}

$$-4x^{m+1} {}_2F_1(1, m+1; m+2; iax) + \frac{4x^{m+1}}{1-iax} + \frac{x^{m+1}}{m+1}$$

Antiderivative was successfully verified.

[In] Int[E^((4*I)*ArcTan[a*x])*x^m,x]

[Out] $x^{(1+m)}/(1+m) + (4*x^{(1+m)})/(1-I*a*x) - 4*x^{(1+m)}*\text{Hypergeometric2F1}[1, 1+m, 2+m, I*a*x]$

Rule 66

Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[c^n*((b*x)^(m+1)/(b*(m+1)))*Hypergeometric2F1[-n, m+1, m+2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0]))

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n+1)*((e + f*x)^(p+1)/(d*f*(n+p+2))), x] + Dist[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1)))/(d*f*(n+p+2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n+p+2, 0]

Rule 91

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[(b*c - a*d)^2*(c + d*x)^(n+1)*((e + f*x)^(p+1)/(d^2*(d*e - c*f)*(n+1))), x] - Dist[1/(d^2*(d*e - c*f)*(n+1)), Int[(c + d*x)^(n+1)*(e + f*x)^p*Simp[a^2*d^2*f*(n+p+2) + b^2*c*(d*e*(n+1) + c*f*(p+1)) - 2*a*b*d*(d*e*(n+1) + c*f*(p+1)) - b^2*d*(d*e - c*f)*(n+1)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n+p+3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p,

1])))

Rule 5170

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
 \int e^{4i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 + iax)^2}{(1 - iax)^2} dx \\
 &= \frac{4x^{1+m}}{1 - iax} + \frac{\int \frac{x^m (-a^2(3+4m) - ia^3x)}{1 - iax} dx}{a^2} \\
 &= \frac{x^{1+m}}{1 + m} + \frac{4x^{1+m}}{1 - iax} - (4(1 + m)) \int \frac{x^m}{1 - iax} dx \\
 &= \frac{x^{1+m}}{1 + m} + \frac{4x^{1+m}}{1 - iax} - 4x^{1+m} {}_2F_1(1, 1 + m; 2 + m; iax)
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 58, normalized size = 1.16

$$\frac{x^{1+m}(5i + 4im + ax - 4(1 + m)(i + ax) {}_2F_1(1, 1 + m; 2 + m; iax))}{(1 + m)(i + ax)}$$

Antiderivative was successfully verified.

[In] Integrate[E^((4*I)*ArcTan[a*x])*x^m,x]

[Out] (x^(1 + m)*(5*I + (4*I)*m + a*x - 4*(1 + m)*(I + a*x)*Hypergeometric2F1[1, 1 + m, 2 + m, I*a*x]))/((1 + m)*(I + a*x))

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 5.

time = 0.10, size = 417, normalized size = 8.34

method	result
meijerg	$\frac{(a^2)^{-\frac{1}{2} - \frac{m}{2}} \left(\frac{2x^{1+m}(a^2)^{\frac{1}{2} + \frac{m}{2}}}{2a^2x^2 + 2} + \frac{2x^{1+m}(a^2)^{\frac{1}{2} + \frac{m}{2}} \left(-\frac{m^2}{4} + \frac{1}{4} \right) \Phi(-a^2x^2, 1, \frac{1}{2} + \frac{m}{2})}{1+m} \right)}{2} + \frac{2i(a^2)^{-\frac{m}{2}} \left(\frac{x^m(a^2)^{\frac{m}{2}}(-2-m)}{(2+m)(a^2x^2+1)} + \frac{x^m(a^2)^{\frac{m}{2}}m}{a} \right)}{a}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^4/(a^2*x^2+1)^2*x^m,x,method=_RETURNVERBOSE)

```
[Out] 1/2*(a^2)^(-1/2-1/2*m)*(2*x^(1+m)*(a^2)^(1/2+1/2*m)/(2*a^2*x^2+2)+2/(1+m)*x
^(1+m)*(a^2)^(1/2+1/2*m)*(-1/4*m^2+1/4)*LerchPhi(-a^2*x^2,1,1/2+1/2*m))+2*I
/a*(a^2)^(-1/2*m)*(1/(2+m)*x^m*(a^2)^(1/2*m)*(-2-m)/(a^2*x^2+1)+1/2*x^m*(a^
2)^(1/2*m)*m*LerchPhi(-a^2*x^2,1,1/2*m))-3*(a^2)^(-1/2-1/2*m)*(1/(m+3)*x^(1
+m)*(a^2)^(3/2+1/2*m)*(-m-3)/a^2/(a^2*x^2+1)+1/2*x^(1+m)*(a^2)^(3/2+1/2*m)*
(1+m)/a^2*LerchPhi(-a^2*x^2,1,1/2+1/2*m))-2*I*(a^2)^(-1/2*m)/a*(x^m*(a^2)^(
1/2*m)*(2*a^2*x^2+m+2)/(a^2*x^2+1)/m-1/2*x^m*(a^2)^(1/2*m)*(2+m)*LerchPhi(-
a^2*x^2,1,1/2*m))+1/2*(a^2)^(-1/2-1/2*m)*(x^(1+m)*(a^2)^(1/2*m+5/2)*(2*a^2*
x^2+m+3)/(a^2*x^2+1)/a^4/(1+m)-1/2*x^(1+m)*(a^2)^(1/2*m+5/2)*(m+3)/a^4*Lerc
hPhi(-a^2*x^2,1,1/2+1/2*m))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^m,x, algorithm="maxima")
```

```
[Out] integrate((I*a*x + 1)^4*x^m/(a^2*x^2 + 1)^2, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^m,x, algorithm="fricas")
```

```
[Out] integral((a^2*x^2 - 2*I*a*x - 1)*x^m/(a^2*x^2 + 2*I*a*x - 1), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m(ax - i)^4}{(a^2x^2 + 1)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2*x**m,x)
```

```
[Out] Integral(x**m*(a*x - I)**4/(a**2*x**2 + 1)**2, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^m,x, algorithm="giac")

[Out] integrate((I*a*x + 1)^4*x^m/(a^2*x^2 + 1)^2, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^m (1 + a x i)^4}{(a^2 x^2 + 1)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a*x*1i + 1)^4)/(a^2*x^2 + 1)^2,x)

[Out] int((x^m*(a*x*1i + 1)^4)/(a^2*x^2 + 1)^2, x)

3.136 $\int e^{2i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=39

$$-\frac{x^{1+m}}{1+m} + \frac{2x^{1+m} {}_2F_1(1, 1+m; 2+m; iax)}{1+m}$$

[Out] $-x^{(1+m)}/(1+m)+2*x^{(1+m)}*\text{hypergeom}([1, 1+m], [2+m], I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5170, 81, 66}

$$-\frac{x^{m+1}}{m+1} + \frac{2x^{m+1} {}_2F_1(1, m+1; m+2; iax)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}*x^m, x]$

[Out] $-(x^{(1+m)}/(1+m)) + (2*x^{(1+m)}*\text{Hypergeometric2F1}[1, 1+m, 2+m, I*a*x])/ (1+m)$

Rule 66

$\text{Int}[(b_.)*(x_)^m*((c_) + (d_.)*(x_)^n), x_Symbol] :> \text{Simp}[c^n*((b*x)^{m+1}/(b*(m+1)))*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(x/c)], x] /; \text{FreeQ}\{b, c, d, m, n\}, x \ \&\& \ \text{IntegerQ}[m] \ \&\& \ (\text{IntegerQ}[n] \ || \ (\text{GtQ}[c, 0] \ \&\& \ !(\text{EqQ}[n, -2^{(-1)}] \ \&\& \ \text{EqQ}[c^2 - d^2, 0] \ \&\& \ \text{GtQ}[-d/(b*c), 0])))$

Rule 81

$\text{Int}[(a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^n)*((e_.) + (f_.)*(x_)^p), x_Symbol] :> \text{Simp}[b*(c + d*x)^{n+1}*((e + f*x)^{p+1}/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x \ \&\& \ \text{NeQ}[n+p+2, 0]$

Rule 5170

$\text{Int}[E^{\text{ArcTan}[a_.)*(x_)]*(n_.)*(x_)^m, x_Symbol] :> \text{Int}[x^m*((1 - I*a*x)^{I*(n/2)})/(1 + I*a*x)^{I*(n/2)}], x] /; \text{FreeQ}\{a, m, n\}, x \ \&\& \ \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
\int e^{2i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m(1+iax)}{1-iax} dx \\
&= -\frac{x^{1+m}}{1+m} + 2 \int \frac{x^m}{1-iax} dx \\
&= -\frac{x^{1+m}}{1+m} + \frac{2x^{1+m} {}_2F_1(1, 1+m; 2+m; iax)}{1+m}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 29, normalized size = 0.74

$$\frac{x^{1+m}(-1 + 2 {}_2F_1(1, 1+m; 2+m; iax))}{1+m}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a*x])*x^m,x]``[Out] (x^(1+m)*(-1 + 2*Hypergeometric2F1[1, 1+m, 2+m, I*a*x]))/(1+m)`**Maple [C]** Result contains higher order function than in optimal. Order 9 vs. order 5.

time = 0.10, size = 175, normalized size = 4.49

method	result
meijerg	$\frac{x^{1+m} \left(\frac{1}{2} + \frac{m}{2}\right) \Phi(-a^2 x^2, 1, \frac{1}{2} + \frac{m}{2})}{1+m} + \frac{i(a^2)^{-\frac{m}{2}} \left(\frac{2x^m (a^2)^{\frac{m}{2}}}{m} + \frac{x^m (a^2)^{\frac{m}{2}} (-2-m) \Phi(-a^2 x^2, 1, \frac{m}{2})}{2+m} \right)}{a} - \frac{(a^2)^{-\frac{1}{2} - \frac{m}{2}} \left(\frac{2x^{1+m} (a^2)^{\frac{3}{2} + \frac{m}{2}}}{(1+m)a^2} \right)}{a}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^2/(a^2*x^2+1)*x^m,x,method=_RETURNVERBOSE)`

```
[Out] 1/(1+m)*x^(1+m)*(1/2+1/2*m)*LerchPhi(-a^2*x^2,1,1/2+1/2*m)+I/a*(a^2)^(-1/2*m)*(2*x^m*(a^2)^(1/2*m)/m+1/(2+m)*x^m*(a^2)^(1/2*m)*(-2-m)*LerchPhi(-a^2*x^2,1,1/2*m))-1/2*(a^2)^(-1/2-1/2*m)*(2*x^(1+m)*(a^2)^(3/2+1/2*m)/(1+m)/a^2+1/(m+3)*x^(1+m)*(a^2)^(3/2+1/2*m)*(-m-3)/a^2*LerchPhi(-a^2*x^2,1,1/2+1/2*m))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^m,x, algorithm="maxima")`

[Out] integrate((I*a*x + 1)^2*x^m/(a^2*x^2 + 1), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^m,x, algorithm="fricas")

[Out] integral(-(a*x - I)*x^m/(a*x + I), x)

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 129 vs. $2(27) = 54$.

time = 2.05, size = 129, normalized size = 3.31

$$\frac{i a m x^2 x^m \Phi\left(a x e^{\frac{i \pi}{2}}, 1, m+2\right) \Gamma(m+2)}{\Gamma(m+3)} + \frac{2 i a x^2 x^m \Phi\left(a x e^{\frac{i \pi}{2}}, 1, m+2\right) \Gamma(m+2)}{\Gamma(m+3)} + \frac{m x x^m \Phi\left(a x e^{\frac{i \pi}{2}}, 1, m+1\right) \Gamma(m+1)}{\Gamma(m+2)} + \frac{x x^m \Phi\left(a x e^{\frac{i \pi}{2}}, 1, m+1\right) \Gamma(m+1)}{\Gamma(m+2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)*x**m,x)

[Out] I*a*m*x**2*x**m*lerchphi(a*x*exp_polar(I*pi/2), 1, m + 2)*gamma(m + 2)/gamma(m + 3) + 2*I*a*x**2*x**m*lerchphi(a*x*exp_polar(I*pi/2), 1, m + 2)*gamma(m + 2)/gamma(m + 3) + m*x*x**m*lerchphi(a*x*exp_polar(I*pi/2), 1, m + 1)*gamma(m + 1)/gamma(m + 2) + x*x**m*lerchphi(a*x*exp_polar(I*pi/2), 1, m + 1)*gamma(m + 1)/gamma(m + 2)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^m,x, algorithm="giac")

[Out] integrate((I*a*x + 1)^2*x^m/(a^2*x^2 + 1), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{x^m (1 + a x i)^2}{a^2 x^2 + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a*x*1i + 1)^2)/(a^2*x^2 + 1),x)

[Out] int((x^m*(a*x*1i + 1)^2)/(a^2*x^2 + 1), x)

3.137 $\int e^{-2i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=39

$$-\frac{x^{1+m}}{1+m} + \frac{2x^{1+m} {}_2F_1(1, 1+m; 2+m; -iax)}{1+m}$$

[Out] $-x^{(1+m)}/(1+m)+2*x^{(1+m)}*\text{hypergeom}([1, 1+m], [2+m], -I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 39, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5170, 81, 66}

$$-\frac{x^{m+1}}{m+1} + \frac{2x^{m+1} {}_2F_1(1, m+1; m+2; -iax)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^m/E^{((2*I)*\text{ArcTan}[a*x])}, x]$

[Out] $-(x^{(1+m)}/(1+m)) + (2*x^{(1+m)}*\text{Hypergeometric2F1}[1, 1+m, 2+m, (-I)*a*x])/(1+m)$

Rule 66

$\text{Int}[(b_*)(x_)^{(m_*)}((c_*) + (d_*)(x_))^{(n_*)}, x_Symbol] \rightarrow \text{Simp}[c^{n*}((b*x)^{(m+1})/(b*(m+1)))*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(x/c)], x] /;$ FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))

Rule 81

$\text{Int}[(a_*) + (b_*)(x_)*((c_*) + (d_*)(x_))^{(n_*)}((e_*) + (f_*)(x_))^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[b*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1})/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1)))/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n+p+2, 0]

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_*)(x_)]*(n_*)}*(x_)^{(m_*)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned}
\int e^{-2i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m(1-iax)}{1+iax} dx \\
&= -\frac{x^{1+m}}{1+m} + 2 \int \frac{x^m}{1+iax} dx \\
&= -\frac{x^{1+m}}{1+m} + \frac{2x^{1+m} {}_2F_1(1, 1+m; 2+m; -iax)}{1+m}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 29, normalized size = 0.74

$$\frac{x^{1+m}(-1 + 2 {}_2F_1(1, 1+m; 2+m; -iax))}{1+m}$$

Antiderivative was successfully verified.

[In] Integrate[x^m/E^((2*I)*ArcTan[a*x]),x]**[Out]** (x^(1+m)*(-1 + 2*Hypergeometric2F1[1, 1+m, 2+m, (-I)*a*x]))/(1+m)**Maple [C]** Result contains higher order function than in optimal. Order 9 vs. order 5.

time = 0.08, size = 158, normalized size = 4.05

method	result
meijerg	$\frac{i(ia)^{-m} \left(\frac{x^m (ia)^m (-a^2 m x^2 - iamx - 2iax - m^2 - 3m - 2)}{(1+m)m(iax+1)} + x^m (ia)^m (2+m) \Phi(-iax, 1, m) \right)}{a} - \frac{i(ia)^{-m} \left(\frac{x^m (ia)^m (-1-m)}{(1+m)(iax+1)} + x^m (ia)^m m \Phi \right)}{a}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m/(1+I*a*x)^2*(a^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] I*(I*a)^(-m)/a*(x^m*(I*a)^m*(-a^2*m*x^2-I*a*m*x-m^2-2*I*a*x-3*m-2)/(1+m)/m/
(1+I*a*x)+x^m*(I*a)^m*(2+m)*LerchPhi(-I*a*x,1,m))-I*(I*a)^(-m)/a*(1/(1+m)*x
^m*(I*a)^m*(-1-m)/(1+I*a*x)+x^m*(I*a)^m*m*LerchPhi(-I*a*x,1,m))

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/(1+I*a*x)^2*(a^2*x^2+1),x, algorithm="maxima")**[Out]** integrate((a^2*x^2 + 1)*x^m/(I*a*x + 1)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($x^m/(1+I*a*x)^2*(a^2*x^2+1)$,x, algorithm="fricas")**[Out]** integral($-(a*x + I)*x^m/(a*x - I)$, x)**Sympy [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 136 vs. $2(29) = 58$.

time = 2.63, size = 136, normalized size = 3.49

$$-\frac{ia mx^2 x^m \Phi\left(ax e^{\frac{3i\pi}{2}}, 1, m+2\right) \Gamma(m+2)}{\Gamma(m+3)} - \frac{2ia x^2 x^m \Phi\left(ax e^{\frac{3i\pi}{2}}, 1, m+2\right) \Gamma(m+2)}{\Gamma(m+3)} + \frac{m x x^m \Phi\left(ax e^{\frac{3i\pi}{2}}, 1, m+1\right) \Gamma(m+1)}{\Gamma(m+2)} + \frac{x x^m \Phi\left(ax e^{\frac{3i\pi}{2}}, 1, m+1\right) \Gamma(m+1)}{\Gamma(m+2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($x**m/(1+I*a*x)**2*(a**2*x**2+1)$,x)

[Out] $-I*a*m*x**2*x**m*lerchphi(a*x*exp_polar(3*I*pi/2), 1, m + 2)*gamma(m + 2)/gamma(m + 3) - 2*I*a*x**2*x**m*lerchphi(a*x*exp_polar(3*I*pi/2), 1, m + 2)*gamma(m + 2)/gamma(m + 3) + m*x*x**m*lerchphi(a*x*exp_polar(3*I*pi/2), 1, m + 1)*gamma(m + 1)/gamma(m + 2) + x*x**m*lerchphi(a*x*exp_polar(3*I*pi/2), 1, m + 1)*gamma(m + 1)/gamma(m + 2)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($x^m/(1+I*a*x)^2*(a^2*x^2+1)$,x, algorithm="giac")**[Out]** integrate($(a^2*x^2 + 1)*x^m/(I*a*x + 1)^2$, x)**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{x^m (a^2 x^2 + 1)}{(1 + a x i)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int($(x^m*(a^2*x^2 + 1))/(a*x*1i + 1)^2$,x)**[Out]** int($(x^m*(a^2*x^2 + 1))/(a*x*1i + 1)^2$, x)

3.138 $\int e^{-4i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=50

$$\frac{x^{1+m}}{1+m} + \frac{4x^{1+m}}{1+iax} - 4x^{1+m} {}_2F_1(1, 1+m; 2+m; -iax)$$

[Out] $x^{(1+m)/(1+m)+4*x^{(1+m)/(1+I*a*x)}-4*x^{(1+m)*\text{hypergeom}([1, 1+m], [2+m], -I*a*x)}$

Rubi [A]

time = 0.03, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5170, 91, 81, 66}

$$-4x^{m+1} {}_2F_1(1, m+1; m+2; -iax) + \frac{4x^{m+1}}{1+iax} + \frac{x^{m+1}}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^m/E^{((4*I)*\text{ArcTan}[a*x])}, x]$

[Out] $x^{(1+m)/(1+m)} + (4*x^{(1+m)})/(1+I*a*x) - 4*x^{(1+m)*\text{Hypergeometric2F1}[1, 1+m, 2+m, (-I)*a*x]}$

Rule 66

$\text{Int}(((b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.)}, x_Symbol] :> \text{Simp}[c^{n*}((b*x)^{(m+1})/(b*(m+1)))*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(x/c)], x] /; \text{FreeQ}\{b, c, d, m, n\}, x\} \&\& \text{IntegerQ}[m] \&\& (\text{IntegerQ}[n] \mid\mid (\text{GtQ}[c, 0] \&\& \text{!(EqQ}[n, -2^{(-1)}] \&\& \text{EqQ}[c^2 - d^2, 0] \&\& \text{GtQ}[-d/(b*c), 0])))$

Rule 81

$\text{Int}(((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^{(n_.)*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] :> \text{Simp}[b*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1})/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1)))/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x\} \&\& \text{NeQ}[n+p+2, 0]$

Rule 91

$\text{Int}(((a_.) + (b_.)*(x_))^{2*}((c_.) + (d_.)*(x_))^{(n_.)*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] :> \text{Simp}[(b*c - a*d)^2*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1})/(d^2*(d*e - c*f)*(n+1))), x] - \text{Dist}[1/(d^2*(d*e - c*f)*(n+1)), \text{Int}[(c + d*x)^{(n+1)}*(e + f*x)^p*\text{Simp}[a^2*d^2*f*(n+p+2) + b^2*c*(d*e*(n+1) + c*f*(p+1)) - 2*a*b*d*(d*e*(n+1) + c*f*(p+1)) - b^2*d*(d*e - c*f)*(n+1)*x, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n, p\}, x\} \&\& (\text{LtQ}[n, -1] \mid\mid$

```
(EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1]))
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned}
 \int e^{-4i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 - iax)^2}{(1 + iax)^2} dx \\
 &= \frac{4x^{1+m}}{1 + iax} + \frac{\int \frac{x^m (-a^2(3+4m) + ia^3x)}{1 + iax} dx}{a^2} \\
 &= \frac{x^{1+m}}{1 + m} + \frac{4x^{1+m}}{1 + iax} - (4(1 + m)) \int \frac{x^m}{1 + iax} dx \\
 &= \frac{x^{1+m}}{1 + m} + \frac{4x^{1+m}}{1 + iax} - 4x^{1+m} {}_2F_1(1, 1 + m; 2 + m; -iax)
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 58, normalized size = 1.16

$$\frac{x^{1+m}(-5i - 4im + ax - 4(1 + m)(-i + ax)) {}_2F_1(1, 1 + m; 2 + m; -iax)}{(1 + m)(-i + ax)}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^m/E^((4*I)*ArcTan[a*x]), x]
```

```
[Out] (x^(1 + m)*(-5*I - (4*I)*m + a*x - 4*(1 + m)*(-I + a*x)*Hypergeometric2F1[1, 1 + m, 2 + m, (-I)*a*x]))/((1 + m)*(-I + a*x))
```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 5.

time = 0.10, size = 428, normalized size = 8.56

method	result
meijerg	$ \frac{i(ia)^{-m} \left(\frac{6a^4 x^4 m + 6ia^3 x^3 m + a^2 x^2 m^4 + 24ia^3 x^3 + 11a^2 x^2 m^3 - 2iax m^4 + 46a^2 m^2 x^2 - 21iax m^3 + 90a^2 m x^2 - 79iax m^2 + 72a^2 x^2 - 126ia}{(1+m)m(iax+1)^3} \right)}{6a} $

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m/(1+I*a*x)^4*(a^2*x^2+1)^2,x,method=_RETURNVERBOSE)
```

```
[Out] -1/6*I*(I*a)^(-m)/a*(x^m*(I*a)^m*(a^2*x^2*m^4+6*a^4*x^4*m+11*a^2*x^2*m^3-72
*I*a*x-21*I*a*x*m^3+46*a^2*m^2*x^2+24*I*a^3*x^3-m^4-79*I*a*x*m^2+90*a^2*m*x
^2+6*I*a^3*x^3*m-10*m^3+72*a^2*x^2-2*I*a*x*m^4-35*m^2-126*I*a*m*x-50*m-24)/
(1+m)/m/(1+I*a*x)^3+x^m*(I*a)^m*(m^3+9*m^2+26*m+24)*LerchPhi(-I*a*x,1,m))+1
/3*I*(I*a)^(-m)/a*(-x^m*(I*a)^m*(-a^2*m^2*x^2-4*a^2*m*x^2+2*I*a*x*m^2-6*a^2
*x^2+7*I*a*m*x+m^2+6*I*a*x+3*m+2)/(1+I*a*x)^3+x^m*(I*a)^m*m*(m^2+3*m+2)*Ler
chPhi(-I*a*x,1,m))-1/6*I*(I*a)^(-m)/a*(-x^m*(I*a)^m*(-a^2*m^2*x^2+2*a^2*m*x
^2+2*I*a*x*m^2-5*I*a*m*x+m^2-3*m+2)/(1+I*a*x)^3+x^m*(I*a)^m*(m^2-3*m+2)*m*L
erchPhi(-I*a*x,1,m))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^m/(1+I*a*x)^4*(a^2*x^2+1)^2,x, algorithm="maxima")
```

```
[Out] integrate((a^2*x^2 + 1)^2*x^m/(I*a*x + 1)^4, x)
```

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^m/(1+I*a*x)^4*(a^2*x^2+1)^2,x, algorithm="fricas")
```

```
[Out] integral((a^2*x^2 + 2*I*a*x - 1)*x^m/(a^2*x^2 - 2*I*a*x - 1), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m (a^2 x^2 + 1)^2}{(ax - i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**m/(1+I*a*x)**4*(a**2*x**2+1)**2,x)
```

```
[Out] Integral(x**m*(a**2*x**2 + 1)**2/(a*x - I)**4, x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/(1+I*a*x)^4*(a^2*x^2+1)^2,x, algorithm="giac")

[Out] integrate((a^2*x^2 + 1)^2*x^m/(I*a*x + 1)^4, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^m (a^2 x^2 + 1)^2}{(1 + a x i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a^2*x^2 + 1)^2)/(a*x*1i + 1)^4,x)

[Out] int((x^m*(a^2*x^2 + 1)^2)/(a*x*1i + 1)^4, x)

3.139 $\int e^{-6i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=115

$$-\frac{x^{1+m}(1-iax)^2}{(1+m)(1+iax)^2} + \frac{4ix^{1+m}(i(1+m)^2 - a(3+3m+m^2)x)}{(1+m)(1+iax)^2} + \frac{2(3+4m+2m^2)x^{1+m} {}_2F_1(1, 1+m; 2+m, -iax)}{1+m}$$

[Out] $-x^{1+m}*(1-I*a*x)^2/(1+m)/(1+I*a*x)^2+4*I*x^{1+m}*(I*(1+m)^2-a*(m^2+3*m+3)*x)/(1+m)/(1+I*a*x)^2+2*(2*m^2+4*m+3)*x^{1+m}*hypergeom([1, 1+m], [2+m], -I*a*x)/(1+m)$

Rubi [A]

time = 0.06, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5170, 102, 150, 66}

$$\frac{2(2m^2+4m+3)x^{m+1} {}_2F_1(1, m+1; m+2; -iax)}{m+1} + \frac{4ix^{m+1}(-a(m^2+3m+3)x+i(m+1)^2)}{(m+1)(1+iax)^2} - \frac{(1-iax)^2x^{m+1}}{(m+1)(1+iax)^2}$$

Antiderivative was successfully verified.

[In] Int[x^m/E^((6*I)*ArcTan[a*x]), x]

[Out] $-((x^{1+m}*(1-I*a*x)^2)/((1+m)*(1+I*a*x)^2)) + ((4*I)*x^{1+m}*(I*(1+m)^2 - a*(3+3*m+m^2)*x))/((1+m)*(1+I*a*x)^2) + (2*(3+4*m+2*m^2)*x^{1+m}*Hypergeometric2F1[1, 1+m, 2+m, (-I)*a*x])/((1+m))$

Rule 66

Int[((b_.)*(x_))^(m_)*((c_.)+(d_.)*(x_))^(n_), x_Symbol] := Simp[c^n*((b*x)^(m+1)/(b*(m+1)))*Hypergeometric2F1[-n, m+1, m+2, (-d)*(x/c)], x] /; FreeQ[{b, c, d, m, n}, x] && !IntegerQ[m] && (IntegerQ[n] || (GtQ[c, 0] && !(EqQ[n, -2^(-1)] && EqQ[c^2 - d^2, 0] && GtQ[-d/(b*c), 0])))

Rule 102

Int[((a_.)+(b_.)*(x_))^(m_)*((c_.)+(d_.)*(x_))^(n_)*((e_.)+(f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a+b*x)^(m-1)*(c+d*x)^(n+1)*((e+f*x)^(p+1)/(d*f*(m+n+p+1))), x] + Dist[1/(d*f*(m+n+p+1)), Int[(a+b*x)^(m-2)*(c+d*x)^n*(e+f*x)^p*Simp[a^2*d*f*(m+n+p+1) - b*(b*c*e*(m-1) + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(2*m+n+p) - b*(d*e*(m+n) + c*f*(m+p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m+n+p+1, 0] && IntegerQ[m]

Rule 150

Int[((a_.)+(b_.)*(x_))^(m_)*((c_.)+(d_.)*(x_))^(n_)*((e_.)+(f_.)*(x_))*((g_.)+(h_.)*(x_)), x_Symbol] := Simp[((b^3*c*e*g*(m+2) - a^3*d*f*h*(

```

n + 2) - a^2*b*(c*f*h*m - d*(f*g + e*h)*(m + n + 3)) - a*b^2*(c*(f*g + e*h)
+ d*e*g*(2*m + n + 4)) + b*(a^2*d*f*h*(m - n) - a*b*(2*c*f*h*(m + 1) - d*(
f*g + e*h)*(n + 1)) + b^2*(c*(f*g + e*h)*(m + 1) - d*e*g*(m + n + 2)))*x)/(
b^2*(b*c - a*d)^2*(m + 1)*(m + 2))*(a + b*x)^(m + 1)*(c + d*x)^(n + 1), x]
+ Dist[f*(h/b^2) - (d*(m + n + 3)*(a^2*d*f*h*(m - n) - a*b*(2*c*f*h*(m + 1)
) - d*(f*g + e*h)*(n + 1)) + b^2*(c*(f*g + e*h)*(m + 1) - d*e*g*(m + n + 2)
)))/(b^2*(b*c - a*d)^2*(m + 1)*(m + 2)), Int[(a + b*x)^(m + 2)*(c + d*x)^n,
x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && (LtQ[m, -2] || (EqQ[
m + n + 3, 0] && !LtQ[n, -2]))

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Integ
rQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{-6i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 - iax)^3}{(1 + iax)^3} dx \\
&= -\frac{x^{1+m} (1 - iax)^2}{(1 + m)(1 + iax)^2} - \frac{i \int \frac{x^m (1 - iax) (2ia(1+m) + 2a^2(3+m)x)}{(1 + iax)^3} dx}{a(1 + m)} \\
&= -\frac{x^{1+m} (1 - iax)^2}{(1 + m)(1 + iax)^2} + \frac{4ix^{1+m} (i(1 + m)^2 - a(3 + 3m + m^2)x)}{(1 + m)(1 + iax)^2} + (2(3 + 4m + 2m^2)) \\
&= -\frac{x^{1+m} (1 - iax)^2}{(1 + m)(1 + iax)^2} + \frac{4ix^{1+m} (i(1 + m)^2 - a(3 + 3m + m^2)x)}{(1 + m)(1 + iax)^2} + \frac{2(3 + 4m + 2m^2)}{(1 + m)(1 + iax)^2}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 94, normalized size = 0.82

$$\frac{x^{1+m} (5 + 10iax - a^2 x^2 + 4m(2 + 3iax) + m^2(4 + 4iax) + 2(3 + 4m + 2m^2) (-i + ax)^2 {}_2F_1(1, 1 + m; 2 + m; -iax))}{(1 + m)(-i + ax)^2}$$

Antiderivative was successfully verified.

```
[In] Integrate[x^m/E^((6*I)*ArcTan[a*x]), x]
```

```
[Out] (x^(1 + m)*(5 + (10*I)*a*x - a^2*x^2 + 4*m*(2 + (3*I)*a*x) + m^2*(4 + (4*I)
*a*x) + 2*(3 + 4*m + 2*m^2)*(-I + a*x)^2*Hypergeometric2F1[1, 1 + m, 2 + m,
(-I)*a*x]))/((1 + m)*(-I + a*x)^2)

```

Maple [C] Result contains higher order function than in optimal. Order 9 vs. order 5.

time = 0.17, size = 1196, normalized size = 10.40

method	result
meijerg	$i^{(ia)^{-m}} \left(\frac{x^m (ia)^m (-a^4 x^4 m^6 - 120 a^6 x^6 m - 22 a^4 x^4 m^5 + 4 i a^3 x^3 m^6 - 120 i a^5 x^5 m - 197 a^4 x^4 m^4 + 87 i a^3 x^3 m^5 - 720 i a^5 x^5 - 932 a^4 x^4 m^3 + 764 i a^3 x^3 m^2 - 197 a^4 x^4 m^6 - 120 a^6 x^6 m + 129 a^2 x^2 m^5 - 720 i a^5 x^5 + 7200 i a^3 x^3 - 3600 i a x + 11722 a^2 m^2 x^2 + 1112 a^2 x^2 m^4 + 4911 a^2 x^2 m^3 - 3600 a^4 x^4 - 175 m^4 + 4 i a^3 x^3 m^6 + 87 i a^3 x^3 m^5 - 120 i a^5 x^5 m + 3483 i a^3 x^3 m^3 - 85 i a x m^5 + 8802 i a^3 x^3 m^2 - 4200 a^4 x^4 m - 7076 i a x m^2 - 8100 i a m x + 764 i a^3 x^3 m^4 - 4 i a x m^6 - a^4 x^4 m^6 + 7200 a^2 x^2 - 720 i a x m^4 + 12000 i a^3 x^3 m - 3095 i a x m^3) / (1+m) / m / (1+I a x)^5 + x^m (I a)^m (m^5 + 20 m^4 + 155 m^3 + 580 m^2 + 1044 m + 720) * \text{LerchPhi}(-I a x, 1, m) - 1/40 * I * (I a)^{-m} / a * (-x^m (I a)^m (24 + 35 m^2 + 50 m - 392 a^2 m x^2 + a^4 x^4 m^4 + 11 a^4 x^4 m^3 + 46 a^4 x^4 m^2 + 10 m^3 - 240 i a^3 x^3 + 120 i a x - 239 a^2 m^2 x^2 - 6 a^2 x^2 m^4 - 63 a^2 x^2 m^3 + 120 a^4 x^4 - 312 i a^3 x^3 m + 41 i a x m^3 + 149 i a x m^2 + 226 i a m x + m^4 - 43 i a^3 x^3 m^3 - 171 i a^3 x^3 m^2 + 4 i a x m^4 - 4 i a^3 x^3 m^4 + 96 a^4 x^4 m - 240 a^2 x^2) / (1+I a x)^5 + x^m (I a)^m m (m^4 + 10 m^3 + 35 m^2 + 50 m + 24) * \text{LerchPhi}(-I a x, 1, m) + 1/40 * I * (I a)^{-m} / a * (-x^m (I a)^m (a^4 x^4 m^4 + a^4 x^4 m^3 - 21 i a x m^2 - 4 a^4 x^4 m^2 - 4 i a m x - 6 a^2 x^2 m^4 - 4 a^4 x^4 m + 20 i a x - 3 a^2 x^2 m^3 - 3 i a^3 x^3 m^3 + 18 i a^3 x^3 m + 31 a^2 m^2 x^2 + 4 i a x m^4 + m^4 + 18 a^2 m x^2 - 4 i a^3 x^3 m^4 - 40 a^2 x^2 + 19 i a^3 x^3 m^2 - 5 m^2 + i a x m^3 + 4) / (1+I a x)^5 + x^m (I a)^m (m^2 - 3 m + 2) * m (m^2 + 3 m + 2) * \text{LerchPhi}(-I a x, 1, m) - 1/120 * I * (I a)^{-m} / a * (-x^m (I a)^m (a^4 x^4 m^4 - 9 a^4 x^4 m^3 + 129 i a x m^2 + 26 a^4 x^4 m^2 - 154 i a m x - 6 a^2 x^2 m^4 - 24 a^4 x^4 m + 37 i a^3 x^3 m^3 + 57 a^2 x^2 m^3 + 108 i a^3 x^3 m + 4 i a x m^4 - 179 a^2 m^2 x^2 - 4 i a^3 x^3 m^4 + m^4 + 188 a^2 m x^2 - 111 i a^3 x^3 m^2 - 10 m^3 - 39 i a x m^3 + 35 m^2 - 50 m + 24) / (1+I a x)^5 + x^m (I a)^m (m^4 - 10 m^3 + 35 m^2 - 50 m + 24) * m * \text{LerchPhi}(-I a x, 1, m) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^m/(1+I*a*x)^6*(a^2*x^2+1)^3,x,method=_RETURNVERBOSE)`

[Out] $1/120 * I * (I a)^{-m} / a * (x^m (I a)^m (-720 - 1624 m^2 - 1764 m + 14400 a^2 m x^2 - 22 a^4 x^4 m^5 - 197 a^4 x^4 m^4 - 932 a^4 x^4 m^3 - 2556 a^4 x^4 m^2 - m^6 - 735 m^3 - 21 m^5 + 6 a^2 x^2 m^6 - 120 a^6 x^6 m + 129 a^2 x^2 m^5 - 720 i a^5 x^5 + 7200 i a^3 x^3 - 3600 i a x + 11722 a^2 m^2 x^2 + 1112 a^2 x^2 m^4 + 4911 a^2 x^2 m^3 - 3600 a^4 x^4 - 175 m^4 + 4 i a^3 x^3 m^6 + 87 i a^3 x^3 m^5 - 120 i a^5 x^5 m + 3483 i a^3 x^3 m^3 - 85 i a x m^5 + 8802 i a^3 x^3 m^2 - 4200 a^4 x^4 m - 7076 i a x m^2 - 8100 i a m x + 764 i a^3 x^3 m^4 - 4 i a x m^6 - a^4 x^4 m^6 + 7200 a^2 x^2 - 720 i a x m^4 + 12000 i a^3 x^3 m - 3095 i a x m^3) / (1+m) / m / (1+I a x)^5 + x^m (I a)^m (m^5 + 20 m^4 + 155 m^3 + 580 m^2 + 1044 m + 720) * \text{LerchPhi}(-I a x, 1, m) - 1/40 * I * (I a)^{-m} / a * (-x^m (I a)^m (24 + 35 m^2 + 50 m - 392 a^2 m x^2 + a^4 x^4 m^4 + 11 a^4 x^4 m^3 + 46 a^4 x^4 m^2 + 10 m^3 - 240 i a^3 x^3 + 120 i a x - 239 a^2 m^2 x^2 - 6 a^2 x^2 m^4 - 63 a^2 x^2 m^3 + 120 a^4 x^4 - 312 i a^3 x^3 m + 41 i a x m^3 + 149 i a x m^2 + 226 i a m x + m^4 - 43 i a^3 x^3 m^3 - 171 i a^3 x^3 m^2 + 4 i a x m^4 - 4 i a^3 x^3 m^4 + 96 a^4 x^4 m - 240 a^2 x^2) / (1+I a x)^5 + x^m (I a)^m m (m^4 + 10 m^3 + 35 m^2 + 50 m + 24) * \text{LerchPhi}(-I a x, 1, m) + 1/40 * I * (I a)^{-m} / a * (-x^m (I a)^m (a^4 x^4 m^4 + a^4 x^4 m^3 - 21 i a x m^2 - 4 a^4 x^4 m^2 - 4 i a m x - 6 a^2 x^2 m^4 - 4 a^4 x^4 m + 20 i a x - 3 a^2 x^2 m^3 - 3 i a^3 x^3 m^3 + 18 i a^3 x^3 m + 31 a^2 m^2 x^2 + 4 i a x m^4 + m^4 + 18 a^2 m x^2 - 4 i a^3 x^3 m^4 - 40 a^2 x^2 + 19 i a^3 x^3 m^2 - 5 m^2 + i a x m^3 + 4) / (1+I a x)^5 + x^m (I a)^m (m^2 - 3 m + 2) * m (m^2 + 3 m + 2) * \text{LerchPhi}(-I a x, 1, m) - 1/120 * I * (I a)^{-m} / a * (-x^m (I a)^m (a^4 x^4 m^4 - 9 a^4 x^4 m^3 + 129 i a x m^2 + 26 a^4 x^4 m^2 - 154 i a m x - 6 a^2 x^2 m^4 - 24 a^4 x^4 m + 37 i a^3 x^3 m^3 + 57 a^2 x^2 m^3 + 108 i a^3 x^3 m + 4 i a x m^4 - 179 a^2 m^2 x^2 - 4 i a^3 x^3 m^4 + m^4 + 188 a^2 m x^2 - 111 i a^3 x^3 m^2 - 10 m^3 - 39 i a x m^3 + 35 m^2 - 50 m + 24) / (1+I a x)^5 + x^m (I a)^m (m^4 - 10 m^3 + 35 m^2 - 50 m + 24) * m * \text{LerchPhi}(-I a x, 1, m)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^m/(1+I*a*x)^6*(a^2*x^2+1)^3,x, algorithm="maxima")`

[Out] `integrate((a^2*x^2 + 1)^3*x^m/(I*a*x + 1)^6, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/(1+I*a*x)^6*(a^2*x^2+1)^3,x, algorithm="fricas")

[Out] integral(-(a^3*x^3 + 3*I*a^2*x^2 - 3*a*x - I)*x^m/(a^3*x^3 - 3*I*a^2*x^2 - 3*a*x + I), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{x^m}{a^6 x^6 - 6ia^5 x^5 - 15a^4 x^4 + 20ia^3 x^3 + 15a^2 x^2 - 6iax - 1} dx - \int \frac{3a^2 x^2 x^m}{a^6 x^6 - 6ia^5 x^5 - 15a^4 x^4 + 20ia^3 x^3 + 15a^2 x^2 - 6iax - 1} dx - \int \frac{3a^4 x^4 x^m}{a^6 x^6 - 6ia^5 x^5 - 15a^4 x^4 + 20ia^3 x^3 + 15a^2 x^2 - 6iax - 1} dx - \int \frac{a^6 x^6 x^m}{a^6 x^6 - 6ia^5 x^5 - 15a^4 x^4 + 20ia^3 x^3 + 15a^2 x^2 - 6iax - 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**m/(1+I*a*x)**6*(a**2*x**2+1)**3,x)

[Out] -Integral(x**m/(a**6*x**6 - 6*I*a**5*x**5 - 15*a**4*x**4 + 20*I*a**3*x**3 + 15*a**2*x**2 - 6*I*a*x - 1), x) - Integral(3*a**2*x**2*x**m/(a**6*x**6 - 6*I*a**5*x**5 - 15*a**4*x**4 + 20*I*a**3*x**3 + 15*a**2*x**2 - 6*I*a*x - 1), x) - Integral(3*a**4*x**4*x**m/(a**6*x**6 - 6*I*a**5*x**5 - 15*a**4*x**4 + 20*I*a**3*x**3 + 15*a**2*x**2 - 6*I*a*x - 1), x) - Integral(a**6*x**6*x**m/(a**6*x**6 - 6*I*a**5*x**5 - 15*a**4*x**4 + 20*I*a**3*x**3 + 15*a**2*x**2 - 6*I*a*x - 1), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/(1+I*a*x)^6*(a^2*x^2+1)^3,x, algorithm="giac")

[Out] integrate((a^2*x^2 + 1)^3*x^m/(I*a*x + 1)^6, x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m (a^2 x^2 + 1)^3}{(1 + a x i)^6} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a^2*x^2 + 1)^3)/(a*x*1i + 1)^6,x)

[Out] int((x^m*(a^2*x^2 + 1)^3)/(a*x*1i + 1)^6, x)

3.140 $\int e^{3i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=159

$$-\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} - \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m} + \frac{4x^{1+m} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} + \frac{4iax^{2+m} {}_2F_1\left(\frac{3}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m}$$

[Out] $-3*x^{(1+m)}*\text{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], -a^2*x^2)/(1+m) - I*a*x^{(2+m)}*\text{hypergeom}([1/2, 1+1/2*m], [2+1/2*m], -a^2*x^2)/(2+m) + 4*x^{(1+m)}*\text{hypergeom}([3/2, 1/2+1/2*m], [3/2+1/2*m], -a^2*x^2)/(1+m) + 4*I*a*x^{(2+m)}*\text{hypergeom}([3/2, 1+1/2*m], [2+1/2*m], -a^2*x^2)/(2+m)$

Rubi [A]

time = 0.66, antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5168, 6874, 371, 864, 822}

$$-\frac{3x^{m+1} {}_2F_1\left(\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; -a^2x^2\right)}{m+1} + \frac{4x^{m+1} {}_2F_1\left(\frac{3}{2}, \frac{m+1}{2}; \frac{m+3}{2}; -a^2x^2\right)}{m+1} - \frac{iax^{m+2} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{2}; \frac{m+4}{2}; -a^2x^2\right)}{m+2} + \frac{4iax^{m+2} {}_2F_1\left(\frac{3}{2}, \frac{m+2}{2}; \frac{m+4}{2}; -a^2x^2\right)}{m+2}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a*x])*x^m,x]

[Out] $(-3*x^{(1+m)}*\text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, -(a^2*x^2)]/(1+m) - (I*a*x^{(2+m)}*\text{Hypergeometric2F1}[1/2, (2+m)/2, (4+m)/2, -(a^2*x^2)]/(2+m) + (4*x^{(1+m)}*\text{Hypergeometric2F1}[3/2, (1+m)/2, (3+m)/2, -(a^2*x^2)]/(1+m) + ((4*I)*a*x^{(2+m)}*\text{Hypergeometric2F1}[3/2, (2+m)/2, (4+m)/2, -(a^2*x^2)]/(2+m)$

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[a^p * ((c*x)^(m+1)/(c*(m+1))) * Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 822

Int[((e_.)*(x_))^(m_.)*((f_) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[f, Int[(e*x)^m*(a+c*x^2)^p, x], x] + Dist[g/e, Int[(e*x)^(m+1)*(a+c*x^2)^p, x], x] /; FreeQ[{a, c, e, f, g, p}, x] && !RationalQ[m] && !IGtQ[p, 0]

Rule 864

Int[((x_)^(n_.)*((a_) + (c_.)*(x_)^2)^(p_))/((d_) + (e_.)*(x_)), x_Symbol] := Int[x^n*(a/d + c*(x/e))*(a+c*x^2)^(p-1), x] /; FreeQ[{a, c, d, e, n}, x]

```
p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && ( !IntegerQ[n] || !IntegerQ[2*p] || IGtQ[n, 2] || (GtQ[p, 0] && NeQ[n, 2]))
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]]
```

Rubi steps

$$\begin{aligned}
 \int e^{3i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 + iax)^2}{(1 - iax) \sqrt{1 + a^2 x^2}} dx \\
 &= \int \left(-\frac{3x^m}{\sqrt{1 + a^2 x^2}} - \frac{iax^{1+m}}{\sqrt{1 + a^2 x^2}} + \frac{4x^m}{(1 - iax) \sqrt{1 + a^2 x^2}} \right) dx \\
 &= -\left(3 \int \frac{x^m}{\sqrt{1 + a^2 x^2}} dx \right) + 4 \int \frac{x^m}{(1 - iax) \sqrt{1 + a^2 x^2}} dx - (ia) \int \frac{x^{1+m}}{\sqrt{1 + a^2 x^2}} dx \\
 &= -\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m} - \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2 x^2\right)}{2+m} + 4 \int \frac{x^m (1 + iax)}{(1 + a^2 x^2)^{3/2}} dx \\
 &= -\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m} - \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2 x^2\right)}{2+m} + 4 \int \frac{x^m}{(1 + a^2 x^2)^{3/2}} dx \\
 &= -\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m} - \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2 x^2\right)}{2+m} + \frac{4x^{1+m} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.06, size = 113, normalized size = 0.71

$$\frac{ix^{1+m} \sqrt{1-iax} \sqrt{-i+ax} \left(F_1\left(1+m; -\frac{1}{2}, \frac{1}{2}; 2+m; -iax, iax\right) - 2F_1\left(1+m; -\frac{1}{2}, \frac{3}{2}; 2+m; -iax, iax\right) \right)}{(1+m) \sqrt{1+iax} \sqrt{i+ax}}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[E^((3*I)*ArcTan[a*x])*x^m,x]
```

[Out] $((-I)*x^{(1+m)}*\text{Sqrt}[1-I*a*x]*\text{Sqrt}[-I+a*x]*(\text{AppellF1}[1+m, -1/2, 1/2, 2+m, (-I)*a*x, I*a*x] - 2*\text{AppellF1}[1+m, -1/2, 3/2, 2+m, (-I)*a*x, I*a*x]))/((1+m)*\text{Sqrt}[1+I*a*x]*\text{Sqrt}[I+a*x])$

Maple [A]

time = 0.08, size = 146, normalized size = 0.92

method	result
meijerg	$\frac{x^{1+m} \text{hypergeom}\left(\left[\frac{3}{2}, \frac{1}{2} + \frac{m}{2}\right], \left[\frac{3}{2} + \frac{m}{2}\right], -a^2 x^2\right)}{1+m} + \frac{3ia x^{2+m} \text{hypergeom}\left(\left[\frac{3}{2}, 1 + \frac{m}{2}\right], \left[2 + \frac{m}{2}\right], -a^2 x^2\right)}{2+m} - \frac{3a^2 x^{m+3} \text{hypergeom}\left(\left[\frac{3}{2}, \frac{3}{2}\right], \left[\frac{3}{2} + m\right], -a^2 x^2\right)}{m+3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^m,x,method=_RETURNVERBOSE)`

[Out] $x^{(1+m)}*\text{hypergeom}\left(\left[\frac{3}{2}, \frac{1}{2} + \frac{1}{2}m\right], \left[\frac{3}{2} + \frac{1}{2}m\right], -a^2*x^2\right)/(1+m) + 3*I*a/(2+m)*x^{(2+m)}*\text{hypergeom}\left(\left[\frac{3}{2}, 1 + \frac{1}{2}m\right], \left[2 + \frac{1}{2}m\right], -a^2*x^2\right) - 3*a^2/(m+3)*x^{(m+3)}*\text{hypergeom}\left(\left[\frac{3}{2}, \frac{3}{2} + \frac{1}{2}m\right], \left[\frac{1}{2}m + \frac{5}{2}\right], -a^2*x^2\right) - I*a^3/(4+m)*x^{(4+m)}*\text{hypergeom}\left(\left[\frac{3}{2}, 2 + \frac{1}{2}m\right], \left[\frac{1}{2}m + 3\right], -a^2*x^2\right)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^m,x, algorithm="maxima")`

[Out] `integrate((I*a*x + 1)^3*x^m/(a^2*x^2 + 1)^(3/2), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^m,x, algorithm="fricas")`

[Out] `integral(sqrt(a^2*x^2 + 1)*(-I*a*x - 1)*x^m/(a^2*x^2 + 2*I*a*x - 1), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i\left(\int \frac{ix^m}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ax^m}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}}\right) dx + \int \frac{a^3x^3x^m}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx + \int \left(-\frac{3ia^2x^2x^m}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}}\right) dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)*x**m,x)`

```
[Out] -I*(Integral(I*x**m/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)),
x) + Integral(-3*a*x*x**m/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 +
1))), x) + Integral(a**3*x**3*x**m/(a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a*
*2*x**2 + 1)), x) + Integral(-3*I*a**2*x**2*x**m/(a**2*x**2*sqrt(a**2*x**2
+ 1) + sqrt(a**2*x**2 + 1)), x))
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^m,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m (1 + a x i)^3}{(a^2 x^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^m*(a*x*i + 1)^3)/(a^2*x^2 + 1)^(3/2),x)
```

```
[Out] int((x^m*(a*x*i + 1)^3)/(a^2*x^2 + 1)^(3/2), x)
```

3.141 $\int e^{i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=79

$$\frac{x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} + \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m}$$

[Out] $x^{(1+m)} \text{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], -a^2*x^2)/(1+m) + I*a*x^{(2+m)} \text{hypergeom}([1/2, 1+1/2*m], [2+1/2*m], -a^2*x^2)/(2+m)$

Rubi [A]

time = 0.03, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5168, 822, 371}

$$\frac{x^{m+1} {}_2F_1\left(\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; -a^2x^2\right)}{m+1} + \frac{iax^{m+2} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{2}; \frac{m+4}{2}; -a^2x^2\right)}{m+2}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])*x^m,x]

[Out] $(x^{(1+m)} \text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, -(a^2*x^2)])/(1+m) + (I*a*x^{(2+m)} \text{Hypergeometric2F1}[1/2, (2+m)/2, (4+m)/2, -(a^2*x^2)])/(2+m)$

Rule 371

Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p * ((c*x)^(m+1)/(c*(m+1))) * Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 822

Int[((e_.)*(x_))^(m_.)*((f_) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Dist[f, Int[(e*x)^m*(a+c*x^2)^p, x], x] + Dist[g/e, Int[(e*x)^(m+1)*(a+c*x^2)^p, x], x] /; FreeQ[{a, c, e, f, g, p}, x] && !RationalQ[m] && !IGtQ[p, 0]

Rule 5168

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a*x)^(I*n+1/2)/((1 + I*a*x)^(I*n-1/2)*Sqrt[1 + a^2*x^2])), x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n-1)/2]

Rubi steps

$$\begin{aligned}
\int e^{i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m(1+iax)}{\sqrt{1+a^2x^2}} dx \\
&= (ia) \int \frac{x^{1+m}}{\sqrt{1+a^2x^2}} dx + \int \frac{x^m}{\sqrt{1+a^2x^2}} dx \\
&= \frac{x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} + \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.03, size = 85, normalized size = 1.08

$$\frac{ix^{1+m} \sqrt{1-iax} \sqrt{-i+ax} F_1\left(1+m; -\frac{1}{2}, \frac{1}{2}; 2+m; -iax, iax\right)}{(1+m) \sqrt{1+iax} \sqrt{i+ax}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a*x])*x^m,x]

[Out] (I*x^(1+m)*Sqrt[1-I*a*x]*Sqrt[-I+a*x]*AppellF1[1+m, -1/2, 1/2, 2+m, (-I)*a*x, I*a*x])/((1+m)*Sqrt[1+I*a*x]*Sqrt[I+a*x])

Maple [A]

time = 0.06, size = 71, normalized size = 0.90

method	result	size
meijerg	$\frac{x^{1+m} \operatorname{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2} + \frac{m}{2}\right], \left[\frac{3}{2} + \frac{m}{2}\right], -a^2x^2\right)}{1+m} + \frac{ia x^{2+m} \operatorname{hypergeom}\left(\left[\frac{1}{2}, 1 + \frac{m}{2}\right], \left[2 + \frac{m}{2}\right], -a^2x^2\right)}{2+m}$	71

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^m,x,method=_RETURNVERBOSE)

[Out] x^(1+m)*hypergeom([1/2, 1/2+1/2*m], [3/2+1/2*m], -a^2*x^2)/(1+m)+I*a*x^(2+m)*hypergeom([1/2, 1+1/2*m], [2+1/2*m], -a^2*x^2)/(2+m)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^m,x, algorithm="maxima")

[Out] integrate((I*a*x + 1)*x^m/sqrt(a^2*x^2 + 1), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^m,x, algorithm="fricas")

[Out] integral(I*sqrt(a^2*x^2 + 1)*x^m/(a*x + I), x)

Sympy [A]

time = 1.79, size = 95, normalized size = 1.20

$$\frac{iax^2x^m\Gamma\left(\frac{m}{2}+1\right) {}_2F_1\left(\frac{1}{2}, \frac{m}{2}+1 \middle| \frac{m}{2}+2; a^2x^2e^{i\pi}\right)}{2\Gamma\left(\frac{m}{2}+2\right)} + \frac{xx^m\Gamma\left(\frac{m}{2}+\frac{1}{2}\right) {}_2F_1\left(\frac{1}{2}, \frac{m}{2}+\frac{1}{2} \middle| \frac{m}{2}+\frac{3}{2}; a^2x^2e^{i\pi}\right)}{2\Gamma\left(\frac{m}{2}+\frac{3}{2}\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)*x**m,x)

[Out] I*a*x**2*x**m*gamma(m/2 + 1)*hyper((1/2, m/2 + 1), (m/2 + 2,), a**2*x**2*exp_polar(I*pi))/(2*gamma(m/2 + 2)) + x*x**m*gamma(m/2 + 1/2)*hyper((1/2, m/2 + 1/2), (m/2 + 3/2,), a**2*x**2*exp_polar(I*pi))/(2*gamma(m/2 + 3/2))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^m,x, algorithm="giac")

[Out] integrate((I*a*x + 1)*x^m/sqrt(a^2*x^2 + 1), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m (1 + a x i)}{\sqrt{a^2 x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a*x*1i + 1))/(a^2*x^2 + 1)^(1/2),x)

[Out] int((x^m*(a*x*1i + 1))/(a^2*x^2 + 1)^(1/2), x)

3.142 $\int e^{-i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=79

$$\frac{x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} - \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m}$$

[Out] $x^{(1+m)} \cdot \text{hypergeom}\left(\left[\frac{1}{2}, \frac{1}{2}+1/2*m\right], \left[\frac{3}{2}+1/2*m\right], -a^2*x^2\right) / (1+m) - I*a*x^{(2+m)} \cdot \text{hypergeom}\left(\left[\frac{1}{2}, 1+1/2*m\right], \left[2+1/2*m\right], -a^2*x^2\right) / (2+m)$

Rubi [A]

time = 0.03, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5168, 822, 371}

$$\frac{x^{m+1} {}_2F_1\left(\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; -a^2x^2\right)}{m+1} - \frac{iax^{m+2} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{2}; \frac{m+4}{2}; -a^2x^2\right)}{m+2}$$

Antiderivative was successfully verified.

[In] `Int[x^m/E^(I*ArcTan[a*x]),x]`

[Out] $(x^{(1+m)} \cdot \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(1+m)}{2}, \frac{(3+m)}{2}, -(a^2*x^2)\right]) / (1+m) - (I*a*x^{(2+m)} \cdot \text{Hypergeometric2F1}\left[\frac{1}{2}, \frac{(2+m)}{2}, \frac{(4+m)}{2}, -(a^2*x^2)\right]) / (2+m)$

Rule 371

`Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] :> Simp[a^p * ((c*x)^(m+1)/(c*(m+1))) * Hypergeometric2F1[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /; FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])`

Rule 822

`Int[((e_.)*(x_))^(m_.)*((f_) + (g_.)*(x_))*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Dist[f, Int[(e*x)^m*(a+c*x^2)^p, x], x] + Dist[g/e, Int[(e*x)^(m+1)*(a+c*x^2)^p, x], x] /; FreeQ[{a, c, e, f, g, p}, x] && !RationalQ[m] && !IGtQ[p, 0]`

Rule 5168

`Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a*x)^(n*(m+1)/2)/((1 + I*a*x)^(n*(m-1)/2)*Sqrt[1 + a^2*x^2])], x] /; FreeQ[{a, m}, x] && IntegerQ[(I*n - 1)/2]`

Rubi steps

$$\begin{aligned}
\int e^{-i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m(1-iax)}{\sqrt{1+a^2x^2}} dx \\
&= -\left((ia) \int \frac{x^{1+m}}{\sqrt{1+a^2x^2}} dx \right) + \int \frac{x^m}{\sqrt{1+a^2x^2}} dx \\
&= \frac{x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} - \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.03, size = 85, normalized size = 1.08

$$-\frac{ix^{1+m} \sqrt{1+iax} \sqrt{i+ax} F_1\left(1+m; \frac{1}{2}, -\frac{1}{2}; 2+m; -iax, iax\right)}{(1+m) \sqrt{1-iax} \sqrt{-i+ax}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[x^m/E^(I*ArcTan[a*x]),x]

[Out] ((-I)*x^(1+m)*Sqrt[1+I*a*x]*Sqrt[I+a*x]*AppellF1[1+m, 1/2, -1/2, 2+m, (-I)*a*x, I*a*x])/((1+m)*Sqrt[1-I*a*x]*Sqrt[-I+a*x])

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{x^m \sqrt{a^2x^2+1}}{iax+1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m/(1+I*a*x)*(a^2*x^2+1)^(1/2),x)

[Out] int(x^m/(1+I*a*x)*(a^2*x^2+1)^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a^2*x^2+1)*x^m/(I*a*x+1),x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^m/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="fricas")``[Out] integral(-I*sqrt(a^2*x^2 + 1)*x^m/(a*x - I), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^m \sqrt{a^2 x^2 + 1}}{ax - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x**m/(1+I*a*x)*(a**2*x**2+1)**(1/2),x)``[Out] -I*Integral(x**m*sqrt(a**2*x**2 + 1)/(a*x - I), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^m/(1+I*a*x)*(a^2*x^2+1)^(1/2),x, algorithm="giac")``[Out] integrate(sqrt(a^2*x^2 + 1)*x^m/(I*a*x + 1), x)`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m \sqrt{a^2 x^2 + 1}}{1 + a x i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((x^m*(a^2*x^2 + 1)^(1/2))/(a*x*1i + 1),x)``[Out] int((x^m*(a^2*x^2 + 1)^(1/2))/(a*x*1i + 1), x)`

3.143 $\int e^{-3i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=159

$$-\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} + \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m} + \frac{4x^{1+m} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2x^2\right)}{1+m} - \frac{4iax^{2+m} {}_2F_1\left(\frac{3}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2x^2\right)}{2+m}$$

[Out] $-3*x^{(1+m)}*\text{hypergeom}([1/2, 1/2+1/2*m], [3/2+1/2*m], -a^2*x^2)/(1+m)+I*a*x^{(2+m)}*\text{hypergeom}([1/2, 1+1/2*m], [2+1/2*m], -a^2*x^2)/(2+m)+4*x^{(1+m)}*\text{hypergeom}([3/2, 1/2+1/2*m], [3/2+1/2*m], -a^2*x^2)/(1+m)-4*I*a*x^{(2+m)}*\text{hypergeom}([3/2, 1+1/2*m], [2+1/2*m], -a^2*x^2)/(2+m)$

Rubi [A]

time = 0.59, antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5168, 6874, 371, 864, 822}

$$-\frac{3x^{m+1} {}_2F_1\left(\frac{1}{2}, \frac{m+1}{2}; \frac{m+3}{2}; -a^2x^2\right)}{m+1} + \frac{4x^{m+1} {}_2F_1\left(\frac{3}{2}, \frac{m+1}{2}; \frac{m+3}{2}; -a^2x^2\right)}{m+1} + \frac{iax^{m+2} {}_2F_1\left(\frac{1}{2}, \frac{m+2}{2}; \frac{m+4}{2}; -a^2x^2\right)}{m+2} - \frac{4iax^{m+2} {}_2F_1\left(\frac{3}{2}, \frac{m+2}{2}; \frac{m+4}{2}; -a^2x^2\right)}{m+2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^m/E^{((3*I)*\text{ArcTan}[a*x])}, x]$

[Out] $(-3*x^{(1+m)}*\text{Hypergeometric2F1}[1/2, (1+m)/2, (3+m)/2, -(a^2*x^2)]/(1+m) + (I*a*x^{(2+m)}*\text{Hypergeometric2F1}[1/2, (2+m)/2, (4+m)/2, -(a^2*x^2)]/(2+m) + (4*x^{(1+m)}*\text{Hypergeometric2F1}[3/2, (1+m)/2, (3+m)/2, -(a^2*x^2)]/(1+m) - ((4*I)*a*x^{(2+m)}*\text{Hypergeometric2F1}[3/2, (2+m)/2, (4+m)/2, -(a^2*x^2)]/(2+m)$

Rule 371

$\text{Int}[(c_.*x_*)^{(m_*)}*((a_*) + (b_.*x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[a^p * ((c*x)^{(m+1})/(c*(m+1))) * \text{Hypergeometric2F1}[-p, (m+1)/n, (m+1)/n+1, (-b)*(x^n/a)], x] /;$ FreeQ[{a, b, c, m, n, p}, x] && !IGtQ[p, 0] && (ILtQ[p, 0] || GtQ[a, 0])

Rule 822

$\text{Int}[(e_.*x_*)^{(m_*)}*((f_*) + (g_.*x_*)^{(n_*)})^{(p_*)}, x_Symbol] \rightarrow \text{Dist}[f, \text{Int}[(e*x)^m*(a+c*x^2)^p, x], x] + \text{Dist}[g/e, \text{Int}[(e*x)^{(m+1)}*(a+c*x^2)^p, x], x] /;$ FreeQ[{a, c, e, f, g, p}, x] && !RationalQ[m] && !IGtQ[p, 0]

Rule 864

$\text{Int}[(x_*)^{(n_*)}*((a_*) + (c_.*x_*)^{(p_*)})^{(q_*)}/((d_*) + (e_.*x_*)^{(r_*)}), x_Symbol] \rightarrow \text{Int}[x^n*(a/d + c*(x/e))*(a+c*x^2)^{(p-1)}, x] /;$ FreeQ[{a, c, d, e, n}, x]

```
p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && ( !IntegerQ[n] || !In
tegerQ[2*p] || IGtQ[n, 2] || (GtQ[p, 0] && NeQ[n, 2]))
```

Rule 5168

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*
x)^((I*n + 1)/2)/((1 + I*a*x)^((I*n - 1)/2)*Sqrt[1 + a^2*x^2]), x] /; Free
Q[{a, m}, x] && IntegerQ[(I*n - 1)/2]
```

Rule 6874

```
Int[u_, x_Symbol] := With[{v = ExpandIntegrand[u, x]}, Int[v, x] /; SumQ[v]
]
```

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 - iax)^2}{(1 + iax) \sqrt{1 + a^2 x^2}} dx \\
&= \int \left(-\frac{3x^m}{\sqrt{1 + a^2 x^2}} + \frac{iax^{1+m}}{\sqrt{1 + a^2 x^2}} + \frac{4x^m}{(1 + iax) \sqrt{1 + a^2 x^2}} \right) dx \\
&= -\left(3 \int \frac{x^m}{\sqrt{1 + a^2 x^2}} dx \right) + 4 \int \frac{x^m}{(1 + iax) \sqrt{1 + a^2 x^2}} dx + (ia) \int \frac{x^{1+m}}{\sqrt{1 + a^2 x^2}} dx \\
&= -\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m} + \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2 x^2\right)}{2+m} + 4 \int \frac{x^m (1 - iax)^2}{(1 + a^2 x^2)^{3/2}} dx \\
&= -\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m} + \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2 x^2\right)}{2+m} + 4 \int \frac{x^m}{(1 + a^2 x^2)^{3/2}} dx \\
&= -\frac{3x^{1+m} {}_2F_1\left(\frac{1}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{1+m} + \frac{iax^{2+m} {}_2F_1\left(\frac{1}{2}, \frac{2+m}{2}; \frac{4+m}{2}; -a^2 x^2\right)}{2+m} + \frac{4x^{1+m} {}_2F_1\left(\frac{3}{2}, \frac{1+m}{2}; \frac{3+m}{2}; -a^2 x^2\right)}{2+m}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 6 vs. order 5 in optimal.

time = 0.05, size = 113, normalized size = 0.71

$$\frac{ix^{1+m} \sqrt{1+iax} \sqrt{i+ax} \left(F_1\left(1+m; \frac{1}{2}, -\frac{1}{2}; 2+m; -iax, iax\right) - 2F_1\left(1+m; \frac{3}{2}, -\frac{1}{2}; 2+m; -iax, iax\right) \right)}{(1+m) \sqrt{1-iax} \sqrt{-i+ax}}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[x^m/E^((3*I)*ArcTan[a*x]), x]
```

[Out] $(I*x^{(1+m)}*Sqrt[1+I*a*x]*Sqrt[I+a*x]*(AppellF1[1+m, 1/2, -1/2, 2+m, (-I)*a*x, I*a*x] - 2*AppellF1[1+m, 3/2, -1/2, 2+m, (-I)*a*x, I*a*x]) / ((1+m)*Sqrt[1-I*a*x]*Sqrt[-I+a*x])$

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{x^m (a^2 x^2 + 1)^{\frac{3}{2}}}{(iax + 1)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^m/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x)`

[Out] `int(x^m/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^m/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out] `integrate((a^2*x^2 + 1)^(3/2)*x^m/(I*a*x + 1)^3, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^m/(1+I*a*x)^3*(a^2*x^2+1)^(3/2),x, algorithm="fricas")`

[Out] `integral(sqrt(a^2*x^2 + 1)*(I*a*x - 1)*x^m/(a^2*x^2 - 2*I*a*x - 1), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{x^m \sqrt{a^2 x^2 + 1}}{a^3 x^3 - 3ia^2 x^2 - 3ax + i} dx + \int \frac{a^2 x^2 x^m \sqrt{a^2 x^2 + 1}}{a^3 x^3 - 3ia^2 x^2 - 3ax + i} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**m/(1+I*a*x)**3*(a**2*x**2+1)**(3/2),x)`

[Out] `I*(Integral(x**m*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x) + Integral(a**2*x**2*x**m*sqrt(a**2*x**2 + 1)/(a**3*x**3 - 3*I*a**2*x**2 - 3*a*x + I), x))`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/(1+I*a*x)³*(a²*x²+1)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m (a^2 x^2 + 1)^{3/2}}{(1 + a x i)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*(a²*x² + 1)^(3/2))/(a*x*1i + 1)³,x)

[Out] int((x^m*(a²*x² + 1)^(3/2))/(a*x*1i + 1)³, x)

3.144 $\int e^{\frac{5}{2}i \operatorname{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; \frac{5}{4}, -\frac{5}{4}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, -5/4, 5/4, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; \frac{5}{4}, -\frac{5}{4}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((5*I)/2)*\operatorname{ArcTan}[a*x]}*x^m, x]$

[Out] $(x^{(1+m)} \operatorname{AppellF1}[1+m, 5/4, -5/4, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\operatorname{Int}[(b_*)^{(x_*)^{(m_*)}}*((c_*) + (d_*)^{(x_*)^{(n_*)}}*((e_*) + (f_*)^{(x_*)^{(p_*)}}), x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[c^n * e^p * ((b*x)^{(m+1)}) / (b*(m+1))] * \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5170

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*)^{(x_*)}] * (n_*))^{(x_*)^{(m_*)}}, x_{\text{Symbol}}] \rightarrow \operatorname{Int}[x^m * ((1 - I*a*x)^{(I*(n/2)}) / (1 + I*a*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{\frac{5}{2}i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1+iax)^{5/4}}{(1-iax)^{5/4}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; \frac{5}{4}, -\frac{5}{4}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.22, size = 0, normalized size = 0.00

$$\int e^{\frac{5}{2}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^(((5*I)/2)*ArcTan[a*x])*x^m,x]**[Out]** Integrate[E^(((5*I)/2)*ArcTan[a*x])*x^m, x]**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{5}{2}} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^m,x)**[Out]** int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^m,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^m,x, algorithm="maxima")**[Out]** integrate(x^m*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^m,x, algorithm="fricas")**[Out]** integral(-(a*x - I)*x^m*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/(a*x + I), x)**Sympy [F(-1)]** Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2)*x**m,x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2)*x^m,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.03
```

$$\int x^m \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{5/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)
```

```
[Out] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

3.145 $\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; \frac{3}{4}, -\frac{3}{4}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, -3/4, 3/4, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; \frac{3}{4}, -\frac{3}{4}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)/2)*\text{ArcTan}[a*x]}*x^m, x]$

[Out] $(x^{(1+m)} \text{AppellF1}[1+m, 3/4, -3/4, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\text{Int}[(b_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}*((e_*) + (f_*)*(x_*)^{(p_*)}), x_Symbol] \rightarrow \text{Simp}[c^n * e^p * ((b*x)^{(m+1})/(b*(m+1)))*\text{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; \text{FreeQ}[\{b, c, d, e, f, m, n, p\}, x] \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c, 0] \& \& (\text{IntegerQ}[p] \parallel \text{GtQ}[e, 0])$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_*)] * (n_*))} * (x_*)^{(m_*)}, x_Symbol] \rightarrow \text{Int}[x^m * ((1 - I*a*x)^{(I*(n/2))} / (1 + I*a*x)^{(I*(n/2))}), x] /; \text{FreeQ}[\{a, m, n\}, x] \& \& \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{\frac{3}{2}i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1+iax)^{3/4}}{(1-iax)^{3/4}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; \frac{3}{4}, -\frac{3}{4}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.17, size = 0, normalized size = 0.00

$$\int e^{\frac{3}{2}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^(((3*I)/2)*ArcTan[a*x])*x^m, x]

[Out] Integrate[E^(((3*I)/2)*ArcTan[a*x])*x^m, x]

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{3}{2}} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^m, x)

[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^m, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^m, x, algorithm="maxima")

[Out] integrate(x^m*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^m, x, algorithm="fricas")

[Out] integral(I*sqrt(a^2*x^2 + 1)*x^m*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/(a*x + I), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2)*x**m,x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2)*x^m,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.03
```

$$\int x^m \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)
```

3.146 $\int e^{\frac{1}{2}i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; \frac{1}{4}, -\frac{1}{4}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, -1/4, 1/4, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; \frac{1}{4}, -\frac{1}{4}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] Int[E^((I/2)*ArcTan[a*x])*x^m,x]

[Out] (x^(1+m)*AppellF1[1+m, 1/4, -1/4, 2+m, I*a*x, (-I)*a*x])/(1+m)

Rule 138

Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[c^n*e^p*((b*x)^(m+1)/(b*(m+1)))*AppellF1[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{\frac{1}{2}i\text{tan}^{-1}(ax)} x^m dx &= \int \frac{x^m \sqrt[4]{1+iax}}{\sqrt[4]{1-iax}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; \frac{1}{4}, -\frac{1}{4}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.17, size = 0, normalized size = 0.00

$$\int e^{\frac{1}{2}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

`[In] Integrate[E^((I/2)*ArcTan[a*x])*x^m,x]``[Out] Integrate[E^((I/2)*ArcTan[a*x])*x^m, x]`**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^m,x)``[Out] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^m,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^m,x, algorithm="maxima")``[Out] integrate(x^m*sqrt((I*a*x + 1)/sqrt(a^2*x^2 + 1)), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^m,x, algorithm="fricas")``[Out] integral(x^m*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I)), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x^m \sqrt{\frac{i(ax - i)}{\sqrt{a^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2)*x**m,x)
```

```
[Out] Integral(x**m*sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2)*x^m,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int x^m \sqrt{\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)
```

```
[Out] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)
```

3.147 $\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; -\frac{1}{4}, \frac{1}{4}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, 1/4, -1/4, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; -\frac{1}{4}, \frac{1}{4}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^m/E^{((I/2)*\text{ArcTan}[a*x])}, x]$

[Out] $(x^{(1+m)} \text{AppellF1}[1+m, -1/4, 1/4, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\text{Int}[(b_*) (x_*)^{(m_*)} ((c_*) + (d_*) (x_*))^{(n_*)} ((e_*) + (f_*) (x_*))^{(p_*)}, x_Symbol] \rightarrow \text{Simp}[c^n e^p ((b*x)^{(m+1})/(b*(m+1))) * \text{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; \text{FreeQ}[\{b, c, d, e, f, m, n, p\}, x] \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c, 0] \& \& (\text{IntegerQ}[p] \parallel \text{GtQ}[e, 0])$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_*) (x_*)] * (n_*))} (x_*)^{(m_*)}, x_Symbol] \rightarrow \text{Int}[x^m * ((1 - I*a*x)^{(I*(n/2)}) / (1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}[\{a, m, n\}, x] \& \& \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{-\frac{1}{2}i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m \sqrt[4]{1-iax}}{\sqrt[4]{1+iax}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; -\frac{1}{4}, \frac{1}{4}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.19, size = 0, normalized size = 0.00

$$\int e^{-\frac{1}{2}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[x^m/E^{((I/2)*ArcTan[a*x])}, x]**[Out]** Integrate[x^m/E^{((I/2)*ArcTan[a*x])}, x]**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m}{\sqrt{\frac{iax + 1}{\sqrt{a^2x^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m/((1+I*a*x)/(a²*x²+1)^(1/2))^(1/2), x)**[Out]** int(x^m/((1+I*a*x)/(a²*x²+1)^(1/2))^(1/2), x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/((1+I*a*x)/(a²*x²+1)^(1/2))^(1/2), x, algorithm="maxima")**[Out]** integrate(x^m/sqrt((I*a*x + 1)/sqrt(a²*x² + 1)), x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/((1+I*a*x)/(a²*x²+1)^(1/2))^(1/2), x, algorithm="fricas")**[Out]** integral(-I*sqrt(a²*x² + 1)*x^m*sqrt(I*sqrt(a²*x² + 1)/(a*x + I))/(a*x - I), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m}{\sqrt{\frac{i(ax-i)}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x**m/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/2),x)``[Out] Integral(x**m/sqrt(I*(a*x - I)/sqrt(a**2*x**2 + 1)), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/2),x, algorithm="giac")``[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{x^m}{\sqrt{\frac{1+ax\text{li}}{\sqrt{a^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^m/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2),x)``[Out] int(x^m/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/2), x)`

$$3.148 \quad \int e^{-\frac{3}{2}i \operatorname{ArcTan}(ax)} x^m dx$$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; -\frac{3}{4}, \frac{3}{4}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 3/4, -3/4, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; -\frac{3}{4}, \frac{3}{4}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[x^m/E^{((3*I)/2)*\operatorname{ArcTan}[a*x]}, x]$

[Out] $(x^{(1+m)} \operatorname{AppellF1}[1+m, -3/4, 3/4, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\operatorname{Int}[(b_*)^m * ((c_*) + (d_*) * (x_*))^{(n_*)} * ((e_*) + (f_*) * (x_*))^{(p_*)}, x_Symbol] \rightarrow \operatorname{Simp}[c_*^{n_*} e^{p_*} * ((b_* x)^{(m+1)}) / (b_*^{(m+1)}) * \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d_*) * (x/c), (-f_*) * (x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] & & !IntegerQ[m] & & !IntegerQ[n] & & GtQ[c, 0] & & (IntegerQ[p] || GtQ[e, 0])

Rule 5170

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*) * (x_*)]) * (n_*)} * (x_*)^{(m_*)}, x_Symbol] \rightarrow \operatorname{Int}[x^m * ((1 - I*a*x)^{(I*(n/2)}) / (1 + I*a*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, m, n}, x] & & !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-\frac{3}{2}i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 - iax)^{3/4}}{(1 + iax)^{3/4}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; -\frac{3}{4}, \frac{3}{4}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.19, size = 0, normalized size = 0.00

$$\int e^{-\frac{3}{2}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

`[In] Integrate[x^m/E^(((3*I)/2)*ArcTan[a*x]),x]``[Out] Integrate[x^m/E^(((3*I)/2)*ArcTan[a*x]), x]`**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)``[Out] int(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="maxima")``[Out] integrate(x^m/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(3/2), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="fricas")``[Out] integral(-(a*x + I)*x^m*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/(a*x - I), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m}{\left(\frac{i(ax-i)}{\sqrt{a^2x^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**m/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(3/2),x)
```

```
[Out] Integral(x**m/(I*(a*x - I)/sqrt(a**2*x**2 + 1))**(3/2), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int \frac{x^m}{\left(\frac{1+ax \operatorname{li}}{\sqrt{a^2 x^2 + 1}}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(x^m/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(3/2), x)
```

3.149 $\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; -\frac{5}{4}, \frac{5}{4}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, 5/4, -5/4, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; -\frac{5}{4}, \frac{5}{4}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^m/E^{((5*I)/2)*\text{ArcTan}[a*x]}, x]$

[Out] $(x^{(1+m)} \text{AppellF1}[1+m, -5/4, 5/4, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\text{Int}[(b_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}*((e_*) + (f_*)*(x_*)^{(p_*)}), x_Symbol] \rightarrow \text{Simp}[c^n * e^p * ((b*x)^{(m+1})/(b*(m+1))) * \text{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; \text{FreeQ}\{b, c, d, e, f, m, n, p\}, x] \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c, 0] \& \& (\text{IntegerQ}[p] \parallel \text{GtQ}[e, 0])$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_*)] * (n_*))} * (x_*)^{(m_*)}, x_Symbol] \rightarrow \text{Int}[x^m * ((1 - I*a*x)^{(I*(n/2))}) / (1 + I*a*x)^{(I*(n/2))}], x] /; \text{FreeQ}\{a, m, n\}, x] \& \& \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{-\frac{5}{2}i \tan^{-1}(ax)} x^m dx &= \int \frac{x^m (1 - iax)^{5/4}}{(1 + iax)^{5/4}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; -\frac{5}{4}, \frac{5}{4}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.22, size = 0, normalized size = 0.00

$$\int e^{-\frac{5}{2}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[x^m/E^(((5*I)/2)*ArcTan[a*x]),x]

[Out] Integrate[x^m/E^(((5*I)/2)*ArcTan[a*x]), x]

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m}{\left(\frac{iax+1}{\sqrt{a^2x^2+1}}\right)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

[Out] int(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="maxima")

[Out] integrate(x^m/((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(5/2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="fricas")

[Out] integral(sqrt(a^2*x^2 + 1)*(I*a*x - 1)*x^m*sqrt(I*sqrt(a^2*x^2 + 1)/(a*x + I))/(a^2*x^2 - 2*I*a*x - 1), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x**m/((1+I*a*x)/(a**2*x**2+1)**(1/2))**(5/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3006 deep
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^m/((1+I*a*x)/(a^2*x^2+1)^(1/2))^(5/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.03
```

$$\int \frac{x^m}{\left(\frac{1+axi}{\sqrt{a^2x^2+1}}\right)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2),x)
```

```
[Out] int(x^m/((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(5/2), x)
```

$$3.150 \quad \int e^{\frac{2\text{ArcTan}(x)}{3}} x^m dx$$

Optimal. Leaf size=38

$$\frac{x^{1+m} F_1\left(1+m; -\frac{i}{3}, \frac{i}{3}; 2+m; ix, -ix\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, 1/3*I, -1/3*I, 2+m, -I*x, I*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; -\frac{i}{3}, \frac{i}{3}; m+2; ix, -ix\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*\text{ArcTan}[x])/3)*x^m}, x]$

[Out] $(x^{(1+m)} \text{AppellF1}[1+m, -1/3*I, I/3, 2+m, I*x, (-I)*x])/(1+m)$

Rule 138

$\text{Int}[(b_*)^{(x_*)^{(m_*)} * ((c_*) + (d_*)^{(x_*)^{(n_*)} * ((e_*) + (f_*)^{(x_*)^{(p_*)})})}, x_Symbol] \rightarrow \text{Simp}[c^n * e^p * ((b*x)^{(m+1}) / (b*(m+1))) * \text{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; \text{FreeQ}\{b, c, d, e, f, m, n, p\}, x] \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c, 0] \& \& (\text{IntegerQ}[p] \mid \mid \text{GtQ}[e, 0])$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_*)^{(x_*)} * (n_*)^{(x_*)^{(m_*)})}], x_Symbol] \rightarrow \text{Int}[x^m * ((1 - I*a*x)^{(I*(n/2)}) / (1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, m, n\}, x] \& \& \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{\frac{2}{3} \tan^{-1}(x)} x^m dx &= \int (1 - ix)^{\frac{i}{3}} (1 + ix)^{-\frac{i}{3}} x^m dx \\ &= \frac{x^{1+m} F_1\left(1+m; -\frac{i}{3}, \frac{i}{3}; 2+m; ix, -ix\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.15, size = 0, normalized size = 0.00

$$\int e^{\frac{2\text{ArcTan}(x)}{3}} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^((2*ArcTan[x])/3)*x^m,x]

[Out] Integrate[E^((2*ArcTan[x])/3)*x^m, x]

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{\frac{2\arctan(x)}{3}} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2/3*arctan(x))*x^m,x)

[Out] int(exp(2/3*arctan(x))*x^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2/3*arctan(x))*x^m,x, algorithm="maxima")

[Out] integrate(x^m*e^(2/3*arctan(x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2/3*arctan(x))*x^m,x, algorithm="fricas")

[Out] integral(x^m*e^(2/3*arctan(x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^m e^{\frac{2\text{atan}(x)}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(2/3*atan(x))*x**m,x)`

[Out] `Integral(x**m*exp(2*atan(x)/3), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(2/3*arctan(x))*x^m,x, algorithm="giac")`

[Out] `integrate(x^m*e^(2/3*arctan(x)), x)`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int x^m e^{\frac{2 \operatorname{atan}(x)}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^m*exp((2*atan(x))/3),x)`

[Out] `int(x^m*exp((2*atan(x))/3), x)`

3.151 $\int e^{\frac{\text{ArcTan}(x)}{3}} x^m dx$

Optimal. Leaf size=38

$$\frac{x^{1+m} F_1\left(1+m; -\frac{i}{6}, \frac{i}{6}; 2+m; ix, -ix\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, 1/6*I, -1/6*I, 2+m, -I*x, I*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; -\frac{i}{6}, \frac{i}{6}; m+2; ix, -ix\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(\text{ArcTan}[x]/3)} * x^m, x]$

[Out] $(x^{(1+m)} \text{AppellF1}[1+m, -1/6*I, I/6, 2+m, I*x, (-I)*x])/(1+m)$

Rule 138

$\text{Int}[(b_*)*(x_)^{(m_*)}*((c_*) + (d_*)*(x_))^{(n_*)}*((e_*) + (f_*)*(x_))^{(p_*)}, x_ \text{Symbol}] \rightarrow \text{Simp}[c^{n*} e^{p*} ((b*x)^{(m+1})/(b*(m+1)))*\text{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; \text{FreeQ}\{b, c, d, e, f, m, n, p\}, x] \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c, 0] \& \& (\text{IntegerQ}[p] \text{ || } \text{GtQ}[e, 0])$

Rule 5170

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_)]*(n_*)}*(x_)^{(m_*)}, x_ \text{Symbol}] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2)})/(1 + I*a*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, m, n\}, x] \& \& \text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{\frac{1}{3} \tan^{-1}(x)} x^m dx &= \int (1 - ix)^{\frac{i}{6}} (1 + ix)^{-\frac{i}{6}} x^m dx \\ &= \frac{x^{1+m} F_1\left(1+m; -\frac{i}{6}, \frac{i}{6}; 2+m; ix, -ix\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.15, size = 0, normalized size = 0.00

$$\int e^{\frac{\text{ArcTan}(x)}{3}} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^(ArcTan[x]/3)*x^m,x]

[Out] Integrate[E^(ArcTan[x]/3)*x^m, x]

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{\frac{\arctan(x)}{3}} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(1/3*arctan(x))*x^m,x)

[Out] int(exp(1/3*arctan(x))*x^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(1/3*arctan(x))*x^m,x, algorithm="maxima")

[Out] integrate(x^m*e^(1/3*arctan(x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(1/3*arctan(x))*x^m,x, algorithm="fricas")

[Out] integral(x^m*e^(1/3*arctan(x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^m e^{\frac{\text{atan}(x)}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(1/3*atan(x))*x**m,x)

[Out] Integral(x**m*exp(atan(x)/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(1/3*arctan(x))*x^m,x, algorithm="giac")

[Out] integrate(x^m*e^(1/3*arctan(x)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.03

$$\int x^m e^{\frac{\arctan(x)}{3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m*exp(atan(x)/3),x)

[Out] int(x^m*exp(atan(x)/3), x)

3.152 $\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=36

$$\frac{x^{1+m} F_1\left(1+m; \frac{1}{8}, -\frac{1}{8}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \text{AppellF1}(1+m, -1/8, 1/8, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 36, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; \frac{1}{8}, -\frac{1}{8}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] Int[E^((I/4)*ArcTan[a*x])*x^m,x]

[Out] (x^(1+m)*AppellF1[1+m, 1/8, -1/8, 2+m, I*a*x, (-I)*a*x])/(1+m)

Rule 138

Int[((b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[c^n*e^p*((b*x)^(m+1)/(b*(m+1)))*AppellF1[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5170

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*(x_)^(m_.), x_Symbol] :> Int[x^m*((1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{\frac{1}{4}i\text{tan}^{-1}(ax)} x^m dx &= \int \frac{x^m \sqrt[8]{1+iax}}{\sqrt[8]{1-iax}} dx \\ &= \frac{x^{1+m} F_1\left(1+m; \frac{1}{8}, -\frac{1}{8}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.17, size = 0, normalized size = 0.00

$$\int e^{\frac{1}{4}i\text{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^((I/4)*ArcTan[a*x])*x^m,x]**[Out]** Integrate[E^((I/4)*ArcTan[a*x])*x^m, x]**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \left(\frac{iax + 1}{\sqrt{a^2x^2 + 1}} \right)^{\frac{1}{4}} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^m,x)**[Out]** int(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^m,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^m,x, algorithm="maxima")**[Out]** integrate(x^m*((I*a*x + 1)/sqrt(a^2*x^2 + 1))^(1/4), x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^m,x, algorithm="fricas")**[Out]** integral(x^m*(I*sqrt(a^2*x^2 + 1)/(a*x + I))^(1/4), x)**Sympy [F(-1)]** Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a**2*x**2+1)**(1/2))**(1/4)*x**m,x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*a*x)/(a^2*x^2+1)^(1/2))^(1/4)*x^m,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.03
```

$$\int x^m \left(\frac{1 + a x i}{\sqrt{a^2 x^2 + 1}} \right)^{1/4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4),x)
```

```
[Out] int(x^m*((a*x*1i + 1)/(a^2*x^2 + 1)^(1/2))^(1/4), x)
```

3.153 $\int e^{in \operatorname{ArcTan}(ax)} x^m dx$

Optimal. Leaf size=40

$$\frac{x^{1+m} F_1\left(1+m; \frac{n}{2}, -\frac{n}{2}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, -1/2*n, 1/2*n, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.02, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {5170, 138}

$$\frac{x^{m+1} F_1\left(m+1; \frac{n}{2}, -\frac{n}{2}; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*n*\operatorname{ArcTan}[a*x])} * x^m, x]$

[Out] $(x^{(1+m)} \operatorname{AppellF1}[1+m, n/2, -1/2*n, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\operatorname{Int}[(b_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}*((e_*) + (f_*)*(x_*)^{(p_*)}), x_Symbol] \rightarrow \operatorname{Simp}[c^n e^p ((b*x)^{(m+1})/(b*(m+1))) * \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /; \operatorname{FreeQ}\{b, c, d, e, f, m, n, p\}, x] \& \& \operatorname{IntegerQ}[m] \& \& \operatorname{IntegerQ}[n] \& \& \operatorname{GtQ}[c, 0] \& \& (\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[e, 0])$

Rule 5170

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*)*(x_*)] * (n_*)} * (x_*)^{(m_*)}, x_Symbol] \rightarrow \operatorname{Int}[x^m * ((1 - I*a*x)^{(I*(n/2)}) / (1 + I*a*x)^{(I*(n/2)})), x] /; \operatorname{FreeQ}\{a, m, n\}, x] \& \& \operatorname{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{in \tan^{-1}(ax)} x^m dx &= \int x^m (1 - iax)^{-n/2} (1 + iax)^{n/2} dx \\ &= \frac{x^{1+m} F_1\left(1+m; \frac{n}{2}, -\frac{n}{2}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.17, size = 0, normalized size = 0.00

$$\int e^{in \operatorname{ArcTan}(ax)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^(I*n*ArcTan[a*x])*x^m,x]

[Out] Integrate[E^(I*n*ArcTan[a*x])*x^m, x]

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{in \arctan(ax)} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(I*n*arctan(a*x))*x^m,x)

[Out] int(exp(I*n*arctan(a*x))*x^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^m,x, algorithm="maxima")

[Out] integrate(x^m*e^(I*n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))*x^m,x, algorithm="fricas")

[Out] integral(x^m/(-(a*x + I)/(a*x - I))^(1/2*n), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^m e^{in \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))*x**m,x)

[Out] Integral(x**m*exp(I*n*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^m,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int x^m e^{n \operatorname{atan}(ax) 1i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^m*exp(n*atan(a*x)*1i),x)

[Out] int(x^m*exp(n*atan(a*x)*1i), x)

3.154 $\int e^{in\text{ArcTan}(ax)} x^3 dx$

Optimal. Leaf size=171

$$\frac{x^2(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}}{4a^2} - \frac{(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}(6+n^2+2ianx)}{24a^4} - \frac{2^{-2+\frac{n}{2}}n(8+n^2)(1-iax)^{1-\frac{n}{2}}}{3a^4}$$

[Out] $\frac{1}{4}x^2(1-Iax)^{(1-1/2*n)}(1+Iax)^{(1+1/2*n)}/a^2-1/24*(1-Iax)^{(1-1/2*n)}*(1+Iax)^{(1+1/2*n)}*(6+n^2+2*I*a*n*x)/a^4-1/3*2^{(-2+1/2*n)}*n*(n^2+8)*(1-Iax)^{(1-1/2*n)}*\text{hypergeom}([-1/2*n, 1-1/2*n], [2-1/2*n], 1/2-1/2*Iax)/a^4/(2-n)$

Rubi [A]

time = 0.08, antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$,

Rules used = {5170, 102, 152, 71}

$$\frac{2^{\frac{n}{2}-2}n(n^2+8)(1-iax)^{1-\frac{n}{2}}{}_2F_1(1-\frac{n}{2}, -\frac{n}{2}; 2-\frac{n}{2}; \frac{1}{2}(1-iax))}{3a^4(2-n)} - \frac{(1+iax)^{\frac{n+2}{2}}(2ianx+n^2+6)(1-iax)^{1-\frac{n}{2}}}{24a^4} + \frac{x^2(1+iax)^{\frac{n+2}{2}}(1-iax)^{1-\frac{n}{2}}}{4a^2}$$

Antiderivative was successfully verified.

[In] Int[E^(I*n*ArcTan[a*x])*x^3,x]

[Out] $(x^2*(1-Iax)^{(1-n/2)}*(1+Iax)^{((2+n)/2)})/(4*a^2) - ((1-Iax)^{(1-n/2)}*(1+Iax)^{((2+n)/2)}*(6+n^2+(2*I)*a*n*x))/(24*a^4) - (2^{(-2+n/2)}*n*(8+n^2)*(1-Iax)^{(1-n/2)}*\text{Hypergeometric2F1}[1-n/2, -1/2*n, 2-n/2, (1-Iax)/2])/(3*a^4*(2-n))$

Rule 71

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 102

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]

Rule 152

```

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))
)*(g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x))*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n},
x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^m, x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int e^{in \tan^{-1}(ax)} x^3 dx &= \int x^3 (1 - iax)^{-n/2} (1 + iax)^{n/2} dx \\
&= \frac{x^2(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{4a^2} + \frac{\int x(1 - iax)^{-n/2}(1 + iax)^{n/2}(-2 - ianx) dx}{4a^2} \\
&= \frac{x^2(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{4a^2} - \frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}(6 + n^2 + 2ianx)}{24a^4} + \frac{(in(8 + n^2) - 2ianx)}{24a^4} \\
&= \frac{x^2(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{4a^2} - \frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}(6 + n^2 + 2ianx)}{24a^4} - \frac{2^{-2+\frac{n}{2}}n(8 + n^2 - 2ianx)}{24a^4}
\end{aligned}$$

Mathematica [A]

time = 0.61, size = 158, normalized size = 0.92

$$\frac{e^{in \operatorname{ArcTan}(ax)} \left(-e^{2i \operatorname{ArcTan}(ax)} n^2 (8 + n^2) {}_2F_1\left(1, 1 + \frac{n}{2}; 2 + \frac{n}{2}; -e^{2i \operatorname{ArcTan}(ax)}\right) + (2 + n) \left(ian(8 + n^2)x - (12 + n^2 + 2ianx)(1 + a^2x^2) + 6(1 + a^2x^2)^2 + n(8 + n^2) {}_2F_1\left(1, \frac{n}{2}; 1 + \frac{n}{2}; -e^{2i \operatorname{ArcTan}(ax)}\right) \right) \right)}{24a^4(2 + n)}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(I*n*ArcTan[a*x])*x^3,x]
```

```
[Out] (E^(I*n*ArcTan[a*x])*(-(E^((2*I)*ArcTan[a*x]))*n^2*(8 + n^2)*Hypergeometric2
F1[1, 1 + n/2, 2 + n/2, -E^((2*I)*ArcTan[a*x])]) + (2 + n)*(I*a*n*(8 + n^2)
*x - (12 + n^2 + (2*I)*a*n*x)*(1 + a^2*x^2) + 6*(1 + a^2*x^2)^2 + n*(8 + n^
2)*Hypergeometric2F1[1, n/2, 1 + n/2, -E^((2*I)*ArcTan[a*x])])))/(24*a^4*(2
+ n))

```


Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{in \arctan(ax)} x^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(I*n*arctan(a*x))*x^3,x)

[Out] int(exp(I*n*arctan(a*x))*x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^3,x, algorithm="maxima")

[Out] integrate(x^3*e^(I*n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^3,x, algorithm="fricas")

[Out] integral(x^3/(-(a*x + I)/(a*x - I))^(1/2*n), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 e^{in \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))*x**3,x)

[Out] Integral(x**3*exp(I*n*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^3,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^3 e^{n \operatorname{atan}(ax) i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^3*exp(n*atan(a*x)*1i),x)

[Out] int(x^3*exp(n*atan(a*x)*1i), x)

3.155 $\int e^{in\text{ArcTan}(ax)} x^2 dx$

Optimal. Leaf size=159

$$-\frac{in(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}}{6a^3} + \frac{x(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}}{3a^2} - \frac{i2^{n/2}(2+n^2)(1-iax)^{1-\frac{n}{2}} {}_2F_1\left(1-\frac{n}{2}, -\frac{n}{2}; 2-\frac{n}{2}; 2-\frac{n}{2}\right)}{3a^3(2-n)}$$

[Out] $-1/6*I*n*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(1+1/2*n)}/a^3+1/3*x*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(1+1/2*n)}/a^2-1/3*I*2^{(1/2*n)}*(n^2+2)*(1-I*a*x)^{(1-1/2*n)}*\text{hypergeom}([-1/2*n, 1-1/2*n], [2-1/2*n], 1/2-1/2*I*a*x)/a^3/(2-n)$

Rubi [A]

time = 0.06, antiderivative size = 159, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {5170, 92, 81, 71}

$$-\frac{i2^{n/2}(n^2+2)(1-iax)^{1-\frac{n}{2}} {}_2F_1\left(1-\frac{n}{2}, -\frac{n}{2}; 2-\frac{n}{2}; \frac{1}{2}(1-iax)\right)}{3a^3(2-n)} - \frac{in(1+iax)^{\frac{n+2}{2}}(1-iax)^{1-\frac{n}{2}}}{6a^3} + \frac{x(1+iax)^{\frac{n+2}{2}}(1-iax)^{1-\frac{n}{2}}}{3a^2}$$

Antiderivative was successfully verified.

[In] Int[E^(I*n*ArcTan[a*x])*x^2,x]

[Out] $((-1/6*I)*n*(1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((2+n)/2)}/a^3 + (x*(1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((2+n)/2)})/(3*a^2) - ((I/3)*2^{(n/2)}*(2+n^2)*(1-I*a*x)^{(1-n/2)}*\text{Hypergeometric2F1}[1-n/2, -1/2*n, 2-n/2, (1-I*a*x)/2])/(a^3*(2-n))$

Rule 71

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 81

Int[((a_) + (b_)*(x_))*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92

Int[((a_) + (b_)*(x_))^(2)*((c_) + (d_)*(x_))^(n_)*((e_) + (f_)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(

```
d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)
^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b
*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ
[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int e^{in \tan^{-1}(ax)} x^2 dx &= \int x^2 (1 - iax)^{-n/2} (1 + iax)^{n/2} dx \\ &= \frac{x(1 - iax)^{1-\frac{n}{2}} (1 + iax)^{\frac{2+n}{2}}}{3a^2} + \frac{\int (1 - iax)^{-n/2} (1 + iax)^{n/2} (-1 - ianx) dx}{3a^2} \\ &= -\frac{in(1 - iax)^{1-\frac{n}{2}} (1 + iax)^{\frac{2+n}{2}}}{6a^3} + \frac{x(1 - iax)^{1-\frac{n}{2}} (1 + iax)^{\frac{2+n}{2}}}{3a^2} - \frac{(2 + n^2) \int (1 - iax)^{-n/2} dx}{6a^2} \\ &= -\frac{in(1 - iax)^{1-\frac{n}{2}} (1 + iax)^{\frac{2+n}{2}}}{6a^3} + \frac{x(1 - iax)^{1-\frac{n}{2}} (1 + iax)^{\frac{2+n}{2}}}{3a^2} - \frac{i2^{n/2}(2 + n^2)(1 - iax)^{-n/2}}{6a^2} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 116, normalized size = 0.73

$$\frac{(1 - iax)^{-n/2} (i + ax) ((-2 + n)(1 + iax)^{n/2} (-i + ax) (-in + 2ax) + 2^{1+\frac{n}{2}} (2 + n^2) {}_2F_1(1 - \frac{n}{2}, -\frac{n}{2}; 2 - \frac{n}{2}, \frac{1}{2}(1 - iax)))}{6a^3(-2 + n)}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(I*n*ArcTan[a*x])*x^2,x]
```

```
[Out] ((I + a*x)*((-2 + n)*(1 + I*a*x)^(n/2)*(-I + a*x)*((-I)*n + 2*a*x) + 2^(1 +
n/2)*(2 + n^2)*Hypergeometric2F1[1 - n/2, -1/2*n, 2 - n/2, (1 - I*a*x)/2])
)/(6*a^3*(-2 + n)*(1 - I*a*x)^(n/2))
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{in \arctan(ax)} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(I*n*arctan(a*x))*x^2,x)`

[Out] `int(exp(I*n*arctan(a*x))*x^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*arctan(a*x))*x^2,x, algorithm="maxima")`

[Out] `integrate(x^2*e^(I*n*arctan(a*x)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*arctan(a*x))*x^2,x, algorithm="fricas")`

[Out] `integral(x^2/(-(a*x + I)/(a*x - I))^(1/2*n), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 e^{in \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*atan(a*x))*x**2,x)`

[Out] `Integral(x**2*exp(I*n*atan(a*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*arctan(a*x))*x^2,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x^2 e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^2*exp(n*atan(a*x)*1i),x)
```

```
[Out] int(x^2*exp(n*atan(a*x)*1i), x)
```

3.156 $\int e^{in \operatorname{ArcTan}(ax)} x dx$

Optimal. Leaf size=107

$$\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{2a^2} + \frac{2^{n/2}n(1 - iax)^{1-\frac{n}{2}} {}_2F_1\left(1 - \frac{n}{2}, -\frac{n}{2}; 2 - \frac{n}{2}; \frac{1}{2}(1 - iax)\right)}{a^2(2 - n)}$$

[Out] $1/2*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(1+1/2*n)}/a^2+2^{(1/2*n)*n}*(1-I*a*x)^{(1-1/2*n)}*\operatorname{hypergeom}([-1/2*n, 1-1/2*n], [2-1/2*n], 1/2-1/2*I*a*x)/a^2/(2-n)$

Rubi [A]

time = 0.03, antiderivative size = 107, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 13, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {5170, 81, 71}

$$\frac{2^{n/2}n(1 - iax)^{1-\frac{n}{2}} {}_2F_1\left(1 - \frac{n}{2}, -\frac{n}{2}; 2 - \frac{n}{2}; \frac{1}{2}(1 - iax)\right)}{a^2(2 - n)} + \frac{(1 + iax)^{\frac{n+2}{2}}(1 - iax)^{1-\frac{n}{2}}}{2a^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*n*\operatorname{ArcTan}[a*x])}*x, x]$

[Out] $((1 - I*a*x)^{(1 - n/2)}*(1 + I*a*x)^{((2 + n)/2)})/(2*a^2) + (2^{(n/2)*n}*(1 - I*a*x)^{(1 - n/2)}*\operatorname{Hypergeometric2F1}[1 - n/2, -1/2*n, 2 - n/2, (1 - I*a*x)/2])/(a^2*(2 - n))$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)})], x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& \operatorname{IntegerQ}[m] \&\& \operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}\{b/(b*c - a*d), 0\} \&\& (\operatorname{RationalQ}[m] \mid\mid \operatorname{!(RationalQ}[n] \&\& \operatorname{GtQ}\{-d/(b*c - a*d), 0\}))$

Rule 81

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(c_+)}*((d_+)*(x_+))^{(n_+)}*((e_+ + (f_+)*(x_+))^{(p_+)})], x_Symbol] \rightarrow \operatorname{Simp}[b*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 2))), x] + \operatorname{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(d*f*(n + p + 2)), \operatorname{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x \&\& \operatorname{NeQ}[n + p + 2, 0]$

Rule 5170

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a_+]*(x_+))^{(n_+)}}*(x_+)^{(m_+)}, x_Symbol] \rightarrow \operatorname{Int}[x^m*((1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}), x] /; \operatorname{FreeQ}\{a, m, n\}, x \&\& \operatorname{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned}
\int e^{in \tan^{-1}(ax)} x dx &= \int x(1 - iax)^{-n/2}(1 + iax)^{n/2} dx \\
&= \frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{2a^2} - \frac{(in) \int (1 - iax)^{-n/2}(1 + iax)^{n/2} dx}{2a} \\
&= \frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{2a^2} + \frac{2^{n/2}n(1 - iax)^{1-\frac{n}{2}} {}_2F_1\left(1 - \frac{n}{2}, -\frac{n}{2}; 2 - \frac{n}{2}; \frac{1}{2}(1 - iax)\right)}{a^2(2 - n)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 105, normalized size = 0.98

$$\frac{(1 - iax)^{-n/2}(i + ax) \left((-2 + n)(1 + iax)^{n/2}(-i + ax) + i2^{1+\frac{n}{2}}n {}_2F_1\left(1 - \frac{n}{2}, -\frac{n}{2}; 2 - \frac{n}{2}; \frac{1}{2}(1 - iax)\right) \right)}{2a^2(-2 + n)}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*n*ArcTan[a*x])*x,x]`

```
[Out] ((I + a*x)*((-2 + n)*(1 + I*a*x)^(n/2)*(-I + a*x) + I*2^(1 + n/2)*n*Hypergeometric2F1[1 - n/2, -1/2*n, 2 - n/2, (1 - I*a*x)/2]))/(2*a^2*(-2 + n)*(1 - I*a*x)^(n/2))
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{in \arctan(ax)} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(I*n*arctan(a*x))*x,x)``[Out] int(exp(I*n*arctan(a*x))*x,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*arctan(a*x))*x,x, algorithm="maxima")``[Out] integrate(x*e^(I*n*arctan(a*x)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*arctan(a*x))*x,x, algorithm="fricas")``[Out] integral(x/(-(a*x + I)/(a*x - I))^(1/2*n), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x e^{in \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*atan(a*x))*x,x)``[Out] Integral(x*exp(I*n*atan(a*x)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*arctan(a*x))*x,x, algorithm="giac")``[Out] sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int x e^{n \operatorname{atan}(ax) 1i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x*exp(n*atan(a*x)*1i),x)``[Out] int(x*exp(n*atan(a*x)*1i), x)`

3.157 $\int e^{in \operatorname{ArcTan}(ax)} dx$

Optimal. Leaf size=71

$$\frac{i2^{1+\frac{n}{2}}(1-iax)^{1-\frac{n}{2}} {}_2F_1\left(1-\frac{n}{2}, -\frac{n}{2}; 2-\frac{n}{2}; \frac{1}{2}(1-iax)\right)}{a(2-n)}$$

[Out] $I*2^{(1+1/2*n)}*(1-I*a*x)^{(1-1/2*n)}*\operatorname{hypergeom}([-1/2*n, 1-1/2*n], [2-1/2*n], 1/2-1/2*I*a*x)/a/(2-n)$

Rubi [A]

time = 0.01, antiderivative size = 71, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 11, $\frac{\text{number of rules}}{\text{integrand size}} = 0.182$, Rules used = {5169, 71}

$$\frac{i2^{\frac{n}{2}+1}(1-iax)^{1-\frac{n}{2}} {}_2F_1\left(1-\frac{n}{2}, -\frac{n}{2}; 2-\frac{n}{2}; \frac{1}{2}(1-iax)\right)}{a(2-n)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*n*\operatorname{ArcTan}[a*x])}, x]$

[Out] $(I*2^{(1+n/2)}*(1-I*a*x)^{(1-n/2)}*\operatorname{Hypergeometric2F1}[1-n/2, -1/2*n, 2-n/2, (1-I*a*x)/2])/(a*(2-n))$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] := \operatorname{Simp}[(a + b*x)^{(m + 1)} / (b*(m + 1)*(b*(b*c - a*d))^{(n)}) * \operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x]$
 $\&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{!IntegerQ}[m] \&\& \operatorname{!IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] \operatorname{||} \operatorname{!(RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5169

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_+)*(x_+)]*(n_+))}, x_Symbol] := \operatorname{Int}[(1 - I*a*x)^{(I*(n/2))} / (1 + I*a*x)^{(I*(n/2))}, x] /;$ $\operatorname{FreeQ}\{a, n\}, x] \&\& \operatorname{!IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{in \tan^{-1}(ax)} dx &= \int (1-iax)^{-n/2} (1+iax)^{n/2} dx \\ &= \frac{i2^{1+\frac{n}{2}}(1-iax)^{1-\frac{n}{2}} {}_2F_1\left(1-\frac{n}{2}, -\frac{n}{2}; 2-\frac{n}{2}; \frac{1}{2}(1-iax)\right)}{a(2-n)} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 53, normalized size = 0.75

$$\frac{4ie^{i(2+n)\text{ArcTan}(ax)} {}_2F_1\left(2, 1 + \frac{n}{2}; 2 + \frac{n}{2}; -e^{2i\text{ArcTan}(ax)}\right)}{a(2+n)}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*n*ArcTan[a*x]), x]``[Out] ((-4*I)*E^(I*(2+n)*ArcTan[a*x])*Hypergeometric2F1[2, 1 + n/2, 2 + n/2, -E^((2*I)*ArcTan[a*x])])/(a*(2+n))`**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{in \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(I*n*arctan(a*x)), x)``[Out] int(exp(I*n*arctan(a*x)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*arctan(a*x)), x, algorithm="maxima")``[Out] integrate(e^(I*n*arctan(a*x)), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*arctan(a*x)), x, algorithm="fricas")``[Out] integral(1/((-a*x + I)/(a*x - I))^(1/2*n), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{in \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x)),x)

[Out] Integral(exp(I*n*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x)),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax) \operatorname{li}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x)*1i),x)

[Out] int(exp(n*atan(a*x)*1i), x)

3.158 $\int \frac{e^{in \operatorname{ArcTan}(ax)}}{x} dx$

Optimal. Leaf size=125

$$\frac{2(1-iax)^{-n/2}(1+iax)^{n/2} {}_2F_1\left(1, -\frac{n}{2}; 1 - \frac{n}{2}, \frac{1-iax}{1+iax}\right)}{n} - \frac{2^{1+\frac{n}{2}}(1-iax)^{-n/2} {}_2F_1\left(-\frac{n}{2}, -\frac{n}{2}; 1 - \frac{n}{2}; \frac{1}{2}(1-iax)\right)}{n}$$

[Out] $2*(1+I*a*x)^{(1/2*n)}*\operatorname{hypergeom}([1, -1/2*n], [1-1/2*n], (1-I*a*x)/(1+I*a*x))/n/((1-I*a*x)^{(1/2*n)})-2^{(1+1/2*n)}*\operatorname{hypergeom}([-1/2*n, -1/2*n], [1-1/2*n], 1/2-1/2*I*a*x)/n/((1-I*a*x)^{(1/2*n)})$

Rubi [A]

time = 0.04, antiderivative size = 125, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.267$, Rules used = {5170, 132, 71, 133}

$$\frac{2(1-iax)^{-n/2}(1+iax)^{n/2} {}_2F_1\left(1, -\frac{n}{2}; 1 - \frac{n}{2}, \frac{1-iax}{1+iax}\right)}{n} - \frac{2^{\frac{n}{2}+1}(1-iax)^{-n/2} {}_2F_1\left(-\frac{n}{2}, -\frac{n}{2}; 1 - \frac{n}{2}; \frac{1}{2}(1-iax)\right)}{n}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*n*\operatorname{ArcTan}[a*x])}/x, x]$

[Out] $(2*(1 + I*a*x)^{(n/2)}*\operatorname{Hypergeometric2F1}[1, -1/2*n, 1 - n/2, (1 - I*a*x)/(1 + I*a*x)])/(n*(1 - I*a*x)^{(n/2)}) - (2^{(1 + n/2)}*\operatorname{Hypergeometric2F1}[-1/2*n, -1/2*n, 1 - n/2, (1 - I*a*x)/2])/(n*(1 - I*a*x)^{(n/2)})$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)})^{(p_+)}, x_Symbol] := \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& !\operatorname{IntegerQ}\{m\} \&\& !\operatorname{IntegerQ}\{n\} \&\& \operatorname{GtQ}\{b/(b*c - a*d), 0\} \&\& (\operatorname{RationalQ}\{m\} || !(\operatorname{RationalQ}\{n\} \&\& \operatorname{GtQ}\{-d/(b*c - a*d), 0\}))$

Rule 132

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)})^{(p_+)}, x_Symbol] := \operatorname{Dist}[b*d^{(m + n)}*f^p, \operatorname{Int}[(a + b*x)^{(m - 1)}/(c + d*x)^m, x] + \operatorname{Int}[(a + b*x)^{(m - 1)}*((e + f*x)^p/(c + d*x)^m)*\operatorname{ExpandToSum}[(a + b*x)*(c + d*x)^{(-p - 1)} - (b*d^{(-p - 1)}*f^p)/(e + f*x)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, n\}, x \&\& \operatorname{EqQ}\{m + n + p + 1, 0\} \&\& \operatorname{ILtQ}\{p, 0\} \&\& (\operatorname{GtQ}\{m, 0\} || \operatorname{SumSimplerQ}\{m, -1\} || !(\operatorname{GtQ}\{n, 0\} || \operatorname{SumSimplerQ}\{n, -1\}))$

Rule 133

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)})^{(p_+)}, x_Symbol] := \operatorname{Simp}[(b*c - a*d)^n*((a + b*x)^{(m + 1)}/((m + 1)*(b*e -$

```
a*f)^(n + 1)*(e + f*x)^(m + 1))*Hypergeometric2F1[m + 1, -n, m + 2, (-(d*
e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x))), x] /; FreeQ[{a, b, c, d, e,
f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1]
|| !SumSimplerQ[p, 1]) && !ILtQ[m, 0]
```

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Integ
rQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{in \tan^{-1}(ax)}}{x} dx &= \int \frac{(1 - iax)^{-n/2} (1 + iax)^{n/2}}{x} dx \\ &= - \left((ia) \int (1 - iax)^{-1 - \frac{n}{2}} (1 + iax)^{n/2} dx \right) + \int \frac{(1 - iax)^{-1 - \frac{n}{2}} (1 + iax)^{n/2}}{x} dx \\ &= \frac{2(1 - iax)^{-n/2} (1 + iax)^{n/2} {}_2F_1\left(1, -\frac{n}{2}; 1 - \frac{n}{2}; \frac{1 - iax}{1 + iax}\right)}{n} - \frac{2^{1 + \frac{n}{2}} (1 - iax)^{-n/2} {}_2F_1\left(-\frac{n}{2}, -\frac{n}{2}; 1 - \frac{n}{2}; \frac{1 - iax}{1 + iax}\right)}{n} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 106, normalized size = 0.85

$$\frac{2(1 - iax)^{-n/2} \left((1 + iax)^{n/2} {}_2F_1\left(1, -\frac{n}{2}; 1 - \frac{n}{2}; \frac{i+ax}{i-ax}\right) - 2^{n/2} {}_2F_1\left(-\frac{n}{2}, -\frac{n}{2}; 1 - \frac{n}{2}; \frac{1}{2}(1 - iax)\right) \right)}{n}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(I*n*ArcTan[a*x])/x,x]
```

```
[Out] (2*((1 + I*a*x)^(n/2)*Hypergeometric2F1[1, -1/2*n, 1 - n/2, (I + a*x)/(I -
a*x)] - 2^(n/2)*Hypergeometric2F1[-1/2*n, -1/2*n, 1 - n/2, (1 - I*a*x)/2]))
/(n*(1 - I*a*x)^(n/2))
```

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \arctan(ax)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(I*n*arctan(a*x))/x,x)
```

[Out] `int(exp(I*n*arctan(a*x))/x,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*arctan(a*x))/x,x, algorithm="maxima")`

[Out] `integrate(e^(I*n*arctan(a*x))/x, x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*arctan(a*x))/x,x, algorithm="fricas")`

[Out] `integral(1/(x*(-(a*x + I)/(a*x - I))^(1/2*n)), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \operatorname{atan}(ax)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*atan(a*x))/x,x)`

[Out] `Integral(exp(I*n*atan(a*x))/x, x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(I*n*arctan(a*x))/x,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax) i}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*atan(a*x)*1i)/x,x)`

[Out] `int(exp(n*atan(a*x)*1i)/x, x)`

$$3.159 \quad \int \frac{e^{in \operatorname{ArcTan}(ax)}}{x^2} dx$$

Optimal. Leaf size=79

$$-\frac{4ia(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{1}{2}(-2+n)} {}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}; \frac{1-iax}{1+iax}\right)}{2-n}$$

[Out] $-4*I*a*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(-1+1/2*n)}*hypergeom([2, 1-1/2*n], [2-1/2*n], (1-I*a*x)/(1+I*a*x))/(2-n)$

Rubi [A]

time = 0.02, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.133$, Rules used = {5170, 133}

$$-\frac{4ia(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{n-2}{2}} {}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}; \frac{1-iax}{1+iax}\right)}{2-n}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I*n*ArcTan[a*x])}/x^2, x]$

[Out] $((-4*I)*a*(1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((-2+n)/2)}*Hypergeometric2F1[2, 1-n/2, 2-n/2, (1-I*a*x)/(1+I*a*x)])/(2-n)$

Rule 133

$\text{Int}[(a_. + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[(b*c - a*d)^n*((a + b*x)^{(m+1)})/((m+1)*(b*e - a*f)^{(n+1)}*(e + f*x)^{(m+1)})]*Hypergeometric2F1[m+1, -n, m+2, (-d*e - c*f)*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /;$ FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

Rule 5170

$\text{Int}[E^{(ArcTan[(a_.)*(x_)]*(n_.))}*(x_)^{(m_.)}, x_Symbol] \rightarrow \text{Int}[x^m*((1 - I*a*x)^{(I*(n/2)})/(1 + I*a*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int \frac{e^{in \tan^{-1}(ax)}}{x^2} dx &= \int \frac{(1-iax)^{-n/2}(1+iax)^{n/2}}{x^2} dx \\ &= -\frac{4ia(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{1}{2}(-2+n)} {}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}; \frac{1-iax}{1+iax}\right)}{2-n} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 82, normalized size = 1.04

$$-\frac{2ia(1-iax)^{1-\frac{n}{2}}(1+iax)^{-1+\frac{n}{2}}{}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}; -\frac{1-iax}{-1-iax}\right)}{1-\frac{n}{2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*n*ArcTan[a*x])/x^2,x]**[Out]** ((-2*I)*a*(1-I*a*x)^(1-n/2)*(1+I*a*x)^(-1+n/2)*Hypergeometric2F1[2, 1-n/2, 2-n/2, -(1-I*a*x)/(-1-I*a*x)])/(1-n/2)**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \arctan(ax)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(I*n*arctan(a*x))/x^2,x)**[Out]** int(exp(I*n*arctan(a*x))/x^2,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^2,x, algorithm="maxima")**[Out]** integrate(e^(I*n*arctan(a*x))/x^2, x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^2,x, algorithm="fricas")**[Out]** integral(1/(x^2*(-(a*x + I)/(a*x - I))^(1/2*n)), x)**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \operatorname{atan}(ax)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))/x**2,x)

[Out] Integral(exp(I*n*atan(a*x))/x**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^2,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax) \operatorname{li}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x)*1i)/x^2,x)

[Out] int(exp(n*atan(a*x)*1i)/x^2, x)

3.160 $\int \frac{e^{in \operatorname{ArcTan}(ax)}}{x^3} dx$

Optimal. Leaf size=120

$$-\frac{(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}}{2x^2} + \frac{2a^2n(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{1}{2}(-2+n)} {}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}, \frac{1-iax}{1+iax}\right)}{2-n}$$

[Out] $-1/2*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(1+1/2*n)}/x^2+2*a^2*n*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(-1+1/2*n)}*hypergeom([2, 1-1/2*n], [2-1/2*n], (1-I*a*x)/(1+I*a*x))/(2-n)$

Rubi [A]

time = 0.04, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5170, 98, 133}

$$\frac{2a^2n(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{n-2}{2}} {}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}, \frac{1-iax}{1+iax}\right)}{2-n} - \frac{(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{n+2}{2}}}{2x^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I*n*ArcTan[a*x])}/x^3, x]$

[Out] $-1/2*((1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((2+n)/2)})/x^2 + (2*a^2*n*(1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((-2+n)/2)}*Hypergeometric2F1[2, 1-n/2, 2-n/2, (1-I*a*x)/(1+I*a*x)]/(2-n)$

Rule 98

$\text{Int}[(a_. + (b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.)*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] :> \text{Simp}[b*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/((m + 1)*(b*c - a*d)*(b*e - a*f))], x] + \text{Dist}[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), \text{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{EqQ}[\text{Simplify}[m + n + p + 3], 0] \&\& (\text{LtQ}[m, -1] || \text{SumSimplerQ}[m, 1])$

Rule 133

$\text{Int}[(a_. + (b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.)*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] :> \text{Simp}[(b*c - a*d)^n*((a + b*x)^{(m + 1)})/((m + 1)*(b*e - a*f)^{(n + 1)}*(e + f*x)^{(m + 1)})]*Hypergeometric2F1[m + 1, -n, m + 2, -(d*e - c*f)*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, p\}, x] \&\& \text{EqQ}[m + n + p + 2, 0] \&\& \text{ILtQ}[n, 0] \&\& (\text{SumSimplerQ}[m, 1] || !\text{SumSimplerQ}[p, 1]) \&\& !\text{ILtQ}[m, 0]$

Rule 5170

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a*x)^(I*(n/2)))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !IntegerQ[(I*n - 1)/2]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{in \tan^{-1}(ax)}}{x^3} dx &= \int \frac{(1 - iax)^{-n/2}(1 + iax)^{n/2}}{x^3} dx \\ &= -\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{2x^2} + \frac{1}{2}(ian) \int \frac{(1 - iax)^{-n/2}(1 + iax)^{n/2}}{x^2} dx \\ &= -\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{2x^2} + \frac{2a^2n(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{1}{2}(-2+n)} {}_2F_1(2, 1 - \frac{n}{2}; 2 - \frac{n}{2}; \frac{1-i+iax}{1+iax})}{2-n} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 114, normalized size = 0.95

$$\frac{(1 - iax)^{-n/2}(1 + iax)^{n/2}(i + ax) \left(-((-2 + n)(-i + ax)^2) + 4a^2nx^2 {}_2F_1\left(2, 1 - \frac{n}{2}; 2 - \frac{n}{2}; \frac{i+ax}{i-ax}\right) \right)}{2(-2 + n)x^2(-i + ax)}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(I*n*ArcTan[a*x])/x^3,x]
```

```
[Out] ((1 + I*a*x)^(n/2)*(I + a*x)*(-((-2 + n)*(-I + a*x)^2) + 4*a^2*n*x^2*Hypergeometric2F1[2, 1 - n/2, 2 - n/2, (I + a*x)/(I - a*x)]))/(2*(-2 + n)*x^2*(1 - I*a*x)^(n/2)*(-I + a*x))
```

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \arctan(ax)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(I*n*arctan(a*x))/x^3,x)
```

```
[Out] int(exp(I*n*arctan(a*x))/x^3,x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^3,x, algorithm="maxima")

[Out] integrate(e^(I*n*arctan(a*x))/x^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^3,x, algorithm="fricas")

[Out] integral(1/(x^3*(-(a*x + I)/(a*x - I))^(1/2*n)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \operatorname{atan}(ax)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))/x**3,x)

[Out] Integral(exp(I*n*atan(a*x))/x**3, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^3,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax) 1i}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x)*1i)/x^3,x)

[Out] int(exp(n*atan(a*x)*1i)/x^3, x)

$$3.161 \quad \int \frac{e^{in \operatorname{ArcTan}(ax)}}{x^4} dx$$

Optimal. Leaf size=171

$$\frac{(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}}{3x^3} - \frac{ian(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{2+n}{2}}}{6x^2} + \frac{2ia^3(2+n^2)(1-iax)^{1-\frac{n}{2}}(1+iax)^{\frac{1}{2}(-2+n)} {}_2F_1\left(\begin{matrix} 2, 1-1/2*n \\ 2-1/2*n \end{matrix}, \frac{(1-iax)}{(1+iax)}\right)}{3(2-n)}$$

[Out] $-1/3*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(1+1/2*n)}/x^3-1/6*I*a*n*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(1+1/2*n)}/x^2+2/3*I*a^3*(n^2+2)*(1-I*a*x)^{(1-1/2*n)}*(1+I*a*x)^{(-1+1/2*n)}*\operatorname{hypergeom}\left([2, 1-1/2*n], [2-1/2*n], (1-I*a*x)/(1+I*a*x)\right)/(2-n)$

Rubi [A]

time = 0.05, antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 15, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5170, 105, 156, 12, 133}

$$\frac{2ia^3(n^2+2)(1+iax)^{\frac{n-2}{2}}(1-iax)^{1-\frac{n}{2}} {}_2F_1\left(2, 1-\frac{n}{2}; 2-\frac{n}{2}; \frac{1-iax}{1+iax}\right)}{3(2-n)} - \frac{(1+iax)^{\frac{n+2}{2}}(1-iax)^{1-\frac{n}{2}}}{3x^3} - \frac{ian(1+iax)^{\frac{n+2}{2}}(1-iax)^{1-\frac{n}{2}}}{6x^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*n*\operatorname{ArcTan}[a*x])}/x^4, x]$

[Out] $-1/3*((1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((2+n)/2)})/x^3 - ((I/6)*a*n*(1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((2+n)/2)})/x^2 + (((2*I)/3)*a^3*(2+n^2)*(1-I*a*x)^{(1-n/2)}*(1+I*a*x)^{((-2+n)/2)}*\operatorname{Hypergeometric2F1}[2, 1-n/2, 2-n/2, (1-I*a*x)/(1+I*a*x)])/(2-n)$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[u, (b_*)(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 105

$\operatorname{Int}[(a_*) + (b_*)(x_*)^{(m_*)}*((c_*) + (d_*)(x_*)^{(n_*)}*((e_*) + (f_*)(x_*)^{(p_*)}))^{(p_*)}, x_Symbol] := \operatorname{Simp}[b*(a + b*x)^{(m+1)}*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)})/((m+1)*(b*c - a*d)*(b*e - a*f)), x] + \operatorname{Dist}[1/((m+1)*(b*c - a*d)*(b*e - a*f)), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*(e + f*x)^p*\operatorname{Simp}[a*d*f*(m+1) - b*(d*e*(m+n+2) + c*f*(m+p+2)) - b*d*f*(m+n+p+3)*x, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \operatorname{ILtQ}[m, -1] \ \&\& \ (\operatorname{IntegerQ}[n] \ || \ \operatorname{IntegersQ}[2*n, 2*p] \ || \ \operatorname{ILtQ}[m+n+p+3, 0])$

Rule 133

$\operatorname{Int}[(a_*) + (b_*)(x_*)^{(m_*)}*((c_*) + (d_*)(x_*)^{(n_*)}*((e_*) + (f_*)(x_*)^{(p_*)}))^{(p_*)}, x_Symbol] := \operatorname{Simp}[(b*c - a*d)^n*(a + b*x)^{(m+1)}/((m+1)*(b*e -$

```

a*f)^(n + 1)*(e + f*x)^(m + 1))*Hypergeometric2F1[m + 1, -n, m + 2, (-d*
e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x))), x] /; FreeQ[{a, b, c, d, e,
f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1]
|| !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
)^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 5170

```

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.), x_Symbol] := Int[x^m*((1 - I*a
*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2))), x] /; FreeQ[{a, m, n}, x] && !Intege
rQ[(I*n - 1)/2]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{in \tan^{-1}(ax)}}{x^4} dx &= \int \frac{(1 - iax)^{-n/2}(1 + iax)^{n/2}}{x^4} dx \\
&= -\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{3x^3} - \frac{1}{3} \int \frac{(1 - iax)^{-n/2}(1 + iax)^{n/2}(-ian + a^2x)}{x^3} dx \\
&= -\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{3x^3} - \frac{ian(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{6x^2} - \frac{1}{6} \int \frac{a^2(2 + n^2)(1 - iax)}{x} dx \\
&= -\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{3x^3} - \frac{ian(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{6x^2} - \frac{1}{6}(a^2(2 + n^2)) \int \frac{(1 - iax)}{x} dx \\
&= -\frac{(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{3x^3} - \frac{ian(1 - iax)^{1-\frac{n}{2}}(1 + iax)^{\frac{2+n}{2}}}{6x^2} + \frac{2ia^3(2 + n^2)(1 - iax)^{1-\frac{n}{2}}}{6(-2 + n)x^3}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 119, normalized size = 0.70

$$-\frac{(1 - iax)^{-n/2}(1 + iax)^{\frac{1}{2}(-2+n)}(i + ax) \left(-((-2 + n)(-i + ax)^2(-2i + anx)) + 4a^3(2 + n^2)x^3 {}_2F_1\left(2, 1 - \frac{n}{2}; 2 - \frac{n}{2}; \frac{i+ax}{i-ax}\right) \right)}{6(-2 + n)x^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*n*ArcTan[a*x])/x^4,x]

[Out] $-1/6*((1 + I*a*x)^{((-2 + n)/2)}*(I + a*x)*(-((-2 + n)*(-I + a*x)^2*(-2*I + a*n*x)) + 4*a^3*(2 + n^2)*x^3*Hypergeometric2F1[2, 1 - n/2, 2 - n/2, (I + a*x)/(I - a*x)]))/((-2 + n)*x^3*(1 - I*a*x)^{(n/2)})$

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \arctan(ax)}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(I*n*arctan(a*x))/x^4,x)

[Out] int(exp(I*n*arctan(a*x))/x^4,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^4,x, algorithm="maxima")

[Out] integrate(e^(I*n*arctan(a*x))/x^4, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))/x^4,x, algorithm="fricas")

[Out] integral(1/(x^4*(-(a*x + I)/(a*x - I))^(1/2*n)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{in \operatorname{atan}(ax)}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))/x**4,x)

[Out] Integral(exp(I*n*atan(a*x))/x**4, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(I*n*arctan(a*x))/x^4,x, algorithm="giac")``[Out] sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax) 1i}}{x^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*atan(a*x)*1i)/x^4,x)``[Out] int(exp(n*atan(a*x)*1i)/x^4, x)`

3.162 $\int e^{i\text{ArcTan}(a+bx)} x^4 dx$

Optimal. Leaf size=276

$$\frac{(3i + 12a - 24ia^2 - 16a^3 + 8ia^4) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{8b^5} - \frac{(i + 8a)x^2 \sqrt{1 - ia - ibx} (1 + ia + ibx)^{3/2}}{20b^3}$$

[Out] $\frac{1}{8}*(3-12*I*a-24*a^2+16*I*a^3+8*a^4)*\text{arcsinh}(b*x+a)/b^5-1/20*(I+8*a)*x^2*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^3+1/5*x^3*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^2+1/120*(1+I*a+I*b*x)^{(3/2)}*(19*I+114*a-86*I*a^2-96*a^3-2*(13-14*I*a-36*a^2)*b*x)*(1-I*a-I*b*x)^{(1/2)}/b^5+1/8*(3*I+12*a-24*I*a^2-16*a^3+8*I*a^4)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^5$

Rubi [A]

time = 0.15, antiderivative size = 276, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 102, 158, 152, 52, 55, 633, 221}

$$\frac{\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}(-96a^3-2(-36a^2-14ia+13)bx-86ia^2+114a+19i)}{120b^5} + \frac{(8a^4-16a^3-24a^2+12a+3)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{8b^3} + \frac{(8a^4+16ia^2-24a^2-12ia+3)\text{sinh}^{-1}(a+bx)}{8b^3} - \frac{(8a+i)x^2\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{20b^3} + \frac{x^2\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{5b^2}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a + b*x])*x^4, x]

[Out] $((3*I + 12*a - (24*I)*a^2 - 16*a^3 + (8*I)*a^4)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/(8*b^5) - ((I + 8*a)*x^2*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)})/(20*b^3) + (x^3*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)})/(5*b^2) + (\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)}*(19*I + 114*a - (86*I)*a^2 - 96*a^3 - 2*(13 - (14*I)*a - 36*a^2)*b*x))/(120*b^5) + ((3 - (12*I)*a - 24*a^2 + (16*I)*a^3 + 8*a^4)*\text{ArcSinh}[a + b*x])/(8*b^5)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] :> Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]

Rule 102

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)), Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 158

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p + 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
```

`I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]`

Rubi steps

$$\begin{aligned}
 \int e^{i \tan^{-1}(a+bx)} x^4 dx &= \int \frac{x^4 \sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx \\
 &= \frac{x^3 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{5b^2} + \frac{\int \frac{x^2 \sqrt{1+ia+ibx} (-3(1+a^2)-(i+8a)bx)}{\sqrt{1-ia-ibx}} dx}{5b^2} \\
 &= -\frac{(i+8a)x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{20b^3} + \frac{x^3 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{5b^2} + \dots \\
 &= -\frac{(i+8a)x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{20b^3} + \frac{x^3 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{5b^2} + \dots \\
 &= \frac{(3i+12a-24ia^2-16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{(i+8a)x^2 \sqrt{1-ia-ibx}}{8b^5} \\
 &= \frac{(3i+12a-24ia^2-16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{(i+8a)x^2 \sqrt{1-ia-ibx}}{8b^5} \\
 &= \frac{(3i+12a-24ia^2-16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{(i+8a)x^2 \sqrt{1-ia-ibx}}{8b^5} \\
 &= \frac{(3i+12a-24ia^2-16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{(i+8a)x^2 \sqrt{1-ia-ibx}}{8b^5}
 \end{aligned}$$

Mathematica [A]

time = 0.18, size = 157, normalized size = 0.57

$$\frac{\sqrt{1+a^2+2abx+b^2x^2}(64i+275a-332ia^2-250a^3+24ia^4+(-45+116ia+130a^2-24ia^3)bx+2i(-16+35ia+12a^2)b^2x^2+6(5-4ia)b^3x^3+24ib^4x^4)+15(3-12ia-24a^2+16ia^3+8a^4)\sinh^{-1}(a+bx)}{120b^5}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*ArcTan[a + b*x])*x^4,x]`

`[Out] (Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]*(64*I + 275*a - (332*I)*a^2 - 250*a^3 + (24*I)*a^4 + (-45 + (116*I)*a + 130*a^2 - (24*I)*a^3)*b*x + (2*I)*(-16 + (35*I)*a + 12*a^2)*b^2*x^2 + 6*(5 - (4*I)*a)*b^3*x^3 + (24*I)*b^4*x^4) + 15*(3 - (12*I)*a - 24*a^2 + (16*I)*a^3 + 8*a^4)*ArcSinh[a + b*x]]/(120*b^5)`

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1259 vs. $2(225) = 450$.
time = 0.11, size = 1260, normalized size = 4.57

method	result
risch	$\frac{i(24x^4b^4 - 24ab^3x^3 - 30ib^3x^3 + 24a^2b^2x^2 + 70ia b^2x^2 - 24a^3bx - 130ix a^2b + 24a^4 + 250ia^3 - 32b^2x^2 + 116abx + 45ibx - 332a^2 - 275ia + 64)}{120b^5}$

default

$$ib \frac{x^4 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{5b^2}$$

$$9a \frac{x^3 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4b^2}$$

$$7a \frac{x^2 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{3b^2}$$

5a

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^4,x,method=_RETURNVERBOSE)
[Out] I*b*(1/5*x^4/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-9/5*a/b*(1/4*x^3/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-7/4*a/b*(1/3*x^2/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/3*a/b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-1/2*(a^2+1)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-2/3*(a^2+1)/b^2*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-3/4*(a^2+1)/b^2*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-1/2*(a^2+1)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-4/5*(a^2+1)/b^2*(1/3*x^2/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/3*a/b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-1/2*(a^2+1)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-2/3*(a^2+1)/b^2*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)))+(1+I*a)*(1/4*x^3/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-7/4*a/b*(1/3*x^2/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/3*a/b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-1/2*(a^2+1)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-2/3*(a^2+1)/b^2*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-3/4*(a^2+1)/b^2*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-1/2*(a^2+1)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)))
```

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 749 vs. $2(200) = 400$.

time = 0.27, size = 749, normalized size = 2.71

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^4,x, algorithm="maxima")
[Out] 1/5*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*x^4/b - 9/20*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*a*x^3/b^2 - 1/4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*a - 1)*x^3/b^2 + 21/20*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*a^2*x^2/b^3 - 7/12*sqrt
```

$$(b^2x^2 + 2abx + a^2 + 1) * (Ia + 1) * x^2 / b^3 - 63/8 * Ia^5 * \operatorname{arcsinh}(2 * (b^2x + a * b) / \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}) / b^5 + 35/8 * a^4 * (Ia + 1) * \operatorname{arcsinh}(2 * (b^2x + a * b) / \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}) / b^5 - 21/8 * I * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * a^3 * x / b^4 + 35/24 * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * a^2 * (Ia + 1) * x / b^4 - 4/15 * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * (Ia^2 + I) * x^2 / b^3 + 35/4 * I * (a^2 + 1) * a^3 * \operatorname{arcsinh}(2 * (b^2x + a * b) / \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}) / b^5 - 15/4 * (a^2 + 1) * a^2 * (Ia + 1) * \operatorname{arcsinh}(2 * (b^2x + a * b) / \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}) / b^5 + 63/8 * I * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * a^4 / b^5 - 35/8 * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * a^3 * (Ia + 1) / b^5 + 161/120 * I * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * (a^2 + 1) * a * x / b^4 - 3/8 * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * (a^2 + 1) * (Ia + 1) * x / b^4 - 15/8 * I * (a^2 + 1)^2 * a * \operatorname{arcsinh}(2 * (b^2x + a * b) / \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}) / b^5 - 3/8 * (a^2 + 1)^2 * (-Ia - 1) * \operatorname{arcsinh}(2 * (b^2x + a * b) / \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}) / b^5 - 49/8 * I * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * (a^2 + 1) * a^2 / b^5 + 55/24 * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * (a^2 + 1) * a * (Ia + 1) / b^5 + 8/15 * I * \sqrt{(b^2x^2 + 2abx + a^2 + 1)} * (a^2 + 1)^2 / b^5$$

Fricas [A]

time = 2.38, size = 177, normalized size = 0.64

$$\frac{186i a^5 - 1345 a^4 - 1730i a^3 + 1320 a^2 - 120(8a^4 + 16i a^3 - 24a^2 - 12i a + 3) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 8(-24i b^4 x^4 + 6(4i a - 5)b^3 x^3 + 2(-12i a^2 + 35a + 16i)b^2 x^2 - 24i a^4 + 250a^3 + (24i a^3 - 130a^2 - 116i a + 45)bx + 332i a - 275a - 64i) \sqrt{b^2x^2 + 2abx + a^2 + 1} + 300i a}{960 b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^4,x, algorithm="fricas")

[Out] 1/960*(186*I*a^5 - 1345*a^4 - 1730*I*a^3 + 1320*a^2 - 120*(8*a^4 + 16*I*a^3 - 24*a^2 - 12*I*a + 3)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 8*(-24*I*b^4*x^4 + 6*(4*I*a - 5)*b^3*x^3 + 2*(-12*I*a^2 + 35*a + 16*I)*b^2*x^2 - 24*I*a^4 + 250*a^3 + (24*I*a^3 - 130*a^2 - 116*I*a + 45)*b*x + 332*I*a^2 - 275*a - 64*I)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 300*I*a)/b^5

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \left(-\frac{ix^4}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} \right) dx + \int \frac{ax^4}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx + \int \frac{bx^5}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)*x**4,x)

[Out] I*(Integral(-I*x**4/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(a*x**4/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(b*x**5/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x))

Giac [A]

time = 0.47, size = 205, normalized size = 0.74

$$-\frac{1}{120} \sqrt{(bx+a)^2+1} \left(2 \left(3x \left(-\frac{4ix}{b} - \frac{4iab^7+5b^7}{b^8} \right) - \frac{12ia^2b^6-35ab^6-16ib^6}{b^8} x - \frac{-24ia^3b^5+130a^2b^5+116iab^5-45b^5}{b^8} x - \frac{24ia^4b^4-250a^3b^4-332ia^2b^4+275ab^4+64ib^4}{b^8} \right) - \frac{(8a^4+16ia^3-24a^2-12ia+3) \log\left(-ab - \left(x|b| - \sqrt{(bx+a)^2+1}\right)|b|\right)}{8b^4|b|} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^4,x, algorithm="giac")

[Out]
$$-1/120*\sqrt{(b*x + a)^2 + 1}*((2*(3*x*(-4*I*x/b - (-4*I*a*b^7 + 5*b^7)/b^9) - (12*I*a^2*b^6 - 35*a*b^6 - 16*I*b^6)/b^9)*x - (-24*I*a^3*b^5 + 130*a^2*b^5 + 116*I*a*b^5 - 45*b^5)/b^9)*x - (24*I*a^4*b^4 - 250*a^3*b^4 - 332*I*a^2*b^4 + 275*a*b^4 + 64*I*b^4)/b^9) - 1/8*(8*a^4 + 16*I*a^3 - 24*a^2 - 12*I*a + 3)*\log(-a*b - (x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*\text{abs}(b))/(b^4*\text{abs}(b))$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^4 (1 + a \operatorname{li} + b x \operatorname{li})}{\sqrt{(a + b x)^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*(a*li + b*x*li + 1))/((a + b*x)^2 + 1)^(1/2),x)

[Out] int((x^4*(a*li + b*x*li + 1))/((a + b*x)^2 + 1)^(1/2), x)

3.163 $\int e^{i\text{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=201

$$\frac{(3 - 12ia - 12a^2 + 8ia^3) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{8b^4} + \frac{x^2 \sqrt{1 - ia - ibx} (1 + ia + ibx)^{3/2}}{4b^2} - \frac{\sqrt{1 - ia - i}}$$

[Out] 1/8*(3*I+12*a-12*I*a^2-8*a^3)*arcsinh(b*x+a)/b^4+1/4*x^2*(1+I*a+I*b*x)^(3/2)*(1-I*a-I*b*x)^(1/2)/b^2-1/24*(1+I*a+I*b*x)^(3/2)*(7-10*I*a-18*a^2+2*(I+6*a)*b*x)*(1-I*a-I*b*x)^(1/2)/b^4-1/8*(3-12*I*a-12*a^2+8*I*a^3)*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/b^4

Rubi [A]

time = 0.14, antiderivative size = 201, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 102, 152, 52, 55, 633, 221}

$$\frac{\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}(-18a^2+2(6a+i)bx-10ia+7)}{24b^4} - \frac{(8ia^3-12a^2-12ia+3)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{8b^4} + \frac{(-8a^3-12ia^2+12a+3i)\sinh^{-1}(a+bx)}{8b^4} + \frac{x^2\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{4b^2}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a + b*x])*x^3,x]

[Out] -1/8*((3 - (12*I)*a - 12*a^2 + (8*I)*a^3)*Sqrt[1 - I*a - I*b*x]*Sqrt[1 + I*a + I*b*x])/b^4 + (x^2*Sqrt[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^(3/2))/(4*b^2) - (Sqrt[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^(3/2)*(7 - (10*I)*a - 18*a^2 + 2*(I + 6*a)*b*x))/(24*b^4) + ((3*I + 12*a - (12*I)*a^2 - 8*a^3)*ArcSinh[a + b*x])/(8*b^4)

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(
b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ
[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 55

```
Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[
1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b
+ d, 0] && GtQ[a + c, 0]
```

Rule 102

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x
```

```
)^(p + 1)/(d*f*(m + n + p + 1)), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a
+ b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b
*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))] + b*(a*d*f*(2*m + n + p) - b*
(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p
}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_
))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{i \tan^{-1}(a+bx)} x^3 dx &= \int \frac{x^3 \sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx \\
&= \frac{x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} + \frac{\int \frac{x \sqrt{1+ia+ibx} (-2(1+a^2)-(i+6a)bx)}{\sqrt{1-ia-ibx}} dx}{4b^2} \\
&= \frac{x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} - \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2} (7-10ia-18a^2)}{24b^4} \\
&= -\frac{(3-12ia-12a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} \\
&= -\frac{(3-12ia-12a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} \\
&= -\frac{(3-12ia-12a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} \\
&= -\frac{(3-12ia-12a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2}
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 121, normalized size = 0.60

$$\frac{-\sqrt{1+a^2+2abx+b^2x^2}(16-39ia-44a^2+6ia^3+(9i+20a-6ia^2)bx+2i(4i+3a)b^2x^2-6ib^3x^3)+3(3i+12a-12ia^2-8a^3)\sinh^{-1}(a+bx)}{24b^4}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a + b*x])*x^3,x]

[Out] $(-\text{Sqrt}[1+a^2+2*a*b*x+b^2*x^2]*(16-(39*I)*a-44*a^2+(6*I)*a^3+(9*I+20*a-(6*I)*a^2)*b*x+(2*I)*(4*I+3*a)*b^2*x^2-(6*I)*b^3*x^3))+3*(3*I+12*a-(12*I)*a^2-8*a^3)*\text{ArcSinh}[a+b*x])/(24*b^4)$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 748 vs. $2(163) = 326$.

time = 0.11, size = 749, normalized size = 3.73

method	result
risch	$ -\frac{i(-6b^3x^3+6ab^2x^2+8ib^2x^2-6a^2bx-20iabx+6a^3+44ia^2+9bx-39a-16i)\sqrt{b^2x^2+2abx+a^2+1}}{24b^4} - \frac{3i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}}+\sqrt{b^2}\right)}{24b^4} $

default	ib	$\frac{x^3 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4b^2}$	$7a$	$\frac{x^2 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{3b^2}$	$5a$	$\frac{x \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{2b^2}$	$3a$
---------	------	--	------	--	------	--	------

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^3,x,method=_RETURNVERBOSE)`

[Out] $I*b*(1/4*x^3/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-7/4*a/b*(1/3*x^2/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-5/3*a/b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-1/2*(a^2+1)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-2/3*(a^2+1)/b^2*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}))-3/4*(a^2+1)/b^2*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-1/2*(a^2+1)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}))+1+I*a)*(1/3*x^2/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-5/3*a/b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-1/2*(a^2+1)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}))+1+I*a)$

$$(b^2x+ab)/(b^2)^{(1/2)}+(b^2x^2+2abx+a^2+1)^{(1/2)}/(b^2)^{(1/2)}-1/2*(a^2+1)/b^2*\ln((b^2x+ab)/(b^2)^{(1/2)}+(b^2x^2+2abx+a^2+1)^{(1/2)}/(b^2)^{(1/2)})-2/3*(a^2+1)/b^2*(1/b^2*(b^2x^2+2abx+a^2+1)^{(1/2)}-a/b*\ln((b^2x+ab)/(b^2)^{(1/2)}+(b^2x^2+2abx+a^2+1)^{(1/2)}/(b^2)^{(1/2)}))$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 529 vs. $2(145) = 290$.
time = 0.28, size = 529, normalized size = 2.63

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^3,x, algorithm="maxima")

[Out] $1/4*I*\sqrt{b^2x^2 + 2abx + a^2 + 1}*x^3/b - 7/12*I*\sqrt{b^2x^2 + 2abx + a^2 + 1}*x^2/b^2 - 1/3*\sqrt{b^2x^2 + 2abx + a^2 + 1}*(-Ia - 1)*x^2/b^2 + 35/8*I*a^4*\operatorname{arcsinh}(2*(b^2x + ab)/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2})/b^4 - 5/2*a^3*(Ia + 1)*\operatorname{arcsinh}(2*(b^2x + ab)/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2})/b^4 + 35/24*I*\sqrt{b^2x^2 + 2abx + a^2 + 1}*a^2*x/b^3 - 5/6*\sqrt{b^2x^2 + 2abx + a^2 + 1}*a*(Ia + 1)*x/b^3 - 15/4*I*(a^2 + 1)*a^2*\operatorname{arcsinh}(2*(b^2x + ab)/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2})/b^4 + 3/2*(a^2 + 1)*a*(Ia + 1)*\operatorname{arcsinh}(2*(b^2x + ab)/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2})/b^4 - 35/8*I*\sqrt{b^2x^2 + 2abx + a^2 + 1}*a^3/b^4 + 5/2*\sqrt{b^2x^2 + 2abx + a^2 + 1}*a^2*(Ia + 1)/b^4 - 3/8*\sqrt{b^2x^2 + 2abx + a^2 + 1}*(Ia^2 + I)*x/b^3 + 3/8*I*(a^2 + 1)^2*\operatorname{arcsinh}(2*(b^2x + ab)/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2})/b^4 + 55/24*I*\sqrt{b^2x^2 + 2abx + a^2 + 1}*(a^2 + 1)*a/b^4 - 2/3*\sqrt{b^2x^2 + 2abx + a^2 + 1}*(a^2 + 1)*(Ia + 1)/b^4$

Fricas [A]

time = 2.36, size = 139, normalized size = 0.69

$$\frac{-45i a^4 + 224 a^3 + 192i a^2 + 24(8 a^3 + 12i a^2 - 12a - 3i) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 8(-6i b^3x^3 + 2(3i a - 4)b^2x^2 + 6i a^3 + (-6i a^2 + 20a + 9i)bx - 44a^2 - 39i a + 16)\sqrt{b^2x^2 + 2abx + a^2 + 1} - 72a}{192b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^3,x, algorithm="fricas")

[Out] $1/192*(-45*I*a^4 + 224*a^3 + 192*I*a^2 + 24*(8*a^3 + 12*I*a^2 - 12*a - 3*I)*\log(-b*x - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 8*(-6*I*b^3*x^3 + 2*(3*I*a - 4)*b^2*x^2 + 6*I*a^3 + (-6*I*a^2 + 20*a + 9*I)*b*x - 44*a^2 - 39*I*a + 16)*\sqrt{b^2x^2 + 2abx + a^2 + 1} - 72*a)/b^4$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i\left(\int\left(-\frac{ix^3}{\sqrt{a^2+2abx+b^2x^2+1}}\right)dx + \int\frac{ax^3}{\sqrt{a^2+2abx+b^2x^2+1}}dx + \int\frac{bx^4}{\sqrt{a^2+2abx+b^2x^2+1}}dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)*x**3,x)

[Out] I*(Integral(-I*x**3/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(a*x**3/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(b*x**4/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x))

Giac [A]

time = 0.45, size = 155, normalized size = 0.77

$$-\frac{1}{24} \sqrt{(bx+a)^2+1} \left(\left(2x \left(-\frac{3ix}{b} - \frac{-3iab^5+4b^5}{b^7} \right) - \frac{6ia^2b^4-20ab^4-9ib^4}{b^7} \right) x - \frac{-6ia^3b^3+44a^2b^3+39iab^3-16b^3}{b^7} \right) + \frac{(8a^3+12ia^2-12a-3i) \log\left(-ab - \left(x|b| - \sqrt{(bx+a)^2+1}\right)|b|\right)}{8b^3|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^3,x, algorithm="giac")

[Out] -1/24*sqrt((b*x + a)^2 + 1)*((2*x*(-3*I*x/b - (-3*I*a*b^5 + 4*b^5)/b^7) - (6*I*a^2*b^4 - 20*a*b^4 - 9*I*b^4)/b^7)*x - (-6*I*a^3*b^3 + 44*a^2*b^3 + 39*I*a*b^3 - 16*b^3)/b^7) + 1/8*(8*a^3 + 12*I*a^2 - 12*a - 3*I)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^3*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3 (1 + a \operatorname{li} + b x \operatorname{li})}{\sqrt{(a + b x)^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a*1i + b*x*1i + 1))/((a + b*x)^2 + 1)^(1/2),x)

[Out] int((x^3*(a*1i + b*x*1i + 1))/((a + b*x)^2 + 1)^(1/2), x)

3.164 $\int e^{i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=171

$$\frac{(i+2a-2ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} - \frac{(i+4a)\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}}{6b^3} + \frac{x\sqrt{1-ia-ibx}}{3}$$

[Out] $-1/2*(1-2*I*a-2*a^2)*\text{arcsinh}(b*x+a)/b^3-1/6*(I+4*a)*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^3+1/3*x*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^2-1/2*(I+2*a-2*I*a^2)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3$

Rubi [A]

time = 0.09, antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 92, 81, 52, 55, 633, 221}

$$\frac{(-2ia^2+2a+i)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{2b^3} - \frac{(-2a^2-2ia+1)\sinh^{-1}(a+bx)}{2b^3} - \frac{(4a+i)\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{6b^3} + \frac{x\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{3b^2}$$

Antiderivative was successfully verified.

[In] `Int[E^(I*ArcTan[a + b*x])*x^2,x]`

[Out] $-1/2*((I+2*a-(2*I)*a^2)*\text{Sqrt}[1-I*a-I*b*x]*\text{Sqrt}[1+I*a+I*b*x])/b^3 - ((I+4*a)*\text{Sqrt}[1-I*a-I*b*x]*(1+I*a+I*b*x)^{(3/2)})/(6*b^3) + (x*\text{Sqrt}[1-I*a-I*b*x]*(1+I*a+I*b*x)^{(3/2)})/(3*b^2) - ((1-(2*I)*a-2*a^2)*\text{ArcSinh}[a+b*x])/(2*b^3)$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 55

`Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`

Rule 81

`Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f}`

, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92

Int[((a_.) + (b_.)*(x_))²((c_.) + (d_.)*(x_))^(n_.)((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)ⁿ(e + f*x)^p*Simp[a²*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)²], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)²)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b² - 4*a*c)))^p), Subst[Int[Simp[1 - x²/(b² - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b²/c, 0]

Rule 5203

Int[E^{(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)}((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m((1 - I*a*c - I*b*c*x)^{(I*(n/2))}/(1 + I*a*c + I*b*c*x)^{(I*(n/2))}), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2 \sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx \\
&= \frac{x \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{3b^2} + \frac{\int \frac{\sqrt{1+ia+ibx} (-1-a^2-(i+4a)bx)}{\sqrt{1-ia-ibx}} dx}{3b^2} \\
&= -\frac{(i+4a) \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{6b^3} + \frac{x \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{3b^2} - \frac{(1-ia-ibx)^{3/2}}{3b^2} \\
&= -\frac{(i+2a-2ia^2) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} - \frac{(i+4a) \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{6b^3} \\
&= -\frac{(i+2a-2ia^2) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} - \frac{(i+4a) \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{6b^3} \\
&= -\frac{(i+2a-2ia^2) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} - \frac{(i+4a) \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{6b^3} \\
&= -\frac{(i+2a-2ia^2) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} - \frac{(i+4a) \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{6b^3} \\
&= -\frac{(i+2a-2ia^2) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} - \frac{(i+4a) \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{6b^3}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 135, normalized size = 0.79

$$\frac{\sqrt{1+a^2+2abx+b^2x^2}(-4i+2ia^2+3bx+2ib^2x^2+a(-9-2ibx))}{6b^3} + \frac{\sqrt[4]{-1}(-1+2ia+2a^2)\sqrt{-ib}\sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{b^{7/2}}$$

Warning: Unable to verify antiderivative.

`[In] Integrate[E^(I*ArcTan[a + b*x])*x^2,x]`

```
[Out] (Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]*(-4*I + (2*I)*a^2 + 3*b*x + (2*I)*b^2*x^2 + a*(-9 - (2*I)*b*x)))/(6*b^3) + ((-1)^(1/4)*(-1 + (2*I)*a + 2*a^2)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/b^(7/2)
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 435 vs. $2(135) = 270$.

time = 0.09, size = 436, normalized size = 2.55

method	result
--------	--------

risch	$\frac{i(2b^2x^2 - 2abx - 3ibx + 2a^2 + 9ia - 4)\sqrt{b^2x^2 + 2abx + a^2 + 1}}{6b^3} + \frac{i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)^a}{b^2\sqrt{b^2}} + \frac{\ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)^a}{b^2\sqrt{b^2}}$
default	$ib \left(\frac{x^2\sqrt{b^2x^2 + 2abx + a^2 + 1}}{3b^2} - \frac{5a \left(\frac{x\sqrt{b^2x^2 + 2abx + a^2 + 1}}{2b^2} - \frac{3a \left(\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b^2} - \frac{a \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{b^2\sqrt{b^2}} \right)}{b^2} \right)}{b^2} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^2,x,method=_RETURNVERBOSE)`

[Out] $I*b*(1/3*x^2/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-5/3*a/b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-1/2*(a^2+1)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}))-2/3*(a^2+1)/b^2*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)})))+(1+I*a)*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}))-1/2*(a^2+1)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)})/(b^2)^{(1/2)})$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 351 vs. $2(119) = 238$.
time = 0.26, size = 351, normalized size = 2.05

$$\frac{\sqrt{b^2x^2+2abx+a^2+1}}{3b} - \frac{5a^2 \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^3} - \frac{3a^2(n+1) \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^3} - \frac{5i \sqrt{b^2x^2+2abx+a^2+1} \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{6b^3} - \frac{\sqrt{b^2x^2+2abx+a^2+1} \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^3} - \frac{3i(n+1) \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^3} - \frac{(n^2+1)(n+1) \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^3} - \frac{5i \sqrt{b^2x^2+2abx+a^2+1}}{2b^3} - \frac{3 \sqrt{b^2x^2+2abx+a^2+1} \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^3} - \frac{2 \sqrt{b^2x^2+2abx+a^2+1} \operatorname{arcsinh}\left(\frac{2b^2x+a}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{3b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^2,x, algorithm="maxima")`

[Out] $1/3*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*x^2/b - 5/2*I*a^3*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^3 + 3/2*a^2*(I*a + 1)*\operatorname{arcsinh}($

$$2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^3 - 5/6*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1)*a*x/b^2 - 1/2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*a - 1)*x/b^2 + 3/2*I*(a^2 + 1)*a*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^3 - 1/2*(a^2 + 1)*(I*a + 1)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^3 + 5/2*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1)*a^2/b^3 - 3/2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1)*a*(I*a + 1)/b^3 - 2/3*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1)*(I*a^2 + I)/b^3}$$

Fricas [A]

time = 1.37, size = 106, normalized size = 0.62

$$\frac{7ia^3 - 21a^2 - 12(2a^2 + 2ia - 1)\log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 4\sqrt{b^2x^2 + 2abx + a^2 + 1}(-2ib^2x^2 + (2ia - 3)bx - 2ia^2 + 9a + 4i) - 9ia}{24b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^2,x, algorithm="fricas")

[Out] 1/24*(7*I*a^3 - 21*a^2 - 12*(2*a^2 + 2*I*a - 1)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-2*I*b^2*x^2 + (2*I*a - 3)*b*x - 2*I*a^2 + 9*a + 4*I) - 9*I*a)/b^3

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i\left(\int\left(-\frac{ix^2}{\sqrt{a^2 + 2abx + b^2x^2 + 1}}\right)dx + \int\frac{ax^2}{\sqrt{a^2 + 2abx + b^2x^2 + 1}}dx + \int\frac{bx^3}{\sqrt{a^2 + 2abx + b^2x^2 + 1}}dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)*x**2,x)

[Out] I*(Integral(-I*x**2/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(a*x**2/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(b*x**3/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x))

Giac [A]

time = 0.44, size = 113, normalized size = 0.66

$$-\frac{1}{6}\sqrt{(bx+a)^2+1}\left(x\left(-\frac{2ix}{b}-\frac{-2iab^3+3b^3}{b^5}\right)-\frac{2ia^2b^2-9ab^2-4ib^2}{b^5}\right)-\frac{(2a^2+2ia-1)\log\left(-ab-\left(x|b|-\sqrt{(bx+a)^2+1}\right)|b|\right)}{2b^2|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x^2,x, algorithm="giac")

[Out] -1/6*sqrt((b*x + a)^2 + 1)*(x*(-2*I*x/b - (-2*I*a*b^3 + 3*b^3)/b^5) - (2*I*a^2*b^2 - 9*a*b^2 - 4*I*b^2)/b^5) - 1/2*(2*a^2 + 2*I*a - 1)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^2*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^2 (1 + a + b x)}{\sqrt{(a + b x)^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a+bx+1))/((a+bx)^2+1)^(1/2),x)

[Out] int((x^2*(a+bx+1))/((a+bx)^2+1)^(1/2), x)

3.165 $\int e^{i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=110

$$\frac{(1-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} + \frac{\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}}{2b^2} - \frac{(i+2a)\sinh^{-1}(a+bx)}{2b^2}$$

[Out] $-1/2*(I+2*a)*\text{arcsinh}(b*x+a)/b^2+1/2*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^2+1/2*(1-2*I*a)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2$

Rubi [A]

time = 0.05, antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5203, 81, 52, 55, 633, 221}

$$\frac{\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{2b^2} + \frac{(1-2ia)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{2b^2} - \frac{(2a+i)\sinh^{-1}(a+bx)}{2b^2}$$

Antiderivative was successfully verified.

[In] `Int[E^(I*ArcTan[a + b*x])*x,x]`

[Out] `((1 - (2*I)*a)*Sqrt[1 - I*a - I*b*x]*Sqrt[1 + I*a + I*b*x])/(2*b^2) + (Sqrt[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^(3/2))/(2*b^2) - ((I + 2*a)*ArcSinh[a + b*x])/(2*b^2)`

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 55

`Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] :> Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`

Rule 81

`Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]`

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{i \tan^{-1}(a+bx)} x \, dx &= \int \frac{x \sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} \, dx \\
 &= \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2b^2} - \frac{(i+2a) \int \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} \, dx}{2b} \\
 &= \frac{(1-2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} + \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2b^2} - \frac{(i+2a)}{2b} \\
 &= \frac{(1-2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} + \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2b^2} - \frac{(i+2a)}{2b} \\
 &= \frac{(1-2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} + \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2b^2} - \frac{(i+2a)}{2b} \\
 &= \frac{(1-2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} + \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2b^2} - \frac{(i+2a)}{2b}
 \end{aligned}$$

Mathematica [A]

time = 0.09, size = 108, normalized size = 0.98

$$\frac{(2-ia+ibx) \sqrt{1+a^2+2abx+b^2x^2}}{2b^2} + \frac{(-1)^{3/4} (i+2a) \sinh^{-1} \left(\frac{(\frac{1}{2}+\frac{i}{2}) \sqrt{b} \sqrt{-i(i+ax)}}{\sqrt{-ib}} \right)}{\sqrt{-ib} b^{3/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a + b*x])*x,x]

[Out] $((2 - I*a + I*b*x)*\text{Sqrt}[1 + a^2 + 2*a*b*x + b^2*x^2])/(2*b^2) + ((-1)^{(3/4)} * (I + 2*a)*\text{ArcSinh}[(1/2 + I/2)*\text{Sqrt}[b]*\text{Sqrt}[(-I)*(I + a + b*x)])/\text{Sqrt}[(-I)*b]) / (\text{Sqrt}[(-I)*b]*b^{(3/2)})$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 237 vs. 2(86) = 172.

time = 0.11, size = 238, normalized size = 2.16

method	result
risch	$\frac{i(-bx+a+2i)\sqrt{b^2x^2+2abx+a^2+1}}{2b^2} - \frac{i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{2b\sqrt{b^2}} - \frac{a \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{b\sqrt{b^2}}$
default	$ib \left(\frac{x\sqrt{b^2x^2+2abx+a^2+1}}{2b^2} - \frac{3a \left(\frac{\sqrt{b^2x^2+2abx+a^2+1}}{b^2} - \frac{a \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{b\sqrt{b^2}} \right)}{2b} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x,x,method=_RETURNVERBOSE)

[Out] $I*b*(1/2*x/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} - 3/2*a/b*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} - a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)}) - 1/2*(a^2+1)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)}) + (1+I*a)*(1/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} - a/b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 209 vs. 2(76) = 152.

time = 0.26, size = 209, normalized size = 1.90

$$\frac{3i a^2 \operatorname{arsinh}\left(\frac{2(b^2x+ab)}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^2} - \frac{a(i a + 1) \operatorname{arsinh}\left(\frac{2(b^2x+ab)}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{b^2} + \frac{i \sqrt{b^2x^2+2abx+a^2+1} x}{2b} - \frac{(i a^2 + i) \operatorname{arsinh}\left(\frac{2(b^2x+ab)}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{2b^2} - \frac{3i \sqrt{b^2x^2+2abx+a^2+1} a}{2b^2} + \frac{\sqrt{b^2x^2+2abx+a^2+1} (i a + 1)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x,x, algorithm="maxima")

[Out] $3/2*I*a^2*\operatorname{arcsinh}(2*(b^2*x + a*b)/\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^2 - a*(I*a + 1)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^2 + 1/2*I*\operatorname{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*x/b - 1/2*(I*a^2 + I)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^2$

$$\frac{(b^2x + ab) \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2}}{b^2} - \frac{3}{2} I \sqrt{b^2x^2 + 2abx + a^2 + 1} + \frac{2abx + a^2 + 1}{b^2} + \frac{\sqrt{b^2x^2 + 2abx + a^2 + 1} (Ia + 1)}{b^2}$$

Fricas [A]

time = 2.60, size = 79, normalized size = 0.72

$$\frac{-3ia^2 + 4(2a + i) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 4\sqrt{b^2x^2 + 2abx + a^2 + 1}(-ibx + ia - 2) + 4a}{8b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x,x, algorithm="fricas")

[Out] 1/8*(-3*I*a^2 + 4*(2*a + I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*b*x + I*a - 2) + 4*a)/b^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \left(-\frac{ix}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} \right) dx + \int \frac{ax}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx + \int \frac{bx^2}{\sqrt{a^2 + 2abx + b^2x^2 + 1}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)*x,x)

[Out] I*(Integral(-I*x/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(a*x/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x) + Integral(b*x**2/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1), x))

Giac [A]

time = 0.46, size = 75, normalized size = 0.68

$$-\frac{1}{2} \sqrt{(bx + a)^2 + 1} \left(-\frac{ix}{b} + \frac{iab - 2b}{b^3} \right) + \frac{(2a + i) \log \left(-ab - \left(x|b| - \sqrt{(bx + a)^2 + 1} \right) |b| \right)}{2b|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)*x,x, algorithm="giac")

[Out] -1/2*sqrt((b*x + a)^2 + 1)*(-I*x/b + (I*a*b - 2*b)/b^3) + 1/2*(2*a + I)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x(1 + a \operatorname{li} + b x \operatorname{li})}{\sqrt{(a + b x)^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*(a*li + b*x*li + 1))/((a + b*x)^2 + 1)^(1/2),x)

[Out] int((x*(a*li + b*x*li + 1))/((a + b*x)^2 + 1)^(1/2), x)

3.166 $\int e^{i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=52

$$\frac{i\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{b} + \frac{\sinh^{-1}(a+bx)}{b}$$

[Out] arcsinh(b*x+a)/b+I*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/b

Rubi [A]

time = 0.03, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5201, 52, 55, 633, 221}

$$\frac{\sinh^{-1}(a+bx)}{b} + \frac{i\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{b}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a + b*x]),x]

[Out] (I*Sqrt[1 - I*a - I*b*x]*Sqrt[1 + I*a + I*b*x])/b + ArcSinh[a + b*x]/b

Rule 52

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*(b*c - a*d)/(
b*(m + n + 1)), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b,
c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[
m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n
+ 2, 0] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 55

```
Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[
1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b
+ d, 0] && GtQ[a + c, 0]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
```

+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5201

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.)), x_Symbol] :> Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{i \tan^{-1}(a+bx)} dx &= \int \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx \\
 &= \frac{i\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{b} + \int \frac{1}{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}} dx \\
 &= \frac{i\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{b} + \int \frac{1}{\sqrt{(1-ia)(1+ia)+2abx+b^2x^2}} dx \\
 &= \frac{i\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{b} + \frac{\text{Subst} \left(\int \frac{1}{\sqrt{1+\frac{x^2}{4b^2}}} dx, x, 2ab+2b^2x \right)}{2b^2} \\
 &= \frac{i\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{b} + \frac{\sinh^{-1}(a+bx)}{b}
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 28, normalized size = 0.54

$$\frac{i\sqrt{1+(a+bx)^2} + \sinh^{-1}(a+bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a + b*x]),x]

[Out] (I*Sqrt[1 + (a + b*x)^2] + ArcSinh[a + b*x])/b

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 163 vs. 2(43) = 86.

time = 0.08, size = 164, normalized size = 3.15

method	result
--------	--------

risch	$\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b} + \frac{\ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{\sqrt{b^2}}$
default	$\frac{\ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{\sqrt{b^2}} + \frac{ia \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{\sqrt{b^2}} + ib \left(\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b^2} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\ln((b^2x+a*b)/(b^2)^{(1/2)}+(b^2x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)}+I*a*\ln((b^2x+a*b)/(b^2)^{(1/2)}+(b^2x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)}+I*b*(1/b^2*(b^2x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*\ln((b^2x+a*b)/(b^2)^{(1/2)}+(b^2x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})$

Maxima [A]

time = 0.29, size = 62, normalized size = 1.19

$$\frac{\operatorname{arsinh}\left(\frac{2(b^2x+ab)}{\sqrt{-4a^2b^2+4(a^2+1)b^2}}\right)}{b} + \frac{i\sqrt{b^2x^2+2abx+a^2+1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2),x, algorithm="maxima")`

[Out] $\operatorname{arcsinh}(2*(b^2x+a*b)/\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})/b+I*\sqrt{b^2x^2+2*a*b*x+a^2+1}/b$

Fricas [A]

time = 1.54, size = 60, normalized size = 1.15

$$\frac{ia + 2i\sqrt{b^2x^2 + 2abx + a^2 + 1} - 2 \log\left(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2),x, algorithm="fricas")`

[Out] $1/2*(I*a + 2*I*\sqrt{b^2x^2 + 2*a*b*x + a^2 + 1} - 2*\log(-b*x - a + \sqrt{b^2x^2 + 2*a*b*x + a^2 + 1}))/b$

Sympy [A]

time = 1.76, size = 36, normalized size = 0.69

$$\begin{cases} \frac{i\sqrt{(a+bx)^2+1} + \operatorname{asinh}(a+bx)}{b} & \text{for } b \neq 0 \\ \frac{x(ia+1)}{\sqrt{a^2+1}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2),x)

[Out] Piecewise(((I*sqrt((a + b*x)**2 + 1) + asinh(a + b*x))/b, Ne(b, 0)), (x*(I*a + 1)/sqrt(a**2 + 1), True))

Giac [A]

time = 0.44, size = 51, normalized size = 0.98

$$-\frac{\log\left(-ab - \left(x|b| - \sqrt{(bx+a)^2 + 1}\right)|b|\right)}{|b|} + \frac{i\sqrt{(bx+a)^2 + 1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2),x, algorithm="giac")

[Out] -log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/abs(b) + I*sqrt((b*x + a)^2 + 1)/b

Mupad [B]

time = 1.09, size = 97, normalized size = 1.87

$$\frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{b} \operatorname{li} + \frac{\operatorname{asinh}(a + bx)}{b} + \frac{a \operatorname{asinh}(a + bx)}{b} \operatorname{li} - \frac{ab^2 \ln\left(\sqrt{a^2 + 2abx + b^2x^2 + 1} + \frac{xb^2 + ab}{\sqrt{b^2}}\right)}{(b^2)^{3/2}} \operatorname{li}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2),x)

[Out] ((a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2)*1i)/b + asinh(a + b*x)/b + (a*asinh(a + b*x)*1i)/b - (a*b^2*log((a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2) + (a*b + b^2*x)/(b^2)^(1/2))*1i)/(b^2)^(3/2)

$$3.167 \quad \int \frac{e^{i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=89

$$i \sinh^{-1}(a+bx) - \frac{2\sqrt{i-a} \tanh^{-1}\left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}}\right)}{\sqrt{i+a}}$$

[Out] I*arcsinh(b*x+a)-2*arctanh((I+a)^(1/2)*(1+I*a+I*b*x)^(1/2)/(I-a)^(1/2)/(1-I*a-I*b*x)^(1/2))*(I-a)^(1/2)/(I+a)^(1/2)

Rubi [A]

time = 0.07, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 132, 55, 633, 221, 12, 95, 214}

$$i \sinh^{-1}(a+bx) - \frac{2\sqrt{-a+i} \tanh^{-1}\left(\frac{\sqrt{a+i} \sqrt{ia+ibx+1}}{\sqrt{-a+i} \sqrt{-ia-ibx+1}}\right)}{\sqrt{a+i}}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a + b*x])/x,x]

[Out] I*ArcSinh[a + b*x] - (2*Sqrt[I - a]*ArcTanh[(Sqrt[I + a]*Sqrt[1 + I*a + I*b*x])/(Sqrt[I - a]*Sqrt[1 - I*a - I*b*x])])/Sqrt[I + a]

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 55

Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 132

```

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))

```

Rule 214

```

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

```

Rule 221

```

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

```

Rule 633

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

```

Rule 5203

```

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{\sqrt{1+ia+ibx}}{x\sqrt{1-ia-ibx}} dx \\
&= -\left((-1-ia) \int \frac{1}{x\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx \right) + (ib) \int \frac{1}{\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx \\
&= (2(1+ia)) \text{Subst} \left(\int \frac{1}{-1-ia - (-1+ia)x^2} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} \right) + (ib) \int \frac{1}{\sqrt{(1-ia-ibx)(1+ia+ibx)}} dx \\
&= -\frac{2\sqrt{i-a} \tanh^{-1} \left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}} \right)}{\sqrt{i+a}} + \frac{i \text{Subst} \left(\int \frac{1}{\sqrt{1+\frac{x^2}{4b^2}}} dx, x, 2ab+2b^2 \right)}{2b} \\
&= i \sinh^{-1}(a+bx) - \frac{2\sqrt{i-a} \tanh^{-1} \left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}} \right)}{\sqrt{i+a}}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 142, normalized size = 1.60

$$\frac{2(-1)^{3/4}\sqrt{-ib} \sinh^{-1} \left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b} \sqrt{-i(i+a+bx)}}{\sqrt{-ib}} \right)}{\sqrt{b}} - \frac{2\sqrt{-1-ia} \tanh^{-1} \left(\frac{\sqrt{-1-ia} \sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}} \right)}{\sqrt{-1+ia}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*ArcTan[a + b*x])/x,x]`

```
[Out] (2*(-1)^(3/4)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/Sqrt[b] - (2*Sqrt[-1 - I*a]*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[-1 + I*a]])/Sqrt[-1 + I*a]
```

Maple [A]

time = 0.10, size = 107, normalized size = 1.20

method	result	s
default	$ \frac{ib \ln \left(\frac{b^2 x + ab}{\sqrt{b^2}} + \sqrt{b^2 x^2 + 2abx + a^2 + 1} \right)}{\sqrt{b^2}} - \frac{(ia+1) \ln \left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x} \right)}{\sqrt{a^2+1}} $	1

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x,x,method=_RETURNVERBOSE)`

[Out] $I*b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)}-(1+I*a)/(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x)$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 233 vs. 2(59) = 118.

time = 0.27, size = 233, normalized size = 2.62

$$- \frac{i a \operatorname{arsinh}\left(\frac{2 a b x}{\sqrt{-4 a^2 b^2+4\left(a^2+1\right) b^2}}\right)+\sqrt{-4 a^2 b^2+4\left(a^2+1\right) b^2}}{\sqrt{a^2+1}}-\frac{\operatorname{arsinh}\left(\frac{2 a b x}{\sqrt{-4 a^2 b^2+4\left(a^2+1\right) b^2}}\right)+\sqrt{-4 a^2 b^2+4\left(a^2+1\right) b^2}}{\sqrt{a^2+1}}+i \operatorname{arsinh}\left(\frac{2\left(b^2 x+a b\right)}{\sqrt{-4 a^2 b^2+4\left(a^2+1\right) b^2}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x,x, algorithm="maxima")`

[Out] $-I*a*\operatorname{arcsinh}(2*a*b*x/(\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})*\operatorname{abs}(x))+2*a^2/(\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})*\operatorname{abs}(x)+2/(\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})*\operatorname{abs}(x))/\sqrt{a^2+1}-\operatorname{arcsinh}(2*a*b*x/(\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})*\operatorname{abs}(x))+2*a^2/(\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})*\operatorname{abs}(x)+2/(\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})*\operatorname{abs}(x))/\sqrt{a^2+1}+I*\operatorname{arcsinh}(2*(b^2*x+a*b)/\sqrt{-4*a^2*b^2+4*(a^2+1)*b^2})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(59) = 118.

time = 2.50, size = 144, normalized size = 1.62

$$\sqrt{\frac{a-i}{a+i}} \log\left(-bx+(ia-1)\sqrt{\frac{a-i}{a+i}}+\sqrt{b^2x^2+2abx+a^2+1}\right)-\sqrt{\frac{a-i}{a+i}} \log\left(-bx+(-ia+1)\sqrt{\frac{a-i}{a+i}}+\sqrt{b^2x^2+2abx+a^2+1}\right)-i \log\left(-bx-a+\sqrt{b^2x^2+2abx+a^2+1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)/x,x, algorithm="fricas")`

[Out] $\sqrt{-(a-I)/(a+I)}*\log(-b*x+(I*a-1)*\sqrt{-(a-I)/(a+I)}+\sqrt{b^2*x^2+2*a*b*x+a^2+1})-\sqrt{-(a-I)/(a+I)}*\log(-b*x+(-I*a+1)*\sqrt{-(a-I)/(a+I)}+\sqrt{b^2*x^2+2*a*b*x+a^2+1})-I*\log(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2+1})$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i\left(\int \frac{b}{\sqrt{a^2+2abx+b^2x^2+1}} dx + \int \left(-\frac{i}{x\sqrt{a^2+2abx+b^2x^2+1}}\right) dx + \int \frac{a}{x\sqrt{a^2+2abx+b^2x^2+1}} dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)/x,x)`

[Out] $I*(\operatorname{Integral}(b/\sqrt{a^2+2*a*b*x+b^2*x^2+1}),x)+\operatorname{Integral}(-I/(x*\sqrt{a^2+2*a*b*x+b^2*x^2+1}),x)+\operatorname{Integral}(a/(x*\sqrt{a^2+2*a*b*x+b^2*x^2+1}),x)$

Giac [A]

time = 0.49, size = 112, normalized size = 1.26

$$\frac{(-ia - 1) \log \left(\frac{-2x|b|+2\sqrt{(bx+a)^2+1} - 2\sqrt{a^2+1}}{-2x|b|+2\sqrt{(bx+a)^2+1} + 2\sqrt{a^2+1}} \right)}{\sqrt{a^2+1}} - \frac{ib \log \left(-ab - \left(x|b| - \sqrt{(bx+a)^2+1} \right) |b| \right)}{|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x,x, algorithm="giac")

[Out] $-(I*a - 1) \log(\text{abs}(-2*x*\text{abs}(b) + 2*\text{sqrt}((b*x + a)^2 + 1) - 2*\text{sqrt}(a^2 + 1)) / \text{abs}(-2*x*\text{abs}(b) + 2*\text{sqrt}((b*x + a)^2 + 1) + 2*\text{sqrt}(a^2 + 1))) / \text{sqrt}(a^2 + 1) - I*b*\log(-a*b - (x*\text{abs}(b) - \text{sqrt}((b*x + a)^2 + 1))*\text{abs}(b)) / \text{abs}(b)$

Mupad [B]

time = 1.15, size = 118, normalized size = 1.33

$$\text{asinh}(a + bx) \text{ li} - \frac{\ln \left(ab + \frac{a^2+1}{x} + \frac{\sqrt{a^2+1} \sqrt{a^2+2abx+b^2x^2+1}}{x} \right)}{\sqrt{a^2+1}} - \frac{a \ln \left(ab + \frac{a^2+1}{x} + \frac{\sqrt{a^2+1} \sqrt{a^2+2abx+b^2x^2+1}}{x} \right)}{\sqrt{a^2+1}} \text{ li}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)/(x*((a + b*x)^2 + 1)^(1/2)),x)

[Out] $\text{asinh}(a + b*x) * 1i - \log(a*b + (a^2 + 1)/x + ((a^2 + 1)^(1/2) * (a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2)) / x) / (a^2 + 1)^(1/2) - (a * \log(a*b + (a^2 + 1)/x + ((a^2 + 1)^(1/2) * (a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2)) / x) * 1i) / (a^2 + 1)^(1/2)$

$$3.168 \quad \int \frac{e^{i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=130

$$-\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1-ia)x} + \frac{2ib \tanh^{-1} \left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}} \right)}{\sqrt{i-a} (i+a)^{3/2}}$$

[Out] $2*I*b*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)/(1-I*a-I*b*x)^{(1/2)}})/(I+a)^{(3/2)/(I-a)^{(1/2)}-(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(1-I*a)})/x$

Rubi [A]

time = 0.05, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5203, 96, 95, 214}

$$\frac{2ib \tanh^{-1} \left(\frac{\sqrt{a+i} \sqrt{ia+ibx+1}}{\sqrt{-a+i} \sqrt{-ia-ibx+1}} \right)}{\sqrt{-a+i} (a+i)^{3/2}} - \frac{\sqrt{-ia-ibx+1} \sqrt{ia+ibx+1}}{(1-ia)x}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a + b*x])/x^2,x]

[Out] $-((\operatorname{Sqrt}[1-I*a-I*b*x]*\operatorname{Sqrt}[1+I*a+I*b*x])/((1-I*a)*x)) + ((2*I)*b*\operatorname{ArcTanH}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])])/(\operatorname{Sqrt}[I-a]*(I+a)^{(3/2)})$

Rule 95

Int[(((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.))/((e_.) + (f_.)*(x_.)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.)*((e_.) + (f_.)*(x_.))^(p_.), x_Symbol] :> Simp[(a + b*x)^(m+1)*(c + d*x)^n*(e + f*x)^(p+1)/((m+1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m+1)*(b*e - a*f))], Int[(a + b*x)^(m+1)*(c + d*x)^(n-1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{\sqrt{1+ia+ibx}}{x^2 \sqrt{1-ia-ibx}} dx \\
 &= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1-ia)x} - \frac{b \int \frac{1}{x \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}} dx}{i+a} \\
 &= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1-ia)x} - \frac{(2b) \text{Subst}\left(\int \frac{1}{-1-ia-(-1+ia)x^2} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}}\right)}{i+a} \\
 &= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1-ia)x} + \frac{2ib \tanh^{-1}\left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}}\right)}{\sqrt{i-a} (i+a)^{3/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.06, size = 120, normalized size = 0.92

$$-i \left(\frac{\sqrt{1+a^2+2abx+b^2x^2}}{ix+ax} + \frac{2b \tanh^{-1}\left(\frac{\sqrt{-1-ia} \sqrt{-i(i+a+bx)}}{\sqrt{-1+ia} \sqrt{1+ia+ibx}}\right)}{\sqrt{-1-ia} (-1+ia)^{3/2}} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a + b*x])/x^2,x]

[Out] (-I)*(Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]/(I*x + a*x) + (2*b*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x])])/(Sqrt[-1 - I*a]*(-1 + I*a)^(3/2)))

Maple [A]

time = 0.11, size = 152, normalized size = 1.17

method	result
risch	$-\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{(i+a)x} + \frac{b \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2 + 2abx + a^2 + 1}}{x}\right)}{(i+a)\sqrt{a^2+1}}$
default	$-\frac{ib \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2 + 2abx + a^2 + 1}}{x}\right)}{\sqrt{a^2+1}} + (ia+1) \left(-\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{(a^2+1)x} + \dots \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^2,x,method=_RETURNVERBOSE)`

[Out]
$$-I*b/(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x)+(1+I*a)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b/(a^2+1)^{(3/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x))$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 239 vs. $2(86) = 172$.

time = 0.29, size = 239, normalized size = 1.84

$$\frac{a(i+1)b \operatorname{arsinh}\left(\frac{2abx}{\sqrt{-4a^2b^2+4(a^2+1)b^2|x|}} + \frac{2a^2}{\sqrt{-4a^2b^2+4(a^2+1)b^2|x|}} + \frac{2}{\sqrt{-4a^2b^2+4(a^2+1)b^2|x|}}\right) + ib \operatorname{arsinh}\left(\frac{2abx}{\sqrt{-4a^2b^2+4(a^2+1)b^2|x|}} + \frac{2a^2}{\sqrt{-4a^2b^2+4(a^2+1)b^2|x|}} + \frac{2}{\sqrt{-4a^2b^2+4(a^2+1)b^2|x|}}\right)}{(a^2+1)^{3/2}} + \frac{\sqrt{b^2x^2+2abx+a^2+1}(-i-a-1)}{(a^2+1)x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^2,x, algorithm="maxima")`

[Out]
$$a*(I*a + 1)*b*\operatorname{arcsinh}(2*a*b*x/(\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*\operatorname{abs}(x))) + 2*a^2/(\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*\operatorname{abs}(x)) + 2/(\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*\operatorname{abs}(x)))/(a^2 + 1)^{(3/2)} - I*b*\operatorname{arcsinh}(2*a*b*x/(\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*\operatorname{abs}(x))) + 2*a^2/(\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*\operatorname{abs}(x)) + 2/(\operatorname{sqrt}(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*\operatorname{abs}(x)))/\operatorname{sqrt}(a^2 + 1) + \operatorname{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*a - 1)/((a^2 + 1)*x)$$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 224 vs. $2(86) = 172$.

time = 1.53, size = 224, normalized size = 1.72

$$\frac{(a+i)\sqrt{\frac{b^2}{a^4+2ia^3+2ia-1}}x \log\left(\frac{b^2x-\sqrt{b^2x^2+2abx+a^2+1}b^{(a^3+ia^2+ia+1)}}{b}\sqrt{\frac{b^2}{a^4+2ia^3+2ia-1}}\right) - (a+i)\sqrt{\frac{b^2}{a^4+2ia^3+2ia-1}}x \log\left(\frac{b^2x-\sqrt{b^2x^2+2abx+a^2+1}b^{-(a^3+ia^2+ia+1)}}{b}\sqrt{\frac{b^2}{a^4+2ia^3+2ia-1}}\right) + ibx + i\sqrt{b^2x^2+2abx+a^2+1}}{(a+i)x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^2,x, algorithm="fricas")`

[Out]
$$-((a + I)*\operatorname{sqrt}(b^2/(a^4 + 2*I*a^3 + 2*I*a - 1))*x*\log(-(b^2*x - \operatorname{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1))*b + (a^3 + I*a^2 + a + I)*\operatorname{sqrt}(b^2/(a^4 + 2*I*a^3 + \dots)))$$

$(2Ia - 1))/b - (a + I)\sqrt{b^2/(a^4 + 2Ia^3 + 2Ia - 1)}x \log(-b^2 x - \sqrt{b^2 x^2 + 2a b x + a^2 + 1}b - (a^3 + I a^2 + a + I)\sqrt{b^2/(a^4 + 2Ia^3 + 2Ia - 1)})/b + I b x + I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}/((a + I)x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \left(-\frac{i}{x^2 \sqrt{a^2 + 2abx + b^2 x^2 + 1}} \right) dx + \int \frac{a}{x^2 \sqrt{a^2 + 2abx + b^2 x^2 + 1}} dx + \int \frac{b}{x \sqrt{a^2 + 2abx + b^2 x^2 + 1}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)/x**2,x)

[Out] I*(Integral(-I/(x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(a/(x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(b/(x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x))

Giac [A]

time = 0.49, size = 145, normalized size = 1.12

$$\frac{b \log \left(\frac{\left| \begin{array}{l} 2x|b|-2 \sqrt{(bx+a)^2 + 1} - 2 \sqrt{a^2 + 1} \\ 2x|b|-2 \sqrt{(bx+a)^2 + 1} + 2 \sqrt{a^2 + 1} \end{array} \right|}{\sqrt{a^2 + 1} (a + i)} \right)}{\left(\left(x|b| - \sqrt{(bx+a)^2 + 1} \right) ab + a^2 |b| + |b| \right) \left(\left(x|b| - \sqrt{(bx+a)^2 + 1} \right)^2 - a^2 - 1 \right) (ia - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^2,x, algorithm="giac")

[Out] b*log(abs(2*x*abs(b) - 2*sqrt((b*x + a)^2 + 1) - 2*sqrt(a^2 + 1))/abs(2*x*abs(b) - 2*sqrt((b*x + a)^2 + 1) + 2*sqrt(a^2 + 1)))/(sqrt(a^2 + 1)*(a + I)) - 2*((x*abs(b) - sqrt((b*x + a)^2 + 1))*a*b + a^2*abs(b) + abs(b))/((x*abs(b) - sqrt((b*x + a)^2 + 1))^2 - a^2 - 1)*(I*a - 1)

Mupad [B]

time = 1.68, size = 218, normalized size = 1.68

$$\frac{a \operatorname{batanh} \left(\frac{a^2 + b x + 1}{\sqrt{a^2 + 1} \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} \right)}{(a^2 + 1)^{3/2}} - \frac{\sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x (a^2 + 1)} - \frac{b \ln \left(a b + \frac{a^2 + 1}{x} + \frac{\sqrt{a^2 + 1} \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} \right)}{\sqrt{a^2 + 1}} + \frac{a^2 \operatorname{batanh} \left(\frac{a^2 + b x + 1}{\sqrt{a^2 + 1} \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} \right)}{(a^2 + 1)^{3/2}} - \frac{a \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x (a^2 + 1)} \operatorname{li}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a**1i + b*x**1i + 1)/(x**2*((a + b*x)^2 + 1)^(1/2)),x)

[Out] (a^2*b*atanh((a^2 + a*b*x + 1)/((a^2 + 1)^(1/2)*(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2)))*1i)/(a^2 + 1)^(3/2) - (a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2)/(x*(a^2 + 1)) - (b*log(a*b + (a^2 + 1)/x + ((a^2 + 1)^(1/2)*(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2))/x)*1i)/(a^2 + 1)^(1/2) - (a*(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2)*1i)/(x*(a^2 + 1)) + (a*b*atanh((a^2 + a*b*x + 1)/((a^2 + 1)^(1/2)*(a^2 + b^2*x^2 + 2*a*b*x + 1)^(1/2))))/(a^2 + 1)^(3/2)

$$3.169 \quad \int \frac{e^{i \operatorname{ArcTan}(a+bx)}}{x^3} dx$$

Optimal. Leaf size=201

$$-\frac{(1+2ia)b\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2(i-a)(i+a)^2x} - \frac{\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}}{2(1+a^2)x^2} + \frac{(1+2ia)b^2 \tanh^{-1}\left(\frac{\sqrt{i+a}}{\sqrt{i-a}}\right)}{(i-a)^{3/2}(i+a)}$$

[Out] (1+2*I*a)*b^2*arctanh((I+a)^(1/2)*(1+I*a+I*b*x)^(1/2)/(I-a)^(1/2)/(1-I*a-I*b*x)^(1/2))/(I-a)^(3/2)/(I+a)^(5/2)-1/2*(1+I*a+I*b*x)^(3/2)*(1-I*a-I*b*x)^(1/2)/(a^2+1)/x^2-1/2*(1+2*I*a)*b*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/(I-a)/(I+a)^2/x

Rubi [A]

time = 0.11, antiderivative size = 201, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {5203, 98, 96, 95, 214}

$$-\frac{\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{2(a^2+1)x^2} + \frac{(1+2ia)b^2 \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{3/2}(a+i)^{5/2}} - \frac{(1+2ia)b\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{2(-a+i)(a+i)^2x}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a + b*x])/x^3,x]

[Out] -1/2*((1 + (2*I)*a)*b*Sqrt[1 - I*a - I*b*x]*Sqrt[1 + I*a + I*b*x])/((I - a)*(I + a)^2*x) - (Sqrt[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^(3/2))/(2*(1 + a^2)*x^2) + (((1 + (2*I)*a)*b^2*ArcTanh[(Sqrt[I + a]*Sqrt[1 + I*a + I*b*x])/(Sqrt[I - a]*Sqrt[1 - I*a - I*b*x])])/((I - a)^(3/2)*(I + a)^(5/2))

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{i \tan^{-1}(a+bx)}}{x^3} dx &= \int \frac{\sqrt{1+ia+ibx}}{x^3 \sqrt{1-ia-ibx}} dx \\ &= -\frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2(1+a^2)x^2} + \frac{((i-2a)b) \int \frac{\sqrt{1+ia+ibx}}{x^2 \sqrt{1-ia-ibx}} dx}{2(1+a^2)} \\ &= -\frac{(i-2a)b \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2(1-ia)(1+a^2)x} - \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2(1+a^2)x^2} - \frac{((i-2a)b) \int \frac{\sqrt{1+ia+ibx}}{x \sqrt{1-ia-ibx}} dx}{2(1+a^2)} \\ &= -\frac{(i-2a)b \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2(1-ia)(1+a^2)x} - \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2(1+a^2)x^2} - \frac{((i-2a)b) \int \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx}{2(1+a^2)} \\ &= -\frac{(i-2a)b \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2(1-ia)(1+a^2)x} - \frac{\sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{2(1+a^2)x^2} + \frac{(1+2ia) \int \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx}{2(1+a^2)} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 154, normalized size = 0.77

$$\frac{\frac{i(1+a^2+2ibx-abx)\sqrt{1+a^2+2abx+b^2x^2}}{x^2} + \frac{2(-i+2a)b^2 \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}{\sqrt{-1-ia}\sqrt{-1+ia}}}{2(-i+a)(i+a)^2}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a + b*x])/x^3,x]

[Out] (((-I)*(1 + a^2 + (2*I)*b*x - a*b*x)*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2])/x^2 + (2*(-I + 2*a)*b^2*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)]]/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]))/(Sqrt[-1 - I*a]*Sqrt[-1 + I*a]))/(2*(-I + a)*(I + a)^2)

Maple [A]

time = 0.10, size = 287, normalized size = 1.43

method	result
risch	$-\frac{i(-ab^3x^3+2ib^3x^3-a^2b^2x^2+4iab^2x^2+a^3bx+2ix^2a^2b+a^4+b^2x^2+abx+2ibx+2a^2+1)}{2x^2(i+a)^2(a-i)\sqrt{b^2x^2+2abx+a^2+1}} + \frac{ib^2 \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{2(a^2+1)^{\frac{3}{2}}(i+a)}$
default	$(ia + 1) \left(-\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{2(a^2+1)x^2} - \frac{3ab \left(-\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{(a^2+1)x} + \frac{ab \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{(a^2+1)} \right)}{2(a^2+1)} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^3,x,method=_RETURNVERBOSE)

[Out] (1+I*a)*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b/(a^2+1)^(3/2)*ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))+1/2*b^2/(a^2+1)^(3/2)*ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))+I*b*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b/(a^2+1)^(3/2)*ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 424 vs. 2(135) = 270.

time = 0.28, size = 424, normalized size = 2.11

$\frac{3a^2(x+1)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{-4ab+4}(ax+1)^{3/2}}{\sqrt{-4ab+4}(ax+1)^{3/2}}\right) + \sqrt{-4ab+4}(ax+1)^{3/2}}{2(a^2+1)^2} - \frac{1ab^2 \operatorname{arctanh}\left(\frac{\sqrt{-4ab+4}(ax+1)^{3/2}}{\sqrt{-4ab+4}(ax+1)^{3/2}}\right) + \sqrt{-4ab+4}(ax+1)^{3/2}}{2(a^2+1)^2} - \frac{(-1+1)^{3/2} \operatorname{arctanh}\left(\frac{\sqrt{-4ab+4}(ax+1)^{3/2}}{\sqrt{-4ab+4}(ax+1)^{3/2}}\right) + \sqrt{-4ab+4}(ax+1)^{3/2}}{2(a^2+1)^2} - \frac{3\sqrt{b^2+2abx+2}\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{2(a^2+1)^2} - \frac{ab \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{2(a^2+1)^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^3,x, algorithm="maxima")

[Out]
$$-3/2*a^2*(I*a + 1)*b^2*\operatorname{arcsinh}(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x)) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x))/(a^2 + 1)^{(5/2)} + I*a*b^2*\operatorname{arcsinh}(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x)) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x))/(a^2 + 1)^{(3/2)} - 1/2*(-I*a - 1)*b^2*\operatorname{arcsinh}(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x)) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(x))/(a^2 + 1)^{(3/2)} + 3/2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a*(I*a + 1)*b/((a^2 + 1)^2*x) - I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*b/((a^2 + 1)*x) - 1/2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(I*a + 1)/((a^2 + 1)*x^2)$$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 452 vs. $2(135) = 270$.
time = 2.09, size = 452, normalized size = 2.25

$$\frac{(1 + 2b^2x^2 + \sqrt{b^2x^2 + 2abx + a^2 + 1})^{1/2} (a^2 + 2a^2 + 2b^2 + 6a^2 + 6a^2 - 2a^2 - 2a^2 - 1) \operatorname{arcsinh}\left(\frac{2abx + a^2 + 1}{\sqrt{b^2x^2 + 2abx + a^2 + 1}}\right) + \frac{(4a^2 - 4a - 1) \sqrt{b^2x^2 + 2abx + a^2 + 1}}{2(b^2 + a^2 + 1)^2} (a^2 + 2a^2 + 2b^2 + 6a^2 + 6a^2 - 2a^2 - 2a^2 - 1) \operatorname{arcsinh}\left(\frac{2abx + a^2 + 1}{\sqrt{b^2x^2 + 2abx + a^2 + 1}}\right) - \frac{(4a^2 - 4a - 1) \sqrt{b^2x^2 + 2abx + a^2 + 1}}{2(b^2 + a^2 + 1)^2} (a^2 + 2a^2 + 2b^2 + 6a^2 + 6a^2 - 2a^2 - 2a^2 - 1) \operatorname{arcsinh}\left(\frac{2abx + a^2 + 1}{\sqrt{b^2x^2 + 2abx + a^2 + 1}}\right) + \sqrt{b^2x^2 + 2abx + a^2 + 1} (1 + 2bx - a^2 - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^3,x, algorithm="fricas")

[Out]
$$1/2*((I*a + 2)*b^2*x^2 + \sqrt{(4*a^2 - 4*I*a - 1)*b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1)}*(a^3 + I*a^2 + a + I))*x^2*\log(-((2*a - I)*b^3*x - \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(2*a - I)*b^2 + (a^5 + I*a^4 + 2*a^3 + 2*I*a^2 + a + I)*\sqrt{(4*a^2 - 4*I*a - 1)*b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1)}))/((2*a - I)*b^2)) - \sqrt{(4*a^2 - 4*I*a - 1)*b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1)}*(a^3 + I*a^2 + a + I))*x^2*\log(-((2*a - I)*b^3*x - \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(2*a - I)*b^2 - (a^5 + I*a^4 + 2*a^3 + 2*I*a^2 + a + I)*\sqrt{(4*a^2 - 4*I*a - 1)*b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1)}))/((2*a - I)*b^2)) + \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*((I*a + 2)*b*x - I*a^2 - I))/((a^3 + I*a^2 + a + I)*x^2)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \left(-\frac{i}{x^3 \sqrt{a^2 + 2abx + b^2x^2 + 1}} \right) dx + \int \frac{a}{x^3 \sqrt{a^2 + 2abx + b^2x^2 + 1}} dx + \int \frac{b}{x^2 \sqrt{a^2 + 2abx + b^2x^2 + 1}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)/x**3,x)

[Out] $I \cdot (\text{Integral}(-I/(x^{3}\sqrt{a^{2} + 2abx + b^{2}x^{2} + 1})), x) + \text{Integral}(a/(x^{3}\sqrt{a^{2} + 2abx + b^{2}x^{2} + 1}), x) + \text{Integral}(b/(x^{2}\sqrt{a^{2} + 2abx + b^{2}x^{2} + 1}), x)$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 471 vs. $2(135) = 270$.
time = 0.53, size = 471, normalized size = 2.34

$$\frac{\frac{a^2 - I^2 \log\left(\frac{(a + \sqrt{a^2 + 1} - \sqrt{a^2 + 1})}{(a + \sqrt{a^2 + 1} + \sqrt{a^2 + 1})}\right)}{2(a^2 + 1)\sqrt{a^2 + 1}}}{\frac{(-I(a + \sqrt{a^2 + 1}) - \sqrt{a^2 + 1})^{2a} - 2(I(a + \sqrt{a^2 + 1}) - \sqrt{a^2 + 1})^{2a-1} - 2(I(a + \sqrt{a^2 + 1}) - \sqrt{a^2 + 1})^{2a-2} + \dots + (-I(a + \sqrt{a^2 + 1}) - \sqrt{a^2 + 1})^{-2a}}{(a + \sqrt{a^2 + 1})^{2a-1} - a^{-1}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^3,x, algorithm="giac")`

[Out] $-1/2*(2*a*b^2 - I*b^2)*\log(\text{abs}(2*x*\text{abs}(b) - 2*\sqrt{(b*x + a)^2 + 1}) - 2*\sqrt{a^2 + 1})/\text{abs}(2*x*\text{abs}(b) - 2*\sqrt{(b*x + a)^2 + 1}) + 2*\sqrt{a^2 + 1})/((a^3 + I*a^2 + a + I)*\sqrt{a^2 + 1}) - (4*(-I*x*\text{abs}(b) + I*\sqrt{(b*x + a)^2 + 1}))*a^4*b^2 - 2*I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a^3*b*\text{abs}(b) - 2*I*a^5*b*\text{abs}(b) + 2*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^3*a*b^2 - 2*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*a^3*b^2 + 2*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a^2*b*\text{abs}(b) - 2*a^4*b*\text{abs}(b) - I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^3*b^2 + 5*(-I*x*\text{abs}(b) + I*\sqrt{(b*x + a)^2 + 1})*a^2*b^2 - 2*I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a*b*\text{abs}(b) - 4*I*a^3*b*\text{abs}(b) - 2*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*a*b^2 + 2*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*b*\text{abs}(b) - 4*a^2*b*\text{abs}(b) - (I*x*\text{abs}(b) - I*\sqrt{(b*x + a)^2 + 1})*b^2 - 2*I*a*b*\text{abs}(b) - 2*b*\text{abs}(b))/((a^3 + I*a^2 + a + I)*((x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2 - a^2 - 1)^2)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1 + a \operatorname{li} + b x \operatorname{li}}{x^3 \sqrt{(a + b x)^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*li + b*x*li + 1)/(x^3*((a + b*x)^2 + 1)^(1/2)),x)`

[Out] `int((a*li + b*x*li + 1)/(x^3*((a + b*x)^2 + 1)^(1/2)), x)`

3.170 $\int \frac{e^{i \operatorname{ArcTan}(a+bx)}}{x^4} dx$

Optimal. Leaf size=283

$$-\frac{\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{3(1-ia)x^3} - \frac{(3i-2a)b\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} + \frac{(4+9ia-2a^2)b^2\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x}$$

[Out] $(2a-I*(-2a^2+1))*b^3*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)/(1-I*a-I*b*x)^{(1/2)))/(I-a)^{(5/2)/(I+a)^{(7/2)}-1/3*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(1-I*a)/x^3-1/6*(3I-2a)*b*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(1-I*a)/(a^2+1)/x^2+1/6*(4+9I*a-2a^2)*b^2*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(1-I*a)/(a^2+1)^2/x}$

Rubi [A]

time = 0.16, antiderivative size = 283, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5203, 101, 156, 12, 95, 214}

$$\frac{(2a-i(1-2a^2))b^3 \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{5/2}(a+i)^{7/2}} + \frac{(-2a^2+9ia+4)b^2\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{6(1-ia)(a^2+1)^2x} - \frac{(-2a+3i)b\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{6(1-ia)(a^2+1)x^2} - \frac{\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{3(1-ia)x^3}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(I*\operatorname{ArcTan}[a + b*x])}/x^4, x]$

[Out] $-1/3*(\operatorname{Sqrt}[1 - I*a - I*b*x]*\operatorname{Sqrt}[1 + I*a + I*b*x])/((1 - I*a)*x^3) - ((3*I - 2*a)*b*\operatorname{Sqrt}[1 - I*a - I*b*x]*\operatorname{Sqrt}[1 + I*a + I*b*x])/(6*(1 - I*a)*(1 + a^2)*x^2) + ((4 + (9*I)*a - 2*a^2)*b^2*\operatorname{Sqrt}[1 - I*a - I*b*x]*\operatorname{Sqrt}[1 + I*a + I*b*x])/(6*(1 - I*a)*(1 + a^2)^2*x) + ((2*a - I*(1 - 2*a^2))*b^3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I + a]*\operatorname{Sqrt}[1 + I*a + I*b*x])]/(\operatorname{Sqrt}[I - a]*\operatorname{Sqrt}[1 - I*a - I*b*x]))/((I - a)^{(5/2)}*(I + a)^{(7/2)})$

Rule 12

$\operatorname{Int}[(a_*)*(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{MatchQ}[u, (b_*)*(v_) /; \operatorname{FreeQ}[b, x]]$

Rule 95

$\operatorname{Int}[(((a_.) + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)})/((e_.) + (f_.)*(x_)), x_Symbol] \rightarrow \operatorname{With}[q = \operatorname{Denominator}[m], \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \ \&\& \ \operatorname{EqQ}[m + n + 1, 0] \ \&\& \ \operatorname{RationalQ}[n] \ \&\& \ \operatorname{LtQ}[-1, m, 0] \ \&\& \ \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 101

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1)/((m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m + n + p + 2)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 214

```

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

```

Rule 5203

```

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{i \tan^{-1}(a+bx)}}{x^4} dx &= \int \frac{\sqrt{1+ia+ibx}}{x^4 \sqrt{1-ia-ibx}} dx \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1-ia)x^3} + \frac{\int \frac{(3i-2a)b-2b^2x}{x^3 \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}} dx}{3(1-ia)} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1-ia)x^3} - \frac{(3i-2a)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} - \frac{\int \frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{x^2 \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}} dx}{3(1-ia)} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1-ia)x^3} - \frac{(3i-2a)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} + \frac{(4+9ia)\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1-ia)x^3} - \frac{(3i-2a)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} + \frac{(4+9ia)\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1-ia)x^3} - \frac{(3i-2a)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} + \frac{(4+9ia)\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1-ia)x^3} - \frac{(3i-2a)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2} + \frac{(4+9ia)\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(1-ia)(1+a^2)x^2}
\end{aligned}$$

Mathematica [A]

time = 0.22, size = 247, normalized size = 0.87

$$\frac{2(1-ia)(-i+a)(-i+a+bx)\sqrt{1+a^2+2abx+b^2x^2} + (1+4ia)bx(-i+a+bx)\sqrt{1+a^2+2abx+b^2x^2} + \frac{3(-1-2ia+2a^2)b^2x^2 \left(\sqrt{-1-ia} \sqrt{-1+ia} \sqrt{1+a^2+2abx+b^2x^2} - 2ibx \tanh^{-1} \left(\frac{\sqrt{-1-ia} \sqrt{-i(i+a+bx)}}{\sqrt{-1+ia} \sqrt{1+ia+ibx}} \right) \right)}{6(1+a^2)^2 x^3}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^(I*ArcTan[a + b*x])/x^4,x]

[Out] (2*(1 - I*a)*(-I + a)*(-I + a + b*x)*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2] + (1 + (4*I)*a)*b*x*(-I + a + b*x)*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2] + (3*(-1 - (2*I)*a + 2*a^2)*b^2*x^2*(Sqrt[-1 - I*a]*Sqrt[-1 + I*a]*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2] - (2*I)*b*x*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/ (Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x])]))/(Sqrt[-1 - I*a]*(-1 + I*a)^(3/2)))/(6*(1 + a^2)^2*x^3)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 528 vs. 2(227) = 454.

time = 0.13, size = 529, normalized size = 1.87

method	result
risch	$\frac{i(2a^2b^4x^4 - 9iab^4x^4 + 2a^3b^3x^3 - 15ia^2b^3x^3 - 3ia^3b^2x^2 - 4x^4b^4 + 2a^5bx + 3ia^4bx - 10ab^3x^3 + 3ib^3x^3 + 2a^6 - 2a^2b^2x^2 - 3iab^2x^2 + 4a^3bx)}{6x^3(a-i)^2(i+a)^3\sqrt{b^2x^2 + 2abx + a^2 + 1}}$
default	$(ia + 1) \left(-\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{3(a^2+1)x^3} - \frac{5ab \left(-\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{2(a^2+1)x^2} - \frac{3ab \left(-\frac{\sqrt{b^2x^2 + 2abx + a^2}}{(a^2+1)x} \right)}{3ab} \right)}{3(a^2+1)x^3} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^4,x,method=_RETURNVERBOSE)`

[Out] $(1+I*a)*(-1/3/(a^2+1)/x^3*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/3*a*b/(a^2+1)*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))+1/2*b^2/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-2/3*b^2/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x)))+I*b*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))+1/2*b^2/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 644 vs. 2(198) = 396.

time = 0.27, size = 644, normalized size = 2.28

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^4,x, algorithm="maxima")`

```
[Out] 5/2*a^3*(I*a + 1)*b^3*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*a
bs(x)) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2
*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(7/2) - 3/2*I*a^2*b^3*arcsinh(2*
a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqrt(-4*a^2*b^2
+ 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x))
)/(a^2 + 1)^(5/2) - 3/2*a*(I*a + 1)*b^3*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4
*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x))
+ 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(5/2) + 1/2*I*b
^3*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqr
t(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*
b^2)*abs(x)))/(a^2 + 1)^(3/2) - 5/2*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*a^2*(
I*a + 1)*b^2/((a^2 + 1)^3*x) + 3/2*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*a*b^
2/((a^2 + 1)^2*x) - 2/3*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*a - 1)*b^2/((
a^2 + 1)^2*x) + 5/6*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*a*(I*a + 1)*b/((a^2 +
1)^2*x^2) - 1/2*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b/((a^2 + 1)*x^2) - 1/
3*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(I*a + 1)/((a^2 + 1)*x^3)
```

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 690 vs. $2(198) = 396$.
time = 1.19, size = 690, normalized size = 2.44

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^4,x, algorithm="fricas")
```

```
[Out] 1/6*((-2*I*a^2 - 9*a + 4*I)*b^3*x^3 - 3*sqrt((4*a^4 - 8*I*a^3 - 8*a^2 + 4*I
*a + 1)*b^6/(a^12 + 2*I*a^11 + 4*a^10 + 10*I*a^9 + 5*a^8 + 20*I*a^7 + 20*I*
a^5 - 5*a^4 + 10*I*a^3 - 4*a^2 + 2*I*a - 1))*(a^5 + I*a^4 + 2*a^3 + 2*I*a^2
+ a + I)*x^3*log(-((2*a^2 - 2*I*a - 1)*b^4*x - sqrt(b^2*x^2 + 2*a*b*x + a^
2 + 1)*(2*a^2 - 2*I*a - 1)*b^3 + (a^7 + I*a^6 + 3*a^5 + 3*I*a^4 + 3*a^3 + 3
*I*a^2 + a + I)*sqrt((4*a^4 - 8*I*a^3 - 8*a^2 + 4*I*a + 1)*b^6/(a^12 + 2*I*
a^11 + 4*a^10 + 10*I*a^9 + 5*a^8 + 20*I*a^7 + 20*I*a^5 - 5*a^4 + 10*I*a^3 -
4*a^2 + 2*I*a - 1)))/((2*a^2 - 2*I*a - 1)*b^3)) + 3*sqrt((4*a^4 - 8*I*a^3
- 8*a^2 + 4*I*a + 1)*b^6/(a^12 + 2*I*a^11 + 4*a^10 + 10*I*a^9 + 5*a^8 + 20*
I*a^7 + 20*I*a^5 - 5*a^4 + 10*I*a^3 - 4*a^2 + 2*I*a - 1))*(a^5 + I*a^4 + 2*
a^3 + 2*I*a^2 + a + I)*x^3*log(-((2*a^2 - 2*I*a - 1)*b^4*x - sqrt(b^2*x^2 +
2*a*b*x + a^2 + 1)*(2*a^2 - 2*I*a - 1)*b^3 - (a^7 + I*a^6 + 3*a^5 + 3*I*a^
4 + 3*a^3 + 3*I*a^2 + a + I)*sqrt((4*a^4 - 8*I*a^3 - 8*a^2 + 4*I*a + 1)*b^6
/(a^12 + 2*I*a^11 + 4*a^10 + 10*I*a^9 + 5*a^8 + 20*I*a^7 + 20*I*a^5 - 5*a^4
+ 10*I*a^3 - 4*a^2 + 2*I*a - 1)))/((2*a^2 - 2*I*a - 1)*b^3)) + ((-2*I*a^2
- 9*a + 4*I)*b^2*x^2 - 2*I*a^4 + (2*I*a^3 + 3*a^2 + 2*I*a + 3)*b*x - 4*I*a^
2 - 2*I)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/((a^5 + I*a^4 + 2*a^3 + 2*I*a^2
+ a + I)*x^3)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \left(-\frac{i}{x^4 \sqrt{a^2 + 2abx + b^2x^2 + 1}} \right) dx + \int \frac{a}{x^4 \sqrt{a^2 + 2abx + b^2x^2 + 1}} dx + \int \frac{b}{x^3 \sqrt{a^2 + 2abx + b^2x^2 + 1}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2)/x**4,x)

[Out] I*(Integral(-I/(x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(a/(x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(b/(x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x))

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 884 vs. 2(198) = 396.

time = 0.49, size = 884, normalized size = 3.12

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2)/x^4,x, algorithm="giac")

[Out] 1/2*(2*a^2*b^3 - 2*I*a*b^3 - b^3)*log(abs(2*x*abs(b) - 2*sqrt((b*x + a)^2 + 1) - 2*sqrt(a^2 + 1))/abs(2*x*abs(b) - 2*sqrt((b*x + a)^2 + 1) + 2*sqrt(a^2 + 1)))/((a^5 + I*a^4 + 2*a^3 + 2*I*a^2 + a + I)*sqrt(a^2 + 1)) + 1/3*(8*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*a^5*b^3 + 24*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a^7*b^3 + 24*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a^6*b^2*abs(b) + 8*I*a^8*b^2*abs(b) + 6*(x*abs(b) - sqrt((b*x + a)^2 + 1))^5*a^2*b^3 - 24*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*a^4*b^3 + 18*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^6*b^3 - 12*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a^5*b^2*abs(b) + 12*a^7*b^2*abs(b) - 6*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^5*a*b^3 + 32*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*a^3*b^3 + 54*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a^5*b^3 + 60*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a^4*b^2*abs(b) + 20*I*a^6*b^2*abs(b) - 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^5*b^3 - 24*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*a^2*b^3 + 39*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^4*b^3 - 24*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a^3*b^2*abs(b) + 36*a^5*b^2*abs(b) + 24*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*a*b^3 + 36*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a^3*b^3 + 48*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a^2*b^2*abs(b) + 12*I*a^4*b^2*abs(b) + 24*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^2*b^3 - 12*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b^2*abs(b) + 36*a^3*b^2*abs(b) + 6*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a*b^3 + 12*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b^2*abs(b) - 4*I*a^2*b^2*abs(b) + 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*b^3 + 12*a*b^2*abs(b) - 4*I*b^2*abs(b))/((a^5 + I*a^4 + 2*a^3 + 2*I*a^2 + a + I)*((x*abs(b) - sqrt((b*x + a)^2 + 1))^2 - a^2 - 1)^3)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1 + a x + b x^2}{x^4 \sqrt{(a + b x)^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x + b*x^2 + 1)/(x^4*((a + b*x)^2 + 1)^(1/2)), x)

[Out] int((a*x + b*x^2 + 1)/(x^4*((a + b*x)^2 + 1)^(1/2)), x)

3.171 $\int e^{2i\text{ArcTan}(a+bx)} x^4 dx$

Optimal. Leaf size=92

$$-\frac{2(1-ia)^3x}{b^4} + \frac{i(i+a)^2x^2}{b^3} + \frac{2(1-ia)x^3}{3b^2} + \frac{ix^4}{2b} - \frac{x^5}{5} + \frac{2i(i+a)^4 \log(i+a+bx)}{b^5}$$

[Out] $-2*(1-I*a)^3*x/b^4+I*(I+a)^2*x^2/b^3+2/3*(1-I*a)*x^3/b^2+1/2*I*x^4/b-1/5*x^5+2*I*(I+a)^4*\ln(I+a+b*x)/b^5$

Rubi [A]

time = 0.07, antiderivative size = 92, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$\frac{2i(a+i)^4 \log(a+bx+i)}{b^5} - \frac{2(1-ia)^3x}{b^4} + \frac{i(a+i)^2x^2}{b^3} + \frac{2(1-ia)x^3}{3b^2} + \frac{ix^4}{2b} - \frac{x^5}{5}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a + b*x])}*x^4, x]$

[Out] $(-2*(1 - I*a)^3*x)/b^4 + (I*(I + a)^2*x^2)/b^3 + (2*(1 - I*a)*x^3)/(3*b^2) + ((I/2)*x^4)/b - x^5/5 + ((2*I)*(I + a)^4*\text{Log}[I + a + b*x])/b^5$

Rule 78

$\text{Int}[((a_.) + (b_.)*(x_.)) * ((c_.) + (d_.)*(x_.))^{(n_.)} * ((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) || \text{EqQ}[p, 1] || (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] || \text{LeQ}[9*p + 5*(n + 2), 0] || \text{GeQ}[n + p + 1, 0] || (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5203

$\text{Int}[E^{(\text{ArcTan}[(c_.)*(a_.) + (b_.)*(x_.)])^{(n_.)} * ((d_.) + (e_.)*(x_.))^{(m_.)}, x_Symbol] :> \text{Int}[(d + e*x)^m * ((1 - I*a*c - I*b*c*x)^{(I*(n/2)}) / (1 + I*a*c + I*b*c*x)^{(I*(n/2)})), x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(a+bx)} x^4 dx &= \int \frac{x^4(1+ia+ibx)}{1-ia-ibx} dx \\ &= \int \left(\frac{2(-1+ia)^3}{b^4} + \frac{2i(i+a)^2x}{b^3} + \frac{2(1-ia)x^2}{b^2} + \frac{2ix^3}{b} - x^4 + \frac{2i(i+a)^4}{b^4(i+a+bx)} \right) dx \\ &= -\frac{2(1-ia)^3x}{b^4} + \frac{i(i+a)^2x^2}{b^3} + \frac{2(1-ia)x^3}{3b^2} + \frac{ix^4}{2b} - \frac{x^5}{5} + \frac{2i(i+a)^4 \log(i+a+bx)}{b^5} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 92, normalized size = 1.00

$$-\frac{2(1-ia)^3x}{b^4} + \frac{i(i+a)^2x^2}{b^3} + \frac{2(1-ia)x^3}{3b^2} + \frac{ix^4}{2b} - \frac{x^5}{5} + \frac{2i(i+a)^4 \log(i+a+bx)}{b^5}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a + b*x])*x^4,x]

[Out] (-2*(1 - I*a)^3*x)/b^4 + (I*(I + a)^2*x^2)/b^3 + (2*(1 - I*a)*x^3)/(3*b^2) + ((I/2)*x^4)/b - x^5/5 + ((2*I)*(I + a)^4*Log[I + a + b*x])/b^5

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 227 vs. 2(78) = 156.

time = 0.16, size = 228, normalized size = 2.48

method	result
default	$-\frac{i(-\frac{1}{5}ib^4x^5 - \frac{1}{2}b^3x^4 + \frac{2}{3}ib^2x^3 + \frac{2}{3}ab^2x^3 - 2iabx^2 - a^2bx^2 + 6ia^2x + 2a^3x + x^2b - 2ix - 6ax)}{b^4} + \frac{(2ia^4b - 8a^3b - 12ia^2b + 8ab + 2ib) \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2}$
risch	$-\frac{x^5}{5} - \frac{8i \arctan(bx+a)a}{b^5} + \frac{2x^3}{3b^2} - \frac{2ia^3x}{b^4} - \frac{2ax^2}{b^3} + \frac{8i \arctan(bx+a)a^3}{b^5} + \frac{6a^2x}{b^4} + \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)a^4}{b^5} + \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)}{b^5}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^4,x,method=_RETURNVERBOSE)

[Out] -I/b^4*(-1/5*I*b^4*x^5-1/2*b^3*x^4+2/3*I*b^2*x^3+2/3*a*b^2*x^3-2*I*a*b*x^2-a^2*b*x^2+6*I*a^2*x+2*a^3*x+x^2*b-2*I*x-6*a*x)+1/b^4*(1/2*(2*I*a^4*b-12*I*a^2*b-8*a^3*b+2*I*b+8*a*b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(2*I*a^5-4*I*a^3-6*a^4-6*I*a-4*a^2+2-(2*I*a^4*b-12*I*a^2*b-8*a^3*b+2*I*b+8*a*b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b))

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 149 vs. 2(70) = 140.

time = 0.47, size = 149, normalized size = 1.62

$$\frac{6b^4x^5 - 15ib^3x^4 + 20(ia-1)b^2x^3 + 30(-ia^2 + 2a+ib)x^2 + 60(ia^3 - 3a^2 - 3ia+1)x + 2(a^4 + 4ia^3 - 6a^2 - 4ia+1) \arctan\left(\frac{b^2x+ab}{b}\right) + (ia^4 - 4a^3 - 6ia^2 + 4a+i) \log(b^2x^2 + 2abx + a^2 + 1)}{30b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^4,x, algorithm="maxima")

[Out] -1/30*(6*b^4*x^5 - 15*I*b^3*x^4 + 20*(I*a - 1)*b^2*x^3 + 30*(-I*a^2 + 2*a + I)*b*x^2 + 60*(I*a^3 - 3*a^2 - 3*I*a + 1)*x)/b^4 + 2*(a^4 + 4*I*a^3 - 6*a^2 - 4*I*a + 1)*arctan((b^2*x + a*b)/b)/b^5 + (I*a^4 - 4*a^3 - 6*I*a^2 + 4*a + I)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^5

Fricas [A]

time = 2.57, size = 105, normalized size = 1.14

$$\frac{6b^5x^5 - 15ib^4x^4 + 20(ia - 1)b^3x^3 + 30(-ia^2 + 2a + i)b^2x^2 + 60(ia^3 - 3a^2 - 3ia + 1)bx + 60(-ia^4 + 4a^3 + 6ia^2 - 4a - i)\log\left(\frac{bx+a+i}{b}\right)}{30b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^4,x, algorithm="fricas")

[Out] -1/30*(6*b^5*x^5 - 15*I*b^4*x^4 + 20*(I*a - 1)*b^3*x^3 + 30*(-I*a^2 + 2*a + I)*b^2*x^2 + 60*(I*a^3 - 3*a^2 - 3*I*a + 1)*b*x + 60*(-I*a^4 + 4*a^3 + 6*I*a^2 - 4*a - I)*log((b*x + a + I)/b))/b^5

Sympy [A]

time = 0.25, size = 110, normalized size = 1.20

$$-\frac{x^5}{5} - x^3 \cdot \left(\frac{2ia}{3b^2} - \frac{2}{3b^2}\right) - x^2 \left(-\frac{ia^2}{b^3} + \frac{2a}{b^3} + \frac{i}{b^3}\right) - x \left(\frac{2ia^3}{b^4} - \frac{6a^2}{b^4} - \frac{6ia}{b^4} + \frac{2}{b^4}\right) + \frac{ix^4}{2b} + \frac{2i(a+i)^4 \log(a+bx+i)}{b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**2/(1+(b*x+a)**2)*x**4,x)

[Out] -x**5/5 - x**3*(2*I*a/(3*b**2) - 2/(3*b**2)) - x**2*(-I*a**2/b**3 + 2*a/b**3 + I/b**3) - x*(2*I*a**3/b**4 - 6*a**2/b**4 - 6*I*a/b**4 + 2/b**4) + I*x**4/(2*b) + 2*I*(a + I)**4*log(a + b*x + I)/b**5

Giac [A]

time = 0.43, size = 123, normalized size = 1.34

$$\frac{2(-ia^4 + 4a^3 + 6ia^2 - 4a - i)\log(bx + a + i)}{b^5} - \frac{6b^5x^5 - 15ib^4x^4 + 20iab^3x^3 - 30ia^2b^2x^2 - 20b^3x^3 + 60ia^3bx + 60ab^2x^2 - 180a^2bx + 30ib^2x^2 - 180iabx + 60bx}{30b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^4,x, algorithm="giac")

[Out] -2*(-I*a^4 + 4*a^3 + 6*I*a^2 - 4*a - I)*log(b*x + a + I)/b^5 - 1/30*(6*b^5*x^5 - 15*I*b^4*x^4 + 20*I*a*b^3*x^3 - 30*I*a^2*b^2*x^2 - 20*b^3*x^3 + 60*I*a^3*b*x + 60*a*b^2*x^2 - 180*a^2*b*x + 30*I*b^2*x^2 - 180*I*a*b*x + 60*b*x)/b^5

Mupad [B]

time = 0.58, size = 201, normalized size = 2.18

$$\ln\left(x + \frac{a+1i}{b}\right) \left(\frac{8a-8a^3}{b^5} + \frac{(2a^4-12a^2+2)1i}{b^5}\right) - x^4 \left(\frac{(-1+a1i)1i}{4b} - \frac{(1+a1i)1i}{4b}\right) - \frac{x^5}{5} + \frac{x^2(-1+a1i)^2 \left(\frac{(-1+a1i)1i}{2b^2} - \frac{(1+a1i)1i}{b^5}\right)}{2b^2} - \frac{x^3(-1+a1i) \left(\frac{(-1+a1i)1i}{3b} - \frac{(1+a1i)1i}{b^5}\right)}{3b} + \frac{x(-1+a1i)^3 \left(\frac{(-1+a1i)1i}{b^3} - \frac{(1+a1i)1i}{b^5}\right)}{b^3} 1i$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*(a*1i + b*x*1i + 1)^2)/((a + b*x)^2 + 1),x)

[Out] log(x + (a + 1i)/b)*((8*a - 8*a^3)/b^5 + ((2*a^4 - 12*a^2 + 2)*1i)/b^5) - x^4*(((a*1i - 1)*1i)/(4*b) - ((a*1i + 1)*1i)/(4*b)) - x^5/5 + (x^2*(a*1i - 1)^2*(((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b))/(2*b^2) - (x^3*(a*1i - 1)*((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b)*1i)/(3*b) + (x*(a*1i - 1)^3*(((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b)*1i)/b^3

3.172 $\int e^{2i\text{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=72

$$\frac{2i(i+a)^2x}{b^3} + \frac{(1-ia)x^2}{b^2} + \frac{2ix^3}{3b} - \frac{x^4}{4} - \frac{2(1-ia)^3 \log(i+a+bx)}{b^4}$$

[Out] $2*I*(I+a)^2*x/b^3+(1-I*a)*x^2/b^2+2/3*I*x^3/b-1/4*x^4-2*(1-I*a)^3*\ln(I+a+b*x)/b^4$

Rubi [A]

time = 0.04, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2(1-ia)^3 \log(a+bx+i)}{b^4} + \frac{2i(a+i)^2x}{b^3} + \frac{(1-ia)x^2}{b^2} + \frac{2ix^3}{3b} - \frac{x^4}{4}$$

Antiderivative was successfully verified.

[In] Int[E^((2*I)*ArcTan[a + b*x])*x^3,x]

[Out] $((2*I)*(I+a)^2*x)/b^3 + ((1-I*a)*x^2)/b^2 + (((2*I)/3)*x^3)/b - x^4/4 - (2*(1-I*a)^3*\text{Log}[I+a+b*x])/b^4$

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(a+bx)} x^3 dx &= \int \frac{x^3(1+ia+ibx)}{1-ia-ibx} dx \\ &= \int \left(\frac{2i(i+a)^2}{b^3} + \frac{2(1-ia)x}{b^2} + \frac{2ix^2}{b} - x^3 + \frac{2(-1+ia)^3}{b^3(i+a+bx)} \right) dx \\ &= \frac{2i(i+a)^2x}{b^3} + \frac{(1-ia)x^2}{b^2} + \frac{2ix^3}{3b} - \frac{x^4}{4} - \frac{2(1-ia)^3 \log(i+a+bx)}{b^4} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 72, normalized size = 1.00

$$\frac{2i(i+a)^2x}{b^3} + \frac{(1-ia)x^2}{b^2} + \frac{2ix^3}{3b} - \frac{x^4}{4} - \frac{2(1-ia)^3 \log(i+a+bx)}{b^4}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a + b*x])*x^3,x]

[Out] ((2*I)*(I + a)^2*x)/b^3 + ((1 - I*a)*x^2)/b^2 + (((2*I)/3)*x^3)/b - x^4/4 - (2*(1 - I*a)^3*Log[I + a + b*x])/b^4

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(62) = 124.

time = 0.13, size = 170, normalized size = 2.36

method	result
default	$\frac{i(\frac{1}{4}ib^3x^4 + \frac{2}{3}b^2x^3 - ibx^2 - abx + 2a^2x - 2x)}{b^3} + \frac{(-2ia^3b + 6a^2b + 6iab - 2b) \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2} + \frac{(-2ia^4 + 4a^3 + 2i + 4a - \frac{(-2ia^3b + 6a^2b + 6iab - 2b)}{b^3})}{b^3}$
risch	$-\frac{x^4}{4} + \frac{2ix^3}{3b} + \frac{x^2}{b^2} - \frac{iax^2}{b^2} - \frac{4ax}{b^3} + \frac{2ia^2x}{b^3} - \frac{2ix}{b^3} + \frac{3 \ln(b^2x^2 + 2abx + a^2 + 1)a^2}{b^4} - \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)a^3}{b^4} - \frac{\ln(b^2x^2 + 2abx + a^2 + 1)}{b^4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^3,x,method=_RETURNVERBOSE)

[Out] I/b^3*(1/4*I*b^3*x^4+2/3*b^2*x^3-I*b*x^2-a*b*x^2+4*I*a*x+2*a^2*x-2*x)+1/b^3*(1/2*(-2*I*a^3*b+6*I*a*b+6*a^2*b-2*b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(-2*I*a^4+4*a^3+2*I+4*a-(-2*I*a^3*b+6*I*a*b+6*a^2*b-2*b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b))

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 116 vs. 2(56) = 112.

time = 0.47, size = 116, normalized size = 1.61

$$\frac{3b^3x^4 - 8ib^2x^3 + 12(ia - 1)bx^2 + 24(-ia^2 + 2a + i)x}{12b^3} - \frac{2(a^3 + 3ia^2 - 3a - i) \arctan\left(\frac{b^2x + ab}{b}\right)}{b^4} + \frac{(-ia^3 + 3a^2 + 3ia - 1) \log(b^2x^2 + 2abx + a^2 + 1)}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^3,x, algorithm="maxima")

[Out] -1/12*(3*b^3*x^4 - 8*I*b^2*x^3 + 12*(I*a - 1)*b*x^2 + 24*(-I*a^2 + 2*a + I)*x)/b^3 - 2*(a^3 + 3*I*a^2 - 3*a - I)*arctan((b^2*x + a*b)/b)/b^4 + (-I*a^3 + 3*a^2 + 3*I*a - 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^4

Fricas [A]

time = 2.33, size = 77, normalized size = 1.07

$$\frac{3b^4x^4 - 8ib^3x^3 + 12(ia - 1)b^2x^2 + 24(-ia^2 + 2a + i)bx + 24(ia^3 - 3a^2 - 3ia + 1) \log\left(\frac{bx+a+i}{b}\right)}{12b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^3,x, algorithm="fricas")

[Out] -1/12*(3*b^4*x^4 - 8*I*b^3*x^3 + 12*(I*a - 1)*b^2*x^2 + 24*(-I*a^2 + 2*a + I)*b*x + 24*(I*a^3 - 3*a^2 - 3*I*a + 1)*log((b*x + a + I)/b))/b^4

Sympy [A]

time = 0.19, size = 75, normalized size = 1.04

$$-\frac{x^4}{4} - x^2 \left(\frac{ia}{b^2} - \frac{1}{b^2} \right) - x \left(-\frac{2ia^2}{b^3} + \frac{4a}{b^3} + \frac{2i}{b^3} \right) + \frac{2ix^3}{3b} - \frac{2i(a+i)^3 \log(a+bx+i)}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a)**2/(1+(b*x+a)**2)*x**3,x)

[Out] -x**4/4 - x**2*(I*a/b**2 - 1/b**2) - x*(-2*I*a**2/b**3 + 4*a/b**3 + 2*I/b**3) + 2*I*x**3/(3*b) - 2*I*(a + I)**3*log(a + b*x + I)/b**4

Giac [A]

time = 0.45, size = 83, normalized size = 1.15

$$\frac{2(i a^3 - 3 a^2 - 3 i a + 1) \log(b x + a + i)}{b^4} - \frac{3 b^4 x^4 - 8 i b^3 x^3 + 12 i a b^2 x^2 - 24 i a^2 b x - 12 b^2 x^2 + 48 a b x + 24 i b x}{12 b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^3,x, algorithm="giac")

[Out] -2*(I*a^3 - 3*a^2 - 3*I*a + 1)*log(b*x + a + I)/b^4 - 1/12*(3*b^4*x^4 - 8*I*b^3*x^3 + 12*I*a*b^2*x^2 - 24*I*a^2*b*x - 12*b^2*x^2 + 48*a*b*x + 24*I*b*x)/b^4

Mupad [B]

time = 0.53, size = 153, normalized size = 2.12

$$-x^3 \left(\frac{(-1+a \operatorname{li}) \operatorname{li}}{3b} - \frac{(1+a \operatorname{li}) \operatorname{li}}{3b} \right) - \frac{x^4}{4} + \ln \left(x + \frac{a+1i}{b} \right) \left(\frac{6a^2-2}{b^4} + \frac{(6a-2a^3) \operatorname{li}}{b^4} \right) - \frac{x^2(-1+a \operatorname{li}) \left(\frac{(-1+a \operatorname{li}) \operatorname{li}}{b} - \frac{(1+a \operatorname{li}) \operatorname{li}}{b} \right) \operatorname{li}}{2b} + \frac{x(-1+a \operatorname{li})^2 \left(\frac{(-1+a \operatorname{li}) \operatorname{li}}{b} - \frac{(1+a \operatorname{li}) \operatorname{li}}{b} \right)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*(a*1i + b*x*1i + 1)^2)/((a + b*x)^2 + 1),x)

[Out] log(x + (a + 1i)/b)*(((6*a - 2*a^3)*1i)/b^4 + (6*a^2 - 2)/b^4) - x^4/4 - x^3*(((a*1i - 1)*1i)/(3*b) - ((a*1i + 1)*1i)/(3*b)) - (x^2*(a*1i - 1)*(((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b)*1i)/(2*b) + (x*(a*1i - 1)^2*(((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b))/b^2

3.173 $\int e^{2i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=54

$$\frac{2(1-ia)x}{b^2} + \frac{ix^2}{b} - \frac{x^3}{3} + \frac{2i(i+a)^2 \log(i+a+bx)}{b^3}$$

[Out] $2*(1-I*a)*x/b^2+I*x^2/b-1/3*x^3+2*I*(I+a)^2*\ln(I+a+bx)/b^3$

Rubi [A]

time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$\frac{2i(a+i)^2 \log(a+bx+i)}{b^3} + \frac{2(1-ia)x}{b^2} + \frac{ix^2}{b} - \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] Int[E^((2*I)*ArcTan[a + b*x])*x^2,x]

[Out] $(2*(1 - I*a)*x)/b^2 + (I*x^2)/b - x^3/3 + ((2*I)*(I + a)^2*\text{Log}[I + a + b*x])/b^3$

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2(1+ia+ibx)}{1-ia-ibx} dx \\ &= \int \left(-\frac{2i(i+a)}{b^2} + \frac{2ix}{b} - x^2 + \frac{2i(i+a)^2}{b^2(i+a+bx)} \right) dx \\ &= \frac{2(1-ia)x}{b^2} + \frac{ix^2}{b} - \frac{x^3}{3} + \frac{2i(i+a)^2 \log(i+a+bx)}{b^3} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 54, normalized size = 1.00

$$\frac{2(1-ia)x}{b^2} + \frac{ix^2}{b} - \frac{x^3}{3} + \frac{2i(i+a)^2 \log(i+a+bx)}{b^3}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a + b*x])*x^2,x]`

`[Out] (2*(1 - I*a)*x)/b^2 + (I*x^2)/b - x^3/3 + ((2*I)*(I + a)^2*Log[I + a + b*x])/b^3`

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(47) = 94.

time = 0.11, size = 135, normalized size = 2.50

method	result
default	$\frac{i(\frac{1}{3}ib^2x^3+x^2b-2ix-2ax)}{b^2} + \frac{(2ia^2b-4ab-2ib)\ln(b^2x^2+2abx+a^2+1)}{2b^2} + \frac{\left(2ia^3+2ia-2a^2-2-\frac{(2ia^2b-4ab-2ib)a}{b}\right)\arctan\left(\frac{2b^2x+2ab}{2b}\right)}{b^2}$
risch	$-\frac{x^3}{3} + \frac{ix^2}{b} + \frac{2x}{b^2} - \frac{2iax}{b^2} - \frac{2\ln(b^2x^2+2abx+a^2+1)a}{b^3} + \frac{i\ln(b^2x^2+2abx+a^2+1)a^2}{b^3} - \frac{i\ln(b^2x^2+2abx+a^2+1)}{b^3} + \frac{4i\arctan\left(\frac{2b^2x+2ab}{2b}\right)}{b^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^2,x,method=_RETURNVERBOSE)`

`[Out] I/b^2*(1/3*I*b^2*x^3+x^2*b-2*I*x-2*a*x)+1/b^2*(1/2*(2*I*a^2*b-2*I*b-4*a*b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(2*I*a^3+2*I*a-2*a^2-2-(2*I*a^2*b-2*I*b-4*a*b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b)`

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 87 vs. 2(42) = 84.

time = 0.46, size = 87, normalized size = 1.61

$$-\frac{b^2x^3 - 3ibx^2 + 6(ia - 1)x}{3b^2} + \frac{2(a^2 + 2ia - 1)\arctan\left(\frac{b^2x+ab}{b}\right)}{b^3} + \frac{(ia^2 - 2a - i)\log(b^2x^2 + 2abx + a^2 + 1)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^2,x, algorithm="maxima")`

`[Out] -1/3*(b^2*x^3 - 3*I*b*x^2 + 6*(I*a - 1)*x)/b^2 + 2*(a^2 + 2*I*a - 1)*arctan((b^2*x + a*b)/b)/b^3 + (I*a^2 - 2*a - I)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^3`

Fricas [A]

time = 2.34, size = 53, normalized size = 0.98

$$-\frac{b^3x^3 - 3ib^2x^2 + 6(ia - 1)bx + 6(-ia^2 + 2a + i)\log\left(\frac{bx+a+i}{b}\right)}{3b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^2,x, algorithm="fricas")

[Out] $-1/3*(b^3*x^3 - 3*I*b^2*x^2 + 6*(I*a - 1)*b*x + 6*(-I*a^2 + 2*a + I)*\log((b*x + a + I)/b))/b^3$

Sympy [A]

time = 0.15, size = 46, normalized size = 0.85

$$-\frac{x^3}{3} - x \left(\frac{2ia}{b^2} - \frac{2}{b^2} \right) + \frac{ix^2}{b} + \frac{2i(a+i)^2 \log(a+bx+i)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a)**2/(1+(b*x+a)**2)*x**2,x)

[Out] $-x**3/3 - x*(2*I*a/b**2 - 2/b**2) + I*x**2/b + 2*I*(a + I)**2*\log(a + b*x + I)/b**3$

Giac [A]

time = 0.45, size = 53, normalized size = 0.98

$$-\frac{2(-ia^2 + 2a + i) \log(bx + a + i)}{b^3} - \frac{b^3x^3 - 3ib^2x^2 + 6i abx - 6bx}{3b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x^2,x, algorithm="giac")

[Out] $-2*(-I*a^2 + 2*a + I)*\log(b*x + a + I)/b^3 - 1/3*(b^3*x^3 - 3*I*b^2*x^2 + 6*I*a*b*x - 6*b*x)/b^3$

Mupad [B]

time = 0.51, size = 107, normalized size = 1.98

$$-\ln\left(x + \frac{a+1i}{b}\right) \left(\frac{4a}{b^3} - \frac{(2a^2-2)1i}{b^3}\right) - x^2 \left(\frac{(-1+a1i)1i}{2b} - \frac{(1+a1i)1i}{2b}\right) - \frac{x^3}{3} - \frac{x(-1+a1i) \left(\frac{(-1+a1i)1i}{b} - \frac{(1+a1i)1i}{b}\right) 1i}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*1i + b*x*1i + 1)^2)/((a + b*x)^2 + 1),x)

[Out] $-\log(x + (a + 1i)/b)*((4*a)/b^3 - ((2*a^2 - 2)*1i)/b^3) - x^2*((a*1i - 1)*1i)/(2*b) - ((a*1i + 1)*1i)/(2*b)) - x^3/3 - (x*(a*1i - 1)*((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b)*1i)/b$

3.174 $\int e^{2i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=37

$$\frac{2ix}{b} - \frac{x^2}{2} + \frac{2(1-ia)\log(i+a+bx)}{b^2}$$

[Out] $2*I*x/b - 1/2*x^2 + 2*(1-I*a)*\ln(I+a+b*x)/b^2$

Rubi [A]

time = 0.02, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5203, 78}

$$\frac{2(1-ia)\log(a+bx+i)}{b^2} + \frac{2ix}{b} - \frac{x^2}{2}$$

Antiderivative was successfully verified.

[In] Int[E^((2*I)*ArcTan[a + b*x])*x,x]

[Out] ((2*I)*x)/b - x^2/2 + (2*(1 - I*a)*Log[I + a + b*x])/b^2

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(a+bx)} x dx &= \int \frac{x(1+ia+ibx)}{1-ia-ibx} dx \\ &= \int \left(\frac{2i}{b} - x + \frac{2(1-ia)}{b(i+a+bx)} \right) dx \\ &= \frac{2ix}{b} - \frac{x^2}{2} + \frac{2(1-ia)\log(i+a+bx)}{b^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 37, normalized size = 1.00

$$\frac{2ix}{b} - \frac{x^2}{2} + \frac{2(1-ia)\log(i+a+bx)}{b^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a + b*x])*x,x]

[Out] ((2*I)*x)/b - x^2/2 + (2*(1 - I*a)*Log[I + a + b*x])/b^2

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 98 vs. 2(32) = 64.

time = 0.10, size = 99, normalized size = 2.68

method	result	size
risch	$-\frac{x^2}{2} + \frac{2ix}{b} + \frac{\ln(b^2x^2+2abx+a^2+1)}{b^2} - \frac{2i\arctan(bx+a)}{b^2} - \frac{ia\ln(b^2x^2+2abx+a^2+1)}{b^2} - \frac{2a\arctan(bx+a)}{b^2}$	85
default	$\frac{-\frac{1}{2}x^2b+2ix}{b} + \frac{(-2iab+2b)\ln(b^2x^2+2abx+a^2+1)}{2b^2} + \frac{\left(-2ia^2-2i-\frac{(-2iab+2b)a}{b}\right)\arctan\left(\frac{2b^2x+2ab}{2b}\right)}{b}$	99

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x,x,method=_RETURNVERBOSE)

[Out] 1/b*(-1/2*x^2*b+2*I*x)+1/b*(1/2*(-2*I*a*b+2*b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(-2*I*a^2-2*I-(-2*I*a*b+2*b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b))

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 64 vs. 2(29) = 58.

time = 0.46, size = 64, normalized size = 1.73

$$\frac{bx^2 - 4ix}{2b} - \frac{2(a+i)\arctan\left(\frac{b^2x+ab}{b}\right)}{b^2} + \frac{(-ia+1)\log(b^2x^2+2abx+a^2+1)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x,x, algorithm="maxima")

[Out] -1/2*(b*x^2 - 4*I*x)/b - 2*(a + I)*arctan((b^2*x + a*b)/b)/b^2 + (-I*a + 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^2

Fricas [A]

time = 1.59, size = 35, normalized size = 0.95

$$\frac{b^2x^2 - 4ibx + 4(ia - 1)\log\left(\frac{bx+a+i}{b}\right)}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x,x, algorithm="fricas")

[Out] -1/2*(b^2*x^2 - 4*I*b*x + 4*(I*a - 1)*log((b*x + a + I)/b))/b^2

Sympy [A]

time = 0.10, size = 29, normalized size = 0.78

$$-\frac{x^2}{2} + \frac{2ix}{b} - \frac{2i(a+i)\log(a+bx+i)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x,x)

[Out] -x**2/2 + 2*I*x/b - 2*I*(a + I)*log(a + b*x + I)/b**2

Giac [A]

time = 0.42, size = 35, normalized size = 0.95

$$-\frac{2(i a - 1)\log(bx + a + i)}{b^2} - \frac{b^2 x^2 - 4i b x}{2 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)*x,x, algorithm="giac")

[Out] -2*(I*a - 1)*log(b*x + a + I)/b^2 - 1/2*(b^2*x^2 - 4*I*b*x)/b^2

Mupad [B]

time = 0.13, size = 60, normalized size = 1.62

$$-\ln\left(x + \frac{a + 1i}{b}\right) \left(-\frac{2}{b^2} + \frac{a 2i}{b^2}\right) - x \left(\frac{(-1 + a 1i) 1i}{b} - \frac{(1 + a 1i) 1i}{b}\right) - \frac{x^2}{2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*(a*1i + b*x*1i + 1)^2)/((a + b*x)^2 + 1),x)

[Out] -log(x + (a + 1i)/b)*((a*2i)/b^2 - 2/b^2) - x*((a*1i - 1)*1i)/b - ((a*1i + 1)*1i)/b) - x^2/2

3.175 $\int e^{2i \operatorname{ArcTan}(a+bx)} dx$

Optimal. Leaf size=20

$$-x + \frac{2i \log(i + a + bx)}{b}$$

[Out] $-x+2*I*\ln(I+a+b*x)/b$

Rubi [A]

time = 0.01, antiderivative size = 20, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5201, 45}

$$-x + \frac{2i \log(a + bx + i)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $-x + ((2*I)*\text{Log}[I + a + b*x])/b$

Rule 45

$\text{Int}[(a_. + (b_.)(x_.))^{(m_.)}((c_.) + (d_.)(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x] /; \text{FreeQ}\{a, b, c, d, n, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] \mid\mid (\text{EqQ}[c, 0] \&\& \text{LeQ}[7*m + 4*n + 4, 0]) \mid\mid \text{LtQ}[9*m + 5*(n + 1), 0] \mid\mid \text{GtQ}[m + n + 2, 0])]$

Rule 5201

$\text{Int}[E^{(\text{ArcTan}[(c_.)*((a_.) + (b_.)(x_.)))]*(n_.)}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*c - I*b*c*x)^{I*(n/2)} / (1 + I*a*c + I*b*c*x)^{I*(n/2)}, x] /; \text{FreeQ}\{a, b, c, n, x\}$

Rubi steps

$$\begin{aligned} \int e^{2i \tan^{-1}(a+bx)} dx &= \int \frac{1 + ia + ibx}{1 - ia - ibx} dx \\ &= \int \left(-1 + \frac{2i}{i + a + bx} \right) dx \\ &= -x + \frac{2i \log(i + a + bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 32, normalized size = 1.60

$$-x + \frac{2\text{ArcTan}(a + bx)}{b} + \frac{i \log(1 + (a + bx)^2)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a + b*x]),x]

[Out] -x + (2*ArcTan[a + b*x])/b + (I*Log[1 + (a + b*x)^2])/b

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 50 vs. 2(18) = 36.

time = 0.10, size = 51, normalized size = 2.55

method	result	size
risch	$-x + \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)}{b} + \frac{2 \arctan(bx + a)}{b}$	40
default	$-x + \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)}{b} + \frac{2 \arctan\left(\frac{2b^2x + 2ab}{2b}\right)}{b}$	51

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2),x,method=_RETURNVERBOSE)

[Out] -x+I/b*ln(b^2*x^2+2*a*b*x+a^2+1)+2/b*arctan(1/2*(2*b^2*x+2*a*b)/b)

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 46 vs. 2(16) = 32.

time = 0.48, size = 46, normalized size = 2.30

$$-x + \frac{2 \arctan\left(\frac{b^2x + ab}{b}\right)}{b} + \frac{i \log(b^2x^2 + 2abx + a^2 + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2),x, algorithm="maxima")

[Out] -x + 2*arctan((b^2*x + a*b)/b)/b + I*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/b

Fricas [A]

time = 1.77, size = 22, normalized size = 1.10

$$\frac{bx - 2i \log\left(\frac{bx + a + i}{b}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2),x, algorithm="fricas")

[Out] $-(b*x - 2*I*\log((b*x + a + I)/b))/b$

Sympy [A]

time = 0.07, size = 14, normalized size = 0.70

$$-x + \frac{2i \log(a + bx + i)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))**2/(1+(b*x+a)**2),x)`

[Out] $-x + 2*I*\log(a + b*x + I)/b$

Giac [A]

time = 0.42, size = 16, normalized size = 0.80

$$-x + \frac{2i \log(bx + a + i)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2),x, algorithm="giac")`

[Out] $-x + 2*I*\log(b*x + a + I)/b$

Mupad [B]

time = 0.46, size = 21, normalized size = 1.05

$$-x + \frac{\ln\left(x + \frac{a+1i}{b}\right) 2i}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*1i + b*x*1i + 1)^2/((a + b*x)^2 + 1),x)`

[Out] $(\log(x + (a + 1i)/b)*2i)/b - x$

$$3.176 \quad \int \frac{e^{2i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=38

$$\frac{(i-a) \log(x)}{i+a} - \frac{2 \log(i+a+bx)}{1-ia}$$

[Out] (I-a)*ln(x)/(I+a)-2*ln(I+a+b*x)/(1-I*a)

Rubi [A]

time = 0.02, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$\frac{(-a+i) \log(x)}{a+i} - \frac{2 \log(a+bx+i)}{1-ia}$$

Antiderivative was successfully verified.

[In] Int[E^((2*I)*ArcTan[a + b*x])/x,x]

[Out] ((I - a)*Log[x])/(I + a) - (2*Log[I + a + b*x])/(1 - I*a)

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{1 + ia + ibx}{x(1 - ia - ibx)} dx \\ &= \int \left(\frac{i-a}{(i+a)x} - \frac{2ib}{(i+a)(i+a+bx)} \right) dx \\ &= \frac{(i-a) \log(x)}{i+a} - \frac{2 \log(i+a+bx)}{1-ia} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 31, normalized size = 0.82

$$\frac{(-i + a) \log(x) + 2i \log(i + a + bx)}{i + a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a + b*x])/x,x]**[Out]** -(((-I + a)*Log[x] + (2*I)*Log[I + a + b*x])/(I + a))**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 109 vs. 2(34) = 68.

time = 0.11, size = 110, normalized size = 2.89

method	result	size
risch	$\frac{i \ln(-x)}{i+a} - \frac{\ln(-x)a}{i+a} - \frac{i \ln(b^2x^2+2abx+a^2+1)}{i+a} - \frac{2 \arctan(bx+a)}{i+a}$	69
default	$-\frac{2b \left(\frac{(iab+b) \ln(b^2x^2+2abx+a^2+1)}{2b^2} + \frac{(ia^2-i+2a - \frac{(iab+b)a}{b}) \arctan(\frac{2b^2x+2ab}{2b})}{b} \right)}{a^2+1} + \frac{(-a^2+2ia+1) \ln(x)}{a^2+1}$	110

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x,x,method=_RETURNVERBOSE)**[Out]** -2*b/(a^2+1)*(1/2*(I*b*a+b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(I*a^2-I+2*a-(I*b*a+b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b)+(2*I*a-a^2+1)/(a^2+1)*ln(x)**Maxima [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 78 vs. 2(29) = 58.

time = 0.47, size = 78, normalized size = 2.05

$$-\frac{2(a-i) \arctan\left(\frac{b^2x+ab}{b}\right)}{a^2+1} - \frac{(ia+1) \log(b^2x^2+2abx+a^2+1)}{a^2+1} - \frac{(a^2-2ia-1) \log(x)}{a^2+1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x,x, algorithm="maxima")**[Out]** -2*(a - I)*arctan((b^2*x + a*b)/b)/(a^2 + 1) - (I*a + 1)*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^2 + 1) - (a^2 - 2*I*a - 1)*log(x)/(a^2 + 1)**Fricas [A]**

time = 2.66, size = 27, normalized size = 0.71

$$\frac{(a - i) \log(x) + 2i \log\left(\frac{bx+a+i}{b}\right)}{a + i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x,x, algorithm="fricas")

[Out] -((a - I)*log(x) + 2*I*log((b*x + a + I)/b))/(a + I)

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 100 vs. $2(24) = 48$.

time = 0.45, size = 100, normalized size = 2.63

$$\frac{(a-i)\log\left(-\frac{a^2(a-i)}{a+i} + a^2 - \frac{2ia(a-i)}{a+i} + x(ab-3ib) + \frac{a-i}{a+i} + 1\right)}{a+i} - \frac{2i\log\left(a^2 - \frac{2ia^2}{a+i} + \frac{4a}{a+i} + x(ab-3ib) + 1 + \frac{2i}{a+i}\right)}{a+i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**2/(1+(b*x+a)**2)/x,x)

[Out] -(a - I)*log(-a**2*(a - I)/(a + I) + a**2 - 2*I*a*(a - I)/(a + I) + x*(a*b - 3*I*b) + (a - I)/(a + I) + 1)/(a + I) - 2*I*log(a**2 - 2*I*a**2/(a + I) + 4*a/(a + I) + x*(a*b - 3*I*b) + 1 + 2*I/(a + I))/(a + I)

Giac [A]

time = 0.46, size = 33, normalized size = 0.87

$$-\frac{2ib\log(bx+a+i)}{ab+ib} - \frac{(a-i)\log(|x|)}{a+i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x,x, algorithm="giac")

[Out] -2*I*b*log(b*x + a + I)/(a*b + I*b) - (a - I)*log(abs(x))/(a + I)

Mupad [B]

time = 0.70, size = 32, normalized size = 0.84

$$\ln(x) \left(-1 + \frac{2i}{a+1i}\right) - \frac{\ln(a+bx+1i) 2i}{a+1i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)^2/(x*((a + b*x)^2 + 1)),x)

[Out] log(x)*(2i/(a + 1i) - 1) - (log(a + b*x + 1i)*2i)/(a + 1i)

$$3.177 \quad \int \frac{e^{2i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=55

$$-\frac{i-a}{(i+a)x} - \frac{2ib \log(x)}{(i+a)^2} + \frac{2ib \log(i+a+bx)}{(i+a)^2}$$

[Out] $(-I+a)/(I+a)/x-2*I*b*\ln(x)/(I+a)^2+2*I*b*\ln(I+a+b*x)/(I+a)^2$

Rubi [A]

time = 0.03, antiderivative size = 55, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2ib \log(x)}{(a+i)^2} + \frac{2ib \log(a+bx+i)}{(a+i)^2} - \frac{-a+i}{(a+i)x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a + b*x])/x^2}, x]$

[Out] $-((I - a)/((I + a)*x)) - ((2*I)*b*\text{Log}[x])/(I + a)^2 + ((2*I)*b*\text{Log}[I + a + b*x])/(I + a)^2$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}\{a, b, c, d, e, f\}))))$

Rule 5203

$\text{Int}[E^{(\text{ArcTan}[(c_.)*((a_. + (b_.)*(x_.))]*(n_.))*((d_. + (e_.)*(x_.))^{(m_.)}, x_Symbol] :> \text{Int}[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^{(I*(n/2))}/(1 + I*a*c + I*b*c*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{1 + ia + ibx}{x^2(1 - ia - ibx)} dx \\ &= \int \left(\frac{i-a}{(i+a)x^2} - \frac{2ib}{(i+a)^2 x} + \frac{2ib^2}{(i+a)^2(i+a+bx)} \right) dx \\ &= -\frac{i-a}{(i+a)x} - \frac{2ib \log(x)}{(i+a)^2} + \frac{2ib \log(i+a+bx)}{(i+a)^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 39, normalized size = 0.71

$$\frac{1 + a^2 - 2ibx \log(x) + 2ibx \log(i + a + bx)}{(i + a)^2 x}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a + b*x])/x^2,x]``[Out] (1 + a^2 - (2*I)*b*x*Log[x] + (2*I)*b*x*Log[I + a + b*x])/((I + a)^2*x)`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 160 vs. 2(45) = 90.

time = 0.13, size = 161, normalized size = 2.93

method	result
default	$2b^2 \frac{\left(\frac{(ia^2b+2ab-ib) \ln(b^2x^2+2abx+a^2+1)}{2b^2} + \frac{\left(ia^3-3ia+3a^2-1 - \frac{(ia^2b+2ab-ib)a}{b} \right) \arctan\left(\frac{2b^2x+2ab}{2b} \right)}{b} \right)}{(a^2+1)^2} - \frac{-a^2+2ia+1}{(a^2+1)x} - \frac{2b(ia^2+2a-1)}{(a^2+1)}$
risch	$\frac{a}{(i+a)x} - \frac{i}{(i+a)x} - \frac{b \ln(4a^4b^2x^2+8a^5bx+4a^6+8a^2b^2x^2+16a^3bx+12a^4+4b^2x^2+8abx+12a^2+4)}{ia^2-2a-i} + \frac{2ib \arctan\left(\frac{(2a^2b+2b)x+2a^3+2a^2}{2a^2+2} \right)}{ia^2-2a-i}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^2,x,method=_RETURNVERBOSE)`
`[Out] 2*b^2/(a^2+1)^2*(1/2*(I*b*a^2-I*b+2*a*b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(I*a^3-3*I*a+3*a^2-1-(I*b*a^2-I*b+2*a*b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b))-`
`(2*I*a-a^2+1)/(a^2+1)/x-2*b*(I*a^2-I+2*a)/(a^2+1)^2*ln(x)`
Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 126 vs. 2(38) = 76.

time = 0.48, size = 126, normalized size = 2.29

$$\frac{2(a^2 - 2ia - 1)b \arctan\left(\frac{b^2x+ab}{b}\right)}{a^4 + 2a^2 + 1} + \frac{(ia^2 + 2a - i)b \log(b^2x^2 + 2abx + a^2 + 1)}{a^4 + 2a^2 + 1} - \frac{2(ia^2 + 2a - i)b \log(x)}{a^4 + 2a^2 + 1} + \frac{a^2 - 2ia - 1}{(a^2 + 1)x}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^2,x, algorithm="maxima")`
`[Out] 2*(a^2 - 2*I*a - 1)*b*arctan((b^2*x + a*b)/b)/(a^4 + 2*a^2 + 1) + (I*a^2 +`
`2*a - I)*b*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^4 + 2*a^2 + 1) - 2*(I*a^2 +`
`2*a - I)*b*log(x)/(a^4 + 2*a^2 + 1) + (a^2 - 2*I*a - 1)/((a^2 + 1)*x)`

Fricas [A]

time = 2.12, size = 40, normalized size = 0.73

$$\frac{-2i bx \log(x) + 2i bx \log\left(\frac{bx+a+i}{b}\right) + a^2 + 1}{(a^2 + 2i a - 1)x}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^2,x, algorithm="fricas")``[Out] (-2*I*b*x*log(x) + 2*I*b*x*log((b*x + a + I)/b) + a^2 + 1)/((a^2 + 2*I*a - 1)*x)`**Sympy [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 156 vs. $2(39) = 78$.

time = 0.34, size = 156, normalized size = 2.84

$$-\frac{2ib \log\left(-\frac{2a^3b}{(a+i)^2} - \frac{6ia^2b}{(a+i)^2} + 2ab + \frac{6ab}{(a+i)^2} + 4b^2x + 2ib + \frac{2ib}{(a+i)^2}\right)}{(a+i)^2} + \frac{2ib \log\left(\frac{2a^3b}{(a+i)^2} + \frac{6ia^2b}{(a+i)^2} + 2ab - \frac{6ab}{(a+i)^2} + 4b^2x + 2ib - \frac{2ib}{(a+i)^2}\right)}{(a+i)^2} - \frac{-a+i}{x(a+i)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*(b*x+a)**2/(1+(b*x+a)**2)/x**2,x``[Out] -2*I*b*log(-2*a**3*b/(a + I)**2 - 6*I*a**2*b/(a + I)**2 + 2*a*b + 6*a*b/(a + I)**2 + 4*b**2*x + 2*I*b + 2*I*b/(a + I)**2)/(a + I)**2 + 2*I*b*log(2*a**3*b/(a + I)**2 + 6*I*a**2*b/(a + I)**2 + 2*a*b - 6*a*b/(a + I)**2 + 4*b**2*x + 2*I*b - 2*I*b/(a + I)**2)/(a + I)**2 - (-a + I)/(x*(a + I))`**Giac [A]**

time = 0.45, size = 61, normalized size = 1.11

$$\frac{2b^2 \log(bx + a + i)}{-i a^2 b + 2ab + i b} + \frac{2b \log(|x|)}{i a^2 - 2a - i} + \frac{a^2 + 1}{(a + i)^2 x}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^2,x, algorithm="giac")``[Out] 2*b^2*log(b*x + a + I)/(-I*a^2*b + 2*a*b + I*b) + 2*b*log(abs(x))/(I*a^2 - 2*a - I) + (a^2 + 1)/((a + I)^2*x)`**Mupad [B]**

time = 0.64, size = 98, normalized size = 1.78

$$\frac{a - i}{x(a + 1i)} + \frac{b \operatorname{atanh}\left(\frac{a^2 + a 2i - 1}{(a + 1i)^2} - \frac{x(2a^4 b^2 + 4a^2 b^2 + 2b^2)}{(a + 1i)^2(-ba^3 + 1i b a^2 - ba + b 1i)}\right)}{(a + 1i)^2} 4i$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a*1i + b*x*1i + 1)^2/(x^2*((a + b*x)^2 + 1)),x)``[Out] (a - 1i)/(x*(a + 1i)) + (b*atanh((a*2i + a^2 - 1)/(a + 1i)^2 - (x*(2*b^2 + 4*a^2*b^2 + 2*a^4*b^2))/((a + 1i)^2*(b*1i - a*b + a^2*b*1i - a^3*b)))*4i)/(a + 1i)^2`

$$3.178 \quad \int \frac{e^{2i \operatorname{ArcTan}(a+bx)}}{x^3} dx$$

Optimal. Leaf size=76

$$-\frac{i-a}{2(i+a)x^2} + \frac{2ib}{(i+a)^2x} - \frac{2b^2 \log(x)}{(1-ia)^3} + \frac{2b^2 \log(i+a+bx)}{(1-ia)^3}$$

[Out] $1/2*(-I+a)/(I+a)/x^2+2*I*b/(I+a)^2/x-2*b^2*\ln(x)/(1-I*a)^3+2*b^2*\ln(I+a+b*x)/(1-I*a)^3$

Rubi [A]

time = 0.04, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2b^2 \log(x)}{(1-ia)^3} + \frac{2b^2 \log(a+bx+i)}{(1-ia)^3} + \frac{2ib}{(a+i)^2x} - \frac{-a+i}{2(a+i)x^2}$$

Antiderivative was successfully verified.

[In] Int[E^((2*I)*ArcTan[a + b*x])/x^3,x]

[Out] $-1/2*(I-a)/((I+a)*x^2) + ((2*I)*b)/((I+a)^2*x) - (2*b^2*\log[x])/(1-I*a)^3 + (2*b^2*\log[I+a+b*x])/(1-I*a)^3$

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(a+bx)}}{x^3} dx &= \int \frac{1 + ia + ibx}{x^3(1 - ia - ibx)} dx \\ &= \int \left(\frac{i - a}{(i + a)x^3} - \frac{2ib}{(i + a)^2 x^2} + \frac{2ib^2}{(i + a)^3 x} - \frac{2ib^3}{(i + a)^3(i + a + bx)} \right) dx \\ &= -\frac{i - a}{2(i + a)x^2} + \frac{2ib}{(i + a)^2 x} - \frac{2b^2 \log(x)}{(1 - ia)^3} + \frac{2b^2 \log(i + a + bx)}{(1 - ia)^3} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 63, normalized size = 0.83

$$\frac{(i + a)(1 + a^2 + 4ibx) + 4ib^2 x^2 \log(x) - 4ib^2 x^2 \log(i + a + bx)}{2(i + a)^3 x^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((2*I)*ArcTan[a + b*x])/x^3,x]

[Out] ((I + a)*(1 + a^2 + (4*I)*b*x) + (4*I)*b^2*x^2*Log[x] - (4*I)*b^2*x^2*Log[I + a + b*x])/(2*(I + a)^3*x^2)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 210 vs. 2(65) = 130.

time = 0.14, size = 211, normalized size = 2.78

method	result
default	$-\frac{2b^3 \left(\frac{(ia^3b + 3a^2b - 3iab - b) \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2} + \frac{(ia^4 - 6ia^2 + 4a^3 + i - 4a - \frac{(ia^3b + 3a^2b - 3iab - b)a}{b}) \arctan\left(\frac{2b^2x + 2ab}{2b}\right)}{b} \right)}{(a^2 + 1)^3} - \frac{-a^2 + 2ia}{2(a^2 + 1)}$
risch	$\frac{\frac{2ibx}{a^2 + 2ia - 1} + \frac{a - i}{2i + 2a}}{x^2} - \frac{2b^2 \ln((2a^4b + 4a^2b + 2b)x)}{ia^3 - 3a^2 - 3ia + 1} + \frac{b^2 \ln(4a^8b^2x^2 + 8a^9bx + 4a^{10} + 16a^6b^2x^2 + 32a^7bx + 20a^8 + 24a^4b^2x^2 + 48a^5bx + 4a^6)}{ia^3 - 3a^2 - 3ia + 1}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^3,x,method=_RETURNVERBOSE)

[Out] -2*b^3/(a^2+1)^3*(1/2*(I*b*a^3-3*I*a*b+3*a^2*b-b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(I*a^4-6*I*a^2+4*a^3+I-4*a-(I*b*a^3-3*I*a*b+3*a^2*b-b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b))-1/2*(2*I*a-a^2+1)/(a^2+1)/x^2+2*b*(I*a^2-I+2*a)/(a^2+1)^2/x+2*b^2*(I*a^3-3*I*a+3*a^2-1)/(a^2+1)^3*ln(x)

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 188 vs. 2(58) = 116.

time = 0.48, size = 188, normalized size = 2.47

$$\frac{2(a^3 - 3ia^2 - 3a + i)b^2 \arctan\left(\frac{b^2x+ab}{b}\right) - (ia^3 + 3a^2 - 3ia - 1)b^2 \log(b^2x^2 + 2abx + a^2 + 1) - \frac{2(-ia^3 - 3a^2 + 3ia + 1)b^2 \log(x)}{a^6 + 3a^4 + 3a^2 + 1} + \frac{a^4 - 2ia^3 - 4(-ia^2 - 2a + i)bx - 2ia - 1}{2(a^4 + 2a^2 + 1)x^2}}{a^6 + 3a^4 + 3a^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^3,x, algorithm="maxima")

[Out] -2*(a^3 - 3*I*a^2 - 3*a + I)*b^2*arctan((b^2*x + a*b)/b)/(a^6 + 3*a^4 + 3*a^2 + 1) - (I*a^3 + 3*a^2 - 3*I*a - 1)*b^2*log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^6 + 3*a^4 + 3*a^2 + 1) - 2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b^2*log(x)/(a^6 + 3*a^4 + 3*a^2 + 1) + 1/2*(a^4 - 2*I*a^3 - 4*(-I*a^2 - 2*a + I)*b*x - 2*I*a - 1)/((a^4 + 2*a^2 + 1)*x^2)

Fricas [A]

time = 2.10, size = 69, normalized size = 0.91

$$\frac{4i b^2 x^2 \log(x) - 4i b^2 x^2 \log\left(\frac{bx+a+i}{b}\right) + a^3 - 4(-ia + 1)bx + ia^2 + a + i}{2(a^3 + 3ia^2 - 3a - i)x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^3,x, algorithm="fricas")

[Out] 1/2*(4*I*b^2*x^2*log(x) - 4*I*b^2*x^2*log((b*x + a + I)/b) + a^3 - 4*(-I*a + 1)*b*x + I*a^2 + a + I)/((a^3 + 3*I*a^2 - 3*a - I)*x^2)

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 228 vs. 2(60) = 120.

time = 0.46, size = 228, normalized size = 3.00

$$\frac{2ib^2 \log\left(\frac{-\frac{2a^4b^2}{(a+i)^3} - \frac{8ia^3b^2}{(a+i)^3} + \frac{12a^2b^2}{(a+i)^3} + 2ab^2 + \frac{8iab^2}{(a+i)^3} + 4b^3x + 2ib^2 - \frac{2b^2}{(a+i)^3}\right) - \frac{2ib^2 \log\left(\frac{2a^4b^2}{(a+i)^3} + \frac{8ia^3b^2}{(a+i)^3} - \frac{12a^2b^2}{(a+i)^3} + 2ab^2 - \frac{8iab^2}{(a+i)^3} + 4b^3x + 2ib^2 + \frac{2b^2}{(a+i)^3}\right)}{(a+i)^3} - \frac{-a^2 - 4ibx - 1}{x^2 \cdot (2a^2 + 4ia - 2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x**3,x)

[Out] 2*I*b**2*log(-2*a**4*b**2/(a + I)**3 - 8*I*a**3*b**2/(a + I)**3 + 12*a**2*b**2/(a + I)**3 + 2*a*b**2 + 8*I*a*b**2/(a + I)**3 + 4*b**3*x + 2*I*b**2 - 2*b**2/(a + I)**3)/(a + I)**3 - 2*I*b**2*log(2*a**4*b**2/(a + I)**3 + 8*I*a**3*b**2/(a + I)**3 - 12*a**2*b**2/(a + I)**3 + 2*a*b**2 - 8*I*a*b**2/(a + I)**3 + 4*b**3*x + 2*I*b**2 + 2*b**2/(a + I)**3)/(a + I)**3 - (-a**2 - 4*I*b*x - 1)/(x**2*(2*a**2 + 4*I*a - 2))

Giac [A]

time = 0.42, size = 89, normalized size = 1.17

$$\frac{2b^3 \log(bx + a + i)}{ia^3b - 3a^2b - 3iab + b} + \frac{2b^2 \log(|x|)}{-ia^3 + 3a^2 + 3ia - 1} + \frac{a^3 + ia^2 + 4i(ab + ib)x + a + i}{2(a+i)^3x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^3,x, algorithm="giac")

[Out] $2*b^3*\log(b*x + a + I)/(I*a^3*b - 3*a^2*b - 3*I*a*b + b) + 2*b^2*\log(\text{abs}(x)) / (-I*a^3 + 3*a^2 + 3*I*a - 1) + 1/2*(a^3 + I*a^2 + 4*I*(a*b + I*b)*x + a + I)/((a + I)^3*x^2)$

Mupad [B]

time = 0.69, size = 154, normalized size = 2.03

$$\frac{\frac{a-i}{2(a+1i)} + \frac{bx2i}{(a+1i)^2}}{x^2} + \frac{b^2 \operatorname{atanh}\left(\frac{-a^3 - a^2 3i + 3a + 1i}{(a+1i)^3} + \frac{x(2a^8 b^2 + 8a^6 b^2 + 12a^4 b^2 + 8a^2 b^2 + 2b^2)}{(a+1i)^3(-ba^6 + 2ib a^5 - ba^4 + 4ib a^3 + ba^2 + 2ib a + b)}\right)}{(a+1i)^3} 4i$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)^2/(x^3*((a + b*x)^2 + 1)),x)

[Out] $((a - 1i)/(2*(a + 1i)) + (b*x*2i)/(a + 1i)^2)/x^2 + (b^2*\operatorname{atanh}((3*a - a^2*3i - a^3 + 1i)/(a + 1i)^3 + (x*(2*b^2 + 8*a^2*b^2 + 12*a^4*b^2 + 8*a^6*b^2 + 2*a^8*b^2))/((a + 1i)^3*(b + a*b*2i + a^2*b + a^3*b*4i - a^4*b + a^5*b*2i - a^6*b)))*4i)/(a + 1i)^3$

$$3.179 \quad \int \frac{e^{2i \operatorname{ArcTan}(a+bx)}}{x^4} dx$$

Optimal. Leaf size=93

$$-\frac{i-a}{3(i+a)x^3} + \frac{ib}{(i+a)^2x^2} + \frac{2b^2}{(1-ia)^3x} - \frac{2ib^3 \log(x)}{(i+a)^4} + \frac{2ib^3 \log(i+a+bx)}{(i+a)^4}$$

[Out] $1/3*(-I+a)/(I+a)/x^3+I*b/(I+a)^2/x^2+2*b^2/(1-I*a)^3/x-2*I*b^3*\ln(x)/(I+a)^4+2*I*b^3*\ln(I+a+b*x)/(I+a)^4$

Rubi [A]

time = 0.04, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2ib^3 \log(x)}{(a+i)^4} + \frac{2ib^3 \log(a+bx+i)}{(a+i)^4} + \frac{2b^2}{(1-ia)^3x} + \frac{ib}{(a+i)^2x^2} - \frac{-a+i}{3(a+i)x^3}$$

Antiderivative was successfully verified.

[In] Int[E^((2*I)*ArcTan[a + b*x])/x^4,x]

[Out] $-1/3*(I - a)/((I + a)*x^3) + (I*b)/((I + a)^2*x^2) + (2*b^2)/(((1 - I*a)^3*x) - ((2*I)*b^3*\text{Log}[x])/(I + a)^4 + ((2*I)*b^3*\text{Log}[I + a + b*x])/(I + a)^4$

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\int \frac{e^{2i \tan^{-1}(a+bx)}}{x^4} dx = \int \frac{1 + ia + ibx}{x^4(1 - ia - ibx)} dx$$

$$= \int \left(\frac{i - a}{(i + a)x^4} - \frac{2ib}{(i + a)^2 x^3} + \frac{2ib^2}{(i + a)^3 x^2} - \frac{2ib^3}{(i + a)^4 x} + \frac{2ib^4}{(i + a)^4(i + a + bx)} \right) dx$$

$$= -\frac{i - a}{3(i + a)x^3} + \frac{ib}{(i + a)^2 x^2} + \frac{2b^2}{(1 - ia)^3 x} - \frac{2ib^3 \log(x)}{(i + a)^4} + \frac{2ib^3 \log(i + a + bx)}{(i + a)^4}$$

Mathematica [A]

time = 0.04, size = 88, normalized size = 0.95

$$\frac{(i + a)(i + a + ia^2 + a^3 - 3bx + 3iabx - 6ib^2x^2) - 6ib^3x^3 \log(x) + 6ib^3x^3 \log(i + a + bx)}{3(i + a)^4 x^3}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a + b*x])/x^4, x]`

```
[Out] ((I + a)*(I + a + I*a^2 + a^3 - 3*b*x + (3*I)*a*b*x - (6*I)*b^2*x^2) - (6*I)*b^3*x^3*Log[x] + (6*I)*b^3*x^3*Log[I + a + b*x])/(3*(I + a)^4*x^3)
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 266 vs. 2(79) = 158.

time = 0.16, size = 267, normalized size = 2.87

method	result
default	$2b^4 \left(\frac{(ia^4b + 4a^3b - 6ia^2b - 4ab + ib) \ln(b^2x^2 + 2abx + a^2 + 1)}{2b^2} + \frac{(ia^5 - 10ia^3 + 5a^4 + 5ia - 10a^2 + 1 - \frac{(ia^4b + 4a^3b - 6ia^2b - 4ab + ib)a}{b}) \arctan\left(\frac{2b^2x + 2a}{2b}\right)}{b} \right) \frac{1}{(a^2 + 1)^4}$
risch	$-\frac{2ib^2x^2}{(a^2 + 2ia - 1)(i + a)} + \frac{ibx}{a^2 + 2ia - 1} + \frac{a - i}{3i + 3a} - \frac{b^3 \ln(4a^{12}b^2x^2 + 8a^{13}bx + 4a^{14} + 24a^{10}b^2x^2 + 48a^{11}bx + 28a^{12} + 60a^8b^2x^2 + 120a^9bx + 84a^{10})}{x^3}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^4, x, method=_RETURNVERBOSE)`

```
[Out] 2*b^4/(a^2+1)^4*(1/2*(I*a^4*b-6*I*b*a^2+4*a^3*b+I*b-4*a*b)/b^2*ln(b^2*x^2+2*a*b*x+a^2+1)+(I*a^5-10*I*a^3+5*a^4+5*I*a-10*a^2+1-(I*a^4*b-6*I*b*a^2+4*a^3*b+I*b-4*a*b)*a/b)/b*arctan(1/2*(2*b^2*x+2*a*b)/b))-1/3*(2*I*a-a^2+1)/(a^2+1)/x^3+b*(I*a^2-I+2*a)/(a^2+1)^2/x^2-2*b^2*(I*a^3-3*I*a+3*a^2-1)/(a^2+1)^3/x-2*b^3*(I*a^4-6*I*a^2+4*a^3+I-4*a)/(a^2+1)^4*ln(x)
```

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 263 vs. $2(69) = 138$.
time = 0.49, size = 263, normalized size = 2.83

$$\frac{2(a^4 - 4ia^3 - 6a^2 + 4ia + 1)b^3 \arctan\left(\frac{bx+ia}{b}\right) + (ia^4 + 4a^3 - 6ia^2 - 4a + i)b^3 \log(b^2x^2 + 2abx + a^2 + 1) - 2(ia^4 + 4a^3 - 6ia^2 - 4a + i)b^3 \log(x) + \frac{a^5 - 2ia^4 + 6(-ia^3 - 3a^2 + 3ia + 1)b^2x^2 + a^4 - 4ia^3 + 3(ia^2 + 2a^3 + 2a - i)bx - a^2 - 2ia - 1}{3(a^5 + 3a^4 + 3a^3 + 1)x^3}}{a^5 + 4a^4 + 6a^3 + 4a^2 + 1}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^4,x, algorithm="maxima")

[Out] $2*(a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1)*b^3*\arctan((b^2*x + a*b)/b)/(a^8 + 4*a^6 + 6*a^4 + 4*a^2 + 1) + (I*a^4 + 4*a^3 - 6*I*a^2 - 4*a + I)*b^3*\log(b^2*x^2 + 2*a*b*x + a^2 + 1)/(a^8 + 4*a^6 + 6*a^4 + 4*a^2 + 1) - 2*(I*a^4 + 4*a^3 - 6*I*a^2 - 4*a + I)*b^3*\log(x)/(a^8 + 4*a^6 + 6*a^4 + 4*a^2 + 1) + 1/3*(a^6 - 2*I*a^5 + 6*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b^2*x^2 + a^4 - 4*I*a^3 + 3*(I*a^4 + 2*a^3 + 2*a - I)*b*x - a^2 - 2*I*a - 1)/((a^6 + 3*a^4 + 3*a^2 + 1)*x^3)$

Fricas [A]

time = 2.25, size = 94, normalized size = 1.01

$$\frac{-6ib^3x^3 \log(x) + 6ib^3x^3 \log\left(\frac{bx+a+i}{b}\right) - 6(ia-1)b^2x^2 + a^4 + 2ia^3 - 3(-ia^2 + 2a+i)bx + 2ia - 1}{3(a^4 + 4ia^3 - 6a^2 - 4ia + 1)x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^4,x, algorithm="fricas")

[Out] $1/3*(-6*I*b^3*x^3*\log(x) + 6*I*b^3*x^3*\log((b*x + a + I)/b) - 6*(I*a - 1)*b^2*x^2 + a^4 + 2*I*a^3 - 3*(-I*a^2 + 2*a + I)*b*x + 2*I*a - 1)/((a^4 + 4*I*a^3 - 6*a^2 - 4*I*a + 1)*x^3)$

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 286 vs. $2(73) = 146$.

time = 0.62, size = 286, normalized size = 3.08

$$\frac{2ib^3 \log\left(\frac{-2ia^3b^3 - 10ia^2b^3 + 20a^2b^3 + 20a^2b^3 + 2ab^3 - 10ab^3}{(a+i)^4} + 4b^4x + 2ib^3 - \frac{2ib^3}{(a+i)^2}\right) + \frac{2ib^3 \log\left(\frac{2ia^3b^3 + 10ia^2b^3 - 20a^2b^3 - 20a^2b^3 + 2ab^3 + 10ab^3}{(a+i)^4} + 4b^4x + 2ib^3 + \frac{2ib^3}{(a+i)^2}\right) - a^3 - ia^2 - a + 6ib^2x^2 + x(-3iab + 3b) - i}{(a+i)^4 x^3 (3a^2 + 9ia^2 - 9a - 3i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a)**2/(1+(b*x+a)**2)/x**4,x)

[Out] $-2*I*b**3*\log(-2*a**5*b**3/(a + I)**4 - 10*I*a**4*b**3/(a + I)**4 + 20*a**3*b**3/(a + I)**4 + 20*I*a**2*b**3/(a + I)**4 + 2*a*b**3 - 10*a*b**3/(a + I)**4 + 4*b**4*x + 2*I*b**3 - 2*I*b**3/(a + I)**4)/(a + I)**4 + 2*I*b**3*\log(2*a**5*b**3/(a + I)**4 + 10*I*a**4*b**3/(a + I)**4 - 20*a**3*b**3/(a + I)**4 - 20*I*a**2*b**3/(a + I)**4 + 2*a*b**3 + 10*a*b**3/(a + I)**4 + 4*b**4*x + 2*I*b**3 + 2*I*b**3/(a + I)**4)/(a + I)**4 - (-a**3 - I*a**2 - a + 6*I*b**2*x**2 + x*(-3*I*a*b + 3*b) - I)/(x**3*(3*a**3 + 9*I*a**2 - 9*a - 3*I))$

Giac [A]

time = 0.42, size = 126, normalized size = 1.35

$$\frac{2b^4 \log(bx + a + i)}{-i a^4 b + 4a^3 b + 6i a^2 b - 4ab - ib} + \frac{2b^3 \log(|x|)}{i a^4 - 4a^3 - 6i a^2 + 4a + i} + \frac{a^4 + 2i a^3 - 6i(ab^2 + i b^2)x^2 + 3i(a^2 b + 2i ab - b)x + 2i a - 1}{3(a + i)^4 x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^2/(1+(b*x+a)^2)/x^4,x, algorithm="giac")

[Out] 2*b^4*log(b*x + a + I)/(-I*a^4*b + 4*a^3*b + 6*I*a^2*b - 4*a*b - I*b) + 2*b^3*log(abs(x))/(I*a^4 - 4*a^3 - 6*I*a^2 + 4*a + I) + 1/3*(a^4 + 2*I*a^3 - 6*I*(a*b^2 + I*b^2)*x^2 + 3*I*(a^2*b + 2*I*a*b - b)*x + 2*I*a - 1)/((a + I)^4*x^3)

Mupad [B]

time = 0.72, size = 199, normalized size = 2.14

$$\frac{\frac{a-i}{3(a+i)} - \frac{b^2 x^2 2i}{(a+i)^3} + \frac{b x 1i}{(a+i)^2}}{x^3} + \frac{b^3 \operatorname{atanh}\left(\frac{a^4+a^3 4i-6a^2-a 4i+1}{(a+i)^4} - \frac{x(2a^{12}b^2+12a^{10}b^2+30a^8b^2+40a^6b^2+30a^4b^2+12a^2b^2+2b^2)}{(a+i)^4(-ba^9+3ib a^8+8ib a^6+6ba^5+6iba^4+8ba^3+3ba-b1i)}\right) 4i}{(a + 1i)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)^2/(x^4*((a + b*x)^2 + 1)),x)

[Out] ((a - 1i)/(3*(a + 1i)) - (b^2*x^2*2i)/(a + 1i)^3 + (b*x*1i)/(a + 1i)^2)/x^3 + (b^3*atanh((a^3*4i - 6*a^2 - a*4i + a^4 + 1)/(a + 1i)^4 - (x*(2*b^2 + 12*a^2*b^2 + 30*a^4*b^2 + 40*a^6*b^2 + 30*a^8*b^2 + 12*a^10*b^2 + 2*a^12*b^2))/(a + 1i)^4*(3*a*b - b*1i + 8*a^3*b + a^4*b*6i + 6*a^5*b + a^6*b*8i + a^8*b*3i - a^9*b)))*4i)/(a + 1i)^4

3.180 $\int e^{3i\text{ArcTan}(a+bx)} x^4 dx$

Optimal. Leaf size=324

$$\frac{3(19i + 68a - 88ia^2 - 48a^3 + 8ia^4) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{8b^5} - \frac{2ix^4(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} + \frac{3(17i + 16a)x^2}{b^2}$$

[Out] $-3/8*(19-68*I*a-88*a^2+48*I*a^3+8*a^4)*\text{arcsinh}(b*x+a)/b^5-2*I*x^4*(1+I*a+I*b*x)^{(3/2)}/b/(1-I*a-I*b*x)^{(1/2)}+3/20*(17*I+16*a)*x^2*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^3-11/5*x^3*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^2-1/40*I*(1+I*a+I*b*x)^{(3/2)}*(163-458*I*a-422*a^2+112*I*a^3+2*(61*I+118*a-52*I*a^2)*b*x)*(1-I*a-I*b*x)^{(1/2)}/b^5-3/8*(19*I+68*a-88*I*a^2-48*a^3+8*I*a^4)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^5$

Rubi [A]

time = 0.20, antiderivative size = 324, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 99, 158, 152, 52, 55, 633, 221}

$$\frac{\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}(112a^2+2(-52a^2+118a+61)ix-422a^2-458a+163)-3(8a^4-48a^3-88a^2+68a+19)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}-3(8a^4+48a^3-88a^2-68a+19)\sinh^{-1}(a+bx)+3(16a+17)a^2\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}-11a^2\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}-2ax^4(ia+ibx+1)^{3/2}}{8b^5}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a + b*x])*x^4,x]

[Out] $(-3*(19*I + 68*a - (88*I)*a^2 - 48*a^3 + (8*I)*a^4)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/(8*b^5) - ((2*I)*x^4*(1 + I*a + I*b*x)^{(3/2)})/(b*\text{Sqrt}[1 - I*a - I*b*x]) + (3*(17*I + 16*a)*x^2*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)})/(20*b^3) - (11*x^3*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)})/(5*b^2) - ((I/40)*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)}*(163 - (458*I)*a - 422*a^2 + (112*I)*a^3 + 2*(61*I + 118*a - (52*I)*a^2)*b*x))/b^5 - (3*(19 - (68*I)*a - 88*a^2 + (48*I)*a^3 + 8*a^4)*\text{ArcSinh}[a + b*x])/(8*b^5)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] :> Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b

+ d, 0] && GtQ[a + c, 0]

Rule 99

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p/(b*(m + 1)), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 152

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)), Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]

Rule 158

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p + 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{3i \tan^{-1}(a+bx)} x^4 dx &= \int \frac{x^4(1+ia+ibx)^{3/2}}{(1-ia-ibx)^{3/2}} dx \\
&= -\frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} + \frac{(2i) \int \frac{x^3 \sqrt{1+ia+ibx} (4(1+ia) + \frac{11ibx}{2})}{\sqrt{1-ia-ibx}} dx}{b} \\
&= -\frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} - \frac{11x^3 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{5b^2} + \frac{(2i) \int \frac{x^2 \sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx}{b} \\
&= -\frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} + \frac{3(17i+16a)x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{20b^3} - \frac{11x^3 \sqrt{1+ia+ibx}}{5b^2} \\
&= -\frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} + \frac{3(17i+16a)x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{20b^3} - \frac{11x^3 \sqrt{1+ia+ibx}}{5b^2} \\
&= -\frac{3(19i+68a-88ia^2-48a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
&= -\frac{3(19i+68a-88ia^2-48a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
&= -\frac{3(19i+68a-88ia^2-48a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
&= -\frac{3(19i+68a-88ia^2-48a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
&= -\frac{3(19i+68a-88ia^2-48a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} - \frac{2ix^4(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}}
\end{aligned}$$

Mathematica [A]

time = 0.33, size = 249, normalized size = 0.77

$$\frac{\sqrt{1+ibx} (448i+418ia^4+8a^5+163bx+61ib^2x^2-34b^3x^3-22ib^4x^4+8b^5x^5+14ia^2(121i+8bx)-ia^2(2599-422ibx+52b^2x^2)+a(1763-458ibx+118b^2x^2+32ib^3x^3))}{40b^5 \sqrt{-i(i+bx)}} + \frac{3(-1)^{3/4} (19-68ia-88a^2+48ia^3+8a^4) \sinh^{-1}\left(\frac{(1+i)\sqrt{b} \sqrt{-i(i+bx)}}{\sqrt{-ib}}\right)}{4\sqrt{-ib} b^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])*x^4,x]

[Out]
$$\frac{-1/40 \cdot (\sqrt{1 + I \cdot a + I \cdot b \cdot x}) \cdot (448 \cdot I + (418 \cdot I) \cdot a^4 + 8 \cdot a^5 + 163 \cdot b \cdot x + (61 \cdot I) \cdot b^2 \cdot x^2 - 34 \cdot b^3 \cdot x^3 - (22 \cdot I) \cdot b^4 \cdot x^4 + 8 \cdot b^5 \cdot x^5 + (14 \cdot I) \cdot a^3 \cdot (121 \cdot I + 8 \cdot b \cdot x) - I \cdot a^2 \cdot (2599 - (422 \cdot I) \cdot b \cdot x + 52 \cdot b^2 \cdot x^2) + a \cdot (1763 - (458 \cdot I) \cdot b \cdot x + 118 \cdot b^2 \cdot x^2 + (32 \cdot I) \cdot b^3 \cdot x^3))}{(b^5 \cdot \sqrt{(-I) \cdot (I + a + b \cdot x)})} + \frac{(3 \cdot (-1)^{(3/4)} \cdot (19 - (68 \cdot I) \cdot a - 88 \cdot a^2 + (48 \cdot I) \cdot a^3 + 8 \cdot a^4) \cdot \text{ArcSinh}[\frac{(1/2 + I/2) \cdot \sqrt{b} \cdot \sqrt{(-I) \cdot (I + a + b \cdot x)}}{\sqrt{(-I) \cdot b}}])}{4 \cdot \sqrt{(-I) \cdot b} \cdot b^{(9/2)}}$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 4892 vs. $2(262) = 524$.

time = 0.22, size = 4893, normalized size = 15.10

method	result
risch	$\frac{i(8x^4b^4 - 8ab^3x^3 - 30ib^3x^3 + 8a^2b^2x^2 + 70ia^2b^2x^2 - 8a^3bx - 130ia^2b + 8a^4 + 250ia^3 - 64b^2x^2 + 252abx + 125ibx - 804a^2 - 835ia + 288)}{40b^5}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^4,x,method=_RETURNVERBOSE)

[Out]
$$\begin{aligned} & -I \cdot b^3 \cdot \frac{1}{5} \cdot x^6 / b^2 / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 11/5 \cdot a / b \cdot \frac{1}{4} \cdot x^5 / b^2 / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 9/4 \cdot a / b \cdot \frac{1}{3} \cdot x^4 / b^2 / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 7/3 \cdot a / b \cdot \frac{1}{2} \cdot x^3 / b^2 / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 5/2 \cdot a / b \cdot \frac{x^2}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 3 \cdot a / b \cdot \frac{-x}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - a / b \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} + 1/b^2 \cdot \ln((b^2 \cdot x + a \cdot b) / (b^2)^{(1/2)} + (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)}) / (b^2)^{(1/2)} - 2 \cdot (a^2 + 1) / b^2 \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 3/2 \cdot (a^2 + 1) / b^2 \cdot \frac{-x}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - a / b \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & + 1/b^2 \cdot \ln((b^2 \cdot x + a \cdot b) / (b^2)^{(1/2)} + (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)}) / (b^2)^{(1/2)} - 4/3 \cdot (a^2 + 1) / b^2 \cdot \frac{x^2}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 3 \cdot a / b \cdot \frac{-x}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - a / b \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} + 1/b^2 \cdot \ln((b^2 \cdot x + a \cdot b) / (b^2)^{(1/2)} + (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)}) / (b^2)^{(1/2)} \\ & - 2 \cdot (a^2 + 1) / b^2 \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 5/4 \cdot (a^2 + 1) / b^2 \cdot \frac{1}{2} \cdot x^3 / b^2 / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 5/2 \cdot a / b \cdot \frac{x^2}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 3 \cdot a / b \cdot \frac{-x}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - a / b \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \\ & - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} + 1/b^2 \cdot \ln((b^2 \cdot x + a \cdot b) / (b^2)^{(1/2)} + (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)}) / (b^2)^{(1/2)} \\ & - 2 \cdot (a^2 + 1) / b^2 \cdot \frac{-1}{b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} - 2 \cdot a / b \cdot \frac{2 \cdot b^2 \cdot x + 2 \cdot a \cdot b}{4 \cdot b^2 \cdot (a^2 + 1) - 4 \cdot a^2 \cdot b^2} / (b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1)^{(1/2)} \end{aligned}$$

$$\begin{aligned}
& a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})-3/2*(a^2+1)/ \\
& b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+ \\
& 1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a \\
& ^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} \\
&))/(b^2)^{(1/2)}))-6/5*(a^2+1)/b^2*(1/3*x^4/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} \\
& -7/3*a/b*(1/2*x^3/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-5/2*a/b*(x^2/b^2/(b^2*x \\
& ^2+2*a*b*x+a^2+1)^{(1/2)}-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*(-1 \\
& /b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a \\
& ^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^ \\
& 2*x^2+2*a*b*x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a \\
& *b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+ \\
& 2*a*b*x+a^2+1)^{(1/2)}))-3/2*(a^2+1)/b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} \\
&)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a \\
& ^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1 \\
& /2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}))-4/3*(a^2+1)/b^2*(x^2/b^2 \\
& /b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}- \\
& a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2 \\
& +1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1 \\
& /2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}))-2*(a^2+1)/b^2*(-1/b^2/(b^2* \\
& x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b \\
& ^2*x^2+2*a*b*x+a^2+1)^{(1/2)})))-3*(1+I*a)*b^2*(1/4*x^5/b^2/(b^2*x^2+2*a*b*x \\
& +a^2+1)^{(1/2)}-9/4*a/b*(1/3*x^4/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-7/3*a/b*(1 \\
& /2*x^3/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-5/2*a/b*(x^2/b^2/(b^2*x^2+2*a*b*x+ \\
& a^2+1)^{(1/2)}-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*(-1/b^2/(b^2*x \\
& ^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^ \\
& 2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b \\
& *x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1) \\
& ^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2 \\
& +1)^{(1/2)}))-3/2*(a^2+1)/b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*(-1/b \\
& ^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2 \\
& *b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2* \\
& x^2+2*a*b*x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}))-4/3*(a^2+1)/b^2*(x^2/b^2/(b^2*x^2+2 \\
& *a*b*x+a^2+1)^{(1/2)}-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-a/b*(-1/b^2 \\
& /b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b \\
& ^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^ \\
& 2+2*a*b*x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x \\
& +a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a* \\
& b*x+a^2+1)^{(1/2)})))-5/4*(a^2+1)/b^2*(1/2*x^3/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1 \\
& /2)}-5/2*a/b*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3*a/b*(-x/b^2/(b^2*x^2+2 \\
& *a*b*x+a^2+1)^{(1/2)}-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2* \\
& x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/b^2*\ln(\\
& (b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}))/(b^2)^{(1/2)}))-2*(a^2+ \\
& 1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(\\
& a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}...
\end{aligned}$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3081 vs. $2(230) = 460$.
time = 0.30, size = 3081, normalized size = 9.51

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^4,x, algorithm="maxima")
[Out] -1/5*I*b*x^6/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 11/20*I*a*x^5/sqrt(b^2*x^2
+ 2*a*b*x + a^2 + 1) - 693/4*I*a^7*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x
^2 + 2*a*b*x + a^2 + 1)*b^2) - 33/20*I*a^2*x^4/(sqrt(b^2*x^2 + 2*a*b*x + a
^2 + 1)*b) + 2415/8*I*(a^2 + 1)*a^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x
^2 + 2*a*b*x + a^2 + 1)*b^2) + 231/40*I*a^3*x^3/(sqrt(b^2*x^2 + 2*a*b*x + a
^2 + 1)*b^2) - 2/5*(-I*a^2 - I)*x^4/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) -
3/4*(I*a*b^2 + b^2)*x^5/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2) - 231/8*I*(
a^2 + 1)*a^6/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b
^3) + 945/4*(I*a*b^2 + b^2)*a^6*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 +
2*a*b*x + a^2 + 1)*b^4) - 105*(I*a^2*b + 2*a*b - I*b)*a^5*x/((a^2*b^2 - (a
^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - 2919/20*I*(a^2 + 1)^2
*a^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2) -
15*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x
^2 + 2*a*b*x + a^2 + 1)*b^2) - 231/8*I*a^4*x^2/(sqrt(b^2*x^2 + 2*a*b*x + a
^2 + 1)*b^3) - 111/40*I*(a^2 + 1)*a*x^3/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b
^2) + 9/4*(I*a*b^2 + b^2)*a*x^4/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - (
I*a^2*b + 2*a*b - I*b)*x^4/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2) + 189/5*
I*(a^2 + 1)^2*a^4/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 +
1)*b^3) - 2835/8*(I*a*b^2 + b^2)*(a^2 + 1)*a^4*x/((a^2*b^2 - (a^2 + 1)*b^2
)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^4) + 265/2*(I*a^2*b + 2*a*b - I*b)*(a
^2 + 1)*a^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*
b^3) + 653/40*I*(a^2 + 1)^3*a*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2
*a*b*x + a^2 + 1)*b^2) + 31/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)*a^2*x/
((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2) + 819/40*
I*(a^2 + 1)*a^2*x^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - 63/8*(I*a*b^2
+ b^2)*a^2*x^3/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^4) + 7/2*(I*a^2*b + 2*
a*b - I*b)*a*x^3/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - 1/2*(I*a^3 + 3*a
^2 - 3*I*a - 1)*x^3/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2) + 693/8*I*a^5*a
rcsinh(2*(b^2*x + a*b)/sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^5 + 315/8*(I*a
*b^2 + b^2)*(a^2 + 1)*a^5/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x
+ a^2 + 1)*b^5) - 35/2*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*a^4/((a^2*b^2 - (
a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^4) - 397/40*I*(a^2 + 1)^3
*a^2/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - 5/
2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)*a^3/((a^2*b^2 - (a^2 + 1)*b^2)*sqr
t(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) + 507/4*(I*a*b^2 + b^2)*(a^2 + 1)^2*a^2
*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^4) - 61/2
```

$$\begin{aligned}
&*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)^2*a*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^3) - 3/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^2) + 3 \\
&15/8*(I*a*b^2 + b^2)*a^3*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^5) - 35/2*(I*a^2*b + 2*a*b - I*b)*a^2*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^4) - \\
&8/5*I*(a^2 + 1)^2*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^3) - 5/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^3) + 15/8*(I*a*b^2 + b^2)*(a^2 + 1)*x^3/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^4) - 315 \\
&/4*I*(a^2 + 1)*a^3*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^5 - 42*(I*a*b^2 + b^2)*(a^2 + 1)^2*a^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^5) - 231/4*I*(a^2 + 1)*a^4/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^5) + 29/2*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)^2*a^2/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^4) + 3/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)^2*a/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^3) - 45/8*(I*a*b^2 + b^2)*(a^2 + 1)^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^4) - 147/8*(I*a*b^2 + b^2)*(a^2 + 1)*a*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^5) + 4*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^4) - 945/8*(I*a*b^2 + b^2)*a^4*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^7 + 105/2*(I*a^2*b + 2*a*b - I*b)*a^3*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^6 + 105/8*I*(a^2 + 1)^2*a*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^5 + 15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^5 + 45/8*(I*a*b^2 + b^2)*(a^2 + 1)^3*a/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^5) + 819/20*I*(a^2 + 1)^2*a^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^5) + 315/4*(I*a*b^2 + b^2)*(a^2 + 1)*a^2*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^7 - 45/2*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*a*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^6 - 3/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^5 + 315/4*(I*a*b^2 + b^2)*(a^2 + 1)*a^3/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^7) - 35...
\end{aligned}$$

Fricas [A]

time = 1.72, size = 264, normalized size = 0.81

-62*a^7 + 267*a^6 + 11575*a^5 - 20350*a^4 - (-62*a^6 + 2625*a^5 + 8950*a^4 - 11400*a^3 - 6340*a^2 + 1280)*b*x - 17740*a^2 + 120*(8*a^5 + 56*I*a^4 - 136*a^3 + (8*a^4 + 48*I*a^3 - 88*a^2 - 68*I*a + 19)*b*x - 156*I*a^2 + 87*a + 19*I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 8*(8*I*b^5*x^5 + 22*b^4*x^4 - 2*(16*a + 17*I)*b^3*x^3 + 8*I*a^5 + (52*a^2

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^4,x, algorithm="fricas")

[Out] 1/320*(-62*I*a^6 + 2687*a^5 + 11575*I*a^4 - 20350*a^3 + (-62*I*a^5 + 2625*a^4 + 8950*I*a^3 - 11400*a^2 - 6340*I*a + 1280)*b*x - 17740*I*a^2 + 120*(8*a^5 + 56*I*a^4 - 136*a^3 + (8*a^4 + 48*I*a^3 - 88*a^2 - 68*I*a + 19)*b*x - 156*I*a^2 + 87*a + 19*I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 8*(8*I*b^5*x^5 + 22*b^4*x^4 - 2*(16*a + 17*I)*b^3*x^3 + 8*I*a^5 + (52*a^2

$$+ 118*I*a - 61)*b^2*x^2 - 418*a^4 - 1694*I*a^3 - (112*a^3 + 422*I*a^2 - 458*a - 163*I)*b*x + 2599*a^2 + 1763*I*a - 448)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 7620*a + 1280*I)/(b^6*x + (a + I)*b^5)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2)*x**4,x)

[Out] $-I*(\text{Integral}(I*x**4/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(-3*a*x**4/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(a**3*x**4/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(-3*b*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(-3*I*a**2*x**4/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(-3*I*b**2*x**6/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(3*a*b**2*x**6/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(3*a**2*b*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x) + \text{Integral}(-6*I*a*b*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1})), x)$

Giac [A]

time = 0.46, size = 334, normalized size = 1.03

$$\frac{1}{10} \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \left(\left(\frac{1}{10} \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \right) \left(-4 a^2 b^2 + 20 a^2 b^2 \right) - 4 a^2 b^2 + 20 a^2 b^2 - 20 a^2 b^2 + 120 a^2 b^2 \right) - \frac{12 a^2 b^2 + 20 a^2 b^2 + 80 a^2 b^2 - 40 a^2 b^2}{10 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}} + \frac{12 a^2 b^2 + 20 a^2 b^2 - 40 a^2 b^2 + 120 a^2 b^2}{10 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}} \ln \left(\frac{b x + a + \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{b x + a - \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}} \right) \right) + 2 a^2 b^2 + \left(\frac{1}{10} \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \right) \left(40 b - 40 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \right) \ln \left(\frac{b x + a + \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{b x + a - \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}} \right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^4,x, algorithm="giac")

[Out] -1/40*sqrt((b*x + a)^2 + 1)*((2*(x*(4*I*x/b - (4*I*a*b^17 - 15*b^17)/b^19) - (-4*I*a^2*b^16 + 35*a*b^16 + 32*I*b^16)/b^19)*x - (8*I*a^3*b^15 - 130*a^2*b^15 - 252*I*a*b^15 + 125*b^15)/b^19)*x - (-8*I*a^4*b^14 + 250*a^3*b^14 + 804*I*a^2*b^14 - 835*a*b^14 - 288*I*b^14)/b^19) + 1/8*(8*a^4 + 48*I*a^3 - 8*8*a^2 - 68*I*a + 19)*log(3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b + a^3*b + (x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) + 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^2*abs(b) + 2*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b + 2*I*a^2*b + 4*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a*abs(b) - a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^4*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^4 (1 + a \operatorname{li} + b x \operatorname{li})^3}{((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*(a*1i + b*x*1i + 1)^3)/((a + b*x)^2 + 1)^(3/2),x)

[Out] int((x^4*(a*1i + b*x*1i + 1)^3)/((a + b*x)^2 + 1)^(3/2), x)

3.181 $\int e^{3i\text{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=249

$$\frac{3(17 - 44ia - 36a^2 + 8ia^3) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{8b^4} - \frac{2ix^3(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} - \frac{9x^2\sqrt{1 - ia - ibx}}{4b^2} (1 + ia + ibx)$$

[Out] $-3/8*(17*I+44*a-36*I*a^2-8*a^3)*\text{arcsinh}(b*x+a)/b^4-2*I*x^3*(1+I*a+I*b*x)^(3/2)/b/(1-I*a-I*b*x)^(1/2)-9/4*x^2*(1+I*a+I*b*x)^(3/2)*(1-I*a-I*b*x)^(1/2)/b^2-1/8*I*(1+I*a+I*b*x)^(3/2)*(29*I+54*a-22*I*a^2-2*(11-10*I*a)*b*x)*(1-I*a-I*b*x)^(1/2)/b^4+3/8*(17-44*I*a-36*a^2+8*I*a^3)*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/b^4$

Rubi [A]

time = 0.17, antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 99, 158, 152, 52, 55, 633, 221}

$$\frac{i\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}(-22ia^2-2(11-10ia)bx+54a+29i)}{8b^4} + \frac{3(8ia^3-36a^2-44ia+17)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{8b^4} - \frac{3(-8a^3-36ia^2+44a+17i)\sinh^{-1}(a+bx)}{8b^4} - \frac{9x^2\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{4b^2} - \frac{2ix^3(ia+ibx+1)^{3/2}}{b\sqrt{-ia-ibx+1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)*\text{ArcTan}[a + b*x])}*x^3, x]$

[Out] $(3*(17 - (44*I)*a - 36*a^2 + (8*I)*a^3)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/(8*b^4) - ((2*I)*x^3*(1 + I*a + I*b*x)^(3/2))/(b*\text{Sqrt}[1 - I*a - I*b*x]) - (9*x^2*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^(3/2))/(4*b^2) - ((I/8)*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^(3/2)*(29*I + 54*a - (22*I)*a^2 - 2*(11 - (10*I)*a)*b*x))/b^4 - (3*(17*I + 44*a - (36*I)*a^2 - 8*a^3)*\text{ArcSinh}[a + b*x])/(8*b^4)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^(m + 1)*((c + d*x)^(n)/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

$\text{Int}[1/(\text{Sqrt}[(a_.) + (b_.)*(x_.)]*\text{Sqrt}[(c_.) + (d_.)*(x_.)]), x_Symbol] \rightarrow \text{Int}[1/\text{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /;$ FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]

Rule 99

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)), Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 158

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p + 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
```

$I*b*c*x^{(I*(n/2))}, x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned}
 \int e^{3i \tan^{-1}(a+bx)} x^3 dx &= \int \frac{x^3(1+ia+ibx)^{3/2}}{(1-ia-ibx)^{3/2}} dx \\
 &= -\frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} + \frac{(2i) \int \frac{x^2 \sqrt{1+ia+ibx} (3(1+ia) + \frac{9ibx}{2})}{\sqrt{1-ia-ibx}} dx}{b} \\
 &= -\frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} - \frac{9x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} + \frac{i \int \frac{x \sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} dx}{\sqrt{1-ia-ibx}} \\
 &= -\frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} - \frac{9x^2 \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{4b^2} - \frac{i \sqrt{1-ia-ibx} (1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
 &= \frac{3(17-44ia-36a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} - \frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
 &= \frac{3(17-44ia-36a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} - \frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
 &= \frac{3(17-44ia-36a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} - \frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
 &= \frac{3(17-44ia-36a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} - \frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} \\
 &= \frac{3(17-44ia-36a^2+8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} - \frac{2ix^3(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}}
 \end{aligned}$$

Mathematica [A]

time = 0.19, size = 201, normalized size = 0.81

$$\frac{\sqrt{1+ia+ibx} (80+78ia^3+2a^4-29ibx+11b^2x^2+6ib^3x^3-2b^4x^4+a^2(-233+22ibx)-ia(237-54ibx+10b^2x^2))}{8b^4 \sqrt{-i(i+bx)}} + \frac{3\sqrt{-1}(-17i-44a+36ia^2+8a^3) \sqrt{-ib} \sinh^{-1}\left(\frac{(1+i)\sqrt{b} \sqrt{-i(i+bx)}}{\sqrt{-ib}}\right)}{4b^{9/2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])*x^3,x]

[Out] (Sqrt[1 + I*a + I*b*x]*(80 + (78*I)*a^3 + 2*a^4 - (29*I)*b*x + 11*b^2*x^2 + (6*I)*b^3*x^3 - 2*b^4*x^4 + a^2*(-233 + (22*I)*b*x) - I*a*(237 - (54*I)*b*x + 10*b^2*x^2)))/(8*b^4*Sqrt[(-I)*(I + a + b*x)]) + (3*(-1)^(1/4)*(-17*I -

$$\begin{aligned}
& +a^2+1)^{1/2}-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2*x+2*a* \\
& b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2})+1/b^2*\ln((b^2*x \\
& +a*b)/(b^2)^{1/2}+(b^2*x^2+2*a*b*x+a^2+1)^{1/2})/(b^2)^{1/2})-2*(a^2+1)/b^2 \\
& *(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1) \\
& -4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2})) -3/2*(a^2+1)/b^2*(-x/b^2/(b^2*x^ \\
& 2+2*a*b*x+a^2+1)^{1/2}-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-2*a/b*(2*b \\
& ^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2})+1/b^2* \\
& \ln((b^2*x+a*b)/(b^2)^{1/2}+(b^2*x^2+2*a*b*x+a^2+1)^{1/2})/(b^2)^{1/2})) -4/3 \\
& *(a^2+1)/b^2*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-3*a/b*(-x/b^2/(b^2*x^2+ \\
& 2*a*b*x+a^2+1)^{1/2}-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2 \\
& *x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2})+1/b^2*\ln \\
& ((b^2*x+a*b)/(b^2)^{1/2}+(b^2*x^2+2*a*b*x+a^2+1)^{1/2})/(b^2)^{1/2})-2*(a^2 \\
& +1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2* \\
& (a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}))) +3*I*(1+I*a)^2*b*(1/2*x^ \\
& 3/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-5/2*a/b*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1) \\
&)^{1/2}-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-a/b*(-1/b^2/(b^2*x^2+2* \\
& a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2 \\
& +2*a*b*x+a^2+1)^{1/2})) +1/b^2*\ln((b^2*x+a*b)/(b^2)^{1/2}+(b^2*x^2+2*a*b*x+a^ \\
& 2+1)^{1/2})/(b^2)^{1/2})-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2} \\
&)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2} \\
&)) -3/2*(a^2+1)/b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-a/b*(-1/b^2/(b \\
& ^2*x^2+2*a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2) \\
& / (b^2*x^2+2*a*b*x+a^2+1)^{1/2})) +1/b^2*\ln((b^2*x+a*b)/(b^2)^{1/2}+(b^2*x^2+2 \\
& *a*b*x+a^2+1)^{1/2})/(b^2)^{1/2}))) + (1+I*a)^3*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+ \\
& 1)^{1/2}-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2}-a/b*(-1/b^2/(b^2*x^2+2 \\
& *a*b*x+a^2+1)^{1/2}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^ \\
& 2+2*a*b*x+a^2+1)^{1/2})) +1/b^2*\ln((b^2*x+a*b)/(b^2)^{1/2}+(b^2*x^2+2*a*b*x+a \\
& ^2+1)^{1/2})/(b^2)^{1/2})-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^{1/2} \\
&)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{1/2} \\
&))
\end{aligned}$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2295 vs. $2(175) = 350$.

time = 0.29, size = 2295, normalized size = 9.22

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^3,x, algorithm="maxima")

[Out] $-1/4*I*b*x^5/\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 315/4*I*a^6*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*b) + 3/4*I*a*x^4/\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} - 945/8*I*(a^2 + 1)*a^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*b) - 21/8*I*a^2*x^3/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*b) + 105/8*I*(a^2 + 1)*a^5/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})$

$$\begin{aligned}
& (b^2x^2 + 2abx + a^2 + 1)b^2) - 105*(Iab^2 + b^2)a^5x/((a^2b^2 - \\
& (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^3) + 45*(Ia^2b + 2ab \\
& - I*b)a^4x/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1} \\
& b^2) + 169/4*I*(a^2 + 1)^2a^2x/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + \\
& 2abx + a^2 + 1}b) + 6*(-Ia^3 - 3a^2 + 3Ia + 1)a^3x/((a^2b^2 - (a \\
& ^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b) + 105/8*Ia^3x^2/(\sqrt{b \\
& ^2x^2 + 2abx + a^2 + 1}b^2) - 5/8*(-Ia^2 - I)x^3/(\sqrt{b^2x^2 + 2a \\
& *bx + a^2 + 1}b) - (Iab^2 + b^2)x^4/(\sqrt{b^2x^2 + 2abx + a^2 + 1} \\
& *b^2) - 14*I*(a^2 + 1)^2a^3/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2a \\
& *bx + a^2 + 1}b^2) + 265/2*(Iab^2 + b^2)*(a^2 + 1)a^3x/((a^2b^2 - (a \\
& ^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^3) - 93/2*(Ia^2b + 2ab \\
& - I*b)*(a^2 + 1)a^2x/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + \\
& a^2 + 1}b^2) - 15/8*I*(a^2 + 1)^3x/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^ \\
& 2 + 2abx + a^2 + 1}b) - 5*(-Ia^3 - 3a^2 + 3Ia + 1)*(a^2 + 1)a*x/((\\
& a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b) - 49/8*I*(a^2 \\
& + 1)a*x^2/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^2) + 7/2*(Iab^2 + b^2)*a \\
& *x^3/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^3) - 3/2*(Ia^2b + 2ab - I*b)* \\
& x^3/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^2) - 315/8*Ia^4*\operatorname{arcsinh}(2*(b^2x \\
& + a*b)/\sqrt{-4a^2b^2 + 4*(a^2 + 1)b^2})/b^4 - 35/2*(Iab^2 + b^2)*(a^2 \\
& + 1)a^4/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^4) \\
& + 15/2*(Ia^2b + 2ab - I*b)*(a^2 + 1)a^3/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{ \\
& b^2x^2 + 2abx + a^2 + 1}b^3) + 15/8*I*(a^2 + 1)^3a/((a^2b^2 - (a^2 \\
& + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^2) + (-Ia^3 - 3a^2 + 3Ia \\
& + 1)*(a^2 + 1)a^2/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 \\
& + 1}b^2) - 61/2*(Iab^2 + b^2)*(a^2 + 1)^2a*x/((a^2b^2 - (a^2 + 1)b^2 \\
&)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^3) + 9/2*(Ia^2b + 2ab - I*b)*(a^2 \\
& + 1)^2x/((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^2) \\
& - 35/2*(Iab^2 + b^2)a^2x^2/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^4) + 1 \\
& 5/2*(Ia^2b + 2ab - I*b)a*x^2/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^3) + \\
& (-Ia^3 - 3a^2 + 3Ia + 1)x^2/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^2) + \\
& 105/4*I*(a^2 + 1)a^2*\operatorname{arcsinh}(2*(b^2x + a*b)/\sqrt{-4a^2b^2 + 4*(a^2 + 1) \\
& *b^2})/b^4 + 29/2*(Iab^2 + b^2)*(a^2 + 1)^2a^2/((a^2b^2 - (a^2 + 1)b^ \\
& 2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^4) + 105/4*I*(a^2 + 1)a^3/(\sqrt{b^2 \\
& *x^2 + 2abx + a^2 + 1}b^4) - 9/2*(Ia^2b + 2ab - I*b)*(a^2 + 1)^2a/ \\
& ((a^2b^2 - (a^2 + 1)b^2)*\sqrt{b^2x^2 + 2abx + a^2 + 1}b^3) + 4*(Ia* \\
& b^2 + b^2)*(a^2 + 1)x^2/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^4) + 105/2*(I \\
& *ab^2 + b^2)a^3*\operatorname{arcsinh}(2*(b^2x + a*b)/\sqrt{-4a^2b^2 + 4*(a^2 + 1)b^2 \\
& })/b^6 - 45/2*(Ia^2b + 2ab - I*b)a^2*\operatorname{arcsinh}(2*(b^2x + a*b)/\sqrt{-4a \\
& ^2b^2 + 4*(a^2 + 1)b^2})/b^5 - 15/8*I*(a^2 + 1)^2*\operatorname{arcsinh}(2*(b^2x + a*b) \\
& / \sqrt{-4a^2b^2 + 4*(a^2 + 1)b^2})/b^4 - 3*(-Ia^3 - 3a^2 + 3Ia + 1)a \\
& * \operatorname{arcsinh}(2*(b^2x + a*b)/\sqrt{-4a^2b^2 + 4*(a^2 + 1)b^2})/b^4 - 49/4*I*(\\
& a^2 + 1)^2a/(\sqrt{b^2x^2 + 2abx + a^2 + 1}b^4) - 45/2*(Iab^2 + b^2) \\
& *(a^2 + 1)a*\operatorname{arcsinh}(2*(b^2x + a*b)/\sqrt{-4a^2b^2 + 4*(a^2 + 1)b^2})/b^ \\
& 6 + 9/2*(Ia^2b + 2ab - I*b)*(a^2 + 1)*\operatorname{arcsinh}(2*(b^2x + a*b)/\sqrt{-4a \\
& ^2b^2 + 4*(a^2 + 1)b^2})/b^5 - 35*(Iab^2 + b^2)*(a^2 + 1)a^2/(\sqrt{b^2
\end{aligned}$$

$$*x^2 + 2*a*b*x + a^2 + 1)*b^6) + 15*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*a/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^5) + 2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^4) + 8*(I*a*b^2 + b^2)*(a^2 + 1)^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^6)$$

Fricas [A]

time = 2.82, size = 216, normalized size = 0.87

$\frac{15a^5 - 495a^4 - 1664a^3 + (15a^4 - 480a^3 - 1184a^2 + 968a + 256)bx + 2152a^2 - 24(8a^4 + 44a^3 + (8a^3 + 36a^2 - 44a - 17)bx - 80a^2 - 61a + 17)\log\left(\frac{-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}}{64(b^2x + (a + I)b^4)}\right) - 8(2Ib^4 + 6b^3 - (10a + 11I)b^2 - 2a^4 + 78a^3 + (22a^2 + 54a - 29)bx + 233a^2 - 237a - 80I)\sqrt{b^2x^2 + 2abx + a^2 + 1} + 1224a - 256}{64(b^2x + (a + I)b^4)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^3,x, algorithm="fricas")

[Out] $\frac{1}{64}*(15*I*a^5 - 495*a^4 - 1664*I*a^3 + (15*I*a^4 - 480*a^3 - 1184*I*a^2 + 968*a + 256*I)*b*x + 2152*a^2 - 24*(8*a^4 + 44*I*a^3 + (8*a^3 + 36*I*a^2 - 44*a - 17*I)*b*x - 80*a^2 - 61*I*a + 17)*\log(-b*x - a + \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 8*(2*I*b^4*x^4 + 6*b^3*x^3 - (10*a + 11*I)*b^2*x^2 - 2*I*a^4 + 78*a^3 + (22*a^2 + 54*I*a - 29)*b*x + 233*I*a^2 - 237*a - 80*I)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 1224*I*a - 256)/(b^5*x + (a + I)*b^4)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2)*x**3,x)

[Out] $-I*(\text{Integral}(I*x**3/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*a*x**3/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(a**3*x**3/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*b*x**4/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(b**3*x**6/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*I*a**2*x**3/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*I*b**2*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*I*b**2*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*I*b**2*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \text{Integral}(-3*I*b**2*x**5/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x)$

```
*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a*b**2*x**5/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a**2*b*x**4/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-6*I*a*b*x**4/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x))
```

Giac [A]

time = 0.46, size = 285, normalized size = 1.14

$$\frac{1}{8} \sqrt{(bx+a)^2+1} \left(\frac{1}{2} \left(\frac{12}{x} - \frac{16b^2-4b^4}{b^2} \right) - \frac{2i a^2 b^2 + 20 a b^4 + 19 b^6}{b^2} \right) x - \frac{2i a^2 b^2 - 44 a^2 b^4 - 93 a b^6 + 48 b^8}{b^2} - \frac{(8a^2 + 36a^2 - 44a - 17i) \log \left(\frac{x \left(x \sqrt{(bx+a)^2+1} \right)^2 ab + a^2 b + \left(x \sqrt{(bx+a)^2+1} \right)^2 b + 3 \left(x \sqrt{(bx+a)^2+1} \right) a^2 b + 2i \left(x \sqrt{(bx+a)^2+1} \right)^2 b + 2i a^2 b + 4 \left(x \sqrt{(bx+a)^2+1} \right) a b - \left(x \sqrt{(bx+a)^2+1} \right) b}{8 b^2} \right)}{8 b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^3,x, algorithm="giac")
```

```
[Out] -1/8*sqrt((b*x + a)^2 + 1)*((2*x*(I*x/b - (I*a*b^11 - 4*b^11)/b^13) - (-2*I*a^2*b^10 + 20*a*b^10 + 19*I*b^10)/b^13)*x - (2*I*a^3*b^9 - 44*a^2*b^9 - 93*I*a*b^9 + 48*b^9)/b^13) - 1/8*(8*a^3 + 36*I*a^2 - 44*a - 17*I)*log(3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b + a^3*b + (x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) + 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^2*abs(b) + 2*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b + 2*I*a^2*b + 4*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a*abs(b) - a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/ (b^3*abs(b))
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3 (1 + a \operatorname{li} + b x \operatorname{li})^3}{((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^3*(a*li + b*x*li + 1)^3)/((a + b*x)^2 + 1)^(3/2),x)
```

```
[Out] int((x^3*(a*li + b*x*li + 1)^3)/((a + b*x)^2 + 1)^(3/2), x)
```


3.182 $\int e^{3i \operatorname{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=227

$$\frac{(11i + 18a - 6ia^2) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{2b^3} + \frac{(11i + 18a - 6ia^2) \sqrt{1 - ia - ibx} (1 + ia + ibx)^{3/2}}{6b^3} - i \left(\frac{11i + 18a - 6ia^2}{2b^3} \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx} \right)$$

[Out] $\frac{1}{2}*(11-18*I*a-6*a^2)*\operatorname{arcsinh}(b*x+a)/b^3-I*(I+a)^2*(1+I*a+I*b*x)^{(5/2)}/b^3/(1-I*a-I*b*x)^{(1/2)}+1/6*(11*I+18*a-6*I*a^2)*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^3+1/3*I*(1+I*a+I*b*x)^{(5/2)}*(1-I*a-I*b*x)^{(1/2)}/b^3+1/2*(11*I+18*a-6*I*a^2)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3$

Rubi [A]

time = 0.12, antiderivative size = 227, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 91, 81, 52, 55, 633, 221}

$$\frac{(-6ia^2 + 18a + 11i) \sqrt{-ia - ibx + 1} (ia + ibx + 1)^{3/2}}{6b^3} + \frac{(-6ia^2 + 18a + 11i) \sqrt{-ia - ibx + 1} \sqrt{ia + ibx + 1}}{2b^3} + \frac{(-6a^2 - 18ia + 11) \sinh^{-1}(a + bx)}{2b^3} + \frac{i \sqrt{-ia - ibx + 1} (ia + ibx + 1)^{5/2}}{3b^3} - \frac{i(a + i)^2 (ia + ibx + 1)^{5/2}}{b^5 \sqrt{-ia - ibx + 1}}$$

Antiderivative was successfully verified.

[In] `Int[E^((3*I)*ArcTan[a + b*x])*x^2,x]`

[Out] $((11*I + 18*a - (6*I)*a^2)*\operatorname{Sqrt}[1 - I*a - I*b*x]*\operatorname{Sqrt}[1 + I*a + I*b*x])/(2*b^3) + ((11*I + 18*a - (6*I)*a^2)*\operatorname{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)})/(6*b^3) - (I*(I + a)^2*(1 + I*a + I*b*x)^{(5/2)})/(b^3*\operatorname{Sqrt}[1 - I*a - I*b*x]) + ((I/3)*\operatorname{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(5/2)})/b^3 + ((11 - (18*I)*a - 6*a^2)*\operatorname{ArcSinh}[a + b*x])/(2*b^3)$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 55

`Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`

Rule 81

`Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 1))], x]`

2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 91

Int[((a_.) + (b_.)*(x_))²((c_.) + (d_.)*(x_))^(n_.)((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(b*c - a*d)²(c + d*x)^(n + 1)((e + f*x)^(p + 1)/(d²(d*e - c*f)*(n + 1))), x] - Dist[1/(d²(d*e - c*f)*(n + 1)), Int[(c + d*x)^(n + 1)(e + f*x)^pSimp[a²d²f*(n + p + 2) + b²c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b²d*(d*e - c*f)*(n + 1)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)²], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)²)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b² - 4*a*c)))^p), Subst[Int[Simp[1 - x²/(b² - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b²/c, 0]

Rule 5203

Int[E^{(ArcTan[(c_.)*((a_) + (b_.)*(x_))])}(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m((1 - I*a*c - I*b*c*x)^{(I*(n/2))}/(1 + I*a*c + I*b*c*x)^{(I*(n/2))}), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{3i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2(1+ia+ibx)^{3/2}}{(1-ia-ibx)^{3/2}} dx \\
&= -\frac{i(i+a)^2(1+ia+ibx)^{5/2}}{b^3\sqrt{1-ia-ibx}} - \frac{i \int \frac{(1+ia+ibx)^{3/2}((3-2ia)(i+a)b-b^2x)}{\sqrt{1-ia-ibx}} dx}{b^3} \\
&= -\frac{i(i+a)^2(1+ia+ibx)^{5/2}}{b^3\sqrt{1-ia-ibx}} + \frac{i\sqrt{1-ia-ibx}(1+ia+ibx)^{5/2}}{3b^3} + \frac{(11-18ia-6a^2)}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}}{6b^3} - \frac{i(i+a)^2(1+ia+ibx)^{5/2}}{b^3\sqrt{1-ia-ibx}} + \frac{(11-18ia-6a^2)}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}}{6b^3} \\
&= \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11i+18a-6ia^2)\sqrt{1-ia-ibx}}{6b^3}
\end{aligned}$$

Mathematica [A]

time = 0.17, size = 160, normalized size = 0.70

$$\frac{\sqrt{1+ia+ibx}(52i-53ia^2-2a^3+19bx+7ib^2x^2-2b^3x^3+a(103-16ibx))}{6b^3\sqrt{-i(i+a+bx)}} + \frac{(-1)^{3/4}(-11+18ia+6a^2)\sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{\sqrt{-ib}b^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])*x^2,x]

[Out] (Sqrt[1 + I*a + I*b*x]*(52*I - (53*I)*a^2 - 2*a^3 + 19*b*x + (7*I)*b^2*x^2 - 2*b^3*x^3 + a*(103 - (16*I)*b*x)))/(6*b^3*Sqrt[(-I)*(I + a + b*x)]) + ((-1)^(3/4)*(-11 + (18*I)*a + 6*a^2)*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/(Sqrt[(-I)*b]*b^(5/2))

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1784 vs. 2(179) = 358.

time = 0.16, size = 1785, normalized size = 7.86

method	result
risch	$-\frac{i(2b^2x^2-2abx-9ibx+2a^2+27ia-28)\sqrt{b^2x^2+2abx+a^2+1}}{6b^3} + \frac{11 \ln\left(\frac{b^2x+ab+\sqrt{b^2x^2+2abx+a^2+1}}{\sqrt{b^2}}\right)}{2b^2\sqrt{b^2}}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^2,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & -I*b^3*(1/3*x^4/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-7/3*a/b*(1/2*x^3/b^2/(b^2 \\ & *x^2+2*a*b*x+a^2+1)^(1/2)-5/2*a/b*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3* \\ & a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+ \\ & 1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a \\ & ^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2) \\ &)/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2 \\ & *b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))-3/2 \\ & *(a^2+1)/b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a \\ & *b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+ \\ & 2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2 \\ & +1)^(1/2)))/(b^2)^(1/2))-4/3*(a^2+1)/b^2*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(\\ & 1/2)-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b \\ & *x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2* \\ & a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1 \\ &)^(1/2)))/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2 \\ & *a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2 \\ &))) -3*(1+I*a)*b^2*(1/2*x^3/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/2*a/b*(x^2/ \\ & b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/ \\ & 2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(\\ & a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2) \\ & ^{(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1/b^2/(b \\ & ^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2) \\ & /{(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))-3/2*(a^2+1)/b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a \\ & ^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b) \\ & /{(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a \\ & *b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+3*I*(1+I*a)^2* \\ & b*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2 \\ & +1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(\\ & 4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b \\ &)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1 \\ & /b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a \\ & ^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))+(1+I*a)^3*(-x/b^2/(b^2*x^2+2*a*b*x+ \\ & a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b) \\ &)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+ \end{aligned}$$

$a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}/(b^2)^{(1/2)}$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1608 vs. $2(155) = 310$.

time = 0.29, size = 1608, normalized size = 7.08

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^2,x, algorithm="maxima")
[Out] -35*I*a^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) -
  1/3*I*b*x^4/sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 265/6*I*(a^2 + 1)*a^3*x/((
  a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) + 7/6*I*a*x^3/s
  qrt(b^2*x^2 + 2*a*b*x + a^2 + 1) - 35/6*I*(a^2 + 1)*a^4/((a^2*b^2 - (a^2 +
  1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) - 61/6*I*(a^2 + 1)^2*a*x/((a^2
  *b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 2*(-I*a^3 - 3*a^
  2 + 3*I*a + 1)*a^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^
  2 + 1)) + 45*(I*a*b^2 + b^2)*a^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2
  + 2*a*b*x + a^2 + 1)*b^2) - 18*(I*a^2*b + 2*a*b - I*b)*a^3*x/((a^2*b^2 - (a
  ^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) - 35/6*I*a^2*x^2/(sqrt(b^
  2*x^2 + 2*a*b*x + a^2 + 1)*b) + 29/6*I*(a^2 + 1)^2*a^2/((a^2*b^2 - (a^2 + 1
  )*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) + (-I*a^3 - 3*a^2 + 3*I*a + 1)*
  (a^2 + 1)*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) -
  93/2*(I*a*b^2 + b^2)*(a^2 + 1)*a^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x
  ^2 + 2*a*b*x + a^2 + 1)*b^2) + 15*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*a*x/((a
  ^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) - 4/3*(-I*a^2
  - I)*x^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) - 3/2*(I*a*b^2 + b^2)*x^3/(s
  qrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2) + 35/2*I*a^3*arcsinh(2*(b^2*x + a*b)/
  sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^3 + 15/2*(I*a*b^2 + b^2)*(a^2 + 1)*a^
  3/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - 3*(I*
  a^2*b + 2*a*b - I*b)*(a^2 + 1)*a^2/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2
  + 2*a*b*x + a^2 + 1)*b^2) - (-I*a^3 - 3*a^2 + 3*I*a + 1)*(a^2 + 1)*a/((a^2*
  b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b) + 9/2*(I*a*b^2 +
  b^2)*(a^2 + 1)^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2
  + 1)*b^2) + 15/2*(I*a*b^2 + b^2)*a*x^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b
  ^3) - 3*(I*a^2*b + 2*a*b - I*b)*x^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^2)
  - 15/2*I*(a^2 + 1)*a*arcsinh(2*(b^2*x + a*b)/sqrt(-4*a^2*b^2 + 4*(a^2 + 1)
  *b^2))/b^3 - 9/2*(I*a*b^2 + b^2)*(a^2 + 1)^2*a/((a^2*b^2 - (a^2 + 1)*b^2)*s
  qrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) - 35/3*I*(a^2 + 1)*a^2/(sqrt(b^2*x^2
  + 2*a*b*x + a^2 + 1)*b^3) - 45/2*(I*a*b^2 + b^2)*a^2*arcsinh(2*(b^2*x + a*b
  )/sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^5 + 9*(I*a^2*b + 2*a*b - I*b)*a*arc
  sinh(2*(b^2*x + a*b)/sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2))/b^4 + (-I*a^3 - 3*
  a^2 + 3*I*a + 1)*arcsinh(2*(b^2*x + a*b)/sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)
  )/b^3 + 8/3*I*(a^2 + 1)^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b^3) + 9/2*(I*
```

$$a^2 b^2 + b^2) \cdot (a^2 + 1) \cdot \operatorname{arcsinh}(2 \cdot (b^2 x + a \cdot b) / \sqrt{-4 a^2 b^2 + 4 (a^2 + 1) b^2}) / b^5 + 15 \cdot (I \cdot a \cdot b^2 + b^2) \cdot (a^2 + 1) \cdot a / (\sqrt{b^2 x^2 + 2 a \cdot b \cdot x + a^2 + 1}) \cdot b^5 - 6 \cdot (I \cdot a^2 \cdot b + 2 a \cdot b - I \cdot b) \cdot (a^2 + 1) / (\sqrt{b^2 x^2 + 2 a \cdot b \cdot x + a^2 + 1}) \cdot b^4$$
Fricas [A]

time = 2.11, size = 174, normalized size = 0.77

$$\frac{-7i a^4 + 166 a^3 + (-7i a^3 + 159 a^2 + 249i a - 96) b x + 408i a^2 + 12(6 a^3 + (6 a^2 + 18i a - 11) b x + 24i a^2 - 29 a - 11i) \log\left(\frac{-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 + 1}}{24(b^2 x + (a+i)b^2)}\right) - 4(2i b^3 x^3 + 7 b^2 x^2 + 2i a^3 - (16a + 19i) b x - 53 a^2 - 103i a + 52) \sqrt{b^2 x^2 + 2 abx + a^2 + 1} - 345 a - 96i}{24(b^2 x + (a+i)b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^2,x, algorithm="fricas")

[Out] $\frac{1}{24} \cdot (-7 \cdot I \cdot a^4 + 166 \cdot a^3 + (-7 \cdot I \cdot a^3 + 159 \cdot a^2 + 249 \cdot I \cdot a - 96) \cdot b \cdot x + 408 \cdot I \cdot a^2 + 12 \cdot (6 \cdot a^3 + (6 \cdot a^2 + 18 \cdot I \cdot a - 11) \cdot b \cdot x + 24 \cdot I \cdot a^2 - 29 \cdot a - 11 \cdot I)) \cdot \log(-b \cdot x - a + \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1}) - 4 \cdot (2 \cdot I \cdot b^3 \cdot x^3 + 7 \cdot b^2 \cdot x^2 + 2 \cdot I \cdot a^3 - (16 \cdot a + 19 \cdot I) \cdot b \cdot x - 53 \cdot a^2 - 103 \cdot I \cdot a + 52) \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1} - 345 \cdot a - 96 \cdot I) / (b^4 \cdot x + (a + I) \cdot b^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a)**3/(1+(b*x+a)**2)**(3/2)*x**2,x)

[Out] $-I \cdot (\operatorname{Integral}(I \cdot x^{**2} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(-3 \cdot a \cdot x^{**2} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(a^{**3} \cdot x^{**2} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(-3 \cdot b \cdot x^{**3} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(b^{**3} \cdot x^{**5} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(-3 \cdot I \cdot a^{**2} \cdot x^{**2} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(-3 \cdot I \cdot b^{**2} \cdot x^{**4} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(-3 \cdot I \cdot a^{**2} \cdot x^{**2} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x) + \operatorname{Integral}(-3 \cdot I \cdot b^{**2} \cdot x^{**4} / (a^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + 2 \cdot a \cdot b \cdot x \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + b^{**2} \cdot x^{**2} \cdot \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}) + \sqrt{a^{**2} + 2 \cdot a \cdot b \cdot x + b^{**2} \cdot x^{**2} + 1}), x)$

2 + 1)), x) + Integral(3*a*b**2*x**4/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a**2*b*x**3/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-6*I*a*b*x**3/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x))

Giac [A]

time = 0.47, size = 243, normalized size = 1.07

$$\frac{1}{6} \sqrt{(bx+a)^2+1} \left(x \frac{2bx-2iab^2-9b^3}{b^3} - \frac{-2ia^2b^2+27ab^3+28i6^3}{b^3} \right) + \frac{(6a^2+18ia-11) \log \left(3 \left(x|b| - \sqrt{(bx+a)^2+1} \right)^3 ab + a^3b + \left(x|b| - \sqrt{(bx+a)^2+1} \right)^3 |b| + 3 \left(x|b| - \sqrt{(bx+a)^2+1} \right)^2 a^2|b| + 2a \left(x|b| - \sqrt{(bx+a)^2+1} \right)^3 b + 2ia^2b + 4 \left(x|b| - i\sqrt{(bx+a)^2+1} \right) a|b| - ab - \left(x|b| - \sqrt{(bx+a)^2+1} \right) |b| \right)}{6^3|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x^2,x, algorithm="giac")

[Out] -1/6*sqrt((b*x + a)^2 + 1)*(x*(2*I*x/b - (2*I*a*b^6 - 9*b^6)/b^8) - (-2*I*a^2*b^5 + 27*a*b^5 + 28*I*b^5)/b^8) + 1/6*(6*a^2 + 18*I*a - 11)*log(3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b + a^3*b + (x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) + 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^2*abs(b) + 2*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b + 2*I*a^2*b + 4*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a*abs(b) - a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^2*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2 (1 + a \operatorname{li} + b x \operatorname{li})^3}{((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*li + b*x*li + 1)^3)/((a + b*x)^2 + 1)^(3/2),x)

[Out] int((x^2*(a*li + b*x*li + 1)^3)/((a + b*x)^2 + 1)^(3/2), x)

3.183 $\int e^{3i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=163

$$\frac{3(3-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}}{2b^2} - \frac{(1-ia)(1+ia+ibx)}{b^2\sqrt{1-ia-ibx}}$$

[Out] $3/2*(3*I+2*a)*\text{arcsinh}(b*x+a)/b^2 - (1-I*a)*(1+I*a+I*b*x)^{(5/2)}/b^2 / (1-I*a-I*b*x)^{(1/2)} - 1/2*(3-2*I*a)*(1+I*a+I*b*x)^{(3/2)}*(1-I*a-I*b*x)^{(1/2)}/b^2 - 3/2*(3-2*I*a)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2$

Rubi [A]

time = 0.09, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5203, 79, 52, 55, 633, 221}

$$\frac{(1-ia)(ia+ibx+1)^{5/2}}{b^2\sqrt{-ia-ibx+1}} - \frac{(3-2ia)\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/2}}{2b^2} - \frac{3(3-2ia)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{2b^2} + \frac{3(2a+3i)\sinh^{-1}(a+bx)}{2b^2}$$

Antiderivative was successfully verified.

[In] `Int[E^((3*I)*ArcTan[a + b*x])*x,x]`

[Out] $(-3*(3 - (2*I)*a)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/(2*b^2) - ((3 - (2*I)*a)*\text{Sqrt}[1 - I*a - I*b*x]*(1 + I*a + I*b*x)^{(3/2)})/(2*b^2) - ((1 - I*a)*(1 + I*a + I*b*x)^{(5/2)})/(b^2*\text{Sqrt}[1 - I*a - I*b*x]) + (3*(3*I + 2*a)*\text{ArcSinh}[a + b*x])/(2*b^2)$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 55

`Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] :> Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`

Rule 79

`Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c`


```
*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^(p + 1), x]
, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{3i \tan^{-1}(a+bx)} x dx &= \int \frac{x(1+ia+ibx)^{3/2}}{(1-ia-ibx)^{3/2}} dx \\
&= -\frac{(1-ia)(1+ia+ibx)^{5/2}}{b^2 \sqrt{1-ia-ibx}} + \frac{(3i+2a) \int \frac{(1+ia+ibx)^{3/2}}{\sqrt{1-ia-ibx}} dx}{b} \\
&= -\frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}}{2b^2} - \frac{(1-ia)(1+ia+ibx)^{5/2}}{b^2 \sqrt{1-ia-ibx}} + \frac{(3(3i+2a))}{b} \\
&= -\frac{3(3-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^3}{2b^2} \\
&= -\frac{3(3-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^3}{2b^2} \\
&= -\frac{3(3-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^3}{2b^2} \\
&= -\frac{3(3-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^3}{2b^2} \\
&= -\frac{3(3-2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3-2ia)\sqrt{1-ia-ibx}(1+ia+ibx)^3}{2b^2}
\end{aligned}$$

Mathematica [A]

time = 0.12, size = 132, normalized size = 0.81

$$\frac{\sqrt{1+ia+ibx}(-14+15ia+a^2+5ibx-b^2x^2)}{2b^2\sqrt{-i(i+a+bx)}} + \frac{3\sqrt[4]{-1}(3i+2a)\sqrt{-ib}\sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{b^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])*x,x]

[Out] (Sqrt[1 + I*a + I*b*x]*(-14 + (15*I)*a + a^2 + (5*I)*b*x - b^2*x^2))/(2*b^2 *Sqrt[(-I)*(I + a + b*x)]) + (3*(-1)^(1/4)*(3*I + 2*a)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])]/Sqrt[(-I)*b])/b^(5/2)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1051 vs. 2(129) = 258.

time = 0.15, size = 1052, normalized size = 6.45

method	result
--------	--------

risch	$\frac{i(-bx+a+6i)\sqrt{b^2x^2+2abx+a^2+1}}{2b^2} + \frac{9i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{2b\sqrt{b^2}} + \frac{3a \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{b\sqrt{b^2}}$
default	$-ib^3 \frac{x^3}{2b^2\sqrt{b^2x^2+2abx+a^2+1}} - \frac{5a \frac{x^2}{b^2\sqrt{b^2x^2+2abx+a^2+1}} - \left(\frac{3a}{b^2\sqrt{b^2x^2+2abx+a^2+1}} \right)}{1}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x,x,method=_RETURNVERBOSE)`

[Out]
$$-I*b^3*(1/2*x^3/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/2*a/b*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))-3/2*(a^2+1)/b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2)))-3*(1+I*a)*b^2*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))+3*I*(1+I*a)^2*b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2)))+(1+I*a)^3*(-1/b^2/(b^2$$

$$*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1108 vs. $2(113) = 226$.
time = 0.28, size = 1108, normalized size = 6.80

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x,x, algorithm="maxima")

[Out] $15*I*a^4*b*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 31/2*I*(a^2 + 1)*a^2*b*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 1/2*I*b*x^3/\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 5/2*I*(a^2 + 1)*a^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 6*(I*a^2*b + 2*a*b - I*b)*a^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 18*(I*a*b^2 + b^2)*a^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 3/2*I*(a^2 + 1)^2*b*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + (-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 5/2*I*a*x^2/\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} - 3/2*I*(a^2 + 1)^2*a/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + (-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 3*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 15*(I*a*b^2 + b^2)*(a^2 + 1)*a*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 15/2*I*a^2*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^2 - 3*(I*a*b^2 + b^2)*(a^2 + 1)*a^2/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 3*(I*a^2*b + 2*a*b - I*b)*(a^2 + 1)*a/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 3*(I*a*b^2 + b^2)*x^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 3/2*(-I*a^2 - I)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^2 + 5*I*(a^2 + 1)*a/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 9*(I*a*b^2 + b^2)*a*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^4 - 3*(I*a^2*b + 2*a*b - I*b)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^3 + (I*a^3 + 3*a^2 - 3*I*a - 1)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 6*(I*a*b^2 + b^2)*(a^2 + 1)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b^4$

Fricas [A]

time = 3.04, size = 136, normalized size = 0.83

$$\frac{3i a^3 + (3i a^2 - 44 a - 32i) b x - 47 a^2 - 12((2 a + 3i) b x + 2 a^2 + 5i a - 3) \log\left(\frac{-b x - a + \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{8(b^2 x + (a + i) b^2)}\right) - 4 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} (i b^2 x^2 - i a^2 + 5 b x + 15 a + 14 i) - 76 i a + 32}{8(b^2 x + (a + i) b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x,x, algorithm="fricas")
[Out] 1/8*(3*I*a^3 + (3*I*a^2 - 44*a - 32*I)*b*x - 47*a^2 - 12*((2*a + 3*I)*b*x +
  2*a^2 + 5*I*a - 3)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*s
qrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(I*b^2*x^2 - I*a^2 + 5*b*x + 15*a + 14*I)
- 76*I*a + 32)/(b^3*x + (a + I)*b^2)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2)*x,x)
[Out] -I*(Integral(I*x/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(
a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2
+ 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*a*x/(a**2*s
qrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x
**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a
b*x + b**2*x**2 + 1)), x) + Integral(a**3*x/(a**2*sqrt(a**2 + 2*a*b*x + b**
2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt
(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x
) + Integral(-3*b*x**2/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x
*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**
2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(b**3*x**
4/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x
+ b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a*
**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*I*a**2*x/(a**2*sqrt(a**2 +
2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) +
b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2
*x**2 + 1)), x) + Integral(-3*I*b**2*x**3/(a**2*sqrt(a**2 + 2*a*b*x + b**2*
x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a
**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x)
+ Integral(3*a*b**2*x**3/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b
*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b
**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a**2
*b*x**2/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*
a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + s
qrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-6*I*a*b*x**2/(a**2*sq
rt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2
+ 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*
x + b**2*x**2 + 1)), x))
```

Giac [A]

time = 0.45, size = 209, normalized size = 1.28

$$-\frac{1}{2} \sqrt{(bx+a)^2+1} \left(\frac{ix}{b} - \frac{-ia^2+6b^2}{b^2} \right) - \frac{(2a+3i) \log \left(3 \left(|x| - \sqrt{(bx+a)^2+1} \right)^2 ab + a^2b + \left(|x| - \sqrt{(bx+a)^2+1} \right)^3 |b| + 3 \left(|x| - \sqrt{(bx+a)^2+1} \right)^2 a^2 |b| + 2a \left(|x| - \sqrt{(bx+a)^2+1} \right)^3 b + 2ia^2b + 4 \left(i|x| - i\sqrt{(bx+a)^2+1} \right) a|b| - ab - \left(|x| - \sqrt{(bx+a)^2+1} \right) |b| \right)}{2|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)*x,x, algorithm="giac")
```

```
[Out] -1/2*sqrt((b*x + a)^2 + 1)*(I*x/b + (-I*a*b^2 + 6*b^2)/b^4) - 1/2*(2*a + 3*I)*log(3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b + a^3*b + (x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) + 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^2*abs(b) + 2*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b + 2*I*a^2*b + 4*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a*abs(b) - a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b*abs(b))
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x(1 + a \operatorname{li} + b x \operatorname{li})^3}{((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x*(a*1i + b*x*1i + 1)^3)/((a + b*x)^2 + 1)^(3/2),x)
```

```
[Out] int((x*(a*1i + b*x*1i + 1)^3)/((a + b*x)^2 + 1)^(3/2), x)
```

3.184 $\int e^{3i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=94

$$-\frac{3i\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{b} - \frac{2i(1+ia+ibx)^{3/2}}{b\sqrt{1-ia-ibx}} - \frac{3\sinh^{-1}(a+bx)}{b}$$

[Out] $-3*\text{arcsinh}(b*x+a)/b-2*I*(1+I*a+I*b*x)^{(3/2)}/b/(1-I*a-I*b*x)^{(1/2)}-3*I*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b$

Rubi [A]

time = 0.03, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5201, 49, 52, 55, 633, 221}

$$\frac{2i(ia+ibx+1)^{3/2}}{b\sqrt{-ia-ibx+1}} - \frac{3i\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{b} - \frac{3\sinh^{-1}(a+bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a + b*x]),x]

[Out] $((-3*I)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/b - ((2*I)*(1 + I*a + I*b*x)^{(3/2)})/(b*\text{Sqrt}[1 - I*a - I*b*x]) - (3*\text{ArcSinh}[a + b*x])/b$

Rule 49

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b

+ d, 0] && GtQ[a + c, 0]

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5201

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_.)), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{3i \tan^{-1}(a+bx)} dx &= \int \frac{(1 + ia + ibx)^{3/2}}{(1 - ia - ibx)^{3/2}} dx \\
 &= -\frac{2i(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} - 3 \int \frac{\sqrt{1 + ia + ibx}}{\sqrt{1 - ia - ibx}} dx \\
 &= -\frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - \frac{2i(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} - 3 \int \frac{1}{\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}} dx \\
 &= -\frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - \frac{2i(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} - 3 \int \frac{1}{\sqrt{(1 - ia)(1 + ia) + 2ibx}} dx \\
 &= -\frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - \frac{2i(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} - \frac{3 \operatorname{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{x^2}{4b^2}}} dx, \frac{1 + ia + ibx}{2b} \right)}{2b^2} \\
 &= -\frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - \frac{2i(1 + ia + ibx)^{3/2}}{b\sqrt{1 - ia - ibx}} - \frac{3 \sinh^{-1}(a + bx)}{b}
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 45, normalized size = 0.48

$$\frac{\sqrt{1 + (a + bx)^2} \left(-i + \frac{4}{i + a + bx}\right)}{b} - \frac{3 \sinh^{-1}(a + bx)}{b}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((3*I)*ArcTan[a + b*x]),x]

[Out] (Sqrt[1 + (a + b*x)^2]*(-I + 4/(I + a + b*x)))/b - (3*ArcSinh[a + b*x])/b

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 616 vs. 2(76) = 152.

time = 0.14, size = 617, normalized size = 6.56

method	result
risch	$-\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b} - \frac{3\ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{\sqrt{b^2}} + \frac{4\sqrt{\left(x + \frac{i+a}{b}\right)^2 b^2 - 2ib\left(x + \frac{i+a}{b}\right)}}{b^2\left(x + \frac{i+a}{b}\right)}$
default	$-ib^3 \left(\frac{x^2}{b^2\sqrt{b^2x^2 + 2abx + a^2 + 1}} - \frac{3a}{b^2\sqrt{b^2x^2 + 2abx + a^2 + 1}} - \frac{a}{b^2\sqrt{b^2x^2 + 2abx + a^2 + 1}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out]
$$-I*b^3*(x^2/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a/b*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))-2*(a^2+1)/b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))-3*(1+I*a)*b^2*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))+3*I*(1+I*a)^2*b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+2*(1+I*a)^3*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 736 vs. 2(66) = 132.

time = 0.29, size = 736, normalized size = 7.83

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2),x, algorithm="maxima")

[Out]
$$\begin{aligned} & -6*I*a^3*b^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) \\ & + 5*I*(a^2 + 1)*a*b^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x \\ & + a^2 + 1}) - I*(a^2 + 1)*a^2*b/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + \\ & 2*a*b*x + a^2 + 1}) + 6*(I*a*b^2 + b^2)*a^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{ \\ & b^2*x^2 + 2*a*b*x + a^2 + 1}) - 3*(I*a^2*b + 2*a*b - I*b)*a*b*x/((a^2*b^2 \\ & - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + (I*a^3 + 3*a^2 - 3* \\ & I*a - 1)*b^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) \\ & - I*b*x^2/\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} - 3*(I*a^2*b + 2*a*b - I*b)*a \\ & ^2/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - (-I*a^3 \\ & - 3*a^2 + 3*I*a + 1)*a*b/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x \\ & + a^2 + 1}) - 3*(I*a*b^2 + b^2)*(a^2 + 1)*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{ \\ & b^2*x^2 + 2*a*b*x + a^2 + 1}) + 3*I*a*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{-4*a^2* \\ & b^2 + 4*(a^2 + 1)*b^2})/b + 3*(I*a*b^2 + b^2)*(a^2 + 1)*a/((a^2*b^2 - (a^2 \\ & + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b - 2*(I*a^2 + I)/(\sqrt{b^2*x^2 \\ & + 2*a*b*x + a^2 + 1})*b - 3*(I*a*b^2 + b^2)*\operatorname{arcsinh}(2*(b^2*x + a*b)/\sqrt{ \\ & -4*a^2*b^2 + 4*(a^2 + 1)*b^2})/b^3 + 3*(I*a^2*b + 2*a*b - I*b)/(\sqrt{b^2*x^2 \\ & + 2*a*b*x + a^2 + 1})*b^2) \end{aligned}$$

Fricas [A]

time = 2.49, size = 99, normalized size = 1.05

$$\frac{(-i a + 8) b x - i a^2 + 6 (b x + a + i) \log \left(-b x - a + \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \right) - 2 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} (i b x + i a - 5) + 9 a + 8 i}{2 (b^2 x + (a + i) b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2),x, algorithm="fricas")

[Out]
$$\begin{aligned} & 1/2*((-I*a + 8)*b*x - I*a^2 + 6*(b*x + a + I)*\log(-b*x - a + \sqrt{b^2*x^2 + \\ & 2*a*b*x + a^2 + 1}) - 2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(I*b*x + I*a - 5 \\ &) + 9*a + 8*I)/(b^2*x + (a + I)*b) \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2),x)

[Out]
$$\begin{aligned} & -I*(\operatorname{Integral}(I/(a**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}) + 2*a*b*x*\sqrt{a* \\ & *2 + 2*a*b*x + b**2*x**2 + 1}) + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + \\ & 1} + \sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1}), x) + \operatorname{Integral}(-3*a/(a**2*\sqrt{ \\ & a**2 + 2*a*b*x + b**2*x**2 + 1}) + 2*a*b*x*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + \\ & 1} + b**2*x**2*\sqrt{a**2 + 2*a*b*x + b**2*x**2 + 1} + \sqrt{a**2 + 2*a*b*x \end{aligned}$$

```

+ b**2*x**2 + 1)), x) + Integral(a**3/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2
+ 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2
+ 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + In
tegral(-3*I*a**2/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(
a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2
+ 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*b*x/(a**2*s
qrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x*
**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*
b*x + b**2*x**2 + 1)), x) + Integral(b**3*x**3/(a**2*sqrt(a**2 + 2*a*b*x +
b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*s
qrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1))
, x) + Integral(-3*I*b**2*x**2/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) +
2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b
*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(
3*a*b**2*x**2/(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**
2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 +
1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a**2*b*x/(a**2*
sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x
**2 + 1) + b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a
*b*x + b**2*x**2 + 1)), x) + Integral(-6*I*a*b*x/(a**2*sqrt(a**2 + 2*a*b*x
+ b**2*x**2 + 1) + 2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**2
*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1
)), x))

```

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 180 vs. $2(66) = 132$.

time = 0.46, size = 180, normalized size = 1.91

$$\frac{\log\left(3\left(x|b| - \sqrt{(bx+a)^2+1}\right)^{2ab+a^2b+\left(x|b| - \sqrt{(bx+a)^2+1}\right)^3|b|+3\left(x|b| - \sqrt{(bx+a)^2+1}\right)a^2|b|+2i\left(x|b| - \sqrt{(bx+a)^2+1}\right)^2b+2i a^2b+4\left(i x|b| - i\sqrt{(bx+a)^2+1}\right)a|b|-ab-\left(x|b| - \sqrt{(bx+a)^2+1}\right)|b|\right)}{|b|} - \frac{i\sqrt{(bx+a)^2+1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2),x, algorithm="giac")

[Out] $\log(3*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a*b + a^3*b + (x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^3*\text{abs}(b) + 3*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*a^2*\text{abs}(b) + 2*I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*b + 2*I*a^2*b + 4*(I*x*\text{abs}(b) - I*\sqrt{(b*x + a)^2 + 1})*a*\text{abs}(b) - a*b - (x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1}))*\text{abs}(b))/\text{abs}(b) - I*\sqrt{(b*x + a)^2 + 1}/b$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(1 + a \operatorname{li} + b x \operatorname{li})^3}{((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*1i + b*x*1i + 1)^3/((a + b*x)^2 + 1)^(3/2),x)
```

```
[Out] int((a*1i + b*x*1i + 1)^3/((a + b*x)^2 + 1)^(3/2), x)
```

$$3.185 \quad \int \frac{e^{3i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=134

$$\frac{4\sqrt{1+ia+ibx}}{(1-ia)\sqrt{1-ia-ibx}} - i \sinh^{-1}(a+bx) - \frac{2(i-a)^{3/2} \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i+a)^{3/2}}$$

[Out] $-I*\operatorname{arcsinh}(b*x+a)-2*(I-a)^{(3/2)}*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)/(1-I*a-I*b*x)^{(1/2)})/(I+a)^{(3/2)+4*(1+I*a+I*b*x)^{(1/2)/(1-I*a)/(1-I*a-I*b*x)^{(1/2)})}$

Rubi [A]

time = 0.10, antiderivative size = 134, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 100, 163, 55, 633, 221, 95, 214}

$$\frac{4\sqrt{ia+ibx+1}}{(1-ia)\sqrt{-ia-ibx+1}} - i \sinh^{-1}(a+bx) - \frac{2(-a+i)^{3/2} \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(a+i)^{3/2}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)*\operatorname{ArcTan}[a+b*x])}/x,x]$

[Out] $(4*\operatorname{Sqrt}[1+I*a+I*b*x])/((1-I*a)*\operatorname{Sqrt}[1-I*a-I*b*x]) - I*\operatorname{ArcSinh}[a+b*x] - (2*(I-a)^{(3/2)}*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])])/(I+a)^{(3/2)}$

Rule 55

$\operatorname{Int}[1/(\operatorname{Sqrt}[(a_) + (b_)*(x_)]*\operatorname{Sqrt}[(c_) + (d_)*(x_)]), x_Symbol] \rightarrow \operatorname{Int}[1/\operatorname{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x \ \&\& \operatorname{EqQ}[b + d, 0] \ \&\& \operatorname{GtQ}[a + c, 0]$

Rule 95

$\operatorname{Int}[(((a_) + (b_)*(x_))^{(m_)}*((c_) + (d_)*(x_))^{(n_)})/((e_) + (f_)*(x_))], x_Symbol] \rightarrow \operatorname{With}\{q = \operatorname{Denominator}[m]\}, \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q)}, x], x, (a + b*x)^{(1/q)/(c + d*x)^{(1/q)}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \operatorname{EqQ}[m + n + 1, 0] \ \&\& \operatorname{RationalQ}[n] \ \&\& \operatorname{LtQ}[-1, m, 0] \ \&\& \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 100

$\operatorname{Int}[((a_) + (b_)*(x_))^{(m_)}*((c_) + (d_)*(x_))^{(n_)}*((e_) + (f_)*(x_))^{(p_)}], x_Symbol] \rightarrow \operatorname{Simp}[(b*c - a*d)*(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}$

```

*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m
+ 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(
n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*
(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a,
b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2
*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 163

```

Int[(((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_))*((g_.) + (h_.)*(x_
)))/((a_.) + (b_.)*(x_)), x_Symbol] := Dist[h/b, Int[(c + d*x)^n*(e + f*x)^
p, x], x] + Dist[(b*g - a*h)/b, Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x]
, x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]

```

Rule 214

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

```

Rule 221

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

```

Rule 633

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

```

Rule 5203

```

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_)^(m_.),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{(1+ia+ibx)^{3/2}}{x(1-ia-ibx)^{3/2}} dx \\
&= \frac{4\sqrt{1+ia+ibx}}{(1-ia)\sqrt{1-ia-ibx}} - \frac{2 \int \frac{\frac{1}{2}i(i-a)^2b - \frac{1}{2}(1-ia)b^2x}{x\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx}{(i+a)b} \\
&= \frac{4\sqrt{1+ia+ibx}}{(1-ia)\sqrt{1-ia-ibx}} - \frac{(i-a)^2 \int \frac{1}{x\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx}{1-ia} - (ib) \int \frac{1}{\sqrt{1-ia-ibx}} dx \\
&= \frac{4\sqrt{1+ia+ibx}}{(1-ia)\sqrt{1-ia-ibx}} - \frac{(2(i-a)^2) \text{Subst}\left(\int \frac{1}{-1-ia-(-1+ia)x^2} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}}\right)}{1-ia} \\
&= \frac{4\sqrt{1+ia+ibx}}{(1-ia)\sqrt{1-ia-ibx}} - \frac{2(i-a)^{3/2} \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i+a)^{3/2}} - i \text{Subst}\left(\int \frac{1}{\sqrt{1-ia-ibx}} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}}\right) \\
&= \frac{4\sqrt{1+ia+ibx}}{(1-ia)\sqrt{1-ia-ibx}} - i \sinh^{-1}(a+bx) - \frac{2(i-a)^{3/2} \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i+a)^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.53, size = 196, normalized size = 1.46

$$\frac{2 \left(\frac{2i\sqrt{1+ia+ibx}}{\sqrt{-i(i+a+bx)}} + \frac{\sqrt[4]{-1}^{(i+a)(-ib)^{3/2} \sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}}{b^{3/2}} + \frac{\sqrt{-1-ia}^{(-i+a) \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}}{\sqrt{-1+ia}} \right)}{i+a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((3*I)*ArcTan[a + b*x])/x,x]`

```
[Out] (2*(((2*I)*Sqrt[1 + I*a + I*b*x])/Sqrt[(-I)*(I + a + b*x)] + ((-1)^(1/4)*(I + a)*((-I)*b)^(3/2)*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/b^(3/2) + (Sqrt[-1 - I*a]*(-I + a)*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]])/Sqrt[-1 + I*a]))/(I + a)
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 484 vs. $2(104) = 208$.

time = 0.12, size = 485, normalized size = 3.62

method	result
default	$-ib^3 \left(-\frac{x}{b^2 \sqrt{b^2 x^2 + 2abx + a^2 + 1}} - \frac{a \left(-\frac{1}{b^2 \sqrt{b^2 x^2 + 2abx + a^2 + 1}} - \frac{2a(2b^2 x + 2ab)}{b(4b^2(a^2 + 1) - 4a^2 b^2) \sqrt{b^2 x^2 + 2abx + a^2 + 1}} \right)}{b} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x,x,method=_RETURNVERBOSE)
```

```
[Out] -I*b^3*(-x/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-a/b*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+1/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-3*(1+I*a)*b^2*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))+6*I*(1+I*a)^2*b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+(-I*a^3-3*a^2+3*I*a+1)*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))
```

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 733 vs. $2(90) = 180$.
time = 0.28, size = 733, normalized size = 5.47

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x,x, algorithm="maxima")
```

```
[Out] 2*I*a^2*b^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) + (-I*a^2 - I)*b^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) + (-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)) + I*(a^2 + 1)*a*b^2/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) + (-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*b^2/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)) - 3*(I*a*b^2 + b^2)*a*b*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) + 3*(I*a^2*b + 2*a*b - I*b)*b^2*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 3*(I*a*b^2 + b^2)*a^2/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) + 3*(I*a^2*b + 2*a*b - I*b)*a*b/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - (-I*a^3 - 3*a^2 + 3*I*a + 1)*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(3/2) + (-I*a^3 - 3*a
```


$$\frac{\sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{\sqrt{b^2 x^2 + 2 a b x + a^2 + 1}} + \frac{3(I a b^2 + b^2)}{\sqrt{b^2 x^2 + 2 a b x + a^2 + 1} b^2} - \frac{I \operatorname{arcsinh}(2(b^2 x + a b))}{\sqrt{-4 a^2 b^2 + 4(a^2 + 1)b^2}}$$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 356 vs. $2(90) = 180$.

time = 1.92, size = 356, normalized size = 2.66

$$\frac{((a + 1)(b x^2 + 2 a x - 1) \sqrt{\frac{a^2 - 3 a^2 - 3 a + 1}{a^2 + 3 a^2 - 3 a - 1}} \log\left(\frac{(a - 1) b \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \sqrt{\frac{a^2 - 3 a^2 - 3 a + 1}{a^2 + 3 a^2 - 3 a - 1}}}{a^2 + 3 a^2 - 3 a - 1}\right) - ((a + 1)(b x^2 + 2 a x - 1) \sqrt{\frac{a^2 - 3 a^2 - 3 a + 1}{a^2 + 3 a^2 - 3 a - 1}} \log\left(\frac{(a - 1) b \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} \sqrt{\frac{a^2 - 3 a^2 - 3 a + 1}{a^2 + 3 a^2 - 3 a - 1}}}{a^2 + 3 a^2 - 3 a - 1}\right) + 4 b x + ((-a + 1)(b x - a^2 + 2 a + 1) \log(-b x - a + \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}) + 4 a + 4 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1})}{(a + 1)(b x^2 + 2 a x - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x,x, algorithm="fricas")

[Out] $-\left(\frac{((a + I) b x + a^2 + 2 I a - 1) \sqrt{-(a^3 - 3 I a^2 - 3 a + I)}}{(a^3 + 3 I a^2 - 3 a - I)} \log\left(-\frac{(a - I) b x - \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{(a - I)} - \frac{(I a^2 - 2 a - I) \sqrt{-(a^3 - 3 I a^2 - 3 a + I)}}{(a^3 + 3 I a^2 - 3 a - I)}\right) - \frac{((a + I) b x + a^2 + 2 I a - 1) \sqrt{-(a^3 - 3 I a^2 - 3 a + I)}}{(a^3 + 3 I a^2 - 3 a - I)} \log\left(-\frac{(a - I) b x - \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{(a - I)} - \frac{(-I a^2 + 2 a + I) \sqrt{-(a^3 - 3 I a^2 - 3 a + I)}}{(a^3 + 3 I a^2 - 3 a - I)}\right) + 4 b x + ((-I a + 1) b x - I a^2 + 2 a + I) \log(-b x - a + \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}) + 4 a + 4 \sqrt{b^2 x^2 + 2 a b x + a^2 + 1}}{(a + I) b x + a^2 + 2 I a - 1}\right)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)^(3/2)/x,x)

[Out] $-I \left(\operatorname{Integral}\left(\frac{I}{(a^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} + x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}\right), x\right) + \operatorname{Integral}\left(\frac{-3 a}{(a^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} + b^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}\right), x\right) + \operatorname{Integral}\left(\frac{a^3}{(a^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} + b^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}\right), x\right) + \operatorname{Integral}\left(\frac{-3 I a^2}{(a^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} + b^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}\right), x\right) + \operatorname{Integral}\left(\frac{-3 b x}{(a^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}} + b^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}\right), x\right) + \operatorname{Integral}(b^2 x \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}, x)$

```

3*x**3/(a**2*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**2*sqrt(a**2
+ 2*a*b*x + b**2*x**2 + 1) + b**2*x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)
+ x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*I*b**2*x**2/(a
**2*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**2*sqrt(a**2 + 2*a*b*x
+ b**2*x**2 + 1) + b**2*x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x*sqrt
(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a*b**2*x**2/(a**2*x*sqrt
(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**2*sqrt(a**2 + 2*a*b*x + b**2*x*
*2 + 1) + b**2*x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x*sqrt(a**2 + 2*
a*b*x + b**2*x**2 + 1)), x) + Integral(3*a**2*b*x/(a**2*x*sqrt(a**2 + 2*a*b
*x + b**2*x**2 + 1) + 2*a*b*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**
2*x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x*sqrt(a**2 + 2*a*b*x + b**2*
x**2 + 1)), x) + Integral(-6*I*a*b*x/(a**2*x*sqrt(a**2 + 2*a*b*x + b**2*x**
2 + 1) + 2*a*b*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**3*sqrt(a
**2 + 2*a*b*x + b**2*x**2 + 1) + x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x
))

```

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 252 vs. 2(90) = 180.
time = 0.56, size = 252, normalized size = 1.88

$$\frac{i b \log \left(-3 \left(x |b| - \sqrt{(b x + a)^2 + 1} \right)^2 a b - a^3 b - \left(x |b| - \sqrt{(b x + a)^2 + 1} \right) |b| - 3 \left(x |b| - \sqrt{(b x + a)^2 + 1} \right) a^2 |b| - 2 i \left(x |b| - \sqrt{(b x + a)^2 + 1} \right)^3 b - 2 i a^2 b - 4 \left(i x |b| - i \sqrt{(b x + a)^2 + 1} \right) a |b| + a b + \left(x |b| - \sqrt{(b x + a)^2 + 1} \right) |b| \right)}{3 |b|} \frac{(i a^2 + 2 a - i) \log \left(\frac{-2 x |b| + 2 \sqrt{(b x + a)^2 + 1} - 2 \sqrt{a^2 + 1}}{-2 x |b| + 2 \sqrt{(b x + a)^2 + 1} + 2 \sqrt{a^2 + 1}} \right)}{\sqrt{a^2 + 1} (a + i)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x,x, algorithm="giac")
```

```
[Out] 1/3*I*b*log(-3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b - a^3*b - (x*abs(b)
- sqrt((b*x + a)^2 + 1))^3*abs(b) - 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a
^2*abs(b) - 2*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b - 2*I*a^2*b - 4*(I*x
*abs(b) - I*sqrt((b*x + a)^2 + 1))*a*abs(b) + a*b + (x*abs(b) - sqrt((b*x +
a)^2 + 1))*abs(b))/abs(b) - (I*a^2 + 2*a - I)*log(abs(-2*x*abs(b) + 2*sqrt
((b*x + a)^2 + 1) - 2*sqrt(a^2 + 1))/abs(-2*x*abs(b) + 2*sqrt((b*x + a)^2 +
1) + 2*sqrt(a^2 + 1)))/(sqrt(a^2 + 1)*(a + I))
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(1 + a \operatorname{li} + b x \operatorname{li})^3}{x ((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*1i + b*x*1i + 1)^3/(x*((a + b*x)^2 + 1)^(3/2)),x)
```

```
[Out] int((a*1i + b*x*1i + 1)^3/(x*((a + b*x)^2 + 1)^(3/2)), x)
```

$$3.186 \quad \int \frac{e^{3i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=176

$$-\frac{6ib\sqrt{1+ia+ibx}}{(i+a)^2\sqrt{1-ia-ibx}} - \frac{(1+ia+ibx)^{3/2}}{(1-ia)x\sqrt{1-ia-ibx}} + \frac{6i\sqrt{i-a} b \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i+a)^{5/2}}$$

[Out] $6*I*b*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)/(1-I*a-I*b*x)^{(1/2))})*(I-a)^{(1/2)/(I+a)^{(5/2)}-(1+I*a+I*b*x)^{(3/2)/(1-I*a)/x/(1-I*a-I*b*x)^{(1/2)-6*I*b*(1+I*a+I*b*x)^{(1/2)/(I+a)^2/(1-I*a-I*b*x)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 176, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5203, 96, 95, 214}

$$-\frac{(ia+ibx+1)^{3/2}}{(1-ia)x\sqrt{-ia-ibx+1}} - \frac{6ib\sqrt{ia+ibx+1}}{(a+i)^2\sqrt{-ia-ibx+1}} + \frac{6i\sqrt{-a+i} b \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(a+i)^{5/2}}$$

Antiderivative was successfully verified.

[In] `Int[E^((3*I)*ArcTan[a + b*x])/x^2,x]`

[Out] $((-6*I)*b*\operatorname{Sqrt}[1 + I*a + I*b*x])/((I + a)^2*\operatorname{Sqrt}[1 - I*a - I*b*x]) - (1 + I*a + I*b*x)^{(3/2)/((1 - I*a)*x*\operatorname{Sqrt}[1 - I*a - I*b*x])} + ((6*I)*\operatorname{Sqrt}[I - a]*b*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I + a]*\operatorname{Sqrt}[1 + I*a + I*b*x])/(\operatorname{Sqrt}[I - a]*\operatorname{Sqrt}[1 - I*a - I*b*x])])/(I + a)^{(5/2)}$

Rule 95

`Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

Rule 96

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1)/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_.))*((d_) + (e_)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{3i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{(1+ia+ibx)^{3/2}}{x^2(1-ia-ibx)^{3/2}} dx \\
 &= -\frac{(1+ia+ibx)^{3/2}}{(1-ia)x\sqrt{1-ia-ibx}} - \frac{(3b) \int \frac{\sqrt{1+ia+ibx}}{x(1-ia-ibx)^{3/2}} dx}{i+a} \\
 &= -\frac{6ib\sqrt{1+ia+ibx}}{(i+a)^2\sqrt{1-ia-ibx}} - \frac{(1+ia+ibx)^{3/2}}{(1-ia)x\sqrt{1-ia-ibx}} - \frac{(3(i-a)b) \int \frac{1}{x\sqrt{1-ia-ibx}} dx}{(i+a)^2} \\
 &= -\frac{6ib\sqrt{1+ia+ibx}}{(i+a)^2\sqrt{1-ia-ibx}} - \frac{(1+ia+ibx)^{3/2}}{(1-ia)x\sqrt{1-ia-ibx}} - \frac{(6(i-a)b) \text{Subst}\left(\int \frac{1}{-1-ia-(-1+ibx)} dx\right)}{(i+a)^2} \\
 &= -\frac{6ib\sqrt{1+ia+ibx}}{(i+a)^2\sqrt{1-ia-ibx}} - \frac{(1+ia+ibx)^{3/2}}{(1-ia)x\sqrt{1-ia-ibx}} + \frac{6i\sqrt{i-a} b \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1-ia-ibx}}{\sqrt{i-a}\sqrt{1+ia+ibx}}\right)}{(i+a)^{5/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.10, size = 145, normalized size = 0.82

$$\frac{\frac{\sqrt{1+ia+ibx} (1+a^2-5ibx+abx)}{x\sqrt{-i(i+a+bx)}} + \frac{6i\sqrt{-1-ia} b \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}{\sqrt{-1+ia}}}{(i+a)^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])/x^2,x]

[Out] ((Sqrt[1 + I*a + I*b*x]*(1 + a^2 - (5*I)*b*x + a*b*x))/(x*Sqrt[(-I)*(I + a + b*x)]) + ((6*I)*Sqrt[-1 - I*a]*b*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a

+ b*x)]/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]]]/Sqrt[-1 + I*a])/(I + a)^2

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 630 vs. 2(136) = 272.

time = 0.20, size = 631, normalized size = 3.59

method	result
risch	$\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{(i+a)^2x} (a-i) - \frac{3b\sqrt{a^2 + 1} \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{(a^2+2ia-1)(i+a)} + \dots$
default	$-ib^3\left(-\frac{1}{i^2\sqrt{b^2x^2+2abx+a^2+1}} - \frac{2a(2b^2x+2ab)}{b(4b^2(a^2+1)-4a^2b^2)\sqrt{b^2x^2+2abx+a^2+1}}\right) - \frac{6}{(4b^2(a^2+1)-4a^2b^2)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^2,x,method=_RETURNVERBOSE)

[Out]
$$-I*b^3*(-1/b^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a/b*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))-6*I*a*b^2*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-6*b^2*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*b*(I*a^2-I+2*a)*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))+(-I*a^3-3*a^2+3*I*a+1)*(-1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a*b/(a^2+1)*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-4*b^2/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 992 vs. 2(116) = 232.

time = 0.27, size = 992, normalized size = 5.64

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^2,x, algorithm="maxima")

[Out]
$$-I*a*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 3*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))$$

$$\begin{aligned}
& 2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^2) - I*a^2*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 3*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^3*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)^2) - 3*(I*a^2*b + 2*a*b - I*b)*a*b^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)) + 2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)) - 3*(I*a^2*b + 2*a*b - I*b)*a^2*b^2/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)) + 2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)) + 3*(I*a*b^2 + b^2)*a*b/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) + 3*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b*arcsinh(2*a*b*x/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)))/(a^2 + 1)^(5/2) + I*b/\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} - 3*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)^2) + 3*(I*a^2*b + 2*a*b - I*b)*arcsinh(2*a*b*x/\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)))/(a^2 + 1)^(3/2) - 3*(I*a^2*b + 2*a*b - I*b)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)) - (-I*a^3 - 3*a^2 + 3*I*a + 1)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*(a^2 + 1)*x)
\end{aligned}$$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 389 vs. $2(116) = 232$.
time = 2.39, size = 389, normalized size = 2.21

$$\frac{(-14 - 59a^2 + (-14^2 - 4a - 59a - 3)a^2 + 3a - 13a^2 + (a^2 + 3a^2 - 3a - 10a^2) \sqrt{\frac{a - 10a^2}{a^2 + 3a - 13a^2 - 3a + 1}}) \operatorname{arcsinh}\left(\frac{2abx}{\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)}\right) + 2a^2 \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x) + 2 \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)}{(a^2 + 3a - 13a^2 - 3a + 1) \sqrt{a^2 + 3a - 13a^2 - 3a + 1}} \operatorname{arcsinh}\left(\frac{2abx}{\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)}\right) + 2a^2 \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x) + 2 \sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)}{(a^2 + 3a - 13a^2 - 3a + 1) \sqrt{a^2 + 3a - 13a^2 - 3a + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^2,x, algorithm="fricas")

[Out] $\begin{aligned}
& -((-I*a - 5)*b^2*x^2 + (-I*a^2 - 4*a - 5*I)*b*x - 3*((a^2 + 2*I*a - 1)*b*x^2 + (a^3 + 3*I*a^2 - 3*a - I)*x)*\sqrt{(a - I)*b^2/(a^5 + 5*I*a^4 - 10*a^3 - 10*I*a^2 + 5*a + I)}*\log(-(b^2*x + (a^3 + 3*I*a^2 - 3*a - I)*\sqrt{(a - I)*b^2/(a^5 + 5*I*a^4 - 10*a^3 - 10*I*a^2 + 5*a + I)}) - \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b/b) + 3*((a^2 + 2*I*a - 1)*b*x^2 + (a^3 + 3*I*a^2 - 3*a - I)*x)*\sqrt{(a - I)*b^2/(a^5 + 5*I*a^4 - 10*a^3 - 10*I*a^2 + 5*a + I)}*\log(-(b^2*x - (a^3 + 3*I*a^2 - 3*a - I)*\sqrt{(a - I)*b^2/(a^5 + 5*I*a^4 - 10*a^3 - 10*I*a^2 + 5*a + I)}) - \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1})*b/b) + \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*((-I*a - 5)*b*x - I*a^2 - I)/((a^2 + 2*I*a - 1)*b*x^2 + (a^3 + 3*I*a^2 - 3*a - I)*x)
\end{aligned}$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2)/x**2,x)

[Out] $-I*(\text{Integral}(I/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(-3*a/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(a^{**3}/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(-3*I*a^{**2}/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(-3*b*x/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(b^{**3}*x^{**3}/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(-3*I*b^{**2}*x^{**2}/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(3*a*b^{**2}*x^{**2}/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(3*a^{**2}*b*x/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x) + \text{Integral}(-6*I*a*b*x/(a^{**2}*x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + 2*a*b*x^{**3}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + b^{**2}*x^{**4}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1) + x^{**2}*\text{sqrt}(a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1))), x))$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^2,x, algorithm="giac")

[Out] undef

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(1 + a x + b x^2)^3}{x^2 ((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*x + b*x^2 + 1)^3/(x^2*((a + b*x)^2 + 1)^(3/2)),x)
```

```
[Out] int((a*x + b*x^2 + 1)^3/(x^2*((a + b*x)^2 + 1)^(3/2)), x)
```


$$3.187 \quad \int \frac{e^{3i \operatorname{ArcTan}(a+bx)}}{x^3} dx$$

Optimal. Leaf size=264

$$\frac{3(3i-2a)b^2\sqrt{1+ia+ibx}}{(1+ia)(i+a)^3\sqrt{1-ia-ibx}} + \frac{(3i-2a)b(1+ia+ibx)^{3/2}}{2(1+ia)(i+a)^2x\sqrt{1-ia-ibx}} - \frac{(1+ia+ibx)^{5/2}}{2(1+a^2)x^2\sqrt{1-ia-ibx}} + \frac{3(3+2i)}{2(1+ia)(i+a)^3\sqrt{1-ia-ibx}}$$

[Out] $3*(3+2*I*a)*b^2*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(I-a)^{(1/2)}/(1-I*a-I*b*x)^{(1/2)})/(I+a)^{(7/2)}/(I-a)^{(1/2)}+1/2*(3*I-2*a)*b*(1+I*a+I*b*x)^{(3/2)}/(1+I*a)/(I+a)^2/x/(1-I*a-I*b*x)^{(1/2)}-1/2*(1+I*a+I*b*x)^{(5/2)}/(a^2+1)/x^2/(1-I*a-I*b*x)^{(1/2)}+3*(3*I-2*a)*b^2*(1+I*a+I*b*x)^{(1/2)}/(1+I*a)/(I+a)^3/(1-I*a-I*b*x)^{(1/2)}$

Rubi [A]

time = 0.13, antiderivative size = 264, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {5203, 98, 96, 95, 214}

$$-\frac{(ia+ibx+1)^{5/2}}{2(a^2+1)x^2\sqrt{-ia-ibx+1}} + \frac{3(-2a+3i)b^2\sqrt{ia+ibx+1}}{(1+ia)(a+i)^3\sqrt{-ia-ibx+1}} + \frac{3(3+2ia)b^2 \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{\sqrt{-a+i}(a+i)^{7/2}} + \frac{(-2a+3i)b(ia+ibx+1)^{3/2}}{2(1+ia)(a+i)^2x\sqrt{-ia-ibx+1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)*\operatorname{ArcTan}[a+bx])}/x^3,x]$

[Out] $(3*(3*I-2*a)*b^2*\operatorname{Sqrt}[1+I*a+I*b*x])/((1+I*a)*(I+a)^3*\operatorname{Sqrt}[1-I*a-I*b*x]) + ((3*I-2*a)*b*(1+I*a+I*b*x)^{(3/2)})/(2*(1+I*a)*(I+a)^2*x*\operatorname{Sqrt}[1-I*a-I*b*x]) - (1+I*a+I*b*x)^{(5/2)}/(2*(1+a^2)*x^2*\operatorname{Sqrt}[1-I*a-I*b*x]) + (3*(3+(2*I)*a)*b^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])])/(\operatorname{Sqrt}[I-a]*(I+a)^{(7/2)})$

Rule 95

$\operatorname{Int}[((a_.)+(b_.)*(x_.))^{(m_.)}*((c_.)+(d_.)*(x_.))^{(n_.)}]/((e_.)+(f_.)*(x_.)), x_Symbol] :> \operatorname{With}\{q = \operatorname{Denominator}[m]\}, \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e-a*f-(d*e-c*f)*x^q), x], x, (a+bx)^{(1/q)}/(c+dx)^{(1/q)}], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x \ \&\& \operatorname{EqQ}[m+n+1, 0] \ \&\& \operatorname{RationalQ}[n] \ \&\& \operatorname{LtQ}[-1, m, 0] \ \&\& \operatorname{SimplerQ}[a+bx, c+dx]$

Rule 96

$\operatorname{Int}[((a_.)+(b_.)*(x_.))^{(m_.)}*((c_.)+(d_.)*(x_.))^{(n_.)}*((e_.)+(f_.)*(x_.))^{(p_.)}], x_Symbol] :> \operatorname{Simp}[(a+bx)^{(m+1)}*(c+dx)^n*(e+fx)^{(p+1)}/((m+1)*(b*e-a*f)), x] - \operatorname{Dist}[n*((d*e-c*f)/((m+1)*(b*e-a*f))), \operatorname{Int}[(a+bx)^{(m+1)}*(c+dx)^{(n-1)}*(e+fx)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, p\}, x \ \&\& \operatorname{EqQ}[m+n+p+2, 0] \ \&\& \operatorname{GtQ}[n, 0] \ \&\& (\operatorname{SumSimpler}$

Q[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 98

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])

Rule 214

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{3i \tan^{-1}(a+bx)}}{x^3} dx &= \int \frac{(1 + ia + ibx)^{3/2}}{x^3(1 - ia - ibx)^{3/2}} dx \\
 &= -\frac{(1 + ia + ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 - ia - ibx}} + \frac{((3i - 2a)b) \int \frac{(1 + ia + ibx)^{3/2}}{x^2(1 - ia - ibx)^{3/2}} dx}{2(1 + a^2)} \\
 &= -\frac{(3i - 2a)b(1 + ia + ibx)^{3/2}}{2(1 - ia)(1 + a^2)x\sqrt{1 - ia - ibx}} - \frac{(1 + ia + ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 - ia - ibx}} - \frac{(3(3i - 2a)b^2)}{2(i + a)} \\
 &= -\frac{3(3i - 2a)b^2\sqrt{1 + ia + ibx}}{(i - a)(1 - ia)^3\sqrt{1 - ia - ibx}} - \frac{(3i - 2a)b(1 + ia + ibx)^{3/2}}{2(1 - ia)(1 + a^2)x\sqrt{1 - ia - ibx}} - \frac{(1 + ia)}{2(1 + a^2)x^2} \\
 &= -\frac{3(3i - 2a)b^2\sqrt{1 + ia + ibx}}{(i - a)(1 - ia)^3\sqrt{1 - ia - ibx}} - \frac{(3i - 2a)b(1 + ia + ibx)^{3/2}}{2(1 - ia)(1 + a^2)x\sqrt{1 - ia - ibx}} - \frac{(1 + ia)}{2(1 + a^2)x^2} \\
 &= -\frac{3(3i - 2a)b^2\sqrt{1 + ia + ibx}}{(i - a)(1 - ia)^3\sqrt{1 - ia - ibx}} - \frac{(3i - 2a)b(1 + ia + ibx)^{3/2}}{2(1 - ia)(1 + a^2)x\sqrt{1 - ia - ibx}} - \frac{(1 + ia)}{2(1 + a^2)x^2}
 \end{aligned}$$

Mathematica [A]

time = 0.16, size = 194, normalized size = 0.73

$$\frac{\sqrt{1+ia+ibx} \frac{(i+a+ia^2+a^3-5bx+5iabx+14ib^2x^2-ab^2x^2)}{x^2 \sqrt{-i(i+a+bx)}} - \frac{6i\sqrt{-1-ia} (-3i+2a)b^2 \tanh^{-1}\left(\frac{\sqrt{-1-ia} \sqrt{-i(i+a+bx)}}{\sqrt{-1+ia} \sqrt{1+ia+ibx}}\right)}{\sqrt{-1+ia} (-i+a)}}{2(i+a)^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])/x^3,x]

[Out] ((Sqrt[1 + I*a + I*b*x]*(I + a + I*a^2 + a^3 - 5*b*x + (5*I)*a*b*x + (14*I)*b^2*x^2 - a*b^2*x^2))/(x^2*Sqrt[(-I)*(I + a + b*x)]) - ((6*I)*Sqrt[-1 - I*a]*(-3*I + 2*a)*b^2*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])]/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]))/(Sqrt[-1 + I*a]*(-I + a)))/(2*(I + a)^3)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 945 vs. 2(209) = 418.

time = 0.21, size = 946, normalized size = 3.58

method	result
risch	$\frac{i(-ab^3x^3+6ib^3x^3-a^2b^2x^2+12ia b^2x^2+a^3bx+6ix a^2b+a^4+b^2x^2+abx+6ibx+2a^2+1)}{2x^2(i+a)^3 \sqrt{b^2x^2+2abx+a^2+1}} - \frac{3ib^2 \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1} \sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{2(a^3+3ia^2-3a-i)(i+a)}$
default	$-\frac{2ib^3(2b^2x+2ab)}{(4b^2(a^2+1)-4a^2b^2)\sqrt{b^2x^2+2abx+a^2+1}} + (-ia^3 - 3a^2 + 3ia + 1) \left(-\frac{1}{2(a^2+1)x^2 \sqrt{b^2x^2+2abx+a^2+1}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^3,x,method=_RETURNVERBOSE)

[Out] -2*I*b^3*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+(-I*a^3-3*a^2+3*I*a+1)*(-1/2/(a^2+1)/x^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)

$$\begin{aligned}
&)-5/2*a*b/(a^2+1)*(-1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3*a*b/(a^2+1) \\
&*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4* \\
&b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-1/(a^2+1)^{(3/2)}*\ln((2* \\
&a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x))-4*b^2/(a^2 \\
&+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} \\
&)-3/2*b^2/(a^2+1)*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a*b/(a^2+1)*(2 \\
&*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-1/(a^ \\
&2+1)^{(3/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} \\
&)/x)))-3*(1+I*a)*b^2*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a*b/(a^2+1 \\
&)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-1 \\
&/a^2+1)^{(3/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} \\
&)/x))-3*b*(I*a^2-I+2*a)*(-1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-3* \\
&a*b/(a^2+1)*(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-2*a*b/(a^2+1)*(2*b^2*x \\
&+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}-1/(a^2+1)^{(\\
&3/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x)) \\
&-4*b^2/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a \\
&^2+1)^{(1/2)})
\end{aligned}$$

Maxima [B] Both result and optimal contain **B** but leaf count of result is larger than twice the leaf count of optimal. 1536 vs. $2(182) = 364$.
time = 0.27, size = 1536, normalized size = 5.82

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^3,x, algorithm="maxima")

[Out] $15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^3*b^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^3} + 15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^4*b^4/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^3} + 9*(I*a^2*b + 2*a*b - I*b)*a^2*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^2} + I*b^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)}) - 13/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^2} + 9*(I*a^2*b + 2*a*b - I*b)*a^3*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^2} + I*a*b^4/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)}) - 13/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*b^4/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^2} - 3*(I*a*b^2 + b^2)*a*b^3*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)}) - 6*(I*a^2*b + 2*a*b - I*b)*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)}) - 3*(I*a*b^2 + b^2)*a^2*b^2/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)}) - 6*(I*a^2*b + 2*a*b - I*b)*a*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)}) - 15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*b^2*\operatorname{arcsinh}(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2})*\operatorname{abs}(\sqrt{(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^3}))$

$$\begin{aligned}
& x)) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)))/(a^2 + 1)^{(7/2)} + 15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*b^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^3) - 9*(I*a^2*b + 2*a*b - I*b)*a*b*arcsinh(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x))) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)))/(a^2 + 1)^{(5/2)} + 3/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b^2*arcsinh(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x))) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)))/(a^2 + 1)^{(5/2)} + 9*(I*a^2*b + 2*a*b - I*b)*a*b/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) - 3/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b^2/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) + 3*(I*a*b^2 + b^2)*arcsinh(2*a*b*x/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x))) + 2*a^2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)) + 2/(\sqrt{-4*a^2*b^2 + 4*(a^2 + 1)*b^2}*abs(x)))/(a^2 + 1)^{(3/2)} - 3*(I*a*b^2 + b^2)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)) + 5/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2*x) + 3*(I*a^2*b + 2*a*b - I*b)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)*x) - 1/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)/(\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)*x^2)
\end{aligned}$$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 574 vs. 2(182) = 364.
time = 6.05, size = 574, normalized size = 2.17

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^3,x, algorithm="fricas")

[Out]
$$\begin{aligned}
& 1/2*((-I*a - 14)*b^3*x^3 + (-I*a^2 - 13*a - 14*I)*b^2*x^2 - 3*((a^3 + 3*I*a^2 - 3*a - I)*b*x^3 + (a^4 + 4*I*a^3 - 6*a^2 - 4*I*a + 1)*x^2)*\sqrt{(4*a^2 - 12*I*a - 9)*b^4/(a^8 + 6*I*a^7 - 14*a^6 - 14*I*a^5 - 14*I*a^3 + 14*a^2 + 6*I*a - 1))} \\
& \log(-(2*a - 3*I)*b^3*x - \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(2*a - 3*I)*b^2 + (a^5 + 3*I*a^4 - 2*a^3 + 2*I*a^2 - 3*a - I)*\sqrt{(4*a^2 - 12*I*a - 9)*b^4/(a^8 + 6*I*a^7 - 14*a^6 - 14*I*a^5 - 14*I*a^3 + 14*a^2 + 6*I*a - 1))}) \\
& /((2*a - 3*I)*b^2) + 3*((a^3 + 3*I*a^2 - 3*a - I)*b*x^3 + (a^4 + 4*I*a^3 - 6*a^2 - 4*I*a + 1)*x^2)*\sqrt{(4*a^2 - 12*I*a - 9)*b^4/(a^8 + 6*I*a^7 - 14*a^6 - 14*I*a^5 - 14*I*a^3 + 14*a^2 + 6*I*a - 1))} \\
& \log(-(2*a - 3*I)*b^3*x - \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(2*a - 3*I)*b^2 - (a^5 + 3*I*a^4 - 2*a^3 + 2*I*a^2 - 3*a - I)*\sqrt{(4*a^2 - 12*I*a - 9)*b^4/(a^8 + 6*I*a^7 - 14*a^6 - 14*I*a^5 - 14*I*a^3 + 14*a^2 + 6*I*a - 1))}) \\
& /((2*a - 3*I)*b^2) + ((-I*a - 14)*b^2*x^2 + I*a^3 - 5*(a + I)*b*x - a^2 + I*a - 1)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) \\
& /((a^3 + 3*I*a^2 - 3*a - I)*b*x^3 + (a^4 + 4*I*a^3 - 6*a^2 - 4*I*a + 1)*x^2)
\end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2)/x**3,x)

[Out] $-I \cdot \left(\int \frac{1}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{-3 a}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{a^3}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{-3 I a^2}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{-3 I a b x}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{3 a b^2 x^2}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{3 a^2 b x}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx + \int \frac{-6 I a b x}{(a^2 x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}) + 2 a b x^4 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + b^2 x^5 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1} + x^3 \sqrt{a^2 + 2 a b x + b^2 x^2 + 1}}{x} dx \right)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^3,x, algorithm="giac")

[Out] undef

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(1 + a 1i + b x 1i)^3}{x^3 ((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)^3/(x^3*((a + b*x)^2 + 1)^(3/2)),x)

[Out] int((a*1i + b*x*1i + 1)^3/(x^3*((a + b*x)^2 + 1)^(3/2)), x)

$$3.188 \quad \int \frac{e^{3i \operatorname{ArcTan}(a+bx)}}{x^4} dx$$

Optimal. Leaf size=338

$$\frac{(52 + 51ia - 2a^2)b^3\sqrt{1 + ia + ibx}}{6(i - a)(i + a)^4\sqrt{1 - ia - ibx}} - \frac{(i - a)\sqrt{1 + ia + ibx}}{3(i + a)x^3\sqrt{1 - ia - ibx}} + \frac{7ib\sqrt{1 + ia + ibx}}{6(i + a)^2x^2\sqrt{1 - ia - ibx}} + \frac{(19 + 16ia)b}{6(i - a)(i + a)}$$

[Out] $-(11*I-18*a-6*I*a^2)*b^3*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)})/(1-I*a-I*b*x)^{(1/2)})/(I-a)^{(3/2)/(I+a)^{(9/2)}+1/6*(52+51*I*a-2*a^2)*b^3*(1+I*a+I*b*x)^{(1/2)/(I-a)/(I+a)^4/(1-I*a-I*b*x)^{(1/2)}-1/3*(I-a)*(1+I*a+I*b*x)^{(1/2)/(I+a)/x^3/(1-I*a-I*b*x)^{(1/2)}+7/6*I*b*(1+I*a+I*b*x)^{(1/2)/(I+a)^2/x^2/(1-I*a-I*b*x)^{(1/2)}+1/6*(19+16*I*a)*b^2*(1+I*a+I*b*x)^{(1/2)/(I-a)/(I+a)^3/x/(1-I*a-I*b*x)^{(1/2)}$

Rubi [A]

time = 0.24, antiderivative size = 338, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 100, 156, 157, 12, 95, 214}

$$\frac{(-2a^2 + 51ia + 52)b^3\sqrt{ia + ibx + 1}}{6(-a + i)(a + i)^4\sqrt{-ia - ibx + 1}} - \frac{(-6ia^2 - 18a + 11i)b^3 \tanh^{-1}\left(\frac{\sqrt{-a + i}\sqrt{ia + ibx + 1}}{\sqrt{-a + i}\sqrt{-ia - ibx + 1}}\right)}{(-a + i)^{3/2}(a + i)^{9/2}} + \frac{(19 + 16ia)b^2\sqrt{ia + ibx + 1}}{6(-a + i)(a + i)^3x\sqrt{-ia - ibx + 1}} - \frac{(-a + i)\sqrt{ia + ibx + 1}}{3(a + i)x^3\sqrt{-ia - ibx + 1}} + \frac{7ib\sqrt{ia + ibx + 1}}{6(a + i)^2x^2\sqrt{-ia - ibx + 1}}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a + b*x])/x^4,x]

[Out] $((52 + (51*I)*a - 2*a^2)*b^3*\operatorname{Sqrt}[1 + I*a + I*b*x])/(6*(I - a)*(I + a)^4*\operatorname{Sqrt}[1 - I*a - I*b*x]) - ((I - a)*\operatorname{Sqrt}[1 + I*a + I*b*x])/(3*(I + a)*x^3*\operatorname{Sqrt}[1 - I*a - I*b*x]) + (((7*I)/6)*b*\operatorname{Sqrt}[1 + I*a + I*b*x])/((I + a)^2*x^2*\operatorname{Sqrt}[1 - I*a - I*b*x]) + ((19 + (16*I)*a)*b^2*\operatorname{Sqrt}[1 + I*a + I*b*x])/(6*(I - a)*(I + a)^3*x*\operatorname{Sqrt}[1 - I*a - I*b*x]) - ((11*I - 18*a - (6*I)*a^2)*b^3*\operatorname{ArcTan}[\operatorname{Sqrt}[I + a]*\operatorname{Sqrt}[1 + I*a + I*b*x])/(\operatorname{Sqrt}[I - a]*\operatorname{Sqrt}[1 - I*a - I*b*x])]/((I - a)^{(3/2)}*(I + a)^{(9/2)})$

Rule 12

Int[(a_)*(u_), x_Symbol] :> Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 156

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

Rule 157

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(a+bx)}}{x^4} dx &= \int \frac{(1+ia+ibx)^{3/2}}{x^4(1-ia-ibx)^{3/2}} dx \\
&= -\frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} - \frac{\int \frac{-7(i-a)b+6b^2x}{x^3(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}} dx}{3(1-ia)} \\
&= -\frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} + \frac{7ib\sqrt{1+ia+ibx}}{6(i+a)^2x^2\sqrt{1-ia-ibx}} + \frac{\int \frac{-(19+35ia-16a^2)b^2-14(i-a)}{x^2(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}} dx}{6(1-ia)(1+a^2)} \\
&= -\frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} + \frac{7ib\sqrt{1+ia+ibx}}{6(i+a)^2x^2\sqrt{1-ia-ibx}} + \frac{(19i-16a)b^2\sqrt{1+ia+ibx}}{6(i-a)(1-ia)^3x\sqrt{1-ia-ibx}} \\
&= \frac{(52+51ia-2a^2)b^3\sqrt{1+ia+ibx}}{6(i-a)(i+a)^4\sqrt{1-ia-ibx}} - \frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} + \frac{7ib\sqrt{1+ia+ibx}}{6(i+a)^2x^2\sqrt{1-ia-ibx}} \\
&= \frac{(52+51ia-2a^2)b^3\sqrt{1+ia+ibx}}{6(i-a)(i+a)^4\sqrt{1-ia-ibx}} - \frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} + \frac{7ib\sqrt{1+ia+ibx}}{6(i+a)^2x^2\sqrt{1-ia-ibx}} \\
&= \frac{(52+51ia-2a^2)b^3\sqrt{1+ia+ibx}}{6(i-a)(i+a)^4\sqrt{1-ia-ibx}} - \frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} + \frac{7ib\sqrt{1+ia+ibx}}{6(i+a)^2x^2\sqrt{1-ia-ibx}} \\
&= \frac{(52+51ia-2a^2)b^3\sqrt{1+ia+ibx}}{6(i-a)(i+a)^4\sqrt{1-ia-ibx}} - \frac{(i-a)\sqrt{1+ia+ibx}}{3(i+a)x^3\sqrt{1-ia-ibx}} + \frac{7ib\sqrt{1+ia+ibx}}{6(i+a)^2x^2\sqrt{1-ia-ibx}}
\end{aligned}$$

Mathematica [A]

time = 0.28, size = 282, normalized size = 0.83

$$\frac{2(-1+ia)^{3/2}(1+ia)(i+a)^2(1+ia+ibx)^{5/2} + (3i-4a)(-1+ia)^{5/2}bx(1+ia+ibx)^{5/2} - i(-11-18ia+6a^2)b^2x^2(i\sqrt{-1+ia}\sqrt{1+ia+ibx}(1+a^2-5ibx+abx) - 6\sqrt{-1-ia}bx\sqrt{-i(i+a+bx)}\tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right))}{6(-1+ia)^{5/2}(1+a^2)^2x^3\sqrt{-i(i+a+bx)}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a + b*x])/x^4,x]

[Out] $-1/6*(2*(-1 + I*a)^{(3/2)}*(1 + I*a)*(I + a)^2*(1 + I*a + I*b*x)^{(5/2)} + (3*I - 4*a)*(-1 + I*a)^{(5/2)}*b*x*(1 + I*a + I*b*x)^{(5/2)} - I*(-11 - (18*I)*a + 6*a^2)*b^2*x^2*(I*\text{Sqrt}[-1 + I*a]*\text{Sqrt}[1 + I*a + I*b*x]*(1 + a^2 - (5*I)*b*x + a*b*x) - 6*\text{Sqrt}[-1 - I*a]*b*x*\text{Sqrt}[(-I)*(I + a + b*x)]*\text{ArcTanh}[(\text{Sqrt}[-1 - I*a]*\text{Sqrt}[(-I)*(I + a + b*x)])]/(\text{Sqrt}[-1 + I*a]*\text{Sqrt}[1 + I*a + I*b*x])))/((-1 + I*a)^{(5/2)}*(1 + a^2)^2*x^3*\text{Sqrt}[(-I)*(I + a + b*x)])$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1648 vs. $2(266) = 532$.

time = 0.26, size = 1649, normalized size = 4.88

method	result
risch	$\frac{i(2a^2b^4x^4 - 27ia^2b^4x^4 + 2a^3b^3x^3 - 45ia^2b^3x^3 - 9ia^3b^2x^2 - 28x^4b^4 + 2a^5bx + 9ia^4bx - 58a^3b^3x^3 + 9ib^3x^3 + 2a^6 - 26a^2b^2x^2 - 9ia^2b^2x^2 + 4a^3b)}{6x^3(a-i)(i+a)^4\sqrt{b^2x^2 + 2abx + a^2 + 1}}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^4,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & (-I*a^3-3*a^2+3*I*a+1)*(-1/3/(a^2+1)/x^3/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-7/3* \\ & a*b/(a^2+1)*(-1/2/(a^2+1)/x^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/2*a*b/(a^2+1) \\ & *(-1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a*b/(a^2+1)*(1/(a^2+1)/(b^2* \\ & x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2 \\ & *b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(\\ & a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-4*b^2/(a^2+1)*(2*b^2*x+2*a* \\ & b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))-3/2*b^2/(a^2+1) \\ & *(1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4* \\ & b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2* \\ & a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-4/3*b^2/(\\ & a^2+1)*(-1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a*b/(a^2+1)*(1/(a^2+1) \\ & /b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1) \\ & -4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b \\ & *x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-4*b^2/(a^2+1)*(2*b^2* \\ & x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))-3*b*(I*a \\ & ^2-I+2*a)*(-1/2/(a^2+1)/x^2/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-5/2*a*b/(a^2+1)*(\\ & -1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a*b/(a^2+1)*(1/(a^2+1)/(b^2*x^ \\ & 2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b \\ & ^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^ \\ & 2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-4*b^2/(a^2+1)*(2*b^2*x+2*a*b) \\ & /4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2))-3/2*b^2/(a^2+1)*(\\ & 1/(a^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^ \\ & 2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2*a^ \\ & 2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-I*b^3*(1/(a \\ & ^2+1)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a \\ & ^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2*a^2+2+ \\ & 2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-3*b^2*(1+I*a)*(- \\ & 1/(a^2+1)/x/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-3*a*b/(a^2+1)*(1/(a^2+1)/(b^2*x^2 \\ & +2*a*b*x+a^2+1)^(1/2)-2*a*b/(a^2+1)*(2*b^2*x+2*a*b)/(4*b^2*(a^2+1)-4*a^2*b^ \\ & 2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)-1/(a^2+1)^(3/2)*\ln((2*a^2+2+2*a*b*x+2*(a^ \\ & 2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-4*b^2/(a^2+1)*(2*b^2*x+2*a*b)/ \\ & (4*b^2*(a^2+1)-4*a^2*b^2)/(b^2*x^2+2*a*b*x+a^2+1)^(1/2)) \end{aligned}$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2313 vs. $2(223) = 446$.

time = 0.28, size = 2313, normalized size = 6.84

Too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^4,x, algorithm="maxima")

[Out]
$$\begin{aligned} & -35/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^4*b^6*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^4) - 35/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^5*b^5/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^4) - 45/2*(I*a^2*b + 2*a*b - I*b)*a^3*b^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^3) - I*a*b^6*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)) + 115/6*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^2*b^6*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^3) - 45/2*(I*a^2*b + 2*a*b - I*b)*a^4*b^4/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^3) - I*a^2*b^5/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)) + 115/6*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^3*b^5/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^3) + 9*(I*a*b^2 + b^2)*a^2*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) + 39/2*(I*a^2*b + 2*a*b - I*b)*a*b^5*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) - 8/3*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b^6*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) + 9*(I*a*b^2 + b^2)*a^3*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) + 39/2*(I*a^2*b + 2*a*b - I*b)*a^2*b^4/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) - 8/3*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b^5/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)^2) - 6*(I*a*b^2 + b^2)*b^4*x/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)) + 35/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^3*b^3*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(9/2) - 6*(I*a*b^2 + b^2)*a*b^3/((a^2*b^2 - (a^2 + 1)*b^2)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(a^2 + 1)) - 35/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a^3*b^3/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^4) + 45/2*(I*a^2*b + 2*a*b - I*b)*a^2*b^2*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(7/2) + I*b^3*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x))) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(3/2) - 15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b^3*arcsinh(2*a*b*x/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2*a^2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)) + 2/(sqrt(-4*a^2*b^2 + 4*(a^2 + 1)*b^2)*abs(x)))/(a^2 + 1)^(7/2) - 45/2*(I*a^2*b + 2*a*b - I*b)*a^2*b^2/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)^3) - I*b^3/(sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a^2 + 1)) + 15/2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*a*b^3/(sq$$

$$\begin{aligned} & \text{rt}(b^2x^2 + 2abx + a^2 + 1)(a^2 + 1)^3 - 9(Iab^2 + b^2)ab \operatorname{arcsinh}(2abx/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)) + 2a^2/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x) \\ & + 2/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)))/(a^2 + 1)^{5/2} - 9/2(Ia^2b + 2ab - Ib)b^2 \operatorname{arcsinh}(2abx/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)) \\ & + 2a^2/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x) + 2/\sqrt{-4a^2b^2 + 4(a^2 + 1)b^2} \operatorname{abs}(x)))/(a^2 + 1)^{5/2} \\ & + 9(Iab^2 + b^2)ab/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)^2 + 9/2(Ia^2b + 2ab - Ib)b^2/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)^2 \\ & - 35/6(-Ia^3 - 3a^2 + 3Ia + 1)a^2b^2/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)^3x - 15/2(Ia^2b + 2ab - Ib)ab/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)^2x \\ & + 4/3(-Ia^3 - 3a^2 + 3Ia + 1)b^2/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)^2x + 7/6(-Ia^3 - 3a^2 + 3Ia + 1)ab/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)^2x^2 \\ & + 3(Iab^2 + b^2)/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)x + 3/2(Ia^2b + 2ab - Ib)/\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)x^2 - 1/3(-Ia^3 - 3a^2 + 3Ia + 1) \\ & /(\sqrt{b^2x^2 + 2abx + a^2 + 1}(a^2 + 1)x^3) \end{aligned}$$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 839 vs. $2(223) = 446$.
time = 4.74, size = 839, normalized size = 2.48

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^4,x, algorithm="fricas")`

[Out]
$$\begin{aligned} & 1/6*((2Ia^2 + 51a - 52I)b^4x^4 + (2Ia^3 + 49a^2 - Ia + 52)b^3x^3 + 3\sqrt{(36a^4 - 216Ia^3 - 456a^2 + 396Ia + 121)b^6/(a^{12} + 6Ia^{11} - 12a^{10} - 2Ia^9 - 27a^8 - 36Ia^7 - 36Ia^5 + 27a^4 - 2Ia^3 + 12a^2 + 6Ia - 1)} \\ & ((a^5 + 3Ia^4 - 2a^3 + 2Ia^2 - 3a - I)b^4x^4 + (a^6 + 4Ia^5 - 5a^4 - 5a^2 - 4Ia + 1)x^3) \log(-((6a^2 - 18Ia - 11)b^4x - \sqrt{b^2x^2 + 2abx + a^2 + 1}(6a^2 - 18Ia - 11)b^3 + (a^7 + 3Ia^6 - a^5 + 5Ia^4 - 5a^3 + Ia^2 - 3a - I)\sqrt{(36a^4 - 216Ia^3 - 456a^2 + 396Ia + 121)b^6/(a^{12} + 6Ia^{11} - 12a^{10} - 2Ia^9 - 27a^8 - 36Ia^7 - 36Ia^5 + 27a^4 - 2Ia^3 + 12a^2 + 6Ia - 1)})) / ((6a^2 - 18Ia - 11)b^3) - 3\sqrt{(36a^4 - 216Ia^3 - 456a^2 + 396Ia + 121)b^6/(a^{12} + 6Ia^{11} - 12a^{10} - 2Ia^9 - 27a^8 - 36Ia^7 - 36Ia^5 + 27a^4 - 2Ia^3 + 12a^2 + 6Ia - 1)} \\ & ((a^5 + 3Ia^4 - 2a^3 + 2Ia^2 - 3a - I)b^4x^4 + (a^6 + 4Ia^5 - 5a^4 - 5a^2 - 4Ia + 1)x^3) \log(-((6a^2 - 18Ia - 11)b^4x - \sqrt{b^2x^2 + 2abx + a^2 + 1}(6a^2 - 18Ia - 11)b^3 - (a^7 + 3Ia^6 - a^5 + 5Ia^4 - 5a^3 + Ia^2 - 3a - I)\sqrt{(36a^4 - 216Ia^3 - 456a^2 + 396Ia + 121)b^6/(a^{12} + 6Ia^{11} - 12a^{10} - 2Ia^9 - 27a^8 - 36Ia^7 - 36Ia^5 + 27a^4 - 2Ia^3 + 12a^2 + 6Ia - 1)})) / ((6a^2 - 18Ia - 11)b^3) + ((2Ia^2 + 51a - 52I) \end{aligned}$$

$$) * b^3 x^3 + 2 I a^5 + (16 a^2 - 3 I a + 19) b^2 x^2 - 2 a^4 + 4 I a^3 - 7 (a^3 + I a^2 + a + I) b x - 4 a^2 + 2 I a - 2) \sqrt{b^2 x^2 + 2 a b x + a^2 + 1} / ((a^5 + 3 I a^4 - 2 a^3 + 2 I a^2 - 3 a - I) b x^4 + (a^6 + 4 I a^5 - 5 a^4 - 5 a^2 - 4 I a + 1) x^3)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))**3/(1+(b*x+a)**2)**(3/2)/x**4,x)

[Out] -I*(Integral(I/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*a/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(a**3/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*I*a**2/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*b*x/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(b**3*x**3/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-3*I*b**2*x**2/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a*b**2*x**2/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(3*a**2*b*x/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x) + Integral(-6*I*a*b*x/(a**2*x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + 2*a*b*x**5*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + b**2*x**6*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1) + x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*(b*x+a))^3/(1+(b*x+a)^2)^(3/2)/x^4,x, algorithm="giac")

[Out] undef

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{(1 + a \operatorname{li} + b x \operatorname{li})^3}{x^4 ((a + b x)^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*1i + b*x*1i + 1)^3/(x^4*((a + b*x)^2 + 1)^(3/2)),x)

[Out] int((a*1i + b*x*1i + 1)^3/(x^4*((a + b*x)^2 + 1)^(3/2)), x)

3.189 $\int e^{-i\text{ArcTan}(a+bx)} x^4 dx$

Optimal. Leaf size=276

$$\frac{(3i - 12a - 24ia^2 + 16a^3 + 8ia^4) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{8b^5} + \frac{(i - 8a)x^2(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{20b^3}$$

[Out] $\frac{1}{8}*(3+12*I*a-24*a^2-16*I*a^3+8*a^4)*\text{arcsinh}(b*x+a)/b^5+1/20*(I-8*a)*x^2*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3+1/5*x^3*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2-1/120*(1-I*a-I*b*x)^{(3/2)}*(19*I-114*a-86*I*a^2+96*a^3+2*(13+14*I*a-36*a^2)*b*x)*(1+I*a+I*b*x)^{(1/2)}/b^5-1/8*(3*I-12*a-24*I*a^2+16*a^3+8*I*a^4)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^5$

Rubi [A]

time = 0.16, antiderivative size = 276, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 102, 158, 152, 52, 55, 633, 221}

$$\frac{(-ia - ibx + 1)^{3/2} \sqrt{ia + ibx + 1} (96a^3 + 2(-36a^2 + 14ia + 13)bx - 86ia^2 - 114a + 19)}{120b^5} - \frac{(8ia^4 + 16a^3 - 24ia^2 - 12a + 3) \sqrt{-ia - ibx + 1} \sqrt{ia + ibx + 1}}{8b^5} + \frac{(8a^4 - 16ia^3 - 24a^2 + 12ia + 3) \text{sinh}^{-1}(a + bx)}{8b^5} + \frac{(-8a + i)x^2(-ia - ibx + 1)^{3/2} \sqrt{ia + ibx + 1}}{20b^3} + \frac{x^2(-ia - ibx + 1)^{3/2} \sqrt{ia + ibx + 1}}{5b^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^4/E^{(I*\text{ArcTan}[a + b*x])}, x]$

[Out] $-1/8*((3*I - 12*a - (24*I)*a^2 + 16*a^3 + (8*I)*a^4)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/b^5 + ((I - 8*a)*x^2*(1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x])/(20*b^3) + (x^3*(1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x])/((5*b^2) - ((1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x]*(19*I - 114*a - (86*I)*a^2 + 96*a^3 + 2*(13 + (14*I)*a - 36*a^2)*b*x))/(120*b^5) + ((3 + (12*I)*a - 24*a^2 - (16*I)*a^3 + 8*a^4)*\text{ArcSinh}[a + b*x])/(8*b^5)$

Rule 52

$\text{Int}[(a_.) + (b_.)*(x_.)]^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m + n + 1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0]))) \ \&\& \ !\text{ILtQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 55

$\text{Int}[1/(\text{Sqrt}[(a_.) + (b_.)*(x_.)]*\text{Sqrt}[(c_.) + (d_.)*(x_.)]), x_Symbol] \rightarrow \text{Int}[1/\text{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[b + d, 0] \ \&\& \ \text{GtQ}[a + c, 0]$

Rule 102


```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)), Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 158

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p + 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))]*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c)], x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
```

$I*b*c*x)^{(I*(n/2))}, x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned}
 \int e^{-i \tan^{-1}(a+bx)} x^4 dx &= \int \frac{x^4 \sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} dx \\
 &= \frac{x^3(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2} + \frac{\int \frac{x^2 \sqrt{1-ia-ibx} (-3(1+a^2)+(i-8a)bx)}{\sqrt{1+ia+ibx}} dx}{5b^2} \\
 &= \frac{(i-8a)x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{20b^3} + \frac{x^3(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2} + \int \frac{x \sqrt{1-ia-ibx} (-3(1+a^2)+(i-8a)bx)}{\sqrt{1+ia+ibx}} dx \\
 &= \frac{(i-8a)x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{20b^3} + \frac{x^3(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2} - \frac{\int \frac{x \sqrt{1-ia-ibx} (-3(1+a^2)+(i-8a)bx)}{\sqrt{1+ia+ibx}} dx}{5b^2} \\
 &= -\frac{(3i-12a-24ia^2+16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} + \frac{(i-8a)x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2} \\
 &= -\frac{(3i-12a-24ia^2+16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} + \frac{(i-8a)x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2} \\
 &= -\frac{(3i-12a-24ia^2+16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} + \frac{(i-8a)x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2} \\
 &= -\frac{(3i-12a-24ia^2+16a^3+8ia^4) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^5} + \frac{(i-8a)x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{5b^2}
 \end{aligned}$$

Mathematica [A]

time = 0.18, size = 158, normalized size = 0.57

$$\frac{-\sqrt{1+a^2+2abx+b^2x^2} (64i-275a-332ia^2+250a^3+24ia^4+(45+116ia-130a^2-24ia^3)bx+2(-16i+35a+12ia^2)b^2x^2-6(5+4ia)b^3x^3+24ib^4x^4)+15(3+12ia-24a^2-16ia^3+8a^4)\sinh^{-1}(a+bx)}{120b^5}$$

Antiderivative was successfully verified.

[In] Integrate[x^4/E^(I*ArcTan[a + b*x]),x]

[Out] $(-\text{Sqrt}[1+a^2+2*a*b*x+b^2*x^2]*(64*I-275*a-(332*I)*a^2+250*a^3+(24*I)*a^4+(45+(116*I)*a-130*a^2-(24*I)*a^3)*b*x+2*(-16*I+35*a+(12*I)*a^2)*b^2*x^2-6*(5+(4*I)*a)*b^3*x^3+(24*I)*b^4*x^4))+15*(3+(12*I)*a-24*a^2-(16*I)*a^3+8*a^4)*\text{ArcSinh}[a+b*x])/(120*b^5)$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1151 vs. $2(225) = 450$.
time = 0.17, size = 1152, normalized size = 4.17

method	result
risch	$\frac{i(24x^4b^4 - 24ab^3x^3 + 30ib^3x^3 + 24a^2b^2x^2 - 70ia b^2x^2 - 24a^3bx + 130ix a^2b + 24a^4 - 250ia^3 - 32b^2x^2 + 116abx - 45ibx - 332a^2 + 275ia + \dots)}{120b^5}$ $\left(\frac{x^2(b^2x^2 + 2abx + a^2 + 1)^{\frac{3}{2}}}{5b^2} \right) - \left(\frac{x(b^2x^2 + 2abx + a^2 + 1)^{\frac{3}{2}}}{4b^2} \right) - \left(\frac{(b^2x^2 + 2abx + a^2 + 1)^{\frac{3}{2}}}{3b^2} \right) - \left(\frac{a \left(\frac{(2b^2x + 2ab) \sqrt{b^2x^2 + 2abx + a^2 + 1}}{4b^2} \right)}{a} \right)$
default	$- \frac{x^2(b^2x^2 + 2abx + a^2 + 1)^{\frac{3}{2}}}{5b^2}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^4/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-I/b*(1/5*x^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-7/5*a/b*(1/4*x*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-5/4*a/b*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-a/b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a$

$$\begin{aligned}
& \frac{2b^2}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) - \frac{1}{4} \frac{(a^2+1)}{b^2} \frac{(1/4(2b^2x+2ab)/b^2 + (b^2x^2+2abx+a^2+1)^{1/2})}{(b^2)^{1/2}} \\
& + \frac{1}{8} \frac{(4b^2(a^2+1)-4a^2b^2)}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) - \frac{2}{5} \frac{(a^2+1)}{b^2} \frac{(1/3(b^2x^2+2abx+a^2+1)^{3/2})}{b^2} \\
& - \frac{a}{b} \frac{(1/4(2b^2x+2ab)/b^2 + (b^2x^2+2abx+a^2+1)^{1/2})}{(b^2)^{1/2}} + \frac{1}{8} \frac{(4b^2(a^2+1)-4a^2b^2)}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) \\
& - I \frac{(I-a)}{b^2} \frac{(1/4x(b^2x^2+2abx+a^2+1)^{3/2})}{b^2} - \frac{5}{4} \frac{a}{b} \frac{(1/3(b^2x^2+2abx+a^2+1)^{3/2})}{b^2} - \frac{a}{b} \frac{(1/4(2b^2x+2ab)/b^2 + (b^2x^2+2abx+a^2+1)^{1/2})}{(b^2)^{1/2}} \\
& + \frac{1}{8} \frac{(4b^2(a^2+1)-4a^2b^2)}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) - \frac{1}{4} \frac{(a^2+1)}{b^2} \frac{(1/4(2b^2x+2ab)/b^2 + (b^2x^2+2abx+a^2+1)^{1/2})}{(b^2)^{1/2}} \\
& + \frac{1}{8} \frac{(4b^2(a^2+1)-4a^2b^2)}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) - I \frac{(I-a)^2}{b^3} \frac{(1/3(b^2x^2+2abx+a^2+1)^{3/2})}{b^2} \\
& - \frac{a}{b} \frac{(1/4(2b^2x+2ab)/b^2 + (b^2x^2+2abx+a^2+1)^{1/2})}{(b^2)^{1/2}} + \frac{1}{8} \frac{(4b^2(a^2+1)-4a^2b^2)}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) \\
& - I \frac{(I-a)^3}{b^4} \frac{(1/4(2b^2x+2ab)/b^2 + (b^2x^2+2abx+a^2+1)^{1/2})}{(b^2)^{1/2}} + \frac{1}{8} \frac{(4b^2(a^2+1)-4a^2b^2)}{b^2} \ln\left(\frac{(b^2x+ab)}{(b^2)^{1/2}} + \frac{(b^2x^2+2abx+a^2+1)^{1/2}}{(b^2)^{1/2}}\right) \\
& + (-Ia^4+6Ia^2-4a^3-I+4a)/b^5 \left(\frac{(x-(I-a)/b)^2 b^2+2Ib(x-(I-a)/b)}{(b^2)^{1/2}} + I b \ln\left(\frac{(Ib+(x-(I-a)/b)b^2)}{(b^2)^{1/2}} + \frac{(x-(I-a)/b)^2 b^2+2Ib(x-(I-a)/b)}{(b^2)^{1/2}}\right) \right)
\end{aligned}$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 456 vs. $2(200) = 400$.
time = 0.51, size = 456, normalized size = 1.65

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="maxima")

[Out] $2I\sqrt{b^2x^2+2abx+a^2+1}a^3x/b^4 - 1/5I(b^2x^2+2abx+a^2+1)^{3/2}x^2/b^3 + a^4\operatorname{arcsinh}(bx+a)/b^5 + I\sqrt{b^2x^2+2abx+a^2+1}a^4/b^5 + 3/5I(b^2x^2+2abx+a^2+1)^{3/2}ax/b^4 + 3\sqrt{b^2x^2+2abx+a^2+1}a^2x/b^4 - 2Ia^3\operatorname{arcsinh}(bx+a)/b^5 - 6/5I(b^2x^2+2abx+a^2+1)^{3/2}a^2/b^5 - \sqrt{b^2x^2+2abx+a^2+1}a^3/b^5 + 1/4(b^2x^2+2abx+a^2+1)^{3/2}x/b^4 - 5/2I\sqrt{b^2x^2+2abx+a^2+1}ax/b^4 - 3a^2\operatorname{arcsinh}(bx+a)/b^5 - 13/12(b^2x^2+2abx+a^2+1)^{3/2}a/b^5 + 7/2I\sqrt{b^2x^2+2abx+a^2+1}a^2/b^5 - 5/8\sqrt{b^2x^2+2abx+a^2+1}x/b^4 + 3/2Ia\operatorname{arcsinh}(bx+a)/b^5 + 7/15I(b^2x^2+2abx+a^2+1)^{3/2}/b^5 + 27/8\sqrt{b^2x^2+2abx+a^2+1}a/b^5 + 3/8\operatorname{arcsinh}(bx+a)/b^5 - I\sqrt{b^2x^2+2abx+a^2+1}/b^5$

Fricas [A]

time = 4.00, size = 177, normalized size = 0.64

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="fricas")

[Out] 1/960*(-186*I*a^5 - 1345*a^4 + 1730*I*a^3 + 1320*a^2 - 120*(8*a^4 - 16*I*a^3 - 24*a^2 + 12*I*a + 3)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 8*(24*I*b^4*x^4 + 6*(-4*I*a - 5)*b^3*x^3 + 2*(12*I*a^2 + 35*a - 16*I)*b^2*x^2 + 24*I*a^4 + 250*a^3 + (-24*I*a^3 - 130*a^2 + 116*I*a + 45)*b*x - 332*I*a^2 - 275*a + 64*I)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) - 300*I*a)/b^5

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^4 \sqrt{a^2 + 2abx + b^2x^2 + 1}}{a + bx - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**4/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2),x)

[Out] -I*Integral(x**4*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a + b*x - I), x)

Giac [A]

time = 0.45, size = 205, normalized size = 0.74

$$-\frac{1}{120} \sqrt{(bx+a)^2+1} \left(\left(2 \left(3x \left(\frac{4ix}{b} - \frac{4iab^2+5b^3}{b^3} \right) - \frac{-12ia^2b^2-35ab^3+16ib^4}{b^3} \right) x - \frac{24ia^2b^2+130a^2b^3-116iab^3-45b^4}{b^3} \right) x - \frac{-24ia^2b^2-250a^2b^3+332ia^2b^2+275ab^3-64ib^4}{b^3} \right) - \frac{(8a^4-16ia^3-24a^2+12ia+3) \log \left(\frac{-ab - (x|b| - \sqrt{(bx+a)^2+1})|b|}{8b^4|b|} \right)}{8b^4|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="giac")

[Out] -1/120*sqrt((b*x + a)^2 + 1)*((2*(3*x*(4*I*x/b - (4*I*a*b^7 + 5*b^7)/b^9) - (-12*I*a^2*b^6 - 35*a*b^6 + 16*I*b^6)/b^9)*x - (24*I*a^3*b^5 + 130*a^2*b^5 - 116*I*a*b^5 - 45*b^5)/b^9)*x - (-24*I*a^4*b^4 - 250*a^3*b^4 + 332*I*a^2*b^4 + 275*a*b^4 - 64*I*b^4)/b^9) - 1/8*(8*a^4 - 16*I*a^3 - 24*a^2 + 12*I*a + 3)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^4*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^4 \sqrt{(a+bx)^2+1}}{1+a \operatorname{li}+b x \operatorname{li}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*((a + b*x)^2 + 1)^(1/2))/(a*1i + b*x*1i + 1),x)

[Out] int((x^4*((a + b*x)^2 + 1)^(1/2))/(a*1i + b*x*1i + 1), x)

3.190 $\int e^{-i\text{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=201

$$\frac{(3 + 12ia - 12a^2 - 8ia^3) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{8b^4} + \frac{x^2(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{4b^2} - \frac{(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{4b^2}$$

[Out] $-1/8*(3*I-12*a-12*I*a^2+8*a^3)*\text{arcsinh}(b*x+a)/b^4+1/4*x^2*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2-1/24*(1-I*a-I*b*x)^{(3/2)}*(7+10*I*a-18*a^2-2*(I-6*a)*b*x)*(1+I*a+I*b*x)^{(1/2)}/b^4-1/8*(3+12*I*a-12*a^2-8*I*a^3)*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^4$

Rubi [A]

time = 0.14, antiderivative size = 201, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 102, 152, 52, 55, 633, 221}

$$\frac{(-ia - ibx + 1)^{3/2} \sqrt{ia + ibx + 1} (-18a^2 - 2(-6a + i)bx + 10ia + 7)}{24b^4} - \frac{(-8ia^3 - 12a^2 + 12ia + 3) \sqrt{-ia - ibx + 1} \sqrt{ia + ibx + 1}}{8b^4} - \frac{(8a^3 - 12ia^2 - 12a + 3i) \sinh^{-1}(a + bx)}{8b^4} + \frac{x^2(-ia - ibx + 1)^{3/2} \sqrt{ia + ibx + 1}}{4b^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{(I*\text{ArcTan}[a + b*x])}, x]$

[Out] $-1/8*((3 + (12*I)*a - 12*a^2 - (8*I)*a^3)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/b^4 + (x^2*(1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x])/(4*b^2) - ((1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x]*(7 + (10*I)*a - 18*a^2 - 2*(I - 6*a)*b*x))/(24*b^4) - ((3*I - 12*a - (12*I)*a^2 + 8*a^3)*\text{ArcSinh}[a + b*x])/(8*b^4)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^n/(b*(m + n + 1))}], x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))], \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 55

$\text{Int}[1/(\text{Sqrt}[(a_.) + (b_.)*(x_.)]*\text{Sqrt}[(c_.) + (d_.)*(x_.)]), x_Symbol] \rightarrow \text{Int}[1/\text{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /; \text{FreeQ}\{a, b, c, d\}, x] \&\& \text{EqQ}[b + d, 0] \&\& \text{GtQ}[a + c, 0]$

Rule 102

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a + b*x)^{(m - 1)}*(c + d*x)^{(n + 1)}*(e + f*x)^p, x]$

```
)^(p + 1)/(d*f*(m + n + p + 1)), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a
+ b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b
*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1))] + b*(a*d*f*(2*m + n + p) - b*
(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p
}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_
))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-(a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{-i \tan^{-1}(a+bx)} x^3 dx &= \int \frac{x^3 \sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} dx \\
&= \frac{x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} + \frac{\int \frac{x \sqrt{1-ia-ibx}^{(-2(1+a^2)+(i-6a)bx)}}{\sqrt{1+ia+ibx}} dx}{4b^2} \\
&= \frac{x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} - \frac{(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx} (7+10ia-18a^2)}{24b^4} \\
&= -\frac{(3+12ia-12a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} \\
&= -\frac{(3+12ia-12a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} \\
&= -\frac{(3+12ia-12a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} \\
&= -\frac{(3+12ia-12a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} + \frac{x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2}
\end{aligned}$$

Mathematica [A]

time = 0.11, size = 118, normalized size = 0.59

$$-\frac{\sqrt{1+a^2+2abx+b^2x^2}(16+39ia-44a^2-6ia^3+(-9i+20a+6ia^2)bx-2(4+3ia)b^2x^2+6ib^3x^3)+3(3i-12a-12ia^2+8a^3)\sinh^{-1}(a+bx)}{24b^4}$$

Antiderivative was successfully verified.

[In] Integrate[x^3/E^(I*ArcTan[a + b*x]),x]

[Out] -1/24*(Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]*(16 + (39*I)*a - 44*a^2 - (6*I)*a^3 + (-9*I + 20*a + (6*I)*a^2)*b*x - 2*(4 + (3*I)*a)*b^2*x^2 + (6*I)*b^3*x^3) + 3*(3*I - 12*a - (12*I)*a^2 + 8*a^3)*ArcSinh[a + b*x])/b^4

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 680 vs. 2(163) = 326.

time = 0.12, size = 681, normalized size = 3.39

method	result
risch	$\frac{i(-6b^3x^3+6ab^2x^2-8ib^2x^2-6a^2bx+20iabx+6a^3-44ia^2+9bx-39a+16i)\sqrt{b^2x^2+2abx+a^2+1}}{24b^4} + \frac{3i \ln\left(\frac{b^2x+ab+\sqrt{b^2x^2+2abx+a^2+1}}{\sqrt{b^2x^2+2abx+a^2+1}}\right)}{24b^4}$

default	$i \frac{x \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4b^2} - \frac{5a \left(\frac{b^2 x^2 + 2abx + a^2 + 1}{3b^2} \right)^{\frac{3}{2}} - \left(\frac{(2b^2 x + 2ab) \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4b^2} + \frac{(4b^2(a^2 + 1) - 4a^2 b^2) \ln \left(\frac{b^2 x^2 + 2abx + a^2 + 1}{b^2} \right)}{b} \right)}{4b}$
---------	---

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^3/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-I/b*(1/4*x*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-5/4*a/b*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-a/b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-1/4*(a^2+1)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-I*(I-a)/b^2*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-a/b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))-I*(I-a)^2/b^3*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+ (I*a^3-3*I*a+3*a^2-1)/b^4*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+I*b*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 308 vs. $2(145) = 290$.

time = 0.49, size = 308, normalized size = 1.53

$$\frac{3\sqrt{b^2x^2+2abx+a^2+1}a^2}{2b^3} - \frac{a^2 \operatorname{arcsinh}(bx+a)}{b^3} - \frac{\sqrt{b^2x^2+2abx+a^2+1}a^3}{2b^4} - \frac{(b^2x^2+2abx+a^2+1)^{3/2}}{4b^4} - \frac{3\sqrt{b^2x^2+2abx+a^2+1}ax}{2b^4} - \frac{3a^2 \operatorname{arcsinh}(bx+a)}{2b^4} - \frac{3((b^2x^2+2abx+a^2+1)^{3/2})}{4b^4} - \frac{3\sqrt{b^2x^2+2abx+a^2+1}a^2}{2b^4} - \frac{3a \operatorname{arcsinh}(bx+a)}{2b^4} - \frac{3a \operatorname{arcsinh}(bx+a)}{2b^4} - \frac{10\sqrt{b^2x^2+2abx+a^2+1}a}{8b^4} - \frac{3a \operatorname{arcsinh}(bx+a)}{8b^4} - \frac{\sqrt{b^2x^2+2abx+a^2+1}}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="maxima")`

[Out]
$$-3/2*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^2*x/b^3 - a^3*\operatorname{arcsinh}(b*x + a)/b^4 - 1/2*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^3/b^4 - 1/4*I*(b^2*x^2 + 2*a$$

$$*b*x + a^2 + 1)^{(3/2)}*x/b^3 - 3/2*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a*x/b^3 + 3/2*I*a^2*\text{arcsinh}(b*x + a)/b^4 + 3/4*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a/b^4 + 3/2*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a^2/b^4 + 5/8*I*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*x/b^3 + 3/2*a*\text{arcsinh}(b*x + a)/b^4 + 1/3*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}/b^4 - 19/8*I*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a/b^4 - 3/8*I*\text{arcsinh}(b*x + a)/b^4 - \text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^4$$

Fricas [A]

time = 4.48, size = 139, normalized size = 0.69

$$\frac{45i a^4 + 224 a^3 - 192i a^2 + 24(8 a^3 - 12i a^2 - 12 a + 3i) \log(-bx - a + \sqrt{b^2 x^2 + 2 abx + a^2 + 1}) - 8(6i b^3 x^3 + 2(-3i a - 4)b^2 x^2 - 6i a^3 + (6i a^2 + 20 a - 9i)bx - 44 a^2 + 39i a + 16)\sqrt{b^2 x^2 + 2 abx + a^2 + 1} - 72 a}{192 b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="fricas")

[Out] 1/192*(45*I*a^4 + 224*a^3 - 192*I*a^2 + 24*(8*a^3 - 12*I*a^2 - 12*a + 3*I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 8*(6*I*b^3*x^3 + 2*(-3*I*a - 4)*b^2*x^2 - 6*I*a^3 + (6*I*a^2 + 20*a - 9*I)*b*x - 44*a^2 + 39*I*a + 16)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) - 72*a)/b^4

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^3 \sqrt{a^2 + 2abx + b^2 x^2 + 1}}{a + bx - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2),x)

[Out] -I*Integral(x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a + b*x - I), x)

Giac [A]

time = 0.44, size = 155, normalized size = 0.77

$$-\frac{1}{24} \sqrt{(bx+a)^2+1} \left(\left(2x \left(\frac{3ix}{b} - \frac{3iab^5+4b^5}{b^7} \right) - \frac{-6ia^2b^4-20ab^4+9ib^4}{b^7} \right) x - \frac{6ia^3b^3+44a^2b^3-39iab^3-16b^3}{b^7} \right) + \frac{(8a^3-12ia^2-12a+3i) \log(-ab - (x|b| - \sqrt{(bx+a)^2+1})|b|)}{8b^2|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="giac")

[Out] -1/24*sqrt((b*x + a)^2 + 1)*((2*x*(3*I*x/b - (3*I*a*b^5 + 4*b^5)/b^7) - (-6*I*a^2*b^4 - 20*a*b^4 + 9*I*b^4)/b^7)*x - (6*I*a^3*b^3 + 44*a^2*b^3 - 39*I*a*b^3 - 16*b^3)/b^7) + 1/8*(8*a^3 - 12*I*a^2 - 12*a + 3*I)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^3*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3 \sqrt{(a+bx)^2+1}}{1+a \operatorname{li}+bx \operatorname{li}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^3*((a + b*x)^2 + 1)^(1/2))/(a*x + b*x^2 + 1), x)
```

```
[Out] int((x^3*((a + b*x)^2 + 1)^(1/2))/(a*x + b*x^2 + 1), x)
```

3.191 $\int e^{-i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=171

$$\frac{(i - 2a - 2ia^2) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{2b^3} + \frac{(i - 4a)(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{6b^3} + \frac{x(1 - ia - ibx)^{3/2}}{3b^2}$$

[Out] $-1/2*(1+2*I*a-2*a^2)*\text{arcsinh}(b*x+a)/b^3+1/6*(I-4*a)*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3+1/3*x*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2+1/2*(I-2*a-2*I*a^2)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3$

Rubi [A]

time = 0.10, antiderivative size = 171, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 92, 81, 52, 55, 633, 221}

$$\frac{(-2ia^2 - 2a + i) \sqrt{ia + ibx + 1} \sqrt{-ia - ibx + 1}}{2b^3} - \frac{(-2a^2 + 2ia + 1) \sinh^{-1}(a + bx)}{2b^3} + \frac{(-4a + i) \sqrt{ia + ibx + 1} (-ia - ibx + 1)^{3/2}}{6b^3} + \frac{x \sqrt{ia + ibx + 1} (-ia - ibx + 1)^{3/2}}{3b^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/E^{(I*\text{ArcTan}[a + b*x])}, x]$

[Out] $((I - 2*a - (2*I)*a^2)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/(2*b^3) + ((I - 4*a)*(1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x])/(6*b^3) + (x*(1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x])/(3*b^2) - ((1 + (2*I)*a - 2*a^2)*\text{ArcSinh}[a + b*x])/(2*b^3)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x$ && $\text{NeQ}[b*c - a*d, 0]$ && $\text{GtQ}[n, 0]$ && $\text{NeQ}[m + n + 1, 0]$ && $!(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0])))$ && $!\text{ILtQ}[m + n + 2, 0]$ && $\text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 55

$\text{Int}[1/(\text{Sqrt}[(a_.) + (b_.)*(x_.)]*\text{Sqrt}[(c_.) + (d_.)*(x_.)]), x_Symbol] \rightarrow \text{Int}[1/\text{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x$ && $\text{EqQ}[b + d, 0]$ && $\text{GtQ}[a + c, 0]$

Rule 81

$\text{Int}[(a_. + (b_.)*(x_.))*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[b*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 2))), x] + \text{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /;$ $\text{FreeQ}\{a, b, c, d, e, f$

, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92

Int[((a_.) + (b_.)*(x_))²((c_.) + (d_.)*(x_))^(n_.)((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)ⁿ(e + f*x)^p*Simp[a²*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)²], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)²)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b² - 4*a*c)))^p), Subst[Int[Simp[1 - x²/(b² - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b²/c, 0]

Rule 5203

Int[E^{(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)}((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m((1 - I*a*c - I*b*c*x)^{(I*(n/2))}/(1 + I*a*c + I*b*c*x)^{(I*(n/2))}), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{-i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2 \sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} dx \\
&= \frac{x(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{3b^2} + \frac{\int \frac{\sqrt{1-ia-ibx} (-1-a^2+(i-4a)bx)}{\sqrt{1+ia+ibx}} dx}{3b^2} \\
&= \frac{(i-4a)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{6b^3} + \frac{x(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{3b^2} - \frac{(1+ia+ibx)^{3/2} \sqrt{1+ia+ibx}}{3b^2} \\
&= -\frac{(2a-i(1-2a^2)) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} + \frac{(i-4a)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{6b^3} \\
&= -\frac{(2a-i(1-2a^2)) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} + \frac{(i-4a)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{6b^3} \\
&= -\frac{(2a-i(1-2a^2)) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} + \frac{(i-4a)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{6b^3} \\
&= -\frac{(2a-i(1-2a^2)) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} + \frac{(i-4a)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{6b^3} \\
&= -\frac{(2a-i(1-2a^2)) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^3} + \frac{(i-4a)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{6b^3}
\end{aligned}$$

Mathematica [A]

time = 0.21, size = 162, normalized size = 0.95

$$\frac{i\sqrt{1+ia+ibx}(4+7a^2+2ia^3-7ibx-5b^2x^2+2ib^3x^3+a(5i+8bx))}{6b^3\sqrt{-i(i+a+bx)}} + \frac{\sqrt[4]{-1}(-1-2ia+2a^2)\sqrt{-ib}\sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{b^{7/2}}$$

Antiderivative was successfully verified.

`[In] Integrate[x^2/E^(I*ArcTan[a + b*x]),x]`

```
[Out] ((I/6)*Sqrt[1 + I*a + I*b*x]*(4 + 7*a^2 + (2*I)*a^3 - (7*I)*b*x - 5*b^2*x^2 + (2*I)*b^3*x^3 + a*(5*I + 8*b*x)))/(b^3*Sqrt[(-I)*(I + a + b*x)]) + ((-1)^(1/4)*(-1 - (2*I)*a + 2*a^2)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/b^(7/2)
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 484 vs. 2(135) = 270.

time = 0.10, size = 485, normalized size = 2.84

method	result
--------	--------

risch	$\frac{i(2b^2x^2 - 2abx + 3ibx + 2a^2 - 9ia - 4)\sqrt{b^2x^2 + 2abx + a^2 + 1}}{6b^3} - \frac{i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)a}{b^2\sqrt{b^2}} + \dots$
default	$\frac{\left(\frac{(b^2x^2 + 2abx + a^2 + 1)^{\frac{3}{2}}}{3b^2}\right)^{\frac{3}{2}} - \frac{a \left(\frac{(2b^2x + 2ab)\sqrt{b^2x^2 + 2abx + a^2 + 1}}{4b^2} + \frac{(4b^2(a^2 + 1) - 4a^2b^2) \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{8b^2\sqrt{b^2}} \right)}{b}}{b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out]
$$-I/b^2*(b*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)/b^2-a/b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2)))+I*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))-a*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2)))+(-I*a^2+I-2*a)/b^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+I*b*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))$$

Maxima [A]

time = 0.47, size = 161, normalized size = 0.94

$$\frac{i\sqrt{b^2x^2+2abx+a^2+1}ax}{b^2} + \frac{a^2 \operatorname{arsinh}(bx+a)}{b^2} + \frac{\sqrt{b^2x^2+2abx+a^2+1}x}{2b^2} - \frac{ia \operatorname{arsinh}(bx+a)}{b^3} - \frac{i(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}}{3b^3} - \frac{3\sqrt{b^2x^2+2abx+a^2+1}a}{2b^3} - \frac{\operatorname{arsinh}(bx+a)}{2b^2} + \frac{i\sqrt{b^2x^2+2abx+a^2+1}}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="maxima")`

[Out]
$$I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a*x/b^2 + a^2*\operatorname{arcsinh}(b*x + a)/b^3 + 1/2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*x/b^2 - I*a*\operatorname{arcsinh}(b*x + a)/b^3 - 1/3*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)/b^3 - 3/2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a/b^3 - 1/2*\operatorname{arcsinh}(b*x + a)/b^3 + I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/b^3$$

Fricas [A]

time = 4.60, size = 106, normalized size = 0.62

$$\frac{-7i a^3 - 21 a^2 - 12(2 a^2 - 2i a - 1) \log\left(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right) - 4\sqrt{b^2x^2 + 2abx + a^2 + 1}(2ib^2x^2 + (-2ia - 3)bx + 2ia^2 + 9a - 4i) + 9ia}{24b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="fricas")

[Out] 1/24*(-7*I*a^3 - 21*a^2 - 12*(2*a^2 - 2*I*a - 1)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(2*I*b^2*x^2 + (-2*I*a - 3)*b*x + 2*I*a^2 + 9*a - 4*I) + 9*I*a)/b^3

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^2 \sqrt{a^2 + 2abx + b^2x^2 + 1}}{a + bx - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2),x)

[Out] -I*Integral(x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a + b*x - I), x)

Giac [A]

time = 0.47, size = 113, normalized size = 0.66

$$-\frac{1}{6} \sqrt{(bx+a)^2+1} \left(x \left(\frac{2ix}{b} - \frac{2iab^3+3b^3}{b^5} \right) - \frac{-2ia^2b^2-9ab^2+4ib^2}{b^5} \right) - \frac{(2a^2-2ia-1) \log \left(-ab - \left(x|b| - \sqrt{(bx+a)^2+1} \right) |b| \right)}{2b^2|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="giac")

[Out] -1/6*sqrt((b*x + a)^2 + 1)*(x*(2*I*x/b - (2*I*a*b^3 + 3*b^3)/b^5) - (-2*I*a^2*b^2 - 9*a*b^2 + 4*I*b^2)/b^5) - 1/2*(2*a^2 - 2*I*a - 1)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b^2*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^2 \sqrt{(a+bx)^2+1}}{1+ali+bxli} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*((a + b*x)^2 + 1)^(1/2))/(a*1i + b*x*1i + 1),x)

[Out] int((x^2*((a + b*x)^2 + 1)^(1/2))/(a*1i + b*x*1i + 1), x)

3.192 $\int e^{-i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=110

$$\frac{(1+2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} + \frac{(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{2b^2} + \frac{(i-2a)\sinh^{-1}(a+bx)}{2b^2}$$

[Out] 1/2*(I-2*a)*arcsinh(b*x+a)/b^2+1/2*(1-I*a-I*b*x)^(3/2)*(1+I*a+I*b*x)^(1/2)/b^2+1/2*(1+2*I*a)*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/b^2

Rubi [A]

time = 0.06, antiderivative size = 110, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5203, 81, 52, 55, 633, 221}

$$\frac{\sqrt{ia+ibx+1}(-ia-ibx+1)^{3/2}}{2b^2} + \frac{(1+2ia)\sqrt{ia+ibx+1}\sqrt{-ia-ibx+1}}{2b^2} + \frac{(-2a+i)\sinh^{-1}(a+bx)}{2b^2}$$

Antiderivative was successfully verified.

[In] Int[x/E^(I*ArcTan[a + b*x]),x]

[Out] ((1 + (2*I)*a)*Sqrt[1 - I*a - I*b*x]*Sqrt[1 + I*a + I*b*x])/(2*b^2) + ((1 - I*a - I*b*x)^(3/2)*Sqrt[1 + I*a + I*b*x])/(2*b^2) + ((I - 2*a)*ArcSinh[a + b*x])/(2*b^2)

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]

Rule 81

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{-i \tan^{-1}(a+bx)} x \, dx &= \int \frac{x \sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} \, dx \\
 &= \frac{(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(i - 2a) \int \frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} \, dx}{2b} \\
 &= \frac{(1 + 2ia) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(i - 2a)}{2b} \\
 &= \frac{(1 + 2ia) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(i - 2a)}{2b} \\
 &= \frac{(1 + 2ia) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(i - 2a)}{2b} \\
 &= \frac{(1 + 2ia) \sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(1 - ia - ibx)^{3/2} \sqrt{1 + ia + ibx}}{2b^2} + \frac{(i - 2a)}{2b}
 \end{aligned}$$

Mathematica [A]

time = 0.09, size = 131, normalized size = 1.19

$$\frac{\sqrt{1 + ia + ibx} (2 - ia + a^2 - 3ibx - b^2x^2)}{2b^2 \sqrt{-i(i + a + bx)}} + \frac{(-1)^{3/4} (1 + 2ia) \sqrt{-ib} \sinh^{-1} \left(\frac{(\frac{1}{2} + \frac{i}{2}) \sqrt{b} \sqrt{-i(i + a + bx)}}{\sqrt{-ib}} \right)}{b^{5/2}}$$

Antiderivative was successfully verified.

[In] Integrate[x/E^(I*ArcTan[a + b*x]),x]

[Out] (Sqrt[1 + I*a + I*b*x]*(2 - I*a + a^2 - (3*I)*b*x - b^2*x^2))/(2*b^2*Sqrt[(-I)*(I + a + b*x)]) + ((-1)^(3/4)*(1 + (2*I)*a)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])]/Sqrt[(-I)*b])/b^(5/2)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 236 vs. 2(86) = 172.
time = 0.11, size = 237, normalized size = 2.15

method	result
risch	$\frac{i(-bx+a-2i)\sqrt{b^2x^2+2abx+a^2+1}}{2b^2} + \frac{i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{2b\sqrt{b^2}} - \frac{a \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{b\sqrt{b^2}}$
default	$- \frac{i \left(\frac{(2b^2x+2ab)\sqrt{b^2x^2+2abx+a^2+1}}{4b^2} + \frac{(4b^2(a^2+1)-4a^2b^2) \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{8b^2\sqrt{b^2}} \right)}{b} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x,method=_RETURNVERBOSE)

[Out] -I/b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2)+(1+I*a)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+I*b*ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))

Maxima [A]

time = 0.48, size = 97, normalized size = 0.88

$$-\frac{i\sqrt{b^2x^2+2abx+a^2+1}x}{2b} - \frac{a \operatorname{arsinh}(bx+a)}{b^2} + \frac{i\sqrt{b^2x^2+2abx+a^2+1}a}{2b^2} + \frac{i \operatorname{arsinh}(bx+a)}{2b^2} + \frac{\sqrt{b^2x^2+2abx+a^2+1}}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="maxima")

[Out] -1/2*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*x/b - a*arcsinh(b*x + a)/b^2 + 1/2*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*a/b^2 + 1/2*I*arcsinh(b*x + a)/b^2 + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^2

Fricas [A]

time = 4.73, size = 79, normalized size = 0.72

$$\frac{3i a^2 + 4(2a - i) \log\left(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right) - 4\sqrt{b^2x^2 + 2abx + a^2 + 1}(ibx - ia - 2) + 4a}{8b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="fricas")

[Out] 1/8*(3*I*a^2 + 4*(2*a - I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(I*b*x - I*a - 2) + 4*a)/b^2

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x \sqrt{a^2 + 2abx + b^2x^2 + 1}}{a + bx - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2),x)

[Out] -I*Integral(x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a + b*x - I), x)

Giac [A]

time = 0.45, size = 75, normalized size = 0.68

$$-\frac{1}{2} \sqrt{(bx+a)^2+1} \left(\frac{ix}{b} + \frac{-iab-2b}{b^3} \right) + \frac{(2a-i) \log \left(-ab - \left(x|b| - \sqrt{(bx+a)^2+1} \right) |b| \right)}{2b|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="giac")

[Out] -1/2*sqrt((b*x + a)^2 + 1)*(I*x/b + (-I*a*b - 2*b)/b^3) + 1/2*(2*a - I)*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x \sqrt{(a + bx)^2 + 1}}{1 + a \operatorname{li} + b x \operatorname{li}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*((a + b*x)^2 + 1)^(1/2))/(a*1i + b*x*1i + 1),x)

[Out] int((x*((a + b*x)^2 + 1)^(1/2))/(a*1i + b*x*1i + 1), x)

3.193 $\int e^{-i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=52

$$-\frac{i\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{b} + \frac{\sinh^{-1}(a+bx)}{b}$$

[Out] arcsinh(b*x+a)/b-I*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/b

Rubi [A]

time = 0.02, antiderivative size = 52, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.417$, Rules used = {5201, 52, 55, 633, 221}

$$\frac{\sinh^{-1}(a+bx)}{b} - \frac{i\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{b}$$

Antiderivative was successfully verified.

[In] Int[E^((-I)*ArcTan[a + b*x]),x]

[Out] ((-I)*Sqrt[1 - I*a - I*b*x]*Sqrt[1 + I*a + I*b*x])/b + ArcSinh[a + b*x]/b

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b

+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5201

Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))]*(n_.)), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{-i \tan^{-1}(a+bx)} dx &= \int \frac{\sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} dx \\
 &= -\frac{i\sqrt{1-ia-ibx}}{b} \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} + \int \frac{1}{\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx \\
 &= -\frac{i\sqrt{1-ia-ibx}}{b} \frac{\sqrt{1+ia+ibx}}{\sqrt{(1-ia)(1+ia)+2abx+b^2x^2}} + \int \frac{1}{\sqrt{(1-ia)(1+ia)+2abx+b^2x^2}} dx \\
 &= -\frac{i\sqrt{1-ia-ibx}}{b} \frac{\sqrt{1+ia+ibx}}{\sqrt{1+\frac{x^2}{4b^2}}} + \frac{\text{Subst}\left(\int \frac{1}{\sqrt{1+\frac{x^2}{4b^2}}} dx, x, 2ab+2b^2x\right)}{2b^2} \\
 &= -\frac{i\sqrt{1-ia-ibx}}{b} \frac{\sqrt{1+ia+ibx}}{\sqrt{1+(a+bx)^2}} + \frac{\sinh^{-1}(a+bx)}{b}
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 28, normalized size = 0.54

$$\frac{-i\sqrt{1+(a+bx)^2} + \sinh^{-1}(a+bx)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[E^((-I)*ArcTan[a + b*x]),x]

[Out] ((-I)*Sqrt[1 + (a + b*x)^2] + ArcSinh[a + b*x])/b

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 124 vs. 2(43) = 86.

time = 0.09, size = 125, normalized size = 2.40

method	result	size
--------	--------	------

risch	$-\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b} + \frac{\ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{\sqrt{b^2}}$	69
default	$-\frac{i\left(\sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib\left(x - \frac{i-a}{b}\right)} + \frac{ib\ln\left(\frac{ib + \left(x - \frac{i-a}{b}\right)b^2}{\sqrt{b^2}} + \sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib\left(x - \frac{i-a}{b}\right)}\right)}{\sqrt{b^2}}\right)}{b}$	125

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-I/b*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+I*b*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))$

Maxima [A]

time = 0.47, size = 35, normalized size = 0.67

$$\frac{\operatorname{arsinh}(bx+a)}{b} - \frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="maxima")`

[Out] $\operatorname{arcsinh}(b*x + a)/b - I*\operatorname{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)/b$

Fricas [A]

time = 2.96, size = 60, normalized size = 1.15

$$\frac{-ia - 2i\sqrt{b^2x^2 + 2abx + a^2 + 1} - 2\log\left(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="fricas")`

[Out] $1/2*(-I*a - 2*I*\operatorname{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1) - 2*\log(-b*x - a + \operatorname{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)))/b$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a + bx - i} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2),x)

[Out] -I*Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a + b*x - I), x)

Giac [A]

time = 0.43, size = 51, normalized size = 0.98

$$-\frac{\log\left(-ab - \left(x|b| - \sqrt{(bx+a)^2 + 1}\right)|b|\right)}{|b|} - \frac{i\sqrt{(bx+a)^2 + 1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2),x, algorithm="giac")

[Out] -log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/abs(b) - I*sqrt((b*x + a)^2 + 1)/b

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{(a+bx)^2 + 1}}{1 + a \operatorname{li} + b x \operatorname{li}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)^(1/2)/(a*1i + b*x*1i + 1),x)

[Out] int(((a + b*x)^2 + 1)^(1/2)/(a*1i + b*x*1i + 1), x)

$$3.194 \quad \int \frac{e^{-i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=89

$$-i \sinh^{-1}(a+bx) - \frac{2\sqrt{i+a} \tanh^{-1}\left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}}\right)}{\sqrt{i-a}}$$

[Out] $-I*\operatorname{arcsinh}(b*x+a)-2*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(I-a)^{(1/2)}/(1-I*a-I*b*x)^{(1/2}))*I+a)^{(1/2)}/(I-a)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 132, 55, 633, 221, 12, 95, 214}

$$-i \sinh^{-1}(a+bx) - \frac{2\sqrt{a+i} \tanh^{-1}\left(\frac{\sqrt{a+i} \sqrt{ia+ibx+1}}{\sqrt{-a+i} \sqrt{-ia-ibx+1}}\right)}{\sqrt{-a+i}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(I*ArcTan[a + b*x]))*x],x]`

[Out] $(-I)*\operatorname{ArcSinh}[a + b*x] - (2*\operatorname{Sqrt}[I + a]*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I + a]*\operatorname{Sqrt}[1 + I*a + I*b*x])/(\operatorname{Sqrt}[I - a]*\operatorname{Sqrt}[1 - I*a - I*b*x])])/(\operatorname{Sqrt}[I - a])$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 55

`Int[1/((Sqrt[(a_) + (b_.)*(x_)])*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`

Rule 95

`Int[(((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_)]/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

Rule 132

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{\sqrt{1-ia-ibx}}{x\sqrt{1+ia+ibx}} dx \\
&= -\left((-1+ia) \int \frac{1}{x\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx \right) - (ib) \int \frac{1}{\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx \\
&= (2(1-ia)) \text{Subst} \left(\int \frac{1}{-1-ia-(-1+ia)x^2} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} \right) - (ib) \int \frac{1}{\sqrt{(1-ia-ibx)(1+ia+ibx)}} dx \\
&= \frac{2\sqrt{i+a} \tanh^{-1} \left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}} \right)}{\sqrt{i-a}} - \frac{i \text{Subst} \left(\int \frac{1}{\sqrt{1+\frac{x^2}{4b^2}}} dx, x, 2ab + \dots \right)}{2b} \\
&= -i \sinh^{-1}(a+bx) - \frac{2\sqrt{i+a} \tanh^{-1} \left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}} \right)}{\sqrt{i-a}}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 142, normalized size = 1.60

$$\frac{2\sqrt{-1} (-ib)^{3/2} \sinh^{-1} \left(\frac{(\frac{1}{2} + \frac{i}{2}) \sqrt{b} \sqrt{-i(i+a+bx)}}{\sqrt{-ib}} \right)}{b^{3/2}} - \frac{2\sqrt{-1+ia} \tanh^{-1} \left(\frac{\sqrt{-1-ia} \sqrt{-i(i+a+bx)}}{\sqrt{-1+ia} \sqrt{1+ia+ibx}} \right)}{\sqrt{-1-ia}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^(I*ArcTan[a + b*x]))*x), x]`

```
[Out] (2*(-1)^(1/4)*((-I)*b)^(3/2)*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/b^(3/2) - (2*Sqrt[-1 + I*a]*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]])/Sqrt[-1 - I*a]
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 259 vs. 2(68) = 136.

time = 0.09, size = 260, normalized size = 2.92

method	result
default	$ -\frac{i \left(\sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib \left(x - \frac{i-a}{b}\right)} + \frac{i b \ln \left(\frac{ib + \left(x - \frac{i-a}{b}\right) b^2 + \sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib \left(x - \frac{i-a}{b}\right)}}{\sqrt{b^2}} \right)}{\sqrt{b^2}} \right)}{i-a} + \dots $

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x,x,method=_RETURNVERBOSE)`

[Out]
$$-I/(I-a)*(((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+I*b*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))+I/(I-a)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))$$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: ValueError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x,x, algorithm="maxima")`

[Out] Exception raised: ValueError >> Computation failed since Maxima requested additional constraints; using the 'assume' command before evaluation *may* help (example of legal syntax is 'assume(a-1>0)', see 'assume?' for more details)Is

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 144 vs. 2(59) = 118.

time = 2.53, size = 144, normalized size = 1.62

$$-\sqrt{\frac{a+i}{a-i}} \log\left(-bx+(ia+1)\sqrt{\frac{a+i}{a-i}+\sqrt{b^2x^2+2abx+a^2+1}}\right) + \sqrt{\frac{a+i}{a-i}} \log\left(-bx+(-ia-1)\sqrt{\frac{a+i}{a-i}+\sqrt{b^2x^2+2abx+a^2+1}}\right) + i \log\left(-bx-a+\sqrt{b^2x^2+2abx+a^2+1}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x,x, algorithm="fricas")`

[Out]
$$-\sqrt{-a+I}/(a-I)*\log(-b*x+(I*a+1)*\sqrt{-a+I}/(a-I))+\sqrt{b^2*x^2+2*a*b*x+a^2+1}+\sqrt{-a+I}/(a-I)*\log(-b*x+(-I*a-1)*\sqrt{-a+I}/(a-I))+\sqrt{b^2*x^2+2*a*b*x+a^2+1}+I*\log(-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2+1})$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2+2abx+b^2x^2+1}}{ax+bx^2-ix} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2)/x,x)

[Out] -I*Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a*x + b*x**2 - I*x), x)

Giac [A]

time = 0.46, size = 76, normalized size = 0.85

$$\frac{2(a+i) \arctan\left(\frac{-ix|b|+i\sqrt{(bx+a)^2+1}}{\sqrt{a^2+1}}\right)}{\sqrt{a^2+1}} + \frac{ib \log\left(-ab - \left(x|b| - \sqrt{(bx+a)^2+1}\right)|b|\right)}{|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x,x, algorithm="giac")

[Out] 2*(a + I)*arctan((-I*x*abs(b) + I*sqrt((b*x + a)^2 + 1))/sqrt(a^2 + 1))/sqrt(a^2 + 1) + I*b*log(-a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/abs(b)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{(a+bx)^2+1}}{x(1+ali+bxli)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)^(1/2)/(x*(a*1i + b*x*1i + 1)),x)

[Out] int(((a + b*x)^2 + 1)^(1/2)/(x*(a*1i + b*x*1i + 1)), x)

$$3.195 \quad \int \frac{e^{-i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=130

$$-\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1+ia)x} - \frac{2ib \tanh^{-1} \left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}} \right)}{(i-a)^{3/2} \sqrt{i+a}}$$

[Out] $-2*I*b*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(I-a)^{(1/2)}/(1-I*a-I*b*x)^{(1/2)})/(I-a)^{(3/2)}/(I+a)^{(1/2)}-(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(1+I*a)/x$

Rubi [A]

time = 0.05, antiderivative size = 130, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5203, 96, 95, 214}

$$-\frac{\sqrt{-ia-ibx+1} \sqrt{ia+ibx+1}}{(1+ia)x} - \frac{2ib \tanh^{-1} \left(\frac{\sqrt{a+i} \sqrt{ia+ibx+1}}{\sqrt{-a+i} \sqrt{-ia-ibx+1}} \right)}{(-a+i)^{3/2} \sqrt{a+i}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(I*ArcTan[a + b*x])*x^2),x]`

[Out] $-\left(\frac{\sqrt{1-I*a-I*b*x}*\sqrt{1+I*a+I*b*x}}{(1+I*a)*x}\right) - \left(\frac{(2*I)*b*\operatorname{ArcTanh}\left[\frac{\sqrt{I+a}*\sqrt{1+I*a+I*b*x}}{\sqrt{I-a}*\sqrt{1-I*a-I*b*x}}\right]}{(I-a)^{(3/2)}*\sqrt{I+a}}\right)$

Rule 95

`Int[(((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_)]/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

Rule 96

`Int[((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]`

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_))*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{-i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{\sqrt{1-ia-ibx}}{x^2 \sqrt{1+ia+ibx}} dx \\ &= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1+ia)x} + \frac{b \int \frac{1}{x \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}} dx}{i-a} \\ &= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1+ia)x} + \frac{(2b) \text{Subst} \left(\int \frac{1}{-1-ia-(-1+ia)x^2} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}} \right)}{i-a} \\ &= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{(1+ia)x} - \frac{2ib \tanh^{-1} \left(\frac{\sqrt{i+a} \sqrt{1+ia+ibx}}{\sqrt{i-a} \sqrt{1-ia-ibx}} \right)}{(i-a)^{3/2} \sqrt{i+a}} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 119, normalized size = 0.92

$$i \left(\frac{\sqrt{1+a^2+2abx+b^2x^2}}{(-i+a)x} + \frac{2b \tanh^{-1} \left(\frac{\sqrt{-1-ia} \sqrt{-i(i+a+bx)}}{\sqrt{-1+ia} \sqrt{1+ia+ibx}} \right)}{(-1-ia)^{3/2} \sqrt{-1+ia}} \right)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^(I*ArcTan[a + b*x]))*x^2), x]

[Out] I*(Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]/((-I + a)*x) + (2*b*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x])])/((-1 - I*a)^(3/2)*Sqrt[-1 + I*a]))

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 545 vs. 2(100) = 200.

time = 0.10, size = 546, normalized size = 4.20

method	result
risch	$\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{(a-i)x} + \frac{b \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{(a-i)\sqrt{a^2+1}}$
default	$- \frac{ib \left(\sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib \left(x - \frac{i-a}{b}\right)} + \frac{ib \ln\left(\frac{ib + \left(x - \frac{i-a}{b}\right)b^2 + \sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib \left(x - \frac{i-a}{b}\right)}}{\sqrt{b^2}}\right)}{\sqrt{b^2}} \right)}{(i-a)^2} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^2,x,method=_RETURNVERBOSE)

[Out]
$$-I/(I-a)^2*b*\left(\left(\frac{x-(I-a)}{b}\right)^2*b^2+2*I*b*\left(\frac{x-(I-a)}{b}\right)\right)^{(1/2)}+I*b*\ln\left(\frac{I*b+\left(\frac{x-(I-a)}{b}\right)*b^2}{(b^2)^{(1/2)}+\left(\frac{x-(I-a)}{b}\right)^2*b^2+2*I*b*\left(\frac{x-(I-a)}{b}\right)}\right)^{(1/2)}\right)/(b^2)^{(1/2)}+I*b/(I-a)^2*\left(\frac{(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b*\ln\left(\frac{(b^2*x+a*b)}{(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}\right)}{(b^2)^{(1/2)}-(a^2+1)^{(1/2)}*\ln\left(\frac{2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}\right)}{x}\right)+I/(I-a)*\left(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)}+a*b/(a^2+1)*\left(\frac{(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b*\ln\left(\frac{(b^2*x+a*b)}{(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}\right)}{(b^2)^{(1/2)}-(a^2+1)^{(1/2)}*\ln\left(\frac{2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}\right)}{x}\right)+2*b^2/(a^2+1)*\left(\frac{1}{4}\right)*\frac{(2*b^2*x+2*a*b)}{b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln\left(\frac{(b^2*x+a*b)}{(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}\right)}{(b^2)^{(1/2)})}\right)\right)$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^2,x, algorithm="maxima")

[Out] integrate(sqrt((b*x + a)^2 + 1)/((I*b*x + I*a + 1)*x^2), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 224 vs. 2(86) = 172.

time = 3.84, size = 224, normalized size = 1.72

$$\frac{(a-i)\sqrt{\frac{b^2}{a^2-2ia^3-2ia-1}}x \log\left(\frac{\mu_{x-}\sqrt{b^2x^2+2abx+a^2+1}b^{i(a^2-i^2+a-1)}\sqrt{\frac{b^2}{a^4-2ia^3-2ia-1}}}{b}\right) - (a-i)\sqrt{\frac{b^2}{a^4-2ia^3-2ia-1}}x \log\left(\frac{\mu_{x-}\sqrt{b^2x^2+2abx+a^2+1}b^{-(a^2-i^2+a-1)}\sqrt{\frac{b^2}{a^4-2ia^3-2ia-1}}}{b}\right) - ibx - i\sqrt{b^2x^2+2abx+a^2+1}}{(a-i)x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^2,x, algorithm="fricas")

[Out] $-\left(\frac{(a - I)\sqrt{b^2/(a^4 - 2Ia^3 - 2Ia - 1)} * x \log(-(b^2x - \sqrt{b^2x^2 + 2abx + a^2 + 1})b + (a^3 - Ia^2 + a - I)\sqrt{b^2/(a^4 - 2Ia^3 - 2Ia - 1)})}{b} - \frac{(a - I)\sqrt{b^2/(a^4 - 2Ia^3 - 2Ia - 1)} * x \log(-(b^2x - \sqrt{b^2x^2 + 2abx + a^2 + 1})b - (a^3 - Ia^2 + a - I)\sqrt{b^2/(a^4 - 2Ia^3 - 2Ia - 1)})}{b} - Ibx - I\sqrt{b^2x^2 + 2abx + a^2 + 1}\right) / ((a - I)x)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{ax^2 + bx^3 - ix^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2)/x**2,x)

[Out] $-I * \text{Integral}(\sqrt{a^2 + 2abx + b^2x^2 + 1} / (ax^2 + b^2x^3 - Ix^2), x)$

Giac [A]

time = 0.48, size = 145, normalized size = 1.12

$$\frac{b \log \left(\frac{\left| \begin{matrix} 2x|b|-2\sqrt{(bx+a)^2+1} & -2\sqrt{a^2+1} \\ 2x|b|-2\sqrt{(bx+a)^2+1} & +2\sqrt{a^2+1} \end{matrix} \right|}{\sqrt{a^2+1}(a-i)} \right)}{\left(\left(|x|b| - \sqrt{(bx+a)^2+1} \right)^2 - a^2 - 1 \right) (-ia - 1)} - \frac{2 \left(\left(|x|b| - \sqrt{(bx+a)^2+1} \right) ab + a^2|b| + |b| \right)}{\left(\left(|x|b| - \sqrt{(bx+a)^2+1} \right)^2 - a^2 - 1 \right) (-ia - 1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^2,x, algorithm="giac")

[Out] $b \log(\text{abs}(2x \cdot \text{abs}(b) - 2\sqrt{(bx+a)^2+1}) - 2\sqrt{a^2+1}) / \text{abs}(2x \cdot \text{abs}(b) - 2\sqrt{(bx+a)^2+1}) + 2\sqrt{a^2+1}) / (\sqrt{a^2+1} \cdot (a - I)) - 2 \cdot ((x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1}) \cdot a \cdot b + a^2 \cdot \text{abs}(b) + \text{abs}(b)) / (((x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^2 - a^2 - 1) \cdot (-I \cdot a - 1))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{(a+bx)^2+1}}{x^2(1+alix+bxli)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a + b*x)^2 + 1)^(1/2)/(x^2*(a*1i + b*x*1i + 1)),x)
```

```
[Out] int(((a + b*x)^2 + 1)^(1/2)/(x^2*(a*1i + b*x*1i + 1)), x)
```

$$3.196 \quad \int \frac{e^{-i \operatorname{ArcTan}(a+bx)}}{x^3} dx$$

Optimal. Leaf size=201

$$\frac{(1-2ia)b\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2(i-a)^2(i+a)x} - \frac{(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{2(1+a^2)x^2} + \frac{(1-2ia)b^2 \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1-ia-ibx}}{\sqrt{i-a}\sqrt{1+ia+ibx}}\right)}{(i-a)^{5/2}(i+a)}$$

[Out] $(1-2I*a)*b^2*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(I-a)^{(1/2)}/(1-I*a-I*b*x)^{(1/2)})/(I-a)^{(5/2)}/(I+a)^{(3/2)}-1/2*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/(a^2+1)/x^2+1/2*(1-2I*a)*b*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(I-a)^2/(I+a)/x$

Rubi [A]

time = 0.11, antiderivative size = 201, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {5203, 98, 96, 95, 214}

$$-\frac{(-ia-ibx+1)^{3/2}\sqrt{ia+ibx+1}}{2(a^2+1)x^2} + \frac{(1-2ia)b^2 \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{5/2}(a+i)^{3/2}} + \frac{(1-2ia)b\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{2(-a+i)^2(a+i)x}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a + b*x]))*x^3], x]

[Out] $((1-(2I)*a)*b*\operatorname{Sqrt}[1-I*a-I*b*x]*\operatorname{Sqrt}[1+I*a+I*b*x])/(2*(I-a)^2*(I+a)*x) - ((1-I*a-I*b*x)^{(3/2)}*\operatorname{Sqrt}[1+I*a+I*b*x])/(2*(1+a^2)*x^2) + (((1-(2I)*a)*b^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])]/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])))/((I-a)^{(5/2)}*(I+a)^{(3/2)})$

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)), x_Symbol] :> Simp[(a + b*x)^(m+1)*(c + d*x)^n*((e + f*x)^(p+1))/((m+1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m+1)*(b*e - a*f))], Int[(a + b*x)^(m+1)*(c + d*x)^(n-1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 98

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, n, p}, x] && EqQ[Simplify[m + n + p + 3], 0] && (LtQ[m, -1] || SumSimplerQ[m, 1])
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(a+bx)}}{x^3} dx &= \int \frac{\sqrt{1-ia-ibx}}{x^3 \sqrt{1+ia+ibx}} dx \\
&= -\frac{(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{2(1+a^2)x^2} - \frac{((i+2a)b) \int \frac{\sqrt{1-ia-ibx}}{x^2 \sqrt{1+ia+ibx}} dx}{2(1+a^2)} \\
&= \frac{(1-2ia)b \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2(i-a)^2(i+a)x} - \frac{(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{2(1+a^2)x^2} + \frac{((i+2a)b) \int \frac{\sqrt{1-ia-ibx}}{x \sqrt{1+ia+ibx}} dx}{2(1+a^2)} \\
&= \frac{(1-2ia)b \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2(i-a)^2(i+a)x} - \frac{(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{2(1+a^2)x^2} + \frac{((i+2a)b) \int \frac{\sqrt{1-ia-ibx}}{x \sqrt{1+ia+ibx}} dx}{2(1+a^2)} \\
&= \frac{(1-2ia)b \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2(i-a)^2(i+a)x} - \frac{(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{2(1+a^2)x^2} + \frac{(1-2ia)b \int \frac{\sqrt{1-ia-ibx}}{x \sqrt{1+ia+ibx}} dx}{2(1+a^2)}
\end{aligned}$$

Mathematica [A]

time = 0.09, size = 154, normalized size = 0.77

$$\frac{i(1+a^2-2ibx-abx)\sqrt{1+a^2+2abx+b^2x^2}}{x^2} + \frac{2(i+2a)b^2 \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}{\sqrt{-1-ia}\sqrt{-1+ia}}$$

$$2(-i+a)^2(i+a)$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^(I*ArcTan[a + b*x])*x^3), x]

[Out] ((I*(1 + a^2 - (2*I)*b*x - a*b*x)*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2])/x^2 + (2*(I + 2*a)*b^2*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x])])/(Sqrt[-1 - I*a]*Sqrt[-1 + I*a]))/(2*(-I + a)^2*(I + a))

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1007 vs. 2(157) = 314.

time = 0.12, size = 1008, normalized size = 5.01

method	result
risch	$\frac{i(-ab^3x^3 - 2ib^3x^3 - a^2b^2x^2 - 4ia^2b^2x^2 + a^3bx - 2ia^2b + a^4 + b^2x^2 + abx - 2ibx + 2a^2 + 1)}{2x^2(i+a)(a-i)^2\sqrt{b^2x^2 + 2abx + a^2 + 1}} - \frac{ib^2 \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2}}{x}\right)}{2(a^2+1)^{\frac{3}{2}}(a-i)}$ $i \left[-\frac{(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}}{2(a^2+1)x^2} - \frac{ab \left[-\frac{(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}}{(a^2+1)x} + \frac{ab \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{\sqrt{b^2}}\right]}{(a^2+1)x} \right]$
default	

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^3, x, method=_RETURNVERBOSE)

[Out] I/(I-a)*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)-1/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b/(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)

$$\begin{aligned}
& 1)^{(1/2)}+a*b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2 \\
&)^{(1/2)}-(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+ \\
& a^2+1)^{(1/2))/x))+2*b^2/(a^2+1)*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a \\
& ^2+1)^{(1/2)}+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b \\
& ^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)))+1/2*b^2/(a^2+1)*((b^2*x^2+2*a*b* \\
& x+a^2+1)^{(1/2)}+a*b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2) \\
&))/(b^2)^{(1/2)}-(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2* \\
& a*b*x+a^2+1)^{(1/2))/x))-I/(I-a)^3*b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b \\
&))^{(1/2)}+I*b*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^{(1/2)}+(x-(I-a)/b)^2*b^2+2*I*b* \\
& (x-(I-a)/b))^{(1/2)))/(b^2)^{(1/2)}+I*b^2/(I-a)^3*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2) \\
&)+a*b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2) \\
&)-(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1) \\
& ^{(1/2))/x))+I*b/(I-a)^2*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)}+a*b/(a^ \\
& 2+1)*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2 \\
& +2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2 \\
& +1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2))/x))+2*b^2/(a^2+1)*(1/4*(2*b^2*x+2* \\
& a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln \\
& ((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2))}
\end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^3,x, algorithm="maxima")

[Out] integrate(sqrt((b*x + a)^2 + 1)/((I*b*x + I*a + 1)*x^3), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 452 vs. $2(135) = 270$.

time = 4.31, size = 452, normalized size = 2.25

$$\frac{(-1 + 2Ia^2 + \sqrt{(4a^2 + 2Ia - 1)I}) \sqrt{(4a^2 + 2Ia - 1)I} \log\left(\frac{(2a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} + (4a^2 + 2Ia - 1)I}{2(a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} + (4a^2 + 2Ia - 1)I}\right) - \sqrt{(4a^2 + 2Ia - 1)I} \log\left(\frac{(2a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} - (4a^2 + 2Ia - 1)I}{2(a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} - (4a^2 + 2Ia - 1)I}\right) + \sqrt{(4a^2 + 2Ia - 1)I} \log\left(\frac{(2a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} + (4a^2 + 2Ia - 1)I}{2(a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} + (4a^2 + 2Ia - 1)I}\right) - \sqrt{(4a^2 + 2Ia - 1)I} \log\left(\frac{(2a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} - (4a^2 + 2Ia - 1)I}{2(a^2 + 2Ia - 1) \sqrt{(4a^2 + 2Ia - 1)I} - (4a^2 + 2Ia - 1)I}\right)}{2(a^2 + 2Ia - 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^3,x, algorithm="fricas")

[Out] $\frac{1}{2} * ((-Ia + 2) * b^2 * x^2 + \sqrt{(4a^2 + 4Ia - 1) * b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1)}) * (a^3 - Ia^2 + a - I) * x^2 * \log(-((2a + I) * b^3 * x - \sqrt{b^2 * x^2 + 2a * b * x + a^2 + 1}) * (2a + I) * b^2 + (a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I) * \sqrt{(4a^2 + 4Ia - 1) * b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1)})) / ((2a + I) * b^2) - \sqrt{b^2 * x^2 + 2a * b * x + a^2 + 1} * (4a^2 + 4Ia - 1) * b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1) * (a^3 - Ia^2 + a - I) * x^2 * \log(-((2a + I) * b^3 * x - \sqrt{b^2 * x^2 + 2a * b * x + a^2 + 1}) * (2a + I) * b^2 - (a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I) * \sqrt{b^2 * x^2 + 2a * b * x + a^2 + 1})) / ((2a + I) * b^2)$

$a - I) \sqrt{(4a^2 + 4Ia - 1)b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1)} / ((2a + I)b^2) + \sqrt{(b^2x^2 + 2abx + a^2 + 1)} \cdot ((-Ia + 2)bx + Ia^2 + I) / ((a^3 - Ia^2 + a - I)x^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{ax^3 + bx^4 - ix^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2)/x**3,x)

[Out] -I*Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a*x**3 + b*x**4 - I*x**3), x)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 471 vs. $2(135) = 270$.

time = 0.48, size = 471, normalized size = 2.34

$$\frac{(a^2 + b^2) \log\left(\frac{(a + \sqrt{a^2 + b^2})^2 + 1 - \sqrt{a^2 + b^2}}{(a - \sqrt{a^2 + b^2})^2 + 1 - \sqrt{a^2 + b^2}}\right)}{2(a^2 + b^2) \sqrt{a^2 + b^2}} \cdot \frac{4((a - \sqrt{a^2 + b^2})^2 + 1) \sqrt{a^2 + b^2} + a((a - \sqrt{a^2 + b^2})^2 + 1) \sqrt{a^2 + b^2} + 2((a - \sqrt{a^2 + b^2})^2 + 1) \sqrt{a^2 + b^2} - 2((a - \sqrt{a^2 + b^2})^2 + 1) \sqrt{a^2 + b^2} + 2((a - \sqrt{a^2 + b^2})^2 + 1) \sqrt{a^2 + b^2} - 2((a - \sqrt{a^2 + b^2})^2 + 1) \sqrt{a^2 + b^2}}{(a^2 + b^2) \sqrt{a^2 + b^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^3,x, algorithm="giac")

[Out] $-1/2 * (2ab^2 + Ib^2) * \log(\text{abs}(2x \cdot \text{abs}(b) - 2\sqrt{(bx+a)^2+1}) - 2\sqrt{a^2+1}) / \text{abs}(2x \cdot \text{abs}(b) - 2\sqrt{(bx+a)^2+1}) + 2\sqrt{a^2+1}) / ((a^3 - Ia^2 + a - I) \sqrt{a^2+1}) - (4(Ix \cdot \text{abs}(b) - I \sqrt{(bx+a)^2+1}) * a^4 b^2 + 2I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^2 a^3 b \cdot \text{abs}(b) + 2Ia^5 b \cdot \text{abs}(b) + 2(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^3 a b^2 - 2(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1}) * a^3 b^2 + 2(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^2 a^2 b^2 + 2I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^2 a b \cdot \text{abs}(b) + 4Ia^3 b \cdot \text{abs}(b) - 2(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1}) * a b^2 + 2(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^2 b \cdot \text{abs}(b) - 4a^2 b \cdot \text{abs}(b) - (-Ix \cdot \text{abs}(b) + I \sqrt{(bx+a)^2+1}) * b^2 + 2Ia b \cdot \text{abs}(b) - 2b \cdot \text{abs}(b)) / ((a^3 - Ia^2 + a - I) * ((x \cdot \text{abs}(b) - \sqrt{(bx+a)^2+1})^2 - a^2 - 1)^2)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(a + bx)^2 + 1}}{x^3 (1 + a li + b x li)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)^(1/2)/(x^3*(a*1i + b*x*1i + 1)),x)

[Out] int(((a + b*x)^2 + 1)^(1/2)/(x^3*(a*1i + b*x*1i + 1)), x)

3.197 $\int \frac{e^{-i \operatorname{ArcTan}(a+bx)}}{x^4} dx$

Optimal. Leaf size=283

$$-\frac{\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} + \frac{(4-9ia-2a^2)b^2\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{6(1+ia)(1+a)^2}$$

[Out] $(2*a+I*(-2*a^2+1))*b^3*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)/(1-I*a-I*b*x)^{(1/2)))/(I-a)^{(7/2)/(I+a)^{(5/2)}-1/3*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(1+I*a)/x^3+1/6*(3-2*I*a)*b*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^2/(I+a)/x^2+1/6*(4-9*I*a-2*a^2)*b^2*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(1+I*a)/(a^2+1)^2/x}$

Rubi [A]

time = 0.18, antiderivative size = 282, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.375$, Rules used = {5203, 101, 156, 12, 95, 214}

$$\frac{(-2ia^2+2a+i)b^3 \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{7/2}(a+i)^{5/2}} + \frac{(-2a^2-9ia+4)b^2\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{6(1+ia)(a^2+1)^2x} - \frac{\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{6(-a+i)^2(a+i)x^2}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a + b*x])*x^4),x]

[Out] $-1/3*(\operatorname{Sqrt}[1-I*a-I*b*x]*\operatorname{Sqrt}[1+I*a+I*b*x])/((1+I*a)*x^3) + ((3-(2*I)*a)*b*\operatorname{Sqrt}[1-I*a-I*b*x]*\operatorname{Sqrt}[1+I*a+I*b*x])/((6*(I-a)^2*(I+a)*x^2) + ((4-(9*I)*a-2*a^2)*b^2*\operatorname{Sqrt}[1-I*a-I*b*x]*\operatorname{Sqrt}[1+I*a+I*b*x])/((6*(1+I*a)*(1+a^2)^2*x) + ((I+2*a-(2*I)*a^2)*b^3*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])])/((I-a)^{(7/2)}*(I+a)^{(5/2)})$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 101


```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^ (p_.), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^(p + 1
)/((m + 1)*(b*e - a*f)), x] - Dist[1/((m + 1)*(b*e - a*f)), Int[(a + b*x)^(
m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p*Simp[d*e*n + c*f*(m + p + 2) + d*f*(m
+ n + p + 2)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1
] && GtQ[n, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || Integ
ersQ[p, m + n])

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^ (p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f)),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 214

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

```

Rule 5203

```

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(a+bx)}}{x^4} dx &= \int \frac{\sqrt{1-ia-ibx}}{x^4 \sqrt{1+ia+ibx}} dx \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{\int \frac{-(3i+2a)b-2b^2x}{x^3 \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}} dx}{3(1+ia)} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} - \int \frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{x^2} dx \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} + \frac{(4-9)}{x} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} + \frac{(4-9)}{x} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} + \frac{(4-9)}{x} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} + \frac{(4-9)}{x} \\
&= -\frac{\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{3(1+ia)x^3} + \frac{(3-2ia)b\sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{6(i-a)^2(i+a)x^2} + \frac{(4-9)}{x}
\end{aligned}$$

Mathematica [A]

time = 0.23, size = 247, normalized size = 0.87

$$\frac{2(1+ia)(i+a)(i+ax)\sqrt{1+a^2+2abx+b^2x^2} + (1-4ia)bx(i+ax)\sqrt{1+a^2+2abx+b^2x^2} + \frac{3(-1+2ia+2a^2)^{3/2} \left(\sqrt{-1-ia} \sqrt{-1+ia} \sqrt{1+a^2+2abx+b^2x^2} + 2ibx \tanh^{-1} \left(\frac{\sqrt{-1-ia} \sqrt{-i(i+ax+bx)}}{\sqrt{-1+ia} \sqrt{1+ia+ibx}} \right) \right)}{6(1+a^2)^2 x^3}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^(I*ArcTan[a + b*x]))*x^4),x]

[Out] (2*(1 + I*a)*(I + a)*(I + a + b*x)*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2] + (1 - (4*I)*a)*b*x*(I + a + b*x)*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2] + (3*(-1 + (2*I)*a + 2*a^2)*b^2*x^2*(Sqrt[-1 - I*a]*Sqrt[-1 + I*a]*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2] + (2*I)*b*x*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x])]))/((-1 - I*a)^(3/2)*Sqrt[-1 + I*a])/(6*(1 + a^2)^2*x^3)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1510 vs. 2(226) = 452.

time = 0.14, size = 1511, normalized size = 5.34

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^4,x, algorithm="maxima")

[Out] integrate(sqrt((b*x + a)^2 + 1)/((I*b*x + I*a + 1)*x^4), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 690 vs. $2(194) = 388$.

time = 3.91, size = 690, normalized size = 2.44

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^4,x, algorithm="fricas")

[Out]
$$\frac{1}{6} \left((2Ia^2 - 9a - 4I)b^3x^3 - 3\sqrt{(4a^4 + 8Ia^3 - 8a^2 - 4Ia + 1)b^6/(a^{12} - 2Ia^{11} + 4a^{10} - 10Ia^9 + 5a^8 - 20Ia^7 - 20Ia^5 - 5a^4 - 10Ia^3 - 4a^2 - 2Ia - 1)}(a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I)x^3 \log(-((2a^2 + 2Ia - 1)b^4x - \sqrt{b^2x^2 + 2a*b*x + a^2 + 1})(2a^2 + 2Ia - 1)b^3 + (a^7 - Ia^6 + 3a^5 - 3Ia^4 + 3a^3 - 3Ia^2 + a - I)\sqrt{(4a^4 + 8Ia^3 - 8a^2 - 4Ia + 1)b^6/(a^{12} - 2Ia^{11} + 4a^{10} - 10Ia^9 + 5a^8 - 20Ia^7 - 20Ia^5 - 5a^4 - 10Ia^3 - 4a^2 - 2Ia - 1)})) / ((2a^2 + 2Ia - 1)b^3) \right) + 3\sqrt{(4a^4 + 8Ia^3 - 8a^2 - 4Ia + 1)b^6/(a^{12} - 2Ia^{11} + 4a^{10} - 10Ia^9 + 5a^8 - 20Ia^7 - 20Ia^5 - 5a^4 - 10Ia^3 - 4a^2 - 2Ia - 1)}(a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I)x^3 \log(-((2a^2 + 2Ia - 1)b^4x - \sqrt{b^2x^2 + 2a*b*x + a^2 + 1})(2a^2 + 2Ia - 1)b^3 - (a^7 - Ia^6 + 3a^5 - 3Ia^4 + 3a^3 - 3Ia^2 + a - I)\sqrt{(4a^4 + 8Ia^3 - 8a^2 - 4Ia + 1)b^6/(a^{12} - 2Ia^{11} + 4a^{10} - 10Ia^9 + 5a^8 - 20Ia^7 - 20Ia^5 - 5a^4 - 10Ia^3 - 4a^2 - 2Ia - 1)})) / ((2a^2 + 2Ia - 1)b^3) \right) + ((2Ia^2 - 9a - 4I)b^2x^2 + 2Ia^4 + (-2Ia^3 + 3a^2 - 2Ia + 3)b*x + 4Ia^2 + 2I)\sqrt{b^2x^2 + 2a*b*x + a^2 + 1} / ((a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I)x^3)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{ax^4 + bx^5 - ix^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))*(1+(b*x+a)**2)**(1/2)/x**4,x)

[Out] $-I \cdot \text{Integral}(\sqrt{a^2 + 2abx + b^2x^2 + 1}/(ax^4 + b^5x - Ix^4), x)$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 884 vs. $2(194) = 388$.
time = 0.49, size = 884, normalized size = 3.12

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))*(1+(b*x+a)^2)^(1/2)/x^4,x, algorithm="giac")`

[Out] $\frac{1}{2} \cdot (2a^2b^3 + 2Iab^3 - b^3) \cdot \log(\text{abs}(2x \cdot \text{abs}(b) - 2\sqrt{(bx+a)^2 + 1}) - 2\sqrt{a^2 + 1}) / \text{abs}(2x \cdot \text{abs}(b) - 2\sqrt{(bx+a)^2 + 1}) + 2\sqrt{a^2 + 1}) / ((a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I) \cdot \sqrt{a^2 + 1}) + \frac{1}{3} \cdot (-8I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^3 a^5 b^3 + 24(-Ix \cdot \text{abs}(b) + I \sqrt{(bx+a)^2 + 1}) a^7 b^3 - 24I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 a^6 b^2 \cdot \text{abs}(b) - 8Ia^8 b^2 \cdot \text{abs}(b) + 6(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^5 a^2 b^3 - 24(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^3 a^4 b^3 + 18(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1}) a^6 b^3 - 12(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 a^5 b^2 \cdot \text{abs}(b) + 12a^7 b^2 \cdot \text{abs}(b) + 6I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^5 a b^3 - 32I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^3 a^3 b^3 + 54(-Ix \cdot \text{abs}(b) + I \sqrt{(bx+a)^2 + 1}) a^5 b^3 - 60I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 a^4 b^2 \cdot \text{abs}(b) - 20Ia^6 b^2 \cdot \text{abs}(b) - 3(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^5 b^3 - 24(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^3 a^2 b^3 + 39(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1}) a^4 b^3 - 24(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 a^3 b^2 \cdot \text{abs}(b) + 36a^5 b^2 \cdot \text{abs}(b) - 24I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^3 a b^3 + 36(-Ix \cdot \text{abs}(b) + I \sqrt{(bx+a)^2 + 1}) a^3 b^3 - 48I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 a^2 b^2 \cdot \text{abs}(b) - 12Ia^4 b^2 \cdot \text{abs}(b) + 24(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1}) a^2 b^3 - 12(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 a b^2 \cdot \text{abs}(b) + 36a^3 b^2 \cdot \text{abs}(b) + 6(-Ix \cdot \text{abs}(b) + I \sqrt{(bx+a)^2 + 1}) a b^3 - 12I(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 b^2 \cdot \text{abs}(b) + 4Ia^2 b^2 \cdot \text{abs}(b) + 3(x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1}) b^3 + 12a b^2 \cdot \text{abs}(b) + 4Ib^2 \cdot \text{abs}(b)) / ((a^5 - Ia^4 + 2a^3 - 2Ia^2 + a - I) \cdot ((x \cdot \text{abs}(b) - \sqrt{(bx+a)^2 + 1})^2 - a^2 - 1)^3)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{(a+bx)^2 + 1}}{x^4 (1 + a \operatorname{li} + b x \operatorname{li})} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((a + b*x)^2 + 1)^(1/2)/(x^4*(a*1i + b*x*1i + 1)),x)`

[Out] `int(((a + b*x)^2 + 1)^(1/2)/(x^4*(a*1i + b*x*1i + 1)), x)`

3.198 $\int e^{-2i\text{ArcTan}(a+bx)} x^4 dx$

Optimal. Leaf size=99

$$-\frac{2(1+ia)^3x}{b^4} - \frac{i(i-a)^2x^2}{b^3} + \frac{2(1+ia)x^3}{3b^2} - \frac{ix^4}{2b} - \frac{x^5}{5} - \frac{2i(i-a)^4 \log(i-a-bx)}{b^5}$$

[Out] $-2*(1+I*a)^3*x/b^4 - I*(I-a)^2*x^2/b^3 + 2/3*(1+I*a)*x^3/b^2 - 1/2*I*x^4/b - 1/5*x^5 - 2*I*(I-a)^4*\ln(I-a-b*x)/b^5$

Rubi [A]

time = 0.07, antiderivative size = 99, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2i(-a+i)^4 \log(-a-bx+i)}{b^5} - \frac{2(1+ia)^3x}{b^4} - \frac{i(-a+i)^2x^2}{b^3} + \frac{2(1+ia)x^3}{3b^2} - \frac{ix^4}{2b} - \frac{x^5}{5}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^4/E^{((2*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $(-2*(1 + I*a)^3*x)/b^4 - (I*(I - a)^2*x^2)/b^3 + (2*(1 + I*a)*x^3)/(3*b^2) - ((I/2)*x^4)/b - x^5/5 - ((2*I)*(I - a)^4*\text{Log}[I - a - b*x])/b^5$

Rule 78

$\text{Int}[(a_.) + (b_.)*(x_.)]*((c_.) + (d_.)*(x_.)^{(n_.)})*((e_.) + (f_.)*(x_.)^{(p_.)}), x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

$\text{Int}[E^{(\text{ArcTan}[(c_.)*((a_.) + (b_.)*(x_.)])*(n_.)})*((d_.) + (e_.)*(x_.)^{(m_.)}), x_Symbol] \rightarrow \text{Int}[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^{(I*(n/2)})/(1 + I*a*c + I*b*c*x)^{(I*(n/2)})), x] /;$ FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(a+bx)} x^4 dx &= \int \frac{x^4(1-ia-ibx)}{1+ia+ibx} dx \\ &= \int \left(\frac{2(-1-ia)^3}{b^4} - \frac{2i(-i+a)^2x}{b^3} + \frac{2(1+ia)x^2}{b^2} - \frac{2ix^3}{b} - x^4 - \frac{2i(-i+a)^4}{b^4(-i+a+bx)} \right) dx \\ &= -\frac{2(1+ia)^3x}{b^4} - \frac{i(i-a)^2x^2}{b^3} + \frac{2(1+ia)x^3}{3b^2} - \frac{ix^4}{2b} - \frac{x^5}{5} - \frac{2i(i-a)^4 \log(i-a-bx)}{b^5} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 95, normalized size = 0.96

$$-\frac{2(1+ia)^3x}{b^4} - \frac{i(-i+a)^2x^2}{b^3} + \frac{2(1+ia)x^3}{3b^2} - \frac{ix^4}{2b} - \frac{x^5}{5} - \frac{2i(-i+a)^4 \log(i-a-bx)}{b^5}$$

Antiderivative was successfully verified.

[In] Integrate[x^4/E^((2*I)*ArcTan[a + b*x]),x]

[Out] $(-2*(1 + I*a)^3*x)/b^4 - (I*(-I + a)^2*x^2)/b^3 + (2*(1 + I*a)*x^3)/(3*b^2) - ((I/2)*x^4)/b - x^5/5 - ((2*I)*(-I + a)^4*Log[I - a - b*x])/b^5$

Maple [A]

time = 0.14, size = 125, normalized size = 1.26

method	result
default	$-\frac{i(-\frac{1}{5}ib^4x^5 + \frac{1}{2}b^3x^4 + \frac{2}{3}ib^2x^3 - \frac{2}{3}ab^2x^3 - 2iabx^2 + a^2bx^2 + 6ia^2x - 2a^3x - x^2b - 2ix + 6ax)}{b^4} + \frac{(-2ia^4 - 8a^3 + 12ia^2 + 8a - 2i) \ln(-bx - a)}{b^5}$
risch	$-\frac{x^5}{5} + \frac{2iax^3}{3b^2} + \frac{2x^3}{3b^2} - \frac{6iax}{b^4} - \frac{2ax^2}{b^3} - \frac{ia^2x^2}{b^3} + \frac{6a^2x}{b^4} + \frac{2ia^3x}{b^4} + \frac{ix^2}{b^3} - \frac{2x}{b^4} - \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)}{b^5} - \frac{4 \ln(b^2x^2 + 2abx + a^2 + 1)}{b^5}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^4/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x,method=_RETURNVERBOSE)

[Out] $-I/b^4*(-1/5*I*b^4*x^5 + 1/2*b^3*x^4 + 2/3*I*b^2*x^3 - 2/3*a*b^2*x^3 - 2*I*a*b*x^2 + a^2*b*x^2 + 6*I*a^2*x - 2*a^3*x - x^2*b - 2*I*x + 6*a*x) + (-2*I*a^4 + 12*I*a^2 - 8*a^3 - 2*I*a^2 + 8*a)/b^5*\ln(I-a-b*x)$

Maxima [A]

time = 0.26, size = 105, normalized size = 1.06

$$\frac{6b^4x^5 + 15ib^3x^4 - 20(ia+1)b^2x^3 - 30(-ia^2 - 2a+i)bx^2 - 60(ia^3 + 3a^2 - 3ia - 1)x}{30b^4} - \frac{2(ia^4 + 4a^3 - 6ia^2 - 4a + i) \log(ibx + ia + 1)}{b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="maxima")

[Out] $-1/30*(6*b^4*x^5 + 15*I*b^3*x^4 - 20*(I*a + 1)*b^2*x^3 - 30*(-I*a^2 - 2*a + I)*b*x^2 - 60*(I*a^3 + 3*a^2 - 3*I*a - 1)*x)/b^4 - 2*(I*a^4 + 4*a^3 - 6*I*a^2 - 4*a + I)*\log(I*b*x + I*a + 1)/b^5$

Fricas [A]

time = 3.21, size = 105, normalized size = 1.06

$$\frac{6b^5x^5 + 15ib^4x^4 + 20(-ia-1)b^3x^3 + 30(ia^2 + 2a-i)b^2x^2 + 60(-ia^3 - 3a^2 + 3ia + 1)bx + 60(ia^4 + 4a^3 - 6ia^2 - 4a + i) \log(\frac{bx+a-i}{b})}{30b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="fricas")

[Out] $-1/30*(6*b^5*x^5 + 15*I*b^4*x^4 + 20*(-I*a - 1)*b^3*x^3 + 30*(I*a^2 + 2*a - I)*b^2*x^2 + 60*(-I*a^3 - 3*a^2 + 3*I*a + 1)*b*x + 60*(I*a^4 + 4*a^3 - 6*I*a^2 - 4*a + I)*\log((b*x + a - I)/b))/b^5$

Sympy [A]

time = 0.25, size = 114, normalized size = 1.15

$$-\frac{x^5}{5} - x^3 \left(-\frac{2ia}{3b^2} - \frac{2}{3b^2} \right) - x^2 \left(\frac{ia^2}{b^3} + \frac{2a}{b^3} - \frac{i}{b^3} \right) - x \left(-\frac{2ia^3}{b^4} - \frac{6a^2}{b^4} + \frac{6ia}{b^4} + \frac{2}{b^4} \right) - \frac{ix^4}{2b} - \frac{2i(a-i)^4 \log(a+bx-i)}{b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**4/(1+I*(b*x+a)**2*(1+(b*x+a)**2),x)

[Out] $-x^{**5}/5 - x^{**3}*(-2*I*a/(3*b^{**2}) - 2/(3*b^{**2})) - x^{**2}*(I*a^{**2}/b^{**3} + 2*a/b^{**3} - I/b^{**3}) - x*(-2*I*a^{**3}/b^{**4} - 6*a^{**2}/b^{**4} + 6*I*a/b^{**4} + 2/b^{**4}) - I*x^{**4}/(2*b) - 2*I*(a - I)^{**4}*\log(a + b*x - I)/b^{**5}$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 215 vs. 2(73) = 146.

time = 0.45, size = 215, normalized size = 2.17

$$\frac{i(bx+ia+1)^5 \left(-\frac{15i(2ab-3i)b}{(ibz+ia+1)b} - \frac{20(3a^2b^2-10iab^2-7b^2)}{(ibz+ia+1)^2b^2} + \frac{60i(a^3b^3-6ia^2b^2-9ab^2+4ib^2)}{(ibz+ia+1)^3b^3} + \frac{30(a^4b^4-12ia^3b^3-30a^2b^4+28iab^4+9b^4)}{(ibz+ia+1)^4b^4} + 6 \right) + 2(-ia^4-4a^3+6ia^2+4a-i)\log\left(\frac{1}{\sqrt{(bx+a)^2+1|b|}}\right)}{30b^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="giac")

[Out] $1/30*I*(I*b*x + I*a + 1)^5*(-15*I*(2*a*b - 3*I*b)/((I*b*x + I*a + 1)*b) - 20*(3*a^2*b^2 - 10*I*a*b^2 - 7*b^2)/((I*b*x + I*a + 1)^2*b^2) + 60*I*(a^3*b^3 - 6*I*a^2*b^3 - 9*a*b^3 + 4*I*b^3)/((I*b*x + I*a + 1)^3*b^3) + 30*(a^4*b^4 - 12*I*a^3*b^4 - 30*a^2*b^4 + 28*I*a*b^4 + 9*b^4)/((I*b*x + I*a + 1)^4*b^4) + 6)/b^5 - 2*(-I*a^4 - 4*a^3 + 6*I*a^2 + 4*a - I)*\log(1/(\sqrt{(b*x + a)^2 + 1}*abs(b)))/b^5$

Mupad [B]

time = 0.17, size = 165, normalized size = 1.67

$$\ln\left(x + \frac{a-i}{b}\right) \left(\frac{8a-8a^3}{b^5} - \frac{(2a^4-12a^2+2)1i}{b^5} \right) + x^4 \left(\frac{a-i}{4b} - \frac{a+1i}{4b} \right) - \frac{x^5}{5} + \frac{x^2 \left(\frac{a-i}{b} - \frac{a+1i}{b} \right) (a-i)^2}{2b^2} - \frac{x^3 \left(\frac{a-i}{b} - \frac{a+1i}{b} \right) (a-i)}{3b} - \frac{x \left(\frac{a-i}{b} - \frac{a+1i}{b} \right) (a-i)^3}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*((a + b*x)^2 + 1))/(a*1i + b*x*1i + 1)^2,x)

[Out] $\log(x + (a - 1i)/b)*((8*a - 8*a^3)/b^5 - ((2*a^4 - 12*a^2 + 2)*1i)/b^5) + x^4*((a - 1i)/(4*b) - (a + 1i)/(4*b)) - x^5/5 + (x^2*((a - 1i)/b - (a + 1i)/b)*(a - 1i)^2)/(2*b^2) - (x^3*((a - 1i)/b - (a + 1i)/b)*(a - 1i))/(3*b) - (x*((a - 1i)/b - (a + 1i)/b)*(a - 1i)^3)/b^3$

3.199 $\int e^{-2i\text{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=77

$$-\frac{2i(i-a)^2x}{b^3} + \frac{(1+ia)x^2}{b^2} - \frac{2ix^3}{3b} - \frac{x^4}{4} - \frac{2(1+ia)^3 \log(i-a-bx)}{b^4}$$

[Out] $-2*I*(I-a)^2*x/b^3+(1+I*a)*x^2/b^2-2/3*I*x^3/b-1/4*x^4-2*(1+I*a)^3*\ln(I-a-b*x)/b^4$

Rubi [A]

time = 0.05, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2(1+ia)^3 \log(-a-bx+i)}{b^4} - \frac{2i(-a+i)^2x}{b^3} + \frac{(1+ia)x^2}{b^2} - \frac{2ix^3}{3b} - \frac{x^4}{4}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{((2*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $((-2*I)*(I - a)^2*x)/b^3 + ((1 + I*a)*x^2)/b^2 - (((2*I)/3)*x^3)/b - x^4/4 - (2*(1 + I*a)^3*\text{Log}[I - a - b*x])/b^4$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)}*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& ((\text{ILtQ}[n, 0] \&\& \text{ILtQ}[p, 0]) \|\ \text{EqQ}[p, 1] \|\ (\text{IGtQ}[p, 0] \&\& (!\text{IntegerQ}[n] \|\ \text{LeQ}[9*p + 5*(n + 2), 0] \|\ \text{GeQ}[n + p + 1, 0] \|\ (\text{GeQ}[n + p + 2, 0] \&\& \text{RationalQ}[a, b, c, d, e, f])))$

Rule 5203

$\text{Int}[E^{(\text{ArcTan}[(c_.)*((a_. + (b_.)*(x_.))])*(n_.)}*((d_. + (e_.)*(x_.))^{(m_.)}, x_Symbol] :> \text{Int}[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^{(I*(n/2))}/(1 + I*a*c + I*b*c*x)^{(I*(n/2))}), x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(a+bx)} x^3 dx &= \int \frac{x^3(1-ia-ibx)}{1+ia+ibx} dx \\ &= \int \left(-\frac{2i(-i+a)^2}{b^3} + \frac{2(1+ia)x}{b^2} - \frac{2ix^2}{b} - x^3 + \frac{2(-1-ia)^3}{b^3(-i+a+bx)} \right) dx \\ &= -\frac{2i(i-a)^2x}{b^3} + \frac{(1+ia)x^2}{b^2} - \frac{2ix^3}{3b} - \frac{x^4}{4} - \frac{2(1+ia)^3 \log(i-a-bx)}{b^4} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 77, normalized size = 1.00

$$-\frac{2i(i-a)^2x}{b^3} + \frac{(1+ia)x^2}{b^2} - \frac{2ix^3}{3b} - \frac{x^4}{4} - \frac{2(1+ia)^3 \log(i-a-bx)}{b^4}$$

Antiderivative was successfully verified.

`[In] Integrate[x^3/E^((2*I)*ArcTan[a + b*x]),x]`

```
[Out] ((-2*I)*(I - a)^2*x)/b^3 + ((1 + I*a)*x^2)/b^2 - (((2*I)/3)*x^3)/b - x^4/4 - (2*(1 + I*a)^3*Log[I - a - b*x])/b^4
```

Maple [A]

time = 0.13, size = 85, normalized size = 1.10

method	result
default	$\frac{i(\frac{1}{4}ib^3x^4 - \frac{2}{3}b^2x^3 - ibx^2 + abx^2 + 4iax - 2a^2x + 2x)}{b^3} + \frac{(2ia^3 + 6a^2 - 6ia - 2)\ln(-bx - a + i)}{b^4}$
risch	$-\frac{x^4}{4} - \frac{2ix^3}{3b} + \frac{x^2}{b^2} + \frac{iax^2}{b^2} - \frac{4ax}{b^3} - \frac{2ia^2x}{b^3} + \frac{2ix}{b^3} + \frac{3\ln(b^2x^2 + 2abx + a^2 + 1)a^2}{b^4} + \frac{i\ln(b^2x^2 + 2abx + a^2 + 1)a^3}{b^4} - \frac{\ln(b^2x^2 + 1)}{b^4}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^3/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x,method=_RETURNVERBOSE)`

```
[Out] I/b^3*(1/4*I*b^3*x^4 - 2/3*b^2*x^3 - I*b*x^2 + a*b*x^2 + 4*I*a*x - 2*a^2*x + 2*x) + (2*I*a^3 - 6*I*a + 6*a^2 - 2)/b^4*ln(I - a - b*x)
```

Maxima [A]

time = 0.25, size = 73, normalized size = 0.95

$$\frac{i(-3ib^3x^4 + 8b^2x^3 - 12(a-i)bx^2 + 24(a^2 - 2ia - 1)x)}{12b^3} - \frac{2(-ia^3 - 3a^2 + 3ia + 1)\log(ibx + ia + 1)}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="maxima")`

```
[Out] -1/12*I*(-3*I*b^3*x^4 + 8*b^2*x^3 - 12*(a - I)*b*x^2 + 24*(a^2 - 2*I*a - 1)*x)/b^3 - 2*(-I*a^3 - 3*a^2 + 3*I*a + 1)*log(I*b*x + I*a + 1)/b^4
```

Fricas [A]

time = 2.90, size = 77, normalized size = 1.00

$$\frac{3b^4x^4 + 8ib^3x^3 + 12(-ia - 1)b^2x^2 + 24(ia^2 + 2a - i)bx + 24(-ia^3 - 3a^2 + 3ia + 1)\log\left(\frac{bx+a-i}{b}\right)}{12b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x^3/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="fricas")`

[Out] $-1/12*(3*b^4*x^4 + 8*I*b^3*x^3 + 12*(-I*a - 1)*b^2*x^2 + 24*(I*a^2 + 2*a - I)*b*x + 24*(-I*a^3 - 3*a^2 + 3*I*a + 1)*\log((b*x + a - I)/b))/b^4$

Sympy [A]

time = 0.20, size = 76, normalized size = 0.99

$$-\frac{x^4}{4} - x^2 \left(-\frac{ia}{b^2} - \frac{1}{b^2} \right) - x \left(\frac{2ia^2}{b^3} + \frac{4a}{b^3} - \frac{2i}{b^3} \right) - \frac{2ix^3}{3b} + \frac{2i(a-i)^3 \log(a+bx-i)}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**3/(1+I*(b*x+a))**2*(1+(b*x+a)**2), x)`

[Out] $-x^{**4}/4 - x^{**2}*(-I*a/b^{**2} - 1/b^{**2}) - x*(2*I*a^{**2}/b^{**3} + 4*a/b^{**3} - 2*I/b^{**3}) - 2*I*x^{**3}/(3*b) + 2*I*(a - I)^{**3}*\log(a + b*x - I)/b^{**4}$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 158 vs. 2(59) = 118.

time = 0.43, size = 158, normalized size = 2.05

$$\frac{(ibx+ia+1)^4 \left(-\frac{4i(3ab-5ib)}{(ibx+ia+1)b} - \frac{18(a^2b^2-4iab^2-3b^2)}{(ibx+ia+1)^2b^2} + \frac{12i(a^3b^3-9ia^2b^3-15ab^3+7ib^3)}{(ibx+ia+1)^3b^3} + 3 \right)}{12b^4} - \frac{2(i a^3 + 3 a^2 - 3i a - 1) \log \left(\frac{1}{\sqrt{(bx+a)^2 + 1} |b|} \right)}{b^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^3/(1+I*(b*x+a))^2*(1+(b*x+a)^2), x, algorithm="giac")`

[Out] $-1/12*(I*b*x + I*a + 1)^4*(-4*I*(3*a*b - 5*I*b)/((I*b*x + I*a + 1)*b) - 18*(a^2*b^2 - 4*I*a*b^2 - 3*b^2)/((I*b*x + I*a + 1)^2*b^2) + 12*I*(a^3*b^3 - 9*I*a^2*b^3 - 15*a*b^3 + 7*I*b^3)/((I*b*x + I*a + 1)^3*b^3) + 3)/b^4 - 2*(I*a^3 + 3*a^2 - 3*I*a - 1)*\log(1/(\sqrt{(b*x + a)^2 + 1}*\text{abs}(b)))/b^4$

Mupad [B]

time = 0.54, size = 129, normalized size = 1.68

$$x^3 \left(\frac{a-i}{3b} - \frac{a+1i}{3b} \right) - \frac{x^4}{4} - \ln \left(x + \frac{a-i}{b} \right) \left(-\frac{6a^2-2}{b^4} + \frac{(6a-2a^3)1i}{b^4} \right) - \frac{x^2 \left(\frac{a-i}{b} - \frac{a+1i}{b} \right) (a-i)}{2b} + \frac{x \left(\frac{a-i}{b} - \frac{a+1i}{b} \right) (a-i)^2}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^3*((a+b*x)^2+1))/(a*1i+b*x*1i+1)^2,x)`

[Out] $x^3*((a-1i)/(3*b) - (a+1i)/(3*b)) - x^4/4 - \log(x + (a-1i)/b)*(((6*a - 2*a^3)*1i)/b^4 - (6*a^2 - 2)/b^4) - (x^2*((a-1i)/b - (a+1i)/b)*(a-1i))/b^2 + (x*((a-1i)/b - (a+1i)/b)*(a-1i)^2)/b^2$

3.200 $\int e^{-2i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=59

$$\frac{2(1+ia)x}{b^2} - \frac{ix^2}{b} - \frac{x^3}{3} - \frac{2i(i-a)^2 \log(i-a-bx)}{b^3}$$

[Out] $2*(1+I*a)*x/b^2 - I*x^2/b - 1/3*x^3 - 2*I*(I-a)^2*\ln(I-a-b*x)/b^3$

Rubi [A]

time = 0.04, antiderivative size = 59, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2i(-a+i)^2 \log(-a-bx+i)}{b^3} + \frac{2(1+ia)x}{b^2} - \frac{ix^2}{b} - \frac{x^3}{3}$$

Antiderivative was successfully verified.

[In] Int[x^2/E^((2*I)*ArcTan[a + b*x]),x]

[Out] $(2*(1 + I*a)*x)/b^2 - (I*x^2)/b - x^3/3 - ((2*I)*(I - a)^2*\text{Log}[I - a - b*x])/b^3$

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2(1-ia-ibx)}{1+ia+ibx} dx \\ &= \int \left(\frac{2i(-i+a)}{b^2} - \frac{2ix}{b} - x^2 - \frac{2i(-i+a)^2}{b^2(-i+a+bx)} \right) dx \\ &= \frac{2(1+ia)x}{b^2} - \frac{ix^2}{b} - \frac{x^3}{3} - \frac{2i(i-a)^2 \log(i-a-bx)}{b^3} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 55, normalized size = 0.93

$$\frac{bx(6 + 6ia - 3ibx - b^2x^2) - 6i(-i + a)^2 \log(i - a - bx)}{3b^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^((2*I)*ArcTan[a + b*x]), x]

[Out] (b*x*(6 + (6*I)*a - (3*I)*b*x - b^2*x^2) - (6*I)*(-I + a)^2*Log[I - a - b*x])/ (3*b^3)

Maple [A]

time = 0.09, size = 59, normalized size = 1.00

method	result
default	$\frac{i(\frac{1}{3}ib^2x^3 - x^2b - 2ix + 2ax)}{b^2} + \frac{(-2ia^2 - 4a + 2i) \ln(-bx - a + i)}{b^3}$
risch	$-\frac{x^3}{3} - \frac{ix^2}{b} + \frac{2x}{b^2} + \frac{2iax}{b^2} - \frac{2 \ln(b^2x^2 + 2abx + a^2 + 1)a}{b^3} - \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)a^2}{b^3} + \frac{i \ln(b^2x^2 + 2abx + a^2 + 1)}{b^3} - \frac{4i \arctan(\frac{bx+a-i}{b})}{b^3}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*(b*x+a))^2*(1+(b*x+a)^2), x, method=_RETURNVERBOSE)

[Out] I/b^2*(1/3*I*b^2*x^3 - x^2*b - 2*I*x + 2*a*x) + (-2*I*a^2 + 2*I - 4*a)/b^3*ln(I - a - b*x)

Maxima [A]

time = 0.29, size = 53, normalized size = 0.90

$$\frac{b^2x^3 + 3ibx^2 + 6(-ia - 1)x}{3b^2} - \frac{2(i a^2 + 2a - i) \log(ibx + ia + 1)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*(b*x+a))^2*(1+(b*x+a)^2), x, algorithm="maxima")

[Out] -1/3*(b^2*x^3 + 3*I*b*x^2 + 6*(-I*a - 1)*x)/b^2 - 2*(I*a^2 + 2*a - I)*log(I*b*x + I*a + 1)/b^3

Fricas [A]

time = 3.32, size = 53, normalized size = 0.90

$$\frac{b^3x^3 + 3ib^2x^2 + 6(-ia - 1)bx + 6(i a^2 + 2a - i) \log\left(\frac{bx+a-i}{b}\right)}{3b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*(b*x+a))^2*(1+(b*x+a)^2), x, algorithm="fricas")

[Out] $-1/3*(b^3*x^3 + 3*I*b^2*x^2 + 6*(-I*a - 1)*b*x + 6*(I*a^2 + 2*a - I)*\log((b*x + a - I)/b))/b^3$

Sympy [A]

time = 0.16, size = 49, normalized size = 0.83

$$-\frac{x^3}{3} - x \left(-\frac{2ia}{b^2} - \frac{2}{b^2} \right) - \frac{ix^2}{b} - \frac{2i(a-i)^2 \log(a+bx-i)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2/(1+I*(b*x+a))**2*(1+(b*x+a)**2),x)`

[Out] $-x**3/3 - x*(-2*I*a/b**2 - 2/b**2) - I*x**2/b - 2*I*(a - I)**2*\log(a + b*x - I)/b**3$

Giac [B] Both result and optimal contain **B** complex but leaf count of result is larger than twice the leaf count of optimal. 109 vs. $2(45) = 90$.

time = 0.44, size = 109, normalized size = 1.85

$$\frac{i(ibx+ia+1)^3 \left(-\frac{3i(ab-2ib)}{(ibx+ia+1)b} - \frac{3(a^2b^2-6iab^2-5b^2)}{(ibx+ia+1)^2b^2} + 1 \right)}{3b^3} - \frac{2(-ia^2-2a+i) \log \left(\frac{1}{\sqrt{(bx+a)^2+1|b|}} \right)}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="giac")`

[Out] $-1/3*I*(I*b*x + I*a + 1)^3*(-3*I*(a*b - 2*I*b)/((I*b*x + I*a + 1)*b) - 3*(a^2*b^2 - 6*I*a*b^2 - 5*b^2)/((I*b*x + I*a + 1)^2*b^2) + 1)/b^3 - 2*(-I*a^2 - 2*a + I)*\log(1/(\sqrt{(b*x + a)^2 + 1}*abs(b)))/b^3$

Mupad [B]

time = 0.54, size = 90, normalized size = 1.53

$$-\ln \left(x + \frac{a-i}{b} \right) \left(\frac{4a}{b^3} + \frac{(2a^2-2)1i}{b^3} \right) + x^2 \left(\frac{a-i}{2b} - \frac{a+1i}{2b} \right) - \frac{x^3}{3} - \frac{x \left(\frac{a-i}{b} - \frac{a+1i}{b} \right) (a-i)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2*((a+b*x)^2+1))/(a*1i+b*x*1i+1)^2,x)`

[Out] $x^2*((a-1i)/(2*b) - (a+1i)/(2*b)) - \log(x + (a-1i)/b)*((4*a)/b^3 + ((2*a^2-2)*1i)/b^3) - x^3/3 - (x*((a-1i)/b - (a+1i)/b)*(a-1i))/b$

3.201 $\int e^{-2i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=40

$$-\frac{2ix}{b} - \frac{x^2}{2} + \frac{2(1+ia)\log(i-a-bx)}{b^2}$$

[Out] $-2*I*x/b-1/2*x^2+2*(1+I*a)*\ln(I-a-b*x)/b^2$

Rubi [A]

time = 0.03, antiderivative size = 40, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5203, 78}

$$\frac{2(1+ia)\log(-a-bx+i)}{b^2} - \frac{2ix}{b} - \frac{x^2}{2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x/E^{((2*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $((-2*I)*x)/b - x^2/2 + (2*(1 + I*a)*\text{Log}[I - a - b*x])/b^2$

Rule 78

$\text{Int}[(a_.) + (b_.)*(x_)]*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

$\text{Int}[E^{\text{ArcTan}[(c_.)*((a_.) + (b_.)*(x_))]}*(n_.)*((d_.) + (e_.)*(x_))^{(m_.)}, x_Symbol] \rightarrow \text{Int}[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^{(I*(n/2)})/(1 + I*a*c + I*b*c*x)^{(I*(n/2)}), x] /;$ FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(a+bx)} x dx &= \int \frac{x(1-ia-ibx)}{1+ia+ibx} dx \\ &= \int \left(-\frac{2i}{b} - x + \frac{2(1+ia)}{b(-i+a+bx)} \right) dx \\ &= -\frac{2ix}{b} - \frac{x^2}{2} + \frac{2(1+ia)\log(i-a-bx)}{b^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 40, normalized size = 1.00

$$-\frac{2ix}{b} - \frac{x^2}{2} + \frac{2(1+ia)\log(i-a-bx)}{b^2}$$

Antiderivative was successfully verified.

`[In] Integrate[x/E^((2*I)*ArcTan[a + b*x]),x]``[Out] ((-2*I)*x)/b - x^2/2 + (2*(1 + I*a)*Log[I - a - b*x])/b^2`**Maple [A]**

time = 0.08, size = 39, normalized size = 0.98

method	result	size
default	$-\frac{\frac{1}{2}x^2b+2ix}{b} + \frac{(2ia+2)\ln(-bx-a+i)}{b^2}$	39
risch	$-\frac{x^2}{2} - \frac{2ix}{b} + \frac{\ln(b^2x^2+2abx+a^2+1)}{b^2} + \frac{2i\arctan(bx+a)}{b^2} + \frac{ia\ln(b^2x^2+2abx+a^2+1)}{b^2} - \frac{2a\arctan(bx+a)}{b^2}$	85

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x,method=_RETURNVERBOSE)``[Out] -1/b*(1/2*x^2*b+2*I*x)+(2*I*a+2)/b^2*ln(I-a-b*x)`**Maxima [A]**

time = 0.26, size = 36, normalized size = 0.90

$$\frac{i(ibx^2 - 4x)}{2b} - \frac{2(-ia - 1)\log(ibx + ia + 1)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="maxima")``[Out] 1/2*I*(I*b*x^2 - 4*x)/b - 2*(-I*a - 1)*log(I*b*x + I*a + 1)/b^2`**Fricas [A]**

time = 4.04, size = 35, normalized size = 0.88

$$-\frac{b^2x^2 + 4ibx + 4(-ia - 1)\log\left(\frac{bx+a-i}{b}\right)}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="fricas")``[Out] -1/2*(b^2*x^2 + 4*I*b*x + 4*(-I*a - 1)*log((b*x + a - I)/b))/b^2`

Sympy [A]

time = 0.12, size = 29, normalized size = 0.72

$$-\frac{x^2}{2} - \frac{2ix}{b} + \frac{2i(a-i)\log(a+bx-i)}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))**2*(1+(b*x+a)**2),x)**[Out]** -x**2/2 - 2*I*x/b + 2*I*(a - I)*log(a + b*x - I)/b**2**Giac [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 72 vs. 2(32) = 64.

time = 0.42, size = 72, normalized size = 1.80

$$\frac{i \left(\frac{(ibx+ia+1)^2 \left(-\frac{2i(iab+3b)}{(ibx+ia+1)b} + i \right)}{b} + \frac{4(a-i)\log\left(\frac{1}{\sqrt{(bx+a)^2+1|b|}}\right)}{b} \right)}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="giac")**[Out]** -1/2*I*((I*b*x + I*a + 1)^2*(-2*I*(I*a*b + 3*b)/((I*b*x + I*a + 1)*b) + I)/b + 4*(a - I)*log(1/(sqrt((b*x + a)^2 + 1)*abs(b)))/b/b**Mupad [B]**

time = 0.50, size = 51, normalized size = 1.28

$$\ln\left(x + \frac{a-i}{b}\right) \left(\frac{2}{b^2} + \frac{a2i}{b^2}\right) - \frac{x^2}{2} + x \left(\frac{a-i}{b} - \frac{a+1i}{b}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*((a + b*x)^2 + 1))/(a*1i + b*x*1i + 1)^2,x)**[Out]** log(x + (a - 1i)/b)*((a*2i)/b^2 + 2/b^2) - x^2/2 + x*((a - 1i)/b - (a + 1i)/b)

3.202 $\int e^{-2i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=23

$$-x - \frac{2i \log(i - a - bx)}{b}$$

[Out] $-x-2*I*\ln(I-a-b*x)/b$

Rubi [A]

time = 0.01, antiderivative size = 23, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5201, 45}

$$-x - \frac{2i \log(-a - bx + i)}{b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((-2*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $-x - ((2*I)*\text{Log}[I - a - b*x])/b$

Rule 45

$\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x] /; \text{FreeQ}\{a, b, c, d, n, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] \mid\mid (\text{EqQ}[c, 0] \&\& \text{LeQ}[7*m + 4*n + 4, 0]) \mid\mid \text{LtQ}[9*m + 5*(n + 1), 0] \mid\mid \text{GtQ}[m + n + 2, 0])]$

Rule 5201

$\text{Int}[E^{(\text{ArcTan}[(c_.)*((a_.) + (b_.)*(x_.))])^{(n_.)}}, x_Symbol] \rightarrow \text{Int}[(1 - I*a*c - I*b*c*x)^{I*(n/2)} / (1 + I*a*c + I*b*c*x)^{I*(n/2)}, x] /; \text{FreeQ}\{a, b, c, n, x\}$

Rubi steps

$$\begin{aligned} \int e^{-2i \tan^{-1}(a+bx)} dx &= \int \frac{1 - ia - ibx}{1 + ia + ibx} dx \\ &= \int \left(-1 - \frac{2i}{-i + a + bx} \right) dx \\ &= -x - \frac{2i \log(i - a - bx)}{b} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 32, normalized size = 1.39

$$-x + \frac{2\text{ArcTan}(a + bx)}{b} - \frac{i \log(1 + (a + bx)^2)}{b}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((-2*I)*ArcTan[a + b*x]),x]``[Out] -x + (2*ArcTan[a + b*x])/b - (I*Log[1 + (a + b*x)^2])/b`**Maple [A]**

time = 0.07, size = 22, normalized size = 0.96

method	result	size
default	$-x - \frac{2i \ln(-bx - a + i)}{b}$	22
risch	$-x - \frac{i \ln(b^2 x^2 + 2abx + a^2 + 1)}{b} + \frac{2 \arctan(bx + a)}{b}$	40

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x,method=_RETURNVERBOSE)``[Out] -x-2*I*ln(I-a-b*x)/b`**Maxima [A]**

time = 0.26, size = 19, normalized size = 0.83

$$-x - \frac{2i \log(i bx + i a + 1)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="maxima")``[Out] -x - 2*I*log(I*b*x + I*a + 1)/b`**Fricas [A]**

time = 2.55, size = 22, normalized size = 0.96

$$-\frac{bx + 2i \log\left(\frac{bx+a-i}{b}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="fricas")``[Out] -(b*x + 2*I*log((b*x + a - I)/b))/b`

Sympy [A]

time = 0.07, size = 15, normalized size = 0.65

$$-x - \frac{2i \log(a + bx - i)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))**2*(1+(b*x+a)**2),x)

[Out] -x - 2*I*log(a + b*x - I)/b

Giac [A]

time = 0.45, size = 37, normalized size = 1.61

$$\frac{i(ibx + ia + 1)}{b} + \frac{2i \log\left(\frac{1}{\sqrt{(bx + a)^2 + 1} |b|}\right)}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2),x, algorithm="giac")

[Out] I*(I*b*x + I*a + 1)/b + 2*I*log(1/(sqrt((b*x + a)^2 + 1)*abs(b)))/b

Mupad [B]

time = 0.06, size = 21, normalized size = 0.91

$$-x - \frac{\ln\left(x + \frac{a-i}{b}\right) 2i}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)/(a*1i + b*x*1i + 1)^2,x)

[Out] - x - (log(x + (a - 1i)/b)*2i)/b

$$3.203 \quad \int \frac{e^{-2i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=41

$$\frac{(i+a) \log(x)}{i-a} - \frac{2 \log(i-a-bx)}{1+ia}$$

[Out] (I+a)*ln(x)/(I-a)-2*ln(I-a-b*x)/(1+I*a)

Rubi [A]

time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$\frac{(a+i) \log(x)}{-a+i} - \frac{2 \log(-a-bx+i)}{1+ia}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((2*I)*ArcTan[a + b*x])*x),x]

[Out] ((I + a)*Log[x])/(I - a) - (2*Log[I - a - b*x])/(1 + I*a)

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{1 - ia - ibx}{x(1 + ia + ibx)} dx \\ &= \int \left(\frac{-i - a}{(-i + a)x} + \frac{2ib}{(-i + a)(-i + a + bx)} \right) dx \\ &= \frac{(i + a) \log(x)}{i - a} - \frac{2 \log(i - a - bx)}{1 + ia} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 34, normalized size = 0.83

$$\frac{-((i + a) \log(x)) + 2i \log(i - a - bx)}{-i + a}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a + b*x])*x),x]``[Out] (-((I + a)*Log[x]) + (2*I)*Log[I - a - b*x])/(-I + a)`**Maple [A]**

time = 0.10, size = 42, normalized size = 1.02

method	result	size
default	$-\frac{2i \ln(-bx-a+i)}{i-a} + \frac{(-a^2-1) \ln(x)}{(i-a)^2}$	42
risch	$-\frac{i \ln(b^2x^2+2abx+a^2+1)}{i-a} + \frac{2 \arctan(bx+a)}{i-a} + \frac{i \ln(x)}{i-a} + \frac{\ln(x)a}{i-a}$	72

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x,x,method=_RETURNVERBOSE)``[Out] -2*I/(I-a)*ln(I-a-b*x)+(-a^2-1)/(I-a)^2*ln(x)`**Maxima [A]**

time = 0.26, size = 47, normalized size = 1.15

$$\frac{2(-ia-1) \log(ibx+ia+1)}{a^2-2ia-1} - \frac{(a^2+1) \log(x)}{a^2-2ia-1}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x,x, algorithm="maxima")``[Out] -2*(-I*a - 1)*log(I*b*x + I*a + 1)/(a^2 - 2*I*a - 1) - (a^2 + 1)*log(x)/(a^2 - 2*I*a - 1)`**Fricas [A]**

time = 3.91, size = 27, normalized size = 0.66

$$\frac{(a + i) \log(x) - 2i \log\left(\frac{bx+a-i}{b}\right)}{a - i}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x,x, algorithm="fricas")``[Out] -((a + I)*log(x) - 2*I*log((b*x + a - I)/b))/(a - I)`

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(24) = 48$.
time = 0.47, size = 99, normalized size = 2.41

$$\frac{(a+i) \log\left(a^2 - \frac{a^2(a+i)}{a-i} + \frac{2ia(a+i)}{a-i} + x(ab+3ib) + 1 + \frac{a+i}{a-i}\right)}{a-i} + \frac{2i \log\left(a^2 + \frac{2ia^2}{a-i} + \frac{4a}{a-i} + x(ab+3ib) + 1 - \frac{2i}{a-i}\right)}{a-i}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))**2*(1+(b*x+a)**2)/x,x)

[Out] $-(a + I) \log(a^2 - a^2(a + I)/(a - I) + 2I*a*(a + I)/(a - I) + x*(a*b + 3*I*b) + 1 + (a + I)/(a - I))/(a - I) + 2*I \log(a^2 + 2I*a^2/(a - I) + 4*a/(a - I) + x*(a*b + 3*I*b) + 1 - 2I/(a - I))/(a - I)$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 70 vs. $2(32) = 64$.
time = 0.45, size = 70, normalized size = 1.71

$$-ib \left(\frac{(-ia + 1) \log\left(\frac{a}{ibx+ia+1} - \frac{i}{ibx+ia+1} + i\right)}{ab - ib} + \frac{i \log\left(\frac{1}{\sqrt{(bx+a)^2 + 1|b|}}\right)}{b} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x,x, algorithm="giac")

[Out] $-I*b*((-I*a + 1)*\log(a/(I*b*x + I*a + 1) - I/(I*b*x + I*a + 1) + I)/(a*b - I*b) + I*\log(1/\sqrt{(b*x + a)^2 + 1}*abs(b)))/b$

Mupad [B]

time = 0.72, size = 34, normalized size = 0.83

$$-\frac{2 \ln(a + bx - i)}{1 + a \operatorname{li}} + \ln(x) \left(\frac{2}{1 + a \operatorname{li}} - 1 \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)/(x*(a*1i + b*x*1i + 1)^2),x)

[Out] $\log(x)*(2/(a*1i + 1) - 1) - (2*\log(a + b*x - 1i))/(a*1i + 1)$

3.204 $\int \frac{e^{-2i \operatorname{ArcTan}(a+bx)}}{x^2} dx$

Optimal. Leaf size=62

$$-\frac{i+a}{(i-a)x} + \frac{2ib \log(x)}{(i-a)^2} - \frac{2ib \log(i-a-bx)}{(i-a)^2}$$

[Out] $(-I-a)/(I-a)/x+2*I*b*\ln(x)/(I-a)^2-2*I*b*\ln(I-a-b*x)/(I-a)^2$

Rubi [A]

time = 0.03, antiderivative size = 62, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$\frac{2ib \log(x)}{(-a+i)^2} - \frac{2ib \log(-a-bx+i)}{(-a+i)^2} - \frac{a+i}{(-a+i)x}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((2*I)*\text{ArcTan}[a + b*x])}*x^2), x]$

[Out] $-((I + a)/((I - a)*x)) + ((2*I)*b*\text{Log}[x])/(I - a)^2 - ((2*I)*b*\text{Log}[I - a - b*x])/(I - a)^2$

Rule 78

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol)] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

$\text{Int}[E^{\text{ArcTan}[(c_.)*((a_. + (b_.)*(x_.)))]*(n_.)*((d_. + (e_.)*(x_.))^{(m_.)}, x_Symbol)] :> \text{Int}[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^{(I*(n/2)})/(1 + I*a*c + I*b*c*x)^{(I*(n/2)}), x] /;$ FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{1 - ia - ibx}{x^2(1 + ia + ibx)} dx \\ &= \int \left(\frac{-i - a}{(-i + a)x^2} + \frac{2ib}{(-i + a)^2 x} - \frac{2ib^2}{(-i + a)^2(-i + a + bx)} \right) dx \\ &= -\frac{i + a}{(i - a)x} + \frac{2ib \log(x)}{(i - a)^2} - \frac{2ib \log(i - a - bx)}{(i - a)^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 0.68

$$\frac{1 + a^2 + 2ibx \log(x) - 2ibx \log(i - a - bx)}{(-i + a)^2 x}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((2*I)*ArcTan[a + b*x]))*x^2, x]

[Out] (1 + a^2 + (2*I)*b*x*Log[x] - (2*I)*b*x*Log[I - a - b*x])/((-I + a)^2*x)

Maple [A]

time = 0.11, size = 69, normalized size = 1.11

method	result
default	$-\frac{-a^2-1}{x(i-a)^2} - \frac{2b(ia+1)\ln(x)}{(i-a)^3} + \frac{2b(ia+1)\ln(-bx-a+i)}{(i-a)^3}$
risch	$\frac{i}{(a-i)x} + \frac{a}{(a-i)x} - \frac{2b\ln((-2a^2b-2b)x)}{ia^2+2a-i} + \frac{b\ln(4a^4b^2x^2+8a^5bx+4a^6+8a^2b^2x^2+16a^3bx+12a^4+4b^2x^2+8abx+12a^2+4)}{ia^2+2a-i}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^2, x, method=_RETURNVERBOSE)

[Out] -(-a^2-1)/x/(I-a)^2-2*b*(1+I*a)/(I-a)^3*ln(x)+2*b*(1+I*a)/(I-a)^3*ln(I-a-b*x)

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 110 vs. 2(41) = 82.

time = 0.27, size = 110, normalized size = 1.77

$$-\frac{2(a-i)b \log(ibx+ia+1)}{-ia^3-3a^2+3ia+1} + \frac{2(a-i)b \log(x)}{-ia^3-3a^2+3ia+1} + \frac{a^3+(a^2+1)bx-ia^2+a-i}{(a^2-2ia-1)bx^2+(a^3-3ia^2-3a+i)x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^2, x, algorithm="maxima")

[Out] -2*(a - I)*b*log(I*b*x + I*a + 1)/(-I*a^3 - 3*a^2 + 3*I*a + 1) + 2*(a - I)*b*log(x)/(-I*a^3 - 3*a^2 + 3*I*a + 1) + (a^3 + (a^2 + 1)*b*x - I*a^2 + a - I)/((a^2 - 2*I*a - 1)*b*x^2 + (a^3 - 3*I*a^2 - 3*a + I)*x)

Fricas [A]

time = 4.08, size = 40, normalized size = 0.65

$$\frac{2ibx \log(x) - 2ibx \log\left(\frac{bx+a-i}{b}\right) + a^2 + 1}{(a^2 - 2ia - 1)x}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^2,x, algorithm="fricas")`

[Out] $(2*I*b*x*\log(x) - 2*I*b*x*\log((b*x + a - I)/b) + a^2 + 1)/((a^2 - 2*I*a - 1)*x)$

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 158 vs. $2(41) = 82$.

time = 0.35, size = 158, normalized size = 2.55

$$\frac{2ib \log\left(-\frac{2a^3b}{(a-i)^2} + \frac{6ia^2b}{(a-i)^2} + 2ab + \frac{6ab}{(a-i)^2} + 4b^2x - 2ib - \frac{2ib}{(a-i)^2}\right)}{(a-i)^2} - \frac{2ib \log\left(\frac{2a^3b}{(a-i)^2} - \frac{6ia^2b}{(a-i)^2} + 2ab - \frac{6ab}{(a-i)^2} + 4b^2x - 2ib + \frac{2ib}{(a-i)^2}\right)}{(a-i)^2} - \frac{-a-i}{x(a-i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^2,x)`

[Out] $2*I*b*\log(-2*a**3*b/(a - I)**2 + 6*I*a**2*b/(a - I)**2 + 2*a*b + 6*a*b/(a - I)**2 + 4*b**2*x - 2*I*b - 2*I*b/(a - I)**2)/(a - I)**2 - 2*I*b*\log(2*a**3*b/(a - I)**2 - 6*I*a**2*b/(a - I)**2 + 2*a*b - 6*a*b/(a - I)**2 + 4*b**2*x - 2*I*b + 2*I*b/(a - I)**2)/(a - I)**2 - (-a - I)/(x*(a - I))$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(41) = 82$.

time = 0.44, size = 95, normalized size = 1.53

$$\frac{2b^2 \log\left(-\frac{ia}{ibx+ia+1} - \frac{1}{ibx+ia+1} + 1\right)}{-ia^2b - 2ab + ib} - \frac{ab + ib}{(a-i)^2\left(\frac{ia}{ibx+ia+1} + \frac{1}{ibx+ia+1} - 1\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^2,x, algorithm="giac")`

[Out] $2*b^2*\log(-I*a/(I*b*x + I*a + 1) - 1/(I*b*x + I*a + 1) + 1)/(-I*a^2*b - 2*a*b + I*b) - (a*b + I*b)/((a - I)^2*(I*a/(I*b*x + I*a + 1) + 1/(I*b*x + I*a + 1) - 1))$

Mupad [B]

time = 0.65, size = 100, normalized size = 1.61

$$\frac{-1 + a \operatorname{li}}{x(1 + a \operatorname{li})} - \frac{4b \operatorname{atan}\left(\frac{a^2 \operatorname{li} + 2a - i}{(a-i)^2} + \frac{x(2a^4 b^2 + 4a^2 b^2 + 2b^2)}{(a-i)^2(-1ib a^3 + b a^2 - 1ib a + b)}\right)}{(a-i)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(((a + b*x)^2 + 1)/(x^2*(a*1i + b*x*1i + 1)^2),x)`

[Out] $(a*1i - 1)/(x*(a*1i + 1)) - (4*b*\operatorname{atan}((2*a + a^2*1i - 1i)/(a - 1i)^2 + (x*(2*b^2 + 4*a^2*b^2 + 2*a^4*b^2))/(a - 1i)^2*(b - a*b*1i + a^2*b - a^3*b*1i)))/(a - 1i)^2$

$$3.205 \quad \int \frac{e^{-2i \operatorname{ArcTan}(a+bx)}}{x^3} dx$$

Optimal. Leaf size=83

$$\frac{-i-a}{2(i-a)x^2} - \frac{2ib}{(i-a)^2x} - \frac{2b^2 \log(x)}{(1+ia)^3} + \frac{2b^2 \log(i-a-bx)}{(1+ia)^3}$$

[Out] 1/2*(-I-a)/(I-a)/x^2-2*I*b/(I-a)^2/x-2*b^2*ln(x)/(1+I*a)^3+2*b^2*ln(I-a-b*x)/(1+I*a)^3

Rubi [A]

time = 0.04, antiderivative size = 81, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$-\frac{2b^2 \log(x)}{(1+ia)^3} + \frac{2b^2 \log(-a-bx+i)}{(1+ia)^3} - \frac{2ib}{(-a+i)^2x} - \frac{a+i}{2(-a+i)x^2}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((2*I)*ArcTan[a + b*x]))*x^3],x]

[Out] -1/2*(I + a)/((I - a)*x^2) - ((2*I)*b)/((I - a)^2*x) - (2*b^2*Log[x])/(1 + I*a)^3 + (2*b^2*Log[I - a - b*x])/(1 + I*a)^3

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\int \frac{e^{-2i \tan^{-1}(a+bx)}}{x^3} dx = \int \frac{1 - ia - ibx}{x^3(1 + ia + ibx)} dx$$

$$= \int \left(\frac{-i - a}{(-i + a)x^3} + \frac{2ib}{(-i + a)^2 x^2} - \frac{2ib^2}{(-i + a)^3 x} + \frac{2ib^3}{(-i + a)^3(-i + a + bx)} \right) dx$$

$$= -\frac{i + a}{2(i - a)x^2} - \frac{2ib}{(i - a)^2 x} - \frac{2b^2 \log(x)}{(1 + ia)^3} + \frac{2b^2 \log(i - a - bx)}{(1 + ia)^3}$$

Mathematica [A]

time = 0.03, size = 66, normalized size = 0.80

$$\frac{(-i + a)(1 + a^2 - 4ibx) - 4ib^2 x^2 \log(x) + 4ib^2 x^2 \log(i - a - bx)}{2(-i + a)^3 x^2}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a + b*x]))*x^3),x]``[Out] ((-I + a)*(1 + a^2 - (4*I)*b*x) - (4*I)*b^2*x^2*Log[x] + (4*I)*b^2*x^2*Log[I - a - b*x])/(2*(-I + a)^3*x^2)`**Maple [A]**

time = 0.13, size = 109, normalized size = 1.31

method	result
default	$-\frac{-a^4 + 2ia^3 + 2ia + 1}{2(i-a)^4 x^2} - \frac{2b^2(ia+1)\ln(x)}{(i-a)^4} - \frac{2b(ia^2+2a-i)}{(i-a)^4 x} + \frac{2b^2(ia+1)\ln(-bx-a+i)}{(i-a)^4}$
risch	$-\frac{2ibx}{a^2-2ia-1} + \frac{i+a}{2a-2i} + \frac{2b^2 \ln((2a^4b+4a^2b+2b)x)}{ia^3+3a^2-3ia-1} - \frac{b^2 \ln(4a^8b^2x^2+8a^9bx+4a^{10}+16a^6b^2x^2+32a^7bx+20a^8+24a^4b^2x^2+48a^5bx+4a^6b^2x^2+48a^5bx+4a^6b^2x^2+48a^5bx+4a^6b^2x^2)}{ia^3+3a^2-3ia-1}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^3,x,method=_RETURNVERBOSE)``[Out] -1/2/(I-a)^4*(2*I*a^3-a^4+2*I*a+1)/x^2-2*b^2*(1+I*a)/(I-a)^4*ln(x)-2*b*(I*a^2-I+2*a)/(I-a)^4/x+2*b^2*(1+I*a)/(I-a)^4*ln(I-a-b*x)`**Maxima [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 160 vs. $2(61) = 122$.

time = 0.28, size = 160, normalized size = 1.93

$$-\frac{2(-ia-1)b^2 \log(ibx+ia+1)}{a^4-4ia^3-6a^2+4ia+1} - \frac{2(ia+1)b^2 \log(x)}{a^4-4ia^3-6a^2+4ia+1} + \frac{4(-ia-1)b^2x^2+a^4-2ia^3+(a^3-5ia^2-7a+3i)bx-2ia-1}{2((a^3-3ia^2-3a+i)bx^3+(a^4-4ia^3-6a^2+4ia+1)x^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^3,x, algorithm="maxima")

[Out] $-2*(-I*a - 1)*b^2*\log(I*b*x + I*a + 1)/(a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1) - 2*(I*a + 1)*b^2*\log(x)/(a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1) + 1/2*(4*(-I*a - 1)*b^2*x^2 + a^4 - 2*I*a^3 + (a^3 - 5*I*a^2 - 7*a + 3*I)*b*x - 2*I*a - 1)/((a^3 - 3*I*a^2 - 3*a + I)*b*x^3 + (a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1)*x^2)$

Fricas [A]

time = 2.30, size = 69, normalized size = 0.83

$$\frac{-4i b^2 x^2 \log(x) + 4i b^2 x^2 \log\left(\frac{bx+a-i}{b}\right) + a^3 - 4(i a + 1)bx - i a^2 + a - i}{2(a^3 - 3i a^2 - 3a + i)x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^3,x, algorithm="fricas")

[Out] $1/2*(-4*I*b^2*x^2*\log(x) + 4*I*b^2*x^2*\log((b*x + a - I)/b) + a^3 - 4*(I*a + 1)*b*x - I*a^2 + a - I)/((a^3 - 3*I*a^2 - 3*a + I)*x^2)$

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 226 vs. $2(61) = 122$.

time = 0.51, size = 226, normalized size = 2.72

$$\frac{2ib^2 \log\left(\frac{-2a^4b^2}{(a-i)^3} + \frac{8ia^3b^2}{(a-i)^2} + \frac{12a^2b^2}{(a-i)} + 2ab^2 - \frac{8iab^2}{(a-i)^2} + 4b^3x - 2ib^2 - \frac{2b^2}{(a-i)^2}\right)}{(a-i)^3} + \frac{2ib^2 \log\left(\frac{2a^4b^2}{(a-i)^3} - \frac{8ia^3b^2}{(a-i)^2} - \frac{12a^2b^2}{(a-i)} + 2ab^2 + \frac{8iab^2}{(a-i)^2} + 4b^3x - 2ib^2 + \frac{2b^2}{(a-i)^2}\right)}{(a-i)^3} - \frac{-a^2 + 4ibx - 1}{x^2 \cdot (2a^2 - 4ia - 2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a)**2*(1+(b*x+a)**2)/x**3,x)

[Out] $-2*I*b**2*\log(-2*a**4*b**2/(a - I)**3 + 8*I*a**3*b**2/(a - I)**3 + 12*a**2*b**2/(a - I)**3 + 2*a*b**2 - 8*I*a*b**2/(a - I)**3 + 4*b**3*x - 2*I*b**2 - 2*b**2/(a - I)**3)/(a - I)**3 + 2*I*b**2*\log(2*a**4*b**2/(a - I)**3 - 8*I*a**3*b**2/(a - I)**3 - 12*a**2*b**2/(a - I)**3 + 2*a*b**2 + 8*I*a*b**2/(a - I)**3 + 4*b**3*x - 2*I*b**2 + 2*b**2/(a - I)**3)/(a - I)**3 - (-a**2 + 4*I*b*x - 1)/(x**2*(2*a**2 - 4*I*a - 2))$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 142 vs. $2(61) = 122$.

time = 0.44, size = 142, normalized size = 1.71

$$\frac{2b^3 \log\left(-\frac{ia}{ibx+ia+1} - \frac{1}{ibx+ia+1} + 1\right)}{ia^3b + 3a^2b - 3iab - b} + \frac{\frac{iab^2-5b^2}{-ia-1} + \frac{2i(ab^3+3ib^3)}{(ibx+ia+1)b}}{2(a-i)^2\left(\frac{ia}{ibx+ia+1} + \frac{1}{ibx+ia+1} - 1\right)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^3,x, algorithm="giac")

[Out] $2*b^3*\log(-I*a/(I*b*x + I*a + 1) - 1/(I*b*x + I*a + 1) + 1)/(I*a^3*b + 3*a^2*b - 3*I*a*b - b) + 1/2*((I*a*b^2 - 5*b^2)/(-I*a - 1) + 2*I*(a*b^3 + 3*I*b^3)/((I*b*x + I*a + 1)*b))/((a - I)^2*(I*a/(I*b*x + I*a + 1) + 1/(I*b*x + I*a + 1) - 1)^2)$

Mupad [B]

time = 0.70, size = 156, normalized size = 1.88

$$\frac{\frac{a+1i}{2(a-i)} - \frac{bx2i}{(a-i)^2}}{x^2} - \frac{b^2 \operatorname{atanh}\left(\frac{-a^3+a^2 3i+3a-i}{(a-i)^3} - \frac{x(2a^8b^2+8a^6b^2+12a^4b^2+8a^2b^2+2b^2)}{(a-i)^3(ba^6+2iba^5+ba^4+4iba^3-ba^2+2iba-b)}\right)}{(a-i)^3} 4i$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\operatorname{int}(((a + b*x)^2 + 1)/(x^3*(a*1i + b*x*1i + 1)^2), x)$

[Out] $((a + 1i)/(2*(a - 1i)) - (b*x*2i)/(a - 1i)^2)/x^2 - (b^2*\operatorname{atanh}((3*a + a^2*3i - a^3 - 1i)/(a - 1i)^3 - (x*(2*b^2 + 8*a^2*b^2 + 12*a^4*b^2 + 8*a^6*b^2 + 2*a^8*b^2))/((a - 1i)^3*(a*b*2i - b - a^2*b + a^3*b*4i + a^4*b + a^5*b*2i + a^6*b))))*4i)/(a - 1i)^3$

$$3.206 \quad \int \frac{e^{-2i \operatorname{ArcTan}(a+bx)}}{x^4} dx$$

Optimal. Leaf size=104

$$\frac{-i-a}{3(i-a)x^3} - \frac{ib}{(i-a)^2x^2} + \frac{2b^2}{(1+ia)^3x} + \frac{2ib^3 \log(x)}{(i-a)^4} - \frac{2ib^3 \log(i-a-bx)}{(i-a)^4}$$

[Out] $1/3*(-I-a)/(I-a)/x^3-I*b/(I-a)^2/x^2+2*b^2/(1+I*a)^3/x+2*I*b^3*\ln(x)/(I-a)^4-2*I*b^3*\ln(I-a-b*x)/(I-a)^4$

Rubi [A]

time = 0.05, antiderivative size = 102, normalized size of antiderivative = 0.98, number of steps used = 3, number of rules used = 2, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5203, 78}

$$\frac{2ib^3 \log(x)}{(-a+i)^4} - \frac{2ib^3 \log(-a-bx+i)}{(-a+i)^4} + \frac{2b^2}{(1+ia)^3x} - \frac{ib}{(-a+i)^2x^2} - \frac{a+i}{3(-a+i)x^3}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((2*I)*ArcTan[a + b*x])*x^4),x]

[Out] $-1/3*(I+a)/((I-a)*x^3) - (I*b)/((I-a)^2*x^2) + (2*b^2)/((1+I*a)^3*x) + ((2*I)*b^3*\log[x])/(I-a)^4 - ((2*I)*b^3*\log[I-a-b*x])/(I-a)^4$

Rule 78

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)*(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n}, x] && NeQ[b*c - a*d, 0] && ((ILtQ[n, 0] && ILtQ[p, 0]) || EqQ[p, 1] || (IGtQ[p, 0] && (!IntegerQ[n] || LeQ[9*p + 5*(n + 2), 0] || GeQ[n + p + 1, 0] || (GeQ[n + p + 2, 0] && RationalQ[a, b, c, d, e, f])))

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\int \frac{e^{-2i \tan^{-1}(a+bx)}}{x^4} dx = \int \frac{1 - ia - ibx}{x^4(1 + ia + ibx)} dx$$

$$= \int \left(\frac{-i - a}{(-i + a)x^4} + \frac{2ib}{(-i + a)^2 x^3} - \frac{2ib^2}{(-i + a)^3 x^2} + \frac{2ib^3}{(-i + a)^4 x} - \frac{2ib^4}{(-i + a)^4(-i + a + bx)} \right) dx$$

$$= -\frac{i + a}{3(i - a)x^3} - \frac{ib}{(i - a)^2 x^2} + \frac{2b^2}{(1 + ia)^3 x} + \frac{2ib^3 \log(x)}{(i - a)^4} - \frac{2ib^3 \log(i - a - bx)}{(i - a)^4}$$

Mathematica [A]

time = 0.04, size = 91, normalized size = 0.88

$$\frac{(-i + a)(-i + a - ia^2 + a^3 - 3bx - 3iabx + 6ib^2x^2) + 6ib^3x^3 \log(x) - 6ib^3x^3 \log(i - a - bx)}{3(-i + a)^4 x^3}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a + b*x]))*x^4),x]`

```
[Out] ((-I + a)*(-I + a - I*a^2 + a^3 - 3*b*x - (3*I)*a*b*x + (6*I)*b^2*x^2) + (6*I)*b^3*x^3*Log[x] - (6*I)*b^3*x^3*Log[I - a - b*x])/(3*(-I + a)^4*x^3)
```

Maple [A]

time = 0.15, size = 150, normalized size = 1.44

method	result
default	$-\frac{a^5 - 3ia^4 - 2a^3 - 2ia^2 - 3a + i}{3(i-a)^5 x^3} + \frac{b(ia^3 + 3a^2 - 3ia - 1)}{(i-a)^5 x^2} - \frac{2b^3(ia+1)\ln(x)}{(i-a)^5} - \frac{2b^2(ia^2 + 2a - i)}{(i-a)^5 x} + \frac{2b^3(ia+1)\ln(-bx-a+i)}{(i-a)^5}$
risch	$\frac{2ib^2 x^2}{(a^2 - 2ia - 1)(a-i)} - \frac{ibx}{a^2 - 2ia - 1} + \frac{i+a}{3a-3i} - \frac{2b^3 \ln((-2a^6 b - 6a^4 b - 6a^2 b - 2b)x)}{ia^4 + 4a^3 - 6ia^2 - 4a + i} + \frac{b^3 \ln(4a^{12} b^2 x^2 + 8a^{13} bx + 4a^{14} + 24a^{10} b^2 x^2 + 48a^{11} bx)}{ia^4 + 4a^3 - 6ia^2 - 4a + i}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^4,x,method=_RETURNVERBOSE)`

```
[Out] -1/3/(I-a)^5*(-3*I*a^4+a^5-2*I*a^2-2*a^3+I-3*a)/x^3+b*(I*a^3-3*I*a+3*a^2-1)/(I-a)^5/x^2-2*b^3*(1+I*a)/(I-a)^5*ln(x)-2*b^2*(I*a^2-I+2*a)/(I-a)^5/x+2*b^3*(1+I*a)/(I-a)^5*ln(I-a-b*x)
```

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 218 vs. $2(72) = 144$.

time = 0.28, size = 218, normalized size = 2.10

$$\frac{2(a-i)b^3 \log(ibx+ia+1)}{ia^5+5a^4-10ia^3-10a^2+5ia+1} - \frac{2(a-i)b^3 \log(x)}{ia^5+5a^4-10ia^3-10a^2+5ia+1} + \frac{6(a-i)b^3 x^3 - ia^5 + 3(a^2 - 2ia - 1)b^2 x^2 - 3a^4 + 2ia^3 - (ia^4 + 5a^3 - 9ia^2 - 7a + 2i)bx - 2a^2 + 3ia + 1}{3((-ia^4 - 4a^3 + 6ia^2 + 4a - i)bx^4 + (-ia^5 - 5a^4 + 10ia^3 + 10a^2 - 5ia - 1)x^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^4,x, algorithm="maxima")

[Out] $2*(a - I)*b^3*\log(I*b*x + I*a + 1)/(I*a^5 + 5*a^4 - 10*I*a^3 - 10*a^2 + 5*I*a + 1) - 2*(a - I)*b^3*\log(x)/(I*a^5 + 5*a^4 - 10*I*a^3 - 10*a^2 + 5*I*a + 1) + 1/3*(6*(a - I)*b^3*x^3 - I*a^5 + 3*(a^2 - 2*I*a - 1)*b^2*x^2 - 3*a^4 + 2*I*a^3 - (I*a^4 + 5*a^3 - 9*I*a^2 - 7*a + 2*I)*b*x - 2*a^2 + 3*I*a + 1)/((-I*a^4 - 4*a^3 + 6*I*a^2 + 4*a - I)*b*x^4 + (-I*a^5 - 5*a^4 + 10*I*a^3 + 10*a^2 - 5*I*a - 1)*x^3)$

Fricas [A]

time = 2.68, size = 94, normalized size = 0.90

$$\frac{6i b^3 x^3 \log(x) - 6i b^3 x^3 \log\left(\frac{bx+a-i}{b}\right) - 6(-ia-1)b^2 x^2 + a^4 - 2i a^3 - 3(i a^2 + 2a - i)bx - 2i a - 1}{3(a^4 - 4i a^3 - 6a^2 + 4i a + 1)x^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^4,x, algorithm="fricas")

[Out] $1/3*(6*I*b^3*x^3*\log(x) - 6*I*b^3*x^3*\log((b*x + a - I)/b) - 6*(-I*a - 1)*b^2*x^2 + a^4 - 2*I*a^3 - 3*(I*a^2 + 2*a - I)*b*x - 2*I*a - 1)/((a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1)*x^3)$

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 286 vs. $2(75) = 150$.

time = 0.67, size = 286, normalized size = 2.75

$$\frac{2ib^3 \log\left(-\frac{2a^2i^2}{(a-i)^2} + \frac{10ia^2i^2}{(a-i)^2} + \frac{20a^2i^2}{(a-i)^2} - \frac{20a^2i^2}{(a-i)^2} + 2ab^3 - \frac{10ab^2}{(a-i)} + 4b^4x - 2ib^3 + \frac{2ib^3}{(a-i)^2}\right) - \frac{2ib^3 \log\left(\frac{2a^2i^2}{(a-i)^2} - \frac{10ia^2i^2}{(a-i)^2} - \frac{20a^2i^2}{(a-i)^2} + \frac{20a^2i^2}{(a-i)^2} + 2ab^3 + \frac{10ab^2}{(a-i)} + 4b^4x - 2ib^3 - \frac{2ib^3}{(a-i)^2}\right)}{(a-i)^4} - \frac{-a^3 + ia^2 - a - 6ib^2x^2 + x(3iab + 3b) + i}{x^3 \cdot (3a^3 - 9ia^2 - 9a + 3i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a)**2*(1+(b*x+a)**2)/x**4,x)

[Out] $2*I*b**3*\log(-2*a**5*b**3/(a - I)**4 + 10*I*a**4*b**3/(a - I)**4 + 20*a**3*b**3/(a - I)**4 - 20*I*a**2*b**3/(a - I)**4 + 2*a*b**3 - 10*a*b**3/(a - I)**4 + 4*b**4*x - 2*I*b**3 + 2*I*b**3/(a - I)**4)/(a - I)**4 - 2*I*b**3*\log(2*a**5*b**3/(a - I)**4 - 10*I*a**4*b**3/(a - I)**4 - 20*a**3*b**3/(a - I)**4 + 20*I*a**2*b**3/(a - I)**4 + 2*a*b**3 + 10*a*b**3/(a - I)**4 + 4*b**4*x - 2*I*b**3 - 2*I*b**3/(a - I)**4)/(a - I)**4 - (-a**3 + I*a**2 - a - 6*I*b**2*x**2 + x*(3*I*a*b + 3*b) + I)/(x**3*(3*a**3 - 9*I*a**2 - 9*a + 3*I))$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 183 vs. $2(72) = 144$.

time = 0.44, size = 183, normalized size = 1.76

$$\frac{2b^4 \log\left(-\frac{ia}{ibx+ia+1} - \frac{1}{ibx+ia+1} + 1\right)}{-ia^4b - 4a^3b + 6ia^2b + 4ab - ib} + \frac{-iab^3+10b^3}{ia+1} + \frac{3i(ab^4+8ib^4)}{(ibx+ia+1)b} + \frac{3(a^2b^5+4iab^5+5b^5)}{(ibx+ia+1)^2b^2}}{3(a-i)^3\left(\frac{ia}{ibx+ia+1} + \frac{1}{ibx+ia+1} - 1\right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^2*(1+(b*x+a)^2)/x^4,x, algorithm="giac")

[Out] $2*b^4*\log(-I*a/(I*b*x + I*a + 1) - 1/(I*b*x + I*a + 1) + 1)/(-I*a^4*b - 4*a^3*b + 6*I*a^2*b + 4*a*b - I*b) + 1/3*((-I*a*b^3 + 10*b^3)/(I*a + 1) + 3*I*(a*b^4 + 8*I*b^4)/((I*b*x + I*a + 1)*b) + 3*(a^2*b^5 + 4*I*a*b^5 + 5*b^5)/((I*b*x + I*a + 1)^2*b^2))/((a - I)^3*(I*a/(I*b*x + I*a + 1) + 1/(I*b*x + I*a + 1) - 1)^3)$

Mupad [B]

time = 0.77, size = 199, normalized size = 1.91

$$\frac{\frac{a+1i}{3(a-i)} + \frac{b^2 x^2 2i}{(a-i)^3} - \frac{b x 1i}{(a-i)^2}}{x^3} - \frac{4b^3 \operatorname{atan}\left(\frac{(a^4 - a^3 4i - 6a^2 + a 4i + 1) 1i}{(a-i)^4} + \frac{x(2a^{12}b^2 + 12a^{10}b^2 + 30a^8b^2 + 40a^6b^2 + 30a^4b^2 + 12a^2b^2 + 2b^2)}{(a-i)^4(-1i b a^9 + 3b a^8 + 8b a^6 + 6i b a^5 + 6b a^4 + 8i b a^3 + 3i b a - b)}\right)}{(a-i)^4}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)/(x^4*(a*1i + b*x*1i + 1)^2),x)

[Out] $((a + 1i)/(3*(a - 1i)) + (b^2*x^2*2i)/(a - 1i)^3 - (b*x*1i)/(a - 1i)^2)/x^3 - (4*b^3*\operatorname{atan}(((a*4i - 6*a^2 - a^3*4i + a^4 + 1)*1i)/(a - 1i)^4 + (x*(2*b^2 + 12*a^2*b^2 + 30*a^4*b^2 + 40*a^6*b^2 + 30*a^8*b^2 + 12*a^{10}*b^2 + 2*a^{12}*b^2))/((a - 1i)^4*(a*b*3i - b + a^3*b*8i + 6*a^4*b + a^5*b*6i + 8*a^6*b + 3*a^8*b - a^9*b*1i))))/(a - 1i)^4)$

3.207 $\int e^{-3i\text{ArcTan}(a+bx)} x^4 dx$

Optimal. Leaf size=324

$$\frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(19i-68a-88ia^2+48a^3+8ia^4)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{8b^5} - \frac{3(17i-16a)x^2}{8b^5}$$

[Out] $-3/8*(19+68*I*a-88*a^2-48*I*a^3+8*a^4)*\text{arcsinh}(b*x+a)/b^5+2*I*x^4*(1-I*a-I*b*x)^{(3/2)}/b/(1+I*a+I*b*x)^{(1/2)}-3/20*(17*I-16*a)*x^2*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3-11/5*x^3*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2+1/40*I*(1-I*a-I*b*x)^{(3/2)}*(163+458*I*a-422*a^2-112*I*a^3-2*(61*I-118*a-52*I*a^2)*b*x)*(1+I*a+I*b*x)^{(1/2)}/b^5+3/8*(19*I-68*a-88*I*a^2+48*a^3+8*I*a^4)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^5$

Rubi [A]

time = 0.20, antiderivative size = 324, normalized size of antiderivative = 1.00, number of steps used = 9, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 99, 158, 152, 52, 55, 633, 221}

$\frac{(-ia-ibx+1)^{3/2}\sqrt{ia+ibx+1}(-112ia^3-2(-52ia^2-118a+61)bx-422a^2+458a+163)}{40b^5} + \frac{3(8a^4+48a^3-88a^2-68a+19)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{80b^5} - \frac{3(8a^4-48a^3-88a^2+68a+19)\text{sinh}^{-1}(a+bx)}{80b^5} - \frac{3(-16a+17i)x^2(-ia-ibx+1)^{3/2}\sqrt{ia+ibx+1}}{20b^5} - \frac{11x^2(-ia-ibx+1)^{3/2}\sqrt{ia+ibx+1}}{5b^2} + \frac{2ix^4(-ia-ibx+1)^{3/2}}{4b\sqrt{ia+ibx+1}}$

Antiderivative was successfully verified.

[In] Int[x^4/E^((3*I)*ArcTan[a + b*x]),x]

[Out] $((2*I)*x^4*(1-I*a-I*b*x)^{(3/2)})/(b*\text{Sqrt}[1+I*a+I*b*x]) + (3*(19*I-68*a-(88*I)*a^2+48*a^3+(8*I)*a^4)*\text{Sqrt}[1-I*a-I*b*x]*\text{Sqrt}[1+I*a+I*b*x])/(8*b^5) - (3*(17*I-16*a)*x^2*(1-I*a-I*b*x)^{(3/2)}*\text{Sqrt}[1+I*a+I*b*x])/(20*b^3) - (11*x^3*(1-I*a-I*b*x)^{(3/2)}*\text{Sqrt}[1+I*a+I*b*x])/(5*b^2) + ((I/40)*(1-I*a-I*b*x)^{(3/2)}*\text{Sqrt}[1+I*a+I*b*x]*(163+(458*I)*a-422*a^2-(112*I)*a^3-2*(61*I-118*a-(52*I)*a^2)*b*x))/b^5 - (3*(19+(68*I)*a-88*a^2-(48*I)*a^3+8*a^4)*\text{ArcSinh}[a+b*x])/(8*b^5)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b

+ d, 0] && GtQ[a + c, 0]

Rule 99

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^p/(b*(m + 1))), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

Rule 152

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)), Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]

Rule 158

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p + 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(a+bx)} x^4 dx &= \int \frac{x^4(1-ia-ibx)^{3/2}}{(1+ia+ibx)^{3/2}} dx \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{(2i) \int \frac{x^3 \sqrt{1-ia-ibx} (4(1-ia) - \frac{11ibx}{2})}{\sqrt{1+ia+ibx}} dx}{b} \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{11x^3(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{5b^2} - \frac{(2i) \int x^2 \sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} dx \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{3(17i-16a)x^2(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{20b^3} - \frac{11x^3(1-ia-ibx)^{3/2}}{5b^2} \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{3(17i-16a)x^2(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{20b^3} - \frac{11x^3(1-ia-ibx)^{3/2}}{5b^2} \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(19i-68a-88ia^2+48a^3+8ia^4)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{8b^5} \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(19i-68a-88ia^2+48a^3+8ia^4)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{8b^5} \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(19i-68a-88ia^2+48a^3+8ia^4)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{8b^5} \\
&= \frac{2ix^4(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(19i-68a-88ia^2+48a^3+8ia^4)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{8b^5}
\end{aligned}$$

Mathematica [A]

time = 0.37, size = 299, normalized size = 0.92

$$\frac{448i+8ia^6+2853a+22487x^2+9593x^3-5614x^4-3085x^5+816x^6+a^7(410+816x)+2a^8(-6384+2053a)+a^9(-905-200448x+6087x^2)-a^{10}(8366+26355a+356487x^2+2092x^3)+a^{11}(-1315+146888a-51582x^2+116487x^3+1085x^4+8165x^5)}{\sqrt{1+a^2+2abx+b^2x^2}} + \frac{30\sqrt{-1}(19i-68a-88ia^2+48a^3+8ia^4)\sqrt{b}\operatorname{arctanh}\left(\frac{(1+i)\sqrt{b}\sqrt{-i(a+bx)}}{\sqrt{-ib}}\right)}{\sqrt{-ib}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[x^4/E^((3*I)*ArcTan[a + b*x]),x]

[Out]
$$\begin{aligned} & ((448*I + (8*I)*a^6 + 285*b*x + (224*I)*b^2*x^2 + 95*b^3*x^3 - (56*I)*b^4*x^4 \\ & - 30*b^5*x^5 + (8*I)*b^6*x^6 + a^5*(410 + (8*I)*b*x) + 2*a^4*(-638*I + 2 \\ & 65*b*x) + a^3*(-905 - (2004*I)*b*x + 60*b^2*x^2) - a^2*(836*I + 2635*b*x + \\ & (356*I)*b^2*x^2 + 20*b^3*x^3) + a*(-1315 + (1468*I)*b*x - 515*b^2*x^2 + (11 \\ & 6*I)*b^3*x^3 + 10*b^4*x^4 + (8*I)*b^5*x^5))/\text{Sqrt}[1 + a^2 + 2*a*b*x + b^2*x^2] \\ & + (30*(-1)^{(1/4)}*(19*I - 68*a - (88*I)*a^2 + 48*a^3 + (8*I)*a^4)*\text{Sqrt}[b] \\ & *\text{ArcSinh}[(1/2 + I/2)*\text{Sqrt}[b]*\text{Sqrt}[(-I)*(I + a + b*x)])/ \text{Sqrt}[(-I)*b])/ \text{Sqrt} \\ & [(-I)*b)]/(40*b^5) \end{aligned}$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1361 vs. $2(262) = 524$.

time = 0.22, size = 1362, normalized size = 4.20

method	result
risch	$\frac{i(8x^4b^4 - 8ab^3x^3 + 30ib^3x^3 + 8a^2b^2x^2 - 70iab^2x^2 - 8a^3bx + 130ix a^2b + 8a^4 - 250ia^3 - 64b^2x^2 + 252abx - 125ibx - 804a^2 + 835ia + 288)\sqrt{b^5}}{40b^5}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^4/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out]
$$\begin{aligned} & I/b^4*(b*(1/5*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)/b^2-a/b*(1/8*(2*b^2*x+2*a*b)/b^2 \\ & *(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2* \\ & b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2 \\ &)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2) \\ &))+3*I*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+3/16*(4*b^2* \\ & (a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2) \\ & +1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2* \\ & a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))-3*a*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a \\ & *b*x+a^2+1)^(3/2)+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b \\ & ^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2* \\ & x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))) + 6/b^5*(I*a \\ & ^2-I+2*a)*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(\\ & I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((\\ & I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2) \\ &))/(b^2)^(1/2))+I*a^4-6*I*a^2+4*a^3+I-4*a)/b^7*(I/b/(x-(I-a)/b)^3*((x-(I- \\ & a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)-2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b) \\ & ^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a) \\ &)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b \\ & *(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2 \\ & *b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))))+4*(-I*a^3-3*a^2+3*I*a+1)/b^6 \\ & *(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3 \end{aligned}$$

$$\left(\left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{3/2} + Ib \left(\frac{1}{4} \left(2 \left(\frac{x-(I-a)}{b} \right) b^2 + 2Ib \right) / b^2 \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{1/2} + \frac{1}{2} \ln \left(\frac{Ib + \left(\frac{x-(I-a)}{b} \right) b^2}{(b^2)^{1/2}} + \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{1/2} \right) / (b^2)^{1/2} \right)$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1368 vs. $2(230) = 460$.
time = 0.51, size = 1368, normalized size = 4.22

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($x^4/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^{3/2}$, x, algorithm="maxima")

[Out] $I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a^4/(b^7*x^2 + 2*a*b^6*x + a^2*b^5 - 2*I*b^6*x - 2*I*a*b^5 - b^5) + 4*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a^3/(b^7*x^2 + 2*a*b^6*x + a^2*b^5 - 2*I*b^6*x - 2*I*a*b^5 - b^5) + 4*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a^3/(2*I*b^6*x + 2*I*a*b^5 + 2*b^5) + 6*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^4/(I*b^6*x + I*a*b^5 + b^5) - 6*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a^2/(b^7*x^2 + 2*a*b^6*x + a^2*b^5 - 2*I*b^6*x - 2*I*a*b^5 - b^5) - 12*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a^2/(2*I*b^6*x + 2*I*a*b^5 + 2*b^5) + 24*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^3/(I*b^6*x + I*a*b^5 + b^5) - 4*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a/(b^7*x^2 + 2*a*b^6*x + a^2*b^5 - 2*I*b^6*x - 2*I*a*b^5 - b^5) - 12*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a/(2*I*b^6*x + 2*I*a*b^5 + 2*b^5) - 36*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^2/(I*b^6*x + I*a*b^5 + b^5) + I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}/(b^7*x^2 + 2*a*b^6*x + a^2*b^5 - 2*I*b^6*x - 2*I*a*b^5 - b^5) + 4*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}/(2*I*b^6*x + 2*I*a*b^5 + 2*b^5) - 24*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a/(I*b^6*x + I*a*b^5 + b^5) - 3*a^4*\operatorname{arcsinh}(b*x + a)/b^5 + 6*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(I*b^6*x + I*a*b^5 + b^5) - I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a*x/b^4 - 3*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}*a^2*x/b^4 + 18*I*a^3*\operatorname{arcsinh}(b*x + a)/b^5 + I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a^2/b^5 + 6*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^3/b^5 - 3*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}*a^3/b^5 - 3/4*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*x/b^4 - 3/2*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a*x/b^4 + 6*I*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}*a*x/b^4 + 3*a^2*\operatorname{arcsin}(I*b*x + I*a + 2)/b^5 + 36*a^2*\operatorname{arcsinh}(b*x + a)/b^5 + 1/5*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{5/2}/b^5 + 13/4*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}*a/b^5 - 39/2*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^2/b^5 + 12*I*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}*a^2/b^5 - 9/8*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*x/b^4 + 3*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}*x/b^4 - 6*I*a*\operatorname{arcsin}(I*b*x + I*a + 2)/b^5 - 63/2*I*a*\operatorname{arcsinh}(b*x + a)/b^5 - 2*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{3/2}/b^5 - 153/8*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a/b^5 + 15*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}*a/b^5 - 3*\operatorname{arcsin}(I*b*x + I*a + 2)/b^5 - 81/$

$8*\operatorname{arcsinh}(b*x + a)/b^5 + 6*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/b^5 - 6*I*\sqrt{-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3}/b^5$

Fricas [A]

time = 2.30, size = 264, normalized size = 0.81

$\frac{62*a^6 + 2687*a^5 - 11575*a^4 - 20350*a^3 + 6252*a^2 - 8950*a - 11400}{320*b^6*x + (a - I)*b^5} + \frac{6340*a^5 + 2625*a^4 - 8950*a^3 - 11400*a^2 + 6340*a + 1280}{320*b^6*x + (a - I)*b^5} * b*x + \frac{17740*a^4 + 120*(8*a^5 - 56*I*a^4 - 136*a^3 + (8*a^4 - 48*I*a^3 - 88*a^2 + 68*I*a + 19)*b*x + 156*I*a^2 + 87*a - 19*I)*\log(-b*x - a + \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 8*(-8*I*b^5*x^5 + 22*b^4*x^4 - 2*(16*a - 17*I)*b^3*x^3 - 8*I*a^5 + (52*a^2 - 118*I*a - 61)*b^2*x^2 - 418*a^4 + 1694*I*a^3 - (112*a^3 - 422*I*a^2 - 458*a + 163*I)*b*x + 2599*a^2 - 1763*I*a - 448)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 7620*a - 1280*I}{320*b^6*x + (a - I)*b^5}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{320}*(62*I*a^6 + 2687*a^5 - 11575*I*a^4 - 20350*a^3 + (62*I*a^5 + 2625*a^4 - 8950*I*a^3 - 11400*a^2 + 6340*I*a + 1280)*b*x + 17740*I*a^2 + 120*(8*a^5 - 56*I*a^4 - 136*a^3 + (8*a^4 - 48*I*a^3 - 88*a^2 + 68*I*a + 19)*b*x + 156*I*a^2 + 87*a - 19*I)*\log(-b*x - a + \sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}) - 8*(-8*I*b^5*x^5 + 22*b^4*x^4 - 2*(16*a - 17*I)*b^3*x^3 - 8*I*a^5 + (52*a^2 - 118*I*a - 61)*b^2*x^2 - 418*a^4 + 1694*I*a^3 - (112*a^3 - 422*I*a^2 - 458*a + 163*I)*b*x + 2599*a^2 - 1763*I*a - 448)*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1} + 7620*a - 1280*I)/(b^6*x + (a - I)*b^5)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**4/(1+I*(b*x+a))**3*(1+(b*x+a)**2)**(3/2),x)

[Out] Timed out

Giac [A]

time = 0.46, size = 336, normalized size = 1.04

$\frac{-\frac{1}{320}\sqrt{b^2*x^2+1}\left(\frac{1}{3}\left(\frac{62}{b^6}+\frac{2687}{b^5}+\frac{11575}{b^4}+\frac{20350}{b^3}+\frac{6252}{b^2}+\frac{8950}{b}+11400\right)+\frac{6340}{b^5}+\frac{2625}{b^4}+\frac{8950}{b^3}+\frac{11400}{b^2}+\frac{6340}{b}+1280\right)*b*x+\frac{17740}{b^4}+\frac{120}{b^5}\left(8*a^5-56*I*a^4-136*a^3+(8*a^4-48*I*a^3-88*a^2+68*I*a+19)*b*x+156*I*a^2+87*a-19*I\right)*\log\left(\frac{-b*x-a+\sqrt{b^2*x^2+2*a*b*x+a^2+1}}{b}\right)-8\left(-8*I*b^5*x^5+22*b^4*x^4-2*(16*a-17*I)*b^3*x^3-8*I*a^5+(52*a^2-118*I*a-61)*b^2*x^2-418*a^4+1694*I*a^3-(112*a^3-422*I*a^2-458*a+163*I)*b*x+2599*a^2-1763*I*a-448\right)*\sqrt{b^2*x^2+2*a*b*x+a^2+1}+7620*a-1280*I}{320*b^6*x+(a-I)*b^5}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^4/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="giac")

[Out] $-\frac{1}{40}*\sqrt{(b*x + a)^2 + 1}*((2*(x*(-4*I*x/b - (-4*I*a*b^17 - 15*b^17)/b^19) - (4*I*a^2*b^16 + 35*a*b^16 - 32*I*b^16)/b^19)*x - (-8*I*a^3*b^15 - 130*a^2*b^15 + 252*I*a*b^15 + 125*b^15)/b^19)*x - (8*I*a^4*b^14 + 250*a^3*b^14 - 804*I*a^2*b^14 - 835*a*b^14 + 288*I*b^14)/b^19 + 1/8*(8*a^4 - 48*I*a^3 - 88*a^2 + 68*I*a + 19)*\log(3*I*(x*\operatorname{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a*b + I*a^3*b + I*(x*\operatorname{abs}(b) - \sqrt{(b*x + a)^2 + 1})^3*\operatorname{abs}(b) - 3*(-I*x*\operatorname{abs}(b) + I*\sqrt{(b*x + a)^2 + 1})*a^2*\operatorname{abs}(b) + 2*(x*\operatorname{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*$

$b + 2*a^2*b + 4*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a*abs(b) - I*a*b + (-I*x*abs(b) + I*sqrt((b*x + a)^2 + 1))*abs(b))/(b^4*abs(b))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^4 ((a + bx)^2 + 1)^{3/2}}{(1 + a li + b x li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^4*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3,x)

[Out] int((x^4*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3, x)

3.208 $\int e^{-3i\text{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=249

$$\frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{8b^4} - \frac{9x^2(1-ia-ibx)^{3/2}\sqrt{1-ia-ibx}}{4b^2}$$

[Out] $\frac{3}{8}*(17*I-44*a-36*I*a^2+8*a^3)*\text{arcsinh}(b*x+a)/b^4+2*I*x^3*(1-I*a-I*b*x)^(3/2)/b/(1+I*a+I*b*x)^(1/2)-9/4*x^2*(1-I*a-I*b*x)^(3/2)*(1+I*a+I*b*x)^(1/2)/b^2-1/8*I*(1-I*a-I*b*x)^(3/2)*(29*I-54*a-22*I*a^2+2*(11+10*I*a)*b*x)*(1+I*a+I*b*x)^(1/2)/b^4+3/8*(17+44*I*a-36*a^2-8*I*a^3)*(1-I*a-I*b*x)^(1/2)*(1+I*a+I*b*x)^(1/2)/b^4$

Rubi [A]

time = 0.17, antiderivative size = 249, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 99, 158, 152, 52, 55, 633, 221}

$$\frac{i(-ia-ibx+1)^{3/2}\sqrt{ia+ibx+1}(-22ia^2+2(11+10ia)bx-54a+29i)}{8b^4} + \frac{3(-8ia^3-36a^2+44ia+17)\sqrt{-ia-ibx+1}\sqrt{ia+ibx+1}}{8b^4} + \frac{3(8a^3-36ia^2-44a+17i)\sinh^{-1}(a+bx)}{8b^4} - \frac{9x^2(-ia-ibx+1)^{3/2}\sqrt{ia+ibx+1}}{4b^2} + \frac{2ix^3(-ia-ibx+1)^{3/2}}{b\sqrt{ia+ibx+1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^3/E^{((3*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $((2*I)*x^3*(1-I*a-I*b*x)^(3/2))/(b*\text{Sqrt}[1+I*a+I*b*x]) + (3*(17+(4*4*I)*a-36*a^2-(8*I)*a^3)*\text{Sqrt}[1-I*a-I*b*x]*\text{Sqrt}[1+I*a+I*b*x])/(8*b^4) - (9*x^2*(1-I*a-I*b*x)^(3/2)*\text{Sqrt}[1+I*a+I*b*x])/(4*b^2) - ((I/8)*(1-I*a-I*b*x)^(3/2)*\text{Sqrt}[1+I*a+I*b*x]*(29*I-54*a-(22*I)*a^2+2*(11+(10*I)*a)*b*x))/b^4 + (3*(17*I-44*a-(36*I)*a^2+8*a^3)*\text{ArcSinh}[a+b*x])/(8*b^4)$

Rule 52

$\text{Int}[(a_.) + (b_.)*(x_.))^(m_.)*((c_.) + (d_.)*(x_.))^(n_.), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m + n + 1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0]))) \ \&\& \ !\text{ILtQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 55

$\text{Int}[1/(\text{Sqrt}[(a_.) + (b_.)*(x_.)]*\text{Sqrt}[(c_.) + (d_.)*(x_.)]), x_Symbol] \rightarrow \text{Int}[1/\text{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /; \text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{EqQ}[b + d, 0] \ \&\& \ \text{GtQ}[a + c, 0]$

Rule 99

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p/(b*(m + 1)), x] - Dist[1/(b*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^(p - 1)*Simp[d*e*n + c*f*p + d*f*(n + p)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f}, x] && LtQ[m, -1] && GtQ[n, 0] && GtQ[p, 0] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m + 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)), Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 158

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[h*(a + b*x)^m*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 2))), x] + Dist[1/(d*f*(m + n + p + 2)), Int[(a + b*x)^(m - 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*g*(m + n + p + 2) - h*(b*c*e*m + a*(d*e*(n + 1) + c*f*(p + 1))) + (b*d*f*g*(m + n + p + 2) + h*(a*d*f*m - b*(d*e*(m + n + 1) + c*f*(m + p + 1)))*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && GtQ[m, 0] && NeQ[m + n + p + 2, 0] && IntegerQ[m]
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*c/(b^2 - 4*a*c)))^p, Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
```

$\int e^{-3i \tan^{-1}(a+bx)} x^3 dx$ /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{-3i \tan^{-1}(a+bx)} x^3 dx &= \int \frac{x^3(1-ia-ibx)^{3/2}}{(1+ia+ibx)^{3/2}} dx \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{(2i) \int \frac{x^2 \sqrt{1-ia-ibx} (3(1-ia) - \frac{9ibx}{2})}{\sqrt{1+ia+ibx}} dx}{b} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{9x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} - \frac{i \int \frac{x \sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} dx}{b} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} - \frac{9x^2(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{4b^2} - \frac{i(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{b} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4} \\
 &= \frac{2ix^3(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3(17+44ia-36a^2-8ia^3) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{8b^4}
 \end{aligned}$$

Mathematica [A]

time = 0.28, size = 244, normalized size = 0.98

$$\frac{80 - 2ia^5 - 51ibx + 40b^2x^2 - 17ib^3x^3 - 8b^4x^4 + 2ib^5x^5 + a^4(-76 - 2ibx) - 5a^3(-31i + 20bx) + a^2(4 + 265ibx - 12b^2x^2) + a(157i + 212bx + 53ib^2x^2 + 4b^3x^3 + 2ib^4x^4)}{8b^4\sqrt{1+a^2+2abx+b^2x^2}} + \frac{3\sqrt{-1}(17i-44a-36ia^2+8a^3)\sqrt{-ib}\sinh^{-1}\left(\frac{(1+i)\sqrt{b}\sqrt{-i(a+bx)}}{\sqrt{-ib}}\right)}{4b^{9/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[x^3/E^((3*I)*ArcTan[a + b*x]),x]

[Out] (80 - (2*I)*a^5 - (51*I)*b*x + 40*b^2*x^2 - (17*I)*b^3*x^3 - 8*b^4*x^4 + (2*I)*b^5*x^5 + a^4*(-76 - (2*I)*b*x) - 5*a^3*(-31*I + 20*b*x) + a^2*(4 + (265*I)*b*x - 12*b^2*x^2) + a*(157*I + 212*b*x + (53*I)*b^2*x^2 + 4*b^3*x^3 +

$$(2*I)*b^4*x^4)/(8*b^4*sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]) + (3*(-1)^(1/4)*(17*I - 44*a - (36*I)*a^2 + 8*a^3)*sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*sqrt[b]*sqrt[(-I)*(I + a + b*x)])]/sqrt[(-I)*b])/(4*b^(9/2))$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 982 vs. 2(200) = 400.
 time = 0.19, size = 983, normalized size = 3.95

method	result
risch	$\frac{i(-2b^3x^3+2ab^2x^2-8ib^2x^2-2a^2bx+20iabx+2a^3-44ia^2+19bx-93a+48i)\sqrt{b^2x^2+2abx+a^2+1}}{8b^4} - \frac{27i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{b^3}$
default	$i \left(\frac{(2b^2x+2ab)(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}}{8b^2} + \frac{3(4b^2(a^2+1)-4a^2b^2) \left(\frac{(2b^2x+2ab)\sqrt{b^2x^2+2abx+a^2+1}}{4b^2} + \frac{(4b^2(a^2+1)-4a^2b^2) \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{16b^2} \right)}{16b^2} \right)$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^3/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x,method=_RETURNVERBOSE)
[Out] I/b^3*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/(b^2)^(1/2))+3*(-I*a-1)/b^4*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))+(-I*a^3-3*a^2+3*I*a+1)/b^6*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)-2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2)))+3/b^5*(I*a^2-I+2*a)*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2)))/b^2)^(1/2))
```

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 979 vs. 2(175) = 350.
 time = 0.48, size = 979, normalized size = 3.93

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="maxima")

[Out] $-I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a^3/(b^6*x^2 + 2*a*b^5*x + a^2*b^4 - 2*I*b^5*x - 2*I*a*b^4 - b^4) - 3*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a^2/(b^6*x^2 + 2*a*b^5*x + a^2*b^4 - 2*I*b^5*x - 2*I*a*b^4 - b^4) - 3*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a^2/(2*I*b^5*x + 2*I*a*b^4 + 2*b^4) - 6*I*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a^3/(I*b^5*x + I*a*b^4 + b^4) + 3*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a/(b^6*x^2 + 2*a*b^5*x + a^2*b^4 - 2*I*b^5*x - 2*I*a*b^4 - b^4) + 6*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a/(2*I*b^5*x + 2*I*a*b^4 + 2*b^4) - 18*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a^2/(I*b^5*x + I*a*b^4 + b^4) + (b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}/(b^6*x^2 + 2*a*b^5*x + a^2*b^4 - 2*I*b^5*x - 2*I*a*b^4 - b^4) + 3*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}/(2*I*b^5*x + 2*I*a*b^4 + 2*b^4) + 18*I*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a/(I*b^5*x + I*a*b^4 + b^4) + 3*a^3*\text{arcsinh}(b*x + a)/b^4 + 6*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)/(I*b^5*x + I*a*b^4 + b^4) + 1/4*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*x/b^3 + 3/2*\text{sqrt}(-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3)*a*x/b^3 - 27/2*I*a^2*\text{arcsinh}(b*x + a)/b^4 - 3/4*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}*a/b^4 - 9/2*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a^2/b^4 + 3/2*\text{sqrt}(-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3)*a^2/b^4 + 3/8*I*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*x/b^3 - 3/2*I*\text{sqrt}(-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3)*x/b^3 - 3/2*a*\text{arcsin}(I*b*x + I*a + 2)/b^4 - 18*a*\text{arcsinh}(b*x + a)/b^4 - (b^2*x^2 + 2*a*b*x + a^2 + 1)^{(3/2)}/b^4 + 75/8*I*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)*a/b^4 - 9/2*I*\text{sqrt}(-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3)*a/b^4 + 3/2*I*\text{arcsin}(I*b*x + I*a + 2)/b^4 + 63/8*I*\text{arcsinh}(b*x + a)/b^4 + 9/2*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)/b^4 - 3*\text{sqrt}(-b^2*x^2 - 2*a*b*x - a^2 + 4*I*b*x + 4*I*a + 3)/b^4$

Fricas [A]

time = 3.92, size = 216, normalized size = 0.87

$$\frac{-15a^5 - 495a^4 + 1664a^3 + (-15a^4 - 480a^3 + 1184a^2 + 968a - 256)bx + 2152a^2 - 24(8a^4 - 44a^3 + (8a^3 - 36a^2 - 4a + 17)bx - 80a^2 + 61a + 17)\log\left(\frac{-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}}{64(b^5x + (a - I)b^4)}\right) - 8(-2Ib^4x^4 + 6b^3x^3 - (10a - 11I)b^2x^2 + 2Ia^4 + 78a^3 + (22a^2 - 54Ia - 29)bx - 233Ia^2 - 237a + 80I)\text{sqrt}(b^2x^2 + 2abx + a^2 + 1) - 1224a - 256}{64(b^5x + (a - I)b^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="fricas")

[Out] $1/64*(-15*I*a^5 - 495*a^4 + 1664*I*a^3 + (-15*I*a^4 - 480*a^3 + 1184*I*a^2 + 968*a - 256*I)*b*x + 2152*a^2 - 24*(8*a^4 - 44*I*a^3 + (8*a^3 - 36*I*a^2 - 44*a + 17*I)*b*x - 80*a^2 + 61*I*a + 17)*\log(-b*x - a + \text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 8*(-2*I*b^4*x^4 + 6*b^3*x^3 - (10*a - 11*I)*b^2*x^2 + 2*I*a^4 + 78*a^3 + (22*a^2 - 54*I*a - 29)*b*x - 233*I*a^2 - 237*a + 80*I)*\text{sqrt}(b^2*x^2 + 2*a*b*x + a^2 + 1) - 1224*I*a - 256)/(b^5*x + (a - I)*b^4)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**3/(1+I*(b*x+a))**3*(1+(b*x+a)**2)**(3/2), x)

[Out] Timed out

Giac [A]

time = 0.45, size = 285, normalized size = 1.14

$$\frac{1}{8} \sqrt{bx+a^2+1} \left(\left(\frac{x}{a} - \frac{-10b^2-4b^2}{32a^2} \right) - \frac{2i^2b^3+20ab^3-19b^3}{32a^2} x - \frac{-2i^2b^3-44a^2b^3+33ab^3+45b^3}{32a^2} \right) - \frac{(8a^2-36a^2-44a+17) \log\left(3\left(x\sqrt{bx+a^2+1}\right)^2+2^2b^2+2\left(x\sqrt{bx+a^2+1}\right)\right) + 3\left(x\sqrt{bx+a^2+1}\right)^2-2\left(x\sqrt{bx+a^2+1}\right)^2-2i^2b^2+4\left(-i^2b^2+1\sqrt{bx+a^2+1}\right)ab-ab-\left(x\sqrt{bx+a^2+1}\right)b}{8b^2|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^3/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2), x, algorithm="giac")

[Out] $-1/8*\sqrt{(b*x + a)^2 + 1}*((2*x*(-I*x/b - (-I*a*b^{11} - 4*b^{11})/b^{13}) - (2*I*a^2*b^{10} + 20*a*b^{10} - 19*I*b^{10})/b^{13})*x - (-2*I*a^3*b^9 - 44*a^2*b^9 + 93*I*a*b^9 + 48*b^9)/b^{13}) - 1/8*(8*a^3 - 36*I*a^2 - 44*a + 17*I)*\log(3*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a*b + a^3*b + (x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^3*\text{abs}(b) + 3*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*a^2*\text{abs}(b) - 2*I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*b - 2*I*a^2*b + 4*(-I*x*\text{abs}(b) + I*\sqrt{(b*x + a)^2 + 1})*a*\text{abs}(b) - a*b - (x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*\text{abs}(b)))/(b^3*\text{abs}(b))$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3 ((a + bx)^2 + 1)^{3/2}}{(1 + a li + b x li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^3*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3,x)

[Out] int((x^3*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3, x)

3.209 $\int e^{-3i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=229

$$\frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} - \frac{(11i-18a-6ia^2)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} - \frac{(11i-18a-6ia^2)(1-ia-ibx)^{5/2}}{6b^3}$$

[Out] $\frac{1}{2}*(11+18*I*a-6*a^2)*\text{arcsinh}(b*x+a)/b^3+I*(I-a)^2*(1-I*a-I*b*x)^{(5/2)}/b^3/(1+I*a+I*b*x)^{(1/2)}-1/6*(11*I-18*a-6*I*a^2)*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3-1/3*I*(1-I*a-I*b*x)^{(5/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3-1/2*(11*I-18*a-6*I*a^2)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^3$

Rubi [A]

time = 0.12, antiderivative size = 229, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 91, 81, 52, 55, 633, 221}

$$\frac{(-6ia^2 - 18a + 11i)\sqrt{ia+ibx+1}(-ia-ibx+1)^{3/2}}{6b^3} - \frac{(-6ia^2 - 18a + 11i)\sqrt{ia+ibx+1}\sqrt{-ia-ibx+1}}{2b^3} + \frac{(-6a^2 + 18ia + 11)\sinh^{-1}(a+bx)}{2b^3} - \frac{i\sqrt{ia+ibx+1}(-ia-ibx+1)^{5/2}}{3b^3} + \frac{i(-a+i)^2(-ia-ibx+1)^{5/2}}{b^3\sqrt{ia+ibx+1}}$$

Antiderivative was successfully verified.

[In] `Int[x^2/E^((3*I)*ArcTan[a + b*x]),x]`

[Out] $(I*(I-a)^2*(1-I*a-I*b*x)^{(5/2)})/(b^3*\text{Sqrt}[1+I*a+I*b*x]) - ((11*I-18*a-(6*I)*a^2)*\text{Sqrt}[1-I*a-I*b*x]*\text{Sqrt}[1+I*a+I*b*x])/(2*b^3) - ((11*I-18*a-(6*I)*a^2)*(1-I*a-I*b*x)^{(3/2)}*\text{Sqrt}[1+I*a+I*b*x])/(6*b^3) - ((I/3)*(1-I*a-I*b*x)^{(5/2)}*\text{Sqrt}[1+I*a+I*b*x])/b^3 + ((11+(18*I)*a-6*a^2)*\text{ArcSinh}[a+b*x])/(2*b^3)$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 55

`Int[1/(Sqrt[(a_.) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] := Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]`

Rule 81

`Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)/(d*f*(n + p + 1)), x]`

2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 91

Int[((a_.) + (b_.)*(x_))^(c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(b*c - a*d)^2*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d^2*(d*e - c*f)*(n + 1))), x] - Dist[1/(d^2*(d*e - c*f)*(n + 1)), Int[(c + d*x)^(n + 1)*(e + f*x)^p*Simp[a^2*d^2*f*(n + p + 2) + b^2*c*(d*e*(n + 1) + c*f*(p + 1)) - 2*a*b*d*(d*e*(n + 1) + c*f*(p + 1)) - b^2*d*(d*e - c*f)*(n + 1)*x, x], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && (LtQ[n, -1] || (EqQ[n + p + 3, 0] && NeQ[n, -1] && (SumSimplerQ[n, 1] || !SumSimplerQ[p, 1])))

Rule 221

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2(1-ia-ibx)^{3/2}}{(1+ia+ibx)^{3/2}} dx \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{i \int \frac{(1-ia-ibx)^{3/2}(-i-a)(3+2ia)b-b^2x}{\sqrt{1+ia+ibx}} dx}{b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} - \frac{i(1-ia-ibx)^{5/2}\sqrt{1+ia+ibx}}{3b^3} + \frac{(11+18ia-6a^2)}{3b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{(18a-i(11-6a^2))(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{6b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{(18a-i(11-6a^2))\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11+18ia-6a^2)}{3b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{(18a-i(11-6a^2))\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11+18ia-6a^2)}{3b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{(18a-i(11-6a^2))\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11+18ia-6a^2)}{3b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{(18a-i(11-6a^2))\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11+18ia-6a^2)}{3b^3} \\
&= \frac{i(i-a)^2(1-ia-ibx)^{5/2}}{b^3\sqrt{1+ia+ibx}} + \frac{(18a-i(11-6a^2))\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^3} + \frac{(11+18ia-6a^2)}{3b^3}
\end{aligned}$$

Mathematica [A]

time = 0.24, size = 198, normalized size = 0.86

$$\frac{2ia^4 + a^3(51 + 2ibx) + a^2(-50i + 69bx) + a(51 - 106ibx + 9b^2x^2 + 2ib^3x^3) + i(-52 + 33ibx - 26b^2x^2 + 9ib^3x^3 + 2b^4x^4)}{6b^3\sqrt{1+a^2+2abx+b^2x^2}} + \frac{\sqrt[3]{-1}(11+18ia-6a^2)\sqrt{-ib}\sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{b^{7/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[x^2/E^((3*I)*ArcTan[a + b*x]),x]

[Out] ((2*I)*a^4 + a^3*(51 + (2*I)*b*x) + a^2*(-50*I + 69*b*x) + a*(51 - (106*I)*b*x + 9*b^2*x^2 + (2*I)*b^3*x^3) + I*(-52 + (33*I)*b*x - 26*b^2*x^2 + (9*I)*b^3*x^3 + 2*b^4*x^4))/(6*b^3*sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]) + ((-1)^(1/4)*(11 + (18*I)*a - 6*a^2)*sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*sqrt[b]*sqrt[(-I)*(I + a + b*x)])/sqrt[(-I)*b]])/b^(7/2)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 798 vs. 2(181) = 362.

time = 0.17, size = 799, normalized size = 3.49

method	result
risch	$\frac{i(2b^2x^2 - 2abx + 9ibx + 2a^2 - 27ia - 28)\sqrt{b^2x^2 + 2abx + a^2 + 1}}{6b^3} + \frac{11 \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2 + 2abx + a^2 + 1}\right)}{2b^2\sqrt{b^2}} + \dots$
default	$2i(i-a) \left(-\frac{i\left(\left(x-\frac{i-a}{b}\right)^2 b^2 + 2ib\left(x-\frac{i-a}{b}\right)\right)^{\frac{5}{2}}}{b\left(x-\frac{i-a}{b}\right)^2} + 3ib \left(\frac{\left(\left(x-\frac{i-a}{b}\right)^2 b^2 + 2ib\left(x-\frac{i-a}{b}\right)\right)^{\frac{3}{2}}}{3} + ib \left(\frac{2\left(x-\frac{i-a}{b}\right)b^2 + 2ib}{4b^2} \sqrt{\left(x-\frac{i-a}{b}\right)^2 b^2 + \dots} \right) \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$2*I*(I-a)/b^4*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2)))+I/b^3*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2)))+I*(I-a)^2/b^5*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)-2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2))))$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 624 vs. $2(155) = 310$.
time = 0.48, size = 624, normalized size = 2.72

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="maxima")`

[Out]
$$I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)*a^2/(b^5*x^2 + 2*a*b^4*x + a^2*b^3 - 2*I*b^4*x - 2*I*a*b^3 - b^3) + 2*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)*a/(b^5*x^2 + 2*a*b^4*x + a^2*b^3 - 2*I*b^4*x - 2*I*a*b^3 - b^3) + 2*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)*a/(2*I*b^4*x + 2*I*a*b^3 + 2*b^3) + 6*I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a^2/(I*b^4*x + I*a*b^3 + b^3) - I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)/(b^5*x^2 + 2*a*b^4*x + a^2*b^3 - 2*I*b^4*x - 2*I*a*b^3 - b^3) - 2*I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)/(2*I*b^4*x + 2*I*a*b^3 + 2*b^3) + 12*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*a/(I*b^4*x + I*a*b^3 + b^3) - 3*a$$

$$\begin{aligned} & \frac{2 \operatorname{arcsinh}(bx + a)}{b^3} - \frac{6 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{(b^4 x + Iab^3 + b^3)} - \frac{1}{2} \sqrt{-b^2 x^2 - 2abx - a^2 + 4Ibx + 4Ia + 3} \frac{x}{b^2} \\ & + \frac{9Ia \operatorname{arcsinh}(bx + a)}{b^3} + \frac{1}{3} I (b^2 x^2 + 2abx + a^2 + 1)^{3/2} \frac{1}{b^3} + 3 \sqrt{b^2 x^2 + 2abx + a^2 + 1} \frac{a}{b^3} \\ & - \frac{1}{2} \sqrt{-b^2 x^2 - 2abx - a^2 + 4Ibx + 4Ia + 3} \frac{a}{b^3} + \frac{1}{2} \arcsin(Ibx + Ia + 2) \frac{1}{b^3} \\ & + \frac{6 \operatorname{arcsinh}(bx + a)}{b^3} - \frac{3 \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{b^3} + \sqrt{-b^2 x^2 - 2abx - a^2 + 4Ibx + 4Ia + 3} \frac{1}{b^3} \end{aligned}$$

Fricas [A]

time = 2.99, size = 174, normalized size = 0.76

$$\frac{7i a^4 + 166 a^3 + (7i a^3 + 159 a^2 - 249i a - 96)bx - 408i a^2 + 12(6a^3 + (6a^2 - 18i a - 11)bx - 24i a^2 - 29a + 11i) \log(-bx - a + \sqrt{b^2 x^2 + 2abx + a^2 + 1}) - 4(-2i b^3 x^3 + 7b^2 x^2 - 2i a^3 - (16a - 19i)bx - 53a^2 + 103i a + 52) \sqrt{b^2 x^2 + 2abx + a^2 + 1} - 345a + 96i}{24(b^4 x + (a - i)b^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{24} (7Ia^4 + 166a^3 + (7Ia^3 + 159a^2 - 249Ia - 96)bx - 408Ia^2 + 12(6a^3 + (6a^2 - 18Ia - 11)bx - 24Ia^2 - 29a + 11I) \log(-bx - a + \sqrt{b^2 x^2 + 2abx + a^2 + 1}) - 4(-2Ib^3 x^3 + 7b^2 x^2 - 2Ia^3 - (16a - 19I)bx - 53a^2 + 103Ia + 52) \sqrt{b^2 x^2 + 2abx + a^2 + 1} - 345a + 96I) / (b^4 x + (a - I)b^3)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\int \frac{x^2 \sqrt{a^2 + 2abx + b^2 x^2 + 1}}{a^3 + 3a^2 bx - 3a^2 + 3ab^2 x^2 - 6iabx - 3a + b^3 x^3 - 3i b^2 x - 3ix + 1} dx + \int \frac{a^2 x^2 \sqrt{a^2 + 2abx + b^2 x^2 + 1}}{a^3 + 3a^2 bx - 3a^2 + 3ab^2 x^2 - 6iabx - 3a + b^3 x^3 - 3i b^2 x - 3ix + 1} dx + \int \frac{b^2 x^4 \sqrt{a^2 + 2abx + b^2 x^2 + 1}}{a^3 + 3a^2 bx - 3a^2 + 3ab^2 x^2 - 6iabx - 3a + b^3 x^3 - 3i b^2 x - 3ix + 1} dx + \int \frac{2abx^2 \sqrt{a^2 + 2abx + b^2 x^2 + 1}}{a^3 + 3a^2 bx - 3a^2 + 3ab^2 x^2 - 6iabx - 3a + b^3 x^3 - 3i b^2 x - 3ix + 1} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*(b*x+a))**3*(1+(b*x+a)**2)**(3/2),x)

[Out] $I * (\text{Integral}(x^{**2} \sqrt{a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1} / (a^{**3} + 3*a^{**2}*b*x - 3*I*a^{**2} + 3*a*b^{**2}*x^{**2} - 6*I*a*b*x - 3*a + b^{**3}*x^{**3} - 3*I*b^{**2}*x^{**2} - 3*b*x + I), x) + \text{Integral}(a^{**2}*x^{**2} \sqrt{a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1} / (a^{**3} + 3*a^{**2}*b*x - 3*I*a^{**2} + 3*a*b^{**2}*x^{**2} - 6*I*a*b*x - 3*a + b^{**3}*x^{**3} - 3*I*b^{**2}*x^{**2} - 3*b*x + I), x) + \text{Integral}(b^{**2}*x^{**4} \sqrt{a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1} / (a^{**3} + 3*a^{**2}*b*x - 3*I*a^{**2} + 3*a*b^{**2}*x^{**2} - 6*I*a*b*x - 3*a + b^{**3}*x^{**3} - 3*I*b^{**2}*x^{**2} - 3*b*x + I), x) + \text{Integral}(2*a*b*x^{**3} \sqrt{a^{**2} + 2*a*b*x + b^{**2}*x^{**2} + 1} / (a^{**3} + 3*a^{**2}*b*x - 3*I*a^{**2} + 3*a*b^{**2}*x^{**2} - 6*I*a*b*x - 3*a + b^{**3}*x^{**3} - 3*I*b^{**2}*x^{**2} - 3*b*x + I), x))$

Giac [A]

time = 0.46, size = 243, normalized size = 1.06

$$-\frac{1}{6} \sqrt{(bx+a)^2+1} \left(x \left(\frac{2ix}{b} + \frac{2iab^2+9b^3}{-6a} \right) + \frac{-2ia^2b-27ab^3+28b^4}{6} \right) + \frac{(6a^2-18a-11) \log \left(\frac{3 \left(|x| - \sqrt{(bx+a)^2+1} \right) ab + ia^2b + i \left(|x| - \sqrt{(bx+a)^2+1} \right) |x| - 3 \left(-ix|x| + i \sqrt{(bx+a)^2+1} \right) a^2 |x| + 2 \left(|x| - \sqrt{(bx+a)^2+1} \right) b + 2a^2b + 4 \left(|x| - \sqrt{(bx+a)^2+1} \right) a |x| - iab + \left(-ix|x| + i \sqrt{(bx+a)^2+1} \right) |x| \right)}{6|b|^3|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="giac")

[Out]
$$-1/6*\sqrt{(b*x + a)^2 + 1}*(x*(-2*I*x/b + (2*I*a*b^6 + 9*b^6)/b^8) + (-2*I*a^2*b^5 - 27*a*b^5 + 28*I*b^5)/b^8) + 1/6*(6*a^2 - 18*I*a - 11)*\log(3*I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*a*b + I*a^3*b + I*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^3*\text{abs}(b) - 3*(-I*x*\text{abs}(b) + I*\sqrt{(b*x + a)^2 + 1})*a^2*\text{abs}(b) + 2*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})^2*b + 2*a^2*b + 4*(x*\text{abs}(b) - \sqrt{(b*x + a)^2 + 1})*a*\text{abs}(b) - I*a*b + (-I*x*\text{abs}(b) + I*\sqrt{(b*x + a)^2 + 1})*\text{abs}(b))/(b^2*\text{abs}(b))$$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2 ((a + bx)^2 + 1)^{3/2}}{(1 + a li + b x li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3,x)

[Out] int((x^2*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3, x)

3.210 $\int e^{-3i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=163

$$\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2\sqrt{1+ia+ibx}} - \frac{3(3+2ia)\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{2b^2} - \frac{(3+2ia)(1-ia-ibx)^{3/2}\sqrt{1+ia+ibx}}{2b^2}$$

[Out] $-3/2*(3*I-2*a)*\text{arcsinh}(b*x+a)/b^2-(1+I*a)*(1-I*a-I*b*x)^{(5/2)}/b^2/(1+I*a+I*b*x)^{(1/2)}-1/2*(3+2*I*a)*(1-I*a-I*b*x)^{(3/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2-3/2*(3+2*I*a)*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b^2$

Rubi [A]

time = 0.08, antiderivative size = 163, normalized size of antiderivative = 1.00, number of steps used = 7, number of rules used = 6, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.429$, Rules used = {5203, 79, 52, 55, 633, 221}

$$-\frac{(1+ia)(-ia-ibx+1)^{5/2}}{b^2\sqrt{ia+ibx+1}} - \frac{(3+2ia)\sqrt{ia+ibx+1}(-ia-ibx+1)^{3/2}}{2b^2} - \frac{3(3+2ia)\sqrt{ia+ibx+1}\sqrt{-ia-ibx+1}}{2b^2} - \frac{3(-2a+3i)\sinh^{-1}(a+bx)}{2b^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x/E^{((3*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $-(((1 + I*a)*(1 - I*a - I*b*x)^{(5/2)})/(b^2*\text{Sqrt}[1 + I*a + I*b*x])) - (3*(3 + (2*I)*a)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/(2*b^2) - ((3 + (2*I)*a)*(1 - I*a - I*b*x)^{(3/2)}*\text{Sqrt}[1 + I*a + I*b*x])/(2*b^2) - (3*(3*I - 2*a)*\text{ArcSinh}[a + b*x])/(2*b^2)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_. + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^n/(b*(m + n + 1))}, x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))], \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{GtQ}[n, 0] \&\& \text{NeQ}[m + n + 1, 0] \&\& !(\text{IGtQ}[m, 0] \&\& (!\text{IntegerQ}[n] || (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0]))) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 55

$\text{Int}[1/(\text{Sqrt}[(a_. + (b_.)*(x_.)]*\text{Sqrt}[(c_. + (d_.)*(x_.))], x_Symbol] :> \text{Int}[1/\text{Sqrt}[a*c - b*(a - c)*x - b^2*x^2], x] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{EqQ}[b + d, 0] \&\& \text{GtQ}[a + c, 0]$

Rule 79

$\text{Int}[(a_. + (b_.)*(x_.))*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)*((e + f*x)^{(p + 1)}/(f*(p + 1)*(c*f - d*e))}, x] - \text{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c$

```
*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^(p + 1), x]
, x] /; FreeQ[{a, b, c, d, e, f, n}, x] && LtQ[p, -1] && (!LtQ[n, -1] || IntegerQ[p] || !(IntegerQ[n] || !(EqQ[e, 0] || !(EqQ[c, 0] || LtQ[p, n]))))
```

Rule 221

```
Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] :> Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]
```

Rule 633

```
Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] :> Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{-3i \tan^{-1}(a+bx)} x dx &= \int \frac{x(1-ia-ibx)^{3/2}}{(1+ia+ibx)^{3/2}} dx \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{(3i-2a) \int \frac{(1-ia-ibx)^{3/2}}{\sqrt{1+ia+ibx}} dx}{b} \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{(3+2ia)(1-ia-ibx)^{3/2} \sqrt{1+ia+ibx}}{2b^2} - \frac{(3(3i-2a))}{b^2} \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{3(3+2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} - \frac{(3+2ia)}{b^2} \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{3(3+2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} - \frac{(3+2ia)}{b^2} \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{3(3+2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} - \frac{(3+2ia)}{b^2} \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{3(3+2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} - \frac{(3+2ia)}{b^2} \\
&= -\frac{(1+ia)(1-ia-ibx)^{5/2}}{b^2 \sqrt{1+ia+ibx}} - \frac{3(3+2ia) \sqrt{1-ia-ibx} \sqrt{1+ia+ibx}}{2b^2} - \frac{(3+2ia)}{b^2}
\end{aligned}$$

Mathematica [A]

time = 0.23, size = 157, normalized size = 0.96

$$\frac{i(14i - a^3 + 9bx + 6ib^2x^2 + b^3x^3 + a^2(14i - bx) + a(-1 + 20ibx + b^2x^2))}{2b^2 \sqrt{1 + a^2 + 2abx + b^2x^2}} + \frac{3\sqrt{-1}(-3i + 2a)\sqrt{-ib} \sinh^{-1}\left(\frac{(\frac{1}{2} + \frac{i}{2})\sqrt{b} \sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{b^{5/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[x/E^((3*I)*ArcTan[a + b*x]),x]

[Out] ((I/2)*(14*I - a^3 + 9*b*x + (6*I)*b^2*x^2 + b^3*x^3 + a^2*(14*I - b*x) + a*(-1 + (20*I)*b*x + b^2*x^2)))/(b^2*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2]) + (3*(-1)^(1/4)*(-3*I + 2*a)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])]/Sqrt[(-I)*b])/b^(5/2)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 594 vs. 2(129) = 258.

time = 0.16, size = 595, normalized size = 3.65

method	result
--------	--------

risch	$\frac{i(-bx+a-6i)\sqrt{b^2x^2+2abx+a^2+1}}{2b^2} - \frac{9i \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{2b\sqrt{b^2}} + \frac{3a \ln\left(\frac{b^2x+ab}{\sqrt{b^2}} + \sqrt{b^2x^2+2abx+a^2+1}\right)}{b^3}$
default	$i \left(-\frac{i \left(\left(x - \frac{i-a}{b} \right)^2 b^2 + 2ib \left(x - \frac{i-a}{b} \right) \right)^{\frac{5}{2}}}{b \left(x - \frac{i-a}{b} \right)^2} + 3ib \left(\frac{\left(\left(x - \frac{i-a}{b} \right)^2 b^2 + 2ib \left(x - \frac{i-a}{b} \right) \right)^{\frac{3}{2}}}{3} + ib \left(\frac{\left(\left(x - \frac{i-a}{b} \right)^2 b^2 + 2ib \left(x - \frac{i-a}{b} \right) \right)^{\frac{1}{2}}}{4b^2} \right) \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x,method=_RETURNVERBOSE)`

[Out]
$$\frac{I}{b^3} \left(-\frac{I}{b} \left(\frac{x-(I-a)}{b} \right)^2 \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{5}{2}} + 3Ib \left(\frac{1}{3} \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{3}{2}} + Ib \left(\frac{1}{4} \left(2 \left(\frac{x-(I-a)}{b} \right) b^2 + 2Ib \right) / b^2 \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{1}{2}} + \frac{1}{2} \ln \left(\frac{Ib + \left(\frac{x-(I-a)}{b} \right) b^2}{\left(b^2 \right)^{\frac{1}{2}} + \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right)} \right)^{\frac{1}{2}} \right) + I \left(\frac{I-a}{b} \right)^4 \left(\frac{I}{b} \left(\frac{x-(I-a)}{b} \right)^3 \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{5}{2}} - 2Ib \left(\frac{x-(I-a)}{b} \right)^2 \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{5}{2}} + 3Ib \left(\frac{1}{3} \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{3}{2}} + Ib \left(\frac{1}{4} \left(2 \left(\frac{x-(I-a)}{b} \right) b^2 + 2Ib \right) / b^2 \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right) \right)^{\frac{1}{2}} + \frac{1}{2} \ln \left(\frac{Ib + \left(\frac{x-(I-a)}{b} \right) b^2}{\left(b^2 \right)^{\frac{1}{2}} + \left(\frac{x-(I-a)}{b} \right)^2 b^2 + 2Ib \left(\frac{x-(I-a)}{b} \right)} \right)^{\frac{1}{2}} \right)$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 293 vs. $2(113) = 226$.

time = 0.47, size = 293, normalized size = 1.80

$$\frac{i(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}a}{b^2x^2+2ab^2x+a^2b^2-2iab^2-b^2} - \frac{(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}}{b^2x^2+2ab^2x+a^2b^2-2iab^2-b^2} - \frac{(b^2x^2+2abx+a^2+1)^{\frac{3}{2}}}{2ib^2x+2iab^2+2b^2} - \frac{6i\sqrt{b^2x^2+2abx+a^2+1}a}{ib^2x+iab^2+b^2} + \frac{3a \operatorname{arsinh}(bx+a)}{b^2} - \frac{6\sqrt{b^2x^2+2abx+a^2+1}}{ib^2x+iab^2+b^2} - \frac{9i \operatorname{arsinh}(bx+a)}{2b^2} - \frac{3\sqrt{b^2x^2+2abx+a^2+1}}{2b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="maxima")`

[Out]
$$-I(b^2x^2+2a*b*x+a^2+1)^{\frac{3}{2}}*a/(b^4*x^2+2*a*b^3*x+a^2*b^2-2*I*b^3*x-2*I*a*b^2-b^2) - (b^2*x^2+2*a*b*x+a^2+1)^{\frac{3}{2}}/(b^4*x^2+2*a*b^3*x+a^2*b^2-2*I*b^3*x-2*I*a*b^2-b^2) - (b^2*x^2+2*a*b*x+a^2+1)^{\frac{3}{2}}/(2*I*b^3*x+2*I*a*b^2+2*b^2) - 6*I*\operatorname{sqrt}(b^2*x^2+2*a*b*x+a^2+1)*a/(I*b^3*x+I*a*b^2+b^2) + 3*a*\operatorname{arcsinh}(b*x+a)/b^2 - 6*\operatorname{sqrt}(b^2*x^2+2*a*b*x+a^2+1)/(I*b^3*x+I*a*b^2+b^2) - 9/2*I*\operatorname{arcsinh}(b*x+a)/b^2 - 3/2*\operatorname{sqrt}(b^2*x^2+2*a*b*x+a^2+1)/b^2$$

Fricas [A]

time = 1.90, size = 136, normalized size = 0.83

$$\frac{-3ia^3 + (-3ia^2 - 44a + 32i)bx - 47a^2 - 12((2a - 3i)bx + 2a^2 - 5ia - 3) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 4\sqrt{b^2x^2 + 2abx + a^2 + 1}(-ib^2x^2 + ia^2 + 5bx + 15a - 14i) + 76ia + 32}{8(b^3x + (a - i)b^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="fricas")

[Out] 1/8*(-3*I*a^3 + (-3*I*a^2 - 44*a + 32*I)*b*x - 47*a^2 - 12*((2*a - 3*I)*b*x + 2*a^2 - 5*I*a - 3)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*b^2*x^2 + I*a^2 + 5*b*x + 15*a - 14*I) + 76*I*a + 32)/(b^3*x + (a - I)*b^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\int \frac{x\sqrt{a^2+2bx+b^2x^2+1}}{a^3+3a^2bx-3a^2+3ab^2x^2-6iabx-3a+b^3x^3-3ib^2x-3bx+1} dx + \int \frac{a^2x\sqrt{a^2+2bx+b^2x^2+1}}{a^3+3a^2bx-3a^2+3ab^2x^2-6iabx-3a+b^3x^3-3ib^2x-3bx+1} dx + \int \frac{b^2x^2\sqrt{a^2+2bx+b^2x^2+1}}{a^3+3a^2bx-3a^2+3ab^2x^2-6iabx-3a+b^3x^3-3ib^2x-3bx+1} dx + \int \frac{2abx^2\sqrt{a^2+2bx+b^2x^2+1}}{a^3+3a^2bx-3a^2+3ab^2x^2-6iabx-3a+b^3x^3-3ib^2x-3bx+1} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a)**3*(1+(b*x+a)**2)**(3/2),x)

[Out] I*(Integral(x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x) + Integral(a**2*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x) + Integral(b**2*x**3*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x) + Integral(2*a*b*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x))

Giac [A]

time = 0.46, size = 210, normalized size = 1.29

$$-\frac{1}{2}\sqrt{(bx+a)^2+1}\left(\frac{ix}{b}-\frac{-iab^2-6b^2}{b^4}\right)-\frac{(2a-3i)\log\left(3\left(|x|-\sqrt{(bx+a)^2+1}\right)ab+a^2b+\left(|x|-\sqrt{(bx+a)^2+1}\right)^3\right)+3\left(|x|-\sqrt{(bx+a)^2+1}\right)a^2|b|-2i\left(|x|-\sqrt{(bx+a)^2+1}\right)b-2ia^2b+4\left(-ix|b|+i\sqrt{(bx+a)^2+1}\right)a|b|-ab-\left(|x|-\sqrt{(bx+a)^2+1}\right)|b|}{2|b|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="giac")

[Out] -1/2*sqrt((b*x + a)^2 + 1)*(-I*x/b - (-I*a*b^2 - 6*b^2)/b^4) - 1/2*(2*a - 3*I)*log(3*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b + a^3*b + (x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) + 3*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a^2*abs(b) - 2*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b - 2*I*a^2*b + 4*(-I*x*abs(b) + I*sqrt((b*x + a)^2 + 1))*a*abs(b) - a*b - (x*abs(b) - sqrt((b*x + a)^2 + 1))*abs(b))/(b*abs(b))

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x \left((a + bx)^2 + 1 \right)^{3/2}}{(1 + a li + bx li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3,x)
```

```
[Out] int((x*((a + b*x)^2 + 1)^(3/2))/(a*1i + b*x*1i + 1)^3, x)
```

3.211 $\int e^{-3i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=94

$$\frac{2i(1-ia-ibx)^{3/2}}{b\sqrt{1+ia+ibx}} + \frac{3i\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}}{b} - \frac{3\sinh^{-1}(a+bx)}{b}$$

[Out] $-3*\text{arcsinh}(b*x+a)/b+2*I*(1-I*a-I*b*x)^{(3/2)}/b/(1+I*a+I*b*x)^{(1/2)}+3*I*(1-I*a-I*b*x)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/b$

Rubi [A]

time = 0.03, antiderivative size = 94, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5201, 49, 52, 55, 633, 221}

$$\frac{2i(-ia-ibx+1)^{3/2}}{b\sqrt{ia+ibx+1}} + \frac{3i\sqrt{ia+ibx+1}\sqrt{-ia-ibx+1}}{b} - \frac{3\sinh^{-1}(a+bx)}{b}$$

Antiderivative was successfully verified.

[In] Int[E^((-3*I)*ArcTan[a + b*x]),x]

[Out] $((2*I)*(1 - I*a - I*b*x)^{(3/2)})/(b*\text{Sqrt}[1 + I*a + I*b*x]) + ((3*I)*\text{Sqrt}[1 - I*a - I*b*x]*\text{Sqrt}[1 + I*a + I*b*x])/b - (3*\text{ArcSinh}[a + b*x])/b$

Rule 49

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(ILeQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 55

Int[1/(Sqrt[(a_) + (b_.)*(x_)]*Sqrt[(c_.) + (d_.)*(x_)]), x_Symbol] :> Int[1/Sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b

+ d, 0] && GtQ[a + c, 0]

Rule 221

Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 633

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b + 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

Rule 5201

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{-3i \tan^{-1}(a+bx)} dx &= \int \frac{(1 - ia - ibx)^{3/2}}{(1 + ia + ibx)^{3/2}} dx \\
 &= \frac{2i(1 - ia - ibx)^{3/2}}{b\sqrt{1 + ia + ibx}} - 3 \int \frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} dx \\
 &= \frac{2i(1 - ia - ibx)^{3/2}}{b\sqrt{1 + ia + ibx}} + \frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - 3 \int \frac{1}{\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}} dx \\
 &= \frac{2i(1 - ia - ibx)^{3/2}}{b\sqrt{1 + ia + ibx}} + \frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - 3 \int \frac{1}{\sqrt{(1 - ia)(1 + ia) + 2ibx}} dx \\
 &= \frac{2i(1 - ia - ibx)^{3/2}}{b\sqrt{1 + ia + ibx}} + \frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - \frac{3 \operatorname{Subst} \left(\int \frac{1}{\sqrt{1 + \frac{x^2}{4b^2}}} dx, \right)}{2b^2} \\
 &= \frac{2i(1 - ia - ibx)^{3/2}}{b\sqrt{1 + ia + ibx}} + \frac{3i\sqrt{1 - ia - ibx} \sqrt{1 + ia + ibx}}{b} - \frac{3 \sinh^{-1}(a + bx)}{b}
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 45, normalized size = 0.48

$$\frac{\sqrt{1 + (a + bx)^2} \left(i + \frac{4}{-i + a + bx} \right)}{b} - \frac{3 \sinh^{-1}(a + bx)}{b}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((-3*I)*ArcTan[a + b*x]),x]

[Out] (Sqrt[1 + (a + b*x)^2]*(I + 4/(-I + a + b*x)))/b - (3*ArcSinh[a + b*x])/b

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 326 vs. 2(76) = 152.

time = 0.14, size = 327, normalized size = 3.48

method	result
risch	$\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{b} - \frac{3 \ln\left(\frac{b^2x+ab + \sqrt{b^2x^2 + 2abx + a^2 + 1}}{\sqrt{b^2}}\right)}{\sqrt{b^2}} + \frac{4\sqrt{\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib\left(x - \frac{i-a}{b}\right)}}{b^2\left(x - \frac{i-a}{b}\right)}$
default	$i\left(\frac{i\left(\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib\left(x - \frac{i-a}{b}\right)\right)^{\frac{5}{2}}}{b\left(x - \frac{i-a}{b}\right)^3} - 2ib\left(-\frac{i\left(\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib\left(x - \frac{i-a}{b}\right)\right)^{\frac{5}{2}}}{b\left(x - \frac{i-a}{b}\right)^2} + 3ib\left(\frac{\left(\left(x - \frac{i-a}{b}\right)^2 b^2 + 2ib\left(x - \frac{i-a}{b}\right)\right)^{\frac{3}{2}}}{3} + ib\left(\frac{2\left(x - \frac{i-a}{b}\right)b^2 + \dots}{b^3}\right)\right)\right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x,method=_RETURNVERBOSE)

[Out] I/b^3*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)-2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^(1/2)))

Maxima [A]

time = 0.46, size = 103, normalized size = 1.10

$$\frac{i(b^2x^2 + 2abx + a^2 + 1)^{\frac{3}{2}}}{b^3x^2 + 2ab^2x + a^2b - 2ib^2x - 2iab - b} - \frac{3 \operatorname{arsinh}(bx + a)}{b} + \frac{6i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{ib^2x + iab + b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="maxima")

[Out] I*(b^2*x^2 + 2*a*b*x + a^2 + 1)^(3/2)/(b^3*x^2 + 2*a*b^2*x + a^2*b - 2*I*b^2*x - 2*I*a*b - b) - 3*arcsinh(b*x + a)/b + 6*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(I*b^2*x + I*a*b + b)

Fricas [A]

time = 2.16, size = 99, normalized size = 1.05

$$\frac{(ia + 8)bx + ia^2 + 6(bx + a - i) \log(-bx - a + \sqrt{b^2x^2 + 2abx + a^2 + 1}) - 2\sqrt{b^2x^2 + 2abx + a^2 + 1}(-ibx - ia - 5) + 9a - 8i}{2(b^2x + (a - i)b)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="fricas")

[Out] 1/2*((I*a + 8)*b*x + I*a^2 + 6*(b*x + a - I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 2*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-I*b*x - I*a - 5) + 9*a - 8*I)/(b^2*x + (a - I)*b)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3 + 3a^2bx - 3a^2 + 3ab^2x^2 - 6abx - 3a + b^3x^3 - 3b^2x^2 - 3bx + I} dx + \int \frac{a^2\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3 + 3a^2bx - 3a^2 + 3ab^2x^2 - 6abx - 3a + b^3x^3 - 3b^2x^2 - 3bx + I} dx + \int \frac{b^2x^2\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3 + 3a^2bx - 3a^2 + 3ab^2x^2 - 6abx - 3a + b^3x^3 - 3b^2x^2 - 3bx + I} dx + \int \frac{2abx\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3 + 3a^2bx - 3a^2 + 3ab^2x^2 - 6abx - 3a + b^3x^3 - 3b^2x^2 - 3bx + I} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x)

[Out] I*(Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x) + Integral(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x) + Integral(b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x) + Integral(2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3 + 3*a**2*b*x - 3*I*a**2 + 3*a*b**2*x**2 - 6*I*a*b*x - 3*a + b**3*x**3 - 3*I*b**2*x**2 - 3*b*x + I), x))

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 182 vs. 2(66) = 132.

time = 0.49, size = 182, normalized size = 1.94

$$\frac{\log\left(3i\left(x|b| - \sqrt{(bx+a)^2 + 1}\right)^2 ab + i a^2 b + i\left(x|b| - \sqrt{(bx+a)^2 + 1}\right)^3 |b| - 3\left(-ix|b| + i\sqrt{(bx+a)^2 + 1}\right) a^2 |b| + 2\left(x|b| - \sqrt{(bx+a)^2 + 1}\right)^2 b + 2a^2 b + 4\left(x|b| - \sqrt{(bx+a)^2 + 1}\right) a |b| - i ab + \left(-ix|b| + i\sqrt{(bx+a)^2 + 1}\right) |b|\right)}{i\sqrt{(bx+a)^2 + 1}} \Big|_x + \frac{i\sqrt{(bx+a)^2 + 1}}{b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2),x, algorithm="giac")

[Out] log(3*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b + I*a^3*b + I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) - 3*(-I*x*abs(b) + I*sqrt((b*x + a)^2 + 1))*a^2*abs(b) + 2*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b + 2*a^2*b + 4*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a*abs(b) - I*a*b + (-I*x*abs(b) + I*sqrt((b*x + a)^2 + 1))*abs(b))/abs(b) + I*sqrt((b*x + a)^2 + 1)/b

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{((a + bx)^2 + 1)^{3/2}}{(1 + a li + bx li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a + b*x)^2 + 1)^(3/2)/(a*1i + b*x*1i + 1)^3,x)
```

```
[Out] int(((a + b*x)^2 + 1)^(3/2)/(a*1i + b*x*1i + 1)^3, x)
```


$$3.212 \quad \int \frac{e^{-3i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=134

$$\frac{4\sqrt{1-ia-ibx}}{(1+ia)\sqrt{1+ia+ibx}} + i \sinh^{-1}(a+bx) - \frac{2(i+a)^{3/2} \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i-a)^{3/2}}$$

[Out] I*arcsinh(b*x+a)-2*(I+a)^(3/2)*arctanh((I+a)^(1/2)*(1+I*a+I*b*x)^(1/2)/(I-a)^(1/2)/(1-I*a-I*b*x)^(1/2))/(I-a)^(3/2)+4*(1-I*a-I*b*x)^(1/2)/(1+I*a)/(1+I*a+I*b*x)^(1/2)

Rubi [A]

time = 0.08, antiderivative size = 134, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 8, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.500$, Rules used = {5203, 100, 163, 55, 633, 221, 95, 214}

$$\frac{4\sqrt{-ia-ibx+1}}{(1+ia)\sqrt{ia+ibx+1}} + i \sinh^{-1}(a+bx) - \frac{2(a+i)^{3/2} \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a + b*x]))*x], x]

[Out] (4*sqrt[1 - I*a - I*b*x])/((1 + I*a)*sqrt[1 + I*a + I*b*x]) + I*ArcSinh[a + b*x] - (2*(I + a)^(3/2)*ArcTanh[(sqrt[I + a]*sqrt[1 + I*a + I*b*x])/(sqrt[I - a]*sqrt[1 - I*a - I*b*x])])/(I - a)^(3/2)

Rule 55

Int[1/(sqrt[(a_) + (b_.)*(x_)]*sqrt[(c_) + (d_.)*(x_)]), x_Symbol] := Int[1/sqrt[a*c - b*(a - c)*x - b^2*x^2], x] /; FreeQ[{a, b, c, d}, x] && EqQ[b + d, 0] && GtQ[a + c, 0]

Rule 95

Int[(((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_))/((e_) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q], x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)

```

*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m
+ 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(
n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*
(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a,
b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2
*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])

```

Rule 163

```

Int[(((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p_))*((g_.) + (h_.)*(x_
)))/((a_.) + (b_.)*(x_)), x_Symbol] := Dist[h/b, Int[(c + d*x)^n*(e + f*x)^
p, x], x] + Dist[(b*g - a*h)/b, Int[(c + d*x)^n*((e + f*x)^p/(a + b*x)), x]
, x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x]

```

Rule 214

```

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

```

Rule 221

```

Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt
[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

```

Rule 633

```

Int[((a_.) + (b_.)*(x_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Dist[1/(2*c*(-4*
(c/(b^2 - 4*a*c)))^p), Subst[Int[Simp[1 - x^2/(b^2 - 4*a*c), x]^p, x], x, b
+ 2*c*x], x] /; FreeQ[{a, b, c, p}, x] && GtQ[4*a - b^2/c, 0]

```

Rule 5203

```

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_)^(m_.),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{(1-ia-ibx)^{3/2}}{x(1+ia+ibx)^{3/2}} dx \\
&= \frac{4\sqrt{1-ia-ibx}}{(1+ia)\sqrt{1+ia+ibx}} + \frac{2 \int \frac{-\frac{1}{2}i(i+a)^2b - \frac{1}{2}(1+ia)b^2x}{x\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx}{(i-a)b} \\
&= \frac{4\sqrt{1-ia-ibx}}{(1+ia)\sqrt{1+ia+ibx}} - \frac{(i+a)^2 \int \frac{1}{x\sqrt{1-ia-ibx}\sqrt{1+ia+ibx}} dx}{1+ia} + (ib) \int \frac{1}{\sqrt{1-ia-ibx}} dx \\
&= \frac{4\sqrt{1-ia-ibx}}{(1+ia)\sqrt{1+ia+ibx}} - \frac{(2(i+a)^2) \text{Subst}\left(\int \frac{1}{-1-ia-(-1+ia)x^2} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}}\right)}{1+ia} \\
&= \frac{4\sqrt{1-ia-ibx}}{(1+ia)\sqrt{1+ia+ibx}} - \frac{2(i+a)^{3/2} \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i-a)^{3/2}} + i \text{Subst}\left(\int \frac{1}{\sqrt{1-ia-ibx}} dx, x, \frac{\sqrt{1+ia+ibx}}{\sqrt{1-ia-ibx}}\right) \\
&= \frac{4\sqrt{1-ia-ibx}}{(1+ia)\sqrt{1+ia+ibx}} + i \sinh^{-1}(a+bx) - \frac{2(i+a)^{3/2} \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i-a)^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.46, size = 189, normalized size = 1.41

$$\frac{2(-1)^{3/4}\sqrt{-ib} \sinh^{-1}\left(\frac{(\frac{1}{2}+\frac{i}{2})\sqrt{b}\sqrt{-i(i+a+bx)}}{\sqrt{-ib}}\right)}{\sqrt{b}} + \frac{2\left(-\frac{2\sqrt{1+a^2+2abx+b^2x^2}}{-i+a+bx} + \frac{\sqrt{-1+ia} \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}{\sqrt{-1-ia}}\right)}{-i+a}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((3*I)*ArcTan[a + b*x]))*x, x]

[Out] (2*(-1)^(3/4)*Sqrt[(-I)*b]*ArcSinh[((1/2 + I/2)*Sqrt[b]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[(-I)*b]])/Sqrt[b] + (2*((-2*Sqrt[1 + a^2 + 2*a*b*x + b^2*x^2])/(-I + a + b*x) + (Sqrt[-1 + I*a]*(I + a)*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)])/Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]])/Sqrt[-1 - I*a]))/(-I + a)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1066 vs. 2(104) = 208.

time = 0.12, size = 1067, normalized size = 7.96

method	result
default	$i \left(\frac{\left(\left(x - \frac{i-a}{b} \right)^2 b^2 + 2ib \left(x - \frac{i-a}{b} \right) \right)^{\frac{5}{2}}}{b \left(x - \frac{i-a}{b} \right)^2} + 3ib \left(\frac{\left(\left(x - \frac{i-a}{b} \right)^2 b^2 + 2ib \left(x - \frac{i-a}{b} \right) \right)^{\frac{3}{2}}}{3} + ib \left(\frac{2 \left(x - \frac{i-a}{b} \right) b^2 + 2ib}{4b^2} \sqrt{\left(x - \frac{i-a}{b} \right)^2 b^2 + 2ib} \right) \right) - \frac{}{(i-a)^2 b}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x,x,method=_RETURNVERBOSE)

[Out]
$$-I/(I-a)^2/b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(1/2)))+I/(I-a)^3*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(1/2)))+I/(I-a)/b^2*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)-2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(1/2))))-I/(I-a)^3*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x,x, algorithm="maxima")

[Out] integrate(((b*x + a)^2 + 1)^(3/2)/((I*b*x + I*a + 1)^3*x), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 356 vs. 2(90) = 180.

time = 1.90, size = 356, normalized size = 2.66

$$\frac{((a-1)bx+a^2-2(a-1)\sqrt{\frac{a^2+3a^2-3a-1}{a^2-3a^2-3a+1}} \log\left(\frac{(a+1)bx+\sqrt{b^2+2ab+a^2+1} \operatorname{arctan}\left(\frac{a^2+3a^2-3a-1}{a^2-3a^2-3a+1}\right)}{a^2-3a^2-3a+1}\right) - ((a-1)bx+a^2-2(a-1)\sqrt{\frac{a^2+3a^2-3a-1}{a^2-3a^2-3a+1}} \log\left(\frac{(a+1)bx+\sqrt{b^2+2ab+a^2+1} \operatorname{arctan}\left(\frac{a^2+3a^2-3a-1}{a^2-3a^2-3a+1}\right)}{a^2-3a^2-3a+1}\right) - 4bx - ((a+1)bx+ia^2+2a-i) \log(-bx-a+\sqrt{b^2+2ab+a^2+1}) - 4a-4\sqrt{b^2+2ab+a^2+1}+4i}{(a-1)bx+a^2-2(a-1)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x,x, algorithm="fricas")

[Out] (((a - I)*b*x + a^2 - 2*I*a - 1)*sqrt(-(a^3 + 3*I*a^2 - 3*a - I)/(a^3 - 3*I*a^2 - 3*a + I))*log(-((a + I)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a + I) - (I*a^2 + 2*a - I)*sqrt(-(a^3 + 3*I*a^2 - 3*a - I)/(a^3 - 3*I*a^2 - 3*a + I))))/(a + I) - ((a - I)*b*x + a^2 - 2*I*a - 1)*sqrt(-(a^3 + 3*I*a^2 - 3*a - I)/(a^3 - 3*I*a^2 - 3*a + I))*log(-((a + I)*b*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(a + I) - (-I*a^2 - 2*a + I)*sqrt(-(a^3 + 3*I*a^2 - 3*a - I)/(a^3 - 3*I*a^2 - 3*a + I))))/(a + I) - 4*b*x - ((I*a + 1)*b*x + I*a^2 + 2*a - I)*log(-b*x - a + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)) - 4*a - 4*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1) + 4*I)/((a - I)*b*x + a^2 - 2*I*a - 1)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^2x + 3a^2bx^2 - 3a^2x + 3a^2b^2x^2 - 6abx^2 - 3ax + b^2x^2 - 3bx^2 + 1x} dx + \int \frac{a^2\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^2x + 3a^2bx^2 - 3a^2x + 3a^2b^2x^2 - 6abx^2 - 3ax + b^2x^2 - 3bx^2 + 1x} dx + \int \frac{b^2\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^2x + 3a^2bx^2 - 3a^2x + 3a^2b^2x^2 - 6abx^2 - 3ax + b^2x^2 - 3bx^2 + 1x} dx + \int \frac{2abx\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^2x + 3a^2bx^2 - 3a^2x + 3a^2b^2x^2 - 6abx^2 - 3ax + b^2x^2 - 3bx^2 + 1x} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a)**3*(1+(b*x+a)**2)**(3/2)/x,x)

[Out] I*(Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x + 3*a**2*b*x**2 - 3*I*a**2*x + 3*a*b**2*x**3 - 6*I*a*b*x**2 - 3*a*x + b**3*x**4 - 3*I*b**2*x**3 - 3*b*x**2 + I*x), x) + Integral(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x + 3*a**2*b*x**2 - 3*I*a**2*x + 3*a*b**2*x**3 - 6*I*a*b*x**2 - 3*a*x + b**3*x**4 - 3*I*b**2*x**3 - 3*b*x**2 + I*x), x) + Integral(b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x + 3*a**2*b*x**2 - 3*I*a**2*x + 3*a*b**2*x**3 - 6*I*a*b*x**2 - 3*a*x + b**3*x**4 - 3*I*b**2*x**3 - 3*b*x**2 + I*x), x) + Integral(2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x + 3*a**2*b*x**2 - 3*I*a**2*x + 3*a*b**2*x**3 - 6*I*a*b*x**2 - 3*a*x + b**3*x**4 - 3*I*b**2*x**3 - 3*b*x**2 + I*x), x))

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(90) = 180.

time = 0.53, size = 253, normalized size = 1.89

$$\frac{i b \log \left(-3 \left(|x| - \sqrt{(bx+a)^2 + 1} \right) ab - i a^2 b - i \left(|x| - \sqrt{(bx+a)^2 + 1} \right) |x| - 3 \left(|x| - i \sqrt{(bx+a)^2 + 1} \right) a^2 |x| - 2 \left(|x| - \sqrt{(bx+a)^2 + 1} \right) b - 2 a^2 b - 4 \left(|x| - \sqrt{(bx+a)^2 + 1} \right) a |x| + i ab + \left(|x| - i \sqrt{(bx+a)^2 + 1} \right) |x| \right)}{3 |x|} + \frac{(i a^2 - 2 a - i) \log \left(\frac{-|x| + i \sqrt{(bx+a)^2 + 1} - i \sqrt{a^2 + 1}}{-|x| + i \sqrt{(bx+a)^2 + 1} + i \sqrt{a^2 + 1}} \right)}{\sqrt{a^2 + 1} (a - i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x,x, algorithm="giac")

[Out] -1/3*I*b*log(-3*I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*a*b - I*a^3*b - I*(x*abs(b) - sqrt((b*x + a)^2 + 1))^3*abs(b) - 3*(I*x*abs(b) - I*sqrt((b*x + a)^2 + 1))*a^2*abs(b) - 2*(x*abs(b) - sqrt((b*x + a)^2 + 1))^2*b - 2*a^2*b - 4*(x*abs(b) - sqrt((b*x + a)^2 + 1))*a*abs(b) + I*a*b + (I*x*abs(b) - I*sq

```
rt((b*x + a)^2 + 1)*abs(b))/abs(b) + (I*a^2 - 2*a - I)*log(abs(-2*x*abs(b)
+ 2*sqrt((b*x + a)^2 + 1) - 2*sqrt(a^2 + 1))/abs(-2*x*abs(b) + 2*sqrt((b*x
+ a)^2 + 1) + 2*sqrt(a^2 + 1)))/(sqrt(a^2 + 1)*(a - I))
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{((a + bx)^2 + 1)^{3/2}}{x(1 + a \operatorname{li} + b x \operatorname{li})^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)^(3/2)/(x*(a*1i + b*x*1i + 1)^3), x)

[Out] int(((a + b*x)^2 + 1)^(3/2)/(x*(a*1i + b*x*1i + 1)^3), x)

$$3.213 \quad \int \frac{e^{-3i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=178

$$\frac{6ib\sqrt{1-ia-ibx}}{(i-a)^2\sqrt{1+ia+ibx}} - \frac{(1-ia-ibx)^{3/2}}{(1+ia)x\sqrt{1+ia+ibx}} - \frac{6i\sqrt{i+a} b \tanh^{-1}\left(\frac{\sqrt{i+a}\sqrt{1+ia+ibx}}{\sqrt{i-a}\sqrt{1-ia-ibx}}\right)}{(i-a)^{5/2}}$$

[Out] $-6*I*b*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)}/(I-a)^{(1/2)}/(1-I*a-I*b*x)^{(1/2}))*I+a)^{(1/2)}/(I-a)^{(5/2)}-(1-I*a-I*b*x)^{(3/2)}/(1+I*a)/x/(1+I*a+I*b*x)^{(1/2)}+6*I*b*(1-I*a-I*b*x)^{(1/2)}/(I-a)^2/(1+I*a+I*b*x)^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 178, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5203, 96, 95, 214}

$$-\frac{(-ia-ibx+1)^{3/2}}{(1+ia)x\sqrt{ia+ibx+1}} + \frac{6ib\sqrt{-ia-ibx+1}}{(-a+i)^2\sqrt{ia+ibx+1}} - \frac{6i\sqrt{a+i} b \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a + b*x])*x^2),x]

[Out] $((6*I)*b*\operatorname{Sqrt}[1-I*a-I*b*x])/((I-a)^2*\operatorname{Sqrt}[1+I*a+I*b*x]) - (1-I*a-I*b*x)^{(3/2)}/((1+I*a)*x*\operatorname{Sqrt}[1+I*a+I*b*x]) - ((6*I)*\operatorname{Sqrt}[I+a]*b*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])])/((I-a)^{(5/2)})$

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m+1)*(c + d*x)^n*((e + f*x)^(p+1))/((m+1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m+1)*(b*e - a*f))], Int[(a + b*x)^(m+1)*(c + d*x)^(n-1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_))*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-3i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{(1 - ia - ibx)^{3/2}}{x^2(1 + ia + ibx)^{3/2}} dx \\
 &= -\frac{(1 - ia - ibx)^{3/2}}{(1 + ia)x\sqrt{1 + ia + ibx}} + \frac{(3b) \int \frac{\sqrt{1 - ia - ibx}}{x(1 + ia + ibx)^{3/2}} dx}{i - a} \\
 &= \frac{6ib\sqrt{1 - ia - ibx}}{(i - a)^2\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{3/2}}{(1 + ia)x\sqrt{1 + ia + ibx}} + \frac{(3(i + a)b) \int \frac{1}{x\sqrt{1 - ia - ibx}} dx}{(i - a)^2} \\
 &= \frac{6ib\sqrt{1 - ia - ibx}}{(i - a)^2\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{3/2}}{(1 + ia)x\sqrt{1 + ia + ibx}} + \frac{(6(i + a)b) \text{Subst}\left(\int \frac{1}{-1 - ia - (-1 + ibx)} dx\right)}{(i - a)^2} \\
 &= \frac{6ib\sqrt{1 - ia - ibx}}{(i - a)^2\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{3/2}}{(1 + ia)x\sqrt{1 + ia + ibx}} - \frac{6i\sqrt{i + a} b \tanh^{-1}\left(\frac{\sqrt{i + a} \sqrt{1 + ia + ibx}}{\sqrt{i - a} \sqrt{1 + ia + ibx}}\right)}{(i - a)^{5/2}}
 \end{aligned}$$

Mathematica [A]

time = 0.13, size = 145, normalized size = 0.81

$$\frac{\sqrt{-i(i + a + bx)} (1 + a^2 + 5ibx + abx)}{x\sqrt{1 + ia + ibx}} - \frac{6i\sqrt{-1 + ia} b \tanh^{-1}\left(\frac{\sqrt{-1 - ia} \sqrt{-i(i + a + bx)}}{\sqrt{-1 + ia} \sqrt{1 + ia + ibx}}\right)}{\sqrt{-1 - ia}}}{(-i + a)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((3*I)*ArcTan[a + b*x])*x^2), x]

[Out] ((Sqrt[(-I)*(I + a + b*x)]*(1 + a^2 + (5*I)*b*x + a*b*x))/(x*Sqrt[1 + I*a + I*b*x]) - ((6*I)*Sqrt[-1 + I*a]*b*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a

$$+ b*x)]/(\text{Sqrt}[-1 + I*a]*\text{Sqrt}[1 + I*a + I*b*x]])/\text{Sqrt}[-1 - I*a]/(-I + a)^2$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 1542 vs. 2(138) = 276.

time = 0.19, size = 1543, normalized size = 8.67

method	result
risch	$-\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{(a-i)^2x} + \frac{3b \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)a^2}{(a^2-2ia-1)(i-a)\sqrt{a^2+1}} + \frac{3b \ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{(a^2-2ia-1)(i-a)\sqrt{a^2+1}}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^2,x,method=_RETURNVERBOSE)`

[Out]
$$\begin{aligned} & I/(I-a)^2/b*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)- \\ & 2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b \\ & *(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b \\ & ^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I \\ & -a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(\\ & (1/2))))+3*I/(I-a)^4*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I* \\ & b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(\\ & (1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x- \\ & (I-a)/b))^(1/2)))/(b^2)^(1/2))-3*I/(I-a)^4*b*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(\\ & 3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2* \\ & (a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(\\ & 1/2)))/(b^2)^(1/2)+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b \\ &)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((\\ & 2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x))-I/(I-a) \\ & ^3*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+3*a*b/(a^2+1)*(1/3*(b^2*x^2+ \\ & 2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(\\ & 1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+ \\ & 2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a \\ & *b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a \\ & ^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/ \\ & 2))/x)))+4*b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/ \\ & 2)+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a \\ & *b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1 \\ & /2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))))-2*I/(I-a)^3*(-I/b/(x-(I-a) \\ &)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^ \\ & 2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(\\ & I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(\\ & 1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(1/2)))) \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^2,x, algorithm="maxima")
```

```
[Out] integrate(((b*x + a)^2 + 1)^(3/2)/((I*b*x + I*a + 1)^3*x^2), x)
```

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 389 vs. $2(116) = 232$.

time = 4.20, size = 389, normalized size = 2.19

$$\frac{(a-5)Ia^2 + (a^2-4a+5)(a-3)(a^2-2a-1)(a^2-3a^2-3a+5)}{a^2-5a^2-10a^2+5a-1} \log\left(\frac{(a+I)\sqrt{a^2+2ab+I}}{a^2-5a^2-10a^2+5a-1}\right) + 3I(a^2-2a-1)(a^2-3a^2-3a+5) \frac{(a+I)\sqrt{a^2+2ab+I}}{a^2-5a^2-10a^2+5a-1} \log\left(\frac{(a+I)\sqrt{a^2+2ab+I}}{a^2-5a^2-10a^2+5a-1}\right) + \sqrt{I^2+2ab+I}(a-5)(a^2+5)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^2,x, algorithm="fricas")
```

```
[Out] -((I*a - 5)*b^2*x^2 + (I*a^2 - 4*a + 5*I)*b*x - 3*((a^2 - 2*I*a - 1)*b*x^2 + (a^3 - 3*I*a^2 - 3*a + I)*x)*sqrt((a + I)*b^2/(a^5 - 5*I*a^4 - 10*a^3 + 10*I*a^2 + 5*a - I))*log(-(b^2*x + (a^3 - 3*I*a^2 - 3*a + I)*sqrt((a + I)*b^2/(a^5 - 5*I*a^4 - 10*a^3 + 10*I*a^2 + 5*a - I)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b)/b) + 3*((a^2 - 2*I*a - 1)*b*x^2 + (a^3 - 3*I*a^2 - 3*a + I)*x)*sqrt((a + I)*b^2/(a^5 - 5*I*a^4 - 10*a^3 + 10*I*a^2 + 5*a - I))*log(-(b^2*x - (a^3 - 3*I*a^2 - 3*a + I)*sqrt((a + I)*b^2/(a^5 - 5*I*a^4 - 10*a^3 + 10*I*a^2 + 5*a - I)) - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*b)/b) + sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*((I*a - 5)*b*x + I*a^2 + I)/((a^2 - 2*I*a - 1)*b*x^2 + (a^3 - 3*I*a^2 - 3*a + I)*x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\frac{\sqrt{a^2+2ab+I}}{a^2+3a^2b^3-3a^2+3a^2b^2-6a^2b^3-3a^2+I} + \int \frac{a^2\sqrt{a^2+2ab+I}}{a^2+3a^2b^3-3a^2+3a^2b^2-6a^2b^3-3a^2+I} dx + \int \frac{I^2\sqrt{a^2+2ab+I}}{a^2+3a^2b^3-3a^2+3a^2b^2-6a^2b^3-3a^2+I} dx + \int \frac{2ab\sqrt{a^2+2ab+I}}{a^2+3a^2b^3-3a^2+3a^2b^2-6a^2b^3-3a^2+I} dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a)**3*(1+(b*x+a)**2)**(3/2)/x**2,x)
```

```
[Out] I*(Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**2 + 3*a**2*b*x**3 - 3*I*a**2*x**2 + 3*a*b**2*x**4 - 6*I*a*b*x**3 - 3*a*x**2 + b**3*x**5 - 3*I*b**2*x**4 - 3*b*x**3 + I*x**2), x) + Integral(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**2 + 3*a**2*b*x**3 - 3*I*a**2*x**2 + 3*a*b**2*x**4 - 6*I*a*b*x**3 - 3*a*x**2 + b**3*x**5 - 3*I*b**2*x**4 - 3*b*x**3 + I*x**2), x) + Integral(b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**2 + 3*a**2*b*x**3 - 3*I*a**2*x**2 + 3*a*b**2*x**4 - 6*I*a*b*x**3 - 3*a*x**2 + b
```

```
*3*x**5 - 3*I*b**2*x**4 - 3*b*x**3 + I*x**2), x) + Integral(2*a*b*x*sqrt(a*
*2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**2 + 3*a**2*b*x**3 - 3*I*a**2*x**2 +
3*a*b**2*x**4 - 6*I*a*b*x**3 - 3*a*x**2 + b**3*x**5 - 3*I*b**2*x**4 - 3*b*x
**3 + I*x**2), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^2,x, algorithm="giac")
```

```
[Out] undef
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{((a + bx)^2 + 1)^{3/2}}{x^2 (1 + a li + b x li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a + b*x)^2 + 1)^(3/2)/(x^2*(a*1i + b*x*1i + 1)^3),x)
```

```
[Out] int(((a + b*x)^2 + 1)^(3/2)/(x^2*(a*1i + b*x*1i + 1)^3), x)
```

$$3.214 \quad \int \frac{e^{-3i \operatorname{ArcTan}(a+bx)}}{x^3} dx$$

Optimal. Leaf size=264

$$-\frac{3(3i+2a)b^2\sqrt{1-ia-ibx}}{(1+ia)^3(i+a)\sqrt{1+ia+ibx}} + \frac{(3-2ia)b(1-ia-ibx)^{3/2}}{2(i-a)^2(i+a)x\sqrt{1+ia+ibx}} - \frac{(1-ia-ibx)^{5/2}}{2(1+a^2)x^2\sqrt{1+ia+ibx}} + \frac{3(3-2ia)b^2\sqrt{1-ia-ibx}}{(1+ia)^3(i+a)\sqrt{1+ia+ibx}}$$

[Out] $3*(3-2*I*a)*b^2*\operatorname{arctanh}((I+a)^{(1/2)}*(1+I*a+I*b*x)^{(1/2)/(I-a)^{(1/2)/(1-I*a-I*b*x)^{(1/2)})/(I-a)^{(7/2)/(I+a)^{(1/2)+1/2*(3-2*I*a)*b*(1-I*a-I*b*x)^{(3/2)/(I-a)^2/(I+a)/x/(1+I*a+I*b*x)^{(1/2)-1/2*(1-I*a-I*b*x)^{(5/2)/(a^2+1)/x^2/(1+I*a+I*b*x)^{(1/2)-3*(3*I+2*a)*b^2*(1-I*a-I*b*x)^{(1/2)/(1+I*a)^3/(I+a)/(1+I*a+I*b*x)^{(1/2)}}$

Rubi [A]

time = 0.12, antiderivative size = 264, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 5, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.312$, Rules used = {5203, 98, 96, 95, 214}

$$-\frac{(-ia-ibx+1)^{5/2}}{2(a^2+1)x^2\sqrt{ia+ibx+1}} - \frac{3(2a+3i)b^2\sqrt{-ia-ibx+1}}{(1+ia)^3(a+i)\sqrt{ia+ibx+1}} + \frac{3(3-2ia)b^2 \tanh^{-1}\left(\frac{\sqrt{a+i}\sqrt{ia+ibx+1}}{\sqrt{-a+i}\sqrt{-ia-ibx+1}}\right)}{(-a+i)^{7/2}\sqrt{a+i}} + \frac{(3-2ia)b(-ia-ibx+1)^{3/2}}{2(-a+i)^2(a+i)x\sqrt{ia+ibx+1}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a + b*x]))*x^3), x]

[Out] $(-3*(3*I+2*a)*b^2*\operatorname{Sqrt}[1-I*a-I*b*x])/((1+I*a)^3*(I+a)*\operatorname{Sqrt}[1+I*a+I*b*x]) + ((3-(2*I)*a)*b*(1-I*a-I*b*x)^{(3/2)})/(2*(I-a)^2*(I+a)*x*\operatorname{Sqrt}[1+I*a+I*b*x]) - (1-I*a-I*b*x)^{(5/2)}/(2*(1+a^2)*x^2*\operatorname{Sqrt}[1+I*a+I*b*x]) + (3*(3-(2*I)*a)*b^2*\operatorname{ArcTanh}[(\operatorname{Sqrt}[I+a]*\operatorname{Sqrt}[1+I*a+I*b*x])/(\operatorname{Sqrt}[I-a]*\operatorname{Sqrt}[1-I*a-I*b*x])])/((I-a)^{(7/2)}*\operatorname{Sqrt}[I+a])$

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m+1)-1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 96

Int[((a_.) + (b_.)*(x_))^(m_))*((c_.) + (d_.)*(x_))^(n_))*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m+1)*(c + d*x)^n*((e + f*x)^(p+1))/((m+1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m+1)*(b*e - a*f))], Int[(a + b*x)^(m+1)*(c + d*x)^(n-1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimpler

$Q[m, 1] \parallel !\text{SumSimplerQ}[p, 1]) \&\& \text{NeQ}[m, -1]$

Rule 98

$\text{Int}[(a_.) + (b_.)(x_)^m)((c_.) + (d_.)(x_)^n)((e_.) + (f_.)(x_)^p), x_Symbol] \rightarrow \text{Simp}[b*(a + b*x)^{m+1}*(c + d*x)^{n+1}*((e + f*x)^{p+1}/((m+1)*(b*c - a*d)*(b*e - a*f))), x] + \text{Dist}[(a*d*f*(m+1) + b*c*f*(n+1) + b*d*e*(p+1))/((m+1)*(b*c - a*d)*(b*e - a*f)), \text{Int}[(a + b*x)^{m+1}*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& \text{EqQ}[\text{Simplify}[m + n + p + 3], 0] \&\& (\text{LtQ}[m, -1] \parallel \text{SumSimplerQ}[m, 1])$

Rule 214

$\text{Int}[(a_.) + (b_.)(x_)^2]^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b\}, x] \&\& \text{NegQ}[a/b]$

Rule 5203

$\text{Int}[E^{\text{ArcTan}[(c_.)*(a_.) + (b_.)(x_)]*(n_.)}*((d_.) + (e_.)(x_)^m), x_Symbol] \rightarrow \text{Int}[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^{I*(n/2)})/(1 + I*a*c + I*b*c*x)^{I*(n/2)}], x] /; \text{FreeQ}\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned} \int \frac{e^{-3i \tan^{-1}(a+bx)}}{x^3} dx &= \int \frac{(1 - ia - ibx)^{3/2}}{x^3(1 + ia + ibx)^{3/2}} dx \\ &= -\frac{(1 - ia - ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 + ia + ibx}} - \frac{((3i + 2a)b) \int \frac{(1 - ia - ibx)^{3/2}}{x^2(1 + ia + ibx)^{3/2}} dx}{2(1 + a^2)} \\ &= \frac{(3 - 2ia)b(1 - ia - ibx)^{3/2}}{2(i - a)^2(i + a)x\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 + ia + ibx}} + \frac{(3(3i + 2a)b^2) \int \frac{(1 - ia - ibx)^{3/2}}{x\sqrt{1 + ia + ibx}} dx}{2(i - a)^2} \\ &= -\frac{3(3 - 2ia)b^2\sqrt{1 - ia - ibx}}{(i - a)^3(i + a)\sqrt{1 + ia + ibx}} + \frac{(3 - 2ia)b(1 - ia - ibx)^{3/2}}{2(i - a)^2(i + a)x\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 + ia + ibx}} \\ &= -\frac{3(3 - 2ia)b^2\sqrt{1 - ia - ibx}}{(i - a)^3(i + a)\sqrt{1 + ia + ibx}} + \frac{(3 - 2ia)b(1 - ia - ibx)^{3/2}}{2(i - a)^2(i + a)x\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 + ia + ibx}} \\ &= -\frac{3(3 - 2ia)b^2\sqrt{1 - ia - ibx}}{(i - a)^3(i + a)\sqrt{1 + ia + ibx}} + \frac{(3 - 2ia)b(1 - ia - ibx)^{3/2}}{2(i - a)^2(i + a)x\sqrt{1 + ia + ibx}} - \frac{(1 - ia - ibx)^{5/2}}{2(1 + a^2)x^2\sqrt{1 + ia + ibx}} \end{aligned}$$

Mathematica [A]

time = 0.18, size = 194, normalized size = 0.73

$$\frac{\sqrt{-i(i+a+bx)} \frac{(-i+a-ia^2+a^3-5bx-5iabx-14ib^2x^2-ab^2x^2)}{x^2\sqrt{1+ia+ibx}} + \frac{6i\sqrt{-1+ia} (3i+2a)b^2 \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}{\sqrt{-1-ia} (i+a)}}{2(-i+a)^3}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((3*I)*ArcTan[a + b*x])*x^3), x]

[Out] ((Sqrt[(-I)*(I + a + b*x)]*(-I + a - I*a^2 + a^3 - 5*b*x - (5*I)*a*b*x - (14*I)*b^2*x^2 - a*b^2*x^2))/(x^2*Sqrt[1 + I*a + I*b*x]) + ((6*I)*Sqrt[-1 + I*a]*(3*I + 2*a)*b^2*ArcTanh[(Sqrt[-1 - I*a]*Sqrt[(-I)*(I + a + b*x)]]/(Sqrt[-1 + I*a]*Sqrt[1 + I*a + I*b*x]))/(Sqrt[-1 - I*a]*(I + a)))/(2*(-I + a)^3)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 2327 vs. 2(209) = 418.

time = 0.22, size = 2328, normalized size = 8.82

method	result
risch	$-\frac{i(-ab^3x^3-6ib^3x^3-a^2b^2x^2-12ia^2b^2x^2+a^3bx-6ia^2b+a^4+b^2x^2+abx-6ibx+2a^2+1)}{2x^2(a-i)^3\sqrt{b^2x^2+2abx+a^2+1}} - \frac{3ib^2\ln\left(\frac{2a^2+2+2abx+2\sqrt{a^2+1}\sqrt{b^2x^2+2abx+a^2+1}}{x}\right)}{2(a^3-3ia^2-3a+i)(i-a)}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^3,x,method=_RETURNVERBOSE)

[Out]
$$-3*I*b/(I-a)^4*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(1/2)))+I/(I-a)^3*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)-2*I*b*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(5/2)+3*I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2)+I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^(1/2)+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2)))/(b^2)^(1/2)))-I/(I-a)^3*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+1/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+3*a*b/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x^2+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/b^2)$$

$$\begin{aligned} & \wedge(1/2))/b^2)^{(1/2)} - (a^2+1)^{(1/2)} * \ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x)) + 4*b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)} + 3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + 1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2 * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})) + 3/2 * b^2/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)} + a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + 1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2 * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})) + (a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + a*b * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)} - (a^2+1)^{(1/2)} * \ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x))) + 6*I/(I-a)^5*b^2*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(3/2) + I*b*(1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2) + 1/2 * \ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^{(1/2)} + ((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^(1/2))/(b^2)^{(1/2)})) - 6*I/(I-a)^5*b^2*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)} + a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + 1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2 * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})) + (a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + a*b * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)} - (a^2+1)^{(1/2)} * \ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x))) - 3*I/(I-a)^4*b*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(5/2) + 3*a*b/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)} + a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + 1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2 * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})) + (a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + a*b * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)} - (a^2+1)^{(1/2)} * \ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/x))) + 4*b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)} + 3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)} + 1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2 * \ln((b^2*x+a*b)/(b^2)^{(1/2)} + (b^2*x^2+2*a*b*x+a^2+1)^{(1/2)})/(b^2)^{(1/2)})) \wedge(1/2))) \end{aligned}$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^3,x, algorithm="maxima")

[Out] integrate(((b*x + a)^2 + 1)^(3/2)/((I*b*x + I*a + 1)^3*x^3), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 574 vs. 2(180) = 360.

time = 1.96, size = 574, normalized size = 2.17

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^3,x, algorithm="fricas")
[Out] 1/2*((I*a - 14)*b^3*x^3 + (I*a^2 - 13*a + 14*I)*b^2*x^2 - 3*((a^3 - 3*I*a^2 - 3*a + I)*b*x^3 + (a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1)*x^2)*sqrt((4*a^2 + 12*I*a - 9)*b^4/(a^8 - 6*I*a^7 - 14*a^6 + 14*I*a^5 + 14*I*a^3 + 14*a^2 - 6*I*a - 1))*log(-((2*a + 3*I)*b^3*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(2*a + 3*I)*b^2 + (a^5 - 3*I*a^4 - 2*a^3 - 2*I*a^2 - 3*a + I)*sqrt((4*a^2 + 12*I*a - 9)*b^4/(a^8 - 6*I*a^7 - 14*a^6 + 14*I*a^5 + 14*I*a^3 + 14*a^2 - 6*I*a - 1))))/((2*a + 3*I)*b^2)) + 3*((a^3 - 3*I*a^2 - 3*a + I)*b*x^3 + (a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1)*x^2)*sqrt((4*a^2 + 12*I*a - 9)*b^4/(a^8 - 6*I*a^7 - 14*a^6 + 14*I*a^5 + 14*I*a^3 + 14*a^2 - 6*I*a - 1))*log(-((2*a + 3*I)*b^3*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(2*a + 3*I)*b^2 - (a^5 - 3*I*a^4 - 2*a^3 - 2*I*a^2 - 3*a + I)*sqrt((4*a^2 + 12*I*a - 9)*b^4/(a^8 - 6*I*a^7 - 14*a^6 + 14*I*a^5 + 14*I*a^3 + 14*a^2 - 6*I*a - 1))))/((2*a + 3*I)*b^2)) + ((I*a - 14)*b^2*x^2 - I*a^3 - 5*(a - I)*b*x - a^2 - I*a - 1)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/((a^3 - 3*I*a^2 - 3*a + I)*b*x^3 + (a^4 - 4*I*a^3 - 6*a^2 + 4*I*a + 1)*x^2)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\int \frac{\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3x^3 + 3a^2bx^2 + 3ab^2x + b^3} dx + \int \frac{a^2\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3x^3 + 3a^2bx^2 + 3ab^2x + b^3} dx + \int \frac{a^2\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3x^3 + 3a^2bx^2 + 3ab^2x + b^3} dx + \int \frac{2abx\sqrt{a^2 + 2abx + b^2x^2 + 1}}{a^3x^3 + 3a^2bx^2 + 3ab^2x + b^3} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a)**3*(1+(b*x+a)**2)**(3/2)/x**3,x)
[Out] I*(Integral(sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**3 + 3*a**2*b*x**4 - 3*I*a**2*x**3 + 3*a*b**2*x**5 - 6*I*a*b*x**4 - 3*a*x**3 + b**3*x**6 - 3*I*b**2*x**5 - 3*b*x**4 + I*x**3), x) + Integral(a**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**3 + 3*a**2*b*x**4 - 3*I*a**2*x**3 + 3*a*b**2*x**5 - 6*I*a*b*x**4 - 3*a*x**3 + b**3*x**6 - 3*I*b**2*x**5 - 3*b*x**4 + I*x**3), x) + Integral(b**2*x**2*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**3 + 3*a**2*b*x**4 - 3*I*a**2*x**3 + 3*a*b**2*x**5 - 6*I*a*b*x**4 - 3*a*x**3 + b**3*x**6 - 3*I*b**2*x**5 - 3*b*x**4 + I*x**3), x) + Integral(2*a*b*x*sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)/(a**3*x**3 + 3*a**2*b*x**4 - 3*I*a**2*x**3 + 3*a*b**2*x**5 - 6*I*a*b*x**4 - 3*a*x**3 + b**3*x**6 - 3*I*b**2*x**5 - 3*b*x**4 + I*x**3), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^3,x, algorithm="giac")

[Out] undef

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{((a + bx)^2 + 1)^{3/2}}{x^3 (1 + a li + bx li)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)^(3/2)/(x^3*(a*1i + b*x*1i + 1)^3),x)

[Out] int(((a + b*x)^2 + 1)^(3/2)/(x^3*(a*1i + b*x*1i + 1)^3), x)

$$3.215 \quad \int \frac{e^{-3i \operatorname{ArcTan}(a+bx)}}{x^4} dx$$

Optimal. Leaf size=339

$$\frac{(52 - 51ia - 2a^2) b^3 \sqrt{1 - ia - ibx}}{6(i - a)^4 (i + a) \sqrt{1 + ia + ibx}} - \frac{(i + a) \sqrt{1 - ia - ibx}}{3(i - a) x^3 \sqrt{1 + ia + ibx}} - \frac{7ib \sqrt{1 - ia - ibx}}{6(i - a)^2 x^2 \sqrt{1 + ia + ibx}} + \frac{(19 - 16ia)}{6(i - a)^3 (i + a)}$$

[Out] (11*I+18*a-6*I*a^2)*b^3*arctanh((I+a)^(1/2)*(1+I*a+I*b*x)^(1/2)/(I-a)^(1/2)/(1-I*a-I*b*x)^(1/2))/(I-a)^(9/2)/(I+a)^(3/2)-1/6*(52-51*I*a-2*a^2)*b^3*(1-I*a-I*b*x)^(1/2)/(I-a)^4/(I+a)/(1+I*a+I*b*x)^(1/2)-1/3*(I+a)*(1-I*a-I*b*x)^(1/2)/(I-a)/x^3/(1+I*a+I*b*x)^(1/2)-7/6*I*b*(1-I*a-I*b*x)^(1/2)/(I-a)^2/x^2/(1+I*a+I*b*x)^(1/2)+1/6*(19-16*I*a)*b^2*(1-I*a-I*b*x)^(1/2)/(I-a)^3/(I+a)/x/(1+I*a+I*b*x)^(1/2)

Rubi [A]

time = 0.25, antiderivative size = 339, normalized size of antiderivative = 1.00, number of steps used = 8, number of rules used = 7, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.438$, Rules used = {5203, 100, 156, 157, 12, 95, 214}

$$\frac{(-2a^2 - 51ia + 52) b^3 \sqrt{-ia - ibx + 1}}{6(-a + i)^4 (a + i) \sqrt{ia + ibx + 1}} + \frac{(-6ia^2 + 18a + 11i) b^3 \tanh^{-1}\left(\frac{\sqrt{a+i} \sqrt{ia+ibx+1}}{\sqrt{-a+i} \sqrt{-ia-ibx+1}}\right)}{(-a+i)^{9/2} (a+i)^{3/2}} + \frac{(19-16ia) b^2 \sqrt{-ia-ibx+1}}{6(-a+i)^3 (a+i) x \sqrt{ia+ibx+1}} - \frac{(a+i) \sqrt{-ia-ibx+1}}{3(-a+i)^2 x^2 \sqrt{ia+ibx+1}} - \frac{7ib \sqrt{-ia-ibx+1}}{6(-a+i)^2 x^2 \sqrt{ia+ibx+1}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a + b*x])*x^4), x]

[Out] -1/6*((52 - (51*I)*a - 2*a^2)*b^3*Sqrt[1 - I*a - I*b*x])/((I - a)^4*(I + a)*Sqrt[1 + I*a + I*b*x]) - ((I + a)*Sqrt[1 - I*a - I*b*x])/(3*(I - a)*x^3*Sqrt[1 + I*a + I*b*x]) - (((7*I)/6)*b*Sqrt[1 - I*a - I*b*x])/((I - a)^2*x^2*Sqrt[1 + I*a + I*b*x]) + ((19 - (16*I)*a)*b^2*Sqrt[1 - I*a - I*b*x])/(6*(I - a)^3*(I + a)*x*Sqrt[1 + I*a + I*b*x]) + ((11*I + 18*a - (6*I)*a^2)*b^3*ArcTanh[(Sqrt[I + a]*Sqrt[1 + I*a + I*b*x])/(Sqrt[I - a]*Sqrt[1 - I*a - I*b*x])])/((I - a)^(9/2)*(I + a)^(3/2))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 95

Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] := With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]

Rule 100

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(b*c - a*d)*(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*((e + f*x)^(p + 1)/(b*(b*e - a*f)*(m + 1))), x] + Dist[1/(b*(b*e - a*f)*(m + 1)), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 2)*(e + f*x)^p*Simp[a*d*(d*e*(n - 1) + c*f*(p + 1)) + b*c*(d*e*(m - n + 2) - c*f*(m + p + 2)) + d*(a*d*f*(n + p) + b*(d*e*(m + 1) - c*f*(m + n + p + 1)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, p}, x] && LtQ[m, -1] && GtQ[n, 1] && (IntegersQ[2*m, 2*n, 2*p] || IntegersQ[m, n + p] || IntegersQ[p, m + n])
```

Rule 156

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

Rule 157

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g - a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && LtQ[m, -1] && IntegersQ[2*m, 2*n, 2*p]
```

Rule 214

```
Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(a+bx)}}{x^4} dx &= \int \frac{(1-ia-ibx)^{3/2}}{x^4(1+ia+ibx)^{3/2}} dx \\
&= -\frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{\int \frac{7(i+a)b+6b^2x}{x^3\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}} dx}{3(1+ia)} \\
&= -\frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{7ib\sqrt{1-ia-ibx}}{6(i-a)^2x^2\sqrt{1+ia+ibx}} + \frac{\int \frac{-(19-35ia-16a^2)b^2+14(i+a)b^2x}{x^2\sqrt{1-ia-ibx}(1+ia+ibx)^{3/2}} dx}{6(1+ia)(1+a^2)} \\
&= -\frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{7ib\sqrt{1-ia-ibx}}{6(i-a)^2x^2\sqrt{1+ia+ibx}} + \frac{(19i+16a)b^2\sqrt{1-ia-ibx}}{6(1+ia)^3(i+a)x\sqrt{1+ia+ibx}} \\
&= -\frac{(52-51ia-2a^2)b^3\sqrt{1-ia-ibx}}{6(i-a)^4(i+a)\sqrt{1+ia+ibx}} - \frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{7ib\sqrt{1-ia-ibx}}{6(i-a)^2x^2\sqrt{1+ia+ibx}} \\
&= -\frac{(52-51ia-2a^2)b^3\sqrt{1-ia-ibx}}{6(i-a)^4(i+a)\sqrt{1+ia+ibx}} - \frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{7ib\sqrt{1-ia-ibx}}{6(i-a)^2x^2\sqrt{1+ia+ibx}} \\
&= -\frac{(52-51ia-2a^2)b^3\sqrt{1-ia-ibx}}{6(i-a)^4(i+a)\sqrt{1+ia+ibx}} - \frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{7ib\sqrt{1-ia-ibx}}{6(i-a)^2x^2\sqrt{1+ia+ibx}} \\
&= -\frac{(52-51ia-2a^2)b^3\sqrt{1-ia-ibx}}{6(i-a)^4(i+a)\sqrt{1+ia+ibx}} - \frac{(i+a)\sqrt{1-ia-ibx}}{3(i-a)x^3\sqrt{1+ia+ibx}} - \frac{7ib\sqrt{1-ia-ibx}}{6(i-a)^2x^2\sqrt{1+ia+ibx}}
\end{aligned}$$

Mathematica [A]

time = 0.29, size = 275, normalized size = 0.81

$$\frac{-2(-1-ia)^{7/2}(1-ia)(-i(i+a+bx))^{5/2} - (-1-ia)^{5/2}(3i+4a)bx(-i(i+a+bx))^{3/2} + i(-11+18ia+6a^2)b^2x^2 \left(\frac{-i\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}} \right) (1+a^2+5ibx+abx) - 6\sqrt{-1+ia}bx\sqrt{1+ia+ibx} \tanh^{-1}\left(\frac{\sqrt{-1-ia}\sqrt{-i(i+a+bx)}}{\sqrt{-1+ia}\sqrt{1+ia+ibx}}\right)}{6(-1-ia)^{5/2}(1+a^2)^2x^3\sqrt{1+ia+ibx}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((3*I)*ArcTan[a + b*x]))*x^4),x]

[Out]
$$\begin{aligned}
& -1/6*(-2*(-1-I*a)^{(7/2)}*(1-I*a)*((-I)*(I+a+b*x))^{(5/2)} - (-1-I*a)^{(5/2)}*(3*I+4*a)*b*x*((-I)*(I+a+b*x))^{(5/2)} + I*(-11+(18*I)*a+6*a^2)*b^2*x^2*((-I)*\text{Sqrt}[-1-I*a]*\text{Sqrt}[(-I)*(I+a+b*x)]*(1+a^2+(5*I)*b*x+a*b*x) - 6*\text{Sqrt}[-1+I*a]*b*x*\text{Sqrt}[1+I*a+I*b*x]*\text{ArcTanh}[(\text{Sqrt}[-1-I*a]*\text{Sqrt}[(-I)*(I+a+b*x)])]/(\text{Sqrt}[-1+I*a]*\text{Sqrt}[1+I*a+I*b*x])]) / ((-1-I*a)^{(5/2)}*(1+a^2)^2*x^3*\text{Sqrt}[1+I*a+I*b*x])
\end{aligned}$$

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 3636 vs. $2(267) = 534$.

time = 0.30, size = 3637, normalized size = 10.73

method	result
risch	$\frac{i(2a^2b^4x^4+27iab^4x^4+2a^3b^3x^3+45ia^2b^3x^3+9ia^3b^2x^2-28x^4b^4+2a^5bx-9ia^4bx-58ab^3x^3-9ib^3x^3+2a^6-26a^2b^2x^2+9iab^2x^2+4a^6)}{6x^3(i+a)(a-i)^4\sqrt{b^2x^2+2abx+a^2+1}}$
default	Expression too large to display

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^4,x,method=_RETURNVERBOSE)`

[Out]
$$-I/(I-a)^3*(-1/3/(a^2+1)/x^3*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)-1/3*a*b/(a^2+1)*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+1/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+3*a*b/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x)))+4*b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)))+3/2*b^2/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x)))+2/3*b^2/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+3*a*b/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x)))+4*b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)))+3*I/(I-a)^4*b*(-1/2/(a^2+1)/x^2*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+1/2*a*b/(a^2+1)*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^(5/2)+3*a*b/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^(3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^(1/2)+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2))+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^(1/2)+a*b*\ln((b^2*x+a*b)/(b^2)^(1/2)+(b^2*x^2+2*a*b*x+a^2+1)^(1/2)))/(b^2)^(1/2)-(a^2+1)^(1/2)*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^(1/2)*(b^2*x^2+2*a*b*x+a^2+1)^(1/2))/x)))+4$$

$$\begin{aligned}
& *b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)}+3/16*(4 \\
& *b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1 \\
&)^{(1/2)}+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x \\
& ^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)})))+3/2*b^2/(a^2+1)*(1/3*(b^2*x^2+2*a*b \\
& *x+a^2+1)^{(3/2)}+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+ \\
& 1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b \\
& *x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b*\ln \\
& ((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-(a^2+1) \\
& ^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2))/x \\
&))))+10*I*b^3/(I-a)^6*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^{(3/2)}+I*b* \\
& (1/4*(2*(x-(I-a)/b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^{(1 \\
& /2)}+1/2*\ln((I*b+(x-(I-a)/b)*b^2)/(b^2)^{(1/2)}+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I \\
& -a)/b))^{(1/2)))/(b^2)^{(1/2)}))-10*I/(I-a)^6*b^3*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^ \\
& (3/2)+a*b*(1/4*(2*b^2*x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/8*(4*b^2 \\
& *(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1) \\
& ^{(1/2)))/(b^2)^{(1/2)}+(a^2+1)*((b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b*\ln((b^2*x+a* \\
& b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}-(a^2+1)^{(1/2)}*\ln(\\
& (2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2))/x)))-4*I*b^ \\
& 2/(I-a)^5*(-I/b/(x-(I-a)/b)^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^{(5/2)}+3 \\
& *I*b*(1/3*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^{(3/2)}+I*b*(1/4*(2*(x-(I-a)/ \\
& b)*b^2+2*I*b)/b^2*((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^{(1/2)}+1/2*\ln((I*b+(\\
& x-(I-a)/b)*b^2)/(b^2)^{(1/2)}+((x-(I-a)/b)^2*b^2+2*I*b*(x-(I-a)/b))^{(1/2)))/(b \\
& ^2)^{(1/2)}))-6*I/(I-a)^5*b^2*(-1/(a^2+1)/x*(b^2*x^2+2*a*b*x+a^2+1)^{(5/2)}+3* \\
& a*b/(a^2+1)*(1/3*(b^2*x^2+2*a*b*x+a^2+1)^{(3/2)}+a*b*(1/4*(2*b^2*x+2*a*b)/b^2 \\
& *(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*\ln((b^2*x+ \\
& a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)}+(a^2+1)*((b^2* \\
& x^2+2*a*b*x+a^2+1)^{(1/2)}+a*b*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^ \\
& 2+1)^{(1/2)))/(b^2)^{(1/2)}-(a^2+1)^{(1/2)}*\ln((2*a^2+2+2*a*b*x+2*(a^2+1)^{(1/2)}*(\\
& b^2*x^2+2*a*b*x+a^2+1)^{(1/2))/x)))+4*b^2/(a^2+1)*(1/8*(2*b^2*x+2*a*b)/b^2*(\\
& b^2*x^2+2*a*b*x+a^2+1)^{(3/2)}+3/16*(4*b^2*(a^2+1)-4*a^2*b^2)/b^2*(1/4*(2*b^2 \\
& *x+2*a*b)/b^2*(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)}+1/8*(4*b^2*(a^2+1)-4*a^2*b^2)/b \\
& ^2*\ln((b^2*x+a*b)/(b^2)^{(1/2)}+(b^2*x^2+2*a*b*x+a^2+1)^{(1/2)))/(b^2)^{(1/2)})) \\
& +I*b/(I-a)^4*(I/b/(x-(I-a)/b)^3*((x-(I-a)/b)^2*...
\end{aligned}$$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^4,x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 839 vs. 2(223) = 446.

time = 2.53, size = 839, normalized size = 2.47

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^4,x, algorithm="fricas")
[Out] 1/6*((-2*I*a^2 + 51*a + 52*I)*b^4*x^4 + (-2*I*a^3 + 49*a^2 + I*a + 52)*b^3*x^3 + 3*sqrt((36*a^4 + 216*I*a^3 - 456*a^2 - 396*I*a + 121)*b^6/(a^12 - 6*I*a^11 - 12*a^10 + 2*I*a^9 - 27*a^8 + 36*I*a^7 + 36*I*a^5 + 27*a^4 + 2*I*a^3 + 12*a^2 - 6*I*a - 1))*((a^5 - 3*I*a^4 - 2*a^3 - 2*I*a^2 - 3*a + I)*b*x^4 + (a^6 - 4*I*a^5 - 5*a^4 - 5*a^2 + 4*I*a + 1)*x^3)*log(-((6*a^2 + 18*I*a - 11)*b^4*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(6*a^2 + 18*I*a - 11)*b^3 + (a^7 - 3*I*a^6 - a^5 - 5*I*a^4 - 5*a^3 - I*a^2 - 3*a + I)*sqrt((36*a^4 + 216*I*a^3 - 456*a^2 - 396*I*a + 121)*b^6/(a^12 - 6*I*a^11 - 12*a^10 + 2*I*a^9 - 27*a^8 + 36*I*a^7 + 36*I*a^5 + 27*a^4 + 2*I*a^3 + 12*a^2 - 6*I*a - 1))))/(6*a^2 + 18*I*a - 11)*b^3)) - 3*sqrt((36*a^4 + 216*I*a^3 - 456*a^2 - 396*I*a + 121)*b^6/(a^12 - 6*I*a^11 - 12*a^10 + 2*I*a^9 - 27*a^8 + 36*I*a^7 + 36*I*a^5 + 27*a^4 + 2*I*a^3 + 12*a^2 - 6*I*a - 1))*((a^5 - 3*I*a^4 - 2*a^3 - 2*I*a^2 - 3*a + I)*b*x^4 + (a^6 - 4*I*a^5 - 5*a^4 - 5*a^2 + 4*I*a + 1)*x^3)*log(-((6*a^2 + 18*I*a - 11)*b^4*x - sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(6*a^2 + 18*I*a - 11)*b^3 - (a^7 - 3*I*a^6 - a^5 - 5*I*a^4 - 5*a^3 - I*a^2 - 3*a + I)*sqrt((36*a^4 + 216*I*a^3 - 456*a^2 - 396*I*a + 121)*b^6/(a^12 - 6*I*a^11 - 12*a^10 + 2*I*a^9 - 27*a^8 + 36*I*a^7 + 36*I*a^5 + 27*a^4 + 2*I*a^3 + 12*a^2 - 6*I*a - 1))))/(6*a^2 + 18*I*a - 11)*b^3)) + ((-2*I*a^2 + 51*a + 52*I)*b^3*x^3 - 2*I*a^5 + (16*a^2 + 3*I*a + 19)*b^2*x^2 - 2*a^4 - 4*I*a^3 - 7*(a^3 - I*a^2 + a - I)*b*x - 4*a^2 - 2*I*a - 2)*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/((a^5 - 3*I*a^4 - 2*a^3 - 2*I*a^2 - 3*a + I)*b*x^4 + (a^6 - 4*I*a^5 - 5*a^4 - 5*a^2 + 4*I*a + 1)*x^3)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*(b*x+a))**3*(1+(b*x+a)**2)**(3/2)/x**4,x)
```

```
[Out] Timed out
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*(b*x+a))^3*(1+(b*x+a)^2)^(3/2)/x^4,x, algorithm="giac")

[Out] undef

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{((a + bx)^2 + 1)^{3/2}}{x^4 (1 + a^2 + b^2 x^2)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a + b*x)^2 + 1)^(3/2)/(x^4*(a^2 + b*x^2 + 1)^3),x)

[Out] int(((a + b*x)^2 + 1)^(3/2)/(x^4*(a^2 + b*x^2 + 1)^3), x)

3.216 $\int e^{\frac{1}{2}i \text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=494

$$\frac{(3i + 4a - 8ia^2)(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{8b^3} - \frac{(i + 8a)(1 - ia - ibx)^{3/4}(1 + ia + ibx)^{5/4}}{12b^3} + \frac{x(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{12b^3}$$

[Out] $-1/8*(3*I+4*a-8*I*a^2)*(1-I*a-I*b*x)^(3/4)*(1+I*a+I*b*x)^(1/4)/b^3-1/12*(I+8*a)*(1-I*a-I*b*x)^(3/4)*(1+I*a+I*b*x)^(5/4)/b^3+1/3*x*(1-I*a-I*b*x)^(3/4)*(1+I*a+I*b*x)^(5/4)/b^2+1/16*(3*I+4*a-8*I*a^2)*\arctan(1-(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4))/b^3-1/16*(3*I+4*a-8*I*a^2)*\arctan(1+(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4))/b^3-1/32*(3*I+4*a-8*I*a^2)*\ln(1-(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4)+(1-I*a-I*b*x)^(1/2)/(1+I*a+I*b*x)^(1/2))/b^3-1/32*(3*I+4*a-8*I*a^2)*\ln(1+(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4)+(1-I*a-I*b*x)^(1/2)/(1+I*a+I*b*x)^(1/2))/b^3$

Rubi [A]

time = 0.30, antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5203, 92, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{(-8a^2 + 4a + 3) \text{ArcTan}\left(\frac{1 - \sqrt{2} \sqrt{1 + ia + ibx}}{2a + ibx + 1}\right)}{8\sqrt{2}b^3} - \frac{(-8a^2 + 4a + 3) \text{ArcTan}\left(\frac{1 + \sqrt{2} \sqrt{1 + ia + ibx}}{2a + ibx + 1}\right)}{8\sqrt{2}b^3} - \frac{(-8a^2 + 4a + 3)(-in - 1)^{3/4} \sqrt[4]{1 + ia + ibx}}{16\sqrt{2}b^3} - \frac{(-8a^2 + 4a + 3) \log\left(\frac{\sqrt{2} \sqrt{1 + ia + ibx} + 1}{\sqrt{1 + ia + ibx}}\right)}{16\sqrt{2}b^3} - \frac{(-8a^2 + 4a + 3) \log\left(\frac{\sqrt{2} \sqrt{1 + ia + ibx} - 1}{\sqrt{1 + ia + ibx}}\right)}{16\sqrt{2}b^3} - \frac{(8a + 1)(-in - ibx + 1)^{3/4}(1 + ia + ibx)^{5/4}}{12b^3} - \frac{x(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{12b^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(I/2)*\text{ArcTan}[a + b*x]}*x^2, x]$

[Out] $-1/8*((3*I + 4*a - (8*I)*a^2)*(1 - I*a - I*b*x)^(3/4)*(1 + I*a + I*b*x)^(1/4))/b^3 - ((I + 8*a)*(1 - I*a - I*b*x)^(3/4)*(1 + I*a + I*b*x)^(5/4))/(12*b^3) + (x*(1 - I*a - I*b*x)^(3/4)*(1 + I*a + I*b*x)^(5/4))/(3*b^2) + ((3*I + 4*a - (8*I)*a^2)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/ (8*\text{Sqrt}[2]*b^3) - ((3*I + 4*a - (8*I)*a^2)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/ (8*\text{Sqrt}[2]*b^3) - ((3*I + 4*a - (8*I)*a^2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/ (16*\text{Sqrt}[2]*b^3) + ((3*I + 4*a - (8*I)*a^2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/ (16*\text{Sqrt}[2]*b^3)$

Rule 52

$\text{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] := \text{Simp}[(a + b*x)^{m+1} * (c + d*x)^n / (b*(m+n+1)), x] + \text{Dist}[n*(b*c - a*d) / (b*(m+n+1)), \text{Int}[(a + b*x)^m * (c + d*x)^{n-1}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ

$[m, 0] \&\& (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0])) \ \&\& \ !\text{ILtQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \ :> \ \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p(m+1)-1)}(c - a(d/b) + d(x^{p/b}))^n, x], x, (a + b*x)^{(1/p)}], x]] \ /; \ \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 81

$\text{Int}[(a_.) + (b_.)(x_)^{(c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_Symbol] \ :> \ \text{Simp}[b*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 2, 0]$

Rule 92

$\text{Int}[(a_.) + (b_.)(x_)^2((c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_Symbol] \ :> \ \text{Simp}[b*(a + b*x)*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+3))), x] + \text{Dist}[1/(d*f*(n+p+3)), \text{Int}[(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(n+p+3) - b*(b*c*e + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(n+p+4) - b*(d*e*(n+2) + c*f*(p+2))]*x, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 3, 0]$

Rule 210

$\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \ :> \ \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \ /; \ \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 303

$\text{Int}[(x_)^2/((a_) + (b_.)(x_)^4), x_Symbol] \ :> \ \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*s), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Dist}[1/(2*s), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] \ /; \ \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 338

$\text{Int}[(x_)^{(m_.)}((a_) + (b_.)(x_)^{(n_.)})^{(p_)}, x_Symbol] \ :> \ \text{Dist}[a^{(p+(m+1)/n)}, \text{Subst}[\text{Int}[x^m/(1 - b*x^n)^{(p+(m+1)/n+1)}, x], x, x/(a + b*x^n)]$

$^{(1/n)}$, x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
 $^{(-1)}$] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
 Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
 /(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
 & EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
 eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_.))*((d_) + (e_)*(x_)^(m_.),
 x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
 I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{2}i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2 \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx \\
&= \frac{x(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{3b^2} + \frac{\int \frac{\sqrt[4]{1+ia+ibx} (-1-a^2-\frac{1}{2}(i+8a)bx)}{\sqrt[4]{1-ia-ibx}} dx}{3b^2} \\
&= -\frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} + \frac{x(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{3b^2} - \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3} \\
&= -\frac{(3i+4a-8ia^2)(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{8b^3} - \frac{(i+8a)(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{12b^3}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 121, normalized size = 0.24

$$\frac{(-i(i+a+bx))^{3/4} \left(-i\sqrt[4]{1+ia+ibx} (1+8a^2+5ibx-4b^2x^2+a(-7i+4bx)) + 2i\sqrt[4]{2} (-3+4ia+8a^2) {}_2F_1\left(-\frac{1}{4}, \frac{3}{4}; \frac{7}{4}; -\frac{1}{2}i(i+a+bx)\right) \right)}{12b^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a + b*x])*x^2,x]

[Out] (((-I)*(I + a + b*x))^(3/4))*((-I)*(1 + I*a + I*b*x)^(1/4))*(1 + 8*a^2 + (5*I)*b*x - 4*b^2*x^2 + a*(-7*I + 4*b*x)) + (2*I)*2^(1/4)*(-3 + (4*I)*a + 8*a^2)*Hypergeometric2F1[-1/4, 3/4, 7/4, (-1/2*I)*(I + a + b*x)]/(12*b^3)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x^2,x)

[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate


Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x^2,x, algorithm="maxima")

[Out] integrate(x^2*sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1)), x)

Fricas [A]

time = 4.16, size = 554, normalized size = 1.12



Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x^2,x, algorithm="fricas")

[Out] $\frac{1}{48} \cdot (3 \cdot b^3 \cdot \sqrt{(64 \cdot I \cdot a^4 - 64 \cdot a^3 - 64 \cdot I \cdot a^2 + 24 \cdot a + 9 \cdot I) / b^6}) \cdot \log((I \cdot b^3 \cdot \sqrt{(64 \cdot I \cdot a^4 - 64 \cdot a^3 - 64 \cdot I \cdot a^2 + 24 \cdot a + 9 \cdot I) / b^6}) + (8 \cdot a^2 + 4 \cdot I \cdot a - 3) \cdot \sqrt{I \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1}} / (b \cdot x + a + I)) / (8 \cdot a^2 + 4 \cdot I \cdot a - 3)) - 3 \cdot b^3 \cdot \sqrt{(64 \cdot I \cdot a^4 - 64 \cdot a^3 - 64 \cdot I \cdot a^2 + 24 \cdot a + 9 \cdot I) / b^6}) \cdot \log((-I \cdot b^3 \cdot \sqrt{(64 \cdot I \cdot a^4 - 64 \cdot a^3 - 64 \cdot I \cdot a^2 + 24 \cdot a + 9 \cdot I) / b^6}) + (8 \cdot a^2 + 4 \cdot I \cdot a - 3) \cdot \sqrt{I \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1}} / (b \cdot x + a + I)) / (8 \cdot a^2 + 4 \cdot I \cdot a - 3)) + 3 \cdot b^3 \cdot \sqrt{(-64 \cdot I \cdot a^4 + 64 \cdot a^3 + 64 \cdot I \cdot a^2 - 24 \cdot a - 9 \cdot I) / b^6}) \cdot \log((I \cdot b^3 \cdot \sqrt{(-64 \cdot I \cdot a^4 + 64 \cdot a^3 + 64 \cdot I \cdot a^2 - 24 \cdot a - 9 \cdot I) / b^6}) + (8 \cdot a^2 + 4 \cdot I \cdot a - 3) \cdot \sqrt{I \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1}} / (b \cdot x + a + I)) / (8 \cdot a^2 + 4 \cdot I \cdot a - 3)) - 3 \cdot b^3 \cdot \sqrt{(-64 \cdot I \cdot a^4 + 64 \cdot a^3 + 64 \cdot I \cdot a^2 - 24 \cdot a - 9 \cdot I) / b^6}) \cdot \log((-I \cdot b^3 \cdot \sqrt{(-64 \cdot I \cdot a^4 + 64 \cdot a^3 + 64 \cdot I \cdot a^2 - 24 \cdot a - 9 \cdot I) / b^6}) + (8 \cdot a^2 + 4 \cdot I \cdot a - 3) \cdot \sqrt{I \cdot \sqrt{b^2 \cdot x^2 + 2 \cdot a \cdot b \cdot x + a^2 + 1}} / (b \cdot x + a + I)) / (8 \cdot a^2 + 4 \cdot I \cdot a - 3)) + 2 \cdot (8 \cdot b^3 \cdot x^3 - 2 \cdot I \cdot b^2 \cdot x^2 + 8 \cdot a^3 + (8 \cdot I \cdot a - 1) \cdot b \cdot x +$

$34*I*a^2 - 37*a - 11*I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)))/b^3$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \sqrt{\frac{i(a + bx - i)}{\sqrt{a^2 + 2abx + b^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2)*x**2,x)

[Out] Integral(x**2*sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \sqrt{\frac{1 + a \operatorname{li} + b x \operatorname{li}}{\sqrt{(a + b x)^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2),x)

[Out] int(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)

3.217 $\int e^{\frac{1}{2}i \text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=410

$$\frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(1-4ia) \text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2}b^2}$$

[Out] $1/4*(1-4*I*a)*(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(1/4)}/b^2+1/2*(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(5/4)}/b^2-1/8*(1-4*I*a)*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^2+1/8*(1-4*I*a)*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^2+1/16*(1-4*I*a)*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)}/b^2-1/16*(1-4*I*a)*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)}/b^2$

Rubi [A]

time = 0.22, antiderivative size = 410, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {5203, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{(1-4ia)\text{ArcTan}\left(1-\frac{\sqrt{2}\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}}\right)}{4\sqrt{2}b^2} + \frac{(1-4ia)\text{ArcTan}\left(1+\frac{\sqrt{2}\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}}\right)}{4\sqrt{2}b^2} + \frac{(-ia-ibx+1)^{3/4}(ia+ibx+1)^{5/4}}{2b^2} + \frac{(1-4ia)(-ia-ibx+1)^{3/4}\sqrt{ia+ibx+1}}{4b^2} + \frac{(1-4ia)\log\left(\frac{\sqrt{-ia-ibx+1}-\sqrt{2}\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}}+1\right)}{8\sqrt{2}b^2} - \frac{(1-4ia)\log\left(\frac{\sqrt{-ia-ibx+1}+\sqrt{2}\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}}+1\right)}{8\sqrt{2}b^2}$$

Antiderivative was successfully verified.

[In] Int[E^((I/2)*ArcTan[a + b*x])*x,x]

[Out] $((1-(4*I)*a)*(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(1/4)})/(4*b^2) + ((1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(5/4)})/(2*b^2) - ((1-(4*I)*a)*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(4*\text{Sqrt}[2]*b^2) + ((1-(4*I)*a)*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(4*\text{Sqrt}[2]*b^2) + ((1-(4*I)*a)*\text{Log}[1+\text{Sqrt}[1-I*a-I*b*x]/\text{Sqrt}[1+I*a+I*b*x] - (\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(8*\text{Sqrt}[2]*b^2) - ((1-(4*I)*a)*\text{Log}[1+\text{Sqrt}[1-I*a-I*b*x]/\text{Sqrt}[1+I*a+I*b*x] + (\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(8*\text{Sqrt}[2]*b^2)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
```


e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_.)*((d_) + (e_)*(x_)^(m_.)), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{\frac{1}{2}i \tan^{-1}(a+bx)} x dx &= \int \frac{x \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx \\
&= \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b} \\
&= \frac{(1-4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} + \frac{(1-ia-ibx)^{3/4}(1+ia+ibx)^{5/4}}{2b^2} - \frac{(i+4a) \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 81, normalized size = 0.20

$$\frac{(-i(i+a+bx))^{3/4} \left(3(1+ia+ibx)^{5/4} + 2\sqrt[4]{2} (1-4ia) {}_2F_1\left(-\frac{1}{4}, \frac{3}{4}; \frac{7}{4}; -\frac{1}{2}i(i+a+bx)\right) \right)}{6b^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a + b*x])*x,x]

[Out] (((-I)*(I + a + b*x))^(3/4)*(3*(1 + I*a + I*b*x)^(5/4) + 2*2^(1/4)*(1 - (4*I)*a)*Hypergeometric2F1[-1/4, 3/4, 7/4, (-1/2*I)*(I + a + b*x)]))/ (6*b^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x,x)**[Out]** int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x,x, algorithm="maxima")**[Out]** integrate(x*sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1)), x)**Fricas [A]**

time = 4.34, size = 415, normalized size = 1.01

$$\frac{b^2 \sqrt{\frac{16Ia^2 - 8a - I}{b^4}} \log\left(\frac{\sqrt{\frac{16Ia^2 - 8a - I}{b^4}} \sqrt{\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + I}}}{\sqrt{\frac{16Ia^2 - 8a - I}{b^4}}}\right) - b^2 \sqrt{\frac{16Ia^2 - 8a - I}{b^4}} \log\left(\frac{\sqrt{\frac{16Ia^2 - 8a - I}{b^4}} \sqrt{\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + I}}}{\sqrt{\frac{16Ia^2 - 8a - I}{b^4}}}\right) + b^2 \sqrt{\frac{-16Ia^2 + 8a + I}{b^4}} \log\left(\frac{\sqrt{\frac{-16Ia^2 + 8a + I}{b^4}} \sqrt{\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + I}}}{\sqrt{\frac{-16Ia^2 + 8a + I}{b^4}}}\right) - b^2 \sqrt{\frac{-16Ia^2 + 8a + I}{b^4}} \log\left(\frac{\sqrt{\frac{-16Ia^2 + 8a + I}{b^4}} \sqrt{\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + I}}}{\sqrt{\frac{-16Ia^2 + 8a + I}{b^4}}}\right) - 2(2b^2x^2 - 2a^2 - Ibx - 5Ia + 3) \sqrt{\frac{\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + I}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x,x, algorithm="fricas")

[Out] $-1/8*(b^2*\sqrt{(16*I*a^2 - 8*a - I)/b^4})*\log((I*b^2*\sqrt{(16*I*a^2 - 8*a - I)/b^4} + (4*a + I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a + I)) - b^2*\sqrt{(16*I*a^2 - 8*a - I)/b^4}*\log((-I*b^2*\sqrt{(16*I*a^2 - 8*a - I)/b^4} + (4*a + I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a + I)) + b^2*\sqrt{(-16*I*a^2 + 8*a + I)/b^4}*\log((I*b^2*\sqrt{(-16*I*a^2 + 8*a + I)/b^4} + (4*a + I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a + I)) - b^2*\sqrt{(-16*I*a^2 + 8*a + I)/b^4}*\log((-I*b^2*\sqrt{(-16*I*a^2 + 8*a + I)/b^4} + (4*a + I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a + I)) - 2*(2*b^2*x^2 - 2*a^2 - I*b*x - 5*I*a + 3)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)}/b^2$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x \sqrt{\frac{i(a + bx - i)}{\sqrt{a^2 + 2abx + b^2x^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2)*x,x)
```

```
[Out] Integral(x*sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)*x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x \sqrt{\frac{1 + a \operatorname{li} + b x \operatorname{li}}{\sqrt{(a + b x)^2 + 1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2),x)
```

```
[Out] int(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)
```

3.218 $\int e^{\frac{1}{2}i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=338

$$\frac{i(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{b} - \frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b}$$

[Out] $I*(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(1/4)}/b-1/2*I*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}+1/2*I*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}+1/4*I*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}-1/4*I*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 338, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {5201, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} + \frac{i(-ia-ibx+1)^{3/4}\sqrt[4]{ia+ibx+1}}{b} + \frac{i\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b} - \frac{i\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((I/2)*\text{ArcTan}[a + b*x])}, x]$

[Out] $(I*(1 - I*a - I*b*x)^{(3/4)}*(1 + I*a + I*b*x)^{(1/4)})/b - (I*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(I*\text{Sqrt}[2]*b) + (I*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(I*\text{Sqrt}[2]*b) + ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(I*\text{Sqrt}[2]*b) - ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(I*\text{Sqrt}[2]*b)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

$\text{Int}[(a_. + (b_.)*(x_))^{(m_)}*((c_.) + (d_.)*(x_))^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) +$

```
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b
)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5201

```
Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]
```

Rubi steps

$$\begin{aligned}
 \int e^{\frac{1}{2}i \tan^{-1}(a+bx)} dx &= \int \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}} dx \\
 &= \frac{i(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{b} + \frac{1}{2} \int \frac{1}{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}} dx \\
 &= \frac{i(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{b} + \frac{(2i) \text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-ia-ibx}\right)}{b} \\
 &= \frac{i(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{b} + \frac{(2i) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{b} \\
 &= \frac{i(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{b} - \frac{i \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{b} + \frac{i \text{Subst}\left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{2b} + \frac{i \log\left(1 + \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{2\sqrt{2} b} \\
 &= \frac{i(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{b} - \frac{i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2} b} + \frac{i \tan^{-1}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2} b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 45, normalized size = 0.13

$$\frac{8ie^{\frac{5}{2}i \text{ArcTan}(a+bx)} {}_2F_1\left(\frac{5}{4}, 2; \frac{9}{4}; -e^{2i \text{ArcTan}(a+bx)}\right)}{5b}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a + b*x]),x]

[Out] (((-8*I)/5)*E^(((5*I)/2)*ArcTan[a + b*x])*Hypergeometric2F1[5/4, 2, 9/4, -E^((2*I)*ArcTan[a + b*x])]/b

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x)

[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1)), x)

Fricas [A]

time = 3.15, size = 255, normalized size = 0.75

$$\frac{b\sqrt{\frac{I}{b^2}} \log\left(i b\sqrt{\frac{I}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - b\sqrt{\frac{I}{b^2}} \log\left(-i b\sqrt{\frac{I}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) + b\sqrt{\frac{-I}{b^2}} \log\left(i b\sqrt{\frac{-I}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - b\sqrt{\frac{-I}{b^2}} \log\left(-i b\sqrt{\frac{-I}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) + 2(bx + a + i)\sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="fricas")

[Out] 1/2*(b*sqrt(I/b^2)*log(I*b*sqrt(I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(I/b^2)*log(-I*b*sqrt(I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + b*sqrt(-I/b^2)*log(I*b*sqrt(-I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(-I/b^2)*log(-I*b*sqrt(-I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 2*(b*x + a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{\frac{i(a+bx)+1}{\sqrt{(a+bx)^2+1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2), x)``[Out] Integral(sqrt((I*(a + b*x) + 1)/sqrt((a + b*x)**2 + 1)), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2), x, algorithm="giac")``[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \sqrt{\frac{1+a1i+bx1i}{\sqrt{(a+bx)^2+1}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)``[Out] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)`

$$3.219 \quad \int \frac{e^{\frac{1}{2}i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=395

$$\frac{2\sqrt[4]{i-a} \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}\right)}{\sqrt[4]{i+a}} - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}}\right)$$

[Out] $-2*(I-a)^{(1/4)}*\arctan((I+a)^{(1/4)}*(1+I*(b*x+a))^{(1/4)} / (I-a)^{(1/4)} / (1-I*(b*x+a))^{(1/4)}) / (I+a)^{(1/4)} - 2*(I-a)^{(1/4)}*\operatorname{arctanh}((I+a)^{(1/4)}*(1+I*(b*x+a))^{(1/4)} / (I-a)^{(1/4)} / (1-I*(b*x+a))^{(1/4)}) / (I+a)^{(1/4)} - 1/2*\ln(1-(1+I*(b*x+a))^{(1/4)})*2^{(1/2)} / (1-I*(b*x+a))^{(1/4)} + (1+I*(b*x+a))^{(1/2)} / (1-I*(b*x+a))^{(1/2)})*2^{(1/2)} + 1/2*\ln(1+(1+I*(b*x+a))^{(1/4)})*2^{(1/2)} / (1-I*(b*x+a))^{(1/4)} + (1+I*(b*x+a))^{(1/2)} / (1-I*(b*x+a))^{(1/2)})*2^{(1/2)} - \arctan(1-(1+I*(b*x+a))^{(1/4)})*2^{(1/2)} / (1-I*(b*x+a))^{(1/4)})*2^{(1/2)} + \arctan(1+(1+I*(b*x+a))^{(1/4)})*2^{(1/2)} / (1-I*(b*x+a))^{(1/4)})*2^{(1/2)}$

Rubi [A]

time = 0.21, antiderivative size = 395, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5202, 456, 492, 217, 1179, 642, 1176, 631, 210, 218, 214, 211}

$$\frac{2\sqrt{-a+i} \operatorname{ArcTan}\left(\frac{\sqrt[4]{a+i} \sqrt[4]{1+i(a+bx)}}{\sqrt{-a+i} \sqrt[4]{1-i(a+bx)}}\right)}{\sqrt[4]{a+i}} - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}}\right) - \frac{\log\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}} - \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} + 1\right)}{\sqrt{2}} + \frac{\log\left(\frac{\sqrt{1+i(a+bx)}}{\sqrt{1-i(a+bx)}} + \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} + 1\right)}{\sqrt{2}} - \frac{2\sqrt{-a+i} \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{a+i} \sqrt[4]{1+i(a+bx)}}{\sqrt{-a+i} \sqrt[4]{1-i(a+bx)}}\right)}{\sqrt[4]{a+i}}$$

Antiderivative was successfully verified.

[In] Int[E^((I/2)*ArcTan[a + b*x])/x,x]

[Out] $(-2*(I-a)^{(1/4)}*\operatorname{ArcTan}(((I+a)^{(1/4)}*(1+I*(a+b*x))^{(1/4)}) / ((I-a)^{(1/4)}*(1-I*(a+b*x))^{(1/4)})) / (I+a)^{(1/4)} - \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1+I*(a+b*x))^{(1/4)}) / (1-I*(a+b*x))^{(1/4)}] + \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1+I*(a+b*x))^{(1/4)}) / (1-I*(a+b*x))^{(1/4)}] - (2*(I-a)^{(1/4)}*\operatorname{ArcTanh}(((I+a)^{(1/4)}*(1+I*(a+b*x))^{(1/4)}) / ((I-a)^{(1/4)}*(1-I*(a+b*x))^{(1/4)})) / (I+a)^{(1/4)} - \operatorname{Log}[1 - (\operatorname{Sqrt}[2]*(1+I*(a+b*x))^{(1/4)}) / (1-I*(a+b*x))^{(1/4)}] + \operatorname{Sqrt}[1+I*(a+b*x)] / \operatorname{Sqrt}[1-I*(a+b*x)] / \operatorname{Sqrt}[2] + \operatorname{Log}[1 + (\operatorname{Sqrt}[2]*(1+I*(a+b*x))^{(1/4)}) / (1-I*(a+b*x))^{(1/4)}] + \operatorname{Sqrt}[1+I*(a+b*x)] / \operatorname{Sqrt}[1-I*(a+b*x)] / \operatorname{Sqrt}[2]$

Rule 210

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 217

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 456

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_)*((c_) + (d_)*(x_)^(n_))^(q_), x_Symbol] := Int[x^(m + n*(p + q))*(b + a/x^n)^p*(d + c/x^n)^q, x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && IntegersQ[p, q] && NegQ[n]

Rule 492

Int[((e_)*(x_)^(m_))/(((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_))), x_Symbol] := Dist[(-a)*(e^n/(b*c - a*d)), Int[(e*x)^(m - n)/(a + b*x^n), x], x] + Dist[c*(e^n/(b*c - a*d)), Int[(e*x)^(m - n)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LeQ[n, m, 2*n - 1]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5202

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_))*(x_)^(m_), x_Symbol] := Dist
[4/(I^m*n*b^(m + 1)*c^(m + 1)), Subst[Int[x^(2/(I*n))*((1 - I*a*c - (1 + I*
a*c)*x^(2/(I*n)))^m/(1 + x^(2/(I*n)))^(m + 2)), x], x, (1 - I*c*(a + b*x))^
(I*(n/2))/(1 + I*c*(a + b*x))^(I*(n/2)], x] /; FreeQ[{a, b, c}, x] && ILtQ
[m, 0] && LtQ[-1, I*n, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{1}{2}i \tan^{-1}(a+bx)}}{x} dx &= 8 \text{Subst} \left(\int \frac{1}{\left(1 + \frac{1}{x^4}\right) \left(1 - ia - \frac{1+ia}{x^4}\right) x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) \\
&= 8 \text{Subst} \left(\int \frac{x^4}{(1+x^4)(-1-ia+(1-ia)x^4)} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) \\
&= 4 \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) + (4(1+ia)) \text{Subst} \left(\int \frac{1}{-1-ia+(1-ia)x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) \\
&= 2 \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) + 2 \text{Subst} \left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) \\
&= -\frac{2\sqrt[4]{i-a} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}} \right)}{\sqrt[4]{i+a}} - \frac{2\sqrt[4]{i-a} \tanh^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}} \right)}{\sqrt[4]{i+a}} \\
&= -\frac{2\sqrt[4]{i-a} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}} \right)}{\sqrt[4]{i+a}} - \frac{2\sqrt[4]{i-a} \tanh^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}} \right)}{\sqrt[4]{i+a}} \\
&= -\frac{2\sqrt[4]{i-a} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}} \right)}{\sqrt[4]{i+a}} - \sqrt{2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 124, normalized size = 0.31

$$\frac{2}{3}(-i(i+a+bx))^{3/4} \left(-\sqrt[4]{2} {}_2F_1 \left(\frac{3}{4}, \frac{3}{4}; \frac{7}{4}; -\frac{1}{2}i(i+a+bx) \right) + \frac{2(-i+a) {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx} \right)}{(i+a)(1+ia+ibx)^{3/4}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a + b*x])/x,x]

[Out] (2*((-I)*(I + a + b*x))^(3/4)*(-(2^(1/4))*Hypergeometric2F1[3/4, 3/4, 7/4, (-1/2*I)*(I + a + b*x)]) + (2*(-I + a)*Hypergeometric2F1[3/4, 1, 7/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)])/((I + a)*(1 + I*a + I*b*x)^(3/4)))/3

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x)
```

```
[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x, algorithm="maxima")
```

```
[Out] integrate(sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))/x, x)
```

Fricas [A]

time = 2.81, size = 414, normalized size = 1.05

```
1/2*sqrt(4*I)*log(1/2*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(4*I)*log(-1/2*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 1/2*sqrt(-4*I)*log(1/2*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(-4*I)*log(-1/2*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - I*(-(a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + I*(-(a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - I*(-(a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))
```

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x, algorithm="fricas")
```

```
[Out] 1/2*sqrt(4*I)*log(1/2*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(4*I)*log(-1/2*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 1/2*sqrt(-4*I)*log(1/2*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(-4*I)*log(-1/2*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - I*(-(a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + I*(-(a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - I*(-(a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((-a - I)/(a + I))^(1/4)*log(sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{i(a+bx-i)}{\sqrt{a^2+2abx+b^2x^2+1}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2)/x,x)**[Out]** Integral(sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1))/x, x)**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x, algorithm="giac")**[Out]** Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{1+a\,1i+b\,x\,1i}{\sqrt{(a+bx)^2+1}}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2)/x,x)**[Out]** int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2)/x, x)

$$3.220 \quad \int \frac{e^{\frac{1}{2}i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=205

$$-\frac{(i+a+bx)\sqrt[4]{1+i(a+bx)}}{(i+a)x\sqrt[4]{1-i(a+bx)}} + \frac{ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a}\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a}\sqrt[4]{1-i(a+bx)}}\right)}{(i-a)^{3/4}(i+a)^{5/4}} + \frac{ib \tanh^{-1}\left(\frac{\sqrt[4]{i+a}\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a}\sqrt[4]{1-i(a+bx)}}\right)}{(i-a)^{3/4}(i+a)^{5/4}}$$

[Out] $-(I+a+b*x)*(1+I*(b*x+a))^{(1/4)}/(I+a)/x/(1-I*(b*x+a))^{(1/4)}+I*b*\arctan((I+a)^{(1/4)*(1+I*(b*x+a))^{(1/4)}/(I-a)^{(1/4)/(1-I*(b*x+a))^{(1/4)})/(I-a)^{(3/4)/(I+a)^{(5/4)}+I*b*\operatorname{arctanh}((I+a)^{(1/4)*(1+I*(b*x+a))^{(1/4)}/(I-a)^{(1/4)/(1-I*(b*x+a))^{(1/4)})/(I-a)^{(3/4)/(I+a)^{(5/4)}$

Rubi [A]

time = 0.11, antiderivative size = 205, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5202, 269, 294, 218, 214, 211}

$$\frac{ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{a+i}\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{-a+i}\sqrt[4]{1-i(a+bx)}}\right)}{(-a+i)^{3/4}(a+i)^{5/4}} - \frac{\sqrt[4]{1+i(a+bx)}(a+bx+i)}{(a+i)x\sqrt[4]{1-i(a+bx)}} + \frac{ib \tanh^{-1}\left(\frac{\sqrt[4]{a+i}\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{-a+i}\sqrt[4]{1-i(a+bx)}}\right)}{(-a+i)^{3/4}(a+i)^{5/4}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((I/2)*\operatorname{ArcTan}[a + b*x])}/x^2, x]$

[Out] $-\left(\frac{(I+a+b*x)*(1+I*(a+b*x))^{(1/4)}}{(I+a)*x*(1-I*(a+b*x))^{(1/4)}}\right) + \left(\frac{I*b*\operatorname{ArcTan}\left[\frac{(I+a)^{(1/4)*(1+I*(a+b*x))^{(1/4)}}{(I-a)^{(1/4)*(1-I*(a+b*x))^{(1/4)}}$

Rule 211

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[a/b, 2])/a]*\operatorname{ArcTan}[x/\operatorname{Rt}[a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ \operatorname{PosQ}[a/b]$

Rule 214

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(\operatorname{Rt}[-a/b, 2])/a]*\operatorname{ArcTanh}[x/\operatorname{Rt}[-a/b, 2]], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ \operatorname{NegQ}[a/b]$

Rule 218

$\operatorname{Int}[(a_+ + (b_+)*(x_+)^4)^{-1}, x_Symbol] \rightarrow \operatorname{With}\{r = \operatorname{Numerator}[\operatorname{Rt}[-a/b, 2]], s = \operatorname{Denominator}[\operatorname{Rt}[-a/b, 2]]\}, \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r - s*x^2), x], x] + \operatorname{Dist}[r/(2*a), \operatorname{Int}[1/(r + s*x^2), x], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ \operatorname{!GtQ}[a/b]$

, 0]

Rule 269

Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Int[x^(m + n*p)*
(b + a/x^n)^p, x] /; FreeQ[{a, b, m, n}, x] && IntegerQ[p] && NegQ[n]

Rule 294

Int[((c_)*(x_))^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]

Rule 5202

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*(x_)^(m_), x_Symbol] := Dist
[4/(I^m*n*b^(m + 1)*c^(m + 1)), Subst[Int[x^(2/(I*n))*((1 - I*a*c - (1 + I*
a*c)*x^(2/(I*n)))^m/(1 + x^(2/(I*n)))^(m + 2)), x], x, (1 - I*c*(a + b*x))^
(I*(n/2))/(1 + I*c*(a + b*x)^(I*(n/2))], x] /; FreeQ[{a, b, c}, x] && ILtQ
[m, 0] && LtQ[-1, I*n, 1]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{\frac{1}{2}i \tan^{-1}(a+bx)}}{x^2} dx &= (8ib) \text{Subst} \left(\int \frac{1}{(1 - ia - \frac{1+ia}{x^4})^2 x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) \\
 &= (8ib) \text{Subst} \left(\int \frac{x^4}{(-1 - ia + (1 - ia)x^4)^2} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right) \\
 &= -\frac{(i+a+bx)\sqrt[4]{1+i(a+bx)}}{(i+a)x\sqrt[4]{1-i(a+bx)}} - \frac{(2b) \text{Subst} \left(\int \frac{1}{-1-ia+(1-ia)x^4} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right)}{i+a} \\
 &= -\frac{(i+a+bx)\sqrt[4]{1+i(a+bx)}}{(i+a)x\sqrt[4]{1-i(a+bx)}} + \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{i-a}-\sqrt{i+a} x^2} dx, x, \frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right)}{\sqrt{i-a}(1-ia)} \\
 &= -\frac{(i+a+bx)\sqrt[4]{1+i(a+bx)}}{(i+a)x\sqrt[4]{1-i(a+bx)}} + \frac{ib \tan^{-1} \left(\frac{\sqrt[4]{i+a}\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{i-a}\sqrt[4]{1-i(a+bx)}} \right)}{(i-a)^{3/4}(i+a)^{5/4}} + \frac{ib \tanh^{-1} \left(\frac{\sqrt[4]{1+i(a+bx)}}{\sqrt[4]{1-i(a+bx)}} \right)}{(i-a)^{3/4}(i+a)^{5/4}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 110, normalized size = 0.54

$$\frac{(-i(i+a+bx))^{3/4} \left(3(i+a)(-i+a+bx) + 2ibx {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right) \right)}{3(i+a)^2 x (1+ia+ibx)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((I/2)*ArcTan[a + b*x])/x^2,x]

[Out] (((-I)*(I + a + b*x))^(3/4)*(3*(I + a)*(-I + a + b*x) + (2*I)*b*x*Hypergeometric2F1[3/4, 1, 7/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)]))/(3*(I + a)^2*x*(1 + I*a + I*b*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x)

[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x, algorithm="maxima")

[Out] integrate(sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))/x^2, x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 598 vs. 2(141) = 282.

time = 3.26, size = 598, normalized size = 2.92

$$\frac{(-i(i+a+bx))^{3/4} \left(3(i+a)(-i+a+bx) + 2ibx {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right) \right)}{3(i+a)^2 x (1+ia+ibx)^{3/4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x, algorithm="fricas")

```
[Out] 1/2*((-b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1))
^(1/4)*(-I*a + 1)*x*log((b*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/(b*x +
a + I)) + (-b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a
- 1))^(1/4)*(a^2 + 1))/b) + (-b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^
3 - 2*a^2 + 2*I*a - 1))^(1/4)*(I*a - 1)*x*log((b*sqrt(I*sqrt(b^2*x^2 + 2*a*
b*x + a^2 + 1))/(b*x + a + I)) - (-b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*
I*a^3 - 2*a^2 + 2*I*a - 1))^(1/4)*(a^2 + 1))/b) - (-b^4/(a^8 + 2*I*a^7 + 2*
a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1))^(1/4)*(a + I)*x*log((b*sqrt(I
*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/(b*x + a + I)) - (-b^4/(a^8 + 2*I*a^7 +
2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1))^(1/4)*(I*a^2 + I))/b) + (-b
^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1))^(1/4)*(
a + I)*x*log((b*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/(b*x + a + I)) - (
-b^4/(a^8 + 2*I*a^7 + 2*a^6 + 6*I*a^5 + 6*I*a^3 - 2*a^2 + 2*I*a - 1))^(1/4)
*(-I*a^2 - I))/b) - 2*(b*x + a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1
))/(b*x + a + I))/((a + I)*x)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2)/x**2,x)
```

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x, algorithm="giac"
)
```

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\sqrt{\frac{1 + a \operatorname{li} + b x \operatorname{li}}{\sqrt{(a + b x)^2 + 1}}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2)/x^2,x)
```

```
[Out] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2)/x^2, x)
```

3.221 $\int e^{\frac{3}{2}i \operatorname{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=494

$$\frac{(17i + 36a - 24ia^2) \sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{24b^3} - \frac{(3i + 8a) \sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{7/4}}{12b^3} + \frac{x \sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b^3}$$

[Out] $-1/24*(17*I+36*a-24*I*a^2)*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}/b^3-1/12*(3*I+8*a)*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(7/4)}/b^3+1/3*x*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(7/4)}/b^2+1/16*(17*I+36*a-24*I*a^2)*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^3-1/16*(17*I+36*a-24*I*a^2)*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^3-1/32*(17*I+36*a-24*I*a^2)*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)}/b^3-1/32*(17*I+36*a-24*I*a^2)*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)}/b^3$

Rubi [A]

time = 0.29, antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5203, 92, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{(-24a^2 + 36a + 17i) \operatorname{ArcTan}\left(\frac{1 - \sqrt{2} \sqrt{1 - ia - ibx}}{2 + ia + ibx}\right)}{8\sqrt{2}b^3} - \frac{(-24a^2 + 36a + 17i) \operatorname{ArcTan}\left(\frac{1 + \sqrt{2} \sqrt{1 - ia - ibx}}{2 + ia + ibx}\right)}{8\sqrt{2}b^3} - \frac{(-24a^2 + 36a + 17i) \sqrt{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{24b^3} - \frac{(-24a^2 + 36a + 17i) \ln\left(\frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} + 1\right)}{16\sqrt{2}b^3} - \frac{(-24a^2 + 36a + 17i) \ln\left(\frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} - 1\right)}{16\sqrt{2}b^3} - \frac{(3i + 8a) \sqrt{1 - ia - ibx} (1 + ia + ibx)^{7/4}}{12b^3} + \frac{x^2 \sqrt{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b^3}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)/2)*\operatorname{ArcTan}[a + b*x]}*x^2, x]$

[Out] $-1/24*((17*I + 36*a - (24*I)*a^2)*(1 - I*a - I*b*x)^{(1/4)}*(1 + I*a + I*b*x)^{(3/4)}/b^3 - ((3*I + 8*a)*(1 - I*a - I*b*x)^{(1/4)}*(1 + I*a + I*b*x)^{(7/4)})/(12*b^3) + (x*(1 - I*a - I*b*x)^{(1/4)}*(1 + I*a + I*b*x)^{(7/4)})/(3*b^2) + ((17*I + 36*a - (24*I)*a^2)*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(8*\operatorname{Sqrt}[2]*b^3) - ((17*I + 36*a - (24*I)*a^2)*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(8*\operatorname{Sqrt}[2]*b^3) + ((17*I + 36*a - (24*I)*a^2)*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a - I*b*x]/\operatorname{Sqrt}[1 + I*a + I*b*x] - (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(16*\operatorname{Sqrt}[2]*b^3) - ((17*I + 36*a - (24*I)*a^2)*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a - I*b*x]/\operatorname{Sqrt}[1 + I*a + I*b*x] + (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(16*\operatorname{Sqrt}[2]*b^3)$

Rule 52

$\operatorname{Int}[(a + b*x)^m * ((c + d*x)^n)^m, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{m+1} * ((c + d*x)^n / (b*(m + n + 1))), x] + \operatorname{Dist}[n * ((b*c - a*d) / (b*(m + n + 1))), \operatorname{Int}[(a + b*x)^m * (c + d*x)^{n-1}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ

$[m, 0] \&\& (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0])) \ \&\& \ !\text{ILtQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \ :> \ \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)}(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}], x]] \ /; \ \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 81

$\text{Int}[(a_.) + (b_.)(x_)^{(c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_Symbol] \ :> \ \text{Simp}[b*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 2, 0]$

Rule 92

$\text{Int}[(a_.) + (b_.)(x_)^2((c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_Symbol] \ :> \ \text{Simp}[b*(a + b*x)*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+3))), x] + \text{Dist}[1/(d*f*(n+p+3)), \text{Int}[(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(n+p+3) - b*(b*c*e + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(n+p+4) - b*(d*e*(n+2) + c*f*(p+2))]*x, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 3, 0]$

Rule 210

$\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \ :> \ \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \ /; \ \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 217

$\text{Int}[(a_) + (b_.)(x_)^4)^{-1}, x_Symbol] \ :> \ \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*r), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x]] \ /; \ \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 246

$\text{Int}[(a_) + (b_.)(x_)^{(n_)}^{(p_)}, x_Symbol] \ :> \ \text{Dist}[a^{(p+1/n)}, \text{Subst}[\text{Int}[1/(1 - b*x^n)^{(p+1/n+1)}, x], x, x/(a + b*x^n)^{(1/n)}], x] \ /; \ \text{FreeQ}[\{a,$

b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_.))*((d_) + (e_)*(x_)^(m_.)), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{\frac{3}{2}i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx \\
&= \frac{x^4 \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{3b^2} + \frac{\int \frac{(1+ia+ibx)^{3/4} (-1-a^2-\frac{1}{2}(3i+8a)bx)}{(1-ia-ibx)^{3/4}} dx}{3b^2} \\
&= -\frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{12b^3} + \frac{x^4 \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{3b^2} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3} \\
&= -\frac{(17i+36a-24ia^2) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{24b^3} - \frac{(3i+8a) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{12b^3}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 121, normalized size = 0.24

$$\frac{\sqrt[4]{-i(i+a+bx)} (-i(1+ia+ibx)^{3/4} (3+8a^2+7ibx-4b^2x^2+a(-5i+4bx)) + 2i2^{3/4}(-17+36ia+24a^2) {}_2F_1(-\frac{3}{4}, \frac{1}{4}, \frac{5}{4}, -\frac{1}{2}i(i+a+bx))}{12b^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a + b*x])*x^2,x]

[Out] (((-I)*(I + a + b*x))^(1/4))*((-I)*(1 + I*a + I*b*x)^(3/4))*(3 + 8*a^2 + (7*I)*b*x - 4*b^2*x^2 + a*(-5*I + 4*b*x)) + (2*I)*2^(3/4)*(-17 + (36*I)*a + 24*a^2)*Hypergeometric2F1[-3/4, 1/4, 5/4, (-1/2*I)*(I + a + b*x)]/(12*b^3)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}} \right)^{\frac{3}{2}} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x^2,x)``[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x^2,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x^2,x, algorithm="maxima")``[Out] integrate(x^2*((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2), x)`**Fricas [A]**

time = 3.02, size = 561, normalized size = 1.14

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x^2,x, algorithm="fricas")`

```
[Out] 1/48*(3*b^3*sqrt((576*I*a^4 - 1728*a^3 - 2112*I*a^2 + 1224*a + 289*I)/b^6)*
log((b^3*sqrt((576*I*a^4 - 1728*a^3 - 2112*I*a^2 + 1224*a + 289*I)/b^6) + (
24*a^2 + 36*I*a - 17)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I
)))/(24*a^2 + 36*I*a - 17)) - 3*b^3*sqrt((576*I*a^4 - 1728*a^3 - 2112*I*a^2
+ 1224*a + 289*I)/b^6)*log(-(b^3*sqrt((576*I*a^4 - 1728*a^3 - 2112*I*a^2 +
1224*a + 289*I)/b^6) - (24*a^2 + 36*I*a - 17)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*
x + a^2 + 1)/(b*x + a + I)))/(24*a^2 + 36*I*a - 17)) - 3*b^3*sqrt((-576*I*a
^4 + 1728*a^3 + 2112*I*a^2 - 1224*a - 289*I)/b^6)*log((b^3*sqrt((-576*I*a^4
+ 1728*a^3 + 2112*I*a^2 - 1224*a - 289*I)/b^6) + (24*a^2 + 36*I*a - 17)*sq
rt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(24*a^2 + 36*I*a - 1
7)) + 3*b^3*sqrt((-576*I*a^4 + 1728*a^3 + 2112*I*a^2 - 1224*a - 289*I)/b^6)
*log(-(b^3*sqrt((-576*I*a^4 + 1728*a^3 + 2112*I*a^2 - 1224*a - 289*I)/b^6)
- (24*a^2 + 36*I*a - 17)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a
```

+ I)))/(24*a^2 + 36*I*a - 17)) + 2*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(8*I*b^2*x^2 - 2*(4*I*a - 7)*b*x + 8*I*a^2 - 46*a - 23*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/b^3

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2)*x**2,x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3004 deep

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 \left(\frac{1 + a \operatorname{li} + b x \operatorname{li}}{\sqrt{(a + b x)^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2),x)

[Out] int(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)

3.222 $\int e^{\frac{3}{2}i \operatorname{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=410

$$\frac{(3-4ia)\sqrt[4]{1-ia-ibx}(1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx}(1+ia+ibx)^{7/4}}{2b^2} - \frac{3(3-4ia)\operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2}b^2}$$

[Out] $\frac{1}{4}(3-4Ia)(1-Ia-Ib*x)^{1/4}(1+Ia+Ib*x)^{3/4}/b^2 + \frac{1}{2}(1-Ia-Ib*x)^{1/4}(1+Ia+Ib*x)^{7/4}/b^2 - \frac{3}{8}(3-4Ia)\operatorname{arctan}\left(\frac{1-(1-Ia-Ib*x)^{1/4}}{(1+Ia+Ib*x)^{1/4}}\right)/b^2 + \frac{3}{8}(3-4Ia)\operatorname{arctan}\left(\frac{1+(1-Ia-Ib*x)^{1/4}}{(1+Ia+Ib*x)^{1/4}}\right)/b^2 - \frac{3}{16}(3-4Ia)\ln\left(\frac{1-(1-Ia-Ib*x)^{1/4}}{(1+Ia+Ib*x)^{1/4}}\right)/b^2 + \frac{3}{16}(3-4Ia)\ln\left(\frac{1+(1-Ia-Ib*x)^{1/4}}{(1+Ia+Ib*x)^{1/4}}\right)/b^2$

Rubi [A]

time = 0.21, antiderivative size = 410, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {5203, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{3(3-4ia)\operatorname{ArcTan}\left(\frac{1-\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2}b^2} + \frac{3(3-4ia)\operatorname{ArcTan}\left(\frac{1+\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2}b^2} + \frac{\sqrt[4]{1-ia-ibx}(1+ia+ibx)^{3/4}}{2b^2} + \frac{(3-4ia)\sqrt[4]{1-ia-ibx}(1+ia+ibx)^{7/4}}{4b^2} - \frac{3(3-4ia)\log\left(\frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} + \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} + 1\right)}{8\sqrt{2}b^2} + \frac{3(3-4ia)\log\left(\frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} - \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} + 1\right)}{8\sqrt{2}b^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3I)/2)\operatorname{ArcTan}[a+bx]}x, x]$

[Out] $((3-(4I)a)(1-Ia-Ib*x)^{1/4}(1+Ia+Ib*x)^{3/4})/(4b^2) + ((1-Ia-Ib*x)^{1/4}(1+Ia+Ib*x)^{7/4})/(2b^2) - (3(3-(4I)a)\operatorname{ArcTan}[1-(\sqrt{2}(1-Ia-Ib*x)^{1/4})/(1+Ia+Ib*x)^{1/4}])/(4\sqrt{2}b^2) + (3(3-(4I)a)\operatorname{ArcTan}[1+(\sqrt{2}(1-Ia-Ib*x)^{1/4})/(1+Ia+Ib*x)^{1/4}])/(4\sqrt{2}b^2) - (3(3-(4I)a)\operatorname{Log}[1+\sqrt{2}(1-Ia-Ib*x)^{1/4}/(1+Ia+Ib*x)^{1/4}])/(8\sqrt{2}b^2) + (3(3-(4I)a)\operatorname{Log}[1+\sqrt{2}(1-Ia-Ib*x)^{1/4}/(1+Ia+Ib*x)^{1/4}])/(8\sqrt{2}b^2)$

Rule 52

$\operatorname{Int}[(a_. + (b_.)(x_.))^{(m_.)}((c_.) + (d_.)(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + bx)^{(m+1)}((c + dx)^n/(b(m+n+1))), x] + \operatorname{Dist}[n((bc - ad)/(b(m+n+1))), \operatorname{Int}[(a + bx)^m(c + dx)^{(n-1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
```

e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_.)*((d_) + (e_)*(x_)^m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{\frac{3}{2}i \tan^{-1}(a+bx)} x dx &= \int \frac{x(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx \\
&= \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} - \frac{(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} - \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} + \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} + \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} + \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} + \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} + \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} - \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b} \\
&= \frac{(3-4ia)\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{7/4}}{2b^2} - \frac{(3(3i+4a) \int \frac{(1+ia+ibx)^{3/4}}{(1-ia-ibx)^{3/4}} dx)}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 79, normalized size = 0.19

$$\frac{\sqrt[4]{-i(i+a+bx)} \left((1+ia+ibx)^{7/4} + 2 \cdot 2^{3/4} (3-4ia) {}_2F_1\left(-\frac{3}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{1}{2}i(i+a+bx)\right) \right)}{2b^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a + b*x])*x,x]

[Out] (((-I)*(I + a + b*x))^(1/4)*((1 + I*a + I*b*x)^(7/4) + 2*2^(3/4)*(3 - (4*I)*a)*Hypergeometric2F1[-3/4, 1/4, 5/4, (-1/2*I)*(I + a + b*x)]))/(2*b^2)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}} \right)^{\frac{3}{2}} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x,x)

[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x,x, algorithm="maxima")

[Out] integrate(x*((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2), x)

Fricas [A]

time = 2.95, size = 431, normalized size = 1.05

$$\frac{3b^2 \sqrt{24a^2 - 24a - 9I} \log\left(\frac{\sqrt{24a^2 - 24a - 9I} \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 3I}\right) - 3b^2 \sqrt{24a^2 - 24a - 9I} \log\left(\frac{\sqrt{24a^2 - 24a - 9I} \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 3I}\right) - 3b^2 \sqrt{24a^2 - 24a - 9I} \log\left(\frac{\sqrt{24a^2 - 24a - 9I} \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 3I}\right) + 3b^2 \sqrt{24a^2 - 24a - 9I} \log\left(\frac{\sqrt{24a^2 - 24a - 9I} \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 3I}\right) + 2\sqrt{24a^2 - 24a - 9I} \sqrt{b^2 x^2 + 2abx + a^2 + 1} \sqrt{2Ibx - 2Ia + 5} \sqrt{b^2 x^2 + 2abx + a^2 + 1}}{b^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x,x, algorithm="fricas")

[Out] 1/8*(3*b^2*sqrt(-(16*I*a^2 - 24*a - 9*I)/b^4)*log((b^2*sqrt(-(16*I*a^2 - 24*a - 9*I)/b^4) + (4*a + 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a + 3*I)) - 3*b^2*sqrt(-(16*I*a^2 - 24*a - 9*I)/b^4)*log(-(b^2*sqrt(-(16*I*a^2 - 24*a - 9*I)/b^4) - (4*a + 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a + 3*I)) - 3*b^2*sqrt(-(-16*I*a^2 + 24*a + 9*I)/b^4)*log((b^2*sqrt(-(-16*I*a^2 + 24*a + 9*I)/b^4) + (4*a + 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a + 3*I)) + 3*b^2*sqrt(-(-16*I*a^2 + 24*a + 9*I)/b^4)*log(-(b^2*sqrt(-(-16*I*a^2 + 24*a + 9*I)/b^4) - (4*a + 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a + 3*I)) + 2*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(2*I*b*x - 2*I*a + 5)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b^2

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2)*x,x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)*x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int x \left(\frac{1 + a \operatorname{li} + b x \operatorname{li}}{\sqrt{(a + b x)^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)
```


3.223 $\int e^{\frac{3}{2}i \operatorname{ArcTan}(a+bx)} dx$

Optimal. Leaf size=338

$$\frac{i\sqrt[4]{1-ia-ibx}(1+ia+ibx)^{3/4}}{b} - \frac{3i \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b} + \frac{3i \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b}$$

[Out] $I*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}/b-3/2*I*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}+3/2*I*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}-3/4*I*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}+3/4*I*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 338, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$,

Rules used = {5201, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{3i \operatorname{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} + \frac{3i \operatorname{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} + \frac{i\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/4}}{b} - \frac{3i \log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b} + \frac{3i \log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] `Int[E^(((3*I)/2)*ArcTan[a + b*x]),x]`

[Out] $(I*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)})/b - ((3*I)*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*b) + ((3*I)*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*b) - (((3*I)/2)*\operatorname{Log}[1 + \operatorname{Sqrt}[1-I*a-I*b*x]/\operatorname{Sqrt}[1+I*a+I*b*x] - (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*b) + (((3*I)/2)*\operatorname{Log}[1 + \operatorname{Sqrt}[1-I*a-I*b*x]/\operatorname{Sqrt}[1+I*a+I*b*x] + (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}])/(\operatorname{Sqrt}[2]*b)$

Rule 52

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +`

```
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5201

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]

Rubi steps

$$\begin{aligned}
 \int e^{\frac{3}{2}i \tan^{-1}(a+bx)} dx &= \int \frac{(1 + ia + ibx)^{3/4}}{(1 - ia - ibx)^{3/4}} dx \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} + \frac{3}{2} \int \frac{1}{(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}} dx \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} + \frac{(6i) \text{Subst}\left(\int \frac{1}{\sqrt[4]{2 - x^4}} dx, x, \sqrt[4]{1 - ia - ibx}\right)}{b} \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} + \frac{(6i) \text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{b} \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} + \frac{(3i) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{b} + \frac{(3i)}{b} \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} + \frac{(3i) \text{Subst}\left(\int \frac{1}{1-\sqrt{2}xx^2} dx, x, \frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{2b} \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} - \frac{3i \log\left(1 + \frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} - \frac{\sqrt{2} \sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{2\sqrt{2} b} \\
 &= \frac{i\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{b} - \frac{3i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{\sqrt{2} b} + \frac{3i \tan^{-1}\left(\frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{b}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 45, normalized size = 0.13

$$\frac{8ie^{\frac{7}{2}i \text{ArcTan}(a+bx)} {}_2F_1\left(\frac{7}{4}, 2; \frac{11}{4}; -e^{2i \text{ArcTan}(a+bx)}\right)}{7b}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a + b*x]),x]

[Out] (((-8*I)/7)*E^(((7*I)/2)*ArcTan[a + b*x])*Hypergeometric2F1[7/4, 2, 11/4, -E^((2*I)*ArcTan[a + b*x])]/b

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \left(\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}} \right)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x)

[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="maxima")

[Out] integrate(((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2), x)

Fricas [A]

time = 3.58, size = 268, normalized size = 0.79

$$\frac{b\sqrt{\frac{9i}{b^2}} \log\left(\frac{1}{3}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - b\sqrt{\frac{9i}{b^2}} \log\left(-\frac{1}{3}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - b\sqrt{\frac{9i}{b^2}} \log\left(\frac{1}{3}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) + b\sqrt{\frac{9i}{b^2}} \log\left(-\frac{1}{3}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) + 2i\sqrt{b^2x^2 + 2abx + a^2 + 1} \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="fricas")

[Out] 1/2*(b*sqrt(9*I/b^2)*log(1/3*b*sqrt(9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(9*I/b^2)*log(-1/3*b*sqrt(9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(-9*I/b^2)*log(1/3*b*sqrt(-9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + b*sqrt(-9*I/b^2)*log(-1/3*b*sqrt(-9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 2*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2),x)
```

```
[Out] Timed out
```

Giac [F(-2)]

```
time = 0.00, size = 0, normalized size = 0.00
```

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

```
time = 0.00, size = -1, normalized size = -0.00
```

$$\int \left(\frac{1 + a i + b x i}{\sqrt{(a + b x)^2 + 1}} \right)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2),x)
```

```
[Out] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)
```

$$3.224 \quad \int \frac{e^{\frac{3}{2}i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=427

$$\frac{2(i-a)^{3/4} \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{(i+a)^{3/4}} + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)$$

[Out] $2*(I-a)^{(3/4)}*\arctan((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4)))/(I+a)^{(3/4)}-2*(I-a)^{(3/4)}*\operatorname{arctanh}((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4)))/(I+a)^{(3/4)}+1/2*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)})/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2))*2^{(1/2)}-1/2*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)})/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2))*2^{(1/2)}+\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)})/(1+I*a+I*b*x)^{(1/4)}*2^{(1/2)}-\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)})/(1+I*a+I*b*x)^{(1/4)}*2^{(1/2)}$

Rubi [A]

time = 0.21, antiderivative size = 427, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 15, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5203, 132, 65, 246, 217, 1179, 642, 1176, 631, 210, 12, 95, 304, 211, 214}

$$\frac{2(-a+i)^{3/4} \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ibx+1}}{\sqrt[4]{i-a} \sqrt[4]{1-ibx+1}}\right)}{(a+i)^{3/4}} + \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{i-a-ibx+1}}{\sqrt[4]{i+ibx+1}}\right) - \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{i-a-ibx+1}}{\sqrt[4]{i+ibx+1}}\right) + \frac{\log\left(\frac{\sqrt[4]{i-a-ibx+1}}{\sqrt[4]{i+ibx+1}} - \frac{\sqrt{2} \sqrt[4]{i-a-ibx+1}}{\sqrt[4]{i+ibx+1}} + 1\right)}{\sqrt{2}} - \frac{\log\left(\frac{\sqrt[4]{i-a-ibx+1}}{\sqrt[4]{i+ibx+1}} + \frac{\sqrt{2} \sqrt[4]{i-a-ibx+1}}{\sqrt[4]{i+ibx+1}} + 1\right)}{\sqrt{2}} - \frac{2(-a+i)^{3/4} \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ibx+1}}{\sqrt[4]{i-a} \sqrt[4]{1-ibx+1}}\right)}{(a+i)^{3/4}}$$

Antiderivative was successfully verified.

[In] Int[E^(((3*I)/2)*ArcTan[a + b*x])/x,x]

[Out] $(2*(I-a)^{(3/4)}*\operatorname{ArcTan}(((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)))/((I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4))})/(I+a)^{(3/4)} + \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}] - \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}] - (2*(I-a)^{(3/4)}*\operatorname{ArcTan}(((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)))/((I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4))})/(I+a)^{(3/4)} + \operatorname{Log}[1 + \operatorname{Sqrt}[1-I*a-I*b*x]/\operatorname{Sqrt}[1+I*a+I*b*x] - (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}]/\operatorname{Sqrt}[2] - \operatorname{Log}[1 + \operatorname{Sqrt}[1-I*a-I*b*x]/\operatorname{Sqrt}[1+I*a+I*b*x] + (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}]/\operatorname{Sqrt}[2]$

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 65

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b) +

$d*(x^p/b)^n, x, (a + b*x)^{1/p}, x] /; \text{FreeQ}\{a, b, c, d, x\} \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 95

$\text{Int}[(((a_.) + (b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.))}/((e_.) + (f_.)*(x_))), x_Symbol] \rightarrow \text{With}\{q = \text{Denominator}[m]\}, \text{Dist}[q, \text{Subst}[\text{Int}[x^{q*(m+1)-1}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{1/q}/(c + d*x)^{1/q}], x] /; \text{FreeQ}\{a, b, c, d, e, f, x\} \ \&\& \ \text{EqQ}[m + n + 1, 0] \ \&\& \ \text{RationalQ}[n] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{SimplerQ}[a + b*x, c + d*x]$

Rule 132

$\text{Int}[((a_.) + (b_.)*(x_))^{(m_.)*((c_.) + (d_.)*(x_))^{(n_.)*((e_.) + (f_.)*(x_))^{(p_.)}}, x_Symbol] \rightarrow \text{Dist}[b*d^{(m+n)}*f^p, \text{Int}[(a + b*x)^{(m-1)}/(c + d*x)^m, x], x] + \text{Int}[(a + b*x)^{(m-1)}*(e + f*x)^p/(c + d*x)^m*\text{ExpandToSum}[(a + b*x)*(c + d*x)^{-(p-1)} - (b*d^{-(p-1)}*f^p)/(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, m, n\}, x\} \ \&\& \ \text{EqQ}[m + n + p + 1, 0] \ \&\& \ \text{ILtQ}[p, 0] \ \&\& \ (\text{GtQ}[m, 0] \ || \ \text{SumSimplerQ}[m, -1] \ || \ !(\text{GtQ}[n, 0] \ || \ \text{SumSimplerQ}[n, -1]))$

Rule 210

$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-(\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 211

$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[a/b, 2]/a)*\text{ArcTan}[x/\text{Rt}[a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{PosQ}[a/b]$

Rule 214

$\text{Int}[((a_) + (b_.)*(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(\text{Rt}[-a/b, 2]/a)*\text{ArcTanh}[x/\text{Rt}[-a/b, 2]], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ \text{NegQ}[a/b]$

Rule 217

$\text{Int}[((a_) + (b_.)*(x_)^4)^{-1}, x_Symbol] \rightarrow \text{With}\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*r), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] /; \text{FreeQ}\{a, b, x\} \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]
```

Rule 304

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_)^(m_.)), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\frac{3}{2}i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{(1+ia+ibx)^{3/4}}{x(1-ia-ibx)^{3/4}} dx \\
&= -\left((-1-ia) \int \frac{1}{x(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}} dx \right) + (ib) \int \frac{1}{(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}} dx \\
&= -\left(4\text{Subst}\left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-ia-ibx} \right) \right) + (4(1+ia))\text{Subst}\left(\int \frac{1}{-1-ia-\dots} \right) \\
&= -\left(4\text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} \right) \right) - \frac{(2(i-a))\text{Subst}\left(\int \frac{1}{\sqrt{i-a}-\sqrt{i+a}} \right)}{\sqrt{i+a}} \\
&= \frac{2(i-a)^{3/4} \tan^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i+a)^{3/4}} - \frac{2(i-a)^{3/4} \tanh^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia}}{\sqrt[4]{i-a} \sqrt[4]{1-ia}} \right)}{(i+a)^{3/4}} \\
&= \frac{2(i-a)^{3/4} \tan^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i+a)^{3/4}} - \frac{2(i-a)^{3/4} \tanh^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia}}{\sqrt[4]{i-a} \sqrt[4]{1-ia}} \right)}{(i+a)^{3/4}} \\
&= \frac{2(i-a)^{3/4} \tan^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i+a)^{3/4}} - \frac{2(i-a)^{3/4} \tanh^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia}}{\sqrt[4]{i-a} \sqrt[4]{1-ia}} \right)}{(i+a)^{3/4}} \\
&= \frac{2(i-a)^{3/4} \tan^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i+a)^{3/4}} + \sqrt{2} \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 122, normalized size = 0.29

$$2\sqrt[4]{-i(i+a+bx)} \left(-2^{3/4} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{1}{2}i(i+a+bx)\right) + \frac{2(-i+a) {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right)}{(i+a)\sqrt[4]{1+ia+ibx}} \right)$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a + b*x])/x,x]

[Out] 2*((-I)*(I + a + b*x))^(1/4)*(-(2^(3/4)*Hypergeometric2F1[1/4, 1/4, 5/4, (-1/2*I)*(I + a + b*x)]) + (2*(-I + a)*Hypergeometric2F1[1/4, 1, 5/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)]))/((I + a)*(1 + I*a + I*b*x)^(1/4)))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}} \right)^{\frac{3}{2}}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x)**[Out]** int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x, algorithm="maxima")**[Out]** integrate(((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2)/x, x)**Fricas [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 690 vs. 2(284) = 568.

time = 2.65, size = 690, normalized size = 1.62

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x, algorithm="fricas")

[Out] 1/2*sqrt(4*I)*log(1/2*I*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(4*I)*log(-1/2*I*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(-4*I)*log(1/2*I*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 1/2*sqrt(-4*I)*log(-1/2*I*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - ((a^3 - 3*I*a^2 - 3*a + I)/(a^3 + 3*I*a^2 - 3*a - I))^(1/4)*log(((a^2 + 2*I*a - 1)*(-(a^3 - 3*I*a^2 - 3*a + I)/(a^3 + 3*I*a^2 - 3*a - I)))^(3/4) + (a^2 - 2*I*a - 1)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(a^2 - 2*I*a - 1) + ((-a^3 - 3*I*a^2 - 3*a + I)/(a^3 + 3*I*a^2 - 3*a - I))^(1/4)*log(-((a^2 + 2*I*a - 1)*(-(a^3 - 3*I*a^2 - 3*a + I)/(a^3 + 3*I*a^2 - 3*a - I)))^(3/4) - (a^2 - 2*I*a - 1)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x

+ a² + 1)/(b*x + a + I)))/(a² - 2*I*a - 1)) + I*(-(a³ - 3*I*a² - 3*a + I)/(a³ + 3*I*a² - 3*a - I))^(1/4)*log(((I*a² - 2*a - I)*(-(a³ - 3*I*a² - 3*a + I)/(a³ + 3*I*a² - 3*a - I))^(3/4) + (a² - 2*I*a - 1)*sqrt(I*sqrt(b²*x² + 2*a*b*x + a² + 1)/(b*x + a + I)))/(a² - 2*I*a - 1)) - I*(-(a³ - 3*I*a² - 3*a + I)/(a³ + 3*I*a² - 3*a - I))^(1/4)*log(((I*a² - 2*a - I)*(-(a³ - 3*I*a² - 3*a + I)/(a³ + 3*I*a² - 3*a - I))^(3/4) + (a² - 2*I*a - 1)*sqrt(I*sqrt(b²*x² + 2*a*b*x + a² + 1)/(b*x + a + I)))/(a² - 2*I*a - 1))

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2)/x,x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+ai+bx}{\sqrt{(a+bx)^2+1}} \right)^{3/2}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*i + b*x*i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2)/x,x)

[Out] int(((a*i + b*x*i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2)/x, x)

3.225 $\int \frac{e^{\frac{3}{2}i \operatorname{ArcTan}(a+bx)}}{x^2} dx$

Optimal. Leaf size=211

$$\frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{(1-ia)x} - \frac{3ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{\sqrt[4]{i-a} (i+a)^{7/4}} + \frac{3ib \tanh^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{\sqrt[4]{i-a} (i+a)^{7/4}}$$

[Out] $-(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}/(1-I*a)/x-3*I*b*\arctan((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4)})/(I-a)^{(1/4)}/(I+a)^{(7/4)}+3*I*b*\operatorname{arctanh}((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4)})/(I-a)^{(1/4)}/(I+a)^{(7/4)}$

Rubi [A]

time = 0.12, antiderivative size = 211, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5203, 96, 95, 304, 211, 214}

$$-\frac{3ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{a+i} \sqrt[4]{ia+ibx+1}}{\sqrt[4]{-a+i} \sqrt[4]{-ia-ibx+1}}\right)}{\sqrt[4]{-a+i} (a+i)^{7/4}} - \frac{\sqrt[4]{-ia-ibx+1} (ia+ibx+1)^{3/4}}{(1-ia)x} + \frac{3ib \tanh^{-1}\left(\frac{\sqrt[4]{a+i} \sqrt[4]{ia+ibx+1}}{\sqrt[4]{-a+i} \sqrt[4]{-ia-ibx+1}}\right)}{\sqrt[4]{-a+i} (a+i)^{7/4}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((3*I)/2)*\operatorname{ArcTan}[a + b*x]}/x^2, x]$

[Out] $-\left(\frac{(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}}{(1-I*a)*x}\right) - \left(\frac{3*I*b*\operatorname{ArcTan}\left[\frac{(I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}}{(I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4)}}\right]}{(I-a)^{(1/4)}*(I+a)^{(7/4)}} + \frac{3*I*b*\operatorname{ArcTanh}\left[\frac{(I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}}{(I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4)}}\right]}{(I-a)^{(1/4)}*(I+a)^{(7/4)}}\right)$

Rule 95

$\operatorname{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{With}[q = \operatorname{Denominator}[m], \operatorname{Dist}[q, \operatorname{Subst}[\operatorname{Int}[x^{(q*(m+1)-1)}/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^{(1/q)}/(c + d*x)^{(1/q)}], x]] /; \operatorname{FreeQ}\{a, b, c, d, e, f\}, x] \&\& \operatorname{EqQ}[m + n + 1, 0] \&\& \operatorname{RationalQ}[n] \&\& \operatorname{LtQ}[-1, m, 0] \&\& \operatorname{SimplerQ}[a + b*x, c + d*x]$

Rule 96

$\operatorname{Int}[((a_.) + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*(c + d*x)^n*((e + f*x)^{(p+1)})/((m+1)*(b*e - a*f)), x] - \operatorname{Dist}[n*((d*e - c*f)/((m+1)*(b*e - a*f))), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}*(e + f*x)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, p\}, x] \&\& \operatorname{EqQ}[m + n + p + 2, 0] \&\& \operatorname{GtQ}[n, 0] \&\& \operatorname{SumSimpler}$

Q[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 304

Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[s/(2*b), Int[1/(r + s*x^2), x], x] - Dist[s/(2*b), Int[1/(r - s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{\frac{3}{2}i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{(1+ia+ibx)^{3/4}}{x^2(1-ia-ibx)^{3/4}} dx \\
 &= -\frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{(1-ia)x} - \frac{(3b) \int \frac{1}{x(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}} dx}{2(i+a)} \\
 &= -\frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{(1-ia)x} - \frac{(6b) \text{Subst}\left(\int \frac{x^2}{-1-ia-(-1+ia)x^4} dx, x, \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}}\right)}{i+a} \\
 &= -\frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{(1-ia)x} + \frac{(3ib) \text{Subst}\left(\int \frac{1}{\sqrt{i-a}-\sqrt{i+a} x^2} dx, x, \frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}}\right)}{(i+a)^{3/2}} \\
 &= -\frac{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{(1-ia)x} - \frac{3ib \tan^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{\sqrt[4]{i-a} (i+a)^{7/4}} + \frac{3ib \tanh^{-1}\left(\frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1-ia-ibx}}\right)}{\sqrt[4]{i-a} (i+a)^{7/4}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 106, normalized size = 0.50

$$\frac{\sqrt[4]{-i(i+a+bx)} \left(1 + a^2 + ibx + abx + 6ibx {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right)\right)}{(i+a)^2 x \sqrt[4]{1+ia+ibx}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((3*I)/2)*ArcTan[a + b*x])/x^2,x]

[Out] (((-I)*(I + a + b*x))^(1/4)*(1 + a^2 + I*b*x + a*b*x + (6*I)*b*x*Hypergeometric2F1[1/4, 1, 5/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)])) / ((I + a)^2*x*(1 + I*a + I*b*x)^(1/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}\right)^{\frac{3}{2}}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x)

[Out] int(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x, algorithm="maxima")

[Out] integrate(((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2)/x^2, x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 694 vs. 2(137) = 274.

time = 3.97, size = 694, normalized size = 3.29

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x, algorithm="fricas")

[Out] $\frac{1}{2} \cdot (3 \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{1/4} \cdot (-Ia + 1) \cdot x \cdot \log((b^3 \cdot \sqrt{I \cdot \sqrt{b^2x^2 + 2a \cdot bx + a^2 + 1}} / (bx + a + I)) + (a^6 + 4Ia^5 - 5a^4 - 5a^2 - 4Ia + 1) \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{3/4}) / b^3) + 3 \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{1/4} \cdot (Ia - 1) \cdot x \cdot \log((b^3 \cdot \sqrt{I \cdot \sqrt{b^2x^2 + 2a \cdot bx + a^2 + 1}} / (bx + a + I)) - (a^6 + 4Ia^5 - 5a^4 - 5a^2 - 4Ia + 1) \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{3/4}) / b^3) + 3 \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{1/4} \cdot (a + I) \cdot x \cdot \log((b^3 \cdot \sqrt{I \cdot \sqrt{b^2x^2 + 2a \cdot bx + a^2 + 1}} / (bx + a + I)) - (Ia^6 - 4a^5 - 5Ia^4 - 5Ia^2 + 4a + I) \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{3/4}) / b^3) - 3 \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{1/4} \cdot (a + I) \cdot x \cdot \log((b^3 \cdot \sqrt{I \cdot \sqrt{b^2x^2 + 2a \cdot bx + a^2 + 1}} / (bx + a + I)) - (-Ia^6 + 4a^5 + 5Ia^4 + 5Ia^2 - 4a - I) \cdot (-b^4/(a^8 + 6Ia^7 - 14a^6 - 14Ia^5 - 14Ia^3 + 14a^2 + 6Ia - 1))^{3/4}) / b^3) - 2 \cdot I \cdot \sqrt{b^2x^2 + 2a \cdot bx + a^2 + 1} \cdot \sqrt{I \cdot \sqrt{b^2x^2 + 2a \cdot bx + a^2 + 1}} / (bx + a + I)) / ((a + I) \cdot x)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2)/x**2,x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{\left(\frac{1+ai+bx}{\sqrt{(a+bx)^2+1}} \right)^{3/2}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2)/x^2,x)

[Out] int(((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2)/x^2, x)

3.226 $\int e^{-\frac{1}{2}i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=494

$$\frac{(3i - 4a - 8ia^2) \sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}}{8b^3} + \frac{(i - 8a)(1 - ia - ibx)^{5/4} (1 + ia + ibx)^{3/4}}{12b^3} + \frac{x(1 - ia - ibx)^{5/4} (1 + ia + ibx)^{3/4}}{12b^3}$$

[Out] $\frac{1}{8}*(3*I-4*a-8*I*a^2)*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}/b^3+1/12*(I-8*a)*(1-I*a-I*b*x)^{(5/4)}*(1+I*a+I*b*x)^{(3/4)}/b^3+1/3*x*(1-I*a-I*b*x)^{(5/4)}*(1+I*a+I*b*x)^{(3/4)}/b^2+1/16*(3*I-4*a-8*I*a^2)*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^3*2^{(1/2)}-1/16*(3*I-4*a-8*I*a^2)*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^3*2^{(1/2)}+1/32*(3*I-4*a-8*I*a^2)*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b^3*2^{(1/2)}-1/32*(3*I-4*a-8*I*a^2)*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b^3*2^{(1/2)}$

Rubi [A]

time = 0.29, antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5203, 92, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{(-8a^2 - 4a + 3)\text{ArcTan}\left(\frac{1 - \sqrt{2}\sqrt{1 - ia - ibx}}{2a + ibx + 1}\right)}{8\sqrt{2}b} - \frac{(-8a^2 - 4a + 3)\text{ArcTan}\left(\frac{1 + \sqrt{2}\sqrt{1 - ia - ibx}}{2a + ibx + 1}\right)}{8\sqrt{2}b} - \frac{(-8a^2 - 4a + 3)(ia + ibx + 1)^{3/4}\sqrt{1 - ia - ibx}}{32b^3} - \frac{(-8a^2 - 4a + 3)\log\left(\frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} - \frac{\sqrt{2}\sqrt{1 - ia - ibx}}{2a + ibx + 1}\right)}{16\sqrt{2}b} - \frac{(-8a^2 - 4a + 3)\log\left(\frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} + \frac{\sqrt{2}\sqrt{1 - ia - ibx}}{2a + ibx + 1}\right)}{16\sqrt{2}b} - \frac{(-8a + 1)(ia + ibx + 1)^{3/4}(ia - ibx + 1)^{3/4}}{12b^3} + \frac{x(ia + ibx + 1)^{3/4}(ia - ibx + 1)^{3/4}}{12b^3}$$

Antiderivative was successfully verified.

[In] Int[x^2/E^((I/2)*ArcTan[a + b*x]),x]

[Out] $((3*I - 4*a - (8*I)*a^2)*(1 - I*a - I*b*x)^{(1/4)}*(1 + I*a + I*b*x)^{(3/4)})/(8*b^3) + ((I - 8*a)*(1 - I*a - I*b*x)^{(5/4)}*(1 + I*a + I*b*x)^{(3/4)})/(12*b^3) + (x*(1 - I*a - I*b*x)^{(5/4)}*(1 + I*a + I*b*x)^{(3/4)})/(3*b^2) + ((3*I - 4*a - (8*I)*a^2)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4})])/(8*\text{Sqrt}[2]*b^3) - ((3*I - 4*a - (8*I)*a^2)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4})])/(8*\text{Sqrt}[2]*b^3) + ((3*I - 4*a - (8*I)*a^2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4})])/(16*\text{Sqrt}[2]*b^3) - ((3*I - 4*a - (8*I)*a^2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4})])/(16*\text{Sqrt}[2]*b^3)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ

$[m, 0] \&\& (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0])) \ \&\& \ !\text{ILtQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \ :> \ \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m+1) - 1)}(c - a*(d/b) + d*(x^{p/b})^n, x], x, (a + b*x)^{(1/p)}], x]] \ /; \ \text{FreeQ}[\{a, b, c, d\}, x] \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{LtQ}[-1, m, 0] \ \&\& \ \text{LeQ}[-1, n, 0] \ \&\& \ \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 81

$\text{Int}[(a_.) + (b_.)(x_)^{(n_)}((c_.) + (d_.)(x_)^{(n_)}((e_.) + (f_.)(x_)^{(p_)}), x_Symbol] \ :> \ \text{Simp}[b*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 2, 0]$

Rule 92

$\text{Int}[(a_.) + (b_.)(x_)^{2*((c_.) + (d_.)(x_)^{(n_)}((e_.) + (f_.)(x_)^{(p_)}), x_Symbol] \ :> \ \text{Simp}[b*(a + b*x)*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+3))), x] + \text{Dist}[1/(d*f*(n+p+3)), \text{Int}[(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(n+p+3) - b*(b*c*e + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(n+p+4) - b*(d*e*(n+2) + c*f*(p+2))]*x, x], x] \ /; \ \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \text{NeQ}[n + p + 3, 0]$

Rule 210

$\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \ :> \ \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] \ /; \ \text{FreeQ}[\{a, b\}, x] \ \&\& \ \text{PosQ}[a/b] \ \&\& \ (\text{LtQ}[a, 0] \ || \ \text{LtQ}[b, 0])$

Rule 217

$\text{Int}[(a_) + (b_.)(x_)^4)^{-1}, x_Symbol] \ :> \ \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*r), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x] + \text{Dist}[1/(2*r), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x]] \ /; \ \text{FreeQ}[\{a, b\}, x] \ \&\& \ (\text{GtQ}[a/b, 0] \ || \ (\text{PosQ}[a/b] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \ \&\& \ \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 246

$\text{Int}[(a_) + (b_.)(x_)^{(n_)}^{(p_)}, x_Symbol] \ :> \ \text{Dist}[a^{(p+1/n)}, \text{Subst}[\text{Int}[1/(1 - b*x^n)^{(p+1/n+1)}, x], x, x/(a + b*x^n)^{(1/n)}], x] \ /; \ \text{FreeQ}[\{a,$

b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*((d_) + (e_)*(x_)^(m_)), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{-\frac{1}{2}i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2 \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} dx \\
&= \frac{x(1-ia-ibx)^{5/4}(1+ia+ibx)^{3/4}}{3b^2} + \frac{\int \frac{\sqrt[4]{1-ia-ibx} (-1-a^2+\frac{1}{2}(i-8a)bx)}{\sqrt[4]{1+ia+ibx}} dx}{3b^2} \\
&= \frac{(i-8a)(1-ia-ibx)^{5/4}(1+ia+ibx)^{3/4}}{12b^3} + \frac{x(1-ia-ibx)^{5/4}(1+ia+ibx)^{3/4}}{3b^2} - \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3} \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3} \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3} \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3} \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3} \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3} \\
&= -\frac{(4a-i(3-8a^2)) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{8b^3} + \frac{(i-8a)(1-ia-ibx)^{5/4}(1-ia-ibx)^{3/4}}{12b^3}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 99, normalized size = 0.20

$$\frac{(-i(i+a+bx))^{5/4} (5(1+ia+ibx)^{3/4}(i-8a+4bx) + 3 \cdot 2^{3/4}(-3i+4a+8ia^2) {}_2F_1(\frac{1}{4}, \frac{5}{4}, \frac{9}{4}; -\frac{1}{2}i(i+a+bx)))}{60b^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^((I/2)*ArcTan[a + b*x]), x]

[Out] $((-I)(I + a + b*x))^{5/4} * (5*(1 + I*a + I*b*x))^{3/4} * (I - 8*a + 4*b*x) + 3*2^{3/4} * (-3*I + 4*a + (8*I)*a^2) * \text{Hypergeometric2F1}[1/4, 5/4, 9/4, (-1/2*I) * (I + a + b*x)] / (60*b^3)$

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x)`

[Out] `int(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="maxima")`

[Out] `integrate(x^2/sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1)), x)`

Fricas [A]

time = 2.55, size = 561, normalized size = 1.14

$$\frac{1}{48} \left(3b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} \log\left(\frac{b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} + (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} - (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}})}\right) - 3b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} \log\left(\frac{-b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} - (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} + (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}})}\right) - 3b^3 \sqrt{\frac{-64Ia^4 - 64a^3 + 64Ia^2 + 24a - 9I}{b^6}} \log\left(\frac{b^3 \sqrt{\frac{-64Ia^4 - 64a^3 + 64Ia^2 + 24a - 9I}{b^6}} + (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(b^3 \sqrt{\frac{-64Ia^4 - 64a^3 + 64Ia^2 + 24a - 9I}{b^6}} - (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}})}\right) \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="fricas")`

[Out] $\frac{1}{48} \left(3b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} \log\left(\frac{b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} + (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} - (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}})}\right) - 3b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} \log\left(\frac{-b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} - (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(b^3 \sqrt{\frac{64Ia^4 + 64a^3 - 64Ia^2 - 24a + 9I}{b^6}} + (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}})}\right) - 3b^3 \sqrt{\frac{-64Ia^4 - 64a^3 + 64Ia^2 + 24a - 9I}{b^6}} \log\left(\frac{b^3 \sqrt{\frac{-64Ia^4 - 64a^3 + 64Ia^2 + 24a - 9I}{b^6}} + (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(b^3 \sqrt{\frac{-64Ia^4 - 64a^3 + 64Ia^2 + 24a - 9I}{b^6}} - (8a^2 - 4Ia - 3) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}})}\right) \right)$

- 3)) + 3*b^3*sqrt((-64*I*a^4 - 64*a^3 + 64*I*a^2 + 24*a - 9*I)/b^6)*log(-(b^3*sqrt((-64*I*a^4 - 64*a^3 + 64*I*a^2 + 24*a - 9*I)/b^6) - (8*a^2 - 4*I*a - 3)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(8*a^2 - 4*I*a - 3)) + 2*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*(-8*I*b^2*x^2 - 2*(-4*I*a - 5)*b*x - 8*I*a^2 - 26*a + 11*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b^3

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\sqrt{\frac{i(a+bx-i)}{\sqrt{a^2+2abx+b^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2),x)

[Out] Integral(x**2/sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx);OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2}{\sqrt{\frac{1+a\,i+b\,x\,i}{\sqrt{(a+b\,x)^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/((a*i + b*x*i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2),x)

[Out] int(x^2/((a*i + b*x*i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)

3.227 $\int e^{-\frac{1}{2}i\text{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=410

$$\frac{(1+4ia)\sqrt{1-ia-ibx}(1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4}(1+ia+ibx)^{3/4}}{2b^2} + \frac{(1+4ia)\text{ArcTan}\left(1-\frac{\sqrt{2}\sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}}\right)}{4\sqrt{2}b^2}$$

[Out] $1/4*(1+4*I*a)*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}/b^2+1/2*(1-I*a-I*b*x)^{(5/4)}*(1+I*a+I*b*x)^{(3/4)}/b^2+1/8*(1+4*I*a)*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^2+2^{(1/2)}-1/8*(1+4*I*a)*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b^2+2^{(1/2)}+1/16*(1+4*I*a)*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b^2+2^{(1/2)}-1/16*(1+4*I*a)*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b^2+2^{(1/2)}$

Rubi [A]

time = 0.22, antiderivative size = 410, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {5203, 81, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{(1+4ia)\text{ArcTan}\left(1-\frac{\sqrt{2}\sqrt{1-ia-ibx+1}}{\sqrt{1+ia+ibx+1}}\right)}{4\sqrt{2}b^2} - \frac{(1+4ia)\text{ArcTan}\left(1+\frac{\sqrt{2}\sqrt{1-ia-ibx+1}}{\sqrt{1+ia+ibx+1}}\right)}{4\sqrt{2}b^2} + \frac{(ia+ibx+1)^{5/4}(1-ia-ibx+1)^{3/4}}{2b^2} + \frac{(1+4ia)(ia+ibx+1)^{3/4}\sqrt{1-ia-ibx+1}}{4b^2} + \frac{(1+4ia)\log\left(\frac{\sqrt{1-ia-ibx+1}}{\sqrt{1+ia+ibx+1}}-\frac{\sqrt{2}\sqrt{1-ia-ibx+1}}{\sqrt{1+ia+ibx+1}}+1\right)}{8\sqrt{2}b^2} - \frac{(1+4ia)\log\left(\frac{\sqrt{1-ia-ibx+1}}{\sqrt{1+ia+ibx+1}}+\frac{\sqrt{2}\sqrt{1-ia-ibx+1}}{\sqrt{1+ia+ibx+1}}+1\right)}{8\sqrt{2}b^2}$$

Antiderivative was successfully verified.

[In] Int[x/E^((I/2)*ArcTan[a + b*x]),x]

[Out] $((1+(4*I)*a)*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)})/(4*b^2) + ((1-I*a-I*b*x)^{(5/4)}*(1+I*a+I*b*x)^{(3/4)})/(2*b^2) + ((1+(4*I)*a)*\text{ArcTan}[1-(\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(4*\text{Sqrt}[2]*b^2) - ((1+(4*I)*a)*\text{ArcTan}[1+(\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(4*\text{Sqrt}[2]*b^2) + ((1+(4*I)*a)*\text{Log}[1+\text{Sqrt}[1-I*a-I*b*x]/\text{Sqrt}[1+I*a+I*b*x] - (\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(8*\text{Sqrt}[2]*b^2) - ((1+(4*I)*a)*\text{Log}[1+\text{Sqrt}[1-I*a-I*b*x]/\text{Sqrt}[1+I*a+I*b*x] + (\text{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)})]/(8*\text{Sqrt}[2]*b^2)$

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^(n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))), Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^(n), x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p
_.), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
```


e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_.)*((d_) + (e_)*(x_)^m_), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{-\frac{1}{2}i \tan^{-1}(a+bx)} x dx &= \int \frac{x^4 \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} dx \\
&= \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} + \frac{(i-4a) \int \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} dx}{4b} \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} + \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} - \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} - \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} - \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} - \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} - \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} + \dots \\
&= \frac{(1+4ia) \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{4b^2} + \frac{(1-ia-ibx)^{5/4} (1+ia+ibx)^{3/4}}{2b^2} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 84, normalized size = 0.20

$$\frac{i(-i(i+a+bx))^{5/4} (5i(1+ia+ibx)^{3/4} + 2^{3/4}(-i+4a) {}_2F_1(\frac{1}{4}, \frac{5}{4}; \frac{9}{4}; -\frac{1}{2}i(i+a+bx)))}{10b^2}$$

Antiderivative was successfully verified.

[In] Integrate[x/E^((I/2)*ArcTan[a + b*x]),x]

[Out] ((-1/10*I)*((-I)*(I + a + b*x))^(5/4)*((5*I)*(1 + I*a + I*b*x)^(3/4) + 2^(3/4)*(-I + 4*a)*Hypergeometric2F1[1/4, 5/4, 9/4, (-1/2*I)*(I + a + b*x)]))/b^2

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x}{\sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2), x)

[Out] int(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2), x, algorithm="maxima")

[Out] integrate(x/sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1)), x)

Fricas [A]

time = 2.77, size = 421, normalized size = 1.03

$$\frac{r^2 \sqrt{\frac{8a^2 + 8a - 1}{b}} \log\left(\frac{r^2 \sqrt{\frac{16a^2 + 8a - 1}{b}} \sqrt{\frac{\sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 1}}}{\sqrt{\frac{16a^2 + 8a - 1}{b}}}\right) - r^2 \sqrt{\frac{8a^2 + 8a - 1}{b}} \log\left(\frac{r^2 \sqrt{\frac{16a^2 + 8a - 1}{b}} \sqrt{\frac{\sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 1}}}{\sqrt{\frac{16a^2 + 8a - 1}{b}}}\right) - r^2 \sqrt{\frac{16a^2 - 8a + 1}{b}} \log\left(\frac{r^2 \sqrt{\frac{16a^2 - 8a + 1}{b}} \sqrt{\frac{\sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 1}}}{\sqrt{\frac{16a^2 - 8a + 1}{b}}}\right) + r^2 \sqrt{\frac{16a^2 - 8a + 1}{b}} \log\left(\frac{r^2 \sqrt{\frac{16a^2 - 8a + 1}{b}} \sqrt{\frac{\sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 1}}}{\sqrt{\frac{16a^2 - 8a + 1}{b}}}\right) - 2 \sqrt{b^2 x^2 + 2abx + a^2 + 1} (-3bx + 3a + 3) \sqrt{\frac{\sqrt{b^2 x^2 + 2abx + a^2 + 1}}{4a + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2), x, algorithm="fricas")

[Out] $-1/8*(b^2*\sqrt{(16*I*a^2 + 8*a - I)/b^4})*\log((b^2*\sqrt{(16*I*a^2 + 8*a - I)/b^4} + (4*a - I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a - I)) - b^2*\sqrt{(16*I*a^2 + 8*a - I)/b^4}*\log(-(b^2*\sqrt{(16*I*a^2 + 8*a - I)/b^4} - (4*a - I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a - I)) - b^2*\sqrt{(-16*I*a^2 - 8*a + I)/b^4}*\log((b^2*\sqrt{(-16*I*a^2 - 8*a + I)/b^4} + (4*a - I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a - I)) + b^2*\sqrt{(-16*I*a^2 - 8*a + I)/b^4}*\log(-(b^2*\sqrt{(-16*I*a^2 - 8*a + I)/b^4} - (4*a - I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)})/(4*a - I)) - 2*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*(-2*I*b*x + 2*I*a + 3)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)}/b^2$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x}{\sqrt{\frac{i(a+bx-i)}{\sqrt{a^2+2abx+b^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2),x)``[Out] Integral(x/sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1)), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="giac")``[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x}{\sqrt{\frac{1+a\,i+b\,x\,i}{\sqrt{(a+bx)^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x/((a*i + b*x*i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2),x)``[Out] int(x/((a*i + b*x*i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)`

3.228 $\int e^{-\frac{1}{2}i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=338

$$\frac{i\sqrt[4]{1-ia-ibx}(1+ia+ibx)^{3/4}}{b} - \frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b}$$

[Out] $-I*(1-I*a-I*b*x)^{(1/4)}*(1+I*a+I*b*x)^{(3/4)}/b-1/2*I*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}+1/2*I*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}-1/4*I*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}+1/4*I*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 338, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$, Rules used = {5201, 52, 65, 246, 217, 1179, 642, 1176, 631, 210}

$$\frac{i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} + \frac{i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} - \frac{i\sqrt{-ia-ibx+1}(ia+ibx+1)^{3/4}}{b} - \frac{i\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b} + \frac{i\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((-1/2*I)*\text{ArcTan}[a + b*x])}, x]$

[Out] $((-I)*(1 - I*a - I*b*x)^{(1/4)}*(1 + I*a + I*b*x)^{(3/4)})/b - (I*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(Sqrt[2]*b) + (I*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(Sqrt[2]*b) - ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(Sqrt[2]*b) + ((I/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(Sqrt[2]*b)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ[m, 0] && (!IntegerQ[n] || (GtQ[m, 0] && LtQ[m - n, 0]))) && !ILtQ[m + n + 2, 0] && IntLinearQ[a, b, c, d, m, n, x]

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) +$

```
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 217

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]
], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4),
x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b
}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &&
AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 246

```
Int[((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + 1/n), Subst[Int
[1/(1 - b*x^n)^(p + 1/n + 1), x], x, x/(a + b*x^n)^(1/n)], x] /; FreeQ[{a,
b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2^(-1)] && IntegerQ[p + 1/
n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5201

```
Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{1}{2}i \tan^{-1}(a+bx)} dx &= \int \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} dx \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} + \frac{1}{2} \int \frac{1}{(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}} dx \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} + \frac{(2i) \text{Subst}\left(\int \frac{1}{\sqrt[4]{2-x^4}} dx, x, \sqrt[4]{1-ia-ibx}\right)}{b} \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} + \frac{(2i) \text{Subst}\left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{b} \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} + \frac{i \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{b} + \frac{i \text{Subst}\left(\int \frac{1}{1-x^2} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{b} \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} + \frac{i \text{Subst}\left(\int \frac{1}{1-\sqrt{2}x+x^2} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{2b} \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} - \frac{i \log\left(1 + \frac{\sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}} - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{2\sqrt{2}b} \\
&= -\frac{i\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}}{b} - \frac{i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b} + \frac{i \tan^{-1}\left(\frac{\sqrt{1-ia-ibx}}{\sqrt{1+ia+ibx}}\right)}{b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 45, normalized size = 0.13

$$\frac{8ie^{\frac{3}{2}i \text{ArcTan}(a+bx)} {}_2F_1\left(\frac{3}{4}, 2; \frac{7}{4}; -e^{2i \text{ArcTan}(a+bx)}\right)}{3b}$$

Antiderivative was successfully verified.

[In] Integrate[E^((-1/2*I)*ArcTan[a + b*x]),x]

[Out] (((-8*I)/3)*E^(((3*I)/2)*ArcTan[a + b*x])*Hypergeometric2F1[3/4, 2, 7/4, -E^((2*I)*ArcTan[a + b*x])]/b

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x)

[Out] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="maxima")

[Out] integrate(1/sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1)), x)

Fricas [A]

time = 3.80, size = 266, normalized size = 0.79

$$\frac{b\sqrt{\frac{i}{b^2}} \log\left(b\sqrt{\frac{i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - b\sqrt{\frac{1}{b^2}} \log\left(-b\sqrt{\frac{1}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - b\sqrt{\frac{i}{b^2}} \log\left(b\sqrt{\frac{i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) + b\sqrt{\frac{i}{b^2}} \log\left(-b\sqrt{\frac{i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}\right) - 2i\sqrt{b^2x^2 + 2abx + a^2 + 1} \sqrt{\frac{i\sqrt{b^2x^2 + 2abx + a^2 + 1}}{bx + a + i}}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="fricas")

[Out] 1/2*(b*sqrt(I/b^2)*log(b*sqrt(I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(I/b^2)*log(-b*sqrt(I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(-I/b^2)*log(b*sqrt(-I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + b*sqrt(-I/b^2)*log(-b*sqrt(-I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 2*I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{i(a+bx)+1}{\sqrt{(a+bx)^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2),x)``[Out] Integral(1/sqrt((I*(a + b*x) + 1)/sqrt((a + b*x)**2 + 1)), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2),x, algorithm="giac")``[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\sqrt{\frac{1+a\text{li}+bx\text{li}}{\sqrt{(a+bx)^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2),x)``[Out] int(1/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2), x)`

$$3.229 \quad \int \frac{e^{-\frac{1}{2}i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=395

$$\frac{2\sqrt[4]{i+a} \operatorname{ArcTan}\left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}}\right)}{\sqrt[4]{i-a}} - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}}\right)$$

[Out] $-2*(I+a)^{(1/4)}*\arctan((I-a)^{(1/4)}*(1-I*(b*x+a))^{(1/4)}/(I+a)^{(1/4)/(1+I*(b*x+a))^{(1/4))}/(I-a)^{(1/4)}-2*(I+a)^{(1/4)}*\operatorname{arctanh}((I-a)^{(1/4)}*(1-I*(b*x+a))^{(1/4)}/(I+a)^{(1/4)/(1+I*(b*x+a))^{(1/4))}/(I-a)^{(1/4)}-1/2*\ln(1-(1-I*(b*x+a))^{(1/4)})*2^{(1/2)}/(1+I*(b*x+a))^{(1/4)}+1/(1+I*(b*x+a))^{(1/2)}*(1-I*(b*x+a))^{(1/2)})*2^{(1/2)}+1/2*\ln(1+(1-I*(b*x+a))^{(1/4)})*2^{(1/2)}/(1+I*(b*x+a))^{(1/4)}+1/(1+I*(b*x+a))^{(1/2)}*(1-I*(b*x+a))^{(1/2)})*2^{(1/2)}-\arctan(1-(1-I*(b*x+a))^{(1/4)})*2^{(1/2)}/(1+I*(b*x+a))^{(1/4)})*2^{(1/2)}+\operatorname{arctan}(1+(1-I*(b*x+a))^{(1/4)})*2^{(1/2)}/(1+I*(b*x+a))^{(1/4)})*2^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 395, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.611$, Rules used = {5202, 492, 217, 1179, 642, 1176, 631, 210, 218, 214, 211}

$$\frac{2\sqrt[4]{a+i} \operatorname{ArcTan}\left(\frac{\sqrt[4]{-a+i} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{a+i} \sqrt[4]{1+i(a+bx)}}\right)}{\sqrt[4]{-a+i}} - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}}\right) - \frac{\log\left(\frac{\sqrt{1-i(a+bx)}}{\sqrt{1+i(a+bx)}} - \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt{2} \sqrt[4]{1+i(a+bx)}} + 1\right)}{\sqrt{2}} + \frac{\log\left(\frac{\sqrt{1-i(a+bx)}}{\sqrt{1+i(a+bx)}} + \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt{2} \sqrt[4]{1+i(a+bx)}} + 1\right)}{\sqrt{2}} - \frac{2\sqrt[4]{a+i} \operatorname{tanh}^{-1}\left(\frac{\sqrt[4]{-a+i} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{a+i} \sqrt[4]{1+i(a+bx)}}\right)}{\sqrt[4]{-a+i}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^((I/2)*ArcTan[a + b*x]))*x], x]`

[Out] $(-2*(I+a)^{(1/4)}*\operatorname{ArcTan}(((I-a)^{(1/4)}*(1-I*(a+b*x))^{(1/4)})/((I+a)^{(1/4)}*(1+I*(a+b*x))^{(1/4)})))/(I-a)^{(1/4)} - \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1-I*(a+b*x))^{(1/4)})/(1+I*(a+b*x))^{(1/4)}] + \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1-I*(a+b*x))^{(1/4)})/(1+I*(a+b*x))^{(1/4)}] - (2*(I+a)^{(1/4)}*\operatorname{ArcTan}h(((I-a)^{(1/4)}*(1-I*(a+b*x))^{(1/4)})/((I+a)^{(1/4)}*(1+I*(a+b*x))^{(1/4)})))/(I-a)^{(1/4)} - \operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*(a+b*x)]]/\operatorname{Sqrt}[1 + I*(a+b*x)] - (\operatorname{Sqrt}[2]*(1-I*(a+b*x))^{(1/4)})/(1+I*(a+b*x))^{(1/4)}/\operatorname{Sqrt}[2] + \operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*(a+b*x)]]/\operatorname{Sqrt}[1 + I*(a+b*x)] + (\operatorname{Sqrt}[2]*(1-I*(a+b*x))^{(1/4)})/(1+I*(a+b*x))^{(1/4)}/\operatorname{Sqrt}[2]$

Rule 210

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (LtQ[a, 0] || LtQ[b, 0])`

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 217

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[a/b, 2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*r), Int[(r - s*x^2)/(a + b*x^4), x], x] + Dist[1/(2*r), Int[(r + s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a, b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] && AtomQ[SplitProduct[SumBaseQ, b]]))

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 492

Int[((e_)*(x_)^(m_))/(((a_) + (b_)*(x_)^(n_))*((c_) + (d_)*(x_)^(n_))), x_Symbol] := Dist[(-a)*(e^n/(b*c - a*d)), Int[(e*x)^(m - n)/(a + b*x^n), x], x] + Dist[c*(e^n/(b*c - a*d)), Int[(e*x)^(m - n)/(c + d*x^n), x], x] /; FreeQ[{a, b, c, d, e, m}, x] && NeQ[b*c - a*d, 0] && IGtQ[n, 0] && LeQ[n, m, 2*n - 1]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*Simplify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; FreeQ[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := Simp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d, e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[
e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5202

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_)*(x_)^(m_), x_Symbol] := Dist
[4/(I^m*n*b^(m + 1)*c^(m + 1)), Subst[Int[x^(2/(I*n))*((1 - I*a*c - (1 + I*
a*c)*x^(2/(I*n)))^m/(1 + x^(2/(I*n)))^(m + 2)), x], x, (1 - I*c*(a + b*x))^
(I*(n/2))/(1 + I*c*(a + b*x)^(I*(n/2))], x] /; FreeQ[{a, b, c}, x] && ILtQ
[m, 0] && LtQ[-1, I*n, 1]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{1}{2}i \tan^{-1}(a+bx)}}{x} dx &= - \left(8 \text{Subst} \left(\int \frac{x^4}{(1+x^4)(1-ia-(1+ia)x^4)} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right) \right) \\
&= 4 \text{Subst} \left(\int \frac{1}{1+x^4} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right) - (4(1-ia)) \text{Subst} \left(\int \frac{1}{1-ia+(-1-ia)x^4} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right) \\
&= 2 \text{Subst} \left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right) + 2 \text{Subst} \left(\int \frac{1+x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right) \\
&= - \frac{2\sqrt[4]{i+a} \tan^{-1} \left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}} \right)}{\sqrt[4]{i-a}} - \frac{2\sqrt[4]{i+a} \tanh^{-1} \left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}} \right)}{\sqrt[4]{i-a}} \\
&= - \frac{2\sqrt[4]{i+a} \tan^{-1} \left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}} \right)}{\sqrt[4]{i-a}} - \frac{2\sqrt[4]{i+a} \tanh^{-1} \left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}} \right)}{\sqrt[4]{i-a}} \\
&= - \frac{2\sqrt[4]{i+a} \tan^{-1} \left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}} \right)}{\sqrt[4]{i-a}} - \sqrt{2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 126, normalized size = 0.32

$$\frac{2\sqrt[4]{-i(i+a+bx)} \left(2^{3/4}\sqrt[4]{1+ia+ibx} {}_2F_1\left(\frac{1}{4}, \frac{1}{4}; \frac{5}{4}; -\frac{1}{2}i(i+a+bx)\right) - 2 {}_2F_1\left(\frac{1}{4}, 1; \frac{5}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right) \right)}{\sqrt[4]{1+ia+ibx}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a + b*x]))*x, x]

[Out] (2*((-I)*(I + a + b*x))^(1/4)*(2^(3/4)*(1 + I*a + I*b*x))^(1/4)*Hypergeometric2F1[1/4, 1/4, 5/4, (-1/2*I)*(I + a + b*x)] - 2*Hypergeometric2F1[1/4, 1, 5/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)])/(1 + I*a + I*b*x)^(1/4)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x)

[Out] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x, algorithm="maxima")

[Out] integrate(1/(x*sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))), x)

Fricas [A]

time = 5.51, size = 470, normalized size = 1.19

$$\frac{1}{2} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}} \sqrt{\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x, algorithm="fricas")

```
[Out] -1/2*sqrt(4*I)*log(1/2*I*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 1/2*sqrt(4*I)*log(-1/2*I*sqrt(4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + 1/2*sqrt(-4*I)*log(1/2*I*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 1/2*sqrt(-4*I)*log(-1/2*I*sqrt(-4*I) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + ((-a + I)/(a - I))^(1/4)*log(((a - I)*(-a + I)/(a - I))^(3/4) + (a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(a + I) - ((-a + I)/(a - I))^(1/4)*log(-((a - I)*(-a + I)/(a - I))^(3/4) - (a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(a + I) - I*(-(a + I)/(a - I))^(1/4)*log(((I*a + 1)*(-a + I)/(a - I))^(3/4) + (a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(a + I) + I*(-(a + I)/(a - I))^(1/4)*log((-I*a - 1)*(-a + I)/(a - I))^(3/4) + (a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(a + I)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x \sqrt{\frac{i(a+bx-i)}{\sqrt{a^2+2abx+b^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2)/x,x)
```

```
[Out] Integral(1/(x*sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1))), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x,x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x \sqrt{\frac{1 + a li + b x li}{\sqrt{(a + b x)^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2)))^(1/2),x)
```

```
[Out] int(1/(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2)))^(1/2), x)
```

$$3.230 \quad \int \frac{e^{-\frac{1}{2}i \operatorname{ArcTan}(a+bx)}}{x^2} dx$$

Optimal. Leaf size=210

$$\frac{(i-a-bx)\sqrt[4]{1-i(a+bx)}}{(i-a)x\sqrt[4]{1+i(a+bx)}} - \frac{ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{i-a}\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a}\sqrt[4]{1+i(a+bx)}}\right)}{(i-a)^{5/4}(i+a)^{3/4}} - \frac{ib \tanh^{-1}\left(\frac{\sqrt[4]{i-a}\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a}\sqrt[4]{1+i(a+bx)}}\right)}{(i-a)^{5/4}(i+a)^{3/4}}$$

[Out] $-(I-a-b*x)*(1-I*(b*x+a))^{(1/4)}/(I-a)/x/(1+I*(b*x+a))^{(1/4)}-I*b*\arctan((I-a)^{(1/4)}*(1-I*(b*x+a))^{(1/4)}/(I+a)^{(1/4)}/(1+I*(b*x+a))^{(1/4)})/(I-a)^{(5/4)}/(I+a)^{(3/4)}-I*b*\operatorname{arctanh}((I-a)^{(1/4)}*(1-I*(b*x+a))^{(1/4)}/(I+a)^{(1/4)}/(1+I*(b*x+a))^{(1/4)})/(I-a)^{(5/4)}/(I+a)^{(3/4)}$

Rubi [A]

time = 0.10, antiderivative size = 210, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.278$, Rules used = {5202, 294, 218, 214, 211}

$$-\frac{ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{-a+i}\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{a+i}\sqrt[4]{1+i(a+bx)}}\right)}{(-a+i)^{5/4}(a+i)^{3/4}} - \frac{\sqrt[4]{1-i(a+bx)}(-a-bx+i)}{(-a+i)x\sqrt[4]{1+i(a+bx)}} - \frac{ib \tanh^{-1}\left(\frac{\sqrt[4]{-a+i}\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{a+i}\sqrt[4]{1+i(a+bx)}}\right)}{(-a+i)^{5/4}(a+i)^{3/4}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^((I/2)*ArcTan[a + b*x])*x^2), x]`

[Out] $-\left(\frac{(I-a-b*x)*(1-I*(a+b*x))^{(1/4)}}{(I-a)*x*(1+I*(a+b*x))^{(1/4)}}\right) - \left(\frac{I*b*\operatorname{ArcTan}\left[\frac{(I-a)^{(1/4)}*(1-I*(a+b*x))^{(1/4)}}{(I+a)^{(1/4)}*(1+I*(a+b*x))^{(1/4)}}\right]}{(I-a)^{(5/4)}*(I+a)^{(3/4)}} - \frac{I*b*\operatorname{ArcTanh}\left[\frac{(I-a)^{(1/4)}*(1-I*(a+b*x))^{(1/4)}}{(I+a)^{(1/4)}*(1+I*(a+b*x))^{(1/4)}}\right]}{(I-a)^{(5/4)}*(I+a)^{(3/4)}}\right)$

Rule 211

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]`

Rule 214

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]`

Rule 218

`Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b]`

, 0]

Rule 294

```
Int[((c_.)*(x_))^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Simp[c^(
n - 1)*(c*x)^(m - n + 1)*((a + b*x^n)^(p + 1)/(b*n*(p + 1))), x] - Dist[c^n
*((m - n + 1)/(b*n*(p + 1))), Int[(c*x)^(m - n)*(a + b*x^n)^(p + 1), x], x]
/; FreeQ[{a, b, c}, x] && IGtQ[n, 0] && LtQ[p, -1] && GtQ[m + 1, n] && !
LtQ[(m + n*(p + 1) + 1)/n, 0] && IntBinomialQ[a, b, c, n, m, p, x]
```

Rule 5202

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_)*(x_)^(m_), x_Symbol] := Dist
[4/(I^m*n*b^(m + 1)*c^(m + 1)), Subst[Int[x^(2/(I*n))*((1 - I*a*c - (1 + I*
a*c)*x^(2/(I*n)))^m/(1 + x^(2/(I*n)))^(m + 2)), x], x, (1 - I*c*(a + b*x))^
(I*(n/2))/(1 + I*c*(a + b*x)^(I*(n/2))], x] /; FreeQ[{a, b, c}, x] && ILtQ
[m, 0] && LtQ[-1, I*n, 1]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{-\frac{1}{2}i \tan^{-1}(a+bx)}}{x^2} dx &= - \left((8ib) \text{Subst} \left(\int \frac{x^4}{(1-ia - (1+ia)x^4)^2} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right) \right) \\ &= - \frac{(i-a-bx) \sqrt[4]{1-i(a+bx)}}{(i-a)x \sqrt[4]{1+i(a+bx)}} - \frac{(2b) \text{Subst} \left(\int \frac{1}{1-ia+(-1-ia)x^4} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right)}{i-a} \\ &= - \frac{(i-a-bx) \sqrt[4]{1-i(a+bx)}}{(i-a)x \sqrt[4]{1+i(a+bx)}} - \frac{b \text{Subst} \left(\int \frac{1}{\sqrt{i+a} - \sqrt{i-a} x^2} dx, x, \frac{\sqrt[4]{1-i(a+bx)}}{\sqrt[4]{1+i(a+bx)}} \right)}{(1+ia)\sqrt{i+a}} \\ &= - \frac{(i-a-bx) \sqrt[4]{1-i(a+bx)}}{(i-a)x \sqrt[4]{1+i(a+bx)}} - \frac{ib \tan^{-1} \left(\frac{\sqrt[4]{i-a} \sqrt[4]{1-i(a+bx)}}{\sqrt[4]{i+a} \sqrt[4]{1+i(a+bx)}} \right)}{(i-a)^{5/4}(i+a)^{3/4}} - \frac{ib \tanh^{-1}}{\dots} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 107, normalized size = 0.51

$$- \frac{\sqrt[4]{-i(i+a+bx)} \left(1 + a^2 + ibx + abx - 2ibx {}_2F_1 \left(\frac{1}{4}, 1; \frac{5}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx} \right) \right)}{(1+a^2)x \sqrt[4]{1+ia+ibx}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((I/2)*ArcTan[a + b*x])*x^2),x]

[Out] -((((-I)*(I + a + b*x))^(1/4)*(1 + a^2 + I*b*x + a*b*x - (2*I)*b*x*Hypergeometric2F1[1/4, 1, 5/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)]))/((1 + a^2)*x*(1 + I*a + I*b*x)^(1/4)))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\sqrt{\frac{1 + i(bx + a)}{\sqrt{1 + (bx + a)^2}} x^2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x)

[Out] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x, algorithm="maxima")

[Out] integrate(1/(x^2*sqrt((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 707 vs. 2(141) = 282.

time = 8.00, size = 707, normalized size = 3.37

$$\frac{(-b^4/(a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{1/4} \cdot (-Ia - 1) \cdot x \cdot \log((b^3 \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}} / (b x + a + I)) + (a^6 - 2Ia^5 + a^4 - 4Ia^3 - a^2 - 2Ia - 1) \cdot (-b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{3/4}) / b^3 + (-b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{1/4} \cdot (Ia + 1) \cdot x \cdot \log((b^3 \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}} / (b x + a + I)) - (a^6 - 2Ia^5 + a^4 - 4Ia^3 - a^2 - 2Ia - 1) \cdot (-b^4 / (a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{3/4}) / b^3}{b^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x, algorithm="fricas")

[Out] 1/2*((-b^4/(a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^(1/4)*(-Ia - 1)*x*log((b^3*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/(b*x + a + I)) + (a^6 - 2*I*a^5 + a^4 - 4*I*a^3 - a^2 - 2*I*a - 1)*(-b^4/(a^8 - 2*I*a^7 + 2*a^6 - 6*I*a^5 - 6*I*a^3 - 2*a^2 - 2*I*a - 1))^(3/4))/b^3) + (-b^4/(a^8 - 2*I*a^7 + 2*a^6 - 6*I*a^5 - 6*I*a^3 - 2*a^2 - 2*I*a - 1))^(1/4)*(I*a + 1)*x*log((b^3*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1))/(b*x + a + I)) - (a^6 - 2*I*a^5 + a^4 - 4*I*a^3 - a^2 - 2*I*a - 1)*(-b^4/(a^8 - 2*I*a^7 + 2*a^6 - 6*I*a^5 - 6*I*a^3 - 2*a^2 - 2*I*a - 1))^(3/4))/b^3)

$$2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{3/4})/b^3) + (-b^4/(a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{1/4}*(a - I)*x*\log((b^3*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}}/(b*x + a + I)) - (I*a^6 + 2*a^5 + I*a^4 + 4*a^3 - I*a^2 + 2*a - I)*(-b^4/(a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{3/4})/b^3) - (-b^4/(a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{1/4}*(a - I)*x*\log((b^3*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}}/(b*x + a + I)) - (-I*a^6 - 2*a^5 - I*a^4 - 4*a^3 + I*a^2 - 2*a + I)*(-b^4/(a^8 - 2Ia^7 + 2a^6 - 6Ia^5 - 6Ia^3 - 2a^2 - 2Ia - 1))^{3/4})/b^3) + 2I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}}/(b*x + a + I))/((a - I)*x)$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{x^2 \sqrt{\frac{i(a+bx-i)}{\sqrt{a^2+2abx+b^2x^2+1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(1/2)/x**2,x)

[Out] Integral(1/(x**2*sqrt(I*(a + b*x - I)/sqrt(a**2 + 2*a*b*x + b**2*x**2 + 1))), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(1/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x^2 \sqrt{\frac{1 + a \operatorname{li} + b x \operatorname{li}}{\sqrt{(a + b x)^2 + 1}}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2)),x)
```

```
[Out] int(1/(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(1/2)), x)
```

3.231 $\int e^{-\frac{3}{2}i\text{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=494

$$\frac{(17i - 36a - 24ia^2)(1 - ia - ibx)^{3/4}\sqrt[4]{1 + ia + ibx}}{24b^3} + \frac{(3i - 8a)(1 - ia - ibx)^{7/4}\sqrt[4]{1 + ia + ibx}}{12b^3} + \frac{x(1 - ia - ibx)^{3/4}\sqrt[4]{1 + ia + ibx}}{12b^3}$$

[Out] 1/24*(17*I-36*a-24*I*a^2)*(1-I*a-I*b*x)^(3/4)*(1+I*a+I*b*x)^(1/4)/b^3+1/12*(3*I-8*a)*(1-I*a-I*b*x)^(7/4)*(1+I*a+I*b*x)^(1/4)/b^3+1/3*x*(1-I*a-I*b*x)^(7/4)*(1+I*a+I*b*x)^(1/4)/b^2+1/16*(17*I-36*a-24*I*a^2)*arctan(1-(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4))/b^3*2^(1/2)-1/16*(17*I-36*a-24*I*a^2)*arctan(1+(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4))/b^3*2^(1/2)-1/32*(17*I-36*a-24*I*a^2)*ln(1-(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4)+(1-I*a-I*b*x)^(1/2)/(1+I*a+I*b*x)^(1/2))/b^3*2^(1/2)+1/32*(17*I-36*a-24*I*a^2)*ln(1+(1-I*a-I*b*x)^(1/4)*2^(1/2)/(1+I*a+I*b*x)^(1/4)+(1-I*a-I*b*x)^(1/2)/(1+I*a+I*b*x)^(1/2))/b^3*2^(1/2)

Rubi [A]

time = 0.29, antiderivative size = 494, normalized size of antiderivative = 1.00, number of steps used = 15, number of rules used = 12, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.667$, Rules used = {5203, 92, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{(-24a^2 - 36a + 17i)\text{ArcTan}\left(\frac{1 - \sqrt{2}\sqrt{1 + ia + ibx}}{\sqrt{1 + ia + ibx}}\right)}{8\sqrt{2}b^3} - \frac{(-24a^2 - 36a + 17i)\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt{1 + ia + ibx}}{\sqrt{1 + ia + ibx}}\right)}{8\sqrt{2}b^3} + \frac{(-24a^2 - 36a + 17i)\sqrt{1 + ia + ibx}\left(\cos - \text{ArcTan}\right)}{24b^3} - \frac{(-24a^2 - 36a + 17i)\log\left(\frac{\sqrt{1 + ia + ibx}}{\sqrt{1 + ia + ibx}}\right)}{16\sqrt{2}b^3} + \frac{(-24a^2 - 36a + 17i)\log\left(\frac{\sqrt{1 + ia + ibx}}{\sqrt{1 + ia + ibx}} + 1\right)}{16\sqrt{2}b^3} + \frac{(-8a + 30i)\sqrt{1 + ia + ibx}\left(\cos - \text{ArcTan}\right)}{12b^3} + \frac{e^{2i\text{ArcTan}\left(\frac{1 - \sqrt{2}\sqrt{1 + ia + ibx}}{\sqrt{1 + ia + ibx}}\right)}}{3b^3}$$

Antiderivative was successfully verified.

[In] Int[x^2/E^(((3*I)/2)*ArcTan[a + b*x]),x]

[Out] ((17*I - 36*a - (24*I)*a^2)*(1 - I*a - I*b*x)^(3/4)*(1 + I*a + I*b*x)^(1/4))/(24*b^3) + ((3*I - 8*a)*(1 - I*a - I*b*x)^(7/4)*(1 + I*a + I*b*x)^(1/4))/(12*b^3) + (x*(1 - I*a - I*b*x)^(7/4)*(1 + I*a + I*b*x)^(1/4))/(3*b^2) + ((17*I - 36*a - (24*I)*a^2)*ArcTan[1 - (Sqrt[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/(8*Sqrt[2]*b^3) - ((17*I - 36*a - (24*I)*a^2)*ArcTan[1 + (Sqrt[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/(8*Sqrt[2]*b^3) - ((17*I - 36*a - (24*I)*a^2)*Log[1 + Sqrt[1 - I*a - I*b*x]/Sqrt[1 + I*a + I*b*x] - (Sqrt[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/(16*Sqrt[2]*b^3) + ((17*I - 36*a - (24*I)*a^2)*Log[1 + Sqrt[1 - I*a - I*b*x]/Sqrt[1 + I*a + I*b*x] + (Sqrt[2]*(1 - I*a - I*b*x)^(1/4))/(1 + I*a + I*b*x)^(1/4)])/(16*Sqrt[2]*b^3)

Rule 52

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + n + 1))), x] + Dist[n*((b*c - a*d)/(b*(m + n + 1))], Int[(a + b*x)^m*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && NeQ[m + n + 1, 0] && !(IGtQ

$[m, 0] \&\& (!\text{IntegerQ}[n] \mid\mid (\text{GtQ}[m, 0] \&\& \text{LtQ}[m - n, 0])) \&\& !\text{ILtQ}[m + n + 2, 0] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_.) + (b_.)(x_)^{(m_)}((c_.) + (d_.)(x_)^{(n_)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p(m+1)-1)}(c - a(d/b) + d(x^{p/b}))^n, x], x, (a + b*x)^{(1/p)}], x]] /; \text{FreeQ}[\{a, b, c, d\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[-1, m, 0] \&\& \text{LeQ}[-1, n, 0] \&\& \text{LeQ}[\text{Denominator}[n], \text{Denominator}[m]] \&\& \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 81

$\text{Int}[(a_.) + (b_.)(x_)^{(c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[b*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+2))), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \&\& \text{NeQ}[n + p + 2, 0]$

Rule 92

$\text{Int}[(a_.) + (b_.)(x_)^2((c_.) + (d_.)(x_)^{(n_.)}((e_.) + (f_.)(x_)^{(p_.)}, x_Symbol] \rightarrow \text{Simp}[b*(a + b*x)*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)}/(d*f*(n+p+3))), x] + \text{Dist}[1/(d*f*(n+p+3)), \text{Int}[(c + d*x)^n*(e + f*x)^p*\text{Simp}[a^2*d*f*(n+p+3) - b*(b*c*e + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(n+p+4) - b*(d*e*(n+2) + c*f*(p+2))]*x, x], x] /; \text{FreeQ}[\{a, b, c, d, e, f, n, p\}, x] \&\& \text{NeQ}[n + p + 3, 0]$

Rule 210

$\text{Int}[(a_) + (b_.)(x_)^2)^{-1}, x_Symbol] \rightarrow \text{Simp}[(-\text{Rt}[-a, 2]*\text{Rt}[-b, 2])^{-1})*\text{ArcTan}[\text{Rt}[-b, 2]*(x/\text{Rt}[-a, 2])], x] /; \text{FreeQ}[\{a, b\}, x] \&\& \text{PosQ}[a/b] \&\& (\text{LtQ}[a, 0] \mid\mid \text{LtQ}[b, 0])$

Rule 303

$\text{Int}[x^2/((a_) + (b_.)(x_)^4), x_Symbol] \rightarrow \text{With}[\{r = \text{Numerator}[\text{Rt}[a/b, 2]], s = \text{Denominator}[\text{Rt}[a/b, 2]]\}, \text{Dist}[1/(2*s), \text{Int}[(r + s*x^2)/(a + b*x^4), x], x] - \text{Dist}[1/(2*s), \text{Int}[(r - s*x^2)/(a + b*x^4), x], x]] /; \text{FreeQ}[\{a, b\}, x] \&\& (\text{GtQ}[a/b, 0] \mid\mid (\text{PosQ}[a/b] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, a]] \&\& \text{AtomQ}[\text{SplitProduct}[\text{SumBaseQ}, b]]))$

Rule 338

$\text{Int}[x^{(m_.)}((a_) + (b_.)(x_)^{(n_.)})^{(p_)}, x_Symbol] \rightarrow \text{Dist}[a^{(p + (m + 1)/n)}, \text{Subst}[\text{Int}[x^m/(1 - b*x^n)^{(p + (m + 1)/n + 1)}], x], x, x/(a + b*x^n)]$

$^{(1/n)}$, x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
 $^{(-1)}$] && IntegersQ[m, p + (m + 1)/n]

Rule 631

Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
 implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
 Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]

Rule 642

Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
 imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
 e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
 2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
 /(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
 & EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
 -2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
 x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
 eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_.))*((d_) + (e_)*(x_)^(m_.),
 x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
 I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{-\frac{3}{2}i \tan^{-1}(a+bx)} x^2 dx &= \int \frac{x^2(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx \\
&= \frac{x(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{3b^2} + \frac{\int \frac{(1-ia-ibx)^{3/4}(-1-a^2+\frac{1}{2}(3i-8a)bx)}{(1+ia+ibx)^{3/4}} dx}{3b^2} \\
&= \frac{(3i-8a)(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{12b^3} + \frac{x(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{3b^2} - (1 \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3} \\
&= -\frac{(36a-i(17-24a^2))(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{24b^3} + \frac{(3i-8a)(1-ia-ibx)^{7/4}}{12b^3}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.06, size = 98, normalized size = 0.20

$$\frac{(-i(i+a+bx))^{7/4} \left(7\sqrt[4]{1+ia+ibx} (3i-8a+4bx) + \sqrt[4]{2} (-17i+36a+24ia^2) {}_2F_1\left(\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; -\frac{1}{2}i(i+a+bx)\right) \right)}{84b^3}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/E^(((3*I)/2)*ArcTan[a + b*x]),x]

[Out] (((-I)*(I + a + b*x))^(7/4)*(7*(1 + I*a + I*b*x)^(1/4)*(3*I - 8*a + 4*b*x) + 2^(1/4)*(-17*I + 36*a + (24*I)*a^2)*Hypergeometric2F1[3/4, 7/4, 11/4, (-1/2*I)*(I + a + b*x)]))/(84*b^3)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{x^2}{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x)**[Out]** int(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="maxima")**[Out]** integrate(x^2/((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2), x)**Fricas [A]**

time = 5.35, size = 555, normalized size = 1.12

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="fricas")

[Out] $\frac{1}{48} \cdot (3b^3 \sqrt{(576Ia^4 + 1728a^3 - 2112Ia^2 - 1224a + 289I)}/b^6) \cdot \log\left(\frac{Ib^3 \sqrt{(576Ia^4 + 1728a^3 - 2112Ia^2 - 1224a + 289I)}/b^6 + (24a^2 - 36Ia - 17) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I)}{(24a^2 - 36Ia - 17)} - 3b^3 \sqrt{(576Ia^4 + 1728a^3 - 2112Ia^2 - 1224a + 289I)}/b^6\right) \cdot \log\left(\frac{-Ib^3 \sqrt{(576Ia^4 + 1728a^3 - 2112Ia^2 - 1224a + 289I)}/b^6 + (24a^2 - 36Ia - 17) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I)}{(24a^2 - 36Ia - 17)} + 3b^3 \sqrt{(-576Ia^4 - 1728a^3 + 2112Ia^2 + 1224a - 289I)}/b^6\right) \cdot \log\left(\frac{Ib^3 \sqrt{(-576Ia^4 - 1728a^3 + 2112Ia^2 + 1224a - 289I)}/b^6 + (24a^2 - 36Ia - 17) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I)}{(24a^2 - 36Ia - 17)} - 3b^3 \sqrt{(-576Ia^4 - 1728a^3 + 2112Ia^2 + 1224a - 289I)}/b^6\right) \cdot \log\left(\frac{-Ib^3 \sqrt{(-576Ia^4 - 1728a^3 + 2112Ia^2 + 1224a - 289I)}/b^6 + (24a^2 - 36Ia - 17) \sqrt{I \sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I)}{(24a^2 - 36Ia - 17)} - 3b^3 \sqrt{(-576Ia^4 - 1728a^3 + 2112Ia^2 + 1224a - 289I)}/b^6\right)$

$I/b^6) + (24*a^2 - 36*I*a - 17)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I))/(24*a^2 - 36*I*a - 17)) - 2*(8*b^3*x^3 + 22*I*b^2*x^2 + 8*a^3 - (40*I*a + 37)*b*x - 38*I*a^2 + 23*a - 23*I)*\sqrt{I*\sqrt{b^2*x^2 + 2*a*b*x + a^2 + 1}/(b*x + a + I)))/b^3$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2),x)`

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2}{\left(\frac{1+ali+bxli}{\sqrt{(a+bx)^2+1}}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2),x)`

[Out] `int(x^2/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)`

3.232 $\int e^{-\frac{3}{2}i \operatorname{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=410

$$\frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} + \frac{3(3+4ia) \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2} b^2}$$

[Out] $\frac{1}{4}(3+4Ia)(1-Ia-Ib*x)^{3/4}(1+Ia+Ib*x)^{1/4}/b^2 + \frac{1}{2}(1-Ia-Ib*x)^{7/4}(1+Ia+Ib*x)^{1/4}/b^2 + \frac{3}{8}(3+4Ia)*\arctan(1-(1-Ia-Ib*x)^{1/4}) * 2^{1/2}/(1+Ia+Ib*x)^{1/4} - \frac{3}{8}(3+4Ia)*\arctan(1+(1-Ia-Ib*x)^{1/4}) * 2^{1/2}/(1+Ia+Ib*x)^{1/4} - \frac{3}{16}(3+4Ia)*\ln(1-(1-Ia-Ib*x)^{1/4}) * 2^{1/2}/(1+Ia+Ib*x)^{1/4} + \frac{3}{16}(3+4Ia)*\ln(1+(1-Ia-Ib*x)^{1/4}) * 2^{1/2}/(1+Ia+Ib*x)^{1/4} + \frac{3}{16}(3+4Ia)*\ln(1+(1-Ia-Ib*x)^{1/4}) * 2^{1/2}/(1+Ia+Ib*x)^{1/4} - \frac{3}{16}(3+4Ia)*\ln(1-(1-Ia-Ib*x)^{1/4}) * 2^{1/2}/(1+Ia+Ib*x)^{1/4}$

Rubi [A]

time = 0.21, antiderivative size = 410, normalized size of antiderivative = 1.00, number of steps used = 14, number of rules used = 11, integrand size = 16, $\frac{\text{number of rules}}{\text{integrand size}} = 0.688$, Rules used = {5203, 81, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{3(3+4ia) \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2} b^2} - \frac{3(3+4ia) \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{4\sqrt{2} b^2} + \frac{\sqrt[4]{1+ia+ibx} (-ia-ibx+1)^{3/4}}{2b^2} + \frac{(3+4ia) \sqrt[4]{1+ia+ibx} (-ia-ibx+1)^{3/4}}{4b^2} - \frac{3(3+4ia) \log\left(\frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}} - \frac{\sqrt{2} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}} + 1\right)}{8\sqrt{2} b^2} + \frac{3(3+4ia) \log\left(\frac{\sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}} + \frac{\sqrt{2} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{1+ia+ibx}} + 1\right)}{8\sqrt{2} b^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[x/E^{((3*I)/2)*\operatorname{ArcTan}[a + b*x]}, x]$

[Out] $((3 + (4*I)*a)*(1 - I*a - I*b*x)^{3/4}(1 + I*a + I*b*x)^{1/4})/(4*b^2) + ((1 - I*a - I*b*x)^{7/4}(1 + I*a + I*b*x)^{1/4})/(2*b^2) + (3*(3 + (4*I)*a)*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{1/4})/(1 + I*a + I*b*x)^{1/4}])/(4*\operatorname{Sqrt}[2]*b^2) - (3*(3 + (4*I)*a)*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{1/4})/(1 + I*a + I*b*x)^{1/4}])/(4*\operatorname{Sqrt}[2]*b^2) - (3*(3 + (4*I)*a)*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a - I*b*x]/\operatorname{Sqrt}[1 + I*a + I*b*x] - (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{1/4})/(1 + I*a + I*b*x)^{1/4}])/(8*\operatorname{Sqrt}[2]*b^2) + (3*(3 + (4*I)*a)*\operatorname{Log}[1 + \operatorname{Sqrt}[1 - I*a - I*b*x]/\operatorname{Sqrt}[1 + I*a + I*b*x] + (\operatorname{Sqrt}[2]*(1 - I*a - I*b*x)^{1/4})/(1 + I*a + I*b*x)^{1/4}])/(8*\operatorname{Sqrt}[2]*b^2)$

Rule 52

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^n/(b*(m+n+1))), x] + \operatorname{Dist}[n*((b*c - a*d)/(b*(m+n+1))), \operatorname{Int}[(a + b*x)^m*(c + d*x)^{(n-1)}, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d\}, x \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& \operatorname{GtQ}[n, 0] \&\& \operatorname{NeQ}[m+n+1, 0] \&\& \operatorname{!(IGtQ}[m, 0] \&\& \operatorname{!(IntegerQ}[n] || (\operatorname{GtQ}[m, 0] \&\& \operatorname{LtQ}[m-n, 0]))) \&\& \operatorname{!ILtQ}[m+n+2, 0] \&\& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

```
Int[((a_.) + (b_.)*(x_)^(m_))*((c_.) + (d_.)*(x_)^(n_)), x_Symbol] := With[
{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m + 1) - 1)*(c - a*(d/b) +
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 81

```
Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_)^(n_))*((e_.) + (f_.)*(x_)^(p
_.)), x_Symbol] := Simp[b*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p +
2))), x] + Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(d*f*(
n + p + 2)), Int[(c + d*x)^n*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f
, n, p}, x] && NeQ[n + p + 2, 0]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
```

e}, x] && EqQ[2*c*d - b*e, 0]

Rule 1176

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]

Rule 1179

Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] :> With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_.)*((d_) + (e_)*(x_)^m_), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
\int e^{-\frac{3}{2}i \tan^{-1}(a+bx)} x dx &= \int \frac{x(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx \\
&= \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} + \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} + \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} - \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} - \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} + \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} - \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} + \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} - \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b} \\
&= \frac{(3+4ia)(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{4b^2} + \frac{(1-ia-ibx)^{7/4} \sqrt[4]{1+ia+ibx}}{2b^2} + \frac{(3i-4a) \int \frac{(1-ia-ibx)^{3/4}}{(1+ia+ibx)^{3/4}} dx}{4b}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.03, size = 84, normalized size = 0.20

$$\frac{i(-i(i+a+bx))^{7/4} \left(7i \sqrt[4]{1+ia+ibx} + \sqrt[4]{2} (-3i+4a) {}_2F_1\left(\frac{3}{4}, \frac{7}{4}; \frac{11}{4}; -\frac{1}{2}i(i+a+bx)\right) \right)}{14b^2}$$

Antiderivative was successfully verified.

[In] Integrate[x/E^(((3*I)/2)*ArcTan[a + b*x]),x]

[Out] ((-1/14*I)*((-I)*(I + a + b*x))^(7/4)*((7*I)*(1 + I*a + I*b*x)^(1/4) + 2^(1/4)*(-3*I + 4*a)*Hypergeometric2F1[3/4, 7/4, 11/4, (-1/2*I)*(I + a + b*x)])/b^2

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{x}{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2), x)**[Out]** int(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2), x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2), x, algorithm="maxima")**[Out]** integrate(x/((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2), x)**Fricas [A]**

time = 6.18, size = 433, normalized size = 1.06

$$\frac{1}{8} \sqrt{\frac{16I^2a^2 - 24a - 9I}{b^4}} \log\left(\frac{-Ib^2\sqrt{16I^2a^2 - 24a - 9I} + 24a - 9I}{b^4}\right) - \frac{(4a - 3I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(bx + a + I)(4a - 3I)} - 3b^2\sqrt{\frac{16I^2a^2 - 24a - 9I}{b^4}} \log\left(\frac{-Ib^2\sqrt{16I^2a^2 - 24a - 9I} - (4a - 3I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}}{b^4}\right) + 3b^2\sqrt{\frac{-16I^2a^2 - 24a + 9I}{b^4}} \log\left(\frac{-Ib^2\sqrt{-16I^2a^2 - 24a + 9I} - (4a - 3I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}}{b^4}\right) - (4a - 3I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I) - 3b^2\sqrt{\frac{-16I^2a^2 - 24a + 9I}{b^4}} \log\left(\frac{-Ib^2\sqrt{-16I^2a^2 - 24a + 9I} - (-16I^2a^2 - 24a + 9I)}{b^4}\right) - (4a - 3I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I) - 2(2b^2x^2 - 2a^2 + 7Ibx + 3Ia - 5)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}/(bx + a + I)/b^2$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2), x, algorithm="fricas")

[Out] 1/8*(3*b^2*sqrt(-(16*I*a^2 + 24*a - 9*I)/b^4)*log(-I*b^2*sqrt(-(16*I*a^2 + 24*a - 9*I)/b^4) - (4*a - 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a - 3*I) - 3*b^2*sqrt(-(16*I*a^2 + 24*a - 9*I)/b^4)*log(-(-I*b^2*sqrt(-(16*I*a^2 + 24*a - 9*I)/b^4) - (4*a - 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a - 3*I) + 3*b^2*sqrt(-16*I*a^2 - 24*a + 9*I)/b^4)*log(-I*b^2*sqrt(-16*I*a^2 - 24*a + 9*I)/b^4) - (4*a - 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a - 3*I) - 3*b^2*sqrt(-16*I*a^2 - 24*a + 9*I)/b^4)*log(-(-I*b^2*sqrt(-16*I*a^2 - 24*a + 9*I)/b^4) - (4*a - 3*I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I)))/(4*a - 3*I) - 2*(2*b^2*x^2 - 2*a^2 + 7*I*b*x + 3*I*a - 5)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b^2

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2),x)

[Out] Timed out

Giac [F(-2)]
time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
the root of a polynomial with parameters. This might be wrong.The choice wa
s done

Mupad [F]
time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x}{\left(\frac{1+ai+bx\ i}{\sqrt{(a+bx)^2+1}}\right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2),x)

[Out] int(x/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)

3.233 $\int e^{-\frac{3}{2}i\text{ArcTan}(a+bx)} dx$

Optimal. Leaf size=338

$$\frac{i(1-ia-ibx)^{3/4}\sqrt[4]{1+ia+ibx}}{b} - \frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b} + \frac{3i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)}{\sqrt{2}b}$$

[Out] $-I*(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(1/4)}/b-3/2*I*\arctan(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}+3/2*I*\arctan(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})/b*2^{(1/2)}+3/4*I*\ln(1-(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}-3/4*I*\ln(1+(1-I*a-I*b*x)^{(1/4)}*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})/b*2^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 338, normalized size of antiderivative = 1.00, number of steps used = 13, number of rules used = 10, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.714$,

Rules used = {5201, 52, 65, 338, 303, 1176, 631, 210, 1179, 642}

$$\frac{3i\text{ArcTan}\left(1 - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} + \frac{3i\text{ArcTan}\left(1 + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}}\right)}{\sqrt{2}b} - \frac{i(-ia-ibx+1)^{3/4}\sqrt[4]{ia+ibx+1}}{b} + \frac{3i\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} - \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b} - \frac{3i\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} + \frac{\sqrt{2}\sqrt[4]{-ia-ibx+1}}{\sqrt[4]{ia+ibx+1}} + 1\right)}{2\sqrt{2}b}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((-3*I)/2)*\text{ArcTan}[a + b*x]}, x]$

[Out] $((-I)*(1 - I*a - I*b*x)^{(3/4)}*(1 + I*a + I*b*x)^{(1/4)})/b - ((3*I)*\text{ArcTan}[1 - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(\text{Sqrt}[2]*b) + ((3*I)*\text{ArcTan}[1 + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(\text{Sqrt}[2]*b) + (((3*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] - (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(\text{Sqrt}[2]*b) - (((3*I)/2)*\text{Log}[1 + \text{Sqrt}[1 - I*a - I*b*x]/\text{Sqrt}[1 + I*a + I*b*x] + (\text{Sqrt}[2]*(1 - I*a - I*b*x)^{(1/4)})/(1 + I*a + I*b*x)^{(1/4)}])/(\text{Sqrt}[2]*b)$

Rule 52

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}*((c + d*x)^n/(b*(m + n + 1))), x] + \text{Dist}[n*((b*c - a*d)/(b*(m + n + 1))), \text{Int}[(a + b*x)^m*(c + d*x)^{(n - 1)}, x], x] /;$ $\text{FreeQ}\{a, b, c, d\}, x \ \&\& \ \text{NeQ}[b*c - a*d, 0] \ \&\& \ \text{GtQ}[n, 0] \ \&\& \ \text{NeQ}[m + n + 1, 0] \ \&\& \ !(\text{IGtQ}[m, 0] \ \&\& \ (!\text{IntegerQ}[n] \ || \ (\text{GtQ}[m, 0] \ \&\& \ \text{LtQ}[m - n, 0]))) \ \&\& \ !\text{ILtQ}[m + n + 2, 0] \ \&\& \ \text{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 65

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{With}[\{p = \text{Denominator}[m]\}, \text{Dist}[p/b, \text{Subst}[\text{Int}[x^{(p*(m + 1) - 1)}*(c - a*(d/b) +$

```
d*(x^p/b))^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 303

```
Int[(x_)^2/((a_) + (b_.)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_.)*((a_) + (b_.)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_.)*(x_) + (c_.)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_.)*(x_))/((a_.) + (b_.)*(x_) + (c_.)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_.)*(x_)^2)/((a_) + (c_.)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x], x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x] /; FreeQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5201

```
Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_), x_Symbol] := Int[(1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2)), x] /; FreeQ[{a, b, c, n}, x]
```

Rubi steps

$$\begin{aligned}
\int e^{-\frac{3}{2}i \tan^{-1}(a+bx)} dx &= \int \frac{(1 - ia - ibx)^{3/4}}{(1 + ia + ibx)^{3/4}} dx \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} + \frac{3}{2} \int \frac{1}{\sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}} dx \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} + \frac{(6i) \text{Subst}\left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1 - ia - ibx}\right)}{b} \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} + \frac{(6i) \text{Subst}\left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{b} \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} - \frac{(3i) \text{Subst}\left(\int \frac{1-x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{b} + \dots \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} + \frac{(3i) \text{Subst}\left(\int \frac{1}{1-\sqrt{2} x+x^2} dx, x, \frac{\sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{2b} \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} + \frac{3i \log\left(1 + \frac{\sqrt{1 - ia - ibx}}{\sqrt{1 + ia + ibx}} - \frac{\sqrt{2} \sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{2\sqrt{2} b} \\
&= -\frac{i(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{b} - \frac{3i \tan^{-1}\left(1 - \frac{\sqrt{2} \sqrt[4]{1 - ia - ibx}}{\sqrt[4]{1 + ia + ibx}}\right)}{\sqrt{2} b} + \dots
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 43, normalized size = 0.13

$$-\frac{8ie^{\frac{1}{2}i \text{ArcTan}(a+bx)} {}_2F_1\left(\frac{1}{4}, 2; \frac{5}{4}; -e^{2i \text{ArcTan}(a+bx)}\right)}{b}$$

Antiderivative was successfully verified.

[In] Integrate[E^(((−3*I)/2)*ArcTan[a + b*x]),x]

[Out] ((−8*I)*E^((I/2)*ArcTan[a + b*x])*Hypergeometric2F1[1/4, 2, 5/4, −E^((2*I)*ArcTan[a + b*x])])/b

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x)

[Out] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="maxima")

[Out] integrate(((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(-3/2), x)

Fricas [A]

time = 4.29, size = 255, normalized size = 0.75

$$\frac{b\sqrt{\frac{9i}{b^2}} \log\left(\frac{1}{2}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2+2abx+a^2+1}}{bx+a+i}}\right) - b\sqrt{\frac{9i}{b^2}} \log\left(-\frac{1}{2}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2+2abx+a^2+1}}{bx+a+i}}\right) + b\sqrt{\frac{9i}{b^2}} \log\left(\frac{1}{2}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2+2abx+a^2+1}}{bx+a+i}}\right) - b\sqrt{\frac{9i}{b^2}} \log\left(-\frac{1}{2}b\sqrt{\frac{9i}{b^2}} + \sqrt{\frac{i\sqrt{b^2x^2+2abx+a^2+1}}{bx+a+i}}\right) - 2(bx+a+i)\sqrt{\frac{i\sqrt{b^2x^2+2abx+a^2+1}}{bx+a+i}}}{2b}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2),x, algorithm="fricas")

[Out] 1/2*(b*sqrt(9*I/b^2)*log(1/3*I*b*sqrt(9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(9*I/b^2)*log(-1/3*I*b*sqrt(9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) + b*sqrt(-9*I/b^2)*log(1/3*I*b*sqrt(-9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - b*sqrt(-9*I/b^2)*log(-1/3*I*b*sqrt(-9*I/b^2) + sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))) - 2*(b*x + a + I)*sqrt(I*sqrt(b^2*x^2 + 2*a*b*x + a^2 + 1)/(b*x + a + I))/b

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{i(a+bx)+1}{\sqrt{(a+bx)^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2), x)``[Out] Integral(((I*(a + b*x) + 1)/sqrt((a + b*x)**2 + 1))**(-3/2), x)`**Giac [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2), x, algorithm="giac")`

`[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
 UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for
 the root of a polynomial with parameters. This might be wrong.The choice wa
 s done`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{\left(\frac{1+a1i+bx1i}{\sqrt{(a+bx)^2+1}}\right)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)``[Out] int(1/((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2))^(3/2), x)`

$$3.234 \quad \int \frac{e^{-\frac{3}{2}i \operatorname{ArcTan}(a+bx)}}{x} dx$$

Optimal. Leaf size=427

$$-\frac{2(i+a)^{3/4} \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{(i-a)^{3/4}} - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}}\right)$$

[Out] $-2*(I+a)^{(3/4)}*\arctan((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4)})/(I-a)^{(3/4)}-2*(I+a)^{(3/4)}*\operatorname{arctanh}((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4)})/(I-a)^{(3/4)}+1/2*\ln(1-(1-I*a-I*b*x)^{(1/4)})*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})*2^{(1/2)}-1/2*\ln(1+(1-I*a-I*b*x)^{(1/4)})*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)}+(1-I*a-I*b*x)^{(1/2)}/(1+I*a+I*b*x)^{(1/2)})*2^{(1/2)}-\arctan(1-(1-I*a-I*b*x)^{(1/4)})*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})*2^{(1/2)}+\operatorname{arctan}(1+(1-I*a-I*b*x)^{(1/4)})*2^{(1/2)}/(1+I*a+I*b*x)^{(1/4)})*2^{(1/2)}$

Rubi [A]

time = 0.20, antiderivative size = 427, normalized size of antiderivative = 1.00, number of steps used = 18, number of rules used = 15, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.833$, Rules used = {5203, 132, 65, 338, 303, 1176, 631, 210, 1179, 642, 12, 95, 218, 214, 211}

$$\frac{2(a+i)^{3/4} \operatorname{ArcTan}\left(\frac{\sqrt{a+1} \sqrt{ia+ibx+1}}{\sqrt{-a+1} \sqrt{-ia-ibx+1}}\right)}{(-a+i)^{3/4}} - \sqrt{2} \operatorname{ArcTan}\left(1 - \frac{\sqrt{2} \sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}}\right) + \sqrt{2} \operatorname{ArcTan}\left(1 + \frac{\sqrt{2} \sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}}\right) + \frac{\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} - \frac{\sqrt{2} \sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} + 1\right)}{\sqrt{2}} - \frac{\log\left(\frac{\sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} + \frac{\sqrt{2} \sqrt{-ia-ibx+1}}{\sqrt{ia+ibx+1}} + 1\right)}{\sqrt{2}} - \frac{2(a+i)^{3/4} \operatorname{tanh}^{-1}\left(\frac{\sqrt{a+1} \sqrt{ia+ibx+1}}{\sqrt{-a+1} \sqrt{-ia-ibx+1}}\right)}{(-a+i)^{3/4}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(((3*I)/2)*ArcTan[a + b*x]))*x], x]`

[Out] $(-2*(I+a)^{(3/4)}*\operatorname{ArcTan}(((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)})/((I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4)})))/(I-a)^{(3/4)} - \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 - (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}] + \operatorname{Sqrt}[2]*\operatorname{ArcTan}[1 + (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}] - (2*(I+a)^{(3/4)}*\operatorname{ArcTan}(((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)})/((I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4)})))/(I-a)^{(3/4)} + \operatorname{Log}[1 + \operatorname{Sqrt}[1-I*a-I*b*x]/\operatorname{Sqrt}[1+I*a+I*b*x] - (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}]/\operatorname{Sqrt}[2] - \operatorname{Log}[1 + \operatorname{Sqrt}[1-I*a-I*b*x]/\operatorname{Sqrt}[1+I*a+I*b*x] + (\operatorname{Sqrt}[2]*(1-I*a-I*b*x)^{(1/4)})/(1+I*a+I*b*x)^{(1/4)}]/\operatorname{Sqrt}[2]$

Rule 12

`Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]`

Rule 65

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := With[{p = Denominator[m]}, Dist[p/b, Subst[Int[x^(p*(m+1)-1)*(c-a*(d/b) +`

```
d*(x^p/b)^n, x], x, (a + b*x)^(1/p)], x]] /; FreeQ[{a, b, c, d}, x] && NeQ
[b*c - a*d, 0] && LtQ[-1, m, 0] && LeQ[-1, n, 0] && LeQ[Denominator[n], Den
ominator[m]] && IntLinearQ[a, b, c, d, m, n, x]
```

Rule 95

```
Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x
_)), x_Symbol] :=> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1)
- 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)
], x]] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n]
&& LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]
```

Rule 132

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x
_))^(p_), x_Symbol] :=> Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x
)^(m), x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(
a + b*x)*(c + d*x)^(-p - 1) - (b*d^(-p - 1)*f^p)/(e + f*x)^p, x], x] /; Fre
eQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (
GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))
```

Rule 210

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :=> Simp[(-(Rt[-a, 2]*Rt[-b, 2])^(
-1))*ArcTan[Rt[-b, 2]*(x/Rt[-a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] &
& (LtQ[a, 0] || LtQ[b, 0])
```

Rule 211

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :=> Simp[(Rt[a/b, 2]/a)*ArcTan[x/R
t[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]
```

Rule 214

```
Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :=> Simp[(Rt[-a/b, 2]/a)*ArcTanh[x
/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]
```

Rule 218

```
Int[((a_) + (b_.)*(x_)^4)^(-1), x_Symbol] :=> With[{r = Numerator[Rt[-a/b, 2
]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x]
+ Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b
, 0]
```

Rule 303

```
Int[(x_)^2/((a_) + (b_)*(x_)^4), x_Symbol] := With[{r = Numerator[Rt[a/b,
2]], s = Denominator[Rt[a/b, 2]]}, Dist[1/(2*s), Int[(r + s*x^2)/(a + b*x^4
), x], x] - Dist[1/(2*s), Int[(r - s*x^2)/(a + b*x^4), x], x]] /; FreeQ[{a,
b}, x] && (GtQ[a/b, 0] || (PosQ[a/b] && AtomQ[SplitProduct[SumBaseQ, a]] &
& AtomQ[SplitProduct[SumBaseQ, b]]))
```

Rule 338

```
Int[(x_)^(m_)*((a_) + (b_)*(x_)^(n_))^(p_), x_Symbol] := Dist[a^(p + (m +
1)/n), Subst[Int[x^m/(1 - b*x^n)^(p + (m + 1)/n + 1), x], x, x/(a + b*x^n)
^(1/n)], x] /; FreeQ[{a, b}, x] && IGtQ[n, 0] && LtQ[-1, p, 0] && NeQ[p, -2
^(-1)] && IntegersQ[m, p + (m + 1)/n]
```

Rule 631

```
Int[((a_) + (b_)*(x_) + (c_)*(x_)^2)^(-1), x_Symbol] := With[{q = 1 - 4*S
implify[a*(c/b^2)]}, Dist[-2/b, Subst[Int[1/(q - x^2), x], x, 1 + 2*c*(x/b)
], x] /; RationalQ[q] && (EqQ[q^2, 1] || !RationalQ[b^2 - 4*a*c])] /; Free
Q[{a, b, c}, x] && NeQ[b^2 - 4*a*c, 0]
```

Rule 642

```
Int[((d_) + (e_)*(x_))/((a_) + (b_)*(x_) + (c_)*(x_)^2), x_Symbol] := S
imp[d*(Log[RemoveContent[a + b*x + c*x^2, x]]/b), x] /; FreeQ[{a, b, c, d,
e}, x] && EqQ[2*c*d - b*e, 0]
```

Rule 1176

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
2*(d/e), 2]}, Dist[e/(2*c), Int[1/Simp[d/e + q*x + x^2, x], x], x] + Dist[e
/(2*c), Int[1/Simp[d/e - q*x + x^2, x], x], x]] /; FreeQ[{a, c, d, e}, x] &
& EqQ[c*d^2 - a*e^2, 0] && PosQ[d*e]
```

Rule 1179

```
Int[((d_) + (e_)*(x_)^2)/((a_) + (c_)*(x_)^4), x_Symbol] := With[{q = Rt[
-2*(d/e), 2]}, Dist[e/(2*c*q), Int[(q - 2*x)/Simp[d/e + q*x - x^2, x], x],
x] + Dist[e/(2*c*q), Int[(q + 2*x)/Simp[d/e - q*x - x^2, x], x], x]] /; Fre
eQ[{a, c, d, e}, x] && EqQ[c*d^2 - a*e^2, 0] && NegQ[d*e]
```

Rule 5203

```
Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))]*(n_))*((d_) + (e_)*(x_))^(m_),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```


Rubi steps

$$\begin{aligned}
\int \frac{e^{-\frac{3}{2}i \tan^{-1}(a+bx)}}{x} dx &= \int \frac{(1-ia-ibx)^{3/4}}{x(1+ia+ibx)^{3/4}} dx \\
&= -\left((-1+ia) \int \frac{1}{x^4 \sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}} dx \right) - (ib) \int \frac{1}{\sqrt[4]{1-ia-ibx} (1+ia+ibx)^{3/4}} dx \\
&= 4\text{Subst} \left(\int \frac{x^2}{(2-x^4)^{3/4}} dx, x, \sqrt[4]{1-ia-ibx} \right) + (4(1-ia))\text{Subst} \left(\int \frac{1}{-1-ia-(1+ia+ibx)^{3/4}} dx, x, \sqrt[4]{1-ia-ibx} \right) \\
&= 4\text{Subst} \left(\int \frac{x^2}{1+x^4} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} \right) - \frac{(2(i+a))\text{Subst} \left(\int \frac{1}{\sqrt{i-a}-\sqrt{i+a}x} dx, x, \frac{\sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} \right)}{\sqrt{i-a}} \\
&= -\frac{2(i+a)^{3/4} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} - \frac{2(i+a)^{3/4} \tanh^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} \\
&= -\frac{2(i+a)^{3/4} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} - \frac{2(i+a)^{3/4} \tanh^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} \\
&= -\frac{2(i+a)^{3/4} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} - \frac{2(i+a)^{3/4} \tanh^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} \\
&= -\frac{2(i+a)^{3/4} \tan^{-1} \left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}} \right)}{(i-a)^{3/4}} - \sqrt{2} \tan^{-1} \left(1 - \frac{\sqrt{2} \sqrt[4]{1-ia-ibx}}{\sqrt[4]{1+ia+ibx}} \right)
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 128, normalized size = 0.30

$$\frac{2(-i(i+a+bx))^{3/4} \left(\sqrt[4]{2} (1+ia+ibx)^{3/4} {}_2F_1 \left(\frac{3}{4}, \frac{3}{4}, \frac{7}{4}; -\frac{1}{2}i(i+a+bx) \right) - 2 {}_2F_1 \left(\frac{3}{4}, 1; \frac{7}{4}, \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx} \right) \right)}{3(1+ia+ibx)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a + b*x]))*x, x]

[Out] (2*((-I)*(I + a + b*x))^(3/4)*(2^(1/4)*(1 + I*a + I*b*x)^(3/4)*Hypergeometric2F1[3/4, 3/4, 7/4, (-1/2*I)*(I + a + b*x)] - 2*Hypergeometric2F1[3/4, 1, 7/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)]))/((3*(1 + I*a + I*b*x)^(3/4)))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}\right)^{\frac{3}{2}} x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x)**[Out]** int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x, algorithm="maxima")**[Out]** integrate(1/(x*((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2)), x)**Fricas [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 629 vs. 2(284) = 568.

time = 7.02, size = 629, normalized size = 1.47

$$\frac{1}{2} \sqrt{4I} \log\left(\frac{1}{2} \sqrt{4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}\right) + \frac{1}{2} \sqrt{4I} \log\left(-\frac{1}{2} \sqrt{4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}\right) - \frac{1}{2} \sqrt{-4I} \log\left(\frac{1}{2} \sqrt{-4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}\right) + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}} \log\left(-\frac{1}{2} \sqrt{-4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}\right) + \left(-a^3 + 3I a^2 - 3a - I\right) \left(a^3 - 3I a^2 - 3a + I\right)^{-1/4} \log\left(\left(a + I\right) \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}\right) + (a - I) \left(-a^3 + 3I a^2 - 3a - I\right) \left(a^3 - 3I a^2 - 3a + I\right)^{-1/4} \left(a + I\right) - \left(-a^3 + 3I a^2 - 3a - I\right) \left(a^3 - 3I a^2 - 3a + I\right)^{-1/4} \log\left(\left(a + I\right) \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}\right) - (a - I) \left(-a^3 + 3I a^2 - 3a - I\right) \left(a^3 - 3I a^2 - 3a + I\right)^{-1/4} \left(a + I\right) + I \left(-a^3 + 3I a^2 - 3a - I\right) \left(a^3 - 3I a^2 - 3a + I\right)^{-1/4} \left(a + I\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x, algorithm="fricas")

[Out] $-1/2 \sqrt{4I} \log(1/2 \sqrt{4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}) / (b x + a + I) + 1/2 \sqrt{4I} \log(-1/2 \sqrt{4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}) / (b x + a + I) - 1/2 \sqrt{-4I} \log(1/2 \sqrt{-4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}) / (b x + a + I) + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}} \log(-1/2 \sqrt{-4I} + \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}) / (b x + a + I) + (-a^3 + 3I a^2 - 3a - I) / (a^3 - 3I a^2 - 3a + I)^{1/4} \log((a + I) \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}) / (b x + a + I) + (a - I) (-a^3 + 3I a^2 - 3a - I) / (a^3 - 3I a^2 - 3a + I)^{1/4} / (a + I) - (-a^3 + 3I a^2 - 3a - I) / (a^3 - 3I a^2 - 3a + I)^{1/4} \log((a + I) \sqrt{I \sqrt{b^2 x^2 + 2a b x + a^2 + 1}}) / (b x + a + I) - (a - I) (-a^3 + 3I a^2 - 3a - I) / (a^3 - 3I a^2 - 3a + I)^{1/4} / (a + I) + I (-a^3 + 3I a^2 - 3a - I) / (a^3 - 3I a^2 - 3a + I)^{1/4} / (a + I)$

$$\frac{3a - I}{(a^3 - 3Ia^2 - 3a + I)^{1/4}} \log\left(\frac{(a + I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(bx + a + I)} + \frac{Ia + 1}{(a^3 - 3Ia^2 - 3a + I)^{1/4}} \frac{-(a^3 + 3Ia^2 - 3a - I)}{(a + I)} - I \frac{-(a^3 + 3Ia^2 - 3a - I)}{(a^3 - 3Ia^2 - 3a + I)^{1/4}} \log\left(\frac{(a + I)\sqrt{I\sqrt{b^2x^2 + 2abx + a^2 + 1}}}{(bx + a + I)} + \frac{-Ia - 1}{(a^3 - 3Ia^2 - 3a + I)^{1/4}} \frac{-(a^3 + 3Ia^2 - 3a - I)}{(a + I)}\right)\right)$$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2)/x,x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x \left(\frac{1+ali+bxli}{\sqrt{(a+bx)^2+1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2)))^(3/2),x)

[Out] int(1/(x*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2)))^(3/2), x)

3.235 $\int \frac{e^{-\frac{3}{2}i \operatorname{ArcTan}(a+bx)}}{x^2} dx$

Optimal. Leaf size=211

$$\frac{(1-ia-ibx)^{3/4} \sqrt[4]{1+ia+ibx}}{(1+ia)x} - \frac{3ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{(i-a)^{7/4} \sqrt[4]{i+a}} - \frac{3ib \tanh^{-1}\left(\frac{\sqrt[4]{i+a} \sqrt[4]{1+ia+ibx}}{\sqrt[4]{i-a} \sqrt[4]{1-ia-ibx}}\right)}{(i-a)^{7/4} \sqrt[4]{i+a}}$$

[Out] $-(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(1/4)}/(1+I*a)/x-3*I*b*\arctan((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4))}/(I-a)^{(7/4)}/(I+a)^{(1/4)}-3*I*b*\operatorname{arctanh}((I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}/(I-a)^{(1/4)}/(1-I*a-I*b*x)^{(1/4))}/(I-a)^{(7/4)}/(I+a)^{(1/4)}$

Rubi [A]

time = 0.11, antiderivative size = 211, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 18, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5203, 96, 95, 218, 214, 211}

$$\frac{3ib \operatorname{ArcTan}\left(\frac{\sqrt[4]{a+i} \sqrt[4]{ia+ibx+1}}{\sqrt[4]{-a+i} \sqrt[4]{-ia-ibx+1}}\right)}{(-a+i)^{7/4} \sqrt[4]{a+i}} - \frac{(-ia-ibx+1)^{3/4} \sqrt[4]{ia+ibx+1}}{(1+ia)x} - \frac{3ib \tanh^{-1}\left(\frac{\sqrt[4]{a+i} \sqrt[4]{ia+ibx+1}}{\sqrt[4]{-a+i} \sqrt[4]{-ia-ibx+1}}\right)}{(-a+i)^{7/4} \sqrt[4]{a+i}}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^(((3*I)/2)*ArcTan[a + b*x]))*x^2),x]`

[Out] $-\left(\frac{(1-I*a-I*b*x)^{(3/4)}*(1+I*a+I*b*x)^{(1/4)}}{(1+I*a)*x}\right) - \left(\frac{3*I*b*\operatorname{ArcTan}\left[\frac{(I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}}{(I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4)}}\right]}{(I-a)^{(7/4)}*(I+a)^{(1/4)}} - \frac{3*I*b*\operatorname{ArcTanh}\left[\frac{(I+a)^{(1/4)}*(1+I*a+I*b*x)^{(1/4)}}{(I-a)^{(1/4)}*(1-I*a-I*b*x)^{(1/4)}}\right]}{(I-a)^{(7/4)}*(I+a)^{(1/4)}}\right)$

Rule 95

`Int[(((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_))/((e_.) + (f_.)*(x_)), x_Symbol] :> With[{q = Denominator[m]}, Dist[q, Subst[Int[x^(q*(m + 1) - 1)/(b*e - a*f - (d*e - c*f)*x^q), x], x, (a + b*x)^(1/q)/(c + d*x)^(1/q)], x] /; FreeQ[{a, b, c, d, e, f}, x] && EqQ[m + n + 1, 0] && RationalQ[n] && LtQ[-1, m, 0] && SimplerQ[a + b*x, c + d*x]`

Rule 96

`Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(a + b*x)^(m + 1)*(c + d*x)^n*((e + f*x)^(p + 1))/((m + 1)*(b*e - a*f)), x] - Dist[n*((d*e - c*f)/((m + 1)*(b*e - a*f))], Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1)*(e + f*x)^p, x], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && GtQ[n, 0] && (SumSimpler`

Q[m, 1] || !SumSimplerQ[p, 1]) && NeQ[m, -1]

Rule 211

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[a/b, 2]/a)*ArcTan[x/Rt[a/b, 2]], x] /; FreeQ[{a, b}, x] && PosQ[a/b]

Rule 214

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(Rt[-a/b, 2]/a)*ArcTanh[x/Rt[-a/b, 2]], x] /; FreeQ[{a, b}, x] && NegQ[a/b]

Rule 218

Int[((a_) + (b_)*(x_)^4)^(-1), x_Symbol] := With[{r = Numerator[Rt[-a/b, 2]], s = Denominator[Rt[-a/b, 2]]}, Dist[r/(2*a), Int[1/(r - s*x^2), x], x] + Dist[r/(2*a), Int[1/(r + s*x^2), x], x]] /; FreeQ[{a, b}, x] && !GtQ[a/b, 0]

Rule 5203

Int[E^(ArcTan[(c_)*((a_) + (b_)*(x_))])*(n_)*((d_) + (e_)*(x_))^(m_), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-\frac{3}{2}i \tan^{-1}(a+bx)}}{x^2} dx &= \int \frac{(1 - ia - ibx)^{3/4}}{x^2(1 + ia + ibx)^{3/4}} dx \\
 &= -\frac{(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{(1 + ia)x} + \frac{(3b) \int \frac{1}{x \sqrt[4]{1 - ia - ibx} (1 + ia + ibx)^{3/4}} dx}{2(i - a)} \\
 &= -\frac{(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{(1 + ia)x} + \frac{(6b) \text{Subst} \left(\int \frac{1}{-1 - ia - (-1 + ia)x^4} dx, x, \frac{\sqrt[4]{1 + ia + ibx}}{\sqrt[4]{1 - ia - ibx}} \right)}{i - a} \\
 &= -\frac{(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{(1 + ia)x} - \frac{(3ib) \text{Subst} \left(\int \frac{1}{\sqrt{i - a} - \sqrt{i + a} x^2} dx, x, \frac{\sqrt[4]{1 + ia + ibx}}{\sqrt[4]{1 - ia - ibx}} \right)}{(i - a)^{3/2}} \\
 &= -\frac{(1 - ia - ibx)^{3/4} \sqrt[4]{1 + ia + ibx}}{(1 + ia)x} - \frac{3ib \tan^{-1} \left(\frac{\sqrt[4]{i + a} \sqrt[4]{1 + ia + ibx}}{\sqrt[4]{i - a} \sqrt[4]{1 - ia - ibx}} \right)}{(i - a)^{7/4} \sqrt[4]{i + a}} - \frac{3ib \tan^{-1} \left(\frac{\sqrt[4]{i + a} \sqrt[4]{1 + ia + ibx}}{\sqrt[4]{i - a} \sqrt[4]{1 - ia - ibx}} \right)}{(i - a)^{7/4} \sqrt[4]{i + a}}
 \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 107, normalized size = 0.51

$$-\frac{(-i(i+a+bx))^{3/4} \left(1+a^2+ibx+abx-2ibx {}_2F_1\left(\frac{3}{4}, 1; \frac{7}{4}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right)\right)}{(1+a^2)x(1+ia+ibx)^{3/4}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(((3*I)/2)*ArcTan[a + b*x]))*x^2, x]

[Out] -((((-I)*(I + a + b*x))^(3/4)*(1 + a^2 + I*b*x + a*b*x - (2*I)*b*x*Hypergeometric2F1[3/4, 1, 7/4, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)])))/((1 + a^2)*x*(1 + I*a + I*b*x)^(3/4))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{1}{\left(\frac{1+i(bx+a)}{\sqrt{1+(bx+a)^2}}\right)^{\frac{3}{2}} x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2, x)

[Out] int(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2, x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2, x, algorithm="maxima")

[Out] integrate(1/(x^2*((I*b*x + I*a + 1)/sqrt((b*x + a)^2 + 1))^(3/2)), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 613 vs. 2(137) = 274.

time = 5.54, size = 613, normalized size = 2.91

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x, algorithm="fricas")

[Out] $\frac{1}{2} \cdot (3 \cdot (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (-Ia - 1) \cdot x \cdot \log((b \cdot \sqrt{I \sqrt{b^2x^2 + 2a \cdot b \cdot x + a^2 + 1}})/(b \cdot x + a + I)) + (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (a^2 - 2Ia - 1))/b + 3 \cdot (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (Ia + 1) \cdot x \cdot \log((b \cdot \sqrt{I \sqrt{b^2x^2 + 2a \cdot b \cdot x + a^2 + 1}})/(b \cdot x + a + I)) - (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (a^2 - 2Ia - 1))/b - 3 \cdot (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (a - I) \cdot x \cdot \log((b \cdot \sqrt{I \sqrt{b^2x^2 + 2a \cdot b \cdot x + a^2 + 1}})/(b \cdot x + a + I)) - (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (Ia^2 + 2a - I))/b + 3 \cdot (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (a - I) \cdot x \cdot \log((b \cdot \sqrt{I \sqrt{b^2x^2 + 2a \cdot b \cdot x + a^2 + 1}})/(b \cdot x + a + I)) - (-b^4/(a^8 - 6Ia^7 - 14a^6 + 14Ia^5 + 14Ia^3 + 14a^2 - 6Ia - 1))^{1/4} \cdot (-Ia^2 - 2a + I))/b + 2 \cdot (b \cdot x + a + I) \cdot \sqrt{I \sqrt{b^2x^2 + 2a \cdot b \cdot x + a^2 + 1}}/(b \cdot x + a + I)))/((a - I) \cdot x)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)**2)**(1/2))**(3/2)/x**2,x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/((1+I*(b*x+a))/(1+(b*x+a)^2)^(1/2))^(3/2)/x^2,x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, need to choose a branch for the root of a polynomial with parameters. This might be wrong.The choice was done

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{1}{x^2 \left(\frac{1+a1i+b x 1i}{\sqrt{(a+b x)^2+1}} \right)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2)))^(3/2),x)

[Out] int(1/(x^2*((a*1i + b*x*1i + 1)/((a + b*x)^2 + 1)^(1/2)))^(3/2), x)

3.236 $\int e^{n \operatorname{ArcTan}(a+bx)} x^m dx$

Optimal. Leaf size=140

$$\frac{x^{1+m}(1-ia-ibx)^{\frac{in}{2}}(1+ia+ibx)^{-\frac{in}{2}}\left(1-\frac{bx}{i-a}\right)^{\frac{in}{2}}\left(1+\frac{bx}{i+a}\right)^{-\frac{in}{2}}F_1\left(1+m;-\frac{in}{2},\frac{in}{2};2+m;-\frac{bx}{i+a},\frac{bx}{i-a}\right)}{1+m}$$

[Out] $x^{(1+m)}*(1-I*a-I*b*x)^{(1/2*I*n)}*(1-b*x/(I-a))^{(1/2*I*n)}*AppellF1(1+m,1/2*I*n,-1/2*I*n,2+m,b*x/(I-a),-b*x/(I+a))/(1+m)/((1+I*a+I*b*x)^{(1/2*I*n)})/((1+b*x/(I+a))^{(1/2*I*n)})$

Rubi [A]

time = 0.06, antiderivative size = 140, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5203, 140, 138}

$$\frac{x^{m+1}(-ia-ibx+1)^{\frac{in}{2}}(ia+ibx+1)^{-\frac{in}{2}}\left(1-\frac{bx}{-a+i}\right)^{\frac{in}{2}}\left(1+\frac{bx}{a+i}\right)^{-\frac{in}{2}}F_1\left(m+1;-\frac{in}{2},\frac{in}{2};m+2;-\frac{bx}{a+i},\frac{bx}{i-a}\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(n*\text{ArcTan}[a + b*x])}*x^m,x]$

[Out] $(x^{(1+m)}*(1-I*a-I*b*x)^{((I/2)*n)}*(1-(b*x)/(I-a))^{((I/2)*n)}*AppellF1[1+m,(-1/2*I)*n,(I/2)*n,2+m,-((b*x)/(I+a)),(b*x)/(I-a)]/((1+m)*(1+I*a+I*b*x)^{((I/2)*n)}*(1+(b*x)/(I+a))^{((I/2)*n)})$

Rule 138

$\text{Int}[(b_.*(x_))^{(m_)}*((c_)+(d_.*(x_))^{(n_)}*((e_)+(f_.*(x_))^{(p_)}),x_Symbol] :> \text{Simp}[c^n*e^p*((b*x)^{(m+1)}/(b*(m+1)))*AppellF1[m+1,-n,-p,m+2,(-d)*(x/c),(-f)*(x/e)],x] /; \text{FreeQ}\{b,c,d,e,f,m,n,p\},x\} \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c,0] \& \& (\text{IntegerQ}[p] \mid \mid \text{GtQ}[e,0])$

Rule 140

$\text{Int}[(b_.*(x_))^{(m_)}*((c_)+(d_.*(x_))^{(n_)}*((e_)+(f_.*(x_))^{(p_)}),x_Symbol] :> \text{Dist}[c^{\text{IntPart}[n]}*((c+d*x)^{\text{FracPart}[n]}/(1+d*(x/c))^{\text{FracPart}[n]}],\text{Int}[(b*x)^m*(1+d*(x/c))^n*(e+f*x)^p,x],x] /; \text{FreeQ}\{b,c,d,e,f,m,n,p\},x\} \& \& \text{IntegerQ}[m] \& \& \text{IntegerQ}[n] \& \& \text{GtQ}[c,0]$

Rule 5203

$\text{Int}[E^{(\text{ArcTan}[(c_)*((a_)+(b_.*(x_))])*(n_))}*((d_)+(e_.*(x_))^{(m_)}),x_Symbol] :> \text{Int}[(d+e*x)^m*((1-I*a*c-I*b*c*x)^{(I*(n/2))}/(1+I*a*c+I*b*c*x)^{(I*(n/2)}),x] /; \text{FreeQ}\{a,b,c,d,e,m,n\},x]$

Rubi steps

$$\begin{aligned}
\int e^{n \tan^{-1}(a+bx)} x^m dx &= \int x^m (1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} dx \\
&= \left((1 - ia - ibx)^{\frac{in}{2}} \left(1 - \frac{ibx}{1 - ia} \right)^{-\frac{in}{2}} \right) \int x^m (1 + ia + ibx)^{-\frac{in}{2}} \left(1 - \frac{ibx}{1 - ia} \right)^{\frac{in}{2}} dx \\
&= \left((1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} \left(1 - \frac{ibx}{1 - ia} \right)^{-\frac{in}{2}} \left(1 + \frac{ibx}{1 + ia} \right)^{\frac{in}{2}} \right) \int x^m \left(1 - \frac{ibx}{1 + ia} \right)^{\frac{in}{2}} dx \\
&= \frac{x^{1+m} (1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} \left(1 - \frac{bx}{i-a} \right)^{\frac{in}{2}} \left(1 + \frac{bx}{i+a} \right)^{-\frac{in}{2}} F_1 \left(1 + m; -\frac{in}{2}, \frac{in}{2}; 2 \right)}{1 + m}
\end{aligned}$$

Mathematica [F]

time = 0.68, size = 0, normalized size = 0.00

$$\int e^{n \text{ArcTan}(a+bx)} x^m dx$$

Verification is not applicable to the result.

[In] Integrate[E^(n*ArcTan[a + b*x])*x^m,x]

[Out] Integrate[E^(n*ArcTan[a + b*x])*x^m, x]

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{n \arctan(bx+a)} x^m dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(b*x+a))*x^m,x)

[Out] int(exp(n*arctan(b*x+a))*x^m,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))*x^m,x, algorithm="maxima")

[Out] integrate(x^m*e^(n*arctan(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(b*x+a))*x^m,x, algorithm="fricas")``[Out] integral(x^m*e^(n*arctan(b*x + a)), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int x^m e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*atan(b*x+a))*x**m,x)``[Out] Integral(x**m*exp(n*atan(a + b*x)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(b*x+a))*x^m,x, algorithm="giac")``[Out] sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int x^m e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(x^m*exp(n*atan(a + b*x)),x)``[Out] int(x^m*exp(n*atan(a + b*x)), x)`

3.237 $\int e^{n \operatorname{ArcTan}(a+bx)} x^3 dx$

Optimal. Leaf size=260

$$\frac{x^2(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}}{4b^2} - \frac{(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}(6-18a^2-10an-n^2+2b(6a+n)x)}{24b^4}$$

[Out] $1/4*x^2*(1-I*a-I*b*x)^(1+1/2*I*n)*(1+I*a+I*b*x)^(1-1/2*I*n)/b^2-1/24*(1-I*a-I*b*x)^(1+1/2*I*n)*(1+I*a+I*b*x)^(1-1/2*I*n)*(6-18*a^2-10*a*n-n^2+2*b*(6*a+n)*x)/b^4+1/3*2^(-2-1/2*I*n)*(24*a^3+36*a^2*n-12*a*(-n^2+2)-n*(-n^2+8))*(1-I*a-I*b*x)^(1+1/2*I*n)*hypergeom([1/2*I*n, 1+1/2*I*n],[2+1/2*I*n],1/2-1/2*I*a-1/2*I*b*x)/b^4/(2*I-n)$

Rubi [A]

time = 0.14, antiderivative size = 260, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5203, 102, 152, 71}

$$\frac{(-ia-ibx+1)^{1+\frac{in}{2}}(-18a^2+2bx(6a+n)-10an-n^2+6)(ia+ibx+1)^{1-\frac{in}{2}} + 2^{-2-\frac{in}{2}}(24a^3+36a^2n-12a(2-n^2)-n(8-n^2))(-ia-ibx+1)^{1+\frac{in}{2}} {}_2F_1\left(\frac{in}{2}+1, \frac{in}{2}; \frac{in}{2}+2; \frac{1}{2}(-ia-ibx+1)\right) + x^2(-ia-ibx+1)^{1+\frac{in}{2}}(ia+ibx+1)^{1-\frac{in}{2}}}{24b^4 + 3b^4(-n+2i)}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a + b*x])*x^3,x]

[Out] $(x^2*(1-I*a-I*b*x)^(1+(I/2)*n)*(1+I*a+I*b*x)^(1-(I/2)*n))/(4*b^2) - ((1-I*a-I*b*x)^(1+(I/2)*n)*(1+I*a+I*b*x)^(1-(I/2)*n)*(6-18*a^2-10*a*n-n^2+2*b*(6*a+n)*x))/(24*b^4) + (2^(-2-(I/2)*n)*(24*a^3+36*a^2*n-12*a*(2-n^2)-n*(8-n^2))*(1-I*a-I*b*x)^(1+(I/2)*n)*Hypergeometric2F1[1+(I/2)*n,(I/2)*n,2+(I/2)*n,(1-I*a-I*b*x)/2])/(3*b^4*(2*I-n))$

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 102

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)^(m - 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(m + n + p + 1))), x] + Dist[1/(d*f*(m + n + p + 1)), Int[(a + b*x)^(m - 2)*(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(m + n + p + 1) - b*(b*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1)) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}

`}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]`

Rule 152

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))
*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(-a*d*f*h*(n + 2) + b*c*f*h*(m
+ 2) - b*d*(f*g + e*h)*(m + n + 3) - b*d*f*h*(m + n + 2)*x)*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d^2*(m + n + 2)*(m + n + 3))), x] + Dist[(a^2*d
^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m
+ n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n +
3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d^2*(m + n + 2)*(m + n + 3)),
Int[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}
, x] && NeQ[m + n + 2, 0] && NeQ[m + n + 3, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_.) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(a+bx)} x^3 dx &= \int x^3 (1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} dx \\ &= \frac{x^2(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{4b^2} + \frac{\int x(1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} (-2 - ibx) dx}{4b^2} \\ &= \frac{x^2(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{4b^2} - \frac{(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}} (6 - 2ibx)}{24b^4} \\ &= \frac{x^2(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{4b^2} - \frac{(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}} (6 - 2ibx)}{24b^4} \end{aligned}$$

Mathematica [A]

time = 1.14, size = 243, normalized size = 0.93

$$\frac{e^{n \operatorname{ArcTan}(a+bx)} \left((24a^3 + 36a^2n + n(-8 + n^2) + 12a(-2 + n^2))(a+bx) - (-12 + 36a^2 + 12an + n^2)(1 + a^2 + 2abx + b^2x^2) + 2(12a + n)(a+bx)(1 + (a+bx)^2) - 6(1 + (a+bx)^2)^2 + \frac{(24a^3 + 36a^2n + n(-8 + n^2) + 12a(-2 + n^2))(-e^{n \operatorname{ArcTan}(a+bx)} \operatorname{Erfi}\left(\frac{1 - \frac{a+bx}{1 + (a+bx)^2}}{1 + (a+bx)^2}\right) + (24a^3 + 36a^2n + n(-8 + n^2) + 12a(-2 + n^2))e^{n \operatorname{ArcTan}(a+bx)} \operatorname{Erfi}\left(\frac{1 - \frac{a+bx}{1 + (a+bx)^2}}{1 + (a+bx)^2}\right))}{-24b^4} \right)}{24b^4}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(n*ArcTan[a + b*x])*x^3,x]`

`[Out] -1/24*(E^(n*ArcTan[a + b*x]))*((24*a^3 + 36*a^2*n + n*(-8 + n^2) + 12*a*(-2 + n^2))*(a + b*x) - (-12 + 36*a^2 + 12*a*n + n^2)*(1 + a^2 + 2*a*b*x + b^2*`

$$x^2) + 2*(12*a + n)*(a + b*x)*(1 + (a + b*x)^2) - 6*(1 + (a + b*x)^2)^2 + (24*a^3 + 36*a^2*n + n*(-8 + n^2) + 12*a*(-2 + n^2))*(-E^((2*I)*ArcTan[a + b*x])*n*Hypergeometric2F1[1, 1 - (I/2)*n, 2 - (I/2)*n, -E^((2*I)*ArcTan[a + b*x])]) + (2*I + n)*Hypergeometric2F1[1, (-1/2*I)*n, 1 - (I/2)*n, -E^((2*I)*ArcTan[a + b*x])]))/(-2 + I*n))/b^4$$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(bx+a)} x^3 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(b*x+a))*x^3,x)

[Out] int(exp(n*arctan(b*x+a))*x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))*x^3,x, algorithm="maxima")

[Out] integrate(x^3*e^(n*arctan(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(b*x+a))*x**3,x, algorithm="fricas")

[Out] integral(x**3*e^(n*atan(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^3 e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(b*x+a))*x**3,x)

[Out] Integral(x**3*exp(n*atan(a + b*x)), x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(b*x+a))*x^3,x, algorithm="giac")`

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^3 e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^3*exp(n*atan(a + b*x)),x)`

[Out] `int(x^3*exp(n*atan(a + b*x)), x)`

3.238 $\int e^{n \operatorname{ArcTan}(a+bx)} x^2 dx$

Optimal. Leaf size=220

$$\frac{(4a+n)(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}}{6b^3} + \frac{x(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}}{3b^2} + \frac{2^{-\frac{in}{2}}(2-6a^2-6an)}{3b^2}$$

[Out] $-1/6*(4*a+n)*(1-I*a-I*b*x)^{(1+1/2*I*n)}*(1+I*a+I*b*x)^{(1-1/2*I*n)}/b^3+1/3*x*(1-I*a-I*b*x)^{(1+1/2*I*n)}*(1+I*a+I*b*x)^{(1-1/2*I*n)}/b^2+1/3*(-6*a^2-6*a*n-n^2+2)*(1-I*a-I*b*x)^{(1+1/2*I*n)}*\operatorname{hypergeom}([1/2*I*n, 1+1/2*I*n], [2+1/2*I*n], 1/2-1/2*I*a-1/2*I*b*x)/(2^{(1/2*I*n)})/b^3/(2*I-n)$

Rubi [A]

time = 0.11, antiderivative size = 220, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.286$, Rules used = {5203, 92, 81, 71}

$$\frac{2^{-\frac{in}{2}}(-6a^2-6an-n^2+2)(-ia-ibx+1)^{1+\frac{in}{2}}{}_2F_1\left(\frac{in}{2}+1, \frac{in}{2}; \frac{in}{2}+2; \frac{1}{2}(-ia-ibx+1)\right)}{3b^3(-n+2i)} - \frac{(4a+n)(-ia-ibx+1)^{1+\frac{in}{2}}(ia+ibx+1)^{1-\frac{in}{2}}}{6b^3} + \frac{x(-ia-ibx+1)^{1+\frac{in}{2}}(ia+ibx+1)^{1-\frac{in}{2}}}{3b^2}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a + b*x])}*x^2, x]$

[Out] $-1/6*((4*a + n)*(1 - I*a - I*b*x)^{(1 + (I/2)*n)}*(1 + I*a + I*b*x)^{(1 - (I/2)*n)})/b^3 + (x*(1 - I*a - I*b*x)^{(1 + (I/2)*n)}*(1 + I*a + I*b*x)^{(1 - (I/2)*n)})/(3*b^2) + ((2 - 6*a^2 - 6*a*n - n^2)*(1 - I*a - I*b*x)^{(1 + (I/2)*n)}*Hypergeometric2F1[1 + (I/2)*n, (I/2)*n, 2 + (I/2)*n, (1 - I*a - I*b*x)/2])/3*2^{((I/2)*n)}*b^3*(2*I - n)$

Rule 71

$\operatorname{Int}[(a_ + (b_)*(x_))^{(m_)}*((c_ + (d_)*(x_))^{(n_)}), x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 81

$\operatorname{Int}[(a_ + (b_)*(x_))*((c_ + (d_)*(x_))^{(n_)}*((e_ + (f_)*(x_))^{(p_)}), x_Symbol] \rightarrow \operatorname{Simp}[b*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 2))), x] + \operatorname{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(d*f*(n + p + 2)), \operatorname{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /;$ FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 92


```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 5203

```
Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.))*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] := Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]
```

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(a+bx)} x^2 dx &= \int x^2 (1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} dx \\ &= \frac{x(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{3b^2} + \frac{\int (1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} (-1 - ibx) dx}{3b^2} \\ &= -\frac{(4a + n)(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{6b^3} + \frac{x(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{3b^2} \\ &= -\frac{(4a + n)(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{6b^3} + \frac{x(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{3b^2} \end{aligned}$$

Mathematica [A]

time = 0.12, size = 160, normalized size = 0.73

$$\frac{(-i(i + a + bx))^{1+\frac{in}{2}} \left(-((4a + n)(1 + ia + ibx)^{1-\frac{in}{2}}) + 2bx(1 + ia + ibx)^{1-\frac{in}{2}} + \frac{2^{1-\frac{in}{2}}(-2+6a^2+6an+n^2) {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; 2+\frac{in}{2}; -\frac{1}{2}i(i+a+bx)\right)}{-2i+n} \right)}{6b^3}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(n*ArcTan[a + b*x])*x^2, x]
```

```
[Out] (((-I)*(I + a + b*x))^(1 + (I/2)*n))*(-((4*a + n)*(1 + I*a + I*b*x)^(1 - (I/2)*n)) + 2*b*x*(1 + I*a + I*b*x)^(1 - (I/2)*n) + (2^(1 - (I/2)*n)*(-2 + 6*a^2 + 6*a*n + n^2)*Hypergeometric2F1[1 + (I/2)*n, (I/2)*n, 2 + (I/2)*n, (-1/2*I)*(I + a + b*x)]/(-2*I + n)))/(6*b^3)
```

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{n \arctan(bx+a)} x^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*arctan(b*x+a))*x^2,x)`

[Out] `int(exp(n*arctan(b*x+a))*x^2,x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(b*x+a))*x^2,x, algorithm="maxima")`

[Out] `integrate(x^2*e^(n*arctan(b*x + a)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(b*x+a))*x^2,x, algorithm="fricas")`

[Out] `integral(x^2*e^(n*arctan(b*x + a)), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(b*x+a))*x**2,x)`

[Out] `Integral(x**2*exp(n*atan(a + b*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(b*x+a))*x^2,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(x^2*exp(n*atan(a + b*x)),x)`

[Out] `int(x^2*exp(n*atan(a + b*x)), x)`

3.239 $\int e^{n \operatorname{ArcTan}(a+bx)} x dx$

Optimal. Leaf size=147

$$\frac{(1 - ia - ibx)^{1+\frac{in}{2}}(1 + ia + ibx)^{1-\frac{in}{2}}}{2b^2} + \frac{2^{-\frac{in}{2}}(2a + n)(1 - ia - ibx)^{1+\frac{in}{2}} {}_2F_1\left(1 + \frac{in}{2}, \frac{in}{2}; 2 + \frac{in}{2}; \frac{1}{2}(1 - ia - ibx)\right)}{b^2(2i - n)}$$

[Out] $1/2*(1-I*a-I*b*x)^{(1+1/2*I*n)}*(1+I*a+I*b*x)^{(1-1/2*I*n)}/b^2+(2*a+n)*(1-I*a-I*b*x)^{(1+1/2*I*n)}*hypergeom([1/2*I*n, 1+1/2*I*n], [2+1/2*I*n], 1/2-1/2*I*a-1/2*I*b*x)/(2^{(1/2*I*n)})/b^2/(2*I-n)$

Rubi [A]

time = 0.04, antiderivative size = 147, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 12, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5203, 81, 71}

$$\frac{2^{-\frac{in}{2}}(2a + n)(-ia - ibx + 1)^{1+\frac{in}{2}} {}_2F_1\left(\frac{in}{2} + 1, \frac{in}{2}; \frac{in}{2} + 2; \frac{1}{2}(-ia - ibx + 1)\right)}{b^2(-n + 2i)} + \frac{(-ia - ibx + 1)^{1+\frac{in}{2}}(ia + ibx + 1)^{1-\frac{in}{2}}}{2b^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(n \operatorname{ArcTan}[a + b*x])} * x, x]$

[Out] $((1 - I*a - I*b*x)^{(1 + (I/2)*n)}*(1 + I*a + I*b*x)^{(1 - (I/2)*n)})/(2*b^2) + ((2*a + n)*(1 - I*a - I*b*x)^{(1 + (I/2)*n)}*Hypergeometric2F1[1 + (I/2)*n, (I/2)*n, 2 + (I/2)*n, (1 - I*a - I*b*x)/2])/(2^{((I/2)*n)}*b^2*(2*I - n))$

Rule 71

$\text{Int}[(a + b*x)^{(m+1)}/(b*(m+1)*(b*(b*c - a*d))^n) * \text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !RationalQ[n] && GtQ[-d/(b*c - a*d), 0])

Rule 81

$\text{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n+1)}*(e + f*x)^{(p+1)}/(d*f*(n+p+2)), x] + \text{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))]/(d*f*(n+p+2)), \text{Int}[(c + d*x)^n*(e + f*x)^p, x] /;$ FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0]

Rule 5203

$\text{Int}[E^{(\operatorname{ArcTan}[(c + d*x)/(a + b*x)])} * (d + e*x)^m * ((1 - I*a*c - I*b*c*x)^{(I*(n/2)})/(1 + I*a*c + I*b*c*x))^{(n/2)}, x]$

$I*b*c*x)^{(I*(n/2))}, x] /; \text{FreeQ}[\{a, b, c, d, e, m, n\}, x]$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(a+bx)} x dx &= \int x(1-ia-ibx)^{\frac{in}{2}}(1+ia+ibx)^{-\frac{in}{2}} dx \\ &= \frac{(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}}{2b^2} - \frac{(2a+n) \int (1-ia-ibx)^{\frac{in}{2}}(1+ia+ibx)^{-\frac{in}{2}}}{2b} \\ &= \frac{(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}}{2b^2} + \frac{2^{-\frac{in}{2}}(2a+n)(1-ia-ibx)^{1+\frac{in}{2}} {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; -2-in, \frac{in}{2}\right)}{b^2(2i-n)} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 128, normalized size = 0.87

$$\frac{i(-i(i+a+bx))^{1+\frac{in}{2}} \left((1+ia+ibx)^{-\frac{in}{2}}(-i+a+bx) + \frac{2^{1-\frac{in}{2}}(2a+n) {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; 2+\frac{in}{2}; -\frac{1}{2}i(i+a+bx)\right)}{-2-in} \right)}{2b^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a + b*x])*x,x]

[Out] $((I/2)*((-I)*(I+a+b*x))^{(1+(I/2)*n)}*((-I+a+b*x)/(1+I*a+I*b*x))^{((I/2)*n)} + (2^{(1-(I/2)*n)}*(2*a+n)*\text{Hypergeometric2F1}[1+(I/2)*n, (I/2)*n, 2+(I/2)*n, (-1/2*I)*(I+a+b*x)])/(-2-I*n)))/b^2$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(bx+a)} x dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(b*x+a))*x,x)

[Out] int(exp(n*arctan(b*x+a))*x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))*x,x, algorithm="maxima")

[Out] integrate(x*e^(n*arctan(b*x + a)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))*x,x, algorithm="fricas")

[Out] integral(x*e^(n*arctan(b*x + a)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(b*x+a))*x,x)

[Out] Integral(x*exp(n*atan(a + b*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))*x,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int x e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x*exp(n*atan(a + b*x)),x)

[Out] int(x*exp(n*atan(a + b*x)), x)

3.240 $\int e^{n \operatorname{ArcTan}(a+bx)} dx$

Optimal. Leaf size=91

$$-\frac{2^{1-\frac{in}{2}}(1-ia-ibx)^{1+\frac{in}{2}} {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; 2+\frac{in}{2}; \frac{1}{2}(1-ia-ibx)\right)}{b(2i-n)}$$

[Out] $-2^{(1-1/2*I*n)*(1-I*a-I*b*x)^{(1+1/2*I*n)}*\operatorname{hypergeom}([1/2*I*n, 1+1/2*I*n], [2+1/2*I*n], 1/2-1/2*I*a-1/2*I*b*x)/b/(2*I-n)$

Rubi [A]

time = 0.01, antiderivative size = 91, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 10, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5201, 71}

$$-\frac{2^{1-\frac{in}{2}}(-ia-ibx+1)^{1+\frac{in}{2}} {}_2F_1\left(\frac{in}{2}+1, \frac{in}{2}; \frac{in}{2}+2; \frac{1}{2}(-ia-ibx+1)\right)}{b(-n+2i)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a + b*x])}, x]$

[Out] $-((2^{(1-(I/2)*n)}*(1-I*a-I*b*x)^{(1+(I/2)*n)}*\operatorname{Hypergeometric2F1}[1+(I/2)*n, (I/2)*n, 2+(I/2)*n, (1-I*a-I*b*x)/2])/(b*(2*I-n))$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}/(b*(m+1)*(b/(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x$
 $\&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] \mid\mid !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5201

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(c_+)*((a_+) + (b_+)*(x_+))])^{(n_+)}, x_Symbol] \rightarrow \operatorname{Int}[(1 - I*a*c - I*b*c*x)^{(I*(n/2))}/(1 + I*a*c + I*b*c*x)^{(I*(n/2))}, x] /;$ $\operatorname{FreeQ}\{a, b, c, n\}, x]$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(a+bx)} dx &= \int (1-ia-ibx)^{\frac{in}{2}} (1+ia+ibx)^{-\frac{in}{2}} dx \\ &= -\frac{2^{1-\frac{in}{2}}(1-ia-ibx)^{1+\frac{in}{2}} {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; 2+\frac{in}{2}; \frac{1}{2}(1-ia-ibx)\right)}{b(2i-n)} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 60, normalized size = 0.66

$$\frac{4e^{(2i+n)\text{ArcTan}(a+bx)} {}_2F_1\left(2, 1 - \frac{in}{2}; 2 - \frac{in}{2}; -e^{2i\text{ArcTan}(a+bx)}\right)}{b(2i+n)}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(n*ArcTan[a + b*x]), x]``[Out] (4*E^((2*I + n)*ArcTan[a + b*x])*Hypergeometric2F1[2, 1 - (I/2)*n, 2 - (I/2)*n, -E^((2*I)*ArcTan[a + b*x])])/(b*(2*I + n))`**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{n \arctan(bx+a)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(b*x+a)), x)``[Out] int(exp(n*arctan(b*x+a)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(b*x+a)), x, algorithm="maxima")``[Out] integrate(e^(n*arctan(b*x + a)), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(b*x+a)), x, algorithm="fricas")``[Out] integral(e^(n*arctan(b*x + a)), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(b*x+a)),x)`

[Out] `Integral(exp(n*atan(a + b*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(b*x+a)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(a+bx)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*atan(a + b*x)),x)`

[Out] `int(exp(n*atan(a + b*x)), x)`

3.241 $\int \frac{e^{n \operatorname{ArcTan}(a+bx)}}{x} dx$

Optimal. Leaf size=191

$$\frac{2i(1-ia-ibx)^{\frac{in}{2}}(1+ia+ibx)^{-\frac{in}{2}} {}_2F_1\left(1, \frac{in}{2}; 1 + \frac{in}{2}; \frac{(i-a)(1-ia-ibx)}{(i+a)(1+ia+ibx)}\right)}{n} - \frac{i2^{1-\frac{in}{2}}(1-ia-ibx)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; 1 + \frac{in}{2}; \frac{(i-a)(1-ia-ibx)}{(i+a)(1+ia+ibx)}\right)}{n}$$

[Out] 2*I*(1-I*a-I*b*x)^(1/2*I*n)*hypergeom([1, 1/2*I*n], [1+1/2*I*n], (I-a)*(1-I*a-I*b*x)/(I+a)/(1+I*a+I*b*x))/n/((1+I*a+I*b*x)^(1/2*I*n))-I*2^(1-1/2*I*n)*(1-I*a-I*b*x)^(1/2*I*n)*hypergeom([1/2*I*n, 1/2*I*n], [1+1/2*I*n], 1/2-1/2*I*a-1/2*I*b*x)/n

Rubi [A]

time = 0.07, antiderivative size = 191, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.357$, Rules used = {5203, 132, 71, 12, 133}

$$\frac{2i(-ia-ibx+1)^{\frac{in}{2}}(ia+ibx+1)^{-\frac{in}{2}} {}_2F_1\left(1, \frac{in}{2}; \frac{in}{2} + 1; \frac{(i-a)(-ia-ibx+1)}{(a+i)(ia+ibx+1)}\right)}{n} - \frac{i2^{1-\frac{in}{2}}(-ia-ibx+1)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; \frac{in}{2} + 1; \frac{1}{2}(-ia-ibx+1)\right)}{n}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a + b*x])/x,x]

[Out] ((2*I)*(1 - I*a - I*b*x)^((I/2)*n)*Hypergeometric2F1[1, (I/2)*n, 1 + (I/2)*n, ((I - a)*(1 - I*a - I*b*x))/((I + a)*(1 + I*a + I*b*x))])/n*(1 + I*a + I*b*x)^((I/2)*n) - (I*2^(1 - (I/2)*n)*(1 - I*a - I*b*x)^((I/2)*n)*Hypergeometric2F1[(I/2)*n, (I/2)*n, 1 + (I/2)*n, (1 - I*a - I*b*x)/2])/n

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_) /; FreeQ[b, x]]

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 132

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Dist[b*d^(m + n)*f^p, Int[(a + b*x)^(m - 1)/(c + d*x)^m, x], x] + Int[(a + b*x)^(m - 1)*((e + f*x)^p/(c + d*x)^m)*ExpandToSum[(

$a + b*x)*(c + d*x)^{-p - 1} - (b*d^{-p - 1}*f^p)/(e + f*x)^p, x]$, x] /; FreeQ[{a, b, c, d, e, f, m, n}, x] && EqQ[m + n + p + 1, 0] && ILtQ[p, 0] && (GtQ[m, 0] || SumSimplerQ[m, -1] || !(GtQ[n, 0] || SumSimplerQ[n, -1]))

Rule 133

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/((m + 1)*(b*e - a*f)^(n + 1)*(e + f*x)^(m + 1)))*Hypergeometric2F1[m + 1, -n, m + 2, -(d*e - c*f)/(a + b*x)/((b*c - a*d)*(e + f*x))], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))])*(n_.)*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2))/(1 + I*a*c + I*b*c*x)^(I*(n/2))), x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(a+bx)}}{x} dx &= \int \frac{(1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}}}{x} dx \\ &= - \left((-1 + ia) \int \frac{(1 - ia - ibx)^{-1 + \frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}}}{x} dx \right) - (ib) \int (1 - ia - ibx)^{-1} dx \\ &= \frac{2i(1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}} {}_2F_1\left(1, \frac{in}{2}; 1 + \frac{in}{2}; \frac{(i-a)(1-ia-ibx)}{(i+a)(1+ia+ibx)}\right)}{n} - \frac{i2^{1-\frac{in}{2}}(1 - ia)}{n} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 170, normalized size = 0.89

$$\frac{2i(1 + ia + ibx)^{-\frac{in}{2}} (-i(i + a + bx))^{\frac{in}{2}} \left({}_2F_1\left(1, \frac{in}{2}; 1 + \frac{in}{2}; \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right) - 2^{-\frac{in}{2}} (1 + ia + ibx)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; 1 + \frac{in}{2}; -\frac{1}{2}i(i + a + bx)\right) \right)}{n}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a + b*x])/x,x]

[Out] ((2*I)*((-I)*(I + a + b*x))^((I/2)*n)*Hypergeometric2F1[1, (I/2)*n, 1 + (I/2)*n, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)] - ((1 + I*a + I*b*x)^((I/2)*n)*Hypergeometric2F1[(I/2)*n, (I/2)*n, 1 + (I/2)*n, (-1/2*I)*(I + a + b*x)]/2^((I/2)*n))/(n*(1 + I*a + I*b*x)^((I/2)*n))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(bx+a)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(b*x+a))/x,x)

[Out] int(exp(n*arctan(b*x+a))/x,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x,x, algorithm="maxima")

[Out] integrate(e^(n*arctan(b*x + a))/x, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x,x, algorithm="fricas")

[Out] integral(e^(n*arctan(b*x + a))/x, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(a+bx)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(b*x+a))/x,x)

[Out] Integral(exp(n*atan(a + b*x))/x, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*arctan(b*x+a))/x,x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(a+bx)}}{x} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(n*atan(a + b*x))/x,x)
```

```
[Out] int(exp(n*atan(a + b*x))/x, x)
```

3.242 $\int \frac{e^{n \operatorname{ArcTan}(a+bx)}}{x^2} dx$

Optimal. Leaf size=128

$$\frac{4b(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{-1-\frac{in}{2}} {}_2F_1\left(2, 1+\frac{in}{2}; 2+\frac{in}{2}, \frac{(i-a)(1-ia-ibx)}{(i+a)(1+ia+ibx)}\right)}{(i+a)^2(2i-n)}$$

[Out] -4*b*(1-I*a-I*b*x)^(1+1/2*I*n)*(1+I*a+I*b*x)^(-1-1/2*I*n)*hypergeom([2, 1+1/2*I*n], [2+1/2*I*n], (I-a)*(1-I*a-I*b*x)/(I+a)/(1+I*a+I*b*x))/(I+a)^2/(2*I-n)

Rubi [A]

time = 0.03, antiderivative size = 128, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5203, 133}

$$\frac{4b(-ia-ibx+1)^{1+\frac{in}{2}}(ia+ibx+1)^{-1-\frac{in}{2}} {}_2F_1\left(2, \frac{in}{2}+1; \frac{in}{2}+2; \frac{(i-a)(-ia-ibx+1)}{(a+i)(ia+ibx+1)}\right)}{(a+i)^2(-n+2i)}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a + b*x])/x^2,x]

[Out] (-4*b*(1 - I*a - I*b*x)^(1 + (I/2)*n)*(1 + I*a + I*b*x)^(-1 - (I/2)*n)*Hypergeometric2F1[2, 1 + (I/2)*n, 2 + (I/2)*n, ((I - a)*(1 - I*a - I*b*x))/((I + a)*(1 + I*a + I*b*x))]/((I + a)^2*(2*I - n))

Rule 133

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/((m + 1)*(b*e - a*f)^(n + 1)*(e + f*x)^(m + 1)))*Hypergeometric2F1[m + 1, -n, m + 2, (-(d*e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.), x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c + I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\int \frac{e^{n \tan^{-1}(a+bx)}}{x^2} dx = \int \frac{(1-ia-ibx)^{\frac{in}{2}}(1+ia+ibx)^{-\frac{in}{2}}}{x^2} dx$$

$$= -\frac{4b(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{-1-\frac{in}{2}} {}_2F_1\left(2, 1+\frac{in}{2}; 2+\frac{in}{2}, \frac{(i-a)(1-ia-ibx)}{(i+a)(1+ia+ibx)}\right)}{(i+a)^2(2i-n)}$$

Mathematica [A]

time = 0.02, size = 125, normalized size = 0.98

$$\frac{4ib(1+ia+ibx)^{-\frac{in}{2}}(-i(i+a+bx))^{1+\frac{in}{2}} {}_2F_1\left(2, 1+\frac{in}{2}; 2+\frac{in}{2}, \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right)}{(i+a)^2(-2i+n)(-i+a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a + b*x])/x^2,x]

[Out] ((-4*I)*b*((-I)*(I + a + b*x))^(1 + (I/2)*n)*Hypergeometric2F1[2, 1 + (I/2)*n, 2 + (I/2)*n, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)]/((I + a)^2*(-2*I + n)*(1 + I*a + I*b*x)^((I/2)*n)*(-I + a + b*x))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(bx+a)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(b*x+a))/x^2,x)**[Out]** int(exp(n*arctan(b*x+a))/x^2,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x^2,x, algorithm="maxima")**[Out]** integrate(e^(n*arctan(b*x + a))/x^2, x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x^2,x, algorithm="fricas")

[Out] integral(e^(n*arctan(b*x + a))/x^2, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(a+bx)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(b*x+a))/x**2,x)

[Out] Integral(exp(n*atan(a + b*x))/x**2, x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x^2,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(a+bx)}}{x^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a + b*x))/x^2,x)

[Out] int(exp(n*atan(a + b*x))/x^2, x)

3.243 $\int \frac{e^{n \operatorname{ArcTan}(a+bx)}}{x^3} dx$

Optimal. Leaf size=207

$$\frac{(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{1-\frac{in}{2}}}{2(1+a^2)x^2} - \frac{2b^2(2a-n)(1-ia-ibx)^{1+\frac{in}{2}}(1+ia+ibx)^{-1-\frac{in}{2}} {}_2F_1\left(2, 1+\frac{in}{2}; 2, (i-a)(i+a)^3(2i-n)\right)}{(i-a)(i+a)^3(2i-n)}$$

[Out] $-1/2*(1-I*a-I*b*x)^{(1+1/2*I*n)}*(1+I*a+I*b*x)^{(1-1/2*I*n)}/(a^2+1)/x^2-2*b^2*(2*a-n)*(1-I*a-I*b*x)^{(1+1/2*I*n)}*(1+I*a+I*b*x)^{(-1-1/2*I*n)}*\operatorname{hypergeom}([2, 1+1/2*I*n], [2+1/2*I*n], (I-a)*(1-I*a-I*b*x)/(I+a)/(1+I*a+I*b*x))/(I-a)/(I+a)^3/(2*I-n)$

Rubi [A]

time = 0.09, antiderivative size = 207, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 14, $\frac{\text{number of rules}}{\text{integrand size}} = 0.214$, Rules used = {5203, 98, 133}

$$\frac{(-ia-ibx+1)^{1+\frac{in}{2}}(ia+ibx+1)^{1-\frac{in}{2}}}{2(a^2+1)x^2} - \frac{2b^2(2a-n)(-ia-ibx+1)^{1+\frac{in}{2}}(ia+ibx+1)^{-1-\frac{in}{2}} {}_2F_1\left(2, \frac{in}{2}+1, \frac{in}{2}+2; \frac{(i-a)(-ia-ibx+1)}{(a+i)(ia+ibx+1)}\right)}{(-a+i)(a+i)^3(-n+2i)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n \operatorname{ArcTan}[a + b*x])}/x^3, x]$

[Out] $-1/2*((1-I*a-I*b*x)^{(1+(I/2)*n)}*(1+I*a+I*b*x)^{(1-(I/2)*n)})/((1+a^2)*x^2)-(2*b^2*(2*a-n)*(1-I*a-I*b*x)^{(1+(I/2)*n)}*(1+I*a+I*b*x)^{(-1-(I/2)*n)}*\operatorname{Hypergeometric2F1}[2, 1+(I/2)*n, 2+(I/2)*n, ((I-a)*(1-I*a-I*b*x))/((I+a)*(1+I*a+I*b*x))]/((I-a)*(I+a)^3*(2*I-n))$

Rule 98

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[b*(a + b*x)^{(m+1)}*(c + d*x)^{(n+1)}*((e + f*x)^{(p+1)})/((m+1)*(b*c - a*d)*(b*e - a*f)), x] + \operatorname{Dist}[(a*d*f*(m+1) + b*c*f*(n+1) + b*d*e*(p+1))/((m+1)*(b*c - a*d)*(b*e - a*f)), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^n*(e + f*x)^p, x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& \operatorname{EqQ}[\operatorname{Simplify}[m + n + p + 3], 0] \&\& (\operatorname{LtQ}[m, -1] || \operatorname{SumSimplerQ}[m, 1])$

Rule 133

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(b*c - a*d)^n*((a + b*x)^{(m+1)})/((m+1)*(b*e - a*f)^{(n+1)}*(e + f*x)^{(m+1)})*\operatorname{Hypergeometric2F1}[m+1, -n, m+2, -(d*(e - c*f))/((a + b*x)/((b*c - a*d)*(e + f*x)))] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, p\}, x] \&\& \operatorname{EqQ}[m + n + p + 2, 0] \&\& \operatorname{ILtQ}[n, 0] \&\& (\operatorname{SumSimplerQ}[m, 1])$

|| !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

Rule 5203

Int[E^(ArcTan[(c_.)*((a_) + (b_.)*(x_))]*(n_.))*((d_.) + (e_.)*(x_))^(m_.),
x_Symbol] :> Int[(d + e*x)^m*((1 - I*a*c - I*b*c*x)^(I*(n/2)))/(1 + I*a*c +
I*b*c*x)^(I*(n/2))], x] /; FreeQ[{a, b, c, d, e, m, n}, x]

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(a+bx)}}{x^3} dx &= \int \frac{(1 - ia - ibx)^{\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}}}{x^3} dx \\ &= -\frac{(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{2(1+a^2)x^2} - \frac{(b(2a-n)) \int \frac{(1-ia-ibx)^{\frac{in}{2}} (1+ia+ibx)^{-\frac{in}{2}}}{x^2} dx}{2(1+a^2)} \\ &= -\frac{(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{1-\frac{in}{2}}}{2(1+a^2)x^2} + \frac{2b^2(2a-n)(1 - ia - ibx)^{1+\frac{in}{2}} (1 + ia + ibx)^{-\frac{in}{2}}}{(i+a)^2(1+a^2)} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 173, normalized size = 0.84

$$\frac{i(1+ia+ibx)^{-\frac{in}{2}}(-i(i+a+bx))^{1+\frac{in}{2}} \left((i+a)^2(-2i+n)(-i+a+bx)^2 + 4b^2(-2a+n)x^2 {}_2F_1\left(2, 1+\frac{in}{2}; 2+\frac{in}{2}, \frac{1+a^2-ibx+abx}{1+a^2+ibx+abx}\right) \right)}{2(-i+a)(i+a)^3(-2i+n)x^2(-i+a+bx)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a + b*x])/x^3,x]

[Out] ((-1/2*I)*((-I)*(I + a + b*x))^(1 + (I/2)*n))*((I + a)^2*(-2*I + n)*(-I + a + b*x)^2 + 4*b^2*(-2*a + n)*x^2*Hypergeometric2F1[2, 1 + (I/2)*n, 2 + (I/2)*n, (1 + a^2 - I*b*x + a*b*x)/(1 + a^2 + I*b*x + a*b*x)])/((-I + a)*(I + a)^3*(-2*I + n)*x^2*(1 + I*a + I*b*x)^((I/2)*n)*(-I + a + b*x))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(bx+a)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(b*x+a))/x^3,x)

[Out] int(exp(n*arctan(b*x+a))/x^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x^3,x, algorithm="maxima")

[Out] integrate(e^(n*arctan(b*x + a))/x^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x^3,x, algorithm="fricas")

[Out] integral(e^(n*arctan(b*x + a))/x^3, x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(a+bx)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(b*x+a))/x**3,x)

[Out] Integral(exp(n*atan(a + b*x))/x**3, x)

Giac [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(b*x+a))/x^3,x, algorithm="giac")

[Out] Timed out

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{e^{n \operatorname{atan}(a+bx)}}{x^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a + b*x))/x^3,x)

[Out] int(exp(n*atan(a + b*x))/x^3, x)

3.244 $\int e^{\text{ArcTan}(ax)} (c + a^2 cx^2)^p dx$

Optimal. Leaf size=102

$$\frac{i2^{(1-\frac{i}{2})+p}(1-iax)^{(1+\frac{i}{2})+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1(\frac{i}{2}-p, (1+\frac{i}{2})+p; (2+\frac{i}{2})+p; \frac{1}{2}(1-iax))}{a((2+i)+2p)}$$

[Out] $I*2^{(1-1/2*I+p)}*(1-I*a*x)^{(1+1/2*I+p)}*(a^2*c*x^2+c)^p*\text{hypergeom}([1/2*I-p, 1+1/2*I+p], [2+1/2*I+p], 1/2-1/2*I*a*x)/a/(2+I+2*p)/((a^2*x^2+1)^p)$

Rubi [A]

time = 0.06, antiderivative size = 102, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.158$, Rules used = {5184, 5181, 71}

$$\frac{i2^{p+(1-\frac{i}{2})}(1-iax)^{p+(1+\frac{i}{2})}(a^2x^2+1)^{-p}(a^2cx^2+c)^p {}_2F_1(\frac{i}{2}-p, p+(1+\frac{i}{2}); p+(2+\frac{i}{2}); \frac{1}{2}(1-iax))}{a(2p+(2+i))}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{\text{ArcTan}[a*x]}*(c + a^2*c*x^2)^p, x]$

[Out] $(I*2^{((1-I/2)+p)}*(1-I*a*x)^{((1+I/2)+p)}*(c+a^2*c*x^2)^p*\text{Hypergeometric2F1}[I/2-p, (1+I/2)+p, (2+I/2)+p, (1-I*a*x)/2])/a*((2+I)+2*p)*(1+a^2*x^2)^p)$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+)^2)^{(n_+)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] :> \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] :> \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int e^{\tan^{-1}(ax)}(c+a^2cx^2)^p dx &= \left((1+a^2x^2)^{-p} (c+a^2cx^2)^p \right) \int e^{\tan^{-1}(ax)}(1+a^2x^2)^p dx \\
&= \left((1+a^2x^2)^{-p} (c+a^2cx^2)^p \right) \int (1-iax)^{\frac{i}{2}+p} (1+iax)^{-\frac{i}{2}+p} dx \\
&= \frac{i2^{(1-\frac{i}{2})+p} (1-iax)^{(1+\frac{i}{2})+p} (1+a^2x^2)^{-p} (c+a^2cx^2)^p {}_2F_1\left(\frac{i}{2}-p, \left(1+\frac{i}{2}\right)+p; \left(2+\frac{i}{2}\right)+p; \frac{1}{2}(1-iax)\right)}{a((2+i)+2p)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 102, normalized size = 1.00

$$\frac{i2^{-\frac{i}{2}+p} (1-iax)^{(1+\frac{i}{2})+p} (1+a^2x^2)^{-p} (c+a^2cx^2)^p {}_2F_1\left(\frac{i}{2}-p, \left(1+\frac{i}{2}\right)+p; \left(2+\frac{i}{2}\right)+p; \frac{1}{2}(1-iax)\right)}{a\left(\left(1+\frac{i}{2}\right)+p\right)}$$

Antiderivative was successfully verified.

`[In] Integrate[E^ArcTan[a*x]*(c + a^2*c*x^2)^p,x]`

```
[Out] (I*2^(-1/2*I + p)*(1 - I*a*x)^((1 + I/2) + p)*(c + a^2*c*x^2)^p*Hypergeometric2F1[I/2 - p, (1 + I/2) + p, (2 + I/2) + p, (1 - I*a*x)/2])/(a*((1 + I/2) + p)*(1 + a^2*x^2)^p)
```

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{\arctan(ax)}(a^2cx^2+c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(arctan(a*x))*(a^2*c*x^2+c)^p,x)``[Out] int(exp(arctan(a*x))*(a^2*c*x^2+c)^p,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="maxima")``[Out] integrate((a^2*c*x^2 + c)^p*e^(arctan(a*x)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="fricas")
```

```
[Out] integral((a^2*c*x^2 + c)^p*e^(arctan(a*x)), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^p e^{\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(atan(a*x))*(a**2*c*x**2+c)**p,x)
```

```
[Out] Integral((c*(a**2*x**2 + 1))**p*exp(atan(a*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{\operatorname{atan}(ax)} (ca^2x^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(atan(a*x))*(c + a^2*c*x^2)^p,x)
```

```
[Out] int(exp(atan(a*x))*(c + a^2*c*x^2)^p, x)
```

3.245 $\int e^{\text{ArcTan}(ax)}(c + a^2cx^2)^2 dx$

Optimal. Leaf size=63

$$\frac{\left(\frac{1}{37} + \frac{6i}{37}\right) 2^{3-\frac{i}{2}} c^2 (1-iax)^{3+\frac{i}{2}} {}_2F_1\left(-2 + \frac{i}{2}, 3 + \frac{i}{2}; 4 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

[Out] $(1/37+6/37*I)*2^{(3-1/2*I)}*c^2*(1-I*a*x)^{(3+1/2*I)}*\text{hypergeom}([3+1/2*I, -2+1/2*I], [4+1/2*I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.03, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5181, 71}

$$\frac{\left(\frac{1}{37} + \frac{6i}{37}\right) 2^{3-\frac{i}{2}} c^2 (1-iax)^{3+\frac{i}{2}} {}_2F_1\left(-2 + \frac{i}{2}, 3 + \frac{i}{2}; 4 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{\text{ArcTan}[a*x]}*(c + a^2*c*x^2)^2, x]$

[Out] $((1/37 + (6*I)/37)*2^{(3 - I/2)}*c^2*(1 - I*a*x)^{(3 + I/2)}*\text{Hypergeometric2F1}[-2 + I/2, 3 + I/2, 4 + I/2, (1 - I*a*x)/2])/a$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[m] \&\& !\text{IntegerQ}[n] \&\& \text{GtQ}[b/(b*c - a*d), 0] \&\& (\text{RationalQ}[m] || !(\text{RationalQ}[n] \&\& \text{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[a_+)*(x_+)]*(n_+)}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] := \text{Dist}[c^{p_+}, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x \&\& \text{EqQ}[d, a^2*c] \&\& (\text{IntegerQ}[p] || \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{\tan^{-1}(ax)}(c + a^2cx^2)^2 dx &= c^2 \int (1-iax)^{2+\frac{i}{2}}(1+iax)^{2-\frac{i}{2}} dx \\ &= \frac{\left(\frac{1}{37} + \frac{6i}{37}\right) 2^{3-\frac{i}{2}} c^2 (1-iax)^{3+\frac{i}{2}} {}_2F_1\left(-2 + \frac{i}{2}, 3 + \frac{i}{2}; 4 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 63, normalized size = 1.00

$$\frac{\left(\frac{1}{37} + \frac{6i}{37}\right) 2^{3-\frac{i}{2}} c^2 (1 - iax)^{3+\frac{i}{2}} {}_2F_1\left(-2 + \frac{i}{2}, 3 + \frac{i}{2}; 4 + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]*(c + a^2*c*x^2)^2,x]

[Out] ((1/37 + (6*I)/37)*2^(3 - I/2)*c^2*(1 - I*a*x)^(3 + I/2)*Hypergeometric2F1[-2 + I/2, 3 + I/2, 4 + I/2, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{\arctan(ax)} (a^2 c x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))*(a^2*c*x^2+c)^2,x)

[Out] int(exp(arctan(a*x))*(a^2*c*x^2+c)^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^2*e^(arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] integral((a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2)*e^(arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c^2 \left(\int 2a^2 x^2 e^{\operatorname{atan}(ax)} dx + \int a^4 x^4 e^{\operatorname{atan}(ax)} dx + \int e^{\operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(atan(a*x))*(a**2*c*x**2+c)**2,x)
```

```
[Out] c**2*(Integral(2*a**2*x**2*exp(atan(a*x)), x) + Integral(a**4*x**4*exp(atan(a*x)), x) + Integral(exp(atan(a*x)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{\operatorname{atan}(ax)} (ca^2x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(atan(a*x))*(c + a^2*c*x^2)^2,x)
```

```
[Out] int(exp(atan(a*x))*(c + a^2*c*x^2)^2, x)
```

3.246 $\int e^{\text{ArcTan}(ax)}(c + a^2cx^2) dx$

Optimal. Leaf size=61

$$\frac{\left(\frac{1}{17} + \frac{4i}{17}\right) 2^{2-\frac{i}{2}} c (1-iax)^{2+\frac{i}{2}} {}_2F_1\left(-1 + \frac{i}{2}, 2 + \frac{i}{2}; 3 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

[Out] (1/17+4/17*I)*2^(2-1/2*I)*c*(1-I*a*x)^(2+1/2*I)*hypergeom([2+1/2*I, -1+1/2*I], [3+1/2*I], 1/2-1/2*I*a*x)/a

Rubi [A]

time = 0.02, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 17, $\frac{\text{number of rules}}{\text{integrand size}} = 0.118$, Rules used = {5181, 71}

$$\frac{\left(\frac{1}{17} + \frac{4i}{17}\right) 2^{2-\frac{i}{2}} c (1-iax)^{2+\frac{i}{2}} {}_2F_1\left(-1 + \frac{i}{2}, 2 + \frac{i}{2}; 3 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]*(c + a^2*c*x^2),x]

[Out] ((1/17 + (4*I)/17)*2^(2 - I/2)*c*(1 - I*a*x)^(2 + I/2)*Hypergeometric2F1[-1 + I/2, 2 + I/2, 3 + I/2, (1 - I*a*x)/2])/a

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{\tan^{-1}(ax)}(c + a^2cx^2) dx &= c \int (1-iax)^{1+\frac{i}{2}}(1+iax)^{1-\frac{i}{2}} dx \\ &= \frac{\left(\frac{1}{17} + \frac{4i}{17}\right) 2^{2-\frac{i}{2}} c (1-iax)^{2+\frac{i}{2}} {}_2F_1\left(-1 + \frac{i}{2}, 2 + \frac{i}{2}; 3 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 61, normalized size = 1.00

$$\frac{\left(\frac{1}{17} + \frac{4i}{17}\right) 2^{2-\frac{i}{2}} c (1 - iax)^{2+\frac{i}{2}} {}_2F_1\left(-1 + \frac{i}{2}, 2 + \frac{i}{2}; 3 + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]*(c + a^2*c*x^2), x]

[Out] ((1/17 + (4*I)/17)*2^(2 - I/2)*c*(1 - I*a*x)^(2 + I/2)*Hypergeometric2F1[-1 + I/2, 2 + I/2, 3 + I/2, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{\arctan(ax)} (a^2 c x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))*(a^2*c*x^2+c), x)

[Out] int(exp(arctan(a*x))*(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)*e^(arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c), x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)*e^(arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c \left(\int a^2 x^2 e^{\operatorname{atan}(ax)} dx + \int e^{\operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))*(a**2*c*x**2+c),x)

[Out] c*(Integral(a**2*x**2*exp(atan(a*x)), x) + Integral(exp(atan(a*x)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{\operatorname{atan}(ax)} (ca^2x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))*(c + a^2*c*x^2),x)

[Out] int(exp(atan(a*x))*(c + a^2*c*x^2), x)

3.247 $\int e^{\text{ArcTan}(ax)} dx$

Optimal. Leaf size=60

$$\frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-\frac{i}{2}} (1-iax)^{1+\frac{i}{2}} {}_2F_1\left(\frac{i}{2}, 1 + \frac{i}{2}; 2 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

[Out] (1/5+2/5*I)*2^(1-1/2*I)*(1-I*a*x)^(1+1/2*I)*hypergeom([1/2*I, 1+1/2*I], [2+1/2*I], 1/2-1/2*I*a*x)/a

Rubi [A]

time = 0.01, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 6, $\frac{\text{number of rules}}{\text{integrand size}} = 0.333$, Rules used = {5169, 71}

$$\frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-\frac{i}{2}} (1-iax)^{1+\frac{i}{2}} {}_2F_1\left(\frac{i}{2}, 1 + \frac{i}{2}; 2 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x], x]

[Out] ((1/5 + (2*I)/5)*2^(1 - I/2)*(1 - I*a*x)^(1 + I/2)*Hypergeometric2F1[I/2, 1 + I/2, 2 + I/2, (1 - I*a*x)/2])/a

Rule 71

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n)*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5169

Int[E^(ArcTan[(a_)*(x_)])*(n_.), x_Symbol] := Int[(1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{\tan^{-1}(ax)} dx &= \int (1-iax)^{\frac{i}{2}} (1+iax)^{-\frac{i}{2}} dx \\ &= \frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-\frac{i}{2}} (1-iax)^{1+\frac{i}{2}} {}_2F_1\left(\frac{i}{2}, 1 + \frac{i}{2}; 2 + \frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 45, normalized size = 0.75

$$\frac{\left(\frac{4}{5} - \frac{8i}{5}\right) e^{(1+2i)\text{ArcTan}(ax)} {}_2F_1\left(1 - \frac{i}{2}, 2; 2 - \frac{i}{2}; -e^{2i\text{ArcTan}(ax)}\right)}{a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^ArcTan[a*x], x]`

```
[Out] ((4/5 - (8*I)/5)*E^((1 + 2*I)*ArcTan[a*x])*Hypergeometric2F1[1 - I/2, 2, 2 - I/2, -E^((2*I)*ArcTan[a*x])])/a
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(arctan(a*x)), x)``[Out] int(exp(arctan(a*x)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(arctan(a*x)), x, algorithm="maxima")``[Out] integrate(e^(arctan(a*x)), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(arctan(a*x)), x, algorithm="fricas")``[Out] integral(e^(arctan(a*x)), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{\text{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(atan(a*x)),x)`

[Out] `Integral(exp(atan(a*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arctan(a*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(atan(a*x)),x)`

[Out] `int(exp(atan(a*x)), x)`

$$3.248 \quad \int \frac{e^{\text{ArcTan}(ax)}}{c+a^2cx^2} dx$$

Optimal. Leaf size=13

$$\frac{e^{\text{ArcTan}(ax)}}{ac}$$

[Out] exp(arctan(a*x))/a/c

Rubi [A]

time = 0.02, antiderivative size = 13, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.053$, Rules used = {5179}

$$\frac{e^{\text{ArcTan}(ax)}}{ac}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c + a^2*c*x^2),x]

[Out] E^ArcTan[a*x]/(a*c)

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\int \frac{e^{\tan^{-1}(ax)}}{c + a^2cx^2} dx = \frac{e^{\tan^{-1}(ax)}}{ac}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.01, size = 35, normalized size = 2.69

$$\frac{(1 - iax)^{\frac{i}{2}}(1 + iax)^{-\frac{i}{2}}}{ac}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2),x]

[Out] (1 - I*a*x)^(I/2)/(a*c*(1 + I*a*x)^(I/2))

Maple [A]

time = 0.07, size = 13, normalized size = 1.00

method	result	size
gospers	$\frac{e^{\arctan(ax)}}{ac}$	13
risch	$\frac{(-iax+1)^{\frac{i}{2}}(iax+1)^{-\frac{i}{2}}}{ac}$	28

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(arctan(a*x))/(a^2*c*x^2+c),x,method=_RETURNVERBOSE)
```

```
[Out] exp(arctan(a*x))/a/c
```

Maxima [A]

time = 0.49, size = 12, normalized size = 0.92

$$\frac{e^{\arctan(ax)}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c),x, algorithm="maxima")
```

```
[Out] e^(arctan(a*x))/(a*c)
```

Fricas [A]

time = 2.99, size = 12, normalized size = 0.92

$$\frac{e^{\arctan(ax)}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c),x, algorithm="fricas")
```

```
[Out] e^(arctan(a*x))/(a*c)
```

Sympy [A]

time = 0.51, size = 12, normalized size = 0.92

$$\begin{cases} \frac{e^{\arctan(ax)}}{ac} & \text{for } a \neq 0 \\ \frac{x}{c} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c),x)
```

```
[Out] Piecewise((exp(atan(a*x))/(a*c), Ne(a, 0)), (x/c, True))
```

Giac [A]

time = 0.43, size = 12, normalized size = 0.92

$$\frac{e^{\arctan(ax)}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c),x, algorithm="giac")

[Out] e^(arctan(a*x))/(a*c)

Mupad [B]

time = 0.53, size = 12, normalized size = 0.92

$$\frac{e^{\operatorname{atan}(ax)}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2),x)

[Out] exp(atan(a*x))/(a*c)

$$3.249 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$$

Optimal. Leaf size=50

$$\frac{2e^{\text{ArcTan}(ax)}}{5ac^2} + \frac{e^{\text{ArcTan}(ax)}(1+2ax)}{5ac^2(1+a^2x^2)}$$

[Out] $2/5*\exp(\arctan(a*x))/a/c^2+1/5*\exp(\arctan(a*x))*(2*a*x+1)/a/c^2/(a^2*x^2+1)$

Rubi [A]

time = 0.04, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5178, 5179}

$$\frac{(2ax+1)e^{\text{ArcTan}(ax)}}{5ac^2(a^2x^2+1)} + \frac{2e^{\text{ArcTan}(ax)}}{5ac^2}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c+a^2*c*x^2)^2,x]

[Out] $(2*E^{\text{ArcTan}[a*x]})/(5*a*c^2) + (E^{\text{ArcTan}[a*x]}*(1+2*a*x))/(5*a*c^2*(1+a^2*x^2))$

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.))/((c_.)+(d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n-2*a*(p+1)*x)*(c+d*x^2)^(p+1)*(E^(n*ArcTan[a*x])/(a*c*(n^2+4*(p+1)^2))), x] + Dist[2*(p+1)*((2*p+3)/(c*(n^2+4*(p+1)^2))), Int[(c+d*x^2)^(p+1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2+4*(p+1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)])*(n_.))/((c_.)+(d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned} \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx &= \frac{e^{\tan^{-1}(ax)}(1+2ax)}{5ac^2(1+a^2x^2)} + \frac{2 \int \frac{e^{\tan^{-1}(ax)}}{c+a^2cx^2} dx}{5c} \\ &= \frac{2e^{\tan^{-1}(ax)}}{5ac^2} + \frac{e^{\tan^{-1}(ax)}(1+2ax)}{5ac^2(1+a^2x^2)} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.02, size = 60, normalized size = 1.20

$$\frac{(1 - iax)^{\frac{i}{2}}(1 + iax)^{-\frac{i}{2}}(3 + 2ax + 2a^2x^2)}{5c^2(a + a^3x^2)}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2)^2,x]

[Out] ((1 - I*a*x)^(I/2)*(3 + 2*a*x + 2*a^2*x^2))/(5*c^2*(1 + I*a*x)^(I/2)*(a + a^3*x^2))

Maple [A]

time = 0.09, size = 39, normalized size = 0.78

method	result	size
gosper	$\frac{e^{\arctan(ax)}(2a^2x^2+2ax+3)}{5(a^2x^2+1)ac^2}$	39

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)

[Out] 1/5*exp(arctan(a*x))*(2*a^2*x^2+2*a*x+3)/(a^2*x^2+1)/a/c^2

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^2, x)

Fricas [A]

time = 3.35, size = 39, normalized size = 0.78

$$\frac{(2a^2x^2 + 2ax + 3)e^{\arctan(ax)}}{5(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] 1/5*(2*a^2*x^2 + 2*a*x + 3)*e^(arctan(a*x))/(a^3*c^2*x^2 + a*c^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(42) = 84$.

time = 1.43, size = 95, normalized size = 1.90

$$\begin{cases} \frac{2a^2x^2e^{\operatorname{atan}(ax)}}{5a^3c^2x^2+5ac^2} + \frac{2axe^{\operatorname{atan}(ax)}}{5a^3c^2x^2+5ac^2} + \frac{3e^{\operatorname{atan}(ax)}}{5a^3c^2x^2+5ac^2} & \text{for } a \neq 0 \\ \frac{x}{c^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**2,x)

[Out] Piecewise((2*a**2*x**2*exp(atan(a*x))/(5*a**3*c**2*x**2 + 5*a*c**2) + 2*a*x*exp(atan(a*x))/(5*a**3*c**2*x**2 + 5*a*c**2) + 3*exp(atan(a*x))/(5*a**3*c**2*x**2 + 5*a*c**2), Ne(a, 0)), (x/c**2, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.56, size = 44, normalized size = 0.88

$$\frac{e^{\operatorname{atan}(ax)} \left(\frac{3}{5a^3c^2} + \frac{2x}{5a^2c^2} + \frac{2x^2}{5ac^2} \right)}{\frac{1}{a^2} + x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^2,x)

[Out] (exp(atan(a*x))*(3/(5*a^3*c^2) + (2*x)/(5*a^2*c^2) + (2*x^2)/(5*a*c^2)))/(1/a^2 + x^2)

$$3.250 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$$

Optimal. Leaf size=83

$$\frac{24e^{\text{ArcTan}(ax)}}{85ac^3} + \frac{e^{\text{ArcTan}(ax)}(1+4ax)}{17ac^3(1+a^2x^2)^2} + \frac{12e^{\text{ArcTan}(ax)}(1+2ax)}{85ac^3(1+a^2x^2)}$$

[Out] 24/85*exp(arctan(a*x))/a/c^3+1/17*exp(arctan(a*x))*(4*a*x+1)/a/c^3/(a^2*x^2+1)^2+12/85*exp(arctan(a*x))*(2*a*x+1)/a/c^3/(a^2*x^2+1)

Rubi [A]

time = 0.06, antiderivative size = 83, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5178, 5179}

$$\frac{12(2ax+1)e^{\text{ArcTan}(ax)}}{85ac^3(a^2x^2+1)} + \frac{(4ax+1)e^{\text{ArcTan}(ax)}}{17ac^3(a^2x^2+1)^2} + \frac{24e^{\text{ArcTan}(ax)}}{85ac^3}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c+a^2*c*x^2)^3,x]

[Out] (24*E^ArcTan[a*x])/(85*a*c^3) + (E^ArcTan[a*x]*(1+4*a*x))/(17*a*c^3*(1+a^2*x^2)^2) + (12*E^ArcTan[a*x]*(1+2*a*x))/(85*a*c^3*(1+a^2*x^2))

Rule 5178

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]
```

Rule 5179

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx &= \frac{e^{\tan^{-1}(ax)}(1+4ax)}{17ac^3(1+a^2x^2)^2} + \frac{12 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{17c} \\
&= \frac{e^{\tan^{-1}(ax)}(1+4ax)}{17ac^3(1+a^2x^2)^2} + \frac{12e^{\tan^{-1}(ax)}(1+2ax)}{85ac^3(1+a^2x^2)} + \frac{24 \int \frac{e^{\tan^{-1}(ax)}}{c+a^2cx^2} dx}{85c^2} \\
&= \frac{24e^{\tan^{-1}(ax)}}{85ac^3} + \frac{e^{\tan^{-1}(ax)}(1+4ax)}{17ac^3(1+a^2x^2)^2} + \frac{12e^{\tan^{-1}(ax)}(1+2ax)}{85ac^3(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.21, size = 114, normalized size = 1.37

$$\frac{(12-24i)(1-iax)^{-1+\frac{i}{2}}(1+iax)^{-1-\frac{i}{2}} + \frac{24(1-iax)^{\frac{i}{2}}(1+iax)^{-\frac{i}{2}}((1-i)+ax)}{-i+ax} + \frac{5e^{\text{ArcTan}(ax)}(1+4ax)}{(1+a^2x^2)^2}}{85ac^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2)^3, x]

[Out] ((12 - 24*I)/((1 - I*a*x)^(1 - I/2)*(1 + I*a*x)^(1 + I/2)) + (24*(1 - I*a*x)^(I/2)*((1 - I) + a*x))/((1 + I*a*x)^(I/2)*(-I + a*x)) + (5*E^ArcTan[a*x]*(1 + 4*a*x))/(1 + a^2*x^2)^2)/(85*a*c^3)

Maple [A]

time = 0.08, size = 55, normalized size = 0.66

method	result	size
gospers	$\frac{e^{\arctan(ax)}(24a^4x^4+24a^3x^3+60a^2x^2+44ax+41)}{85(a^2x^2+1)^2c^3a}$	55

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)

[Out] 1/85*exp(arctan(a*x))*(24*a^4*x^4+24*a^3*x^3+60*a^2*x^2+44*a*x+41)/(a^2*x^2+1)^2/c^3/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^3, x)

Fricas [A]

time = 3.72, size = 66, normalized size = 0.80

$$\frac{(24 a^4 x^4 + 24 a^3 x^3 + 60 a^2 x^2 + 44 a x + 41) e^{\arctan(ax)}}{85 (a^5 c^3 x^4 + 2 a^3 c^3 x^2 + a c^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^3,x, algorithm="fricas")

[Out] 1/85*(24*a^4*x^4 + 24*a^3*x^3 + 60*a^2*x^2 + 44*a*x + 41)*e^(arctan(a*x))/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 223 vs. 2(75) = 150.

time = 4.12, size = 223, normalized size = 2.69

$$\begin{cases} \frac{24a^4x^4e^{\arctan(ax)}}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} + \frac{24a^3x^3e^{\arctan(ax)}}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} + \frac{60a^2x^2e^{\arctan(ax)}}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} + \frac{44axe^{\arctan(ax)}}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} + \frac{41e^{\arctan(ax)}}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} & \text{for } a \neq 0 \\ \frac{x}{c^3} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**3,x)

[Out] Piecewise((24*a**4*x**4*exp(atan(a*x))/(85*a**5*c**3*x**4 + 170*a**3*c**3*x**2 + 85*a*c**3) + 24*a**3*x**3*exp(atan(a*x))/(85*a**5*c**3*x**4 + 170*a**3*c**3*x**2 + 85*a*c**3) + 60*a**2*x**2*exp(atan(a*x))/(85*a**5*c**3*x**4 + 170*a**3*c**3*x**2 + 85*a*c**3) + 44*a*x*exp(atan(a*x))/(85*a**5*c**3*x**4 + 170*a**3*c**3*x**2 + 85*a*c**3) + 41*exp(atan(a*x))/(85*a**5*c**3*x**4 + 170*a**3*c**3*x**2 + 85*a*c**3), Ne(a, 0)), (x/c**3, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^3,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.62, size = 74, normalized size = 0.89

$$\frac{24 e^{\arctan(ax)}}{85 a c^3} + \frac{12 e^{\arctan(ax)} (2 a x + 1)}{85 a c^3 (a^2 x^2 + 1)} + \frac{e^{\arctan(ax)} (4 a x + 1)}{17 a c^3 (a^2 x^2 + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^3,x)

[Out] (24*exp(atan(a*x)))/(85*a*c^3) + (12*exp(atan(a*x))*(2*a*x + 1))/(85*a*c^3*(a^2*x^2 + 1)) + (exp(atan(a*x))*(4*a*x + 1))/(17*a*c^3*(a^2*x^2 + 1)^2)

$$3.251 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$$

Optimal. Leaf size=116

$$\frac{144e^{\text{ArcTan}(ax)}}{629ac^4} + \frac{e^{\text{ArcTan}(ax)}(1+6ax)}{37ac^4(1+a^2x^2)^3} + \frac{30e^{\text{ArcTan}(ax)}(1+4ax)}{629ac^4(1+a^2x^2)^2} + \frac{72e^{\text{ArcTan}(ax)}(1+2ax)}{629ac^4(1+a^2x^2)}$$

[Out] 144/629*exp(arctan(a*x))/a/c^4+1/37*exp(arctan(a*x))*(6*a*x+1)/a/c^4/(a^2*x^2+1)^3+30/629*exp(arctan(a*x))*(4*a*x+1)/a/c^4/(a^2*x^2+1)^2+72/629*exp(arctan(a*x))*(2*a*x+1)/a/c^4/(a^2*x^2+1)

Rubi [A]

time = 0.08, antiderivative size = 116, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5178, 5179}

$$\frac{72(2ax+1)e^{\text{ArcTan}(ax)}}{629ac^4(a^2x^2+1)} + \frac{30(4ax+1)e^{\text{ArcTan}(ax)}}{629ac^4(a^2x^2+1)^2} + \frac{(6ax+1)e^{\text{ArcTan}(ax)}}{37ac^4(a^2x^2+1)^3} + \frac{144e^{\text{ArcTan}(ax)}}{629ac^4}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c + a^2*c*x^2)^4,x]

[Out] (144*E^ArcTan[a*x]/(629*a*c^4) + (E^ArcTan[a*x]*(1 + 6*a*x))/(37*a*c^4*(1 + a^2*x^2)^3) + (30*E^ArcTan[a*x]*(1 + 4*a*x))/(629*a*c^4*(1 + a^2*x^2)^2) + (72*E^ArcTan[a*x]*(1 + 2*a*x))/(629*a*c^4*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.))/((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x]))/(a*c*(n^2 + 4*(p + 1)^2)), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)])*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^4} dx &= \frac{e^{\tan^{-1}(ax)}(1+6ax)}{37ac^4(1+a^2x^2)^3} + \frac{30 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx}{37c} \\
&= \frac{e^{\tan^{-1}(ax)}(1+6ax)}{37ac^4(1+a^2x^2)^3} + \frac{30e^{\tan^{-1}(ax)}(1+4ax)}{629ac^4(1+a^2x^2)^2} + \frac{360 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{629c^2} \\
&= \frac{e^{\tan^{-1}(ax)}(1+6ax)}{37ac^4(1+a^2x^2)^3} + \frac{30e^{\tan^{-1}(ax)}(1+4ax)}{629ac^4(1+a^2x^2)^2} + \frac{72e^{\tan^{-1}(ax)}(1+2ax)}{629ac^4(1+a^2x^2)} + \frac{144 \int \frac{e^{\tan^{-1}(ax)}}{c+a^2cx^2} dx}{629c^3} \\
&= \frac{144e^{\tan^{-1}(ax)}}{629ac^4} + \frac{e^{\tan^{-1}(ax)}(1+6ax)}{37ac^4(1+a^2x^2)^3} + \frac{30e^{\tan^{-1}(ax)}(1+4ax)}{629ac^4(1+a^2x^2)^2} + \frac{72e^{\tan^{-1}(ax)}(1+2ax)}{629ac^4(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.20, size = 123, normalized size = 1.06

$$\frac{17ce^{\text{ArcTan}(ax)}(1+6ax) + 6(c+a^2cx^2) \left(5e^{\text{ArcTan}(ax)}(1+4ax) + 12(1-iax)^{\frac{1}{2}}(1+iax)^{-\frac{1}{2}}(-i+ax)(i+ax)(3+2ax+2a^2x^2) \right)}{629ac^2(c+a^2cx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2)^4,x]

[Out] (17*c*E^ArcTan[a*x]*(1 + 6*a*x) + 6*(c + a^2*c*x^2)*(5*E^ArcTan[a*x]*(1 + 4*a*x) + (12*(1 - I*a*x)^(I/2)*(-I + a*x)*(I + a*x)*(3 + 2*a*x + 2*a^2*x^2)))/(1 + I*a*x)^(I/2)))/(629*a*c^2*(c + a^2*c*x^2)^3)

Maple [A]

time = 0.07, size = 71, normalized size = 0.61

method	result	size
gospers	$\frac{e^{\arctan(ax)}(144a^6x^6 + 144a^5x^5 + 504a^4x^4 + 408a^3x^3 + 606a^2x^2 + 366ax + 263)}{629(a^2x^2 + 1)^3c^4a}$	71

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^4,x,method=_RETURNVERBOSE)

[Out] 1/629*exp(arctan(a*x))*(144*a^6*x^6+144*a^5*x^5+504*a^4*x^4+408*a^3*x^3+606*a^2*x^2+366*a*x+263)/(a^2*x^2+1)^3/c^4/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="maxima")

[Out] integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^4, x)

Fricas [A]

time = 2.61, size = 93, normalized size = 0.80

$$\frac{(144 a^6 x^6 + 144 a^5 x^5 + 504 a^4 x^4 + 408 a^3 x^3 + 606 a^2 x^2 + 366 a x + 263) e^{\arctan(ax)}}{629 (a^7 c^4 x^6 + 3 a^5 c^4 x^4 + 3 a^3 c^4 x^2 + a c^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="fricas")

[Out] 1/629*(144*a^6*x^6 + 144*a^5*x^5 + 504*a^4*x^4 + 408*a^3*x^3 + 606*a^2*x^2 + 366*a*x + 263)*e^(arctan(a*x))/(a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 398 vs. $2(107) = 214$.

time = 10.80, size = 398, normalized size = 3.43

$$\left(\frac{144 a^6 x^6 \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} + \frac{144 a^5 x^5 \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} + \frac{504 a^4 x^4 \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} + \frac{408 a^3 x^3 \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} + \frac{606 a^2 x^2 \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} + \frac{366 a x \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} + \frac{263 \exp(\arctan(ax))}{629 a^7 c^4 x^6 + 1887 a^5 c^4 x^4 + 1887 a^3 c^4 x^2 + 629 a c^4} \right) \text{ for } a \neq 0$$

otherwise

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**4,x)

[Out] Piecewise((144*a**6*x**6*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4) + 144*a**5*x**5*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4) + 504*a**4*x**4*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4) + 408*a**3*x**3*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4) + 606*a**2*x**2*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4) + 366*a*x*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4) + 263*exp(atan(a*x))/(629*a**7*c**4*x**6 + 1887*a**5*c**4*x**4 + 1887*a**3*c**4*x**2 + 629*a*c**4), Ne(a, 0)), (x/c**4, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.69, size = 104, normalized size = 0.90

$$\frac{144 e^{\operatorname{atan}(ax)}}{629 a c^4} + \frac{72 e^{\operatorname{atan}(ax)} (2 a x + 1)}{629 a c^4 (a^2 x^2 + 1)} + \frac{30 e^{\operatorname{atan}(ax)} (4 a x + 1)}{629 a c^4 (a^2 x^2 + 1)^2} + \frac{e^{\operatorname{atan}(ax)} (6 a x + 1)}{37 a c^4 (a^2 x^2 + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^4,x)

[Out] (144*exp(atan(a*x)))/(629*a*c^4) + (72*exp(atan(a*x))*(2*a*x + 1))/(629*a*c^4*(a^2*x^2 + 1)) + (30*exp(atan(a*x))*(4*a*x + 1))/(629*a*c^4*(a^2*x^2 + 1)^2) + (exp(atan(a*x))*(6*a*x + 1))/(37*a*c^4*(a^2*x^2 + 1)^3)

$$3.252 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^5} dx$$

Optimal. Leaf size=149

$$\frac{8064e^{\text{ArcTan}(ax)}}{40885ac^5} + \frac{e^{\text{ArcTan}(ax)}(1+8ax)}{65ac^5(1+a^2x^2)^4} + \frac{56e^{\text{ArcTan}(ax)}(1+6ax)}{2405ac^5(1+a^2x^2)^3} + \frac{336e^{\text{ArcTan}(ax)}(1+4ax)}{8177ac^5(1+a^2x^2)^2} + \frac{4032e^{\text{ArcTan}(ax)}(1+2ax)}{40885ac^5(1+a^2x^2)}$$

[Out] 8064/40885*exp(arctan(a*x))/a/c^5+1/65*exp(arctan(a*x))*(8*a*x+1)/a/c^5/(a^2*x^2+1)^4+56/2405*exp(arctan(a*x))*(6*a*x+1)/a/c^5/(a^2*x^2+1)^3+336/8177*exp(arctan(a*x))*(4*a*x+1)/a/c^5/(a^2*x^2+1)^2+4032/40885*exp(arctan(a*x))*(2*a*x+1)/a/c^5/(a^2*x^2+1)

Rubi [A]

time = 0.11, antiderivative size = 149, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5178, 5179}

$$\frac{4032(2ax+1)e^{\text{ArcTan}(ax)}}{40885ac^5(a^2x^2+1)} + \frac{336(4ax+1)e^{\text{ArcTan}(ax)}}{8177ac^5(a^2x^2+1)^2} + \frac{56(6ax+1)e^{\text{ArcTan}(ax)}}{2405ac^5(a^2x^2+1)^3} + \frac{(8ax+1)e^{\text{ArcTan}(ax)}}{65ac^5(a^2x^2+1)^4} + \frac{8064e^{\text{ArcTan}(ax)}}{40885ac^5}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c + a^2*c*x^2)^5, x]

[Out] (8064*E^ArcTan[a*x])/(40885*a*c^5) + (E^ArcTan[a*x]*(1 + 8*a*x))/(65*a*c^5*(1 + a^2*x^2)^4) + (56*E^ArcTan[a*x]*(1 + 6*a*x))/(2405*a*c^5*(1 + a^2*x^2)^3) + (336*E^ArcTan[a*x]*(1 + 4*a*x))/(8177*a*c^5*(1 + a^2*x^2)^2) + (4032*E^ArcTan[a*x]*(1 + 2*a*x))/(40885*a*c^5*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^5} dx &= \frac{e^{\tan^{-1}(ax)}(1+8ax)}{65ac^5(1+a^2x^2)^4} + \frac{56 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^4} dx}{65c} \\
&= \frac{e^{\tan^{-1}(ax)}(1+8ax)}{65ac^5(1+a^2x^2)^4} + \frac{56e^{\tan^{-1}(ax)}(1+6ax)}{2405ac^5(1+a^2x^2)^3} + \frac{336 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx}{481c^2} \\
&= \frac{e^{\tan^{-1}(ax)}(1+8ax)}{65ac^5(1+a^2x^2)^4} + \frac{56e^{\tan^{-1}(ax)}(1+6ax)}{2405ac^5(1+a^2x^2)^3} + \frac{336e^{\tan^{-1}(ax)}(1+4ax)}{8177ac^5(1+a^2x^2)^2} + \frac{4032 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{8177c^3} \\
&= \frac{e^{\tan^{-1}(ax)}(1+8ax)}{65ac^5(1+a^2x^2)^4} + \frac{56e^{\tan^{-1}(ax)}(1+6ax)}{2405ac^5(1+a^2x^2)^3} + \frac{336e^{\tan^{-1}(ax)}(1+4ax)}{8177ac^5(1+a^2x^2)^2} + \frac{4032e^{\tan^{-1}(ax)}(1+2ax)}{40885ac^5(1+a^2x^2)} \\
&= \frac{8064e^{\tan^{-1}(ax)}}{40885ac^5} + \frac{e^{\tan^{-1}(ax)}(1+8ax)}{65ac^5(1+a^2x^2)^4} + \frac{56e^{\tan^{-1}(ax)}(1+6ax)}{2405ac^5(1+a^2x^2)^3} + \frac{336e^{\tan^{-1}(ax)}(1+4ax)}{8177ac^5(1+a^2x^2)^2} + \frac{4032e^{\tan^{-1}(ax)}(1+2ax)}{40885ac^5(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.20, size = 153, normalized size = 1.03

$$\frac{629e^{\text{ArcTan}(ax)}(1+8ax) + \frac{56(c+a^2cx^2)(17ce^{\text{ArcTan}(ax)}(1+6ax)+6(c+a^2cx^2)(5e^{\text{ArcTan}(ax)}(1+4ax)+12(1-iax)^{\frac{1}{2}}(1+iax)^{-\frac{1}{2}}(-i+ax)(i+ax)(3+2ax+2a^2x^2)))}{c^2}}{40885ac(c+a^2cx^2)^4}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2)^5, x]

[Out] (629*E^ArcTan[a*x]*(1 + 8*a*x) + (56*(c + a^2*c*x^2)*(17*c*E^ArcTan[a*x]*(1 + 6*a*x) + 6*(c + a^2*c*x^2)*(5*E^ArcTan[a*x]*(1 + 4*a*x) + (12*(1 - I*a*x)^(I/2)*(-I + a*x)*(I + a*x)*(3 + 2*a*x + 2*a^2*x^2))/(1 + I*a*x)^(I/2)))))/c^2)/(40885*a*c*(c + a^2*c*x^2)^4)

Maple [A]

time = 0.09, size = 87, normalized size = 0.58

method	result	size
gospers	$\frac{e^{\arctan(ax)}(8064a^8x^8+8064a^7x^7+36288a^6x^6+30912a^5x^5+62160a^4x^4+43344a^3x^3+48664a^2x^2+25528ax+15357)}{40885(a^2x^2+1)^4c^5a}$	87

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^5, x, method=_RETURNVERBOSE)

[Out] 1/40885*exp(arctan(a*x))*(8064*a^8*x^8+8064*a^7*x^7+36288*a^6*x^6+30912*a^5*x^5+62160*a^4*x^4+43344*a^3*x^3+48664*a^2*x^2+25528*a*x+15357)/(a^2*x^2+1)^4/c^5/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^5,x, algorithm="maxima")**[Out]** integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^5, x)**Fricas [A]**

time = 2.53, size = 120, normalized size = 0.81

$$\frac{(8064 a^8 x^8 + 8064 a^7 x^7 + 36288 a^6 x^6 + 30912 a^5 x^5 + 62160 a^4 x^4 + 43344 a^3 x^3 + 48664 a^2 x^2 + 25528 a x + 15357) e^{\arctan(ax)}}{40885 (a^9 c^5 x^8 + 4 a^7 c^5 x^6 + 6 a^5 c^5 x^4 + 4 a^3 c^5 x^2 + a c^5)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^5,x, algorithm="fricas")

[Out] 1/40885*(8064*a^8*x^8 + 8064*a^7*x^7 + 36288*a^6*x^6 + 30912*a^5*x^5 + 62160*a^4*x^4 + 43344*a^3*x^3 + 48664*a^2*x^2 + 25528*a*x + 15357)*e^(arctan(a*x))/(a^9*c^5*x^8 + 4*a^7*c^5*x^6 + 6*a^5*c^5*x^4 + 4*a^3*c^5*x^2 + a*c^5)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 620 vs. 2(139) = 278.

time = 25.66, size = 620, normalized size = 4.16

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**5,x)

[Out] Piecewise((8064*a**8*x**8*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 8064*a**7*x**7*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 36288*a**6*x**6*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 30912*a**5*x**5*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 62160*a**4*x**4*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 43344*a**3*x**3*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 48664*a**2*x**2*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 25528*a*x*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 15357*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5))

```
5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5) + 15357*exp(atan(a*x))/(40885*a**9*c**5*x**8 + 163540*a**7*c**5*x**6 + 245310*a**5*c**5*x**4 + 163540*a**3*c**5*x**2 + 40885*a*c**5), Ne(a, 0)), (x/c**5, True))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^5,x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [B]

time = 0.74, size = 134, normalized size = 0.90

$$e^{\operatorname{atan}(ax)} \left(\frac{15357}{40885 a^9 c^5} + \frac{25528x}{40885 a^8 c^5} + \frac{8064x^2}{40885 a^7 c^5} + \frac{8064x^3}{40885 a^6 c^5} + \frac{36288x^4}{40885 a^5 c^5} + \frac{30912x^5}{40885 a^4 c^5} + \frac{336x^6}{221 a^3 c^5} + \frac{43344x^7}{40885 a^2 c^5} + \frac{48664x^8}{40885 a c^5} \right) \frac{1}{\frac{1}{a^8} + x^8 + \frac{4x^6}{a^2} + \frac{6x^4}{a^4} + \frac{4x^2}{a^6}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^5,x)
```

```
[Out] (exp(atan(a*x))*(15357/(40885*a^9*c^5) + (25528*x)/(40885*a^8*c^5) + (8064*x^2)/(40885*a^7*c^5) + (8064*x^3)/(40885*a^6*c^5) + (36288*x^4)/(40885*a^5*c^5) + (30912*x^5)/(40885*a^4*c^5) + (336*x^6)/(221*a^3*c^5) + (43344*x^7)/(40885*a^2*c^5) + (48664*x^8)/(40885*a*c^5)))/(1/a^8 + x^8 + (4*x^6)/a^2 + (6*x^4)/a^4 + (4*x^2)/a^6)
```


3.253 $\int e^{\text{ArcTan}(ax)}(c + a^2cx^2)^{3/2} dx$

Optimal. Leaf size=98

$$\frac{\left(\frac{1}{13} + \frac{5i}{13}\right) 2^{\frac{3}{2} - \frac{i}{2}} c (1 - iax)^{\frac{5}{2} + \frac{i}{2}} \sqrt{c + a^2cx^2} {}_2F_1\left(-\frac{3}{2} + \frac{i}{2}, \frac{5}{2} + \frac{i}{2}; \frac{7}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2x^2}}$$

[Out] (1/13+5/13*I)*2^(3/2-1/2*I)*c*(1-I*a*x)^(5/2+1/2*I)*hypergeom([5/2+1/2*I, -3/2+1/2*I], [7/2+1/2*I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^(1/2)/a/(a^2*x^2+1)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{1}{13} + \frac{5i}{13}\right) 2^{\frac{3}{2} - \frac{i}{2}} c (1 - iax)^{\frac{5}{2} + \frac{i}{2}} \sqrt{a^2cx^2 + c} {}_2F_1\left(-\frac{3}{2} + \frac{i}{2}, \frac{5}{2} + \frac{i}{2}; \frac{7}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{a^2x^2 + 1}}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]*(c + a^2*c*x^2)^(3/2), x]

[Out] ((1/13 + (5*I)/13)*2^(3/2 - I/2)*c*(1 - I*a*x)^(5/2 + I/2)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-3/2 + I/2, 5/2 + I/2, 7/2 + I/2, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && Eq

$Q[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{\tan^{-1}(ax)} (c + a^2cx^2)^{3/2} dx &= \frac{\left(c\sqrt{c + a^2cx^2}\right) \int e^{\tan^{-1}(ax)} (1 + a^2x^2)^{3/2} dx}{\sqrt{1 + a^2x^2}} \\ &= \frac{\left(c\sqrt{c + a^2cx^2}\right) \int (1 - iax)^{\frac{3}{2} + \frac{i}{2}} (1 + iax)^{\frac{3}{2} - \frac{i}{2}} dx}{\sqrt{1 + a^2x^2}} \\ &= \frac{\left(\frac{1}{13} + \frac{5i}{13}\right) 2^{\frac{3}{2} - \frac{i}{2}} c (1 - iax)^{\frac{5}{2} + \frac{i}{2}} \sqrt{c + a^2cx^2} {}_2F_1\left(-\frac{3}{2} + \frac{i}{2}, \frac{5}{2} + \frac{i}{2}; \frac{7}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2x^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 98, normalized size = 1.00

$$\frac{\left(\frac{1}{13} + \frac{5i}{13}\right) 2^{\frac{3}{2} - \frac{i}{2}} c (1 - iax)^{\frac{5}{2} + \frac{i}{2}} \sqrt{c + a^2cx^2} {}_2F_1\left(-\frac{3}{2} + \frac{i}{2}, \frac{5}{2} + \frac{i}{2}; \frac{7}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]*(c + a^2*c*x^2)^(3/2), x]

[Out] ((1/13 + (5*I)/13)*2^(3/2 - I/2)*c*(1 - I*a*x)^(5/2 + I/2)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-3/2 + I/2, 5/2 + I/2, 7/2 + I/2, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{\arctan(ax)} (a^2cx^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))*(a^2*c*x^2+c)^(3/2), x)

[Out] int(exp(arctan(a*x))*(a^2*c*x^2+c)^(3/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(3/2)*e^(arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(3/2)*e^(arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^{\frac{3}{2}} e^{\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))*(a**2*c*x**2+c)**(3/2),x)

[Out] Integral((c*(a**2*x**2 + 1))**(3/2)*exp(atan(a*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{\operatorname{atan}(ax)} (ca^2x^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))*(c + a^2*c*x^2)^(3/2),x)

[Out] int(exp(atan(a*x))*(c + a^2*c*x^2)^(3/2), x)

3.254 $\int e^{\text{ArcTan}(ax)} \sqrt{c + a^2cx^2} dx$

Optimal. Leaf size=97

$$\frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{\frac{1}{2} - \frac{i}{2}} (1 - iax)^{\frac{3}{2} + \frac{i}{2}} \sqrt{c + a^2cx^2} {}_2F_1\left(-\frac{1}{2} + \frac{i}{2}, \frac{3}{2} + \frac{i}{2}; \frac{5}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2x^2}}$$

[Out] (1/5+3/5*I)*2^(1/2-1/2*I)*(1-I*a*x)^(3/2+1/2*I)*hypergeom([3/2+1/2*I, -1/2+1/2*I], [5/2+1/2*I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^(1/2)/a/(a^2*x^2+1)^(1/2)

Rubi [A]

time = 0.05, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{\frac{1}{2} - \frac{i}{2}} (1 - iax)^{\frac{3}{2} + \frac{i}{2}} \sqrt{a^2cx^2 + c} {}_2F_1\left(-\frac{1}{2} + \frac{i}{2}, \frac{3}{2} + \frac{i}{2}; \frac{5}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{a^2x^2 + 1}}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2],x]

[Out] ((1/5 + (3*I)/5)*2^(1/2 - I/2)*(1 - I*a*x)^(3/2 + I/2)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-1/2 + I/2, 3/2 + I/2, 5/2 + I/2, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Rule 71

```
Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^(n)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))
```

Rule 5181

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rule 5184

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int e^{\tan^{-1}(ax)} \sqrt{c + a^2 cx^2} dx &= \frac{\sqrt{c + a^2 cx^2} \int e^{\tan^{-1}(ax)} \sqrt{1 + a^2 x^2} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\sqrt{c + a^2 cx^2} \int (1 - iax)^{\frac{1}{2} + \frac{i}{2}} (1 + iax)^{\frac{1}{2} - \frac{i}{2}} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{\frac{1}{2} - \frac{i}{2}} (1 - iax)^{\frac{3}{2} + \frac{i}{2}} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} + \frac{i}{2}, \frac{3}{2} + \frac{i}{2}; \frac{5}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 97, normalized size = 1.00

$$\frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{\frac{1}{2} - \frac{i}{2}} (1 - iax)^{\frac{3}{2} + \frac{i}{2}} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} + \frac{i}{2}, \frac{3}{2} + \frac{i}{2}; \frac{5}{2} + \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2], x]

[Out] $\left(\left(\frac{1}{5} + \frac{3I}{5}\right) 2^{\frac{1}{2} - \frac{I}{2}} (1 - I a x)^{\frac{3}{2} + \frac{I}{2}} \sqrt{c + a^2 c x^2} \text{Hypergeometric2F1}\left[-\frac{1}{2} + \frac{I}{2}, \frac{3}{2} + \frac{I}{2}, \frac{5}{2} + \frac{I}{2}, \frac{(1 - I a x)}{2}\right]\right) / (a \sqrt{1 + a^2 x^2})$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{\arctan(ax)} \sqrt{a^2 c x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(arctan(a*x))*(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(arctan(a*x))*(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(sqrt(a^2*c*x^2 + c)*e^(arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(a^2*c*x^2 + c)*e^(arctan(a*x)), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c(a^2x^2 + 1)} e^{\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(atan(a*x))*(a**2*c*x**2+c)**(1/2),x)
```

```
[Out] Integral(sqrt(c*(a**2*x**2 + 1))*exp(atan(a*x)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{\operatorname{atan}(ax)} \sqrt{ca^2x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(atan(a*x))*(c + a^2*c*x^2)^(1/2),x)
```

```
[Out] int(exp(atan(a*x))*(c + a^2*c*x^2)^(1/2), x)
```

$$3.255 \quad \int \frac{e^{\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$$

Optimal. Leaf size=93

$$\frac{(1+i)2^{-\frac{1}{2}-\frac{i}{2}}(1-iax)^{\frac{1}{2}+\frac{i}{2}}\sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}+\frac{i}{2}, \frac{1}{2}+\frac{i}{2}; \frac{3}{2}+\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

[Out] (1+I)*2^(-1/2-1/2*I)*(1-I*a*x)^(1/2+1/2*I)*hypergeom([1/2+1/2*I, 1/2+1/2*I], [3/2+1/2*I], 1/2-1/2*I*a*x)*(a^2*x^2+1)^(1/2)/a/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.05, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$,

Rules used = {5184, 5181, 71}

$$\frac{(1+i)2^{-\frac{1}{2}-\frac{i}{2}}(1-iax)^{\frac{1}{2}+\frac{i}{2}}\sqrt{a^2x^2+1} {}_2F_1\left(\frac{1}{2}+\frac{i}{2}, \frac{1}{2}+\frac{i}{2}; \frac{3}{2}+\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/Sqrt[c + a^2*c*x^2], x]

[Out] ((1 + I)*(1 - I*a*x)^(1/2 + I/2)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 + I/2, 1/2 + I/2, 3/2 + I/2, (1 - I*a*x)/2])/(2^(1/2 + I/2)*a*Sqrt[c + a^2*c*x^2])

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{\tan^{-1}(ax)}}{\sqrt{c+a^2cx^2}} dx &= \frac{\sqrt{1+a^2x^2} \int \frac{e^{\tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx}{\sqrt{c+a^2cx^2}} \\
&= \frac{\sqrt{1+a^2x^2} \int (1-iax)^{-\frac{1}{2}+\frac{i}{2}} (1+iax)^{-\frac{1}{2}-\frac{i}{2}} dx}{\sqrt{c+a^2cx^2}} \\
&= \frac{(1+i)2^{-\frac{1}{2}-\frac{i}{2}} (1-iax)^{\frac{1}{2}+\frac{i}{2}} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}+\frac{i}{2}, \frac{1}{2}+\frac{i}{2}; \frac{3}{2}+\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 93, normalized size = 1.00

$$\frac{(1+i)2^{-\frac{1}{2}-\frac{i}{2}} (1-iax)^{\frac{1}{2}+\frac{i}{2}} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}+\frac{i}{2}, \frac{1}{2}+\frac{i}{2}; \frac{3}{2}+\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^ArcTan[a*x]/Sqrt[c + a^2*c*x^2], x]`

```
[Out] ((1 + I)*(1 - I*a*x)^(1/2 + I/2)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 + I/2, 1/2 + I/2, 3/2 + I/2, (1 - I*a*x)/2])/(2^(1/2 + I/2)*a*Sqrt[c + a^2*c*x^2])
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{\arctan(ax)}}{\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)``[Out] int(exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2), x, algorithm="maxima")`

[Out] integrate(e^(arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(e^(arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(exp(atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{\operatorname{atan}(ax)}}{\sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^(1/2),x)

[Out] int(exp(atan(a*x))/(c + a^2*c*x^2)^(1/2), x)

$$3.256 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=35

$$\frac{e^{\text{ArcTan}(ax)}(1+ax)}{2ac\sqrt{c+a^2cx^2}}$$

[Out] 1/2*exp(arctan(a*x))*(a*x+1)/a/c/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {5177}

$$\frac{(ax+1)e^{\text{ArcTan}(ax)}}{2ac\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c+a^2*c*x^2)^(3/2),x]

[Out] (E^ArcTan[a*x]*(1+a*x))/(2*a*c*Sqrt[c+a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.)+(d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n+a*x)*(E^(n*ArcTan[a*x]))/(a*c*(n^2+1)*Sqrt[c+d*x^2]), x] /; FreeQ[{a,c,d,n},x] && EqQ[d,a^2*c] && !IntegerQ[I*n]

Rubi steps

$$\int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx = \frac{e^{\tan^{-1}(ax)}(1+ax)}{2ac\sqrt{c+a^2cx^2}}$$

Mathematica [A]

time = 0.01, size = 35, normalized size = 1.00

$$\frac{e^{\text{ArcTan}(ax)}(1+ax)}{2ac\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^ArcTan[a*x]/(c+a^2*c*x^2)^(3/2),x]

[Out] (E^ArcTan[a*x]*(1+a*x))/(2*a*c*Sqrt[c+a^2*c*x^2])

Maple [A]

time = 0.08, size = 37, normalized size = 1.06

method	result	size
gosper	$\frac{(a^2x^2+1)(ax+1)e^{\arctan(ax)}}{2a(a^2cx^2+c)^{\frac{3}{2}}}$	37

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)`[Out] $1/2*(a^2*x^2+1)*(a*x+1)*\exp(\arctan(a*x))/a/(a^2*c*x^2+c)^{(3/2)}$ **Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`[Out] `integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^(3/2), x)`**Fricas [A]**

time = 3.11, size = 42, normalized size = 1.20

$$\frac{\sqrt{a^2cx^2 + c} (ax + 1)e^{\arctan(ax)}}{2(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`[Out] `1/2*sqrt(a^2*c*x^2 + c)*(a*x + 1)*e^(arctan(a*x))/(a^3*c^2*x^2 + a*c^2)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(atan(a*x))/(a**2*c*x**2+c)**(3/2),x)`[Out] `Integral(exp(atan(a*x))/(c*(a**2*x**2 + 1))**(3/2), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B]

time = 0.60, size = 33, normalized size = 0.94

$$\frac{e^{\operatorname{atan}(ax)} \left(\frac{x}{2c} + \frac{1}{2ac} \right)}{\sqrt{ca^2x^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(atan(a*x))/(c + a^2*c*x^2)^(3/2),x)`

[Out] `(exp(atan(a*x))*(x/(2*c) + 1/(2*a*c)))/(c + a^2*c*x^2)^(1/2)`

$$3.257 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$$

Optimal. Leaf size=72

$$\frac{e^{\text{ArcTan}(ax)}(1+3ax)}{10ac(c+a^2cx^2)^{3/2}} + \frac{3e^{\text{ArcTan}(ax)}(1+ax)}{10ac^2\sqrt{c+a^2cx^2}}$$

[Out] 1/10*exp(arctan(a*x))*(3*a*x+1)/a/c/(a^2*c*x^2+c)^(3/2)+3/10*exp(arctan(a*x))*(a*x+1)/a/c^2/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 72, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5177}

$$\frac{3(ax+1)e^{\text{ArcTan}(ax)}}{10ac^2\sqrt{a^2cx^2+c}} + \frac{(3ax+1)e^{\text{ArcTan}(ax)}}{10ac(a^2cx^2+c)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c+a^2*c*x^2)^(5/2),x]

[Out] (E^ArcTan[a*x]*(1+3*a*x))/(10*a*c*(c+a^2*c*x^2)^(3/2)) + (3*E^ArcTan[a*x]*(1+a*x))/(10*a*c^2*Sqrt[c+a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.)+(d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n+a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2+1)*Sqrt[c+d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.)+(d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n-2*a*(p+1)*x)*(c+d*x^2)^(p+1)*(E^(n*ArcTan[a*x])/(a*c*(n^2+4*(p+1)^2))), x] + Dist[2*(p+1)*((2*p+3)/(c*(n^2+4*(p+1)^2))), Int[(c+d*x^2)^(p+1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2+4*(p+1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\int \frac{e^{\tan^{-1}(ax)}}{(c + a^2cx^2)^{5/2}} dx = \frac{e^{\tan^{-1}(ax)}(1 + 3ax)}{10ac(c + a^2cx^2)^{3/2}} + \frac{3 \int \frac{e^{\tan^{-1}(ax)}}{(c + a^2cx^2)^{3/2}} dx}{5c}$$

$$= \frac{e^{\tan^{-1}(ax)}(1 + 3ax)}{10ac(c + a^2cx^2)^{3/2}} + \frac{3e^{\tan^{-1}(ax)}(1 + ax)}{10ac^2\sqrt{c + a^2cx^2}}$$

Mathematica [A]

time = 0.03, size = 60, normalized size = 0.83

$$\frac{e^{\text{ArcTan}(ax)}(4 + 6ax + 3a^2x^2 + 3a^3x^3)}{10c^2(a + a^3x^2)\sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2)^(5/2), x]``[Out] (E^ArcTan[a*x]*(4 + 6*a*x + 3*a^2*x^2 + 3*a^3*x^3))/(10*c^2*(a + a^3*x^2)*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.08, size = 54, normalized size = 0.75

method	result	size
gospers	$\frac{(a^2x^2+1)(3a^3x^3+3a^2x^2+6ax+4)e^{\arctan(ax)}}{10a(a^2cx^2+c)^{\frac{5}{2}}}$	54

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, method=_RETURNVERBOSE)``[Out] 1/10*(a^2*x^2+1)*(3*a^3*x^3+3*a^2*x^2+6*a*x+4)*exp(arctan(a*x))/a/(a^2*c*x^2+c)^(5/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, algorithm="maxima")``[Out] integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^(5/2), x)`

Fricas [A]

time = 3.34, size = 70, normalized size = 0.97

$$\frac{(3a^3x^3 + 3a^2x^2 + 6ax + 4)\sqrt{a^2cx^2 + c} e^{\arctan(ax)}}{10(a^5c^3x^4 + 2a^3c^3x^2 + ac^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="fricas")

[Out] 1/10*(3*a^3*x^3 + 3*a^2*x^2 + 6*a*x + 4)*sqrt(a^2*c*x^2 + c)*e^(arctan(a*x))/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{\operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**(5/2),x)

[Out] Integral(exp(atan(a*x))/(c*(a**2*x**2 + 1))**(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.65, size = 78, normalized size = 1.08

$$\frac{e^{\operatorname{atan}(ax)} \left(\frac{2}{5a^3c^2} + \frac{3x^3}{10c^2} + \frac{3x}{5a^2c^2} + \frac{3x^2}{10ac^2} \right)}{\frac{\sqrt{ca^2x^2 + c}}{a^2} + x^2 \sqrt{ca^2x^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^(5/2),x)

[Out] (exp(atan(a*x))*(2/(5*a^3*c^2) + (3*x^3)/(10*c^2) + (3*x)/(5*a^2*c^2) + (3*x^2)/(10*a*c^2)))/((c + a^2*c*x^2)^(1/2)/a^2 + x^2*(c + a^2*c*x^2)^(1/2))

$$3.258 \quad \int \frac{e^{\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$$

Optimal. Leaf size=108

$$\frac{e^{\text{ArcTan}(ax)}(1+5ax)}{26ac(c+a^2cx^2)^{5/2}} + \frac{e^{\text{ArcTan}(ax)}(1+3ax)}{13ac^2(c+a^2cx^2)^{3/2}} + \frac{3e^{\text{ArcTan}(ax)}(1+ax)}{13ac^3\sqrt{c+a^2cx^2}}$$

[Out] 1/26*exp(arctan(a*x))*(5*a*x+1)/a/c/(a^2*c*x^2+c)^(5/2)+1/13*exp(arctan(a*x))*(3*a*x+1)/a/c^2/(a^2*c*x^2+c)^(3/2)+3/13*exp(arctan(a*x))*(a*x+1)/a/c^3/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.08, antiderivative size = 108, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5177}

$$\frac{3(ax+1)e^{\text{ArcTan}(ax)}}{13ac^3\sqrt{a^2cx^2+c}} + \frac{(3ax+1)e^{\text{ArcTan}(ax)}}{13ac^2(a^2cx^2+c)^{3/2}} + \frac{(5ax+1)e^{\text{ArcTan}(ax)}}{26ac(a^2cx^2+c)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[E^ArcTan[a*x]/(c+a^2*c*x^2)^(7/2),x]

[Out] (E^ArcTan[a*x]*(1+5*a*x))/(26*a*c*(c+a^2*c*x^2)^(5/2)) + (E^ArcTan[a*x]*(1+3*a*x))/(13*a*c^2*(c+a^2*c*x^2)^(3/2)) + (3*E^ArcTan[a*x]*(1+a*x))/(13*a*c^3*Sqrt[c+a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.)+(d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(n+a*x)*(E^(n*ArcTan[a*x]))/(a*c*(n^2+1)*Sqrt[c+d*x^2]), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.)+(d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n-2*a*(p+1)*x)*(c+d*x^2)^(p+1)*(E^(n*ArcTan[a*x]))/(a*c*(n^2+4*(p+1)^2)), x] + Dist[2*(p+1)*((2*p+3)/(c*(n^2+4*(p+1)^2))), Int[(c+d*x^2)^(p+1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2+4*(p+1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\begin{aligned}
\int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^{7/2}} dx &= \frac{e^{\tan^{-1}(ax)}(1+5ax)}{26ac(c+a^2cx^2)^{5/2}} + \frac{10 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^{5/2}} dx}{13c} \\
&= \frac{e^{\tan^{-1}(ax)}(1+5ax)}{26ac(c+a^2cx^2)^{5/2}} + \frac{e^{\tan^{-1}(ax)}(1+3ax)}{13ac^2(c+a^2cx^2)^{3/2}} + \frac{6 \int \frac{e^{\tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx}{13c^2} \\
&= \frac{e^{\tan^{-1}(ax)}(1+5ax)}{26ac(c+a^2cx^2)^{5/2}} + \frac{e^{\tan^{-1}(ax)}(1+3ax)}{13ac^2(c+a^2cx^2)^{3/2}} + \frac{3e^{\tan^{-1}(ax)}(1+ax)}{13ac^3\sqrt{c+a^2cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 79, normalized size = 0.73

$$\frac{e^{\text{ArcTan}(ax)}(9 + 17ax + 14a^2x^2 + 18a^3x^3 + 6a^4x^4 + 6a^5x^5)}{26ac^3(1 + a^2x^2)^2\sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^ArcTan[a*x]/(c + a^2*c*x^2)^(7/2), x]``[Out] (E^ArcTan[a*x]*(9 + 17*a*x + 14*a^2*x^2 + 18*a^3*x^3 + 6*a^4*x^4 + 6*a^5*x^5))/(26*a*c^3*(1 + a^2*x^2)^2*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.07, size = 70, normalized size = 0.65

method	result	size
gospers	$\frac{(a^2x^2+1)(6a^5x^5+6a^4x^4+18a^3x^3+14a^2x^2+17ax+9)e^{\arctan(ax)}}{26a(a^2cx^2+c)^{\frac{7}{2}}}$	70

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, method=_RETURNVERBOSE)``[Out] 1/26*(a^2*x^2+1)*(6*a^5*x^5+6*a^4*x^4+18*a^3*x^3+14*a^2*x^2+17*a*x+9)*exp(arctan(a*x))/a/(a^2*c*x^2+c)^(7/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, algorithm="maxima")`

[Out] integrate(e^(arctan(a*x))/(a^2*c*x^2 + c)^(7/2), x)

Fricas [A]

time = 3.07, size = 97, normalized size = 0.90

$$\frac{(6a^5x^5 + 6a^4x^4 + 18a^3x^3 + 14a^2x^2 + 17ax + 9)\sqrt{a^2cx^2 + c}e^{\arctan(ax)}}{26(a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2),x, algorithm="fricas")

[Out] 1/26*(6*a^5*x^5 + 6*a^4*x^4 + 18*a^3*x^3 + 14*a^2*x^2 + 17*a*x + 9)*sqrt(a^2*c*x^2 + c)*e^(arctan(a*x))/(a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(atan(a*x))/(a**2*c*x**2+c)**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.67, size = 120, normalized size = 1.11

$$\frac{e^{\operatorname{atan}(ax)} \left(\frac{9}{26a^5c^3} + \frac{3x^5}{13c^3} + \frac{17x}{26a^4c^3} + \frac{3x^4}{13ac^3} + \frac{9x^3}{13a^2c^3} + \frac{7x^2}{13a^3c^3} \right)}{\sqrt{ca^2x^2 + c} + x^4\sqrt{ca^2x^2 + c} + \frac{2x^2\sqrt{ca^2x^2 + c}}{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(atan(a*x))/(c + a^2*c*x^2)^(7/2),x)

[Out] (exp(atan(a*x))*(9/(26*a^5*c^3) + (3*x^5)/(13*c^3) + (17*x)/(26*a^4*c^3) + (3*x^4)/(13*a*c^3) + (9*x^3)/(13*a^2*c^3) + (7*x^2)/(13*a^3*c^3)))/((c + a^2*c*x^2)^(1/2)/a^4 + x^4*(c + a^2*c*x^2)^(1/2) + (2*x^2*(c + a^2*c*x^2)^(1/2))/a^2)

3.259 $\int e^{2\text{ArcTan}(ax)}(c + a^2cx^2)^p dx$

Optimal. Leaf size=90

$$\frac{i2^{-i+p}(1-iax)^{(1+i)+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1(i-p, (1+i)+p; (2+i)+p; \frac{1}{2}(1-iax))}{a((1+i)+p)}$$

[Out] $I*2^{(-I+p)}*(1-I*a*x)^{(1+I+p)}*(a^2*c*x^2+c)^p*\text{hypergeom}([I-p, 1+I+p], [2+I+p], 1/2-1/2*I*a*x)/a/(1+I+p)/((a^2*x^2+1)^p)$

Rubi [A]

time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5184, 5181, 71}

$$\frac{i2^{p-i}(1-iax)^{p+(1+i)}(a^2x^2+1)^{-p}(a^2cx^2+c)^p {}_2F_1(i-p, p+(1+i); p+(2+i); \frac{1}{2}(1-iax))}{a(p+(1+i))}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(2*\text{ArcTan}[a*x])}*(c + a^2*c*x^2)^p, x]$

[Out] $(I*2^{(-I+p)}*(1-I*a*x)^{((1+I)+p)}*(c+a^2*c*x^2)^p*\text{Hypergeometric2F1}[I-p, (1+I)+p, (2+I)+p, (1-I*a*x)/2])/a*((1+I)+p)*(1+a^2*x^2)^p)$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+)^2)^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}/(b*(m+1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[a_+]*(x_+))^{(n_+)}}*((c_+) + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[a_+]*(x_+))^{(n_+)}}*((c_+) + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int e^{2 \tan^{-1}(ax)} (c + a^2 cx^2)^p dx &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int e^{2 \tan^{-1}(ax)} (1 + a^2 x^2)^p dx \\
&= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int (1 - iax)^{i+p} (1 + iax)^{-i+p} dx \\
&= \frac{i 2^{-i+p} (1 - iax)^{(1+i)+p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p {}_2F_1(i - p, (1 + i) + p; (2 + i) + p; \frac{1}{2}(1 - iax))}{a((1 + i) + p)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 90, normalized size = 1.00

$$\frac{i 2^{-i+p} (1 - iax)^{(1+i)+p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p {}_2F_1(i - p, (1 + i) + p; (2 + i) + p; \frac{1}{2}(1 - iax))}{a((1 + i) + p)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^p,x]

[Out] (I*2^(-I + p)*(1 - I*a*x)^((1 + I) + p)*(c + a^2*c*x^2)^p*Hypergeometric2F1[I - p, (1 + I) + p, (2 + I) + p, (1 - I*a*x)/2])/(a*((1 + I) + p)*(1 + a^2*x^2)^p)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{2 \arctan(ax)} (a^2 cx^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^p,x)

[Out] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^p,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^p*e^(2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^p*e^(2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^p e^{2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))*(a**2*c*x**2+c)**p,x)

[Out] Integral((c*(a**2*x**2 + 1))**p*exp(2*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{2\operatorname{atan}(ax)} (ca^2x^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^p,x)

[Out] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^p, x)

3.260 $\int e^{2\text{ArcTan}(ax)}(c + a^2cx^2)^2 dx$

Optimal. Leaf size=53

$$\frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{1-i} c^2 (1 - iax)^{3+i} {}_2F_1\left(-2 + i, 3 + i; 4 + i; \frac{1}{2}(1 - iax)\right)}{a}$$

[Out] (1/5+3/5*I)*2^(1-I)*c^2*(1-I*a*x)^(3+I)*hypergeom([3+I, -2+I], [4+I], 1/2-1/2*I*a*x)/a

Rubi [A]

time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5181, 71}

$$\frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{1-i} c^2 (1 - iax)^{3+i} {}_2F_1\left(-2 + i, 3 + i; 4 + i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^2,x]

[Out] ((1/5 + (3*I)/5)*2^(1 - I)*c^2*(1 - I*a*x)^(3 + I)*Hypergeometric2F1[-2 + I, 3 + I, 4 + I, (1 - I*a*x)/2])/a

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{2\text{tan}^{-1}(ax)}(c + a^2cx^2)^2 dx &= c^2 \int (1 - iax)^{2+i}(1 + iax)^{2-i} dx \\ &= \frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{1-i} c^2 (1 - iax)^{3+i} {}_2F_1\left(-2 + i, 3 + i; 4 + i; \frac{1}{2}(1 - iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 53, normalized size = 1.00

$$\frac{\left(\frac{1}{5} + \frac{3i}{5}\right) 2^{1-i} c^2 (1 - iax)^{3+i} {}_2F_1\left(-2 + i, 3 + i; 4 + i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^2,x]

[Out] ((1/5 + (3*I)/5)*2^(1 - I)*c^2*(1 - I*a*x)^(3 + I)*Hypergeometric2F1[-2 + I, 3 + I, 4 + I, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{2 \arctan(ax)} (a^2 c x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^2,x)

[Out] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^2*e^(2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] integral((a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2)*e^(2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c^2 \left(\int 2a^2 x^2 e^{2 \operatorname{atan}(ax)} dx + \int a^4 x^4 e^{2 \operatorname{atan}(ax)} dx + \int e^{2 \operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))*(a**2*c*x**2+c)**2,x)

[Out] c**2*(Integral(2*a**2*x**2*exp(2*atan(a*x)), x) + Integral(a**4*x**4*exp(2*atan(a*x)), x) + Integral(exp(2*atan(a*x)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{2\operatorname{atan}(ax)} (ca^2x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^2,x)

[Out] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^2, x)

3.261 $\int e^{2\text{ArcTan}(ax)}(c + a^2cx^2) dx$

Optimal. Leaf size=51

$$\frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-i} c (1 - iax)^{2+i} {}_2F_1\left(-1 + i, 2 + i; 3 + i; \frac{1}{2}(1 - iax)\right)}{a}$$

[Out] $(1/5+2/5*I)*2^{(1-I)}*c*(1-I*a*x)^{(2+I)}*\text{hypergeom}([-1+I, 2+I], [3+I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.02, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5181, 71}

$$\frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-i} c (1 - iax)^{2+i} {}_2F_1\left(-1 + i, 2 + i; 3 + i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(2*\text{ArcTan}[a*x])}*(c + a^2*c*x^2), x]$

[Out] $((1/5 + (2*I)/5)*2^{(1 - I)}*c*(1 - I*a*x)^{(2 + I)}*\text{Hypergeometric2F1}[-1 + I, 2 + I, 3 + I, (1 - I*a*x)/2])/a$

Rule 71

$\text{Int}[(a_ + (b_)*(x_))^{(m_)}*((c_ + (d_)*(x_))^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_)*(x_)]*(n_))}*((c_ + (d_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{2\tan^{-1}(ax)}(c + a^2cx^2) dx &= c \int (1 - iax)^{1+i}(1 + iax)^{1-i} dx \\ &= \frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-i} c (1 - iax)^{2+i} {}_2F_1\left(-1 + i, 2 + i; 3 + i; \frac{1}{2}(1 - iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 51, normalized size = 1.00

$$\frac{\left(\frac{1}{5} + \frac{2i}{5}\right) 2^{1-i} c (1 - iax)^{2+i} {}_2F_1\left(-1 + i, 2 + i; 3 + i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])*(c + a^2*c*x^2), x]

[Out] ((1/5 + (2*I)/5)*2^(1 - I)*c*(1 - I*a*x)^(2 + I)*Hypergeometric2F1[-1 + I, 2 + I, 3 + I, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{2\arctan(ax)} (a^2cx^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))*(a^2*c*x^2+c), x)

[Out] int(exp(2*arctan(a*x))*(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)*e^(2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c), x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)*e^(2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c\left(\int a^2x^2e^{2\operatorname{atan}(ax)} dx + \int e^{2\operatorname{atan}(ax)} dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))*(a**2*c*x**2+c), x)

[Out] c*(Integral(a**2*x**2*exp(2*atan(a*x)), x) + Integral(exp(2*atan(a*x)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c), x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{2\operatorname{atan}(ax)} (ca^2x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))*(c + a^2*c*x^2), x)

[Out] int(exp(2*atan(a*x))*(c + a^2*c*x^2), x)

3.262 $\int e^{2\text{ArcTan}(ax)} dx$

Optimal. Leaf size=46

$$\frac{(1+i)2^{-1-i}(1-iax)^{1+i} {}_2F_1\left(i, 1+i; 2+i; \frac{1}{2}(1-iax)\right)}{a}$$

[Out] (1+I)*2^(-1-I)*(1-I*a*x)^(1+I)*hypergeom([I, 1+I], [2+I], 1/2-1/2*I*a*x)/a

Rubi [A]

time = 0.01, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5169, 71}

$$\frac{(1+i)2^{-1-i}(1-iax)^{1+i} {}_2F_1\left(i, 1+i; 2+i; \frac{1}{2}(1-iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x]), x]

[Out] ((1 + I)*(1 - I*a*x)^(1 + I)*Hypergeometric2F1[I, 1 + I, 2 + I, (1 - I*a*x)/2])/(2^(1 + I)*a)

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5169

Int[E^(ArcTan[(a_.)*(x_)]*(n_.)), x_Symbol] :> Int[(1 - I*a*x)^(I*(n/2))/(1 + I*a*x)^(I*(n/2)), x] /; FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{2 \tan^{-1}(ax)} dx &= \int (1 - iax)^i (1 + iax)^{-i} dx \\ &= \frac{(1+i)2^{-1-i}(1-iax)^{1+i} {}_2F_1\left(i, 1+i; 2+i; \frac{1}{2}(1-iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 37, normalized size = 0.80

$$\frac{(1-i)e^{(2+2i)\text{ArcTan}(ax)} {}_2F_1(1-i, 2; 2-i; -e^{2i\text{ArcTan}(ax)})}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x]), x]

[Out] ((1 - I)*E^((2 + 2*I)*ArcTan[a*x])*Hypergeometric2F1[1 - I, 2, 2 - I, -E^((2*I)*ArcTan[a*x])])/a

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{2\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x)), x)

[Out] int(exp(2*arctan(a*x)), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x)), x, algorithm="maxima")

[Out] integrate(e^(2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x)), x, algorithm="fricas")

[Out] integral(e^(2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{2\text{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(2*atan(a*x)),x)`

[Out] `Integral(exp(2*atan(a*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(2*arctan(a*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(2*atan(a*x)),x)`

[Out] `int(exp(2*atan(a*x)), x)`

$$3.263 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{c+a^2cx^2} dx$$

Optimal. Leaf size=18

$$\frac{e^{2\text{ArcTan}(ax)}}{2ac}$$

[Out] 1/2*exp(2*arctan(a*x))/a/c

Rubi [A]

time = 0.02, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {5179}

$$\frac{e^{2\text{ArcTan}(ax)}}{2ac}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2),x]

[Out] E^(2*ArcTan[a*x])/(2*a*c)

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\int \frac{e^{2\text{tan}^{-1}(ax)}}{c + a^2cx^2} dx = \frac{e^{2\text{tan}^{-1}(ax)}}{2ac}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.01, size = 34, normalized size = 1.89

$$\frac{(1 - iax)^i(1 + iax)^{-i}}{2ac}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2),x]

[Out] (1 - I*a*x)^I/(2*a*c*(1 + I*a*x)^I)

Maple [A]

time = 0.06, size = 16, normalized size = 0.89

method	result	size
gospers	$\frac{e^{2 \arctan(ax)}}{2ac}$	16
risch	$\frac{(-iax+1)^i (iax+1)^{-i}}{2ac}$	29

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c),x,method=_RETURNVERBOSE)
```

```
[Out] 1/2*exp(2*arctan(a*x))/a/c
```

Maxima [A]

time = 0.48, size = 15, normalized size = 0.83

$$\frac{e^{(2 \arctan(ax))}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="maxima")
```

```
[Out] 1/2*e^(2*arctan(a*x))/(a*c)
```

Fricas [A]

time = 2.41, size = 15, normalized size = 0.83

$$\frac{e^{(2 \arctan(ax))}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="fricas")
```

```
[Out] 1/2*e^(2*arctan(a*x))/(a*c)
```

Sympy [A]

time = 0.55, size = 15, normalized size = 0.83

$$\begin{cases} \frac{e^{2 \operatorname{atan}(ax)}}{2ac} & \text{for } a \neq 0 \\ \frac{x}{c} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*atan(a*x))/(a**2*c*x**2+c),x)
```

```
[Out] Piecewise((exp(2*atan(a*x))/(2*a*c), Ne(a, 0)), (x/c, True))
```


Giac [A]

time = 0.41, size = 15, normalized size = 0.83

$$\frac{e^{(2 \arctan(ax))}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="giac")

[Out] 1/2*e^(2*arctan(a*x))/(a*c)

Mupad [B]

time = 0.53, size = 15, normalized size = 0.83

$$\frac{e^{2 \operatorname{atan}(ax)}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))/(c + a^2*c*x^2),x)

[Out] exp(2*atan(a*x))/(2*a*c)

3.264

$$\int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$$

Optimal. Leaf size=53

$$\frac{e^{2\text{ArcTan}(ax)}}{8ac^2} + \frac{e^{2\text{ArcTan}(ax)}(1+ax)}{4ac^2(1+a^2x^2)}$$

[Out] 1/8*exp(2*arctan(a*x))/a/c^2+1/4*exp(2*arctan(a*x))*(a*x+1)/a/c^2/(a^2*x^2+1)

Rubi [A]

time = 0.04, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$\frac{(ax+1)e^{2\text{ArcTan}(ax)}}{4ac^2(a^2x^2+1)} + \frac{e^{2\text{ArcTan}(ax)}}{8ac^2}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^2,x]

[Out] E^(2*ArcTan[a*x])/(8*a*c^2) + (E^(2*ArcTan[a*x])*(1 + a*x))/(4*a*c^2*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_.)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] & & EqQ[d, a^2*c] & & LtQ[p, -1] & & !IntegerQ[I*n] & & NeQ[n^2 + 4*(p + 1)^2, 0] & & IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_.)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] & & EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned} \int \frac{e^{2\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx &= \frac{e^{2\tan^{-1}(ax)}(1+ax)}{4ac^2(1+a^2x^2)} + \frac{\int \frac{e^{2\tan^{-1}(ax)}}{c+a^2cx^2} dx}{4c} \\ &= \frac{e^{2\tan^{-1}(ax)}}{8ac^2} + \frac{e^{2\tan^{-1}(ax)}(1+ax)}{4ac^2(1+a^2x^2)} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.01, size = 55, normalized size = 1.04

$$\frac{(1 - iax)^i(1 + iax)^{-i}(3 + 2ax + a^2x^2)}{8c^2(a + a^3x^2)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^2,x]

[Out] ((1 - I*a*x)^I*(3 + 2*a*x + a^2*x^2))/(8*c^2*(1 + I*a*x)^I*(a + a^3*x^2))

Maple [A]

time = 0.08, size = 40, normalized size = 0.75

method	result	size
gospers	$\frac{e^{2 \arctan(ax)}(a^2x^2 + 2ax + 3)}{8(a^2x^2 + 1)ac^2}$	40

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)

[Out] 1/8*exp(2*arctan(a*x))*(a^2*x^2+2*a*x+3)/(a^2*x^2+1)/a/c^2

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(e^(2*arctan(a*x))/(a^2*c*x^2 + c)^2, x)

Fricas [A]

time = 2.76, size = 40, normalized size = 0.75

$$\frac{(a^2x^2 + 2ax + 3)e^{(2 \arctan(ax))}}{8(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] 1/8*(a^2*x^2 + 2*a*x + 3)*e^(2*arctan(a*x))/(a^3*c^2*x^2 + a*c^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 99 vs. $2(42) = 84$.

time = 1.49, size = 99, normalized size = 1.87

$$\begin{cases} \frac{a^2 x^2 e^{2 \operatorname{atan}(ax)}}{8a^3 c^2 x^2 + 8ac^2} + \frac{2axe^{2 \operatorname{atan}(ax)}}{8a^3 c^2 x^2 + 8ac^2} + \frac{3e^{2 \operatorname{atan}(ax)}}{8a^3 c^2 x^2 + 8ac^2} & \text{for } a \neq 0 \\ \frac{x}{c^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))/(a**2*c*x**2+c)**2,x)

[Out] Piecewise((a**2*x**2*exp(2*atan(a*x))/(8*a**3*c**2*x**2 + 8*a*c**2) + 2*a*x*exp(2*atan(a*x))/(8*a**3*c**2*x**2 + 8*a*c**2) + 3*exp(2*atan(a*x))/(8*a**3*c**2*x**2 + 8*a*c**2), Ne(a, 0)), (x/c**2, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.58, size = 46, normalized size = 0.87

$$\frac{e^{2 \operatorname{atan}(ax)} \left(\frac{3}{8a^3 c^2} + \frac{x}{4a^2 c^2} + \frac{x^2}{8ac^2} \right)}{\frac{1}{a^2} + x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))/(c + a^2*c*x^2)^2,x)

[Out] (exp(2*atan(a*x))*(3/(8*a^3*c^2) + x/(4*a^2*c^2) + x^2/(8*a*c^2)))/(1/a^2 + x^2)

$$3.265 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$$

Optimal. Leaf size=88

$$\frac{3e^{2\text{ArcTan}(ax)}}{40ac^3} + \frac{e^{2\text{ArcTan}(ax)}(1+2ax)}{10ac^3(1+a^2x^2)^2} + \frac{3e^{2\text{ArcTan}(ax)}(1+ax)}{20ac^3(1+a^2x^2)}$$

[Out] 3/40*exp(2*arctan(a*x))/a/c^3+1/10*exp(2*arctan(a*x))*(2*a*x+1)/a/c^3/(a^2*x^2+1)^2+3/20*exp(2*arctan(a*x))*(a*x+1)/a/c^3/(a^2*x^2+1)

Rubi [A]

time = 0.06, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$\frac{3(ax+1)e^{2\text{ArcTan}(ax)}}{20ac^3(a^2x^2+1)} + \frac{(2ax+1)e^{2\text{ArcTan}(ax)}}{10ac^3(a^2x^2+1)^2} + \frac{3e^{2\text{ArcTan}(ax)}}{40ac^3}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^3,x]

[Out] (3*E^(2*ArcTan[a*x]))/(40*a*c^3) + (E^(2*ArcTan[a*x])*(1 + 2*a*x))/(10*a*c^3*(1 + a^2*x^2)^2) + (3*E^(2*ArcTan[a*x])*(1 + a*x))/(20*a*c^3*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2)), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx &= \frac{e^{2 \tan^{-1}(ax)}(1+2ax)}{10ac^3(1+a^2x^2)^2} + \frac{3 \int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{5c} \\
&= \frac{e^{2 \tan^{-1}(ax)}(1+2ax)}{10ac^3(1+a^2x^2)^2} + \frac{3e^{2 \tan^{-1}(ax)}(1+ax)}{20ac^3(1+a^2x^2)} + \frac{3 \int \frac{e^{2 \tan^{-1}(ax)}}{c+a^2cx^2} dx}{20c^2} \\
&= \frac{3e^{2 \tan^{-1}(ax)}}{40ac^3} + \frac{e^{2 \tan^{-1}(ax)}(1+2ax)}{10ac^3(1+a^2x^2)^2} + \frac{3e^{2 \tan^{-1}(ax)}(1+ax)}{20ac^3(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.10, size = 86, normalized size = 0.98

$$\frac{4e^{2\text{ArcTan}(ax)}(1+2ax) + 3(1-iax)^i(1+iax)^{-i}(1+a^2x^2)(3+2ax+a^2x^2)}{40ac^3(1+a^2x^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^3,x]

[Out] (4*E^(2*ArcTan[a*x])*(1 + 2*a*x) + (3*(1 - I*a*x)^I*(1 + a^2*x^2)*(3 + 2*a*x + a^2*x^2))/(1 + I*a*x)^I)/(40*a*c^3*(1 + a^2*x^2)^2)

Maple [A]

time = 0.08, size = 57, normalized size = 0.65

method	result	size
gospers	$\frac{e^{2 \arctan(ax)}(3a^4x^4 + 6a^3x^3 + 12a^2x^2 + 14ax + 13)}{40(a^2x^2 + 1)^2c^3a}$	57

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)

[Out] 1/40*exp(2*arctan(a*x))*(3*a^4*x^4+6*a^3*x^3+12*a^2*x^2+14*a*x+13)/(a^2*x^2+1)^2/c^3/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] integrate($e^{(2*\arctan(ax))}/(a^2*c*x^2 + c)^3, x$)

Fricas [A]

time = 2.51, size = 68, normalized size = 0.77

$$\frac{(3a^4x^4 + 6a^3x^3 + 12a^2x^2 + 14ax + 13)e^{(2\arctan(ax))}}{40(a^5c^3x^4 + 2a^3c^3x^2 + ac^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($\exp(2*\arctan(ax))/(a^2*c*x^2+c)^3, x$, algorithm="fricas")

[Out] $1/40*(3*a^4*x^4 + 6*a^3*x^3 + 12*a^2*x^2 + 14*a*x + 13)*e^{(2*\arctan(ax))}/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)$

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 231 vs. 2(78) = 156.

time = 4.11, size = 231, normalized size = 2.62

$$\begin{cases} \frac{3a^4x^4e^{2\arctan(ax)}}{40a^5c^3x^4+80a^3c^3x^2+40ac^3} + \frac{6a^3x^3e^{2\arctan(ax)}}{40a^5c^3x^4+80a^3c^3x^2+40ac^3} + \frac{12a^2x^2e^{2\arctan(ax)}}{40a^5c^3x^4+80a^3c^3x^2+40ac^3} + \frac{14axe^{2\arctan(ax)}}{40a^5c^3x^4+80a^3c^3x^2+40ac^3} + \frac{13e^{2\arctan(ax)}}{40a^5c^3x^4+80a^3c^3x^2+40ac^3} & \text{for } a \neq 0 \\ \frac{x}{c^3} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($\exp(2*\atan(ax))/(a**2*c*x**2+c)**3, x$)

[Out] Piecewise(($3*a**4*x**4*\exp(2*\atan(ax))/(40*a**5*c**3*x**4 + 80*a**3*c**3*x**2 + 40*a*c**3) + 6*a**3*x**3*\exp(2*\atan(ax))/(40*a**5*c**3*x**4 + 80*a**3*c**3*x**2 + 40*a*c**3) + 12*a**2*x**2*\exp(2*\atan(ax))/(40*a**5*c**3*x**4 + 80*a**3*c**3*x**2 + 40*a*c**3) + 14*a*x*\exp(2*\atan(ax))/(40*a**5*c**3*x**4 + 80*a**3*c**3*x**2 + 40*a*c**3) + 13*\exp(2*\atan(ax))/(40*a**5*c**3*x**4 + 80*a**3*c**3*x**2 + 40*a*c**3)$, Ne(a, 0)), (x/c**3, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($\exp(2*\arctan(ax))/(a^2*c*x^2+c)^3, x$, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.62, size = 79, normalized size = 0.90

$$\frac{3e^{2\arctan(ax)}}{40ac^3} + \frac{3e^{2\arctan(ax)}(ax+1)}{20ac^3(a^2x^2+1)} + \frac{e^{2\arctan(ax)}(2ax+1)}{10ac^3(a^2x^2+1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int($\exp(2*\atan(ax))/(c + a^2*c*x^2)^3, x$)

[Out] $(3*\exp(2*\atan(ax)))/(40*a*c^3) + (3*\exp(2*\atan(ax))*(ax + 1))/(20*a*c^3*(a^2*x^2 + 1)) + (\exp(2*\atan(ax))*(2*ax + 1))/(10*a*c^3*(a^2*x^2 + 1)^2)$

$$3.266 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$$

Optimal. Leaf size=123

$$\frac{9e^{2\text{ArcTan}(ax)}}{160ac^4} + \frac{e^{2\text{ArcTan}(ax)}(1+3ax)}{20ac^4(1+a^2x^2)^3} + \frac{3e^{2\text{ArcTan}(ax)}(1+2ax)}{40ac^4(1+a^2x^2)^2} + \frac{9e^{2\text{ArcTan}(ax)}(1+ax)}{80ac^4(1+a^2x^2)}$$

[Out] 9/160*exp(2*arctan(a*x))/a/c^4+1/20*exp(2*arctan(a*x))*(3*a*x+1)/a/c^4/(a^2*x^2+1)^3+3/40*exp(2*arctan(a*x))*(2*a*x+1)/a/c^4/(a^2*x^2+1)^2+9/80*exp(2*arctan(a*x))*(a*x+1)/a/c^4/(a^2*x^2+1)

Rubi [A]

time = 0.09, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$\frac{9(ax+1)e^{2\text{ArcTan}(ax)}}{80ac^4(a^2x^2+1)} + \frac{3(2ax+1)e^{2\text{ArcTan}(ax)}}{40ac^4(a^2x^2+1)^2} + \frac{(3ax+1)e^{2\text{ArcTan}(ax)}}{20ac^4(a^2x^2+1)^3} + \frac{9e^{2\text{ArcTan}(ax)}}{160ac^4}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^4,x]

[Out] (9*E^(2*ArcTan[a*x]))/(160*a*c^4) + (E^(2*ArcTan[a*x])*(1 + 3*a*x))/(20*a*c^4*(1 + a^2*x^2)^3) + (3*E^(2*ArcTan[a*x])*(1 + 2*a*x))/(40*a*c^4*(1 + a^2*x^2)^2) + (9*E^(2*ArcTan[a*x])*(1 + a*x))/(80*a*c^4*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^4} dx &= \frac{e^{2 \tan^{-1}(ax)}(1+3ax)}{20ac^4(1+a^2x^2)^3} + \frac{3 \int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx}{4c} \\
&= \frac{e^{2 \tan^{-1}(ax)}(1+3ax)}{20ac^4(1+a^2x^2)^3} + \frac{3e^{2 \tan^{-1}(ax)}(1+2ax)}{40ac^4(1+a^2x^2)^2} + \frac{9 \int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{20c^2} \\
&= \frac{e^{2 \tan^{-1}(ax)}(1+3ax)}{20ac^4(1+a^2x^2)^3} + \frac{3e^{2 \tan^{-1}(ax)}(1+2ax)}{40ac^4(1+a^2x^2)^2} + \frac{9e^{2 \tan^{-1}(ax)}(1+ax)}{80ac^4(1+a^2x^2)} + \frac{9 \int \frac{e^{2 \tan^{-1}(ax)}}{c+a^2cx^2} dx}{80c^3} \\
&= \frac{9e^{2 \tan^{-1}(ax)}}{160ac^4} + \frac{e^{2 \tan^{-1}(ax)}(1+3ax)}{20ac^4(1+a^2x^2)^3} + \frac{3e^{2 \tan^{-1}(ax)}(1+2ax)}{40ac^4(1+a^2x^2)^2} + \frac{9e^{2 \tan^{-1}(ax)}(1+ax)}{80ac^4(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.19, size = 122, normalized size = 0.99

$$\frac{8ce^{2\text{ArcTan}(ax)}(1+3ax) + 3(c+a^2cx^2)(4e^{2\text{ArcTan}(ax)}(1+2ax) + 3(1-iax)^i(1+iax)^{-i}(-i+ax)(i+ax)(3+2ax+a^2x^2))}{160ac^2(c+a^2cx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^4, x]

[Out] (8*c*E^(2*ArcTan[a*x])*(1 + 3*a*x) + 3*(c + a^2*c*x^2)*(4*E^(2*ArcTan[a*x])*(1 + 2*a*x) + (3*(1 - I*a*x)^I*(-I + a*x)*(I + a*x)*(3 + 2*a*x + a^2*x^2))/(1 + I*a*x^I)))/(160*a*c^2*(c + a^2*c*x^2)^3)

Maple [A]

time = 0.08, size = 73, normalized size = 0.59

method	result	size
gosper	$\frac{e^{2 \arctan(ax)}(9a^6x^6 + 18a^5x^5 + 45a^4x^4 + 60a^3x^3 + 75a^2x^2 + 66ax + 47)}{160(a^2x^2 + 1)^3c^4a}$	73

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^4, x, method=_RETURNVERBOSE)

[Out] 1/160*exp(2*arctan(a*x))*(9*a^6*x^6+18*a^5*x^5+45*a^4*x^4+60*a^3*x^3+75*a^2*x^2+66*a*x+47)/(a^2*x^2+1)^3/c^4/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="maxima")

[Out] integrate(e^(2*arctan(a*x))/(a^2*c*x^2 + c)^4, x)

Fricas [A]

time = 2.74, size = 95, normalized size = 0.77

$$\frac{(9a^6x^6 + 18a^5x^5 + 45a^4x^4 + 60a^3x^3 + 75a^2x^2 + 66ax + 47)e^{(2 \arctan(ax))}}{160(a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="fricas")

[Out] 1/160*(9*a^6*x^6 + 18*a^5*x^5 + 45*a^4*x^4 + 60*a^3*x^3 + 75*a^2*x^2 + 66*a*x + 47)*e^(2*arctan(a*x))/(a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 410 vs. 2(112) = 224.

time = 10.73, size = 410, normalized size = 3.33

$$\left\{ \begin{array}{l} \frac{9a^6x^6 + 18a^5x^5 + 45a^4x^4 + 60a^3x^3 + 75a^2x^2 + 66ax + 47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \\ \frac{18a^5x^5 + 45a^4x^4 + 60a^3x^3 + 75a^2x^2 + 66ax + 47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \\ \frac{45a^4x^4 + 60a^3x^3 + 75a^2x^2 + 66ax + 47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \\ \frac{60a^3x^3 + 75a^2x^2 + 66ax + 47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \\ \frac{75a^2x^2 + 66ax + 47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \\ \frac{66ax + 47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \\ \frac{47}{160a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4} e^{2 \arctan(ax)} \end{array} \right. \text{ for } a \neq 0 \\ \frac{1}{c^4} \text{ otherwise}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))/(a**2*c*x**2+c)**4,x)

[Out] Piecewise((9*a**6*x**6*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4) + 18*a**5*x**5*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4) + 45*a**4*x**4*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4) + 60*a**3*x**3*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4) + 75*a**2*x**2*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4) + 66*a*x*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4) + 47*exp(2*atan(a*x))/(160*a**7*c**4*x**6 + 480*a**5*c**4*x**4 + 480*a**3*c**4*x**2 + 160*a*c**4), N e(a, 0)), (x/c**4, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.67, size = 111, normalized size = 0.90

$$\frac{9 e^{2 \operatorname{atan}(a x)}}{160 a c^4} + \frac{9 e^{2 \operatorname{atan}(a x)} (a x + 1)}{80 a c^4 (a^2 x^2 + 1)} + \frac{3 e^{2 \operatorname{atan}(a x)} (2 a x + 1)}{40 a c^4 (a^2 x^2 + 1)^2} + \frac{e^{2 \operatorname{atan}(a x)} (3 a x + 1)}{20 a c^4 (a^2 x^2 + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(2*atan(a*x))/(c + a^2*c*x^2)^4,x)`

[Out] `(9*exp(2*atan(a*x)))/(160*a*c^4) + (9*exp(2*atan(a*x))*(a*x + 1))/(80*a*c^4*(a^2*x^2 + 1)) + (3*exp(2*atan(a*x))*(2*a*x + 1))/(40*a*c^4*(a^2*x^2 + 1)^2) + (exp(2*atan(a*x))*(3*a*x + 1))/(20*a*c^4*(a^2*x^2 + 1)^3)`

$$3.267 \quad \int e^{2\text{ArcTan}(ax)} (c + a^2cx^2)^{3/2} dx$$

Optimal. Leaf size=88

$$\frac{\left(\frac{2}{29} + \frac{5i}{29}\right) 2^{\frac{5}{2}-i} c(1-iax)^{\frac{5}{2}+i} \sqrt{c+a^2cx^2} {}_2F_1\left(-\frac{3}{2}+i, \frac{5}{2}+i; \frac{7}{2}+i; \frac{1}{2}(1-iax)\right)}{a\sqrt{1+a^2x^2}}$$

[Out] (2/29+5/29*I)*2^(5/2-I)*c*(1-I*a*x)^(5/2+I)*hypergeom([5/2+I, -3/2+I], [7/2+I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^(1/2)/a/(a^2*x^2+1)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{2}{29} + \frac{5i}{29}\right) 2^{\frac{5}{2}-i} c(1-iax)^{\frac{5}{2}+i} \sqrt{a^2cx^2+c} {}_2F_1\left(-\frac{3}{2}+i, \frac{5}{2}+i; \frac{7}{2}+i; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2),x]

[Out] ((2/29 + (5*I)/29)*2^(5/2 - I)*c*(1 - I*a*x)^(5/2 + I)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-3/2 + I, 5/2 + I, 7/2 + I, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{2 \tan^{-1}(ax)} (c + a^2 cx^2)^{3/2} dx &= \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int e^{2 \tan^{-1}(ax)} (1 + a^2 x^2)^{3/2} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int (1 - iax)^{\frac{3}{2}+i} (1 + iax)^{\frac{3}{2}-i} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\left(\frac{2}{29} + \frac{5i}{29}\right) 2^{\frac{5}{2}-i} c (1 - iax)^{\frac{5}{2}+i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{3}{2} + i, \frac{5}{2} + i; \frac{7}{2} + i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 88, normalized size = 1.00

$$\frac{\left(\frac{2}{29} + \frac{5i}{29}\right) 2^{\frac{5}{2}-i} c (1 - iax)^{\frac{5}{2}+i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{3}{2} + i, \frac{5}{2} + i; \frac{7}{2} + i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2), x]

[Out] ((2/29 + (5*I)/29)*2^(5/2 - I)*c*(1 - I*a*x)^(5/2 + I)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-3/2 + I, 5/2 + I, 7/2 + I, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{2 \arctan(ax)} (a^2 cx^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(3/2), x)

[Out] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(3/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(3/2), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(3/2)*e^(2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(3/2)*e^(2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^{\frac{3}{2}} e^{2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))*(a**2*c*x**2+c)**(3/2),x)

[Out] Integral((c*(a**2*x**2 + 1))**(3/2)*exp(2*atan(a*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{2\operatorname{atan}(ax)} (ca^2x^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^(3/2),x)

[Out] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^(3/2), x)

3.268 $\int e^{2\text{ArcTan}(ax)} \sqrt{c + a^2cx^2} dx$

Optimal. Leaf size=87

$$\frac{\left(\frac{2}{13} + \frac{3i}{13}\right) 2^{\frac{3}{2}-i} (1-iax)^{\frac{3}{2}+i} \sqrt{c+a^2cx^2} {}_2F_1\left(-\frac{1}{2}+i, \frac{3}{2}+i; \frac{5}{2}+i; \frac{1}{2}(1-iax)\right)}{a\sqrt{1+a^2x^2}}$$

[Out] (2/13+3/13*I)*2^(3/2-I)*(1-I*a*x)^(3/2+I)*hypergeom([-1/2+I, 3/2+I], [5/2+I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^(1/2)/a/(a^2*x^2+1)^(1/2)

Rubi [A]

time = 0.05, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$,

Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{2}{13} + \frac{3i}{13}\right) 2^{\frac{3}{2}-i} (1-iax)^{\frac{3}{2}+i} \sqrt{a^2cx^2+c} {}_2F_1\left(-\frac{1}{2}+i, \frac{3}{2}+i; \frac{5}{2}+i; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])*Sqrt[c + a^2*c*x^2], x]

[Out] (((2/13 + (3*I)/13)*2^(3/2 - I)*(1 - I*a*x)^(3/2 + I)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-1/2 + I, 3/2 + I, 5/2 + I, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2]))

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[(((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int e^{2 \tan^{-1}(ax)} \sqrt{c + a^2 cx^2} dx &= \frac{\sqrt{c + a^2 cx^2} \int e^{2 \tan^{-1}(ax)} \sqrt{1 + a^2 x^2} dx}{\sqrt{1 + a^2 x^2}} \\
&= \frac{\sqrt{c + a^2 cx^2} \int (1 - iax)^{\frac{1}{2}+i} (1 + iax)^{\frac{1}{2}-i} dx}{\sqrt{1 + a^2 x^2}} \\
&= \frac{\left(\frac{2}{13} + \frac{3i}{13}\right) 2^{\frac{3}{2}-i} (1 - iax)^{\frac{3}{2}+i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} + i, \frac{3}{2} + i; \frac{5}{2} + i; \frac{1}{2}(1 - iax)\right)}{a \sqrt{1 + a^2 x^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 87, normalized size = 1.00

$$\frac{\left(\frac{2}{13} + \frac{3i}{13}\right) 2^{\frac{3}{2}-i} (1 - iax)^{\frac{3}{2}+i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} + i, \frac{3}{2} + i; \frac{5}{2} + i; \frac{1}{2}(1 - iax)\right)}{a \sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(2*ArcTan[a*x])*Sqrt[c + a^2*c*x^2],x]`

```
[Out] ((2/13 + (3*I)/13)*2^(3/2 - I)*(1 - I*a*x)^(3/2 + I)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-1/2 + I, 3/2 + I, 5/2 + I, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{2 \arctan(ax)} \sqrt{a^2 cx^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x)``[Out] int(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")``[Out] integrate(sqrt(a^2*c*x^2 + c)*e^(2*arctan(a*x)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(a^2*c*x^2 + c)*e^(2*arctan(a*x)), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c(a^2x^2 + 1)} e^{2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*atan(a*x))*(a**2*c*x**2+c)**(1/2),x)
```

```
[Out] Integral(sqrt(c*(a**2*x**2 + 1))*exp(2*atan(a*x)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{2\operatorname{atan}(ax)} \sqrt{ca^2x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^(1/2),x)
```

```
[Out] int(exp(2*atan(a*x))*(c + a^2*c*x^2)^(1/2), x)
```

$$3.269 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\left(\frac{2}{5} + \frac{i}{5}\right) 2^{\frac{1}{2}-i} (1-iax)^{\frac{1}{2}+i} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2} + i, \frac{1}{2} + i; \frac{3}{2} + i; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

[Out] (2/5+1/5*I)*2^(1/2-I)*(1-I*a*x)^(1/2+I)*hypergeom([1/2+I, 1/2+I],[3/2+I],1/2-1/2*I*a*x)*(a^2*x^2+1)^(1/2)/a/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{2}{5} + \frac{i}{5}\right) 2^{\frac{1}{2}-i} (1-iax)^{\frac{1}{2}+i} \sqrt{a^2x^2+1} {}_2F_1\left(\frac{1}{2} + i, \frac{1}{2} + i; \frac{3}{2} + i; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/Sqrt[c + a^2*c*x^2],x]

[Out] ((2/5 + I/5)*2^(1/2 - I)*(1 - I*a*x)^(1/2 + I)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 + I, 1/2 + I, 3/2 + I, (1 - I*a*x)/2])/(a*Sqrt[c + a^2*c*x^2])

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || ! (RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{2 \tan^{-1}(ax)}}{\sqrt{c+a^2cx^2}} dx &= \frac{\sqrt{1+a^2x^2} \int \frac{e^{2 \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx}{\sqrt{c+a^2cx^2}} \\ &= \frac{\sqrt{1+a^2x^2} \int (1-iax)^{-\frac{1}{2}+i} (1+iax)^{-\frac{1}{2}-i} dx}{\sqrt{c+a^2cx^2}} \\ &= \frac{\left(\frac{2}{5} + \frac{i}{5}\right) 2^{\frac{1}{2}-i} (1-iax)^{\frac{1}{2}+i} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}+i, \frac{1}{2}+i; \frac{3}{2}+i; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 87, normalized size = 1.00

$$\frac{\left(\frac{2}{5} + \frac{i}{5}\right) 2^{\frac{1}{2}-i} (1-iax)^{\frac{1}{2}+i} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}+i, \frac{1}{2}+i; \frac{3}{2}+i; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]

[Out] ((2/5 + I/5)*2^(1/2 - I)*(1 - I*a*x)^(1/2 + I)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 + I, 1/2 + I, 3/2 + I, (1 - I*a*x)/2])/(a*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{2 \arctan(ax)}}{\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(e^(2*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")``[Out] integral(e^(2*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{2 \operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(2*atan(a*x))/(a**2*c*x**2+c)**(1/2),x)``[Out] Integral(exp(2*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")``[Out] sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{2 \operatorname{atan}(ax)}}{\sqrt{c a^2 x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(2*atan(a*x))/(c + a^2*c*x^2)^(1/2),x)``[Out] int(exp(2*atan(a*x))/(c + a^2*c*x^2)^(1/2), x)`

$$3.270 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=37

$$\frac{e^{2\text{ArcTan}(ax)}(2+ax)}{5ac\sqrt{c+a^2cx^2}}$$

[Out] 1/5*exp(2*arctan(a*x))*(a*x+2)/a/c/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 37, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {5177}

$$\frac{(ax+2)e^{2\text{ArcTan}(ax)}}{5ac\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]

[Out] (E^(2*ArcTan[a*x])*(2 + a*x))/(5*a*c*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.) + (d_.)*(x_)^2)^(3/2), x_Symbol] :>
Simp[(n + a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 1)*Sqrt[c + d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rubi steps

$$\int \frac{e^{2\tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx = \frac{e^{2\tan^{-1}(ax)}(2+ax)}{5ac\sqrt{c+a^2cx^2}}$$

Mathematica [A]

time = 0.01, size = 37, normalized size = 1.00

$$\frac{e^{2\text{ArcTan}(ax)}(2+ax)}{5ac\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]

[Out] (E^(2*ArcTan[a*x])*(2 + a*x))/(5*a*c*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.06, size = 39, normalized size = 1.05

method	result	size
gosper	$\frac{(a^2x^2+1)(ax+2)e^{2\arctan(ax)}}{5a(a^2cx^2+c)^{\frac{3}{2}}}$	39

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)
```

```
[Out] 1/5*(a^2*x^2+1)*(a*x+2)*exp(2*arctan(a*x))/a/(a^2*c*x^2+c)^(3/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate(e^(2*arctan(a*x))/(a^2*c*x^2 + c)^(3/2), x)
```

Fricas [A]

time = 1.76, size = 44, normalized size = 1.19

$$\frac{\sqrt{a^2cx^2 + c} (ax + 2)e^{2\arctan(ax)}}{5(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")
```

```
[Out] 1/5*sqrt(a^2*c*x^2 + c)*(a*x + 2)*e^(2*arctan(a*x))/(a^3*c^2*x^2 + a*c^2)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{2\operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*atan(a*x))/(a**2*c*x**2+c)**(3/2),x)
```

```
[Out] Integral(exp(2*atan(a*x))/(c*(a**2*x**2 + 1))**(3/2), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [B]

time = 0.61, size = 35, normalized size = 0.95

$$\frac{e^{2\operatorname{atan}(ax)} \left(\frac{x}{5c} + \frac{2}{5ac} \right)}{\sqrt{ca^2x^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(2*atan(a*x))/(c + a^2*c*x^2)^(3/2),x)`

[Out] `(exp(2*atan(a*x))*(x/(5*c) + 2/(5*a*c)))/(c + a^2*c*x^2)^(1/2)`

$$3.271 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$$

Optimal. Leaf size=76

$$\frac{e^{2\text{ArcTan}(ax)}(2+3ax)}{13ac(c+a^2cx^2)^{3/2}} + \frac{6e^{2\text{ArcTan}(ax)}(2+ax)}{65ac^2\sqrt{c+a^2cx^2}}$$

[Out] 1/13*exp(2*arctan(a*x))*(3*a*x+2)/a/c/(a^2*c*x^2+c)^(3/2)+6/65*exp(2*arctan(a*x))*(a*x+2)/a/c^2/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 76, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {5178, 5177}

$$\frac{6(ax+2)e^{2\text{ArcTan}(ax)}}{65ac^2\sqrt{a^2cx^2+c}} + \frac{(3ax+2)e^{2\text{ArcTan}(ax)}}{13ac(a^2cx^2+c)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^(5/2), x]

[Out] (E^(2*ArcTan[a*x])*(2 + 3*a*x))/(13*a*c*(c + a^2*c*x^2)^(3/2)) + (6*E^(2*ArcTan[a*x])*(2 + a*x))/(65*a*c^2*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n + a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 1)*Sqrt[c + d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\int \frac{e^{2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{5/2}} dx = \frac{e^{2 \tan^{-1}(ax)}(2 + 3ax)}{13ac(c + a^2 cx^2)^{3/2}} + \frac{6 \int \frac{e^{2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx}{13c}$$

$$= \frac{e^{2 \tan^{-1}(ax)}(2 + 3ax)}{13ac(c + a^2 cx^2)^{3/2}} + \frac{6e^{2 \tan^{-1}(ax)}(2 + ax)}{65ac^2 \sqrt{c + a^2 cx^2}}$$

Mathematica [A]

time = 0.02, size = 62, normalized size = 0.82

$$\frac{e^{2 \operatorname{ArcTan}(ax)}(22 + 21ax + 12a^2 x^2 + 6a^3 x^3)}{65c^2 (a + a^3 x^2) \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^(5/2), x]

[Out] (E^(2*ArcTan[a*x])*(22 + 21*a*x + 12*a^2*x^2 + 6*a^3*x^3))/(65*c^2*(a + a^3*x^2)*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.10, size = 56, normalized size = 0.74

method	result	size
gosper	$\frac{(a^2 x^2 + 1)(6a^3 x^3 + 12a^2 x^2 + 21ax + 22)e^{2 \arctan(ax)}}{65a(a^2 cx^2 + c)^{5/2}}$	56

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, method=_RETURNVERBOSE)

[Out] 1/65*(a^2*x^2+1)*(6*a^3*x^3+12*a^2*x^2+21*a*x+22)*exp(2*arctan(a*x))/a/(a^2*c*x^2+c)^(5/2)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, algorithm="maxima")

[Out] integrate(e^(2*arctan(a*x))/(a^2*c*x^2 + c)^(5/2), x)

Fricas [A]

time = 3.27, size = 72, normalized size = 0.95

$$\frac{(6a^3x^3 + 12a^2x^2 + 21ax + 22)\sqrt{a^2cx^2 + c}e^{(2\arctan(ax))}}{65(a^5c^3x^4 + 2a^3c^3x^2 + ac^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="fricas")

[Out] 1/65*(6*a^3*x^3 + 12*a^2*x^2 + 21*a*x + 22)*sqrt(a^2*c*x^2 + c)*e^(2*arctan(a*x))/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{2\operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*atan(a*x))/(a**2*c*x**2+c)**(5/2),x)

[Out] Integral(exp(2*atan(a*x))/(c*(a**2*x**2 + 1))**(5/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.22, size = 80, normalized size = 1.05

$$\frac{e^{2\operatorname{atan}(ax)} \left(\frac{22}{65a^3c^2} + \frac{6x^3}{65c^2} + \frac{21x}{65a^2c^2} + \frac{12x^2}{65ac^2} \right)}{\frac{\sqrt{ca^2x^2 + c}}{a^2} + x^2 \sqrt{ca^2x^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(2*atan(a*x))/(c + a^2*c*x^2)^(5/2),x)

[Out] (exp(2*atan(a*x))*(22/(65*a^3*c^2) + (6*x^3)/(65*c^2) + (21*x)/(65*a^2*c^2) + (12*x^2)/(65*a*c^2)))/((c + a^2*c*x^2)^(1/2)/a^2 + x^2*(c + a^2*c*x^2)^(1/2))

$$3.272 \quad \int \frac{e^{2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$$

Optimal. Leaf size=114

$$\frac{e^{2\text{ArcTan}(ax)}(2+5ax)}{29ac(c+a^2cx^2)^{5/2}} + \frac{20e^{2\text{ArcTan}(ax)}(2+3ax)}{377ac^2(c+a^2cx^2)^{3/2}} + \frac{24e^{2\text{ArcTan}(ax)}(2+ax)}{377ac^3\sqrt{c+a^2cx^2}}$$

[Out] 1/29*exp(2*arctan(a*x))*(5*a*x+2)/a/c/(a^2*c*x^2+c)^(5/2)+20/377*exp(2*arctan(a*x))*(3*a*x+2)/a/c^2/(a^2*c*x^2+c)^(3/2)+24/377*exp(2*arctan(a*x))*(a*x+2)/a/c^3/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 114, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {5178, 5177}

$$\frac{24(ax+2)e^{2\text{ArcTan}(ax)}}{377ac^3\sqrt{a^2cx^2+c}} + \frac{20(3ax+2)e^{2\text{ArcTan}(ax)}}{377ac^2(a^2cx^2+c)^{3/2}} + \frac{(5ax+2)e^{2\text{ArcTan}(ax)}}{29ac(a^2cx^2+c)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^(7/2), x]

[Out] (E^(2*ArcTan[a*x])*(2 + 5*a*x))/(29*a*c*(c + a^2*c*x^2)^(5/2)) + (20*E^(2*ArcTan[a*x])*(2 + 3*a*x))/(377*a*c^2*(c + a^2*c*x^2)^(3/2)) + (24*E^(2*ArcTan[a*x])*(2 + a*x))/(377*a*c^3*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n + a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 1)*Sqrt[c + d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\begin{aligned}
\int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^{7/2}} dx &= \frac{e^{2 \tan^{-1}(ax)}(2+5ax)}{29ac(c+a^2cx^2)^{5/2}} + \frac{20 \int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^{5/2}} dx}{29c} \\
&= \frac{e^{2 \tan^{-1}(ax)}(2+5ax)}{29ac(c+a^2cx^2)^{5/2}} + \frac{20e^{2 \tan^{-1}(ax)}(2+3ax)}{377ac^2(c+a^2cx^2)^{3/2}} + \frac{120 \int \frac{e^{2 \tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx}{377c^2} \\
&= \frac{e^{2 \tan^{-1}(ax)}(2+5ax)}{29ac(c+a^2cx^2)^{5/2}} + \frac{20e^{2 \tan^{-1}(ax)}(2+3ax)}{377ac^2(c+a^2cx^2)^{3/2}} + \frac{24e^{2 \tan^{-1}(ax)}(2+ax)}{377ac^3\sqrt{c+a^2cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 81, normalized size = 0.71

$$\frac{e^{2 \operatorname{ArcTan}(ax)}(114 + 149ax + 136a^2x^2 + 108a^3x^3 + 48a^4x^4 + 24a^5x^5)}{377ac^3(1 + a^2x^2)^2 \sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(2*ArcTan[a*x])/(c + a^2*c*x^2)^(7/2), x]`

```
[Out] (E^(2*ArcTan[a*x])*(114 + 149*a*x + 136*a^2*x^2 + 108*a^3*x^3 + 48*a^4*x^4 + 24*a^5*x^5))/(377*a*c^3*(1 + a^2*x^2)^2*Sqrt[c + a^2*c*x^2])
```

Maple [A]

time = 0.07, size = 72, normalized size = 0.63

method	result	size
gospers	$\frac{(a^2x^2+1)(24a^5x^5+48a^4x^4+108a^3x^3+136a^2x^2+149ax+114)e^{2 \arctan(ax)}}{377a(a^2cx^2+c)^{\frac{7}{2}}}$	72

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, method=_RETURNVERBOSE)`

```
[Out] 1/377*(a^2*x^2+1)*(24*a^5*x^5+48*a^4*x^4+108*a^3*x^3+136*a^2*x^2+149*a*x+114)*exp(2*arctan(a*x))/a/(a^2*c*x^2+c)^(7/2)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, algorithm="maxima")`

[Out] integrate($e^{(2*\arctan(ax))}/(a^2*c*x^2 + c)^{(7/2)}$, x)

Fricas [A]

time = 2.10, size = 99, normalized size = 0.87

$$\frac{(24 a^5 x^5 + 48 a^4 x^4 + 108 a^3 x^3 + 136 a^2 x^2 + 149 a x + 114) \sqrt{a^2 c x^2 + c} e^{(2 \arctan(ax))}}{377 (a^7 c^4 x^6 + 3 a^5 c^4 x^4 + 3 a^3 c^4 x^2 + a c^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($\exp(2*\arctan(ax))/(a^2*c*x^2+c)^{(7/2)}$, x, algorithm="fricas")

[Out] $1/377*(24*a^5*x^5 + 48*a^4*x^4 + 108*a^3*x^3 + 136*a^2*x^2 + 149*a*x + 114) * \sqrt{a^2*c*x^2 + c} * e^{(2*\arctan(ax))} / (a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($\exp(2*\operatorname{atan}(ax))/(a**2*c*x**2+c)**(7/2)$, x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate($\exp(2*\arctan(ax))/(a^2*c*x^2+c)^{(7/2)}$, x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.69, size = 122, normalized size = 1.07

$$\frac{e^{2 \operatorname{atan}(ax)} \left(\frac{114}{377 a^5 c^3} + \frac{24 x^5}{377 c^3} + \frac{149 x}{377 a^4 c^3} + \frac{48 x^4}{377 a c^3} + \frac{108 x^3}{377 a^2 c^3} + \frac{136 x^2}{377 a^3 c^3} \right)}{\frac{\sqrt{c a^2 x^2 + c}}{a^4} + x^4 \sqrt{c a^2 x^2 + c} + \frac{2 x^2 \sqrt{c a^2 x^2 + c}}{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int($\exp(2*\operatorname{atan}(ax))/(c + a^2*c*x^2)^{(7/2)}$, x)

[Out] $(\exp(2*\operatorname{atan}(ax))*(114/(377*a^5*c^3) + (24*x^5)/(377*c^3) + (149*x)/(377*a^4*c^3) + (48*x^4)/(377*a*c^3) + (108*x^3)/(377*a^2*c^3) + (136*x^2)/(377*a^3*c^3)))/((c + a^2*c*x^2)^{(1/2)}/a^4 + x^4*(c + a^2*c*x^2)^{(1/2)} + (2*x^2*(c + a^2*c*x^2)^{(1/2)})/a^2)$

3.273 $\int e^{-\text{ArcTan}(ax)}(c + a^2cx^2)^p dx$

Optimal. Leaf size=101

$$\frac{2^{(1+\frac{i}{2})+p}(1-iax)^{(1-\frac{i}{2})+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1(-\frac{i}{2}-p, (1-\frac{i}{2})+p; (2-\frac{i}{2})+p; \frac{1}{2}(1-iax))}{a((-1-2i)-2ip)}$$

[Out] $2^{(1+1/2*I+p)}*(1-I*a*x)^{(1-1/2*I+p)}*(a^2*c*x^2+c)^p*\text{hypergeom}([-1/2*I-p, 1-1/2*I+p], [2-1/2*I+p], 1/2-1/2*I*a*x)/a/(-1-2*I-2*I*p)/((a^2*x^2+1)^p)$

Rubi [A]

time = 0.06, antiderivative size = 101, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5184, 5181, 71}

$$\frac{2^{p+(1+\frac{i}{2})}(1-iax)^{p+(1-\frac{i}{2})}(a^2x^2+1)^{-p}(a^2cx^2+c)^p {}_2F_1(-p-\frac{i}{2}, p+(1-\frac{i}{2}); p+(2-\frac{i}{2}); \frac{1}{2}(1-iax))}{a(-2ip-(1+2i))}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)^p/E^{\text{ArcTan}[a*x]}, x]$

[Out] $(2^{((1 + I/2) + p)}*(1 - I*a*x)^{((1 - I/2) + p)}*(c + a^2*c*x^2)^p*\text{Hypergeometric2F1}[-1/2*I - p, (1 - I/2) + p, (2 - I/2) + p, (1 - I*a*x)/2])/ (a*((-1 - 2*I) - (2*I)*p)*(1 + a^2*x^2)^p)$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+)^2)^{(n_+)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] :> \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] :> \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int e^{-\tan^{-1}(ax)}(c+a^2cx^2)^p dx &= \left((1+a^2x^2)^{-p} (c+a^2cx^2)^p \right) \int e^{-\tan^{-1}(ax)}(1+a^2x^2)^p dx \\
&= \left((1+a^2x^2)^{-p} (c+a^2cx^2)^p \right) \int (1-iax)^{-\frac{i}{2}+p}(1+iax)^{\frac{i}{2}+p} dx \\
&= \frac{2^{(1+\frac{i}{2})+p}(1-iax)^{(1-\frac{i}{2})+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1\left(-\frac{i}{2}-p, (1-\frac{i}{2})+p; (1-\frac{i}{2})+p; \frac{1}{2}(1-iax)\right)}{a((-1-2i)-2ip)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 102, normalized size = 1.01

$$\frac{i2^{\frac{i}{2}+p}(1-iax)^{(1-\frac{i}{2})+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1\left(-\frac{i}{2}-p, (1-\frac{i}{2})+p; (2-\frac{i}{2})+p; \frac{1}{2}(1-iax)\right)}{a\left((1-\frac{i}{2})+p\right)}$$

Antiderivative was successfully verified.

`[In] Integrate[(c + a^2*c*x^2)^p/E^ArcTan[a*x], x]`

```
[Out] (I*2^(I/2 + p)*(1 - I*a*x)^((1 - I/2) + p)*(c + a^2*c*x^2)^p*Hypergeometric
2F1[-1/2*I - p, (1 - I/2) + p, (2 - I/2) + p, (1 - I*a*x)/2])/(a*((1 - I/2)
+ p)*(1 + a^2*x^2)^p)
```

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int (a^2cx^2 + c)^p e^{-\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a^2*c*x^2+c)^p/exp(arctan(a*x)), x)``[Out] int((a^2*c*x^2+c)^p/exp(arctan(a*x)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((a^2*c*x^2+c)^p/exp(arctan(a*x)), x, algorithm="maxima")``[Out] integrate((a^2*c*x^2 + c)^p*e^(-arctan(a*x)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^p/exp(arctan(a*x)),x, algorithm="fricas")
```

```
[Out] integral((a^2*c*x^2 + c)^p*e^(-arctan(a*x)), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^p e^{-\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a**2*c*x**2+c)**p/exp(atan(a*x)),x)
```

```
[Out] Integral((c*(a**2*x**2 + 1))**p*exp(-atan(a*x)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^p/exp(arctan(a*x)),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{-\operatorname{atan}(ax)} (ca^2x^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-atan(a*x))*(c + a^2*c*x^2)^p,x)
```

```
[Out] int(exp(-atan(a*x))*(c + a^2*c*x^2)^p, x)
```


3.274 $\int e^{-\text{ArcTan}(ax)}(c + a^2cx^2)^2 dx$

Optimal. Leaf size=63

$$-\frac{\left(\frac{1}{37} - \frac{6i}{37}\right) 2^{3+\frac{i}{2}} c^2 (1 - iax)^{3-\frac{i}{2}} {}_2F_1\left(-2 - \frac{i}{2}, 3 - \frac{i}{2}; 4 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

[Out] $(-1/37+6/37*I)*2^{(3+1/2*I)}*c^2*(1-I*a*x)^{(3-1/2*I)}*\text{hypergeom}([3-1/2*I, -2-1/2*I], [4-1/2*I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.03, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5181, 71}

$$-\frac{\left(\frac{1}{37} - \frac{6i}{37}\right) 2^{3+\frac{i}{2}} c^2 (1 - iax)^{3-\frac{i}{2}} {}_2F_1\left(-2 - \frac{i}{2}, 3 - \frac{i}{2}; 4 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)^2/E^{\text{ArcTan}[a*x]}, x]$

[Out] $((-1/37 + (6*I)/37)*2^{(3 + I/2)}*c^2*(1 - I*a*x)^{(3 - I/2)}*\text{Hypergeometric2F1}[-2 - I/2, 3 - I/2, 4 - I/2, (1 - I*a*x)/2])/a$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[a_+)*(x_+)]*(n_+)}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] := \text{Dist}[c^{(p)}, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{-\tan^{-1}(ax)}(c + a^2cx^2)^2 dx &= c^2 \int (1 - iax)^{2-\frac{i}{2}}(1 + iax)^{2+\frac{i}{2}} dx \\ &= -\frac{\left(\frac{1}{37} - \frac{6i}{37}\right) 2^{3+\frac{i}{2}} c^2 (1 - iax)^{3-\frac{i}{2}} {}_2F_1\left(-2 - \frac{i}{2}, 3 - \frac{i}{2}; 4 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 63, normalized size = 1.00

$$\frac{\left(\frac{1}{37} - \frac{6i}{37}\right) 2^{3+\frac{i}{2}} c^2 (1 - iax)^{3-\frac{i}{2}} {}_2F_1\left(-2 - \frac{i}{2}, 3 - \frac{i}{2}; 4 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[(c + a^2*c*x^2)^2/E^ArcTan[a*x], x]

[Out] ((-1/37 + (6*I)/37)*2^(3 + I/2)*c^2*(1 - I*a*x)^(3 - I/2)*Hypergeometric2F1[-2 - I/2, 3 - I/2, 4 - I/2, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (a^2 c x^2 + c)^2 e^{-\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)^2/exp(arctan(a*x)), x)

[Out] int((a^2*c*x^2+c)^2/exp(arctan(a*x)), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^2/exp(arctan(a*x)), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^2*e^(-arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^2/exp(arctan(a*x)), x, algorithm="fricas")

[Out] integral((a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2)*e^(-arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c^2 \left(\int 2a^2 x^2 e^{-\operatorname{atan}(ax)} dx + \int a^4 x^4 e^{-\operatorname{atan}(ax)} dx + \int e^{-\operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a**2*c*x**2+c)**2/exp(atan(a*x)),x)
```

```
[Out] c**2*(Integral(2*a**2*x**2*exp(-atan(a*x)), x) + Integral(a**4*x**4*exp(-atan(a*x)), x) + Integral(exp(-atan(a*x)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^2/exp(arctan(a*x)),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{-\operatorname{atan}(ax)} (ca^2x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-atan(a*x))*(c + a^2*c*x^2)^2,x)
```

```
[Out] int(exp(-atan(a*x))*(c + a^2*c*x^2)^2, x)
```

3.275 $\int e^{-\text{ArcTan}(ax)}(c + a^2cx^2) dx$

Optimal. Leaf size=61

$$-\frac{\left(\frac{1}{17} - \frac{4i}{17}\right) 2^{2+\frac{i}{2}} c (1 - iax)^{2-\frac{i}{2}} {}_2F_1\left(-1 - \frac{i}{2}, 2 - \frac{i}{2}; 3 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

[Out] $(-1/17+4/17*I)*2^{(2+1/2*I)}*c*(1-I*a*x)^{(2-1/2*I)}*\text{hypergeom}([-1-1/2*I, 2-1/2*I], [3-1/2*I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.02, antiderivative size = 61, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5181, 71}

$$-\frac{\left(\frac{1}{17} - \frac{4i}{17}\right) 2^{2+\frac{i}{2}} c (1 - iax)^{2-\frac{i}{2}} {}_2F_1\left(-1 - \frac{i}{2}, 2 - \frac{i}{2}; 3 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)/E^{\text{ArcTan}[a*x]}, x]$

[Out] $((-1/17 + (4*I)/17)*2^{(2 + I/2)}*c*(1 - I*a*x)^{(2 - I/2)}*\text{Hypergeometric2F1}[-1 - I/2, 2 - I/2, 3 - I/2, (1 - I*a*x)/2])/a$

Rule 71

$\text{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}*((c_.) + (d_.)*(x_.)^{(n_.)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)])*(n_.)}*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{-\tan^{-1}(ax)}(c + a^2cx^2) dx &= c \int (1 - iax)^{1-\frac{i}{2}}(1 + iax)^{1+\frac{i}{2}} dx \\ &= -\frac{\left(\frac{1}{17} - \frac{4i}{17}\right) 2^{2+\frac{i}{2}} c (1 - iax)^{2-\frac{i}{2}} {}_2F_1\left(-1 - \frac{i}{2}, 2 - \frac{i}{2}; 3 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 61, normalized size = 1.00

$$\frac{\left(\frac{1}{17} - \frac{4i}{17}\right) 2^{2+\frac{i}{2}} c (1 - iax)^{2-\frac{i}{2}} {}_2F_1\left(-1 - \frac{i}{2}, 2 - \frac{i}{2}; 3 - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[(c + a^2*c*x^2)/E^ArcTan[a*x], x]

[Out] ((-1/17 + (4*I)/17)*2^(2 + I/2)*c*(1 - I*a*x)^(2 - I/2)*Hypergeometric2F1[-1 - I/2, 2 - I/2, 3 - I/2, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (a^2 c x^2 + c) e^{-\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)/exp(arctan(a*x)), x)

[Out] int((a^2*c*x^2+c)/exp(arctan(a*x)), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)/exp(arctan(a*x)), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)*e^(-arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)/exp(arctan(a*x)), x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)*e^(-arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c \left(\int a^2 x^2 e^{-\operatorname{atan}(ax)} dx + \int e^{-\operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**2*c*x**2+c)/exp(atan(a*x)),x)

[Out] c*(Integral(a**2*x**2*exp(-atan(a*x)), x) + Integral(exp(-atan(a*x)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)/exp(arctan(a*x)),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{-\operatorname{atan}(ax)} (ca^2x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))*(c + a^2*c*x^2),x)

[Out] int(exp(-atan(a*x))*(c + a^2*c*x^2), x)

3.276 $\int e^{-\text{ArcTan}(ax)} dx$

Optimal. Leaf size=60

$$-\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+\frac{i}{2}} (1-iax)^{1-\frac{i}{2}} {}_2F_1\left(-\frac{i}{2}, 1-\frac{i}{2}; 2-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

[Out] $(-1/5+2/5*I)*2^{(1+1/2*I)}*(1-I*a*x)^{(1-1/2*I)}*\text{hypergeom}([-1/2*I, 1-1/2*I], [2-1/2*I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.01, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5169, 71}

$$-\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+\frac{i}{2}} (1-iax)^{1-\frac{i}{2}} {}_2F_1\left(-\frac{i}{2}, 1-\frac{i}{2}; 2-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(-\text{ArcTan}[a*x])}, x]$

[Out] $((-1/5 + (2*I)/5)*2^{(1 + I/2)}*(1 - I*a*x)^{(1 - I/2)}*\text{Hypergeometric2F1}[-1/2*I, 1 - I/2, 2 - I/2, (1 - I*a*x)/2])/a$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)} / (b*(m + 1)*(b/(b*c - a*d))^{(n)}) * \text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5169

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}, x_Symbol] := \text{Int}[(1 - I*a*x)^{(I*(n/2))} / (1 + I*a*x)^{(I*(n/2))}, x] /;$ FreeQ[{a, n}, x] && !IntegerQ[(I*n - 1)/2]

Rubi steps

$$\begin{aligned} \int e^{-\tan^{-1}(ax)} dx &= \int (1-iax)^{-\frac{i}{2}} (1+iax)^{\frac{i}{2}} dx \\ &= -\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+\frac{i}{2}} (1-iax)^{1-\frac{i}{2}} {}_2F_1\left(-\frac{i}{2}, 1-\frac{i}{2}; 2-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 45, normalized size = 0.75

$$-\frac{\left(\frac{4}{5} + \frac{8i}{5}\right) e^{(-1+2i)\text{ArcTan}(ax)} {}_2F_1\left(1 + \frac{i}{2}, 2; 2 + \frac{i}{2}; -e^{2i\text{ArcTan}(ax)}\right)}{a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(-ArcTan[a*x]), x]``[Out] ((-4/5 - (8*I)/5)*Hypergeometric2F1[1 + I/2, 2, 2 + I/2, -E^((2*I)*ArcTan[a*x])])/(a*E^((1 - 2*I)*ArcTan[a*x]))`**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{-\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(-arctan(a*x)), x)``[Out] int(exp(-arctan(a*x)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(-arctan(a*x)), x, algorithm="maxima")``[Out] integrate(e^(-arctan(a*x)), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(-arctan(a*x)), x, algorithm="fricas")``[Out] integral(e^(-arctan(a*x)), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{-\text{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(-atan(a*x)),x)`

[Out] `Integral(exp(-atan(a*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(-arctan(a*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{-\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(-atan(a*x)),x)`

[Out] `int(exp(-atan(a*x)), x)`

$$3.277 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{c+a^2cx^2} dx$$

Optimal. Leaf size=16

$$-\frac{e^{-\text{ArcTan}(ax)}}{ac}$$

[Out] -1/a/c/exp(arctan(a*x))

Rubi [A]

time = 0.02, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {5179}

$$-\frac{e^{-\text{ArcTan}(ax)}}{ac}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)),x]

[Out] -(1/(a*c*E^ArcTan[a*x]))

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\int \frac{e^{-\tan^{-1}(ax)}}{c+a^2cx^2} dx = -\frac{e^{-\tan^{-1}(ax)}}{ac}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.01, size = 36, normalized size = 2.25

$$-\frac{(1-iax)^{-\frac{i}{2}}(1+iax)^{\frac{i}{2}}}{ac}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)),x]

[Out] -((1 + I*a*x)^(I/2)/(a*c*(1 - I*a*x)^(I/2)))

Maple [A]

time = 0.06, size = 16, normalized size = 1.00

method	result	size
gospers	$-\frac{e^{-\arctan(ax)}}{ac}$	16
risch	$-\frac{(-iax+1)^{-\frac{i}{2}}(iax+1)^{\frac{i}{2}}}{ac}$	33

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/exp(arctan(a*x))/(a^2*c*x^2+c),x,method=_RETURNVERBOSE)`

[Out] $-1/a/c/\exp(\arctan(a*x))$

Maxima [A]

time = 0.28, size = 23, normalized size = 1.44

$$-\frac{2e^{(-\arctan(ax))}}{a^3cx^2 + ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c),x,algorithm="maxima")`

[Out] $-2*e^{(-\arctan(a*x))/(a^3*c*x^2 + a*c)}$

Fricas [A]

time = 1.46, size = 15, normalized size = 0.94

$$-\frac{e^{(-\arctan(ax))}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c),x,algorithm="fricas")`

[Out] $-e^{(-\arctan(a*x))/(a*c)}$

Sympy [A]

time = 4.05, size = 15, normalized size = 0.94

$$\begin{cases} -\frac{e^{-\operatorname{atan}(ax)}}{ac} & \text{for } a \neq 0 \\ \frac{x}{c} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(atan(a*x))/(a**2*c*x**2+c),x)`

[Out] `Piecewise((-exp(-atan(a*x))/(a*c), Ne(a, 0)), (x/c, True))`

Giac [A]

time = 0.43, size = 15, normalized size = 0.94

$$-\frac{e^{(-\arctan(ax))}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c),x, algorithm="giac")

[Out] -e^(-arctan(a*x))/(a*c)

Mupad [B]

time = 0.53, size = 15, normalized size = 0.94

$$-\frac{e^{-\operatorname{atan}(ax)}}{ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a^2*c*x^2),x)

[Out] -exp(-atan(a*x))/(a*c)

$$3.278 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$$

Optimal. Leaf size=54

$$-\frac{2e^{-\text{ArcTan}(ax)}}{5ac^2} - \frac{e^{-\text{ArcTan}(ax)}(1-2ax)}{5ac^2(1+a^2x^2)}$$

[Out] -2/5/a/c^2/exp(arctan(a*x))+1/5*(2*a*x-1)/a/c^2/exp(arctan(a*x))/(a^2*x^2+1)

Rubi [A]

time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$-\frac{(1-2ax)e^{-\text{ArcTan}(ax)}}{5ac^2(a^2x^2+1)} - \frac{2e^{-\text{ArcTan}(ax)}}{5ac^2}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^2), x]

[Out] -2/(5*a*c^2*E^ArcTan[a*x]) - (1 - 2*a*x)/(5*a*c^2*E^ArcTan[a*x]*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.))/((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)])*(n_.))/((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned} \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx &= -\frac{e^{-\tan^{-1}(ax)}(1-2ax)}{5ac^2(1+a^2x^2)} + \frac{2 \int \frac{e^{-\tan^{-1}(ax)}}{c+a^2cx^2} dx}{5c} \\ &= -\frac{2e^{-\tan^{-1}(ax)}}{5ac^2} - \frac{e^{-\tan^{-1}(ax)}(1-2ax)}{5ac^2(1+a^2x^2)} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.02, size = 60, normalized size = 1.11

$$-\frac{(1-iax)^{-\frac{i}{2}}(1+iax)^{\frac{i}{2}}(3-2ax+2a^2x^2)}{5c^2(a+a^3x^2)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^2), x]

[Out] -1/5*((1 + I*a*x)^(I/2)*(3 - 2*a*x + 2*a^2*x^2))/(c^2*(1 - I*a*x)^(I/2)*(a + a^3*x^2))

Maple [A]

time = 0.10, size = 41, normalized size = 0.76

method	result	size
gosper	$-\frac{(2a^2x^2-2ax+3)e^{-\arctan(ax)}}{5(a^2x^2+1)c^2a}$	41

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)

[Out] -1/5*(2*a^2*x^2-2*a*x+3)/(a^2*x^2+1)/c^2/exp(arctan(a*x))/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(e^(-arctan(a*x))/(a^2*c*x^2 + c)^2, x)

Fricas [A]

time = 2.26, size = 41, normalized size = 0.76

$$-\frac{(2a^2x^2 - 2ax + 3)e^{-\arctan(ax)}}{5(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] -1/5*(2*a^2*x^2 - 2*a*x + 3)*e^(-arctan(a*x))/(a^3*c^2*x^2 + a*c^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 116 vs. $2(44) = 88$.

time = 22.63, size = 116, normalized size = 2.15

$$\begin{cases} -\frac{2a^2x^2}{5a^3c^2x^2e^{\operatorname{atan}(ax)}+5ac^2e^{\operatorname{atan}(ax)}} + \frac{2ax}{5a^3c^2x^2e^{\operatorname{atan}(ax)}+5ac^2e^{\operatorname{atan}(ax)}} - \frac{3}{5a^3c^2x^2e^{\operatorname{atan}(ax)}+5ac^2e^{\operatorname{atan}(ax)}} & \text{for } a \neq 0 \\ \frac{x}{c^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**2,x)

[Out] Piecewise((-2*a**2*x**2/(5*a**3*c**2*x**2*exp(atan(a*x)) + 5*a*c**2*exp(atan(a*x))) + 2*a*x/(5*a**3*c**2*x**2*exp(atan(a*x)) + 5*a*c**2*exp(atan(a*x))) - 3/(5*a**3*c**2*x**2*exp(atan(a*x)) + 5*a*c**2*exp(atan(a*x))), Ne(a, 0)), (x/c**2, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.56, size = 47, normalized size = 0.87

$$-\frac{e^{-\operatorname{atan}(ax)} \left(\frac{3}{5a^3c^2} - \frac{2x}{5a^2c^2} + \frac{2x^2}{5ac^2} \right)}{\frac{1}{a^2} + x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a^2*c*x^2)^2,x)

[Out] -(exp(-atan(a*x))*(3/(5*a^3*c^2) - (2*x)/(5*a^2*c^2) + (2*x^2)/(5*a*c^2)))/(1/a^2 + x^2)

$$3.279 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$$

Optimal. Leaf size=89

$$-\frac{24e^{-\text{ArcTan}(ax)}}{85ac^3} - \frac{e^{-\text{ArcTan}(ax)}(1-4ax)}{17ac^3(1+a^2x^2)^2} - \frac{12e^{-\text{ArcTan}(ax)}(1-2ax)}{85ac^3(1+a^2x^2)}$$

[Out] -24/85/a/c^3/exp(arctan(a*x))+1/17*(4*a*x-1)/a/c^3/exp(arctan(a*x))/(a^2*x^2+1)^2-12/85*(-2*a*x+1)/a/c^3/exp(arctan(a*x))/(a^2*x^2+1)

Rubi [A]

time = 0.06, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$-\frac{(1-4ax)e^{-\text{ArcTan}(ax)}}{17ac^3(a^2x^2+1)^2} - \frac{12(1-2ax)e^{-\text{ArcTan}(ax)}}{85ac^3(a^2x^2+1)} - \frac{24e^{-\text{ArcTan}(ax)}}{85ac^3}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c+a^2*c*x^2)^3),x]

[Out] -24/(85*a*c^3*E^ArcTan[a*x]) - (1-4*a*x)/(17*a*c^3*E^ArcTan[a*x]*(1+a^2*x^2)^2) - (12*(1-2*a*x))/(85*a*c^3*E^ArcTan[a*x]*(1+a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_.)+(d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n-2*a*(p+1)*x)*(c+d*x^2)^(p+1)*(E^(n*ArcTan[a*x]))/(a*c*(n^2+4*(p+1)^2)), x] + Dist[2*(p+1)*((2*p+3)/(c*(n^2+4*(p+1)^2))), Int[(c+d*x^2)^(p+1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] & & EqQ[d, a^2*c] & & LtQ[p, -1] & & !IntegerQ[I*n] & & NeQ[n^2+4*(p+1)^2, 0] & & IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.)+(d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] & & EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx &= -\frac{e^{-\tan^{-1}(ax)}(1-4ax)}{17ac^3(1+a^2x^2)^2} + \frac{12 \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{17c} \\
&= -\frac{e^{-\tan^{-1}(ax)}(1-4ax)}{17ac^3(1+a^2x^2)^2} - \frac{12e^{-\tan^{-1}(ax)}(1-2ax)}{85ac^3(1+a^2x^2)} + \frac{24 \int \frac{e^{-\tan^{-1}(ax)}}{c+a^2cx^2} dx}{85c^2} \\
&= -\frac{24e^{-\tan^{-1}(ax)}}{85ac^3} - \frac{e^{-\tan^{-1}(ax)}(1-4ax)}{17ac^3(1+a^2x^2)^2} - \frac{12e^{-\tan^{-1}(ax)}(1-2ax)}{85ac^3(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.12, size = 91, normalized size = 1.02

$$\frac{5e^{-\text{ArcTan}(ax)}(-1+4ax) - 12(1-iax)^{-\frac{i}{2}}(1+iax)^{\frac{i}{2}}(1+a^2x^2)(3-2ax+2a^2x^2)}{85ac^3(1+a^2x^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^3), x]

[Out] ((5*(-1 + 4*a*x))/E^ArcTan[a*x] - (12*(1 + I*a*x)^(I/2)*(1 + a^2*x^2)*(3 - 2*a*x + 2*a^2*x^2))/(1 - I*a*x)^(I/2))/(85*a*c^3*(1 + a^2*x^2)^2)

Maple [A]

time = 0.08, size = 57, normalized size = 0.64

method	result	size
gosper	$-\frac{(24a^4x^4 - 24a^3x^3 + 60a^2x^2 - 44ax + 41)e^{-\arctan(ax)}}{85(a^2x^2 + 1)^2c^3a}$	57

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)

[Out] -1/85*(24*a^4*x^4-24*a^3*x^3+60*a^2*x^2-44*a*x+41)/(a^2*x^2+1)^2/c^3/exp(arctan(a*x))/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(e^{(-arctan(a*x))}/(a²*c*x² + c)³, x)

Fricas [A]

time = 1.48, size = 68, normalized size = 0.76

$$\frac{(24 a^4 x^4 - 24 a^3 x^3 + 60 a^2 x^2 - 44 a x + 41) e^{-\arctan(ax)}}{85 (a^5 c^3 x^4 + 2 a^3 c^3 x^2 + a c^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a²*c*x²+c)³,x, algorithm="fricas")

[Out] -1/85*(24*a⁴*x⁴ - 24*a³*x³ + 60*a²*x² - 44*a*x + 41)*e^{(-arctan(a*x))} / (a⁵*c³*x⁴ + 2*a³*c³*x² + a*c³)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 291 vs. 2(76) = 152.

time = 103.57, size = 291, normalized size = 3.27

$$\left\{ \begin{array}{l} -\frac{24a^4x^4}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} + \frac{24a^3x^3}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} - \frac{60a^2x^2}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} + \frac{44ax}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} - \frac{41}{85a^5c^3x^4+170a^3c^3x^2+85ac^3} \end{array} \right. \begin{array}{l} \text{for } a \neq 0 \\ \text{otherwise} \end{array}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**3,x)

[Out] Piecewise((-24*a**4*x**4/(85*a**5*c**3*x**4*exp(atan(a*x)) + 170*a**3*c**3*x**2*exp(atan(a*x)) + 85*a*c**3*exp(atan(a*x))) + 24*a**3*x**3/(85*a**5*c**3*x**4*exp(atan(a*x)) + 170*a**3*c**3*x**2*exp(atan(a*x)) + 85*a*c**3*exp(atan(a*x))) - 60*a**2*x**2/(85*a**5*c**3*x**4*exp(atan(a*x)) + 170*a**3*c**3*x**2*exp(atan(a*x)) + 85*a*c**3*exp(atan(a*x))) + 44*a*x/(85*a**5*c**3*x**4*exp(atan(a*x)) + 170*a**3*c**3*x**2*exp(atan(a*x)) + 85*a*c**3*exp(atan(a*x))) - 41/(85*a**5*c**3*x**4*exp(atan(a*x)) + 170*a**3*c**3*x**2*exp(atan(a*x)) + 85*a*c**3*exp(atan(a*x))), Ne(a, 0)), (x/c**3, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a²*c*x²+c)³,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.62, size = 80, normalized size = 0.90

$$\frac{12 e^{-\operatorname{atan}(ax)} (2ax - 1)}{85 a c^3 (a^2 x^2 + 1)} - \frac{24 e^{-\operatorname{atan}(ax)}}{85 a c^3} + \frac{e^{-\operatorname{atan}(ax)} (4ax - 1)}{17 a c^3 (a^2 x^2 + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-atan(a*x))/(c + a^2*c*x^2)^3,x)
```

```
[Out] (12*exp(-atan(a*x))*(2*a*x - 1))/(85*a*c^3*(a^2*x^2 + 1)) - (24*exp(-atan(a*x)))/(85*a*c^3) + (exp(-atan(a*x))*(4*a*x - 1))/(17*a*c^3*(a^2*x^2 + 1)^2)
```

$$3.280 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$$

Optimal. Leaf size=124

$$-\frac{144e^{-\text{ArcTan}(ax)}}{629ac^4} - \frac{e^{-\text{ArcTan}(ax)}(1-6ax)}{37ac^4(1+a^2x^2)^3} - \frac{30e^{-\text{ArcTan}(ax)}(1-4ax)}{629ac^4(1+a^2x^2)^2} - \frac{72e^{-\text{ArcTan}(ax)}(1-2ax)}{629ac^4(1+a^2x^2)}$$

[Out] -144/629/a/c^4/exp(arctan(a*x))+1/37*(6*a*x-1)/a/c^4/exp(arctan(a*x))/(a^2*x^2+1)^3-30/629*(-4*a*x+1)/a/c^4/exp(arctan(a*x))/(a^2*x^2+1)^2-72/629*(-2*a*x+1)/a/c^4/exp(arctan(a*x))/(a^2*x^2+1)

Rubi [A]

time = 0.09, antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$-\frac{(1-6ax)e^{-\text{ArcTan}(ax)}}{37ac^4(a^2x^2+1)^3} - \frac{72(1-2ax)e^{-\text{ArcTan}(ax)}}{629ac^4(a^2x^2+1)} - \frac{30(1-4ax)e^{-\text{ArcTan}(ax)}}{629ac^4(a^2x^2+1)^2} - \frac{144e^{-\text{ArcTan}(ax)}}{629ac^4}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^4),x]

[Out] -144/(629*a*c^4*E^ArcTan[a*x]) - (1 - 6*a*x)/(37*a*c^4*E^ArcTan[a*x]*(1 + a^2*x^2)^3) - (30*(1 - 4*a*x))/(629*a*c^4*E^ArcTan[a*x]*(1 + a^2*x^2)^2) - (72*(1 - 2*a*x))/(629*a*c^4*E^ArcTan[a*x]*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))/((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^4} dx &= -\frac{e^{-\tan^{-1}(ax)}(1-6ax)}{37ac^4(1+a^2x^2)^3} + \frac{30 \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx}{37c} \\
&= -\frac{e^{-\tan^{-1}(ax)}(1-6ax)}{37ac^4(1+a^2x^2)^3} - \frac{30e^{-\tan^{-1}(ax)}(1-4ax)}{629ac^4(1+a^2x^2)^2} + \frac{360 \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{629c^2} \\
&= -\frac{e^{-\tan^{-1}(ax)}(1-6ax)}{37ac^4(1+a^2x^2)^3} - \frac{30e^{-\tan^{-1}(ax)}(1-4ax)}{629ac^4(1+a^2x^2)^2} - \frac{72e^{-\tan^{-1}(ax)}(1-2ax)}{629ac^4(1+a^2x^2)} + \frac{144 \int \frac{e^{-\tan^{-1}(ax)}}{c+a^2cx^2} dx}{629c} \\
&= -\frac{144e^{-\tan^{-1}(ax)}}{629ac^4} - \frac{e^{-\tan^{-1}(ax)}(1-6ax)}{37ac^4(1+a^2x^2)^3} - \frac{30e^{-\tan^{-1}(ax)}(1-4ax)}{629ac^4(1+a^2x^2)^2} - \frac{72e^{-\tan^{-1}(ax)}(1-2ax)}{629ac^4(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.26, size = 127, normalized size = 1.02

$$\frac{17ce^{-\text{ArcTan}(ax)}(-1+6ax) - 6(c+a^2cx^2) \left(5e^{-\text{ArcTan}(ax)}(1-4ax) + 12(1-iax)^{-\frac{1}{2}}(1+iax)^{\frac{1}{2}}(-i+ax)(i+ax)(3-2ax+2a^2x^2) \right)}{629ac^2(c+a^2cx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^4), x]

[Out] ((17*c*(-1 + 6*a*x))/E^ArcTan[a*x] - 6*(c + a^2*c*x^2)*((5*(1 - 4*a*x))/E^ArcTan[a*x] + (12*(1 + I*a*x)^(I/2)*(-I + a*x)*(I + a*x)*(3 - 2*a*x + 2*a^2*x^2))/(1 - I*a*x)^(I/2)))/(629*a*c^2*(c + a^2*c*x^2)^3)

Maple [A]

time = 0.08, size = 73, normalized size = 0.59

method	result	size
gospers	$-\frac{(144a^6x^6 - 144a^5x^5 + 504a^4x^4 - 408a^3x^3 + 606a^2x^2 - 366ax + 263)e^{-\arctan(ax)}}{629(a^2x^2 + 1)^3c^4a}$	73

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^4, x, method=_RETURNVERBOSE)

[Out] -1/629*(144*a^6*x^6-144*a^5*x^5+504*a^4*x^4-408*a^3*x^3+606*a^2*x^2-366*a*x+263)/(a^2*x^2+1)^3/c^4/exp(arctan(a*x))/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="maxima")

[Out] integrate(e^(-arctan(a*x))/(a^2*c*x^2 + c)^4, x)

Fricas [A]

time = 2.41, size = 95, normalized size = 0.77

$$\frac{(144 a^6 x^6 - 144 a^5 x^5 + 504 a^4 x^4 - 408 a^3 x^3 + 606 a^2 x^2 - 366 a x + 263) e^{(-\arctan(ax))}}{629 (a^7 c^4 x^6 + 3 a^5 c^4 x^4 + 3 a^3 c^4 x^2 + a c^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="fricas")

[Out] -1/629*(144*a^6*x^6 - 144*a^5*x^5 + 504*a^4*x^4 - 408*a^3*x^3 + 606*a^2*x^2 - 366*a*x + 263)*e^(-arctan(a*x))/(a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**4,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.68, size = 112, normalized size = 0.90

$$\frac{72 e^{-\operatorname{atan}(ax)} (2ax - 1)}{629 a c^4 (a^2 x^2 + 1)} - \frac{144 e^{-\operatorname{atan}(ax)}}{629 a c^4} + \frac{30 e^{-\operatorname{atan}(ax)} (4ax - 1)}{629 a c^4 (a^2 x^2 + 1)^2} + \frac{e^{-\operatorname{atan}(ax)} (6ax - 1)}{37 a c^4 (a^2 x^2 + 1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a^2*c*x^2)^4,x)

[Out] (72*exp(-atan(a*x))*(2*a*x - 1))/(629*a*c^4*(a^2*x^2 + 1)) - (144*exp(-atan(a*x)))/(629*a*c^4) + (30*exp(-atan(a*x))*(4*a*x - 1))/(629*a*c^4*(a^2*x^2 + 1)^2) + (exp(-atan(a*x))*(6*a*x - 1))/(37*a*c^4*(a^2*x^2 + 1)^3)

$$3.281 \quad \int e^{-\text{ArcTan}(ax)}(c + a^2cx^2)^{3/2} dx$$

Optimal. Leaf size=98

$$-\frac{\left(\frac{1}{13} - \frac{5i}{13}\right) 2^{\frac{3}{2} + \frac{i}{2}} c (1 - iax)^{\frac{5}{2} - \frac{i}{2}} \sqrt{c + a^2cx^2} {}_2F_1\left(-\frac{3}{2} - \frac{i}{2}, \frac{5}{2} - \frac{i}{2}; \frac{7}{2} - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2x^2}}$$

[Out] $(-1/13 + 5/13*I)*2^{(3/2 + 1/2*I)}*c*(1 - I*a*x)^{(5/2 - 1/2*I)}*\text{hypergeom}([5/2 - 1/2*I, -3/2 - 1/2*I], [7/2 - 1/2*I], 1/2 - 1/2*I*a*x)*(a^2*c*x^2 + c)^{(1/2)}/a/(a^2*x^2 + 1)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 98, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$-\frac{\left(\frac{1}{13} - \frac{5i}{13}\right) 2^{\frac{3}{2} + \frac{i}{2}} c (1 - iax)^{\frac{5}{2} - \frac{i}{2}} \sqrt{a^2cx^2 + c} {}_2F_1\left(-\frac{3}{2} - \frac{i}{2}, \frac{5}{2} - \frac{i}{2}; \frac{7}{2} - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{a^2x^2 + 1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)^{(3/2)}/E^{\text{ArcTan}[a*x]}, x]$

[Out] $((-1/13 + (5*I)/13)*2^{(3/2 + I/2)}*c*(1 - I*a*x)^{(5/2 - I/2)}*\text{Sqrt}[c + a^2*c*x^2]*\text{Hypergeometric2F1}[-3/2 - I/2, 5/2 - I/2, 7/2 - I/2, (1 - I*a*x)/2])/(a*\text{Sqrt}[1 + a^2*x^2])$

Rule 71

$\text{Int}[(a_) + (b_)*(x_)^{(m_)}*((c_) + (d_)*(x_)^{(n_)}), x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{\text{ArcTan}[(a_)*(x_)]*(n_)}*((c_) + (d_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{\text{ArcTan}[(a_)*(x_)]*(n_)}*((c_) + (d_)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && Eq

`Q[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])`

Rubi steps

$$\begin{aligned} \int e^{-\tan^{-1}(ax)}(c+a^2cx^2)^{3/2} dx &= \frac{\left(c\sqrt{c+a^2cx^2}\right) \int e^{-\tan^{-1}(ax)}(1+a^2x^2)^{3/2} dx}{\sqrt{1+a^2x^2}} \\ &= \frac{\left(c\sqrt{c+a^2cx^2}\right) \int (1-iax)^{\frac{3}{2}-\frac{i}{2}}(1+iax)^{\frac{3}{2}+\frac{i}{2}} dx}{\sqrt{1+a^2x^2}} \\ &= -\frac{\left(\frac{1}{13}-\frac{5i}{13}\right) 2^{\frac{3}{2}+\frac{i}{2}} c(1-iax)^{\frac{5}{2}-\frac{i}{2}} \sqrt{c+a^2cx^2} {}_2F_1\left(-\frac{3}{2}-\frac{i}{2}, \frac{5}{2}-\frac{i}{2}, \frac{7}{2}-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{1+a^2x^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 98, normalized size = 1.00

$$-\frac{\left(\frac{1}{13}-\frac{5i}{13}\right) 2^{\frac{3}{2}+\frac{i}{2}} c(1-iax)^{\frac{5}{2}-\frac{i}{2}} \sqrt{c+a^2cx^2} {}_2F_1\left(-\frac{3}{2}-\frac{i}{2}, \frac{5}{2}-\frac{i}{2}, \frac{7}{2}-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{1+a^2x^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[(c + a^2*c*x^2)^(3/2)/E^ArcTan[a*x], x]`

`[Out] ((-1/13 + (5*I)/13)*2^(3/2 + I/2)*c*(1 - I*a*x)^(5/2 - I/2)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-3/2 - I/2, 5/2 - I/2, 7/2 - I/2, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])`

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (a^2cx^2 + c)^{\frac{3}{2}} e^{-\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a^2*c*x^2+c)^(3/2)/exp(arctan(a*x)), x)`

`[Out] int((a^2*c*x^2+c)^(3/2)/exp(arctan(a*x)), x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(3/2)/exp(arctan(a*x)),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(3/2)*e^(-arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(3/2)/exp(arctan(a*x)),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(3/2)*e^(-arctan(a*x)), x)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**2*c*x**2+c)**(3/2)/exp(atan(a*x)),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(3/2)/exp(arctan(a*x)),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{-\operatorname{atan}(ax)} (ca^2x^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))*(c + a^2*c*x^2)^(3/2),x)

[Out] int(exp(-atan(a*x))*(c + a^2*c*x^2)^(3/2), x)

3.282 $\int e^{-\text{ArcTan}(ax)} \sqrt{c + a^2cx^2} dx$

Optimal. Leaf size=97

$$\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{\frac{1}{2} + \frac{i}{2}} (1 - iax)^{\frac{3}{2} - \frac{i}{2}} \sqrt{c + a^2cx^2} {}_2F_1\left(-\frac{1}{2} - \frac{i}{2}, \frac{3}{2} - \frac{i}{2}; \frac{5}{2} - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2x^2}}$$

[Out] $(-1/5+3/5*I)*2^{(1/2+1/2*I)}*(1-I*a*x)^{(3/2-1/2*I)}*\text{hypergeom}([3/2-1/2*I, -1/2-1/2*I], [5/2-1/2*I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^{(1/2)}/a/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 97, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$,

Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{\frac{1}{2} + \frac{i}{2}} (1 - iax)^{\frac{3}{2} - \frac{i}{2}} \sqrt{a^2cx^2 + c} {}_2F_1\left(-\frac{1}{2} - \frac{i}{2}, \frac{3}{2} - \frac{i}{2}; \frac{5}{2} - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{a^2x^2 + 1}}$$

Antiderivative was successfully verified.

[In] Int[Sqrt[c + a^2*c*x^2]/E^ArcTan[a*x],x]

[Out] $((-1/5 + (3*I)/5)*2^{(1/2 + I/2)}*(1 - I*a*x)^{(3/2 - I/2)}*\text{Sqrt}[c + a^2*c*x^2]*\text{Hypergeometric2F1}[-1/2 - I/2, 3/2 - I/2, 5/2 - I/2, (1 - I*a*x)/2])/(a*\text{Sqrt}[1 + a^2*x^2])$

Rule 71

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^(n)))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_)*(x_)]*(n_.))*((c_) + (d_)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_)*(x_)]*(n_.))*((c_) + (d_)*(x_)^2)^(p_), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{-\tan^{-1}(ax)} \sqrt{c + a^2 cx^2} dx &= \frac{\sqrt{c + a^2 cx^2} \int e^{-\tan^{-1}(ax)} \sqrt{1 + a^2 x^2} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\sqrt{c + a^2 cx^2} \int (1 - iax)^{\frac{1}{2} - \frac{i}{2}} (1 + iax)^{\frac{1}{2} + \frac{i}{2}} dx}{\sqrt{1 + a^2 x^2}} \\ &= -\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{\frac{1}{2} + \frac{i}{2}} (1 - iax)^{\frac{3}{2} - \frac{i}{2}} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} - \frac{i}{2}, \frac{3}{2} - \frac{i}{2}; \frac{5}{2} - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 97, normalized size = 1.00

$$-\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{\frac{1}{2} + \frac{i}{2}} (1 - iax)^{\frac{3}{2} - \frac{i}{2}} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} - \frac{i}{2}, \frac{3}{2} - \frac{i}{2}; \frac{5}{2} - \frac{i}{2}; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + a^2*c*x^2]/E^ArcTan[a*x], x]

[Out] ((-1/5 + (3*I)/5)*2^(1/2 + I/2)*(1 - I*a*x)^(3/2 - I/2)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-1/2 - I/2, 3/2 - I/2, 5/2 - I/2, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \sqrt{a^2 cx^2 + c} e^{-\arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)^(1/2)/exp(arctan(a*x)), x)

[Out] int((a^2*c*x^2+c)^(1/2)/exp(arctan(a*x)), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(1/2)/exp(arctan(a*x)), x, algorithm="maxima")

[Out] integrate(sqrt(a^2*c*x^2 + c)*e^(-arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^(1/2)/exp(arctan(a*x)),x, algorithm="fricas")
```

```
[Out] integral(sqrt(a^2*c*x^2 + c)*e^(-arctan(a*x)), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c(a^2x^2 + 1)} e^{-\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a**2*c*x**2+c)**(1/2)/exp(atan(a*x)),x)
```

```
[Out] Integral(sqrt(c*(a**2*x**2 + 1))*exp(-atan(a*x)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^(1/2)/exp(arctan(a*x)),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{-\operatorname{atan}(ax)} \sqrt{ca^2x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-atan(a*x))*(c + a^2*c*x^2)^(1/2),x)
```

```
[Out] int(exp(-atan(a*x))*(c + a^2*c*x^2)^(1/2), x)
```

$$3.283 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$$

Optimal. Leaf size=93

$$\frac{(1-i)2^{-\frac{1}{2}+\frac{i}{2}}(1-iax)^{\frac{1}{2}-\frac{i}{2}}\sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, \frac{1}{2}-\frac{i}{2}; \frac{3}{2}-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

[Out] (-1+I)*2^(-1/2+1/2*I)*(1-I*a*x)^(1/2-1/2*I)*hypergeom([1/2-1/2*I, 1/2-1/2*I], [3/2-1/2*I], 1/2-1/2*I*a*x)*(a^2*x^2+1)^(1/2)/a/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 93, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$,

Rules used = {5184, 5181, 71}

$$\frac{(1-i)2^{-\frac{1}{2}+\frac{i}{2}}(1-iax)^{\frac{1}{2}-\frac{i}{2}}\sqrt{a^2x^2+1} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, \frac{1}{2}-\frac{i}{2}; \frac{3}{2}-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2]), x]

[Out] ((-1 + I)*(1 - I*a*x)^(1/2 - I/2)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 - I/2, 1/2 - I/2, 3/2 - I/2, (1 - I*a*x)/2])/(2^(1/2 - I/2)*a*Sqrt[c + a^2*c*x^2])

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n)*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\tan^{-1}(ax)}}{\sqrt{c+a^2cx^2}} dx &= \frac{\sqrt{1+a^2x^2} \int \frac{e^{-\tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx}{\sqrt{c+a^2cx^2}} \\
&= \frac{\sqrt{1+a^2x^2} \int (1-iax)^{-\frac{1}{2}-\frac{i}{2}}(1+iax)^{-\frac{1}{2}+\frac{i}{2}} dx}{\sqrt{c+a^2cx^2}} \\
&= -\frac{(1-i)2^{-\frac{1}{2}+\frac{i}{2}}(1-iax)^{\frac{1}{2}-\frac{i}{2}}\sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, \frac{1}{2}-\frac{i}{2}; \frac{3}{2}-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 93, normalized size = 1.00

$$-\frac{(1-i)2^{-\frac{1}{2}+\frac{i}{2}}(1-iax)^{\frac{1}{2}-\frac{i}{2}}\sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}-\frac{i}{2}, \frac{1}{2}-\frac{i}{2}; \frac{3}{2}-\frac{i}{2}; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2]),x]`

```
[Out] ((-1 + I)*(1 - I*a*x)^(1/2 - I/2)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 - I/2, 1/2 - I/2, 3/2 - I/2, (1 - I*a*x)/2])/(2^(1/2 - I/2)*a*Sqrt[c + a^2*c*x^2])
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{-\arctan(ax)}}{\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2),x)``[Out] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2),x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] integrate(e^{-arctan(a*x)}/sqrt(a²*c*x² + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a²*c*x²+c)^(1/2),x, algorithm="fricas")

[Out] integral(e^{-arctan(a*x)}/sqrt(a²*c*x² + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{-\operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(exp(-atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a²*c*x²+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{-\operatorname{atan}(ax)}}{\sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a²*c*x²)^(1/2),x)

[Out] int(exp(-atan(a*x))/(c + a²*c*x²)^(1/2), x)

$$3.284 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=38

$$-\frac{e^{-\text{ArcTan}(ax)}(1-ax)}{2ac\sqrt{c+a^2cx^2}}$$

[Out] 1/2*(a*x-1)/a/c/exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {5177}

$$-\frac{(1-ax)e^{-\text{ArcTan}(ax)}}{2ac\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^(3/2)),x]

[Out] -1/2*(1 - a*x)/(a*c*E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))/((c_) + (d_.)*(x_)^2)^(3/2), x_Symbol] :=
Simp[(n + a*x)*(E^(n*ArcTan[a*x]))/(a*c*(n^2 + 1)*Sqrt[c + d*x^2]), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rubi steps

$$\int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx = -\frac{e^{-\tan^{-1}(ax)}(1-ax)}{2ac\sqrt{c+a^2cx^2}}$$

Mathematica [A]

time = 0.02, size = 37, normalized size = 0.97

$$\frac{e^{-\text{ArcTan}(ax)}(-1+ax)}{2ac\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^(3/2)),x]

[Out] (-1 + a*x)/(2*a*c*E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.09, size = 39, normalized size = 1.03

method	result	size
gospers	$\frac{(a^2x^2+1)(ax-1)e^{-\arctan(ax)}}{2a(a^2cx^2+c)^{\frac{3}{2}}}$	39

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)`[Out] $1/2*(a^2*x^2+1)*(a*x-1)/a/\exp(\arctan(a*x))/(a^2*c*x^2+c)^{(3/2)}$ **Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`[Out] `integrate(e^(-arctan(a*x))/(a^2*c*x^2 + c)^(3/2), x)`**Fricas [A]**

time = 2.66, size = 44, normalized size = 1.16

$$\frac{\sqrt{a^2cx^2 + c} (ax - 1)e^{(-\arctan(ax))}}{2(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`[Out] `1/2*sqrt(a^2*c*x^2 + c)*(a*x - 1)*e^(-arctan(a*x))/(a^3*c^2*x^2 + a*c^2)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{-\operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**(3/2),x)`[Out] `Integral(exp(-atan(a*x))/(c*(a**2*x**2 + 1))**(3/2), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.63, size = 35, normalized size = 0.92

$$\frac{e^{-\operatorname{atan}(ax)} \left(\frac{x}{2c} - \frac{1}{2ac} \right)}{\sqrt{ca^2x^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a^2*c*x^2)^(3/2),x)

[Out] (exp(-atan(a*x))*(x/(2*c) - 1/(2*a*c)))/(c + a^2*c*x^2)^(1/2)

$$3.285 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$$

Optimal. Leaf size=77

$$-\frac{e^{-\text{ArcTan}(ax)}(1-3ax)}{10ac(c+a^2cx^2)^{3/2}} - \frac{3e^{-\text{ArcTan}(ax)}(1-ax)}{10ac^2\sqrt{c+a^2cx^2}}$$

[Out] 1/10*(3*a*x-1)/a/c/exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2)-3/10*(-a*x+1)/a/c^2/exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {5178, 5177}

$$-\frac{3(1-ax)e^{-\text{ArcTan}(ax)}}{10ac^2\sqrt{a^2cx^2+c}} - \frac{(1-3ax)e^{-\text{ArcTan}(ax)}}{10ac(a^2cx^2+c)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^(5/2)), x]

[Out] -1/10*(1 - 3*a*x)/(a*c*E^ArcTan[a*x]*(c + a^2*c*x^2)^(3/2)) - (3*(1 - a*x))/(10*a*c^2*E^ArcTan[a*x]*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n + a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 1)*Sqrt[c + d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^{5/2}} dx = -\frac{e^{-\tan^{-1}(ax)}(1-3ax)}{10ac(c+a^2cx^2)^{3/2}} + \frac{3 \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx}{5c}$$

$$= -\frac{e^{-\tan^{-1}(ax)}(1-3ax)}{10ac(c+a^2cx^2)^{3/2}} - \frac{3e^{-\tan^{-1}(ax)}(1-ax)}{10ac^2\sqrt{c+a^2cx^2}}$$

Mathematica [A]

time = 0.03, size = 62, normalized size = 0.81

$$\frac{e^{-\text{ArcTan}(ax)}(-4 + 6ax - 3a^2x^2 + 3a^3x^3)}{10c^2(a + a^3x^2)\sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^(5/2)), x]``[Out] (-4 + 6*a*x - 3*a^2*x^2 + 3*a^3*x^3)/(10*c^2*E^ArcTan[a*x]*(a + a^3*x^2)*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.09, size = 56, normalized size = 0.73

method	result	size
gospers	$\frac{(a^2x^2+1)(3a^3x^3-3a^2x^2+6ax-4)e^{-\arctan(ax)}}{10a(a^2cx^2+c)^{\frac{5}{2}}}$	56

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, method=_RETURNVERBOSE)``[Out] 1/10*(a^2*x^2+1)*(3*a^3*x^3-3*a^2*x^2+6*a*x-4)/a/exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, algorithm="maxima")``[Out] integrate(e^(-arctan(a*x))/(a^2*c*x^2 + c)^(5/2), x)`

Fricas [A]

time = 2.78, size = 72, normalized size = 0.94

$$\frac{(3a^3x^3 - 3a^2x^2 + 6ax - 4)\sqrt{a^2cx^2 + c} e^{(-\arctan(ax))}}{10(a^5c^3x^4 + 2a^3c^3x^2 + ac^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="fricas")

[Out] 1/10*(3*a^3*x^3 - 3*a^2*x^2 + 6*a*x - 4)*sqrt(a^2*c*x^2 + c)*e^(-arctan(a*x))/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3006 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.65, size = 81, normalized size = 1.05

$$-\frac{e^{-\operatorname{atan}(ax)} \left(\frac{2}{5a^3c^2} - \frac{3x^3}{10c^2} - \frac{3x}{5a^2c^2} + \frac{3x^2}{10ac^2} \right)}{\frac{\sqrt{ca^2x^2+c}}{a^2} + x^2 \sqrt{ca^2x^2+c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a^2*c*x^2)^(5/2),x)

[Out] -(exp(-atan(a*x))*(2/(5*a^3*c^2) - (3*x^3)/(10*c^2) - (3*x)/(5*a^2*c^2) + (3*x^2)/(10*a*c^2)))/((c + a^2*c*x^2)^(1/2)/a^2 + x^2*(c + a^2*c*x^2)^(1/2))

$$3.286 \quad \int \frac{e^{-\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$$

Optimal. Leaf size=115

$$-\frac{e^{-\text{ArcTan}(ax)}(1-5ax)}{26ac(c+a^2cx^2)^{5/2}} - \frac{e^{-\text{ArcTan}(ax)}(1-3ax)}{13ac^2(c+a^2cx^2)^{3/2}} - \frac{3e^{-\text{ArcTan}(ax)}(1-ax)}{13ac^3\sqrt{c+a^2cx^2}}$$

[Out] 1/26*(5*a*x-1)/a/c/exp(arctan(a*x))/(a^2*c*x^2+c)^(5/2)+1/13*(3*a*x-1)/a/c^2/exp(arctan(a*x))/(a^2*c*x^2+c)^(3/2)-3/13*(-a*x+1)/a/c^3/exp(arctan(a*x))/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.09, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {5178, 5177}

$$-\frac{3(1-ax)e^{-\text{ArcTan}(ax)}}{13ac^3\sqrt{a^2cx^2+c}} - \frac{(1-3ax)e^{-\text{ArcTan}(ax)}}{13ac^2(a^2cx^2+c)^{3/2}} - \frac{(1-5ax)e^{-\text{ArcTan}(ax)}}{26ac(a^2cx^2+c)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^ArcTan[a*x]*(c+a^2*c*x^2)^(7/2)),x]

[Out] -1/26*(1-5*a*x)/(a*c*E^ArcTan[a*x]*(c+a^2*c*x^2)^(5/2)) - (1-3*a*x)/(13*a*c^2*E^ArcTan[a*x]*(c+a^2*c*x^2)^(3/2)) - (3*(1-a*x))/(13*a*c^3*E^ArcTan[a*x]*Sqrt[c+a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.)+(d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(n+a*x)*(E^(n*ArcTan[a*x]))/(a*c*(n^2+1)*Sqrt[c+d*x^2]), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_.)+(d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n-2*a*(p+1)*x)*(c+d*x^2)^(p+1)*(E^(n*ArcTan[a*x]))/(a*c*(n^2+4*(p+1)^2)), x] + Dist[2*(p+1)*((2*p+3)/(c*(n^2+4*(p+1)^2))), Int[(c+d*x^2)^(p+1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2+4*(p+1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^{7/2}} dx &= -\frac{e^{-\tan^{-1}(ax)}(1-5ax)}{26ac(c+a^2cx^2)^{5/2}} + \frac{10 \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^{5/2}} dx}{13c} \\
&= -\frac{e^{-\tan^{-1}(ax)}(1-5ax)}{26ac(c+a^2cx^2)^{5/2}} - \frac{e^{-\tan^{-1}(ax)}(1-3ax)}{13ac^2(c+a^2cx^2)^{3/2}} + \frac{6 \int \frac{e^{-\tan^{-1}(ax)}}{(c+a^2cx^2)^{3/2}} dx}{13c^2} \\
&= -\frac{e^{-\tan^{-1}(ax)}(1-5ax)}{26ac(c+a^2cx^2)^{5/2}} - \frac{e^{-\tan^{-1}(ax)}(1-3ax)}{13ac^2(c+a^2cx^2)^{3/2}} - \frac{3e^{-\tan^{-1}(ax)}(1-ax)}{13ac^3\sqrt{c+a^2cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.04, size = 81, normalized size = 0.70

$$\frac{e^{-\text{ArcTan}(ax)}(-9 + 17ax - 14a^2x^2 + 18a^3x^3 - 6a^4x^4 + 6a^5x^5)}{26ac^3(1 + a^2x^2)^2\sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^ArcTan[a*x]*(c + a^2*c*x^2)^(7/2)), x]``[Out] (-9 + 17*a*x - 14*a^2*x^2 + 18*a^3*x^3 - 6*a^4*x^4 + 6*a^5*x^5)/(26*a*c^3*E^ArcTan[a*x]*(1 + a^2*x^2)^2*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.07, size = 72, normalized size = 0.63

method	result	size
gospers	$\frac{(a^2x^2+1)(6a^5x^5-6a^4x^4+18a^3x^3-14a^2x^2+17ax-9)e^{-\arctan(ax)}}{26a(a^2cx^2+c)^{\frac{7}{2}}}$	72

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, method=_RETURNVERBOSE)``[Out] 1/26*(a^2*x^2+1)*(6*a^5*x^5-6*a^4*x^4+18*a^3*x^3-14*a^2*x^2+17*a*x-9)/a/exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, algorithm="maxima")`

[Out] integrate(e^{-arctan(a*x)}/(a²*c*x² + c)^(7/2), x)

Fricas [A]

time = 2.12, size = 99, normalized size = 0.86

$$\frac{(6a^5x^5 - 6a^4x^4 + 18a^3x^3 - 14a^2x^2 + 17ax - 9)\sqrt{a^2cx^2 + c}e^{-\arctan(ax)}}{26(a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a²*c*x²+c)^(7/2),x, algorithm="fricas")

[Out] 1/26*(6*a⁵*x⁵ - 6*a⁴*x⁴ + 18*a³*x³ - 14*a²*x² + 17*a*x - 9)*sqrt(a²*c*x² + c)*e^{-arctan(a*x)}/(a⁷*c⁴*x⁶ + 3*a⁵*c⁴*x⁴ + 3*a³*c⁴*x² + a*c⁴)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(atan(a*x))/(a**2*c*x**2+c)**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(arctan(a*x))/(a²*c*x²+c)^(7/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.67, size = 123, normalized size = 1.07

$$-\frac{e^{-\operatorname{atan}(ax)}\left(\frac{9}{26a^5c^3} - \frac{3x^5}{13c^3} - \frac{17x}{26a^4c^3} + \frac{3x^4}{13ac^3} - \frac{9x^3}{13a^2c^3} + \frac{7x^2}{13a^3c^3}\right)}{\frac{\sqrt{ca^2x^2+c}}{a^4} + x^4\sqrt{ca^2x^2+c} + \frac{2x^2\sqrt{ca^2x^2+c}}{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-atan(a*x))/(c + a²*c*x²)^(7/2),x)

[Out] -(exp(-atan(a*x))*(9/(26*a⁵*c³) - (3*x⁵)/(13*c³) - (17*x)/(26*a⁴*c³) + (3*x⁴)/(13*a*c³) - (9*x³)/(13*a²*c³) + (7*x²)/(13*a³*c³)))/((c + a²*c*x²)^(1/2)/a⁴ + x⁴*(c + a²*c*x²)^(1/2) + (2*x²*(c + a²*c*x²)^(1/2))/a²)

3.287 $\int e^{-2\text{ArcTan}(ax)}(c + a^2cx^2)^p dx$

Optimal. Leaf size=90

$$\frac{i2^{i+p}(1-iax)^{(1-i)+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1(-i-p, (1-i)+p; (2-i)+p; \frac{1}{2}(1-iax))}{a((1-i)+p)}$$

[Out] $I*2^{(I+p)}*(1-I*a*x)^{(1-I+p)}*(a^2*c*x^2+c)^p*\text{hypergeom}([-I-p, 1-I+p], [2-I+p], 1/2-1/2*I*a*x)/a/(1-I+p)/((a^2*x^2+1)^p)$

Rubi [A]

time = 0.05, antiderivative size = 90, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5184, 5181, 71}

$$\frac{i2^{p+i}(1-iax)^{p+(1-i)}(a^2x^2+1)^{-p}(a^2cx^2+c)^p {}_2F_1(-p-i, p+(1-i); p+(2-i); \frac{1}{2}(1-iax))}{a(p+(1-i))}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)^p/E^{(2*\text{ArcTan}[a*x])}, x]$

[Out] $(I*2^{(I+p)}*(1-I*a*x)^{((1-I)+p)}*(c+a^2*c*x^2)^p*\text{Hypergeometric2F1}[-I-p, (1-I)+p, (2-I)+p, (1-I*a*x)/2])/(a*((1-I)+p)*(1+a^2*x^2)^p)$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+)^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m+1)}/(b*(m+1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \&\& \text{NeQ}\{b*c - a*d, 0\} \&\& !\text{IntegerQ}[m] \&\& !\text{IntegerQ}[n] \&\& \text{GtQ}\{b/(b*c - a*d), 0\} \&\& (\text{RationalQ}[m] \mid\mid !(\text{RationalQ}[n] \&\& \text{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)])^{(n_+)}}*((c_+) + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x \&\& \text{EqQ}[d, a^2*c] \&\& (\text{IntegerQ}[p] \mid\mid \text{GtQ}[c, 0])$

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)])^{(n_+)}}*((c_+) + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x \&\& \text{EqQ}[d, a^2*c] \&\& !(\text{IntegerQ}[p] \mid\mid \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned}
\int e^{-2 \tan^{-1}(ax)} (c + a^2 cx^2)^p dx &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int e^{-2 \tan^{-1}(ax)} (1 + a^2 x^2)^p dx \\
&= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int (1 - iax)^{-i+p} (1 + iax)^{i+p} dx \\
&= \frac{i 2^{i+p} (1 - iax)^{(1-i)+p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p {}_2F_1(-i - p, (1 - i) + p; (2 - i) + p; \frac{1}{2}(1 - iax))}{a((1 - i) + p)}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 90, normalized size = 1.00

$$\frac{i 2^{i+p} (1 - iax)^{(1-i)+p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p {}_2F_1(-i - p, (1 - i) + p; (2 - i) + p; \frac{1}{2}(1 - iax))}{a((1 - i) + p)}$$

Antiderivative was successfully verified.

[In] Integrate[(c + a^2*c*x^2)^p/E^(2*ArcTan[a*x]),x]

[Out] (I*2^(I + p)*(1 - I*a*x)^((1 - I) + p)*(c + a^2*c*x^2)^p*Hypergeometric2F1[-I - p, (1 - I) + p, (2 - I) + p, (1 - I*a*x)/2])/(a*((1 - I) + p)*(1 + a^2*x^2)^p)

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int (a^2 c x^2 + c)^p e^{-2 \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)^p/exp(2*arctan(a*x)),x)

[Out] int((a^2*c*x^2+c)^p/exp(2*arctan(a*x)),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^p/exp(2*arctan(a*x)),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^p*e^(-2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^p/exp(2*arctan(a*x)),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^p*e^(-2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^p e^{-2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**2*c*x**2+c)**p/exp(2*atan(a*x)),x)

[Out] Integral((c*(a**2*x**2 + 1))**p*exp(-2*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^p/exp(2*arctan(a*x)),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{-2\operatorname{atan}(ax)} (ca^2x^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^p,x)

[Out] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^p, x)

$$3.288 \quad \int e^{-2\text{ArcTan}(ax)}(c + a^2cx^2)^2 dx$$

Optimal. Leaf size=53

$$-\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{1+i} c^2 (1 - iax)^{3-i} {}_2F_1\left(-2 - i, 3 - i; 4 - i; \frac{1}{2}(1 - iax)\right)}{a}$$

[Out] $(-1/5+3/5*I)*2^{(1+I)}*c^2*(1-I*a*x)^{(3-I)}*\text{hypergeom}([3-I, -2-I], [4-I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.03, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5181, 71}

$$-\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{1+i} c^2 (1 - iax)^{3-i} {}_2F_1\left(-2 - i, 3 - i; 4 - i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Int[(c + a^2*c*x^2)^2/E^(2*ArcTan[a*x]),x]

[Out] $((-1/5 + (3*I)/5)*2^{(1 + I)}*c^2*(1 - I*a*x)^{(3 - I)}*\text{Hypergeometric2F1}[-2 - I, 3 - I, 4 - I, (1 - I*a*x)/2])/a$

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{-2\text{tan}^{-1}(ax)}(c + a^2cx^2)^2 dx &= c^2 \int (1 - iax)^{2-i}(1 + iax)^{2+i} dx \\ &= -\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{1+i} c^2 (1 - iax)^{3-i} {}_2F_1\left(-2 - i, 3 - i; 4 - i; \frac{1}{2}(1 - iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 53, normalized size = 1.00

$$\frac{\left(\frac{1}{5} - \frac{3i}{5}\right) 2^{1+i} c^2 (1 - iax)^{3-i} {}_2F_1\left(-2 - i, 3 - i; 4 - i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[(c + a^2*c*x^2)^2/E^(2*ArcTan[a*x]),x]

[Out] ((-1/5 + (3*I)/5)*2^(1 + I)*c^2*(1 - I*a*x)^(3 - I)*Hypergeometric2F1[-2 - I, 3 - I, 4 - I, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (a^2 c x^2 + c)^2 e^{-2 \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)^2/exp(2*arctan(a*x)),x)

[Out] int((a^2*c*x^2+c)^2/exp(2*arctan(a*x)),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^2/exp(2*arctan(a*x)),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^2*e^(-2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^2/exp(2*arctan(a*x)),x, algorithm="fricas")

[Out] integral((a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2)*e^(-2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c^2 \left(\int 2a^2 x^2 e^{-2 \operatorname{atan}(ax)} dx + \int a^4 x^4 e^{-2 \operatorname{atan}(ax)} dx + \int e^{-2 \operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**2*c*x**2+c)**2/exp(2*atan(a*x)),x)

[Out] c**2*(Integral(2*a**2*x**2*exp(-2*atan(a*x)), x) + Integral(a**4*x**4*exp(-2*atan(a*x)), x) + Integral(exp(-2*atan(a*x)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^2/exp(2*arctan(a*x)),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{-2\operatorname{atan}(ax)} (ca^2x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^2,x)

[Out] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^2, x)

$$3.289 \quad \int e^{-2\text{ArcTan}(ax)}(c + a^2cx^2) dx$$

Optimal. Leaf size=51

$$-\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+i} c (1 - iax)^{2-i} {}_2F_1\left(-1 - i, 2 - i; 3 - i; \frac{1}{2}(1 - iax)\right)}{a}$$

[Out] $(-1/5+2/5*I)*2^{(1+I)*c*(1-I*a*x)^{(2-I)*\text{hypergeom}([2-I, -1-I], [3-I], 1/2-1/2*I*a*x)/a}$

Rubi [A]

time = 0.02, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5181, 71}

$$-\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+i} c (1 - iax)^{2-i} {}_2F_1\left(-1 - i, 2 - i; 3 - i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Int[(c + a^2*c*x^2)/E^(2*ArcTan[a*x]),x]

[Out] $((-1/5 + (2*I)/5)*2^{(1 + I)*c*(1 - I*a*x)^{(2 - I)*\text{Hypergeometric2F1}[-1 - I, 2 - I, 3 - I, (1 - I*a*x)/2]})/a$

Rule 71

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_)*(x_)])*(n_)*((c_) + (d_)*(x_)^2)^(p_), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{-2\text{tan}^{-1}(ax)}(c + a^2cx^2) dx &= c \int (1 - iax)^{1-i}(1 + iax)^{1+i} dx \\ &= -\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+i} c (1 - iax)^{2-i} {}_2F_1\left(-1 - i, 2 - i; 3 - i; \frac{1}{2}(1 - iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 51, normalized size = 1.00

$$-\frac{\left(\frac{1}{5} - \frac{2i}{5}\right) 2^{1+i} c (1 - iax)^{2-i} {}_2F_1\left(-1 - i, 2 - i; 3 - i; \frac{1}{2}(1 - iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[(c + a^2*c*x^2)/E^(2*ArcTan[a*x]),x]

[Out] ((-1/5 + (2*I)/5)*2^(1 + I)*c*(1 - I*a*x)^(2 - I)*Hypergeometric2F1[-1 - I, 2 - I, 3 - I, (1 - I*a*x)/2])/a

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (a^2 c x^2 + c) e^{-2 \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)/exp(2*arctan(a*x)),x)

[Out] int((a^2*c*x^2+c)/exp(2*arctan(a*x)),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)/exp(2*arctan(a*x)),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)*e^(-2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)/exp(2*arctan(a*x)),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)*e^(-2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c \left(\int a^2 x^2 e^{-2 \operatorname{atan}(ax)} dx + \int e^{-2 \operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a**2*c*x**2+c)/exp(2*atan(a*x)),x)
```

```
[Out] c*(Integral(a**2*x**2*exp(-2*atan(a*x)), x) + Integral(exp(-2*atan(a*x)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)/exp(2*arctan(a*x)),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{-2 \operatorname{atan}(ax)} (ca^2 x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-2*atan(a*x))*(c + a^2*c*x^2),x)
```

```
[Out] int(exp(-2*atan(a*x))*(c + a^2*c*x^2), x)
```

3.290 $\int e^{-2\text{ArcTan}(ax)} dx$

Optimal. Leaf size=46

$$\frac{(1-i)2^{-1+i}(1-iax)^{1-i} {}_2F_1\left(-i, 1-i; 2-i; \frac{1}{2}(1-iax)\right)}{a}$$

[Out] $(-1+I)*2^{(-1+I)}*(1-I*a*x)^{(1-I)}*\text{hypergeom}([-I, 1-I], [2-I], 1/2-1/2*I*a*x)/a$

Rubi [A]

time = 0.01, antiderivative size = 46, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5169, 71}

$$\frac{(1-i)2^{-1+i}(1-iax)^{1-i} {}_2F_1\left(-i, 1-i; 2-i; \frac{1}{2}(1-iax)\right)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{-2*\text{ArcTan}[a*x]}, x]$

[Out] $((-1 + I)*(1 - I*a*x)^{(1 - I)}*\text{Hypergeometric2F1}[-I, 1 - I, 2 - I, (1 - I*a*x)/2])/(2^{(1 - I)*a})$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \&\& \text{NeQ}[b*c - a*d, 0] \&\& !\text{IntegerQ}[m] \&\& !\text{IntegerQ}[n] \&\& \text{GtQ}[b/(b*c - a*d), 0] \&\& (\text{RationalQ}[m] || !(\text{RationalQ}[n] \&\& \text{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5169

$\text{Int}[E^{\text{ArcTan}[(a_+)*(x_+)]*(n_+)}, x_Symbol] :> \text{Int}[(1 - I*a*x)^{(I*(n/2))}/(1 + I*a*x)^{(I*(n/2))}, x] /; \text{FreeQ}\{a, n\}, x \&\& !\text{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{-2 \tan^{-1}(ax)} dx &= \int (1 - iax)^{-i} (1 + iax)^i dx \\ &= \frac{(1-i)2^{-1+i}(1-iax)^{1-i} {}_2F_1\left(-i, 1-i; 2-i; \frac{1}{2}(1-iax)\right)}{a} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 37, normalized size = 0.80

$$\frac{(1+i)e^{(-2+2i)\text{ArcTan}(ax)} {}_2F_1(1+i, 2; 2+i; -e^{2i\text{ArcTan}(ax)})}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(-2*ArcTan[a*x]), x]

[Out] ((-1 - I)*Hypergeometric2F1[1 + I, 2, 2 + I, -E^((2*I)*ArcTan[a*x])])/(a*E^((2 - 2*I)*ArcTan[a*x]))

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{-2 \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*arctan(a*x)), x)

[Out] int(exp(-2*arctan(a*x)), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(-2*arctan(a*x)), x, algorithm="maxima")

[Out] integrate(e^(-2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(-2*arctan(a*x)), x, algorithm="fricas")

[Out] integral(e^(-2*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{-2 \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(-2*atan(a*x)),x)`

[Out] `Integral(exp(-2*atan(a*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(-2*arctan(a*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int e^{-2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(-2*atan(a*x)),x)`

[Out] `int(exp(-2*atan(a*x)), x)`

$$3.291 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{c+a^2cx^2} dx$$

Optimal. Leaf size=18

$$-\frac{e^{-2\text{ArcTan}(ax)}}{2ac}$$

[Out] -1/2/a/c/exp(2*arctan(a*x))

Rubi [A]

time = 0.02, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {5179}

$$-\frac{e^{-2\text{ArcTan}(ax)}}{2ac}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)),x]

[Out] -1/2*1/(a*c*E^(2*ArcTan[a*x]))

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\int \frac{e^{-2 \tan^{-1}(ax)}}{c + a^2 cx^2} dx = -\frac{e^{-2 \tan^{-1}(ax)}}{2ac}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.01, size = 34, normalized size = 1.89

$$-\frac{(1 - iax)^{-i}(1 + iax)^i}{2ac}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)),x]

[Out] -1/2*(1 + I*a*x)^I/(a*c*(1 - I*a*x)^I)

Maple [A]

time = 0.06, size = 18, normalized size = 1.00

method	result	size
gospers	$-\frac{e^{-2 \arctan(ax)}}{2ac}$	18
risch	$-\frac{(-iax+1)^{-i}(iax+1)^i}{2ac}$	33

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c),x,method=_RETURNVERBOSE)
```

```
[Out] -1/2/a/c/exp(2*arctan(a*x))
```

Maxima [A]

time = 0.27, size = 23, normalized size = 1.28

$$-\frac{e^{(-2 \arctan(ax))}}{a^3cx^2 + ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="maxima")
```

```
[Out] -e^(-2*arctan(a*x))/(a^3*c*x^2 + a*c)
```

Fricas [A]

time = 3.82, size = 15, normalized size = 0.83

$$-\frac{e^{(-2 \arctan(ax))}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="fricas")
```

```
[Out] -1/2*e^(-2*arctan(a*x))/(a*c)
```

Sympy [A]

time = 8.60, size = 19, normalized size = 1.06

$$\begin{cases} -\frac{e^{-2 \operatorname{atan}(ax)}}{2ac} & \text{for } a \neq 0 \\ \frac{x}{c} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c),x)
```

```
[Out] Piecewise((-exp(-2*atan(a*x))/(2*a*c), Ne(a, 0)), (x/c, True))
```

Giac [A]

time = 0.40, size = 15, normalized size = 0.83

$$-\frac{e^{(-2 \arctan(ax))}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="giac")

[Out] -1/2*e^(-2*arctan(a*x))/(a*c)

Mupad [B]

time = 0.53, size = 15, normalized size = 0.83

$$-\frac{e^{-2 \operatorname{atan}(ax)}}{2ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2),x)

[Out] -exp(-2*atan(a*x))/(2*a*c)

$$3.292 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^2} dx$$

Optimal. Leaf size=54

$$-\frac{e^{-2\text{ArcTan}(ax)}}{8ac^2} - \frac{e^{-2\text{ArcTan}(ax)}(1-ax)}{4ac^2(1+a^2x^2)}$$

[Out] -1/8/a/c^2/exp(2*arctan(a*x))+1/4*(a*x-1)/a/c^2/exp(2*arctan(a*x))/(a^2*x^2+1)

Rubi [A]

time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$-\frac{(1-ax)e^{-2\text{ArcTan}(ax)}}{4ac^2(a^2x^2+1)} - \frac{e^{-2\text{ArcTan}(ax)}}{8ac^2}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^2), x]

[Out] -1/8*1/(a*c^2*E^(2*ArcTan[a*x])) - (1 - a*x)/(4*a*c^2*E^(2*ArcTan[a*x])*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2)), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned} \int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^2} dx &= -\frac{e^{-2 \tan^{-1}(ax)}(1 - ax)}{4ac^2(1 + a^2x^2)} + \frac{\int \frac{e^{-2 \tan^{-1}(ax)}}{c + a^2 cx^2} dx}{4c} \\ &= -\frac{e^{-2 \tan^{-1}(ax)}}{8ac^2} - \frac{e^{-2 \tan^{-1}(ax)}(1 - ax)}{4ac^2(1 + a^2x^2)} \end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.02, size = 55, normalized size = 1.02

$$-\frac{(1-iax)^{-i}(1+iax)^i(3-2ax+a^2x^2)}{8c^2(a+a^3x^2)}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(2*ArcTan[a*x]))*(c + a^2*c*x^2)^2, x]

[Out] -1/8*((1 + I*a*x)^I*(3 - 2*a*x + a^2*x^2))/(c^2*(1 - I*a*x)^I*(a + a^3*x^2))

Maple [A]

time = 0.09, size = 42, normalized size = 0.78

method	result	size
gospers	$-\frac{(a^2x^2-2ax+3)e^{-2\arctan(ax)}}{8(a^2x^2+1)c^2a}$	42

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x,method=_RETURNVERBOSE)

[Out] -1/8*(a^2*x^2-2*a*x+3)/(a^2*x^2+1)/c^2/exp(2*arctan(a*x))/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(e^(-2*arctan(a*x))/(a^2*c*x^2 + c)^2, x)

Fricas [A]

time = 2.69, size = 40, normalized size = 0.74

$$-\frac{(a^2x^2 - 2ax + 3)e^{(-2\arctan(ax))}}{8(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] -1/8*(a^2*x^2 - 2*a*x + 3)*e^(-2*arctan(a*x))/(a^3*c^2*x^2 + a*c^2)

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 124 vs. $2(46) = 92$.

time = 44.95, size = 124, normalized size = 2.30

$$\begin{cases} -\frac{a^2 x^2}{8a^3 c^2 x^2 e^{2 \operatorname{atan}(ax)} + 8ac^2 e^{2 \operatorname{atan}(ax)}} + \frac{2ax}{8a^3 c^2 x^2 e^{2 \operatorname{atan}(ax)} + 8ac^2 e^{2 \operatorname{atan}(ax)}} - \frac{3}{8a^3 c^2 x^2 e^{2 \operatorname{atan}(ax)} + 8ac^2 e^{2 \operatorname{atan}(ax)}} & \text{for } a \neq 0 \\ \frac{x}{c^2} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c)**2,x)

[Out] Piecewise((-a**2*x**2/(8*a**3*c**2*x**2*exp(2*atan(a*x)) + 8*a*c**2*exp(2*atan(a*x))) + 2*a*x/(8*a**3*c**2*x**2*exp(2*atan(a*x)) + 8*a*c**2*exp(2*atan(a*x))) - 3/(8*a**3*c**2*x**2*exp(2*atan(a*x)) + 8*a*c**2*exp(2*atan(a*x))), Ne(a, 0)), (x/c**2, True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.57, size = 47, normalized size = 0.87

$$-\frac{e^{-2 \operatorname{atan}(ax)} \left(\frac{3}{8a^3 c^2} - \frac{x}{4a^2 c^2} + \frac{x^2}{8ac^2} \right)}{\frac{1}{a^2} + x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^2,x)

[Out] -(exp(-2*atan(a*x))*(3/(8*a^3*c^2) - x/(4*a^2*c^2) + x^2/(8*a*c^2)))/(1/a^2 + x^2)

$$3.293 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^3} dx$$

Optimal. Leaf size=89

$$-\frac{3e^{-2\text{ArcTan}(ax)}}{40ac^3} - \frac{e^{-2\text{ArcTan}(ax)}(1-2ax)}{10ac^3(1+a^2x^2)^2} - \frac{3e^{-2\text{ArcTan}(ax)}(1-ax)}{20ac^3(1+a^2x^2)}$$

[Out] $-3/40/a/c^3/\exp(2*\arctan(a*x))+1/10*(2*a*x-1)/a/c^3/\exp(2*\arctan(a*x))/(a^2*x^2+1)^2-3/20*(-a*x+1)/a/c^3/\exp(2*\arctan(a*x))/(a^2*x^2+1)$

Rubi [A]

time = 0.07, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$-\frac{(1-2ax)e^{-2\text{ArcTan}(ax)}}{10ac^3(a^2x^2+1)^2} - \frac{3(1-ax)e^{-2\text{ArcTan}(ax)}}{20ac^3(a^2x^2+1)} - \frac{3e^{-2\text{ArcTan}(ax)}}{40ac^3}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^3), x]

[Out] $-3/(40*a*c^3*E^(2*ArcTan[a*x])) - (1 - 2*a*x)/(10*a*c^3*E^(2*ArcTan[a*x])*(1 + a^2*x^2)^2) - (3*(1 - a*x))/(20*a*c^3*E^(2*ArcTan[a*x])*(1 + a^2*x^2))$

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x]))/(a*c*(n^2 + 4*(p + 1)^2)), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-2 \tan^{-1}(ax)}}{(c+a^2cx^2)^3} dx &= -\frac{e^{-2 \tan^{-1}(ax)}(1-2ax)}{10ac^3(1+a^2x^2)^2} + \frac{3 \int \frac{e^{-2 \tan^{-1}(ax)}}{(c+a^2cx^2)^2} dx}{5c} \\
&= -\frac{e^{-2 \tan^{-1}(ax)}(1-2ax)}{10ac^3(1+a^2x^2)^2} - \frac{3e^{-2 \tan^{-1}(ax)}(1-ax)}{20ac^3(1+a^2x^2)} + \frac{3 \int \frac{e^{-2 \tan^{-1}(ax)}}{c+a^2cx^2} dx}{20c^2} \\
&= -\frac{3e^{-2 \tan^{-1}(ax)}}{40ac^3} - \frac{e^{-2 \tan^{-1}(ax)}(1-2ax)}{10ac^3(1+a^2x^2)^2} - \frac{3e^{-2 \tan^{-1}(ax)}(1-ax)}{20ac^3(1+a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.12, size = 85, normalized size = 0.96

$$\frac{e^{-2 \operatorname{ArcTan}(ax)}(-4+8ax) - 3(1-iax)^{-i}(1+iax)^i(1+a^2x^2)(3-2ax+a^2x^2)}{40ac^3(1+a^2x^2)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(2*ArcTan[a*x])*(c+a^2*c*x^2)^3),x]

[Out] ((-4+8*a*x)/E^(2*ArcTan[a*x]) - (3*(1+I*a*x)^I*(1+a^2*x^2)*(3-2*a*x+a^2*x^2))/(1-I*a*x)^I)/(40*a*c^3*(1+a^2*x^2)^2)

Maple [A]

time = 0.08, size = 59, normalized size = 0.66

method	result	size
gospers	$-\frac{(3a^4x^4-6a^3x^3+12a^2x^2-14ax+13)e^{-2 \arctan(ax)}}{40(a^2x^2+1)^2c^3a}$	59

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)

[Out] -1/40*(3*a^4*x^4-6*a^3*x^3+12*a^2*x^2-14*a*x+13)/(a^2*x^2+1)^2/c^3/exp(2*arctan(a*x))/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] integrate($e^{(-2*\arctan(ax))}/(a^2*c*x^2 + c)^3, x$)

Fricas [A]

time = 4.05, size = 68, normalized size = 0.76

$$-\frac{(3a^4x^4 - 6a^3x^3 + 12a^2x^2 - 14ax + 13)e^{(-2\arctan(ax))}}{40(a^5c^3x^4 + 2a^3c^3x^2 + ac^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(ax))/(a^2*c*x^2+c)^3,x, algorithm="fricas")

[Out] $-1/40*(3*a^4*x^4 - 6*a^3*x^3 + 12*a^2*x^2 - 14*a*x + 13)*e^{(-2*\arctan(ax))}/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*atan(ax))/(a**2*c*x**2+c)**3,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(ax))/(a^2*c*x^2+c)^3,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.62, size = 79, normalized size = 0.89

$$\frac{3e^{-2\operatorname{atan}(ax)}(ax-1)}{20ac^3(a^2x^2+1)} - \frac{3e^{-2\operatorname{atan}(ax)}}{40ac^3} + \frac{e^{-2\operatorname{atan}(ax)}(2ax-1)}{10ac^3(a^2x^2+1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(ax))/(c + a^2*c*x^2)^3,x)

[Out] $(3*\exp(-2*\operatorname{atan}(ax))*(ax-1))/(20*a*c^3*(a^2*x^2+1)) - (3*\exp(-2*\operatorname{atan}(ax)))/(40*a*c^3) + (\exp(-2*\operatorname{atan}(ax))*(2*ax-1))/(10*a*c^3*(a^2*x^2+1)^2)$

$$3.294 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$$

Optimal. Leaf size=124

$$-\frac{9e^{-2\text{ArcTan}(ax)}}{160ac^4} - \frac{e^{-2\text{ArcTan}(ax)}(1-3ax)}{20ac^4(1+a^2x^2)^3} - \frac{3e^{-2\text{ArcTan}(ax)}(1-2ax)}{40ac^4(1+a^2x^2)^2} - \frac{9e^{-2\text{ArcTan}(ax)}(1-ax)}{80ac^4(1+a^2x^2)}$$

[Out] $-9/160/a/c^4/\exp(2*\arctan(a*x))+1/20*(3*a*x-1)/a/c^4/\exp(2*\arctan(a*x))/(a^2*x^2+1)^3-3/40*(-2*a*x+1)/a/c^4/\exp(2*\arctan(a*x))/(a^2*x^2+1)^2-9/80*(-a*x+1)/a/c^4/\exp(2*\arctan(a*x))/(a^2*x^2+1)$

Rubi [A]

time = 0.09, antiderivative size = 124, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$-\frac{(1-3ax)e^{-2\text{ArcTan}(ax)}}{20ac^4(a^2x^2+1)^3} - \frac{9(1-ax)e^{-2\text{ArcTan}(ax)}}{80ac^4(a^2x^2+1)} - \frac{3(1-2ax)e^{-2\text{ArcTan}(ax)}}{40ac^4(a^2x^2+1)^2} - \frac{9e^{-2\text{ArcTan}(ax)}}{160ac^4}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^4), x]

[Out] $-9/(160*a*c^4*E^(2*ArcTan[a*x])) - (1 - 3*a*x)/(20*a*c^4*E^(2*ArcTan[a*x])*(1 + a^2*x^2)^3) - (3*(1 - 2*a*x))/(40*a*c^4*E^(2*ArcTan[a*x])*(1 + a^2*x^2)^2) - (9*(1 - a*x))/(80*a*c^4*E^(2*ArcTan[a*x])*(1 + a^2*x^2))$

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2)), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_) + (d_.)*(x_)^2), x_Symbol] :> Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^4} dx &= -\frac{e^{-2 \tan^{-1}(ax)}(1 - 3ax)}{20ac^4(1 + a^2x^2)^3} + \frac{3 \int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^3} dx}{4c} \\
&= -\frac{e^{-2 \tan^{-1}(ax)}(1 - 3ax)}{20ac^4(1 + a^2x^2)^3} - \frac{3e^{-2 \tan^{-1}(ax)}(1 - 2ax)}{40ac^4(1 + a^2x^2)^2} + \frac{9 \int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^2} dx}{20c^2} \\
&= -\frac{e^{-2 \tan^{-1}(ax)}(1 - 3ax)}{20ac^4(1 + a^2x^2)^3} - \frac{3e^{-2 \tan^{-1}(ax)}(1 - 2ax)}{40ac^4(1 + a^2x^2)^2} - \frac{9e^{-2 \tan^{-1}(ax)}(1 - ax)}{80ac^4(1 + a^2x^2)} + \frac{9 \int \frac{e^{-2 \tan^{-1}(ax)}}{c + a^2 cx^2} dx}{80c} \\
&= -\frac{9e^{-2 \tan^{-1}(ax)}}{160ac^4} - \frac{e^{-2 \tan^{-1}(ax)}(1 - 3ax)}{20ac^4(1 + a^2x^2)^3} - \frac{3e^{-2 \tan^{-1}(ax)}(1 - 2ax)}{40ac^4(1 + a^2x^2)^2} - \frac{9e^{-2 \tan^{-1}(ax)}(1 - ax)}{80ac^4(1 + a^2x^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.25, size = 121, normalized size = 0.98

$$\frac{8ce^{-2\text{ArcTan}(ax)}(-1 + 3ax) - 3(c + a^2cx^2)(e^{-2\text{ArcTan}(ax)}(4 - 8ax) + 3(1 - iax)^{-i}(1 + iax)^i(-i + ax)(i + ax)(3 - 2ax + a^2x^2))}{160ac^2(c + a^2cx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(2*ArcTan[a*x]))*(c + a^2*c*x^2)^4, x]

[Out] ((8*c*(-1 + 3*a*x))/E^(2*ArcTan[a*x])) - 3*(c + a^2*c*x^2)*((4 - 8*a*x)/E^(2*ArcTan[a*x]) + (3*(1 + I*a*x)^I*(-I + a*x)*(I + a*x)*(3 - 2*a*x + a^2*x^2))/(1 - I*a*x^I))/(160*a*c^2*(c + a^2*c*x^2)^3)

Maple [A]

time = 0.08, size = 75, normalized size = 0.60

method	result	size
gospers	$-\frac{(9a^6x^6 - 18a^5x^5 + 45a^4x^4 - 60a^3x^3 + 75a^2x^2 - 66ax + 47)e^{-2 \arctan(ax)}}{160(a^2x^2 + 1)^3 c^4 a}$	75

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^4, x, method=_RETURNVERBOSE)

[Out] -1/160*(9*a^6*x^6-18*a^5*x^5+45*a^4*x^4-60*a^3*x^3+75*a^2*x^2-66*a*x+47)/(a^2*x^2+1)^3/c^4/exp(2*arctan(a*x))/a

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="maxima")

[Out] integrate(e^(-2*arctan(a*x))/(a^2*c*x^2 + c)^4, x)

Fricas [A]

time = 6.90, size = 95, normalized size = 0.77

$$\frac{(9a^6x^6 - 18a^5x^5 + 45a^4x^4 - 60a^3x^3 + 75a^2x^2 - 66ax + 47)e^{(-2\arctan(ax))}}{160(a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="fricas")

[Out] -1/160*(9*a^6*x^6 - 18*a^5*x^5 + 45*a^4*x^4 - 60*a^3*x^3 + 75*a^2*x^2 - 66*a*x + 47)*e^(-2*arctan(a*x))/(a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c)**4,x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.67, size = 111, normalized size = 0.90

$$\frac{9e^{-2\operatorname{atan}(ax)}(ax-1)}{80ac^4(a^2x^2+1)} - \frac{9e^{-2\operatorname{atan}(ax)}}{160ac^4} + \frac{3e^{-2\operatorname{atan}(ax)}(2ax-1)}{40ac^4(a^2x^2+1)^2} + \frac{e^{-2\operatorname{atan}(ax)}(3ax-1)}{20ac^4(a^2x^2+1)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^4,x)

[Out] (9*exp(-2*atan(a*x))*(a*x - 1))/(80*a*c^4*(a^2*x^2 + 1)) - (9*exp(-2*atan(a*x)))/(160*a*c^4) + (3*exp(-2*atan(a*x))*(2*a*x - 1))/(40*a*c^4*(a^2*x^2 + 1)^2) + (exp(-2*atan(a*x))*(3*a*x - 1))/(20*a*c^4*(a^2*x^2 + 1)^3)

$$3.295 \quad \int e^{-2\text{ArcTan}(ax)}(c + a^2cx^2)^{3/2} dx$$

Optimal. Leaf size=88

$$\frac{\left(\frac{2}{29} - \frac{5i}{29}\right) 2^{\frac{5}{2}+i} c(1-iax)^{\frac{5}{2}-i} \sqrt{c+a^2cx^2} {}_2F_1\left(-\frac{3}{2}-i, \frac{5}{2}-i; \frac{7}{2}-i; \frac{1}{2}(1-iax)\right)}{a\sqrt{1+a^2x^2}}$$

[Out] $(-2/29+5/29*I)*2^{(5/2+I)}*c*(1-I*a*x)^{(5/2-I)}*\text{hypergeom}([-3/2-I, 5/2-I], [7/2-I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^{(1/2)}/a/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{2}{29} - \frac{5i}{29}\right) 2^{\frac{5}{2}+i} c(1-iax)^{\frac{5}{2}-i} \sqrt{a^2cx^2+c} {}_2F_1\left(-\frac{3}{2}-i, \frac{5}{2}-i; \frac{7}{2}-i; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)^{(3/2)}/E^{(2*\text{ArcTan}[a*x])}, x]$

[Out] $((-2/29 + (5*I)/29)*2^{(5/2 + I)}*c*(1 - I*a*x)^{(5/2 - I)}*\text{Sqrt}[c + a^2*c*x^2]*\text{Hypergeometric2F1}[-3/2 - I, 5/2 - I, 7/2 - I, (1 - I*a*x)/2])/(a*\text{Sqrt}[1 + a^2*x^2])$

Rule 71

$\text{Int}[(a + b*x)^{(m+1)}/(b*(m+1)*(b/(b*c - a*d))^n)*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a+b*x)/(b*c - a*d)], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[a*x])*(n)}*((c + d*x^2)^{(p)}), x_Symbol] :=$ Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[a*x])*(n)}*((c + d*x^2)^{(p)}), x_Symbol] :=$ Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int e^{-2 \tan^{-1}(ax)} (c + a^2 cx^2)^{3/2} dx &= \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int e^{-2 \tan^{-1}(ax)} (1 + a^2 x^2)^{3/2} dx}{\sqrt{1 + a^2 x^2}} \\
&= \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int (1 - iax)^{\frac{3}{2}-i} (1 + iax)^{\frac{3}{2}+i} dx}{\sqrt{1 + a^2 x^2}} \\
&= -\frac{\left(\frac{2}{29} - \frac{5i}{29}\right) 2^{\frac{5}{2}+i} c (1 - iax)^{\frac{5}{2}-i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{3}{2} - i, \frac{5}{2} - i; \frac{7}{2} - i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 88, normalized size = 1.00

$$-\frac{\left(\frac{2}{29} - \frac{5i}{29}\right) 2^{\frac{5}{2}+i} c (1 - iax)^{\frac{5}{2}-i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{3}{2} - i, \frac{5}{2} - i; \frac{7}{2} - i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[(c + a^2*c*x^2)^(3/2)/E^(2*ArcTan[a*x]), x]`

```
[Out] ((-2/29 + (5*I)/29)*2^(5/2 + I)*c*(1 - I*a*x)^(5/2 - I)*Sqrt[c + a^2*c*x^2]
*Hypergeometric2F1[-3/2 - I, 5/2 - I, 7/2 - I, (1 - I*a*x)/2])/(a*Sqrt[1 +
a^2*x^2])
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int (a^2 c x^2 + c)^{\frac{3}{2}} e^{-2 \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((a^2*c*x^2+c)^(3/2)/exp(2*arctan(a*x)), x)``[Out] int((a^2*c*x^2+c)^(3/2)/exp(2*arctan(a*x)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((a^2*c*x^2+c)^(3/2)/exp(2*arctan(a*x)), x, algorithm="maxima")`

[Out] integrate((a^2*c*x^2 + c)^(3/2)*e^(-2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(3/2)/exp(2*arctan(a*x)),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(3/2)*e^(-2*arctan(a*x)), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**2*c*x**2+c)**(3/2)/exp(2*atan(a*x)),x)

[Out] Timed out

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(3/2)/exp(2*arctan(a*x)),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{-2\operatorname{atan}(ax)} (ca^2x^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^(3/2),x)

[Out] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^(3/2), x)

3.296 $\int e^{-2\text{ArcTan}(ax)} \sqrt{c + a^2cx^2} dx$

Optimal. Leaf size=87

$$\frac{\left(\frac{2}{13} - \frac{3i}{13}\right) 2^{\frac{3}{2}+i} (1-iax)^{\frac{3}{2}-i} \sqrt{c+a^2cx^2} {}_2F_1\left(-\frac{1}{2}-i, \frac{3}{2}-i; \frac{5}{2}-i; \frac{1}{2}(1-iax)\right)}{a\sqrt{1+a^2x^2}}$$

[Out] $(-2/13+3/13*I)*2^{(3/2+I)}*(1-I*a*x)^{(3/2-I)}*\text{hypergeom}([-1/2-I, 3/2-I], [5/2-I], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^{(1/2)}/a/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$,

Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{2}{13} - \frac{3i}{13}\right) 2^{\frac{3}{2}+i} (1-iax)^{\frac{3}{2}-i} \sqrt{a^2cx^2+c} {}_2F_1\left(-\frac{1}{2}-i, \frac{3}{2}-i; \frac{5}{2}-i; \frac{1}{2}(1-iax)\right)}{a\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[\text{Sqrt}[c + a^2*c*x^2]/E^{(2*\text{ArcTan}[a*x])}, x]$

[Out] $((-2/13 + (3*I)/13)*2^{(3/2 + I)}*(1 - I*a*x)^{(3/2 - I)}*\text{Sqrt}[c + a^2*c*x^2]*\text{Hypergeometric2F1}[-1/2 - I, 3/2 - I, 5/2 - I, (1 - I*a*x)/2])/(a*\text{Sqrt}[1 + a^2*x^2])$

Rule 71

$\text{Int}[(a + b*x)^{(m+1)}/(b*(m+1)*(b*(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a+b*x)/(b*c - a*d)], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_*)])*(n_*)}*((c_*) + (d_*)*(x_*)^2)^{(p_*)}, x_Symbol] :> \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_*)])*(n_*)}*((c_*) + (d_*)*(x_*)^2)^{(p_*)}, x_Symbol] :> \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{-2 \tan^{-1}(ax)} \sqrt{c + a^2 cx^2} dx &= \frac{\sqrt{c + a^2 cx^2} \int e^{-2 \tan^{-1}(ax)} \sqrt{1 + a^2 x^2} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\sqrt{c + a^2 cx^2} \int (1 - iax)^{\frac{1}{2}-i} (1 + iax)^{\frac{1}{2}+i} dx}{\sqrt{1 + a^2 x^2}} \\ &= -\frac{\left(\frac{2}{13} - \frac{3i}{13}\right) 2^{\frac{3}{2}+i} (1 - iax)^{\frac{3}{2}-i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} - i, \frac{3}{2} - i; \frac{5}{2} - i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 87, normalized size = 1.00

$$-\frac{\left(\frac{2}{13} - \frac{3i}{13}\right) 2^{\frac{3}{2}+i} (1 - iax)^{\frac{3}{2}-i} \sqrt{c + a^2 cx^2} {}_2F_1\left(-\frac{1}{2} - i, \frac{3}{2} - i; \frac{5}{2} - i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[Sqrt[c + a^2*c*x^2]/E^(2*ArcTan[a*x]), x]

[Out] ((-2/13 + (3*I)/13)*2^(3/2 + I)*(1 - I*a*x)^(3/2 - I)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[-1/2 - I, 3/2 - I, 5/2 - I, (1 - I*a*x)/2])/(a*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \sqrt{a^2 cx^2 + c} e^{-2 \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)^(1/2)/exp(2*arctan(a*x)), x)

[Out] int((a^2*c*x^2+c)^(1/2)/exp(2*arctan(a*x)), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^(1/2)/exp(2*arctan(a*x)), x, algorithm="maxima")

[Out] integrate(sqrt(a^2*c*x^2 + c)*e^(-2*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^(1/2)/exp(2*arctan(a*x)),x, algorithm="fricas")
```

```
[Out] integral(sqrt(a^2*c*x^2 + c)*e^(-2*arctan(a*x)), x)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c(a^2x^2 + 1)} e^{-2\operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a**2*c*x**2+c)**(1/2)/exp(2*atan(a*x)),x)
```

```
[Out] Integral(sqrt(c*(a**2*x**2 + 1))*exp(-2*atan(a*x)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((a^2*c*x^2+c)^(1/2)/exp(2*arctan(a*x)),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{-2\operatorname{atan}(ax)} \sqrt{ca^2x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^(1/2),x)
```

```
[Out] int(exp(-2*atan(a*x))*(c + a^2*c*x^2)^(1/2), x)
```

$$3.297 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{\sqrt{c + a^2cx^2}} dx$$

Optimal. Leaf size=87

$$\frac{\left(\frac{2}{5} - \frac{i}{5}\right) 2^{\frac{1}{2}+i} (1 - iax)^{\frac{1}{2}-i} \sqrt{1 + a^2x^2} {}_2F_1\left(\frac{1}{2} - i, \frac{1}{2} - i; \frac{3}{2} - i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{c + a^2cx^2}}$$

[Out] $(-2/5+1/5*I)*2^{(1/2+I)}*(1-I*a*x)^{(1/2-I)}*\text{hypergeom}([1/2-I, 1/2-I], [3/2-I], 1/2-1/2*I*a*x)*(a^2*x^2+1)^{(1/2)}/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 87, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{\left(\frac{2}{5} - \frac{i}{5}\right) 2^{\frac{1}{2}+i} (1 - iax)^{\frac{1}{2}-i} \sqrt{a^2x^2 + 1} {}_2F_1\left(\frac{1}{2} - i, \frac{1}{2} - i; \frac{3}{2} - i; \frac{1}{2}(1 - iax)\right)}{a\sqrt{a^2cx^2 + c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{(2*\text{ArcTan}[a*x])})*\text{Sqrt}[c + a^2*c*x^2]], x]$

[Out] $((-2/5 + I/5)*2^{(1/2 + I)}*(1 - I*a*x)^{(1/2 - I)}*\text{Sqrt}[1 + a^2*x^2]*\text{Hypergeometric2F1}[1/2 - I, 1/2 - I, 3/2 - I, (1 - I*a*x)/2])/(a*\text{Sqrt}[c + a^2*c*x^2])$

Rule 71

$\text{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+)^2)^{(n_+)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\text{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x \&\& \text{NeQ}\{b*c - a*d, 0\} \&\& \text{IntegerQ}\{m\} \&\& \text{IntegerQ}\{n\} \&\& \text{GtQ}\{b/(b*c - a*d), 0\} \&\& (\text{RationalQ}\{m\} \mid\mid \text{!(RationalQ}\{n\} \&\& \text{GtQ}\{-d/(b*c - a*d), 0\}))$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^{(p)}, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x \&\& \text{EqQ}\{d, a^2*c\} \&\& (\text{IntegerQ}\{p\} \mid\mid \text{GtQ}\{c, 0\})$

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^{(\text{IntPart}[p])}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x \&\& \text{EqQ}\{d, a^2*c\} \&\& \text{!(IntegerQ}\{p\} \mid\mid \text{GtQ}\{c, 0\})$

Rubi steps

$$\begin{aligned}
\int \frac{e^{-2 \tan^{-1}(ax)}}{\sqrt{c+a^2cx^2}} dx &= \frac{\sqrt{1+a^2x^2} \int \frac{e^{-2 \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx}{\sqrt{c+a^2cx^2}} \\
&= \frac{\sqrt{1+a^2x^2} \int (1-iax)^{-\frac{1}{2}-i} (1+iax)^{-\frac{1}{2}+i} dx}{\sqrt{c+a^2cx^2}} \\
&= -\frac{\left(\frac{2}{5}-\frac{i}{5}\right) 2^{\frac{1}{2}+i} (1-iax)^{\frac{1}{2}-i} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}-i, \frac{1}{2}-i; \frac{3}{2}-i; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 87, normalized size = 1.00

$$-\frac{\left(\frac{2}{5}-\frac{i}{5}\right) 2^{\frac{1}{2}+i} (1-iax)^{\frac{1}{2}-i} \sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}-i, \frac{1}{2}-i; \frac{3}{2}-i; \frac{1}{2}(1-iax)\right)}{a\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(2*ArcTan[a*x])*Sqrt[c + a^2*c*x^2]), x]

[Out] ((-2/5 + I/5)*2^(1/2 + I)*(1 - I*a*x)^(1/2 - I)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 - I, 1/2 - I, 3/2 - I, (1 - I*a*x)/2])/(a*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{-2 \arctan(ax)}}{\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)

[Out] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(e^(-2*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")``[Out] integral(e^(-2*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{-2\operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c)**(1/2),x)``[Out] Integral(exp(-2*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")``[Out] sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{-2\operatorname{atan}(ax)}}{\sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^(1/2),x)``[Out] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^(1/2), x)`

$$3.298 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=38

$$-\frac{e^{-2\text{ArcTan}(ax)}(2-ax)}{5ac\sqrt{c+a^2cx^2}}$$

[Out] 1/5*(a*x-2)/a/c/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.043$, Rules used = {5177}

$$-\frac{(2-ax)e^{-2\text{ArcTan}(ax)}}{5ac\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)),x]

[Out] -1/5*(2 - a*x)/(a*c*E^(2*ArcTan[a*x])*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n + a*x)*(E^(n*ArcTan[a*x]))/(a*c*(n^2 + 1)*Sqrt[c + d*x^2]), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rubi steps

$$\int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx = -\frac{e^{-2 \tan^{-1}(ax)}(2 - ax)}{5ac\sqrt{c + a^2 cx^2}}$$

Mathematica [A]

time = 0.02, size = 37, normalized size = 0.97

$$\frac{e^{-2\text{ArcTan}(ax)}(-2+ax)}{5ac\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)),x]

[Out] (-2 + a*x)/(5*a*c*E^(2*ArcTan[a*x])*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.08, size = 41, normalized size = 1.08

method	result	size
gospers	$\frac{(a^2x^2+1)(ax-2)e^{-2\arctan(ax)}}{5a(a^2cx^2+c)^{\frac{3}{2}}}$	41

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)`[Out] $1/5*(a^2*x^2+1)*(a*x-2)/a/\exp(2*\arctan(a*x))/(a^2*c*x^2+c)^{(3/2)}$ **Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`[Out] `integrate(e^(-2*arctan(a*x))/(a^2*c*x^2 + c)^(3/2), x)`**Fricas [A]**

time = 4.47, size = 44, normalized size = 1.16

$$\frac{\sqrt{a^2cx^2 + c} (ax - 2)e^{-2\arctan(ax)}}{5(a^3c^2x^2 + ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`[Out] `1/5*sqrt(a^2*c*x^2 + c)*(a*x - 2)*e^(-2*arctan(a*x))/(a^3*c^2*x^2 + a*c^2)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{-2\operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c)**(3/2),x)`[Out] `Integral(exp(-2*atan(a*x))/(c*(a**2*x**2 + 1))**(3/2), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.18, size = 35, normalized size = 0.92

$$\frac{e^{-2\operatorname{atan}(ax)} \left(\frac{x}{5c} - \frac{2}{5ac} \right)}{\sqrt{ca^2x^2 + c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^(3/2),x)

[Out] (exp(-2*atan(a*x))*(x/(5*c) - 2/(5*a*c)))/(c + a^2*c*x^2)^(1/2)

$$3.299 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{5/2}} dx$$

Optimal. Leaf size=77

$$\frac{e^{-2\text{ArcTan}(ax)}(2-3ax)}{13ac(c+a^2cx^2)^{3/2}} - \frac{6e^{-2\text{ArcTan}(ax)}(2-ax)}{65ac^2\sqrt{c+a^2cx^2}}$$

[Out] 1/13*(3*a*x-2)/a/c/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2)-6/65*(-a*x+2)/a/c^2/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 77, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {5178, 5177}

$$\frac{6(2-ax)e^{-2\text{ArcTan}(ax)}}{65ac^2\sqrt{a^2cx^2+c}} - \frac{(2-3ax)e^{-2\text{ArcTan}(ax)}}{13ac(a^2cx^2+c)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c+a^2*c*x^2)^(5/2)),x]

[Out] -1/13*(2-3*a*x)/(a*c*E^(2*ArcTan[a*x])*(c+a^2*c*x^2)^(3/2)) - (6*(2-a*x))/(65*a*c^2*E^(2*ArcTan[a*x])*Sqrt[c+a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)/((c_.)+(d_.)*(x_)^2)^(3/2), x_Symbol] :> Simp[(n+a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2+1)*Sqrt[c+d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.)+(d_.)*(x_)^2)^(p_), x_Symbol] :> Simp[(n-2*a*(p+1)*x)*(c+d*x^2)^(p+1)*(E^(n*ArcTan[a*x])/(a*c*(n^2+4*(p+1)^2))), x] + Dist[2*(p+1)*((2*p+3)/(c*(n^2+4*(p+1)^2))), Int[(c+d*x^2)^(p+1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2+4*(p+1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{5/2}} dx = -\frac{e^{-2 \tan^{-1}(ax)}(2 - 3ax)}{13ac(c + a^2 cx^2)^{3/2}} + \frac{6 \int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx}{13c}$$

$$= -\frac{e^{-2 \tan^{-1}(ax)}(2 - 3ax)}{13ac(c + a^2 cx^2)^{3/2}} - \frac{6e^{-2 \tan^{-1}(ax)}(2 - ax)}{65ac^2 \sqrt{c + a^2 cx^2}}$$

Mathematica [A]

time = 0.03, size = 62, normalized size = 0.81

$$\frac{e^{-2 \text{ArcTan}(ax)}(-22 + 21ax - 12a^2x^2 + 6a^3x^3)}{65c^2(a + a^3x^2)\sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(5/2)), x]``[Out] (-22 + 21*a*x - 12*a^2*x^2 + 6*a^3*x^3)/(65*c^2*E^(2*ArcTan[a*x])*(a + a^3*x^2)*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.07, size = 58, normalized size = 0.75

method	result	size
gospers	$\frac{(a^2x^2+1)(6a^3x^3-12a^2x^2+21ax-22)e^{-2 \arctan(ax)}}{65a(a^2cx^2+c)^{\frac{5}{2}}}$	58

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, method=_RETURNVERBOSE)``[Out] 1/65*(a^2*x^2+1)*(6*a^3*x^3-12*a^2*x^2+21*a*x-22)/a/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2), x, algorithm="maxima")``[Out] integrate(e^(-2*arctan(a*x))/(a^2*c*x^2 + c)^(5/2), x)`

Fricas [A]

time = 4.92, size = 72, normalized size = 0.94

$$\frac{(6a^3x^3 - 12a^2x^2 + 21ax - 22)\sqrt{a^2cx^2 + c}e^{-2\arctan(ax)}}{65(a^5c^3x^4 + 2a^3c^3x^2 + ac^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="fricas")

[Out] 1/65*(6*a^3*x^3 - 12*a^2*x^2 + 21*a*x - 22)*sqrt(a^2*c*x^2 + c)*e^(-2*arctan(a*x))/(a^5*c^3*x^4 + 2*a^3*c^3*x^2 + a*c^3)

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c)**(5/2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 4371 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.63, size = 81, normalized size = 1.05

$$-\frac{e^{-2\operatorname{atan}(ax)}\left(\frac{22}{65a^3c^2} - \frac{6x^3}{65c^2} - \frac{21x}{65a^2c^2} + \frac{12x^2}{65ac^2}\right)}{\frac{\sqrt{ca^2x^2+c}}{a^2} + x^2\sqrt{ca^2x^2+c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^(5/2),x)

[Out] -(exp(-2*atan(a*x))*(22/(65*a^3*c^2) - (6*x^3)/(65*c^2) - (21*x)/(65*a^2*c^2) + (12*x^2)/(65*a*c^2)))/((c + a^2*c*x^2)^(1/2)/a^2 + x^2*(c + a^2*c*x^2)^(1/2))

$$3.300 \quad \int \frac{e^{-2\text{ArcTan}(ax)}}{(c+a^2cx^2)^{7/2}} dx$$

Optimal. Leaf size=115

$$-\frac{e^{-2\text{ArcTan}(ax)}(2-5ax)}{29ac(c+a^2cx^2)^{5/2}} - \frac{20e^{-2\text{ArcTan}(ax)}(2-3ax)}{377ac^2(c+a^2cx^2)^{3/2}} - \frac{24e^{-2\text{ArcTan}(ax)}(2-ax)}{377ac^3\sqrt{c+a^2cx^2}}$$

[Out] 1/29*(5*a*x-2)/a/c/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(5/2)-20/377*(-3*a*x+2)/a/c^2/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(3/2)-24/377*(-a*x+2)/a/c^3/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.10, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.087$, Rules used = {5178, 5177}

$$-\frac{24(2-ax)e^{-2\text{ArcTan}(ax)}}{377ac^3\sqrt{a^2cx^2+c}} - \frac{20(2-3ax)e^{-2\text{ArcTan}(ax)}}{377ac^2(a^2cx^2+c)^{3/2}} - \frac{(2-5ax)e^{-2\text{ArcTan}(ax)}}{29ac(a^2cx^2+c)^{5/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(7/2)), x]

[Out] -1/29*(2 - 5*a*x)/(a*c*E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(5/2)) - (20*(2 - 3*a*x))/(377*a*c^2*E^(2*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)) - (24*(2 - a*x))/(377*a*c^3*E^(2*ArcTan[a*x])*Sqrt[c + a^2*c*x^2])

Rule 5177

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2)^(3/2), x_Symbol] := Simp[(n + a*x)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 1)*Sqrt[c + d*x^2])), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && !IntegerQ[I*n]

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rubi steps

$$\begin{aligned}
\int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{7/2}} dx &= -\frac{e^{-2 \tan^{-1}(ax)}(2 - 5ax)}{29ac(c + a^2 cx^2)^{5/2}} + \frac{20 \int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{5/2}} dx}{29c} \\
&= -\frac{e^{-2 \tan^{-1}(ax)}(2 - 5ax)}{29ac(c + a^2 cx^2)^{5/2}} - \frac{20e^{-2 \tan^{-1}(ax)}(2 - 3ax)}{377ac^2(c + a^2 cx^2)^{3/2}} + \frac{120 \int \frac{e^{-2 \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx}{377c^2} \\
&= -\frac{e^{-2 \tan^{-1}(ax)}(2 - 5ax)}{29ac(c + a^2 cx^2)^{5/2}} - \frac{20e^{-2 \tan^{-1}(ax)}(2 - 3ax)}{377ac^2(c + a^2 cx^2)^{3/2}} - \frac{24e^{-2 \tan^{-1}(ax)}(2 - ax)}{377ac^3 \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 81, normalized size = 0.70

$$\frac{e^{-2 \operatorname{ArcTan}(ax)}(-114 + 149ax - 136a^2x^2 + 108a^3x^3 - 48a^4x^4 + 24a^5x^5)}{377ac^3(1 + a^2x^2)^2 \sqrt{c + a^2cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^(2*ArcTan[a*x]))*(c + a^2*c*x^2)^(7/2), x]``[Out] (-114 + 149*a*x - 136*a^2*x^2 + 108*a^3*x^3 - 48*a^4*x^4 + 24*a^5*x^5)/(377*a*c^3*E^(2*ArcTan[a*x]))*(1 + a^2*x^2)^2*Sqrt[c + a^2*c*x^2]`**Maple [A]**

time = 0.07, size = 74, normalized size = 0.64

method	result	size
gospers	$\frac{(a^2x^2+1)(24a^5x^5-48a^4x^4+108a^3x^3-136a^2x^2+149ax-114)e^{-2 \arctan(ax)}}{377a(a^2cx^2+c)^{\frac{7}{2}}}$	74

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, method=_RETURNVERBOSE)``[Out] 1/377*(a^2*x^2+1)*(24*a^5*x^5-48*a^4*x^4+108*a^3*x^3-136*a^2*x^2+149*a*x-114)/a/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2), x, algorithm="maxima")`

[Out] integrate($e^{-2\arctan(ax)}/(a^2cx^2 + c)^{7/2}$, x)

Fricas [A]

time = 4.17, size = 99, normalized size = 0.86

$$\frac{(24a^5x^5 - 48a^4x^4 + 108a^3x^3 - 136a^2x^2 + 149ax - 114)\sqrt{a^2cx^2 + c}e^{-2\arctan(ax)}}{377(a^7c^4x^6 + 3a^5c^4x^4 + 3a^3c^4x^2 + ac^4)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2),x, algorithm="fricas")

[Out] 1/377*(24*a^5*x^5 - 48*a^4*x^4 + 108*a^3*x^3 - 136*a^2*x^2 + 149*a*x - 114)*sqrt(a^2*c*x^2 + c)*e^(-2*arctan(a*x))/(a^7*c^4*x^6 + 3*a^5*c^4*x^4 + 3*a^3*c^4*x^2 + a*c^4)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*atan(a*x))/(a**2*c*x**2+c)**(7/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/exp(2*arctan(a*x))/(a^2*c*x^2+c)^(7/2),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.66, size = 123, normalized size = 1.07

$$-\frac{e^{-2\operatorname{atan}(ax)}\left(\frac{114}{377a^5c^3} - \frac{24x^5}{377c^3} - \frac{149x}{377a^4c^3} + \frac{48x^4}{377ac^3} - \frac{108x^3}{377a^2c^3} + \frac{136x^2}{377a^3c^3}\right)}{\sqrt{\frac{ca^2x^2+c}{a^4}} + x^4\sqrt{ca^2x^2+c} + \frac{2x^2\sqrt{ca^2x^2+c}}{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(-2*atan(a*x))/(c + a^2*c*x^2)^(7/2),x)

[Out] -(exp(-2*atan(a*x))*(114/(377*a^5*c^3) - (24*x^5)/(377*c^3) - (149*x)/(377*a^4*c^3) + (48*x^4)/(377*a*c^3) - (108*x^3)/(377*a^2*c^3) + (136*x^2)/(377*a^3*c^3)))/((c + a^2*c*x^2)^(1/2)/a^4 + x^4*(c + a^2*c*x^2)^(1/2) + (2*x^2*(c + a^2*c*x^2)^(1/2))/a^2)

$$3.301 \quad \int \frac{e^{5i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=50

$$-\frac{2i}{a(1-iax)^2} + \frac{4i}{a(1-iax)} + \frac{i \log(i+ax)}{a}$$

[Out] $-2*I/a/(1-I*a*x)^2+4*I/a/(1-I*a*x)+I*\ln(I+a*x)/a$

Rubi [A]

time = 0.04, antiderivative size = 50, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 45}

$$\frac{4i}{a(1-iax)} - \frac{2i}{a(1-iax)^2} + \frac{i \log(ax+i)}{a}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((5*I)*\text{ArcTan}[a*x])}/\text{Sqrt}[1+a^2*x^2],x]$

[Out] $(-2*I)/(a*(1-I*a*x)^2) + (4*I)/(a*(1-I*a*x)) + (I*\text{Log}[I+a*x])/a$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{5i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{(1+iax)^2}{(1-iax)^3} dx \\ &= \int \left(\frac{4}{(1-iax)^3} - \frac{4}{(1-iax)^2} + \frac{1}{1-iax} \right) dx \\ &= -\frac{2i}{a(1-iax)^2} + \frac{4i}{a(1-iax)} + \frac{i \log(i+ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 42, normalized size = 0.84

$$\frac{i(-2 + 4iax + (i + ax)^2 \log(i + ax))}{a(i + ax)^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((5*I)*ArcTan[a*x])/Sqrt[1 + a^2*x^2], x]

[Out] (I*(-2 + (4*I)*a*x + (I + a*x)^2*Log[I + a*x]))/(a*(I + a*x)^2)

Maple [A]

time = 0.08, size = 34, normalized size = 0.68

method	result
default	$\frac{-4x - \frac{2i}{a}}{(ax+i)^2} + \frac{i \ln(ax+i)}{a}$
risch	$\frac{-4x - \frac{2i}{a}}{(ax+i)^2} + \frac{i \ln(a^2x^2+1)}{2a} + \frac{\arctan(ax)}{a}$
meijerg	$\frac{x\sqrt{a^2} (3a^2x^2+5) + \sqrt[3]{a^2} \arctan(ax)}{2(a^2x^2+1)^2} + \frac{5iax^2(a^2x^2+2)}{4(a^2x^2+1)^2} - \frac{5 \left(-\frac{x(a^2)^{\frac{3}{2}}(-3a^2x^2+3)}{6a^2(a^2x^2+1)^2} + \frac{(a^2)^{\frac{3}{2}} \arctan(ax)}{2a^3} \right)}{2\sqrt{a^2}} - \frac{5ia^3x^4}{2(a^2x^2+1)^2} + \dots$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^5/(a^2*x^2+1)^3,x,method=_RETURNVERBOSE)

[Out] (-4*x-2*I/a)/(I+a*x)^2+I*ln(I+a*x)/a

Maxima [A]

time = 0.48, size = 63, normalized size = 1.26

$$-\frac{2(2a^3x^3 - 3ia^2x^2 - i)}{a^5x^4 + 2a^3x^2 + a} + \frac{\arctan(ax)}{a} + \frac{i \log(a^2x^2 + 1)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^3,x, algorithm="maxima")

[Out] -2*(2*a^3*x^3 - 3*I*a^2*x^2 - I)/(a^5*x^4 + 2*a^3*x^2 + a) + arctan(a*x)/a + 1/2*I*log(a^2*x^2 + 1)/a

Fricas [A]

time = 3.26, size = 53, normalized size = 1.06

$$-\frac{4ax - (ia^2x^2 - 2ax - i) \log\left(\frac{ax+i}{a}\right) + 2i}{a^3x^2 + 2ia^2x - a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^3,x, algorithm="fricas")

[Out] $-(4ax - (Ia^2x^2 - 2ax - I)\log((ax + I)/a) + 2I)/(a^3x^2 + 2Ia^2x - a)$

Sympy [A]

time = 0.16, size = 36, normalized size = 0.72

$$\frac{-4ax - 2i}{a^3x^2 + 2ia^2x - a} + \frac{i \log(ax + i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**5/(a**2*x**2+1)**3,x)

[Out] $(-4ax - 2I)/(a^3x^2 + 2Ia^2x - a) + I\log(ax + I)/a$

Giac [A]

time = 0.41, size = 30, normalized size = 0.60

$$\frac{i \log(ax + i)}{a} - \frac{2(2ax + i)}{(ax + i)^2 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^3,x, algorithm="giac")

[Out] $I\log(ax + I)/a - 2*(2ax + I)/((ax + I)^2 a)$

Mupad [B]

time = 0.13, size = 49, normalized size = 0.98

$$\frac{\ln\left(x + \frac{1i}{a}\right) 1i}{a} - \frac{\frac{4x}{a^2} + \frac{2i}{a^3}}{x^2 - \frac{1}{a^2} + \frac{x2i}{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^5/(a^2*x^2 + 1)^3,x)

[Out] $(\log(x + 1i/a)*1i)/a - ((4*x)/a^2 + 2i/a^3)/((x*2i)/a - 1/a^2 + x^2)$

$$3.302 \quad \int \frac{e^{4i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=73

$$\frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \frac{2i(1+iax)^{3/2}}{3a(1-iax)^{3/2}} + \frac{\sinh^{-1}(ax)}{a}$$

[Out] $-2/3*I*(1+I*a*x)^{(3/2)}/a/(1-I*a*x)^{(3/2)}+\operatorname{arcsinh}(a*x)/a+2*I*(1+I*a*x)^{(1/2)}/a/(1-I*a*x)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5181, 49, 41, 221}

$$-\frac{2i(1+iax)^{3/2}}{3a(1-iax)^{3/2}} + \frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} + \frac{\sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((4*I)*\operatorname{ArcTan}[a*x])}/\operatorname{Sqrt}[1+a^2*x^2],x]$

[Out] $((2*I)*\operatorname{Sqrt}[1+I*a*x])/(a*\operatorname{Sqrt}[1-I*a*x]) - (((2*I)/3)*(1+I*a*x)^{(3/2)})/(a*(1-I*a*x)^{(3/2)}) + \operatorname{ArcSinh}[a*x]/a$

Rule 41

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] :> \operatorname{Int}[(a*c + b*d*x^2)^m, x] /;$ FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))

Rule 49

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] :> \operatorname{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^n/(b*(m+1))), x] - \operatorname{Dist}[d*(n/(b*(m+1))), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}, x], x] /;$ FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(IleQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_+ + (b_+)*(x_+)^2], x_Symbol] :> \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /;$ FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]

Rule 5181

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :>
Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /
; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{(1+iax)^{3/2}}{(1-iax)^{5/2}} dx \\ &= -\frac{2i(1+iax)^{3/2}}{3a(1-iax)^{3/2}} - \int \frac{\sqrt{1+iax}}{(1-iax)^{3/2}} dx \\ &= \frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \frac{2i(1+iax)^{3/2}}{3a(1-iax)^{3/2}} + \int \frac{1}{\sqrt{1-iax}\sqrt{1+iax}} dx \\ &= \frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \frac{2i(1+iax)^{3/2}}{3a(1-iax)^{3/2}} + \int \frac{1}{\sqrt{1+a^2x^2}} dx \\ &= \frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \frac{2i(1+iax)^{3/2}}{3a(1-iax)^{3/2}} + \frac{\sinh^{-1}(ax)}{a} \end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.01, size = 48, normalized size = 0.66

$$-\frac{4i\sqrt{2} {}_2F_1\left(-\frac{3}{2}, -\frac{3}{2}; -\frac{1}{2}; \frac{1}{2}(1-iax)\right)}{3a(1-iax)^{3/2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((4*I)*ArcTan[a*x])/Sqrt[1 + a^2*x^2], x]

[Out] (((-4*I)/3)*Sqrt[2]*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 - I*a*x)/2])/(a*(1 - I*a*x)^(3/2))

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 223 vs. 2(57) = 114.

time = 0.09, size = 224, normalized size = 3.07

method	result
meijerg	$\frac{x(2a^2x^2+3)}{3(a^2x^2+1)^{\frac{3}{2}}} + \frac{8i\left(\frac{\sqrt{\pi}}{2} - \frac{\sqrt{\pi}}{2(a^2x^2+1)^{\frac{3}{2}}}\right)}{3a\sqrt{\pi}} - \frac{2a^2x^3}{(a^2x^2+1)^{\frac{3}{2}}} - \frac{8i\left(\sqrt{\pi} - \frac{\sqrt{\pi}(12a^2x^2+8)}{8(a^2x^2+1)^{\frac{3}{2}}}\right)}{3a\sqrt{\pi}} + \frac{-\sqrt{\pi}x(a^2)^{\frac{5}{2}}(20a^2x^2+15)}{15a^4(a^2x^2+1)^{\frac{3}{2}}\sqrt{\pi}}$

default	$\frac{x}{3(a^2x^2+1)^{\frac{3}{2}}} + \frac{2x}{3\sqrt{a^2x^2+1}} + a^4 \left(-\frac{x^3}{3a^2(a^2x^2+1)^{\frac{3}{2}}} + \frac{-\frac{x}{a^2\sqrt{a^2x^2+1}} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{a^2\sqrt{a^2}}}{a^2} \right) - 4ia^3$
---------	--

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^4/(a^2*x^2+1)^(5/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{3}x/(a^2x^2+1)^{(3/2)} + \frac{2}{3}x/(a^2x^2+1)^{(1/2)} + a^4 \left(-\frac{1}{3}x^3/a^2/(a^2x^2+1)^{(3/2)} + \frac{1}{a^2} \left(-\frac{x}{a^2\sqrt{a^2x^2+1}} + \frac{\ln(a^2x/(a^2)^{(1/2)} + (a^2x^2+1)^{(1/2)})}{(a^2)^{(1/2)}} \right) - \frac{4Ia^3(-x^2/a^2/(a^2x^2+1)^{(3/2)} - 2/3/a^4/(a^2x^2+1)^{(3/2)})}{(a^2)^{(1/2)}} - \frac{6a^2(-1/2*x/a^2/(a^2x^2+1)^{(3/2)} + 1/2/a^2*(1/3*x/(a^2x^2+1)^{(3/2)} + 2/3*x/(a^2x^2+1)^{(1/2)})}{(a^2)^{(1/2)}} - \frac{4}{3}I/a/(a^2x^2+1)^{(3/2)} \right)$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 112 vs. $2(51) = 102$.

time = 0.27, size = 112, normalized size = 1.53

$$-\frac{1}{3}a^4x \left(\frac{3x^2}{(a^2x^2+1)^{\frac{3}{2}}a^2} + \frac{2}{(a^2x^2+1)^{\frac{3}{2}}a^4} \right) + \frac{4iax^2}{(a^2x^2+1)^{\frac{3}{2}}} - \frac{5x}{3\sqrt{a^2x^2+1}} + \frac{\operatorname{arsinh}(ax)}{a} + \frac{7x}{3(a^2x^2+1)^{\frac{3}{2}}} + \frac{4i}{3(a^2x^2+1)^{\frac{3}{2}}a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^4/(a^2*x^2+1)^(5/2),x, algorithm="maxima")`

[Out] $-1/3*a^4*x*(3*x^2/((a^2*x^2+1)^{(3/2)}*a^2) + 2/((a^2*x^2+1)^{(3/2)}*a^4)) + 4*I*a*x^2/(a^2*x^2+1)^{(3/2)} - 5/3*x/\operatorname{sqrt}(a^2*x^2+1) + \operatorname{arcsinh}(a*x)/a + 7/3*x/(a^2*x^2+1)^{(3/2)} + 4/3*I/((a^2*x^2+1)^{(3/2)}*a)$

Fricas [A]

time = 2.73, size = 86, normalized size = 1.18

$$\frac{8a^2x^2 + 16iax + 3(a^2x^2 + 2iax - 1) \log(-ax + \sqrt{a^2x^2 + 1}) + 4\sqrt{a^2x^2 + 1}(2ax + i) - 8}{3(a^3x^2 + 2ia^2x - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^4/(a^2*x^2+1)^(5/2),x, algorithm="fricas")`

[Out] $-1/3*(8*a^2*x^2 + 16*I*a*x + 3*(a^2*x^2 + 2*I*a*x - 1)*\log(-a*x + \operatorname{sqrt}(a^2*x^2 + 1))) + 4*\operatorname{sqrt}(a^2*x^2 + 1)*(2*a*x + I) - 8)/(a^3*x^2 + 2*I*a^2*x - a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(ax - i)^4}{(a^2x^2 + 1)^{\frac{5}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**(5/2),x)

[Out] Integral((a*x - I)**4/(a**2*x**2 + 1)**(5/2), x)

Giac [A]

time = 0.46, size = 24, normalized size = 0.33

$$-\frac{\log\left(-x|a| + \sqrt{a^2x^2 + 1}\right)}{|a|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^(5/2),x, algorithm="giac")

[Out] -log(-x*abs(a) + sqrt(a^2*x^2 + 1))/abs(a)

Mupad [B]

time = 0.53, size = 92, normalized size = 1.26

$$\frac{\operatorname{asinh}\left(x\sqrt{a^2}\right)}{\sqrt{a^2}} - \frac{8\sqrt{a^2x^2+1}}{3\left(x\sqrt{a^2} + \frac{\sqrt{a^2}1i}{a}\right)\sqrt{a^2}} + \frac{a\sqrt{a^2x^2+1}4i}{3(a^4x^2 + a^3x2i - a^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(a^2*x^2 + 1)^(5/2),x)

[Out] asinh(x*(a^2)^(1/2))/(a^2)^(1/2) - (8*(a^2*x^2 + 1)^(1/2))/(3*(((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2)) + (a*(a^2*x^2 + 1)^(1/2)*4i)/(3*(a^3*x*2i - a^2 + a^4*x^2))

$$3.303 \quad \int \frac{e^{3i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=30

$$\frac{2}{a(i+ax)} - \frac{i \log(i+ax)}{a}$$

[Out] 2/a/(I+a*x)-I*ln(I+a*x)/a

Rubi [A]

time = 0.03, antiderivative size = 30, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 45}

$$\frac{2}{a(ax+i)} - \frac{i \log(ax+i)}{a}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a*x])/Sqrt[1 + a^2*x^2],x]

[Out] 2/(a*(I + a*x)) - (I*Log[I + a*x])/a

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])]

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{3i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{1+iax}{(1-iax)^2} dx \\ &= \int \left(-\frac{2}{(i+ax)^2} - \frac{i}{i+ax} \right) dx \\ &= \frac{2}{a(i+ax)} - \frac{i \log(i+ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 30, normalized size = 1.00

$$\frac{2}{a(i+ax)} - \frac{i \log(i+ax)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x])/Sqrt[1 + a^2*x^2], x]

[Out] 2/(a*(I + a*x)) - (I*Log[I + a*x])/a

Maple [A]

time = 0.08, size = 28, normalized size = 0.93

method	result	size
default	$\frac{2}{a(ax+i)} - \frac{i \ln(ax+i)}{a}$	28
risch	$\frac{2}{a(ax+i)} - \frac{i \ln(a^2x^2+1)}{2a} - \frac{\arctan(ax)}{a}$	40
meijerg	$\frac{\frac{2x\sqrt{a^2}}{2a^2x^2+2} + \frac{\sqrt{a^2}}{a} \arctan(ax)}{2\sqrt{a^2}} + \frac{3iax^2}{2(a^2x^2+1)} - \frac{3\left(-\frac{x(a^2)^{\frac{3}{2}}}{a^2(a^2x^2+1)} + \frac{(a^2)^{\frac{3}{2}} \arctan(ax)}{a^3}\right)}{2\sqrt{a^2}} - \frac{i\left(-\frac{a^2x^2}{a^2x^2+1} + \ln(a^2x^2+1)\right)}{2a}$	140

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^2,x,method=_RETURNVERBOSE)

[Out] 2/a/(I+a*x)-I*ln(I+a*x)/a

Maxima [A]

time = 0.46, size = 43, normalized size = 1.43

$$\frac{2(ax-i)}{a^3x^2+a} - \frac{\arctan(ax)}{a} - \frac{i \log(a^2x^2+1)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^2,x, algorithm="maxima")

[Out] 2*(a*x - I)/(a^3*x^2 + a) - arctan(a*x)/a - 1/2*I*log(a^2*x^2 + 1)/a

Fricas [A]

time = 3.59, size = 31, normalized size = 1.03

$$\frac{(-i ax + 1) \log\left(\frac{ax+i}{a}\right) + 2}{a^2x + ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^2,x, algorithm="fricas")

[Out] ((-I*a*x + 1)*log((a*x + I)/a) + 2)/(a^2*x + I*a)

Sympy [A]

time = 0.08, size = 19, normalized size = 0.63

$$\frac{2}{a^2x + ia} - \frac{i \log(ax + i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**2,x)

[Out] 2/(a**2*x + I*a) - I*log(a*x + I)/a

Giac [A]

time = 0.41, size = 24, normalized size = 0.80

$$-\frac{i \log(ax + i)}{a} + \frac{2}{(ax + i)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^2,x, algorithm="giac")

[Out] -I*log(a*x + I)/a + 2/((a*x + I)*a)

Mupad [B]

time = 0.49, size = 28, normalized size = 0.93

$$\frac{2}{x a^2 + a \operatorname{li}} - \frac{\ln(ax + \operatorname{li}) \operatorname{li}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^3/(a^2*x^2 + 1)^2,x)

[Out] 2/(a*1i + a^2*x) - (log(a*x + 1i)*1i)/a

$$3.304 \quad \int \frac{e^{2i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=41

$$-\frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \frac{\sinh^{-1}(ax)}{a}$$

[Out] $-\operatorname{arcsinh}(a*x)/a-2*I*(1+I*a*x)^{(1/2)}/a/(1-I*a*x)^{(1/2)}$

Rubi [A]

time = 0.02, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5181, 49, 41, 221}

$$-\frac{\sinh^{-1}(ax)}{a} - \frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((2*I)*\operatorname{ArcTan}[a*x])}/\operatorname{Sqrt}[1+a^2*x^2],x]$

[Out] $((-2*I)*\operatorname{Sqrt}[1+I*a*x])/(a*\operatorname{Sqrt}[1-I*a*x]) - \operatorname{ArcSinh}[a*x]/a$

Rule 41

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Int}[(a*c + b*d*x^2)^m, x] /; \operatorname{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \operatorname{EqQ}[b*c + a*d, 0] \ \&\& (\operatorname{IntegerQ}[m] \ || \ (\operatorname{GtQ}[a, 0] \ \&\& \operatorname{GtQ}[c, 0]))$

Rule 49

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^n/(b*(m+1))), x] - \operatorname{Dist}[d*(n/(b*(m+1))), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}, x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{GtQ}[n, 0] \ \&\& \operatorname{LtQ}[m, -1] \ \&\& !(\operatorname{IntegerQ}[n] \ \&\& !\operatorname{IntegerQ}[m]) \ \&\& !(I\operatorname{LeQ}[m+n+2, 0] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n+m+1, 0])) \ \& \ \& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_+) + (b_+)*(x_+)^2], x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b]$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_+)*(x_+)])^{(n_+)}}*((c_+) + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1 - I*a*x)^{(p+I*(n/2))}*(1 + I*a*x)^{(p-I*(n/2))}, x], x] /$

; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
 \int \frac{e^{2i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{\sqrt{1+iax}}{(1-iax)^{3/2}} dx \\
 &= -\frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \int \frac{1}{\sqrt{1-iax}\sqrt{1+iax}} dx \\
 &= -\frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \int \frac{1}{\sqrt{1+a^2x^2}} dx \\
 &= -\frac{2i\sqrt{1+iax}}{a\sqrt{1-iax}} - \frac{\sinh^{-1}(ax)}{a}
 \end{aligned}$$

Mathematica [A]

time = 0.02, size = 52, normalized size = 1.27

$$\frac{2i \left(\frac{\sqrt{1+iax}}{\sqrt{1-iax}} + \text{ArcSin} \left(\frac{\sqrt{1-iax}}{\sqrt{2}} \right) \right)}{a}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((2*I)*ArcTan[a*x])/Sqrt[1 + a^2*x^2], x]

[Out] ((-2*I)*(Sqrt[1 + I*a*x]/Sqrt[1 - I*a*x] + ArcSin[Sqrt[1 - I*a*x]/Sqrt[2]]))/a

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 86 vs. 2(34) = 68.

time = 0.07, size = 87, normalized size = 2.12

method	result	size
default	$ \frac{x}{\sqrt{a^2x^2+1}} - a^2 \left(-\frac{x}{a^2\sqrt{a^2x^2+1}} + \frac{\ln\left(\frac{a^2x}{\sqrt{a^2}} + \sqrt{a^2x^2+1}\right)}{a^2\sqrt{a^2}} \right) - \frac{2i}{a\sqrt{a^2x^2+1}} $	87
meijerg	$ \frac{x}{\sqrt{a^2x^2+1}} + \frac{2i \left(\sqrt{\pi} - \frac{\sqrt{\pi}}{\sqrt{a^2x^2+1}} \right)}{a\sqrt{\pi}} - \frac{\sqrt{\pi} x (a^2)^{\frac{3}{2}}}{a^2\sqrt{a^2x^2+1}} + \frac{\sqrt{\pi} (a^2)^{\frac{3}{2}} \text{arcsinh}(ax)}{a^3\sqrt{\pi}\sqrt{a^2}} $	96

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^2/(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $x/(a^2x^2+1)^{1/2}-a^2(-x/a^2/(a^2x^2+1)^{1/2}+1/a^2\ln(a^2x/(a^2)^{1/2})+(a^2x^2+1)^{1/2})/(a^2)^{1/2})-2*I/a/(a^2x^2+1)^{1/2}$

Maxima [A]

time = 0.25, size = 40, normalized size = 0.98

$$\frac{2x}{\sqrt{a^2x^2+1}} - \frac{\operatorname{arsinh}(ax)}{a} - \frac{2i}{\sqrt{a^2x^2+1}a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^2/(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out] $2*x/\sqrt{a^2*x^2+1} - \operatorname{arcsinh}(a*x)/a - 2*I/(\sqrt{a^2*x^2+1}*a)$

Fricas [A]

time = 4.77, size = 54, normalized size = 1.32

$$\frac{2ax + (ax + i) \log(-ax + \sqrt{a^2x^2 + 1}) + 2\sqrt{a^2x^2 + 1} + 2i}{a^2x + ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^2/(a^2*x^2+1)^(3/2),x, algorithm="fricas")`

[Out] $(2*a*x + (a*x + I)*\log(-a*x + \sqrt{a^2*x^2 + 1}) + 2*\sqrt{a^2*x^2 + 1} + 2*I)/(a^2*x + I*a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{a^2x^2}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx - \int \left(-\frac{2iax}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx - \int \left(-\frac{1}{a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**2/(a**2*x**2+1)**(3/2),x)`

[Out] $-\operatorname{Integral}(a**2*x**2/(a**2*x**2*\sqrt{a**2*x**2+1} + \sqrt{a**2*x**2+1}), x) - \operatorname{Integral}(-2*I*a*x/(a**2*x**2*\sqrt{a**2*x**2+1} + \sqrt{a**2*x**2+1}), x) - \operatorname{Integral}(-1/(a**2*x**2*\sqrt{a**2*x**2+1} + \sqrt{a**2*x**2+1}), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)^(3/2),x, algorithm="giac")

[Out] undef

Mupad [B]

time = 0.49, size = 55, normalized size = 1.34

$$-\frac{\operatorname{asinh}\left(x\sqrt{a^2}\right)}{\sqrt{a^2}} + \frac{2\sqrt{a^2x^2+1}}{\left(x\sqrt{a^2} + \frac{\sqrt{a^2}1i}{a}\right)\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/(a^2*x^2 + 1)^(3/2),x)

[Out] (2*(a^2*x^2 + 1)^(1/2))/((((a^2)^(1/2)*1i)/a + x*(a^2)^(1/2))*(a^2)^(1/2))
- asinh(x*(a^2)^(1/2))/(a^2)^(1/2)

$$3.305 \quad \int \frac{e^{i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=15

$$\frac{i \log(i+ax)}{a}$$

[Out] I*ln(I+a*x)/a

Rubi [A]

time = 0.02, antiderivative size = 15, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 31}

$$\frac{i \log(ax+i)}{a}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])/Sqrt[1 + a^2*x^2],x]

[Out] (I*Log[I + a*x])/a

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{1}{1-iax} dx \\ &= \frac{i \log(i+ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 15, normalized size = 1.00

$$\frac{i \log(i+ax)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a*x])/Sqrt[1 + a^2*x^2],x]

[Out] (I*Log[I + a*x])/a

Maple [A]

time = 0.06, size = 26, normalized size = 1.73

method	result	size
default	$\frac{i \ln(a^2 x^2 + 1)}{2a} + \frac{\arctan(ax)}{a}$	26
meijerg	$\frac{i \ln(a^2 x^2 + 1)}{2a} + \frac{\arctan(ax)}{a}$	26
risch	$\frac{i \ln(a^2 x^2 + 1)}{2a} + \frac{\arctan(ax)}{a}$	26

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)/(a^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] 1/2*I/a*ln(a^2*x^2+1)+arctan(a*x)/a

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 24 vs. 2(11) = 22.

time = 0.46, size = 24, normalized size = 1.60

$$\frac{\arctan(ax)}{a} + \frac{i \log(a^2 x^2 + 1)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1),x, algorithm="maxima")

[Out] arctan(a*x)/a + 1/2*I*log(a^2*x^2 + 1)/a

Fricas [A]

time = 3.08, size = 15, normalized size = 1.00

$$\frac{i \log\left(\frac{ax+i}{a}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1),x, algorithm="fricas")

[Out] I*log((a*x + I)/a)/a

Sympy [A]

time = 0.01, size = 8, normalized size = 0.53

$$\frac{i \log(ax + i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1),x)

[Out] I*log(a*x + I)/a

Giac [A]

time = 0.45, size = 12, normalized size = 0.80

$$\frac{i \log(-i a x + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1),x, algorithm="giac")

[Out] I*log(-I*a*x + 1)/a

Mupad [B]

time = 0.47, size = 15, normalized size = 1.00

$$\frac{\ln\left(x + \frac{1i}{a}\right) 1i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)/(a^2*x^2 + 1),x)

[Out] (log(x + 1i/a)*1i)/a

$$3.306 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=16

$$-\frac{i \log(i - ax)}{a}$$

[Out] -I*ln(I-a*x)/a

Rubi [A]

time = 0.02, antiderivative size = 16, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 31}

$$-\frac{i \log(-ax + i)}{a}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a*x])*Sqrt[1 + a^2*x^2]),x]

[Out] ((-I)*Log[I - a*x])/a

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{1}{1+iax} dx \\ &= -\frac{i \log(i - ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 16, normalized size = 1.00

$$-\frac{i \log(i - ax)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(I*ArcTan[a*x])*Sqrt[1 + a^2*x^2]),x]

[Out] ((-I)*Log[I - a*x])/a

Maple [A]

time = 0.07, size = 15, normalized size = 0.94

method	result	size
default	$-\frac{i \ln(iax+1)}{a}$	15
meijerg	$-\frac{i \ln(iax+1)}{a}$	15
risch	$-\frac{i \ln(a^2x^2+1)}{2a} + \frac{\arctan(ax)}{a}$	26

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x),x,method=_RETURNVERBOSE)

[Out] -I*ln(1+I*a*x)/a

Maxima [A]

time = 0.25, size = 12, normalized size = 0.75

$$-\frac{i \log(iax + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x),x, algorithm="maxima")

[Out] -I*log(I*a*x + 1)/a

Fricas [A]

time = 3.22, size = 15, normalized size = 0.94

$$-\frac{i \log\left(\frac{ax-i}{a}\right)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x),x, algorithm="fricas")

[Out] -I*log((a*x - I)/a)/a

Sympy [A]

time = 0.01, size = 10, normalized size = 0.62

$$-\frac{i \log(ax - i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x),x)

[Out] -I*log(a*x - I)/a

Giac [A]

time = 0.42, size = 12, normalized size = 0.75

$$-\frac{i \log(i a x + 1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x),x, algorithm="giac")

[Out] -I*log(I*a*x + 1)/a

Mupad [B]

time = 0.48, size = 15, normalized size = 0.94

$$-\frac{\ln\left(x - \frac{1i}{a}\right) 1i}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(a*x*1i + 1),x)

[Out] -(log(x - 1i/a)*1i)/a

$$3.307 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=41

$$\frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} - \frac{\sinh^{-1}(ax)}{a}$$

[Out] $-\operatorname{arcsinh}(a*x)/a+2*I*(1-I*a*x)^{(1/2)}/a/(1+I*a*x)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 41, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5181, 49, 41, 221}

$$-\frac{\sinh^{-1}(ax)}{a} + \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(E^{((2*I)*\operatorname{ArcTan}[a*x])})*\operatorname{Sqrt}[1+a^2*x^2]),x]$

[Out] $((2*I)*\operatorname{Sqrt}[1-I*a*x])/(a*\operatorname{Sqrt}[1+I*a*x]) - \operatorname{ArcSinh}[a*x]/a$

Rule 41

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Int}[(a*c + b*d*x^2)^m, x] /; \operatorname{FreeQ}\{a, b, c, d, m\}, x] \ \&\& \operatorname{EqQ}[b*c + a*d, 0] \ \&\& (\operatorname{IntegerQ}[m] \ || \ (\operatorname{GtQ}[a, 0] \ \&\& \operatorname{GtQ}[c, 0]))$

Rule 49

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+) + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)}*((c + d*x)^n/(b*(m+1))), x] - \operatorname{Dist}[d*(n/(b*(m+1))), \operatorname{Int}[(a + b*x)^{(m+1)}*(c + d*x)^{(n-1)}, x], x] /; \operatorname{FreeQ}\{a, b, c, d\}, x] \ \&\& \operatorname{NeQ}[b*c - a*d, 0] \ \&\& \operatorname{GtQ}[n, 0] \ \&\& \operatorname{LtQ}[m, -1] \ \&\& !(\operatorname{IntegerQ}[n] \ \&\& !\operatorname{IntegerQ}[m]) \ \&\& !(I\operatorname{LeQ}[m+n+2, 0] \ \&\& (\operatorname{FractionQ}[m] \ || \operatorname{GeQ}[2*n+m+1, 0])) \ \& \ \& \operatorname{IntLinearQ}[a, b, c, d, m, n, x]$

Rule 221

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_+) + (b_+)*(x_+)^2], x_Symbol] \rightarrow \operatorname{Simp}[\operatorname{ArcSinh}[\operatorname{Rt}[b, 2]*(x/\operatorname{Sqrt}[a])]/\operatorname{Rt}[b, 2], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{GtQ}[a, 0] \ \&\& \operatorname{PosQ}[b]$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_+)*(x_+)])^{(n_+)}}*((c_+) + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1 - I*a*x)^{(p+I*(n/2))}*(1 + I*a*x)^{(p-I*(n/2))}, x], x] /$

; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{\sqrt{1-iax}}{(1+iax)^{3/2}} dx \\ &= \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} - \int \frac{1}{\sqrt{1-iax}\sqrt{1+iax}} dx \\ &= \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} - \int \frac{1}{\sqrt{1+a^2x^2}} dx \\ &= \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} - \frac{\sinh^{-1}(ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 56, normalized size = 1.37

$$\frac{2\left(\sqrt{1+a^2x^2} + (-1-iax)\text{ArcSin}\left(\frac{\sqrt{1-iax}}{\sqrt{2}}\right)\right)}{a(-i+ax)}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((2*I)*ArcTan[a*x])*Sqrt[1 + a^2*x^2]),x]

[Out] (2*(Sqrt[1 + a^2*x^2] + (-1 - I*a*x)*ArcSin[Sqrt[1 - I*a*x]/Sqrt[2]]))/(a*(-I + a*x))

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 148 vs. 2(34) = 68.

time = 0.08, size = 149, normalized size = 3.63

method	result
default	$-\frac{i\left(\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)\right)^{\frac{3}{2}}}{a\left(x-\frac{i}{a}\right)^2} - ia \left(\sqrt{\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)} + \frac{ia \ln\left(\frac{ia + \left(x-\frac{i}{a}\right)a^2 + \sqrt{\left(x-\frac{i}{a}\right)^2 a^2 + 2ia\left(x-\frac{i}{a}\right)}}{\sqrt{a^2}}\right)}{\sqrt{a^2}} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)

[Out] $-1/a^2*(I/a/(x-I/a)^2*((x-I/a)^{2*a^2+2*I*a*(x-I/a)})^{(3/2)}-I*a*((x-I/a)^{2*a^2+2*I*a*(x-I/a)})^{(1/2)}+I*a*\ln((I*a+(x-I/a)*a^2)/(a^2)^{(1/2)}+((x-I/a)^{2*a^2+2*I*a*(x-I/a)})^{(1/2)}))/(a^2)^{(1/2))}$

Maxima [A]

time = 0.47, size = 33, normalized size = 0.80

$$-\frac{\operatorname{arsinh}(ax)}{a} + \frac{2i\sqrt{a^2x^2+1}}{ia^2x+a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)^(1/2),x, algorithm="maxima")`

[Out] $-\operatorname{arcsinh}(a*x)/a + 2*I*\sqrt{a^2*x^2 + 1}/(I*a^2*x + a)$

Fricas [A]

time = 3.30, size = 54, normalized size = 1.32

$$\frac{2ax + (ax - i)\log(-ax + \sqrt{a^2x^2 + 1}) + 2\sqrt{a^2x^2 + 1} - 2i}{a^2x - ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)^(1/2),x, algorithm="fricas")`

[Out] $(2*a*x + (a*x - I)*\log(-a*x + \sqrt{a^2*x^2 + 1}) + 2*\sqrt{a^2*x^2 + 1} - 2*I)/(a^2*x - I*a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{\sqrt{a^2x^2+1}}{a^2x^2-2iax-1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)**2*(a**2*x**2+1)**(1/2),x)`

[Out] $-\operatorname{Integral}(\sqrt{a**2*x**2 + 1}/(a**2*x**2 - 2*I*a*x - 1), x)$

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)^(1/2),x, algorithm="giac")`

[Out] undef

Mupad [B]

time = 0.49, size = 56, normalized size = 1.37

$$-\frac{\operatorname{asinh}\left(x\sqrt{a^2}\right)}{\sqrt{a^2}} - \frac{2\sqrt{a^2x^2+1}}{\left(-x\sqrt{a^2} + \frac{\sqrt{a^2}i}{a}\right)\sqrt{a^2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(1/2)/(a*x*i + 1)^2,x)

[Out] - asinh(x*(a^2)^(1/2))/(a^2)^(1/2) - (2*(a^2*x^2 + 1)^(1/2))/(((a^2)^(1/2)*i)/a - x*(a^2)^(1/2))*(a^2)^(1/2))

$$3.308 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=32

$$-\frac{2}{a(i-ax)} + \frac{i \log(i-ax)}{a}$$

[Out] -2/a/(I-a*x)+I*ln(I-a*x)/a

Rubi [A]

time = 0.03, antiderivative size = 32, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 45}

$$\frac{i \log(-ax+i)}{a} - \frac{2}{a(-ax+i)}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a*x])*Sqrt[1+a^2*x^2]),x]

[Out] -2/(a*(I-a*x))+(I*Log[I-a*x])/a

Rule 45

Int[((a_.)+(b_.)*(x_))^(m_.)*((c_.)+(d_.)*(x_))^(n_.),x_Symbol]>Int[ExpandIntegrand[(a+b*x)^m*(c+d*x)^n,x],x]/;FreeQ[{a,b,c,d,n},x]&&NeQ[b*c-a*d,0]&&IGtQ[m,0]&&(IntegerQ[n]||EqQ[c,0]&&LeQ[7*m+4*n+4,0])||LtQ[9*m+5*(n+1),0]||GtQ[m+n+2,0]

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.)+(d_.)*(x_)^2)^(p_.),x_Symbol]>Dist[c^p,Int[(1-I*a*x)^(p+I*(n/2))*(1+I*a*x)^(p-I*(n/2)),x],x]/;FreeQ[{a,c,d,n,p},x]&&EqQ[d,a^2*c]&&(IntegerQ[p]||GtQ[c,0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-3i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{1-iax}{(1+iax)^2} dx \\ &= \int \left(-\frac{2}{(-i+ax)^2} + \frac{i}{-i+ax} \right) dx \\ &= -\frac{2}{a(i-ax)} + \frac{i \log(i-ax)}{a} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 32, normalized size = 1.00

$$-\frac{2}{a(i-ax)} + \frac{i \log(i-ax)}{a}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x])*Sqrt[1 + a^2*x^2]),x]

[Out] -2/(a*(I - a*x)) + (I*Log[I - a*x])/a

Maple [A]

time = 0.06, size = 30, normalized size = 0.94

method	result	size
default	$-\frac{2}{a(-ax+i)} + \frac{i \ln(-ax+i)}{a}$	30
risch	$\frac{2}{a(ax-i)} + \frac{i \ln(a^2x^2+1)}{2a} - \frac{\arctan(ax)}{a}$	40
meijerg	$\frac{i \left(-\frac{ix a(9iax+6)}{3(iax+1)^2} + 2 \ln(iax+1) \right)}{2a} + \frac{x(iax+2)}{2(iax+1)^2}$	59

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^3*(a^2*x^2+1),x,method=_RETURNVERBOSE)

[Out] -2/a/(I-a*x)+I*ln(I-a*x)/a

Maxima [A]

time = 0.25, size = 41, normalized size = 1.28

$$-\frac{4(-iax-1)}{2ia^3x^2+4a^2x-2ia} + \frac{i \log(iax+1)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1),x, algorithm="maxima")

[Out] -4*(-I*a*x - 1)/(2*I*a^3*x^2 + 4*a^2*x - 2*I*a) + I*log(I*a*x + 1)/a

Fricas [A]

time = 3.00, size = 31, normalized size = 0.97

$$\frac{(iax+1) \log\left(\frac{ax-i}{a}\right) + 2}{a^2x - ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1),x, algorithm="fricas")

[Out] $((I*a*x + 1)*\log((a*x - I)/a) + 2)/(a^2*x - I*a)$

Sympy [A]

time = 0.08, size = 19, normalized size = 0.59

$$\frac{2}{a^2x - ia} + \frac{i \log(ax - i)}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)**3*(a**2*x**2+1),x)`

[Out] $2/(a**2*x - I*a) + I*\log(a*x - I)/a$

Giac [A]

time = 0.43, size = 24, normalized size = 0.75

$$\frac{i \log(ax - i)}{a} + \frac{2}{(ax - i)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^3*(a^2*x^2+1),x, algorithm="giac")`

[Out] $I*\log(a*x - I)/a + 2/((a*x - I)*a)$

Mupad [B]

time = 0.49, size = 29, normalized size = 0.91

$$-\frac{2}{-a^2x + a \operatorname{li}} + \frac{\ln(ax - i) \operatorname{li}}{a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a^2*x^2 + 1)/(a*x*1i + 1)^3,x)`

[Out] $(\log(a*x - 1i)*1i)/a - 2/(a*1i - a^2*x)$

$$3.309 \quad \int \frac{e^{-4i \operatorname{ArcTan}(ax)}}{\sqrt{1+a^2x^2}} dx$$

Optimal. Leaf size=73

$$\frac{2i(1-iax)^{3/2}}{3a(1+iax)^{3/2}} - \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} + \frac{\sinh^{-1}(ax)}{a}$$

[Out] $2/3*I*(1-I*a*x)^{(3/2)}/a/(1+I*a*x)^{(3/2)} + \operatorname{arcsinh}(a*x)/a - 2*I*(1-I*a*x)^{(1/2)}/a/(1+I*a*x)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 73, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5181, 49, 41, 221}

$$\frac{2i(1-iax)^{3/2}}{3a(1+iax)^{3/2}} - \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} + \frac{\sinh^{-1}(ax)}{a}$$

Antiderivative was successfully verified.

[In] `Int[1/(E^((4*I)*ArcTan[a*x])*Sqrt[1 + a^2*x^2]),x]`

[Out] `((2*I)/3)*(1 - I*a*x)^(3/2)/(a*(1 + I*a*x)^(3/2)) - ((2*I)*Sqrt[1 - I*a*x])/ (a*Sqrt[1 + I*a*x]) + ArcSinh[a*x]/a`

Rule 41

`Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[(a*c + b*d*x^2)^m, x] /; FreeQ[{a, b, c, d, m}, x] && EqQ[b*c + a*d, 0] && (IntegerQ[m] || (GtQ[a, 0] && GtQ[c, 0]))`

Rule 49

`Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Simp[(a + b*x)^(m + 1)*((c + d*x)^n/(b*(m + 1))), x] - Dist[d*(n/(b*(m + 1))), Int[(a + b*x)^(m + 1)*(c + d*x)^(n - 1), x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && GtQ[n, 0] && LtQ[m, -1] && !(IntegerQ[n] && !IntegerQ[m]) && !(IntegerQ[m + n + 2, 0] && (FractionQ[m] || GeQ[2*n + m + 1, 0])) && IntLinearQ[a, b, c, d, m, n, x]`

Rule 221

`Int[1/Sqrt[(a_) + (b_)*(x_)^2], x_Symbol] := Simp[ArcSinh[Rt[b, 2]*(x/Sqrt[a])]/Rt[b, 2], x] /; FreeQ[{a, b}, x] && GtQ[a, 0] && PosQ[b]`

Rule 5181

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :=
Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /
; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-4i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx &= \int \frac{(1-iax)^{3/2}}{(1+iax)^{5/2}} dx \\
&= \frac{2i(1-iax)^{3/2}}{3a(1+iax)^{3/2}} - \int \frac{\sqrt{1-iax}}{(1+iax)^{3/2}} dx \\
&= \frac{2i(1-iax)^{3/2}}{3a(1+iax)^{3/2}} - \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} + \int \frac{1}{\sqrt{1-iax}\sqrt{1+iax}} dx \\
&= \frac{2i(1-iax)^{3/2}}{3a(1+iax)^{3/2}} - \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} + \int \frac{1}{\sqrt{1+a^2x^2}} dx \\
&= \frac{2i(1-iax)^{3/2}}{3a(1+iax)^{3/2}} - \frac{2i\sqrt{1-iax}}{a\sqrt{1+iax}} + \frac{\sinh^{-1}(ax)}{a}
\end{aligned}$$

Mathematica [A]

time = 0.05, size = 82, normalized size = 1.12

$$\frac{2i \left(\frac{2\sqrt{1+iax} (1+iax+2a^2x^2)}{\sqrt{1-iax} (-i+ax)^2} + 3\text{ArcSin} \left(\frac{\sqrt{1-iax}}{\sqrt{2}} \right) \right)}{3a}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((4*I)*ArcTan[a*x])*Sqrt[1 + a^2*x^2]), x]

[Out] (((2*I)/3)*((2*Sqrt[1 + I*a*x]*(1 + I*a*x + 2*a^2*x^2))/(Sqrt[1 - I*a*x]*(-I + a*x)^2) + 3*ArcSin[Sqrt[1 - I*a*x]/Sqrt[2]]))/a

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 304 vs. 2(57) = 114.

time = 0.09, size = 305, normalized size = 4.18

method	result
default	$ \frac{i \left(\left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{5}{2}} \right)}{3a \left(x - \frac{i}{a} \right)^4} - 2ia \left(-\frac{i \left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{5}{2}}}{a \left(x - \frac{i}{a} \right)^2} + 3ia \frac{\left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{3}{2}}}{3} + i \right) $

a^4

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)^4*(a^2*x^2+1)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{a^4} \left(\frac{1}{3} \frac{I/a}{(x-I/a)^4} \left(\frac{(x-I/a)^2 a^2 + 2 I a (x-I/a)}{(x-I/a)^2 a^2 + 2 I a (x-I/a)} \right)^{5/2} - \frac{1}{3} I a \left(\frac{I/a}{(x-I/a)^3} \left(\frac{(x-I/a)^2 a^2 + 2 I a (x-I/a)}{(x-I/a)^2 a^2 + 2 I a (x-I/a)} \right)^{5/2} - 2 I a \left(\frac{-I/a}{(x-I/a)^2} \left(\frac{(x-I/a)^2 a^2 + 2 I a (x-I/a)}{(x-I/a)^2 a^2 + 2 I a (x-I/a)} \right)^{5/2} + 3 I a \left(\frac{1}{3} \left(\frac{(x-I/a)^2 a^2 + 2 I a (x-I/a)}{(x-I/a)^2 a^2 + 2 I a (x-I/a)} \right)^{3/2} + I a \left(\frac{1}{4} \left(2 \left(\frac{(x-I/a)^2 a^2 + 2 I a (x-I/a)}{(x-I/a)^2 a^2 + 2 I a (x-I/a)} \right)^{1/2} + \frac{1}{2} \ln \left(\frac{I a + (x-I/a) a^2}{(a^2)^{1/2} + \left(\frac{(x-I/a)^2 a^2 + 2 I a (x-I/a)}{(x-I/a)^2 a^2 + 2 I a (x-I/a)} \right)^{1/2} \right) \right) \right) \right)$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 107 vs. $2(51) = 102$.

time = 0.47, size = 107, normalized size = 1.47

$$\frac{i(a^2x^2 + 1)^{\frac{3}{2}}}{-3i a^4 x^3 - 9 a^3 x^2 + 9i a^2 x + 3a} + \frac{\operatorname{arsinh}(ax)}{a} - \frac{2i \sqrt{a^2x^2 + 1}}{3(a^3x^2 - 2i a^2x - a)} - \frac{7i \sqrt{a^2x^2 + 1}}{3i a^2x + 3a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^(3/2),x, algorithm="maxima")`

[Out] $I*(a^2*x^2 + 1)^{3/2}/(-3*I*a^4*x^3 - 9*a^3*x^2 + 9*I*a^2*x + 3*a) + \operatorname{arcsinh}(a*x)/a - 2/3*I*\sqrt{a^2*x^2 + 1}/(a^3*x^2 - 2*I*a^2*x - a) - 7*I*\sqrt{a^2*x^2 + 1}/(3*I*a^2*x + 3*a)$

Fricas [A]

time = 5.12, size = 86, normalized size = 1.18

$$\frac{8a^2x^2 - 16i ax + 3(a^2x^2 - 2i ax - 1) \log(-ax + \sqrt{a^2x^2 + 1}) + 4\sqrt{a^2x^2 + 1}(2ax - i) - 8}{3(a^3x^2 - 2i a^2x - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^(3/2),x, algorithm="fricas")`

[Out] $-1/3*(8*a^2*x^2 - 16*I*a*x + 3*(a^2*x^2 - 2*I*a*x - 1)*\log(-a*x + \sqrt{a^2*x^2 + 1})) + 4*\sqrt{a^2*x^2 + 1}*(2*a*x - I) - 8)/(a^3*x^2 - 2*I*a^2*x - a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a^2x^2 + 1)^{\frac{3}{2}}}{(ax - i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**4*(a**2*x**2+1)**(3/2),x)

[Out] Integral((a**2*x**2 + 1)**(3/2)/(a*x - I)**4, x)

Giac [A]

time = 0.46, size = 24, normalized size = 0.33

$$-\frac{\log\left(-x|a| + \sqrt{a^2x^2 + 1}\right)}{|a|}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^(3/2),x, algorithm="giac")

[Out] -log(-x*abs(a) + sqrt(a^2*x^2 + 1))/abs(a)

Mupad [B]

time = 0.10, size = 93, normalized size = 1.27

$$\frac{\operatorname{asinh}\left(x\sqrt{a^2}\right)}{\sqrt{a^2}} + \frac{8\sqrt{a^2x^2+1}}{3\left(-x\sqrt{a^2} + \frac{\sqrt{a^2}}{a}\right)\sqrt{a^2}} + \frac{a\sqrt{a^2x^2+1}4i}{3\left(-a^4x^2 + a^3x2i + a^2\right)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/(a*x*1i + 1)^4,x)

[Out] asinh(x*(a^2)^(1/2))/(a^2)^(1/2) + (8*(a^2*x^2 + 1)^(1/2))/(3*(((a^2)^(1/2)*1i)/a - x*(a^2)^(1/2))*(a^2)^(1/2)) + (a*(a^2*x^2 + 1)^(1/2)*4i)/(3*(a^3*x*2i + a^2 - a^4*x^2))

$$3.310 \quad \int \frac{e^{5i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=131

$$-\frac{2i\sqrt{1+a^2x^2}}{a(1-iax)^2\sqrt{c+a^2cx^2}} + \frac{4i\sqrt{1+a^2x^2}}{a(1-iax)\sqrt{c+a^2cx^2}} + \frac{i\sqrt{1+a^2x^2} \log(i+ax)}{a\sqrt{c+a^2cx^2}}$$

[Out] $-2*I*(a^2*x^2+1)^{(1/2)}/a/(1-I*a*x)^2/(a^2*c*x^2+c)^{(1/2)}+4*I*(a^2*x^2+1)^{(1/2)}/a/(1-I*a*x)/(a^2*c*x^2+c)^{(1/2)}+I*\ln(I+a*x)*(a^2*x^2+1)^{(1/2)}/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 131, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 45}

$$\frac{4i\sqrt{a^2x^2+1}}{a(1-iax)\sqrt{a^2cx^2+c}} - \frac{2i\sqrt{a^2x^2+1}}{a(1-iax)^2\sqrt{a^2cx^2+c}} + \frac{i\sqrt{a^2x^2+1} \log(ax+i)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(5*I)*\text{ArcTan}[a*x]}/\text{Sqrt}[c + a^2*c*x^2], x]$

[Out] $((-2*I)*\text{Sqrt}[1 + a^2*x^2])/(a*(1 - I*a*x)^2*\text{Sqrt}[c + a^2*c*x^2]) + ((4*I)*\text{Sqrt}[1 + a^2*x^2])/(a*(1 - I*a*x)*\text{Sqrt}[c + a^2*c*x^2]) + (I*\text{Sqrt}[1 + a^2*x^2]*\text{Log}[I + a*x])/(a*\text{Sqrt}[c + a^2*c*x^2])$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LtQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p*\text{IntPart}[p]*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
 \int \frac{e^{5i \tan^{-1}(ax)}}{\sqrt{c+a^2cx^2}} dx &= \frac{\sqrt{1+a^2x^2} \int \frac{e^{5i \tan^{-1}(ax)}}{\sqrt{1+a^2x^2}} dx}{\sqrt{c+a^2cx^2}} \\
 &= \frac{\sqrt{1+a^2x^2} \int \frac{(1+iax)^2}{(1-iax)^3} dx}{\sqrt{c+a^2cx^2}} \\
 &= \frac{\sqrt{1+a^2x^2} \int \left(\frac{4}{(1-iax)^3} - \frac{4}{(1-iax)^2} + \frac{1}{1-iax} \right) dx}{\sqrt{c+a^2cx^2}} \\
 &= -\frac{2i\sqrt{1+a^2x^2}}{a(1-iax)^2\sqrt{c+a^2cx^2}} + \frac{4i\sqrt{1+a^2x^2}}{a(1-iax)\sqrt{c+a^2cx^2}} + \frac{i\sqrt{1+a^2x^2} \log(i+ax)}{a\sqrt{c+a^2cx^2}}
 \end{aligned}$$

Mathematica [A]

time = 0.03, size = 69, normalized size = 0.53

$$\frac{i\sqrt{1+a^2x^2}(-2+4iax+(i+ax)^2 \log(i+ax))}{a(i+ax)^2\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((5*I)*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]

[Out] (I*Sqrt[1 + a^2*x^2]*(-2 + (4*I)*a*x + (I + a*x)^2*Log[I + a*x]))/(a*(I + a*x)^2*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.10, size = 84, normalized size = 0.64

method	result	size
risch	$\frac{\sqrt{a^2x^2+1}(-4x-\frac{2i}{a})}{\sqrt{c(a^2x^2+1)}(ax+i)^2} + \frac{i\sqrt{a^2x^2+1} \ln(ax+i)}{\sqrt{c(a^2x^2+1)}a}$	82
default	$\frac{\sqrt{c(a^2x^2+1)}(i \ln(ax+i)a^2x^2 - 2 \ln(ax+i)ax - i \ln(ax+i) - 4ax - 2i)}{\sqrt{a^2x^2+1}ca(ax+i)^2}$	84

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(1/2), x, method=_RETURNVERBOSE)

[Out] 1/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)*(I*ln(I+a*x)*a^2*x^2-2*ln(I+a*x)*a*x-I*ln(I+a*x)-4*a*x-2*I)/c/a/(I+a*x)^2

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")
```

```
[Out] integrate((I*a*x + 1)^5/(sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 1)^(5/2)), x)
```

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 364 vs. $2(107) = 214$.

time = 4.25, size = 364, normalized size = 2.78

$$\frac{-4\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}ax^2+(ia^4cx^4-2a^3cx^3-2acx-ic)\sqrt{\frac{1}{a^2c}}\log\left(\frac{(ia^4x^4-2a^3x^3-2a^2x^2+ic)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(ia^4cx^4-2a^3cx^3+ia^2cx^2-2a^2cx)\sqrt{\frac{1}{a^2c}}}{8(a^2x^2+1)(a^2cx^2+c)}\right)+(-ia^4cx^4+2a^3cx^3+2acx+ic)\sqrt{\frac{1}{a^2c}}\log\left(\frac{(ia^4x^4-2a^3x^3-2a^2x^2+ic)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(-ia^4cx^4+2a^3cx^3-ia^2cx^2+2a^2cx)\sqrt{\frac{1}{a^2c}}}{8(a^2x^2+1)(a^2cx^2+c)}\right)}{2(a^4cx^4+2ia^3cx^3+2iacx-c)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/2*(-4*I*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)*a*x^2 + (I*a^4*c*x^4 - 2*a^3*c*x^3 - 2*a*c*x - I*c)*sqrt(1/(a^2*c))*log(1/8*((I*a^6*x^2 - 2*a^5*x - 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (I*a^9*c*x^4 - 2*a^8*c*x^3 + I*a^7*c*x^2 - 2*a^6*c*x)*sqrt(1/(a^2*c))))/(a^3*x^3 + I*a^2*x^2 + a*x + I)) + (-I*a^4*c*x^4 + 2*a^3*c*x^3 + 2*a*c*x + I*c)*sqrt(1/(a^2*c))*log(1/8*((I*a^6*x^2 - 2*a^5*x - 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (-I*a^9*c*x^4 + 2*a^8*c*x^3 - I*a^7*c*x^2 + 2*a^6*c*x)*sqrt(1/(a^2*c))))/(a^3*x^3 + I*a^2*x^2 + a*x + I)))/(a^4*c*x^4 + 2*I*a^3*c*x^3 + 2*I*a*c*x - c)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{((\frac{1}{2}(-4I\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}ax^2+(ia^4cx^4-2a^3cx^3-2acx-ic)\sqrt{\frac{1}{a^2c}}\log(\frac{(ia^4x^4-2a^3x^3-2a^2x^2+ic)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(ia^4cx^4-2a^3cx^3+ia^2cx^2-2a^2cx)\sqrt{\frac{1}{a^2c}}}{8(a^2x^2+1)(a^2cx^2+c)})))+(-ia^4cx^4+2a^3cx^3+2acx+ic)\sqrt{\frac{1}{a^2c}}\log(\frac{(ia^4x^4-2a^3x^3-2a^2x^2+ic)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(-ia^4cx^4+2a^3cx^3-ia^2cx^2+2a^2cx)\sqrt{\frac{1}{a^2c}}}{8(a^2x^2+1)(a^2cx^2+c)}))}{2(a^4cx^4+2ia^3cx^3+2iacx-c)}}{2(a^4cx^4+2ia^3cx^3+2iacx-c)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)**5/(a**2*x**2+1)**(5/2)/(a**2*c*x**2+c)**(1/2),x)
```

```
[Out] I*(Integral(-I/(a**4*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(5*a*x/(a**4*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-10*a**3*x**3/(a**4*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c))
```

), x) + Integral(a**5*x**5/(a**4*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(10*I*a**2*x**2/(a**4*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-5*I*a**4*x**4/(a**4*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate((I*a*x + 1)^5/(sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 1)^(5/2)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(1 + a x i)^5}{\sqrt{c a^2 x^2 + c} (a^2 x^2 + 1)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^5/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^(5/2)),x)

[Out] int((a*x*1i + 1)^5/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^(5/2)), x)

$$3.311 \quad \int \frac{e^{4i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=96

$$-\frac{2ic(1+iax)^3}{3a(c+a^2cx^2)^{3/2}} + \frac{2i(1+iax)}{a\sqrt{c+a^2cx^2}} + \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{c+a^2cx^2}}\right)}{a\sqrt{c}}$$

[Out] $-2/3*I*c*(1+I*a*x)^3/a/(a^2*c*x^2+c)^{(3/2)}+\operatorname{arctanh}(a*x*c^{(1/2)}/(a^2*c*x^2+c)^{(1/2)})/a/c^{(1/2)}+2*I*(1+I*a*x)/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5183, 683, 667, 223, 212}

$$-\frac{2ic(1+iax)^3}{3a(a^2cx^2+c)^{3/2}} + \frac{2i(1+iax)}{a\sqrt{a^2cx^2+c}} + \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{a^2cx^2+c}}\right)}{a\sqrt{c}}$$

Antiderivative was successfully verified.

[In] `Int[E^((4*I)*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]`

[Out] `(((-2*I)/3)*c*(1 + I*a*x)^3)/(a*(c + a^2*c*x^2)^(3/2)) + ((2*I)*(1 + I*a*x))/(a*Sqrt[c + a^2*c*x^2]) + ArcTanh[(a*Sqrt[c]*x)/Sqrt[c + a^2*c*x^2]]/(a*Sqrt[c])`

Rule 212

`Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[-b, 2]))*ArcTanh[Rt[-b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && NegQ[a/b] && (GtQ[a, 0] || LtQ[b, 0])`

Rule 223

`Int[1/Sqrt[(a_) + (b_.)*(x_)^2], x_Symbol] := Subst[Int[1/(1 - b*x^2), x], x, x/Sqrt[a + b*x^2]] /; FreeQ[{a, b}, x] && !GtQ[a, 0]`

Rule 667

`Int[((d_) + (e_.)*(x_))^(2*((a_) + (c_.)*(x_)^2)^(p_)), x_Symbol] := Simp[e*(d + e*x)*((a + c*x^2)^(p + 1)/(c*(p + 1))), x] - Dist[e^2*((p + 2)/(c*(p + 1))), Int[(a + c*x^2)^(p + 1), x], x] /; FreeQ[{a, c, d, e, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && LtQ[p, -1]`

Rule 683

```
Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
e*(d + e*x)^(m - 1)*((a + c*x^2)^(p + 1)/(c*(p + 1))), x] - Dist[e^2*((m +
p)/(c*(p + 1))), Int[(d + e*x)^(m - 2)*(a + c*x^2)^(p + 1), x], x] /; FreeQ
[{a, c, d, e}, x] && EqQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && GtQ[m, 1] && In
tegerQ[2*p]
```

Rule 5183

```
Int[E^(ArcTan[(a_.)*(x_)])*(n_)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Di
st[1/c^(I*(n/2)), Int[(c + d*x^2)^(p + I*(n/2))/(1 + I*a*x)^(I*n), x], x] /
; FreeQ[{a, c, d, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0]) &
& ILtQ[I*(n/2), 0]
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{4i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= c^2 \int \frac{(1 + iax)^4}{(c + a^2 cx^2)^{5/2}} dx \\
&= -\frac{2ic(1 + iax)^3}{3a(c + a^2 cx^2)^{3/2}} - c \int \frac{(1 + iax)^2}{(c + a^2 cx^2)^{3/2}} dx \\
&= -\frac{2ic(1 + iax)^3}{3a(c + a^2 cx^2)^{3/2}} + \frac{2i(1 + iax)}{a\sqrt{c + a^2 cx^2}} + \int \frac{1}{\sqrt{c + a^2 cx^2}} dx \\
&= -\frac{2ic(1 + iax)^3}{3a(c + a^2 cx^2)^{3/2}} + \frac{2i(1 + iax)}{a\sqrt{c + a^2 cx^2}} + \text{Subst}\left(\int \frac{1}{1 - a^2 cx^2} dx, x, \frac{x}{\sqrt{c + a^2 cx^2}}\right) \\
&= -\frac{2ic(1 + iax)^3}{3a(c + a^2 cx^2)^{3/2}} + \frac{2i(1 + iax)}{a\sqrt{c + a^2 cx^2}} + \frac{\tanh^{-1}\left(\frac{a\sqrt{c} x}{\sqrt{c + a^2 cx^2}}\right)}{a\sqrt{c}}
\end{aligned}$$

Mathematica [C] Result contains higher order function than in optimal. Order 5 vs. order 3 in optimal.

time = 0.02, size = 71, normalized size = 0.74

$$-\frac{4i\sqrt{2 + 2a^2x^2} {}_2F_1\left(-\frac{3}{2}, -\frac{3}{2}; -\frac{1}{2}; \frac{1}{2}(1 - iax)\right)}{3a(1 - iax)^{3/2}\sqrt{c + a^2cx^2}}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[E^((4*I)*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]
```

```
[Out] (((-4*I)/3)*Sqrt[2 + 2*a^2*x^2]*Hypergeometric2F1[-3/2, -3/2, -1/2, (1 - I*
a*x)/2])/(a*(1 - I*a*x)^(3/2)*Sqrt[c + a^2*c*x^2])
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 525 vs. 2(80) = 160.
time = 0.14, size = 526, normalized size = 5.48

method	result
default	$\frac{\ln\left(\frac{a^2cx}{\sqrt{a^2c}} + \sqrt{a^2cx^2 + c}\right)}{\sqrt{a^2c}} + \frac{2\left(i\sqrt{-a^2} - a\right)\sqrt{\left(x + \frac{\sqrt{-a^2}}{a^2}\right)^2 a^2c - 2c\sqrt{-a^2}\left(x + \frac{\sqrt{-a^2}}{a^2}\right)}}{a^3c\left(x + \frac{\sqrt{-a^2}}{a^2}\right)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)
[Out] ln(a^2*c*x/(a^2*c)^(1/2)+(a^2*c*x^2+c)^(1/2))/(a^2*c)^(1/2)+2/a^3*(I*(-a^2)^(1/2)-a)/c/(x+(-a^2)^(1/2)/a^2)*((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)-2/a^3*(I*(-a^2)^(1/2)+a)*(-1/3/c/(-a^2)^(1/2)/(x-(-a^2)^(1/2)/a^2)^2*((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2)-1/3/c/(x-(-a^2)^(1/2)/a^2)*((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2))-2/a^3*(I*(-a^2)^(1/2)+a)/c/(x-(-a^2)^(1/2)/a^2)*((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2)+2/a^3*(I*(-a^2)^(1/2)-a)*(1/3/c/(-a^2)^(1/2)/(x+(-a^2)^(1/2)/a^2)^2*((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)-1/3/c/(x+(-a^2)^(1/2)/a^2)*((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")
[Out] integrate((I*a*x + 1)^4/(sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 1)^2), x)
```

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 186 vs. 2(75) = 150.
time = 2.81, size = 186, normalized size = 1.94

$$\frac{3(a^3cx^2 + 2ia^2cx - ac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(a^2cx + \sqrt{a^2cx^2 + c}\right)\sqrt{\frac{1}{a^2c}}}{x}\right) - 3(a^3cx^2 + 2ia^2cx - ac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(a^2cx - \sqrt{a^2cx^2 + c}\right)\sqrt{\frac{1}{a^2c}}}{x}\right) - 8\sqrt{a^2cx^2 + c}(2ax + i)}{6(a^3cx^2 + 2ia^2cx - ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] 1/6*(3*(a^3*c*x^2 + 2*I*a^2*c*x - a*c)*sqrt(1/(a^2*c))*log(2*(a^2*c*x + sqrt(a^2*c*x^2 + c)*a^2*c*sqrt(1/(a^2*c))))/x - 3*(a^3*c*x^2 + 2*I*a^2*c*x - a*c)*sqrt(1/(a^2*c))*log(2*(a^2*c*x - sqrt(a^2*c*x^2 + c)*a^2*c*sqrt(1/(a^2*c))))/x - 8*sqrt(a^2*c*x^2 + c)*(2*a*x + I)/(a^3*c*x^2 + 2*I*a^2*c*x - a*c)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(ax - i)^4}{\sqrt{c(a^2x^2 + 1)} (a^2x^2 + 1)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral((a*x - I)**4/(sqrt(c*(a**2*x**2 + 1))*(a**2*x**2 + 1)**2), x)

Giac [A]

time = 0.51, size = 132, normalized size = 1.38

$$\frac{\log\left(\left|-\sqrt{a^2c}x + \sqrt{a^2cx^2 + c}\right|\right)}{a\sqrt{c}} - \frac{8\left(3\left(\sqrt{a^2c}x - \sqrt{a^2cx^2 + c}\right)^2 + 3i\left(\sqrt{a^2c}x - \sqrt{a^2cx^2 + c}\right)\sqrt{c} - 2c\right)}{3\left(i\sqrt{a^2c}x - i\sqrt{a^2cx^2 + c} - \sqrt{c}\right)^3 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] -log(abs(-sqrt(a^2*c)*x + sqrt(a^2*c*x^2 + c)))/(a*sqrt(c)) - 8/3*(3*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c))^2 + 3*I*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c)))*sqrt(c) - 2*c)/((I*sqrt(a^2*c)*x - I*sqrt(a^2*c*x^2 + c) - sqrt(c))^3*a)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(1 + ax \operatorname{li})^4}{\sqrt{ca^2x^2 + c} (a^2x^2 + 1)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*li + 1)^4/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^2),x)

[Out] int((a*x*li + 1)^4/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^2), x)

$$3.312 \quad \int \frac{e^{3i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=84

$$\frac{2\sqrt{1+a^2x^2}}{a(i+ax)\sqrt{c+a^2cx^2}} - \frac{i\sqrt{1+a^2x^2} \log(i+ax)}{a\sqrt{c+a^2cx^2}}$$

[Out] $2*(a^2*x^2+1)^{(1/2)}/a/(I+a*x)/(a^2*c*x^2+c)^{(1/2)}-I*\ln(I+a*x)*(a^2*x^2+1)^{(1/2)}/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 45}

$$\frac{2\sqrt{a^2x^2+1}}{a(ax+i)\sqrt{a^2cx^2+c}} - \frac{i\sqrt{a^2x^2+1} \log(ax+i)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[E^((3*I)*ArcTan[a*x])/Sqrt[c + a^2*c*x^2],x]

[Out] $(2*\text{Sqrt}[1 + a^2*x^2])/(a*(I + a*x)*\text{Sqrt}[c + a^2*c*x^2]) - (I*\text{Sqrt}[1 + a^2*x^2]*\text{Log}[I + a*x])/(a*\text{Sqrt}[c + a^2*c*x^2])$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{3i \tan^{-1}(ax)}}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \frac{1+iax}{(1-iax)^2} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \left(-\frac{2}{(i+ax)^2} - \frac{i}{i+ax} \right) dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{2\sqrt{1 + a^2 x^2}}{a(i + ax)\sqrt{c + a^2 cx^2}} - \frac{i\sqrt{1 + a^2 x^2} \log(i + ax)}{a\sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 55, normalized size = 0.65

$$\frac{\sqrt{1 + a^2 x^2} \left(\frac{2}{i+ax} - i \log(i + ax) \right)}{a\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x])/Sqrt[c + a^2*c*x^2],x]

[Out] (Sqrt[1 + a^2*x^2]*(2/(I + a*x) - I*Log[I + a*x]))/(a*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.09, size = 61, normalized size = 0.73

method	result	size
default	$\frac{(-i \ln(ax+i)ax + \ln(ax+i) + 2) \sqrt{c(a^2x^2 + 1)}}{\sqrt{a^2x^2 + 1} ca(ax+i)}$	61
risch	$\frac{2\sqrt{a^2x^2 + 1}}{\sqrt{c(a^2x^2 + 1)} a(ax+i)} - \frac{i\sqrt{a^2x^2 + 1} \ln(ax+i)}{\sqrt{c(a^2x^2 + 1)} a}$	76

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] (-I*ln(I+a*x)*a*x+ln(I+a*x)+2)/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)/c/a/(I+a*x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate((I*a*x + 1)^3/(sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 1)^(3/2)), x)

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 357 vs. 2(70) = 140.

time = 2.81, size = 357, normalized size = 4.25

$$\frac{(-i a^2 c x^3 + a^2 c x^2 - i a c x + c) \sqrt{\frac{1}{a^2 c}} \log\left(\frac{(i a^2 x^2 - 2 a^2 c x - 2 i a^2 c) \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1} + (i a^2 c x^2 - 2 a^2 c x - 2 i a^2 c) \sqrt{\frac{1}{a^2 c}}}{8(a^2 x^2 + 1)a^2 c x + 1}\right) + (i a^2 c x^3 - a^2 c x^2 + i a c x - c) \sqrt{\frac{1}{a^2 c}} \log\left(\frac{(i a^2 x^2 - 2 a^2 c x - 2 i a^2 c) \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1} + (-i a^2 c x^2 + 2 a^2 c x - i a^2 c) \sqrt{\frac{1}{a^2 c}}}{8(a^2 x^2 + 1)a^2 c x + 1}\right)}{2(a^2 c x^3 + i a^2 c x^2 + a c x + i c)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] 1/2*((-I*a^3*c*x^3 + a^2*c*x^2 - I*a*c*x + c)*sqrt(1/(a^2*c))*log(1/8*((I*a^6*x^2 - 2*a^5*x - 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (I*a^9*c*x^4 - 2*a^8*c*x^3 + I*a^7*c*x^2 - 2*a^6*c*x)*sqrt(1/(a^2*c)))/(a^3*x^3 + I*a^2*x^2 + a*x + I)) + (I*a^3*c*x^3 - a^2*c*x^2 + I*a*c*x - c)*sqrt(1/(a^2*c))*log(1/8*((I*a^6*x^2 - 2*a^5*x - 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (-I*a^9*c*x^4 + 2*a^8*c*x^3 - I*a^7*c*x^2 + 2*a^6*c*x)*sqrt(1/(a^2*c)))/(a^3*x^3 + I*a^2*x^2 + a*x + I)) + 4*I*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)*x/(a^3*c*x^3 + I*a^2*c*x^2 + a*c*x + I*c)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \left(\int \frac{i}{a^2 x^2 \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c} + \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c}} dx + \int \left(-\frac{3ax}{a^2 x^2 \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c} + \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c}} \right) dx + \int \frac{a^2 x^3}{a^2 x^2 \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c} + \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c}} dx + \int \left(-\frac{3a^2 x^2}{a^2 x^2 \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c} + \sqrt{a^2 x^2 + 1} \sqrt{a^2 c x^2 + c}} \right) dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)/(a**2*c*x**2+c)**(1/2),x)

[Out] -I*(Integral(I/(a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-3*a*x/(a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a**3*x**3/(a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-3*I*a**2*x**2/(a**2*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((I*a*x + 1)^3/(sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 1)^(3/2)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(1 + a x i)^3}{\sqrt{c a^2 x^2 + c} (a^2 x^2 + 1)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*x*1i + 1)^3/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^(3/2)),x)
```

```
[Out] int((a*x*1i + 1)^3/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^(3/2)), x)
```

$$3.313 \quad \int \frac{e^{2i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=63

$$-\frac{2i(1+iax)}{a\sqrt{c+a^2cx^2}} - \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{c+a^2cx^2}}\right)}{a\sqrt{c}}$$

[Out] $-\operatorname{arctanh}(a*x*c^{(1/2)}/(a^2*c*x^2+c)^{(1/2)})/a/c^{(1/2)}-2*I*(1+I*a*x)/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5183, 667, 223, 212}

$$-\frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{a^2cx^2+c}}\right)}{a\sqrt{c}} - \frac{2i(1+iax)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{((2*I)*\operatorname{ArcTan}[a*x])}/\operatorname{Sqrt}[c + a^2*c*x^2], x]$

[Out] $((-2*I)*(1 + I*a*x))/(a*\operatorname{Sqrt}[c + a^2*c*x^2]) - \operatorname{ArcTanh}[(a*\operatorname{Sqrt}[c]*x)/\operatorname{Sqrt}[c + a^2*c*x^2]]/(a*\operatorname{Sqrt}[c])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_)*(x_)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 667

$\operatorname{Int}[(d_ + (e_)*(x_)^2)^2*((a_ + (c_)*(x_)^2)^{p_}), x_Symbol] \rightarrow \operatorname{Simp}[e*(d + e*x)*((a + c*x^2)^{(p+1)}/(c*(p+1))), x] - \operatorname{Dist}[e^2*((p+2)/(c*(p+1))), \operatorname{Int}[(a + c*x^2)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, c, d, e, p\}, x \ \&\& \operatorname{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\operatorname{IntegerQ}[p] \ \&\& \operatorname{LtQ}[p, -1]$

Rule 5183

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Di
st[1/c^(I*(n/2)), Int[(c + d*x^2)^(p + I*(n/2))/(1 + I*a*x)^(I*n), x], x] /
; FreeQ[{a, c, d, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0]) &
& ILtQ[I*(n/2), 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= c \int \frac{(1 + iax)^2}{(c + a^2 cx^2)^{3/2}} dx \\ &= -\frac{2i(1 + iax)}{a\sqrt{c + a^2 cx^2}} - \int \frac{1}{\sqrt{c + a^2 cx^2}} dx \\ &= -\frac{2i(1 + iax)}{a\sqrt{c + a^2 cx^2}} - \text{Subst}\left(\int \frac{1}{1 - a^2 cx^2} dx, x, \frac{x}{\sqrt{c + a^2 cx^2}}\right) \\ &= -\frac{2i(1 + iax)}{a\sqrt{c + a^2 cx^2}} - \frac{\tanh^{-1}\left(\frac{a\sqrt{c} x}{\sqrt{c + a^2 cx^2}}\right)}{a\sqrt{c}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 91, normalized size = 1.44

$$-\frac{2i\sqrt{1 + a^2 x^2} \left(\sqrt{1 + iax} + \sqrt{1 - iax} \text{ArcSin}\left(\frac{\sqrt{1 - iax}}{\sqrt{2}}\right) \right)}{a\sqrt{1 - iax} \sqrt{c + a^2 cx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((2*I)*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]

[Out] ((-2*I)*Sqrt[1 + a^2*x^2]*(Sqrt[1 + I*a*x] + Sqrt[1 - I*a*x]*ArcSin[Sqrt[1 - I*a*x]/Sqrt[2]]))/(a*Sqrt[1 - I*a*x]*Sqrt[c + a^2*c*x^2])

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 203 vs. 2(53) = 106.

time = 0.08, size = 204, normalized size = 3.24

method	result
default	$-\frac{\ln\left(\frac{a^2 cx}{\sqrt{a^2 c}} + \sqrt{a^2 c x^2 + c}\right)}{\sqrt{a^2 c}} - \frac{(i\sqrt{-a^2} - a) \sqrt{\left(x + \frac{\sqrt{-a^2}}{a^2}\right)^2 a^2 c - 2c\sqrt{-a^2} \left(x + \frac{\sqrt{-a^2}}{a^2}\right)}}{a^3 c \left(x + \frac{\sqrt{-a^2}}{a^2}\right)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $-\ln(a^2*c*x/(a^2*c)^(1/2)+(a^2*c*x^2+c)^(1/2))/(a^2*c)^(1/2)-1/a^3*(I*(-a^2)^(1/2)-a)/c/(x+(-a^2)^(1/2)/a^2)*((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)+1/a^3*(I*(-a^2)^(1/2)+a)/c/(x-(-a^2)^(1/2)/a^2)*((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2)$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate((I*a*x + 1)^2/(sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 1)), x)`

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 152 vs. $2(50) = 100$.

time = 2.74, size = 152, normalized size = 2.41

$$\frac{(a^2cx + iac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(a^2cx + \sqrt{a^2cx^2 + c}\right)a^2c\sqrt{\frac{1}{a^2c}}}{x}\right) - (a^2cx + iac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(a^2cx - \sqrt{a^2cx^2 + c}\right)a^2c\sqrt{\frac{1}{a^2c}}}{x}\right) - 4\sqrt{a^2cx^2 + c}}{2(a^2cx + iac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

[Out] $-1/2*((a^2*c*x + I*a*c)*\sqrt{1/(a^2*c)}*\log(2*(a^2*c*x + \sqrt{a^2*c*x^2 + c})*a^2*c*\sqrt{1/(a^2*c)})/x) - (a^2*c*x + I*a*c)*\sqrt{1/(a^2*c)}*\log(2*(a^2*c*x - \sqrt{a^2*c*x^2 + c})*a^2*c*\sqrt{1/(a^2*c)})/x) - 4*\sqrt{a^2*c*x^2 + c})/(a^2*c*x + I*a*c)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{a^2x^2}{a^2x^2\sqrt{a^2cx^2 + c} + \sqrt{a^2cx^2 + c}} dx - \int \left(-\frac{2iax}{a^2x^2\sqrt{a^2cx^2 + c} + \sqrt{a^2cx^2 + c}}\right) dx - \int \left(-\frac{1}{a^2x^2\sqrt{a^2cx^2 + c} + \sqrt{a^2cx^2 + c}}\right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)/(a**2*c*x**2+c)**(1/2),x)

[Out] -Integral(a**2*x**2/(a**2*x**2*sqrt(a**2*c*x**2 + c) + sqrt(a**2*c*x**2 + c)), x) - Integral(-2*I*a*x/(a**2*x**2*sqrt(a**2*c*x**2 + c) + sqrt(a**2*c*x**2 + c)), x) - Integral(-1/(a**2*x**2*sqrt(a**2*c*x**2 + c) + sqrt(a**2*c*x**2 + c)), x)

Giac [A]

time = 0.46, size = 70, normalized size = 1.11

$$\frac{\log\left(\left|-\sqrt{a^2c}x + \sqrt{a^2cx^2 + c}\right|\right)}{a\sqrt{c}} - \frac{4}{\left(i\sqrt{a^2c}x - i\sqrt{a^2cx^2 + c} - \sqrt{c}\right)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] log(abs(-sqrt(a^2*c)*x + sqrt(a^2*c*x^2 + c)))/(a*sqrt(c)) - 4/((I*sqrt(a^2*c)*x - I*sqrt(a^2*c*x^2 + c) - sqrt(c))*a)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{(1 + axi)^2}{\sqrt{ca^2x^2 + c} (a^2x^2 + 1)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)),x)

[Out] int((a*x*1i + 1)^2/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)), x)

$$3.314 \quad \int \frac{e^{i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=42

$$\frac{i\sqrt{1 + a^2 x^2} \log(i + ax)}{a\sqrt{c + a^2 cx^2}}$$

[Out] I*ln(I+a*x)*(a^2*x^2+1)^(1/2)/a/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.05, antiderivative size = 42, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 31}

$$\frac{i\sqrt{a^2 x^2 + 1} \log(ax + i)}{a\sqrt{a^2 cx^2 + c}}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])/Sqrt[c + a^2*c*x^2],x]

[Out] (I*Sqrt[1 + a^2*x^2]*Log[I + a*x])/(a*Sqrt[c + a^2*c*x^2])

Rule 31

Int[((a_) + (b_.)*(x_))^(n_), x_Symbol] := Simp[Log[RemoveContent[a + b*x, x]]/b, x] /; FreeQ[{a, b}, x]

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{i \tan^{-1}(ax)}}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{1 - iax} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{i\sqrt{1 + a^2 x^2} \log(i + ax)}{a\sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 42, normalized size = 1.00

$$\frac{i\sqrt{1 + a^2 x^2} \log(i + ax)}{a\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]``[Out] (I*Sqrt[1 + a^2*x^2]*Log[I + a*x])/(a*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.08, size = 53, normalized size = 1.26

method	result	size
risch	$\frac{i\sqrt{a^2 x^2 + 1} \ln(ax+i)}{\sqrt{c(a^2 x^2 + 1)} a}$	38
default	$\frac{\sqrt{c(a^2 x^2 + 1)} (i \ln(a^2 x^2 + 1) + 2 \arctan(ax))}{2\sqrt{a^2 x^2 + 1} ca}$	53

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2), x, method=_RETURNVERBOSE)``[Out] 1/2/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)*(I*ln(a^2*x^2+1)+2*arctan(a*x))/c/a`**Maxima [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 253 vs. 2(34) = 68.
time = 3.62, size = 253, normalized size = 6.02

$$\frac{1}{2}i\sqrt{\frac{1}{a^2c}}\log\left(\frac{(ia^6x^2-2a^5x-2ia^4)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(ia^6cx^4-2a^8cx^3+ia^7cx^2-2a^6cx)\sqrt{\frac{1}{a^2c}}}{8(a^3x^3+ia^2x^2+ax+i)}\right)-\frac{1}{2}i\sqrt{\frac{1}{a^2c}}\log\left(\frac{(ia^6x^2-2a^5x-2ia^4)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(-ia^6cx^4+2a^8cx^3-ia^7cx^2+2a^6cx)\sqrt{\frac{1}{a^2c}}}{8(a^3x^3+ia^2x^2+ax+i)}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] 1/2*I*sqrt(1/(a^2*c))*log(1/8*((I*a^6*x^2 - 2*a^5*x - 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (I*a^9*c*x^4 - 2*a^8*c*x^3 + I*a^7*c*x^2 - 2*a^6*c*x)*sqrt(1/(a^2*c)))/(a^3*x^3 + I*a^2*x^2 + a*x + I)) - 1/2*I*sqrt(1/(a^2*c))*log(1/8*((I*a^6*x^2 - 2*a^5*x - 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (-I*a^9*c*x^4 + 2*a^8*c*x^3 - I*a^7*c*x^2 + 2*a^6*c*x)*sqrt(1/(a^2*c)))/(a^3*x^3 + I*a^2*x^2 + a*x + I))

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i\left(\int\left(-\frac{i}{\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}}\right)dx+\int\frac{ax}{\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}}dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/(a**2*c*x**2+c)**(1/2),x)

[Out] I*(Integral(-I/(sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a*x/(sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] integrate((I*a*x + 1)/(sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)), x)

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{1 + a x \operatorname{li}}{\sqrt{c a^2 x^2 + c} \sqrt{a^2 x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^(1/2)),x)

[Out] int((a*x*1i + 1)/((c + a^2*c*x^2)^(1/2)*(a^2*x^2 + 1)^(1/2)), x)

$$3.315 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=43

$$-\frac{i\sqrt{1+a^2x^2} \log(i-ax)}{a\sqrt{c+a^2cx^2}}$$

[Out] $-I*\ln(I-a*x)*(a^2*x^2+1)^{(1/2)}/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 43, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 31}

$$-\frac{i\sqrt{a^2x^2+1} \log(-ax+i)}{a\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{(I*\text{ArcTan}[a*x])*\text{Sqrt}[c + a^2*c*x^2]}), x]$

[Out] $((-I)*\text{Sqrt}[1 + a^2*x^2]*\text{Log}[I - a*x])/(a*\text{Sqrt}[c + a^2*c*x^2])$

Rule 31

$\text{Int}[(a_ + (b_)*(x_))^{(-1)}, x_Symbol] \rightarrow \text{Simp}[\text{Log}[\text{RemoveContent}[a + b*x, x]]/b, x] /; \text{FreeQ}\{a, b\}, x]$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_)*(x_)]*(n_))}*((c_ + (d_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x] \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_)*(x_)]*(n_))}*((c_ + (d_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]} / (1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x] \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int \frac{e^{-i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-i \tan^{-1}(ax)}}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{1+iax} dx}{\sqrt{c + a^2 cx^2}} \\ &= -\frac{i\sqrt{1 + a^2 x^2} \log(i - ax)}{a\sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 43, normalized size = 1.00

$$-\frac{i\sqrt{1 + a^2 x^2} \log(i - ax)}{a\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^(I*ArcTan[a*x]))*Sqrt[c + a^2*c*x^2]),x]

[Out] ((-I)*Sqrt[1 + a^2*x^2]*Log[I - a*x])/(a*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.09, size = 42, normalized size = 0.98

method	result	size
risch	$-\frac{i\sqrt{a^2 x^2 + 1} \ln(-ax+i)}{\sqrt{c(a^2 x^2 + 1)} a}$	39
default	$-\frac{i\sqrt{c(a^2 x^2 + 1)} \ln(iax+1)}{\sqrt{a^2 x^2 + 1} ca}$	42

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] -I/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)/c*ln(1+I*a*x)/a

Maxima [A]

time = 0.28, size = 15, normalized size = 0.35

$$-\frac{i \log(i a x + 1)}{a\sqrt{c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] -I*log(I*a*x + 1)/(a*sqrt(c))

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 253 vs. $2(35) = 70$.
time = 2.80, size = 253, normalized size = 5.88

$$\frac{1}{2}i\sqrt{\frac{1}{a^2c}}\log\left(\frac{(-ia^6x^2-2a^5x+2ia^4)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(ia^9cx^4+2a^8cx^3+ia^7cx^2+2a^6cx)\sqrt{\frac{1}{a^2c}}}{8(a^3x^3-ia^2x^2+ax-i)}}\right)-\frac{1}{2}i\sqrt{\frac{1}{a^2c}}\log\left(\frac{(-ia^6x^2-2a^5x+2ia^4)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(-ia^9cx^4-2a^8cx^3-ia^7cx^2-2a^6cx)\sqrt{\frac{1}{a^2c}}}{8(a^3x^3-ia^2x^2+ax-i)}}\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] $\frac{1}{2}I\sqrt{\frac{1}{a^2c}}\log\left(\frac{1}{8}\left((-Ia^6x^2-2a^5x+2Ia^4)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(Ia^9cx^4+2a^8cx^3+Ia^7cx^2+2a^6cx)\sqrt{\frac{1}{a^2c}}\right)\right)/(a^3x^3-Ia^2x^2+ax-I)-\frac{1}{2}I\sqrt{\frac{1}{a^2c}}\log\left(\frac{1}{8}\left((-Ia^6x^2-2a^5x+2Ia^4)\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}+(-Ia^9cx^4-2a^8cx^3-Ia^7cx^2-2a^6cx)\sqrt{\frac{1}{a^2c}}\right)\right)/(a^3x^3-Ia^2x^2+ax-I)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2x^2+1}}{ax\sqrt{a^2cx^2+c}-i\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/(a**2*c*x**2+c)**(1/2),x)

[Out] -I*Integral(sqrt(a**2*x**2 + 1)/(a*x*sqrt(a**2*c*x**2 + c) - I*sqrt(a**2*c*x**2 + c)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx)::OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{\sqrt{a^2 x^2 + 1}}{\sqrt{c a^2 x^2 + c} (1 + a x i)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(1/2)/((c + a^2*c*x^2)^(1/2)*(a*x*i + 1)), x)

[Out] int((a^2*x^2 + 1)^(1/2)/((c + a^2*c*x^2)^(1/2)*(a*x*i + 1)), x)

$$3.316 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=63

$$\frac{2i(1 - iax)}{a\sqrt{c + a^2 cx^2}} - \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{c + a^2 cx^2}}\right)}{a\sqrt{c}}$$

[Out] $-\operatorname{arctanh}(a*x*c^{(1/2)}/(a^2*c*x^2+c)^{(1/2)})/a/c^{(1/2)}+2*I*(1-I*a*x)/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 63, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5182, 667, 223, 212}

$$-\frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{a^2 cx^2 + c}}\right)}{a\sqrt{c}} + \frac{2i(1 - iax)}{a\sqrt{a^2 cx^2 + c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(E^{((2*I)*\operatorname{ArcTan}[a*x])}*\operatorname{Sqrt}[c + a^2*c*x^2]), x]$

[Out] $((2*I)*(1 - I*a*x))/(a*\operatorname{Sqrt}[c + a^2*c*x^2]) - \operatorname{ArcTanh}[(a*\operatorname{Sqrt}[c]*x)/\operatorname{Sqrt}[c + a^2*c*x^2]]/(a*\operatorname{Sqrt}[c])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_)*(x_)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /; \operatorname{FreeQ}\{a, b\}, x \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 667

$\operatorname{Int}[(d_ + (e_)*(x_)^2)^2*((a_ + (c_)*(x_)^2)^{p_}), x_Symbol] \rightarrow \operatorname{Simp}[e*(d + e*x)*((a + c*x^2)^{(p + 1)}/(c*(p + 1))), x] - \operatorname{Dist}[e^2*((p + 2)/(c*(p + 1))), \operatorname{Int}[(a + c*x^2)^{(p + 1)}, x], x] /; \operatorname{FreeQ}\{a, c, d, e, p\}, x \ \&\& \operatorname{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\operatorname{IntegerQ}[p] \ \&\& \operatorname{LtQ}[p, -1]$

Rule 5182

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] := Di
st[c^(I*(n/2)), Int[(c + d*x^2)^(p - I*(n/2))*(1 - I*a*x)^(I*n), x], x] /;
FreeQ[{a, c, d, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0]) &&
IGtQ[I*(n/2), 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{\sqrt{c + a^2cx^2}} dx &= c \int \frac{(1 - iax)^2}{(c + a^2cx^2)^{3/2}} dx \\ &= \frac{2i(1 - iax)}{a\sqrt{c + a^2cx^2}} - \int \frac{1}{\sqrt{c + a^2cx^2}} dx \\ &= \frac{2i(1 - iax)}{a\sqrt{c + a^2cx^2}} - \text{Subst}\left(\int \frac{1}{1 - a^2cx^2} dx, x, \frac{x}{\sqrt{c + a^2cx^2}}\right) \\ &= \frac{2i(1 - iax)}{a\sqrt{c + a^2cx^2}} - \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{c + a^2cx^2}}\right)}{a\sqrt{c}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 117, normalized size = 1.86

$$\frac{2\sqrt{1 + a^2x^2} \left((1 - iax)\sqrt{1 + iax} - i\sqrt{1 - iax}(-i + ax)\text{ArcSin}\left(\frac{\sqrt{1 - iax}}{\sqrt{2}}\right) \right)}{a\sqrt{1 - iax}(-i + ax)\sqrt{c + a^2cx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((2*I)*ArcTan[a*x])*Sqrt[c + a^2*c*x^2]),x]

[Out] (2*Sqrt[1 + a^2*x^2]*((1 - I*a*x)*Sqrt[1 + I*a*x] - I*Sqrt[1 - I*a*x]*(-I + a*x)*ArcSin[Sqrt[1 - I*a*x]/Sqrt[2]]))/(a*Sqrt[1 - I*a*x]*(-I + a*x)*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.08, size = 87, normalized size = 1.38

method	result	size
default	$-\frac{\ln\left(\frac{a^2cx}{\sqrt{a^2c}} + \sqrt{a^2cx^2 + c}\right)}{\sqrt{a^2c}} + \frac{2\sqrt{\left(x - \frac{i}{a}\right)^2 a^2c + 2iac\left(x - \frac{i}{a}\right)}}{a^2c\left(x - \frac{i}{a}\right)}$	87

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] $-\ln(a^2cx/(a^2c)^{(1/2)+(a^2cx^2+c)^{(1/2)})/(a^2c)^{(1/2)+2/a^2/c/(x-I/a)} * ((x-I/a)^2a^2c+2Ia*c*(x-I/a))^{(1/2)}$

Maxima [A]

time = 0.46, size = 40, normalized size = 0.63

$$\frac{2i\sqrt{a^2cx^2+c}}{ia^2cx+ac} - \frac{\operatorname{arsinh}(ax)}{a\sqrt{c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] $2*I*\sqrt{a^2*c*x^2+c}/(I*a^2*c*x+a*c) - \operatorname{arcsinh}(a*x)/(a*\sqrt{c})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 152 vs. $2(50) = 100$.

time = 3.01, size = 152, normalized size = 2.41

$$\frac{(a^2cx - iac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(\frac{a^2cx + \sqrt{a^2cx^2+c}}{x} + a^2c\sqrt{\frac{1}{a^2c}}\right)}{x}\right) - (a^2cx - iac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(\frac{a^2cx - \sqrt{a^2cx^2+c}}{x} + a^2c\sqrt{\frac{1}{a^2c}}\right)}{x}\right) - 4\sqrt{a^2cx^2+c}}{2(a^2cx - iac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

[Out] $-1/2*((a^2cx - I*a*c)*\sqrt{1/(a^2c)}*\log(2*(a^2cx + \sqrt{a^2cx^2+c})*a^2c*\sqrt{1/(a^2c)}))/x - (a^2cx - I*a*c)*\sqrt{1/(a^2c)}*\log(2*(a^2cx - \sqrt{a^2cx^2+c})*a^2c*\sqrt{1/(a^2c)}))/x - 4*\sqrt{a^2cx^2+c}/(a^2cx - I*a*c)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{a^2x^2}{a^2x^2\sqrt{a^2cx^2+c} - 2iax\sqrt{a^2cx^2+c} - \sqrt{a^2cx^2+c}} dx - \int \frac{1}{a^2x^2\sqrt{a^2cx^2+c} - 2iax\sqrt{a^2cx^2+c} - \sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)**2*(a**2*x**2+1)/(a**2*c*x**2+c)**(1/2),x)`

[Out] $-\operatorname{Integral}(a**2*x**2/(a**2*x**2*\sqrt{a**2*c*x**2+c} - 2*I*a*x*\sqrt{a**2*c*x**2+c} - \sqrt{a**2*c*x**2+c}), x) - \operatorname{Integral}(1/(a**2*x**2*\sqrt{a**2*c*x**2+c} - 2*I*a*x*\sqrt{a**2*c*x**2+c} - \sqrt{a**2*c*x**2+c}), x)$

Giac [A]

time = 0.44, size = 70, normalized size = 1.11

$$\frac{\log\left(\left|-\sqrt{a^2c}x + \sqrt{a^2cx^2 + c}\right|\right)}{a\sqrt{c}} - \frac{4}{\left(-i\sqrt{a^2c}x + i\sqrt{a^2cx^2 + c} - \sqrt{c}\right)a}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
[Out] log(abs(-sqrt(a^2*c)*x + sqrt(a^2*c*x^2 + c)))/(a*sqrt(c)) - 4/((-I*sqrt(a^2*c)*x + I*sqrt(a^2*c*x^2 + c) - sqrt(c))*a)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{a^2 x^2 + 1}{\sqrt{c a^2 x^2 + c} (1 + a x i)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a^2*x^2 + 1)/((c + a^2*c*x^2)^(1/2)*(a*x*1i + 1)^2),x)
```

```
[Out] int((a^2*x^2 + 1)/((c + a^2*c*x^2)^(1/2)*(a*x*1i + 1)^2), x)
```

$$3.317 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=86

$$-\frac{2\sqrt{1+a^2x^2}}{a(i-ax)\sqrt{c+a^2cx^2}} + \frac{i\sqrt{1+a^2x^2} \log(i-ax)}{a\sqrt{c+a^2cx^2}}$$

[Out] $-2*(a^2*x^2+1)^{(1/2)}/a/(I-a*x)/(a^2*c*x^2+c)^{(1/2)}+I*\ln(I-a*x)*(a^2*x^2+1)^{(1/2)}/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 45}

$$\frac{i\sqrt{a^2x^2+1} \log(-ax+i)}{a\sqrt{a^2cx^2+c}} - \frac{2\sqrt{a^2x^2+1}}{a(-ax+i)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a*x])*Sqrt[c + a^2*c*x^2]),x]

[Out] $(-2*\text{Sqrt}[1 + a^2*x^2])/(a*(I - a*x)*\text{Sqrt}[c + a^2*c*x^2]) + (I*\text{Sqrt}[1 + a^2*x^2]*\text{Log}[I - a*x])/(a*\text{Sqrt}[c + a^2*c*x^2])$

Rule 45

Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_.), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{-3i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-3i \tan^{-1}(ax)}}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \frac{1 - iax}{(1 + iax)^2} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \left(-\frac{2}{(-i+ax)^2} + \frac{i}{-i+ax} \right) dx}{\sqrt{c + a^2 cx^2}} \\
&= -\frac{2\sqrt{1 + a^2 x^2}}{a(i - ax)\sqrt{c + a^2 cx^2}} + \frac{i\sqrt{1 + a^2 x^2} \log(i - ax)}{a\sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 60, normalized size = 0.70

$$\frac{\sqrt{1 + a^2 x^2} \left(-\frac{2}{a(i - ax)} + \frac{i \log(i - ax)}{a} \right)}{\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x])*Sqrt[c + a^2*c*x^2]),x]

[Out] (Sqrt[1 + a^2*x^2]*(-2/(a*(I - a*x)) + (I*Log[I - a*x])/a))/Sqrt[c + a^2*c*x^2]

Maple [A]

time = 0.10, size = 66, normalized size = 0.77

method	result	size
default	$\frac{(-i \ln(-ax+i)ax - \ln(-ax+i) - 2) \sqrt{c(a^2x^2 + 1)}}{\sqrt{a^2x^2 + 1} c(-ax+i)a}$	66
risch	$\frac{2\sqrt{a^2x^2 + 1}}{\sqrt{c(a^2x^2 + 1)} a(ax-i)} + \frac{i\sqrt{a^2x^2 + 1} \ln(ax-i)}{\sqrt{c(a^2x^2 + 1)} a}$	76

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)

[Out] (-I*ln(I-a*x)*a*x - ln(I-a*x) - 2)/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)/c/(I-a*x)/a

Maxima [A]

time = 0.26, size = 35, normalized size = 0.41

$$\frac{i \log(i a x + 1)}{a \sqrt{c}} + \frac{2}{a^2 \sqrt{c} x - i a \sqrt{c}}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")
```

```
[Out] I*log(I*a*x + 1)/(a*sqrt(c)) + 2/(a^2*sqrt(c)*x - I*a*sqrt(c))
```

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 357 vs. 2(71) = 142.

time = 4.54, size = 357, normalized size = 4.15

$$\frac{(-i a^3 c x^3 - a^2 c x^2 - i a c x - c) \sqrt{\frac{1}{a^2 c}} \log\left(\frac{(-i a^3 c x^3 - a^2 c x^2 - i a c x - c) \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1} + (i a^3 c x^3 + a^2 c x^2 + i a c x + c) \sqrt{\frac{1}{a^2 c}}}{8(a^3 c x^3 - i a^2 c x^2 + a c x - i c)}\right) + (i a^3 c x^3 + a^2 c x^2 + i a c x + c) \sqrt{\frac{1}{a^2 c}} \log\left(\frac{(-i a^3 c x^3 - a^2 c x^2 - i a c x - c) \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1} + (i a^3 c x^3 + a^2 c x^2 + i a c x + c) \sqrt{\frac{1}{a^2 c}}}{8(a^3 c x^3 - i a^2 c x^2 + a c x - i c)}\right) - 4i \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1} x}{2(a^3 c x^3 - i a^2 c x^2 + a c x - i c)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")
```

```
[Out] 1/2*((-I*a^3*c*x^3 - a^2*c*x^2 - I*a*c*x - c)*sqrt(1/(a^2*c))*log(1/8*((-I*a^6*x^2 - 2*a^5*x + 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (I*a^9*c*x^4 + 2*a^8*c*x^3 + I*a^7*c*x^2 + 2*a^6*c*x)*sqrt(1/(a^2*c))))/(a^3*x^3 - I*a^2*x^2 + a*x - I) + (I*a^3*c*x^3 + a^2*c*x^2 + I*a*c*x + c)*sqrt(1/(a^2*c))*log(1/8*((-I*a^6*x^2 - 2*a^5*x + 2*I*a^4)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1) + (-I*a^9*c*x^4 - 2*a^8*c*x^3 - I*a^7*c*x^2 - 2*a^6*c*x)*sqrt(1/(a^2*c))))/(a^3*x^3 - I*a^2*x^2 + a*x - I) - 4*I*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)*x/(a^3*c*x^3 - I*a^2*c*x^2 + a*c*x - I*c)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \frac{\sqrt{a^2 x^2 + 1}}{a^3 x^3 \sqrt{a^2 c x^2 + c} - 3i a^2 x^2 \sqrt{a^2 c x^2 + c} - 3a x \sqrt{a^2 c x^2 + c} + i \sqrt{a^2 c x^2 + c}} dx + \int \frac{a^2 x^2 \sqrt{a^2 x^2 + 1}}{a^3 x^3 \sqrt{a^2 c x^2 + c} - 3i a^2 x^2 \sqrt{a^2 c x^2 + c} - 3a x \sqrt{a^2 c x^2 + c} + i \sqrt{a^2 c x^2 + c}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/(a**2*c*x**2+c)**(1/2),x)
```

```
[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**3*x**3*sqrt(a**2*c*x**2 + c) - 3*I*a**2*x**2*sqrt(a**2*c*x**2 + c) - 3*a*x*sqrt(a**2*c*x**2 + c) + I*sqrt(a**2*c*x**2 + c)), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**3*x**3*sqrt(a**2*c*x**2 + c) - 3*I*a**2*x**2*sqrt(a**2*c*x**2 + c) - 3*a*x*sqrt(a**2*c*x**2 + c) + I*sqrt(a**2*c*x**2 + c)), x))
```


Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
[Out] integrate((a^2*x^2 + 1)^(3/2)/(sqrt(a^2*c*x^2 + c)*(I*a*x + 1)^3), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(a^2 x^2 + 1)^{3/2}}{\sqrt{c a^2 x^2 + c} (1 + a x i)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a^2*x^2 + 1)^(3/2)/((c + a^2*c*x^2)^(1/2)*(a*x*i + 1)^3),x)
```

```
[Out] int((a^2*x^2 + 1)^(3/2)/((c + a^2*c*x^2)^(1/2)*(a*x*i + 1)^3), x)
```

$$3.318 \quad \int \frac{e^{-4i \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=96

$$\frac{2ic(1-iax)^3}{3a(c+a^2cx^2)^{3/2}} - \frac{2i(1-iax)}{a\sqrt{c+a^2cx^2}} + \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{c+a^2cx^2}}\right)}{a\sqrt{c}}$$

[Out] $2/3*I*c*(1-I*a*x)^3/a/(a^2*c*x^2+c)^{(3/2)}+\operatorname{arctanh}(a*x*c^{(1/2)}/(a^2*c*x^2+c)^{(1/2)})/a/c^{(1/2)}-2*I*(1-I*a*x)/a/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 96, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.200$, Rules used = {5182, 683, 667, 223, 212}

$$\frac{2ic(1-iax)^3}{3a(a^2cx^2+c)^{3/2}} - \frac{2i(1-iax)}{a\sqrt{a^2cx^2+c}} + \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{a^2cx^2+c}}\right)}{a\sqrt{c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[1/(E^{((4*I)*\operatorname{ArcTan}[a*x])}*\operatorname{Sqrt}[c + a^2*c*x^2]), x]$

[Out] $((2*I)/3)*c*(1 - I*a*x)^3/(a*(c + a^2*c*x^2)^{(3/2)}) - ((2*I)*(1 - I*a*x))/(a*\operatorname{Sqrt}[c + a^2*c*x^2]) + \operatorname{ArcTanh}[(a*\operatorname{Sqrt}[c]*x)/\operatorname{Sqrt}[c + a^2*c*x^2]]/(a*\operatorname{Sqrt}[c])$

Rule 212

$\operatorname{Int}[(a_ + (b_)*(x_)^2)^{-1}, x_Symbol] \rightarrow \operatorname{Simp}[(1/(\operatorname{Rt}[a, 2]*\operatorname{Rt}[-b, 2]))*\operatorname{ArcTanh}[\operatorname{Rt}[-b, 2]*(x/\operatorname{Rt}[a, 2])], x] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \operatorname{NegQ}[a/b] \ \&\& (\operatorname{GtQ}[a, 0] \ || \ \operatorname{LtQ}[b, 0])$

Rule 223

$\operatorname{Int}[1/\operatorname{Sqrt}[(a_ + (b_)*(x_)^2)], x_Symbol] \rightarrow \operatorname{Subst}[\operatorname{Int}[1/(1 - b*x^2), x], x, x/\operatorname{Sqrt}[a + b*x^2]] /; \operatorname{FreeQ}\{a, b\}, x] \ \&\& \ !\operatorname{GtQ}[a, 0]$

Rule 667

$\operatorname{Int}[(d_ + (e_)*(x_))^2*((a_ + (c_)*(x_)^2)^{p_}), x_Symbol] \rightarrow \operatorname{Simp}[e*(d + e*x)*((a + c*x^2)^{(p+1)}/(c*(p+1))), x] - \operatorname{Dist}[e^2*((p+2)/(c*(p+1))), \operatorname{Int}[(a + c*x^2)^{(p+1)}, x], x] /; \operatorname{FreeQ}\{a, c, d, e, p\}, x] \ \&\& \operatorname{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\operatorname{IntegerQ}[p] \ \&\& \operatorname{LtQ}[p, -1]$

Rule 683

```
Int[((d_) + (e_.)*(x_))^(m_)*((a_) + (c_.)*(x_)^2)^(p_), x_Symbol] := Simp[
e*(d + e*x)^(m - 1)*((a + c*x^2)^(p + 1)/(c*(p + 1))), x] - Dist[e^2*((m +
p)/(c*(p + 1))), Int[(d + e*x)^(m - 2)*(a + c*x^2)^(p + 1), x], x] /; FreeQ
[{a, c, d, e}, x] && EqQ[c*d^2 + a*e^2, 0] && LtQ[p, -1] && GtQ[m, 1] && In
tegerQ[2*p]
```

Rule 5182

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Di
st[c^(I*(n/2)), Int[(c + d*x^2)^(p - I*(n/2))*(1 - I*a*x)^(I*n), x], x] /;
FreeQ[{a, c, d, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0]) &&
IGtQ[I*(n/2), 0]
```

Rubi steps

$$\begin{aligned} \int \frac{e^{-4i \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= c^2 \int \frac{(1 - iax)^4}{(c + a^2 cx^2)^{5/2}} dx \\ &= \frac{2ic(1 - iax)^3}{3a(c + a^2 cx^2)^{3/2}} - c \int \frac{(1 - iax)^2}{(c + a^2 cx^2)^{3/2}} dx \\ &= \frac{2ic(1 - iax)^3}{3a(c + a^2 cx^2)^{3/2}} - \frac{2i(1 - iax)}{a\sqrt{c + a^2 cx^2}} + \int \frac{1}{\sqrt{c + a^2 cx^2}} dx \\ &= \frac{2ic(1 - iax)^3}{3a(c + a^2 cx^2)^{3/2}} - \frac{2i(1 - iax)}{a\sqrt{c + a^2 cx^2}} + \text{Subst}\left(\int \frac{1}{1 - a^2 cx^2} dx, x, \frac{x}{\sqrt{c + a^2 cx^2}}\right) \\ &= \frac{2ic(1 - iax)^3}{3a(c + a^2 cx^2)^{3/2}} - \frac{2i(1 - iax)}{a\sqrt{c + a^2 cx^2}} + \frac{\tanh^{-1}\left(\frac{a\sqrt{c}x}{\sqrt{c + a^2 cx^2}}\right)}{a\sqrt{c}} \end{aligned}$$

Mathematica [A]

time = 0.06, size = 132, normalized size = 1.38

$$\frac{2\sqrt{1 + a^2 x^2} \left(2i\sqrt{1 + iax} (1 + iax + 2a^2 x^2) + 3i\sqrt{1 - iax} (-i + ax)^2 \text{ArcSin}\left(\frac{\sqrt{1 - iax}}{\sqrt{2}}\right) \right)}{3a\sqrt{1 - iax} (-i + ax)^2 \sqrt{c + a^2 cx^2}}$$

Warning: Unable to verify antiderivative.

```
[In] Integrate[1/(E^((4*I)*ArcTan[a*x])*Sqrt[c + a^2*c*x^2]),x]
```

```
[Out] (2*Sqrt[1 + a^2*x^2]*((2*I)*Sqrt[1 + I*a*x]*(1 + I*a*x + 2*a^2*x^2) + (3*I)
*Sqrt[1 - I*a*x]*(-I + a*x)^2*ArcSin[Sqrt[1 - I*a*x]/Sqrt[2]]))/(3*a*Sqrt[1
- I*a*x]*(-I + a*x)^2*Sqrt[c + a^2*c*x^2])
```

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 187 vs. 2(80) = 160.
time = 0.09, size = 188, normalized size = 1.96

method	result
default	$\frac{\ln\left(\frac{a^2cx}{\sqrt{a^2c}} + \sqrt{a^2cx^2 + c}\right)}{\sqrt{a^2c}} - \frac{4\sqrt{\left(x - \frac{i}{a}\right)^2 a^2c + 2iac\left(x - \frac{i}{a}\right)}}{a^2c\left(x - \frac{i}{a}\right)} - \frac{4\left(\frac{i\sqrt{\left(x - \frac{i}{a}\right)^2 a^2c + 2iac\left(x - \frac{i}{a}\right)}}{3ac\left(x - \frac{i}{a}\right)^2}\right)}{4}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x,method=_RETURNVERBOSE)`

[Out] $\ln(a^2cx/(a^2c)^{(1/2)} + (a^2cx^2+c)^{(1/2)})/(a^2c)^{(1/2)} - 4/a^2c/(x-I/a) * ((x-I/a)^2 a^2c + 2Iac(x-I/a))^{(1/2)} - 4/a^2 * (1/3I/a/c/(x-I/a)^2 * ((x-I/a)^2 a^2c + 2Iac(x-I/a))^{(1/2)} - 1/3c/(x-I/a) * ((x-I/a)^2 a^2c + 2Iac(x-I/a))^{(1/2)})$

Maxima [A]

time = 0.46, size = 76, normalized size = 0.79

$$-\frac{4i\sqrt{a^2cx^2+c}}{3(a^3cx^2-2ia^2cx-ac)} - \frac{8i\sqrt{a^2cx^2+c}}{3ia^2cx+3ac} + \frac{\operatorname{arsinh}(ax)}{a\sqrt{c}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] $-4/3I*\sqrt{a^2cx^2+c}/(a^3cx^2-2Ia^2cx-ac) - 8I*\sqrt{a^2cx^2+c}/(3Ia^2cx+3ac) + \operatorname{arcsinh}(ax)/(a*\sqrt{c})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 186 vs. 2(75) = 150.
time = 2.27, size = 186, normalized size = 1.94

$$\frac{3(a^3cx^2-2ia^2cx-ac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(a^2cx+\sqrt{a^2cx^2+c}a^2c\sqrt{\frac{1}{a^2c}}\right)}{x}\right) - 3(a^3cx^2-2ia^2cx-ac)\sqrt{\frac{1}{a^2c}} \log\left(\frac{2\left(a^2cx-\sqrt{a^2cx^2+c}a^2c\sqrt{\frac{1}{a^2c}}\right)}{x}\right) - 8\sqrt{a^2cx^2+c}(2ax-i)}{6(a^3cx^2-2ia^2cx-ac)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

[Out] $\frac{1}{6} * (3 * (a^3 * c * x^2 - 2 * I * a^2 * c * x - a * c) * \sqrt{1 / (a^2 * c)}) * \log(2 * (a^2 * c * x + \sqrt{a^2 * c * x^2 + c}) * a^2 * c * \sqrt{1 / (a^2 * c)}) / x - 3 * (a^3 * c * x^2 - 2 * I * a^2 * c * x - a * c) * \sqrt{1 / (a^2 * c)}) * \log(2 * (a^2 * c * x - \sqrt{a^2 * c * x^2 + c}) * a^2 * c * \sqrt{1 / (a^2 * c)}) / x - 8 * \sqrt{a^2 * c * x^2 + c} * (2 * a * x - I) / (a^3 * c * x^2 - 2 * I * a^2 * c * x - a * c)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a^2 x^2 + 1)^2}{\sqrt{c(a^2 x^2 + 1)} (ax - i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**4*(a**2*x**2+1)**2/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral((a**2*x**2 + 1)**2/(sqrt(c*(a**2*x**2 + 1))*(a*x - I)**4), x)

Giac [A]

time = 0.47, size = 132, normalized size = 1.38

$$\frac{\log\left(\left|-\sqrt{a^2 c} x + \sqrt{a^2 c x^2 + c}\right|\right)}{a \sqrt{c}} - \frac{8\left(3\left(\sqrt{a^2 c} x - \sqrt{a^2 c x^2 + c}\right)^2 - 3i\left(\sqrt{a^2 c} x - \sqrt{a^2 c x^2 + c}\right)\sqrt{c} - 2c\right)}{3\left(-i\sqrt{a^2 c} x + i\sqrt{a^2 c x^2 + c} - \sqrt{c}\right)^3 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] $-\log(\text{abs}(-\sqrt{a^2 * c} * x + \sqrt{a^2 * c * x^2 + c})) / (a * \sqrt{c}) - 8 / 3 * (3 * (\sqrt{a^2 * c} * x - \sqrt{a^2 * c * x^2 + c})^2 - 3 * I * (\sqrt{a^2 * c} * x - \sqrt{a^2 * c * x^2 + c}) * \sqrt{c} - 2 * c) / ((-I * \sqrt{a^2 * c} * x + I * \sqrt{a^2 * c * x^2 + c} - \sqrt{c})^3 * a)$

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{(a^2 x^2 + 1)^2}{\sqrt{c a^2 x^2 + c} (1 + a x i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^2/((c + a^2*c*x^2)^(1/2)*(a*x*I + 1)^4),x)

[Out] int((a^2*x^2 + 1)^2/((c + a^2*c*x^2)^(1/2)*(a*x*I + 1)^4), x)

$$3.319 \quad \int \frac{e^{5i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=35

$$-\frac{2}{3a(i+ax)^3} - \frac{i}{2a(i+ax)^2}$$

[Out] $-2/3/a/(I+a*x)^3-1/2*I/a/(I+a*x)^2$

Rubi [A]

time = 0.03, antiderivative size = 35, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 45}

$$-\frac{i}{2a(ax+i)^2} - \frac{2}{3a(ax+i)^3}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((5*I)*\text{ArcTan}[a*x])}/(1 + a^2*x^2)^{(3/2)}, x]$

[Out] $-2/(3*a*(I + a*x)^3) - (I/2)/(a*(I + a*x)^2)$

Rule 45

$\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{5i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1+iax}{(1-iax)^4} dx \\ &= \int \left(\frac{2}{(i+ax)^4} + \frac{i}{(i+ax)^3} \right) dx \\ &= -\frac{2}{3a(i+ax)^3} - \frac{i}{2a(i+ax)^2} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 24, normalized size = 0.69

$$-\frac{1 + 3iax}{6a(i + ax)^3}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((5*I)*ArcTan[a*x])/(1 + a^2*x^2)^(3/2), x]``[Out] -1/6*(1 + (3*I)*a*x)/(a*(I + a*x)^3)`**Maple [A]**

time = 0.09, size = 20, normalized size = 0.57

method	result
default	$-\frac{\frac{ix}{2} - \frac{1}{6a}}{(ax+i)^3}$
risch	$-\frac{\frac{ix}{2} - \frac{1}{6a}}{(ax+i)^3}$
norman	$\frac{x + \frac{5}{2}iax^2 + \frac{1}{6}ia^5x^6 - \frac{5}{3}a^2x^3}{(a^2x^2+1)^3}$
meijerg	$\frac{x\sqrt{a^2} \left(\frac{15a^4x^4 + 40a^2x^2 + 33}{4(a^2x^2+1)^3} + \frac{15\sqrt{a^2} \arctan(ax)}{4a} \right) + \frac{5iax^2(a^4x^4 + 3a^2x^2 + 3)}{6(a^2x^2+1)^3} - \frac{5 \left(-\frac{x(a^2)^{\frac{3}{2}}(-3a^4x^4 - 8a^2x^2 + 3)}{4a^2(a^2x^2+1)^3} + \frac{3(a^2)^{\frac{3}{2}} \arctan(ax)}{4a^3} \right)}{6\sqrt{a^2}}}{12\sqrt{a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^5/(a^2*x^2+1)^4,x,method=_RETURNVERBOSE)``[Out] (-1/2*I*x-1/6/a)/(I+a*x)^3`**Maxima [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 59 vs. 2(25) = 50.

time = 0.48, size = 59, normalized size = 1.69

$$-\frac{3ia^4x^4 + 10a^3x^3 - 12ia^2x^2 - 6ax + i}{6(a^7x^6 + 3a^5x^4 + 3a^3x^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^4,x, algorithm="maxima")``[Out] -1/6*(3*I*a^4*x^4 + 10*a^3*x^3 - 12*I*a^2*x^2 - 6*a*x + I)/(a^7*x^6 + 3*a^5*x^4 + 3*a^3*x^2 + a)`**Fricas [A]**

time = 5.41, size = 35, normalized size = 1.00

$$\frac{-3iax - 1}{6(a^4x^3 + 3ia^3x^2 - 3a^2x - ia)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^4,x, algorithm="fricas")

[Out] 1/6*(-3*I*a*x - 1)/(a^4*x^3 + 3*I*a^3*x^2 - 3*a^2*x - I*a)

Sympy [A]

time = 0.14, size = 39, normalized size = 1.11

$$\frac{-3iax - 1}{6a^4x^3 + 18ia^3x^2 - 18a^2x - 6ia}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**5/(a**2*x**2+1)**4,x)

[Out] (-3*I*a*x - 1)/(6*a**4*x**3 + 18*I*a**3*x**2 - 18*a**2*x - 6*I*a)

Giac [A]

time = 0.40, size = 18, normalized size = 0.51

$$-\frac{3iax + 1}{6(ax + i)^3a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^4,x, algorithm="giac")

[Out] -1/6*(3*I*a*x + 1)/((a*x + I)^3*a)

Mupad [B]

time = 0.10, size = 21, normalized size = 0.60

$$-\frac{3ax - i}{6a(-1 + ax1i)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^5/(a^2*x^2 + 1)^4,x)

[Out] -(3*a*x - 1i)/(6*a*(a*x*1i - 1)^3)

$$3.320 \quad \int \frac{e^{4i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=67

$$-\frac{i(1+iax)^{3/2}}{5a(1-iax)^{5/2}} - \frac{i(1+iax)^{3/2}}{15a(1-iax)^{3/2}}$$

[Out] $-1/5*I*(1+I*a*x)^{(3/2)}/a/(1-I*a*x)^{(5/2)}-1/15*I*(1+I*a*x)^{(3/2)}/a/(1-I*a*x)^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5181, 47, 37}

$$-\frac{i(1+iax)^{3/2}}{15a(1-iax)^{3/2}} - \frac{i(1+iax)^{3/2}}{5a(1-iax)^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((4*I)*\text{ArcTan}[a*x])}/(1 + a^2*x^2)^{(3/2)}, x]$

[Out] $((-1/5*I)*(1 + I*a*x)^{(3/2)})/(a*(1 - I*a*x)^{(5/2)}) - ((I/15)*(1 + I*a*x)^{(3/2)})/(a*(1 - I*a*x)^{(3/2)})$

Rule 37

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1)}/((b*c - a*d)*(m + 1)))}, x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[m + n + 2, 0] \&\& \text{NeQ}[m, -1]$

Rule 47

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1)}/((b*c - a*d)*(m + 1)))}, x] - \text{Dist}[d*(\text{Simplify}[m + n + 2]/((b*c - a*d)*(m + 1))), \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]}*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{I} \text{LtQ}[\text{Simplify}[m + n + 2], 0] \&\& \text{NeQ}[m, -1] \&\& !(\text{LtQ}[m, -1] \&\& \text{LtQ}[n, -1] \&\& (\text{EqQ}[a, 0] || (\text{NeQ}[c, 0] \&\& \text{LtQ}[m - n, 0] \&\& \text{IntegerQ}[n]))) \&\& (\text{SumSimplerQ}[m, 1] || !\text{SumSimplerQ}[n, 1])$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /$

; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{\sqrt{1+iax}}{(1-iax)^{7/2}} dx \\ &= -\frac{i(1+iax)^{3/2}}{5a(1-iax)^{5/2}} + \frac{1}{5} \int \frac{\sqrt{1+iax}}{(1-iax)^{5/2}} dx \\ &= -\frac{i(1+iax)^{3/2}}{5a(1-iax)^{5/2}} - \frac{i(1+iax)^{3/2}}{15a(1-iax)^{3/2}} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 47, normalized size = 0.70

$$\frac{(1+iax)^{3/2}(4i+ax)}{15a\sqrt{1-iax}(i+ax)^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((4*I)*ArcTan[a*x])/(1+a^2*x^2)^(3/2), x]

[Out] ((1+I*a*x)^(3/2)*(4*I+a*x))/(15*a*Sqrt[1-I*a*x]*(I+a*x)^2)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 268 vs. 2(49) = 98.

time = 0.09, size = 269, normalized size = 4.01

method	result
trager	$\frac{-a^5x^5-10a^3x^3+20ia^2x^2+15ax-4i}{15(a^2x^2+1)^{\frac{5}{2}}a}$
meijerg	$\frac{x(8a^4x^4+20a^2x^2+15)}{15(a^2x^2+1)^{\frac{5}{2}}} + \frac{16i\left(\frac{3\sqrt{\pi}}{4} - \frac{3\sqrt{\pi}}{4(a^2x^2+1)^{\frac{5}{2}}}\right)}{15a\sqrt{\pi}} - \frac{2a^2x^3(2a^2x^2+5)}{5(a^2x^2+1)^{\frac{5}{2}}} - \frac{16i\left(\frac{\sqrt{\pi}}{2} - \frac{\sqrt{\pi}(20a^2x^2+8)}{16(a^2x^2+1)^{\frac{5}{2}}}\right)}{15a\sqrt{\pi}} + \frac{a^4x^5}{5(a^2x^2+1)}$
default	$\frac{x}{5(a^2x^2+1)^{\frac{5}{2}}} + \frac{4x}{15(a^2x^2+1)^{\frac{3}{2}}} + \frac{8x}{15\sqrt{a^2x^2+1}} + a^4 \left(-\frac{x^3}{2a^2(a^2x^2+1)^{\frac{5}{2}}} + \frac{-\frac{3x}{8a^2(a^2x^2+1)^{\frac{5}{2}}} + \frac{3\left(\frac{x}{5(a^2x^2+1)^{\frac{5}{2}}} + \frac{4x}{15(a^2x^2+1)}\right)}{a^2} \right)$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)^4/(a^2*x^2+1)^(7/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{5}x/(a^2x^2+1)^{5/2} + \frac{4}{15}x/(a^2x^2+1)^{3/2} + \frac{8}{15}x/(a^2x^2+1)^{1/2} + a^4(-\frac{1}{2}x^3/a^2/(a^2x^2+1)^{5/2} + \frac{3}{2}a^2(-\frac{1}{4}x/a^2/(a^2x^2+1)^{5/2} + \frac{1}{4}a^2(1/5x/(a^2x^2+1)^{5/2} + \frac{4}{15}x/(a^2x^2+1)^{3/2} + \frac{8}{15}x/(a^2x^2+1)^{1/2}))) - 4Ia^3(-\frac{1}{3}x^2/a^2/(a^2x^2+1)^{5/2} - \frac{2}{15}a^4/(a^2x^2+1)^{5/2}) - 6a^2(-\frac{1}{4}x/a^2/(a^2x^2+1)^{5/2} + \frac{1}{4}a^2(1/5x/(a^2x^2+1)^{5/2} + \frac{4}{15}x/(a^2x^2+1)^{3/2} + \frac{8}{15}x/(a^2x^2+1)^{1/2})) - \frac{4}{5}I/a/(a^2x^2+1)^{5/2}$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 95 vs. $2(43) = 86$.

time = 0.25, size = 95, normalized size = 1.42

$$-\frac{a^2x^3}{2(a^2x^2+1)^{5/2}} - \frac{x}{15\sqrt{a^2x^2+1}} + \frac{4iax^2}{3(a^2x^2+1)^{5/2}} - \frac{x}{30(a^2x^2+1)^{3/2}} + \frac{11x}{10(a^2x^2+1)^{5/2}} - \frac{4i}{15(a^2x^2+1)^{5/2}}a$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^4/(a^2*x^2+1)^(7/2),x, algorithm="maxima")`

[Out] $-\frac{1}{2}a^2x^3/(a^2x^2+1)^{5/2} - \frac{1}{15}x/\sqrt{a^2x^2+1} + \frac{4}{3}Ia^3x^2/(a^2x^2+1)^{5/2} - \frac{1}{30}x/(a^2x^2+1)^{3/2} + \frac{11}{10}x/(a^2x^2+1)^{5/2} - \frac{4}{15}I/(a^2x^2+1)^{5/2}a$

Fricas [A]

time = 3.28, size = 75, normalized size = 1.12

$$-\frac{a^3x^3 + 3ia^2x^2 - 3ax + (a^2x^2 + 3iax + 4)\sqrt{a^2x^2 + 1} - i}{15(a^4x^3 + 3ia^3x^2 - 3a^2x - ia)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^4/(a^2*x^2+1)^(7/2),x, algorithm="fricas")`

[Out] $-\frac{1}{15}(a^3x^3 + 3Ia^2x^2 - 3a^2x + (a^2x^2 + 3Ia^2x + 4)\sqrt{a^2x^2 + 1} - I)/(a^4x^3 + 3Ia^3x^2 - 3a^2x - Ia)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(ax - i)^4}{(a^2x^2 + 1)^{7/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**4/(a**2*x**2+1)**(7/2),x)`

[Out] `Integral((a*x - I)**4/(a**2*x**2 + 1)**(7/2), x)`

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(43) = 86$.

time = 0.42, size = 111, normalized size = 1.66

$$\frac{2 \left(4a^4 - 25a^2 \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^2 + 15ia \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^3 + 15 \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^4 - 5a^3 \left(i \sqrt{a^2 + \frac{1}{x^2}} - \frac{i}{x} \right) \right)}{15 \left(ia + \sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^(7/2),x, algorithm="giac")

[Out] $\frac{2}{15} * (4a^4 - 25a^2 * (\sqrt{a^2 + 1/x^2} - 1/x)^2 + 15Ia * (\sqrt{a^2 + 1/x^2} - 1/x)^3 + 15 * (\sqrt{a^2 + 1/x^2} - 1/x)^4 - 5a^3 * (I * \sqrt{a^2 + 1/x^2} - I/x)) / (Ia + \sqrt{a^2 + 1/x^2} - 1/x)^5$

Mupad [B]

time = 0.52, size = 41, normalized size = 0.61

$$\frac{\sqrt{a^2 x^2 + 1} (a^2 x^2 i - 3 a x + 4 i)}{15 a (-1 + a x i)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/(a^2*x^2 + 1)^(7/2),x)

[Out] $((a^2 * x^2 + 1)^{(1/2)} * (a^2 * x^2 * 1i - 3 * a * x + 4i)) / (15 * a * (a * x * 1i - 1)^3)$

$$3.321 \quad \int \frac{e^{3i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=19

$$-\frac{i}{2a(1-iax)^2}$$

[Out] $-1/2*I/a/(1-I*a*x)^2$

Rubi [A]

time = 0.03, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 32}

$$-\frac{i}{2a(1-iax)^2}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)*\text{ArcTan}[a*x])}/(1 + a^2*x^2)^{(3/2)}, x]$

[Out] $(-1/2*I)/(a*(1 - I*a*x)^2)$

Rule 32

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] /;$ FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_. + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{3i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1}{(1-iax)^3} dx \\ &= -\frac{i}{2a(1-iax)^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 18, normalized size = 0.95

$$\frac{i}{2a(i+ax)^2}$$

Antiderivative was successfully verified.

[In] Integrate[E^((3*I)*ArcTan[a*x])/(1 + a^2*x^2)^(3/2), x]

[Out] (I/2)/(a*(I + a*x)^2)

Maple [A]

time = 0.08, size = 15, normalized size = 0.79

method	result	s
default	$\frac{i}{2a(ax+i)^2}$	1
risch	$\frac{i}{2a(ax+i)^2}$	1
norman	$\frac{x + \frac{3}{2}iax^2 + \frac{1}{2}ia^3x^4}{(a^2x^2+1)^2}$	3
gospers	$-\frac{(ax+i)(iax+1)^3}{2a(a^2x^2+1)^3}$	3
meijerg	$\frac{x\sqrt{a^2}(3a^2x^2+5)}{2(a^2x^2+1)^2} + \frac{3\sqrt{a^2}\arctan(ax)}{2a} + \frac{3iax^2(a^2x^2+2)}{4(a^2x^2+1)^2} - \frac{3\left(-\frac{x(a^2)^{\frac{3}{2}}(-3a^2x^2+3)}{6a^2(a^2x^2+1)^2} + \frac{(a^2)^{\frac{3}{2}}\arctan(ax)}{2a^3}\right)}{4\sqrt{a^2}} - \frac{ia^3x^4}{4(a^2x^2+1)^2}$	1

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^3,x,method=_RETURNVERBOSE)

[Out] 1/2*I/a/(I+a*x)^2

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 35 vs. $2(13) = 26$.

time = 0.47, size = 35, normalized size = 1.84

$$-\frac{-i a^2 x^2 - 2 a x + i}{2 (a^5 x^4 + 2 a^3 x^2 + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^3,x, algorithm="maxima")

[Out] -1/2*(-I*a^2*x^2 - 2*a*x + I)/(a^5*x^4 + 2*a^3*x^2 + a)

Fricas [A]

time = 1.66, size = 21, normalized size = 1.11

$$\frac{i}{2(a^3x^2 + 2ia^2x - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^3,x, algorithm="fricas")

[Out] $1/2*I/(a^3*x^2 + 2*I*a^2*x - a)$

Sympy [A]

time = 0.09, size = 20, normalized size = 1.05

$$\frac{i}{2a^3x^2 + 4ia^2x - 2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**3/(a**2*x**2+1)**3,x)`

[Out] $I/(2*a**3*x**2 + 4*I*a**2*x - 2*a)$

Giac [A]

time = 0.40, size = 12, normalized size = 0.63

$$\frac{i}{2(ax + i)^2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^3,x, algorithm="giac")`

[Out] $1/2*I/((a*x + I)^2*a)$

Mupad [B]

time = 0.50, size = 24, normalized size = 1.26

$$\frac{1i}{2(a^3x^2 + a^2x2i - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((a*x*1i + 1)^3/(a^2*x^2 + 1)^3,x)`

[Out] $1i/(2*(a^2*x*2i - a + a^3*x^2))$

$$3.322 \quad \int \frac{e^{2i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=67

$$-\frac{i\sqrt{1+iax}}{3a(1-iax)^{3/2}} - \frac{i\sqrt{1+iax}}{3a\sqrt{1-iax}}$$

[Out] $-1/3*I*(1+I*a*x)^{(1/2)}/a/(1-I*a*x)^{(3/2)}-1/3*I*(1+I*a*x)^{(1/2)}/a/(1-I*a*x)^{(1/2)}$

Rubi [A]

time = 0.03, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5181, 47, 37}

$$-\frac{i\sqrt{1+iax}}{3a\sqrt{1-iax}} - \frac{i\sqrt{1+iax}}{3a(1-iax)^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}/(1 + a^2*x^2)^{(3/2)}, x]$

[Out] $((-1/3*I)*\text{Sqrt}[1 + I*a*x])/(a*(1 - I*a*x)^{(3/2)}) - ((I/3)*\text{Sqrt}[1 + I*a*x])/(a*\text{Sqrt}[1 - I*a*x])$

Rule 37

$\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1))/(b*c - a*d)*(m + 1))}, x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[m + n + 2, 0] \&\& \text{NeQ}[m, -1]$

Rule 47

$\text{Int}[(a_.) + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1))/(b*c - a*d)*(m + 1))}, x] - \text{Dist}[d*(\text{Simplify}[m + n + 2]/((b*c - a*d)*(m + 1))), \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]*(c + d*x)^n}, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{LtQ}[\text{Simplify}[m + n + 2], 0] \&\& \text{NeQ}[m, -1] \&\& !(\text{LtQ}[m, -1] \&\& \text{LtQ}[n, -1] \&\& (\text{EqQ}[a, 0] \|\| (\text{NeQ}[c, 0] \&\& \text{LtQ}[m - n, 0] \&\& \text{IntegerQ}[n]))) \&\& (\text{SumSimplerQ}[m, 1] \|\| !\text{SumSimplerQ}[n, 1])$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x\} \&\& \text{EqQ}[d, a^2*c] \&\& (\text{IntegerQ}[p] \|\| \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned}
\int \frac{e^{2i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1}{(1-iax)^{5/2} \sqrt{1+iax}} dx \\
&= -\frac{i\sqrt{1+iax}}{3a(1-iax)^{3/2}} + \frac{1}{3} \int \frac{1}{(1-iax)^{3/2} \sqrt{1+iax}} dx \\
&= -\frac{i\sqrt{1+iax}}{3a(1-iax)^{3/2}} - \frac{i\sqrt{1+iax}}{3a\sqrt{1-iax}}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 48, normalized size = 0.72

$$\frac{(2-iax)\sqrt{1+iax}}{3a\sqrt{1-iax}(i+ax)}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((2*I)*ArcTan[a*x])/(1 + a^2*x^2)^(3/2), x]``[Out] ((2 - I*a*x)*Sqrt[1 + I*a*x])/(3*a*Sqrt[1 - I*a*x]*(I + a*x))`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 103 vs. 2(49) = 98.

time = 0.08, size = 104, normalized size = 1.55

method	result	size
trager	$\frac{a^3x^3+3ax-2i}{3(a^2x^2+1)^{\frac{3}{2}}a}$	31
meijerg	$\frac{x(2a^2x^2+3)}{3(a^2x^2+1)^{\frac{3}{2}}} + \frac{4i\left(\frac{\sqrt{\pi}}{2} - \frac{\sqrt{\pi}}{2(a^2x^2+1)^{\frac{3}{2}}}\right)}{3a\sqrt{\pi}} - \frac{a^2x^3}{3(a^2x^2+1)^{\frac{3}{2}}}$	76
default	$\frac{x}{3(a^2x^2+1)^{\frac{3}{2}}} + \frac{2x}{3\sqrt{a^2x^2+1}} - a^2\left(-\frac{x}{2a^2(a^2x^2+1)^{\frac{3}{2}}} + \frac{\frac{x}{3(a^2x^2+1)^{\frac{3}{2}}} + \frac{2x}{3\sqrt{a^2x^2+1}}}{2a^2}\right) - \frac{2i}{3a(a^2x^2+1)^{\frac{3}{2}}}$	104

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^2/(a^2*x^2+1)^(5/2), x, method=_RETURNVERBOSE)`
`[Out] 1/3*x/(a^2*x^2+1)^(3/2)+2/3*x/(a^2*x^2+1)^(1/2)-a^2*(-1/2*x/a^2/(a^2*x^2+1)^(3/2)+1/2/a^2*(1/3*x/(a^2*x^2+1)^(3/2)+2/3*x/(a^2*x^2+1)^(1/2)))-2/3*I/a/(a^2*x^2+1)^(3/2)`

Maxima [A]

time = 0.26, size = 45, normalized size = 0.67

$$\frac{x}{3\sqrt{a^2x^2+1}} + \frac{2x}{3(a^2x^2+1)^{\frac{3}{2}}} - \frac{2i}{3(a^2x^2+1)^{\frac{3}{2}}a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)^(5/2),x, algorithm="maxima")``[Out] 1/3*x/sqrt(a^2*x^2 + 1) + 2/3*x/(a^2*x^2 + 1)^(3/2) - 2/3*I/((a^2*x^2 + 1)^(3/2)*a)`**Fricas [A]**

time = 2.56, size = 51, normalized size = 0.76

$$\frac{a^2x^2 + 2i ax + \sqrt{a^2x^2 + 1} (ax + 2i) - 1}{3(a^3x^2 + 2i a^2x - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)^2/(a^2*x^2+1)^(5/2),x, algorithm="fricas")``[Out] 1/3*(a^2*x^2 + 2*I*a*x + sqrt(a^2*x^2 + 1)*(a*x + 2*I) - 1)/(a^3*x^2 + 2*I*a^2*x - a)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{a^2x^2}{a^4x^4\sqrt{a^2x^2+1} + 2a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} dx - \int \left(\frac{2iax}{a^4x^4\sqrt{a^2x^2+1} + 2a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx - \int \left(\frac{1}{a^4x^4\sqrt{a^2x^2+1} + 2a^2x^2\sqrt{a^2x^2+1} + \sqrt{a^2x^2+1}} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate((1+I*a*x)**2/(a**2*x**2+1)**(5/2),x)``[Out] -Integral(a**2*x**2/(a**4*x**4*sqrt(a**2*x**2 + 1) + 2*a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) - Integral(-2*I*a*x/(a**4*x**4*sqrt(a**2*x**2 + 1) + 2*a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x) - Integral(-1/(a**4*x**4*sqrt(a**2*x**2 + 1) + 2*a**2*x**2*sqrt(a**2*x**2 + 1) + sqrt(a**2*x**2 + 1)), x)`**Giac [A]**

time = 0.44, size = 67, normalized size = 1.00

$$\frac{2 \left(2a^2 - 3 \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^2 + 3a \left(-i \sqrt{a^2 + \frac{1}{x^2}} + \frac{i}{x} \right) \right)}{3 \left(ia + \sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)^(5/2),x, algorithm="giac")

[Out]
$$\frac{-2/3*(2*a^2 - 3*(\sqrt{a^2 + 1/x^2}) - 1/x)^2 + 3*a*(-I*\sqrt{a^2 + 1/x^2}) + I/x}{(I*a + \sqrt{a^2 + 1/x^2}) - 1/x}^3$$

Mupad [B]

time = 0.50, size = 33, normalized size = 0.49

$$\frac{\sqrt{a^2 x^2 + 1} (-2 + a x i) i}{3 a (-1 + a x i)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*i + 1)^2/(a^2*x^2 + 1)^(5/2),x)

[Out]
$$((a^2*x^2 + 1)^{(1/2)}*(a*x*i - 2)*i)/(3*a*(a*x*i - 1)^2)$$

$$3.323 \quad \int \frac{e^{i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=28

$$\frac{1}{2a(i+ax)} + \frac{\operatorname{ArcTan}(ax)}{2a}$$

[Out] 1/2/a/(I+a*x)+1/2*arctan(a*x)/a

Rubi [A]

time = 0.03, antiderivative size = 28, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5181, 46, 209}

$$\frac{\operatorname{ArcTan}(ax)}{2a} + \frac{1}{2a(ax+i)}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])/(1 + a^2*x^2)^(3/2),x]

[Out] 1/(2*a*(I + a*x)) + ArcTan[a*x]/(2*a)

Rule 46

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] :> Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1}{(1-iax)^2(1+iax)} dx \\
&= \int \left(-\frac{1}{2(i+ax)^2} + \frac{1}{2(1+a^2x^2)} \right) dx \\
&= \frac{1}{2a(i+ax)} + \frac{1}{2} \int \frac{1}{1+a^2x^2} dx \\
&= \frac{1}{2a(i+ax)} + \frac{\tan^{-1}(ax)}{2a}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 21, normalized size = 0.75

$$\frac{\frac{1}{i+ax} + \text{ArcTan}(ax)}{2a}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(I*ArcTan[a*x])/(1 + a^2*x^2)^(3/2), x]``[Out] ((I + a*x)^(-1) + ArcTan[a*x])/(2*a)`**Maple [A]**

time = 0.08, size = 38, normalized size = 1.36

method	result	size
risch	$\frac{1}{2a(ax+i)} + \frac{\arctan(ax)}{2a}$	24
default	$\frac{2a^2x-2ia}{4a^2(a^2x^2+1)} + \frac{\arctan(ax)}{2a}$	38
meijerg	$\frac{\frac{2x\sqrt{a^2}}{2a^2x^2+2} + \frac{\sqrt{a^2}}{a} \arctan(ax)}{2\sqrt{a^2}} + \frac{iax^2}{2a^2x^2+2}$	61

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)/(a^2*x^2+1)^2,x,method=_RETURNVERBOSE)``[Out] 1/4*(2*a^2*x-2*I*a)/a^2/(a^2*x^2+1)+1/2*arctan(a*x)/a`**Maxima [A]**

time = 0.45, size = 28, normalized size = 1.00

$$\frac{ax - i}{2(a^3x^2 + a)} + \frac{\arctan(ax)}{2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^2,x, algorithm="maxima")

[Out] 1/2*(a*x - I)/(a^3*x^2 + a) + 1/2*arctan(a*x)/a

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(22) = 44$.

time = 2.50, size = 49, normalized size = 1.75

$$\frac{(i a x - 1) \log\left(\frac{a x + i}{a}\right) + (-i a x + 1) \log\left(\frac{a x - i}{a}\right) + 2}{4(a^2 x + i a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^2,x, algorithm="fricas")

[Out] 1/4*((I*a*x - 1)*log((a*x + I)/a) + (-I*a*x + 1)*log((a*x - I)/a) + 2)/(a^2*x + I*a)

Sympy [A]

time = 0.11, size = 32, normalized size = 1.14

$$i \left(-\frac{i}{2a^2x + 2ia} + \frac{-\frac{\log(x - \frac{i}{a})}{4} + \frac{\log(x + \frac{i}{a})}{4}}{a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a**2*x**2+1)**2,x)

[Out] I*(-I/(2*a**2*x + 2*I*a) + (-log(x - I/a)/4 + log(x + I/a)/4)/a)

Giac [A]

time = 0.42, size = 36, normalized size = 1.29

$$\frac{i \log(ax + i)}{4a} - \frac{i \log(-i a x - 1)}{4a} + \frac{1}{2(ax + i)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)/(a^2*x^2+1)^2,x, algorithm="giac")

[Out] 1/4*I*log(a*x + I)/a - 1/4*I*log(-I*a*x - 1)/a + 1/2/((a*x + I)*a)

Mupad [B]

time = 0.48, size = 23, normalized size = 0.82

$$\frac{1}{2(x a^2 + a i)} + \frac{\operatorname{atan}(a x)}{2 a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*I + 1)/(a^2*x^2 + 1)^2,x)

[Out] 1/(2*(a*I + a^2*x)) + atan(a*x)/(2*a)

$$3.324 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=29

$$-\frac{1}{2a(i-ax)} + \frac{\operatorname{ArcTan}(ax)}{2a}$$

[Out] -1/2/a/(I-a*x)+1/2*arctan(a*x)/a

Rubi [A]

time = 0.03, antiderivative size = 29, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5181, 46, 209}

$$\frac{\operatorname{ArcTan}(ax)}{2a} - \frac{1}{2a(-ax+i)}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a*x])*(1+a^2*x^2)^(3/2)),x]

[Out] -1/2*1/(a*(I-a*x))+ArcTan[a*x]/(2*a)

Rule 46

Int[((a_) + (b_)*(x_))^(m_)*((c_) + (d_)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5181

Int[E^(ArcTan[(a_)*(x_)])*(n_)*((c_) + (d_)*(x_)^2)^(p_), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1}{(1-iax)(1+iax)^2} dx \\
&= \int \left(-\frac{1}{2(-i+ax)^2} + \frac{1}{2(1+a^2x^2)} \right) dx \\
&= -\frac{1}{2a(i-ax)} + \frac{1}{2} \int \frac{1}{1+a^2x^2} dx \\
&= -\frac{1}{2a(i-ax)} + \frac{\tan^{-1}(ax)}{2a}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 21, normalized size = 0.72

$$\frac{\frac{1}{-i+ax} + \text{ArcTan}(ax)}{2a}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^(I*ArcTan[a*x])*(1 + a^2*x^2)^(3/2)), x]``[Out] ((-I + a*x)^(-1) + ArcTan[a*x])/(2*a)`**Maple [A]**

time = 0.08, size = 43, normalized size = 1.48

method	result	size
risch	$\frac{1}{2a(ax-i)} + \frac{\arctan(ax)}{2a}$	24
default	$-\frac{i \ln(-ax+i)}{4a} - \frac{1}{2a(-ax+i)} + \frac{i \ln(ax+i)}{4a}$	43

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)/(a^2*x^2+1), x, method=_RETURNVERBOSE)``[Out] -1/4*I/a*ln(I-a*x)-1/2/a/(I-a*x)+1/4*I/a*ln(I+a*x)`**Maxima [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)/(a^2*x^2+1), x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: Error executing code in Maxima:
 expt: undefined: 0 to a negative exponent.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 49 vs. $2(22) = 44$.
 time = 3.16, size = 49, normalized size = 1.69

$$\frac{(i a x + 1) \log\left(\frac{a x + i}{a}\right) + (-i a x - 1) \log\left(\frac{a x - i}{a}\right) + 2}{4(a^2 x - i a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)/(a^2*x^2+1),x, algorithm="fricas")

[Out] 1/4*((I*a*x + 1)*log((a*x + I)/a) + (-I*a*x - 1)*log((a*x - I)/a) + 2)/(a^2*x - I*a)

Sympy [A]

time = 0.11, size = 34, normalized size = 1.17

$$-i \left(\frac{i}{2a^2x - 2ia} + \frac{\frac{\log(x - \frac{i}{a})}{4} - \frac{\log(x + \frac{i}{a})}{4}}{a} \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)/(a**2*x**2+1),x)

[Out] -I*(I/(2*a**2*x - 2*I*a) + (log(x - I/a)/4 - log(x + I/a)/4)/a)

Giac [A]

time = 0.41, size = 36, normalized size = 1.24

$$-\frac{i \log(ax - i)}{4a} + \frac{i \log(iax - 1)}{4a} + \frac{1}{2(ax - i)a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)/(a^2*x^2+1),x, algorithm="giac")

[Out] -1/4*I*log(a*x - I)/a + 1/4*I*log(I*a*x - 1)/a + 1/2/((a*x - I)*a)

Mupad [B]

time = 0.48, size = 25, normalized size = 0.86

$$\frac{\operatorname{atan}(ax)}{2a} - \frac{1}{2(-a^2x + a1i)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/((a^2*x^2 + 1)*(a*x*1i + 1)),x)

[Out] atan(a*x)/(2*a) - 1/(2*(a*1i - a^2*x))

$$3.325 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=67

$$\frac{i\sqrt{1-iax}}{3a(1+iax)^{3/2}} + \frac{i\sqrt{1-iax}}{3a\sqrt{1+iax}}$$

[Out] 1/3*I*(1-I*a*x)^(1/2)/a/(1+I*a*x)^(3/2)+1/3*I*(1-I*a*x)^(1/2)/a/(1+I*a*x)^(1/2)

Rubi [A]

time = 0.03, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5181, 47, 37}

$$\frac{i\sqrt{1-iax}}{3a\sqrt{1+iax}} + \frac{i\sqrt{1-iax}}{3a(1+iax)^{3/2}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((2*I)*ArcTan[a*x])*(1 + a^2*x^2)^(3/2)),x]

[Out] ((I/3)*Sqrt[1 - I*a*x])/(a*(1 + I*a*x)^(3/2)) + ((I/3)*Sqrt[1 - I*a*x])/(a*Sqrt[1 + I*a*x])

Rule 37

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp
[(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] /; FreeQ[{
a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && EqQ[m + n + 2, 0] && NeQ[m, -
1]
```

Rule 47

```
Int[((a_.) + (b_.)*(x_))^(m_.)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Simp[
(a + b*x)^(m + 1)*((c + d*x)^(n + 1)/((b*c - a*d)*(m + 1))), x] - Dist[d*(S
implify[m + n + 2]/((b*c - a*d)*(m + 1))), Int[(a + b*x)^Simplify[m + 1]*(c
+ d*x)^n, x], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && I
LtQ[Simplify[m + n + 2], 0] && NeQ[m, -1] && !(LtQ[m, -1] && LtQ[n, -1] &&
(EqQ[a, 0] || (NeQ[c, 0] && LtQ[m - n, 0] && IntegerQ[n]))) && (SumSimpler
Q[m, 1] || !SumSimplerQ[n, 1])
```

Rule 5181

```
Int[E^(ArcTan[(a_.)*(x_)^(n_.)]*(c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :=
Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /
; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-2i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1}{\sqrt{1-iax}(1+iax)^{5/2}} dx \\
&= \frac{i\sqrt{1-iax}}{3a(1+iax)^{3/2}} + \frac{1}{3} \int \frac{1}{\sqrt{1-iax}(1+iax)^{3/2}} dx \\
&= \frac{i\sqrt{1-iax}}{3a(1+iax)^{3/2}} + \frac{i\sqrt{1-iax}}{3a\sqrt{1+iax}}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 48, normalized size = 0.72

$$\frac{\sqrt{1-iax}(2+iax)}{3a\sqrt{1+iax}(-i+ax)}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((2*I)*ArcTan[a*x]))*(1+a^2*x^2)^(3/2),x]``[Out] (Sqrt[1-I*a*x]*(2+I*a*x))/(3*a*Sqrt[1+I*a*x]*(-I+a*x))`**Maple [A]**

time = 0.06, size = 93, normalized size = 1.39

method	result	size
default	$-\frac{i\sqrt{\left(x-\frac{i}{a}\right)^2 a^2+2ia\left(x-\frac{i}{a}\right)}}{3a\left(x-\frac{i}{a}\right)^2} - \frac{\sqrt{\left(x-\frac{i}{a}\right)^2 a^2+2ia\left(x-\frac{i}{a}\right)}}{3\left(x-\frac{i}{a}\right)}$	93

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^2/(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)``[Out] -1/a^2*(1/3*I/a/(x-I/a)^2*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2)-1/3/(x-I/a)*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(1/2))`**Maxima [A]**

time = 0.47, size = 58, normalized size = 0.87

$$-\frac{i\sqrt{a^2x^2+1}}{3(a^3x^2-2ia^2x-a)} + \frac{i\sqrt{a^2x^2+1}}{3ia^2x+3a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2/(a^2*x^2+1)^(1/2),x, algorithm="maxima")

[Out] $-\frac{1}{3}I\sqrt{a^2x^2 + 1}/(a^3x^2 - 2Ia^2x - a) + I\sqrt{a^2x^2 + 1}/(3Ia^2x + 3a)$

Fricas [A]

time = 2.66, size = 51, normalized size = 0.76

$$\frac{a^2x^2 - 2iax + \sqrt{a^2x^2 + 1}(ax - 2i) - 1}{3(a^3x^2 - 2ia^2x - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2/(a^2*x^2+1)^(1/2),x, algorithm="fricas")

[Out] $\frac{1}{3}(a^2x^2 - 2Ia^2x + \sqrt{a^2x^2 + 1})(ax - 2I) - 1)/(a^3x^2 - 2Ia^2x - a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{1}{a^2x^2\sqrt{a^2x^2 + 1} - 2iax\sqrt{a^2x^2 + 1} - \sqrt{a^2x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**2/(a**2*x**2+1)**(1/2),x)

[Out] $-\text{Integral}(1/(a**2*x**2*\sqrt{a**2*x**2 + 1} - 2I*a*x*\sqrt{a**2*x**2 + 1} - \sqrt{a**2*x**2 + 1}), x)$

Giac [A]

time = 0.43, size = 67, normalized size = 1.00

$$\frac{2 \left(2a^2 - 3 \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^2 + 3a \left(i \sqrt{a^2 + \frac{1}{x^2}} - \frac{i}{x} \right) \right)}{3 \left(-ia + \sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2/(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] $-2/3*(2*a^2 - 3*(\sqrt{a^2 + 1/x^2} - 1/x)^2 + 3*a*(I*\sqrt{a^2 + 1/x^2} - I/x))/(-I*a + \sqrt{a^2 + 1/x^2} - 1/x)^3$

Mupad [B]

time = 0.06, size = 31, normalized size = 0.46

$$-\frac{\sqrt{a^2x^2 + 1}(ax - 2i)}{3a(1 + axi)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(1/((a^2*x^2 + 1)^(1/2)*(a*x*1i + 1)^2),x)
```

```
[Out] -((a^2*x^2 + 1)^(1/2)*(a*x - 2i))/(3*a*(a*x*1i + 1)^2)
```

$$3.326 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=19

$$\frac{i}{2a(1+iax)^2}$$

[Out] 1/2*I/a/(1+I*a*x)^2

Rubi [A]

time = 0.02, antiderivative size = 19, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5181, 32}

$$\frac{i}{2a(1+iax)^2}$$

Antiderivative was successfully verified.

[In] Int[1/(E^((3*I)*ArcTan[a*x])*(1 + a^2*x^2)^(3/2)),x]

[Out] (I/2)/(a*(1 + I*a*x)^2)

Rule 32

Int[((a_.) + (b_.)*(x_))^(m_), x_Symbol] := Simp[(a + b*x)^(m + 1)/(b*(m + 1)), x] /; FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-3i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{1}{(1+iax)^3} dx \\ &= \frac{i}{2a(1+iax)^2} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 18, normalized size = 0.95

$$-\frac{i}{2a(-i+ax)^2}$$

Antiderivative was successfully verified.

[In] Integrate[1/(E^((3*I)*ArcTan[a*x]))*(1 + a^2*x^2)^(3/2),x]

[Out] (-1/2*I)/(a*(-I + a*x)^2)

Maple [A]

time = 0.07, size = 16, normalized size = 0.84

method	result	size
risch	$-\frac{i}{2a(ax-i)^2}$	15
default	$\frac{i}{2a(iax+1)^2}$	16
meijerg	$\frac{x(iax+2)}{2(iax+1)^2}$	20
gosper	$\frac{-ax+i}{2a(iax+1)^3}$	22
norman	$\frac{x-\frac{3}{2}iax^2-\frac{1}{2}ia^3x^4}{(a^2x^2+1)^2}$	31

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^3,x,method=_RETURNVERBOSE)

[Out] 1/2*I/a/(1+I*a*x)^2

Maxima [A]

time = 0.25, size = 13, normalized size = 0.68

$$\frac{i}{2(iax+1)^2a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3,x, algorithm="maxima")

[Out] 1/2*I/((I*a*x + 1)^2*a)

Fricas [A]

time = 2.69, size = 21, normalized size = 1.11

$$-\frac{i}{2(a^3x^2 - 2ia^2x - a)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3,x, algorithm="fricas")

[Out] -1/2*I/(a^3*x^2 - 2*I*a^2*x - a)

Sympy [A]

time = 0.09, size = 22, normalized size = 1.16

$$\frac{i}{2a^3x^2 - 4ia^2x - 2a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)**3,x)``[Out] -I/(2*a**3*x**2 - 4*I*a**2*x - 2*a)`**Giac [A]**

time = 0.42, size = 13, normalized size = 0.68

$$\frac{i}{2(iax + 1)^2a}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(1/(1+I*a*x)^3,x, algorithm="giac")``[Out] 1/2*I/((I*a*x + 1)^2*a)`**Mupad [B]**

time = 0.05, size = 24, normalized size = 1.26

$$\frac{1i}{2(-a^3x^2 + a^2x2i + a)}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(a*x*1i + 1)^3,x)``[Out] 1i/(2*(a + a^2*x*2i - a^3*x^2))`

$$3.327 \quad \int \frac{e^{-4i \operatorname{ArcTan}(ax)}}{(1+a^2x^2)^{3/2}} dx$$

Optimal. Leaf size=67

$$\frac{i(1-iax)^{3/2}}{5a(1+iax)^{5/2}} + \frac{i(1-iax)^{3/2}}{15a(1+iax)^{3/2}}$$

[Out] $1/5*I*(1-I*a*x)^{(3/2)}/a/(1+I*a*x)^{(5/2)}+1/15*I*(1-I*a*x)^{(3/2)}/a/(1+I*a*x)^{(3/2)}$

Rubi [A]

time = 0.03, antiderivative size = 67, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5181, 47, 37}

$$\frac{i(1-iax)^{3/2}}{15a(1+iax)^{3/2}} + \frac{i(1-iax)^{3/2}}{5a(1+iax)^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((4*I)*\text{ArcTan}[a*x])*(1+a^2*x^2)^{(3/2))}, x]$

[Out] $((I/5)*(1-I*a*x)^{(3/2)})/(a*(1+I*a*x)^{(5/2)}) + ((I/15)*(1-I*a*x)^{(3/2)})/(a*(1+I*a*x)^{(3/2)})$

Rule 37

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1)})}/((b*c - a*d)*(m + 1)), x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{EqQ}[m + n + 2, 0] \&\& \text{NeQ}[m, -1]$

Rule 47

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] :> \text{Simp}[(a + b*x)^{(m + 1)*((c + d*x)^{(n + 1)})}/((b*c - a*d)*(m + 1)), x] - \text{Dist}[d*(\text{Simplify}[m + n + 2]/((b*c - a*d)*(m + 1))), \text{Int}[(a + b*x)^{\text{Simplify}[m + 1]}*(c + d*x)^n, x], x] /; \text{FreeQ}\{a, b, c, d, m, n\}, x\} \&\& \text{NeQ}[b*c - a*d, 0] \&\& \text{IntegerQ}[m + n + 2] \&\& \text{IntegerQ}[m] \&\& \text{IntegerQ}[n] \&\& (\text{EqQ}[a, 0] || (\text{NeQ}[c, 0] \&\& \text{LtQ}[m - n, 0] \&\& \text{IntegerQ}[n])) \&\& (\text{SumSimplerQ}[m, 1] || !\text{SumSimplerQ}[n, 1])$

Rule 5181

$\text{Int}[E^{\text{ArcTan}[(a_.)*(x_.)]*(n_.)*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] :> \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, n, p\}, x\} \&\& \text{EqQ}[d, a^2*c] \&\& (\text{IntegerQ}[p] || \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned}
\int \frac{e^{-4i \tan^{-1}(ax)}}{(1+a^2x^2)^{3/2}} dx &= \int \frac{\sqrt{1-iax}}{(1+iax)^{7/2}} dx \\
&= \frac{i(1-iax)^{3/2}}{5a(1+iax)^{5/2}} + \frac{1}{5} \int \frac{\sqrt{1-iax}}{(1+iax)^{5/2}} dx \\
&= \frac{i(1-iax)^{3/2}}{5a(1+iax)^{5/2}} + \frac{i(1-iax)^{3/2}}{15a(1+iax)^{3/2}}
\end{aligned}$$

Mathematica [A]

time = 0.01, size = 47, normalized size = 0.70

$$\frac{(1-iax)^{3/2}(-4i+ax)}{15a\sqrt{1+iax}(-i+ax)^2}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((4*I)*ArcTan[a*x]))*(1 + a^2*x^2)^(3/2)),x]``[Out] ((1 - I*a*x)^(3/2)*(-4*I + a*x))/(15*a*Sqrt[1 + I*a*x]*(-I + a*x)^2)`**Maple [A]**

time = 0.08, size = 92, normalized size = 1.37

method	result	size
default	$\frac{i \left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{3}{2}}}{5a \left(x - \frac{i}{a} \right)^4} - \frac{\left(\left(x - \frac{i}{a} \right)^2 a^2 + 2ia \left(x - \frac{i}{a} \right) \right)^{\frac{3}{2}}}{15 \left(x - \frac{i}{a} \right)^3}$	92

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^4*(a^2*x^2+1)^(1/2),x,method=_RETURNVERBOSE)``[Out] 1/a^4*(1/5*I/a/(x-I/a)^4*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2)-1/15/(x-I/a)^3*((x-I/a)^2*a^2+2*I*a*(x-I/a))^(3/2))`**Maxima [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 99 vs. 2(43) = 86.

time = 0.26, size = 99, normalized size = 1.48

$$\frac{2i \sqrt{a^2x^2 + 1}}{-5i a^4x^3 - 15 a^3x^2 + 15i a^2x + 5a} + \frac{i \sqrt{a^2x^2 + 1}}{15(a^3x^2 - 2i a^2x - a)} - \frac{i \sqrt{a^2x^2 + 1}}{15i a^2x + 15a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^(1/2),x, algorithm="maxima")

[Out] $2*I*\sqrt{a^2*x^2 + 1}/(-5*I*a^4*x^3 - 15*a^3*x^2 + 15*I*a^2*x + 5*a) + 1/15 * I*\sqrt{a^2*x^2 + 1}/(a^3*x^2 - 2*I*a^2*x - a) - I*\sqrt{a^2*x^2 + 1}/(15*I*a^2*x + 15*a)$

Fricas [A]

time = 7.37, size = 75, normalized size = 1.12

$$-\frac{a^3x^3 - 3ia^2x^2 - 3ax + (a^2x^2 - 3iax + 4)\sqrt{a^2x^2 + 1} + i}{15(a^4x^3 - 3ia^3x^2 - 3a^2x + ia)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^(1/2),x, algorithm="fricas")

[Out] $-1/15*(a^3*x^3 - 3*I*a^2*x^2 - 3*a*x + (a^2*x^2 - 3*I*a*x + 4)*\sqrt{a^2*x^2 + 1} + I)/(a^4*x^3 - 3*I*a^3*x^2 - 3*a^2*x + I*a)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{\sqrt{a^2x^2 + 1}}{(ax - i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**4*(a**2*x**2+1)**(1/2),x)

[Out] Integral(sqrt(a**2*x**2 + 1)/(a*x - I)**4, x)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 111 vs. $2(43) = 86$.

time = 0.45, size = 111, normalized size = 1.66

$$\frac{2 \left(4a^4 - 25a^2 \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^2 - 15ia \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^3 + 15 \left(\sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^4 - 5a^3 \left(-i \sqrt{a^2 + \frac{1}{x^2}} + \frac{i}{x} \right) \right)}{15 \left(-ia + \sqrt{a^2 + \frac{1}{x^2}} - \frac{1}{x} \right)^5}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^(1/2),x, algorithm="giac")

[Out] $2/15*(4*a^4 - 25*a^2*(\sqrt{a^2 + 1/x^2} - 1/x)^2 - 15*I*a*(\sqrt{a^2 + 1/x^2} - 1/x)^3 + 15*(\sqrt{a^2 + 1/x^2} - 1/x)^4 - 5*a^3*(-I*\sqrt{a^2 + 1/x^2} + I/x))/(-I*a + \sqrt{a^2 + 1/x^2} - 1/x)^5$

Mupad [B]

time = 0.52, size = 40, normalized size = 0.60

$$\frac{\sqrt{a^2 x^2 + 1} (a^2 x^2 - a x 3i + 4) 1i}{15 a (1 + a x 1i)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(1/2)/(a*x*1i + 1)^4,x)**[Out]** ((a^2*x^2 + 1)^(1/2)*(a^2*x^2 - a*x*3i + 4)*1i)/(15*a*(a*x*1i + 1)^3)

$$3.328 \quad \int \frac{e^{5i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=95

$$-\frac{2\sqrt{1+a^2x^2}}{3ac(i+ax)^3\sqrt{c+a^2cx^2}} - \frac{i\sqrt{1+a^2x^2}}{2ac(i+ax)^2\sqrt{c+a^2cx^2}}$$

[Out] $-2/3*(a^2*x^2+1)^{(1/2)}/a/c/(I+a*x)^3/(a^2*c*x^2+c)^{(1/2)}-1/2*I*(a^2*x^2+1)^{(1/2)}/a/c/(I+a*x)^2/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 95, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 45}

$$-\frac{i\sqrt{a^2x^2+1}}{2ac(ax+i)^2\sqrt{a^2cx^2+c}} - \frac{2\sqrt{a^2x^2+1}}{3ac(ax+i)^3\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((5*I)*\text{ArcTan}[a*x])}/(c + a^2*c*x^2)^{(3/2)}, x]$

[Out] $(-2*\text{Sqrt}[1 + a^2*x^2])/(3*a*c*(I + a*x)^3*\text{Sqrt}[c + a^2*c*x^2]) - ((I/2)*\text{Sqrt}[1 + a^2*x^2])/(a*c*(I + a*x)^2*\text{Sqrt}[c + a^2*c*x^2])$

Rule 45

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)}, x_Symbol] := \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n, x], x] /;$ FreeQ[{a, b, c, d, n}, x] && NeQ[b*c - a*d, 0] && IGtQ[m, 0] && (!IntegerQ[n] || (EqQ[c, 0] && LeQ[7*m + 4*n + 4, 0]) || LtQ[9*m + 5*(n + 1), 0] || GtQ[m + n + 2, 0])

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] := \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] := \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{5i \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{5i \tan^{-1}(ax)}}{(1 + a^2 x^2)^{3/2}} dx}{c\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \frac{1 + iax}{(1 - iax)^4} dx}{c\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \left(\frac{2}{(i+ax)^4} + \frac{i}{(i+ax)^3} \right) dx}{c\sqrt{c + a^2 cx^2}} \\
&= -\frac{2\sqrt{1 + a^2 x^2}}{3ac(i + ax)^3 \sqrt{c + a^2 cx^2}} - \frac{i\sqrt{1 + a^2 x^2}}{2ac(i + ax)^2 \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.02, size = 56, normalized size = 0.59

$$-\frac{i(-i + 3ax)\sqrt{1 + a^2 x^2}}{6ac(i + ax)^3 \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((5*I)*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]``[Out] ((-1/6*I)*(-I + 3*a*x)*Sqrt[1 + a^2*x^2])/(a*c*(I + a*x)^3*Sqrt[c + a^2*c*x^2])`**Maple [A]**

time = 0.09, size = 48, normalized size = 0.51

method	result	size
risch	$\frac{\sqrt{a^2 x^2 + 1} \left(-\frac{ix}{2} - \frac{1}{6a}\right)}{c\sqrt{c(a^2 x^2 + 1)}(ax+i)^3}$	47
default	$-\frac{\sqrt{c(a^2 x^2 + 1)}(3iax+1)}{6\sqrt{a^2 x^2 + 1}c^2 a(ax+i)^3}$	48

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(3/2), x, method=_RETURNVERBOSE)``[Out] -1/6/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)*(3*I*a*x+1)/c^2/a/(I+a*x)^3`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((I*a*x + 1)^5/((a^2*c*x^2 + c)^(3/2)*(a^2*x^2 + 1)^(5/2)), x)
```

Fricas [A]

time = 2.98, size = 101, normalized size = 1.06

$$\frac{\sqrt{a^2cx^2 + c} (i a^2x^3 - 3ax^2 - 6ix)\sqrt{a^2x^2 + 1}}{6(a^5c^2x^5 + 3i a^4c^2x^4 - 2a^3c^2x^3 + 2i a^2c^2x^2 - 3ac^2x - ic^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")
```

```
[Out] 1/6*sqrt(a^2*c*x^2 + c)*(I*a^2*x^3 - 3*a*x^2 - 6*I*x)*sqrt(a^2*x^2 + 1)/(a^5*c^2*x^5 + 3*I*a^4*c^2*x^4 - 2*a^3*c^2*x^3 + 2*I*a^2*c^2*x^2 - 3*a*c^2*x - I*c^2)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)**5/(a**2*x**2+1)**(5/2)/(a**2*c*x**2+c)**(3/2),x)
```

```
[Out] I*(Integral(-I/(a**6*c*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(5*a*x/(a**6*c*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-10*a**3*x**3/(a**6*c*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a**5*x**5/(a**6*c*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(10*I*a**2*x**2/(a**6*c*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 3*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-5*I*a**4*x**4/(a**6*c*x**6*sqrt(a**2*x**2 + 1)*sq
```

```
rt(a**2*c*x**2 + c) + 3*a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 +
c) + 3*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*
x**2 + 1)*sqrt(a**2*c*x**2 + c)), x)
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="g
iac")
```

```
[Out] integrate((I*a*x + 1)^5/((a^2*c*x^2 + c)^(3/2)*(a^2*x^2 + 1)^(5/2)), x)
```

Mupad [B]

time = 1.62, size = 48, normalized size = 0.51

$$-\frac{\sqrt{c(a^2x^2+1)}(3ax-i)}{6ac^2\sqrt{a^2x^2+1}(-1+axi)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*x*I + 1)^5/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^(5/2)),x)
```

```
[Out] -((c*(a^2*x^2 + 1))^(1/2)*(3*a*x - I))/(6*a*c^2*(a^2*x^2 + 1)^(1/2)*(a*x*I
I - 1)^3)
```


$$3.329 \quad \int \frac{e^{4i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=69

$$-\frac{ic(1+iax)^4}{3a(c+a^2cx^2)^{5/2}} + \frac{ic(1+iax)^5}{15a(c+a^2cx^2)^{5/2}}$$

[Out] $-1/3*I*c*(1+I*a*x)^4/a/(a^2*c*x^2+c)^{(5/2)}+1/15*I*c*(1+I*a*x)^5/a/(a^2*c*x^2+c)^{(5/2)}$

Rubi [A]

time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$,

Rules used = {5183, 673, 665}

$$\frac{ic(1+iax)^5}{15a(a^2cx^2+c)^{5/2}} - \frac{ic(1+iax)^4}{3a(a^2cx^2+c)^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((4*I)*\text{ArcTan}[a*x])}/(c + a^2*c*x^2)^{(3/2)}, x]$

[Out] $((-1/3*I)*c*(1 + I*a*x)^4)/(a*(c + a^2*c*x^2)^{(5/2)}) + ((I/15)*c*(1 + I*a*x)^5)/(a*(c + a^2*c*x^2)^{(5/2)})$

Rule 665

$\text{Int}[(d + (e_*)*(x_))^{(m_)*((a_*) + (c_*)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*(d + e*x)^m*((a + c*x^2)^{(p + 1)}/(2*c*d*(p + 1))), x] /; \text{FreeQ}\{a, c, d, e, m, p\}, x] \ \&\& \ \text{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{EqQ}[m + 2*p + 2, 0]$

Rule 673

$\text{Int}[(d + (e_*)*(x_))^{(m_)*((a_*) + (c_*)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(-e)*(d + e*x)^m*((a + c*x^2)^{(p + 1)}/(2*c*d*(m + p + 1))), x] + \text{Dist}[\text{Simplify}[m + 2*p + 2]/(2*d*(m + p + 1)), \text{Int}[(d + e*x)^{(m + 1)}*(a + c*x^2)^p, x], x] /; \text{FreeQ}\{a, c, d, e, m, p\}, x] \ \&\& \ \text{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{ILtQ}[\text{Simplify}[m + 2*p + 2], 0]$

Rule 5183

$\text{Int}[E^{\text{ArcTan}[(a_*)*(x_)]*(n_)*((c_*) + (d_*)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[1/c^{(I*(n/2))}, \text{Int}[(c + d*x^2)^{(p + I*(n/2))}/(1 + I*a*x)^{(I*n)}, x], x] /; \text{FreeQ}\{a, c, d, p\}, x] \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0]) \ \&\& \ \text{ILtQ}[I*(n/2), 0]$

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)}}{(c + a^2cx^2)^{3/2}} dx &= c^2 \int \frac{(1 + iax)^4}{(c + a^2cx^2)^{7/2}} dx \\ &= -\frac{ic(1 + iax)^4}{3a(c + a^2cx^2)^{5/2}} - \frac{1}{3}c^2 \int \frac{(1 + iax)^5}{(c + a^2cx^2)^{7/2}} dx \\ &= -\frac{ic(1 + iax)^4}{3a(c + a^2cx^2)^{5/2}} + \frac{ic(1 + iax)^5}{15a(c + a^2cx^2)^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 77, normalized size = 1.12

$$\frac{(1 + iax)^{3/2}(4i + ax)\sqrt{1 + a^2x^2}}{15ac\sqrt{1 - iax}(i + ax)^2\sqrt{c + a^2cx^2}}$$

Warning: Unable to verify antiderivative.

`[In] Integrate[E^((4*I)*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]``[Out] ((1 + I*a*x)^(3/2)*(4*I + a*x)*Sqrt[1 + a^2*x^2])/(15*a*c*Sqrt[1 - I*a*x]*(I + a*x)^2*Sqrt[c + a^2*c*x^2])`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 939 vs. 2(57) = 114.

time = 0.09, size = 940, normalized size = 13.62

method	result
trager	$\frac{(-a^5x^5 - 10a^3x^3 + 20ia^2x^2 + 15ax - 4i)\sqrt{a^2cx^2 + c}}{15c^2(a^2x^2 + 1)^3a}$
default	$\frac{x}{c\sqrt{a^2cx^2 + c}} - \frac{2\left(i\sqrt{-a^2} - a\right) \left(\frac{1}{3c\sqrt{-a^2} \left(x + \frac{\sqrt{-a^2}}{a^2}\right)} \sqrt{\left(x + \frac{\sqrt{-a^2}}{a^2}\right)^2 a^2c - 2c\sqrt{-a^2}} \left(x + \frac{\sqrt{-a^2}}{a}\right) \right)}{a\sqrt{-a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)
[Out] x/c/(a^2*c*x^2+c)^(1/2)-2/a*(I*(-a^2)^(1/2)-a)/(-a^2)^(1/2)*(1/3/c/(-a^2)^(1/2)/(x+(-a^2)^(1/2)/a^2)/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)+1/3/c^2/(-a^2)^(1/2)*(2*a^2*c*(x+(-a^2)^(1/2)/a^2)-2*c*(-a^2)^(1/2))/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2))-2/a^3*(I*(-a^2)^(1/2)+a)*(-1/5/c/(-a^2)^(1/2)/(x-(-a^2)^(1/2)/a^2)^2/((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2)-3/5*a^2/(-a^2)^(1/2)*(-1/3/c/(-a^2)^(1/2)/(x-(-a^2)^(1/2)/a^2)/((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2))-1/3/c^2/(-a^2)^(1/2)*(2*a^2*c*(x-(-a^2)^(1/2)/a^2)+2*c*(-a^2)^(1/2))/((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2))-2/a*(I*(-a^2)^(1/2)+a)/(-a^2)^(1/2)*(-1/3/c/(-a^2)^(1/2)/(x-(-a^2)^(1/2)/a^2)/((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2))-1/3/c^2/(-a^2)^(1/2)*(2*a^2*c*(x-(-a^2)^(1/2)/a^2)+2*c*(-a^2)^(1/2))/((x-(-a^2)^(1/2)/a^2)^2*a^2*c+2*c*(-a^2)^(1/2)*(x-(-a^2)^(1/2)/a^2))^(1/2))+2/a^3*(I*(-a^2)^(1/2)-a)*(1/5/c/(-a^2)^(1/2)/(x+(-a^2)^(1/2)/a^2)^2/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)+3/5*a^2/(-a^2)^(1/2)*(1/3/c/(-a^2)^(1/2)/(x+(-a^2)^(1/2)/a^2)/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)+1/3/c^2/(-a^2)^(1/2)*(2*a^2*c*(x+(-a^2)^(1/2)/a^2)-2*c*(-a^2)^(1/2))/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)))
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")
```

```
[Out] integrate((I*a*x + 1)^4/((a^2*c*x^2 + c)^(3/2)*(a^2*x^2 + 1)^2), x)
```

Fricas [A]

time = 3.43, size = 66, normalized size = 0.96

$$\frac{\sqrt{a^2cx^2 + c} (a^2x^2 + 3iax + 4)}{15(a^4c^2x^3 + 3ia^3c^2x^2 - 3a^2c^2x - iac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")
```

```
[Out] -1/15*sqrt(a^2*c*x^2 + c)*(a^2*x^2 + 3*I*a*x + 4)/(a^4*c^2*x^3 + 3*I*a^3*c^2*x^2 - 3*a^2*c^2*x - I*a*c^2)
```

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(ax - i)^4}{(c(a^2x^2 + 1))^{\frac{3}{2}}(a^2x^2 + 1)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2/(a**2*c*x**2+c)**(3/2),x)**[Out]** Integral((a*x - I)**4/((c*(a**2*x**2 + 1))**(3/2)*(a**2*x**2 + 1)**2), x)**Giac [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(53) = 106.

time = 0.44, size = 134, normalized size = 1.94

$$\frac{2 \left(15 \left(\sqrt{a^2c} x - \sqrt{a^2cx^2 + c} \right)^3 \sqrt{c} - 5i \left(\sqrt{a^2c} x - \sqrt{a^2cx^2 + c} \right)^2 c - 5 \left(\sqrt{a^2c} x - \sqrt{a^2cx^2 + c} \right) c^{\frac{3}{2}} - i c^2 \right)}{15 \left(\sqrt{a^2c} x - \sqrt{a^2cx^2 + c} + i \sqrt{c} \right)^5 ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")**[Out]** 2/15*(15*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c))^3*sqrt(c) - 5*I*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c))^2*c - 5*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c))*c^(3/2) - I*c^2)/((sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c) + I*sqrt(c))^5*a*c)**Mupad [B]**

time = 0.98, size = 46, normalized size = 0.67

$$\frac{\sqrt{c(a^2x^2 + 1)}(a^2x^2 li - 3ax + 4i)}{15ac^2(-1 + ax li)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^4/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^2),x)**[Out]** ((c*(a^2*x^2 + 1))^(1/2)*(a^2*x^2*1i - 3*a*x + 4i))/(15*a*c^2*(a*x*1i - 1)^3)

$$3.330 \quad \int \frac{e^{3i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=49

$$-\frac{i\sqrt{1+a^2x^2}}{2ac(1-iax)^2\sqrt{c+a^2cx^2}}$$

[Out] $-1/2*I*(a^2*x^2+1)^{(1/2)}/a/c/(1-I*a*x)^2/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 32}

$$-\frac{i\sqrt{a^2x^2+1}}{2ac(1-iax)^2\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((3*I)*\text{ArcTan}[a*x])}/(c + a^2*c*x^2)^{(3/2)}, x]$

[Out] $((-1/2*I)*\text{Sqrt}[1 + a^2*x^2])/(a*c*(1 - I*a*x)^2*\text{Sqrt}[c + a^2*c*x^2])$

Rule 32

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] /;$ FreeQ[{a, b, m}, x] && NeQ[m, -1]

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^{\text{IntPart}[p]}*(c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}, \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{3i \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{3i \tan^{-1}(ax)}}{(1 + a^2 x^2)^{3/2}} dx}{c\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{(1 - iax)^3} dx}{c\sqrt{c + a^2 cx^2}} \\
&= -\frac{i\sqrt{1 + a^2 x^2}}{2ac(1 - iax)^2 \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 57, normalized size = 1.16

$$\frac{i\sqrt{1 + a^2 x^2} \sqrt{c + a^2 cx^2}}{2ac^2(-i + ax)(i + ax)^3}$$

Antiderivative was successfully verified.

`[In] Integrate[E^((3*I)*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]``[Out] ((I/2)*Sqrt[1 + a^2*x^2]*Sqrt[c + a^2*c*x^2])/(a*c^2*(-I + a*x)*(I + a*x)^3)`**Maple [A]**

time = 0.08, size = 42, normalized size = 0.86

method	result	size
default	$\frac{i\sqrt{c(a^2x^2 + 1)}}{2\sqrt{a^2x^2 + 1} c^2 a(ax+i)^2}$	42
risch	$\frac{i\sqrt{a^2x^2 + 1}}{2c\sqrt{c(a^2x^2 + 1)} a(ax+i)^2}$	42
gosper	$-\frac{(ax+i)(iax+1)^3}{2a(a^2x^2+1)^{\frac{3}{2}}(a^2cx^2+c)^{\frac{3}{2}}}$	44

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2), x, method=_RETURNVERBOSE)``[Out] 1/2*I/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)/c^2/a/(I+a*x)^2`**Maxima [F(-2)]**

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [A]

time = 7.35, size = 71, normalized size = 1.45

$$\frac{\sqrt{a^2cx^2 + c} \sqrt{a^2x^2 + 1} (iax^2 - 2x)}{2(a^4c^2x^4 + 2ia^3c^2x^3 + 2iac^2x - c^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] 1/2*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)*(I*a*x^2 - 2*x)/(a^4*c^2*x^4 + 2*I*a^3*c^2*x^3 + 2*I*a*c^2*x - c^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\left(\int \frac{1}{\sqrt{a^2cx^2 + c} \sqrt{a^2x^2 + 1}} dx + \int \frac{3ax}{\sqrt{a^2cx^2 + c} \sqrt{a^2x^2 + 1}} dx + \int \frac{a^2x^2}{\sqrt{a^2cx^2 + c} \sqrt{a^2x^2 + 1}} dx + \int \frac{3a^2x^3}{\sqrt{a^2cx^2 + c} \sqrt{a^2x^2 + 1}} dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)/(a**2*c*x**2+c)**(3/2),x)

[Out] -I*(Integral(I/(a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-3*a*x/(a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a**3*x**3/(a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-3*I*a**2*x**2/(a**4*c*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] integrate((I*a*x + 1)^3/((a^2*c*x^2 + c)^(3/2)*(a^2*x^2 + 1)^(3/2)), x)

Mupad [B]

time = 1.25, size = 41, normalized size = 0.84

$$\frac{\sqrt{c(a^2x^2 + 1)} \operatorname{li}}{2ac^2\sqrt{a^2x^2 + 1}(ax + 1)^2}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^3/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^(3/2)),x)

[Out] ((c*(a^2*x^2 + 1))^(1/2)*1i)/(2*a*c^2*(a^2*x^2 + 1)^(1/2)*(a*x + 1i)^2)

$$3.331 \quad \int \frac{e^{2i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=54

$$-\frac{2i(1+iax)}{3a(c+a^2cx^2)^{3/2}} + \frac{x}{3c\sqrt{c+a^2cx^2}}$$

[Out] $-2/3*I*(1+I*a*x)/a/(a^2*c*x^2+c)^{(3/2)}+1/3*x/c/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5183, 667, 197}

$$\frac{x}{3c\sqrt{a^2cx^2+c}} - \frac{2i(1+iax)}{3a(a^2cx^2+c)^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*\text{ArcTan}[a*x])}/(c + a^2*c*x^2)^{(3/2)}, x]$

[Out] $(((-2*I)/3)*(1 + I*a*x))/(a*(c + a^2*c*x^2)^{(3/2)}) + x/(3*c*\text{Sqrt}[c + a^2*c*x^2])$

Rule 197

$\text{Int}[(a_ + (b_)*(x_)^{(n)})^{(p)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^{(p+1)}/a), x] /;$ FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 667

$\text{Int}[(d_ + (e_)*(x_)^2*((a_ + (c_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Simp}[e*(d + e*x)*((a + c*x^2)^{(p+1)}/(c*(p+1))), x] - \text{Dist}[e^2*((p+2)/(c*(p+1))), \text{Int}[(a + c*x^2)^{(p+1)}, x], x] /;$ FreeQ[{a, c, d, e, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && LtQ[p, -1]

Rule 5183

$\text{Int}[E^{(\text{ArcTan}[(a_)*(x_)])*(n_)*((c_ + (d_)*(x_)^2)^{(p_)}), x_Symbol] \rightarrow \text{Dist}[1/c^{(I*(n/2))}, \text{Int}[(c + d*x^2)^{(p + I*(n/2))}/(1 + I*a*x)^{(I*n)}, x], x] /;$ FreeQ[{a, c, d, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0]) && ILtQ[I*(n/2), 0]

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx &= c \int \frac{(1 + iax)^2}{(c + a^2 cx^2)^{5/2}} dx \\ &= -\frac{2i(1 + iax)}{3a(c + a^2 cx^2)^{3/2}} + \frac{1}{3} \int \frac{1}{(c + a^2 cx^2)^{3/2}} dx \\ &= -\frac{2i(1 + iax)}{3a(c + a^2 cx^2)^{3/2}} + \frac{x}{3c\sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 78, normalized size = 1.44

$$\frac{(2 - iax)\sqrt{1 + iax}\sqrt{1 + a^2 x^2}}{3ac\sqrt{1 - iax}(i + ax)\sqrt{c + a^2 cx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[E^((2*I)*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]

[Out] ((2 - I*a*x)*Sqrt[1 + I*a*x]*Sqrt[1 + a^2*x^2])/(3*a*c*Sqrt[1 - I*a*x]*(I + a*x)*Sqrt[c + a^2*c*x^2])

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 397 vs. 2(44) = 88.

time = 0.10, size = 398, normalized size = 7.37

method	result
trager	$\frac{(a^3 x^3 + 3ax - 2i)\sqrt{a^2 c x^2 + c}}{3c^2(a^2 x^2 + 1)^2 a}$
default	$-\frac{x}{c\sqrt{a^2 c x^2 + c}} + \frac{(i\sqrt{-a^2} - a) \left(\frac{1}{3c\sqrt{-a^2} \left(x + \frac{\sqrt{-a^2}}{a^2}\right) \sqrt{\left(x + \frac{\sqrt{-a^2}}{a^2}\right)^2 a^2 c - 2c\sqrt{-a^2}} \left(x + \frac{\sqrt{-a^2}}{a^2}\right)} \right)}{a\sqrt{-a^2}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2), x, method=_RETURNVERBOSE)

[Out] -x/c/(a^2*c*x^2+c)^(1/2)+1/a*(I*(-a^2)^(1/2)-a)/(-a^2)^(1/2)*(1/3/c/(-a^2)^(1/2)/(x+(-a^2)^(1/2)/a^2)/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)+1/3/c^2/(-a^2)^(1/2)*(2*a^2*c*(x+(-a^2)^(1/2)/a^2)-2*c*(-a^2)^(1/2))/((x+(-a^2)^(1/2)/a^2)^2*a^2*c-2*c*(-a^2)^(1/2)*(x+(-a^2)^(1/2)/a^2))^(1/2)

$$2)^{(1/2)/a^2})^{(1/2)}+1/a*(I*(-a^2)^{(1/2)}+a)/(-a^2)^{(1/2)}*(-1/3/c/(-a^2)^{(1/2)}/(x-(-a^2)^{(1/2)/a^2})/((x-(-a^2)^{(1/2)/a^2})^2*a^2*c+2*c*(-a^2)^{(1/2)}*(x-(-a^2)^{(1/2)/a^2}))^{(1/2)}-1/3/c^2/(-a^2)^{(1/2)}*(2*a^2*c*(x-(-a^2)^{(1/2)/a^2})+2*c*(-a^2)^{(1/2)})/((x-(-a^2)^{(1/2)/a^2})^2*a^2*c+2*c*(-a^2)^{(1/2)}*(x-(-a^2)^{(1/2)/a^2}))^{(1/2)})$$

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((I*a*x + 1)^2/((a^2*c*x^2 + c)^(3/2)*(a^2*x^2 + 1)), x)

Fricas [A]

time = 3.63, size = 47, normalized size = 0.87

$$\frac{\sqrt{a^2cx^2 + c} (ax + 2i)}{3 (a^3c^2x^2 + 2ia^2c^2x - ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] 1/3*sqrt(a^2*c*x^2 + c)*(a*x + 2*I)/(a^3*c^2*x^2 + 2*I*a^2*c^2*x - a*c^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{a^2x^2}{a^4cx^4\sqrt{a^2cx^2+c} + 2a^2cx^2\sqrt{a^2cx^2+c} + c\sqrt{a^2cx^2+c}} dx - \int \left(-\frac{2iax}{a^4cx^4\sqrt{a^2cx^2+c} + 2a^2cx^2\sqrt{a^2cx^2+c} + c\sqrt{a^2cx^2+c}} \right) dx - \int \left(-\frac{1}{a^4cx^4\sqrt{a^2cx^2+c} + 2a^2cx^2\sqrt{a^2cx^2+c} + c\sqrt{a^2cx^2+c}} \right) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)/(a**2*c*x**2+c)**(3/2),x)

[Out] -Integral(a**2*x**2/(a**4*c*x**4*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*c*x**2 + c)), x) - Integral(-2*I*a*x/(a**4*c*x**4*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*c*x**2 + c)), x) - Integral(-1/(a**4*c*x**4*sqrt(a**2*c*x**2 + c) + 2*a**2*c*x**2*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*c*x**2 + c)), x)

Giac [A]

time = 0.43, size = 76, normalized size = 1.41

$$\frac{2\sqrt{a^2c} \left(3\sqrt{a^2c} x - 3\sqrt{a^2cx^2 + c} + i\sqrt{c} \right)}{3 \left(\sqrt{a^2c} x - \sqrt{a^2cx^2 + c} + i\sqrt{c} \right)^3 a^2c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] -2/3*sqrt(a^2*c)*(3*sqrt(a^2*c)*x - 3*sqrt(a^2*c*x^2 + c) + I*sqrt(c))/((sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c) + I*sqrt(c))^3*a^2*c)

Mupad [B]

time = 0.65, size = 32, normalized size = 0.59

$$\frac{a^3 x^3 + 3 a x - 2i}{3 a (c (a^2 x^2 + 1))^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a*x*1i + 1)^2/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)),x)

[Out] (3*a*x + a^3*x^3 - 2i)/(3*a*(c*(a^2*x^2 + 1))^(3/2))

$$3.332 \quad \int \frac{e^{i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=88

$$\frac{\sqrt{1+a^2x^2}}{2ac(i+ax)\sqrt{c+a^2cx^2}} + \frac{\sqrt{1+a^2x^2} \operatorname{ArcTan}(ax)}{2ac\sqrt{c+a^2cx^2}}$$

[Out] 1/2*(a^2*x^2+1)^(1/2)/a/c/(I+a*x)/(a^2*c*x^2+c)^(1/2)+1/2*arctan(a*x)*(a^2*x^2+1)^(1/2)/a/c/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.06, antiderivative size = 88, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5184, 5181, 46, 209}

$$\frac{\sqrt{a^2x^2+1} \operatorname{ArcTan}(ax)}{2ac\sqrt{a^2cx^2+c}} + \frac{\sqrt{a^2x^2+1}}{2ac(ax+i)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[E^(I*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]

[Out] Sqrt[1 + a^2*x^2]/(2*a*c*(I + a*x)*Sqrt[c + a^2*c*x^2]) + (Sqrt[1 + a^2*x^2]*ArcTan[a*x])/(2*a*c*Sqrt[c + a^2*c*x^2])

Rule 46

Int[((a_) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 209

Int[((a_) + (b_.)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{i \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{i \tan^{-1}(ax)}}{(1 + a^2 x^2)^{3/2}} dx}{c \sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{(1 - iax)^2 (1 + iax)} dx}{c \sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \left(-\frac{1}{2(i+ax)^2} + \frac{1}{2(1+a^2 x^2)} \right) dx}{c \sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2}}{2ac(i + ax) \sqrt{c + a^2 cx^2}} + \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{1 + a^2 x^2} dx}{2c \sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2}}{2ac(i + ax) \sqrt{c + a^2 cx^2}} + \frac{\sqrt{1 + a^2 x^2} \tan^{-1}(ax)}{2ac \sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 51, normalized size = 0.58

$$\frac{\sqrt{1 + a^2 x^2} \left(\frac{1}{i + ax} + \text{ArcTan}(ax) \right)}{2ac \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*ArcTan[a*x])/(c + a^2*c*x^2)^(3/2), x]

[Out] (Sqrt[1 + a^2*x^2]*((I + a*x)^(-1) + ArcTan[a*x]))/(2*a*c*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.08, size = 58, normalized size = 0.66

method	result	size
default	$-\frac{\sqrt{c(a^2 x^2 + 1)} (-\arctan(ax)a^2 x^2 - ax + i - \arctan(ax))}{2(a^2 x^2 + 1)^{\frac{3}{2}} a c^2}$	58

risch	$\frac{\sqrt{a^2x^2+1}}{2c\sqrt{c(a^2x^2+1)}a(ax+i)} - \frac{i\sqrt{a^2x^2+1}\ln(-iax-1)}{4c\sqrt{c(a^2x^2+1)}a} + \frac{i\sqrt{a^2x^2+1}\ln(iax-1)}{4c\sqrt{c(a^2x^2+1)}a}$	124
-------	--	-----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)`
)

[Out] $-1/2/(a^2*x^2+1)^(3/2)*(c*(a^2*x^2+1))^(1/2)/a*(-\arctan(a*x)*a^2*x^2-a*x+I-\arctan(a*x))/c^2$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(74) = 148$.

time = 5.40, size = 317, normalized size = 3.60

$$\frac{(ia^3c^2x^3 - a^2c^2x^2 + iac^2x - c^2)\sqrt{\frac{1}{a^2c^3}} \log\left(\frac{2\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}e^{ax-(ia^3c^2x^3-ia^2c^2x-iac^2x+c^2)}\sqrt{\frac{1}{a^2c^3}}}{a^2x^2+2a^2x+1}\right) + (-ia^3c^2x^3 + a^2c^2x^2 - iac^2x + c^2)\sqrt{\frac{1}{a^2c^3}} \log\left(\frac{2\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}e^{ax-(-ia^3c^2x^3+ia^2c^2x+iac^2x-c^2)}\sqrt{\frac{1}{a^2c^3}}}{a^2x^2+2a^2x+1}\right) + 4i\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}x}{8(a^3c^2x^3 + ia^2c^2x^2 + ac^2x + ic^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`

[Out] $1/8*((I*a^3*c^2*x^3 - a^2*c^2*x^2 + I*a*c^2*x - c^2)*\sqrt{1/(a^2*c^3)}*\log(2*(2*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^6*x - (I*a^{10}*c^2*x^4 - I*a^6*c^2)*\sqrt{1/(a^2*c^3)}))/(a^4*x^4 + 2*a^2*x^2 + 1)) + (-I*a^3*c^2*x^3 + a^2*c^2*x^2 - I*a*c^2*x + c^2)*\sqrt{1/(a^2*c^3)}*\log(2*(2*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^6*x - (-I*a^{10}*c^2*x^4 + I*a^6*c^2)*\sqrt{1/(a^2*c^3)}))/(a^4*x^4 + 2*a^2*x^2 + 1)) + 4*I*\sqrt{a^2*c*x^2 + c}*\sqrt{a^2*x^2 + 1}*x/(a^3*c^2*x^3 + I*a^2*c^2*x^2 + a*c^2*x + I*c^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i\left(\int\left(-\frac{i}{a^2cx^2\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}+c\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}}\right)dx + \int\frac{ax}{a^2cx^2\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}+c\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}}dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)/(a**2*c*x**2+c)**(3/2),x)
```

```
[Out] I*(Integral(-I/(a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a*x/(a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x))
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{1 + a x \operatorname{li}}{(c a^2 x^2 + c)^{3/2} \sqrt{a^2 x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a*x*1i + 1)/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^(1/2)),x)
```

```
[Out] int((a*x*1i + 1)/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^(1/2)), x)
```


$$3.333 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=89

$$-\frac{\sqrt{1+a^2x^2}}{2ac(i-ax)\sqrt{c+a^2cx^2}} + \frac{\sqrt{1+a^2x^2} \operatorname{ArcTan}(ax)}{2ac\sqrt{c+a^2cx^2}}$$

[Out] $-1/2*(a^2*x^2+1)^{(1/2)}/a/c/(I-a*x)/(a^2*c*x^2+c)^{(1/2)}+1/2*\arctan(a*x)*(a^2*x^2+1)^{(1/2)}/a/c/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 89, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 4, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.160$, Rules used = {5184, 5181, 46, 209}

$$\frac{\sqrt{a^2x^2+1} \operatorname{ArcTan}(ax)}{2ac\sqrt{a^2cx^2+c}} - \frac{\sqrt{a^2x^2+1}}{2ac(-ax+i)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[1/(E^(I*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)), x]

[Out] $-1/2*\operatorname{Sqrt}[1 + a^2*x^2]/(a*c*(I - a*x)*\operatorname{Sqrt}[c + a^2*c*x^2]) + (\operatorname{Sqrt}[1 + a^2*x^2]*\operatorname{ArcTan}[a*x])/(2*a*c*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 46

Int[((a_) + (b_)*(x_)^(m_))*((c_) + (d_)*(x_)^(n_)), x_Symbol] := Int[ExpandIntegrand[(a + b*x)^m*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d}, x] && NeQ[b*c - a*d, 0] && ILtQ[m, 0] && IntegerQ[n] && !(IGtQ[n, 0] && LtQ[m + n + 2, 0])

Rule 209

Int[((a_) + (b_)*(x_)^2)^(-1), x_Symbol] := Simp[(1/(Rt[a, 2]*Rt[b, 2]))*ArcTan[Rt[b, 2]*(x/Rt[a, 2])], x] /; FreeQ[{a, b}, x] && PosQ[a/b] && (GtQ[a, 0] || GtQ[b, 0])

Rule 5181

Int[E^(ArcTan[(a_)*(x_)])*(n_))*((c_) + (d_)*(x_)^2)^(p_), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := D
ist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(
1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && Eq
Q[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{-i \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-i \tan^{-1}(ax)}}{(1 + a^2 x^2)^{3/2}} dx}{c\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{(1 - iax)(1 + iax)^2} dx}{c\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \left(-\frac{1}{2(-i + iax)^2} + \frac{1}{2(1 + a^2 x^2)} \right) dx}{c\sqrt{c + a^2 cx^2}} \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2ac(i - ax)\sqrt{c + a^2 cx^2}} + \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{1 + a^2 x^2} dx}{2c\sqrt{c + a^2 cx^2}} \\
&= -\frac{\sqrt{1 + a^2 x^2}}{2ac(i - ax)\sqrt{c + a^2 cx^2}} + \frac{\sqrt{1 + a^2 x^2} \tan^{-1}(ax)}{2ac\sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 60, normalized size = 0.67

$$\frac{\sqrt{1 + a^2 x^2} \left(-\frac{1}{2a(i - ax)} + \frac{\text{ArcTan}(ax)}{2a} \right)}{c\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

```
[In] Integrate[1/(E^(I*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)), x]
```

```
[Out] (Sqrt[1 + a^2*x^2]*(-1/2*1/(a*(I - a*x)) + ArcTan[a*x]/(2*a)))/(c*Sqrt[c +
a^2*c*x^2])
```

Maple [A]

time = 0.08, size = 86, normalized size = 0.97

method	result	size
default	$\frac{\sqrt{c(a^2 x^2 + 1)} (i \ln(-ax + i) ax - i \ln(ax + i) ax + \ln(-ax + i) - \ln(ax + i) - 2)}{4\sqrt{a^2 x^2 + 1} c^2(-ax + i)a}$	86

risch	$\frac{\sqrt{a^2x^2+1}}{2c\sqrt{c(a^2x^2+1)} a^{(ax-i)}} + \frac{i\sqrt{a^2x^2+1} \ln(iax-1)}{4c\sqrt{c(a^2x^2+1)} a} - \frac{i\sqrt{a^2x^2+1} \ln(-iax-1)}{4c\sqrt{c(a^2x^2+1)} a}$	124
-------	--	-----

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)`

[Out] $\frac{1}{4}(a^2x^2+1)^{1/2}(c(a^2x^2+1))^{1/2}(I\ln(I-ax)*ax-I\ln(I+ax)*ax+\ln(I-ax)-\ln(I+ax)-2)/c^2/(I-ax)/a$

Maxima [A]

time = 0.27, size = 52, normalized size = 0.58

$$\frac{\sqrt{c}}{2(a^2c^2x - iac^2)} - \frac{i \log(ax - i)}{4ac^{\frac{3}{2}}} + \frac{i \log(iax - 1)}{4ac^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`

[Out] $\frac{1}{2}\sqrt{c}/(a^2c^2x - Iac^2) - \frac{1}{4}I\log(ax - I)/(ac^{3/2}) + \frac{1}{4}I\log(Iax - 1)/(ac^{3/2})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 317 vs. $2(74) = 148$.

time = 3.31, size = 317, normalized size = 3.56

$$\frac{(ia^3c^2x^3 + a^2c^2x^2 + iac^2x + c)\sqrt{\frac{1}{a^2c^2}} \log\left(\frac{2\left(2\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}e^{ix-(ia^3c^2x^3-iac^2x-c)}\sqrt{\frac{1}{a^2c^2}}\right)}{a^{2x^2+2a^2x+1}}\right) + (-ia^3c^2x^3 - a^2c^2x^2 - iac^2x - c)\sqrt{\frac{1}{a^2c^2}} \log\left(\frac{2\left(2\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}e^{-ix-(ia^3c^2x^3+iac^2x+c)}\sqrt{\frac{1}{a^2c^2}}\right)}{a^{2x^2+2a^2x+1}}\right) - 4i\sqrt{a^2cx^2+c}\sqrt{a^2x^2+1}x}{8(a^3c^2x^3 - ia^2c^2x^2 + ac^2x - ic^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`

[Out] $\frac{1}{8}((Ia^3c^2x^3 + a^2c^2x^2 + Iac^2x + c^2)\sqrt{1/(a^2c^3)})\log(2*(2*\sqrt{a^2cx^2+c}*\sqrt{a^2x^2+1})a^6x - (Ia^{10}c^2x^4 - Ia^6c^2)\sqrt{1/(a^2c^3)})/(a^4x^4 + 2a^2x^2 + 1) + (-Ia^3c^2x^3 - a^2c^2x^2 - Iac^2x - c^2)\sqrt{1/(a^2c^3)}\log(2*(2*\sqrt{a^2cx^2+c}*\sqrt{a^2x^2+1})a^6x - (-Ia^{10}c^2x^4 + Ia^6c^2)\sqrt{1/(a^2c^3)})/(a^4x^4 + 2a^2x^2 + 1) - 4I*\sqrt{a^2cx^2+c}*\sqrt{a^2x^2+1}x/(a^3c^2x^3 - Ia^2c^2x^2 + ac^2x - Ic^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{\sqrt{a^2x^2+1}}{a^3cx^3\sqrt{a^2cx^2+c} - ia^2cx^2\sqrt{a^2cx^2+c} + acx\sqrt{a^2cx^2+c} - ic\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)*(a**2*x**2+1)**(1/2)/(a**2*c*x**2+c)**(3/2),x)
```

```
[Out] -I*Integral(sqrt(a**2*x**2 + 1)/(a**3*c*x**3*sqrt(a**2*c*x**2 + c) - I*a**2*c*x**2*sqrt(a**2*c*x**2 + c) + a*c*x*sqrt(a**2*c*x**2 + c) - I*c*sqrt(a**2*c*x**2 + c)), x)
```

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")
```

```
[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{\sqrt{a^2 x^2 + 1}}{(c a^2 x^2 + c)^{3/2} (1 + a x i)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a^2*x^2 + 1)^(1/2)/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)),x)
```

```
[Out] int((a^2*x^2 + 1)^(1/2)/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)), x)
```

$$3.334 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=54

$$\frac{2i(1-iax)}{3a(c+a^2cx^2)^{3/2}} + \frac{x}{3c\sqrt{c+a^2cx^2}}$$

[Out] $2/3*I*(1-I*a*x)/a/(a^2*c*x^2+c)^{(3/2)}+1/3*x/c/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.04, antiderivative size = 54, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5182, 667, 197}

$$\frac{x}{3c\sqrt{a^2cx^2+c}} + \frac{2i(1-iax)}{3a(a^2cx^2+c)^{3/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((2*I)*\text{ArcTan}[a*x])*(c+a^2*c*x^2)^{(3/2))}, x]$

[Out] $((2*I)/3)*(1-I*a*x)/(a*(c+a^2*c*x^2)^{(3/2)}+x/(3*c*\text{Sqrt}[c+a^2*c*x^2])$

Rule 197

$\text{Int}[(a_+ + (b_+)*(x_+)^{(n_+))}^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[x*((a + b*x^n)^{(p+1)}/a), x] /;$ FreeQ[{a, b, n, p}, x] && EqQ[1/n + p + 1, 0]

Rule 667

$\text{Int}[(d_+ + (e_+)*(x_+))^{2*((a_+ + (c_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Simp}[e*(d + e*x)*((a + c*x^2)^{(p+1)}/(c*(p+1))), x] - \text{Dist}[e^{2*((p+2)/(c*(p+1))}, \text{Int}[(a + c*x^2)^{(p+1)}, x], x] /;$ FreeQ[{a, c, d, e, p}, x] && EqQ[c*d^2 + a*e^2, 0] && !IntegerQ[p] && LtQ[p, -1]

Rule 5182

$\text{Int}[E^{(\text{ArcTan}[(a_+)*(x_+)]*(n_+))*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \text{Dist}[c^{(I*(n/2))}, \text{Int}[(c + d*x^2)^{(p - I*(n/2))}*(1 - I*a*x)^{(I*n)}, x], x] /;$ FreeQ[{a, c, d, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0]) && IGtQ[I*(n/2), 0]

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)}}{(c + a^2cx^2)^{3/2}} dx &= c \int \frac{(1 - iax)^2}{(c + a^2cx^2)^{5/2}} dx \\ &= \frac{2i(1 - iax)}{3a(c + a^2cx^2)^{3/2}} + \frac{1}{3} \int \frac{1}{(c + a^2cx^2)^{3/2}} dx \\ &= \frac{2i(1 - iax)}{3a(c + a^2cx^2)^{3/2}} + \frac{x}{3c\sqrt{c + a^2cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 78, normalized size = 1.44

$$\frac{\sqrt{1 - iax} (2 + iax) \sqrt{1 + a^2x^2}}{3ac\sqrt{1 + iax} (-i + ax) \sqrt{c + a^2cx^2}}$$

Warning: Unable to verify antiderivative.

[In] Integrate[1/(E^((2*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)), x]

[Out] (Sqrt[1 - I*a*x]*(2 + I*a*x)*Sqrt[1 + a^2*x^2])/(3*a*c*Sqrt[1 + I*a*x]*(-I + a*x)*Sqrt[c + a^2*c*x^2])

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 136 vs. 2(44) = 88.

time = 0.07, size = 137, normalized size = 2.54

method	result
default	$-\frac{x}{c\sqrt{a^2cx^2 + c}} - \frac{2i \left(\frac{i}{3ac(x - \frac{i}{a}) \sqrt{(x - \frac{i}{a})^2 a^2c + 2iac}} + \frac{i(2a^2c(x - \frac{i}{a}) + 2iac)}{3ac^2 \sqrt{(x - \frac{i}{a})^2 a^2c + 2iac}} \right)}{a}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2), x, method=_RETURNVERBOSE)

[Out] -x/c/(a^2*c*x^2+c)^(1/2)-2*I/a*(1/3*I/a/c/(x-I/a)/((x-I/a)^2*a^2*c+2*I*a*c*(x-I/a))^(1/2)+1/3*I/a/c^2*(2*a^2*c*(x-I/a)+2*I*a*c)/((x-I/a)^2*a^2*c+2*I*a*c*(x-I/a))^(1/2))

Maxima [A]

time = 0.27, size = 59, normalized size = 1.09

$$\frac{x}{3\sqrt{a^2cx^2 + c}c} + \frac{2i}{3i\sqrt{a^2cx^2 + c}a^2cx + 3\sqrt{a^2cx^2 + c}ac}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] 1/3*x/(sqrt(a^2*c*x^2 + c)*c) + 2*I/(3*I*sqrt(a^2*c*x^2 + c)*a^2*c*x + 3*sqrt(a^2*c*x^2 + c)*a*c)

Fricas [A]

time = 1.79, size = 47, normalized size = 0.87

$$\frac{\sqrt{a^2cx^2 + c} (ax - 2i)}{3 (a^3c^2x^2 - 2ia^2c^2x - ac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] 1/3*sqrt(a^2*c*x^2 + c)*(a*x - 2*I)/(a^3*c^2*x^2 - 2*I*a^2*c^2*x - a*c^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-\int \frac{a^2x^2}{a^4cx^4\sqrt{a^2cx^2+c} - 2ia^3cx^3\sqrt{a^2cx^2+c} - 2iacx\sqrt{a^2cx^2+c} - c\sqrt{a^2cx^2+c}} dx - \int \frac{1}{a^4cx^4\sqrt{a^2cx^2+c} - 2ia^3cx^3\sqrt{a^2cx^2+c} - 2iacx\sqrt{a^2cx^2+c} - c\sqrt{a^2cx^2+c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**2*(a**2*x**2+1)/(a**2*c*x**2+c)**(3/2),x)

[Out] -Integral(a**2*x**2/(a**4*c*x**4*sqrt(a**2*c*x**2 + c) - 2*I*a**3*c*x**3*sqrt(a**2*c*x**2 + c) - 2*I*a*c*x*sqrt(a**2*c*x**2 + c) - c*sqrt(a**2*c*x**2 + c)), x) - Integral(1/(a**4*c*x**4*sqrt(a**2*c*x**2 + c) - 2*I*a**3*c*x**3*sqrt(a**2*c*x**2 + c) - 2*I*a*c*x*sqrt(a**2*c*x**2 + c) - c*sqrt(a**2*c*x**2 + c)), x)

Giac [A]

time = 0.48, size = 76, normalized size = 1.41

$$\frac{2\sqrt{a^2c} \left(3\sqrt{a^2c} x - 3\sqrt{a^2cx^2 + c} - i\sqrt{c} \right)}{3 \left(\sqrt{a^2c} x - \sqrt{a^2cx^2 + c} - i\sqrt{c} \right)^3 a^2c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] -2/3*sqrt(a^2*c)*(3*sqrt(a^2*c)*x - 3*sqrt(a^2*c*x^2 + c) - I*sqrt(c))/((sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c) - I*sqrt(c))^3*a^2*c)

Mupad [B]

time = 0.65, size = 32, normalized size = 0.59

$$\frac{a^3 x^3 + 3 a x + 2i}{3 a (c (a^2 x^2 + 1))^{3/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)^2),x)

[Out] (3*a*x + a^3*x^3 + 2i)/(3*a*(c*(a^2*x^2 + 1))^(3/2))

$$3.335 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=49

$$\frac{i\sqrt{1+a^2x^2}}{2ac(1+iax)^2\sqrt{c+a^2cx^2}}$$

[Out] $1/2*I*(a^2*x^2+1)^{(1/2)}/a/c/(1+I*a*x)^2/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.05, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$, Rules used = {5184, 5181, 32}

$$\frac{i\sqrt{a^2x^2+1}}{2ac(1+iax)^2\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((3*I)*\text{ArcTan}[a*x])*(c+a^2*c*x^2)^{(3/2))}, x]$

[Out] $((I/2)*\text{Sqrt}[1+a^2*x^2])/(a*c*(1+I*a*x)^2*\text{Sqrt}[c+a^2*c*x^2])$

Rule 32

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}, x_Symbol] \rightarrow \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] /;$ $\text{FreeQ}\{a, b, m, x\} \ \&\& \ \text{NeQ}[m, -1]$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ $\text{FreeQ}\{a, c, d, n, p, x\} \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \rightarrow \text{Dist}[c^{\text{IntPart}[p]}*(c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}, \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ $\text{FreeQ}\{a, c, d, n, p, x\} \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int \frac{e^{-3i \tan^{-1}(ax)}}{(c + a^2 cx^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-3i \tan^{-1}(ax)}}{(1 + a^2 x^2)^{3/2}} dx}{c \sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{1}{(1 + iax)^3} dx}{c \sqrt{c + a^2 cx^2}} \\ &= \frac{i \sqrt{1 + a^2 x^2}}{2ac(1 + iax)^2 \sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 57, normalized size = 1.16

$$\frac{i \sqrt{1 + a^2 x^2} \sqrt{c + a^2 cx^2}}{2ac^2(-i + ax)^3(i + ax)}$$

Antiderivative was successfully verified.

`[In] Integrate[1/(E^((3*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)),x]``[Out] ((-1/2*I)*Sqrt[1 + a^2*x^2]*Sqrt[c + a^2*c*x^2])/(a*c^2*(-I + a*x)^3*(I + a*x))`**Maple [A]**

time = 0.08, size = 43, normalized size = 0.88

method	result	size
risch	$-\frac{i \sqrt{a^2 x^2 + 1}}{2c \sqrt{c(a^2 x^2 + 1)} a(ax-i)^2}$	42
default	$\frac{i \sqrt{c(a^2 x^2 + 1)}}{2 \sqrt{a^2 x^2 + 1} c^2 a(iax+1)^2}$	43
gospers	$\frac{(-ax+i)(a^2 x^2 + 1)^{\frac{3}{2}}}{2a(iax+1)^3(a^2 c x^2 + c)^{\frac{3}{2}}}$	45

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)``[Out] 1/2*I/(a^2*x^2+1)^(1/2)*(c*(a^2*x^2+1))^(1/2)/c^2/a/(1+I*a*x)^2`**Maxima [A]**

time = 0.27, size = 29, normalized size = 0.59

$$\frac{1}{2i a^3 c^{\frac{3}{2}} x^2 + 4 a^2 c^{\frac{3}{2}} x - 2i a c^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] 1/(2*I*a^3*c^(3/2)*x^2 + 4*a^2*c^(3/2)*x - 2*I*a*c^(3/2))

Fricas [A]

time = 2.54, size = 71, normalized size = 1.45

$$\frac{\sqrt{a^2cx^2 + c} \sqrt{a^2x^2 + 1} (-iax^2 - 2x)}{2(a^4c^2x^4 - 2ia^3c^2x^3 - 2iac^2x - c^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] 1/2*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)*(-I*a*x^2 - 2*x)/(a^4*c^2*x^4 - 2*I*a^3*c^2*x^3 - 2*I*a*c^2*x - c^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\left(\int \frac{\sqrt{a^2x^2+1}}{a^4cx^2\sqrt{a^2cx^2+c} - 3ia^4cx^2\sqrt{a^2cx^2+c} - 2a^2c^2\sqrt{a^2cx^2+c} - 2ia^2c^2\sqrt{a^2cx^2+c} - 3acx\sqrt{a^2cx^2+c} + ic\sqrt{a^2cx^2+c}} dx + \int \frac{a^2x^2\sqrt{a^2x^2+1}}{a^4cx^2\sqrt{a^2cx^2+c} - 3ia^4cx^2\sqrt{a^2cx^2+c} - 2a^2c^2\sqrt{a^2cx^2+c} - 2ia^2c^2\sqrt{a^2cx^2+c} - 3acx\sqrt{a^2cx^2+c} + ic\sqrt{a^2cx^2+c}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/(a**2*c*x**2+c)**(3/2),x)

[Out] I*(Integral(sqrt(a**2*x**2 + 1)/(a**5*c*x**5*sqrt(a**2*c*x**2 + c) - 3*I*a**4*c*x**4*sqrt(a**2*c*x**2 + c) - 2*a**3*c*x**3*sqrt(a**2*c*x**2 + c) - 2*I*a**2*c*x**2*sqrt(a**2*c*x**2 + c) - 3*a*c*x*sqrt(a**2*c*x**2 + c) + I*c*sqrt(a**2*c*x**2 + c)), x) + Integral(a**2*x**2*sqrt(a**2*x**2 + 1)/(a**5*c*x**5*sqrt(a**2*c*x**2 + c) - 3*I*a**4*c*x**4*sqrt(a**2*c*x**2 + c) - 2*a**3*c*x**3*sqrt(a**2*c*x**2 + c) - 2*I*a**2*c*x**2*sqrt(a**2*c*x**2 + c) - 3*a*c*x*sqrt(a**2*c*x**2 + c) + I*c*sqrt(a**2*c*x**2 + c)), x))

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INPUT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [B]

time = 1.07, size = 49, normalized size = 1.00

$$\frac{\sqrt{c(a^2x^2 + 1)} \sqrt{a^2x^2 + 1}}{2ac^2(ax + 1i)(1 + ax1i)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*x^2 + 1)^(3/2)/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)^3),x)

[Out] -((c*(a^2*x^2 + 1))^(1/2)*(a^2*x^2 + 1)^(1/2))/(2*a*c^2*(a*x + 1i)*(a*x*1i + 1)^3)

$$3.336 \quad \int \frac{e^{-4i \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=69

$$\frac{ic(1-iax)^4}{3a(c+a^2cx^2)^{5/2}} - \frac{ic(1-iax)^5}{15a(c+a^2cx^2)^{5/2}}$$

[Out] $1/3*I*c*(1-I*a*x)^4/a/(a^2*c*x^2+c)^{(5/2)}-1/15*I*c*(1-I*a*x)^5/a/(a^2*c*x^2+c)^{(5/2)}$

Rubi [A]

time = 0.05, antiderivative size = 69, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 25, $\frac{\text{number of rules}}{\text{integrand size}} = 0.120$,

Rules used = {5182, 673, 665}

$$\frac{ic(1-iax)^4}{3a(a^2cx^2+c)^{5/2}} - \frac{ic(1-iax)^5}{15a(a^2cx^2+c)^{5/2}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[1/(E^{((4*I)*\text{ArcTan}[a*x])*(c+a^2*c*x^2)^{(3/2))},x]$

[Out] $((I/3)*c*(1-I*a*x)^4)/(a*(c+a^2*c*x^2)^{(5/2)}) - ((I/15)*c*(1-I*a*x)^5)/(a*(c+a^2*c*x^2)^{(5/2)})$

Rule 665

$\text{Int}[(d + (e_*)*(x_))^{(m_)*((a_*) + (c_*)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[e*(d + e*x)^m*((a + c*x^2)^{(p+1)})/(2*c*d*(p+1)), x] /; \text{FreeQ}\{a, c, d, e, m, p\}, x] \ \&\& \ \text{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{EqQ}[m + 2*p + 2, 0]$

Rule 673

$\text{Int}[(d + (e_*)*(x_))^{(m_)*((a_*) + (c_*)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Simp}[(-e)*(d + e*x)^m*((a + c*x^2)^{(p+1)})/(2*c*d*(m+p+1)), x] + \text{Dist}[\text{Simplify}[m + 2*p + 2]/(2*d*(m+p+1)), \text{Int}[(d + e*x)^{(m+1)}*(a + c*x^2)^p, x], x] /; \text{FreeQ}\{a, c, d, e, m, p\}, x] \ \&\& \ \text{EqQ}[c*d^2 + a*e^2, 0] \ \&\& \ !\text{IntegerQ}[p] \ \&\& \ \text{ILtQ}[\text{Simplify}[m + 2*p + 2], 0]$

Rule 5182

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_)]*(n_*)))*((c_*) + (d_*)*(x_)^2)^{(p_)}, x_Symbol] \rightarrow \text{Dist}[c^{(I*(n/2))}, \text{Int}[(c + d*x^2)^{(p-I*(n/2))}*(1-I*a*x)^{(I*n)}, x], x] /; \text{FreeQ}\{a, c, d, p\}, x] \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0]) \ \&\& \ \text{IGtQ}[I*(n/2), 0]$

Rubi steps

$$\begin{aligned} \int \frac{e^{-4i \tan^{-1}(ax)}}{(c + a^2cx^2)^{3/2}} dx &= c^2 \int \frac{(1 - iax)^4}{(c + a^2cx^2)^{7/2}} dx \\ &= \frac{ic(1 - iax)^4}{3a(c + a^2cx^2)^{5/2}} - \frac{1}{3}c^2 \int \frac{(1 - iax)^5}{(c + a^2cx^2)^{7/2}} dx \\ &= \frac{ic(1 - iax)^4}{3a(c + a^2cx^2)^{5/2}} - \frac{ic(1 - iax)^5}{15a(c + a^2cx^2)^{5/2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 77, normalized size = 1.12

$$\frac{(1 - iax)^{3/2}(-4i + ax)\sqrt{1 + a^2x^2}}{15ac\sqrt{1 + iax}(-i + ax)^2\sqrt{c + a^2cx^2}}$$

Warning: Unable to verify antiderivative.

`[In] Integrate[1/(E^((4*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)), x]``[Out] ((1 - I*a*x)^(3/2)*(-4*I + a*x)*Sqrt[1 + a^2*x^2])/(15*a*c*Sqrt[1 + I*a*x]*(-I + a*x)^2*Sqrt[c + a^2*c*x^2])`**Maple [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 306 vs. 2(57) = 114.

time = 0.07, size = 307, normalized size = 4.45

method	result
default	$\frac{x}{c\sqrt{a^2cx^2 + c}} - \frac{4 \left(\frac{i}{5ac(x - \frac{i}{a})^2 \sqrt{(x - \frac{i}{a})^2 a^2c + 2iac(x - \frac{i}{a})}} + \frac{3ia \left(\frac{i}{3ac(x - \frac{i}{a}) \sqrt{(x - \frac{i}{a})^2 a^2c + 2iac(x - \frac{i}{a})}} \right)}{a^2} \right)}{a^2}$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2), x, method=_RETURNVERBOSE)``[Out] x/c/(a^2*c*x^2+c)^(1/2)-4/a^2*(1/5*I/a/c/(x-I/a)^2/((x-I/a)^2*a^2*c+2*I*a*c*(x-I/a)^(1/2)+3/5*I*a*(1/3*I/a/c/(x-I/a)/((x-I/a)^2*a^2*c+2*I*a*c*(x-I/a)^(1/2)+1/3*I/a/c^2*(2*a^2*c*(x-I/a)+2*I*a*c)/((x-I/a)^2*a^2*c+2*I*a*c*(x-I`

$$\frac{x}{15\sqrt{a^2cx^2+c}} - \frac{4i}{5\left(\sqrt{a^2cx^2+c}a^3cx^2 - 2i\sqrt{a^2cx^2+c}a^2cx - \sqrt{a^2cx^2+c}ac\right)} - \frac{8i}{15i\sqrt{a^2cx^2+c}a^2cx + 15\sqrt{a^2cx^2+c}ac}$$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 119 vs. 2(53) = 106.

time = 0.27, size = 119, normalized size = 1.72

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] -1/15*x/(sqrt(a^2*c*x^2 + c)*c) - 4/5*I/(sqrt(a^2*c*x^2 + c)*a^3*c*x^2 - 2*I*sqrt(a^2*c*x^2 + c)*a^2*c*x - sqrt(a^2*c*x^2 + c)*a*c) - 8*I/(15*I*sqrt(a^2*c*x^2 + c)*a^2*c*x + 15*sqrt(a^2*c*x^2 + c)*a*c)

Fricas [A]

time = 2.91, size = 66, normalized size = 0.96

$$\frac{\sqrt{a^2cx^2+c}(a^2x^2 - 3iax + 4)}{15(a^4c^2x^3 - 3ia^3c^2x^2 - 3a^2c^2x + iac^2)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] -1/15*sqrt(a^2*c*x^2 + c)*(a^2*x^2 - 3I*a*x + 4)/(a^4*c^2*x^3 - 3I*a^3*c^2*x^2 - 3*a^2*c^2*x + I*a*c^2)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{(a^2x^2 + 1)^2}{(c(a^2x^2 + 1))^{\frac{3}{2}}(ax - i)^4} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(1/(1+I*a*x)**4*(a**2*x**2+1)**2/(a**2*c*x**2+c)**(3/2),x)

[Out] Integral((a**2*x**2 + 1)**2/((c*(a**2*x**2 + 1))**(3/2)*(a*x - I)**4), x)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 134 vs. 2(53) = 106.

time = 0.47, size = 134, normalized size = 1.94

$$\frac{2\left(15\left(\sqrt{a^2c}x - \sqrt{a^2cx^2+c}\right)^3\sqrt{c} + 5i\left(\sqrt{a^2c}x - \sqrt{a^2cx^2+c}\right)^2c - 5\left(\sqrt{a^2c}x - \sqrt{a^2cx^2+c}\right)c^{\frac{3}{2}} + ic^2\right)}{15\left(\sqrt{a^2c}x - \sqrt{a^2cx^2+c} - i\sqrt{c}\right)^5ac}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(1/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^(3/2),x, algorithm="gia
c")
```

```
[Out] 2/15*(15*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c))^3*sqrt(c) + 5*I*(sqrt(a^2*c)
*x - sqrt(a^2*c*x^2 + c))^2*c - 5*(sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c))*c^(
3/2) + I*c^2)/((sqrt(a^2*c)*x - sqrt(a^2*c*x^2 + c) - I*sqrt(c))^5*a*c)
```

Mupad [B]

time = 1.00, size = 45, normalized size = 0.65

$$\frac{\sqrt{c(a^2x^2+1)}(a^2x^2-ax3i+4)li}{15ac^2(1+axli)^3}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((a^2*x^2 + 1)^2/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)^4),x)
```

```
[Out] ((c*(a^2*x^2 + 1))^(1/2)*(a^2*x^2 - a*x*3i + 4)*1i)/(15*a*c^2*(a*x*1i + 1)^
3)
```


3.337 $\int e^{n \operatorname{ArcTan}(ax)} (c + a^2 cx^2)^2 dx$

Optimal. Leaf size=86

$$-\frac{2^{3-\frac{in}{2}} c^2 (1-iax)^{3+\frac{in}{2}} {}_2F_1\left(-2+\frac{in}{2}, 3+\frac{in}{2}; 4+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a(6i-n)}$$

[Out] $-2^{3-1/2*I*n} * c^2 * (1-I*a*x)^{(3+1/2*I*n)} * \operatorname{hypergeom}([-2+1/2*I*n, 3+1/2*I*n], [4+1/2*I*n], 1/2-1/2*I*a*x) / a / (6*I-n)$

Rubi [A]

time = 0.04, antiderivative size = 86, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5181, 71}

$$-\frac{c^2 2^{3-\frac{in}{2}} (1-iax)^{3+\frac{in}{2}} {}_2F_1\left(\frac{in}{2}-2, \frac{in}{2}+3; \frac{in}{2}+4; \frac{1}{2}(1-iax)\right)}{a(-n+6i)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a*x])}*(c + a^2*c*x^2)^2, x]$

[Out] $-((2^{3-(I/2)*n})*c^2*(1-I*a*x)^{(3+(I/2)*n)}*\operatorname{Hypergeometric2F1}[-2+(I/2)*n, 3+(I/2)*n, 4+(I/2)*n, (1-I*a*x)/2])/(a*(6*I-n))$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] := \operatorname{Simp}[(a + b*x)^{(m+1)} / (b*(m+1)*(b/(b*c - a*d))^{(n)}) * \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a+b*x)/(b*c - a*d)], x] / ; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] \mid\mid !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a_+]*(x_+))}*(n_+)*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] := \operatorname{Dist}[c^{(p)}, \operatorname{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] / ; \operatorname{FreeQ}\{a, c, d, n, p\}, x] \&\& \operatorname{EqQ}[d, a^2*c] \&\& (\operatorname{IntegerQ}[p] \mid\mid \operatorname{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} (c + a^2 cx^2)^2 dx &= c^2 \int (1-iax)^{2+\frac{in}{2}} (1+iax)^{2-\frac{in}{2}} dx \\ &= -\frac{2^{3-\frac{in}{2}} c^2 (1-iax)^{3+\frac{in}{2}} {}_2F_1\left(-2+\frac{in}{2}, 3+\frac{in}{2}; 4+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a(6i-n)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 90, normalized size = 1.05

$$\frac{i2^{2-\frac{in}{2}}c^2(1-iax)^{3+\frac{in}{2}}{}_2F_1\left(-2+\frac{in}{2}, 3+\frac{in}{2}; 4+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a\left(3+\frac{in}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*(c + a^2*c*x^2)^2,x]**[Out]** (I*2^(2 - (I/2)*n)*c^2*(1 - I*a*x)^(3 + (I/2)*n)*Hypergeometric2F1[-2 + (I/2)*n, 3 + (I/2)*n, 4 + (I/2)*n, (1 - I*a*x)/2])/(a*(3 + (I/2)*n))**Maple [F]**

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} (a^2 c x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^2,x)**[Out]** int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^2,x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="maxima")**[Out]** integrate((a^2*c*x^2 + c)^2*e^(n*arctan(a*x)), x)**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="fricas")**[Out]** integral((a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2)*e^(n*arctan(a*x)), x)**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$c^2 \left(\int 2a^2 x^2 e^{n \operatorname{atan}(ax)} dx + \int a^4 x^4 e^{n \operatorname{atan}(ax)} dx + \int e^{n \operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*atan(a*x))*(a**2*c*x**2+c)**2,x)
```

```
[Out] c**2*(Integral(2*a**2*x**2*exp(n*atan(a*x)), x) + Integral(a**4*x**4*exp(n*
atan(a*x)), x) + Integral(exp(n*atan(a*x)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^2,x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} (ca^2 x^2 + c)^2 dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^2,x)
```

```
[Out] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^2, x)
```

3.338 $\int e^{n \operatorname{ArcTan}(ax)} (c + a^2 cx^2) dx$

Optimal. Leaf size=84

$$-\frac{2^{2-\frac{in}{2}} c (1-iax)^{2+\frac{in}{2}} {}_2F_1\left(-1+\frac{in}{2}, 2+\frac{in}{2}; 3+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a(4i-n)}$$

[Out] $-2^{2-\frac{1}{2}i n} c (1-I a x)^{2+\frac{1}{2}i n} \operatorname{hypergeom}\left(\left[2+\frac{1}{2}i n, -1+\frac{1}{2}i n\right], \left[3+\frac{1}{2}i n\right], \frac{1}{2}-\frac{1}{2}i a x\right) / a / (4i-n)$

Rubi [A]

time = 0.03, antiderivative size = 84, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 19, $\frac{\text{number of rules}}{\text{integrand size}} = 0.105$, Rules used = {5181, 71}

$$-\frac{c 2^{2-\frac{in}{2}} (1-iax)^{2+\frac{in}{2}} {}_2F_1\left(\frac{in}{2}-1, \frac{in}{2}+2; \frac{in}{2}+3; \frac{1}{2}(1-iax)\right)}{a(-n+4i)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{n \operatorname{ArcTan}[a x]} (c + a^2 c x^2), x\right]$

[Out] $-((2^{2-(I/2)n} c (1-I a x)^{2+(I/2)n} \operatorname{Hypergeometric2F1}[-1+(I/2)n, 2+(I/2)n, 3+(I/2)n, (1-I a x)/2]) / (a(4i-n)))$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)(x_+))^{m_+} ((c_+ + (d_+)(x_+))^{n_+}), x_Symbol] \rightarrow \operatorname{Simp}[(a + b x)^{m+1} / (b(m+1)(b/(b c - a d))^{n+1}) \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)((a + b x)/(b c - a d))], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n, x\}$ && $\operatorname{NeQ}[b c - a d, 0]$ && $\operatorname{IntegerQ}[m]$ && $\operatorname{IntegerQ}[n]$ && $\operatorname{GtQ}[b/(b c - a d), 0]$ && $(\operatorname{RationalQ}[m] \mid \mid \operatorname{IntegerQ}[n] \mid \mid \operatorname{GtQ}[-d/(b c - a d), 0])$

Rule 5181

$\operatorname{Int}\left[E^{(\operatorname{ArcTan}[a_+](x_+))^{n_+}} ((c_+ + (d_+)(x_+)^2)^{p_+}), x_Symbol\right] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1-I a x)^{p+I(n/2)} (1+I a x)^{p-I(n/2)}], x] /;$ $\operatorname{FreeQ}\{a, c, d, n, p, x\}$ && $\operatorname{EqQ}[d, a^2 c]$ && $(\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} (c + a^2 cx^2) dx &= c \int (1-iax)^{1+\frac{in}{2}} (1+iax)^{1-\frac{in}{2}} dx \\ &= -\frac{2^{2-\frac{in}{2}} c (1-iax)^{2+\frac{in}{2}} {}_2F_1\left(-1+\frac{in}{2}, 2+\frac{in}{2}; 3+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a(4i-n)} \end{aligned}$$

Mathematica [A]

time = 0.01, size = 88, normalized size = 1.05

$$\frac{i2^{1-\frac{in}{2}}c(1-iax)^{2+\frac{in}{2}}{}_2F_1\left(-1+\frac{in}{2}, 2+\frac{in}{2}; 3+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a\left(2+\frac{in}{2}\right)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*(c + a^2*c*x^2), x]

[Out] (I*2^(1 - (I/2)*n)*c*(1 - I*a*x)^(2 + (I/2)*n)*Hypergeometric2F1[-1 + (I/2)*n, 2 + (I/2)*n, 3 + (I/2)*n, (1 - I*a*x)/2])/(a*(2 + (I/2)*n))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} (a^2 c x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*(a^2*c*x^2+c), x)

[Out] int(exp(n*arctan(a*x))*(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c), x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c\left(\int a^2 x^2 e^{n \operatorname{atan}(ax)} dx + \int e^{n \operatorname{atan}(ax)} dx\right)$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*(a**2*c*x**2+c),x)

[Out] c*(Integral(a**2*x**2*exp(n*atan(a*x)), x) + Integral(exp(n*atan(a*x)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} (ca^2x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))*(c + a^2*c*x^2),x)

[Out] int(exp(n*atan(a*x))*(c + a^2*c*x^2), x)

3.339 $\int e^{n \operatorname{ArcTan}(ax)} dx$

Optimal. Leaf size=81

$$-\frac{2^{1-\frac{in}{2}}(1-iax)^{1+\frac{in}{2}} {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; 2+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a(2i-n)}$$

[Out] $-2^{(1-1/2*I*n)}*(1-I*a*x)^{(1+1/2*I*n)}*\operatorname{hypergeom}\left(\left[1/2*I*n, 1+1/2*I*n\right], \left[2+1/2*I*n\right], 1/2-1/2*I*a*x\right)/a/(2*I-n)$

Rubi [A]

time = 0.01, antiderivative size = 81, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 8, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5169, 71}

$$-\frac{2^{1-\frac{in}{2}}(1-iax)^{1+\frac{in}{2}} {}_2F_1\left(\frac{in}{2}+1, \frac{in}{2}; \frac{in}{2}+2; \frac{1}{2}(1-iax)\right)}{a(-n+2i)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n*\operatorname{ArcTan}[a*x])}, x\right]$

[Out] $-((2^{(1-(I/2)*n)}*(1-I*a*x)^{(1+(I/2)*n)}*\operatorname{Hypergeometric2F1}\left[1+(I/2)*n, (I/2)*n, 2+(I/2)*n, (1-I*a*x)/2\right])/(a*(2*I-n))$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a_+ + b_+*x_+)^{(m_+ + 1)} / (b_+*(m_+ + 1)*(b_+/(b_+*c_+ - a_+*d_+))^{(n_+)}) * \operatorname{Hypergeometric2F1}[-n_+, m_+ + 1, m_+ + 2, (-d_+)*((a_+ + b_+*x_+)/(b_+*c_+ - a_+*d_+))], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x$
 $\&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] \mid\mid !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5169

$\operatorname{Int}\left[E^{(\operatorname{ArcTan}[(a_+)*(x_+)]*(n_+))}, x_Symbol\right] \rightarrow \operatorname{Int}\left[(1 - I*a*x)^{(I*(n/2))} / (1 + I*a*x)^{(I*(n/2))}, x\right] /;$ $\operatorname{FreeQ}\{a, n\}, x$ $\&\& !\operatorname{IntegerQ}[(I*n - 1)/2]$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} dx &= \int (1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}} dx \\ &= -\frac{2^{1-\frac{in}{2}}(1-iax)^{1+\frac{in}{2}} {}_2F_1\left(1+\frac{in}{2}, \frac{in}{2}; 2+\frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a(2i-n)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 56, normalized size = 0.69

$$\frac{4e^{(2i+n)\text{ArcTan}(ax)} {}_2F_1\left(2, 1 - \frac{in}{2}; 2 - \frac{in}{2}; -e^{2i\text{ArcTan}(ax)}\right)}{a(2i+n)}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(n*ArcTan[a*x]), x]``[Out] (4*E^((2*I + n)*ArcTan[a*x])*Hypergeometric2F1[2, 1 - (I/2)*n, 2 - (I/2)*n, -E^((2*I)*ArcTan[a*x])])/(a*(2*I + n))`**Maple [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(a*x)), x)``[Out] int(exp(n*arctan(a*x)), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x)), x, algorithm="maxima")``[Out] integrate(e^(n*arctan(a*x)), x)`**Fricas [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x)), x, algorithm="fricas")``[Out] integral(e^(n*arctan(a*x)), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x)),x)`

[Out] `Integral(exp(n*atan(a*x)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x)),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*atan(a*x)),x)`

[Out] `int(exp(n*atan(a*x)), x)`

3.340 $\int \frac{e^{n \operatorname{ArcTan}(ax)} x^3}{c+a^2cx^2} dx$

Optimal. Leaf size=131

$$\frac{e^{n \operatorname{ArcTan}(ax)}(2i+n-in^2)}{2a^4cn} - \frac{e^{n \operatorname{ArcTan}(ax)}nx}{2a^3c} + \frac{e^{n \operatorname{ArcTan}(ax)}x^2}{2a^2c} + \frac{ie^{n \operatorname{ArcTan}(ax)}(-2+n^2)}{a^4cn} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; -e^{2i \operatorname{ArcTan}(ax)}\right)$$

[Out] $1/2*\exp(n*\arctan(a*x))*(2*I+n-I*n^2)/a^4/c/n-1/2*\exp(n*\arctan(a*x))*n*x/a^3/c+1/2*\exp(n*\arctan(a*x))*x^2/a^2/c+I*\exp(n*\arctan(a*x))*(n^2-2)*\operatorname{hypergeom}(1, -1/2*I*n), [1-1/2*I*n], -(1+I*a*x)^2/(a^2*x^2+1))/a^4/c/n$

Rubi [A]

time = 0.17, antiderivative size = 206, normalized size of antiderivative = 1.57, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5190, 102, 148, 71}

$$\frac{2^{-1-\frac{in}{2}}(2-n^2)(1-iax)^{1+\frac{in}{2}} {}_2F_1\left(\frac{in}{2}+1, \frac{in}{2}+1; \frac{in}{2}+2; \frac{1}{2}(1-iax)\right)}{a^4c(2+in)} + \frac{i(1+iax)^{-\frac{in}{2}}(ian^2x-n^2-in+2)(1-iax)^{\frac{in}{2}}}{2a^4cn} + \frac{x^2(1+iax)^{-\frac{in}{2}}(1-iax)^{\frac{in}{2}}}{2a^2c}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^3)/(c + a^2*c*x^2), x]$

[Out] $(x^2*(1 - I*a*x)^{((I/2)*n)})/(2*a^2*c*(1 + I*a*x)^{((I/2)*n)}) + ((I/2)*(1 - I*a*x)^{((I/2)*n)}*(2 - I*n - n^2 + I*a*n^2*x))/(a^4*c*n*(1 + I*a*x)^{((I/2)*n)}) + (2^{(-1 - (I/2)*n)}*(2 - n^2)*(1 - I*a*x)^{(1 + (I/2)*n)}*\operatorname{Hypergeometric2F1}[1 + (I/2)*n, 1 + (I/2)*n, 2 + (I/2)*n, (1 - I*a*x)/2])/(a^4*c*(2 + I*n))$

Rule 71

$\operatorname{Int}[(a + b*x)^m * ((c + d*x)^n), x_Symbol] := \operatorname{Simp}[(a + b*x)^{m+1}/(b*(m+1)*(b*(b*c - a*d))^n) * \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*((a + b*x)/(b*c - a*d))], x] /;$ $\operatorname{FreeQ}[\{a, b, c, d, m, n\}, x]$ $\&\& \operatorname{NeQ}[b*c - a*d, 0]$ $\&\& \operatorname{!IntegerQ}[m]$ $\&\& \operatorname{!IntegerQ}[n]$ $\&\& \operatorname{GtQ}[b/(b*c - a*d), 0]$ $\&\& (\operatorname{RationalQ}[m] \mid \mid \operatorname{!RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0])$

Rule 102

$\operatorname{Int}[(a + b*x)^m * ((c + d*x)^n * ((e + f*x)^p)), x_Symbol] := \operatorname{Simp}[b*(a + b*x)^{m-1}*(c + d*x)^{n+1}*(e + f*x)^{p+1}/(d*f*(m+n+p+1)), x] + \operatorname{Dist}[1/(d*f*(m+n+p+1)), \operatorname{Int}[(a + b*x)^{m-2}*(c + d*x)^n*(e + f*x)^p * \operatorname{Simp}[a^2*d*f*(m+n+p+1) - b*(b*c*e*(m-1) + a*(d*e*(n+1) + c*f*(p+1))) + b*(a*d*f*(2*m+n+p) - b*(d*e*(m+n) + c*f*(m+p))]*x, x], x] /;$ $\operatorname{FreeQ}[\{a, b, c, d, e, f, n, p\}, x]$ $\&\& \operatorname{GtQ}[m, 1]$ $\&\& \operatorname{NeQ}[m+n+p+1, 0]$ $\&\& \operatorname{IntegerQ}[m]$

Rule 148

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b^2*d*e*g - a^2*d*f*h*m - a*b*(d
*(f*g + e*h) - c*f*h*(m + 1)) + b*f*h*(b*c - a*d)*(m + 1)*x*(a + b*x)^(m +
1)*((c + d*x)^(n + 1)/(b^2*d*(b*c - a*d)*(m + 1))), x] + Dist[(a*d*f*h*m +
b*(d*(f*g + e*h) - c*f*h*(m + 2)))/(b^2*d), Int[(a + b*x)^(m + 1)*(c + d*x
)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && EqQ[m + n + 2, 0
] && NeQ[m, -1] && (SumSimplerQ[m, 1] || !SumSimplerQ[n, 1])
```

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_
Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(
n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ
[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x^3}{c + a^2 c x^2} dx &= \frac{\int x^3 (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} dx}{c} \\ &= \frac{x^2 (1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{2a^2 c} + \frac{\int x (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} (-2 - anx) dx}{2a^2 c} \\ &= \frac{x^2 (1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{2a^2 c} + \frac{i(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}} (2 - in - n^2 + ian^2 x)}{2a^4 cn} - \frac{i(2 - in - n^2 + ian^2 x)}{2a^4 cn} \\ &= \frac{x^2 (1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{2a^2 c} + \frac{i(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}} (2 - in - n^2 + ian^2 x)}{2a^4 cn} + \frac{2 - in - n^2 + ian^2 x}{2a^4 cn} \end{aligned}$$

Mathematica [A]

time = 0.09, size = 141, normalized size = 1.08

$$\frac{(1 - iax)^{\frac{in}{2}} \left(\frac{(1 + iax)^{-\frac{in}{2}} (2i + n + a^2 n x^2 - n^2 (i + ax))}{n} + \frac{2^{-\frac{in}{2}} (-2 + n^2) (i + ax) {}_2F_1\left(1 + \frac{in}{2}, 1 + \frac{in}{2}; 2 + \frac{in}{2}; \frac{1}{2} (1 - iax)\right)}{-2i + n} \right)}{2a^4 c}$$

Antiderivative was successfully verified.

```
[In] Integrate[(E^(n*ArcTan[a*x])*x^3)/(c + a^2*c*x^2), x]
```

```
[Out] ((1 - I*a*x)^((I/2)*n)*((2*I + n + a^2*n*x^2 - n^2*(I + a*x))/(n*(1 + I*a*x)
)^((I/2)*n)) + ((-2 + n^2)*(I + a*x)*Hypergeometric2F1[1 + (I/2)*n, 1 + (I/
2)*n, 2 + (I/2)*n, (1 - I*a*x)/2])/(2^((I/2)*n)*(-2*I + n)))/(2*a^4*c)
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^3}{a^2 c x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c),x)`

[Out] `int(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c),x, algorithm="maxima")`

[Out] `integrate(x^3*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c),x, algorithm="fricas")`

[Out] `integral(x^3*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{x^3 e^{n \operatorname{atan}(ax)}}{a^2 x^2 + 1} dx}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))*x**3/(a**2*c*x**2+c),x)`

[Out] `Integral(x**3*exp(n*atan(a*x))/(a**2*x**2 + 1), x)/c`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^3 e^{n \operatorname{atan}(ax)}}{c a^2 x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^3*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)`

[Out] `int((x^3*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)`

3.341 $\int \frac{e^{n \operatorname{ArcTan}(ax)} x^2}{c + a^2 c x^2} dx$

Optimal. Leaf size=164

$$-\frac{(1+in)(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{a^3cn} + \frac{x(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{a^2c} + \frac{i2^{1-\frac{in}{2}}(1-iax)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; 1 + \frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a^3c}$$

[Out] $-(1+I*n)*(1-I*a*x)^{(1/2*I*n)}/a^3/c/n/((1+I*a*x)^{(1/2*I*n)})+x*(1-I*a*x)^{(1/2*I*n)}/a^2/c/((1+I*a*x)^{(1/2*I*n)})+I*2^{(1-1/2*I*n)}*(1-I*a*x)^{(1/2*I*n)}*\operatorname{hypergeom}([1/2*I*n, 1/2*I*n], [1+1/2*I*n], 1/2-1/2*I*a*x)/a^3/c$

Rubi [A]

time = 0.10, antiderivative size = 164, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5190, 92, 80, 71}

$$\frac{i2^{1-\frac{in}{2}}(1-iax)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; \frac{in}{2} + 1; \frac{1}{2}(1-iax)\right)}{a^3c} - \frac{(1+in)(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{a^3cn} + \frac{x(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{a^2c}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^2)/(c + a^2 * c * x^2), x]$

[Out] $-\left(\frac{(1+I*n)*(1-I*a*x)^{(I/2)*n}}{a^3*c*n*(1+I*a*x)^{(I/2)*n}}\right) + (x*(1-I*a*x)^{(I/2)*n})/(a^2*c*(1+I*a*x)^{(I/2)*n}) + (I*2^{(1-(I/2)*n)}*(1-I*a*x)^{(I/2)*n}*\operatorname{Hypergeometric2F1}[(I/2)*n, (I/2)*n, 1+(I/2)*n, (1-I*a*x)/2])/(a^3*c)$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)(x_+))^{(m_+)}*((c_+ + (d_+)(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b*(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x \&\& \operatorname{NeQ}\{b*c - a*d, 0\} \&\& \operatorname{IntegerQ}\{m\} \&\& \operatorname{IntegerQ}\{n\} \&\& \operatorname{GtQ}\{b/(b*c - a*d), 0\} \&\& (\operatorname{RationalQ}\{m\} \mid \mid \operatorname{IntegerQ}\{n\} \&\& \operatorname{GtQ}\{-d/(b*c - a*d), 0\})$

Rule 80

$\operatorname{Int}[(a_+ + (b_+)(x_+))*((c_+ + (d_+)(x_+))^{(n_+)})*((e_+ + (f_+)(x_+))^{(p_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(-b*e - a*f)*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(f*(p + 1)*(c*f - d*e))), x] - \operatorname{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(f*(p + 1)*(c*f - d*e)), \operatorname{Int}[(c + d*x)^n*(e + f*x)^p \operatorname{Simplify}[p + 1], x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x \&\& \operatorname{RationalQ}\{p\} \&\& \operatorname{SumSimplerQ}\{p, 1\}$

Rule 92

```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x^2}{c + a^2 c x^2} dx &= \frac{\int x^2 (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} dx}{c} \\ &= \frac{x(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^2 c} + \frac{\int (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} (-1 - anx) dx}{a^2 c} \\ &= -\frac{(1 + in)(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^3 cn} + \frac{x(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^2 c} + \frac{(in) \int (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} dx}{a^2} \\ &= -\frac{(1 + in)(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^3 cn} + \frac{x(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^2 c} + \frac{i2^{1 - \frac{in}{2}} (1 - iax)^{\frac{in}{2}}}{2} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 77, normalized size = 0.47

$$\frac{e^{n \operatorname{ArcTan}(ax)} \left(-2i - n + 4e^{2i \operatorname{ArcTan}(ax)} n {}_2F_1 \left(2, 1 - \frac{in}{2}; 2 - \frac{in}{2}; -e^{2i \operatorname{ArcTan}(ax)} \right) \right)}{a^3 cn (2i + n)}$$

Antiderivative was successfully verified.

```
[In] Integrate[(E^(n*ArcTan[a*x])*x^2)/(c + a^2*c*x^2), x]
```

```
[Out] (E^(n*ArcTan[a*x])*(-2*I - n + 4*E^((2*I)*ArcTan[a*x])*n*Hypergeometric2F1[2, 1 - (I/2)*n, 2 - (I/2)*n, -E^((2*I)*ArcTan[a*x])]))/(a^3*c*n*(2*I + n))
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^2}{a^2 c x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c),x)`

[Out] `int(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c),x, algorithm="maxima")`

[Out] `integrate(x^2*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c),x, algorithm="fricas")`

[Out] `integral(x^2*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{x^2 e^{n \operatorname{atan}(ax)}}{a^2 x^2 + 1} dx}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))*x**2/(a**2*c*x**2+c),x)`

[Out] `Integral(x**2*exp(n*atan(a*x))/(a**2*x**2 + 1), x)/c`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^2 e^{n \operatorname{atan}(ax)}}{c a^2 x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)`

[Out] `int((x^2*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)`

$$3.342 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x}{c + a^2 cx^2} dx$$

Optimal. Leaf size=122

$$\frac{i(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{a^2cn} - \frac{i2^{1-\frac{in}{2}}(1-iax)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; 1 + \frac{in}{2}; \frac{1}{2}(1-iax)\right)}{a^2cn}$$

[Out] I*(1-I*a*x)^(1/2*I*n)/a^2/c/n/((1+I*a*x)^(1/2*I*n))-I*2^(1-1/2*I*n)*(1-I*a*x)^(1/2*I*n)*hypergeom([1/2*I*n, 1/2*I*n],[1+1/2*I*n],1/2-1/2*I*a*x)/a^2/c/n

Rubi [A]

time = 0.06, antiderivative size = 122, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.136$, Rules used = {5190, 80, 71}

$$\frac{i(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{a^2cn} - \frac{i2^{1-\frac{in}{2}}(1-iax)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; \frac{in}{2} + 1; \frac{1}{2}(1-iax)\right)}{a^2cn}$$

Antiderivative was successfully verified.

[In] Int[(E^(n*ArcTan[a*x])*x)/(c + a^2*c*x^2), x]

[Out] (I*(1 - I*a*x)^((I/2)*n))/(a^2*c*n*(1 + I*a*x)^((I/2)*n)) - (I*2^(1 - (I/2)*n)*(1 - I*a*x)^((I/2)*n)*Hypergeometric2F1[(I/2)*n, (I/2)*n, 1 + (I/2)*n, (1 - I*a*x)/2])/(a^2*c*n)

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d))^n))*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 80

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimplerQ[p, 1]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^((m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(

$n/2$)), x], x] /; FreeQ[{ a , c , d , m , n , p }, x] && EqQ[d , a^2*c] && (IntegerQ[p] || GtQ[c , 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x}{c + a^2 c x^2} dx &= \frac{\int x(1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} dx}{c} \\ &= \frac{i(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^2 cn} - \frac{i \int (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-\frac{in}{2}} dx}{ac} \\ &= \frac{i(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}}}{a^2 cn} - \frac{i 2^{1 - \frac{in}{2}} (1 - iax)^{\frac{in}{2}} {}_2F_1\left(\frac{in}{2}, \frac{in}{2}; 1 + \frac{in}{2}; \frac{1}{2}(1 - iax)\right)}{a^2 cn} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 104, normalized size = 0.85

$$\frac{i e^{n \operatorname{ArcTan}(ax)} \left(\frac{{}_2F_1\left(1, 1 - \frac{in}{2}; 2 - \frac{in}{2}; -e^{2i \operatorname{ArcTan}(ax)}\right)}{2i+n} - \frac{{}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; -e^{2i \operatorname{ArcTan}(ax)}\right)}{n} \right)}{a^2 c}$$

Antiderivative was successfully verified.

[In] Integrate[(E^(n*ArcTan[a*x]))*x]/(c + a^2*c*x^2), x]

[Out] ((-I)*E^(n*ArcTan[a*x])*((E^((2*I)*ArcTan[a*x])*Hypergeometric2F1[1, 1 - (I/2)*n, 2 - (I/2)*n, -E^((2*I)*ArcTan[a*x])])/(2*I + n) - Hypergeometric2F1[1, (-1/2*I)*n, 1 - (I/2)*n, -E^((2*I)*ArcTan[a*x])]/n))/(a^2*c)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x}{a^2 c x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x/(a^2*c*x^2+c), x)

[Out] int(exp(n*arctan(a*x))*x/(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x/(a^2*c*x^2+c),x, algorithm="maxima")

[Out] integrate(x*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x/(a^2*c*x^2+c),x, algorithm="fricas")

[Out] integral(x*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x e^{n \operatorname{atan}(ax)}}{a^2 x^2 + 1} dx$$

c

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*x/(a**2*c*x**2+c),x)

[Out] Integral(x*exp(n*atan(a*x))/(a**2*x**2 + 1), x)/c

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x/(a^2*c*x^2+c),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x e^{n \operatorname{atan}(ax)}}{c a^2 x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*exp(n*atan(a*x)))/(c + a^2*c*x^2),x)

[Out] int((x*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)

$$3.343 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{c+a^2cx^2} dx$$

Optimal. Leaf size=18

$$\frac{e^{n \operatorname{ArcTan}(ax)}}{acn}$$

[Out] exp(n*arctan(a*x))/a/c/n

Rubi [A]

time = 0.02, antiderivative size = 18, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.048$, Rules used = {5179}

$$\frac{e^{n \operatorname{ArcTan}(ax)}}{acn}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a*x])/(c + a^2*c*x^2),x]

[Out] E^(n*ArcTan[a*x])/(a*c*n)

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\int \frac{e^{n \tan^{-1}(ax)}}{c + a^2cx^2} dx = \frac{e^{n \tan^{-1}(ax)}}{acn}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.01, size = 42, normalized size = 2.33

$$\frac{(1 - iax)^{\frac{in}{2}}(1 + iax)^{-\frac{in}{2}}}{acn}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(c + a^2*c*x^2),x]

[Out] (1 - I*a*x)^((I/2)*n)/(a*c*n*(1 + I*a*x)^((I/2)*n))

Maple [A]

time = 0.07, size = 18, normalized size = 1.00

method	result	size
gospers	$\frac{e^{n \arctan(ax)}}{acn}$	18
risch	$\frac{(-iax+1)^{\frac{in}{2}}(iax+1)^{-\frac{in}{2}}}{can}$	35

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*arctan(a*x))/(a^2*c*x^2+c),x,method=_RETURNVERBOSE)`

[Out] `exp(n*arctan(a*x))/a/c/n`

Maxima [A]

time = 0.49, size = 17, normalized size = 0.94

$$\frac{e^{(n \arctan(ax))}}{acn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="maxima")`

[Out] `e^(n*arctan(a*x))/(a*c*n)`

Fricas [A]

time = 1.83, size = 17, normalized size = 0.94

$$\frac{e^{(n \arctan(ax))}}{acn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="fricas")`

[Out] `e^(n*arctan(a*x))/(a*c*n)`

Sympy [B] Leaf count of result is larger than twice the leaf count of optimal. 26 vs. 2(12) = 24.

time = 0.60, size = 26, normalized size = 1.44

$$\begin{cases} \frac{x}{c} & \text{for } a = 0 \wedge n = 0 \\ \frac{\operatorname{atan}(ax)}{ac} & \text{for } n = 0 \\ \frac{x}{c} & \text{for } a = 0 \\ \frac{e^{n \operatorname{atan}(ax)}}{acn} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))/(a**2*c*x**2+c),x)`

[Out] Piecewise((x/c, Eq(a, 0) & Eq(n, 0)), (atan(a*x)/(a*c), Eq(n, 0)), (x/c, Eq(a, 0)), (exp(n*atan(a*x))/(a*c*n), True))

Giac [A]

time = 0.41, size = 17, normalized size = 0.94

$$\frac{e^{(n \arctan(ax))}}{acn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c),x, algorithm="giac")

[Out] e^(n*arctan(a*x))/(a*c*n)

Mupad [B]

time = 0.63, size = 17, normalized size = 0.94

$$\frac{e^{n \operatorname{atan}(ax)}}{acn}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(c + a^2*c*x^2),x)

[Out] exp(n*atan(a*x))/(a*c*n)

3.344 $\int \frac{e^{n \operatorname{ArcTan}(ax)}}{x(c+a^2cx^2)} dx$

Optimal. Leaf size=65

$$\frac{ie^{n \operatorname{ArcTan}(ax)}}{cn} - \frac{2ie^{n \operatorname{ArcTan}(ax)} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; e^{2i \operatorname{ArcTan}(ax)}\right)}{cn}$$

[Out] $I*\exp(n*\arctan(a*x))/c/n-2*I*\exp(n*\arctan(a*x))*\operatorname{hypergeom}([1, -1/2*I*n], [1-1/2*I*n], (1+I*a*x)^2/(a^2*x^2+1))/c/n$

Rubi [A]

time = 0.07, antiderivative size = 122, normalized size of antiderivative = 1.88, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5190, 98, 133}

$$\frac{i(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{cn} - \frac{2i(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; \frac{iax+1}{1-iax}\right)}{cn}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a*x])}/(x*(c + a^2*c*x^2)), x]$

[Out] $(I*(1 - I*a*x)^{((I/2)*n)})/(c*n*(1 + I*a*x)^{((I/2)*n)}) - ((2*I)*(1 - I*a*x)^{((I/2)*n)}*\operatorname{Hypergeometric2F1}[1, (-1/2*I)*n, 1 - (I/2)*n, (1 + I*a*x)/(1 - I*a*x)])/(c*n*(1 + I*a*x)^{((I/2)*n)})$

Rule 98

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[b*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + \operatorname{Dist}[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p, x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x\} \&\& \operatorname{EqQ}[\operatorname{Simplify}[m + n + p + 3], 0] \&\& (\operatorname{LtQ}[m, -1] \parallel \operatorname{SumSimplerQ}[m, 1])$

Rule 133

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(b*c - a*d)^n*((a + b*x)^{(m + 1)})/((m + 1)*(b*e - a*f)^{(n + 1)}*(e + f*x)^{(m + 1)})*\operatorname{Hypergeometric2F1}[m + 1, -n, m + 2, (-(d*e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, m, p\}, x\} \&\& \operatorname{EqQ}[m + n + p + 2, 0] \&\& \operatorname{ILtQ}[n, 0] \&\& (\operatorname{SumSimplerQ}[m, 1] \parallel \operatorname{!SumSimplerQ}[p, 1]) \&\& \operatorname{!ILtQ}[m, 0]$

Rule 5190


```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_
Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(
n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ
[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{x(c + a^2cx^2)} dx &= \frac{\int \frac{(1-iax)^{-1+\frac{in}{2}}(1+iax)^{-1-\frac{in}{2}}}{x} dx}{c} \\ &= \frac{i(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{cn} + \frac{\int \frac{(1-iax)^{\frac{in}{2}}(1+iax)^{-1-\frac{in}{2}}}{x} dx}{c} \\ &= \frac{i(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{cn} - \frac{2(1-iax)^{1+\frac{in}{2}}(1+iax)^{-1-\frac{in}{2}} {}_2F_1\left(1, 1 + \frac{in}{2}; 2 + \frac{in}{2}, \frac{1-iax}{1+iax}\right)}{c(2+in)} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 120, normalized size = 1.85

$$\frac{(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}} \left((2+in)(-i+ax) + 2(n-ianx) {}_2F_1\left(1, 1 + \frac{in}{2}; 2 + \frac{in}{2}, \frac{i+ax}{i-ax}\right) \right)}{cn(-2i+n)(-i+ax)}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^(n*ArcTan[a*x])/(x*(c + a^2*c*x^2)), x]
```

```
[Out] ((1 - I*a*x)^((I/2)*n)*((2 + I*n)*(-I + a*x) + 2*(n - I*a*n*x)*Hypergeometr
ic2F1[1, 1 + (I/2)*n, 2 + (I/2)*n, (I + a*x)/(I - a*x)])/(c*n*(-2*I + n)*(
1 + I*a*x)^((I/2)*n)*(-I + a*x))
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{x(a^2cx^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(n*arctan(a*x))/x/(a^2*c*x^2+c), x)
```

```
[Out] int(exp(n*arctan(a*x))/x/(a^2*c*x^2+c), x)
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x/(a^2*c*x^2+c),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/((a^2*c*x^2 + c)*x), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x/(a^2*c*x^2+c),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/(a^2*c*x^3 + c*x), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{e^{n \operatorname{atan}(ax)}}{a^2 x^3 + x} dx}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/x/(a**2*c*x**2+c),x)

[Out] Integral(exp(n*atan(a*x))/(a**2*x**3 + x), x)/c

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x/(a^2*c*x^2+c),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x (ca^2 x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(x*(c + a^2*c*x^2)),x)

[Out] int(exp(n*atan(a*x))/(x*(c + a^2*c*x^2)), x)

$$3.345 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{x^2(c+a^2cx^2)} dx$$

Optimal. Leaf size=90

$$\frac{iae^{n \operatorname{ArcTan}(ax)}(i+n)}{cn} - \frac{e^{n \operatorname{ArcTan}(ax)}}{cx} - \frac{2iae^{n \operatorname{ArcTan}(ax)} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; -1 + \frac{2i}{i+ax}\right)}{c}$$

[Out] $I*a*\exp(n*\arctan(a*x))*(I+n)/c/n-\exp(n*\arctan(a*x))/c/x-2*I*a*\exp(n*\arctan(a*x))*\operatorname{hypergeom}([1, -1/2*I*n], [1-1/2*I*n], -1+2*I/(I+a*x))/c$

Rubi [A]

time = 0.09, antiderivative size = 166, normalized size of antiderivative = 1.84, number of steps used = 5, number of rules used = 5, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.208$, Rules used = {5190, 105, 160, 12, 133}

$$\frac{2ia(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; \frac{iax+1}{1-iax}\right)}{c} - \frac{a(1-in)(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{cn} - \frac{(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{cx}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a*x])}/(x^2*(c+a^2*c*x^2)), x]$

[Out] $-((a*(1-I*n)*(1-I*a*x)^{((I/2)*n)})/(c*n*(1+I*a*x)^{((I/2)*n)})) - (1-I*a*x)^{((I/2)*n)}/(c*x*(1+I*a*x)^{((I/2)*n)}) - ((2*I)*a*(1-I*a*x)^{((I/2)*n)})*\operatorname{Hypergeometric2F1}[1, (-1/2*I)*n, 1 - (I/2)*n, (1+I*a*x)/(1-I*a*x)]/(c*(1+I*a*x)^{((I/2)*n)})$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] \rightarrow \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \ \&\& \ !\operatorname{Match} Q[u, (b_)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 105

$\operatorname{Int}[(a_*)(x_)+(b_*)(x_))^{(m_)*((c_)+(d_)*(x_))^{(n_)*((e_)+(f_)*(x_))^{(p_)}}, x_Symbol] \rightarrow \operatorname{Simp}[b*(a+b*x)^{(m+1)}*(c+d*x)^{(n+1)}*((e+f*x)^{(p+1)}/((m+1)*(b*c-a*d)*(b*e-a*f))), x] + \operatorname{Dist}[1/((m+1)*(b*c-a*d)*(b*e-a*f)), \operatorname{Int}[(a+b*x)^{(m+1)}*(c+d*x)^n*(e+f*x)^p*\operatorname{Simp}[a*d*f*(m+1)-b*(d*e*(m+n+2)+c*f*(m+p+2))-b*d*f*(m+n+p+3)*x, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \ \&\& \ \operatorname{ILtQ}[m, -1] \ \&\& \ (\operatorname{Integer} Q[n] \ || \ \operatorname{Integers} Q[2*n, 2*p] \ || \ \operatorname{ILtQ}[m+n+p+3, 0])$

Rule 133

$\operatorname{Int}[(a_*)(x_)+(b_*)(x_))^{(m_)*((c_)+(d_)*(x_))^{(n_)*((e_)+(f_)*(x_))^{(p_)}}, x_Symbol] \rightarrow \operatorname{Simp}[(b*c-a*d)^n*((a+b*x)^{(m+1)}/((m+1)*(b*e-a*f)^{(n+1)}*(e+f*x)^{(m+1)}))*\operatorname{Hypergeometric2F1}[m+1, -n, m+2, -(d*$

```
e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x))), x] /; FreeQ[{a, b, c, d, e,
f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1]
|| !SumSimplerQ[p, 1]) && !ILtQ[m, 0]
```

Rule 160

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))
)^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1)
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^(n + 1)*((e + f*x)^(p + 1)*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m + n + p + 2,
0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || (!NeQ[n, -1] && SumSimplerQ[n,
1]) && !(NeQ[p, -1] && SumSimplerQ[p, 1]))
```

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_)^(n_.)]*(x_)^(m_.))*((c_.) + (d_.)*(x_)^2)^(p_.), x_
Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(
n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ
[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{x^2 (c + a^2 cx^2)} dx &= \int \frac{(1-iax)^{-1+\frac{in}{2}} (1+iax)^{-1-\frac{in}{2}}}{x^2 c} dx \\ &= -\frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cx} - \int \frac{(1-iax)^{-1+\frac{in}{2}} (1+iax)^{-1-\frac{in}{2}} (-an+a^2x)}{x c} dx \\ &= -\frac{a(1-in)(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cn} - \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cx} + \int \frac{a^2 n^2 (1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{x} dx \\ &= -\frac{a(1-in)(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cn} - \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cx} + \frac{(an) \int \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{x} dx}{c} \\ &= -\frac{a(1-in)(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cn} - \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{cx} - \frac{2an(1-iax)^{1+\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{c} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 142, normalized size = 1.58

$$\frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}} ((-2i+n)(1+iax)(iax+n(i+ax)) + 2an^2x(1-iax) {}_2F_1(1, 1+\frac{in}{2}; 2+\frac{in}{2}, \frac{i+ax}{i-ax}))}{cn(-2i+n)x(-i+ax)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(x^2*(c + a^2*c*x^2)),x]

[Out] $((1 - I*a*x)^{((I/2)*n)}*((-2*I + n)*(1 + I*a*x)*(I*a*x + n*(I + a*x)) + 2*a*n^2*x*(1 - I*a*x)*Hypergeometric2F1[1, 1 + (I/2)*n, 2 + (I/2)*n, (I + a*x)/(I - a*x)]))/(c*n*(-2*I + n)*x*(1 + I*a*x)^{((I/2)*n)}*(-I + a*x))$

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{x^2 (a^2 c x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c),x)

[Out] int(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/((a^2*c*x^2 + c)*x^2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/(a^2*c*x^4 + c*x^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{a^2 x^4 + x^2} dx$$

c

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/x**2/(a**2*c*x**2+c),x)

[Out] `Integral(exp(n*atan(a*x))/(a**2*x**4 + x**2), x)/c`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x^2 (ca^2 x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*atan(a*x))/(x^2*(c + a^2*c*x^2)),x)`

[Out] `int(exp(n*atan(a*x))/(x^2*(c + a^2*c*x^2)), x)`

$$3.346 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{x^3(c+a^2cx^2)} dx$$

Optimal. Leaf size=126

$$\frac{ia^2 e^{n \operatorname{ArcTan}(ax)} (-2 + in + n^2)}{2cn} - \frac{e^{n \operatorname{ArcTan}(ax)}}{2cx^2} - \frac{ae^{n \operatorname{ArcTan}(ax)} n}{2cx} - \frac{ia^2 e^{n \operatorname{ArcTan}(ax)} (-2 + n^2)}{cn} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; e^2\right)$$

[Out] 1/2*I*a^2*exp(n*arctan(a*x))*(-2+I*n+n^2)/c/n-1/2*exp(n*arctan(a*x))/c/x^2-1/2*a*exp(n*arctan(a*x))*n/c/x-I*a^2*exp(n*arctan(a*x))*(n^2-2)*hypergeom([1, -1/2*I*n], [1-1/2*I*n], (1+I*a*x)^2/(a^2*x^2+1))/c/n

Rubi [A]

time = 0.13, antiderivative size = 233, normalized size of antiderivative = 1.85, number of steps used = 6, number of rules used = 6, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.250$, Rules used = {5190, 105, 156, 160, 12, 133}

$$\frac{ia^2(2-n^2)(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}} {}_2F_1\left(1, -\frac{in}{2}; 1 - \frac{in}{2}; \frac{iax+1}{1-iax}\right)}{cn} - \frac{a^2(-in^2+n+2i)(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{2cn} - \frac{(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{2cx^2} - \frac{an(1-iax)^{\frac{in}{2}}(1+iax)^{-\frac{in}{2}}}{2cx}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a*x])/(x^3*(c + a^2*c*x^2)), x]

[Out] -1/2*(a^2*(2*I + n - I*n^2)*(1 - I*a*x)^((I/2)*n))/(c*n*(1 + I*a*x)^((I/2)*n)) - (1 - I*a*x)^((I/2)*n)/(2*c*x^2*(1 + I*a*x)^((I/2)*n)) - (a*n*(1 - I*a*x)^((I/2)*n))/(2*c*x*(1 + I*a*x)^((I/2)*n)) + (I*a^2*(2 - n^2)*(1 - I*a*x)^((I/2)*n)*Hypergeometric2F1[1, (-1/2*I)*n, 1 - (I/2)*n, (1 + I*a*x)/(1 - I*a*x)]/(c*n*(1 + I*a*x)^((I/2)*n))

Rule 12

Int[(a_)*(u_), x_Symbol] := Dist[a, Int[u, x], x] /; FreeQ[a, x] && !MatchQ[u, (b_)*(v_)] /; FreeQ[b, x]

Rule 105

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[b*(a + b*x)^(m + 1)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d*x)^n*(e + f*x)^p*Simp[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && ILtQ[m, -1] && (IntegerQ[n] || IntegersQ[2*n, 2*p] || ILtQ[m + n + p + 3, 0])

Rule 133

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] := Simp[(b*c - a*d)^n*((a + b*x)^(m + 1)/((m + 1)*(b*e -

```

a*f)^(n + 1)*(e + f*x)^(m + 1))*Hypergeometric2F1[m + 1, -n, m + 2, (-(d*
e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /; FreeQ[{a, b, c, d, e,
f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1]
|| !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

```

Rule 156

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]

```

Rule 160

```

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_
))^p_)*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[(b*g - a*h)*(a + b*x)^(m + 1
)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/((m + 1)*(b*c - a*d)*(b*e - a*f))),
x] + Dist[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m + 1)*(c + d
*x)^n*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m + 1) - (b*g
- a*h)*(d*e*(n + 1) + c*f*(p + 1)) - d*f*(b*g - a*h)*(m + n + p + 3)*x, x]
, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m + n + p + 2,
0] && NeQ[m, -1] && (SumSimplerQ[m, 1] || ( !(NeQ[n, -1] && SumSimplerQ[n,
1]) && !(NeQ[p, -1] && SumSimplerQ[p, 1])))

```

Rule 5190

```

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^m_)*((c_) + (d_.)*(x_)^2)^(p_.), x_
Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(
n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ
[p] || GtQ[c, 0])

```

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)}}{x^3 (c + a^2 cx^2)} dx &= \int \frac{(1-iax)^{-1+\frac{in}{2}} (1+iax)^{-1-\frac{in}{2}}}{x^3 c} dx \\
&= -\frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cx^2} - \int \frac{(1-iax)^{-1+\frac{in}{2}} (1+iax)^{-1-\frac{in}{2}} (-an+2a^2x)}{x^2 2c} dx \\
&= -\frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cx^2} - \frac{an(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cx} + \int \frac{(1-iax)^{-1+\frac{in}{2}} (1+iax)^{-1-\frac{in}{2}}}{x 2c} dx \\
&= -\frac{a^2(2i+n-in^2)(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cn} - \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cx^2} - \frac{an(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2c} \\
&= -\frac{a^2(2i+n-in^2)(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cn} - \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cx^2} - \frac{an(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2c} \\
&= -\frac{a^2(2i+n-in^2)(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cn} - \frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2cx^2} - \frac{an(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}}}{2c}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 174, normalized size = 1.38

$$\frac{(1-iax)^{\frac{in}{2}} (1+iax)^{-\frac{in}{2}} (i(-2i+n)(-i+ax)(-2a^2x^2+an^2x(i+ax)+in(1+a^2x^2))+2a^2n(-2+n^2)x^2(1-iax) {}_2F_1(1, 1+\frac{in}{2}; 2+\frac{in}{2}, \frac{i+ax}{i-ax}))}{2cn(-2i+n)x^2(-i+ax)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(x^3*(c + a^2*c*x^2)), x]

[Out] (((1 - I*a*x)^((I/2)*n)*(I*(-2*I + n)*(-I + a*x)*(-2*a^2*x^2 + a*n^2*x*(I + a*x) + I*n*(1 + a^2*x^2)) + 2*a^2*n*(-2 + n^2)*x^2*(1 - I*a*x)*Hypergeometric2F1[1, 1 + (I/2)*n, 2 + (I/2)*n, (I + a*x)/(I - a*x)]))/(2*c*n*(-2*I + n)*x^2*(1 + I*a*x)^((I/2)*n)*(-I + a*x))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{x^3 (a^2 c x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c), x)

[Out] int(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/((a^2*c*x^2 + c)*x^3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/(a^2*c*x^5 + c*x^3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{e^{n \operatorname{atan}(ax)}}{a^2 x^5 + x^3} dx}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/x**3/(a**2*c*x**2+c),x)

[Out] Integral(exp(n*atan(a*x))/(a**2*x**5 + x**3), x)/c

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x^3 (c a^2 x^2 + c)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(x^3*(c + a^2*c*x^2)),x)

[Out] int(exp(n*atan(a*x))/(x^3*(c + a^2*c*x^2)), x)

$$3.347 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^4} dx$$

Optimal. Leaf size=181

$$\frac{720e^{n \operatorname{ArcTan}(ax)}}{ac^4n(4+n^2)(16+n^2)(36+n^2)} + \frac{e^{n \operatorname{ArcTan}(ax)}(n+6ax)}{ac^4(36+n^2)(1+a^2x^2)^3} + \frac{30e^{n \operatorname{ArcTan}(ax)}(n+4ax)}{ac^4(16+n^2)(36+n^2)(1+a^2x^2)^2} + \frac{360e^{n \operatorname{ArcTan}(ax)}(n+2ax)}{ac^4(4+n^2)(16+n^2)(36+n^2)(1+a^2x^2)}$$

[Out] 720*exp(n*arctan(a*x))/a/c^4/n/(n^2+36)/(n^4+20*n^2+64)+exp(n*arctan(a*x))*
(6*a*x+n)/a/c^4/(n^2+36)/(a^2*x^2+1)^3+30*exp(n*arctan(a*x))*(4*a*x+n)/a/c^4/
4/(n^2+16)/(n^2+36)/(a^2*x^2+1)^2+360*exp(n*arctan(a*x))*(2*a*x+n)/a/c^4/(n
^2+36)/(n^4+20*n^2+64)/(a^2*x^2+1)

Rubi [A]

time = 0.13, antiderivative size = 181, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 2, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.095$, Rules used = {5178, 5179}

$$\frac{360(2ax+n)e^{n \operatorname{ArcTan}(ax)}}{ac^4(n^2+4)(n^2+16)(n^2+36)(a^2x^2+1)} + \frac{30(4ax+n)e^{n \operatorname{ArcTan}(ax)}}{ac^4(n^2+16)(n^2+36)(a^2x^2+1)^2} + \frac{(6ax+n)e^{n \operatorname{ArcTan}(ax)}}{ac^4(n^2+36)(a^2x^2+1)^3} + \frac{720e^{n \operatorname{ArcTan}(ax)}}{ac^4n(n^2+4)(n^2+16)(n^2+36)}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a*x])/(c + a^2*c*x^2)^4, x]

[Out] (720*E^(n*ArcTan[a*x]))/(a*c^4*n*(4 + n^2)*(16 + n^2)*(36 + n^2)) + (E^(n*ArcTan[a*x])*(n + 6*a*x))/(a*c^4*(36 + n^2)*(1 + a^2*x^2)^3) + (30*E^(n*ArcTan[a*x])*(n + 4*a*x))/(a*c^4*(16 + n^2)*(36 + n^2)*(1 + a^2*x^2)^2) + (360*E^(n*ArcTan[a*x])*(n + 2*a*x))/(a*c^4*(4 + n^2)*(16 + n^2)*(36 + n^2)*(1 + a^2*x^2))

Rule 5178

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2)^(p_), x_Symbol] := Simp[(n - 2*a*(p + 1)*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x])/(a*c*(n^2 + 4*(p + 1)^2))), x] + Dist[2*(p + 1)*((2*p + 3)/(c*(n^2 + 4*(p + 1)^2))), Int[(c + d*x^2)^(p + 1)*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && LtQ[p, -1] && !IntegerQ[I*n] && NeQ[n^2 + 4*(p + 1)^2, 0] && IntegerQ[2*p]

Rule 5179

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))/((c_.) + (d_.)*(x_)^2), x_Symbol] := Simp[E^(n*ArcTan[a*x])/(a*c*n), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c]

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)}}{(c + a^2 cx^2)^4} dx &= \frac{e^{n \tan^{-1}(ax)}(n + 6ax)}{ac^4(36 + n^2)(1 + a^2 x^2)^3} + \frac{30 \int \frac{e^{n \tan^{-1}(ax)}}{(c + a^2 cx^2)^3} dx}{c(36 + n^2)} \\
&= \frac{e^{n \tan^{-1}(ax)}(n + 6ax)}{ac^4(36 + n^2)(1 + a^2 x^2)^3} + \frac{30e^{n \tan^{-1}(ax)}(n + 4ax)}{ac^4(16 + n^2)(36 + n^2)(1 + a^2 x^2)^2} + \frac{360 \int \frac{e^{n \tan^{-1}(ax)}}{(c + a^2 cx^2)^2} dx}{c^2(16 + n^2)(36 + n^2)} \\
&= \frac{e^{n \tan^{-1}(ax)}(n + 6ax)}{ac^4(36 + n^2)(1 + a^2 x^2)^3} + \frac{30e^{n \tan^{-1}(ax)}(n + 4ax)}{ac^4(16 + n^2)(36 + n^2)(1 + a^2 x^2)^2} + \frac{360e^{n \tan^{-1}(ax)}}{ac^4(4 + n^2)(16 + n^2)} \\
&= \frac{720e^{n \tan^{-1}(ax)}}{ac^4 n(4 + n^2)(16 + n^2)(36 + n^2)} + \frac{e^{n \tan^{-1}(ax)}(n + 6ax)}{ac^4(36 + n^2)(1 + a^2 x^2)^3} + \frac{30e^{n \tan^{-1}(ax)}(n + 4ax)}{ac^4(16 + n^2)(36 + n^2)}
\end{aligned}$$

Mathematica [C] Result contains complex when optimal does not.

time = 0.36, size = 165, normalized size = 0.91

$$\frac{e^{n \operatorname{ArcTan}(ax)}(n + 6ax) + \frac{30(c + a^2 cx^2) \left(e^{n \operatorname{ArcTan}(ax)} n(-2i + n)(2i + n)(n + 4ax) + 12(1 - iax)^{\frac{in}{2}} (1 + iax)^{-\frac{in}{2}} (-i + ax)(i + ax)(2 + n^2 + 2anx + 2a^2 x^2) \right)}{cn(64 + 20n^2 + n^4)}}{ac(36 + n^2)(c + a^2 cx^2)^3}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(c + a^2*c*x^2)^4, x]

[Out] (E^(n*ArcTan[a*x])*(n + 6*a*x) + (30*(c + a^2*c*x^2)*(E^(n*ArcTan[a*x]))*n*(-2*I + n)*(2*I + n)*(n + 4*a*x) + (12*(1 - I*a*x)^((I/2)*n)*(-I + a*x)*(I + a*x)*(2 + n^2 + 2*a*n*x + 2*a^2*x^2))/(1 + I*a*x)^((I/2)*n)))/(c*n*(64 + 20*n^2 + n^4)))/(a*c*(36 + n^2)*(c + a^2*c*x^2)^3)

Maple [A]

time = 0.09, size = 166, normalized size = 0.92

method	result
gospers	$\frac{(720a^6x^6 + 720a^5nx^5 + 360a^4n^2x^4 + 120a^3n^3x^3 + 2160a^4x^4 + 30a^2n^4x^2 + 1920a^3nx^3 + 6an^5x + 840a^2n^2x^2 + n^6 + 240an^3x + 2160a^2x^2 + 720a^2x^2 + 50n^4 + 1584anx + 544n^2 + 720) \exp(n \arctan(ax))}{(a^2x^2 + 1)^3 c^4 an(n^6 + 56n^4 + 784n^2 + 2304)}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^4, x, method=_RETURNVERBOSE)

[Out] (720*a^6*x^6+720*a^5*n*x^5+360*a^4*n^2*x^4+120*a^3*n^3*x^3+2160*a^4*x^4+30*a^2*n^4*x^2+1920*a^3*n*x^3+6*a*n^5*x+840*a^2*n^2*x^2+n^6+240*a*n^3*x+2160*a^2*x^2+50*n^4+1584*a*n*x+544*n^2+720)*exp(n*arctan(a*x))/(a^2*x^2+1)^3/c^4/a/n/(n^6+56*n^4+784*n^2+2304)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="maxima")``[Out] integrate(e^(n*arctan(a*x))/(a^2*c*x^2 + c)^4, x)`**Fricas [A]**

time = 2.79, size = 298, normalized size = 1.65

$$\frac{(720 a^6 x^6 + 720 a^5 n x^5 + n^6 + 360 (a^4 n^2 + 6 a^4) x^4 + 50 n^4 + 120 (a^3 n^3 + 16 a^3 n) x^3 + 30 (a^2 n^4 + 28 a^2 n^2 + 72 a^2) x^2 + 544 n^2 + 6 (a n^5 + 40 a n^3 + 264 a n) x + 720) e^{(n \arctan(ax))}}{a^4 n^7 + 56 a c^4 n^5 + 784 a c^4 n^3 + (a^7 c^4 n^7 + 56 a^7 c^4 n^5 + 784 a^7 c^4 n^3 + 2304 a^7 c^4 n) x^6 + 2304 a c^4 n + 3 (a^5 c^4 n^7 + 56 a^5 c^4 n^5 + 784 a^5 c^4 n^3 + 2304 a^5 c^4 n) x^4 + 3 (a^3 c^4 n^7 + 56 a^3 c^4 n^5 + 784 a^3 c^4 n^3 + 2304 a^3 c^4 n) x^2}$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="fricas")`

```
[Out] (720*a^6*x^6 + 720*a^5*n*x^5 + n^6 + 360*(a^4*n^2 + 6*a^4)*x^4 + 50*n^4 + 1
20*(a^3*n^3 + 16*a^3*n)*x^3 + 30*(a^2*n^4 + 28*a^2*n^2 + 72*a^2)*x^2 + 544*
n^2 + 6*(a*n^5 + 40*a*n^3 + 264*a*n)*x + 720)*e^(n*arctan(a*x))/(a*c^4*n^7
+ 56*a*c^4*n^5 + 784*a*c^4*n^3 + (a^7*c^4*n^7 + 56*a^7*c^4*n^5 + 784*a^7*c^
4*n^3 + 2304*a^7*c^4*n)*x^6 + 2304*a*c^4*n + 3*(a^5*c^4*n^7 + 56*a^5*c^4*n^
5 + 784*a^5*c^4*n^3 + 2304*a^5*c^4*n)*x^4 + 3*(a^3*c^4*n^7 + 56*a^3*c^4*n^5
+ 784*a^3*c^4*n^3 + 2304*a^3*c^4*n)*x^2)
```

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*atan(a*x))/(a**2*c*x**2+c)**4,x)``[Out] Timed out`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^4,x, algorithm="giac")``[Out] sage0*x`

Mupad [B]

time = 0.87, size = 281, normalized size = 1.55

$$e^{n \operatorname{atan}(ax)} \left(\frac{720 x^5}{a^2 c^4 (n^6 + 56 n^4 + 784 n^2 + 2304)} + \frac{n^6 + 50 n^4 + 544 n^2 + 720}{a^7 c^4 n (n^6 + 56 n^4 + 784 n^2 + 2304)} + \frac{720 x^6}{a c^4 n (n^6 + 56 n^4 + 784 n^2 + 2304)} + \frac{6 x (n^4 + 40 n^2 + 264)}{a^6 c^4 (n^6 + 56 n^4 + 784 n^2 + 2304)} + \frac{120 x^3 (n^2 + 16)}{a^3 c^4 (n^6 + 56 n^4 + 784 n^2 + 2304)} + \frac{360 x^4 (n^2 + 6)}{a^5 c^4 n (n^6 + 56 n^4 + 784 n^2 + 2304)} + \frac{30 x^2 (n^4 + 28 n^2 + 72)}{a^5 c^4 n (n^6 + 56 n^4 + 784 n^2 + 2304)} \right) \frac{1}{\frac{1}{a^6} + x^6 + \frac{3x^4}{a^2} + \frac{3x^2}{a^4}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*atan(a*x))/(c + a^2*c*x^2)^4,x)`

[Out] $(\exp(n \operatorname{atan}(ax)) * ((720x^5)/(a^2c^4(784n^2 + 56n^4 + n^6 + 2304)) + (544n^2 + 50n^4 + n^6 + 720)/(a^7c^4n(784n^2 + 56n^4 + n^6 + 2304)) + (720x^6)/(ac^4n(784n^2 + 56n^4 + n^6 + 2304)) + (6*x*(40n^2 + n^4 + 264))/(a^6c^4(784n^2 + 56n^4 + n^6 + 2304)) + (120*x^3*(n^2 + 16))/(a^4c^4(784n^2 + 56n^4 + n^6 + 2304)) + (360*x^4*(n^2 + 6))/(a^3c^4n(784n^2 + 56n^4 + n^6 + 2304)) + (30*x^2*(28n^2 + n^4 + 72))/(a^5c^4n(784n^2 + 56n^4 + n^6 + 2304)))) / (1/a^6 + x^6 + (3*x^4)/a^2 + (3*x^2)/a^4)$

3.348 $\int e^{n \operatorname{ArcTan}(ax)} (c + a^2 cx^2)^{3/2} dx$

Optimal. Leaf size=121

$$\frac{2^{\frac{5}{2}-\frac{in}{2}} c(1-iax)^{\frac{1}{2}(5+in)} \sqrt{c+a^2 cx^2} {}_2F_1\left(\frac{1}{2}(-3+in), \frac{1}{2}(5+in); \frac{1}{2}(7+in); \frac{1}{2}(1-iax)\right)}{a(5i-n)\sqrt{1+a^2 x^2}}$$

[Out] $-2^{(5/2-1/2*I*n)} * c * (1-I*a*x)^{(5/2+1/2*I*n)} * \operatorname{hypergeom}\left(\left[\frac{5}{2}+1/2*I*n, -3/2+1/2*I*n\right], \left[\frac{7}{2}+1/2*I*n\right], 1/2-1/2*I*a*x\right) * (a^2*c*x^2+c)^{(1/2)/a/(5*I-n)/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.08, antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{c2^{\frac{5}{2}-\frac{in}{2}} \sqrt{a^2 cx^2 + c} (1-iax)^{\frac{1}{2}(5+in)} {}_2F_1\left(\frac{1}{2}(in-3), \frac{1}{2}(in+5); \frac{1}{2}(in+7); \frac{1}{2}(1-iax)\right)}{a(-n+5i)\sqrt{a^2 x^2 + 1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n \operatorname{ArcTan}[a*x])} * (c + a^2*c*x^2)^{(3/2)}, x\right]$

[Out] $-((2^{(5/2 - (I/2)*n)} * c * (1 - I*a*x)^{((5 + I*n)/2)} * \operatorname{Sqrt}[c + a^2*c*x^2] * \operatorname{Hypergeometric2F1}\left[\left[-3 + I*n\right]/2, (5 + I*n)/2, (7 + I*n)/2, (1 - I*a*x)/2\right]) / (a * (5*I - n) * \operatorname{Sqrt}[1 + a^2*x^2])$

Rule 71

$\operatorname{Int}\left[\left((a_{.}) + (b_{.}) * (x_{.})\right)^{(m_{.})} * \left((c_{.}) + (d_{.}) * (x_{.})\right)^{(n_{.})}, x_{\text{Symbol}}\right] \rightarrow \operatorname{Simp}\left[\left((a + b*x)^{(m+1)} / (b*(m+1)*(b/(b*c - a*d))^n\right) * \operatorname{Hypergeometric2F1}\left[-n, m+1, m+2, (-d)*((a + b*x)/(b*c - a*d))\right], x\right] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\operatorname{Int}\left[E^{(\operatorname{ArcTan}[a_{.}] * (x_{.}))} * (n_{.}) * \left((c_{.}) + (d_{.}) * (x_{.})^2\right)^{(p_{.})}, x_{\text{Symbol}}\right] \rightarrow \operatorname{Dist}\left[c^p, \operatorname{Int}\left[(1 - I*a*x)^{(p + I*(n/2))} * (1 + I*a*x)^{(p - I*(n/2))}, x\right], x\right] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\operatorname{Int}\left[E^{(\operatorname{ArcTan}[a_{.}] * (x_{.}))} * (n_{.}) * \left((c_{.}) + (d_{.}) * (x_{.})^2\right)^{(p_{.})}, x_{\text{Symbol}}\right] \rightarrow \operatorname{Dist}\left[c^{\operatorname{IntPart}[p]} * \left((c + d*x^2)^{\operatorname{FracPart}[p]} / (1 + a^2*x^2)^{\operatorname{FracPart}[p]}\right), \operatorname{Int}\left[(1 + a^2*x^2)^p * E^{(n \operatorname{ArcTan}[a*x])}, x\right], x\right] /;$ FreeQ[{a, c, d, n, p}, x] && Eq

$Q[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} (c + a^2 cx^2)^{3/2} dx &= \frac{(c\sqrt{c + a^2 cx^2}) \int e^{n \tan^{-1}(ax)} (1 + a^2 x^2)^{3/2} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{(c\sqrt{c + a^2 cx^2}) \int (1 - iax)^{\frac{3}{2} + \frac{in}{2}} (1 + iax)^{\frac{3}{2} - \frac{in}{2}} dx}{\sqrt{1 + a^2 x^2}} \\ &= -\frac{2^{\frac{5}{2} - \frac{in}{2}} c (1 - iax)^{\frac{1}{2}(5+in)} \sqrt{c + a^2 cx^2} {}_2F_1\left(\frac{1}{2}(-3 + in), \frac{1}{2}(5 + in); \frac{1}{2}(7 + in); \frac{1}{2}(1 - iax)\right)}{a(5i - n)\sqrt{1 + a^2 x^2}} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 118, normalized size = 0.98

$$\frac{2^{\frac{5}{2} - \frac{in}{2}} c (1 - iax)^{\frac{5}{2} + \frac{in}{2}} \sqrt{c + a^2 cx^2} {}_2F_1\left(\frac{1}{2}(5 + in), \frac{1}{2}i(3i + n); \frac{1}{2}(7 + in); \frac{1}{2}(1 - iax)\right)}{a(-5i + n)\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2), x]

[Out] (2^(5/2 - (I/2)*n)*c*(1 - I*a*x)^(5/2 + (I/2)*n)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[(5 + I*n)/2, (I/2)*(3*I + n), (7 + I*n)/2, (1 - I*a*x)/2])/(a*(-5*I + n)*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} (a^2 c x^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(3/2), x)

[Out] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(3/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(3/2)*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(3/2)*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^{\frac{3}{2}} e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*(a**2*c*x**2+c)**(3/2),x)

[Out] Integral((c*(a**2*x**2 + 1))**(3/2)*exp(n*atan(a*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} (ca^2x^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^(3/2),x)

[Out] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^(3/2), x)

3.349 $\int e^{n \operatorname{ArcTan}(ax)} \sqrt{c + a^2 cx^2} dx$

Optimal. Leaf size=120

$$\frac{2^{\frac{3}{2}-\frac{in}{2}}(1-iax)^{\frac{1}{2}(3+in)}\sqrt{c+a^2cx^2} {}_2F_1\left(\frac{1}{2}(-1+in), \frac{1}{2}(3+in); \frac{1}{2}(5+in); \frac{1}{2}(1-iax)\right)}{a(3i-n)\sqrt{1+a^2x^2}}$$

[Out] $-2^{(3/2-1/2*I*n)}*(1-I*a*x)^{(3/2+1/2*I*n)}*\operatorname{hypergeom}\left(\left[\frac{3}{2}+1/2*I*n, -1/2+1/2*I*n\right], \left[\frac{5}{2}+1/2*I*n\right], 1/2-1/2*I*a*x\right)*(a^2*c*x^2+c)^{(1/2)}/a/(3*I-n)/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{2^{\frac{3}{2}-\frac{in}{2}}\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(3+in)} {}_2F_1\left(\frac{1}{2}(in-1), \frac{1}{2}(in+3); \frac{1}{2}(in+5); \frac{1}{2}(1-iax)\right)}{a(-n+3i)\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a*x])}*Sqrt[c + a^2*c*x^2], x]$

[Out] $-((2^{(3/2 - (I/2)*n)}*(1 - I*a*x)^{((3 + I*n)/2)}*Sqrt[c + a^2*c*x^2]*\operatorname{Hypergeometric2F1}[-(-1 + I*n)/2, (3 + I*n)/2, (5 + I*n)/2, (1 - I*a*x)/2])/(a*(3*I - n)*Sqrt[1 + a^2*x^2])$

Rule 71

$\operatorname{Int}[(a + b*x)^{(m+1)}/(b*(m+1)*(b*(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x$ && $\operatorname{NeQ}[b*c - a*d, 0]$ && $\operatorname{IntegerQ}[m]$ && $\operatorname{IntegerQ}[n]$ && $\operatorname{GtQ}[b/(b*c - a*d), 0]$ && $(\operatorname{RationalQ}[m] \mid \mid \operatorname{RationalQ}[n] \text{ \&\& } \operatorname{GtQ}[-d/(b*c - a*d), 0])$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a + b*x)]*(n))}*((c + d*(x)^2)^{(p)}), x_Symbol] :>$ $\operatorname{Dist}[c^p, \operatorname{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, n, p\}, x$ && $\operatorname{EqQ}[d, a^2*c]$ && $(\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[c, 0])$

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a + b*x)]*(n))}*((c + d*(x)^2)^{(p)}), x_Symbol] :>$ $\operatorname{Dist}[c^p*\operatorname{IntPart}[p]*((c + d*x^2)^{\operatorname{FracPart}[p]}/(1 + a^2*x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[(1 + a^2*x^2)^p*E^{(n*\operatorname{ArcTan}[a*x])}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, n, p\}, x$ && Eq

$\text{Q}[d, a^2c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} \sqrt{c + a^2cx^2} \, dx &= \frac{\sqrt{c + a^2cx^2} \int e^{n \tan^{-1}(ax)} \sqrt{1 + a^2x^2} \, dx}{\sqrt{1 + a^2x^2}} \\ &= \frac{\sqrt{c + a^2cx^2} \int (1 - iax)^{\frac{1}{2} + \frac{in}{2}} (1 + iax)^{\frac{1}{2} - \frac{in}{2}} \, dx}{\sqrt{1 + a^2x^2}} \\ &= -\frac{2^{\frac{3}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2}(3+in)} \sqrt{c + a^2cx^2} \, {}_2F_1\left(\frac{1}{2}(-1 + in), \frac{1}{2}(3 + in); \frac{1}{2}(5 + in); \frac{1}{2}\right)}{a(3i - n)\sqrt{1 + a^2x^2}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 117, normalized size = 0.98

$$\frac{2^{\frac{3}{2} - \frac{in}{2}} (1 - iax)^{\frac{3}{2} + \frac{in}{2}} \sqrt{c + a^2cx^2} \, {}_2F_1\left(\frac{1}{2}(3 + in), \frac{1}{2}i(i + n); \frac{1}{2}(5 + in); \frac{1}{2}(1 - iax)\right)}{a(-3i + n)\sqrt{1 + a^2x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*Sqrt[c + a^2*c*x^2], x]

[Out] (2^(3/2 - (I/2)*n)*(1 - I*a*x)^(3/2 + (I/2)*n)*Sqrt[c + a^2*c*x^2]*Hypergeometric2F1[(3 + I*n)/2, (I/2)*(I + n), (5 + I*n)/2, (1 - I*a*x)/2])/(a*(-3*I + n)*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} \sqrt{a^2cx^2 + c} \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a^2*c*x^2 + c)*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(a^2*c*x^2 + c)*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt{c(a^2x^2 + 1)} e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(sqrt(c*(a**2*x**2 + 1))*exp(n*atan(a*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} \sqrt{c a^2 x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^(1/2),x)

[Out] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^(1/2), x)

$$3.350 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=120

$$\frac{2^{\frac{1}{2}-\frac{in}{2}}(1-iax)^{\frac{1}{2}(1+in)}\sqrt{1+a^2x^2} {}_2F_1\left(\frac{1}{2}(1+in), \frac{1}{2}(1+in); \frac{1}{2}(3+in); \frac{1}{2}(1-iax)\right)}{a(i-n)\sqrt{c+a^2cx^2}}$$

[Out] $-2^{(1/2-1/2*I*n)}*(1-I*a*x)^{(1/2+1/2*I*n)}*\operatorname{hypergeom}\left([1/2+1/2*I*n, 1/2+1/2*I*n], [3/2+1/2*I*n], 1/2-1/2*I*a*x\right)*(a^2*x^2+1)^{(1/2)}/a/(I-n)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.07, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{2^{\frac{1}{2}-\frac{in}{2}}\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)} {}_2F_1\left(\frac{1}{2}(in+1), \frac{1}{2}(in+1); \frac{1}{2}(in+3); \frac{1}{2}(1-iax)\right)}{a(-n+i)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n*\operatorname{ArcTan}[a*x])}/\operatorname{Sqrt}[c + a^2*c*x^2], x\right]$

[Out] $-((2^{(1/2 - (I/2)*n)}*(1 - I*a*x)^{((1 + I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2]*\operatorname{Hypergeometric2F1}[(1 + I*n)/2, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/2])/(a*(I - n)*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*(a + b*x)/(b*c - a*d)], x] /; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] \mid\mid !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a_+]*(x_+))^{(n_+)}}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \operatorname{Dist}[c^{(p)}, \operatorname{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \operatorname{FreeQ}\{a, c, d, n, p\}, x \&\& \operatorname{EqQ}[d, a^2*c] \&\& (\operatorname{IntegerQ}[p] \mid\mid \operatorname{GtQ}[c, 0])$

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a_+]*(x_+))^{(n_+)}}*((c_+ + (d_+)*(x_+)^2)^{(p_+)}, x_Symbol] \rightarrow \operatorname{Dist}[c^{(\operatorname{IntPart}[p])}*((c + d*x^2)^{\operatorname{FracPart}[p]}/(1 + a^2*x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[(1 + a^2*x^2)^p * E^{(n*\operatorname{ArcTan}[a*x])}, x], x] /; \operatorname{FreeQ}\{a, c, d, n, p\}, x \&\& \operatorname{Eq}$

`Q[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])`

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= -\frac{2^{\frac{1}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2}(1+in)} \sqrt{1 + a^2 x^2} {}_2F_1\left(\frac{1}{2}(1 + in), \frac{1}{2}(1 + in); \frac{1}{2}(3 + in); \frac{1}{2}(1 - iax)\right)}{a(i - n)\sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 117, normalized size = 0.98

$$\frac{2^{\frac{1}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2} + \frac{in}{2}} \sqrt{1 + a^2 x^2} {}_2F_1\left(\frac{1}{2} + \frac{in}{2}, \frac{1}{2} + \frac{in}{2}; \frac{3}{2} + \frac{in}{2}; \frac{1}{2} - \frac{iax}{2}\right)}{a(-i + n)\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(n*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]`

`[Out] (2^(1/2 - (I/2)*n)*(1 - I*a*x)^(1/2 + (I/2)*n)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 + (I/2)*n, 1/2 + (I/2)*n, 3/2 + (I/2)*n, 1/2 - (I/2)*a*x])/(a*(-I + n)*Sqrt[c + a^2*c*x^2])`

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{\sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)`

`[Out] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(exp(n*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{\sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(1/2),x)

[Out] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(1/2), x)

3.351 $\int e^{n \operatorname{ArcTan}(ax)} x^2 (c + a^2 cx^2)^{3/2} dx$

Optimal. Leaf size=283

$$\frac{cn(1-iax)^{\frac{1}{2}(5+in)}(1+iax)^{\frac{1}{2}(5-in)}\sqrt{c+a^2cx^2}}{30a^3\sqrt{1+a^2x^2}} + \frac{cx(1-iax)^{\frac{1}{2}(5+in)}(1+iax)^{\frac{1}{2}(5-in)}\sqrt{c+a^2cx^2}}{6a^2\sqrt{1+a^2x^2}} + 2^{\frac{3}{2}-\frac{in}{2}}c(5$$

[Out] $-1/30*c*n*(1-I*a*x)^{(5/2+1/2*I*n)}*(1+I*a*x)^{(5/2-1/2*I*n)}*(a^2*c*x^2+c)^{(1/2)}/a^3/(a^2*x^2+1)^{(1/2)}+1/6*c*x*(1-I*a*x)^{(5/2+1/2*I*n)}*(1+I*a*x)^{(5/2-1/2*I*n)}*(a^2*c*x^2+c)^{(1/2)}/a^2/(a^2*x^2+1)^{(1/2)}+1/15*2^{(3/2-1/2*I*n)}*c*(-n^2+5)*(1-I*a*x)^{(5/2+1/2*I*n)}*\operatorname{hypergeom}([5/2+1/2*I*n, -3/2+1/2*I*n], [7/2+1/2*I*n], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^{(1/2)}/a^3/(5*I-n)/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.22, antiderivative size = 283, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {5193, 5190, 92, 81, 71}

$$\frac{cx\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(5+in)}(1+iax)^{\frac{1}{2}(5-in)}}{6a^2\sqrt{a^2x^2+1}} + \frac{c2^{\frac{3}{2}-\frac{in}{2}}(5-n^2)\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(5+in)}{}_2F_1(\frac{1}{2}(in-3), \frac{1}{2}(in+5); \frac{1}{2}(in+7); \frac{1}{2}(1-iax))}{15a^3(-n+5i)\sqrt{a^2x^2+1}} - \frac{cn\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(5+in)}(1+iax)^{\frac{1}{2}(5-in)}}{30a^3\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n \operatorname{ArcTan}[a*x])} * x^2 * (c + a^2 * c * x^2)^{(3/2)}, x]$

[Out] $-1/30*(c*n*(1-I*a*x)^{((5+I*n)/2)}*(1+I*a*x)^{((5-I*n)/2)}*\operatorname{Sqrt}[c+a^2*c*x^2])/(a^3*\operatorname{Sqrt}[1+a^2*x^2]) + (c*x*(1-I*a*x)^{((5+I*n)/2)}*(1+I*a*x)^{((5-I*n)/2)}*\operatorname{Sqrt}[c+a^2*c*x^2])/(6*a^2*\operatorname{Sqrt}[1+a^2*x^2]) + (2^{(3/2-(I/2)*n)}*c*(5-n^2)*(1-I*a*x)^{((5+I*n)/2)}*\operatorname{Sqrt}[c+a^2*c*x^2]*\operatorname{Hypergeometric2F1}[(-3+I*n)/2, (5+I*n)/2, (7+I*n)/2, (1-I*a*x)/2])/(15*a^3*(5*I-n)*\operatorname{Sqrt}[1+a^2*x^2])$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)(x_+))^{(m_+)} * ((c_+ + (d_+)(x_+))^{(n_+)}, x_Symbol] \rightarrow \operatorname{Simp}[(a + b*x)^{(m+1)} / (b*(m+1)*(b/(b*c - a*d))^{(n)}) * \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a+b*x)/(b*c - a*d)], x] / ; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] || !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 81

$\operatorname{Int}[(a_+ + (b_+)(x_+)) * ((c_+ + (d_+)(x_+))^{(n_+)} * ((e_+ + (f_+)(x_+))^{(p_+)}, x_Symbol] \rightarrow \operatorname{Simp}[b*(c + d*x)^{(n+1)} * (e + f*x)^{(p+1)} / (d*f*(n+p+2))], x] + \operatorname{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1))] / (d*f*(n+p+2)), \operatorname{Int}[(c + d*x)^n * (e + f*x)^p, x], x] / ; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \operatorname{NeQ}[n+p+2, 0]$

Rule 92

```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rule 5193

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned}
 \int e^{n \tan^{-1}(ax)} x^2 (c + a^2 cx^2)^{3/2} dx &= \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int e^{n \tan^{-1}(ax)} x^2 (1 + a^2 x^2)^{3/2} dx}{\sqrt{1 + a^2 x^2}} \\
 &= \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int x^2 (1 - iax)^{\frac{3}{2} + \frac{in}{2}} (1 + iax)^{\frac{3}{2} - \frac{in}{2}} dx}{\sqrt{1 + a^2 x^2}} \\
 &= \frac{cx(1 - iax)^{\frac{1}{2}(5+in)}(1 + iax)^{\frac{1}{2}(5-in)}\sqrt{c + a^2 cx^2}}{6a^2\sqrt{1 + a^2 x^2}} + \frac{\left(c\sqrt{c + a^2 cx^2}\right) \int (1 - iax)^{\frac{1}{2}(5+in)}(1 + iax)^{\frac{1}{2}(5-in)} dx}{6a^2\sqrt{1 + a^2 x^2}} \\
 &= -\frac{cn(1 - iax)^{\frac{1}{2}(5+in)}(1 + iax)^{\frac{1}{2}(5-in)}\sqrt{c + a^2 cx^2}}{30a^3\sqrt{1 + a^2 x^2}} + \frac{cx(1 - iax)^{\frac{1}{2}(5+in)}(1 + iax)^{\frac{1}{2}(5-in)}}{6a^2\sqrt{1 + a^2 x^2}} \\
 &= -\frac{cn(1 - iax)^{\frac{1}{2}(5+in)}(1 + iax)^{\frac{1}{2}(5-in)}\sqrt{c + a^2 cx^2}}{30a^3\sqrt{1 + a^2 x^2}} + \frac{cx(1 - iax)^{\frac{1}{2}(5+in)}(1 + iax)^{\frac{1}{2}(5-in)}}{6a^2\sqrt{1 + a^2 x^2}}
 \end{aligned}$$

Mathematica [A]

time = 0.19, size = 217, normalized size = 0.77

$$\frac{2^{-1-\frac{n}{2}}c(1 - iax)^{\frac{1}{2}+\frac{in}{2}}(1 + iax)^{-\frac{n}{2}}(i + ax)^2\sqrt{c + a^2 cx^2} \left(2^{\frac{n}{2}}(-5i + n)\sqrt{1 + iax}(-i + ax)^2(-n + 5ax) - 4\sqrt{2}(-5 + n^2)(1 + iax)^{\frac{n}{2}}{}_2F_1\left(\frac{1}{2}(5 + in), \frac{1}{2}i(3i + n); \frac{1}{2}(7 + in); \frac{1}{2}(1 - iax)\right)\right)}{15a^3(-5i + n)\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*x^2*(c + a^2*c*x^2)^(3/2), x]

[Out] (2^(-1 - (I/2)*n)*c*(1 - I*a*x)^(1/2 + (I/2)*n)*(I + a*x)^2*Sqrt[c + a^2*c*x^2]*(2^((I/2)*n)*(-5*I + n)*Sqrt[1 + I*a*x]*(-I + a*x)^2*(-n + 5*a*x) - 4*Sqrt[2]*(-5 + n^2)*(1 + I*a*x)^((I/2)*n)*Hypergeometric2F1[(5 + I*n)/2, (I/2)*(3*I + n), (7 + I*n)/2, (1 - I*a*x)/2]))/(15*a^3*(-5*I + n)*(1 + I*a*x)^((I/2)*n)*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} x^2 (a^2 c x^2 + c)^{\frac{3}{2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(3/2), x)

[Out] int(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(3/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(3/2), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(3/2)*x^2*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(3/2), x, algorithm="fricas")

[Out] integral((a^2*c*x^4 + c*x^2)*sqrt(a^2*c*x^2 + c)*e^(n*arctan(a*x)), x)

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*x**2*(a**2*c*x**2+c)**(3/2),x)

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 e^{n \operatorname{atan}(a x)} (c a^2 x^2 + c)^{3/2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*exp(n*atan(a*x))*(c + a^2*c*x^2)^(3/2),x)

[Out] int(x^2*exp(n*atan(a*x))*(c + a^2*c*x^2)^(3/2), x)

3.352 $\int e^{n \operatorname{ArcTan}(ax)} x^2 \sqrt{c + a^2 cx^2} dx$

Optimal. Leaf size=280

$$\frac{n(1-iax)^{\frac{1}{2}(3+in)}(1+iax)^{\frac{1}{2}(3-in)}\sqrt{c+a^2cx^2}}{12a^3\sqrt{1+a^2x^2}} + \frac{x(1-iax)^{\frac{1}{2}(3+in)}(1+iax)^{\frac{1}{2}(3-in)}\sqrt{c+a^2cx^2}}{4a^2\sqrt{1+a^2x^2}} + \frac{2^{-\frac{1}{2}-\frac{in}{2}}(3-}$$

[Out] $-1/12*n*(1-I*a*x)^{(3/2+1/2*I*n)}*(1+I*a*x)^{(3/2-1/2*I*n)}*(a^2*c*x^2+c)^{(1/2)}/a^3/(a^2*x^2+1)^{(1/2)}+1/4*x*(1-I*a*x)^{(3/2+1/2*I*n)}*(1+I*a*x)^{(3/2-1/2*I*n)}*(a^2*c*x^2+c)^{(1/2)}/a^2/(a^2*x^2+1)^{(1/2)}+1/3*2^{(-1/2-1/2*I*n)}*(-n^2+3)*(1-I*a*x)^{(3/2+1/2*I*n)}*\operatorname{hypergeom}([3/2+1/2*I*n, -1/2+1/2*I*n], [5/2+1/2*I*n], 1/2-1/2*I*a*x)*(a^2*c*x^2+c)^{(1/2)}/a^3/(3*I-n)/(a^2*x^2+1)^{(1/2)}$

Rubi [A]

time = 0.20, antiderivative size = 280, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {5193, 5190, 92, 81, 71}

$$\frac{x\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(3+in)}(1+iax)^{\frac{1}{2}(3-in)}}{4a^2\sqrt{a^2x^2+1}} + \frac{2^{-\frac{1}{2}-\frac{in}{2}}(3-n^2)\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(3+in)}{}_2F_1(\frac{1}{2}(in-1), \frac{1}{2}(in+3); \frac{1}{2}(in+5); \frac{1}{2}(1-iax))}{3a^3(-n+3i)\sqrt{a^2x^2+1}} - \frac{n\sqrt{a^2cx^2+c}(1-iax)^{\frac{1}{2}(3+in)}(1+iax)^{\frac{1}{2}(3-in)}}{12a^3\sqrt{a^2x^2+1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n \operatorname{ArcTan}[a*x])} * x^2 * \operatorname{Sqrt}[c + a^2*c*x^2], x]$

[Out] $-1/12*(n*(1-I*a*x)^{((3+I*n)/2)}*(1+I*a*x)^{((3-I*n)/2)}*\operatorname{Sqrt}[c+a^2*c*x^2])/a^3*\operatorname{Sqrt}[1+a^2*x^2]+(x*(1-I*a*x)^{((3+I*n)/2)}*(1+I*a*x)^{((3-I*n)/2)}*\operatorname{Sqrt}[c+a^2*c*x^2])/(4*a^2*\operatorname{Sqrt}[1+a^2*x^2])+(2^{(-1/2-(I/2)*n)}*(3-n^2)*(1-I*a*x)^{((3+I*n)/2)}*\operatorname{Sqrt}[c+a^2*c*x^2]*\operatorname{Hypergeometric2F1}[(-1+I*n)/2, (3+I*n)/2, (5+I*n)/2, (1-I*a*x)/2])/(3*a^3*(3I-n)*\operatorname{Sqrt}[1+a^2*x^2])$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] :> \operatorname{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)*(b/(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \operatorname{NeQ}[b*c - a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c - a*d), 0] \&\& (\operatorname{RationalQ}[m] || !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0]))$

Rule 81

$\operatorname{Int}[(a_+ + (b_+)*(x_+))*((c_+ + (d_+)*(x_+))^{(n_+)})*((e_+ + (f_+)*(x_+))^{(p_+)}, x_Symbol] :> \operatorname{Simp}[b*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/(d*f*(n + p + 2))), x] + \operatorname{Dist}[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1))]/(d*f*(n + p + 2)), \operatorname{Int}[(c + d*x)^n*(e + f*x)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \operatorname{NeQ}[n + p + 2, 0]$

Rule 92

```
Int[((a_.) + (b_.)*(x_))2*((c_.) + (d_.)*(x_))(n_.)*((e_.) + (f_.)*(x_))(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)(n + 1)*((e + f*x)(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)n*(e + f*x)p*Simp[a2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1)) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 5190

```
Int[E(ArcTan[(a_.)*(x_)]*(n_.))*(x_)(m_.)*((c_.) + (d_.)*(x_)2)(p_.), x_Symbol] := Dist[cp, Int[xm*(1 - I*a*x)(p + I*(n/2))(1 + I*a*x)(p - I*(n/2))), x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rule 5193

```
Int[E(ArcTan[(a_.)*(x_)]*(n_.))*(x_)(m_.)*((c_.) + (d_.)*(x_)2)(p_.), x_Symbol] := Dist[cIntPart[p]*((c + d*x2)FracPart[p]/(1 + a2*x2)FracPart[p]), Int[xm*(1 + a2*x2)p*E(n*ArcTan[a*x])), x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} x^2 \sqrt{c + a^2 cx^2} dx &= \frac{\sqrt{c + a^2 cx^2} \int e^{n \tan^{-1}(ax)} x^2 \sqrt{1 + a^2 x^2} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{\sqrt{c + a^2 cx^2} \int x^2 (1 - iax)^{\frac{1}{2} + \frac{in}{2}} (1 + iax)^{\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{1 + a^2 x^2}} \\ &= \frac{x(1 - iax)^{\frac{1}{2}(3+in)} (1 + iax)^{\frac{1}{2}(3-in)} \sqrt{c + a^2 cx^2}}{4a^2 \sqrt{1 + a^2 x^2}} + \frac{\sqrt{c + a^2 cx^2} \int (1 - iax)^{\frac{1}{2}}}{4a^2} \\ &= -\frac{n(1 - iax)^{\frac{1}{2}(3+in)} (1 + iax)^{\frac{1}{2}(3-in)} \sqrt{c + a^2 cx^2}}{12a^3 \sqrt{1 + a^2 x^2}} + \frac{x(1 - iax)^{\frac{1}{2}(3+in)} (1 + iax)^{\frac{1}{2}(3-in)}}{4a^2 \sqrt{1 + a^2 x^2}} \\ &= -\frac{n(1 - iax)^{\frac{1}{2}(3+in)} (1 + iax)^{\frac{1}{2}(3-in)} \sqrt{c + a^2 cx^2}}{12a^3 \sqrt{1 + a^2 x^2}} + \frac{x(1 - iax)^{\frac{1}{2}(3+in)} (1 + iax)^{\frac{1}{2}(3-in)}}{4a^2 \sqrt{1 + a^2 x^2}} \end{aligned}$$

Mathematica [A]

time = 0.15, size = 214, normalized size = 0.76

$$\frac{2^{-2-\frac{n}{2}}(1 - iax)^{\frac{1}{2}+\frac{n}{2}}(1 + iax)^{-\frac{n}{2}}(i + ax)\sqrt{c + a^2 cx^2} \left(2^{\frac{n}{2}}(-3i + n)\sqrt{1 + iax}(-i + ax)(-n + 3ax) - 2i\sqrt{2}(-3 + n^2)(1 + iax)^{\frac{n}{2}} {}_2F_1\left(\frac{1}{2}(3 + in), \frac{1}{2}(i + n); \frac{1}{2}(5 + in); \frac{1}{2}(1 - iax)\right) \right)}{3a^3(-3i + n)\sqrt{1 + a^2 x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*x^2*Sqrt[c + a^2*c*x^2],x]

[Out] (2^(-2 - (I/2)*n)*(1 - I*a*x)^(1/2 + (I/2)*n)*(I + a*x)*Sqrt[c + a^2*c*x^2] * (2^((I/2)*n)*(-3*I + n)*Sqrt[1 + I*a*x]*(-I + a*x)*(-n + 3*a*x) - (2*I)*Sqrt[2]*(-3 + n^2)*(1 + I*a*x)^((I/2)*n)*Hypergeometric2F1[(3 + I*n)/2, (I/2)*(I + n), (5 + I*n)/2, (1 - I*a*x)/2]))/(3*a^3*(-3*I + n)*(1 + I*a*x)^((I/2)*n)*Sqrt[1 + a^2*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} x^2 \sqrt{a^2 c x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(1/2),x)

[Out] int(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(sqrt(a^2*c*x^2 + c)*x^2*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(a^2*c*x^2 + c)*x^2*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int x^2 \sqrt{c(a^2 x^2 + 1)} e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*x**2*(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(x**2*sqrt(c*(a**2*x**2 + 1))*exp(n*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int x^2 e^{n \operatorname{atan}(ax)} \sqrt{ca^2 x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2*exp(n*atan(a*x))*(c + a^2*c*x^2)^(1/2),x)

[Out] int(x^2*exp(n*atan(a*x))*(c + a^2*c*x^2)^(1/2), x)

3.353 $\int \frac{e^{n \operatorname{ArcTan}(ax)} x^3}{\sqrt{c + a^2 cx^2}} dx$

Optimal. Leaf size=322

$$\frac{x^2(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}\sqrt{1+a^2x^2}}{3a^2\sqrt{c+a^2cx^2}} - \frac{(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}(4-in-n^2+a(1+in)nx)}{6a^4(1+in)\sqrt{c+a^2cx^2}}$$

[Out] $1/3*x^2*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*(a^2*x^2+1)^{(1/2)}/a$
 $^2/(a^2*c*x^2+c)^{(1/2)}-1/6*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*$
 $(4-I*n-n^2+a*(1+I*n)*n*x)*(a^2*x^2+1)^{(1/2)}/a^4/(1+I*n)/(a^2*c*x^2+c)^{(1/2)}$
 $+1/3*2^{(-1/2-1/2*I*n)}*n*(-n^2+5)*(1-I*a*x)^{(3/2+1/2*I*n)}*\operatorname{hypergeom}([3/2+1/2$
 $*I*n, 1/2+1/2*I*n], [5/2+1/2*I*n], 1/2-1/2*I*a*x)*(a^2*x^2+1)^{(1/2)}/a^4/(4*n-$
 $I*(-n^2+3))/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.25, antiderivative size = 322, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {5193, 5190, 102, 151, 71}

$$\frac{x^2\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}}{3a^2\sqrt{a^2cx^2+c}} + \frac{2^{-\frac{1}{2}-\frac{in}{2}}n(5-n^2)\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}{}_2F_1\left(\frac{1}{2}(in+1), \frac{1}{2}(in+3); \frac{1}{2}(in+5); \frac{1}{2}(1-iax)\right)}{3a^4(4n-i(3-n^2))\sqrt{a^2cx^2+c}} - \frac{\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(a(1+in)nx-n^2-in+4)(1+iax)^{\frac{1}{2}(1-in)}}{6a^4(1+in)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])})*x^3]/\operatorname{Sqrt}[c + a^2*c*x^2], x]$

[Out] $(x^2*(1-I*a*x)^{((1+I*n)/2)}*(1+I*a*x)^{((1-I*n)/2)}*\operatorname{Sqrt}[1+a^2*x^2])$
 $/(3*a^2*\operatorname{Sqrt}[c+a^2*c*x^2]) - ((1-I*a*x)^{((1+I*n)/2)}*(1+I*a*x)^{((1-I$
 $n)/2)}*(4-I*n-n^2+a*(1+I*n)*n*x)*\operatorname{Sqrt}[1+a^2*x^2])/(6*a^4*(1+I$
 $n)*\operatorname{Sqrt}[c+a^2*c*x^2]) + (2^{(-1/2-(I/2)*n)}*n*(5-n^2)*(1-I*a*x)^{((3$
 $+I*n)/2)}*\operatorname{Sqrt}[1+a^2*x^2]*\operatorname{Hypergeometric2F1}[(1+I*n)/2, (3+I*n)/2, (5$
 $+I*n)/2, (1-I*a*x)/2])/(3*a^4*(4*n-I*(3-n^2))*\operatorname{Sqrt}[c+a^2*c*x^2])$

Rule 71

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}, x_Symbol] :> \operatorname{Simp}[(a$
 $+ b*x)^{(m+1)}/(b*(m+1)*(b/(b*c-a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m+1$
 $, m+2, (-d)*((a+b*x)/(b*c-a*d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x]
 $\&\& \operatorname{NeQ}[b*c-a*d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b*c-a*d)$
 $, 0] \&\& (\operatorname{RationalQ}[m] || !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b*c-a*d), 0]))$

Rule 102

$\operatorname{Int}[(a_+ + (b_+)*(x_+))^{(m_+)}*((c_+ + (d_+)*(x_+))^{(n_+)}*((e_+ + (f_+)*(x_+)$
 $)^{(p_+)}, x_Symbol] :> \operatorname{Simp}[b*(a+b*x)^{(m-1)}*(c+d*x)^{(n+1)}*((e+f*x)$
 $)^{(p+1)}/(d*f*(m+n+p+1))], x] + \operatorname{Dist}[1/(d*f*(m+n+p+1)), \operatorname{Int}[(a$
 $+ b*x)^{(m-2)}*(c+d*x)^n*(e+f*x)^p*\operatorname{Simp}[a^2*d*f*(m+n+p+1) - b*(b$

*c*e*(m - 1) + a*(d*e*(n + 1) + c*f*(p + 1)) + b*(a*d*f*(2*m + n + p) - b*(d*e*(m + n) + c*f*(m + p)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && GtQ[m, 1] && NeQ[m + n + p + 1, 0] && IntegerQ[m]

Rule 151

Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_) + (f_.)*(x_))*((g_.) + (h_.)*(x_)), x_Symbol] := Simp[((a^2*d*f*h*(n + 2) + b^2*d*e*g*(m + n + 3) + a*b*(c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b*f*h*(b*c - a*d)*(m + 1)*x)/(b^2*d*(b*c - a*d)*(m + 1)*(m + n + 3))*(a + b*x)^(m + 1)*(c + d*x)^(n + 1), x] - Dist[(a^2*d^2*f*h*(n + 1)*(n + 2) + a*b*d*(n + 1)*(2*c*f*h*(m + 1) - d*(f*g + e*h)*(m + n + 3)) + b^2*(c^2*f*h*(m + 1)*(m + 2) - c*d*(f*g + e*h)*(m + 1)*(m + n + 3) + d^2*e*g*(m + n + 2)*(m + n + 3)))/(b^2*d*(b*c - a*d)*(m + 1)*(m + n + 3)), Int[(a + b*x)^(m + 1)*(c + d*x)^n, x], x] /; FreeQ[{a, b, c, d, e, f, g, h, m, n}, x] && ((GeQ[m, -2] && LtQ[m, -1]) || SumSimplerQ[m, 1]) && NeQ[m, -1] && NeQ[m + n + 3, 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)} x^3}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)} x^3}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int x^3 (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{x^2 (1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{3a^2 \sqrt{c + a^2 cx^2}} + \frac{\sqrt{1 + a^2 x^2} \int x (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{3a^2 \sqrt{c + a^2 cx^2}} \\
&= \frac{x^2 (1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{3a^2 \sqrt{c + a^2 cx^2}} - \frac{(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} (4 - i)}{6a^4 (1 + in) \sqrt{c + a^2 cx^2}} \\
&= \frac{x^2 (1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{3a^2 \sqrt{c + a^2 cx^2}} - \frac{(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} (4 - i)}{6a^4 (1 + in) \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.22, size = 248, normalized size = 0.77

$$\frac{2^{-\frac{3}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{in}{2}} \sqrt{1 + a^2 x^2} \left(2^{\frac{1}{2} + \frac{in}{2}} (-3i + n) \sqrt{1 + iax} (-n^2(i + ax) - 2i(-2 + a^2 x^2) + n(1 + iax + 2a^2 x^2)) + 2n(-5 + n^2)(1 + iax)^{\frac{in}{2}} (i + ax) {}_2F_1\left(\frac{1}{2} + \frac{in}{2}, \frac{3}{2} + \frac{in}{2}; \frac{5}{2} + \frac{in}{2}; \frac{1 - iax}{2}\right) \right)}{3a^4 (-3 - 4in + n^2) \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^(n*ArcTan[a*x])*x^3)/Sqrt[c + a^2*c*x^2], x]

[Out] (2^(-3/2 - (I/2)*n)*(1 - I*a*x)^(1/2 + (I/2)*n)*Sqrt[1 + a^2*x^2]*(2^(1/2 + (I/2)*n)*(-3*I + n)*Sqrt[1 + I*a*x]*(-(n^2*(I + a*x)) - (2*I)*(-2 + a^2*x^2) + n*(1 + I*a*x + 2*a^2*x^2)) + 2*n*(-5 + n^2)*(1 + I*a*x)^((I/2)*n)*(I + a*x)*Hypergeometric2F1[1/2 + (I/2)*n, 3/2 + (I/2)*n, 5/2 + (I/2)*n, 1/2 - (I/2)*a*x])/ (3*a^4*(-3 - (4*I)*n + n^2)*(1 + I*a*x)^((I/2)*n)*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^3}{\sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c)^(1/2), x)**[Out]** int(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c)^(1/2), x)**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(x^3*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

[Out] `integral(x^3*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^3 e^{n \operatorname{atan}(ax)}}{\sqrt{c(a^2 x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))*x**3/(a**2*c*x**2+c)**(1/2),x)`

[Out] `Integral(x**3*exp(n*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)`

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^3/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP UT:sage2:=int(sage0,sageVARx):;OUTPUT:sym2poly/r2sym(const gen & e,const in dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^3 e^{n \operatorname{atan}(ax)}}{\sqrt{c a^2 x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^3*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2),x)`

[Out] `int((x^3*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2), x)`

3.354 $\int \frac{e^{n \operatorname{ArcTan}(ax)} x^2}{\sqrt{c + a^2 c x^2}} dx$

Optimal. Leaf size=291

$$\frac{(1 + in)(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{2a^3(i + n)\sqrt{c + a^2cx^2}} + \frac{x(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{2a^2\sqrt{c + a^2cx^2}} - i2^{\frac{1}{2} - \frac{in}{2}}$$

[Out] $-1/2*(1+I*n)*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*(a^2*x^2+1)^{(1/2)}/a^3/(I+n)/(a^2*c*x^2+c)^{(1/2)+1/2*x*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*(a^2*x^2+1)^{(1/2)}/a^2/(a^2*c*x^2+c)^{(1/2)}-I^2^{(1/2-1/2*I*n)}*(-n^2+1)*(1-I*a*x)^{(1/2+1/2*I*n)}*\operatorname{hypergeom}([-1/2+1/2*I*n, 1/2+1/2*I*n], [3/2+1/2*I*n], 1/2-1/2*I*a*x)*(a^2*x^2+1)^{(1/2)}/a^3/(n^2+1)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.25, antiderivative size = 291, normalized size of antiderivative = 1.00, number of steps used = 5, number of rules used = 5, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.192$, Rules used = {5193, 5190, 92, 80, 71}

$$\frac{x\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}}{2a^2\sqrt{a^2cx^2+c}} - \frac{i2^{\frac{1}{2}-\frac{in}{2}}(1-n^2)\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}{}_2F_1(\frac{1}{2}(in-1), \frac{1}{2}(in+1); \frac{1}{2}(in+3); \frac{1}{2}(1-iax))}{a^3(n^2+1)\sqrt{a^2cx^2+c}} - \frac{(1+in)\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}}{2a^3(n+i)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{n \operatorname{ArcTan}[a*x]}) * x^2] / \operatorname{Sqrt}[c + a^2 * c * x^2], x]$

[Out] $-1/2*((1 + I*n)*(1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2])/(a^3*(I + n)*\operatorname{Sqrt}[c + a^2*c*x^2]) + (x*(1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2])/(2*a^2*\operatorname{Sqrt}[c + a^2*c*x^2]) - (I^2^{(1/2 - (I/2)*n)}*(1 - n^2)*(1 - I*a*x)^{((1 + I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2])*\operatorname{Hypergeometric2F1}[(-1 + I*n)/2, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/2]/(a^3*(1 + n^2)*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 71

$\operatorname{Int}[(a + b*x)^m * (c + d*x)^n, x_Symbol] := \operatorname{Simp}[(a + b*x)^{m+1} / (b*(m+1)*(b*c - a*d)^n) * \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x]$ $\&\& \operatorname{NeQ}[b*c - a*d, 0]$ $\&\& \operatorname{IntegerQ}[m]$ $\&\& \operatorname{IntegerQ}[n]$ $\&\& \operatorname{GtQ}[b/(b*c - a*d), 0]$ $\&\& (\operatorname{RationalQ}[m] \mid \mid \operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[-d/(b*c - a*d), 0])$

Rule 80

$\operatorname{Int}[(a + b*x)^m * (c + d*x)^n * (e + f*x)^p, x_Symbol] := \operatorname{Simp}[(-b*e - a*f)*(c + d*x)^{n+1} * (e + f*x)^{p+1} / (f*(p+1)*(c*f - d*e)), x] - \operatorname{Dist}[(a*d*f*(n+p+2) - b*(d*e*(n+1) + c*f*(p+1)) / (f*(p+1)*(c*f - d*e)), \operatorname{Int}[(c + d*x)^n * (e + f*x)^p \operatorname{Simplify}[p+1], x], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x]$ $\&\& \operatorname{IntegerQ}[p]$ $\&\& \operatorname{Sum}$

SimplerQ[p, 1]

Rule 92

```
Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] := Simp[b*(a + b*x)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(d*f*(n + p + 3))), x] + Dist[1/(d*f*(n + p + 3)), Int[(c + d*x)^n*(e + f*x)^p*Simp[a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) + b*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2)))*x, x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 3, 0]
```

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rule 5193

```
Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned}
 \int \frac{e^{n \tan^{-1}(ax)} x^2}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)} x^2}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\
 &= \frac{\sqrt{1 + a^2 x^2} \int x^2 (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{c + a^2 cx^2}} \\
 &= \frac{x(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2a^2 \sqrt{c + a^2 cx^2}} + \frac{\sqrt{1 + a^2 x^2} \int (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{2a^2 \sqrt{c + a^2 cx^2}} \\
 &= -\frac{(1 + in)(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2a^3 (i + n) \sqrt{c + a^2 cx^2}} + \frac{x(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)}}{2a^2 \sqrt{c + a^2 cx^2}} \\
 &= -\frac{(1 + in)(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2a^3 (i + n) \sqrt{c + a^2 cx^2}} + \frac{x(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)}}{2a^2 \sqrt{c + a^2 cx^2}}
 \end{aligned}$$

Mathematica [A]

time = 0.14, size = 206, normalized size = 0.71

$$\frac{2^{-1-\frac{n}{2}}(1-iax)^{\frac{1}{2}+\frac{n}{2}}(1+iax)^{-\frac{n}{2}}\sqrt{1+a^2x^2}\left(2^{\frac{n}{2}}(-i+n)\sqrt{1+iax}(-1+iax+n(-i+ax))+2i\sqrt{2}(-1+n^2)(1+iax)^{\frac{n}{2}}{}_2F_1\left(\frac{1}{2}(1+in),\frac{1}{2}i(i+n);\frac{1}{2}(3+in);\frac{1}{2}(1-iax)\right)\right)}{a^3(1+n^2)\sqrt{c+a^2cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^(n*ArcTan[a*x])*x^2)/Sqrt[c + a^2*c*x^2], x]

[Out] (2^(-1 - (I/2)*n)*(1 - I*a*x)^(1/2 + (I/2)*n)*Sqrt[1 + a^2*x^2]*(2^((I/2)*n)*(-I + n)*Sqrt[1 + I*a*x]*(-1 + I*a*x + n*(-I + a*x)) + (2*I)*Sqrt[2]*(-1 + n^2)*(1 + I*a*x)^((I/2)*n)*Hypergeometric2F1[(1 + I*n)/2, (I/2)*(I + n), (3 + I*n)/2, (1 - I*a*x)/2]))/(a^3*(1 + n^2)*(1 + I*a*x)^((I/2)*n)*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^2}{\sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(x^2*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c)^(1/2), x, algorithm="fricas")

[Out] integral(x^2*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^2 e^{n \operatorname{atan}(ax)}}{\sqrt{c(a^2 x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*x**2/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(x**2*exp(n*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^2/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x^2 e^{n \operatorname{atan}(ax)}}{\sqrt{c a^2 x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2),x)

[Out] int((x^2*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2), x)

$$3.355 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=202

$$\frac{(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{a^2(1 - in)\sqrt{c + a^2cx^2}} - \frac{i2^{\frac{3}{2}-\frac{in}{2}}n(1 - iax)^{\frac{1}{2}(1+in)}\sqrt{1 + a^2x^2}}{a^2(1 + n^2)\sqrt{c + a^2cx^2}} {}_2F_1\left(\frac{1}{2}(-1 + in), \frac{1}{2}(1 + in); \right);$$

[Out] (1-I*a*x)^(1/2+1/2*I*n)*(1+I*a*x)^(1/2-1/2*I*n)*(a^2*x^2+1)^(1/2)/a^2/(1-I*n)/(a^2*c*x^2+c)^(1/2)-I*2^(3/2-1/2*I*n)*n*(1-I*a*x)^(1/2+1/2*I*n)*hypergeom([-1/2+1/2*I*n, 1/2+1/2*I*n],[3/2+1/2*I*n],1/2-1/2*I*a*x)*(a^2*x^2+1)^(1/2)/a^2/(n^2+1)/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.13, antiderivative size = 202, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.167$, Rules used = {5193, 5190, 80, 71}

$$\frac{\sqrt{a^2x^2+1}(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}}{a^2(1 - in)\sqrt{a^2cx^2 + c}} - \frac{i2^{\frac{3}{2}-\frac{in}{2}}n\sqrt{a^2x^2+1}(1 - iax)^{\frac{1}{2}(1+in)}}{a^2(n^2+1)\sqrt{a^2cx^2 + c}} {}_2F_1\left(\frac{1}{2}(in - 1), \frac{1}{2}(in + 1); \frac{1}{2}(in + 3); \frac{1}{2}(1 - iax)\right)$$

Antiderivative was successfully verified.

[In] Int[(E^(n*ArcTan[a*x])*x)/Sqrt[c + a^2*c*x^2], x]

[Out] ((1 - I*a*x)^((1 + I*n)/2)*(1 + I*a*x)^((1 - I*n)/2)*Sqrt[1 + a^2*x^2])/(a^2*(1 - I*n)*Sqrt[c + a^2*c*x^2]) - (I*2^(3/2 - (I/2)*n)*n*(1 - I*a*x)^((1 + I*n)/2)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[(-1 + I*n)/2, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/2])/(a^2*(1 + n^2)*Sqrt[c + a^2*c*x^2])

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] :> Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b*(b*c - a*d)))^n)*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 80

Int[((a_.) + (b_.)*(x_))*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[(-b*e - a*f)*(c + d*x)^(n + 1)*((e + f*x)^(p + 1)/(f*(p + 1)*(c*f - d*e))), x] - Dist[(a*d*f*(n + p + 2) - b*(d*e*(n + 1) + c*f*(p + 1)))/(f*(p + 1)*(c*f - d*e)), Int[(c + d*x)^n*(e + f*x)^Simplify[p + 1], x], x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && !RationalQ[p] && SumSimplerQ[p, 1]

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_
Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(
n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ
[p] || GtQ[c, 0])
```

Rule 5193

```
Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_), x_S
ymbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart
[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d,
m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)} x}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int x(1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{a^2(1 - in) \sqrt{c + a^2 cx^2}} - \frac{\left(n \sqrt{1 + a^2 x^2}\right) \int (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{a(1 - in) \sqrt{c + a^2 cx^2}} \\ &= \frac{(1 - iax)^{\frac{1}{2}(1+in)} (1 + iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{a^2(1 - in) \sqrt{c + a^2 cx^2}} - \frac{i 2^{\frac{3}{2} - \frac{in}{2}} n (1 - iax)^{\frac{1}{2}(1+in)} \sqrt{1 + a^2 x^2} {}_2F_1\left(\frac{1}{2}(1+in), \frac{1}{2}(i+n); \frac{1}{2}(3+in); \frac{1}{2}(1-iax)\right)}{a^2(1 - in) \sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.10, size = 187, normalized size = 0.93

$$\frac{i 2^{-\frac{1}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{in}{2}} \sqrt{1 + a^2 x^2} \left(2^{\frac{1}{2} + \frac{in}{2}} (-i + n) \sqrt{1 + iax} - 4n(1 + iax)^{\frac{in}{2}} {}_2F_1\left(\frac{1}{2}(1+in), \frac{1}{2}(i+n); \frac{1}{2}(3+in); \frac{1}{2}(1-iax)\right) \right)}{a^2(1 + n^2) \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^(n*ArcTan[a*x])*x)/Sqrt[c + a^2*c*x^2], x]

[Out] (I*2^(-1/2 - (I/2)*n)*(1 - I*a*x)^(1/2 + (I/2)*n)*Sqrt[1 + a^2*x^2]*(2^(1/2 + (I/2)*n)*(-I + n)*Sqrt[1 + I*a*x] - 4*n*(1 + I*a*x)^((I/2)*n)*Hypergeometric2F1[(1 + I*n)/2, (I/2)*(I + n), (3 + I*n)/2, (1 - I*a*x)/2])/(a^2*(1 + n^2)*(1 + I*a*x)^((I/2)*n)*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x}{\sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*arctan(a*x))*x/(a^2*c*x^2+c)^(1/2),x)`

[Out] `int(exp(n*arctan(a*x))*x/(a^2*c*x^2+c)^(1/2),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(x*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

[Out] `integral(x*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x e^{n \operatorname{atan}(ax)}}{\sqrt{c(a^2 x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))*x/(a**2*c*x**2+c)**(1/2),x)`

[Out] `Integral(x*exp(n*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{x e^{n \operatorname{atan}(a x)}}{\sqrt{c a^2 x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2), x)

[Out] int((x*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2), x)

$$3.356 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=120

$$\frac{2^{\frac{1}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2}(1+in)} \sqrt{1 + a^2 x^2} {}_2F_1\left(\frac{1}{2}(1 + in), \frac{1}{2}(1 + in); \frac{1}{2}(3 + in); \frac{1}{2}(1 - iax)\right)}{a(i - n) \sqrt{c + a^2 cx^2}}$$

[Out] $-2^{(1/2-1/2*I*n)}*(1-I*a*x)^{(1/2+1/2*I*n)}*\operatorname{hypergeom}\left(\left[\frac{1}{2}+1/2*I*n, \frac{1}{2}+1/2*I*n\right], \left[\frac{3}{2}+1/2*I*n\right], \frac{1}{2}-1/2*I*a*x\right)*(a^2*x^2+1)^{(1/2)}/a/(I-n)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.06, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{2^{\frac{1}{2} - \frac{in}{2}} \sqrt{a^2 x^2 + 1} (1 - iax)^{\frac{1}{2}(1+in)} {}_2F_1\left(\frac{1}{2}(in + 1), \frac{1}{2}(in + 1); \frac{1}{2}(in + 3); \frac{1}{2}(1 - iax)\right)}{a(-n + i) \sqrt{a^2 cx^2 + c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n*\operatorname{ArcTan}[a*x])}/\operatorname{Sqrt}[c + a^2*c*x^2], x\right]$

[Out] $-((2^{(1/2 - (I/2)*n)}*(1 - I*a*x)^{((1 + I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2]*\operatorname{Hypergeometric2F1}[(1 + I*n)/2, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/2])/(a*(I - n)*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 71

$\operatorname{Int}[(a + b*x)^{(m+1)}/(b*(m+1)*(b*(b*c - a*d))^{(n)})*\operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d)*(a + b*x)/(b*c - a*d)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, m, n\}, x$ && $\operatorname{NeQ}[b*c - a*d, 0]$ && $\operatorname{IntegerQ}[m]$ && $\operatorname{IntegerQ}[n]$ && $\operatorname{GtQ}[b/(b*c - a*d), 0]$ && $(\operatorname{RationalQ}[m] \mid \mid \operatorname{IntegerQ}[n] \mid \mid \operatorname{GtQ}[-d/(b*c - a*d), 0])$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a*x])*(n)}*((c + d*x^2)^{(p)}), x_{\text{Symbol}}] :>$ $\operatorname{Dist}[c^p, \operatorname{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, n, p\}, x$ && $\operatorname{EqQ}[d, a^2*c]$ && $(\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[c, 0])$

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a*x])*(n)}*((c + d*x^2)^{(p)}), x_{\text{Symbol}}] :>$ $\operatorname{Dist}[c^p*\operatorname{IntPart}[p]*((c + d*x^2)^{\operatorname{FracPart}[p]}/(1 + a^2*x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[(1 + a^2*x^2)^p*E^{(n*\operatorname{ArcTan}[a*x])}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, n, p\}, x$ && Eq

$Q[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= -\frac{2^{\frac{1}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2} + \frac{in}{2}} \sqrt{1 + a^2 x^2} {}_2F_1\left(\frac{1}{2}(1 + in), \frac{1}{2}(1 + in); \frac{1}{2}(3 + in); \frac{1}{2}(1 - iax)\right)}{a(i - n)\sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 117, normalized size = 0.98

$$\frac{2^{\frac{1}{2} - \frac{in}{2}} (1 - iax)^{\frac{1}{2} + \frac{in}{2}} \sqrt{1 + a^2 x^2} {}_2F_1\left(\frac{1}{2} + \frac{in}{2}, \frac{1}{2} + \frac{in}{2}; \frac{3}{2} + \frac{in}{2}; \frac{1}{2} - \frac{iax}{2}\right)}{a(-i + n)\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/Sqrt[c + a^2*c*x^2], x]

[Out] (2^(1/2 - (I/2)*n)*(1 - I*a*x)^(1/2 + (I/2)*n)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1/2 + (I/2)*n, 1/2 + (I/2)*n, 3/2 + (I/2)*n, 1/2 - (I/2)*a*x])/(a*(-I + n)*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{\sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(exp(n*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{\sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(1/2),x)

[Out] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(1/2), x)

$$3.357 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{x \sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=121

$$\frac{2(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(-1-in)}\sqrt{1 + a^2x^2} {}_2F_1\left(1, \frac{1}{2}(1 + in); \frac{1}{2}(3 + in); \frac{1-iax}{1+iax}\right)}{(1 + in)\sqrt{c + a^2cx^2}}$$

[Out] $-2*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(-1/2-1/2*I*n)}*\operatorname{hypergeom}([1, 1/2+1/2*I*n], [3/2+1/2*I*n], (1-I*a*x)/(1+I*a*x))*(a^2*x^2+1)^{(1/2)}/(1+I*n)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 121, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$,

Rules used = {5193, 5190, 133}

$$\frac{2\sqrt{a^2x^2 + 1}(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(-1-in)} {}_2F_1\left(1, \frac{1}{2}(in + 1); \frac{1}{2}(in + 3); \frac{1-iax}{iax+1}\right)}{(1 + in)\sqrt{a^2cx^2 + c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n \operatorname{ArcTan}[a*x])}/(x*\operatorname{Sqrt}[c + a^2*c*x^2]), x]$

[Out] $(-2*(1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((-1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2]*\operatorname{Hypergeometric2F1}[1, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/(1 + I*a*x)]/((1 + I*n)*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 133

$\operatorname{Int}[(a_. + (b_.)*(x_))^{(m_.)}*((c_.) + (d_.)*(x_))^{(n_.)}*((e_.) + (f_.)*(x_))^{(p_.)}, x_Symbol] \rightarrow \operatorname{Simp}[(b*c - a*d)^n*((a + b*x)^{(m+1)}/((m+1)*(b*e - a*f)^{(n+1)}*(e + f*x)^{(m+1)}))*\operatorname{Hypergeometric2F1}[m+1, -n, m+2, -(d*(e - c*f))/(b*c - a*d)/(e + f*x)], x] /;$ $\operatorname{FreeQ}\{a, b, c, d, e, f, m, p\}, x \ \&\& \operatorname{EqQ}[m + n + p + 2, 0] \ \&\& \operatorname{ILtQ}[n, 0] \ \&\& (\operatorname{SumSimplerQ}[m, 1] \ || \ !\operatorname{SumSimplerQ}[p, 1]) \ \&\& \ !\operatorname{ILtQ}[m, 0]$

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_.)*(x_)]*(n_.))}*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[x^m*(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, m, n, p\}, x \ \&\& \operatorname{EqQ}[d, a^2*c] \ \&\& (\operatorname{IntegerQ}[p] \ || \ \operatorname{GtQ}[c, 0])$

Rule 5193

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_.)*(x_)]*(n_.))}*(x_)^{(m_.)}*((c_.) + (d_.)*(x_)^2)^{(p_.)}, x_Symbol] \rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]}*((c + d*x^2)^{\operatorname{FracPart}[p]}/(1 + a^2*x^2)^{\operatorname{FracPart}[p]})$

[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{x \sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{x \sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{(1-iax)^{-\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}}}{x} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{2(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(-1-in)} \sqrt{1 + a^2 x^2} {}_2F_1\left(1, \frac{1}{2}(1+in); \frac{1}{2}(3+in); \frac{1-iax}{1+iax}\right)}{(1+in)\sqrt{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 120, normalized size = 0.99

$$\frac{2(1-iax)^{\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}} \sqrt{1 + a^2 x^2} {}_2F_1\left(1, \frac{1}{2} + \frac{in}{2}; \frac{3}{2} + \frac{in}{2}; \frac{1+iax}{1-iax}\right)}{(-1-in)\sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(x*Sqrt[c + a^2*c*x^2]), x]

[Out] (2*(1 - I*a*x)^(1/2 + (I/2)*n)*(1 + I*a*x)^(-1/2 - (I/2)*n)*Sqrt[1 + a^2*x^2]*Hypergeometric2F1[1, 1/2 + (I/2)*n, 3/2 + (I/2)*n, (I + a*x)/(I - a*x)]) /((-1 - I*n)*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{x \sqrt{a^2 cx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/x/(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(n*arctan(a*x))/x/(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))/x/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")`

[Out] `integrate(e^(n*arctan(a*x))/(sqrt(a^2*c*x^2 + c)*x), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))/x/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")`

[Out] `integral(sqrt(a^2*c*x^2 + c)*e^(n*arctan(a*x))/(a^2*c*x^3 + c*x), x)`

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x \sqrt{c(a^2 x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))/x/(a**2*c*x**2+c)**(1/2),x)`

[Out] `Integral(exp(n*atan(a*x))/(x*sqrt(c*(a**2*x**2 + 1))), x)`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))/x/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x \sqrt{c a^2 x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int(exp(n*atan(a*x))/(x*(c + a^2*c*x^2)^(1/2)),x)`

[Out] `int(exp(n*atan(a*x))/(x*(c + a^2*c*x^2)^(1/2)), x)`

$$3.358 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{x^2 \sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=196

$$\frac{(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{x\sqrt{c + a^2cx^2}} - \frac{2an(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{(1 + in)\sqrt{c + a^2cx^2}} {}_2F_1\left(1, \frac{1}{2}(1 + in)\right)$$

[Out] $-(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*(a^2*x^2+1)^{(1/2)}/x/(a^2*c*x^2+c)^{(1/2)}-2*a*n*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(-1/2-1/2*I*n)}*\operatorname{hypergeom}([1, 1/2+1/2*I*n], [3/2+1/2*I*n], (1-I*a*x)/(1+I*a*x))*(a^2*x^2+1)^{(1/2)}/(1+I*n)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 196, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 4, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.154$, Rules used = {5193, 5190, 98, 133}

$$\frac{2an\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}{}_2F_1\left(1, \frac{1}{2}(in+1); \frac{1}{2}(in+3); \frac{1-iax}{iax+1}\right)}{(1+in)\sqrt{a^2cx^2+c}} - \frac{\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}}{x\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n \operatorname{ArcTan}[a*x])}/(x^2 \operatorname{Sqrt}[c + a^2*c*x^2]), x]$

[Out] $-(((1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2])/(x*\operatorname{Sqrt}[c + a^2*c*x^2])) - (2*a*n*(1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((-1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2]*\operatorname{Hypergeometric2F1}[1, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/(1 + I*a*x)])/((1 + I*n)*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 98

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \operatorname{Simp}[b*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)}/((m + 1)*(b*c - a*d)*(b*e - a*f))), x] + \operatorname{Dist}[(a*d*f*(m + 1) + b*c*f*(n + 1) + b*d*e*(p + 1))/((m + 1)*(b*c - a*d)*(b*e - a*f)), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, n, p\}, x] \&\& \operatorname{EqQ}[\operatorname{Simplify}[m + n + p + 3], 0] \&\& (\operatorname{LtQ}[m, -1] || \operatorname{SumSimplerQ}[m, 1])$

Rule 133

$\operatorname{Int}[(a_. + (b_.)*(x_.))^{(m_.)}*((c_.) + (d_.)*(x_.))^{(n_.)}*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \operatorname{Simp}[(b*c - a*d)^n*((a + b*x)^{(m + 1)}/((m + 1)*(b*e - a*f)^{(n + 1)}*(e + f*x)^{(m + 1)}))*\operatorname{Hypergeometric2F1}[m + 1, -n, m + 2, -(d*(e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, m, p\}, x] \&\& \operatorname{EqQ}[m + n + p + 2, 0] \&\& (\operatorname{ILtQ}[n, 0] \&\& (\operatorname{SumSimplerQ}[m, 1]$

|| !SumSimplerQ[p, 1]) && !ILtQ[m, 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{x^2 \sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{x^2 \sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{(1-iax)^{-\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}}}{x^2} dx}{\sqrt{c + a^2 cx^2}} \\ &= -\frac{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{x \sqrt{c + a^2 cx^2}} + \frac{\left(an \sqrt{1 + a^2 x^2} \right) \int \frac{(1-iax)^{-\frac{1}{2} + \frac{in}{2}} (1+iax)^{\frac{1}{2} - \frac{in}{2}}}{x}}{\sqrt{c + a^2 cx^2}} \\ &= -\frac{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{x \sqrt{c + a^2 cx^2}} - \frac{2an(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)}}{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)}} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 142, normalized size = 0.72

$$\frac{(1-iax)^{\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}} \sqrt{1 + a^2 x^2} \left(-((-i+n)(-i+ax)) + 2anx {}_2F_1\left(1, \frac{1}{2} + \frac{in}{2}; \frac{3}{2} + \frac{in}{2}; \frac{i+ax}{i-ax}\right) \right)}{(-1-in)x \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(x^2*Sqrt[c + a^2*c*x^2]),x]

[Out] ((1 - I*a*x)^(1/2 + (I/2)*n)*(1 + I*a*x)^(-1/2 - (I/2)*n)*Sqrt[1 + a^2*x^2] *(-((-I + n)*(-I + a*x)) + 2*a*n*x*Hypergeometric2F1[1, 1/2 + (I/2)*n, 3/2 + (I/2)*n, (I + a*x)/(I - a*x)])/((-1 - I*n)*x*Sqrt[c + a^2*c*x^2])

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{x^2 \sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c)^(1/2),x)

[Out] int(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c)^(1/2),x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/(sqrt(a^2*c*x^2 + c)*x^2), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(a^2*c*x^2 + c)*e^(n*arctan(a*x))/(a^2*c*x^4 + c*x^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x^2 \sqrt{c(a^2 x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/x**2/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(exp(n*atan(a*x))/(x**2*sqrt(c*(a**2*x**2 + 1))), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*arctan(a*x))/x^2/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x^2 \sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(n*atan(a*x))/(x^2*(c + a^2*c*x^2)^(1/2)),x)
```

```
[Out] int(exp(n*atan(a*x))/(x^2*(c + a^2*c*x^2)^(1/2)), x)
```

$$3.359 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{x^3 \sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=281

$$\frac{(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{2x^2\sqrt{c + a^2cx^2}} - \frac{an(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}\sqrt{1 + a^2x^2}}{2x\sqrt{c + a^2cx^2}} + \frac{a^2(1 - n^2)(1 - iax)^{\frac{1}{2}(1+in)}(1 + iax)^{\frac{1}{2}(1-in)}}{2x\sqrt{c + a^2cx^2}}$$

[Out] $-1/2*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*(a^2*x^2+1)^{(1/2)}/x^2/(a^2*c*x^2+c)^{(1/2)}-1/2*a*n*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(1/2-1/2*I*n)}*(a^2*x^2+1)^{(1/2)}/x/(a^2*c*x^2+c)^{(1/2)}+a^2*(-n^2+1)*(1-I*a*x)^{(1/2+1/2*I*n)}*(1+I*a*x)^{(-1/2-1/2*I*n)}*\operatorname{hypergeom}([1, 1/2+1/2*I*n], [3/2+1/2*I*n], (1-I*a*x)/(1+I*a*x))*(a^2*x^2+1)^{(1/2)}/(1+I*n)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.18, antiderivative size = 281, normalized size of antiderivative = 1.00, number of steps used = 6, number of rules used = 6, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.231$, Rules used = {5193, 5190, 105, 156, 12, 133}

$$\frac{a^2(1-n^2)\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}{}_2F_1\left(1, \frac{1}{2}(in+1); \frac{1}{2}(in+3); \frac{1-iax}{iax+1}\right)}{(1+in)\sqrt{a^2cx^2+c}} - \frac{an\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}}{2x\sqrt{a^2cx^2+c}} - \frac{\sqrt{a^2x^2+1}(1-iax)^{\frac{1}{2}(1+in)}(1+iax)^{\frac{1}{2}(1-in)}}{2x^2\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n*\operatorname{ArcTan}[a*x])}/(x^3*\operatorname{Sqrt}[c + a^2*c*x^2]), x]$

[Out] $-1/2*((1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2])/(x^2*\operatorname{Sqrt}[c + a^2*c*x^2]) - (a*n*(1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2])/(2*x*\operatorname{Sqrt}[c + a^2*c*x^2]) + (a^2*(1 - n^2)*(1 - I*a*x)^{((1 + I*n)/2)}*(1 + I*a*x)^{((-1 - I*n)/2)}*\operatorname{Sqrt}[1 + a^2*x^2]*\operatorname{Hypergeometric2F1}[1, (1 + I*n)/2, (3 + I*n)/2, (1 - I*a*x)/(1 + I*a*x)])/(1 + I*n)*\operatorname{Sqrt}[c + a^2*c*x^2]$

Rule 12

$\operatorname{Int}[(a_*)(u_), x_Symbol] := \operatorname{Dist}[a, \operatorname{Int}[u, x], x] /; \operatorname{FreeQ}[a, x] \&\& !\operatorname{MatchQ}[u, (b_)*(v_)] /; \operatorname{FreeQ}[b, x]$

Rule 105

$\operatorname{Int}[(a_*)(x_*)^{(m_*)}*((c_*) + (d_*)(x_*))^{(n_*)}*((e_*) + (f_*)(x_*))^{(p_*)}, x_Symbol] := \operatorname{Simp}[b*(a + b*x)^{(m + 1)}*(c + d*x)^{(n + 1)}*((e + f*x)^{(p + 1)})/((m + 1)*(b*c - a*d)*(b*e - a*f)), x] + \operatorname{Dist}[1/((m + 1)*(b*c - a*d)*(b*e - a*f)), \operatorname{Int}[(a + b*x)^{(m + 1)}*(c + d*x)^n*(e + f*x)^p*\operatorname{Simp}[a*d*f*(m + 1) - b*(d*e*(m + n + 2) + c*f*(m + p + 2)) - b*d*f*(m + n + p + 3)*x, x], x] /; \operatorname{FreeQ}\{a, b, c, d, e, f, n, p\}, x] \&\& \operatorname{ILtQ}[m, -1] \&\& (\operatorname{IntegerQ}[n] || \operatorname{IntegersQ}[2*n, 2*p] || \operatorname{ILtQ}[m + n + p + 3, 0])$

Rule 133

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_), x_Symbol] :> Simp[(b*c - a*d)^(n+1)*((a + b*x)^(m+1)/((m+1)*(b*e - a*f)^(n+1)*(e + f*x)^(m+1)))*Hypergeometric2F1[m+1, -n, m+2, (-(d*e - c*f))*((a + b*x)/((b*c - a*d)*(e + f*x)))]], x] /; FreeQ[{a, b, c, d, e, f, m, p}, x] && EqQ[m + n + p + 2, 0] && ILtQ[n, 0] && (SumSimplerQ[m, 1] || !SumSimplerQ[p, 1]) && !ILtQ[m, 0]
```

Rule 156

```
Int[((a_.) + (b_.)*(x_))^(m_)*((c_.) + (d_.)*(x_))^(n_)*((e_.) + (f_.)*(x_))^(p_)*((g_.) + (h_.)*(x_)), x_Symbol] :> Simp[(b*g - a*h)*(a + b*x)^(m+1)*(c + d*x)^(n+1)*(e + f*x)^(p+1)/((m+1)*(b*c - a*d)*(b*e - a*f))], x] + Dist[1/((m+1)*(b*c - a*d)*(b*e - a*f)), Int[(a + b*x)^(m+1)*(c + d*x)^(n+1)*(e + f*x)^p*Simp[(a*d*f*g - b*(d*e + c*f)*g + b*c*e*h)*(m+1) - (b*g - a*h)*(d*e*(n+1) + c*f*(p+1)) - d*f*(b*g - a*h)*(m+n+p+3)*x], x], x] /; FreeQ[{a, b, c, d, e, f, g, h, n, p}, x] && ILtQ[m, -1]
```

Rule 5190

```
Int[E^(ArcTan[(a_.)*(x_)])^(n_.)*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2))], x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])
```

Rule 5193

```
Int[E^(ArcTan[(a_.)*(x_)])^(n_.)*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])
```

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)}}{x^3 \sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{x^3 \sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int \frac{(1-iax)^{-\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}}}{x^3} dx}{\sqrt{c + a^2 cx^2}} \\
&= -\frac{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x^2 \sqrt{c + a^2 cx^2}} - \frac{\sqrt{1 + a^2 x^2} \int \frac{(1-iax)^{-\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}}}{x^2} dx}{2\sqrt{c + a^2 cx^2}} \\
&= -\frac{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x^2 \sqrt{c + a^2 cx^2}} - \frac{an(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x \sqrt{c + a^2 cx^2}} \\
&= -\frac{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x^2 \sqrt{c + a^2 cx^2}} - \frac{an(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x \sqrt{c + a^2 cx^2}} \\
&= -\frac{(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x^2 \sqrt{c + a^2 cx^2}} - \frac{an(1-iax)^{\frac{1}{2}(1+in)} (1+iax)^{\frac{1}{2}(1-in)} \sqrt{1 + a^2 x^2}}{2x \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [A]

time = 0.06, size = 159, normalized size = 0.57

$$\frac{i(1-iax)^{\frac{1}{2} + \frac{in}{2}} (1+iax)^{-\frac{1}{2} - \frac{in}{2}} \sqrt{1 + a^2 x^2} (-(i+n)(-i+ax)(1+anx)) + 2a^2(-1+n^2)x^2 {}_2F_1\left(1, \frac{1}{2} + \frac{in}{2}; \frac{3}{2} + \frac{in}{2}; \frac{i+ax}{i-ax}\right)}{2(-i+n)x^2 \sqrt{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(n*ArcTan[a*x])/(x^3*Sqrt[c + a^2*c*x^2]),x]`

```
[Out] ((I/2)*(1 - I*a*x)^(1/2 + (I/2)*n)*(1 + I*a*x)^(-1/2 - (I/2)*n)*Sqrt[1 + a^2*x^2]*(-((-I + n)*(-I + a*x)*(1 + a*n*x)) + 2*a^2*(-1 + n^2)*x^2*Hypergeometric2F1[1, 1/2 + (I/2)*n, 3/2 + (I/2)*n, (I + a*x)/(I - a*x)]))/((-I + n)*x^2*Sqrt[c + a^2*c*x^2])
```

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{x^3 \sqrt{a^2 cx^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c)^(1/2),x)``[Out] int(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c)^(1/2),x)`

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c)^(1/2),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/(sqrt(a^2*c*x^2 + c)*x^3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")

[Out] integral(sqrt(a^2*c*x^2 + c)*e^(n*arctan(a*x))/(a^2*c*x^5 + c*x^3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x^3 \sqrt{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/x**3/(a**2*c*x**2+c)**(1/2),x)

[Out] Integral(exp(n*atan(a*x))/(x**3*sqrt(c*(a**2*x**2 + 1))), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/x^3/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{x^3 \sqrt{ca^2x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(x^3*(c + a^2*c*x^2)^(1/2)),x)

[Out] int(exp(n*atan(a*x))/(x^3*(c + a^2*c*x^2)^(1/2)), x)

3.360 $\int e^{n \operatorname{ArcTan}(ax)} \sqrt[3]{c + a^2 cx^2} dx$

Optimal. Leaf size=120

$$\frac{3 \cdot 2^{\frac{4}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(8+3in)} \sqrt[3]{c + a^2 cx^2} {}_2F_1\left(\frac{1}{6}(-2 + 3in), \frac{1}{6}(8 + 3in); \frac{1}{6}(14 + 3in); \frac{1}{2}(1 - iax)\right)}{a(8i - 3n) \sqrt[3]{1 + a^2 x^2}}$$

[Out] $-3 \cdot 2^{(4/3 - 1/2 \cdot I \cdot n)} \cdot (1 - I \cdot a \cdot x)^{(4/3 + 1/2 \cdot I \cdot n)} \cdot (a^2 \cdot c \cdot x^2 + c)^{(1/3)} \cdot \operatorname{hypergeom}\left(\frac{4}{3} + \frac{1}{2} \cdot I \cdot n, -\frac{1}{3} + \frac{1}{2} \cdot I \cdot n, \frac{7}{3} + \frac{1}{2} \cdot I \cdot n, \frac{1}{2} - \frac{1}{2} \cdot I \cdot a \cdot x\right) / a / (8 \cdot I - 3 \cdot n) / (a^2 \cdot x^2 + 1)^{(1/3)}$

Rubi [A]

time = 0.08, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{3 \cdot 2^{\frac{4}{3} - \frac{in}{2}} \sqrt[3]{a^2 cx^2 + c} (1 - iax)^{\frac{1}{6}(8+3in)} {}_2F_1\left(\frac{1}{6}(3in - 2), \frac{1}{6}(3in + 8); \frac{1}{6}(3in + 14); \frac{1}{2}(1 - iax)\right)}{a(-3n + 8i) \sqrt[3]{a^2 x^2 + 1}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[E^{(n \cdot \operatorname{ArcTan}[a \cdot x])} \cdot (c + a^2 \cdot c \cdot x^2)^{(1/3)}, x]$

[Out] $(-3 \cdot 2^{(4/3 - (I/2) \cdot n)} \cdot (1 - I \cdot a \cdot x)^{((8 + (3 \cdot I) \cdot n)/6)} \cdot (c + a^2 \cdot c \cdot x^2)^{(1/3)} \cdot \operatorname{Hypergeometric2F1}[-2 + (3 \cdot I) \cdot n/6, (8 + (3 \cdot I) \cdot n)/6, (14 + (3 \cdot I) \cdot n)/6, (1 - I \cdot a \cdot x)/2]) / (a \cdot (8 \cdot I - 3 \cdot n) \cdot (1 + a^2 \cdot x^2)^{(1/3)})$

Rule 71

$\operatorname{Int}[(a + b \cdot x)^{(m)} \cdot ((c + d \cdot x)^{(n)})], x_{\text{Symbol}}] \rightarrow \operatorname{Simp}[(a + b \cdot x)^{(m+1)} / (b \cdot (m+1) \cdot (b \cdot (b \cdot c - a \cdot d))^n) \cdot \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d) \cdot ((a + b \cdot x) / (b \cdot c - a \cdot d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a + b \cdot x) \cdot (x)] \cdot (n))} \cdot ((c + d \cdot x)^2)^{(p)}], x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1 - I \cdot a \cdot x)^{(p + I \cdot (n/2))} \cdot (1 + I \cdot a \cdot x)^{(p - I \cdot (n/2))}], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a + b \cdot x) \cdot (x)] \cdot (n))} \cdot ((c + d \cdot x)^2)^{(p)}], x_{\text{Symbol}}] \rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]} \cdot ((c + d \cdot x^2)^{\operatorname{FracPart}[p]} / (1 + a^2 \cdot x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[(1 + a^2 \cdot x^2)^p \cdot E^{(n \cdot \operatorname{ArcTan}[a \cdot x])}], x] /;$ FreeQ[{a, c, d, n, p}, x] && Eq

$\text{Q}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} \sqrt[3]{c + a^2cx^2} \, dx &= \frac{\sqrt[3]{c + a^2cx^2} \int e^{n \tan^{-1}(ax)} \sqrt[3]{1 + a^2x^2} \, dx}{\sqrt[3]{1 + a^2x^2}} \\ &= \frac{\sqrt[3]{c + a^2cx^2} \int (1 - iax)^{\frac{1}{3} + \frac{in}{2}} (1 + iax)^{\frac{1}{3} - \frac{in}{2}} \, dx}{\sqrt[3]{1 + a^2x^2}} \\ &= -\frac{3 \ 2^{\frac{4}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(8+3in)} \sqrt[3]{c + a^2cx^2} \ {}_2F_1\left(\frac{1}{6}(-2 + 3in), \frac{1}{6}(8 + 3in); \frac{1}{6}(14 + 3in); -\frac{iax}{1 + a^2x^2}\right)}{a(8i - 3n)\sqrt[3]{1 + a^2x^2}} \end{aligned}$$

Mathematica [A]

time = 0.05, size = 120, normalized size = 1.00

$$\frac{3 \ 2^{\frac{4}{3} - \frac{in}{2}} (1 - iax)^{\frac{4}{3} + \frac{in}{2}} \sqrt[3]{c + a^2cx^2} \ {}_2F_1\left(-\frac{1}{3} + \frac{in}{2}, \frac{4}{3} + \frac{in}{2}, \frac{7}{3} + \frac{in}{2}, \frac{1}{2} - \frac{iax}{2}\right)}{a(-8i + 3n)\sqrt[3]{1 + a^2x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])*(c + a^2*c*x^2)^(1/3), x]

[Out] (3*2^(4/3 - (I/2)*n)*(1 - I*a*x)^(4/3 + (I/2)*n)*(c + a^2*c*x^2)^(1/3)*Hypergeometric2F1[-1/3 + (I/2)*n, 4/3 + (I/2)*n, 7/3 + (I/2)*n, 1/2 - (I/2)*a*x])/ (a*(-8*I + 3*n)*(1 + a^2*x^2)^(1/3))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} (a^2cx^2 + c)^{\frac{1}{3}} \, dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/3), x)

[Out] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/3),x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(1/3)*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/3),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(1/3)*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \sqrt[3]{c(a^2x^2 + 1)} e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*(a**2*c*x**2+c)**(1/3),x)

[Out] Integral((c*(a**2*x**2 + 1))**(1/3)*exp(n*atan(a*x)), x)

Giac [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^(1/3),x, algorithm="giac")

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx);;OUTPUT:sym2poly/r2sym(const gen & e,const in
dex_m & i,const vecteur & l) Error: Bad Argument Value

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} (ca^2x^2 + c)^{1/3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^(1/3),x)

[Out] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^(1/3), x)

$$3.361 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{\sqrt[3]{c + a^2 cx^2}} dx$$

Optimal. Leaf size=120

$$\frac{3 \cdot 2^{\frac{2}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(4+3in)} \sqrt[3]{1 + a^2 x^2} {}_2F_1\left(\frac{1}{6}(2 + 3in), \frac{1}{6}(4 + 3in); \frac{1}{6}(10 + 3in); \frac{1}{2}(1 - iax)\right)}{a(4i - 3n) \sqrt[3]{c + a^2 cx^2}}$$

[Out] $-3 \cdot 2^{(2/3 - 1/2 \cdot I \cdot n)} \cdot (1 - I \cdot a \cdot x)^{(2/3 + 1/2 \cdot I \cdot n)} \cdot (a^2 \cdot x^2 + 1)^{(1/3)} \cdot \operatorname{hypergeom}\left(\left[\frac{1}{3} + \frac{1}{2} \cdot I \cdot n, \frac{2}{3} + \frac{1}{2} \cdot I \cdot n\right], \left[\frac{5}{3} + \frac{1}{2} \cdot I \cdot n\right], \frac{1}{2} - \frac{1}{2} \cdot I \cdot a \cdot x\right) / a / (4 \cdot I - 3 \cdot n) / (a^2 \cdot c \cdot x^2 + c)^{(1/3)}$

Rubi [A]

time = 0.08, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{3 \cdot 2^{\frac{2}{3} - \frac{in}{2}} \sqrt[3]{a^2 x^2 + 1} (1 - iax)^{\frac{1}{6}(4+3in)} {}_2F_1\left(\frac{1}{6}(3in + 2), \frac{1}{6}(3in + 4); \frac{1}{6}(3in + 10); \frac{1}{2}(1 - iax)\right)}{a(-3n + 4i) \sqrt[3]{a^2 cx^2 + c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n \cdot \operatorname{ArcTan}[a \cdot x])} / (c + a^2 \cdot c \cdot x^2)^{(1/3)}, x\right]$

[Out] $(-3 \cdot 2^{(2/3 - (I/2) \cdot n)} \cdot (1 - I \cdot a \cdot x)^{((4 + (3 \cdot I) \cdot n)/6)} \cdot (1 + a^2 \cdot x^2)^{(1/3)} \cdot \operatorname{Hypergeometric2F1}\left[\frac{2 + (3 \cdot I) \cdot n}{6}, \frac{4 + (3 \cdot I) \cdot n}{6}, \frac{10 + (3 \cdot I) \cdot n}{6}, \frac{1 - I \cdot a \cdot x}{2}\right]) / (a \cdot (4 \cdot I - 3 \cdot n) \cdot (c + a^2 \cdot c \cdot x^2)^{(1/3)})$

Rule 71

$\operatorname{Int}[(a_ + (b_ \cdot (x_))^m) \cdot ((c_ + (d_ \cdot (x_))^n), x_Symbol] \rightarrow \operatorname{Simp}[(a + b \cdot x)^{(m+1)} / (b \cdot (m+1) \cdot (b/(b \cdot c - a \cdot d))^n) \cdot \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d) \cdot (a + b \cdot x) / (b \cdot c - a \cdot d)], x] / ; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \operatorname{NeQ}\{b \cdot c - a \cdot d, 0\} \&\& !\operatorname{IntegerQ}\{m\} \&\& !\operatorname{IntegerQ}\{n\} \&\& \operatorname{GtQ}\{b/(b \cdot c - a \cdot d), 0\} \&\& (\operatorname{RationalQ}\{m\} \mid \mid !(\operatorname{RationalQ}\{n\} \&\& \operatorname{GtQ}\{-d/(b \cdot c - a \cdot d), 0\}))$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a \cdot x]) \cdot (n \cdot x)} \cdot ((c + (d \cdot x)^2)^p), x_Symbol] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1 - I \cdot a \cdot x)^{(p + I \cdot (n/2))} \cdot (1 + I \cdot a \cdot x)^{(p - I \cdot (n/2))}, x], x] / ; \operatorname{FreeQ}\{a, c, d, n, p\}, x] \&\& \operatorname{EqQ}\{d, a^2 \cdot c\} \&\& (\operatorname{IntegerQ}\{p\} \mid \mid \operatorname{GtQ}\{c, 0\})$

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a \cdot x]) \cdot (n \cdot x)} \cdot ((c + (d \cdot x)^2)^p), x_Symbol] \rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]} \cdot (c + d \cdot x^2)^{\operatorname{FracPart}[p]} / (1 + a^2 \cdot x^2)^{\operatorname{FracPart}[p]}, \operatorname{Int}[(1 + a^2 \cdot x^2)^p \cdot E^{(n \cdot \operatorname{ArcTan}[a \cdot x])}, x], x] / ; \operatorname{FreeQ}\{a, c, d, n, p\}, x] \&\& \operatorname{Eq}$

`Q[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])`

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{\sqrt[3]{c + a^2 cx^2}} dx &= \frac{\sqrt[3]{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{\sqrt[3]{1 + a^2 x^2}} dx}{\sqrt[3]{c + a^2 cx^2}} \\ &= \frac{\sqrt[3]{1 + a^2 x^2} \int (1 - iax)^{-\frac{1}{3} + \frac{in}{2}} (1 + iax)^{-\frac{1}{3} - \frac{in}{2}} dx}{\sqrt[3]{c + a^2 cx^2}} \\ &= -\frac{3 \cdot 2^{\frac{2}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(4+3in)} \sqrt[3]{1 + a^2 x^2} {}_2F_1\left(\frac{1}{6}(2 + 3in), \frac{1}{6}(4 + 3in); \frac{1}{6}(10 + 3in); \frac{1}{2}(1 - iax)\right)}{a(4i - 3n)\sqrt[3]{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 120, normalized size = 1.00

$$\frac{3 \cdot 2^{\frac{2}{3} - \frac{in}{2}} (1 - iax)^{\frac{2}{3} + \frac{in}{2}} \sqrt[3]{1 + a^2 x^2} {}_2F_1\left(\frac{1}{3} + \frac{in}{2}, \frac{2}{3} + \frac{in}{2}; \frac{5}{3} + \frac{in}{2}; \frac{1}{2} - \frac{iax}{2}\right)}{a(-4i + 3n)\sqrt[3]{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(c + a^2*c*x^2)^(1/3), x]

[Out] (3*2^(2/3 - (I/2)*n)*(1 - I*a*x)^(2/3 + (I/2)*n)*(1 + a^2*x^2)^(1/3)*Hypergeometric2F1[1/3 + (I/2)*n, 2/3 + (I/2)*n, 5/3 + (I/2)*n, 1/2 - (I/2)*a*x])/(a*(-4*I + 3*n)*(c + a^2*c*x^2)^(1/3))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{(a^2 c x^2 + c)^{\frac{1}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/3), x)

[Out] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/3),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(1/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/3),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(1/3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{\sqrt[3]{c(a^2x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/(a**2*c*x**2+c)**(1/3),x)

[Out] Integral(exp(n*atan(a*x))/(c*(a**2*x**2 + 1))**(1/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(1/3),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{(ca^2x^2 + c)^{1/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(1/3),x)

[Out] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(1/3), x)

$$3.362 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{2/3}} dx$$

Optimal. Leaf size=120

$$\frac{3 \cdot 2^{\frac{1}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(2+3in)} (1 + a^2x^2)^{2/3} {}_2F_1\left(\frac{1}{6}(2+3in), \frac{1}{6}(4+3in); \frac{1}{6}(8+3in); \frac{1}{2}(1-iax)\right)}{a(2i-3n)(c+a^2cx^2)^{2/3}}$$

[Out] $-3 \cdot 2^{(1/3-1/2 \cdot I \cdot n)} \cdot (1-I \cdot a \cdot x)^{(1/3+1/2 \cdot I \cdot n)} \cdot (a^2 \cdot x^2+1)^{(2/3)} \cdot \operatorname{hypergeom}\left(\left[\frac{1}{3}+1/2 \cdot I \cdot n, \frac{2}{3}+1/2 \cdot I \cdot n\right], \left[\frac{4}{3}+1/2 \cdot I \cdot n\right], \frac{1}{2}-1/2 \cdot I \cdot a \cdot x\right) / a / (2 \cdot I-3 \cdot n) / (a^2 \cdot c \cdot x^2+c)^{(2/3)}$

Rubi [A]

time = 0.08, antiderivative size = 120, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{3 \cdot 2^{\frac{1}{3} - \frac{in}{2}} (a^2x^2 + 1)^{2/3} (1 - iax)^{\frac{1}{6}(2+3in)} {}_2F_1\left(\frac{1}{6}(3in+2), \frac{1}{6}(3in+4); \frac{1}{6}(3in+8); \frac{1}{2}(1-iax)\right)}{a(-3n+2i)(a^2cx^2+c)^{2/3}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n \cdot \operatorname{ArcTan}[a \cdot x])} / (c + a^2 \cdot c \cdot x^2)^{(2/3)}, x\right]$

[Out] $(-3 \cdot 2^{(1/3 - (I/2) \cdot n)} \cdot (1 - I \cdot a \cdot x)^{((2 + (3 \cdot I) \cdot n)/6)} \cdot (1 + a^2 \cdot x^2)^{(2/3)} \cdot \operatorname{Hypergeometric2F1}\left[\frac{2 + (3 \cdot I) \cdot n}{6}, \frac{4 + (3 \cdot I) \cdot n}{6}, \frac{8 + (3 \cdot I) \cdot n}{6}, (1 - I \cdot a \cdot x)/2\right]) / (a \cdot (2 \cdot I - 3 \cdot n) \cdot (c + a^2 \cdot c \cdot x^2)^{(2/3)})$

Rule 71

$\operatorname{Int}[(a + b \cdot x)^{(m)} \cdot ((c + d \cdot x)^{(n)})], x_Symbol] \rightarrow \operatorname{Simp}[(a + b \cdot x)^{(m+1)} / (b \cdot (m+1) \cdot (b \cdot (b \cdot c - a \cdot d))^{(n)}) \cdot \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d) \cdot ((a + b \cdot x) / (b \cdot c - a \cdot d))], x] /;$ FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a \cdot x] \cdot (n))} \cdot ((c + d \cdot x)^2)^{(p)}], x_Symbol] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1 - I \cdot a \cdot x)^{(p + I \cdot (n/2))} \cdot (1 + I \cdot a \cdot x)^{(p - I \cdot (n/2))}], x] /;$ FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a \cdot x] \cdot (n))} \cdot ((c + d \cdot x)^2)^{(p)}], x_Symbol] \rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]} \cdot ((c + d \cdot x^2)^{\operatorname{FracPart}[p]} / (1 + a^2 \cdot x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[(1 + a^2 \cdot x^2)^p \cdot E^{(n \cdot \operatorname{ArcTan}[a \cdot x])}], x] /;$ FreeQ[{a, c, d, n, p}, x] && Eq

Q[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{(c + a^2 cx^2)^{2/3}} dx &= \frac{(1 + a^2 x^2)^{2/3} \int \frac{e^{n \tan^{-1}(ax)}}{(1 + a^2 x^2)^{2/3}} dx}{(c + a^2 cx^2)^{2/3}} \\ &= \frac{(1 + a^2 x^2)^{2/3} \int (1 - iax)^{-\frac{2}{3} + \frac{in}{2}} (1 + iax)^{-\frac{2}{3} - \frac{in}{2}} dx}{(c + a^2 cx^2)^{2/3}} \\ &= -\frac{3 \cdot 2^{\frac{1}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(2+3in)} (1 + a^2 x^2)^{2/3} {}_2F_1\left(\frac{1}{6}(2 + 3in), \frac{1}{6}(4 + 3in); \frac{1}{6}(8 + 3in); \frac{1}{2}(1 - iax)\right)}{a(2i - 3n)(c + a^2 cx^2)^{2/3}} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 120, normalized size = 1.00

$$\frac{3 \cdot 2^{\frac{1}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{3} + \frac{in}{2}} (1 + a^2 x^2)^{2/3} {}_2F_1\left(\frac{1}{3} + \frac{in}{2}, \frac{2}{3} + \frac{in}{2}; \frac{4}{3} + \frac{in}{2}; \frac{1}{2} - \frac{iax}{2}\right)}{a(-2i + 3n)(c + a^2 cx^2)^{2/3}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(c + a^2*c*x^2)^(2/3), x]

[Out] (3*2^(1/3 - (I/2)*n)*(1 - I*a*x)^(1/3 + (I/2)*n)*(1 + a^2*x^2)^(2/3)*Hypergeometric2F1[1/3 + (I/2)*n, 2/3 + (I/2)*n, 4/3 + (I/2)*n, 1/2 - (I/2)*a*x])/(a*(-2*I + 3*n)*(c + a^2*c*x^2)^(2/3))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{(a^2 c x^2 + c)^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(2/3), x)

[Out] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(2/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(2/3),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(2/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(2/3),x, algorithm="fricas")

[Out] integral(e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(2/3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{2}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/(a**2*c*x**2+c)**(2/3),x)

[Out] Integral(exp(n*atan(a*x))/(c*(a**2*x**2 + 1))**(2/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(2/3),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{(ca^2x^2 + c)^{2/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(2/3),x)

[Out] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(2/3), x)

$$3.363 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)}}{(c+a^2cx^2)^{4/3}} dx$$

Optimal. Leaf size=123

$$\frac{3 \cdot 2^{-\frac{1}{3}-\frac{in}{2}} (1-iax)^{\frac{1}{6}(-2+3in)} \sqrt[3]{1+a^2x^2} {}_2F_1\left(\frac{1}{6}(-2+3in), \frac{1}{6}(8+3in); \frac{1}{6}(4+3in); \frac{1}{2}(1-iax)\right)}{ac(2i+3n)\sqrt[3]{c+a^2cx^2}}$$

[Out] $3 \cdot 2^{(-1/3-1/2 \cdot I \cdot n)} \cdot (1-I \cdot a \cdot x)^{(-1/3+1/2 \cdot I \cdot n)} \cdot (a^2 \cdot x^2+1)^{(1/3)} \cdot \operatorname{hypergeom}\left(\left[\frac{4}{3}+1/2 \cdot I \cdot n, -1/3+1/2 \cdot I \cdot n\right], \left[\frac{2}{3}+1/2 \cdot I \cdot n\right], \frac{1}{2}-1/2 \cdot I \cdot a \cdot x\right) / a / c / (2 \cdot I+3 \cdot n) / (a^2 \cdot c \cdot x^2+c)^{(1/3)}$

Rubi [A]

time = 0.09, antiderivative size = 123, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 23, $\frac{\text{number of rules}}{\text{integrand size}} = 0.130$, Rules used = {5184, 5181, 71}

$$\frac{3 \cdot 2^{-\frac{1}{3}-\frac{in}{2}} \sqrt[3]{a^2x^2+1} (1-iax)^{\frac{1}{6}(-2+3in)} {}_2F_1\left(\frac{1}{6}(3in-2), \frac{1}{6}(3in+8); \frac{1}{6}(3in+4); \frac{1}{2}(1-iax)\right)}{ac(3n+2i)\sqrt[3]{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}\left[E^{(n \cdot \operatorname{ArcTan}[a \cdot x])} / (c + a^2 \cdot c \cdot x^2)^{(4/3)}, x\right]$

[Out] $(3 \cdot 2^{(-1/3 - (I/2) \cdot n)} \cdot (1 - I \cdot a \cdot x)^{((-2 + (3 \cdot I) \cdot n)/6)} \cdot (1 + a^2 \cdot x^2)^{(1/3)} \cdot \operatorname{Hypergeometric2F1}\left[(-2 + (3 \cdot I) \cdot n)/6, (8 + (3 \cdot I) \cdot n)/6, (4 + (3 \cdot I) \cdot n)/6, (1 - I \cdot a \cdot x)/2\right]) / (a \cdot c \cdot (2 \cdot I + 3 \cdot n) \cdot (c + a^2 \cdot c \cdot x^2)^{(1/3)})$

Rule 71

$\operatorname{Int}[(a_ + (b_ \cdot (x_))^m) \cdot ((c_ + (d_ \cdot (x_))^n), x_Symbol] \rightarrow \operatorname{Simp}[(a + b \cdot x)^{(m+1)} / (b \cdot (m+1) \cdot (b/(b \cdot c - a \cdot d))^n) \cdot \operatorname{Hypergeometric2F1}[-n, m+1, m+2, (-d) \cdot (a + b \cdot x) / (b \cdot c - a \cdot d)], x] / ; \operatorname{FreeQ}\{a, b, c, d, m, n\}, x] \&\& \operatorname{NeQ}[b \cdot c - a \cdot d, 0] \&\& !\operatorname{IntegerQ}[m] \&\& !\operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[b/(b \cdot c - a \cdot d), 0] \&\& (\operatorname{RationalQ}[m] \mid \mid !(\operatorname{RationalQ}[n] \&\& \operatorname{GtQ}[-d/(b \cdot c - a \cdot d), 0]))$

Rule 5181

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a \cdot x]) \cdot (n \cdot x)} \cdot ((c_ + (d_ \cdot (x_))^2)^{p_}), x_Symbol] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[(1 - I \cdot a \cdot x)^{(p + I \cdot (n/2))} \cdot (1 + I \cdot a \cdot x)^{(p - I \cdot (n/2))}, x], x] / ; \operatorname{FreeQ}\{a, c, d, n, p\}, x] \&\& \operatorname{EqQ}[d, a^2 \cdot c] \&\& (\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[c, 0])$

Rule 5184

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a \cdot x]) \cdot (n \cdot x)} \cdot ((c_ + (d_ \cdot (x_))^2)^{p_}), x_Symbol] \rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]} \cdot (c + d \cdot x^2)^{\operatorname{FracPart}[p]} / (1 + a^2 \cdot x^2)^{\operatorname{FracPart}[p]}, \operatorname{Int}[(1 + a^2 \cdot x^2)^p \cdot E^{(n \cdot \operatorname{ArcTan}[a \cdot x])}, x], x] / ; \operatorname{FreeQ}\{a, c, d, n, p\}, x] \&\& \operatorname{Eq}$

$Q[d, a^2*c] \&\& !(IntegerQ[p] || GtQ[c, 0])$

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)}}{(c + a^2 cx^2)^{4/3}} dx &= \frac{\sqrt[3]{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)}}{(1 + a^2 x^2)^{4/3}} dx}{c \sqrt[3]{c + a^2 cx^2}} \\ &= \frac{\sqrt[3]{1 + a^2 x^2} \int (1 - iax)^{-\frac{4}{3} + \frac{in}{2}} (1 + iax)^{-\frac{4}{3} - \frac{in}{2}} dx}{c \sqrt[3]{c + a^2 cx^2}} \\ &= \frac{3 \cdot 2^{-\frac{1}{3} - \frac{in}{2}} (1 - iax)^{\frac{1}{6}(-2+3in)} \sqrt[3]{1 + a^2 x^2} {}_2F_1\left(\frac{1}{6}(-2 + 3in), \frac{1}{6}(8 + 3in); \frac{1}{6}(4 + 3in); \frac{1}{2}(1 - iax)\right)}{ac(2i + 3n) \sqrt[3]{c + a^2 cx^2}} \end{aligned}$$

Mathematica [A]

time = 0.04, size = 123, normalized size = 1.00

$$\frac{3 \cdot 2^{-\frac{1}{3} - \frac{in}{2}} (1 - iax)^{-\frac{1}{3} + \frac{in}{2}} \sqrt[3]{1 + a^2 x^2} {}_2F_1\left(-\frac{1}{3} + \frac{in}{2}, \frac{4}{3} + \frac{in}{2}; \frac{2}{3} + \frac{in}{2}; \frac{1}{2} - \frac{iax}{2}\right)}{ac(2i + 3n) \sqrt[3]{c + a^2 cx^2}}$$

Antiderivative was successfully verified.

[In] Integrate[E^(n*ArcTan[a*x])/(c + a^2*c*x^2)^(4/3), x]

[Out] (3*2^(-1/3 - (I/2)*n)*(1 - I*a*x)^(-1/3 + (I/2)*n)*(1 + a^2*x^2)^(1/3)*Hypergeometric2F1[-1/3 + (I/2)*n, 4/3 + (I/2)*n, 2/3 + (I/2)*n, 1/2 - (I/2)*a*x])/(a*c*(2*I + 3*n)*(c + a^2*c*x^2)^(1/3))

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)}}{(a^2 c x^2 + c)^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(4/3), x)

[Out] int(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(4/3), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(4/3),x, algorithm="maxima")

[Out] integrate(e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(4/3), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(4/3),x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^(2/3)*e^(n*arctan(a*x))/(a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{atan}(ax)}}{(c(a^2x^2 + 1))^{\frac{4}{3}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))/(a**2*c*x**2+c)**(4/3),x)

[Out] Integral(exp(n*atan(a*x))/(c*(a**2*x**2 + 1))**(4/3), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))/(a^2*c*x^2+c)^(4/3),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{e^{n \operatorname{atan}(ax)}}{(ca^2x^2 + c)^{4/3}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(4/3),x)

[Out] int(exp(n*atan(a*x))/(c + a^2*c*x^2)^(4/3), x)

3.364 $\int e^{n \operatorname{ArcTan}(ax)} x^m (c + a^2 cx^2) dx$

Optimal. Leaf size=49

$$\frac{cx^{1+m} F_1\left(1+m; -1 - \frac{in}{2}, -1 + \frac{in}{2}; 2+m; iax, -iax\right)}{1+m}$$

[Out] $c*x^{(1+m)}*AppellF1(1+m, -1+1/2*I*n, -1-1/2*I*n, 2+m, -I*a*x, I*a*x)/(1+m)$

Rubi [A]

time = 0.05, antiderivative size = 49, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 22, $\frac{\text{number of rules}}{\text{integrand size}} = 0.091$, Rules used = {5190, 138}

$$\frac{cx^{m+1} F_1\left(m+1; -\frac{in}{2} - 1, \frac{in}{2} - 1; m+2; iax, -iax\right)}{m+1}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{(n*\text{ArcTan}[a*x])}*x^m*(c + a^2*c*x^2), x]$

[Out] $(c*x^{(1+m)}*AppellF1[1+m, -1 - (I/2)*n, -1 + (I/2)*n, 2+m, I*a*x, (-I)*a*x])/(1+m)$

Rule 138

$\text{Int}[(b_*)*(x_)^{(m_*)}*((c_*) + (d_*)*(x_)^{(n_*)}*((e_*) + (f_*)*(x_)^{(p_*)}), x_*$
 $\text{Symbol}] \rightarrow \text{Simp}[c^n * e^p * ((b*x)^{(m+1})/(b*(m+1)))*AppellF1[m+1, -n, -p,$
 $m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] &&
 & !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5190

$\text{Int}[E^{(\text{ArcTan}[(a_*)*(x_)]*(n_*))}*(x_)^{(m_*)}*((c_*) + (d_*)*(x_)^2)^{(p_*)}, x_*$
 $\text{Symbol}] \rightarrow \text{Dist}[c^p, \text{Int}[x^m*(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int e^{n \tan^{-1}(ax)} x^m (c + a^2 cx^2) dx &= c \int x^m (1 - iax)^{1 + \frac{in}{2}} (1 + iax)^{1 - \frac{in}{2}} dx \\ &= \frac{cx^{1+m} F_1\left(1+m; -1 - \frac{in}{2}, -1 + \frac{in}{2}; 2+m; iax, -iax\right)}{1+m} \end{aligned}$$

Mathematica [F]

time = 0.35, size = 0, normalized size = 0.00

$$\int e^{n \operatorname{ArcTan}(ax)} x^m (c + a^2 cx^2) dx$$

Verification is not applicable to the result.

[In] Integrate[E^(n*ArcTan[a*x])*x^m*(c + a^2*c*x^2), x]

[Out] Integrate[E^(n*ArcTan[a*x])*x^m*(c + a^2*c*x^2), x]

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} x^m (a^2 cx^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^m*(a^2*c*x^2+c), x)

[Out] int(exp(n*arctan(a*x))*x^m*(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m*(a^2*c*x^2+c), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)*x^m*e^(n*arctan(a*x)), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m*(a^2*c*x^2+c), x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)*x^m*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$c \left(\int x^m e^{n \operatorname{atan}(ax)} dx + \int a^2 x^2 x^m e^{n \operatorname{atan}(ax)} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*atan(a*x))*x**m*(a**2*c*x**2+c),x)
```

```
[Out] c*(Integral(x**m*exp(n*atan(a*x)), x) + Integral(a**2*x**2*x**m*exp(n*atan(a*x)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*arctan(a*x))*x^m*(a^2*c*x^2+c),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int x^m e^{n \operatorname{atan}(ax)} (ca^2 x^2 + c) dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(x^m*exp(n*atan(a*x))*(c + a^2*c*x^2),x)
```

```
[Out] int(x^m*exp(n*atan(a*x))*(c + a^2*c*x^2), x)
```


$$3.365 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{c + a^2 c x^2} dx$$

Optimal. Leaf size=51

$$\frac{x^{1+m} F_1\left(1+m; 1 - \frac{in}{2}, 1 + \frac{in}{2}; 2+m; iax, -iax\right)}{c(1+m)}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 1+1/2*I*n, 1-1/2*I*n, 2+m, -I*a*x, I*a*x)/c/(1+m)$

Rubi [A]

time = 0.07, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5190, 138}

$$\frac{x^{m+1} F_1\left(m+1; 1 - \frac{in}{2}, \frac{in}{2} + 1; m+2; iax, -iax\right)}{c(m+1)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^m)/(c + a^2 * c * x^2), x]$

[Out] $(x^{(1+m)} \operatorname{AppellF1}[1+m, 1 - (I/2)*n, 1 + (I/2)*n, 2+m, I*a*x, (-I)*a*x])/c*(1+m)$

Rule 138

$\operatorname{Int}[(b_*)^m * ((c_*) + (d_*) * (x_*))^{(n_*)} * ((e_*) + (f_*) * (x_*))^{(p_*)}, x_*$
 Symbol] :> $\operatorname{Simp}[c^n * e^p * ((b*x)^{(m+1})/(b*(m+1))) * \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] & & !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a_*] * (x_*))} * (n_*) * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^2)^{(p_*)}, x_*$
 Symbol] :> $\operatorname{Dist}[c^p, \operatorname{Int}[x^m * (1 - I*a*x)^{(p+I*(n/2))} * (1 + I*a*x)^{(p-I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x^m}{c + a^2 c x^2} dx &= \frac{\int x^m (1 - iax)^{-1 + \frac{in}{2}} (1 + iax)^{-1 - \frac{in}{2}} dx}{c} \\ &= \frac{x^{1+m} F_1\left(1+m; 1 - \frac{in}{2}, 1 + \frac{in}{2}; 2+m; iax, -iax\right)}{c(1+m)} \end{aligned}$$

Mathematica [A]

time = 0.12, size = 96, normalized size = 1.88

$$\frac{e^{n \operatorname{ArcTan}(ax)} (1 - e^{2i \operatorname{ArcTan}(ax)})^{-m} (1 + e^{2i \operatorname{ArcTan}(ax)})^m x^m F_1\left(-\frac{in}{2}; m, -m; 1 - \frac{in}{2}; -e^{2i \operatorname{ArcTan}(ax)}, e^{2i \operatorname{ArcTan}(ax)}\right)}{acn}$$

Warning: Unable to verify antiderivative.

[In] Integrate[(E^(n*ArcTan[a*x])*x^m)/(c + a^2*c*x^2), x]

[Out] (E^(n*ArcTan[a*x])*(1 + E^((2*I)*ArcTan[a*x]))^m*x^m*AppellF1[(-1/2*I)*n, m, -m, 1 - (I/2)*n, -E^((2*I)*ArcTan[a*x]), E^((2*I)*ArcTan[a*x])])/(a*c*(1 - E^((2*I)*ArcTan[a*x]))^m*n)

Maple [F]

time = 0.01, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^m}{a^2 c x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c), x)

[Out] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c), x, algorithm="maxima")

[Out] integrate(x^m*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c), x, algorithm="fricas")

[Out] integral(x^m*e^(n*arctan(a*x))/(a^2*c*x^2 + c), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\frac{\int \frac{x^m e^{n \operatorname{atan}(ax)}}{a^2 x^2 + 1} dx}{c}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))*x**m/(a**2*c*x**2+c), x)`

[Out] `Integral(x**m*exp(n*atan(a*x))/(a**2*x**2 + 1), x)/c`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c), x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{c a^2 x^2 + c} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)`

[Out] `int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2), x)`

$$3.366 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{(c+a^2cx^2)^2} dx$$

Optimal. Leaf size=51

$$\frac{x^{1+m} F_1\left(1+m; 2-\frac{in}{2}, 2+\frac{in}{2}; 2+m; iax, -iax\right)}{c^2(1+m)}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 2+1/2*I*n, 2-1/2*I*n, 2+m, -I*a*x, I*a*x)/c^2/(1+m)$

Rubi [A]

time = 0.06, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5190, 138}

$$\frac{x^{m+1} F_1\left(m+1; 2-\frac{in}{2}, \frac{in}{2}+2; m+2; iax, -iax\right)}{c^2(m+1)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^m) / (c + a^2 * c * x^2)^2, x]$

[Out] $(x^{(1+m)} \operatorname{AppellF1}[1+m, 2-(I/2)*n, 2+(I/2)*n, 2+m, I*a*x, (-I)*a*x]) / (c^2 * (1+m))$

Rule 138

$\operatorname{Int}[(b_*) * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^{(n_*)} * ((e_*) + (f_*) * (x_*)^{(p_*)}), x_*$
 Symbol] $\rightarrow \operatorname{Simp}[c^n * e^p * (b*x)^{(m+1)} / (b*(m+1))] * \operatorname{AppellF1}[m+1, -n, -p,$
 $m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] &
 & !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*) * (x_*)] * (n_*))} * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^2)^{(p_*)}, x_*$
 Symbol] $\rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[x^m * (1 - I*a*x)^{(p+I*(n/2))} * (1 + I*a*x)^{(p-I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x^m}{(c+a^2cx^2)^2} dx &= \frac{\int x^m (1-iax)^{-2+\frac{in}{2}} (1+iax)^{-2-\frac{in}{2}} dx}{c^2} \\ &= \frac{x^{1+m} F_1\left(1+m; 2-\frac{in}{2}, 2+\frac{in}{2}; 2+m; iax, -iax\right)}{c^2(1+m)} \end{aligned}$$

Mathematica [F]

time = 0.35, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^2} dx$$

Verification is not applicable to the result.

[In] Integrate[(E^(n*ArcTan[a*x]))*x^m]/(c + a^2*c*x^2)^2,x]

[Out] Integrate[(E^(n*ArcTan[a*x]))*x^m]/(c + a^2*c*x^2)^2, x]

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^m}{(a^2 cx^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^2,x)

[Out] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^2,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^2,x, algorithm="maxima")

[Out] integrate(x^m*e^(n*arctan(a*x))/(a^2*c*x^2 + c)^2, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^2,x, algorithm="fricas")

[Out] integral(x^m*e^(n*arctan(a*x))/(a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{a^4 x^4 + 2a^2 x^2 + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*x**m/(a**2*c*x**2+c)**2,x)

[Out] Integral(x**m*exp(n*atan(a*x))/(a**4*x**4 + 2*a**2*x**2 + 1), x)/c**2

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^2,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{(ca^2x^2 + c)^2} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^2,x)

[Out] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^2, x)

$$3.367 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{(c+a^2cx^2)^3} dx$$

Optimal. Leaf size=51

$$\frac{x^{1+m} F_1\left(1+m; 3-\frac{in}{2}, 3+\frac{in}{2}; 2+m; iax, -iax\right)}{c^3(1+m)}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 3+1/2*I*n, 3-1/2*I*n, 2+m, -I*a*x, I*a*x)/c^3/(1+m)$

Rubi [A]

time = 0.07, antiderivative size = 51, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.083$, Rules used = {5190, 138}

$$\frac{x^{m+1} F_1\left(m+1; 3-\frac{in}{2}, \frac{in}{2}+3; m+2; iax, -iax\right)}{c^3(m+1)}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^m) / (c + a^2 * c * x^2)^3, x]$

[Out] $(x^{(1+m)} \operatorname{AppellF1}[1+m, 3-(I/2)*n, 3+(I/2)*n, 2+m, I*a*x, (-I)*a*x]) / (c^3 * (1+m))$

Rule 138

$\operatorname{Int}[(b_*)^m * ((c_*) + (d_*) * (x_*))^{(n_*)} * ((e_*) + (f_*) * (x_*))^{(p_*)}, x_*$
 Symbol] $\rightarrow \operatorname{Simp}[c^n * e^p * ((b*x)^{(m+1}) / (b*(m+1))) * \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[a_*] * (x_*))} * (n_*) * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^2)^{(p_*)}, x_*$
 Symbol] $\rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[x^m * (1 - I*a*x)^{(p+I*(n/2))} * (1 + I*a*x)^{(p-I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{n \tan^{-1}(ax)} x^m}{(c+a^2cx^2)^3} dx &= \frac{\int x^m (1-iax)^{-3+\frac{in}{2}} (1+iax)^{-3-\frac{in}{2}} dx}{c^3} \\ &= \frac{x^{1+m} F_1\left(1+m; 3-\frac{in}{2}, 3+\frac{in}{2}; 2+m; iax, -iax\right)}{c^3(1+m)} \end{aligned}$$

Mathematica [F]

time = 0.55, size = 0, normalized size = 0.00

$$\int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^3} dx$$

Verification is not applicable to the result.

[In] Integrate[(E^(n*ArcTan[a*x])*x^m)/(c + a^2*c*x^2)^3,x]

[Out] Integrate[(E^(n*ArcTan[a*x])*x^m)/(c + a^2*c*x^2)^3, x]

Maple [F]

time = 0.04, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^m}{(a^2 cx^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^3,x)

[Out] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^3,x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] integrate(x^m*e^(n*arctan(a*x))/(a^2*c*x^2 + c)^3, x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^3,x, algorithm="fricas")

[Out] integral(x^m*e^(n*arctan(a*x))/(a^6*c^3*x^6 + 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 + c^3), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{a^6 x^6 + 3a^4 x^4 + 3a^2 x^2 + 1} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*atan(a*x))*x**m/(a**2*c*x**2+c)**3,x)`

[Out] `Integral(x**m*exp(n*atan(a*x))/(a**6*x**6 + 3*a**4*x**4 + 3*a**2*x**2 + 1), x)/c**3`

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^3,x, algorithm="giac")`

[Out] `sage0*x`

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{(ca^2 x^2 + c)^3} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^3,x)`

[Out] `int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^3, x)`

$$3.368 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{\sqrt{c + a^2 cx^2}} dx$$

Optimal. Leaf size=79

$$\frac{x^{1+m} \sqrt{1 + a^2 x^2} F_1\left(1 + m; \frac{1}{2}(1 - in), \frac{1}{2}(1 + in); 2 + m; iax, -iax\right)}{(1 + m) \sqrt{c + a^2 cx^2}}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 1/2+1/2*I*n, 1/2-1/2*I*n, 2+m, -I*a*x, I*a*x) * (a^2*x^2+1)^{(1/2)/(1+m)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.14, antiderivative size = 79, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {5193, 5190, 138}

$$\frac{\sqrt{a^2 x^2 + 1} x^{m+1} F_1\left(m + 1; \frac{1}{2}(1 - in), \frac{1}{2}(in + 1); m + 2; iax, -iax\right)}{(m + 1) \sqrt{a^2 cx^2 + c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^m) / \operatorname{Sqrt}[c + a^2*c*x^2], x]$

[Out] $(x^{(1 + m)} * \operatorname{Sqrt}[1 + a^2*x^2] * \operatorname{AppellF1}[1 + m, (1 - I*n)/2, (1 + I*n)/2, 2 + m, I*a*x, (-I)*a*x]) / ((1 + m) * \operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 138

$\operatorname{Int}[(b_*) * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^{(n_*)} * ((e_*) + (f_*) * (x_*)^{(p_*)}), x_*$
 $\operatorname{Symbol}] \rightarrow \operatorname{Simp}[c^n * e^p * (b*x)^{(m+1)} / (b*(m+1))] * \operatorname{AppellF1}[m+1, -n, -p,$
 $m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ $\operatorname{FreeQ}\{b, c, d, e, f, m, n, p\}, x] \&\& \operatorname{IntegerQ}[m] \&\& \operatorname{IntegerQ}[n] \&\& \operatorname{GtQ}[c, 0] \&\& (\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[e, 0])$

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*) * (x_*)] * (n_*))} * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^2)^{(p_*)}, x_*$
 $\operatorname{Symbol}] \rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[x^m * (1 - I*a*x)^{(p + I*(n/2))} * (1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, m, n, p\}, x] \&\& \operatorname{EqQ}[d, a^2*c] \&\& (\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[c, 0])$

Rule 5193

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*) * (x_*)] * (n_*))} * (x_*)^{(m_*)} * ((c_*) + (d_*) * (x_*)^2)^{(p_*)}, x_*$
 $\operatorname{symbol}] \rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]} * ((c + d*x^2)^{\operatorname{FracPart}[p]} / (1 + a^2*x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[x^m * (1 + a^2*x^2)^p * E^{(n \operatorname{ArcTan}[a*x])}, x], x] /;$ $\operatorname{FreeQ}\{a, c, d, m, n, p\}, x] \&\& \operatorname{EqQ}[d, a^2*c] \&\& !(\operatorname{IntegerQ}[p] \mid \mid \operatorname{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)} x^m}{\sqrt{c + a^2 cx^2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)} x^m}{\sqrt{1 + a^2 x^2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int x^m (1 - iax)^{-\frac{1}{2} + \frac{in}{2}} (1 + iax)^{-\frac{1}{2} - \frac{in}{2}} dx}{\sqrt{c + a^2 cx^2}} \\
&= \frac{x^{1+m} \sqrt{1 + a^2 x^2} F_1\left(1 + m; \frac{1}{2}(1 - in), \frac{1}{2}(1 + in); 2 + m; iax, -iax\right)}{(1 + m) \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [F]

time = 0.25, size = 0, normalized size = 0.00

$$\int \frac{e^{n \text{ArcTan}(ax)} x^m}{\sqrt{c + a^2 cx^2}} dx$$

Verification is not applicable to the result.

[In] Integrate[(E^(n*ArcTan[a*x]))*x^m/Sqrt[c + a^2*c*x^2], x]

[Out] Integrate[(E^(n*ArcTan[a*x]))*x^m/Sqrt[c + a^2*c*x^2], x]

Maple [F]

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^m}{\sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(1/2), x)

[Out] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(1/2), x)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(1/2), x, algorithm="maxima")

[Out] integrate(x^m*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(1/2),x, algorithm="fricas")``[Out] integral(x^m*e^(n*arctan(a*x))/sqrt(a^2*c*x^2 + c), x)`**Sympy [F]**

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{\sqrt{c(a^2 x^2 + 1)}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*atan(a*x))*x**m/(a**2*c*x**2+c)**(1/2),x)``[Out] Integral(x**m*exp(n*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x)`**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(1/2),x, algorithm="giac")``[Out] sage0*x`**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{\sqrt{c a^2 x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2),x)``[Out] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(1/2), x)`

$$3.369 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=82

$$\frac{x^{1+m} \sqrt{1+a^2x^2} F_1\left(1+m; \frac{1}{2}(3-in), \frac{1}{2}(3+in); 2+m; iax, -iax\right)}{c(1+m)\sqrt{c+a^2cx^2}}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 3/2+1/2*I*n, 3/2-1/2*I*n, 2+m, -I*a*x, I*a*x) * (a^2*x^2+1)^{(1/2)}/c/(1+m)/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {5193, 5190, 138}

$$\frac{\sqrt{a^2x^2+1} x^{m+1} F_1\left(m+1; \frac{1}{2}(3-in), \frac{1}{2}(in+3); m+2; iax, -iax\right)}{c(m+1)\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a*x])} * x^m) / (c + a^2*c*x^2)^{(3/2)}, x]$

[Out] $(x^{(1+m)} * \operatorname{Sqrt}[1 + a^2*x^2] * \operatorname{AppellF1}[1+m, (3-I*n)/2, (3+I*n)/2, 2+m, I*a*x, (-I)*a*x]) / (c*(1+m)*\operatorname{Sqrt}[c + a^2*c*x^2])$

Rule 138

$\operatorname{Int}[(b_*)*(x_*)^{(m_*)}*((c_*) + (d_*)*(x_*)^{(n_*)}*((e_*) + (f_*)*(x_*)^{(p_*)}), x_Symbol] :> \operatorname{Simp}[c^n * e^p * ((b*x)^{(m+1}) / (b*(m+1))) * \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d)*(x/c), (-f)*(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*)*(x_*)])} * (n_*) * (x_*)^{(m_*)} * ((c_*) + (d_*)*(x_*)^2)^{(p_*)}, x_Symbol] :> \operatorname{Dist}[c^p, \operatorname{Int}[x^m * (1 - I*a*x)^{(p+I*(n/2))} * (1 + I*a*x)^{(p-I*(n/2))}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_*)*(x_*)])} * (n_*) * (x_*)^{(m_*)} * ((c_*) + (d_*)*(x_*)^2)^{(p_*)}, x_Symbol] :> \operatorname{Dist}[c^{\operatorname{IntPart}[p]} * ((c + d*x^2)^{\operatorname{FracPart}[p]} / (1 + a^2*x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[x^m * (1 + a^2*x^2)^p * E^{(n \operatorname{ArcTan}[a*x])}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)} x^m}{(c + a^2 cx^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)} x^m}{(1 + a^2 x^2)^{3/2}} dx}{c \sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int x^m (1 - iax)^{-\frac{3}{2} + \frac{in}{2}} (1 + iax)^{-\frac{3}{2} - \frac{in}{2}} dx}{c \sqrt{c + a^2 cx^2}} \\
&= \frac{x^{1+m} \sqrt{1 + a^2 x^2} F_1\left(1 + m; \frac{1}{2}(3 - in), \frac{1}{2}(3 + in); 2 + m; iax, -iax\right)}{c(1 + m) \sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [F]

time = 0.34, size = 0, normalized size = 0.00

$$\int \frac{e^{n \text{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^{3/2}} dx$$

Verification is not applicable to the result.

`[In] Integrate[(E^(n*ArcTan[a*x])*x^m)/(c + a^2*c*x^2)^(3/2), x]``[Out] Integrate[(E^(n*ArcTan[a*x])*x^m)/(c + a^2*c*x^2)^(3/2), x]`**Maple [F]**

time = 0.02, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^m}{(a^2 c x^2 + c)^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(3/2), x)``[Out] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(3/2), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(3/2), x, algorithm="maxima")``[Out] integrate(x^m*e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(3/2), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")

[Out] integral(sqrt(a^2*c*x^2 + c)*x^m*e^(n*arctan(a*x))/(a^4*c^2*x^4 + 2*a^2*c^2*x^2 + c^2), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{(c(a^2 x^2 + 1))^{\frac{3}{2}}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*x**m/(a**2*c*x**2+c)**(3/2),x)

[Out] Integral(x**m*exp(n*atan(a*x))/(c*(a**2*x**2 + 1))**(3/2), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{(c a^2 x^2 + c)^{3/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(3/2),x)

[Out] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(3/2), x)

$$3.370 \quad \int \frac{e^{n \operatorname{ArcTan}(ax)} x^m}{(c+a^2cx^2)^{5/2}} dx$$

Optimal. Leaf size=82

$$\frac{x^{1+m} \sqrt{1+a^2x^2} F_1\left(1+m; \frac{1}{2}(5-in), \frac{1}{2}(5+in); 2+m; iax, -iax\right)}{c^2(1+m) \sqrt{c+a^2cx^2}}$$

[Out] $x^{(1+m)} \operatorname{AppellF1}(1+m, 5/2+1/2 I n, 5/2-1/2 I n, 2+m, -I a x, I a x) (a^2 x^2+1)^{(1/2)} / c^2 / (1+m) / (a^2 c x^2+c)^{(1/2)}$

Rubi [A]

time = 0.15, antiderivative size = 82, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.115$, Rules used = {5193, 5190, 138}

$$\frac{\sqrt{a^2x^2+1} x^{m+1} F_1\left(m+1; \frac{1}{2}(5-in), \frac{1}{2}(in+5); m+2; iax, -iax\right)}{c^2(m+1) \sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\operatorname{Int}[(E^{(n \operatorname{ArcTan}[a x])} x^m) / (c + a^2 c x^2)^{(5/2)}, x]$

[Out] $(x^{(1+m)} \operatorname{Sqrt}[1+a^2 x^2] \operatorname{AppellF1}[1+m, (5-I n)/2, (5+I n)/2, 2+m, I a x, (-I) a x]) / (c^2 (1+m) \operatorname{Sqrt}[c+a^2 c x^2])$

Rule 138

$\operatorname{Int}[(b_.) (x_.)^{(m_.)} ((c_.) + (d_.) (x_.)^{(n_.)}) ((e_.) + (f_.) (x_.)^{(p_.)}), x_]$
 Symbol] $\rightarrow \operatorname{Simp}[c^n e^p ((b x)^{(m+1}) / (b (m+1))) \operatorname{AppellF1}[m+1, -n, -p, m+2, (-d)(x/c), (-f)(x/e)], x] /;$ FreeQ[{b, c, d, e, f, m, n, p}, x] && !IntegerQ[m] && !IntegerQ[n] && GtQ[c, 0] && (IntegerQ[p] || GtQ[e, 0])

Rule 5190

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_.) (x_.)] (n_.)}) (x_.)^{(m_.)} ((c_.) + (d_.) (x_.)^2)^{(p_.)}), x_]$
 Symbol] $\rightarrow \operatorname{Dist}[c^p, \operatorname{Int}[x^m (1-I a x)^{(p+I(n/2))} (1+I a x)^{(p-I(n/2))}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2 c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

$\operatorname{Int}[E^{(\operatorname{ArcTan}[(a_.) (x_.)] (n_.)}) (x_.)^{(m_.)} ((c_.) + (d_.) (x_.)^2)^{(p_.)}), x_]$
 Symbol] $\rightarrow \operatorname{Dist}[c^{\operatorname{IntPart}[p]} ((c+d x^2)^{\operatorname{FracPart}[p]} / (1+a^2 x^2)^{\operatorname{FracPart}[p]}), \operatorname{Int}[x^m (1+a^2 x^2)^p E^{(n \operatorname{ArcTan}[a x])}, x], x] /;$ FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2 c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int \frac{e^{n \tan^{-1}(ax)} x^m}{(c + a^2 cx^2)^{5/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{n \tan^{-1}(ax)} x^m}{(1 + a^2 x^2)^{5/2}} dx}{c^2 \sqrt{c + a^2 cx^2}} \\
&= \frac{\sqrt{1 + a^2 x^2} \int x^m (1 - iax)^{-\frac{5}{2} + \frac{in}{2}} (1 + iax)^{-\frac{5}{2} - \frac{in}{2}} dx}{c^2 \sqrt{c + a^2 cx^2}} \\
&= \frac{x^{1+m} \sqrt{1 + a^2 x^2} F_1\left(1 + m; \frac{1}{2}(5 - in), \frac{1}{2}(5 + in); 2 + m; iax, -iax\right)}{c^2(1 + m)\sqrt{c + a^2 cx^2}}
\end{aligned}$$

Mathematica [F]

time = 0.44, size = 0, normalized size = 0.00

$$\int \frac{e^{n \text{ArcTan}(ax)} x^m}{(c + a^2 cx^2)^{5/2}} dx$$

Verification is not applicable to the result.

`[In] Integrate[(E^(n*ArcTan[a*x]))*x^m/(c + a^2*c*x^2)^(5/2), x]``[Out] Integrate[(E^(n*ArcTan[a*x]))*x^m/(c + a^2*c*x^2)^(5/2), x]`**Maple [F]**

time = 0.03, size = 0, normalized size = 0.00

$$\int \frac{e^{n \arctan(ax)} x^m}{(a^2 c x^2 + c)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(5/2), x)``[Out] int(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(5/2), x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(5/2), x, algorithm="maxima")``[Out] integrate(x^m*e^(n*arctan(a*x))/(a^2*c*x^2 + c)^(5/2), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(5/2),x, algorithm="fricas")
```

```
[Out] integral(sqrt(a^2*c*x^2 + c)*x^m*e^(n*arctan(a*x))/(a^6*c^3*x^6 + 3*a^4*c^3*x^4 + 3*a^2*c^3*x^2 + c^3), x)
```

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*atan(a*x))*x**m/(a**2*c*x**2+c)**(5/2),x)
```

```
[Out] Exception raised: SystemError >> excessive stack use: stack is 3006 deep
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(n*arctan(a*x))*x^m/(a^2*c*x^2+c)^(5/2),x, algorithm="giac")
```

```
[Out] sage0*x
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^m e^{n \operatorname{atan}(ax)}}{(ca^2 x^2 + c)^{5/2}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(5/2),x)
```

```
[Out] int((x^m*exp(n*atan(a*x)))/(c + a^2*c*x^2)^(5/2), x)
```

3.371 $\int e^{n \operatorname{ArcTan}(ax)} (c + a^2 cx^2)^p dx$

Optimal. Leaf size=115

$$\frac{2^{1-\frac{in}{2}+p}(1-iax)^{1+\frac{in}{2}+p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p {}_2F_1\left(\frac{in}{2}-p, 1+\frac{in}{2}+p; 2+\frac{in}{2}+p; \frac{1}{2}(1-iax)\right)}{a(n-2i(1+p))}$$

[Out] $2^{(1-1/2*I*n+p)*(1-I*a*x)^{(1+1/2*I*n+p)*(a^2*c*x^2+c)^p}$ hypergeom([1/2*I*n-p, 1+1/2*I*n+p], [2+1/2*I*n+p], 1/2-1/2*I*a*x)/a/(n-2*I*(1+p))/((a^2*x^2+1)^p)

Rubi [A]

time = 0.06, antiderivative size = 115, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 21, $\frac{\text{number of rules}}{\text{integrand size}} = 0.143$, Rules used = {5184, 5181, 71}

$$\frac{2^{-\frac{in}{2}+p+1}(a^2x^2+1)^{-p}(a^2cx^2+c)^p(1-iax)^{\frac{in}{2}+p+1} {}_2F_1\left(\frac{in}{2}-p, \frac{in}{2}+p+1; \frac{in}{2}+p+2; \frac{1}{2}(1-iax)\right)}{a(n-2i(p+1))}$$

Antiderivative was successfully verified.

[In] Int[E^(n*ArcTan[a*x])*(c + a^2*c*x^2)^p,x]

[Out] $(2^{(1-(I/2)*n+p)*(1-I*a*x)^{(1+(I/2)*n+p)*(c+a^2*c*x^2)^p}$ Hypergeometric2F1[(I/2)*n-p, 1+(I/2)*n+p, 2+(I/2)*n+p, (1-I*a*x)/2])/ (a*(n-(2*I)*(1+p))*(1+a^2*x^2)^p)

Rule 71

Int[((a_) + (b_.)*(x_))^(m_)*((c_) + (d_.)*(x_))^(n_), x_Symbol] := Simp[((a + b*x)^(m + 1)/(b*(m + 1)*(b/(b*c - a*d))^n)*Hypergeometric2F1[-n, m + 1, m + 2, (-d)*((a + b*x)/(b*c - a*d))], x] /; FreeQ[{a, b, c, d, m, n}, x] && NeQ[b*c - a*d, 0] && !IntegerQ[m] && !IntegerQ[n] && GtQ[b/(b*c - a*d), 0] && (RationalQ[m] || !(RationalQ[n] && GtQ[-d/(b*c - a*d), 0]))

Rule 5181

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] := Dist[c^p, Int[(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5184

Int[E^(ArcTan[(a_.)*(x_)])*(n_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] := Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p]/(1 + a^2*x^2)^FracPart[p]), Int[(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
\int e^{n \tan^{-1}(ax)} (c + a^2 cx^2)^p dx &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int e^{n \tan^{-1}(ax)} (1 + a^2 x^2)^p dx \\
&= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int (1 - iax)^{\frac{in}{2}+p} (1 + iax)^{-\frac{in}{2}+p} dx \\
&= \frac{2^{1-\frac{in}{2}+p} (1 - iax)^{1+\frac{in}{2}+p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p {}_2F_1\left(\frac{in}{2} - p, 1 + \frac{in}{2} + p; 2 + \frac{in}{2} + p; \frac{1}{2}(1 - iax)\right)}{a(n - 2i(1 + p))}
\end{aligned}$$

Mathematica [A]

time = 0.03, size = 115, normalized size = 1.00

$$\frac{2^{1-\frac{in}{2}+p} (1 - iax)^{1+\frac{in}{2}+p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p {}_2F_1\left(\frac{in}{2} - p, 1 + \frac{in}{2} + p; 2 + \frac{in}{2} + p; \frac{1}{2}(1 - iax)\right)}{a(n - 2i(1 + p))}$$

Antiderivative was successfully verified.

`[In] Integrate[E^(n*ArcTan[a*x])*(c + a^2*c*x^2)^p,x]`

```
[Out] (2^(1 - (I/2)*n + p)*(1 - I*a*x)^(1 + (I/2)*n + p)*(c + a^2*c*x^2)^p*Hypergeometric2F1[(I/2)*n - p, 1 + (I/2)*n + p, 2 + (I/2)*n + p, (1 - I*a*x)/2])/
(a*(n - (2*I)*(1 + p))*(1 + a^2*x^2)^p)
```

Maple [F]

time = 0.02, size = 0, normalized size = 0.00

$$\int e^{n \arctan(ax)} (a^2 cx^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

`[In] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^p,x)``[Out] int(exp(n*arctan(a*x))*(a^2*c*x^2+c)^p,x)`**Maxima [F]**

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

`[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="maxima")``[Out] integrate((a^2*c*x^2 + c)^p*e^(n*arctan(a*x)), x)`

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="fricas")

[Out] integral((a^2*c*x^2 + c)^p*e^(n*arctan(a*x)), x)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\int (c(a^2x^2 + 1))^p e^{n \operatorname{atan}(ax)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*atan(a*x))*(a**2*c*x**2+c)**p,x)

[Out] Integral((c*(a**2*x**2 + 1))**p*exp(n*atan(a*x)), x)

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(n*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int e^{n \operatorname{atan}(ax)} (ca^2x^2 + c)^p dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^p,x)

[Out] int(exp(n*atan(a*x))*(c + a^2*c*x^2)^p, x)

3.372 $\int e^{-2ip\text{ArcTan}(ax)}(c + a^2cx^2)^p dx$

Optimal. Leaf size=53

$$\frac{i(1 - iax)^{1+2p} (1 + a^2x^2)^{-p} (c + a^2cx^2)^p}{a(1 + 2p)}$$

[Out] $I*(1-I*a*x)^{(1+2*p)}*(a^2*c*x^2+c)^p/a/(1+2*p)/((a^2*x^2+1)^p)$

Rubi [A]

time = 0.05, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5184, 5181, 32}

$$\frac{i(1 - iax)^{2p+1} (a^2x^2 + 1)^{-p} (a^2cx^2 + c)^p}{a(2p + 1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(c + a^2*c*x^2)^p/E^{((2*I)*p*\text{ArcTan}[a*x])}, x]$

[Out] $(I*(1 - I*a*x)^{(1 + 2*p)}*(c + a^2*c*x^2)^p)/(a*(1 + 2*p)*(1 + a^2*x^2)^p)$

Rule 32

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)}, x_Symbol] := \text{Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] /;$ $\text{FreeQ}\{a, b, m\}, x\} \ \&\& \ \text{NeQ}[m, -1]$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))}*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] := \text{Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /;$ $\text{FreeQ}\{a, c, d, n, p\}, x\} \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))}*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] := \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /;$ $\text{FreeQ}\{a, c, d, n, p\}, x\} \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{-2ip \tan^{-1}(ax)} (c + a^2 cx^2)^p dx &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int e^{-2ip \tan^{-1}(ax)} (1 + a^2 x^2)^p dx \\ &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int (1 - iax)^{2p} dx \\ &= \frac{i(1 - iax)^{1+2p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p}{a(1 + 2p)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 39, normalized size = 0.74

$$\frac{e^{-2ip \operatorname{ArcTan}(ax)} (i + ax) (c + a^2 cx^2)^p}{a + 2ap}$$

Antiderivative was successfully verified.

[In] Integrate[(c + a^2*c*x^2)^p/E^((2*I)*p*ArcTan[a*x]), x]**[Out]** ((I + a*x)*(c + a^2*c*x^2)^p)/(E^((2*I)*p*ArcTan[a*x])*(a + 2*a*p))**Maple [A]**

time = 0.14, size = 41, normalized size = 0.77

method	result
gospers	$\frac{(ax+i)(a^2cx^2+c)^p e^{-2ip \arctan(ax)}}{a(1+2p)}$
risch	$\frac{p \left(i \operatorname{csgn}(i(ax-i)(ax+i)) \operatorname{csgn}(ic(ax+i)(ax-i)^2 \pi + ic \operatorname{sgn}(ax-i)^2 \pi + ic \operatorname{sgn}(ax+i)^2 \pi - ic \operatorname{sgn}(ax-i)^2 \operatorname{csgn}(i(ax-i)) \pi - ic \operatorname{sgn}(ax+i) \operatorname{csgn}(i(ax+i)) \right)}{(ax+i)e}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((a^2*c*x^2+c)^p/exp(2*I*p*arctan(a*x)), x, method=_RETURNVERBOSE)**[Out]** (I+a*x)/a/(1+2*p)*(a^2*c*x^2+c)^p/exp(2*I*p*arctan(a*x))**Maxima [A]**

time = 0.28, size = 76, normalized size = 1.43

$$\frac{(ac^p x + i c^p)(a^2 x^2 + 1)^p \cos(2p \arctan(ax)) - (iac^p x - c^p)(a^2 x^2 + 1)^p \sin(2p \arctan(ax))}{2ap + a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^p/exp(2*I*p*arctan(a*x)), x, algorithm="maxima")**[Out]** ((a*c^p*x + I*c^p)*(a^2*x^2 + 1)^p*cos(2*p*arctan(a*x)) - (I*a*c^p*x - c^p)*(a^2*x^2 + 1)^p*sin(2*p*arctan(a*x)))/(2*a*p + a)

Fricas [A]

time = 3.34, size = 42, normalized size = 0.79

$$\frac{(ax + i)(a^2cx^2 + c)^p \left(-\frac{ax+i}{ax-i}\right)^p}{2ap + a}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^p/exp(2*I*p*arctan(a*x)),x, algorithm="fricas")

[Out] (a*x + I)*(a^2*c*x^2 + c)^p*(-(a*x + I)/(a*x - I))^p/(2*a*p + a)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\begin{cases} \frac{x}{\sqrt{c}} & \text{for } a = 0 \wedge p = -\frac{1}{2} \\ c^p x & \text{for } a = 0 \\ \int \frac{e^{i \operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx & \text{for } p = -\frac{1}{2} \\ \frac{ax(a^2cx^2+c)^p}{2ape^{2ip \operatorname{atan}(ax)} + ae^{2ip \operatorname{atan}(ax)}} + \frac{i(a^2cx^2+c)^p}{2ape^{2ip \operatorname{atan}(ax)} + ae^{2ip \operatorname{atan}(ax)}} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a**2*c*x**2+c)**p/exp(2*I*p*atan(a*x)),x)

[Out] Piecewise((x/sqrt(c), Eq(a, 0) & Eq(p, -1/2)), (c**p*x, Eq(a, 0)), (Integral(exp(I*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x), Eq(p, -1/2)), (a*x*(a**2*c*x**2 + c)**p/(2*a*p*exp(2*I*p*atan(a*x)) + a*exp(2*I*p*atan(a*x))) + I*(a**2*c*x**2 + c)**p/(2*a*p*exp(2*I*p*atan(a*x)) + a*exp(2*I*p*atan(a*x))), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((a^2*c*x^2+c)^p/exp(2*I*p*arctan(a*x)),x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.66, size = 54, normalized size = 1.02

$$\left(\frac{x e^{-p \operatorname{atan}(ax) 2i}}{2p + 1} + \frac{e^{-p \operatorname{atan}(ax) 2i} 1i}{a(2p + 1)} \right) (ca^2x^2 + c)^p$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(-p*atan(a*x)*2i)*(c + a^2*c*x^2)^p,x)
```

```
[Out] ((x*exp(-p*atan(a*x)*2i))/(2*p + 1) + (exp(-p*atan(a*x)*2i)*1i)/(a*(2*p + 1))) * (c + a^2*c*x^2)^p
```

3.373 $\int e^{2ip\text{ArcTan}(ax)}(c + a^2cx^2)^p dx$

Optimal. Leaf size=53

$$\frac{i(1+iax)^{1+2p}(1+a^2x^2)^{-p}(c+a^2cx^2)^p}{a(1+2p)}$$

[Out] $-I*(1+I*a*x)^{(1+2*p)}*(a^2*c*x^2+c)^p/a/(1+2*p)/((a^2*x^2+1)^p)$

Rubi [A]

time = 0.04, antiderivative size = 53, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 24, $\frac{\text{number of rules}}{\text{integrand size}} = 0.125$, Rules used = {5184, 5181, 32}

$$\frac{i(1+iax)^{2p+1}(a^2x^2+1)^{-p}(a^2cx^2+c)^p}{a(2p+1)}$$

Antiderivative was successfully verified.

[In] $\text{Int}[E^{((2*I)*p*\text{ArcTan}[a*x])}*(c + a^2*c*x^2)^p, x]$

[Out] $((-I)*(1 + I*a*x)^{(1 + 2*p)}*(c + a^2*c*x^2)^p)/(a*(1 + 2*p)*(1 + a^2*x^2)^p)$

Rule 32

$\text{Int}[(a_.) + (b_.)*(x_.)^{(m_.)}, x_Symbol] \text{ :> Simp}[(a + b*x)^{(m + 1)}/(b*(m + 1)), x] \text{ /; FreeQ}\{a, b, m\}, x\} \ \&\& \ \text{NeQ}\{m, -1\}$

Rule 5181

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))}*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \text{ :> Dist}[c^p, \text{Int}[(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] \text{ /; FreeQ}\{a, c, d, n, p\}, x\} \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ (\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rule 5184

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))}*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] \text{ :> Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] \text{ /; FreeQ}\{a, c, d, n, p\}, x\} \ \&\& \ \text{EqQ}[d, a^2*c] \ \&\& \ !(\text{IntegerQ}[p] \ || \ \text{GtQ}[c, 0])$

Rubi steps

$$\begin{aligned} \int e^{2ip \tan^{-1}(ax)} (c + a^2 cx^2)^p dx &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int e^{2ip \tan^{-1}(ax)} (1 + a^2 x^2)^p dx \\ &= \left((1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p \right) \int (1 + iax)^{2p} dx \\ &= \frac{i(1 + iax)^{1+2p} (1 + a^2 x^2)^{-p} (c + a^2 cx^2)^p}{a(1 + 2p)} \end{aligned}$$

Mathematica [A]

time = 0.02, size = 39, normalized size = 0.74

$$\frac{e^{2ip \operatorname{ArcTan}(ax)} (-i + ax) (c + a^2 cx^2)^p}{a + 2ap}$$

Antiderivative was successfully verified.

```
[In] Integrate[E^((2*I)*p*ArcTan[a*x])*(c + a^2*c*x^2)^p,x]
```

```
[Out] (E^((2*I)*p*ArcTan[a*x])*(-I + a*x)*(c + a^2*c*x^2)^p)/(a + 2*a*p)
```

Maple [A]

time = 0.10, size = 41, normalized size = 0.77

method	result
gospers	$-\frac{(-ax+i)e^{2ip \arctan(ax)} (a^2 cx^2+c)^p}{a(1+2p)}$
risch	$\frac{(ax-i)e^{p \left(i \operatorname{csgn}(i(ax-i)(ax+i)) \operatorname{csgn}(ic(ax+i)(ax-i))^2 \pi - i \operatorname{csgn}(i(ax-i)(ax+i))^3 \pi - i \operatorname{csgn}(ax+i)^2 \pi - i \operatorname{csgn}(ax+i)^2 \operatorname{csgn}(i(ax+i)) \pi + i \operatorname{csgn}(ic(ax+i)) \right)}}{a(1+2p)}$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int(exp(2*I*p*arctan(a*x))*(a^2*c*x^2+c)^p,x,method=_RETURNVERBOSE)
```

```
[Out] -(I-a*x)/a/(1+2*p)*exp(2*I*p*arctan(a*x))*(a^2*c*x^2+c)^p
```

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(exp(2*I*p*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="maxima")
```

```
[Out] integrate((a^2*c*x^2 + c)^p*e^(2*I*p*arctan(a*x)), x)
```

Fricas [A]

time = 14.58, size = 44, normalized size = 0.83

$$\frac{(ax - i)(a^2cx^2 + c)^p}{(2ap + a) \left(-\frac{ax+i}{ax-i}\right)^p}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*I*p*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="fricas")

[Out] (a*x - I)*(a^2*c*x^2 + c)^p/((2*a*p + a)*(-(a*x + I)/(a*x - I))^p)

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$\begin{cases} \frac{x}{\sqrt{c}} & \text{for } a = 0 \wedge p = -\frac{1}{2} \\ c^p x & \text{for } a = 0 \\ \int \frac{e^{-i \operatorname{atan}(ax)}}{\sqrt{c(a^2x^2 + 1)}} dx & \text{for } p = -\frac{1}{2} \\ \frac{ax(a^2cx^2+c)^p e^{2ip \operatorname{atan}(ax)}}{2ap+a} - \frac{i(a^2cx^2+c)^p e^{2ip \operatorname{atan}(ax)}}{2ap+a} & \text{otherwise} \end{cases}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*I*p*atan(a*x))*(a**2*c*x**2+c)**p,x)

[Out] Piecewise((x/sqrt(c), Eq(a, 0) & Eq(p, -1/2)), (c**p*x, Eq(a, 0)), (Integral(exp(-I*atan(a*x))/sqrt(c*(a**2*x**2 + 1)), x), Eq(p, -1/2)), (a*x*(a**2*c*x**2 + c)**p*exp(2*I*p*atan(a*x))/(2*a*p + a) - I*(a**2*c*x**2 + c)**p*exp(2*I*p*atan(a*x))/(2*a*p + a), True))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(2*I*p*arctan(a*x))*(a^2*c*x^2+c)^p,x, algorithm="giac")

[Out] sage0*x

Mupad [B]

time = 0.59, size = 54, normalized size = 1.02

$$\left(\frac{x e^{p \operatorname{atan}(ax) 2i}}{2p + 1} - \frac{e^{p \operatorname{atan}(ax) 2i} 1i}{a (2p + 1)} \right) (ca^2x^2 + c)^p$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(p*atan(a*x)*2i)*(c + a^2*c*x^2)^p,x)

[Out] ((x*exp(p*atan(a*x)*2i))/(2*p + 1) - (exp(p*atan(a*x)*2i)*1i)/(a*(2*p + 1)))*(c + a^2*c*x^2)^p

$$3.374 \quad \int e^{in \operatorname{ArcTan}(ax)} x^2 (c + a^2 cx^2)^{-1 - \frac{n^2}{2}} dx$$

Optimal. Leaf size=60

$$\frac{ie^{in \operatorname{ArcTan}(ax)}(1 - ianx)(c + a^2 cx^2)^{-\frac{n^2}{2}}}{a^3 cn(1 - n^2)}$$

[Out] I*exp(I*n*arctan(a*x))*(1-I*a*n*x)/a^3/c/n/(-n^2+1)/((a^2*c*x^2+c)^(1/2*n^2))

Rubi [A]

time = 0.08, antiderivative size = 60, normalized size of antiderivative = 1.00, number of steps used = 1, number of rules used = 1, integrand size = 35, $\frac{\text{number of rules}}{\text{integrand size}} = 0.029$, Rules used = {5187}

$$\frac{i(1 - ianx)e^{in \operatorname{ArcTan}(ax)}(a^2 cx^2 + c)^{-\frac{n^2}{2}}}{a^3 cn(1 - n^2)}$$

Antiderivative was successfully verified.

[In] Int[E^(I*n*ArcTan[a*x])*x^2*(c + a^2*c*x^2)^(-1 - n^2/2), x]

[Out] (I*E^(I*n*ArcTan[a*x])*(1 - I*a*n*x))/(a^3*c*n*(1 - n^2)*(c + a^2*c*x^2)^(n^2/2))

Rule 5187

Int[E^(ArcTan[(a_.)*(x_.)]*(n_.))*(x_)^2*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Simp[(-1 - a*n*x)*(c + d*x^2)^(p + 1)*(E^(n*ArcTan[a*x]))/(a*d*n*(n^2 + 1)), x] /; FreeQ[{a, c, d, n}, x] && EqQ[d, a^2*c] && EqQ[n^2 - 2*(p + 1), 0] && !IntegerQ[I*n]

Rubi steps

$$\int e^{in \tan^{-1}(ax)} x^2 (c + a^2 cx^2)^{-1 - \frac{n^2}{2}} dx = \frac{ie^{in \tan^{-1}(ax)}(1 - ianx)(c + a^2 cx^2)^{-\frac{n^2}{2}}}{a^3 cn(1 - n^2)}$$

Mathematica [A]

time = 0.02, size = 55, normalized size = 0.92

$$-\frac{e^{in \operatorname{ArcTan}(ax)}(i + anx)(c + a^2 cx^2)^{-\frac{n^2}{2}}}{a^3 cn(-1 + n^2)}$$

Antiderivative was successfully verified.

[In] Integrate[E^(I*n*ArcTan[a*x])*x^2*(c + a^2*c*x^2)^(-1 - n^2/2), x]

[Out] -((E^(I*n*ArcTan[a*x])*(I + a*n*x))/(a^3*c*n*(-1 + n^2)*(c + a^2*c*x^2)^(n^2/2)))

Maple [A]

time = 0.11, size = 62, normalized size = 1.03

method	result	size
gospers	$\frac{(-ax+i)(ax+i)(nax+i)e^{in \arctan(ax)}(a^2cx^2+c)^{-1-\frac{n^2}{2}}}{a^3n(n^2-1)}$	62

Verification of antiderivative is not currently implemented for this CAS.

[In] int(exp(I*n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(-1-1/2*n^2), x, method=_RETURNVERBOSE)

[Out] (I-a*x)*(I+a*x)*(n*a*x+I)*exp(I*n*arctan(a*x))*(a^2*c*x^2+c)^(-1-1/2*n^2)/a^3/n/(n^2-1)

Maxima [F]

time = 0.00, size = 0, normalized size = 0.00

Failed to integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(-1-1/2*n^2), x, algorithm="maxima")

[Out] integrate((a^2*c*x^2 + c)^(-1/2*n^2 - 1)*x^2*e^(I*n*arctan(a*x)), x)

Fricas [A]

time = 7.03, size = 78, normalized size = 1.30

$$\frac{(a^3nx^3 + ia^2x^2 + anx + i)(a^2cx^2 + c)^{-\frac{1}{2}n^2-1}}{(a^3n^3 - a^3n) \left(-\frac{ax+i}{ax-i}\right)^{\frac{1}{2}n}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(-1-1/2*n^2), x, algorithm="fricas")

[Out] -(a^3*n*x^3 + I*a^2*x^2 + a*n*x + I)*(a^2*c*x^2 + c)^(-1/2*n^2 - 1)/((a^3*n^3 - a^3*n)*(-(a*x + I)/(a*x - I))^(1/2*n))

Sympy [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: SystemError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*atan(a*x))*x**2*(a**2*c*x**2+c)**(-1-1/2*n**2),x)

[Out] Exception raised: SystemError >> excessive stack use: stack is 3005 deep

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(exp(I*n*arctan(a*x))*x^2*(a^2*c*x^2+c)^(-1-1/2*n^2),x, algorithm="giac")

[Out] sage0*x

Mupad [F]

time = 0.00, size = -1, normalized size = -0.02

$$\int \frac{x^2 e^{n \operatorname{atan}(ax)} dx}{(c a^2 x^2 + c)^{\frac{n^2}{2} + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*exp(n*atan(a*x)*1i))/(c + a^2*c*x^2)^(n^2/2 + 1),x)

[Out] int((x^2*exp(n*atan(a*x)*1i))/(c + a^2*c*x^2)^(n^2/2 + 1), x)

$$3.375 \quad \int \frac{e^{6i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{19}} dx$$

Optimal. Leaf size=38

$$-\frac{i + 6ax}{210a^3c^{19}(1 - iax)^{21}(1 + iax)^{15}}$$

[Out] 1/210*(-I-6*a*x)/a^3/c^19/(1-I*a*x)^21/(1+I*a*x)^15

Rubi [A]

time = 0.05, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {5190, 82}

$$-\frac{6ax + i}{210a^3c^{19}(1 - iax)^{21}(1 + iax)^{15}}$$

Antiderivative was successfully verified.

[In] Int[(E^((6*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^19,x]

[Out] -1/210*(I + 6*a*x)/(a^3*c^19*(1 - I*a*x)^21*(1 + I*a*x)^15)

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{6i \tan^{-1}(ax)} x^2}{(c + a^2cx^2)^{19}} dx &= \frac{\int \frac{x^2}{(1-iax)^{22}(1+iax)^{16}} dx}{c^{19}} \\ &= -\frac{i + 6ax}{210a^3c^{19}(1 - iax)^{21}(1 + iax)^{15}} \end{aligned}$$

Mathematica [A]

time = 0.78, size = 36, normalized size = 0.95

$$\frac{i + 6ax}{210a^3c^{19}(-i + ax)^{15}(i + ax)^{21}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^((6*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^19,x]

[Out] (I + 6*a*x)/(210*a^3*c^19*(-I + a*x)^15*(I + a*x)^21)

Maple [A]

time = 0.55, size = 34, normalized size = 0.89

method	result	size
default	$\frac{\frac{x}{35a^2} + \frac{i}{210a^3}}{c^{19}(ax+i)^{21}(ax-i)^{15}}$	34
risch	$\frac{\frac{x}{35a^2} + \frac{i}{210a^3}}{c^{19}(ax+i)^{21}(ax-i)^{15}}$	34
gospers	$\frac{(-ax+i)(ax+i)(6ax+i)(iax+1)^6}{210a^3(a^2x^2+1)^{22}c^{19}}$	49

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^6/(a^2*x^2+1)^3*x^2/(a^2*c*x^2+c)^19,x,method=_RETURNVERBOSE)

[Out] 1/c^19*(1/35*x/a^2+1/210*I/a^3)/(I+a*x)^21/(a*x-I)^15

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 292 vs. 2(30) = 60.

time = 0.62, size = 292, normalized size = 7.68

$\frac{6a^6x^7 - 35a^5x^6 - 84a^4x^5 + 105a^3x^4 + 70a^2x^3 - 21a^2x^2 - 1}{210(a^2c^{19}x^2 + 1)^3c^{19}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^6/(a^2*x^2+1)^3*x^2/(a^2*c*x^2+c)^19,x, algorithm="maxima")

[Out] $\frac{1}{210} \cdot (6a^7x^7 - 35Ia^6x^6 - 84a^5x^5 + 105Ia^4x^4 + 70a^3x^3 - 21Ia^2x^2 - I) / (a^{45}c^{19}x^{42} + 21a^{43}c^{19}x^{40} + 210a^{41}c^{19}x^{38} + 1330a^{39}c^{19}x^{36} + 5985a^{37}c^{19}x^{34} + 20349a^{35}c^{19}x^{32} + 54264a^{33}c^{19}x^{30} + 116280a^{31}c^{19}x^{28} + 203490a^{29}c^{19}x^{26} + 293930a^{27}c^{19}x^{24} + 352716a^{25}c^{19}x^{22} + 352716a^{23}c^{19}x^{20} + 293930a^{21}c^{19}x^{18} + 203490a^{19}c^{19}x^{16} + 116280a^{17}c^{19}x^{14} + 54264a^{15}c^{19}x^{12} + 20349a^{13}c^{19}x^{10} + 5985a^{11}c^{19}x^8 + 1330a^9c^{19}x^6 + 210a^7c^{19}x^4 + 21a^5c^{19}x^2 + a^3c^{19})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 379 vs. $2(30) = 60$.
time = 9.67, size = 379, normalized size = 9.97

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)^6/(a^2*x^2+1)^3*x^2/(a^2*c*x^2+c)^19,x, algorithm="fricas")
```

```
[Out] 1/210*(6*a*x + I)/(a^39*c^19*x^36 + 6*I*a^38*c^19*x^35 + 70*I*a^36*c^19*x^33 - 105*a^35*c^19*x^32 + 336*I*a^34*c^19*x^31 - 896*a^33*c^19*x^30 + 720*I*a^32*c^19*x^29 - 3900*a^31*c^19*x^28 - 280*I*a^30*c^19*x^27 - 10752*a^29*c^19*x^26 - 6552*I*a^28*c^19*x^25 - 20020*a^27*c^19*x^24 - 21840*I*a^26*c^19*x^23 - 24960*a^25*c^19*x^22 - 43472*I*a^24*c^19*x^21 - 18018*a^23*c^19*x^20 - 60060*I*a^22*c^19*x^19 - 60060*I*a^20*c^19*x^17 + 18018*a^19*c^19*x^16 - 43472*I*a^18*c^19*x^15 + 24960*a^17*c^19*x^14 - 21840*I*a^16*c^19*x^13 + 20020*a^15*c^19*x^12 - 6552*I*a^14*c^19*x^11 + 10752*a^13*c^19*x^10 - 280*I*a^12*c^19*x^9 + 3900*a^11*c^19*x^8 + 720*I*a^10*c^19*x^7 + 896*a^9*c^19*x^6 + 336*I*a^8*c^19*x^5 + 105*a^7*c^19*x^4 + 70*I*a^6*c^19*x^3 + 6*I*a^4*c^19*x - a^3*c^19)
```

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 439 vs. $2(32) = 64$.
time = 3.69, size = 439, normalized size = 11.55

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)**6/(a**2*x**2+1)**3*x**2/(a**2*c*x**2+c)**19,x)
```

```
[Out] -(-6*a*x - I)/(210*a**39*c**19*x**36 + 1260*I*a**38*c**19*x**35 + 14700*I*a**36*c**19*x**33 - 22050*a**35*c**19*x**32 + 70560*I*a**34*c**19*x**31 - 188160*a**33*c**19*x**30 + 151200*I*a**32*c**19*x**29 - 819000*a**31*c**19*x**28 - 58800*I*a**30*c**19*x**27 - 2257920*a**29*c**19*x**26 - 1375920*I*a**28*c**19*x**25 - 4204200*a**27*c**19*x**24 - 4586400*I*a**26*c**19*x**23 - 5241600*a**25*c**19*x**22 - 9129120*I*a**24*c**19*x**21 - 3783780*a**23*c**19*x**20 - 12612600*I*a**22*c**19*x**19 - 12612600*I*a**20*c**19*x**17 + 3783780*a**19*c**19*x**16 - 9129120*I*a**18*c**19*x**15 + 5241600*a**17*c**19*x**14 - 4586400*I*a**16*c**19*x**13 + 4204200*a**15*c**19*x**12 - 1375920*I*a**14*c**19*x**11 + 2257920*a**13*c**19*x**10 - 58800*I*a**12*c**19*x**9 + 819000*a**11*c**19*x**8 + 151200*I*a**10*c**19*x**7 + 188160*a**9*c**19*x**6 + 70560*I*a**8*c**19*x**5 + 22050*a**7*c**19*x**4 + 14700*I*a**6*c**19*x**3 + 1260*I*a**4*c**19*x - 210*a**3*c**19)
```

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 299 vs. $2(30) = 60$.
time = 0.43, size = 299, normalized size = 7.87

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^6/(a^2*x^2+1)^3*x^2/(a^2*c*x^2+c)^19,x, algorithm="giac")

[Out]
$$\begin{aligned} & -1/901943132160*(358229025*a^{14}*x^{14} - 5340869100*I*a^{13}*x^{13} - 37114698075 \\ & *a^{12}*x^{12} + 159416118225*I*a^{11}*x^{11} + 473088806190*a^{10}*x^{10} - 1026819468 \\ & 675*I*a^9*x^9 - 1682288472150*a^8*x^8 + 2115551402250*I*a^7*x^7 + 205443504 \\ & 6125*a^6*x^6 - 1535397250002*I*a^5*x^5 - 870854759775*a^4*x^4 + 36430753320 \\ & 5*I*a^3*x^3 + 106553746740*a^2*x^2 - 19571887695*I*a*x - 1710785408)/((a*x \\ & - I)^{15}*a^3*c^{19}) + 1/901943132160*(358229025*a^{20}*x^{20} + 7555375800*I*a^{19} \\ & *x^{19} - 75901131600*a^{18}*x^{18} - 483051354975*I*a^{17}*x^{17} + 2184946607340*a^{16} \\ & *x^{16} + 7469205450840*I*a^{15}*x^{15} - 20031221295000*a^{14}*x^{14} - 4317700403 \\ & 7300*I*a^{13}*x^{13} + 76013078916950*a^{12}*x^{12} + 110448380006328*I*a^{11}*x^{11} - \\ & 133277726128008*a^{10}*x^{10} - 133908931763530*I*a^9*x^9 + 111933156213900*a^8 \\ & *x^8 + 77492989590120*I*a^7*x^7 - 44041557267624*a^6*x^6 - 20244576347604* \\ & I*a^5*x^5 + 7349182966545*a^4*x^4 + 2026362494800*I*a^3*x^3 - 396520754280* \\ & a^2*x^2 - 48177926223*I*a*x + 2584181888)/((a*x + I)^{21}*a^3*c^{19}) \end{aligned}$$

Mupad [F(-1)]

time = 0.00, size = -1, normalized size = -0.03

Hanged

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x+1)^6)/((c + a^2*c*x^2)^19*(a^2*x^2 + 1)^3),x)

[Out] \text{Hanged}

$$3.376 \quad \int \frac{e^{4i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^9} dx$$

Optimal. Leaf size=38

$$-\frac{i + 4ax}{60a^3c^9(1 - iax)^{10}(1 + iax)^6}$$

[Out] 1/60*(-I-4*a*x)/a^3/c^9/(1-I*a*x)^10/(1+I*a*x)^6

Rubi [A]

time = 0.06, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {5190, 82}

$$-\frac{4ax + i}{60a^3c^9(1 - iax)^{10}(1 + iax)^6}$$

Antiderivative was successfully verified.

[In] Int[(E^((4*I)*ArcTan[a*x]))*x^2]/(c + a^2*c*x^2)^9,x]

[Out] -1/60*(I + 4*a*x)/(a^3*c^9*(1 - I*a*x)^10*(1 + I*a*x)^6)

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{4i \tan^{-1}(ax)} x^2}{(c + a^2cx^2)^9} dx &= \int \frac{\frac{x^2}{(1-iax)^{11}(1+iax)^7}}{c^9} dx \\ &= -\frac{i + 4ax}{60a^3c^9(1 - iax)^{10}(1 + iax)^6} \end{aligned}$$

Mathematica [A]

time = 0.16, size = 36, normalized size = 0.95

$$-\frac{i + 4ax}{60a^3c^9(-i + ax)^6(i + ax)^{10}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^((4*I)*ArcTan[a*x]))*x^2)/(c + a^2*c*x^2)^9,x]**[Out]** -1/60*(I + 4*a*x)/(a^3*c^9*(-I + a*x)^6*(I + a*x)^10)**Maple [A]**

time = 0.30, size = 35, normalized size = 0.92

method	result	size
risch	$\frac{-\frac{i}{60a^3} - \frac{x}{15a^2}}{(ax+i)^{10}c^9(ax-i)^6}$	34
default	$-\frac{\frac{i}{60a^3} + \frac{x}{15a^2}}{c^9(ax+i)^{10}(ax-i)^6}$	35
gospers	$\frac{(-ax+i)(ax+i)(4ax+i)(iax+1)^4}{60a^3(a^2x^2+1)^{11}c^9}$	49
norman	$\frac{iax^4 + \frac{x^3}{3c} - \frac{a^2x^5}{15c} + \frac{2ia^3x^6}{c} + \frac{7ia^5x^8}{2c} + \frac{21ia^7x^{10}}{5c} + \frac{7ia^9x^{12}}{2c} + \frac{2ia^{11}x^{14}}{c} + \frac{3ia^{13}x^{16}}{4c} + \frac{ia^{15}x^{18}}{6c} + \frac{ia^{17}x^{20}}{60c}}{(a^2x^2+1)^{10}c^8}$	142

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^4/(a^2*x^2+1)^2*x^2/(a^2*c*x^2+c)^9,x,method=_RETURNVERBOSE)**[Out]** -1/c^9*(1/60*I/a^3+1/15*x/a^2)/(I+a*x)^10/(a*x-I)^6**Maxima [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 155 vs. 2(30) = 60.

time = 0.49, size = 155, normalized size = 4.08

$$\frac{4a^5x^5 - 15ia^4x^4 - 20a^3x^3 + 10ia^2x^2 + i}{60(a^{23}c^9x^{20} + 10a^{21}c^9x^{18} + 45a^{19}c^9x^{16} + 120a^{17}c^9x^{14} + 210a^{15}c^9x^{12} + 252a^{13}c^9x^{10} + 210a^{11}c^9x^8 + 120a^9c^9x^6 + 45a^7c^9x^4 + 10a^5c^9x^2 + a^3c^9)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^2/(a^2*c*x^2+c)^9,x, algorithm="maxima")**[Out]** -1/60*(4*a^5*x^5 - 15*I*a^4*x^4 - 20*a^3*x^3 + 10*I*a^2*x^2 + I)/(a^23*c^9*x^20 + 10*a^21*c^9*x^18 + 45*a^19*c^9*x^16 + 120*a^17*c^9*x^14 + 210*a^15*c^9*x^12 + 252*a^13*c^9*x^10 + 210*a^11*c^9*x^8 + 120*a^9*c^9*x^6 + 45*a^7*c^9*x^4 + 10*a^5*c^9*x^2 + a^3*c^9)**Fricas [B]** Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 169 vs. 2(30) = 60.

time = 3.90, size = 169, normalized size = 4.45

$$\frac{4ax + i}{60(a^{19}c^9x^{16} + 4ia^{18}c^9x^{15} + 20ia^{16}c^9x^{13} - 20a^{15}c^9x^{12} + 36ia^{14}c^9x^{11} - 64a^{13}c^9x^{10} + 20ia^{12}c^9x^9 - 90a^{11}c^9x^8 - 20ia^{10}c^9x^7 - 64a^9c^9x^6 - 36ia^8c^9x^5 - 20a^7c^9x^4 - 20ia^6c^9x^3 - 4ia^4c^9x + a^3c^9)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^2/(a^2*c*x^2+c)^9,x, algorithm="fricas")

[Out]
$$-1/60*(4*a*x + I)/(a^{19}*c^9*x^{16} + 4*I*a^{18}*c^9*x^{15} + 20*I*a^{16}*c^9*x^{13} - 20*a^{15}*c^9*x^{12} + 36*I*a^{14}*c^9*x^{11} - 64*a^{13}*c^9*x^{10} + 20*I*a^{12}*c^9*x^9 - 90*a^{11}*c^9*x^8 - 20*I*a^{10}*c^9*x^7 - 64*a^9*c^9*x^6 - 36*I*a^8*c^9*x^5 - 20*a^7*c^9*x^4 - 20*I*a^6*c^9*x^3 - 4*I*a^4*c^9*x + a^3*c^9)$$

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 194 vs. 2(32) = 64.

time = 1.02, size = 194, normalized size = 5.11

$$\frac{-4ax - i}{60a^{19}c^9x^{16} + 240ia^{18}c^9x^{15} + 1200ia^{16}c^9x^{13} - 1200a^{15}c^9x^{12} + 2160ia^{14}c^9x^{11} - 3840a^{13}c^9x^{10} + 1200ia^{12}c^9x^9 - 5400a^{11}c^9x^8 - 1200ia^{10}c^9x^7 - 3840a^9c^9x^6 - 2160ia^8c^9x^5 - 1200a^7c^9x^4 - 1200ia^6c^9x^3 - 240ia^4c^9x + 60a^3c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**4/(a**2*x**2+1)**2*x**2/(a**2*c*x**2+c)**9,x)

[Out]
$$(-4*a*x - I)/(60*a^{19}*c^{**9}*x^{**16} + 240*I*a^{**18}*c^{**9}*x^{**15} + 1200*I*a^{**16}*c^{**9}*x^{**13} - 1200*a^{**15}*c^{**9}*x^{**12} + 2160*I*a^{**14}*c^{**9}*x^{**11} - 3840*a^{**13}*c^{**9}*x^{**10} + 1200*I*a^{**12}*c^{**9}*x^{**9} - 5400*a^{**11}*c^{**9}*x^{**8} - 1200*I*a^{**10}*c^{**9}*x^{**7} - 3840*a^{**9}*c^{**9}*x^{**6} - 2160*I*a^{**8}*c^{**9}*x^{**5} - 1200*a^{**7}*c^{**9}*x^{**4} - 1200*I*a^{**6}*c^{**9}*x^{**3} - 240*I*a^{**4}*c^{**9}*x + 60*a^{**3}*c^{**9})$$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 139 vs. 2(30) = 60.

time = 0.41, size = 139, normalized size = 3.66

$$\frac{-2145a^5x^5 - 12540ia^4x^4 - 30030a^3x^3 + 37080ia^2x^2 + 23841ax - 6476i}{983040(ax-i)^6a^3c^9} + \frac{2145a^9x^9 + 21780ia^8x^8 - 99660a^7x^7 - 270480ia^6x^6 + 481446a^5x^5 + 584920ia^4x^4 - 486220a^3x^3 - 265680ia^2x^2 + 84065ax + 9908i}{983040(ax+i)^{10}a^3c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^4/(a^2*x^2+1)^2*x^2/(a^2*c*x^2+c)^9,x, algorithm="giac")

[Out]
$$-1/983040*(2145*a^5*x^5 - 12540*I*a^4*x^4 - 30030*a^3*x^3 + 37080*I*a^2*x^2 + 23841*a*x - 6476*I)/((a*x - I)^6*a^3*c^9) + 1/983040*(2145*a^9*x^9 + 21780*I*a^8*x^8 - 99660*a^7*x^7 - 270480*I*a^6*x^6 + 481446*a^5*x^5 + 584920*I*a^4*x^4 - 486220*a^3*x^3 - 265680*I*a^2*x^2 + 84065*a*x + 9908*I)/((a*x + I)^{10}*a^3*c^9)$$

Mupad [B]

time = 3.58, size = 160, normalized size = 4.21

$$\frac{4a^5x^5 - a^4x^4 15i - 20a^3x^3 + a^2x^2 10i + 1i}{60a^{23}c^9x^{20} + 600a^{21}c^9x^{18} + 2700a^{19}c^9x^{16} + 7200a^{17}c^9x^{14} + 12600a^{15}c^9x^{12} + 15120a^{13}c^9x^{10} + 12600a^{11}c^9x^8 + 7200a^9c^9x^6 + 2700a^7c^9x^4 + 600a^5c^9x^2 + 60a^3c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] $\text{int}((x^2*(a*x+1)^4)/((c + a^2*c*x^2)^9*(a^2*x^2 + 1)^2),x)$

[Out] $-(a^2*x^2*10i - 20*a^3*x^3 - a^4*x^4*15i + 4*a^5*x^5 + 1i)/(60*a^3*c^9 + 600*a^5*c^9*x^2 + 2700*a^7*c^9*x^4 + 7200*a^9*c^9*x^6 + 12600*a^11*c^9*x^8 + 15120*a^13*c^9*x^10 + 12600*a^15*c^9*x^12 + 7200*a^17*c^9*x^14 + 2700*a^19*c^9*x^16 + 600*a^21*c^9*x^18 + 60*a^23*c^9*x^20)$

$$3.377 \quad \int \frac{e^{2i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^3} dx$$

Optimal. Leaf size=38

$$-\frac{i+2ax}{6a^3c^3(1-iax)^3(1+iax)}$$

[Out] 1/6*(-I-2*a*x)/a^3/c^3/(1-I*a*x)^3/(1+I*a*x)

Rubi [A]

time = 0.05, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {5190, 82}

$$-\frac{2ax+i}{6a^3c^3(1-iax)^3(1+iax)}$$

Antiderivative was successfully verified.

[In] Int[(E^((2*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^3,x]

[Out] -1/6*(I + 2*a*x)/(a^3*c^3*(1 - I*a*x)^3*(1 + I*a*x))

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.)*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{2i \tan^{-1}(ax)} x^2}{(c + a^2cx^2)^3} dx &= \int \frac{\frac{x^2}{(1-iax)^4(1+iax)^2} dx}{c^3} \\ &= -\frac{i+2ax}{6a^3c^3(1-iax)^3(1+iax)} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 36, normalized size = 0.95

$$\frac{i + 2ax}{6a^3c^3(-i + ax)(i + ax)^3}$$

Antiderivative was successfully verified.

[In] Integrate[(E^((2*I)*ArcTan[a*x]))*x^2)/(c + a^2*c*x^2)^3,x]

[Out] (I + 2*a*x)/(6*a^3*c^3*(-I + a*x)*(I + a*x)^3)

Maple [A]

time = 0.11, size = 34, normalized size = 0.89

method	result	size
default	$\frac{\frac{x}{3a^2} + \frac{i}{6a^3}}{c^3(ax+i)^3(ax-i)}$	34
risch	$\frac{\frac{x}{3a^2} + \frac{i}{6a^3}}{c^3(ax+i)^3(ax-i)}$	34
norman	$\frac{\frac{x^3}{3c} + \frac{iax^4}{2c} + \frac{ia^3x^6}{6c}}{(a^2x^2+1)^3c^2}$	47
gospers	$\frac{(-ax+i)(ax+i)(2ax+i)(iax+1)^2}{6a^3(a^2x^2+1)^4c^3}$	49

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^2/(a^2*x^2+1)*x^2/(a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)

[Out] 1/c^3*(1/3*x/a^2+1/6*I/a^3)/(I+a*x)^3/(a*x-I)

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 62 vs. 2(30) = 60.

time = 0.46, size = 62, normalized size = 1.63

$$\frac{2a^3x^3 - 3ia^2x^2 - i}{6(a^9c^3x^6 + 3a^7c^3x^4 + 3a^5c^3x^2 + a^3c^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^2/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] 1/6*(2*a^3*x^3 - 3*I*a^2*x^2 - I)/(a^9*c^3*x^6 + 3*a^7*c^3*x^4 + 3*a^5*c^3*x^2 + a^3*c^3)

Fricas [A]

time = 3.10, size = 49, normalized size = 1.29

$$\frac{2ax + i}{6(a^7c^3x^4 + 2ia^6c^3x^3 + 2ia^4c^3x - a^3c^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^2/(a^2*c*x^2+c)^3,x, algorithm="fricas")

[Out] 1/6*(2*a*x + I)/(a^7*c^3*x^4 + 2*I*a^6*c^3*x^3 + 2*I*a^4*c^3*x - a^3*c^3)

Sympy [A]

time = 0.25, size = 54, normalized size = 1.42

$$-\frac{-2ax - i}{6a^7c^3x^4 + 12ia^6c^3x^3 + 12ia^4c^3x - 6a^3c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)**2/(a**2*x**2+1)*x**2/(a**2*c*x**2+c)**3,x)

[Out] -(-2*a*x - I)/(6*a**7*c**3*x**4 + 12*I*a**6*c**3*x**3 + 12*I*a**4*c**3*x - 6*a**3*c**3)

Giac [A]

time = 0.42, size = 45, normalized size = 1.18

$$-\frac{1}{16(ax - i)a^3c^3} + \frac{3a^2x^2 + 12iax - 5}{48(ax + i)^3a^3c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^2/(a^2*x^2+1)*x^2/(a^2*c*x^2+c)^3,x, algorithm="giac")

[Out] -1/16/((a*x - I)*a^3*c^3) + 1/48*(3*a^2*x^2 + 12*I*a*x - 5)/((a*x + I)^3*a^3*c^3)

Mupad [B]

time = 0.64, size = 47, normalized size = 1.24

$$\frac{\frac{x}{3a^6c^3} + \frac{1i}{6a^7c^3}}{\frac{x^2i}{a^3} - \frac{1}{a^4} + x^4 + \frac{x^3 2i}{a}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1)^2)/((c + a^2*c*x^2)^3*(a^2*x^2 + 1)),x)

[Out] (1i/(6*a^7*c^3) + x/(3*a^6*c^3))/((x^2i)/a^3 - 1/a^4 + x^4 + (x^3*2i)/a)

$$3.378 \quad \int \frac{e^{-2i \operatorname{ArcTan}(ax)} x^2}{(c + a^2 cx^2)^3} dx$$

Optimal. Leaf size=38

$$\frac{i - 2ax}{6a^3c^3(1 - iax)(1 + iax)^3}$$

[Out] 1/6*(I-2*a*x)/a^3/c^3/(1-I*a*x)/(1+I*a*x)^3

Rubi [A]

time = 0.06, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {5190, 82}

$$\frac{-2ax + i}{6a^3c^3(1 - iax)(1 + iax)^3}$$

Antiderivative was successfully verified.

[In] Int[x^2/(E^((2*I)*ArcTan[a*x]))*(c + a^2*c*x^2)^3], x]

[Out] (I - 2*a*x)/(6*a^3*c^3*(1 - I*a*x)*(1 + I*a*x)^3)

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1)))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^ArcTan[(a_.)*(x_)]*(n_.)*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-2i \tan^{-1}(ax)} x^2}{(c + a^2 cx^2)^3} dx &= \int \frac{\frac{x^2}{(1-iax)^2(1+iax)^4} dx}{c^3} \\ &= \frac{i - 2ax}{6a^3c^3(1 - iax)(1 + iax)^3} \end{aligned}$$

Mathematica [A]

time = 0.03, size = 36, normalized size = 0.95

$$\frac{-i + 2ax}{6a^3c^3(-i + ax)^3(i + ax)}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/(E^((2*I)*ArcTan[a*x])*(c + a^2*c*x^2)^3), x]

[Out] (-I + 2*a*x)/(6*a^3*c^3*(-I + a*x)^3*(I + a*x))

Maple [A]

time = 0.11, size = 62, normalized size = 1.63

method	result	size
risch	$\frac{\frac{x}{3a^2} - \frac{i}{6a^3}}{c^3(ax-i)^3(ax+i)}$	34
norman	$\frac{\frac{x^3}{3c} - \frac{iax^4}{2c} - \frac{ia^3x^6}{6c}}{(a^2x^2+1)^3c^2}$	47
gospers	$-\frac{(-2ax+i)(ax+i)(-ax+i)}{6(a^2x^2+1)^2c^3(iax+1)^2a^3}$	49
default	$-\frac{i}{8a^3(-ax+i)^2} - \frac{1}{12a^3(-ax+i)^3} - \frac{1}{16a^3(-ax+i)} - \frac{1}{16a^3(ax+i)}$	62

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^3,x,method=_RETURNVERBOSE)

[Out] 1/c^3*(-1/8*I/a^3/(I-a*x)^2-1/12/a^3/(I-a*x)^3-1/16/a^3/(I-a*x)-1/16/a^3/(I+a*x))

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^3,x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: Error executing code in Maxima: expt: undefined: 0 to a negative exponent.

Fricas [A]

time = 2.64, size = 49, normalized size = 1.29

$$\frac{2ax - i}{6(a^7c^3x^4 - 2ia^6c^3x^3 - 2ia^4c^3x - a^3c^3)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^3,x, algorithm="fricas")

[Out] 1/6*(2*a*x - I)/(a^7*c^3*x^4 - 2*I*a^6*c^3*x^3 - 2*I*a^4*c^3*x - a^3*c^3)

Sympy [A]

time = 0.23, size = 53, normalized size = 1.39

$$\frac{-2ax + i}{6a^7c^3x^4 - 12ia^6c^3x^3 - 12ia^4c^3x - 6a^3c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*a*x)**2*(a**2*x**2+1)/(a**2*c*x**2+c)**3,x)

[Out] -(-2*a*x + I)/(6*a**7*c**3*x**4 - 12*I*a**6*c**3*x**3 - 12*I*a**4*c**3*x - 6*a**3*c**3)

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 80 vs. 2(30) = 60.

time = 0.43, size = 80, normalized size = 2.11

$$\frac{1}{32a^3c^3\left(\frac{2i}{iax+1} - i\right)} - \frac{-\frac{3ia^3c^6}{iax+1} - \frac{6ia^3c^6}{(iax+1)^2} + \frac{4ia^3c^6}{(iax+1)^3}}{48a^6c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^2*(a^2*x^2+1)/(a^2*c*x^2+c)^3,x, algorithm="giac")

[Out] -1/32/(a^3*c^3*(2*I/(I*a*x + 1) - I)) - 1/48*(-3*I*a^3*c^6/(I*a*x + 1) - 6*I*a^3*c^6/(I*a*x + 1)^2 + 4*I*a^3*c^6/(I*a*x + 1)^3)/(a^6*c^9)

Mupad [B]

time = 0.59, size = 65, normalized size = 1.71

$$\frac{2a^3x^3 + a^2x^23i + 1i}{6a^9c^3x^6 + 18a^7c^3x^4 + 18a^5c^3x^2 + 6a^3c^3}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a^2*x^2 + 1))/((c + a^2*c*x^2)^3*(a*x*i + 1)^2),x)

[Out] (a^2*x^2*3i + 2*a^3*x^3 + 1i)/(6*a^3*c^3 + 18*a^5*c^3*x^2 + 18*a^7*c^3*x^4 + 6*a^9*c^3*x^6)

$$3.379 \quad \int \frac{e^{-4i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^9} dx$$

Optimal. Leaf size=38

$$\frac{i - 4ax}{60a^3c^9(1 - iax)^6(1 + iax)^{10}}$$

[Out] 1/60*(I-4*a*x)/a^3/c^9/(1-I*a*x)^6/(1+I*a*x)^10

Rubi [A]

time = 0.06, antiderivative size = 38, normalized size of antiderivative = 1.00, number of steps used = 2, number of rules used = 2, integrand size = 26, $\frac{\text{number of rules}}{\text{integrand size}} = 0.077$, Rules used = {5190, 82}

$$\frac{-4ax + i}{60a^3c^9(1 - iax)^6(1 + iax)^{10}}$$

Antiderivative was successfully verified.

[In] Int[x^2/(E^((4*I)*ArcTan[a*x])*(c + a^2*c*x^2)^9), x]

[Out] (I - 4*a*x)/(60*a^3*c^9*(1 - I*a*x)^6*(1 + I*a*x)^10)

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_.) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-4i \tan^{-1}(ax)} x^2}{(c + a^2cx^2)^9} dx &= \frac{\int \frac{x^2}{(1-iax)^7(1+iax)^{11}} dx}{c^9} \\ &= \frac{i - 4ax}{60a^3c^9(1 - iax)^6(1 + iax)^{10}} \end{aligned}$$

Mathematica [A]

time = 0.15, size = 36, normalized size = 0.95

$$\frac{i - 4ax}{60a^3c^9(-i + ax)^{10}(i + ax)^6}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/(E^((4*I)*ArcTan[a*x]))*(c + a^2*c*x^2)^9],x]

[Out] (I - 4*a*x)/(60*a^3*c^9*(-I + a*x)^10*(I + a*x)^6)

Maple [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 217 vs. 2(33) = 66.

time = 0.32, size = 218, normalized size = 5.74

method	result
risch	$\frac{\frac{i}{60a^3} - \frac{x}{15a^2}}{c^9(ax-i)^{10}(ax+i)^6}$
gospers	$-\frac{(-4ax+i)(ax+i)(-ax+i)}{60(a^2x^2+1)^7c^9(iax+1)^4a^3}$
norman	$-\frac{\frac{iax^4}{c} + \frac{x^3}{3c} - \frac{a^2x^5}{15c} - \frac{2ia^3x^6}{c} - \frac{7ia^5x^8}{2c} - \frac{21ia^7x^{10}}{5c} - \frac{7ia^9x^{12}}{2c} - \frac{2ia^{11}x^{14}}{c} - \frac{3ia^{13}x^{16}}{4c} - \frac{ia^{15}x^{18}}{6c} - \frac{ia^{17}x^{20}}{60c}}{(a^2x^2+1)^{10}c^8}$
default	$\frac{8192a^3(-ax+i)^4 + 1280a^3(-ax+i)^{10} - 1024a^3(-ax+i)^8 - 6144a^3(-ax+i)^6 - 65536a^3(-ax+i)^2 + 768a^3(-ax+i)^9 - 10240a^3(-ax+i)^5 + 4096a^3(-ax+i)^3}{c^9}$

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^9,x,method=_RETURNVERBOSE)

[Out] $1/c^9*(21/8192*I/a^3/(I-a*x)^4+1/1280*I/a^3/(I-a*x)^{10}-1/1024*I/a^3/(I-a*x)^8-7/6144*I/a^3/(I-a*x)^6-165/65536*I/a^3/(I-a*x)^2+1/768/a^3/(I-a*x)^9-21/10240/a^3/(I-a*x)^5+11/4096/a^3/(I-a*x)^3-143/65536/a^3/(I-a*x)+13/16384*I/a^3/(I+a*x)^4-1/12288*I/a^3/(I+a*x)^6-121/65536*I/a^3/(I+a*x)^2-7/20480/a^3/(I+a*x)^5+11/8192/a^3/(I+a*x)^3-143/65536/a^3/(I+a*x))$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^9,x, algorithm="maxima")

[Out] Exception raised: RuntimeError >> ECL says: Error executing code in Maxima: expt: undefined: 0 to a negative exponent.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 169 vs. $2(30) = 60$.
time = 2.67, size = 169, normalized size = 4.45

$$\frac{4ax - i}{60(a^{19}c^9x^{16} - 4ia^{18}c^9x^{15} - 20i a^{16}c^9x^{13} - 20a^{15}c^9x^{12} - 36i a^{14}c^9x^{11} - 64a^{13}c^9x^{10} - 20i a^{12}c^9x^9 - 90a^{11}c^9x^8 + 20i a^{10}c^9x^7 - 64a^9c^9x^6 + 36i a^8c^9x^5 - 20a^7c^9x^4 + 20i a^6c^9x^3 + 4i a^4c^9x + a^3c^9)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^9,x, algorithm="fricas")

[Out] $-1/60*(4*a*x - I)/(a^{19}*c^9*x^{16} - 4*I*a^{18}*c^9*x^{15} - 20*I*a^{16}*c^9*x^{13} - 20*a^{15}*c^9*x^{12} - 36*I*a^{14}*c^9*x^{11} - 64*a^{13}*c^9*x^{10} - 20*I*a^{12}*c^9*x^9 - 90*a^{11}*c^9*x^8 + 20*I*a^{10}*c^9*x^7 - 64*a^9*c^9*x^6 + 36*I*a^8*c^9*x^5 - 20*a^7*c^9*x^4 + 20*I*a^6*c^9*x^3 + 4*I*a^4*c^9*x + a^3*c^9)$

Sympy [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 192 vs. $2(31) = 62$.
time = 0.85, size = 192, normalized size = 5.05

$$\frac{-4ax + i}{60a^{19}c^9x^{16} - 240ia^{18}c^9x^{15} - 1200ia^{16}c^9x^{13} - 1200a^{15}c^9x^{12} - 2160ia^{14}c^9x^{11} - 3840a^{13}c^9x^{10} - 1200ia^{12}c^9x^9 - 5400a^{11}c^9x^8 + 1200ia^{10}c^9x^7 - 3840a^9c^9x^6 + 2160ia^8c^9x^5 - 1200a^7c^9x^4 + 1200ia^6c^9x^3 + 240ia^4c^9x + 60a^3c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*a*x)**4*(a**2*x**2+1)**2/(a**2*c*x**2+c)**9,x)

[Out] $(-4*a*x + I)/(60*a^{19}*c^{9}*x^{16} - 240*I*a^{18}*c^{9}*x^{15} - 1200*I*a^{16}*c^{9}*x^{13} - 1200*a^{15}*c^{9}*x^{12} - 2160*I*a^{14}*c^{9}*x^{11} - 3840*a^{13}*c^{9}*x^{10} - 1200*I*a^{12}*c^{9}*x^9 - 5400*a^{11}*c^{9}*x^8 + 1200*I*a^{10}*c^{9}*x^7 - 3840*a^9*c^{9}*x^6 + 2160*I*a^8*c^{9}*x^5 - 1200*a^7*c^{9}*x^4 + 1200*I*a^6*c^{9}*x^3 + 240*I*a^4*c^{9}*x + 60*a^3*c^9)$

Giac [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 139 vs. $2(30) = 60$.
time = 0.40, size = 139, normalized size = 3.66

$$\frac{2145a^5x^5 + 12540ia^4x^4 - 30030a^3x^3 - 37080ia^2x^2 + 23841ax + 6476i}{983040(ax+i)^5a^3c^9} + \frac{2145a^9x^9 - 21780ia^8x^8 - 99660a^7x^7 + 270480ia^6x^6 + 481446a^5x^5 - 584920ia^4x^4 - 486220a^3x^3 + 265680ia^2x^2 + 84065ax - 9908i}{983040(ax-i)^{10}a^3c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)^4*(a^2*x^2+1)^2/(a^2*c*x^2+c)^9,x, algorithm="giac")

[Out] $-1/983040*(2145*a^5*x^5 + 12540*I*a^4*x^4 - 30030*a^3*x^3 - 37080*I*a^2*x^2 + 23841*a*x + 6476*I)/((a*x + I)^6*a^3*c^9) + 1/983040*(2145*a^9*x^9 - 21780*I*a^8*x^8 - 99660*a^7*x^7 + 270480*I*a^6*x^6 + 481446*a^5*x^5 - 584920*I*a^4*x^4 - 486220*a^3*x^3 + 265680*I*a^2*x^2 + 84065*a*x - 9908*I)/((a*x - I)^{10}*a^3*c^9)$

Mupad [B]

time = 3.40, size = 159, normalized size = 4.18

$$\frac{-4a^5x^5 - a^4x^4 15i + 20a^3x^3 + a^2x^2 10i + 1i}{60a^{23}c^9x^{20} + 600a^{21}c^9x^{18} + 2700a^{19}c^9x^{16} + 7200a^{17}c^9x^{14} + 12600a^{15}c^9x^{12} + 15120a^{13}c^9x^{10} + 12600a^{11}c^9x^8 + 7200a^9c^9x^6 + 2700a^7c^9x^4 + 600a^5c^9x^2 + 60a^3c^9}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2*(a^2*x^2 + 1)^2)/((c + a^2*c*x^2)^9*(a*x*1i + 1)^4),x)`

[Out] `(a^2*x^2*10i + 20*a^3*x^3 - a^4*x^4*15i - 4*a^5*x^5 + 1i)/(60*a^3*c^9 + 600*a^5*c^9*x^2 + 2700*a^7*c^9*x^4 + 7200*a^9*c^9*x^6 + 12600*a^11*c^9*x^8 + 15120*a^13*c^9*x^10 + 12600*a^15*c^9*x^12 + 7200*a^17*c^9*x^14 + 2700*a^19*c^9*x^16 + 600*a^21*c^9*x^18 + 60*a^23*c^9*x^20)`

$$3.380 \quad \int \frac{e^{5i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{27/2}} dx$$

Optimal. Leaf size=65

$$-\frac{(i+5ax)\sqrt{1+a^2x^2}}{120a^3c^{13}(1-iax)^{15}(1+iax)^{10}\sqrt{c+a^2cx^2}}$$

[Out] -1/120*(I+5*a*x)*(a^2*x^2+1)^(1/2)/a^3/c^13/(1-I*a*x)^15/(1+I*a*x)^10/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.14, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {5193, 5190, 82}

$$-\frac{(5ax+i)\sqrt{a^2x^2+1}}{120a^3c^{13}(1-iax)^{15}(1+iax)^{10}\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] Int[(E^((5*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^(27/2), x]

[Out] -1/120*((I + 5*a*x)*Sqrt[1 + a^2*x^2])/(a^3*c^13*(1 - I*a*x)^15*(1 + I*a*x)^10*Sqrt[c + a^2*c*x^2])

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1)))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p])/(1 + a^2*x^2)^FracPart

[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{5i \tan^{-1}(ax)} x^2}{(c + a^2 c x^2)^{27/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{5i \tan^{-1}(ax)} x^2}{(1 + a^2 x^2)^{27/2}} dx}{c^{13} \sqrt{c + a^2 c x^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{x^2}{(1 - iax)^{16} (1 + iax)^{11}} dx}{c^{13} \sqrt{c + a^2 c x^2}} \\ &= -\frac{(i + 5ax) \sqrt{1 + a^2 x^2}}{120 a^3 c^{13} (1 - iax)^{15} (1 + iax)^{10} \sqrt{c + a^2 c x^2}} \end{aligned}$$

Mathematica [A]

time = 0.36, size = 63, normalized size = 0.97

$$\frac{(1 - 5iax) \sqrt{1 + a^2 x^2}}{120 a^3 c^{13} (-i + ax)^{10} (i + ax)^{15} \sqrt{c + a^2 c x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^((5*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^(27/2), x]

[Out] ((1 - (5*I)*a*x)*Sqrt[1 + a^2*x^2])/(120*a^3*c^13*(-I + a*x)^10*(I + a*x)^15*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.26, size = 57, normalized size = 0.88

method	result	size
default	$-\frac{\sqrt{c(a^2 x^2 + 1)} (5iax - 1)}{120 \sqrt{a^2 x^2 + 1} c^{14} a^3 (ax + i)^{15} (-ax + i)^{10}}$	57
gosper	$\frac{(-ax + i)(ax + i)(5ax + i)(iax + 1)^5}{120 a^3 (a^2 x^2 + 1)^{\frac{5}{2}} (a^2 c x^2 + c)^{\frac{27}{2}}}$	58
risch	$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{1}{120 a^3} - \frac{ix}{24 a^2} \right)}{c^{13} \sqrt{c(a^2 x^2 + 1)} (ax + i)^{15} (ax - i)^{10}}$	58

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^5/(a^2*x^2+1)^(5/2)*x^2/(a^2*c*x^2+c)^(27/2), x, method=_RETURNVERBOSE)

[Out] $-1/120/(a^2x^2+1)^{(1/2)}*(c*(a^2x^2+1))^{(1/2)}*(5*I*a*x-1)/c^{14}/a^3/(I+ax)^{15}/(I-ax)^{10}$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)*x^2/(a^2*c*x^2+c)^(27/2),x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 496 vs. $2(53) = 106$.

time = 4.95, size = 496, normalized size = 7.63

(1/120)*((I*a^22*x^25 - 5*a^21*x^24 - 40*a^19*x^22 - 50*I*a^18*x^21 - 126*a^17*x^20 - 280*I*a^16*x^19 - 160*a^15*x^18 - 765*I*a^14*x^17 + 105*a^13*x^16 - 1248*I*a^12*x^15 + 720*a^11*x^14 - 1260*I*a^10*x^13 + 1260*a^9*x^12 - 720*I*a^8*x^11 + 1248*a^7*x^10 - 105*I*a^6*x^9 + 765*a^5*x^8 + 160*I*a^4*x^7 + 280*a^3*x^6 + 126*I*a^2*x^5 + 50*a*x^4 + 40*I*x^3)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)/(a^27*c^14*x^27 + 5*I*a^26*c^14*x^26 + a^25*c^14*x^25 + 45*I*a^24*c^14*x^24 - 50*a^23*c^14*x^23 + 166*I*a^22*c^14*x^22 - 330*a^21*c^14*x^21 + 286*I*a^20*c^14*x^20 - 1045*a^19*c^14*x^19 + 55*I*a^18*c^14*x^18 - 2013*a^17*c^14*x^17 - 825*I*a^16*c^14*x^16 - 2508*a^15*c^14*x^15 - 1980*I*a^14*c^14*x^14 - 1980*a^13*c^14*x^13 - 2508*I*a^12*c^14*x^12 - 825*a^11*c^14*x^11 - 2013*I*a^10*c^14*x^10 + 55*a^9*c^14*x^9 - 1045*I*a^8*c^14*x^8 + 286*a^7*c^14*x^7 - 330*I*a^6*c^14*x^6 + 166*a^5*c^14*x^5 - 50*I*a^4*c^14*x^4 + 45*a^3*c^14*x^3 + I*a^2*c^14*x^2 + 5*a*c^14*x + I*c^14)

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)*x^2/(a^2*c*x^2+c)^(27/2),x, algorithm="fricas")`

[Out] $1/120*(I*a^{22}*x^{25} - 5*a^{21}*x^{24} - 40*a^{19}*x^{22} - 50*I*a^{18}*x^{21} - 126*a^{17}*x^{20} - 280*I*a^{16}*x^{19} - 160*a^{15}*x^{18} - 765*I*a^{14}*x^{17} + 105*a^{13}*x^{16} - 1248*I*a^{12}*x^{15} + 720*a^{11}*x^{14} - 1260*I*a^{10}*x^{13} + 1260*a^9*x^{12} - 720*I*a^8*x^{11} + 1248*a^7*x^{10} - 105*I*a^6*x^9 + 765*a^5*x^8 + 160*I*a^4*x^7 + 280*a^3*x^6 + 126*I*a^2*x^5 + 50*a*x^4 + 40*I*x^3)*sqrt(a^2*c*x^2 + c)*sqrt(a^2*x^2 + 1)/(a^{27}*c^{14}*x^{27} + 5*I*a^{26}*c^{14}*x^{26} + a^{25}*c^{14}*x^{25} + 45*I*a^{24}*c^{14}*x^{24} - 50*a^{23}*c^{14}*x^{23} + 166*I*a^{22}*c^{14}*x^{22} - 330*a^{21}*c^{14}*x^{21} + 286*I*a^{20}*c^{14}*x^{20} - 1045*a^{19}*c^{14}*x^{19} + 55*I*a^{18}*c^{14}*x^{18} - 2013*a^{17}*c^{14}*x^{17} - 825*I*a^{16}*c^{14}*x^{16} - 2508*a^{15}*c^{14}*x^{15} - 1980*I*a^{14}*c^{14}*x^{14} - 1980*a^{13}*c^{14}*x^{13} - 2508*I*a^{12}*c^{14}*x^{12} - 825*a^{11}*c^{14}*x^{11} - 2013*I*a^{10}*c^{14}*x^{10} + 55*a^9*c^{14}*x^9 - 1045*I*a^8*c^{14}*x^8 + 286*a^7*c^{14}*x^7 - 330*I*a^6*c^{14}*x^6 + 166*a^5*c^{14}*x^5 - 50*I*a^4*c^{14}*x^4 + 45*a^3*c^{14}*x^3 + I*a^2*c^{14}*x^2 + 5*a*c^{14}*x + I*c^{14})$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**5/(a**2*x**2+1)**(5/2)*x**2/(a**2*c*x**2+c)**(27/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^5/(a^2*x^2+1)^(5/2)*x^2/(a^2*c*x^2+c)^(27/2),x, algorithm="giac")

[Out] integrate((I*a*x + 1)^5*x^2/((a^2*c*x^2 + c)^(27/2)*(a^2*x^2 + 1)^(5/2)), x)

Mupad [B]

time = 2.21, size = 46, normalized size = 0.71

$$-\frac{c(a x - i)^5 (5 a x + 1i) 1i}{120 a^3 (c (a^2 x^2 + 1))^{29/2} \sqrt{a^2 x^2 + 1}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1)^5)/((c + a^2*c*x^2)^(27/2)*(a^2*x^2 + 1)^(5/2)),x)

[Out] -(c*(a*x - 1i)^5*(5*a*x + 1i)*1i)/(120*a^3*(c*(a^2*x^2 + 1))^(29/2)*(a^2*x^2 + 1)^(1/2))

$$3.381 \quad \int \frac{e^{3i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{11/2}} dx$$

Optimal. Leaf size=65

$$\frac{(i + 3ax)\sqrt{1 + a^2x^2}}{24a^3c^5(1 - iax)^6(1 + iax)^3\sqrt{c + a^2cx^2}}$$

[Out] -1/24*(I+3*a*x)*(a^2*x^2+1)^(1/2)/a^3/c^5/(1-I*a*x)^6/(1+I*a*x)^3/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.15, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {5193, 5190, 82}

$$\frac{(3ax + i)\sqrt{a^2x^2 + 1}}{24a^3c^5(1 - iax)^6(1 + iax)^3\sqrt{a^2cx^2 + c}}$$

Antiderivative was successfully verified.

[In] Int[(E^((3*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^(11/2), x]

[Out] -1/24*((I + 3*a*x)*Sqrt[1 + a^2*x^2])/(a^3*c^5*(1 - I*a*x)^6*(1 + I*a*x)^3*Sqrt[c + a^2*c*x^2])

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1)))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p])/(1 + a^2*x^2)^FracPart

[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x, x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{3i \tan^{-1}(ax)} x^2}{(c + a^2 c x^2)^{11/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{3i \tan^{-1}(ax)} x^2}{(1 + a^2 x^2)^{11/2}} dx}{c^5 \sqrt{c + a^2 c x^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{x^2}{(1 - iax)^7 (1 + iax)^4} dx}{c^5 \sqrt{c + a^2 c x^2}} \\ &= -\frac{(i + 3ax) \sqrt{1 + a^2 x^2}}{24a^3 c^5 (1 - iax)^6 (1 + iax)^3 \sqrt{c + a^2 c x^2}} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 65, normalized size = 1.00

$$\frac{i(i + 3ax) \sqrt{1 + a^2 x^2}}{24a^3 c^5 (-i + ax)^3 (i + ax)^6 \sqrt{c + a^2 c x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^((3*I)*ArcTan[a*x])*x^2)/(c + a^2*c*x^2)^(11/2), x]

[Out] ((I/24)*(I + 3*a*x)*Sqrt[1 + a^2*x^2])/(a^3*c^5*(-I + a*x)^3*(I + a*x)^6*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.13, size = 57, normalized size = 0.88

method	result	size
default	$\frac{\sqrt{c(a^2 x^2 + 1)} (3iax - 1)}{24 \sqrt{a^2 x^2 + 1} c^6 a^3 (ax + i)^6 (-ax + i)^3}$	57
gosper	$\frac{(-ax + i)(ax + i)(3ax + i)(iax + 1)^3}{24a^3(a^2 x^2 + 1)^{\frac{3}{2}}(a^2 c x^2 + c)^{\frac{11}{2}}}$	58
risch	$\frac{\sqrt{a^2 x^2 + 1} \left(\frac{ix}{8a^2} - \frac{1}{24a^3} \right)}{c^5 \sqrt{c(a^2 x^2 + 1)} (ax + i)^6 (ax - i)^3}$	58

Verification of antiderivative is not currently implemented for this CAS.

[In] int((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2/(a^2*c*x^2+c)^(11/2), x, method=_RETURNVERBOSE)

[Out] $-1/24/(a^2*x^2+1)^{(1/2)}*(c*(a^2*x^2+1))^{(1/2)}*(3*I*a*x-1)/c^6/a^3/(I+a*x)^6/(I-a*x)^3$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2/(a^2*c*x^2+c)^(11/2),x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 192 vs. $2(53) = 106$.

time = 3.07, size = 192, normalized size = 2.95

$$\frac{(i a^6 x^9 - 3 a^5 x^8 - 8 a^3 x^6 - 6 i a^2 x^5 - 6 a x^4 - 8 i x^3) \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1}}{24 (a^{11} c^6 x^{11} + 3 i a^{10} c^6 x^{10} + a^9 c^6 x^9 + 11 i a^8 c^6 x^8 - 6 a^7 c^6 x^7 + 14 i a^6 c^6 x^6 - 14 a^5 c^6 x^5 + 6 i a^4 c^6 x^4 - 11 a^3 c^6 x^3 - i a^2 c^6 x^2 - 3 a c^6 x - i c^6)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2/(a^2*c*x^2+c)^(11/2),x, algorithm="fricas")`

[Out] $1/24*(I*a^6*x^9 - 3*a^5*x^8 - 8*a^3*x^6 - 6*I*a^2*x^5 - 6*a*x^4 - 8*I*x^3)*\sqrt{a^2*c*x^2 + c}*\sqrt{a^2*x^2 + 1}/(a^{11}*c^6*x^{11} + 3*I*a^{10}*c^6*x^{10} + a^9*c^6*x^9 + 11*I*a^8*c^6*x^8 - 6*a^7*c^6*x^7 + 14*I*a^6*c^6*x^6 - 14*a^5*c^6*x^5 + 6*I*a^4*c^6*x^4 - 11*a^3*c^6*x^3 - I*a^2*c^6*x^2 - 3*a*c^6*x - I*c^6)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)**3/(a**2*x**2+1)**(3/2)*x**2/(a**2*c*x**2+c)**(11/2),x)`

[Out] $-I*(\text{Integral}(I*x**2/(a**12*c**5*x**12*\sqrt{a**2*x**2 + 1})*\sqrt{a**2*c*x**2 + c} + 6*a**10*c**5*x**10*\sqrt{a**2*x**2 + 1}*\sqrt{a**2*c*x**2 + c} + 15*a**8*c**5*x**8*\sqrt{a**2*x**2 + 1}*\sqrt{a**2*c*x**2 + c} + 20*a**6*c**5*x**6*\sqrt{a**2*x**2 + 1}*\sqrt{a**2*c*x**2 + c} + 15*a**4*c**5*x**4*\sqrt{a**2*x**2 + 1}*\sqrt{a**2*c*x**2 + c} + 6*a**2*c**5*x**2*\sqrt{a**2*x**2 + 1}*\sqrt{a**2*c*x**2 + c} + c**5*\sqrt{a**2*x**2 + 1}*\sqrt{a**2*c*x**2 + c})), x) + \text{Integral}(-3*a*x**3/(a**12*c**5*x**12*\sqrt{a**2*x**2 + 1})*\sqrt{a**2*c*x**2 + c}$

+ 6*a**10*c**5*x**10*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 15*a**8*c**5*x**8*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 20*a**6*c**5*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 15*a**4*c**5*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 6*a**2*c**5*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c**5*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a**3*x**5/(a**12*c**5*x**12*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 6*a**10*c**5*x**10*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 15*a**8*c**5*x**8*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 20*a**6*c**5*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 15*a**4*c**5*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 6*a**2*c**5*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c**5*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(-3*I*a**2*x**4/(a**12*c**5*x**12*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 6*a**10*c**5*x**10*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 15*a**8*c**5*x**8*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 20*a**6*c**5*x**6*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 15*a**4*c**5*x**4*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + 6*a**2*c**5*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c**5*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x))

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate((1+I*a*x)^3/(a^2*x^2+1)^(3/2)*x^2/(a^2*c*x^2+c)^(11/2),x, algorithm="giac")

[Out] integrate((I*a*x + 1)^3*x^2/((a^2*c*x^2 + c)^(11/2)*(a^2*x^2 + 1)^(3/2)), x)

Mupad [B]

time = 1.60, size = 48, normalized size = 0.74

$$\frac{\sqrt{c(a^2x^2 + 1)}(ax - i)^3(3ax + 1i)li}{24a^3c^6(a^2x^2 + 1)^{13/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a*x*1i + 1)^3)/((c + a^2*c*x^2)^(11/2)*(a^2*x^2 + 1)^(3/2)),x)

[Out] ((c*(a^2*x^2 + 1))^(1/2)*(a*x - 1i)^3*(3*a*x + 1i)*1i)/(24*a^3*c^6*(a^2*x^2 + 1)^(13/2))

$$3.382 \quad \int \frac{e^{i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=142

$$-\frac{\sqrt{1+a^2x^2}}{2a^3c(i+ax)\sqrt{c+a^2cx^2}} + \frac{i\sqrt{1+a^2x^2} \log(i-ax)}{4a^3c\sqrt{c+a^2cx^2}} + \frac{3i\sqrt{1+a^2x^2} \log(i+ax)}{4a^3c\sqrt{c+a^2cx^2}}$$

[Out] $-1/2*(a^2*x^2+1)^{(1/2)}/a^3/c/(I+a*x)/(a^2*c*x^2+c)^{(1/2)}+1/4*I*\ln(I-a*x)*(a^2*x^2+1)^{(1/2)}/a^3/c/(a^2*c*x^2+c)^{(1/2)}+3/4*I*\ln(I+a*x)*(a^2*x^2+1)^{(1/2)}/a^3/c/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 142, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {5193, 5190, 90}

$$-\frac{\sqrt{a^2x^2+1}}{2a^3c(ax+i)\sqrt{a^2cx^2+c}} + \frac{i\sqrt{a^2x^2+1} \log(-ax+i)}{4a^3c\sqrt{a^2cx^2+c}} + \frac{3i\sqrt{a^2x^2+1} \log(ax+i)}{4a^3c\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[(E^{(I*\text{ArcTan}[a*x])}*x^2)/(c+a^2*c*x^2)^{(3/2)}, x]$

[Out] $-1/2*\text{Sqrt}[1+a^2*x^2]/(a^3*c*(I+a*x)*\text{Sqrt}[c+a^2*c*x^2]) + ((I/4)*\text{Sqrt}[1+a^2*x^2]*\text{Log}[I-a*x])/(a^3*c*\text{Sqrt}[c+a^2*c*x^2]) + (((3*I)/4)*\text{Sqrt}[1+a^2*x^2]*\text{Log}[I+a*x])/(a^3*c*\text{Sqrt}[c+a^2*c*x^2])$

Rule 90

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_.) + (d_.)*(x_.))^{(n_.)*((e_.) + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, p\}, x] \&\& \text{IntegersQ}[m, n] \&\& (\text{IntegerQ}[p] \parallel (\text{GtQ}[m, 0] \&\& \text{GeQ}[n, -1]))$

Rule 5190

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*x_.^{(m_.)*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] :> \text{Dist}[c^p, \text{Int}[x^m*(1 - I*a*x)^{(p + I*(n/2))}*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, m, n, p\}, x] \&\& \text{EqQ}[d, a^2*c] \&\& (\text{IntegerQ}[p] \parallel \text{GtQ}[c, 0])$

Rule 5193

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*x_.^{(m_.)*((c_.) + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] :> \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[x^m*(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /; \text{FreeQ}\{a, c, d,$

m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
 \int \frac{e^{i \tan^{-1}(ax)} x^2}{(c + a^2 c x^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{i \tan^{-1}(ax)} x^2}{(1 + a^2 x^2)^{3/2}} dx}{c \sqrt{c + a^2 c x^2}} \\
 &= \frac{\sqrt{1 + a^2 x^2} \int \frac{x^2}{(1 - i a x)^2 (1 + i a x)} dx}{c \sqrt{c + a^2 c x^2}} \\
 &= \frac{\sqrt{1 + a^2 x^2} \int \left(\frac{i}{4a^2(-i+ax)} + \frac{1}{2a^2(i+ax)^2} + \frac{3i}{4a^2(i+ax)} \right) dx}{c \sqrt{c + a^2 c x^2}} \\
 &= -\frac{\sqrt{1 + a^2 x^2}}{2a^3 c (i + ax) \sqrt{c + a^2 c x^2}} + \frac{i \sqrt{1 + a^2 x^2} \log(i - ax)}{4a^3 c \sqrt{c + a^2 c x^2}} + \frac{3i \sqrt{1 + a^2 x^2} \log(i + ax)}{4a^3 c \sqrt{c + a^2 c x^2}}
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 74, normalized size = 0.52

$$\frac{\sqrt{1 + a^2 x^2} \left(-\frac{2}{i + ax} + i \log(i - ax) + 3i \log(i + ax) \right)}{4a^3 c \sqrt{c + a^2 c x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[(E^(I*ArcTan[a*x]))*x^2)/(c + a^2*c*x^2)^(3/2), x]

[Out] (Sqrt[1 + a^2*x^2]*(-2/(I + a*x) + I*Log[I - a*x] + (3*I)*Log[I + a*x]))/(4*a^3*c*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.10, size = 87, normalized size = 0.61

method	result	size
default	$\frac{\sqrt{c(a^2 x^2 + 1)} (i \ln(-ax+i)ax + 3i \ln(ax+i)ax - \ln(-ax+i) - 3 \ln(ax+i) - 2)}{4\sqrt{a^2 x^2 + 1} c^2 a^3 (ax+i)}$	87
risch	$-\frac{\sqrt{a^2 x^2 + 1}}{2c \sqrt{c(a^2 x^2 + 1)} a^3 (ax+i)} + \frac{3i \sqrt{a^2 x^2 + 1} \ln(iax-1)}{4c \sqrt{c(a^2 x^2 + 1)} a^3} + \frac{i \sqrt{a^2 x^2 + 1} \ln(-iax-1)}{4c \sqrt{c(a^2 x^2 + 1)} a^3}$	124

Verification of antiderivative is not currently implemented for this CAS.

[In] int(((1+I*a*x)/(a^2*x^2+1)^(1/2))*x^2/(a^2*c*x^2+c)^(3/2), x, method=_RETURNVERBOSE)

[Out] $1/4/(a^2*x^2+1)^{(1/2)}*(c*(a^2*x^2+1))^{(1/2)}*(I*\ln(I-a*x)*a*x+3*I*\ln(I+a*x)*a*x-\ln(I-a*x)-3*\ln(I+a*x)-2)/c^2/a^3/(I+a*x)$

Maxima [F(-2)]

time = 0.00, size = 0, normalized size = 0.00

Exception raised: RuntimeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`

[Out] Exception raised: RuntimeError >> ECL says: THROW: The catch RAT-ERR is undefined.

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`

[Out]
$$\begin{aligned} & -1/8*(3*(I*a^5*c^2*x^3 - a^4*c^2*x^2 + I*a^3*c^2*x - a^2*c^2)*\sqrt{1/(a^6*c^3)} \\ & * \log((I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^3*c*x*\sqrt{1/(a^6*c^3)} \\ & + I*a^2*x^3 + I*x)/(a^3*x^3 + I*a^2*x^2 + a*x + I)) + 3*(-I*a^5*c^2*x^3 + \\ & a^4*c^2*x^2 - I*a^3*c^2*x + a^2*c^2)*\sqrt{1/(a^6*c^3)} * \log((-I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^3*c*x*\sqrt{1/(a^6*c^3)} \\ & + I*a^2*x^3 + I*x)/(a^3*x^3 + I*a^2*x^2 + a*x + I)) - (I*a^5*c^2*x^3 - a^4*c^2*x^2 + I*a^3*c^2*x - \\ & a^2*c^2)*\sqrt{1/(a^6*c^3)} * \log((I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^3*c*x*\sqrt{1/(a^6*c^3)} \\ & - I*a^2*x^3 - I*x)/(a^3*x^3 - I*a^2*x^2 + a*x - I)) \\ & - (-I*a^5*c^2*x^3 + a^4*c^2*x^2 - I*a^3*c^2*x + a^2*c^2)*\sqrt{1/(a^6*c^3)} \\ & * \log((-I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^3*c*x*\sqrt{1/(a^6*c^3)} - \\ & I*a^2*x^3 - I*x)/(a^3*x^3 - I*a^2*x^2 + a*x - I)) + 4*(-I*a^5*c^2*x^3 + a^4 \\ & *c^2*x^2 - I*a^3*c^2*x + a^2*c^2)*\sqrt{1/(a^6*c^3)} * \log((\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^3*c*x*\sqrt{1/(a^6*c^3)} \\ & + a^2*x^3 + x)/(a^2*x^2 + 1)) \\ & + 4*(I*a^5*c^2*x^3 - a^4*c^2*x^2 + I*a^3*c^2*x - a^2*c^2)*\sqrt{1/(a^6*c^3)} \\ & * \log(-(\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1}*a^3*c*x*\sqrt{1/(a^6*c^3)} - a \\ & ^2*x^3 - x)/(a^2*x^2 + 1)) + 4*I*\sqrt{a^2*c*x^2 + c}*\sqrt{a^2*x^2 + 1}*x - \\ & 8*(a^5*c^2*x^3 + I*a^4*c^2*x^2 + a^3*c^2*x + I*a^2*c^2)*\int(1/2*\sqrt{a^2*c*x^2 + c}*\sqrt{a^2*x^2 + 1}*(2*I*a*x + 1)/(a^6*c^2*x^4 + 2*a^4*c^2*x^2 \\ & + a^2*c^2), x)/(a^5*c^2*x^3 + I*a^4*c^2*x^2 + a^3*c^2*x + I*a^2*c^2) \end{aligned}$$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$i \left(\int \left(-\frac{ix^2}{a^2cx^2\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c} + c\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}} \right) dx + \int \frac{ax^3}{a^2cx^2\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c} + c\sqrt{a^2x^2+1}\sqrt{a^2cx^2+c}} dx \right)$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)/(a**2*x**2+1)**(1/2)*x**2/(a**2*c*x**2+c)**(3/2),x)
```

```
[Out] I*(Integral(-I*x**2/(a**2*c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)
+ c*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c)), x) + Integral(a*x**3/(a**2*
c*x**2*sqrt(a**2*x**2 + 1)*sqrt(a**2*c*x**2 + c) + c*sqrt(a**2*x**2 + 1)*sq
rt(a**2*c*x**2 + c)), x))
```

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate((1+I*a*x)/(a^2*x^2+1)^(1/2)*x^2/(a^2*c*x^2+c)^(3/2),x, algorithm=
"giac")
```

```
[Out] integrate((I*a*x + 1)*x^2/((a^2*c*x^2 + c)^(3/2)*sqrt(a^2*x^2 + 1)), x)
```

Mupad [F]

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^2 (1 + a x i)}{(c a^2 x^2 + c)^{3/2} \sqrt{a^2 x^2 + 1}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^2*(a*x*i + 1))/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^(1/2)),x)
```

```
[Out] int((x^2*(a*x*i + 1))/((c + a^2*c*x^2)^(3/2)*(a^2*x^2 + 1)^(1/2)), x)
```

$$3.383 \quad \int \frac{e^{-i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{3/2}} dx$$

Optimal. Leaf size=143

$$\frac{\sqrt{1+a^2x^2}}{2a^3c(i-ax)\sqrt{c+a^2cx^2}} - \frac{3i\sqrt{1+a^2x^2} \log(i-ax)}{4a^3c\sqrt{c+a^2cx^2}} - \frac{i\sqrt{1+a^2x^2} \log(i+ax)}{4a^3c\sqrt{c+a^2cx^2}}$$

[Out] $1/2*(a^2*x^2+1)^{(1/2)}/a^3/c/(I-ax)/(a^2*c*x^2+c)^{(1/2)}-3/4*I*\ln(I-ax)*(a^2*x^2+1)^{(1/2)}/a^3/c/(a^2*c*x^2+c)^{(1/2)}-1/4*I*\ln(I+ax)*(a^2*x^2+1)^{(1/2)}/a^3/c/(a^2*c*x^2+c)^{(1/2)}$

Rubi [A]

time = 0.16, antiderivative size = 143, normalized size of antiderivative = 1.00, number of steps used = 4, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {5193, 5190, 90}

$$\frac{\sqrt{a^2x^2+1}}{2a^3c(-ax+i)\sqrt{a^2cx^2+c}} - \frac{3i\sqrt{a^2x^2+1} \log(-ax+i)}{4a^3c\sqrt{a^2cx^2+c}} - \frac{i\sqrt{a^2x^2+1} \log(ax+i)}{4a^3c\sqrt{a^2cx^2+c}}$$

Antiderivative was successfully verified.

[In] $\text{Int}[x^2/(E^{(I*\text{ArcTan}[a*x])*(c+a^2*c*x^2)^{(3/2)}), x]$

[Out] $\text{Sqrt}[1+a^2*x^2]/(2*a^3*c*(I-a*x)*\text{Sqrt}[c+a^2*c*x^2]) - (((3*I)/4)*\text{Sqrt}[1+a^2*x^2]*\text{Log}[I-a*x])/(a^3*c*\text{Sqrt}[c+a^2*c*x^2]) - ((I/4)*\text{Sqrt}[1+a^2*x^2]*\text{Log}[I+a*x])/(a^3*c*\text{Sqrt}[c+a^2*c*x^2])$

Rule 90

$\text{Int}[(a_. + (b_.)*(x_.))^{(m_.)*((c_. + (d_.)*(x_.))^{(n_.)*((e_. + (f_.)*(x_.))^{(p_.)}, x_Symbol] :> \text{Int}[\text{ExpandIntegrand}[(a + b*x)^m*(c + d*x)^n*(e + f*x)^p, x], x] /; \text{FreeQ}\{a, b, c, d, e, f, p\}, x\} \&\& \text{IntegersQ}\{m, n\} \&\& (\text{IntegerQ}\{p\} \parallel (\text{GtQ}\{m, 0\} \&\& \text{GeQ}\{n, -1\}))$

Rule 5190

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)*((c_. + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] :> \text{Dist}[c^p, \text{Int}[x^m*(1 - I*a*x)^{(p + I*(n/2)))*(1 + I*a*x)^{(p - I*(n/2))}, x], x] /; \text{FreeQ}\{a, c, d, m, n, p\}, x\} \&\& \text{EqQ}\{d, a^2*c\} \&\& (\text{IntegerQ}\{p\} \parallel \text{GtQ}\{c, 0\})$

Rule 5193

$\text{Int}[E^{(\text{ArcTan}[(a_.)*(x_.)]*(n_.))*(x_.)^{(m_.)*((c_. + (d_.)*(x_.)^2)^{(p_.)}, x_Symbol] :> \text{Dist}[c^{\text{IntPart}[p]}*((c + d*x^2)^{\text{FracPart}[p]}/(1 + a^2*x^2)^{\text{FracPart}[p]}), \text{Int}[x^m*(1 + a^2*x^2)^p * E^{(n*\text{ArcTan}[a*x])}, x], x] /; \text{FreeQ}\{a, c, d,$

m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned}
 \int \frac{e^{-i \tan^{-1}(ax)} x^2}{(c + a^2 c x^2)^{3/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-i \tan^{-1}(ax)} x^2}{(1 + a^2 x^2)^{3/2}} dx}{c \sqrt{c + a^2 c x^2}} \\
 &= \frac{\sqrt{1 + a^2 x^2} \int \frac{x^2}{(1 - iax)(1 + iax)^2} dx}{c \sqrt{c + a^2 c x^2}} \\
 &= \frac{\sqrt{1 + a^2 x^2} \int \left(\frac{1}{2a^2(-i+ax)^2} - \frac{3i}{4a^2(-i+ax)} - \frac{i}{4a^2(i+ax)} \right) dx}{c \sqrt{c + a^2 c x^2}} \\
 &= \frac{\sqrt{1 + a^2 x^2}}{2a^3 c (i - ax) \sqrt{c + a^2 c x^2}} - \frac{3i \sqrt{1 + a^2 x^2} \log(i - ax)}{4a^3 c \sqrt{c + a^2 c x^2}} - \frac{i \sqrt{1 + a^2 x^2} \log(i + ax)}{4a^3 c \sqrt{c + a^2 c x^2}}
 \end{aligned}$$

Mathematica [A]

time = 0.04, size = 75, normalized size = 0.52

$$\frac{\sqrt{1 + a^2 x^2} \left(\frac{2}{i - ax} - 3i \log(i - ax) - i \log(i + ax) \right)}{4a^3 c \sqrt{c + a^2 c x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/(E^(I*ArcTan[a*x])*(c + a^2*c*x^2)^(3/2)),x]

[Out] (Sqrt[1 + a^2*x^2]*(2/(I - a*x) - (3*I)*Log[I - a*x] - I*Log[I + a*x]))/(4*a^3*c*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.11, size = 86, normalized size = 0.60

method	result	size
default	$\frac{\sqrt{c(a^2 x^2 + 1)} (3i \ln(-ax+i)ax+i \ln(ax+i)ax+3 \ln(-ax+i)+\ln(ax+i)+2)}{4\sqrt{a^2 x^2 + 1} c^2 a^3 (-ax+i)}$	86
risch	$-\frac{\sqrt{a^2 x^2 + 1}}{2c \sqrt{c(a^2 x^2 + 1)} a^3 (ax-i)} - \frac{3i \sqrt{a^2 x^2 + 1} \ln(-iax-1)}{4c \sqrt{c(a^2 x^2 + 1)} a^3} - \frac{i \sqrt{a^2 x^2 + 1} \ln(iax-1)}{4c \sqrt{c(a^2 x^2 + 1)} a^3}$	124

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x,method=_RETURNVERBOSE)

[Out] $1/4/(a^2*x^2+1)^{(1/2)}*(c*(a^2*x^2+1))^{(1/2)}*(3*I*\ln(I-a*x)*a*x+I*\ln(I+a*x)*a*x+3*\ln(I-a*x)+\ln(I+a*x)+2)/c^2/a^3/(I-a*x)$

Maxima [A]

time = 0.27, size = 54, normalized size = 0.38

$$-\frac{\sqrt{c}}{2(a^4c^2x - ia^3c^2)} - \frac{3i \log(ax - i)}{4a^3c^{\frac{3}{2}}} - \frac{i \log(iax - 1)}{4a^3c^{\frac{3}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="maxima")`

[Out] $-1/2*\sqrt{c}/(a^4*c^2*x - I*a^3*c^2) - 3/4*I*\log(a*x - I)/(a^3*c^{(3/2)}) - 1/4*I*\log(I*a*x - 1)/(a^3*c^{(3/2)})$

Fricas [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="fricas")`

[Out] $1/8*((I*a^5*c^2*x^3 + a^4*c^2*x^2 + I*a^3*c^2*x + a^2*c^2)*\sqrt{1/(a^6*c^3)})*\log((I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*a^3*c*x*\sqrt{1/(a^6*c^3)}) + I*a^2*x^3 + I*x)/(a^3*x^3 + I*a^2*x^2 + a*x + I)) + (-I*a^5*c^2*x^3 - a^4*c^2*x^2 - I*a^3*c^2*x - a^2*c^2)*\sqrt{1/(a^6*c^3)})*\log((-I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*a^3*c*x*\sqrt{1/(a^6*c^3)}) + I*a^2*x^3 + I*x)/(a^3*x^3 + I*a^2*x^2 + a*x + I)) - 3*(I*a^5*c^2*x^3 + a^4*c^2*x^2 + I*a^3*c^2*x + a^2*c^2)*\sqrt{1/(a^6*c^3)})*\log((I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*a^3*c*x*\sqrt{1/(a^6*c^3)}) - I*a^2*x^3 - I*x)/(a^3*x^3 - I*a^2*x^2 + a*x - I)) - 3*(-I*a^5*c^2*x^3 - a^4*c^2*x^2 - I*a^3*c^2*x - a^2*c^2)*\sqrt{1/(a^6*c^3)})*\log((-I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*a^3*c*x*\sqrt{1/(a^6*c^3)}) - I*a^2*x^3 - I*x)/(a^3*x^3 - I*a^2*x^2 + a*x - I)) - 4*(I*a^5*c^2*x^3 + a^4*c^2*x^2 + I*a^3*c^2*x + a^2*c^2)*\sqrt{1/(a^6*c^3)})*\log((\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*a^3*c*x*\sqrt{1/(a^6*c^3)}) + a^2*x^3 + x)/(a^2*x^2 + 1)) - 4*(-I*a^5*c^2*x^3 - a^4*c^2*x^2 - I*a^3*c^2*x - a^2*c^2)*\sqrt{1/(a^6*c^3)})*\log(-(\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*a^3*c*x*\sqrt{1/(a^6*c^3)}) - a^2*x^3 - x)/(a^2*x^2 + 1)) + 4*I*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*x + 8*(a^5*c^2*x^3 - I*a^4*c^2*x^2 + a^3*c^2*x - I*a^2*c^2)*\int(1/2*\sqrt{a^2*c*x^2 + c})*\sqrt{a^2*x^2 + 1})*(-2*I*a*x + 1)/(a^6*c^2*x^4 + 2*a^4*c^2*x^2 + a^2*c^2), x))/(a^5*c^2*x^3 - I*a^4*c^2*x^2 + a^3*c^2*x - I*a^2*c^2)$

Sympy [F]

time = 0.00, size = 0, normalized size = 0.00

$$-i \int \frac{x^2 \sqrt{a^2 x^2 + 1}}{a^3 c x^3 \sqrt{a^2 c x^2 + c} - i a^2 c x^2 \sqrt{a^2 c x^2 + c} + a c x \sqrt{a^2 c x^2 + c} - i c \sqrt{a^2 c x^2 + c}} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x**2/(1+I*a*x)*(a**2*x**2+1)**(1/2)/(a**2*c*x**2+c)**(3/2),x)**[Out]** -I*Integral(x**2*sqrt(a**2*x**2 + 1)/(a**3*c*x**3*sqrt(a**2*c*x**2 + c) - I*a**2*c*x**2*sqrt(a**2*c*x**2 + c) + a*c*x*sqrt(a**2*c*x**2 + c) - I*c*sqrt(a**2*c*x**2 + c)), x)**Giac [F]**

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

[In] integrate(x^2/(1+I*a*x)*(a^2*x^2+1)^(1/2)/(a^2*c*x^2+c)^(3/2),x, algorithm="giac")**[Out]** integrate(sqrt(a^2*x^2 + 1)*x^2/((a^2*c*x^2 + c)^(3/2)*(I*a*x + 1)), x)**Mupad [F]**

time = 0.00, size = -1, normalized size = -0.01

$$\int \frac{x^2 \sqrt{a^2 x^2 + 1}}{(c a^2 x^2 + c)^{3/2} (1 + a x i)} dx$$

Verification of antiderivative is not currently implemented for this CAS.

[In] int((x^2*(a^2*x^2 + 1)^(1/2))/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)),x)**[Out]** int((x^2*(a^2*x^2 + 1)^(1/2))/((c + a^2*c*x^2)^(3/2)*(a*x*1i + 1)), x)

$$3.384 \quad \int \frac{e^{-3i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{11/2}} dx$$

Optimal. Leaf size=65

$$\frac{(i - 3ax)\sqrt{1 + a^2x^2}}{24a^3c^5(1 - iax)^3(1 + iax)^6\sqrt{c + a^2cx^2}}$$

[Out] 1/24*(I-3*a*x)*(a^2*x^2+1)^(1/2)/a^3/c^5/(1-I*a*x)^3/(1+I*a*x)^6/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.15, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {5193, 5190, 82}

$$\frac{(-3ax + i)\sqrt{a^2x^2 + 1}}{24a^3c^5(1 - iax)^3(1 + iax)^6\sqrt{a^2cx^2 + c}}$$

Antiderivative was successfully verified.

[In] Int[x^2/(E^((3*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(11/2)),x]

[Out] ((I - 3*a*x)*Sqrt[1 + a^2*x^2])/(24*a^3*c^5*(1 - I*a*x)^3*(1 + I*a*x)^6*Sqrt[c + a^2*c*x^2])

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1)))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p])/(1 + a^2*x^2)^FracPart

[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x, x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-3i \tan^{-1}(ax)} x^2}{(c + a^2 c x^2)^{11/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-3i \tan^{-1}(ax)} x^2}{(1 + a^2 x^2)^{11/2}} dx}{c^5 \sqrt{c + a^2 c x^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{x^2}{(1 - iax)^4 (1 + iax)^7} dx}{c^5 \sqrt{c + a^2 c x^2}} \\ &= \frac{(i - 3ax) \sqrt{1 + a^2 x^2}}{24a^3 c^5 (1 - iax)^3 (1 + iax)^6 \sqrt{c + a^2 c x^2}} \end{aligned}$$

Mathematica [A]

time = 0.08, size = 65, normalized size = 1.00

$$-\frac{i(-i + 3ax)\sqrt{1 + a^2 x^2}}{24a^3 c^5 (-i + ax)^6 (i + ax)^3 \sqrt{c + a^2 c x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/(E^((3*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(11/2)), x]

[Out] ((-1/24*I)*(-I + 3*a*x)*Sqrt[1 + a^2*x^2])/(a^3*c^5*(-I + a*x)^6*(I + a*x)^3*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.13, size = 57, normalized size = 0.88

method	result	size
risch	$\frac{-\frac{ix}{8a^2} - \frac{1}{24a^3}}{c^5 (a^2 x^2 + 1)^{\frac{5}{2}} \sqrt{c(a^2 x^2 + 1)} (ax - i)^3}$	50
default	$\frac{\sqrt{c(a^2 x^2 + 1)} (3iax + 1)}{24 \sqrt{a^2 x^2 + 1} c^6 a^3 (-ax + i)^6 (ax + i)^3}$	57
gosper	$-\frac{(-ax + i)(ax + i)(-3ax + i)(a^2 x^2 + 1)^{\frac{3}{2}}}{24a^3 (iax + 1)^3 (a^2 c x^2 + c)^{\frac{11}{2}}}$	58

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(11/2), x, method=_RETURN VERBOSE)

[Out] $-1/24/(a^2*x^2+1)^{(1/2)}*(c*(a^2*x^2+1))^{(1/2)}*(3*I*a*x+1)/c^6/a^3/(I-a*x)^6/(I+a*x)^3$

Maxima [A]

time = 0.27, size = 93, normalized size = 1.43

$$\frac{3ax - i}{24i a^{12} c^{\frac{11}{2}} x^9 + 72 a^{11} c^{\frac{11}{2}} x^8 + 192 a^9 c^{\frac{11}{2}} x^6 - 144i a^8 c^{\frac{11}{2}} x^5 + 144 a^7 c^{\frac{11}{2}} x^4 - 192i a^6 c^{\frac{11}{2}} x^3 - 72i a^4 c^{\frac{11}{2}} x - 24 a^3 c^{\frac{11}{2}}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(11/2),x, algorithm="maxima")`

[Out] $(3*a*x - I)/(24*I*a^{12}*c^{(11/2)}*x^9 + 72*a^{11}*c^{(11/2)}*x^8 + 192*a^9*c^{(11/2)}*x^6 - 144*I*a^8*c^{(11/2)}*x^5 + 144*a^7*c^{(11/2)}*x^4 - 192*I*a^6*c^{(11/2)}*x^3 - 72*I*a^4*c^{(11/2)}*x - 24*a^3*c^{(11/2)})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 192 vs. $2(53) = 106$.

time = 3.61, size = 192, normalized size = 2.95

$$\frac{(-i a^6 x^9 - 3 a^5 x^8 - 8 a^3 x^6 + 6i a^2 x^5 - 6 a x^4 + 8i x^3) \sqrt{a^2 c x^2 + c} \sqrt{a^2 x^2 + 1}}{24 (a^{11} c^6 x^{11} - 3i a^{10} c^6 x^{10} + a^9 c^6 x^9 - 11i a^8 c^6 x^8 - 6 a^7 c^6 x^7 - 14i a^6 c^6 x^6 - 14 a^5 c^6 x^5 - 6i a^4 c^6 x^4 - 11 a^3 c^6 x^3 + i a^2 c^6 x^2 - 3 a c^6 x + i c^6)}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(11/2),x, algorithm="fricas")`

[Out] $1/24*(-I*a^6*x^9 - 3*a^5*x^8 - 8*a^3*x^6 + 6*I*a^2*x^5 - 6*a*x^4 + 8*I*x^3)*\sqrt{a^2*c*x^2 + c}*\sqrt{a^2*x^2 + 1}/(a^{11}*c^6*x^{11} - 3*I*a^{10}*c^6*x^{10} + a^9*c^6*x^9 - 11*I*a^8*c^6*x^8 - 6*a^7*c^6*x^7 - 14*I*a^6*c^6*x^6 - 14*a^5*c^6*x^5 - 6*I*a^4*c^6*x^4 - 11*a^3*c^6*x^3 + I*a^2*c^6*x^2 - 3*a*c^6*x + I*c^6)$

Sympy [F(-1)] Timed out

time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2/(1+I*a*x)**3*(a**2*x**2+1)**(3/2)/(a**2*c*x**2+c)**(11/2),x)`

[Out] Timed out

Giac [F]

time = 0.00, size = 0, normalized size = 0.00

could not integrate

Verification of antiderivative is not currently implemented for this CAS.

```
[In] integrate(x^2/(1+I*a*x)^3*(a^2*x^2+1)^(3/2)/(a^2*c*x^2+c)^(11/2),x, algorithm="giac")
```

```
[Out] integrate((a^2*x^2 + 1)^(3/2)*x^2/((a^2*c*x^2 + c)^(11/2)*(I*a*x + 1)^3), x)
```

Mupad [B]

time = 1.59, size = 57, normalized size = 0.88

$$\frac{\sqrt{c(a^2x^2+1)}\sqrt{a^2x^2+1}(1+ax3i)li}{24a^3c^6(ax+1i)^4(1+ax1i)^7}$$

Verification of antiderivative is not currently implemented for this CAS.

```
[In] int((x^2*(a^2*x^2 + 1)^(3/2))/((c + a^2*c*x^2)^(11/2)*(a*x*1i + 1)^3),x)
```

```
[Out] ((c*(a^2*x^2 + 1))^(1/2)*(a^2*x^2 + 1)^(1/2)*(a*x*3i + 1)*1i)/(24*a^3*c^6*(a*x + 1i)^4*(a*x*1i + 1)^7)
```

$$3.385 \quad \int \frac{e^{-5i \operatorname{ArcTan}(ax)} x^2}{(c+a^2cx^2)^{27/2}} dx$$

Optimal. Leaf size=65

$$\frac{(i - 5ax)\sqrt{1 + a^2x^2}}{120a^3c^{13}(1 - iax)^{10}(1 + iax)^{15}\sqrt{c + a^2cx^2}}$$

[Out] 1/120*(I-5*a*x)*(a^2*x^2+1)^(1/2)/a^3/c^13/(1-I*a*x)^10/(1+I*a*x)^15/(a^2*c*x^2+c)^(1/2)

Rubi [A]

time = 0.15, antiderivative size = 65, normalized size of antiderivative = 1.00, number of steps used = 3, number of rules used = 3, integrand size = 28, $\frac{\text{number of rules}}{\text{integrand size}} = 0.107$, Rules used = {5193, 5190, 82}

$$\frac{(-5ax + i)\sqrt{a^2x^2 + 1}}{120a^3c^{13}(1 - iax)^{10}(1 + iax)^{15}\sqrt{a^2cx^2 + c}}$$

Antiderivative was successfully verified.

[In] Int[x^2/(E^((5*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(27/2)),x]

[Out] ((I - 5*a*x)*Sqrt[1 + a^2*x^2])/(120*a^3*c^13*(1 - I*a*x)^10*(1 + I*a*x)^15*Sqrt[c + a^2*c*x^2])

Rule 82

Int[((a_.) + (b_.)*(x_))^(2*((c_.) + (d_.)*(x_))^(n_.))*((e_.) + (f_.)*(x_))^(p_.), x_Symbol] :> Simp[b*(c + d*x)^(n + 1)*(e + f*x)^(p + 1)*((2*a*d*f*(n + p + 3) - b*(d*e*(n + 2) + c*f*(p + 2)) + b*d*f*(n + p + 2)*x)/(d^2*f^2*(n + p + 2)*(n + p + 3)), x] /; FreeQ[{a, b, c, d, e, f, n, p}, x] && NeQ[n + p + 2, 0] && NeQ[n + p + 3, 0] && EqQ[d*f*(n + p + 2)*(a^2*d*f*(n + p + 3) - b*(b*c*e + a*(d*e*(n + 1) + c*f*(p + 1)))) - b*(d*e*(n + 1) + c*f*(p + 1))*(a*d*f*(n + p + 4) - b*(d*e*(n + 2) + c*f*(p + 2))), 0]

Rule 5190

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^p, Int[x^m*(1 - I*a*x)^(p + I*(n/2))*(1 + I*a*x)^(p - I*(n/2)), x], x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && (IntegerQ[p] || GtQ[c, 0])

Rule 5193

Int[E^(ArcTan[(a_.)*(x_)]*(n_.))*(x_)^(m_.)*((c_) + (d_.)*(x_)^2)^(p_.), x_Symbol] :> Dist[c^IntPart[p]*((c + d*x^2)^FracPart[p])/(1 + a^2*x^2)^FracPart

[p]), Int[x^m*(1 + a^2*x^2)^p*E^(n*ArcTan[a*x]), x, x] /; FreeQ[{a, c, d, m, n, p}, x] && EqQ[d, a^2*c] && !(IntegerQ[p] || GtQ[c, 0])

Rubi steps

$$\begin{aligned} \int \frac{e^{-5i \tan^{-1}(ax)} x^2}{(c + a^2 c x^2)^{27/2}} dx &= \frac{\sqrt{1 + a^2 x^2} \int \frac{e^{-5i \tan^{-1}(ax)} x^2}{(1 + a^2 x^2)^{27/2}} dx}{c^{13} \sqrt{c + a^2 c x^2}} \\ &= \frac{\sqrt{1 + a^2 x^2} \int \frac{x^2}{(1 - iax)^{11} (1 + iax)^{16}} dx}{c^{13} \sqrt{c + a^2 c x^2}} \\ &= \frac{(i - 5ax) \sqrt{1 + a^2 x^2}}{120 a^3 c^{13} (1 - iax)^{10} (1 + iax)^{15} \sqrt{c + a^2 c x^2}} \end{aligned}$$

Mathematica [A]

time = 0.39, size = 63, normalized size = 0.97

$$\frac{(1 + 5iax) \sqrt{1 + a^2 x^2}}{120 a^3 c^{13} (-i + ax)^{15} (i + ax)^{10} \sqrt{c + a^2 c x^2}}$$

Antiderivative was successfully verified.

[In] Integrate[x^2/(E^((5*I)*ArcTan[a*x])*(c + a^2*c*x^2)^(27/2)), x]

[Out] ((1 + (5*I)*a*x)*Sqrt[1 + a^2*x^2])/(120*a^3*c^13*(-I + a*x)^15*(I + a*x)^10*Sqrt[c + a^2*c*x^2])

Maple [A]

time = 0.20, size = 57, normalized size = 0.88

method	result	size
risch	$\frac{\frac{1}{120a^3} + \frac{ix}{24a^2}}{c^{13}(a^2x^2+1)^{\frac{19}{2}} \sqrt{c(a^2x^2+1)} (ax-i)^5}$	50
default	$-\frac{\sqrt{c(a^2x^2+1)} (5iax+1)}{120\sqrt{a^2x^2+1} c^{14}a^3(ax+i)^{10}(-ax+i)^{15}}$	57
gospers	$-\frac{(-ax+i)(ax+i)(-5ax+i)(a^2x^2+1)^{\frac{5}{2}}}{120a^3(iax+1)^5(a^2cx^2+c)^{\frac{27}{2}}}$	58

Verification of antiderivative is not currently implemented for this CAS.

[In] int(x^2/(1+I*a*x)^5*(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(27/2), x, method=_RETURNVERBOSE)

[Out] $-1/120/(a^2x^2+1)^{(1/2)}*(c*(a^2x^2+1))^{(1/2)}*(5*I*a*x+1)/c^{14}/a^3/(I+a*x)^{10}/(I-a*x)^{15}$

Maxima [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 274 vs. $2(53) = 106$.
time = 0.34, size = 274, normalized size = 4.22

$\frac{5\sqrt{c}x + \sqrt{c}}{120(a^2x^2+1)^{15}c^{14} - 5a^2c^{14}x^{15} - 40a^2c^{14}x^{14} - 50a^2c^{14}x^{13} - 126a^2c^{14}x^{12} - 280a^2c^{14}x^{11} - 160a^2c^{14}x^{10} - 765a^2c^{14}x^9 + 105a^2c^{14}x^8 - 1248a^2c^{14}x^7 + 720a^2c^{14}x^6 - 1260a^2c^{14}x^5 + 1260a^2c^{14}x^4 - 1248a^2c^{14}x^3 - 160a^2c^{14}x^2 + 160a^2c^{14}x - 1a^2c^{14}}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^5*(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(27/2),x, algorithm="maxima")`

[Out] $1/120*(5*I*a*\sqrt{c}*x + \sqrt{c})/(a^{28}*c^{14}*x^{25} - 5*I*a^{27}*c^{14}*x^{24} - 40*I*a^{25}*c^{14}*x^{22} - 50*a^{24}*c^{14}*x^{21} - 126*I*a^{23}*c^{14}*x^{20} - 280*a^{22}*c^{14}*x^{19} - 160*I*a^{21}*c^{14}*x^{18} - 765*a^{20}*c^{14}*x^{17} + 105*I*a^{19}*c^{14}*x^{16} - 1248*a^{18}*c^{14}*x^{15} + 720*I*a^{17}*c^{14}*x^{14} - 1260*a^{16}*c^{14}*x^{13} + 1260*I*a^{15}*c^{14}*x^{12} - 720*a^{14}*c^{14}*x^{11} + 1248*I*a^{13}*c^{14}*x^{10} - 105*a^{12}*c^{14}*x^9 + 765*I*a^{11}*c^{14}*x^8 + 160*a^{10}*c^{14}*x^7 + 280*I*a^9*c^{14}*x^6 + 126*a^8*c^{14}*x^5 + 50*I*a^7*c^{14}*x^4 + 40*a^6*c^{14}*x^3 + 5*a^4*c^{14}*x - I*a^3*c^{14})$

Fricas [B] Both result and optimal contain complex but leaf count of result is larger than twice the leaf count of optimal. 496 vs. $2(53) = 106$.
time = 3.07, size = 496, normalized size = 7.63

$\frac{(-4a^2c^{14}x^{25} - 5a^2c^{14}x^{24} - 40a^2c^{14}x^{22} + 50a^2c^{14}x^{21} - 126a^2c^{14}x^{20} + 280a^2c^{14}x^{19} - 160a^2c^{14}x^{18} + 765a^2c^{14}x^{17} + 105a^2c^{14}x^{16} - 1248a^2c^{14}x^{15} + 720a^2c^{14}x^{14} + 1260a^2c^{14}x^{13} + 1260a^2c^{14}x^{12} + 720a^2c^{14}x^{11} + 1248a^2c^{14}x^{10} + 105a^2c^{14}x^9 + 765a^2c^{14}x^8 - 160a^2c^{14}x^7 + 280a^2c^{14}x^6 - 126a^2c^{14}x^5 + 50a^2c^{14}x^4 - 40a^2c^{14}x^3 + 5a^2c^{14}x - I*a^2c^{14})\sqrt{a^2c^{14}x^2 + c}\sqrt{a^2x^2 + 1}}{120(a^{28}c^{14}x^{27} - 5Ia^{26}c^{14}x^{26} + a^{25}c^{14}x^{25} - 45Ia^{24}c^{14}x^{24} - 50a^{23}c^{14}x^{23} - 166Ia^{22}c^{14}x^{22} - 330a^{21}c^{14}x^{21} - 286Ia^{20}c^{14}x^{20} - 1045a^{19}c^{14}x^{19} - 55Ia^{18}c^{14}x^{18} - 2013a^{17}c^{14}x^{17} + 825Ia^{16}c^{14}x^{16} - 2508a^{15}c^{14}x^{15} + 1980Ia^{14}c^{14}x^{14} - 1980a^{13}c^{14}x^{13} + 2508Ia^{12}c^{14}x^{12} - 825a^{11}c^{14}x^{11} + 2013Ia^{10}c^{14}x^{10} + 55a^9c^{14}x^9 + 1045Ia^8c^{14}x^8 + 286a^7c^{14}x^7 + 330Ia^6c^{14}x^6 + 166a^5c^{14}x^5 + 50Ia^4c^{14}x^4 + 45a^3c^{14}x^3 - Ia^2c^{14}x^2 + 5a^2c^{14}x - I*c^{14})}$

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^5*(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(27/2),x, algorithm="fricas")`

[Out] $1/120*(-I*a^{22}*x^{25} - 5*a^{21}*x^{24} - 40*a^{19}*x^{22} + 50*I*a^{18}*x^{21} - 126*a^{17}*x^{20} + 280*I*a^{16}*x^{19} - 160*a^{15}*x^{18} + 765*I*a^{14}*x^{17} + 105*a^{13}*x^{16} + 1248*I*a^{12}*x^{15} + 720*a^{11}*x^{14} + 1260*I*a^{10}*x^{13} + 1260*a^9*x^{12} + 720*I*a^8*x^{11} + 1248*a^7*x^{10} + 105*I*a^6*x^9 + 765*a^5*x^8 - 160*I*a^4*x^7 + 280*a^3*x^6 - 126*I*a^2*x^5 + 50*a*x^4 - 40*I*x^3)*\sqrt{a^2*c*x^2 + c}\sqrt{a^2*x^2 + 1}/(a^{27}*c^{14}*x^{27} - 5*I*a^{26}*c^{14}*x^{26} + a^{25}*c^{14}*x^{25} - 45*I*a^{24}*c^{14}*x^{24} - 50*a^{23}*c^{14}*x^{23} - 166*I*a^{22}*c^{14}*x^{22} - 330*a^{21}*c^{14}*x^{21} - 286*I*a^{20}*c^{14}*x^{20} - 1045*a^{19}*c^{14}*x^{19} - 55*I*a^{18}*c^{14}*x^{18} - 2013*a^{17}*c^{14}*x^{17} + 825*I*a^{16}*c^{14}*x^{16} - 2508*a^{15}*c^{14}*x^{15} + 1980*I*a^{14}*c^{14}*x^{14} - 1980*a^{13}*c^{14}*x^{13} + 2508*I*a^{12}*c^{14}*x^{12} - 825*a^{11}*c^{14}*x^{11} + 2013*I*a^{10}*c^{14}*x^{10} + 55*a^9*c^{14}*x^9 + 1045*I*a^8*c^{14}*x^8 + 286*a^7*c^{14}*x^7 + 330*I*a^6*c^{14}*x^6 + 166*a^5*c^{14}*x^5 + 50*I*a^4*c^{14}*x^4 + 45*a^3*c^{14}*x^3 - I*a^2*c^{14}*x^2 + 5*a^2*c^{14}*x - I*c^{14})$

Sympy [F(-1)] Timed out
time = 0.00, size = 0, normalized size = 0.00

Timed out

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x**2/(1+I*a*x)**5*(a**2*x**2+1)**(5/2)/(a**2*c*x**2+c)**(27/2),x)`

[Out] Timed out

Giac [F(-2)]
time = 0.00, size = 0, normalized size = 0.00

Exception raised: TypeError

Verification of antiderivative is not currently implemented for this CAS.

[In] `integrate(x^2/(1+I*a*x)^5*(a^2*x^2+1)^(5/2)/(a^2*c*x^2+c)^(27/2),x, algorithm="giac")`

[Out] Exception raised: TypeError >> An error occurred running a Giac command:INP
UT:sage2:=int(sage0,sageVARx):;OUTPUT:Warning, choosing root of [1,0,%%{-2
, [2,1,2]%%}+%%{-2, [2,0,2]%%}+%%{-2, [0,1,0]%%}+%%{-2, [0,0,0]%%},0,%%
{1, [4,

Mupad [B]
time = 3.09, size = 47, normalized size = 0.72

$$\frac{c^2 \sqrt{a^2 x^2 + 1} (a x + 1i)^5 (1 + a x 5i)}{120 a^3 (c (a^2 x^2 + 1))^{31/2}}$$

Verification of antiderivative is not currently implemented for this CAS.

[In] `int((x^2*(a^2*x^2 + 1)^(5/2))/((c + a^2*c*x^2)^(27/2)*(a*x*1i + 1)^5),x)`

[Out] `(c^2*(a^2*x^2 + 1)^(1/2)*(a*x + 1i)^5*(a*x*5i + 1))/(120*a^3*(c*(a^2*x^2 + 1))^(31/2))`

Chapter 4

Appendix

Local contents

4.1	Download section	1720
4.2	Listing of Grading functions	1720

4.1 Download section

The following zip files contain the raw integrals used in this test.

Mathematica format Mathematica_syntax.zip

Maple and Mupad format Maple_syntax.zip

Sympy format SYMPY_syntax.zip

Sage math format SAGE_syntax.zip

4.2 Listing of Grading functions

The following are the current version of the grading functions used for grading the quality of the antiderivative with reference to the optimal antiderivative included in the test suite.

There is a version for Maple and for Mathematica/Rubi. There is a version for grading Sympy and version for use with Sagemath.

The following are links to the current source code.

The following are the listings of source code of the grading functions.

4.2.1 Mathematica and Rubi grading function

```
(* Original version thanks to Albert Rich emailed on 03/21/2017 *)
(* ::Package:: *)

(* Nasser: April 7, 2022. add second output which gives reason for the grade *)
(*           Small rewrite of logic in main function to make it*)
(*           match Maple's logic. No change in functionality otherwise*)

(* ::Subsection:: *)
(*GradeAntiderivative[result,optimal]*)

(* ::Text:: *)
(*If result and optimal are mathematical expressions, *)
(*           GradeAntiderivative[result,optimal] returns*)
(* "F" if the result fails to integrate an expression that*)
(*           is integrable*)
(* "C" if result involves higher level functions than necessary*)
(* "B" if result is more than twice the size of the optimal*)
(*           antiderivative*)
(* "A" if result can be considered optimal*)
```

```

GradeAntiderivative[result_,optimal_] := Module[{expnResult,expnOptimal,leafCountResult,leafC
  expnResult = ExpnType[result];
  expnOptimal = ExpnType[optimal];
  leafCountResult = LeafCount[result];
  leafCountOptimal = LeafCount[optimal];

  (*Print["expnResult=",expnResult," expnOptimal=",expnOptimal];*)
  If[expnResult<=expnOptimal,
    If[Not[FreeQ[result,Complex]], (*result contains complex*)
      If[Not[FreeQ[optimal,Complex]], (*optimal contains complex*)
        If[leafCountResult<=2*leafCountOptimal,
          finalresult={"A","none"}
          ,(*ELSE*)
          finalresult={"B","Both result and optimal contain complex but leaf count
        ]
        ,(*ELSE*)
        finalresult={"C","Result contains complex when optimal does not."}
      ]
      ,(*ELSE*)(*result does not contains complex*)
      If[leafCountResult<=2*leafCountOptimal,
        finalresult={"A","none"}
        ,(*ELSE*)
        finalresult={"B","Leaf count is larger than twice the leaf count of optimal. $
      ]
    ]
    ,(*ELSE*)(*expnResult>expnOptimal*)
    If[FreeQ[result,Integrate] && FreeQ[result,Int],
      finalresult={"C","Result contains higher order function than in optimal. Order "<
    ,
    finalresult={"F","Contains unresolved integral."}
  ]
];

finalresult
]

(* ::Text:: *)
(*The following summarizes the type number assigned an *)
(*expression based on the functions it involves*)
(*1 = rational function*)
(*2 = algebraic function*)
(*3 = elementary function*)
(*4 = special function*)
(*5 = hyperpergeometric function*)
(*6 = appell function*)
(*7 = rootsum function*)
(*8 = integrate function*)

```

(*9 = unknown function*)

```
ExpnType[expn_] :=
  If[AtomQ[expn],
    1,
    If[ListQ[expn],
      Max[Map[ExpnType, expn]],
      If[Head[expn]===Power,
        If[IntegerQ[expn[[2]]],
          ExpnType[expn[[1]]],
          If[Head[expn[[2]]]===Rational,
            If[IntegerQ[expn[[1]]] || Head[expn[[1]]]===Rational,
              1,
              Max[ExpnType[expn[[1]], 2]],
            Max[ExpnType[expn[[1]], ExpnType[expn[[2]], 3]],
          If[Head[expn]===Plus || Head[expn]===Times,
            Max[ExpnType[First[expn]], ExpnType[Rest[expn]]],
          If[ElementaryFunctionQ[Head[expn]],
            Max[3, ExpnType[expn[[1]]]],
          If[SpecialFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 4]],
          If[HypergeometricFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 5]],
          If[AppellFunctionQ[Head[expn]],
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 6]],
          If[Head[expn]===RootSum,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 7]],
          If[Head[expn]===Integrate || Head[expn]===Int,
            Apply[Max, Append[Map[ExpnType, Apply[List, expn]], 8]],
          9]]]]]]]]]]]
```

```
ElementaryFunctionQ[func_] :=
  MemberQ[{
  Exp, Log,
  Sin, Cos, Tan, Cot, Sec, Csc,
  ArcSin, ArcCos, ArcTan, ArcCot, ArcSec, ArcCsc,
  Sinh, Cosh, Tanh, Coth, Sech, CsCh,
  ArcSinh, ArcCosh, ArcTanh, ArcCoth, ArcSech, ArcCsCh
  }, func]
```

```
SpecialFunctionQ[func_] :=
  MemberQ[{
  Erf, Erfc, Erfi,
  FresnelS, FresnelC,
```

```

ExpIntegralE, ExpIntegralEi, LogIntegral,
SinIntegral, CosIntegral, SinhIntegral, CoshIntegral,
Gamma, LogGamma, PolyGamma,
Zeta, PolyLog, ProductLog,
EllipticF, EllipticE, EllipticPi
},func]

HypergeometricFunctionQ[func_] :=
  MemberQ[{Hypergeometric1F1,Hypergeometric2F1,HypergeometricPFQ},func]

AppellFunctionQ[func_] :=
  MemberQ[{AppellF1},func]

```

4.2.2 Maple grading function

```

# File: GradeAntiderivative.mpl
# Original version thanks to Albert Rich emailed on 03/21/2017

#Nasser 03/22/2017 Use Maple leaf count instead since buildin
#Nasser 03/23/2017 missing 'ln' for ElementaryFunctionQ added
#Nasser 03/24/2017 corrected the check for complex result
#Nasser 10/27/2017 check for leafsize and do not call ExpnType()
#
# if leaf size is "too large". Set at 500,000
#Nasser 12/22/2019 Added debug flag, added 'dilog' to special functions
#
# see problem 156, file Apostol_Problems
#Nasser 4/07/2022 add second output which gives reason for the grade

GradeAntiderivative := proc(result,optimal)
local leaf_count_result,
      leaf_count_optimal,
      ExpnType_result,
      ExpnType_optimal,
      debug:=false;

  leaf_count_result:=leafcount(result);
  #do NOT call ExpnType() if leaf size is too large. Recursion problem
  if leaf_count_result > 500000 then
    return "B","result has leaf size over 500,000. Avoiding possible recursion issues";
  fi;

  leaf_count_optimal := leafcount(optimal);
  ExpnType_result := ExpnType(result);
  ExpnType_optimal := ExpnType(optimal);

```

```

    if debug then
        print("ExpnType_result",ExpnType_result," ExpnType_optimal=",ExpnType_optimal);
    fi;

# If result and optimal are mathematical expressions,
# GradeAntiderivative[result,optimal] returns
# "F" if the result fails to integrate an expression that
#   is integrable
# "C" if result involves higher level functions than necessary
# "B" if result is more than twice the size of the optimal
#   antiderivative
# "A" if result can be considered optimal

#This check below actually is not needed, since I only
#call this grading only for passed integrals. i.e. I check
#for "F" before calling this. But no harm of keeping it here.
#just in case.

if not type(result,freeof('int')) then
    return "F","Result contains unresolved integral";
fi;

if ExpnType_result<=ExpnType_optimal then
    if debug then
        print("ExpnType_result<=ExpnType_optimal");
    fi;
    if is_contains_complex(result) then
        if is_contains_complex(optimal) then
            if debug then
                print("both result and optimal complex");
            fi;
            if leaf_count_result<=2*leaf_count_optimal then
                return "A","";
            else
                return "B",cat("Both result and optimal contain complex but leaf count of r
                    convert(leaf_count_result,string)," vs. $2 (" ,
                    convert(leaf_count_optimal,string)," ) = ",convert(2*leaf_co

        end if
    else #result contains complex but optimal is not
        if debug then
            print("result contains complex but optimal is not");
        fi;
        return "C","Result contains complex when optimal does not.";
    fi;
else # result do not contain complex

```



```

    # this assumes optimal do not as well. No check is needed here.
    if debug then
        print("result do not contain complex, this assumes optimal do not as well")
    fi;
    if leaf_count_result<=2*leaf_count_optimal then
        if debug then
            print("leaf_count_result<=2*leaf_count_optimal");
        fi;
        return "A","";
    else
        if debug then
            print("leaf_count_result>2*leaf_count_optimal");
        fi;
        return "B",cat("Leaf count of result is larger than twice the leaf count of o
                        convert(leaf_count_result,string)," $ vs. $2(",
                        convert(leaf_count_optimal,string),")=",convert(2*leaf_cou

    fi;
    fi;
else #ExpnType(result) > ExpnType(optimal)
    if debug then
        print("ExpnType(result) > ExpnType(optimal)");
    fi;
    return "C",cat("Result contains higher order function than in optimal. Order ",
                  convert(ExpnType_result,string)," vs. order ",
                  convert(ExpnType_optimal,string),".");
fi;

end proc:

#
# is_contains_complex(result)
# takes expressions and returns true if it contains "I" else false
#
#Nasser 032417
is_contains_complex:= proc(expression)
    return (has(expression,I));
end proc:

# The following summarizes the type number assigned an expression
# based on the functions it involves
# 1 = rational function
# 2 = algebraic function
# 3 = elementary function
# 4 = special function
# 5 = hyperpergeometric function
# 6 = appell function
# 7 = rootsum function

```

```

# 8 = integrate function
# 9 = unknown function

ExpnType := proc(expn)
  if type(expn,'atomic') then
    1
  elif type(expn,'list') then
    apply(max,map(ExpnType,expn))
  elif type(expn,'sqrt') then
    if type(op(1,expn),'rational') then
      1
    else
      max(2,ExpnType(op(1,expn)))
    end if
  elif type(expn,'^^') then
    if type(op(2,expn),'integer') then
      ExpnType(op(1,expn))
    elif type(op(2,expn),'rational') then
      if type(op(1,expn),'rational') then
        1
      else
        max(2,ExpnType(op(1,expn)))
      end if
    else
      max(3,ExpnType(op(1,expn)),ExpnType(op(2,expn)))
    end if
  elif type(expn,'+`') or type(expn,'*`') then
    max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
  elif ElementaryFunctionQ(op(0,expn)) then
    max(3,ExpnType(op(1,expn)))
  elif SpecialFunctionQ(op(0,expn)) then
    max(4,apply(max,map(ExpnType,[op(expn)])))
  elif HypergeometricFunctionQ(op(0,expn)) then
    max(5,apply(max,map(ExpnType,[op(expn)])))
  elif AppellFunctionQ(op(0,expn)) then
    max(6,apply(max,map(ExpnType,[op(expn)])))
  elif op(0,expn)='int' then
    max(8,apply(max,map(ExpnType,[op(expn)]))) else
    9
  end if
end proc:

ElementaryFunctionQ := proc(func)
  member(func,[
    exp,log,ln,
    sin,cos,tan,cot,sec,csc,

```

```

    arcsin,arccos,arctan,arccot,arcsec,arccsc,
    sinh,cosh,tanh,coth,sech,csch,
    arcsinh,arccosh,arctanh,arccoth,arcsech,arccsch])
end proc:

SpecialFunctionQ := proc(func)
  member(func, [
    erf,erfc,erfi,
    FresnelS,FresnelC,
    Ei,Ei,Li,Si,Ci,Shi,Chi,
    GAMMA,lnGAMMA,Psi,Zeta,polylog,dilog,LambertW,
    EllipticF,EllipticE,EllipticPi])
end proc:

HypergeometricFunctionQ := proc(func)
  member(func, [Hypergeometric1F1,hypergeom,HypergeometricPFQ])
end proc:

AppellFunctionQ := proc(func)
  member(func, [AppellF1])
end proc:

# u is a sum or product.  rest(u) returns all but the
# first term or factor of u.
rest := proc(u) local v;
  if nops(u)=2 then
    op(2,u)
  else
    apply(op(0,u),op(2..nops(u),u))
  end if
end proc:

#leafcount(u) returns the number of nodes in u.
#Nasser 3/23/17 Replaced by build-in leafCount from package in Maple
leafcount := proc(u)
  MmaTranslator[Mma][LeafCount](u);
end proc:

```

4.2.3 Sympy grading function

```

#Dec 24, 2019. Nasser M. Abbasi:
#      Port of original Maple grading function by
#      Albert Rich to use with Sympy/Python
#Dec 27, 2019 Nasser. Added `RootSum`. See problem 177, Timofeev file
#      added 'exp_polar'
from sympy import *

def leaf_count(expr):
    #sympy do not have leaf count function. This is approximation
    return round(1.7*count_ops(expr))

def is_sqrt(expr):
    if isinstance(expr,Pow):
        if expr.args[1] == Rational(1,2):
            return True
        else:
            return False
    else:
        return False

def is_elementary_function(func):
    return func in [exp,log,ln,sin,cos,tan,cot,sec,csc,
        asin,acos,atan,acot,asec,acsc,sinh,cosh,tanh,coth,sech,csch,
        asinh,acosh,atanh,acoth,asech,acsch
    ]

def is_special_function(func):
    return func in [ erf,erfc,erfi,
        fresnels,fresnelc,Ei,Ei,Li,Si,Ci,Shi,Chi,
        gamma,loggamma,digamma,zeta,polylog,LambertW,
        elliptic_f,elliptic_e,elliptic_pi,exp_polar
    ]

def is_hypergeometric_function(func):
    return func in [hyper]

def is_appell_function(func):
    return func in [appellf1]

def is_atom(expn):
    try:
        if expn.isAtom or isinstance(expn,int) or isinstance(expn,float):
            return True
        else:
            return False

```

```

except AttributeError as error:
    return False

def expnType(expn):
    debug=False
    if debug:
        print("expn=",expn,"type(expn)=",type(expn))

    if is_atom(expn):
        return 1
    elif isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
            return 1
        else:
            return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
    elif isinstance(expn,Pow): #type(expn,'^')
        if isinstance(expn.args[1],Integer): #type(op(2,expn),'integer')
            return expnType(expn.args[0]) #ExpnType(op(1,expn))
        elif isinstance(expn.args[1],Rational): #type(op(2,expn),'rational')
            if isinstance(expn.args[0],Rational): #type(op(1,expn),'rational')
                return 1
            else:
                return max(2,expnType(expn.args[0])) #max(2,ExpnType(op(1,expn)))
        else:
            return max(3,expnType(expn.args[0]),expnType(expn.args[1])) #max(3,ExpnType(op(1,expn)),ExpnT
    elif isinstance(expn,Add) or isinstance(expn,Mul): #type(expn,'+' or type(expn,'*')
        m1 = expnType(expn.args[0])
        m2 = expnType(list(expn.args[1:]))
        return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
    elif is_elementary_function(expn.func): #ElementaryFunctionQ(op(0,expn))
        return max(3,expnType(expn.args[0])) #max(3,ExpnType(op(1,expn)))
    elif is_special_function(expn.func): #SpecialFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(4,m1) #max(4,apply(max,map(ExpnType,[op(expn)])))
    elif is_hypergeometric_function(expn.func): #HypergeometricFunctionQ(op(0,expn))
        m1 = max(map(expnType, list(expn.args)))
        return max(5,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif is_appell_function(expn.func):
        m1 = max(map(expnType, list(expn.args)))
        return max(6,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
    elif isinstance(expn,RootSum):
        m1 = max(map(expnType, list(expn.args))) #Apply[Max,Append[Map[ExpnType,Apply[List,expn]],7]],
        return max(7,m1)
    elif str(expn).find("Integral") != -1:

```

```

    m1 = max(map(expnType, list(expn.args)))
    return max(8,m1)  #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    #print ("Enter grade_antiderivative for sagemath")
    #print("Enter grade_antiderivative, result=",result," optimal=",optimal)

    leaf_count_result = leaf_count(result)
    leaf_count_optimal = leaf_count(optimal)

    #print("leaf_count_result=",leaf_count_result)
    #print("leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

    if str(result).find("Integral") != -1:
        grade = "F"
        grade_annotation = ""
    else:
        if expnType_result <= expnType_optimal:
            if result.has(I):
                if optimal.has(I): #both result and optimal complex
                    if leaf_count_result <= 2*leaf_count_optimal:
                        grade = "A"
                        grade_annotation = ""
                    else:
                        grade = "B"
                        grade_annotation = "Both result and optimal contain complex but leaf count of result is larger"
                else: #result contains complex but optimal is not
                    grade = "C"
                    grade_annotation = "Result contains complex when optimal does not."
            else: # result do not contain complex, this assumes optimal do not as well
                if leaf_count_result <= 2*leaf_count_optimal:
                    grade = "A"
                    grade_annotation = ""
                else:
                    grade = "B"
                    grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_count_result)-str(leaf_count_optimal)
            else:
                grade = "C"
                grade_annotation = "Result contains higher order function than in optimal. Order "+str(ExpnType_result)-str(ExpnType_optimal)

```

```

# print("Before returning. grade=", grade, " grade_annotation=", grade_annotation)

return grade, grade_annotation

```

4.2.4 SageMath grading function

```

# Dec 24, 2019. Nasser: Ported original Maple grading function by
#       Albert Rich to use with Sagemath. This is used to
#       grade Fracas, Giac and Maxima results.
# Dec 24, 2019. Nasser: Added 'exp_integral_e' and 'sng', 'sin_integral'
#       'arctan2', 'floor', 'abs', 'log_integral'
# June 4, 2022 Made default grade_annotation "none" instead of "" due
#       issue later when reading the file.
# July 14, 2022. Added ellipticF. This is until they fix sagemath, then remove it.

from sage.all import *
from sage.symbolic.operators import add_vararg, mul_vararg

debug=False;

def tree_size(expr):
    r"""
    Return the tree size of this expression.
    """
    # print("Enter tree_size, expr is ", expr)

    if expr not in SR:
        # deal with lists, tuples, vectors
        return 1 + sum(tree_size(a) for a in expr)
    expr = SR(expr)
    x, aa = expr.operator(), expr.operands()
    if x is None:
        return 1
    else:
        return 1 + sum(tree_size(a) for a in aa)

def is_sqrt(expr):
    if expr.operator() == operator.pow: # isinstance(expr, Pow):
        if expr.operands()[1] == 1/2: # expr.args[1] == Rational(1,2):
            if debug: print("expr is sqrt")
            return True
        else:
            return False
    else:
        return False

```

```

def is_elementary_function(func):
    #debug=False
    m = func.name() in ['exp','log','ln',
        'sin','cos','tan','cot','sec','csc',
        'arcsin','arccos','arctan','arccot','arcsec','arccsc',
        'sinh','cosh','tanh','coth','sech','csch',
        'arcsinh','arccosh','arctanh','arcoth','arcsech','arccsch','sgn',
        'arctan2','floor','abs'
    ]
    if debug:
        if m:
            print ("func ", func , " is elementary_function")
        else:
            print ("func ", func , " is NOT elementary_function")

    return m

def is_special_function(func):
    #debug=False
    if debug:
        print ("type(func)=", type(func))

    m= func.name() in ['erf','erfc','erfi','fresnel_sin','fresnel_cos','Ei',
        'Ei','Li','Si','sin_integral','Ci','cos_integral','Shi','sinh_integral',
        'Chi','cosh_integral','gamma','log_gamma','psi,zeta',
        'polylog','lambert_w','elliptic_f','elliptic_e','ellipticF',
        'elliptic_pi','exp_integral_e','log_integral']

    if debug:
        print ("m=",m)
        if m:
            print ("func ", func , " is special_function")
        else:
            print ("func ", func , " is NOT special_function")

    return m

def is_hypergeometric_function(func):
    return func.name() in ['hypergeometric','hypergeometric_M','hypergeometric_U']

def is_appell_function(func):
    return func.name() in ['hypergeometric'] #[appellf1] can't find this in sagemath

```



```

def is_atom(expn):

    #debug=False
    if debug:
        print ("Enter is_atom, expn=",expn)

    if not hasattr(expn, 'parent'):
        return False

    #thanks to answer at https://ask.sagemath.org/question/49179/what-is-sagemath-equivalent-to-atomic-try:
    if expn.parent() is SR:
        return expn.operator() is None
    if expn.parent() in (ZZ, QQ, AA, QQbar):
        return expn in expn.parent() # Should always return True
    if hasattr(expn.parent(), "base_ring") and hasattr(expn.parent(), "gens"):
        return expn in expn.parent().base_ring() or expn in expn.parent().gens()

    return False

except AttributeError as error:
    print("Exception,AttributeError in is_atom")
    print ("caught exception" , type(error).__name__ )
    return False

def expnType(expn):

    if debug:
        print (">>>>>Enter expnType, expn=", expn)
        print (">>>>>is_atom(expn)=", is_atom(expn))

    if is_atom(expn):
        return 1
    elif type(expn)==list: #isinstance(expn,list):
        return max(map(expnType, expn)) #apply(max,map(ExpnType,expn))
    elif is_sqrt(expn):
        if type(expn.operands()[0])==Rational: #type(isinstance(expn.args[0],Rational):
            return 1
        else:
            return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
    elif expn.operator() == operator.pow: #isinstance(expn,Pow)
        if type(expn.operands()[1])==Integer: #isinstance(expn.args[1],Integer)
            return expnType(expn.operands()[0]) #expnType(expn.args[0])
        elif type(expn.operands()[1])==Rational: #isinstance(expn.args[1],Rational)
            if type(expn.operands()[0])==Rational: #isinstance(expn.args[0],Rational)

```

```

    return 1
  else:
    return max(2,expnType(expn.operands()[0])) #max(2,expnType(expn.args[0]))
  else:
    return max(3,expnType(expn.operands()[0]),expnType(expn.operands()[1])) #max(3,expnType(expn.op
elif expn.operator() == add_vararg or expn.operator() == mul_vararg: #isinstance(expn,Add) or instan
    m1 = expnType(expn.operands()[0]) #expnType(expn.args[0])
    m2 = expnType(expn.operands()[1:]) #expnType(list(expn.args[1:]))
    return max(m1,m2) #max(ExpnType(op(1,expn)),max(ExpnType(rest(expn))))
elif is_elementary_function(expn.operator()): #is_elementary_function(expn.func)
    return max(3,expnType(expn.operands()[0]))
elif is_special_function(expn.operator()): #is_special_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(4,m1) #max(4,m1)
elif is_hypergeometric_function(expn.operator()): #is_hypergeometric_function(expn.func)
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(5,m1) #max(5,m1)
elif is_appell_function(expn.operator()):
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(6,m1) #max(6,m1)
elif str(expn).find("Integral") != -1: #this will never happen, since it
    #is checked before calling the grading function that is passed.
    #but kept it here.
    m1 = max(map(expnType, expn.operands())) #max(map(expnType, list(expn.args)))
    return max(8,m1) #max(5,apply(max,map(ExpnType,[op(expn)])))
else:
    return 9

#main function
def grade_antiderivative(result,optimal):

    if debug:
        print ("Enter grade_antiderivative for sagemath")
        print("Enter grade_antiderivative, result=",result)
        print("Enter grade_antiderivative, optimal=",optimal)
        print("type(anti)=",type(result))
        print("type(optimal)=",type(optimal))

    leaf_count_result = tree_size(result) #leaf_count(result)
    leaf_count_optimal = tree_size(optimal) #leaf_count(optimal)

    #if debug: print ("leaf_count_result=", leaf_count_result, "leaf_count_optimal=",leaf_count_optimal)

    expnType_result = expnType(result)
    expnType_optimal = expnType(optimal)

```

```

if debug: print ("expnType_result=", expnType_result, "expnType_optimal=",expnType_optimal)

if expnType_result <= expnType_optimal:
    if result.has(I):
        if optimal.has(I): #both result and optimal complex
            if leaf_count_result <= 2*leaf_count_optimal:
                grade = "A"
                grade_annotation = "none"
            else:
                grade = "B"
                grade_annotation = "Both result and optimal contain complex but leaf count of result is larger t
        else: #result contains complex but optimal is not
            grade = "C"
            grade_annotation = "Result contains complex when optimal does not."
    else: # result do not contain complex, this assumes optimal do not as well
        if leaf_count_result <= 2*leaf_count_optimal:
            grade = "A"
            grade_annotation = "none"
        else:
            grade = "B"
            grade_annotation = "Leaf count of result is larger than twice the leaf count of optimal. "+str(leaf_
else:
    grade = "C"
    grade_annotation = "Result contains higher order function than in optimal. Order "+str(expnType_resu

print("Before returning. grade=",grade, " grade_annotation=",grade_annotation)

return grade, grade_annotation

```